

ANTI-SORTILIN ANTIBODIES AND METHODS OF USE THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 62/698,007, filed July 13, 2018, U.S. Provisional Application No. 62/860,184, filed June 11, 2019, and U.S. Provisional Application No. 62/868,849, filed June 28, 2019, each of which is hereby incorporated by reference in its entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0002] This invention was made with government support under Award Number R44AG050363 awarded by the National Institutes of Health. The government has certain rights in the invention.

SUBMISSION OF SEQUENCE LISTING ON ASCII TEXT FILE

[0003] The content of the following submission on ASCII text file is incorporated herein by reference in its entirety: a computer readable form (CRF) of the Sequence Listing (file name: 735022001940SEQLIST.TXT, date recorded: July 10, 2019, size: 153 KB).

FIELD

[0004] This present disclosure relates to anti-Sortilin antibodies, and therapeutic uses of such antibodies.

BACKGROUND

[0005] Sortilin is a Type I transmembrane protein that acts both as a receptor of several ligands, and in the sorting of select cargo from the trans-Golgi network (TGN) to late endosomes and lysosomes for degradation. Sortilin harbors a large extracellular domain that is part of the VPS10 family, homologous to yeast VPS10P, and contains of a 10-blade beta-propeller structure and a cysteine-rich 10CC module (Nykjaer, A *et al.*, (2012) *Trends Neurosci* 35: 261-270; and Zheng, Y *et al.*, (2011) *PLoS One* 6: e21023). A small fraction of Sortilin may be shed by activity of ADAM10 or gamma-Secretase (<5%) (Nykjaer, A *et al.*, (2012) *Trends Neurosci* 35: 261-270; and Willnow, TE *et al.*, (2011) *Curr Opin Lipidol* 22: 79-85).

[0006] Sortilin binds the secreted protein Progranulin (PGRN) and targets it for lysosomal degradation, thus negatively regulating extracellular levels of PGRN (Hu, F *et al.* (2010) *Neuron* 68, 654-667. In line with this, deficiency of Sortilin significantly increases plasma PGRN levels both in mouse models *in vivo* and human cells *in vitro* (Carrasquillo, M.M *et al.*, (2010) *Am J Hum Genet* 87, 890-897; Lee, W.C *et al.*, (2014) *Hum Mol Genet* 23, 1467-1478). Moreover, a polymorphism in Sortilin was

shown to be strongly associated with PGRN serum levels in humans (Carrasquillo MM *et al.*, (2010), *Am J Hum Genet.* 10; 87(6):890-7).

[0007] Progranulin (PGRN) is a secreted, growth factor-like, trophic, and anti-inflammatory protein, which also plays a role as an adipokine involved in diet-induced obesity and insulin resistance (Nguyen DA *et al.*, (2013). *Trends in Endocrinology and Metabolism*, 24, 597- 606). Progranulin deficiency accounts for roughly 25% of all heritable forms of frontotemporal dementia (FTD), an early-onset neurodegenerative disease. Patients with heterozygous loss-of-function mutations in PGRN have ~50% reduced extracellular levels of the protein and they will invariably develop FTD, making PGRN a causal gene for the disease (Baker, M *et al.*, (2006) *Nature* 442, 916-919; Carecchio M *et al.*, (2011) *J Alzheimers Dis* 27, 781-790; Cruts, M *et al.*, (2008) *Trends Genet* 24, 186-194; Galimberti, D *et al.*, (2010) *J Alzheimers Dis* 19, 171-177). In addition, PGRN mutant alleles have been identified in Alzheimer's disease patients (Seelaar, H *et al.*, (2011). *Journal of neurology, neurosurgery, and psychiatry* 82, 476-486). Importantly, PGRN acts protectively in several disease models with increased PGRN levels, accelerating behavioral recovery from ischemia (Tao, J *et al.*, (2012) *Brain Res* 1436, 130-136; Egashira, Y. *et al.*, (2013). *J Neuroinflammation* 10, 105), suppressing locomotor deficits in a Parkinson's disease model (Van Kampen, J.M *et al.* (2014). *PLoS One* 9, e97032), attenuating pathology in a model of amyotrophic lateral sclerosis (Laird, A.S *et al.*, (2010). *PLoS One* 5, e13368.) and arthritis (Tang, W *et al.*, (2011). *Science* 332, 478-484) and preventing memory deficits in an Alzheimer's disease model (Minami, S.S *et al.*, (2014). *Nat Med* 20, 1157-1164).

[0008] Sortilin also binds directly to pro-neurotrophins, such as pro-nerve growth factor (pro-NGF), pro-BDNF, pro-neurotrophin-3, *etc.*, which harbor a pro-domain and are typically pro-apoptotic. Such pro-neurotrophin precursors are released during stress, and Sortilin is involved in regulating their release as well as binding on the receiving cell and stimulation of apoptosis in conjunction with p75NTR (Willnow, TE *et al.*, (2008) *Nat Rev Neurosci* 9: 899-909; Nykjaer, A *et al.*, *Trends Neurosci* 35: 261-270; and Nykjaer, A *et al.*, (2004) *Nature* 427: 843-848; Hiroko Yano *et al.*, (2009) *J Neurosci.*; 29: 14790–14802. Teng H.K., *et al.*, *J. Neurosci.* 25:5455-5463(2005)). Sortilin also binds to p75NTR directly (Skeldal S *et al.*, (2012) *J Biol Chem.*; 287:43798). Sortilin also binds to neurotensin in a region that partially overlaps with Progranulin binding (Quistgaard, EM *et al.*, (2009) *Nat Struct Mol Biol* 16: 96-98; and Zheng, Y *et al.*, *PLoS One* 6: e21023). Sortilin also interacts with the Trk receptors NTRK1, NTRK2, and NTRK3; and can regulate their anterograde axonal transport and signaling (Vaegter, CB *et al.*, (2011) *Nat. Neurosci.* 14:54-61). Sortilin also interacts with and regulates the processing and trafficking of amyloid precursor protein and the resulting production of pathological beta amyloid peptides (Gustafsen C *et al.*, (2013). *J Neurosci.* 2;33(1):64-71).

[0009] Sortilin has also been shown to bind to apolipoproteins and lipoprotein lipase; thus, deficiency leads to reduced VLDL release from liver and reduced cholesterol (Willnow, TE *et al.*, (2011) *Curr Opin Lipidol* 22: 79-85; Kjolby, M *et al.*, (2010) *Cell Metab* 12: 213-223; Nilsson, SK *et al.*, (2007) *Biochemistry* 46: 3896-3904.; Nilsson, SK *et al.*, (2008) *J Biol Chem* 283: 25920-25927; and Klinger, SC *et al.*, (2011) *J Cell Sci* 124: 1095-1105). Recently, Sortilin has also been implicated in binding to APP directly (Gustafsen, C *et al.*, (2013) *J. Neurosc.* 33:64-71) and also to the APP processing enzyme BACE1 (Gustafsen, C *et al.*, (2013) *J. Neurosc.* 33:64-71; and Finan, GM *et al.*, *J Biol Chem* 286: 12602-12616). Sortilin also binds to apolipoprotein E (APOE), to the A beta peptide (Carlo, AS *et al.*, (2013) *J. Neurosc.* 33: 358-370), and to PCSK9 (Gustafsen *et al.*, (2014) *Cell Metab*, 19: 310-318). Sortilin has also been shown to bind to and regulate extracellular levels of PCSK9, which directs low-density lipoprotein receptor for degradation in lysosomes, resulting in increased levels of LDL cholesterol (Gustafsen C *et al.*, (2014). *Cell Metab.* 2014 Feb 4;19(2):310-8).

[0010] When present at intracellular vesicles such as endosomes, the amino-terminal extracellular domain of Sortilin is directed towards the lumen, where cargo of the vesicle is present. The carboxy-terminal intracellular/cytoplasmic domain of Sortilin, however, binds to a series of adaptor proteins, which regulate its trafficking from the surface and within intracellular compartments. These include AP2 (a clathrin adaptor to modulate endocytosis from the cell surface), and the Retromer Complex/API1, which modulate movement from early endosomes to Golgi for recycling; and interaction with GGA (Golgi-localizing, gamma-ear containing, ADP-ribosylation factor binding) family proteins for movement from Golgi directly to early endosomes, usually for subsequent degradation through lysosomes. Thus, Sortilin can bind to ligands at its luminal domain, while engaging the cytoplasmic adaptors that determine its destination to determine intracellular fates, such as degradation for Progranulin and other factors.

[0011] Through its various interactions with proteins, such as Progranulin, Sortilin and its multiple ligands have been shown to be involved in various diseases, disorders, and conditions, such as frontotemporal dementia, amyotrophic lateral sclerosis, amyotrophic lateral sclerosis-frontotemporal dementia phenotypes, Alzheimer's disease, Parkinson's disease, depression, neuropsychiatric disorders, vascular dementia, seizures, retinal dystrophy, age related macular degeneration, glaucoma, traumatic brain injury, aging, seizures, wound healing, stroke, arthritis, and atherosclerotic vascular diseases.

[0012] Accordingly, there is a need for therapeutic antibodies that specifically bind Sortilin proteins and block the binding of Sortilin to its ligands, such as Progranulin, or otherwise modulate the effective concentration of the ligands, in order to treat one or more diseases, disorders, and conditions associated with Sortilin activity.

[0013] All references cited herein, including patents, patent applications and publications, are hereby incorporated by reference in their entirety.

SUMMARY

[0014] The present disclosure is generally directed to compositions that include antibodies, *e.g.*, monoclonal, chimeric, humanized antibodies, antibody fragments, *etc.*, that specifically bind human Sortilin, and to methods of using such compositions.

[0015] In some examples, the present disclosure provides an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region comprises an HVR-H1 comprising SEQ ID NO: 1, an HVR-H2 comprising SEQ ID NO: 2 or SEQ ID NO:3, and an HVR-H3 comprising SEQ ID NO: 5 or SEQ ID NO: 6; and the light chain variable region comprises an HVR-L1 comprising any of SEQ ID NOs: 8-27, an HVR-L2 comprising SEQ ID NO: 29 or SEQ ID NO: 30, and an HVR-L3 comprising SEQ ID NO: 32.

[0016] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYW (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIQQGYGMDV (SEQ ID NO: 5); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0017] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYW (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIQQGYGMDV (SEQ ID NO: 5); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), the HVR-L2 comprising the amino acid sequence LGSNRVS (SEQ ID NO: 30), and the HVR-L3 comprising the amino acid sequence MQQQETPLT (SEQ ID NO: 33).

[0018] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYW (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLES (SEQ ID NO: 3), the HVR-H3 comprising the amino acid sequence ARQGSIQQGYGMDV (SEQ ID NO: 5); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), the HVR-L2 comprising the amino acid sequence LGSNRAS

(SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0019] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0020] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSTGYNYLD (SEQ ID NO: 9), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0021] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSTGYNYLD (SEQ ID NO: 10), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0022] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSTGYNYLD (SEQ ID NO: 21), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0023] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIKQGYYGMDV (SEQ ID NO: 6); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQETPLT (SEQ ID NO: 33).

[0024] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIQQGYYGMDV (SEQ ID NO: 5); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLHSNGYNYLD (SEQ ID NO: 26), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQETPLT (SEQ ID NO: 33).

[0025] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIKQGYYGMDV (SEQ ID NO: 6); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQGLLSNGYNYLD (SEQ ID NO: 27), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0026] In some examples, the present disclosure provides an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), and an HVR-H3 comprising the amino acid sequence ARQGSIKQGYYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0027] In some examples, the present disclosure provides an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein the antibody comprises a heavy chain variable region with an HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), and an HVR-H3 comprising the amino acid sequence ARQGSIQGYYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSTGYNYLD (SEQ ID NO: 9), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0027a] In one aspect, the present disclosure refers to an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein

(a) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence

YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence

TIYHSGSTYYNPSLKS (SEQ ID NO: 2), an HVR-H3 comprising the amino acid sequence

ARQGSIQGYYGMDV (SEQ ID NO: 5); and the light chain variable region comprises an HVR-L1

comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), an HVR-L2

comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the

amino acid sequence MQQQEAPLT (SEQ ID NO: 32);

(b) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence

YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence

TIYHSGSTYYNPSLKS (SEQ ID NO: 2), an HVR-H3 comprising the amino acid sequence

ARQGSIQGYYGMDV (SEQ ID NO: 5); and the light chain variable region comprises an HVR-L1

comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), an HVR-L2

comprising the amino acid sequence LGSNRVS (SEQ ID NO: 30), and an HVR-L3 comprising the

amino acid sequence MQQQETPLT (SEQ ID NO: 33);

(c) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence

YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence

TIYHSGSTYYNPSLES (SEQ ID NO: 3), an HVR-H3 comprising the amino acid sequence

ARQGSIQGYYGMDV (SEQ ID NO: 5); and the light chain variable region comprises an HVR-L1

comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), an HVR-L2

comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the

amino acid sequence MQQQEAPLT (SEQ ID NO: 32);

- (d) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), an HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32);
- (e) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), an HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSTGYNYLD (SEQ ID NO: 9), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32);
- (f) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), an HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQETPLT (SEQ ID NO: 33);
- (g) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), an HVR-H3 comprising the amino acid sequence ARQGSIQGYGMDV (SEQ ID NO: 5); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLHSNGYNYLD (SEQ ID NO: 26), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQETPLT (SEQ ID NO: 33);
- (h) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), an HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQGLLSNGYNYLD (SEQ ID NO: 27), an HVR-L2

comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32);

- (i) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYWYG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), an HVR-H3 comprising the amino acid sequence ARQGSIKQGYYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSSGYNYLD (SEQ ID NO: 10), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32); or
- (j) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYWYG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), an HVR-H3 comprising the amino acid sequence ARQGSIKQGYYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSIGYNYLD (SEQ ID NO: 21), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0027b] In one aspect, the present disclosure refers to an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein: the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYWYG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), and an HVR-H3 comprising the amino acid sequence ARQGSIKQGYYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSTGYNYLD (SEQ ID NO: 9), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0027c] In one aspect, the present disclosure refers to an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein: the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYWYG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), and an HVR-H3 comprising the amino acid sequence ARQGSIKQGYYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), an HVR-L2 comprising

the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0028] In some examples of any of the above aspects, the anti-Sortilin antibody: (a) decreases cell surface levels of Sortilin more than the level of decrease caused by an anti-Sortilin antibody comprising a heavy chain variable region comprising the sequence of SEQ ID NO: 56 and a light chain variable region comprising the sequence of SEQ ID NO: 79; (b) increases extracellular levels of Progranulin more than the level of increase caused by an anti-Sortilin antibody comprising a heavy chain variable region comprising the sequence of SEQ ID NO: 56 and a light chain variable region comprising the sequence of SEQ ID NO: 79; (c) inhibits the interaction between Sortilin and Progranulin more than the level of inhibition caused by an anti-Sortilin antibody comprising a heavy chain variable region comprising the sequence of SEQ ID NO: 56 and a light chain variable region comprising the sequence of SEQ ID NO: 79; or (d) any combination of (a)–(c). In some examples of any of the above aspects, the anti-Sortilin antibody: (a) decreases cell surface levels of Sortilin more than the level of decrease caused by an anti-Sortilin antibody selected from the group consisting of S-60-1, S-60-2, S-60-3, S-60-4, S-60-7, or S-60-8; (b) increases extracellular levels of Progranulin more than the level of increase caused by an anti-Sortilin antibody selected from the group consisting of S-60-1, S-60-2, S-60-3, S-60-4, S-60-7, or S-60-8; (c) inhibits the interaction between Sortilin and Progranulin more than the level of inhibition caused by an anti-Sortilin antibody selected from the group consisting of S-60-1, S-60-2, S-60-3, S-60-4, S-60-7, or S-60-8; or (d) any combination of (a)–(c).

[0029] Certain examples of the present disclosure provide an anti-Sortilin antibody with a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 54-56; and/or a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 57-58, 60-78, or 80.

[0030] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57.

[0031] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 58.

[0032] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 59.

[0033] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 55, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57.

[0034] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 55, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 58.

[0035] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57.

[0036] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 77.

[0037] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 78.

[0038] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 79.

[0039] In some examples, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 80.

[0040] In some examples, the present disclosure provides an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57.

[0041] In certain examples, the present disclosure provides an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 60.

[0042] In certain examples, the present disclosure provides an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 61.

[0043] In certain examples, the present disclosure provides an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 72.

[0043a] In one aspect, the present disclosure refers to an antibody that binds to a Sortilin protein, wherein the antibody comprises:

- (a) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57;
- (b) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 58;
- (c) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 59;
- (d) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 55, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57;
- (e) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 55, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 58;
- (f) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57;
- (g) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 77;
- (h) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 78;
- (i) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 79;
- (j) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 80;
- (k) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 60;

(l) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 61; or

(m) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 72.

[0043b] In one aspect, the present disclosure refers to an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 60.

[0043c] In one aspect, the present disclosure refers to an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57.

[0043d] In one aspect, the present disclosure refers to an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 and a light chain comprising the amino acid sequence of SEQ ID NO: 142.

[0043e] In one aspect, the present disclosure refers to an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 138 and a light chain comprising the amino acid sequence of SEQ ID NO: 142.

[0044] In some examples that may be combined with any of the previous examples, an anti-Sortilin antibody of the present disclosure is of the IgG class, the IgM class, or the IgA class. In certain specific examples, the anti-Sortilin antibody is of the IgG class and has an IgG1, IgG2, IgG3, or IgG4 isotype. In some examples of the present disclosure, provided is an anti-Sortilin antibody of the IgG1, IgG2, IgG3, or IgG4 isotype, wherein (a) the antibody is an IgG1 or IgG2 isotype and the Fc region comprises an amino acid substitution at position P331S, wherein the numbering of the residue position is according to EU numbering; (b) the antibody is an IgG1 isotype and the Fc region comprises amino acid substitutions at positions L234A, L235A, and P331S, wherein the numbering of the residue position is according to EU numbering; (c) the antibody is an IgG1, IgG2, or IgG4 isotype and the Fc region comprises an amino acid substitution at position N297A, wherein the numbering of the residue position is according to EU numbering; or (d) the Fc region comprises an amino acid substitution at positions S267E and L328F, wherein the numbering of the residue position is according to EU numbering. In some examples, the antibody is an IgG1 isotype and the Fc region comprises amino acid substitutions at positions L234A, L235A, and P331S, wherein the numbering of the residue position is according to EU numbering.

[0045] In some examples, the present disclosure provides an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 and a light chain comprising the amino acid sequence of SEQ ID NO: 142.

[0046] In some examples, the present disclosure provides an antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 138 and a light chain comprising the amino acid sequence of SEQ ID NO: 142.

[0047] In certain examples of the present disclosure, the Sortilin protein is a human protein. In some examples, the Sortilin protein is a wild-type protein. In certain examples, the Sortilin protein is a naturally occurring variant. In some examples, an anti-Sortilin antibody of the present disclosure binds specifically to a human Sortilin protein. In some examples that may be combined with any of the previous examples, the antibody is a human antibody, a bispecific antibody, a monoclonal antibody, a multivalent antibody, a conjugated antibody, or a chimeric antibody. In certain examples, an anti-Sortilin antibody of the present disclosure is a monoclonal antibody.

[0048] In some examples, an anti-Sortilin antibody of the present disclosure is a bispecific antibody recognizing a first antigen and a second antigen. In certain examples, bispecific antibodies of the present disclosure recognize both Sortilin and an antigen facilitating transport across the blood-brain-barrier. In some specific examples, the first antigen is Sortilin and the second antigen is Sortilin, transferrin receptor (TR), insulin receptor (HIR), insulin-like growth factor receptor (IGFR), low-density lipoprotein receptor related proteins 1 and 2 (LPR-1 and 2), diphtheria toxin receptor, CRM197, a llama single domain antibody, TMEM 30(A), a protein transduction domain, TAT, Syn-B, penetratin, a poly-arginine peptide, an angiopep peptide, basigin, Glut1, CD98hc, or ANG1005.

[0049] In certain examples of the present disclosure, the anti-Sortilin antibody is an antibody fragment that binds to a human Sortilin protein. In some examples, the anti-Sortilin antibody is an antibody fragment that binds to one or more human proteins selected from the group consisting of human Sortilin, a naturally occurring variant of human Sortilin, and a disease variant of human Sortilin. In certain examples that may be combined with any of the preceding examples, the anti-Sortilin antibody fragment is an Fab, Fab', Fab'-SH, F(ab')₂, Fv, or scFv fragment.

[0050] In certain examples, the present disclosure provides an anti-Sortilin antibody, wherein (a) the anti-Sortilin antibody has increases extracellular levels of Progranulin, decreases cellular levels of Sortilin, inhibits the interaction between Sortilin and Progranulin, or any combination thereof; (b) the anti-Sortilin antibody decreases cell surface levels of Sortilin, increases extracellular levels of Progranulin, inhibits interaction between Sortilin and Progranulin, or any combination thereof; (c) the anti-Sortilin antibody decreases cell surface levels of Sortilin, decreases intracellular levels of Sortilin, decreases total levels of Sortilin, or any combination thereof; (d) the anti-Sortilin antibody induces Sortilin degradation, Sortilin cleavage, Sortilin internalization, Sortilin down regulation, or any combination thereof; (e) the anti-Sortilin antibody decreases cellular levels of Sortilin and inhibits the interaction between Sortilin and Progranulin; (f) the anti-Sortilin antibody decreases cellular levels of Sortilin and increases cellular levels of Progranulin; and/or (g) the anti-Sortilin antibody increases the effective concentration of Progranulin.

[0051] In certain examples, the present disclosure provides an anti-Sortilin antibody, wherein the anti-Sortilin antibody decreases cell surface levels of Sortilin, increases extracellular levels of Progranulin, inhibits interaction between Sortilin and Progranulin, or any combination thereof. In some examples, the anti-Sortilin antibody has a dissociation constant (K_D) for human Sortilin that is up to 2.5-fold lower than an anti-Sortilin antibody comprising a heavy chain variable region comprising the sequence of SEQ ID NO: 56 and a light chain variable region comprising the sequence of SEQ ID NO: 79, wherein the K_D is determined by FACS (*see, e.g., Example 1*). In some examples, the antibody has a dissociation constant (K_D) for human Sortilin that is greater than 1- and up to about 2.1-fold lower

than an anti-Sortilin antibody comprising a heavy chain variable region comprising the sequence of SEQ ID NO: 56 and a light chain variable region comprising the sequence of SEQ ID NO: 79, wherein the K_D is determined by FACS. In some examples, the anti-Sortilin antibody has a dissociation constant (K_D) for human Sortilin that ranges from about $1.10E-8$ M to about $4.68E-10$ M wherein the K_D is determined by FACS (*see, e.g.*, **Example 1**), or about 270 to about 2910 pM wherein the K_D is determined by Bio-layer interferometry (*see, e.g.*, **Example 4**). In some examples, the antibody has a dissociation constant (K_D) for human Sortilin that ranges from about $5.0E-10$ M to about $1.0E-9$ M wherein the K_D is determined by FACS, or about 250-500 pM wherein the K_D is determined by Bio-layer interferometry.

[0052] In some examples that may be combined with any of the previous examples, an anti-Sortilin antibody of the present disclosure (a) reduces cell surface levels of Sortilin with a half maximal effective concentration (EC_{50}) that is less than 150 pM, as measured by flow cytometry; (b) reduces cell surface levels of Sortilin by more than about 50% at 1.25 nM IgG, by more than about 80% at 0.63 nM IgG, or by more than about 69% at 150 nM IgG relative to control, as measured by flow cytometry; increases Progranulin secretion by more than about 1.13 fold over control at 0.63 nM IgG, or by more than about 1.22 fold over control at 50 nM IgG, as measured by standard ELISA; blocks binding of Progranulin to Sortilin with a half maximal effective concentration (EC_{50}) that is less than .325 nM, as measured by flow cytometry; (c) blocks binding of Progranulin to Sortilin by more than about 88% at 50 nM IgG, or by more than about 27.5% at 150 nM IgG relative to control, as measured by flow cytometry; or (f) any combination thereof.

[0053] In some examples, an anti-Sortilin antibody of the present disclosure (a) reduces cell surface levels of Sortilin with a half maximal effective concentration (EC_{50}) that is less than 681 pM, as measured by flow cytometry; (b) reduces cell surface levels of Sortilin by more than about 40% at 1.25 nM IgG, by more than about 29% at 0.6 nM IgG, or by more than about 62% at 150 nM IgG relative to control, as measured by flow cytometry; (c) increases Progranulin secretion by more than about 1.11 fold over control at 0.63 nM IgG, or by more than about 1.75 fold over control at 50 nM IgG, as measured by standard ELISA; (d) blocks binding of Progranulin to Sortilin with a half maximal effective concentration (EC_{50}) that is less than 0.751 nM, as measured by flow cytometry; (e) blocks binding of Progranulin to Sortilin by more than about 90% at 50 nM IgG, or by more than about 95% at 150 nM IgG relative to control, as measured by flow cytometry; or (f) any combination thereof.

[0054] In some examples, an anti-Sortilin antibody of the present disclosure competes with an antibody comprising the heavy chain variable domain and the light chain variable domain of an antibody selected from the group consisting of S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-16, S-60-18, S-60-19, S-60-24, and any combination thereof for binding to Sortilin. In some examples that may be combined with any of the previous examples, an anti-

Sortilin antibody of the present disclosure binds essentially the same Sortilin epitope as an antibody comprising the heavy chain variable domain and the light chain variable domain of an antibody selected from the group consisting of S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-16, S-60-18, S-60-19, and S-60-24.

[0055] In another example, the present disclosure relates to an antibody produced by any of the methods described herein. In some examples, the present disclosure provides an isolated nucleic acid comprising a nucleic acid sequence encoding an anti-Sortilin antibody of any one of the preceding aspects and examples. In one aspect, the present disclosure refers to an isolated nucleic acid comprising a nucleic acid sequence encoding any one of the antibodies disclosed herein. Accordingly, some examples provide a vector containing a nucleic acid encoding an anti-Sortilin antibody, and some examples provide an isolated host cell containing such a vector. In one aspect, the present disclosure refers to a vector comprising the nucleic acid disclosed herein. In one aspect, the present disclosure refers to an isolated host cell comprising the vector disclosed herein. In certain examples, the present disclosure provides a method of producing an anti-Sortilin antibody that binds to Sortilin, comprising culturing such an isolated host cell so that the anti-Sortilin antibody is produced. In one aspect, the present disclosure refers to a method of producing an antibody that binds to Sortilin, comprising culturing the host cell disclosed herein so that the antibody is produced. In certain examples, the method further comprises recovering the antibody produced by the cell. In some examples, provided herein is an isolated antibody that binds to Sortilin and is produced by a method of the present disclosure. In one aspect, the present disclosure refers to an isolated antibody that binds to Sortilin produced by the method disclosed herein.

[0056] In another example, the present disclosure relates to a pharmaceutical composition comprising any of the anti-Sortilin antibodies described herein and a pharmaceutically acceptable carrier. In one aspect, the present disclosure refers to a pharmaceutical composition comprising any one of the antibodies disclosed herein and a pharmaceutically acceptable carrier. In some examples, the present disclosure relates to the use of any of the antibodies or compositions described herein for the preparation or manufacture of a medicament. In some examples, the present disclosure relates to the use of any of the antibodies or compositions described herein for the preparation or manufacture of a medicament for the treatment of frontotemporal dementia, progressive supranuclear palsy, Alzheimer's disease, vascular dementia, seizures, retinal dystrophy, amyotrophic lateral sclerosis, traumatic brain injury, a spinal cord injury, neuropathic pain, dementia, stroke, Parkinson's disease, acute disseminated encephalomyelitis, retinal degeneration, age related macular degeneration, glaucoma, multiple sclerosis, septic shock, bacterial infection, arthritis, or osteoarthritis. Some examples provide an antibody or composition described herein for use in a method of treatment. Some examples provide an antibody or composition described herein for use in a method of treatment of frontotemporal dementia, progressive supranuclear palsy, Alzheimer's disease, vascular dementia, seizures, retinal dystrophy, amyotrophic lateral sclerosis, traumatic brain injury, a spinal cord injury, neuropathic pain, dementia, stroke, Parkinson's disease, acute disseminated encephalomyelitis, retinal degeneration, age related macular degeneration, glaucoma, multiple sclerosis, septic shock, bacterial infection, arthritis, or osteoarthritis.

[0057] In certain examples, provided herein is a method of preventing, reducing risk for, or treating an individual having a disease, disorder, or injury, comprising administering to an individual in need thereof a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure. In one aspect, the present disclosure refers to use of any one the antibodies disclosed herein or the pharmaceutical composition disclosed herein in the manufacture of a medicament for preventing, reducing risk for, or treating a disease, disorder, or injury in an individual. In some

examples, the disease, disorder or injury is selected from the group consisting of frontotemporal dementia, progressive supranuclear palsy, Alzheimer's disease, vascular dementia, seizures, retinal dystrophy, amyotrophic lateral sclerosis, traumatic brain injury, a spinal cord injury, neuropathic pain, dementia, stroke, Parkinson's disease, acute disseminated encephalomyelitis, retinal degeneration, age related macular degeneration, glaucoma, multiple sclerosis, septic shock, bacterial infection, arthritis, and osteoarthritis. In some examples, the disease, disorder, or injury is selected from frontotemporal dementia, progressive supranuclear palsy, Alzheimer's disease, and amyotrophic lateral sclerosis.

[0058] In certain examples, provided herein is a method of inhibiting one or more of neuroinflammation, axonopathy characterized by short axonal outgrowth and aberrant branching, microglial activation, and inflammatory response, comprising administering to the individual a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure. In one aspect, the present disclosure refers to use of any one the antibodies disclosed herein or the pharmaceutical composition disclosed herein in the manufacture of a medicament for inhibiting one or more of neuroinflammation, axonopathy characterized by short axonal outgrowth and aberrant branching, aberrant microglial activation, and inflammatory response in an individual in need thereof.

[0059] In certain examples, provided herein is a method of promoting one or more of wound healing, autophagy, and clearance of aggregate proteins, comprising administering to the individual a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure. In one aspect, the present disclosure refers to use of any one the antibodies disclosed herein or the pharmaceutical composition disclosed herein in the manufacture of a medicament for promoting one or more of wound healing, autophagy, and clearance of aggregate proteins in an individual in need thereof.

[0060] In certain examples, provided herein is a method of preventing, reducing risk, or treating an individual having arthritis, comprising administering to the individual a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure. In one aspect, the present disclosure refers to use of any one the antibodies disclosed herein or the pharmaceutical composition disclosed herein in the manufacture of a medicament for preventing, reducing risk, or treating arthritis in an individual in need thereof.

[0061] In certain examples, provided herein is a method of decreasing expression of one or more pro-inflammatory mediators, comprising administering to the individual a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure. In one aspect, the present disclosure refers to use of any one the antibodies disclosed herein or the pharmaceutical composition disclosed herein in the manufacture of a medicament for decreasing expression of one or more pro-inflammatory mediators in an individual in need thereof. In some examples, the one or more pro-inflammatory mediators are selected from the group consisting of IL-6, IL12p70, IL12p40, IL-1 β , TNF- α , CXCL1, CCL2, CCL3, CCL4, and CCL5.

[0062] In some examples, provided herein is a method of inhibiting the interaction between Sortilin and Progranulin, the method comprising exposing a cell expressing Sortilin to an anti-Sortilin antibody or a pharmaceutical composition of the present disclosure. In one aspect, the present disclosure refers to a method of inhibiting the interaction between Sortilin expressed on a cell and Progranulin, the method comprising exposing a cell expressing Sortilin on the cell surface to any one of the antibodies disclosed herein or the pharmaceutical composition disclosed herein, wherein the cell is *in vitro*. In some examples, the cell is *in vitro*. In some examples, the cell is *in vivo*. In some examples, the method further comprises decreasing the level of Sortilin expressed on the cell surface. In some examples, extracellular levels of Progranulin are increased.

[0063] In some examples, provided herein is a method of increasing levels of Progranulin in an individual in need thereof, the method comprising administering to the individual a therapeutically effective amount of the anti-Sortilin antibody or a pharmaceutical composition of the present disclosure. In one aspect, the present disclosure refers to use of any one the antibodies disclosed herein or the pharmaceutical composition disclosed herein in the manufacture of a medicament for increasing levels of Progranulin in an individual in need thereof. In some examples, levels of Progranulin are increased in plasma. In some examples, levels of Progranulin are increased in cerebrospinal fluid. In some examples, provided herein is a method of decreasing levels of Sortilin in an individual in need thereof, the method comprising administering to the

individual a therapeutically effective amount of an anti-Sortilin antibody or a pharmaceutical composition of the present disclosure. In one aspect, the present disclosure refers to use of any one the antibodies disclosed herein or the pharmaceutical composition disclosed herein in the manufacture of a medicament for decreasing levels of Sortilin in an individual in need thereof. In some examples, the levels of Sortilin are decreased in peripheral white blood cells. In some examples, the individual has one or more mutations in the gene encoding Progranulin. In some the individual is heterozygous for one or more loss-of-function mutations in the gene encoding Progranulin. In some examples, the individual has a c9orf72 hexanucleotide repeat expansion.the individual has or is at risk for frontotemporal dementia, Alzheimer's disease, or amyotrophic lateral sclerosis.

[0064] In some examples that may be combined with any of the previous aspects or examples, the present disclosure provides a method, wherein the anti-Sortilin antibody comprises two or more anti-Sortilin antibodies.

BRIEF DESCRIPTION OF THE DRAWINGS

[0065] **FIGS. 1A-1C** provide pharmacokinetic and pharmacodynamic studies of non-human primates administered single doses of anti-Sortilin antibody S-60-15.1 [N33T] LALAPS (S-60-15.1 [N33T] with huIgG1 with L234A/L235A/P331S mutations). **FIG. 1A** provides the level of SORT1 in peripheral white blood cells as a percentage from baseline at the indicated times after treatment (hours) with the specified anti-Sortilin antibody doses. SORT1 expression decreased with all of the anti-Sortilin antibody doses tested. Higher antibody doses (60 mg/kg, 200 mg/kg) resulted in both an earlier and more prolonged decrease of SORT1 levels compared to lower anti-Sortilin antibody doses (5mg/kg, 20 mg/kg). **FIG. 1B** provides the levels of PGRN in the plasma as a percentage from baseline at the indicated times after treatment (hours) with the specified anti-Sortilin antibody doses. The levels of PGRN increased in a time- and dose-dependent manner. In particular, plasma PGRN levels increased 3- to 4-fold at C_{max} , compared to baseline levels, for all anti-Sortilin antibody doses tested and remained elevated for longer periods of time at the higher antibody doses. **FIG. 1C** provides the levels of PGRN in CSF as a percentage from baseline at the indicated times after treatment (hours) with the specified anti-Sortilin antibody doses. CSF PGRN levels increased 2- to 3-fold above baseline in animals administered either 20 mg/kg, 60 mg/kg, or 200 mg/kg. As observed with plasma PGRN levels (**FIG. 1B**), CSF PGRN levels remained elevated over time in the higher antibody dose groups. For **FIGS. 1A-1C**, n = 3 animals per dose.

[0066] **FIGS. 2A-2C** provide pharmacokinetic and pharmacodynamic studies of non-human primates administered repeat doses of anti-Sortilin antibody S-60-15.1 [N33T] LALAPS. Animals (2 males and 2 females) were administered anti-Sortilin antibody S-60-15.1 [N33T] LALAPS at a dose of 60mg/kg once per week for four weeks. The days on which dosing occurred are represented by the

vertical dashed lines. **FIG. 2A** provides the mean ($\pm$ standard deviation) of the concentration of SORT1 in peripheral white blood cells (WBCs) as a percentage of baseline at the indicated times (days). SORT1 levels in peripheral white blood cells remained decreased throughout the duration of the study. **FIG. 2B** provides the mean ($\pm$ standard deviation) of the concentration of PGRN in plasma as a percentage of baseline (normalized) at the indicated times (days). Plasma PGRN levels increased to 5- to 6-fold above baseline at peak levels. A decrease in plasma PGRN was observed following the fourth and final administration of anti-Sortilin antibody; however, the plasma PGRN levels remained elevated by 2-fold above baseline. **FIG. 2C** provides the mean ($\pm$ standard deviation) of the concentration of PGRN in CSF as a percentage of baseline (normalized) at the indicated times (days). CSF PGRN levels were increased 3- to 4-fold above baseline (**FIG. 2C**).

DETAILED DESCRIPTION

Definitions

[0067] As used herein, the term “*preventing*” includes providing prophylaxis with respect to occurrence or recurrence of a particular disease, disorder, or condition in an individual. An individual may be predisposed to, susceptible to a particular disease, disorder, or condition, or at risk of developing such a disease, disorder, or condition, but has not yet been diagnosed with the disease, disorder, or condition.

[0068] As used herein, an individual “*at risk*” of developing a particular disease, disorder, or condition may or may not have detectable disease or symptoms of disease, and may or may not have displayed detectable disease or symptoms of disease prior to the treatment methods described herein. “At risk” denotes that an individual has one or more risk factors, which are measurable parameters that correlate with development of a particular disease, disorder, or condition, as known in the art. An individual having one or more of these risk factors has a higher probability of developing a particular disease, disorder, or condition than an individual without one or more of these risk factors.

[0069] As used herein, the term “*treatment*” refers to clinical intervention designed to alter the natural course of the individual being treated during the course of clinical pathology. Desirable effects of treatment include decreasing the rate of progression, ameliorating or palliating the pathological state, and remission or improved prognosis of a particular disease, disorder, or condition. An individual is successfully “treated”, for example, if one or more symptoms associated with a particular disease, disorder, or condition are mitigated or eliminated.

[0070] An “*effective amount*” refers to at least an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic or prophylactic result. An effective amount can be provided in one or more administrations. An effective amount herein may vary according to factors such

as the disease state, age, sex, and weight of the individual, and the ability of the treatment to elicit a desired response in the individual. An effective amount is also one in which any toxic or detrimental effects of the treatment are outweighed by the therapeutically beneficial effects. For prophylactic use, beneficial or desired results include results such as eliminating or reducing the risk, lessening the severity, or delaying the onset of the disease, including biochemical, histological and/or behavioral symptoms of the disease, its complications and intermediate pathological phenotypes presenting during development of the disease. For therapeutic use, beneficial or desired results include clinical results such as decreasing one or more symptoms resulting from the disease, increasing the quality of life of those suffering from the disease, decreasing the dose of other medications required to treat the disease, enhancing effect of another medication such as via targeting, delaying the progression of the disease, and/or prolonging survival. An effective amount of drug, compound, or pharmaceutical composition is an amount sufficient to accomplish prophylactic or therapeutic treatment either directly or indirectly. As is understood in the clinical context, an effective amount of a drug, compound, or pharmaceutical composition may or may not be achieved in conjunction with another drug, compound, or pharmaceutical composition. Thus, an “effective amount” may be considered in the context of administering one or more therapeutic agents, and a single agent may be considered to be given in an effective amount if, in conjunction with one or more other agents, a desirable result may be or is achieved

[0071] As used herein, administration “*in conjunction*” with another compound or composition includes simultaneous administration and/or administration at different times. Administration in conjunction also encompasses administration as a co-formulation or administration as separate compositions, including at different dosing frequencies or intervals, and using the same route of administration or different routes of administration.

[0072] An “*individual*” for purposes of treatment, prevention, or reduction of risk refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sport, or pet animals, such as dogs, horses, rabbits, cattle, pigs, hamsters, gerbils, mice, ferrets, rats, cats, and the like. Preferably, the individual is human.

[0073] The terms “Sortilin” or “Sortilin polypeptide” are used interchangeably herein refer herein to any native Sortilin from any mammalian source, including primates (e.g., humans and cynos) and rodents (e.g., mice and rats), unless otherwise indicated. In some embodiments, the term encompasses both wild-type sequences and naturally occurring variant sequences, e.g., splice variants or allelic variants. In some embodiments, the term encompasses “full-length,” unprocessed Sortilin as well as any form of Sortilin that results from processing in the cell. In some embodiments, the Sortilin is human Sortilin. In some embodiments, the amino acid sequence of an exemplary human Sortilin is SEQ ID NO: 81.

[0074] The terms "anti- Sortilin antibody," an "antibody that binds to Sortilin," and "antibody that specifically binds Sortilin" refer to an antibody that is capable of binding Sortilin with sufficient affinity such that the antibody is useful as a diagnostic and/or therapeutic agent in targeting Sortilin. In one embodiment, the extent of binding of an anti- Sortilin antibody to an unrelated, non- Sortilin polypeptide is less than about 10% of the binding of the antibody to Sortilin as measured, e.g., by a radioimmunoassay (RIA). In certain embodiments, an antibody that binds to Sortilin has a dissociation constant (KD) of < 1 μ M, < 100 nM, < 10 nM, < 1 nM, < 0.1 nM, < 0.01 nM, or < 0.001 nM (e.g., 10⁻⁸ M or less, e.g. from 10⁻⁸ M to 10⁻¹³ M, e.g., from 10⁻⁹ M to 10⁻¹³ M). In certain embodiments, an anti-Sortilin antibody binds to an epitope of Sortilin that is conserved among Sortilin from different species.

[0075] The term “immunoglobulin” (Ig) is used interchangeably with “antibody” herein. The term “antibody” herein is used in the broadest sense and specially covers monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies) including those formed from at least two intact antibodies, and antibody fragments so long as they exhibit the desired biological activity.

[0076] “Native antibodies” are usually heterotetrameric glycoproteins of about 150,000 Daltons, composed of two identical Light (“L”) chains and two identical heavy (“H”) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intra-chain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light chain and heavy chain variable domains.

[0077] For the structure and properties of the different classes of antibodies, *see, e.g., Basic and Clinical Immunology*, 8th Ed., Daniel P. Stites, Abba I. Terr and Tristram G. Parslow (eds.), Appleton & Lange, Norwalk, CT, 1994, page 71 and Chapter 6.

[0078] The light chain from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (“ κ ”) and lambda (“ λ ”), based on the amino acid sequences of their constant domains. Depending on the amino acid sequence of the constant domain of their heavy chains (CH), immunoglobulins can be assigned to different classes or isotypes. There are five classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, having heavy chains designated alpha (“ α ”), delta (“ δ ”), epsilon (“ ϵ ”), gamma (“ γ ”), and mu (“ μ ”), respectively. The γ and α classes are further divided into subclasses (isotypes) on the basis of relatively minor differences in the CH sequence and function, e.g., humans express the following subclasses: IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2. The subunit

structures and three-dimensional configurations of different classes of immunoglobulins are well known and described generally in, for example, Abbas *et al.*, Cellular and Molecular Immunology, 4th ed. (W.B. Saunders Co., 2000).

[0079] The “*variable region*” or “*variable domain*” of an antibody, such as an anti-Sortilin antibody of the present disclosure, refers to the amino-terminal domains of the heavy or light chain of the antibody. The variable domains of the heavy chain and light chain may be referred to as “V_H” and “V_L”, respectively. These domains are generally the most variable parts of the antibody (relative to other antibodies of the same class) and contain the antigen binding sites.

[0080] The term “*variable*” refers to the fact that certain segments of the variable domains differ extensively in sequence among antibodies, such as anti-Sortilin antibodies of the present disclosure. The variable domain mediates antigen binding and defines the specificity of a particular antibody for its particular antigen. However, the variability is not evenly distributed across the entire span of the variable domains. Instead, it is concentrated in three segments called hypervariable regions (HVRs) both in the light-chain and the heavy chain variable domains. The more highly conserved portions of variable domains are called the framework regions (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a beta-sheet configuration, connected by three HVRs, which form loops connecting, and in some cases forming part of, the beta-sheet structure. The HVRs in each chain are held together in close proximity by the FR regions and, with the HVRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (*see* Kabat et al., *Sequences of Immunological Interest*, Fifth Edition, National Institute of Health, Bethesda, MD (1991)). The constant domains are not involved directly in the binding of antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent-cellular toxicity.

[0081] An “*isolated*” antibody, such as an anti-Sortilin antibody of the present disclosure, is one that has been identified, separated and/or recovered from a component of its production environment (*e.g.*, naturally or recombinantly). Preferably, the isolated polypeptide is free of association with all other contaminant components from its production environment. Contaminant components from its production environment, such as those resulting from recombinant transfected cells, are materials that would typically interfere with research, diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified: (1) to greater than 95% by weight of antibody as determined by, for example, the Lowry method, and in some embodiments, to greater than 99% by weight; (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within

recombinant T cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, an isolated polypeptide or antibody will be prepared by at least one purification step.

[0082] The term "*monoclonal antibody*" as used herein refers to an antibody, such as a monoclonal anti-Sortilin antibody of the present disclosure, obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations and/or post-translation modifications (*e.g.*, isomerizations, amidations, *etc.*) that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. In contrast to polyclonal antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by a variety of techniques, including, but not limited to one or more of the following methods, immunization methods of animals including, but not limited to rats, mice, rabbits, guinea pigs, hamsters and/or chickens with one or more of DNA(s), virus-like particles, polypeptide(s), and/or cell(s), the hybridoma methods, B-cell cloning methods, recombinant DNA methods, and technologies for producing human or human-like antibodies in animals that have parts or all of the human immunoglobulin loci or genes encoding human immunoglobulin sequences.

[0083] The terms "*full-length antibody*," "*intact antibody*" or "*whole antibody*" are used interchangeably to refer to an antibody, such as an anti-Sortilin antibody of the present disclosure, in its substantially intact form, as opposed to an antibody fragment. Specifically, whole antibodies include those with heavy and light chains including an Fc region. The constant domains may be native sequence constant domains (*e.g.*, human native sequence constant domains) or amino acid sequence variants thereof. In some cases, the intact antibody may have one or more effector functions.

[0084] An "*antibody fragment*" comprises a portion of an intact antibody, preferably the antigen binding and/or the variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂ and Fv fragments; diabodies; linear antibodies (*see* U.S. Patent 5,641,870, Example 2; Zapata *et al.*, *Protein Eng.* 8(10):1057-1062 (1995)); single-chain antibody molecules and multispecific antibodies formed from antibody fragments.

[0085] Papain digestion of antibodies, such as anti-Sortilin antibodies of the present disclosure, produces two identical antigen-binding fragments, called "*Fab*" fragments, and a residual "*Fc*" fragment,

a designation reflecting the ability to crystallize readily. The Fab fragment consists of an entire L chain along with the variable region domain of the H chain (V_H), and the first constant domain of one heavy chain (C_{H1}). Each Fab fragment is monovalent with respect to antigen binding, *i.e.*, it has a single antigen-binding site. Pepsin treatment of an antibody yields a single large $F(ab')_2$ fragment which roughly corresponds to two disulfide linked Fab fragments having different antigen-binding activity and is still capable of cross-linking antigen. Fab' fragments differ from Fab fragments by having a few additional residues at the carboxy terminus of the C_{H1} domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. $F(ab')_2$ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

[0086] The Fc fragment comprises the carboxy-terminal portions of both H chains held together by disulfides. The effector functions of antibodies are determined by sequences in the Fc region, the region which is also recognized by Fc receptors (FcR) found on certain types of cells.

[0087] “Fv” is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This fragment consists of a dimer of one heavy- and one light-chain variable region domain in tight, non-covalent association. From the folding of these two domains emanate six hypervariable loops (3 loops each from the H and L chain) that contribute the amino acid residues for antigen binding and confer antigen binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three HVRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

[0088] “Single-chain Fv” also abbreviated as “sFv” or “scFv” are antibody fragments that comprise the V_H and V_L antibody domains connected into a single polypeptide chain. Preferably, the sFv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of the sFv, *see* Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

[0089] “Functional fragments” of antibodies, such as anti-Sortilin antibodies of the present disclosure, comprise a portion of an intact antibody, generally including the antigen binding or variable region of the intact antibody or the F region of an antibody which retains or has modified FcR binding capability. Examples of antibody fragments include linear antibody, single-chain antibody molecules and multispecific antibodies formed from antibody fragments.

[0090] The term “diabodies” refers to small antibody fragments prepared by constructing sFv fragments (see preceding paragraph) with short linkers (about 5-10) residues) between the V_H and V_L

domains such that inter-chain but not intra-chain pairing of the variable domains is achieved, thereby resulting in a bivalent fragment, *i.e.*, a fragment having two antigen-binding sites. Bispecific diabodies are heterodimers of two “crossover” sFv fragments in which the V_H and V_L domains of the two antibodies are present on different polypeptide chains.

[0091] As used herein, a “chimeric antibody” refers to an antibody (immunoglobulin), such as a chimeric anti-Sortilin antibody of the present disclosure, in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is(are) identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity. Chimeric antibodies of interest herein include PRIMATIZED[®] antibodies wherein the antigen-binding region of the antibody is derived from an antibody produced by, *e.g.*, immunizing macaque monkeys with an antigen of interest. As used herein, “humanized antibody” is used a subset of “chimeric antibodies.”

[0092] “*Humanized*” forms of non-human (*e.g.*, murine) antibodies, such as humanized forms of anti-Sortilin antibodies of the present disclosure, are chimeric antibodies comprising amino acid residues from non-human HVRs and amino acid residues from human FRs. In certain embodiments, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the HVRs (*e.g.*, CDRs) correspond to those of a non-human antibody, and all or substantially all of the FRs correspond to those of a human antibody. A humanized antibody optionally may comprise at least a portion of an antibody constant region derived from a human antibody. A “humanized form” of an antibody, *e.g.*, a non-human antibody, refers to an antibody that has undergone humanization.

[0093] A “*human antibody*” is one that possesses an amino-acid sequence corresponding to that of an antibody, such as an anti-Sortilin antibody of the present disclosure, produced by a human and/or has been made using any of the techniques for making human antibodies as disclosed herein. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues. Human antibodies can be produced using various techniques known in the art, including phage-display libraries and yeast-based platform technologies. Human antibodies can be prepared by administering the antigen to a transgenic animal that has been modified to produce such antibodies in response to antigenic challenge, but whose endogenous loci have been disabled, *e.g.*, immunized xenomice as well as generated via a human B-cell hybridoma technology.

[0094] The term “*hypervariable region*,” “*HVR*,” or “*HV*,” when used herein refers to the regions of an antibody-variable domain, such as that of an anti-Sortilin antibody of the present disclosure, that are

hypervariable in sequence and/or form structurally defined loops. Generally, antibodies comprise six HVRs; three in the V_H (H1, H2, H3), and three in the V_L (L1, L2, L3). In native antibodies, H3 and L3 display the most diversity of the six HVRs, and H3 in particular is believed to play a unique role in conferring fine specificity to antibodies. Naturally occurring camelid antibodies consisting of a heavy chain only are functional and stable in the absence of light chain.

[0095] A number of HVR delineations are in use and are encompassed herein. In some embodiments, the HVRs may be Kabat complementarity-determining regions (CDRs) based on sequence variability and are the most commonly used (Kabat et al., *supra*). In some embodiments, the HVRs may be Chothia CDRs. Chothia refers instead to the location of the structural loops (Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). In some embodiments, the HVRs may be AbM HVRs. The AbM HVRs represent a compromise between the Kabat CDRs and Chothia structural loops, and are used by Oxford Molecular's AbM antibody-modeling software. In some embodiments, the HVRs may be “contact” HVRs. The “contact” HVRs are based on an analysis of the available complex crystal structures. The residues from each of these HVRs are noted below.

Loop	Kabat	AbM	Chothia	Contact
L1	L24-L34	L24-L34	L26-L32	L30-L36
L2	L50-L56	L50-L56	L50-L52	L46-L55
L3	L89-L97	L89-L97	L91-L96	L89-L96
H1	H31-H35B	H26-H35B	H26-H32	H30-H35B (Kabat numbering)
H1	H31-H35	H26-H35	H26-H32	H30-H35 (Chothia numbering)
H2	H50-H65	H50-H58	H53-H55	H47-H58
H3	H95-H102	H95-H102	H96-H101	H93-H101

[0096] HVRs may comprise “extended HVRs” as follows: 24-36 or 24-34 (L1), 46-56 or 50-56 (L2), and 89-97 or 89-96 (L3) in the V_L, and 26-35 (H1), 50-65 or 49-65 (a preferred embodiment) (H2), and 93-102, 94-102, or 95-102 (H3) in the V_H. The variable-domain residues are numbered according to Kabat et al., *supra*, for each of these extended-HVR definitions.

[0097] “Framework” or “FR” residues are those variable-domain residues other than the HVR residues as herein defined.

[0098] An “acceptor human framework” as used herein is a framework comprising the amino acid sequence of a V_L or V_H framework derived from a human immunoglobulin framework or a human consensus framework. An acceptor human framework “derived from” a human immunoglobulin framework or a human consensus framework may comprise the same amino acid sequence thereof, or it may comprise pre-existing amino acid sequence changes. In some embodiments, the number of pre-existing amino acid changes are 10 or less, 9 or less, 8 or less, 7 or less, 6 or less, 5 or less, 4 or less, 3 or

less, or 2 or less. Where pre-existing amino acid changes are present in a V_H, preferable those changes occur at only three, two, or one of positions 71H, 73H and 78H; for instance, the amino acid residues at those positions may be 71A, 73T and/or 78A. In one embodiment, the V_L acceptor human framework is identical in sequence to the V_L human immunoglobulin framework sequence or human consensus framework sequence.

[0099] A “*human consensus framework*” is a framework that represents the most commonly occurring amino acid residues in a selection of human immunoglobulin V_L or V_H framework sequences. Generally, the selection of human immunoglobulin V_L or V_H sequences is from a subgroup of variable domain sequences. Generally, the subgroup of sequences is a subgroup as in Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991). Examples include for the V_L, the subgroup may be subgroup kappa I, kappa II, kappa III or kappa IV as in Kabat *et al.*, *supra*. Additionally, for the V_H, the subgroup may be subgroup I, subgroup II, or subgroup III as in Kabat *et al.*, *supra*.

[0100] An “*amino-acid modification*” at a specified position, *e.g.*, of an anti-Sortilin antibody of the present disclosure, refers to the substitution or deletion of the specified residue, or the insertion of at least one amino acid residue adjacent the specified residue. Insertion “adjacent” to a specified residue means insertion within one to two residues thereof. The insertion may be N-terminal or C-terminal to the specified residue. The preferred amino acid modification herein is a substitution.

[0101] An “*affinity-matured*” antibody, such as an anti-Sortilin antibody of the present disclosure, is one with one or more alterations in one or more HVRs thereof that result in an improvement in the affinity of the antibody for antigen, compared to a parent antibody that does not possess those alteration(s). In one embodiment, an affinity-matured antibody has nanomolar or even picomolar affinities for the target antigen. Affinity-matured antibodies are produced by procedures known in the art. For example, Marks *et al.*, *Bio/Technology* 10:779-783 (1992) describes affinity maturation by V_H- and V_L-domain shuffling. Random mutagenesis of HVR and/or framework residues is described by, for example: Barbas *et al. Proc Nat. Acad. Sci. USA* 91:3809-3813 (1994); Schier *et al. Gene* 169:147-155 (1995); Yelton *et al. J. Immunol.* 155:1994-2004 (1995); Jackson *et al., J. Immunol.* 154(7):3310-9 (1995); and Hawkins *et al, J. Mol. Biol.* 226:889-896 (1992).

[0102] As use herein, the term “*specifically recognizes*” or “*specifically binds*” refers to measurable and reproducible interactions such as attraction or binding between a target and an antibody, such as an anti-Sortilin antibody of the present disclosure, that is determinative of the presence of the target in the presence of a heterogeneous population of molecules including biological molecules. For example, an antibody, such as an anti-Sortilin antibody of the present disclosure, that specifically or preferentially binds to a target or an epitope is an antibody that binds this target or epitope with greater affinity, avidity,

more readily, and/or with greater duration than it binds to other targets or other epitopes of the target. It is also understood by reading this definition that, for example, an antibody (or a moiety) that specifically or preferentially binds to a first target may or may not specifically or preferentially bind to a second target. As such, “*specific binding*” or “*preferential binding*” does not necessarily require (although it can include) exclusive binding. An antibody that specifically binds to a target may have an association constant of at least about 10^3 M^{-1} or 10^4 M^{-1} , sometimes about 10^5 M^{-1} or 10^6 M^{-1} , in other instances about 10^6 M^{-1} or 10^7 M^{-1} , about 10^8 M^{-1} to 10^9 M^{-1} , or about 10^{10} M^{-1} to 10^{11} M^{-1} or higher. A variety of immunoassay formats can be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. *See, e.g.*, Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York, for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity.

[0103] As used herein, an “*interaction*” between a Sortilin protein and a second protein encompasses, without limitation, protein-protein interaction, a physical interaction, a chemical interaction, binding, covalent binding, and ionic binding. As used herein, an antibody “inhibits interaction” between two proteins when the antibody disrupts, reduces, or completely eliminates an interaction between the two proteins. An antibody of the present disclosure, or fragment thereof, “inhibits interaction” between two proteins when the antibody or fragment thereof binds to one of the two proteins.

[0104] An “*agonist*” antibody or an “*activating*” antibody is an antibody, such as an agonist anti-Sortilin antibody of the present disclosure, that induces (*e.g.*, increases) one or more activities or functions of the antigen after the antibody binds the antigen.

[0105] A “*blocking*” antibody, an “*antagonist*” antibody, or an “*inhibitory*” antibody is an antibody, such as an anti-Sortilin antibody of the present disclosure, that inhibits or reduces (*e.g.*, decreases) antigen binding to one or more ligand after the antibody binds the antigen, and/or that inhibits or reduces (*e.g.*, decreases) one or more activities or functions of the antigen after the antibody binds the antigen. In some embodiments, blocking antibodies, antagonist antibodies, or inhibitory antibodies substantially or completely inhibit antigen binding to one or more ligand and/or one or more activities or functions of the antigen.

[0106] Antibody “*effector functions*” refer to those biological activities attributable to the Fc region (a native sequence Fc region or amino acid sequence variant Fc region) of an antibody, and vary with the antibody isotype.

[0107] The term “*Fc region*” herein is used to define a C-terminal region of an immunoglobulin heavy chain, including native-sequence Fc regions and variant Fc regions. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy-chain Fc region is usually

defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The C-terminal lysine (residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accordingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue. Suitable native-sequence Fc regions for use in the antibodies of the present disclosure include human IgG1, IgG2, IgG3 and IgG4.

[0108] A “*native sequence Fc region*” comprises an amino acid sequence identical to the amino acid sequence of an Fc region found in nature. Native sequence human Fc regions include a native sequence human IgG1 Fc region (non-A and A allotypes); native sequence human IgG2 Fc region; native sequence human IgG3 Fc region; and native sequence human IgG4 Fc region as well as naturally occurring variants thereof.

[0109] A “*variant Fc region*” comprises an amino acid sequence which differs from that of a native sequence Fc region by virtue of at least one amino acid modification, preferably one or more amino acid substitution(s). Preferably, the variant Fc region has at least one amino acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, *e.g.* from about one to about ten amino acid substitutions, and preferably from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of the parent polypeptide. The variant Fc region herein will preferably possess at least about 80% homology with a native sequence Fc region and/or with an Fc region of a parent polypeptide, and most preferably at least about 90% homology therewith, more preferably at least about 95% homology therewith.

[0110] “*Fc receptor*” or “*FcR*” describes a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII subclasses, including allelic variants and alternatively spliced forms of these receptors, FcγRII receptors include FcγRIIA (an “activating receptor”) and FcγRIIB (an “inhibiting receptor”), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (“ITAM”) in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (“ITIM”) in its cytoplasmic domain. Other FcRs, including those to be identified in the future, are encompassed by the term “FcR” herein. FcRs can also increase the serum half-life of antibodies. As used herein, “*percent (%) amino acid sequence identity*” and “*homology*” with respect to a peptide, polypeptide or antibody sequence refers to the percentage of amino acid residues in a candidate sequence that are identical with the amino acid

residues in the specific peptide or polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or MEGALIGNTM (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms known in the art needed to achieve maximal alignment over the full length of the sequences being compared.

[0111] An “*isolated*” cell is a molecule or a cell that is identified and separated from at least one contaminant cell with which it is ordinarily associated in the environment in which it was produced. In some embodiments, the isolated cell is free of association with all components associated with the production environment. The isolated cell is in a form other than in the form or setting in which it is found in nature. Isolated cells are distinguished from cells existing naturally in tissues, organs, or individuals. In some embodiments, the isolated cell is a host cell of the present disclosure.

[0112] An “*isolated*” nucleic acid molecule encoding an antibody, such as an anti-Sortilin antibody of the present disclosure, is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the environment in which it was produced. Preferably, the isolated nucleic acid is free of association with all components associated with the production environment. The isolated nucleic acid molecules encoding the polypeptides and antibodies herein is in a form other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from nucleic acid encoding the polypeptides and antibodies herein existing naturally in cells.

[0113] The term “*vector*,” as used herein, is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a “plasmid,” which refers to a circular double stranded DNA into which additional DNA segments may be ligated. Another type of vector is a phage vector. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as “recombinant expression vectors,” or simply, “expression vectors.” In general, expression vectors of utility in recombinant DNA

techniques are often in the form of plasmids. In the present specification, “plasmid” and “vector” may be used interchangeably as the plasmid is the most commonly used form of vector.

[0114] “*Polynucleotide*,” or “*nucleic acid*,” as used interchangeably herein, refer to polymers of nucleotides of any length, and include DNA and RNA. The nucleotides can be deoxyribonucleotides, ribonucleotides, modified nucleotides or bases, and/or their analogs, or any substrate that can be incorporated into a polymer by DNA or RNA polymerase or by a synthetic reaction.

[0115] A “*host cell*” includes an individual cell or cell culture that can be or has been a recipient for vector(s) for incorporation of polynucleotide inserts. Host cells include progeny of a single host cell, and the progeny may not necessarily be completely identical (in morphology or in genomic DNA complement) to the original parent cell due to natural, accidental, or deliberate mutation. A host cell includes cells transfected *in vivo* with a polynucleotide(s) of the present disclosure.

[0116] “Carriers” as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers that are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed.

[0117] The term “*about*” as used herein refers to the usual error range for the respective value readily known to the skilled person in this technical field. Reference to “*about*” a value or parameter herein includes (and describes) embodiments that are directed to that value or parameter *per se*.

[0118] As used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly indicates otherwise. For example, reference to an “antibody” is a reference to from one to many antibodies, such as molar amounts, and includes equivalents thereof known to those skilled in the art, and so forth.

[0119] It is understood that aspect and embodiments of the present disclosure described herein include “comprising,” “consisting,” and “consisting essentially of” aspects and embodiments.

Overview

[0120] The present disclosure relates, in part, to anti-Sortilin antibodies that exhibit one or more improved and/or enhanced functional characteristics relative to an anti-Sortilin antibody, S-60, having a heavy chain variable region and a light chain variable region as described in WO2016164637. Non-limiting improved and/or enhanced functional properties include, for example, antibodies capable of binding Sortilin with higher affinity, reducing cell surface levels of Sortilin, decreasing the half-maximal effective concentration (EC₅₀) to reduce cell surface levels of Sortilin, improving the maximal reduction of cell surface levels of Sortilin, increasing extracellular secretion of a Sortilin ligand: *e.g.* Progranulin (PGRN), decreasing the half-maximal effective concentration (EC₅₀) to block PGRN binding to Sortilin, improving the maximal blocking of PGRN binding to Sortilin, or any combination thereof. Also

contemplated herein are anti-Sortilin antibodies with different Fc variants that exhibit one or more improved and/or enhanced functional characteristics, including decreasing the half-maximal effective concentration (EC_{50}) to reduce cell surface levels of Sortilin, improving the maximal reduction of cell surface levels of Sortilin, increasing extracellular secretion of a Sortilin ligand: *e.g.* Progranulin (PGRN), decreasing the half-maximal effective concentration (EC_{50}) to block PGRN binding to Sortilin, and improving the maximal blocking of PGRN binding to Sortilin.

[0121] In some embodiments, anti-Sortilin antibodies of the present disclosure have higher potencies in reducing cell surface levels of Sortilin relative to an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60. In some embodiments, potency is measured by half maximal effective concentration (EC_{50}) for reduction of cell surface cell levels of SORT1.

[0122] The present disclosure further relates to methods of making and using anti-Sortilin antibodies as described herein; pharmaceutical compositions containing such antibodies; nucleic acids encoding such antibodies; and host cells containing nucleic acids encoding such antibodies.

[0123] In some embodiments, the anti-Sortilin antibodies of the present disclosure may have one or more activities that are due, at least in part, to the ability of the antibodies to reduce levels (*e.g.*, cell surface levels) of Sortilin by inducing degradation, down regulation, cleavage, receptor desensitization, and/or lysosomal targeting of Sortilin. In some embodiments, the anti-Sortilin antibodies exhibit one or more of the following properties: a. have a dissociation constant (K_D) for human Sortilin that is lower than that of an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60; b. decrease cell surface levels of Sortilin (*e.g.*, decrease cell surface levels of Sortilin on engineered cells expressing human Sortilin *in vitro*) with a half-maximal effective concentration (EC_{50}) that is lower than that of an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60; c. have a dissociation constant (K_D) for human Sortilin that may range from about 0.560 nM to about 1.63 nM, for example when the K_D is determined by fluorescent activated cell sorting (FACS); d. have a dissociation constant (K_D) for human Sortilin that may range from about 0.270 nM to about 2.910 nM, for example when the K_D is determined by BioLayer Interferometry; e. decrease cell surface levels of Sortilin (*e.g.*, decreases cell surface levels of Sortilin on engineered cells expressing human Sortilin *in vitro*) with a half-maximal effective concentration (EC_{50}) that may range from about 72.58 pM to about 103.6 pM, for example when the EC_{50} is determined *in vitro* by FACS; f. reduce cell surface levels of Sortilin with a maximum reduction that may range from about 51.33% to 88.76%; g. increase extracellular secretion of PGRN higher than that of an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60; h. block PGRN binding to Sortilin (*e.g.*, block binding of PGRN of Sortilin on

engineered cells expressing human Sortilin *in vitro*) with a half-maximal effective concentration (EC_{50}) that may range from about .325 nM to about 2.27 nM, for example when the EC_{50} is determined *in vitro* by FACS; and/or i. improve the maximal blocking of PGRN binding to Sortilin higher than that of an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60.

[0124] Advantageously, anti-Sortilin antibodies of the present disclosure reduce cell surface levels (*e.g.*, up to approximately 3.02-fold) of Sortilin more potently (*e.g.*, with a lower EC_{50}) as compared to a control anti-Sortilin antibody (*e.g.*, a control anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60) (*See e.g.*, **Example 2**). In some embodiments, anti-Sortilin antibodies of the present disclosure reduce binding of PGRN to Sortilin with a lower EC_{50} (up to approximately 5.36 -fold) more potently as compared to a control anti-Sortilin antibody (*e.g.*, a control anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60) (*See e.g.*, **Example 3**). Moreover, advantageously, anti-Sortilin antibodies of the present disclosure have a higher affinity (*e.g.*, up to approximately 2.79-fold higher affinity) for Sortilin (*e.g.*, a lower K_D value as measured by FACS or BioLayer Interferometry) as compared to a control anti-Sortilin antibody (*e.g.*, a control anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60) (*See e.g.*, **Examples 1 and 4**). Surprisingly, higher affinity for Sortilin does not necessarily correlate with an increase in ability or potency of reduction of cell surface levels of Sortilin (*See, e.g.*, **Examples 2 and 4**) nor an increase in ability or potency of blocking binding of PGRN to Sortilin (*See, e.g.*, **Examples 3 and 4**).

[0125] The present disclosure further relates to anti-Sortilin antibodies with improved stability during manufacturing, storage, and *in vivo* administration. In some embodiments, anti-Sortilin antibodies of the present disclosure have improved stability under various stress conditions (*See, e.g.*, **Example 4**).

Sortilin proteins

[0126] In one aspect, the present disclosure provides antibodies, such as isolated (*e.g.*, monoclonal) antibodies, that interact with or otherwise bind to a region, such as an epitope, within a Sortilin protein of the present disclosure. In some embodiments, the antibodies interact with or otherwise bind to a region, such as an epitope, within a Sortilin protein of the present disclosure with improved/enhanced kinetics (*e.g.*, relative to an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, the antibodies interact with or otherwise bind to a region, such as an epitope, within a human Sortilin protein, with a half-maximal effective concentration (EC_{50}) that is lower than that of a control antibody (*e.g.*, relative to an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure bind to a Sortilin protein and modulate one

or more Sortilin activities after binding to the Sortilin protein, for example, an activity associated with Sortilin expression on a cell. Sortilin proteins of the present disclosure include, without limitation, a mammalian Sortilin protein, human Sortilin protein, mouse Sortilin protein, and rat Sortilin protein.

Exemplary Sortilin protein sequences are shown in **Table 32**.

[0127] Sortilin is variously referred to as sortilin 1, SORT1, 100 kDa NT receptor, glycoprotein 95 (GP95), Progranulin receptor (PGRN-R), and neurotensin receptor 3 (NT-3 or NTR-3). Sortilin is an 831 amino acid protein that encodes a type I membrane receptor. Various Sortilin homologs are known, including without limitation, human Sortilin, rat Sortilin, and mouse Sortilin. The amino acid sequence of human Sortilin is set forth below as SEQ ID NO: 81 (with key amino acid residues predicted to participate in Progranulin binding depicted in bold and, and the predicted pro-NGF binding region underlined):

10	20	30	40	50
MERPWGAADG	LSRWPHGLGL	LLLLQLLP	PPS	TLNQDRLDAP
60	70	80	90	100
SGPIGVSWGL	RAAAAGGAFP	RGGRWRRSAP	GEDEECGRVR	DFVAKLANNT
110	120	130	140	150
HQHVFDLRLG	SVSLSWVGDS	TGVILVLTTF	HVPLVIMTFG	<u>QSKLYRSE</u> DY
160	170	180	190	200
GKNFKDITDL	INNTFIRTEF	GMAIGPENSG	KVVLTAEVSG	GSRGG RI FRS
210	220	230	240	250
SD FA K NF VQ T	DLPFHPLTQM	MYSQNSDYL	LALSTENGLW	VSKNFGGKWE
260	270	280	290	300
EIHKA V CL A K	WGSDNTIFFT	TYANGSCKAD	LGALELWRTS	DLGKSFKTIG
310	320	330	340	350
VKIYS F GLGG	RFL F ASVMAD	KDT T RR I HVS	TDQGD T WSMA	QLPSVGQEQ F
360	370	380	390	400
Y S I LAANDDM	VFMHVDEPGD	TGFGTIFTSD	DRGIVYSKSL	DRHLYTTTGG
410	420	430	440	450
ETDFT N VTSL	RGVYITSVLS	EDNSIQTMIT	FDQGGRWTHL	RKPENSECDA
460	470	480	490	500
TAKNKNECSL	HIHASYSISQ	KLNVPMAPLS	EPNAVGI V IA	HGSVGDAISV
510	520	530	540	550
MVPDVYISDD	GGYSWTKMLE	GPHYYTILDS	GGIIVAIEHS	SRPINVIKFS
560	570	580	590	600
TDEGQCWQTY	TFTRDPIYFT	GLASEPGARS	M NISIWGFTE	<u>S</u> FLTSQWVS Y
610	620	630	640	650
TIDFKDILER	NCEEKDYTIW	LAHSTDPE D Y	EDGCILGYKE	QFLRLRKSSV
660	670	680	690	700
CQNGRDYVVT	KQPSICLCSL	EDFLCDFGY Y	RPENDSKC V E	QPELK G HDL E
710	720	730	740	750
FCLYGREEHL	TTNGYRKIPG	DKCQGGVNPV	REVKDLKKKC	TSNFLSPEKQ
760	770	780	790	800
NSKSNSVPII	LAI V GLMLVT	VVAGVLIVKK	YVCGGRFLVH	RYSVLQ Q HAE
810	820	830		
ANGVDGVDAL	DTASHTNKSG	YHDDSDE D LL	E	

[0128] The amino acid sequence of mouse Sortilin is set forth in SEQ ID NO: 82:

```

MERPRGAADG LLRWPLGLLL LLQLLPAAV GQDRLDAPP PAPLLRWAG PVGVSWGLRA
AAPGGVPVRA GRWRRGAPAE DQDCGRLPDF IAKLTNNTHQ HVFDDLSGSV SLSWVG DSTG
VILVLTTFQV PLVIVSFGQS KLYRSEDY GK NFKDITNLIN NTFIRTEFGM AIGPENSGKV
ILTAEVSGGS RGRVFRSSD FAKNFVQTDL PFHPLTQMMY SPQNSDYLLA LSTENGLWVS
KNFGEKWEEI HKA VCLAKWG PNNIIFFTTH VNGSCKADLG ALELWRTSDL GKTFKTIGVK
IYSFGLGGRF LFASVMADKD TTRRIHVSTD QGDTWSMAQL PSVGQEQFYS ILAANEDMVF
MHVDEPGDTG FGTIFTSDDR GIVYSKSLDR HLYTTTGGET DFTNVTSLRG VYITSTLSED
NSIQSMITFD QGGRWEHLRK PENSKCDATA KNKNECSLHI HASYSISQKL NVPMAPLSEP
NAVGI V IAHG SVGDAISVMV PDVYISDDGG YSWAKMLEGP HYYTILDSGG IIVAIEHSNR
PINVIKFSTD EGQCWQSYVF TQEPYFTGL ASEP G ARSMN ISIWGFTE SF ITRQWVS YTV
DFKDILERN C EEDDYTTWLA HSTD PGDYKD GCILGYKEQF LRLRKSSVCQ NGRDYV VAKQ
PSVCPCSLED FLCDFGYFRP ENASECVEQP ELKGHELEFC LYGKEEHLTT NGYRKIPGDK
CQGGMNPARE VKDLKKKCTS NFLNPTKQNS KSNSVPIILA IVGLMLVTVV AGVLIVKKYV
CGGRFLVHRY SVLQQHAEAD GVEALDSTSH AKSGYHDDSD EDLLE

```

[0129] The amino acid sequence of rat Sortilin is set forth in SEQ ID NO: 83:

```

MERPRGAADG LLRWPLGLLL LLQLLPAAV GQDRLDAPP PAPLLRWAG PVGVSWGLRA
AAPGGVPVRA GRWRRGAPAE DQDCGRLPDF IAKLTNNTHQ HVFDDLSGSV SLSWVG DSTG
VILVLTTFQV PLVIVSFGQS KLYRSEDY GK NFKDITNLIN NTFIRTEFGM AIGPENSGKV
ILTAEVSGGS RGRVFRSSD FAKNFVQTDL PFHPLTQMMY SPQNSDYLLA LSTENGLWVS
KNFGEKWEEI HKA VCLAKWG PNNIIFFTTH VNGSCKADLG ALELWRTSDL GKTFKTIGVK
IYSFGLGGRF LFASVMADKD TTRRIHVSTD QGDTWSMAQL PSVGQEQFYS ILAANDDMVF
MHVDEPGDTG FGTIFTSDDR GIVYSKSLDR HLYTTTGGET DFTNVTSLRG VYITSTLSED
NSIQSMITFD QGGRWEHLQK PENSKCDATA KNKNECSLHI HASYSISQKL NVPMAPLSEP
NAVGI V IAHG SVGDAISVMV PDVYISDDGG YSWAKMLEGP HYYTILDSGG IIVAIEHSNR
PINVIKFSTD EGQCWQSYVF SQEPVYFTGL ASEP G ARSMN ISIWGFTE SF LTRQWVS YTI
DFKDILERN C EENDYTTWLA HSTD PGDYKD GCILGYKEQF LRLRKSSVCQ NGRDYV VAKQ
PSICPCSLED FLCDFGYFRP ENASECVEQP ELKGHELEFC LYGKEEHLTT NGYRKIPGDR
CQGGMNPARE VKDLKKKCTS NFLNPKKQNS KSSSVPIILA IVGLMLVTVV AGVLIVKKYV
CGGRFLVHRY SVLQQHAEAD GVEALDTASH AKSGYHDDSD EDLLE

```

[0130] In some embodiments, the Sortilin is a preprotein that includes a signal sequence. In some embodiments, the Sortilin is a mature protein. In some embodiments, the mature Sortilin protein does not include a signal sequence. In some embodiments, the mature Sortilin protein is expressed on a cell.

[0131] Sortilin proteins of the present disclosure include several domains, including without limitation, a signal sequence, a propeptide, a luminal domain, a Vps10p domain, a 10 CC domain, a transmembrane domain and a cytoplasmic domain. Additionally, proteins of the present disclosure are

expressed at high levels in a number of tissues, including without limitation, the brain, spinal cord, heart and skeletal muscle, thyroid, placenta, and testis.

[0132] Accordingly, as used herein a “Sortilin” protein of the present disclosure includes, without limitation, a mammalian Sortilin protein, human Sortilin protein, primate Sortilin protein, mouse Sortilin protein, and rat Sortilin protein. Additionally, anti-Sortilin antibodies of the present disclosure may bind an epitope within one or more of a mammalian Sortilin protein, human Sortilin protein, primate Sortilin, mouse Sortilin protein, and rat Sortilin protein.

Sortilin protein domains

[0133] Sortilin proteins of the present disclosure contain several domains, such as a Vps10p domain that contains an Asp-box motif, a ten-bladed beta-propeller structure, and a hydrophobic loop; and a 10 CC domain.

[0134] As disclosed herein, interactions between Sortilin proteins of the present disclosure and pro-neurotrophins or neurotrophins are mediated by the Vps10p domain that contains a ten-bladed beta-propeller structure and an Asp-box motif. In certain embodiments, Sortilin proteins of the present disclosure contain a Vps10p domain that includes a ten-bladed beta-propeller structure and is located within amino acid residues 78-611 of human Sortilin (SEQ ID NO: 81) or amino acid residues of a mammalian Sortilin that correspond to amino acid residues 78-611 of SEQ ID NO: 81. In certain embodiments, amino acid residues 190-220 of human Sortilin (SEQ ID NO: 81) or amino acid residues of a mammalian Sortilin that correspond to amino acid residues 190-220 of SEQ ID NO: 81 are located within the Vps10p domain.

[0135] Vps10p domains of the present disclosure may include an Asp-box motif. As used herein, Asp-box motifs have the following sequence: (S/T)-X-(D/N)-X-X-X-X-(W/F/Y) (SEQ ID NO: 84), or X-X-(S/T)-X-(D/N)-X-G-X-(T/S)-(W/F/Y)-X (SEQ ID NO: 85), where X represents any amino acid. In human Sortilin, the Asp-box motif is SSDFAKNF (SEQ ID NO:86), located at amino acid residues 200-207 of human Sortilin. Accordingly, in certain embodiments, an Asp-box motif is located at amino acid residues 200-207 of human Sortilin (SEQ ID NO: 81) or amino acid residues of a mammalian Sortilin that correspond to amino acid residues 200-207 of SEQ ID NO: 81.

[0136] As disclosed herein, interactions between Sortilin proteins of the present disclosure and p75 are mediated by the 10CC domain of the hydrophobic loop of the Vps10p domain.

[0137] In certain embodiments, Sortilin proteins of the present disclosure contain a 10CC domain that is located within amino acid residues 610-757 of human Sortilin (SEQ ID NO: 81) or amino acid residues of a mammalian Sortilin that correspond to amino acid residues 610-757 of SEQ ID NO: 81. In preferred embodiments, amino acid residues 592-593, 610-660, and/or 667-749 of human Sortilin (SEQ

ID NO: 81) or amino acid residues of a mammalian Sortilin that correspond to amino acid residues 592-593, 610-660, and/or 667-749 of SEQ ID NO: 81 are located within the 10CC domain of Sortilin.

[0138] In other embodiments, Sortilin proteins of the present disclosure contain a hydrophobic loop within the Vps10p domain that is located within amino acid residues 130-141 of human Sortilin (SEQ ID NO: 81) or amino acid residues of a mammalian Sortilin that correspond to amino acid residues 130-141 of SEQ ID NO: 81.

[0139] As one of skill in the art will appreciate, the beginning and ending residues of the domains of the present disclosure may vary depending upon the computer modeling program used or the method used for determining the domain.

Sortilin binding partners

[0140] Sortilin proteins of the present disclosure can interact with (*e.g.*, bind to) one or more proteins including, without limitation, Progranulin (PGRN) protein; neurotrophins, such as pro-neurotrophins, pro-neurotrophin-3, neurotrophin-3, pro-neurotrophin-4/5, neurotrophin-4/5, pro-nerve growth factor (Pro-NGF), nerve growth factor (NGF), pro-brain-derived neurotrophic factor (Pro-BDNF), and brain-derived neurotrophic factor (BDNF); neurotensin, p75, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), amyloid precursor protein, A beta peptide, PCSK9, p75NTR, and receptor associated protein (RAP).

Progranulin (PGRN)

[0141] Sortilin proteins of the present disclosure have been shown to interact (*e.g.*, bind to) directly with Progranulin and mediate the degradation of Progranulin (*e.g.*, Zheng, Y *et al.*, (2011) *PLoS ONE* 6(6): e21023).

[0142] Progranulin is variously referred to as PGRN, proepithelin, granulin-epithelin precursor, PC (prostate cancer) cell-derived growth factor (PCDGF), and acrogranin. Progranulin is a 593 amino acid protein that encodes a 68.5 kDa secreted glycoprotein that has 7.5 repeats of smaller granulin (epithelin) motifs, ranging from 6-25 kDa, which can be proteolytically cleaved from the precursor PGRN. Examples of Progranulin cleavage products include, without limitation, granulin A/ Epithelins 1, granulin B Epithelins 2, granulin C, granulins D, granulin E, granulin F, granulin G and any other known peptide products derived from Progranulin.

[0143] Accordingly, anti-Sortilin antibodies of the present disclosure that increase Progranulin levels, decrease cell surface levels of Sortilin, and/or block the interaction (*e.g.*, binding) between Sortilin and Progranulin would be beneficial for preventing, lowering the risk of, or treating conditions and/or diseases associated with decreased levels of Progranulin expression and/or activity, cell death (*e.g.*, neuronal cell death), frontotemporal dementia, Alzheimer's disease, vascular dementia, seizures, retinal dystrophy, a traumatic brain injury, a spinal cord injury, long-term depression, atherosclerotic vascular

diseases, undesirable symptoms of normal aging, dementia, mixed dementia, Creutzfeldt-Jakob disease, normal pressure hydrocephalus, amyotrophic lateral sclerosis, Huntington's disease, taupathy disease, stroke, acute trauma, chronic trauma, lupus, acute and chronic colitis, Crohn's disease, inflammatory bowel disease, ulcerative colitis, malaria, essential tremor, central nervous system lupus, Behcet's disease, Parkinson's disease, dementia with Lewy bodies, multiple system atrophy, intervertebral disc degeneration, Shy-Drager syndrome, progressive supranuclear palsy, cortical basal ganglionic degeneration, acute disseminated encephalomyelitis, granulomatous disorders, Sarcoidosis, diseases of aging, age related macular degeneration, glaucoma, retinitis pigmentosa, retinal degeneration, respiratory tract infection, sepsis, eye infection, systemic infection, inflammatory disorders, arthritis, multiple sclerosis, metabolic disorder, obesity, insulin resistance, type 2 diabetes, tissue or vascular damage, an injury, and/or one or more undesirable symptoms of normal aging. Additionally, anti-Sortilin antibodies of the present disclosure that increase Progranulin levels, decrease cell surface levels of Sortilin, and/or block the interaction (e.g., binding) between Sortilin and Progranulin may inhibit interaction between Sortilin and Progranulin, may induce one or more Progranulin activities, may reduce the endosomal internalization of Progranulin, or fragments thereof, and/or may increase the effective concentration of Progranulin.

[0144] In some embodiments, anti-Sortilin antibodies of the present disclosure that increase Progranulin levels, decrease cell surface levels of Sortilin, and/or block the interaction (e.g., binding) between Sortilin and Progranulin bind to one or more amino acids within amino acid residues 131-138, 175-181, 190-220, 199-220, 190-211, 196-207, 196-199, 200-207, 203-207, 207-231, 207-227, 212-221, 233-243, 237-247, 237-260, 297-317, 314-338, 367-391, 429-443, 623-632, and/or 740-749 of human Sortilin (SEQ ID NO: 81); or within amino acid residues of a mammalian Sortilin that corresponds to amino acid residues 131-138, 175-181, 190-220, 199-220, 190-211, 196-207, 196-199, 200-207, 203-207, 207-231, 207-227, 212-221, 233-243, 237-247, 237-260, 297-317, 314-338, 367-391, 429-443, 623-632, and/or 740-749 of SEQ ID NO: 81. In other embodiments, anti-Sortilin antibodies of the present disclosure that increase Progranulin levels, decrease cell surface levels of Sortilin, and/or block the interaction (e.g., binding) between Sortilin and Progranulin may bind one or more amino acids of amino acid residues His131, Val132, Pro133, Leu134, Val135, Ile136, Met137, Thr138, Arg196, Phe198, Arg199, Phe203, Lys205, Phe207, Thr210, Thr218, Tyr222, Ser223, Ser227, Ser242, Lys243, Lys248, Lys254, Lys260, Ser305, Phe306, Gly307, Arg311, Phe314, Ser316, Arg325, Arg326, Ile327, Phe350, Tyr351, Ser352, Ile353, Asn373, Ser379, Arg382, Tyr386, Ser595, and/or Glu700 of human Sortilin (SEQ ID NO: 81); or of amino acid residues of a mammalian Sortilin that corresponds to one or more amino acid residues His131, Val132, Pro133, Leu134, Val135, Ile136, Met137, Thr138, Arg196, Phe198, Arg199, Phe203, Lys205, Phe207, Thr210, Thr218, Tyr222, Ser223, Ser227, Ser242, Lys243,

Lys248, Lys254, Lys260, Ser305, Phe306, Gly307, Arg311, Phe314, Ser316, Arg325, Arg326, Ile327, Phe350, Tyr351, Ser352, Ile353, Asn373, Ser379, Arg382, Tyr386, Ser595, and/or Glu700 of SEQ ID NO: 81.

Other Sortilin Ligands

[0145] Sortilin proteins of the present disclosure have been shown to interact (e.g., bind) directly with pro-neurotrophins (e.g., pro-NGF), which harbor a pro-domain and are typically pro-apoptotic. This binding may be mediated through a linear epitope on Sortilin that corresponds to amino acid residues 163-174 of SEQ ID NO: 81. Sortilin proteins of the present disclosure also have been shown to interact (e.g., bind) with neurotensin within the beta-propeller structure of Sortilin, and an important contact has been shown to at serine 283 of human Sortilin.

[0146] Sortilin proteins of the present disclosure have also been shown to interact (e.g., bind) with the low affinity nerve growth factor (NGF) receptor (p75) within the 10CC domain of Sortilin or the hydrophobic loop of the Vps10p domain of Sortilin. As disclosed herein, Sortilin proteins of the present disclosure can function as a co-receptor with p75 for pro-neurotrophins, which induce apoptotic signaling. Sortilin proteins of the present disclosure have further been shown to interact (e.g., bind) with amyloid precursor protein (APP).

[0147] Sortilin proteins of the present disclosure have been shown to interact (e.g., bind) with the lipoprotein lipase (LpL). As disclosed herein, Sortilin proteins of the present disclosure bind to and modify the degradation of LpL. Sortilin proteins of the present disclosure have been shown to interact (e.g., bind) with the apolipoprotein AV (APOA5). As disclosed herein, Sortilin proteins of the present disclosure bind to and modify the degradation of APOA5.

[0148] Sortilin proteins of the present disclosure have been shown to interact (e.g., bind) with the apolipoprotein E (APOE, APOE2, APOE3, APOE4). As disclosed herein, Sortilin proteins of the present disclosure bind to and modify the degradation and transport of APOE as well as agents that APOE carries such as the A beta peptide. In addition, Sortilin proteins of the present disclosure have been shown to interact (e.g., bind) with receptor-associate protein (RAP). Further, Sortilin proteins of the present disclosure have been shown to interact (e.g., bind) with Proprotein convertase subtilisin/kexin type 9 (PCSK9), and secrete it to the circulation.

[0149] In some embodiments of any of the anti-Sortilin antibodies, anti-Sortilin antibodies of the present disclosure may also inhibit (e.g., block) the interaction between Sortilin and one or more other Sortilin ligands of the present disclosure. In some embodiments, the one or more other ligands is one or more of pro-neurotrophins, neurotensin, low affinity nerve growth factor (NGF) receptor (p75), amyloid precursor protein, lipoprotein lipase, apolipoprotein AV, apolipoprotein, receptor-associated protein, and/or proprotein convertase subtilisin/kexin type 9. Such antibodies may be beneficial for preventing,

lowering the risk of, or treating conditions and/or diseases associated with decreased levels of one or more other Sortilin ligands expression and/or activity, cell death (e.g., neuronal cell death). In some embodiments, the one or more other ligands is one or more of pro-neurotrophins, neurotensin, low affinity nerve growth factor (NGF) receptor (p75), amyloid precursor protein, lipoprotein lipase, apolipoprotein AV, apolipoprotein, receptor-associated protein, and/or proprotein convertase subtilisin/kexin type 9.

[0150] Anti-Sortilin antibodies of the present disclosure that inhibit (e.g., block) the interaction between Sortilin and one or more other Sortilin ligands of the present disclosure may also prevent cell death (e.g., apoptosis) induced by one or more other Sortilin ligands. In some embodiments, the one or more other ligands is one or more of pro-neurotrophins, neurotensin, low affinity nerve growth factor (NGF) receptor (p75), amyloid precursor protein, lipoprotein lipase, apolipoprotein AV, apolipoprotein, receptor-associated protein, and/or proprotein convertase subtilisin/kexin type 9.

[0151] In some embodiments of any of the anti-Sortilin antibodies, anti-Sortilin antibodies of the present disclosure that inhibit (e.g., block) the interaction between Sortilin and one or more Sortilin ligands of the present disclosure bind one or more amino acids within amino acid residues within amino acid residues 131-138, 175-181, 190-220, 199-220, 190-211, 196-207, 196-199, 200-207, 203-207, 207-231, 207-227, 212-221, 233-243, 237-247, 237-260, 297-317, 314-338, 367-391, 429-443, 623-632, and/or 740-749 of human Sortilin (SEQ ID NO: 81); or within amino acid residues of a mammalian Sortilin that corresponds to amino acid residues 131-138, 175-181, 190-220, 199-220, 190-211, 196-207, 196-199, 200-207, 203-207, 207-231, 207-227, 212-221, 233-243, 237-247, 237-260, 297-317, 314-338, 367-391, 429-443, 623-632, and/or 740-749 of SEQ ID NO: 81. In some embodiments, the one or more other ligands is one or more of pro-neurotrophins, neurotensin, low affinity nerve growth factor (NGF) receptor (p75), amyloid precursor protein, lipoprotein lipase, apolipoprotein AV, apolipoprotein, receptor-associated protein, and/or proprotein convertase subtilisin/kexin type 9.

[0152] In other embodiments of any of the anti-Sortilin antibodies, anti-Sortilin antibodies of the present disclosure that inhibit (e.g., block) the interaction between Sortilin and one or more Sortilin ligands of the present disclosure bind one or more amino acids of amino acid residues His131, Val132, Pro133, Leu134, Val135, Ile136, Met137, Thr138, Arg196, Phe198, Arg199, Phe203, Lys205, Phe207, Thr210, Thr218, Tyr222, Ser223, Ser227, Ser242, Lys243, Lys248, Lys254, Lys260, Ser305, Phe306, Gly307, Arg311, Phe314, Ser316, Arg325, Arg326, Ile327, Phe350, Tyr351, Ser352, Ile353, Asn373, Ser379, Arg382, Tyr386, Ser595, and/or Glu700 of human Sortilin (SEQ ID NO: 81); or of amino acid residues of a mammalian Sortilin that corresponds to one or more amino acid residues His131, Val132, Pro133, Leu134, Val135, Ile136, Met137, Thr138, Arg196, Phe198, Arg199, Phe203, Lys205, Phe207, Thr210, Thr218, Tyr222, Ser223, Ser227, Ser242, Lys243, Lys248, Lys254, Lys260, Ser305, Phe306,

Gly307, Arg311, Phe314, Ser316, Arg325, Arg326, Ile327, Phe350, Tyr351, Ser352, Ile353, Asn373, Ser379, Arg382, Tyr386, Ser595, and/or Glu700 of SEQ ID NO: 81. In some embodiments, the one or more other ligands is one or more of pro-neurotrophins, neurotensin, low affinity nerve growth factor (NGF) receptor (p75), amyloid precursor protein, lipoprotein lipase, apolipoprotein AV, apolipoprotein, receptor-associated protein, and/or proprotein convertase subtilisin/kexin type 9.

Sortilin antibodies

[0153] Certain aspects of the present disclosure relate to anti-Sortilin antibodies comprising one or more improved and/or enhanced functional characteristics. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise one or more improved and/or enhanced functional characteristics relative to an anti-Sortilin antibody, S-60, having a heavy chain variable region and a light chain variable region as described in WO2016164637. In some embodiments, anti-Sortilin antibodies of the present disclosure have an affinity for Sortilin (e.g., human Sortilin) that is higher than that of a control anti-Sortilin antibody (e.g., a control anti-Sortilin antibody comprising a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure decrease cellular levels (e.g., cell surface levels) of Sortilin to a greater degree and with a half-maximal effective concentration (EC50) that is lower than that of a control antibody (e.g., a control anti-Sortilin antibody comprising a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure improve the maximal reduction of cell surface levels of Sortilin relative to an anti-Sortilin antibody comprising a heavy chain variable region and a light chain variable region corresponding to S-60. In some embodiments, anti-Sortilin antibodies of the present disclosure increase the secretion of extracellular Progranulin (PGRN) relative to an anti-Sortilin antibody comprising a heavy chain variable region and a light chain variable region corresponding to S-60. In some embodiments, anti-Sortilin antibodies of the present disclosure blocking binding of PGRN to Sortilin to a greater degree and with a half-maximal effective concentration (EC50) that is lower than that of a control antibody (e.g., a control anti-Sortilin antibody comprising a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure improve the maximal blocking of PGRN binding to Sortilin relative to an anti-Sortilin antibody comprising a heavy chain variable region and a light chain variable region corresponding to S-60.

[0154] Also contemplated herein are anti-Sortilin antibodies with different Fc variants that exhibit one or more improved and/or enhanced functional characteristics relative to an anti-Sortilin antibody comprising a heavy chain variable region and a light chain variable region corresponding to S-60, including decreasing the half-maximal effective concentration (EC50) to reduce cell surface levels of Sortilin, improving the maximal reduction of cell surface levels of Sortilin, increasing extracellular

secretion of PGRN, decreasing the half-maximal effective concentration (EC50) to block PGRN binding to Sortilin, and improving the maximal blocking of PGRN binding to Sortilin.

[0155] In some embodiments, an anti-Sortilin antibody of the present disclosure is a human antibody, a bispecific antibody, a monoclonal antibody, a multivalent antibody, a conjugated antibody, or a chimeric antibody

[0156] In a preferred embodiment, an anti-Sortilin antibody of the present disclosure is a monoclonal antibody.

Anti-Sortilin antibody binding epitope

[0157] In some embodiments, anti-Sortilin antibodies of the present disclosure bind to a Sortilin protein of the present disclosure and/or naturally occurring variants. In some embodiments, anti-Sortilin antibodies of the present disclosure bind to a Sortilin protein, wherein the Sortilin protein is a human protein. In some embodiments, anti-Sortilin antibodies of the present disclosure bind to a Sortilin protein, wherein the Sortilin protein is a wild-type protein. In some embodiments, anti-Sortilin antibodies of the present disclosure bind to a Sortilin protein, wherein the Sortilin protein is a naturally occurring variant.

[0158] In certain preferred embodiments, anti-Sortilin antibodies of the present disclosure bind specifically to a human Sortilin protein.

[0159] Certain aspects of the present disclosure provide anti-Sortilin antibodies that bind a discontinuous Sortilin epitope. In some embodiments, the discontinuous Sortilin epitope comprises two or more peptides, three or more peptides, four or more peptides, five or more peptides, six or more peptides, seven or more peptide, eight or more peptides, nine or more peptides, or 10 or more peptides. In some embodiments, each of the peptides comprise five or more, six or more, seven or more, eight or more, nine or more, 10 or more, 11 or more, 12 or more, 13 or more 14 or more, 15 or more, 16 or more, 17 or more, 18 or more, 19 or more, 20 or more, 21 or more, 22 or more, 23 or more, 24 or more, 25 or more, 26 or more, 27 or more, 28 or more, 29 or more, or 30 or more amino acid residues of the amino acid sequence of SEQ ID NO: 81; or five or more, six or more, seven or more, eight or more, nine or more, 10 or more, 11 or more, 12 or more, 13 or more 14 or more, 15 or more, 16 or more, 17 or more, 18 or more, 19 or more, 20 or more, 21 or more, 22 or more, 23 or more, 24 or more, 25 or more, 26 or more, 27 or more, 28 or more, 29 or more, or 30 or more amino acid residues on a mammalian Sortilin protein corresponding to the amino acid sequence of SEQ ID NO: 81. Other aspects of the present disclosure provide anti-Sortilin antibodies that bind to a conformational epitope of Sortilin.

[0160] Certain aspects of the present disclosure provide anti-Sortilin antibodies that bind to one or more amino acids within amino acid residues 207-231 of human Sortilin (SEQ ID NO: 81); or within amino acid residues on a mammalian Sortilin corresponding to amino acid residues 207-231 of SEQ ID NO: 81. In some embodiments, the anti-Sortilin antibodies bind to one or more amino acids of amino

acid residues Thr218, Tyr222, Ser223, and/or Ser227 of human Sortilin (SEQ ID NO: 81); or of amino acid residues on a mammalian Sortilin corresponding to amino acid residues Thr218, Tyr222, Ser223, and/or Ser227 of SEQ ID NO: 81. In some embodiments, the anti-Sortilin antibodies bind to one or more amino acids of amino acid residues F105, L108, R109, G110, I537, F569, E590, F592, L593, S595, and/or W597 of human Sortilin (SEQ ID NO: 81); or of amino acid residues on a mammalian Sortilin corresponding to amino acid residues F105, L108, R109, G110, I537, F569, E590, F592, L593, S595, and/or W597 of SEQ ID NO: 81. Other aspects of the present disclosure provide anti-Sortilin antibodies that bind to an epitope having amino acid residues (S/T)-X-(D/N)-X-X-X-X-(W/F/Y), where X represents any amino acid (SEQ ID NO: 84).

[0161] In some embodiments, anti-Sortilin antibodies of the present disclosure bind to a Sortilin protein of the present disclosure expressed on the surface of cell and the naked antibodies inhibit interaction (*e.g.*, binding) between the Sortilin protein and a protein selected from Progranulin (PGRN), a pro-neurotrophin, a neurotrophin, pro-neurotrophin-3, neurotrophin-3, pro-neurotrophin-4/5, neurotrophin-4/5, pro-nerve growth factor (pro-NGF), nerve growth factor (NGF), pro-brain-derived neurotrophic factor (pro-BDNF), brain-derived neurotrophic factor (BDNF), neurotensin, p75, Sortilin propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), PCSK9, and receptor associated protein (RAP). In some embodiments, anti-Sortilin antibodies of the present disclosure bind to a Sortilin protein of the present disclosure expressed on the surface of cell and the naked antibodies inhibit interaction (*e.g.*, binding) between the Sortilin protein and PGRN.

[0162] In some embodiments, anti-Sortilin antibodies of the present disclosure that bind to a Sortilin protein of the present disclosure inhibit interaction (*e.g.*, binding) between the Sortilin protein and Progranulin, a pro-neurotrophin, a neurotrophin, pro-neurotrophin-3, neurotrophin-3, pro-neurotrophin-4/5, neurotrophin-4/5, pro-nerve growth factor (pro-NGF), nerve growth factor (NGF), pro-brain-derived neurotrophic factor (pro-BDNF), brain-derived neurotrophic factor (BDNF), neurotensin, p75, Sortilin propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), PCSK9, and receptor associated protein (RAP) by reducing the effective levels of Sortilin that is available to interact with these proteins either on the cell surface or inside the cell.

[0163] In some embodiments, anti-Sortilin antibodies of the present disclosure that bind to a Sortilin protein of the present disclosure inhibit interaction (*e.g.*, binding) between the Sortilin protein and Progranulin, a pro-neurotrophin, a neurotrophin, pro-neurotrophin-3, neurotrophin-3, pro-neurotrophin-4/5, neurotrophin-4/5, pro-nerve growth factor (pro-NGF), nerve growth factor (NGF), pro-brain-derived neurotrophic factor (pro-BDNF), brain-derived neurotrophic factor (BDNF), neurotensin, p75, Sortilin

propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), PCSK9, and receptor associated protein (RAP) by inducing degradation of Sortilin.

[0164] In some embodiments, anti-Sortilin antibodies of the present disclosure bind to a human Sortilin, or a homolog thereof, including without limitation, a mammalian Sortilin protein, or a non-human primate Sortilin protein. In some embodiments, anti-Sortilin antibodies of the present disclosure specifically bind to human Sortilin. In some embodiments, anti-Sortilin antibodies of the present disclosure bind to human Sortilin and are not cross-reactive with Sortilin orthologs or homologs from other species.

Anti-Sortilin antibody competitive binding

[0165] In some embodiments, anti-Sortilin antibodies of the present disclosure competitively inhibit binding of at least one other anti-Sortilin antibody selected from any of the antibodies listed in **Tables 1-29**. In some embodiments, anti-Sortilin antibodies of the present disclosure competitively inhibit binding of at least one antibody selected from: S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24; and any combination thereof.

[0166] In some embodiments, an anti-Sortilin antibody of the present disclosure competes with one or more anti-Sortilin antibodies selected from: S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24 and any combination thereof, for binding to Sortilin when the anti-Sortilin antibody reduces the binding of one or more antibodies selected from: S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24; and any combination thereof to

Sortilin by an amount the ranges from about 50% to 100%, as compared to binding to Sortilin in the absence of the anti-Sortilin antibody.

[0167] In some embodiments, an anti-Sortilin antibody of the present disclosure competes with one or more anti-Sortilin antibodies selected from: S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24, and any combination thereof, for binding to Sortilin when the anti-Sortilin antibody reduces the binding of one or more antibodies selected from: S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24; and any combination thereof to Sortilin by at least 50%, at least 55%, by at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or 100%, as compared to binding to Sortilin in the absence of the anti-Sortilin antibody.

[0168] In some embodiments, an anti-Sortilin antibody of the present disclosure that reduces the binding of one or more antibodies selected from: S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24, and any combination thereof, to Sortilin by 100% indicates that the anti-Sortilin antibody essential completely blocks the binding of one or more antibodies selected from S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24, and any combination thereof, to Sortilin. In some embodiments, the anti-Sortilin antibody and the one or more

antibodies selected from S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24, and any combination thereof, are present in an amount that corresponds to a 10:1 ratio, 9:1 ratio, 8:1 ratio, 7:1 ratio, 6:1 ratio, 5:1 ratio, 4:1 ratio, 3:1 ratio, 2:1 ratio, 1:1 ratio, 0.75:1 ratio, 0.5:1 ratio, 0.25:1 ratio, 0.1:1 ratio, 0.075:1 ratio, 0.050:1 ratio, 0.025:1 ratio, 0.01:1 ratio, 0.0075: ratio, 0.0050:1 ratio, 0.0025:1 ratio, 0.001: ratio, 0.00075:1 ratio, 0.00050:1 ratio, 0.00025:1 ratio, 0.0001: ratio, 1:10 ratio, 1:9 ratio, 1:8 ratio, 1:7 ratio, 1:6 ratio, 1:5 ratio, 1:4 ratio, 1:3 ratio, 1:2 ratio, 1:0.75 ratio, 1:0.5 ratio, 1:0.25 ratio, 1:0.1 ratio, 1:0.075 ratio, 1:0.050 ratio, 1:0.025 ratio, 1:0.01 ratio, 1:0.0075 ratio, 1:0.0050 ratio, 1:0.0025 ratio, 1:0.001 ratio, 1:0.00075 ratio, 1:0.00050 ratio, 1:0.00025 ratio, or 1:0.0001 ratio of anti-Sortilin antibody to one or more antibodies selected from S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24, and any combination thereof. In some embodiments, the anti-Sortilin antibody is present in excess by an amount that ranges from about 1.5-fold to 100-fold, or greater than 100-fold compared to the amount of the one or more antibodies selected from S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24, and any combination thereof. In some embodiments, the anti-Sortilin antibody is present in an amount that is about a 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 15-fold, 20-fold, 25-fold, 30-fold, 35-fold, 40-fold, 45-fold, 50-fold, 55-fold, 60-fold, 65-fold, 70-fold, 75-fold, 80-fold, 85-fold, 90-fold, 95-fold, or 100-fold excess compared to the amount of the one or more antibodies selected from S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-

15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24, and any combination thereof.

[0169] In some embodiments, anti-Sortilin antibodies of the present disclosure bind to an epitope of human Sortilin that is the same as or overlaps with the Sortilin epitope bound by at least one antibody selected from any of the antibodies listed in **Tables 1-29**. In some embodiments, anti-Sortilin antibodies of the present disclosure bind to an epitope of human Sortilin that is the same as or overlaps with the Sortilin epitope bound by at least one antibody selected from: S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24; and any combination thereof. In some embodiments, anti-Sortilin antibodies of the present disclosure bind essentially the same Sortilin epitope bound by at least one antibody selected from any of the antibodies listed in **Tables 1-29**. In some embodiments, anti-Sortilin antibodies of the present disclosure bind essentially the same Sortilin epitope bound by at least one antibody selected from: S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24; and any combination thereof. Detailed exemplary methods for mapping an epitope to which an antibody binds are provided in Morris (1996) "Epitope Mapping Protocols," in *Methods in Molecular Biology* vol. 66 (Humana Press, Totowa, NJ).

[0170] Any suitable competition assay or Sortilin binding assay known in the art, such as BIAcore analysis, ELISA assays, or flow cytometry, may be utilized to determine whether an anti-Sortilin antibody competes with one or more antibodies selected from: S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 (described in WO2016164637); or S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24; and any combination thereof for binding to Sortilin. In an exemplary competition assay, immobilized Sortilin or

cells expressing Sortilin on the cell surface are incubated in a solution comprising a first labeled antibody that binds to Sortilin (e.g., human or non-human primate) and a second unlabeled antibody that is being tested for its ability to compete with the first antibody for binding to Sortilin. The second antibody may be present in a hybridoma supernatant. As a control, immobilized Sortilin or cells expressing Sortilin is incubated in a solution comprising the first labeled antibody but not the second unlabeled antibody. After incubation under conditions permissive for binding of the first antibody to Sortilin, excess unbound antibody is removed, and the amount of label associated with immobilized Sortilin or cells expressing Sortilin is measured. If the amount of label associated with immobilized Sortilin or cells expressing Sortilin is substantially reduced in the test sample relative to the control sample, then that indicates that the second antibody is competing with the first antibody for binding to Sortilin. *See*, Harlow and Lane (1988) *Antibodies: A Laboratory Manual* ch.14 (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY).

[0171] In some embodiments, an anti-Sortilin antibody of the present disclosure competes with an antibody comprising the heavy chain variable domain and the light chain variable domain of an antibody selected from the group consisting of S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-16, S-60-18, S-60-19, S-60-24, and any combination thereof for binding to Sortilin.

[0172] In some embodiments, an anti-Sortilin antibody of the present disclosure binds essentially the same Sortilin epitope as an antibody comprising the heavy chain variable domain and the light chain variable domain of an antibody selected from the group consisting of S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-16, S-60-18, S-60-19, and S-60-24.

[0173] Additional anti-Sortilin antibodies, *e.g.*, antibodies that specifically bind to a Sortilin protein of the present disclosure, may be identified, screened, and/or characterized for their physical/chemical properties and/or biological activities by various assays known in the art.

Anti-Sortilin antibody heavy chain and light chain variable regions

A. Heavy chain HVRs

[0174] In some embodiments, anti-Sortilin antibodies of the present disclosure include a heavy chain variable region comprising one or more (*e.g.*, one or more, two or more, or all three) HVRs selected from HVR-H1, HVR-H2, and HVR-H3 (as shown in **Tables 14-16**). In some embodiments, the heavy chain variable region comprises an HVR-H1, an HVR-H2, and an HVR-H3 (as shown in **Tables 14-16**).

[0175] In some embodiments, the HVR-H1 comprises a sequence of YSISSGYYWG (SEQ ID NO: 1). In some embodiments, the HVR-H2 comprises a sequence according to Formula I: TIYHSGSTYYNPSLX₁S (SEQ ID NO: 4), wherein X₁ is K or E. In some embodiments, the HVR-H2 comprises a sequence selected from SEQ ID NOs: 2-3. In some embodiments, the HVR-H3 comprises a

sequence according to Formula II: ARQGSIX₁QGYYGMDV (SEQ ID NO: 7). In some embodiments, the HVR-H3 comprises a sequence selected from SEQ ID NOs: 5-6.

[0176] In some embodiments, the HVR-H1 comprises an amino acid sequence with at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identity to an amino acid sequence of SEQ ID NO: 1. In some embodiments, the HVR-H1 comprises an amino acid sequence containing substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to an amino acid sequence of SEQ ID NO: 1), but retains the ability to bind to Sortilin. In certain embodiments, up to 1, up to 2, up to 3, up to 4, or up to 5 amino acids been substituted, inserted, and/or deleted in the HVR-H1 amino acid sequence of SEQ ID NO: 1. In some embodiments, the HVR-H2 comprises an amino acid sequence with at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identity to an amino acid sequence selected from SEQ ID NOs: 2-3. In some embodiments, the HVR-H2 comprises an amino acid sequence containing substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to an amino acid sequence selected from SEQ ID NOs: 2-3), but retains the ability to bind to Sortilin. In certain embodiments, up to 1, up to 2, up to 3, up to 4, or up to 5 amino acids been substituted, inserted, and/or deleted in the HVR-H2 amino acid sequence selected from SEQ ID NOs: 2-3. In some embodiments, the HVR-H3 comprises an amino acid sequence with at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identity to an amino acid sequence selected from SEQ ID NOs: 5-6. In some embodiments, the HVR-H3 comprises an amino acid sequence containing substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to an amino acid sequence selected from SEQ ID NOs: 5-6), but retains the ability to bind to Sortilin. In certain embodiments, up to 1, up to 2, up to 3, up to 4, or up to 5 amino acids been substituted, inserted, and/or deleted in the HVR-H3 amino acid sequence selected from SEQ ID NOs: 5-6.

[0177] In some embodiments, the heavy chain variable region comprises an HVR-H1 comprising a sequence of YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising a sequence according to Formula I, and an HVR-H3 comprising a sequence according to Formula II.

[0178] In some embodiments, the heavy chain variable region comprises an HVR-H1 comprising a sequence of SEQ ID NO: 1, an HVR-H2 comprising a sequence selected from SEQ ID NOs: 2-3, and an HVR-H3 comprising a sequence selected from SEQ ID NOs: 5-6.

[0179] In some embodiments, the heavy chain variable region comprises the HVR-H1, HVR-H2, and HVR-H3 of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H],

S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24, or any combination thereof (as shown in **Tables 14-16**).

[0180] In some embodiments, anti-Sortilin antibodies of the present disclosure include a heavy chain variable region, wherein the heavy chain variable region comprises one or more of: (a) an HVR-H1 comprising an amino acid sequence with at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to an HVR-H1 amino acid sequence of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24; (b) an HVR-H2 comprising an amino acid sequence with at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to an HVR-H2 amino acid sequence of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24; and (c) an HVR-H3 comprising an amino acid sequence with at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to an HVR-H3 amino acid sequence of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24.

[0181] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise an HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), and an HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6).

B. Light chain HVRs

[0182] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable region comprising one or more (*e.g.*, one or more, two or more, or all three) HVRs selected from HVR-L1, HVR-L2, and HVR-L3 (as shown in **Tables 17-19**). In some embodiments, the light chain variable region comprises an HVR-L1, an HVR-L2, and an HVR-L3 (as shown in **Tables 17-19**).

[0183] In some embodiments, the HVR-L1 comprises a sequence according to Formula III: RSSQX₁LLX₂SX₃GYNYLD (SEQ ID NO: 28), wherein X₁ is S or G, X₂ is R or H, and X₃ is N, T, S, G, R, D, H, K, Q, Y, E, W, F, I, V, A, M, or L. In some embodiments, the HVR-L1 comprises a sequence selected from SEQ ID NOs: 8-27. In some embodiments, the HVR-L1 comprises a sequence of RSSQSLLRSNGYNYLD (SEQ ID NO:8), RSSQSLLRSTGYNYLD (SEQ ID NO:9), RSSQSLLRSSGYNYLD (SEQ ID NO:10), RSSQSLLRSGGYNYLD (SEQ ID NO:11), RSSQSLLRSRGYNYLD (SEQ ID NO:12), RSSQSLLRSDGYNYLD (SEQ ID NO:13), RSSQSLLRSHGYNYLD (SEQ ID NO:14), RSSQSLLRSKGYNYLD (SEQ ID NO:15), RSSQSLLRSQGYNYLD (SEQ ID NO:16), RSSQSLLRSYGYNYLD (SEQ ID NO:17), RSSQSLLRSEGNYNYLD (SEQ ID NO:18), RSSQSLLRSWGYNYLD (SEQ ID NO:19), RSSQSLLRSFGYNYLD (SEQ ID NO:20), RSSQSLLRSIGYNYLD (SEQ ID NO:21), RSSQSLLRSVGYNYLD (SEQ ID NO:22), RSSQSLLRSAGYNYLD (SEQ ID NO:23), RSSQSLLRSMGYNYLD (SEQ ID NO:24), RSSQSLLRSLGYNYLD (SEQ ID NO:25), RSSQSLLHSNGYNYLD (SEQ ID NO:26), or RSSQGLLSNGYNYLD (SEQ ID NO:27). In one specific embodiment, the HVR-L1 comprises a sequence of RSSQSLLRSNGYNYLD (SEQ ID NO:8). In another specific embodiment, the HVR-L1 comprises a sequence of RSSQSLLRSTGYNYLD (SEQ ID NO:9) (as shown in **Table 19**).

[0184] In some embodiments, the HVR-L2 comprises a sequence according to Formula IV: LGSNRX₁S (SEQ ID NO: 31), wherein X₁ is A or V. In some embodiments, the HVR-L2 comprises a sequence selected from SEQ ID NOs: 29-30.

[0185] In some embodiments, the HVR-L3 comprises a sequence according to Formula V: MQQQEX₁PLT (SEQ ID NO: 34), wherein X₁ is A or T. In some embodiments, the HVR-L3 comprises a sequence selected from SEQ ID NOs: 32-33.

[0186] In some embodiments, the HVR-L1 comprises an amino acid sequence with at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identity to an amino acid sequence selected from SEQ ID NOs: 8-27. In some embodiments, the HVR-L1 comprises an amino acid sequence containing substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to an amino acid sequence selected from SEQ ID NOs: 8-27), but retains the ability to bind to Sortilin. In certain embodiments, up to 1, up to 2, up to 3, up to 4, or up to 5 amino acids been substituted, inserted, and/or deleted in the HVR-L1 amino acid sequence selected from SEQ ID NOs: 8-27. In some

embodiments, the HVR-L2 comprises an amino acid sequence with at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identity to an amino acid sequence selected from SEQ ID NOs: 29-30. In some embodiments, the HVR-L2 comprises an amino acid sequence containing substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to an amino acid sequence selected from SEQ ID NOs: 29-30), but retains the ability to bind to Sortilin. In certain embodiments, up to 1, up to 2, up to 3, up to 4, or up to 5 amino acids been substituted, inserted, and/or deleted in the HVR-L2 amino acid sequence selected from SEQ ID NOs: 29-30. In some embodiments, the HVR-L3 comprises an amino acid sequence with at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identity to an amino acid sequence selected from SEQ ID NOs: 32-33. In some embodiments, the HVR-L3 comprises an amino acid sequence containing substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to an amino acid sequence selected from SEQ ID NOs: 32-33), but retains the ability to bind to Sortilin. In certain embodiments, up to 1, up to 2, up to 3, up to 4, or up to 5 amino acids been substituted, inserted, and/or deleted in the HVR-L3 amino acid sequence selected from SEQ ID NOs: 32-33.

[0187] In some embodiments, the light chain variable region comprises an HVR-L1 comprising a sequence according to Formula III, an HVR-L2 comprising a sequence according to Formula IV, and an HVR-L3 comprising a sequence according to Formula V. In some embodiments, the light chain variable region comprises an HVR-L1 comprising a sequence selected from SEQ ID NOs: 8-27, an HVR-L2 comprising a sequence selected from SEQ ID NOs: 29-30, and an HVR-L3 comprising a sequence selected from SEQ ID NOs: 32-33.

[0188] In some embodiments, the light chain variable region comprises the HVR-L1, HVR-L2, and HVR-L3 of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24, or any combination thereof (as shown in **Tables 17-19**).

[0189] In some embodiments, anti-Sortilin antibodies of the present disclosure include a light chain variable region, wherein the light chain variable region comprises one or more of: (a) an HVR-L1 comprising an amino acid sequence with at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to an HVR-L1 amino acid sequence of antibody S-

60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24; (b) an HVR-L2 comprising an amino acid sequence with at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to an HVR-L2 amino acid sequence of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24; and (c) an HVR-L3 comprising an amino acid sequence with at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to an HVR-L3 amino acid sequence of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24.

[0190] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise an HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0191] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise an HVR-L1 comprising the amino acid sequence RSSQSLLRSTGYNYLD (SEQ ID NO: 9), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

C. Heavy chain HVRs and light chain HVRs

[0192] In some embodiments, anti-Sortilin antibodies of the present disclosure include a heavy chain variable region comprising one or more (*e.g.*, one or more, two or more, or all three) HVRs selected from HVR-H1, HVR-H2, and HVR-H3 (as shown in **Tables 14-16**), and a light chain variable region comprising one or more (*e.g.*, one or more, two or more, or all three) HVRs selected from HVR-L1, HVR-L2, and HVR-L3 (as shown in **Tables 17-19**). In some embodiments, the heavy chain variable

region comprises an HVR-H1, an HVR-H2, and an HVR-H3 (as shown in **Tables 14-16**), and the light chain variable region comprises an HVR-L1, an HVR-L2, and an HVR-L3 (as shown in tables **Tables 17-19**).

[0193] In some embodiments, the heavy chain variable region comprises an HVR-H1 comprising a sequence of YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising a sequence according to Formula I, and an HVR-H3 comprising a sequence according to Formula II, and the light chain variable region comprises an HVR-L1 comprising a sequence according to Formula III, an HVR-L2 comprising a sequence according to Formula IV, and an HVR-L3 comprising a sequence according to Formula V. In some embodiments, the heavy chain variable region comprises an HVR-H1 comprising a sequence of SEQ ID NO: 1, an HVR-H2 comprising a sequence selected from SEQ ID NOs: 2-3, and an HVR-H3 comprising a sequence selected from SEQ ID NOs: 5-6, and the light chain variable region comprises an HVR-L1 comprising a sequence selected from SEQ ID NOs: 8-27, an HVR-L2 comprising a sequence selected from SEQ ID NOs: 29-30, and an HVR-L3 comprising a sequence selected from SEQ ID NOs: 32-33.

[0194] In some aspects, the heavy chain variable region comprises an HVR-H1 comprising a sequence of SEQ ID NO: 1, an HVR-H2 comprising a sequence selected from SEQ ID NOs: 2-3, and an HVR-H3 comprising a sequence selected from SEQ ID NOs: 5-6, and the light chain variable region comprises an HVR-L1 comprising a sequence selected from SEQ ID NOs: 8-27, an HVR-L2 comprising a sequence selected from SEQ ID NOs: 29-30, and an HVR-L3 comprising a sequence of SEQ ID NO: 32.

[0195] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable region comprising the HVR-H1, HVR-H2, and HVR-H3 of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24, or any combination thereof (as shown in **Tables 14-16**); and a light chain variable region comprising the HVR-L1, HVR-L2, and HVR-L3 of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, S-60-24, or any combination thereof (as shown in **Tables 17-19**).

[0196] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable region comprising an HVR-H1, HVR-H2, and HVR-H3 and a light chain variable region comprising an HVR-L1, HVR-L2, and HVR-L3, wherein the antibody comprises the HVR-H1, HVR-H2, HVR-H3, HVR-L1, HVR-L2, and HVR-L3 of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24 (as shown in **Tables 14-19**).

[0197] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region comprises one or more of: (a) an HVR-H1 comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to an HVR-H1 amino acid sequence of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24; (b) an HVR-H2 comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to an HVR-H2 amino acid sequence of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24; and (c) an HVR-H3 comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to an HVR-H3 amino acid sequence of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24; and wherein the light chain variable region comprises one or more of: (a) an HVR-L1 comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least

98%, at least 99%, or 100% identity to an HVR-L1 amino acid sequence of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24; (b) an HVR-L2 comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to an HVR-L2 amino acid sequence of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24; and (c) an HVR-L3 comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to an HVR-L3 amino acid sequence of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24.

[0198] In some embodiments, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIQQGYGMDV (SEQ ID NO: 5); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLSNGYNYLD (SEQ ID NO: 8), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0199] In some embodiments, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIQQGYGMDV (SEQ ID NO: 5); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLSNGYNYLD (SEQ ID NO: 8), the HVR-L2 comprising the amino acid sequence LGSNRVS

(SEQ ID NO: 30), and the HVR-L3 comprising the amino acid sequence MQQQETPLT (SEQ ID NO: 33).

[0200] In some embodiments, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLES (SEQ ID NO: 3), the HVR-H3 comprising the amino acid sequence ARQGSIQGYGMDV (SEQ ID NO: 5); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0201] In some aspects, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIQGYGMDV (SEQ ID NO: 6); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0202] In some aspects, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIQGYGMDV (SEQ ID NO: 6); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSTGYNYLD (SEQ ID NO: 9), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0203] In some embodiments, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIQGYGMDV (SEQ ID NO: 6); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQETPLT (SEQ ID NO: 33).

[0204] In some embodiments, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIQQGYGMDV (SEQ ID NO: 5); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQSLLSNGYNYLD (SEQ ID NO: 26), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQETPLT (SEQ ID NO: 33).

[0205] In some embodiments, an anti-Sortilin antibody of the present disclosure comprises a heavy chain variable region comprising the HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), the HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), the HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and a light chain variable region comprising the HVR-L1 comprising the amino acid sequence RSSQGLLSNGYNYLD (SEQ ID NO: 27), the HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and the HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

D. Heavy chain variable region

[0206] In some embodiments, anti-Sortilin antibodies of the present disclosure include a heavy chain variable region comprising an amino acid sequence selected from SEQ ID NOs: 54-56. In some embodiments, the heavy chain variable region comprises an amino acid sequence with at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identity to an amino acid sequence selected from SEQ ID NOs: 54-56. In some embodiments, the heavy chain variable region comprises an amino acid sequence containing substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to an amino acid sequence selected from SEQ ID NOs: 54-56), but retains the ability to bind to Sortilin. In certain embodiments, up to 1, up to 2, up to 3, up to 4, up to 5, up to 6, up to 7, up to 8, up to 9, or up to 10 amino acids been substituted, inserted, and/or deleted in the heavy chain variable region amino acid sequence selected from SEQ ID NOs: 54-56.

[0207] In some embodiments, the heavy chain variable region comprises the amino acid sequence of SEQ ID NO: 56.

[0208] In some embodiments, anti-Sortilin antibodies of the present disclosure include a heavy chain variable region of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-

15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24 (as shown in **Table 30**).

[0209] In some embodiments, anti-Sortilin antibodies of the present disclosure include a heavy chain variable region comprising an HVR-H1 comprising the amino acid sequence YSISSGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), and an HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6).

E. Light chain variable region

[0210] In some embodiments, anti-Sortilin antibodies of the present disclosure include a light chain variable region comprising an amino acid sequence selected from SEQ ID NOs: 57-80. In some embodiments, the light chain variable region comprises an amino acid sequence with at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identity to an amino acid sequence selected from SEQ ID NOs: 57-80. In some embodiments, the light chain variable region comprises an amino acid sequence containing substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to an amino acid sequence selected from SEQ ID NOs: 57-80), but retains the ability to bind to Sortilin. In certain embodiments, up to 1, up to 2, up to 3, up to 4, up to 5, up to 6, up to 7, up to 8, up to 9, or up to 10 amino acids been substituted, inserted, and/or deleted in the light chain variable region amino acid sequence selected from SEQ ID NOs: 57-80.

[0211] In some embodiments, the light chain variable region includes the amino acid sequence of SEQ ID NO: 57. In some embodiments, the light chain variable region includes the amino acid sequence of SEQ ID NO: 60.

[0212] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable region of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24 (as shown in **Table 31**).

[0213] In some embodiments, anti-Sortilin antibodies of the present disclosure include a light chain variable region comprising an HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

[0214] In some embodiments, anti-Sortilin antibodies of the present disclosure include a light chain variable region comprising an HVR-L1 comprising the amino acid sequence RSSQSLLRSTGYNYLD

(SEQ ID NO: 9), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

F. Heavy chain variable region and light chain variable region

[0215] In some aspects, an anti-Sortilin antibody of the present disclosure includes a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 54-56; and/or a light chain variable region comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 57-80. In some embodiments, the heavy chain variable region comprises an amino acid sequence with at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identity to an amino acid sequence selected from SEQ ID NOs: 54-56, and the light chain variable region comprises an amino acid sequence with at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identity to an amino acid sequence selected from SEQ ID NOs: 57-80. In some embodiments, an anti-Sortilin antibody of the present disclosure includes a heavy chain variable region comprising an amino acid sequence containing substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to an amino acid sequence selected from SEQ ID NOs: 54-56), and a light chain variable region comprising an amino acid sequence containing substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to an amino acid sequence selected from SEQ ID NOs: 57-80), but retains the ability to bind to Sortilin. In certain embodiments, up to 1, up to 2, up to 3, up to 4, up to 5, up to 6, up to 7, up to 8, up to 9, or up to 10 amino acids been substituted, inserted, and/or deleted in the heavy chain variable region amino acid sequence selected from SEQ ID NOs: 54-56; and up to 1, up to 2, up to 3, up to 4, up to 5, up to 6, up to 7, up to 8, up to 9, or up to 10 amino acids been substituted, inserted, and/or deleted in the light chain variable region amino acid sequence selected from SEQ ID NOs: 57-80.

[0216] In some aspects, an anti-Sortilin antibody of the present disclosure includes a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 54-56; and/or a light chain variable region comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 57-58, 60-78, and 80.

[0217] In some embodiments, an anti-Sortilin antibody of the present disclosure binds to a Sortilin protein, wherein the antibody includes a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57; a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 58; a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region

comprising the amino acid sequence of SEQ ID NO: 59; a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 55, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57; a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 55, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 58; a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57; a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 77; a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 78; a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 54, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 79; or a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 80.

[0218] In one aspect, an anti-Sortilin antibody of the present disclosure includes a heavy chain variable region having the amino acid sequence of SEQ ID NO: 56, and a light chain variable region having the amino acid sequence of SEQ ID NO: 57.

[0219] In one aspect, an anti-Sortilin antibody of the present disclosure includes a heavy chain variable region having the amino acid sequence of SEQ ID NO: 56, and a light chain variable region having the amino acid sequence of SEQ ID NO: 60.

[0220] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable region of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24 (as shown in **Table 30**), and a light chain variable region of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24 (as shown in **Table 31**).

[0221] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising an amino acid sequence selected from SEQ ID NOs: 56 and 60. In some embodiments, the antibody comprises a heavy chain variable region of S-60-15 [N33 (wt)] (as shown in

Table 30), and a light chain variable region of antibody S-60-15 [N33 (wt)] (as shown in **Table 31**). In some embodiments, the antibody comprises a heavy chain variable region of S-60-15.1 [N33T] (as shown in **Table 30**), and a light chain variable region of antibody S-60-15.1 [N33T] (as shown in **Table 31**).

Exemplary anti-Sortilin antibodies

[0222] In some embodiments, the anti-Sortilin antibody is an anti-Sortilin monoclonal antibody comprising the heavy chain variable region and the light chain variable region of an antibody selected from S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24. In some embodiments, the anti-Sortilin antibody is an anti-Sortilin monoclonal antibody comprising the heavy chain and the light chain of an antibody selected from S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24.

(1) S-60-10

[0223] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain and a light chain variable domain, wherein the heavy chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-10 or to the amino acid sequence of SEQ ID NO: 54; and/or the light chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-10 or to the amino acid sequence of SEQ ID NO: 57. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-10 or to the amino acid sequence of SEQ ID NO: 54, wherein the heavy chain variable domain comprises the HVR-H1, HVR-H2, and HVR-H3 amino acid sequences of antibody S-60-10. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at

least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-10 or to the amino acid sequence of SEQ ID NO: 57, wherein the light chain variable domain comprises the HVR-L1, HVR-L2, and HVR-L3 amino acid sequences of antibody S-60-10. In some embodiments, the anti-Sortilin antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-10 or to the amino acid sequence of SEQ ID NO: 54 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-10 or the amino acid sequence of SEQ ID NO: 54. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-10 or the amino acid sequence of SEQ ID NO: 54. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VH sequence of antibody S-60-10 or of SEQ ID NO: 54, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) the HVR-H1 amino acid sequence of antibody S-60-10, (b) the HVR-H2 amino acid sequence of antibody S-60-10, and (c) the HVR-H3 amino acid sequence of antibody S-60-10. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain (VL) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-10 or to the amino acid sequence of SEQ ID NO: 57 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-10 or the amino acid sequence of SEQ ID NO: 57. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-10 or the amino acid sequence of SEQ ID NO: 57. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VL sequence of antibody S-60-10 or of SEQ ID NO: 57, including post-translational modifications of that sequence. In a

particular embodiment, the VL comprises one, two or three HVRs selected from: (a) the HVR-L1 amino acid sequence of antibody S-60-10, (b) the HVR-L2 amino acid sequence of antibody S-60-10, and (c) the HVR-L3 amino acid sequence of antibody S-60-10.

[0224] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 105 or SEQ ID NO: 106. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain comprising the amino acid sequence of SEQ ID NO: 139. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 105 or SEQ ID NO: 106 and a light chain comprising the amino acid sequence of SEQ ID NO: 139.

(2) S-60-11

[0225] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain and a light chain variable domain, wherein the heavy chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-11 or to the amino acid sequence of SEQ ID NO: 54; and/or the light chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-11 or to the amino acid sequence of SEQ ID NO: 58. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-11 or to the amino acid sequence of SEQ ID NO: 54, wherein the heavy chain variable domain comprises the HVR-H1, HVR-H2, and HVR-H3 amino acid sequences of antibody S-60-11. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-11 or to the amino acid sequence of SEQ ID NO: 58, wherein the light chain variable domain comprises the HVR-L1, HVR-L2, and HVR-L3 amino acid sequences of antibody S-60-11. In some embodiments, the anti-Sortilin antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-11 or to the amino acid sequence of SEQ ID NO: 54 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the

reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-11 or the amino acid sequence of SEQ ID NO: 54. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-11 or the amino acid sequence of SEQ ID NO: 54. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in in the FR regions. Optionally, the anti-Sortilin antibody comprises the VH sequence of antibody S-60-11 or of SEQ ID NO: 54, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) the HVR-H1 amino acid sequence of antibody S-60-11, (b) the HVR-H2 amino acid sequence of antibody S-60-11, and (c) the HVR-H3 amino acid sequence of antibody S-60-11. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain (VL) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-11 or to the amino acid sequence of SEQ ID NO: 58 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-11 or the amino acid sequence of SEQ ID NO: 58. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-11 or the amino acid sequence of SEQ ID NO: 58. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in in the FR regions. Optionally, the anti-Sortilin antibody comprises the VL sequence of antibody S-60-11 or of SEQ ID NO: 58, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three HVRs selected from: (a) the HVR-L1 amino acid sequence of antibody S-60-11, (b) the HVR-L2 amino acid sequence of antibody S-60-11, and (c) the HVR-L3 amino acid sequence of antibody S-60-11.

[0226] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 105 or SEQ ID NO: 106. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain comprising the amino acid sequence of SEQ ID NO: 140. In some embodiments, anti-Sortilin antibodies of the present

disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 105 or SEQ ID NO: 106 and a light chain comprising the amino acid sequence of SEQ ID NO: 140.

(3) S-60-12

[0227] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain and a light chain variable domain, wherein the heavy chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-12 or to the amino acid sequence of SEQ ID NO: 54; and/or the light chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-12 or to the amino acid sequence of SEQ ID NO: 59. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-12 or to the amino acid sequence of SEQ ID NO: 54, wherein the heavy chain variable domain comprises the HVR-H1, HVR-H2, and HVR-H3 amino acid sequences of antibody S-60-12. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-12 or to the amino acid sequence of SEQ ID NO: 59, wherein the light chain variable domain comprises the HVR-L1, HVR-L2, and HVR-L3 amino acid sequences of antibody S-60-12. In some embodiments, the anti-Sortilin antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-12 or to the amino acid sequence of SEQ ID NO: 54 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-12 or the amino acid sequence of SEQ ID NO: 54. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-12 or the amino acid sequence of SEQ ID NO: 54. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the

substitutions, insertions, or deletions occur in in the FR regions. Optionally, the anti-Sortilin antibody comprises the VH sequence of antibody S-60-12 or of SEQ ID NO: 54, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) the HVR-H1 amino acid sequence of antibody S-60-12, (b) the HVR-H2 amino acid sequence of antibody S-60-12, and (c) the HVR-H3 amino acid sequence of antibody S-60-12. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain (VL) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-12 or to the amino acid sequence of SEQ ID NO: 59 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-12 or the amino acid sequence of SEQ ID NO: 59. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-12 or the amino acid sequence of SEQ ID NO: 59. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in in the FR regions. Optionally, the anti-Sortilin antibody comprises the VL sequence of antibody S-60-12 or of SEQ ID NO: 59, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three HVRs selected from: (a) the HVR-L1 amino acid sequence of antibody S-60-12, (b) the HVR-L2 amino acid sequence of antibody S-60-12, and (c) the HVR-L3 amino acid sequence of antibody S-60-12.

[0228] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 105 or SEQ ID NO: 106. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain comprising the amino acid sequence of SEQ ID NO: 141. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 105 or SEQ ID NO: 106 and a light chain comprising the amino acid sequence of SEQ ID NO: 141.

(4) S-60-13

[0229] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain and a light chain variable domain, wherein the heavy chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-13 or to the amino acid sequence of SEQ ID

NO: 55; and/or the light chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-13 or to the amino acid sequence of SEQ ID NO: 57. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-13 or to the amino acid sequence of SEQ ID NO: 55, wherein the heavy chain variable domain comprises the HVR-H1, HVR-H2, and HVR-H3 amino acid sequences of antibody S-60-13. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-13 or to the amino acid sequence of SEQ ID NO: 57, wherein the light chain variable domain comprises the HVR-L1, HVR-L2, and HVR-L3 amino acid sequences of antibody S-60-13. In some embodiments, the anti-Sortilin antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-13 or to the amino acid sequence of SEQ ID NO: 55 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-13 or the amino acid sequence of SEQ ID NO: 55. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-13 or the amino acid sequence of SEQ ID NO: 55. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VH sequence of antibody S-60-13 or of SEQ ID NO: 55, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) the HVR-H1 amino acid sequence of antibody S-60-13, (b) the HVR-H2 amino acid sequence of antibody S-60-13, and (c) the HVR-H3 amino acid sequence of antibody S-60-13. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain (VL) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain

amino acid sequence of antibody S-60-13 or to the amino acid sequence of SEQ ID NO: 57 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-13 or the amino acid sequence of SEQ ID NO: 57. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-13 or the amino acid sequence of SEQ ID NO: 57. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VL sequence of antibody S-60-13 or of SEQ ID NO: 57, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three HVRs selected from: (a) the HVR-L1 amino acid sequence of antibody S-60-13, (b) the HVR-L2 amino acid sequence of antibody S-60-13, and (c) the HVR-L3 amino acid sequence of antibody S-60-13.

[0230] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 135 or SEQ ID NO: 136. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain comprising the amino acid sequence of SEQ ID NO: 139. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 135 or SEQ ID NO: 136 and a light chain comprising the amino acid sequence of SEQ ID NO: 139.

(5) S-60-14

[0231] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain and a light chain variable domain, wherein the heavy chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-14 or to the amino acid sequence of SEQ ID NO: 55; and/or the light chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-14 or to the amino acid sequence of SEQ ID NO: 58. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-14 or to the amino acid sequence of SEQ ID NO: 55, wherein the heavy chain variable

domain comprises the HVR-H1, HVR-H2, and HVR-H3 amino acid sequences of antibody S-60-14. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-14 or to the amino acid sequence of SEQ ID NO: 58, wherein the light chain variable domain comprises the HVR-L1, HVR-L2, and HVR-L3 amino acid sequences of antibody S-60-14. In some embodiments, the anti-Sortilin antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-14 or to the amino acid sequence of SEQ ID NO: 55 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-14 or the amino acid sequence of SEQ ID NO: 55. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-14 or the amino acid sequence of SEQ ID NO: 55. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in in the FR regions. Optionally, the anti-Sortilin antibody comprises the VH sequence of antibody S-60-14 or of SEQ ID NO: 55, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) the HVR-H1 amino acid sequence of antibody S-60-14, (b) the HVR-H2 amino acid sequence of antibody S-60-14, and (c) the HVR-H3 amino acid sequence of antibody S-60-14. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain (VL) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-14 or to the amino acid sequence of SEQ ID NO: 58 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-14 or the amino acid sequence of SEQ ID NO: 58. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-14 or the amino acid sequence of SEQ ID NO: 58. In certain embodiments, substitutions, insertions, or deletions occur in

regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in in the FR regions. Optionally, the anti-Sortilin antibody comprises the VL sequence of antibody S-60-14 or of SEQ ID NO: 58, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three HVRs selected from: (a) the HVR-L1 amino acid sequence of antibody S-60-14, (b) the HVR-L2 amino acid sequence of antibody S-60-14, and (c) the HVR-L3 amino acid sequence of antibody S-60-14.

[0232] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 135 or SEQ ID NO: 136. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain comprising the amino acid sequence of SEQ ID NO: 140. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 135 or SEQ ID NO: 136 and a light chain comprising the amino acid sequence of SEQ ID NO: 140.

(6) S-60-15

[0233] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain and a light chain variable domain, wherein the heavy chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-15 or to the amino acid sequence of SEQ ID NO: 56; and/or the light chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-15 or to the amino acid sequence of SEQ ID NO: 57. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-15 or to the amino acid sequence of SEQ ID NO: 56, wherein the heavy chain variable domain comprises the HVR-H1, HVR-H2, and HVR-H3 amino acid sequences of antibody S-60-15. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-15 or to the amino acid sequence of SEQ ID NO: 57, wherein the light chain variable domain comprises the HVR-L1, HVR-L2, and HVR-L3 amino acid sequences of antibody S-60-15. In some embodiments, the anti-Sortilin antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at

least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-15 or to the amino acid sequence of SEQ ID NO: 56 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-15 or the amino acid sequence of SEQ ID NO: 56. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-15 or the amino acid sequence of SEQ ID NO: 56. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VH sequence of antibody S-60-15 or of SEQ ID NO: 56, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) the HVR-H1 amino acid sequence of antibody S-60-15, (b) the HVR-H2 amino acid sequence of antibody S-60-15, and (c) the HVR-H3 amino acid sequence of antibody S-60-15. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain (VL) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-15 or to the amino acid sequence of SEQ ID NO: 57 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-15 or the amino acid sequence of SEQ ID NO: 57. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-15 or the amino acid sequence of SEQ ID NO: 57. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VL sequence of antibody S-60-15 or of SEQ ID NO: 57, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three HVRs selected from: (a) the HVR-L1 amino acid sequence of antibody S-60-15, (b) the HVR-L2 amino acid sequence of antibody S-60-15, and (c) the HVR-L3 amino acid sequence of antibody S-60-15.

[0234] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 or SEQ ID NO: 138. In some

embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain comprising the amino acid sequence of SEQ ID NO: 139. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 or SEQ ID NO: 138 and a light chain comprising the amino acid sequence of SEQ ID NO: 139.

(7) S-60-15.1

[0235] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain and a light chain variable domain, wherein the heavy chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-15.1 or to the amino acid sequence of SEQ ID NO: 56; and/or the light chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-15.1 or to the amino acid sequence of SEQ ID NO: 60. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-15.1 or to the amino acid sequence of SEQ ID NO: 56, wherein the heavy chain variable domain comprises the HVR-H1, HVR-H2, and HVR-H3 amino acid sequences of antibody S-60-15.1. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-15.1 or to the amino acid sequence of SEQ ID NO: 60, wherein the light chain variable domain comprises the HVR-L1, HVR-L2, and HVR-L3 amino acid sequences of antibody S-60-15.1. In some embodiments, the anti-Sortilin antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-15.1 or to the amino acid sequence of SEQ ID NO: 56 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-15.1 or the amino acid sequence of SEQ ID NO: 56. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the heavy chain variable domain amino acid sequence of

antibody S-60-15.1 or the amino acid sequence of SEQ ID NO: 56. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in in the FR regions. Optionally, the anti-Sortilin antibody comprises the VH sequence of antibody S-60-15.1 or of SEQ ID NO: 56, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) the HVR-H1 amino acid sequence of antibody S-60-15.1, (b) the HVR-H2 amino acid sequence of antibody S-60-15.1, and (c) the HVR-H3 amino acid sequence of antibody S-60-15.1. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain (VL) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-15.1 or to the amino acid sequence of SEQ ID NO: 60 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-15.1 or the amino acid sequence of SEQ ID NO: 60. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-15.1 or the amino acid sequence of SEQ ID NO: 60. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in in the FR regions. Optionally, the anti-Sortilin antibody comprises the VL sequence of antibody S-60-15.1 or of SEQ ID NO: 60, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three HVRs selected from: (a) the HVR-L1 amino acid sequence of antibody S-60-15.1, (b) the HVR-L2 amino acid sequence of antibody S-60-15.1, and (c) the HVR-L3 amino acid sequence of antibody S-60-15.1.

[0236] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 or SEQ ID NO: 138. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain comprising the amino acid sequence of SEQ ID NO: 142. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 or SEQ ID NO: 138 and a light chain comprising the amino acid sequence of SEQ ID NO: 142.

(8) S-60-16

[0237] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain and a light chain variable domain, wherein the heavy chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least

94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-16 or to the amino acid sequence of SEQ ID NO: 56; and/or the light chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-16 or to the amino acid sequence of SEQ ID NO: 77. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-16 or to the amino acid sequence of SEQ ID NO: 56, wherein the heavy chain variable domain comprises the HVR-H1, HVR-H2, and HVR-H3 amino acid sequences of antibody S-60-16. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-16 or to the amino acid sequence of SEQ ID NO: 77, wherein the light chain variable domain comprises the HVR-L1, HVR-L2, and HVR-L3 amino acid sequences of antibody S-60-16. In some embodiments, the anti-Sortilin antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-16 or to the amino acid sequence of SEQ ID NO: 56 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-16 or the amino acid sequence of SEQ ID NO: 56. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-16 or the amino acid sequence of SEQ ID NO: 56. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VH sequence of antibody S-60-16 or of SEQ ID NO: 56, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) the HVR-H1 amino acid sequence of antibody S-60-16, (b) the HVR-H2 amino acid sequence of antibody S-60-16, and (c) the HVR-H3 amino acid sequence of antibody S-60-16. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain

(VL) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-16 or to the amino acid sequence of SEQ ID NO: 77 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-16 or the amino acid sequence of SEQ ID NO: 77. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-16 or the amino acid sequence of SEQ ID NO: 77. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VL sequence of antibody S-60-16 or of SEQ ID NO: 77, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three HVRs selected from: (a) the HVR-L1 amino acid sequence of antibody S-60-16, (b) the HVR-L2 amino acid sequence of antibody S-60-16, and (c) the HVR-L3 amino acid sequence of antibody S-60-16.

[0238] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 or SEQ ID NO: 138. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain comprising the amino acid sequence of SEQ ID NO: 131. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 or SEQ ID NO: 138 and a light chain comprising the amino acid sequence of SEQ ID NO: 131.

(9) S-60-18

[0239] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain and a light chain variable domain, wherein the heavy chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-18 or to the amino acid sequence of SEQ ID NO: 56; and/or the light chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-18 or to the amino acid sequence of SEQ ID NO: 78. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%,

at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-18 or to the amino acid sequence of SEQ ID NO: 56, wherein the heavy chain variable domain comprises the HVR-H1, HVR-H2, and HVR-H3 amino acid sequences of antibody S-60-18. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-18 or to the amino acid sequence of SEQ ID NO: 78, wherein the light chain variable domain comprises the HVR-L1, HVR-L2, and HVR-L3 amino acid sequences of antibody S-60-18. In some embodiments, the anti-Sortilin antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-18 or to the amino acid sequence of SEQ ID NO: 56 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-18 or the amino acid sequence of SEQ ID NO: 56. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-18 or the amino acid sequence of SEQ ID NO: 56. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VH sequence of antibody S-60-18 or of SEQ ID NO: 56, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) the HVR-H1 amino acid sequence of antibody S-60-18, (b) the HVR-H2 amino acid sequence of antibody S-60-18, and (c) the HVR-H3 amino acid sequence of antibody S-60-18. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain (VL) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-18 or to the amino acid sequence of SEQ ID NO: 78 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-18 or the amino acid sequence of SEQ ID NO: 78. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or

deleted in the light chain variable domain amino acid sequence of antibody S-60-18 or the amino acid sequence of SEQ ID NO: 78. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VL sequence of antibody S-60-18 or of SEQ ID NO: 78, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three HVRs selected from: (a) the HVR-L1 amino acid sequence of antibody S-60-18, (b) the HVR-L2 amino acid sequence of antibody S-60-18, and (c) the HVR-L3 amino acid sequence of antibody S-60-18.

[0240] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 or SEQ ID NO: 138. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain comprising the amino acid sequence of SEQ ID NO: 132. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 or SEQ ID NO: 138 and a light chain comprising the amino acid sequence of SEQ ID NO: 132.

(10) S-60-19

[0241] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain and a light chain variable domain, wherein the heavy chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-19 or to the amino acid sequence of SEQ ID NO: 54; and/or the light chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-19 or to the amino acid sequence of SEQ ID NO: 79. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-19 or to the amino acid sequence of SEQ ID NO: 54, wherein the heavy chain variable domain comprises the HVR-H1, HVR-H2, and HVR-H3 amino acid sequences of antibody S-60-19. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-19 or to the amino acid sequence of SEQ ID NO: 79, wherein the light chain variable domain comprises the HVR-L1, HVR-L2, and HVR-L3 amino

acid sequences of antibody S-60-19. In some embodiments, the anti-Sortilin antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-19 or to the amino acid sequence of SEQ ID NO: 54 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-19 or the amino acid sequence of SEQ ID NO: 54. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-19 or the amino acid sequence of SEQ ID NO: 54. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in in the FR regions. Optionally, the anti-Sortilin antibody comprises the VH sequence of antibody S-60-19 or of SEQ ID NO: 54, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) the HVR-H1 amino acid sequence of antibody S-60-19, (b) the HVR-H2 amino acid sequence of antibody S-60-19, and (c) the HVR-H3 amino acid sequence of antibody S-60-19. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain (VL) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-19 or to the amino acid sequence of SEQ ID NO: 79 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-19 or the amino acid sequence of SEQ ID NO: 79. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-19 or the amino acid sequence of SEQ ID NO: 79. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in in the FR regions. Optionally, the anti-Sortilin antibody comprises the VL sequence of antibody S-60-19 or of SEQ ID NO: 79, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three HVRs selected from: (a) the HVR-L1 amino acid sequence of antibody S-60-19, (b) the HVR-L2 amino acid sequence of antibody S-60-19, and (c) the HVR-L3 amino acid sequence of antibody S-60-19.

[0242] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 105 or SEQ ID NO: 106. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain comprising the amino acid sequence of SEQ ID NO: 133. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 105 or SEQ ID NO: 106 and a light chain comprising the amino acid sequence of SEQ ID NO: 133.

(11) S-60-24

[0243] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain and a light chain variable domain, wherein the heavy chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-24 or to the amino acid sequence of SEQ ID NO: 56; and/or the light chain variable domain comprises an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-24 or to the amino acid sequence of SEQ ID NO: 80. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-24 or to the amino acid sequence of SEQ ID NO: 56, wherein the heavy chain variable domain comprises the HVR-H1, HVR-H2, and HVR-H3 amino acid sequences of antibody S-60-24. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain comprising an amino acid sequence with at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-24 or to the amino acid sequence of SEQ ID NO: 80, wherein the light chain variable domain comprises the HVR-L1, HVR-L2, and HVR-L3 amino acid sequences of antibody S-60-24. In some embodiments, the anti-Sortilin antibody comprises a heavy chain variable domain (VH) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a heavy chain variable domain amino acid sequence of antibody S-60-24 or to the amino acid sequence of SEQ ID NO: 56 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-24 or the amino acid

sequence of SEQ ID NO: 56. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the heavy chain variable domain amino acid sequence of antibody S-60-24 or the amino acid sequence of SEQ ID NO: 56. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VH sequence of antibody S-60-24 or of SEQ ID NO: 56, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three HVRs selected from: (a) the HVR-H1 amino acid sequence of antibody S-60-24, (b) the HVR-H2 amino acid sequence of antibody S-60-24, and (c) the HVR-H3 amino acid sequence of antibody S-60-24. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable domain (VL) sequence having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a light chain variable domain amino acid sequence of antibody S-60-24 or to the amino acid sequence of SEQ ID NO: 80 and contains substitutions (*e.g.*, conservative substitutions, insertions, or deletions relative to the reference sequence), but the anti-Sortilin antibody comprising that sequence retains the ability to bind to Sortilin. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-24 or the amino acid sequence of SEQ ID NO: 80. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in the light chain variable domain amino acid sequence of antibody S-60-24 or the amino acid sequence of SEQ ID NO: 80. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FR regions). In some embodiments, the substitutions, insertions, or deletions occur in the FR regions. Optionally, the anti-Sortilin antibody comprises the VL sequence of antibody S-60-24 or of SEQ ID NO: 80, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three HVRs selected from: (a) the HVR-L1 amino acid sequence of antibody S-60-24, (b) the HVR-L2 amino acid sequence of antibody S-60-24, and (c) the HVR-L3 amino acid sequence of antibody S-60-24.

[0244] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 or SEQ ID NO: 138. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain comprising the amino acid sequence of SEQ ID NO: 134. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 or SEQ ID NO: 138 and a light chain comprising the amino acid sequence of SEQ ID NO: 134.

[0245] In some embodiments, an anti-Sortilin antibody of the present disclosure binds essentially the same Sortilin epitope as an antibody comprising the heavy chain variable domain and the light chain

variable domain of an antibody selected from the group consisting of S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-16, S-60-18, S-60-19, and S-60-24.

[0246] In some embodiments, the anti-Sortilin antibody is anti-Sortilin monoclonal antibody S-60-10. In some embodiments, the anti-Sortilin antibody is an isolated antibody which binds essentially the same Sortilin epitope as S-60-10. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region of monoclonal antibody S-60-10. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the light chain variable region of monoclonal antibody S-60-10. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region and the light chain variable region of monoclonal antibody S-60-10.

[0247] In some embodiments, the anti-Sortilin antibody is anti-Sortilin monoclonal antibody S-60-11. In some embodiments, the anti-Sortilin antibody is an isolated antibody which binds essentially the same Sortilin epitope as S-60-11. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region of monoclonal antibody S-60-11. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the light chain variable region of monoclonal antibody S-60-11. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region and the light chain variable region of monoclonal antibody S-60-11.

[0248] In some embodiments, the anti-Sortilin antibody is anti-Sortilin monoclonal antibody S-60-12. In some embodiments, the anti-Sortilin antibody is an isolated antibody which binds essentially the same Sortilin epitope as S-60-12. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region of monoclonal antibody S-60-12. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the light chain variable region of monoclonal antibody S-60-12. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region and the light chain variable region of monoclonal antibody S-60-12.

[0249] In some embodiments, the anti-Sortilin antibody is anti-Sortilin monoclonal antibody S-60-13. In some embodiments, the anti-Sortilin antibody is an isolated antibody which binds essentially the same Sortilin epitope as S-60-13. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region of monoclonal antibody S-60-13. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the light chain variable region of monoclonal antibody S-60-13. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region and the light chain variable region of monoclonal antibody S-60-13.

[0250] In some embodiments, the anti-Sortilin antibody is anti-Sortilin monoclonal antibody S-60-14. In some embodiments, the anti-Sortilin antibody is an isolated antibody which binds essentially the same Sortilin epitope as S-60-14. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region of monoclonal antibody S-60-14. In some embodiments, the

anti-Sortilin antibody is an isolated antibody comprising the light chain variable region of monoclonal antibody S-60-14. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region and the light chain variable region of monoclonal antibody S-60-14.

[0251] In some embodiments, the anti-Sortilin antibody is anti-Sortilin monoclonal antibody S-60-15. In some embodiments, the anti-Sortilin antibody is an isolated antibody which binds essentially the same Sortilin epitope as S-60-15. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region of monoclonal antibody S-60-15. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the light chain variable region of monoclonal antibody S-60-15. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region and the light chain variable region of monoclonal antibody S-60-15.

[0252] In some embodiments, the anti-Sortilin antibody is anti-Sortilin monoclonal antibody S-60-15.1. In some embodiments, the anti-Sortilin antibody is an isolated antibody which binds essentially the same Sortilin epitope as S-60-15.1. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region of monoclonal antibody S-60-15.1. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the light chain variable region of monoclonal antibody S-60-15.1. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region and the light chain variable region of monoclonal antibody S-60-15.1.

[0253] In some embodiments, the anti-Sortilin antibody is anti-Sortilin monoclonal antibody S-60-16. In some embodiments, the anti-Sortilin antibody is an isolated antibody which binds essentially the same Sortilin epitope as S-60-16. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region of monoclonal antibody S-60-16. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the light chain variable region of monoclonal antibody S-60-16. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region and the light chain variable region of monoclonal antibody S-60-16.

[0254] In some embodiments, the anti-Sortilin antibody is anti-Sortilin monoclonal antibody S-60-18. In some embodiments, the anti-Sortilin antibody is an isolated antibody which binds essentially the same Sortilin epitope as S-60-18. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region of monoclonal antibody S-60-18. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the light chain variable region of monoclonal antibody S-60-18. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region and the light chain variable region of monoclonal antibody S-60-18.

[0255] In some embodiments, the anti-Sortilin antibody is anti-Sortilin monoclonal antibody S-60-19. In some embodiments, the anti-Sortilin antibody is an isolated antibody which binds essentially the

same Sortilin epitope as S-60-19. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region of monoclonal antibody S-60-19. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the light chain variable region of monoclonal antibody S-60-19. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region and the light chain variable region of monoclonal antibody S-60-19.

[0256] In some embodiments, the anti-Sortilin antibody is anti-Sortilin monoclonal antibody S-60-24. In some embodiments, the anti-Sortilin antibody is an isolated antibody which binds essentially the same Sortilin epitope as S-60-24. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region of monoclonal antibody S-60-24. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the light chain variable region of monoclonal antibody S-60-24. In some embodiments, the anti-Sortilin antibody is an isolated antibody comprising the heavy chain variable region and the light chain variable region of monoclonal antibody S-60-24.

[0257] In certain embodiments, the anti-Sortilin antibody is an antagonist antibody. In certain embodiments, the anti-Sortilin antibody is an agonist antibody. In some embodiments, anti-Sortilin antibodies of the present disclosure are of the IgG class the IgM class, or the IgA class. In some embodiments, anti-Sortilin antibodies of the present disclosure are of the IgG class and have an IgG1, IgG2, IgG3, or IgG4 isotype.

[0258] Additional anti-Sortilinantibodies, *e.g.*, antibodies that specifically bind to a Sortilin protein of the present disclosure, may be identified, screened, and/or characterized for their physical/chemical properties and/or biological activities by various assays known in the art.

[0259] Certain aspects of the present disclosure relate to the use of two or more anti-Sortilin antibodies that when utilized together display additive or synergistic effects, as compared to utilization of a corresponding single anti-Sortilin antibody.

[0260] In some embodiments, an anti-Sortilin antibody of the present disclosure is an antibody fragment that binds to a human Sortilin protein.

[0261] In some embodiments, an anti-Sortilin antibody of the present disclosure is an antibody fragment that binds to one or more human proteins selected from the group consisting of human Sortilin, a naturally occurring variant of human Sortilin, and a disease variant of human Sortilin.

[0262] In some embodiments, an anti-Sortilin antibody of the present disclosure is antibody fragment, wherein the antibody fragment is an Fab, Fab', Fab'-SH, F(ab')₂, Fv, or scFv fragment.

Antibody frameworks

[0263] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable region comprising one or more (*e.g.*, one or more, two or more, three or more, or all four) framework regions selected from VH FR1, VH FR2, VH FR3, and VH FR4 (as shown in **Tables 20-23**).

In some embodiments, the VH FR1 comprises a sequence of QVQLQESGPGGLVKPSETLSL TCAVSG (SEQ ID NO: 35). In some embodiments, the VH FR2 comprises a sequence of WIRQPPGKGLEWIG (SEQ ID NO: 36). In some embodiments, the VH FR3 comprises the sequence according to Formula VI: $X_1VTISVDTSKNQFSLX_2LSSVTAADTAVYYC$ (SEQ ID NO: 39), wherein X_1 is Q or R, and X_2 is E or K. In some embodiments, VH FR3 comprises a sequence selected from the group consisting of SEQ ID NOs: 37-38. In some embodiments, VH FR4 comprises a sequence of WGQGTTVTVSS (SEQ ID NO: 40). In some embodiments, an antibody comprises a heavy chain variable region comprising a VH FR1 comprising the sequence of SEQ ID NO: 35, a VH FR2 comprising the sequence of SEQ ID NO: 36, a VH FR3 according to Formula VI, and a VH FR4 comprising the sequence of SEQ ID NO: 40.

[0264] In some embodiments, an antibody comprises a heavy chain variable region comprising a VH FR1 comprising the sequence of SEQ ID NO: 35, a VH FR2 comprising the sequence of SEQ ID NO: 36, a VH FR3 comprising the sequence selected from SEQ ID NOs: 37-38, and a VH FR4 comprising the sequence of SEQ ID NO: 40.

[0265] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable region comprising a VH FR1, a VH FR2, a VH FR3, and VH FR4 of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24 (as shown in **Tables 20-23**).

[0266] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable region comprising one or more (*e.g.*, one or more, two or more, three or more, or all four) framework regions selected from VL FR1, VL FR2, VL FR3, and VL FR4 (as shown in **Tables 24-27**).

In some embodiments, the VL FR1 comprises a sequence according to Formula VII:

$DIVMTQSPLSLPVTGPX_1X_2ASISC$ (SEQ ID NO: 44), wherein X_1 is E or G, and X_2 is P or S. In some embodiments, VL FR1 comprises a sequence selected from the group consisting of SEQ ID NOs: 41-43.

In some embodiments, the VL FR2 comprises a sequence according to Formula VIII:

$WYLQKPGQX_1PQLLIY$ (SEQ ID NO: 47), wherein X_1 is S or P. In some embodiments, VL FR2 comprises a sequence selected from the group consisting of SEQ ID NOs: 45-46. In some embodiments, the VL FR3 comprises a sequence according to Formula IX: $GVPDRX_1SGSGSGT$

$DFTLKISR X_2EAEDVGX_3YYC$ (SEQ ID NO: 52), wherein X_1 is F or L, X_2 is A or V, and X_3 is V or A.

In some embodiments, VL FR3 comprises a sequence selected from the group consisting of SEQ ID NOs: 48-51. In some embodiments, the VL FR4 comprises a sequence of FGGGTKVEIK (SEQ ID NO: 53). In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable

region comprising a VL FR1 comprising the sequence according to Formula VII, a VL FR2 comprising the sequence according to Formula VIII, a VL FR3 comprising the sequence according to Formula IX, and a VL FR4 comprising the sequence of SEQ ID NO: 53.

[0267] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable region comprising a VL FR1 comprising the sequence selected from SEQ ID NOs: 41-43, a VL FR2 comprising the sequence selected from SEQ ID NOs: 45-46, a VL FR3 comprising the sequence selected from SEQ ID NOs: 48-51, and a VL FR4 comprising the sequence of SEQ ID NO: 53.

[0268] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a light chain variable region comprising a VL FR1, a VL FR2, a VL FR3, and VL FR4 of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24 (as shown in **Tables 24-27**).

[0269] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable region comprising one or more (*e.g.*, one or more, two or more, three or more, or all four) framework regions selected from VH FR1, VH FR2, VH FR3, and VH FR4 (as shown in **Tables 20-23**), and a light chain variable region comprising one or more (*e.g.*, one or more, two or more, three or more, or all four) framework regions selected from VL FR1, VL FR2, VL FR3, and VL FR4 (as shown in **Tables 24-27**). In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable region comprising a

[0270] VH FR1 comprising the sequence of SEQ ID NO: 35, a VH FR2 comprising the sequence of SEQ ID NO: 36, a VH FR3 according to Formula VI, and a VH FR4 comprising the sequence of SEQ ID NO: 40; and a light chain variable region comprising a VL FR1 comprising the sequence according to Formula VII, a VL FR2 comprising the sequence according to Formula VIII, a VL FR3 comprising the sequence according to Formula IX, and a VL FR4 comprising the sequence of SEQ ID NO: 53. In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable region comprising a VH FR1 comprising the sequence of SEQ ID NO: 35, a VH FR2 comprising the sequence of SEQ ID NO: 36, a VH FR3 comprising the sequence selected from SEQ ID NOs: 37-38, and a VH FR4 comprising the sequence of SEQ ID NO: 40; a light chain variable region comprising a VL FR1 comprising the sequence selected from SEQ ID NOs: 41-43, a VL FR2 comprising the sequence selected from SEQ ID NOs: 45-46, a VL FR3 comprising the sequence selected from SEQ ID NOs: 48-51, and a VL FR4 comprising the sequence of SEQ ID NO: 53.

[0271] In some embodiments, anti-Sortilin antibodies of the present disclosure comprise a heavy chain variable region comprising a VH FR1, a VH FR2, a VH FR3, and VH FR4 of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24 (as shown in **Tables 20-23**), and a light chain variable region comprising a VL FR1, a VL FR2, a VL FR3, and VL FR4 of antibody S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16; S-60-18, S-60-19, or S-60-24 (as shown in **Tables 24-27**).

Anti-Sortilin Antibody Activities

[0272] In certain aspects of any of the anti-Sortilin antibodies, anti-Sortilin antibodies of the present disclosure can inhibit one or more activities of a Sortilin protein, including, but not limited to, decreasing cellular levels of Sortilin (*e.g.*, cell surface levels of Sortilin, intracellular levels of Sortilin, and/or total levels of Sortilin); increasing Progranulin levels (*e.g.*, extracellular levels of Progranulin and/or cellular levels of Progranulin); and inhibiting the interaction (*e.g.*, binding) between Progranulin and Sortilin. As contemplated herein, anti-Sortilin antibodies of the present disclosure may inhibit additional activities of a Sortilin protein, including but not limited to inhibiting interaction (*e.g.*, binding) with one or more of pro-neurotrophins of the present disclosure (pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, *etc.*), neurotrophins of the present disclosure (neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, Sortilin propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), and receptor associated protein (RAP), decreasing secretion of PCSK9, decreasing production of beta amyloid peptide..

[0273] In certain embodiments, the present disclosure provides an anti-Sortilin antibody, wherein (a) the anti-Sortilin antibody increases extracellular levels of Progranulin, decreases cellular levels of Sortilin, inhibits interaction between Sortilin and Progranulin, or any combination thereof; (b) the anti-Sortilin antibody decreases cell surface levels of Sortilin, increases extracellular levels of Progranulin, inhibits interaction between Sortilin and Progranulin, or any combination thereof; (c) the anti-Sortilin antibody decreases cell surface levels of Sortilin, decreases intracellular levels of Sortilin, decreases total levels of Sortilin, or any combination thereof; (d) the anti-Sortilin antibody induces Sortilin degradation, Sortilin cleavage, Sortilin internalization, Sortilin down regulation, or any combination thereof; (e)

the anti-Sortilin antibody decreases cellular levels of Sortilin and inhibits the interaction between Sortilin and Progranulin; (f) the anti-Sortilin antibody decreases cellular levels of Sortilin and increases cellular levels of Progranulin; and/or (g) the anti-Sortilin antibody increases the effective concentration of Progranulin.

[0274] In certain embodiments, the present disclosure provides an anti-Sortilin antibody, wherein the anti-Sortilin antibody decreases cell surface levels of Sortilin, increases extracellular levels of Progranulin, inhibits interaction between Sortilin and Progranulin, or any combination thereof.

[0275] In some embodiments, an anti-Sortilin antibody of the present disclosure (a) reduces cell surface levels of Sortilin with a half maximal effective concentration (EC_{50}) that is less than 150 pM, as measured by flow cytometry; (b) reduces cell surface levels of Sortilin by more than about 50% at 1.25 nM IgG, by more than about 80% at 0.63 nM IgG, or by more than about 69% at 150 nM IgG relative to control, as measured by flow cytometry; increases Progranulin secretion by more than about 1.13 fold over control at 0.63 nM IgG, or by more than about 1.22 fold over control at 50 nM IgG, as measured by standard ELISA; blocks binding of Progranulin to Sortilin with a half maximal effective concentration (EC_{50}) that is less than .325 nM, as measured by flow cytometry; (e) blocks binding of Progranulin to Sortilin by more than about 88% at 50 nM IgG, or by more than about 27.5% at 150 nM IgG relative to control, as measured by flow cytometry; or (f) any combination thereof (*see, e.g., Examples 2-4*).

[0276] In some embodiments, an anti-Sortilin antibody of the present disclosure (a) reduces cell surface levels of Sortilin with a half maximal effective concentration (EC_{50}) that is less than 681 pM, as measured by flow cytometry; (b) reduces cell surface levels of Sortilin by more than about 40% at 1.25 nM IgG, by more than about 29% at 0.6 nM IgG, or by more than about 62% at 150 nM IgG relative to control, as measured by flow cytometry; (c) increases Progranulin secretion by more than about 1.11 fold over control at 0.63 nM IgG, or by more than about 1.75 fold over control at 50 nM IgG, as measured by standard ELISA; (d) blocks binding of Progranulin to Sortilin with a half maximal effective concentration (EC_{50}) that is less than 0.751 nM, as measured by flow cytometry; (e) blocks binding of Progranulin to Sortilin by more than about 90% at 50 nM IgG, or by more than about 95% at 150 nM IgG relative to control, as measured by flow cytometry; or (f) any combination thereof (*see, e.g., Examples 2-4*).

Decreasing Sortilin levels

[0277] In some embodiments, anti-Sortilin antibodies of the present disclosure bind to a Sortilin protein of the present disclosure expressed on the surface of a cell and modulate (e.g., induce or inhibit) one or more Sortilin activities of the present disclosure after binding to the surface-expressed Sortilin protein.

[0278] In some embodiments, anti-Sortilin antibodies of the present disclosure decrease cellular levels of Sortilin *in vitro*. In some embodiments, anti-Sortilin antibodies of the present disclosure may

decrease cellular levels of Sortilin *in vivo* (e.g., in the brain, and/or peripheral organs of an individual). In some embodiments, a decrease in cellular levels of Sortilin comprises a decrease in cell surface levels of Sortilin. As used herein, an anti-Sortilin antibody decreases cell surface levels of Sortilin if it induces a decrease at saturating antibody concentrations (e.g., 0.6 nM, 0.63 nM, 1.25 nM, 50 nM or 150 nM) and/or relative to a control antibody (e.g. an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60) in cell surface levels of Sortilin as measured by any *in vitro* cell-based assays or suitable *in vivo* model described herein or known in the art. In some embodiments, a decrease in cellular levels of Sortilin comprises a decrease in intracellular levels of Sortilin. As contemplated herein, an anti-Sortilin antibody decreases intracellular levels of Sortilin if it induces a decrease at saturating antibody concentrations and/or relative to a control antibody (e.g. an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60) in intracellular levels of Sortilin as measured by any *in vitro* cell-based assays or suitable *in vivo* model described herein or known in the art. In some embodiments, a decrease in cellular levels of Sortilin comprises a decrease in total levels of Sortilin. As contemplated herein, an anti-Sortilin antibody decreases total levels of Sortilin if it induces a decrease at saturating antibody concentrations and/or relative to a control antibody (e.g. an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60) in total levels of Sortilin as measured by any *in vitro* cell-based assays or suitable *in vivo* model described herein or known in the art.

[0279] As used herein, levels of Sortilin may refer to expression levels of the gene encoding Sortilin; to expression levels of one or more transcripts encoding Sortilin; to expression levels of Sortilin protein; and/or to the amount of Sortilin protein present within cells and/or on the cell surface. Any methods known in the art for measuring levels of gene expression, transcription, translation, and/or protein abundance or localization may be used to determine the levels of Sortilin.

[0280] Cellular levels of Sortilin may refer to, without limitation, cell surface levels of Sortilin, intracellular levels of Sortilin, and total levels of Sortilin. In some embodiments, a decrease in cellular levels of Sortilin comprises decrease in cell surface levels of Sortilin. In some embodiments, anti-Sortilin antibodies of the present disclosure that decrease cellular levels of Sortilin (e.g., cell surface levels of Sortilin) have one or more of the following characteristics: (1) inhibits or reduces one or more Sortilin activities; (2) the ability to inhibit or reduce binding of a Sortilin to one or more of its ligands; (3) the ability to reduce Sortilin expression in Sortilin-expressing cells; (4) the ability to interact, bind, or recognize a Sortilin protein; (5) the ability to specifically interact with or bind to a Sortilin protein; and (6) the ability to treat, ameliorate, or prevent any aspect of a disease or disorder described or contemplated herein.

[0281] In some embodiments, an isolated anti-Sortilin antibody of the present disclosure induces downregulation of Sortilin. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure induces cleavage of Sortilin. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure induces internalization of Sortilin. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure induces shedding of Sortilin. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure induces degradation of Sortilin. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure induces desensitization of Sortilin. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure acts as a ligand mimetic to transiently activate Sortilin. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure acts as a ligand mimetic and transiently activates Sortilin before inducing a decrease in cellular levels of Sortilin and/or inhibition of interaction (*e.g.*, binding) between Sortilin and one or more Sortilin ligands. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure acts as a ligand mimetic and transiently activates Sortilin before inducing degradation of Sortilin. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure acts as a ligand mimetic and transiently activates Sortilin before inducing cleavage of Sortilin. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure acts as a ligand mimetic and transiently activates Sortilin before inducing internalization of Sortilin. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure acts as a ligand mimetic and transiently activates Sortilin before inducing shedding of Sortilin. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure acts as a ligand mimetic and transiently activates Sortilin before inducing downregulation of Sortilin expression. In some embodiments, an isolated anti-Sortilin antibody of the present disclosure acts as a ligand mimetic and transiently activates Sortilin before inducing desensitization of Sortilin.

[0282] In certain embodiments, anti-Sortilin antibodies of the present disclosure may decrease cellular levels of Sortilin (*e.g.*, cell surface levels of Sortilin, intracellular levels of Sortilin, and/or total levels of Sortilin) by inducing Sortilin degradation. Accordingly, in some embodiments, anti-Sortilin antibodies of the present disclosure induce Sortilin degradation.

[0283] Anti-Sortilin antibodies of the present disclosure may decrease cellular levels (*e.g.*, cell surface levels) of Sortilin with a half-maximal effective concentration (EC_{50}) (*e.g.*, when measured *in vitro*) in the picomolar range. In certain embodiments, the EC_{50} of the antibody is less than about 680.9 pM. In certain embodiments, the EC_{50} of the antibody is about 72.58 pM to about 680.9 nM. In certain embodiments, the EC_{50} of the antibody is about 103.6 pM to about 680.9 nM. In certain embodiments, the EC_{50} of the antibody is less than about 600 pM, 500 pM, 400 pM, 300 pM, 200 pM, 100 pM, 50 pM, 40 pM, 30 pM, 20 pM, 10 pM, 1 pM, or 0.5 pM.

[0284] In some embodiments, the EC₅₀ of the antibody is less than about or equal to about 675 pM, 650 pM, 625 pM, 600 pM, 575 pM, 550 pM, 525 pM, 500 pM, 475 pM, 450 pM, 425 pM, 400 pM, 375 pM, 350pM, 325 pM, 300 pM, 275 pM, 250 pM, 225 pM, 200 pM, 175 pM, 150 pM, 125 pM, 100 pM, 90 pM, 80 pM, 70 pM, 60 pM, 50 pM, 40 pM, 30 pM, 20 pM, 10 pM, 9 pM, 8 pM, 7 pM, 6 pM, 5 pM, 4 pM, 3 pM, 2 pM, 1 pM, or 0.5 pM.

[0285] In some embodiments, the EC₅₀ of the antibody is less than about 680.9 pM. In some embodiments, the EC₅₀ of the antibody is greater than about or equal to about 0.1 pM, 0.5pM, 1 pM, 10 pM, 20 pM, 30 pM, 40 pM, 50 pM, 60 pM, 70 pM, 80 pM, 90 pM, 100 pM, 125 pM, 150 pM, 175 pM, 200 pM, 225 pM, 250 pM, 275 pM, 300 pM, 325 pM, 350 pM, 375 pM, 400 pM, 425 pM, 450 pM, 475 pM, 500 pM, 525 pM, 550 pM, 575 pM, 600 pM, 625 pM, 650 pM, 675 pM. That is, the EC₅₀ of the antibody can be any of a range having an upper limit of about 675 pM, 650 nM, 650 pM, 625 pM, 600 pM, 575 pM, 550 pM, 525 pM, 500 pM, 475 pM, 450 pM, 425 pM, 400 pM, 375 pM, 350pM, 325 pM, 300 pM, 275 pM, 250 pM, 225 pM, 200 pM, 175 pM, 150 pM, 125 pM, 100 pM, 90 pM, 80 pM, 70 pM, 60 pM, 50 pM, 40 pM, 30 pM, 20 pM, 10 pM, 1 pM, or 0.5 pM, and an independently selected lower limit of about 0.1 pM, 0.5pM, 1 pM, 10 pM, 20 pM, 30 pM, 40 pM, 50 pM, 60 pM, 70 pM, 80 pM, 90 pM, 100 pM, 125 pM, 150 pM, 175 pM, 200 pM, 225 pM, 250 pM, 275 pM, 300 pM, 325 pM, 350 pM, 375 pM, 400 pM, 425 pM, 450 pM, 475 pM, 500 pM, 525 pM, 550 pM, 575 pM, 600 pM, 625 pM, 650 pM, or 675 pM, wherein the lower limit is less than the upper limit. In some embodiments, the EC₅₀ of the antibody is any of about 1 pM, 2 pM, 3 pM, 4 pM, 5 pM, 6 pM, 7 pM, 8 pM, 9 pM, 10 pM, 15 pM, 20 pM, 25 pM, 30 pM, 35 pM, 40 pM, 45 pM, 50 pM, 55 pM, 60 pM, 65 pM, 70 pM, 75 pM, 80 pM, 85 pM, 90 pM, 95 pM, 100 pM, 105 pM, 110 pM, 115 pM, 120 pM, 125 pM, 130 pM, 135 pM, 140 pM, 145 pM, 150 pM, 155 pM, 160 pM, 165 pM, 170 pM, 175 pM, 180 pM, 185 pM, 190 pM, 195 pM, or 200 pM.

[0286] In some embodiments, an anti-Sortilin antibody of the present disclosure reduces cell surface levels of Sortilin with a half maximal effective concentration (EC₅₀) that is less than 150 pM, as measured by flow cytometry. In some embodiments, the EC₅₀ of an anti-Sortilin antibody of the present disclosure is about 103.6 pM. In some embodiments, the EC₅₀ of an anti-Sortilin antibody of the present disclosure is about 72.58 pM.

[0287] In some embodiments, an anti-Sortilin antibody of the present disclosure reduces cell surface levels of Sortilin by more than about 40% at 1.25 nM IgG or by more than about 80% at 0.63 nM IgG, as measured by flow cytometry. In some embodiments, an anti-Sortilin antibody of the present disclosure reduces cell surface levels of Sortilin by about 60.92% at 1.25 nM IgG, as measured by flow cytometry. In some embodiments, an anti-Sortilin antibody of the present disclosure reduces cell surface levels of Sortilin by about 69.3% at 150 nM IgG, as measured by flow cytometry. In some embodiments, an anti-

Sortilin antibody of the present disclosure reduces cell surface levels of Sortilin by about 70.3% at 150 nM IgG, as measured by flow cytometry.

[0288] Various methods of measuring antibody EC₅₀ values are known in the art, including, for example, by flow cytometry (*See e.g.*, **Example 2**). In some embodiments, the EC₅₀ is measured *in vitro* using cells engineered to express human Sortilin. In some embodiments, the EC₅₀ is measured at a temperature of approximately 4°C. In some embodiments, the EC₅₀ is measured at a temperature of approximately 25°C. In some embodiments, the EC₅₀ is measured at a temperature of approximately 35°C. In some embodiments, the EC₅₀ is measured at a temperature of approximately 37°C. In some embodiments, the EC₅₀ is determined using a monovalent antibody (*e.g.*, a Fab) or a full-length antibody in a monovalent form. In some embodiments, the EC₅₀ is determined using antibodies containing constant regions that demonstrate enhanced Fc receptor binding. In some embodiments, the EC₅₀ is determined using antibodies containing constant regions that demonstrate reduced Fc receptor binding.

[0289] In some embodiments, anti-Sortilin antibodies of the present disclosure have higher potencies in reducing cell surface levels of Sortilin relative to a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure decrease cellular levels (*e.g.*, cell surface levels) of Sortilin with a lower EC₅₀ (*e.g.*, as measured *in vitro*) than a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure decrease cellular levels (*e.g.*, cell surface levels) of Sortilin with an EC₅₀ that is at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99% lower than the EC₅₀ of a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure decrease cellular levels (*e.g.*, cell surface levels) of Sortilin with an EC₅₀ that is at least about 1-fold, at least about 1.1-fold, at least about 1.5-fold, at least about 2-fold, at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, at least about 10-fold, at least about 12.5-fold, at least about 15-fold, at least about 17.5-fold, at least about 20-fold, at least about 22.5-fold, at least about 25-fold, at least about 27.5-fold, at least about 30-fold, at least about 50-fold, or at least about 100-fold lower than the EC₅₀ of a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60).

[0290] In some embodiments, anti-Sortilin antibodies of the present disclosure have an EC₅₀ that is at least 1.5-fold lower than control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure have an EC₅₀ that is at least 1.1-fold lower than control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60).

[0291] In some embodiments, an anti-Sortilin antibody of the present disclosure (a) reduces cell surface levels of Sortilin with a half maximal effective concentration (EC₅₀) that is less than 681 pM, as measured by flow cytometry; (b) reduces cell surface levels of Sortilin by more than about 40% at 1.25 nM IgG, by more than about 29% at 0.6 nM IgG, or by more than about 62% at 150 nM IgG relative to control, as measured by flow cytometry; (c) increases Progranulin secretion by more than about 1.11 fold over control at 0.63 nM IgG, or by more than about 1.75 fold over control at 50 nM IgG, as measured by standard ELISA; (d) blocks binding of Progranulin to Sortilin with a half maximal effective concentration (EC₅₀) that is less than 0.751 nM, as measured by flow cytometry; (e) blocks binding of Progranulin to Sortilin by more than about 90% at 50 nM IgG, or by more than about 95% at 150 nM IgG relative to control, as measured by flow cytometry; or (f) any combination thereof (*see, e.g.*, **Examples 2-4**).

Increasing Progranulin levels

[0292] In some embodiments, anti-Sortilin antibodies of the present disclosure increase extracellular levels of Progranulin *in vitro*. In some embodiments, anti-Sortilin antibodies of the present disclosure may increase cellular levels of Progranulin or *in vivo* (*e.g.*, in the brain, blood, and/or peripheral organs of an individual). As used herein, an anti-Sortilin antibody increases extracellular levels of Progranulin if it induces an increase at saturating antibody concentrations (*e.g.*, 0.6 nM, 0.63 nM, 1.25 nM, 50 nM or 150 nM) and/or relative to a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60) in extracellular levels of Progranulin as measured by any *in vitro* cell-based assays or in tissue-based (such as brain tissue-based) assays described herein or known in the art. As contemplated herein, an anti-Sortilin antibody increases cellular levels of Progranulin if it induces an increase at saturating antibody concentrations (*e.g.*, 0.6 nM, 0.63 nM, 1.25 nM, 50 nM or 150 nM) and/or relative to a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60) in cellular levels of Progranulin as measured by any *in vitro* cell-based assays or in tissue-based (such as brain tissue-based) assays described herein or known in the art.

[0293] As used herein, levels of Progranulin may refer to expression levels of the gene encoding Progranulin; to expression levels of one or more transcripts encoding Progranulin; to expression levels of Progranulin protein; and/or to the amount of Progranulin protein secreted from cells and/or present within

cells. Any methods known in the art for measuring levels of gene expression, transcription, translation, protein abundance, protein secretion, and/or protein localization may be used to determine the levels of Progranulin.

[0294] As used herein, Progranulin levels may refer to, without limitation, extracellular levels of Progranulin, intracellular levels of Progranulin, and total levels of Progranulin. In some embodiments, an increase in levels of Progranulin comprises an increase in extracellular levels of Progranulin.

[0295] In some embodiments, an anti-Sortilin antibody of the present disclosure increases Progranulin secretion by more than about 1.11 fold over control at 0.63 nM IgG, as measured by standard ELISA. In some embodiments, an anti-Sortilin antibody of the present disclosure increases Progranulin secretion by about 1.42 fold over control at 0.63 nM IgG, as measured by standard ELISA. In some embodiments, an anti-Sortilin antibody of the present disclosure increases Progranulin secretion by more than about 1.75 fold over control at 50 nM IgG, as measured by standard ELISA. In some embodiments, an anti-Sortilin antibody of the present disclosure increases Progranulin secretion by about 1.97 fold over control at 50 nM IgG, as measured by standard ELISA. In some embodiments, an anti-Sortilin antibody of the present disclosure increases Progranulin secretion by about 2.29 fold over control at 50 nM IgG, as measured by standard ELISA.

[0296] Various methods of measuring Progranulin secretion are known in the art, including, for example, by ELISA (*See e.g., Examples 2 and 4*). In some embodiments, the EC_{50} is measured *in vitro* using cells expressing human Sortilin. In some embodiments, Progranulin secretion is determined using a monovalent antibody (*e.g.*, a Fab) or a full-length antibody in a monovalent form. In some embodiments, Progranulin secretion is determined using antibodies containing constant regions that demonstrate enhanced Fc receptor binding. In some embodiments, Progranulin secretion is determined using antibodies containing constant regions that demonstrate reduced Fc receptor binding.

[0297] In some embodiments, anti-Sortilin antibodies of the present disclosure have higher potencies in increasing levels of Progranulin relative to a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure increase levels (*e.g.*, extracellular levels) of Progranulin with a lower EC_{50} (*e.g.*, as measured *in vitro*) than a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure increase levels (*e.g.*, extracellular levels) of Progranulin by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99%

than a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure increase levels (*e.g.*, extracellular levels) of Progranulin by about 1-fold, at least about 1.1-fold, at least about 1.5-fold, at least about 2-fold, at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, at least about 10-fold, at least about 12.5-fold, at least about 15-fold, at least about 17.5-fold, at least about 20-fold, at least about 22.5-fold, at least about 25-fold, at least about 27.5-fold, at least about 30-fold, at least about 50-fold, or at least about 100-fold higher than a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60).

[0298] In some embodiments, anti-Sortilin antibodies of the present disclosure increase Progranulin levels by about 1.1-fold higher than a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure increase Progranulin levels by about 1.3-fold higher than a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60).

[0299] In some embodiments, anti-Sortilin antibodies of the present disclosure increase the effective concentration of Progranulin. The effective concentration of Progranulin refers to the concentration of Progranulin in plasma or cerebrospinal fluid. In some embodiments, an increase in the effective concentration of Progranulin is an increase of greater than 1.5 fold. In some embodiments, the effective concentration of Progranulin is increased for 7-28 days.

Decreasing interaction between Sortilin and binding partner

[0300] In certain embodiments, the anti-Sortilin antibodies inhibit interaction (*e.g.*, binding) between a Sortilin protein of the present disclosure and one or more proteins selected from Progranulin, a pro-neurotrophin, a neurotrophin, pro-neurotrophin-3, neurotrophin-3, pro-neurotrophin-4/5, neurotrophin-4/5, pro-nerve growth factor (pro-NGF), nerve growth factor (NGF), pro-brain-derived neurotrophic factor (pro-BDNF), brain-derived neurotrophic factor (BDNF), neurotensin, p75, Sortilin propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), PCSK9, and receptor associated protein (RAP), and/or naturally occurring variants. In a specific embodiment, anti-Sortilin antibodies of the present disclosure inhibit interaction (*e.g.*, binding) between a Sortilin protein of the present disclosure and Progranulin.

[0301] In some embodiments, the anti-Sortilin antibodies may inhibit interaction (*e.g.*, binding) between a Sortilin protein of the present disclosure and a neurotrophin of the present disclosure, such as a pro-neurotrophin, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, and BDNF. In other embodiments, the anti-Sortilin antibodies may inhibit

interaction (*e.g.*, binding) between a Sortilin protein of the present disclosure and neurotensin. In other embodiments, the anti-Sortilin antibodies inhibit interaction (*e.g.*, binding) between a Sortilin protein of the present disclosure and p75. In other embodiments, the anti-Sortilin antibodies may inhibit interaction (*e.g.*, binding) between a Sortilin protein of the present disclosure and a Sort-pro. In other embodiments, the anti-Sortilin antibodies may inhibit interaction (*e.g.*, binding) between a Sortilin protein of the present disclosure and APP. In other embodiments, the anti-Sortilin antibodies may inhibit the production of the A beta peptide. In other embodiments, the anti-Sortilin antibodies may inhibit the transport and secretion of PCSK9. In other embodiments, the anti-Sortilin antibodies may inhibit interaction (*e.g.*, binding) between a Sortilin protein of the present disclosure and LpL. In other embodiments, the anti-Sortilin antibodies may inhibit interaction (*e.g.*, binding) between a Sortilin protein of the present disclosure and APOA5. In other embodiments, the anti-Sortilin antibodies may inhibit interaction (*e.g.*, binding) between a Sortilin protein of the present disclosure and APOE. In other embodiments, the anti-Sortilin antibodies may inhibit interaction (*e.g.*, binding) between a Sortilin protein of the present disclosure and RAP.

[0302] In some embodiments, anti-Sortilin antibodies of the present disclosure bind to a Sortilin protein of the present disclosure expressed on the surface of cell and the naked antibodies inhibit interaction (*e.g.*, binding) between the Sortilin protein and one or more Sortilin ligands. In some embodiments, anti-Sortilin antibodies of the present disclosure that bind to a Sortilin protein of the present inhibit interaction (*e.g.*, binding) between the Sortilin protein and one or more Sortilin ligands by reducing the effective levels of Sortilin that is available to interact with these proteins either on the cell surface or inside the cell. In some embodiments, anti-Sortilin antibodies of the present disclosure that bind to a Sortilin protein of the present inhibit interaction (*e.g.*, binding) between the Sortilin protein and one or more Sortilin ligands by inducing degradation of Sortilin.

A. Decreasing interaction between Sortilin and Progranulin

[0303] In some embodiments, anti-Sortilin antibodies of the present disclosure increase Progranulin levels and/or decrease cellular levels of Sortilin while blocking (*e.g.* inhibiting) the interaction (*e.g.*, binding) between Sortilin and Progranulin. Accordingly, in some embodiments, anti-Sortilin antibodies of the present disclosure block the interaction (*e.g.*, binding) between Sortilin and Progranulin. As used herein, an anti-Sortilin antibody blocks the interaction (*e.g.*, binding) between Sortilin and Progranulin if it decreases Progranulin binding to Sortilin relative to a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60) at saturating antibody concentrations (*e.g.*, 0.6 nM, 0.63 nM, 1.25 nM, 50 nM or 150 nM) in any *in vitro* assay or cell-based culture assay described herein or known in the art.

[0304] Anti-Sortilin antibodies of the present disclosure may decrease Progranulin binding to Sortilin with a half-maximal effective concentration (EC_{50}) (*e.g.*, when measured *in vitro*) in the picomolar range. In certain embodiments, the EC_{50} of the antibody is less than about 2.2 nM. In certain embodiments, the EC_{50} of the antibody is less than about 1.22 nM. In certain embodiments, the EC_{50} of the antibody is less than about 751 pM. In certain embodiments, the EC_{50} of the antibody is about 325 pM to about 751 nM. In certain embodiments, the EC_{50} of the antibody is about 405 pM to about 751 nM. In certain embodiments, the EC_{50} of the antibody is about 588 pM to about 751 nM. In certain embodiments, the EC_{50} of the antibody is less than about 2.2 nM, 2.1 nM, 2.0 nM, 1.9 nM, 1.8 nM, 1.7 nM, 1.6 nM, 1.5 nM, 1.4 nM, 1.3 nM, 1.2 nM, 1.1 nM, 1.0 nM, 900 pM, 800 pM, 700 pM, 600 pM, 500 pM, 400 pM, 300 pM, 200 pM, 100 pM, 50 pM, 40 pM, 30 pM, 20 pM, 10 pM, 1 pM, or 0.5 pM.

[0305] In some embodiments, the EC_{50} of the antibody for decreasing Progranulin binding to Sortilin is less than about or equal to about 2.2 nM, 2.1 nM, 2.0 nM, 1.9 nM, 1.8 nM, 1.7 nM, 1.6 nM, 1.5 nM, 1.4 nM, 1.3 nM, 1.2 nM, 1.1 nM, 1.0 nM, 900 pM, 800 pM, 700 pM, 600 pM, 500 pM, 475 pM, 450 pM, 425 pM, 400 pM, 375 pM, 350 pM, 325 pM, 300 pM, 275 pM, 250 pM, 225 pM, 200 pM, 175 pM, 150 pM, 125 pM, 100 pM, 90 pM, 80 pM, 70 pM, 60 pM, 50 pM, 40 pM, 30 pM, 20 pM, 10 pM, 9 pM, 8 pM, 7 pM, 6 pM, 5 pM, 4 pM, 3 pM, 2 pM, 1 pM, or 0.5 pM.

[0306] In some embodiments, the EC_{50} of an anti-Sortilin antibody of the present disclosure is about 1.22 nM. In some embodiments, the EC_{50} of an anti-Sortilin antibody of the present disclosure is about 588 pM. In some embodiments, the EC_{50} of an anti-Sortilin antibody of the present disclosure is about 405 pM. In some embodiments, the EC_{50} of an anti-Sortilin antibody of the present disclosure is about 325 pM.

[0307] Various methods of measuring antibody EC_{50} values are known in the art, including, for example, by flow cytometry (*See e.g.*, **Example 3**). In some embodiments, the EC_{50} for decreasing Progranulin binding to Sortilin is measured *in vitro* using cells expressing human Sortilin. In some embodiments, the EC_{50} is measured at a temperature of approximately 4°C. In some embodiments, the EC_{50} is measured at a temperature of approximately 25°C. In some embodiments, the EC_{50} is measured at a temperature of approximately 35°C. In some embodiments, the EC_{50} is measured at a temperature of approximately 37°C. In some embodiments, the EC_{50} for decreasing Progranulin binding to Sortilin is determined using a monovalent antibody (*e.g.*, a Fab) or a full-length antibody in a monovalent form. In some embodiments, the EC_{50} is determined using antibodies containing constant regions that demonstrate enhanced Fc receptor binding. In some embodiments, the EC_{50} for decreasing Progranulin binding to Sortilin is determined using antibodies containing constant regions that demonstrate reduced Fc receptor binding.

[0308] In some embodiments, anti-Sortilin antibodies of the present disclosure have higher potencies in reducing Progranulin binding to Sortilin relative to a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure decrease Progranulin binding to Sortilin with a lower EC₅₀ (*e.g.*, as measured *in vitro*) than a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure decrease Progranulin binding to Sortilin with an EC₅₀ that is at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99% lower than the EC₅₀ of a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure decrease Progranulin binding to Sortilin with an EC₅₀ that is at least about 1-fold, at least about 1.1-fold, at least about 1.5-fold, at least about 2-fold, at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, at least about 10-fold, at least about 12.5-fold, at least about 15-fold, at least about 17.5-fold, at least about 20-fold, at least about 22.5-fold, at least about 25-fold, at least about 27.5-fold, at least about 30-fold, at least about 50-fold, or at least about 100-fold lower than the EC₅₀ of a control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60).

[0309] In some embodiments, anti-Sortilin antibodies of the present disclosure have an EC₅₀ that is at least 1.3-fold lower than control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure have an EC₅₀ that is at least 1.8-fold lower than control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure have an EC₅₀ that is at least 1.9-fold lower than control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60). In some embodiments, anti-Sortilin antibodies of the present disclosure have an EC₅₀ that is at least 2.3-fold lower than control antibody (*e.g.* an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60).

[0310] Any *in vitro* cell-based assays or suitable *in vivo* model described herein or known in the art may be used to measure inhibition or reduction of interaction (*e.g.*, binding) between Sortilin and one or

more Sortilin ligands. In some embodiments, anti-Sortilin antibodies of the present disclosure inhibit or reduce interaction (*e.g.*, binding) between Sortilin and one or more Sortilin ligands by reducing Sortilin expression (*e.g.*, by reducing cell surface levels of Sortilin). In some embodiments, anti-Sortilin antibodies of the present disclosure inhibit or reduce interaction (*e.g.*, binding) between Sortilin and one or more Sortilin ligands by at least 21%, at least 22%, at least 23%, at least 24%, at least 25%, at least 26%, at least 27%, at least 28%, at least 29%, at least 30%, at least 31%, at least 32%, at least 33%, at least 34%, at least 35%, at least 36%, at least 37%, at least 38%, at least 39%, at least 40%, at least 41%, at least 42%, at least 43%, at least 44%, at least 45%, at least 46%, at least 47%, at least 48%, at least 49%, at least 50%, at least 51%, at least 52%, at least 53%, at least 54%, at least 55%, at least 56%, at least 57%, at least 58%, at least 59%, at least 60%, at least 61%, at least 62%, at least 63%, at least 64%, at least 65%, at least 66%, at least 67%, at least 68%, at least 69%, at least 70%, at least 71%, at least 72%, at least 73%, at least 74%, at least 75%, at least 76%, at least 77%, at least 78%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or more at saturating antibody concentrations utilizing any *in vitro* assay or cell-based culture assay described herein or known in the art..

[0311] In some embodiments, an anti-Sortilin antibody of the present disclosure blocks Progranulin binding to Sortilin by more than about 90% at 50 nM IgG or by more than about 96% at 150 nM IgG, as measured by flow cytometry. In some embodiments, an anti-Sortilin antibody of the present disclosure blocks Progranulin binding to Sortilin by about 90.74% at 50 nM IgG, as measured by flow cytometry. In some embodiments, an anti-Sortilin antibody of the present disclosure blocks Progranulin binding to Sortilin by about 96.5% at 150 nM IgG, as measured by flow cytometry. In some embodiments, an anti-Sortilin antibody of the present disclosure blocks Progranulin binding to Sortilin by about 96.9% at 150 nM IgG, as measured by flow cytometry.

Decreasing expression of pro-inflammatory mediators

[0312] In some embodiments, anti-Sortilin antibodies of the present disclosure may decrease the expression of pro-inflammatory mediators after binding to a Sortilin protein expressed in a cell.

[0313] As used herein, pro-inflammatory mediators are proteins involved either directly or indirectly (*e.g.*, by way of pro-inflammatory signaling pathways) in a mechanism that induces, activates, promotes, or otherwise decreases an inflammatory response. Any method known in the art for identifying and characterizing pro-inflammatory mediators may be used.

[0314] Examples of pro-inflammatory mediators include, without limitation, cytokines, such as type I and II interferons, IL-6, IL12p70, IL12p40, IL-1 β , TNF- α , IL-8, CRP, IL-20 family members, IL-33, LIF, OSM, CNTF, GM-CSF, IL-11, IL-12, IL-17, IL-18, and CRP. Further examples of pro-

inflammatory mediators include, without limitation, chemokines, such as CXCL1, CCL2, CCL3, CCL4, and CCL5.

[0315] In some embodiments, the anti-Sortilin antibodies of the present disclosure may decrease functional expression and/or secretion of pro-inflammatory mediators, IL-6, IL12p70, IL12p40, IL-1 β , TNF- α , CXCL1, CCL2, CCL3, CCL4, and CCL5. In certain embodiments, decreased expression of the pro-inflammatory mediators occurs in macrophages, dendritic cells, monocytes, osteoclasts, Langerhans cells of skin, Kupffer cells, T cells, and/or microglial cells. Decreased expression may include, without limitation, a decrease in gene expression, a decrease in transcriptional expression, or a decrease in protein expression. Any method known in the art for determining gene, transcript (*e.g.*, mRNA), and/or protein expression may be used. For example, Northern blot analysis may be used to determine pro-inflammatory mediator gene expression levels, RT-PCR may be used to determine the level of pro-inflammatory mediator transcription, and Western blot analysis may be used to determine pro-inflammatory mediator protein levels.

[0316] As used herein, a pro-inflammatory mediator may have decreased expression if its expression in one or more cells of a subject treated with a Sortilin agent, such as an agonist anti-Sortilin antibody of the present disclosure is more than the expression of the same pro-inflammatory mediator expressed in one or more cells of a corresponding subject that is not treated with the agonist anti-Sortilin antibody. In some embodiments, the anti-Sortilin antibody of the present disclosure may decrease pro-inflammatory mediator expression in one or more cells of a subject by at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 100%, at least 110%, at least 115%, at least 120%, at least 125%, at least 130%, at least 135%, at least 140%, at least 145%, at least 150%, at least 160%, at least 170%, at least 180%, at least 190%, or at least 200% for example, as compared to pro-inflammatory mediator expression in one or more cells of a corresponding subject that is not treated with the anti-Sortilin antibody. In other embodiments, the anti-Sortilin antibody may decrease pro-inflammatory mediator expression in one or more cells of a subject by at least 1.5 fold, at least 1.6 fold, at least 1.7 fold, at least 1.8 fold, at least 1.9 fold, at least 2.0 fold, at least 2.1 fold, at least 2.15 fold, at least 2.2 fold, at least 2.25 fold, at least 2.3 fold, at least 2.35 fold, at least 2.4 fold, at least 2.45 fold, at least 2.5 fold, at least 2.55 fold, at least 3.0 fold, at least 3.5 fold, at least 4.0 fold, at least 4.5 fold, at least 5.0 fold, at least 5.5 fold, at least 6.0 fold, at least 6.5 fold, at least 7.0 fold, at least 7.5 fold, at least 8.0 fold, at least 8.5 fold, at least 9.0 fold, at least 9.5 fold, or at least 10 fold, for example, as compared to pro-inflammatory mediator expression in one or more cells of a corresponding subject that is not treated with the anti-Sortilin antibody.

In some embodiments, an anti-Sortilin antibody according to any of the above embodiments may incorporate any of the features, singly or in combination, as described in Sections 1-7 below:

(1) Anti-Sortilin antibody binding affinity

[0317] In some embodiments of any of the antibodies provided herein, the antibody has a dissociation constant (K_D) of $< 1 \mu\text{M}$, $< 100 \text{ nM}$, $< 10 \text{ nM}$, $< 1 \text{ nM}$, $< 0.1 \text{ nM}$, $< 0.01 \text{ nM}$, or $< 0.001 \text{ nM}$ (*e.g.*, 10^{-8} M or less, *e.g.*, from 10^{-8} M to 10^{-13} M , *e.g.*, from 10^{-9} M to 10^{-13} M).

[0318] Anti-Sortilin antibodies of the present disclosure may have nanomolar or even picomolar affinities for the target antigen (*e.g.*, human Sortilin or mammalian Sortilin). In certain embodiments, the binding affinity of an anti-Sortilin antibody of the present disclosure for target antigen (*e.g.*, human Sortilin or mammalian Sortilin) is measured by the dissociation constant, K_D . Dissociation constants may be determined through any analytical technique, including any biochemical or biophysical technique such as fluorescent activated cell sorting (FACS), flow cytometry, enzyme-linked immunosorbent assay (ELISA), surface plasmon resonance (SPR), BioLayer interferometry (*see, e.g.*, Octet System by ForteBio), meso scale discover (*see, e.g.*, MSD-SET), isothermal titration calorimetry (ITC), differential scanning calorimetry (DSC), circular dichroism (CD), stopped-flow analysis, and colorimetric or fluorescent protein melting analyses; or a cell binding assay. In some embodiments, the K_D for Sortilin is determined at a temperature of approximately 25°C . In some embodiments, the dissociation constant (K_D) may be measured at 4°C or room temperature utilizing, for example, FACS or BioLayer interferometry assay.

[0319] In some embodiments, the K_D for Sortilin is determined at a temperature of approximately 4°C . In some embodiments, the K_D is determined using a monovalent antibody (*e.g.*, a Fab) or a full-length antibody in a monovalent form. In some embodiments, the K_D is determined using a bivalent antibody and monomeric recombinant Sortilin protein.

[0320] In certain embodiments, the K_D of an anti-Sortilin antibody of the present disclosure for human Sortilin, mammalian Sortilin, or both, is measured using FACS as described herein (*see, e.g.*, **Examples 1 and 4**). In certain embodiments, the K_D of an anti-Sortilin antibody of the present disclosure for human Sortilin, mammalian Sortilin, or both, is measured using BioLayer Interferometry as described herein (*see, e.g.*, **Example 4**).

[0321] In some embodiments, the anti-Sortilin antibody has a dissociation constant (K_D) for human Sortilin that is up to 2.5-fold lower than an anti-Sortilin antibody comprising a heavy chain variable region comprising the sequence of SEQ ID NO: 56 and a light chain variable region comprising the sequence of SEQ ID NO: 79, wherein the K_D is determined by FACS (*see, e.g.*, **Example 1**). In some embodiments, the anti-Sortilin antibody has a dissociation constant (K_D) for human Sortilin that ranges from about $1.10\text{E-}8 \text{ M}$ to about $4.68\text{E-}10 \text{ M}$ wherein the K_D is determined by FACS (*see, e.g.*, **Example**

1), or about 270 to about 2910 pM wherein the K_D is determined by Bio-layer interferometry (*see, e.g.,*

Example 4).

[0322] In certain embodiments, the K_D of an anti-Sortilin antibody of the present disclosure for human Sortilin, mammalian Sortilin, or both, may be less than 100 nM, less than 90 nM, less than 80 nM, less than 70 nM, less than 60 nM, less than 50 nM, less than 40 nM, less than 30 nM, less than 20 nM, less than 10 nM, less than 9 nM, less than 8 nM, less than 7 nM, less than 6 nM, less than 5 nM, less than 4 nM, less than 3 nM, less than 2 nM, less than 1 nM, less than 0.5 nM, less than 0.1 nM, less than 0.09 nM, less than 0.08 nM, less than 0.07 nM, less than 0.06 nM, less than 0.05 nM, less than 0.04 nM, less than 0.03 nM, less than 0.02 nM, less than 0.01 nM, less than 0.009 nM, less than 0.008 nM, less than 0.007 nM, less than 0.006 nM, less than 0.005 nM, less than 0.004 nM, less than 0.003 nM, less than 0.002 nM, less than 0.001 nM, or less than 0.001 nM.

[0323] The dissociation constants (K_D) of anti-Sortilin antibodies for human Sortilin, mammalian Sortilin, or both, may be less than 10 nM, less than 9.5 nM, less than 9 nM, less than 8.5 nM, less than 8 nM, less than 7.5 nM, less than 7 nM, less than 6.9 nM, less than 6.8 nM, less than 6.7 nM, less than 6.6 nM, less than 6.5 nM, less than 6.4 nM, less than 6.3 nM, less than 6.2 nM, less than 6.1 nM, less than 6 nM, less than 5.5 nM, less than 5 nM, less than 4.5 nM, less than 4 nM, less than 3.5 nM, less than 3 nM, less than 2.5 nM, less than 2 nM, less than 1.5 nM, less than 1 nM, less than 0.95 nM, less than 0.9 nM, less than 0.89 nM, less than 0.88 nM, less than 0.87 nM, less than 0.86 nM, less than 0.85 nM, less than 0.84 nM, less than 0.83 nM, less than 0.82 nM, less than 0.81 nM, less than 0.8 nM, less than 0.75 nM, less than 0.7 nM, less than 0.65 nM, less than 0.64 nM, less than 0.63 nM, less than 0.62 nM, less than 0.61 nM, less than 0.6 nM, less than 0.55 nM, less than 0.5 nM, less than 0.45 nM, less than 0.4 nM, less than 0.35 nM, less than 0.3 nM, less than 0.29 nM, less than 0.28 nM, less than 0.27 nM, less than 0.26 nM, less than 0.25 nM, less than 0.24 nM, less than 0.23 nM, less than 0.22 nM, less than 0.21 nM, less than 0.2 nM, less than 0.15 nM, less than 0.1 nM, less than 0.09 nM, less than 0.08 nM, less than 0.07 nM, less than 0.06 nM, less than 0.05 nM, less than 0.04 nM, less than 0.03 nM, less than 0.02 nM, less than 0.01 nM, less than 0.009 nM, less than 0.008 nM, less than 0.007 nM, less than 0.006 nM, less than 0.005 nM, less than 0.004 nM, less than 0.003 nM, less than 0.002 nM, or less than 0.001 nM.

[0324] In certain embodiments, the dissociation constant (K_D) of the antibody for Sortilin is from about 0.560 nM to about 1.63 nM, for example when the K_D is determined by FACS. In certain embodiments, the dissociation constant (K_D) of the antibody for Sortilin is from about 0.270 nM to about 2.910 nM, for example when the K_D is determined by BioLayer Interferometry. In some embodiments, the antibody has a dissociation constant (K_D) for human Sortilin, mouse Sortilin, or both, that ranges from about 0.36 nM to about 0.43 nM, or less than 1.02 nM. In some embodiments, the dissociation constant is

less than 1.02 nM. In some embodiments, an anti-Sortilin antibody of the present disclosure has a dissociation constant for human Sortilin of .560 nM or less.

[0325] In one specific embodiment, an anti-Sortilin antibody of the present disclosure has a dissociation constant for human Sortilin of about .560 nM. In one specific embodiment, an anti-Sortilin antibody of the present disclosure has a dissociation constant for human Sortilin of about .423 nM. In one specific embodiment, an anti-Sortilin antibody of the present disclosure has a dissociation constant for human Sortilin of about .365 nM. In one specific embodiment, an anti-Sortilin antibody of the present disclosure has a dissociation constant for human Sortilin of about .344 nM. In one specific embodiment, an anti-Sortilin antibody of the present disclosure has a dissociation constant for human Sortilin of about .298 nM. In one specific embodiment, an anti-Sortilin antibody of the present disclosure has a dissociation constant for human Sortilin of about .270 nM. In another specific embodiment, an anti-Sortilin antibody of the present disclosure has a dissociation constant for human Sortilin of about .260 nM.

[0326] In some embodiments, anti-Sortilin antibodies of the present disclosure have a lower dissociation constant (K_D) for Sortilin than a control anti-Sortilin antibody (*e.g.*, a control anti-Sortilin antibody comprising a heavy chain variable region and a light chain variable region corresponding to S-60. In some embodiments, anti-Sortilin antibodies of the present disclosure have a K_D for a target (*e.g.*, human Sortilin) that is at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99% lower than the K_D of a control anti-Sortilin antibody for the target (*e.g.*, a control anti-Sortilin antibody comprising a heavy chain variable region and a light chain variable region corresponding to S-60. In some embodiments, anti-Sortilin antibodies of the present disclosure have a K_D for a target (*e.g.*, human Sortilin) that is at least about 1-fold, at least about 1.1-fold, at least about 1.5-fold, at least about 2-fold, at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, at least about 10-fold, at least about 12.5-fold, at least about 15-fold, at least about 17.5-fold, at least about 20-fold, at least about 22.5-fold, at least about 25-fold, at least about 27.5-fold, at least about 30-fold, at least about 50-fold, at least about 100-fold, at least about 200-fold, at least about 300-fold, at least about 400-fold, at least about 500-fold, at least about 600-fold, at least about 700-fold, at least about 800-fold, at least about 900-fold, or at least about 1000-fold lower than the K_D of a control anti-Sortilin antibody for the target (*e.g.*, a control anti-Sortilin antibody comprising a heavy chain variable region and a light chain variable region corresponding to S-60.

[0327] In some embodiments, anti-Sortilin antibodies of the present disclosure have a K_D for human Sortilin that is at least 100-fold lower than an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60. In some embodiments, anti-Sortilin antibodies of the present disclosure have a K_D for human Sortilin that is at least 50-fold lower than an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60. In some embodiments, anti-Sortilin antibodies of the present disclosure have a K_D for human Sortilin that is at least 10-fold lower than an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60. In some embodiments, anti-Sortilin antibodies of the present disclosure have a K_D for human Sortilin that is at least 5-fold lower than an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60. In some embodiments, anti-Sortilin antibodies of the present disclosure have a K_D for human Sortilin that is at least 2-fold lower than an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60.

[0328] In a specific embodiment, an anti-Sortilin antibody of the present disclosure has a K_D for human Sortilin that is about 2.79-fold lower than an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60. In another specific embodiment, an anti-Sortilin antibody of the present disclosure has a K_D for human Sortilin that is about 2.05-fold lower than an anti-Sortilin antibody having a heavy chain variable region and a light chain variable region corresponding to S-60.

(2) Antibody fragments

[0329] In some embodiments of any of the antibodies provided herein, the antibody is an antibody fragment. Antibody fragments include, but are not limited to, Fab, Fab', Fab'-SH, F(ab')₂, Fv, and scFv fragments, and other fragments described below. For a review of certain antibody fragments, *see* Hudson et al. *Nat. Med.* 9:129-134 (2003). For a review of scFv fragments, *see, e.g.*, WO 93/16185; and U.S. Patent Nos. 5571894 and 5587458. For discussion of Fab and F(ab')₂ fragments comprising salvage receptor binding epitope residues and having increased *in vivo* half-life, *see* U.S. Patent No. 5869046.

[0330] Diabodies are antibody fragments with two antigen-binding sites that may be bivalent or bispecific. *See, for example*, EP404097; WO 1993/01161; Hudson et al. *Nat. Med.* 9:129-134 (2003). Triabodies and tetrabodies are also described in Hudson et al. *Nat. Med.* 9:129-134 (2003). Single-domain antibodies are antibody fragments comprising all or a portion of the heavy chain variable domain or all or a portion of the light chain variable domain of an antibody. In certain embodiments, a single-domain antibody is a human single-domain antibody (*see, e.g.*, U.S. Patent No. 6248516).

[0331] Antibody fragments can be made by various techniques, including but not limited to proteolytic digestion of an intact antibody as well as production by recombinant host cells (*e.g.*, *E. coli* or phage), as described herein.

[0332] In some embodiments, the antibody fragment is used in combination with a second Sortilin antibody and/or with one or more antibodies that specifically bind a disease-causing protein selected from: amyloid beta or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, proline-arginine (PR) repeat peptides, and any combination thereof.

(3) Chimeric and Humanized antibodies

[0333] In some embodiments of any of the antibodies provided herein, the antibody is a chimeric antibody. Certain chimeric antibodies are described, *e.g.*, in U.S. Patent No. 4816567. In one example, a chimeric antibody comprises a non-human variable region (*e.g.*, a variable region derived from a mouse, rat, hamster, rabbit, or non-human primate, such as a monkey) and a human constant region. In a further example, a chimeric antibody is a "class switched" antibody in which the class or subclass has been changed from that of the parent antibody. Chimeric antibodies include antigen-binding fragments thereof.

[0334] In some embodiments of any of the antibodies provided herein, the antibody is a humanized antibody. Typically, a non-human antibody is humanized to reduce immunogenicity to humans, while retaining the specificity and affinity of the parental non-human antibody. In certain embodiments, a humanized antibody is substantially non-immunogenic in humans. In certain embodiments, a humanized antibody has substantially the same affinity for a target as an antibody from another species from which the humanized antibody is derived. *See, e.g.*, U.S. Pat. No. 5530101, 5693761; 5693762; and 5585089. In certain embodiments, amino acids of an antibody variable domain that can be modified without diminishing the native affinity of the antigen binding domain while reducing its immunogenicity are identified. *See, e.g.*, U.S. Pat. Nos. 5766886 and 5869619. Generally, a humanized antibody comprises one or more variable domains in which HVRs (or portions thereof) are derived from a non-human antibody, and FRs (or portions thereof) are derived from human antibody sequences. A humanized antibody optionally will also comprise at least a portion of a human constant region. In some embodiments, some FR residues in a humanized antibody are substituted with corresponding residues

from a non-human antibody (*e.g.*, the antibody from which the HVR residues are derived), for example, to restore or improve antibody specificity or affinity.

[0335] Humanized antibodies and methods of making them are reviewed, for example, in Almagro et al. *Front. Biosci.* 13:1619-1633 (2008), and are further described, *e.g.*, in US Patent Nos. 5821337, 7527791, 6982321, and 7087409. Human framework regions that may be used for humanization include but are not limited to: framework regions selected using the "best-fit" method (*see, e.g.*, Sims et al. *J. Immunol.* 151:2296 (1993)); framework regions derived from the consensus sequence of human antibodies of a particular subgroup of light or heavy chain variable regions (*see, e.g.*, Carter et al. *Proc. Natl. Acad. Sci. USA* 89:4285 (1992); and Presta et al., *J. Immunol.* 151:2623 (1993)); human mature (somatically mutated) framework regions or human germline framework regions (*see, e.g.*, Almagro and Fransson *Front. Biosci.* 13:1619-1633 (2008)); and framework regions derived from screening FR libraries (*see, e.g.*, Baca et al. *J. Biol. Chem.* 272:10678-10684 (1997) and Rosok et al. *J. Biol. Chem.* 271:22611-22618 (1996)).

(4) Human Antibodies

[0336] In some embodiments of any of the antibodies provided herein, the antibody is a human antibody. Human antibodies can be produced using various techniques known in the art. Human antibodies are described generally in van Dijk et al. *Curr. Opin. Pharmacol.* 5:368-74 (2001) and Lonberg *Curr. Opin. Immunol.* 20:450-459 (2008).

[0337] Human antibodies may be prepared by administering an immunogen to a transgenic animal that has been modified to produce intact human antibodies or intact antibodies with human variable regions in response to antigenic challenge. One can engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig loci in anticipation that such mice would produce human antibodies in the absence of mouse antibodies. Large human Ig fragments can preserve the large variable gene diversity as well as the proper regulation of antibody production and expression. By exploiting the mouse machinery for antibody diversification and selection and the lack of immunological tolerance to human proteins, the reproduced human antibody repertoire in these mouse strains can yield high affinity fully human antibodies against any antigen of interest, including human antigens. Using the hybridoma technology, antigen-specific human MAbs with the desired specificity can be produced and selected. Certain exemplary methods are described in U.S. Pat. No. 5545807, EP 546073, and EP 546073. *See also*, for example, U.S. Patent Nos. 6075181 and 6150584 describing XENOMOUSE™ technology; U.S. Patent No. 5770429 describing HUMAB® technology; U.S. Patent No. 7041870 describing K-MOUSE® technology, and U.S. Patent Application Publication No. US 2007/0061900, describing VELOCIMOUSE® technology. Human variable regions from intact antibodies generated by such animals may be further modified, *e.g.*, by combining with a different human constant region.

[0338] Human antibodies can also be made by hybridoma-based methods. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies have been described. (See, e.g., Kozbor *J. Immunol.* 133:3001 (1984) and Boerner et al. *J. Immunol.* 147:86 (1991)). Human antibodies generated via human B-cell hybridoma technology are also described in Li et al. *Proc. Natl. Acad. Sci. USA*, 103:3557-3562 (2006). Additional methods include those described, for example, in U.S. Patent No. 7189826 (describing production of monoclonal human IgM antibodies from hybridoma cell lines). Human hybridoma technology (Trioma technology) is also described in Vollmers et al. *Histology and Histopathology* 20(3):927-937 (2005) and Vollmers et al. *Methods and Findings in Experimental and Clinical Pharmacology* 27(3):185-91 (2005). Human antibodies may also be generated by isolating Fv clone variable domain sequences selected from human-derived phage display libraries. Such variable domain sequences may then be combined with a desired human constant domain. Techniques for selecting human antibodies from antibody libraries are described below.

[0339] In some embodiments of any of the antibodies provided herein, the antibody is a human antibody isolated by *in vitro* methods and/or screening combinatorial libraries for antibodies with the desired activity or activities. Suitable examples include but are not limited to phage display (CAT, Morphosys, Dyax, Biosite/Medarex, Xoma, Symphogen, Alexion (formerly Proliferon), Affimed) ribosome display (CAT), yeast-based platforms (Adimab), and the like. In certain phage display methods, repertoires of VH and VL genes are separately cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be screened for antigen-binding phage as described in Winter et al. *Ann. Rev. Immunol.* 12: 433-455 (1994). For example, a variety of methods are known in the art for generating phage display libraries and screening such libraries for antibodies possessing the desired binding characteristics. See also Sidhu et al. *J. Mol. Biol.* 338(2): 299-310, 2004; Lee et al. *J. Mol. Biol.* 340(5): 1073-1093, 2004; Fellouse *Proc. Natl. Acad. Sci. USA* 101(34):12467-12472 (2004); and Lee et al. *J. Immunol. Methods* 284(-2):1 19-132 (2004). Phage typically display antibody fragments, either as single-chain Fv (scFv) fragments or as Fab fragments. Libraries from immunized sources provide high-affinity antibodies to the immunogen without the requirement of constructing hybridomas. Alternatively, the naive repertoire can be cloned (e.g., from human) to provide a single source of antibodies to a wide range of non-self and also self-antigens without any immunization as described by Griffiths et al. *EMBO J.* 12: 725-734 (1993). Finally, naive libraries can also be made synthetically by cloning unrearranged V-gene segments from stem cells, and using PCR primers comprising random sequence to encode the highly variable HVR3 regions and to accomplish rearrangement *in vitro*, as described by Hoogenboom et al. *J. Mol. Biol.*, 227: 381-388, 1992. Patent publications describing human antibody phage libraries include, for example: US Patent No. 5750373,

and US Patent Publication Nos. 2007/0292936 and 2009/0002360. Antibodies isolated from human antibody libraries are considered human antibodies or human antibody fragments herein.

(5) Constant Regions including Fc regions

[0340] In some embodiments of any of the antibodies provided herein, the antibody comprises an Fc. In some embodiments, the Fc is a human IgG1, IgG2, IgG3, and/or IgG4 isotype. In some embodiments, the antibody is of the IgG class, the IgM class, or the IgA class.

[0341] In certain embodiments of any of the antibodies provided herein, the antibody has an IgG2 isotype. In some embodiments, the antibody contains a human IgG2 constant region. In some embodiments, the human IgG2 constant region includes an Fc region. In some embodiments, the antibody induces the one or more Sortilin activities or independently of binding to an Fc receptor. In some embodiments, the antibody binds an inhibitory Fc receptor. In certain embodiments, the inhibitory Fc receptor is inhibitory Fc-gamma receptor IIB (FcγIIB).

[0342] In certain embodiments of any of the antibodies provided herein, the antibody has an IgG1 isotype. In some embodiments, the antibody contains a mouse IgG1 constant region. In some embodiments, the antibody contains a human IgG1 constant region. In some embodiments, the human IgG1 constant region includes an Fc region. In some embodiments, the antibody binds an inhibitory Fc receptor. In certain embodiments, the inhibitory Fc receptor is inhibitory Fc-gamma receptor IIB (FcγIIB).

[0343] In certain embodiments of any of the antibodies provided herein, the antibody has an IgG4 isotype. In some embodiments, the antibody contains a human IgG4 constant region. In some embodiments, the human IgG4 constant region includes an Fc region. In some embodiments, the antibody binds an inhibitory Fc receptor. In certain embodiments, the inhibitory Fc receptor is inhibitory Fc-gamma receptor IIB (FcγIIB).

[0344] In certain embodiments of any of the antibodies provided herein, the antibody has a hybrid IgG2/4 isotype. In some embodiments, the antibody includes an amino acid sequence comprising amino acids 118 to 260 according to EU numbering of human IgG2 and amino acids 261-447 according to EU numbering of human IgG4 (WO 1997/11971; WO 2007/106585).

[0345] In some embodiments, the Fc region increases clustering without activating complement as compared to a corresponding antibody comprising an Fc region that does not comprise the amino acid substitutions. In some embodiments, the antibody induces one or more activities of a target specifically bound by the antibody. In some embodiments, the antibody binds to Sortilin.

[0346] It may also be desirable to modify an anti-Sortilin antibody of the present disclosure to modify effector function and/or to increase serum half-life of the antibody. For example, the Fc receptor binding site on the constant region may be modified or mutated to remove or reduce binding affinity to

certain Fc receptors, such as FcγRI, FcγRII, and/or FcγRIII to reduce Antibody-dependent cell-mediated cytotoxicity. In some embodiments, the effector function is impaired by removing N-glycosylation of the Fc region (*e.g.*, in the CH2 domain of IgG) of the antibody. In some embodiments, the effector function is impaired by modifying regions such as 233-236, 297, and/or 327-331 of human IgG as described in WO 99/58572 and Armour et al. *Molecular Immunology* 40: 585-593 (2003); Reddy et al. *J. Immunology* 164:1925-1933 (2000). In other embodiments, it may also be desirable to modify an anti-Sortilin antibody of the present disclosure to modify effector function to increase finding selectivity toward the ITIM-containing FcγRIIb (CD32b) to increase clustering of Sortilin antibodies on adjacent cells without activating humoral responses including Antibody-dependent cell-mediated cytotoxicity and antibody-dependent cellular phagocytosis.

[0347] To increase the serum half-life of the antibody, one may incorporate a salvage receptor binding epitope into the antibody (especially an antibody fragment) as described in U.S. Patent 5739277, for example. As used herein, the term “*salvage receptor binding epitope*” refers to an epitope of the Fc region of an IgG molecule (*e.g.*, IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule. Other amino acid sequence modifications.

(6) Multispecific Antibodies

[0348] Multispecific are antibodies that have binding specificities for at least two different epitopes, including those on the same or another polypeptide (*e.g.*, one or more Sortilin polypeptides of the present disclosure). In some embodiments, the multispecific antibody can be a bispecific antibody. In some embodiments, the multispecific antibody can be a trispecific antibody. In some embodiments, the multispecific antibody can be a tetraspecific antibody. Such antibodies can be derived from full-length antibodies or antibody fragments (*e.g.*, F(ab')₂ bispecific antibodies). In some embodiments, the multispecific antibody comprises a first antigen binding region which binds to first site on Sortilin and comprises a second antigen binding region which binds to a second site on Sortilin. In some embodiment, the multispecific antibodies comprises a first antigen binding region which binds to Sortilin and a second antigen binding region that binds to a second polypeptide.

[0349] Provided herein are multispecific antibodies comprises a first antigen binding region, wherein the first antigen binding region comprises the six HVRs of an antibody described herein, which binds to Sortilin and a second antigen binding region that binds to a second polypeptide. In some embodiments, the first antigen binding region comprises the V_H or V_L of an antibody described herein.

[0350] In some embodiments of any of the multispecific antibodies, the second polypeptide is a) an antigen facilitating transport across the blood-brain-barrier; (b) an antigen facilitating transport across the blood-brain-barrier selected from transferrin receptor (TR), insulin receptor (HIR), insulin-like growth factor receptor (IGFR), low-density lipoprotein receptor related proteins 1 and 2 (LPR-1 and 2),

diphtheria toxin receptor, CRM197, a llama single domain antibody, TMEM 30(A), a protein transduction domain, TAT, Syn-B, penetratin, a poly-arginine peptide, an angiopep peptide, and ANG1005; (c) a disease-causing protein selected from amyloid beta, oligomeric amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), c9RAN protein, prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides; (d) ligands and/or proteins expressed on immune cells, wherein the ligands and/or proteins selected from CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA-4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, BTLA, KIR, GAL9, TIM3, A2AR, LAG-3, and phosphatidylserine; and/or (e) a protein, lipid, polysaccharide, or glycolipid expressed on one or more tumor cells and any combination thereof.

[0351] Numerous antigens are known in the art that facilitate transport across the blood-brain barrier (*see, e.g.,* Gabathuler R. *Neurobiol. Dis.* 37:48-57 (2010)). Such second antigens include, without limitation, transferrin receptor (TR), insulin receptor (HIR), Insulin-like growth factor receptor (IGFR), low-density lipoprotein receptor related proteins 1 and 2 (LPR-1 and 2), diphtheria toxin receptor, including CRM197 (a non-toxic mutant of diphtheria toxin), llama single domain antibodies such as TMEM 30(A) (Flippase), protein transduction domains such as TAT, Syn-B, or penetratin, poly-arginine or generally positively charged peptides, Angiopep peptides such as ANG1005 (*see, e.g.,* Gabathuler, 2010), and other cell surface proteins that are enriched on blood-brain barrier endothelial cells (*see, e.g.,* Daneman et al. *PLoS One* 5(10):e13741 (2010)).

[0352] The multivalent antibodies may recognize the Sortilin antigen as well as without limitation additional antigens A β peptide, antigen or an α -synuclein protein antigen or, Tau protein antigen or, TDP-43 protein antigen or, prion protein antigen or, huntingtin protein antigen, or RAN, translation Products antigen, including the DiPeptide Repeats,(DPRs peptides) composed of glycine-alanine (GA), glycine-proline (GP), glycine-arginine (GR), proline-alanine (PA), or proline-arginine (PR), Insulin receptor, insulin like growth factor receptor. Transferrin receptor or any other antigen that facilitate antibody transfer across the blood brain barrier. In some embodiments, the second polypeptide is transferrin. In some embodiments, the second polypeptide is Tau. In some embodiments, the second

polypeptide is A β . In some embodiments, the second polypeptide is TREM2. In some embodiments, the second polypeptide is α -synuclein.

[0353] The multivalent antibody contains at least one polypeptide chain (and preferably two polypeptide chains), wherein the polypeptide chain or chains comprise two or more variable domains. For instance, the polypeptide chain or chains may comprise VD1-(X1)_n-VD2-(X2)_n-Fc, wherein VD1 is a first variable domain, VD2 is a second variable domain, Fc is one polypeptide chain of an Fc region, X1 and X2 represent an amino acid or polypeptide, and n is 0 or 1. Similarly, the polypeptide chain or chains may comprise V_H-C_{H1}-flexible linker-V_H-C_{H1}-Fc region chain; or V_H-C_{H1}-V_H-C_{H1}-Fc region chain. The multivalent antibody herein preferably further comprises at least two (and preferably four) light chain variable domain polypeptides. The multivalent antibody herein may, for instance, comprise from about two to about eight light chain variable domain polypeptides. The light chain variable domain polypeptides contemplated here comprise a light chain variable domain and, optionally, further comprise a CL domain.

[0354] Techniques for making multispecific antibodies include, but are not limited to, recombinant co-expression of two immunoglobulin heavy chain- light chain pairs having different specificities (*see* Milstein and Cuello *Nature* 305: 537 (1983), WO 93/08829, and Traunecker et al. *EMBO J.* 10:3655 (1991)), and "knob-in-hole" engineering (*see, e.g.*, U.S. Patent No. 5731168). *See also* WO 2013/026833 (CrossMab). Multi-specific antibodies may also be made by engineering electrostatic steering effects for making antibody Fc- heterodimeric molecules (WO 2009/089004A1); cross-linking two or more antibodies (*see, e.g.*, US Patent No. 4676980); using leucine; using "diabody" technology for making bispecific antibody fragments (*see, e.g.*, Hollinger et al. *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993)); and using single-chain Fv (scFv) dimers (*see, e.g.*, Gruber et al. *J. Immunol.* 152:5368 (1994)); and preparing trispecific antibodies as described, *e.g.*, in Tutt et al. *J. Immunol.* 147: 60 (1991).

[0355] Engineered antibodies with three or more functional antigen binding sites, including "Octopus antibodies," are also included herein (*see, e.g.*, US 2006/0025576). The antibody herein also includes a "Dual Acting FAb" or "DAF" comprising an antigen binding site that binds to multiple Sortilin (*see*, US 2008/0069820, for example).

(7) *Antibodies with improved stability*

[0356] Amino acid sequence modifications of anti-Sortilin antibodies of the present disclosure, or antibody fragments thereof to improve stability during manufacturing, storage, and *in vivo* administration, are also contemplated. For example, it may be desirable to reduce degradation of the antibodies or antibody fragments of the present disclosure through multiple pathways, including without limitation, oxidation and deamidation. Amino acid sequence variants of the antibodies or antibody fragments are prepared by introducing appropriate nucleotide changes into the nucleic acid encoding the antibodies or antibody fragments, or by peptide synthesis. Such modifications include, for example, deletions from,

and/or insertions into and/or substitutions of, residues within the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution can be made to arrive at the final construct, provided that the final construct possesses the desired characteristics (*i.e.*, reduced susceptibility to degradation).

[0357] In some embodiments, the asparagine (N33) site in the HVR-L1 region of an anti-Sortilin antibody of the present disclosure may be susceptible to degradation by means of deamidation. In certain embodiments, the asparagine (N33) site in the HVR-L1 region of S-60-15 (SEQ ID NO:8) may be susceptible to deamidation. Upon deamidation, the asparagine (N33) site in the HVR-L1 region of S-60-15 results in an Asn to Asp/IsoAsp change. In certain embodiments, the asparagine (N33) site in the HVR-L1 region of S-60-15 may be substituted to prevent or reduce deamidation. Non-limiting exemplary amino acid sequence variants of S-60-15 having amino acid substitutions in the asparagine (N33) site of the HVR-L1 region include S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], or S-60-15.17 [N33L].

(8) *Antibody Variants*

[0358] In some embodiments of any of the antibodies provided herein, amino acid sequence variants of the antibodies are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody.

(i) *Substitution, Insertion, and Deletion Variants*

[0359] In some embodiments of any of the antibodies provided herein, antibody variants having one or more amino acid substitutions are provided. Amino acid sequence variants of an antibody may be prepared by introducing appropriate modifications into the nucleotide sequence encoding the antibody, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of residues within the amino acid sequences of the antibody. **TABLE 37: Amino Acid Substitutions**

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp, Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn

Original Residue	Exemplary Substitutions	Preferred Substitutions
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala
His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu
Phe (F)	Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Ser	Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

[0360] Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;
- (2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;
- (3) acidic: Asp, Glu;
- (4) basic: His, Lys, Arg;
- (5) residues that influence chain orientation: Gly, Pro; and
- (6) aromatic: Trp, Tyr, Phe.

[0361] For example, non-conservative substitutions can involve the exchange of a member of one of these classes for a member from another class. Such substituted residues can be introduced, for example, into regions of a human antibody that are homologous with non-human antibodies, or into the non-homologous regions of the molecule.

[0362] In making changes to the polypeptide or antibody described herein, according to certain embodiments, the hydropathic index of amino acids can be considered. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics. They are:

isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (−0.4); threonine (−0.7); serine (−0.8); tryptophan (−0.9); tyrosine (−1.3); proline (−1.6); histidine (−3.2); glutamate (−3.5); glutamine (−3.5); aspartate (−3.5); asparagine (−3.5); lysine (−3.9); and arginine (−4.5).

[0363] The importance of the hydropathic amino acid index in conferring interactive biological function on a protein is understood in the art. Kyte et al. *J. Mol. Biol.*, 157:105-131 (1982). It is known that certain amino acids can be substituted for other amino acids having a similar hydropathic index or score and still retain a similar biological activity. In making changes based upon the hydropathic index, in certain embodiments, the substitution of amino acids whose hydropathic indices are within ± 2 is included. In certain embodiments, those which are within ± 1 are included, and in certain embodiments, those within ± 0.5 are included.

[0364] It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity, particularly where the biologically functional protein or peptide thereby created is intended for use in immunological embodiments, as in the present case. In certain embodiments, the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigenicity, i.e., with a biological property of the protein.

[0365] The following hydrophilicity values have been assigned to these amino acid residues: arginine (+3.0); lysine (+3.0 $\pm$ 1); aspartate (+3.0 $\pm$ 1); glutamate (+3.0 $\pm$ 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (−0.4); proline (−0.5 $\pm$ 1); alanine (−0.5); histidine (−0.5); cysteine (−1.0); methionine (−1.3); valine (−1.5); leucine (−1.8); isoleucine (−1.8); tyrosine (−2.3); phenylalanine (−2.5) and tryptophan (−3.4). In making changes based upon similar hydrophilicity values, in certain embodiments, the substitution of amino acids whose hydrophilicity values are within ± 2 is included, in certain embodiments, those which are within ± 1 are included, and in certain embodiments, those within ± 0.5 are included. One can also identify epitopes from primary amino acid sequences on the basis of hydrophilicity. These regions are also referred to as “epitopic core regions”.

[0366] In certain embodiments, substitutions, insertions, or deletions may occur within one or more HVRs so long as such alterations do not substantially reduce the ability of the antibody to bind antigen. For example, conservative alterations (e.g., conservative substitutions as provided herein) that do not substantially reduce binding affinity may be made in HVRs. Such alterations may, for example, be outside of antigen contacting residues in the HVRs. In certain embodiments of the variant VH and VL sequences provided above, each HVR either is unaltered, or contains no more than one, two or three amino acid substitutions.

[0367] Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides comprising a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody to an enzyme (*e.g.*, for ADEPT) or a polypeptide which increases the serum half-life of the antibody.

[0368] Any cysteine residue not involved in maintaining the proper conformation of the antibody also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment, such as an Fv fragment).

(ii) Glycosylation variants

[0369] In some embodiments of any of the antibodies provided herein, the antibody is altered to increase or decrease the extent to which the antibody is glycosylated. Addition or deletion of glycosylation sites to an antibody may be conveniently accomplished by altering the amino acid sequence such that one or more glycosylation sites is created or removed.

[0370] Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

[0371] Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

[0372] Where the antibody comprises an Fc region, the carbohydrate attached thereto may be altered. Native antibodies produced by mammalian cells typically comprise a branched, biantennary oligosaccharide that is generally attached by an N-linkage to Asn297 according to Kabat numbering of the CH2 domain of the Fc region. The oligosaccharide may include various carbohydrates, for example, mannose, N-acetyl glucosamine (GlcNAc), galactose, and sialic acid, as well as a fucose attached to a GlcNAc in the "stem" of the biantennary oligosaccharide structure. In some embodiments, modifications

of the oligosaccharide in an antibody of the invention may be made in order to create antibody variants with certain improved properties.

[0373] In one embodiment, antibody variants are provided having a carbohydrate structure that lacks fucose attached (directly or indirectly) to an Fc region. *See, e.g.*, US Patent Publication Nos. 2003/0157108 and 2004/0093621. Examples of publications related to "defucosylated" or "fucose-deficient" antibody variants include: US 2003/0157108; US 2003/0115614; US 2002/0164328; US 2004/0093621; US 2004/0132140; US 2004/0110704; US 2004/0110282; US 2004/0109865; Okazaki et al. *J. Mol. Biol.* 336:1239-1249 (2004); Yamane-Ohnuki et al. *Biotech. Bioeng.* 87:614 (2004). Examples of cell lines capable of producing defucosylated antibodies include Led 3 CHO cells deficient in protein fucosylation (Ripka et al. *Arch. Biochem. Biophys.* 249:533-545 (1986); US 2003/0157108), and knockout cell lines, such as alpha-1,6-fucosyltransferase gene, FUT8, knockout CHO cells (*see, e.g.*, Yamane-Ohnuki et al. *Biotech. Bioeng.* 87: 614 (2004) and Kanda et al. *Biotechnol. Bioeng.* 94(4):680-688 (2006)).

(iii) Modified Constant regions

[0374] In some embodiments of any of the antibodies provided herein, the antibody Fc is an antibody Fc isotypes and/or modifications. In some embodiments, the antibody Fc isotype and/or modification is capable of binding to Fc gamma receptor.

[0375] In some embodiments of any of the antibodies provided herein, the modified antibody Fc is an IgG1 modified Fc. In some embodiments, the IgG1 modified Fc comprises one or more modifications. For example, in some embodiments, the IgG1 modified Fc comprises one or more amino acid substitutions (*e.g.*, relative to a wild-type Fc region of the same isotype). In some embodiments, the one or more amino acid substitutions are selected from N297A (Bolt S et al. (1993) *Eur J Immunol* 23:403-411), D265A (Shields et al. (2001) *R. J. Biol. Chem.* 276, 6591-6604), L234A, L235A (Hutchins et al. (1995) *Proc Natl Acad Sci USA*, 92:11980-11984; Alegre et al., (1994) *Transplantation* 57:1537-1543. 31; Xu et al., (2000) *Cell Immunol*, 200:16-26), G237A (Alegre et al. (1994) *Transplantation* 57:1537-1543. 31; Xu et al. (2000) *Cell Immunol*, 200:16-26), C226S, C229S, E233P, L234V, L234F, L235E (McEarchern et al., (2007) *Blood*, 109:1185-1192), P331S (Sazinsky et al., (2008) *Proc Natl Acad Sci USA* 2008, 105:20167-20172), S267E, L328F, A330L, M252Y, S254T, and/or T256E, where the amino acid position is according to the EU numbering convention. In some embodiments of any of the antibodies provided herein, the antibody is an IgG1 isotype and the Fc region comprises amino acid substitutions at positions L234A, L235A, and P331S, wherein the numbering of the residue position is according to EU numbering.

[0376] In some embodiments of any of the IgG1 modified Fc, the Fc comprises N297A mutation according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises

D265A and N297A mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises D270A mutations according to EU numbering. In some embodiments, the IgG1 modified Fc comprises L234A and L235A mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises L234A and G237A mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises L234A, L235A and G237A mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises one or more (including all) of P238D, L328E, E233, G237D, H268D, P271G and A330R mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises one or more of S267E/L328F mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises P238D, L328E, E233D, G237D, H268D, P271G and A330R mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises P238D, L328E, G237D, H268D, P271G and A330R mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises P238D, S267E, L328E, E233D, G237D, H268D, P271G and A330R mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises P238D, S267E, L328E, G237D, H268D, P271G and A330R mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises C226S, C229S, E233P, L234V, and L235A mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises L234F, L235E, and P331S mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises S267E and L328F mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises S267E mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises a substitute of the constant heavy 1 (CH1) and hinge region of IgG1 with CH1 and hinge region of IgG2 (amino acids 118-230 of IgG2 according to EU numbering) with a Kappa light chain.

[0377] In some embodiments of any of the IgG1 modified Fc, the Fc includes two or more amino acid substitutions that increase antibody clustering without activating complement as compared to a corresponding antibody having an Fc region that does not include the two or more amino acid substitutions. Accordingly, in some embodiments of any of the IgG1 modified Fc, the IgG1 modified Fc is an antibody comprising an Fc region, where the antibody comprises an amino acid substitution at position E430G and one or more amino acid substitutions in the Fc region at a residue position selected from: L234F, L235A, L235E, S267E, K322A, L328F, A330S, P331S, and any combination thereof according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G, L243A, L235A, and P331S according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G and P331S

according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G and K322A according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G, A330S, and P331S according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G, K322A, A330S, and P331S according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G, K322A, and A330S according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G, K322A, and P331S according to EU numbering.

[0378] In some embodiments of any of the IgG1 modified Fc, the IgG1 modified Fc may further comprise herein may be combined with an A330L mutation (Lazar et al. *Proc Natl Acad Sci USA*, 103:4005-4010 (2006)), or one or more of L234F, L235E, and/or P331S mutations (Sazinsky et al. *Proc Natl Acad Sci USA*, 105:20167-20172 (2008)), according to the EU numbering convention, to eliminate complement activation. In some embodiments of any of the IgG1 modified Fc, the IgG1 modified Fc may further comprise one or more of A330L, A330S, L234F, L235E, and/or P331S according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the IgG1 modified Fc may further comprise one or more mutations to enhance the antibody half-life in human serum (*e.g.*, one or more (including all) of M252Y, S254T, and T256E mutations according to the EU numbering convention). In some embodiments of any of the IgG1 modified Fc, the IgG1 modified Fc may further comprise one or more of E430G, E430S, E430F, E430T, E345K, E345Q, E345R, E345Y, S440Y, and/or S440W according to EU numbering.

[0379] Other aspects of the present disclosure relate to antibodies having modified constant regions (*i.e.*, Fc regions). An antibody dependent on binding to FcγR receptor to activate targeted receptors may lose its agonist activity if engineered to eliminate FcγR binding (*see, e.g.*, Wilson et al. *Cancer Cell* 19:101-113 (2011); Armour et al. *Immunology* 40:585-593 (2003); and White et al. *Cancer Cell* 27:138-148 (2015)). As such, it is thought that an anti-Sortlin antibody of the present disclosure with the correct epitope specificity can activate the target antigen, with minimal adverse effects, when the antibody has an Fc domain from a human IgG2 isotype (CH1 and hinge region) or another type of Fc domain that is capable of preferentially binding the inhibitory FcγRIIB r receptors, or a variation thereof.

[0380] In some embodiments of any of the antibodies provided herein, the modified antibody Fc is an IgG2 modified Fc. In some embodiments, the IgG2 modified Fc comprises one or more modifications. For example, in some embodiments, the IgG2 modified Fc comprises one or more amino acid substitutions (*e.g.*, relative to a wild-type Fc region of the same isotype). In some embodiments of any of the IgG2 modified Fc, the one or more amino acid substitutions are selected from V234A (Alegre et al. *Transplantation* 57:1537-1543 (1994); Xu et al. *Cell Immunol*, 200:16-26 (2000)); G237A (Cole et al.

Transplantation, 68:563-571 (1999)); H268Q, V309L, A330S, P331S (US 2007/0148167; Armour et al. *Eur J Immunol* 29: 2613-2624 (1999); Armour et al. *The Haematology Journal* 1(Suppl.1):27 (2000); Armour et al. *The Haematology Journal* 1(Suppl.1):27 (2000)), C219S, and/or C220S (White et al. *Cancer Cell* 27, 138-148 (2015)); S267E, L328F (Chu et al. *Mol Immunol*, 45:3926-3933 (2008)); and M252Y, S254T, and/or T256E according to the EU numbering convention. In some embodiments of any of the IgG2 modified Fc, the Fc comprises an amino acid substitution at positions V234A and G237A according to EU numbering. In some embodiments of any of the IgG2 modified Fc, the Fc comprises an amino acid substitution at positions C219S or C220S according to EU numbering. In some embodiments of any of the IgG2 modified Fc, the Fc comprises an amino acid substitution at positions A330S and P331S according to EU numbering. In some embodiments of any of the IgG2 modified Fc, the Fc comprises an amino acid substitution at positions S267E and L328F according to EU numbering.

[0381] In some embodiments of any of the IgG2 modified Fc, the Fc comprises a C127S amino acid substitution according to the EU numbering convention (White et al., (2015) *Cancer Cell* 27, 138-148; Lightle et al. *Protein Sci.* 19:753-762 (2010); and WO 2008/079246). In some embodiments of any of the IgG2 modified Fc, the antibody has an IgG2 isotype with a Kappa light chain constant domain that comprises a C214S amino acid substitution according to the EU numbering convention (White et al. *Cancer Cell* 27:138-148 (2015); Lightle et al. *Protein Sci.* 19:753-762 (2010); and WO 2008/079246).

[0382] In some embodiments of any of the IgG2 modified Fc, the Fc comprises a C220S amino acid substitution according to the EU numbering convention. In some embodiments of any of the IgG2 modified Fc, the antibody has an IgG2 isotype with a Kappa light chain constant domain that comprises a C214S amino acid substitution according to the EU numbering convention.

[0383] In some embodiments of any of the IgG2 modified Fc, the Fc comprises a C219S amino acid substitution according to the EU numbering convention. In some embodiments of any of the IgG2 modified Fc, the antibody has an IgG2 isotype with a Kappa light chain constant domain that comprises a C214S amino acid substitution according to the EU numbering convention.

[0384] In some embodiments of any of the IgG2 modified Fc, the Fc includes an IgG2 isotype heavy chain constant domain 1(CH1) and hinge region (White et al. *Cancer Cell* 27:138-148 (2015)). In certain embodiments of any of the IgG2 modified Fc, the IgG2 isotype CH1 and hinge region comprise the amino acid sequence of 118-230 according to EU numbering. In some embodiments of any of the IgG2 modified Fc, the antibody Fc region comprises a S267E amino acid substitution, a L328F amino acid substitution, or both, and/or a N297A or N297Q amino acid substitution according to the EU numbering convention.

[0385] In some embodiments of any of the IgG2 modified Fc, the Fc further comprises one or more amino acid substitution at positions E430G, E430S, E430F, E430T, E345K, E345Q, E345R, E345Y,

S440Y, and S440W according to EU numbering. In some embodiments of any of the IgG2 modified Fc, the Fc may further comprise one or more mutations to enhance the antibody half-life in human serum (e.g., one or more (including all) of M252Y, S254T, and T256E mutations according to the EU numbering convention). In some embodiments of any of the IgG2 modified Fc, the Fc may further comprise A330S and P331S.

[0386] In some embodiments of any of the IgG2 modified Fc, the Fc is an IgG2/4 hybrid Fc. In some embodiments, the IgG2/4 hybrid Fc comprises IgG2 aa 118 to 260 and IgG4 aa 261 to 447. In some embodiments of any IgG2 modified Fc, the Fc comprises one or more amino acid substitutions at positions H268Q, V309L, A330S, and P331S according to EU numbering.

[0387] In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises one or more additional amino acid substitutions selected from A330L, L234F; L235E, or P331S according to EU numbering; and any combination thereof.

[0388] In certain embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises one or more amino acid substitutions at a residue position selected from C127S, L234A, L234F, L235A, L235E, S267E, K322A, L328F, A330S, P331S, E345R, E430G, S440Y, and any combination thereof according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G, L243A, L235A, and P331S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G and P331S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G and K322A according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G, A330S, and P331S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G, K322A, A330S, and P331S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G, K322A, and A330S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G, K322A, and P331S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions S267E and L328F according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at position C127S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E345R, E430G and S440Y according to EU numbering.

[0389] In some embodiments of any of the antibodies provided herein, the modified antibody Fc is an IgG4 modified Fc. In some embodiments, the IgG4 modified Fc comprises one or more modifications. For example, in some embodiments, the IgG4 modified Fc comprises one or more amino acid substitutions (*e.g.*, relative to a wild-type Fc region of the same isotype). In some embodiments of any of the IgG4 modified Fc, the one or more amino acid substitutions are selected from L235A, G237A, S229P, L236E (Reddy et al. *J Immunol* 164:1925-1933(2000)), S267E, E318A, L328F, M252Y, S254T, and/or T256E according to the EU numbering convention. In some embodiments of any of the IgG4 modified Fc, the Fc may further comprise L235A, G237A, and E318A according to the EU numbering convention. In some embodiments of any of the IgG4 modified Fc, the Fc may further comprise S228P and L235E according to the EU numbering convention. In some embodiments of any of the IgG4 modified Fc, the IgG4 modified Fc may further comprise S267E and L328F according to the EU numbering convention.

[0390] In some embodiments of any of the IgG4 modified Fc, the IgG4 modified Fc comprises may be combined with an S228P mutation according to the EU numbering convention (Angal et al. *Mol Immunol.* 30:105-108 (1993)) and/or with one or more mutations described in (Peters et al. *J Biol Chem.* 287(29):24525-33 (2012)) to enhance antibody stabilization.

[0391] In some embodiments of any of the IgG4 modified Fc, the IgG4 modified Fc may further comprise one or more mutations to enhance the antibody half-life in human serum (*e.g.*, one or more (including all) of M252Y, S254T, and T256E mutations according to the EU numbering convention).

[0392] In some embodiments of any of the IgG4 modified Fc, the Fc comprises L235E according to EU numbering. In certain embodiments of any of the IgG4 modified Fc, the Fc comprises one or more amino acid substitutions at a residue position selected from C127S, F234A, L235A, L235E, S267E, K322A, L328F, E345R, E430G, S440Y, and any combination thereof, according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at positions E430G, L243A, L235A, and P331S according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at positions E430G and P331S according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at positions E430G and K322A according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at position E430 according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc region comprises an amino acid substitution at positions E430G and K322A according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at positions S267E and L328F according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at position C127S according to EU numbering.

In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at positions E345R, E430G and S440Y according to EU numbering.

(9) *Other antibody modifications*

[0393] In some embodiments of any of the antibodies, the antibody is a derivative. The term “derivative” refers to a molecule that includes a chemical modification other than an insertion, deletion, or substitution of amino acids (or nucleic acids). In certain embodiments, derivatives comprise covalent modifications, including, but not limited to, chemical bonding with polymers, lipids, or other organic or inorganic moieties. In certain embodiments, a chemically modified antigen binding protein can have a greater circulating half-life than an antigen binding protein that is not chemically modified. In certain embodiments, a chemically modified antigen binding protein can have improved targeting capacity for desired cells, tissues, and/or organs. In some embodiments, a derivative antigen binding protein is covalently modified to include one or more water soluble polymer attachments, including, but not limited to, polyethylene glycol, polyoxyethylene glycol, or polypropylene glycol. *See, e.g.*, U.S. Pat. Nos. 4640835, 4496689, 4301144, 4670417, 4791192 and 4179337. In certain embodiments, a derivative antigen binding protein comprises one or more polymer, including, but not limited to, monomethoxy-polyethylene glycol, dextran, cellulose, , copolymers of ethylene glycol/propylene glycol, carboxymethylcellulose, polyvinyl pyrrolidone, poly-1, 3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), poly-(N-vinyl pyrrolidone)-polyethylene glycol, propylene glycol homopolymers, a polypropylene oxide/ethylene oxide co-polymer, polyoxyethylated polyols (*e.g.*, glycerol) and polyvinyl alcohol, as well as mixtures of such polymers.

[0394] In certain embodiments, a derivative is covalently modified with polyethylene glycol (PEG) subunits. In certain embodiments, one or more water-soluble polymer is bonded at one or more specific position, for example at the amino terminus, of a derivative. In certain embodiments, one or more water-soluble polymer is randomly attached to one or more side chains of a derivative. In certain embodiments, PEG is used to improve the therapeutic capacity for an antigen binding protein. In certain embodiments, PEG is used to improve the therapeutic capacity for a humanized antibody. Certain such methods are discussed, for example, in U.S. Pat. No. 6133426, which is hereby incorporated by reference for any purpose.

[0395] Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed “peptide mimetics” or “peptidomimetics.” Fauchere, *J. Adv. Drug Res.*, 15:29 (1986); and Evans et al. *J. Med. Chem.*, 30:1229 (1987), which are incorporated herein by reference for any purpose. Such compounds are often developed with the aid of computerized molecular modeling. Peptide mimetics that

are structurally similar to therapeutically useful peptides can be used to produce a similar therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (*i.e.*, a polypeptide that has a biochemical property or pharmacological activity), such as human antibody, but have one or more peptide linkages optionally replaced by a linkage selected from: -CH₂NH-, -CH₂S-, -CH₂-CH₂-, -CH=CH-(*cis* and *trans*), -COCH₂-, -CH(OH)CH₂-, and -CH₂SO-, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (*e.g.*, D-lysine in place of L-lysine) can be used in certain embodiments to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation can be generated by methods known in the art (Rizo and Gierasch *Ann. Rev. Biochem.*, 61:387 (1992), incorporated herein by reference for any purpose); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

[0396] Drug conjugation involves coupling of a biological active cytotoxic (anticancer) payload or drug to an antibody that specifically targets a certain tumor marker (*e.g.* a polypeptide that, ideally, is only to be found in or on tumor cells). Antibodies track these proteins down in the body and attach themselves to the surface of cancer cells. The biochemical reaction between the antibody and the target protein (antigen) triggers a signal in the tumor cell, which then absorbs or internalizes the antibody together with the cytotoxin. After the ADC is internalized, the cytotoxic drug is released and kills the cancer. Due to this targeting, ideally the drug has lower side effects and gives a wider therapeutic window than other chemotherapeutic agents. Techniques to conjugate antibodies are disclosed are known in the art (*see, e.g.*, Jane de Lartigue OncLive July 5, 2012; ADC Review on antibody-drug conjugates; and Ducry et al. *Bioconjugate Chemistry* 21 (1):5-13 (2010).

Binding assays and other assays

[0397] Anti-Sortilin antibodies of the present disclosure may be tested for antigen binding activity, *e.g.*, by known methods such as ELISA, surface plasmon resonance (SPR), Western blot, flow cytometry, FACS, Bio-layer interferometry *etc.*

[0398] In some embodiments, competition assays may be used to identify an antibody that competes with any of the antibodies described herein. In some embodiments, competition assays may be used to identify an antibody that competes with any of the antibodies listed in **Tables 1-30**, or comprising the heavy chain variable region and the light chain variable region of an antibody selected from S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-7, S-60-8, S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-

15.16 [N33M], S-60-15.17 [N33L], S-60-16, S-60-18, S-60-19, and S-60-24 for binding to Sortilin. In certain embodiments, such a competing antibody binds to the same epitope (e.g., a linear or a conformational epitope) that is bound by any of the antibodies listed in **Tables 1-30**, or comprising the heavy chain variable region and the light chain variable region of an antibody selected from S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-7, S-60-8, S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16, S-60-18, S-60-19, and S-60-24. Detailed exemplary methods for mapping an epitope to which an antibody binds are provided in Morris (1996) "Epitope Mapping Protocols," in *Methods in Molecular Biology* vol. 66 (Humana Press, Totowa, NJ).

[0399] In an exemplary competition assay, immobilized Sortilin or cells expressing Sortilin on a cell surface are incubated in a solution comprising a first labeled antibody that binds to Sortilin (e.g., human or non-human primate) and a second unlabeled antibody that is being tested for its ability to compete with the first antibody for binding to Sortilin. The second antibody may be present in a hybridoma supernatant. As a control, immobilized Sortilin or cells expressing Sortilin is incubated in a solution comprising the first labeled antibody but not the second unlabeled antibody. After incubation under conditions permissive for binding of the first antibody to Sortilin, excess unbound antibody is removed, and the amount of label associated with immobilized Sortilin or cells expressing Sortilin is measured. If the amount of label associated with immobilized Sortilin or cells expressing Sortilin is substantially reduced in the test sample relative to the control sample, then that indicates that the second antibody is competing with the first antibody for binding to Sortilin. See, Harlow and Lane (1988) *Antibodies: A Laboratory Manual* ch.14 (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY).

Ligand binding assays

[0400] Further provided herein are methods of screening for anti-Sortilin antibodies that bind His131, Val132, Pro133, Leu134, Val135, Ile136, Met137, Thr138, Arg196, Phe198, Arg199, Phe203, Lys205, Phe207, Thr210, Thr218, Tyr222, Ser223, Ser227, Ser242, Lys243, Lys248, Lys254, Lys260, Ser305, Phe306, Gly307, Arg311, Phe314, Ser316, Arg325, Arg326, Ile327, Phe350, Tyr351, Ser352, Ile353, Asn373, Ser379, Arg382, Tyr386, Ser595, and/or Glu700 of human Sortilin (SEQ ID NO: 81); or to amino acid residues of a mammalian Sortilin that corresponds to one or more amino acid residues His131, Val132, Pro133, Leu134, Val135, Ile136, Met137, Thr138, Arg196, Phe198, Arg199, Phe203, Lys205, Phe207, Thr210, Thr218, Tyr222, Ser223, Ser227, Ser242, Lys243, Lys248, Lys254, Lys260, Ser305, Phe306, Gly307, Arg311, Phe314, Ser316, Arg325, Arg326, Ile327, Phe350, Tyr351, Ser352, Ile353, Asn373, Ser379, Arg382, Tyr386, Ser595, and/or Glu700 of SEQ ID NO: 81, thereby blocking

the interactions between Sortilin and a Sortilin ligand (*e.g.*, Progranulin, pro-neurotrophin, pro-NGF, pro-BDNF, pro-NT3, p75, APP, LpL, APOA5, APOE). In some embodiments, a peptide library can be synthesized in which a Sortilin protein is dissected into consecutive 15-mer and 25-mer peptides separated by one amino acid residue and subsequently spotted onto filters. Binding of a Sortilin ligand can then be tested for its ability to interact with the receptor peptide or with peptides that are, for example, mutated at His131, Val132, Pro133, Leu134, Val135, Ile136, Met137, Thr138, Arg196, Phe198, Arg199, Phe203, Lys205, Phe207, Thr210, Thr218, Tyr222, Ser223, Ser227, Ser242, Lys243, Lys248, Lys254, Lys260, Ser305, Phe306, Gly307, Arg311, Phe314, Ser316, Arg325, Arg326, Ile327, Phe350, Tyr351, Ser352, Ile353, Asn373, Ser379, Arg382, Tyr386, Ser595, and/or Glu700 of human Sortilin libraries in the presence or absence of the anti-Sortilin antibodies by SPOT binding analysis (*e.g.*, Frank, R and Overwin, H (1996) *Methods. Mol. Biol.* 66, 149-169; Reineke, U *et al.*, (2002) *J. Immunol. Methods* 267, 13-26; and Andersen, OS *et al.*, (2010) *J. Biological Chemistry* 285, 12210-12222).

[0401] Further provided herein are methods of screening for anti-Sortilin antibodies that block interactions (*e.g.*, binding) Sortilin and a Sortilin ligand (*e.g.*, Progranulin, pro-neurotrophin, pro-NGF, pro-BDNF, pro-NT3, p75, APP, LpL, APOA5, APOE). In some embodiments, the interaction between Sortilin and Sortilin ligands (*e.g.*, Progranulin, pro-neurotrophin, pro-NGF, pro-BDNF, pro-NT3, p75, APP, LpL, APOA5, APOE) may be characterized using surface plasmon resonance analysis (*e.g.*, Skeldal, S *et al.*, (2012) *J Biol Chem.*, 287:43798; and Andersen, OS *et al.*, (2010) *The Journal Of Biological Chemistry*, 285,12210-12222, a pulldown assay (*e.g.*, Andersen, OS *et al.*, (2010) *The Journal Of Biological Chemistry*, 285,12210-12222, cellulose-bound proteins (*e.g.*, Andersen, OS *et al.*, (2010) *The Journal Of Biological Chemistry*, 285,12210-12222), a proximity ligation assay (*e.g.*, Gustafsen, C *et al.*, (2013) *The Journal of Neuroscience*, 33:64-71), and/or alkaline phosphatase-tagged ligands in cell binding assays (*e.g.*, Hu, F *et al.*, (2010) *Neuron* 68, 654-667).

Cell-based assays

[0402] Further provided herein are methods of screening for a Sortilin binding antagonist, such as an anti-Sortilin antibody, that include contacting an agent (*e.g.*, an anti-Sortilin antibody) with a cell expressing a Sortilin protein on its cell surface. In some embodiments, the agent and cell are further contacted with a Sortilin ligand of the present disclosure. In some embodiments, the cell itself expresses a Sortilin ligand of the present disclosure. The cell-based methods are particularly suited for screening and validating Sortilin binding antagonists (*e.g.*, anti-Sortilin antibodies) by assessing the effect on the interaction between Sortilin and a Sortilin ligand in the context of a cell.

[0403] Accordingly, certain aspects of the present disclosure relate to a cell expressing a Sortilin protein of the present disclosure on its cell surface. In some embodiments, the cell endogenously expresses a Sortilin protein of the present disclosure. In some embodiments, the cell is recombinantly

engineered to express a Sortilin protein of the present disclosure. In any of these embodiments, the Sortilin protein of the present disclosure (whether endogenous or recombinant) encoded by the polynucleotide will preferably include at least protein domains required for post-translational processing, membrane translocation, and targeting to the cell surface, including without limitation a signal peptide and a transmembrane domain. In some embodiments, the signal peptide and/or transmembrane domain may refer to the endogenous Sortilin signal peptide and/or transmembrane domain. In other embodiments, the signal peptide and/or transmembrane domain may refer to an exogenous signal peptide and/or transmembrane domain known to promote cell surface expression in the desired host cell. In preferred embodiments, the Sortilin protein will also contain a domain sufficient for binding a Sortilin ligand of the present disclosure.

[0404] In these embodiments, any cell that expresses a Sortilin protein of the present disclosure on its cell surface may be used. In some embodiments, the cell endogenously expresses a Sortilin protein of the present disclosure on its cell surface. In some embodiments, the cell is recombinantly engineered to express a Sortilin protein of the present disclosure on its cell surface. Any suitable Sortilin ligand of the present disclosure may be used, such that it retains the ability to bind to the Sortilin protein expressed on the cell surface. The Sortilin ligand need not be fluorescently labeled. Levels of Sortilin ligand may be detected by any assay known in the art, including without limitation ELISA, SPR, Western blotting, mass spectrometry, immunoprecipitation, peptide microarray, and so forth.

[0405] In some embodiments, the methods disclosed herein involve culturing a cell that expresses both a Sortilin protein on its cell surface and a Sortilin ligand in a media under conditions in which the Sortilin protein and the Sortilin ligand are expressed and the Sortilin ligand is released into the media; contacting the cell with an agent (*e.g.*, an anti-Sortilin antibody) under conditions in which the Sortilin protein is capable of binding to the Sortilin ligand; and detecting an increase in the level of the Sortilin ligand in the media, as compared to the level of the Sortilin ligand in the media in the absence of the agent. An increase in the level of the Sortilin ligand indicates that the agent is a Sortilin binding antagonist. Without wishing to be bound to theory, it is thought that the interaction between the Sortilin protein expressed on the cell surface and the secreted Sortilin ligand will result in endocytosis and lysosomal degradation of the Sortilin ligand. Therefore, it is thought that decreasing this interaction (*e.g.*, by addition of a Sortilin binding antagonist of the present disclosure) leads to an increase in the level of the Sortilin ligand in the media over time.

[0406] In these embodiments, any cell that expresses a Sortilin protein of the present disclosure on its cell surface and expresses and secretes a Sortilin ligand of the present disclosure may be used. In some embodiments, the cell may endogenously express a Sortilin protein of the present disclosure on its cell surface. In some embodiments, the cell may endogenously express and secrete a Sortilin ligand of the

present disclosure. In some embodiments, the cell is a U-251 cell, and the Sortilin ligand is a Progranulin protein. In some embodiments, the cell may be recombinantly engineered to express a Sortilin protein of the present disclosure on its cell surface. In some embodiments, the cell may be recombinantly engineered to express and secrete a Sortilin ligand of the present disclosure.

[0407] In any of the cell-based assays described herein, a Sortilin ligand of the present disclosure may be used. In some embodiments, the Sortilin ligand is a Progranulin protein. The Sortilin ligand may be a full-length protein, or it may be a Sortilin-binding peptide fragment thereof.

Nucleic acids, vectors, and host cells

[0408] Anti-Sortilin antibodies of the present disclosure may be produced using recombinant methods and compositions, *e.g.*, as described in U.S. Patent No. 4816567. In some embodiments, isolated nucleic acids having a nucleotide sequence encoding any of the anti-Sortilin antibodies of the present disclosure are provided. Such nucleic acids may encode an amino acid sequence comprising the V_L and/or an amino acid sequence comprising the V_H of the anti-Sortilin antibody (*e.g.*, the light and/or heavy chains of the antibody). In some embodiments, one or more vectors (*e.g.*, expression vectors) comprising such nucleic acids are provided. In some embodiments, a host cell comprising such nucleic acid is also provided. In some embodiments, the host cell comprises (*e.g.*, has been transduced with): (1) a vector comprising a nucleic acid that encodes an amino acid sequence comprising the V_L of the antibody and an amino acid sequence comprising the V_H of the antibody, or (2) a first vector comprising a nucleic acid that encodes an amino acid sequence comprising the V_L of the antibody and a second vector comprising a nucleic acid that encodes an amino acid sequence comprising the V_H of the antibody. In some embodiments, the host cell is eukaryotic, *e.g.*, a Chinese Hamster Ovary (CHO) cell or lymphoid cell (*e.g.*, Y0, NS0, Sp20 cell). Host cells of the present disclosure also include, without limitation, isolated cells, *in vitro* cultured cells, and *ex vivo* cultured cells.

[0409] Methods of making an anti-Sortilin antibody of the present disclosure are provided. In some embodiments, the method includes culturing a host cell of the present disclosure comprising a nucleic acid encoding the anti-Sortilin antibody, under conditions suitable for expression of the antibody. In some embodiments, the antibody is subsequently recovered from the host cell (or host cell culture medium).

[0410] For recombinant production of an anti-Sortilin antibody of the present disclosure, a nucleic acid encoding the anti-Sortilin antibody is isolated and inserted into one or more vectors for further cloning and/or expression in a host cell. Such nucleic acid may be readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody).

[0411] Suitable vectors comprising a nucleic acid sequence encoding any of the anti-Sortilin antibodies of the present disclosure, or cell-surface expressed fragments or polypeptides thereof

polypeptides (including antibodies) described herein include, without limitation, cloning vectors and expression vectors. Suitable cloning vectors can be constructed according to standard techniques, or may be selected from a large number of cloning vectors available in the art. While the cloning vector selected may vary according to the host cell intended to be used, useful cloning vectors generally have the ability to self-replicate, may possess a single target for a particular restriction endonuclease, and/or may carry genes for a marker that can be used in selecting clones comprising the vector. Suitable examples include plasmids and bacterial viruses, *e.g.*, pUC18, pUC19, Bluescript (*e.g.*, pBS SK+) and its derivatives, mpl8, mpl9, pBR322, pMB9, ColE1, pCR1, RP4, phage DNAs, and shuttle vectors such as pSA3 and pAT28. These and many other cloning vectors are available from commercial vendors such as BioRad, Strategene, and Invitrogen.

[0412] Suitable host cells for cloning or expression of antibody-encoding vectors include prokaryotic or eukaryotic cells. For example, anti-Sortilin antibodies of the present disclosure may be produced in bacteria, in particular when glycosylation and Fc effector function are not needed. For expression of antibody fragments and polypeptides in bacteria (*e.g.*, U.S. Patent Nos. 5648237, 5789199, and 5840523. After expression, the antibody may be isolated from the bacterial cell paste in a soluble fraction and can be further purified.

[0413] In addition to prokaryotes, eukaryotic microorganisms, such as filamentous fungi or yeast, are also suitable cloning or expression hosts for antibody-encoding vectors, including fungi and yeast strains whose glycosylation pathways have been “humanized,” resulting in the production of an antibody with a partially or fully human glycosylation pattern (*e.g.*, Gerngross *Nat. Biotech.* 22:1409-1414 (2004); and Li et al. *Nat. Biotech.* 24:210-215 (2006)).

[0414] Suitable host cells for the expression of glycosylated antibody can also be derived from multicellular organisms (invertebrates and vertebrates). Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains have been identified which may be used in conjunction with insect cells, particularly for transfection of *Spodoptera frugiperda* cells. Plant cell cultures can also be utilized as hosts (*e.g.*, U.S. Patent Nos. 5959177, 6040498, 6420548, 7125978, and 6417429, describing PLANTIBODIES™ technology for producing antibodies in transgenic plants).

[0415] Vertebrate cells may also be used as hosts. For example, mammalian cell lines that are adapted to grow in suspension may be useful. Other examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7); human embryonic kidney line (293 or 293 cells as described, *e.g.*, in Graham et al. *J. Gen Virol.* 36:59 (1977)); baby hamster kidney cells (BHK); mouse sertoli cells (TM4 cells as described, *e.g.*, in Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1); African green monkey kidney cells (VERO-76); human cervical carcinoma cells (HELA); canine kidney cells (MDCK; buffalo rat liver cells (BRL 3A); human lung cells (W138); human liver

cells (Hep G2); mouse mammary tumor (MMT 060562); TRI cells, as described, *e.g.*, in Mather et al. *Annals N.Y. Acad. Sci.* 383:44-68 (1982); MRC 5 cells; and FS4 cells. Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including DHFR- CHO cells (Urlaub et al. *Proc. Natl. Acad. Sci. USA* 77:4216 (1980)); and myeloma cell lines such as Y0, NS0 and Sp2/0. For a review of certain mammalian host cell lines suitable for antibody production, *see, e.g.*, Yazaki and Wu, *Methods in Molecular Biology, Vol. 248* (B.K.C. Lo, ed., Humana Press, Totowa, NJ), pp. 255-268 (2003).

Pharmaceutical compositions

[0416] Provided herein are pharmaceutical compositions and/or pharmaceutical formulations comprising the anti-Sortilin antibodies of the present disclosure and a pharmaceutically acceptable carrier.

[0417] In some embodiments, pharmaceutically acceptable carrier preferably are nontoxic to recipients at the dosages and concentrations employed. The antibodies described herein may be formulated into preparations in solid, semi-solid, liquid or gaseous forms. Examples of such formulations include, without limitation, tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants, gels, microspheres, and aerosols. Pharmaceutically acceptable carriers can include, depending on the formulation desired, pharmaceutically-acceptable, non-toxic carriers or diluents, which are vehicles commonly used to formulate pharmaceutical compositions for animal or human administration. In certain embodiments, the pharmaceutical composition can comprise formulation materials for modifying, maintaining or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption or penetration of the composition.

[0418] In certain embodiments, pharmaceutically acceptable carriers include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine or lysine); antimicrobials; antioxidants (such as ascorbic acid, sodium sulfite or sodium hydrogen-sulfite); buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates or other organic acids); bulking agents (such as mannitol or glycine); chelating agents (such as ethylenediamine tetraacetic acid (EDTA)); complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin); fillers; monosaccharides; disaccharides; and other carbohydrates (such as glucose, mannose or dextrans); proteins (such as serum albumin, gelatin or immunoglobulins); coloring, flavoring and diluting agents; emulsifying agents; hydrophilic polymers (such as polyvinylpyrrolidone); low molecular weight polypeptides; salt-forming counterions (such as sodium); preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid or hydrogen peroxide); solvents (such as glycerin, propylene glycol or polyethylene glycol); sugar alcohols (such as mannitol or sorbitol); suspending agents; surfactants or wetting agents (such as pluronics, PEG, sorbitan esters, polysorbates such as polysorbate 20, polysorbate 80, triton, tromethamine, lecithin, cholesterol,

tyloxapal); stability enhancing agents (such as sucrose or sorbitol); tonicity enhancing agents (such as alkali metal halides, preferably sodium or potassium chloride, mannitol sorbitol); delivery vehicles; diluents; excipients and/or pharmaceutical adjuvants. Further examples of formulations that are suitable for various types of administration can be found in *Remington: The Science and Practice of Pharmacy*, Pharmaceutical Press 22nd ed. (2013). For a brief review of methods for drug delivery, *see*, Langer, *Science* 249:1527-1533 (1990).

[0419] Formulations suitable for parenteral administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can comprise antioxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives.

[0420] Formulations may be optimized for retention and stabilization in the brain or central nervous system. When the agent is administered into the cranial compartment, it is desirable for the agent to be retained in the compartment, and not to diffuse or otherwise cross the blood brain barrier. Stabilization techniques include cross-linking, multimerizing, or linking to groups such as polyethylene glycol, polyacrylamide, neutral protein carriers, *etc.* in order to achieve an increase in molecular weight.

[0421] Other strategies for increasing retention include the entrapment of the antibody, such as an anti-Sortilin antibody of the present disclosure, in a biodegradable or bioerodible implant. The rate of release of the therapeutically active agent is controlled by the rate of transport through the polymeric matrix, and the biodegradation of the implant. Implants may be particles, sheets, patches, plaques, fibers, microcapsules and the like and may be of any size or shape compatible with the selected site of insertion. Biodegradable polymeric compositions which may be employed may be organic esters or ethers, which when degraded result in physiologically acceptable degradation products, including the monomers. Anhydrides, amides, orthoesters or the like, by themselves or in combination with other monomers, may find use. The polymers will be condensation polymers. The polymers may be cross-linked or non-cross-linked. Of particular interest are polymers of hydroxyaliphatic carboxylic acids, either homo- or copolymers, and polysaccharides. Included among the polyesters of interest are polymers of D-lactic acid, L-lactic acid, racemic lactic acid, glycolic acid, polycaprolactone, and combinations thereof. Among the polysaccharides of interest are calcium alginate, and functionalized celluloses, particularly carboxymethylcellulose esters characterized by being water insoluble, a molecular weight of about 5 kD to 500 kD, *etc.* Biodegradable hydrogels may also be employed in the implants of the subject invention. Hydrogels are typically a copolymer material, characterized by the ability to imbibe a liquid.

Pharmaceutical dosages

[0422] An antibody provided herein (and any additional therapeutic agent) can be administered by any suitable means, including parenteral, intrapulmonary, intranasal, intralesional administration, intracerebrospinal, intracranial, intraspinal, intrasynovial, intrathecal, oral, topical, or inhalation routes. Parenteral infusions include intramuscular, intravenous administration as a bolus or by continuous infusion over a period of time, intraarterial, intra-articular, intraperitoneal, or subcutaneous administration. In some embodiments, the administration is intravenous administration. In some embodiments, the administration is subcutaneous. Dosing can be by any suitable route, *e.g.* by injections, such as intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic. Various dosing schedules including but not limited to single or multiple administrations over various time-points, bolus administration, and pulse infusion are contemplated herein.

[0423] Antibodies provided herein would be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The antibody need not be, but is optionally formulated with one or more agents currently used to prevent or treat the disorder in question. The effective amount of such other agents depends on the amount of antibody present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as described herein, or about from 1 to 99% of the dosages described herein, or in any dosage and by any route that is empirically/clinically determined to be appropriate.

[0424] Dosages for a particular anti-Sortilin antibody may be determined empirically in individuals who have been given one or more administrations of the anti-Sortilin antibody. Individuals are given incremental doses of an anti-Sortilin antibody. To assess efficacy of an anti-Sortilin antibody, a clinical symptom of any of the diseases, disorders, or conditions of the present disclosure (*e.g.*, frontotemporal dementia, Alzheimer's disease, vascular dementia, seizures, retinal dystrophy, a traumatic brain injury, a spinal cord injury, long-term depression, atherosclerotic vascular diseases, and undesirable symptoms of normal aging) can be monitored.

[0425] For the prevention or treatment of disease, the appropriate dosage of an antibody of the invention (when used alone or in combination with one or more other additional therapeutic agents) will depend on the type of disease to be treated, the type of antibody, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the

patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments.

[0426] Depending on the type and severity of the disease, about 1 µg/kg to 15 mg/kg (*e.g.*, 0.1 mg/kg-10 mg/kg) of antibody can be an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. One typical daily dosage might range from about 1 µg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment would generally be sustained until a desired suppression of disease symptoms occurs. One exemplary dosage of the antibody would be in the range from about 0.05 mg/kg to about 10 mg/kg. Thus, one or more doses of about 0.5 mg/kg, 2.0 mg/kg, 4.0 mg/kg or 10 mg/kg (or any combination thereof) may be administered to the patient. Such doses may be administered intermittently, *e.g.*, every week or every three weeks (*e.g.*, such that the patient receives from about two to about twenty, or *e.g.*, about six doses of the antibody). In certain embodiments, dosing frequency is three times per day, twice per day, once per day, once every other day, once weekly, once every two weeks, once every four weeks, once every five weeks, once every six weeks, once every seven weeks, once every eight weeks, once every nine weeks, once every ten weeks, or once monthly, once every two months, once every three months, or longer. An initial higher loading dose, followed by one or more lower doses may be administered. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

Therapeutic uses

[0427] In certain aspects, anti-Sortilin antibodies of the present disclosure can be used for preventing, reducing risk for, or treating an individual having a disease, disorder, or injury. Anti-Sortilin antibodies of the present disclosure can be used to prevent, reduce risk of, or treat cell death (*e.g.*, neuronal cell death), frontotemporal dementia, Alzheimer's disease, vascular dementia, seizures, retinal dystrophy, a traumatic brain injury, a spinal cord injury, long-term depression, atherosclerotic vascular diseases, undesirable symptoms of normal aging, dementia, mixed dementia, Creutzfeldt-Jakob disease, normal pressure hydrocephalus, amyotrophic lateral sclerosis, Huntington's disease, tauopathy disease, stroke, acute trauma, chronic trauma, lupus, acute and chronic colitis, Crohn's disease, inflammatory bowel disease, ulcerative colitis, malaria, essential tremor, central nervous system lupus, Behcet's disease, Parkinson's disease, dementia with Lewy bodies, multiple system atrophy, intervertebral disc degeneration, Shy-Drager syndrome, progressive supranuclear palsy, cortical basal ganglionic degeneration, acute disseminated encephalomyelitis, granulomatous disorders, Sarcoidosis, diseases of aging, age related macular degeneration, glaucoma, retinitis pigmentosa, retinal degeneration, respiratory tract infection, sepsis, eye infection, systemic infection, inflammatory disorders, arthritis, multiple

sclerosis, metabolic disorder, obesity, insulin resistance, type 2 diabetes, tissue or vascular damage, an injury, and/or one or more undesirable symptoms of normal aging.

[0428] In certain aspects, provided herein is a method of preventing, reducing risk for, or treating an individual having a disease, disorder, or injury, comprising administering to an individual in need thereof a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure. In some embodiments, the disease, disorder or injury is selected from the group consisting of frontotemporal dementia, progressive supranuclear palsy, Alzheimer's disease, vascular dementia, seizures, retinal dystrophy, amyotrophic lateral sclerosis, traumatic brain injury, a spinal cord injury, dementia, stroke, Parkinson's disease, acute disseminated encephalomyelitis, retinal degeneration, age related macular degeneration, glaucoma, multiple sclerosis, septic shock, bacterial infection, arthritis, and osteoarthritis.

[0429] Certain aspects of the present disclosure provide methods of increasing Progranulin levels in an individual in need thereof, such as in the brain, blood, and/or peripheral organs of the individual, by administering to the individual a therapeutically effective amount of one or more anti-Sortilin antibodies of the present disclosure. Other aspects of the present disclosure provide methods of increasing extracellular levels of Progranulin, by contacting one or more cells with one or more anti-Sortilin antibodies of the present disclosure. In some embodiments, levels of Progranulin are increased without decreasing cellular levels of Sortilin. Other aspects of the present disclosure provide methods of decreasing cellular levels of Sortilin in an individual in need thereof, such as in the brain and/or peripheral organs of the individual, by administering to the individual a therapeutically effective amount of one or more anti-Sortilin antibodies of the present disclosure. Other aspects of the present disclosure provide methods of decreasing cellular levels of Sortilin of one or more cells, comprising contacting one or more cells with one or more anti-Sortilin antibodies of the present disclosure.

[0430] Further aspects of the present disclosure provide methods for increasing the effective concentrations of Progranulin and/or reducing the effective concentrations of a neurotrophin of the present disclosure (*e.g.*, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-neurotrophins, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, Sortilin propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), PCSK9, and receptor associated protein (RAP) in an individual in need thereof, by administering to the individual a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure to inhibit the interaction between Sortilin and Progranulin, a neurotrophin of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, Sortilin propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), and/or receptor associated protein (RAP).

[0431] The present disclosure also provides methods of inhibiting the interaction between Sortilin and Progranulin, a neurotrophin of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, Sortilin propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), PCSK9, and/or receptor associated protein (RAP); as well as one or more activities of Sortilin, Progranulin, a neurotrophin of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, Sortilin propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), and/or receptor associated protein (RAP) in an individual by administering to the individual a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure.

[0432] In certain aspects, provided herein is a method of preventing, reducing risk for, or treating an individual having a disease, disorder, or injury, comprising administering to an individual in need thereof a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure.

[0433] As disclosed herein, anti-Sortilin antibodies of the present disclosure may be used for preventing, reducing risk, or treating frontotemporal dementia, Alzheimer's disease, vascular dementia, seizures, retinal dystrophy, a traumatic brain injury, a spinal cord injury, long-term depression, atherosclerotic vascular diseases, undesirable symptoms of normal aging, dementia, mixed dementia, Creutzfeldt-Jakob disease, normal pressure hydrocephalus, amyotrophic lateral sclerosis, Huntington's disease, taupathy disease, stroke, acute trauma, chronic trauma, lupus, acute and chronic colitis, Crohn's disease, inflammatory bowel disease, ulcerative colitis, malaria, essential tremor, central nervous system lupus, Behcet's disease, Parkinson's disease, dementia with Lewy bodies, multiple system atrophy, intervertebral disc degeneration, Shy-Drager syndrome, progressive supranuclear palsy, cortical basal ganglionic degeneration, acute disseminated encephalomyelitis, granulomatous disorders, Sarcoidosis, diseases of aging, age related macular degeneration, glaucoma, retinitis pigmentosa, retinal degeneration, respiratory tract infection, sepsis, eye infection, systemic infection, inflammatory disorders, arthritis, multiple sclerosis, metabolic disorder, obesity, insulin resistance, type 2 diabetes, tissue or vascular damage, an injury, and one or more undesirable symptoms of normal aging.

[0434] In some embodiments, the present disclosure provides methods of preventing, reducing risk, or treating an individual having frontotemporal dementia, Alzheimer's disease, vascular dementia, seizures, retinal dystrophy, a traumatic brain injury, a spinal cord injury, long-term depression, atherosclerotic vascular diseases, undesirable symptoms of normal aging, dementia, mixed dementia, Creutzfeldt-Jakob disease, normal pressure hydrocephalus, amyotrophic lateral sclerosis, Huntington's

disease, taupathy disease, stroke, acute trauma, chronic trauma, lupus, acute and chronic colitis, Crohn's disease, inflammatory bowel disease, ulcerative colitis, malaria, essential tremor, central nervous system lupus, Behcet's disease, Parkinson's disease, dementia with Lewy bodies, multiple system atrophy, intervertebral disc degeneration, Shy-Drager syndrome, progressive supranuclear palsy, cortical basal ganglionic degeneration, acute disseminated encephalomyelitis, granulomatous disorders, Sarcoidosis, diseases of aging, age related macular degeneration, glaucoma, retinitis pigmentosa, retinal degeneration, respiratory tract infection, sepsis, eye infection, systemic infection, inflammatory disorders, arthritis, multiple sclerosis, metabolic disorder, obesity, insulin resistance, type 2 diabetes, tissue or vascular damage, an injury, and one or more undesirable symptoms of normal aging, in an individual in need thereof by administering to the individual a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure to: (i) inhibit the interaction between Sortilin and Progranulin, a neurotrophin of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, Sortilin propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), and/or receptor associated protein (RAP); and/or (ii) inhibit one or more activities of Sortilin, Progranulin, a neurotrophin of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, Sortilin propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), and/or receptor associated protein (RAP). In some embodiments, the present disclosure provides methods of inducing wound healing in an individual in need thereof by administering to the individual a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure.

[0435] The present disclosure also provides methods of promoting cell survival, such as neuronal cell survival, by administering an anti-Sortilin antibody of the present disclosure to inhibit the interaction between Sortilin and Progranulin, a neurotrophin of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, amyloid precursor protein (APP), and the A beta peptide,. The anti-Sortilin antibody may be administered to cells *in vitro* to promote cell survival. Alternatively, the anti-Sortilin antibody may be administered *in vivo* (*e.g.*, by administering the antibody to an individual) to promote cell survival.

[0436] In certain aspects, provided herein is a method of inhibiting one or more of neuroinflammation, axonopathy characterized by short axonal outgrowth and aberrant branching, microglial activation, and inflammatory response, comprising administering to the individual a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure.

[0437] The present disclosure provides methods of inhibiting neuroinflammation, axonopathy characterized by short axonal outgrowth and aberrant branching, microglial activation, and inflammatory response and promoting wound healing, autophagy and the clearance of aggregate proteins by administering an anti-Sortilin antibody of the present disclosure to inhibit the interaction between Sortilin and Progranulin, a neurotrophin of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, amyloid precursor protein (APP), and the A beta peptide. The anti-Sortilin antibody may be administered to cells *in vitro*. Alternatively, the anti-Sortilin antibody may be administered *in vivo* (*e.g.*, by administering the antibody to an individual).

[0438] In certain aspects, provided herein is a method of promoting one or more of wound healing, autophagy, and clearance of aggregate proteins, comprising administering to the individual a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure.

[0439] In certain aspects, provided herein is a method of preventing, reducing risk, or treating an individual having arthritis, comprising administering to the individual a therapeutically effective amount of an anti-Sortilin antibody of the present disclosure.

[0440] The present disclosure also provides methods of decreasing expression of one or more pro-inflammatory mediators by administering to an individual in need thereof an anti-Sortilin antibody of the present disclosure. In some embodiments, the one or more pro-inflammatory mediators are selected from IL-6, IL12p70, IL12p40, IL-1 β , TNF- α , CXCL1, CCL2, CCL3, CCL4, and CCL5.

[0441] In some embodiments, a method of the present disclosure includes an anti-Sortilin antibody comprising two or more anti-Sortilin antibodies.

Dementia

[0442] Dementia is a non-specific syndrome (*i.e.*, a set of signs and symptoms) that presents as a serious loss of global cognitive ability in a previously unimpaired person, beyond what might be expected from normal ageing. Dementia may be static as the result of a unique global brain injury. Alternatively, dementia may be progressive, resulting in long-term decline due to damage or disease in the body. While dementia is much more common in the geriatric population, it can also occur before the age of 65. Cognitive areas affected by dementia include, without limitation, memory, attention span, language, and problem solving. Generally, symptoms must be present for at least six months to before an individual is diagnosed with dementia.

[0443] Exemplary forms of dementia include, without limitation, frontotemporal dementia, Alzheimer's disease, vascular dementia, semantic dementia, and dementia with Lewy bodies.

[0444] Without wishing to be bound by theory, it is believed that administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat dementia. In some

embodiments, administering an anti-Sortilin antibody may induce one or more Progranulin activities in an individual having dementia (e.g., neurotrophic and/or survival activity on neurons, and anti-inflammatory activity).

Frontotemporal dementia

[0445] Frontotemporal dementia (FTD) is a condition resulting from the progressive deterioration of the frontal lobe of the brain. Over time, the degeneration may advance to the temporal lobe. Second only to Alzheimer's disease (AD) in prevalence, FTD accounts for 20% of pre-senile dementia cases. The clinical features of FTD include memory deficits, behavioral abnormalities, personality changes, and language impairments (Cruts, M. & Van Broeckhoven, C., *Trends Genet.* 24:186-194 (2008); Neary, D., *et al.*, *Neurology* 51:1546-1554 (1998); Ratnavalli, E., Brayne, C., Dawson, K. & Hodges, J. R., *Neurology* 58:1615-1621 (2002)).

[0446] A substantial portion of FTD cases are inherited in an autosomal dominant fashion, but even in one family, symptoms can span a spectrum from FTD with behavioral disturbances, to Primary Progressive Aphasia, to Cortico-Basal Ganglionic Degeneration. FTD, like most neurodegenerative diseases, can be characterized by the pathological presence of specific protein aggregates in the diseased brain. Historically, the first descriptions of FTD recognized the presence of intraneuronal accumulations of hyperphosphorylated Tau protein in neurofibrillary tangles or Pick bodies. A causal role for the microtubule associated protein Tau was supported by the identification of mutations in the gene encoding the Tau protein in several families (Hutton, M., *et al.*, *Nature* 393:702-705 (1998). However, the majority of FTD brains show no accumulation of hyperphosphorylated Tau but do exhibit immunoreactivity to ubiquitin (Ub) and TAR DNA binding protein (TDP43) (Neumann, M., *et al.*, *Arch. Neurol.* 64:1388-1394 (2007)). A majority of those FTD cases with Ub inclusions (FTD-U) were shown to carry mutations in the Progranulin gene.

[0447] Progranulin mutations result in haploinsufficiency and are known to be present in nearly 50% of familial FTD cases, making Progranulin mutation a major genetic contributor to FTD. Without wishing to be bound by theory, it is believed that the loss-of-function heterozygous character of Progranulin mutations indicates that in healthy individuals, Progranulin expression plays a dose-dependent, critical role in protecting healthy individuals from the development of FTD. Accordingly, increasing levels of Progranulin by inhibiting the interaction between Sortilin and Progranulin, can prevent, reduce the risk, and/or treat FTD.

[0448] In some embodiments, administering an anti-Sortilin antibody of the present disclosure, can prevent, reduce the risk, and/or treat FTD. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having FTD.

Alzheimer's disease

[0449] Alzheimer's disease (AD) is the most common form of dementia. There is no cure for the disease, which worsens as it progresses, and eventually leads to death. Most often, AD is diagnosed in people over 65 years of age. However, the less-prevalent early-onset Alzheimer's can occur much earlier.

[0450] Common symptoms of Alzheimer's disease include, behavioral symptoms, such as difficulty in remembering recent events; cognitive symptoms, confusion, irritability and aggression, mood swings, trouble with language, and long-term memory loss. As the disease progresses bodily functions are lost, ultimately leading to death. Alzheimer's disease develops for an unknown and variable amount of time before becoming fully apparent, and it can progress undiagnosed for years.

[0451] It has been shown that Sortilin binds to amyloid precursor protein (APP) and the APP processing enzyme BACE1. Without wishing to be bound by theory, it is believed that these interactions are involved in Alzheimer's disease. Accordingly, and without wishing to be bound by theory, it is believed that anti-Sortilin antibodies of the present disclosure can be utilized to inhibit such interactions and prevent, reduce the risk of, or treat Alzheimer's disease in individuals in need thereof.

[0452] In some embodiments, and without wishing to be bound by theory, it is believed that anti-Sortilin antibodies of the present disclosure that inhibit the interaction between Sortilin and neurotrophins of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), p75, amyloid precursor protein (APP), and/or the A beta peptide, or that inhibit one or more activities of Sortilin can be utilized to prevent, reduce the risk of, or treat Alzheimer's disease in individuals in need thereof.

[0453] In some embodiments, administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat Alzheimer's disease. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having Alzheimer's disease.

Vascular Dementia

[0454] Vascular dementia (VaD) is a subtly progressive worsening of memory and other cognitive functions that is believed to be due to cerebrovascular disease (vascular disease within the brain). Cerebrovascular disease is the progressive change in our blood vessels (vasculature) in the brain (cerebrum). The most common vascular change associated with age is the accumulation of cholesterol and other substances in the blood vessel walls. This results in the thickening and hardening of the walls, as well as narrowing of the vessels, which can result in a reduction or even a complete stopping of blood flow to brain regions supplied by the affected artery. Vascular dementia patients often present with similar symptoms to Alzheimer's disease (AD) patients. However, the related changes in the brain are not due to AD pathology but to chronic reduced blood flow in the brain, eventually resulting in dementia. VaD is considered one of the most common types of dementia in older adults. Symptoms of VaD include

difficulties with memory, difficulty with organization and solving complex problems, slowed thinking, distraction or "absent mindedness," difficulty retrieving words from memory, changes in mood or behavior such as depression, irritability, or apathy, and hallucinations or delusions.

[0455] Without wishing to be bound by theory, it is believed that one or more activities of Sortilin, or one or more interactions between Sortilin and Progranulin, neurotrophins of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, lipoprotein lipase, apolipoprotein AV, and/or receptor-associated protein are involved in vascular dementia. Accordingly, and without wishing to be bound by theory, it is believed that anti-Sortilin antibodies of the present disclosure that inhibit the interaction between Sortilin and neurotrophins of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, Sortilin propeptide (Sort-pro), amyloid precursor protein (APP), the A beta peptide, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), and/or receptor associated protein (RAP); or that inhibit one or more activities of Sortilin can be utilized to prevent, reduce the risk of, or treat vascular dementia in individuals in need thereof.

[0456] In some embodiments, administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat VaD. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having VaD.

Seizures, retinal dystrophy, traumatic brain injuries, spinal cord injuries, and long-term depression

[0457] As used herein, retinal dystrophy refers to any disease or condition that involves the degeneration of the retinal. Such diseases or conditions may lead to loss of vision or complete blindness.

[0458] As used herein, seizures also include epileptic seizures, and refer to a transient symptom of abnormal excessive or synchronous neuronal activity in the brain. The outward effect can be as dramatic as a wild thrashing movement or as mild as a brief loss of awareness. Seizures can manifest as an alteration in mental state, tonic or clonic movements, convulsions, and various other psychic symptoms.

[0459] Traumatic brain injuries (TBI), may also be known as intracranial injuries. Traumatic brain injuries occur when an external force traumatically injures the brain. Traumatic brain injuries can be classified based on severity, mechanism (closed or penetrating head injury), or other features (*e.g.*, occurring in a specific location or over a widespread area).

[0460] Spinal cord injuries (SCI) include any injury to the spinal cord that is caused by trauma instead of disease. Depending on where the spinal cord and nerve roots are damaged, the symptoms can vary widely, from pain to paralysis to incontinence. Spinal cord injuries are described at various levels of

"incomplete", which can vary from having no effect on the patient to a "complete" injury which means a total loss of function.

[0461] Long-term depression (LTD) is an activity-dependent reduction in the efficacy of neuronal synapses lasting hours or longer following a long patterned stimulus. Long-term depression can occur in many areas of the central nervous system with varying mechanisms depending upon brain region and developmental progress. Long-term depression can occur in the hippocampus, cerebellum, and in different types of neurons that release various neurotransmitters. Without wishing to be bound by theory, it is believed that long-term depression may be associated with neurodegeneration, dementia, and Alzheimer's disease.

[0462] It has been shown that pro-neurotrophins (*e.g.*, pro- neurotrophin-4/5, neurotrophin-4/5, pro-NGF, pro-BDNF, *etc.*) play a role in seizures, retinal dystrophy, traumatic brain injury, spinal cord injury, and long-term depression.

[0463] Accordingly, and without wishing to be bound by theory, it is believed that anti-Sortilin antibodies of the present disclosure that inhibit the interaction between Sortilin and neurotrophins of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*); or that inhibit one or more activities of Sortilin can be utilized to prevent, reduce the risk of, or treat seizures, retinal dystrophy, traumatic brain injuries, spinal cord injuries, and/or long-term depression in individuals in need thereof.

[0464] In some embodiments, administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat seizures, retinal dystrophy, traumatic brain injuries, spinal cord injuries, and/or long-term depression. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having seizures, retinal dystrophy, traumatic brain injuries, spinal cord injuries, and/or long-term depression.

Atherosclerotic vascular diseases

[0465] As used herein, "atherosclerotic vascular disease," "ASVD," and "atherosclerosis" are used interchangeably and refer to any condition in which an artery wall thickens as a result of the accumulation of fatty materials such as cholesterol, lipids, and triglyceride. Atherosclerotic vascular diseases include, without limitation, any ASVD-associated condition, disorder, or disease, including without limitation, thromboembolism, stroke, ischemia, infarctions, coronary thrombosis, myocardial infarction (*e.g.*, heart attack), and claudication.

[0466] As disclosed herein, Sortilin proteins of the present disclosure are involved in lipid regulation, by binding lipid-associated proteins, such as receptor associated protein, lipoprotein lipase and apolipoproteins APOA5 and APOE .

[0467] Accordingly, and without wishing to be bound by theory, it is believed that anti-Sortilin antibodies of the present disclosure that inhibit the interaction between Sortilin and lipoprotein lipase (LpL), apolipoprotein AV (APOA5), apolipoprotein E (APOE), and/or receptor associated protein (RAP); or that inhibit one or more activities of Sortilin can be utilized to prevent, reduce the risk of, or treat one or more atherosclerotic vascular disease in individuals in need thereof.

[0468] In some embodiments, administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat atherosclerotic vascular disease. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having atherosclerotic vascular disease.

Undesirable symptoms of aging

[0469] As used herein, undesirable symptoms of aging include, without limitation, memory loss, behavioral changes, dementia, Alzheimer's disease, retinal degeneration, atherosclerotic vascular diseases, hearing loss, and cellular break-down.

[0470] In some embodiments, and without wishing to be bound by theory, it is believed that anti-Sortilin antibodies of the present disclosure that inhibit the interaction between Sortilin and Progranulin, neurotrophins of the present disclosure (e.g., pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, etc.), neurotensin, p75, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), and/or receptor associated protein (RAP); or that inhibit one or more activities of Sortilin can be utilized to prevent, reduce the risk of, or treat one or more undesirable symptoms of aging.

[0471] In some embodiments, administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat one or more undesirable symptoms of aging. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having one or more undesirable symptoms of aging.

Amyotrophic lateral sclerosis (ALS)

[0472] As used herein, amyotrophic lateral sclerosis (ALS) or, motor neuron disease or, Lou Gehrig's disease are used interchangeably and refer to a debilitating disease with varied etiology characterized by rapidly progressive weakness, muscle atrophy and fasciculations, muscle spasticity, difficulty speaking (dysarthria), difficulty swallowing (dysphagia), and difficulty breathing (dyspnea).

[0473] PGRN haploinsufficiency due to heterozygous loss-of-function mutations in the *GRN* gene results in a reduction of CSF PGRN levels and is causal for the development of frontotemporal dementia (FTD) with TDP-43 pathology (Sleegers *et al.*, (2009) *Ann Neurol* 65:603; Smith *et al.*, (2012) *Am J Hum Genet* 90:1102). TDP-43 has also been identified as a major pathological protein in ALS, suggesting a similarity between ALS and FTD.

[0474] For example, over twenty dominant mutations in TDP-43 have been identified in sporadic and familial ALS patients (Lagier-Tourenne *et al.*, (2009) Cell 136:1001) and TDP-43 positive aggregates are found in approximately 95% of ALS cases (Prasad *et al.*, (2019) Front Mol Neurosci 12:25). Furthermore, ALS risk genes, such as MOBP, C9ORF72, MOBKL2B, NSF and FUS, can also cause FTD (Karch *et al.*, (2018) JAMA Neurol 75:860). In addition, both PGRN and C9ORF72 mutations are associated with abnormal microglial activation, which appears to be another common pathology of FTD and ALS (Haukedal *et al.*, (2019) J Mol Biol 431:1818). Other evidence also suggests that ALS and FTD are closely related conditions with overlapping genetic, neuropathological, and clinical features (Weishaupt *et al.*, (2016) Trends Mol Med 22:769; McCauley *et al.*, (2018) Acta Neuropathol 137:715). Taken together, these results suggest that both diseases could benefit from shared treatments and that PGRN genetic variability acts as a modifier of the course of ALS.

[0475] In some embodiments, and without wishing to be bound by theory, it is believed that anti-Sortilin antibodies of the present disclosure that inhibit the interaction between Sortilin and Progranulin, neurotrophins of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), and/or receptor associated protein (RAP); or that inhibit one or more activities of Sortilin can be utilized to prevent, or treat one or more undesirable symptoms of ALS

[0476] In some embodiments, administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat ALS. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having ALS.

Depression

[0477] As used herein, depression or, major depressive disorder (MDD), clinical depression, major depression, unipolar depression, unipolar disorder, recurrent depression or, dysthymia, are used interchangeably and refer to a mental disorder characterized by episodes of all-encompassing low mood accompanied by low self-esteem and loss of interest or pleasure in normally enjoyable activities.

[0478] Accordingly, and without wishing to be bound by theory, it is believed that anti-Sortilin antibodies of the present disclosure that inhibit the interaction between Sortilin and Progranulin, neurotrophins of the present disclosure (*e.g.*, pro-neurotrophins, pro-neurotrophin-3, pro-neurotrophin-4/5, pro-NGF, pro-BDNF, neurotrophin-3, neurotrophin-4/5, NGF, BDNF, *etc.*), neurotensin, p75, lipoprotein lipase (LpL), apolipoprotein AV (APOA5), and/or receptor associated protein (RAP); or that inhibit one or more activities of Sortilin can be utilized to prevent, or treat one or more undesirable symptoms of depression.

[0479] In some embodiments, administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat depression. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having depression.

Parkinson's disease

[0480] Parkinson's disease, which may be referred to as idiopathic or primary parkinsonism, hypokinetic rigid syndrome (HRS), or paralysis agitans, is a neurodegenerative brain disorder that affects motor system control. The progressive death of dopamine-producing cells in the brain leads to the major symptoms of Parkinson's. Most often, Parkinson's disease is diagnosed in people over 50 years of age. Parkinson's disease is idiopathic (having no known cause) in most people. However, genetic factors also play a role in the disease.

[0481] Symptoms of Parkinson's disease include, without limitation, tremors of the hands, arms, legs, jaw, and face, muscle rigidity in the limbs and trunk, slowness of movement (bradykinesia), postural instability, difficulty walking, neuropsychiatric problems, changes in speech or behavior, depression, anxiety, pain, psychosis, dementia, hallucinations, and sleep problems.

[0482] In some embodiments, administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat Parkinson's disease. In some embodiments, administering an anti-Sortilin antibody may induce one or more Progranulin activities in an individual having Parkinson's disease. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having Parkinson's disease.

Huntington's disease

[0483] Huntington's disease (HD) is an inherited neurodegenerative disease caused by an autosomal dominant mutation in the Huntingtin gene (HTT). Expansion of a cytokine-adenine-guanine (CAG) triplet repeat within the Huntingtin gene results in production of a mutant form of the Huntingtin protein (Htt) encoded by the gene. This mutant Huntingtin protein (mHtt) is toxic and contributes to neuronal death. Symptoms of Huntington's disease most commonly appear between the ages of 35 and 44, although they can appear at any age.

[0484] Symptoms of Huntington's disease, include, without limitation, motor control problems, jerky, random movements (chorea), abnormal eye movements, impaired balance, seizures, difficulty chewing, difficulty swallowing, cognitive problems, altered speech, memory deficits, thinking difficulties, insomnia, fatigue, dementia, changes in personality, depression, anxiety, and compulsive behavior.

[0485] In some embodiments, administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat Huntington's disease. In some embodiments, administering an anti-Sortilin antibody may induce one or more Progranulin activities in an individual having Huntington's

disease. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having Huntington's disease.

Taupathy disease

[0486] Taupathy diseases, or Tauopathies, are a class of neurodegenerative disease caused by aggregation of the microtubule-associated protein tau within the brain. Alzheimer's disease (AD) is the most well-known taupathy disease, and involves an accumulation of tau protein within neurons in the form of insoluble neurofibrillary tangles (NFTs). Other taupathy diseases and disorders include progressive supranuclear palsy, dementia pugilistica (chronic traumatic encephalopathy), frontotemporal dementia and parkinsonism linked to chromosome 17, LYTIC-Bodig disease (Parkinson-dementia complex of Guam), Tangle-predominant dementia, Ganglioglioma and gangliocytoma, Meningioangiomas, Subacute sclerosing panencephalitis, lead encephalopathy, tuberous sclerosis, Hallervorden-Spatz disease, lipofuscinosis, Pick's disease, corticobasal degeneration, Argyrophilic grain disease (AGD), Huntington's disease, and frontotemporal lobar degeneration.

[0487] In some embodiments, administering an anti-Sortilin antibody of the present disclosure, can prevent, reduce the risk, and/or treat taupathy disease. In some embodiments, administering an anti-Sortilin antibody may induce one or more Progranulin activities in an individual having a taupathy disease. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having a taupathy disease.

Multiple sclerosis

[0488] Multiple sclerosis (MS) can also be referred to as disseminated sclerosis or encephalomyelitis disseminata. MS is an inflammatory disease in which the fatty myelin sheaths around the axons of the brain and spinal cord are damaged, leading to demyelination and scarring as well as a broad spectrum of signs and symptoms.

[0489] Symptoms of MS include, without limitation, changes in sensation, such as loss of sensitivity or tingling; pricking or numbness, such as hypoesthesia and paresthesia; muscle weakness; clonus; muscle spasms; difficulty in moving; difficulties with coordination and balance, such as ataxia; problems in speech, such as dysarthria, or in swallowing, such as dysphagia; visual problems, such as nystagmus, optic neuritis including phosphenes, and diplopia; fatigue; acute or chronic pain; and bladder and bowel difficulties; cognitive impairment of varying degrees; emotional symptoms of depression or unstable mood; Uhthoff's phenomenon, which is an exacerbation of extant symptoms due to an exposure to higher than usual ambient temperatures; and Lhermitte's sign, which is an electrical sensation that runs down the back when bending the neck.

[0490] In some embodiments, administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat multiple sclerosis. In some embodiments, administering an anti-

Sortilin antibody may induce one or more Progranulin activities in an individual having multiple sclerosis. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having multiple sclerosis.

Glaucoma and macular degeneration

[0491] Glaucoma describes, without limitation, a group of diseases that are characterized by a damaged optic nerve, resulting in vision loss and blindness. Glaucoma is usually caused by increased fluid pressure (= intraocular pressure) in the anterior chamber underneath the cornea. Glaucoma results in the successive loss of retinal ganglion cells that are important for vision. Age-related macular degeneration usually affects older people and primarily causes loss of vision in the macula, the central field of vision. Macular degeneration causes, without limitation, drusen, pigmentary changes, distorted vision, hemorrhages of the eye, atrophy, reduced visual acuity, blurred vision, central scotomas, reduced color vision and reduced contrast sensitivity.

[0492] Without wishing to be bound by theory, it is believed that administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat glaucoma and macular degeneration. In some embodiments, administering an anti-Sortilin antibody may induce one or more Progranulin activities in an individual having glaucoma or macular degeneration. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having glaucoma or macular degeneration.

Degenerative disc disease (DDD)

[0493] Degenerative disc disease (DDD) describes, without limitation, a group of diseases in which intervertebral disc (IVD) undergoes extensive morphological as well as biomechanical changes, and usually manifests clinically in patients with lower back pain. Degenerative discs typically show degenerative fibrocartilage and clusters of chondrocytes, suggestive of repair. Inflammation may or may not be present. The pathologic findings in DDD include protrusion, spondylolysis, and/or subluxation of vertebrae (spondylolisthesis) and spinal stenosis.

[0494] Without wishing to be bound by theory, it is believed that administering an anti-Sortilin antibody of the present disclosure can prevent, reduce the risk, and/or treat DDD. In some embodiments, administering an anti-Sortilin antibody may induce one or more Progranulin activities in an individual having DDD. In some embodiments, administering an anti-Sortilin antibody may modulate one or more Sortilin activities in an individual having DDD.

Pain

[0495] Pain describes, without limitation, neuropathic pain arising from nerve injury, e.g., from trauma or disease. Such injury may include injury to peripheral nerves and/or the spinal cord. Sortilin

antagonists have been shown to alleviate neuropathic pain. (Richner M *et al.* (2019) *Sci. Adv.* 5: eaav9946.)

Kits/Articles of Manufacture

[0496] Provided herein are articles of manufacture (*e.g.*, kit) comprising an anti-Sortilin antibody described herein. Article of manufacture may include one or more containers comprising an antibody described herein. Containers may be any suitable packaging including, but is not limited to, vials, bottles, jars, flexible packaging (*e.g.*, sealed Mylar or plastic bags), and the like. The containers may be unit doses, bulk packages (*e.g.*, multi-dose packages) or sub-unit doses.

[0497] In some embodiments, the kits may further include a second agent. In some embodiments, the second agent is a pharmaceutically-acceptable buffer or diluting agent including, but not limited to, such as bacteriostatic water for injection (BWFI), phosphate- buffered saline, Ringer's solution and dextrose solution. In some embodiments, the second agent is a pharmaceutically active agent.

[0498] In some embodiments of any of the articles of manufacture, the article of manufactures further include instructions for use in accordance with the methods of this disclosure. The instructions generally include information as to dosage, dosing schedule, and route of administration for the intended treatment. In some embodiments, these instructions comprise a description of administration of the isolated antibody of the present disclosure (*e.g.*, an anti-Sortilin antibody described herein) to prevent, reduce risk, or treat an individual having a disease, disorder, or injury selected from dementia, frontotemporal dementia, Alzheimer's disease, gauché's disease, vascular dementia, seizures, retinal dystrophy, a traumatic brain injury, a spinal cord injury, atherosclerotic vascular diseases, undesirable symptoms of normal aging, amyotrophic lateral sclerosis (ALS), long-term depression, Parkinson's disease, Huntington's disease, Taupathy disease, multiple sclerosis, age related macular degeneration, glaucoma, degenerative disc disease (DDD), Creutzfeldt-Jakob disease, normal pressure hydrocephalus, Nasu-Hakola disease, stroke, acute trauma, chronic trauma, lupus, acute and chronic colitis, Crohn's disease, inflammatory bowel disease, ulcerative colitis, malaria, essential tremor, central nervous system lupus, Behcet's disease, mixed dementia, dementia with Lewy bodies, multiple system atrophy, Shy-Drager syndrome, progressive supranuclear palsy, cortical basal ganglionic degeneration, acute disseminated encephalomyelitis, granulomatous disorders, sarcoidosis, diseases of aging, retinitis pigmentosa, retinal degeneration, respiratory tract infection, sepsis, eye infection, systemic infection, lupus, arthritis, and wound healing, according to any methods of this disclosure. In some embodiments, the disease, disorder, or injury is frontotemporal dementia. In some embodiments, the instructions include instructions for use of the anti-Sortilin antibody and the second agent (*e.g.*, second pharmaceutically active agent).

Diagnostic uses

[0499] The isolated antibodies of the present disclosure (*e.g.*, an anti-Sortilin antibody described herein) also have diagnostic utility. This disclosure therefore provides for methods of using the antibodies of this disclosure, or functional fragments thereof, for diagnostic purposes, such as the detection of a Sortilin protein in an individual or in tissue samples derived from an individual.

[0500] In some embodiments, the individual is a human. In some embodiments, the individual is a human patient suffering from, or at risk for developing a disease, disorder, or injury of the present disclosure. In some embodiments, the diagnostic methods involve detecting a Sortilin protein in a biological sample, such as a biopsy specimen, a tissue, or a cell. An anti-Sortilin antibody described herein is contacted with the biological sample and antigen-bound antibody is detected. For example, a biopsy specimen may be stained with an anti-Sortilin antibody described herein in order to detect and/or quantify disease-associated cells. The detection method may involve quantification of the antigen-bound antibody. Antibody detection in biological samples may occur with any method known in the art, including immunofluorescence microscopy, immunocytochemistry, immunohistochemistry, ELISA, FACS analysis, immunoprecipitation, or micro-positron emission tomography. In certain embodiments, the antibody is radiolabeled, for example with ^{18}F and subsequently detected utilizing micro-positron emission tomography analysis. Antibody-binding may also be quantified in a patient by non-invasive techniques such as positron emission tomography (PET), X-ray computed tomography, single-photon emission computed tomography (SPECT), computed tomography (CT), and computed axial tomography (CAT).

[0501] In other embodiments, an isolated antibody of the present disclosure (*e.g.*, an anti-Sortilin antibody described herein) may be used to detect and/or quantify, for example, microglia in a brain specimen taken from a preclinical disease model (*e.g.*, a non-human disease model). As such, an isolated antibody of the present disclosure (*e.g.*, an anti-Sortilin antibody described herein) may be useful in evaluating therapeutic response after treatment in a model for a nervous system disease or injury such as frontotemporal dementia, Alzheimer's disease, vascular dementia, seizures, retinal dystrophy, atherosclerotic vascular diseases, Nasu-Hakola disease, or multiple sclerosis, as compared to a control.

[0502] The present disclosure will be more fully understood by reference to the following Examples. They should not, however, be construed as limiting the scope of the present disclosure. All citations throughout the disclosure are hereby expressly incorporated by reference.

EXAMPLES

[0503] Examples 1-3 describe the generation of affinity-matured variants of an anti-human Sortilin antibody, S-60, and the characterization of those variants with respect to not only their binding affinity for

SORT1 but also their biological activity in down-regulating SORT1 expression, increasing PGRN secretion, and blocking PGRN-SORT1 binding. Surprisingly, it was found that improved biological activity did not necessarily correlate with increased affinity.

Example 1: Generation of S-60 Variants and Measurement of SORT1 Binding Affinity

[0504] The purpose of the following Example was to generate affinity-matured variants of an anti-human Sortilin antibody, S-60, and to characterize the binding of the affinity-matured antibodies to human Sortilin (SORT1).

S-60 affinity matured variants

[0505] Affinity-matured antibodies against human Sortilin (SORT1) were generated and their binding to SORT1 measured. The anti-SORT1 antibody, S-60, as well as S-60-1, S-60-2, S-60-3, S-60-4, S-60-5, S-60-6, S-60-7, S-60-8, and S-60-9 have been described in WO2016164637. The V_H and V_L sequences for S-60-9 and S60 in WO2016164637 are identical, and S-60 will be used herein to refer to the antibody. The V_H and V_L sequences for S-60-5, S-60-6, and S-60-7 in WO2016164637 are identical, and S-60-7 will be used herein to refer to the antibody. S-60 was found to be both a potent downregulator of SORT1 levels and a potent blocker of binding between SORT1 and PGRN (*see, e.g.*, Tables 8-10 and FIGs 6A and 10D in WO2016164637). Therefore, S-60 was selected for further affinity maturation.

[0506] The anti-SORT1 variants S-60-10, S-60-11, S-60-12, S-60-13, S-60-14, S-60-15, S-60-16, S-60-18, S-60-19, and S-60-24 were generated as described below.

Production of S-60 affinity matured variants

[0507] The anti-SORT1 antibody S-60 was affinity-matured. Briefly, diversified antibody libraries were created in yeast for each of the starting parental antibodies. The diversity in the first round of affinity maturation was created by utilizing standard molecular cloning techniques to combine the parental heavy chain HVR-H3 and light chain (LC) with pre-existing genetic diversity in the HVR-H1 and HVR-H2 regions of the heavy chain (HC) (termed "H1/H2" optimization). For the second round of affinity maturation, both heavy chain variable region (V_H) and light chain variable region (V_L) sequences were optimized, with a particular focus on HVR-H3.

[0508] Selection pressures used for screening the libraries included human SORT1 antigen equilibrium titration, parental antibody Fab competition kinetics, and the use of polyspecificity reagent deselection. FACS flow cytometry was then employed to visualize and select antibodies, using standard techniques (*see, e.g.*, Chao et al. *Nature Protocols*, 2006). The desired population was then carried forward into additional selection rounds.

[0509] The affinity-matured anti-SORT1 antibodies were purified as follows: clones were grown to saturation and then induced for 48 h at 30°C with shaking. After induction, yeast cells were pelleted and the supernatants were harvested for purification. Immunoglobulins were purified using a Protein A

column and eluted with acetic acid, pH 2.0. Fab fragments were generated by papain digestion and purified over CaptureSelect IgG-CH1 affinity matrix (LifeTechnologies).

Measurement of binding affinity to SORT1

[0510] The anti-SORT1 antibody S-60 and affinity-matured variants thereof were tested and compared in a SORT1 binding assay. For the binding assay, stable HEK293T cells expressing human SORT1 were harvested by trypsinization, washed in PBS, counted and plated on 96-well u-bottom plates at 1×10^5 cells/ well. The plates were spun at 1,400 rpm for 3 minutes and primary anti-SORT1 or control antibodies were added in FACS buffer (PBS + 2 FBS) and incubated on ice for one hour. Cells were subsequently centrifuged as before and washed thrice with FACS buffer. Cells were then incubated with anti-human APC conjugated secondary antibody (BD Biosciences) in FACS buffer for 30 minutes on ice. Cells were again washed three times with FACS buffer and analyzed on a BD FACS Canto or an Intellicyt Flow Cytometer. Binding was measured as median fluorescent intensity (MFI) in the PE channel of the GFP positive cell population.

Results

[0511] The binding affinities of the anti-SORT1 antibodies to SORT1 are shown below in **Table 1**. All antibodies were tested on the huIgG1 backbone.

Table 1: Binding Affinity to SORT1.

Antibody ID	K _D (M)	Fold change from S-60
S-60	1.15E-09*	
S-60-1	7.221E-09	0.16
S-60-2	1.101E-08	0.10
S-60-3	8.56E-10*	1.34
S-60-4	6.34E-10	1.81
S-60-7	7.87E-10	1.46
S-60-8	4.68E-10	2.46
S-60-10	6.5E-10	1.77
S-60-11	6.5E-10	1.77
S-60-12	8.9E-10	1.29
S-60-13	7.5E-10	1.53
S-60-14	1.05E-09	1.10
S-60-15	5.60E-10	2.05
S-60-16	7.80E-10	1.47
S-60-18	1.63E-09	0.71
S-60-19	8.00E-10	1.44
S-60-24	7.55E-10*	1.52

*Note: K_D values represent the average of two experiments.

[0512] Anti-SORT1 antibodies S-60-3, S-60-4, S-60-7, S-60-8, S-60-10, S-60-11, S-60-13, S-60-15, S-60-16, S-60-19, and S-60-24 showed significantly improved affinity for binding to SORT1 compared to S-60. In particular, S-60-8 and S-60-15 demonstrated the highest binding affinities to SORT1 of the tested variants, having a 2.46- and 2.05-increased fold change, respectively, compared to the binding affinity of the parental antibody, S-60.

Example 2: Effect of S-60 Variants on SORT1 Expression and PGRN Secretion

[0513] The purpose of the following Example was to characterize the effect of affinity-matured variants of an anti-human SORT1 antibody, S-60, on SORT1 expression and extracellular levels of Progranulin (PGRN).

SORT1 Expression and PGRN Secretion Assays

[0514] Affinity matured variants of S-60 were generated as described in Example 1 and were subsequently screened by FACS for their abilities to downregulate SORT1 and to elevate extracellular levels of PGRN. In order to assay the effect of the anti-SORT1 antibodies on SORT1 expression and PGRN secretion, U251 cells that endogenously express SORT1 and secrete PGRN were incubated with anti-SORT1 antibody S-60 and affinity-matured variants. Cell surface levels of SORT1 at different antibody concentrations were measured by FACS as follows: anti-SORT1 antibody S-60 and affinity-matured variants were added to U251 cells seeded at 3×10^3 cells/well in 96 well plates with a serial dilution from 0.023nM to 150nM. After 72 h, cells were harvested with Trypsin, washed in PBS and labeled with Dylight-650 conjugated anti-SORT1 antibody S-2-11 (described in WO2016164637) to quantify the levels of SORT1 protein remaining. After cells were incubated with 5 μ g/ml S-2-11 conjugated to DyLight-650 (Invitrogen) for one hour on ice, cells were washed three times in PBS+ 2 FBS and binding was quantified using a FACSCanto™ or Intellicyt Flow cytometer as median fluorescence intensity (MFI) of APC.

[0515] The amount of PGRN secreted into the cell supernatant over the course of the 72h was measured by standard ELISA as follows: cells were harvested and the media was collected; the concentration of PGRN in the media samples was measured using an R&D Systems human PGRN DuoSet ELISA kit, according to manufacturer's instructions. Data were analyzed in Microsoft Excel and GraphPad Prism.

Results

[0516] The half maximal effective concentration (EC_{50}) for down-regulation (DR) of cell surface levels of SORT1 for each of the tested S-60 variants is shown in **Table 2**. SORT1 DR was quantified for each S-60 variant tested and expressed as a percent of the control (untreated cells) (**Tables 2-4**). Additionally, the level of extracellular PGRN secretion was quantified and expressed as a fold change

relative to the untreated control as shown in **Tables 2-4**. Independent experiments are represented in separate tables (**Tables 2-4**). All antibodies were tested on the huIgG1 WT backbone.

Table 2: Percent SORT1 Downregulation (DR) and PGRN Secretion for S-60, S-60-1, S-60-2, S-60-3, S-60-4, S-60-7, and S-60-8

Antibody ID	DR EC ₅₀ (M)	DR at 0.6 nM antibody (IgG)	PGRN secretion (fold over control) 0.62 or 0.63 nM antibody (IgG)
S-60	6.81E-10	29.3	1.38
S-60-1	2.73E-10	50.4	1.60
S-60-2	4.20E-10	40.3	1.75
S-60-3	2.69E-10	43.0	1.77
S-60-4	4.18E-10	39.6	1.61
S-60-7	4.47E-10	32.6	1.40
S-60-8	5.76E-10	32.0	1.54

Table 3: Percent SORT1 Downregulation (DR) and PGRN Secretion for S-60, S-60-3, S-60-24, S-60-15, S-60-16, S-60-18, and S-60-19.

Antibody ID	DR at 1.25nM antibody (IgG)	PGRN secretion (fold over control), 0.63nM antibody (IgG)
S-60	40.17	1.11
S-60-3	61.50	1.32
S-60-24	65.20	1.35
S-60-15	60.92	1.42
S-60-16	70.14	1.42
S-60-18	51.33	1.13
S-60-19	55.46	1.35

Table 4: Percent SORT1 Downregulation (DR) and PGRN Secretion for S-60-24, S-60-10, S-60-11, S-60-12, S-60-13, and S-60-14.

Antibody ID	DR at 0.63nM antibody (IgG)	PGRN secretion (fold over control), 0.63nM antibody (IgG)
S-60-24	81.48	1.62
S-60-10	85.41	1.85
S-60-11	88.76	1.89
S-60-12	87.38	1.89
S-60-13	82.33	1.77
S-60-14	80.47	1.81

[0517] In a first experiment analyzing the SORT1 DR EC₅₀, the percent of SORT1 DR using a saturating antibody concentration of 0.6 nM IgG, and PGRN secretion using a saturating antibody concentration of 0.62 or 0.63 nM IgG, the following variants were tested: S-60, S-60-1, S-60-2, S-60-3,

S-60-4, S-60-7, and S-60-8. As shown in **Table 2**, S-60-3 demonstrated the lowest DR EC₅₀ (.269 nM) and the highest PGRN secretion, a 1.77-fold increase relative to the control, and an improvement over S-60, which showed a 1.38-fold increase over the control. Using 0.6 nM IgG, S-60-1 exhibited the most substantial effect on SORT1 expression of the variants tested, down-regulating SORT1 by 50.4% compared to the control, whereas S-60 down-regulated SORT1 by 29.3%.

[0518] In a subsequent experiment, the percent of DR using a saturating antibody concentration of 1.25 nM IgG, and PGRN secretion using a saturating antibody concentration of 0.63 nM IgG was analyzed for the following variants: S-60, S-60-3, S-60-24, S-60-15, S-60-16, S-60-18, and S-60-19. As shown in **Table 3**, both S-60-15 and S-60-16 demonstrated the highest levels of extracellular PGRN secretion with a 1.42-fold increase compared to the control, an improvement over S-60 which showed a 1.11-fold increase over the control. Indeed, these two variants also significantly reduced SORT1 expression compared to S-60, down-regulating SORT1 by 60.92% and 70.14%, respectively, relative to the control, whereas S-60 downregulated SORT1 by only 40.17%.

[0519] In a third experiment, the percent of DR using a saturating antibody concentration of 0.63 nM IgG, and PGRN secretion using a saturating antibody concentration of 0.63 nM IgG was analyzed for the following variants: S-60-24, S-60-10, S-60-11, S-60-12, S-60-13, and S-60-14. As shown in **Table 4**, S-60-11 and S-60-12 displayed the highest amount of extracellular PGRN secretion with a 1.89-fold increase compared to the control. S-60-11 also demonstrated the highest SORT1 DR at 0.63 nM IgG, down-regulating SORT1 by 88.76% relative to the control.

Conclusion

[0520] As shown in **Table 1** in **Example 1**, S-60-8 and S-60-15 demonstrated the highest affinities for binding to SORT1 compared to S-60 and the other S-60 variants tested with the lowest K_D values of 4.68E-10M and 5.60E-10, respectively. However, when compared to a specific variant, S-60-3, having a higher K_D value of 8.56E-10M (**Table 1**), S-60-8 displayed a smaller increase in PGRN secretion (**Table 2**) while S-60-15 displayed a higher fold increase in PGRN secretion than did S-60-3 (**Table 3**). Additionally, S-60-8 (at a saturating concentration of 0.6 nM IgG) demonstrated a smaller percent reduction of SORT1 expression (**Table 2**) compared to S-60-3 and S-60-15 (at a saturating concentration of 1.25 nM IgG), which down-regulated SORT1 by 61.50% and 60.92%, respectively (**Table 3**). Thus, S-60-15 performed better than S-60-3 with respect to increasing PGRN secretion and exhibited a similar effect compared to S-60-3 in down-regulating cell surface levels of SORT1, whereas S-60-8 performed worse than S-60-3 with respect to increasing PGRN secretion and down-regulating cell surface levels of SORT1.

[0521] The results of this study indicate that the relative efficacy of anti-SORT1 S-60 antibody variants with respect to down-regulating SORT1 expression or increasing PGRN secretion was not predicted by their relative binding affinities for Sortilin.

[0522] Furthermore, S-60-15 showed the greatest increase in PGRN secretion compared to S-60 and certain S-60 variants (Table 3, last column), including S-60-3. S-60-3, in turn, showed increased PGRN secretion relative to S-60-1, -2, -4, -7, and -8 (Table 2, last column). Therefore S-60-15 is an effective antibody for increasing PGRN secretion relative to those S-60 variants. The ability to increase PGRN secretion is an important property in selecting a therapeutic antibody for treatment of diseases such as FTD and other pathologies in which PGRN deficiency plays a direct causal role. Such an antibody would be useful in restoring PGRN levels to, or closer to, wild-type levels, thereby treating the disease. Moreover, this property is an important pharmacodynamic parameter in assessing antibody activity *in vivo* (see below, Example 5).

Example 3: S-60 variants block binding of PGRN to SORT1

[0523] The purpose of the following Example was to characterize the effect of affinity-matured variants of an anti-human SORT1 antibody, S-60, on the binding interaction between SORT1 and a natural ligand, PGRN.

SORT1-PGRN Blocking Assay

[0524] The ability of S-60 variants to block the binding of PGRN to SORT1 was also tested. Recombinant human PGRN (PGRN) (Adipogen) was biotinylated with an EZ-Link Micro NHS-PEG4 kit from ThermoScientific/Pierce according to the manufacturer's instructions. A stable cell line expressing full-length untagged human SORT1 (SORT1) was established by viral infection of HEK293T cells, and positive selection with hygromycin (Genscript custom project). As control cells, parental HEK293T cells were utilized.

[0525] SORT1-expressing cells or control cells were harvested and washed in PBS. Biotinylated human PGRN was added in PBS+ 2% FBS with or without a titration of anti-SORT1 antibodies or control human IgG1 isotype antibodies and incubated on ice for 2h. After washing cells 3 times in PBS+ 2% FBS, cells were incubated in Streptavidin-APC (BD Biosciences, 1:100) on ice for 30 min. Then cells were washed again, resuspended in PBS+ 2% FBS and analyzed on a FACSCanto™ flow cytometer (BD Biosciences, Mississauga, ON). PGRN binding was measured as the median fluorescence intensity (MFI) of APC of the SORT1 expressing cell population. Binding of biotinylated PGRN to SORT1 expressed on HEK293T cells was measured by FACS in absence or presence of S-60 and its variant antibodies.

Results

[0526] The half maximal effective concentration (EC_{50}) for blocking PGRN binding to SORT1 for each of the tested S-60 variants is shown in **Tables 5-7**. Additionally, the maximum level of blocking PGRN to SORT1 was quantified and expressed as a percentage of the maximum block achieved using a saturating concentration of 50 nM IgG or 150 nM IgG (**Table 5-7**). Independent experiments are shown in separate tables below.

Table 5: Blocking of PGRN binding to SORT1 expressed on HEK293T cells for S-60-1, S-60-2, S-60-3, S-60-4, S-60-7, S-60-8, and S-60.

Antibody ID	PGRN binding block EC_{50} (nM)	PGRN binding block (% max at 150nM antibody (IgG))
S-60-1	5.986	95.2
S-60-2	7.509	94.5
S-60-3	0.733	95.5
S-60-4	0.584	95.7
S-60-7	0.944	96.5
S-60-8	0.985	95.6
S-60	0.751	95.9

Table 6: Blocking of PGRN binding to SORT1 expressed on HEK293T cells for S-60, S-60-3, S-60-24, S-60-15, S-60-16, S-60-18, and S-60-19.

Antibody ID	PGRN binding block EC_{50} (M)	PGRN binding block (% max at 50nM antibody (IgG))
S-60	2.20E-09	90.13
S-60-3	1.76E-09	90.44
S-60-24	1.11E-09	88.94
S-60-15	1.22E-09	90.74
S-60-16	1.20E-09	90.44
S-60-18	2.27E-09	89.33
S-60-19	1.53E-09	90.15

Table 7: Blocking of PGRN binding to SORT1 expressed on HEK293T cells for S-60-24, S-60-10, S-60-11, S-60-12, S-60-13, and S-60-14.

Antibody ID	PGRN binding block EC_{50} (M)	PGRN binding block (% max at 50nM antibody (IgG))
S-60-24	4.70E-10	95.64
S-60-10	4.60E-10	93.84
S-60-11	4.10E-10	94.97
S-60-12	6.10E-10	93.26
S-60-13	5.60E-10	93.83
S-60-14	6.00E-10	93.67

[0527] In a first experiment analyzing the PGRN binding block EC_{50} and the percentage of PGRN binding block to SORT1 with a saturating concentration of 150 nM IgG, the following variants were tested: S-60-1, S-60-2, S-60-3, S-60-4, S-60-7, S-60-8, and S-60. As shown in **Table 5**, S-60-4 demonstrated the lowest EC_{50} value (0.584 nM), achieving a 95.7% block of PGRN binding to SORT1. Of the variants analyzed, S-60-7 displayed the greatest reduction in the PGRN binding, blocking 96.5% of PGRN binding to SORT1, though it displayed lower efficiency at blocking PGRN binding (higher EC_{50} value) than S-60 and variants S-60-3 and S-60-4.

[0528] In a subsequent experiment analyzing the PGRN binding block EC_{50} and the percentage of PGRN binding block to SORT1 with a saturating concentration of 50 nM IgG, the following variants were tested: S-60, S-60-3, S-60-24, S-60-15, S-60-16, S-60-18, and S-60-19. As shown in **Table 6**, S-60-24, S-60-15, and S-60-16 displayed the lowest EC_{50} values compared to S-60, ranging from 1.11 nM to 1.22 nM. Furthermore, the S-60-15 variant demonstrated the strongest potency compared to S-60, achieving a 90.74% block of PGRN binding to SORT1.

[0529] In a third experiment, the following variants were tested: S-60-24, S-60-10, S-60-11, S-60-12, S-60-13, and S-60-14. As shown in **Table 7**, S-60-11 demonstrated the lowest EC_{50} (0.410 nM), achieving a 94.97% block of PGRN binding to SORT1. Of the variants analyzed, S-60-24 displayed the greatest reduction in the PGRN binding, blocking 95.64% of PGRN binding to SORT1.

Conclusion

[0530] As shown in **Table 1** in **Example 1**, both S-60-8 and S-60-15 demonstrated the highest affinities for binding to SORT1 relative to S-60 and the other S-60 variants tested, with S-60-8 having the highest affinity for SORT1. However, when compared to S-60, S-60-8 displayed a lower efficiency at blocking binding to PGRN (higher EC_{50} value relative to S-60) (**Table 5**) whereas S-60-15 displayed a higher efficiency at blocking binding to PGRN (reduced EC_{50} value relative to S-60) (**Table 6**). Moreover, S-60-8 demonstrated a smaller percent maximal blocking of PGRN binding to SORT1 compared to S-60 (**Table 5**) whereas S-60-15 demonstrated a greater percent maximal blocking of PGRN binding to SORT1 relative to S-60 (**Table 6**). Thus, S-60-15 performed better than S-60-8 when compared to S-60 with respect to EC_{50} value in blocking PGRN binding to SORT1 and the maximum percentage of blocking PGRN binding to SORT1, despite the fact that S-60-8 showed higher affinity binding to SORT1 compared to S-60-15.

[0531] Thus, the results of this study indicate, unexpectedly, that the relative efficacy of anti-SORT1 S-60 antibody variants with respect to blocking the degree of PGRN binding to SORT1 was not predicted by their relative binding affinities for SORT1.

Example 4: S-60-15 Stability and Stress Testing Analysis

[0532] The purpose of the following Example was to characterize S-60-15, an affinity-matured variant of S-60 described in previous Examples, for stability under various stress testing conditions.

[0533] Stability of a therapeutic antibody is important for clinical efficacy. During manufacturing, storage, and *in vivo* administration, therapeutic antibodies are at risk for degradation via multiple pathways. The factors that are responsible for such degradation remain poorly understood. Given that S-60-15 demonstrated a high binding affinity for SORT1 (*see Example 1*), effectively reduced cell surface levels of SORT1 and increased extracellular PGRN secretion (*see Example 2*), as well substantially blocked PGRN binding to SORT1 (*see Example 3*), and unexpectedly performed these functional tests better than S-60-8 which showed higher affinity for SORT1, S-60-15 was further evaluated for stability.

pH and Temperature Stress Testing

[0534] In order to measure the stability of the amino acid residues within S-60-15, samples of S-60-15 were subjected to pH and temperature stress conditions to emulate stress conditions that occur during manufacturing, storage, and *in vivo* administration. Briefly, individual samples of S-60-15 were subjected to stress testing at pH 3.5 or pH 5.0 and at 40°C or 50°C. The samples were tested at various time points – 0, 1, 3 and 5 days for the two pH conditions, and 0, 7, 14 and 30 days for the high temperature stress conditions – by treatment with DTT and IAA followed by trypsin digestion. The digested samples were analyzed by Liquid Chromatography with Mass Spectroscopy detection using a Waters ACQUITY UPLC coupled to a Xevo G2-XS QTOF mass spectrometer using a BEH C18 column. Results are shown in **Table 8**.

Table 8. Stoichiometry Percentage Table (% of each peptide with the modifier). Deamid (deamidation); Oxid (oxidation); CAM (carbamido methyl).

S-60-15 Peptide	Modifier	Non-stress	pH5 d0	pH5 d1	pH5 d3	pH5 d5	pH3.5 d0	pH3.5 d1	pH3.5 d3	pH3.5 d5	T40 d7	T40 d14	T40 d30	T50 d7	T50 d14	T50 d30
NQFSLK (SEQ ID NO:87)	Deamid	1.3	1.5	1.5	1.5	0.9	1.5	1.4	1.5	1.4	1.3	1.3	1.3	1.3	1.3	1.4
QGYYGMDVWVGQTTVTSSASTK (SEQ ID NO:88)	Oxid M(1)	6.2	6.8	6.2	6.6	8.8	6.2	6.8	6.6	6.4	5.7	5.1	5.6	5.1	5.4	1.3
QGYYGMDVWVGQTTVTSSASTK (SEQ ID NO:88)	Deamid Q(1)	0.5	0.6	0.6	0.6	0.7	0.6	0.6	0.6	0.6	0.8	1.0	1.0	1.4	2.3	3.6
DYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTV PSSSLGTK (SEQ ID NO:89)	Deamid N(1) Oxid W(1)	3.3	3.3	3.3	3.3	3.8	3.4	3.4	3.4	3.2	3.2	3.1	0.0	3.2	3.1	3.2
DYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTV PSSSLGTK (SEQ ID NO:89)	Deamid N(1)	0.3	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.4	0.5	0.8	0.7	1.0	1.5
DTLMISR (SEQ ID NO:90)	Oxid M(1)	3.1	3.6	3.4	3.5	3.5	3.6	3.1	3.4	3.3	4.0	4.4	5.8	5.2	7.2	12.2
TPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNNAK (SEQ ID NO:91)	Deamid N(1)	1.4	1.1	1.2	1.3	1.7	1.5	1.2	1.4	1.3	0.2	0.3	1.3	1.1	0.2	5.2
VVSVLTVLHQDWLNGK (SEQ ID NO:92)	Deamid N(1)	7.8	7.2	6.7	6.9	6.5	7.0	6.9	7.0	6.8	8.1	8.4	9.2	9.3	12.6	12.0
EPQVYTLPPSQEEMTK (SEQ ID NO:93)	Oxid. M(1)	0.3	0.3	0.0	0.0	1.5	0.2	0.4	0.4	0.5	0.8	0.0	0.0	0.8	1.3	1.1
NQVSLTCLVK (SEQ ID NO:94)	Deamid N(1) CAM C(1)	1.9	2.1	2.0	2.0	1.7	2.0	2.0	2.0	1.9	1.9	2.1	1.9	2.1	2.5	2.0
GFYPSDIAVEWESNGQPENNYK (SEQ ID NO:95)	Deamid N(1)	5.2	4.9	5.1	5.0	6.4	5.0	5.0	5.2	4.9	7.0	8.9	9.6	11.1	19.0	45.9
WQEGNVFSCSVMHEALH_NHYTQK (SEQ ID NO:96)	Deamid N(1) CAM C(1)	3.8	4.0	4.0	4.0	4.4	3.9	3.0	4.0	3.9	3.7	3.7	3.9	3.9	4.4	6.7
WQEGNVFSCSVMHEALH_NHYTQK (SEQ ID NO:96)	CAM C(1), Oxid. M(1)	1.6	1.8	1.8	1.8	1.9	1.8	1.8	1.9	1.8	1.8	1.9	2.6	2.2	2.6	5.1
DIVMTQSPLSLPVTPGEPASISCR (SEQ ID NO:97)	CAM C(1), Oxid. M(1)	0.7	0.8	0.8	0.8	0.8	0.8	0.7	0.8	0.7	0.8	0.8	1.1	0.8	0.8	2.0
SSQSLLR (SEQ ID NO:98)	Deamid Q(1)	6.8	6.1	5.2	5.4	6.5	6.0	5.4	6.7	5.8	7.3	6.9	7.0	6.8	7.2	6.6

S-60-15 Peptide	Modifier	Non-stress	pH5 d0	pH5 d1	pH5 d3	pH5 d5	pH3.5 d0	pH3.5 d1	pH3.5 d3	pH3.5 d5	T40 d7	T40 d14	T40 d30	T50 d7	T50 d14	T50 d30
SNGYNYLDWYLQKPGQSP QLLIYLGSR (SEQ ID NO:99)	Deamid N(1)	11.2	9.3	9.4	9.3	12.4	9.8	9.7	9.8	9.6	15.7	20.3	21.9	30.7	45.4	65.6
AEAEDVGYYCMQQQEAP LTFGGGTK (SEQ ID NO:100)	CAM C(1), Oxid. M(1)	0.5	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.5	0.6	1.1	0.8	1.1	2.1
SGTASVVCLLNNFYPR (SEQ ID NO:101)	Deamid N(1)	2.6	2.8	2.9	2.8	3.1	2.8	2.7	2.8	2.7	2.6	2.7	2.5	2.6	3.0	3.4
VDNALQSGNSQESVTEQDSK (SEQ ID NO:102)	Deamid N(1)	0.3	0.4	0.4	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.5	0.5	0.6	0.8	1.3

[0535] Amino acids having the potential to incur post-translational modifications (PTMs) upon stress were identified, and the corresponding peptide fragments generated from the stressed samples were tracked kinetically. The results confirm that two asparagine-glycine sites, one in the Fc region of the heavy chain (SEQ ID NO:11) and one in the Fab region of the light chain (SEQ ID NO:15) of S-60-15, were the most susceptible to PTM, exhibiting 45.9% and 65.6% deamidation, respectively, after being subjected to 50°C storage for 30 days. No other PTM or cleavage site exhibited notable PTM or degradation upon stress testing.

Deamidation of the asparagine (N33) site

[0536] Any deamidation in the Fc region of the heavy chain of S-60-15 should not affect the binding affinity of S-60-15 to SORT1, as that region does not directly interact with SORT1. However, the asparagine (N33) site in the Fab region of S-60-15 is in the HVR-L1 (SEQ ID NO:8), the complementarity determining region of the light chain, which participates in antigen binding. Thus, deamidation of the asparagine (N33) site in the Fab region of S-60-15 may affect the binding affinity of S-60-15 to SORT1. Upon deamidation, the asparagine (N33) site in the Fab region of S-60-15 undergoes an Asn to Asp/IsoAsp change.

[0537] Site-directed mutagenesis was utilized to test the effects of the deamidation of the asparagine (N33) site in the Fab region of S-60-15 and the potential amino acid substitution to reduce the likelihood of degradation and deamidation and resulting manufacturing liability. Seventeen different point mutations to the HVR-L1 N33 position of S-60-15 huIgG1 were produced and sequenced using standard procedures. The G1m3 or G1m(f) allotype with L234A/L235A/P331S (“LALAPS” motif) mutations was used as a background (Jefferis R and Lefranc M-P, *Mabs*, 2009 Jul-Aug; 1(4): 332–338). The LALAPS point mutations are intended to improve overall safety by minimizing effector functions such as Fc gamma receptor binding, complement activation, and antibody dependent cell mediated cytotoxicity (ADCC). Reduction of Fc gamma receptor binding, complement activation and ADCC as a result of LALAPS mutations have been observed in various experiments (data not shown). Bio-layer interferometry (BLI) data was collected at a rate of 5 Hz at 25°C on a Pall ForteBio Octet RED96 instrument. Data analysis was performed using ForteBio Data Analysis Software, version 9.0. Standard kinetic buffer (PBS, 0.1 BSA, 0.02 Tween-20, pH 7.2) was used for the assay and for preparing reagents. For all assays, sensor tips were equilibrated in buffer prior to analysis.

[0538] For the initial off-rate screen, S-60-15 LALAPS N33 and N33X mutants (1 µg/mL, 300 s loading time) were captured on Anti-Human IgG Fc Capture Dip and Read Biosensors (Pall ForteBio, Menlo Park, CA). Twenty nM histidine-tagged human SORT1 (R&D Systems, Minneapolis, MN) was then bound to the captured anti-SORT1 surface (200 s association time, 1200 s dissociation time). The resulting BLI signal was obtained as the difference in response from the reference (1 µg/mL S60-15 + 0 nM SORT1) sensor. A zero-ligand control (0 µg/mL IgG + 20 nM SORT1) showed no measurable non-specific binding of antigen to the sensor tip surface. Full, local fitting using a 1:1 interaction model was

applied to extract association and dissociation rate constants (k_a and k_d , respectively) for each N33 mutant. Affinity constants (K_D) were calculated from the ratio k_d/k_a . Different amino acid substitutions were made at N33 and tested for binding to SORT1 in an off- rate screen (under non-stress conditions) as shown in **Table 9**.

Table 9: Extracted k_a , k_d , and K_D for S-60-15 N33X Mutants binding to SORT1

S-60-15 Variants	k_a (Ms) ⁻¹	k_d (s ⁻¹)	K_D (pM)
S-60-15 [N33 (wt)]	5.12E+05	1.41E-04	270
S-60-15.1 [N33T]	4.7E+05	1.67E-04	344
S-60-15.2 [N33S]	5.63E+05	2.47E-04	440
S-60-15.3 [N33G]	4.98E+05	4.11E-04	825
S-60-15.4 [N33R]	5.24E+05	4.60E-04	880
S-60-15.5 [N33D]	5.90E+05	6.10E-04	1000
S-60-15.6 [N33H]	5.48E+05	4.60E-04	823
S-60-15.7 [N33K]	5.63E+05	5.47E-04	965
S-60-15.8 [N33Q]	6.09E+05	6.53E-04	905
S-60-15.9 [N33Y]	5.11E+05	6.75E-04	1285
S-60-15.10 [N33E]	6.76E+05	4.30E-03	2007
S-60-15.11 [N33W]	6.01E+05	5.79E-04	960
S-60-15.12 [N33F]	2.80E+05	8.16E-04	2910
S-60-15.13 [N33I]	4.91E+05	2.28E-04	460
S-60-15.14 [N33V]	4.45E+05	3.06E-04	690
S-60-15.15 [N33A]	4.93E+05	3.15E-04	625
S-60-15.16 [N33M]	4.95E+05	4.23E-04	850
S-60-15.17 [N33L]	4.12E+05	4.25E-04	1030

Note: k_a , k_d , and K_D represent an average value from multiple experiments.

[0539] Screening this N33X mutant panel showed that the S-60-15.5 antibody (N33D), which mimics deamidation of N33, led to a significant reduction in binding, with a K_D of 1000 pM compared to 270 pM for N33 wild-type (wt) (**Table 9**). In contrast, the S-60-15.1 antibody (N33T) had the best retention of S-60-15 affinity to SORT1, displaying the smallest K_D difference from wild-type in the panel (**Table 9**).

[0540] In addition, complete kinetic analysis using multiple concentrations of SORT1 was performed for S-60-15.1 [N33T] as huIgG1 and huIgG1 LALAPS in an independent experiment. The kinetic data are displayed in **Table 10**. Kinetic data demonstrate that S60-15.1 [N33T] is a high affinity antibody both as huIgG1 and huIgG1 LALAPS.

Table 10: SORT1-binding kinetic data for S-60-15.1 [N33T] as huIgG1 and huIgG1 LALAPS.

Antibody ID	k_a (Ms) ⁻¹	k_d (s ⁻¹)	K_D (pM)
S-60-15.1 [N33T]	5.99E+05	1.79E-04	298
S-60-15.1 [N33T] LALAPS	4.79E+05	1.27E-04	260

[0541] S-60-15.1 [N33T], both as huIgG1 and huIgG1 LALAPS, and S-60-15.5 [N33D] were subsequently tested for cell binding and functionality as described below.

[0542] Affinity matured antibodies S-60, S-60-15.1 [N33T] and S-60-15.5 [N33D] were tested side-by-side for binding to human SORT1 expressed on HEK293T cells as described earlier. The results are shown in **Table 11**. S-60-15.1 [N33T] as huIgG1 or huIgG1 LALAPS showed a higher affinity than S-60 (**Table 11**), with a lower K_d and a higher B_{max} . The Fc variant did not appear to have an effect on affinity, having only marginal differences in K_d and B_{max} of S-60-15.1 [N33T] huIgG1 vs. huIgG1 LALAPS. The completely deamidated version of S-60-15.5 [N33D] did not show any significant binding (**Table 11**, N.B. (No Binding)) to human SORT1 expressing cells, suggesting that amidation of the N33 residue is essential for target binding.

Table 11: Cell binding affinity values as measured by FACS.

Antibody ID	K_d (nM)	B_{max} (MFI)
S60	1.02	758381
S-60-15.5 [N33D]	N.B.	N.B.
S-60-15.1 [N33T]	0.3651	791996
S-60-15.1 [N33T] LALAPS	0.4231	792791

[0543] U251 cells that endogenously express SORT1 and secrete PGRN were incubated with anti-SORT1 antibody S-60, S-60-15.1 [N33T] or S-60-15.5 [N33D] at different concentrations, and cell surface levels of SORT1 were measured by FACS as described in **Example 2**. The control was untreated cells. In order to quantify the percentage of Sortilin down-regulation, a saturating antibody concentration of 150 nM IgG was used. For quantification of PGRN secretion, a saturating antibody concentration of 50 nM IgG was utilized. Results are shown in **Table 12**. S-60-15.1 [N33T] as huIgG1 or huIgG1 LALAPS resulted in significant downregulation of SORT1 and an increase in PGRN secretion. Both the level of downregulation of SORT1 and the level of increase in PGRN secretion were significantly improved in S-60-15.1 [N33T] as huIgG1 or huIgG1 LALAPS as compared to S-60. S-60-15.1 [N33T] as huIgG1 displayed only a minor reduction in EC_{50} and percent SORT1 down-regulation, having a minor fold increase in PGRN secretion compared to S-60-15.1 [N33T] as huIgG1 LALAPS. In contrast, the completely deamidated version of S-60-15.5 [N33D] performed worse than S-60 in each of these functional assays; it did not show significant downregulation of SORT1 and had only minimal change in secreted PGRN levels.

Table 12: Downregulation (DR) of cell surface SORT1 by S60 antibody variants.

Antibody ID	EC ₅₀ (pM)	% DR at 150nM antibody (IgG)	PGRN secretion, fold over control at 50nM antibody (IgG)
S-60	680.9	62.5	1.75
S-60-15.5 [N33D]	n.d.	3.4	1.22
S-60-15.1 [N33T]	72.58	69.3	2.29
S-60-15.1 [N33T] LALAPS	103.6	70.3	1.97

[0544] Binding of biotinylated PGRN to SORT1 expressed on HEK293T cells was measured by FACS in the absence or presence of anti-SORT1 antibody S-60, S-60-15.1 [N33T] and S-60-15.5 [N33D] as described in **Example 3**. Results are shown in **Table 13**. A saturating antibody concentration of 150 nM IgG was used for quantifying the percentage of blocking of PGRN binding to SORT1.

Table 13: Block of PGRN binding to SORT1 expressed on HEK293T cells.

Antibody ID	EC ₅₀ (nM)	Percentage block at 150nM antibody (IgG)
S60	0.751	95.9
S-60-15.5 [N33D]	0.588	27.5
S-60-15.1 [N33T]	0.325	96.5
S-60-15.1 [N33T] LALAPS	0.405	96.9

[0545] The S-60-15.1 [N33T] variants, both with and without LALAPS, displayed the lowest EC₅₀ values and the highest level of blocking PGRN binding to SORT1 as compared to S-60-15.5 as well as compared to S-60. Thus, S-60-15.1 [N33T] performed better than S-60 and S-60-15.5 in blocking PGRN binding to SORT1.

[0546] In summary, S-60-15 was found to undergo extensive deamidation at N33 under storage stress (see **Table 8**), which increases the likelihood of degradation and resulting manufacturing liability. The previously described anti-SORT1 antibody S-60 and variants S-60-1, S-60-2, S-60-3, S-60-4, S-60-7, and S-60-8 also contain an asparagine at position 33 in the V_L. In particular, the location of the N33 deamidation site within the HVR-L1 site is likely to affect the binding of S-60-15 to SORT1. S-60-15.1 [N33T], one of several tested S-60-15 variants, retained significant binding affinity to SORT1 as compared to S-60-15 (see **Table 9**) and showed significantly improved binding affinity to SORT1 as compared to S-60 (see **Table 11**; compare to S-60-15 WT in **Table 1**). This high retention of binding affinity was unexpected for an antibody having a non-preferred substitution within HVR-L1 because the HVR-L1 residues may contact antigen. Also, unexpectedly, the S-60-15.1 [N33T] variant retained increased downregulation of SORT1, increased secretion of PGRN, and increased blocking of PGRN binding to SORT1 as compared to S-60 (see **Tables 12 and 13**). These results were also unexpected because a non-preferred substitution in an HVR sequence would likewise have been expected to negatively affect the functional characteristics of an antibody. Instead, S-60-15.1 [N33T] retains the desirable properties of downregulating cell-surface SORT1, increasing PGRN secretion, and blocking

binding of PGRN to SORT1 in addition to being more stable than previously described antibodies by removal of a manufacturing liability at position 33 in the V_L.

Example 5: Anti-Sortilin Antibody PK and PD in Non-Human Primates

[0547] In this Example, the pharmacokinetics (PK) and pharmacodynamics (PD) of intravenously (IV) administered anti-Sortilin antibody S-60-15.1 [N33T] LALAPS were determined in non-human primates.

Materials and Methods

Single dose pharmacokinetic and pharmacodynamic studies

[0548] For single dose pharmacokinetic studies, cynomolgus monkeys were administered anti-Sortilin antibody by single IV dose of 5mg/kg, 20mg/kg, 60mg/kg, or 200mg/kg on Day 0 (n = 3 animals per dose). Blood and CSF were drawn from the animals at multiple time-points thereafter to obtain anti-Sortilin antibody concentrations in plasma and cerebrospinal fluid (CSF), which are measurements of anti-Sortilin antibody pharmacokinetics. Progranulin (PGRN) concentration and the levels of Sortilin (SORT1) on white blood cells (WBCs), which are measurements of pharmacodynamics, were also determined.

[0549] Anti-Sortilin antibody concentrations were assayed using an ELISA assay with anti-Sortilin antibody -specific anti-idiotypic antibodies. PGRN concentrations were assayed with a commercially-available ELISA kit. Levels of SORT1 on white blood cells were assayed using an ELISA assay, and normalized to protein concentration.

Results

[0550] **Table 14** provides the plasma mean C_{max}, mean AUC, and t_{1/2} for each of the tested anti-Sortilin antibody doses.

Table 14. C_{max}, mean AUC, and t_{1/2} for the indicated anti-Sortilin antibody doses (n=3 for each dose).

Antibody Dose	Mean C_{max} (µg/ml)	Mean AUC (µg x hr/ml)	t ½ hours
5mg/kg	156	2,870	4.7
20mg/kg	697	26,500	13.3
60mg/kg	2,570	118,000	42
200mg/kg	7,910	366,000	73.6

[0551] As shown in **FIG. 1A**, SORT1 expression levels in peripheral white blood cells decreased after treatment of non-human primates with any of the anti-Sortilin antibody doses tested. The higher anti-Sortilin antibody doses (60 mg/kg, 200 mg/kg) resulted in both an earlier and more prolonged decrease of

SORT1 levels in peripheral white blood cells compared to lower anti-Sortilin antibody doses (5mg/kg, 20 mg/kg).

[0552] The levels of PGRN increased in the plasma of non-human primates administered a single IV injection of anti-Sortilin antibody in a time- and dose-dependent manner (**FIG. 1B**). In particular, plasma PGRN levels increased 3- to 4-fold at C_{max} , compared to baseline levels, for all anti-Sortilin antibody doses tested. Plasma PGRN levels remained elevated for longer periods of time at the higher antibody doses. Additionally, increased plasma PGRN levels were correlated with decreased expression levels of SORT1 in peripheral white blood cells.

[0553] The levels of PGRN in CSF were also increased in non-human primates administered a single IV injection of anti-Sortilin antibody. As shown in **FIG. 1C**, CSF PGRN levels increased 2- to 3-fold above baseline in animals administered either 20 mg/kg, 60 mg/kg, or 200 mg/kg. As observed with plasma PGRN levels, CSF PGRN levels remained elevated over time in the higher antibody dose groups.

[0554] **Table 15** provides the CSF mean C_{max} , mean AUC, and $t_{1/2}$ for each of the tested anti-Sortilin antibody doses in non-human primates. Anti-Sortilin antibody CSF concentrations were on average around 0.1% the amount observed in plasma.

TABLE 15. Anti-Sortilin antibody CSF PK parameters and estimated half-life in non-human primates.

[1] Level	Dose	C_{max} (µg/mL)	AUC_{all} (h*µg/mL)	CL (mL/h/kg)	$t_{1/2}$ hours (days)
5mg/kg	20	184	20692	32.3 (1.34)	
20mg/kg	2243	35717	745	23.8 (1)	
60mg/kg	6842	113573	623	38.3 (1.6)	
200mg/kg	4595	349187	1037	72.4 (3.02)	

Repeat dose pharmacokinetic and pharmacodynamic studies

[0555] Further pharmacokinetic and pharmacodynamic studies were performed in non-human primates administered anti-Sortilin antibody following a repeat-dose regimen. In these studies, animals (2 males and 2 females) were administered anti-Sortilin antibody at a dose of 60mg/kg once per week for four weeks. At various timepoints thereafter, SORT1 expression levels in peripheral white blood cells were determined. In addition, plasma and CSF levels of the anti-Sortilin antibody, were determined.

[0556] As shown in **FIG. 2A**, SORT1 levels in peripheral white blood cells remained decreased throughout the duration of the study. Plasma PGRN levels increased to 5- to 6-fold above baseline at peak levels (**FIG. 2B**). A decrease in plasma PGRN was observed following the fourth and final administration of anti-Sortilin antibody; however, the plasma PGRN levels remained elevated by 2-fold above baseline. Additionally, CSF PGRN levels were increased 3- to 4-fold above baseline (**FIG. 2C**).

[0557] The systemic anti-Sortilin antibody exposure, assessed by mean C_{max} and AUC₀₋₁₆₈, was 2100 µg/mL and 114,000 µg/mL × hr on Day 1, and 3020 µg/mL and 174,000 µg/mL × hr on Day 22. These results showed that exposure was higher on Day 22 compared to Day 1, indicating some accumulation of the antibody.

[0558] CSF concentration of anti-Sortilin antibody in these animals ranged from 0.03% to 0.12% of that observed in plasma, consistent with the distribution of other antibodies in the CSF (Pestalozzi *et al.*, (2000) J Clin Oncol 18(11):2349-51; Petereit *et al.*, (2009) Mult Scler 15(2):189-92).

[0559] Taken together, these results indicate that the S-60-15.1 [N33T] variant, in spite of having a relatively short half-life, has sustained activity *in vivo*, decreasing SORT1 levels in peripheral white blood cells and increasing plasma and CSF PGRN levels.

[0560] Moreover, IV administration of S-60-15.1 [N33T] to cynomolgous monkeys in both a 4-week and a 26-week repeat dose toxicology study was well tolerated at up to 200 mg/kg weekly, and there were no adverse findings that would preclude the conduct of clinical studies in humans based on those and other toxicology studies which included assessments of cytokine release, mortality, body weight, respiratory rate and depth, and local tolerability at injection site. Therefore, S-60-15.1 [N33T] is suitable for testing in human clinical studies of FTD and other indications as contemplated herein.

[0561] Lack of adverse findings with respect to toxicity from administration of an anti-Sortilin antibody was surprising in view of the fact that Sortilin acts as a receptor of several ligands and has multiple roles in cellular transport and signaling (Nykjaer, A *et al.*, (2012) *Trends Neurosci* 35: 261-270). Due to Sortilin's multiple ligands and functional roles, administration of an anti-Sortilin antibody might have been expected to cause off-target effects; however, the S-60-15.1 [N33T] antibody was well tolerated at the administered doses.

Sequences of the V_H regions of the S-60 antibody variants, alignment to S-60, and HVR locations

```

S-60      (SEQ ID NO: 56)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-1    (SEQ ID NO: 107) QVQLQESGPGLVKPSSETLSLTCAVSGYSISSVRYWGWIRQPPGKGLEWIGSIYHSGSTYY 60
S-60-2    (SEQ ID NO: 108) QVQLQESGPGLVKPSSETLSLTCAVSGYSISSVRYWGWIRQPPGKGLEWIGAIYPSGSTYY 60
S-60-3    (SEQ ID NO: 109) QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-4    (SEQ ID NO: 110) QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-7    (SEQ ID NO: 111) QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-8    (SEQ ID NO: 112) QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-10   (SEQ ID NO: 54)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-11   (SEQ ID NO: 54)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-12   (SEQ ID NO: 54)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-13   (SEQ ID NO: 55)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-14   (SEQ ID NO: 55)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-15   (SEQ ID NO: 56)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-15.1 (SEQ ID NO: 56)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-16   (SEQ ID NO: 56)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-18   (SEQ ID NO: 56)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-19   (SEQ ID NO: 54)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60-24   (SEQ ID NO: 56)  QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGWIRQPPGKGLEWIGTIYHSGSTYY 60
S-60 Alignment      *****
S-60 HVR H1 (SEQ ID NO: 1) -----YSISSGYYWG-----
S-60 HVR H2 (SEQ ID NO: 2) -----TIYHSGSTYY

S-60      (SEQ ID NO: 56 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60-1    (SEQ ID NO: 107 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60-2    (SEQ ID NO: 108 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60-3    (SEQ ID NO: 109 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60-4    (SEQ ID NO: 110 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60-7    (SEQ ID NO: 111 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60-8    (SEQ ID NO: 112 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60-10   (SEQ ID NO: 54 cont.) NPSLKSQVTISVDTSKNQFSLELSSVTAADTAVYYCARQGSIQQGYGMDVWGQGT122TVTVSS
S-60-11   (SEQ ID NO: 54 cont.) NPSLKSQVTISVDTSKNQFSLELSSVTAADTAVYYCARQGSIQQGYGMDVWGQGT122TVTVSS
S-60-12   (SEQ ID NO: 54 cont.) NPSLKSQVTISVDTSKNQFSLELSSVTAADTAVYYCARQGSIQQGYGMDVWGQGT122TVTVSS
S-60-13   (SEQ ID NO: 55 cont.) NPSLESRV122TSVDTSKNQFSLKLSSVTAADTAVYYCARQGSIQQGYGMDVWGQGT122TVTVSS
S-60-14   (SEQ ID NO: 55 cont.) NPSLESRV122TSVDTSKNQFSLKLSSVTAADTAVYYCARQGSIQQGYGMDVWGQGT122TVTVSS
S-60-15   (SEQ ID NO: 56 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60-15.1 (SEQ ID NO: 56 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60-16   (SEQ ID NO: 56 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60-18   (SEQ ID NO: 56 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60-19   (SEQ ID NO: 54 cont.) NPSLKSQVTISVDTSKNQFSLELSSVTAADTAVYYCARQGSIQQGYGMDVWGQGT122TVTVSS
S-60-24   (SEQ ID NO: 56 cont.) NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMDVWGQGT122TVTVSS
S-60 Alignment      *****:*****:*****:*****
S-60 HVR H2 (SEQ ID NO: 2 cont.) NPSLKS-----
S-60 HVR H3 (SEQ ID NO: 6)      -----ARQGSIKQGYGMDV-----

```

Sequences of the V_L regions of the S-60 antibody variants, alignment to S-60, and HVR locations

S-60 (SEQ ID NO: 79) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLHSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-1 (SEQ ID NO: 113) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLHSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-2 (SEQ ID NO: 114) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLHSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-3 (SEQ ID NO: 115) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLRSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-4 (SEQ ID NO: 116) DIVMTQSPLSLPVTTPGESASISCRSSQGLLRNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-7 (SEQ ID NO: 117) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLRSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-8 (SEQ ID NO: 118) DIVMTQSPLSLPVTTPGGPASISCRSSQSLLRSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-10 (SEQ ID NO: 57) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLRSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-11 (SEQ ID NO: 58) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLRSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-12 (SEQ ID NO: 59) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLRSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-13 (SEQ ID NO: 57) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLRSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-14 (SEQ ID NO: 58) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLRSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-15 (SEQ ID NO: 57) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLRSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-15.1 (SEQ ID NO: 60) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLRSTGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-16 (SEQ ID NO: 77) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLRSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-18 (SEQ ID NO: 78) DIVMTQSPLSLPVTTPGGPASISCRSSQSLLRSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-19 (SEQ ID NO: 79) DIVMTQSPLSLPVTTPGEPASISCRSSQSLLHSNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60-24 (SEQ ID NO: 80) DIVMTQSPLSLPVTTPGESASISCRSSQGLLRNGYNYLDWYLQKPGQSPQQLLIYLGSNRA 60
S-60 Alignment *****.***:*****
S-60 HVR L1 (SEQ ID NO: 27) -----RSSQSLLHSNGYNYLD-----
S-60 HVR L2 (SEQ ID NO: 29) -----LGSNRA

S-60 (SEQ ID NO: 79 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-1 (SEQ ID NO: 113 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-2 (SEQ ID NO: 114 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-3 (SEQ ID NO: 115 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-4 (SEQ ID NO: 116 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-7 (SEQ ID NO: 117 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-8 (SEQ ID NO: 118 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-10 (SEQ ID NO: 57 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-11 (SEQ ID NO: 58 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-12 (SEQ ID NO: 59 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-13 (SEQ ID NO: 57 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-14 (SEQ ID NO: 58 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-15 (SEQ ID NO: 57 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-15.1 (SEQ ID NO: 60 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-16 (SEQ ID NO: 77 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-18 (SEQ ID NO: 78 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-19 (SEQ ID NO: 79 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60-24 (SEQ ID NO: 80 cont.) SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQQQETPLTFGGGTKVEIK 112
S-60 Align. *****.*****.*****.*****.*****
S-60 HVR L2 (SEQ ID NO: 29 cont.) S-----
S-60 HVR L3 (SEQ ID NO: 33) -----MQQQETPLT-----

Table 16: Heavy chain HVR H1 sequences of anti-SORT1 antibodies

Ab(s)	HVR H1	SEQ ID NO:
S-60; S-60-10; S-60-11; S-60-12; S-60-13; S-60-14; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]; S-60-16; S-60-18; S-60-19; S-60-24	YSSSGYYWG	1

Table 17: Heavy chain HVR H2 sequences of anti-SORT1 antibodies

Ab(s)	HVR H2	SEQ ID NO:
S-60; S-60-10; S-60-11; S-60-12; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]; S-60-16; S-60-18; S-60-19; S-60-24	TIYHSGSTYYNPSL KS	2
S-60-13; S-60-14	TIYHSGSTYYNPSL ES	3
Formula I	TIYHSGSTYYNPSL X ₁ S X ₁ is K or E	4

Table 18: Heavy chain HVR H3 sequences of anti-SORT1 antibodies

Ab(s)	HVR H3	SEQ ID NO:
S-60-10; S-60-11; S-60-12; S-60-13; S-60-14; S-60-19	ARQGSIQQGYGM DV	5
S-60; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]; S-60-16; S-60-18; S-60-24	ARQGSIQQGYGM DV	6
Formula II	ARQGSIX ₁ QGYGM DV X ₁ is Q or K	7

Table 19: Light chain HVR L1 sequences of anti-SORT1 antibodies

Ab(s)	HVR L1	SEQ ID NO:
S-60-10; S-60-11; S-60-12; S-60-13; S-60-14; S-60-15 [N33 (wt)]; S-60-16; S-60-18	RSSQSLLRSNGYNY LD	8
S-60-15.1 [N33T]	RSSQSLLRSTGYNYL D	9
S-60-15.2 [N33S]	RSSQSLLRSGYNYL D	10
S-60-15.3 [N33G]	RSSQSLLRSGGYNY	11

Ab(s)	HVR L1	SEQ ID NO:
	LD	
S-60-15.4 [N33R]	RSSQSLLRSRGYNY LD	12
S-60-15.5 [N33D]	RSSQSLLRSDGYNY LD	13
S-60-15.6 [N33H]	RSSQSLLRSHGYNY LD	14
S-60-15.7 [N33K]	RSSQSLLRSKGYNY LD	15
S-60-15.8 [N33Q]	RSSQSLLRSQGYNY LD	16
S-60-15.9 [N33Y]	RSSQSLLRSYGYNY LD	17
S-60-15.10 [N33E]	RSSQSLLRSEGYNYL D	18
S-60-15.11 [N33W]	RSSQSLLRSWGYNY LD	19
S-60-15.12 [N33F]	RSSQSLLRSFGYNYL D	20
S-60-15.13 [N33I]	RSSQSLLRSIGYNYL D	21
S-60-15.14 [N33V]	RSSQSLLRSVGNYNY LD	22
S-60-15.15 [N33A]	RSSQSLLRSAGYNY LD	23
S-60-15.16 [N33M]	RSSQSLLRSMGYNY LD	24
S-60-15.17 [N33L]	RSSQSLLRSLGYNYL D	25
S-60; S-60-19	RSSQSLLHSNGYNY LD	26
S-60-24	RSSQGLLRNNGYNY LD	27
Formula III	RSSQX ₁ LLX ₂ SX ₃ GYN YLD X ₁ is S or G X ₂ is R or H X ₃ is N, T, S, G, R, D, H, K, Q, Y, E, W, F, I, V, A, M, or L	28

Table 20: Light chain HVR L2 sequences of anti-SORT1 antibodies

Ab(s)	HVR L2	SEQ ID NO:
S-60; S-60-10; S-60-11; S-60-13; S-60-14; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]; S-60-16; S-60-18; S-60-19; S-60-24	LGSNRAS	29
S-60-12	LGSNRVS	30

Ab(s)	HVR L2	SEQ ID NO:
Formula IV	LGSNRX ₁ S X ₁ is A or V	31

Table 21: Light chain HVR L3 sequences of anti-SORT1 antibodies

Ab(s)	HVR L3	SEQ ID NO:
S-60-10; S-60-11; S-60-13; S-60-14; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]; S-60-16; S-60-24	MQQQEAPLT	32
S-60; S-60-12; S-60-18; S-60-19	MQQQETPLT	33
Formula V	MQQQEX ₁ PLT X ₁ is A or T	34

Table 22: Heavy chain framework 1 sequences of anti-SORT1 antibodies

Ab(s)	VH FR1	SEQ ID NO:
S-60-10; S-60-11; S-60-12; S-60-13; S-60-14; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]; S-60-16; S-60-18; S-60-19; S-60-24	QVQLQESGPGLVKPS SETLSLTCAVSG	35

Table 23: Heavy chain framework 2 sequences of anti-SORT1 antibodies

Ab(s)	VH FR2	SEQ ID NO:
S-60-10; S-60-11; S-60-12; S-60-13; S-60-14; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]; S-60-16; S-60-18; S-60-19; S-60-24	WIRQPPGKGLEWIG	36

Table 24: Heavy chain framework 3 sequences of anti-SORT1 antibodies

Ab(s)	VH FR3	SEQ ID NO:
S-60-10; S-60-11; S-60-12; S-60-19	QVTISVDTSKNQFSL ELSSVTAADTAVYY C	37
S-60-13; S-60-14; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-	RVTISVDTSKNQFSL KLSSVTAADTAVYY C	38

Ab(s)	VH FR3	SEQ ID NO:
15.17 [N33L]; S-60-16; S-60-18; S-60-24		
Formula VI	X ₁ VTISVDTSKNQFS LX ₂ LSSVTAADTAVY YC X ₁ is Q or R X ₂ is E or K	39

Table 25: Heavy chain framework 4 sequences of anti-SORT1 antibodies

Ab(s)	VH FR4	SEQ ID NO:
S-60-10; S-60-11; S-60-12; S-60-13; S-60-14; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]; S-60-16; S-60-18; S-60-19; S-60-24	WGQGTTVTVSS	40

Table 26: Light chain framework 1 sequences of anti-SORT1 antibodies

Ab(s)	VL FR1	SEQ ID NO:
S-60-10; S-60-11; S-60-12; S-60-13; S-60-14; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]; S-60-16; S-60-19	DIVMTQSPLSLPVTP GEPASISC	41
S-60-18	DIVMTQSPLSLPVTP GGPASISC	42
S-60-24	DIVMTQSPLSLPVTP GESASISC	43
Formula VII	DIVMTQSPLSLPVTP GX ₁ X ₂ ASISC X ₁ is E or G X ₂ is P or S	44

Table 27: Light chain framework 2 sequences of anti-SORT1 antibodies

Ab(s)	VL FR2	SEQ ID NO:
S-60-10; S-60-11; S-60-13; S-60-14; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]; S-60-16; S-60-18; S-60-19; S-60-24	WYLQKPGQSPQLLI Y	45
S-60-12	WYLQKPGQPPQLLI Y	46
Formula VIII	WYLQKPGQX ₁ PQLLI	47

Ab(s)	VL FR2	SEQ ID NO:
	Y X ₁ is S or P	

Table 28: Light chain framework 3 sequences of anti-SORT1 antibodies

Ab(s)	VL FR3	SEQ ID NO:
S-60-10; S-60-13; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]	GVPDRFSGSGSGTD FTLKISRRAEADVGV YYC	48
S-60-11; S-60-12; S-60-14; S-60-19; S-60-24	GVPDRFSGSGSGTD FTLKISRVEADVGV YYC	49
S-60-16	GVPDRFSGSGSGTD FTLKISRVEADVGA YYC	50
S-60-18	GVPDRLSGSGSGTD FTLKISRVEADVGV YYC	51
Formula IX	GVPDRX ₁ SGSGSGTD FTLKISRX ₂ EAEDVG X ₃ YYC X ₁ is F or L X ₂ is A or V X ₃ is V or A	52

Table 29: Light chain framework 4 sequences of anti-SORT1 antibodies

Ab(s)	VL FR4	SEQ ID NO:
S-60-10; S-60-11; S-60-12; S-60-13; S-60-14; S-60-15 [N33 (wt)]; S-60-15.1 [N33T]; S-60-15.2 [N33S]; S-60-15.3 [N33G]; S-60-15.4 [N33R]; S-60-15.5 [N33D]; S-60-15.6 [N33H]; S-60-15.7 [N33K]; S-60-15.8 [N33Q]; S-60-15.9 [N33Y]; S-60-15.10 [N33E]; S-60-15.11 [N33W]; S-60-15.12 [N33F]; S-60-15.13 [N33I]; S-60-15.14 [N33V]; S-60-15.15 [N33A]; S-60-15.16 [N33M]; S-60-15.17 [N33L]; S-60-16; S-60-18; S-60-19; S-60-24	FGGGTKVEIK	53

Table 30: Heavy chain variable region sequences of anti-SORT1 antibodies

Ab(s)	HCVR	SEQ ID NO:
S-60-10, S-60-11, S-60-12, S-60-19	QVQLQESGPGLVKPSSETLSLTCAVSG YSISSGYYWGWIRPPGKGLEWIGTIY HSGSTYYNPSLKSQVTISVDTSKNQFS LELSSVTAADTAVYYCARQGSIQQGY YGMDVWGQGTITVTVSS	54
S-30-13, S-60-14	QVQLQESGPGLVKPSSETLSLTCAVSG YSISSGYYWGWIRPPGKGLEWIGTIY HSGSTYYNPSLESRTISVDTSKNQFS LKLSSVTAADTAVYYCARQGSIQQGY	55

Ab(s)	HCVR	SEQ ID NO:
	YGMDVWGQGTITVTVSS	
S-60, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16, S-60-18, S-60-24	QVQLQESGPGGLVVKPSETLSLTCAVSG YISSGGYYWGWIROPKGLGWIGTIY HSGSTYYNPSLKSRTISVDTSKNQFS LKLSSVTAADTAVYYCARQGSIKQGY YGMDVWGQGTITVTVSS	56

Table 31: Light chain variable region sequences of anti-SORT1 antibodies

Ab(s)	LCVR	SEQ ID NO:
S-60-10; S-60-13; S-60-15 [N33 (wt)]	DIVMTQSPLSLPVTTPGEPASISCRSSQS LLRSGYNYLDWYLQKPGQSPQLLIY LGSNRRASGVPDRFSGSGSGTDFTLKIS RAEAEDVGVYYCMQQQEAPLTFGGG TKVEIK	57
S-60-11; S-60-14	DIVMTQSPLSLPVTTPGEPASISCRSSQS LLRSGYNYLDWYLQKPGQSPQLLIY LGSNRRASGVPDRFSGSGSGTDFTLKIS RVEAEDVGVYYCMQQQEAPLTFGGG TKVEIK	58
S-60-12	DIVMTQSPLSLPVTTPGEPASISCRSSQS LLRSGYNYLDWYLQKPGQSPQLLIY LGSNRRVSGVPDRFSGSGSGTDFTLKIS RVEAEDVGVYYCMQQQETPLTFGGG TKVEIK	59
S-60-15.1 [N33T]	DIVMTQSPLSLPVTTPGEPASISCRSSQS LLRSTGYNYLDWYLQKPGQSPQLLIY LGSNRRASGVPDRFSGSGSGTDFTLKIS RAEAEDVGVYYCMQQQEAPLTFGGG TKVEIK	60
S-60-15.2 [N33S]	DIVMTQSPLSLPVTTPGEPASISCRSSQS LLRSSGYNYLDWYLQKPGQSPQLLIY LGSNRRASGVPDRFSGSGSGTDFTLKIS RAEAEDVGVYYCMQQQEAPLTFGGG TKVEIK	61
S-60-15.3 [N33G]	DIVMTQSPLSLPVTTPGEPASISCRSSQS LLRSGGYNYLDWYLQKPGQSPQLLIY LGSNRRASGVPDRFSGSGSGTDFTLKIS RAEAEDVGVYYCMQQQEAPLTFGGG TKVEIK	62
S-60-15.4 [N33R]	DIVMTQSPLSLPVTTPGEPASISCRSSQS LLRSRGYNYLDWYLQKPGQSPQLLIY LGSNRRASGVPDRFSGSGSGTDFTLKIS RAEAEDVGVYYCMQQQEAPLTFGGG TKVEIK	63
S-60-15.5 [N33D]	DIVMTQSPLSLPVTTPGEPASISCRSSQS LLRSDGYNYLDWYLQKPGQSPQLLIY LGSNRRASGVPDRFSGSGSGTDFTLKIS	64

Ab(s)	LCVR	SEQ ID NO:
	RAEAEDVGVIYCMQQQEAPLTFGGG TKVEIK	
S-60-15.6 [N33H]	DIVMTQSPLSLPVTGPGEPAISCRSSQS LLRSHGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RAEAEDVGVIYCMQQQEAPLTFGGG TKVEIK	65
S-60-15.7 [N33K]	DIVMTQSPLSLPVTGPGEPAISCRSSQS LLRSKGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RAEAEDVGVIYCMQQQEAPLTFGGG TKVEIK	66
S-60-15.8 [N33Q]	DIVMTQSPLSLPVTGPGEPAISCRSSQS LLRSQGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RAEAEDVGVIYCMQQQEAPLTFGGG TKVEIK	67
S-60-15.9 [N33Y]	DIVMTQSPLSLPVTGPGEPAISCRSSQS LLRSYGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RAEAEDVGVIYCMQQQEAPLTFGGG TKVEIK	68
S-60-15.10 [N33E]	DIVMTQSPLSLPVTGPGEPAISCRSSQS LLRSEGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RAEAEDVGVIYCMQQQEAPLTFGGG TKVEIK	69
S-60-15.11 [N33W]	DIVMTQSPLSLPVTGPGEPAISCRSSQS LLRSWGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS SRAEAEDVGVIYCMQQQEAPLTFGGG GTKVEIK	70
S-60-15.12 [N33F]	DIVMTQSPLSLPVTGPGEPAISCRSSQS LLRSFGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RAEAEDVGVIYCMQQQEAPLTFGGG TKVEIK	71
S-60-15.13 [N33I]	DIVMTQSPLSLPVTGPGEPAISCRSSQS LLRSIGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RAEAEDVGVIYCMQQQEAPLTFGGG TKVEIK	72
S-60-15.14 [N33V]	DIVMTQSPLSLPVTGPGEPAISCRSSQS LLRSVGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RAEAEDVGVIYCMQQQEAPLTFGGG TKVEIK	73
S-60-15.15 [N33A]	DIVMTQSPLSLPVTGPGEPAISCRSSQS LLRSAGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RAEAEDVGVIYCMQQQEAPLTFGGG TKVEIK	74
S-60-15.16 [N33M]	DIVMTQSPLSLPVTGPGEPAISCRSSQS	75

Ab(s)	LCVR	SEQ ID NO:
	LLRSMGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RAEAEDVGVYYCMQQQEAPLTFGGG TKVEIK	
S-60-15.17 [N33L]	DIVMTQSPLSLPVTTPGEPASISCRSSQS LLRSLGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RAEAEDVGVYYCMQQQEAPLTFGGG TKVEIK	76
S-60-16	DIVMTQSPLSLPVTTPGEPASISCRSSQS LLRSLGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RVEAEDVGAYYCMQQQEAPLTFGGG TKVEIK	77
S-60-18	DIVMTQSPLSLPVTTPGGPASISCRSSQS LLRSLGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RVEAEDVGVYYCMQQQETPLTFGGG TKVEIK	78
S-60, S-60-19	DIVMTQSPLSLPVTTPGEPASISCRSSQS LLHSLGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RVEAEDVGVYYCMQQQETPLTFGGG TKVEIK	79
S-60-24	DIVMTQSPLSLPVTTPGESASISCRSSQG LLRSLGYNYLDWYLQKPGQSPQLLIY LGSNRASGVDPDRFSGSGSGTDFTLKIS RVEAEDVGVYYCMQQQEAPLTFGGG TKVEIK	80

Table 32: Sortilin amino acid sequences

Description	Sequence	SEQ ID NO
Human Sortilin	10 20 30 40 50 MERPWGAADG LSRWPHGLGL LLLLQLLPPS TLSQDRLDAP PPPAAPLPRW 60 70 80 90 100 SGPIGVSWGL RAAAAGGAFP RGGRWRRSAP GEDEECGRVR DFLAKLANNT 110 120 130 140 150 HQHVFDDLRL SVSLSWVGDS TGVILVLTTF HVPLVIMTFG QSKLYRSEDY 160 170 180 190 200 GKNFKDITDL INNTFIRTEF GMAIGPENSG KVLTAEVSG GSRGGRIFRS 210 220 230 240 250 SDFAKNFVQT DLPFHPLTQM MYSPQNSDYL LALSTENGLW VSKNFGGKWE 260 270 280 290 300 EIHKAVCLAK WGSNTIFFT TYANGSCKAD LGALELWRTS DLGKSFKTIG 310 320 330 340 350 VKIYSFGLGG RFLFASVMAD KDTTRRIHVS TDQGDWTSMA QLPSVGQEQF 360 370 380 390 400 YSILAANDDM VFMHVDEPGD TGFGTIFTSD DRGIVYSKSL DRHLYTTTGG 410 420 430 440 450 ETDFTNVTSL RGVYITSVLS EDNSIQTMIT FDQGGRWTHL RKPENSECDA 460 470 480 490 500 TAKNKNECSL HIHASYSISQ KLNVPMAPLS EPNAVGIIVIA HGSVGDAISV 510 520 530 540 550 MVPDVYISDD GGYSWTKMLE GPHYTTILDS GGIIVAIEHS SRPINVIKFS	81

Description	Sequence	SEQ ID NO
	560 570 580 590 600 TDEGQCWQTY TFTRDPIYFT GLASEPGARS MNISIWGFTE SFLTSQWVSY 610 620 630 640 650 TIDFKDILER NCEEKDYTIW LAHSTDPEY EDGCILGYKE QFLRLRKSSV 660 670 680 690 700 CQNGRDYVVT KQPSICLCSL EDFLCDFGY RPENDSKCVE QPELKGHDLE 710 720 730 740 750 FCLYGREEHL TTNGYRKIPG DKCQGGVNPV REVKDLKKKC TSNFLSPEKQ 760 770 780 790 800 NSKSNSVPII LAIVGLMLVT VVAGVLIVKK YVCGGRFLVH RYSVLQQHAE 810 820 830 ANGVDGVDAL DTASHTNKSG YHDDSDDLL E	
Mouse Sortilin	MERPRGAADG LLRWPLGLLL LLQLLPAAV GQDRLDAPPP PAPPLLRWAG PVGVSWSGLRA AAPGGPVPRA GRWRRGAPAE DQDCGRLPDF IAKLTNNTHQ HVFDDLSGSV SLSWVGDSGT VILVLTTFQV PLVIVSFGQS KLYRSEDYK NFKDITNLIN NTFIRTEFGM AIGPENSGKV ILTAEVSGGS RGGRVFRSSD FAKNFVQTDL PFHPLTQMMY SPQNSDYLLA LSTENGLWVS KNFGEKWEEI HKAVCLAKWG PNNIIFFTTH VNGSCKADLG ALELWRTSDL GKTFKTIGVK IYSFGLGGRF LFASVMADKD TTRRIHVSTD QGDTWSMAQL PSVGQEQFY ILAANEDMVF MHVDEPGDTG FGTIFTSDDR GIVYSKSLDR HLYTTTGGET DFTNVTSLRG VYITSTLSED NSIQSMITFD QGGRWEHLRK PENSKCDATA KNKNECSLHI HASYSISQKL NVPMAPLSEP NAVGIVIAHG SVGDAISVMV PDVYISDDGG YSWAKMLEGP HYYTILDSGG IIVAIEHSNR PINVIKFSTD EGQCWQSYVF TQEPYFTGL ASEPGARSMN ISIWGFTESE ITRQWVSYTV DFKDILERN EEDDYTTWLA HSTDPGDYKD GCILGYKEQF LRLRKSSVCQ NGRDYVVAQ PSVCPCSLD FLCDFGYFRP ENASECVEQP ELKGHELEFC LYGKEEHLTT NGYRKIPGDK CQGGMNPARE VKDLKKKCTS NFLNPTKQNS KSNSVPIILA IVGLMLVTVV AGVLIVKKYV CGGRFLVHRY SVLQQHAEAD GVEALDSTSH AKSGYHDDSD EDLLE	82
Rat Sortilin	MERPRGAADG LLRWPLGLLL LLQLLPAAV GQDRLDAPPP PAPPLLRWAG PVGVSWSGLRA AAPGGPVPRA GRWRRGAPAE DQDCGRLPDF IAKLTNNTHQ HVFDDLSGSV SLSWVGDSGT VILVLTTFQV PLVIVSFGQS KLYRSEDYK NFKDITNLIN NTFIRTEFGM AIGPENSGKV ILTAEVSGGS RGGRVFRSSD FAKNFVQTDL PFHPLTQMMY SPQNSDYLLA LSTENGLWVS KNFGEKWEEI HKAVCLAKWG PNNIIFFTTH VNGSCKADLG ALELWRTSDL GKTFKTIGVK IYSFGLGGRF LFASVMADKD TTRRIHVSTD QGDTWSMAQL PSVGQEQFY ILAANDDMVF MHVDEPGDTG FGTIFTSDDR GIVYSKSLDR HLYTTTGGET DFTNVTSLRG VYITSTLSED NSIQSMITFD QGGRWEHLQK PENSKCDATA KNKNECSLHI HASYSISQKL NVPMAPLSEP NAVGIVIAHG SVGDAISVMV PDVYISDDGG YSWAKMLEGP HYYTILDSGG IIVAIEHSNR PINVIKFSTD EGQCWQSYVF SQEPVYFTGL ASEPGARSMN ISIWGFTESE LTRQWVSYTI DFKDILERN EENDYTTWLA HSTDPGDYKD GCILGYKEQF LRLRKSSVCQ NGRDYVVAQ PSICPCSLD FLCDFGYFRP ENASECVEQP ELKGHELEFC LYGKEEHLTT NGYRKIPGDR CQGGMNPARE VKDLKKKCTS NFLNPKKQNS KSSSVPIILA IVGLMLVTVV AGVLIVKKYV CGGRFLVHRY SVLQQHAEAD GVEALDTASH AKSGYHDDSD EDLLE	83

Table 33: S-60-15 peptide sequences

Description	Sequence	SEQ ID NO
Asp-box motif	(S/T) -X- (D/N) -X-X-X-X- (W/F/Y) X is any amino acid	84
Asp-box motif	X-X- (S/T) -X- (D/N) -X-G-X- (T/S) - (W/F/Y) -X X is any amino acid	85
Asp-box motif in human Sortilin	SSDFAKNF	86

(residues 200-207)		
S-60-15 Peptide	NQFSLK	87
S-60-15 Peptide	QGYYGMDVWGQGTTVTVSSASTK	88
S-60-15 Peptide	DYFPEPVTVS WN SGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTK	89
S-60-15 Peptide	DTL M ISR	90
S-60-15 Peptide	TPEVTCVVVDVSQEDPEVQFNWYVDGVEVH NAK	91
S-60-15 Peptide	VVSVLTVLHQDWLNG K	92
S-60-15 Peptide	EPQVYTLPPSQEEM MTK	93
S-60-15 Peptide	NQVSLTCLVK	94
S-60-15 Peptide	GFYPSDIAVEWESN G QPENNYK	95
S-60-15 Peptide	WQEGNVFSCSV M HEALHNHYTQK	96
S-60-15 Peptide	DIV MT QSPLSLPVTGPGEPAISCR	97
S-60-15 Peptide	SSQ S LLR	98
S-60-15 Peptide	S N GYNYLDWYLQKPGQSPQLLIYLGSR	99
S-60-15 Peptide	AEAEDVGYYC M QQQEAPLTFGGGTK	100
S-60-15 Peptide	SGTASVVCLL N NFYPR	101
S-60-15 Peptide	VD N ALQSGNSQESVTEQDSK	102

Table 34: Fc domain amino acid sequences

Description	Sequence	SEQ ID NO
huIgG1 LALAPS Fc	ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKV DKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPE VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV VSVLTVLHQDWLNGKEYKCKVSNKALPASIEKTISKAKGQPREPQV YTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQK SLSLSPGK	103
huIgG1 LALAPS Fc without C- terminal lysine	ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKV DKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPE VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV VSVLTVLHQDWLNGKEYKCKVSNKALPASIEKTISKAKGQPREPQV YTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQK	104

Description	Sequence	SEQ ID NO
	SLSLSPG	

Table 35: Full-length heavy chain amino acid sequences

Description	Sequence	SEQ ID NO
S-60-10, S-60-11, S-60-12, S-60-19 with Fc LALAPS with C-terminal Lysine	QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGW IRQPPGKGLEWIGTIYHSGSTYYNPSLKSQVTISVDT KNQFSLELSSVTAADTAVYYCARQGSIQGGYYGMDV WGQGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAALG CLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLY SLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEP KSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTK PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPASIEKTISKAKGQPREPQVYTLPPSRDELTKNQV SLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD DGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNH YTQKSLSLSPGK	105
S-60-10, S-60-11, S-60-12, S-60-19 with Fc LALAPS without C-terminal Lysine	QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGW IRQPPGKGLEWIGTIYHSGSTYYNPSLKSQVTISVDT KNQFSLELSSVTAADTAVYYCARQGSIQGGYYGMDV WGQGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAALG CLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLY SLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEP KSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTK PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPASIEKTISKAKGQPREPQVYTLPPSRDELTKNQV SLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD DGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNH YTQKSLSLSPG	106
S-60-13, S-60-14 with Fc LALAPS with C-terminal Lysine	QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGW IRQPPGKGLEWIGTIYHSGSTYYNPSLESRTISVDTSK NQFSLKLSSVTAADTAVYYCARQGSIQGGYYGMDV WGQGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAALG CLVKDYFPEPTVSWNSGALTSGVHTFPAVLQSSGLY SLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEP KSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTK PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPASIEKTISKAKGQPREPQVYTLPPSRDELTKNQV SLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD DGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNH YTQKSLSLSPGK	135
S-60-13, S-60-14 with Fc LALAPS without C-terminal Lysine	QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWGW IRQPPGKGLEWIGTIYHSGSTYYNPSLESRTISVDTSK NQFSLKLSSVTAADTAVYYCARQGSIQGGYYGMDV WGQGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAALG	136

Description	Sequence	SEQ ID NO
	CLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY SLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEP KSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTK PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPASIEKTISKAKGQPREPQVYTLPPSRDELTKNQV SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD DGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNH YTQKSLSLSPG	
S-60, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16, S-60-18, S-60-24 <u>with Fc LALAPS with C-terminal Lysine</u>	QVQLQESGPGLVKPSETLSLTCAVSGYSISSGYYWGW IRQPPGKGLEWIGTIYHSGSTYYNPSLKSRTISVDT KNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMD VWGQGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAAL GCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGL YSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVE PKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMIS RTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVS NKALPASIEKTISKAKGQPREPQVYTLPPSRDELTKNQ VSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNH YTQKSLSLSPGK	137
S-60, S-60-15 [N33 (wt)], S-60-15.1 [N33T], S-60-15.2 [N33S], S-60-15.3 [N33G], S-60-15.4 [N33R], S-60-15.5 [N33D], S-60-15.6 [N33H], S-60-15.7 [N33K], S-60-15.8 [N33Q], S-60-15.9 [N33Y], S-60-15.10 [N33E], S-60-15.11 [N33W], S-60-15.12 [N33F], S-60-15.13 [N33I], S-60-15.14 [N33V], S-60-15.15 [N33A], S-60-15.16 [N33M], S-60-15.17 [N33L], S-60-16, S-60-18, S-60-24 <u>with Fc LALAPS without C-terminal Lysine</u>	QVQLQESGPGLVKPSETLSLTCAVSGYSISSGYYWGW IRQPPGKGLEWIGTIYHSGSTYYNPSLKSRTISVDT KNQFSLKLSSVTAADTAVYYCARQGSIKQGYGMD VWGQGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAAL GCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGL YSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVE PKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMIS RTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVS NKALPASIEKTISKAKGQPREPQVYTLPPSRDELTKNQ VSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNH YTQKSLSLSPG	138

Table 36: Full-length light chain amino acid sequences

Description	Sequence	SEQ ID NO
S-60-10; S-60-13; S-60-15 [N33 (wt)]	DIVMTQSPLSLPVTPGEPASIS CRSSQSLLRNGYNYLDWYL QKPGQSPQLLIYLGSRASGV	139

Description	Sequence	SEQ ID NO
	PDRFSGSGSGTDFTLKISRAEA EDVGVIYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRRGEC	
S-60-11; S-60-14	DIVMTQSPLSLPVTGPGEASIS CRSSQSLRSNGYNYLDWYL QKPGQSPQLLIYLGSRASGV PDRFSGSGSGTDFTLKISRVEA EDVGVIYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRRGEC	140
S-60-12	DIVMTQSPLSLPVTGPGEASIS CRSSQSLRSNGYNYLDWYL QKPGQSPQLLIYLGSRVSGV PDRFSGSGSGTDFTLKISRVEA EDVGVIYCMQQQETPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRRGEC	141
S-60-15.1 [N33T]	DIVMTQSPLSLPVTGPGEASIS CRSSQSLRSTGYNYLDWYL QKPGQSPQLLIYLGSRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVIYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRRGEC	142
S-60-15.2 [N33S]	DIVMTQSPLSLPVTGPGEASIS CRSSQSLRSGYNYLDWYL QKPGQSPQLLIYLGSRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVIYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRRGEC	143
S-60-15.3 [N33G]	DIVMTQSPLSLPVTGPGEASIS	144

Description	Sequence	SEQ ID NO
	CRSSQSLLRSGGNYLDWYL QKPGQSPQLLIYLGSRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRRGEC	
S-60-15.4 [N33R]	DIVMTQSPLSLPVTGPGEASIS CRSSQSLLRSGGNYLDWYL QKPGQSPQLLIYLGSRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRRGEC	145
S-60-15.5 [N33D]	DIVMTQSPLSLPVTGPGEASIS CRSSQSLLRSDGNYLDWYL QKPGQSPQLLIYLGSRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRRGEC	146
S-60-15.6 [N33H]	DIVMTQSPLSLPVTGPGEASIS CRSSQSLLRSHGNYLDWYL QKPGQSPQLLIYLGSRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRRGEC	119
S-60-15.7 [N33K]	DIVMTQSPLSLPVTGPGEASIS CRSSQSLLRSGGNYLDWYL QKPGQSPQLLIYLGSRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS	120

Description	Sequence	SEQ ID NO
	PVTKSFNRGEC	
S-60-15.8 [N33Q]	DIVMTQSPLSLPVTTPGEPASIS CRSSQSLLRSQGYNYLDWYL QKPGQSPQLLIYLGSNRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRGEC	121
S-60-15.9 [N33Y]	DIVMTQSPLSLPVTTPGEPASIS CRSSQSLLRSYGYNYLDWYL QKPGQSPQLLIYLGSNRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRGEC	122
S-60-15.10 [N33E]	DIVMTQSPLSLPVTTPGEPASIS CRSSQSLLRSEGYNYLDWYL QKPGQSPQLLIYLGSNRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRGEC	123
S-60-15.11 [N33W]	DIVMTQSPLSLPVTTPGEPASIS CRSSQSLLRSWGYNYLDWYL QKPGQSPQLLIYLGSNRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLSK ADYEKHKVYACEVTHQGLSS PVTKSFNRGEC	124
S-60-15.12 [N33F]	DIVMTQSPLSLPVTTPGEPASIS CRSSQSLLRSFGYNYLDWYL QKPGQSPQLLIYLGSNRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES	125

Description	Sequence	SEQ ID NO
	VTEQDSKDYSLSTLTLTK ADYEKHKVYACEVTHQGLSS PVTKSFNREGC	
S-60-15.13 [N33I]	DIVMTQSPLSLPVTGPGEASIS CRSSQSLLRSIGYNYLDWYLQ KPGQSPQLLIYLGSNRASGVP DRFSGSGSGTDFTLKISRAEAE DVGVIYCMQQQEAPLTFGGG TKVEIKRTVAAPSVFIFPPSDE QLKSGTASVVCLLNNFYPREA KVQWKVDNALQSGNSQESVT EQDSKDYSLSTLTLTKAD YEKHKVYACEVTHQGLSSPV TKSFNREGC	126
S-60-15.14 [N33V]	DIVMTQSPLSLPVTGPGEASIS CRSSQSLLRSVGYNYLDWYL QKPGQSPQLLIYLGSNRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVIYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLTK ADYEKHKVYACEVTHQGLSS PVTKSFNREGC	127
S-60-15.15 [N33A]	DIVMTQSPLSLPVTGPGEASIS CRSSQSLLRSAGYNYLDWYL QKPGQSPQLLIYLGSNRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVIYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLTK ADYEKHKVYACEVTHQGLSS PVTKSFNREGC	128
S-60-15.16 [N33M]	DIVMTQSPLSLPVTGPGEASIS CRSSQSLLRSMGYNYLDWYL QKPGQSPQLLIYLGSNRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVIYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLTK ADYEKHKVYACEVTHQGLSS PVTKSFNREGC	129
S-60-15.17 [N33L]	DIVMTQSPLSLPVTGPGEASIS CRSSQSLLRSLGYNYLDWYL QKPGQSPQLLIYLGSNRASGV PDRFSGSGSGTDFTLKISRAEA EDVGVIYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD	130

Description	Sequence	SEQ ID NO
	EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLTK ADYEKHKVYACEVTHQGLSS PVTKSFNRGEC	
S-60-16	DIVMTQSPLSLPVTGPGEASIS CRSSQSLLRSNGYNYLDWYL QKPGQSPQLLIYLGSRASGV PDRFSGSGSGTDFTLKISRVEA EDVGAYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLTK ADYEKHKVYACEVTHQGLSS PVTKSFNRGEC	131
S-60-18	DIVMTQSPLSLPVTGGPASIS CRSSQSLLRSNGYNYLDWYL QKPGQSPQLLIYLGSRASGV PDRLSGSGSGTDFTLKISRVEA EDVGYYCMQQQETPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLTK ADYEKHKVYACEVTHQGLSS PVTKSFNRGEC	132
S-60, S-60-19	DIVMTQSPLSLPVTGPGEASIS CRSSQSLLHSNGYNYLDWYL QKPGQSPQLLIYLGSRASGV PDRFSGSGSGTDFTLKISRVEA EDVGYYCMQQQETPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLTK ADYEKHKVYACEVTHQGLSS PVTKSFNRGEC	133
S-60-24	DIVMTQSPLSLPVTGPGEASIS CRSSQGLLRNGYNYLDWYL QKPGQSPQLLIYLGSRASGV PDRFSGSGSGTDFTLKISRVEA EDVGYYCMQQQEAPLTFGG GTKVEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQES VTEQDSKDYSLSTLTLTK ADYEKHKVYACEVTHQGLSS PVTKSFNRGEC	134

CLAIMS:

1. An antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein:
 - (a) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYW (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), and an HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSNGYNYLD (SEQ ID NO: 8), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32); or
 - (b) the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISSGYW (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), and an HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSTGYNYLD (SEQ ID NO: 9), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).
2. The antibody of claim 1, wherein the antibody:
 - (a) decreases cell surface levels of Sortilin more than the level of decrease caused by an anti-Sortilin antibody comprising a heavy chain variable region comprising the sequence of SEQ ID NO: 56 and a light chain variable region comprising the sequence of SEQ ID NO: 79;
 - (b) increases extracellular levels of Progranulin more than the level of increase caused by an anti-Sortilin antibody comprising a heavy chain variable region comprising the sequence of SEQ ID NO: 56 and a light chain variable region comprising the sequence of SEQ ID NO: 79;
 - (c) inhibits the interaction between Sortilin expressed on a cell and Progranulin more than the level of inhibition caused by an anti-Sortilin antibody

comprising a heavy chain variable region comprising the sequence of SEQ ID NO: 56 and a light chain variable region comprising the sequence of SEQ ID NO: 79; or

(d) any combination of (a)–(c).

3. The antibody of claim 1 or claim 2, wherein the antibody has a dissociation constant (K_D) for human Sortilin that is at least 1.1-fold lower than an anti-Sortilin antibody comprising a heavy chain variable region comprising the sequence of SEQ ID NO: 56 and a light chain variable region comprising the sequence of SEQ ID NO: 79, wherein the K_D is determined by FACS.
4. The antibody of any one of claims 1-3, wherein the antibody has a dissociation constant (K_D) for human Sortilin that ranges from 5.0E-10 M to 1.0E-9 M wherein the K_D is determined by FACS, or 250-500 pM wherein the K_D is determined by Bio-layer interferometry.
5. An antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein: the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISGGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), and an HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence RSSQSLLRSTGYNLYD (SEQ ID NO: 9), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).
6. An antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein: the heavy chain variable region comprises an HVR-H1 comprising the amino acid sequence YSISGGYYWG (SEQ ID NO: 1), an HVR-H2 comprising the amino acid sequence TIYHSGSTYYNPSLKS (SEQ ID NO: 2), and an HVR-H3 comprising the amino acid sequence ARQGSIKQGYGMDV (SEQ ID NO: 6); and the light chain variable region comprises an HVR-L1 comprising the amino acid sequence

RSSQSLLRSNGYNYLD (SEQ ID NO: 8), an HVR-L2 comprising the amino acid sequence LGSNRAS (SEQ ID NO: 29), and an HVR-L3 comprising the amino acid sequence MQQQEAPLT (SEQ ID NO: 32).

7. An antibody that binds to a Sortilin protein, wherein the antibody comprises:
 - (a) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57; or
 - (b) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 60.
8. An antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 60.
9. An antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 56 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 57.
10. An antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 137 and a light chain comprising the amino acid sequence of SEQ ID NO: 142.
11. An antibody that binds to a Sortilin protein, wherein the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 138 and a light chain comprising the amino acid sequence of SEQ ID NO: 142.
12. The antibody of any one of claims 1-9, wherein the antibody is of the IgG class, the IgM class, or the IgA class.
13. The antibody of claim 12, wherein the antibody is of the IgG class and has an IgG1, IgG2, IgG3, or IgG4 isotype.
14. The antibody of claim 13, wherein:
 - (a) the antibody is an IgG1 or IgG2 isotype and the Fc region comprises an amino acid substitution at position P331S, wherein the numbering of the residue position is according to EU numbering;

(b) the antibody is an IgG1 isotype and the Fc region comprises amino acid substitutions at positions L234A, L235A, and P331S, wherein the numbering of the residue position is according to EU numbering;

(c) the antibody is an IgG1, IgG2, or IgG4 isotype and the Fc region comprises an amino acid substitution at position N297A, wherein the numbering of the residue position is according to EU numbering; or

(d) the Fc region comprises an amino acid substitution at positions S267E and L328F, wherein the numbering of the residue position is according to EU numbering.

15. The antibody of claim 14, wherein the antibody is an IgG1 isotype and the Fc region comprises amino acid substitutions at positions L234A, L235A, and P331S, wherein the numbering of the residue position is according to EU numbering.
16. The antibody of any one of claims 1-15, wherein the Sortilin protein is a human Sortilin protein.
17. The antibody of claim 16, wherein the human Sortilin protein comprises the amino acid sequence of SEQ ID NO: 81.
18. The antibody of any one of claims 1-16, wherein the Sortilin protein is a wild-type protein, a naturally occurring variant, or a disease variant.
19. The antibody of any one of claims 1-18, wherein the antibody is a human antibody, a bispecific antibody, a monoclonal antibody, a multivalent antibody, or a conjugated antibody.
20. The antibody of claim 19, wherein the antibody is a bispecific antibody recognizing a first antigen and a second antigen.
21. The antibody of claim 20, wherein the first antigen is Sortilin and the second antigen is an antigen facilitating transport across the blood-brain-barrier.
22. The antibody of claim 21, wherein the second antigen is selected from the group consisting of transferrin receptor (TR), insulin receptor (HIR), insulin-like growth factor receptor (IGFR), low-density lipoprotein receptor related proteins 1 and 2 (LPR-1 and 2), diphtheria toxin receptor, CRM197, a llama single domain antibody,

TMEM 30(A), a protein transduction domain, TAT, Syn-B, penetratin, a poly-arginine peptide, an angiopep peptide, basigin, Glut1, CD98hc, and ANG1005.

23. The antibody of any one of claims 1-9 and 12-22, wherein the antibody is an antibody fragment that binds to a human Sortilin protein.
24. The antibody of claim 23, wherein the antibody fragment is an Fab, Fab', Fab'-SH, F(ab')₂, Fv, or scFv fragment.
25. An isolated nucleic acid comprising a nucleic acid sequence encoding the antibody of any one of claims 1-24.
26. A vector comprising the nucleic acid of claim 25.
27. An isolated host cell comprising the vector of claim 26.
28. A method of producing an antibody that binds to Sortilin, comprising culturing the host cell of claim 27 so that the antibody is produced.
29. The method of claim 28, further comprising recovering the antibody produced by the host cell.
30. An isolated antibody that binds to Sortilin produced by the method of claim 28 or claim 29.
31. A pharmaceutical composition comprising the antibody of any one of claims 1-24 and claim 30 and a pharmaceutically acceptable carrier.
32. Use of the antibody of any one of claims 1-24 and claim 30 or the pharmaceutical composition of claim 31 in the manufacture of a medicament for preventing, reducing risk for, or treating a disease, disorder, or injury in an individual, wherein the disease, disorder, or injury is selected from the group consisting of frontotemporal dementia, Alzheimer's disease, vascular dementia, seizures, retinal dystrophy, amyotrophic lateral sclerosis, traumatic brain injury, a spinal cord injury, dementia, stroke, Parkinson's disease, retinal degeneration, arthritis, and osteoarthritis.
33. The use of claim 32, wherein the disease, disorder, or injury is selected from the group consisting of frontotemporal dementia, Parkinson's disease, Alzheimer's disease, and amyotrophic lateral sclerosis.

34. The antibody of any one of claims 1-24 and claim 30 or the pharmaceutical composition of claim 31 for use in inhibiting one or more of neuroinflammation, axonopathy characterized by short axonal outgrowth and aberrant branching, aberrant microglial activation, and inflammatory response in an individual in need thereof.
35. The antibody of any one of claims 1-24 and claim 30 or the pharmaceutical composition of claim 31 for use in promoting one or more of wound healing, autophagy, and clearance of aggregate proteins in an individual in need thereof.
36. Use of the antibody of any one of claims 1-24 and claim 30 or the pharmaceutical composition of claim 31 in the manufacture of a medicament for preventing, reducing risk of, or treating arthritis in an individual in need thereof.
37. The antibody of any one of claims 1-24 and claim 30 or the pharmaceutical composition of claim 31 for use in decreasing expression of one or more pro-inflammatory mediators in an individual in need thereof.
38. The antibody or pharmaceutical composition of claim 37, wherein the one or more pro-inflammatory mediators are selected from the group consisting of IL-6, IL12p70, IL12p40, IL-1 β , TNF- α , CXCL1, CCL2, CCL3, CCL4, and CCL5.
39. A method of inhibiting the interaction between Sortilin expressed on a cell and Progranulin, the method comprising exposing a cell expressing Sortilin on the cell surface to the antibody of any one of claims 1-24 and claim 30 or the pharmaceutical composition of claim 31, wherein the cell is *in vitro*.
40. The method of claim 39, further comprising decreasing the level of Sortilin expressed on the cell surface.
41. The method of claim 39 or claim 40, wherein extracellular levels of Progranulin are increased.
42. The antibody of any one of claims 1-24 and claim 30 or the pharmaceutical composition of claim 31 for use in increasing levels of Progranulin in an individual in need thereof.
43. The antibody or pharmaceutical composition of claim 42, wherein levels of Progranulin are increased in plasma.

44. The antibody or pharmaceutical composition of claim 42, wherein levels of Progranulin are increased in cerebrospinal fluid.
45. The antibody of any one of claims 1-24 and claim 30 or the pharmaceutical composition of claim 31 for use in decreasing levels of Sortilin in an individual in need thereof.
46. The antibody or pharmaceutical composition of claim 45, wherein the levels of Sortilin are decreased in peripheral white blood cells.
47. The use of any one of claims 32-33 and 36, or the antibody or pharmaceutical composition of any one of claims 34-35, 37-38 and 42-46, wherein the individual has one or more mutations in the gene encoding Progranulin.
48. The use, antibody, or pharmaceutical composition of claim 47, wherein the individual is heterozygous for one or more loss-of-function mutations in the gene encoding Progranulin.
49. The use, antibody, or pharmaceutical composition of claim 47 or claim 48, wherein the individual has or is at risk for frontotemporal dementia, Alzheimer's disease, or amyotrophic lateral sclerosis.
50. The use of any one of claims 32-33, 36, and 47-49, the antibody or pharmaceutical composition of any one of claims 34-35, 37-38 and 42-49, or the method of any one of claims 39-41, wherein the antibody comprises two or more antibodies that bind to a Sortilin protein.