

SEQUENCE LISTING

<110> TAKEDA PHARMACEUTICAL COMPANY LIMITED

<120> Cationic Compounds

<130> PT38-9027W0

<150> JP 2017-254667

<151> 2017-12-28

<160> 4

<170> PatentIn version 3.5

<210> 1

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Colla1 siRNA (Sense)

<220>

<221> misc_feature

<222> (1)..(21)

<223> The sequence comprises both RNA and DNA bases, some of which are modified bases.

<220>

<221> modified_base

<222> (2)..(2)

<223> The "u" stands for 2'-O-methyluridine (um)

<220>

<221> modified_base

<222> (3)..(3)

<223> The "c" stands for 2'-O-methylcytidine (cm)

<220>

<221> modified_base

<222> (4)..(4)

<223> The "u" stands for 2'-O-methyluridine (um)

<220>

<221> modified_base

<222> (8)..(8)

<223> The "c" stands for 2'-O-methylcytidine (cm)

<220>
 <221> modified_base
 <222> (10)..(10)
 <223> The "u" stands for 2'-O-methyluridine (um)

 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> The "u" stands for 2'-O-methyluridine (um)

 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> The "c" stands for 2'-O-methylcytidine (cm)

 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> The "u" stands for 2'-O-methyluridine (um)

 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> The "c" stands for 2'-O-methylcytidine (cm)

 <220>
 <221> modified_base
 <222> (17)..(17)
 <223> The "c" stands for 2'-O-methylcytidine (cm)

 <220>
 <221> modified_base
 <222> (18)..(18)
 <223> The "u" stands for 2'-O-methyluridine (um)

 <220>
 <221> misc_feature
 <222> (20)..(21)
 <223> The nucleosides (t and t) are bound with phosphorothioate.

<400> 1
 gucuagacau guucagcuut t

21

<210> 2
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
<223> Col1a1 siRNA (Antisense)

<220>
<221> misc_feature
<222> (1)..(21)
<223> The sequence comprises both RNA and DNA bases, some of which are modified bases.

<220>
<221> modified_base
<222> (9)..(9)
<223> The "c" stands for 2'-O-methylcytidine (cm)

<220>
<221> modified_base
<222> (15)..(15)
<223> The "u" stands for 2'-O-methyluridine (um)

<220>
<221> misc_feature
<222> (20)..(21)
<223> The nucleosides (t and t) are bound with phosphorothioate.

<400> 2
aagcugaaca ugucuagact t

21

<210> 3
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Factor VII siRNA (Sense)

<220>
<221> misc_feature
<222> (1)..(21)
<223> The sequence comprises both RNA and DNA bases, some of which are modified bases.

<220>
<221> modified_base
<222> (1)..(1)
<223> The "c" stands for 2'-O-methylcytidine (cm)

<220>
 <221> modified_base
 <222> (2)..(2)
 <223> The "u" stands for 2'-O-methyluridine (um)

<220>
 <221> modified_base
 <222> (4)..(4)
 <223> The "c" stands for 2'-O-methylcytidine (cm)

<220>
 <221> modified_base
 <222> (10)..(10)
 <223> The "c" stands for 2'-O-methylcytidine (cm)

<220>
 <221> modified_base
 <222> (12)..(12)
 <223> The "u" stands for 2'-O-methyluridine (um)

<220>
 <221> modified_base
 <222> (13)..(13)
 <223> The "c" stands for 2'-O-methylcytidine (cm)

<220>
 <221> modified_base
 <222> (14)..(14)
 <223> The "c" stands for 2'-O-methylcytidine (cm)

<220>
 <221> modified_base
 <222> (15)..(15)
 <223> The "u" stands for 2'-O-methyluridine (um)

<220>
 <221> modified_base
 <222> (16)..(16)
 <223> The "u" stands for 2'-O-methyluridine (um)

<220>
 <221> modified_base
 <222> (17)..(17)
 <223> The "c" stands for 2'-O-methylcytidine (cm)

<220>
 <221> modified_base
 <222> (19)..(19)

<223> The "u" stands for 2'-O-methyluridine (um)

<220>

<221> misc_feature

<222> (20)..(21)

<223> The nucleosides (t and t) are bound with phosphorothioate.

<400> 3

cuacgaaagc auccuucagt t

21

<210> 4

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Factor VII siRNA (Antisense)

<220>

<221> misc_feature

<222> (1)..(21)

<223> The sequence comprises both RNA and DNA bases, some of which are modified bases.

<220>

<221> modified_base

<222> (17)..(17)

<223> The "u" stands for 2'-O-methyluridine (um)

<220>

<221> misc_feature

<222> (20)..(21)

<223> The nucleosides (t and t) are bound with phosphorothioate.

<400> 4

augaaggaug cuuucguagt t

21