
SUMMARY

[0007] Provided herein are compounds and compositions that selectively inhibit Bcr-Abl tyrosine kinases and that are useful for treating disorders mediated by Bcr-Abl tyrosine kinases.

[0007a] In an aspect of the present disclosure, there is provided a compound of formula (I-A):



(I-A)

or a pharmaceutically acceptable salt thereof, wherein:

X is $NR^{3'}$ or CR^3,

Y is NR^2 or CR^4,

wherein when X is $NR^{3'}$ then Y is CR^4, Y has a double bond to CR^5, and X has a single bond to CR^5; or when X is CR^3 then Y is NR^2, Y has a single bond to CR^5, and X has a double bond to CR^5;

m is an integer from 0 to 3;

each $R^1$is independently -D, -F, C_1-C_3 alkyl, C_1-C_3 alkylene-NR^7R^8, C_1-C_3 alkylene-$NR^{7'}R^{8'}$, C_1-C_3 alkylene-OH, C_1-C_3 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-OH, C_1-C_2 alkylene-(4- to 8-membered heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(4- to 8-membered heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_7 heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, or C_1-C_2 alkylene-(C_3-C_7 heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, wherein the alkyl, alkylene, cycloalkylene, and heterocycloalkylene moieties in R^1

are optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

R^2 is -H, C_1-C_3 alkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^3 is -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, halogen, or -CN;

$R^{3'}$ is -H, C_1-C_3 alkyl, -C_3-C_6 cycloalkyl, or -CN;

R^4 is -H, C_1-C_3 alkyl, or halogen, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^5 is 5-to-10-membered heteroaryl, wherein said 5-to-10-membered heteroaryl is optionally substituted with 1-5 R^9 groups;

each R^7 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;

each R^8 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;


each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 3- to 8-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each $R^9$is independently halogen, -OR^{10}, -NR^7R^8, C_1-C_3 alkyl, C_1-C_3 haloalkyl, C_3-C_6 cycloalkyl, -CN, $S(O)_nC_1$-C_3 alkyl, or $S(O)_nC_3$-C_6 cycloalkyl,

wherein n is an integer from 0 to 2; and

each R^{10} is independently -H, C_1-C_3 alkyl, C_1-C_3 haloalkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with hydroxyl, C_1-C_3 alkoxy, and/or 1-6 deuterium atoms.

[0007b] In another aspect of the present disclosure, there is provided a compound of formula (I-A'):



(I-A')

or a pharmaceutically acceptable salt thereof, wherein:

X is $NR^{3'}$ or CR^3,

Y is NR^2 or CR^4,

wherein when X is $NR^{3'}$ then Y is CR^4, Y has a double bond to CR^5, and X has a single bond to CR^5; or when X is CR^3 then Y is NR^2, Y has a single bond to CR^5, and X has a double bond to CR^5;

m is an integer from 0 to 3;

each R^1 is independently -D, -F, C_1-C_3 alkyl, C_1-C_3 alkylene-NR^7R^8, C_1-C_3 alkylene-$NR^{7'}R^{8'}$, C_1-C_3 alkylene-OH, C_1-C_3 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-OH, C_1-C_2 alkylene-(C_4-C_6 heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, or C_1-C_2 alkylene-(C_4-C_6 heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, wherein the alkyl, alkylene, cycloalkylene, and heterocycloalkylene moieties in R^1 are optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

R^2 is -H, C_1-C_3 alkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^3 is -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, halogen, or -CN;

$R^{3'}$ is -H, C_1-C_3 alkyl, -C_3-C_6 cycloalkyl, or -CN;

R^4 is -H, C_1-C_3 alkyl, or halogen, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^5 is 5-to-10-membered heteroaryl, wherein said said 5-to-10-membered heteroaryl is optionally substituted with 1-5 R^9 groups;

each R^7 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;

each R^8 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;


each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 4-to-6-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^9 is independently halogen, $-OR^{10}$, $-NR^7R^8$, C_1-C_3 alkyl, C_1-C_3 haloalkyl, C_3-C_6 cycloalkyl, -CN, $S(O)_nC_1$-C_3 alkyl, or $S(O)_nC_3$-C_6 cycloalkyl,

wherein n is an integer from 0 to 2; and

each R^{10} is independently -H, C_1-C_3 alkyl, C_1-C_3 haloalkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with hydroxyl, C_1-C_3 alkoxy, and/or 1-6 deuterium atoms.

[0007c] In another aspect of the present disclosure, there is provided a compound of formula (I-B):



(I-B)

or a pharmaceutically acceptable salt thereof, wherein:

X is $NR^{3'}$ or CR^3,

Y is NR^2 or CR^4,

wherein when X is $NR^{3'}$ then Y is CR^4, Y has a double bond to CR^5, and X has a single bond to CR^5; or when X is CR^3 then Y is NR^2, Y has a single bond to CR^5, and X has a double bond to CR^5;

m is an integer from 0 to 3;

each R^1 is independently -D, -F, C_1-C_3 alkyl, C_1-C_3 alkylene-NR^7R^8, C_1-C_3 alkylene-$NR^{7'}R^{8'}$, C_1-C_3 alkylene-OH, C_1-C_3 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-OH, C_1-C_2 alkylene-(4- to 8-membered heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(4- to 8-membered heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_7 heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, or C_1-C_2 alkylene-(C_3-C_7 heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, wherein the alkyl, alkylene, cycloalkylene, and heterocycloalkylene moieties in R^1 are optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

R^2 is -H, C_1-C_3 alkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^3 is -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, halogen, or -CN;

$R^{3'}$ is -H, C_1-C_3 alkyl, -C_3-C_6 cycloalkyl, or -CN;

R^4 is -H, C_1-C_3 alkyl, or halogen, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^5 is C_6-C_{14} aryl or 5-to-10-membered heteroaryl, wherein said C_6-C_{14} aryl or said 5-to-10-membered heteroaryl is optionally substituted with 1-5 R^9 groups;

each R^7 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;

each R^8 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;

each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 3- to 8-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally


substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^9 is independently halogen, -OR^{10}, -NR^7R^8, C_1-C_3 alkyl, C_1-C_3 haloalkyl, C_3-C_6 cycloalkyl, -CN, $S(O)_nC_1$-C_3 alkyl, or $S(O)_nC_3$-C_6 cycloalkyl,

wherein n is an integer from 0 to 2; and

each R^{10} is independently -H, C_1-C_3 alkyl, C_1-C_3 haloalkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with hydroxyl, C_1-C_3 alkoxy, and/or 1-6 deuterium atoms.

[0007d] In another aspect of the present disclosure, there is provided a compound of formula (I-B'):



(I-B')

or a pharmaceutically acceptable salt thereof, wherein:

X is $NR^{3'}$ or CR^3,

Y is NR^2 or CR^4,

wherein when X is $NR^{3'}$ then Y is CR^4, Y has a double bond to CR^5, and X has a single bond to CR^5; or when X is CR^3 then Y is NR^2, Y has a single bond to CR^5, and X has a double bond to CR^5;

m is an integer from 0 to 3;

R^2 is -H, C_1-C_3 alkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^3 is -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, halogen, or -CN;

$R^{3'}$ is -H, C_1-C_3 alkyl, -C_3-C_6 cycloalkyl, or -CN;

R^4 is -H, C_1-C_3 alkyl, or halogen, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^5 is C_6-C_{14} aryl or 5-to-10-membered heteroaryl, wherein said C_6-C_{14} aryl or said 5-to-10-membered heteroaryl is optionally substituted with 1-5 R^9 groups;

R^6 is -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 alkylene-NR^7R^8, C_1-C_6 alkylene-$NR^{7'}R^{8'}$, C_1-C_6 alkylene-OH, C_1-C_6 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-OH, C_1-C_2 alkylene-(C_4-C_6 heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, or C_1-C_2 alkylene-(C_4-C_6 heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, wherein the alkyl, alkylene, cycloalkyl, cycloalkylene, and heterocycloalkylene moieties in R^6 are optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^7 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;

each R^8 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;


each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 4-to-6-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;


each R^9 is independently halogen, $-OR^{10}$, $-NR^7R^8$, C_1-C_3 alkyl, C_1-C_3 haloalkyl, C_3-C_6 cycloalkyl, -CN, $S(O)_nC_1$-C_3 alkyl, or $S(O)_nC_3$-C_6 cycloalkyl,

wherein n is an integer from 0 to 2; and


each R^{10} is independently -H, C_1-C_3 alkyl, C_1-C_3 haloalkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with hydroxyl, C_1-C_3 alkoxy, and/or 1-6 deuterium atoms.


[0007e] In a further aspect of the present disclosure, there is provided a compound selected from the group consisting of:





;




;







;




;




;

;




;

;














;

;


;

;


;

;




;

;




;

;




;

;

;

;

;

;

;

;

;

;

;

;



;



;




F
O
N
H
N
N
NH
;

F
O
N
H
N
N
O
CD3
;














F
O
N
H
N
N
O
;

F
O
N
H
N
N
O
;


F
O
N
H
N
O
F
N
O
;

F
O
N
H
N
O
N
N
O
;


N
O
N
H
N
O
;

O
N
H
N
N
O
;




O
N
H
N
F
;

F
O
N
H
N
O
;




F
O
N
H
N
F
O
;

O
N
H
N
F
;




;

;

;

;

;

;

;

;

;

;

;

;



;

;


;

;




;

;




;

;




;

;



;



;




;

;




;

;




;

;


;

;


;

;










;

;


;

;


;

;

;

,














;

;

;

;

;

;

;

;

;

;











;










;


;

;

;

;

;

;

;

;

;

;

;

;




;



;




;

;

;

;

;

;

;

;

;

;


;

;

;

;

;

;

;

;

;



;

;

;

;

;


;

;




;

;



and




;

,



or a pharmaceutically acceptable salt thereof.



[0007d] In another aspect of the present disclosure, there is provided a compound selected from the group consisting of:




;

;




;

;




O
N
N
H
O
O
;

O
N
N
H
NC
;




OH
O
N
N
H
;

O
N
N
H
OH
;














O—
O
N
N
H
;

O
N
N
H
O
;


O
N
N
H
—O
;

O
N
N
H
F
;


O
N
N
H
NH
;

O
N
N
H
NH
;














O
N
N
H
NH
;

O
N
N
H
NH
N
;














O
N
N
H
NH
N
;

O
N
N
H
N
NH
;


;

;

;

;

;

;

;

;

;

;

;

;

;












;


;

;

;

;

;

;

;

;

;

;

;

;



;

;

;

;

;

;

;

;

;

;

;

;











;

;


;

;

;

;

;

;

;

;

;

;

;

;

;

;











;

;

;

;

;

;

;

;

;

;

;


;











;

;


;

;

;

;

;

,

;

;

;













;

;

;


;

;

;

;

;

;

;

;

;

;











;











;


;

;

;

;

;

;

;

;

;

;

;

;



;













;


D
D
D
CD3O
O
N
N
F
N
H
N
O
D
D
D
;

O
N
N
F
(S)
(S)
N
H
N
;

O
N
N
F
(S)
(S)
N
H
N
O
;

O
N
N
F
(S)
(S)
N
H
N
O
;

O
N
N
F
(S)
(S)
N
H
N
O
;

O
O
N
N
F
(S)
(S)
N
H
N
;

O
O
N
N
F
(S)
(S)
N
H
N
;

O
O
N
N
N
N
O
;

O
O
N
N
F
(S)
(S)
N
H
N
O
;

O
N
N
O
N
N
O
;


;

;

;

;

;

;

;

;

;

;


;


;

;

;


;

;

;

;

;

;

;

;

;

;

;

;

;


;



;


;


and



,

or a pharmaceutically acceptable salt thereof.


[0007e] In a further aspect of the present disclosure, there is provided a compound having the structure



,

or a pharmaceutically acceptable salt thereof.


[0007f] In another aspect of the present disclosure, there is provided a compound having the structure



,

or a pharmaceutically acceptable salt thereof.

[0007g] In a further aspect of the present disclosure, there is provided a compound having the structure



or a pharmaceutically acceptable salt thereof.


[0007h] In another aspect of the present disclosure, there is provided a pharmaceutical composition comprising the compound disclosed herein, or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable excipients.

[0007i] In a further aspect of the present disclosure, there is provided a use of a compound disclosed herein, or a pharmaceutically acceptable salt thereof, or the pharmaceutical composition disclosed herein, in the manufacture of a medicament for inhibiting Bcr-Abl enzymatic activity in a cell.

[0007j] In another aspect of the present disclosure, there is provided a use of a compound disclosed herein, or a pharmaceutically acceptable salt thereof, or the pharmaceutical composition disclosed herein, in the manufacture of a medicament for treating chronic myeloid leukemia (CML), acute myeloid leukemia (AML), acute lymphoblastic leukemia (ALL), or a mixed phenotype acute leukemia, in a human in need thereof.

[0007k] In a further aspect of the present disclosure, there is provided a compound disclosed herein, or a pharmaceutically acceptable salt thereof, or the pharmaceutical composition disclosed herein, for use in therapy.

[0008] In one example, provided herein is a compound of formula (I),





(I)

or a pharmaceutically acceptable salt, solvate, hydrate, or co-crystal thereof, or a mixture of any of the foregoing, wherein:

X is $NR^{3'}$ or CR^3,

Y is NR^2 or CR^4,

wherein when X is $NR^{3'}$ then Y is CR^4, Y has a double bond to CR^5, and X has a single bond to CR^5; or when X is CR^3 then Y is NR^2, Y has a single bond to CR^5, and X has a double bond to CR^5;

R^0 is a group  or ;


m is an integer from 0 to 3;

each R^1 is independently -D, -F, C_1-C_3 alkyl, C_1-C_3 alkylene-NR^7R^8, C_1-C_3 alkylene-$NR^{7'}R^{8'}$, C_1-C_3 alkylene-OH, C_1-C_3 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-OH, C_1-C_2 alkylene-(4- to 8-membered heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(4- to 8-membered heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_7 heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, or C_1-C_2 alkylene-(C_3-C_7 heterocycloalkylene)-(C_0-C_2 alkylene)-

CLAIMS

What is claimed:

Claim 1. A compound of formula (I-A):



(I-A)

or a pharmaceutically acceptable salt thereof, wherein:

X is $NR^{3'}$ or CR^3,

Y is NR^2 or CR^4,

wherein when X is $NR^{3'}$ then Y is CR^4, Y has a double bond to CR^5, and X has a single bond to CR^5; or when X is CR^3 then Y is NR^2, Y has a single bond to CR^5, and X has a double bond to CR^5;

m is an integer from 0 to 3;

each R^1 is independently -D, -F, C_1-C_3 alkyl, C_1-C_3 alkylene-NR^7R^8, C_1-C_3 alkylene-$NR^{7'}R^{8'}$, C_1-C_3 alkylene-OH, C_1-C_3 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-OH, C_1-C_2 alkylene-(4- to 8-membered heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(4- to 8-membered heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_7 heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, or C_1-C_2 alkylene-(C_3-C_7 heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, wherein the alkyl, alkylene, cycloalkylene, and heterocycloalkylene moieties in R^1 are optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

R^2 is -H, C_1-C_3 alkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^3 is -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, halogen, or -CN;

$R^{3'}$ is -H, C_1-C_3 alkyl, -C_3-C_6 cycloalkyl, or -CN;

R^4 is -H, C_1-C_3 alkyl, or halogen, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^5 is 5-to-10-membered heteroaryl, wherein said 5-to-10-membered heteroaryl is optionally substituted with 1-5 R^9 groups;

each R^7 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;

each R^8 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;


each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 3- to 8-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^9 is independently halogen, -OR^{10}, -NR^7R^8, C_1-C_3 alkyl, C_1-C_3 haloalkyl, C_3-C_6 cycloalkyl, -CN, $S(O)_nC_1$-C_3 alkyl, or $S(O)_nC_3$-C_6 cycloalkyl,

wherein n is an integer from 0 to 2; and

each R^{10} is independently -H, C_1-C_3 alkyl, C_1-C_3 haloalkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with hydroxyl, C_1-C_3 alkoxy, and/or 1-6 deuterium atoms.

Claim 2. A compound of formula (I-A'):



(I-A')

or a pharmaceutically acceptable salt thereof, wherein:

X is $NR^{3'}$ or CR^3,

Y is NR^2 or CR^4,

wherein when X is $NR^{3'}$ then Y is CR^4, Y has a double bond to CR^5, and X has a single bond to CR^5; or when X is CR^3 then Y is NR^2, Y has a single bond to CR^5, and X has a double bond to CR^5;

m is an integer from 0 to 3;

each R^1 is independently -D, -F, C_1-C_3 alkyl, C_1-C_3 alkylene-NR^7R^8, C_1-C_3 alkylene-$NR^{7'}R^{8'}$, C_1-C_3 alkylene-OH, C_1-C_3 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-OH, C_1-C_2 alkylene-(C_4-C_6 heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, or C_1-C_2 alkylene-(C_4-C_6 heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, wherein the alkyl, alkylene, cycloalkylene, and heterocycloalkylene moieties in R^1 are optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

R^2 is -H, C_1-C_3 alkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^3 is -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, halogen, or -CN;

$R^{3'}$ is -H, C_1-C_3 alkyl, -C_3-C_6 cycloalkyl, or -CN;

R^4 is -H, C_1-C_3 alkyl, or halogen, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^5 is 5-to-10-membered heteroaryl, wherein said 5-to-10-membered heteroaryl is optionally substituted with 1-5 R^9 groups;

each R^7 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;

each R^8 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;


each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 4-to-6-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^9 is independently halogen, -OR^{10}, -NR^7R^8, C_1-C_3 alkyl, C_1-C_3 haloalkyl, C_3-C_6 cycloalkyl, -CN, $S(O)_nC_1$-C_3 alkyl, or $S(O)_nC_3$-C_6 cycloalkyl,


wherein n is an integer from 0 to 2; and

each R^{10} is independently -H, C_1-C_3 alkyl, C_1-C_3 haloalkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with hydroxyl, C_1-C_3 alkoxy, and/or 1-6 deuterium atoms.

Claim 3. The compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of formula (I) is a compound of formula (I-A-i) or formula (I-A-ii):



(I-A-i), or



(I-A-ii),

wherein

m is an integer 0 or 2;

each R^1 is independently -F, C_1-C_3 alkyl, C_1-C_3 alkylene-NR^7R^8, C_1-C_3 alkylene-$NR^{7'}R^{8'}$, C_1-C_3 alkylene-OH, C_1-C_3 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_3 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$,C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-OH, C_1-C_2 alkylene-(C_4-C_6 heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, or C_1-C_2 alkylene-(C_4-C_6 heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, wherein the alkyl, alkylene, cycloalkylene, and heterocycloalkylene in R^1 are optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;


R^2 is -H, -CH_3, CD_3, -CHF_2, or -CH_2CH_3;

R^3 is -H, C_1-C_3 alkyl, C_3-cycloalkyl, halogen, or -CN;

$R^{3'}$ is -H, C_1-C_3 alkyl, C_3-cycloalkyl, or -CN;

R^4 is -H, -CH_3, -CD_3, -CHF_2, -CH_2CH_3, or halogen;

R^5 is 5-to-10-membered heteroaryl, wherein said 5-to-10-membered heteroaryl is selected from the group consisting of:



wherein ⎓ indicates a single or double bond, and wherein said 5-to-10-membered heteroaryl is optionally substituted with 1-5 R^9 groups;

each R^7 is independently -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkyl-CN, or C_2-C_3 heteroalkyl;

each R^8 is independently -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkyl-CN, or C_2-C_3 heteroalkyl;

each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 4-to-6-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally

substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^9 is independently halogen, -OR^{10}, C_1-C_3 alkyl, -CF_2H, -CF_3, C_3-C_6 cycloalkyl, or
-CN, and

each R^{10} is independently -H, C_1-C_3 alkyl, -CD_3, -CF_2H, -CF_3, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with hydroxyl and/or C_1-C_3 alkoxy and/or 1-6 deuterium atoms.


Claim 4. The compound of any one of claims 1 to 3, or a pharmaceutically acceptable salt thereof, wherein

each R^1 is independently -F, C_1-C_3 alkylene-$NR^{7'}R^{8'}$, or C_1-C_3 alkylene-OH; and

wherein each pair of $R^{7'}$ and $R^{8'}$ of R^1 taken together with the nitrogen atom to which they are attached independently form a 4-to-6-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N and O, and wherein the nitrogen atom of any primary or secondary amine present in the heterocyclic ring is optionally substituted by -H or C_1-C_3 alkyl.

Claim 5. The compound of any one of claims 1 to 4, or a pharmaceutically acceptable salt thereof, wherein

R^5 is 5-to-10-membered heteroaryl, wherein said 5-to-10-membered heteroaryl is selected from the group consisting of:



wherein ⎓ indicates a single or double bond, and wherein said 5-to-10-membered heteroaryl is optionally substituted with 1-3 R^9 groups.

Claim 6. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein

R^2 is $-CH_3$, $-CD_3$, or $-CH_2CH_3$;

R^3 is -H, -F, $-CH_3$, or -CN;

$R^{3'}$ is -H or $-CH_3$; and


R^4 is -H, -F or $-CH_3$.

Claim 7. The compound of any one of claims 1 to 6, or a pharmaceutically acceptable salt thereof, wherein


each R^9 is independently -F, -Cl, $-OR^{10}$, $-CH_3$, or -CN, and

each R^{10} is independently -H, $-CH_3$, $-CD_3$, or $-CH_2CH_3$, wherein said $-CH_3$ or said $-CH_2CH_3$ is optionally substituted with hydroxyl and/or $-OCH_3$.

Claim 8. The compound of any one of claims 1 to 7, or a pharmaceutically acceptable salt thereof, wherein the compound of formula (I) is a compound of formula (I-A-i) or formula (I-A-ii):



(I-A-i), or



(I-A-ii),

wherein

m is an integer 0 or 1;

R^1 is -F, C_1-C_3 alkylene-$NR^{7'}R^{8'}$, or C_1-C_3 alkylene-OH;


R^2 is $-CH_3$, $-CD_3$, or $-CH_2CH_3$;

R^3 is -H, -F, $-CH_3$, or -CN;

$R^{3'}$ is -H or $-CH_3$;

R^4 is -H, -F or $-CH_3$;

R^5 is 5-to-10-membered heteroaryl, wherein said 5-to-10-membered heteroaryl is selected from the group consisting of:




wherein ⁻⁻⁻ indicates a single or double bond, and wherein said 5-to-10-membered heteroaryl is optionally substituted with 1-3 R^9 groups;

each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 4-to-6-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N and O, and wherein the nitrogen atom of any primary or secondary amine present in the heterocyclic ring is optionally substituted by -H or C_1-C_3 alkyl;


each R^9 is independently -F, -Cl, $-OR^{10}$, $-CH_3$, or -CN, and

each R^{10} is independently -H, $-CH_3$, $-CD_3$, or $-CH_2CH_3$, wherein said $-CH_3$ or said $-CH_2CH_3$ is optionally substituted with hydroxyl and/or $-OCH_3$.

Claim 9. The compound of any one of claims 1 to 8, or a pharmaceutically acceptable salt thereof, wherein the compound of formula (I) is a compound of formula (I-A-i) or formula (I-A-ii):



(I-A-i), or



(I-A-ii),

wherein

m is an integer 0 or 1;

R^1 is -F;


R^2 is $-CH_3$;

R^3 is -H or $-CH_3$;


$R^{3'}$ is -H or $-CH_3$;

R^4 is $-CH_3$;

R^5 is a 5-to-10-membered heteroaryl, wherein said 5-to-10-membered heteroaryl is selected from the group consisting of:



and




,

wherein said 5-to-10-membered heteroaryl is optionally substituted with 1-3 R^9 groups;

each R^9 is independently -F or $-OR^{10}$, and

each R^{10} is independently -H or $-CH_3$.

Claim 10. A compound of formula (I-B):



(I-B)

or a pharmaceutically acceptable salt thereof, wherein:

X is $NR^{3'}$ or CR^3,

Y is NR^2 or CR^4,

wherein when X is $NR^{3'}$ then Y is CR^4, Y has a double bond to CR^5, and X has a single bond to CR^5; or when X is CR^3 then Y is NR^2, Y has a single bond to CR^5, and X has a double bond to CR^5;

m is an integer from 0 to 3;

each R^1 is independently -D, -F, C_1-C_3 alkyl, C_1-C_3 alkylene-NR^7R^8, C_1-C_3 alkylene-$NR^{7'}R^{8'}$, C_1-C_3 alkylene-OH, C_1-C_3 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-OH, C_1-C_2 alkylene-(4- to 8-membered heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(4- to 8-membered heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_7 heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, or C_1-C_2 alkylene-(C_3-C_7 heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, wherein the alkyl, alkylene, cycloalkylene, and heterocycloalkylene moieties in R^1 are optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

R^2 is -H, C_1-C_3 alkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^3 is -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, halogen, or -CN;

$R^{3'}$ is -H, C_1-C_3 alkyl, -C_3-C_6 cycloalkyl, or -CN;

R^4 is -H, C_1-C_3 alkyl, or halogen, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^5 is C_6-C_{14} aryl or 5-to-10-membered heteroaryl, wherein said C_6-C_{14} aryl or said 5-to-10-membered heteroaryl is optionally substituted with 1-5 R^9 groups;

each R^7 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;

each R^8 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;


each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 3- to 8-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^9 is independently halogen, -OR^{10}, -NR^7R^8, C_1-C_3 alkyl, C_1-C_3 haloalkyl, C_3-C_6 cycloalkyl, -CN, $S(O)_nC_1$-C_3 alkyl, or $S(O)_nC_3$-C_6 cycloalkyl,

wherein n is an integer from 0 to 2; and

each R^{10} is independently -H, C_1-C_3 alkyl, C_1-C_3 haloalkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with hydroxyl, C_1-C_3 alkoxy, and/or 1-6 deuterium atoms.

Claim 11. A compound of formula (I-B’):



(I-B’)

or a pharmaceutically acceptable salt thereof, wherein:

X is $NR^{3'}$ or CR^3,

Y is NR^2 or CR^4,

wherein when X is $NR^{3'}$ then Y is CR^4, Y has a double bond to CR^5, and X has a single bond to CR^5; or when X is CR^3 then Y is NR^2, Y has a single bond to CR^5, and X has a double bond to CR^5;

m is an integer from 0 to 3;

R^2 is -H, C_1-C_3 alkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^3 is -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, halogen, or -CN;

$R^{3'}$ is -H, C_1-C_3 alkyl, -C_3-C_6 cycloalkyl, or -CN;

R^4 is -H, C_1-C_3 alkyl, or halogen, wherein said C_1-C_3 alkyl is optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms;

R^5 is C_6-C_{14} aryl or 5-to-10-membered heteroaryl, wherein said C_6-C_{14} aryl or said 5-to-10-membered heteroaryl is optionally substituted with 1-5 R^9 groups;

R^6 is -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 alkylene-NR^7R^8, C_1-C_6 alkylene-$NR^{7'}R^{8'}$, C_1-C_6 alkylene-OH, C_1-C_6 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-OH, C_1-C_2 alkylene-(C_4-C_6 heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, or C_1-C_2 alkylene-(C_4-C_6 heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, wherein the alkyl, alkylene, cycloalkyl, cycloalkylene, and heterocycloalkylene moieties in R^6 are optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^7 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;

each R^8 is independently -H, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_1-C_6 haloalkyl, C_1-C_6 alkylene-CN, or C_1-C_6 heteroalkyl;


each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 4-to-6-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^9 is independently halogen, -OR^{10}, -NR^7R^8, C_1-C_3 alkyl, C_1-C_3 haloalkyl, C_3-C_6 cycloalkyl, -CN, $S(O)_nC_1$-C_3 alkyl, or $S(O)_nC_3$-C_6 cycloalkyl,


wherein n is an integer from 0 to 2; and

each R^{10} is independently -H, C_1-C_3 alkyl, C_1-C_3 haloalkyl, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with hydroxyl, C_1-C_3 alkoxy, and/or 1-6 deuterium atoms.

Claim 12. The compound of claim 10 or 11, or a pharmaceutically acceptable salt thereof, wherein the compound of formula (I) is a compound of formula (I-B-i) or formula (I-B-ii):



(I-B-i), or



(I-B-ii),

wherein


R^2 is -H, -CH_3, CD_3, -CHF_2, or -CH_2CH_3;

R^3 is -H, C_1-C_3 alkyl, C_3-cycloalkyl, halogen, or -CN;

$R^{3'}$ is -H, C_1-C_3 alkyl, C_3-cycloalkyl, or -CN;

R^4 is -H, -CH_3, -CD_3, -CHF_2, -CH_2CH_3, or halogen;

R^5 is C_6-C_{14} aryl or 5-to-10-membered heteroaryl, wherein said 5-to-10-membered heteroaryl is selected from the group consisting of:



wherein ⎓ indicates a single or double bond, and wherein said C_6-C_{14} aryl or said 5-to-10-membered heteroaryl is optionally substituted with 1-5 R^9 groups;

R^6 is C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_1-C_3 alkylene-NR^7R^8, C_1-C_3 alkylene-$NR^{7'}R^{8'}$, C_1-C_3 alkylene-OH, C_1-C_3 alkylene-CN, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, C_1-C_2 alkylene-(C_3-C_6 cycloalkylene)-(C_0-C_2 alkylene)-OH, C_1-C_2 alkylene-(C_4-C_6

heterocycloalkylene)-(C_0-C_2 alkylene)-NR^7R^8, or C_1-C_2 alkylene-(C_4-C_6 heterocycloalkylene)-(C_0-C_2 alkylene)-$NR^{7'}R^{8'}$, wherein the alkyl, alkylene, cycloalkyl, cycloalkylene, and heterocycloalkylene moieties in R^6 are optionally substituted with 1-3 fluorine atoms and/or 1-6 deuterium atoms, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^7 is independently -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkyl-CN, or C_2-C_3 heteroalkyl;

each R^8 is independently -H, C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkyl-CN, or C_2-C_3 heteroalkyl;

each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 4-to-6-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^9 is independently halogen, $-OR^{10}$, C_1-C_3 alkyl, $-CF_2H$, $-CF_3$, C_3-C_6 cycloalkyl, or -CN, and

each R^{10} is independently -H, C_1-C_3 alkyl, $-CD_3$, $-CF_2H$, $-CF_3$, or C_3-C_6 cycloalkyl, wherein said C_1-C_3 alkyl is optionally substituted with hydroxyl and/or C_1-C_3 alkoxy.


Claim 13. The compound of any one of claims 1, 2 and 10 to 12, or a pharmaceutically acceptable salt thereof, wherein

R^6 is C_1-C_3 alkyl or C_1-C_3 alkylene-$NR^{7'}R^{8'}$; and

wherein each pair of $R^{7'}$ and $R^{8'}$ of R^6 taken together with the nitrogen atom to which they are attached independently form a 4-to-6-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl.

Claim 14. The compound of any one of claims 1, 2 and 10 to 13, or a pharmaceutically acceptable salt thereof, wherein

R^5 is phenyl or 5-to-10-membered heteroaryl, wherein said 5-to-10-membered heteroaryl is selected from the group consisting of:



,

wherein said phenyl or said 5-to-10-membered heteroaryl is optionally substituted with 1-3 R^9 groups.

Claim 15. The compound of any one of claims 1, 2 and 10 to 14, or a pharmaceutically acceptable salt thereof, wherein

each R^9 is independently -F, $-OR^{10}$, or $-CH_3$, and

each R^{10} is independently -H, $-CH_3$, $-CD_3$, $-CF_2H$, or $-CF_3$.

Claim 16. The compound of any one of claims 1, 2 and 10 to 15, or a pharmaceutically acceptable salt thereof, wherein


R^2 is -H or $-CH_3$;

R^3 is -H;


$R^{3'}$ is -H; and

R^4 is -H or $-CH_3$.

Claim 17. The compound of any one of claims 1, 2 and 10 to 16, or a pharmaceutically acceptable salt thereof, wherein the compound of formula (I) is a compound of formula (I-B-i) or formula (I-B-ii):



(I-B-i), or



(I-B-ii),


wherein

R^2 is -H or -CH_3;

R^3 is -H;

$R^{3'}$ is -H;

R^4 is -H or -CH_3;

R^5 is phenyl or 5-to-10-membered heteroaryl, wherein said 5-to-10-membered heteroaryl is selected from the group consisting of: 

, wherein said phenyl or said 5-to-10-membered heteroaryl is optionally substituted with 1-3 R^9 groups;


R^6 is C_1-C_3 alkyl or C_1-C_3 alkylene-$NR^{7'}R^{8'}$;


each pair of $R^{7'}$ and $R^{8'}$ taken together with the nitrogen atom to which they are attached independently form a 4-to-6-membered heterocyclic ring, wherein the heterocyclic ring optionally contains an additional 1-2 heteroatoms selected from the group consisting of N, O, and S, and wherein each heterocyclic nitrogen atom, if present, is independently optionally substituted with C_1-C_3 alkyl, C_3-C_6 cycloalkyl, C_2-C_3 haloalkyl, C_2-C_3 alkylene-CN, or C_2-C_3 heteroalkyl;

each R^9 is independently -F, -OR^{10}, or -CH_3, and


each R^{10} is independently -H, -CH_3, -CD_3, -CF_2H, or -CF_3.


Claim 18. A compound selected from the group consisting of:




;



;




;



;




;



;




;



;

















;


;















;


;


;

;

;

;

;

;

;

;

;

;

;

;

;

;













;

;




;

;

;

;

;

;

;

;

;

;

;

;













;

;


;

;

;

;

;

;

;

;

;

;

;

;



;



;

;

;




;

;

;

;

;

;

;

;

;

;

;

;



;



;




;

;












;

;


;

;


;

;














;

;












;

;












;

,

;

;

;

;

;

;

;

;


;


;


D3C
O
N
N
O
F
O
N
H
N
O
;

CD3
O
N
N
O
F
O
N
H
N
O
CD3
;

O
N
N
O
CHF2
F
O
N
H
N
O
;

O
N
N
O
F
O
N
H
N
—O
;

H
N
O
N
N
O
F
O
N
H
N
O
;

O
N
N
O
F
O
N
H
N
O
;

O
N
N
F
O
F
O
N
H
N
—O
;

O
N
N
O
F
O
N
H
N
O
O
;

O
N
N
O
NH2
F
O
N
H
N
O
;

O
N
N
O
N
F
O
N
H
N
O
;

O
N
N
O
NH
F
O
N
H
N
O
;

O
N
N
O
F
O
N
H
N
HO
;


;

;

;

;

;

;

;

;

;

;




;

;

;

;

;

;

;

;

;

;



;

;



;

;

;



;


;

;


;

;




;

;













and
















;

,

or a pharmaceutically acceptable salt thereof.

Claim 19. A compound selected from the group consisting of:





;

;











;

;

;

;

;

;

;

;

;

;

;

;




;

;

;

;

;

;

;

;

;

;

;

;

;

;




;













;














;

;

;











;

;

;

;

;

;

;

;

;

;

;

;











;

;

;

;














F
O
N
F
(R)
(R)
N
H
N
O
;

O
N
F
(S)
(S)
N
H
N
O
N
;

F
O
N
F
(S)
(S)
N
H
N
O
N
;

F
O
N
N
H
N
;

N
O
N
F
(S)
(S)
N
H
N
O
;

F
N
O
N
F
(S)
(S)
N
H
N
O
;

N
O
N
F
(S)
(S)
N
H
N
O
;

F
N
O
N
F
(S)
(S)
N
H
N
O
;

N
O
N
F
(S)
(S)
N
H
N
O
;

CD3
N
O
N
F
(S)
(S)
N
H
N
O
;

CD3
N
F
O
N
F
(S)
(S)
N
H
N
O
;

Cl
N
O
N
F
(S)
(S)
N
H
N
O
;














;

;

;

;

;

;

;

;

;

;

;

;














;

;

;

;

;

,

;

;

;

;


;

;

;

;

;

;

;

;

;

;













F
(S)
(S)
O
N
H
N
H
O
N
N
O
;

F
O
N
H
N
N
O
N
N
O
;

F
O
N
H
N
N
O
F
N
O
;

F
O
N
H
N
N
O
N
N
O
O
;

F
O
N
H
N
N
O
N
N
NH2
O
;

F
O
N
H
N
N
O
N
N
N
O
;

F
O
N
H
N
N
O
N
N
NH
O
;

F
O
N
H
N
N
O
N
HO
;

F
O
N
H
N
N
O
N
N
O
;

F
O
N
H
N
N
O
N
N
OH
O
;

F
O
NH
N
N
NH
;

F
O
N
H
N
N
O
N
HN
;


;

;

;

;

;

;

;

;

;

;

;

;



;

;

;

;

;

;

;

;

;

;

;

;


;

;


;

;

;

;

;

;

;

;

;

;

;

;


;

;

and 

,

or a pharmaceutically acceptable salt thereof.


Claim 20. A compound having the structure



,

or a pharmaceutically acceptable salt thereof.


Claim 21. A compound having the structure



,

or a pharmaceutically acceptable salt thereof.

Claim 22. A compound having the structure



,

or a pharmaceutically acceptable salt thereof.

Claim 23. A pharmaceutical composition comprising the compound of any one of claims 1 to 22, or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable excipients.

Claim 24. A compound of any one of claims 1 to 22, or a pharmaceutically acceptable salt thereof, or the pharmaceutical composition of claim 23, for use in inhibiting Bcr-Abl enzymatic activity in a cell.

Claim 25. Use of a compound of any one of claims 1 to 22, or a pharmaceutically acceptable salt thereof, or the pharmaceutical composition of claim 23, in the manufacture of a medicament for treating chronic myeloid leukemia (CML), acute myeloid leukemia (AML), acute lymphoblastic leukemia (ALL), or a mixed phenotype acute leukemia, in a human in need thereof.

Claim 26. The use of claim 25, wherein the leukemia is refractory leukemia.

Claim 27. The use of claim 26, wherein the refractory leukemia is associated with a mutation in the Bcr-Abl tyrosine kinase gene resulting in specific amino acid substitutions selected from the group consisting of M244V, L248V, G250E, G250A, Q252H, Q252R, Y253F, Y253H, E255K, E255V, D276G, F311L, T315N, T315A, F317V, F317L, M343T, M351T, E355G, F359A, F359V, V379I, F382L, L387M, H396P, H396R, S417Y, E459K, F486S, and T315I.

Claim 28. The use of claim 27, wherein the refractory leukeumia is associated with a mutation in the Bcr-Abl tyrosine kinase gene resulting in specific amino acid substitution T315I.

Claim 29. The use of claim 26, wherein the human having refractory leukemia has one or more mutations in the Bcr-Abl tyrosine kinase gene resulting in specific amino acid substitutions selected from the group consisting of M244V, L248V, G250E, G250A, Q252H, Q252R, Y253F, Y253H, E255K, E255V, D276G, F311L, T315N, T315A, F317V, F317L, M343T, M351T, E355G, F359A, F359V, V379I, F382L, L387M, H396P, H396R, S417Y, E459K, F486S, and T315I.

Claim 30. The use of claim 29, wherein the human having refractory leukemia has a mutation in the Bcr-Abl tyrosine kinase gene resulting in specific amino acid substitution T315I.

Claim 31. The compound for use of claim 24, or the use of any one of claims 25 to 30, wherein the compound, or a pharmaceutically acceptable salt thereof, or the pharmaceutical composition is to be administered, and wherein one or more pharmaceutical agents including

anti-microtubular therapies, topoisomerase inhibitors, alkylating agents, nucleotide synthesis inhibitors, DNA synthesis inhibtiors, protein synthesis inhibitors, developmental signaling pathway inhibitors, pro-apoptotic agents, Abl myristoyl-pocket binding inhibitors, MEK1/2 inhibitors, AKT inhibitors, PI3K inhibitors and/or radiation are to be further administered.

Claim 32. The compound for use or use of claim 31, wherein the one or more pharmaceutical agents include Abl myristoyl-pocket binding inhibitors.

Claim 33. The compound for use or use of claim 32, wherein the Abl myristoyl-pocket binding inhibitors is asciminib.

Claim 34. A compound of any one of claims 1 to 22, or a pharmaceutically acceptable salt thereof, or the pharmaceutical composition of claim 23, for use in therapy.