




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 277/48, 417/12, 417/14, 471/10, 491/10, A61K 31/426, 31/427, C07D 471/04 // (C07D 471/10, 235:00, 221:00) (C07D 491/10, 317:00, 221:00) (C07D 471/04, 235:00, 221:00)	A1	(11) International Publication Number: WO 00/26203 (43) International Publication Date: 11 May 2000 (11.05.00)
(21) International Application Number: PCT/EP99/08307 (22) International Filing Date: 27 October 1999 (27.10.99) (30) Priority Data: 9823873.6 30 October 1998 (30.10.98) GB (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN S.P.A. [IT/IT]; Via Robert Koch, 1.2, I-20152 Milano (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): PEVARELLO, Paolo [IT/IT]; Piazza San Pietro in Ciel d'Oro, 7/A, I-27100 Pavia (IT). AMICI, Raffaella [IT/IT]; Via N. Rocca, 11, I-29100 Piacenza (IT). TRAQUANDI, Gabriella [IT/IT]; Via F. Cilea, 106, I-20151 Milano (IT). VILLA, Manuela [IT/IT]; Via San Bernardino, 12, I-22040 Lurago d'Erba (IT). VULPETTI, Anna [IT/IT]; Via Voltumo Portici/2 80, I-20047 Brugherio (IT). ISACCHI, Antonella [IT/IT]; Via Montecatini, 14, I-20144 Milano (IT).		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: 2-UREIDO-THIAZOLE DERIVATIVES, PROCESS FOR THEIR PREPARATION, AND THEIR USE AS ANTITUMOR AGENTS  (I) (57) Abstract Compounds which are 2-ureido-1, 3-thiazole derivatives of formula (I) wherein R is a halogen atom, a nitro group, an optionally substituted amino group or it is a group, optionally further substituted, selected from: i) straight or branched C₁-C₆ alkyl; ii) C₃-C₆ cycloalkyl; iii) aryl or arylalkyl with from 1 to 6 carbon atoms within the straight or branched alkyl chain; R₁ is an optionally substituted group selected from: i) straight or branched C₁-C₆ alkyl; ii) 3 to 6 membered carbocycle or 5 to 7 membered heterocycle ring; iii) aryl or arylcarbonyl; iv) arylalkyl with from 1 to 6 carbon atoms within the straight or branched alkyl chain; R₂ is hydrogen, a straight or branched C₁-C₄ alkyl or C₂-C₄ alkenyl or alkynyl group; or, taken together with the nitrogen atom to which they are bonded, R₁ and R₂ form a substituted or unsubstituted group selected from: i) an optionally benzocondensed or bridged 5 to 7 membered heterocycle; or ii) a 9 to 11 membered spiro-heterocyclic compound; or a pharmaceutically acceptable salt thereof; are useful for treating cell proliferative disorders associated with an altered cell dependent kinase activity.		