

TRICYCLIC QUINOLINE AND QUINOXALINE DERIVATIVES

FIELD OF THE INVENTION

The present invention relates to tricyclic quinoline and quinoxaline derivatives, to a pharmaceutical composition containing such compounds, to their use as modulators, especially agonists or partial agonists, of the 5-HT_{2C} receptor, their use for preparing a medicament for the prevention or treatment of conditions and disorders which respond to the modulation of 5-HT_{2C} receptor, to a method for preventing or treating conditions and disorders which respond to the modulation of 5-HT_{2C} receptor, and processes for preparing such compounds and compositions.

BACKGROUND OF THE INVENTION

Diseases, disorders and conditions where 5-HT_{2C} modulation is desired are for example depression, anxiety, schizophrenia, bipolar disorder, obsessive compulsive disorder, migraine, pain, epilepsy, substance abuse, eating disorders, obesity, diabetes, erectile dysfunction and others.

Serotonin (5-hydroxytryptamine, 5-HT), a monoamine neurotransmitter and local hormone, is formed by the hydroxylation and decarboxylation of tryptophan. The greatest concentration is found in the enterochromaffin cells of the gastrointestinal tract, the remainder being predominantly present in platelets and in the Central Nervous System (CNS). 5-HT is implicated in a vast array of physiological and pathophysiological pathways. In the periphery, it contracts a number of smooth muscles and induces endothelium-dependent vasodilation. In the CNS, it is believed to be involved in a wide range of functions, including the control of appetite, mood, anxiety, hallucinations, sleep, vomiting and pain perception.

Neurons that secrete 5-HT are termed serotonergic. The function of 5-HT is exerted upon its interaction with specific (serotonergic) neurons. Seven types of 5-HT receptors have been identified: 5-HT₁ (with subtypes 5-HT_{1A}, 5-HT_{1B}, 5-HT_{1D}, 5-HT_{1E} and 5-HT_{1F}), 5-HT₂ (with subtypes 5-HT_{2A}, 5-HT_{2B} and 5-HT_{2C}), 5-HT₃, 5-HT₄, 5-HT₅ (with subtypes 5-HT_{5A} and 5-HT_{5B}), 5-HT₆ and 5-HT₇. Most of these receptors are coupled to G-proteins that affect the activities of adenylate cyclase or phospholipase C γ .

Alterations in the activity of multiple neurotransmitter receptor systems (dopamine, serotonin, glutamate, GABA, acetylcholine) have been implicated in the manifestation of the symptoms of schizophrenia. The most widely accepted "Dopamine Hypothesis of Schizophrenia" in its simplest form states that the positive symptoms of this pathology relate to a functional hyperactivity of the mesolimbic dopaminergic system, while the negative and cognitive aspects can be traced to a functional hypoactivity of the mesocortical dopaminergic projections. Atypical antipsychotics block the mesolimbic dopaminergic neurotransmission, thereby controlling positive symptoms, with little or no effect on the nigrostriatal system, leading to less induction of extrapyramidal side effects (EPS).

Primary negative and cognitive symptoms of schizophrenia reflect a dysfunction of the frontal cortex ("hypofrontality"), which is thought to be induced by a decreased tone in the mesocortical dopaminergic projection field [Davis KL, Kahn RS, Ko G and Davidson M (1991). Dopamine in schizophrenia: a review and re-conceptualization. *Am J Psychiatry* 148: 1474 – 86. Weinberger DR and Berman KF (1996). Prefrontal function in schizophrenia: confounds and controversies. *Philos Trans R Soc Lond B Biol Sci* 351: 1495 - 503]. Agents that selectively enhance dopamine levels in the cortex have the potential to address the negative symptoms of this disorder. Atypical antipsychotics lack robust efficacy against negative and cognitive components of the schizophrenic syndrome.

The schizophrenic symptomatology is further complicated by the occurrence of drug-induced so-called secondary negative symptoms and cognitive impairment, which are difficult to distinguish from primary negative and cognitive symptoms [Remington G and Kapur S (2000). Atypical antipsychotics: are some more atypical than others? *Psychopharmacol* 148: 3 – 15]. The occurrence of secondary negative symptoms not only limits therapeutic efficacy but also, together with these side effects, negatively affects patient compliance.

It may thus be hypothesized that a novel mechanistic approach that blocks dopaminergic neurotransmission in the limbic system but does not affect the striatal and pituitary projection fields, and stimulates frontocortical projection fields, would provide an efficacious treatment for all parts of the schizophrenic pathology, including its positive, negative and cognitive symptoms. Moreover, a selective compound that is substantially

free of the ancillary pharmacology that characterizes current agents would be expected to avoid a variety of off-target side effects that plague current treatments such as extrapyramidal side effects (EPS) and weight gain.

The 5-HT_{2C} receptor, previously named 5-HT_{1C}, is a G-protein-coupled receptor, which couples to multiple cellular effector systems including the phospholipase C, A and D pathways. It is found primarily in the brain and its distribution is particularly high in the plexus choroideus, where it is assumed to control cerebrospinal fluid production [Kaufman MJ, Hirata F (1996) Cyclic GMP inhibits phosphoinositide turnover in choroid plexus: evidence for interactions between second messengers concurrently triggered by 5-HT_{2C} receptors. *Neurosci Lett* 206:153-156]. Very high levels were also found in the retrosplenial, piriform and entorhinal cortex, anterior olfactory nucleus, lateral septal nucleus, subthalamic nucleus, amygdala, subiculum and ventral part of CA3, lateral habenula, substantia nigra pars compacta, several brainstem nuclei and the whole grey matter of the spinal cord [Pompeiano M, Palacios JM, Mengod G (1994). Distribution of the serotonin 5-HT₂ receptor family mRNAs: comparison between 5-HT_{2A} and 5-HT_{2C} receptors. *Brain Res Mol Brain Res* 23:163-178]. A comparison of the distribution of 5-HT_{2C} mRNA with that of 5-HT_{2C} protein in monkey and human brains has revealed both pre- and postsynaptic localization [Lopez-Gimenez JF, Mengod G, Palacios JM, Vilaro MT (2001) Regional distribution and cellular localization of 5-HT_{2C} receptor mRNA in monkey brain: comparison with [³H]mesulergine binding sites and choline acetyltransferase mRNA. *Synapse* 42:12-26].

It is anticipated that modulation of the 5-HT_{2C} receptor will improve disorders such as depression, anxiety, schizophrenia, cognitive deficits of schizophrenia, obsessive compulsive disorder, bipolar disorder, migraine, epilepsy, substance abuse, eating disorders, obesity, diabetes, sexual dysfunction/erectile dysfunction, sleep disorders, psoriasis, Parkinson's disease, pain conditions and disorders, and spinal cord injury, smoking cessation, ocular hypertension and Alzheimer's disease. Modulators of the 5-HT_{2C} receptor are also shown to be useful in the modulation of bladder function, including the prevention or treatment of urinary incontinence.

There is an ongoing need for providing compounds having high affinity and selectivity for the 5-HT_{2C} receptor. In particular the compounds should have low affinity to adrenergic receptors, such as the α_1 -adrenergic receptor, histamine receptors, such as


the H₁-receptor, and dopaminergic receptors, such as the D₂-receptor, in order to avoid or reduce side effects associated with modulation of these receptors, such as postural hypotension, reflex tachycardia, potentiation of the antihypertensive effect of prazosin, terazosin, doxazosin and labetalol or dizziness associated with the blockade of the α_1 -adrenergic receptor, weight gain, sedation, drowsiness or potentiation of central depressant drugs associated with the blockade of the H₁-receptor, or extrapyramidal movement disorder, such as dystonia, parkinsonism, akathisia, tardive dyskinesia or rabbit syndrome, or endocrine effects, such as prolactin elevation (galactorrhea, gynecomastia, menstrual changes, sexual dysfunction in males), associated with the blockade of the D₂-receptor.

The present invention provides compounds which have an affinity for the 5-HT_{2C}, thus allowing the treatment of disorders related to or affected by the 5-HT_{2C} receptor.

SUMMARY OF THE INVENTION

The invention is directed to tricyclic quinoline and quinoxaline derivatives, compositions comprising such compounds, their use as modulators, especially agonists or partial agonists, of the 5-HT_{2C} receptor, their use for preparing a medicament for the prevention or treatment of conditions and disorders which respond to the modulation of 5-HT_{2C} receptor, to a method for preventing or treating conditions and disorders which respond to the modulation of 5-HT_{2C} receptor, and processes for preparing such compounds and compositions.

In one aspect, the present invention relates to compounds of the formula (I):



wherein

G is $(CR^{3a}R^{3b})_n$;

X is NR^6 or CR^7R^8 ;

R^1 is selected from the group consisting of hydrogen, cyano, $\text{C}_1\text{-C}_6\text{-alkyl}$, fluorinated $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_2\text{-C}_6\text{-alkenyl}$, fluorinated $\text{C}_2\text{-C}_6\text{-alkenyl}$, $\text{C}_2\text{-C}_6\text{-alkynyl}$, fluorinated $\text{C}_2\text{-C}_6\text{-alkynyl}$, $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, fluorinated $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, $\text{C}_1\text{-C}_6\text{-alkoxy}$, fluorinated $\text{C}_1\text{-C}_6\text{-alkoxy}$, $-\text{C}(=\text{O})\text{R}^{10}$, phenyl, phenyl- $\text{C}_1\text{-C}_2\text{-alkyl}$ and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO_2 and optionally also 1 or 2 $\text{C}=\text{O}$ and/or $\text{C}=\text{S}$ groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R^{11} ;

each R^2 is independently selected from the group consisting of cyano, nitro, $\text{C}_1\text{-C}_6\text{-alkyl}$, fluorinated $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-hydroxyalkyl}$, $\text{C}_2\text{-C}_6\text{-alkenyl}$, fluorinated $\text{C}_2\text{-C}_6\text{-alkenyl}$, $\text{C}_2\text{-C}_6\text{-alkynyl}$, fluorinated $\text{C}_2\text{-C}_6\text{-alkynyl}$, $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, fluorinated $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, $\text{C}_1\text{-C}_6\text{-alkoxy}$, fluorinated $\text{C}_1\text{-C}_6\text{-alkoxy}$, $\text{C}_1\text{-C}_6\text{-alkoxy-}\text{C}_1\text{-C}_4\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-hydroxyalkoxy}$, $\text{C}_1\text{-C}_6\text{-alkoxy-}\text{C}_1\text{-C}_4\text{-alkoxy}$, $\text{C}_1\text{-C}_6\text{-alkylthio}$, fluorinated $\text{C}_1\text{-C}_6\text{-alkylthio}$, $\text{C}_1\text{-C}_6\text{-alkylsulfinyl}$, fluorinated $\text{C}_1\text{-C}_6\text{-alkylsulfinyl}$, $\text{C}_1\text{-C}_6\text{-alkylsulfonyl}$, fluorinated $\text{C}_1\text{-C}_6\text{-alkylsulfonyl}$, $-\text{NR}^{12a}\text{R}^{12b}$, $-\text{CH}_2\text{NR}^{12a}\text{R}^{12b}$, $-\text{NR}^{12a}\text{C}(\text{O})\text{R}^{10}$, $-\text{C}(=\text{O})\text{R}^{10}$, $\text{SO}_2\text{NR}^{12a}\text{R}^{12b}$, $\text{C}_1\text{-C}_6\text{-alkylcarbonyloxy}$, fluorinated $\text{C}_1\text{-C}_6\text{-alkylcarbonyloxy}$, phenyl, phenyl- $\text{C}_1\text{-C}_2\text{-alkyl}$, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO_2 , $\text{C}=\text{O}$ and $\text{C}=\text{S}$ as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R^{11} ;

R^{3a} and R^{3b} , independently of each other, are selected from the group consisting of hydrogen, cyano, nitro, $\text{C}_1\text{-C}_6\text{-alkyl}$, fluorinated $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-hydroxyalkyl}$, $\text{C}_2\text{-C}_6\text{-alkenyl}$, fluorinated $\text{C}_2\text{-C}_6\text{-alkenyl}$, $\text{C}_2\text{-C}_6\text{-alkynyl}$, fluorinated $\text{C}_2\text{-C}_6\text{-alkynyl}$, $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, fluorinated $\text{C}_3\text{-C}_8\text{-cycloalkyl}$, $\text{C}_1\text{-C}_6\text{-alkoxy}$, fluorinated

C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹;

R^{4a} and R^{4b}, independently of each other, are selected from the group consisting of hydrogen, cyano, nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹; or R^{4a} and R^{4b} form together a group =O or =S;

R^{5a} and R^{5b}, independently of each other, are selected from the group consisting of hydrogen, deuterium, halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-

alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹; where R^{5a} and R^{5b} are not simultaneously hydroxy; or

R^{5a} and R^{5b}, together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring (i.e. a spiro ring), where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹; or

R^{5a} and R⁶, together with the atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹; or

R^{5a} and R⁷, together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹;

R^6 is selected from the group consisting of hydrogen, cyano, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_3 - C_8 -cycloalkyl- C_1 - C_4 -alkyl, fluorinated C_3 - C_8 -cycloalkyl- C_1 - C_4 -alkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, $-C(=O)R^{10}$, $-SO_2R^{10}$, phenyl, phenyl- C_1 - C_2 -alkyl and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO_2 and optionally also 1 or 2 $C=O$ and/or $C=S$ groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R^{11} ;

R^7 and R^8 , independently of each other, are selected from the group consisting of deuterium, halogen, cyano, nitro, hydroxyl, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, fluorinated C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, fluorinated C_1 - C_6 -alkylsulfonyl, $-NR^{12a}R^{12b}$, $-CH_2NR^{12a}R^{12b}$, $-NR^{12a}C(O)R^{10}$, $-C(=O)R^{10}$, $SO_2NR^{12a}R^{12b}$, C_1 - C_6 -alkylcarbonyloxy, fluorinated C_1 - C_6 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_2 -alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO_2 , $C=O$ and $C=S$ as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R^{11} ; where R^7 and R^8 are not simultaneously hydroxyl; and where R^7 is not hydroxyl if R^8 is C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, phenoxy or benzyloxy; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring (i.e. a spiro ring), where the ring may contain 1, 2, 3 or 4 heteroatoms or het-

eroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹;

each R⁹ is independently selected from the group consisting of halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₃-C₈-cycloalkenyl, fluorinated C₃-C₈-cycloalkenyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹; or two radicals R⁹ bound on neighbouring carbon atoms, together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹;

each R¹⁰ is independently selected from the group consisting of hydrogen, cyano, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or

maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the five last-mentioned radicals may be substituted with one or more substituents R¹¹;

each R¹¹ is independently selected from the group consisting of halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -COOH, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyl, fluorinated C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, fluorinated C₁-C₆-alkoxycarbonyl, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy and fluorinated C₁-C₆-alkylcarbonyloxy; or two radicals R¹¹, together with the atom(s) they are bound to, form a saturated, partially unsaturated or maximally unsaturated 3-, 4-, 5-, 6- or 7-membered carbocyclic or heterocyclic ring, where the heterocyclic ring contains 1, 2 or 3 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members;

R^{12a} and R^{12b}, independently of each other and independently of each occurrence, are selected from the group consisting of hydrogen, cyano, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkylcarbonyl, fluorinated C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, fluorinated C₁-C₆-alkoxycarbonyl, phenyl and benzyl, where the phenyl moieties in the two last-mentioned radicals may carry 1, 2 or 3 substituents selected from halogen, cyano, nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy; or, if R^{12a} and R^{12b} are bound to the same nitrogen atom, together with this nitrogen atom may form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated

or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents selected from halogen, cyano nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy;

a is 0, 1 or 2;

b is 0, 1, 2 or 3; and

n is 1 or 2;

and the N-oxides, tautomeric forms, stereoisomers and pharmaceutically acceptable salts thereof, and the compound of the general formula I, wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope.

In another aspect, the invention relates to compounds of formula I as defined above, wherein

G is (CR^{3a}R^{3b})_n;

X is NR⁶ or CR⁷R⁸;

R¹ is selected from the group consisting of hydrogen, cyano, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, -C(=O)R¹⁰, phenyl, phenyl-C₁-C₂-alkyl and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO₂ and optionally also 1 or 2 C=O and/or C=S groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R¹¹;

each R^2 is independently selected from the group consisting of cyano, nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, fluorinated C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, fluorinated C_1 - C_6 -alkylsulfonyl, $-NR^{12a}R^{12b}$, $-CH_2NR^{12a}R^{12b}$, $-NR^{12a}C(O)R^{10}$, $-C(=O)R^{10}$, $SO_2NR^{12a}R^{12b}$, C_1 - C_6 -alkylcarbonyloxy, fluorinated C_1 - C_6 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_2 -alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO_2 , C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R^{11} ;

R^{3a} and R^{3b} , independently of each other, are selected from the group consisting of hydrogen, cyano, nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, fluorinated C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, fluorinated C_1 - C_6 -alkylsulfonyl, $-NR^{12a}R^{12b}$, $-CH_2NR^{12a}R^{12b}$, $-NR^{12a}C(O)R^{10}$, $-C(=O)R^{10}$, $SO_2NR^{12a}R^{12b}$, C_1 - C_6 -alkylcarbonyloxy, fluorinated C_1 - C_6 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_2 -alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO_2 , C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R^{11} ;

R^{4a} and R^{4b} , independently of each other, are selected from the group consisting of hydrogen, cyano, nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, fluorinated C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, fluorinated C_1 - C_6 -alkylsulfonyl, $-NR^{12a}R^{12b}$, $-CH_2NR^{12a}R^{12b}$, $-NR^{12a}C(O)R^{10}$, $-C(=O)R^{10}$, $SO_2NR^{12a}R^{12b}$, C_1 - C_6 -alkylcarbonyloxy, fluorinated C_1 - C_6 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_2 -alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO_2 , C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R^{11} ; or R^{4a} and R^{4b} form together a group =O or =S;

R^{5a} and R^{5b} , independently of each other, are selected from the group consisting of hydrogen, deuterium, halogen, cyano, nitro, hydroxy, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, fluorinated C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, fluorinated C_1 - C_6 -alkylsulfonyl, $-NR^{12a}R^{12b}$, $-CH_2NR^{12a}R^{12b}$, $-NR^{12a}C(O)R^{10}$, $-C(=O)R^{10}$, $SO_2NR^{12a}R^{12b}$, C_1 - C_6 -alkylcarbonyloxy, fluorinated C_1 - C_6 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_2 -alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO_2 , C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R^{11} ; or

R^{5a} and R^{5b} , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ; or R^{5a} and R^6 , together with the atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ; or

R^{5a} and R^7 , together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ;

R^6 is selected from the group consisting of hydrogen, cyano, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-C₁-C₄-alkyl, fluorinated C₃-C₈-cycloalkyl-C₁-C₄-alkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, -C(=O) R^{10} , -SO₂ R^{10} , phenyl, phenyl-C₁-C₂-alkyl and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO₂ and optionally also 1 or 2 C=O and/or C=S groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R^{11} ;

R^7 and R^8 , independently of each other, are selected from the group consisting of deuterium, halogen, cyano, nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy,

fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹; or R⁷ and R⁸, together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹;

each R⁹ is independently selected from the group consisting of halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₃-C₈-cycloalkenyl, fluorinated C₃-C₈-cycloalkenyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹; or

two radicals R^9 bound on neighbouring carbon atoms, together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ;

each R^{10} is independently selected from the group consisting of hydrogen, cyano, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the five last-mentioned radicals may be substituted with one or more substituents R^{11} ;

each R^{11} is independently selected from the group consisting of halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -COOH, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyl, fluorinated C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, fluorinated C₁-C₆-alkoxycarbonyl, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy and fluorinated C₁-C₆-alkylcarbonyloxy;

R^{12a} and R^{12b} , independently of each other and independently of each occurrence, are selected from the group consisting of hydrogen, cyano, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated

C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkylcarbonyl, fluorinated C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, fluorinated C₁-C₆-alkoxycarbonyl, phenyl and benzyl, where the phenyl moieties in the two last-mentioned radicals may carry 1, 2 or 3 substituents selected from halogen, cyano nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy; or, if R^{12a} and R^{12b} are bound to the same nitrogen atom, together with this nitrogen atom may form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents selected from halogen, cyano nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy;

a is 0, 1 or 2;


b is 0, 1, 2 or 3; and

n is 1 or 2;

and the N-oxides, tautomeric forms, stereoisomers and pharmaceutically acceptable salts thereof, and the compound of the general formula I, wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope.

Preferably, R^{5a} and R^{5b} are not simultaneously hydroxyl.

In another aspect, the invention relates to compounds of formula I



wherein

G is $(\text{CR}^{3a}\text{R}^{3b})_n$;

X is NR^6 or CR^7R^8 ;

R^1 is selected from the group consisting of hydrogen, cyano, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, $-\text{C}(=\text{O})\text{R}^{10}$, phenyl, phenyl- C_1 - C_2 -alkyl and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO_2 and optionally also 1 or 2 $\text{C}=\text{O}$ and/or $\text{C}=\text{S}$ groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R^{11} ;

each R^2 is independently selected from the group consisting of cyano, nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, fluorinated C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, fluorinated C_1 - C_6 -alkylsulfonyl, $-\text{NR}^{12a}\text{R}^{12b}$, $-\text{CH}_2\text{NR}^{12a}\text{R}^{12b}$, $-\text{NR}^{12a}\text{C}(\text{O})\text{R}^{10}$, $-\text{C}(=\text{O})\text{R}^{10}$, $\text{SO}_2\text{NR}^{12a}\text{R}^{12b}$, C_1 - C_6 -alkylcarbonyloxy, fluorinated C_1 - C_6 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_2 -alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or

maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹;

R^{3a} and R^{3b}, independently of each other, are selected from the group consisting of hydrogen, cyano, nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹;

R^{4a} and R^{4b}, independently of each other, are selected from the group consisting of hydrogen, cyano, nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered

saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹; or R^{4a} and R^{4b} form together a group =O or =S;

R^{5a} and R^{5b}, independently of each other, are selected from the group consisting of hydrogen, deuterium, halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹; or R^{5a} and R^{5b}, together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹; or R^{5a} and R⁶, together with the atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹; or

R^{5a} and R^7 , together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ;

R^6 is selected from the group consisting of hydrogen, cyano, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, -C(=O) R^{10} , -SO₂ R^{10} , phenyl, phenyl-C₁-C₂-alkyl and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO₂ and optionally also 1 or 2 C=O and/or C=S groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R^{11} ;

R^7 and R^8 , independently of each other, are selected from the group consisting of deuterium, halogen, cyano, nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O) R^{10} , -C(=O) R^{10} , SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R^{11} ; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ;

each R^9 is independently selected from the group consisting of halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₃-C₈-cycloalkenyl, fluorinated C₃-C₈-cycloalkenyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R^{11} ; or two radicals R^9 bound on neighbouring carbon atoms, together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ;

each R^{10} is independently selected from the group consisting of hydrogen, cyano, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-

alkoxy, $-\text{NR}^{12a}\text{R}^{12b}$, $-\text{CH}_2\text{NR}^{12a}\text{R}^{12b}$, phenyl, phenyl- C_1 - C_2 -alkyl, phenoxy, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO_2 , $\text{C}=\text{O}$ and $\text{C}=\text{S}$ as ring members, where the cyclic moieties in the five last-mentioned radicals may be substituted with one or more substituents R^{11} ;

each R^{11} is independently selected from the group consisting of halogen, cyano, nitro, hydroxy, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, fluorinated C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, fluorinated C_1 - C_6 -alkylsulfonyl, $-\text{COOH}$, $-\text{NR}^{12a}\text{R}^{12b}$, $-\text{CH}_2\text{NR}^{12a}\text{R}^{12b}$, C_1 - C_6 -alkylcarbonyl, fluorinated C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -alkoxycarbonyl, fluorinated C_1 - C_6 -alkoxycarbonyl, $\text{SO}_2\text{NR}^{12a}\text{R}^{12b}$, C_1 - C_6 -alkylcarbonyloxy and fluorinated C_1 - C_6 -alkylcarbonyloxy;

R^{12a} and R^{12b} , independently of each other and independently of each occurrence, are selected from the group consisting of hydrogen, cyano, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkylcarbonyl, fluorinated C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -alkoxycarbonyl, fluorinated C_1 - C_6 -alkoxycarbonyl, phenyl and benzyl, where the phenyl moieties in the two last-mentioned radicals may carry 1, 2 or 3 substituents selected from halogen, cyano, nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy and fluorinated C_1 - C_6 -alkoxy; or, if R^{12a} and R^{12b} are bound to the same nitrogen atom, together with this nitrogen atom may form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO_2 , $\text{C}=\text{O}$ and $\text{C}=\text{S}$ as ring members, and where the ring may be substituted with

one or more substituents selected from halogen, cyano nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy;

a is 0, 1 or 2;

b is 0, 1, 2 or 3; and

n is 1 or 2;

and the N-oxides, tautomeric forms, stereoisomers and pharmaceutically acceptable salts thereof, and the compound of the general formula I, wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope.

Preferably, R^{5a} and R^{5b} are not simultaneously hydroxyl.

In another aspect, the invention relates to a pharmaceutical composition comprising a therapeutically effective amount of at least one compound of formula I or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof, or comprising at least one compound as defined above or below wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope, preferably wherein at least one hydrogen atom has been replaced by a deuterium atom, in combination with at least one pharmaceutically acceptable carrier and/or auxiliary substance.

In yet another aspect, the invention relates to a compound of formula I or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for use as a medicament.

In yet another aspect, the invention relates to a compound of formula I or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for the treatment of disorders which responds to the modulation of the 5-HT_{2c} receptor.

In yet another aspect, the invention relates to the use of a compound of formula I or of an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of disorders which respond to the modulation of the 5-HT_{2c} receptor.

In yet another aspect, the invention relates to the use of a compound of formula I or of an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of disorders selected from the group consisting of damage of the central nervous system, disorders of the central nervous system, eating disorders, ocular hypertension, cardiovascular disorders, gastrointestinal disorders and diabetes, and especially from the group consisting of bipolar disorder, depression, atypical depression, mood episodes, adjustment disorders, anxiety, panic disorders, post-traumatic syndrome, psychoses, schizophrenia, cognitive deficits of schizophrenia, memory loss, dementia of aging, Alzheimer's disease, behavioral disorders associated with dementia, social phobia, mental disorders in childhood, attention deficit hyperactivity disorder, organic mental disorders, autism, mutism, disruptive behavior disorder, impulse control disorder, borderline personality disorder, obsessive compulsive disorder, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, seizure disorders, epilepsy, substance use disorders, alcohol abuse, cocaine abuse, tobacco abuse, smoking cessation, sexual dysfunction/erectile dysfunction in males, sexual dysfunction in females, premenstrual syndrome, late luteal phase syndrome, chronic fatigue syndrome, sleep disorders, sleep apnoea, chronic fatigue syndrome, psoriasis, Parkinson's disease, spinal cord injury, trauma, stroke, pain, bladder dysfunction/urinary incontinence, encephalitis, meningitis, eating disorders, obesity, bulimia, weight loss, anorexia nervosa, ocular hypertension, cardiovascular disorders, gastrointestinal disorders, diabetes insipidus, diabetes mellitus, type I diabetes, type II diabetes, type III diabetes, diabetes secondary to pancreatic diseases, diabetes related to steroid use, diabetes complications, hyperglycemia and insulin resistance.

In yet another aspect, the invention relates to a method for treating disorders which respond to the modulation of the 5-HT_{2c} receptor, which method comprises administering to a subject in need thereof at least one compound of formula I or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof.

In yet another aspect, the invention relates to a method for treating disorders selected from the group consisting of damage of the central nervous system, disorders of the central nervous system, eating disorders, ocular hypertension, cardiovascular disorders, gastrointestinal disorders and diabetes, and especially from the group consisting of

bipolar disorder, depression, atypical depression, mood episodes, adjustment disorders, anxiety, panic disorders, post-traumatic syndrome, psychoses, schizophrenia, cognitive deficits of schizophrenia, memory loss, dementia of aging, Alzheimer's disease, behavioral disorders associated with dementia, social phobia, mental disorders in childhood, attention deficit hyperactivity disorder, organic mental disorders, autism, mutism, disruptive behavior disorder, impulse control disorder, borderline personality disorder, obsessive compulsive disorder, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, seizure disorders, epilepsy, substance use disorders, alcohol abuse, cocaine abuse, tobacco abuse, smoking cessation, sexual dysfunction/erectile dysfunction in males, sexual dysfunction in females, premenstrual syndrome, late luteal phase syndrome, chronic fatigue syndrome, sleep disorders, sleep apnoea, chronic fatigue syndrome, psoriasis, Parkinson's disease, spinal cord injury, trauma, stroke, pain, bladder dysfunction/urinary incontinence, encephalitis, meningitis, eating disorders, obesity, bulimia, weight loss, anorexia nervosa, ocular hypertension, cardiovascular disorders, gastrointestinal disorders, diabetes insipidus, diabetes mellitus, type I diabetes, type II diabetes, type III diabetes, diabetes secondary to pancreatic diseases, diabetes related to steroid use, diabetes complications, hyperglycemia and insulin resistance, which method comprises administering to a subject in need thereof at least one compound of formula I or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof.


DETAILED DESCRIPTION

The compounds of the formula I may exist in different spatial arrangements. For example, if the compounds possess one or more centers of asymmetry, polysubstituted rings or double bonds, or as different tautomers, the present invention contemplates the possible use of enantiomeric mixtures, in particular racemates, diastereomeric mixtures and tautomeric mixtures, such as the respective essentially pure enantiomers, diastereomers and tautomers of the compounds of formula I and/or their salts.

It is likewise possible to use physiologically tolerated salts of the compounds of the formula I, especially acid addition salts with physiologically tolerated acids. Examples of suitable physiologically tolerated organic and inorganic acids are hydrochloric

acid, hydrobromic acid, phosphoric acid, sulfuric acid, acetic acid, trifluoroacetic acid, C₁-C₄-alkylsulfonic acids, such as methanesulfonic acid, aromatic sulfonic acids, such as benzenesulfonic acid and toluenesulfonic acid, oxalic acid, maleic acid, fumaric acid, lactic acid, tartaric acid, adipic acid and benzoic acid. Other utilizable acids are described in Fortschritte der Arzneimittelforschung [Advances in drug research], Volume 10, pages 224 et seq., Birkhäuser Verlag, Basel and Stuttgart, 1966.

The compounds of formula I may also be present in the form of tautomers. In one aspect, tautomerism may be present in compounds I wherein R^{4a} and R^{4b} form together a group =O and R^{5a} or R^{5b} is H. For example, the compounds of formula I may have the following tautomeric formulae:



In another aspect, tautomerism may be present in compounds I containing rings which have one or more C=O groups as ring members which are neighboured to a CH₂ group.

The organic moieties mentioned in the above definitions of the variables are, like the term halogen, collective terms for individual listings of the individual group members. The prefix C_n-C_m indicates in each case the possible number of carbon atoms in the group.

The term "halogen" denotes in each case fluorine, bromine, chlorine or iodine. In one aspect, the halogen may be fluorine, chlorine or bromine.

The term "alkyl" as used herein and in the alkyl moieties of alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylcarbonyl, alkoxycarbonyl and the like refers to saturated straight-chain or branched hydrocarbon radicals having 1 to 2 ("C₁-C₂-alkyl"), 1 to 3 ("C₁-C₃-alkyl"), 1 to 4 ("C₁-C₄-alkyl") or 1 to 6 ("C₁-C₆-alkyl") carbon atoms. C₁-C₂-

Alkyl is methyl or ethyl. C₁-C₃-Alkyl is additionally propyl and isopropyl. C₁-C₄-Alkyl is additionally butyl, 1-methylpropyl (sec-butyl), 2-methylpropyl (isobutyl) or 1,1-dimethylethyl (tert-butyl). C₁-C₆-Alkyl is additionally also, for example, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, or 1-ethyl-2-methylpropyl.

The term "fluorinated alkyl" as used herein refers to straight-chain or branched alkyl groups having 1 to 2 ("fluorinated C₁-C₂-alkyl"), 1 to 3 ("fluorinated C₁-C₃-alkyl"), 1 to 4 ("fluorinated C₁-C₄-alkyl") or 1 to 6 ("fluorinated C₁-C₆-alkyl") carbon atoms (as mentioned above), where some or all of the hydrogen atoms in these groups are replaced by fluorine atoms. Fluorinated C₁-C₂-alkyl is an alkyl group having 1 or 2 carbon atoms (as mentioned above), where at least one of the hydrogen atoms, e.g. 1, 2, 3, 4 or 5 hydrogen atoms in these groups are replaced by fluorine atoms, such as difluoromethyl, trifluoromethyl, 1-fluoroethyl, (R)-1-fluoroethyl, (S)-1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, or pentafluoroethyl. Fluorinated C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms (as mentioned above), where at least one of the hydrogen atoms, e.g. 1, 2, 3, 4 or 5 hydrogen atoms in these groups are replaced by fluorine atoms. Examples are, apart those listed above for C₁-C₂-fluoroalkyl, 1-fluoropropyl, (R)-1-fluoropropyl, (S)-1-fluoropropyl, 2-fluoropropyl, (R)-2-fluoropropyl, (S)-2-fluoropropyl, 3-fluoropropyl, 1,1-difluoropropyl, 2,2-difluoropropyl, 1,2-difluoropropyl, 2,3-difluoropropyl, 1,3-difluoropropyl, 3,3-difluoropropyl, 1,1,2-trifluoropropyl, 1,2,2-trifluoropropyl, 1,2,3-trifluoropropyl, 2,2,3-trifluoropropyl, 3,3,3-trifluoropropyl, 1,1,1-trifluoroprop-2-yl, 2-fluoro-1-methylethyl, (R)-2-fluoro-1-methylethyl, (S)-2-fluoro-1-methylethyl, 2,2-difluoro-1-methylethyl, (R)-2,2-difluoro-1-methylethyl, (S)-2,2-difluoro-1-methylethyl, 1,2-difluoro-1-methylethyl, (R)-1,2-difluoro-1-methylethyl, (S)-1,2-difluoro-1-methylethyl, 2,2,2-trifluoro-1-methylethyl, (R)-2,2,2-trifluoro-1-methylethyl, (S)-2,2,2-trifluoro-1-methylethyl, 2-fluoro-1-(fluoromethyl)ethyl, 1-(difluoromethyl)-2,2-difluoroethyl, 1-(trifluoromethyl)-2,2,2-trifluoroethyl, 1-(trifluoromethyl)-1,2,2,2-

tetrafluoroethyl, 1-fluorobutyl, (R)-1-fluorobutyl, (S)-1-fluorobutyl, 2-fluorobutyl, (R)-2-fluorobutyl, (S)-2-fluorobutyl, 3-fluorobutyl, (R)-3-fluorobutyl, (S)-3-fluorobutyl, 4-fluorobutyl, 1,1-difluorobutyl, 2,2-difluorobutyl, 3,3-difluorobutyl, 4,4-difluorobutyl, 4,4,4-trifluorobutyl and the like. Fluorinated C₁-C₆-alkyl is a straight-chain or branched alkyl group having 1 to 6 carbon atoms (as mentioned above), where at least one of the hydrogen atoms, e.g. 1, 2, 3, 4 or 5 hydrogen atoms in these groups are replaced by fluorine atoms. Additionally examples include for C₁-C₄-fluoroalkyl, 1-fluoropentyl, (R)-1-fluoropentyl, (S)-1-fluoropentyl, 2-fluoropentyl, (R)-2-fluoropentyl, (S)-2-fluoropentyl, 3-fluoropentyl, (R)-3-fluoropentyl, (S)-3-fluoropentyl, 4-fluoropentyl, (R)-4-fluoropentyl, (S)-4-fluoropentyl, 5-fluoropentyl, (R)-5-fluoropentyl, (S)-5-fluoropentyl, 1-fluorohexyl, (R)-1-fluorohexyl, (S)-1-fluorohexyl, 2-fluorohexyl, (R)-2-fluorohexyl, (S)-2-fluorohexyl, 3-fluorohexyl, (R)-3-fluorohexyl, (S)-3-fluorohexyl, 4-fluorohexyl, (R)-4-fluorohexyl, (S)-4-fluorohexyl, 5-fluorohexyl, (R)-5-fluorohexyl, (S)-5-fluorohexyl, 6-fluorohexyl, (R)-6-fluorohexyl, (S)-6-fluorohexyl, and the like.

The term "alkenyl" as used herein refers to monounsaturated straight-chain or branched hydrocarbon radicals having 2 to 3 ("C₂-C₃-alkenyl"), 2 to 4 ("C₂-C₄-alkenyl") or 2 to 6 ("C₂-C₆-alkenyl") carbon atoms and a double bond in any position, such as C₂-C₃-alkenyl, such as ethenyl, 1-propenyl, 2-propenyl or 1-methylethenyl; C₂-C₄-alkenyl, such as ethenyl, 1-propenyl, 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl or 2-methyl-2-propenyl; and C₂-C₆-alkenyl, such as ethenyl, 1-propenyl, 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-1-butenyl, 2-methyl-1-butenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-1-propenyl, 1-ethyl-2-propenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-1-pentenyl, 2-methyl-1-pentenyl, 3-methyl-1-pentenyl, 4-methyl-1-pentenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-

butenyl, 1,2-dimethyl-1-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-1-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-1-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 3,3-dimethyl-1-butenyl, 3,3-dimethyl-2-butenyl, 1-ethyl-1-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-1-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl, 1-ethyl-2-methyl-1-propenyl, 1-ethyl-2-methyl-2-propenyl and the like.

The term "fluorinated alkenyl" as used herein refers to unsaturated straight-chain or branched hydrocarbon radicals having 2 to 3 ("fluorinated C₂-C₃-alkenyl"), 2 to 4 ("fluorinated C₂-C₄-alkenyl") or 2 to 6 ("fluorinated C₂-C₆-alkenyl") carbon atoms and a double bond in any position (as mentioned above), where some or all of the hydrogen atoms in these groups are replaced by fluorine atoms, such as, fluorovinyl, fluoroallyl and the like.

The term "alkynyl" as used herein refers to straight-chain or branched hydrocarbon groups having 2 to 3 ("C₂-C₃-alkynyl"), 2 to 4 ("C₂-C₄-alkynyl") or 2 to 6 ("C₂-C₆-alkynyl") carbon atoms and one or two triple bonds in any position, such as C₂-C₃-alkynyl, such as ethynyl, 1-propynyl or 2-propynyl; C₂-C₄-alkynyl, such as ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 1-methyl-2-propynyl and the like, and C₂-C₆-alkynyl, such as ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 1-methyl-2-propynyl, 1-pentyne, 2-pentyne, 3-pentyne, 4-pentyne, 1-methyl-2-butyne, 1-methyl-3-butyne, 2-methyl-3-butyne, 3-methyl-1-butyne, 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 1-methyl-2-pentyne, 1-methyl-3-pentyne, 1-methyl-4-pentyne, 2-methyl-3-pentyne, 2-methyl-4-pentyne, 3-methyl-1-pentyne, 3-methyl-4-pentyne, 4-methyl-1-pentyne, 4-methyl-2-pentyne, 1,1-dimethyl-2-butyne, 1,1-dimethyl-3-butyne, 1,2-dimethyl-3-butyne, 2,2-dimethyl-3-butyne, 3,3-dimethyl-1-butyne, 1-ethyl-2-butyne, 1-ethyl-3-butyne, 2-ethyl-3-butyne, 1-ethyl-1-methyl-2-propynyl and the like.

The term "fluorinated alkynyl" as used herein refers to unsaturated straight-chain or branched hydrocarbon radicals having 2 to 3 ("fluorinated C₂-C₃-alkynyl"), 3 to 4 ("fluorinated C₃-C₄-alkynyl") or 2 to 6 ("fluorinated C₂-C₆-alkynyl") carbon atoms and

one or two triple bonds in any position (as mentioned above), where some or all of the hydrogen atoms in these groups are replaced by fluorine atoms.

The term "cycloalkyl" as used herein refers to mono- or bicyclic saturated hydrocarbon radicals having 3 to 8 ("C₃-C₈-cycloalkyl"), in particular 3 to 6 carbon atoms ("C₃-C₆-cycloalkyl") or 3 to 5 carbon atoms ("C₃-C₅-cycloalkyl") or 3 or 4 carbon atoms ("C₃-C₄-cycloalkyl"). In one aspect, C₃-C₄-cycloalkyl, C₃-C₅-cycloalkyl and C₃-C₆-cycloalkyl are monocyclic. Examples for C₃-C₄-cycloalkyl are cyclopropyl and cyclobutyl. Examples of monocyclic radicals having 3 to 5 carbon atoms are cyclopropyl, cyclobutyl and cyclopentyl. Examples of monocyclic radicals having 3 to 6 carbon atoms are cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. Examples of monocyclic radicals having 3 to 8 carbon atoms are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl. Examples of bicyclic radicals having 7 or 8 carbon atoms comprise bicyclo[2.2.1]heptyl, bicyclo[3.1.1]heptyl, bicyclo[2.2.2]octyl and bicyclo[3.2.1]octyl. In one aspect, the term cycloalkyl denotes a monocyclic saturated hydrocarbon radical.

The term "fluorinated cycloalkyl" as used herein refers to mono- or bicyclic saturated hydrocarbon groups having 3 to 8 ("C₃-C₈-halocycloalkyl") or preferably 3 to 6 ("C₃-C₆-halocycloalkyl") or 3 to 5 ("C₃-C₅-halocycloalkyl") carbon ring members (as mentioned above) in which some or all of the hydrogen atoms are replaced by fluorine atoms. Examples include 1-fluorocyclopropyl, 2-fluorocyclopropyl, (S)- and (R)-2,2-difluorocyclopropyl, 1,2-difluorocyclopropyl, 2,3-difluorocyclopropyl, pentafluorocyclopropyl, 1-fluorocyclobutyl, 2-fluorocyclobutyl, 3-fluorocyclobutyl, 2,2-difluorocyclobutyl, 3,3-difluorocyclobutyl, 1,2-difluorocyclobutyl, 1,3-difluorocyclobutyl, 2,3-difluorocyclobutyl, 2,4-difluorocyclobutyl, 1,2,2-trifluorocyclobutyl, 1-fluorocycloheptyl, 2-fluorocycloheptyl, 3-fluorocycloheptyl, 4-fluorocycloheptyl, 1,2-difluorocycloheptyl, 1,3-difluorocycloheptyl, 1,4-difluorocycloheptyl, 2,2-difluorocycloheptyl, 2,3-difluorocycloheptyl, 2,4-difluorocycloheptyl, 2,5-difluorocycloheptyl, 2,6-difluorocycloheptyl, 2,7-difluorocycloheptyl, 3,3-difluorocycloheptyl, 3,4-difluorocycloheptyl, 3,5-difluorocycloheptyl, 3,6-difluorocycloheptyl, 4,4-difluorocycloheptyl, 4,5-difluorocycloheptyl, and the like.

The term "cycloalkenyl" as used herein refers to monocyclic partially unsaturated, non-aromatic hydrocarbon radicals having 3 to 8 ("C₃-C₈-cycloalkenyl"), in particular 5

to 7 carbon atoms ("C₅-C₇-cycloalkenyl") or 5 or 6 carbon atoms ("C₅-C₆-cycloalkenyl") and one or more non-cumulative, preferably one, C-C double bonds in the ring. Examples for C₅-C₆-cycloalkenyl are cyclopent-1-en-1-yl, cyclopent-1-en-3-yl, cyclopent-1-en-4-yl, cyclopenta-1,3-dien-1-yl, cyclopenta-1,3-dien-2-yl, cyclopenta-1,3-dien-5-yl, cyclohex-1-en-1-yl, cyclohex-1-en-3-yl, cyclohex-1-en-4-yl, cyclohexa-1,3-dien-1-yl, cyclohexa-1,3-dien-2-yl, cyclohexa-1,3-dien-5-yl, cyclohexa-1,4-dien-1-yl and cyclohexa-1,4-dien-3-yl. Examples of C₅-C₇-cycloalkenyl are, apart those mentioned above, include for C₅-C₆-cycloalkenyl, cyclohept-1-en-1-yl, cyclohept-1-en-3-yl, cyclohept-1-en-4-yl, cyclohept-1-en-5-yl, cyclohepta-1,3-dien-1-yl, cyclohepta-1,3-dien-2-yl, cyclohepta-1,3-dien-5-yl, cyclohepta-1,3-dien-6-yl, cyclohepta-1,4-dien-1-yl, cyclohepta-1,4-dien-2-yl, cyclohepta-1,4-dien-3-yl and cyclohepta-1,4-dien-6-yl. Examples of C₃-C₈-cycloalkenyl are, apart those mentioned above for C₅-C₆-cycloalkenyl, cycloprop-1-en-1-yl, cycloprop-1-en-3-yl, cyclobut-1-en-1-yl, cyclobut-1-en-3-yl, cyclooct-1-en-1-yl, cyclooct-1-en-3-yl, cyclooct-1-en-4-yl, cyclooct-1-en-5-yl, cycloocta-1,3-dien-1-yl, cycloocta-1,3-dien-2-yl, cycloocta-1,3-dien-5-yl, cycloocta-1,3-dien-6-yl, cycloocta-1,4-dien-1-yl, cycloocta-1,4-dien-2-yl, cycloocta-1,4-dien-3-yl, cycloocta-1,4-dien-6-yl, cycloocta-1,4-dien-7-yl, cycloocta-1,5-dien-1-yl, and cycloocta-1,5-dien-3-yl.

The term "fluorinated cycloalkenyl" as used herein refers to monocyclic partially unsaturated, non-aromatic hydrocarbon radicals having 3 to 8 ("fluorinated C₃-C₈-cycloalkenyl"), in particular 5 to 7 carbon atoms ("fluorinated C₅-C₇-cycloalkenyl") or 5 or 6 carbon atoms ("fluorinated C₅-C₆-cycloalkenyl") and one or more non-cumulative, preferably one, C-C double bonds in the ring and in which some or all of the hydrogen atoms are replaced by fluorine atoms.

The term "cycloalkyl-C₁-C₄-alkyl" refers to a C₃-C₈-cycloalkyl group ("C₃-C₈-cycloalkyl-C₁-C₄-alkyl"), preferably a C₃-C₆-cycloalkyl group ("C₃-C₆-cycloalkyl-C₁-C₄-alkyl") as defined above which is bound to the remainder of the molecule via a C₁-C₄-alkyl group, as defined above. The term "cycloalkyl-C₁-C₂-alkyl" refers to a C₃-C₈-cycloalkyl group ("C₃-C₈-cycloalkyl-C₁-C₂-alkyl"), preferably a C₃-C₆-cycloalkyl group ("C₃-C₆-cycloalkyl-C₁-C₂-alkyl") as defined above which is bound to the remainder of the molecule via a C₁-C₂-alkyl group, as defined above. Examples for C₃-C₆-cycloalkyl-C₁-C₂-alkyl are cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclobutylethyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl and cyclohexylethyl. Ex-

amples for C₃-C₆-cycloalkyl-C₁-C₄-alkyl, apart those mentioned for C₃-C₆-cycloalkyl-C₁-C₂-alkyl, are cyclopropylpropyl, cyclopropylbutyl, cyclobutylpropyl, cyclobutylbutyl, cyclopentylpropyl, cyclopentylbutyl, cyclohexylpropyl and cyclohexylbutyl. Examples for C₃-C₈-cycloalkyl-C₁-C₂-alkyl, apart those mentioned for C₃-C₆-cycloalkyl-C₁-C₂-alkyl, are cycloheptylmethyl, cycloheptylethyl, cyclooctylmethyl and cyclooctylethyl. Examples for C₃-C₈-cycloalkyl-C₁-C₄-alkyl, apart those mentioned for C₃-C₈-cycloalkyl-C₁-C₂-alkyl, are cycloheptylpropyl, cycloheptylbutyl, cyclooctylpropyl and cyclooctylbutyl.

The term "fluorinated cycloalkyl-C₁-C₄-alkyl" refers to a fluorinated C₃-C₈-cycloalkyl group ("fluorinated C₃-C₈-cycloalkyl-C₁-C₄-alkyl"), preferably a fluorinated C₃-C₆-cycloalkyl group ("fluorinated C₃-C₆-cycloalkyl-C₁-C₄-alkyl") as defined above which is bound to the remainder of the molecule via a C₁-C₄-alkyl group, as defined above.

The term "fluorinated cycloalkyl-C₁-C₂-alkyl" refers to a fluorinated C₃-C₈-cycloalkyl group ("fluorinated C₃-C₈-cycloalkyl-C₁-C₂-alkyl"), preferably a fluorinated C₃-C₆-cycloalkyl group ("fluorinated C₃-C₆-cycloalkyl-C₁-C₂-alkyl") as defined above which is bound to the remainder of the molecule via a C₁-C₂-alkyl group, as defined above.

The term "C₁-C₂-alkoxy" is a C₁-C₂-alkyl group, as defined above, attached via an oxygen atom. The term "C₁-C₃-alkoxy" is a C₁-C₃-alkyl group, as defined above, attached via an oxygen atom. The term "C₁-C₄-alkoxy" is a C₁-C₄-alkyl group, as defined above, attached via an oxygen atom. The term "C₁-C₆-alkoxy" is a C₁-C₆-alkyl group, as defined above, attached via an oxygen atom. C₁-C₂-Alkoxy is methoxy or ethoxy. C₁-C₃-Alkoxy is additionally, for example, n-propoxy and 1-methylethoxy (isopropoxy). C₁-C₄-Alkoxy is additionally, for example, butoxy, 1-methylpropoxy (sec-butoxy), 2-methylpropoxy (isobutoxy) or 1,1-dimethylethoxy (tert-butoxy). C₁-C₆-Alkoxy is additionally, for example, pentoxy, 1-methylbutoxy, 2-methylbutoxy, 3-methylbutoxy, 1,1-dimethylpropoxy, 1,2-dimethylpropoxy, 2,2-dimethylpropoxy, 1-ethylpropoxy, hexoxy, 1-methylpentoxy, 2-methylpentoxy, 3-methylpentoxy, 4-methylpentoxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy, 1,3-dimethylbutoxy, 2,2-dimethylbutoxy, 2,3-dimethylbutoxy, 3,3-dimethylbutoxy, 1-ethylbutoxy, 2-ethylbutoxy, 1,1,2-trimethylpropoxy, 1,2,2-trimethylpropoxy, 1-ethyl-1-methylpropoxy or 1-ethyl-2-

methylpropoxy. C₁-C₈-Alkoxy is additionally, for example, heptyloxy, octyloxy, 2-ethylhexyloxy and positional isomers thereof.

The term "fluorinated C₁-C₂-alkoxy" is a fluorinated C₁-C₂-alkyl group, as defined above, attached via an oxygen atom. The term "fluorinated C₁-C₃-alkoxy" is a fluorinated C₁-C₃-alkyl group, as defined above, attached via an oxygen atom. The term "fluorinated C₁-C₆-haloalkoxy" is a fluorinated C₁-C₆-alkyl group, as defined above, attached via an oxygen atom. Fluorinated C₁-C₂-alkoxy is, for example, OCH₂F, OCHF₂, OCF₃, 1-fluoroethoxy, (R)-1-fluoroethoxy, (S)-1-fluoroethoxy, 2-fluoroethoxy, 1,1-difluoroethoxy, 1,2-difluoroethoxy, 2,2-difluoroethoxy, 1,1,2-trifluoroethoxy, 1,2,2-trifluoroethoxy, 2,2,2-trifluoroethoxy, 1,1,2,2-tetrafluoroethoxy or OC₂F₅. Fluorinated C₁-C₃-alkoxy is additionally, for example, 1-fluoropropoxy, (R)-1-fluoropropoxy, (S)-1-fluoropropoxy, 2-fluoropropoxy, (R)-2-fluoropropoxy, (S)-2-fluoropropoxy, 3-fluoropropoxy, 1,1-difluoropropoxy, 2,2-difluoropropoxy, 2,3-difluoropropoxy, 3,3-difluoropropoxy, 3,3,3-trifluoropropoxy, (R)-2-fluoro-1-methylethoxy, (S)-2-fluoro-1-methylethoxy, (R)-2,2-difluoro-1-methylethoxy, (S)-2,2-difluoro-1-methylethoxy, (R)-1,2-difluoro-1-methylethoxy, (S)-1,2-difluoro-1-methylethoxy, (R)-2,2,2-trifluoro-1-methylethoxy, (S)-2,2,2-trifluoro-1-methylethoxy, 2-fluoro-1-(fluoromethyl)ethoxy, 1-(difluoromethyl)-2,2-difluoroethoxy, OCH₂-C₂F₅, OCF₂-C₂F₅ or 1-(CH₂F)-2-fluoroethoxy. Fluorinated C₁-C₄-alkoxy is additionally, for example, 1-fluorobutoxy, (R)-1-fluorobutoxy, (S)-1-fluorobutoxy, 2-fluorobutoxy, 3-fluorobutoxy, 4-fluorobutoxy, 1,1-difluorobutoxy, 2,2-difluorobutoxy, 3,3-difluorobutoxy, 4,4-difluorobutoxy, 4,4,4-trifluorobutoxy or nonafluorobutoxy. Fluorinated C₁-C₆-alkoxy is additionally, for example, 5-fluoropentoxy, undecafluoropentoxy, 6-fluorohexoxy or tridecafluorohexoxy.

The term "C₁-C₄-alkoxy-C₁-C₄-alkyl" as used herein, refers to a straight-chain or branched alkyl group having 1 to 4 carbon atoms, as defined above, where one hydrogen atom is replaced by a C₁-C₄-alkoxy group, as defined above. The term "C₁-C₆-alkoxy-C₁-C₄-alkyl" as used herein, refers to a straight-chain or branched alkyl group having 1 to 4 carbon atoms, as defined above, where one hydrogen atom is replaced by a C₁-C₆-alkoxy group, as defined above. Examples are methoxymethyl, ethoxymethyl, propoxymethyl, isopropoxymethyl, n-butoxymethyl, sec-butoxymethyl, isobutoxymethyl, tert-butoxymethyl, 1-methoxyethyl, 1-ethoxyethyl, 1-propoxyethyl,

1-isopropoxyethyl, 1-n-butoxyethyl, 1-sec-butoxyethyl, 1-isobutoxyethyl, 1-tert-butoxyethyl, 2-methoxyethyl, 2-ethoxyethyl, 2-propoxyethyl, 2-isopropoxyethyl, 2-n-butoxyethyl, 2-sec-butoxyethyl, 2-isobutoxyethyl, 2-tert-butoxyethyl, 1-methoxypropyl, 1-ethoxypropyl, 1-propoxypropyl, 1-isopropoxypropyl, 1-n-butoxypropyl, 1-sec-butoxypropyl, 1-isobutoxypropyl, 1-tert-butoxypropyl, 2-methoxypropyl, 2-ethoxypropyl, 2-propoxypropyl, 2-isopropoxypropyl, 2-n-butoxypropyl, 2-sec-butoxypropyl, 2-isobutoxypropyl, 2-tert-butoxypropyl, 3-methoxypropyl, 3-ethoxypropyl, 3-propoxypropyl, 3-isopropoxypropyl, 3-n-butoxypropyl, 3-sec-butoxypropyl, 3-isobutoxypropyl, 3-tert-butoxypropyl and the like.

The term "hydroxy-C₁-C₄-alkyl" as used herein, refers to a straight-chain or branched alkyl group having 1 to 4 carbon atoms, as defined above, where one hydrogen atom is replaced by a hydroxy group. The term "hydroxy-C₁-C₆-alkyl" as used herein, refers to a straight-chain or branched alkyl group having 1 to 6 carbon atoms, as defined above, where one hydrogen atom is replaced by a hydroxy group. Examples for hydroxy-C₁-C₄-alkyl include hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 1-hydroxypropyl, 2-hydroxypropyl, 3-hydroxypropyl, 1-hydroxyprop-2-yl, 2-hydroxyprop-2-yl, 1-hydroxybutyl, 2-hydroxybutyl, 3-hydroxybutyl, 4-hydroxybutyl, 1-hydroxybut-2-yl, 2-hydroxybut-2-yl, 3-hydroxybut-2-yl, 4-hydroxybut-2-yl, hydroxy-tert-butyl and the like. Examples for hydroxy-C₁-C₆-alkyl are, apart those mentioned for hydroxy-C₁-C₄-alkyl, include, 1-hydroxypentyl, 2-hydroxypentyl, 3-hydroxypentyl, 4-hydroxypentyl, 5-hydroxypentyl, 1-hydroxyhexyl, 2-hydroxyhexyl, 3-hydroxyhexyl, 4-hydroxyhexyl, 5-hydroxyhexyl, 6-hydroxyhexyl and the like.

The term "hydroxy-C₁-C₄-alkoxy" as used herein, refers to a C₁-C₄-alkoxy group, as defined above, where one hydrogen atom is replaced by a hydroxy group. The term "hydroxy-C₁-C₆-alkoxy" as used herein, refers to a C₁-C₆-alkoxy group, as defined above, where one hydrogen atom is replaced by a hydroxy group. Examples for hydroxy-C₁-C₄-alkoxy include hydroxymethoxy, 1-hydroxyethoxy, 2-hydroxyethoxy, 1-hydroxypropoxy, 2-hydroxypropoxy, 3-hydroxypropoxy, 1-hydroxy-2-propoxy, 2-hydroxy-2-propoxy, 1-hydroxybutoxy, 2-hydroxybutoxy, 3-hydroxybutoxy, 4-hydroxybutoxy, 1-hydroxy-2-butoxy, 2-hydroxy-2-butoxy, 3-hydroxy-2-butoxy, 4-hydroxy-2-butoxy, hydroxy-tert-butoxy and the like. Examples for hydroxy-C₁-C₆-alkoxy include, apart those mentioned for hydroxy-C₁-C₄-alkoxy, 1-hydroxypentoxy, 2-

hydroxypentoxy, 3-hydroxypentoxy, 4-hydroxypentoxy, 5-hydroxypentoxy, 1-hydroxyhexoxy, 2-hydroxyhexoxy, 3-hydroxyhexoxy, 4-hydroxyhexoxy, 5-hydroxyhexoxy, 6-hydroxyhexoxy and the like.

The term "C₁-C₄-alkoxy-C₁-C₄-alkoxy" as used herein, refers to a C₁-C₄-alkoxy group, as defined above, where one hydrogen atom is replaced by a C₁-C₄-alkoxy group, as defined above. The term "C₁-C₆-alkoxy-C₁-C₄-alkoxy" as used herein, refers to a C₁-C₄-alkoxy group, as defined above, where one hydrogen atom is replaced by a C₁-C₆-alkoxy group, as defined above. Examples are methoxymethoxy, ethoxymethoxy, propoxymethoxy, isopropoxymethoxy, butoxymethoxy, sec-butoxymethoxy, isobutoxymethoxy, tert-butoxymethoxy, 1-methoxyethoxy, 1-ethoxyethoxy, 1-propoxyethoxy, 1-isopropoxyethoxy, 1-butoxyethoxy, 1-sec-butoxyethoxy, 1-isobutoxyethoxy, 1-tert-butoxyethoxy, 2-methoxyethoxy, 2-ethoxyethoxy, 2-propoxyethoxy, 2-isopropoxyethoxy, 2-butoxyethoxy, 2-sec-butoxyethoxy, 2-isobutoxyethoxy, 2-tert-butoxyethoxy, 1-methoxypropoxy, 1-ethoxypropoxy, 1-propoxypropoxy, 1-isopropoxypropoxy, 1-butoxypropoxy, 1-sec-butoxypropoxy, 1-isobutoxypropoxy, 1-tert-butoxypropoxy, 2-methoxypropoxy, 2-ethoxypropoxy, 2-propoxypropoxy, 2-isopropoxypropoxy, 2-butoxypropoxy, 2-sec-butoxypropoxy, 2-isobutoxypropoxy, 2-tert-butoxypropoxy, 3-methoxypropoxy, 3-ethoxypropoxy, 3-propoxypropoxy, 3-isopropoxypropoxy, 3-butoxypropoxy, 3-sec-butoxypropoxy, 3-isobutoxypropoxy, 3-tert-butoxypropoxy and the like.

The term "C₁-C₂-alkylthio" is a C₁-C₂-alkyl group, as defined above, attached via a sulfur atom. The term "C₁-C₃-alkylthio" refers to a C₁-C₃-alkyl group, as defined above, attached via a sulfur atom. The term "C₁-C₄-alkylthio" is a C₁-C₄-alkyl group, as defined above, attached via a sulfur atom. The term "C₁-C₆-alkylthio" refers to a C₁-C₆-alkyl group, as defined above, attached via a sulfur atom. The term "C₁-C₁₀-alkylthio" refers to a C₁-C₁₀-alkyl group, as defined above, attached via a sulfur atom. C₁-C₂-Alkylthio is methylthio or ethylthio. C₁-C₃-Alkylthio is additionally, for example, n-propylthio or 1-methylethylthio (isopropylthio). C₁-C₄-Alkylthio is additionally, for example, butylthio, 1-methylpropylthio (sec-butylthio), 2-methylpropylthio (isobutylthio) or 1,1-dimethylethylthio (tert-butylthio). C₁-C₆-Alkylthio is additionally, for example, pentylthio, 1-methylbutylthio, 2-methylbutylthio, 3-methylbutylthio, 1,1-dimethylpropylthio, 1,2-dimethylpropylthio, 2,2-dimethylpropylthio, 1-ethylpropylthio,

hexylthio, 1-methylpentylthio, 2-methylpentylthio, 3-methylpentylthio, 4-methylpentylthio, 1,1-dimethylbutylthio, 1,2-dimethylbutylthio, 1,3-dimethylbutylthio, 2,2-dimethylbutylthio, 2,3-dimethylbutylthio, 3,3-dimethylbutylthio, 1-ethylbutylthio, 2-ethylbutylthio, 1,1,2-trimethylpropylthio, 1,2,2-trimethylpropylthio, 1-ethyl-1-methylpropylthio or 1-ethyl-2-methylpropylthio.

The term "fluorinated C₁-C₂-alkylthio" refers to a fluorinated C₁-C₂-alkyl group, as defined above, attached via a sulfur atom. The term "fluorinated C₁-C₃-alkylthio" refers to a fluorinated C₁-C₃-alkyl group, as defined above, attached via a sulfur atom. The term "fluorinated C₁-C₄-alkylthio" refers to a fluorinated C₁-C₄-alkyl group, as defined above, attached via a sulfur atom. The term "fluorinated C₁-C₆-alkylthio" refers to a fluorinated C₁-C₆-alkyl group, as defined above, attached via a sulfur atom. Fluorinated C₁-C₂-alkylthio refers to, for example, SCH₂F, SCHF₂, SCF₃, 2-fluoroethylthio, 2,2-difluoroethylthio, 2,2,2-trifluoroethylthio, or SC₂F₅. Fluorinated C₁-C₃-alkylthio may additionally, for example, include 2-fluoropropylthio, 3-fluoropropylthio, 2,2-difluoropropylthio, 2,3-difluoropropylthio, 3,3,3-trifluoropropylthio, SCH₂-C₂F₅, SCF₂-C₂F₅ or 1-(CH₂F)-2-fluoroethylthio. Fluorinated C₁-C₄-alkylthio may additionally, for example, include 4-fluorobutylthio or nonafluorobutylthio. Fluorinated C₁-C₆-alkylthio is additionally, for example, 5-fluoropentylthio, undecafluoropentylthio, 6-fluorohexylthio or dodecafluorohexylthio.

The term "C₁-C₂-alkylsulfinyl" refers to a C₁-C₂-alkyl group, as defined above, attached via a sulfinyl [S(O)] group. The term "C₁-C₄-alkylsulfinyl" is a C₁-C₄-alkyl group, as defined above, attached via a sulfinyl [S(O)] group. The term "C₁-C₆-alkylsulfinyl" is a C₁-C₆-alkyl group, as defined above, attached via a sulfinyl [S(O)] group. C₁-C₂-Alkylsulfinyl is methylsulfinyl or ethylsulfinyl. C₁-C₄-Alkylsulfinyl is additionally, for example, n-propylsulfinyl, 1-methylethylsulfinyl (isopropylsulfinyl), butylsulfinyl, 1-methylpropylsulfinyl (sec-butylsulfinyl), 2-methylpropylsulfinyl (isobutylsulfinyl) or 1,1-dimethylethylsulfinyl (tert-butylsulfinyl). C₁-C₆-Alkylsulfinyl is additionally, for example, pentylsulfinyl, 1-methylbutylsulfinyl, 2-methylbutylsulfinyl, 3-methylbutylsulfinyl, 1,1-dimethylpropylsulfinyl, 1,2-dimethylpropylsulfinyl, 2,2-dimethylpropylsulfinyl, 1-ethylpropylsulfinyl, hexylsulfinyl, 1-methylpentylsulfinyl, 2-methylpentylsulfinyl, 3-methylpentylsulfinyl, 4-methylpentylsulfinyl, 1,1-dimethylbutylsulfinyl, 1,2-dimethylbutylsulfinyl,

1,3-dimethylbutylsulfinyl, 2,2-dimethylbutylsulfinyl, 2,3-dimethylbutylsulfinyl, 3,3-dimethylbutylsulfinyl, 1-ethylbutylsulfinyl, 2-ethylbutylsulfinyl, 1,1,2-trimethylpropylsulfinyl, 1,2,2-trimethylpropylsulfinyl, 1-ethyl-1-methylpropylsulfinyl or 1-ethyl-2-methylpropylsulfinyl.

The term "fluorinated C₁-C₂-alkylsulfinyl" refers to a fluorinated C₁-C₂-alkyl group, as defined above, attached via a sulfinyl [S(O)] group. The term "fluorinated C₁-C₃-alkylsulfinyl" refers to a fluorinated C₁-C₃-alkyl group, as defined above, attached via a sulfinyl [S(O)] group. The term "fluorinated C₁-C₄-alkylsulfinyl" refers to a fluorinated C₁-C₄-alkyl group, as defined above, attached via a sulfinyl [S(O)] group. The term "fluorinated C₁-C₆-alkylsulfinyl" refers to a fluorinated C₁-C₆-alkyl group, as defined above, attached via a sulfinyl [S(O)] group. Fluorinated C₁-C₂-alkylsulfinyl is, for example, S(O)CH₂F, S(O)CHF₂, S(O)CF₃, 2-fluoroethylsulfinyl, 2,2-difluoroethylsulfinyl, 2,2,2-trifluoroethylsulfinyl, or S(O)C₂F₅. Fluorinated C₁-C₃-alkylsulfinyl may additionally, for example, include 2-fluoropropylsulfinyl, 3-fluoropropylsulfinyl, 2,2-difluoropropylsulfinyl, 2,3-difluoropropylsulfinyl, 3,3,3-trifluoropropylsulfinyl, S(O)CH₂-C₂F₅, S(O)CF₂-C₂F₅ or 1-(CH₂F)-2-fluoroethylsulfinyl. Fluorinated C₁-C₄-alkylsulfinyl may additionally, for example, include 4-fluorobutylsulfinyl or nonafluorobutylsulfinyl. Fluorinated C₁-C₆-alkylsulfinyl may additionally, for example, include 5-fluoropentylsulfinyl, undecafluoropentylsulfinyl, 6-fluorohexylsulfinyl or dodecafluorohexylsulfinyl.

The term "C₁-C₂-alkylsulfonyl" refers to a C₁-C₂-alkyl group, as defined above, attached via a sulfonyl [S(O)₂] group. The term "C₁-C₄-alkylsulfonyl" refers to a C₁-C₄-alkyl group, as defined above, attached via a sulfonyl [S(O)₂] group. The term "C₁-C₆-alkylsulfonyl" is a C₁-C₆-alkyl group, as defined above, attached via a sulfonyl [S(O)₂] group. C₁-C₂-Alkylsulfonyl refers to a methylsulfonyl or ethylsulfonyl. C₁-C₄-Alkylsulfonyl is additionally, for example, n-propylsulfonyl, 1-methylethylsulfonyl (isopropylsulfonyl), butylsulfonyl, 1-methylpropylsulfonyl (sec-butylsulfonyl), 2-methylpropylsulfonyl (isobutylsulfonyl) or 1,1-dimethylethylsulfonyl (tert-butylsulfonyl). C₁-C₆-Alkylsulfonyl is additionally, for example, pentylsulfonyl, 1-methylbutylsulfonyl, 2-methylbutylsulfonyl, 3-methylbutylsulfonyl, 1,1-dimethylpropylsulfonyl, 1,2-dimethylpropylsulfonyl, 2,2-dimethylpropylsulfonyl, 1-ethylpropylsulfonyl, hexylsulfonyl, 1-methylpentylsulfonyl, 2-methylpentylsulfonyl,

3-methylpentylsulfonyl, 4-methylpentylsulfonyl, 1,1-dimethylbutylsulfonyl, 1,2-dimethylbutylsulfonyl, 1,3-dimethylbutylsulfonyl, 2,2-dimethylbutylsulfonyl, 2,3-dimethylbutylsulfonyl, 3,3-dimethylbutylsulfonyl, 1-ethylbutylsulfonyl, 2-ethylbutylsulfonyl, 1,1,2-trimethylpropylsulfonyl, 1,2,2-trimethylpropylsulfonyl, 1-ethyl-1-methylpropylsulfonyl or 1-ethyl-2-methylpropylsulfonyl.

The term "fluorinated C₁-C₂-alkylsulfonyl" refers to a fluorinated C₁-C₂-alkyl group, as defined above, attached via a sulfonyl [S(O)₂] group. The term "fluorinated C₁-C₃-alkylsulfonyl" refers to a fluorinated C₁-C₃-alkyl group, as defined above, attached via a sulfonyl [S(O)₂] group. The term "fluorinated C₁-C₄-alkylsulfonyl" refers to a fluorinated C₁-C₄-alkyl group, as defined above, attached via a sulfonyl [S(O)₂] group. The term "fluorinated C₁-C₆-alkylsulfonyl" refers to a fluorinated C₁-C₆-alkyl group, as defined above, attached via a sulfonyl [S(O)₂] group. Fluorinated C₁-C₂-alkylsulfonyl is, for example, S(O)₂CH₂F, S(O)₂CHF₂, S(O)₂CF₃, 2-fluoroethylsulfonyl, 2,2-difluoroethylsulfonyl, 2,2,2-trifluoroethylsulfonyl, or S(O)₂C₂F₅. Fluorinated C₁-C₃-alkylsulfonyl is additionally, for example, 2-fluoropropylsulfonyl, 3-fluoropropylsulfonyl, 2,2-difluoropropylsulfonyl, 2,3-difluoropropylsulfonyl, 3,3,3-trifluoropropylsulfonyl, S(O)₂CH₂-C₂F₅, S(O)₂CF₂-C₂F₅ or 1-(CH₂F)-2-fluoroethylsulfonyl. Fluorinated C₁-C₄-alkylsulfonyl is additionally, for example, 4-fluorobutylsulfonyl or nonafluorobutylsulfonyl. Fluorinated C₁-C₆-alkylsulfonyl is additionally, for example, 5-fluoropentylsulfonyl, undecafluoropentylsulfonyl, 6-fluorohexylsulfonyl or dodecafluorohexylsulfonyl.

C₁-C₄-Alkylcarbonyl refers to a straight-chain or branched alkyl group having from 1 to 4 carbon atoms), which is bound to the remainder of the molecule via a carbonyl group (CO), such as in acetyl, propionyl, isopropylcarbonyl, butylcarbonyl, sec-butylcarbonyl, isobutylcarbonyl, and tert-butylcarbonyl. C₁-C₆-Alkylcarbonyl is a straight-chain or branched alkyl group having from 1 to 6 carbon atoms, which is bound to the remainder of the molecule via a carbonyl group (CO). Examples include, apart from those listed above for C₁-C₄-alkylcarbonyl, pentylcarbonyl, hexylcarbonyl and the constitutional isomers thereof.

Fluorinated C₁-C₄-alkylcarbonyl refers to a straight-chain or branched fluorinated alkyl group having from 1 to 4 carbon atoms as defined above, which is bound to the remainder of the molecule via a carbonyl group (CO). Fluorinated C₁-C₆-alkylcarbonyl

is a straight-chain or branched fluorinated alkyl group having from 1 to 6 carbon atoms as defined above, which is bound to the remainder of the molecule via a carbonyl group (CO). Examples include trifluoromethylcarbonyl, 2,2,2-trifluoroethylcarbonyl and the like.

C₃-C₆-cycloalkylcarbonyl relates to a C₃-C₆-cycloalkyl group as defined above which is bound to the remainder of the molecule via a carbonyl group (CO), such as in cyclopropylcarbonyl, cyclobutylcarbonyl, cyclopentylcarbonyl and cyclohexylcarbonyl.

C₁-C₆-Alkoxycarbonyl refers to a straight-chain or branched alkoxy group having from 1 to 6, especially 1 to 4 carbon atoms (= C₁-C₄-alkoxycarbonyl), in particular 1 to 3 carbon atoms (= C₁-C₃-alkoxycarbonyl), which is bound to the remainder of the molecule via a carbonyl group (CO), such as in methoxycarbonyl, ethoxycarbonyl, propyloxycarbonyl, and isopropyloxycarbonyl.

Fluorinated C₁-C₆-alkoxycarbonyl refers to a straight-chain or branched fluorinated alkoxy group having from 1 to 6, especially 1 to 4 carbon atoms (=fluorinated C₁-C₄-alkoxycarbonyl), in particular 1 to 3 carbon atoms (=fluorinated C₁-C₃-alkoxycarbonyl) as defined above, which is bound to the remainder of the molecule via a carbonyl group (CO). Examples include trifluoromethoxycarbonyl, 2,2,2-trifluoroethoxycarbonyl and the like.

C₁-C₄-Alkylcarbonyloxy refers to a straight-chain or branched alkyl group having from 1 to 4 carbon atoms, which is bound to the remainder of the molecule via a carbonyloxy group [C(O)-O-], such as in acet(yl)oxy, propionyloxy, isopropylcarbonyloxy, butylcarbonyloxy, sec-butylcarbonyloxy, isobutylcarbonyloxy, and tert-butylcarbonyloxy. C₁-C₆-Alkylcarbonyloxy is a straight-chain or branched alkyl group having from 1 to 6 carbon atoms, which is bound to the remainder of the molecule via a carbonyloxy group [C(O)-O-]. Examples include, apart those listed above for C₁-C₄-alkylcarbonyloxy, pentylcarbonyloxy, hexylcarbonyloxy and the constitutional isomers thereof.

Fluorinated C₁-C₄-alkylcarbonyloxy refers to a straight-chain or branched fluorinated alkyl group having from 1 to 4 carbon atoms as defined above, which is bound to the remainder of the molecule via a carbonyloxy group [C(O)-O-]. Fluorinated C₁-C₆-alkylcarbonyloxy is a straight-chain or branched fluorinated alkyl group having from 1 to 6 carbon atoms as defined above, which is bound to the remainder of the molecule

via a carbonyloxy group [C(O)-O-]. Examples include trifluoromethylcarbonyloxy, 2,2,2-trifluoroethylcarbonyloxy and the like.

Phenyl-C₁-C₂-alkyl is a phenyl group bound to the remainder of the molecule via a C₁-C₂-alkyl group. Examples are benzyl, 1-phenylethyl and 2-phenylethyl (phenethyl).

The term "3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO₂ and optionally also 1 or 2 C=O and/or C=S groups as ring members" denotes a 3-, 4-, 5-, 6-, 7- or 8-membered, preferably a 3-, 4-, 5-, 6- or 7-membered saturated, partially unsaturated or maximum unsaturated heteromonocyclic ring containing 1, 2, 3 or 4 (preferably 1, 2 or 3) heteroatoms or heteroatom groups selected from N, O, S, SO and SO₂ and optionally also 1 or 2 C=O and/or C=S groups as ring members.

Unsaturated rings contain at least one C-C and/or C-N and/or N-N double bond(s). Maximally unsaturated rings contain as many conjugated C-C and/or C-N and/or N-N double bonds as allowed by the ring size. Maximally unsaturated 5- or 6-membered heterocyclic rings are aromatic. 7- and 8-membered rings cannot be aromatic. They are homoaromatic (7-membered ring, 3 double bonds) or have 4 double bonds (8-membered ring). Partially unsaturated rings contain less than the maximum number of C-C and/or C-N and/or N-N double bond(s) allowed by the ring size. The heterocyclic ring may be attached to the remainder of the molecule via a carbon ring member or via a nitrogen ring member. As a matter of course, the heterocyclic ring contains at least one carbon ring atom. If the ring contains more than one O ring atom, these are not adjacent.

Examples of a 3-, 4-, 5-, 6- or 7-membered saturated heterocyclic ring include: Oxiranyl, thiiranyl, aziridinyl, oxetanyl, thietanyl, azetidiny, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrofuran-2-on-3-yl, tetrahydrofuran-2-on-4-yl, tetrahydrofuran-2-on-5-yl, tetrahydrofuran-2-thion-3-yl, tetrahydrofuran-2-thion-4-yl, tetrahydrofuran-2-thion-5-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, tetrahydrothien-2-on-3-yl, tetrahydrothien-2-on-4-yl, tetrahydrothien-2-on-5-yl, tetrahydrothien-2-thion-3-yl, tetrahydrothien-2-thion-4-yl, tetrahydrothien-2-thion-5-yl, pyrrolidin-1-yl, pyrrolidine-2-on-1-yl, pyrrolidine-2,5-dion-1-yl, pyrrolidine-2-thion-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, pyrrolidine-2-on-3-yl, pyrrolidine-2-on-4-yl, pyrrolidine-2-on-5-yl, pyrrolidine-2,5-dion-3-yl, pyrrolidine-2-thion-3-yl, pyrrolidine-2-thion-4-yl, pyrrolidine-2-thion-5-

yl, pyrazolidin-1-yl, pyrazolidin-3-yl, pyrazolidin-4-yl, pyrazolidin-5-yl, imidazolidin-1-yl, imidazolidin-2-on-1-yl, imidazolidin-2-thion-1-yl, imidazolidin-2-yl, imidazolidin-4-yl, imidazolidin-2-on-4-yl, imidazolidin-2-thion-4-yl, oxazolidin-2-yl, oxazolidin-3-yl, oxazolidin-4-yl, oxazolidin-5-yl, isoxazolidin-2-yl, isoxazolidin-3-yl, isoxazolidin-4-yl, isoxazolidin-5-yl, thiazolidin-2-yl, thiazolidin-3-yl, thiazolidin-4-yl, thiazolidin-5-yl, isothiazolidin-2-yl, isothiazolidin-3-yl, isothiazolidin-4-yl, isothiazolidin-5-yl, 1,2,4-oxadiazolidin-3-yl, 1,2,4-oxadiazolidin-5-yl, 1,2,4-thiadiazolidin-3-yl, 1,2,4-thiadiazolidin-5-yl, 1,2,4-triazolidin-3-yl, 1,3,4-oxadiazolidin-2-yl, 1,3,4-thiadiazolidin-2-yl, 1,3,4-triazolidin-1-yl, 1,3,4-triazolidin-2-yl, 2-tetrahydropyranyl, 4-tetrahydropyranyl, 1,3-dioxan-5-yl, 1,4-dioxan-2-yl, piperidin-1-yl, piperidin-2-on-1-yl, piperidin-2,5-dion-1-yl, piperidine-2-thion-1-yl, piperidin-2-yl, piperidin-3-yl, piperidin-2-on-3-yl, piperidin-2,5-dion-3-yl, piperidin-2-thion-3-yl, piperidin-4-yl, hexahydropyridazin-3-yl, hexahydropyridazin-4-yl, hexahydropyrimidin-2-yl, hexahydropyrimidin-4-yl, hexahydropyrimidin-5-yl, piperazin-1-yl, piperazin-2-yl, 1,3,5-hexahydrotriazin-1-yl, 1,3,5-hexahydrotriazin-2-yl and 1,2,4-hexahydrotriazin-3-yl, morpholin-2-yl, morpholin-3-yl, morpholin-4-yl, thiomorpholin-2-yl, thiomorpholin-3-yl, thiomorpholin-4-yl, 1-oxothiomorpholin-2-yl, 1-oxothiomorpholin-3-yl, 1-oxothiomorpholin-4-yl, 1,1-dioxothiomorpholin-2-yl, 1,1-dioxothiomorpholin-3-yl, 1,1-dioxothiomorpholin-4-yl, azepan-1-, -2-, -3- or -4-yl, oxepan-2-, -3-, -4- or -5-yl, hexahydro-1,3-diazepinyl, hexahydro-1,4-diazepinyl, hexahydro-1,3-oxazepinyl, hexahydro-1,4-oxazepinyl, hexahydro-1,3-dioxepinyl, hexahydro-1,4-dioxepinyl and the like.

Examples of a 3-, 4-, 5-, 6-, 7- or 8-membered saturated heterocyclic ring further include oxocane, thiocane, azocane, [1,3]diazocane, [1,4]diazocane, [1,5]diazocane, [1,5]oxazocane and the like.

Examples of a 3-, 4-, 5-, 6- or 7-membered partially unsaturated heterocyclic ring include: 2,3-dihydrofur-2-yl, 2,3-dihydrofur-3-yl, 2,4-dihydrofur-2-yl, 2,4-dihydrofur-3-yl, 2,3-dihydrothien-2-yl, 2,3-dihydrothien-3-yl, 2,4-dihydrothien-2-yl, 2,4-dihydrothien-3-yl, 2-pyrrolin-2-yl, 2-pyrrolin-3-yl, 3-pyrrolin-2-yl, 3-pyrrolin-3-yl, 2-isoxazolin-3-yl, 3-isoxazolin-3-yl, 4-isoxazolin-3-yl, 2-isoxazolin-4-yl, 3-isoxazolin-4-yl, 4-isoxazolin-4-yl, 2-isoxazolin-5-yl, 3-isoxazolin-5-yl, 4-isoxazolin-5-yl, 2-isothiazolin-3-yl, 3-isothiazolin-3-yl, 4-isothiazolin-3-yl, 2-isothiazolin-4-yl, 3-isothiazolin-4-yl, 4-isothiazolin-4-yl, 2-isothiazolin-5-yl, 3-isothiazolin-5-yl, 4-

isothiazolin-5-yl, 2,3-dihydropyrazol-1-yl, 2,3-dihydropyrazol-2-yl, 2,3-dihydropyrazol-3-yl, 2,3-dihydropyrazol-4-yl, 2,3-dihydropyrazol-5-yl, 3,4-dihydropyrazol-1-yl, 3,4-dihydropyrazol-3-yl, 3,4-dihydropyrazol-4-yl, 3,4-dihydropyrazol-5-yl, 4,5-dihydropyrazol-1-yl, 4,5-dihydropyrazol-3-yl, 4,5-dihydropyrazol-4-yl, 4,5-dihydropyrazol-5-yl, 2,3-dihydrooxazol-2-yl, 2,3-dihydrooxazol-3-yl, 2,3-dihydrooxazol-4-yl, 2,3-dihydrooxazol-5-yl, 3,4-dihydrooxazol-2-yl, 3,4-dihydrooxazol-3-yl, 3,4-dihydrooxazol-4-yl, 3,4-dihydrooxazol-5-yl, 3,4-dihydrooxazol-2-yl, 3,4-dihydrooxazol-3-yl, 3,4-dihydrooxazol-4-yl, 2-, 3-, 4-, 5- or 6-di- or tetrahydropyridinyl, 3-di- or tetrahydropyridazinyl, 4-di- or tetrahydropyridazinyl, 2-di- or tetrahydropyrimidinyl, 4-di- or tetrahydropyrimidinyl, 5-di- or tetrahydropyrimidinyl, di- or tetrahydropyrazinyl, 1,3,5-di- or tetrahydrotriazin-2-yl, 1,2,4-di- or tetrahydrotriazin-3-yl, 2,3,4,5-tetrahydro[1H]azepin-1-, -2-, -3-, -4-, -5-, -6- or -7-yl, 3,4,5,6-tetrahydro[2H]azepin-2-, -3-, -4-, -5-, -6- or -7-yl, 2,3,4,7-tetrahydro[1H]azepin-1-, -2-, -3-, -4-, -5-, -6- or -7-yl, 2,3,6,7-tetrahydro[1H]azepin-1-, -2-, -3-, -4-, -5-, -6- or -7-yl, tetrahydrooxepinyl, such as 2,3,4,5-tetrahydro[1H]oxepin-2-, -3-, -4-, -5-, -6- or -7-yl, 2,3,4,7-tetrahydro[1H]oxepin-2-, -3-, -4-, -5-, -6- or -7-yl, 2,3,6,7-tetrahydro[1H]oxepin-2-, -3-, -4-, -5-, -6- or -7-yl, tetrahydro-1,3-diazepinyl, tetrahydro-1,4-diazepinyl, tetrahydro-1,3-oxazepinyl, tetrahydro-1,4-oxazepinyl, tetrahydro-1,3-dioxepinyl and tetrahydro-1,4-dioxepinyl.

Examples of a 3-, 4-, 5-, 6-, 7- or 8-membered partially unsaturated heterocyclic ring further include 1,2,3,4,5,6-hexahydroazocine, 2,3,4,5,6,7-hexahydroazocine, 1,2,3,4,5,8-hexahydroazocine, 1,2,3,4,7,8-hexahydroazocine, 1,2,3,4,5,6-hexahydro-[1,5]diazocine, 1,2,3,4,7,8-hexahydro-[1,5]diazocine and the like.

Examples of a 3-, 4-, 5-, 6- or 7-membered maximally unsaturated (including aromatic) heterocyclic ring include 5- or 6-membered heteroaromatic rings, such as 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 1,3,4-triazol-1-yl, 1,3,4-triazol-2-yl, 2-pyridinyl, 3-pyridinyl, 4-pyridinyl, 1-oxypyridin-2-yl, 1-oxypyridin-3-yl, 1-oxypyridin-4-yl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl and 2-pyrazinyl, and also homoaromatic radicals, such as 1H-azepine, 1H-[1,3]-diazepine and 1H-[1,4]-diazepine.

Examples of a 3-, 4-, 5-, 6-, 7- or 8-membered maximally unsaturated heterocyclic ring further include [1,3]diazocine, [1,5]diazocine and [1,5]diazocine.

A 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members is either saturated, partially unsaturated and carbocyclic (if it contains only C=O and/or C=S as heteroatom group and no further heteroatoms or heteroatom groups) or saturated, partially unsaturated or maximally unsaturated heterocyclic. Examples are, in addition to the heterocyclic rings mentioned above, carbocyclic rings, such as cyclopropanonyl, cyclobutanonyl, cyclopentanonyl, cyclohexanonyl, cyclohexandionyl, cycloheptanonyl, cyclooctanonyl, cyclopropanthionyl, cyclobutanthionyl, cyclopentanthionyl, cyclohexanthionyl, cyclohexandithionyl, cycloheptanthionyl, cyclooctanthionyl, cyclopropenonyl, cyclopentenonyl, cyclohexenonyl and the like.

The remarks made above and in the following with respect to preferred aspects of the invention, e.g. to preferred meanings of the variables R¹, R², R^{3a}, R^{3b}, R^{4a}, R^{4b}, R^{5a}, R^{5b}, R⁶, R⁷, R⁸, R⁹, a, b and n of compounds I, to preferred compounds I and to preferred embodiments of the method or the use according to the invention, apply in each case on their own or in particular to combinations thereof.

In a preferred embodiment, R¹ is selected from hydrogen, cyano, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, -C(=O)R¹⁰, phenyl and benzyl, where the phenyl moiety in the two last-mentioned radicals may carry 1, 2 or 3 radicals R¹¹, where R¹⁰ and R¹¹ have one of the general meanings given above, or, in particular, one of the preferred meanings given below.

In the above group -C(=O)R¹⁰ as a meaning for R¹, R¹⁰ is preferably selected from C₁-C₄-alkyl and C₁-C₄-alkoxy and more preferably from C₁-C₂-alkyl and tert-butoxy.

More preferably, R¹ is selected from hydrogen and C₁-C₆-alkyl, in particular from hydrogen and methyl, and is specifically hydrogen.

In a preferred embodiment, R² is selected from cyano, nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy, more preferably from

C₁-C₆-alkyl, in particular from methyl, ethyl, propyl and isopropyl, more particularly from methyl and ethyl, and is specifically methyl.

In a preferred embodiment, R^{3a} and R^{3b}, independently of each other, are selected from hydrogen, cyano, nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy. More preferably, R^{3a} is selected from hydrogen, cyano, nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy, and R^{3b} is hydrogen. Even more preferably, R^{3a} is selected from hydrogen and methyl and R^{3b} is hydrogen. In particular, both R^{3a} and R^{3b} are hydrogen.

In a preferred embodiment, R^{4a} and R^{4b}, independently of each other, are selected from hydrogen, C₁-C₆-alkyl and fluorinated C₁-C₆-alkyl or form together a group =O. More preferably, they are hydrogen or form together a group =O. In another embodiment, one of R^{4a} and R^{4b} is hydrogen and the other is methyl; especially if X is CR⁷R⁸. In particular, both R^{4a} and R^{4b} are hydrogen.

In a preferred embodiment,

R^{5a} is selected from hydrogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy; and in case that X is CR⁷R⁸ is further selected from halogen; or

R^{5a} and R⁶, together with the atoms they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹; or R^{5a} and R⁷, together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹; and

R^{5b} is selected from hydrogen and deuterium and is preferably hydrogen;

where R^{11} has one of the general meanings given above, or, in particular, one of the preferred meanings given below.

More preferably,

R^{5a} is selected from hydrogen, C_1 - C_6 -alkyl and fluorinated C_1 - C_6 -alkyl; and in case that X is CR^7R^8 is further selected from halogen; and is preferably selected from hydrogen and methyl, or

R^{5a} and R^6 form together a group $(CH_2)_r$, where r is 2, 3, 4 or 5, preferably 2, 3 or 4; or

R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 2, 3, 4 or 5, preferably 2, 3 or 4; and

R^{5b} is selected from hydrogen and deuterium and is preferably hydrogen.

In an alternative more preferred embodiment,

R^{5a} is selected from hydrogen, C_1 - C_6 -alkyl and fluorinated C_1 - C_6 -alkyl; and in case that X is CR^7R^8 is further selected from halogen; and is preferably selected from hydrogen and methyl, or

R^{5a} and R^6 form together a group $(CH_2)_r$, where r is 2, 3, 4 or 5, where two hydrogen atoms bound to adjacent CH_2 groups may be replaced by two radicals R^{11} , where the two radicals R^{11} form together a group $(CH_2)_t$, where t is 1, 2, 3, 4 or 5, preferably 1, 2 or 3; or

R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 2, 3, 4 or 5, preferably 2, 3 or 4; and

R^{5b} is selected from hydrogen and deuterium and is preferably hydrogen.

In particular,

R^{5a} is hydrogen; and in case that X is CR^7R^8 , is selected from hydrogen and methyl and is in particular hydrogen; or

R^{5a} and R^6 form together a group $(CH_2)_r$, where r is 3, 4 or 5, where two hydrogen atoms bound to adjacent CH_2 groups may be replaced by two radicals R^{11} , where the two radicals R^{11} form together a group $(CH_2)_t$, where t is 1, 2, 3, 4 or 5, preferably 1, 2 or 3 and specifically 1; or

R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 2, 3 or 4, in particular 3; and R^{5b} is hydrogen.

In an alternatively preferred embodiment, R^{5a} and R^{5b} , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ; and form in particular a 5- or 6-membered saturated or partially unsaturated carbocyclic ring, and where the ring may be substituted with one or more substituents R^{11} .

In one embodiment, X is NR^6 , where R^6 has one of the general meanings given above, or, in particular, one of the preferred meanings given below.

Preferably,

R^6 is selected from the group consisting of hydrogen, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, -C(=O) R^{10} , -SO₂ R^{10} , phenyl and benzyl, where the phenyl moiety in the two last-mentioned radicals may carry 1, 2 or 3 radicals R^{11} ; or

R^{5a} and R^6 , together with the atoms they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ; where R^{10} and R^{11} have one of the general meanings given above, or, in particular, one of the preferred meanings given below.

In the above group -C(=O) R^{10} , R^6 , R^{10} is preferably selected from C₁-C₄-alkyl and C₁-C₄-alkoxy and more preferably from C₁-C₂-alkyl and tert-butoxy.

In the above group -SO₂ R^{10} , R^6 , R^{10} is preferably selected from C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl and phenyl, where phenyl may be substituted with 1, 2, 3, 4 or 5, preferably 1, 2 or 3, in particular 1, radicals R^{11} , where R^{11} has one of the general meanings given above, or, in particular, one of the preferred meanings given below.

In one more preferred embodiment, R^6 is $-SO_2R^{10}$, where R^{10} has one of the general meanings given above or is preferably selected from C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl and phenyl, where phenyl may be substituted with 1, 2, 3, 4 or 5, preferably 1, 2 or 3, in particular 1, radicals R^{11} , where R^{11} has one of the general meanings given above, or, in particular, one of the preferred meanings given below.

In another preferred embodiment, R^6 is selected from hydrogen, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkyl- C_1 - C_4 -alkyl, fluorinated C_3 - C_6 -cycloalkyl- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, fluorinated C_1 - C_4 -alkoxy, $-C(=O)R^{10}$, where R^{10} is selected from C_1 - C_6 -alkyl and C_3 - C_6 -cycloalkyl; phenyl, phenyl- C_1 - C_2 -alkyl and a 3-, 4-, 5- or 6-membered saturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO_2 and optionally also 1 or 2 $C=O$ and/or $C=S$ groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R^{11} .

In another preferred embodiment, R^6 is selected from C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkyl- C_1 - C_2 -alkyl, fluorinated C_3 - C_6 -cycloalkyl- C_1 - C_2 -alkyl, phenyl, phenyl- C_1 - C_2 -alkyl and a 3-, 4-, 5- or 6-membered saturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO_2 and optionally also 1 or 2 $C=O$ and/or $C=S$ groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R^{11} .

In a more preferred embodiment, R^6 is phenyl- C_1 - C_2 -alkyl, and in particular benzyl.

In another more preferred embodiment, R^6 is selected from C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkyl- C_1 - C_2 -alkyl and fluorinated C_3 - C_6 -cycloalkyl- C_1 - C_2 -alkyl, and in particular from C_1 - C_4 -

alkyl, fluorinated C₁-C₄-alkyl, C₃-C₆-cycloalkyl, fluorinated C₃-C₆-cycloalkyl, C₃-C₆-cycloalkyl-methyl and fluorinated C₃-C₆-cycloalkyl-methyl.

In another more preferred embodiment, R⁶ is a 3-, 4-, 5- or 6-membered saturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO₂ and optionally also 1 or 2 C=O and/or C=S groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R¹¹. Even more preferably, R⁶ is a 3-, 4- or 5-membered saturated heterocyclic ring containing 1 or 2 heteroatoms independently selected from N, O and S, where the heterocyclic ring may be substituted with one or two substituents selected from halogen and C₁-C₄-alkyl, and is in particular oxetanyl which may carry one or two substituents selected from halogen and C₁-C₄-alkyl and especially from F.

In another more preferred embodiment, R⁶ is selected from hydrogen, C₁-C₄-alkyl, fluorinated C₁-C₄-alkyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkyl-C₁-C₂-alkyl, C₃-C₆-cycloalkylcarbonyl, C₁-C₄-alkoxy, fluorinated C₁-C₄-alkoxy, phenyl-C₁-C₂-alkyl and a 3-, 4-, 5- or 6-membered saturated heterocyclic ring containing 1 or 2 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO₂ as ring members; and is in particular selected from C₁-C₄-alkyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkyl-C₁-C₂-alkyl, and a 3-, 4-, 5- or 6-membered saturated heterocyclic ring containing 1 or 2 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO₂ as ring members, where the saturated heterocyclic ring preferably contains 1 heteroatom selected from O and S as ring member. Specifically, R⁶ is selected from C₁-C₄-alkyl, C₃-C₆-cycloalkyl and oxetanyl and very specifically from ethyl, n-propyl, cyclopropyl, cyclobutyl and 3-oxetanyl.

In another more preferred embodiment, R^{5a} and R⁶, together with the atoms they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and

C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ;

where R^{11} has one of the general meanings given above, or, in particular, one of the preferred meanings given below.

In particular, R^{5a} and R^6 form together a group $(CH_2)_r$, where r is 2, 3, 4 or 5, preferably 2, 3 or 4 or, in particular, 3, 4 or 5. Alternatively, R^{5a} and R^6 form together a group $(CH_2)_r$, where r is 2, 3, 4 or 5 and in particular 3, 4 or 5, where two hydrogen atoms bound to adjacent CH_2 groups may be replaced by two radicals R^{11} , where the two radicals R^{11} form together a group $(CH_2)_t$, where t is 1, 2, 3, 4 or 5, in particular 1, 2 or 3 and specifically 1.

In an alternative embodiment, X is CR^7R^8 , where R^7 and R^8 have one of the general meanings given above, or, in particular, one of the preferred meanings given below.

In case that R^7 and R^8 , together with the carbon atom they are bound to, form a ring, this ring is spiro-bound to the carbon atom carrying R^7 and R^8 .

Preferably,

R^7 and R^8 , independently of each other, are selected from the group consisting of halogen, cyano, nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, fluorinated C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, fluorinated C_1 - C_6 -alkylsulfonyl, $-NR^{12a}R^{12b}$, $-CH_2NR^{12a}R^{12b}$, $-NR^{12a}C(O)R^{10}$, $-C(=O)R^{10}$, $SO_2NR^{12a}R^{12b}$, C_1 - C_6 -alkylcarbonyloxy, fluorinated C_1 - C_6 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_2 -alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO_2 , C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R^{11} ; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ; or

R^{5a} and R^7 , together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ;

in which case R^8 has one of the above meanings or is deuterium;

where R^{11} has one of the general meanings given above, or, in particular, one of the preferred meanings given below.

More preferably,

R^7 and R^8 , independently of each other, are selected from halogen, cyano, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, phenyl, phenyl-C₁-C₂-alkyl, benzyloxy and a 3-, 4-, 5- or 6-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the four last-mentioned radicals may be substituted with one or more substituents R^{11} ; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ; or

R^{5a} and R^7 , together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ;

in which case R^8 has one of the above meanings or is deuterium; and
 R^{5b} is selected from hydrogen and deuterium and is preferably hydrogen;
where R^{11} has one of the general meanings given above, or, in particular, one of the preferred meanings given below.

Even more preferably,

R^7 and R^8 , independently of each other, are selected from halogen, cyano, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl and phenyl, or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ; or

R^5 and R^7 form together a group $(CH_2)_s$, where s is 2, 3, 4 or 5, preferably 2, 3 or 4;

in which case R^8 has one of the above meanings or is deuterium; and

R^{5b} is selected from hydrogen and deuterium and is preferably hydrogen;

where R^{11} has one of the general meanings given above, or, in particular, one of the preferred meanings given below.

In particular,

R^8 is selected from halogen, cyano, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl and phenyl; and

R^7 is selected from C_1 - C_6 -alkyl and fluorinated C_1 - C_6 -alkyl, preferably from methyl and CF_3 and is in particular methyl; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ; or

R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 2, 3, 4 or 5, preferably 2, 3 or 4;

in which case R^8 has one of the above meanings or is deuterium; and

R^{5b} is selected from hydrogen and deuterium and is preferably hydrogen;

where R^{11} has one of the general meanings given above, or, in particular, one of the preferred meanings given below.

In an alternative particular embodiment,

R^7 is C_1 - C_6 -alkyl and is in particular methyl;

R^8 is selected from hydroxyl, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy and phenyl and in particular from C_1 - C_4 -alkyl, fluorinated C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, fluorinated C_1 - C_4 -alkoxy and phenyl; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated or partially unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ; or

R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 2, 3 or 4 and where R^8 is methyl; and

R^{5b} is hydrogen.

Specifically,

R^8 is selected from C_1 - C_4 -alkyl, fluorinated C_1 - C_2 -alkyl and phenyl and specifically from methyl, CF_3 and phenyl; and

R^7 is methyl; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5- or 6-membered saturated or partially unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ; or

R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 3 or 4, specifically 3;

in which case R^8 is selected from methyl and deuterium; and

R^{5b} is selected from hydrogen and deuterium and is specifically hydrogen;

where R^{11} has one of the general meanings given above, or, in particular, one of the preferred meanings given below.

In an alternative specific embodiment,

R^7 is methyl;

R^8 is selected from hydroxyl, methyl, ethyl, CF_3 , methoxy and phenyl; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5- or 6-membered saturated or partially unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ; or

R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 2, 3 or 4 and in particular and where R^8 is methyl; and

R^{5b} is hydrogen.

Very specifically,

R^8 is selected from methyl, CF_3 and phenyl; and

R^7 is methyl; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 4, 5- or 6-membered saturated or partially unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ; or form a 3-membered saturated or partially unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ; or

R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 3 or 4, specifically 3;

in which case R^8 is selected from methyl and deuterium and in particular methyl; and

R^{5b} is selected from hydrogen and deuterium and is specifically hydrogen;

where R^{11} has one of the general meanings given above, or, in particular, one of the preferred meanings given below.

In one particular embodiment, the ring formed by R^7 and R^8 together with the carbon atom they are bound to is a carbocyclic ring, preferably a saturated or partially unsaturated carbocyclic ring, in particular a saturated carbocyclic ring.

If R^7 and R^8 , together with the carbon atom they are bound to, form a carbocyclic ring, this is particularly preferably a 4- or 5-membered saturated carbocyclic ring (i.e. R^7 and R^8 form together a group $-(CH_2)_3-$ or $-(CH_2)_4-$) and in particular a 4-membered saturated carbocyclic ring (i.e. R^7 and R^8 form together a group $-(CH_2)_3-$), where the ring may be substituted with one or more substituents R^{11} . Specifically, the ring is not substituted. Alternatively, the carbocyclic ring is 3-membered (i.e. R^7 and R^8 form together a group $-(CH_2)_2-$), where the ring may be substituted with one or more substituents R^{11} . Specifically, the ring is not substituted.

In an alternative particular embodiment, the ring formed by R^7 and R^8 together with the carbon atom they are bound to is a heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO and SO_2 as ring members, preferably containing 1 or 2 heteroatoms or heteroatom-containing groups selected from O, S, N, SO and SO_2 as ring members. Preferably, the heterocyclic ring is

3-, 4-, 5- or 6-membered and saturated and may carry one or more substituents R^{11} . In particular, the heterocyclic ring is oxetanyl which may carry one or more substituents R^{11} .

Specifically, R^7 and R^8 , together with the carbon atom they are bound to, form a 3- or 4-membered saturated carbocyclic ring (i.e. R^7 and R^8 form together a group $-(CH_2)_2-$ or $-(CH_2)_3-$) and in particular a 4-membered saturated carbocyclic ring (i.e. R^7 and R^8 form together a group $-(CH_2)_3-$); the ring not being substituted.

In a preferred embodiment, each R^9 is independently selected from halogen, cyano, nitro, hydroxy, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy and a 3-, 4-, 5- or 6-membered saturated heterocyclic ring containing 1 or 2 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO_2 as ring members, where the heterocyclic ring may be substituted with one or more substituents R^{11} . The heterocyclic ring is specifically oxetanyl.

More preferably, each R^9 is independently selected from halogen, cyano, nitro, hydroxy, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy and fluorinated C_1 - C_6 -alkoxy, even more preferably from halogen, cyano, C_1 - C_4 -alkyl, fluorinated C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy and fluorinated C_1 - C_4 -alkoxy, and is in particular selected from halogen, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl and C_1 - C_6 -alkoxy, more particularly from halogen, C_1 - C_6 -alkyl and C_1 - C_6 -alkoxy, and is specifically halogen, especially F or Cl, and more specifically F.

In case that X is NR^6 , in alternatively more preferred embodiment, each R^9 is independently selected from halogen, cyano, nitro, hydroxy, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_1 - C_6 -alkoxy and fluorinated C_1 - C_6 -alkoxy, even more preferably from halogen, C_1 - C_6 -alkyl and C_3 - C_6 -cycloalkyl, and in particular from halogen and C_1 - C_6 -alkyl.

In a preferred embodiment, R^{10} is selected from hydrogen, hydroxy, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, $-NR^{12a}R^{12b}$, $-CH_2NR^{12a}R^{12b}$, phenyl, phenyl- C_1 - C_2 -alkyl, phenoxy, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the five last-mentioned radicals may be substituted with one or more substituents R^{11} , where R^{11} , R^{12a} and R^{12b} have one of the general meanings given above, or, in particular, one of the preferred meanings given below. More preferably, R^{10} is selected from hydrogen, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy and fluorinated C_1 - C_6 -alkoxy, and even more preferably from C_1 - C_4 -alkyl and C_1 - C_4 -alkoxy. In particular, R^{10} is selected from C_1 - C_2 -alkyl and tert-butoxy.

In a preferred embodiment, each R^{11} is independently selected from halogen, cyano, nitro, hydroxy, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy and fluorinated C_1 - C_6 -alkoxy, and is in particular halogen or C_1 - C_4 -alkyl. Alternatively, two radicals R^{11} bound on adjacent ring atoms form together a group $(CH_2)_t$, where t is 1, 2, 3, 4 or 5 and in particular 1, 2 or 3.

In a preferred embodiment, R^{12a} and R^{12b} , independently of each other and independently of each occurrence, are selected from hydrogen, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -alkylcarbonyl, fluorinated C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -alkoxycarbonyl, fluorinated C_1 - C_6 -alkoxycarbonyl, phenyl and benzyl, where the phenyl moieties in the two last-mentioned radicals may carry 1, 2 or 3 substituents selected from halogen, cyano nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy and fluorinated C_1 - C_6 -alkoxy; or, if R^{12a} and R^{12b} are bound to the same nitrogen atom, together with this nitrogen atom may form a 3-, 4-, 5-, 6- or 7-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1 or 2 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents selected

from halogen, cyano nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy.

More preferably, R^{12a} and R^{12b}, independently of each other and independently of each occurrence, are selected from hydrogen, C₁-C₆-alkyl, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl and benzyl, where the phenyl moiety in the last-mentioned radical may carry 1, 2 or 3 substituents selected from halogen, cyano nitro, C₁-C₄-alkyl, fluorinated C₁-C₄-alkyl, C₁-C₄-alkoxy and fluorinated C₁-C₄-alkoxy; or, if R^{12a} and R^{12b} are bound to the same nitrogen atom, together with this nitrogen atom may form a 5- or 6-membered saturated or aromatic heterocyclic ring, where the ring may further contain 1 or 2 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂ and C=O as ring members, and where the ring may be substituted with 1 or 2 substituents selected from halogen, cyano nitro, C₁-C₄-alkyl, fluorinated C₁-C₄-alkyl, C₁-C₄-alkoxy and fluorinated C₁-C₄-alkoxy.

In particular, R^{12a} and R^{12b}, independently of each other and independently of each occurrence, are selected from hydrogen and C₁-C₆-alkyl.


In a preferred embodiment, a is 0 or 1 and in particular 0.

In a particular embodiment, a is 1 and R² is bound in β-position to the nitrogen ring atom carrying R¹ (this is for the example the position of R^{2b} in below formula I.2).

In a preferred embodiment, b is 0, 1 or 2 and in particular 0 or 1.


In a preferred embodiment, n is 1.

In a particular embodiment, the compound of formula I is a compound of formula I.1



wherein X, R^{5a}, R^{5b}, R⁹ and b have one of the above general, or, in particular, one of the above preferred meanings.


In a specific embodiment, the compound of formula I is a compound of formula I.1.1



wherein

R^{9a} is selected from H, halogen, C₁-C₄-alkyl, fluorinated C₁-C₄-alkyl, C₃-C₆-cycloalkyl and fluorinated C₃-C₆-cycloalkyl, especially from H or F; and R^{5a}, R^{5b}, R⁶, R⁹ and b have one of the above general, or, in particular, one of the above preferred meanings.

In another specific embodiment, the compound of formula I is a compound of formula I.1.2



wherein

R⁷ is selected from C₁-C₄-alkyl and fluorinated C₁-C₄-alkyl;

R⁸ is selected from C₁-C₄-alkyl, fluorinated C₁-C₄-alkyl, C₁-C₄-alkoxy, fluorinated C₁-C₄-alkoxy and phenyl; or

R⁷ and R⁸, together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7-membered, in particular a 3- or 4-membered saturated or partially unsaturated ring, especially a saturated carbocyclic ring, where the ring may be substituted

with one or more substituents R^{11} , and form especially a 3- or 4-membered saturated carbocyclic ring;

R^{9a} is H, Cl, F or methyl, especially H, Cl or F; and

R^9 and b have one of the above general, or, in particular, one of the above preferred meanings.

In an alternative embodiment, in compounds I.1.2,

R^7 is selected from C_1 - C_4 -alkyl and fluorinated C_1 - C_4 -alkyl, preferably from methyl and CF_3 , and is in particular methyl;


R^8 is selected from C_1 - C_4 -alkyl and fluorinated C_1 - C_4 -alkyl, and preferably from methyl and CF_3 ; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated or partially unsaturated ring, where the ring may be substituted with one or more substituents R^{11} , and form preferably a 5- or 6-membered saturated or partially unsaturated ring;

R^{9a} is H or F; and


R^9 and b have one of the above general, or, in particular, one of the above preferred meanings.

In another specific embodiment, the compound of formula I is a compound of formula I.1.3



wherein R^8 is selected from deuterium, F, Cl, CN and CH_3 , in particular from deuterium, F, Cl and CN.


Among these, preference is given to the trans compounds, i.e. of formula I.1.3-trans:



(I.1.3-trans)

wherein R^8 is selected from deuterium, F, Cl, CN and CH_3 , in particular from deuterium, F, Cl and CN.


In another particular embodiment, the compound of formula I is a compound of formula I.2



(I.2)

wherein X, R^{5a} , R^{5b} , R^9 and b have one of the above general, or, in particular, one of the above preferred meanings.

In another particular embodiment, the compound of formula I is a compound of formula I.2.1



(I.2.1)

wherein

R^{2a} , R^{2b} and R^{3a} , independently of each other, are selected from hydrogen and methyl;

and where in particular at most one of R^{2a} , R^{2b} and R^{3a} is methyl;

R^{9a} is selected from H, halogen, C_1 - C_4 -alkyl, fluorinated C_1 - C_4 -alkyl, C_3 - C_6 -cycloalkyl and fluorinated C_3 - C_6 -cycloalkyl; and
 R^{5a} , R^{5b} , R^6 , R^9 and b have one of the above general, or, in particular, one of the above preferred meanings.


In particular, in compounds I.2.1,

R^{9a} is H or F; and

R^{5a} , R^{5b} , R^6 , R^9 and b have one of the above general, or, in particular, one of the above preferred meanings.

In particular, in compounds I.2.1, R^{2b} is methyl and R^{2a} and R^{3a} are hydrogen.

In another particular embodiment, the compound of formula I is a compound of formula I.2.2



wherein

R^{2a} , R^{2b} , R^{3a} and R^{5a} , independently of each other, are selected from hydrogen, methyl and ethyl, in particular from hydrogen and methyl;

R^7 is selected from C_1 - C_4 -alkyl and fluorinated C_1 - C_4 -alkyl, in particular from methyl and CF_3 , and is especially methyl;

R^8 is selected from C_1 - C_4 -alkyl, fluorinated C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, fluorinated C_1 - C_4 -alkoxy and phenyl; and in particular from methyl and CF_3 ; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7-membered, in particular a 3- or 4-membered saturated or partially unsaturated, especially a saturated carbocyclic ring, where the ring may be substituted with one or more substituents R^{11} , and form especially a 3- or 4-membered saturated carbocyclic ring;

R^{9a} is H, Cl, F or methyl, especially H, Cl or F; and
 R^9 and b have one of the above general, or, in particular, one of the above preferred meanings.

In particular, in compound I.2.2,

R^{2a} , R^{2b} , R^{3a} and R^{5a} , independently of each other, are selected from hydrogen and methyl;

R^7 is selected from C_1 - C_4 -alkyl and fluorinated C_1 - C_4 -alkyl, preferably from methyl and CF_3 , and is in particular methyl;

R^8 is selected from C_1 - C_4 -alkyl and fluorinated C_1 - C_4 -alkyl, and preferably from methyl and CF_3 ; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated or partially unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ;

R^{9a} is H or F; and

R^9 and b have one of the above general, or, in particular, one of the above preferred meanings.


In particular, in compound I.2.2, R^{2b} is methyl and R^{2a} , R^{3a} and R^{5a} are hydrogen.

In the above formulae I.1.1, I.1.2, I.2.1 and I.2.2, (b-1) is preferably 0.


Preferably, in the above formulae I.2, I.2.1 and I.2.2, R^{2a} , R^{2b} , R^{3a} and R^{5a} are selected from hydrogen and methyl, with the proviso that at most two, preferably at most one, of the substituents R^{2a} , R^{2b} , R^{3a} and R^{5a} are methyl. In particular, R^{2b} is hydrogen and R^{2a} , R^{3a} and R^{5a} are selected from hydrogen and methyl, with the proviso that at most two, preferably at most one, of the substituents R^{2a} , R^{3a} and R^{5a} are methyl. However, more particularly, R^{2b} is hydrogen or methyl, especially methyl, and R^{2a} , R^{3a} and R^{5a} are hydrogen (if the latter does not form a ring together with R^6 or R^7 and the atoms they are bound to).

Preferably, in the above formulae I.2, I.2.1 and I.2.2, R^7 and R^8 , together with the carbon atom they are bound to, form a 3, 4-, 5- or 6-membered saturated or partially unsaturated carbocyclic ring, preferably a 3, 4- or 5-membered saturated carbocyclic ring, in particular a 4-membered saturated carbocyclic ring, where the ring may be substituted with one or more substituents R^{11} .


Examples of preferred compounds are compounds of the following formulae I.a to I.ii, where the variables have one of the general or preferred meanings given above. Examples of preferred compounds are the individual compounds compiled in the tables 1 to 36 below. Moreover, the meanings mentioned below for the individual variables in the tables are per se, independently of the combination in which they are mentioned, a particularly preferred embodiment of the substituents in question.




I.a




I.b




I.c




I.d




I.e




I.f




I.g




I.h




I.i




I.j




I.k




I.l




I.m




I.n




I.o




I.p




I.q




I.r




I.s




I.t




I.u




I.v




I.w




I.x




I.y




I.z




I.zz




I.aa




I.bb




I.cc




I.dd




I.ee




I.ff



I.gg



I.hh



I.ii

Table 1

Compounds of the formula I.a in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 2

Compounds of the formula I.b in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 3

Compounds of the formula I.c in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 4

Compounds of the formula I.d in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 5

Compounds of the formula I.e in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 6

Compounds of the formula I.f in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 7

Compounds of the formula I.g in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 8

Compounds of the formula I.h in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 9

Compounds of the formula I.i in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 10

Compounds of the formula I.j in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 11

Compounds of the formula I.k in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 12

Compounds of the formula I.l in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 13

Compounds of the formula I.m in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 14

Compounds of the formula I.n in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 15

Compounds of the formula I.o in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 16

Compounds of the formula I.p in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 17

Compounds of the formula I.q in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 18

Compounds of the formula I.r in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 19

Compounds of the formula I.s in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 20

Compounds of the formula I.t in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 21

Compounds of the formula I.u in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 22

Compounds of the formula I.v in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 23

Compounds of the formula I.w in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 24

Compounds of the formula I.x in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 25

Compounds of the formula I.y in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 26

Compounds of the formula I.z in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 27

Compounds of the formula I.zz in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} for a compound corresponds in each case to one row of Table B.

Table 28

Compounds of the formula I.aa in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 29

Compounds of the formula I.bb in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 30

Compounds of the formula I.cc in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 31

Compounds of the formula I.dd in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 32

Compounds of the formula I.ee in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 33

Compounds of the formula I.ff in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 34

Compounds of the formula I.gg in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 35

Compounds of the formula I.hh in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table 36

Compounds of the formula I.ii in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} for a compound corresponds in each case to one row of Table A.

Table A

No.	R^{9a}	R^{5b}	R^{5a}	R^6
A-1	H	H	H	-SO ₂ -CH ₃
A-2	F	H	H	-SO ₂ -CH ₃
A-3	Cl	H	H	-SO ₂ -CH ₃
A-4	Br	H	H	-SO ₂ -CH ₃
A-5	CH ₃	H	H	-SO ₂ -CH ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-6	CF ₃	H	H	-SO ₂ -CH ₃
A-7	OCH ₃	H	H	-SO ₂ -CH ₃
A-8	OCF ₃	H	H	-SO ₂ -CH ₃
A-9	cyclopropyl	H	H	-SO ₂ -CH ₃
A-10	cyclobutyl	H	H	-SO ₂ -CH ₃
A-11	cyclopentyl	H	H	-SO ₂ -CH ₃
A-12	cyclohexyl	H	H	-SO ₂ -CH ₃
A-13	H	CH ₃	H	-SO ₂ -CH ₃
A-14	F	CH ₃	H	-SO ₂ -CH ₃
A-15	Cl	CH ₃	H	-SO ₂ -CH ₃
A-16	Br	CH ₃	H	-SO ₂ -CH ₃
A-17	CH ₃	CH ₃	H	-SO ₂ -CH ₃
A-18	CF ₃	CH ₃	H	-SO ₂ -CH ₃
A-19	OCH ₃	CH ₃	H	-SO ₂ -CH ₃
A-20	OCF ₃	CH ₃	H	-SO ₂ -CH ₃
A-21	cyclopropyl	CH ₃	H	-SO ₂ -CH ₃
A-22	cyclobutyl	CH ₃	H	-SO ₂ -CH ₃
A-23	cyclopentyl	CH ₃	H	-SO ₂ -CH ₃
A-24	cyclohexyl	CH ₃	H	-SO ₂ -CH ₃
A-25	H	CH ₃	CH ₃	-SO ₂ -CH ₃
A-26	F	CH ₃	CH ₃	-SO ₂ -CH ₃
A-27	Cl	CH ₃	CH ₃	-SO ₂ -CH ₃
A-28	Br	CH ₃	CH ₃	-SO ₂ -CH ₃
A-29	CH ₃	CH ₃	CH ₃	-SO ₂ -CH ₃
A-30	CF ₃	CH ₃	CH ₃	-SO ₂ -CH ₃
A-31	OCH ₃	CH ₃	CH ₃	-SO ₂ -CH ₃
A-32	OCF ₃	CH ₃	CH ₃	-SO ₂ -CH ₃
A-33	cyclopropyl	CH ₃	CH ₃	-SO ₂ -CH ₃
A-34	cyclobutyl	CH ₃	CH ₃	-SO ₂ -CH ₃
A-35	cyclopentyl	CH ₃	CH ₃	-SO ₂ -CH ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-36	cyclohexyl	CH ₃	CH ₃	-SO ₂ -CH ₃
A-37	H	H	H	-SO ₂ -CH ₂ CH ₃
A-38	F	H	H	-SO ₂ -CH ₂ CH ₃
A-39	Cl	H	H	-SO ₂ -CH ₂ CH ₃
A-40	Br	H	H	-SO ₂ -CH ₂ CH ₃
A-41	CH ₃	H	H	-SO ₂ -CH ₂ CH ₃
A-42	CF ₃	H	H	-SO ₂ -CH ₂ CH ₃
A-43	OCH ₃	H	H	-SO ₂ -CH ₂ CH ₃
A-44	OCF ₃	H	H	-SO ₂ -CH ₂ CH ₃
A-45	cyclopropyl	H	H	-SO ₂ -CH ₂ CH ₃
A-46	cyclobutyl	H	H	-SO ₂ -CH ₂ CH ₃
A-47	cyclopentyl	H	H	-SO ₂ -CH ₂ CH ₃
A-48	cyclohexyl	H	H	-SO ₂ -CH ₂ CH ₃
A-49	H	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-50	F	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-51	Cl	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-52	Br	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-53	CH ₃	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-54	CF ₃	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-55	OCH ₃	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-56	OCF ₃	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-57	cyclopropyl	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-58	cyclobutyl	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-59	cyclopentyl	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-60	cyclohexyl	CH ₃	H	-SO ₂ -CH ₂ CH ₃
A-61	H	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃
A-62	F	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃
A-63	Cl	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃
A-64	Br	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃
A-65	CH ₃	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-66	CF ₃	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃
A-67	OCH ₃	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃
A-68	OCF ₃	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃
A-69	cyclopropyl	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃
A-70	cyclobutyl	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃
A-71	cyclopentyl	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃
A-72	cyclohexyl	CH ₃	CH ₃	-SO ₂ -CH ₂ CH ₃
A-73	H	H	H	-SO ₂ -CF ₃
A-74	F	H	H	-SO ₂ -CF ₃
A-75	Cl	H	H	-SO ₂ -CF ₃
A-76	Br	H	H	-SO ₂ -CF ₃
A-77	CH ₃	H	H	-SO ₂ -CF ₃
A-78	CF ₃	H	H	-SO ₂ -CF ₃
A-79	OCH ₃	H	H	-SO ₂ -CF ₃
A-80	OCF ₃	H	H	-SO ₂ -CF ₃
A-81	cyclopropyl	H	H	-SO ₂ -CF ₃
A-82	cyclobutyl	H	H	-SO ₂ -CF ₃
A-83	cyclopentyl	H	H	-SO ₂ -CF ₃
A-84	cyclohexyl	H	H	-SO ₂ -CF ₃
A-85	H	CH ₃	H	-SO ₂ -CF ₃
A-86	F	CH ₃	H	-SO ₂ -CF ₃
A-87	Cl	CH ₃	H	-SO ₂ -CF ₃
A-88	Br	CH ₃	H	-SO ₂ -CF ₃
A-89	CH ₃	CH ₃	H	-SO ₂ -CF ₃
A-90	CF ₃	CH ₃	H	-SO ₂ -CF ₃
A-91	OCH ₃	CH ₃	H	-SO ₂ -CF ₃
A-92	OCF ₃	CH ₃	H	-SO ₂ -CF ₃
A-93	cyclopropyl	CH ₃	H	-SO ₂ -CF ₃
A-94	cyclobutyl	CH ₃	H	-SO ₂ -CF ₃
A-95	cyclopentyl	CH ₃	H	-SO ₂ -CF ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-96	cyclohexyl	CH ₃	H	-SO ₂ -CF ₃
A-97	H	CH ₃	CH ₃	-SO ₂ -CF ₃
A-98	F	CH ₃	CH ₃	-SO ₂ -CF ₃
A-99	Cl	CH ₃	CH ₃	-SO ₂ -CF ₃
A-100	Br	CH ₃	CH ₃	-SO ₂ -CF ₃
A-101	CH ₃	CH ₃	CH ₃	-SO ₂ -CF ₃
A-102	CF ₃	CH ₃	CH ₃	-SO ₂ -CF ₃
A-103	OCH ₃	CH ₃	CH ₃	-SO ₂ -CF ₃
A-104	OCF ₃	CH ₃	CH ₃	-SO ₂ -CF ₃
A-105	cyclopropyl	CH ₃	CH ₃	-SO ₂ -CF ₃
A-106	cyclobutyl	CH ₃	CH ₃	-SO ₂ -CF ₃
A-107	cyclopentyl	CH ₃	CH ₃	-SO ₂ -CF ₃
A-108	cyclohexyl	CH ₃	CH ₃	-SO ₂ -CF ₃
A-109	H	H	H	-SO ₂ -C ₆ H ₅ *
A-110	F	H	H	-SO ₂ -C ₆ H ₅
A-111	Cl	H	H	-SO ₂ -C ₆ H ₅
A-112	Br	H	H	-SO ₂ -C ₆ H ₅
A-113	CH ₃	H	H	-SO ₂ -C ₆ H ₅
A-114	CF ₃	H	H	-SO ₂ -C ₆ H ₅
A-115	OCH ₃	H	H	-SO ₂ -C ₆ H ₅
A-116	OCF ₃	H	H	-SO ₂ -C ₆ H ₅
A-117	cyclopropyl	H	H	-SO ₂ -C ₆ H ₅
A-118	cyclobutyl	H	H	-SO ₂ -C ₆ H ₅
A-119	cyclopentyl	H	H	-SO ₂ -C ₆ H ₅
A-120	cyclohexyl	H	H	-SO ₂ -C ₆ H ₅
A-121	H	CH ₃	H	-SO ₂ -C ₆ H ₅
A-122	F	CH ₃	H	-SO ₂ -C ₆ H ₅
A-123	Cl	CH ₃	H	-SO ₂ -C ₆ H ₅
A-124	Br	CH ₃	H	-SO ₂ -C ₆ H ₅
A-125	CH ₃	CH ₃	H	-SO ₂ -C ₆ H ₅

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-126	CF ₃	CH ₃	H	-SO ₂ -C ₆ H ₅
A-127	OCH ₃	CH ₃	H	-SO ₂ -C ₆ H ₅
A-128	OCF ₃	CH ₃	H	-SO ₂ -C ₆ H ₅
A-129	cyclopropyl	CH ₃	H	-SO ₂ -C ₆ H ₅
A-130	cyclobutyl	CH ₃	H	-SO ₂ -C ₆ H ₅
A-131	cyclopentyl	CH ₃	H	-SO ₂ -C ₆ H ₅
A-132	cyclohexyl	CH ₃	H	-SO ₂ -C ₆ H ₅
A-133	H	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-134	F	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-135	Cl	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-136	Br	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-137	CH ₃	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-138	CF ₃	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-139	OCH ₃	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-140	OCF ₃	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-141	cyclopropyl	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-142	cyclobutyl	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-143	cyclopentyl	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-144	cyclohexyl	CH ₃	CH ₃	-SO ₂ -C ₆ H ₅
A-145	H	H	H	-SO ₂ -C ₆ H ₄ CH ₃ **
A-146	F	H	H	-SO ₂ -C ₆ H ₄ CH ₃
A-147	Cl	H	H	-SO ₂ -C ₆ H ₄ CH ₃
A-148	Br	H	H	-SO ₂ -C ₆ H ₄ CH ₃
A-149	CH ₃	H	H	-SO ₂ -C ₆ H ₄ CH ₃
A-150	CF ₃	H	H	-SO ₂ -C ₆ H ₄ CH ₃
A-151	OCH ₃	H	H	-SO ₂ -C ₆ H ₄ CH ₃
A-152	OCF ₃	H	H	-SO ₂ -C ₆ H ₄ CH ₃
A-153	cyclopropyl	H	H	-SO ₂ -C ₆ H ₄ CH ₃
A-154	cyclobutyl	H	H	-SO ₂ -C ₆ H ₄ CH ₃
A-155	cyclopentyl	H	H	-SO ₂ -C ₆ H ₄ CH ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-156	cyclohexyl	H	H	-SO ₂ -C ₆ H ₄ CH ₃
A-157	H	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-158	F	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-159	Cl	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-160	Br	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-161	CH ₃	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-162	CF ₃	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-163	OCH ₃	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-164	OCF ₃	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-165	cyclopropyl	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-166	cyclobutyl	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-167	cyclopentyl	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-168	cyclohexyl	CH ₃	H	-SO ₂ -C ₆ H ₄ CH ₃
A-169	H	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-170	F	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-171	Cl	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-172	Br	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-173	CH ₃	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-174	CF ₃	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-175	OCH ₃	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-176	OCF ₃	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-177	cyclopropyl	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-178	cyclobutyl	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-179	cyclopentyl	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-180	cyclohexyl	CH ₃	CH ₃	-SO ₂ -C ₆ H ₄ CH ₃
A-181	H	H	H	benzyl
A-182	F	H	H	benzyl
A-183	Cl	H	H	benzyl
A-184	Br	H	H	benzyl
A-185	CH ₃	H	H	benzyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-186	CF ₃	H	H	benzyl
A-187	OCH ₃	H	H	benzyl
A-188	OCF ₃	H	H	benzyl
A-189	cyclopropyl	H	H	benzyl
A-190	cyclobutyl	H	H	benzyl
A-191	cyclopentyl	H	H	benzyl
A-192	cyclohexyl	H	H	benzyl
A-193	H	CH ₃	H	benzyl
A-194	F	CH ₃	H	benzyl
A-195	Cl	CH ₃	H	benzyl
A-196	Br	CH ₃	H	benzyl
A-197	CH ₃	CH ₃	H	benzyl
A-198	CF ₃	CH ₃	H	benzyl
A-199	OCH ₃	CH ₃	H	benzyl
A-200	OCF ₃	CH ₃	H	benzyl
A-201	cyclopropyl	CH ₃	H	benzyl
A-202	cyclobutyl	CH ₃	H	benzyl
A-203	cyclopentyl	CH ₃	H	benzyl
A-204	cyclohexyl	CH ₃	H	benzyl
A-205	H	CH ₃	CH ₃	benzyl
A-206	F	CH ₃	CH ₃	benzyl
A-207	Cl	CH ₃	CH ₃	benzyl
A-208	Br	CH ₃	CH ₃	benzyl
A-209	CH ₃	CH ₃	CH ₃	benzyl
A-210	CF ₃	CH ₃	CH ₃	benzyl
A-211	OCH ₃	CH ₃	CH ₃	benzyl
A-212	OCF ₃	CH ₃	CH ₃	benzyl
A-213	cyclopropyl	CH ₃	CH ₃	benzyl
A-214	cyclobutyl	CH ₃	CH ₃	benzyl
A-215	cyclopentyl	CH ₃	CH ₃	benzyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-216	cyclohexyl	CH ₃	CH ₃	benzyl
A-217	H	H	H	H
A-218	F	H	H	H
A-219	Cl	H	H	H
A-220	Br	H	H	H
A-221	CH ₃	H	H	H
A-222	CF ₃	H	H	H
A-223	OCH ₃	H	H	H
A-224	OCF ₃	H	H	H
A-225	cyclopropyl	H	H	H
A-226	cyclobutyl	H	H	H
A-227	cyclopentyl	H	H	H
A-228	cyclohexyl	H	H	H
A-229	H	CH ₃	H	H
A-230	F	CH ₃	H	H
A-231	Cl	CH ₃	H	H
A-232	Br	CH ₃	H	H
A-233	CH ₃	CH ₃	H	H
A-234	CF ₃	CH ₃	H	H
A-235	OCH ₃	CH ₃	H	H
A-236	OCF ₃	CH ₃	H	H
A-237	cyclopropyl	CH ₃	H	H
A-238	cyclobutyl	CH ₃	H	H
A-239	cyclopentyl	CH ₃	H	H
A-240	cyclohexyl	CH ₃	H	H
A-241	H	CH ₃	CH ₃	H
A-242	F	CH ₃	CH ₃	H
A-243	Cl	CH ₃	CH ₃	H
A-244	Br	CH ₃	CH ₃	H
A-245	CH ₃	CH ₃	CH ₃	H

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-246	CF ₃	CH ₃	CH ₃	H
A-247	OCH ₃	CH ₃	CH ₃	H
A-248	OCF ₃	CH ₃	CH ₃	H
A-249	cyclopropyl	CH ₃	CH ₃	H
A-250	cyclobutyl	CH ₃	CH ₃	H
A-251	cyclopentyl	CH ₃	CH ₃	H
A-252	cyclohexyl	CH ₃	CH ₃	H
A-253	H	H	H	CH ₃
A-254	F	H	H	CH ₃
A-255	Cl	H	H	CH ₃
A-256	Br	H	H	CH ₃
A-257	CH ₃	H	H	CH ₃
A-258	CF ₃	H	H	CH ₃
A-259	OCH ₃	H	H	CH ₃
A-260	OCF ₃	H	H	CH ₃
A-261	cyclopropyl	H	H	CH ₃
A-262	cyclobutyl	H	H	CH ₃
A-263	cyclopentyl	H	H	CH ₃
A-264	cyclohexyl	H	H	CH ₃
A-265	H	CH ₃	H	CH ₃
A-266	F	CH ₃	H	CH ₃
A-267	Cl	CH ₃	H	CH ₃
A-268	Br	CH ₃	H	CH ₃
A-269	CH ₃	CH ₃	H	CH ₃
A-270	CF ₃	CH ₃	H	CH ₃
A-271	OCH ₃	CH ₃	H	CH ₃
A-272	OCF ₃	CH ₃	H	CH ₃
A-273	cyclopropyl	CH ₃	H	CH ₃
A-274	cyclobutyl	CH ₃	H	CH ₃
A-275	cyclopentyl	CH ₃	H	CH ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-276	cyclohexyl	CH ₃	H	CH ₃
A-277	H	CH ₃	CH ₃	CH ₃
A-278	F	CH ₃	CH ₃	CH ₃
A-279	Cl	CH ₃	CH ₃	CH ₃
A-280	Br	CH ₃	CH ₃	CH ₃
A-281	CH ₃	CH ₃	CH ₃	CH ₃
A-282	CF ₃	CH ₃	CH ₃	CH ₃
A-283	OCH ₃	CH ₃	CH ₃	CH ₃
A-284	OCF ₃	CH ₃	CH ₃	CH ₃
A-285	cyclopropyl	CH ₃	CH ₃	CH ₃
A-286	cyclobutyl	CH ₃	CH ₃	CH ₃
A-287	cyclopentyl	CH ₃	CH ₃	CH ₃
A-288	cyclohexyl	CH ₃	CH ₃	CH ₃
A-289	H	H	H	C ₂ H ₅
A-290	F	H	H	C ₂ H ₅
A-291	Cl	H	H	C ₂ H ₅
A-292	Br	H	H	C ₂ H ₅
A-293	CH ₃	H	H	C ₂ H ₅
A-294	CF ₃	H	H	C ₂ H ₅
A-295	OCH ₃	H	H	C ₂ H ₅
A-296	OCF ₃	H	H	C ₂ H ₅
A-297	cyclopropyl	H	H	C ₂ H ₅
A-298	cyclobutyl	H	H	C ₂ H ₅
A-299	cyclopentyl	H	H	C ₂ H ₅
A-300	cyclohexyl	H	H	C ₂ H ₅
A-301	H	CH ₃	H	C ₂ H ₅
A-302	F	CH ₃	H	C ₂ H ₅
A-303	Cl	CH ₃	H	C ₂ H ₅
A-304	Br	CH ₃	H	C ₂ H ₅
A-305	CH ₃	CH ₃	H	C ₂ H ₅

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-306	CF ₃	CH ₃	H	C ₂ H ₅
A-307	OCH ₃	CH ₃	H	C ₂ H ₅
A-308	OCF ₃	CH ₃	H	C ₂ H ₅
A-309	cyclopropyl	CH ₃	H	C ₂ H ₅
A-310	cyclobutyl	CH ₃	H	C ₂ H ₅
A-311	cyclopentyl	CH ₃	H	C ₂ H ₅
A-312	cyclohexyl	CH ₃	H	C ₂ H ₅
A-313	H	CH ₃	CH ₃	C ₂ H ₅
A-314	F	CH ₃	CH ₃	C ₂ H ₅
A-315	Cl	CH ₃	CH ₃	C ₂ H ₅
A-316	Br	CH ₃	CH ₃	C ₂ H ₅
A-317	CH ₃	CH ₃	CH ₃	C ₂ H ₅
A-318	CF ₃	CH ₃	CH ₃	C ₂ H ₅
A-319	OCH ₃	CH ₃	CH ₃	C ₂ H ₅
A-320	OCF ₃	CH ₃	CH ₃	C ₂ H ₅
A-321	cyclopropyl	CH ₃	CH ₃	C ₂ H ₅
A-322	cyclobutyl	CH ₃	CH ₃	C ₂ H ₅
A-323	cyclopentyl	CH ₃	CH ₃	C ₂ H ₅
A-324	cyclohexyl	CH ₃	CH ₃	C ₂ H ₅
A-325	H	H	H	n-propyl
A-326	F	H	H	n-propyl
A-327	Cl	H	H	n-propyl
A-328	Br	H	H	n-propyl
A-329	CH ₃	H	H	n-propyl
A-330	CF ₃	H	H	n-propyl
A-331	OCH ₃	H	H	n-propyl
A-332	OCF ₃	H	H	n-propyl
A-333	cyclopropyl	H	H	n-propyl
A-334	cyclobutyl	H	H	n-propyl
A-335	cyclopentyl	H	H	n-propyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-336	cyclohexyl	H	H	n-propyl
A-337	H	CH ₃	H	n-propyl
A-338	F	CH ₃	H	n-propyl
A-339	Cl	CH ₃	H	n-propyl
A-340	Br	CH ₃	H	n-propyl
A-341	CH ₃	CH ₃	H	n-propyl
A-342	CF ₃	CH ₃	H	n-propyl
A-343	OCH ₃	CH ₃	H	n-propyl
A-344	OCF ₃	CH ₃	H	n-propyl
A-345	cyclopropyl	CH ₃	H	n-propyl
A-346	cyclobutyl	CH ₃	H	n-propyl
A-347	cyclopentyl	CH ₃	H	n-propyl
A-348	cyclohexyl	CH ₃	H	n-propyl
A-349	H	CH ₃	CH ₃	n-propyl
A-350	F	CH ₃	CH ₃	n-propyl
A-351	Cl	CH ₃	CH ₃	n-propyl
A-352	Br	CH ₃	CH ₃	n-propyl
A-353	CH ₃	CH ₃	CH ₃	n-propyl
A-354	CF ₃	CH ₃	CH ₃	n-propyl
A-355	OCH ₃	CH ₃	CH ₃	n-propyl
A-356	OCF ₃	CH ₃	CH ₃	n-propyl
A-357	cyclopropyl	CH ₃	CH ₃	n-propyl
A-358	cyclobutyl	CH ₃	CH ₃	n-propyl
A-359	cyclopentyl	CH ₃	CH ₃	n-propyl
A-360	cyclohexyl	CH ₃	CH ₃	n-propyl
A-361	H	H	H	isopropyl
A-362	F	H	H	isopropyl
A-363	Cl	H	H	isopropyl
A-364	Br	H	H	isopropyl
A-365	CH ₃	H	H	isopropyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-366	CF ₃	H	H	isopropyl
A-367	OCH ₃	H	H	isopropyl
A-368	OCF ₃	H	H	isopropyl
A-369	cyclopropyl	H	H	isopropyl
A-370	cyclobutyl	H	H	isopropyl
A-371	cyclopentyl	H	H	isopropyl
A-372	cyclohexyl	H	H	isopropyl
A-373	H	CH ₃	H	isopropyl
A-374	F	CH ₃	H	isopropyl
A-375	Cl	CH ₃	H	isopropyl
A-376	Br	CH ₃	H	isopropyl
A-377	CH ₃	CH ₃	H	isopropyl
A-378	CF ₃	CH ₃	H	isopropyl
A-379	OCH ₃	CH ₃	H	isopropyl
A-380	OCF ₃	CH ₃	H	isopropyl
A-381	cyclopropyl	CH ₃	H	isopropyl
A-382	cyclobutyl	CH ₃	H	isopropyl
A-383	cyclopentyl	CH ₃	H	isopropyl
A-384	cyclohexyl	CH ₃	H	isopropyl
A-385	H	CH ₃	CH ₃	isopropyl
A-386	F	CH ₃	CH ₃	isopropyl
A-387	Cl	CH ₃	CH ₃	isopropyl
A-388	Br	CH ₃	CH ₃	isopropyl
A-389	CH ₃	CH ₃	CH ₃	isopropyl
A-390	CF ₃	CH ₃	CH ₃	isopropyl
A-391	OCH ₃	CH ₃	CH ₃	isopropyl
A-392	OCF ₃	CH ₃	CH ₃	isopropyl
A-393	cyclopropyl	CH ₃	CH ₃	isopropyl
A-394	cyclobutyl	CH ₃	CH ₃	isopropyl
A-395	cyclopentyl	CH ₃	CH ₃	isopropyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-396	cyclohexyl	CH ₃	CH ₃	isopropyl
A-397	H	H	H	n-butyl
A-398	F	H	H	n-butyl
A-399	Cl	H	H	n-butyl
A-400	Br	H	H	n-butyl
A-401	CH ₃	H	H	n-butyl
A-402	CF ₃	H	H	n-butyl
A-403	OCH ₃	H	H	n-butyl
A-404	OCF ₃	H	H	n-butyl
A-405	cyclopropyl	H	H	n-butyl
A-406	cyclobutyl	H	H	n-butyl
A-407	cyclopentyl	H	H	n-butyl
A-408	cyclohexyl	H	H	n-butyl
A-409	H	CH ₃	H	n-butyl
A-410	F	CH ₃	H	n-butyl
A-411	Cl	CH ₃	H	n-butyl
A-412	Br	CH ₃	H	n-butyl
A-413	CH ₃	CH ₃	H	n-butyl
A-414	CF ₃	CH ₃	H	n-butyl
A-415	OCH ₃	CH ₃	H	n-butyl
A-416	OCF ₃	CH ₃	H	n-butyl
A-417	cyclopropyl	CH ₃	H	n-butyl
A-418	cyclobutyl	CH ₃	H	n-butyl
A-419	cyclopentyl	CH ₃	H	n-butyl
A-420	cyclohexyl	CH ₃	H	n-butyl
A-421	H	CH ₃	CH ₃	n-butyl
A-422	F	CH ₃	CH ₃	n-butyl
A-423	Cl	CH ₃	CH ₃	n-butyl
A-424	Br	CH ₃	CH ₃	n-butyl
A-425	CH ₃	CH ₃	CH ₃	n-butyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-426	CF ₃	CH ₃	CH ₃	n-butyl
A-427	OCH ₃	CH ₃	CH ₃	n-butyl
A-428	OCF ₃	CH ₃	CH ₃	n-butyl
A-429	cyclopropyl	CH ₃	CH ₃	n-butyl
A-430	cyclobutyl	CH ₃	CH ₃	n-butyl
A-431	cyclopentyl	CH ₃	CH ₃	n-butyl
A-432	cyclohexyl	CH ₃	CH ₃	n-butyl
A-433	H	H	H	cyclopropyl
A-434	F	H	H	cyclopropyl
A-435	Cl	H	H	cyclopropyl
A-436	Br	H	H	cyclopropyl
A-437	CH ₃	H	H	cyclopropyl
A-438	CF ₃	H	H	cyclopropyl
A-439	OCH ₃	H	H	cyclopropyl
A-440	OCF ₃	H	H	cyclopropyl
A-441	cyclopropyl	H	H	cyclopropyl
A-442	cyclobutyl	H	H	cyclopropyl
A-443	cyclopentyl	H	H	cyclopropyl
A-444	cyclohexyl	H	H	cyclopropyl
A-445	H	CH ₃	H	cyclopropyl
A-446	F	CH ₃	H	cyclopropyl
A-447	Cl	CH ₃	H	cyclopropyl
A-448	Br	CH ₃	H	cyclopropyl
A-449	CH ₃	CH ₃	H	cyclopropyl
A-450	CF ₃	CH ₃	H	cyclopropyl
A-451	OCH ₃	CH ₃	H	cyclopropyl
A-452	OCF ₃	CH ₃	H	cyclopropyl
A-453	cyclopropyl	CH ₃	H	cyclopropyl
A-454	cyclobutyl	CH ₃	H	cyclopropyl
A-455	cyclopentyl	CH ₃	H	cyclopropyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-456	cyclohexyl	CH ₃	H	cyclopropyl
A-457	H	CH ₃	CH ₃	cyclopropyl
A-458	F	CH ₃	CH ₃	cyclopropyl
A-459	Cl	CH ₃	CH ₃	cyclopropyl
A-460	Br	CH ₃	CH ₃	cyclopropyl
A-461	CH ₃	CH ₃	CH ₃	cyclopropyl
A-462	CF ₃	CH ₃	CH ₃	cyclopropyl
A-463	OCH ₃	CH ₃	CH ₃	cyclopropyl
A-464	OCF ₃	CH ₃	CH ₃	cyclopropyl
A-465	cyclopropyl	CH ₃	CH ₃	cyclopropyl
A-466	cyclobutyl	CH ₃	CH ₃	cyclopropyl
A-467	cyclopentyl	CH ₃	CH ₃	cyclopropyl
A-468	cyclohexyl	CH ₃	CH ₃	cyclopropyl
A-469	H	H	H	cyclobutyl
A-470	F	H	H	cyclobutyl
A-471	Cl	H	H	cyclobutyl
A-472	Br	H	H	cyclobutyl
A-473	CH ₃	H	H	cyclobutyl
A-474	CF ₃	H	H	cyclobutyl
A-475	OCH ₃	H	H	cyclobutyl
A-476	OCF ₃	H	H	cyclobutyl
A-477	cyclopropyl	H	H	cyclobutyl
A-478	cyclobutyl	H	H	cyclobutyl
A-479	cyclopentyl	H	H	cyclobutyl
A-480	cyclohexyl	H	H	cyclobutyl
A-481	H	CH ₃	H	cyclobutyl
A-482	F	CH ₃	H	cyclobutyl
A-483	Cl	CH ₃	H	cyclobutyl
A-484	Br	CH ₃	H	cyclobutyl
A-485	CH ₃	CH ₃	H	cyclobutyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-486	CF ₃	CH ₃	H	cyclobutyl
A-487	OCH ₃	CH ₃	H	cyclobutyl
A-488	OCF ₃	CH ₃	H	cyclobutyl
A-489	cyclopropyl	CH ₃	H	cyclobutyl
A-490	cyclobutyl	CH ₃	H	cyclobutyl
A-491	cyclopentyl	CH ₃	H	cyclobutyl
A-492	cyclohexyl	CH ₃	H	cyclobutyl
A-493	H	CH ₃	CH ₃	cyclobutyl
A-494	F	CH ₃	CH ₃	cyclobutyl
A-495	Cl	CH ₃	CH ₃	cyclobutyl
A-496	Br	CH ₃	CH ₃	cyclobutyl
A-497	CH ₃	CH ₃	CH ₃	cyclobutyl
A-498	CF ₃	CH ₃	CH ₃	cyclobutyl
A-499	OCH ₃	CH ₃	CH ₃	cyclobutyl
A-500	OCF ₃	CH ₃	CH ₃	cyclobutyl
A-501	cyclopropyl	CH ₃	CH ₃	cyclobutyl
A-502	cyclobutyl	CH ₃	CH ₃	cyclobutyl
A-503	cyclopentyl	CH ₃	CH ₃	cyclobutyl
A-504	cyclohexyl	CH ₃	CH ₃	cyclobutyl
A-505	H	H	H	cyclopentyl
A-506	F	H	H	cyclopentyl
A-507	Cl	H	H	cyclopentyl
A-508	Br	H	H	cyclopentyl
A-509	CH ₃	H	H	cyclopentyl
A-510	CF ₃	H	H	cyclopentyl
A-511	OCH ₃	H	H	cyclopentyl
A-512	OCF ₃	H	H	cyclopentyl
A-513	cyclopropyl	H	H	cyclopentyl
A-514	cyclobutyl	H	H	cyclopentyl
A-515	cyclopentyl	H	H	cyclopentyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-516	cyclohexyl	H	H	cyclopentyl
A-517	H	CH ₃	H	cyclopentyl
A-518	F	CH ₃	H	cyclopentyl
A-519	Cl	CH ₃	H	cyclopentyl
A-520	Br	CH ₃	H	cyclopentyl
A-521	CH ₃	CH ₃	H	cyclopentyl
A-522	CF ₃	CH ₃	H	cyclopentyl
A-523	OCH ₃	CH ₃	H	cyclopentyl
A-524	OCF ₃	CH ₃	H	cyclopentyl
A-525	cyclopropyl	CH ₃	H	cyclopentyl
A-526	cyclobutyl	CH ₃	H	cyclopentyl
A-527	cyclopentyl	CH ₃	H	cyclopentyl
A-528	cyclohexyl	CH ₃	H	cyclopentyl
A-529	H	CH ₃	CH ₃	cyclopentyl
A-530	F	CH ₃	CH ₃	cyclopentyl
A-531	Cl	CH ₃	CH ₃	cyclopentyl
A-532	Br	CH ₃	CH ₃	cyclopentyl
A-533	CH ₃	CH ₃	CH ₃	cyclopentyl
A-534	CF ₃	CH ₃	CH ₃	cyclopentyl
A-535	OCH ₃	CH ₃	CH ₃	cyclopentyl
A-536	OCF ₃	CH ₃	CH ₃	cyclopentyl
A-537	cyclopropyl	CH ₃	CH ₃	cyclopentyl
A-538	cyclobutyl	CH ₃	CH ₃	cyclopentyl
A-539	cyclopentyl	CH ₃	CH ₃	cyclopentyl
A-540	cyclohexyl	CH ₃	CH ₃	cyclopentyl
A-541	H	H	H	cyclohexyl
A-542	F	H	H	cyclohexyl
A-543	Cl	H	H	cyclohexyl
A-544	Br	H	H	cyclohexyl
A-545	CH ₃	H	H	cyclohexyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-546	CF ₃	H	H	cyclohexyl
A-547	OCH ₃	H	H	cyclohexyl
A-548	OCF ₃	H	H	cyclohexyl
A-549	cyclopropyl	H	H	cyclohexyl
A-550	cyclobutyl	H	H	cyclohexyl
A-551	cyclopentyl	H	H	cyclohexyl
A-552	cyclohexyl	H	H	cyclohexyl
A-553	H	CH ₃	H	cyclohexyl
A-554	F	CH ₃	H	cyclohexyl
A-555	Cl	CH ₃	H	cyclohexyl
A-556	Br	CH ₃	H	cyclohexyl
A-557	CH ₃	CH ₃	H	cyclohexyl
A-558	CF ₃	CH ₃	H	cyclohexyl
A-559	OCH ₃	CH ₃	H	cyclohexyl
A-560	OCF ₃	CH ₃	H	cyclohexyl
A-561	cyclopropyl	CH ₃	H	cyclohexyl
A-562	cyclobutyl	CH ₃	H	cyclohexyl
A-563	cyclopentyl	CH ₃	H	cyclohexyl
A-564	cyclohexyl	CH ₃	H	cyclohexyl
A-565	H	CH ₃	CH ₃	cyclohexyl
A-566	F	CH ₃	CH ₃	cyclohexyl
A-567	Cl	CH ₃	CH ₃	cyclohexyl
A-568	Br	CH ₃	CH ₃	cyclohexyl
A-569	CH ₃	CH ₃	CH ₃	cyclohexyl
A-570	CF ₃	CH ₃	CH ₃	cyclohexyl
A-571	OCH ₃	CH ₃	CH ₃	cyclohexyl
A-572	OCF ₃	CH ₃	CH ₃	cyclohexyl
A-573	cyclopropyl	CH ₃	CH ₃	cyclohexyl
A-574	cyclobutyl	CH ₃	CH ₃	cyclohexyl
A-575	cyclopentyl	CH ₃	CH ₃	cyclohexyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-576	cyclohexyl	CH ₃	CH ₃	cyclohexyl
A-577	H	H	H	oxetan-3-yl
A-578	F	H	H	oxetan-3-yl
A-579	Cl	H	H	oxetan-3-yl
A-580	Br	H	H	oxetan-3-yl
A-581	CH ₃	H	H	oxetan-3-yl
A-582	CF ₃	H	H	oxetan-3-yl
A-583	OCH ₃	H	H	oxetan-3-yl
A-584	OCF ₃	H	H	oxetan-3-yl
A-585	cyclopropyl	H	H	oxetan-3-yl
A-586	cyclobutyl	H	H	oxetan-3-yl
A-587	cyclopentyl	H	H	oxetan-3-yl
A-588	cyclohexyl	H	H	oxetan-3-yl
A-589	H	CH ₃	H	oxetan-3-yl
A-590	F	CH ₃	H	oxetan-3-yl
A-591	Cl	CH ₃	H	oxetan-3-yl
A-592	Br	CH ₃	H	oxetan-3-yl
A-593	CH ₃	CH ₃	H	oxetan-3-yl
A-594	CF ₃	CH ₃	H	oxetan-3-yl
A-595	OCH ₃	CH ₃	H	oxetan-3-yl
A-596	OCF ₃	CH ₃	H	oxetan-3-yl
A-597	cyclopropyl	CH ₃	H	oxetan-3-yl
A-598	cyclobutyl	CH ₃	H	oxetan-3-yl
A-599	cyclopentyl	CH ₃	H	oxetan-3-yl
A-600	cyclohexyl	CH ₃	H	oxetan-3-yl
A-601	H	CH ₃	CH ₃	oxetan-3-yl
A-602	F	CH ₃	CH ₃	oxetan-3-yl
A-603	Cl	CH ₃	CH ₃	oxetan-3-yl
A-604	Br	CH ₃	CH ₃	oxetan-3-yl
A-605	CH ₃	CH ₃	CH ₃	oxetan-3-yl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-606	CF ₃	CH ₃	CH ₃	oxetan-3-yl
A-607	OCH ₃	CH ₃	CH ₃	oxetan-3-yl
A-608	OCF ₃	CH ₃	CH ₃	oxetan-3-yl
A-609	cyclopropyl	CH ₃	CH ₃	oxetan-3-yl
A-610	cyclobutyl	CH ₃	CH ₃	oxetan-3-yl
A-611	cyclopentyl	CH ₃	CH ₃	oxetan-3-yl
A-612	cyclohexyl	CH ₃	CH ₃	oxetan-3-yl
A-613	H	H	H	-CH ₂ -cyclopropyl
A-614	F	H	H	-CH ₂ -cyclopropyl
A-615	Cl	H	H	-CH ₂ -cyclopropyl
A-616	Br	H	H	-CH ₂ -cyclopropyl
A-617	CH ₃	H	H	-CH ₂ -cyclopropyl
A-618	CF ₃	H	H	-CH ₂ -cyclopropyl
A-619	OCH ₃	H	H	-CH ₂ -cyclopropyl
A-620	OCF ₃	H	H	-CH ₂ -cyclopropyl
A-621	cyclopropyl	H	H	-CH ₂ -cyclopropyl
A-622	cyclobutyl	H	H	-CH ₂ -cyclopropyl
A-623	cyclopentyl	H	H	-CH ₂ -cyclopropyl
A-624	cyclohexyl	H	H	-CH ₂ -cyclopropyl
A-625	H	CH ₃	H	-CH ₂ -cyclopropyl
A-626	F	CH ₃	H	-CH ₂ -cyclopropyl
A-627	Cl	CH ₃	H	-CH ₂ -cyclopropyl
A-628	Br	CH ₃	H	-CH ₂ -cyclopropyl
A-629	CH ₃	CH ₃	H	-CH ₂ -cyclopropyl
A-630	CF ₃	CH ₃	H	-CH ₂ -cyclopropyl
A-631	OCH ₃	CH ₃	H	-CH ₂ -cyclopropyl
A-632	OCF ₃	CH ₃	H	-CH ₂ -cyclopropyl
A-633	cyclopropyl	CH ₃	H	-CH ₂ -cyclopropyl
A-634	cyclobutyl	CH ₃	H	-CH ₂ -cyclopropyl
A-635	cyclopentyl	CH ₃	H	-CH ₂ -cyclopropyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-636	cyclohexyl	CH ₃	H	-CH ₂ -cyclopropyl
A-637	H	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-638	F	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-639	Cl	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-640	Br	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-641	CH ₃	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-642	CF ₃	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-643	OCH ₃	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-644	OCF ₃	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-645	cyclopropyl	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-646	cyclobutyl	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-647	cyclopentyl	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-648	cyclohexyl	CH ₃	CH ₃	-CH ₂ -cyclopropyl
A-649	H	H	H	-CH ₂ -cyclobutyl
A-650	F	H	H	-CH ₂ -cyclobutyl
A-651	Cl	H	H	-CH ₂ -cyclobutyl
A-652	Br	H	H	-CH ₂ -cyclobutyl
A-653	CH ₃	H	H	-CH ₂ -cyclobutyl
A-654	CF ₃	H	H	-CH ₂ -cyclobutyl
A-655	OCH ₃	H	H	-CH ₂ -cyclobutyl
A-656	OCF ₃	H	H	-CH ₂ -cyclobutyl
A-657	cyclopropyl	H	H	-CH ₂ -cyclobutyl
A-658	cyclobutyl	H	H	-CH ₂ -cyclobutyl
A-659	cyclopentyl	H	H	-CH ₂ -cyclobutyl
A-660	cyclohexyl	H	H	-CH ₂ -cyclobutyl
A-661	H	CH ₃	H	-CH ₂ -cyclobutyl
A-662	F	CH ₃	H	-CH ₂ -cyclobutyl
A-663	Cl	CH ₃	H	-CH ₂ -cyclobutyl
A-664	Br	CH ₃	H	-CH ₂ -cyclobutyl
A-665	CH ₃	CH ₃	H	-CH ₂ -cyclobutyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-666	CF ₃	CH ₃	H	-CH ₂ -cyclobutyl
A-667	OCH ₃	CH ₃	H	-CH ₂ -cyclobutyl
A-668	OCF ₃	CH ₃	H	-CH ₂ -cyclobutyl
A-669	cyclopropyl	CH ₃	H	-CH ₂ -cyclobutyl
A-670	cyclobutyl	CH ₃	H	-CH ₂ -cyclobutyl
A-671	cyclopentyl	CH ₃	H	-CH ₂ -cyclobutyl
A-672	cyclohexyl	CH ₃	H	-CH ₂ -cyclobutyl
A-673	H	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-674	F	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-675	Cl	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-676	Br	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-677	CH ₃	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-678	CF ₃	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-679	OCH ₃	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-680	OCF ₃	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-681	cyclopropyl	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-682	cyclobutyl	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-683	cyclopentyl	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-684	cyclohexyl	CH ₃	CH ₃	-CH ₂ -cyclobutyl
A-685	H	H	H	-CH ₂ -cyclopentyl
A-686	F	H	H	-CH ₂ -cyclopentyl
A-687	Cl	H	H	-CH ₂ -cyclopentyl
A-688	Br	H	H	-CH ₂ -cyclopentyl
A-689	CH ₃	H	H	-CH ₂ -cyclopentyl
A-690	CF ₃	H	H	-CH ₂ -cyclopentyl
A-691	OCH ₃	H	H	-CH ₂ -cyclopentyl
A-692	OCF ₃	H	H	-CH ₂ -cyclopentyl
A-693	cyclopropyl	H	H	-CH ₂ -cyclopentyl
A-694	cyclobutyl	H	H	-CH ₂ -cyclopentyl
A-695	cyclopentyl	H	H	-CH ₂ -cyclopentyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-696	cyclohexyl	H	H	-CH ₂ -cyclopentyl
A-697	H	CH ₃	H	-CH ₂ -cyclopentyl
A-698	F	CH ₃	H	-CH ₂ -cyclopentyl
A-699	Cl	CH ₃	H	-CH ₂ -cyclopentyl
A-700	Br	CH ₃	H	-CH ₂ -cyclopentyl
A-701	CH ₃	CH ₃	H	-CH ₂ -cyclopentyl
A-702	CF ₃	CH ₃	H	-CH ₂ -cyclopentyl
A-703	OCH ₃	CH ₃	H	-CH ₂ -cyclopentyl
A-704	OCF ₃	CH ₃	H	-CH ₂ -cyclopentyl
A-705	cyclopropyl	CH ₃	H	-CH ₂ -cyclopentyl
A-706	cyclobutyl	CH ₃	H	-CH ₂ -cyclopentyl
A-707	cyclopentyl	CH ₃	H	-CH ₂ -cyclopentyl
A-708	cyclohexyl	CH ₃	H	-CH ₂ -cyclopentyl
A-709	H	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-710	F	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-711	Cl	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-712	Br	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-713	CH ₃	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-714	CF ₃	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-715	OCH ₃	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-716	OCF ₃	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-717	cyclopropyl	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-718	cyclobutyl	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-719	cyclopentyl	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-720	cyclohexyl	CH ₃	CH ₃	-CH ₂ -cyclopentyl
A-721	H	H	H	-CH ₂ -cyclohexyl
A-722	F	H	H	-CH ₂ -cyclohexyl
A-723	Cl	H	H	-CH ₂ -cyclohexyl
A-724	Br	H	H	-CH ₂ -cyclohexyl
A-725	CH ₃	H	H	-CH ₂ -cyclohexyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-726	CF ₃	H	H	-CH ₂ -cyclohexyl
A-727	OCH ₃	H	H	-CH ₂ -cyclohexyl
A-728	OCF ₃	H	H	-CH ₂ -cyclohexyl
A-729	cyclopropyl	H	H	-CH ₂ -cyclohexyl
A-730	cyclobutyl	H	H	-CH ₂ -cyclohexyl
A-731	cyclopentyl	H	H	-CH ₂ -cyclohexyl
A-732	cyclohexyl	H	H	-CH ₂ -cyclohexyl
A-733	H	CH ₃	H	-CH ₂ -cyclohexyl
A-734	F	CH ₃	H	-CH ₂ -cyclohexyl
A-735	Cl	CH ₃	H	-CH ₂ -cyclohexyl
A-736	Br	CH ₃	H	-CH ₂ -cyclohexyl
A-737	CH ₃	CH ₃	H	-CH ₂ -cyclohexyl
A-738	CF ₃	CH ₃	H	-CH ₂ -cyclohexyl
A-739	OCH ₃	CH ₃	H	-CH ₂ -cyclohexyl
A-740	OCF ₃	CH ₃	H	-CH ₂ -cyclohexyl
A-741	cyclopropyl	CH ₃	H	-CH ₂ -cyclohexyl
A-742	cyclobutyl	CH ₃	H	-CH ₂ -cyclohexyl
A-743	cyclopentyl	CH ₃	H	-CH ₂ -cyclohexyl
A-744	cyclohexyl	CH ₃	H	-CH ₂ -cyclohexyl
A-745	H	CH ₃	CH ₃	-CH ₂ -cyclohexyl
A-746	F	CH ₃	CH ₃	-CH ₂ -cyclohexyl
A-747	Cl	CH ₃	CH ₃	-CH ₂ -cyclohexyl
A-748	Br	CH ₃	CH ₃	-CH ₂ -cyclohexyl
A-749	CH ₃	CH ₃	CH ₃	-CH ₂ -cyclohexyl
A-750	CF ₃	CH ₃	CH ₃	-CH ₂ -cyclohexyl
A-751	OCH ₃	CH ₃	CH ₃	-CH ₂ -cyclohexyl
A-752	OCF ₃	CH ₃	CH ₃	-CH ₂ -cyclohexyl
A-753	cyclopropyl	CH ₃	CH ₃	-CH ₂ -cyclohexyl
A-754	cyclobutyl	CH ₃	CH ₃	-CH ₂ -cyclohexyl
A-755	cyclopentyl	CH ₃	CH ₃	-CH ₂ -cyclohexyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-756	cyclohexyl	CH ₃	CH ₃	-CH ₂ -cyclohexyl
A-757	H	H	H	-C(O)-cyclopropyl
A-758	F	H	H	-C(O)-cyclopropyl
A-759	Cl	H	H	-C(O)-cyclopropyl
A-760	Br	H	H	-C(O)-cyclopropyl
A-761	CH ₃	H	H	-C(O)-cyclopropyl
A-762	CF ₃	H	H	-C(O)-cyclopropyl
A-763	OCH ₃	H	H	-C(O)-cyclopropyl
A-764	OCF ₃	H	H	-C(O)-cyclopropyl
A-765	cyclopropyl	H	H	-C(O)-cyclopropyl
A-766	cyclobutyl	H	H	-C(O)-cyclopropyl
A-767	cyclopentyl	H	H	-C(O)-cyclopropyl
A-768	cyclohexyl	H	H	-C(O)-cyclopropyl
A-769	H	CH ₃	H	-C(O)-cyclopropyl
A-770	F	CH ₃	H	-C(O)-cyclopropyl
A-771	Cl	CH ₃	H	-C(O)-cyclopropyl
A-772	Br	CH ₃	H	-C(O)-cyclopropyl
A-773	CH ₃	CH ₃	H	-C(O)-cyclopropyl
A-774	CF ₃	CH ₃	H	-C(O)-cyclopropyl
A-775	OCH ₃	CH ₃	H	-C(O)-cyclopropyl
A-776	OCF ₃	CH ₃	H	-C(O)-cyclopropyl
A-777	cyclopropyl	CH ₃	H	-C(O)-cyclopropyl
A-778	cyclobutyl	CH ₃	H	-C(O)-cyclopropyl
A-779	cyclopentyl	CH ₃	H	-C(O)-cyclopropyl
A-780	cyclohexyl	CH ₃	H	-C(O)-cyclopropyl
A-781	H	CH ₃	CH ₃	-C(O)-cyclopropyl
A-782	F	CH ₃	CH ₃	-C(O)-cyclopropyl
A-783	Cl	CH ₃	CH ₃	-C(O)-cyclopropyl
A-784	Br	CH ₃	CH ₃	-C(O)-cyclopropyl
A-785	CH ₃	CH ₃	CH ₃	-C(O)-cyclopropyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-786	CF ₃	CH ₃	CH ₃	-C(O)-cyclopropyl
A-787	OCH ₃	CH ₃	CH ₃	-C(O)-cyclopropyl
A-788	OCF ₃	CH ₃	CH ₃	-C(O)-cyclopropyl
A-789	cyclopropyl	CH ₃	CH ₃	-C(O)-cyclopropyl
A-790	cyclobutyl	CH ₃	CH ₃	-C(O)-cyclopropyl
A-791	cyclopentyl	CH ₃	CH ₃	-C(O)-cyclopropyl
A-792	cyclohexyl	CH ₃	CH ₃	-C(O)-cyclopropyl
A-793	H	H	H	-C(O)-cyclobutyl
A-794	F	H	H	-C(O)-cyclobutyl
A-795	Cl	H	H	-C(O)-cyclobutyl
A-796	Br	H	H	-C(O)-cyclobutyl
A-797	CH ₃	H	H	-C(O)-cyclobutyl
A-798	CF ₃	H	H	-C(O)-cyclobutyl
A-799	OCH ₃	H	H	-C(O)-cyclobutyl
A-800	OCF ₃	H	H	-C(O)-cyclobutyl
A-801	cyclopropyl	H	H	-C(O)-cyclobutyl
A-802	cyclobutyl	H	H	-C(O)-cyclobutyl
A-803	cyclopentyl	H	H	-C(O)-cyclobutyl
A-804	cyclohexyl	H	H	-C(O)-cyclobutyl
A-805	H	CH ₃	H	-C(O)-cyclobutyl
A-806	F	CH ₃	H	-C(O)-cyclobutyl
A-807	Cl	CH ₃	H	-C(O)-cyclobutyl
A-808	Br	CH ₃	H	-C(O)-cyclobutyl
A-809	CH ₃	CH ₃	H	-C(O)-cyclobutyl
A-810	CF ₃	CH ₃	H	-C(O)-cyclobutyl
A-811	OCH ₃	CH ₃	H	-C(O)-cyclobutyl
A-812	OCF ₃	CH ₃	H	-C(O)-cyclobutyl
A-813	cyclopropyl	CH ₃	H	-C(O)-cyclobutyl
A-814	cyclobutyl	CH ₃	H	-C(O)-cyclobutyl
A-815	cyclopentyl	CH ₃	H	-C(O)-cyclobutyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-816	cyclohexyl	CH ₃	H	-C(O)-cyclobutyl
A-817	H	CH ₃	CH ₃	-C(O)-cyclobutyl
A-818	F	CH ₃	CH ₃	-C(O)-cyclobutyl
A-819	Cl	CH ₃	CH ₃	-C(O)-cyclobutyl
A-820	Br	CH ₃	CH ₃	-C(O)-cyclobutyl
A-821	CH ₃	CH ₃	CH ₃	-C(O)-cyclobutyl
A-822	CF ₃	CH ₃	CH ₃	-C(O)-cyclobutyl
A-823	OCH ₃	CH ₃	CH ₃	-C(O)-cyclobutyl
A-824	OCF ₃	CH ₃	CH ₃	-C(O)-cyclobutyl
A-825	cyclopropyl	CH ₃	CH ₃	-C(O)-cyclobutyl
A-826	cyclobutyl	CH ₃	CH ₃	-C(O)-cyclobutyl
A-827	cyclopentyl	CH ₃	CH ₃	-C(O)-cyclobutyl
A-828	cyclohexyl	CH ₃	CH ₃	-C(O)-cyclobutyl
A-829	H	H	H	-C(O)-cyclopentyl
A-830	F	H	H	-C(O)-cyclopentyl
A-831	Cl	H	H	-C(O)-cyclopentyl
A-832	Br	H	H	-C(O)-cyclopentyl
A-833	CH ₃	H	H	-C(O)-cyclopentyl
A-834	CF ₃	H	H	-C(O)-cyclopentyl
A-835	OCH ₃	H	H	-C(O)-cyclopentyl
A-836	OCF ₃	H	H	-C(O)-cyclopentyl
A-837	cyclopropyl	H	H	-C(O)-cyclopentyl
A-838	cyclobutyl	H	H	-C(O)-cyclopentyl
A-839	cyclopentyl	H	H	-C(O)-cyclopentyl
A-840	cyclohexyl	H	H	-C(O)-cyclopentyl
A-841	H	CH ₃	H	-C(O)-cyclopentyl
A-842	F	CH ₃	H	-C(O)-cyclopentyl
A-843	Cl	CH ₃	H	-C(O)-cyclopentyl
A-844	Br	CH ₃	H	-C(O)-cyclopentyl
A-845	CH ₃	CH ₃	H	-C(O)-cyclopentyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-846	CF ₃	CH ₃	H	-C(O)-cyclopentyl
A-847	OCH ₃	CH ₃	H	-C(O)-cyclopentyl
A-848	OCF ₃	CH ₃	H	-C(O)-cyclopentyl
A-849	cyclopropyl	CH ₃	H	-C(O)-cyclopentyl
A-850	cyclobutyl	CH ₃	H	-C(O)-cyclopentyl
A-851	cyclopentyl	CH ₃	H	-C(O)-cyclopentyl
A-852	cyclohexyl	CH ₃	H	-C(O)-cyclopentyl
A-853	H	CH ₃	CH ₃	-C(O)-cyclopentyl
A-854	F	CH ₃	CH ₃	-C(O)-cyclopentyl
A-855	Cl	CH ₃	CH ₃	-C(O)-cyclopentyl
A-856	Br	CH ₃	CH ₃	-C(O)-cyclopentyl
A-857	CH ₃	CH ₃	CH ₃	-C(O)-cyclopentyl
A-858	CF ₃	CH ₃	CH ₃	-C(O)-cyclopentyl
A-859	OCH ₃	CH ₃	CH ₃	-C(O)-cyclopentyl
A-860	OCF ₃	CH ₃	CH ₃	-C(O)-cyclopentyl
A-861	cyclopropyl	CH ₃	CH ₃	-C(O)-cyclopentyl
A-862	cyclobutyl	CH ₃	CH ₃	-C(O)-cyclopentyl
A-863	cyclopentyl	CH ₃	CH ₃	-C(O)-cyclopentyl
A-864	cyclohexyl	CH ₃	CH ₃	-C(O)-cyclopentyl
A-865	H	H	H	-C(O)-cyclohexyl
A-866	F	H	H	-C(O)-cyclohexyl
A-867	Cl	H	H	-C(O)-cyclohexyl
A-868	Br	H	H	-C(O)-cyclohexyl
A-869	CH ₃	H	H	-C(O)-cyclohexyl
A-870	CF ₃	H	H	-C(O)-cyclohexyl
A-871	OCH ₃	H	H	-C(O)-cyclohexyl
A-872	OCF ₃	H	H	-C(O)-cyclohexyl
A-873	cyclopropyl	H	H	-C(O)-cyclohexyl
A-874	cyclobutyl	H	H	-C(O)-cyclohexyl
A-875	cyclopentyl	H	H	-C(O)-cyclohexyl

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-876	cyclohexyl	H	H	-C(O)-cyclohexyl
A-877	H	CH ₃	H	-C(O)-cyclohexyl
A-878	F	CH ₃	H	-C(O)-cyclohexyl
A-879	Cl	CH ₃	H	-C(O)-cyclohexyl
A-880	Br	CH ₃	H	-C(O)-cyclohexyl
A-881	CH ₃	CH ₃	H	-C(O)-cyclohexyl
A-882	CF ₃	CH ₃	H	-C(O)-cyclohexyl
A-883	OCH ₃	CH ₃	H	-C(O)-cyclohexyl
A-884	OCF ₃	CH ₃	H	-C(O)-cyclohexyl
A-885	cyclopropyl	CH ₃	H	-C(O)-cyclohexyl
A-886	cyclobutyl	CH ₃	H	-C(O)-cyclohexyl
A-887	cyclopentyl	CH ₃	H	-C(O)-cyclohexyl
A-888	cyclohexyl	CH ₃	H	-C(O)-cyclohexyl
A-889	H	CH ₃	CH ₃	-C(O)-cyclohexyl
A-890	F	CH ₃	CH ₃	-C(O)-cyclohexyl
A-891	Cl	CH ₃	CH ₃	-C(O)-cyclohexyl
A-892	Br	CH ₃	CH ₃	-C(O)-cyclohexyl
A-893	CH ₃	CH ₃	CH ₃	-C(O)-cyclohexyl
A-894	CF ₃	CH ₃	CH ₃	-C(O)-cyclohexyl
A-895	OCH ₃	CH ₃	CH ₃	-C(O)-cyclohexyl
A-896	OCF ₃	CH ₃	CH ₃	-C(O)-cyclohexyl
A-897	cyclopropyl	CH ₃	CH ₃	-C(O)-cyclohexyl
A-898	cyclobutyl	CH ₃	CH ₃	-C(O)-cyclohexyl
A-899	cyclopentyl	CH ₃	CH ₃	-C(O)-cyclohexyl
A-900	cyclohexyl	CH ₃	CH ₃	-C(O)-cyclohexyl
A-901	H	H	-(CH ₂) ₃ -	
A-902	F	H	-(CH ₂) ₃ -	
A-903	Cl	H	-(CH ₂) ₃ -	
A-904	Br	H	-(CH ₂) ₃ -	
A-905	CH ₃	H	-(CH ₂) ₃ -	

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-906	CF ₃	H		-(CH ₂) ₃ -
A-907	OCH ₃	H		-(CH ₂) ₃ -
A-908	OCF ₃	H		-(CH ₂) ₃ -
A-909	cyclopropyl	H		-(CH ₂) ₃ -
A-910	cyclobutyl	H		-(CH ₂) ₃ -
A-911	cyclopentyl	H		-(CH ₂) ₃ -
A-912	cyclohexyl	H		-(CH ₂) ₃ -
A-913	H	CH ₃		-(CH ₂) ₃ -
A-914	F	CH ₃		-(CH ₂) ₃ -
A-915	Cl	CH ₃		-(CH ₂) ₃ -
A-916	Br	CH ₃		-(CH ₂) ₃ -
A-917	CH ₃	CH ₃		-(CH ₂) ₃ -
A-918	CF ₃	CH ₃		-(CH ₂) ₃ -
A-919	OCH ₃	CH ₃		-(CH ₂) ₃ -
A-920	OCF ₃	CH ₃		-(CH ₂) ₃ -
A-921	cyclopropyl	CH ₃		-(CH ₂) ₃ -
A-922	cyclobutyl	CH ₃		-(CH ₂) ₃ -
A-923	cyclopentyl	CH ₃		-(CH ₂) ₃ -
A-924	cyclohexyl	CH ₃		-(CH ₂) ₃ -
A-925	H	D		-(CH ₂) ₃ -
A-926	F	D		-(CH ₂) ₃ -
A-927	Cl	D		-(CH ₂) ₃ -
A-928	Br	D		-(CH ₂) ₃ -
A-929	CH ₃	D		-(CH ₂) ₃ -
A-930	CF ₃	D		-(CH ₂) ₃ -
A-931	OCH ₃	D		-(CH ₂) ₃ -
A-932	OCF ₃	D		-(CH ₂) ₃ -
A-933	cyclopropyl	D		-(CH ₂) ₃ -
A-934	cyclobutyl	D		-(CH ₂) ₃ -
A-935	cyclopentyl	D		-(CH ₂) ₃ -

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-936	cyclohexyl	D		-(CH ₂) ₃ -
A-937	H	H		-(CH ₂) ₄ -
A-938	F	H		-(CH ₂) ₄ -
A-939	Cl	H		-(CH ₂) ₄ -
A-940	Br	H		-(CH ₂) ₄ -
A-941	CH ₃	H		-(CH ₂) ₄ -
A-942	CF ₃	H		-(CH ₂) ₄ -
A-943	OCH ₃	H		-(CH ₂) ₄ -
A-944	OCF ₃	H		-(CH ₂) ₄ -
A-945	cyclopropyl	H		-(CH ₂) ₄ -
A-946	cyclobutyl	H		-(CH ₂) ₄ -
A-947	cyclopentyl	H		-(CH ₂) ₄ -
A-948	cyclohexyl	H		-(CH ₂) ₄ -
A-949	H	CH ₃		-(CH ₂) ₄ -
A-950	F	CH ₃		-(CH ₂) ₄ -
A-951	Cl	CH ₃		-(CH ₂) ₄ -
A-952	Br	CH ₃		-(CH ₂) ₄ -
A-953	CH ₃	CH ₃		-(CH ₂) ₄ -
A-954	CF ₃	CH ₃		-(CH ₂) ₄ -
A-955	OCH ₃	CH ₃		-(CH ₂) ₄ -
A-956	OCF ₃	CH ₃		-(CH ₂) ₄ -
A-957	cyclopropyl	CH ₃		-(CH ₂) ₄ -
A-958	cyclobutyl	CH ₃		-(CH ₂) ₄ -
A-959	cyclopentyl	CH ₃		-(CH ₂) ₄ -
A-960	cyclohexyl	CH ₃		-(CH ₂) ₄ -
A-961	H	D		-(CH ₂) ₄ -
A-962	F	D		-(CH ₂) ₄ -
A-963	Cl	D		-(CH ₂) ₄ -
A-964	Br	D		-(CH ₂) ₄ -
A-965	CH ₃	D		-(CH ₂) ₄ -

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-966	CF ₃	D		-(CH ₂) ₄ -
A-967	OCH ₃	D		-(CH ₂) ₄ -
A-968	OCF ₃	D		-(CH ₂) ₄ -
A-969	cyclopropyl	D		-(CH ₂) ₄ -
A-970	cyclobutyl	D		-(CH ₂) ₄ -
A-971	cyclopentyl	D		-(CH ₂) ₄ -
A-972	cyclohexyl	D		-(CH ₂) ₄ -
A-973	H	H		-(CH ₂) ₅ -
A-974	F	H		-(CH ₂) ₅ -
A-975	Cl	H		-(CH ₂) ₅ -
A-976	Br	H		-(CH ₂) ₅ -
A-977	CH ₃	H		-(CH ₂) ₅ -
A-978	CF ₃	H		-(CH ₂) ₅ -
A-979	OCH ₃	H		-(CH ₂) ₅ -
A-980	OCF ₃	H		-(CH ₂) ₅ -
A-981	cyclopropyl	H		-(CH ₂) ₅ -
A-982	cyclobutyl	H		-(CH ₂) ₅ -
A-983	cyclopentyl	H		-(CH ₂) ₅ -
A-984	cyclohexyl	H		-(CH ₂) ₅ -
A-985	H	CH ₃		-(CH ₂) ₅ -
A-986	F	CH ₃		-(CH ₂) ₅ -
A-987	Cl	CH ₃		-(CH ₂) ₅ -
A-988	Br	CH ₃		-(CH ₂) ₅ -
A-989	CH ₃	CH ₃		-(CH ₂) ₅ -
A-990	CF ₃	CH ₃		-(CH ₂) ₅ -
A-991	OCH ₃	CH ₃		-(CH ₂) ₅ -
A-992	OCF ₃	CH ₃		-(CH ₂) ₅ -
A-993	cyclopropyl	CH ₃		-(CH ₂) ₅ -
A-994	cyclobutyl	CH ₃		-(CH ₂) ₅ -
A-995	cyclopentyl	CH ₃		-(CH ₂) ₅ -


No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-996	cyclohexyl	CH ₃		-(CH ₂) ₅ -
A-997	H	D		-(CH ₂) ₅ -
A-998	F	D		-(CH ₂) ₅ -
A-999	Cl	D		-(CH ₂) ₅ -
A-1000	Br	D		-(CH ₂) ₅ -
A-1001	CH ₃	D		-(CH ₂) ₅ -
A-1002	CF ₃	D		-(CH ₂) ₅ -
A-1003	OCH ₃	D		-(CH ₂) ₅ -
A-1004	OCF ₃	D		-(CH ₂) ₅ -
A-1005	cyclopropyl	D		-(CH ₂) ₅ -
A-1006	cyclobutyl	D		-(CH ₂) ₅ -
A-1007	cyclopentyl	D		-(CH ₂) ₅ -
A-1008	cyclohexyl	D		-(CH ₂) ₅ -
A-1009	F	H		-(CH ₂ -CH-(μ-CH ₂)-CH)-***
A-1010	Cl	H		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1011	Br	H		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1012	CH ₃	H		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1013	CF ₃	H		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1014	OCH ₃	H		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1015	OCF ₃	H		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1016	cyclopropyl	H		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1017	cyclobutyl	H		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1018	cyclopentyl	H		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1019	cyclohexyl	H		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1020	H	CH ₃		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1021	F	CH ₃		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1022	Cl	CH ₃		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1023	Br	CH ₃		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1024	CH ₃	CH ₃		-(CH ₂ -CH-(μ-CH ₂)-CH)-
A-1025	CF ₃	CH ₃		-(CH ₂ -CH-(μ-CH ₂)-CH)-

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁶
A-1026	OCH ₃	CH ₃	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1027	OCF ₃	CH ₃	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1028	cyclopropyl	CH ₃	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1029	cyclobutyl	CH ₃	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1030	cyclopentyl	CH ₃	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1031	cyclohexyl	CH ₃	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1032	H	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1033	F	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1034	Cl	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1035	Br	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1036	CH ₃	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1037	CF ₃	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1038	OCH ₃	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1039	OCF ₃	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1040	cyclopropyl	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1041	cyclobutyl	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1042	cyclopentyl	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	
A-1043	cyclohexyl	D	-(CH ₂ -CH-(μ-CH ₂)-CH)-	

* C₆H₅ = phenyl

** C₆H₄CH₃ = 4-methylphenyl (p-tolyl)

*** -(CH₂-CH-(μ-CH₂)-CH)- =



(# = attachment point to the atoms carrying R⁶ (left #) and R^{5a} (right #))

Table B

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-1	H	H	H	CH ₃	CH ₃
B-2	F	H	H	CH ₃	CH ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-3	Cl	H	H	CH ₃	CH ₃
B-4	Br	H	H	CH ₃	CH ₃
B-5	CH ₃	H	H	CH ₃	CH ₃
B-6	CF ₃	H	H	CH ₃	CH ₃
B-7	OCH ₃	H	H	CH ₃	CH ₃
B-8	OCF ₃	H	H	CH ₃	CH ₃
B-9	H	F	H	CH ₃	CH ₃
B-10	F	F	H	CH ₃	CH ₃
B-11	Cl	F	H	CH ₃	CH ₃
B-12	Br	F	H	CH ₃	CH ₃
B-13	CH ₃	F	H	CH ₃	CH ₃
B-14	CF ₃	F	H	CH ₃	CH ₃
B-15	OCH ₃	F	H	CH ₃	CH ₃
B-16	OCF ₃	F	H	CH ₃	CH ₃
B-17	H	Cl	H	CH ₃	CH ₃
B-18	F	Cl	H	CH ₃	CH ₃
B-19	Cl	Cl	H	CH ₃	CH ₃
B-20	Br	Cl	H	CH ₃	CH ₃
B-21	CH ₃	Cl	H	CH ₃	CH ₃
B-22	CF ₃	Cl	H	CH ₃	CH ₃
B-23	OCH ₃	Cl	H	CH ₃	CH ₃
B-24	OCF ₃	Cl	H	CH ₃	CH ₃
B-25	H	CH ₃	H	CH ₃	CH ₃
B-26	F	CH ₃	H	CH ₃	CH ₃
B-27	Cl	CH ₃	H	CH ₃	CH ₃
B-28	Br	CH ₃	H	CH ₃	CH ₃
B-29	CH ₃	CH ₃	H	CH ₃	CH ₃
B-30	CF ₃	CH ₃	H	CH ₃	CH ₃
B-31	OCH ₃	CH ₃	H	CH ₃	CH ₃
B-32	OCF ₃	CH ₃	H	CH ₃	CH ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-33	H	CH ₃	CH ₃	CH ₃	CH ₃
B-34	F	CH ₃	CH ₃	CH ₃	CH ₃
B-35	Cl	CH ₃	CH ₃	CH ₃	CH ₃
B-36	Br	CH ₃	CH ₃	CH ₃	CH ₃
B-37	CH ₃	CH ₃	CH ₃	CH ₃	CH ₃
B-38	CF ₃	CH ₃	CH ₃	CH ₃	CH ₃
B-39	OCH ₃	CH ₃	CH ₃	CH ₃	CH ₃
B-40	OCF ₃	CH ₃	CH ₃	CH ₃	CH ₃
B-41	H	H	H	CH ₃	CH ₂ CH ₃
B-42	F	H	H	CH ₃	CH ₂ CH ₃
B-43	Cl	H	H	CH ₃	CH ₂ CH ₃
B-44	Br	H	H	CH ₃	CH ₂ CH ₃
B-45	CH ₃	H	H	CH ₃	CH ₂ CH ₃
B-46	CF ₃	H	H	CH ₃	CH ₂ CH ₃
B-47	OCH ₃	H	H	CH ₃	CH ₂ CH ₃
B-48	OCF ₃	H	H	CH ₃	CH ₂ CH ₃
B-49	H	F	H	CH ₃	CH ₂ CH ₃
B-50	F	F	H	CH ₃	CH ₂ CH ₃
B-51	Cl	F	H	CH ₃	CH ₂ CH ₃
B-52	Br	F	H	CH ₃	CH ₂ CH ₃
B-53	CH ₃	F	H	CH ₃	CH ₂ CH ₃
B-54	CF ₃	F	H	CH ₃	CH ₂ CH ₃
B-55	OCH ₃	F	H	CH ₃	CH ₂ CH ₃
B-56	OCF ₃	F	H	CH ₃	CH ₂ CH ₃
B-57	H	Cl	H	CH ₃	CH ₂ CH ₃
B-58	F	Cl	H	CH ₃	CH ₂ CH ₃
B-59	Cl	Cl	H	CH ₃	CH ₂ CH ₃
B-60	Br	Cl	H	CH ₃	CH ₂ CH ₃
B-61	CH ₃	Cl	H	CH ₃	CH ₂ CH ₃
B-62	CF ₃	Cl	H	CH ₃	CH ₂ CH ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-63	OCH ₃	Cl	H	CH ₃	CH ₂ CH ₃
B-64	OCF ₃	Cl	H	CH ₃	CH ₂ CH ₃
B-65	H	CH ₃	H	CH ₃	CH ₂ CH ₃
B-66	F	CH ₃	H	CH ₃	CH ₂ CH ₃
B-67	Cl	CH ₃	H	CH ₃	CH ₂ CH ₃
B-68	Br	CH ₃	H	CH ₃	CH ₂ CH ₃
B-69	CH ₃	CH ₃	H	CH ₃	CH ₂ CH ₃
B-70	CF ₃	CH ₃	H	CH ₃	CH ₂ CH ₃
B-71	OCH ₃	CH ₃	H	CH ₃	CH ₂ CH ₃
B-72	OCF ₃	CH ₃	H	CH ₃	CH ₂ CH ₃
B-73	H	CH ₃	CH ₃	CH ₃	CH ₂ CH ₃
B-74	F	CH ₃	CH ₃	CH ₃	CH ₂ CH ₃
B-75	Cl	CH ₃	CH ₃	CH ₃	CH ₂ CH ₃
B-76	Br	CH ₃	CH ₃	CH ₃	CH ₂ CH ₃
B-77	CH ₃	CH ₃	CH ₃	CH ₃	CH ₂ CH ₃
B-78	CF ₃	CH ₃	CH ₃	CH ₃	CH ₂ CH ₃
B-79	OCH ₃	CH ₃	CH ₃	CH ₃	CH ₂ CH ₃
B-80	OCF ₃	CH ₃	CH ₃	CH ₃	CH ₂ CH ₃
B-81	H	H	H	CH ₃	CF ₃
B-82	F	H	H	CH ₃	CF ₃
B-83	Cl	H	H	CH ₃	CF ₃
B-84	Br	H	H	CH ₃	CF ₃
B-85	CH ₃	H	H	CH ₃	CF ₃
B-86	CF ₃	H	H	CH ₃	CF ₃
B-87	OCH ₃	H	H	CH ₃	CF ₃
B-88	OCF ₃	H	H	CH ₃	CF ₃
B-89	H	F	H	CH ₃	CF ₃
B-90	F	F	H	CH ₃	CF ₃
B-91	Cl	F	H	CH ₃	CF ₃
B-92	Br	F	H	CH ₃	CF ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-93	CH ₃	F	H	CH ₃	CF ₃
B-94	CF ₃	F	H	CH ₃	CF ₃
B-95	OCH ₃	F	H	CH ₃	CF ₃
B-96	OCF ₃	F	H	CH ₃	CF ₃
B-97	H	Cl	H	CH ₃	CF ₃
B-98	F	Cl	H	CH ₃	CF ₃
B-99	Cl	Cl	H	CH ₃	CF ₃
B-100	Br	Cl	H	CH ₃	CF ₃
B-101	CH ₃	Cl	H	CH ₃	CF ₃
B-102	CF ₃	Cl	H	CH ₃	CF ₃
B-103	OCH ₃	Cl	H	CH ₃	CF ₃
B-104	OCF ₃	Cl	H	CH ₃	CF ₃
B-105	H	CH ₃	H	CH ₃	CF ₃
B-106	F	CH ₃	H	CH ₃	CF ₃
B-107	Cl	CH ₃	H	CH ₃	CF ₃
B-108	Br	CH ₃	H	CH ₃	CF ₃
B-109	CH ₃	CH ₃	H	CH ₃	CF ₃
B-110	CF ₃	CH ₃	H	CH ₃	CF ₃
B-111	OCH ₃	CH ₃	H	CH ₃	CF ₃
B-112	OCF ₃	CH ₃	H	CH ₃	CF ₃
B-113	H	CH ₃	CH ₃	CH ₃	CF ₃
B-114	F	CH ₃	CH ₃	CH ₃	CF ₃
B-115	Cl	CH ₃	CH ₃	CH ₃	CF ₃
B-116	Br	CH ₃	CH ₃	CH ₃	CF ₃
B-117	CH ₃	CH ₃	CH ₃	CH ₃	CF ₃
B-118	CF ₃	CH ₃	CH ₃	CH ₃	CF ₃
B-119	OCH ₃	CH ₃	CH ₃	CH ₃	CF ₃
B-120	OCF ₃	CH ₃	CH ₃	CH ₃	CF ₃
B-121	H	H	H	CF ₃	CF ₃
B-122	F	H	H	CF ₃	CF ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-123	Cl	H	H	CF ₃	CF ₃
B-124	Br	H	H	CF ₃	CF ₃
B-125	CH ₃	H	H	CF ₃	CF ₃
B-126	CF ₃	H	H	CF ₃	CF ₃
B-127	OCH ₃	H	H	CF ₃	CF ₃
B-128	OCF ₃	H	H	CF ₃	CF ₃
B-129	H	F	H	CF ₃	CF ₃
B-130	F	F	H	CF ₃	CF ₃
B-131	Cl	F	H	CF ₃	CF ₃
B-132	Br	F	H	CF ₃	CF ₃
B-133	CH ₃	F	H	CF ₃	CF ₃
B-134	CF ₃	F	H	CF ₃	CF ₃
B-135	OCH ₃	F	H	CF ₃	CF ₃
B-136	OCF ₃	F	H	CF ₃	CF ₃
B-137	H	Cl	H	CF ₃	CF ₃
B-138	F	Cl	H	CF ₃	CF ₃
B-139	Cl	Cl	H	CF ₃	CF ₃
B-140	Br	Cl	H	CF ₃	CF ₃
B-141	CH ₃	Cl	H	CF ₃	CF ₃
B-142	CF ₃	Cl	H	CF ₃	CF ₃
B-143	OCH ₃	Cl	H	CF ₃	CF ₃
B-144	OCF ₃	Cl	H	CF ₃	CF ₃
B-145	H	CH ₃	H	CF ₃	CF ₃
B-146	F	CH ₃	H	CF ₃	CF ₃
B-147	Cl	CH ₃	H	CF ₃	CF ₃
B-148	Br	CH ₃	H	CF ₃	CF ₃
B-149	CH ₃	CH ₃	H	CF ₃	CF ₃
B-150	CF ₃	CH ₃	H	CF ₃	CF ₃
B-151	OCH ₃	CH ₃	H	CF ₃	CF ₃
B-152	OCF ₃	CH ₃	H	CF ₃	CF ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-153	H	CH ₃	CH ₃	CF ₃	CF ₃
B-154	F	CH ₃	CH ₃	CF ₃	CF ₃
B-155	Cl	CH ₃	CH ₃	CF ₃	CF ₃
B-156	Br	CH ₃	CH ₃	CF ₃	CF ₃
B-157	CH ₃	CH ₃	CH ₃	CF ₃	CF ₃
B-158	CF ₃	CH ₃	CH ₃	CF ₃	CF ₃
B-159	OCH ₃	CH ₃	CH ₃	CF ₃	CF ₃
B-160	OCF ₃	CH ₃	CH ₃	CF ₃	CF ₃
B-161	H	H	H	CH ₃	C ₆ H ₅ *
B-162	F	H	H	CH ₃	C ₆ H ₅
B-163	Cl	H	H	CH ₃	C ₆ H ₅
B-164	Br	H	H	CH ₃	C ₆ H ₅
B-165	CH ₃	H	H	CH ₃	C ₆ H ₅
B-166	CF ₃	H	H	CH ₃	C ₆ H ₅
B-167	OCH ₃	H	H	CH ₃	C ₆ H ₅
B-168	OCF ₃	H	H	CH ₃	C ₆ H ₅
B-169	H	F	H	CH ₃	C ₆ H ₅
B-170	F	F	H	CH ₃	C ₆ H ₅
B-171	Cl	F	H	CH ₃	C ₆ H ₅
B-172	Br	F	H	CH ₃	C ₆ H ₅
B-173	CH ₃	F	H	CH ₃	C ₆ H ₅
B-174	CF ₃	F	H	CH ₃	C ₆ H ₅
B-175	OCH ₃	F	H	CH ₃	C ₆ H ₅
B-176	OCF ₃	F	H	CH ₃	C ₆ H ₅
B-177	H	Cl	H	CH ₃	C ₆ H ₅
B-178	F	Cl	H	CH ₃	C ₆ H ₅
B-179	Cl	Cl	H	CH ₃	C ₆ H ₅
B-180	Br	Cl	H	CH ₃	C ₆ H ₅
B-181	CH ₃	Cl	H	CH ₃	C ₆ H ₅
B-182	CF ₃	Cl	H	CH ₃	C ₆ H ₅

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-183	OCH ₃	Cl	H	CH ₃	C ₆ H ₅
B-184	OCF ₃	Cl	H	CH ₃	C ₆ H ₅
B-185	H	CH ₃	H	CH ₃	C ₆ H ₅
B-186	F	CH ₃	H	CH ₃	C ₆ H ₅
B-187	Cl	CH ₃	H	CH ₃	C ₆ H ₅
B-188	Br	CH ₃	H	CH ₃	C ₆ H ₅
B-189	CH ₃	CH ₃	H	CH ₃	C ₆ H ₅
B-190	CF ₃	CH ₃	H	CH ₃	C ₆ H ₅
B-191	OCH ₃	CH ₃	H	CH ₃	C ₆ H ₅
B-192	OCF ₃	CH ₃	H	CH ₃	C ₆ H ₅
B-193	H	CH ₃	CH ₃	CH ₃	C ₆ H ₅
B-194	F	CH ₃	CH ₃	CH ₃	C ₆ H ₅
B-195	Cl	CH ₃	CH ₃	CH ₃	C ₆ H ₅
B-196	Br	CH ₃	CH ₃	CH ₃	C ₆ H ₅
B-197	CH ₃	CH ₃	CH ₃	CH ₃	C ₆ H ₅
B-198	CF ₃	CH ₃	CH ₃	CH ₃	C ₆ H ₅
B-199	OCH ₃	CH ₃	CH ₃	CH ₃	C ₆ H ₅
B-200	OCF ₃	CH ₃	CH ₃	CH ₃	C ₆ H ₅
B-201	H	H	H	CH ₃	OCH ₃
B-202	F	H	H	CH ₃	OCH ₃
B-203	Cl	H	H	CH ₃	OCH ₃
B-204	Br	H	H	CH ₃	OCH ₃
B-205	CH ₃	H	H	CH ₃	OCH ₃
B-206	CF ₃	H	H	CH ₃	OCH ₃
B-207	OCH ₃	H	H	CH ₃	OCH ₃
B-208	OCF ₃	H	H	CH ₃	OCH ₃
B-209	H	F	H	CH ₃	OCH ₃
B-210	F	F	H	CH ₃	OCH ₃
B-211	Cl	F	H	CH ₃	OCH ₃
B-212	Br	F	H	CH ₃	OCH ₃

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-213	CH ₃	F	H	CH ₃	OCH ₃
B-214	CF ₃	F	H	CH ₃	OCH ₃
B-215	OCH ₃	F	H	CH ₃	OCH ₃
B-216	OCF ₃	F	H	CH ₃	OCH ₃
B-217	H	Cl	H	CH ₃	OCH ₃
B-218	F	Cl	H	CH ₃	OCH ₃
B-219	Cl	Cl	H	CH ₃	OCH ₃
B-220	Br	Cl	H	CH ₃	OCH ₃
B-221	CH ₃	Cl	H	CH ₃	OCH ₃
B-222	CF ₃	Cl	H	CH ₃	OCH ₃
B-223	OCH ₃	Cl	H	CH ₃	OCH ₃
B-224	OCF ₃	Cl	H	CH ₃	OCH ₃
B-225	H	CH ₃	H	CH ₃	OCH ₃
B-226	F	CH ₃	H	CH ₃	OCH ₃
B-227	Cl	CH ₃	H	CH ₃	OCH ₃
B-228	Br	CH ₃	H	CH ₃	OCH ₃
B-229	CH ₃	CH ₃	H	CH ₃	OCH ₃
B-230	CF ₃	CH ₃	H	CH ₃	OCH ₃
B-231	OCH ₃	CH ₃	H	CH ₃	OCH ₃
B-232	OCF ₃	CH ₃	H	CH ₃	OCH ₃
B-233	H	CH ₃	CH ₃	CH ₃	OCH ₃
B-234	F	CH ₃	CH ₃	CH ₃	OCH ₃
B-235	Cl	CH ₃	CH ₃	CH ₃	OCH ₃
B-236	Br	CH ₃	CH ₃	CH ₃	OCH ₃
B-237	CH ₃	CH ₃	CH ₃	CH ₃	OCH ₃
B-238	CF ₃	CH ₃	CH ₃	CH ₃	OCH ₃
B-239	OCH ₃	CH ₃	CH ₃	CH ₃	OCH ₃
B-240	OCF ₃	CH ₃	CH ₃	CH ₃	OCH ₃
B-241	H	H	H	CH ₃	OH
B-242	F	H	H	CH ₃	OH

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-243	Cl	H	H	CH ₃	OH
B-244	Br	H	H	CH ₃	OH
B-245	CH ₃	H	H	CH ₃	OH
B-246	CF ₃	H	H	CH ₃	OH
B-247	OCH ₃	H	H	CH ₃	OH
B-248	OCF ₃	H	H	CH ₃	OH
B-249	H	F	H	CH ₃	OH
B-250	F	F	H	CH ₃	OH
B-251	Cl	F	H	CH ₃	OH
B-252	Br	F	H	CH ₃	OH
B-253	CH ₃	F	H	CH ₃	OH
B-254	CF ₃	F	H	CH ₃	OH
B-255	OCH ₃	F	H	CH ₃	OH
B-256	OCF ₃	F	H	CH ₃	OH
B-257	H	Cl	H	CH ₃	OH
B-258	F	Cl	H	CH ₃	OH
B-259	Cl	Cl	H	CH ₃	OH
B-260	Br	Cl	H	CH ₃	OH
B-261	CH ₃	Cl	H	CH ₃	OH
B-262	CF ₃	Cl	H	CH ₃	OH
B-263	OCH ₃	Cl	H	CH ₃	OH
B-264	OCF ₃	Cl	H	CH ₃	OH
B-265	H	CH ₃	H	CH ₃	OH
B-266	F	CH ₃	H	CH ₃	OH
B-267	Cl	CH ₃	H	CH ₃	OH
B-268	Br	CH ₃	H	CH ₃	OH
B-269	CH ₃	CH ₃	H	CH ₃	OH
B-270	CF ₃	CH ₃	H	CH ₃	OH
B-271	OCH ₃	CH ₃	H	CH ₃	OH
B-272	OCF ₃	CH ₃	H	CH ₃	OH

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-273	H	CH ₃	CH ₃	CH ₃	OH
B-274	F	CH ₃	CH ₃	CH ₃	OH
B-275	Cl	CH ₃	CH ₃	CH ₃	OH
B-276	Br	CH ₃	CH ₃	CH ₃	OH
B-277	CH ₃	CH ₃	CH ₃	CH ₃	OH
B-278	CF ₃	CH ₃	CH ₃	CH ₃	OH
B-279	OCH ₃	CH ₃	CH ₃	CH ₃	OH
B-280	OCF ₃	CH ₃	CH ₃	CH ₃	OH
B-281	H	H	H	-(CH ₂) ₂ -	
B-282	F	H	H	-(CH ₂) ₂ -	
B-283	Cl	H	H	-(CH ₂) ₂ -	
B-284	Br	H	H	-(CH ₂) ₂ -	
B-285	CH ₃	H	H	-(CH ₂) ₂ -	
B-286	CF ₃	H	H	-(CH ₂) ₂ -	
B-287	OCH ₃	H	H	-(CH ₂) ₂ -	
B-288	OCF ₃	H	H	-(CH ₂) ₂ -	
B-289	H	F	H	-(CH ₂) ₂ -	
B-290	F	F	H	-(CH ₂) ₂ -	
B-291	Cl	F	H	-(CH ₂) ₂ -	
B-292	Br	F	H	-(CH ₂) ₂ -	
B-293	CH ₃	F	H	-(CH ₂) ₂ -	
B-294	CF ₃	F	H	-(CH ₂) ₂ -	
B-295	OCH ₃	F	H	-(CH ₂) ₂ -	
B-296	OCF ₃	F	H	-(CH ₂) ₂ -	
B-297	H	Cl	H	-(CH ₂) ₂ -	
B-298	F	Cl	H	-(CH ₂) ₂ -	
B-299	Cl	Cl	H	-(CH ₂) ₂ -	
B-300	Br	Cl	H	-(CH ₂) ₂ -	
B-301	CH ₃	Cl	H	-(CH ₂) ₂ -	
B-302	CF ₃	Cl	H	-(CH ₂) ₂ -	

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-303	OCH ₃	Cl	H		-(CH ₂) ₂ -
B-304	OCF ₃	Cl	H		-(CH ₂) ₂ -
B-305	H	CH ₃	H		-(CH ₂) ₂ -
B-306	F	CH ₃	H		-(CH ₂) ₂ -
B-307	Cl	CH ₃	H		-(CH ₂) ₂ -
B-308	Br	CH ₃	H		-(CH ₂) ₂ -
B-309	CH ₃	CH ₃	H		-(CH ₂) ₂ -
B-310	CF ₃	CH ₃	H		-(CH ₂) ₂ -
B-311	OCH ₃	CH ₃	H		-(CH ₂) ₂ -
B-312	OCF ₃	CH ₃	H		-(CH ₂) ₂ -
B-313	H	CH ₃	CH ₃		-(CH ₂) ₂ -
B-314	F	CH ₃	CH ₃		-(CH ₂) ₂ -
B-315	Cl	CH ₃	CH ₃		-(CH ₂) ₂ -
B-316	Br	CH ₃	CH ₃		-(CH ₂) ₂ -
B-317	CH ₃	CH ₃	CH ₃		-(CH ₂) ₂ -
B-318	CF ₃	CH ₃	CH ₃		-(CH ₂) ₂ -
B-319	OCH ₃	CH ₃	CH ₃		-(CH ₂) ₂ -
B-320	OCF ₃	CH ₃	CH ₃		-(CH ₂) ₂ -
B-321	H	H	H		-(CH ₂) ₃ -
B-322	F	H	H		-(CH ₂) ₃ -
B-323	Cl	H	H		-(CH ₂) ₃ -
B-324	Br	H	H		-(CH ₂) ₃ -
B-325	CH ₃	H	H		-(CH ₂) ₃ -
B-326	CF ₃	H	H		-(CH ₂) ₃ -
B-327	OCH ₃	H	H		-(CH ₂) ₃ -
B-328	OCF ₃	H	H		-(CH ₂) ₃ -
B-329	H	F	H		-(CH ₂) ₃ -
B-330	F	F	H		-(CH ₂) ₃ -
B-331	Cl	F	H		-(CH ₂) ₃ -
B-332	Br	F	H		-(CH ₂) ₃ -

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-333	CH ₃	F	H		-(CH ₂) ₃ -
B-334	CF ₃	F	H		-(CH ₂) ₃ -
B-335	OCH ₃	F	H		-(CH ₂) ₃ -
B-336	OCF ₃	F	H		-(CH ₂) ₃ -
B-337	H	Cl	H		-(CH ₂) ₃ -
B-338	F	Cl	H		-(CH ₂) ₃ -
B-339	Cl	Cl	H		-(CH ₂) ₃ -
B-340	Br	Cl	H		-(CH ₂) ₃ -
B-341	CH ₃	Cl	H		-(CH ₂) ₃ -
B-342	CF ₃	Cl	H		-(CH ₂) ₃ -
B-343	OCH ₃	Cl	H		-(CH ₂) ₃ -
B-344	OCF ₃	Cl	H		-(CH ₂) ₃ -
B-345	H	CH ₃	H		-(CH ₂) ₃ -
B-346	F	CH ₃	H		-(CH ₂) ₃ -
B-347	Cl	CH ₃	H		-(CH ₂) ₃ -
B-348	Br	CH ₃	H		-(CH ₂) ₃ -
B-349	CH ₃	CH ₃	H		-(CH ₂) ₃ -
B-350	CF ₃	CH ₃	H		-(CH ₂) ₃ -
B-351	OCH ₃	CH ₃	H		-(CH ₂) ₃ -
B-352	OCF ₃	CH ₃	H		-(CH ₂) ₃ -
B-353	H	CH ₃	CH ₃		-(CH ₂) ₃ -
B-354	F	CH ₃	CH ₃		-(CH ₂) ₃ -
B-355	Cl	CH ₃	CH ₃		-(CH ₂) ₃ -
B-356	Br	CH ₃	CH ₃		-(CH ₂) ₃ -
B-357	CH ₃	CH ₃	CH ₃		-(CH ₂) ₃ -
B-358	CF ₃	CH ₃	CH ₃		-(CH ₂) ₃ -
B-359	OCH ₃	CH ₃	CH ₃		-(CH ₂) ₃ -
B-360	OCF ₃	CH ₃	CH ₃		-(CH ₂) ₃ -
B-361	H	H	H		-(CH ₂) ₄ -
B-362	F	H	H		-(CH ₂) ₄ -

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-363	Cl	H	H		-(CH ₂) ₄ -
B-364	Br	H	H		-(CH ₂) ₄ -
B-365	CH ₃	H	H		-(CH ₂) ₄ -
B-366	CF ₃	H	H		-(CH ₂) ₄ -
B-367	OCH ₃	H	H		-(CH ₂) ₄ -
B-368	OCF ₃	H	H		-(CH ₂) ₄ -
B-369	H	F	H		-(CH ₂) ₄ -
B-370	F	F	H		-(CH ₂) ₄ -
B-371	Cl	F	H		-(CH ₂) ₄ -
B-372	Br	F	H		-(CH ₂) ₄ -
B-373	CH ₃	F	H		-(CH ₂) ₄ -
B-374	CF ₃	F	H		-(CH ₂) ₄ -
B-375	OCH ₃	F	H		-(CH ₂) ₄ -
B-376	OCF ₃	F	H		-(CH ₂) ₄ -
B-377	H	Cl	H		-(CH ₂) ₄ -
B-378	F	Cl	H		-(CH ₂) ₄ -
B-379	Cl	Cl	H		-(CH ₂) ₄ -
B-380	Br	Cl	H		-(CH ₂) ₄ -
B-381	CH ₃	Cl	H		-(CH ₂) ₄ -
B-382	CF ₃	Cl	H		-(CH ₂) ₄ -
B-383	OCH ₃	Cl	H		-(CH ₂) ₄ -
B-384	OCF ₃	Cl	H		-(CH ₂) ₄ -
B-385	H	CH ₃	H		-(CH ₂) ₄ -
B-386	F	CH ₃	H		-(CH ₂) ₄ -
B-387	Cl	CH ₃	H		-(CH ₂) ₄ -
B-388	Br	CH ₃	H		-(CH ₂) ₄ -
B-389	CH ₃	CH ₃	H		-(CH ₂) ₄ -
B-390	CF ₃	CH ₃	H		-(CH ₂) ₄ -
B-391	OCH ₃	CH ₃	H		-(CH ₂) ₄ -
B-392	OCF ₃	CH ₃	H		-(CH ₂) ₄ -

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-393	H	CH ₃	CH ₃		-(CH ₂) ₄ -
B-394	F	CH ₃	CH ₃		-(CH ₂) ₄ -
B-395	Cl	CH ₃	CH ₃		-(CH ₂) ₄ -
B-396	Br	CH ₃	CH ₃		-(CH ₂) ₄ -
B-397	CH ₃	CH ₃	CH ₃		-(CH ₂) ₄ -
B-398	CF ₃	CH ₃	CH ₃		-(CH ₂) ₄ -
B-399	OCH ₃	CH ₃	CH ₃		-(CH ₂) ₄ -
B-400	OCF ₃	CH ₃	CH ₃		-(CH ₂) ₄ -
B-401	H	H	H		-(CH ₂) ₅ -
B-402	F	H	H		-(CH ₂) ₅ -
B-403	Cl	H	H		-(CH ₂) ₅ -
B-404	Br	H	H		-(CH ₂) ₅ -
B-405	CH ₃	H	H		-(CH ₂) ₅ -
B-406	CF ₃	H	H		-(CH ₂) ₅ -
B-407	OCH ₃	H	H		-(CH ₂) ₅ -
B-408	OCF ₃	H	H		-(CH ₂) ₅ -
B-409	H	F	H		-(CH ₂) ₅ -
B-410	F	F	H		-(CH ₂) ₅ -
B-411	Cl	F	H		-(CH ₂) ₅ -
B-412	Br	F	H		-(CH ₂) ₅ -
B-413	CH ₃	F	H		-(CH ₂) ₅ -
B-414	CF ₃	F	H		-(CH ₂) ₅ -
B-415	OCH ₃	F	H		-(CH ₂) ₅ -
B-416	OCF ₃	F	H		-(CH ₂) ₅ -
B-417	H	Cl	H		-(CH ₂) ₅ -
B-418	F	Cl	H		-(CH ₂) ₅ -
B-419	Cl	Cl	H		-(CH ₂) ₅ -
B-420	Br	Cl	H		-(CH ₂) ₅ -
B-421	CH ₃	Cl	H		-(CH ₂) ₅ -
B-422	CF ₃	Cl	H		-(CH ₂) ₅ -

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-423	OCH ₃	Cl	H		-(CH ₂) ₅ -
B-424	OCF ₃	Cl	H		-(CH ₂) ₅ -
B-425	H	CH ₃	H		-(CH ₂) ₅ -
B-426	F	CH ₃	H		-(CH ₂) ₅ -
B-427	Cl	CH ₃	H		-(CH ₂) ₅ -
B-428	Br	CH ₃	H		-(CH ₂) ₅ -
B-429	CH ₃	CH ₃	H		-(CH ₂) ₅ -
B-430	CF ₃	CH ₃	H		-(CH ₂) ₅ -
B-431	OCH ₃	CH ₃	H		-(CH ₂) ₅ -
B-432	OCF ₃	CH ₃	H		-(CH ₂) ₅ -
B-433	H	CH ₃	CH ₃		-(CH ₂) ₅ -
B-434	F	CH ₃	CH ₃		-(CH ₂) ₅ -
B-435	Cl	CH ₃	CH ₃		-(CH ₂) ₅ -
B-436	Br	CH ₃	CH ₃		-(CH ₂) ₅ -
B-437	CH ₃	CH ₃	CH ₃		-(CH ₂) ₅ -
B-438	CF ₃	CH ₃	CH ₃		-(CH ₂) ₅ -
B-439	OCH ₃	CH ₃	CH ₃		-(CH ₂) ₅ -
B-440	OCF ₃	CH ₃	CH ₃		-(CH ₂) ₅ -
B-441	H	H	H		-(CH ₂) ₆ -
B-442	F	H	H		-(CH ₂) ₆ -
B-443	Cl	H	H		-(CH ₂) ₆ -
B-444	Br	H	H		-(CH ₂) ₆ -
B-445	CH ₃	H	H		-(CH ₂) ₆ -
B-446	CF ₃	H	H		-(CH ₂) ₆ -
B-447	OCH ₃	H	H		-(CH ₂) ₆ -
B-448	OCF ₃	H	H		-(CH ₂) ₆ -
B-449	H	F	H		-(CH ₂) ₆ -
B-450	F	F	H		-(CH ₂) ₆ -
B-451	Cl	F	H		-(CH ₂) ₆ -
B-452	Br	F	H		-(CH ₂) ₆ -

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-453	CH ₃	F	H		-(CH ₂) ₆ -
B-454	CF ₃	F	H		-(CH ₂) ₆ -
B-455	OCH ₃	F	H		-(CH ₂) ₆ -
B-456	OCF ₃	F	H		-(CH ₂) ₆ -
B-457	H	Cl	H		-(CH ₂) ₆ -
B-458	F	Cl	H		-(CH ₂) ₆ -
B-459	Cl	Cl	H		-(CH ₂) ₆ -
B-460	Br	Cl	H		-(CH ₂) ₆ -
B-461	CH ₃	Cl	H		-(CH ₂) ₆ -
B-462	CF ₃	Cl	H		-(CH ₂) ₆ -
B-463	OCH ₃	Cl	H		-(CH ₂) ₆ -
B-464	OCF ₃	Cl	H		-(CH ₂) ₆ -
B-465	H	CH ₃	H		-(CH ₂) ₆ -
B-466	F	CH ₃	H		-(CH ₂) ₆ -
B-467	Cl	CH ₃	H		-(CH ₂) ₆ -
B-468	Br	CH ₃	H		-(CH ₂) ₆ -
B-469	CH ₃	CH ₃	H		-(CH ₂) ₆ -
B-470	CF ₃	CH ₃	H		-(CH ₂) ₆ -
B-471	OCH ₃	CH ₃	H		-(CH ₂) ₆ -
B-472	OCF ₃	CH ₃	H		-(CH ₂) ₆ -
B-473	H	CH ₃	CH ₃		-(CH ₂) ₆ -
B-474	F	CH ₃	CH ₃		-(CH ₂) ₆ -
B-475	Cl	CH ₃	CH ₃		-(CH ₂) ₆ -
B-476	Br	CH ₃	CH ₃		-(CH ₂) ₆ -
B-477	CH ₃	CH ₃	CH ₃		-(CH ₂) ₆ -
B-478	CF ₃	CH ₃	CH ₃		-(CH ₂) ₆ -
B-479	OCH ₃	CH ₃	CH ₃		-(CH ₂) ₆ -
B-480	OCF ₃	CH ₃	CH ₃		-(CH ₂) ₆ -
B-481	H	H	H		-CH=CH-CH ₂ -CH ₂ -
B-482	F	H	H		-CH=CH-CH ₂ -CH ₂ -

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-483	Cl	H	H		-CH=CH-CH ₂ -CH ₂ -
B-484	Br	H	H		-CH=CH-CH ₂ -CH ₂ -
B-485	CH ₃	H	H		-CH=CH-CH ₂ -CH ₂ -
B-486	CF ₃	H	H		-CH=CH-CH ₂ -CH ₂ -
B-487	OCH ₃	H	H		-CH=CH-CH ₂ -CH ₂ -
B-488	OCF ₃	H	H		-CH=CH-CH ₂ -CH ₂ -
B-489	H	F	H		-CH=CH-CH ₂ -CH ₂ -
B-490	F	F	H		-CH=CH-CH ₂ -CH ₂ -
B-491	Cl	F	H		-CH=CH-CH ₂ -CH ₂ -
B-492	Br	F	H		-CH=CH-CH ₂ -CH ₂ -
B-493	CH ₃	F	H		-CH=CH-CH ₂ -CH ₂ -
B-494	CF ₃	F	H		-CH=CH-CH ₂ -CH ₂ -
B-495	OCH ₃	F	H		-CH=CH-CH ₂ -CH ₂ -
B-496	OCF ₃	F	H		-CH=CH-CH ₂ -CH ₂ -
B-497	H	Cl	H		-CH=CH-CH ₂ -CH ₂ -
B-498	F	Cl	H		-CH=CH-CH ₂ -CH ₂ -
B-499	Cl	Cl	H		-CH=CH-CH ₂ -CH ₂ -
B-500	Br	Cl	H		-CH=CH-CH ₂ -CH ₂ -
B-501	CH ₃	Cl	H		-CH=CH-CH ₂ -CH ₂ -
B-502	CF ₃	Cl	H		-CH=CH-CH ₂ -CH ₂ -
B-503	OCH ₃	Cl	H		-CH=CH-CH ₂ -CH ₂ -
B-504	OCF ₃	Cl	H		-CH=CH-CH ₂ -CH ₂ -
B-505	H	CH ₃	H		-CH=CH-CH ₂ -CH ₂ -
B-506	F	CH ₃	H		-CH=CH-CH ₂ -CH ₂ -
B-507	Cl	CH ₃	H		-CH=CH-CH ₂ -CH ₂ -
B-508	Br	CH ₃	H		-CH=CH-CH ₂ -CH ₂ -
B-509	CH ₃	CH ₃	H		-CH=CH-CH ₂ -CH ₂ -
B-510	CF ₃	CH ₃	H		-CH=CH-CH ₂ -CH ₂ -
B-511	OCH ₃	CH ₃	H		-CH=CH-CH ₂ -CH ₂ -
B-512	OCF ₃	CH ₃	H		-CH=CH-CH ₂ -CH ₂ -

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-513	H	CH ₃	CH ₃	-CH=CH-CH ₂ -CH ₂ -	
B-514	F	CH ₃	CH ₃	-CH=CH-CH ₂ -CH ₂ -	
B-515	Cl	CH ₃	CH ₃	-CH=CH-CH ₂ -CH ₂ -	
B-516	Br	CH ₃	CH ₃	-CH=CH-CH ₂ -CH ₂ -	
B-517	CH ₃	CH ₃	CH ₃	-CH=CH-CH ₂ -CH ₂ -	
B-518	CF ₃	CH ₃	CH ₃	-CH=CH-CH ₂ -CH ₂ -	
B-519	OCH ₃	CH ₃	CH ₃	-CH=CH-CH ₂ -CH ₂ -	
B-520	OCF ₃	CH ₃	CH ₃	-CH=CH-CH ₂ -CH ₂ -	
B-521	H	H	-(CH ₂) ₃ -		CH ₃
B-522	F	H	-(CH ₂) ₃ -		CH ₃
B-523	Cl	H	-(CH ₂) ₃ -		CH ₃
B-524	Br	H	-(CH ₂) ₃ -		CH ₃
B-525	CH ₃	H	-(CH ₂) ₃ -		CH ₃
B-526	CF ₃	H	-(CH ₂) ₃ -		CH ₃
B-527	OCH ₃	H	-(CH ₂) ₃ -		CH ₃
B-528	OCF ₃	H	-(CH ₂) ₃ -		CH ₃
B-529	H	D	-(CH ₂) ₃ -		CH ₃
B-530	F	D	-(CH ₂) ₃ -		CH ₃
B-531	Cl	D	-(CH ₂) ₃ -		CH ₃
B-532	Br	D	-(CH ₂) ₃ -		CH ₃
B-533	CH ₃	D	-(CH ₂) ₃ -		CH ₃
B-534	CF ₃	D	-(CH ₂) ₃ -		CH ₃
B-535	OCH ₃	D	-(CH ₂) ₃ -		CH ₃
B-536	OCF ₃	D	-(CH ₂) ₃ -		CH ₃
B-537	H	H	-(CH ₂) ₃ -		D
B-538	F	H	-(CH ₂) ₃ -		D
B-539	Cl	H	-(CH ₂) ₃ -		D
B-540	Br	H	-(CH ₂) ₃ -		D
B-541	CH ₃	H	-(CH ₂) ₃ -		D
B-542	CF ₃	H	-(CH ₂) ₃ -		D

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-543	OCH ₃	H	-(CH ₂) ₃ -		D
B-544	OCF ₃	H	-(CH ₂) ₃ -		D
B-545	H	D	-(CH ₂) ₃ -		D
B-546	F	D	-(CH ₂) ₃ -		D
B-547	Cl	D	-(CH ₂) ₃ -		D
B-548	Br	D	-(CH ₂) ₃ -		D
B-549	CH ₃	D	-(CH ₂) ₃ -		D
B-550	CF ₃	D	-(CH ₂) ₃ -		D
B-551	OCH ₃	D	-(CH ₂) ₃ -		D
B-552	OCF ₃	D	-(CH ₂) ₃ -		D
B-553	H	H	-(CH ₂) ₄ -		CH ₃
B-554	F	H	-(CH ₂) ₄ -		CH ₃
B-555	Cl	H	-(CH ₂) ₄ -		CH ₃
B-556	Br	H	-(CH ₂) ₄ -		CH ₃
B-557	CH ₃	H	-(CH ₂) ₄ -		CH ₃
B-558	CF ₃	H	-(CH ₂) ₄ -		CH ₃
B-559	OCH ₃	H	-(CH ₂) ₄ -		CH ₃
B-560	OCF ₃	H	-(CH ₂) ₄ -		CH ₃
B-561	H	D	-(CH ₂) ₄ -		CH ₃
B-562	F	D	-(CH ₂) ₄ -		CH ₃
B-563	Cl	D	-(CH ₂) ₄ -		CH ₃
B-564	Br	D	-(CH ₂) ₄ -		CH ₃
B-565	CH ₃	D	-(CH ₂) ₄ -		CH ₃
B-566	CF ₃	D	-(CH ₂) ₄ -		CH ₃
B-567	OCH ₃	D	-(CH ₂) ₄ -		CH ₃
B-568	OCF ₃	D	-(CH ₂) ₄ -		CH ₃
B-569	H	H	-(CH ₂) ₄ -		D
B-570	F	H	-(CH ₂) ₄ -		D
B-571	Cl	H	-(CH ₂) ₄ -		D
B-572	Br	H	-(CH ₂) ₄ -		D

No.	R ^{9a}	R ^{5b}	R ^{5a}	R ⁷	R ⁸
B-573	CH ₃	H		-(CH ₂) ₄ -	D
B-574	CF ₃	H		-(CH ₂) ₄ -	D
B-575	OCH ₃	H		-(CH ₂) ₄ -	D
B-576	OCF ₃	H		-(CH ₂) ₄ -	D
B-577	H	D		-(CH ₂) ₄ -	D
B-578	F	D		-(CH ₂) ₄ -	D
B-579	Cl	D		-(CH ₂) ₄ -	D
B-580	Br	D		-(CH ₂) ₄ -	D
B-581	CH ₃	D		-(CH ₂) ₄ -	D
B-582	CF ₃	D		-(CH ₂) ₄ -	D
B-583	OCH ₃	D		-(CH ₂) ₄ -	D
B-584	OCF ₃	D		-(CH ₂) ₄ -	D

* C₆H₅ = phenyl


In a specific embodiment, the invention relates to compounds I selected from the compounds of the examples, either in form of free bases or of any pharmaceutically acceptable salt thereof or a stereoisomer thereof.

The compounds of the present invention can be prepared by using routine techniques familiar to a skilled person. In particular, the compounds of the formula I can be prepared according to the following schemes, wherein the variables, if not stated otherwise, are as defined above.

Compounds of formula I wherein X is CR⁷R⁸ and R^{5b} is H (= compounds I') can be synthesized as described in scheme 1 below. The protected tetrahydrobenzodiazepine or tetrahydroquinoxaline **1**, wherein Z is a hydrogen or a halogen atom, such as Cl, Br or I and PG is a common protective group, such as a carbamate, especially boc, is acylated with the acrylic acid derivative **2**, wherein LG is an appropriate leaving group, such as Cl or an anhydride or a chloroformate, in the presence of a base, such as triethylamine or Hünig's base, in an organic solvent, such as ether or methylene chloride. Reaction of **3** with a Lewis acid or a Brönstedt acid HA or irradiation with a suitable wave-

length commonly derived from a mercury lamp in an adequate solvent, such as acetone or toluene, in a common photoreactor yields cyclization to **4**. Reduction of the carbonyl group with common reduction agents like borohydrides such as sodium borohydride or borane-tetrahydrofuran-complex yields **5**, which is deprotected using suitable reagents such as strong bases or acids to I', wherein R^{4a} and R^{4b} and R^{5b} are H. Compounds I wherein R^{4a} and R^{4b} form together =O can be obtained by skipping the reduction step to **5** and deprotecting **4**.

Scheme 1




If desired, substituents R^1 different from hydrogen can be introduced for example via alkylation under typical conditions such as stirring in an appropriate solvent in the presence of an alkylhalide and a base or via other common substitution reactions.

Compounds I wherein R^{5b} is different from H can be prepared, for example, by reacting compound **4** with a compound $LG-R^{5b}$ in the presence of a base, wherein LG is an appropriate leaving group, such as Cl or Br.

Compounds I wherein R^{5b} is different from H and R^8 is CH_3 (= compounds I'') can be prepared, for example, as outlined in scheme 2 below. Acylation of **1** with the allylic acid derivative **6**, wherein LG is an appropriate leaving group, such as Cl or an anhydride or a chloroformate, in the presence of a base, such as triethylamine or Hünig's base, in an organic solvent, such as ether or methylene chloride, yields **7**, which is

reacted in a Heck-type reaction, e.g. employing palladium acetate in the presence of a base, such as potassium carbonate, to yield the cyclization product **8**. Reduction of the carbonyl group with common reduction agents like borohydrides such as sodium borohydride or borane-tetrahydrofuran-complex yields **9**, which is deprotected using suitable reagents such as strong bases or acids to I'', wherein R^{4a} and R^{4b} are H. Compounds I wherein R^{4a} and R^{4b} form together =O can be obtained by skipping the reduction step to **9** and deprotecting **8**. If desired, substituents R¹ different from hydrogen can be introduced for example via alkylation under typical conditions such as stirring in an appropriate solvent in the presence of an alkylhalide and a base or via other common substitution reactions.


Scheme 2



Compounds I wherein X is NR⁶ (= compounds I''') can be prepared as outlined in scheme 3 below. Acylation of **1** with the β -amino acid derivative **10**, wherein PG' is a protection group different from PG and LG is an appropriate leaving group, such as Cl or an anhydride or a chloroformate, in the presence of a base, such as triethylamine or Hünig's base, in an organic solvent, such as ether or methylene chloride, yields **11**. This is first selectively deprotected at the amino group NR⁶PG'. The partially deprotected compound is then reacted in a cyclization reaction to **12** under Buchwald-Hartwig reaction conditions such as the use of a Pd catalyst, e.g. tetrakis(triphenylphosphine) palladium in the presence of as base, such as sodium tert-butyrate and potassium carbonate. Reduction of the carbonyl group of **12** with common reduction agents like borohydrides

such as sodium borohydride or borane-tetrahydrofuran-complex yields **13**, which is deprotected using suitable reagents such as strong bases or acids to I''', wherein R^{4a} and R^{4b} are H. Compounds I wherein R^{4a} and R^{4b} form together =O can be obtained by skipping the reduction step to **13** and deprotecting **12**. If desired, substituents R¹ different from hydrogen can be introduced for example via alkylation under typical conditions such as stirring in an appropriate solvent in the presence of an alkylhalide and a base or via other common substitution reactions.


Scheme 3



Alternatively to the method depicted in scheme 1, compounds I wherein X is CR⁷R⁸ and R^{5b} is H (= compounds I') can be synthesized as described in scheme 4 below. Readily available anilines **14** are derivatized with carbonyl moieties **15** by acylation procedures employing appropriate leaving groups LG, such as chlorides or anhydrides, in the presence of a base such as triethylamine or Hünig's base, in an organic solvent, such as diethyl ether or methylene chloride to yield **16**. Cyclization products **17** are received by irradiation with a suitable wavelength commonly derived from a mercury lamp in an adequate solvent such as acetone or toluene in a common photoreactor known to those skilled in the art. Reduction with common reduction agents like borohydrides such as sodium borohydride or borane-tetrahydrofuran-complex yield suitable intermediates **18** that are alkylated with alkylamides **19** employing suitable leaving groups LG such as chlorides or bromides. After reduction to the corresponding amines

20 cyclization is performed in a suitable solvent like acetonitrile or methanol at room temperature or higher temperature by addition of an acid and substituted ketones or aldehydes to obtain compounds **I'**.


Scheme 4



Alternatively to the method depicted in scheme 3, compounds **I** wherein X is NR^6 (= compounds **I'''**) can be prepared as outlined in scheme 5 below.

Readily available starting materials **22** are alkylated with alkylamides **19** employing suitable leaving groups LG such as chlorides or bromides to give **23**. After reduction to the corresponding amines **24** cyclization is performed in a suitable solvent like acetonitrile or methanol at room temperature or higher temperature by addition of an acid and substituted ketones or aldehydes **21** to obtain compounds **I'''**.

Scheme 5



Compounds I wherein R^{4a} and R^{4b} are not H or do not form together a group =O can be prepared by standard derivatization methods of compounds wherein R^{4a} and R^{4b} form together a group =O. For instance, compounds wherein R^{4a} and R^{4b} form together a group =S may be prepared by reaction with a sulfurization agent, such as Lawesson's reagent or P_2S_5 . Alkyl and related groups as radicals R^{4a} and R^{4b} may be introduced via Grignard reduction. Amino and related groups may be introduced via reductive amination. Hydroxyl group R^{4a} or R^{4b} may be introduced by reducing the carbonyl group. This may be alkylated to yield alkoxy and related groups R^{4a} and R^{4b} or substituted by diverse groups.

If not otherwise indicated, the above-described reactions are generally carried out in a solvent at temperatures between room temperature and the boiling temperature of the solvent employed. Alternatively, the activation energy which is required for the reaction can be introduced into the reaction mixture using microwaves, something which has proved to be of value, in particular, in the case of the reactions catalyzed by transition metals (with regard to reactions using microwaves, see Tetrahedron 2001, 57, p. 9199 ff. p. 9225 ff. and also, in a general manner, "Microwaves in Organic Synthesis", André Loupy (Ed.), Wiley-VCH 2002).

The acid addition salts of compounds I are prepared in a customary manner by mixing the free base with a corresponding acid, where appropriate in solution in an organic solvent, for example a lower alcohol, such as methanol, ethanol or propanol, an ether, such as methyl tert-butyl ether or diisopropyl ether, a ketone, such as acetone or methyl ethyl ketone, or an ester, such as ethyl acetate.

Routine experimentations, including appropriate manipulation of the reaction conditions, reagents and sequence of the synthetic route, protection of any chemical functionality that may not be compatible with the reaction conditions, and deprotection at a suitable point in the reaction sequence of the preparation methods are within routine techniques.

Suitable protecting groups and the methods for protecting and deprotecting different substituents using such suitable protecting groups are well known to those skilled in the art; examples of which may be found in T. Greene and P. Wuts, *Protective Groups in Organic Synthesis* (3rd ed.), John Wiley & Sons, NY (1999), which is herein incorporated by reference in its entirety. Synthesis of the compounds of the invention may be accomplished by methods analogous to those described in the synthetic schemes described hereinabove and in specific examples.

Starting materials, if not commercially available, may be prepared by procedures selected from standard organic chemical techniques, techniques that are analogous to the synthesis of known, structurally similar compounds, or techniques that are analogous to the above described schemes or the procedures described in the synthetic examples section.

When an optically active form of a compound of the invention is required, it may be obtained by carrying out one of the procedures described herein using an optically active starting material (prepared, for example, by asymmetric induction of a suitable reaction step), or by resolution of a mixture of the stereoisomers of the compound or intermediates using a standard procedure (such as chromatographic separation, recrystallization or enzymatic resolution).

Similarly, when a pure geometric isomer of a compound of the invention is required, it may be obtained by carrying out one of the above procedures using a pure geometric isomer as a starting material, or by resolution of a mixture of the geometric

isomers of the compound or intermediates using a standard procedure such as chromatographic separation.

Moreover, the present invention relates to compounds of formula I as defined above, wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope (e.g., hydrogen by deuterium, ^{12}C by ^{13}C , ^{14}N by ^{15}N , ^{16}O by ^{18}O) and preferably wherein at least one hydrogen atom has been replaced by a deuterium atom.

Additionally, the compounds according to the invention include more of the respective isotope than occurs naturally and which is present in compounds I.

Stable isotopes (e.g., deuterium, ^{13}C , ^{15}N , ^{18}O) are nonradioactive isotopes which contain one additional neutron than the normally abundant isotope of the respective atom. Deuterated compounds have been used in pharmaceutical research to investigate the in vivo metabolic fate of the compounds by evaluation of the mechanism of action and metabolic pathway of the non deuterated parent compound (Blake et al. *J. Pharm. Sci.* 64, 3, 367-391 (1975)). Such metabolic studies are important in the design of safe, effective therapeutic drugs, either because the in vivo active compound administered to the patient or because the metabolites produced from the parent compound prove to be toxic or carcinogenic (Foster et al., *Advances in Drug Research* Vol. 14, pp. 2-36, Academic press, London, 1985; Kato et al., *J. Labelled Comp. Radiopharmaceut.*, 36(10):927-932 (1995); Kushner et al., *Can. J. Physiol. Pharmacol.*, 77, 79-88 (1999)).

Incorporation of a heavy atom, particularly substitution of deuterium for hydrogen, can give rise to an isotope effect that could alter the pharmacokinetics of the drug.

Stable isotope labeling of a drug can alter its physico-chemical properties such as pK_a and lipid solubility. These changes may influence the fate of the drug at different steps along its passage through the body. Absorption, distribution, metabolism or excretion can be changed. Absorption and distribution are processes that depend primarily on the molecular size and the lipophilicity of the substance. These effects and alterations can affect the pharmacodynamic response of the drug molecule if the isotopic substitution affects a region involved in a ligand-receptor interaction.

Drug metabolism can give rise to large isotopic effect if the breaking of a chemical bond to a deuterium atom is the rate limiting step in the process. While some of the physical properties of a stable isotope-labeled molecule are different from those of the unlabeled one, the chemical and biological properties are the same, with one important

exception: because of the increased mass of the heavy isotope, any bond involving the heavy isotope and another atom will be stronger than the same bond between the light isotope and that atom. In any reaction in which the breaking of this bond is the rate limiting step, the reaction will proceed slower for the molecule with the heavy isotope due to "kinetic isotope effect". A reaction involving breaking a C--D bond can be up to 700 percent slower than a similar reaction involving breaking a C--H bond. If the C--D bond is not involved in any of the steps leading to the metabolite, there may not be any effect to alter the behavior of the drug. If a deuterium is placed at a site involved in the metabolism of a drug, an isotope effect will be observed only if breaking of the C--D bond is the rate limiting step. There is evidence to suggest that whenever cleavage of an aliphatic C--H bond occurs, usually by oxidation catalyzed by a mixed-function oxidase, replacement of the hydrogen by deuterium will lead to observable isotope effect. It is also important to understand that the incorporation of deuterium at the site of metabolism slows its rate to the point where another metabolite produced by attack at a carbon atom not substituted by deuterium becomes the major pathway a process called "metabolic switching".

Deuterium tracers, such as deuterium-labeled drugs and doses, in some cases repeatedly, of thousands of milligrams of deuterated water, are also used in healthy humans of all ages, including neonates and pregnant women, without reported incident (e.g. Pons G and Rey E, *Pediatrics* 1999 104: 633; Coward W A et al., *Lancet* 1979 7: 13; Schwarcz H P, *Control. Clin. Trials* 1984 5(4 Suppl): 573; Rodewald L E et al., *J. Pediatr.* 1989 114: 885; Butte N F et al. *Br. J. Nutr.* 1991 65: 3; MacLennan A H et al. *Am. J. Obstet Gynecol.* 1981 139: 948). Thus, it is clear that any deuterium released, for instance, during the metabolism of compounds of this invention poses no health risk.

The weight percentage of hydrogen in a mammal (approximately 9%) and natural abundance of deuterium (approximately 0.015%) indicates that a 70 kg human normally contains nearly a gram of deuterium. Furthermore, replacement of up to about 15% of normal hydrogen with deuterium has been effected and maintained for a period of days to weeks in mammals, including rodents and dogs, with minimal observed adverse effects (Czajka D M and Finkel A J, *Ann. N.Y. Acad. Sci.* 1960 84: 770; Thomson J F, *Ann. New York Acad. Sci* 1960 84: 736; Czajka D M et al., *Am. J. Physiol.* 1961 201: 357). Higher deuterium concentrations, usually in excess of 20%, can be toxic in ani-

mals. However, acute replacement of as high as 15%-23% of the hydrogen in humans' fluids with deuterium was found not to cause toxicity (Blagojevic N et al. in "Dosimetry & Treatment Planning for Neutron Capture Therapy", Zamenhof R, Solares G and Harling O Eds. 1994. Advanced Medical Publishing, Madison Wis. pp.125-134; Diabetes Metab. 23: 251 (1997)).

Increasing the amount of deuterium present in a compound above its natural abundance is called enrichment or deuterium-enrichment. Examples of the amount of enrichment include from about 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 16, 21, 25, 29, 33, 37, 42, 46, 50, 54, 58, 63, 67, 71, 75, 79, 84, 88, 92, 96, to about 100 mol %.

The hydrogens present on a particular organic compound have different capacities for exchange with deuterium. Certain hydrogen atoms are easily exchangeable under physiological conditions and, if replaced by deuterium atoms, it is expected that they will readily exchange for protons after administration to a patient. Certain hydrogen atoms may be exchanged for deuterium atoms by the action of a deuterio acid such as D_2SO_4/D_2O . Alternatively, deuterium atoms may be incorporated in various combinations during the synthesis of compounds of the invention. Certain hydrogen atoms are not easily exchangeable for deuterium atoms. However, deuterium atoms at the remaining positions may be incorporated by the use of deuterated starting materials or intermediates during the construction of compounds of the invention.

Deuterated and deuterium-enriched compounds of the invention can be prepared by using known methods described in the literature. Such methods can be carried out utilizing corresponding deuterated and optionally, other isotope-containing reagents and/or intermediates to synthesize the compounds delineated herein, or invoking standard synthetic protocols known in the art for introducing isotopic atoms to a chemical structure. Relevant procedures and intermediates are disclosed, for instance in Lizondo, J et al., *Drugs Fut*, 21(11), 1116 (1996); Brickner, S J et al., *J Med Chem*, 39(3), 673 (1996); Mallesham, B et al., *Org Lett*, 5(7), 963 (2003); PCT publications WO1997010223, WO2005099353, WO1995007271, WO2006008754; US Patent Nos. 7538189; 7534814; 7531685; 7528131; 7521421; 7514068; 7511013; and US Patent Application Publication Nos. 20090137457; 20090131485; 20090131363; 20090118238; 20090111840; 20090105338; 20090105307; 20090105147;

20090093422; 20090088416; 20090082471, the methods are hereby incorporated by reference.

The present invention further relates to a pharmaceutical composition comprising a therapeutically effective amount of at least one compound I as defined above or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof, or comprising at least one compound as defined in any of the preceding claims wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope, preferably wherein at least one hydrogen atom has been replaced by a deuterium atom, in combination with at least one pharmaceutically acceptable carrier and/or auxiliary substance.

The present invention further relates to a compound I as defined above or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for use as a medicament.

The present invention also relates to a compound I as defined above or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for the treatment of disorders which respond to the modulation of the 5-HT_{2c} receptor.

The present invention also relates to the use of a compound I as defined above or of an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of disorders which respond to the modulation of the 5-HT_{2c} receptor, and to a method for treating disorders which respond to the modulation of the 5-HT_{2c} receptor, which method comprises administering to a subject in need thereof at least one compound I as defined above or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof.

The compounds of the present invention are modulators of the 5-HT_{2c} receptor. Specifically, the compounds of formula I are agonists or partial agonists of the 5-HT_{2c} receptor. Thus, in a specific embodiment, the invention relates to a compound I as defined above or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for the treatment of disorders which respond to 5-HT_{2c} receptor agonists, further to the use of a compound I as defined above or of an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for the manu-

facture of a medicament for the treatment of disorders which respond to 5-HT_{2c} receptor agonists, and to a method for treating disorders which respond to 5-HT_{2c} receptor agonists, which method comprises administering to a subject in need thereof at least one compound I as defined above or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof.

Within the meaning of the invention, the term "disorder" denotes disturbances and/or anomalies which are as a rule regarded as being pathological conditions or functions and which can manifest themselves in the form of particular signs, symptoms and/or malfunctions. While the treatment according to the invention can be directed toward individual disorders, i.e. anomalies or pathological conditions, it is also possible for several anomalies, which may be causatively linked to each other, to be combined into patterns, i.e. syndromes, which can be treated in accordance with the invention.

In one aspect of the invention, the diseases to be treated are disorders are damage of the central nervous system, disorders of the central nervous system, eating disorders, ocular hypertension, cardiovascular disorders, gastrointestinal disorders and diabetes.

Disorders or diseases of the central nervous system are understood as meaning disorders which affect the spinal cord and, in particular, the brain. These are, for example, cognitive dysfunction, attention deficit disorder/hyperactivity syndrome and cognitive deficits related with schizophrenia, attention deficit/hyperactivity syndrome, personality disorders, affective disorders, motion or motor disorders, pain, migraine, sleep disorders (including disturbances of the Circadian rhythm), feeding disorders, diseases associated with neurodegeneration, addiction diseases, obesity or psoriasis.

Examples of cognitive dysfunction are deficits in memory, cognition, and learning, Alzheimer's disease, age-related cognitive decline, and mild cognitive impairment, or any combinations thereof. Examples of personality disorders are schizophrenia and cognitive deficits related to schizophrenia. Examples of affective disorders are depression, anxiety, bipolar disorder and obsessive compulsive disorders, or any combination thereof. Examples of motion or motor disorders are Parkinson's disease and epilepsy. Examples of feeding disorders are obesity, bulimia, weight loss and anorexia, especially anorexia nervosa. Examples of diseases associated with neurodegeneration are stroke, spinal or head trauma, and head injuries, such as hydrocephalus.

Pain condition includes nociceptive pain, neuropathic pain or a combination thereof. Such pain conditions or disorders can include, but are not limited to, post-operative pain, osteoarthritis pain, pain due to inflammation, rheumatoid arthritis pain, musculoskeletal pain, burn pain (including sunburn), ocular pain, the pain associated with dental conditions (such as dental caries and gingivitis), post-partum pain, bone fracture, herpes, HIV, traumatic nerve injury, stroke, post-ischemia, fibromyalgia, reflex sympathetic dystrophy, complex regional pain syndrome, spinal cord injury, sciatica, phantom limb pain, diabetic neuropathy, hyperalgesia and cancer.

In certain other embodiments, the disease condition is bladder dysfunction, including urinary incontinence.

Diabetes includes diabetes insipidus, diabetes mellitus, type I diabetes, type II diabetes, type III diabetes, diabetes secondary to pancreatic diseases, diabetes related to steroid use, diabetes complications, hyperglycemia and insulin resistance.

The addiction diseases include psychiatric disorders and behavioral disturbances which are caused by the abuse of psychotropic substances, such as pharmaceuticals or narcotics, and also other addiction diseases, such as addiction to gaming (impulse control disorders not elsewhere classified). Examples of addictive substances are: opioids (e.g. morphine, heroin and codeine), cocaine; nicotine; alcohol; substances which interact with the GABA chloride channel complex, sedatives, hypnotics and tranquilizers, for example benzodiazepines; LSD; cannabinoids; psychomotor stimulants, such as 3,4-methylenedioxy-*N*-methamphetamine (ecstasy); amphetamine and amphetamine-like substances such as methylphenidate, other stimulants including caffeine and nicotine. Addictive substances which come particularly into consideration are opioids, cocaine, amphetamine or amphetamine-like substances, nicotine and alcohol. Especially, addiction disorders include alcohol abuse, cocaine abuse, tobacco abuse and smoking cessation.

With regard to the treatment of addiction diseases, particular preference is given to those compounds according to the invention of the formula (I) which themselves do not possess any psychotropic effect. This can also be observed in a test using rats, which, after having been administered compounds which can be used in accordance with the invention, reduce their self administration of psychotropic substances, for example cocaine.

Examples of gastrointestinal disorders are irritable bowel syndrome.

Preferably, the disorders are selected from the group consisting of bipolar disorder, depression, atypical depression, mood episodes, adjustment disorders, anxiety, panic disorders, post-traumatic syndrome, psychoses, schizophrenia, cognitive deficits of schizophrenia, memory loss, dementia of aging, Alzheimer's disease, behavioral disorders associated with dementia, social phobia, mental disorders in childhood, attention deficit hyperactivity disorder, organic mental disorders, autism, mutism, disruptive behavior disorder, impulse control disorder, borderline personality disorder, obsessive compulsive disorder, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, seizure disorders, epilepsy, substance use disorders, alcohol abuse, cocaine abuse, tobacco abuse, smoking cessation, sexual dysfunction/erectile dysfunction in males, sexual dysfunction in females, premenstrual syndrome, late luteal phase syndrome, chronic fatigue syndrome, sleep disorders, sleep apnoea, chronic fatigue syndrome, psoriasis, Parkinson's disease, spinal cord injury, trauma, stroke, pain, bladder dysfunction/urinary incontinence, encephalitis, meningitis, eating disorders, obesity, bulimia, weight loss, anorexia nervosa, ocular hypertension, cardiovascular disorders, gastrointestinal disorders, diabetes insipidus, diabetes mellitus, type I diabetes, type II diabetes, type III diabetes, diabetes secondary to pancreatic diseases, diabetes related to steroid use, diabetes complications, hyperglycemia and insulin resistance, and are specifically schizophrenia, depression, bipolar disorders, obesity or substance use disorders.

The compounds of the invention may be used for a preventive treatment (prophylaxis), in particular as relapse prophylaxis or phase prophylaxis, but are preferably used for a treatment in its proper sense, i.e. for the treatment of acute or chronic signs, symptoms and/or malfunctions. The treatment can be orientated symptomatically, for example as the suppression of symptoms. It can be effected over a short period, be orientated over the medium term or can be a long-term treatment, for example within the context of a maintenance therapy.

In another embodiment, the present invention relates to the use of a compound I as defined above or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for preparing a medicament for preventing (the development of) a disease condition as described above and to a method for preventing (the development

of) a disease condition as described above comprises administering to the subject in need of treatment thereof (e.g., a mammal, such as a human) a therapeutically effective amount of a compound I as defined above or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof. As used herein, the term “prevent” a disease condition by administration of any of the compounds described herein means that the detectable physical characteristics or symptoms of the disease or condition do not develop following the administration of the compound described herein. Alternatively, the method comprises administering to the subject a therapeutically effective amount of a compound I as defined above or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof, in combination with a therapeutically effective amount of at least one cognitive enhancing drug.

In yet another embodiment, the present invention relates to the use a compound I as defined above or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for preparing a medicament for preventing the progression (e.g., worsening) of a disease condition and to a method for preventing the progression (e.g., worsening) of a disease condition, which method comprises administering to the subject in need of treatment thereof (e.g., a mammal, such as a human) a therapeutically effective amount of a compound I as defined above or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof.

There are several lines of evidence suggesting that 5-HT_{2C} agonists or partial agonists would have therapeutic use in a variety of diseases, disorders and conditions.

Knockout mice models lacking the 5-HT_{2C} receptor exhibit hyperphagia, obesity and are more prone to seizures and sudden death [Tecott LH, Sun LM, Akana SF, Strack AM, Lowenstein DH, Dallman MF, Julius D (1995) Eating disorder and epilepsy in mice lacking 5-HT_{2C} serotonin receptors. *Nature* 374:542-546]. They also exhibit compulsive-like behavior [Chou-Green JM, Holscher TD, Dallman MF, Akana SF (2003). Compulsive behavior in the 5-HT_{2C} receptor knockout mouse. *Phys. Behav.* 78:641-649], hyperresponsiveness to repeated stress [Chou-Green JM, Holscher TD, Dallman MF, Akana SF (2003). Repeated stress in young and old 5-HT_{2C} receptor knockout mouse. *Phys. Behav.* 79:217-226], wakefulness [Frank MG, Stryker MP, Tecott LH (2002). Sleep and sleep homeostasis in mice lacking the 5-HT_{2C} receptor. *Neuropsychopharmacology* 27:869-873], hyperactivity and drug dependence [Rocha

BA, Goulding EH, O'Dell LE, Mead AN, Coufal NG, Parsons LH, Tecott LH (2002). Enhanced locomotor, reinforcing and neurochemical effects of cocaine in serotonin 5-hydroxytryptamine 2C receptor mutant mice. *J. Neurosci.* 22:10039-10045].

5-HT_{2C} is unique among other G-protein-coupled receptors (GPCRs) in that its pre-mRNA is a substrate for base modification via hydrolytic deamination of adenosines to yield inosines. Five adenosines, located within a sequence encoding the putative second intracellular domain can be converted to inosines. This editing can alter the coding potential of the triplet codons and allows for the generation of multiple different receptor isoforms. The edited receptor isoforms were shown to have reduced ability to interact with G-proteins in the absence of agonist stimulation [Werry, TD, Loiacono R, Sexton PA, Christopoulos A (2008). RNA editing of the serotonin 5-HT_{2C} receptor and its effects on cell signaling, pharmacology and brain function. *Pharmac. Therap.* 119:7-23].

Edited 5-HT_{2C} isoforms with reduced function are significantly expressed in the brains of depressed suicide victims [Schmauss C (2003) Serotonin 2C receptors: suicide, serotonin, and runaway RNA editing. *Neuroscientist* 9:237-242. Iwamoto K, Kato T (2003). RNA editing of serotonin 2C receptor in human postmortem brains of major mental disorders. *Neurosci. Lett.* 346:169-172] and in the learned helplessness rats (a well established animal model of depression) [Iwamoto K, Nakatanib N, Bundoa M, Yoshikawab T, Katoa T (2005). Altered RNA editing of serotonin 2C receptor in a rat model of depression. *Neurosci. Res.* 53: 69-76] suggesting a link between 5-HT_{2C} function and depression. There are also implications of edited 5-HT_{2C} isoforms and spatial memory [Du Y, Stasko M, Costa AC, Davisson MT, Gardiner KJ (2007). Editing of the serotonin 2C receptor pre-mRNA Effects of the Morris Water Maze. *Gene* 391:186-197]. In addition, fully edited isoforms of the human 5-HT_{2C} receptor display a striking reduction in sensitivity to lysergic acid diethylamide (LSD) and to atypical antipsychotic drugs clozapine and loxapine, suggesting a possible role of the receptor in the etiology and pharmacology of schizophrenia [Niswender CM, Herrick-Davis K, Dilley GE, Meltzer HY, Overholser JC, Stockmeier CA, Emeson RB, Sanders-Bush E (2001). RNA Editing of the Human Serotonin 5-HT_{2C} Receptor: Alterations in Suicide and Implications for Serotonergic Pharmacotherapy. *Neuropsychopharm.* 24:478-491].

Recently, the availability of potent and selective 5-HT_{2C} receptor agonists made it possible to directly investigate the effects of 5-HT_{2C} agonists and their therapeutic potential. Thus recent studies demonstrated that selective 5-HT_{2C} agonists resulted in decreased food intake and body weight gain in normal and obese rats [Smith BM, et al. (2008). Discovery and structure-activity relationship of (1*R*)-8-chloro-2,3,4,5-tetrahydro-1-methyl-1*H*-3-benzazepine (Lorcaserin), a selective serotonin 5-HT_{2C} receptor agonist for the treatment of obesity. *J Med Chem* 51:305-313. Thomsen WJ, Grottick AJ, Menzaghi F, Reyes-Saldana H, Espitia S, Yuskin D, Whelan K, Martin M, Morgan M, Chen W, Al-Shama H, Smith B, Chalmers D, Behan D (2008) Lorcaserin, A Novel Selective Human 5-HT_{2C} Agonist: In Vitro and In Vivo Pharmacological Characterization. *J Pharmacol Exp Ther.* 325:577-587. Rosenzweig-Lipson S, Zhang J, Mazandarani H, Harrison BL, Sabb A, Sabalski J, Stack G, Welmaker G, Barrett JE, Dunlop J (2006) Antiobesity-like effects of the 5-HT_{2C} receptor agonist WAY-161503. *Brain Res.* 1073-1074:240-251. Dunlop J, Sabb AL, Mazandarani H, Zhang J, Kalgaonker S, Shukhina E, Sukoff S, Vogel RL, Stack G, Schechter L, Harrison BL, Rosenzweig-Lipson S (2005). WAY-163909 [97*bR*, 10*aR*)-1,2,3,4,8,9,10,10*a*-octahydro-7*bH*-cyclopenta-[*b*][1,4]diazepino[6,7,1*hi*]indole], a novel 5-hydroxytryptamine 2C receptor –selective agonist with anorectic activity. *J Pharmacol Exp Ther.* 313:862-869.].

Furthermore, selective 5-HT_{2C} receptor agonists produce antidepressant effects in animal models of depression comparable to those of SSRIs but with a much faster onset of action and a therapeutic window that avoids antidepressant-induced sexual dysfunction. These agonists were also effective in animal models of compulsive behavior such as scheduled induced polydipsia and they also exhibited decreased hyperactivity and aggression in rodents [Rosenzweig-Lipson S, Sabb A, Stack G, Mitchell P, Lucki I, Malberg JE, Grauer S, Brennan J, Cryan JF, Sukoff Rizzo SJ, Dunlop J, Barrett JE, Marquis KL (2007) Antidepressant-like effects of the novel, selective, 5-HT_{2C} receptor agonist WAY-163909 in rodents. *Psychopharmacology* (Berlin) 192:159-170. Rosenzweig-Lipson S, Dunlop J, Marquis KL (2007) 5-HT_{2C} receptor agonists as an innovative approach for psychiatric disorders. *Drug news Perspect*, 20: 565-571. Cryan, JF, Lucki I (2000). Antidepressant-like behavioral effects mediated by 5-Hydroxytryptamine 2C receptors. *J. Pharm. Exp. Ther.* 295:1120-1126.].

Acute or chronic administration of 5-HT_{2C} agonists decreases the firing rate of ventral tegmental area dopamine neurons but not that of substantia nigra. In addition 5-HT_{2C} agonists reduce dopamine levels in the nucleus accumbens but not in the striatum (the region of the brain mostly associated with extrapyramidal side effects) [Di Matteo, V., Di Giovanni, G., Di Mascio, M., & Esposito, E. (1999). SB 242084, a selective serotonin 2C receptor antagonist, increases dopaminergic transmission in the mesolimbic system. *Neuropharmacology* 38, 1195 – 1205. Di Giovanni, G., Di Matteo, V., Di Mascio, M., & Esposito, E. (2000). Preferential modulation of mesolimbic vs. nigrostriatal dopaminergic function by serotonin 2C/2B receptor agonists: a combined in vivo electrophysiological and microdialysis study. *Synapse* 35, 53 – 61. Marquis KL, Sabb AL, Logue SF, Brennan JA, Piesla MJ, Comery TA, Grauer SM, Ashby CR, Jr., Nguyen HQ, Dawson LA, Barrett JE, Stack G, Meltzer HY, Harrison BL, Rosenzweig-Lipson S (2007) WAY-163909 [(7bR,10aR)-1,2,3,4,8,9,10,10a-octahydro-7bH-cyclopenta-[b][1,4]diazepino[6,7,1hi]indole]: A novel 5-hydroxytryptamine 2C receptor-selective agonist with preclinical antipsychotic-like activity. *J Pharmacol Exp Ther* 320:486-496.]. Therefore it is expected that 5-HT_{2C} receptor agonists will selectively decrease mesolimbic dopamine levels without affecting the nigrostriatal pathway thus avoiding the EPS side effects of typical antipsychotics. Several 5-HT_{2C} receptor agonists have shown antipsychotic activity in animal models of schizophrenia without EPS based on the lack of effect in catalepsy [Marquis KL, Sabb AL, Logue SF, Brennan JA, Piesla MJ, Comery TA, Grauer SM, Ashby CR, Jr., Nguyen HQ, Dawson LA, Barrett JE, Stack G, Meltzer HY, Harrison BL, Rosenzweig-Lipson S (2007) WAY-163909 [(7bR,10aR)-1,2,3,4,8,9,10,10a-octahydro-7bH-cyclopenta-[b][1,4]diazepino[6,7,1hi]indole]: A novel 5-hydroxytryptamine 2C receptor-selective agonist with preclinical antipsychotic-like activity. *J Pharmacol Exp Ther* 320:486-496. Siuciak JA, Chapin DS, McCarthy SA, Guanowsky V, Brown J, Chiang P, Marala R, Patterson T, Seymour PA, Swick A, Iredale PA (2007) CP-809,101, a selective 5-HT_{2C} agonist, shows activity in animal models of antipsychotic activity. *Neuropharmacology* 52:279-290]. The antipsychotic activity of 5-HT_{2C} receptor agonists without EPS coupled with their beneficial effects in mood disorders and cognition and their antiobesity like effects render 5-HT_{2C} receptor agonists as unique agents to treat schizophrenia [Rosenzweig-Lipson S, Dunlop J, Marquis KL (2007) 5-HT_{2C} receptor agonists as an innovative ap-

proach for psychiatric disorders. *Drug news Perspect*, 20: 565-571. Dunlop J, Marquis KL, Lim HK, Leung L, Kao J, Cheesman C, Rosenzweig-Lipson S (2006). Pharmacological profile of the 5-HT_{2C} receptor agonist WAY-163909; therapeutic potential in multiple indications. *CNS Dug Rev*. 12:167-177.].

In addition 5-HT_{2C} modulation has been implicated in epilepsy [Isaac M (2005). Serotonergic 5-HT_{2C} receptors as a potential therapeutic target for the antiepileptic drugs. *Curr. Topics Med. Chem.* 5:59:67], psoriasis [Thorslund K, Nordlind K (2007). Serotonergic drugs-a possible role in the treatment of psoriasis? *Drug News Perspect* 20:521-525], Parkinson's disease and related motor disorders [Esposito E, Di Matteo V, Pierucci M, Benigno A, Di Giavanni, G (2007). Role of central 5-HT_{2C} receptor in the control of basal ganglia functions. *The Basal Ganglia Pathophysiology: Recent Advances* 97-127], behavioral deficits [Barr AM, Lahmann-Masten V, Paulus M, Gainetdinov RP, Caron MG, Geyer MA (2004). The selective serotonin-2A receptor antagonist M100907 reverses behavioral deficits in dopamine transporter knockout mice. *Neuropsychopharmacology* 29:221-228], anxiety [Dekeyne A, Mannoury la Cour C, Gobert A, Brocco M, Lejuene F, Serres F, Sharp T, Daszuta A, Soumier A, Papp M, Rivet JM, Flik G, Cremers TI, Muller O, Lavielle G, Millan MJ (2208). S32006, a novel 5-HT_{2C} receptor antagonists displaying broad-based antidepressant and anxiolytic properties in rodent models. *Psychopharmacology* 199:549-568. Nunes-de-Souza V, Nunes-de-Souza RL, Rodgers RJ, Canto-de-Souza A (2008). 5-HT₂ receptor activation in the midbrain periaqueductal grey (PAG) reduces anxiety-like behavior in mice. *Behav. Brain Res.* 187:72-79.], migraine [Leone M, Rigamonti A, D'Amico D, Grazzi L, Usai S, Bussone G (2001). The serotonergic system in migraine. *Journal of Headache and Pain* 2(Suppl. 1):S43-S46], Alzheimer's disease [Arjona AA, Pooler AM, Lee RK, Wurtman RJ (2002). Effect of a 5-HT_{2C} serotonin agonist, dexnorfenfluramine, on amyloid precursor protein metabolism in guinea pigs. *Brain Res.* 951:135-140], pain and spinal cord injury [Nakae A, Nakai K, Tanaka T, Hagihira S, Shibata M, Ueda K, Masimo T (2008). The role of RNA editing of the serotonin 2C receptor in a rat model of oro-facial neuropathic pain. *The European Journal of Neuroscience* 27:2373-2379. Nakae A, Nakai K, Tanaka T, Takashina M, Hagihira S, Shibata M, Ueda K, Mashimo T (2008). Serotonin 2C receptor mRNA editing in neuropathic pain model. *Neurosci. Res.* 60:228-231. Kao T, Shumsky JS, Jacob-Vadakot S, Timothy HB, Murray M,

Moxon, KA (2006). Role of the 5-HT_{2C} receptor in improving weight-supported stepping in adult rats spinalized as neonates. *Brain Res.* 1112:159-168.], sexual dysfunction [Motofei IG (2008). A dual physiological character for sexual function: the role of serotonergic receptors. *BJU International* 101:531-534. Shimada I, Maeno K, Kondoh Y, Kaku H, Sugasawa K, Kimura Y, Hatanaka K,; Naitou Y, Wanibuchi F, Sakamoto S,; Tsukamoto S (2008). Synthesis and structure-activity relationships of a series of benzazepine derivatives as 5-HT_{2C} receptor agonists. *Bioorg. Med. Chem.* 16:3309-3320.], smoking cessation [Fletcher PJ, Le AD, Higgins GA (2008). Serotonin receptors as potential targets for modulation of nicotine use and dependence. *Progress Brain Res.* 172:361-83], substance dependence [Bubar MJ, Cunningham KA (2008). Prospects for serotonin 5-HT_{2R} pharmacotherapy in psychostimulant abuse. *Progress Brain Res.* 172:319-46], and ocular hypertension [Sharif NA, McLaughlin MA, Kelly CR (2006). AL-34662: a potent, selective, and efficacious ocular hypotensive serotonin-2 receptor agonist. *J Ocul Pharmacol Ther.* 23:1-13].

Further, 5HT modulation can be useful in the treatment of pain, both neuropathic and nociceptive pain, see for example U.S. Patent application publication US2007/0225277. Obata, Hideaki; Ito, Naomi; Sasaki, Masayuki; Saito, Shigeru; Goto, Fumio. Possible involvement of spinal noradrenergic mechanisms in the antiallodynic effect of intrathecally administered 5-HT_{2C} receptor agonists in the rats with peripheral nerve injury. *European Journal of Pharmacology* (2007), 567(1-2), 89-94. Serotonin_{2C} receptor mRNA editing in neuropathic pain model. Nakae, Aya; Nakai, Kunihiro; Tanaka, Tatsuya; Takashina, Masaki; Hagihira, Satoshi; Shibata, Masahiko; Ueda, Koichi; Mashimo, Takashi. Department of Anesthesiology & Intensive Care Medicine, Graduate School of Medicine, Osaka University, *Neuroscience Research* (Amsterdam, Netherlands) (2008), 60(2), 228-231. Antiallodynic effects of intrathecally administered 5-HT_{2C} receptor agonists in rats with nerve injury. Obata, Hideaki; Saito, Shigeru; Sakurazawa, Shinobu; Sasaki, Masayuki; Usui, Tadashi; Goto, Fumio. Department of Anesthesiology, Gunma University Graduate School of Medicine, Maebashi, Gunma, Japan. *Pain* (2004), 108(1-2), 163-169. Influence of 5,7-dihydroxytryptamine (5,7-DHT) on the antinociceptive effect of serotonin (5-HT) 5-HT_{2C} receptor agonist in male and female rats. Brus, Ryszard; Kasperska, Alicja;

Oswiecimska, Joanna; Szkilnik, Ryszard. Department of Pharmacology, Silesian Medical University, Zabrze, Pol. *Medical Science Monitor* (1997), 3(5), 654-656.

Modulation of 5HT₂ receptors may be beneficial in the treatment of conditions related to bladder function, in particular, urinary incontinence. [Discovery of a novel azepine series of potent and selective 5-HT_{2C} agonists as potential treatments for urinary incontinence. Brennan, Paul E.; Whitlock, Gavin A.; Ho, Danny K. H.; Conlon, Kelly; McMurray, Gordon. *Bioorganic & Medicinal Chemistry Letters* (2009), 19(17), 4999-5003. Investigation of the role of 5-HT₂ receptor subtypes in the control of the bladder and the urethra in the anesthetized female rat. Mbaki, Y.; Ramage, A. G. Department of Pharmacology, University College London, London, UK. *British Journal of Pharmacology* (2008), 155(3), 343-356.] In particular, compounds with agonist activity at 5-HT_{2C} have been shown to be useful in treating urinary incontinence, see for example U.S. Patent application publications US2008/0146583 and US 2007/0225274.

In the use and the method of the invention, an effective quantity of one or more compounds, as a rule formulated in accordance with pharmaceutical and veterinary practice, is administered to the individual to be treated, preferably a mammal, in particular a human being, productive animal or domestic animal. Whether such a treatment is indicated, and in which form it is to take place, depends on the individual case and is subject to medical assessment (diagnosis) which takes into consideration signs, symptoms and/or malfunctions which are present, the risks of developing particular signs, symptoms and/or malfunctions, and other factors.

Actual dosage levels of active ingredients in the pharmaceutical compositions of the present invention can be varied so as to obtain an amount of the active compound(s) that is effective to achieve the desired therapeutic response for a particular subject (e.g., a mammal, preferably, a human (patient)), compositions and mode of administration. The selected dosage level will depend upon the activity of the particular compound, the route of administration, the severity of the condition being treated and the condition and prior medical history of the patient being treated. However, it is within the skill of the art to start doses of the compound at levels lower than required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

Compounds of the present invention can also be administered to a subject as a pharmaceutical composition comprising the compounds of interest in combination with at least one pharmaceutically acceptable carriers. The phrase "therapeutically effective amount" of the compound of the present invention means a sufficient amount of the compound to treat disorders, at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well-known in the medical arts. For example, it is well within the skill of the art to start doses of the compound at levels lower than required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

The total daily dose of the compounds of this invention administered to a subject (namely, a mammal, such as a human) ranges from about 0.01 mg/kg body weight to about 100 mg/kg body weight. More preferable doses can be in the range of from about 0.01 mg/kg body weight to about 30 mg/kg body weight. If desired, the effective daily dose can be divided into multiple doses for purposes of administration. Consequently, single dose compositions may contain such amounts or submultiples thereof to make up the daily dose.

In one aspect, the present invention provides pharmaceutical compositions. The pharmaceutical compositions of the present invention comprise the compounds of the present invention or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt or solvate thereof. The pharmaceutical compositions of the present invention comprise compounds of the present invention that can be formulated together with at least one non-toxic pharmaceutically acceptable carrier.

In yet another embodiment, the present invention provides a pharmaceutical composition comprising compounds of the present invention or an N-oxide, a tautomer-

ic form, a stereoisomer or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable carriers, alone or in combination with one or more compounds that are not the compounds of the present invention. Examples of one or more compounds that can be combined with the compounds of the present invention in pharmaceutical compositions, include, but are not limited to, one or more cognitive enhancing drugs.

The pharmaceutical compositions of this present invention can be administered to a subject (e.g., a mammal, such as a human) orally, rectally, parenterally, intracisterally, intravaginally, intraperitoneally, topically (as by powders, ointments or drops), buccally or as an oral or nasal spray. The term "parenterally" as used herein, refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

The term "pharmaceutically acceptable carrier" as used herein, means a non-toxic, inert solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. Some examples of materials which can serve as pharmaceutically acceptable carriers are sugars such as, but not limited to, lactose, glucose and sucrose; starches such as, but not limited to, corn starch and potato starch; cellulose and its derivatives such as, but not limited to, sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as, but not limited to, cocoa butter and suppository waxes; oils such as, but not limited to, peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols; such a propylene glycol; esters such as, but not limited to, ethyl oleate and ethyl laurate; agar; buffering agents such as, but not limited to, magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as, but not limited to, sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

Pharmaceutical compositions of the present invention for parenteral injection comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile

injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol and the like), vegetable oils (such as olive oil), injectable organic esters (such as ethyl oleate) and suitable mixtures thereof. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms can be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of the drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This can be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound may be mixed with at least one inert, pharmaceutically acceptable excipient or carrier, such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol and silicic acid; b) binders such as carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose and acacia; c) humectants such as glycerol; d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates and sodium carbonate; e) solution retarding agents such as paraffin; f) absorption accelerators such as quaternary ammonium compounds; g) wetting agents such as cetyl alcohol and glycerol monostearate; h) absorbents such as kaolin and bentonite clay and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such carriers as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The solid dosage forms of tablets, dragees, capsules, pills and granules can be prepared with coatings and shells such as enteric coatings and other coatings well-known in the pharmaceutical formulating art. They may optionally contain opacifying agents and may also be of a composition such that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

The active compounds can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned carriers.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydro-

furfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan and mixtures thereof.

Besides inert diluents, the oral compositions may also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, tragacanth and mixtures thereof.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating carriers or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

Compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals which are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients and the like. The preferred lipids are natural and synthetic phospholipids and phosphatidyl cholines (lecithins) used separately or together.

Methods to form liposomes are known in the art. See, for example, Prescott, Ed., *Methods in Cell Biology*, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 et seq.

Dosage forms for topical administration of a compound of the present invention include powders, sprays, ointments and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives, buffers or propellants which may be required. Ophthalmic formulations, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

The compounds of the present invention can be used in the form of pharmaceutically acceptable salts derived from inorganic or organic acids. The phrase "pharmaceutically acceptable salt" means those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like and are commensurate with a reasonable benefit/risk ratio.

Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al. describe pharmaceutically acceptable salts in detail in (*J. Pharmaceutical Sciences*, 1977, 66: 1 et seq.). The salts can be prepared in situ during the final isolation and purification of the compounds of the invention or separately by reacting a free base function with a suitable organic acid. Representative acid addition salts include, but are not limited to acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate (isothionate), lactate, malate, maleate, methanesulfonate, nicotine, 2-naphthalenesulfonate, oxalate, palmitate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, phosphate, glutamate, bicarbonate, p-toluenesulfonate and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides such as, but not limited to, methyl, ethyl, propyl, and butyl chlorides, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl and diamyl sulfates; long chain halides such as, but not limited to, decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; arylalkyl halides like benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products are thereby obtained. Examples of acids which can be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, hydrobromic acid, sulfuric acid, and phosphoric acid and such organic acids as acetic acid, fumaric acid, maleic acid, 4-methylbenzenesulfonic acid, succinic acid and citric acid.

Basic addition salts can be prepared in situ during the final isolation and purification of compounds of this invention by reacting a carboxylic acid-containing moiety with a suitable base such as, but not limited to, the hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation or with ammonia or an organic primary,

secondary or tertiary amine. Pharmaceutically acceptable salts include, but are not limited to, cations based on alkali metals or alkaline earth metals such as, but not limited to, lithium, sodium, potassium, calcium, magnesium and aluminum salts and the like and nontoxic quaternary ammonia and amine cations including ammonium, tetramethylammonium, tetraethylammonium, methylammonium, dimethylammonium, trimethylammonium, triethylammonium, diethylammonium, ethylammonium and the like. Other representative organic amines useful for the formation of base addition salts include ethylenediamine, ethanolamine, diethanolamine, piperidine, piperazine and the like.

The compounds of the present invention can exist in unsolvated as well as solvated forms, including hydrated forms, such as hemi-hydrates. In general, the solvated forms, with pharmaceutically acceptable solvents such as water and ethanol among others are equivalent to the unsolvated forms for the purposes of the invention.

The following examples serve to explain the invention without limiting it.

Examples

The compounds were either characterized via proton-NMR in d_6 -dimethylsulfoxide or d -chloroform or d_4 -methanol on a 400 MHz or 500 MHz NMR instrument (Bruker AVANCE), or by mass spectrometry, generally recorded via HPLC-MS in a fast gradient on C18-material (electrospray-ionisation (ESI) mode).

The magnetic nuclear resonance spectral properties (NMR) refer to the chemical shifts (δ) expressed in parts per million (ppm). The relative area of the shifts in the ^1H -NMR spectrum corresponds to the number of hydrogen atoms for a particular functional type in the molecule. The nature of the shift, as regards multiplicity, is indicated as singlet (s), broad singlet (s. br.), doublet (d), broad doublet (d br.), triplet (t), broad triplet (t br.), quartet (q), quintet (quint.) and multiplet (m).

Enantiomers were separated / purified either by chiral supercritical fluid chromatography (SFC) (method A) or by chiral HPLC (method B).

Method A - Chiral Analytical SFC

Analytical SFC was performed on an Aurora A5 SFC Fusion and Agilent 1100 system running under Agilent Chemstation software control. The SFC system included a 10-way column switcher, CO₂ pump, modifier pump, oven, and backpressure regulator. The mobile phase comprised of supercritical CO₂ supplied by a beverage-grade CO₂ cylinder with a modifier mixture of methanol at a flow rate of 3 mL/min. Oven temperature was at 35 °C and the outlet pressure at 150 bar. The UV detector was set to collect at wavelengths of 220nm and 254 nm. The mobile phase gradient started with 5% modifier and held it for 0.1 minutes at a flow rate of 1 mL/min, then the flow rate was ramped up to 3 mL/min and held for 0.4 min. The modifier was ramped from 5% to 15% over the next 8 minutes at 3 mL/min then held for 1 minute at 15% modifier (3 mL/min). The gradient was ramped down from 15% to 5% modifier over 0.5 min (3 mL/min). The instrument was fitted with a Chiralpak AS-H column with dimensions of 4.6 mm i.d. x 150 mm length with 5 µm particles.

Method B - Chiral HPLC

System: KNAUER preparative HPLC

Pump: Preparative pump 1800

Detector: Smartline UV detector 2600 257 nm

Sample pump: Knauer HPLC-Pump K-120

Fractional collector: Smartline Valves Drive S6

Software: ChromGate® V3.1.7, KNAUER Instrument Control

Column: Welk O 4.6 mm ID x 250 mm

Column temperature: 25°C

Mobile phase: hexane/isopropylamine 95/5

Flow rate: 1 ml/min

I. Preparation Examples

Example 1

8,8-Dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline (compound of the formula I.g, I.h or I.i in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-1 of Table B)

1.1 Preparation of tert-butyl 1-(3-methylbut-2-enoyl)-2,3-dihydro-1H-benzo[e][1,4]diazepine-4(5H)-carboxylate

1g (4.04 mmol) of tert-butyl 2,3-dihydro-1H-benzo[e][1,4]diazepine-4(5H)-carboxylate was dissolved in 20 mL of dichloromethane and treated with 1.54 mL (8.86 mL) of triethylamine followed by 0.53 mL (4.83 mmol) of 3-methylbut-2-enoyl chloride. The reaction mixture was stirred over night at room temperature, poured onto water (50 mL) and extracted three times with 50 mL of dichloromethane each. The organic phases were combined, washed with a saturated solution of sodium chloride, dried over magnesium sulfate and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 5-10% methanol in dichloromethane) to yield 1.27 g of the title compound as a white solid.

ESI-MS: m/z (%): 275.10 (100, $[M-C_4H_9+H]^+$).

1.2 Preparation of tert-butyl 8,8-dimethyl-6-oxo-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline-2(6H)-carboxylate

A solution of 0.33 g (1.0 mmol) of tert-butyl 1-(3-methylbut-2-enoyl)-2,3-dihydro-1H-benzo[e][1,4]diazepine-4(5H)-carboxylate of step 1.1 was irradiated with a 150 W mercury lamp in an immersion well reactor with a pyrex filter in 20 mL of acetone until completion of the reaction monitored by liquid chromatography. The solution was concentrated in vacuo and the residue was purified by column chromatography on silica (eluent: 10-30% ethyl acetate in heptane) to yield 202 mg of the title compound as a beige solid.

ESI-MS: m/z (%): 275.10 (100, $[M-C_4H_9+H]^+$).

1.3 Preparation of tert-butyl 8,8-dimethyl-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline-2(6H)-carboxylate

A solution of 38 mg (0.115 mmol) tert-butyl 8,8-dimethyl-6-oxo-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline-2(6H)-carboxylate of step 1.2 in 1 mL of tetrahydrofuran was treated with 0.46 mL of 1 molar solution of borohydride-tetrahydrofuran complex in tetrahydrofuran. The mixture was stirred over night, and then quenched with water and diluted hydrochloric acid. The pH was adjusted to pH 9 by addition of aqueous sodium hydroxide solution and the mixture extracted three times with 10 mL of dichloromethane each. The solution was dried over magnesium sulfate and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 20-30% ethyl acetate in heptane) to yield 21 mg of the title compound as a clear oil.

ESI-MS: m/z (%): 317.20 (100, $[M+H]^+$).

1.4 Preparation of 8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline

287 mg (0.9 mmol) of tert-butyl 8,8-dimethyl-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline-2(6H)-carboxylate of step 1.3 were dissolved in 5 mL of dichloromethane and treated with 2.5 mL of trifluoroacetic acid. The mixture was stirred over night at room temperature and then extracted once with water (10 mL). The organic phase was then extracted twice with a diluted solution of sodium hydroxide (10 mL each), dried over magnesium sulfate and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 10-30% methanol in dichloromethane) to yield 141 mg of the title compound as a yellow oil.

ESI-MS: m/z (%): 217.15 (100, $[M+H]^+$).

$^1\text{H-NMR}$ (500 MHz, CDCl_3): δ = 7.15 (d, 1H), 6.85 (d, 1H), 6.75 (m, 1H), 3.80 (s, 2H), 3.15 (m, 2H), 3.00 (s, 4H), 1.65 (m, 2H), 1.20 (s, 6H).

Example 2

1,2,3,4-Tetrahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopent[2]en]-6(7H)-one 2,2,2-trifluoroacetate (compound of the formula I.j, I.k or I.l in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-481 of Table B)

- 2.1 Preparation of tert-butyl 6-oxo-3,4,6,7-tetrahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopent[2]ene]-2(1H)-carboxylate (compound of the formula I.g, I.h or I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-401 of Table B, wherein however R¹ is not H, but Boc)

A solution of 50 mg (0.128 mmol) tert-butyl 9-chloro-1-(2-cyclopentenylacetyl)-2,3-dihydro-1H-benzo[e][1,4]diazepine-4(5H)-carboxylate was dissolved in 1 mL of degassed dry acetonitrile under argon, treated with 1.4 mg (0.006 mmol) of palladium acetate, 6.10 mg (0.013 mmol) dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine and 53 mg (0.384 mmol) potassium carbonate. The mixture was stirred for 10 hours at 120°C in a synthesis microwave system. After cooling to room temperature the mixture was filtered over celite, diluted with dichloromethane and then extracted twice with a diluted solution of sodium chloride (10 mL each), dried over magnesium sulfate and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 10-30% ethyl acetate in heptane) to yield 40 mg of the title compound as a yellow oil.

ESI-MS: *m/z* (%): 299.10 (100, [M-C₄H₉+H]⁺).

- 2.2 Preparation of 1,2,3,4-tetrahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopent[2]en]-6(7H)-one 2,2,2-trifluoroacetate

Boc-deprotection of the compound obtained in step 2.1 analogously to example 1.4 yielded the title compound.

ESI-MS: *m/z* (%): 255 (100, [M+H]⁺).

Example 3

8-Methyl-8-phenyl-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-6(2H)-one, hydrochloride (compound of the formula I.j, I.k or I.l in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-161 of Table B)

392 mg (1 mmol) of tert-butyl 1-(3-phenylbut-2-enoyl)-2,3-dihydro-1H-benzo[e][1,4]diazepine-4(5H)-carboxylate were treated with 1 g of polyphosphoric acid. The mixture was heated to 90°C for 2 hours and then quenched with a saturated solution of potassium carbonate at 0°C. The mixture was extracted three times with ethyl acetate (20 mL each), the combined organic fractions dried over magnesium sulfate and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 10-30% methanol in dichloromethane) to yield 86 mg of the title compound as a yellow foam.

ESI-MS: m/z (%): 293.1 (100, $[M+H]^+$).

¹H-NMR (500 MHz, DMSO-*d*₆): δ = 10.05 (bs, 1H), 9.85 (bs, 1H), 7.40 (d, 1H), 7.35 (d, 1H), 7.25 (m, 2H), 7.20 (m, 2H), 7.10 (m, 2H), 4.55 (m, 1H), 4.35 (m, 1H), 4.15 (m, 1H), 3.55 (m, 1H), 3.25 (m, 2H), 3.10 (m, 1H), 2.70 (m, 1H), 1.60 (s, 3H).

The following example was prepared analogously to example 1.

Example 4

8,8-Dimethyl-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-*ij*]quinolin-6(2H)-one (compound of the formula I.j, I.k or I.l in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-1 of Table B)

ESI-MS: m/z (%): 231 (100, $[M+H]^+$).

¹H-NMR (500 MHz, CDCl₃): δ = 7.20 (d, 1H), 7.00 (m, 2H), 4.10 (s, 4H), 3.20 (m, 2H), 2.50 (s, 2H), 1.30 (s, 6H).

The following examples 5 and 6 were prepared analogously to example 2.

Example 5

1,2,3,4-Tetrahydrospiro[[1,4]diazepino[6,7,1-*ij*]quinoline-8,1'-cyclopentan]-6(7H)-one (compound of the formula I.j, I.k or I.l in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-361 of Table B)

ESI-MS: m/z (%): 257 (100, $[M+H]^+$).

Example 6

1,2,3,4,6,7-Hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopentane] (compound of the formula I.g, I.h or I.i in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-361 of Table B)

ESI-MS: m/z (%): 243 (100, $[M+H]^+$).

1H -NMR (500 MHz, $CDCl_3$): δ = 7.15 (d, 1H), 6.95 (d, 1H), 6.80 (m, 1H), 3.90 (s, 2H), 3.20 (m, 2H), 3.05 (m, 4H), 1.95-1.65 (m, 10H).

The following examples 7 to 13 were prepared analogously to example 1.

Example 7

12a-Methyl-4,5,6,7,10,11,12,12a-octahydrocyclopenta[c][1,4]diazepino[6,7,1-ij]quinolin-9(9aH)-one (racemic cis diastereomer of the compound of the formula I.j, I.k or I.l in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-521 of Table B)

ESI-MS: m/z (%): 257 (100, $[M+H]^+$).

Example 8

8-Methyl-8-(trifluoromethyl)-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline (single enantiomer of the compound of the formula I.g, I.h or I.i in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-81 of Table B)

ESI-MS: m/z (%): 271 (100, $[M+H]^+$).

The retention time according to method A is 1.396 min.

Example 9

8-Methyl-8-(trifluoromethyl)-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline (single enantiomer of the compound of the formula I.g, I.h or I.i in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-81 of Table B)

ESI-MS: m/z (%): 271 (100, $[M+H]^+$).

The retention time according to method A is 1.543 min.

Example 10

12a-Methyl-4,5,6,7,10,11,12,12a-octahydrocyclopenta[c][1,4]diazepino[6,7,1-ij]quinolin-9(9aH)-one (racemic trans diastereomer of the compound of the formula I.j, I.k or I.l in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-521 of Table B)

ESI-MS: m/z (%): 257 (100, $[M+H]^+$).

Example 11

12a-Methyl-4,5,6,7,9,9a,10,11,12,12a-decahydrocyclopenta[c][1,4]diazepino[6,7,1-ij]quinoline (racemic cis diastereomer of the compound of the formula I.g, I.h or I.i in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-521 of Table B)

ESI-MS: m/z (%): 243 (100, $[M+H]^+$).

1H -NMR (500 MHz, $CDCl_3$): δ = 7.15 (d, 1H), 6.90 (d, 1H), 6.80 (m, 1H), 3.90 (m, 2H), 3.15 (m, 1H), 3.05 (m, 3H), 2.95 (m, 1H), 2.80 (m, 1H), 1.95 (m, 1H), 1.90 (m, 1H), 1.80 (m, 2H), 1.60 (m, 1H), 1.55 (m, 1H), 1.30 (m, 1H), 1.20 (s, 3H).

Example 12

8-Methyl-8-(trifluoromethyl)-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-6(2H)-one (racemic mixture of compound of the formula I.j, I.k or I.l in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-81 of Table B)

ESI-MS: m/z (%): 285 (100, $[M+H]^+$).

Example 13

12a-Methyl-4,5,6,7,9,9a,10,11,12,12a-decahydrocyclopenta[c][1,4]diazepino-[6,7,1-ij]quinoline hydrochloride (racemic trans diastereomer of the compound of the formula I.g, I.h or I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-521 of Table B)

ESI-MS: *m/z* (%): 243 (100, [M+H]⁺).

¹H-NMR (500 MHz, CDCl₃): δ = 9.45 (bs, 1H), 9.05 (bs, 1H), 7.05 (d, 1H), 7.00 (d, 1H), 6.70 (m, 1H), 4.30 (m, 1H), 3.95 (m, 1H), 3.55 (m, 2H), 3.35 (m, 1H), 3.25 (m, 2H), 2.95 (m, 1H), 1.95 (m, 1H), 1.85 (m, 1H), 1.80 (m, 1H), 1.70 (m, 1H), 1.55 (m, 1H), 1.40 (m, 1H), 1.25 (m, 1H), 0.90 (s, 3H).

The following examples 14 and 15 were prepared analogously to example 3.

Example 14

8-Methyl-8-phenyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline hydrochloride (racemic mixture in form of the salt of the compound of the formula I.g, I.h or I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-161 of Table B)

ESI-MS: *m/z* (%): 279 (100, [M+H]⁺)

¹H-NMR (500 MHz, CDCl₃): δ = 7.22 - 7.27 (m, 2H), 7.11 - 7.19 (m, 3H), 6.95 - 7.04 (m, 2H), 6.78 (dd, *J* = 8.0, 7.5 Hz, 1H), 3.99 (d, *J* = 14.4 Hz, 1H), 3.87 99 (d, *J* = 14.5 Hz, 1H), 3.20 - 3.26 (m, 1H), 3.01 - 3.08 (m, 4H), 2.87 - 2.94 (m, 1H), 2.09 - 2.15 (m, 1H), 1.92-1.99 (m, 1H), 1.73 (s, 3H) ppm.

Example 15

8-Methyl-8-phenyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline (racemic mixture of compound of the formula I.g, I.h or I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-161 of Table B)

ESI-MS: *m/z* (%): 279 (100, [M+H]⁺).

¹H NMR (CDCl₃, 500MHz): [ppm]: 7.25 (m, 2H), 7.15 (m, 3H), 7.05 (d, 1H), 6.95 (d, 1H), 6.80 (m, 1H), 4.00 (m, 1H), 3.90 (m, 1H), 3.25 (m, 1H), 3.05 (m, 4H), 2.90 (m, 1H), 2.10 (m, 1H), 1.95 (m, 1H), 1.75 (s, 3H).

Example 16

(R)-5,6,7,9,9a,10,11,12-Octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline (2,2,2-trifluoroacetate) (R-enantiomer of the compound of the formula I.a, I.b or I.c in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row A-865 of Table A)

16.1 Preparation of (R)-tert-butyl 9-oxo-6,7,9a,10,11,12-hexahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline-5(9H)-carboxylate

65 mg of (R)-tert-butyl 9-chloro-1-(pyrrolidine-2-carbonyl)-2,3-dihydro-1H-benzo[e][1,4]diazepine-4(5H)-carboxylate was stirred together with 1,8 mg of 2'-(di-tert-butylphosphino)-N,N-dimethylbiphenyl-2-amine, 2.5 mg Pd₂(dba)₃ and 19 mg sodium 2-methylpropan-2-olate at 100°C over 10 hours in toluene. The reaction mixture was then cooled to room temperature and evaporated. The residue was purified by column chromatography on silica (eluent: 10-25% ethyl acetate in heptane) to yield 20 mg of the title compound as a beige solid.

ESI-MS: *m/z* (%): 344.20 (100, [M+H]⁺).

16.2 Preparation of (R)-tert-butyl 6,7,9a,10,11,12-hexahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline-5(9H)-carboxylate (R-enantiomer of the compound of the formula I.a, I.b or I.c in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row A-121 of Table A, wherein however R¹ is not H, but Boc).

Reduction of the compound obtained in step 16.1 analogously to example 1.3 yielded the title compound.

ESI-MS: *m/z* (%): 330.20 (100, [M+H]⁺).

16.3 Preparation of (R)-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline bis(2,2,2-trifluoroacetate) (R-enantiomer of the compound of the formula I.a, I.b or I.c in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row A-121 of Table A)

Boc-deprotection of the compound obtained in step 16.2 analogously to example 1.4 yielded the title compound.

ESI-MS: *m/z* (%): 230.15 (100, [M+H]⁺).

Example 17 was prepared analogously to example 16 employing (S)-tert-butyl 9-chloro-1-(pyrrolidine-2-carbonyl)-2,3-dihydro-1H-benzo[e][1,4]diazepine-4(5H)-carboxylate as starting material which was derived from (R)-1-(((9H-fluoren-9-yl)methoxy)carbonyl)pyrrolidine-2-carboxylic acid.

Example 17

(S)-5,6,7,9,9a,10,11,12-Octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline (2,2,2-trifluoroacetate) (S-enantiomer of the compound of the formula I.a, I.b or I.c in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row A-865 of Table A)

ESI-MS: *m/z* (%): 230 (100, [M+H]⁺).

The following examples 18 to 23 were prepared analogously to example 1.

Example 18

9-Chloro-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline, (2,2,2-trifluoroacetate) (compound of the formula I.g in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-3 of Table B)

ESI-MS: *m/z* (%): 251 (100, [M+H]⁺).

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 8.95 (bs, 2H), 7.35 (d, 1H), 7.25 (d, 1H), 4.15 (s, 2H), 3.30-3.20 (m, 6H), 1.65 (m, 2H), 1.25 (s, 6H).

Example 19

10-Chloro-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline (2,2,2-trifluoroacetate) (compound of the formula I.h in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-3 of Table B)

ESI-MS: *m/z* (%): 251 (100, [M+H]⁺).

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 9.10 (bs, 2H), 7.35 (m, 1H), 7.20 (m, 1H), 4.15 (s, 2H), 3.30-3.20 (m, 6H), 1.65 (m, 2H), 1.25 (s, 6H).

Example 20

9-Chloro-8,8-dimethyl-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-6(2H)-one 2,2,2-trifluoroacetate (compound of the formula I.j in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-3 of Table B)

ESI-MS: *m/z* (%): 265 (100, [M+H]⁺).

Example 21

10-Chloro-8,8-dimethyl-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-6(2H)-one 2,2,2-trifluoroacetate (compound of the formula I.k in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-3 of Table B)

ESI-MS: *m/z* (%): 265 (100, [M+H]⁺).

Example 22

9-Fluoro-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline (compound of the formula I.g in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-2 of Table B)

ESI-MS: *m/z* (%): 235 (100, [M+H]⁺).

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 6.95 (m, 1H), 6.55 (m, 1H), 3.95 (s, 2H), 3.20-3.10 (m, 6H), 1.70 (m, 2H), 1.40 (s, 6H).

Example 23

10-Fluoro-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline (2,2,2-trifluoroacetate) (compound of the formula I.h in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-2 of Table B)

ESI-MS: *m/z* (%): 235 (100, [M+H]⁺).

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 9.05 (bs, 2H), 7.05 (d, 1H), 6.80 (m, 1H), 4.20 (s, 2H), 3.35 (m, 2H), 3.30 (m, 3.25), 3.25 (m, 2H), 1.70 (m, 2H), 1.30 (s, 6H).

The following examples 24 to 26 were prepared analogously to example 2.

Example 24

1,2,3,4,6,7-Hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclohexane] (2,2,2-trifluoroacetate) (compound of the formula I.g, I.h or I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-401 of Table B)

ESI-MS: *m/z* (%): 257.2 (100, [M+H]⁺).

¹H NMR (CDCl₃, 500MHz): [ppm]: 8.85 (bs, 2H), 7.45 (d, 1H), 7.15 (d, 1H), 6.90 (m, 1H), 4.15 (s, 2H), 3.25 (m, 4H), 3.15 (m, 2H), 1.80 (m, 2H), 1.70 (m, 4H), 1.55 (m, 4H), 1.30 (m, 2H).

Example 25

1',2',3',4',6',7'-Hexahydrospiro[cyclobutane-1,8'-[1,4]diazepino[6,7,1-ij]quinoline], hydrochloride
(compound of the formula I.g, I.h or I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-321 of Table B)

ESI-MS [M+H]⁺ = 229.20

¹H NMR (CDCl₃, 500MHz): [ppm]: 8.80 (bs, 2H), 7.50 (d, 1H), 6.95 (d, 1H), 6.90 (m, 1H), 4.10 (s, 2H), 3.25 (m, 2H), 3.20 (m, 2H), 3.10 (m, 2H), 2.35 (m, 2H), 2.00 (m, 2H), 1.90 (m, 4H).

Example 26

9'-Fluoro-1',2',3',4',6',7'-hexahydrospiro[cyclobutane-1,8'-[1,4]diazepino[6,7,1-*ij*]quinoline], hydrochloride

(compound of the formula I.g in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-322 of Table B)

ESI-MS [M+H⁺] = 247.15

¹H NMR (DMSO-d₆, 500MHz): [ppm]: 9.00 (bs, 2H), 7.25 (m, 1H), 6.80 (m, 1H), 4.10 (s, 2H), 3.25 (m, 2H), 3.20 (m, 2H), 3.10 (m, 2H), 2.65 (m, 2H), 2.00 (m, 4H), 1.85 (m, 2H).

The following examples 27 and 28 were prepared analogously to example 1.

Example 27

Enantiomer of 7,8,8-trimethyl-1,2,3,4,7,8-hexahydro-6*H*-[1,4]diazepino[6,7,1-*ij*]quinoline]

(enantiomer of the compound of the formula I.g, I.h or I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-25 of Table B)

ESI-MS [M+H⁺] = 231.20

¹H NMR (CDCl₃, 500MHz): [ppm]: 7.25 (d, 1H), 7.00 (d, 1H), 6.85 (dd, 1H), 4.00 (m, 1H), 3.90 (m, 1H), 3.15 (m, 4H), 3.05 (m, 2H), 1.85 (m, 1H), 1.30 (s, 3H), 1.10 (s, 3H), 0.95 (d, 3H).

The retention time of the Boc-protected precursor according to method B is 10.363.

Example 28

Enantiomer of 7,8,8-trimethyl-1,2,3,4,7,8-hexahydro-6*H*-[1,4]diazepino[6,7,1-*ij*]quinoline]

(enantiomer of the compound of the formula I.g, I.h or I.i in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-25 of Table B)

ESI-MS $[M+H]^+ = 231.20$

1H NMR ($CDCl_3$, 500MHz): [ppm]: 7.25 (d, 1H), 7.00 (d, 1H), 6.85 (dd, 1H), 4.00 (m, 1H), 3.90 (m, 1H), 3.15 (m, 4H), 3.05 (m, 2H), 1.85 (m, 1H), 1.30 (s, 3H), 1.10 (s, 3H), 0.95 (d, 3H).

The retention time of the Boc-protected precursor according to method B is 11.410 min.

The following example 29 was prepared analogously to example 1.

Example 29

3-Methyl-1',2',3',4',6',7'-hexahydrospiro[cyclobutane-1,8'-[1,4]diazepino[6,7,1-*ij*]quinoline], hydrochloride

(compound of the formula I.p, I.q or I.r in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-321 of Table B)

ESI-MS $[M+H]^+ = 243.20$

1H NMR ($DMSO-d_6$, 500MHz): [ppm]: 9.30 (bs, 1H), 8.55 (bs, 1H), 7.60 (d, 1H), 7.20 (d, 1H), 6.95 (dd, 1H), 4.20 (m, 1H), 4.05 (m, 1H), 3.35 (m, 1H), 3.20 (m, 2H), 2.90 (m, 1H), 2.40 (m, 1H), 2.25 (m, 1H), 2.05 (m, 1H), 1.95 (m, 6H), 1.2 (d, 3H).

Example 30

3-Benzyl-2',3',7',8',9',10'-hexahydro-1H-3',8',10a'-triazacyclohepta[de]naphthalene

(compound of the formula I.a, I.b or I.c in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row A-181 of Table A)

30.1 Preparation of 2-(4-benzyl-3,4-dihydro-2H-quinoxalin-1-yl)-acetamide

A solution of 1 g (4.46 mmol) 1-benzyl-1,2,3,4-tetrahydroquinoline in 10 ml of dimethylformamide was treated with 3.14 ml (18.15 mmol) of N-Ethyl-N-isopropylpropan-2-amine, and subsequently 863 mg (9.23 mmol) 2-chloroacetamide

was added. The reaction mixture was heated in a microwave system at 100°C for four hours. The resulting mixture was quenched with water and extracted once with ethylacetate. The aqueous phase was set to pH 10 with sodium hydroxide solution and extracted 3x with ethylacetate. The combined organic extracts were dried with magnesium sulfate, evaporated till dryness and directly purified by column chromatography on silica (eluent: starting with heptane and then up to 100 % ethylacetate) to yield 1.224 g of the title compound.

ESI-MS: m/z (%): 282.10 (100, [M+H]⁺).

30.2 Preparation of 2-(4-Benzyl-3,4-dihydro-2H-quinoxalin-1-yl)-ethylamine

To 1.224 g (4.35 mmol) of 2-(4-benzyl-3,4-dihydro-2H-quinoxalin-1-yl)-acetamide were added 10.88 ml (21.75 mmol) borane dimethylsulfide THF solution and subsequently heated to 60°C for six hours in a microwave system. The mixture was quenched with 1 molar hydrochloric acid and methanol and heated for 15 minutes at 60°C in a microwave system. The reaction mixture was diluted with ethylacetate and extracted 3x with 1 molar hydrochlorid acid. The combined aqueous phases were set to pH 10 with sodium hydroxide solution and extracted 3x with dichloromethane. The combined organic phases were dried and evaporated to yield 1.089 g of the title compound as an oil.

ESI-MS: m/z (%): 268.15 (100, [M+H]⁺).

30.3 Preparation of 3-benzyl-2,3,7,8,9,10-hexahydro-1H-3,8,10a-triaza-cyclohepta[de]-naphthalene

A solution of 1.089 g (4.07 mmol) of 2-(4-benzyl-3,4-dihydro-2H-quinoxalin-1-yl)-ethylamine and 20 ml ethanol was treated with 122 mg (4.07 mmol) of formaldehyde and 511 mg (4.48 mmol) of trifluoroacetic acid and stirred over night at room temperature. The solvent was evaporated and dichloromethane was added to the crude mixture. The organic phase was washed with 1 molar sodium hydroxide solution. The organic phase was dried and purified by column chromatography on silica (eluent:

starting with dichloromethane and then up to 100 % methanol) to yield 448 mg of the title compound.

ESI-MS $[M+H]^+ = 280.10$ (100, $[M+H]^+$).

^1H NMR (CDCl_3 , 500MHz): [ppm]: 7.35 (m, 3H), 7.30 (m, 2H), 6.80 (dd, 1H), 6.65 (d, 1H), 6.60 (d, 1H), 4.55 (s, 2H), 4.10 (s, 2H), 3.30 (m, 4H), 3.25 (m, 2H), 3.10 (m, 2H).

Example 31

9-Fluoro-1-methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane], trifluoroacetic acid salt

(compound of the formula I.m in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-322 of Table B)

31.1 Preparation of 2-cyclobutylidene-N-(3-fluorophenyl)-acetamide

1.5 g (13.50 mmol) of 3-fluoroaniline were dissolved in 20 ml of dichloromethane and treated with 5.19 g (29.7 mmol) of N-ethyl-N-isopropylpropan-2-amine. Within 10 min 12.82 g (16.20 mmol) of 2-cyclobutylideneacetyl chloride were added at 0°C and the solution was stirred overnight at room temperature. The mixture was poured on ice water and extracted 2x with dichloromethane. The combined organic phases were washed once with saturated sodium chloride solution, dried with MgSO_4 and evaporated in vacuo. The crude oil was treated with heptane and decanted twice, redissolved in ether-dichloromethane and treated with heptane till crystallization, yielding 2.2g (77%) of a beige solid.

ESI-MS: m/z (%): 206 (100, $[M+H]^+$).

31.2 Preparation of 5'-fluoro-1'H-spiro[cyclobutane-1,4'-quinolin]-2'(3'H)-one

A solution of 400 mg (1.94 mmol) of 2-cyclobutylidene-N-(3-fluorophenyl)-acetamide obtained in step 31.1 was dissolved in 19.5 ml of toluene and irradiated with a Hg middle pressure lamp in a suitable device till completion of the reaction indicated by liquid chromatography. The crude solution was concentrated in vacuo and the residue

was purified by column chromatography on silica (eluent: 10-30% ethyl acetate in heptane) to yield 55 mg (14%) of the title compound as a beige solid.

ESI-MS: m/z (%): 206 (100, [M+H]⁺).

31.3 Preparation of 5'-fluoro-2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinoline]

A solution of 50 mg (0.244 mmol) of 5'-fluoro-1'H-spiro[cyclobutane-1,4'-quinolin]-2'(3'H)-one obtained in step 31.2 in 1 mL of tetrahydrofuran was treated with 0.73 mL of 1 M solution of borohydride-tetrahydrofuran complex in tetrahydrofuran. The mixture was stirred at reflux for 2h, quenched with water and evaporated till dryness. The crude compound was used in the subsequent step without purification.

ESI-MS: m/z (%): 192 (100, [M+H]⁺).

31.4 Preparation of 2-(5'-fluoro-2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinolin]-1'-yl)-acetamide

A solution of 50 mg (0.26 mmol) of 5'-fluoro-2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinoline] obtained in step 31.3 in 1 ml of dimethylformamide was treated with 6.2 mg (0.26 mmol) of sodium hydride followed by 72 mg (0.52 mmol) 2-bromoacetamide and heated in a microwave system at 100°C for one hour. The resulting mixture was quenched with water, evaporated till dryness and directly purified by column chromatography on silica (eluent: 10% methanol in dichloromethane) to yield 70 mg of the title compound with about 80% purity determined by LCMS.

ESI-MS: m/z (%): 249 (100, [M+H]⁺).

31.5 Preparation of 2-(5'-fluoro-2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinolin]-1'-yl)ethanamine

A solution of 40 mg (0.16 mmol) of 2-(5'-fluoro-2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinolin]-1'-yl)-acetamide obtained in step 31.4 in 1 ml of tetrahydrofurane was treated with 38 µl (0.38 mmol) of borane dimethylsulfide and heated to 70°C for two hours. The mixture was quenched with water and hydrochloric acid, diluted with

dichloromethane and extracted twice with water. The combined aqueous phases were set to pH 10 with sodium hydroxide solution and extracted 3x with dichloromethane. The combined organic phases were dried and evaporated to yield 10 mg of the title compound as an oil.

ESI-MS: m/z (%): 235 (100, [M+H]⁺).

31.6 Preparation of 9-fluoro-1-methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane], trifluoroacetic acid salt

A solution of 10 mg (0.043 mmol) of 2-(5'-fluoro-2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinolin]-1'-yl)ethan-amine obtained in step 31.5 in 0.4 ml of acetonitrile/methanol was treated with 3.8 mg (0.085 mmol) of acetaldehyde and 6.5 μ l (0.085 mmol) of trifluoroacetic acid and stirred over night at room temperature. The solvent was evaporated and the crude mixture purified with reversed phase liquid chromatography to yield 1.5 mg of the title compound as beige solid.

ESI-MS: m/z (%): 261 (100, [M+H]⁺).

¹H-NMR (500 MHz, pyridine-d₅): δ = 7.05 (dd, 1H), 6.80 (dd, 1H), 4.20 (q, 1H), 3.05 (m, 2H), 3.00 (m, 2H), 2.85 (m, 2H), 2.75 (m, 2H), 2.05 (m, 2H), 1.90-1.70 (m, 4H), 1.50 (d, 3H) ppm.

Example 32

2,3,5,6,7,8-Hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline

(compound of the formula I.a, I.b or I.c in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-217 of Table A)

32.1 Preparation of tert-butyl 1-benzyl-2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate

To a solution of 388 mg (1.39 mmol) of 1-benzyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline (compound of example 30) in 15 ml of dichloromethane was treated with 0.49 ml (2.78 mmol) of N,N-diisopropylethylamine.

333 mg (1.53 mmol) of di-tert-butyl dicarbonate were dissolved in 5 ml of dichloromethane, slowly added to the reaction mixture and the solution was then stirred overnight at room temperature. The solution was diluted with dichloromethane and treated twice with 10% aqueous solution of citric acid. The organic phase was washed once with saturated sodium chloride solution, dried with MgSO_4 and concentrated in vacuo, yielding 559 mg of the crude product. The crude compound was used in the subsequent step without purification.

ESI-MS $[\text{M}+\text{H}^+] = 380.20$

32.2 Preparation of tert-butyl 2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate

559 mg (1.47 mmol) of tert-butyl 1-benzyl-2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate (559 mg, 1,473 μmol) (from step 32.1) were dissolved in 10 ml of methanol. The solution was hydrogenated with a 10% Pd/C cartridge using the H-Cube apparatus (from Thales Nano) at 80 bar and 70°C with a flow rate of 0.5 ml/min for 6 hours. The solution was concentrated in vacuo yielding 320 mg of the crude product. The crude compound was used in the subsequent step without purification.

ESI-MS $[\text{M}+\text{H}^+] = 290.20$

32.3 Preparation of 2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline

30 mg (0.104 mmol) of tert-butyl 2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate (from step 32.2) were dissolved in 2 ml of dichloromethane and treated with 0.20 ml (2.59 mmol) of trifluoroacetic acid at 0°C. The solution was allowed to warm up to room temperature and stirred for 2 h. The reaction mixture was diluted with dichloromethane and extracted once with a 1 molar aqueous solution of sodium hydroxide. The aqueous phase was extracted 3x with ethyl acetate. The combined organic phases were dried with magnesium sulfate, and concentrated in vacuo yielding 14 mg (0.076 mmol) of the title compound.

ESI-MS $[M+H]^+ = 190.10$

1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 6.75 (m, 1H), 6.55 (d, 1H), 6.50 (d, 1H), 3.90 (s, 2H), 3.25 (m, 4H), 3.10 (m, 2H), 2.95 (m, 2H).

Example 33

1-Cyclobutyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline 2,2,2-trifluoroacetate

(compound of the formula I.a, I.b or I.c in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} is as in row A-433 of Table A)

33.1 Preparation of tert-butyl 1-cyclobutyl-2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate

A solution of 79 mg (0.237 mmol) of tert-butyl 2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate (from step 32.2) in 2 ml THF was treated with 38.3 mg (0.546 mmol) of cyclobutanone and molecular sieves (4Å). The reaction mixture was stirred at room temperature overnight. 231 mg (1.09 mmol) of sodium triacetoxyborohydride was added and the reaction mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate and extracted once with a 1 molar aqueous solution of sodium hydroxide. The organic phase was dried with magnesium sulfate, concentrated in vacuo and the residue was purified by column chromatography on silica (eluent: 0-10% methanol in dichloromethane) yielding 78 mg (0.205 mmol) of the title compound.

ESI-MS $[M+H]^+ = 344.20$

33.2 Preparation of 1-cyclobutyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline 2,2,2-trifluoroacetate

78.4 mg (0.228 mmol) of tert-butyl 1-cyclobutyl-2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate (from step 33.1) were dissolved

in 1 ml of dichloromethane and treated with 0.53 ml (6.85 mmol) of trifluoroacetic acid at 0°C. The solution was stirred for 2 h and then concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-30% methanol in dichloromethane) to yield 69 mg of crude product, which was purified again via preparative HPLC to yield 46 mg of the title compound.

ESI-MS $[M+H]^+ = 244.20$

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 8.70 (bs, 2H), 6.80 (m, 1H), 6.75 (d, 1H), 6.70 (d, 1H), 4.10 (m, 2H), 4.05 (m, 1H), 3.25 (m, 2H), 3.20 (m, 2H), 3.10 (m, 4H), 2.20 (m, 2H), 2.10 (m, 2H), 1.65 (m, 2H).

Example 34

1-Methyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline

(compound of the formula I.a, I.b or I.c in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-253 of Table A)

The title compound was prepared in analogy to example 33, using however formaldehyde instead of cyclobutanone.

ESI-MS $[M+H]^+ = 204.10$

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 6.85 (m, 1H), 6.65 (d, 1H), 6.55 (d, 1H), 3.90 (s, 2H), 3.35 (m, 2H), 3.15 (m, 2H), 3.10 (m, 2H), 2.95 (m, 5H).

Example 35

1-(Oxetan-3-yl)-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline 2,2,2-trifluoroacetate

(compound of the formula I.a, I.b or I.c in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-541 of Table A)

The title compound was prepared in analogy to example 33, using however oxetan-3-one instead of cyclobutanone.

ESI-MS $[M+H^+] = 246.15$

^1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 8.80 (m, 2H), 6.75 (m, 2H), 6.30 (d, 1H), 4.80 (m, 2H), 4.65 (m, 3H), 4.10 (s, 2H), 3.35 (m, 2H), 3.20 (m, 4H), 3.10 (m, 2H).

Example 36

1-(Cyclopropylmethyl)-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline (compound of the formula I.a, I.b or I.c in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} is as in row A-577 of Table A)

The title compound was prepared in analogy to example 33, using however cyclopropanecarboxaldehyde instead of cyclobutanone.

ESI-MS $[M+H^+] = 244.20$

^1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 6.85 (m, 1H), 6.75 (m, 1H), 6.55 (m, 1H), 4.00 (m, 2H), 3.35-3.15 (m, 8H), 3.00 (m, 2H), 1.05 (m, 1H), 0.50 (m, 2H), 0.20 (m, 2H).

Example 37

1-(Cyclopentylmethyl)-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline (compound of the formula I.a, I.b or I.c in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} is as in row A-649 of Table A)

The title compound was prepared in analogy to example 33, using however cyclopentanecarboxaldehyde instead of cyclobutanone.

ESI-MS $[M+H^+] = 272.20$

^1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 6.80 (m, 1H), 6.65 (d, 1H), 6.50 (d, 1H), 3.95 (m, 2H), 3.30-3.20 (m, 6H), 3.15 (m, 2H), 2.95 (m, 2H), 2.30 (m, 1H), 1.90-1.55 (m, 6H), 1.25 (m, 2H).

Example 38

Cyclopropyl(2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxalin-1-yl)methanone

(compound of the formula I.a, I.b or I.c in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-721 of Table A)

38.1 Preparation of tert-butyl 1-(cyclopropanecarbonyl)-2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate

To 50 mg (0.173 mmol) of tert-butyl 2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate dissolved in 1 ml of dichloromethane were added 0.77 ml (4.43 mmol) of N,N-diisopropylethylamine in an inert atmosphere at 0°C. 19 µl (0.21 mmol) of cyclopropanecarbonyl chloride was dissolved in 1 ml of dichloromethane and slowly added to the reaction mixture. The reaction mixture was stirred at room temperature for 5 h. The solution was diluted with dichloromethane and treated twice with 10% aqueous solution of citric acid. The organic phase was washed once with saturated sodium chloride solution, dried with MgSO₄ and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-10% methanol in dichloromethane) yielding 35 mg (0.098 mmol) of the title compound.

ESI-MS [M-tert-butyl+H⁺] = 302.10

38.2 Preparation of cyclopropyl(2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxalin-1-yl)methanone

35 mg (0.098 mmol) of tert-butyl 1-(cyclopropanecarbonyl)-2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate were dissolved in 2 ml of dichloromethane and treated with 150 µl (1.96 mmol) of trifluoroacetic acid at 0°C. The solution was stirred overnight at room temperature. 100 µl (1.31 mmol) of trifluoroacetic acid were added and the reaction mixture was stirred for 1 h. The reaction mixture was treated 3x with acidified water. The combined aqueous phase was basified with 1 N sodium hydroxide solution and extracted 5x with dichloromethane.

The combined organic phase was dried with MgSO_4 and concentrated in vacuo to yield 23 mg of the title compound.

ESI-MS $[\text{M}+\text{H}^+] = 258.20$

^1H NMR ($\text{DMSO}-d_6$, 500MHz): δ [ppm]: 7.30 (m, 1H), 7.05 (m, 1H), 6.80 (m, 1H), 4.10 (s, 2H), 3.90 (m, 2H), 3.45 (m, 4H), 3.30 (m, 2H), 2.10 (m, 1H), 1.15 (m, 2H), 0.85 (m, 2H).

Example 39

Cyclopentyl(2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxalin-1-yl)methanone

(compound of the formula I.a, I.b or I.c in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} is as in row A-793 of Table A)

The title compound was prepared in analogy to example 38 using cyclopentanecarbonyl chloride.

ESI-MS $[\text{M}+\text{H}^+] = 286.20$

^1H NMR ($\text{DMSO}-d_6$, 500MHz): δ [ppm]: 7.00 (m, 2H), 6.80 (m, 1H), 4.10 (s, 2H), 3.90 (m, 2H), 3.45-3.20 (m, 7H), 1.80 (m, 6H), 1.55 (m, 2H).

Example 40

1-Cyclopropyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline

(compound of the formula I.a, I.b or I.c in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} is as in row A-397 of Table A)

40.1 Preparation of 2-(4-cyclopropyl-3,4-dihydroquinoxalin-1(2H)-yl)acetamide

A solution of 500 mg (2.87 mmol) of 1-cyclopropyl-1,2,3,4-tetrahydroquinoxaline in 5 ml dimethylformamide was treated with 2.0 ml (11.68 mmol) of N-ethyl-N-isopropylpropan-2-amine, and subsequently 555 mg (5.94 mmol) 2-chloroacetamide were added. The reaction mixture was heated in a microwave system at 100°C for 12

hours. The reaction mixture was quenched with water and basified with 1 N sodium hydroxide solution. The aqueous phase was extracted 3x with dichloromethane. The combined organic extracts were dried with magnesium sulfate and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-20% methanol in dichloromethane) to yield 620 mg of the title compound.

ESI-MS $[M+H^+] = 232.20$

40.2 Preparation of 2-(4-cyclopropyl-3,4-dihydroquinoxalin-1(2H)-yl)ethanamine

To 620 mg (2.68 mmol) of 2-(4-cyclopropyl-3,4-dihydroquinoxalin-1(2H)-yl)acetamide in 2 ml of THF was added 5.4 ml (10.80 mmol) of 2 molar borane dimethylsulfide THF solution and subsequently heated for 7 hours at 60°C in a microwave system. An additional 2.7 ml (5.40 mmol) of 2 molar borane dimethylsulfide THF solution was added to the reaction mixture and heated for 10 hours at 60°C in a microwave system. To the reaction mixture were added 2 ml of methanol, acidified with 2 molar hydrochloric acid and heated for 15 minutes at 60°C in a microwave system. The reaction mixture was diluted with ethyl acetate and extracted twice with 1 molar hydrochloric acid. The combined aqueous phases were set to pH 10 with sodium hydroxide solution and extracted 3x with dichloromethane. The combined organic phases were dried and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-15% methanol in dichloromethane) to yield 347 mg of the title compound.

ESI-MS $[M+H^+] = 218.20$

40.3 Preparation of 1-Cyclopropyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]-quinoxaline

A solution of 347 mg (1.6 mmol) of 2-(4-cyclopropyl-3,4-dihydroquinoxalin-1(2H)-yl)ethanamine in 5 ml of ethanol were treated with 119 μ l (1.6 mmol) of 37% aqueous formaldehyde solution and 135 μ l (1.76 mmol) of trifluoroacetic acid. The reaction mixture was stirred for 36 h at room temperature. Water was added and the reaction mixture

was basified using 2 N aqueous sodium hydroxide solution. The aqueous phase was extracted three times with dichloromethane. The combined organic phases were dried and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-20% methanol in dichloromethane) to yield 177 mg of the title compound.

ESI-MS $[M+H]^+ = 230.20$

1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 7.10 (d, 1H), 6.80 (m, 1H), 6.60 (d, 1H), 3.85 (s, 2H), 3.20 (m, 2H), 3.10 (m, 2H), 3.00 (m, 4H), 2.25 (m, 1H), 0.80 (m, 2H), 0.55 (m, 2H).

Example 41

1-Cyclopentyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline
(compound of the formula I.a, I.b or I.c in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} is as in row A-469 of Table A)

The title compound was prepared in analogy to example 40 using 1-cyclopentyl-1,2,3,4-tetrahydroquinoxaline as starting material.

ESI-MS $[M+H]^+ = 258.20$

1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 6.80 (d, 1H), 6.70 (m, 1H), 6.45 (d, 1H), 4.25 (m, 1H), 3.75 (s, 2H), 3.15 (m, 2H), 3.05 (m, 2H), 2.90 (m, 2H), 2.85 (m, 2H), 1.80 (m, 2H), 1.65 (m, 2H), 1.55 (m, 4H).

Example 42

1-Cyclopropyl-5-methyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline
(compound of the formula I.gg, I.hh or I.ii in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} is as in row A-397 of Table A)

42.1 Preparation of 2-(4-cyclopropyl-3,4-dihydroquinoxalin-1(2H)-yl)propanamide

A solution of 300 mg (1.72 mmol) of 1-cyclopropyl-1,2,3,4-tetrahydroquinoxaline in 3 ml dimethylformamide was treated with 1.2 ml (7.0 mmol) of N-ethyl-N-isopropylpropan-2-amine, 387 mg (2.58 mmol) of sodium iodide and subsequently 523 mg (3.44 mmol) of 2-bromopropanamide were added. The reaction mixture was heated in a microwave system at 100°C for 8 hours. The reaction mixture was quenched with water and basified with 1 N sodium hydroxide solution. The aqueous phase was extracted 3x with dichloromethane. The combined organic extracts were dried with magnesium sulfate and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-20% methanol in dichloromethane) to yield 453 mg of the title compound.

ESI-MS $[M+H^+] = 246.20$

42.2 Preparation of 2-(4-cyclopropyl-3,4-dihydroquinoxalin-1(2H)-yl)propan-1-amine

To 453 mg (1.85 mmol) of 2-(4-cyclopropyl-3,4-dihydroquinoxalin-1(2H)-yl)propanamide in 2 ml THF were added 4.3 ml (8.60 mmol) of 2 molar borane dimethylsulfide THF solution and subsequently heated for 9 hours at 60°C and 2 hours at 70°C in a microwave system. To the reaction mixture was added 2 ml of methanol, acidified with 2 molar hydrochloric acid and heated for 15 minutes at 60°C in a microwave system. The reaction mixture was diluted with ethyl acetate and extracted twice with 1 molar hydrochloric acid. The combined aqueous phases were set to pH 10 with sodium hydroxide solution and extracted 3x with dichloromethane. The combined organic phases were dried and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-15% methanol in dichloromethane) to yield 209 mg of the title compound.

ESI-MS $[M+H^+] = 232.20$

42.3 Preparation of 1-cyclopropyl-5-methyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino-[1,7,6-de]quinoxaline

A solution of 209 mg (0.9 mmol) of 2-(4-cyclopropyl-3,4-dihydroquinoxalin-1(2H)-yl)propan-1-amine in 3 ml of ethanol were treated with 67 μ l (0.9 mmol) of 37% aqueous formaldehyde solution and 76 μ l (0.99 mmol) of trifluoroacetic acid. The reaction mixture was stirred for 36 h at room temperature. Water was added and the reaction mixture was basified using 2 N aqueous sodium hydroxide solution. The aqueous phase was extracted three times with dichloromethane. The combined organic phases were dried and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-20% methanol in dichloromethane) to yield 136 mg of the title compound.

ESI-MS $[M+H]^+ = 244.20$

^1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 7.00 (d, 1H), 6.65 (m, 1H), 6.50 (d, 1H), 3.75 (d, 1H), 3.45 (d, 1H), 3.30-3.15 (m, 4H), 3.05 (m, 1H), 2.80 (d, 1H), 2.65 (d, 1H), 2.20 (m, 1H), 0.90 (d, 3H), 0.80 (m, 2H), 0.50 (m, 2H).

Example 43

1-Cyclopentyl-5-methyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline (compound of the formula I.gg, I.hh or I.ii in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} is as in row A-469 of Table A)

The title compound was prepared in analogy to example using however 1-cyclopentyl-1,2,3,4-tetrahydroquinoxaline instead of 1-cyclopropyl-1,2,3,4-tetrahydroquinoxaline.

ESI-MS $[M+H]^+ = 272.20$

^1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 6.70 (d, 1H), 6.65 (m, 1H), 6.40 (d, 1H), 4.20 (m, 1H), 3.75 (d, 1H), 3.50 (d, 1H), 3.25-3.10 (m, 3H), 3.00 (m, 2H), 2.85 (d, 1H), 2.70 (d, 1H), 1.80 (m, 2H), 1.70-1.55 (m, 5H), 1.50 (m, 1H), 0.90 (d, 3H).

Example 44

5,6,7,9,9a,10,11,12,13,14-decahydro-4H-azepino[1,2-a][1,4]diazepino[1,7,6-de]quinoxaline

(compound of the formula I.a, I.b or I.c in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-937 of Table A)

44.1 Preparation of ethyl 1-(2-nitrophenyl)azepane-2-carboxylate

A solution of 450 mg (3.19 mmol) of 1-fluoro-2-nitrobenzene, 546 mg (3.19 mmol) of ethyl azepane-2-carboxylate and 0.89 ml (6.38 mmol) of triethylamine in 10 ml acetonitrile was stirred overnight at reflux. The reaction mixture was concentrated in vacuo. To the residue was added aqueous sodium bicarbonate solution and extracted twice with ethyl acetate. The combined organic extracts were dried with sodium sulfate and concentrated in vacuo to yield 780 mg of the crude title compound. The crude product was used without further purification in the next step.

ESI-MS [M+H⁺] = 293.10

44.2 Preparation of 6a,7,8,9,10,11-hexahydroazepino[1,2-a]quinoxalin-6(5H)-one

780 mg (2.67 mmol) of ethyl 1-(2-nitrophenyl)azepane-2-carboxylate were dissolved in 30 ml methanol and treated with 284 mg of 10% Pd/C in a hydrogen atmosphere. The solution was stirred at room temperature for 1 h. The catalyst was filtered off and the solution was concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-50% ethyl acetate in heptane) to yield 159 mg of the title compound.

ESI-MS [M+H⁺] = 217.10

44.3 Preparation of 5,6,6a,7,8,9,10,11-octahydroazepino[1,2-a]quinoxaline

To 105 mg (0.485 mmol) 6a,7,8,9,10,11-hexahydroazepino[1,2-a]quinoxalin-6(5H)-one in 4 ml of THF (tetrahydrofuran) were added 0.97 ml (0.97 mmol) of 1 molar borane dimethylsulfide THF solution and subsequently heated for 1 h at 80°C in a microwave system. The reaction mixture was poured on ice water, acidified with hydrochloric acid

solution to pH = 5 and extracted twice with ethyl acetate. The combined organic phases were dried and concentrated in vacuo. The residue was acidified with hydrochloric acid solution and extracted once with ether. The aqueous phase was basified with aqueous sodium bicarbonate solution and extracted 3x with dichloromethane. The combined organic phases were dried and concentrated in vacuo to yield 85 mg of the crude title compound.

ESI-MS $[M+H^+] = 203.20$

44.4 Preparation of 2-(6a,7,8,9,10,11-hexahydroazepino[1,2-a]quinoxalin-5(6H)-yl)-acetamide

A solution of 85 mg (0.40 mmol) of 5,6,6a,7,8,9,10,11-octahydroazepino[1,2-a]quinoxaline in 2 ml of dimethylformamide was treated with 16 mg (0.40 mmol) of sodium hydride (60% in mineral oil) and stirred for 15 min at room temperature. 112 mg (1.2 mmol) of 2-chloroacetamide were added and the reaction mixture was heated in a microwave system at 100°C for 4 hours. 209 μ l (1.2 mmol) of N-ethyl-N-isopropylpropan-2-amine were added to the reaction mixture and heated at 100°C for 2 hours. 110 mg (0.8 mmol) of 2-bromoacetamide were added and the reaction mixture was heated in a microwave system at 100°C for 2 hours. The reaction mixture was basified with sodium bicarbonate solution and extracted 2x with ethyl acetate. The combined organic extracts were dried with magnesium sulfate and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: : 0-50% ethyl acetate in heptane) to yield 42 mg of the title compound.

ESI-MS $[M+H^+] = 260.20$

44.5 Preparation of 2-(6a,7,8,9,10,11-hexahydroazepino[1,2-a]quinoxalin-5(6H)-yl)-ethanamine

To 42 mg (0.146 mmol) of 2-(6a,7,8,9,10,11-hexahydroazepino[1,2-a]quinoxalin-5(6H)-yl)acetamide in 1 ml of THF were added 292 μ l (8.60 mmol) of 2 molar borane

dimethylsulfide THF solution and subsequently heated for 6 hours at 60°C in a microwave system. Additional 292 μ l (8.60 mmol) of 2 molar borane dimethylsulfide THF solution were added to the reaction mixture and heated for 4 hours at 60°C in a microwave system. To the reaction mixture was added 2 ml of methanol, acidified with 2 molar aqueous hydrochloric acid solution and heated for 15 minutes at 60°C in a microwave system. The reaction mixture was diluted with ethyl acetate and extracted twice with 1 molar hydrochloric acid. The combined aqueous phases were set to pH 10 with sodium hydroxide solution and extracted 3x with dichloromethane. The combined organic phases were dried and concentrated in vacuo to yield 36 mg of the title compound.

ESI-MS $[M+H]^+ = 246.20$

44.6 Preparation of 5,6,7,9,9a,10,11,12,13,14-decahydro-4H-azepino[1,2-a][1,4]-diazepino[1,7,6-de]quinoxaline

A solution of 36 mg (0.147 mmol) of 2-(6a,7,8,9,10,11-hexahydroazepino[1,2-a]quinoxalin-5(6H)-yl)ethanamine in 2 ml of ethanol were treated with 11 μ l (0.147 mmol) of 37% aqueous formaldehyde solution and 12 μ l (0.161 mmol) of trifluoroacetic acid. The reaction mixture was stirred overnight at room temperature. The reaction mixture was concentrated in vacuo, treated with aqueous sodium bicarbonate solution and extracted twice with dichloromethane. The combined organic phases were dried and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: : 0-20% methanol in dichloromethane) to yield 12.7 mg of the title compound.

ESI-MS $[M+H]^+ = 258.20$

^1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 6.65 (m, 1H), 6.50 (d, 1H), 6.35 (d, 1H), 3.70 (m, 1H), 3.60 (m, 1H), 3.20 (m, 2H), 3.10 (m, 1H), 3.00 (m, 3H), 2.95 (m, 1H), 2.85 (m, 1H), 2.80 (m, 1H), 1.80 (m, 2H), 1.65-1.50 (m, 4H), 1.40 (m, 2H).

Example 45

7-Methyl-5,6,7,9,9^a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

(compound of the formula I.gg, I.hh or I.ii in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-865 of Table A)

The title compound was prepared in analogy to example 42, using however 1,2,3,3a,4,5-hexahydropyrrolo[1,2-a]quinoxaline and 2-bromopropanamide.

ESI-MS [M+H⁺] = 244.20

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 6.80 (m, 1H), 6.50 (d, 1H), 6.45 (d, 1H), 3.95 (d, 1H), 3.70 (d, 1H), 3.50 (m, 2H), 3.15 (m, 2H), 2.85 (m, 2H), 2.75 (m, 1H), 2.20 (m, 1H), 2.10 (m, 1H), 2.00 (m, 1H), 1.90 (m, 1H), 1.35 (m, 1H), 1.20 (d, 3H).

Example 46

1-Fluoro-7-methyl-5,6,7,9,9^a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

(compound of the formula I.gg in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-866 of Table A)

The title compound was prepared in analogy to example 42, using however 9-fluoro-1,2,3,3a,4,5-hexahydropyrrolo[1,2-a]quinoxaline and 2-bromopropanamide.

ESI-MS [M+H⁺] = 262.20

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 6.60 (m, 1H), 6.55 (m, 1H), 3.90 (d, 1H), 3.80 (m, 1H), 3.55 (d, 1H), 3.20 (d, 1H), 3.00 (m, 2H), 2.80 (m, 2H), 2.55 (m, 1H), 2.15 (m, 2H), 1.80 (m, 2H), 1.50 (m, 1H), 1.25 (d, 3H).

Example 47

1,2,3,4,6,7-Hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopropane]

(compound of the formula I.g, I.h or I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-281 of Table B)

47.1 Preparation of 2-(2',3'-dihydro-1'H-spiro[cyclopropane-1,4'-quinolin]-1'-yl)-acetamide

A solution of 500 mg (3.14 mmol) of 2',3'-dihydro-1'H-spiro[cyclopropane-1,4'-quinoline] in 5 ml of dimethylformamide was treated with 2.2 ml (12.8 mmol) of N-ethyl-N-isopropylpropan-2-amine and 706 mg (4.71 mmol) of sodium iodide; and subsequently 670 mg (7.16 mmol) of 2-chloroacetamide were added. The reaction mixture was heated in a microwave system at 100°C for 10 hours. The reaction mixture was quenched with water and basified with 1 N sodium hydroxide solution. The aqueous phase was extracted 3x with dichloromethane. The combined organic extracts were dried with magnesium sulfate and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-20% methanol in dichloromethane) to yield 624 mg of the title compound.

ESI-MS $[M+H^+] = 217.20$

47.2 Preparation of 2-(2',3'-dihydro-1'H-spiro[cyclopropane-1,4'-quinolin]-1'-yl)ethan-amine

To 624 mg (2.89 mmol) of 2-(2',3'-dihydro-1'H-spiro[cyclopropane-1,4'-quinolin]-1'-yl)acetamide in 2 ml of THF were added 5.77 ml (11.54 mmol) of 2 molar borane dimethylsulfide THF solution and subsequently heated for 9 hours at 60°C in a microwave system. To the reaction mixture were added 2 ml of methanol, acidified with 2 molar hydrochloric acid and heated for 15 minutes at 60°C in a microwave system. The reaction mixture was diluted with ethyl acetate and extracted twice with 1 molar hydrochloric acid. The combined aqueous phases were set to pH 10 with sodium hydroxide solution and extracted 3x with dichloromethane. The combined organic phases were dried and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: : 0-20% methanol in dichloromethane) to yield 299 mg of the title compound.

ESI-MS $[M+H^+] = 203.20$

47.3 Preparation of 1,2,3,4,6,7-Hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopropane]

A solution of 299.6 mg (1.48 mmol) of 2-(2',3'-dihydro-1'H-spiro[cyclopropane-1,4'-quinolin]-1'-yl)ethanamine in 3 ml of ethanol was treated with 110 μ l (1.48 mmol) of 37% aqueous formaldehyde solution and 126 μ l (1.63 mmol) of trifluoroacetic acid. The reaction mixture was stirred overnight at room temperature. Water was added and the reaction mixture was basified using 2 N aqueous sodium hydroxide solution. The aqueous phase was extracted three times with dichloromethane. The combined organic phases were dried and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-20% methanol in dichloromethane) to yield 131 mg of the title compound.

ESI-MS $[M+H]^+ = 215.20$

1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 6.90 (d, 1H), 6.70 (m, 1H), 6.55 (d, 1H), 3.75 (s, 2H), 3.25 (m, 2H), 3.00 (m, 2H), 2.90 (m, 2H), 1.60 (m, 2H), 0.95 (s, 2H), 0.80 (s, 2H).

Example 48

4-Methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopropane]

(compound of the formula I.s, I.t or I.u in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-281 of Table B)

The title compound was prepared in analogy to example 47, using however 2',3'-dihydro-1'H-spiro[cyclopropane-1,4'-quinoline] and 2-bromopropanamide.

ESI-MS $[M+H]^+ = 229.20$

1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 6.85 (d, 1H), 6.65 (m, 1H), 6.55 (d, 1H), 3.80 (d, 1H), 3.50 (d, 1H), 3.25 (m, 3H), 2.85 (d, 1H), 2.70 (d, 1H), 1.70 (m, 1H), 1.60 (m, 1H), 1.05 (m, 1H), 0.90 (m, 3H), 0.85 (m, 1H), 0.75 (m, 2H).

Example 49

11-Fluoro-8,8-Dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline
(compound of the formula I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as
in row B-2 of Table B)

The title compound was prepared in analogy to example 47, using however 7-fluoro-
4,4-dimethyl-1,2,3,4-tetrahydroquinoline and 2-bromoacetamide.

ESI-MS [M+H⁺] = 235.20

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 7.15 (dd, 1H), 6.55 (dd, 1H), 3.80 (s, 2H),
3.20 (m, 2H), 3.10 (m, 2H), 2.90 (m, 2H), 1.60 (m, 2H), 1.20 (s, 6H).

Example 50

10-Methoxy-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline,
trifluoroacetic acid
(compound of the formula I.h in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as
in row B-7 of Table B)

The title compound was prepared in analogy to example 1, using however 7-methoxy-
2,3-dihydro-1H-benzo[c][1,4]diazepine-4(5H)-carboxylate and 3-methylbut-2-enoyl
chloride.

ESI-MS [M+H⁺] = 247.20

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 8.75 (bs, 2H), 6.95 (s, 1H), 6.85 (s, 1H), 4.15
(m, 2H), 3.70 (s, 3H), 3.20 (m, 2H), 3.15 (m, 4H), 1.65 (m, 2H), 1.25 (s, 6H).

Example 51

4,8,8-Trimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline 2,2,2-
trifluoroacetate
(compound of the formula I.s, I.t or I.u in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and
R^{9a} is as in row B-1 of Table B)

The title compound was prepared in analogy to example 47, using however 4,4-dimethyl-1,2,3,4-tetrahydroquinoline and 2-bromopropanamide.

ESI-MS $[M+H]^+ = 231.20$

1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 9.90 (bs, 1H), 9.20 (bs, 1H); 7.35 (d, 1H), 7.10 (d, 1H), 6.90 (m, 1H), 4.30 (d, 1H), 4.05 (d, 1H), 3.45 (m, 1H), 3.25 (m, 3H), 3.15 (m, 1H), 1.70 (m, 2H), 1.30 (m, 9H).

Example 52

4-Ethyl-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline
(compound of the formula I.v, I.w or I.x in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-1 of Table B)

The title compound was prepared in analogy to example 47, using however 4,4-dimethyl-1,2,3,4-tetrahydroquinoline and 2-bromobutanamide.

ESI-MS $[M+H]^+ = 245.20$

1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 7.15 (d, 1H), 6.85 (d, 1H), 6.70 (m, 1H), 3.85 (d, 1H), 3.50 (d, 1H), 3.20 (m, 2H), 3.00 (d, 1H), 2.95 (m, 1H), 2.75 (d, 1H), 1.60 (m, 2H), 1.40 (m, 2H), 1.25 (s, 3H), 1.30 (s, 3H), 0.80 (m, 3H).

Example 53

6,8,8-Trimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline
(compound of the formula I.y, I.z or I.zz in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-1 of Table B)

The title compound was prepared in analogy to example 47, using however 2,4,4-trimethyl-1,2,3,4-tetrahydroquinoline and 2-chloroacetamide.

ESI-MS $[M+H]^+ = 231.20$

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 7.05 (d, 1H), 6.90 (d, 1H), 6.65 (m, 1H), 3.85 (d, 1H), 3.45 (d, 1H), 3.30 (m, 3H), 3.00 (m, 1H), 2.70 (m, 1H), 1.65 (m, 1H), 1.40 (m, 1H), 1.25 (s, 3H), 1.15 (s, 6H).

Example 54

One enantiomer of 8-ethyl-8-methyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]-quinoline, trifluoroacetic acid

(enantiomer of the compound of the formula I.g, I.h or I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-41 of Table B)

The title compound was prepared in analogy to example 1, using however tert-butyl 2,3-dihydro-1H-benzo[e][1,4]diazepine-4(5H)-carboxylate and (Z)-3-methylpent-2-enoyl chloride.

ESI-MS [M+H⁺] = 231.20

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 8.90 (bs, 1H), 8.65 (bs, 1H), 7.30 (d, 1H), 7.15 (d, 1H), 6.90 (m, 1H), 4.20 (m, 1H), 4.10 (m, 1H), 3.25 (m, 6H), 1.85 (m, 1H), 1.70 (m, 1H), 1.60 (m, 1H), 1.45 (m, 1H), 1.20 (s, 3H), 0.70 (m, 3H).

Example 55

Other enantiomer of 8-ethyl-8-methyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline, trifluoroacetic acid

(enantiomer of the compound of the formula I.g, I.h or I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-41 of Table B)

The title compound was prepared in analogy to example 1, using however tert-butyl 2,3-dihydro-1H-benzo[e][1,4]diazepine-4(5H)-carboxylate and (Z)-3-methylpent-2-enoyl chloride.

ESI-MS [M+H⁺] = 231.10

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 8.90 (bs, 1H), 8.65 (bs, 1H), 7.30 (d, 1H), 7.15 (d, 1H), 6.90 (m, 1H), 4.20 (m, 1H), 4.10 (m, 1H), 3.20 (m, 6H), 1.85 (m, 1H), 1.70 (m, 1H), 1.60 (m, 1H), 1.45 (m, 1H), 1.20 (s, 3H), 0.70 (m, 3H).

Example 56

4-Methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane]

(compound of the formula I.s, I.t or I.u in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-321 of Table B)

56.1 Preparation of 2-cyclobutylidene-N-phenylacetamide

3.35 g (36 mmol) of aniline were dissolved in 60 ml of dichloromethane and treated with 13.83 ml (79 mmol) of N-ethyl-N-isopropylpropan-2-amine. Within 10 min 19.80 ml of a 1 molar solution of 2-cyclobutylideneacetyl chloride in dichloromethane was added at 0°C and the solution was stirred overnight at room temperature. The reaction mixture was poured on ice water and extracted 2x with dichloromethane. The combined organic phases were washed once with saturated sodium chloride solution, dried with magnesium sulfate and concentrated in vacuo. The crude oil was dissolved in dichloromethane and treated with ether till crystallization, yielding 5.5 g of the title compound.

ESI-MS [M+H⁺] = 188.10

56.2 Preparation of 1'H-spiro[cyclobutane-1,4'-quinolin]-2'(3'H)-one

A solution of 2.0 g (10.68 mmol) of 2-cyclobutylidene-N-phenylacetamide in 214 ml of toluene was irradiated with a 700 W mercury lamp in a flow reactor (FEP-tubing, pyrex filter) with a flow rate of 1 ml per min at a temperature range of 0°C to 20°C with a residence time of 5 to 7 min. The reaction mixture was concentrated in vacuo and the residue was purified by column chromatography on silica (eluent: 0-55% ethyl acetate in cyclohexane) to yield 940 mg of the title compound.

ESI-MS $[M+H^+] = 188.10$

56.3 Preparation of 2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinoline]

To 850 mg (4.54 mmol) of 1'H-spiro[cyclobutane-1,4'-quinolin]-2'(3'H)-one in 5 ml of THF were added 9.08 ml (9.08 mmol) of 1 molar borane dimethylsulfide THF solution and subsequently heated for 30 min at 80°C in a microwave system. The reaction mixture was acidified with 6 molar hydrochloric acid solution and treated with ethyl acetate. The organic phase was extracted twice with 1 molar hydrochloric acid solution. The combined aqueous phases were set to pH 10 with sodium bicarbonate and extracted 3x with ethyl acetate. The combined organic phases were dried and concentrated in vacuo to yield 565 mg of the crude title compound.

ESI-MS $[M+H^+] = 174.10$

56.4 Preparation of 2-(2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinolin]-1'-yl)propanamide

ESI-MS $[M+H^+] = 245.20$

A solution of 300 mg (1.73 mmol) of 2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinoline] in 10 ml dimethylformamide was treated with 69 mg (1.73 mmol) of sodium hydride (60% in mineral oil) and stirred for 15 min at 50°C, then transferred to a microwave vial and stirred for 5 min at 70°C in a microwave system. Subsequently 790 mg (5.19 mmol) of 2-bromoacetamide were added and the reaction mixture was heated in a microwave system at 100°C for 7 hours. Additional 263 mg (1.73 mmol) of 2-bromoacetamide were added and the reaction mixture was heated in a microwave system at 100°C for 2 hours. The reaction mixture was quenched with ice water, acidified to pH = 5 and extracted with ether. The aqueous phase was basified with sodium bicarbonate solution and extracted 3x with dichloromethane. The combined organic extracts were dried with magnesium sulfate and concentrated in vacuo. The residue was purified

by column chromatography on silica (eluent: 0-55% ethyl acetate in heptane) to yield 258 mg of the title compound.

56.5 Preparation of 2-(2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinolin]-1'-yl)propan-1-amine

To 258 mg (0.348 mmol) of 2-(2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinolin]-1'-yl)propanamide in 2 ml THF were added 1.05 ml (2.10 mmol) of 2 molar borane dimethylsulfide THF solution and subsequently heated for 6 hours at 60°C in a microwave system. Subsequently 348 µl (0.70 mmol) of 2 molar borane dimethylsulfide THF solution was added to the reaction mixture and heated for 2 hours at 60°C in a microwave system. Subsequently 697 µl (1.39 mmol) of 2 molar borane dimethylsulfide THF solution was added to the reaction mixture and heated for 3 hours at 60°C in a microwave system. To the reaction mixture was added 2 ml of methanol, acidified with 2 molar hydrochloric acid and heated for 5 minutes at 60°C in a microwave system. The reaction mixture was diluted with ethyl acetate and extracted twice with 1 molar hydrochloric acid. The combined aqueous phases were set to pH 10 with sodium bicarbonate solution and extracted 3x with dichloromethane. The combined organic phases were dried and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-20% methanol in dichloromethane) to yield 90 mg of the title compound.

ESI-MS $[M+H]^+ = 231.20$

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 7.35 (d, 1H), 6.95 (dd, 1H), 6.70 (d, 1H), 6.55 (dd, 1H), 3.80 (m, 1H), 3.05 (m, 1H), 3.00 (m, 1H), 2.70 (m, 1H), 2.55 (m, 1H), 2.35 (m, 1H), 2.25 (m, 1H), 2.00 (m, 1H), 1.90 (m, 5H), 1.05 (d, 3H).

56.6 Preparation of 4-methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]-quinoline-8,1'-cyclobutane]

A solution of 90 mg (0.313 mmol) of 2-(2',3'-dihydro-1'H-spiro[cyclobutane-1,4'-quinolin]-1'-yl)propan-1-amine in 2 ml of ethanol was treated with 23 µl (0.313 mmol) of 37% aqueous formaldehyde solution and 26 µl (0.344 mmol) of trifluoroacetic acid.

The reaction mixture was stirred overnight at room temperature. Further 12 μ l (0.156 mmol) of 37% aqueous formaldehyde solution and 12 μ l (0.156 mmol) of trifluoroacetic acid were added to the reaction mixture and stirring was continued overnight at room temperature. The reaction mixture was concentrated in vacuo and to the residue was added aqueous sodium bicarbonate solution. The aqueous phase was extracted twice with dichloromethane. The combined organic phases were dried and concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-10% methanol in dichloromethane) to yield 38 mg of the title compound.

ESI-MS $[M+H^+] = 243.20$

1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 7.35 (d, 1H), 6.90 (d, 1H), 6.75 (d, 1H), 3.80 (m, 1H), 3.40 (m, 1H), 3.20 (m, 2H), 3.10 (m, 1H), 2.80 (m, 1H), 2.70 (m, 1H), 2.40 (m, 1H), 2.15 (m, 1H), 2.00 (m, 4H), 1.85 (m, 1H), 1.70 (m, 1H), 0.80 (s, 3H).

Example 57

1-Fluoro-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline, trifluoroacetic acid

(compound of the formula I.a in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} is as in row A-866 of Table A)

57.1 Preparation of methyl pyrrolidine-2-carboxylate

To a solution of pyrrolidine-2-carboxylic acid (75 g, 651 mmol) in methanol (750 ml), $SOCl_2$ (250 ml) was added dropwise at 0 °C, and the reaction mixture was stirred for 16 hrs at 20 °C. One additional vial was set up in the same way. Both reaction mixtures were combined and concentrated under reduced pressure to give the title compound (90 g, 543 mmol, yield 50%).

1H NMR (400 MHz, DMSO- d_6): δ [ppm]: 9.83 (s, 1H), 4.32 (t, $J = 7.6$ Hz, 1H), 3.72 (s, 3H), 3.21-3.15 (m, 2H), 2.22-2.20 (m, 1H), 1.96-1.87 (m, 3H)

57.2 Preparation of methyl 1-(2-fluoro-6-nitro-phenyl)-pyrrolidine-2-carboxylate

To a mixture of methyl pyrrolidine-2-carboxylate (17.18 g, 104 mmol) in acetonitril (500 mL), triethylamine (26.3 mL, 189 mmol) and 1,2-difluoro-3-nitrobenzene (15 g, 94 mmol) were added at 20 °C, and the reaction mixture was stirred for 12 h at 80°C. Then the mixture was cooled to 20°C, concentrated under reduced pressure, and the residue was purified by column chromatography on silica gel (petrol ether:ethyl acetate = 20:1 to 10:1) to give the title compound (20 g, 74.6 mmol, yield 79%).

¹H NMR (400MHz, CDCl₃): δ [ppm]: 7.36 (d, *J* = 8 Hz, 1H), 7.15-7.11 (m, 1H), 6.99-6.96 (m, 1H), 4.35-4.31 (m, 1H), 3.55 (s, 3H), 3.52-3.49 (m, 1H), 3.06-3.04 (m, 1H), 2.29-2.26 (m, 1H), 2.06-1.89 (m, 3H)

57.3 Preparation of 9-fluoro-2,3,3a,5,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-4-one

To a solution of methyl 1-(2-fluoro-6-nitro-phenyl)-pyrrolidine-2-carboxylate (10 g, 37.3 mmol) in methanol (200 ml), Pd/C (3.97 g, 37.3 mmol) was added under Ar protection, then the reaction mixture was stirred for 12 h at 20°C at 15 psi under H₂ atmosphere. One additional vial was set up as described above. The two reaction mixtures were combined, filtered, and the filtrate was concentrated under reduced pressure to give the title compound (10 g, 48.5 mmol, yield 65%).

¹H NMR (400MHz, CDCl₃): δ [ppm]: 9.33 (s, 1H), 6.73-6.63 (m, 2H), 6.55-6.53 (m, 1H), 3.89-3.86 (m, 1H), 3.72-3.68 (m, 1H), 3.40-3.39 (m, 1H), 2.30-2.22 (m, 2H), 1.92-1.84 (m, 2H)

57.4 Preparation of 9-fluoro-1,2,3,3a,4,5,5a,6-octahydropyrrolo[1,2-a]quinoxaline

To a solution of 9-fluoro-2,3,3a,5,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-4-one (10 g, 48.5 mmol) in THF (200 ml), a solution of BH₃.THF (12.5 g, 145 mmol) was added at 0 °C, then the reaction solution was stirred 12 h at 70°C. After cooling to 0°C, methanol (100 ml) was added dropwise at 0°C, and the reaction solution was stirred for

30 minutes at 20°C, concentrated under reduced pressure, and the residue was purified by column chromatography on silica gel (petrol ether:ethyl acetate = 20 :1 to 10:1) to give the title compound (5 g, 26 mmol, yield 53.6%).

¹H NMR (400MHz, CDCl₃): δ [ppm]: 6.54-6.49 (m, 1H), 6.40-6.37 (m, 1H), 6.28 (d, *J* = 8 Hz, 1H), 3.92 (s, 1H), 3.78-3.76 (m, 1H), 3.34-3.20 (m, 1H), 3.20-3.17 (m, 1H), 2.99-2.97 (m, 1H), 2.66-2.61 (m, 1H), 2.13-2.11 (m, 1H), 1.86-1.78 (m, 2H), 1.50-1.47 (m, 1H)

57.5 Preparation of 2-(9-fluoro-2,3,3a,4,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-5-yl)acetamide

To a solution of 9-fluoro-1,2,3,3a,4,5,5a,6-octahydropyrrolo[1,2-a]quinoxaline (5.6 g, 29.1 mmol), diisopropylethyl amine (30.5 mL, 175 mmol) in dimethylformamide (DMF) (100 ml), 2-bromoacetamide (16.08 g, 117 mmol) was added and the reaction solution was stirred 12 h at 100°C. After cooling to 20°C, water (300 ml) was added, and the mixture was extracted with ethyl acetate (3 × 200 ml); the organic layer was dried over anhydrous Na₂SO₄, concentrated under reduced pressure, and the residue was purified by column chromatography on silica gel (dichloromethane: methanol = 100:1 to 50:1) to give the title compound (3 g, 12.03 mmol, yield 41.3%).

LCMS (ESI⁺): *m/z* 250 (M+H)⁺

57.6 Preparation of 2-(9-fluoro-2,3,3a,4,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-5-yl)ethanamine

To a solution of 2-(9-fluoro-2,3,3a,4,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-5-yl)acetamide (3 g, 12.03 mmol) in THF (20 ml), a solution of BH₃.THF (3.1 g, 36.1 mmol) was added at 0°C, then the reaction solution was stirred for 12 h at 70°C. After cooling to 0°C, methanol (100 ml) was added at 0 °C, and the reaction solution was stirred for 30 minutes at 20°C. Then the solution was concentrated under reduced pressure, and the residue was purified by column chromatography on silica gel (petrol

ether/ethyl acetate = 20:1 to 10:1) to give title compound (1.3 g, 5.52 mmol, yield 45.9%).

¹H NMR (400MHz, CD₃OD): δ [ppm]: 6.63-6.59 (m, 1H), 6.48 (d, *J* = 8.4 Hz, 1H), 6.34-6.29 (m, 1H), 3.73-3.70 (m, 1H), 3.34-3.25 (m, 3H), 3.11-3.08 (m, 1H), 2.84-2.79 (m, 3H), 2.64-2.61 (m, 1H), 2.13-2.09 (m, 1H), 1.80-1.76 (m, 2H), 1.55-1.51 (m, 1H)

57.7 Preparation of 1-fluoro-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

To a solution of 2-(9-fluoro-2,3,3a,4,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-5-yl)ethanamine (1.1 g, 4.67 mmol) and formaldehyde (2.106 g, 70.1 mmol) in ethanol (20 ml), trifluoroacetic acid (TFA) (2.88 mL, 37.4 mmol) was added, and the reaction solution was stirred for 12 h at 90°C. After cooling to 40°C, the solution was concentrated under reduced pressure, and the residue was purified by prep-HPLC to give the title compound as TFA salt (1.087 g, yield 94%).

Prep-HPLC Method:

Instrument: Shimadzu LC-20AP preparative HPLC

Column: Luna(2) C18 250*50mm i.d. 10u

Mobile phase: A for H₂O(0.09%TFA) and B for CH₃CN

Gradient: B from 0% to 20% in 20min

Flow rate: 80ml/min

Wavelength: 220 & 254nm

Injection amount: 1.1 g per injection

¹H NMR (400MHz, CD₃OD): δ [ppm]: 6.79-6.71 (m, 2H), 4.23(s, 2H), 4.01-3.96 (m, 1H), 3.36-3.32 (m, 3H), 3.29-3.27(m, 2H), 3.15-3.12 (m, 2H), 2.49-2.43 (m, 1H), 2.29-2.21 (m, 1H), 1.95-1.90 (m, 2H), 1.58-1.54 (m, 1H)

Example 58

1-Bromo-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

(compound of the formula I.a in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-868 of Table A)

58.1 Preparation of 1-(2-bromo-6-nitro-phenyl)-pyrrolidine-2-carboxylate

A mixture of 1-bromo-2-fluoro-3-nitrobenzene (60 g, 273 mmol), methyl pyrrolidine-2-carboxylate (54.2 g, 327 mmol; see example 57.1) and triethylamine (83 g, 818 mmol) was heated at 70°C for 16 h. Then the mixture was cooled, diluted with ethyl acetate (1000 ml), washed with 2N HCl (500 ml), aqueous K₂CO₃ (300 ml) and brine (300 ml) sequentially, and the aqueous phase was extracted with ethyl acetate (500 ml) again. The ethyl acetate layer was dried over Na₂SO₄, filtered, concentrated, and the residue was purified by column chromatography on silica gel (eluted with petrol ether/ethyl acetate = 50:1 to 10:1) to afford the title compound (66 g, yield 74%) as yellow solid.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 7.80-7.78 (m, 1H), 7.64-7.62 (m, 1H), 7.19 (t, *J* = 8.2 Hz, 1H), 4.26-4.16 (m, 1H), 3.70-3.65 (m, 1H), 3.59 (s, 3H), 3.36-3.23 (m, 1H), 2.40-2.23 (m, 2H), 2.15-2.04 (m, 2H)

58.2 Preparation of 9-bromo-2,3,3a,5,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-4-one

To a solution of 1-(2-bromo-6-nitro-phenyl)-pyrrolidine-2-carboxylate (45 g, 137 mmol) in methanol (225 ml) was added a solution of NH₄Cl (65.8 g, 1230 mmol) in water (225 ml) and Fe powder (53.4 g, 957 mmol) at 23°C, and the resulting mixture was heated at 85 °C for 1.5 h. Then the reaction was cooled, diluted with methanol (1000 ml), and the resulting mixture was stirred for 10 minutes, filtered, and the filter cake was washed with methanol (500 ml). The filtrate was concentrated under reduced pressure, the residue was put into water, extracted with ethyl acetate (3 × 800 ml), the extracts were washed with brine (400 ml), dried over Na₂SO₄, filtered, and concentrated to afford the title compound (33 g, yield 90%) as a light yellow solid.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 9.78 (s, 1H), 7.27-7.17 (m, 1H), 6.91-6.75 (m, 2H), 4.10-3.90 (m, 2H), 3.09-3.06 (m, 1H), 2.86-2.68 (m, 1H), 2.37-2.20 (m, 1H), 2.04-2.02 (m, 1H), 1.88-1.68 (m, 1H)

58.3 Preparation of 9-bromo-1,2,3,3a,4,5,5a,6-octahydropyrrolo[1,2-a]quinoxaline

To a solution of 9-bromo-2,3,3a,5,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-4-one (24 g, 90 mmol) in THF (360 ml) was added dropwise BH₃·DMS (50 ml, 500 mmol, 10 M in dimethylsulfide (DMS)) at 23°C, and the resulting solution was stirred at 70°C for 2 h. The reaction was then cooled, and quenched with methanol. One additional reaction was set up and quenched as described above. The two resulting mixtures were combined, concentrated to about one third of volume, 6N HCl (250 mL) was added, and the resulting solution was heated at 70°C for 20 minutes. Then the solution was cooled, adjusted to pH 9 by addition of saturated aqueous K₂CO₃, extracted with ethyl acetate (3 × 400 mL), the extracts were washed with brine, dried over Na₂SO₄, concentrated to give the residue, which was purified by column chromatography on silica gel (eluted with petrol ether/ethyl acetate = 20:1 to 5:1) to afford the title compound (40 g, yield 88%) as a white solid.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 6.93-6.91 (m, 1H), 6.66 (t, *J* = 7.7 Hz, 1H), 6.54-6.51 (m, 1H), 4.24-4.20 (m, 1H), 4.08 (s, 1H), 3.37-3.35 (m, 1H), 3.14-3.11 (m, 1H), 2.89-2.76 (m, 1H), 2.76-2.62 (m, 1H), 2.37-2.22 (m, 1H), 1.96-1.93 (m, 1H), 1.89-1.77 (m, 1H), 1.75-1.63 (m, 1H).

58.4 Preparation of 2-(9-bromo-2,3,3a,4,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-5-yl)acetamide

A mixture of 9-bromo-1,2,3,3a,4,5,5a,6-octahydropyrrolo[1,2-a]quinoxaline (20 g, 79 mmol), 2-bromoacetamide (32.7 g, 237 mmol), and DIEA (51.1 g, 395 mmol) in DMF (200 mL) was stirred at 100 °C for 15 h. The reaction was cooled, diluted with water (600 ml), and extracted with ethyl acetate (3 × 400 ml). The organic layers were washed with brine (200 ml), dried over Na₂SO₄ and filtered, concentrated to give a residue,

which was recrystallized from ethyl acetate to afford the title compound (22 g, yield 90%) as a light yellow solid.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 7.04 (d, *J* = 7.9 Hz, 1H), 6.75 (t, *J* = 7.9 Hz, 1H), 6.55 (d, *J* = 7.9 Hz, 1H), 6.39 (s, 1H), 5.69 (s, 1H), 4.28-4.25 (m, 1H), 3.96-3.75 (m, 2H), 3.55-3.42 (m, 1H), 3.13-3.04 (m, 1H), 3.03-2.91 (m, 1H), 2.77-2.65 (m, 1H), 2.39-2.23 (m, 1H), 2.04-1.90 (m, 1H), 1.90-1.78 (m, 1H), 1.71-1.67 (m, 1H)

58.5 Preparation of 2-(9-bromo-2,3,3a,4,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-5-yl)ethanamine

To a solution of 2-(9-bromo-2,3,3a,4,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-5-yl)acetamide (11 g, 35.5 mmol) in THF (165 ml) was added dropwise BH₃·DMS (18 ml, 180 mmol, 10 M in DMS) at 23°C, and the resulting solution was heated at 70°C for 4 h. The reaction was then quenched with methanol. One additional reaction was set up and quenched as described above. The two resulting mixtures were combined, concentrated to about one third of volume, 6N HCl (200 mL) was added, and the resulting solution was heated at 70°C for 20 minutes. The solution was then cooled, adjusted to pH 9 by addition of saturated aqueous K₂CO₃, and extracted with ethyl acetate (3 × 300 mL). The extracts were washed with brine, dried over Na₂SO₄ and concentrated to give a residue, which was purified by column chromatography on silica gel (eluted with dichloromethane/methanol = 60:1 to 15:1) to afford the title compound (15 g, yield 71%) as a light yellow oil.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 6.94-6.87 (m, 1H), 6.78-6.67 (m, 2H), 4.25-4.21 (m, 1H), 3.51-3.40 (m, 1H), 3.39-3.28 (m, 2H), 3.07-3.05 (m, 1H), 3.02-2.84 (m, 3H), 2.67-2.57 (m, 1H), 2.29-2.27 (m, 1H), 2.00-1.87 (m, 1H), 1.87-1.74 (m, 1H), 1.71-1.60 (m, 1H), 1.26 (s, 2H)

58.6 Preparation of 1-bromo-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

To a solution of 2-(9-bromo-2,3,3a,4,5a,6-hexahydro-1H-pyrrolo[1,2-a]quinoxalin-5-yl)ethanamine (13 g, 43.9 mmol) in ethanol (390 ml) was added formaldehyde (7.1 g, 88 mmol, 37% aqueous) and TFA (15 g, 132 mmol), and the resulting solution was heated at 80°C for 2 h. Then the reaction was concentrated, the residue was dissolved in ethyl acetate (300 ml), and washed with saturated aqueous K₂CO₃ (150 ml), and brine (100 ml). The ethyl acetate layer was dried over Na₂SO₄, filtered, and concentrated to afford the title compound (13 g, crude) as a yellow solid.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 7.17-7.08 (m, 1H), 6.69-6.60 (m, 1H), 4.40-4.27 (m, 1H), 3.96-3.78 (m, 2H), 3.24-3.14 (m, 1H), 3.13-3.02 (m, 2H), 3.01-2.88 (m, 3H), 2.79-2.70 (m, 1H), 2.66-2.58 (m, 1H), 2.36-2.28 (m, 1H), 2.23 (s, 1H), 1.99-1.88 (m, 1H), 1.83-1.71 (m, 1H), 1.71-1.61 (m, 1H)

Analytical Method: The gradient was 1-90% B in 3.4 min, 90-100% B in 0.45 min, 100-1% B in 0.01 min, and then held at 1% B for 0.65 min (0.8 mL/min flow rate). Mobile phase A was 0.0375% TFA in water, mobile phase B was 0.018% TFA in acetonitril. The column used for the chromatography was a 2.1 x 50 mm Venusil XBP-C18 column (5 µm particles). Detection methods are diode array (DAD) and evaporative light scattering (ELSD) detection as well as positive/negative electrospray ionization.).

Example 59

1-Methyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline, trifluoroacetic acid

(compound of the formula I.a in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-869 of Table A)

59.1 Preparation of boc-protected 1-bromo-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

To a solution of 1-bromo-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline (13 g, 42.2 mmol) in dichloromethane (DCM) (260 ml) were added triethylamine (6.4 g, 63.3 mmol) and a solution of Boc₂O (10.1 g, 46.4 mmol) in DCM (20 ml) at 0°C, then it was allowed to warm to 23°C and stirred for 16 h.

The reaction was diluted with DCM (250 ml), washed with 2 N HCl (200 ml), saturated aqueous K₂CO₃ (150 ml) and brine (150 ml), and the water phase was extracted with DCM (200 mL) again. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated, and the residue was purified by column chromatography on silica gel (eluted with petrol ether/ethyl acetate = 30:1 to 4:1) to give the boc-protected compound (6 g, yield 35 %, 2 steps).

LCMS (ESI+): m/z 408 (M+H)⁺

¹H NMR (400 MHz; CD₃OD): δ [ppm]: 7.11 (d, *J* = 7.50 Hz, 1H), 6.73 (d, *J* = 7.94 Hz, 1H), 4.67-4.50 (m, 1H), 4.36-4.24 (m, 1H), 4.18-4.01 (m, 1H), 3.91 (d, *J* = 11.91 Hz, 1H), 3.27 (s, 1H), 3.20-2.90 (m, 4H), 2.76-2.55 (m, 2H), 2.27-2.25 (m, 1H), 1.97-1.66(m, 3H), 1.46 (m, 9H)

59.2 Preparation of boc-protected 1-methyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

To a mixture of boc-protected 1-bromo-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline (500 mg, 1.2 mmol), tricyclohexylphosphine (34.3 mg, 0.12 mmol), and Pd(OAc)₂ (13.7 mg, 0.06 mmol) in toluene (20 ml) under N₂ atmosphere was added methylboronic acid (366 mg, 6.1 mmol), K₃PO₄ (1.04g, 4.9 mmol) and Water (2 mL). The reaction mixture was heated at 100°C for 16 h. After cooling, the reaction mixture was diluted with ethyl acetate (100 ml) and washed with brine. The organic layer was dried over Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography on silica gel (eluted with petrol ether/ethyl acetate = 25:1 to 4:1) to afford the title compound (0.35 g, yield 73%) as white solid.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 6.95-6.49 (m, 2H), 4.81-4.48 (m, 1H), 4.20-3.68 (m, 3H), 3.52-3.17 (m, 2H), 3.11-2.81 (m, 3H), 2.78-2.51 (m, 2H), 2.32 (s, 3H), 2.01-1.86 (m, 1H), 1.86-1.71 (m, 1H), 1.71-1.60 (m, 1H), 1.44 (s, 9H)

59.3 Preparation of 1-methyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

To a solution of boc-protected 1-methyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline (350 mg, 0.9 mmol) in DCM (6 ml) was added TFA (3 ml), and the resulting solution was stirred at 20°C for 5 h. The reaction was then concentrated and the residue was purified by Prep-HPLC and lyophilized to afford the title compound (242 mg, yield 92%, TFA salt) as a light gray solid.

¹H NMR (400 MHz; CD₃OD): δ [ppm]: 7.12 (d, *J* = 7.5 Hz, 1H), 6.97-6.86 (m, 1H), 4.34-4.22 (m, 2H), 4.16-4.11 (m, 1H), 3.69-3.66 (m, 1H), 3.54-3.43 (m, 1H), 3.39-3.33 (m, 2H), 3.30-3.23 (m, 2H), 3.10-2.97 (m, 2H), 2.43 (s, 3H), 2.41-2.31 (m, 1H), 2.13-1.98 (m, 2H), 1.95-1.85 (m, 1H)

Example 60

1-Cyclopropyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline, trifluoroacetic acid

(compound of the formula I.a in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-873 of Table A)

The compound was prepared in analogy to example 59.

LCMS (ESI⁺): *m/z* 270 (M+H)⁺

¹H NMR (400 MHz; CD₃OD): δ [ppm]: 7.10 (d, *J* = 7.9 Hz, 1H), 6.67 (d, *J* = 7.9 Hz, 1H), 4.41-4.38 (m, 1H), 4.26 (s, 2H), 3.67-3.66 (m, 1H), 3.52-3.42 (m, 1H), 3.39-3.33 (m, 2H), 3.29-3.21 (m, 2H), 3.15-3.09 (m, 1H), 3.00-2.87 (m, 1H), 2.46-2.31 (m, 1H), 2.17-1.94 (m, 3H), 1.93-1.82 (m, 1H), 1.20-1.17 (m, 1H), 1.07-0.93 (m, 2H), 0.69-0.54 (m, 1H).

Example 61

1-Cyclobutyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

(compound of the formula I.a in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-874 of Table A)

61.1 Preparation of boc-protected 1-(1-hydroxy-cyclobut-1-yl)-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

To a solution of boc-protected 1-bromo-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline (800 mg, 1.96 mmol) in THF (16 ml) was added dropwise n-BuLi (1.6 ml, 3.92 mmol, 2.5 M in hexane) at -70°C with stirring under N₂ atmosphere, and the resulting solution was stirred at -70°C for 1 hour. A solution of cyclobutanone (275 mg, 3.92 mmol) in THF (0.5 ml) was added dropwise at -70°C, and the reaction solution was stirred for 2 h at same temperature. Then it was allowed to warm slowly to 20°C and stirred for 16 h. The reaction was quenched by saturated aqueous NH₄Cl, extracted with ethyl acetate (2 × 100 ml), the organic layers were dried over Na₂SO₄, concentrated, and the residue was purified by column chromatography on silica gel (eluted with petrol ether/ethyl acetate = 10:1 to 2:1) to afford the title compound (0.5 g, yield 64%) as white solid.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 7.07-6.90 (m, 2H), 4.64-4.48 (m, 1H), 4.23 (d, *J* = 14.1 Hz, 1H), 3.86 (d, *J* = 11.5 Hz, 1H), 3.68-3.67 (m, 1H), 3.52-3.31 (m, 1H), 3.25-3.11 (m, 2H), 3.10-2.94 (m, 2H), 2.94-2.76 (m, 2H), 2.60-2.47 (m, 2H), 2.46-2.33 (m, 2H), 2.29-2.16 (m, 1H), 2.01-1.82 (m, 3H), 1.82-1.68 (m, 1H), 1.68-1.54 (m, 1H), 1.41 (s, 9H).

61.2 Preparation of 1-cyclobutyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

To a solution of boc-protected 1-(1-hydroxy-cyclobut-1-yl)-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline (380 mg, 0.95 mmol) in DCM (3 ml) was added triethylsilane (1.66 g, 14.3 mmol), and the resulting solution was stirred for 0.5 h. Then TFA (10 ml) was added, and the resulting solution was stirred at 60°C overnight. The solvent was removed under reduced pressure. The residue

was dissolved in 3 N HCl (50 ml), and washed with DCM (3 × 50 ml). The aqueous phase was adjusted to pH 9 by addition of saturated aqueous K₂CO₃, the resulting mixture was extracted with DCM (2 × 100 ml), the DCM layers were washed with brine (30 ml), dried over Na₂SO₄, filtered, and concentrated to afford the title compound (160 mg, yield 60%) as a gray solid.

LCMS (ESI+): m/z 284(M+H)⁺

¹H NMR (400 MHz; CD₃OD): δ [ppm]: 6.91 (d, *J* = 7.9 Hz, 1H), 6.79 (d, *J* = 7.5 Hz, 1H), 3.89-3.71 (m, 3H), 3.69-3.60 (m, 1H), 3.17-3.05 (m, 1H), 3.05-2.84 (m, 5H), 2.68-2.48 (m, 2H), 2.48-2.36 (m, 1H), 2.33-2.18 (m, 2H), 2.17-1.93 (m, 3H), 1.93-1.74 (m, 3H), 1.68-1.54 (m, 1H)

Example 62

1-Cyclopentyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline, trifluoroacetic acid

(compound of the formula I.a in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-875 of Table A)

62.1 Preparation of boc-protected 1-(1-hydroxy-cyclopent-1-yl)-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

To a solution of boc-protected 1-bromo-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline (100 mg, 0.25 mmol) in THF (2 ml) was added dropwise n-BuLi (0.2 ml, 0.5 mmol, 2.5 M in hexane) at -70°C with stirring under N₂ atmosphere, and the resulting solution was stirred at -70°C for 1 hour. A solution of cyclopentanone (40 mg, 0.5 mmol) in THF (0.2 ml) was added dropwise at -70°C, and the reaction solution was stirred for 8 h at same temperature. It was then allowed to warm slowly to 20°C and stirred for 16 h. The reaction was quenched by saturated aqueous NH₄Cl, extracted with ethyl acetate (2 × 50 ml), the organic layers were dried over Na₂SO₄, concentrated, and the residue was purified by Prep-TLC (petrol ether/ethyl acetate = 2:1) to afford the title compound (16 mg, yield 22%) as white solid.

¹H NMR (400 MHz; DMSO-*d*₆ (T = 273+80 K)): δ [ppm]: 7.50 (s, 1H), 6.81-6.75 (m, 1H), 6.74-6.68 (m, 1H), 4.46-4.24 (m, 2H), 3.75-3.66 (m, 2H), 3.42-3.26 (m, 2H), 3.24-3.12 (m, 2H), 2.93 (d, *J* = 9.7 Hz, 1H), 2.63-2.51 (m, 2H), 2.15-2.03 (m, 4H), 1.99-1.90 (m, 1H), 1.83-1.82 (m, 5H), 1.72-1.55 (m, 3H), 1.34 (s, 9H)

62.2 Preparation of boc-protected 1-cyclopent-1-enyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

A mixture of boc-protected 1-(1-hydroxy-cyclopent-1-yl)-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline (100 mg, 0.25 mmol), 2-(cyclopent-1-en-1-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (95 mg, 0.49 mmol), Pd(PPh₃)₂Cl₂ (12 mg, 0.017 mmol) and Cs₂CO₃ (160 mg, 0.49 mmol) in dioxane (2 ml) and water (0.6 ml) was placed in a microwave tube under N₂ atmosphere. The reaction mixture was heated at 100 °C for 1 hour in microwave reactor. Six additional reactions were set up as described above. All seven reaction mixtures were combined, was put into water, extracted with ethyl acetate (2 × 100 ml); the extracts were washed with brine, dried over Na₂SO₄, and concentrated to give crude product, which was purified by column chromatography on silica gel (eluted with petrol ether/ethyl acetate = 25:1 to 5:1) to afford the title compound (400 mg, yield 58%) as light yellow solid.

¹H NMR (400 MHz; CDCl₃): δ[ppm]: 6.93-6.68 (m, 2H), 5.94-5.78 (m, 1H), 4.81-4.52 (m, 1H), 4.15 (d, *J* = 13.7 Hz, 1H), 4.08- 3.78 (m, 1H), 3.71-3.58 (m, 1H), 3.45-3.17 (m, 2H), 3.15-2.85 (m, 4H), 2.69-2.41 (m, 4H), 2.38-2.18 (m, 2H), 2.02-1.96 (m, 2H), 1.86-1.66 (m, 2H), 1.59-1.51 (m, 1H), 1.44 (s, 9H)

62.3 Preparation of boc-protected 1-cyclopentyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

A mixture of 1-cyclopent-1-enyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline (0.3 g, 0.76 mmol), and Pd/C (0.09 g, 5%) in methanol (20 ml) was stirred at 20 °C under a H₂ balloon for 16 h. Then the

reaction was filtered, and concentrated to afford the title compound (0.29 g, yield 96%) as light yellow solid.

LCMS (ESI+): m/z 398 (M+H)⁺

62.4 Preparation of 1-cyclopentyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline

To a solution of boc-protected 1-cyclopentyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline (0.29 g, 0.74 mmol) in DCM (6 ml) was added TFA (3 ml), and the resulting solution was stirred for 3 h at 25°C. The reaction was concentrated and lyophilized to afford the title compound (351 mg, yield 90%, TFA salt) as brown oil.

¹H NMR (400 MHz; CD₃OD): δ[ppm]: 7.34 (d, *J* = 8.4 Hz, 1H), 7.13 (d, *J* = 7.9 Hz, 1H), 4.37-4.25 (m, 2H), 4.24-4.14 (m, 1H), 3.96-3.87 (m, 1H), 3.62-3.52 (m, 1H), 3.44-3.32 (m, 4H), 3.30-3.19 (m, 3H), 2.48-2.35 (m, 1H), 2.25-2.04 (m, 4H), 2.04-1.85 (m, 3H), 1.85-1.70 (m, 2H), 1.70-1.56 (m, 2H)

Example 63

8-Methyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-8-ol
(compound of the formula I.g, I.h or I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-241 of Table B)

63.1 Preparation of boc-protected ethyl 3-(2,3,4,5-tetrahydro-1,4-benzodiazepin-1-yl)propanoate

To a solution of 4-boc-protected 2,3,4,5-tetrahydro-1H-1,4-benzodiazepine (7 g, 28.2 mmol) in acetic acid (14 ml) was added ethyl acrylate (14.1 g, 141 mmol) and the reaction mixture was heated at 100°C for 16 hrs. Then it was cooled, concentrated, and the residue was purified by column chromatography on silica gel (eluted with petrol

ether/ethyl acetate = 25:1 to 5:1) to give the title compound (5 g, yield 51%) as light yellow oil.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 7.32-7.14 (m, 2H), 7.03-6.88 (m, 2H), 4.43-4.26 (m, 2H), 4.20-4.08 (m, 2H), 3.64-3.51 (m, 4H), 3.10-2.99 (m, 2H), 2.63-2.53 (m, 2H), 1.49-1.38 (m, 9H), 1.30-1.19 (m, 3H)

63.2 Preparation of boc-protected 3-(2,3,4,5-tetrahydro-1,4-benzodiazepin-1-yl)propanoic acid

To a solution of boc-protected ethyl 3-(2,3,4,5-tetrahydro-1,4-benzodiazepin-1-yl)propanoate (7.5 g, 21.5 mmol) in THF (70 mL), H₂O (50 mL) and methanol (23 mL) was added a solution of NaOH (2.6 g, 64.6 mmol) in H₂O (20 mL) at 5°C, and the resulting solution was stirred at 25°C for 3 h. Then the solution was cooled to 5°C, adjusted to pH 3 by addition of 1N HCl, and the resulting mixture was extracted with ethyl acetate three times. The ethyl acetate layers were combined, dried over Na₂SO₄, filtered, and concentrated to give the title compound (6.7 g, yield 97%) as yellow oil.

¹H NMR (400 MHz; CD₃OD): δ [ppm]: δ 7.23-7.06 (m, 2H), 6.98-6.85 (m, 2H), 4.39-4.21 (m, 2H), 3.65-3.41 (m, 4H), 3.05-2.91 (m, 2H), 2.56 (t, *J* = 6.6 Hz, 2H), 1.40-1.31 (m, 9H).

63.3 Preparation of boc-protected 3-(2,3,4,5-tetrahydro-1,4-benzodiazepin-1-yl)propanoyl chloride

To a solution of boc-protected 3-(2,3,4,5-tetrahydro-1,4-benzodiazepin-1-yl)propanoic acid (6.5 g, 20.3 mmol) in dichloromethane (300 mL) were added dimethylformamide (0.15 g, 2 mmol) and dropwise a solution of (COCl)₂ (7.7 g, 60.9 mmol) in dichloromethane (30 mL) at 0°C. The reaction was stirred for 1 hour at 25°C, and concentrated under reduced pressure below 30°C to give the title compound as a residue which was used directly in the next step without further purification.

63.4 Preparation of boc-protected 1,2,3,4,6,7-hexahydro[1,4]diazepino[6,7,1-ij]-quinolin-8-one

To a solution of boc-protected 3-(2,3,4,5-tetrahydro-1,4-benzodiazepin-1-yl)propanoyl chloride (6.6 g, crude, ~20 mmol) in dichloroethane (400 ml) was added AlCl_3 (10.4 g, 78 mmol) in portions at 0~5 °C, then the resulting mixture was stirred at 65 °C for 7 hrs. The reaction was quenched with water (100 ml), adjusted to pH 9 by addition of aqueous saturated K_2CO_3 , then Boc_2O (6.4 g, 29.2 mmol) was added, and the resulting mixture was stirred for 2 hrs. The mixture was extracted with dichloromethane twice, the organic phase was washed with brine, dried over Na_2SO_4 , concentrated, and the residue was purified by column chromatography on silica gel (eluted with petrol ether/ethyl acetate = 10:1 to 2:1) to give the title compound (4.2 g, yield 71%) as yellow solid.

$^1\text{H NMR}$ (400 MHz, CDCl_3): δ [ppm]: 7.85 (d, $J = 7.5$ Hz, 1H), 7.40-7.20 (m, 1H), 6.83 (t, $J = 7.5$ Hz, 1H), 4.58-4.40 (m, 2H), 3.74 (d, $J = 3.5$ Hz, 2H), 3.60-3.50 (m, 2H), 3.44-3.37 (m, 2H), 2.72-2.62 (m, 2H), 1.47-1.29 (m, 9H).

63.5 Preparation of 1,2,3,4,6,7-hexahydro[1,4]diazepino[6,7,1-ij]-quinolin-8-one

To a solution of boc-protected 1,2,3,4,6,7-hexahydro[1,4]diazepino[6,7,1-ij]-quinolin-8-one (500 mg, 1.7 mmol) in dichloromethane (10 ml) was added trifluoroacetic acid (5 ml), and then the resulting mixture was stirred at 25 °C for 3 hrs. This was concentrated and lyophilized to give the title compound (500 mg, yield 100 %, trifluoroacetic acid salt) as yellow solid.

$^1\text{H NMR}$ (400 MHz; CD_3OD): δ [ppm]: 7.92-7.90 (m, 1H), 7.49 (d, $J = 6.6$ Hz, 1H), 6.97 (t, $J = 7.7$ Hz, 1H), 4.40 (s, 2H), 3.71-3.63 (m, 4H), 3.55-3.47 (m, 2H), 2.75-2.67 (m, 2H).

63.6 Preparation of Cbz-protected 1,2,3,4,6,7-hexahydro[1,4]diazepino[6,7,1-ij]-quinolin-8-one

To a solution of 1,2,3,4,6,7-hexahydro[1,4]diazepino[6,7,1-ij]-quinolin-8-one (2 g, 6.92 mmol) (TFA salt) in DCM (30 ml) were added triethylamine (1.4 g, 13.8 mmol) and a solution of CbzCl (Cbz = benzyloxycarbonyl; 1.5 g, 9 mmol) in DCM (5 mL) at 0~5 °C, then the reaction mixture was stirred at 25 °C for 2 hrs. Then it was diluted with DCM (100 ml), washed with aqueous K₂CO₃ (50 mL) and brine (50 ml); the water phase was extracted with DCM again, the DCM layer was dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give the title compound (1.5 g, yield 64%) as brown oil.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 7.79 (d, *J* = 7.6 Hz, 1H), 7.30-7.11 (m, 6H), 6.76-6.72 (m, 1H), 5.04-5.01 (m, 2H), 4.55-4.48 (m, 2H), 3.75-3.74 (m, 2H), 3.48-3.38 (m, 4H), 2.60-2.55 (m, 2H)

63.7 Preparation of Cbz-protected 8-methyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-8-ol

To a solution of Cbz-protected 1,2,3,4,6,7-hexahydro[1,4]diazepino[6,7,1-ij]-quinolin-8-one (1.5 g, 4.5 mmol) in THF (45 ml) was added dropwise methyllithium (3.7 ml, 5.8 mmol, 1.6 M in ether) at -70°C, and the resulting solution was stirred at -70°C for 2 h. Then it was quenched with aqueous NH₄Cl (20 ml), extracted with ethyl acetate (50 ml) three times, the ethyl acetate layers were washed with brine (30 ml), dried over Na₂SO₄, filtered, concentrated, and the residue was purified by column chromatography on silica gel (eluted with DCM/methanol = 150:1 to 50:1) to give the title compound (1 g, yield 64%) as light yellow oil.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 7.47 (d, *J* = 7.5 Hz, 1H), 7.39-7.25 (m, 5H), 7.25-6.99 (m, 1H), 6.93-6.80 (m, 1H), 5.18-5.02 (m, 2H), 4.62-4.35 (m, 2H), 3.79 s, 1H), 3.73-3.59 (m, 1H), 3.45-3.34 (m, 1H), 3.29-3.15 (m, 3H), 2.00-1.89 (m, 2H), 1.59 (s, 3H)

63.8 Preparation of 8-methyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-8-ol

A mixture of Cbz-protected 8-hydroxy-8-methyl-1,2,3,4,7,8-hexahydro-6H-[1,4]diazepino[6,7,1-ij]quinoline (0.5 g, 1.42 mmol), $\text{NH}_3\cdot\text{H}_2\text{O}$ (1 ml) and Pd/C (168 mg, 10%) in THF (30 ml) was stirred under a H_2 balloon for 2 h. Then it was filtered, concentrated, and the residue was purified by Prep-HPC (basic method) and lyophilized to give the title compound (214 mg, yield 69%) as white solid.

LCMS (ESI+): m/z 219 ($\text{M}+\text{H}$)⁺

¹H NMR (400 MHz; CDCl_3): δ [ppm]: 7.44 (d, $J = 7.5$ Hz, 1H), 7.01 (d, $J = 7.1$ Hz, 1H), 6.83 (t, $J = 7.5$ Hz, 1H), 3.92-3.81 (m, 2H), 3.43-3.32 (m, 1H), 3.29-3.17 (m, 1H), 3.14-3.00 (m, 4H), 1.99-1.91 (m, 2H), 1.59 (s, 3H)

Example 64

8-Methoxy-8-methyl-2,3,4,6,7,8-hexahydro-6H-[1,4]diazepino[6,7,1-ij]quinoline (compound of the formula I.g, I.h or I.i in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-201 of Table B)

64.1 Preparation of Cbz-protected 8-methoxy-8-methyl-1,2,3,4,7,8-hexahydro-6H-[1,4]diazepino[6,7,1-ij]quinoline

To a solution of Cbz-protected 8-methyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-8-ol (0.5 mg, 1.42 mmol; see example 63.2) in DMF (10 ml) was added NaH (113 mg, 2.84 mmol) in portions at 0~5 °C, and then the resulting mixture was stirred at 25°C for 0.5 hrs. A solution of MeI (403 mg, 2.84 mmol) in DMF (2 ml) was added dropwise at 0~5 °C, then the reaction mixture was stirred at 25°C for 16 h. It was quenched with water (30 ml), and extracted with ethyl acetate (60 ml) twice. The ethyl acetate layer was washed with brine, dried over Na_2SO_4 , and concentrated to give a residue, which was purified by column chromatography on silica gel (eluted with petrol ether/ ethyl acetate =15:1 to 4:1) to give the title compound (200 mg, yield 39%) as light yellow oil.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 7.41-7.24 (m, 6H), 7.23-7.01 (m, 1H), 6.93-6.79 (m, 1H), 5.12 (s, 2H), 4.64-4.49 (m, 1H), 4.44-4.34 (m, 1H), 3.88-3.75 (m, 1H), 3.69-3.58 (m, 1H), 3.40-3.26 (m, 2H), 3.23-3.12 (m, 2H), 3.08 (s, 3H), 2.28-2.19 (m, 1H), 1.72-1.63 (m, 1H), 1.53 (s, 3H)

64.2 Preparation of 8-methoxy-8-methyl-1,2,3,4,7,8-hexahydro-6H-[1,4]diazepino[6,7,1-ij]quinoline

A mixture of Cbz-protected 8-methoxy-8-methyl-1,2,3,4,7,8-hexahydro-6H-[1,4]diazepino[6,7,1-ij]quinoline (160 mg, 0.44 mmol), NH₃·H₂O (0.5 ml) and Pd/C (50 mg, 10%) in THF (20 ml) was stirred under a H₂ balloon at 25 °C for 2 hrs. Then it was filtered, concentrated, and the residue was purified by Prep-HPC (basic method) and lyophilized to give the title compound (55 mg, yield 54%) as yellow oil.

LCMS (ESI⁺): m/z 233(M+H)⁺

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 7.30-7.27 (m, 1H), 7.02 (d, *J* = 7.1 Hz, 1H), 6.83 (t, *J* = 7.5 Hz, 1H), 3.94-3.84 (m, 2H), 3.38-3.24 (m, 2H), 3.16-3.01 (m, 7H), 2.27-2.22 (m, 1H), 1.69-1.63 (m, 1H), 1.54 (s, 3H)

Example 65

2,8-Dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-8-ol

A mixture of Cbz-protected 8-methyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-8-ol (50 mg, 0.14 mmol; see example 63.2), formaldehyde (21 mg, 0.28 mmol, 40% in water) and Pd/C (17 mg, 10%) in methanol (5 ml) was stirred under a H₂ balloon for 6 hrs. Then it was filtered, concentrated, and the residue was purified by Prep-HPC (basic method) and lyophilized to give the title compound (12 mg, yield 36%) as light yellow solid.

¹H NMR (400 MHz; CDCl₃): δ [ppm]: 7.44-7.42 (m, 1H), 7.07-7.00 (m, 1H), 6.83 (t, *J* = 7.5 Hz, 1H), 3.71-3.57 (m, 2H), 3.40-3.31 (m, 1H), 3.25-3.17 (m, 1H), 3.17-3.04 (m, 2H), 2.82 (t, *J* = 5.1 Hz, 2H), 2.37 (s, 3H), 1.94 (t, *J* = 6.0 Hz, 2H), 1.58 (s, 3H)

Example 66

11-Methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane], 2,2,2-trifluoroacetic acid

(compound of the formula I.i in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-325 of Table B)

The title compound was prepared in analogy to example 1, using however tert-butyl 6-methyl-2,3,4,5-tetrahydro-1H-1,4-benzodiazepine-4-carboxylate and 2-cyclobutylideneacetic acid as starting materials.

ESI-MS [M+H⁺] = 243.20

¹H NMR (DMSO-d₆, 500MHz) [ppm]: 7.28 (d, *J* = 7.9 Hz, 1H), 6.71 (d, *J* = 7.9 Hz, 1H), 3.74 (s, 2H), 3.10-3.03 (m, 2H), 2.99-2.94 (m, 2H), 2.90-2.84 (m, 2H), 2.34-2.25 (m, 2H), 2.22 (s, 3H), 2.09-1.96 (m, 1H), 1.92-1.83 (m, 5H).

Example 67

9-Methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane], trifluoroacetic acid (compound of the formula I.g in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-325 in Table B)

The intermediate tert-butyl 9-chloro-3,4,6,7-tetrahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane]-2(1H)-carboxylate was prepared in analogy to example 1, steps 1.1 to 1.3 using tert-butyl 8-chloro-2,3-dihydro-1H-benzo[e][1,4]diazepine-4(5H)-carboxylate and 2-cyclobutylideneacetyl chloride as starting materials.

67.1 Preparation of tert-butyl 9-methyl-3,4,6,7-tetrahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane]-2(1H)-carboxylate

75 mg (0.21 mmol) of tert-butyl 9-chloro-3,4,6,7-tetrahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane]-2(1H)-carboxylate were dissolved in 0.5 ml of tetrahydrofuran, 0.5 ml toluene and 5 µl water. To this solution 35 µl (0.25 mmol) trimethyl-

boroxine, 0.2 mg (0.4 μmol) of dicyclohexyl(2',4',6'-triisopropyl-[1,1'-biphenyl]-2-yl)phosphine and 65.8 mg (0.31 mmol) of potassium phosphate were added and the solution was degassed with Argon. 0.1 mg (0.4 μmol) of palladium(II)acetate were added and the reaction mixture was heated at 100°C for 10 minutes in a microwave unit. The reaction mixture was concentrated in vacuo. The residue was purified by column chromatography on silica (eluent: 0-30% methanol in dichloromethane) to yield 40 mg of the title compound.

ESI-MS $[\text{M}+\text{H}^+] = 343.20$

67.2 Preparation of 9-methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]-quinoline-8,1'-cyclobutane], trifluoroacetic acid


To 40 mg (0.12 mmol) of tert-butyl 9-methyl-3,4,6,7-tetrahydrospiro[[1,4]diazepino-[6,7,1-ij]quinoline-8,1'-cyclobutane]-2(1H)-carboxylate in 3 ml dichloromethane were added 90 μl (1.2 mmol) of trifluoroacetic acid at 0°C. The solution was stirred for 2 h at room temperature. The reaction mixture was concentrated in vacuo. The residue was purified by preparative HPLC to yield 4.6 mg of the title compound.

ESI-MS $[\text{M}+\text{H}^+] = 243.20$

^1H NMR (CDCl_3 , 500MHz): δ [ppm]: 9.15 (bs, 2H), 6.95 (d, 1H), 6.80 (d, 1H), 4.15 (m, 2H), 3.30 (m, 2H), 3.25 (m, 2H), 3.10 (m, 2H), 2.85 (m, 2H), 2.70 (s, 3H), 2.20 (m, 2H), 2.00 (m, 2H), 1.85 (m, 2H).

Example 68

4,5,6,7,9,9a,9b,10,10a,11-Decahydrocyclopropa[3,4]pyrrolo[1,2-a][1,4]diazepino[1,7,6-de]quinoxaline



The title compound was prepared in analogy to example 44 using however 1-fluoro-2-nitrobenzene and ethyl 3-azabicyclo[3.1.0]hexane-2-carboxylate as starting materials.

ESI-MS $[M+H]^+ = 242.20$

1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 6.65 (m, 1H), 6.35 (d, 1H), 6.30 (d, 1H), 3.65 (m, 1H), 3.60 (m, 1H), 3.35 (m, 4H), 3.30 (m, 1H), 2.90 (m, 3H), 2.55 (m, 1H), 1.65 (m, 1H), 1.60 (m, 1H), 0.50 (m, 1H), 0.15 (m, 1H).

Example 69

1-Ethyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline (compound of the formula I.a, I.b or I.c in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} is as in row A-289 in Table A)

The title compound was prepared in analogy to example 33, using however acetaldehyde instead of cyclobutanone.

ESI-MS $[M+H]^+ = 218.20$

1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 6.70 (m, 1H), 6.60 (d, 1H), 6.40 (d, 1H), 3.65 (s, 2H), 3.35 (m, 2H), 3.15 (m, 2H), 3.10 (m, 2H), 2.90 (m, 2H), 2.80 (m, 2H), 1.05 (t, 3H).

Example 70

1-Propyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline (compound of the formula I.a, I.b or I.c in which the combination of R^{5a} , R^{5b} , R^6 and R^{9a} is as in row A-325 in Table A)

The title compound was prepared in analogy to example 33, using however propionaldehyde instead of cyclobutanone.

ESI-MS $[M+H]^+ = 232.20$

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 6.70 (m, 1H), 6.60 (d, 1H), 6.40 (d, 1H), 3.65 (s, 2H), 3.25 (m, 2H), 3.15 (m, 4H), 2.90 (m, 2H), 2.80 (m, 2H), 1.50 (m, 2H), 0.90 (t, 3H).

Example 71

1-Cyclobutyl-5-methyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline (compound of the formula I.gg, I.hh or I.ii in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-433 in Table A)

The title compound was prepared in analogy to example 33, using tert-butyl 5-methyl-2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate and cyclobutanone. tert-Butyl 5-methyl-2,3,5,6-tetrahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline-7(8H)-carboxylate was prepared in analogy to example 30 however using 2-bromopropanamide instead of 2-chloroacetamide and in analogy to example 32.

ESI-MS [M+H⁺] = 258.20

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 6.65 (m, 1H), 6.55 (d, 1H), 6.45 (d, 1H), 3.90 (m, 1H), 3.80 (m, 1H), 3.55 (m, 1H), 3.30-3.15 (m, 3H), 3.10 (m, 1H), 2.95 (m, 1H), 2.85 (m, 1H), 2.75 (m, 1H), 2.20 (m, 1H), 2.15 (m, 1H), 2.05 (m, 2H), 1.65 (m, 2H), 0.9 (d, 3H).

Example 72

5-Methyl-1-(oxetan-3-yl)-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline (compound of the formula I.gg, I.hh or I.ii in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-541 in Table A)

The title compound was prepared in analogy to example 71, using however oxetane-3-one instead of cyclobutanone.

ESI-MS [M+H⁺] = 260.20

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 6.60 (m, 1H), 6.55 (d, 1H), 6.05 (d, 1H), 4.80 (m, 1H), 4.75 (m, 1H), 4.65 (m, 1H), 4.60 (m, 1H), 4.45 (m, 1H), 3.80 (d, 1H),

3.50 (d, 1H), 3.35 (m, 2H), 3.15 (m, 2H), 2.90 (m, 1H), 2.85 (m, 1H), 2.75 (m, 1H), 0.9 (d, 3H).

Example 73

1-Ethyl-5-methyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline (compound of the formula I.gg, I.hh or I.ii in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-289 in Table A)

The title compound was prepared in analogy to example 71, using however acetaldehyde instead of cyclobutanone.

ESI-MS [M+H⁺] = 232.20

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 6.80 (m, 1H), 6.65 (d, 1H), 6.50 (d, 1H), 3.95 (d, 1H), 3.80 (d, 1H), 3.45-3.05 (m, 8H), 2.85 (m, 1H), 1.15 (t, 3H), 1.00 (d, 3H).

Example 74

5-Methyl-1-propyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline (compound of the formula I.gg, I.hh or I.ii in which the combination of R^{5a}, R^{5b}, R⁶ and R^{9a} is as in row A-325 in Table A)

The title compound was prepared in analogy to example 71, using however propionaldehyde instead of cyclobutanone.

ESI-MS [M+H⁺] = 246.20

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 6.80 (m, 1H), 6.60 (d, 1H), 6.50 (d, 1H), 3.95 (d, 1H), 3.80 (d, 1H), 3.30 (m, 4H), 3.20 (m, 2H), 3.10 (m, 1H), 3.00 (m, 1H), 2.80 (m, 1H), 1.60 (m, 2H), 0.95 (d, 3H), 0.90 (t, 3H).

Example 75

9-Chloro-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane], trifluoroacetic acid (compound of the formula I.g in which the combination of R^{5a}, R^{5b}, R⁷, R⁸ and R^{9a} is as in row B-323 in Table B)

The title compound was prepared in analogy to example 1 using tert-butyl 8-chloro-2,3-dihydro-1H-benzo[e][1,4]diazepine-4(5H)-carboxylate and 2-cyclobutylideneacetyl chloride as starting materials.

ESI-MS $[M+H^+] = 236.10$

^1H NMR (CDCl_3 , 500MHz): δ [ppm]: 9.60 (bs, 2H), 7.00 (m, 2H), 4.15 (s, 2H), 3.25 (m, 6H), 3.10 (m, 2H), 2.15 (m, 2H), 2.05 (m, 2H), 1.75 (m, 2H).

Example 76

10-Methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane], trifluoroacetic acid (compound of the formula I.h in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-325 in Table B)

ESI-MS $[M+H^+] = 243.20$

^1H NMR (CDCl_3 , 500MHz): δ [ppm]: 9.35 (bs, 2H), 7.40 (s, 1H), 6.90 (s, 1H), 4.15 (m, 2H), 3.35 (m, 2H), 3.25 (m, 2H), 3.15 (m, 2H), 2.40 (m, 2H), 2.30 (s, 3H), 2.15-1.90 (m, 6H).

Example 77

10-Chloro-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane], trifluoroacetic acid (compound of the formula I.h in which the combination of R^{5a} , R^{5b} , R^7 , R^8 and R^{9a} is as in row B-323 in Table B)

ESI-MS $[M+H^+] = 236.10$

^1H NMR (CDCl_3 , 500MHz): δ [ppm]: 9.65 (bs, 2H), 7.55 (s, 1H), 7.05 (s, 1H), 4.15 (s, 2H), 3.30 (m, 4H), 3.20 (m, 2H), 2.35 (m, 2H), 2.15-1.90 (m, 6H).

II. Biological tests

Functional activity

The functional activity of compounds of formula I was assayed by incubation with U2OS_HTR_{2C}_β-Arrestin cells (DiscoverX, 93-0289C3) to induce beta-arrestin2 recruitment to the 5-HT_{2C} receptor. The agonist-induced recruitment and proximity of the receptor and beta-arrestin2 leads to complementation and formation of active β-galactosidase. The enzyme complementation results in enzyme activity, which is measured following the termination of the agonist incubation using DiscoverX's detection reagent, which contains a chemiluminescent substrate which produces a high intensity signal. Cells were plated and a medium-change to a 1% serum containing medium was performed 24h later. The next day, test compounds were added and incubated for 1.5 h before addition of detection reagent.

The response produced was measured and compared with the response produced by 10 [μ]M 5-HT or the maximal effect induced by 5-HT (defined as 100%) to which it was expressed as a percentage response (relative efficacy). Dose response curves were constructed using Graphpad Prism (Graph Software Inc.) or using in house adapted software using a 4 parameter dose response model with variable slope ($\text{fit} = (\text{Bottom} + (\text{Top}-\text{Bottom})/(1+10^{((\text{LogEC50}-x)*\text{HillSlope}))})$ res = (y-fit)). Results are compiled in the table below.

Table

#	Potency (EC50) in functional assay	% efficacy
1	++	71
6	++	63
8	++	40
9	+++	61
16	+	55
17	++	84
19	++	25
22	++	69
25	++	90
26	+++	107
27	+	87

#	Potency (EC50) in functional assay	% efficacy
28	++	55
33	+	78
35	++	112
36	+	31
37	+	21
40	++	131
41	+	75
42	+++	163
43	+	59
44	++	55
45	++	77
46	+	71
47	++	59
48	++	105
49	+	78
51	++	102
54	+	24
55	+	26
56	+++	124
57	+	35
59	++	51
66	+	53
67	+++	116
68	++	76
69	+	54
70	++	63
76	+++	98


Potency (EC50):

+ from 200 nM to < 1µM

++	from 20 nM to < 200 nM
+++	< 20 nM

We claim:

1. A compound of the formula I



wherein

G is $(CR^{3a}R^{3b})_n$;

X is NR^6 or CR^7R^8 ;

R^1 is selected from the group consisting of hydrogen, cyano, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, $-C(=O)R^{10}$, phenyl, phenyl- C_1 - C_2 -alkyl and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO_2 and optionally also 1 or 2 $C=O$ and/or $C=S$ groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R^{11} ;

each R^2 is independently selected from the group consisting of cyano, nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl,

fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹;

R^{3a} and R^{3b}, independently of each other, are selected from the group consisting of hydrogen, cyano, nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹;

R^{4a} and R^{4b}, independently of each other, are selected from the group consisting of hydrogen, cyano, nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl,

fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹; or R^{4a} and R^{4b} form together a group =O or =S;

R^{5a} and R^{5b}, independently of each other, are selected from the group consisting of hydrogen, deuterium, halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹; where R^{5a} and R^{5b} are not simultaneously hydroxy; or

R^{5a} and R^{5b} , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ; or

R^{5a} and R^6 , together with the atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ; or

R^{5a} and R^7 , together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ;

R^6 is selected from the group consisting of hydrogen, cyano, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-C₁-C₄-alkyl, fluorinated C₃-C₈-cycloalkyl-C₁-C₄-alkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, -C(=O) R^{10} , -SO₂ R^{10} , phenyl, phenyl-C₁-C₂-alkyl and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO₂ and optionally also 1 or 2 C=O and/or C=S groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R^{11} ;

R^7 and R^8 , independently of each other, are selected from the group consisting of deuterium, halogen, cyano, nitro, hydroxyl, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, fluorinated C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, fluorinated C_1 - C_6 -alkylsulfonyl, $-NR^{12a}R^{12b}$, $-CH_2NR^{12a}R^{12b}$, $-NR^{12a}C(O)R^{10}$, $-C(=O)R^{10}$, $SO_2NR^{12a}R^{12b}$, C_1 - C_6 -alkylcarbonyloxy, fluorinated C_1 - C_6 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_2 -alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO_2 , C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R^{11} ; where R^7 and R^8 are not simultaneously hydroxyl; and where R^7 is not hydroxyl if R^8 is C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, phenoxy or benzyloxy; or R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO_2 , C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ;

each R^9 is independently selected from the group consisting of halogen, cyano, nitro, hydroxy, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_3 - C_8 -cycloalkenyl, fluorinated C_3 - C_8 -cycloalkenyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy-

C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, -NR^{12a}C(O)R¹⁰, -C(=O)R¹⁰, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy, fluorinated C₁-C₆-alkylcarbonyloxy, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R¹¹; or two radicals R⁹ bound on neighboring carbon atoms, together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹;

each R¹⁰ is independently selected from the group consisting of hydrogen, cyano, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, phenyl, phenyl-C₁-C₂-alkyl, phenoxy, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the five last-mentioned radicals may be substituted with one or more substituents R¹¹;

each R¹¹ is independently selected from the group consisting of halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-hydroxyalkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-

alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₄-alkyl, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₁-C₆-alkylthio, fluorinated C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, fluorinated C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, fluorinated C₁-C₆-alkylsulfonyl, -COOH, -NR^{12a}R^{12b}, -CH₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyl, fluorinated C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, fluorinated C₁-C₆-alkoxycarbonyl, SO₂NR^{12a}R^{12b}, C₁-C₆-alkylcarbonyloxy and fluorinated C₁-C₆-alkylcarbonyloxy;

or two radicals R¹¹, together with the atom(s) they are bound to, form a saturated, partially unsaturated or maximally unsaturated 3-, 4-, 5-, 6- or 7-membered carbocyclic or heterocyclic ring, where the heterocyclic ring contains 1, 2 or 3 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members;

R^{12a} and R^{12b}, independently of each other and independently of each occurrence, are selected from the group consisting of hydrogen, cyano, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₂-C₆-alkenyl, fluorinated C₂-C₆-alkenyl, C₂-C₆-alkynyl, fluorinated C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, fluorinated C₃-C₈-cycloalkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, C₁-C₆-alkylcarbonyl, fluorinated C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, fluorinated C₁-C₆-alkoxycarbonyl, phenyl and benzyl, where the phenyl moieties in the two last-mentioned radicals may carry 1, 2 or 3 substituents selected from halogen, cyano nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy; or,

if R^{12a} and R^{12b} are bound to the same nitrogen atom, together with this nitrogen atom may form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents selected from halogen, cyano nitro, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy;

a is 0, 1 or 2;

b is 0, 1, 2 or 3; and

n is 1 or 2;

and the N-oxides, tautomeric forms, stereoisomers and pharmaceutically acceptable salts thereof, and the compound of the general formula I, wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope.

2. The compound as claimed in claim 1, wherein at least one hydrogen atom has been replaced by a deuterium atom.
3. The compound as claimed in any of the preceding claims, where R^1 is selected from hydrogen and C_1 - C_6 -alkyl and is preferably hydrogen.
4. The compound as claimed in any of the preceding claims, where R^2 is selected from cyano, nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy and fluorinated C_1 - C_6 -alkoxy, in particular from C_1 - C_6 -alkyl and especially from methyl.
5. The compound as claimed in any of the preceding claims, where R^{3a} and R^{3b} , independently of each other, are selected from hydrogen, cyano, nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy and fluorinated C_1 - C_6 -alkoxy.
6. The compound as claimed in claim 5, where R^{3a} is selected from hydrogen, cyano, nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy and fluorinated C_1 - C_6 -alkoxy, and R^{3b} is hydrogen.
7. The compound as claimed in claim 6, where R^{3a} is selected from hydrogen and methyl, and R^{3b} is hydrogen.

8. The compound as claimed in claim 7, where R^{3a} and R^{3b} are hydrogen.
9. The compound as claimed in any of the preceding claims, where R^{4a} and R^{4b} , independently of each other, are selected from hydrogen, C_1 - C_6 -alkyl and fluorinated C_1 - C_6 -alkyl or form together a group =O, and are in particular hydrogen or methyl or form together a group =O, and are especially hydrogen.
10. The compound as claimed in any of the preceding claims, where
 R^{5a} is selected from hydrogen, cyano, nitro, hydroxy, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy and fluorinated C_1 - C_6 -alkoxy; and in case that X is CR^7R^8 is further selected from halogen; or
 R^{5a} and R^6 , together with the atoms they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ; or
 R^{5a} and R^7 , together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R^{11} ; and
 R^{5b} is selected from hydrogen and deuterium.
11. The compound as claimed in claim 10, where
 R^{5a} is selected from hydrogen, C_1 - C_6 -alkyl and fluorinated C_1 - C_6 -alkyl; and in case that X is CR^7R^8 is further selected from halogen; and is in particular hydrogen; or
 R^{5a} and R^6 form together a group $(CH_2)_r$, where r is 2, 3, 4 or 5; or
 R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 2, 3, 4 or 5; and
 R^{5b} is selected from hydrogen and deuterium.

12. The compound as claimed in claim 10, where
R^{5a} is selected from hydrogen, C₁-C₆-alkyl and fluorinated C₁-C₆-alkyl; and in case that X is CR⁷R⁸ is further selected from halogen; and is in particular hydrogen; or
R^{5a} and R⁶ form together a group (CH₂)_r, where r is 2, 3, 4 or 5, where two hydrogen atoms bound to adjacent CH₂ groups may be replaced by two radicals R¹¹, where the two radicals R¹¹ form together a group (CH₂)_t, where t is 1, 2, 3, 4 or 5; or
R^{5a} and R⁷ form together a group (CH₂)_s, where s is 2, 3, 4 or 5; and
R^{5b} is selected from hydrogen and deuterium.
13. The compound as claimed in any of claims 1 to 9, where R^{5a} and R^{5b}, together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹; and form preferably a 5- or 6-membered saturated or partially unsaturated carbocyclic ring, and where the ring may be substituted with one or more substituents R¹¹.
14. The compound as claimed in any of the preceding claims, where X is NR⁶, where R⁶ is as defined in claim 1.
15. The compound as claimed in claim 14, where X is NR⁶, where R⁶ is -SO₂R¹⁰, where R¹⁰ is as defined in claim 1.
16. The compound as claimed in claim 15, where R¹⁰ is selected from C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl and phenyl, where phenyl may be substituted with 1, 2, 3, 4 or 5 radicals R¹¹.

17. The compound as claimed in claim 14, where X is NR^6 , where R^6 is selected from hydrogen, $\text{C}_1\text{-C}_6\text{-alkyl}$, fluorinated $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_3\text{-C}_6\text{-cycloalkyl}$, fluorinated $\text{C}_3\text{-C}_6\text{-cycloalkyl}$, $\text{C}_3\text{-C}_6\text{-cycloalkyl-C}_1\text{-C}_4\text{-alkyl}$, fluorinated $\text{C}_3\text{-C}_6\text{-cycloalkyl-C}_1\text{-C}_4\text{-alkyl}$, $\text{C}_1\text{-C}_4\text{-alkoxy}$, fluorinated $\text{C}_1\text{-C}_4\text{-alkoxy}$, $-\text{C}(=\text{O})\text{R}^{10}$, where R^{10} is selected from $\text{C}_1\text{-C}_6\text{-alkyl}$ and $\text{C}_3\text{-C}_6\text{-cycloalkyl}$; phenyl, phenyl- $\text{C}_1\text{-C}_2\text{-alkyl}$ and a 3-, 4-, 5- or 6-membered saturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO_2 and optionally also 1 or 2 $\text{C}=\text{O}$ and/or $\text{C}=\text{S}$ groups as ring members, where the cyclic moieties in the three last-mentioned radicals may be substituted with one or more substituents R^{11} .
18. The compound as claimed in claim 17, where R^6 is selected from hydrogen, $\text{C}_1\text{-C}_4\text{-alkyl}$, fluorinated $\text{C}_1\text{-C}_4\text{-alkyl}$, $\text{C}_3\text{-C}_6\text{-cycloalkyl}$, $\text{C}_3\text{-C}_6\text{-cycloalkyl-C}_1\text{-C}_2\text{-alkyl}$, $\text{C}_3\text{-C}_6\text{-cycloalkylcarbonyl}$, $\text{C}_1\text{-C}_4\text{-alkoxy}$, fluorinated $\text{C}_1\text{-C}_4\text{-alkoxy}$, phenyl- $\text{C}_1\text{-C}_2\text{-alkyl}$ and a 3-, 4-, 5- or 6-membered saturated heterocyclic ring containing 1 or 2 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO_2 as ring members; and is in particular selected from $\text{C}_1\text{-C}_4\text{-alkyl}$, $\text{C}_3\text{-C}_6\text{-cycloalkyl}$, $\text{C}_3\text{-C}_6\text{-cycloalkyl-C}_1\text{-C}_2\text{-alkyl}$, and a 3-, 4-, 5- or 6-membered saturated heterocyclic ring containing 1 or 2 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO_2 as ring members.
19. The compound as claimed in claim 18, where R^6 is selected from $\text{C}_1\text{-C}_4\text{-alkyl}$, $\text{C}_3\text{-C}_6\text{-cycloalkyl}$ and oxetanyl.
20. The compound as claimed in claim 14, where X is NR^6 , where R^{5a} and R^6 , together with the atoms they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated, partially unsaturated or maximally unsaturated heterocyclic ring, where the ring may further contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO_2 , $\text{C}=\text{O}$ and $\text{C}=\text{S}$ as ring members, and where the ring may be substituted with one or more substituents R^{11} .

21. The compound as claimed in claim 20, where R^{5a} and R^6 form together a group $(CH_2)_r$, where r is 2, 3, 4 or 5 and in particular 3, 4 or 5.
22. The compound as claimed in claim 20, where R^{5a} and R^6 form together a group $(CH_2)_r$, where r is 2, 3, 4 or 5 and in particular 3, 4 or 5, where two hydrogen atoms bound to adjacent CH_2 groups may be replaced by two radicals R^{11} , where the two radicals R^{11} form together a group $(CH_2)_t$, where t is 1, 2, 3, 4 or 5 and in particular 1, 2 or 3.
23. The compound as claimed in any of claims 1 to 13, where X is CR^7R^8 , where R^7 and R^8 are as defined in claim 1.
24. The compound as claimed in claim 23, where R^7 and R^8 , independently of each other, are selected from the group consisting of halogen, cyano, nitro, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, C_2 - C_6 -alkenyl, fluorinated C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, fluorinated C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, fluorinated C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -alkoxy- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylthio, fluorinated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, fluorinated C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, fluorinated C_1 - C_6 -alkylsulfonyl, $-NR^{12a}R^{12b}$, $-CH_2NR^{12a}R^{12b}$, $-NR^{12a}C(O)R^{10}$, $-C(=O)R^{10}$, $SO_2NR^{12a}R^{12b}$, C_1 - C_6 -alkylcarbonyloxy, fluorinated C_1 - C_6 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_2 -alkyl, phenoxy, phenylsulfonyl, benzyloxy and a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO_2 , C=O and C=S as ring members, where the cyclic moieties in the six last-mentioned radicals may be substituted with one or more substituents R^{11} ; or R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or het-

eroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹; or

R^{5a} and R⁷, together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6-, 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹; in which case R⁸ has one of the above meanings or is deuterium.

25. The compound as claimed in claim 22, where

R⁷ and R⁸, independently of each other, are selected from halogen, cyano, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy, fluorinated C₁-C₆-alkoxy, phenyl, phenyl-C₁-C₂-alkyl, benzyloxy and a 3-, 4-, 5- or 6-membered saturated, partially unsaturated or maximally unsaturated ring containing 1, 2, 3 or 4 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO, SO₂, C=O and C=S as ring members, where the cyclic moieties in the four last-mentioned radicals may be substituted with one or more substituents R¹¹; or

R⁷ and R⁸, together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹; or

R^{5a} and R⁷, together with the carbon atoms they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may contain 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO, SO₂, C=O and C=S as ring members, and where the ring may be substituted with one or more substituents R¹¹; in which case R⁸ has one of the above meanings or is deuterium; and

- R^{5b} is selected from hydrogen and deuterium.
26. The compound as claimed in claim 23, where
 R^7 and R^8 , independently of each other, are selected from halogen, cyano, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl and phenyl, or
 R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ; or
 R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 2, 3, 4 or 5; in which case R^8 has one of the above meanings or is deuterium; and
 R^{5b} is selected from hydrogen and deuterium.
27. The compound as claimed in claim 24, where
 R^8 is selected from halogen, cyano, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl and phenyl; and
 R^7 is selected from C_1 - C_6 -alkyl and fluorinated C_1 - C_6 -alkyl, preferably from methyl and CF_3 and is in particular methyl; or
 R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7- or 8-membered saturated, partially unsaturated or maximally unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ; or
 R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 2, 3 or 4; in which case R^8 has one of the above meanings or is deuterium; and
 R^{5b} is selected from hydrogen and deuterium.
28. The compound as claimed in claim 22, where
 R^7 is C_1 - C_6 -alkyl and is in particular methyl;
 R^8 is selected from hydroxyl, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_1 - C_6 -alkoxy, fluorinated C_1 - C_6 -alkoxy and phenyl and in particular from C_1 - C_4 -alkyl, fluorinated C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, fluorinated C_1 - C_4 -alkoxy and phenyl; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7-membered saturated or partially unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ; or


R^{5a} and R^7 form together a group $(CH_2)_s$, where s is 2, 3 or 4 and where R^8 is methyl; and

R^{5b} is hydrogen.

29. The compound as claimed in any of claims 20 to 26, wherein the ring formed by R^7 and R^8 together with the carbon atom they are bound to is a carbocyclic ring, in particular a saturated carbocyclic ring.
30. The compound as claimed in claim 27, where R^7 and R^8 , together with the carbon atom they are bound to, form a 3, 4-, 5- or 6-membered saturated or partially unsaturated carbocyclic ring, preferably a 3, 4- or 5-membered saturated carbocyclic ring, in particular a 4-membered saturated carbocyclic ring, where the ring may be substituted with one or more substituents R^{11} .
31. The compound as claimed in any of claims 20 to 26, wherein the ring formed by R^7 and R^8 together with the carbon atom they are bound to is a heterocyclic ring containing 1, 2, 3 or 4 heteroatoms or heteroatom-containing groups selected from O, S, N, SO and SO_2 as ring members preferably containing 1 or 2 heteroatoms selected from O, S and N as ring members.
32. The compound as claimed in claim 29, wherein the heterocyclic ring is a saturated 3-, 4- or 5-membered heterocyclic ring containing one heteroatom selected from O, S and N as ring member, where the ring may be substituted with one or more substituents R^{11} , and is in particular oxetanyl which may be substituted with one or more substituents R^{11} .
33. The compound as claimed in any of the preceding claims, where each R^9 is independently selected from halogen, cyano, nitro, hydroxy, C_1 - C_6 -alkyl, fluorinated C_1 - C_6 -alkyl, C_3 - C_6 -cycloalkyl, fluorinated C_3 - C_6 -cycloalkyl, C_1 - C_6 -alkoxy, fluori-


nated C₁-C₆-alkoxy and a 3-, 4-, 5- or 6-membered saturated heterocyclic ring containing 1 or 2 heteroatoms or heteroatom groups independently selected from N, O, S, NO, SO and SO₂ as ring members, where the heterocyclic ring may be substituted with one or more substituents R¹¹, in particular from halogen, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₃-C₆-cycloalkyl and C₁-C₆-alkoxy, and is especially halogen.

34. The compound as claimed in any of the preceding claims, where each R¹¹ is independently selected from halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, fluorinated C₁-C₆-alkyl, C₁-C₆-alkoxy and fluorinated C₁-C₆-alkoxy, and is in particular halogen or C₁-C₄-alkyl.
35. The compound as claimed in any of the preceding claims, where a is 0 or 1, in particular 0.
36. The compound as claimed in claim 33, where a is 1 and R² is bound in β -position to the nitrogen ring atom carrying R¹.
37. The compound as claimed in any of the preceding claims, where b is 0 or 1.
38. The compound as claimed in any of the preceding claims, where n is 1.
39. The compound as claimed in any of the preceding claims, of formula I.1



wherein X, R^{5a}, R^{5b}, R⁹ and b are as defined in any of claims 1 and 10 to 34.

40. The compound as claimed in claim 37, of formula I.1.1




wherein

R^{9a} is selected from H, halogen, C_1 - C_4 -alkyl, fluorinated C_1 - C_4 -alkyl, C_3 - C_6 -cycloalkyl and fluorinated C_3 - C_6 -cycloalkyl; and

R^{5a} , R^{5b} , R^6 , R^9 and b are as defined in any of claims 1 and 10 to 35.

41. The compound as claimed in claim 37, of formula I.1.2



wherein

R^7 is selected from C_1 - C_4 -alkyl and fluorinated C_1 - C_4 -alkyl, in particular from methyl and CF_3 , and is especially methyl;


R^8 is selected from C_1 - C_4 -alkyl, fluorinated C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, fluorinated C_1 - C_4 -alkoxy and phenyl; and in particular from methyl and CF_3 ; or

R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7-membered, in particular a 3- or 4-membered saturated or partially unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ;

R^{9a} is H, Cl, F or methyl; and


R^9 and b are as defined in any of claims 1, 31, 32, 34 and 35.

42. The compound as claimed in claim 37, of formula I.1.3




wherein R^8 is selected from deuterium, F, Cl, CN and CH_3 .

43. The compound as claimed in any of claims 1 to 36, of formula I.2



wherein X, R^{5a} , R^{5b} , R^9 and b are as defined in any of claims 1 and 10 to 35.

44. The compound as claimed in claim 41 of formula I.2.1




wherein

R^{2a} , R^{2b} and R^{3a} , independently of each other, are selected from hydrogen and methyl; and where in particular at most one of R^{2a} , R^{2b} and R^{3a} is methyl;

R^{9a} is selected from H, halogen, C_1 - C_4 -alkyl, fluorinated C_1 - C_4 -alkyl, C_3 - C_6 -cycloalkyl and fluorinated C_3 - C_6 -cycloalkyl; and

R^{5a} , R^{5b} , R^6 , R^9 and b are as defined in any of claims 1 and 10 to 19, 30, 31 and 32.

45. The compound as claimed in claim 42, where R^{2b} is methyl and R^{2a} and R^{3a} are hydrogen.
46. The compound as claimed in claim 41, of formula I.2.2



wherein

- R^{2a} , R^{2b} , R^{3a} and R^{5a} , independently of each other, are selected from hydrogen, methyl and ethyl, in particular from hydrogen and methyl;
- R^7 is selected from C_1 - C_4 -alkyl and fluorinated C_1 - C_4 -alkyl, in particular from methyl and CF_3 , and is especially methyl;
- R^8 is selected from C_1 - C_4 -alkyl, fluorinated C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, fluorinated C_1 - C_4 -alkoxy and phenyl; and in particular from methyl and CF_3 ; or
- R^7 and R^8 , together with the carbon atom they are bound to, form a 3-, 4-, 5-, 6- or 7-membered, in particular a 3- or 4-membered saturated or partially unsaturated ring, where the ring may be substituted with one or more substituents R^{11} ;
- R^{9a} is H, Cl, F or methyl; and
- R^9 and b are as defined in any of claims 1, 31, 32 and 34.

47. The compound as claimed in claim 44, wherein R^{2a} , R^{2b} , R^{3a} and R^{5a} are selected from hydrogen and methyl, with the proviso that at most 2, preferably at most 1, of R^{2a} , R^{2b} , R^{3a} and R^{5a} are methyl.
48. The compound as claimed in claim 45, where R^{2b} is methyl and R^{2a} , R^{3a} and R^{5a} are hydrogen.

49. The compound as claimed in any of claims 44 to 46, where R⁷ and R⁸, together with the carbon atom they are bound to, form a 3, 4-, 5- or 6-membered saturated or partially unsaturated carbocyclic ring, preferably a 3, 4- or 5-membered saturated carbocyclic ring, in particular a 4-membered saturated carbocyclic ring, where the ring may be substituted with one or more substituents R¹¹.
50. A compound of formula I as claimed in any of the preceding claims, selected from
- 8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinolone;
- 1,2,3,4-tetrahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopent[2]en]-6(7H)-one;
- 8-methyl-8-phenyl-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-6(2H)-one;
- 8,8-dimethyl-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-6(2H)-one;
- 1,2,3,4-tetrahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopentan]-6(7H)-one;
- 1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopentane];
- 12a-methyl-4,5,6,7,10,11,12,12a-octahydrocyclopenta[c][1,4]diazepino[6,7,1-ij]quinolin-9(9aH)-one;
- 8-methyl-8-(trifluoromethyl)-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinolone;
- 12a-methyl-4,5,6,7,10,11,12,12a-octahydrocyclopenta[c][1,4]diazepino[6,7,1-ij]quinolin-9(9aH)-one;
- 12a-methyl-4,5,6,7,9,9a,10,11,12,12a-decahydrocyclopenta[c][1,4]diazepino[6,7,1-ij]quinoline;
- 8-methyl-8-(trifluoromethyl)-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-6(2H)-one;
- 12a-methyl-4,5,6,7,9,9a,10,11,12,12a-decahydrocyclopenta[c][1,4]diazepino[6,7,1-ij]quinoline;
- 8-methyl-8-phenyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
- (R)-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline;

(S)-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline;
9-chloro-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
10-chloro-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
9-chloro-8,8-dimethyl-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-6(2H)-one;
10-chloro-8,8-dimethyl-3,4,7,8-tetrahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-6(2H)-one;
9-fluoro-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
10-fluoro-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclohexane];
1',2',3',4',6',7'-hexahydrospiro[cyclobutane-1,8'-[1,4]diazepino[6,7,1-ij]quinoline];
9'-fluoro-1',2',3',4',6',7'-hexahydrospiro[cyclobutane-1,8'-[1,4]diazepino[6,7,1-ij]quinoline];
7,8,8-trimethyl-1,2,3,4,7,8-hexahydro-6H-[1,4]diazepino[6,7,1-ij]quinoline];
9-fluoro-1-methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane];
3-methyl-1',2',3',4',6',7'-hexahydrospiro[cyclobutane-1,8'-[1,4]diazepino[6,7,1-ij]quinoline];
3-benzyl-2',3',7',8',9',10'-hexahydro-1H-3',8',10a'-triazacyclohepta[de]naphthalene;
2,3,5,6,7,8-Hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;
1-Cyclobutyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;
1-Methyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;
1-(Oxetan-3-yl)-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;
1-(Cyclopropylmethyl)-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;

1-(Cyclopentylmethyl)-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;
Cyclopropyl(2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxalin-1-yl)methanone;
Cyclopentyl(2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxalin-1-yl)methanone;
1-Cyclopropyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;
1-Cyclopentyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;
1-Cyclopropyl-5-methyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;
6,6-Dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
5,6,7,9,9a,10,11,12,13,14-decahydro-4H-azepino[1,2-a][1,4]diazepino[1,7,6-de]quinoxaline;
7-Methyl-5,6,7,9,9^a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline;
1-Fluoro-7-methyl-5,6,7,9,9^a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline;
1,2,3,4,6,7-Hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopropane];
4-Methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclopropane];
11-Fluoro-8,8-Dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
10-Methoxy-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
4,8,8-Trimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
4-Ethyl-8,8-dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
6,8,8-Trimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
8-ethyl-8-methyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
(R)-8-ethyl-8-methyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
(S)-8-ethyl-8-methyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinoline;
4-Methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane];

1-Fluoro-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline;

1-Bromo-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline;

1-Methyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline;

1-Cyclopropyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline;

1-Cyclobutyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline;

1-Cyclopentyl-5,6,7,9,9a,10,11,12-octahydro-4H-[1,4]diazepino[1,7,6-de]pyrrolo[1,2-a]quinoxaline;

8-Methyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-8-ol;

8-Methoxy-8-methyl-2,3,4,6,7,8-hexahydro-6H-[1,4]diazepino[6,7,1-ij]quinoline;

2,8-Dimethyl-2,3,4,6,7,8-hexahydro-1H-[1,4]diazepino[6,7,1-ij]quinolin-8-ol;

11-Methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane];

9-Methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane];

4,5,6,7,9,9a,9b,10,10a,11-Decahydrocyclopropa[3,4]pyrrolo[1,2-a][1,4]diazepino[1,7,6-de]quinoxaline;

1-Ethyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;

1-Propyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;

1-Cyclobutyl-5-methyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;

5-Methyl-1-(oxetan-3-yl)-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;

1-Ethyl-5-methyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;

5-Methyl-1-propyl-2,3,5,6,7,8-hexahydro-1H-[1,4]diazepino[1,7,6-de]quinoxaline;

9-Chloro-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane];

10-Methyl-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane];

10-Chloro-1,2,3,4,6,7-hexahydrospiro[[1,4]diazepino[6,7,1-ij]quinoline-8,1'-cyclobutane];

and the stereoisomers and pharmaceutically acceptable salts thereof.

51. A pharmaceutical composition comprising a therapeutically effective amount of at least one compound as claimed in any of the preceding claims or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof, or comprising at least one compound as defined in any of the preceding claims wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope, preferably wherein at least one hydrogen atom has been replaced by a deuterium atom, in combination with at least one pharmaceutically acceptable carrier and/or auxiliary substance.
52. The compound as claimed in any of claims 1 to 50 or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for use as a medicament.
53. The compound as claimed in any of claims 1 to 50 or an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for the treatment of disorders which respond to the modulation of the 5-HT_{2c} receptor.
54. The use of a compound as claimed in any of claims 1 to 50 or of an N-oxide, a tautomeric form, a stereoisomer or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of disorders which respond to the modulation of the 5-HT_{2c} receptor.
55. A method for treating disorders which respond to the modulation of the 5-HT_{2c} receptor, which method comprises administering to a subject in need thereof at least one compound as defined in any of claims 1 to 50 or an N-oxide, a tautomer-

ic form, a stereoisomer or a pharmaceutically acceptable salt thereof.

56. The compound as claimed in claim 53 or the use as claimed in claim 54 or the method as claimed in claim 55, where the disorders are selected from the group consisting of damage of the central nervous system, disorders of the central nervous system, eating disorders, ocular hypertension, cardiovascular disorders, gastrointestinal disorders and diabetes.
57. The compound or the use or the method as claimed in claim 56, where the disorders are selected from the group consisting of bipolar disorder, depression, atypical depression, mood episodes, adjustment disorders, anxiety, panic disorders, post-traumatic syndrome, psychoses, schizophrenia, cognitive deficits of schizophrenia, memory loss, dementia of aging, Alzheimer's disease, behavioral disorders associated with dementia, social phobia, mental disorders in childhood, attention deficit hyperactivity disorder, organic mental disorders, autism, mutism, disruptive behavior disorder, impulse control disorder, borderline personality disorder, obsessive compulsive disorder, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, seizure disorders, epilepsy, substance use disorders, alcohol abuse, cocaine abuse, tobacco abuse, smoking cessation, sexual dysfunction/erectile dysfunction in males, sexual dysfunction in females, premenstrual syndrome, late luteal phase syndrome, chronic fatigue syndrome, sleep disorders, sleep apnoea, chronic fatigue syndrome, psoriasis, Parkinson's disease, spinal cord injury, trauma, stroke, pain, bladder dysfunction/urinary incontinence, encephalitis, meningitis, eating disorders, obesity, bulimia, weight loss, anorexia nervosa, ocular hypertension, cardiovascular disorders, gastrointestinal disorders, diabetes insipidus, diabetes mellitus, type I diabetes, type II diabetes, type III diabetes, diabetes secondary to pancreatic diseases, diabetes related to steroid use, diabetes complications, hyperglycemia and insulin resistance.
58. The compound or the use or the method as claimed in claim 57, where the disorders are selected from schizophrenia, depression, bipolar disorders, obesity and substance use disorders.