

METHODS OF DOSING OF APICAL SODIUM-DEPENDENT BILE ACID TRANSPORTER INHIBITORS (ASBTIs)

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application No. 63/271,916, filed October 26, 2021, U.S. Provisional Application No. 63/280,470, filed November 17, 2021, and U.S. Provisional Application No. 63/354,424, filed June 22, 2022, the disclosures of which are herein incorporated by reference in their entirety.

FIELD OF THE INVENTION

[0002] The present invention relates generally to methods for reducing, minimizing, preventing, ameliorating, or eliminating one or more side effects associated with administration of an apical sodium-dependent bile acid transporter inhibitor (ASBTI) in a subject in need thereof by administering a therapeutically effective amount of the ASBTI to the subject before ingestion of food. The present invention relates also to methods of treating cholestatic liver disease in a subject comprising administering a therapeutically effective amount of an ASBTI to the subject before ingestion of food and/or in a fasted state.

BACKGROUND

[0003] Hypercholelism and cholestatic liver diseases are liver diseases associated with impaired bile secretion (i.e., cholestasis), associated with and often secondary to the intracellular accumulation of bile acids/salts in the hepatocyte. Hypercholelism is characterized by increased serum concentration of bile acid or bile salt. Cholestasis can be categorized clinicopathologically into two principal categories of obstructive, often extrahepatic, cholestasis, and nonobstructive, or intrahepatic, cholestasis. Nonobstructive intrahepatic cholestasis can further be classified into two principal subgroups of primary intrahepatic cholestasis that result from constitutively defective bile secretion, and secondary intrahepatic cholestasis that result from hepatocellular injury. Primary intrahepatic cholestasis includes diseases such as benign recurrent intrahepatic cholestasis, which is predominantly an adult form with similar clinical symptoms, and progressive familial intrahepatic cholestasis (PFIC) types 1, 2, and 3, which are diseases that affect children.

[0004] Pediatric cholestatic liver diseases affect a small percentage of children, but therapy results in significant healthcare costs each year. Currently, many of the pediatric cholestatic liver diseases require invasive and costly treatments such as liver transplantation and surgery. An effective and less invasive treatment at a dosing with minimal gastrointestinal adverse effects that is suitable for the pediatric population is not available.

[0005] In recent years apical sodium-dependent bile acid transporter inhibitors (ASBTIs) have evolved as an important novel class of therapeutic agents capable of reducing serum and/or hepatic bile acids and thus reducing cholestasis. There exists a further urgent need to reduce ASBTI adverse effects and/or to develop methods of treating cholestasis and cholestatic liver disease with reduced adverse effects.

SUMMARY OF THE INVENTION

[0006] Various non-limiting aspects and embodiments of the invention are described below.

[0007] In one aspect, the present invention provides a method for reducing, minimizing, preventing, ameliorating, or eliminating one or more side effects associated with administration of an apical sodium-dependent bile acid transporter inhibitor (ASBTI) in a subject in need thereof, the method comprising administering a therapeutically effective amount of the ASBTI to the subject before ingestion of food.

[0008] In certain embodiments, the one or more side effects associated with administration of the ASBTI is reduced, minimized, prevented, ameliorated or eliminated as compared to the side effects when the ASBTI is administered after ingestion of food, at the same time as food, or mixed with food.

[0009] In another aspect, the present invention provides a method of improving gastrointestinal (GI) tolerability of an ASBTI in a subject in need thereof, the method comprising administering a therapeutically effective amount of the ASBTI to the subject before ingestion of food.

[0010] In some embodiments, the improvement of GI tolerability comprises a reduction, minimization, prevention, amelioration, or elimination of one or more GI adverse events.

[0011] In some embodiments, the improvement of GI tolerability comprises a reduction, minimization, prevention, amelioration, or elimination of one or more of diarrhea, loose stools, nausea, abdominal pain, and anorectal discomfort.

[0012] In some embodiments, the GI tolerability is improved as compared to the GI tolerability when the ASBTI is administered at mealtime or immediately after food intake.

[0013] In some embodiments, the GI tolerability is improved by at least 10% as compared to the GI tolerability when the ASBTI is administered at mealtime or immediately after food intake.

[0014] In some embodiments, the GI tolerability is improved by at least 20% as compared to the GI tolerability when the ASBTI is administered at mealtime or immediately after food intake.

[0015] In some embodiments, the GI tolerability is improved by at least 50% as compared to the GI tolerability when the ASBTI is administered at mealtime or immediately after food intake.

[0016] In one aspect, the present invention provides method of treating cholestatic liver disease in a subject in need thereof, the method comprising administering a therapeutically effective amount of an ASBTI to the subject before ingestion of food, wherein the subject experiences a reduction in frequency and/or severity of one or more side effects associated with the administration of the ASBTI.

[0017] In certain embodiments, the frequency and/or severity of side effects is reduced as compared to the side effects when the ASBTI is administered after ingestion of food, at the same time as food, or mixed with food.

[0018] In certain embodiments, the cholestatic liver disease is a pediatric cholestatic liver disease. In certain embodiments, the cholestatic liver disease is an adult cholestatic liver disease. In certain embodiments, the cholestatic liver disease is non-obstructive cholestasis, extrahepatic cholestasis, intrahepatic cholestasis, primary intrahepatic cholestasis, secondary intrahepatic cholestasis, progressive familial intrahepatic cholestasis (PFIC), PFIC type 1, PFIC type 2, PFIC type 3, benign recurrent intrahepatic cholestasis (BRIC), BRIC type 1, BRIC type 2, BRIC type 3, total parenteral nutrition associated cholestasis, paraneoplastic cholestasis, Stauffer syndrome, intrahepatic cholestasis of pregnancy, contraceptive-associated cholestasis, drug-associated cholestasis, infection-associated cholestasis, Dubin-Johnson Syndrome, primary biliary cirrhosis (PBC), primary sclerosing cholangitis (PSC), gallstone disease, Alagille syndrome, biliary atresia, post-Kasai biliary atresia, post-liver transplantation biliary atresia, post-liver transplantation cholestasis, post-liver transplantation associated liver disease, intestinal failure associated liver disease, bile acid mediated liver injury, MRP2 deficiency syndrome, or neonatal sclerosing cholangitis. In certain embodiments, the cholestatic liver disease is Alagille syndrome, PFIC, BRIC, PSC, PBC, or biliary atresia.


[0019] In certain embodiments, the ASBTI is administered to the subject in a fasted state. In certain embodiments, the ASBTI is administered less than about 60 minutes before ingestion of

food. In certain embodiments, the ASBTI is administered less than about 30 minutes before ingestion of food. In certain embodiments, the ASBTI is administered immediately prior to the ingestion of food. In certain embodiments, the ASBTI is administered at least 4 hours after the last meal.


[0020] In certain embodiments, the ASBTI is administered once daily. In certain embodiments, the ASBTI is administered twice daily.

[0021] In certain embodiments, the ASBTI is administered in an amount of about 0.1 mg to about 100 mg per dose. In certain embodiments, the ASBTI is administered in an amount of about 10 mg to about 100 mg per dose. In certain embodiments, the ASBTI is administered in an amount of about 20 mg to about 80 mg per dose. In certain embodiments, the ASBTI is administered in an amount of about 100 ug/kg/day to 1400 ug/kg/day. In certain embodiments, the ASBTI is administered in an amount of about 400 ug/kg/day to 800 ug/kg/day.

[0022] In certain embodiments, the ASBTI is selected from




(maralixibat) and



(volixibat) or a

pharmaceutically acceptable salt thereof. In certain embodiments, the ASBTI is



(maralixibat chloride). In certain embodiments, the ASBTI is

volixibat or a pharmaceutically acceptable salt thereof. In certain embodiments, the ASBTI is volixibat potassium.

[0023] In certain embodiments, the subject has not ingested food for about 0.5 to about 16 hours before the administration of the ASBTI.

[0024] In certain embodiments, the subject is a pediatric subject. In certain embodiments, the pediatric subject is 0 to 18 years of age.

[0025] In certain embodiments, the ASBTI is administered orally.

[0026] In certain embodiments, less than 10% of the ASBTI is systemically absorbed. In certain embodiments, less than 30% of the ASBTI is systemically absorbed.

BRIEF DESCRIPTION OF THE DRAWINGS

[0027] **Figure 1** depicts study designs, drugs (MRX = maralixibat, VLX = volixibat), doses and at mealtime/fasted timelines for certain embodiments according to the disclosure. *Fasting was started ≥ 10 -12 hours before MRX dosing. †Patients were randomized 1:1 to two cohorts and received sequentially fasted then at mealtime dosing, or at mealtime then fasted dosing.

[0028] **Figure 2** is pie charts showing the proportion of healthy participants who experienced gastrointestinal (GI) treatment-emergent adverse events (AEs) following ASBTI administration at mealtime versus in the fasted state for each of the three studies.

[0029] **Figure 3** shows the effect of maralixibat (MRX) on fecal bile acid (fBA) excretion in dogs: study drug, and dosing/mealtime schedule.

[0030] **Figure 4** is a bar graph of the effect of maralixibat (MRX) on fecal bile acid (fBA) in dogs: change in fBA excretion from pre-treatment to day 7.

DETAILED DESCRIPTION

[0031] Detailed embodiments of the present invention are disclosed herein; however, it is to be understood that the disclosed embodiments are merely illustrative of the invention that may be embodied in various forms. In addition, each of the examples given in connection with the various embodiments of the invention is intended to be illustrative, and not restrictive. Therefore, specific structural and functional details disclosed herein are not to be interpreted as limiting, but merely as a representative basis for teaching one skilled in the art to variously employ the present invention.

[0032] Bile acids/salts play a critical role in activating digestive enzymes and solubilizing fats and fat-soluble vitamins and are involved in liver, biliary, and intestinal disease. Bile acids are synthesized in the liver by a multistep, multiorganelle pathway. Hydroxyl groups are added to specific sites on the steroid structure, the double bond of the cholesterol B ring is reduced, and

the hydrocarbon chain is shortened by three carbon atoms resulting in a carboxyl group at the end of the chain. The most common bile acids are cholic acid and chenodeoxycholic acid (the "primary bile acids"). Before exiting the hepatocytes and forming bile, the bile acids are conjugated to either glycine (to produce glycocholic acid or glycochenodeoxycholic acid) or taurine (to produce taurocholic acid or taurochenodeoxycholic acid). The conjugated bile acids are called bile salts and their amphipathic nature makes them more efficient detergents than bile acids. Bile salts, not bile acids, are found in bile.

[0033] Bile salts are excreted by the hepatocytes into the canaliculi to form bile. The canaliculi drain into the right and left hepatic ducts and the bile flows to the gallbladder. Bile is released from the gallbladder and travels to the duodenum, where it contributes to the metabolism and degradation of fat. The bile salts are reabsorbed in the terminal ileum and transported back to the liver via the portal vein. Bile salts often undergo multiple enterohepatic circulations before being excreted via feces. A small percentage of bile salts may be reabsorbed in the proximal intestine by either passive or carrier-mediated transport processes. Most bile salts are reclaimed in the distal ileum by a sodium-dependent apically located bile acid transporter referred to as apical sodium-dependent bile acid transporter (ASBT). At the basolateral surface of the enterocyte, a truncated version of ASBT is involved in vectorial transfer of bile acids/salts into the portal circulation. Completion of the enterohepatic circulation occurs at the basolateral surface of the hepatocyte by a transport process that is primarily mediated by a sodium-dependent bile acid transporter. Intestinal bile acid transport plays a key role in the enterohepatic circulation of bile salts. Molecular analysis of this process has recently led to important advances in understanding of the biology, physiology and pathophysiology of intestinal bile acid transport.

[0034] Within the intestinal lumen, bile acid concentrations vary, with the bulk of the reuptake occurring in the distal intestine. Described herein are certain compositions and methods that control bile acid concentrations in the intestinal lumen, thereby controlling the hepatocellular damage caused by bile acid accumulation in the liver and dosing in the fasted state for minimal gastrointestinal adverse effects.

[0035] The presently disclosed subject matter is based, at least in part, on the discovery that administration of an ASBTI to a subject in need thereof before ingestion of food unexpectedly leads to reduction, minimization, prevention, amelioration, and/or elimination of one or more side effects associated with the administration of the ASBTI.

Classes of Cholestasis and Cholestatic Liver Disease

[0036] As used herein, "cholestasis" means the disease or symptoms comprising impairment of bile formation and/or bile flow. As used herein, "cholestatic liver disease" means a liver disease associated with cholestasis. Cholestatic liver diseases are often associated with jaundice, fatigue, and pruritis. Biomarkers of cholestatic liver disease include elevated serum bile acid concentrations, elevated serum alkaline phosphatase (AP), elevated gamma-glutamyltranspeptidase, elevated conjugated hyperbilirubinemia, and elevated serum cholesterol.

[0037] Cholestatic liver disease can be sorted clinicopathologically between two principal categories of obstructive, often extrahepatic, cholestasis, and nonobstructive, or intrahepatic, cholestasis. In the former, cholestasis results when bile flow is mechanically blocked, as by gallstones or tumor, or as in extrahepatic biliary atresia.

[0038] The latter group who has nonobstructive intrahepatic cholestasis in turn fall into two principal subgroups. In the first subgroup, cholestasis results when processes of bile secretion and modification, or of synthesis of constituents of bile, are caught up secondarily in hepatocellular injury so severe that nonspecific impairment of many functions can be expected, including those subserving bile formation. In the second subgroup, no presumed cause of hepatocellular injury can be identified. Cholestasis in such patients appears to result when one of the steps in bile secretion or modification, or of synthesis of constituents of bile, is constitutively damaged. Such cholestasis is considered primary.

[0039] Accordingly, provided herein are methods and compositions for stimulating epithelial proliferation and/or regeneration of intestinal lining and/or enhancement of the adaptive processes in the intestine in individuals with hypercholeemia and/or a cholestatic liver disease. In some of such embodiments, the methods comprise increasing bile acid concentrations and/or GLP-2 concentrations in the intestinal lumen.

[0040] Increased levels of bile acids, and elevated levels of AP (alkaline phosphatase), LAP (leukocyte alkaline phosphatase), gamma GT (gamma-glutamyl transpeptidase), and 5'-nucleotidase are biochemical hallmarks of cholestasis and cholestatic liver disease. Accordingly, provided herein are methods and compositions for stimulating epithelial proliferation and/or regeneration of intestinal lining and/or enhancement of the adaptive processes in the intestine in individuals with hypercholeemia, and elevated levels of AP (alkaline phosphatase), LAP

(leukocyte alkaline phosphatase), gamma GT (gamma-glutamyl transpeptidase or GGT), and/or 5'-nucleotidase. In some of such embodiments, the methods comprise increasing bile acid concentrations in the intestinal lumen. Further provided herein, are methods and compositions for reducing hypercholeemia, and elevated levels of AP (alkaline phosphatase), LAP (leukocyte alkaline phosphatase), gamma GT (gamma-glutamyl transpeptidase), and 5'-nucleotidase comprising reducing overall serum bile acid load by excreting bile acid in the feces.

[0041] Pruritus is often associated with hypercholeemia and cholestatic liver diseases. It has been suggested that pruritus results from bile salts acting on peripheral pain afferent nerves. The degree of pruritus varies with the individual (i.e., some individuals are more sensitive to elevated levels of bile acids/salts).

[0042] Administration of agents that reduce serum bile acid concentrations has been shown to reduce pruritus in certain individuals. Accordingly, provided herein are methods and compositions for stimulating epithelial proliferation and/or regeneration of intestinal lining and/or enhancement of the adaptive processes in the intestine in individuals with pruritus. In some of such embodiments, the methods comprise increasing bile acid concentrations in the intestinal lumen. Further provided herein, are methods and compositions for treating pruritus comprising reducing overall serum bile acid load by excreting bile acid in the feces.

[0043] Another symptom of hypercholeemia and cholestatic liver disease is the increase in serum concentration of conjugated bilirubin. Elevated serum concentrations of conjugated bilirubin result in jaundice and dark urine. The magnitude of elevation is not diagnostically important as no relationship has been established between serum levels of conjugated bilirubin and the severity of hypercholeemia and cholestatic liver disease. Conjugated bilirubin concentration rarely exceeds 30 mg/dL. Accordingly, provided herein are methods and compositions for stimulating epithelial proliferation and/or regeneration of intestinal lining and/or enhancement of the adaptive processes in the intestine in individuals with elevated serum concentrations of conjugated bilirubin. In some of such embodiments, the methods comprise increasing bile acid concentrations in the intestinal lumen. Further provided herein, are methods and compositions for treating elevated serum concentrations of conjugated bilirubin comprising reducing overall serum bile acid load by excreting bile acid in the feces.

[0044] Increased serum concentration of nonconjugated bilirubin is also considered diagnostic of hypercholeemia and cholestatic liver disease. Portions of serum bilirubin and covalently bound

to albumin (delta bilirubin or biliprotein). This fraction may account for a large proportion of total bilirubin in patients with cholestatic jaundice. The presence of large quantities of delta bilirubin indicates long-standing cholestasis. Delta bilirubin in cord blood or the blood of a newborn is indicative of cholestasis/cholestatic liver disease that antedates birth. Accordingly, provided herein are methods and compositions for stimulating epithelial proliferation and/or regeneration of intestinal lining and/or enhancement of the adaptive processes in the intestine in individuals with elevated serum concentrations of nonconjugated bilirubin or delta bilirubin. In some of such embodiments, the methods comprise increasing bile acid concentrations in the intestinal lumen. Further provided herein, are methods and compositions for treating elevated serum concentrations of nonconjugated bilirubin and delta bilirubin comprising reducing overall serum bile acid load by excreting bile acid in the feces.

[0045] Cholestasis and cholestatic liver disease results in hypercholelism. During metabolic cholestasis, the hepatocytes retains bile salts. Bile salts are regurgitated from the hepatocyte into the serum, which results in an increase in the concentration of bile salts in the peripheral circulation. Furthermore, the uptake of bile salts entering the liver in portal vein blood is inefficient, which results in spillage of bile salts into the peripheral circulation. Accordingly, provided herein are methods and compositions for stimulating epithelial proliferation and/or regeneration of intestinal lining and/or enhancement of the adaptive processes in the intestine in individuals with hypercholelism. In some of such embodiments, the methods comprise increasing bile acid concentrations in the intestinal lumen. Further provided herein, are methods and compositions for treating hypercholelism comprising reducing overall serum bile acid load by excreting bile acid in the feces.

[0046] Hyperlipidemia is characteristic of some but not all cholestatic diseases. Serum cholesterol is elevated in cholestasis due to the decrease in circulating bile salts which contribute to the metabolism and degradation of cholesterol. Cholesterol retention is associated with an increase in membrane cholesterol content and a reduction in membrane fluidity and membrane function. Furthermore, as bile salts are the metabolic products of cholesterol, the reduction in cholesterol metabolism results in a decrease in bile acid/salt synthesis. Serum cholesterol observed in children with cholestasis ranges between about 1,000 mg/dL and about 4,000 mg/dL. Accordingly, provided herein are methods and compositions for stimulating epithelial proliferation and/or regeneration of intestinal lining and/or enhancement of the adaptive

processes in the intestine in individuals with hyperlipidemia. In some of such embodiments, the methods comprise increasing bile acid concentrations in the intestinal lumen. Further provided herein, are methods and compositions for treating hyperlipidemia comprising reducing overall serum bile acid load by excreting bile acid in the feces.

[0047] In individuals with hypercholeemia and cholestatic liver diseases, xanthomas develop from the deposition of excess circulating cholesterol into the dermis. The development of xanthomas is more characteristic of obstructive cholestasis than of hepatocellular cholestasis. Planar xanthomas first occur around the eyes and then in the creases of the palms and soles, followed by the neck. Tuberous xanthomas are associated with chronic and long-term cholestasis. Accordingly, provided herein are methods and compositions for stimulating epithelial proliferation and/or regeneration of intestinal lining and/or enhancement of the adaptive processes in the intestine in individuals with xanthomas. In some of such embodiments, the methods comprise increasing bile acid concentrations in the intestinal lumen. Further provided herein, are methods and compositions for treating xanthomas comprising reducing overall serum bile acid load by excreting bile acid in the feces.

[0048] In children with chronic cholestasis, one of the major consequences of hypercholeemia and cholestatic liver disease is failure to thrive. Failure to thrive is a consequence of reduced delivery of bile salts to the intestine, which contributes to inefficient digestion and absorption of fats, and reduced uptake of vitamins (vitamins E, D, K, and A are all malabsorbed in cholestasis). Furthermore, the delivery of fat into the colon can result in colonic secretion and diarrhea. Treatment of failure to thrive involves dietary substitution and supplementation with long-chain triglycerides, medium-chain triglycerides, and vitamins. Ursodeoxycholic acid, which is used to treat some cholestatic conditions, does not form mixed micelles and has no effect on fat absorption. Accordingly, provided herein are methods and compositions for stimulating epithelial proliferation and/or regeneration of intestinal lining and/or enhancement of the adaptive processes in the intestine in individuals (e.g., children) with failure to thrive. In some of such embodiments, the methods comprise increasing bile acid concentrations in the intestinal lumen. Further provided herein, are methods and compositions for treating failure to thrive comprising reducing overall serum bile acid load by excreting bile acid in the feces.

Primary Biliary Cirrhosis (PBC)

[0049] Primary biliary cirrhosis is an autoimmune disease of the liver characterized by the destruction of the bile canaliculi. Damage to the bile canaliculi results in the build-up of bile in the liver (i.e., cholestasis). The retention of bile in the liver damages liver tissue and may lead to scarring, fibrosis, and cirrhosis. PBC usually presents in adulthood (e.g., ages 40 and over). Individuals with PBC often present with fatigue, pruritus, and/or jaundice. PBC is diagnosed if the individual has elevated AP concentrations for at least 6 months, elevated gammaGT levels, antimitochondrial antibodies (AMA) in the serum (>1 :40), and florid bile duct lesions. Serum ALT and serum AST and conjugated bilirubin may also be elevated, but these are not considered diagnostic. Cholestasis associated with PBC has been treated or ameliorated by administration of ursodeoxycholic acid (UDCA or Ursodiol). Corticosteroids (e.g., prednisone and budesonide) and immunosuppressive agents (e.g., azathioprine, cyclosporin A, methotrexate, chlorambucil and mycophenolate) have been used to treat cholestasis associated with PBC. Sulindac, bezafibrate, tamoxifen, and lamivudine have also been shown to treat or ameliorate cholestasis associated with PBC.

Progressive Familial Intrahepatic Cholestasis (PFIC)

PFIC is a rare genetic disorder that causes progressive liver disease typically leading to liver failure. In people with PFIC, liver cells are less able to secrete bile. The resulting buildup of bile causes liver disease in affected individuals. Signs and symptoms of PFIC typically begin in infancy. Patients experience severe itching, jaundice, failure to grow at the expected rate (failure to thrive), and an increasing inability of the liver to function (liver failure). The disease is estimated to affect one in every 50,000 to 100,000 births in the United States and Europe. Six types of PFIC have been genetically identified, all of which are similarly characterized by impaired bile flow and progressive liver disease.

PFIC 1

[0050] PFIC 1 (also known as, Byler disease or FIC1 deficiency) is associated with mutations in the ATP8B1 gene (also designated as FIC1). This gene, which encodes a P-type ATPase, is located on human chromosome 18 and is also mutated in the milder phenotype, benign recurrent intrahepatic cholestasis type 1 (BRIO) and in Greenland familial cholestasis. FIC1 protein is located on the canalicular membrane of the hepatocyte but within the liver it is mainly expressed in cholangiocytes. P-type ATPase appears to be an aminophospholipid transporter responsible

for maintaining the enrichment of phosphatidylserine and phosphatidylethanolamine on the inner leaflet of the plasma membrane in comparison of the outer leaflet. The asymmetric distribution of lipids in the membrane bilayer plays a protective role against high bile salt concentrations in the canalicular lumen. The abnormal protein function may indirectly disturb the biliary secretion of bile acids. The anomalous secretion of bile acids/salts leads to hepatocyte bile acid overload.

[0051] PFIC 1 typically presents in infants (e.g., age 6-18 months). The infants may show signs of pruritus, jaundice, abdominal distension, diarrhea, malnutrition, and shortened stature. Biochemically, individuals with PFIC 1 have elevated serum transaminases, elevated bilirubin, elevated serum bile acid levels, and low levels of gammaGT. The individual may also have liver fibrosis. Individuals with PFIC 1 typically do not have bile duct proliferation. Most individuals with PFIC 1 will develop end-stage liver disease by 10 years of age. No medical treatments have proven beneficial for the long-term treatment of PFIC 1. In order to reduce extrahepatic symptoms (e.g., malnutrition and failure to thrive), children are often administered medium chain triglycerides and fat-soluble vitamins. Ursodiol has not been demonstrated as effective in individuals with PFIC 1.

PFIC 2

[0052] PFIC 2 (also known as, Byler Syndrome or BSEP deficiency) is associated with mutations in the ABCB11 gene (also designated BSEP). The ABCB11 gene encodes the ATP-dependent canalicular bile salt export pump (BSEP) of human liver and is located on human chromosome 2. BSEP protein, expressed at the hepatocyte canalicular membrane, is the major exporter of primary bile acids/salts against extreme concentration gradients. Mutations in this protein responsible for the decreased biliary bile salt secretion described in affected patients, leading to decreased bile flow and accumulation of bile salts inside the hepatocyte with ongoing severe hepatocellular damage.

[0053] PFIC 2 typically presents in infants (e.g., age 6-18 months). The infants may show signs of pruritus. Biochemically, individuals with PFIC 2 have elevated serum transaminases, elevated bilirubin, elevated serum bile acid levels, and low levels of gammaGT. The individual may also have portal inflammation and giant cell hepatitis. Further, individuals often develop hepatocellular carcinoma. No medical treatments have proven beneficial for the long-term treatment of PFIC 2. In order to reduce extrahepatic symptoms (e.g., malnutrition and failure to

thrive), children are often administered medium chain triglycerides and fat- soluble vitamins. The PFIC 2 patient population accounts for approximately 60% of the PFIC population.

PFIC 3

[0054] PFIC 3 (also known as MDR3 deficiency) is caused by a genetic defect in the ABCB4 gene (also designated MDR3) located on chromosome 7. Class III Multidrug Resistance (MDR3) P-glycoprotein (P-gp), is a phospholipid translocator involved in biliary phospholipid (phosphatidylcholine) excretion in the canicular membrane of the hepatocyte. PFIC 3 results from the toxicity of bile in which detergent bile salts are not inactivated by phospholipids, leading to bile canaliculi and biliary epithelium injuries.

[0055] PFIC 3 also presents in early childhood. As opposed to PFIC 1 and PFIC 2, individuals have elevated gammaGT levels. Individuals also have portal inflammation, fibrosis, cirrhosis, and massive bile duct proliferation. Individuals may also develop intrahepatic gallstone disease. Ursodiol has been effective in treating or ameliorating PFIC 3.

Benign Recurrent Intrahepatic Cholestasis (BRIC)

BRIC 1

[0056] BRIC1 is caused by a genetic defect of the FIC1 protein in the canicular membrane of hepatocytes. BRIC1 is typically associated with normal serum cholesterol and γ -glutamyltranspeptidase levels, but elevated serum bile salts. Residual FIC1 expression and function is associated with BRIC1. Despite recurrent attacks of cholestasis or cholestatic liver disease, there is no progression to chronic liver disease in a majority of patients. During the attacks, the patients are severely jaundiced and have pruritis, steatorrhea, and weight loss. Some patients also have renal stones, pancreatitis, and diabetes.

BRIC 2

[0057] BRIC2 is caused by mutations in ABCB11, leading to defective BSEP expression and/or function in the canicular membrane of hepatocytes.

BRIC 3

[0058] BRIC3 is related to the defective expression and/or function of MDR3 in the canicular membrane of hepatocytes. Patients with MDR3 deficiency usually display elevated serum γ -glutamyltranspeptidase levels in the presence of normal or slightly elevated bile acid levels.

Dubin-Johnson Syndrome (DJS)

[0059] DJS is characterized by conjugated hyperbilirubinemia due to inherited dysfunction of MRP2. Hepatic function is preserved in affected patients. Several different mutations have been associated with this condition, resulting either in the complete absence of immunohistochemically detectable MRP2 in affected patients or impaired protein maturation and sorting.

*Acquired Cholestatic Disease*Primary Biliary Cirrhosis (PBC)

[0060] PBC is a chronic inflammatory hepatic disorder slowly progressing to end stage liver failure in most of the affected patients. In PBC, the inflammatory process affects predominantly the small bile ducts.

Primary Sclerosing Cholangitis (PSC)

[0061] PSC is a chronic inflammatory hepatic disorder slowly progressing to end stage liver failure in most of the affected patients. In PSC inflammation, fibrosis and obstruction of large and medium sized intra- and extrahepatic ductuli is predominant.

[0062] PSC is characterized by progressive cholestasis. Cholestasis can often lead to severe pruritus which significantly impairs quality of life.

Intrahepatic Cholestasis of Pregnancy (ICP)

[0063] ICP is characterized by occurrence of transient cholestasis or cholestatic liver disease in pregnant women typically occurring in the third trimester of pregnancy, when the circulating levels of estrogens are high. ICP is associated with pruritis and biochemical cholestasis or cholestatic liver disease of varying severity and constitutes a risk factor for prematurity and intrauterine fetal death. A genetic predisposition has been suspected based upon the strong regional clustering, the higher prevalence in female family members of patients with ICP and the susceptibility of ICP patients to develop intrahepatic cholestasis or cholestatic liver disease under other hormonal challenges such as oral contraception. A heterogeneous state for an MDR3 gene defect may represent a genetic predisposition.

Gallstone disease

[0064] Gallstone disease is one of the most common and costly of all digestive diseases with a prevalence of up to 17% in Caucasian women. Cholesterol containing gallstones are the major form of gallstones and supersaturation of bile with cholesterol is therefore a prerequisite for

gallstone formation. ABCB4 mutations may be involved in the pathogenesis of cholesterol gallstone disease.

Drug induced cholestasis

[0065] Inhibition of BSEP function by drugs is an important mechanism of drug-induced cholestasis, leading to the hepatic accumulation of bile salts and subsequent liver cell damage. Several drugs have been implicated in BSEP inhibition. Most of these drugs, such as rifampicin, cyclosporine, glibenclamide, or troglitazone directly cis-inhibit ATP-dependent taurocholate transport in a competitive manner, while estrogen and progesterone metabolites indirectly trans-inhibits BSEP after secretion into the bile canaliculus by Mrp2. Alternatively, drug-mediated stimulation of MRP2 can promote cholestasis or cholestatic liver disease by changing bile composition.

Total parenteral nutrition associated cholestasis

[0066] TPNAC is one of the most serious clinical scenarios where cholestasis or cholestatic liver disease occurs rapidly and is highly linked with early death. Infants, who are usually premature and who have had gut resections are dependent upon TPN for growth and frequently develop cholestasis or cholestatic liver disease that rapidly progresses to fibrosis, cirrhosis, and portal hypertension, usually before 6 months of life. The degree of cholestasis or cholestatic liver disease and chance of survival in these infants have been linked to the number of septic episodes, likely initiated by recurrent bacterial translocation across their gut mucosa. Although there are also cholestatic effects from the intravenous formulation in these infants, septic mediators likely contribute the most to altered hepatic function.

Total Parenteral Nutrition Associated Cholestasis

[0067] TPNAC is one of the most serious clinical scenarios where cholestasis or cholestatic liver disease occurs rapidly and is highly linked with early death. Infants, who are usually premature and who have had gut resections are dependent upon TPN for growth and frequently develop cholestasis or cholestatic liver disease that rapidly progresses to fibrosis, cirrhosis, and portal hypertension, usually before 6 months of life. The degree of cholestasis or cholestatic liver disease and chance of survival in these infants have been linked to the number of septic episodes, likely initiated by recurrent bacterial translocation across their gut mucosa. Although there are also cholestatic effects from the intravenous formulation in these infants, septic mediators likely contribute the most to altered hepatic function.

Alagille syndrome (ALGS)

[0068] Alagille syndrome is a genetic disorder that affects the liver and other organs. ALGS is also known as syndromic intrahepatic bile duct paucity or arteriohepatic dysplasia. ALGS is a rare genetic disorder in which bile ducts are abnormally narrow, malformed, and reduced in number, which leads to bile accumulation in the liver and ultimately progressive liver disease. ALGS is autosomal dominant, caused by mutations in JAG1 (> 90% of cases) or NOTCH2. The estimated incidence of ALGS is one in every 30,000 or 50,000 births in the United States and Europe. In patients with ALGS, multiple organ systems may be affected by the mutation, including the liver, heart, kidneys and central nervous system. The accumulation of bile acids prevents the liver from working properly to eliminate waste from the bloodstream and leads to progressive liver disease that ultimately requires liver transplantation in 15% to 47% of patients. Signs and symptoms arising from liver damage in ALGS may include jaundice, pruritus and xanthomas, and decreased growth. The pruritus experienced by patients with ALGS is among the most severe in any chronic liver disease and is present in most affected children by the third year of life.

[0069] ALGS often presents during infancy (e.g., age 6-18 months) through early childhood (e.g., age 3-5 years) and may stabilize after the age of 10. Symptoms may include chronic progressive cholestasis, ductopenia, jaundice, pruritus, xanthomas, congenital heart problems, paucity of intrahepatic bile ducts, poor linear growth, hormone resistance, posterior embryotoxon, Axenfeld anomaly, retinitis pigmentosa, pupillary abnormalities, cardiac murmur, atrial septal defect, ventricular septal defect, patent ductus arteriosus, and Tetralogy of Fallot. Individuals diagnosed with Alagille syndrome have been treated with ursodiol, hydroxyzine, cholestyramine, rifampicin, and phenobarbital. Due to a reduced ability to absorb fat-soluble vitamins, individuals with Alagille Syndrome are further administered high dose multivitamins.

Biliary atresia

[0070] Biliary atresia is a life-threatening condition in infants in which the bile ducts inside or outside the liver do not have normal openings. With biliary atresia, bile becomes trapped, builds up, and damages the liver. The damage leads to scarring, loss of liver tissue, and cirrhosis. Without treatment, the liver eventually fails, and the infant needs a liver transplant to stay alive. The two types of biliary atresia are fetal and perinatal. Fetal biliary atresia appears while the

baby is in the womb. Perinatal biliary atresia is much more common and does not become evident until 2 to 4 weeks after birth.

Post-Kasai biliary atresia

[0071] Biliary atresia is treated with surgery called the Kasai procedure or a liver transplant. The Kasai procedure is usually the first treatment for biliary atresia. During a Kasai procedure, the pediatric surgeon removes the infant's damaged bile ducts and brings up a loop of intestine to replace them. While the Kasai procedure can restore bile flow and correct many problems caused by biliary atresia, the surgery doesn't cure biliary atresia. If the Kasai procedure is not successful, infants usually need a liver transplant within 1 to 2 years. Even after a successful surgery, most infants with biliary atresia slowly develop cirrhosis over the years and require a liver transplant by adulthood. Possible complications after the Kasai procedure include ascites, bacterial cholangitis, portal hypertension, and pruritis.

Post liver transplantation biliary atresia

[0072] If the atresia is complete, liver transplantation is the only option. Although liver transplantation is generally successful at treating biliary atresia, liver transplantation may have complications such as organ rejection. Also, a donor liver may not become available. Further, in some patients, liver transplantation may not be successful at curing biliary atresia.

Xanthoma

[0073] Xanthoma is a skin condition associated with cholestatic liver diseases, in which certain fats build up under the surface of the skin. Cholestasis results in several disturbances of lipid metabolism resulting in formation of an abnormal lipid particle in the blood called lipoprotein X. Lipoprotein X is formed by regurgitation of bile lipids into the blood from the liver and does not bind to the LDL receptor to deliver cholesterol to cells throughout the body as does normal LDL. Lipoprotein X increases liver cholesterol production by fivefold and blocks normal removal of lipoprotein particles from the blood by the liver.

General Definitions

[0074] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

[0075] As used in this specification and the appended claims, the singular forms “a”, “an”, and “the” include plural references unless the context clearly dictates otherwise. Thus, for example, a

reference to “a method” includes one or more methods, and/or steps of the type described herein and/or which will become apparent to those persons skilled in the art upon reading this disclosure.

[0076] The term “baseline” or “pre-administration baseline,” as used herein, refers to information gathered at the beginning of a study or an initial known value which is used for comparison with later data. A baseline is an initial measurement of a measurable condition that is taken at an early time point and used for comparison over time to look for changes in the measurable condition. For example, serum bile acid concentration in a patient before administration of a drug (baseline) and after administration of the drug. Baseline is an observation or value that represents the normal or beginning level of a measurable quality, used for comparison with values representing response to intervention or an environmental stimulus. The baseline is time “zero”, before participants in a study receive an experimental agent or intervention, or negative control. For example, “baseline” may refer in some instances 1) to the state of a measurable quantity just prior to the initiation of a clinical study or 2) the state of a measurable quantity just prior to altering a dosage level or composition administered to a patient from a first dosage level or composition to a second dosage level or composition.

[0077] The terms “level” and “concentration,” as used herein, are used interchangeably. For example, “high serum levels of bilirubin” may alternatively be phrased “high serum concentrations of bilirubin.”

[0078] The terms “normalized” or “normal range,” as used herein, indicates age-specific values that are within a range corresponding to a healthy individual (i.e., normal or normalized values). For example, the phrase “serum bilirubin concentrations were normalized within three weeks” means that serum bilirubin concentrations fell within a range known in the art to correspond to that of a healthy individual (i.e., within a normal and not e.g. an elevated range) within three weeks. In various embodiments, a normalized serum bilirubin concentration is from about 0.1 mg/dL to about 1.2 mg/dL. In various embodiments, a normalized serum bile acid concentration is from about 0 $\mu\text{mol/L}$ to about 25 $\mu\text{mol/L}$.

[0079] The terms “ITCHRO(OBS)” and “ITCHRO” (alternatively, “ItchRO(Pt)”) as used herein, are used interchangeably with the qualification that the ITCHRO(OBS) scale is used to measure severity of pruritus in children under the age of 18 and the ITCHRO scale is used to measure severity of pruritus in adults of at least 18 years of age. Therefore, where

ITCHRO(OBS) scale is mentioned with regard to an adult patient, the ITCHRO scale is the scale being indicated. Similarly, whenever the ITCHRO scale is mentioned with regard to a pediatric patient, the ITCHRO(OBS) scale is usually the scale being indicated (some older children were permitted to report their own scores as ITCHRO scores. The ITCHRO(OBS) scale ranges from 0 to 4 and the ITCHRO scale ranges from 0 to 10.

[0080] The term "bile acid" or "bile acids," as used herein, includes steroid acids (and/or the carboxylate anion thereof), and salts thereof, found in the bile of an animal (e.g., a human), including, by way of non-limiting example, cholic acid, cholate, deoxycholic acid, deoxycholate, hyodeoxycholic acid, hyodeoxycholate, glycocholic acid, glycocholate, taurocholic acid, taurocholate, chenodeoxycholic acid, ursodeoxycholic acid, ursodiol, a tauroursodeoxycholic acid, a glyoursodeoxycholic acid, a 7-B-methyl cholic acid, a methyl lithocholic acid, chenodeoxycholate, lithocholic acid, lithocate, and the like. Taurocholic acid and/or taurocholate are referred to herein as TCA. Any reference to a bile acid used herein includes reference to a bile acid, one and only one bile acid, one or more bile acids, or to at least one bile acid. Therefore, the terms "bile acid," "bile salt," "bile acid/salt," "bile acids," "bile salts," and "bile acids/salts" are, unless otherwise indicated, utilized interchangeably herein. Any reference to a bile acid used herein includes reference to a bile acid or a salt thereof. Furthermore, pharmaceutically acceptable bile acid esters are optionally utilized as the "bile acids" described herein, e.g., bile acids/salts conjugated to an amino acid (e.g., glycine or taurine). Other bile acid esters include, e.g., substituted or unsubstituted alkyl ester, substituted or unsubstituted heteroalkyl esters, substituted or unsubstituted aryl esters, substituted or unsubstituted heteroaryl esters, or the like. For example, the term "bile acid" includes cholic acid conjugated with either glycine or taurine: glycocholate and taurocholate, respectively (and salts thereof). Any reference to a bile acid used herein includes reference to an identical compound naturally or synthetically prepared. Furthermore, it is to be understood that any singular reference to a component (bile acid or otherwise) used herein includes reference to one and only one, one or more, or at least one of such components. Similarly, any plural reference to a component used herein includes reference to one and only one, one or more, or at least one of such components, unless otherwise noted.

[0081] The term "subject", "patient", "participant", or "individual" are used interchangeably herein and refer to mammals and non-mammals, e.g., suffering from a disorder described herein.

Examples of mammals include, but are not limited to, any member of the mammalian class: humans, non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. Examples of non-mammals include, but are not limited to, birds, fish and the like. In one embodiment of the methods and compositions provided herein, the mammal is a human.

[0082] The term "about," as used herein, includes any value that is within 10% of the described value.

[0083] The term "composition," as used herein includes the disclosure of both a composition and a composition administered in a method as described herein. Furthermore, in some embodiments, the composition of the present invention is or comprises a "formulation," an oral dosage form or a rectal dosage form as described herein.

[0084] The terms "treat," "treating" or "treatment," and other grammatical equivalents as used herein, include alleviating, inhibiting or reducing symptoms, reducing or inhibiting severity of, reducing incidence of, reducing or inhibiting recurrence of, delaying onset of, delaying recurrence of, abating or ameliorating a disease or condition symptoms, ameliorating the underlying causes of symptoms, inhibiting the disease or condition, e.g., arresting the development of the disease or condition, relieving the disease or condition, causing regression of the disease or condition, relieving a condition caused by the disease or condition, or stopping the symptoms of the disease or condition. The terms further include achieving a therapeutic benefit. By therapeutic benefit is meant eradication or amelioration of the underlying disorder being treated, and/or the eradication or amelioration of one or more of the physiological symptoms associated with the underlying disorder such that an improvement is observed in the patient.

[0085] The terms "effective amount" or "therapeutically effective amount" as used herein, refer to a sufficient amount of at least one agent (e.g., a therapeutically active agent) being administered which achieve a desired result in a subject or individual, e.g., to relieve to some extent one or more symptoms of a disease or condition being treated. In certain instances, the result is a reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. In certain instances, an "effective amount" for therapeutic uses is the amount of the composition comprising an agent as set forth herein required to provide a clinically significant decrease in a disease. An appropriate "effective"

amount in any individual case is determined using any suitable technique, such as a dose escalation study. In some embodiments, a "therapeutically effective amount," or an "effective amount" of an ASBTI refers to a sufficient amount of an ASBTI to treat cholestasis or a cholestatic liver disease in a subject or individual.

[0086] The terms "administer," "administering," "administration," and the like, as used herein, refer to the methods that may be used to enable delivery of agents or compositions to the desired site of biological action. These methods include, but are not limited to oral routes, intraduodenal routes, parenteral injection (including intravenous, subcutaneous, intraperitoneal, intramuscular, intravascular or infusion), topical and rectal administration. Administration techniques that are optionally employed with the agents and methods described herein are found in sources e.g., Goodman and Gilman, *The Pharmacological Basis of Therapeutics*, current ed.; Pergamon; and Remington's, *Pharmaceutical Sciences* (current edition), Mack Publishing Co., Easton, Pa, all of which are incorporated herein by reference in their entirety for all purposes. In certain embodiments, the agents and compositions described herein are administered orally.

[0087] The term "ASBT inhibitor" refers to a compound that inhibits apical sodium-dependent bile transport or any recuperative bile salt transport. The term Apical Sodium-dependent Bile Transporter (ASBT) is used interchangeably with the term Ileal Bile Acid Transporter (IBAT).

[0088] The phrase "pharmaceutically acceptable", as used in connection with compositions of the invention, refers to molecular entities and other ingredients of such compositions that are physiologically tolerable and do not typically produce untoward reactions when administered to a mammal (e.g., a human). Preferably, as used herein, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in mammals, and more particularly in humans.

[0089] In various embodiments, pharmaceutically acceptable salts described herein include, by way of non-limiting example, a nitrate, chloride, bromide, phosphate, sulfate, acetate, hexafluorophosphate, citrate, gluconate, benzoate, propionate, butyrate, subsalicylate, maleate, laurate, malate, fumarate, succinate, tartrate, amsonate, pamoate, p-toluenesulfonate, mesylate and the like. Furthermore, pharmaceutically acceptable salts include, by way of non-limiting example, alkaline earth metal salts (e.g., calcium or magnesium), alkali metal salts (e.g., sodium-dependent or potassium), ammonium salts and the like.

[0090] As used herein the term “fasted state” is defined as a state one in which a subject has completely digested and absorbed the last meal, and the subject’s insulin levels are at a low or baseline level. In some embodiments, a fasted state is defined as a state of not consuming any food for at least 4 hours for a subject 18 years or over. In some embodiments, a fasted state is defined as a state of not consuming any food for at least 2 hours for a pediatric subject. In some embodiments, a fasted state is defined as a state of about 30 minutes before a meal.

[0091] As used herein, a fasted patient is defined as a patient who has not eaten any food, i.e., has fasted for at least 4 hours before the administration of the ASBTI (for a subject 18 years or over) or at least 2 hours before the administration of the ASBTI (for a pediatric subject), and at least 30 minutes after the administration of the ASBTI. The ASBTI is optionally administered with water during the fasting period, and water can be allowed ad libitum.

[0092] As used herein, tolerability refers to the degree to which adverse effects of a drug can be tolerated by a patient. In certain embodiments, gastrointestinal (GI) tolerability refers to the degree to which a patient can tolerate GI adverse effects. In some embodiments, an improvement in GI tolerability comprises a reduction, minimization, prevention, amelioration, or elimination of one or more GI adverse effects.

Bile Acid

[0093] Bile contains water, electrolytes and a numerous organic molecules including bile acids, cholesterol, phospholipids and bilirubin. Bile is secreted from the liver and stored in the gall bladder, and upon gall bladder contraction, due to ingestion of a fatty meal, bile passes through the bile duct into the intestine. Bile acids/salts are critical for digestion and absorption of fats and fat-soluble vitamins in the small intestine. Adult humans produce 400 to 800 mL of bile daily. The secretion of bile can be considered to occur in two stages. Initially, hepatocytes secrete bile into canaliculi, from which it flows into bile ducts and this hepatic bile contains large quantities of bile acids, cholesterol and other organic molecules. Then, as bile flows through the bile ducts, it is modified by addition of a watery, bicarbonate -rich secretion from ductal epithelial cells. Bile is concentrated, typically five-fold, during storage in the gall bladder.

[0094] The flow of bile is lowest during fasting, and a majority of that is diverted into the gallbladder for concentration. When chyme from an ingested meal enters the small intestine, acid and partially digested fats and proteins stimulate secretion of cholecystokinin and secretin, both of which are important for secretion and flow of bile. Cholecystokinin (cholecysto = gallbladder

and kinin = movement) is a hormone which stimulates contractions of the gallbladder and common bile duct, resulting in delivery of bile into the gut. The most potent stimulus for release of cholecystokinin is the presence of fat in the duodenum. Secretin is a hormone secreted in response to acid in the duodenum, and it stimulates biliary duct cells to secrete bicarbonate and water, which expands the volume of bile and increases its flow out into the intestine.

[0095] Bile acids/salts are derivatives of cholesterol. Cholesterol, ingested as part of the diet or derived from hepatic synthesis, are converted into bile acids/salts in the hepatocyte. Examples of such bile acids/salts include cholic and chenodeoxycholic acids, which are then conjugated to an amino acid (such as glycine or taurine) to yield the conjugated form that is actively secreted into canaliculi. The most abundant of the bile salts in humans are cholate and deoxycholate, and they are normally conjugated with either glycine or taurine to give glycocholate or taurocholate respectively.

[0096] Free cholesterol is virtually insoluble in aqueous solutions, however in bile it is made soluble by the presence of bile acids/salts and lipids. Hepatic synthesis of bile acids/salts accounts for the majority of cholesterol breakdown in the body. In humans, roughly 500 mg of cholesterol are converted to bile acids/salts and eliminated in bile every day. Therefore, secretion into bile is a major route for elimination of cholesterol. Large amounts of bile acids/salts are secreted into the intestine every day, but only relatively small quantities are lost from the body. This is because approximately 95% of the bile acids/salts delivered to the duodenum are absorbed back into blood within the ileum, by a process known as "Enterohepatic Recirculation".

[0097] Venous blood from the ileum goes straight into the portal vein, and hence through the sinusoids of the liver. Hepatocytes extract bile acids/salts very efficiently from sinusoidal blood, and little escapes the healthy liver into systemic circulation. Bile acids/salts are then transported across the hepatocytes to be resecreted into canaliculi. The net effect of this enterohepatic recirculation is that each bile salt molecule is reused about 20 times, often two or three times during a single digestive phase. Bile biosynthesis represents the major metabolic fate of cholesterol, accounting for more than half of the approximate 800 mg/day of cholesterol that an average adult uses up in metabolic processes. In comparison, steroid hormone biosynthesis consumes only about 50 mg of cholesterol per day. Much more than 400 mg of bile salts is required and secreted into the intestine per day, and this is achieved by re-cycling the bile salts.

Most of the bile salts secreted into the upper region of the small intestine are absorbed along with the dietary lipids that they emulsified at the lower end of the small intestine. They are separated from the dietary lipid and returned to the liver for re-use. Recycling thus enables 20-30 g of bile salts to be secreted into the small intestine each day.

[0098] Bile acids/salts are amphipathic, with the cholesterol-derived portion containing both hydrophobic (lipid soluble) and polar (hydrophilic) moieties while the amino acid conjugate is generally polar and hydrophilic. This amphipathic nature enables bile acids/salts to carry out two important functions: emulsification of lipid aggregates and solubilization and transport of lipids in an aqueous environment. Bile acids/salts have detergent action on particles of dietary fat which causes fat globules to break down or to be emulsified. Emulsification is important since it greatly increases the surface area of fat available for digestion by lipases which cannot access the inside of lipid droplets. Furthermore, bile acids/salts are lipid carriers and are able to solubilize many lipids by forming micelles and are critical for transport and absorption of the fat-soluble vitamins.

[0099] The term “non-systemic” or “minimally absorbed,” as used herein, refers to low systemic bioavailability and/or absorption of an administered compound. In some embodiments a non-systemic compound is a compound that is substantially not absorbed systemically. In some embodiments, ASBTI compositions described herein deliver the ASBTI to the distal ileum, colon, and/or rectum and not systemically (e.g., a substantial portion of the ASBTI is not systemically absorbed. In some embodiments, the systemic absorption of a non-systemic compound is <0.1%, <0.3%, <0.5%, <0.6%, <0.7%, <0.8%, <0.9%, <1%, <1.5%, <2%, <3%, or < 5 % of the administered dose (wt. % or mol %). In some embodiments, the systemic absorption of a non-systemic compound is < 10 % of the administered dose. In some embodiments, the systemic absorption of a non-systemic compound is < 15 % of the administered dose. In some embodiments, the systemic absorption of a non-systemic compound is < 25% of the administered dose. In an alternative approach, a non-systemic ASBTI is a compound that has lower systemic bioavailability relative to the systemic bioavailability of a systemic ASBTI (e.g., compound 100A, 100C). In some embodiments, the bioavailability of a non-systemic ASBTI described herein is < 30%, < 40%, < 50%, < 60%, or < 70% of the bioavailability of a systemic ASBTI (e.g., compound 100A, 100C).

[00100] In another alternative approach, compositions described herein are formulated to deliver < 10 % of the administered dose of the ASBTI systemically. In some embodiments, the compositions described herein are formulated to deliver < 20 % of the administered dose of the ASBTI systemically. In some embodiments, the compositions described herein are formulated to deliver < 30 % of the administered dose of the ASBTI systemically. In some embodiments, the compositions described herein are formulated to deliver < 40 % of the administered dose of the ASBTI systemically. In some embodiments, the compositions described herein are formulated to deliver < 50 % of the administered dose of the ASBTI systemically. In some embodiments, the compositions described herein are formulated to deliver < 60 % of the administered dose of the ASBTI systemically. In some embodiments, the compositions described herein are formulated to deliver < 70 % of the administered dose of the ASBTI systemically. In some embodiments, systemic absorption is determined in any suitable manner, including the total circulating amount, the amount cleared after administration, or the like.


ASBT Inhibitors

[00101] In various embodiments of methods of the present invention, ASBT inhibitors are administered to a subject. ASBT inhibitors (ASBTIs) reduce or inhibit bile acid recycling in the distal gastrointestinal (GI) tract, including the distal ileum, the colon and/or the rectum. Inhibition of the apical sodium-dependent bile acid transport interrupts the enterohepatic circulation of bile acids and results in more bile acids being excreted in the feces, leading to lower levels of bile acids systemically, thereby reducing bile acid mediated liver damage and related effects and complications. In certain embodiments, the ASBTIs are systemically absorbed. In certain embodiments, the ASBTIs are not systemically absorbed. In some embodiments, ASBTIs described herein are modified or substituted to be non-systemic.


[00102] In certain embodiments, compounds described herein have one or more chiral centers. As such, all stereoisomers are envisioned herein. In various embodiments, compounds described herein are present in optically active or racemic forms. It is to be understood that the compounds of the present invention encompass racemic, optically-active, regioisomeric and stereoisomeric forms, or combinations thereof that possess the therapeutically useful properties described herein. Preparation of optically active forms is achieved in any suitable manner, including by way of non-limiting example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic

separation using a chiral stationary phase. In some embodiments, mixtures of one or more isomer is utilized as the therapeutic compound described herein. In certain embodiments, compounds described herein contains one or more chiral centers. These compounds are prepared by any means, including enantioselective synthesis and/or separation of a mixture of enantiomers and/or diastereomers. Resolution of compounds and isomers thereof is achieved by any means including, by way of non-limiting example, chemical processes, enzymatic processes, fractional crystallization, distillation, chromatography, and the like.

[00103] In some embodiments, the ASBTI is




[00104] In some embodiments, the ASBTI is



[00105] (maralixibat chloride, LUM-001, SHP625, lopixibat chloride), or an alternative pharmaceutically acceptable salt thereof.



[00106] In some embodiments, the ASBTI is



[00107] (volixibat, (2R,3R,4S,5R,6R)-4-benzyloxy-6-{3-[3-((3S,4R,5R)-3-butyl-7-dimethylamino-3-ethyl-4-hydroxy-1,1-dioxo-2,3,4,5-tetrahydro-1H-


benzo[b]thiepin-5-yl)-phenyl]-ureido}-3,5-dihydroxy-tetrahydro-pyran-2-ylmethyl) hydrogen sulfate), or a pharmaceutically acceptable salt thereof.

[00108] In some embodiments, the ASBTI is




[00109] or

(LUM-002; SHP626; SAR548304; volixibat potassium), or an alternative pharmaceutically acceptable salt thereof.



[00110] In various embodiments the ASBTI is


(odevixibat; AZD8294; WHO10706; AR-H064974; SCHEMBL946468; A4250; 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(N-((R)-a-[N-((S)-1-carboxypropyl) carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine), or a pharmaceutically acceptable salt thereof.



[00111] In some embodiments, the ASBTI is

(elobixibat; 2-[[[(2R)-2-[[2-[(3,3-dibutyl-7-methylsulfanyl-1,1-dioxo-5-phenyl-2,4-dihydro-

1λ6,5-benzothiazepin-8-yl)oxy]acetyl]amino]-2-phenylacetyl]amino]acetic acid), or a pharmaceutically acceptable salt thereof.



[00112] In some embodiments, the ASBTI is (GSK2330672; linerixibat; 3-(((3R,5R)-3-butyl-3-ethyl-7-(methoxy)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,4-benzothiazepin-8-yl)methyl)amino)pentanedioic acid), or a pharmaceutically acceptable salt thereof.

[00113] In some embodiments, the ASBTI used in the methods or compositions of the present invention is maralixibat (e.g., as maralixibat chloride), volixibat (e.g., as volixibat potassium), or odeixibat (A4250), or a pharmaceutically acceptable salt thereof.

[00114] In some embodiments, the ASBTI used in the methods or compositions of the present invention is maralixibat, or a pharmaceutically acceptable salt thereof.

[00115] In some embodiments, the ASBTI used in the methods or compositions of the present invention is volixibat, or a pharmaceutically acceptable salt thereof.

[00116] In some embodiments, the ASBTI used in the methods or compositions of the present invention is odeixibat, or a pharmaceutically acceptable salt thereof.

[00117] In some embodiments, the ASBTI used in the methods or compositions of the present invention is elobixibat, or a pharmaceutically acceptable salt thereof.

[00118] In some embodiments, the ASBTI used in the methods or compositions of the present invention is GSK2330672, or a pharmaceutically acceptable salt thereof.

[00119] In some embodiments, the ASBTI may comprise a mixture of different ASBTIs; for example, the ASBTI may be a composition comprising maralixibat, volixibat, odeixibat, GSK2330672, elobixibat, or various combinations thereof.

Methods for treating cholestasis and minimizing gastrointestinal adverse effects

[00120] Provided herein is a method for treating cholestasis in a subject having a liver disease. The method includes administering to a subject in need of treatment an Apical Sodium-dependent Bile Acid Transporter Inhibitor (ASBTI). The ASBTI is maralixibat or volixibat, or a

pharmaceutically acceptable salt thereof. The ASBTI is administered in an amount of from about 100 $\mu\text{g/kg/day}$ to about 1400 $\mu\text{g/kg/day}$.

[00121] Provided herein is a method for treating cholestatic liver disease in a subject in need thereof, the method comprising administering a therapeutically effective amount of an ASBTI to the subject before ingestion of food, wherein the subject experiences a reduction in frequency and/or severity of one or more side effects associated with the administration of the ASBTI. The method includes administering to a subject in need of treatment an ASBTI before ingestion of food. In certain embodiments, the ASBTI is maralixibat or volixibat, or a pharmaceutically acceptable salt thereof. The ASBTI is administered in an amount of from about 100 $\mu\text{g/kg/day}$ to about 1400 $\mu\text{g/kg/day}$.

[00122] Provided herein is a method for reducing, minimizing, preventing, ameliorating, or eliminating one or more side effects associated with administration of an ASBTI in a subject in need thereof, the method comprising administering a therapeutically effective amount of the ASBTI to the subject before ingestion of food. In certain embodiments, the one or more side effects associated with administration of the ASBTI is reduced, minimized, prevented, ameliorated or eliminated as compared to the side effects when the ASBTI is administered after ingestion of food, at the same time as food, or mixed with food.

[00123] In certain embodiments, the one or more side effects is diarrhea, loose stools, nausea, gastrointestinal pain, abdominal pain, cramping, anorectal discomfort, or a combination thereof.

[00124] In certain embodiments, the ASBTI is administered to the subject in a fasted state. In certain embodiments, the ASBTI is administered less than about 1 minute, less than about 5 minutes, less than about 10 minutes, less than about 15 minutes, less than about 20 minutes, less than about 30 minutes or less than about 60 minutes before ingestion of food. In certain embodiments, the ASBTI is administered immediately prior to the ingestion of food.

[00125] In various embodiments, the liver disease is a cholestatic liver disease. In some embodiments, the liver disease is PFIC, ALGS, PSC, biliary atresia, intrahepatic cholestasis of pregnancy, PBC, any of the cholestatic liver diseases discussed above, or various combinations thereof.

[00126] In certain embodiments, the cholestatic liver disease is progressive familial intrahepatic cholestasis (PFIC), PFIC type 1, PFIC type 2, PFIC type 3, Alagille syndrome, Dubin-Johnson Syndrome, biliary atresia, post-Kasai biliary atresia, post-liver transplantation biliary atresia,

post-liver transplantation cholestasis, post-liver transplantation associated liver disease, intestinal failure associated liver disease, bile acid mediated liver injury, pediatric primary sclerosing cholangitis, MRP2 deficiency syndrome, neonatal sclerosing cholangitis, a pediatric obstructive cholestasis, a pediatric non-obstructive cholestasis, a pediatric extrahepatic cholestasis, a pediatric intrahepatic cholestasis, a pediatric primary intrahepatic cholestasis, a pediatric secondary intrahepatic cholestasis, benign recurrent intrahepatic cholestasis (BRIC), BRIP type 1, BRIC type 2, BRIC type 3, total parenteral nutrition associated cholestasis, paraneoplastic cholestasis, Stauffer syndrome, drug-associated cholestasis, infection-associated cholestasis, or gallstone disease. In some embodiments, the cholestatic liver disease is a pediatric form of liver disease. In some embodiments, the subject has intrahepatic cholestasis of pregnancy (ICP).

[00127] In certain embodiments, a cholestatic liver disease is characterized by one or more symptoms selected from jaundice, pruritis, cirrhosis, hypercholelismia, neonatal respiratory distress syndrome, lung pneumonia, increased serum concentration of bile acids, increased hepatic concentration of bile acids, increased serum concentration of bilirubin, hepatocellular injury, liver scarring, liver failure, hepatomegaly, xanthomas, malabsorption, splenomegaly, diarrhea, pancreatitis, hepatocellular necrosis, giant cell formation, hepatocellular carcinoma, gastrointestinal bleeding, portal hypertension, hearing loss, fatigue, loss of appetite, anorexia, peculiar smell, dark urine, light stools, steatorrhea, failure to thrive, and/or renal failure.

[00128] In various embodiments the liver disease is PFIC 2 and the subject has a non-truncating mutation in the *ABCB11* gene. In various embodiments the non-truncating mutation in the *ABCB11* gene is a missense mutation. In various embodiments the missense mutation may be selected from one of those mutations listed in Byrne, et al., “Missense Mutations and Single Nucleotide Polymorphisms in *ABCB11* Impair Bile Salt Export Pump Processing and Function or Disrupt Pre-Messenger RNA Splicing,” *Hepatology*, 49:553-567 (2009), which is incorporated herein by reference in its entirety for all purposes.

[00129] In various embodiments the subject has a condition associated with, caused by or caused in part by a BSEP deficiency. In certain embodiments, the condition associated with, caused by or caused in part by the BSEP deficiency is neonatal hepatitis, primary biliary cirrhosis (PBC), primary sclerosing cholangitis (PSC), PFIC 2, benign recurrent intrahepatic cholestasis (BRIC), intrahepatic cholestasis of pregnancy (ICP), drug-induced cholestasis, oral-contraceptive-induced cholestasis, biliary atresia, or a combination thereof.

[00130] In various embodiments, the patient is a pediatric patient under the age of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or 18 years old. In certain embodiments, the pediatric subject is a newborn, a pre-term newborn, an infant, a toddler, a preschooler, a school-age child, a pre-pubescent child, post-pubescent child, an adolescent, or a teenager under the age of eighteen. In some embodiments, the pediatric subject is a newborn, a pre-term newborn, an infant, a toddler, a preschooler, or a school-age child. In some embodiments, the pediatric subject is a newborn, a pre-term newborn, an infant, a toddler, or a preschooler. In some embodiments, the pediatric subject is a newborn, a pre-term newborn, an infant, or a toddler. In some embodiments, the pediatric subject is a newborn, a pre-term newborn, or an infant. In some embodiments, the pediatric subject is a newborn. In some embodiments, the pediatric subject is an infant. In some embodiments, the pediatric subject is a toddler. In various embodiments, the pediatric patient has PFIC 2, PFIC 1, or ALGS. In some embodiments, the patient is an adult over the age of 18, 20, 30, 40, 50, 60, or 70. In some patients, the adult patient has PSC. In some embodiments, the pediatric patient has any pediatric cholestatic condition resulting in below normal growth, height, or weight.

[00131] In certain embodiments, methods of the present invention comprise non-systemic administration of a therapeutically effective amount of an ASBTI. In certain embodiments, the methods comprise contacting the gastrointestinal tract, including the distal ileum and/or the colon and/or the rectum, of an individual in need thereof with an ASBTI. In various embodiments, the methods of the present invention cause a reduction in intraenterocyte bile acids, or a reduction in damage to hepatocellular or intestinal architecture caused by cholestasis or a cholestatic liver disease.

[00132] In various embodiments, methods of the present invention comprise delivering to ileum or colon of the individual a therapeutically effective amount of any ASBTI described herein.

[00133] In various embodiments, methods of the present invention comprise reducing damage to hepatocellular or intestinal architecture or cells from cholestasis or a cholestatic liver disease comprising administration of a therapeutically effective amount of an ASBTI. In certain embodiments, the methods of the present invention comprise reducing intraenterocyte bile acids/salts through administration of a therapeutically effective amount of an ASBTI to an individual in need thereof.

[00134] In some embodiments, methods of the present invention provide for inhibition of bile salt recycling upon administration of any of the compounds described herein to an individual. In some embodiments, an ASBTI described herein is systemically absorbed upon administration. In some embodiments, an ASBTI described herein is not absorbed systemically. In some embodiments, an ASBTI herein is administered to the individual orally. In some embodiments, an ASBTI described herein is delivered and/or released in the distal ileum of an individual.

[00135] In various embodiments, contacting the distal ileum of an individual with an ASBTI (e.g., any ASBTI described herein) inhibits bile acid reuptake and increases the concentration of bile acids/salts in the vicinity of L-cells in the distal ileum and/or colon and/or rectum, thereby reducing intraenterocyte bile acids, reducing serum and/or hepatic bile acid levels, reducing overall serum bile acid load, and/or reducing damage to ileal architecture caused by cholestasis or a cholestatic liver disease. Without being limited to any particular theory, reducing serum and/or hepatic bile acid levels ameliorates hypercholeemia and/or cholestatic disease.

[00136] Administration of a compound described herein may be achieved in any suitable manner including, by way of non-limiting example, by oral, enteric, parenteral (e.g., intravenous, subcutaneous, intramuscular), intranasal, buccal, topical, rectal, or transdermal administration routes. Any compound or composition described herein may be administered in a method or formulation appropriate to treat a newborn or an infant. Any compound or composition described herein may be administered in an oral formulation (e.g., solid or liquid) to treat a newborn or an infant. Any compound or composition described herein may be administered prior to ingestion of food, with food or after ingestion of food.

[00137] In certain embodiments, a compound or a composition comprising a compound described herein is administered for prophylactic and/or therapeutic treatments. In therapeutic applications, the compositions are administered to an individual already suffering from a disease or condition, in an amount sufficient to cure or at least partially arrest the symptoms of the disease or condition. In various instances, amounts effective for this use depend on the severity and course of the disease or condition, previous therapy, the individual's health status, weight, and response to the drugs, and the judgment of the treating physician.

[00138] In prophylactic applications, compounds or compositions containing compounds described herein may be administered to an individual susceptible to or otherwise at risk of a particular disease, disorder or condition. In certain embodiments of this use, the precise amounts

of compound administered depend on the individual's state of health, weight, and the like. Furthermore, in some instances, when a compound or composition described herein is administered to an individual, effective amounts for this use depend on the severity and course of the disease, disorder or condition, previous therapy, the individual's health status and response to the drugs, and the judgment of the treating physician.

[00139] In certain embodiments of the methods of the present invention, wherein following administration of a selected dose of a compound or composition described herein, an individual's condition does not improve, upon the doctor's discretion the administration of a compound or composition described herein is optionally administered chronically, that is, for an extended period of time, including throughout the duration of the individual's life in order to ameliorate or otherwise control or limit the symptoms of the individual's disorder, disease or condition.

[00140] In certain embodiments of the methods of the present invention, an effective amount of a given agent varies depending upon one or more of a number of factors such as the particular compound, disease or condition and its severity, the identity (e.g., weight) of the subject or host in need of treatment, and is determined according to the particular circumstances surrounding the case, including, e.g., the specific agent being administered, the route of administration, the condition being treated, and the subject or host being treated. In some embodiments, doses administered include those up to the maximum tolerable dose. In some embodiments, doses administered include those up to the maximum tolerable dose by a newborn or an infant.

[00141] In various embodiments of the methods of the present invention, a desired dose is conveniently presented in a single dose or in divided doses administered simultaneously (or over a short period of time) or at appropriate intervals, for example as two, three, four or more sub-doses per day. In various embodiments, a single dose of an ASBTI is administered every 6 hours, every 12 hours, every 24 hours, every 48 hours, every 72 hours, every 96 hours, every 5 days, every 6 days, or once a week. In some embodiments the total single dose of an ASBTI is in a range described below.

[00142] In various embodiments of methods of the present invention, in the case wherein the patient's status does improve, upon the doctor's discretion an ASBTI is optionally given continuously; alternatively, the dose of drug being administered is temporarily reduced or temporarily suspended for a certain length of time (i.e., a "drug holiday"). The length of the drug holiday optionally varies between 2 days and 1 year, including by way of example only, 2 days,

3 days, 4 days, 5 days, 6 days, 7 days, 10 days, 12 days, 15 days, 20 days, 28 days, 35 days, 50 days, 70 days, 100 days, 120 days, 150 days, 180 days, 200 days, 250 days, 280 days, 300 days, 320 days, 350 days, or 365 days. The dose reduction during a drug holiday includes from 10%-100% of the original dose, including, by way of example only, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% of the original dose. In some embodiments the total single dose of an ASBTI is in a range described below.

[00143] Once improvement of the patient's conditions has occurred, a maintenance dose is administered if necessary. Subsequently, the dosage or the frequency of administration, or both, is reduced, as a function of the symptoms, to a level at which the improved disease, disorder or condition is retained. In some embodiments, patients require intermittent treatment on a long-term basis upon any recurrence of symptoms.

[00144] In certain instances, there are a large number of variables in regard to an individual treatment regime, and considerable excursions from these recommended values are considered within the scope described herein. Dosages described herein are optionally altered depending on a number of variables such as, by way of non-limiting example, the activity of the compound used, the disease or condition to be treated, the mode of administration, the requirements of the individual subject, the severity of the disease or condition being treated, and the judgment of the practitioner.

[00145] Toxicity and therapeutic efficacy of such therapeutic regimens are optionally determined by pharmaceutical procedures in cell cultures or experimental animals, including, but not limited to, the determination of the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between the toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD₅₀ and ED₅₀. Compounds exhibiting high therapeutic indices are preferred. In certain embodiments, data obtained from cell culture assays and animal studies are used in formulating a range of dosage for use in human. In specific embodiments, the dosage of compounds described herein lies within a range of circulating concentrations that include the ED₅₀ with minimal toxicity. The dosage optionally varies within this range depending upon the dosage form employed and the route of administration utilized.

[00146] In certain embodiments, the composition used or administered comprises an absorption inhibitor, a carrier, and one or more of a cholesterol absorption inhibitor, an enteroendocrine peptide, a peptidase inhibitor, a spreading agent, and a wetting agent.

[00147] In some embodiments of methods of the present invention, the composition used to prepare an oral dosage form or administered orally comprises an absorption inhibitor, an orally suitable carrier, an optional cholesterol absorption inhibitor, an optional enteroendocrine peptide, an optional peptidase inhibitor, an optional spreading agent, and an optional wetting agent. In certain embodiments, the orally administered compositions evoke an anorectal response. In specific embodiments, the anorectal response is an increase in secretion of one or more enteroendocrine by cells in the colon and/or rectum (e.g., in L-cells the epithelial layer of the colon, ileum, rectum, or a combination thereof). In some embodiments, the anorectal response persists for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 or 24 hours. In other embodiments the anorectal response persists for a period between 24 hours and 48 hours, while in other embodiments the anorectal response persists for a period greater than 48 hours.

Dosages

[00148] In various embodiments the ASBTI is maralixibat or volixibat, or a pharmaceutically acceptable salt thereof.

[00149] In various embodiments the ASBTI is administered to a subject before ingestion of food.

[00150] In various embodiments, efficacy and safety of ASBTI administration to the patient is monitored by measuring serum levels of 7α -hydroxy-4-cholesten-3-one (7α C4), sBA concentration, a ratio of 7α C4 to sBA (7α C4:sBA), serum conjugated bilirubin concentration, serum autotaxin concentration, serum bilirubin concentration, serum total cholesterol concentration, serum LDL-C concentration, serum ALT concentration, serum AST concentration, or a combination thereof. In various embodiments, efficacy of ASBTI administration is measured by monitoring observer-reported itch reported outcome (ITCHRO(OBS)) score, a HRQoL (e.g., PedsQL) score, a CSS score, a xanthoma score, a height Z-score, a weight Z-score, or various combinations thereof. In various embodiments, the method includes monitoring serum levels of 7α -hydroxy-4-cholesten-3-one (7α C4), sBA concentration, a ratio of 7α C4 to sBA (7α C4:sBA), serum conjugated bilirubin concentration, serum total

cholesterol concentration, serum LDL-C concentration, serum autotaxin concentration, serum bilirubin concentration, serum ALT concentration, serum AST concentration, or a combination thereof. In various embodiments, the method includes monitoring observer-reported itch reported outcome (ITCHRO(OBS)) score, a weight Z-score, a HRQoL (e.g., PedsQL) score, a xanthoma score, a CSS score, a height Z-score, or various combinations thereof.

[00151] In some embodiments, the ASBTI is administered at a dose of about or at least about 0.5 µg/kg, 1 µg/kg, 2 µg/kg, 3 µg/kg, 4 µg/kg, 5 µg/kg, 6 µg/kg, 7 µg/kg, 8 µg/kg, 9 µg/kg, 10 µg/kg, 15 µg/kg, 20 µg/kg, 25 µg/kg, 30 µg/kg, 35 µg/kg, 40 µg/kg, 45 µg/kg, 50 µg/kg, 55 µg/kg, 60 µg/kg, 65 µg/kg, 70 µg/kg, 75 µg/kg, 80 µg/kg, 85 µg/kg, 90 µg/kg, 100 µg/kg, 140 µg/kg, 150 µg/kg, 200 µg/kg, 240 µg/kg, 250 µg/kg, 280 µg/kg, 300 µg/kg, 360 µg/kg, 380 µg/kg, 400 µg/kg, 500 µg/kg, 560 µg/kg, 600 µg/kg, 700 µg/kg, 800 µg/kg, 880 µg/kg, 900 µg/kg, 1,000 µg/kg, 1,100 µg/kg, 1,200 µg/kg, 1,300 µg/kg, 1,400 µg/kg, 1,500 µg/kg, 1,600 µg/kg, 1,700 µg/kg, 1,800 µg/kg, 1,900 µg/kg, or 2,000 µg/kg. In various embodiments, the ASBTI is administered at a dose not exceeding about 1 µg/kg, 2 µg/kg, 3 µg/kg, 4 µg/kg, 5 µg/kg, 6 µg/kg, 7 µg/kg, 8 µg/kg, 9 µg/kg, 10 µg/kg, 15 µg/kg, 20 µg/kg, 25 µg/kg, 30 µg/kg, 35 µg/kg, 40 µg/kg, 45 µg/kg, 50 µg/kg, 55 µg/kg, 60 µg/kg, 65 µg/kg, 70 µg/kg, 75 µg/kg, 80 µg/kg, 85 µg/kg, 90 µg/kg, 100 µg/kg, 140 µg/kg, 150 µg/kg, 200 µg/kg, 240 µg/kg, 250 µg/kg, 280 µg/kg, 300 µg/kg, 360 µg/kg, 380 µg/kg, 400 µg/kg, 500 µg/kg, 560 µg/kg, 600 µg/kg, 700 µg/kg, 800 µg/kg, 880 µg/kg, 900 µg/kg, 1,000 µg/kg, 1,100 µg/kg, 1,200 µg/kg, 1,300 µg/kg, 1,400 µg/kg, 1,500 µg/kg, 1,600 µg/kg, 1,700 µg/kg, 1,800 µg/kg, 1,900 µg/kg, 2,000, or 2,100 µg/kg. In various embodiments, the ASBTI is administered at a dose of about or of at least about 0.5 mg/day, 1 mg/day, 2 mg/day, 3 mg/day, 4 mg/day, 5 mg/day, 6 mg/day, 7 mg/day, 8 mg/day, 9 mg/day, 10 mg/day, 11 mg/day, 12 mg/day, 13 mg/day, 14 mg/day, 15 mg/day, 16 mg/day, 17 mg/day, 18 mg/day, 19 mg/day, 20 mg/day, 30 mg/day, 40 mg/day, 50 mg/day, 60 mg/day, 70 mg/day, 80 mg/day, 90 mg/day, 100 mg/day, 150 mg/day, 200 mg/day, 300 mg/day, 500 mg/day, 600 mg/day, 700 mg/day, 800 mg/day, 900 mg/day, 1000 mg/day. In various embodiments, the ASBTI is administered at a dose of not more than about 1 mg/day, 2 mg/day, 3 mg/day, 4 mg/day, 5 mg/day, 6 mg/day, 7 mg/day, 8 mg/day, 9 mg/day, 10 mg/day, 11 mg/day, 12 mg/day, 13 mg/day, 14 mg/day, 15 mg/day, 16 mg/day, 17 mg/day, 18 mg/day, 19 mg/day, 20 mg/day, 30 mg/day, 40 mg/day, 50 mg/day, 60 mg/day, 70 mg/day, 80 mg/day, 90 mg/day, 100 mg/day, 150

mg/day, 200 mg/day, 300 mg/day, 500 mg/day, 600 mg/day, 700 mg/day, 800 mg/day, 900 mg/day, 1,000 mg/day, 1,100 mg/day.

[00152] In some embodiments, the ASBTI is administered at a dose of from about 140 $\mu\text{g/kg/day}$ to about 1400 $\mu\text{g/kg/day}$. In various embodiments, the ASBTI is administered at a dose of about or at least about 0.5 $\mu\text{g/kg/day}$, 1 $\mu\text{g/kg/day}$, 2 $\mu\text{g/kg/day}$, 3 $\mu\text{g/kg/day}$, 4 $\mu\text{g/kg/day}$, 5 $\mu\text{g/kg/day}$, 6 $\mu\text{g/kg/day}$, 7 $\mu\text{g/kg/day}$, 8 $\mu\text{g/kg/day}$, 9 $\mu\text{g/kg/day}$, 10 $\mu\text{g/kg/day}$, 15 $\mu\text{g/kg/day}$, 20 $\mu\text{g/kg/day}$, 25 $\mu\text{g/kg/day}$, 30 $\mu\text{g/kg/day}$, 35 $\mu\text{g/kg/day}$, 40 $\mu\text{g/kg/day}$, 45 $\mu\text{g/kg/day}$, 50 $\mu\text{g/kg/day}$, 100 $\mu\text{g/kg/day}$, 140 $\mu\text{g/kg/day}$, 150 $\mu\text{g/kg/day}$, 200 $\mu\text{g/kg/day}$, 240 $\mu\text{g/kg/day}$, 280 $\mu\text{g/kg/day}$, 300 $\mu\text{g/kg/day}$, 250 $\mu\text{g/kg/day}$, 280 $\mu\text{g/kg/day}$, 300 $\mu\text{g/kg/day}$, 360 $\mu\text{g/kg/day}$, 380 $\mu\text{g/kg/day}$, 400 $\mu\text{g/kg/day}$, 500 $\mu\text{g/kg/day}$, 560 $\mu\text{g/kg/day}$, 600 $\mu\text{g/kg/day}$, 700 $\mu\text{g/kg/day}$, 800 $\mu\text{g/kg/day}$, 880 $\mu\text{g/kg/day}$, 900 $\mu\text{g/kg/day}$, 1,000 $\mu\text{g/kg/day}$, 1,100 $\mu\text{g/kg/day}$, 1,200 $\mu\text{g/kg/day}$, or 1,300 $\mu\text{g/kg/day}$. In various embodiments, the ASBTI is administered at a dose not exceeding about 1 $\mu\text{g/kg/day}$, 2 $\mu\text{g/kg/day}$, 3 $\mu\text{g/kg/day}$, 4 $\mu\text{g/kg/day}$, 5 $\mu\text{g/kg/day}$, 6 $\mu\text{g/kg/day}$, 7 $\mu\text{g/kg/day}$, 8 $\mu\text{g/kg/day}$, 9 $\mu\text{g/kg/day}$, 10 $\mu\text{g/kg/day}$, 15 $\mu\text{g/kg/day}$, 20 $\mu\text{g/kg/day}$, 25 $\mu\text{g/kg/day}$, 30 $\mu\text{g/kg/day}$, 35 $\mu\text{g/kg/day}$, 40 $\mu\text{g/kg/day}$, 45 $\mu\text{g/kg/day}$, 50 $\mu\text{g/kg/day}$, 100 $\mu\text{g/kg/day}$, 140 $\mu\text{g/kg/day}$, 150 $\mu\text{g/kg/day}$, 200 $\mu\text{g/kg/day}$, 240 $\mu\text{g/kg/day}$, 280 $\mu\text{g/kg/day}$, 300 $\mu\text{g/kg/day}$, 250 $\mu\text{g/kg/day}$, 280 $\mu\text{g/kg/day}$, 300 $\mu\text{g/kg/day}$, 360 $\mu\text{g/kg/day}$, 380 $\mu\text{g/kg/day}$, 400 $\mu\text{g/kg/day}$, 500 $\mu\text{g/kg/day}$, 560 $\mu\text{g/kg/day}$, 600 $\mu\text{g/kg/day}$, 700 $\mu\text{g/kg/day}$, 800 $\mu\text{g/kg/day}$, 880 $\mu\text{g/kg/day}$, 900 $\mu\text{g/kg/day}$, 1,000 $\mu\text{g/kg/day}$, 1,100 $\mu\text{g/kg/day}$, 1,200 $\mu\text{g/kg/day}$, 1,300 $\mu\text{g/kg/day}$, or 1,400 $\mu\text{g/kg/day}$. In various embodiments, the ASBTI is administered at a dose of from about 0.5 $\mu\text{g/kg/day}$ to about 500 $\mu\text{g/kg/day}$, from about 0.5 $\mu\text{g/kg/day}$ to about 250 $\mu\text{g/kg/day}$, from about 1 $\mu\text{g/kg/day}$ to about 100 $\mu\text{g/kg/day}$, from about 10 $\mu\text{g/kg/day}$ to about 50 $\mu\text{g/kg/day}$, from about 10 $\mu\text{g/kg/day}$ to about 100 $\mu\text{g/kg/day}$, from about 0.5 $\mu\text{g/kg/day}$ to about 2000 $\mu\text{g/kg/day}$, from about 280 $\mu\text{g/kg/day}$ to about 1400 $\mu\text{g/kg/day}$, from about 420 $\mu\text{g/kg/day}$ to about 1400 $\mu\text{g/kg/day}$, from about 250 to about 550 $\mu\text{g/kg/day}$, from about 560 $\mu\text{g/kg/day}$ to about 1400 $\mu\text{g/kg/day}$, from 700 $\mu\text{g/kg/day}$ to about 1400 $\mu\text{g/kg/day}$, from about 560 $\mu\text{g/kg/day}$ to about 1200 $\mu\text{g/kg/day}$, from about 700 $\mu\text{g/kg/day}$ to about 1200 $\mu\text{g/kg/day}$, from about 560 $\mu\text{g/kg/day}$ to about 1000 $\mu\text{g/kg/day}$, from about 700 $\mu\text{g/kg/day}$ to about 1000 $\mu\text{g/kg/day}$, from about 800 $\mu\text{g/kg/day}$ to about 1000 $\mu\text{g/kg/day}$, from about 200 $\mu\text{g/kg/day}$ to about 600 $\mu\text{g/kg/day}$, from about 300 $\mu\text{g/kg/day}$ to about 600 $\mu\text{g/kg/day}$, from about 400 $\mu\text{g/kg/day}$ to about 500 $\mu\text{g/kg/day}$, from about 400 $\mu\text{g/kg/day}$ to about 600 $\mu\text{g/kg/day}$, from about 400 $\mu\text{g/kg/day}$ to about 700

μg/kg/day, from about 400 μg/kg/day to about 800 μg/kg/day, from about 500 μg/kg/day to about 800 μg/kg/day, from about 500 μg/kg/day to about 900 μg/kg/day, from about 600 μg/kg/day to about 900 μg/kg/day, from about 700 μg/kg/day to about 900 μg/kg/day, from about 200 μg/kg/day to about 600 μg/kg/day, from about 800 μg/kg/day to about 900 μg/kg/day, from about 100 μg/kg/day to about 1500 μg/kg/day, from about 300 μg/kg/day to about 2,000 μg/kg/day, or from about 400 μg/kg/day to about 2000 μg/kg/day.

[00153] In some embodiments, the ASBTI is administered at a dose of from about 30 μg/kg to about 1400 μg/kg per dose. In some embodiments, the ASBTI is administered at a dose of from about 0.5 μg/kg to about 2000 μg/kg per dose, from about 0.5 μg/kg to about 1500 μg/kg per dose, from about 100 μg/kg to about 700 μg/kg per dose, from about 5 μg/kg to about 100 μg/kg per dose, from about 10 μg/kg to about 500 μg/kg per dose, from about 50 μg/kg to about 1400 μg/kg per dose, from about 300 μg/kg to about 2,000 μg/kg per dose, from about 60 μg/kg to about 1200 μg/kg per dose, from about 70 μg/kg to about 1000 μg/kg per dose, from about 70 μg/kg to about 700 μg/kg per dose, from 80 μg/kg to about 1000 μg/kg per dose, from 80 μg/kg to about 800 μg/kg per dose, from 100 μg/kg to about 800 μg/kg per dose, from 100 μg/kg to about 600 μg/kg per dose, from 150 μg/kg to about 700 μg/kg per dose, from 150 μg/kg to about 500 μg/kg per dose, from 200 μg/kg to about 400 μg/kg per dose, from 200 μg/kg to about 300 μg/kg per dose, or from 300 μg/kg to about 400 μg/kg per dose.

[00154] In some embodiments, the ASBTI is administered at a dose of from about 0.5 mg/day to about 550 mg/day. In various embodiments, the ASBTI is administered at a dose of from about 1 mg/day to about 500 mg/day, from about 1 mg/day to about 300 mg/day, from about 1 mg/day to about 200 mg/day, from about 2 mg/day to about 300 mg/day, from about 2 mg/day to about 200 mg/day, from about 4 mg/day to about 300 mg/day, from about 4 mg/day to about 200 mg/day, from about 4 mg/day to about 150 mg/day, from about 5 mg/day to about 150 mg/day, from about 5 mg/day to about 100 mg/day, from about 5 mg/day to about 80 mg/day, from about 5 mg/day to about 50 mg/day, from about 5 mg/day to about 40 mg/day, from about 5 mg/day to about 30 mg/day, from about 5 mg/day to about 20 mg/day, from about 5 mg/day to about 15 mg/day, from about 10 mg/day to about 100 mg/day, from about 10 mg/day to about 80 mg/day, from about 10 mg/day to about 50 mg/day, from about 10 mg/day to about 40 mg/day, from about 10 mg/day to about 20 mg/day, from about 20 mg/day to about 100 mg/day, from about 20

mg/day to about 80 mg/day, from about 20 mg/day to about 50 mg/day, or from about 20 mg/day to about 40 mg/day, or from about 20 mg/day to about 30 mg/day.

[00155] In some embodiments, the ASBTI is administered twice daily (BID) in an amount of about 200 µg/kg to about 400 µg/kg per dose. In some embodiments, the ASBTI is administered in an amount of about 280 µg/kg/day to about 1400 µg/kg/day. In some embodiments, the ASBTI is administered in an amount of about 400 µg/kg/day to about 800 µg/kg/day. In some embodiments, the ASBTI is administered in an amount of about 20 mg/day to about 50 mg/day. In some embodiments, the ASBTI is administered in an amount of from about 5 mg/day to about 15 mg/day. In some embodiments, the ASBTI is administered in an amount of from about 560 µg/kg/day to about 1,400 µg/kg/day. In some embodiments, the ASBTI is administered in an amount of from about 700 µg/kg/day to about 1,400 µg/kg/day. In some embodiments, the ASBTI is administered in an amount of from about 400 µg/kg/day to about 800 µg/kg/day. In some embodiments, the ASBTI is administered in an amount of from about 700 µg/kg/day to about 900 µg/kg/day. In some embodiments, the ASBTI is administered in an amount of from about 560 µg/kg/day to about 1400 µg/kg/day. In some embodiments, the ASBTI is administered in an amount from 700 µg/kg/day to about 1400 µg/kg/day. In some embodiments, the ASBTI is administered in an amount of from about 200 µg/kg/day to about 600 µg/kg/day. In some embodiments, the ASBTI is administered in an amount of from about 400 µg/kg/day to about 600 µg/kg/day.

[00156] In various embodiments, the dose of the ASBTI is a first dose level. In various embodiments, the dose of the ASBTI is a second dose level. In some embodiments, the second dose level is greater than the first dose level. In some embodiments, the second dose level is about or at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90 or 100 times or fold greater than the first dose level. In some embodiments, the second dose level is not in excess of about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, or 150 times or fold greater than the first dose level.

[00157] In various embodiments, the ASBTI is administered once daily (QD) at one of the above doses or within one of the above dose ranges. In various embodiments, the ASBTI is administered twice daily (BID) at one of the above doses or within one of the above dose ranges. In various embodiments, an ASBTI dose is administered daily, every other day, twice a week, or once a week.

[00158] In various embodiments, the ASBTI is administered regularly for a period of about or of at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 48, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 600, 700, or 800 weeks. In various embodiments, the ASBTI is administered for not more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 48, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, or 1000 weeks. In various embodiments, the ASBTI is administered regularly for a period of about or of at least about 0.5, 1, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, or 10 years. In various embodiments, the ASBTI is administered regularly for a period not in excess of about 0.5, 1, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 15 years.

Reduction in symptoms or a change in a disease-relevant laboratory measures of cholestatic liver disease

[00159] In various embodiments of the above methods of the invention, administration of the ASBTI results in a reduction in a symptom or a change in a disease-relevant laboratory measure of the cholestatic liver disease (i.e., improvement in the patient's condition) that is maintained for about or for at least about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 17 weeks, 18 weeks, 19 weeks, 20 weeks, 21 weeks, 22 weeks, 23 weeks, 24 weeks, 6 months, 25 weeks, 26 weeks, 27 weeks, 28 weeks, 29 weeks, 30 weeks, 31 weeks, 32 weeks, 33 weeks, 34 weeks, 35 weeks, 36 weeks, 37 weeks, 38 weeks, 39 weeks, 40 weeks, 41 weeks, 42 weeks, 43 weeks, 44 weeks, 45 weeks, 46 weeks, 47 weeks, 48 weeks, 49 weeks, 50 weeks, 51 weeks, 52 weeks, 1 year, 13 months, 14 months, 15 months, 16 months, 17 months, 18 months, 19 months, 20 months, 21 months, 22 months, 23 months, 24 months, 2 years, 2.5 years, 3 years, 3.5 years, 4 years, 4.5 years, 5 years, 5.5 years, 6 years, 6.5 years, 7 years, 8 years, 9 years, or 10 years. In various embodiments, the reduction in the symptom or a change in a disease-relevant laboratory measure comprises a reduction in sBA concentration, an increase in serum 7αC4 concentration, an increase in the 7αC4:sBA ratio, an increase in fBA excretion, a reduction in pruritis, a decrease in serum total cholesterol concentration, a decrease in serum LDL-C cholesterol concentration, a reduction in ALT levels, an increase in a quality of life inventory score, an increase in a quality of life inventory score related to fatigue, a reduction in a xanthoma score, a reduction in serum autotaxin concentration, an increase in growth, or a combination thereof. In various embodiments, the reduction in the symptom or a change in a disease-relevant laboratory measure is determined relative to a

baseline level. That is, the reduction in the symptom or a change in a disease-relevant laboratory measure is determined relative to a measurement of the symptom or a change in a disease-relevant laboratory measure prior to 1) changing a dose level of the ASBTI administered to the patient, 2) changing a dosing regimen followed for the patient, 3) commencing administration of the ASBTI, or 4) any other of various alterations made with the intention of reducing the symptom or a change in a disease-relevant laboratory measure in the patient. In various embodiments, the reduction in symptom or a change in a disease-relevant laboratory measure is a statistically significant reduction.

[00160] In various embodiments, the reduction in a symptom or a change in a disease-relevant laboratory measure of the cholestatic liver disease is measured as a progressive decrease in the symptom or a change in a disease-relevant laboratory measure for about or for at least about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 17 weeks, 18 weeks, 19 weeks, 20 weeks, 21 weeks, 22 weeks, 23 weeks, 24 weeks, 25 weeks, 26 weeks, 27 weeks, 28 weeks, 29 weeks, 30 weeks, 31 weeks, 32 weeks, 33 weeks, 34 weeks, 35 weeks, 36 weeks, 37 weeks, 38 weeks, 39 weeks, 40 weeks, 41 weeks, 42 weeks, 43 weeks, 44 weeks, 45 weeks, 46 weeks, 47 weeks, 48 weeks, 49 weeks, 50 weeks, 51 weeks, 52 weeks, 1 year, 13 months, 14 months, 15 months, 16 months, 17 months, 18 months, 19 months, 20 months, 21 months, 22 months, 23 months, 23 months, 2 years, 2.5 years, 3 years, 3.5 years, 4 years, 4.5 years, 5 years, 5.5 years, 6 years, 6.5 years, 7 years, 8 years, 9 years, or 10 years.

[00161] In some embodiments, the patient is the pediatric patient and the reduction in symptom or a change in a disease-relevant laboratory measure comprises an increase or improvement in growth. In some embodiments, the increase in growth is measured relative to baseline. In various embodiments, increase in growth is measured as an increase in height Z-score or in weight Z-score. In various embodiments, the increase in height Z-score or in weight Z-score is statistically significant. In various embodiments, the height Z-score, the weight Z-score, or both is increased by at least 0.1, 0.11, 0.12, 0.13, 0.14, 0.15, 0.16, 0.17, 0.18, 0.19, 0.2, 0.21, 0.22, 0.23, 0.24, 0.25, 0.26, 0.27, 0.28, 0.29, 0.3, 0.31, 0.32, 0.33, 0.34, 0.35, 0.36, 0.37, 0.38, 0.39, 0.4, 0.41, 0.42, 0.43, 0.44, 0.45, 0.46, 0.47, 0.48, 0.49, 0.5, 0.51, 0.52, 0.53, 0.54, 0.55, 0.56, 0.57, 0.58, 0.59, 0.6, 0.7, 0.8, or 0.9 relative to baseline. In some embodiments, the height Z-score, the weight Z-

score, or both progressively increases during administration of the ASBTI for a period of about or of at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 48, 50, 60, 70, or 72 weeks.

[00162] In various embodiments, the administration of the ASBTI results in an increase in serum 7 α C4 concentration. In various embodiments, the serum 7 α C4 concentration is increased by about or at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, or 500 times or fold relative to baseline. In various embodiments the serum 7 α C4 concentration is increased about or at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 150%, 200%, 300%, 400%, 500%, 600%, 700%, 800%, 900%, 1,000%, or 10,000% relative to baseline.

[00163] In various embodiments, the administration of the ASBTI results in an increase in the 7 α C4:sBA ratio to about or by at least about 1.25, 1.5, 1.75, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 500, 750, 1,000, 2,000, 3,000, 4,000, 5,000 or 10,000-fold relative to baseline.

[00164] In various embodiments, the administration of the ASBTI results in an increase in fBA excretion. In some embodiments, the administration of the ASBTI results in an increase in fBA excretion of about or of at least about 100%, 110%, 115%, 120%, 130%, 150%, 200%, 250%, 275%, 300%, 400%, 500%, 600%, 700%, 800%, 1,000%, 5,000%, 10,000% or 15,000% relative to baseline. In various embodiments, fBA excretion is increased by about or by at least about 1, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100 fold or times relative to baseline. In some embodiments, fBA excretion is increased by about or by at least about 100 μ mol, 150 μ mol, 200 μ mol, 250 μ mol, 300 μ mol, 400 μ mol, 500 μ mol, 600 μ mol, 700 μ mol, 800 μ mol, 900 μ mol, 1,000 μ mol, or 1,500 μ mol relative to baseline. In various embodiments, administration of the ASBTI results in a dose-dependent increase in fBA excretion so that administration of a higher dose of the ASBTI results in a corresponding higher level of fBA excretion. In various embodiments, the ASBTI is administered at a dose sufficient to result in an increase in bile acid secretion relative to baseline of at least about or of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100 fold or times relative to baseline.

[00165] In various embodiments, the administration of the ASBTI results in a decrease in sBA concentration of about or of at least about 5%, 10%, 15%, 20%, 25%, 30%, 31%, 35%, 40%, 45%, 50%, 55%, 57%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% relative to baseline.

[00166] In some embodiments, the administration of the ASBTI results in a reduction in severity of pruritus. In various embodiments, the severity of pruritus is measured using an ITCHRO(OBS) score, an ITCHRO score, a CSS score, or a combination thereof. In various embodiments, the administration of the ASBTI results in a reduction in the ITCHRO(OBS) score on a scale of 1 to 4 of about or of at least about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.25, 2.5, or 3 relative to baseline. In various embodiments, the administration of the ASBTI results in a reduction in the ITCHRO score on a scale of 1 to 10 of about or of at least about 0.1, 0.2, 0.3, 0.4, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, or 10. In various embodiments, the administration of the ASBTI results in a reduction of the ITCHRO(OBS) score, the ITCHRO score, or both to zero. In various embodiments, the administration of the ASBTI results in a reduction of the ITCHRO(OBS) score or ITCHRO score to 1.0 or lower. In various embodiments, the administration of the ASBTI results in a reduction of the CSS score by about or of at least about 0.1, 0.2, 0.3, 0.4, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.25, 2.5, or 3 relative to baseline. In various embodiments, the administration of the ASBTI results in a reduction of the CSS score to zero. In various embodiments, the administration of the ASBTI results in a reduction in the CSS score, the ITCHRO(OBS) score, the ITCHRO score, or a combination thereof by about or by at least about 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% relative to baseline. In various embodiments, a reduced value relative to baseline of the CSS score, the ITCHRO(OBS) score, the ITCHRO score, or a combination thereof is observed on 10%, 20%, 30%, 40%, 50%, 60%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% of days.

[00167] In some embodiments, patients with a higher baseline ITCHRO(OBS) score demonstrate a greater reduction in the symptom or a change in a disease-relevant laboratory measure than patients having a lower baseline ITCHRO(OBS) score. In some embodiments, patients with a baseline ITCHRO(OBS) score of at least 2, 3, or 4 or an ITCHRO score of at least 4, 5, 6, 7, 8, 9, or 10 have a greater reduction in the symptom or a change in a disease-relevant laboratory measure relative to baseline than a lower reduction in patients having a lower baseline severity of pruritus score. In various embodiments, patients having PSC and baseline ITCHRO scores of at least 4 demonstrate a greater reduction in the symptom or a change in a disease-relevant laboratory measure than patients having a baseline ITCHRO score of less than

4. In various embodiments, the method includes predicting that a patient will have a greater reduction in the symptom or a change in a disease-relevant laboratory measure if a baseline ITCHRO score of the patient is at least 4 as compared to a patient having a baseline ITCHRO score of less than 4. In various embodiments the lower reduction is about or less than about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, or 60% the greater reduction. In various embodiments a difference in the reduction in the symptom or a change in a disease-relevant laboratory measure (i.e., between the greater reduction and the lower reduction) between patients having an ITCHRO score of at least 4 at baseline and patients having an ITCHRO score of less than 4 at baseline is measured at about or at least about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 17 weeks, 18 weeks, 19 weeks, 20 weeks, 21 weeks, 22 weeks, 23 weeks, 24 weeks, 6 months, 25 weeks, 26 weeks, 27 weeks, 28 weeks, 29 weeks, 30 weeks, 31 weeks, 32 weeks, 33 weeks, 34 weeks, 35 weeks, 36 weeks, 37 weeks, 38 weeks, 39 weeks, 40 weeks, 41 weeks, 42 weeks, 43 weeks, 44 weeks, 45 weeks, 46 weeks, 47 weeks, 48 weeks, 49 weeks, 50 weeks, 51 weeks, 52 weeks, 1 year, 13 months, 14 months, 15 months, 16 months, 17 months, 18 months, 19 months, 20 months, 21 months, 22 months, 23 months, 23 months, 2 years, 2.5 years, 3 years, 3.5 years, 4 years, 4.5 years, 5 years, 5.5 years, 6 years, 6.5 years, 7 years, 8 years, 9 years, or 10 years following first administration of the ASBTI at the first dose or at the second dose.

[00168] In various embodiments, reduction in severity of pruritus caused by administration of the ASBTI to the patient is positively correlated with a reduction in sBA concentration in the patient. In various embodiments, a greater reduction in sBA concentration in the patient correlates with a corresponding greater reduction in severity of pruritus.

[00169] In various embodiments, the administration of the ASBTI results in a reduction in serum LDL-C concentration relative to baseline. In some embodiments the serum LDL-C concentration is reduced by about or by at least about 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% relative to baseline.

[00170] In some embodiments, the administration of the ASBTI results in a reduction in serum total cholesterol concentration relative to baseline. In some embodiments, the administration of the ASBTI results in a reduction in serum LDL-C levels relative to baseline. In some embodiments the serum total cholesterol concentration, the serum LDL-C levels, or both is

reduced by about or by at least about 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% relative to baseline. In various embodiments, the administration of the ASBTI results in a reduction in serum total cholesterol concentration, of serum LDL-C levels, or both of about or of at least about 1 mg/dL, 2 mg/dL, 3 mg/dL, 4 mg/dL, 5 mg/dL, 10 mg/dL, 12.5 mg/dL, 15 mg/dL, 20 mg/dL, 30 mg/dL, 40 mg/dL or 50 mg/dL relative to baseline.

[00171] In various embodiments, the administration of the ASBTI results in a decrease in serum autotaxin concentration. In some embodiments, the administration of the ASBTI results in a reduction in autotaxin concentration of about or of at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% relative to baseline.

[00172] In various embodiments, administration of the ASBTI results in an increase in a quality of life inventory score or in a quality of life inventory score related to fatigue. The quality of life inventory score can be a health-related quality of life (HRQoL) score. In some embodiments, the HRQoL score is a PedsQL score. In various embodiments, the administration of the ASBTI results an increase in the PedsQL score or in a PedsQL score related to fatigue of about or of at least about 5%, 10%, 15%, 20%, 25%, 30%, 45%, or 50% relative to baseline.

[00173] In various embodiments, administration of the ASBTI results in a decrease in a xanthoma score relative to baseline. In some embodiments, the xanthoma score is reduced by about or by at least about 2.5%, 5%, 10%, 15%, 20%, 35%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% relative to baseline.

[00174] In various embodiments, the administration of the ASBTI results in the reduction in the symptom or a change in a disease-relevant laboratory measure by about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, 12, days, 13 days, 14 days, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 17 weeks, 18 weeks, 19 weeks, 20 weeks, 21 weeks, 22 weeks, 23 weeks, 24 weeks, 25 weeks, 26 weeks, 27 weeks, 28 weeks, 29 weeks, 30 weeks, 31 weeks, 32 weeks, 33 weeks, 34 weeks, 35 weeks, 36 weeks, 37 weeks, 38 weeks, 39 weeks, 40 weeks, 41 weeks, 42 weeks, 43 weeks, 44 weeks, 45 weeks, 46 weeks, 47 weeks, 48 weeks, 49 weeks, 50 weeks, 51 weeks, 52 weeks, or 1 year.

[00175] In various embodiments, serum bilirubin concentration is at pre-administration baseline levels or at normal levels at about or by about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 1

week, 2 weeks, 3 weeks, 4 weeks, 1 month, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 2 months, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 4 months, 17 weeks, 18 weeks, 19 weeks, 20 weeks, 21 weeks, 22 weeks, 23 weeks, 24 weeks, 25 weeks, 26 weeks, 27 weeks, 28 weeks, 29 weeks, 30 weeks, 31 weeks, 32 weeks, 33 weeks, 34 weeks, 35 weeks, 36 weeks, 37 weeks, 38 weeks, 39 weeks, 40 weeks, 41 weeks, 42 weeks, 43 weeks, 44 weeks, 45 weeks, 46 weeks, 47 weeks, 48 weeks, 49 weeks, 50 weeks, 51 weeks, 52 weeks, or 1 year.

[00176] In various embodiments, serum ALT concentration is at pre-administration baseline levels or at normal levels at about or by about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 4 months, 17 weeks, 18 weeks, 19 weeks, 20 weeks, 21 weeks, 22 weeks, 23 weeks, 24 weeks, 25 weeks, 26 weeks, 27 weeks, 28 weeks, 29 weeks, 30 weeks, 31 weeks, 32 weeks, 33 weeks, 34 weeks, 35 weeks, 36 weeks, 37 weeks, 38 weeks, 39 weeks, 40 weeks, 41 weeks, 42 weeks, 43 weeks, 44 weeks, 45 weeks, 46 weeks, 47 weeks, 48 weeks, 49 weeks, 50 weeks, 51 weeks, 52 weeks, or 1 year. In some embodiments, the administration of the ASBTI results in a reduction in ALT levels relative to baseline of about or of at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, or 15%.

[00177] In various embodiments, serum ALT concentration, serum AST concentration, serum bilirubin concentration, serum conjugated bilirubin concentration, or various combinations thereof are within normal range or at pre-administration baseline levels at about or by about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 4 months, 17 weeks, 18 weeks, 19 weeks, 20 weeks, 21 weeks, 22 weeks, 23 weeks, 24 weeks, 25 weeks, 26 weeks, 27 weeks, 28 weeks, 29 weeks, 30 weeks, 31 weeks, 32 weeks, 33 weeks, 34 weeks, 35 weeks, 36 weeks, 37 weeks, 38 weeks, 39 weeks, 40 weeks, 41 weeks, 42 weeks, 43 weeks, 44 weeks, 45 weeks, 46 weeks, 47 weeks, 48 weeks, 49 weeks, 50 weeks, 51 weeks, 52 weeks, or 1 year. In various embodiments, the administration of the ASBTI does not result in a statistically significant change from baseline in serum bilirubin concentration, serum AST concentration, serum ALT concentration, serum alkaline phosphatase concentration, or some combination thereof for a period of at least about or of about 1 day, 2

days, 3 days, 4 days, 5 days, 6 days, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 17 weeks, 18 weeks, 19 weeks, 20 weeks, 21 weeks, 22 weeks, 23 weeks, 24 weeks, 25 weeks, 26 weeks, 27 weeks, 28 weeks, 29 weeks, 30 weeks, 31 weeks, 32 weeks, 33 weeks, 34 weeks, 35 weeks, 36 weeks, 37 weeks, 38 weeks, 39 weeks, 40 weeks, 41 weeks, 42 weeks, 43 weeks, 44 weeks, 45 weeks, 46 weeks, 47 weeks, 48 weeks, 49 weeks, 50 weeks, 51 weeks, 52 weeks, or 1 year. In various embodiments, for adult patients with an ITCHRO score of at least 4 at baseline, the administration of the ASBTI does not result in a significant change from baseline in serum conjugated bilirubin concentration for a period of at least about or of about 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 11 weeks, 12 weeks, 13 weeks, 14 weeks, 15 weeks, 16 weeks, 17 weeks, 18 weeks, 19 weeks, 20 weeks, 21 weeks, 22 weeks, 23 weeks, 24 weeks, 25 weeks, 26 weeks, 27 weeks, 28 weeks, 29 weeks, 30 weeks, 31 weeks, 32 weeks, 33 weeks, 34 weeks, 35 weeks, 36 weeks, 37 weeks, 38 weeks, 39 weeks, 40 weeks, 41 weeks, 42 weeks, 43 weeks, 44 weeks, 45 weeks, 46 weeks, 47 weeks, 48 weeks, 49 weeks, 50 weeks, 51 weeks, 52 weeks, or 1 year.

[00178] In various embodiments of the above methods of the invention, administration of the ASBTI results in reduction, prevention, amelioration, or elimination of one or more side effects associated with administration of the ASBTI in a subject in need thereof. In various embodiments, the frequency and/or severity of side effects is reduced as compared to the side effects when the ASBTI is administered after ingestion of food, at the same time as food, or mixed with food. In various embodiments, the one or more side effects is diarrhea, loose stools, nausea, gastrointestinal pain, abdominal pain, cramping, anorectal discomfort, or a combination thereof.

[00179] In various embodiments of the above methods of the invention, administration of the ASBTI results in an improvement of the GI tolerability of the ASBTI. In some embodiments, the improvement is as compared to the GI tolerability when the ASBTI is administered at mealtime or immediately after food intake.

[00180] In some embodiments, the GI tolerability is improved by at least 10%, or at least 15%, or at least 20%, or at least 25%, or at least 30%, or at least 35%, or at least 40%, or at least 45%,

or at least 50%, or at least 60%, or at least 70% as compared to the GI tolerability when the ASBTI is administered at mealtime or immediately after food intake.

Dose Modulation

[00181] In various embodiments, the method includes modulating a dosage of the ASBTI administered to the patient. The modulation includes determining the 7 α C4:sBA ratio for the patient at a baseline (e.g., prior to administration of the ASBTI or prior to modulating (e.g., increasing) a dosage of the ASBTI), and further determining the 7 α C4:sBA ratio after administering the ASBTI at a first dose or modulating (e.g., increasing) a dosage amount of the ASBTI to a second dose. If the 7 α C4:sBA ratio does not increase by at least 1, 1.25, 1.5, 1.75, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 500, 750, 1,000, 2,000, 3,000, 4,000, 5,000 or 10,000-fold from baseline, the dose of the ASBTI is increased until the ratio increases at least about 1.25, 1.5, 1.75, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 500, 750, 1,000, 2,000, 3,000, 4,000, 5,000 or 10,000-fold relative to baseline. In various embodiments, the dose of the ASBTI is increased or decreased to achieve and maintain a particular 7 α C4:sBA ratio.

[00182] In various embodiments, the modulating includes increasing a dose of the ASBTI from a first dose level to a second dose level greater than the first dose level if the 7 α C4:sBA ratio initially increases by at least 1, 1.25, 1.5, 1.75, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 500, 750, 1,000, 2,000, 3,000, 4,000, 5,000 or 10,000-fold from baseline and then begins to decrease or decreases to less than 1, 1.25, 1.5, 1.75, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 500, 750, 1,000, 2,000, 3,000, 4,000, 5,000 or 10,000-fold or greater higher than baseline. The dose level is increased until the 7 α C4:sBA ratio increases to at least 1, 1.25, 1.5, 1.75, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 500, 750, 1,000, 2,000, 3,000, 4,000, 5,000 or 10,000-fold from the baseline.

[00183] In some embodiments, the modulation includes administering a first dose of the ASBTI to the patient. If the 7 α C4:sBA ratio does not increase or increase by at least 1, 1.25, 1.5, 1.75, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 500, 750, 1,000, 2,000, 3,000, 4,000, 5,000 or 10,000-fold from baseline, the patient is then administered a second dose of the ASBTI higher than the first dose. The dose administered to the patient continues to be increased until the 7 α C4:sBA ratio increases by at least 1, 1.25, 1.5, 1.75, 2, 2.5, 3, 4, 5, 6, 7,

8, 9, 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 500, 750, 1,000, 2,000, 3,000, 4,000, 5,000 or 10,000-fold from baseline.

[00184] In various embodiments, the 7 α C4:sBA ratio is measured about daily, bi-weekly, weekly, bi-monthly, monthly, every two months, every three months, every four months, every five months, every six months, or annually, and the dose of the ASBTI is modulated as necessary each time the ratio is measured.

Pharmaceutical Compositions

[00185] In some embodiments, the ASBTI is administered as a pharmaceutical composition comprising an ASBTI (the composition or the pharmaceutical composition). Any composition described herein can be formulated for ileal, rectal and/or colonic delivery. In more specific embodiments, the composition is formulated for non-systemic or local delivery to the rectum and/or colon. It is to be understood that, as used herein, delivery to the colon includes delivery to sigmoid colon, transverse colon, and/or ascending colon. In still more specific embodiments, the composition is formulated for non-systemic or local delivery to the rectum and/or colon is administered rectally. In other specific embodiments, the composition is formulated for non-systemic or local delivery to the rectum and/or colon is administered orally.

[00186] Provided herein, in certain embodiments, is a pharmaceutical composition comprising a therapeutically effective amount of any compound described herein. In certain instances, the pharmaceutical composition comprises an ASBT inhibitor (e.g., any ASBTI described herein).

[00187] In certain embodiments, pharmaceutical compositions are formulated in a conventional manner using one or more physiologically acceptable carriers including, e.g., excipients and auxiliaries which facilitate processing of the active compounds into preparations which are suitable for pharmaceutical use. In certain embodiments, proper formulation is dependent upon the route of administration chosen. A summary of pharmaceutical compositions described herein is found, for example, in Remington: *The Science and Practice of Pharmacy*, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms*, Mareel Decker, New York, N.Y., 1980; and *Pharmaceutical Dosage Forms and Drug Delivery Systems*, Seventh Ed. (Lippincott Williams & Wilkins 1999), all of which references are incorporated herein in their entirety for all purposes.

[00188] A pharmaceutical composition, as used herein, refers to a mixture of a compound described herein, with other chemical components, such as carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, and/or excipients. In certain instances, the pharmaceutical composition facilitates administration of the compound to an individual or cell. In certain embodiments of practicing the methods of treatment or use provided herein, therapeutically effective amounts of compounds described herein are administered in a pharmaceutical composition to an individual having a disease, disorder, or condition to be treated. In specific embodiments, the individual is a human. As discussed herein, the compounds described herein are either utilized singly or in combination with one or more additional therapeutic agents.

[00189] In certain embodiments, the pharmaceutical formulations described herein are administered to an individual in any manner, including one or more of multiple administration routes, such as, by way of non-limiting example, oral, parenteral (e.g., intravenous, subcutaneous, intramuscular), intranasal, buccal, topical, rectal, or transdermal administration routes.

[00190] In certain embodiments, a pharmaceutical compositions described herein includes one or more compound described herein as an active ingredient in free-acid or free-base form, or in a pharmaceutically acceptable salt form. In some embodiments, the compounds described herein are utilized as an N-oxide or in a crystalline or amorphous form (i.e., a polymorph). In some situations, a compound described herein exists as tautomers. All tautomers are included within the scope of the compounds presented herein. In certain embodiments, a compound described herein exists in an unsolvated or solvated form, wherein solvated forms comprise any pharmaceutically acceptable solvent, e.g., water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be described herein.

[00191] A “carrier” includes, in some embodiments, a pharmaceutically acceptable excipient and is selected on the basis of compatibility with compounds described herein, such as, compounds of any of Formula I-VI, and the release profile properties of the desired dosage form. Exemplary carrier materials include, e.g., binders, suspending agents, disintegration agents, filling agents, surfactants, solubilizers, stabilizers, lubricants, wetting agents, diluents, and the like. See, e.g., *Remington: The Science and Practice of Pharmacy*, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., *Remington's Pharmaceutical Sciences*,

Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H. A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms*, Mareel Decker, New York, N.Y., 1980; and *Pharmaceutical Dosage Forms and Drug Delivery Systems*, Seventh Ed. (Lippincott Williams & Wilkins 1999), all of which references are incorporated herein in their entirety for all purposes.

[00192] Moreover, in certain embodiments, the pharmaceutical compositions described herein are formulated as a dosage form. As such, in some embodiments, provided herein is a dosage form comprising a compound described herein, suitable for administration to an individual. In certain embodiments, suitable dosage forms include, by way of non-limiting example, aqueous oral dispersions, liquids, gels, syrups, elixirs, slurries, suspensions, solid oral dosage forms, aerosols, controlled release formulations, fast melt formulations, effervescent formulations, lyophilized formulations, tablets, powders, pills, dragees, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate release and controlled release formulations.

[00193] In some embodiments, provided herein is a composition comprising an enteroendocrine peptide secretion enhancing agent and, optionally, a pharmaceutically acceptable carrier for alleviating symptoms of cholestasis or a cholestatic liver disease in an individual.

[00194] In certain embodiments, the composition comprises an enteroendocrine peptide secretion enhancing agent and an absorption inhibitor. In specific embodiments, the absorption inhibitor is an inhibitor that inhibits the absorption of the (or at least one of the) specific enteroendocrine peptide secretion enhancing agent with which it is combined. In some embodiments, the composition comprises an enteroendocrine peptide secretion enhancing agent, an absorption inhibitor and a carrier (e.g., an orally suitable carrier or a rectally suitable carrier, depending on the mode of intended administration). In certain embodiments, the composition comprises an enteroendocrine peptide secretion enhancing agent, an absorption inhibitor, a carrier, and one or more of a cholesterol absorption inhibitor, an enteroendocrine peptide, a peptidase inhibitor, a spreading agent, and a wetting agent.

[00195] In other embodiments, the compositions described herein are administered orally for non-systemic delivery of the ASBTI to the rectum and/or colon, including the sigmoid colon, transverse colon, and/or ascending colon. In specific embodiments, compositions formulated for oral administration are, by way of non-limiting example, enterically coated or formulated oral dosage forms, such as, tablets and/or capsules.

Absorption Inhibitors

[00196] In certain embodiments, the composition described herein as being formulated for the non-systemic delivery of ASBTI further includes an absorption inhibitor. As used herein, an absorption inhibitor includes an agent or group of agents that inhibit absorption of a bile acid/salt.

[00197] Suitable bile acid absorption inhibitors (also described herein as absorption inhibiting agents) may include, by way of non-limiting example, anionic exchange matrices, polyamines, quaternary amine containing polymers, quaternary ammonium salts, polyallylamine polymers and copolymers, colesvelam, colesvelam hydrochloride, CholestaGel (N,N,N-trimethyl-6-(2-propenylamino)-1-hexanaminium chloride polymer with (chloromethyl)oxirane, 2-propen-1-amine and N-2-propenyl-1-decanamine hydrochloride), cyclodextrins, chitosan, chitosan derivatives, carbohydrates which bind bile acids, lipids which bind bile acids, proteins and proteinaceous materials which bind bile acids, and antibodies and albumins which bind bile acids. Suitable cyclodextrins include those that bind bile acids/salts such as, by way of non-limiting example, β -cyclodextrin and hydroxypropyl- β -cyclodextrin. Suitable proteins, include those that bind bile acids/salts such as, by way of non-limiting example, bovine serum albumin, egg albumin, casein, α -acid glycoprotein, gelatin, soy proteins, peanut proteins, almond proteins, and wheat vegetable proteins.

[00198] In certain embodiments the absorption inhibitor is cholestyramine. In specific embodiments, cholestyramine is combined with a bile acid. Cholestyramine, an ion exchange resin, is a styrene polymer containing quaternary ammonium groups crosslinked by divinylbenzene. In other embodiments, the absorption inhibitor is colestipol. In specific embodiments, colestipol is combined with a bile acid. Colestipol, an ion exchange resin, is a copolymer of diethylenetriamine and 1-chloro-2,3-epoxypropane.

[00199] In certain embodiments of the compositions and methods described herein the ASBTI is linked to an absorption inhibitor, while in other embodiments the ASBTI and the absorption inhibitor are separate molecular entities.

Cholesterol Absorption Inhibitors

[00200] In certain embodiments, a composition described herein optionally includes at least one cholesterol absorption inhibitor. Suitable cholesterol absorption inhibitors include, by way of non-limiting example, ezetimibe (SCH 58235), ezetimibe analogs, ACT inhibitors, stigmastanyl

phosphorylcholine, stigmastanyl phosphorylcholine analogues, β -lactam cholesterol absorption inhibitors, sulfate polysaccharides, neomycin, plant sponins, plant sterols, phytostanol preparation FM-VP4, Sitostanol, β -sitosterol, acyl-CoA:cholesterol-O-acyltransferase (ACAT) inhibitors, Avasimibe, Implitapide, steroidal glycosides and the like. Suitable enzetimibe analogs include, by way of non-limiting example, SCH 48461, SCH 58053 and the like. Suitable ACT inhibitors include, by way of non-limiting example, trimethoxy fatty acid anilides such as Cl-976, 3-[decyldimethylsilyl]-N-[2-(4-methylphenyl)-1-phenylethyl]-propanamide, melinamide and the like. β -lactam cholesterol absorption inhibitors include, by way of non-limiting example, β R-4S)-1,4-bis-(4-methoxyphenyl)-3- β -phenylpropyl)-2-azetidinone and the like.

Peptidase Inhibitors

[00201] In some embodiments, the compositions described herein optionally include at least one peptidase inhibitor. Such peptidase inhibitors include, but are not limited to, dipeptidyl peptidase-4 inhibitors (DPP-4), neutral endopeptidase inhibitors, and converting enzyme inhibitors. Suitable dipeptidyl peptidase-4 inhibitors (DPP-4) include, by way of non-limiting example, Vildagliptin, 2.S)-1-{2-[β -hydroxy-1-adamantyl)amino]acetyl}pyrrolidine-2-carbonitrile, Sitagliptin, β R)-3-amino-1-[9-(trifluoromethyl)-1,4,7,8-tetrazabicyclo[4.3.0]nona-6,8-dien-4-yl]-4-(2,4,5-trifluorophenyl)butan-1-one, Saxagliptin, and (1S,3S,5S)-2-[(2S)-2-amino-2- β -hydroxy-1-adamantyl)acetyl]-2-azabicyclo[3.1.0]hexane-3-carbonitrile. Such neutral endopeptidase inhibitors include, but are not limited to, Candoxatrilat and Ecadotril.

Spreading Agents/Wetting Agents

[00202] In certain embodiments, the composition described herein optionally comprises a spreading agent. In some embodiments, a spreading agent is utilized to improve spreading of the composition in the colon and/or rectum. Suitable spreading agents include, by way of non-limiting example, hydroxyethylcellulose, hydroxypropylmethyl cellulose, polyethylene glycol, colloidal silicon dioxide, propylene glycol, cyclodextrins, microcrystalline cellulose, polyvinylpyrrolidone, polyoxyethylated glycerides, polycarbophil, di-n-octyl ethers, CetiolTMOE, fatty alcohol polyalkylene glycol ethers, AethoxalTMB), 2-ethylhexyl palmitate, CegesoftTMC 24), and isopropyl fatty acid esters.

[00203] In some embodiments, the compositions described herein optionally comprise a wetting agent. In some embodiments, a wetting agent is utilized to improve wettability of the composition in the colon and rectum. Suitable wetting agents include, by way of non-limiting

example, surfactants. In some embodiments, surfactants are selected from, by way of non-limiting example, polysorbate (e.g., 20 or 80), stearyl heptanoate, caprylic/capric fatty acid esters of saturated fatty alcohols of chain length C₁₂-C₁₈, isostearyl diglycerol isostearic acid, sodium dodecyl sulphate, isopropyl myristate, isopropyl palmitate, and isopropyl myristate/isopropyl stearate/isopropyl palmitate mixture.

Vitamins

[00204] In some embodiments, the methods provided herein further comprise administering one or more vitamins.

[00205] In some embodiments, the vitamin is vitamin A, B1, B2, B3, B5, B6, B7, B9, B12, C, D, E, K, folic acid, pantothenic acid, niacin, riboflavin, thiamine, retinol, beta carotene, pyridoxine, ascorbic acid, cholecalciferol, cyanocobalamin, tocopherols, phylloquinone, menaquinone.

[00206] In some embodiments, the vitamin is a fat-soluble vitamin such as vitamin A, D, E, K, retinol, beta carotene, cholecalciferol, tocopherols, phylloquinone. In a preferred embodiment, the fat-soluble vitamin is tocopherol polyethylene glycol succinate (TPGS).

Bile Acid Sequestrants/Binders

[00207] In some embodiments, a labile bile acid sequestrant is an enzyme dependent bile acid sequestrant. In certain embodiments, the enzyme is a bacterial enzyme. In some embodiments, the enzyme is a bacterial enzyme found in high concentration in human colon or rectum relative to the concentration found in the small intestine. Examples of micro-flora activated systems include dosage forms comprising pectin, galactomannan, and/or Azo hydrogels and/or glycoside conjugates (e.g., conjugates of D-galactoside, β -D-xylopyranoside or the like) of the active agent. Examples of gastrointestinal micro-flora enzymes include bacterial glycosidases such as, for example, D-galactosidase, β -D-glucosidase, α -L-arabinofuranosidase, β -D-xylopyranosidase or the like.

[00208] In certain embodiments, a labile bile acid sequestrant is a time-dependent bile acid sequestrant. In some embodiments, a labile bile acid sequestrant releases a bile acid or is degraded after 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 seconds of sequestration. In some embodiments, a labile bile acid sequestrant releases a bile acid or is degraded after 15, 20, 25, 30, 35, 40, 45, 50, or 55 seconds of sequestration. In some embodiments, a labile bile acid sequestrant releases a bile acid or is degraded after 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 minutes of sequestration. In some

embodiments, a labile bile acid sequestrant releases a bile acid or is degraded after about 15, 20, 25, 30, 35, 45, 50, or 55 minutes of sequestration. In some embodiments, a labile bile acid sequestrant releases a bile acid or is degraded after about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 hours of sequestration. In some embodiments, a labile bile acid sequestrant releases a bile acid or is degraded after 1, 2, or 3 days of sequestration.

[00209] In some embodiments, the labile bile acid sequestrant has a low affinity for bile acid. In certain embodiments, the labile bile acid sequestrant has a high affinity for a primary bile acid and a low affinity for a secondary bile acid.

[00210] In some embodiments, the labile bile acid sequestrant is a pH dependent bile acid sequestrant. In certain embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 6 or below and a low affinity for bile acid at a pH above 6. In certain embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 6.5 or below and a low affinity for bile acid at a pH above 6.5. In certain embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 7 or below and a low affinity for bile acid at a pH above 7. In certain embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 7.1 or below and a low affinity for bile acid at a pH above 7.1. In certain embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 7.2 or below and a low affinity for bile acid at a pH above 7.2. In certain embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 7.3 or below and a low affinity for bile acid at a pH above 7.3. In certain embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 7.4 or below and a low affinity for bile acid at a pH above 7.4. In certain embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 7.5 or below and a low affinity for bile acid at a pH above 7.5. In certain embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 7.6 or below and a low affinity for bile acid at a pH above 7.6. In certain embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 7.7 or below and a low affinity for bile acid at a pH above 7.7. In certain embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 7.8 or below and a low affinity for bile acid at a pH above 7.8. In some embodiments, the pH dependent bile acid sequestrant degrades at a pH above 6. In some embodiments, the pH dependent bile acid sequestrant degrades at a pH above 6.5. In some embodiments, the pH dependent bile acid

sequestrant degrades at a pH above 7. In some embodiments, the pH dependent bile acid sequestrant degrades at a pH above 7.1. In some embodiments, the pH dependent bile acid sequestrant degrades at a pH above 7.2. In some embodiments, the pH dependent bile acid sequestrant degrades at a pH above 7.3. In some embodiments, the pH dependent bile acid sequestrant degrades at a pH above 7.4. In some embodiments, the pH dependent bile acid sequestrant degrades at a pH above 7.5. In some embodiments, the pH dependent bile acid sequestrant degrades at a pH above 7.6. In some embodiments, the pH dependent bile acid sequestrant degrades at a pH above 7.7. In some embodiments, the pH dependent bile acid sequestrant degrades at a pH above 7.8. In some embodiments, the pH dependent bile acid sequestrant degrades at a pH above 7.9.

[00211] In certain embodiments, the labile bile acid sequestrant is lignin or a modified lignin. In some embodiments, the labile bile acid sequestrant is a polycationic polymer or copolymer. In certain embodiments, the labile bile acid sequestrant is a polymer or copolymer comprising one or more N-alkenyl-N-alkylamine residues; one or more N,N,N-trialkyl-N-(N'-alkenylamino)alkyl-azanium residues; one or more N,N,N-trialkyl-N-alkenyl-azanium residues; one or more alkenyl-amine residues; or a combination thereof. In some embodiments, the bile acid binder is cholestyramine, and various compositions including cholestyramine, which are described, for example, in U. S. Pat. Nos. 3,383,281; 3,308,020; 3,769,399; 3,846,541; 3,974,272; 4,172,120; 4,252,790; 4,340,585; 4,814,354; 4,874,744; 4,895,723; 5,695,749; and 6,066, 336, all of which are incorporated herein by reference in their entirety for all purposes. In some embodiments, the bile acid binder is cholestipol or cholesevelam.

Routes of administration, dosage forms, and dosing regimens

[00212] In some embodiments, the compositions described herein, and the compositions administered in the methods described herein are formulated to inhibit bile acid reuptake or reduce serum or hepatic bile acid levels. In certain embodiments, the compositions described herein are formulated for rectal or oral administration. In some embodiments, such formulations are administered rectally or orally, respectively. In some embodiments, the compositions described herein are combined with a device for local delivery of the compositions to the rectum and/or colon (sigmoid colon, transverse colon, or ascending colon). In certain embodiments, for rectal administration the composition described herein are formulated as enemas, rectal gels, rectal foams, rectal aerosols, suppositories, jelly suppositories, or retention enemas. In some

embodiments, for oral administration the compositions described herein are formulated for oral administration and enteric delivery to the colon.

[00213] In certain embodiments, the compositions or methods described herein are non-systemic. In some embodiments, compositions described herein deliver the ASBTI to the distal ileum, colon, and/or rectum and not systemically (e.g., a substantial portion of the enteroendocrine peptide secretion enhancing agent is not systemically absorbed). In some embodiments, oral compositions described herein deliver the ASBTI to the distal ileum, colon, and/or rectum and not systemically (e.g., a substantial portion of the enteroendocrine peptide secretion enhancing agent is not systemically absorbed). In some embodiments, rectal compositions described herein deliver the ASBTI to the distal ileum, colon, and/or rectum and not systemically (e.g., a substantial portion of the enteroendocrine peptide secretion enhancing agent is not systemically absorbed). In certain embodiments, non-systemic compositions described herein deliver less than 90% w/w of the ASBTI systemically. In certain embodiments, non-systemic compositions described herein deliver less than 80% w/w of the ASBTI systemically. In certain embodiments, non-systemic compositions described herein deliver less than 70% w/w of the ASBTI systemically. In certain embodiments, non-systemic compositions described herein deliver less than 60% w/w of the ASBTI systemically. In certain embodiments, non-systemic compositions described herein deliver less than 50% w/w of the ASBTI systemically. In certain embodiments, non-systemic compositions described herein deliver less than 40% w/w of the ASBTI systemically. In certain embodiments, non-systemic compositions described herein deliver less than 30% w/w of the ASBTI systemically. In certain embodiments, non-systemic compositions described herein deliver less than 25% w/w of the ASBTI systemically. In certain embodiments, non-systemic compositions described herein deliver less than 20% w/w of the ASBTI systemically. In certain embodiments, non-systemic compositions described herein deliver less than 15% w/w of the ASBTI systemically. In certain embodiments, non-systemic compositions described herein deliver less than 10% w/w of the ASBTI systemically. In certain embodiments, non-systemic compositions described herein deliver less than 5% w/w of the ASBTI systemically. In some embodiments, systemic absorption is determined in any suitable manner, including the total circulating amount, the amount cleared after administration, or the like.

[00214] In certain embodiments, the compositions and/or formulations described herein are administered at least once a day. In certain embodiments, the formulations containing the ASBTI are administered at least twice a day, while in other embodiments the formulations containing the ASBTI are administered at least three times a day. In certain embodiments, the formulations containing the ASBTI are administered up to five times a day. It is to be understood that in certain embodiments, the dosage regimen of composition containing the ASBTI described herein is determined by considering various factors such as the patient's age, sex, and diet.

[00215] The concentration of the ASBTI administered in the formulations described herein ranges from about 1 mM to about 1 M. In certain embodiments the concentration of the ASBTI administered in the formulations described herein ranges from about 1 mM to about 750 mM. In certain embodiments the concentration of the ASBTI administered in the formulations described herein ranges from about 1 mM to about 500 mM. In certain embodiments the concentration of the ASBTI administered in the formulations described herein ranges from about 5 mM to about 500 mM. In certain embodiments the concentration of the ASBTI administered in the formulations described herein ranges from about 10 mM to about 500 mM. In certain embodiments the concentration of the administered in the formulations described herein ranges from about 25 mM to about 500 mM. In certain embodiments the concentration of the ASBTI administered in the formulations described herein ranges from about 50 mM to about 500 mM. In certain embodiments the concentration of the ASBTI administered in the formulations described herein ranges from about 100 mM to about 500 mM. In certain embodiments the concentration of the ASBTI administered in the formulations described herein ranges from about 200 mM to about 500 mM.

[00216] In certain embodiments, by targeting the distal gastrointestinal tract (e.g., distal ileum, colon, and/or rectum), compositions and methods described herein provide efficacy (e.g., in reducing microbial growth and/or alleviating symptoms of cholestasis or a cholestatic liver disease) with a reduced dose of enteroendocrine peptide secretion enhancing agent (e.g., as compared to an oral dose that does not target the distal gastrointestinal tract).

Oral Administration for Colonic Delivery

[00217] In certain aspects, the composition or formulation containing one or more compounds described herein is orally administered for local delivery of an ASBTI, or a compound described herein to the colon and/or rectum. Unit dosage forms of such compositions include a pill, tablet

or capsules formulated for enteric delivery to colon. In certain embodiments, such pills, tablets or capsule contain the compositions described herein entrapped or embedded in microspheres. In some embodiments, microspheres include, by way of non-limiting example, chitosan microcores HPMC capsules and cellulose acetate butyrate (CAB) microspheres. In certain embodiments, oral dosage forms are prepared using conventional methods known to those in the field of pharmaceutical formulation. For example, in certain embodiments, tablets are manufactured using standard tablet processing procedures and equipment. An exemplary method for forming tablets is by direct compression of a powdered, crystalline or granular composition containing the active agent(s), alone or in combination with one or more carriers, additives, or the like. In alternative embodiments, tablets are prepared using wet-granulation or dry-granulation processes. In some embodiments, tablets are molded rather than compressed, starting with a moist or otherwise tractable material.

[00218] In certain embodiments, tablets prepared for oral administration contain various excipients, including, by way of non-limiting example, binders, diluents, lubricants, disintegrants, fillers, stabilizers, surfactants, preservatives, coloring agents, flavoring agents and the like. In some embodiments, binders are used to impart cohesive qualities to a tablet, ensuring that the tablet remains intact after compression. Suitable binder materials include, by way of non-limiting example, starch (including corn starch and pregelatinized starch), gelatin, sugars (including sucrose, glucose, dextrose and lactose), polyethylene glycol, propylene glycol, waxes, and natural and synthetic gums, e.g., acacia sodium alginate, polyvinylpyrrolidone, cellulosic polymers (including hydroxypropyl cellulose, hydroxypropyl methylcellulose, methyl cellulose, ethyl cellulose, hydroxyethyl cellulose, and the like), Veegum, and combinations thereof. In certain embodiments, diluents are utilized to increase the bulk of the tablet so that a practical size tablet is provided. Suitable diluents include, by way of non-limiting example, dicalcium phosphate, calcium sulfate, lactose, cellulose, kaolin, mannitol, sodium chloride, dry starch, powdered sugar and combinations thereof. In certain embodiments, lubricants are used to facilitate tablet manufacture; examples of suitable lubricants include, by way of non-limiting example, vegetable oils such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil, and oil of theobroma, glycerin, magnesium stearate, calcium stearate, stearic acid and combinations thereof. In some embodiments, disintegrants are used to facilitate disintegration of the tablet, and include, by way of non-limiting example, starches, clays, celluloses, algin, gums, crosslinked

polymers and combinations thereof. Fillers include, by way of non-limiting example, materials such as silicon dioxide, titanium dioxide, alumina, talc, kaolin, powdered cellulose and microcrystalline cellulose, as well as soluble materials such as mannitol, urea, sucrose, lactose, dextrose, sodium chloride and sorbitol. In certain embodiments, stabilizers are used to inhibit or retard drug decomposition reactions that include, by way of example, oxidative reactions. In certain embodiments, surfactants are anionic, cationic, amphoteric or nonionic surface active agents.

[00219] In some embodiments, ASBTIs, or other compounds described herein are orally administered in association with a carrier suitable for delivery to the distal gastrointestinal tract (e.g., distal ileum, colon, and/or rectum).

[00220] In certain embodiments, a composition described herein comprises an ASBTI, or other compounds described herein in association with a matrix (e.g., a matrix comprising hypermellose) that allows for controlled release of an active agent in the distal part of the ileum and/or the colon. In some embodiments, a composition comprises a polymer that is pH sensitive (e.g., a MMX™ matrix from Cosmo Pharmaceuticals) and allows for controlled release of an active agent in the distal part of the ileum. Examples of such pH sensitive polymers suitable for controlled release include and are not limited to polyacrylic polymers (e.g., anionic polymers of methacrylic acid and/or methacrylic acid esters, e.g., Carbopol® polymers) that comprise acidic groups (e.g., —COOH, —SO₃H) and swell in basic pH of the intestine (e.g., pH of about 7 to about 8). In some embodiments, a composition suitable for controlled release in the distal ileum comprises microparticulate active agent (e.g., micronized active agent). In some embodiments, a non-enzymatically degrading poly(dl-lactide-co-glycolide) (PLGA) core is suitable for delivery of an enteroendocrine peptide secretion enhancing agent to the distal ileum. In some embodiments, a dosage form comprising an enteroendocrine peptide secretion enhancing agent is coated with an enteric polymer (e.g., Eudragit® S-100, cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropylmethylcellulose phthalate, anionic polymers of methacrylic acid, methacrylic acid esters or the like) for site specific delivery to the distal ileum and/or the colon. In some embodiments, bacterially activated systems are suitable for targeted delivery to the distal part of the ileum. Examples of micro-flora activated systems include dosage forms comprising pectin, galactomannan, and/or Azo hydrogels and/or glycoside conjugates (e.g., conjugates of D-galactoside, β-D-xylopyranoside or the like) of the active agent. Examples

of gastrointestinal micro-flora enzymes include bacterial glycosidases such as, for example, D-galactosidase, β -D-glucosidase, α -L-arabinofuranosidase, β -D-xylopyranosidase or the like.

[00221] The pharmaceutical composition described herein optionally include an additional therapeutic compound described herein and one or more pharmaceutically acceptable additives such as a compatible carrier, binder, filling agent, suspending agent, flavoring agent, sweetening agent, disintegrating agent, dispersing agent, surfactant, lubricant, colorant, diluent, solubilizer, moistening agent, plasticizer, stabilizer, penetration enhancer, wetting agent, anti-foaming agent, antioxidant, preservative, or one or more combination thereof. In some embodiments, using standard coating procedures, such as those described in Remington's Pharmaceutical Sciences, 20th Edition (2000), a film coating is provided around the formulation of the compound of Formula I. In one embodiment, a compound described herein is in the form of a particle and some or all of the particles of the compound are coated. In certain embodiments, some or all of the particles of a compound described herein are microencapsulated. In some embodiments, the particles of the compound described herein are not microencapsulated and are uncoated.

[00222] In further embodiments, a tablet or capsule comprising an ASBTI or other compounds described herein is film-coated for delivery to targeted sites within the gastrointestinal tract. Examples of enteric film coats include and are not limited to hydroxypropylmethylcellulose, polyvinyl pyrrolidone, hydroxypropyl cellulose, polyethylene glycol 3350, 4500, 8000, methyl cellulose, pseudoethylcellulose, amylopectin and the like.

Pediatric Dosage Formulations and Compositions

[00223] Provided herein, in certain embodiments, is a pediatric dosage formulation or composition comprising a therapeutically effective amount of any compound described herein. In certain instances, the pharmaceutical composition comprises an ASBT inhibitor (e.g., any ASBTI described herein).

[00224] In certain embodiments, suitable dosage forms for the pediatric dosage formulation or composition include, by way of non-limiting example, aqueous or non-aqueous oral dispersions, liquids, gels, syrups, elixirs, slurries, suspensions, solutions, controlled release formulations, fast melt formulations, effervescent formulations, lyophilized formulations, chewable tablets, gummy candy, orally disintegrating tablets, powders for reconstitution as suspension or solution, sprinkle oral powder or granules, dragees, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate release and

controlled release formulations. In some embodiments, provided herein is a pharmaceutical composition wherein the pediatric dosage form is selected from a solution, syrup, suspension, elixir, powder for reconstitution as suspension or solution, dispersible/effervescent tablet, chewable tablet, gummy candy, lollipop, freezer pops, troches, oral thin strips, orally disintegrating tablet, orally disintegrating strip, sachet, and sprinkle oral powder or granules.

[00225] In another aspect, provided herein is a pharmaceutical composition wherein at least one excipient is a flavoring agent or a sweetener. In some embodiments, provided herein is a coating. In some embodiments, provided herein is a taste-masking technology selected from coating of drug particles with a taste-neutral polymer by spray-drying, wet granulation, fluidized bed, and microencapsulation; coating with molten waxes of a mixture of molten waxes and other pharmaceutical adjuvants; entrapment of drug particles by complexation, flocculation or coagulation of an aqueous polymeric dispersion; adsorption of drug particles on resin and inorganic supports; and solid dispersion wherein a drug and one or more taste neutral compounds are melted and cooled, or co-precipitated by a solvent evaporation. In some embodiments, provided herein is a delayed or sustained release formulation comprising drug particles or granules in a rate controlling polymer or matrix.

[00226] Suitable sweeteners include sucrose, glucose, fructose or intense sweeteners, i.e. agents with a high sweetening power when compared to sucrose (e.g. at least 10 times sweeter than sucrose). Suitable intense sweeteners comprise aspartame, saccharin, sodium or potassium or calcium saccharin, acesulfame potassium, sucralose, alitame, xylitol, cyclamate, neomate, neohesperidine dihydrochalcone or mixtures thereof, thaumatin, palatinit, stevioside, rebaudioside, Magnasweet®. The total concentration of the sweeteners may range from effectively zero to about 300 mg/ml based on the liquid composition upon reconstitution.

[00227] In order to increase the palatability of the liquid composition upon reconstitution with an aqueous medium, one or more taste-making agents may be added to the composition in order to mask the taste of the ASBT inhibitor. A taste-masking agent can be a sweetener, a flavoring agent or a combination thereof. The taste-masking agents typically provide up to about 0.1% or 5% by weight of the total pharmaceutical composition. In a preferred embodiment of the present invention, the composition contains both sweetener(s) and flavor(s).

[00228] A flavoring agent herein is a substance capable of enhancing taste or aroma of a composition. Suitable natural or synthetic flavoring agents can be selected from standard

reference books, for example Fenaroli's Handbook of Flavor Ingredients, 3rd edition (1995). Non-limiting examples of flavoring agents and/or sweeteners useful in the formulations described herein, include, e.g., acacia syrup, acesulfame K, alitame, anise, apple, aspartame, banana, Bavarian cream, berry, black currant, butterscotch, calcium citrate, camphor, caramel, cherry, cherry cream, chocolate, cinnamon, bubble gum, citrus, citrus punch, citrus cream, cotton candy, cocoa, cola, cool cherry, cool citrus, cyclamate, cylamate, dextrose, *eucalyptus*, eugenol, fructose, fruit punch, ginger, glycyrrhetinate, *glycyrrhiza* (licorice) syrup, grape, grapefruit, honey, isomalt, lemon, lime, lemon cream, monoammonium glycyrrhizinate (MagnaSweet®), maltol, mannitol, maple, marshmallow, menthol, mint cream, mixed berry, neohesperidine DC, neotame, orange, pear, peach, peppermint, peppermint cream, Prosweet® Powder, raspberry, root beer, rum, saccharin, safrole, sorbitol, spearmint, spearmint cream, strawberry, strawberry cream, *stevia*, sucralose, sucrose, sodium saccharin, saccharin, aspartame, acesulfame potassium, mannitol, talin, sylitol, sucralose, sorbitol, Swiss cream, tagatose, tangerine, thaumatin, tutti frutti, vanilla, walnut, watermelon, wild cherry, wintergreen, xylitol, or any combination of these flavoring ingredients, e.g., anise-menthol, cherry-anise, cinnamon-orange, cherry-cinnamon, chocolate-mint, honey-lemon, lemon-lime, lemon-mint, menthol-*eucalyptus*, orange-cream, vanilla-mint, and mixtures thereof. Flavoring agents can be used singly or in combinations of two or more. In some embodiments, the aqueous liquid dispersion comprises a sweetening agent or flavoring agent in a concentration ranging from about 0.001% to about 5.0% the volume of the aqueous dispersion. In one embodiment, the aqueous liquid dispersion comprises a sweetening agent or flavoring agent in a concentration ranging from about 0.001% to about 1.0% the volume of the aqueous dispersion. In another embodiment, the aqueous liquid dispersion comprises a sweetening agent or flavoring agent in a concentration ranging from about 0.005% to about 0.5% the volume of the aqueous dispersion. In yet another embodiment, the aqueous liquid dispersion comprises a sweetening agent or flavoring agent in a concentration ranging from about 0.01% to about 1.0% the volume of the aqueous dispersion. In yet another embodiment, the aqueous liquid dispersion comprises a sweetening agent or flavoring agent in a concentration ranging from about 0.01% to about 0.5% the volume of the aqueous dispersion.

[00229] In certain embodiments, a pediatric pharmaceutical composition described herein includes one or more compound described herein as an active ingredient in free-acid or free-base form, or in a pharmaceutically acceptable salt form. In some embodiments, the compounds

described herein are utilized as an N-oxide or in a crystalline or amorphous form (i.e., a polymorph). In some situations, a compound described herein exists as tautomers. All tautomers are included within the scope of the compounds presented herein. In certain embodiments, a compound described herein exists in an unsolvated or solvated form, wherein solvated forms comprise any pharmaceutically acceptable solvent, e.g., water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be described herein.

[00230] A “carrier” for pediatric pharmaceutical compositions includes, in some embodiments, a pharmaceutically acceptable excipient and is selected on the basis of compatibility with compounds described herein, such as, compounds of any of Formula I-VI, and the release profile properties of the desired dosage form. Exemplary carrier materials include, e.g., binders, suspending agents, disintegration agents, filling agents, surfactants, solubilizers, stabilizers, lubricants, wetting agents, diluents, and the like. See, e.g., *Remington: The Science and Practice of Pharmacy*, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pa. 1975; Liberman, H. A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms*, Marcel Decker, New York, N.Y., 1980; and *Pharmaceutical Dosage Forms and Drug Delivery Systems*, Seventh Ed. (Lippincott Williams & Wilkins 1999), all of which references are incorporated herein by reference in their entirety for all purposes.

[00231] Moreover, in certain embodiments, the pediatric pharmaceutical compositions described herein are formulated as a dosage form. As such, in some embodiments, provided herein is a dosage form comprising a compound described herein, suitable for administration to an individual. In certain embodiments, suitable dosage forms include, by way of non-limiting example, aqueous oral dispersions, liquids, gels, syrups, elixirs, slurries, suspensions, solid oral dosage forms, aerosols, controlled release formulations, fast melt formulations, effervescent formulations, lyophilized formulations, tablets, powders, pills, dragees, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate release and controlled release formulations.

[00232] In certain aspects, the pediatric composition or formulation containing one or more compounds described herein is orally administered for local delivery of an ASBTI, or a compound described herein to the colon and/or rectum. Unit dosage forms of such compositions include a pill, tablet or capsules formulated for enteric delivery to colon.

[00233] In some embodiments, ASBTIs, or other compounds described herein are orally administered in association with a carrier suitable for delivery to the distal gastrointestinal tract (e.g., distal ileum, colon, and/or rectum).

[00234] In certain embodiments, a pediatric composition described herein comprises an ASBTI, or other compounds described herein in association with a matrix (e.g., a matrix comprising hypermellose) that allows for controlled release of an active agent in the distal part of the ileum and/or the colon. In some embodiments, a composition comprises a polymer that is pH sensitive (e.g., a MMXTM matrix from Cosmo Pharmaceuticals) and allows for controlled release of an active agent in the distal part of the ileum. Examples of such pH sensitive polymers suitable for controlled release include and are not limited to polyacrylic polymers (e.g., anionic polymers of methacrylic acid and/or methacrylic acid esters, e.g., Carbopol® polymers) that comprise acidic groups (e.g., —COOH, —SO₃H) and swell in basic pH of the intestine (e.g., pH of about 7 to about 8). In some embodiments, a composition suitable for controlled release in the distal ileum comprises microparticulate active agent (e.g., micronized active agent). In some embodiments, a non-enzymatically degrading poly(dl-lactide-co-glycolide) (PLGA) core is suitable for delivery of an enteroendocrine peptide secretion enhancing agent to the distal ileum. In some embodiments, a dosage form comprising an enteroendocrine peptide secretion enhancing agent is coated with an enteric polymer (e.g., Eudragit® S-100, cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropylmethylcellulose phthalate, anionic polymers of methacrylic acid, methacrylic acid esters or the like) for site specific delivery to the distal ileum and/or the colon. In some embodiments, bacterially activated systems are suitable for targeted delivery to the distal part of the ileum. Examples of micro-flora activated systems include dosage forms comprising pectin, galactomannan, and/or Azo hydrogels and/or glycoside conjugates (e.g., conjugates of D-galactoside, β-D-xylopyranoside or the like) of the active agent. Examples of gastrointestinal micro-flora enzymes include bacterial glycosidases such as, for example, D-galactosidase, β-D-glucosidase, α-L-arabinofuranosidase, β-D-xylopyranosidase or the like.

[00235] The pediatric pharmaceutical composition described herein optionally include an additional therapeutic compound described herein and one or more pharmaceutically acceptable additives such as a compatible carrier, binder, filling agent, suspending agent, flavoring agent, sweetening agent, disintegrating agent, dispersing agent, surfactant, lubricant, colorant, diluent, solubilizer, moistening agent, plasticizer, stabilizer, penetration enhancer, wetting agent, anti-

foaming agent, antioxidant, preservative, or one or more combination thereof. In some aspects, using standard coating procedures, such as those described in *Remington's Pharmaceutical Sciences*, 20th Edition (2000), a film coating is provided around the formulation of the compound of Formula I. In one embodiment, a compound described herein is in the form of a particle and some or all of the particles of the compound are coated. In certain embodiments, some or all of the particles of a compound described herein are microencapsulated. In some embodiments, the particles of the compound described herein are not microencapsulated and are uncoated.

Liquid Dosage Forms

[00236] The pharmaceutical liquid dosage forms of the invention may be prepared according to techniques well-known in the art of pharmacy.

[00237] A solution refers to a liquid pharmaceutical formulation wherein the active ingredient is dissolved in the liquid. Pharmaceutical solutions of the invention include syrups and elixirs. A suspension refers to a liquid pharmaceutical formulation wherein the active ingredient is in a precipitate in the liquid.

[00238] In a liquid dosage form, it is desirable to have a particular pH and/or to be maintained within a specific pH range. In order to control the pH, a suitable buffer system can be used. In addition, the buffer system should have sufficient capacity to maintain the desired pH range. Examples of the buffer system useful in the present invention include but are not limited to, citrate buffers, phosphate buffers, or any other suitable buffer known in the art. Preferably the buffer system include sodium citrate, potassium citrate, sodium bicarbonate, potassium bicarbonate, sodium dihydrogen phosphate and potassium dihydrogen phosphate, etc. The concentration of the buffer system in the final suspension varies according to factors such as the strength of the buffer system and the pH/pH ranges required for the liquid dosage form. In one embodiment, the concentration is within the range of 0.005 to 0.5 w/v % in the final liquid dosage form.

[00239] The pharmaceutical composition comprising the liquid dosage form of the present invention can also include a suspending/stabilizing agent to prevent settling of the active material. Over time the settling could lead to caking of the active to the inside walls of the product pack, leading to difficulties with redispersion and accurate dispensing. Suitable stabilizing agents include but are not limited to, the polysaccharide stabilizers such as xanthan,

guar and tragacanth gums as well as the cellulose derivatives HPMC (hydroxypropyl methylcellulose), methyl cellulose and Avicel RC-591 (microcrystalline cellulose/sodium carboxymethyl cellulose). In another embodiment, polyvinylpyrrolidone (PVP) can also be used as a stabilizing agent.

[00240] In addition to the aforementioned components, the ASBTI oral suspension form can also optionally contain other excipients commonly found in pharmaceutical compositions such as alternative solvents, taste-masking agents, antioxidants, fillers, acidifiers, enzyme inhibitors and other components as described in Handbook of Pharmaceutical Excipients, Rowe et al., Eds., 4th Edition, Pharmaceutical Press (2003), which is hereby incorporated by reference in its entirety for all purposes.

[00241] Addition of an alternative solvent may help increase solubility of an active ingredient in the liquid dosage form, and consequently the absorption and bioavailability inside the body of a subject. Preferably the alternative solvents include methanol, ethanol or propylene glycol and the like.

[00242] In another aspect, the present invention provides a process for preparing the liquid dosage form. The process comprises steps of bringing ASBTI or its pharmaceutically acceptable salts thereof into mixture with the components including glycerol or syrup or the mixture thereof, a preservative, a buffer system and a suspending/stabilizing agent, etc., in a liquid medium. In general, the liquid dosage form is prepared by uniformly and intimately mixing these various components in the liquid medium. For example, the components such as glycerol or syrup or the mixture thereof, a preservative, a buffer system and a suspending/stabilizing agent, etc., can be dissolved in water to form the aqueous solution, then the active ingredient can be then dispersed in the aqueous solution to form a suspension.

[00243] In some embodiments, the liquid dosage form provided herein can be in a volume of between about 0.1 ml to about 50 ml. In some embodiments, the liquid dosage form provided herein can be in a volume of between about 0.2 ml to about 40 ml. In some embodiments, the liquid dosage form provided herein can be in a volume of between about 0.5 ml to about 30 ml. In some embodiments, the liquid dosage form provided herein can be in a volume of between about 1 ml to about 20 ml. In some embodiments, the liquid dosage form provided herein can be in a volume of between about 0.1 ml to about 20 ml. In some embodiments, the liquid dosage form provided herein can be in a volume of about 0.1 ml to about 20 ml. In some embodiments,

the ASBTI can be in an amount ranging from about 0.001% to about 90% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 0.01% to about 80% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 0.1% to about 70% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 1% to about 60% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 1% to about 50% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 1% to about 40% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 1% to about 30% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 1% to about 20% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 1% to about 10% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 5% to about 70% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 5% to about 60% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 5% to about 50% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 5% to about 40% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 5% to about 30% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 5% to about 20% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 5% to about 10% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 10% to about 50% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 10% to about 40% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 10% to about 30% of the total volume. In some embodiments, the ASBTI can be in an amount ranging from about 10% to about 20% of the total volume. In one embodiment, the resulted liquid dosage form can be in a liquid volume of 0. ml to 30 ml, preferably 0.1 ml to 20 ml, and the active ingredient can be in an amount ranging from about 0.001 mg/ml to about 16 mg/ml, or from about 0.025 mg/ml to about 8 mg/ml, or from about 0.1 mg/ml to about 4 mg/ml, or about 0.25 mg/ml, or about 0.5 mg/ml, or about 1 mg/ml, or about 2 mg/ml, or about 4 mg/ml, or about 5 mg/ml, or about 8 mg/ml, or about 9 mg/ml, or about 10 mg/ml, or about 12 mg/ml, or about 14 mg/ml or about 16 mg/ml.

Bile Acid Sequestrant

[00244] In certain embodiments, an oral formulation for use in any method described herein is, e.g., an ASBTI in association with a labile bile acid sequestrant. A labile bile acid sequestrant is a bile acid sequestrant with a labile affinity for bile acids. In certain embodiments, a bile acid sequestrant described herein is an agent that sequesters (e.g., absorbs or is charged with) bile acid, and/or the salts thereof.

[00245] In specific embodiments, the labile bile acid sequestrant is an agent that sequesters (e.g., absorbs or is charged with) bile acid, and/or the salts thereof, and releases at least a portion of the absorbed or charged bile acid, and/or salts thereof in the distal gastrointestinal tract (e.g., the colon, ascending colon, sigmoid colon, distal colon, rectum, or any combination thereof). In certain embodiments, the labile bile acid sequestrant is an enzyme dependent bile acid sequestrant. In specific embodiments, the enzyme is a bacterial enzyme. In some embodiments, the enzyme is a bacterial enzyme found in high concentration in human colon or rectum relative to the concentration found in the small intestine. Examples of micro-flora activated systems include dosage forms comprising pectin, galactomannan, and/or Azo hydrogels and/or glycoside conjugates (e.g., conjugates of D-galactoside, β -D-xylopyranoside or the like) of the active agent. Examples of gastrointestinal micro-flora enzymes include bacterial glycosidases such as, for example, D-galactosidase, β -D-glucosidase, α -L-arabinofuranosidase, β -D-xylopyranosidase or the like. In some embodiments, the labile bile acid sequestrant is a time dependent bile acid sequestrant (i.e., the bile acid sequesters the bile acid and/or salts thereof and after a time releases at least a portion of the bile acid and/or salts thereof). In some embodiments, a time dependent bile acid sequestrant is an agent that degrades in an aqueous environment over time. In certain embodiments, a labile bile acid sequestrant described herein is a bile acid sequestrant that has a low affinity for bile acid and/or salts thereof, thereby allowing the bile acid sequestrant to continue to sequester bile acid and/or salts thereof in an environ where the bile acids/salts and/or salts thereof are present in high concentration and release them in an environ wherein bile acids/salts and/or salts thereof are present in a lower relative concentration. In some embodiments, the labile bile acid sequestrant has a high affinity for a primary bile acid and a low affinity for a secondary bile acid, allowing the bile acid sequestrant to sequester a primary bile acid or salt thereof and subsequently release a secondary bile acid or salt thereof as the primary bile acid or salt thereof is converted (e.g., metabolized) to the secondary bile acid or salt thereof.

In some embodiments, the labile bile acid sequestrant is a pH dependent bile acid sequestrant. In some embodiments, the pH dependent bile acid sequestrant has a high affinity for bile acid at a pH of 6 or below and a low affinity for bile acid at a pH above 6. In certain embodiments, the pH dependent bile acid sequestrant degrades at a pH above 6.

[00246] In some embodiments, labile bile acid sequestrants described herein include any compound, e.g., a macro-structured compound, that can sequester bile acids/salts and/or salts thereof through any suitable mechanism. For example, in certain embodiments, bile acid sequestrants sequester bile acids/salts and/or salts thereof through ionic interactions, polar interactions, static interactions, hydrophobic interactions, lipophilic interactions, hydrophilic interactions, steric interactions, or the like. In certain embodiments, macrostructured compounds sequester bile acids/salts and/or sequestrants by trapping the bile acids/salts and/or salts thereof in pockets of the macrostructured compounds and, optionally, other interactions, such as those described above. In some embodiments, bile acid sequestrants (e.g., labile bile acid sequestrants) include, by way of non-limiting example, lignin, modified lignin, polymers, polycationic polymers and copolymers, polymers and/or copolymers comprising anyone one or more of N-alkenyl-N-alkylaminc residues; one or more N,N,N-trialkyl-N-(N'-alkenylamino)alkyl-azanium residues; one or more N,N,N-trialkyl-N-alkenyl-azanium residues; one or more alkenyl-amine residues; or a combination thereof, or any combination thereof.

Covalent Linkage of the Drug with a Carrier

[00247] In some embodiments, strategies used for colon targeted delivery include, by way of non-limiting example, covalent linkage of the ASBTI or other compounds described herein to a carrier, coating the dosage form with a pH-sensitive polymer for delivery upon reaching the pH environment of the colon, using redox sensitive polymers, using a time released formulation, utilizing coatings that are specifically degraded by colonic bacteria, using bioadhesive system and using osmotically controlled drug delivery systems.

[00248] In certain embodiments of such oral administration of a composition containing an ASBTI or other compounds described herein involves covalent linking to a carrier wherein upon oral administration the linked moiety remains intact in the stomach and small intestine. Upon entering the colon, the covalent linkage is broken by the change in pH, enzymes, and/or degradation by intestinal microflora. In certain embodiments, the covalent linkage between the ASBTI and the carrier includes, by way of non-limiting example, azo linkage, glycoside

conjugates, glucuronide conjugates, cyclodextrin conjugates, dextran conjugates, and amino-acid conjugates (high hydrophilicity and long chain length of the carrier amino acid).

Coating with Polymers: pH-Sensitive Polymers

[00249] In some embodiments, the oral dosage forms described herein are coated with an enteric coating to facilitate the delivery of an ASBTI or other compounds described herein to the colon and/or rectum. In certain embodiments, an enteric coating is one that remains intact in the low pH environment of the stomach, but readily dissolved when the optimum dissolution pH of the particular coating is reached which depends upon the chemical composition of the enteric coating. The thickness of the coating will depend upon the solubility characteristics of the coating material. In certain embodiments, the coating thicknesses used in such formulations described herein range from about 25 μm to about 200 μm .

[00250] In certain embodiments, the compositions or formulations described herein are coated such that an ASBTI or other compounds described herein of the composition or formulation is delivered to the colon and/or rectum without absorbing at the upper part of the intestine. In a specific embodiment, specific delivery to the colon and/or rectum is achieved by coating of the dosage form with polymers that degrade only in the pH environment of the colon. In alternative embodiments, the composition is coated with an enteric coat that dissolves in the pH of the intestines and an outer layer matrix that slowly erodes in the intestine. In some of such embodiments, the matrix slowly erodes until only a core composition comprising an enteroendocrine peptide secretion enhancing agent (and, in some embodiments, an absorption inhibitor of the agent) is left and the core is delivered to the colon and/or rectum.

[00251] In certain embodiments, pH-dependent systems exploit the progressively increasing pH along the human gastrointestinal tract (GIT) from the stomach (pH 1-2 which increases to 4 during digestion), small intestine (pH 6-7) at the site of digestion and it to 7-8 in the distal ileum. In certain embodiments, dosage forms for oral administration of the compositions described herein are coated with pH-sensitive polymer(s) to provide delayed release and protect the enteroendocrine peptide secretion enhancing agents from gastric fluid. In certain embodiments, such polymers are able to withstand the lower pH values of the stomach and of the proximal part of the small intestine but disintegrate at the neutral or slightly alkaline pH of the terminal ileum and/or ileocecal junction. Thus, in certain embodiments, provided herein is an oral dosage form comprising a coating, the coating comprising a pH-sensitive polymer. In some

embodiments, the polymers used for colon and/or rectum targeting include, by way of non-limiting example, methacrylic acid copolymers, methacrylic acid and methyl methacrylate copolymers, Eudragit L100, Eudragit S100, Eudragit L-30D, Eudragit FS-30D, Eudragit L100-55, polyvinylacetate phthalate, hydroxypropyl ethyl cellulose phthalate, hydroxypropyl methyl cellulose phthalate 50, hydroxypropyl methyl cellulose phthalate 55, cellulose acetate trimellitate, cellulose acetate phthalate and combinations thereof.

[00252] In certain embodiments, oral dosage forms suitable for delivery to the colon and/or rectum comprise a coating that has a biodegradable and/or bacteria degradable polymer or polymers that are degraded by the microflora (bacteria) in the colon. In such biodegradable systems suitable polymers include, by way of non-limiting example, azo polymers, linear-type-segmented polyurethanes containing azo groups, polygalactomannans, pectin, glutaraldehyde crosslinked dextran, polysaccharides, amylose, guar gum, pectin, chitosan, inulin, cyclodextrins, chondroitin sulphate, dextrans, locust bean gum, chondroitin sulphate, chitosan, poly (-caprolactone), polylactic acid and poly(lactic-co-glycolic acid).

[00253] In certain embodiments of such oral administration of compositions containing one or more ASBTIs or other compounds described herein, the compositions are delivered to the colon without absorbing at the upper part of the intestine by coating of the dosage forms with redox sensitive polymers that are degraded by the microflora (bacteria) in the colon. In such biodegradable systems such polymers include, by way of non-limiting example, redox-sensitive polymers containing an azo and/or a disulfide linkage in the backbone.

[00254] In some embodiments, compositions formulated for delivery to the colon and/or rectum are formulated for time-release. In some embodiments, time release formulations resist the acidic environment of the stomach, thereby delaying the release of the enteroendocrine peptide secretion enhancing agents until the dosage form enters the colon and/or rectum.

[00255] In certain embodiments the time released formulations described herein comprise a capsule (comprising an enteroendocrine peptide secretion enhancing agent and an optional absorption inhibitor) with hydrogel plug. In certain embodiments, the capsule and hydrogel plug are covered by a water-soluble cap and the whole unit is coated with an enteric polymer. When the capsule enters the small intestine the enteric coating dissolves and the hydrogels plug swells and dislodges from the capsule after a period of time and the composition is released from the capsule. The amount of hydrogel is used to adjust the period of time to the release the contents.

[00256] In some embodiments, provided herein is an oral dosage form comprising a multi-layered coat, wherein the coat comprises different layers of polymers having different pH-sensitivities. As the coated dosage form moves along GIT the different layers dissolve depending on the pH encountered. Polymers used in such formulations include, by way of non-limiting example, polymethacrylates with appropriate pH dissolution characteristics, Eudragit® RL and Eudragit® RS (inner layer), and Eudragit® FS (outer layer). In other embodiments the dosage form is an enteric coated tablets having an outer shell of hydroxypropylcellulose or hydroxypropylmethylcellulose acetate succinate (HPMCAS).

[00257] In some embodiments, provided herein is an oral dosage form that comprises coat with cellulose butyrate phthalate, cellulose hydrogen phthalate, cellulose propionate phthalate, polyvinyl acetate phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate, dioxypopyl methylcellulose succinate, carboxymethyl ethylcellulose, hydroxypropyl methylcellulose acetate succinate, polymers and copolymers formed from acrylic acid, methacrylic acid, and combinations thereof.

Combination Therapy

[00258] In some embodiments, the methods provided herein comprise administering a compound (e.g., an ASBTI) or composition described herein in combination with one or more additional agents. In some embodiments, the present invention also provides a composition comprising a compound (e.g., an ASBTI) with one or more additional agents.

Fat Soluble Vitamins

[00259] In some embodiments, the methods provided herein further comprise administering one or more vitamins. In some embodiments, the vitamin is vitamin A, B1, B2, B3, B5, B6, B7, B9, B12, C, D, E, K, folic acid, pantothenic acid, niacin, riboflavin, thiamine, retinol, beta carotene, pyridoxine, ascorbic acid, cholecalciferol, cyanocobalamin, tocopherols, phylloquinone, menaquinone.

[00260] In some embodiments, the vitamin is a fat soluble vitamin such as vitamin A, D, E, K, retinol, beta carotene, cholecalciferol, tocopherols, phylloquinone. In a preferred embodiment, the fat soluble vitamin is tocopherol polyethylene glycol succinate (TPGS).

ASBTIs and PPAR agonists

[00261] In various embodiments, the present invention provides methods of use of combinations combinations of ASBTIs with PPAR (peroxisome proliferator-activated receptor) agonists. In various embodiments, the PPAR agonist is a fibrate drug. In some embodiments, the fibrate drug is clofibrate, gemfibrozil, ciprofibrate, benzaifibrate, fenofibrate, or various combinations thereof. In various embodiments, the PPAR agonist is aleglitazar, muraglitazar, tesaglitazar, saroglitazar, GW501516, GW-9662, a thiazolidinedione (TZD), a NSAID (e.g., IBUPROFEN), an indole, or various combinations thereof.

ASBTIs and FXR drugs

[00262] In various embodiments, the present invention provides methods of use of combinations of ASBTIs with farnesoid X receptor (FXR) targeting drugs. In various embodiments, the FXR targeting drug is avermectin B1a, bepridil, fluticasone propionate, GW4064, gliquidone, nicardipine, triclosan, CDCA, ivermectin, chlorotrianisene, tribenoside, mometasone furoate, miconazole, amiodarone, butoconazole, bromocryptine mesylate, pizotifen malate, or various combinations thereof.

Partial External Biliary Diversion (PEBD)

[00263] In some embodiments, the methods provided herein further comprise using partial external biliary diversion as a treatment for patients who have not yet developed cirrhosis. This treatment helps reduce the circulation of bile acids/salts in the liver in order to reduce complications and prevent the need for early transplantation in many patients.

[00264] This surgical technique involves isolating a segment of intestine 10 cm long for use as a biliary conduit (a channel for the passage of bile) from the rest of the intestine. One end of the conduit is attached to the gallbladder and the other end is brought out to the skin to form a stoma (a surgically constructed opening to permit the passage of waste). Partial external biliary diversion may be used for patients who are unresponsive to all medical therapy, especially older, larger patients. This procedure may not be of help to young patients such as infants. Partial external biliary diversion may decrease the intensity of the itching and abnormally low levels of cholesterol in the blood.

ASBTI and Ursodiol

[00265] In some embodiments, an ASBTI is administered in combination with ursodiol or ursodeoxycholic acid, chenodeoxycholic acid, cholic acid, taurocholic acid, ursocholic acid, glycocholic acid, glycodeoxycholic acid, taurodeoxycholic acid, taurocholate, glycochenodeoxycholic acid, tauroursodeoxycholic acid. In some instances, an increase in the concentration of bile acids/salts in the distal intestine induces intestinal regeneration, attenuating intestinal injury, reducing bacterial translocation, inhibiting the release of free radical oxygen, inhibiting production of proinflammatory cytokines, or any combination thereof or any combination thereof.

[00266] In certain embodiments, the patient is administered ursodiol at a daily dose of about or of at least about 5 mg, 10 mg, 15 mg, 20 mg, 25 mg, 30 mg, 35 mg, 36 mg, 40 mg, 45 mg, 50 mg, 55 mg, 60 mg, 65 mg, 70 mg, 75 mg, 80 mg, 85 mg, 90 mg, 95 mg, 100 mg, 150 mg, 200 mg, 250 mg, 300 mg, 350 mg, 400 mg, 450 mg, 500 mg, 550 mg, 600 mg, 650 mg, 700 mg, 750 mg, 800 mg, 850 mg, 900 mg, 950 mg, 1,000 mg, 1,250 mg, 1,500 mg, 1,750 mg, 2,000 mg, 2,250 mg, 2,500 mg, 2,750 mg, or 3,000 mg. In certain embodiments, the patient is administered ursodiol at a daily dose of about or of no more than about 10 mg, 15 mg, 20 mg, 25 mg, 30 mg, 35 mg, 36 mg, 40 mg, 45 mg, 50 mg, 55 mg, 60 mg, 65 mg, 70 mg, 75 mg, 80 mg, 85 mg, 90 mg, 95 mg, 100 mg, 150 mg, 200 mg, 250 mg, 300 mg, 350 mg, 400 mg, 450 mg, 500 mg, 550 mg, 600 mg, 650 mg, 700 mg, 750 mg, 800 mg, 850 mg, 900 mg, 950 mg, 1,000 mg, 1,250 mg, 1,500 mg, 1,750 mg, 2,000 mg, 2,250 mg, 2,500 mg, 2,750 mg, 3,000 mg, or 3,500 mg. In various embodiments, the patient is administered ursodiol at a daily dose of about or of at least about 3 mg to about 300 mg, about 30 mg to about 250 mg, from about 36 mg to about 200 mg, from about 10 mg to about 3000 mg, from about 1000 mg to about 2000 mg, or from about 1500 to about 1900 mg.

[00267] In various embodiments the ursodiol is administered as a tablet. In various embodiments, the ursodiol is administered as a suspension. In various embodiments, the concentration of ursodiol in the suspension is from about 10 mg/mL to about 200 mg/mL, from about 50 mg/mL to about 150 mg/mL, from about 10 mg/mL to about 500 mg/mL, or from about 40 mg/mL to about 60 mg/mL. In various embodiments, the concentration of ursodiol in suspension is about or is at least about 20 mg/mL, 25 mg/mL, 30 mg/mL, 35 mg/mL, 40 mg/mL, 45 mg/mL, 50 mg/mL, 55 mg/mL, 60 mg/mL, 65 mg/mL, 70 mg/mL, 75 mg/mL, or 80

mg/mL. In various embodiments, the concentration of ursodiol in suspension is no more than about 25 mg/mL, 30 mg/mL, 35 mg/mL, 40 mg/mL, 45 mg/mL, 50 mg/mL, 55 mg/mL, 60 mg/mL, 65 mg/mL, 70 mg/mL, 75 mg/mL, 80 mg/mL, or 85 mg/mL.

[00268] An ASBTI and a second active ingredient are used such that the combination is present in a therapeutically effective amount. That therapeutically effective amount arises from the use of a combination of an ASBTI and the other active ingredient (e.g., ursodiol) wherein each is used in a therapeutically effective amount, or by virtue of additive or synergistic effects arising from the combined use, each can also be used in a subclinical therapeutically effective amount, i.e., an amount that, if used alone, provides for reduced effectiveness for the therapeutic purposes noted herein, provided that the combined use is therapeutically effective. In some embodiments, the use of a combination of an ASBTI and any other active ingredient as described herein encompasses combinations where the ASBTI or the other active ingredient is present in a therapeutically effective amount, and the other is present in a subclinical therapeutically effective amount, provided that the combined use is therapeutically effective owing to their additive or synergistic effects. As used herein, the term “additive effect” describes the combined effect of two (or more) pharmaceutically active agents that is equal to the sum of the effect of each agent given alone. A synergistic effect is one in which the combined effect of two (or more) pharmaceutically active agents is greater than the sum of the effect of each agent given alone. Any suitable combination of an ASBTI with one or more of the aforementioned other active ingredients and optionally with one or more other pharmacologically active substances is contemplated as being within the scope of the methods described herein.

[00269] In some embodiments, the particular choice of compounds depends upon the diagnosis of the attending physicians and their judgment of the condition of the individual and the appropriate treatment protocol. The compounds are optionally administered concurrently (e.g., simultaneously, essentially simultaneously or within the same treatment protocol) or sequentially, depending upon the nature of the disease, disorder, or condition, the condition of the individual, and the actual choice of compounds used. In certain instances, the determination of the order of administration, and the number of repetitions of administration of each therapeutic agent during a treatment protocol, is based on an evaluation of the disease being treated and the condition of the individual.

[00270] In some embodiments, therapeutically-effective dosages vary when the drugs are used in treatment combinations. Methods for experimentally determining therapeutically-effective dosages of drugs and other agents for use in combination treatment regimens are described in the literature.

[00271] In some embodiments of the combination therapies described herein, dosages of the co-administered compounds vary depending on the type of co-drug employed, on the specific drug employed, on the disease or condition being treated and so forth. In addition, when co-administered with one or more biologically active agents, the compound provided herein is optionally administered either simultaneously with the biologically active agent(s), or sequentially. In certain instances, if administered sequentially, the attending physician will decide on the appropriate sequence of therapeutic compound described herein in combination with the additional therapeutic agent.

[00272] The multiple therapeutic agents (at least one of which is a therapeutic compound described herein) are optionally administered in any order or even simultaneously. If simultaneously, the multiple therapeutic agents are optionally provided in a single, unified form, or in multiple forms (by way of example only, either as a single pill or as two separate pills). In certain instances, one of the therapeutic agents is optionally given in multiple doses. In other instances, both are optionally given as multiple doses. If not simultaneous, the timing between the multiple doses is any suitable timing; e.g, from more than zero weeks to less than four weeks. In addition, the combination methods, compositions and formulations are not to be limited to the use of only two agents; the use of multiple therapeutic combinations are also envisioned (including two or more compounds described herein).

[00273] In certain embodiments, a dosage regimen to treat, prevent, or ameliorate the condition(s) for which relief is sought, is modified in accordance with a variety of factors. These factors include the disorder from which the subject suffers, as well as the age, weight, sex, diet, and medical condition of the subject. Thus, in various embodiments, the dosage regimen actually employed varies and deviates from the dosage regimens set forth herein.

[00274] In some embodiments, the pharmaceutical agents which make up the combination therapy described herein are provided in a combined dosage form or in separate dosage forms intended for substantially simultaneous administration. In certain embodiments, the pharmaceutical agents that make up the combination therapy are administered sequentially, with

either therapeutic compound being administered by a regimen calling for two-step administration. In some embodiments, two-step administration regimen calls for sequential administration of the active agents or spaced-apart administration of the separate active agents. In certain embodiments, the time period between the multiple administration steps varies, by way of non-limiting example, from a few minutes to several hours, depending upon the properties of each pharmaceutical agent, such as potency, solubility, bioavailability, plasma half-life and kinetic profile of the pharmaceutical agent.

[00275] In certain embodiments, provided herein are combination therapies. In certain embodiments, the compositions described herein comprise an additional therapeutic agent. In some embodiments, the methods described herein comprise administration of a second dosage form comprising an additional therapeutic agent. In certain embodiments, combination therapies the compositions described herein are administered as part of a regimen. Therefore, additional therapeutic agents and/or additional pharmaceutical dosage form can be applied to a patient either directly or indirectly, and concomitantly or sequentially, with the compositions and formulations described herein.

Kits

[00276] In another aspect, provided herein are kits containing a device for oral administration and a pharmaceutical composition as described herein. In certain embodiments the kits include prefilled sachet or bottle for oral administration. In certain embodiments the kits include prefilled syringes for administration of oral enemas.

Release in Distal Ileum and/or Colon

[00277] In certain embodiments, a dosage form comprises a matrix (e.g., a matrix comprising hypermellose) that allows for controlled release of an active agent in the distal jejunum, proximal ileum, distal ileum and/or the colon. In some embodiments, a dosage form comprises a polymer that is pH sensitive (e.g., a MMX™ matrix from Cosmo Pharmaceuticals) and allows for controlled release of an active agent in the ileum and/or the colon. Examples of such pH sensitive polymers suitable for controlled release include and are not limited to polyacrylic polymers (e.g., anionic polymers of methacrylic acid and/or methacrylic acid esters, e.g., Carbopol® polymers) that comprise acidic groups (e.g., —COOH, —SO₃H) and swell in basic pH of the intestine (e.g., pH of about 7 to about 8). In some embodiments, a dosage form suitable for controlled release in the distal ileum comprises microparticulate active agent (e.g.,

micronized active agent). In some embodiments, a non-enzymatically degrading poly(dl-lactide-co-glycolide) (PLGA) core is suitable for delivery of an ASBTI to the distal ileum. In some embodiments, a dosage form comprising an ASBTI is coated with an enteric polymer (e.g., Eudragit® S-100, cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropylmethylcellulose phthalate, anionic polymers of methacrylic acid, methacrylic acid esters or the like) for site specific delivery to the ileum and/or the colon. In some embodiments, bacterially activated systems are suitable for targeted delivery to the ileum. Examples of micro-flora activated systems include dosage forms comprising pectin, galactomannan, and/or Azo hydrogels and/or glycoside conjugates (e.g., conjugates of D-galactoside, β -D-xylopyranoside or the like) of the active agent. Examples of gastrointestinal micro-flora enzymes include bacterial glycosidases such as, for example, D-galactosidase, β -D-glucosidase, α -L-arabinofuranosidase, β -D-xylopyranosidase or the like.

[00278] The pharmaceutical solid dosage forms described herein optionally include an additional therapeutic compound described herein and one or more pharmaceutically acceptable additives such as a compatible carrier, binder, filling agent, suspending agent, flavoring agent, sweetening agent, disintegrating agent, dispersing agent, surfactant, lubricant, colorant, diluent, solubilizer, moistening agent, plasticizer, stabilizer, penetration enhancer, wetting agent, anti-foaming agent, antioxidant, preservative, or one or more combination thereof. In some aspects, using standard coating procedures, such as those described in *Remington's Pharmaceutical Sciences*, 20th Edition (2000), a film coating is provided around the formulation of the ASBTI. In one embodiment, a compound described herein is in the form of a particle and some or all of the particles of the compound are coated. In certain embodiments, some or all of the particles of a compound described herein are microencapsulated. In some embodiments, the particles of the compound described herein are not microencapsulated and are uncoated.

[00279] An ASBT inhibitor may be used in the preparation of medicaments for the prophylactic and/or therapeutic treatment of cholestasis or a cholestatic liver disease. A method for treating any of the diseases or conditions described herein in an individual in need of such treatment, may involve administration of pharmaceutical compositions containing at least one ASBT inhibitor described herein, or a pharmaceutically acceptable salt, pharmaceutically acceptable N-oxide, pharmaceutically active metabolite, pharmaceutically acceptable prodrug, or pharmaceutically acceptable solvate thereof, in therapeutically effective amounts to said individual.

EXAMPLES

[00280] The following examples are provided to further describe some of the embodiments disclosed herein. The examples are intended to illustrate, not to limit, the disclosed embodiments.

Example 1. Dosing apical sodium-dependent bile acid transporter inhibitor (ASBTI) in the fasted state minimizes gastrointestinal adverse effects while maintaining pharmacodynamic effect

[00281] Apical sodium-dependent bile acid transporter inhibitor (ASBTI), also known as ileal bile acid transporter inhibitors (IBATi), decrease enterohepatic circulation of bile acids (BAs) by reducing their reabsorption and increasing fecal BA (fBA) excretion. ASBTI, including maralixibat (MRX; recently approved for the treatment of cholestatic pruritus in patients with Alagille syndrome [ALGS] >1 year of age) and volixibat (VLX), decrease the toxic accumulation of BAs in the liver and mitigate cholestasis. GI adverse effects (AEs; diarrhea, abdominal pain) are a potential side effect of ASBTI, but MRX and VLX can be taken in the fasted state 30 mins before a meal which may minimize GI AEs. The aim of this analysis was to understand the impact of the timing of ASBTI dosing relative to mealtime on the pharmacodynamic (PD) effect and GI AEs.

[00282] AE data from 3 separate phase 1 clinical studies on MRX and VLX in healthy participants were compiled to assess the impact of ASBTI dosing vs mealtime on GI AEs (Table 1, **Figure 1**). In each clinical study, the ASBTI was administered in both a fed and fasted state, and rates of GI AEs were compared. Separately, AE data from placebo (PBO)-controlled trials in patients with ALGS, where MRX was dosed 30 mins prior to a meal, allows a comparison of GI AE rates for MRX vs PBO.

[00283] Across the 3 phase 1 clinical studies there was a lower rate of GI adverse events (AEs) when dosing ASBTI in the fasted state (0%, 0%, and 50% reported a GI AE in Study 1, 2, and 3 respectively) compared to the fed state or at mealtime (75%, 33%, and 100% in Study 1, 2, and 3 respectively), **Figure 2**. In PBO-controlled trials in patients with ALGS where MRX was dosed in the fasted state 30 mins prior to a meal, diarrhea was reported at similar frequencies in patients on drug and PBO (43.6% for MRX vs. 44.4% for PBO).

[00284] The PD impact of ASBTI dosing time to mealtime was investigated in healthy dogs (**Figure 3**). In healthy dogs, MRX significantly increased fBA excretion ($p < 0.01$ vs pretreatment

baseline by paired t-test) regardless of dosing time relative to the daily meal. Maximal increases in fBA excretion were seen when dosing 30 minutes prior to the meal (231% increase) to 4 hours after the meal (229% increase), indicating flexibility in the timing of ASBTI dosing versus mealtime to maintain maximal PD effect (**Figure 4**). $P < 0.01$ versus pre-treatment, by one-tail paired t-test. Data are presented as $\pm$ SEM ($n = 7-8$). % = change compared with pre-treatment value. Groups were dosed at the indicated times. Fecal samples were taken 48 hours before treatment initiation and for the last 72 hours of the 7-day treatment period and analyzed for bile acid content.

[00285] Animal PD data demonstrated herein, showed that there is flexibility in dosing ASBTI relative to meals to increase fBA excretion. Optimal GI tolerability of ASBTI administration in healthy individuals is achieved by dosing in a fasted state. Rates of GI AEs in patients with ALGS dosed with MRX in the fasted state were similar to PBO. Future studies may allow a more detailed elucidation of the relationship between food, ASBTI dosing, GI AEs, and efficacy.

[00286] These data demonstrate that GI tolerability is improved when dosing an ASBTI in the fasted state, versus dosing at mealtime or immediately after food intake. Animal data show that PD effect is maintained regardless of dosing time relative to mealtime, suggesting that efficacy can be maintained while minimizing GI effects.

Table 1: Comparison of the Incidence of GI TEAEs Following ASBTI Administration in the Fed Versus Fasted State

	Volixibat 75 mg		Maralixibat 10 mg		Maralixibat 30 mg	
	Fasted (n=8)	Fed (n=8)	Fasted (n=6)	Fed (n=6)	Fasted (n=12)	Fed (n=12)
Any GI TEAE	0 (0%)	6 (75%)	0 (0%)	2 (33%)	6 (50%)	12 (100%)
Diarrhea	-	4 (50%)	-	1 (17%)	6 (50%)	11 (92%)
Loose stools	-	2 (25%)	-	-	-	-
Nausea	-	-	-	1 (17%)	-	3 (25%)
Abdominal pain	-	-	-	-	1 (8%)	5 (42%)
Anorectal discomfort	-	-	-	-	-	1 (8%)

* * *

[00287] All references cited anywhere within this specification are incorporated herein by reference in their entirety for all purposes.

[00288] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only.

[00289] Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range and each endpoint, unless otherwise indicated herein, and each separate value and endpoint is incorporated into the specification as if it were individually recited herein.

[00290] Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

WHAT IS CLAIMED IS:

1. A method for reducing, minimizing, preventing, ameliorating, or eliminating one or more side effects associated with administration of an apical sodium-dependent bile acid transporter inhibitor (ASBTI) in a subject in need thereof, the method comprising administering a therapeutically effective amount of the ASBTI to the subject before ingestion of food.
2. The method of claim 1 wherein the one or more side effects associated with administration of the ASBTI is reduced, minimized, prevented, ameliorated or eliminated as compared to the side effects when the ASBTI is administered after ingestion of food, at the same time as food, or mixed with food.
3. A method of improving gastrointestinal (GI) tolerability of an ASBTI in a subject in need thereof, the method comprising administering a therapeutically effective amount of the ASBTI to the subject before ingestion of food.
4. The method of claim 3, wherein the improvement of GI tolerability comprises a reduction, minimization, prevention, amelioration, or elimination of one or more GI adverse events.
5. The method of claim 3 or 4, wherein the improvement of GI tolerability comprises a reduction, minimization, prevention, amelioration, or elimination of one or more of diarrhea, loose stools, nausea, abdominal pain, and anorectal discomfort.
6. The method of any one of claims 3-5, wherein the GI tolerability is improved as compared to the GI tolerability when the ASBTI is administered at mealtime or immediately after food intake.
7. The method of any one of claims 3-6, wherein the GI tolerability is improved by at least 10% as compared to the GI tolerability when the ASBTI is administered at mealtime or immediately after food intake.
8. The method of any one of claims 3-7, wherein the GI tolerability is improved by at least

20% as compared to the GI tolerability when the ASBTI is administered at mealtime or immediately after food intake.

9. The method of any one of claims 3-8, wherein the GI tolerability is improved by at least 50% as compared to the GI tolerability when the ASBTI is administered at mealtime or immediately after food intake.

10. A method of treating cholestatic liver disease in a subject in need thereof, the method comprising administering a therapeutically effective amount of an ASBTI to the subject before ingestion of food, wherein the subject experiences a reduction in frequency and/or severity of one or more side effects associated with the administration of the ASBTI.

11. The method of claim 10, wherein the frequency and/or severity of side effects is reduced as compared to the side effects when the ASBTI is administered after ingestion of food, at the same time as food, or mixed with food.

12. The method of claim 10 or 11, wherein the cholestatic liver disease is a pediatric cholestatic liver disease.

13. The method of claim 10 or 11, wherein the cholestatic liver disease is an adult cholestatic liver disease.

14. The method of any one of claims 10-13, wherein the cholestatic liver disease is non-obstructive cholestasis, extrahepatic cholestasis, intrahepatic cholestasis, primary intrahepatic cholestasis, secondary intrahepatic cholestasis, progressive familial intrahepatic cholestasis (PFIC), PFIC type 1, PFIC type 2, PFIC type 3, benign recurrent intrahepatic cholestasis (BRIC), BRIC type 1, BRIC type 2, BRIC type 3, total parenteral nutrition associated cholestasis, paraneoplastic cholestasis, Stauffer syndrome, intrahepatic cholestasis of pregnancy, contraceptive-associated cholestasis, drug-associated cholestasis, infection-associated cholestasis, Dubin-Johnson Syndrome, primary biliary cirrhosis (PBC), primary sclerosing cholangitis (PSC),

gallstone disease, Alagille syndrome, biliary atresia, post-Kasai biliary atresia, post-liver transplantation biliary atresia, post-liver transplantation cholestasis, post-liver transplantation associated liver disease, intestinal failure associated liver disease, bile acid mediated liver injury, MRP2 deficiency syndrome, or neonatal sclerosing cholangitis

15. The method of any one of claims 10-14, wherein the cholestatic liver disease is Alagille syndrome, PFIC, BRIC, PSC, PBC, or biliary atresia.

16. The method of any one of claims 1, 2, and 10-15, wherein the one of more side effects is diarrhea, loose stools, nausea, gastrointestinal pain, abdominal pain, cramping, anorectal discomfort, or a combination thereof.

17. The method of any one of claims claim 1-16, wherein the ASBTI is administered to the subject in a fasted state.

18. The method of any one of claims claim 1-16, wherein the ASBTI is administered less than about 60 minutes before ingestion of food.


19. The method of any one of claims 1-18, wherein the ASBTI is administered less than about 30 minutes before ingestion of food.

20. The method of any one of claims 1-19, wherein the ASBTI is administered immediately prior to the ingestion of food.


21. The method of any one of claims 1-20, wherein the ASBTI is administered at least 4 hours after the last meal.

22. The method of any one of claims 1-21, wherein the ASBTI is administered once daily.

23. The method of any one of claims 1-23, wherein the ASBTI is administered twice daily.
24. The method of any one of claims 1-23, wherein the ASBTI is administered in an amount of about 0.1 mg to about 100 mg per dose.
25. The method of claim 24, wherein the ASBTI is administered in an amount of about 10 mg to about 100 mg per dose.
26. The method of claims 24 or 25, wherein the ASBTI is administered in an amount of about 20 mg to about 80 mg per dose.
27. The method of any one of claims 1-26, wherein the ASBTI is administered in an amount of about 100 ug/kg/day to 1400 ug/kg/day.
28. The method of claim 27, wherein the ASBTI is administered in an amount of about 400 ug/kg/day to about 800 ug/kg/day.
29. The method of any one of claims 1-28, wherein the ASBTI is selected from




(maralixibat) and



(volixibat) or a

30. The method of claim 29, wherein the ASBTI is



(maralixibat chloride).

31. The method of claim 29, wherein the ASBTI is volixibat or a pharmaceutically acceptable salt thereof.
32. The method of any of claims 1-31, wherein the subject has not ingested food for about 0.5 to about 16 hours before the administration of the ASBTI.
33. The method of any of claims 1-32, wherein the subject is a pediatric subject.
34. The method of claim 33, wherein the pediatric subject is 0 to 18 years of age.
35. The method of any of claims 1-26, wherein the ASBTI is administered orally.
36. The method of any of claims 1-35, wherein less than 10% of the ASBTI is systemically absorbed.
37. The method of any of claims 1-35, wherein less than 30% of the ASBTI is systemically absorbed.