





Fig. 2


Fullscale: CD16a-Val158

Human FcγRIIIa-Val158 Binding of
αβTCR HEBE1 IgG mutants




zoomed:

Human FcγRIIIa-Val158 Binding of
αβTCR HEBE1 IgG mutants



Fullscale: CD16a-Phe158

Human FcγRIIIa-Phe158 Binding of
αβTCR HEBE1 IgG mutants



zoomed:

Human FcγRIIIa-Phe158 Binding of
αβTCR HEBE1 IgG mutants




Fig. 3




Fig. 4




Fig. 5

6/84




Fig. 6






Fig. 7

8/84



Western Blot



B² **Biacore**

Unknown	Conc ($\mu\text{g/mL}$)	Relative Response (RU)	Calc Conc (M)	Calc Conc ($\mu\text{g/mL}$)
2C3 in media	3.00	855.75	2.45E-08	3.669
2C3 in media	3.00	860.96	2.47E-08	3.710
2C3 in media	3.00	866.89	2.51E-08	3.758
A114N		146.5	1.60E-09	0.240
Mock		87.84	8.62E-10	0.129
NGT		124.52	1.30E-09	0.196
S298N/Y300S		112.14	1.15E-09	0.172
S440N		146.36	1.60E-09	0.240
S442N		121.21	1.26E-09	0.189
WT (Katya)		148.41	1.63E-09	0.244
WT (Tim)		148.08	1.62E-09	0.244
Y436S		158.72	1.78E-09	0.267
Y436T		84.27	8.23E-10	0.123
Incubated media		87.38	8.57E-10	0.129

Fig. 6

10/84






Fig. 10




Lane	Mutant	mg/ml
1	wt	2.66
2	A114N	2.99
3	Y436S	1.61
4	S298N/Y300S	0.99
5	Y436T	0.41
6	S440N	1.21
7	S442N	1.62
8	NGT	2.21



Fig. 11

**Fig. 12**



13/84

**Fig. 13**

14/84

A.**B.****Fig. 14**

15/84

A.**B.****Fig. 15**

16/84

Sample	k_a ($\times 10^6/\text{Ms}$)	k_d ($\times 10^{-2}/\text{s}$)	R_{max} (RU)	K_D (nM)
GLD52	7.0	1.7	67.0	2.44
WT 2C3	6.0	1.1	64.2	1.75
A114N	4.7	1.1	59.5	2.45
Y436S	5.9	1.0	66.9	1.73
S298N/Y300S	5.7	1.0	63.3	1.80
Y436T	4.8	0.9	65.7	1.95
S440N	5.8	1.1	66.8	1.84
S442N	5.7	1.1	66.2	1.85
NGT	7.9	1.1	70.2	1.35

Fig. 16

17/84

Sample	K_{on} ($\times 10^6 M^{-1} s^{-1}$)	K_{off} ($\times 10^{-2} s^{-1}$)	K_D (nM)
WT 2C3	5.2	1.1	2.1
A114N	5.3	1.3	2.4

Fig. 17






Fig. 18







Fig. 18


21/84

A.


Sample	lot #	Octet Conc ($\mu\text{g/mL}$)
Mock media	11/23/2009	too low
wt 2C3	11/23/2009	2.54
A114N	11/23/2009	2.83
S298N/Y300S	11/23/2009	1.36
S440N	11/23/2009	1.32
S442N	11/23/2009	1.21
Y436S	11/23/2009	1.92
Y436T	11/23/2009	0.34
NGT	11/23/2009	1.90

B.**Fig. 19**

22/84

**Fig. 20**

HIC Analysis of Purified A114N Herceptin



10% Tris-glycine SDS-PAGE




Fig. 21

24/84




Fig. 22

Max. 1.8e4 cps.








Fig. 23

**Fig. 24**


27/84

A.

**Fig. 25**

B.**Aminoxy-Cys-MC-VC-PABC-MMAE**

Molecular Weight: 1510

Aminoxy-Cys-MC-VC-PABC-PEG8-Dol10

Molecular Weight: 1987

Fig. 25




Fig. 25

30/84

Synthesis of aminoxy-Cys-MC-VC-PABC-MMAE and aminoxy-Cys-MC-VC-PABC-PEG8-Dol10




Fig. 26




Fig. 27






Fig. 28

33/84

**Fig. 29**



MC-VC-PABC-MMAE





Fig. 30




Figure 31: please align the legend so that symbols and words line up





Fig. 31

36/84

**Fig. 32**




Fig. 33

LC/MS of S298N/T299A/Y300S (NNAS) heavy chain




Fig. 34

39/84





Fig. 35

40/84

**Fig. 36**




Fig. 37




Fig. 38






Fig. 39

44/84

Lactose aminooxy for native glycans (SAM or GAM):



Exact Mass: 472.19
Molecular Weight: 472.44

**Fig. 40**

45/84

Fig. 41


**bisM6P hexamannose aminoxy
(SAM on Fc glycans):**




Sulfo SMCC

Voyager Spec #1[BP-1565.4, 4504]

bisM6P maleimide



**bisM6P hexamannose maleimide
(Targeting hinge cyteines):**



46/84





Fig. 42

47/84

**Fig. 43**

48/84





Fig. 44



50/84

Samples	MW (kDa)	# Glycan
Control	145.42 $\pm$ 0.24	0.00
M6P conjugate	147.95 $\pm$ 0.09	1.92
Lactose conjugate	146.35 $\pm$ 0.36	1.96

Fig. 46




Fig. 47

Galactose quantitation:




Fig. 48






Fig. 49

54/84

Control:




Lactose maleimide conjugate:


**Fig. 50**

55/84


Control:



M6P conjugate:

**Fig. 51**

56/84



	mol per mol sialic acid
NNAS	2.4 ± 0
Sialylated NNAS	3.7 ± 0.2
Desialylated and galactosylated NNAS	0 ± 0

Fig. 52




Fig. 53







Fig. 54


bisM6P hexasaccharide aminoxy
(SAM):



M6P conjugate at Fc (pAb, 3 glycans)	*		
mAb 3F11		*	
M6P conjugate at Fc (mAb, 1.5 glycans)			*
pAb			*



Fig. 55

60/84



**Fig. 56**


61/84

	mol galactose per mol AB	mol glycopeptide per mol Ab
NNAS	1.4 + 0	0
NNAS desialylated/galactosylated	7.4 + 0.3	0
NNAS conjugate	9.4 + 0	3.1


Fig. 57**MALDI-TOF:****Dionex HPLC:****Fig. 58**

62/84



A**4-12% NuPAGE:****B****Sialic acid analysis:****Fig. 59**

**Fig. 60**


64/84

**Fig. 61**


65/84

A.**4-12% NuPAGE:****B.****Sialic acid analysis:****Fig. 62**


66/84

**Fig. 63**


67/84

**Fig. 64**


68/84

***Fig. 65***


69/84

**Fig. 66**

70/84

***Fig. 67***


71/84



Glycopeptide aminooxy: lactose₃-Cys₃-Gly₄

Fig. 68

72/84

**Fig. 69**





Fig. 70

74/84

	Mol per mol, Mean $\pm$ SD
Wild-type	0.1 $\pm$ 0
A114N	0.5 $\pm$ 0
NNAS	2.3 $\pm$ 0
A114N/NNAS	2.7 $\pm$ 0

Fig. 71

**Fig. 72**

76/84

	Mol per mol, Mean + SD
Wild-type	1.4
A114N	2.7
NNAS	3.4
A114N/NNAS	4.6

Fig. 73

Polyclonal antibody uptake into HepG2 cells






Fig. 74

78/84

**Fig. 75**










Fig. 76


80/84

**Trivalent GalNAc containing glycan****Trivalent Galactose containing glycopeptide****Fig. 77**

Control enzyme: SAM conjugate (~0.8 GalNAc) SAM conjugate (~1.6 GalNAc)



81/84




Trivalent GalNAc containing glycan


Trivalent Galactose containing glycopeptide

Fig. 78

82/84




83/84



Trivalent GalNAc glycan C12

Intact protein MALDI analysis

**Fig. 80**






Fig. 81