


ANTIVIRAL COMPOUNDS WITH A FUSED TRICYCLIC RING

FIELD OF THE INVENTION:

The present invention is related to novel compounds of the general formula I,



their tautomeric forms, their stereoisomers, their analogs, their prodrugs, their isotopes, their N-oxides, their metabolites, their pharmaceutically acceptable salts, polymorphs, solvates, optical isomers, clathrates, co-crystals, combinations with suitable medicament, pharmaceutical compositions containing them, methods of making of the above compounds, and their use as antiviral candidate, more specifically as anti-HCV.

BACKGROUND OF THE INVENTION

Persistent hepatitis C virus (HCV) infection is a major health problem globally affecting ~3% of the world population and is an important contributor to chronic liver disease culminating with liver cirrhosis, hepatocellular carcinoma and liver failure [Szabo E, Lotz G, et al., *Pathol. Oncol. Res.* 2003, 9, 215-221; Hoofnagle JH., *Hepatology* 1997, 26 15S-20]. An estimated 170 million chronic carriers worldwide are at risk of developing liver disease. In the United States alone ~3 million are chronically infected with HCV and the number of HCV related deaths is increasing significantly over the years [Barnes E., WHO factsheet 2010.

http://www.who.int/vaccine_research/diseases/viral_cancers/en/index2.html]. Clinically, chronic infection is often asymptomatic with latent periods lasting for decades before manifestation by which time extensive liver damage has occurred. HCV is spread primarily by unscreened blood transfusions and use of contaminated needles and syringes; the highest risk groups are intravenous drug users and people who received blood transfusions (mainly haemophiliacs) before 1990 when screening for HCV was introduced. Factors that have been reported to influence the rate of HCV disease progression include age (increasing age is associated with more rapid progression), gender (males have more rapid disease progression than females), alcohol consumption (associated with an

increased rate of disease progression), HIV co-infection (associated with a markedly increased rate of disease progression), and fatty liver.

Despite significant efforts, no vaccine exists for HCV and until a year ago the standard therapy for HCV was a combination of pegylated interferon (PEG-IFN) α and weight based ribavirin (RBV), which was inadequate for majority of the patients and therapy associated side effects such as pancytopenia, flu-like symptoms or depression were commonly observed leading to early treatment discontinuation [Fried MW, *et al.*, *N Engl J Med.* 2002, 347, 975-982]. The approval of two direct acting agents (DAA) i.e. 1st generation protease inhibitors, Incivek and Victrelis in May 2011 ushered in the era of specifically targeted HCV therapy [Jesudian AB, Gambarin-Gelwan M and Jacobson IM., *Gastroenterology Hepatol.* **2012**, 8, 91-101].


The combination of above mentioned DAAs, Peg-IFN and RBV (triple therapy) substantially increases the rate of sustained virologic response in treatment naïve and experienced patients. However, a number of issue restrict the usage of these drugs – i) complex treatment algorithms issued by the regulatory bodies; ii) they are restricted to genotype 1; iii) low barrier to resistance mutations and iv) increased cost of therapy leading to limited access to care. Hence, there exists a need for alternative therapeutic strategies that ensure broader genotype coverage, better efficacy, better tolerance and limited selection of resistant HCV variants.

The sequence diversity of HCV is complex with the virus organized into 6 distinct genotypes and over 100 subtypes. Additionally, HCV exists as many closely related viral sequences, termed as quasi-species, in the infected individual, making specific pharmaceutical targeting of HCV proteins challenging due to the rapid evolution of escape mutants. It is increasingly evident that a broad collection of specific, pan genotypic anti-viral drugs targeting multiple essential viral functions, in addition to the current viral therapies will be required for effective global control of HCV.

SUMMARY OF THE INVENTION

According to one aspect of the present invention there is provided a novel compound of the general formula (I), its tautomeric forms, its stereoisomers, its analogs, its prodrugs, its

isotopes, its N-oxides, its metabolites, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates, or its co-crystals,



wherein, R^{1a} , R^1 to R^7 , m , n , Y , ring A , ring D and ring E are as defined hereinbelow.


In another aspect, the present invention provides a pharmaceutical composition, containing the compound of the general formula (I) as defined herein, its tautomeric forms, its stereoisomers, its analogs, its prodrugs, its isotopes, its N-oxides, its metabolites, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates, or its co-crystals in combination with the usual pharmaceutically employed carriers, diluents and the like are useful for the treatment of HCV infection.

DESCRIPTION OF THE INVENTION

HCV is a member of the *Flaviviridae* family of enveloped, positive stranded RNA viruses belonging to the genus *Hepacivirus*. The genome is a single ~9.6kb strand of RNA and consists of one open reading frame that encodes for a polyprotein of ~3000 amino acids flanked by untranslated regions at both 5' and 3' ends. This precursor polyprotein is then processed by viral and cellular proteases to yield 10 separate mature viral proteins critical for replication and assembly of progeny viral particles. The organization of the structural and non-structural proteins in the HCV polyprotein is as follows: C-E1-E2-P7-NS2-NS3-NS4a-NS4b-NS5a-NS5b. The three structural proteins C, E1 and E2 are involved in packaging of the virus and the infectivity cycle. The function of the p7 protein is unknown. Of the non-structural proteins, NS2 is a zinc dependent metalloproteinase that functions in conjunction with a part of NS3 protein. NS3 protein has two catalytic activities associated with it: a serine protease at the N-terminal end which requires NS4A as a cofactor, and an ATPase dependent helicase activity at the C-terminal end. NS5A is a membrane anchored phosphoprotein that is present in basally phosphorylated (56kDa) and hyperphosphorylated (58kDa) forms. Its precise role has not been determined but it has been shown to play a role in RNA binding, multiple host protein interactions, and interferon

resistance. Additionally recent evidence suggests that NS5A plays an important role in replication and infectivity of HCV. The NS5B protein encodes an RNA dependent RNA polymerase activity, key to the generation of progeny viruses. While the pathology of HCV infection mainly affects the liver, the virus is found in other cell types in the body including peripheral blood lymphocytes [Thomson BJ et al., *Clin Microbial Infect.* 2005, 11, 86-94; Moriishi K et al., *Antivir. Chem. Chemother.* 2003, 14, 285-297]. Characterization of the replicase machinery required for HCV RNA synthesis has defined the protease/helicase NS3 protein, the NS4A cofactor, the NS4B integral membrane protein, the NS5A protein and the RNA dependent RNA polymerase NS5B as being its essential components.

Hence, one of the aspects of the present invention is provision of novel compounds of the general formula I,



their tautomeric forms, their isomers, their isotopes, their metabolites, their prodrugs, their pharmaceutically acceptable salts, pharmaceutical compositions containing them, methods of making of the above compounds, and their use as antiviral compounds;

wherein,

Ring 'A' is a 5 to 10 membered saturated carbocycle, the said carbocycle may be a monocyclic system, or a fused- or bridged- or a spiro- bicyclic system;

ring D is selected from substituted- or unsubstituted- 5 to 10 membered carbocycles, substituted- or unsubstituted- 5 to 10 membered heterocycles containing 1 to 3 heteroatoms/groups selected from N(R⁸), S(O)_p, O or C(=O), substituted- or unsubstituted- aromatic carbocycle, and 5 to 6 membered substituted- or unsubstituted- aromatic heterocycle containing heteroatoms selected from N, S or O;

Ring 'E' is selected from 5 to 10 membered carbocycle and 5 to 10 membered heterocycle, the ring 'E' may be monocyclic, fused bicyclic, bridged bicyclic or spiro bicyclic;

Y is selected from CH(R^{2a}) and N(R²)

R^1 is selected from the group consisting of $R^{8a}C(=O)N(R^9)-$, $R^8OC(=O)N(R^9)-$, $R^8(R^9)N-$, $R^8(R^9)NC(=O)N(R^{10})-$, $R^8SO_2N(R^9)-$, $R^{8a}OC(=O)N(R^9)CR^a(R^b)C(=O)N(R^{10})-$, $R^{8a}OC(=O)N(R^9)CR^a(R^b)C(R^c)(R^d)C(=O)N(R^{10})-$, $R^8(R^{9a})NC(=O)N(R^9)CR^a(R^b)C(=O)N(R^{10})-$, and $R^8(R^{9a})NC(=O)N(R^9)CR^a(R^b)C(R^c)(R^d)C(=O)N(R^{10})-$;

R^{1a} is selected independently at each occurrence from the group consisting of halogen, substituted- or unsubstituted- C_{1-6} alkyl, $R^{8a}C(=O)-$, $R^{8b}O-$, $R^{8a}OC(=O)-$, $R^{8a}C(=O)O-$, and $R^8(R^9)NC(=O)-$;

R^2 is selected independently at each occurrence from the group consisting of substituted- or unsubstituted- C_{1-6} alkyl, $R^{8a}C(=O)-$, $R^{8a}SO_2-$, $R^{8a}OC(=O)-$, $R^8(R^9)NC(=O)-$, $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)C(=O)-$, $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)C(R^c)(R^d)C(=O)-$, $R^8(R^9)NC(=O)N(R^{10})C(R^a)(R^b)C(=O)-$, $R^8(R^9)NC(=O)N(R^{10})C(R^a)(R^b)C(R^c)(R^d)C(=O)-$, $R^8SO_2N(R^9)C(R^a)(R^b)C(=O)-$, $R^8SO_2N(R^9)C(R^a)(R^b)C(R^c)(R^d)C(=O)-$, and $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)SO_2-$;

R^{2a} is selected independently at each occurrence from the group consisting of $R^{8a}C(=O)N(R^9)-$, $R^8OC(=O)N(R^9)-$, $R^8(R^9)N-$, $R^8(R^9)NC(=O)N(R^{10})-$, $R^{8a}SO_2N(R^9)-$, $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)C(=O)N(R^{10})-$, $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)C(R^c)(R^d)C(=O)N(R^{10})-$, $R^8(R^{9a})NC(=O)N(R^9)C(R^a)(R^b)C(=O)N(R^{10})-$, and $R^8(R^{9a})NC(=O)N(R^9)CR^a(R^b)C(R^c)(R^d)C(=O)N(R^{10})-$;

R^3 is selected from O and $N(R^{11})$;

R^4 is selected independently at each occurrence from $CR^a(R^b)$, O, $N(R^{11})$, substituted- or unsubstituted carbocycle, substituted- or unsubstituted- heterocycle, substituted- or unsubstituted- arylene and substituted- or unsubstituted- heteroarylene, wherein, substitutions on carbocycle, heterocycle, arylene, and heteroarylene are selected from the group consisting of halogen, substituted- or unsubstituted- C_{1-6} alkyl, and alkyl-O-;

R^5 is selected from hydrogen and substituted- or unsubstituted- alkyl;

R^6 and R^7 are independently selected from the group consisting of hydrogen, halogen, substituted- or unsubstituted- C_{1-6} alkyl, $R^{8b}O-$, and $R^8(R^9)N-$;

wherein, R^8 , R^9 , R^{9a} and R^{10} are independently selected from the group consisting of hydrogen, substituted- or unsubstituted- C_{1-6} alkyl, substituted- or unsubstituted- aryl,

substituted- or unsubstituted- heteroaryl, substituted- or unsubstituted- cycloalkyl, and substituted- or unsubstituted- heterocyclyl;

R^{8a} is selected from the group consisting of substituted- or unsubstituted- C_{1-6} alkyl, substituted- or unsubstituted- aryl, substituted- or unsubstituted- heteroaryl, substituted- or unsubstituted- cycloalkyl, and substituted- or unsubstituted- heterocyclyl;

R^{8b} is selected from the group consisting of hydrogen, substituted- or unsubstituted- C_{1-6} alkyl, perhaloalkyl, substituted- or unsubstituted- aryl, substituted- or unsubstituted- heteroaryl, substituted- or unsubstituted- cycloalkyl, and substituted- or unsubstituted- heterocyclyl;

R^{11} is selected from hydrogen and substituted- or unsubstituted- alkyl group;

R^a , R^b , R^c and R^d , are independently selected from the group consisting of hydrogen, halogen, substituted- or unsubstituted- C_{1-6} alkyl, substituted- or unsubstituted- aryl, substituted- or unsubstituted- heteroaryl, substituted- or unsubstituted- cycloalkyl, and substituted- or unsubstituted- heterocyclyl; R^a , R^b , R^c and R^d together with the carbon atom to which they are attached may take part in forming substituted- or unsubstituted- carbocycle, or substituted- or unsubstituted- heterocycle containing 3 to 10 atoms;

m is an integer ranging between 0 to 2, selected independently at each occurrence;

n is an integer ranging between 0 and 2;

p is an integer ranging between 0 and 2;

when $n = 2$ and R^4 is selected as $CR^a(R^b)$ for both the occurrences, two R^a 's together can form a bond to form an alkenylene linkage or two R^a 's and two R^b 's together can form bonds to form an alkynylene linkage;

'alkyl' may be substituted with 1 to 4 substituents selected from the group consisting of oxo, halogen, cyano, aryl, heteroaryl, cycloalkyl, $R^{12a}SO_2-$, $R^{12b}O-$, $R^{12a}OC(=O)-$, $R^{12a}C(=O)O-$, $(R^{12})(H)NC(=O)-$, $(R^{12})(alkyl)NC(=O)-$, $R^{12a}C(=O)N(H)-$, $R^{12}(H)N-$, $R^{12}(alkyl)N-$, $R^{12}(H)NC(=O)N(H)-$, and $R^{12}(alkyl)NC(=O)N(H)-$;

'cycloalkyl' and 'carbocycle' may be substituted with 1 to 2 substituents selected from the group consisting of oxo, halogen, cyano, aryl, heteroaryl, alkyl, haloalkyl, $R^{12a}C(=O)-$,

$R^{12a}SO_2-$, $R^{12b}O-$, $R^{12a}OC(=O)-$, $R^{12a}C(=O)O-$, $R^{12}(H)NC(=O)-$, $R^{12}(alkyl)NC(=O)-$, $R^{12a}C(=O)N(H)-$, $R^{12}(H)N-$, $R^{12}(alkyl)N-$, $R^{12}(H)NC(=O)N(H)-$, and $R^{12}(alkyl)NC(=O)N(H)-$;

'aryl' or 'aromatic carbocycle', may be substituted with 1 to 2 substituents selected from the group consisting of halogen, nitro, cyano, hydroxy, C_1 to C_6 alkyl, C_3 to C_6 cycloalkyl, 3- to 6- membered heterocycle, C_1 to C_6 perhaloalkyl, alkyl-O-, perhaloalkyl-O-, alkyl-N(alkyl)-, alkyl-N(H)-, H_2N- , alkyl-SO₂-, perhaloalkyl-SO₂-, alkyl-C(=O)N(alkyl)-, alkyl-C(=O)N(H)-, alkyl-N(alkyl)C(=O)-, alkyl-N(H)C(=O)-, $H_2NC(=O)-$, alkyl-N(alkyl)SO₂-, alkyl-N(H)SO₂-, H_2NSO_2- ;

'heteroaryl' or 'aromatic heterocycle' may be substituted with 1 to 2 substituents selected from the group consisting of halogen, nitro, cyano, hydroxy, C_1 to C_6 alkyl, C_3 to C_6 cycloalkyl, 3- to 6- membered heterocycle, C_1 to C_6 perhaloalkyl, alkyl-O-, perhaloalkyl-O-, alkyl-N(alkyl)-, alkyl-N(H)-, H_2N- , alkyl-SO₂-, perhaloalkyl-SO₂-, alkyl-C(=O)N(alkyl)-, alkyl-C(=O)N(H)-, alkyl-N(alkyl)C(=O)-, alkyl-N(H)C(=O)-, $H_2NC(=O)-$, alkyl-N(alkyl)SO₂-, and alkyl-N(H)SO₂-, H_2NSO_2- ;

ring carbons of 'heterocyclyl' and 'heterocycle' may be substituted with 1 to 2 substituents selected from the group consisting of oxo, halogen, nitro, cyano, aryl, heteroaryl, alkyl, $R^{12b}O-$, $R^{12a}OC(=O)-$, $R^{12a}C(=O)O-$, $R^{12}(H)NC(=O)-$, $R^{12}(alkyl)NC(=O)-$, $R^{12a}C(=O)N(H)-$, $R^{12}(H)N-$, $R^{12}(alkyl)N-$, $R^{12}(H)NC(=O)N(H)-$, and $R^{12}(alkyl)NC(=O)N(H)-$; the substituents on ring nitrogen(s) of 'heterocyclyl' and 'heterocycle' are selected from the group consisting of aryl, heteroaryl, alkyl, $R^{12a}C(=O)-$, $R^{12a}SO_2-$, $R^{12a}OC(=O)-$, $R^{12}(H)NC(=O)-$, $R^{12}(alkyl)NC(=O)-$;

R^{12} is selected from hydrogen and alkyl;

R^{12a} is selected as alkyl, perhaloalkyl;

R^{12b} is selected from hydrogen, alkyl, perhaloalkyl.

Whenever a range of the number of atoms in a structure is indicated (e.g., a C_{1-12} , C_{1-8} , C_{1-6} , or C_{1-4} alkyl, alkylamino, etc.), it is specifically contemplated that any sub-range or individual number of carbon atoms falling within the indicated range also can be used. Thus, for instance, the recitation of a range of 1-8 carbon atoms (e.g., C_1-C_8), 1-6 carbon atoms (e.g., C_1-C_6), 1-4 carbon atoms (e.g., C_1-C_4), 1-3 carbon atoms (e.g., C_1-C_3), or 2-8 carbon atoms (e.g., C_2-C_8) as used with respect to any chemical group (e.g., alkyl, alkylamino, etc.) referenced herein encompasses and specifically describes 1, 2, 3, 4, 5, 6,

7, 8, 9, 10, 11, and/or 12 carbon atoms, as appropriate, as well as any sub-range thereof (e.g., 1-2 carbon atoms, 1-3 carbon atoms, 1-4 carbon atoms, 1-5 carbon atoms, 1-6 carbon atoms, 1-7 carbon atoms, 1-8 carbon atoms, 1-9 carbon atoms, 1-10 carbon atoms, 1-11 carbon atoms, 1-12 carbon atoms, 2-3 carbon atoms, 2-4 carbon atoms, 2-5 carbon atoms, 2-6 carbon atoms, 2-7 carbon atoms, 2-8 carbon atoms, 2-9 carbon atoms, 2-10 carbon atoms, 2-11 carbon atoms, 2-12 carbon atoms, 3-4 carbon atoms, 3-5 carbon atoms, 3-6 carbon atoms, 3-7 carbon atoms, 3-8 carbon atoms, 3-9 carbon atoms, 3-10 carbon atoms, 3-11 carbon atoms, 3-12 carbon atoms, 4-5 carbon atoms, 4-6 carbon atoms, 4-7 carbon atoms, 4-8 carbon atoms, 4-9 carbon atoms, 4-10 carbon atoms, 4-11 carbon atoms, and/or 4-12 carbon atoms, etc., as appropriate).

One of the embodiments of the present invention is compound of formula (I) as described above, wherein, ring A is selected as cyclopentane;


In any of the embodiments of the invention described above, R^1 is particularly selected as $R^{8a}OC(=O)N(R^9)CR^a(R^b)C(=O)N(R^{10})-$;

In any of the embodiments of the invention described above, m is particularly selected as 0 at all the occurrences;

In any of the embodiments of the invention described above, Y is particularly selected as $N(R^2)$;

In any of the embodiments of the invention described above, R^2 is particularly selected as $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)C(=O)-$;

In any of the embodiments of the invention described above, ring E is particularly selected as -




In any of the embodiments of the invention described above, R^3 is particularly selected as NH;

In any of the embodiments of the invention described above, R^4 is particularly selected as phenylene, and n is particularly selected as 1;

In any of the embodiments of the invention described above, R^5 , R^6 and R^7 are particularly selected as hydrogen;

In any of the embodiments of the invention described above, Ring D is particularly selected from a six membered carbocycle and seven membered heterocycle containing one heteroatom;

In any of the embodiments of the invention described above, ring A is selected as cyclopentane; R^1 is particularly selected as $R^{8a}OC(=O)N(R^9)CR^a(R^b)C(=O)N(R^{10})-$; m is particularly selected as 0 at all the occurrences; ring E is particularly selected as -



R^3 is particularly selected as NH; R^4 is particularly selected as phenylene, n is particularly selected as 1; R^5 , R^6 and R^7 are particularly selected as hydrogen; Ring D is particularly selected from a six membered carbocycle and seven membered heterocycle containing one heteroatom;

General terms used in formula can be defined as follows; however, the meaning stated should not be interpreted as limiting the scope of the term *per se*.

The term "**alkyl**", as used herein, means a straight chain or branched hydrocarbon containing from 1 to 20 carbon atoms. Preferably the alkyl chain may contain 1 to 10 carbon atoms. More preferably alkyl chain may contain up to 6 carbon atoms. Representative examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, and n-hexyl.

'Alkyl' as defined hereinabove may be substituted with one or more substituents selected independently from the group comprising of oxo, halogen, cyano, aryl, heteroaryl, cycloalkyl, $R^{12a}SO_2-$, $R^{12b}O-$, $R^{12a}OC(=O)-$, $R^{12a}C(=O)O-$, $(R^{12})(H)NC(=O)-$,

(R¹²)(alkyl)NC(=O)-, R^{12a}C(=O)N(H)-, R¹²(H)N-, R¹²(alkyl)N-, R¹²(H)NC(=O)N(H)-, and R¹²(alkyl)NC(=O)N(H)-; wherein R¹² is selected from hydrogen and alkyl; R^{12a} is selected from alkyl and perhaloalkyl; and R^{12b} is selected from hydrogen, alkyl and perhaloalkyl.

The term “**haloalkyl**” used herein means an alkyl group as defined hereinabove wherein at least one of the hydrogen atoms of the said alkyl group is substituted with halogen. The haloalkyl group is exemplified by monofluoromethyl, 1,2-dichloroethyl and the like. The term “**perhaloalkyl**” means an alkyl group as defined hereinabove wherein all the hydrogen atoms of the said alkyl group are substituted with halogen. The perhaloalkyl group is exemplified by trifluoromethyl, pentafluoroethyl and the like.

The term “**cycloalkyl**” as used herein, means a monocyclic, bicyclic, or tricyclic non-aromatic ring system containing from 3 to 14 carbon atoms, preferably monocyclic cycloalkyl ring containing 3 to 6 carbon atoms. Examples of monocyclic ring systems include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Bicyclic ring systems are also exemplified by a bridged monocyclic ring system in which two non-adjacent carbon atoms of the monocyclic ring are linked by an alkylene bridge. Representative examples of bicyclic ring systems include, but are not limited to, bicyclo[3.1.1]heptane, bicyclo[2.2.1]heptane, bicyclo[2.2.2]octane, bicyclo[3.2.2]nonane, bicyclo[3.3.1]nonane, and bicyclo[4.2.1]nonane, bicyclo[3.3.2]decane, bicyclo[3.1.0]hexane, bicyclo[4.1.0]heptane, bicyclo[3.2.0]heptanes, octahydro-1H-indene. Tricyclic ring systems are also exemplified by a bicyclic ring system in which two non-adjacent carbon atoms of the bicyclic ring are linked by a bond or an alkylene bridge. Representative examples of tricyclic-ring systems include, but are not limited to, tricyclo[3.3.1.0^{3,7}]nonane and tricyclo[3.3.1.1^{3,7}]decane (adamantane). The term cycloalkyl also include spiro systems wherein one of the ring is annulated on a single carbon atom such ring systems are exemplified by spiro[2.5]octane, spiro[4.5]decane, spiro[bicyclo[4.1.0]heptane-2,1'-cyclopentane], hexahydro-2'H-spiro[cyclopropane-1,1'-pentalene].

The term “**cycloalkenyl**” as used herein, means a cycloalkyl group as defined above containing at least one double bond.

The term "**carbocycle**" as used herein, means a cyclic system made up of carbon atoms, which includes cycloalkane, cycloalkenes and aromatic systems.

Cycloalkyl, cycloalkenyl and carbocycle as defined hereinabove may be substituted with one or more substituents selected independently from the group comprising of oxo, halogen, cyano, aryl, heteroaryl, alkyl, haloalkyl, $R^{12a}C(=O)-$, $R^{12a}SO_2-$, $R^{12b}O-$, $R^{12a}OC(=O)-$, $R^{12a}C(=O)O-$, $R^{12}(H)NC(=O)-$, $R^{12}(alkyl)NC(=O)-$, $R^{12a}C(=O)N(H)-$, $R^{12}(H)N-$, $R^{12}(alkyl)N-$, $R^{12}(H)NC(=O)N(H)-$, and $R^{12}(alkyl)NC(=O)N(H)-$; wherein R^{12} is selected from hydrogen and alkyl; R^{12a} is selected from alkyl and perhaloalkyl; and R^{12b} is selected from hydrogen, alkyl and perhaloalkyl.

The term "**aryl**" refers to a monovalent monocyclic, bicyclic or tricyclic aromatic hydrocarbon ring system. Examples of aryl groups include phenyl, naphthyl, anthracenyl, fluorenyl, indenyl, azulenyl, and the like. Aryl group also include partially saturated bicyclic and tricyclic aromatic hydrocarbons such as tetrahydro-naphthalene. The said aryl group also includes aryl rings fused with heteroaryl or heterocyclic rings such as 2,3-dihydro-benzo[1,4]dioxin-6-yl, 2,3-dihydro-benzo[1,4]dioxin-5-yl, 2,3-dihydro-benzofuran-5-yl, 2,3-dihydro-benzofuran-4-yl, 2,3-dihydro-benzofuran-6-yl, 2,3-dihydro-benzofuran-6-yl, 2,3-dihydro-1H-indol-5-yl, 2,3-dihydro-1H-indol-4-yl, 2,3-dihydro-1H-indol-6-yl, 2,3-dihydro-1H-indol-7-yl, benzo[1,3]dioxol-4-yl, benzo[1,3]dioxol-5-yl, 1,2,3,4-tetrahydroquinoliny, 1,2,3,4-tetrahydroisoquinoliny, 2,3-dihydrobenzothien-4-yl, 2-oxoindolin-5-yl.

Aryl as defined hereinabove may be substituted with one or more substituents selected independently from the group comprising of halogen, nitro, cyano, hydroxy, C_1 to C_6 alkyl, C_3 to C_6 cycloalkyl, 3- to 6- membered heterocycle, C_1 to C_6 perhaloalkyl, alkyl-O-, perhaloalkyl-O-, alkyl-N(alkyl)-, alkyl-N(H)-, H_2N- , alkyl-SO₂-, perhaloalkyl-SO₂-, alkyl-C(=O)N(alkyl)-, alkyl-C(=O)N(H)-, alkyl-N(alkyl)C(=O)-, alkyl-N(H)C(=O)-, $H_2NC(=O)-$, alkyl-N(alkyl)SO₂-, alkyl-N(H)SO₂-, H_2NSO_2- .

The term "**heteroaryl**" refers to a 5-14 membered monocyclic, bicyclic, or tricyclic ring system having 1-4 ring heteroatoms selected from O, N, or S, and the remainder ring atoms being carbon (with appropriate hydrogen atoms unless otherwise indicated), wherein at least one ring in the ring system is aromatic. Heteroaryl groups may be optionally substituted with one or more substituents. In one embodiment, 0, 1, 2, 3, or 4

atoms of each ring of a heteroaryl group may be substituted by a substituent. Examples of heteroaryl groups include pyridyl, 1-oxo-pyridyl, furanyl, thienyl, pyrrolyl, oxazolyl, oxadiazolyl, imidazolyl, thiazolyl, isoxazolyl, quinolinyl, pyrazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, triazolyl, thiadiazolyl, isoquinolinyl, benzoxazolyl, benzofuranyl, indoliziny, imidazopyridyl, tetrazolyl, benzimidazolyl, benzothiazolyl, benzothiadiazolyl, benzoxadiazolyl, indolyl, azaindolyl, imidazopyridyl, quinazolinyl, purinyl, pyrrolo[2,3]pyrimidinyl, pyrazolo[3,4]pyrimidinyl, and benzo(b)thienyl, 2,3-thiadiazolyl, 1H-pyrazolo[5,1-c]-1,2,4-triazolyl, pyrrolo[3,4-d]-1,2,3-triazolyl, cyclopentatriazolyl, 3H-pyrrolo[3,4-c] isoxazolyl and the like.

heteroaryl as defined hereinabove may be substituted with one or more substituents selected independently from the group comprising of halogen, nitro, cyano, hydroxy, C₁ to C₆ alkyl, C₃ to C₆ cycloalkyl, 3- to 6- membered heterocycle, C₁ to C₆ perhaloalkyl, alkyl-O-, perhaloalkyl-O-, alkyl-N(alkyl)-, alkyl-N(H)-, H₂N-, alkyl-SO₂-, perhaloalkyl-SO₂-, alkyl-C(=O)N(alkyl)-, alkyl-C(=O)N(H)-, alkyl-N(alkyl)C(=O)-, alkyl-N(H)C(=O)-, H₂NC(=O)-, alkyl-N(alkyl)SO₂-, and alkyl-N(H)SO₂-, H₂NSO₂-.

The term "**heterocycle**" or "heterocyclic" as used herein, means a 'cycloalkyl' group wherein one or more of the carbon atoms replaced by -O-, -S-, -S(O₂)-, -S(O)-, -N(R^m)-, -Si(R^m)Rⁿ-, wherein, R^m and Rⁿ are independently selected from hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, and heterocyclyl. The heterocycle may be connected to the parent molecular moiety through any carbon atom or any nitrogen atom contained within the heterocycle. Representative examples of monocyclic heterocycle include, but are not limited to, azetidiny, azepanyl, aziridinyl, diazepanyl, 1,3-dioxanyl, 1,3-dioxolanyl, 1,3-dithiolanyl, 1,3-dithianyl, imidazoliny, imidazolidiny, isothiazoliny, isothiazolidiny, isoxazoliny, isoxazolidiny, morpholiny, oxadiazoliny, oxadiazolidiny, oxazoliny, oxazolidiny, piperaziny, piperidinyl, pyranyl, pyrazoliny, pyrazolidiny, pyrroliny, pyrrolidinyl, tetrahydrofurany, tetrahydrothienyl, thiadiazoliny, thiadiazolidiny, thiazoliny, thiazolidiny, thiomorpholiny, 1.1-dioxidothiomorpholiny (thiomorpholine sulfone), thiopyranyl, and trithianyl. Representative examples of bicyclic heterocycle include, but are not limited to 1,3-benzodioxolyl, 1,3-benzodithiolyl, 2,3-dihydro-1,4-benzodioxiny, 2,3-dihydro-1-benzofurany, 2,3-dihydro-1-benzothienyl, 2,3-dihydro-1 H-indolyl and 1,2,3,4-tetrahydroquinoliny. The term heterocycle also include bridged heterocyclic systems such as azabicyclo[3.2.1]octane, azabicyclo[3.3.1]nonane and the like.

Heterocyclyl group may be substituted on ring carbons with one or more substituents selected independently from the group comprising of oxo, halogen, nitro, cyano, aryl, hereroaryl, alkyl, $R^{12b}O-$, $R^{12a}OC(=O)-$, $R^{12a}C(=O)O-$, $R^{12}(H)NC(=O)-$, $R^{12}(alkyl)NC(O)-$, $R^{12a}C(=O)N(H)-$, $R^{12}(H)N-$, $R^{12}(alkyl)N-$, $R^{12}(H)NC(=O)N(H)-$, and $R^{12}(alkyl)NC(=O)N(H)-$; wherein R^{12} is selected from hydrogen and alkyl; R^{12a} is selected from alkyl and perhaloalkyl; and R^{12b} is selected from hydrogen, alkyl and perhaloalkyl.

Heterocyclyl group may further be substituted on ring nitrogen(s) with substituents selected from the group comprising of aryl, hereroaryl, alkyl, $R^{12a}C(=O)-$, $R^{12a}SO_2-$, $R^{12a}OC(=O)-$, $R^{12}(H)NC(=O)-$, $R^{12}(alkyl)NC(=O)-$; wherein R^{12} is selected from hydrogen and alkyl; R^{12a} is selected from alkyl and perhaloalkyl; and R^{12b} is selected from hydrogen, alkyl and perhaloalkyl.

The term '**oxo**' means a divalent oxygen ($=O$) attached to the parent group. For example oxo attached to carbon forms a carbonyl, oxo substituted on cyclohexane forms a cyclohexanone, and the like.

The term '**annulated**' means the ring system under consideration is either annulated with another ring at a carbon atom of the cyclic system or across a bond of the cyclic system as in the case of fused or spiro ring systems.

The term '**bridged**' means the ring system under consideration contain an alkylene bridge having 1 to 4 methylene units joining two non adjacent ring atoms.

A compound its stereoisomers, racemates, pharmaceutically acceptable salt thereof as described hereinabove wherein the compound of general formula I is selected from:

1. methyl ((S)-1-(((1R,2S)-2-(5-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-3H-benzo[2,3]oxepino[4,5-d]imidazol-8-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate.


2. methyl ((S)-1-(((1S,2R)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate.
3. methyl ((S)-1-(((1S,2S)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate.
4. methyl ((S)-1-(((1S,2S)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-1H-benzo[2,3]oxepino[4,5-d]imidazol-8-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate.
5. methyl ((S)-1-(((1S,2S)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate.
6. methyl ((S)-1-(((1S,2R)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate.

According to a feature of the present invention, the compounds of general formula I where all the symbols are as defined earlier, can be prepared by methods given in Schemes given below or example. Representative procedures are shown below, however; the disclosure should not be construed to limit the scope of the invention arriving at compound of formula I as disclosed hereinabove.

Scheme 1: The amide coupling between 4-halobenzene-1,2-diamine **1** and 2-amino cyclopentanecarboxylic acid **2** (individual isomers *RR*, *SS*, *RS* and *SR* could be made by

methods known in the literature, Tet. Lett., 2002, 43, 5401-5404; J. Am. Chem. Soc., 1999, 121, 7574-7581; Tet. Asymm., 2008, 19, 2796-2803) might result in the formation of a mixture of amides **3a** and **3b** which can be converted to the benzimidazole compound **4** by cyclization with either a dehydrating agent or under basic or acidic conditions. Intermediate **11** could be synthesized either from aldehyde **5** or from phenol **8** (ref. J. Med. Chem., 2000, 43, 2049-2063; *ibid*, 2005, 48, 7351-7362; WO2009102633 A1). Wittig reaction on **5** using the appropriate base and solvent might lead to the formation of **6** which on reduction with Pt-based reagents amongst others may lead to the formation of the acid **7** which under acidic conditions may lead to the formation of **11**. Alternatively, **11** could be also synthesized from the phenol **8** which on alkylation might lead to the formation of **9** which on acidic/basic hydrolysis could lead to the synthesis of the acid **10**. Cyclization of **10** under acidic conditions might lead to the formation of **11**. On treatment with isoamyl nitrite in an appropriate solvent, oxime **12** might be generated from **11** which upon reaction with Boc-L-prolinal under basic conditions may lead to formation of **13**. Treatment of **13** with triethyl phosphite might lead to the formation of **14**. One-pot Pd-catalyzed C-C coupling of 1,4-diphenylboronic acid/boronate with **14** and **4** or sequential coupling of 1,4-diphenylboronic acid/boronate with **14** and **4** followed by de-protection and amide formation with (S)-2-((methoxycarbonyl)amino)-3-methylbutanoic acid or similar moieties according to coupling methods known in the art might lead to the formation of **16** or compounds of formula I. Similarly, compounds with oxazole fused tricyclic system could be prepared starting from α -halogenated **11** followed by O-alkylation with boc-protected L-proline and cyclization using ammonium acetate.


Similar compounds with general formula as depicted below could be synthesized by following the above protocol.



Scheme 1

Scheme 2: Substituted diamine **17**, prepared by methods well known in the art, could be coupled to **2** by methods known in the art and cyclized (similar to Scheme 1) to generate **18**. De-protection of hydroxyl group of **18** and alkylation under basic conditions with synthon **14** might lead to the synthesis of **19**. De-protection of **19** under acidic conditions and sequential coupling with (S)-2-((methoxycarbonyl)amino)-3-methylbutanoic acid or similar moieties might lead to the formation of **20** or compounds of formula I.


Similar compounds with general formula as depicted below could be synthesized by following the above protocol or similar.




Scheme 2

Scheme 3: Diazotization followed by Sandmeyer reaction on synthon **21** may lead to the formation of **22**. Synthesis of **23** may be accomplished by halogenation of **22** by suitable halogenating agents. O-alkylation of **23** with Boc-L-proline using an appropriate base like triethylamine, 4-methylmorpholine, diisopropylethylamine and solvent like DMF, 1,4-dioxane, THF, acetonitrile might lead to the formation of **24**. Cyclization of intermediate **24** using ammonium acetate might lead to the formation of synthon **25**. Following a similar synthetic sequence as in Scheme 2, formation of **27** might be achieved starting from **18** and **25**.

Similar compounds with general formula as depicted below could be synthesized by following the above protocol.



Compounds of the present invention were prepared using synthetic scheme I provided below:



Scheme I

A further embodiment of the present invention includes pharmaceutical compositions comprising any single compound, a combination of two or more compounds delineated herein, or a pharmaceutically acceptable salt thereof, with a pharmaceutically acceptable carrier or excipient.

Yet a further embodiment of the present invention is a pharmaceutical composition comprising any single compound or a combination of two or more compounds delineated herein, or a pharmaceutically acceptable salt thereof, in combination with one or more agents known in the art, with a pharmaceutically acceptable carrier or excipient.

It will be further appreciated that compounds of the present invention can be administered as the sole active pharmaceutical agent, or used in combination with one or more agents to treat or prevent hepatitis C infections or the symptoms associated with HCV infection. Other agents to be administered in combination with a compound or combination of compounds of the present invention include therapies for diseases caused by HCV infection that suppresses HCV viral replication by direct or indirect mechanisms. These agents include, but not limited to, host immune modulators (for example, interferon-alpha, pegylated interferon-alpha, consensus interferon, interferon-beta, interferon-gamma, CpG oligonucleotides and the like); antiviral compounds that inhibit host cellular functions such as inosine monophosphate dehydrogenase (for example, ribavirin and the like); cytokines that modulate immune function (for example, interleukin 2, interleukin 6, and interleukin 12); a compound that enhances the development of type 1 helper T cell response; interfering RNA; anti-sense RNA; vaccines comprising HCV antigens or antigen adjuvant combinations directed against HCV; agents that interact with host cellular components to block viral protein synthesis by inhibiting the internal ribosome entry site (IRES) initiated translation step of HCV viral replication or to block viral particle maturation and release with agents targeted toward the viroporin family of membrane proteins such as, for example, HCV P7 and the like; and any agent or combination of agents that inhibit the replication of HCV by targeting other proteins of the viral genome involved in the viral replication and/or interfere with the function of other viral targets, such as inhibitors of NS3/NS4A protease, NS3 helicase, NS5B polymerase, NS4A protein and NS5A protein.

According to yet another embodiment, the pharmaceutical compositions of the present invention may further comprise inhibitor(s) of other targets in the HCV life cycle, including, but not limited to, helicase, polymerase, metalloprotease, NS4A protein, NS5A protein, and internal ribosome entry site (IRES).

Accordingly, one embodiment of the present invention is directed to a method for treating or preventing an infection caused by an RNA-containing virus comprising co-administering to a patient in need of such treatment one or more agents selected from the group consisting of a host immune modulator and a second or more antiviral agents, or a combination thereof, with a therapeutically effective amount of a compound or combination of compounds of the present invention, or a pharmaceutically acceptable salt thereof. Examples of the host immune modulator are, but not limited to, interferon-alpha, pegylated-interferon-alpha, interferon-beta, interferon-gamma, a cytokine, a vaccine, and a

vaccine comprising an antigen and an adjuvant, and said second antiviral agent inhibits replication of HCV either by inhibiting host cellular functions associated with viral replication or by targeting proteins of the viral genome. Example of the RNA-containing virus includes, but not limited to, hepatitis C virus (HCV).

A further embodiment of the present invention is directed to a method of treating or preventing infection caused by an RNA-containing virus comprising co-administering to a patient in need of such treatment an agent or combination of agents that treat or alleviate symptoms of HCV infection including cirrhosis and inflammation of the liver, with a therapeutically effective amount of a compound or combination of compounds of the present invention, or a pharmaceutically acceptable salt thereof. Example of the RNA-containing virus includes, but not limited to, hepatitis C virus (HCV).

Yet another embodiment of the present invention provides a method of treating or preventing infection caused by an RNA-containing virus comprising co-administering to a patient in need of such treatment one or more agents that treat patients for disease caused by hepatitis B (HBV) infection, with a therapeutically effective amount of a compound or a combination of compounds of the present invention, or a pharmaceutically acceptable salt thereof. An agent that treats patients for disease caused by hepatitis B (HBV) infection may be for example, but not limited thereto, L-deoxythymidine, adefovir, lamivudine or tenfovir, or any combination thereof. Example of the RNA-containing virus includes, but not limited to, hepatitis C virus (HCV).

A further embodiment of the present invention provides a method of treating or preventing infection caused by an RNA-containing virus comprising co-administering to a patient in need of such treatment one or more agents that treat patients for disease caused by human immunodeficiency virus (HIV) infection, with a therapeutically effective amount of a compound or a combination of compounds of the present invention, or a pharmaceutically acceptable salt thereof. The agent that treats patients for disease caused by human immunodeficiency virus (HIV) infection may include, but is not limited thereto, ritonavir, lopinavir, indinavir, nelfmavir, saquinavir, amprenavir, atazanavir, tipranavir, TMC-114, fosamprenavir, zidovudine, lamivudine, didanosine, stavudine, tenofovir, zalcitabine, abacavir, efavirenz, nevirapine, delavirdine, TMC-125, L-870812, S-1360, enfuvirtide (T-20) or T-1249, or any combination thereof. Example of the RNA-containing virus includes, but not limited to, hepatitis C virus (HCV).

It can occur that a patient may be co-infected with hepatitis C virus and one or more other viruses, including but not limited to human immunodeficiency virus (HIV), hepatitis A virus (HAV) and hepatitis B virus (HBV). Thus also contemplated is combination therapy to treat such co-infections by co-administering a compound according to the present invention with at least one of an HIV inhibitor, an HAV inhibitor and an HBV inhibitor.

In addition, the present invention provides the use of a compound or a combination of compounds of the invention, or a therapeutically acceptable salt thereof, and one or more agents selected from the group consisting of a host immune modulator and a second or more antiviral agents, or a combination thereof, to prepare a medicament for the treatment of an infection caused by an RNA-containing virus in a patient, particularly hepatitis C virus. Examples of the host immune modulators include, but are not limited to, interferon-alpha, pegylated-interferon-alpha, interferon-beta, interferon-gamma, a cytokine, a vaccine, and a vaccine comprising an antigen and an adjuvant, and said second antiviral agent inhibits replication of HCV either by inhibiting host cellular functions associated with viral replication or by targeting proteins of the viral genome.

When used in the above or other treatments, combination of compound or compounds of the present invention, together with one or more agents as defined herein above, can be employed in pure form or, where such forms exist, in pharmaceutically acceptable salt thereof. Alternatively, such combination of therapeutic agents can be administered as a pharmaceutical composition containing a therapeutically effective amount of the compound or combination of compounds of interest, or their pharmaceutically acceptable salt thereof, in combination with one or more agents as defined hereinabove, and a pharmaceutically acceptable carrier. Such pharmaceutical compositions can be used for inhibiting the replication of an RNA-containing virus, particularly Hepatitis C virus (HCV), by contacting said virus with said pharmaceutical composition. In addition, such compositions are useful for the treatment or prevention of an infection caused by an RNA-containing virus, particularly Hepatitis C virus (HCV).

Hence, a still further embodiment of the invention is directed to a method of treating or preventing infection caused by an RNA-containing virus, particularly a hepatitis C virus (HCV), comprising administering to a patient in need of such treatment a pharmaceutical composition comprising a compound or combination of compounds of the invention or a

pharmaceutically acceptable salt thereof, and one or more agents as defined hereinabove, with a pharmaceutically acceptable carrier.

When administered as a combination, the therapeutic agents can be formulated as separate compositions which are given at the same time or within a predetermined period of time, or the therapeutic agents can be given as a single unit dosage form.

Antiviral agents contemplated for use in such combination therapy include agents (compounds or biologicals) that are effective to inhibit the formation and/or replication of a virus in a mammal, including but not limited to agents that interfere with either host or viral mechanisms necessary for the formation and/or replication of a virus in a mammal. Such agents can be selected from another anti-HCV agent; an HIV inhibitor; an HAV inhibitor; and an HBV inhibitor.

Other agents to be administered in combination with a compound of the present invention include a cytochrome P450 monooxygenase inhibitor, which is expected to inhibit metabolism of the compounds of the invention. Therefore, the cytochrome P450 monooxygenase inhibitor would be in an amount effective to inhibit metabolism of the compounds of the present invention. Accordingly, the CYP inhibitor is administered in an amount such that the bioavailability of the compounds of the present invention is increased in comparison to the bioavailability in the absence of the CYP inhibitor.

The term 'room temperature' used in the specification denotes any temperature ranging between about 20°C to about 40°C, except and otherwise it is specifically mentioned in the specification.

The intermediates and the compounds of the present invention may be obtained in pure form in a manner known *per se*, for example, by distilling off the solvent in vacuum and re-crystallizing the residue obtained from a suitable solvent, such as pentane, diethyl ether, isopropyl ether, chloroform, dichloromethane, ethyl acetate, acetone or their combinations or subjecting it to one of the purification methods, such as column chromatography (e.g., flash chromatography) on a suitable support material such as alumina or silica gel using eluent such as dichloromethane, ethyl acetate, hexane, methanol, acetone and their combinations. Preparative LC-MS method is also used for the purification of molecules described herein.

Salts of compound of formula I can be obtained by dissolving the compound in a suitable solvent, for example in a chlorinated hydrocarbon, such as methyl chloride or chloroform or a low molecular weight aliphatic alcohol, for example, ethanol or isopropanol, which was then treated with the desired acid or base as described in Berge S.M. et al. "Pharmaceutical Salts, a review article in Journal of Pharmaceutical sciences volume 66, page 1-19 (1977)" and in handbook of pharmaceutical salts properties, selection, and use by P.H.Einrich Stahland Camille G.wermuth, Wiley- VCH (2002). Lists of suitable salts can also be found in *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing Company, Easton, PA, 1990, p. 1445, and *Journal of Pharmaceutical Science*, 66, 2-19 (1977). For example, they can be a salt of an alkali metal (e.g., sodium or potassium), alkaline earth metal (e.g., calcium), or ammonium of salt.

The compound of the invention or a composition thereof can potentially be administered as a pharmaceutically acceptable acid-addition, base neutralized or addition salt, formed by reaction with inorganic acids, such as hydrochloric acid, hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, and phosphoric acid, and organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, and fumaric acid, or by reaction with an inorganic base, such as sodium hydroxide, potassium hydroxide. The conversion to a salt is accomplished by treatment of the base compound with at least a stoichiometric amount of an appropriate acid. Typically, the free base is dissolved in an inert organic solvent such as diethyl ether, ethyl acetate, chloroform, ethanol, methanol, and the like, and the acid is added in a similar solvent. The mixture is maintained at a suitable temperature (e.g., between 0 °C and 50 °C). The resulting salt precipitates spontaneously or can be brought out of solution with a less polar solvent.

The stereoisomers of the compounds of formula I of the present invention may be prepared by stereospecific syntheses or resolution of the achiral compound using an optically active amine, acid or complex forming agent, and separating the diastereomeric salt/complex by fractional crystallization or by column chromatography.

The term "prodrug" denotes a derivative of a compound, which derivative, when administered to warm-blooded animals, e.g. humans, is converted into the compound (drug). The enzymatic and/or chemical hydrolytic cleavage of the compounds of the

present invention occurs in such a manner that the proven drug form (parent carboxylic acid drug) is released, and the moiety or moieties split off remain nontoxic or are metabolized so that nontoxic metabolic products are produced. For example, a carboxylic acid group can be esterified, e.g., with a methyl group or ethyl group to yield an ester. When an ester is administered to a subject, the ester is cleaved, enzymatically or non-enzymatically, reductively, oxidatively, or hydrolytically, to reveal the anionic group. An anionic group can be esterified with moieties (e.g., acyloxymethyl esters) which are cleaved to reveal an intermediate compound which subsequently decomposes to yield the active compound.

The prodrugs can be prepared in situ during the isolation and purification of the compounds, or by separately reacting the purified compound with a suitable derivatizing agent. For example, hydroxy groups can be converted into esters via treatment with a carboxylic acid in the presence of a catalyst. Examples of cleavable alcohol prodrug moieties include substituted or unsubstituted, branched or unbranched lower alkyl ester moieties, e.g., ethyl esters, di-lower alkylamino lower-alkyl esters, e.g., dimethylaminoethyl ester, acylamino lower alkyl esters, acyloxy lower alkyl esters (e.g., pivaloyloxymethyl ester), aryl esters, e.g., phenyl ester, aryl-lower alkyl esters, e.g., benzyl ester, optionally substituted, e.g., with methyl, halo, or methoxy substituents aryl and aryl-lower alkyl esters, amides, lower-alkyl amides, di-lwer alkyl amides, and hydroxy amides.

Thus the present invention further provides a pharmaceutical composition, containing the compounds of the general formula (I) as defined above, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates and its co-crystals in combination with the usual pharmaceutically acceptable carriers, diluents and the like.

The pharmaceutically acceptable carrier (or excipient) is preferably one that is chemically inert to the compound of the invention and one that has no detrimental side effects or toxicity under the conditions of use. Such pharmaceutically acceptable carriers preferably include saline (e.g., 0.9% saline), Cremophor EL (which is a derivative of castor oil and ethylene oxide available from Sigma Chemical Co., St. Louis, MO) (e.g., 5% Cremophor EL/5% ethanol/90% saline, 10% Cremophor EL/90% saline, or 50% Cremophor EL/50%

ethanol), propylene glycol (e.g., 40% propylene glycol/10% ethanol/50% water), polyethylene glycol (e.g., 40% PEG 400/60% saline), and alcohol (e.g., 40% ethanol/60% water). A preferred pharmaceutical carrier is polyethylene glycol, such as PEG 400, and particularly a composition comprising 40% PEG 400 and 60% water or saline. The choice of carrier will be determined in part by the particular compound chosen, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of the pharmaceutical composition of the present invention.

The following formulations for oral, aerosol, parenteral, subcutaneous, intravenous, intraarterial, intramuscular, interperitoneal, rectal, and vaginal administration are merely exemplary and are in no way limiting.

The pharmaceutical compositions can be administered parenterally, e.g., intravenously, intraarterially, subcutaneously, intradermally, intrathecally, or intramuscularly. Thus, the invention provides compositions for parenteral administration that comprise a solution of the compound of the invention dissolved or suspended in an acceptable carrier suitable for parenteral administration, including aqueous and non-aqueous, isotonic sterile injection solutions.

Overall, the requirements for effective pharmaceutical carriers for parenteral compositions are well known to those of ordinary skill in the art. See *Pharmaceutics and Pharmacy Practice*, J.B. Lippincott Company, Philadelphia, PA, Banker and Chalmers, eds., pages 238-250 (1982), and *ASHP Handbook on Injectable Drugs*, Toissel, 4th ed., pages 622-630 (1986). Such compositions include solutions containing anti-oxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The compound can be administered in a physiologically acceptable diluent in a pharmaceutical carrier, such as a sterile liquid or mixture of liquids, including water, saline, aqueous dextrose and related sugar solutions, an alcohol, such as ethanol, isopropanol (for example in topical applications), or hexadecyl alcohol, glycols, such as propylene glycol or polyethylene glycol, dimethylsulfoxide, glycerol ketals, such as 2,2-dimethyl-1,3-dioxolane-4-methanol, ethers, such as poly(ethyleneglycol) 400, an oil, a fatty acid, a fatty acid ester or glyceride, or an acetylated fatty acid glyceride with or without the addition of a

pharmaceutically acceptable surfactant, such as a soap or a detergent, suspending agent, such as pectin, carbomers, methylcellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agents and other pharmaceutical adjuvants.

Oils useful in parenteral formulations include petroleum, animal, vegetable, and synthetic oils. Specific examples of oils useful in such formulations include peanut, soybean, sesame, cottonseed, corn, olive, petrolatum, and mineral oil. Suitable fatty acids for use in parenteral formulations include oleic acid, stearic acid, and isostearic acid. Ethyl oleate and isopropyl myristate are examples of suitable fatty acid esters.

Suitable soaps for use in parenteral formulations include fatty alkali metal, ammonium, and triethanolamine salts, and suitable detergents include (a) cationic detergents such as, for example, dimethyl dialkyl ammonium halides, and alkyl pyridinium halides, (b) anionic detergents such as, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates, (c) nonionic detergents such as, for example, fatty amine oxides, fatty acid alkanolamides, and polyoxyethylene polypropylene copolymers, (d) amphoteric detergents such as, for example, alkyl- β -aminopropionates, and 2-alkyl-imidazoline quaternary ammonium salts, and (e) mixtures thereof.

The parenteral formulations typically will contain from about 0.5% or less to about 25% or more by weight of a compound of the invention in solution. Preservatives and buffers can be used. In order to minimize or eliminate irritation at the site of injection, such compositions can contain one or more nonionic surfactants having a hydrophile-lipophile balance (HLB) of from about 12 to about 17. The quantity of surfactant in such formulations will typically range from about 5% to about 15% by weight. Suitable surfactants include polyethylene sorbitan fatty acid esters, such as sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol. The parenteral formulations can be presented in unit-dose or multi-dose sealed containers, such as ampoules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid excipient, for example, water, for injections, immediately prior to use. Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules, and tablets.

Topical formulations, including those that are useful for transdermal drug release, are well known to those of skill in the art and are suitable in the context of the present invention for application to skin.

Formulations suitable for oral administration can consist of (a) liquid solutions, such as an effective amount of a compound of the invention dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a pre-determined amount of the compound of the invention, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e) suitable emulsions. Liquid formulations can include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent. Capsule forms can be of the ordinary hard- or soft-shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium phosphate, and cornstarch. Tablet forms can include one or more of lactose, sucrose, mannitol, corn starch, potato starch, alginic acid, microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants, diluents, buffering agents, disintegrating agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible excipients. Lozenge forms can comprise the compound ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles comprising a compound of the invention in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like containing, in addition to the compound of the invention, such excipients as are known in the art.

An compound of the present invention, alone or in combination with other suitable components, can be made into aerosol formulations to be administered via inhalation. A compound or epimer of the invention is preferably supplied in finely divided form along with a surfactant and propellant. Typical percentages of the compounds of the invention can be about 0.01% to about 20% by weight, preferably about 1% to about 10% by weight. The surfactant must, of course, be nontoxic, and preferably soluble in the propellant. Representative of such surfactants are the esters or partial esters of fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic

anhydride. Mixed esters, such as mixed or natural glycerides can be employed. The surfactant can constitute from about 0.1% to about 20% by weight of the composition, preferably from about 0.25% to about 5%. The balance of the composition is ordinarily propellant. A carrier can also be included as desired, e.g., lecithin, for intranasal delivery. These aerosol formulations can be placed into acceptable pressurized propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like. They also can be formulated as pharmaceuticals for non-pressured preparations, such as in a nebulizer or an atomizer. Such spray formulations can be used to spray mucosa.

Additionally, the compound of the invention can be made into suppositories by mixing with a variety of bases, such as emulsifying bases or water-soluble bases. Formulations suitable for vaginal administration can be presented as pessaries, tampons, creams, gels, pastes, foams, or spray formulas containing, in addition to the compound ingredient, such carriers as are known in the art to be appropriate.

The concentration of the compound in the pharmaceutical formulations can vary, e.g., from less than about 1% to about 10%, to as much as 20% to 50% or more by weight, and can be selected primarily by fluid volumes, and viscosities, in accordance with the particular mode of administration selected.

For example, a typical pharmaceutical composition for intravenous infusion could be made up to contain 250 ml of sterile Ringer's solution, and 100 mg of at least one compound of the invention. Actual methods for preparing parenterally administrable compounds of the invention will be known or apparent to those skilled in the art and are described in more detail in, for example, *Remington's Pharmaceutical Science* (17th ed., Mack Publishing Company, Easton, PA, 1985).

It will be appreciated by one of ordinary skill in the art that, in addition to the aforescribed pharmaceutical compositions, the compound of the invention can be formulated as inclusion complexes, such as cyclodextrin inclusion complexes, or liposomes. Liposomes can serve to target a compound of the invention to a particular tissue, such as lymphoid tissue or cancerous hepatic cells. Liposomes can also be used to increase the half-life of a compound of the invention. Many methods are available for

preparing liposomes, as described in, for example, Szoka et al., *Ann. Rev. Biophys. Bioeng.*, 9, 467 (1980) and U.S. Patents 4,235,871, 4,501,728, 4,837,028, and 5,019,369.

The present invention also provides a pharmaceutical composition, containing the compounds of the general formula (I) as defined above, its tautomeric forms, its stereoisomers, its analogs, its prodrugs, its isotopes, its metabolites, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates and its co-crystals in combination with the usual pharmaceutically employed carriers, diluents and the like, and for use in any of the methods described herein.

The compounds of the invention can be administered in a dose sufficient to treat the disease, condition or disorder. Such doses are known in the art (see, for example, the *Physicians' Desk Reference* (2004)). The compounds can be administered using techniques such as those described in, for example, Wasserman et al., *Cancer*, 36, pp. 1258-1268 (1975) and *Physicians' Desk Reference*, 58th ed., Thomson PDR (2004).

Suitable doses and dosage regimens can be determined by conventional range-finding techniques known to those of ordinary skill in the art. Generally, treatment is initiated with smaller dosages that are less than the optimum dose of the compound of the present invention. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. The present method can involve the administration of about 0.1 µg to about 50 mg of at least one compound of the invention per kg body weight of the individual. For a 70 kg patient, dosages of from about 10 µg to about 200 mg of the compound of the invention would be more commonly used, depending on a patient's physiological response.

By way of example and not intending to limit the invention, the dose of the pharmaceutically active agent(s) described herein for methods of treating or preventing a disease or condition as described above can be about 0.001 to about 1 mg/kg body weight of the subject per day, for example, about 0.001 mg, 0.002 mg, 0.005 mg, 0.010 mg, 0.015 mg, 0.020 mg, 0.025 mg, 0.050 mg, 0.075 mg, 0.1 mg, 0.15 mg, 0.2 mg, 0.25 mg, 0.5 mg, 0.75 mg, or 1 mg/kg body weight per day. The dose of the pharmaceutically active agent(s) described herein for the described methods can be about 1 to about 1000 mg/kg body weight of the subject being treated per day, for example, about 1 mg, 2 mg, 5 mg, 10

mg, 15 mg, 0.020 mg, 25 mg, 50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 250 mg, 500 mg, 750 mg, or 1000 mg/kg body weight per day.

In accordance with embodiments, the present invention provides methods of treating, preventing, ameliorating, and/or inhibiting a hepatitis C virus infection comprising administering a compound of formula (I) or a salt thereof.

The terms "treat," "prevent," "ameliorate," and "inhibit," as well as words stemming therefrom, as used herein, do not necessarily imply 100% or complete treatment, prevention, amelioration, or inhibition. Rather, there are varying degrees of treatment, prevention, amelioration, and inhibition of which one of ordinary skill in the art recognizes as having a potential benefit or therapeutic effect. In this respect, the inventive methods can provide any amount of any level of treatment, prevention, amelioration, or inhibition of the disorder in a mammal. For example, a disorder, including symptoms or conditions thereof, may be reduced by, for example, 100%, 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, or 10%. Furthermore, the treatment, prevention, amelioration, or inhibition provided by the inventive method can include treatment, prevention, amelioration, or inhibition of one or more conditions or symptoms of the disorder, e.g., cancer. Also, for purposes herein, "treatment," "prevention," "amelioration," or "inhibition" can encompass delaying the onset of the disorder, or a symptom or condition thereof.

In accordance with the invention, the term subject includes an "animal" which in turn includes a mammal such as, without limitation, the order Rodentia, such as mice, and the order Lagomorpha, such as rabbits. It is preferred that the mammals are from the order Carnivora, including Felines (cats) and Canines (dogs). It is more preferred that the mammals are from the order Artiodactyla, including Bovines (cows) and Swine (pigs) or of the order Perssodactyla, including Equines (horses). It is most preferred that the mammals are of the order Primates, Ceboids, or Simoids (monkeys) or of the order Anthropoids (humans and apes). An especially preferred mammal is the human.

The term "viral infection" refers to the introduction of a virus into cells or tissues, e.g., hepatitis C virus (HCV). In general, the introduction of a virus is also associated with replication. Viral infection may be determined by measuring virus antibody titer in samples of a biological fluid, such as blood, using, e.g., enzyme immunoassay. Other suitable

diagnostic methods include molecular based techniques, such as RT-PCR, direct hybrid capture assay, nucleic acid sequence based amplification, and the like. A virus may infect an organ, e.g., liver, and cause disease, e.g., hepatitis, cirrhosis, chronic liver disease and hepatocellular carcinoma.

The term "immune modulator" refers to any substance meant to alter the working of the humoral or cellular immune system of a subject. Such immune modulators include inhibitors of mast cell-mediated inflammation, interferons, interleukins, prostaglandins, steroids, cortico-steroids, colony-stimulating factors, chemotactic factors, etc.

It should also be understood that the compounds of the present invention can inhibit multiple genotypes of HCV. In one of the embodiment compound of the present invention are active against the 1a, 1b, 2a, 2b, 3a, 4a and 5a genotypes.

Following are the abbreviations used and meaning thereof in the specification:

DCM: dichloromethane. TFA: trifluoroacetic acid. Boc: *tert*-butoxycarbonyl. CDCl₃: Deuteriochloroform. DMSO-*d*₆: Dimethyl sulfoxide-*d*₆. DMF: N, N-dimethylformamide. THF: Tetrahydrofuran. DME: 1, 2-dimethoxyethane. EtOAc: Ethyl acetate. DIPEA: N, N-diisopropylethylamine. Pd(PPh₃)₄: Tetrakis(triphenylphosphine)palladium(0). Na₂SO₄: sodium sulphate. EDCI: *N*-(3-Dimethylaminopropyl)-*N*'-ethylcarbodiimide. HOBt: 1-Hydroxybenzotriazole hydrate. HATU: *O*-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate. DDQ: 2,3-Dichloro-5,6-dicyano-*p*-benzoquinone. MeOD: Methanol-*d*₄. NaHCO₃: sodium bicarbonate. HCl: hydrochloric acid. ACN: Acetonitrile. pet-ether: petroleum ether. HPLC: High Performance Liquid Chromatography.


The following examples are provided to further illustrate the present invention and therefore should not be construed in any way to limit the scope of the present invention. All ¹HNMR spectra were determined in the solvents indicated and chemical shifts are reported in δ units downfield from the internal standard tetramethylsilane (TMS) and interproton coupling constants are reported in Hertz (Hz). In case of mixture of the isomers, the peak values given are for the dominant isomer (rotamer/tautomer).

Example 1: Preparation of methyl ((S)-1-(((1R,2S)-2-(5-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-3H-benzo[2,3]oxepino[4,5-d]imidazol-8-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 1):

Step 1: Mixture of tert-butyl ((1R,2S)-2-((2-amino-4-bromophenyl)carbamoyl)cyclopentyl) carbamate AND tert-butyl ((1R,2S)-2-((2-amino-5-bromophenyl)carbamoyl)cyclopentyl) carbamate (1a):

(1S,2R)-2-((tert-butoxycarbonyl)amino)cyclopentanecarboxylic acid [synthesized using procedures described in *Tetrahedron: Asymmetry*, 2008, 19, 2796–2803] (1.52 g, 6.62 mmol) was dissolved in DMF (12 mL) and Hunig's base (1.73 mL, 9.93 mmol) was added followed by EDCI hydrochloride (1.91 g, 9.93 mmol). To this stirred mixture at room temperature was added HOBT (1.01 g, 6.62 mmol) after 10 min. followed by 4-bromobenzene-1,2-diamine (1.36 g, 7.28 mmol) after another 10 min. The brown mixture was allowed to stir overnight after which ice cold water was added and the organics were extracted with ethyl acetate (100 mL) twice. The combined ethyl acetate extracts were dried over Na₂SO₄ and concentrated under reduced pressure to obtain a brown mass which was purified by column chromatography by gradient elution with ethyl acetate-hexane (20-30%) to obtain a mixture of two regio-isomers (1.16 g, 46%). ¹H NMR (400 MHz, DMSO-D₆): δ 8.93 (br s, 2H), 7.13-6.97 (m, 2H), 6.74-6.71 (m, 2H), 6.68-6.60 (m, 4H), 5.15-5.02 (m, 2H), 4.18-4.13 (m, 2H), 3.01-2.93 (m, 2H), 1.95-1.34 (m, 12H), 1.26 (s, 18H). *m/z* 398.10 (M⁺) and 400.10 (M⁺+2).


Step 2: tert-butyl ((1R,2S)-2-(5-bromo-1H-benzo[d]imidazol-2-yl)cyclopentyl)carbamate (1b):



The mixture of two regioisomers (1.2 g, 3.26 mmol) as synthesized in Step 1 (1a) were taken in acetic acid (10 mL) and heated at 60 °C for 10 h. The reaction was cooled and most of the acetic acid was removed under reduced pressure and the contents were neutralized with saturated NaHCO₃ solution. The organics were extracted with ethyl


acetate (100 mL) twice. The combined ethyl acetate extracts were dried over Na_2SO_4 and concentrated under reduced pressure to obtain a brown mass which was purified by column chromatography by gradient elution with ethyl acetate-hexane (30-40%) to obtain the title compound, (0.93 g, 81%). ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 12.1 (brs, 1H), 7.72-7.51 (m, 1H), 7.48-7.29 (m, 1H), 7.25-7.21 (m, 1H), 6.42 (brs, 1H), 4.19-4.15 (m, 1H), 3.48-3.42 (m, 1H), 2.28-1.54 (m, 6H), 1.21 (s, 9H). m/z 380.2 (M^+) and 382.2 (M^++2).

Step 3: tert-butyl ((1R,2S)-2-(5-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)carbamate (1c):




1,4-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzene [synthesized using the procedures described in WO2003/105237A1] (1.3 g, 3.9 mmol) and tert-butyl ((1R,2S)-2-(5-bromo-1H-benzo[d]imidazol-2-yl)cyclopentyl)carbamate (1b) from the previous step (0.6 g, 1.58 mmol) was added to a degassed mixture of DME- H_2O (3:1, 4 mL) in a microwave vial followed by Na_2CO_3 (0.84 g, 7.9 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (7 mol%). The mixture was irradiated with microwaves at 120 $^\circ\text{C}$ for 40 min and the reaction mass was extracted with ethyl acetate and washed with brine. The ethyl acetate extracts were dried over Na_2SO_4 , concentrated under reduced pressure and purified by column chromatography (60% ethyl acetate-hexane) to obtain the title compound (0.25 g, 32%). ^1H NMR (400 MHz, CD_3OD): 7.82 (d, $J = 8.4$ Hz, 2H), 7.76-7.62 (m, 4H), 7.52 (s, 1H), 4.42-4.31 (m, 1H), 3.62-3.51 (m, 1H), 2.43-1.56 (m, 6H), 1.42 (s, 12H), 1.24-1.20 (m, 9H). m/z 504.2 (M^++1).

Step 4: (S)-tert-butyl 2-(8-(4-(2-((1S,2R)-2-((tert-butoxycarbonyl)amino)cyclopentyl)-1H-benzo[d]imidazol-5-yl)phenyl)-4,5-dihydro-3H-benzo[2,3]oxepino[4,5-d]imidazol-2-yl)pyrrolidine-1-carboxylate (1d):



A solution of (S)-tert-butyl 2-(8-bromo-4,5-dihydro-3H-benzo[2,3]oxepino[4,5-d]imidazol-2-yl)pyrrolidine-1-carboxylate [synthesized using the procedures described in WO2009102633] (0.10 g, 0.23 mmol) and tert-butyl ((1R,2S)-2-(5-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)carbamate (1c) (0.14 g, 0.28 mmol) from the previous step in a mixture of DME-H₂O (3:1, 4 mL) was degassed using nitrogen gas for 15 min. Sodium carbonate (0.12 g, 1.15 mmol) and Pd(PPh₃)₄ (0.03 g, 0.023 mmol) were added and irradiated using microwaves at 120 °C for 40 min. The contents were taken up in ethyl acetate and washed with water. The organic layer was dried over Na₂SO₄ and concentrated under reduced pressure to obtain the crude compound which was purified by flash chromatography (60% EtOAc/hexane) to get a yellow solid (0.1 g, 60%). ¹H-NMR (400 MHz, CDCl₃) δ 7.83-7.35 (m, 10H), 5.07-4.90 (m, 2H), 4.40-4.20 (m, 3H), 3.70-3.10 (m, 5H), 2.95-2.90 (m, 1H), 2.45-1.82 (m, 8H), 1.65-1.51 (m, 18H), *m/z* 731.40 (M⁺+1).

Step 5: methyl ((S)-1-(((1R,2S)-2-(5-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-3H-benzo[2,3]oxepino[4,5-d]imidazol-8-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 1):




The intermediate (1d) from the previous step (0.1 g, 0.14 mmol) was dissolved in DCM (5 mL) and TFA (0.2 mL) was added drop wise to the reaction mixture at 5-10 °C. After completion of addition, the mixture was stirred at room temperature for 2 h. The mixture was evaporated to dryness and triturated with diethyl ether to obtain the product as TFA salt (0.07 g) which was taken to next step without any further purification. The crude TFA

salt was dissolved in anhydrous DMF (3 mL), cooled to 0 °C and DIPEA (0.09 g, 0.66 mmol) was added to it and the contents were stirred for 5 min under nitrogen atmosphere. (S)-2-((methoxycarbonyl)amino)-3-methylbutanoic acid (0.05 g, 0.26 mmol) followed by HATU (0.1 g, 0.26 mmol) were added to it and the reaction mixture was gradually warmed to rt and stirred overnight. Crushed ice was added to the reaction mixture and the precipitate was filtered, washed with n-pentane and purified by preparative HPLC (0.022 g, 20 %). ¹H NMR (DMSO-*d*₆, 400 MHz), δ 12.20-11.80 (m, 2H), 8.30-8.10 (m, 1H), 7.85-7.30 (m, 10H), 7.11-7.00 (m, 2H), 5.10-5.02 (m, 2H), 4.40-4.25 (m, 3H), 4.10-4.05 (m, 2H), 3.88-3.82 (m, 2H), 3.60-3.48 (m, 6H), 3.10-3.07 (m, 2H), 2.30-1.60 (m, 12H), 0.97-0.68 (m, 12H), *m/z* 845.40 (M⁺+1). m.p.: 253 °C.


Example 2: Preparation of methyl ((S)-1-(((1S,2R)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 2):

Step 1: tert-butyl ((1S,2R)-2-(5-bromo-1H-benzo[d]imidazol-2-yl)cyclopentyl)carbamate (2a):




Title compound was synthesized by following the procedure as described in Steps 1 and 2 of Example 1, using (1R,2S)-2-((tert-butoxycarbonyl)amino)cyclopentanecarboxylic acid [synthesized using the procedures described in *Tetrahedron: Asymmetry*, 2008, 19, 2796–2803] and 4-bromobenzene-1,2-diamine as the starting materials. ¹H-NMR (400 MHz, CDCl₃) δ 7.80-7.70 (m, 1H), 7.50-7.45 (m, 1H), 7.35-7.30 (m, 1H), 4.97-4.90 (m, 1H), 4.80-4.70 (m, 1H), 4.50-4.40 (m, 1H), 3.60-3.50 (m, 1H), 3.40-3.30 (m, 1H), 2.40-1.60 (m, 4H), 1.30 (s, 9H); *m/z* 381.97 (M⁺+1).

Step 2: tert-butyl ((1S,2R)-2-(5-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)carbamate (2b):




Title compound was synthesized by following the procedure as described in Step 3 of Example 1, using the intermediate (2a) from previous step. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.90-7.80 (m, 2H), 7.60-7.55 (m, 4H), 7.50-7.40 (m, 2H), 4.97-4.80 (m, 1H), 4.30-4.20 (m, 1H), 3.40-3.30 (m, 1H), 2.40-1.60 (m, 6H), 1.50 (s, 12H), 1.30 (s, 9H); m/z 503.69 ($\text{M}^+ + 1$).

Step 3: (S)-tert-butyl 2-(7-(4-(2-((1R,2S)-2-((tert-butoxycarbonyl)amino)cyclopentyl)-1H-benzo[d]imidazol-6-yl)phenyl)-4,5-dihydro-1H-naphtho[1,2-d]imidazol-2-yl)pyrrolidine-1-carboxylate (2c):



Title compound was synthesized by following the procedure as described in Step 4 of Example 1, using the intermediate (2b) from the previous step and (S)-tert-butyl 2-(7-bromo-4,5-dihydro-1H-naphtho[1,2-d]imidazol-2-yl)pyrrolidine-1-carboxylate [synthesized using procedures described in WO2009102633] as the starting materials. $^1\text{H NMR}$ (DMSO-D_6 , 400 MHz) δ 12.10 (bs, 1H), 11.80 (bs, 1H), 7.83-7.50 (m, 9H), 6.85-6.80 (m, 1H), 4.82-4.73 (m, 2H), 4.40-4.20 (m, 2H), 3.55-3.50 (m, 1H), 3.10-3.01 (m, 3H), 2.90-2.70 (m, 3H), 2.40-1.70 (m, 8H), 1.60-1.51 (m, 18H). m/z 715.40 ($\text{M}^+ + 1$).


Step 4: methyl ((S)-1-(((1S,2R)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 2):



Title compound was synthesized by following the procedure as described in Step 5 of Example 1, using the intermediate (2c) from the previous step. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz), δ 12.20-11.80 (m, 2H), 7.78-7.74 (m, 4H), 7.56-7.49 (m, 6H), 7.30-7.28 (m, 2H), 5.06-5.02 (m, 1H), 4.08-4.04 (m, 2H), 3.80-3.75 (m, 4H), 3.51 (s, 6H), 3.05-3.00 (m, 4H), 2.21-1.80 (m, 12H), 0.97-0.68 (m, 12H), m/z 830.6 (M^++1). m.p.: 235-238 $^\circ\text{C}$.


Example 3: Preparation of methyl ((S)-1-(((1S,2S)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 3):

Step 1: tert-butyl ((1S,2S)-2-(5-bromo-1H-benzo[d]imidazol-2-yl)cyclopentyl)carbamate (3a):




Title compound was synthesized by following the procedure as described in Steps 1 and 2 of Example 1, using (1S,2S)-2-((tert-butoxycarbonyl)amino)cyclopentanecarboxylic acid [synthesized using procedures described in *J. Org. Chem.*, 2001, 66 (16), 5629-5632] and 4-bromobenzene-1,2-diamine as starting materials. ^1H -NMR (400 MHz, CDCl_3) δ 7.80-7.70 (m, 1H), 7.50-7.45 (m, 1H), 7.35-7.30 (m, 1H), 4.97-4.90 (m, 1H), 4.80-4.70 (m, 1H), 4.50-4.40 (m, 1H), 3.60-3.50 (m, 1H), 3.40-3.30 (m, 1H), 2.40-1.60 (m, 4H), 1.30 (s, 9H); m/z 381.97 (M^++1).

Step 2: tert-butyl ((1S,2S)-2-(5-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)carbamate (3b):




Title compound was synthesized by following the procedure as described in Step 3 of Example 1, using the intermediate (3a) from previous step as starting material. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.95-7.76 (m, 2H), 7.65-7.58 (m, 4H), 7.56-7.50 (m, 2H), 5.01-4.85 (m, 1H), 4.25-4.20 (m, 1H), 3.40-3.30 (m, 1H), 2.30-1.60 (m, 6H), 1.45 (s, 12H), 1.35 (s, 9H). m/z 503.69 (M^++1).

Step 3: (S)-tert-butyl 2-(7-(4-(2-((1S,2S)-2-((tert-butoxycarbonyl)amino)cyclopentyl)-1H-benzo[d]imidazol-6-yl)phenyl)-4,5-dihydro-1H-naphtho[1,2-d]imidazol-2-yl)pyrrolidine-1-carboxylate (3c):



Title compound was synthesized by following the procedure as described in Step 4 of Example 1, using the intermediate (3b) from the previous step and (S)-tert-butyl 2-(7-bromo-4,5-dihydro-1H-naphtho[1,2-d]imidazol-2-yl)pyrrolidine-1-carboxylate [synthesized using the procedures described in WO2009102633] as the starting materials. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.80-7.40 (m, 9H), 6.95-6.90 (m, 1H), 4.95-4.83 (m, 2H), 4.30-4.15 (m, 2H), 3.55-3.40 (m, 3H), 3.10-3.05 (m, 2H), 2.90-2.80 (m, 2H), 2.40-1.70 (m, 8H), 1.60-1.51 (m, 18H), m/z 715.40 (M^++1).


Step 4: methyl ((S)-1-(((1S,2S)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 3):



Title compound was synthesized by following the procedure as described in Step 5 of Example 1, using the intermediate (3c) from the previous step. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz), δ 12.20-11.80 (m, 2H), 7.78-7.74 (m, 4H), 7.56-7.49 (m, 6H), 7.30-7.28 (m, 2H), 5.06-5.02 (m, 1H), 4.08-4.04 (m, 2H), 3.80-3.75 (m, 4H), 3.51 (s, 6H), 3.05-3.00 (m, 4H), 2.21-1.80 (m, 12H), 0.97-0.68 (m, 12H), m/z 830.6 ($\text{M}^+ + 1$). m.p.: 218-220 $^\circ\text{C}$.

Example 4: Preparation of methyl ((S)-1-(((1S,2S)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-1H-benzo[2,3]oxepino[4,5-d]imidazol-8-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 4):


Step 1: (S)-tert-butyl 2-(8-(4-(2-((1S,2S)-2-((tert-butoxycarbonyl)amino)cyclopentyl)-1H-benzo[d]imidazol-6-yl)phenyl)-4,5-dihydro-1H-benzo[2,3]oxepino[4,5-d]imidazol-2-yl)pyrrolidine-1-carboxylate (4a):



Title compound was synthesized by following the procedure as described in Step 4 of Example 1, using intermediate from Step 2 of Example 3 (3b) and (S)-tert-butyl 2-(8-bromo-4,5-dihydro-3H-benzo[2,3]oxepino[4,5-d]imidazol-2-yl)pyrrolidine-1-carboxylate [synthesized using the procedures described in WO2009102633] as the starting materials. ^1H -NMR (400 MHz, CDCl_3) δ 7.83-7.35 (m, 10H), 5.07-4.90 (m, 2H), 4.40-4.20 (m, 3H), 3.70-3.10 (m, 5H), 2.95-2.90 (m, 1H), 2.45-1.82 (m, 8H), 1.65-1.51 (m, 18H), m/z 731.50 ($\text{M}^+ + 1$).

Step 2: methyl ((S)-1-(((1S,2S)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-4,5-dihydro-1H-benzo[2,3]oxepino[4,5-d]imidazol-8-


yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 4):



Title compound was synthesized by following the procedure as described in Step 5 of Example 1, using the intermediate (4a) from the previous step. ^1H NMR (DMSO- d_6 , 400 MHz), δ 12.20-11.80 (m, 2H), 8.30-8.10 (m, 1H), 7.85-7.30 (m, 10H), 7.11-7.00 (m, 2H), 5.12-4.90 (m, 2H), 4.40-4.15 (m, 3H), 4.10-4.05 (m, 2H), 3.88-3.82 (m, 2H), 3.60-3.48 (m, 6H), 3.10-3.07 (m, 2H), 2.30-1.60 (m, 12H), 0.97-0.68 (m, 12H), m/z 845.50 ($M^+ + 1$). m.p.: 240-242 $^\circ\text{C}$.


Example 5: Preparation of methyl ((S)-1-(((1S,2S)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 5):

Step 1: (S)-tert-butyl 2-(7-bromo-3H-naphtho[1,2-d]imidazol-2-yl)pyrrolidine-1-carboxylate (5a):




DDQ (0.41 g, 1.79 mmol) was added to the solution of (S)-tert-butyl 2-(7-bromo-4,5-dihydro-3H-naphtho[1,2-d]imidazol-2-yl)pyrrolidine-1-carboxylate [synthesized using the procedures described in WO2009102633] (0.5 g, 1.19 mmol) in benzene (10 mL) and stirred at 80 $^\circ\text{C}$ for 3 h. Benzene was evaporated under reduced pressure and the residue was purified by column chromatography (60% EtOAc/hexane) to afford the title compound (0.3 g, 60%). ^1H -NMR (400 MHz, CDCl_3): δ 8.05-8.03 (m, 1H), 7.62-7.50 (m, 4H), 5.24-5.23 (m, 1H), 3.61-3.50 (m, 1H), 3.15-3.10 (m, 1H), 2.29-2.19 (m, 2H), 2.09-2.01 (m, 2H), 1.54-1.34 (m, 9H), m/z 416.09 ($M^+ + 1$).

Step 2: (S)-tert-butyl 2-(7-(4-(2-((1S,2S)-2-((tert-butoxycarbonyl)amino)cyclopentyl)-1H-benzo[d]imidazol-6-yl)phenyl)-1H-naphtho[1,2-d]imidazol-2-yl)pyrrolidine-1-carboxylate (5b):



Title compound was synthesized by following the procedure as described in Step 4 of Example 1 using the intermediates from the previous step (5a) and Step 2 of Example 3 (3b). $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.10-7.45 (m, 12H), 5.40-5.30 (m, 1H), 5.02-4.80 (m, 1H), 4.20-4.10 (m, 1H), 3.75-3.35 (m, 2H), 3.25-2.95 (m, 2H), 2.55-1.90 (m, 8H), 1.60-1.50 (m, 18H), m/z 713.40 ($\text{M}^+ + 1$).


Step 3: methyl ((S)-1-(((1S,2S)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 5):



Title compound was synthesized by following the procedure as described in Step 5 of Example 1, using the intermediate (5b) from the previous step. $^1\text{H NMR}$ ($\text{DMSO}-d_6$, 400 MHz), δ 13.10 (bs, 1H), 12.25 (bs, 1H), 8.45-8.20 (m, 3H), 8.10-7.43 (m, 9H), 7.35-7.33 (m, 1H), 7.04-7.02 (m, 1H), 5.34-5.29 (m, 1H), 4.45-4.40 (m, 2H), 4.25-4.20 (m, 2H), 3.95-3.75 (m, 2H), 3.55-3.51 (m, 6H), 2.45-1.55 (m, 12H), 0.97-0.68 (m, 12H), m/z 827.40 ($\text{M}^+ + 1$). m.p.: 200-202 $^\circ\text{C}$.


Example 6: Preparation of methyl ((S)-1-(((1S,2R)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 6):

Step 1: (S)-tert-butyl 2-(7-(4-(2-((1R,2S)-2-((tert-butoxycarbonyl)amino)cyclopentyl)-1H-benzo[d]imidazol-6-yl)phenyl)-1H-naphtho[1,2-d]imidazol-2-yl)pyrrolidine-1-carboxylate (6a):



Title compound was synthesized by following the procedure as described in Step 4 of Example 1 using the intermediates from Step 1 of Example 5, (5a) and Step 2 of Example 2 (2b) as starting materials. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 11.70 (bs, 1H), 11.05 (bs, 1H), 8.30-7.45 (m, 12H), 5.30-5.20 (m, 1H), 4.95-4.80 (m, 1H), 4.55-4.40 (m, 1H), 3.60-3.05 (m, 4H), 2.45-1.90 (m, 8H), 1.60-1.50 (m, 18H), m/z 713.50 ($\text{M}^+ + 1$).

Step 2: methyl ((S)-1-(((1S,2R)-2-(6-(4-(2-((S)-1-((S)-2-(methoxycarbonyl)amino-3-methylbutanoyl)pyrrolidin-2-yl)-1H-naphtho[1,2-d]imidazol-7-yl)phenyl)-1H-benzo[d]imidazol-2-yl)cyclopentyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (Compound 6):



Title compound was synthesized by following the procedure as described in Step 5 of Example 1, using the intermediate (6a) from the previous step. $^1\text{H NMR}$ ($\text{DMSO}-d_6$, 400 MHz), δ 13.10 (bs, 1H), 12.25 (bs, 1H), 8.45-8.20 (m, 3H), 8.10-7.43 (m, 9 H), 7.35-7.33 (m, 1H), 7.04-7.02 (m, 1H), 5.34-5.29 (m, 1H), 4.45-4.40 (m, 1H), 4.25-4.20 (m, 1H), 3.95-3.75 (m, 4H), 3.55-3.51 (m, 6H), 2.45-1.55 (m, 12H), 0.97-0.39 (m, 12H), m/z 827.50 ($\text{M}^+ + 1$). m.p.: 257-259 $^\circ\text{C}$.

Example 7: Biological Activity

Anti-viral activity of the compounds of the invention was monitored using an HCV replicon assay. The Huh7.5/ Con1/SG-Neo(I)hRluc2aUb cell line persistently expressing a bicistronic genotype 1b replicon in Huh 7.5 cells was obtained from Apath LLC. This cell line was used to test inhibition of replicon levels by test compound using *Renilla* luciferase enzyme activity readout as a measure of viral replication efficiency.

Briefly, 7000 cells were seeded in 96 well black clear bottom plates and allowed to adhere overnight. The next day each compound was added in triplicate to the cells at the desired concentration with a final DMSO concentration of 0.5%. Cells in media alone and cells incubated without drug with 0.5% DMSO served as controls. The plates were incubated for 72h at 37°C prior to running the luciferase assay. Enzyme activity was measured using *Renilla-Glo* Luciferase Assay kit from Promega as per the manufacturer's instructions. The following equation was used to generate the percent inhibition value for each test concentration.

$$\% \text{ Inhibition} = \frac{\text{Average Control (cells alone +0.5\% DMSO)} - \text{Average compound value(cells + drug)}}{\text{Average Control (cells alone+0.5\% DMSO)}} \times 100$$

The EC₅₀ value was determined using GraphPad Prism and the following equation:


$$Y = \text{Bottom} + (\text{Top} - \text{Bottom}) / (1 + 10^{((\text{LogIC}_{50} - X) * \text{HillSlope}))}$$

EC₅₀ values/% inhibitions of compounds were determined 2-3 times in the replicon assay.

It is found that the compounds of the present invention exhibit EC₅₀ value between 20 pM to 2 nM for genotype 1b replicon.

CLAIMS

1. A compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals,



wherein, in the compound of formula I,

Ring 'A' is a 5 to 10 membered saturated carbocycle, the said carbocycle may be a monocyclic system, or a fused- or bridged- or a spiro- bicyclic system;

ring D is selected from substituted- or unsubstituted- 5 to 10 membered carbocycles, substituted- or unsubstituted- 5 to 10 membered heterocycles containing 1 to 3 heteroatoms/groups selected from N(R⁸), S(O)_p, O or C(=O), substituted- or unsubstituted- aromatic carbocycle, and 5 to 6 membered substituted- or unsubstituted- aromatic heterocycle containing heteroatoms selected from N, S or O;

Ring 'E' is selected from 5 to 10 membered carbocycle and 5 to 10 membered heterocycle, the ring 'E' may be monocyclic, fused bicyclic, bridged bicyclic or spiro bicyclic;;

Y is selected from CH(R^{2a}) and N(R²)

R¹ is selected from the group consisting of R^{8a}C(=O)N(R⁹)-, R⁸OC(=O)N(R⁹)-, R⁸(R⁹)N-, R⁸(R⁹)NC(=O)N(R¹⁰)-, R⁸SO₂N(R⁹)-, R^{8a}OC(=O)N(R⁹)CR^a(R^b)C(=O)N(R¹⁰)-, R^{8a}OC(=O)N(R⁹)CR^a(R^b)C(R^c)(R^d)C(=O)N(R¹⁰)-, R⁸(R^{9a})NC(=O)N(R⁹)CR^a(R^b)C(=O)N(R¹⁰)-, and R⁸(R^{9a})NC(=O)N(R⁹)CR^a(R^b)C(R^c)(R^d)C(=O)N(R¹⁰)-;

R^{1a} is selected independently at each occurrence from the group consisting of halogen, substituted- or unsubstituted- C_{1-6} alkyl, $R^{8a}C(=O)-$, $R^{8b}O-$, $R^{8a}OC(=O)-$, $R^{8a}C(=O)O-$, and $R^8(R^9)NC(=O)-$;

R^2 is selected independently at each occurrence from the group consisting of substituted- or unsubstituted- C_{1-6} alkyl, $R^{8a}C(=O)-$, $R^{8a}SO_2-$, $R^{8a}OC(=O)-$, $R^8(R^9)NC(=O)-$, $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)C(=O)-$, $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)C(R^c)(R^d)C(=O)-$, $R^8(R^9)NC(=O)N(R^{10})C(R^a)(R^b)C(=O)-$, $R^8(R^9)NC(=O)N(R^{10})C(R^a)(R^b)C(R^c)(R^d)C(=O)-$, $R^8SO_2N(R^9)C(R^a)(R^b)C(=O)-$, $R^8SO_2N(R^9)C(R^a)(R^b)C(R^c)(R^d)C(=O)-$, $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)SO_2-$, and $R^8SO_2N(R^9)C(=O)C(R^a)(R^b)N(R^{10})C(=O)-$;

R^{2a} is selected independently at each occurrence from the group consisting of $R^{8a}C(=O)N(R^9)-$, $R^8OC(=O)N(R^9)-$, $R^8(R^9)N-$, $R^8(R^9)NC(=O)N(R^{10})-$, $R^{8a}SO_2N(R^9)-$, $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)C(=O)N(R^{10})-$, $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)C(R^c)(R^d)C(=O)N(R^{10})-$, $R^8(R^{9a})NC(=O)N(R^9)C(R^a)(R^b)C(=O)N(R^{10})-$, and $R^8(R^{9a})NC(=O)N(R^9)C(R^a)(R^b)C(R^c)(R^d)C(=O)N(R^{10})-$;

R^3 is selected from O and $N(R^{11})$;

R^4 is selected independently at each occurrence from $CR^a(R^b)$, O, $N(R^{11})$, substituted- or unsubstituted carbocycle, substituted- or unsubstituted- heterocycle, substituted- or unsubstituted- arylene and substituted- or unsubstituted- heteroarylene, wherein, substitutions on carbocycle, heterocycle, arylene, and heteroarylene are selected from the group consisting of halogen, substituted- or unsubstituted- C_{1-6} alkyl, and alkyl-O-;

R^5 is selected from hydrogen and substituted- or unsubstituted- alkyl;

R^6 and R^7 are independently selected from the group consisting of hydrogen, halogen, substituted- or unsubstituted- C_{1-6} alkyl, $R^{8b}O-$, and $R^8(R^9)N-$;

wherein, R^8 , R^9 , R^{9a} and R^{10} are independently selected from the group consisting of hydrogen, substituted- or unsubstituted- C_{1-6} alkyl, substituted- or unsubstituted- aryl, substituted- or unsubstituted- heteroaryl, substituted- or unsubstituted- cycloalkyl, and substituted- or unsubstituted- heterocyclyl;

R^{8a} is selected from the group consisting of substituted- or unsubstituted- C_{1-6} alkyl, substituted- or unsubstituted- aryl, substituted- or unsubstituted- heteroaryl, substituted- or unsubstituted- cycloalkyl, and substituted- or unsubstituted- heterocyclyl;

R^{8b} is selected from the group consisting of hydrogen, substituted- or unsubstituted- C_{1-6} alkyl, perhaloalkyl, substituted- or unsubstituted- aryl, substituted- or unsubstituted- heteroaryl, substituted- or unsubstituted- cycloalkyl, and substituted- or unsubstituted- heterocyclyl;

R^{11} is selected from hydrogen and substituted- or unsubstituted- alkyl group;

R^a , R^b , R^c and R^d , are independently selected from the group consisting of hydrogen, halogen, substituted- or unsubstituted- C_{1-6} alkyl, substituted- or unsubstituted- aryl, substituted- or unsubstituted- heteroaryl, substituted- or unsubstituted- cycloalkyl, and substituted- or unsubstituted- heterocyclyl; R^a , R^b , R^c and R^d together with the carbon atom to which they are attached may take part in forming substituted- or unsubstituted- carbocycle, or substituted- or unsubstituted- heterocycle containing 3 to 10 atoms;

m is an integer ranging between 0 to 2, selected independently at each occurrences;

n is an integer ranging between 0 and 2;

p is an integer ranging between 0 and 2;

when $n = 2$ and R^4 is selected as $CR^a(R^b)$ for both the occurrences, two R^a s together can form a bond to form a alkenylene linkage or two R^a s and two R^b s together can form bonds to form alkynylene linkage;

'alkyl' may be substituted with 1 to 4 substituents selected from the group consisting of oxo, halogen, cyano, aryl, hereroaryl, cycloalkyl, $R^{12a}SO_2-$, $R^{12b}O-$, $R^{12a}OC(=O)-$, $R^{12a}C(=O)O-$, $(R^{12})(H)NC(=O)-$, $(R^{12})(alkyl)NC(=O)-$, $R^{12a}C(=O)N(H)-$, $R^{12}(H)N-$, $R^{12}(alkyl)N-$, $R^{12}(H)NC(=O)N(H)-$, and $R^{12}(alkyl)NC(=O)N(H)-$;

'cycloalkyl' and 'carbocycle' may be substituted with 1 to 2 substituents selected from the group consisting of oxo, halogen, cyano, aryl, hereroaryl, alkyl, haloalkyl, $R^{12a}C(=O)-$, $R^{12a}SO_2-$, $R^{12b}O-$, $R^{12a}OC(=O)-$, $R^{12a}C(=O)O-$, $R^{12}(H)NC(=O)-$,

$R^{12}(\text{alkyl})\text{NC}(=\text{O})-$, $R^{12a}\text{C}(=\text{O})\text{N}(\text{H})-$, $R^{12}(\text{H})\text{N}-$, $R^{12}(\text{alkyl})\text{N}-$, $R^{12}(\text{H})\text{NC}(=\text{O})\text{N}(\text{H})-$, and $R^{12}(\text{alkyl})\text{NC}(=\text{O})\text{N}(\text{H})-$;

'aryl' or 'aromatic carbocycle', may be substituted with 1 to 2 substituents selected from the group consisting of halogen, nitro, cyano, hydroxy, C_1 to C_6 alkyl, C_3 to C_6 cycloalkyl, 3- to 6- membered heterocycle, C_1 to C_6 perhaloalkyl, alkyl-O-, perhaloalkyl-O-, alkyl-N(alkyl)-, alkyl-N(H)-, $\text{H}_2\text{N}-$, alkyl-SO₂-, perhaloalkyl-SO₂-, alkyl-C(=O)N(alkyl)-, alkyl-C(=O)N(H)-, alkyl-N(alkyl)C(=O)-, alkyl-N(H)C(=O)-, $\text{H}_2\text{NC}(=\text{O})-$, alkyl-N(alkyl)SO₂-, alkyl-N(H)SO₂-, H_2NSO_2- ;

'heteroaryl' or 'aromatic heterocycle' may be substituted with 1 to 2 substituents selected from the group consisting of halogen, nitro, cyano, hydroxy, C_1 to C_6 alkyl, C_3 to C_6 cycloalkyl, 3- to 6- membered heterocycle, C_1 to C_6 perhaloalkyl, alkyl-O-, perhaloalkyl-O-, alkyl-N(alkyl)-, alkyl-N(H)-, $\text{H}_2\text{N}-$, alkyl-SO₂-, perhaloalkyl-SO₂-, alkyl-C(=O)N(alkyl)-, alkyl-C(=O)N(H)-, alkyl-N(alkyl)C(=O)-, alkyl-N(H)C(=O)-, $\text{H}_2\text{NC}(=\text{O})-$, alkyl-N(alkyl)SO₂-, and alkyl-N(H)SO₂-, H_2NSO_2- ;

ring carbons of 'heterocyclyl' and 'heterocycle' may be substituted with 1 to 2 substituents selected from the group consisting of oxo, halogen, nitro, cyano, aryl, heteroaryl, alkyl, $R^{12b}\text{O}-$, $R^{12a}\text{OC}(=\text{O})-$, $R^{12a}\text{C}(=\text{O})\text{O}-$, $R^{12}(\text{H})\text{NC}(=\text{O})-$, $R^{12}(\text{alkyl})\text{NC}(\text{O})-$, $R^{12a}\text{C}(=\text{O})\text{N}(\text{H})-$, $R^{12}(\text{H})\text{N}-$, $R^{12}(\text{alkyl})\text{N}-$, $R^{12}(\text{H})\text{NC}(=\text{O})\text{N}(\text{H})-$, and $R^{12}(\text{alkyl})\text{NC}(=\text{O})\text{N}(\text{H})-$; the substituents on ring nitrogen(s) of 'heterocyclyl' and 'heterocycle' are selected from the group consisting of aryl, heteroaryl, alkyl, $R^{12a}\text{C}(=\text{O})-$, $R^{12a}\text{SO}_2-$, $R^{12a}\text{OC}(=\text{O})-$, $R^{12}(\text{H})\text{NC}(=\text{O})-$, $R^{12}(\text{alkyl})\text{NC}(=\text{O})-$;


R^{12} is selected from hydrogen and alkyl;

R^{12a} is selected as alkyl, perhaloalkyl;


R^{12b} is selected from hydrogen, alkyl, perhaloalkyl.

2. The compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals, as claimed in claim 1, wherein ring A is selected as cyclopentane.

3. The compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals, as claimed in claim 1 or 2, wherein R^1 is selected as $R^{8a}OC(=O)N(R^9)CR^a(R^b)C(=O)N(R^{10})-$.
4. The compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals, as claimed in any one of claims 1-3, wherein m is selected as 0 at all the occurrences.
5. The compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals, as claimed in any one of claims 1-4, wherein ring Y is selected as $N(R^2)$.
6. The compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals, as claimed in any one of claims 1-5, wherein R^2 is selected as $R^{8a}OC(=O)N(R^9)C(R^a)(R^b)C(=O)-$, and q is selected as 1.
7. The compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals, as claimed in any one of claims 1-6, wherein ring E is selected as-



8. The compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals, as claimed in any one of claims 1-7, wherein R^3 is selected as NH.
9. The compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals, as claimed in any one of claims 1-8, wherein R^4 is selected as phenylene, and n is selected as 1.
10. The compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals, as claimed in any one of claims 1-9, wherein R^5 , R^6 and R^7 are selected as hydrogen.
11. The compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals, as claimed in any one of claims 1-10, wherein ring D is selected from a six membered carbocycle and seven membered heterocycle containing one heteroatom.
12. The compound of formula I, its tautomeric forms, its stereoisomers, its analogues, its prodrugs, its isotopically substituted analogues, its metabolites, its sulfoxides, its N-oxides, its pharmaceutically acceptable salts, its polymorphs, its solvates, its optical isomers, its clathrates or its co-crystals, as claimed in any one of claims 1-11, wherein ring A is selected as cyclopentane; R^1 is selected as $R^{8a}OC(=O)N(R^9)CR^a(R^b)C(=O)N(R^{10})-$; m is selected as 0 at all the occurrences;



ring E is selected as- ; R^3 is selected as NH; R^4 is selected as phenylene, n is selected as 1; R^5 , R^6 and R^7 are selected as hydrogen; Ring D is selected from a six membered carbocycle and seven membered heterocycle containing one heteroatom.

13. A pharmaceutical composition comprising a compound or a combination of compounds according to any one of claims 1-12 or a pharmaceutically acceptable salt thereof, in combination with a pharmaceutically acceptable carrier or excipient.
14. A method of inhibiting the replication of an RNA-containing virus comprising contacting said virus with a therapeutically effective amount of a compound or combination of compounds of any one of claims 1-12, or a pharmaceutically acceptable salt thereof.
15. A method of treating or preventing infection caused by an RNA-containing virus comprising administering to a patient in need of such treatment a therapeutically effective amount of a compound or combination of compounds of any one of claims 1-12, or a pharmaceutically acceptable salt thereof.
16. The method of claim 15, wherein the RNA-containing virus is hepatitis C virus.
17. The method of claim 15, further comprising the step of co-administering one or more agents selected from the group consisting of a host immune modulator and an antiviral agent, or a combination thereof.

18. The method of claim 17, wherein the host immune modulator is selected from the group consisting of interferon-alpha, pegylated-interferon-alpha, interferon-beta, interferon-gamma, consensus interferon, a cytokine, and a vaccine.
19. The method of claim 17, wherein the antiviral agent inhibits replication of HCV by inhibiting host cellular functions associated with viral replication.
20. The method of claim 17, wherein the antiviral agent inhibits the replication of HCV by targeting proteins of the viral genome.
21. The method of claim 17, wherein said antiviral agent is an inhibitor of a HCV viral protein, a replication process or a combination thereof, wherein said targeting protein or replication process is selected from the group consisting of helicase, protease, polymerase, metalloprotease, NS4A, NS4B, NS5A, assembly, entry, and IRES.
22. The method of claim 15, further comprising the step of co-administering an agent or combination of agents that treat or alleviate symptoms of HCV infection selected from cirrhosis and inflammation of the liver.
23. The method of claim 15, further comprising the step of co-administering one or more agents that treat patients for disease caused by hepatitis B (HBV) infection.
24. The method of claim 15, further comprising the step of co-administering one or more agents that treat patients for disease caused by human immunodeficiency virus (HIV) infection.

25. The pharmaceutical composition of claim 13, further comprising an agent selected from interferon, pegylated interferon, ribavirin, amantadine, an HCV protease inhibitor, an HCV polymerase inhibitor, an HCV helicase inhibitor, or an internal ribosome entry site inhibitor.
26. The composition of claim 13, further comprising a cytochrome P450 monooxygenase inhibitor or a pharmaceutically acceptable salt thereof.
27. A method of treating hepatitis C infection in a subject in need thereof comprising co-administering to said subject a cytochrome P450 monooxygenase inhibitor or a pharmaceutically acceptable salt thereof, and a compound of any one of claims 1-15 or a pharmaceutically acceptable salt thereof.