
CLAIMS

1. A combination product comprising:

(i) CVC or an analogue, pharmaceutically acceptable salt or solvate thereof, and

(ii) at least one PPAR agonist of formula (I),



in which:

Y1 represents a halogen, a Ra, or Ga-Ra group;

A represents a CH=CH or a CH2-CH2 group;

Y2 represents a Gb-Rb group;

Ga and Gb, identical or different, represent an atom of oxygen or sulfur;

Ra represents a hydrogen atom, an unsubstituted (C1-C6)alkyl group, a (C6-C14)aryl group or a (C1-C6)alkyl group that is substituted by one or more halogen atoms, a (C1-C6)alkoxy or a (C1-C6)alkylthio group, (C3-C14)cycloalkyl groups, (C3-C14)cycloalkylthio groups or heterocyclic groups;

Rb represents a (C1-C6)alkyl group substituted by at least a –COORc group, wherein Rc represents a hydrogen atom, or a (C1-C6)alkyl group that is substituted or not by one or more halogen atoms, (C3-C14)cycloalkyl groups, or heterocyclic groups; and

Y4 and Y5, identical or different, representing a (C1-C6)alkyl group that is substituted or not by one or more halogen atoms, (C3-C14)cycloalkyl groups or heterocyclic groups.

2. The combination product according to claim 1, comprising:

(i) CVC or a pharmaceutically acceptable salt thereof; and

(ii) Elafibranor or a pharmaceutically acceptable salt thereof.

3. The combination product according to claim 1 or 2, wherein the combination product is a composition comprising components (i) and (ii) and a pharmaceutically acceptable carrier.

4. The combination product according to claim 1 or 2, wherein the combination product is a kit of parts comprising components (i) and (ii), for sequential, separate or simultaneous use.

5. The combination product according to any one of claims 1 to 4, wherein components (i) and (ii) are formulated in an injectable suspension, a gel, an oil, a pill, a tablet, a suppository, a powder, a capsule, an aerosol, an oinment, a cream, a patch, or means of galenic forms for a prolonged and/or slow release.

6. The combination product according to any one of claims 1 to 5, further comprising at least one therapeutically active agent with known antifibrotic activity selected from pirfenidone or receptor tyrosine kinase inhibitors (RTKIs) such as Nintedanib, Sorafenib and other RTKIs, or angiotensin II (AT1) receptor blockers, or CTGF inhibitor, or any antifibrotic compound susceptible to interfere with the TGFβ and BMP-activated pathways including activators of the latent TGFβ complex such as MMP2, MMP9, THBS1 or cell-surface integrins, TGFβ receptors type I (TGFBRI) or type II (TGFBRII) and their ligands such as TGFβ, Activin, inhibin, Nodal, anti-Müllerian hormone, GDFs or BMPs, auxiliary co-receptors (also known as type III receptors), or components of the SMAD-dependent canonical pathway including regulatory or inhibitory SMAD proteins, or members of the SMAD-independent or non-canonical pathways including various branches of MAPK signaling, TAK1, Rho-like GTPase signaling pathways, phosphatidylinositol-3 kinase/AKT pathways, TGFβ-induced EMT process, or canonical and non-canonical Hedgehog signaling pathways including Hh ligands or target genes, or any members of the WNT, or Notch pathways which are susceptible to influence TGFβ.

7. The combination product according to any one of claims 1 to 6, further comprising at least one therapeutically active agent selected from JAK/STAT inhibitors and other anti-inflammatory and/or immunosuppressant agents.

8. The combination product according to claim 7, wherein the therapeutically active agent is selected from glucocorticoids, NSAIDS, cyclophosphamide, nitrosoureas, folic acid analogs, purine analogs, pyrimidine analogs, methotrexate, azathioprine, mercaptopurine, ciclosporin, myriocin, tacrolimus, sirolimus, mycophenolic acid derivatives, fingolimod and other sphingosine-1-phosphate receptor modulators, monoclonal and/or polyclonal antibodies against such targets as proinflammatory cytokines and proinflammatory cytokine receptors, T-cell receptor and integrins.

9. The combination product according to any one of claims 1 to 8, for use as a medicament.

10. The combination product according to any one of claims 1 to 9, for use in a method for treating an inflammatory, metabolic, fibrotic and cholestatic disease.

11. The combination product for use according to claim 10, wherein the fibrotic disorder is selected in the group consisting of liver, kidney, skin, epidermis, endodermis, muscle, tendon, cartilage, heart, pancreas, lung, uterus, nervous system, testis, ovary, adrenal gland, artery, vein, colon, intestine (e.g. small intestine), biliary tract, soft tissue (e.g. mediastinum or retroperitoneum), bone marrow, joint and stomach fibrosis.

12. The combination product for use according to claim 10 or 11, wherein the fibrotic disorder is selected in the group consisting of liver, gut, lung, heart, kidney, muscle, skin, soft tissue, bone marrow, intestinal, and joint fibrosis.

13. The combination product for use according to any one of claims 10 to 12, wherein the disease is selected in the group consisting of metabolic liver diseases, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), drug-induced liver diseases, alcohol-induced liver diseases, infectious agent induced liver diseases, inflammatory liver diseases, immune system dysfunction-mediated liver diseases, dyslipidemia, cardiovascular diseases, restenosis, syndrome X, metabolic syndrome, diabetes, obesity, hypertension, chronic cholangiopathies such as Primary Sclerosing Cholangitis (PSC), Primary Biliary Cholangitis (PBC), biliary atresia, familial intrahepatic cholestasis type 3 (PFIC3), inflammatory bowel diseases, Crohn's disease, ulcerative colitis, keloid, old myocardial infarction, scleroderma/systemic sclerosis, inflammatory diseases, neurodegenerative diseases, cancers, liver cancer, hepatocallular carcinoma, gastrointestinal cancer, gastric cancer, meningioma associated with neurofibromatosis, pancreatic neuroendocrine tumors, pancreatic exocrine tumors, leukemia, myeloproliferative/myelodisplastic diseases, mastocytosis, dermatofibrosarcoma, solid tumors including breast, lung, thyroid or colorectal cancer, a prostate cancer, liver fibrosis or cirrhosis of any origin, metabolic disease-induced liver fibrosis or cirrhosis, NAFLD-induced fibrosis or cirrhosis, NASH-induced fibrosis or cirrhosis, alcohol-induced liver fibrosis or cirrhosis, drug-induced liver fibrosis or cirrhosis, infectious agent-induced liver fibrosis or cirrhosis, parasite infection-induced liver fibrosis or cirrhosis, bacterial infection-induced liver fibrosis or cirrhosis, viral infection-induced fibrosis

or cirrhosis, HBV-infection induced liver fibrosis or cirrhosis, HCV-infection induced liver fibrosis or cirrhosis, HIV-infection induced liver fibrosis or cirrhosis, dual HCV and HIV-infection induced liver fibrosis or cirrhosis, radiation- or chemotherapy-induced fibrosis or cirrhosis, biliary tract fibrosis, liver fibrosis or cirrhosis due to any chronic cholestatic disease, gut fibrosis of any etiology, Crohn's disease-induced fibrosis, ulcerative colitis-induced fibrosis, intestine (e.g. small intestine) fibrosis, colon fibrosis, stomach fibrosis, skin fibrosis, epidermis fibrosis, endodermis fibrosis, skin fibrosis due to scleroderma/systemic sclerosis, lung fibrosis, lung fibrosis consecutive to chronic inflammatory airway diseases, such as COPD, asthma, emphysema, smoker's lung,tuberculosis, pulmonary fibrosis, idiopathic pulmonary fibrosis (IPF), heart fibrosis, kidney fibrosis, nephrogenic systemic fibrosis, muscle fibrosis, soft tissue (e.g. mediastinum or retroperitoneum) fibrosis, bone marrow fibrosis, joint fibrosis, tendon fibrosis, cartilage fibrosis, pancreas fibrosis, uterus fibrosis, nervous system fibrosis, testis fibrosis, ovary fibrosis, adrenal gland fibrosis, artery fibrosis, vein fibrosis, eye fibrosis, endomyocardial fibrosis, mediastinal fibrosis, myelofibrosis, retroperitoneal fibrosis, progressive massive fibrosis (a complication of coal workers' pneumoconiosis), proliferative fibrosis, neoplastic fibrosis, peri-implantational fibrosis and asbestosis, arthrofibrosis, adhesive capsulitis.

14. The combination product for use according to any one of claims 10 to 12, wherein the disease is selected in the group consisting of, the disease is selected in the group consisting of metabolic liver diseases, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), drug-induced liver diseases, alcohol-induced liver diseases, infectious agent induced liver diseases, inflammatory liver diseases, immune system dysfunction-mediated liver diseases, dyslipidemia, cardiovascular diseases, restenosis, syndrome X, metabolic syndrome, diabetes, obesity, hypertension, chronic cholangiopathies such as Primary Sclerosing Cholangitis (PSC), Primary Biliary Cholangitis (PBC), biliary atresia, familial intrahepatic cholestasis type 3 (PFIC3), inflammatory bowel diseases, Crohn's disease, ulcerative colitis, liver cancer, hepatocallular carcinoma, gastrointestinal cancer, gastric cancer, colorectal cancer, metabolic disease-induced liver fibrosis or cirrhosis, NAFLD-induced fibrosis or cirrhosis, NASH-induced fibrosis or cirrhosis, alcohol-induced liver fibrosis or cirrhosis, drug-induced liver fibrosis or cirrhosis, infectious agent-induced liver fibrosis or cirrhosis, parasite infection-induced liver fibrosis or cirrhosis, bacterial infection-induced liver fibrosis or cirrhosis, viral infection-induced fibrosis or cirrhosis, HBV-infection induced liver fibrosis or cirrhosis, HCV-infection induced liver fibrosis

or cirrhosis, HIV-infection induced liver fibrosis or cirrhosis, dual HCV and HIV-infection induced liver fibrosis or cirrhosis, radiation- or chemotherapy-induced fibrosis or cirrhosis, biliary tract fibrosis, liver fibrosis or cirrhosis due to any chronic cholestatic disease, gut fibrosis of any etiology, Crohn's disease-induced fibrosis, ulcerative colitis-induced fibrosis, intestine (e.g. small intestine) fibrosis, colon fibrosis, stomach fibrosis, lung fibrosis, lung fibrosis consecutive to chronic inflammatory airway diseases, such as COPD, asthma, emphysema, smoker's lung, tuberculosis, pulmonary fibrosis, idiopathic pulmonary fibrosis (IPF).

15. Use of:

(i) CVC or an analogue, pharmaceutically acceptable salt or solvate thereof, and

(ii) at least one PPAR agonist of formula (I),

in which:

Y1 represents a halogen, a Ra, or Ga-Ra group;

A represents a CH=CH or a CH2-CH2 group;

Y2 represents a Gb-Rb group;

Ga and Gb, identical or different, represent an atom of oxygen or sulfur;

Ra represents a hydrogen atom, an unsubstituted (C1-C6)alkyl group, a (C6-C14)aryl group or a (C1-C6)alkyl group that is substituted by one or more halogen atoms, a (C1-C6)alkoxy or a (C1-C6)alkylthio group, (C3-C14)cycloalkyl groups, (C3-C14)cycloalkylthio groups or heterocyclic groups;

Rb represents a (C1-C6)alkyl group substituted by at least a –COORc group, wherein Rc represents a hydrogen atom, or a (C1-C6)alkyl group that is substituted or not by one or more halogen atoms, (C3-C14)cycloalkyl groups, or heterocyclic groups; and

Y4 and Y5, identical or different, representing a (C1-C6)alkyl group that is substituted or not by one or more halogen atoms, (C3-C14)cycloalkyl groups or heterocyclic groups;

in the preparation of a medicament for the treatment of an inflammatory, metabolic, fibrotic and cholestatic disease.

16. The use according to claim 15, comprising:

(i) CVC or a pharmaceutically acceptable salt thereof; and

(ii) Elafibranor or a pharmaceutically acceptable salt thereof.

17. The use according to claim 15 or claim 16, further comprising at least one therapeutically active agent with known antifibrotic activity selected from pirfenidone or receptor tyrosine kinase inhibitors (RTKIs) such as Nintedanib, Sorafenib and other RTKIs, or angiotensin II (AT1) receptor blockers, or CTGF inhibitor, or any antifibrotic compound susceptible to interfere with the TGFβ and BMP-activated pathways including activators of the latent TGFβ complex such as MMP2, MMP9, THBS1 or cell-surface integrins, TGFβ receptors type I (TGFBRI) or type II (TGFBRII) and their ligands such as TGFβ, Activin, inhibin, Nodal, anti-Müllerian hormone, GDFs or BMPs, auxiliary co-receptors (also known as type III receptors), or components of the SMAD-dependent canonical pathway including regulatory or inhibitory SMAD proteins, or members of the SMAD-independent or non-canonical pathways including various branches of MAPK signaling, TAK1, Rho-like GTPase signaling pathways, phosphatidylinositol-3 kinase/AKT pathways, TGFβ-induced EMT process, or canonical and non-canonical Hedgehog signaling pathways including Hh ligands or target genes, or any members of the WNT, or Notch pathways which are susceptible to influence TGFβ.

18. The use according to any one of claims 15 to 17, further comprising at least one therapeutically active agent selected from JAK/STAT inhibitors and other anti-inflammatory and/or immunosuppressant agents.

19. Use of the combination product according to any one of claims 1 to 8, in the preparation of a medicament for the treatment of an inflammatory, metabolic, fibrotic and cholestatic disease.

20. The use according to any one of claims 15 to 19, wherein the fibrotic disorder is selected in the group consisting of liver, kidney, skin, epidermis, endodermis, muscle, tendon, cartilage, heart, pancreas, lung, uterus, nervous system, testis, ovary, adrenal gland, artery, vein, colon, intestine (e.g. small intestine), biliary tract, soft tissue (e.g. mediastinum or retroperitoneum), bone marrow, joint and stomach fibrosis.

21. The use according to any one of claims 15 to 20, wherein the fibrotic disorder is selected in the group consisting of liver, gut, lung, heart, kidney, muscle, skin, soft tissue, bone marrow, intestinal, and joint fibrosis.

22. The use according to any one of claims 15 to 21, wherein the disease is selected in the group consisting of metabolic liver diseases, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), drug-induced liver diseases, alcohol-induced liver diseases, infectious agent induced liver diseases, inflammatory liver diseases, immune system dysfunction-mediated liver diseases, dyslipidemia, cardiovascular diseases, restenosis, syndrome X, metabolic syndrome, diabetes, obesity, hypertension, chronic cholangiopathies such as Primary Sclerosing Cholangitis (PSC), Primary Biliary Cholangitis (PBC), biliary atresia, familial intrahepatic cholestasis type 3 (PFIC3), inflammatory bowel diseases, Crohn's disease, ulcerative colitis, keloid, old myocardial infarction, scleroderma/systemic sclerosis, inflammatory diseases, neurodegenerative diseases, cancers, liver cancer, hepatocallular carcinoma, gastrointestinal cancer, gastric cancer, meningioma associated with neurofibromatosis, pancreatic neuroendocrine tumors, pancreatic exocrine tumors, leukemia, myeloproliferative/myelodisplastic diseases, mastocytosis, dermatofibrosarcoma, solid tumors including breast, lung, thyroid or colorectal cancer, a prostate cancer, liver fibrosis or cirrhosis of any origin, metabolic disease-induced liver fibrosis or cirrhosis, NAFLD-induced fibrosis or cirrhosis, NASH-induced fibrosis or cirrhosis, alcohol-induced liver fibrosis or cirrhosis, drug-induced liver fibrosis or cirrhosis, infectious agent-induced liver fibrosis or cirrhosis, parasite infection-induced liver fibrosis or cirrhosis, bacterial infection-induced liver fibrosis or cirrhosis, viral infection-induced fibrosis or cirrhosis, HBV-infection induced liver fibrosis or cirrhosis, HCV-infection induced liver fibrosis or cirrhosis, HIV-infection induced liver fibrosis or cirrhosis, dual HCV and HIV-infection induced liver fibrosis or cirrhosis, radiation- or chemotherapy-induced fibrosis or cirrhosis, biliary tract fibrosis, liver fibrosis or cirrhosis due to any chronic cholestatic disease, gut fibrosis of any etiology, Crohn's disease-induced fibrosis, ulcerative colitis-induced fibrosis, intestine (e.g. small intestine) fibrosis, colon fibrosis, stomach fibrosis, skin fibrosis, epidermis fibrosis, endodermis fibrosis, skin fibrosis due to scleroderma/systemic sclerosis, lung fibrosis, lung fibrosis consecutive to chronic inflammatory airway diseases, such as COPD, asthma, emphysema, smoker's lung,tuberculosis, pulmonary fibrosis, idiopathic pulmonary fibrosis (IPF), heart fibrosis, kidney fibrosis, nephrogenic systemic fibrosis, muscle fibrosis, soft tissue (e.g. mediastinum or retroperitoneum) fibrosis, bone marrow fibrosis, joint fibrosis, tendon fibrosis, cartilage fibrosis, pancreas fibrosis, uterus fibrosis, nervous system fibrosis, testis fibrosis, ovary fibrosis, adrenal gland fibrosis, artery fibrosis, vein fibrosis, eye fibrosis, endomyocardial fibrosis, mediastinal fibrosis, myelofibrosis, retroperitoneal fibrosis, progressive massive fibrosis (a complication of coal workers' pneumoconiosis), proliferative

fibrosis, neoplastic fibrosis, peri-implantational fibrosis and asbestosis, arthrofibrosis, adhesive capsulitis.

23. The use according to any one of claims 15 to 21, wherein the disease is selected in the group consisting of, the disease is selected in the group consisting of metabolic liver diseases, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), drug-induced liver diseases, alcohol-induced liver diseases, infectious agent induced liver diseases, inflammatory liver diseases, immune system dysfunction-mediated liver diseases, dyslipidemia, cardiovascular diseases, restenosis, syndrome X, metabolic syndrome, diabetes, obesity, hypertension, chronic cholangiopathies such as Primary Sclerosing Cholangitis (PSC), Primary Biliary Cholangitis (PBC), biliary atresia, familial intrahepatic cholestasis type 3 (PFIC3), inflammatory bowel diseases, Crohn's disease, ulcerative colitis, liver cancer, hepatocallular carcinoma, gastrointestinal cancer, gastric cancer, colorectal cancer, metabolic disease-induced liver fibrosis or cirrhosis, NAFLD-induced fibrosis or cirrhosis, NASH-induced fibrosis or cirrhosis, alcohol-induced liver fibrosis or cirrhosis, drug-induced liver fibrosis or cirrhosis, infectious agent-induced liver fibrosis or cirrhosis, parasite infection-induced liver fibrosis or cirrhosis, bacterial infection-induced liver fibrosis or cirrhosis, viral infection-induced fibrosis or cirrhosis, HBV-infection induced liver fibrosis or cirrhosis, HCV-infection induced liver fibrosis or cirrhosis, HIV-infection induced liver fibrosis or cirrhosis, dual HCV and HIV-infection induced liver fibrosis or cirrhosis, radiation- or chemotherapy-induced fibrosis or cirrhosis, biliary tract fibrosis, liver fibrosis or cirrhosis due to any chronic cholestatic disease, gut fibrosis of any etiology, Crohn's disease-induced fibrosis, ulcerative colitis-induced fibrosis, intestine (e.g. small intestine) fibrosis, colon fibrosis, stomach fibrosis, lung fibrosis, lung fibrosis consecutive to chronic inflammatory airway diseases, such as COPD, asthma, emphysema, smoker's lung, tuberculosis, pulmonary fibrosis, idiopathic pulmonary fibrosis (IPF).