

Plasma concentration of TMC114 (ng/ml)
Time after administration (hr)
Plasma concentration of TMC114 (ng/ml)
Time after administration (hr)

Figure 1



Medication=mean
Plasma_concentration_of_TMC114 (ng/ml)
Time_after_administration (hr)
Medication=mean
Plasma_concentration_of_TMC114 (ng/ml)
Time_after_administration (hr)

Figure 2



[RTV] (μM)
0,09
0,08
0,07
0,06
0,05
0,04
0,03
0,02
0,01
0
0 0,002 0,004 0,006 0,008 0,01
[TMC114] (μM)
[IDV] (μM)
0,05
0,045
0,04
0,035
0,03
0,025
0,02
0,015
0,01
0,005
0
0 0,002 0,004 0,006 0,008 0,01
[TMC114] (μM)
[NFV] (μM)
0,08
0,07
0,06
0,05
0,04
0,03
0,02
0,01
0
0 0,002 0,004 0,006 0,008 0,01
[TMC114] (μM)
[SQV] (μM)
0,03
0,025
0,02
0,015
0,01
0,005
0
0 0,002 0,004 0,006 0,008 0,01
[TMC114] (μM)
[APV] (μM)
0,08
0,07
0,06
0,05
0,04
0,03
0,02
0,01
0
0 0,002 0,004 0,006 0,008
[TMC114] (μM)

Figure 3




Session I
Session II
10000
1000
100
10
1
conc (ng/ml)
0
12
24
36
48
60
72
time (h)

Figure 4


400 mg b.i.d.
800 mg b.i.d.
800 mg t.i.d
1200 mg t.i.d
conc (ng/ml)
time (day)



I 200/100 o.d.
II 400/100 o.d.
III 300/100 b.i.d.
IV 600/200 o.d.
V 1200/200 o.d.
Conc. (ng/ml)
Time (day)

Figure 5



Placebo
Compound of formula (4)
Compound of formula (4)/RTV
0
10
20
30
40
50
60
70
80
90
100
%
Diarrhea
Rash
Pruritis
Hyperbilirubinemia
Headache
Abdominal pain
Vomiting
Nausea
Bowel sounds
Flatulence
Appetite decrease
Fatigue

Figure 6