

NUCLEOSIDE PHOSPHORAMIDATES

Priority

This application is claims priority to U.S. Patent Application No.
5 61/319,513, filed March 31, 2010, U.S. Patent Application No. 61/319,548, filed
March 31, 2010, and U.S. Patent Application No. 12/783,680, filed May 20, 2010,
the subject matter of which are incorporated by reference in their entirety.

Field of the Invention

10

Disclosed herein are nucleoside phosphoramidates and their use as agents for
treating viral diseases. These compounds are inhibitors of RNA-dependent RNA
viral replication and are useful as inhibitors of HCV NS5B polymerase, as inhibitors
of HCV replication and for treatment of hepatitis C infection in mammals.

15

Background

Hepatitis C virus (HCV) infection is a major health problem that leads to
chronic liver disease, such as cirrhosis and hepatocellular carcinoma, in a substantial
20 number of infected individuals, estimated to be 2-15% of the world's population.
There are an estimated 4.5 million infected people in the United States alone,
according to the U.S. Center for Disease Control. According to the World Health
Organization, there are more than 200 million infected individuals worldwide, with
at least 3 to 4 million people being infected each year. Once infected, about 20% of
25 people clear the virus, but the rest can harbor HCV the rest of their lives. Ten to
twenty percent of chronically infected individuals eventually develop liver-
destroying cirrhosis or cancer. The viral disease is transmitted parenterally by
contaminated blood and blood products, contaminated needles, or sexually and
vertically from infected mothers or carrier mothers to their offspring. Current
30 treatments for HCV infection, which are restricted to immunotherapy with
recombinant interferon- α alone or in combination with the nucleoside analog
ribavirin, are of limited clinical benefit. Moreover, there is no established vaccine
for HCV. Consequently, there is an urgent need for improved therapeutic agents
that effectively combat chronic HCV infection.

The HCV virion is an enveloped positive-strand RNA virus with a single oligoribonucleotide genomic sequence of about 9600 bases which encodes a polyprotein of about 3,010 amino acids. The protein products of the HCV gene consist of the structural proteins C, E1, and E2, and the non-structural proteins NS2, NS3, NS4A and NS4B, and NS5A and NS5B. The nonstructural (NS) proteins are believed to provide the catalytic machinery for viral replication. The NS3 protease releases NS5B, the RNA-dependent RNA polymerase from the polyprotein chain. HCV NS5B polymerase is required for the synthesis of a double-stranded RNA from a single-stranded viral RNA that serves as a template in the replication cycle of HCV. Therefore, NS5B polymerase is considered to be an essential component in the HCV replication complex (K. Ishi, et al, *Hepatology*, 1999, 29: 1227-1235; V. Lohmann, et al., *Virology*, 1998, 249: 108-118). Inhibition of HCV NS5B polymerase prevents formation of the double-stranded HCV RNA and therefore constitutes an attractive approach to the development of HCV-specific antiviral therapies.

HCV belongs to a much larger family of viruses that share many common features.

Flaviviridae Viruses

The *Flaviviridae* family of viruses comprises at least three distinct genera: *pestiviruses*, which cause disease in cattle and pigs; *flaviviruses*, which are the primary cause of diseases such as dengue fever and yellow fever; and *hepaciviruses*, whose sole member is HCV. The flavivirus genus includes more than 68 members separated into groups on the basis of serological relatedness (Calisher et al., *J. Gen. Virol*, 1993,70,37-43). Clinical symptoms vary and include fever, encephalitis and hemorrhagic fever (*Fields Virology*, Editors: Fields, B. N., Knipe, D. M., and Howley, P. M., Lippincott-Raven Publishers, Philadelphia, PA, 1996, Chapter 31, 931-959). Flaviviruses of global concern that are associated with human disease include the Dengue Hemorrhagic Fever viruses (DHF), yellow fever virus, shock syndrome and Japanese encephalitis virus (Halstead, S. B., *Rev. Infect. Dis.*, 1984, 6, 251-264; Halstead, S. B., *Science*, 239:476-481, 1988; Monath, T. P., *New Eng. J. Med*, 1988, 319, 641-643).

The pestivirus genus includes bovine viral diarrhea virus (BVDV), classical swine fever virus (CSFV, also called hog cholera virus) and border disease virus (BDV) of sheep (Moennig, V. et al. *Adv. Vir. Res.* 1992, 41, 53-98). Pestivirus

infections of domesticated livestock (cattle, pigs and sheep) cause significant economic losses worldwide. BVDV causes mucosal disease in cattle and is of significant economic importance to the livestock industry (Meyers, G. and Thiel, H.J., *Advances in Virus Research*, 1996, 47, 53-118; Moennig V., et al, *Adv. Vir. Res.* 1992, 41, 53-98). Human pestiviruses have not been as extensively characterized as the animal pestiviruses. However, serological surveys indicate considerable pestivirus exposure in humans.

Pestiviruses and hepaciviruses are closely related virus groups within the *Flaviviridae* family. Other closely related viruses in this family include the GB virus A, GB virus A-like agents, GB virus-B and GB virus-C (also called hepatitis G virus, HGV). The hepacivirus group (hepatitis C virus; HCV) consists of a number of closely related but genotypically distinguishable viruses that infect humans. There are at least 6 HCV genotypes and more than 50 subtypes. Due to the similarities between pestiviruses and hepaciviruses, combined with the poor ability of hepaciviruses to grow efficiently in cell culture, bovine viral diarrhea virus (BVDV) is often used as a surrogate to study the HCV virus.

The genetic organization of pestiviruses and hepaciviruses is very similar. These positive stranded RNA viruses possess a single large open reading frame (ORF) encoding all the viral proteins necessary for virus replication. These proteins are expressed as a polyprotein that is co- and post-translationally processed by both cellular and virus-encoded proteinases to yield the mature viral proteins. The viral proteins responsible for the replication of the viral genome RNA are located within approximately the carboxy-terminal. Two-thirds of the ORF are termed nonstructural (NS) proteins. The genetic organization and polyprotein processing of the nonstructural protein portion of the ORF for pestiviruses and hepaciviruses is very similar. For both the pestiviruses and hepaciviruses, the mature nonstructural (NS) proteins, in sequential order from the amino-terminus of the nonstructural protein coding region to the carboxy-terminus of the ORF, consist of p7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.

The NS proteins of pestiviruses and hepaciviruses share sequence domains that are characteristic of specific protein functions. For example, the NS3 proteins of viruses in both groups possess amino acid sequence motifs characteristic of serine proteinases and of helicases (Gorbalenya et al., *Nature*, 1988, 333, 22; Bazan and Fletterick *Virology*, 1989, 171, 637-639; Gorbalenya et al., *Nucleic Acid Res.*, 1989,

17, 3889-3897). Similarly, the NS5B proteins of pestiviruses and hepaciviruses have the motifs characteristic of RNA-directed RNA polymerases (Koonin, E.V. and Dolja, V.V., *Crit. Rev. Biochem. Molec. Biol.* 1993, 28, 375-430).

The actual roles and functions of the NS proteins of pestiviruses and
5 hepaciviruses in the lifecycle of the viruses are directly analogous. In both cases, the NS3 serine proteinase is responsible for all proteolytic processing of polyprotein precursors downstream of its position in the ORF (Wiskerchen and Collett, *Virology*, 1991, 184, 341-350; Bartenschlager et al., *J. Virol.* 1993, 67, 3835-3844; Eckart et al. *Biochem. Biophys. Res. Comm.* 1993, 192, 399-406; Grakoui et al., *J.*
10 *Virol.* 1993, 67, 2832-2843; Grakoui et al., *Proc. Natl. Acad. Sci. USA* 1993, 90, 10583-10587; Hijikata et al., *J. Virol.* 1993, 67, 4665-4675; Tome et al., *J. Virol.*, 1993, 67, 4017-4026). The NS4A protein, in both cases, acts as a cofactor with the NS3 serine protease (Bartenschlager et al., *J. Virol.* 1994, 68, 5045-5055; Failla et al., *J. Virol.* 1994, 68, 3753-3760; Xu et al., *J. Virol.*, 1997, 71:53 12-5322). The
15 NS3 protein of both viruses also functions as a helicase (Kim et al., *Biochem. Biophys. Res. Comm.*, 1995, 215, 160-166; Jin and Peterson, *Arch. Biochem. Biophys.*, 1995, 323, 47-53; Warrenner and Collett, *J. Virol.* 1995, 69, 1720-1726). Finally, the NS5B proteins of pestiviruses and hepaciviruses have the predicted RNA-directed RNA polymerases activity (Behrens et al., *EMBO*, 1996, 15, 12-22;
20 Lechmann et al., *J. Virol.*, 1997, 71, 8416-8428; Yuan et al., *Biochem. Biophys. Res. Comm.* 1997, 232, 231-235; Hagedorn, PCT WO 97/12033; Zhong et al, *J. Virol.*, 1998, 72, 9365-9369).

Currently, there are limited treatment options for individuals infected with hepatitis C virus. The current approved therapeutic option is the use of
25 immunotherapy with recombinant interferon- α alone or in combination with the nucleoside analog ribavirin. This therapy is limited in its clinical effectiveness and only 50% of treated patients respond to therapy. Therefore, there is significant need for more effective and novel therapies to address the unmet medical need posed by HCV infection.

30 A number of potential molecular targets for drug development of direct acting antivirals as anti-HCV therapeutics have now been identified including, but not limited to, the NS2-NS3 autoprotease, the NS3 protease, the NS3 helicase and the NS5B polymerase. The RNA-dependent RNA polymerase is absolutely essential for

replication of the single-stranded, positive sense, RNA genome and this enzyme has elicited significant interest among medicinal chemists.

Inhibitors of HCV NS5B as potential therapies for HCV infection have been reviewed: Tan, S.-L., et al., *Nature Rev. Drug Discov.*, 2002, 1, 867-881; Walker, M.P. et al., *Exp. Opin. Investigational Drugs*, 2003, 12, 1269-1280; Ni, Z.-J., et al., *Current Opinion in Drug Discovery and Development*, 2004, 7, 446-459; Beaulieu, P. L., et al., *Current Opinion in Investigational Drugs*, 2004, 5, 838-850; Wu, J., et al., *Current Drug Targets-Infectious Disorders*, 2003, 3, 207-219; Griffith, R.C., et al., *Annual Reports in Medicinal Chemistry*, 2004, 39, 223-237; Carrol, S., et al., *Infectious Disorders-Drug Targets*, 2006, 6, 17-29. The potential for the emergence of resistant HCV strains and the need to identify agents with broad genotype coverage supports the need for continuing efforts to identify novel and more effective nucleosides as HCV NS5B inhibitors.

Nucleoside inhibitors of NS5B polymerase can act either as a non-natural substrate that results in chain termination or as a competitive inhibitor which competes with nucleotide binding to the polymerase. To function as a chain terminator the nucleoside analog must be taken up by the cell and converted *in vivo* to a triphosphate to compete for the polymerase nucleotide binding site. This conversion to the triphosphate is commonly mediated by cellular kinases which imparts additional structural requirements on a potential nucleoside polymerase inhibitor. Unfortunately, this limits the direct evaluation of nucleosides as inhibitors of HCV replication to cell-based assays capable of *in situ* phosphorylation.


In some cases, the biological activity of a nucleoside is hampered by its poor substrate characteristics for one or more of the kinases needed to convert it to the active triphosphate form. Formation of the monophosphate by a nucleoside kinase is generally viewed as the rate limiting step of the three phosphorylation events. To circumvent the need for the initial phosphorylation step in the metabolism of a nucleoside to the active triphosphate analog, the preparation of stable phosphate prodrugs has been reported. Nucleoside phosphoramidate prodrugs have been shown to be precursors of the active nucleoside triphosphate and to inhibit viral replication when administered to viral infected whole cells (McGuigan, C., et al., *J. Med. Chem.*, 1996, 39, 1748-1753; Valette, G., et al., *J. Med. Chem.*, 1996, 39, 1981-1990; Balzarini, J., et al., *Proc. National Acad Sci USA*, 1996, 93, 7295-7299; Siddiqui, A. Q., et al., *J. Med. Chem.*, 1999, 42, 4122-4128; Eisenberg, E. J., et al.,

Nucleosides, Nucleotides and Nucleic Acids, 2001, 20, 1091-1098; Lee, W.A., et al., *Antimicrobial Agents and Chemotherapy*, 2005, 49, 1898); US 2006/0241064; and WO 2007/095269.


Also limiting the utility of nucleosides as viable therapeutic agents is their sometimes poor physicochemical and pharmacokinetic properties. These poor properties can limit the intestinal absorption of an agent and limit uptake into the target tissue or cell. To improve on their properties prodrugs of nucleosides have been employed. It has been demonstrated that preparation of nucleoside phosphoramidates improves the systemic absorption of a nucleoside and furthermore, the phosphoramidate moiety of these "pronucleotides" is masked with neutral lipophilic groups to obtain a suitable partition coefficient to optimize uptake and transport into the cell dramatically enhancing the intracellular concentration of the nucleoside monophosphate analog relative to administering the parent nucleoside alone. Enzyme-mediated hydrolysis of the phosphate ester moiety produces a nucleoside monophosphate wherein the rate limiting initial phosphorylation is unnecessary. To this end, U.S. Patent Application 12/053,015, which corresponds to WO 2008/121634 and US 2010/0016251, discloses a number of phosphoramidate nucleoside prodrugs, many of which show activity in an HCV assay. Several compounds disclosed in US 2010/0016251 were tested as a potential clinical candidate for approval by the FDA.

Summary of the Invention


Disclosed herein is a compound represented by formula 4 and its respective phosphorus-based diastereomers represented by formulas *S_P*-4 and *R_P*-4.



4



S_P-4



R_P-4

Brief Description of the Drawings

- 5
- Figure 1.** High resolution XRPD diffractogram of **4**.
- Figure 2.** High resolution XRPD diffractogram of *R_P-4*.
- 10 **Figure 3.** High resolution XRPD diffractogram of *S_P-4* (Form 1).
- Figure 4.** High resolution XRPD diffractogram of *S_P-4* (Form 1).
- Figure 5.** High resolution XRPD diffractogram of *S_P-4*·CH₂Cl₂ (Form 2).
- 15 **Figure 6.** High resolution XRPD diffractogram of *S_P-4*·CHCl₃ (Form 3).
- Figure 7.** High resolution XRPD diffractogram of *S_P-4* (Form 4).
- 20 **Figure 8.** High resolution XRPD diffractogram of *S_P-4* (Form 5).
- Figure 9.** High resolution XRPD diffractogram of *S_P-4* (amorphous).
- Figure 10.** X-Ray Crystal Structure for *S_P-4* (Form 1)
- 25 **Figure 11.** X-Ray Crystal (Isotropic) Structure for *S_P-4*·CH₂Cl₂ (Form 2)
- Figure 12.** X-Ray Crystal (Anisotropic) Structure for *S_P-4*·CH₂Cl₂ (Form 2)
- 30 **Figure 13.** X-Ray Crystal Structure for *S_P-4*·CHCl₃ (Form 3)
- Figure 14.** FT-IR spectrum of **4**.
- Figure 15.** FT-IR spectrum of *R_P-4*.
- 35 **Figure 16.** FT-IR spectrum of *S_P-4*

Figure 17. TGA and DSC analysis of **4**.

Figure 18. TGA and DSC analysis of *R_P*-**4**.

5 **Figure 19.** TGA and DSC analysis of *S_P*-**4**.

Figure 20A. X-Ray Crystal Structure for **8** (*S_P*-isomer) (molecule no. 1 of the asymmetric unit).

10 **Figure 20B.** X-Ray Crystal Structure for **8** (*S_P*-isomer) (molecule no. 2 of the asymmetric unit).

Figure 21. High resolution XRPD diffractogram of *S_P*-**4** (Form 6).

15 **Figure 22A.** X-Ray Crystal Structure for (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate (molecule no. 1 of the asymmetric unit).

20 **Figure 22B.** X-Ray Crystal Structure for (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate (molecule no. 2 of the asymmetric unit).

Detailed Description of the Invention

25 *Definitions*

The phrase "a" or "an" entity as used herein refers to one or more of that entity; for example, a compound refers to one or more compounds or at least one compound. As such, the terms "a" (or "an"), "one or more", and "at least one" can be used interchangeably herein.

30 The terms "optional" or "optionally" as used herein means that a subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "optional bond" means that the bond may or may not be present, and that the description includes single, double, or triple bonds.

35 The term "P*" means that the phosphorus atom is chiral and that it has a corresponding Cahn-Ingold-Prelog designation of "R" or "S" which have their accepted plain meanings.

The term "purified," as described herein, refers to the purity of a given compound. For example, a compound is "purified" when the given compound is a
40 major component of the composition, i.e., at least 50% w/w pure. Thus, "purified"

embraces at least 50% w/w purity, at least 60% w/w purity, at least 70% purity, at least 80% purity, at least 85% purity, at least 90% purity, at least 92% purity, at least 94% purity, at least 96% purity, at least 97% purity, at least 98% purity, at least 99% purity, at least 99.5% purity, and at least 99.9% purity, wherein "substantially pure" embraces at least 97% purity, at least 98% purity, at least 99% purity, at least 99.5% purity, and at least 99.9% purity

The term "metabolite," as described herein, refers to a compound produced in vivo after administration to a subject in need thereof.

The term "about" (also represented by ~) means that the recited numerical value is part of a range that varies within standard experimental error.

The expression "substantially as shown in..." a specified XRPD pattern means that the peak positions shown in the XRPD pattern are substantially the same, within visual inspection or resort to selected peak listings ($\pm 0.2^\circ 2\theta$). One of ordinary skill understands that the intensities can vary depending on the sample.

The term "substantially anhydrous" means that a substance contains at most 10% by weight of water, preferably at most 1% by weight of water, more preferably at most 0.5% by weight of water, and most preferably at most 0.1% by weight of water.

A solvent or anti-solvent (as used in reactions, crystallization, etc. or lattice and/or adsorbed solvents) includes at least one of a C₁ to C₈ alcohol, a C₂ to C₈ ether, a C₃ to C₇ ketone, a C₃ to C₇ ester, a C₁ to C₂ chlorocarbon, a C₂ to C₇ nitrile, a miscellaneous solvent, a C₅ to C₁₂ saturated hydrocarbon, and a C₆ to C₁₂ aromatic hydrocarbon.

The C₁ to C₈ alcohol refers to a straight/branched and/or cyclic/acyclic alcohol having such number of carbons. The C₁ to C₈ alcohol includes, but is not limited to, methanol, ethanol, n-propanol, isopropanol, isobutanol, hexanol, and cyclohexanol.

The C₂ to C₈ ether refers to a straight/branched and/or cyclic/acyclic ether having such number of carbons. The C₂ to C₈ ether includes, but is not limited to, dimethyl ether, diethyl ether, di-isopropyl ether, di-n-butyl ether, methyl-t-butyl ether (MTBE), tetrahydrofuran, and dioxane

The C₃ to C₇ ketone refers to a straight/branched and/or cyclic/acyclic ketone having such number of carbons. The C₃ to C₇ ketone includes, but is not limited to,

acetone, methyl ethyl ketone, propanone, butanone, methyl isobutyl ketone, methyl butyl ketone, and cyclohexanone.

The C₃ to C₇ ester refers to a straight/branched and/or cyclic/acyclic ester having such number of carbons. The C₃ to C₇ ester includes, but is not limited to, ethyl acetate, propyl acetate, n-butyl acetate, etc.

The C₁ to C₂ chlorocarbon refers to a chlorocarbon having such number of carbons. The C₁ to C₂ chlorocarbon includes, but is not limited to, chloroform, methylene chloride (DCM), carbon tetrachloride, 1,2-dichloroethane, and tetrachloroethane.

A C₂ to C₇ nitrile refers to a nitrile have such number of carbons. The C₂ to C₇ nitrile includes, but is not limited to, acetonitrile, propionitrile, etc.

A miscellaneous solvent refers to a solvent commonly employed in organic chemistry, which includes, but is not limited to, diethylene glycol, diglyme (diethylene glycol dimethyl ether), 1,2-dimethoxy-ethane, dimethylformamide, dimethylsulfoxide, ethylene glycol, glycerin, hexamethylphosphoramide, hexamethylphosphorous triamide, N-methyl-2-pyrrolidinone, nitromethane, pyridine, triethyl amine, and acetic acid.

The term C₅ to C₁₂ saturated hydrocarbon refers to a straight/branched and/or cyclic/acyclic hydrocarbon. The C₅ to C₁₂ saturated hydrocarbon includes, but is not limited to, n-pentane, petroleum ether (ligroine), n-hexane, n-heptane, cyclohexane, and cycloheptane.

The term C₆ to C₁₂ aromatic refers to substituted and unsubstituted hydrocarbons having a phenyl group as their backbone. Preferred hydrocarbons include benzene, xylene, toluene, chlorobenzene, o-xylene, m-xylene, p-xylene, xylenes, with toluene being more preferred.

The term "halo" or "halogen" as used herein, includes chloro, bromo, iodo and fluoro.

The term "blocking group" refers to a chemical group which exhibits the following characteristics. The "group" is derived from a "protecting compound." Groups that are selective for primary hydroxyls over secondary hydroxyls that can be put on under conditions consistent with the stability of the phosphoramidate (pH 2-8) and impart on the resulting product substantially different physical properties allowing for an easier separation of the 3'-phosphoramidate-5'-new group product from the unreacted desired compound. The group must react selectively in good

yield to give a protected substrate that is stable to the projected reactions (see Protective Groups in Organic Synthesis, 3rd ed. T. W. Greene and P. G. M. Wuts, John Wiley & Sons, New York, N.Y., 1999). Examples of groups include, but are not limited to: benzoyl, acetyl, phenyl-substituted benzoyl, tetrahydropyranyl, trityl, DMT (4,4'-dimethoxytrityl), MMT (4-monomethoxytrityl), trimethoxytrityl, pixyl (9-phenylxanthen-9-yl) group, thiopixyl (9-phenylthioxanthen-9-yl) or 9-(p-methoxyphenyl)xanthine-9-yl (MOX), etc.; C(O)-alkyl, C(O)Ph, C(O)aryl, CH₂O-alkyl, CH₂O-aryl, SO₂-alkyl, SO₂-aryl, *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl. Acetals, such as MOM or THP and the like are considered possible groups. Fluorinated compounds are also contemplated in so far that they can be attached to the compound and can be selectively removed by passing through a fluorous solid phase extraction media (FluoroFlash[®]). A specific example includes a fluorinated trityl analog, trityl analog 1-[4-(1*H*,1*H*,2*H*,2*H*-perfluorodecyl)phenyl]-1,1-diphenylmethanol. Other fluorinated analogs of trityl, BOC, Fmoc, CBz, etc. are also contemplated. Sulfonyl chlorides like p-toluenesulfonyl chloride can react selectively on the 5' position. Esters could be formed selectively such as acetates and benzoates. Dicarboxylic anhydrides such as succinic anhydride and its derivatives can be used to generate an ester linkage with a free carboxylic acid, such examples include, but are not limited to oxalyl, malonyl, succinyl, glutaryl, adipyl, pimelyl, superyl, azeloyl, sebacoyl, phthalyl, isophthalyl, terephthalyl, etc. The free carboxylic acid increases the polarity dramatically and can also be used as a handle to extract the reaction product into mildly basic aqueous phases such as sodium bicarbonate solutions. The phosphoramidate group is relatively stable in acidic media, so groups requiring acidic reaction conditions, such as, tetrahydropyranyl, could also be used.

The term "protecting group" which is derived from a "protecting compound," has its plain and ordinary meaning, i.e., at least one protecting or blocking group is bound to at least one functional group (e.g., -OH, -NH₂, etc.) that allows chemical modification of at least one other functional group. Examples of protecting groups, include, but are not limited to, benzoyl, acetyl, phenyl-substituted benzoyl, tetrahydropyranyl, trityl, DMT (4,4'-dimethoxytrityl), MMT (4-monomethoxytrityl), trimethoxytrityl, pixyl (9-phenylxanthen-9-yl) group, thiopixyl (9-phenylthioxanthen-9-yl) or 9-(p-methoxyphenyl)xanthine-9-yl (MOX), etc.; C(O)-alkyl, C(O)Ph, C(O)aryl, C(O)O(lower alkyl), C(O)O(lower alkylene)aryl (e.g., -

C(O)OCH₂Ph), C(O)O-aryl, CH₂O-alkyl, CH₂O-aryl, SO₂-alkyl, SO₂-aryl, a protecting group comprising at least one silicon atom, such as, *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl, Si(lower alkyl)₂OSi(lower alkyl)₂OH (such as, –Si(^{*i*}Pr)₂OSi(^{*i*}Pr)₂OH).

5 The term "protecting compound," as used herein and unless otherwise defined, refers to a compound that contains a "protecting group" and that is capable of reacting with a compound that contains functional groups that are capable of being protected.

 The term "leaving group", as used herein, has the same meaning to the
10 skilled artisan (Advanced Organic Chemistry: reactions, mechanisms and structure—Fourth Edition by Jerry March, John Wiley and Sons Ed.; 1992 pages 351-357) and represents a group which is part of and attached to a substrate molecule; in a reaction where the substrate molecule undergoes a displacement reaction (with for example a nucleophile), the leaving group is then displaced. Examples of leaving groups
15 include, but are not limited to: halogen (F, Cl, Br, and I), preferably Cl, Br, or I; tosylate, mesylate, triflate, acetate, camphorsulfonate, aryloxy, and aryloxy substituted with at least one electron withdrawing group (e.g., *p*-nitrophenoxide, 2-chlorophenoxide, 4-chlorophenoxide, 2,4-dinitrophenoxide, pentafluorophenoxide, etc.), etc. The term "electron withdrawing group" is accorded its plain meaning
20 here. Examples of electron withdrawing groups include, but are not limited to, a halogen, –NO₂, –C(O)(lower alkyl), –C(O)(aryl), –C(O)O(lower alkyl), –C(O)O(aryl), etc.

 The term "basic reagent", as used herein, means a compound that is capable of deprotonating a hydroxyl group. Examples of basic reagents include, but are not
25 limited to, a (lower alk)oxide ((lower alkyl)OM) in combination with an alcoholic solvent, where (lower alk)oxides include, but are not limited to, MeO[–], EtO[–], ^{*n*}PrO[–], ^{*i*}PrO[–], ^{*t*}BuO[–], ^{*i*}AmO[–] (iso-amyloxy), etc., and where M is an alkali metal cation, such as Li⁺, Na⁺, K⁺, etc. Alcoholic solvents include (lower alkyl)OH, such as, for example, MeOH, EtOH, ^{*n*}PrOH, ^{*i*}PrOH, ^{*t*}BuOH, ^{*i*}AmOH, etc. Non-alkoxy bases can
30 also be used such as sodium hydride, sodium hexamethyldisilazane, lithium hexamethyldisilazane, lithium diisopropylamide, calcium hydride, sodium carbonate, potassium carbonate, cesium carbonate, DBU, DBN, Grignard reagents, such as (lower alkyl)Mg(halogen), which include but are not limited to MeMgCl, MeMgBr, ^{*t*}BuMgCl, ^{*t*}BuMgBr, etc.

The term "base" embraces the term "basic reagent" and is meant to be a compound that is capable of deprotonating a proton containing compound, i.e., a Brønsted base. In addition to the examples recited above, further examples of a base include, but are not limited to pyridine, collidine, 2,6-(lower alkyl)-pyridine, dimethyl-aniline, imidazole, N-methyl-imidazole, pyrazole, N-methyl-pyrazole, triethylamine, di-isopropylethylamine, etc.

The term "non-nucleophilic base" means a compound that is capable of acting as a Brønsted base, but has low nucleophilicity. Examples of non-nucleophilic bases include, but are not limited to, potassium carbonate, cesium carbonate, di-isopropylamine, di-isopropylethylamine, triethylamine, quinuclidine, naphthalene-1,8-diamine, 2,2,6,6-tetramethylpiperidine, 1,8-diazabicycloundec-7-ene, 4-dimethylamino-pyridine, pyridine, a 2,6-di-C1-6-alkyl-pyridine, a 2,4,6-tri-C1-6-alkyl-pyridine, 1,5-diazabicyclo[4.3.0]non-5-ene, and 1,4-diazabicyclo[2.2.2]octane.

The term "electron withdrawing group" is accorded its plain meaning. Examples of electron withdrawing groups include, but are not limited to, a halogen (F, Cl, Br, or I), $-\text{NO}_2$, $-\text{C}(\text{O})(\text{lower alkyl})$, $-\text{C}(\text{O})(\text{aryl})$, $-\text{C}(\text{O})\text{O}(\text{lower alkyl})$, $-\text{C}(\text{O})\text{O}(\text{aryl})$, etc.

The term "co-crystallates" include co-crystallates of **4**, R_P -**4**, or S_P -**4** in combination with salts, which embraces pharmaceutically acceptable salts.

The term "salts," as described herein, refers to a compound comprising a cation and an anion, which can be produced by the protonation of a proton-accepting moiety and/or deprotonation of a proton-donating moiety. It should be noted that protonation of the proton-accepting moiety results in the formation of a cationic species in which the charge is balanced by the presence of a physiological anion, whereas deprotonation of the proton-donating moiety results in the formation of an anionic species in which the charge is balanced by the presence of a physiological cation.

The phrase "pharmaceutically acceptable salt" means a salt that is pharmaceutically acceptable. Examples of pharmaceutically acceptable salts include, but are not limited to: (1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as glycolic acid, pyruvic acid, lactic acid, malonic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric

acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, lauryl

5 sulfuric acid, gluconic acid, glutamic acid, salicylic acid, muconic acid, and the like or (2) basic addition salts formed with the conjugate bases of any of the inorganic acids listed above, wherein the conjugate bases comprise a cationic component selected from among Na^+ , K^+ , Mg^{2+} , Ca^{2+} , NH_gR^{g+} , in which R^g is a C_{1-3} alkyl and g is a number selected from among 0, 1, 2, 3, or 4. It should be understood that all

10 references to pharmaceutically acceptable salts include solvent addition forms (solvates) or crystal forms (polymorphs) as defined herein, of the same acid addition salt.

The term "alkyl" refers to an unbranched or branched chain, saturated, monovalent hydrocarbon residue containing 1 to 30 carbon atoms. The term " C_{1-M} alkyl" refers to an alkyl comprising 1 to M carbon atoms, where M is an integer

15 having the following values: 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30. The term " C_{1-4} alkyl" refers to an alkyl containing 1 to 4 carbon atoms. The term "lower alkyl" denotes a straight or branched chain hydrocarbon residue comprising 1 to 6 carbon atoms. " C_{1-20} alkyl" as used herein refers to an alkyl comprising 1 to 20 carbon atoms. " C_{1-10} alkyl" as used herein refers to an alkyl comprising 1 to 10 carbons. Examples of alkyl groups include, but are not limited to, lower alkyl groups include methyl, ethyl, propyl, *i*-propyl, *n*-butyl, *i*-butyl, *t*-butyl or pentyl, isopentyl, neopentyl, hexyl, heptyl, and octyl. The term (ar)alkyl or (heteroaryl)alkyl indicate the alkyl group is optionally

20 substituted by an aryl or a heteroaryl group respectively.

The term "alkenyl" refers to an unsubstituted hydrocarbon chain radical having from 2 to 10 carbon atoms having one or two olefinic double bonds, preferably one olefinic double bond. The term " C_{2-N} alkenyl" refers to an alkenyl comprising 2 to N carbon atoms, where N is an integer having the following values:

30 3, 4, 5, 6, 7, 8, 9, or 10. The term " C_{2-10} alkenyl" refers to an alkenyl comprising 2 to 10 carbon atoms. The term " C_{2-4} alkenyl" refers to an alkenyl comprising 2 to 4 carbon atoms. Examples include, but are not limited to, vinyl, 1-propenyl, 2-propenyl (allyl) or 2-butenyl (crotyl).

The term "aryl," as used herein, and unless otherwise specified, refers to substituted or unsubstituted phenyl (Ph), biphenyl, or naphthyl, preferably the term aryl refers to substituted or unsubstituted phenyl. The aryl group can be substituted with one or more moieties selected from among hydroxyl, F, Cl, Br, I, amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, and phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in T.W. Greene and P.G. M. Wuts, "Protective Groups in Organic Synthesis," 3rd ed., John Wiley & Sons, 1999.

The term "aryloxide," as used herein, and unless otherwise specified, refers to substituted or unsubstituted phenoxide (PhO⁻), p-phenyl-phenoxide (p-Ph-PhO⁻), or naphthoxide, preferably the term aryloxide refers to substituted or unsubstituted phenoxide. The aryloxide group can be substituted with one or more moieties selected from among hydroxyl, F, Cl, Br, I, -C(O)(lower alkyl), -C(O)O(lower alkyl), amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, and phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in T.W. Greene and P.G. M. Wuts, "Protective Groups in Organic Synthesis," 3rd ed., John Wiley & Sons, 1999.

The term "preparation" or "dosage form" is intended to include both solid and liquid formulations of the active compound and one skilled in the art will appreciate that an active ingredient can exist in different preparations depending on the desired dose and pharmacokinetic parameters.

The term "excipient" as used herein refers to a compound that is used to prepare a pharmaceutical composition, and is generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes excipients that are acceptable for veterinary use as well as human pharmaceutical use.

The term "crystalline" refers to a situation where a solid sample of either *S_P-4* or *R_P-4* has crystalline characteristics when determined by X-ray powder diffraction or a single crystal X-ray technique.


The term "crystal-like" refers to a situation where a solid sample of either *S_P-4* or *R_P-4* has crystalline characteristics when determined by one means, e.g., visually or by optical or polarizing microscopy, but does not have crystalline characteristics when determined by another means, e.g., x-ray powder diffraction.

Methods of visually determining the crystallinity of a solid sample by visual or by optical or by polarizing microscopy are disclosed in USP <695> and <776>, both of which are incorporated by reference. A solid sample of either *S_P*-**4** or *R_P*-**4** that is "crystal-like" may be crystalline under certain conditions but may become non-crystalline when subjected to other conditions.

The term "amorphous" refers to a situation where a solid sample of either *S_P*-**4** or *R_P*-**4** is neither crystalline nor crystal-like.

Embodiments


A first embodiment is directed to a compound represented by formula **4**:



4

wherein P* represents a chiral phosphorus atom. Due to the chiral phosphorus atom, the compound represented by formula **4** comprises two diastereomers designated as *R_P*-**4** and *S_P*-**4**. The compound represented by formula **4** can also be part of a solvate, a hydrate, or a mixed solvate/hydrate. The solvate is designated as **4**·n*S*, while the hydrate is designated as **4**·mH₂O, where *S* is a lattice solvent, n varies by an integer or non-integer amount from about 0 to about 3 and m varies by an integer or non-integer amount from about 0 to about 5. Finally, the compound represented by formula **4** might not exist as a solvate or hydrate, but have a certain advantageous amount of adsorbed solvent (*S*) or water. In which case, the amount of *S* or water can vary from about 0 wt. % to about 10 wt.% based on the weight of the compound represented by formula **4**. The compound represented by formula **4** and its solvates and hydrates thereof is crystalline, crystal-like, or amorphous.

A second embodiment is directed to a compound represented by formula *R_P*-**4**:

*R_P-4*

The compound represented by formula *R_P-4* can also be part of a solvate, a hydrate, or a mixed solvate/hydrate. The solvate is designated as *R_P-4*·*nS*, while the hydrate is designated as *S_P-4*·*mH₂O*, where *S* is a lattice solvent, *n* varies by an integer or non-integer amount from about 0 to about 3 and *m* varies by an integer or non-integer amount from about 0 to about 5. Finally, the compound represented by formula *R_P-4* might not exist as a solvate, hydrate, or mixed solvate/hydrate, but have a certain advantageous amount of adsorbed solvent (*S*), water, or both *S* and water. In which case, the amount of *S* or water can vary from about 0 wt. % to about 10 wt.% based on the weight of the compound represented by formula *R_P-4*. The compound represented by formula *R_P-4* and its solvates and hydrates thereof is crystalline, crystal-like, or amorphous.

A first aspect of the second embodiment is directed to crystalline *R_P-4*.

A second aspect of the second embodiment is directed to crystalline *R_P-4* having XRPD 2θ-reflections (°) at about: 6.6, 7.1, 9.0, 11.6, 17.9, 20.7, 24.1, 24.4, and 26.2.

A third aspect of the second embodiment is directed to a crystalline *R_P-4* having XRPD 2θ-reflections (°) at about: 6.6, 7.1, 9.0, 11.0, 11.6, 12.0, 16.0, 17.9, 19.6, 20.7, 21.0, 21.7, 21.9, 22.2, 23.1, 24.1, 24.4, 26.1, 27.3, 27.7, and 28.2.

A fourth aspect of the second embodiment is directed to crystalline *R_P-4* having an XRPD diffraction pattern substantially as that shown in Fig. 2.

A fifth aspect of the second embodiment is directed to *R_P-4* having the following FT-IR peaks (cm⁻¹): 1742, 1713, 1679, 1460, 1377, 1259, 1157, and 1079.


A sixth aspect of the second embodiment is directed to *R_P-4* having an FT-IR spectrum substantially as that shown in Fig. 15.

A seventh aspect of the second embodiment is directed to substantially pure *R_P-4*.

An eighth aspect of the second embodiment is directed to substantially pure crystalline *R_P-4*.

A ninth aspect of the second embodiment is directed to substantially pure amorphous *R_P-4*.

5 A third embodiment is directed to a compound represented by formula *S_P-4*:



S_P-4

The compound represented by formula *S_P-4* can also be part of a solvate, a hydrate, or a mixed solvate/hydrate. The solvate is designated as *S_P-4·nS*, while the hydrate is designated as *S_P-4·mH₂O*, where *S* is a lattice solvent, *n* varies in an integer or non-integer amount from about 0 to about 3 and *m* varies in an integer or non-integer amount from about 0 to about 5. Finally, the compound represented by formula *S_P-4* might not exist as a solvate or hydrate, but have a certain advantageous amount of adsorbed solvent (*S*) or water. In which case, the amount of *S* or water can vary from about 0 wt. % to about 10 wt.% based on the weight of the compound represented by formula *S_P-4*. The compound represented by formula *S_P-4* and its solvates and hydrates thereof is crystalline, crystal-like, or amorphous.

A first aspect of the third embodiment is directed to crystalline *S_P-4*.

A second aspect of the third embodiment is directed to a monoclinic crystalline *S_P-4*, preferably having the following unit cell parameters *a* ~ 12.88 Å, *b* ~ 6.17 Å, *c* ~ 17.73 Å, and β ~ 92.05°.

A third aspect of the third embodiment is directed to a monoclinic crystalline *S_P-4*, preferably having the following unit cell parameters *a* ~ 20.09 Å, *b* ~ 6.10 Å, *c* ~ 23.01 Å, and β ~ 112.29°.

A fourth aspect of the third embodiment is directed to a monoclinic crystalline *S_P-4*, preferably having the following unit cell parameters *a* ~ 12.83 Å, *b* ~ 6.15 Å, *c* ~ 17.63 Å, and β ~ 91.75°.

A fifth aspect of the third embodiment is directed to a monoclinic crystalline *Sp-4*, preferably having the following unit cell parameters $a \sim 12.93 \text{ \AA}$, $b \sim 6.18 \text{ \AA}$, $c \sim 18.01 \text{ \AA}$, and $\beta \sim 96.40^\circ$.

5 A sixth aspect of the third embodiment is directed to a crystalline *Sp-4* having XRPD 2 θ -reflections ($^\circ$) at about: 5.2, 7.5, 9.6, 16.7, 18.3, 22.2.

A seventh aspect of the third embodiment is directed to a crystalline *Sp-4* having XRPD 2 θ -reflections ($^\circ$) at about: 5.0, 7.3, 9.4, and 18.1.

An eighth aspect of the third embodiment is directed to a crystalline *Sp-4* having XRPD 2 θ -reflections ($^\circ$) at about: 4.9, 6.9, 9.8, 19.8, 20.6, 24.7, and 26.1.

10 A ninth aspect of the third embodiment is directed to a crystalline *Sp-4* having XRPD 2 θ -reflections ($^\circ$) at about: 6.9, 9.8, 19.7, 20.6, and 24.6.

A ninth aspect of the third embodiment is directed to a crystalline *Sp-4* having XRPD 2 θ -reflections ($^\circ$) at about: 5.0, 6.8, 19.9, 20.6, 20.9, and 24.9.

15 A tenth aspect of the third embodiment is directed to a crystalline *Sp-4* having XRPD 2 θ -reflections ($^\circ$) at about: 5.2, 6.6, 7.1, 15.7, 19.1, and 25.0.

An eleventh aspect of the third embodiment is directed to crystalline *Sp-4* having XRPD 2 θ -reflections ($^\circ$) at about: 6.1, 8.2, 10.4, 12.7, 17.2, 17.7, 18.0, 18.8, 19.4, 19.8, 20.1, 20.8, 21.8, and 23.3.

20 A twelfth aspect of the third embodiment is directed to crystalline *Sp-4* having an XRPD diffraction pattern substantially as that shown in any one of Fig. 3, Fig. 4, Fig. 5, Fig. 6, Fig. 7, Fig. 8, and Fig 21.

A thirteenth aspect of the third embodiment is directed to *Sp-4* having the following FT-IR peaks (cm^{-1}) at about: 1743, 1713, 1688, 1454, 1378, 1208, and 1082.

25 A fourteenth aspect of the third embodiment is directed to *Sp-4* having an FT-IR spectrum substantially as that shown in Fig. 7.

A fifteenth aspect of the third embodiment is directed to substantially pure *Sp-4*.

30 A sixteenth aspect of the third embodiment is directed to substantially pure crystalline *Sp-4*.

A seventeenth aspect of the third embodiment is directed to substantially pure amorphous *Sp-4*.

Dosage, Administration, and Use

A fourth embodiment is directed to a composition for the treatment and/or prophylaxis of any of the viral agents using any of compounds **4**, *R_P-4*, or *S_P-4*.

Possible viral agents include, but are not limited to: hepatitis C virus, hepatitis B virus, Hepatitis A virus, West Nile virus, yellow fever virus, dengue virus,
5 rhinovirus, polio virus, bovine viral diarrhea virus, Japanese encephalitis virus, or those viruses belonging to the groups of pestiviruses, hepaciviruses, or flaviviruses.

An aspect of this embodiment is directed to a composition for the treatment of any of the viral agents disclosed herein said composition comprising a
10 pharmaceutically acceptable medium selected from among an excipient, carrier, diluent, and equivalent medium and any of compounds **4**, *R_P-4*, or *S_P-4*, that is intended to include its hydrates, solvates, and any crystalline forms of any of compounds **4**, *R_P-4*, or *S_P-4* or its hydrates and solvates thereof.

The compounds **4**, *R_P-4*, or *S_P-4* may be independently formulated in a wide
15 variety of oral administration dosage forms and carriers. Oral administration can be in the form of tablets, coated tablets, hard and soft gelatin capsules, solutions, emulsions, syrups, or suspensions. The compounds **4**, *R_P-4*, or *S_P-4* are efficacious when administered by suppository administration, among other routes of administration. The most convenient manner of administration is generally oral
20 using a convenient daily dosing regimen which can be adjusted according to the severity of the disease and the patient's response to the antiviral medication.

The compounds **4**, *R_P-4*, or *S_P-4* together with one or more conventional excipients, carriers, or diluents, may be placed into the form of pharmaceutical compositions and unit dosages. The pharmaceutical compositions and unit dosage
25 forms may be comprised of conventional ingredients in conventional proportions, with or without additional active compounds and the unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed. The pharmaceutical compositions may be employed as solids, such as tablets or filled capsules, semisolids, powders,
30 sustained release formulations, or liquids such as suspensions, emulsions, or filled capsules for oral use; or in the form of suppositories for rectal or vaginal administration. A typical preparation will contain from about 5% to about 95% active compound or compounds (w/w).

The compounds 4, *R*_P-4, or *S*_P-4 can be administered alone but will generally be administered in admixture with one or more suitable pharmaceutical excipients, diluents or carriers selected with regard to the intended route of administration and standard pharmaceutical practice.

5 Solid form preparations include, for example, powders, tablets, pills, capsules, suppositories, and dispersible granules. A solid carrier may be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material. In powders, the carrier generally is a finely divided solid
10 which is a mixture with the finely divided active component. In tablets, the active component generally is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted in the shape and size desired. Suitable carriers include but are not limited to magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose,
15 sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. Solid form preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like. Examples of solid formulations are exemplified in EP 0524579; US 2002/0142050; US 2004/0224917; US 2005/0048116; US
20 2005/0058710; US 2006/0034937; US 2006/0057196; US 2006/0188570; US 2007/0026073; US 2007/0059360; US 2007/0077295; US 2007/0099902; US 2008/0014228; US 6,267,985; US 6,294,192; US 6,383,471; US 6,395,300; US 6,569,463; US 6,635,278; US 6,645,528; US 6,923,988; US 6,932,983; US 7,060,294; and US 7,462,608, each of which is incorporated by reference.

25 Liquid formulations also are suitable for oral administration include liquid formulation including emulsions, syrups, elixirs and aqueous suspensions. These include solid form preparations which are intended to be converted to liquid form preparations shortly before use. Examples of liquid formulation are exemplified in U.S. Patent Nos. 3,994,974; 5,695,784; and 6,977,257. Emulsions may be prepared
30 in solutions, for example, in aqueous propylene glycol solutions or may contain emulsifying agents such as lecithin, sorbitan monooleate, or acacia. Aqueous suspensions can be prepared by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins,

methycellulose, sodium carboxymethylcellulose, and other well known suspending agents.

The compounds **4**, *R_P-4*, or *S_P-4* may be independently formulated for administration as suppositories. A low melting wax, such as a mixture of fatty acid glycerides or cocoa butter is first melted and the active component is dispersed homogeneously, for example, by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and to solidify.

The compounds **4**, *R_P-4*, or *S_P-4* may be independently formulated for vaginal administration. Pessaries, tampons, creams, gels, pastes, foams or sprays containing in addition to the active ingredient such carriers as are known in the art to be appropriate. Certain of these formulations may also be used in conjunction with a condom with or without a spermicidal agent.

Suitable formulations along with pharmaceutical carriers, diluents and excipients are described in *Remington: The Science and Practice of Pharmacy 1995*, edited by E. W. Martin, Mack Publishing Company, 19th edition, Easton, Pennsylvania, which is hereby incorporated by reference. A skilled formulation scientist may modify the formulations within the teachings of the specification to provide numerous formulations for a particular route of administration without rendering compositions containing the compounds contemplated herein unstable or compromising their therapeutic activity.

Additionally, the purified compounds **4**, *R_P-4*, or *S_P-4* may be independently formulated in conjunction with liposomes or micelles. As to liposomes, it is contemplated that the purified compounds can be formulated in a manner as disclosed in U.S. Patent Nos. 4,797,285; 5,013,556; 5,077,056; 5,077,057; 5,154,930; 5,192,549; 5,213,804; 5,225,212; 5,277,914; 5,316,771; 5,376,380; 5,549,910; 5,567,434; 5,736,155; 5,827,533; 5,882,679; 5,891,468; 6,060,080; 6,132,763; 6,143,321; 6,180,134; 6,200,598; 6,214,375; 6,224,903; 6,296,870; 6,653,455; 6,680,068; 6,726,925; 7,060,689; and 7,070,801, each of which is incorporated by reference. As to micelles, it is contemplated that the purified compounds can be formulated in a manner as disclosed in U.S. Patent Nos. 5,145,684 and 5,091,188, both of which are incorporated by reference.

The fifth embodiment is directed to a use of any of compounds **4**, *R_P-4*, or *S_P-4* in the manufacture of a medicament for the treatment of any condition the result of an infection by any one of the following viral agents: hepatitis C virus,

West Nile virus, yellow fever virus, dengue virus, rhinovirus, polio virus, hepatitis A virus, bovine viral diarrhea virus and Japanese encephalitis virus.

5 The term "medicament" means a substance used in a method of treatment and/or prophylaxis of a subject in need thereof, wherein the substance includes, but is not limited to, a composition, a formulation, a dosage form, and the like, comprising any of compounds **4**, *R_P-4*, or *S_P-4*. It is contemplated that the use of any of compounds **4**, *R_P-4*, or *S_P-4* in the manufacture of a medicament, for the treatment of any of the antiviral conditions disclosed herein, either alone or in combination with another compound disclosed herein. A medicament includes, but is not limited to, any one of the compositions contemplated by the fourth embodiment disclosed herein.

A sixth embodiment is directed to a method of treatment and/or prophylaxis in a subject in need thereof said method comprises administering a therapeutically effective amount of any of compounds **4**, *R_P-4*, or *S_P-4* to the subject.

15 It is intended that a subject in need thereof is one that has any condition the result of an infection by any of the viral agents disclosed herein, which includes, but is not limited to, hepatitis C virus, West Nile virus, yellow fever virus, dengue virus, rhinovirus, polio virus, hepatitis A virus, bovine viral diarrhea virus or Japanese encephalitis virus, flaviviruses or pestiviruses or hepaciviruses or a viral agent causing symptoms equivalent or comparable to any of the above-listed viruses.

20 The term "subject" means a mammal, which includes, but is not limited to, cattle, pigs, sheep, chicken, turkey, buffalo, llama, ostrich, dogs, cats, and humans, preferably the subject is a human. It is contemplated that in the method of treating a subject thereof of the ninth embodiment can be any of the compounds contemplated herein, either alone or in combination with another compound disclosed herein.

25 The term "therapeutically effective amount" as used herein means an amount required to reduce symptoms of the disease in an individual. The dose will be adjusted to the individual requirements in each particular case. That dosage can vary within wide limits depending upon numerous factors such as the severity of the disease to be treated, the age and general health condition of the patient, other medicaments with which the patient is being treated, the route and form of administration and the preferences and experience of the medical practitioner involved. For oral administration, a daily dosage of between about 0.001 and about 10 g, including all values in between, such as 0.001, 0.0025, 0.005, 0.0075, 0.01,

0.025, 0.050, 0.075, 0.1, 0.125, 0.150, 0.175, 0.2, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, and 9.5, per day should be appropriate in monotherapy and/or in combination therapy. A particular daily dosage is between about 0.01 and about 1 g per day, including all incremental values of 0.01 g (i.e., 10 mg) in between, a preferred daily dosage about 0.01 and about 0.8 g per day, more preferably about 0.01 and about 0.6 g per day, and most preferably about 0.01 and about 0.25 g per day, each of which including all incremental values of 0.01 g in between. Generally, treatment is initiated with a large initial "loading dose" to rapidly reduce or eliminate the virus following by a decreasing the dose to a level sufficient to prevent resurgence of the infection. One of ordinary skill in treating diseases described herein will be able, without undue experimentation and in reliance on knowledge, experience and the disclosures of this application, to ascertain a therapeutically effective amount of the compound disclosed herein for a given disease and patient.

Therapeutic efficacy can be ascertained from tests of liver function including, but not limited to protein levels such as serum proteins (e.g., albumin, clotting factors, alkaline phosphatase, aminotransferases (e.g., alanine transaminase, aspartate transaminase), 5'-nucleosidase, γ -glutamyltranspeptidase, etc.), synthesis of bilirubin, synthesis of cholesterol, and synthesis of bile acids; a liver metabolic function, including, but not limited to, carbohydrate metabolism, amino acid and ammonia metabolism. Alternatively the therapeutic effectiveness may be monitored by measuring HCV-RNA. The results of these tests will allow the dose to be optimized.

A first aspect of the sixth embodiment is directed to a method of treatment and/or prophylaxis in a subject in need thereof said method comprises administering to the subject a therapeutically effective amount of a compound represented by any of compounds 4, *R_p*-4, or *S_p*-4 and a therapeutically effective amount of another antiviral agent; wherein the administration is concurrent or alternative. It is understood that the time between alternative administration can range between 1-24 hours, which includes any sub-range in between including, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, and 23 hours.

Examples of "another antiviral agent" include, but are not limited to: HCV NS3 protease inhibitors (see EP 1881001, US 2003187018, US 2005267018, WO 2003006490, WO 200364456, WO 2004094452, WO 2005028502, WO

2005037214, WO 2005095403, WO 2007014920, WO 2007014921, WO 2007014922, WO 2007014925, WO 2007014926, WO 2007015824, WO 2008010921, and WO 2008010921); HCV NS5B Inhibitors (see US 2004229840, US 2005154056, US 2005-98125, US 20060194749, US 20060241064, US 20060293306, US 2006040890, US 2006040927, US 2006166964, US 2007275947, US 6784166, US20072759300, WO 2002057287, WO 2002057425, WO 2003010141, WO 2003037895, WO 2003105770, WO 2004000858, WO 2004002940, WO 2004002944, WO 2004002977, WO 2004003138, WO 2004041201, WO 2004065367, WO 2004096210, WO 2005021568, WO 2005103045, WO 2005123087, WO 2006012078, WO 2006020082, WO 2006065335, WO 2006065590, WO 2006093801, WO 200702602, WO 2007039142, WO 2007039145, WO 2007076034, WO 2007088148, WO 2007092000, and WO2007095269); HCV NS4 Inhibitors (see WO 2005067900 and WO 2007070556); HCV NS5a Inhibitors (see US 2006276511, WO 2006035061, WO 2006100310, WO 2006120251, and WO 2006120252); Toll-like receptor agonists (see WO 2007093901); and other inhibitors (see WO 2000006529, WO 2003101993, WO 2004009020, WO 2004014313, WO 2004014852, and WO 2004035571); and compounds disclosed in U.S. Patent Application No. 12/053,015, filed March 21, 2008 (US 2010/0016251) (the contents of which are incorporated by reference), interferon- α , interferon- β , pegylated interferon- α , ribavirin, levovirin, viramidine, another nucleoside HCV polymerase inhibitor, a HCV non-nucleoside polymerase inhibitor, a HCV protease inhibitor, a HCV helicase inhibitor or a HCV fusion inhibitor.


When any of compounds 4, R_p -4, or S_p -4 are administered in combination with another antiviral agent the activity may be increased over the parent compound. When the treatment is combination therapy, such administration may be concurrent or sequential with respect to that of the nucleoside derivatives. "Concurrent administration" as used herein thus includes administration of the agents at the same time or at different times. Administration of two or more agents at the same time can be achieved by a single formulation containing two or more active ingredients or by substantially simultaneous administration of two or more dosage forms with a single active agent.

It will be understood that references herein to treatment extend to prophylaxis as well as to the treatment of existing conditions. Furthermore, the term

"treatment" of a HCV infection, as used herein, also includes treatment or prophylaxis of a disease or a condition associated with or mediated by HCV infection, or the clinical symptoms thereof.

Preparation

- 5 A seventh embodiment is directed to a process for preparing any one of compounds **4**, R_P -**4**, or S_P -**4**, which comprises: a) reacting an isopropyl-alanate, **A**, a di-LG-phenylphosphate, **B**, 2'-deoxy-2'-fluoro-2'-C-methyluridine, **3**, and a base to obtain a first mixture comprising at least one of S_P -**4** and R_P -**4**



- 10 wherein X is a conjugate base of an acid, n is 0 or 1, and LG is a leaving group; b) reacting the first mixture with a protecting compound to obtain a second mixture comprising at least one of protected S_P -**4** and protected R_P -**4**; and c) optionally subjecting the second mixture to crystallization, chromatography, or extraction in order to obtain **4**, S_P -**4**, or R_P -**4**.


- 15 In a first aspect of the seventh embodiment, the isopropyl alanate is present as its hydrochloric acid salt, which is preferably, substantially anhydrous.

 In a second aspect of the seventh embodiment, the base is N-methylimidazole.

- 20 In a third aspect of the seventh embodiment, the mole ratio of **A**-to-**B**-to-**3** is about 1.6-to-1.3-to-1.

 In a fourth aspect of the seventh embodiment, the protecting compound is t-butyldimethylsilyl chloride.

- 25 An eighth embodiment is directed to a process for preparing S_P -**4** or R_P -**4**, which comprises: a) reacting an isopropyl-alanate, **A**, a di-LG-phenylphosphate, **B**, 2'-deoxy-2'-fluoro-2'-C-methyluridine, **3**, and a base to obtain a first mixture comprising at least one of S_P -**4** and R_P -**4**



wherein X is a conjugate base of an acid, n is 0 or 1, and LG is a leaving group; and
 b) optionally subjecting the second mixture to crystallization, chromatography, or extraction in order to obtain purified S_{P-4} or R_{P-4} .

5 A first aspect of the eighth embodiment for preparing R_{P-4} additionally includes further purifying the second mixture or the purified R_{P-4} by dissolving or suspending the second mixture or the purified R_{P-4} mixture in a solvent; optionally followed by seeding with crystalline R_{P-4} ; and adding sufficient anti-solvent to obtain crystalline R_{P-4} .


10 A second aspect of the eighth embodiment for preparing S_{P-4} additionally includes further purifying the second mixture or the purified S_{P-4} by d) dissolving or suspending the second mixture or the purified S_{P-4} in a solvent followed by seeding with crystalline S_{P-4} at about room temperature; collecting a first solid the majority of which comprises S_{P-4} ; dissolving the first solid in a solvent at its reflux
 15 temperature; and cooling or adding an anti-solvent to obtain a second solid.

 A third aspect of the eighth embodiment for the preparation of S_{P-4} , additionally includes further purifying S_{P-4} by d) dissolving or suspending the second mixture or the purified S_{P-4} mixture in a first solvent followed by adding an anti-solvent so as to obtain a first composition in which the residual solvent/anti-
 20 solvent is removed by decanting to obtain a residue; treating the residue with a solution containing the first solvent and anti-solvent to yield a second composition whereby upon reducing the pressure affords a first solid; dissolving or suspending the first solid using a second solvent so as to obtain a third composition; adding seed crystals of S_{P-4} to the third composition; collecting a second solid; dissolving or
 25 suspending the second solid in a third solvent, optionally heated to the reflux temperature of the third solvent to obtain a fourth composition, and, if necessary, cooling the fourth composition to obtain a third solid comprising S_{P-4} which is collected by filtration.


In a fourth aspect of the eighth embodiment for the preparation of *S_P-4*, *S_P-4* is further purified by the second mixture or the purified *S_P-4* by d) adding silica gel to the second mixture or the purified *S_P-4* followed by solvent evaporation to afford a dry slurry; stirring the dry slurry in a first solvent/anti-solvent combination to
5 obtain a first wet slurry; decanting the first solvent/anti-solvent combination from the first wet slurry to obtain a second wet slurry and a first composition; adding to the second wet slurry a second solvent/anti-solvent combination followed by stirring; decanting the second solvent/anti-solvent combination from the second wet slurry to obtain a third wet slurry and a second composition; optionally repeating
10 steps g)–h) on the third wet slurry or additional wet slurries; evaporating the solvent from the second composition, and optionally any additional composition obtained from optional step i) to obtain a first solid; dissolving or suspending the first solid in a solution containing a third solvent and optionally a fourth solvent to obtain a third composition; optionally adding seed crystals of *S_P-4* to the third composition;
15 obtaining from the third composition a second solid comprising *S_P-4*; and optionally recrystallizing the second solid using a third solvent to obtain a third solid comprising *S_P-4*.

One of ordinary skill will appreciate that the compounds can be separated by traditional extraction, traditional crystallization or traditional chromatographic
20 techniques. Traditional chromatographic techniques include, but are not limited to, chromatography on silica gel (using, e.g., 3-5% methanol in DCM or 4-6% isopropanol in DCM) to produce enhanced levels of one isomer (50-100%) and then crystallize it. Alternatively, one could use reversed phase chromatography (using, e.g., 1-30% acetonitrile-aqueous mobile phase). Furthermore the compounds can be
25 isolated by supercritical fluid chromatography SFC with carbon dioxide as the main solvent and alcohols such as methanol as a modifier, preferably using the appropriate chiral media, such as, Daicel Chiralpack IA. Alternatively, SMB chromatography may be employed using the appropriate chiral media, such as, Daicel ChiralPack IA, using a mixture of solvents such as hexanes/isopropanol or
30 single solvents such as ethyl acetate.

A ninth embodiment is directed to a process for preparing *S_P-4*, which comprises: a) reacting an isopropyl-alanyl-phosphoramidate with a 3'-O-protected or unprotected **3**, and a basic reagent to obtain a composition comprising protected or unprotected *S_P-4*



wherein the isopropyl-alanyl-phosphoramidate is comprised of a mixture of diastereomers represented by the following structures:



5 wherein the ratio of **C:C'** is about 1:1.


In a first aspect, the basic reagent is t-butylmagnesium chloride and the ratio of **C:C'** is greater than or equal to about 1:1.

In a second aspect, the basic reagent is t-butylmagnesium chloride and the ratio of **C:C'**; is greater than about 1:1.


10 In a third aspect, the basic reagent is t-butylmagnesium chloride and the ratio of **C:C'** is at least about 1.5:1, about 2.3:1, about 4:1, about 5.7:1, about 9:1, about 19:1, about 32.3:1, about 49:1, or about 99:1.

In a fourth aspect the LG' is selected from among, 2,4-dinitrophenoxide, 4-nitrophenoxide, 2-nitrophenoxide, 2-chloro-4-nitrophenoxide, 2,4-dichlorophenoxide, and pentafluorophenoxide, the basic reagent is t-butylmagnesium chloride, and the ratio of **C:C'** is at least about 1.5:1, about 2.3:1, about 4:1, about 5.7:1, about 9:1, about 19:1, about 32.3:1, about 49:1, or about 99:1.

20 A fifth aspect for preparing **S_P-4**, comprises: a) reacting an isopropyl-alanyl-phosphoramidate (**C**) with a 3'-O-protected or unprotected **3**, and a basic reagent to obtain a composition comprising protected or unprotected **S_P-4**




- wherein Z is a protecting group or hydrogen; LG' is a leaving group; and b)
- optionally subjecting the obtained protected or unprotected S_P -**4** to chromatography,
- 5 extraction, or crystallization in order to obtain purified protected or unprotected S_P -**4**. In a sub-embodiment, LG' is tosylate, camphorsulfonate, or an aryloxy substituted with at least one electron withdrawing group; more preferably, LG' is selected from among 2,4-dinitrophenoxide, 4-nitrophenoxide, 2-nitrophenoxide, 2-chloro-4-nitrophenoxide, 2,4-dichlorophenoxide, or pentafluorophenoxide. In a
- 10 further sub-embodiment, when S_P -**4** is protected, i.e., Z is not hydrogen, the process of the ninth embodiment is further directed to deprotecting protected S_P -**4**. In a further sub-embodiment, the reaction is conducted in a polar aprotic solvent, such as, tetrahydrofuran or another ethereal solvent either one being alone or in combination with each other or with a C_2 to C_7 nitrile, such as acetonitrile.
- 15 The process of the ninth embodiment further comprises 1) reacting $(LG')P(O)(LG)_2$, wherein LG, independent of LG', is a leaving group, with (i) isopropyl-alanate and a first base to obtain $(LG')P(O)(LG)(NHAla-^iPr)$ followed by reacting $(LG')P(O)(LG)(NHAla-^iPr)$ with phenol and a second base to obtain a mixture comprising **C** and **C'**, (ii) phenol and a first base to obtain
- 20 $(LG')P(O)(LG)(OPh)$ followed by reacting $(LG')P(O)(LG)(OPh)$ with isopropyl-alanate and a second base to obtain a mixture comprising **C** and **C'**, or (iii) combining isopropyl-alanate, phenol, and at least one base to obtain a mixture comprising **C** and **C'**; or 2) reacting $(PhO)P(O)(LG)_2$, wherein LG is a leaving group, with (i) isopropyl-alanate and a first base to obtain $(PhO)P(O)(LG)(NHAla-^iPr)$ followed by reacting $(PhO)P(O)(LG)(NHAla-^iPr)$ with a leaving group
- 25 precursor (LG'H) and a second base to obtain a mixture comprising **C** and **C'**,




and subjecting the mixture to chromatography or crystallizing the mixture to obtain **C**. In an aspect of the ninth embodiment, the isopropyl alanate is present as its hydrochloric acid salt, which is preferably, substantially anhydrous.

- 5 A tenth embodiment is directed to a process for preparing *R_p*-**4**, which comprises: a) reacting an isopropyl-alanyl-phosphoramidate with a 3'-O-protected or unprotected **3**, and a basic reagent to obtain a composition comprising protected or unprotected *R_p*-**4**



- 10 wherein the isopropyl-alanyl-phosphoramidate is comprised of a mixture of diastereomers represented by the following structures:



wherein the ratio of **C'**:**C** is about 1:1.


- 15 In a first aspect, the basic reagent is t-butylmagnesium chloride and the ratio of **C'**:**C** is greater than or equal to about 1:1.

In a second aspect, the basic reagent is t-butylmagnesium chloride and the ratio of **C'**:**C**; is greater than about 1:1.

- 20 In a third aspect, the basic reagent is t-butylmagnesium chloride and the ratio of **C'**:**C** is at least about 1.5:1, about 2.3:1, about 4:1, about 5.7:1, about 9:1, about 19:1, about 32.3:1, about 49:1, or about 99:1.

A fourth aspect the LG' is p-nitrophenoxide, the basic reagent is t-butylmagnesium chloride, and the ratio of **C'**:**C** is at least about 1.5:1, about 2.3:1, about 4:1, about 5.7:1, about 9:1, about 19:1, about 32.3:1, about 49:1, or about 99:1.

A fifth aspect for preparing R_P -4, comprises: a) reacting an isopropyl-alanyl-phosphoramidate (C') with a 3'-O-protected or unprotected **3**, and a basic reagent to obtain a composition comprising protected or unprotected R_P -4



5

wherein Z is a protecting group or hydrogen; LG' is a leaving group; and b) optionally subjecting the obtained protected or unprotected R_P -4 to chromatography, extraction, or crystallization in order to obtain purified protected or unprotected R_P -4. In a sub-embodiment, LG' is tosylate, camphorsulfonate, or an aryloxy substituted with at least one electron withdrawing group; more preferably, LG' is selected from among p-nitrophenoxide, 2,4-dinitrophenoxide, and pentafluorophenoxide. In a further sub-embodiment, when R_P -4 is protected, i.e., Z is not hydrogen, the process of the ninth embodiment is further directed to deprotecting protected R_P -4. In a further sub-embodiment, the reaction is conducted

10 substituted with at least one electron withdrawing group; more preferably, LG' is selected from among p-nitrophenoxide, 2,4-dinitrophenoxide, and pentafluorophenoxide. In a further sub-embodiment, when R_P -4 is protected, i.e., Z is not hydrogen, the process of the ninth embodiment is further directed to deprotecting protected R_P -4. In a further sub-embodiment, the reaction is conducted


15 in a polar aprotic solvent, such as, tetrahydrofuran or another ethereal solvent either one being alone or in combination with each other or with a C_2 to C_7 nitrile, such as acetonitrile.

The process of the tenth embodiment further comprises 1) reacting $(LG')P(O)(LG)_2$, wherein LG, independent of LG' , is a leaving group, with (i)

20 isopropyl-alanate and a first base to obtain $(LG')P(O)(LG)(NHAla-iPr)$ followed by reacting $(LG')P(O)(LG)(NHAla-iPr)$ with phenol and a second base to obtain a mixture comprising **C** and C' , (ii) phenol and a first base to obtain $(LG')P(O)(LG)(OPh)$ followed by reacting $(LG')P(O)(LG)(OPh)$ with isopropyl-alanate and a second base to obtain a mixture comprising **C** and C' , or (iii)

25 combining isopropyl-alanate, phenol, and at least one base to obtain a mixture comprising **C** and C' ; or 2) reacting $(PhO)P(O)(LG)_2$, wherein LG' , independent of LG, is a leaving group, with (i) isopropyl-alanate and a first base to obtain $(PhO)P(O)(LG)(NHAla-iPr)$ followed by reacting $(PhO)P(O)(LG)(NHAla-iPr)$ with


a leaving group precursor and a second base to obtain a mixture comprising **C** and **C'**,



and subjecting the mixture to chromatography or crystallizing the mixture to obtain **C'**. In an aspect of the ninth embodiment, the isopropyl alanate is present as its hydrochloric acid salt, which is preferably, substantially anhydrous.


An eleventh embodiment is directed to a composition obtained by the processes recited in the seventh embodiment, the eighth embodiment, the ninth embodiment or the tenth embodiment as well as their respective aspects. An aspect of the eleventh embodiment is directed to a composition obtained by any one of the exemplified embodiments disclosed below. The so obtained composition can be crystalline, crystal-like, amorphous, or a combination thereof.

A twelfth embodiment is directed to a compound **3**



wherein Z is a protecting group or hydrogen; which is useful for the preparation of R_P -4 or S_P -4.

A first aspect of the twelfth embodiment is selected from among a compound having the following structure




3a: Z = -C(O)CH₂CH₂C(O)CH₃

3b: Z = -C(O)OCH₂Ph

3c: Z = -Si(Me)₂^tBu

3d: Z = -Si(ⁱPr)₂OSi(ⁱPr)₂OH

A thirteenth embodiment is directed to a compound, its salt, hydrate, solvate, or combination thereof, represented by the following structures



where LG' is a leaving group, which is useful for the preparation of *R_P*-4 or *S_P*-4.

In a first aspect of the thirteenth embodiment, LG' is tosylate, camphorsulfonate, an aryloxy, or an aryloxy substituted with at least one electron withdrawing group.

In a second aspect of the thirteenth embodiment, LG' is selected from among 2,4-dinitrophenoxide, 4-nitrophenoxide, 2-nitrophenoxide, 2-chloro-4-nitrophenoxide, 2,4-dichlorophenoxide, or pentafluorophenoxide.

In a third aspect of the thirteenth embodiment, LG' is pentafluorophenoxide or 4-nitro-phenoxide.

A fourth aspect of the thirteenth embodiment is directed to compound C, wherein LG' is 2,4-dinitrophenoxide, 4-nitrophenoxide, 2-nitrophenoxide, 2-chloro-4-nitrophenoxide, 2,4-dichlorophenoxide, or pentafluorophenoxide.

A fifth aspect of the thirteenth embodiment is directed to compound C, wherein LG' is 4-nitrophenoxide or pentafluorophenoxide.


A sixth aspect of the thirteenth embodiment is directed to compound C, wherein LG' is 4-nitrophenoxide.

A seventh aspect of the thirteenth embodiment is directed to crystalline compound C, wherein LG' is 4-nitrophenoxide.

An eighth aspect of the thirteenth embodiment is directed to compound C, wherein LG' is pentafluorophenoxide.

A ninth aspect of the thirteenth embodiment is directed to crystalline compound C, wherein LG' is pentafluorophenoxide.

A fourteenth embodiment is directed to a process for preparing a compound represented by the structural formula



crystallizing the compound from a composition, comprising

- a) a first composition;
- b) a second leaving group precursor;
- c) a non-nucleophilic base; and
- d) a liquid composition;

5 wherein the first composition comprises the compound and its corresponding P-based diastereomer.

 In a first aspect of the fourteenth embodiment, the mole amount of the compound and the mole amount of its P-based diastereomer are the same or different.

10 In a second aspect of the fourteenth embodiment, the mole amount of the compound is greater than the mole amount of its corresponding P-based diastereomer or vice versa.

 In a third aspect of the fourteenth embodiment, the second leaving group precursor is 2,4-dinitrophenol, 4-nitrophenol, 2-nitrophenol, 2-chloro-4-nitrophenol,
15 2,4-dichlorophenol, or pentafluorophenol.

 In a fourth aspect of the fourteenth embodiment, LG' is pentafluorophenoxide. In a first sub-aspect, the second leaving group precursor is pentafluorophenol. In a second sub-aspect, the amount of pentafluorophenol ranges from about 0.01 mole equivalents to about 10 mole equivalents relative to the mole
20 amount of the compound and its P-based diastereomer and all mole equivalents in between. In a third sub-aspect, the amount of pentafluorophenol ranges from about 0.1 mole equivalents to about 1 mole equivalents relative to the mole amount of the compound and its P-based diastereomer and all mole equivalents in between.

 In a fifth aspect of the fourteenth embodiment, the crystallizing occurs at a
25 temperature that ranges from about -10°C to about +40°C and all temperature values in between. In a first sub-aspect, the crystallizing occurs at about room temperature.

 In a sixth aspect of the fourteenth embodiment, the non-nucleophilic base is selected from among potassium carbonate, cesium carbonate, di-isopropylamine, di-isopropylethylamine, triethylamine, quinuclidine, naphthalene-1,8-diamine, 2,2,6,6-tetramethylpiperidine, 1,8-diazabicycloundec-7-ene, 4-dimethylamino-pyridine,
30 pyridine, a 2,6-di-C₁₋₆-alkyl-pyridine, a 2,4,6-tri-C₁₋₆-alkyl-pyridine, and mixtures thereof. In a first sub-aspect, the non-nucleophilic base is triethylamine or 1,8-diazabicycloundec-7-ene. In a second sub-aspect, the non-nucleophilic base is triethylamine.

In a seventh aspect of the fourteenth embodiment, the non-nucleophilic base is present in an amount that ranges from about 0.01 equivalents mol to about 10 mol equivalents, and all mole equivalents in between, relative to the total mole amount of the compound and its P-based diastereomer. In a first sub-aspect, the non-

5 nucleophilic base is present in an amount that ranges from about 0.1 mol equivalents to about 1 mol equivalents, and all mole equivalents in between, relative to the total mole amount of the compound and its P-based diastereomer .

In an eighth aspect of the fourteenth embodiment, the solubility of the compound is less than the solubility of its corresponding P-based diastereomer in the

10 liquid composition or vice versa.

In a ninth aspect of the fourteenth embodiment, the liquid composition comprises at least one of a solvent and an anti-solvent. In a first sub-aspect, the liquid composition comprises at least one of a C₁ to C₈ alcohol, a C₂ to C₈ ether, a C₃ to C₇ ketone, a C₃ to C₇ ester, a C₁ to C₂ chlorocarbon, a C₂ to C₇ nitrile, a C₅ to C₁₂

15 saturated hydrocarbon, and a C₆ to C₁₂ aromatic hydrocarbon. In a second sub-aspect, the liquid composition comprises at least one of a C₂ to C₈ ether, a C₃ to C₇ ester, a C₅ to C₁₂ saturated hydrocarbon, and a C₆ to C₁₂ aromatic hydrocarbon. In a third sub-aspect, the liquid composition comprises at least one of a C₂ to C₈ ether, a C₃ to C₇ ester, and a C₅ to C₁₂ saturated hydrocarbon. In a fourth sub-aspect, the

20 liquid composition comprises at least one of ethyl acetate, t-butyl-methylether, and hexane. In a fifth sub-aspect, the liquid composition comprises ethyl acetate and hexane. In a sixth sub-aspect, the liquid composition comprises t-butyl-methylether and hexane.

In a tenth aspect of the fourteenth embodiment, the amount of liquid

25 composition ranges from about 1 mL to about 10 mL for every gram of the first composition and all mL/g values in between.

An eleventh aspect of the fourteenth embodiment further comprises adding crystalline compound to the composition. A first sub-aspect further comprises adding about 0.1 to about 1 wt. %, and all wt. % values in between, of crystalline

30 compound to the first composition.

A twelfth aspect of the fourteenth embodiment further comprises

a) reacting PhOP(O)(LG)_2 and $^i\text{Pr-Ala-NH}_2 \cdot \text{HCl}$ in the presence of a first base to obtain $(\text{PhO})\text{P(O)(LG)(NHAla-}^i\text{Pr)}$;

b) reacting (PhO)P(O)(LG)(NHAla-ⁱPr) with a first leaving group precursor (LG'H) in the presence of a second base to obtain the composition comprising the compound and its P-based diastereomer;


wherein LG and LG', independent of each other, are leaving groups;

5 wherein the first leaving group precursor and the second leaving group precursor are the same or different; and

wherein the first base and the second base are the same or different.

A fifteenth embodiment is directed to a process for preparing crystalline (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate

10 having the following structure,



which comprises:

crystallizing (S)-isopropyl 2-(((S)-

(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate from a second

15 composition comprising

- a) a first composition;
- b) pentafluorophenol;
- c) a non-nucleophilic base; and
- d) a liquid composition;

20 wherein the first composition comprises (S)-isopropyl 2-(((S)-
(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate and (S)-
isopropyl 2-(((R)-
(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate.

In a first aspect of the fifteenth embodiment, the mole amount of the (S)-
25 isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate and
the mole amount of the (S)-isopropyl 2-(((R)-
(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate are the same or
different.

In a second aspect of the fifteenth embodiment, the mole amount of the (S)-
30 isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate is

greater than the mole amount of the (S)-isopropyl 2-(((R)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate.

In a third aspect of the fifteenth embodiment, the amount of pentafluorophenol ranges from about 0.01 mole equivalents to about 10 mole equivalents (and all mole equivalent values in between) relative to the mole amount of (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate and (S)-isopropyl 2-(((R)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate. In a first sub-aspect, the amount of pentafluorophenol ranges from about 0.1 mole equivalents to about 1 mole equivalents (and all mole equivalent values in between) relative to the mole amount of (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate and (S)-isopropyl 2-(((R)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate.

In a fourth aspect of the fifteenth embodiment, the crystallizing occurs at a temperature that ranges from about -10°C to about +40°C and all temperature values in between. In a first sub-aspect, the crystallizing occurs at about room temperature.

In a fifth aspect of the fifteenth embodiment, the non-nucleophilic base is selected from among potassium carbonate, cesium carbonate, di-isopropylamine, di-isopropylethylamine, triethylamine, quinuclidine, naphthalene-1,8-diamine, 2,2,6,6-tetramethylpiperidine, 1,8-diazabicycloundec-7-ene, 4-dimethylamino-pyridine, pyridine, a 2,6-di-C₁₋₆-alkyl-pyridine, a 2,4,6-tri-C₁₋₆-alkyl-pyridine, and mixtures thereof. In a first sub-aspect, the non-nucleophilic base is triethylamine or 1,8-diazabicycloundec-7-ene. In a second sub-aspect, the non-nucleophilic base is triethylamine.

In a sixth aspect of the fifteenth embodiment, the non-nucleophilic base is present in an amount that ranges from about 0.1 to about 1 mol equivalents (and all mole equivalent values in between) relative to the total mole amount of (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate and (S)-isopropyl 2-(((R)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate.

In a seventh aspect of the fifteenth embodiment, the solubility of (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate is less than the solubility of (S)-isopropyl 2-(((R)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate in the liquid composition.

In an eighth aspect of the fifteenth embodiment, the liquid composition comprises at least one of a solvent and an anti-solvent. In a first sub-aspect, the liquid composition comprises at least one of a C₁ to C₈ alcohol, a C₂ to C₈ ether, a C₃ to C₇ ketone, a C₃ to C₇ ester, a C₁ to C₂ chlorocarbon, a C₂ to C₇ nitrile, a C₅ to C₁₂ saturated hydrocarbon, and a C₆ to C₁₂ aromatic hydrocarbon. In a second sub-aspect, the liquid composition comprises at least one of a C₂ to C₈ ether, a C₃ to C₇ ester, a C₅ to C₁₂ saturated hydrocarbon, and a C₆ to C₁₂ aromatic hydrocarbon. In a third sub-aspect, the liquid composition comprises at least one of a C₂ to C₈ ether, a C₃ to C₇ ester, and a C₅ to C₁₂ saturated hydrocarbon. In a fourth sub-aspect, the liquid composition comprises at least one of ethyl acetate, t-butyl-methylether, and hexane. In a fifth sub-aspect, the liquid composition comprises ethyl acetate and hexane. In a sixth sub-aspect, the liquid composition comprises t-butyl-methylether and hexane.

In a ninth aspect of the fifteenth embodiment, the amount of liquid composition ranges from about 1 to about 10 mL for every gram (and all mL/g values in between) of the first composition.

A tenth aspect of the fifteenth embodiment further comprises adding crystalline (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate to the second composition.

An eleventh aspect of the fifteenth embodiment further comprises adding about 0.1 to about 1 wt. % (and all wt.% values in between) of crystalline (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate based on the total weight of (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate in the first composition.

A sixteenth embodiment is directed to crystalline (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate obtained by the process of the fifteenth embodiment.

A seventeenth embodiment is directed to a process for preparing (S)-isopropyl 2-(((S)-((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate, which comprises:

crystallizing (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate from a second composition comprising

- a) a first composition;
- 5 b) pentafluorophenol;
- c) a non-nucleophilic base; and
- d) a liquid composition;

wherein the first composition comprises (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate and (S)-
 10 isopropyl 2-(((R)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate.

A first aspect of the seventeenth embodiment is directed to a process for preparing (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate, which comprises:

15 contacting (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate with a product obtained by reacting a t-butylmagnesium halide with 1-((2R,3R,4R,5R)-3-fluoro-4-hydroxy-5-(hydroxymethyl)-3-methyltetrahydrofuran-2-yl)pyrimidine-2,4(1H,3H)-dione with
 20 a t-butylmagnesium halide.

In a second aspect of the seventeenth embodiment, the contacting occurs in a medium having a temperature that ranges from about 0°C to about 40°C and all temperature values in between.

In a third aspect of the seventeenth embodiment, the contacting occurs in a
 25 medium having a temperature that ranges from about 0°C to about 30°C and all temperature values in between.

In a fourth aspect of the seventeenth embodiment, the mole ratio of t-butylmagnesium halide to 1-((2R,3R,4R,5R)-3-fluoro-4-hydroxy-5-(hydroxymethyl)-3-methyltetrahydrofuran-2-yl)pyrimidine-2,4(1H,3H)-dione ranges
 30 from about 2 to about 2.2. In a first sub-aspect, the mole ratio of t-butylmagnesium halide to 1-((2R,3R,4R,5R)-3-fluoro-4-hydroxy-5-(hydroxymethyl)-3-methyltetrahydrofuran-2-yl)pyrimidine-2,4(1H,3H)-dione is about 2.1.


In a fifth aspect of the seventeenth embodiment, the t-butylmagnesium halide is t-butylmagnesium chloride.


An eighteenth embodiment is directed to a process for preparing substantially pure (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate, which comprises:

5 obtaining (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate according to any of the relevant embodiments disclosed herein, and

10 crystallizing the so-formed (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate.

A nineteenth embodiment is directed to an isotopically-labeled analog of *R_P-4* or *S_P-4*. The term "isotopically-labeled" analog refers to an analog of *R_P-4* or *S_P-4* that is a "deuterated analog", a "¹³C-labeled analog," or a "deuterated/¹³C-labeled analog." The term "deuterated analog" means a compound described herein, whereby a ¹H-isotope, i.e., hydrogen (H), is substituted by a ²H-isotope, i.e., deuterium (D). Deuterium substitution can be partial or complete. Partial deuterium substitution means that at least one hydrogen is substituted by at least one deuterium. For instance, for *R_P-4* or *S_P-4*, one of ordinary skill can contemplate at least the following partial deuterated analogs (where "*d_n*" represents n-number of deuterium atoms, such as, for an isopropyl group n = 1-7, while for a phenyl group, n = 1-5), as well as those depicted below.






Although the methyl groups depicted above are shown as being completely deuterated, one will recognize that partial-deuterated variations are also possible, such as, $-CD_2H$ and $-CD_2H$. Isotopic labels on the furanose and base are also contemplated. Likewise, the terms " ^{13}C -labeled analog" and "deuterated/ ^{13}C -labeled analog" refers to a compound described herein, whereby carbon atom is enriched with a ^{13}C -isotope meaning that the extent of enrichment exceeds the usual natural abundance of about 1.1%.

Examples

Not to be limited by way of example, the following examples serve to facilitate a better understanding of the disclosure.

Synthetic Aspects




In order to prepare the uridine nucleoside, one could take advantage of an advanced tribenzoylated cytidine intermediate in the synthesis of certain 3',5'-diacylated analogs of **3** (see below) already produced efficiently on a pilot plant scale (see WO 2006/031725 or US 2006/0122146, both of which are incorporated by reference in their entirety). The following method was found to be scalable and cost-efficient.



3',5'-O-dibenzo-2'-deoxy-2'-fluoro-2'-C-methyl- N^4 -benzoylcytidine (**1**) is obtained by a method disclosed in WO 2006/031725 and WO 2008/045419 both of

which are hereby incorporated by reference in their entirety. **1** is treated with 70% aqueous acetic acid to form 3',5'-O-dibenzoyl-2'-deoxy-2'-fluoro-2'-C-methyl-uridine (**2**). The benzoyl esters can be hydrolyzed by a number of methods as well, e.g., alkoxides in alcoholic solvent, such as sodium methoxide in methanol, potassium carbonate in methanol, or ethanol analogs, alkylamines such as methylamine in methanol, butylamine etc. Methanolic ammonia was chosen for the larger scale work. The uridine product (**3**) can be purified by crystallization to afford a 70% yield from the tribenzoylated cytidine (**1**).

Numerous literature procedures detail different routes and conditions to make phosphoramidates using several fold equivalents of reagents. See, for example, McGuigan et al. *J. Med. Chem.* 2005, 48, 3504-3515 and McGuigan et al. *J. Med. Chem.* 2006, 49, 7215. For process scale work, there is only one presently known example, which is disclosed in Lehsten et al., *Org. Process Res. Dev.* 2002, 6, 819-822 ("Lehsten"). In this reference, the authors introduce the concept of a "one-pot procedure" in which an amino acid hydrochloride salt and phenyl dichlorophosphate are reacted together with N-methylimidazole in dichloromethane. Later the nucleoside is added to form the desired 5'-O-phosphoramidate product, which in the present case would yield a compound represented by formula **4**. Unfortunately, the Lehsten procedure suffered from drawbacks. For example, the Lehsten procedure utilized a far larger excess of reagents than was necessary which added to the cost and difficulty of chromatographic purification. Furthermore, Lehsten suggested that one could control the reaction selectivity on the 5'-hydroxyl over the 3'-hydroxyl compared to a literature reference through using lower temperatures and slow addition of the nucleoside.



Using the Lehn procedure for the compounds disclosed herein provided
for about 1-5% of mono-substituted 3'-O-phosphoramidate diastereomers (**5**) and
about 10-30% of the bis-substituted product (**6**). As the polarity of the 3'-
diastereomers was very similar to the desired 5'-diastereomers (**4**), chromatographic
separation was very challenging. Scaling up the process was nearly impossible
without discarding a substantial portion of the less polar 5'-diastereomers (**4**) or
accepting a higher level of contamination of the 3'-diastereomers (**5**). In an initial 50
g scale-up, the resultant product contained a 3'-diastereomer (**5**) contamination of
about 3%, which co-eluted with the less polar of the 5'-diastereromer (**4**).

Disclosed herein are reaction conditions which use lesser amounts of reagents and a method to selectively remove the impurity 3'-O-phosphoramidate diastereomers (**5**) with an easier chromatographic separation thereby affording the 15 desired 5'-O-phosphoramidate diastereomers in much higher purity (**4**).

For the reagent stoichiometry, a study was made in which the stoichiometry of the reagents was systematically changed and the results were monitored by phosphorus NMR of the crude reaction as Lehsten had reported. In the more
20 successful runs, the isolated yield and purity of the desired product were compared. It was observed that the primary 5'-hydroxyl reacts at a faster rate than the secondary 3'-hydroxyl. This creates a competing situation between the reaction progress of consuming all the starting nucleoside and converting 5'- and 3'-monosubstituted products (**4** and **5**) to the 5',3'-bis substituted products (**6**). The 3'-monosubstituted
25 product converts to the bis product at a faster rate than the 5'-monosubstituted product, so it is possible to reduce the 3'-diastereomer contamination level by

pushing the reaction more to the bis-substituted products. However, with an effective way to remove the 3'-diastereomers, the reaction can be optimized to produce more of the desired 5'-diastereomer without having to sacrifice as much of the 5'-diastereomer being converted to the bis-substituted (**6**). It was also observed that the amino acid hydrochloride is very hygroscopic. As any water present would consume an equivalent amount of the phenyl dichlorophosphate reagent, care must be taken to keep the amino acid substantially anhydrous or it should be made substantially anhydrous prior to use. In short, Lehsten had reported that the optimum ratio of amino acid to phenyl dichlorophosphate to nucleoside was 3.5:2.5:1 respectively. It was found that the optimum ratio of amino acid to phenyl dichlorophosphate to nucleoside of about 1.6 to about 1.3 to about 1 is optimal under conditions in which the 3'-diastereomer can be efficiently removed and when the amino acid hydrochloride is substantially anhydrous. By using a smaller amount of the reagents, a cost savings is realized coupled with a simplification of the chromatographic separation of the desired product from reagent by-products and from the reduced level of bis diastereomers.

In one alternative procedure, a 3'-hydroxy-blocked derivative of **3** was prepared using a t-butyldimethylsilyl blocking group in two steps. This was then converted to its 5'-phosphoramidate derivative. The desire being that the silyl group could then be removed and there would be no 3' isomers (**5**) or 3',5'-bis phosphoramidates (**6**). A similar approach was demonstrated by Borch and Fries (U.S. Patent 5,233,031) in a low overall yield on an alkyl phosphoramidate.


Another alternative approach was to use the direct synthesis and then use chemistry to help differentiate the 3'-diastereomer impurities **5** from the desired 5'-diastereomers **4** to help the separation. A group was desired that would selectively react with the free primary hydroxyl of the 3'-O-phosphoramidate impurity **5** over the free secondary hydroxyl of the desired 5'-O-phosphoramidate **4**. It was also desired that the blocking group significantly change the polarity of the resulting 5'-O-blocked 3'-O-phosphoramidate product from the desired 5'-O-phosphoramidate **4**. There would be no extra step needed to remove the blocking group as the desired 5'-diastereomers **4** would not be changed. The chemically altered 3'-diastereomers would then allow easier chromatographic separation or separation by special scavenging supports or by extractions.


Specifically, the blocking group tert-butyldimethylsilyl (tBDMS) met these criteria and was the first one to be demonstrated and subsequently used on a multi-kilogram scale. Under certain conditions such as in pyridine as solvent and base, the tBDMS group reacts with high selectivity at the primary hydroxyl position over the 3' secondary hydroxyl position. The phosphoramidate reaction uses N-methylimidazole (NMI) as a base. In the presence of NMI, the silylation is less selective. Preferably, the amount of NMI should be reduced. This can be accomplished easily after the phosphoramidate reaction by washing the reaction solution with 1 N hydrochloric acid. The NMI and the remaining starting nucleoside are removed, leaving a crude mixture of mono and bis substituted products and reagent by-products. This is then dissolved in pyridine and treated with tert-butyldimethylsilyl chloride. The 3'-monosubstituted product **5** is converted in a few hours or less to the 5'-O-tBDMS-3'-O-phosphoramidate **7**. The reaction progress can be monitored by HPLC. The polarity of this silylated product **7** is less than the bis-phosphoramidate **6** and is readily removed by chromatography. Using this method, it was possible to reduce the level of 3'-monophosphoramidate **5** to less than 0.1% of the 5'-product **4** compared to 1-3% without the silyl treatment. Similarly, treatment with dimethoxytriphenylmethyl chloride (DMT-Cl) under the same conditions worked just as well. It was also easier to identify the DMT reaction product by TLC as DMT containing molecules stain bright orange on heating or exposure to acid. One can also envision many other blocking groups, as noted above.

Both the reaction conditions and the scavenging of the 3'-impurity are general methods and could be applied to most nucleoside phosphoramidates with a free 3' hydroxyl. The phosphoramidate moiety could be any combination of amino acid ester and aromatic alcohol. The nucleoside moiety could be any nucleoside in which a 5' phosphoramidate would lead to a 5'-monophosphate and could be further metabolized to the 5'-triphosphate form.

The following scheme is the main reaction scheme illustrated for making isopropyl L-alanate phenyl phosphoramidate of 2'-deoxy-2'-fluoro-2'-C-methyluridine with the major product as the desired 5'-O-phosphoramidate (**4**, two diastereomers) and the minor product as the 3'-O-phosphoramidate (**5**, two diastereomers) and the 3',5' -bis-O-phosphoramidate (**6**, four diastereomers). The reagents are added in the stoichiometric ratios as described in the method of

preparation section. The reaction is allowed to proceed until about 5% of the starting material remains as judged by UV visualization on thin layer chromatography (TLC). Also UPLC/MS showed approximately 10% of the 3',5' bis-phosphoramidate **6** had formed compared to the desired 5'-product. After quenching and an acidic aqueous workup, the crude residue from the organic layer was prepared for the silylation. Under the described reaction conditions, the silyl group preferentially reacted with the free 5'-hydroxyl of the 3'-O-phosphoramidate to form **7**. The reaction was continued until the 3'-O-phosphoramidate was no longer detectable by UPLC/MS.





After working up the silylation reaction, the desired product is subjected to chromatography on silica gel and is eluted with a gradient of methanol in dichloromethane (1-4%). The desired 5'-monophosphoramidate **4** elutes last.

Method of Preparation

10 **Example 1. Preparation of 2'-deoxy-2'-fluoro-2'-C-methyluridine (3)**

In a 10 L flask, was added 3', 5'-O-dibenzozyl-2'-deoxy-2'-fluoro-2'-C-methyl-N⁴-benzoylcytidine (500 g, 0.874 mol) and 70% aqueous acetic acid (7.5 L). The solution was heated to reflux (110°C) for 20 h. TLC indicated a complete reaction (R_f 0.6 in 5% methanol in dichloromethane (DCM)). The mixture was cooled to ambient temperature and diluted with water (2 L). After stirring for 2 h, the resulting precipitate was collected by filtration and the solid was rinsed with water (5 L) and dried in the atmosphere at ambient temperature for 12 h to afford 360 g (88%). This dibenzoyluridine intermediate was used directly in the next step by adding it all to freshly prepared methanolic ammonia (5.4 L, ca 25%) at 0°C. This temperature was maintained for 3 h and then allowed to warm to 15°C for 24 h. TLC indicated a complete reaction (R_f 0.4 in 10% methanol in DCM). The reaction mixture was filtered through a Celite bed and concentrated under reduced pressure to give the crude product (216 g). The crude product was stirred with ethyl acetate (325 mL) for 3 h at ambient temperature. The resulting solid was collected by filtration and washed with ethyl acetate (216 mL). The solid was dried under vacuum at ambient temperature for 4h to afford 160 g (78%) of the desired product

in 98.7% HPLC purity. ¹H-NMR (DMSO-*d*₆) δ 11.44 (br s, 1H, NH), 7.95 (d, 1H, C-6H), 5.97 (d, 1H, C-1'H), 5.64 (d, 1H, C-5H), 3.84-3.77 (m, 3H, C-5'-Ha, C-3'H, C-4'H), 3.63-3.60 (m, 1H, C5'-Hb), 1.23 (d, 3H, C-2'-CH₃). ES-MS M-1 259.

5 **Example 2. Preparation of (S)-2-[(1R,4R,5R)-5-(2,4-Dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-4-(R)-fluoro-3-hydroxy-4-methyl-tetrahydro-furan-2-ylmethoxy]-phenoxy-phosphorylamino}-propionic acid isopropyl ester (4)**

Synonym: 5'-O-(Isopropyl-L-alanate, phenyl phosphoramidyl)-2'-deoxy-2'-
10 fluoro-2'-C-methyl-uridine diastereomeric mixture.

A 5 L 3-necked flask was fitted with a mechanical stirrer, brine ice bath, internal thermometer, and a nitrogen atmosphere. The flask was charged with L-alanine isopropyl ester hydrochloride (82.0 g, 0.490 moles) and anhydrous dichloromethane (0.80 L). While this was stirring, phenyl dichlorophosphate (85.0 g,
15 0.40 moles) was added in one lot and stirred. While maintaining the internal temperature between -5 to 5°C, a solution of N-methylimidazole (NMI, 250 g, 3.07 moles) in dichloromethane (250 mL) was added over a period of a half hour. The solution was allowed to stir for 1 h in this temperature range. 2'-Deoxy-2'-fluoro-2'-C-methyl-uridine (3, 80.0 g, 0.307 moles) was added at 0°C in one portion and then
20 the reaction flask was allowed to warm up slowly in the brine bath. At 1 h, the internal temperature was up to -2°C. TLC (5% Methanol in DCM) at 1 h showed that more than 50% of nucleoside was consumed. The bath was removed and the reaction flask reached ambient temperature over 1 h more. TLC after 3 h and at 5 h total showed 95% of the starting nucleoside was consumed. The reaction mixture
25 was quenched by adding methanol (100 mL) and stirring the reaction for 5 minutes.

The reaction mixture was washed with 1N HCl (2 X 500 mL) followed by saturated sodium bicarbonate solution (2 X 500 mL). The separated organic layer was dried over anhydrous sodium sulfate (50 g) and filtered. The solution was evaporated under reduced pressure and then under high vacuum to dryness to give
30 the crude product as a viscous oil (170 g). NMRs of the crude product (³¹P and ¹H) were taken. The ³¹P-NMR indicated about 1% of the total phosphorus integration was due to the presence of the 3' isomer 5.

To the crude product was added anhydrous pyridine (1700 mL). The solvent was evaporated under reduced pressure and then under high vacuum in order to

reduce the water content of the crude mixture through co-evaporation. The resulting oil was re-dissolved in anhydrous pyridine (500 ml) and then was added excess *t*-butyldimethylsilyl chloride (9.0 g, 60mM). The reaction was stirred at ambient temperature. Reaction progress was monitored by UPLC/MS. After 3 hours, the 3' impurity **5** could no longer be detected and the reaction was quenched by the addition of methanol (50 mL).

The reaction was evaporated under reduced pressure to an oil. The residue was dissolved in ethyl acetate (1.5 L) and washed with 1N HCl (2X 500 mL), followed by saturated sodium bicarbonate solution (2 X 500 mL). The organic layer was dried over anhydrous sodium sulfate (50 g), filtered and evaporated under reduced pressure to give the crude product as a pale yellow oil.

The crude oil was diluted with the same volume of dichloromethane and loaded onto a 2.5 Kg silica gel cartridge in a radial compression module at 100 psi of air pressure. Using a gradient pump at 60 psi and a flow rate of 400 mL/min, the cartridge was washed with methylene chloride (4L) followed by a gradient 1-4% methanol in methylene chloride (48 L). Most of the major impurities (di-(isopropylalanyl)phenyl phosphate, 3',5'-bis phosphoramidate (**6**), 3'-phosphoramidate-5'-TBDMS adduct (**7**)) eluted with ~3% gradient. The desired product eluted between 3 and 4% methanol. The product containing fractions were sorted into two lots. The first contained small amounts of upper impurities and the latter was pure product. The first set of fractions contained small amounts of less polar impurities (upper impurities) such as the 3',5'-bis phosphoramidate and the di-alanylphenyl phosphate and a mostly the *R_P* diastereomer and required a second column purification. (The relative terminology, upper vs. lower refers to the elution on normal-phase silica-gel chromatography, where the "upper isomer" means the first eluting isomer.) The second set of fractions did not have a significant amount of impurities –just the remaining *R_P* and mostly the *S_P* diastereomers. It was later recombined with the twice-columned fractions. The solvent was evaporated under reduced pressure and the resulting white foam was further dried (0.20 mmHg) for 1 h to give 42 g of the impure lot (4:1 upper vs lower isomer based of ³¹P-NMR) and 38 g of the pure lot (1:3 upper vs lower isomer). The impure lot was recolumned in a similar manner to give 3.8 g of 97% pure upper isomer (fraction set aside) and 36 g of pure product in a 4:1 ratio. The two main lots were dissolved in DCM, combined, evaporated under reduced pressure and dried (50°C, 0.2 mmHg, 24 h) to

get 74 g (45.7%) of pure product **4** with a diastereomeric ratio of 48: 51, as a white foam, mp about 75-85°C.

In order to produce an amorphous solid of the diastereomeric mixture, 74 g of the white foam was stirred in with t-butyl methyl ether (750 mL) resulting in a partial solution and a gummy solid residue. While stirring, heptanes (750 mL) was added slowly and the suspension was mechanically stirred for 1 hour until most of the gum was converted to a white solid. The solid was scraped up with a spatula and the resulting slurry was filtered. The solid was washed with heptanes (4 X 50 mL) and dried under vacuum (50°C, 0.2 mmHg, 24 h) to give a white, amorphous powder (64 g) with a broad melting range of ca 70-80°C. ¹H and ³¹P NMR conformed to structure and HPLC showed a purity of 99.8% with a diastereomeric ratio of 46:54 (also confirmed by ³¹P NMR).

Alternative method to make solid mixture of **4**. After chromatography, the residue was co-evaporated with dichloromethane twice (5 mL/g) and dried for 24 h at 35-40°C at 35-45 mTorr. The foam residue was sieved through a 250 micron screen and further dried under the same conditions until the residual dichloromethane fell below 400 ppm as measured by headspace GC. The resulting fine off-white to white amorphous powder has a glass transition temperature range of 53.7 to 63.5°C.

Characterization of the mixture of isomers (**4**): ¹H-NMR (CDCl₃) δ 10.05 (br s, 1H, NH, *S_P*), 10.00 (br s, 1H, NH, *R_P*), 7.49 (d, 1H, C6-H, *S_P*), 7.36 (m, 5H, C6-H, *R_P*, aromatic), 7.23-7.14 (m, 6H, *R_P/S_P*, aromatic), 6.18 (br d, 2H, C1'-H, *R_P/S_P*), 5.63 (d, 1H, C5-H, *S_P*), 5.58 (d, 1H, C5-H, *R_P*), 5.01 (m, 2H, CH-(CH₃)₂, *R_P/S_P*), 4.46-4.33 (m, 8H, C-5'-H₂, ala-NH, C3'-OH, *R_P/S_P*), 4.12 (m, 2 H, ala-CH-CH₃, *R_P/S_P*), 4.01-3.85 (m, 4H, C3'-H, C4'-H, *R_P/S_P*), 1.39-1.22 (m, 12H, all CH₃, *R_P/S_P*).


³¹P-NMR (CDCl₃) δ 3.60 (*R_P*), 3.20 (*S_P*) relative to triphenylphosphate at -17.80 ppm. ES-MS M+1 530.2. Elemental Analysis: Calculated % (including 0.29% water as found by Karl Fisher analysis) C, 49.75; H, 5.54; N, 7.90, F, 3.58, P, 5.84. Found %: C, 49.50; H, 5.44; N, 7.85; F, 3.62; P, 6.05.

Discussion on separation of isomers


Compound **4** due to the chirality at phosphorus is comprised of two diastereomers, which are designated as *S_P*-**4** and *R_P*-**4**. The stereochemical

assignment was made based on single crystal X-ray analysis of *S_p*-**4**. Both *R_p*-**4** and *S_p*-**4** gave crystalline product.

The procedures for crystallization are outlined below.



S_p-**4**



R_p-**4**

Example 3. Crystallization of the *R_p*-**4** isomer. The chromatographed fraction of containing the first eluting, less polar *R_p*-**4** isomer (3.8 g, 97% pure) was dissolved in isopropanol (36 g) and diluted with heptanes until cloudy (72 g). The solution was seeded and stirred at ambient temperature for 5 h. The resulting solid was collected by vacuum filtration, washed with heptanes (2x20 mL) and dried (50°C, 0.2 mm, 24 h) to 2.3 g of very small white needles mp 136.2-137.8°C. HPLC purity of the resultant material was found to be 99.02%.

R_p-**4**: ¹H-NMR (CDCl₃) δ 9.10 (br s, 1H, NH), 7.36 (m, 2H, *o*-aromatic), 7.26-7.16 (m, 4 H, C6-H, *m,p*-aromatic), 6.16 (br d, 1H, C1'-H), 5.58 (d, 1H, C5-H), 5.01 (sept, 1H, CH-(CH₃)₂), 4.52-4.47 (m, 2H, C-5'-H₂), 4.10 (d, 1H, C3'-H), 4.02-3.76 (m, 4H, ala-NH, C3'-OH, C4'-H, ala-CH-CH₃), 1.37-1.20 (m, 12H, all CH₃).

Example 4. Preparation and crystallization of *S_p*-**4**.

Method 1: Direct precipitation from crude **4**: To a stirred solution of L-alanine isopropyl ester hydrochloride (10.5g, 61.5mmol, azeotropically dried, two times, with 50 mL of toluene each time) in dichloromethane (100 mL) was added phenyl dichlorophosphate (7.5 mL, 50 mmol) at room temperature. The mixture was cooled to -10°C and then was added a solution of NMI (30.5 mL, 384.3mmol) in 30 mL of dichloromethane over a period of 30 min. After completion of the addition, the mixture was stirred between -10 and -15°C for 1h. To the above mixture was added 2'-deoxy-2'-fluoro-2'-C-methyluridine (**3**) (10g, 38.4mmol) in one lot and the mixture was stirred below -10°C for 3h and then slowly allowed to warm to 20°C (6h). The mixture was stirred at this temperature over night (15h) and then quenched

with 10 mL of methanol. The solvent was evaporated and the residue was re-dissolved in EtOAc (200 mL). The EtOAc layer was washed with water (100mL), 1N HCl (3x75 mL), 2 % aqueous NaHCO₃ solution (50 mL) and brine (50mL). The organic layer was dried over Na₂SO₄, filtered and concentrated. The residue was
5 dried under high vacuum for 2h to give white foam (22 g).

The above foam was dissolved in 33 mL of DCM and then was added 65 mL of IPE (isopropyl ether) to give a saturated solution. The solution was filtered through a small pad of Celite and the filtrate was stirred with S_P-4 seeds for 72h at ambient temperature (about 22°C –note that cooling the suspension to 0°C led to
10 oiling out the crude product). The white solid was filtered, washed with IPE (20mL) and dried to give 4.58g (~85:15 mixture of S_P-4:R_P-4 respectively as determined by ³¹P NMR) of a white powder. The above solid was suspended in 23 mL of DCM and then refluxed for 3h. The mixture was cooled to room temperature and stirred for 15h. The white solid was filtered, washed with 4.5 mL of cold DCM and dried
15 under high vacuum at 45°C to give pure S_P-4, mp 93.9-104.7°C, HPLC purity 99.74% (3.11g, 15.2 % from the uridine nucleoside).

S_P-4 ¹H-NMR (CDCl₃) δ 8.63 (br s, 1H, NH), 7.47 (d, 1H, C6-H), 7.30 (m, 2H, *o*-aromatic), 7.26-7.18 (m, 3H, *m,p*-aromatic), 6.18 (br d, 1H, C1'-H), 5.70 (d, 1H, C5-H), 5.02 (sept, CH-(CH₃)₂), 4.53 (m, 2H, C-5'-H₂), 4.11 (d, 1H, C3'-H),
20 3.97 (m, 3H, C3'-OH, C4'-H, ala-CH-CH₃), 3.77 (br s, 1H, ala-NH), 1.39 (d, 3H, C2'-CH₃), 1.37 (d, 3H, ala-CH₃), 1.24 (d, 6H, CH-(CH₃)₂).

Method 2: Oiling out from crude 4: To a stirred solution of L-alanine isopropyl ester hydrochloride (20.6g, 123mmol, azeotropically dried, two times, with 75 mL of toluene each time) in dichloromethane (200 mL) was added phenyl
25 dichlorophosphate (14.9 mL, 100mmol) at room temperature. The mixture was cooled to -10°C and then was added a solution of NMI (61.3 mL, 769mmol) in 60 mL of dichloromethane over a period of 30 min. After completion of the addition, the mixture was stirred between -10°C and -15°C for 1h. To the above mixture was added 2'-deoxy-2'-fluoro-2'-C-methyluridine (**3**) (20g, 76.9mmol) in one lot and the
30 mixture was stirred below -10°C for 3h and then slowly allowed to warm to 20°C (6h). The mixture was stirred at this temperature over night (15h) and then quenched with 10 mL of methanol. The solvent was evaporated and the residue was re-dissolved in EtOAc (400 mL). The EtOAc layer was washed with water (200mL), 1N HCl (3x100 mL), 2 %

aqueous NaHCO₃ solution (100 mL) and brine (50mL). The organic layer was dried over Na₂SO₄, filtered and concentrated. The residue was dried under high vacuum for 2h to give white foam (43 g). The above foam was dissolved in 86 mL of EtOAc in a two neck round bottom flask equipped with a mechanical stirrer. While stirring, 100 mL of heptane was added slowly and the suspension was stirred for 1h. The top layer was decanted and the residue was again stirred with 50 mL of 2:3 EtOAc/heptane solutions for 10 min and then decanted. The residue was dried under high vacuum to give white foam (31g).


The above foam was dissolved in 46 mL of DCM and then was added 95 mL of IPE to give a saturated solution. The solution was filtered through a small pad of Celite and the filtrate was stirred with S_P-4 seeds for 72h at ambient temperature. The white solid was filtered, washed with IPE (30mL) and dried to give 7.33g (~85:15 mixture of S_P-4 : R_P-4 respectively as determined by ³¹P NMR) of white powder. The above solid was suspended in 36 mL of DCM and then refluxed for 3h. The mixture was cooled to room temperature and stirred for 15h. The white solid was filtered, washed with 7.5 mL of cold DCM and dried under high vacuum at 45°C to give >99% pure S_P-4, (4.78g, 11.6 % from the uridine nucleoside).

Method 3: Silica gel loading of crude 4: 5.0 g of crude 4 was produced as in the same manner as the mixture of diastereomers just before the column chromatography step starting with approximately 2.5 g of 2'-deoxy-2'-fluoro-2'-C-methyluridine (3). The crude was dissolved in 10 mL of DCM and 10 g of silica gel was added to the solution. The solvent was evaporated to give dry slurry. The slurry was stirred with 40 mL of 50% EtOAc/hexanes for 15 min and then filtered. The silica gel was washed with additional 10 mL of 50% EtOAc/hexanes. The silica gel was then washed with 15% MeOH/DCM (100 mL) and collected separately. The solvent was evaporated and dried under high vacuum to give 4.0 g of residue (foam). The residue was dissolved in DCM (6mL) and then was added ~9mL of IPE to make a saturated solution. The mixture was then gently stirred overnight with S_P-4 seeds at ambient temperature. The white solid was filtered and washed with IPE (5 mL) to give 1.28 g of product. ³¹P NMR revealed that the above product contains 77:23 mixture of S_P-4 : R_P-4 respectively. This was recrystallized from 20 mL of DCM to obtain 0.75 g of >99% pure S_P-4 (about 12% from the uridine nucleoside). This preparation of S_P-4 does not require the silylation step as done for the mixture,

so the entire reaction procedure is shown above. Aspects of single crystalline and polymorphic forms of *S_P*-4 are presented below.

Method 4: 40.0 g of 1:1 mixture of 4 was dissolved in 90 mL of dichloromethane. Diisopropylether (70 mL) was added to the above solution to give a saturated solution. (The quantity of diisopropyl ether may vary based on the purity of the product.) The solution was seeded with pure *S_P*-4 (> 99%) and the mixture was gently stirred with a stir bar at room temperature for 20h (formation of solid was observed after 2h). The solid was filtered, washed with 40 mL of the mixture of diisopropylether/dichloromethane (1:1) and dried to give white solid (16.6 g, 89.35 % pure *S_P*-4 by NMR). This solid was suspended in 83 mL dichloromethane and refluxed for 3h. The suspension was cooled to room temperature and stirred over night. The solid was filtered and washed with 10mL of cold DCM. The solid was dried under vacuum to give *S_P*-4 (13.1 g, 99.48 % pure by HPLC). 11g of this solid was redissolved in 330 mL of DCM under hot conditions. The solution was cooled to room temperature and left at this temperature over night. The crystalline product was filtered and dried to give 10.5 g of *S_P*-4 (99.74 % by HPLC).


Compounds *S_P*-4 and *R_P*-4 may alternatively be prepared, in accordance with the ninth or tenth embodiment, by reacting nucleoside (protected or unprotected) 3 with an isopropyl-alanyl-phosphoramidate (mixture of C and C', C or C'), as shown in the following equation.



P.D. Howes et al. *Nucleosides, Nucleotides & Nucleic Acids* 2003, Vol. 22, Nos. 5-8, pp. 687-689 ("Howes") discloses 2'- and 5'-phosphoramidates obtained by a reaction with t-butylmagnesium chloride. There, Howes discloses that when a 3'-deoxy-cytidine nucleoside is reacted with (S)-2-[chloro-phenoxy-phosphorylamino] propionic acid methyl ester in the presence of 1.2 equivalents of t-butylmagnesium chloride, selective phosphorylation on the 2'-position occurred, but that with an additional equivalent of t-butylmagnesium chloride selective phosphorylation on the


5'-position occurred. This disclosure should be contrasted to that which is disclosed in Howes' Scheme 1.

Example 5-1. Preparation of (S)-2-[(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester



To a stirred solution of 4-nitrophenyl phosphorodichloridate 12.8g, 50 mmol) in dichloromethane (100 mL) was added a solution of phenol and triethylamine (7.7 mL, 55 mmol) in dichloromethane (100 mL) at -78°C over a period of 20min. The mixture was stirred at this temperature for 30min and then transferred to another round bottom flask containing L-alanine isopropyl ester hydrochloride (8.38g, 50mmol) in dichloromethane (100 mL) at 0°C. To the mixture was added second portion of triethylamine (14.6 mL, 105 mmol) over a period of 15min. The mixture was stirred at 0°C for 1h and then the solvent was evaporated. The residue was triturated with ethyl acetate (150 mL) and the white solid was filtered off. The filtrate was concentrated under reduced pressure to give pale yellow oil. The crude compound was chromatographed using 0-20% ethyl acetate/hexanes gradient to give product (17g, 83 %yield) as a mixture of diastereomers in about 1:1 ratio. ³¹P NMR (162 MHz, DMSO-d₆): δ -0.31, -0.47; ¹H NMR (400 MHz, DMSO-d₆): δ 8.31-8.27 (m, 2H), 7.51-7.37(m, 4H), 7.27-7.19(m, 3H), 6.70-6.63(m, 1H), 4.85-4.78(m, 1H), 3.97-3.86(m, 1H), 1.21-1.19(m, 3H), 1.11-1.09 (m, 6H); MS (ESI) m/z 407 (M-1)⁺. ³¹P NMR (162 MHz, CDCl₃): δ -2.05, -2.10; ¹H NMR (400 MHz, CDCl₃): δ 8.22 (d, J = 9.2Hz, 2H), 7.41-7.33(m, 4H), 7.26-7.18(m, 3H), 5.05-4.96(m, 1H), 4.14-4.05(m, 1H), 3.93-3.88(m, 1H), 1.38(d, J = 6.8Hz, 3H), 1.22 (dd, J = 6.2 & 3.0Hz, 6H); MS (ESI) m/z 407 (M-1)⁺.

Example 5-2. Preparation of S_P-4/R_P-4.




To a stirred solution of 1-((2R,3R,4R,5R)-3-Fluoro-4-hydroxy-5-hydroxymethyl-3-methyl-tetrahydro-furan-2-yl)-1H-pyrimidine-2,4-dione (130mg, 0.5mmol) in dry THF (1.5mL) was added a 1.0M solution of tert-butylmagnesium chloride (1.05mL, 1.05mmol, 2.1 equiv)) at room temperature over a period of 5min.

- 5 After 30min, a solution of (S)-2-[(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester (1:1 mixture of isomers, 408mg, 1mmol) in THF (1.5mL) was added drop-wise over a period of 5min. The mixture was allowed to stir at room temperature for 48h and then quenched with saturated aqueous NH_4Cl (20mL). The mixture was partitioned between ethyl acetate (50mL) and water (20mL). The combined organic extract was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a pale yellow residue. Column chromatography of the residue using 0-2% MeOH/dichloromethane gradient gave a white foamy solid (125mg, 47% yield, mixture of S_P -4/ R_P -4 in about 3.05:1.0 ratio).

15

Example 6. Preparation and non-chromatographic isolation of (S)-2-[(S)-(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester



- 20 L-alanine isopropyl ester hydrochloride (330 g, 1.97 mol) was pre-dried by co-evaporation with toluene (2 x 400 mL) under reduced pressure and then dried in a vacuum oven (50°C, 0.2 mmHg, 17 h). To a stirred solution of 4-nitrophenyl phosphorodichloridate (500.0 g, 1.953 mol) in anhydrous dichloromethane (3.0 L) was added a solution of phenol (183.8 g, 1.953 mol) and triethylamine (300 mL, 2.15 mol) in dichloromethane (900 mL) at -60°C internal temperature over a period of 3 hours. The mixture was stirred at this temperature for additional 30 min and then allowed to warm up to -5°C over 2.5 hours. The pre-dried amino acid ester was added at -5~0°C under an atmosphere of nitrogen over 10 mins. The residue of aminoester salt in the addition flask was transferred to the reaction mixture via
- 25 rinsing with dichloromethane (2 x 100 mL). The mixture was stirred at 0°C for 40 mins and a second portion of triethylamine (571 mL, 4.10 mol) was added over a period of 40 mins at 0°C. The mixture was stirred at 0~10°C for 3 h and then the

white solid (triethylamine hydrochloride) was filtered off and rinsed with dichloromethane (3 x 300 mL). The filtrate was concentrated under reduced pressure and the residue was triturated with methyl t-butyl ether (MTBE, 4 L). The additional solid salt thus formed was filtered off and rinsed with MTBE (3 x 150 mL). The filtrate was concentrated under reduced pressure to give clear light brown color oil. The residue was co-evaporated with hexanes (2 x 140 mL) to remove any residual MTBE and further dried under vacuum at 40°C for 2 hours. The dry residue was mixed with diisopropyl ether (IPE, 1.1 L) and stirred at 5 °C in an ice-water bath. Small amount of crystal seeds of the desired *S_P*-isomer of the product was added to the solution and the mixture was stirred at 5°C for over 22 h to form a medium thick slurry. This was allowed to stand in a freezer (-10°C) for 44 h. The precipitated product was collected via filtration and rinsed with pre-cooled mixed solvents of IPE and hexanes (1:1, 3 x 190 mL). The solid was dried under vacuum (0.5 mm Hg) at ambient temperature until a constant weight was obtained to give 227.23 g (yield: 28.5%) as a white powder solid. The ratio of two diastereomers *S_P*:*R_P* was 9.65/1 based on ³¹P NMR (162 MHz, DMSO-*d*₆, δ -0.31 (*S_P*), -0.47). The product was recrystallized by dissolving in IPE (840 mL) while heating in a 60°C bath. The above solution was stirred at room temperature for 1 h and then a small amount of crystal *S_P* isomer seeds was added. White powder solid was formed within 2 hours and the flask was stored in a freezer (-10°C) for 16 hours. A white and fine crystalline solid obtained was filtered, washed with pre-cooled IPE (3x 50 mL) and dried under vacuum (ambient, 0.5 mm Hg) to a constant weight to give white fluffy solid (177.7 g, 22% overall yield or 44% overall yield based on theoretical yield of the *S_P* isomer) with diastereomeric ratio of 48/1 based on ³¹P-NMR. Mp 62-66°C.

³¹P NMR (162 MHz, DMSO-*d*₆): δ -0.31; **¹H NMR (400 MHz, DMSO-*d*₆):** δ 8.30-8.27 (m, 2H), 7.49(d, J=8.8Hz, 2H), 7.41-7.37(m, 2H), 7.23-7.19 (m, 3H), 6.66 (dd, J=13.6, 10.0Hz, 1H), 4.86-4.78 (m, 1H), 3.97-3.86 (m, 1H), 1.19 (d, J=7.2Hz, 3H), 1.10(d, J=6.4Hz, 6H);

³¹P NMR (162 MHz, CDCl₃): δ -2.05; (162 MHz, DMSO-*d*₆): δ -0.31; **¹H NMR (400 MHz, CDCl₃):** δ 8.22 (d, J=9.2Hz, 2H), 7.41-7.33(m, 4H), 7.26-7.18(m, 3H), 5.05-4.96(m, 1H), 4.14-4.05(m, 1H), 3.93-3.88(m, 1H), 1.38(d, J=6.8Hz, 3H), 1.22 (dd, J=6.2 & 3.0Hz, 6H); **¹H NMR (400 MHz, DMSO-*d*₆):** δ 8.30-8.27 (m, 2H), 7.49(d, J=8.8Hz, 2H), 7.41-7.37(m, 2H), 7.23-7.19 (m, 3H), 6.66 (dd, J=13.6,

10.0Hz, 1H), 4.86-4.78 (m, 1H), 3.97-3.86 (m, 1H), 1.19 (d, J=7.2Hz, 3H), 1.10(d, J=6.4Hz, 6H)

MS (ESI) m/z 407 (M-1)⁺.


The stereochemistry of **8** (*S_P*-isomer) has been confirmed by single crystal X-ray crystallography, see details provided below.

Example 7. Separation of the diastereomeric mixture (S)-2-[(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester by SFC

A sample of the mixture of diastereomers (4.8 g) enriched with the *R_P*-isomer was subjected to SFC using a ChiralPak AD-H (2x15 cm) column and eluted with 35% isopropanol in carbon dioxide at 100 bar. An injection loading of 4 mL of sample at a concentration of 17 mg/mL of methanol was used. The *R_P*-isomer [(S)-2-[(R)-(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester] eluted first. The appropriate fractions of the multiple runs were combined and concentrated under reduced pressure to give 2.9 g of the *R_P*-isomer [(S)-2-[(R)-(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester] as a light yellow viscous oil and 1.9 g of the *S_P*-isomer [(S)-2-[(S)-(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester] as a white solid. Analytical data of *R_P*-isomer is similar to the product isolated by the above crystallization method.

Analytical Data for (S)-2-[(R)-(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester (8**, *R_P*-isomer):** ³¹P NMR (162 MHz, DMSO-d₆): δ -0.47; ¹H NMR (400 MHz, DMSO-d₆): δ 8.30-8.27 (m, 2H), 7.46-7.38 (m, 4H), 7.27-7.20 (m, 3H), 6.68 (dd, J=13.8, 10.2Hz, 1H), 4.86-4.77 (m, 1H), 3.97-3.86 (m, 1H), 1.20 (d, J=7.2Hz, 3H), 1.10(dd, J=6.2, 2.2Hz, 6H); MS (ESI) m/z 407 (M-1)⁺.

Example 8-1. Preparation of racemic 2-[(4-chloro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester (±):




To a stirred solution of 4-chloro-phenyl phosphorodichloridate (2.45g, 10.0 mmol) in dichloromethane (20 mL) was added a solution of phenol (0.94 g, 10 mmol) and triethylamine (1.56 mL, 11 mmol) in dichloromethane (20mL) at -78°C over a period of 20 min. The mixture was stirred at this temperature for 30min and then transferred to another round bottom flask containing L-alanine isopropyl ester hydrochloride (1.67 g, 10 mmol) in dichloromethane (50 mL) at 0°C. To the mixture was added second lot of triethylamine (2.92 mL, 21 mmol) over a period of 15min. The mixture was stirred at 0°C for 1h and then the solvent was evaporated. The residue was triturated with ethyl acetate (30mL) and the white solid was filtered off. The filtrate was concentrated under reduced pressure to give pale yellow oil. The crude compound was chromatographed using 10-20% ethyl acetate/hexanes gradient to give product (2.0 g, 50% yield) as a mixture of diastereomers in about 1:1 ratio. ³¹P NMR (162 MHz, CDCl₃): δ -1.58, -1.62; ¹H NMR (400 MHz, CDCl₃): δ 7.06-7.51 (m, 8H), 7.15-7.28 (m, 2H), 7.29-7.47(m, 2H), 4.0-4.10(m, 1H), 3.82-3.88 (m, 3H), 1.35-1.36 (dd, 6H); 1.19-1.22 (m, 3H). MS (ESI) *m/z* 398 (M-1)⁺. The resultant product is purified by extraction, crystallization, or chromatography, as noted above.

Example 8-2. Preparation of (S)-Isopropyl 2-((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)-phosphorylamino)propanoate (4).

To a stirred solution of 1-((2R,3R,4R,5R)-3-Fluoro-4-hydroxy-5-hydroxymethyl-3-methyl-tetrahydro-furan-2-yl)-1H-pyrimidine-2,4-dione (**3**, 2.6 g, 10 mmol) in dry THF (50 mL) was added a 1.7 M solution of tert-butyilmagnesium chloride (12.4 mL, 21 mmol, 2.1 equiv)) at room temperature over a period of 15 min. After 30 min, a solution of racemic (2-[(4-chloro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester (4.08g, 10mmol) in THF (15mL) was added drop wise over a period of 10min. The mixture was allowed to stir at room temperature for 72. TLC co-spot with authentic product showed around 5% of the desired product had formed compared to the starting nucleoside.

Example 9-1. Preparation of racemic 2-[(2-chloro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester ($\pm$).



5 To a stirred solution of 2-chloro-phenyl phosphorodichloridate (9.8 g, 40 mmol) in dichloromethane (80 mL) was added a solution of phenol (3.76 g, 40 mmol) and triethylamine (6.16 mL, 44 mmol) in dichloromethane (80 mL) at -78°C over a period of 20 min. The mixture was stirred at this temperature for 30 min and then transferred to another round bottom flask containing L-alanine isopropyl ester

10 hydrochloride (6.7 g, 40 mmol) in dichloromethane (150 mL) at 0°C. To the mixture was added second portion of triethylamine (11.6 mL, 84 mmol) over a period of 15 min. The mixture was stirred at 0°C for 1h and then the solvent was evaporated. The residue was triturated with ethyl acetate (100mL) and the white solid was filtered off. The filtrate was concentrated under reduced pressure to give

15 pale yellow oil. The crude compound was chromatographed using 10-20% ethyl acetate/hexanes gradient to give product (11.3g, 72 % yield) as a mixture of diastereomers in about 1:1 ratio. ³¹P NMR (162 MHz, CDCl₃): δ -1.58, -1.61; ¹H NMR (400 MHz, CDCl₃): δ 7.06-7.51 (m, 8H), 5.02-5.94 (m, 1H), 4.10-4.16(m, 1H), 3.31-3.94(m, 1H), 1.18-1.35(m, 3H), 1.38-1.40 (dd, 6H); MS (ESI) m/z 398 (M-1)⁺. The resultant product is purified by extraction, crystallization, or

20 chromatography, as noted above.

Example 9-2. Preparation of (S)-isopropyl 2-((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)- phosphorylamino)propanoate.

25


To a stirred solution of 1-((2R,3R,4R,5R)-3-Fluoro-4-hydroxy-5-hydroxymethyl-3-methyl-tetrahydro-furan-2-yl)-1H-pyrimidine-2,4-dione (**3**, 2.6 g, 10 mmol) in dry THF (50 mL) was added a 1.7 M solution of tert-butyilmagnesium chloride (12.4 mL, 21 mmol, 2.1 equiv)) at room temperature over a period of 15

30 min. After 30 min, a solution of (2-[(2-chloro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester (racemic 4.08 g, 10 mmol) in

THF (15mL) was added drop wise over a period of 10min. The mixture was allowed to stir at room temperature for 72h. TLC co-spot with authentic product showed around 5-10% of the desired product had formed compared to the starting nucleoside.


5

Example 10-1. Preparation of racemic 2-[(2,3,4,5,6-pentafluoro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester ($\pm$).



- 10 To a stirred solution of pentafluorophenyl phosphorodichloridate (6.0 g, 20 mmol) in dichloromethane (40 mL) was added a solution of phenol and triethylamine (3.08 mL, 22 mmol) in dichloromethane (40 mL) at -78°C over a period of 20 min. The mixture was stirred at this temperature for 30 min and then transferred to another round bottom flask containing L-alanine isopropyl ester
- 15 hydrochloride (3.35 g, 20 mmol) in dichloromethane (100 mL) at 0 °C. To the mixture was added second lot of triethylamine (5.84 mL, 42 mmol) over a period of 15 min. The mixture was stirred at 0°C for 1h and then the solvent was evaporated. The residue was triturated with ethyl acetate (60mL) and the white solid was filtered off. The filtrate was concentrated under reduced pressure to give pale yellow oil as a
- 20 mixture of diastereomers in about 1:1 ratio. ³¹P NMR (162 MHz, CDCl₃): δ -0.49, -0.58. The resultant product is purified by extraction, crystallization, or chromatography, as noted above.

- Example 10-2. Preparation of diastereomeric mixture (S)-2-[(2,3,4,5,6-pentafluoro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester and isolation of single diastereomer (S)-2-[(S)-(2,3,4,5,6-pentafluoro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester via**
- 5 crystallization-induced dynamic resolution with multiple crops.**



- To a 2 L of three-necked round bottom flask fitted with a mechanic stirrer and low temperature thermometer were added 60 g (284 mmol) of phenyl
- 10 dichlorophosphate and 300 mL of anhydrous dichloromethane. The solution was cooled to 0°C under nitrogen atmosphere and *iso*-propyl alanate hydrochloride salt (oven dried *in vacuo*, 47.7 g, 284 mmol) was added quickly as a solid. The mixture was stirred and cooled to -55°C in a dry ice-acetone bath. A solution of 60.32 g of triethylamine (596 mmol) in 300 mL of dichloromethane was added through an
- 15 addition funnel over 70 minutes. The white cloudy mixture was stirred at -55°C for half hour and then the temperature was raised to -5°C slowly over 1.5 h. A pre-cooled (room temperature) mixture of pentafluorophenol (52.42 g, 284 mmol) and triethylamine (32.11 g, 317 mmol) in 180 mL of dichloromethane was added to the mixture via an addition funnel over 1 h
- 20 at 0°C and the resulting mixture was stirred at 0°C for 4 hours. The white precipitate (TEA•HCl) was filtered out and rinsed with dichloromethane (3x 50 mL). The filtrate was concentrated under reduced pressure and white solid residue was triturated in 880 mL of *t*-butyl methyl ether (TBME) at room temperature for one hour. White suspension was filtered and solid was rinsed with TBME (3x 150 mL).
- 25 Solid was distributed in a mixture of ethyl acetate (600 mL) and water (150 mL).

Organic layer was separated and washed with water (3 x 100 mL). Organic layer was dried over MgSO_4 and concentrated to afford 29.92 g (66 mmol) of product (S_P -isomer, as confirmed by X-ray crystallography, *infra*) as a white feather solid.

The filtrate from above TBME trituration was concentrated under reduced pressure to a white solid residue and the solid was triturated in 450 mL of mixed ethyl acetate and hexanes (20:80, v/v) at room temperature for 75 minutes. Solid (solid 1) was collected by filtration and rinsed with 20% ethyl acetate in hexanes (75 mL, 2x 30 mL). The mother liquor was concentrated to furnish an off-white solid that was triturated in 20% ethyl acetate in hexanes (185 mL) at room temperature for 17 hours. A white solid (solid 2) was collected by filtration and rinsed with 20% ethyl acetate in hexanes (2 x 10 mL). Solid 1 and solid 2 were combined and dissolved in 1.2 L of ethyl acetate. The solution was washed with water (3x 150 mL), brine (50 mL) and dried over MgSO_4 . The solution was concentrated under reduced pressure to afford 72.8 g (161 mmol) of pure product. Total amount of product was 102.72 g (226 mmol, 80%). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.38-7.33 (m, 2 H), 7.27-7.24 (m, 2 H), 7.23-7.19 (m, 1 H), 5.04 (hept, 1 H), 4.18-4.09 (m, 1 H), 4.01-3.96 (m, 1 H), 1.45 (d, 3 H), 1.25 (dd, 6 H). ^{31}P NMR (CDCl_3 , 162 MHz) δ : -0.50.

Example 10-3: Preparation of diastereomeric mixture (S)-2-[(2,3,4,5,6-pentafluoro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester and isolation of single diastereomer (S)-2-[(S)-(2,3,4,5,6-pentafluoro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester via crystallization-induced dynamic resolution in a single crop.

To a 1 L of dry three-necked flask fitted with a low-temperature thermometer and a mechanical stirrer was loaded phenyl phosphorodichloridate (25 g, 118.5 mmol). Anhydrous dichloromethane (125 mL) was added and the solution was cooled to 0°C. The alanine ester salt (oven dried) (19.86 g, 1 eq) was added quickly under N_2 while agitated. The solution was cooled to ca -50°C (internal temperature (in an acetone/dry ice bath under N_2)). A solution of triethylamine (25.2 g, 2.1 eq) in DCM (125 mL) was added dropwise via an addition funnel over 0.5 h at -50°C and the resulting white slurry was stirred at about -50°C for 0.5 h. The mixture was allowed to warm up to 0°C over 1.5 h and then a pre-mixed cooled solution of

pentafluorophenol (21.82 g, 1 eq) and TEA (13.2 g, 1.1 eq) (caution: heat released while mixing pentafluorophenol and TEA) in 75 mL of DCM was added over 0.5 h at 0°C via an addition funnel. The mixture was stirred at 0°C for additional 4 h.

The mixture was filtered through a Buchner funnel and the collected solid
5 triethylamine hydrochloride was rinsed with DCM (3 x 40 mL). The filtrate was checked by ³¹P-NMR (ratio ca 1.14:1 favored the *S_P*-diastereomer – downfield peak) and was divided into two parts of equal weight. One of them was concentrated under reduced pressure. The white solid residue (31 g) was triturated in a mixture of EtOAc and hexanes (150 mL, 20:80, v/v) at RT for 17 h allowing time for dynamic
10 resolution of the less soluble *S_P* isomer. The white slurry was filtered and solid was rinsed with 20% EtOAc in hexanes (2x 25 mL). The solid (22.58 g) was checked by ¹H-NMR and ³¹P-NMR and it contained product as one isomer contaminated with triethylamine hydrochloride salt. The solid was dissolved and partitioned in 310 mL of EtOAc and 100 mL of water. After separation of the organic layer, the aqueous
15 layer was back-extracted with EtOAc (50 mL). The combined organic layer was washed with water (3x 80 mL), brine (50 mL) and dried over MgSO₄. The solution was concentrated under reduced pressure and then dried under high vacuum at RT to a constant weight to furnish 17.36 g of product as a white solid from the one half of the reaction.. The yield is 64%. The mother liquor from above was concentrated to
20 a gummy residue (7.89 g) that contained the reagents with a ratio of 1:1.2 (desired/undesired) based on ³¹P-NMR.

Example 10-4. Preparation of (S)-2-[(2,3,4,5,6-pentafluoro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester

25 DCM (11.5 L) was charged into the clean and dry glass reactor. Phenylldichlorophosphate (2.3 kg, 10.9 mol) was charged to the reactor under nitrogen. The solution was then cooled to 0°C. L-Alanine isopropylester hydrochloride (1.83 kg , 10.9 mol) was then added in one portion and continued stirring for 30 min. The reaction mass was cooled to -50°C internal temperature by
30 using a dry ice/acetone bath. A mixture of TEA (2.1 eq, 3.17 L) in DCM (11.5 L) was added to the above reaction solution slowly over a period of 8 h to keep the internal temperature between -40 to -50°C. After completion of addition, the

reaction was maintained at the same temperature range for about 1h. The mixture was allowed to warm to 0°C over about 4h.

In the meantime, in another reactor, DCM (6.9 L) was charged, and then pentafluorophenol (2.0 Kg, 10.9 mol) was added under nitrogen. The solution was cooled to 0°C, then TEA (1.1 eq, 1.65 L) was added to the pentafluorophenol solution (exothermic) over a period of about 2 h. The resulting solution in turn was slowly added to the first solution containing the phenyl dichlorophosphate and amino acid ester while keeping the temperature between 0 to 5°C over a period of about 7 h. After the completion of the addition, stirring was continued at that temperature range for about 4 h. The reaction progress was monitored by HPLC. When less than 5% of pentafluorophenol remained, the reaction was stopped. Note that chiral HPLC indicated an even mixture of diastereomers of the product at this point.

The reaction suspension was filtered through a Nutsche filter to remove most of the suspended triethylamine hydrochloride salt. The salt cake was washed with excess amount of DCM (9 L) and this wash was added to the main filtrate. The filtrate was concentrated under reduced pressure at 35°C to give a solid residue. The solid residue was co-evaporated with hexane (4 L) to further reduce levels of residual DCM. To this residual solid was added 20% MTBE/Hexane 6 L and the suspension was stirred for about 17 hr at ambient temperature and monitored by HPLC. The pH of the solution remained basic due to residual TEA. During this time, a dynamic resolution took place in which the precipitated solid was the desired S_p-diastereomer isomer and the supernatant remained in equilibrium between S_p-diastereomer and R_p-diastereomer.


The suspension was passed through a Nutsche filter and the desired product solid, still contaminated with TEA hydrochloride, was washed with 5% MTBE/Hexane (1 L). The solid was dissolved in ethyl acetate (35 L) and the solution was washed with water (3 x 35 L) and brine (10 L) and then the solution was dried over solid sodium sulfate, filtered concentrated under reduced pressure keeping the reactor temp below 44°C. The solid residue was co-evaporated with hexane (4 L). The reactor was brought to ambient temperature and 5% MTBE/Hexane (5 L) was added. The thick suspension was stirred for 15 min and then the solid was collected by filtration. The collected solid was washed with of hexane (2.5 L) and dried under high vacuum at ambient temperature to a constant

weight to give the final product (S_P -diastereomer) as a white solid, 2.6 kg (53%); 99.5% pure by HPLC, 0.4% of other R_P -diastereomer.

Example 10-5. Preparation of (S)-isopropyl 2-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)-phosphorylamino)propanoate.

To a stirred solution of 1-((2R,3R,4R,5R)-3-Fluoro-4-hydroxy-5-hydroxymethyl-3-methyl-tetrahydro-furan-2-yl)-1H-pyrimidine-2,4-dione (**3**, 2.6g, 10mmol) in dry THF (50mL) was added a 1.7M solution of *tert*-butylmagnesium chloride (12.4mL, 21mmol, 2.1 equiv)) at room temperature over a period of 15min. After 30min, a solution of crude racemic (**2**-[(2,3,4,5,6-pentafluoro phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester (4.08g, 10mmol) in THF (15mL) was added drop wise over a period of 10min. The mixture was allowed to stir at room temperature for 72h. TLC co-spot with authentic product showed around 40-50% of the desired product had formed compared to the starting nucleoside.


Example 10-6. Preparation of (S)-Isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphorylamino)propanoate (S_P -4) using (S)-2-[(S)-(2,3,4,5,6-pentafluoro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester and purification by crystallization only.



To a stirred solution of **3** (10g, 38.46mmol, dried under vacuum at 50°C for 20h) in dry THF (165mL) was added a 1.7M solution of *tert*-butylmagnesium chloride in THF (47.5 mL, 80.77 mmol) over a period of 20min while keeping the

flask in a cold water bath (5°C) under a nitrogen atmosphere. After completion of the addition, the cold bath was removed and the white suspension was stirred at room temperature (20°C) for 30 min. A solution of (*S*)-2-[(*S*)-(2,3,4,5,6-pentafluorophenoxy)-phenoxyphosphorylamino] propionic acid isopropyl ester (20.9 g, 46.11 mmol) in anhydrous THF (165 mL) was then added to the reaction mixture over a period of 30 min. The mixture was stirred at room temperature (20°C) for 3.5 h. Stirring was continued for additional 1.5 h, at which stage TLC indicated >95% conversion and no significant difference in 3',5'-*bis*-phosphoramidate impurity intensity from 2h. The reaction mixture was quenched with saturated aq. NH₄Cl (10 mL) and then the solvent was evaporated at 25°C. The residue was partitioned between ethyl acetate (400 mL) and saturated ammonium chloride (60 mL)/water (20 mL) mixture. The organic layer was separated, washed with saturated ammonium chloride (80 mL) and water (3 x 60 mL). The aqueous layer up to this point was kept separately. The organic layer was washed with 5% aq. sodium carbonate (3 x 50 mL) and water (2 x 60 mL). The first aqueous layer was extracted with additional ethyl acetate (100 mL), washed with water (2 x 20 mL) and then the aqueous layer obtained from the sodium carbonate washings was extracted with the same ethyl acetate extract. The organic layer was washed with water (2 x 20 mL) and combined with the main lot. The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated to give a foamy solid (19.32 g). The residue was dissolved in 60 mL of dichloromethane (a white solid was precipitated out and a cake was formed in about five minutes) and then was added 25 mL of IPE. The suspension was stirred at room temperature for 2 h. The white solid was filtered, washed with 1:1 mixture of cold (0°C) IPE/dichloromethane (20 mL) and dried to give 11.77 g (58% yield) of the product as an amorphous white solid. The above solid was re-dissolved in dichloromethane (350 mL), filtered and evaporated under atmospheric pressure (45°C bath temperature) to a volume of ~120 mL. The solution was allowed to stand at room temperature (21°C) for 20 h. The white crystalline solid (dichloromethane solvate) was collected by filtration, washed with cold (0°C) dichloromethane (10 mL) and dried under high vacuum for 4 h at ambient temperature to give pure non-solvated product (10.62 g, 52% yield) as white needles. HPLC purity 99.8%. Spectral properties match those values reported herein.

Example 10-7. Preparation of (S)-Isopropyl 2-((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphorylamino)propanoate (*S_P*-4) using (S)-2-[(S)-(2,3,4,5,6-pentafluoro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester, modified reaction conditions and workup and purification by crystallization only.



10


To a stirred suspension of 1-((2R,3R,4R,5R)-3-fluoro-4-hydroxy-5-(hydroxymethyl)-3-methyltetrahydrofuran-2-yl)pyrimidine-2,4(1H,3H)-dione (**3**, 5.0 g, 19.1 mmol, dried under vacuum at 50°C for 20 h) in dry THF (75 mL) was added a 1.7 M solution of tert-butylmagnesium chloride in THF (23.7 mL, 40.35 mmol) using an addition funnel over a period of 30 min at -5°C. The white suspension was stirred at this temperature for 30 min and then warmed to ambient temperature (20°C) at which temperature it was stirred for additional 30 min. The reaction mixture was cooled to 5°C and then was added a solution of (S)-2-[(S)-(2,3,4,5,6-pentafluorophenoxy)-phenoxyphosphorylamino] propionic acid isopropyl ester (10.45 g, 23.06 mmol) in THF (50 mL) over a period of 30 min. The mixture was stirred at 5°C for 18 h, cooled to -5°C and then quenched with 2N HCl (25 mL). Toluene (100 mL) was added to the mixture and warmed to room temperature. After 20 min the layers were separated. The organic layer was washed with 1N HCl (2 x 10mL), water (10 mL), 5% aqueous Na₂CO₃ (4 x 10mL), water (2 x 10mL) and brine (10 mL). All the aqueous layers were re-extracted with toluene (20 mL), washed with 5% aqueous Na₂CO₃ (2 x 5 mL), water (10 mL) and brine (5 mL). The combined organic layer was dried over anhydrous sodium sulfate, filtered and evaporated to an approximate volume of 20 mL. Dichloromethane (20 mL) was added to the solution and the mixture was stirred at room temperature for 18 h. The solid was filtered, washed with 1:1 MTBE/DCM mixture (2 x 10mL) and dried under high vacuum to give white solid (7.7 g). HPLC of the solid at this indicated

25

30

98.21% *S_P*-**4**, 0.18% of un-reacted **3** and 0.67% of 3',5'-bis-phosphoramidate impurity. The above solid of *S_P*-**4** was re-dissolved in dichloromethane (77 mL, heated in a pressure vessel at 55°C) and allowed to stand at room temperature for 20 h. The crystalline solid was filtered washed with cold dichloromethane (5 mL, 0°C) and dried under high vacuum to give pure product as a white solid (6.9 g, 68% yield, 99.79% pure by HPLC).

The preparation and purification of **C** or **C'** provides for direct access to either *S_P*-**4** or *R_P*-**4**, as illustrated in the following examples.



Example 11. Preparation of *S_P*-4**** (32 mg-scale): To a stirred solution of 1-((2R,3R,4R,5R)-3-Fluoro-4-hydroxy-5-hydroxymethyl-3-methyl-tetrahydro-furan-2-yl)-1H-pyrimidine-2,4-dione **3** (32 mg, 0.12 mmol) in dry THF (1mL) was added a 1M solution of tButylmagnesium chloride (0.26 mL, 0.26 mmol, 2.1 equiv)) at room temperature over a period of 3 min. After 30 min, a solution of (S)-2-[(S)-(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester (**8**, *S_P*-isomer) in THF (0.5mL) was added drop wise over a period of 3min. The mixture was allowed to stir at room temperature for 42h and then quenched with saturated aqueous NH₄Cl (10mL). The mixture was partitioned between ethyl acetate and water. The combined organic extract was dried over anhydrous sodium sulfate and concentrated. The residue was chromatographed using 0-4% methanol/dichloromethane gradient to give *S_P*-**4** as foamy solid (29mg, 44.5%yield). ¹H and ³¹P NMR agree to that which is disclosed herein.

Example 12. Preparation of *S_P*-4**** (2.6 g-scale, without chromatography): To a stirred solution of 1-((2R,3R,4R,5R)-3-Fluoro-4-hydroxy-5-hydroxymethyl-3-methyl-tetrahydro-furan-2-yl)-1H-pyrimidine-2,4-dione (2.6 g, 10 mmol) in dry THF (50 mL) was added a 1.7 M solution of tert-butylmagnesium chloride (12.4 mL, 21 mmol, 2.1 equiv)) at room temperature over a period of 15 min. After 30

min, a solution of (S)-2-[(S)-(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester (**8**, *S_P*-isomer, 4.08g, 10mmol) in THF (15 mL) was added drop wise over a period of 10min. The mixture was allowed to stir at room temperature for 6 0h and then quenched with saturated aqueous NH₄Cl (20mL). The mixture was partitioned between ethyl acetate (150 mL) and sequentially, 10% aqueous Na₂CO₃ (3 x 20 mL) and water (20 mL). The combined organic extract was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a pale yellow residue (3.8 g). The residue was dissolved in dichloromethane (7.6 mL) and then stirred for 20h at room temperature. The white solid was filtered, washed with 1:1 IPE/dichloromethane (5 mL) and dried under vacuum to give pure product as white solid (1.85 g, 35%yield).


Example 13. Preparation of *S_P*-4 using NaHMDS: To a stirred solution of 1-((2R,3R,4R,5R)-3-Fluoro-4-hydroxy-5-hydroxymethyl-3-methyl-tetrahydro-furan-2-yl) -1H-pyrimidine-2,4-dione (71 mg, 0.27 mmol) in dry THF (2.0 mL) was added a 2.0 M solution of sodium bis(trimethylsilyl)amide (NaHMDS) in THF (270 μ L, 0.54 mmol) at -78°C over a period of 2 min. After 30 min, a solution of (S)-2-[(S)-(4-Nitro-phenoxy)-phenoxy-phosphorylamino]-propionic acid isopropyl ester (**8**, *S_P*-isomer, 111 mg, 0.27 mmol) in THF (1 mL) was added to the mixture. The reaction mixture was allowed stir at this temperature for 2h and then warmed to -20°C at which temperature it was stirred for additional 20h. TLC indicated ~30% of unreacted nucleoside starting material. Hence, additional 0.5 equivalents of the reagent (55 mg, 0.14 mmol) in THF (0.5 mL) was added to the reaction mixture and stirred for another 6 h. The reaction mixture was quenched with saturated aqueous ammonium chloride solution and then partitioned between ethyl acetate and water. The combined organic extract was dried over anhydrous sodium sulfate and concentrated to give a light brown residue. Column chromatography of the crude product using 0-5% methanol/dichloromethane gradient gave *S_P*-4 (22 mg, 15% yield), 3'-phosphoramidate (**5**, *S_P*-isomer, 11.5 mg, 16% yield) and bis phosphoramidate (**6**, *S_P*, *S_P*-isomer, 12.6mg).

Example 14. Preparation of *R_P*-4 (260 mg-scale): To a stirred solution of 1-((2R,3R,4R,5R)-3-Fluoro-4-hydroxy-5-hydroxymethyl-3-methyl-tetrahydro-furan-2-

yl)-1H-pyrimidine-2,4-dione (260 mg, 1 mmol) in dry THF (6 mL) was added a 1.7 M solution of tert-butylmagnesium chloride (1.23 mL, 2.1 mmol, 2.1 equiv)) at room temperature over a period of 5 min. After 30 min, a solution of (S)-2-[(R)-(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester (**8**, *R_P*-isomer) in THF (3 mL) was added drop wise over a period of 3 min. The mixture was allowed to stir at room temperature for 96 h and then quenched with saturated aqueous NH₄Cl (10 mL). The mixture was partitioned between ethyl acetate (50 mL) and water (2 x 20 mL). The combined organic extract was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a pale yellow residue (490 mg). The residue was chromatographed using 0-5% methanol/dichloromethane gradient to give product as a white solid (160 mg, 30% yield).

The preparation of *S_P*-**4** or *R_P*-**4** may also be achieved by reacting 3'-protected **3** with the appropriate reagent **C** or **C'** or a mixture containing **C** and **C'**, as illustrated in the following examples.

Example 15. Preparation of *S_P*-4** with **3a** as a Synthetic Intermediate**



Example 15-1. Synthesis of 5'-O-tert-Butyldimethylsilyl-2'-deoxy-2'-fluoro-2'-C-methyluridine (9):

To a stirred solution of 2'-deoxy-2'-fluoro-2'-C-methyluridine (**3**, 81.1 g, 312 mmol) in dry pyridine (750 mL) was added drop-wise a solution of TBDMSCl (103.19 g, 685.6 mmol) in dry pyridine (500 mL) over a period of 45 min at ambient temperature. The reaction was allowed to stir at ambient temperature for 24 h. Methanol (85 mL) was added to the reaction mixture and it was allowed to stir for 10 min and then the solvents were distilled off under reduced pressure. Hot water (45°C) (1 L) was added to the reaction mass and the mixture extracted with ethyl acetate (2 x 500 mL), washed with water (1 x 500 mL). The organic layer was dried over anhydrous sodium sulfate. Ethyl acetate was distilled off and the residue obtained was co-evaporated with toluene (2 x 500 mL) to give crude **9** as a white foam. Yield = 116.9 g (quantitative). ¹H NMR (CDCl₃, 300 MHz): δ 0.1 (s, 6H), 0.91 (s, 9H), 1.22 (d, 3H, J = 21 Hz), 2.50 (s, 2H), 3.75-4.05 (m, 4H), 5.54 (d, 1H, J

= 9 Hz), 5.73 (s, 1H), 6.0 (d, 1H, J = 18 Hz), 7.81 (d, 1H, J = 9 Hz), 8.57 (br, s, 1H), 11.1 (s, 1H).

Example 15-2. Synthesis of 5'-O -(tert -Butyldimethylsilyl)-3'-O-levulinyl-2'-deoxy-2'-fluoro 2'-C-methyl-uridine (10):

To a stirred solution of nucleoside **9** (116.9 g, 312.1 mmol) in DCM (1 L) was added DMAP (30.5 g, 249.7 mmol) and this was allowed to stir at RT for 20 min. A soln. of levulinic anhydride (133.6 g, 642.3 mmol) in DCM (200 mL) was added to the mixture and allowed to stir for 24 h. TLC of the mixture indicated completion of reaction. Cold water (500 mL) was added and the mixture stirred for 20 min. Layers were separated and the organic layer was washed with sat. sodium bicarbonate solution (2 x 250 mL), dried over anhydrous sodium sulfate and then the solvent was distilled under reduced pressure to give yellow oil. Crude yield: 197.6 g (135 %). The material was used as is for the next step. ¹H NMR (CDCl₃, 300 MHz) δ 0.11 (s, 6H), 0.94 (s, 9H), 1.34 (d, 3H, J = 21 Hz), 2.22 (s, 3H), 2.6-2.89 (m, 4H), 3.72 (m, 1H), 4.01 (d, 1H, J = 12 Hz), 4.23 (d, 1H, J = 9 Hz), 5.33 (dd, 1H, J = 15 Hz), 5.73 (d, 1H, J = 6 Hz), 6.26 (d, 1H, J = 15 Hz), 8.12 (d, 1H, J = 12 Hz), 8.72 (br, s, 1H).

Example 15-3. Synthesis of 3'-O-levulinyl-2'-deoxy-2'-fluoro 2'-C-methyl-uridine (3a):

Crude **10** (197.6 g, ~312.1 mmol) was dissolved in DCM (1 L) to which was added TEA.3HF (50.3 g, 312.1 mmol) and allowed to stir overnight at ambient temperature. TLC of the mixture indicated about 50 % completion of reaction. Another equivalent of TEA.3HF (50.3 g, 312.1 mmol) was added and the reaction mixture was allowed to stir for 6 h. TLC at this point indicated about 10 % of unreacted starting material. Another 0.25 eq of TEA.3HF (12.5 g, 78.0 mmol) was added and the reaction mixture was allowed to stir overnight. Reaction mixture was concentrated to dryness to give yellow oil. Crude from all the batches was purified by column chromatography on silica gel (0-2% MeOH in DCM) to give 124.1 g of 3'-levulinate as a white foam solid (90% purified yield over three steps from 2'-deoxy-2'-fluoro-2'-C-methyluridine). ¹H NMR: (CDCl₃, 400 MHz) δ 1.55 (d, 3H, CH₃, J = 20 Hz), 2.36 (s, 3H, CH₃), 2.8-3.03 (m, 5H, CH₂CH₃), 3.91-3.96 (dd, 1H,

CH"), 4.2-4.25 (m, 1H, CH'), 4.34 (dd, 1H, CH, J = 8 Hz), 5.25 (dd, 1H, J = 16 Hz), 5.93 (d, 1H, J = 8 Hz), 8.20 (d, 1H, J = 8 Hz), 9.18 (s, 1H).

Example 15-4. Stereoselective synthesis of (S)-2-[(1R,4R,5R)-5-(2,4-Dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-4-(R)-fluoro-3-(4-oxopentanoyl)-4-methyl-tetrahydro-furan-2-ylmethoxy]-phenoxy-phosphorylamino}-propionic acid (S)-isopropyl ester (11):

To a solution of the nucleoside (**3a**, 1.00 mmol, 358 mg) in 5 ml anhydrous THF that was cooled to 0°C was added tBuMgCl (1.7 M in THF, 2 eq) and allowed it to warm to ambient temperature and stirred for half hour. To this mixture was added the reagent (ca. 97% chiral purity) (S)-2-[(S)-(4-nitro-phenoxy)-phenoxy-phosphorylamino] propionic acid isopropyl ester (**8**, *S_p*-isomer) (408 mg, 1.00 mmol, 1.00 eq.) in one lot and allowed it to stir at rt. After 16 h, there was ~30% starting material left. The reaction mixture was quenched with saturated NH₄Cl solution 10 ml, and the aqueous phase was extracted with ethyl acetate (3 x 25 ml). The combined organic layer was washed with brine and dried over anhydrous sodium sulfate and evaporated to dryness to give a pale yellow foam (500 mg). This was purified by silica gel chromatography using 2-5% methanol in methylene chloride to give the product as a white foam (275 mg) of about 97% P chiral purity and unreacted starting material (162 mg). Based on consumed starting material, the yield was 76%. ³¹P NMR (CDCl₃, 162 MHz): 3.7 ppm; ¹H NMR (CDCl₃, 400 MHz): δ 1.22 (dd, 6H, J = 6.4 Hz), 1.37 (s, 3H), 1.58 (s, 3H), 2.18 (s, 3H), 2.63-2.9 (m, 4H), 4.0 (d, 1H, J = 8 Hz), 4.2-4.33 (m, 1H), 4.57 (d, 1H, J = 8Hz), 4.96-5.00 (sept, 1H), 5.2 (dd, 1H, J = 9 Hz), 5.42 (d, 1H, J = 8Hz), 6.19 (d, 1H, J = 18Hz), 7.15-7.35 (m, 5H), 7.5 (d, 1H, J = 5.6 Hz), 8.2 (br, s, 1H).

Example 15-5. Synthesis of (S)-2-[(1R,4R,5R)-5-(2,4-Dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-4-(R)-fluoro-3-hydroxy-4-methyl-tetrahydro-furan-2-ylmethoxy]-phenoxy-phosphorylamino}-propionic acid (S)-isopropyl ester (*S_P*-4)


5 A solution of sodium sulfite was prepared by adding Na₂S₂O₃ (1.51 g) and Na₂S₂O₅ (0.57 g) in water (25 mL). To a solution of the levulinate (**11**, 250 mg, 0.40 mmol) in anhydrous THF (2.5 mL) was added 1.0 ml of the sodium sulfite solution. This was allowed to stir at room temperature for 4 h. The reaction mixture was poured in to water (15 mL) and extracted with ethyl acetate (3x25 mL) dried
10 and evaporated to give quantitatively a white solid product with about 97% P chiral purity which matched the physical and spectral properties of *S_P*-4 produced directly from the unprotected nucleoside.

Example 16. Alternative procedure for preparing *S_P*-4 from 3a.

15 To a stirred solution of 4-oxo-pentanoic acid (2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-4-fluoro-2-hydroxymethyl-4-methyl-tetrahydro-furan-3-yl ester (**3a**, 210 mg, 0.59 mmol) in dry THF (1.5 mL) was added a 1.7 M solution of tert-butylmagnesium chloride (1.07 mL, 1.82 mmol) at room temperature over a period of 2 min. Initially, a white precipitate was observed and after 10 min
20 the reaction mixture turned to dark yellow solution. After 30min, a solution of (S)-2-[(S)-(4-nitrophenoxy)-phenoxy-phosphorylamino]-propionic acid isopropyl ester (**8** (*S_P*-isomer), 382mg, 0.94mmol) in THF (1.5mL) was added drop wise over a period of 3min. The mixture was heated at 40°C for 5h at which time TLC and ¹H NMR indicated less than 2% of unreacted starting material. The reaction was quenched
25 with saturated aqueous ammonium chloride and then partitioned between ethyl acetate and water. The combined organic layer was washed with 10% aqueous Na₂CO₃ solution (3 x 10 mL), followed by water. The organic layer was dried over anhydrous sodium sulfate and concentrated to give brown color residue (410 mg). The crude product was dissolved in tetrahydrofuran (1.0 mL) and then was added an
30 aqueous solution of the mixture of sodium sulfite (37 mg, 0.295 mmol) and sodium metabisulfite (224 mg, 1.18 mmol) in 1mL of water. The mixture was heated at 45°C for 20 h at which stage only about 10% conversion was observed by TLC, hence the additional sodium sulfite (74 mg) and sodium metabisulfite (448 mg) was

added and the heating was continued for additional 52 h. At this time, about 40% conversion observed by TLC. The reaction mixture was partitioned between water and ethyl acetate. The combined organic layer was dried over anhydrous sodium sulfate and concentrated to give a brown residue (210 mg). Column chromatography of the residue using 0-5% MeOH/DCM gradient gave unreacted starting material (89mg) and *S_P*-4 (57mg, 18% yield, 24% based on recovered starting material).

Example 17. Preparation of *S_P*-4 with 3c as a Synthetic Intermediate



Example 17-1. Preparation of 1-[(2*R*,3*R*,4*R*,5*R*)-4-(tert-butyldimethylsilyloxy)-3-fluoro-5-hydroxymethyl-3-methyl-tetrahydrofuran-2-yl]-1*H*-pyrimidine-2,4-dione, 12.

To a solution of **3** (10.0g, 38.43 mmol) in pyridine (50 mL) were added dichloromethane (50 mL). The solution was cooled to 0°C. To the solution was added 4,4'-dimethoxytrityl chloride (14.32 g, 42.27 mmol) and the solution was stirred at 0°C for 5 h. Methanol (5 mL) was added to quench the reaction. The solution was concentrated to dryness under reduced pressure and the residue was partitioned between ethyl acetate (500 mL) and water (50 mL). The organic solution was washed with brine (50 mL) and dried (sodium sulfate, 4 g). The solvent was removed under reduced pressure and the residue was dissolved in dichloromethane (100 mL). To the solution were added imidazole (7.83 g, 115 mmol) and *t*-butyldimethylsilyl chloride (8.68 g, 57.6 mmol). The solution was stirred at ambient

temperature for 16 h. Methanol was added to quench the reaction (5 mL) and the solvent was removed under reduced pressure and the residue was partitioned between ethyl acetate (500 mL) and water (50 mL). The organic solution was dried (sodium sulfate, 4 g) and evaporated under reduced pressure. The residue was
5 purified by column chromatography (10-40% EtOAc in Hexane) to give 5'-O-DMT-3'-O-tBDMS intermediate product. This is turn was treated with 1% trifluoroacetic acid in dichloromethane (200 mL). The solution was stirred at ambient temperature for 1h. Water (20 mL) was added and the solution was stirred at ambient for another 1h. Methanol (5 mL) was slowly added and the solution was stirred at ambient for
10 another 1h. Ammonium hydroxide was added to adjust the solution pH to 7. The organic solution was separated, dried (sodium sulfate, 4 g) and evaporated to dryness under reduced pressure. The residue was purified by silica gel column chromatography (1-5% methanol in dichloromethane) to give **12** as a white solid 7.5g in 50% yield over the three steps. ¹H NMR (DMSO-d₆) δ (ppm) 11.48 (br s, 1H, NH), 7.94 (d, 1H, H-6), 6.00 (d, 1H, H-1'), 5.69 (d, 1H, H-5), 4.06 (dd, 1H, 3'-H), 3.85 (m, 2H, H-5'a, H-4'), 3.58 (br d, 1H, H-5'b), 1.27 (d, 3 H, 2-CH₃), 0.89 (s, 9H, C(CH₃)₃), 0.12 (s, 6H, Si(CH₃)₂).

Example 17-2. Preparation of S_P-4 using 1-[(2R,3R,4R,5R)-4-(tert-butyltrimethylsilyloxy)-3-fluoro-5-hydroxymethyl-3-methyl-tetrahydro-furan-2-yl]-1H-pyrimidine-2,4-dione (3c).


To a stirred solution of 1-[(2R,3R,4R,5R)-4-(tert-butyltrimethylsilyloxy)-3-fluoro-5-hydroxymethyl-3-methyl-tetrahydro-furan-2-yl]-1H-pyrimidine-2,4-dione (**12**, 374 mg, 1 mmol) in dry THF (3 mL) was added a 1.7 M solution of tert-butylmagnesium chloride (1.8 mL, 3.1 mmol) at room temperature over a period of
25 2 min. Initially, a white precipitate was observed and after 10min the reaction mixture turned to clear dark yellow solution. After 30 min, a solution of (S)-2-[(S)-(4-nitrophenoxy)-phenoxy-phosphorylamino]-propionic acid isopropyl ester (**8**, S_P-isomer, 653 mg, 1.6 mmol) in THF (2.5 mL) was added drop wise over a period of 3
30 min. The mixture was heated at 40°C for 20 h at which time TLC and ¹H NMR indicated less than 5% of unreacted starting material. The reaction mixture was quenched with saturated aqueous ammonium chloride and then partitioned between ethyl acetate and water. The organic layer was washed with 10% aqueous Na₂CO₃

solution (3 x 10 mL), followed by water (20 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated to give brown residue containing **3c** (850 mg). The crude product was dissolved in tetrahydrofuran (2 mL) and was added 0.8 mL of 80% aqueous formic acid at room temperature. The reaction mixture was

5 heated at 50°C for 96h. About 70% conversion was observed by TLC. The reaction mixture was poured into cold saturated aqueous sodium bicarbonate and then partitioned between ethyl acetate and water. The combined organic layer was dried over anhydrous sodium sulfate and concentrated to give brown residue (220 mg). Column chromatography of the residue using 0-5% MeOH/DCM gradient gave

10 unreacted starting material (21 mg) and *S_P*-**4** (77 mg, 35% yield, 39% yield based on recovered starting material).

Example 18. Preparation of *S_P*-**4** with **3d** as a Synthetic Intermediate



Example 18-1. Preparation of 3d

To a stirred solution of **3** in pyridine (20 mL) at 0°C was added TIPDS-Cl drop-wise over a period of 15 min. The mixture was slowly allowed to warm to room temperature at which temperature it was stirred for 16 h. The pyridine was
5 evaporated and the residue was co-evaporated with toluene (50 mL). The residue was then triturated with hexanes and the white precipitate was filtered off using a pad of Celite. The filtrate was concentrated under reduced pressure to give a foamy solid (12.97 g). The crude product (**13**) was redissolved in tetrahydrofuran (75mL) and was added an aqueous solution of TFA (75mL, 1:1 TFA/water) at 0°C over a
10 period of 20 min. The mixture was stirred at this temperature for 6 h. TLC indicated ~5% of starting material. The reaction mixture was quenched with saturated aqueous NaHCO₃ until pH 8 and then extracted with ethyl acetate. The combined organic extract was washed with water, dried and concentrated to give white crystalline solid. Further trituration of the solid with hexanes (30 mL) gave white solid which
15 was filtered and dried under high vacuum to give **3d** (10.1 g, 84 % yield over 2 steps). ¹H NMR (400 MHz, CDCl₃): δ 8.83 (bs, 1H), 7.94 (bd, J=6.0Hz, 1H), 6.10(bd, J=18.4Hz, 1H), 5.71 (d, J=8.2Hz, 1H), 4.43 (bs, 1H), 4.36 (dd, J=22.6, 9.0Hz, 1H), 4.27 (bs, 1H), 4.10(d, J=13.2Hz, 1H), 4.03 (d, J=9.2Hz, 1H), 3.92 (d, J=13.2Hz, 1H), 1.39 (d, J=22.0Hz, 3H), 1.11-0.92 (m, 28H).

20


Example 18-2. Preparation of S_P-4

To a stirred solution of **3d** (520 mg, 1 mmol) in dry THF (5 mL) was added a 1.7M solution of tert-butyilmagnesium chloride (1.8 mL, 3.1 mmol, 3.1 equiv)) at room temperature over a period of 15 min. After 30 min, a solution of (S)-2-[(S)-(4-nitro-phenoxy)-phenoxyphosphorylamino] propionic acid isopropyl ester (**8**, S_P-
25 isomer, 653 mg, 1.6 mmol) in THF (1 mL) was added drop wise over a period of 3 min. The mixture was allowed to stir at room temperature for 60 h. ¹H and ³¹P NMR of the crude sample indicated mixture of diastereomers in about 1:0.76. The reaction mixture was quenched with saturated aqueous NH₄Cl (20 mL). The mixture was
30 partitioned between ethyl acetate (150 mL) and sequentially, 10% aqueous Na₂CO₃ (3 x 20mL) and water (20 mL). The combined organic extract was dried over

anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a pale yellow residue (**14**, 878 mg).

The above compound, **14**, was redissolved in tetrahydrofuran (3 mL) and then was added 80% aqueous formic acid. The mixture was heated at 55°C for 20h. The reaction mixture was cooled to 0°C, and then quenched with saturated aqueous sodium bicarbonate (pH 7.0). The reaction mixture was then partitioned between ethyl acetate and water. The combined organic layer was dried over sodium sulfate and concentrated to give 560mg of the residue. The residue was chromatographed using 0-5% methanol/dichloromethane gradient to give unreacted starting material (**14**, 242mg) and *S_P*-**4** (80mg, 15% yield) as a white solid.

Example 19. Preparation of Isotopically Labeled *S_P*-**4**



Example 19-1. Preparation of 1-((6aR,8R,9R,9aS)-9-hydroxy-2,2,4,4-tetraisopropyltetrahydro-6H-furo[3,2-f][1,3,5,2,4]trioxadisilocin-8-yl)pyrimidine-2,4(1H,3H)-dione, 16

Uridine (**15**, 100.0 g, 409.5 mmol) was co-evaporated to dryness with
5 anhydrous pyridine (600 mL) and re-suspended in anhydrous pyridine (700 mL). To
this stirred fine suspension was added 1,3-dichloro-1,1,3,3-tetraisopropylidisiloxane
(135.7 g, 482.5 mmol) over 60 min at ambient temperature. After stirring the fine
suspension for 17 h at ambient temperature, the reaction was quenched by adding
methanol (20 mL) and then concentrated under reduced pressure. The residue was
10 partitioned between ethyl acetate (1.5 L) and water (2 L). The organic layer was
further washed with 5% hydrochloric acid (2 x 1 L), brine (500 mL), dried over solid
sodium sulfate (50 g), filtered and concentrated under reduced pressure to the crude
product, ca 250 g. The residue was subjected to a filtration column using silica gel
(1.75 kg) and a gradient of ethyl acetate in hexanes 20-65%. The pure product
15 fractions as judged by a homogenous TLC (R_f 0.55 in 1:1 hexanes-ethyl acetate)
were combined and concentrated under reduced pressure and dried (40°C, 0.2 mm
Hg, 24 h) to afford 145.5 g (76%) of **16** as a white foam solid. An additional
fraction (35 g) of slightly impure **16** was also collected. ^1H NMR (DMSO- d_6) δ
(ppm) 11.35 (s, 1H, NH), 7.66 (d, 1H, J = 7.6 Hz, H -6), 5.57 (d, 1H, J = 4.8 Hz, 2'-
20 OH), 5.50-5.49 (m, 2H, 1'- H and H -5), 4.14-4.18 (m, 3H, 2', 3', 4'- H), 3.97-3.87 (m,
2H, 5'- H_a and H_b), 1.02-0.95 (m, 28H, $\text{CH}(\text{CH}_3)_2$).

Example 19-2. Preparation of 1-((6aR,8R,9aR)-2,2,4,4-tetraisopropyl-9-oxotetrahydro-6H-furo[3,2-f][1,3,5,2,4]trioxadisilocin-8-yl)pyrimidine-2,4(1H,3H)-dione, 17

To a dry three-necked round flask were added anhydrous DCM (600 mL)
and DMSO (30.82 g, 394.5 mmol). The solution was cooled to -78°C in a dry
ice/acetone bath under an atmosphere of nitrogen. Trifluoroacetic anhydride (neat,
30 77.7 g, 369.8 mmol) was added via a syringe over 40 mins and afforded a cloudy
mixture. To the mixture a solution of uridine derivative **16** in DCM (600 mL) was
added dropwise over 75 mins at -78°C via an addition funnel. The heterogeneous
mixture was stirred for 2 h at -78~-65°C and then anhydrous triethylamine (92 mL)

was added via a syringe quickly to form a clear light yellow solution. After 1 h at low temperature, the reaction was complete as shown by TLC (30% EtOAc in hexanes). The cooling bath was removed and the reaction mixture was warmed up slowly to ambient temperature over 1 h. The reaction was quenched by addition of
5 sat. NH_4Cl (180 mL). Water (200 mL) was added and organic layer was separated. The aqueous layer was extracted again with DCM (300 mL). The combined organic layer was washed with water (3x 400 mL), brine (150 mL), and dried over Na_2SO_4 . Removal of solvent afforded a sticky brown residue.

The crude oil residue (contained trace of DCM) was stored overnight in the
10 freezer. After overnight, some crystal solid was observed in the oil. The oil was dissolved in 500 ml hexanes at ambient temperature. The solution was stored in the freezer for 24 hours and more solid was formed. Solid was collected via filtration and rinsed with cold 10% DCM in hexanes (1 L) to remove most of the orange color. The solid (**17**) was dried under vacuum for 2 h and then air dried for 24 h.
15 The solid weighed 21 g after dried at 50 °C under vacuum. The filtrate was concentrated and the residue was purified via column chromatography (10-70% ethyl acetate in hexanes) to afford an additional 37 g (combined yield of 97%) of **17** as a light orange solid.

20 **Example 19-3. Preparation of 1-((2R,3S,4R,5R)-3,4-dihydroxy-5-(hydroxymethyl)-3- ^{13}C -perdeuteriomethyltetrahydrofuran-2-yl)pyrimidine-2,4(1H,3H)-dione, **18****

Magnesium (3.53 g, 147 mmol), washed with 5% aqueous hydrochloric acid and dried (50°C, 0.2mm Hg, 24 h), was put into a two neck round bottomed flask
25 equipped with a magnetic stirrer and a condenser. The flask was filled with argon gas and then anhydrous ether (80 mL) was added. To the magnesium in ether was added slowly perdeuterio- ^{13}C methyl iodide (15.06 g, 110.3 mmol), which generated an exothermic reaction. After the reaction mixture was cooled down, the supernatant was transferred to a solution of dried compound **17** (50°C, 0.2mm Hg,
30 15 h) (10.0 g, 20.63 mmol) in anhydrous THF (1 L) at -50°C over 20 min. The temperature was allowed to rise to -40°C and the mixture was stirred at between -40 to -25°C for 4h. Upon completion of reaction, the mixture was diluted with EtOAc (1L) at -50 °C and then brine (300

mL) was added slowly. The organic layer was separated and then washed with sat'd ammonium chloride solution (300 mL x 2) and dried with sodium sulfate. After filtration and concentration under reduced pressure, the residue was dissolved in MeOH (250 mL). Ammonium fluoride (12 g) and TBAF (400 mg) were added.

- 5 The resulting mixture was stirred at 90°C for 7h and then concentrated with silica gel (20 g) under reduced pressure. After thorough vacuum drying, the obtained residue was purified by flash silica gel column chromatography (MeOH:CH₂Cl₂ = 1:20 to 1:10) give compound **18** (5 g, 46%) as a white solid. ¹H NMR (DMSO-*d*₆) δ (ppm) 11.26 (s, 1H, NH), 7.65 (d, 1H, *J* = 8.4 Hz, H-6), 5.77 (d, 1H, *J* = 2.4 Hz, H-1'), 5.57 (d, 1H, *J* = 8.0 Hz, H-5), 5.46 (d, 1H, *J* = 5.2 Hz, HO-3'), 5.24 (d, 1H, *J* = 2.4 Hz, HO-2'), 5.14 (t, 1H, *J* = 5.6 Hz, HO-5'), 3.74-3.56 (m, 4H, H-3', 4', 5', 5'').

Example 19-4. Preparation of ((2R,3R,4S,5R)-3-acetoxy-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-hydroxy-4-¹³C-perdeuteriomethyltetrahydrofuran-2-yl)methyl acetate, **19**

- To a solution of compound **18** (5.00 g, 19.1 mmol) in anhydrous pyridine (100 mL) was added acetic anhydride (3 mL) at ambient temperature. The resulting mixture was stirred at ambient temperature for 15h, diluted with EtOAc (250 mL),
20 washed with water (50 mL x 3), and dried with sodium sulfate. After filtration and concentration, the residue was purified by flash column chromatography (MeOH 0 to 5% in CH₂Cl₂) to give compound **19** (4.0 g, 68%) as a gray solid.

Example 19-5. Preparation of ((2R,3R,4R,5R)-3-acetoxy-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-4-¹³C-perdeuteriomethyltetrahydrofuran-2-yl)methyl acetate, **20**


- To a solution of compound **19** (2.33 g, 6.73 mmol) in anhydrous CH₂Cl₂ (60 mL) was added DAST (1.33 mL, 10.1 mmol) at -78°C slowly. The resulting mixture was stirred for 30 min after exposed to ambient temperature. An additional
30 two 2.33 g scale reactions and one 1.00 g scale reaction were conducted exactly the same way. All four reaction mixtures were combined, diluted with CH₂Cl₂ (300 mL), and washed with ice-water (100 mL x 2) and then cold aqueous NaHCO₃ solution (100 mL x 2). After drying, filtration, and concentration, the residue was

purified by flash silica gel column chromatography (EtOAc 0% to 50% in hexanes, compound came out at around 48%) to give compound **20** (2.0 g, 24% from total 7.99 g of compound **19**) as a white solid. ¹H NMR (CDCl₃) δ (ppm) 8.27 (s, 1H, NH), 7.55 (d, 1H, *J* = 8.4 Hz, H-6), 6.17 (d, 1H, *J* = 18.8 Hz, H-1'), 5.78 (dd, 1H, *J* = 1.2, 8.4 Hz, H-5), 5.12 (dd, 1H, *J* = 9.6, 21.6 Hz, H-3'), 4.40-4.31 (m, 3H, H-4', 5', 5''), 2.19 (s, 3H, CH₃), 2.15 (s, 3H, CH₃).

Example 19-6. Preparation of 1-((2R,3R,4R,5R)-3-fluoro-4-hydroxy-5-(hydroxymethyl)-3-¹³C-perdeuteriomethyltetrahydrofuran-2-yl)pyrimidine-2,4(1H,3H)-dione, **21**

To a solution of compound **20** (2 g, 5.74 mmol) in methanol (20 mL) was added n-butylamine (6 mL). The resulting mixture was stirred at rt for 15h and concentrated with silica gel *in vacuo*. The obtained residue was purified by flash silica gel column chromatography (MeOH 0 to 10% in CH₂Cl₂) to give compound **21** (1.3 g, 85%) as a white solid. ¹H NMR (CD₃OD) δ (ppm) 8.08 (d, 1H, *J* = 8.0 Hz, H-6), 6.13 (d, 1H, *J* = 18.4 Hz, H-1'), 5.70 (d, 1H, *J* = 8.0 Hz, H-5), 3.99 (d, 1H, *J* = 13.6 Hz, H-5'), 3.97-3.91 (m, 2H, H-3', 4'), 3.80 (dd, 1H, *J* = 2.0, 12.8 Hz, H-5''), ESMS (M+1) estimated 265, observed 265.

20



Example 19-7. Preparation of (S)-Isopropyl 2-((((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-¹³C-perdeuteriomethyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphorylamino)propanoate, **22**

To a solution of the unprotected nucleoside **21** (207 mg, 0.783 mmol) and *N*-methylimidazole (0.4 mL, 5 mmol) in THF (4 mL) was added the pre-made phosphorochloridate in THF (1.0 M, 2.35 mL, 2.35 mmol) at 0°C dropwise. The reaction was slowly warmed to ambient temperature over 1 h and then water (1 mL)

and EtOAc (5 mL) were added. The organic solution was washed with sat. aq. mono basic sodium citrate (2 x 2 ml), sat. aq. NaHCO₃ (1 x 2 ml), dried (MgSO₄) and concentrated under reduced pressure. The crude was purified by silica column chromatography using 0 to 5% ⁱPrOH in CH₂Cl₂ as eluents to give the

- 5 phosphoramidate, **22** (216 mg, 52%, 1:1 mixture of P-diastereomers) as a white solid: ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.54 (s, 1H), 7.56 (d, *J* = 6.8 Hz, 1H), 7.40-7.35 (m, 2H), 7.23-7.18 (m, 3 H), 6.14-5.96 (m, 2H), 5.89 (dd, *J* = 5.6, 25.6 Hz, 1H), 5.55 (t, *J* = 8.4 Hz, 1H), 4.85 (dq, *J* = 1.6, 6.0 Hz, 1H), 4.44-4.32 (m, 1H), 4.25 (m, 1H), 4.06-3.98 (m, 1H), 3.86-3.70 (m, 2H), 1.30-1.08 (m, 9H); ³¹P NMR (162
10 MHz, DMSO-*d*₆) δ 4.90, 4.77; LRMS (ESI) [M + H]⁺ calculated for C₂₁¹³CH₂₇D₃FN₃O₉P 534.5, found 534.4.

- Example 19-8. Preparation of (2S)-2-((((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-¹³C-perdeuteriomethyltetrahydrofuran-2-yl)methoxy)(hydroxy)phosphoryl)amino)propanoic acid, **23****
15

- Phosphoramidate **22** (147 mg, 0.276 mmol) was suspended in triethylamine (2 mL) and water (0.5 mL), and heated at 60°C for 30 h. Then the volatile
20 components were evaporated under reduced pressure. The crude was purified by silica column chromatography by eluting with 50-70% ⁱPrOH in CH₂Cl₂ and then, 0 to 20% NH₄OH in ⁱPrOH to give **23** as a white solid (95 mg, 83%): ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.00 (d, *J* = 8.4 Hz, 1H), 5.98 (d, *J* = 19.2 Hz, 1H), 5.52 (d, *J* = 8.4 Hz, 1H), 4.02-3.81 (m, 4H), 1.10 (d, *J* = 6.8 Hz, 3H); ³¹P NMR (162 MHz,
25 DMSO-*d*₆) δ 8.12; LRMS (ESI) [M + H]⁺ calculated for C₁₂¹³CH₁₇D₃FN₃O₉P 416.3, found 416.4.

Properties of Samples of R_p-4, 4, and S_p-4

- 30 Samples of R_p-4, 4, and S_p-4 were analyzed by X-Ray Powder Diffraction (XRPD), Nuclear Magnetic Resonance (NMR) spectrometry, Fourier Transform Infrared (FT-IR) spectroscopy, Differential Scanning Calorimetry (DSC), Thermal Gravimetric Analysis (TGA), Gravimetric Vapor Sorption (GVS), Thermodynamic Aqueous Solubility, and High Performance Liquid Chromatography (HPLC).

35

Example 20. X-Ray Powder Diffraction

Samples of *R_p-4*, *4*, and *S_p-4* were analyzed by X-Ray Powder Diffraction (XRPD) under the following regimen.

5 **a. Bruker AXS/Siemens D5000**

X-Ray Powder Diffraction patterns were collected on a Siemens D5000 diffractometer using Cu K α radiation (40kV, 40mA), θ - θ goniometer, divergence of V20 and receiving slits, a graphite secondary monochromator and a scintillation
10 counter. The instrument is performance checked using a certified Corundum standard (NIST 1976). The software used for data collection was Diffrac Plus XRPD Commander v2.3.1 and the data were analyzed and presented using Diffrac *Plus* EVA v 11.0.0.2 or v 13.0.0.2.

15 Ambient conditions

Samples run under ambient conditions were prepared as flat plate specimens using powder as received. Approximately 35 mg of the sample was gently packed into a cavity cut into polished, zero-background (510) silicon wafer. The sample was
20 rotated in its own plane during analysis. The details of the data collection are: angular range: 2 to 42°2 θ ; step size: 0.05°2 θ ; and collection time: 4 s.step⁻¹.

b. Bruker AXS C2 GADDS

25 X-Ray Powder Diffraction patterns were collected on a Bruker AXS C2 GADDS diffractometer using Cu K α radiation (40 kV, 40 mA), automated XYZ stage, laser video microscope for auto-sample positioning and a HiStar 2-dimensional area detector. X-ray optics consists of a single Göbel multilayer mirror coupled with a pinhole collimator of 0.3 mm.

30

The beam divergence, i.e. the effective size of the X-ray beam on the sample, was approximately 4 mm. A θ - θ continuous scan mode was employed with a sample - detector distance of 20 cm which gives an effective 2 θ range of 3.2°– 29.7°. Typically the sample would be exposed to the X-ray beam for 120 seconds. The

software used for data collection was GADDS for WNT 4.1.16 and the data were analyzed and presented using *DiffraC Plus* EVA v 9.0.0.2 or v 13.0.0.2.

Ambient conditions

5

Samples run under ambient conditions were prepared as flat plate specimens using powder as received without grinding. Approximately 1-2 mg of the sample was lightly pressed on a glass slide to obtain a flat surface.

10 X-Ray Powder Diffraction (XRPD)

4 was found by XRPD to be amorphous (see **Fig. 1**). High resolution XRPD analysis of *R_P-4* prepared according to Example 3 confirmed a crystalline solid exhibiting a different powder pattern to that of *S_P-4* (prepared according to Example 4, Method 4), which was also confirmed to be a crystalline solid. The XRPD results table for *R_P-4* and *S_P-4* are shown in **Table 1** with all peaks exhibiting an intensity of $\leq 5\%$ (*R_P-4*) and $\leq 3\%$ (*S_P-4*) excluded.

Table 1. XRPD Data for *R_P-4* and *S_P-4*.

XRPD data for <i>R_P-4</i>		XRPD data for <i>S_P-4</i> (Form 1)	
Angle 2-Theta °	Intensity %	Angle 2-Theta °	Intensity %
6.616	51.1	4.900	6.8
7.106	40.5	5.190	19.8
8.980	30.0	7.501	100.0
11.020	21.7	8.355	4.1
11.559	77.1	8.965	7.7
11.950	12.8	9.619	21.2
13.023	5.2	10.145	3.6
14.099	6.2	14.393	4.9
15.121	5.7	16.300	7.0
15.624	5.4	16.688	10.6
16.003	17.8	17.408	5.5
17.882	100.0	17.820	8.2
18.567	8.8	18.262	31.5
19.564	22.7	18.600	6.3
20.280	5.6	18.900	7.3
20.728	42.5	19.150	6.1
21.047	19.9	19.696	4.8
21.671	22.0	20.398	4.4
21.943	23.3	20.710	6.9
22.214	18.9	21.950	6.1

XRPD data for <i>R_p</i> -4		XRPD data for <i>S_p</i> -4(Form 1)	
Angle 2-Theta °	Intensity %	Angle 2-Theta °	Intensity %
23.074	28.5	22.175	12.2
24.145	30.3	22.511	5.6
24.355	39.1	22.924	3.1
25.366	7.6	23.360	6.5
26.146	36.2	23.538	7.1
27.000	9.0	23.910	7.4
27.313	15.6	24.873	3.7
27.677	22.7	25.123	4.9
28.219	12.8	25.649	4.2
28.661	6.2	26.748	5.2
29.450	6.8	27.339	3.7
29.735	9.4	27.646	3.5
31.372	8.2	28.066	3.1
31.967	10.9	29.050	3.0
32.295	6.4	29.541	3.6
33.001	11.4	30.178	3.8
33.774	11.8	31.648	3.1
34.385	6.6	32.721	3.5
34.734	6.5	33.154	3.0
35.600	7.3	33.923	3.5
35.965	13.1	34.341	3.1
36.409	14.7	35.465	3.5
36.880	7.0	36.923	3.1
37.509	5.9	37.760	3.4
37.870	6.0	38.404	3.3
38.313	5.8	40.416	3.1
38.943	8.4		
40.093	6.6		
40.511	7.8		
41.429	6.5		

A sample of *S_p*-4 was ground with a pestle and mortar, and then successively passed through 500 and 250 µm sieves to yield the sample as a fine powder. This sample was reanalyzed by high resolution XRPD, confirming no form change had occurred.

Example 21. Crystallization Studies for *S_p*-4.

Crystalline *S_p*-4 exhibits polymorphism. Thus, an aspect is directed to crystalline *S_p*-4 and its individual polymorphic forms. *S_p*-4 can exist in at least five polymorphic forms, designated as Forms 1-5. Furthermore, amorphous *S_p*-4 can also be prepared. A typical crystallization provides for dissolving about 100 mg of

5 *S_P-4* in an appropriate volume of crystallization solvent (acetonitrile (5 vol), chloroform (5 vol), n-butyl acetate (7 vol), dichloromethane (50 vol), anisole (7 vol), and 1:1 MTBE/heptane (50 vol)) and then allowing for evaporation of the solution at 5 °C. Various crystalline forms were obtained, but each form, upon filtration and/or drying, afforded Form 1.

Forms 1, 2 and 3 are a non-solvated form, 1:1 DCM solvate and 1:1 chloroform solvate, respectively, as was confirmed by single crystal X-ray and XRPD analysis. Forms 4 and 5 were obtained from crystallization of *S_P-4* from solutions of acetonitrile and anisole, respectively. Sufficient data could not be collected to determine whether Forms 4 and 5 are unsolvated, hydrated or solvated since single crystals of sufficient quality were not obtained. Forms 4 and 5 transform to Form 1 on filtration. Two additional crystalline forms are obtained upon crystallization of *S_P-4* from n-butyl acetate (*n*BuAc) and a solution containing methyl-*t*-butyl ether (MTBE) and heptane; upon filtration both of these crystalline forms convert to Form 1. Forms 2 and 3 also transform to Form 1 on isolation. Form 1 is a non-solvated form that exhibits a broad melting endotherm with an onset temperature of 94.3°C and ΔH_{fus} of 24.0 kJ mol⁻¹. An additional XRPD pattern of *S_P-4* Form 1 is depicted in **Figure 4**.

20 **Conversion of *S_P-4*, Form 1 to *S_P-4*, Form 6**

Form 1 can be converted to Form 6 in at least two ways.

First, by allowing fine crystals of Form 1 to be exposed to atmospheric humidity for several days, a monohydrate of Form 1 is produced with an appearance of a solidified gum. After grinding the monohydrate solid to a fine powder, the XRPD pattern remains consistent with Form 1. Upon standing in an open vessel for 6-10 weeks, the ground material slowly changes to Form 6 as an anhydrous solid. Form 1 is stable for at least 2 years in a sealed container.

Alternatively, Form 1 can be suspended in water 5-50 mg/mL at ambient temperature and over a few hours be transformed into Form 6. The water transformation process efficiency can be improved by heating the water to a point to dissolve more Form 1 and to increase the fluidity of the immiscible portion of *S_P-4* from a stiff gum to a suspended oil at about or above. Over time, Form 6 can begin to crystallize out at 50°C. Further cooling the suspension to 0-10°C leads to a higher

recovery of solid. Crystallization from water also removes more polar trace impurities, leading to improved overall purity.

Redissolving Form 6 into an organic solvent such as dichloromethane or acetonitrile followed by crystallization provides Form 1, even when seeding with crystalline Form 6.

To a dry 100 mL one-neck round bottom flask equipped with an rubber septum and magnetic stir bar was charged 1.04 grams of *Sp*-4, Form 1. HPLC purity 99.7%. Charged 40 mL of DI water. Began stirring suspension vigorously while heating to 50°C. Once the temperature reached 50°C, the mostly homogenous solution was held for 60 mins, during which time solids began to precipitate from solution, forming a thin slurry. The slurry was cooled to 20°C over 90 mins and held for 16 hours at 20°C, followed by cooling further to 0-5°C over 30 mins and held at 0-5°C for 2.5 hours. The slurry was filtered on a medium-porosity glass fritted funnel and washed with 10 mL of ice-chilled water. The wet cake was suction dried for 2 hours before drying in a vacuum oven overnight for 23 hours at 50°C. Isolated 0.88 g (84.6% recovery) of *Sp*-4, Form 6.

Form 6 has an observed melting point of about 124.5-126°C.

Example 21-1. *Sp*-4 Form 1

A peak listing of *Sp*-4 Form 1 is presented in Table 2.

Angle	Intensity %
2-Theta °	%
5.0	74.8
7.3	100.0
7.8	2.2
8.2	6.8
8.8	9.3
9.4	23.5
10.0	8.4
11.4	4.2
13.3	3.0
14.2	6.1
14.9	3.5
16.1	7.9
16.6	13.2

Angle 2-Theta °	Intensity %
17.3	15.4
17.7	10.1
18.1	42.6
18.4	7.6
18.7	11.4
18.9	5.7
19.3	5.0
19.6	2.9
20.2	8.5
20.5	11.5
20.8	3.6
21.8	7.2
22.0	14.5
22.4	9.6
23.2	5.3
23.4	5.8
23.5	4.6
23.8	7.4
24.0	3.1
24.7	2.5
25.0	13.0
25.5	3.1
26.6	4.5
27.2	3.2
27.5	2.2
28.1	2.9
30.0	3.2

Example 21-2. *S_P*-4 Form 2

An XRPD pattern of *S_P*-4 Form 2 is depicted in **Figure 5**.

5

A peak listing of *S_P*-4 Form 2 is presented in **Table 3**.

Angle 2-Theta °	Intensity %
4.9	44.1
5.1	19.1

Angle 2-Theta °	Intensity %
6.9	62.1
8.7	6.8
9.8	28.6
10.1	10.4
13.7	7.0
16.7	3.1
19.5	8.9
19.8	45.5
20.1	18.6
20.4	3.6
20.6	25.6
20.9	15.9
21.1	10.9
22.1	3.4
24.6	38.7
24.7	100.0
25.1	61.2
26.1	53.3
39.0	6.3

Example 21-3. Sp-4 Form 3

An XRPD pattern of Sp-4 Form 3 is depicted in **Figure 6**.

5

A peak listing of Sp-4 Form 3 is presented in **Table 4**.

Angle 2-Theta °	Intensity %
5.0	10.0
6.9	23.3
9.8	22.6
19.7	34.8
20.6	100.0
21.8	10.5
24.6	65.3
34.7	4.1

Example 21-4. Sp-5 Form 4

An XRPD pattern of Sp-4 Form 4 is depicted in **Figure 7**.

- 5 A peak listing of Sp-4 Form 4 is presented in **Table 5**.

Angle	Intensity %
2-Theta °	%
5.0	29.8
6.8	100.0
8.2	4.8
8.7	5.2
9.9	3.8
13.7	1.7
14.9	4.8
19.9	22.5
20.4	2.1
20.6	20.0
20.9	20.0
24.7	3.4
24.9	29.9
25.1	1.5
36.8	3.1

Example 21-5. Sp-4 Form 5

- 10 An XRPD pattern of Sp-4 Form 5 is depicted in **Figure 8**.

A peak listing of Sp-4 Form 5 is presented in **Table 6**.

Angle	Intensity %
2-Theta °	%
5.2	52.9
6.6	100.0
7.1	25.9
9.7	12.1
10.4	16.4
13.4	11.4

Angle 2-Theta °	Intensity %
15.7	25.8
19.1	31.1
19.9	12.9
20.0	9.0
21.3	3.5
25.0	22.3
25.6	2.3
26.3	5.9
26.9	2.0
31.7	2.1

Example 21-5. *S_P*-4 Form 6

An XRPD pattern of *S_P*-4 Form 6 is depicted in Figure 21.

5

A peak listing of *S_P*-4 Form 6 is presented in following table.

Angle 2-Theta °	d-spacing Å	Intensity %
6.08	14.51	66.7
8.2	10.77	62.1
10.38	8.52	29.8
10.85	8.14	10.4
12.17	7.26	12.0
12.7	6.96	66.4
13.73	6.44	14.9
14.1	6.27	13.8
15.91	5.57	3.1
16.83	5.26	8.7
17.17	5.16	19.7
17.66	5.01	56.2
17.95	4.93	37.7
18.79	4.72	59.0
19.1	4.64	14.3
19.41	4.57	37.2
19.8	4.48	46.0
20.11	4.41	68.8
20.82	4.26	100.0
21.81	4.07	36.8
22.03	4.03	7.4
23.03	3.86	14.2

Angle 2-Theta °	d-spacing Å	Intensity %
23.26	3.82	21.6
23.64	3.76	6.3
23.89	3.72	7.0
24.73	3.6	3.3

Example 21-7. *S_P*-4 (Amorphous)

An XRPD pattern for amorphous *S_P*-4 is depicted in **Figure 9**.

5

Example 22. Single Crystal X-ray Crystallography of *S_P*-4 and its solvates

Example 22-1. Single Crystal X-Ray Crystallography of *S_P*-4 (Form 1)

10 Figure 10 shows an X-ray crystal structure for *S_P*-4 Form 1. There, the figure shows a view of molecules of Form 1 from the crystal structure showing the numbering scheme employed. Anisotropic atomic displacement ellipsoids for the non-hydrogen atoms are shown at the 50% probability level. Hydrogen atoms are displayed with an arbitrarily small radius.

15 The structure solution was obtained by direct methods, full-matrix least-squares refinement on F^2 with weighting $w^{-1} = \sigma^2(F_o^2) + (0.0592P)^2 + (0.6950P)$, where $P = (F_o^2 + 2F_c^2)/3$, anisotropic displacement parameters, empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm. Final $wR^2 = \{\Sigma[w(F_o^2 - F_c^2)^2]/\Sigma[w(F_o^2)^2]\}^{1/2} = 0.0871$ for all data,
 20 conventional $R_1 = 0.0329$ on F values of 7090 reflections with $F_o > 4\sigma(F_o)$, $S = 1.016$ for all data and 870 parameters. Final $\Delta/\sigma(\text{max})$ 0.001, $\Delta/\sigma(\text{mean})$, 0.000. Final difference map between +0.534 and -0.36 e Å⁻³.

Table 7. Single Crystal Parameters of Form 1

25

Molecular formula	C ₂₂ H ₂₉ F ₁ N ₃ O ₉ P ₁				
Molecular weight	529.45				
Crystal system	Monoclinic				
Space group	$P2_1$	a	20.0898(5)Å,	α	90°,
		b	6.10290(10)Å,	β	112.290(3)°,
		c	23.0138(6)Å,	γ	90°
V	2610.79(10)Å ³				

Z	4
D_c	1.347g.cm^{-1}
μ	1.475mm^{-1}
Source, λ	Cu K α , $1.54178\text{\AA}$
$F(000)$	1112
T	100(1)K
Crystal	Colorless plate, $0.12 \times 0.09 \times 0.03\text{mm}$
Data truncated to	$0.80\text{\AA}$
θ_{max}	74.48°
Completeness	99.4%
Reflections	14854
Unique reflections	7513
R_{int}	0.0217

Example 22-2. Single Crystal X-Ray Crystallography of S_P -4 (Form 2)

Figure 11 shows an X-ray crystal structure for S_P -4 Form 2. There, this figure shows a view of molecules of Form 2 from the crystal structure showing the numbering scheme employed. The heteroatoms were resolved isotropically due to very weak data. Hydrogen atoms are not displayed.

The structure solution was obtained by direct methods, full-matrix least-squares refinement on F^2 with weighting $w^{-1} = \sigma^2(F_o^2) + (0.0975P)^2 + (10.6969P)$, where $P = (F_o^2 + 2F_c^2)/3$, anisotropic displacement parameters, empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm. Final $wR^2 = \{\Sigma[w(F_o^2 - F_c^2)^2]/\Sigma[w(F_o^2)^2]\}^{1/2} = 0.1883$ for all data, conventional $R_1 = 0.0741$ on F values of 2525 reflections with $F_o > 4\sigma(F_o)$, $S = 1.05$ for all data and 158 parameters. Final $\Delta/\sigma(\text{max})$ 0.000, $\Delta/\sigma(\text{mean})$, 0.000. Final difference map between $+1.388$ and $-0.967\text{ e \AA}^{-3}$.

Table 8. Single Crystal Parameters of Form 2

Molecular formula	$\text{C}_{23}\text{H}_{31}\text{Cl}_2\text{FN}_3\text{O}_9\text{P}$				
Molecular weight	614.38				
Crystal system	Monoclinic				
Space group	$P2_1$	a	$12.8315(3)\text{\AA}$,	α	90° ,
		b	$6.14530(10)\text{\AA}$,	β	$91.752(2)^\circ$,
		c	$17.6250(4)\text{\AA}$,	γ	90°
V	$1389.14(5)\text{\AA}^3$				
Z	2				

D_c	1.469g.cm ⁻¹
μ	3.196mm ⁻¹
Source, λ	Cu-K, 1.54178Å
$F(000)$	640
T	293(2)K
Data truncated to	0.80 Å
$\theta_{\max}$	62.23°
Completeness	91.1%
Reflections	3528
Unique reflections	2562
R_{int}	0.0227

Example 22-3. Single Crystal X-Ray Crystallography of *S_P-4* (Form 2)

Figure 12 depicts an X-ray Crystal Structure (ORTEP – anisotropic) *S_P-4* (Form 2). A crystal structure of the methylene chloride solvate of *S_P-4* (Form 2), C₂₃H₃₁N₃PO₉FCl₂, yields a monoclinic space group P2₁ (systematic absences 0k0: k=odd) with a=12.8822(14) Å, b=6.1690(7) Å, c=17.733(2) Å, β =92.045(3)°, V=1408.4(3)Å³, Z=2 and d_{calc} =1.449 g/cm³. X-ray intensity data were collected on a Rigaku Mercury CCD area detector employing graphite-monochromated Mo-K α radiation (λ =0.71073 Å) at a temperature of 143K. Preliminary indexing was performed from a series of twelve 0.5° rotation images with exposures of 30 seconds. A total of 648 rotation images were collected with a crystal to detector distance of 35 mm, a 2 θ swing angle of -12°, rotation widths of 0.5° and exposures of 30 seconds: scan no. 1 was a ϕ -scan from 315° to 525° at ω = 10° and χ = 20°; scan no. 2 was an ω -scan from -20° to 5° at χ = -90° and ϕ = 315°; scan no. 3 was an ω -scan from -20° to 4° at χ = -90° and ϕ = 135°; scan no. 4 was an ω -scan from -20° to 5° at χ = -90° and ϕ = 225°; scan no. 5 was an ω -scan from -20° to 20° at χ = -90° and ϕ = 45°. Rotation images were processed using CrystalClear (CrystalClear: Rigaku Corporation, 1999), producing a listing of unaveraged F^2 and $\sigma(F^2)$ values which were then passed to the CrystalStructure (CrystalStructure: Crystal Structure Analysis Package, Rigaku Corp. Rigaku/MS (2002)) program package for further processing and structure solution on a Dell Pentium III computer. A total of 7707 reflections were measured over the ranges $5.48 \leq 2\theta \leq 50.04^\circ$, $-14 \leq h \leq 15$, $-7 \leq k \leq 6$, $-19 \leq l \leq 21$ yielding 4253 unique reflections (R_{int} = 0.0180). The intensity data

were corrected for Lorentz and polarization effects and for absorption using REQAB (minimum and maximum transmission 0.824, 1.000).

The structure was solved by direct methods (SIR97, SIR97: Altomare, A., M. Burla, M. Camalli, G. Cascarano, C. Giacovazzo, A. Guagliardi, A. Moliterni, G. Polidori & R. Spagna (1999). *J. Appl. Cryst.*, **32**, 115-119). Refinement was by full-matrix least squares based on F^2 using SHELXL-97 (SHELXL-97: Sheldrick, G.M. (2008) *Acta Cryst.*, **A64**, 112-122). All reflections were used during refinement. The weighting scheme used was $w=1/[\sigma^2(F_o^2) + 0.0472P^2 + 0.4960P]$ where $P = (F_o^2 + 2F_c^2)/3$. Non-hydrogen atoms were refined anisotropically and hydrogen atoms were refined using a "riding" model. Refinement converged to $R_1=0.0328$ and $wR_2=0.0817$ for 4046 reflections for which $F > 4\sigma(F)$ and $R_1=0.0348$, $wR_2=0.0838$ and $GOF = 1.056$ for all 4253 unique, non-zero reflections and 358 variables ($R_1 = \sum ||F_o| - |F_c|| / \sum |F_o|$; $wR_2 = \{ \sum w (F_o^2 - F_c^2)^2 / \sum w(F_o^2)^2 \}^{1/2}$; $GOF = \{ \sum w (F_o^2 - F_c^2)^2 / (n - p) \}^{1/2}$; where n = the number of reflections and p = the number of parameters refined). The maximum Δ/σ in the final cycle of least squares was 0.000 and the two most prominent peaks in the final difference Fourier were +0.312 and -0.389 e/Å³. The Flack absolute structure parameter refined to -0.06(6) thus corroborating the stereochemistry of the title compound.

Table 1 lists cell information, data collection parameters, and refinement data. Final positional and equivalent isotropic thermal parameters are given in Table 2. Anisotropic thermal parameters are in Table 3. ("ORTEP-II: A Fortran Thermal Ellipsoid Plot Program for Crystal Structure Illustrations". C.K. Johnson (1976) ORNL-5138.) representation of the molecule with 30% probability thermal ellipsoids displayed.

25

Table 9. Summary of Structure Determination of Compound $S_p\text{-4}\cdot\text{CH}_2\text{Cl}_2$.

Formula:	$\text{C}_{23}\text{H}_{31}\text{N}_3\text{PO}_9\text{FCl}_2$
Formula weight:	614.38
Crystal class:	monoclinic
Space group:	$P2_1$ (#4)
Z	2
Cell constants:	
a	12.8822(14) Å

b	6.1690(7) Å
c	17.733(2) Å
β	92.045(3)°
V	1408.4(3) Å ³
μ	3.48 cm ⁻¹
crystal size, mm	0.42 x 0.12 x 0.10
D _{calc}	1.449 g/cm ³
F(000)	640
Radiation:	Mo-K α (λ =0.71073 Å)
2 θ range	5.48 – 50.04 °
hkl collected:	-14 $\leq$ h $\leq$ 15; -7 $\leq$ k $\leq$ 6; -19 $\leq$ l $\leq$ 21
No. reflections measured:	7707
No. unique reflections:	4253 (R _{int} =0.0180)
No. observed reflections	4046 (F>4 σ)
No. reflections used in refinement	4253
No. parameters	358
R indices (F>4 σ)	R ₁ =0.0328 wR ₂ =0.0817
R indices (all data)	R ₁ =0.0348 wR ₂ =0.0838
GOF:	1.056
Final Difference Peaks, e/Å ³	+0.312, -0.389

Example 22-4. Single Crystal X-Ray Crystallography of Sp-4 (Form 3)

5 Figure 13 shows an X-ray crystal structure for Sp-4 Form 3. There, this figure shows a view of molecules of Form 3 from the crystal structure showing the numbering scheme employed. Anisotropic atomic displacement ellipsoids for the non-hydrogen atoms are shown at the 50% probability level. Hydrogen atoms are displayed with an arbitrarily small radius.

10 The structure solution was obtained by direct methods, full-matrix least-squares refinement on F^2 with weighting $w^{-1} = \sigma^2(F_o^2) + (0.0512P)^2 + (0.6810P)$, where $P = (F_o^2 + 2F_c^2)/3$, anisotropic displacement parameters, empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm. Final $wR^2 = \{\Sigma[w(F_o^2 - F_c^2)^2]/\Sigma[w(F_o^2)^2]\}^{1/2} = 0.0796$ for all data,

15 conventional $R_1 = 0.0294$ on F values of 2486 reflections with $F_o > 4\sigma(F_o)$, $S = 1.068$ for all data and 377 parameters. Final $\Delta/\sigma(\text{max})$ 0.001, $\Delta/\sigma(\text{mean})$, 0.000. Final difference map between +0.211 and -0.334 e Å⁻³.

Table 10. Single Crystal Parameters of Form 3

Molecular formula	$C_{23}H_{30}Cl_3F_1N_3O_9P_1$				
Molecular weight	648.82				
Crystal system	Monoclinic				
Space group	$P2_1$	a	12.9257(4)Å,	α	90°,
		b	6.18080(10)Å,	β	96.399(2)°,
		c	18.0134(4)Å,	γ	90°
V	1430.15(6)Å ³				
Z	2				
D_c	1.507g.cm ⁻¹				
μ	3.977mm ⁻¹				
Source, λ	Cu K α , 1.54178Å				
$F(000)$	672				
T	100(1)K				
Crystal	Colorless needle, 0.22 x 0.03 x 0.02mm				
Data truncated to	0.80 Å				
θ_{max}	74.41°				
Completeness	69.1%				
Reflections	3062				
Unique reflections	2607				
R_{int}	0.0198				

5 Example 23. Stability at elevated temperatures and relative humidity

A sample of **Rp-4** was stored in a humidity chamber at 40°C and 75% relative humidity for one week, and the sample was reanalyzed by XRPD. The powder pattern obtained for **Rp-4** showed no substantial change during the course of the experiment, meaning that no change in solid form was observed. This should be contrasted to a sample of **4**, which deliquesced within about 16 hours upon storage at 40°C and 75 % relative humidity. Indeed, an illustration of the deliquescent nature of **4** is illustrated by the following. A sample of **4** was passed through a 250 μ m sieve then samples were stored at 40°C / 75 % RH and 25°C / 53 % relative humidity and visual observations were taken at regular intervals. The results are given in **Table 4**.

Table 11. Stability of **4** to elevated relative humidity.

Conditions	t = 1.5 h	t = 4.5 h	t = 6.5 h	t = 8.5 h	t = 73 h
40°C / 75%RH	Deliquescence	-	-	-	-
25°C / 53%RH	No deliquescence	Sticky solid	Partial deliquescence	Almost complete deliquescence	Deliquescence

Upon storage at 40 °C and 75 % relative humidity a sample of *S_P-4* deliquesced inside 16 hours. For instance, a sample of *S_P-4* was ground with a pestle and mortar, and then successively passed through 500 and 250 μm sieves to yield the sample as a fine powder. Samples of this material were stored at 40°C and 75% relative humidity and 25°C and 53 % RH and visual observations were taken at regular intervals. The results are given in **Table 5**.

10

Table 12. Stability of *S_P-4* to elevated relative humidity.

Conditions	t = 1.5 h	t = 4.5 h	t = 104 h
40 °C / 75 % RH	No deliquescence	Deliquescence	-
25 °C / 53 % RH	No deliquescence	No deliquescence	No deliquescence

XRPD analysis of the sample after storage at 25°C and 53% RH for 104 hours showed no significant changes in the diffractograms produced indicating that no form change had occurred.

Example 24. Fourier Transform – Infrared (FT-IR) Spectrometry

Data were collected on a Perkin-Elmer Spectrum One fitted with a universal Attenuated Total Reflectance (ATR) sampling accessory. The data were collected and analyzed using Spectrum v5.0.1 software.

The IR spectrum obtained for **4**, *R_P-4*, and *S_P-4* are shown in Figs. 5–7, respectively. Selected peaks, in wavenumbers (cm^{-1}) are recited below:

4: ~1680, ~1454, ~1376, ~1205, ~1092, ~1023 (Fig. 14);
R_P-4: ~1742, ~1713, ~1679, ~1460, ~1377, ~1259, ~1157, ~1079 (Fig. 15);
 and

Sp-4 (Form 1): ~1743, ~1713, ~1688, ~1454, ~1378, ~1208, ~1082 (Fig. 16).

Example 25. Differential Scanning Calorimetry (DSC) Thermo-Gravimetric Analysis (TGA)

DSC data were collected on a TA Instruments Q2000 equipped with a 50 position auto-sampler. The calibration for thermal capacity was carried out using sapphire and the calibration for energy and temperature was carried out using certified indium.

Modulated temperature DSC was carried out on typically 0.8-1.2 mg of each sample, in a pin-holed aluminum pan, using an underlying heating rate of $2^{\circ}\text{C}.\text{min}^{-1}$ and temperature modulation parameters of $\pm 0.2^{\circ}\text{C}.\text{min}^{-1}$ and 40 seconds. A purge of dry nitrogen at $50 \text{ ml}.\text{min}^{-1}$ was maintained over the sample.

The instrument control software was Advantage for Q Series v2.8.0.392 and Thermal Advantage v4.8.3 and the data were analyzed using Universal Analysis v4.3A.

DSC data were collected on a Mettler DSC 823e equipped with a 34 position auto-sampler. The instrument was calibrated for energy and temperature using certified indium. Typically 0.8-1.2 mg of each sample, in a pin-holed aluminum pan, was heated at $10^{\circ}\text{C}.\text{min}^{-1}$ from 25°C to 250°C . A nitrogen purge at $50 \text{ ml}.\text{min}^{-1}$ was maintained over the sample. The instrument control and data analysis software was STARe v9.20.

DSC data for Sp-4 (Form 6) were collected using a DSC instrument (TA Q2000), using a heating rate of $10^{\circ}\text{C}/\text{min}$ under a continuous flow of dry nitrogen gas ($100 \text{ ml}/\text{min}$). Approximately 2.2 mg of sample was accurately weighed and heated in a non-hermetically sealed, 'Tzero' pan with a loose-fitting lid. The instrument was calibrated (enthalpy and temperature) with an indium standard and (heat capacity) with a sapphire standard. Uncertainties are estimated to be $\pm 0.1^{\circ}\text{C}$ for temperatures and $\pm 5\%$ for measured enthalpies. TA Universal Analysis software was employed to measure onset temperatures.

TGA data were collected on a Mettler TGA/SDTA 851e equipped with a 34 position auto-sampler. The instrument was temperature calibrated using certified indium. Typically 8-12 mg of each sample was loaded onto a pre-weighed aluminum

crucible and was heated at $10^{\circ}\text{C}.\text{min}^{-1}$ from ambient temperature to 350°C . A nitrogen purge at $50 \text{ ml}.\text{min}^{-1}$ was maintained over the sample. The instrument control and data analysis software was STARe v9.20.

DSC analysis of **4** showed a single broad endotherm with an onset of 58.7°C ($\Delta H 14 \text{ J}.\text{g}^{-1}$) confirmed to be due to molecular relaxation during the glass transition by further modulated DSC analysis (Fig. 17). TGA analysis of **4** showed no weight loss before decomposition above 240°C , confirming the material to be non-solvated. As the XRPD analysis of **4** confirmed the material to be amorphous, modulated DSC analysis was undertaken in an attempt to calculate the glass transition temperature, which was found to be 57°C .

DSC analysis showed a single sharp endotherm with an onset of 136.2°C ($\Delta H 76 \text{ J}.\text{g}^{-1}$) confirmed to be a melt by hot stage microscopy. See Fig. 18. TGA analysis of *Rp-4* showed no weight loss before decomposition above 240°C , confirming the material to be non-solvated.

DSC analysis of *Sp-4* showed a single broad endotherm with an onset of 93.9°C ($\Delta H 43 \text{ J}.\text{g}^{-1}$) confirmed to a melt by hot stage microscopy. See Fig. 19. TGA analysis of *Sp-4* showed no weight loss before decomposition above 240°C , confirming the material to be non-solvated.

DSC analysis of *Sp-4* (Form 6) showed a broad endotherm with an onset of 120.7°C ($\Delta H 79 \text{ J}.\text{g}^{-1}$).

Example 26. Gravimetric Vapour Sorption (GVS)

SMS DVS Intrinsic

Sorption isotherms were obtained using a SMS DVS Intrinsic moisture sorption analyzer, controlled by SMS Analysis Suite software. The sample temperature was maintained at 25°C by the instrument controls. The humidity was controlled by mixing streams of dry and wet nitrogen, with a total flow rate of $200 \text{ ml}.\text{min}^{-1}$. The relative humidity was measured by a calibrated Rotronic probe (dynamic range of 1.0-100 %RH), located near the sample. The weight change, (mass relaxation) of the sample as a function of %RH was constantly monitored by the microbalance (accuracy $\pm 0.005 \text{ mg}$).

Typically 5-20 mg of sample was placed in a tared mesh stainless steel basket under ambient conditions. The sample was loaded and unloaded at 40 %RH

and 25°C (typical room conditions). A moisture sorption isotherm was performed as outlined below (2 scans giving 1 complete cycle). The standard isotherm was performed at 25°C at 10 %RH intervals over a 0.5-90 %RH range.

5 **Table 13. Method Parameters for SMS DVS Intrinsic Experiments**

Parameters	Values
Adsorption - Scan 1	40 - 90
Desorption / Adsorption - Scan 2	90 - 0, 0 - 40
Intervals (%RH)	10
Number of Scans	2
Flow rate (ml.min ⁻¹)	200
Temperature (°C)	25
Stability (°C.min ⁻¹)	0.2
Sorption Time (hours)	6 hour time out

The sample was recovered after completion of the isotherm and re-analyzed by XRPD.

10 GVS analysis showed *R_p-4* to be non-hygroscopic exhibiting reversible uptake of approximately 0.2 wt % of water from 0 to 90 % relative humidity. Re-analysis of the sample by XRPD after the GVS experiment showed no change in form.

15 A sample of *S_p-4* was ground with a pestle and mortar, and then successively passed through 500 and 250 µm sieves to yield the sample as a fine powder that was then analyzed using a modified single cycle method. The sample was taken from 40 % RH (approximately ambient) to 60 % RH, instead of 90 % for the standard method, and then cycled to 0 % and back to 40 % RH. This analysis showed *S_p-4* to be non-hygroscopic up to 60 % RH, with reversible uptake of ~0.2 % by weight of
20 water from 0 to 60 % RH.

Example 27. Thermodynamic Aqueous Solubility

25 Aqueous solubility was determined by suspending a sufficient amount of compound in water to give a maximum final concentration of ≥ 10 mg.ml⁻¹ of the parent free-form of the compound. The suspension was equilibrated at 25 °C for 24 hours then the pH was measured. The suspension was then filtered through a glass fiber C filter into a 96 well plate. The filtrate was then diluted by a factor of 101. Quantitation was by HPLC with reference to a standard solution of approximately

0.1 mg.ml⁻¹ in DMSO. Different volumes of the standard, diluted and undiluted sample solutions were injected. The solubility was calculated using the peak areas determined by integration of the peak found at the same retention time as the principal peak in the standard injection.

5

Table 14. HPLC Method Parameters for Solubility Measurements

Type of method:	Reverse phase with gradient elution		
Column:	Phenomenex Luna, C18 (2) 5µm 50 x 4.6 mm		
Column Temperature (°C):	25		
Standard Injections (µl):	1, 2, 3, 5, 7, 10		
Test Injections (µl):	1, 2, 3, 10, 20, 50		
Detection: Wavelength, Bandwidth (nm) :	260,80		
Flow Rate (ml.min ⁻¹):	2		
Phase A:	0.1% TFA in water		
Phase B:	0.085% TFA in acetonitrile		
Timetable:	Time (min)	% Phase A	% Phase B
	0.0	95	5
	1.0	80	20
	2.3	5	95
	3.3	5	95
	3.5	95	5
	4.4	95	5

Analysis was performed under the above-noted conditions on an Agilent
 10 HP1100 series system equipped with a diode array detector and using ChemStation
 software vB.02.01-SR1.

Table 15. Aqueous solubility result for *R_P*-4, 4, and *S_P*-4.

Sample ID	pH of Unfiltered mixture	Solubility ₁ / mg.ml ⁻¹	Comments
<i>R_P</i> -4	7.12	1.58	Suspension
4	7.03	6.11	Residual solid
<i>S_P</i> -4	6.88	5.65	Residual solid

15

Example 28. Chemical Purity Determination by HPLC

Various HPLC conditions can be used to determine the chemical purity of the compounds disclosed herein. One such example is disclosed above in relation to the thermodynamic aqueous solubility studies. Another example is disclosed below.

HPLC Conditions:

LC: Waters Alliance 2695 Separations Module, Waters 2996 PDA detector and Waters Empower 2 Software (Version 6.00)

10 **Column:** Phenomenex Luna C18(2); 4.6 x 50mm; 3 μ m

Flow rate: 1.2 mL/min

Injection Volume: 10 μ L

Mobile phase: Solvent A: 95% Water with 5% Methanol and 10mM Ammonium Acetate; pH~5.3

15 **Gradient:** Solvent B: MeOH with 10mM Ammonium Acetate

hold at 0%B	3 min
0-47%B	3-4 min
hold at 47 %B	4-10 min
47%-74%B	10-11min
20 hold at 74%B	11-13.5min
return to 0%B	13.5-13.6 min
hold at 0%B	13.6-15.5 min

Under these conditions, the purity of **4**, *R_p*-**4**, and *S_p*-**4** was determined to be ~99.6, ~99%, and ~99.5%, respectively. It is noted that higher purities can be realized by optimizing the methods disclosed above.

Inspection of the XRPD diffractograms shows that the two crystalline single diastereoisomers gave clearly different XRPD patterns. Additionally, there was a clear difference in the melting point of the two crystalline diastereoisomers, with *R_p*-**4** having a considerably higher onset than *S_p*-**4** (136°C vs. 94°C).

Example 29. Additional Separation Methods

The following SFC separation (conditions listed below) yielded adequate separation of a mixture of the diastereomers, *R_p*-**4** and *S_p*-**4**.

Preparative Method:	Analytical Method:
Chiralpak AS-H (2 x 25 cm) SN# 07-8656	Chiralpak AS-H (25 x 0.46cm)
20% methanol/CO ₂ (100 bar)	20% methanol/CO ₂ (100 bar)
50 ml/min, 220 nm.	3 ml/min, 220 nm.
Conc.: 260mg/30ml methanol, inj vol.: 1.5ml	

The following SFC separation (conditions listed below) yielded adequate separation of a mixture of the diastereomers, *R_P*-4 and *S_P*-4.

Preparative Method:	Analytical Method:
Chiralpak IA(2 x 15 cm) 802091	Chiralpak IA(15 x 0.46 cm)
30% isopropanol(0.1% DEA)/CO ₂ , 100 bar	40% methanol(DEA)/CO ₂ , 100 bar
60 mL/min, 220 nm.	3 mL/min, 220 nm.
inj vol.: 2 mL, 20 mg/mL methanol	

5 **Table 16. Summary of results from the batch characterization of *R_P*-4, 4, and *S_P*-4.**

Analysis	<i>R_P</i> -4	4	<i>S_P</i> -4
Proton NMR	Single diastereoisomer	1:1 Mixture of diastereoisomers	Single diastereoisomer
XRPD	Crystalline – different from <i>S_P</i> -4	Amorphous	Crystalline – different from <i>R_P</i> -4
DSC	Endotherm; melt - 136°C	Endotherm; 59°C	Endotherm; melt - 94°C
TGA	No wt loss, decomposition >240°C	No wt loss, decomposition >240°C	No wt loss, decomposition >240°C
IR	See above	See above	See above
Aq Solubility (mg.mL ⁻¹)	1.58	6.11	5.65
HPLC Purity	96.9 %	99.6 %	99.5 %
40°C / 75 % RH	No form change	Deliquescence inside 1.5 h	Deliquescence inside 4.5 h
25°C / 53 % RH	-	Deliquescence	No form change
GVS	Non-hygroscopic up to 90 % RH	-	Non-hygroscopic up to 60 % RH

10 **Example 30. X-Ray Crystallography of 8 (*S_P*-isomer)**

Compound **8** (*S_P*-isomer), C₁₈H₂₁N₂PO₇, crystallizes in the monoclinic space group P2₁ (systematic absences 0k0: k=odd) with a=5.3312(4)Å, b=15.3388(8)Å, c=23.7807(13)Å, β=92.891(3)°, V=1942.2(2)Å³, Z=4, and d_{calc}=1.397 g/cm³. X-ray intensity data were collected on a Bruker APEXII CCD area detector employing
 15 graphite-monochromated Mo-Kα radiation (λ=0.71073 Å) at a temperature of 100(1)K. Figures 20A and 20B show molecules numbered 1 and 2, respectively, of the asymmetric unit.

Preliminary indexing was performed from a series of thirty-six 0.5° rotation frames with exposures of 30 seconds. A total of 3608 frames were collected with a
 20 crystal to detector distance of 70.00 mm, rotation widths of 0.5° and exposures of 20 seconds:

scan type	2 θ	ω	φ	χ	frames
φ	-35.50	279.40	27.32	48.96	725
φ	24.50	22.31	35.56	69.08	692
ω	-13.00	321.68	247.79	69.08	95
φ	34.50	204.08	28.21	-92.80	293
φ	-30.50	310.60	214.10	54.21	361
φ	32.00	304.67	24.47	50.72	722
φ	-35.50	122.14	316.59	-78.84	720

Rotation frames were integrated using SAINT (Bruker (2009) SAINT.

Bruker AXS Inc., Madison, Wisconsin, USA.) producing a listing of unaveraged F^2 and $\sigma(F^2)$ values which were then passed to the SHELXTL (Bruker (2009)

- 5 SHELXTL. Bruker AXS Inc., Madison, Wisconsin, USA.) program package for further processing and structure solution on a Dell Pentium 4 computer. A total of 6909 reflections were measured over the ranges $1.58 \leq \theta \leq 25.09^\circ$, $-6 \leq h \leq 6$, $-18 \leq k \leq 18$, $-28 \leq l \leq 28$ yielding 6909 unique reflections ($R_{int} = 0.0581$). The intensity data were corrected for Lorentz and polarization effects and for absorption using
- 10 SADABS (Sheldrick, G.M. (2007) SADABS. University of Gottingen, Germany.) (minimum and maximum transmission 0.6093, 0.7452).

The structure was solved by direct methods (SHELXS-97 (Sheldrick, G.M. (2008) *Acta Cryst.* A64,112-122.)). Refinement was by full-matrix least squares based on F^2 using SHELXL-97 (Sheldrick, G.M. (2008) *Acta Cryst.* A64, 112-122.).

- 15 All reflections were used during refinement. The weighting scheme used was $w=1/[\sigma^2(F_o^2) + (0.0000P)^2 + 14.0738P]$ where $P = (F_o^2 + 2F_c^2)/3$. Non-hydrogen atoms were refined anisotropically and hydrogen atoms were refined using a riding model. Refinement converged to $R1 = 0.0847$ and $wR2 = 0.1899$ for 6173 observed reflections for which $F > 4\sigma(F)$ and $R1=0.0963$ and $wR2=0.1963$ and $GOF=1.119$
- 20 for all 6909 unique, non-zero reflections and 512 variables ($R1 = \sum ||F_o| - |F_c|| / \sum |F_o|$; $wR2 = [\sum w(F_o^2 - F_c^2)^2 / \sum w(F_o^2)^2]^{1/2}$; $GOF = [\sum w(F_o^2 - F_c^2)^2 / (n - p)]^{1/2}$; where n = the number of reflections and p = the number of parameters refined). The maximum Δ/σ in the final cycle of least squares was 0.000 and the two most prominent peaks in the final difference Fourier were +0.402 and -0.559 $e/\text{\AA}^3$. **Figs 20A and 20B** are
- 25 ORTEP (30% probability thermal ellipsoids) of molecules 1 and 2 of the asymmetric unit.

Table 17. Summary of Structure Determination of Compound 8 (*S_p*-isomer)

Empirical formula	C ₁₈ H ₂₁ N ₂ PO ₇
Formula weight	408.34
Temperature	100(1) K
Wavelength	0.71073 Å
Crystal system	monoclinic
Space group	P2 ₁
Cell constants:	
a	5.3312(4) Å
b	15.3388(8) Å
c	23.7807(13) Å
β	92.891(3)°
Volume	1942.2(2) Å ³
Z	4
Density (calculated)	1.397 Mg/m ³
Absorption coefficient	0.185 mm ⁻¹
F(000)	856
Crystal size	0.40 x 0.10 x 0.08 mm ³
Theta range for data collection	1.58 to 25.09°
Index ranges	-6 ≤ h ≤ 6, -18 ≤ k ≤ 18, -28 ≤ l ≤ 28
Reflections collected	6909
Independent reflections	6909 [R(int) = 0.0581]
Completeness to theta = 25.09°	99.6 %
Absorption correction	Semi-empirical from equivalents
Max. and min. transmission	0.7452 and 0.6093
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	6909 / 1 / 512
Goodness-of-fit on F ²	1.119
Final R indices [I > 2σ(I)]	R1 = 0.0847, wR2 = 0.1899
R indices (all data)	R1 = 0.0963, wR2 = 0.1963
Absolute structure parameter	0.1(2)
Largest diff. peak and hole	0.402 and -0.559 e.Å ⁻³

Example 32. X-Ray Crystallography of (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate

(S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)-
 5 amino)propanoate, C₁₈H₁₇NPO₅F₅, crystallizes in the triclinic space group P1 with a
 = 5.2641(6) Å, b = 12.0548(13) Å, c = 16.4307(15) Å, α = 74.960(4)°, β =
 83.959(4)°, γ = 80.275(4)°, V = 990.40(18) Å³, Z = 2, and d_{calc} = 1.520 g/cm³. X-ray
 intensity data were collected on a Bruker APEXII CCD area detector employing
 graphite-monochromated Mo-Kα radiation (λ = 0.71073 Å) at a temperature of
 10 143(1)K. Preliminary indexing was performed from a series of thirty-six 0.5°
 rotation frames with exposures of 20 seconds. A total of 3593 frames were collected
 with a crystal to detector distance of 37.600 mm, rotation widths of 0.5° and
 exposures of 20 seconds:

scan type	2θ	ω	φ	χ	frames
φ	-15.50	258.48	-351.72	19.46	739
φ	-20.50	-17.45	-37.67	-73.06	725
ω	-10.50	-53.05	-87.93	99.72	80
φ	19.50	-32.21	-88.94	36.30	219
ω	-10.50	-14.33	80.80	-60.33	122
ω	17.00	-38.90	-41.64	83.36	116
ω	17.00	-37.89	-175.56	82.07	114
φ	19.50	59.55	-11.29	-26.26	739
φ	-10.50	318.39	-335.56	52.47	739

15

Rotation frames were integrated using SAINT (Bruker (2009) SAINT. Bruker AXS
 Inc., Madison, Wisconsin, USA.), producing a listing of unaveraged F² and σ(F²)
 values which were then passed to the SHELXTL (Bruker (2009) SHELXTL. Bruker
 AXS Inc., Madison, Wisconsin, USA.) program package for further processing and
 20 structure solution on a Dell Pentium 4 computer. A total of 17880 reflections were
 measured over the ranges 1.77 ≤ θ ≤ 25.12°, -6 ≤ h ≤ 6, -14 ≤ k ≤ 14, -19 ≤ l ≤ 19
 yielding 6897 unique reflections (Rint = 0.0212). The intensity data were corrected
 for Lorentz and polarization effects and for absorption using SADABS (Sheldrick,
 G.M. (2007) SADABS. University of Gottingen, Germany.) (minimum and
 25 maximum transmission 0.6887, 0.7452).

The structure was solved by direct methods (SHELXS-97(Sheldrick, G.M. (2008)
Acta Cryst. A64,112-122.)). Refinement was by full-matrix least squares based on

- F^2 using SHELXL-97 (Sheldrick, G.M. (2008) *Acta Cryst.* A64,112-122.). All reflections were used during refinement. The weighting scheme used was $w=1/[\sigma^2(F_o^2) + (0.0344P)^2 + 0.1102P]$ where $P = (F_o^2 + 2F_c^2)/3$. Non-hydrogen atoms were refined anisotropically and hydrogen atoms were refined using a riding model. Refinement converged to $R1=0.0259$ and $wR2=0.0609$ for 6527 observed reflections for which $F > 4\sigma(F)$ and $R1 = 0.0284$ and $wR2 = 0.0621$ and $GOF = 1.040$ for all 6897 unique, non-zero reflections and 548 variables. ($R1 = \Sigma||F_o| - |F_c|| / \Sigma |F_o|$; $wR2 = [\Sigma w(F_o^2 - F_c^2)^2 / \Sigma w(F_o^2)^2]^{1/2}$; $GOF = [\Sigma w(F_o^2 - F_c^2)^2 / (n - p)]^{1/2}$; where n = the number of reflections and p = the number of parameters refined.) The maximum Δ/σ in the final cycle of least squares was 0.001 and the two most prominent peaks in the final difference Fourier were +0.254 and -0.236 e/Å³. **Figs. 22A and 22B** are ORTEP (30% probability thermal ellipsoids) of molecules 1 and 2 of the asymmetric unit.
- Table 18. Summary of Structure Determination of (S)-isopropyl 2-(((S)-(perfluorophenoxy)(phenoxy)phosphoryl)amino)propanoate**

Empirical formula	C ₁₈ H ₁₇ NPO ₅ F ₅
Formula weight	453.30
Temperature	143(1) K
Wavelength	0.71073 Å
Crystal system	triclinic
Space group	P1
Cell constants:	
a	5.2641(6) Å
b	12.0548(13) Å
c	16.4307(15) Å
α	74.960(4)°
β	83.959(4)°
γ	80.275(4)°
Volume	990.40(18) Å ³
Z	2
Density (calculated)	1.520 Mg/m ³
Absorption coefficient	0.216 mm ⁻¹
F(000)	464

Crystal size	0.45 x 0.05 x 0.04 mm ³
Theta range for data collection	1.77 to 25.12°
Index ranges	-6 ≤ h ≤ 6, -14 ≤ k ≤ 14, -19 ≤ l ≤ 19
Reflections collected	17880
Independent reflections	6897 [R(int) = 0.0212]
Completeness to theta = 25.12°	99.5 %
Absorption correction	Semi-empirical from equivalents
Max. and min. transmission	0.7452 and 0.6887
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	6897 / 3 / 548
Goodness-of-fit on F ²	1.040
Final R indices [I > 2σ(I)]	R1 = 0.0259, wR2 = 0.0609
R indices (all data)	R1 = 0.0284, wR2 = 0.0621
Absolute structure parameter	-0.01(5)
Largest diff. peak and hole	0.254 and -0.236 e.Å ⁻³

Example 33. Biological Activity

- 5 Replicon containing cells were seeded at either 3,000 cells/well (50 μL) in 96-well white/opaque plates, or 1,500 cells/well (25 μL) in 384-well white/opaque plates. 50 μL of 2X compound were added in the 96 well plate or 25 μL of 2X compound were added in the 384 well plate. The plates were incubated at 37°C in a humidified 5% CO₂ atmosphere for 4 days. After incubation, Bright-Glo reagent (50
10 μL for 96-well plate, or 25 μL for 384-well plate) was added to measure the firefly luciferase reporter for HCV replication. Percent inhibition was calculated against the no drug control.

Compound	HCV Replicon Activity (μM)
4	0.58
<i>R_p</i> - 4	2.87
<i>S_p</i> - 4	0.13

- 15 *R_p*-**4** and *S_p*-**4** have been demonstrated to have broad genotype coverage. For example, both have been shown to be active against hepatitis C virus, genotypes 1-4.

This application is a continuation-in-part of U.S. Patent Application No. 12/783,680 filed on May 20, 2010, which claims priority to U.S. Provisional Patent


Application Nos. 61/179,923, filed May 20, 2009, and 61/319,513, filed March 31, 2010, the subject matter of which is incorporated by reference in its entirety.

The subject matter of U.S. Patent Application Nos. 12/783,680 and 12/053,015 and U.S. Provisional Patent Application Nos. 61/179,923, filed May 20, 5 2009, and 61/319,513, filed March 31, 2010, is hereby incorporated by reference in its entirety. The subject matter of all cited references is hereby incorporated by reference. In the event that the meaning of an incorporated term conflicts with the meaning of a term defined herein, the meaning of the terms contained in the present disclosure control over the meaning of the incorporated terms.

10

CLAIMS

1. Crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate represented by the formula S_{P-4} :



having XRPD 2 θ -reflections ($\pm 0.2^\circ$) at 6.1 and 12.7.

2. The crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate of claim 1 further having XRPD 2 θ -reflections ($\pm 0.2^\circ$) at 8.2, 10.4, 17.2, 17.7, 18.0, 18.8, 19.4, 19.8, 20.1, 20.8, 21.8, and 23.3.

3. The crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate of claim 1 or 2 having XRPD 2 θ -reflections ($\pm 0.2^\circ$) at 6.08, 8.2, 10.38, 10.85, 12.17, 12.7, 13.73, 14.1, 15.91, 16.83, 17.17, 17.66, 17.95, 18.79, 19.1, 19.41, 19.8, 20.11, 20.82, 21.81, 22.03, 23.03, 23.26, 23.64, 23.89, and 24.73.

4. The crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate of claim 3, wherein the XRPD 2 θ -reflections ($\pm 0.2^\circ$) at 6.08, 8.2, 10.38, 10.85, 12.17, 12.7, 13.73, 14.1, 15.91, 16.83, 17.17, 17.66, 17.95, 18.79, 19.1, 19.41, 19.8, 20.11, 20.82, 21.81, 22.03, 23.03, 23.26, 23.64, 23.89, and 24.73 have intensities (%) of 66.7, 62.1, 29.8, 10.4, 12.0, 66.4, 14.9, 13.8, 3.1, 8.7, 19.7, 56.2, 37.7, 59.0, 14.3, 37.2, 46.0, 68.8, 100.0, 36.8, 7.4, 14.2, 21.6, 6.3, 7.0, and 3.3, respectively.

5. A composition comprising crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate of any one of claims 1-4.

6. A pharmaceutical composition comprising crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate of any one of claims 1-4 and a pharmaceutically acceptable medium.

7. Crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate of any one of claims 1-4 for use in the treatment of a hepatitis C virus infection.

8. Crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate of any one of claims 1-4 for use in the treatment of a hepatitis C virus infection in a human.

9. A pharmaceutical composition comprising crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate of any one of claims 1-4, and another antiviral agent.

10. The pharmaceutical composition of claim 9, wherein the another antiviral agent is an HCV NS3 inhibitor.

11. The pharmaceutical composition of claim 9, wherein the another antiviral agent is an HCV NS5A inhibitor.


12. Crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-

yl)methoxy)(phenoxy)phosphoryl)amino)propanoate of any one of claims 1-4 for use in the treatment of a hepatitis C virus infection in combination with another antiviral agent.

13. Crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate of any one of claims 1-4 for use in the treatment of a hepatitis C virus infection in combination with another antiviral agent, wherein the another antiviral agent is an HCV NS3 inhibitor.

14. Crystalline (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino)propanoate of any one of claims 1-4 for use in the treatment of a hepatitis C virus infection in combination with another antiviral agent, wherein the another antiviral agent is an HCV NS5A inhibitor.

15. A tablet comprising Sp-4:



wherein at least 70% of the Sp-4 is a crystalline Sp-4 having XRPD 2 θ -reflections ($\pm 0.2^\circ$) at 6.1 and 12.7.

16. The tablet according to claim 15 wherein at least 80% of the Sp-4 is a crystalline Sp-4 having XRPD 2 θ -reflections ($\pm 0.2^\circ$) at 6.1 and 12.7.

17. The tablet according to claim 15 or 16 wherein at least 90% of the Sp-4 is a crystalline Sp-4 having XRPD 2 θ -reflections ($\pm 0.2^\circ$) at 6.1 and 12.7.
18. The tablet according to any one of claims 15-17 wherein the crystalline Sp-4 further has XRPD 2 θ -reflections ($\pm 0.2^\circ$) at 8.2, 10.4, 17.2, 17.7, 18.0, 18.8, 19.4, 19.8, 20.1, 20.8, 21.8 and 23.3.
19. Use of a tablet according to any one of claims 15 to 18 for the treatment of a hepatitis C virus infection.
20. Use of a tablet according to any one of claims 15 to 18 and at least one other antiviral agent for the treatment of a hepatitis C virus infection.
21. Use according to claim 20, wherein the other antiviral agent is a HCV NS3 protease inhibitor.
22. Use according to claim 20, wherein the other antiviral agent is a HCV NS5A inhibitor.
23. Use according to claim 20, wherein the other antiviral agent is a HCV NS3 protease inhibitor and a HCV NS5A inhibitor.