

TREM COMPOSITIONS AND METHODS RELATING THERETO

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Provisional Application No. 63/031,941, filed on
5 May 29, 2020, the entire contents of which is hereby incorporated by reference.

BACKGROUND

Transfer RNAs (tRNAs) are complex, naturally occurring RNA molecules that possess a
number of functions including initiation and elongation of proteins.

10

SUMMARY

The present disclosure features modified tRNA-based effector molecules (TREMs, e.g., a
TREM, TREM core fragment, or TREM fragment), as well as related compositions and uses
thereof. A TREM or a related composition thereof can be used, *inter alia*, to modulate a
15 production parameter (e.g., an expression parameter and/or a signaling parameter) of an RNA
corresponding to, or a polypeptide encoded by, a nucleic acid sequence comprising an
endogenous open reading frame (ORF) having a premature termination codon (PTC).
Accordingly, in an aspect, the present disclosure provides a method of modulating a production
parameter of an mRNA corresponding to, or polypeptide encoded by, an endogenous open
20 reading frame (ORF) in a cell, which ORF comprises a codon having a first sequence,
comprising contacting the cell with a TREM composition comprising a TREM, a TREM core
fragment, or a TREM fragment disclosed herein in an amount and/or for a time sufficient to
modulate the production parameter of the mRNA or polypeptide, thereby modulating the
production parameter in the cell. In an embodiment, the TREM, TREM core fragment, or
25 TREM fragment has an anticodon that pairs with the codon having the first sequence.

In another aspect, provided herein is method of modulating a production parameter of an
mRNA corresponding to, or polypeptide encoded by, an endogenous open reading frame (ORF)
in a subject, which ORF comprises a codon having a first sequence, comprising contacting the
subject with a TREM composition comprising a TREM, a TREM core fragment, or a TREM
30 fragment disclosed herein in an amount and/or for a time sufficient to modulate the production
parameter of the mRNA or polypeptide, wherein the TREM, TREM core fragment or TREM

fragment has an anticodon that pairs with the codon having the first sequence, thereby modulating the production parameter in the subject. In an embodiment, the production parameter comprises a signaling parameter and/or an expression parameter, e.g., as described herein.

5 In another aspect, provided herein is a method of modulating expression of a protein in a cell, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a codon having a first sequence, comprising contacting the cell with a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein in an amount and/or for a time sufficient to modulate expression of the encoded protein, wherein the TREM, TREM core fragment or TREM fragment has an anticodon
10 that pairs with the codon having the first sequence, thereby modulating expression of the protein in the cell.

In yet another aspect, provided herein is a method of modulating expression of a protein in a subject, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a codon having a first sequence, comprising
15 contacting the subject with a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein in an amount and/or for a time sufficient to modulate expression of the encoded protein, wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence, thereby modulating expression of the protein in the subject.

20 In an aspect, the disclosure provides, a method of treating a subject having an endogenous open reading frame (ORF) which comprises a codon having a first sequence, comprising providing a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein wherein the TREM comprises a tRNA moiety having an anticodon that pairs with the codon of the ORF having the first sequence; contacting the subject
25 with the composition comprising a TREM, TREM core fragment or TREM fragment in an amount and/or for a time sufficient to treat the subject, thereby treating the subject.

In one aspect, provided herein is a method of modulating a production parameter of an mRNA corresponding to, or polypeptide encoded by, an endogenous open reading frame (ORF) in a subject, which ORF comprises a premature termination codon (PTC), contacting the subject
30 with a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein in an amount and/or for a time sufficient to modulate the production parameter

of the mRNA or polypeptide, wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence, thereby modulating the production parameter in the subject. In an embodiment, the production parameter comprises a signaling parameter and/or an expression parameter, e.g., as described herein.

5 In an aspect, the disclosure provides a method of treating a subject having an endogenous open reading frame (ORF) which comprises a premature termination codon (PTC), comprising providing a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein, wherein the TREM comprises a tRNA moiety having an anticodon that pairs with the PTC in the ORF; contacting the subject with the composition comprising a
10 TREM, TREM core fragment or TREM fragment in an amount and/or for a time sufficient to treat the subject, thereby treating the subject. In an embodiment, the PTC comprises UAA, UGA or UAG.

In yet another aspect, disclosed herein is a method of modulating expression of a protein in a cell, wherein the protein is encoded by a nucleic acid comprising an endogenous open
15 reading frame (ORF), which ORF comprises a premature termination codon (PTC), comprising contacting the cell with a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein in an amount and/or for a time sufficient to modulate expression of the encoded protein, wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the PTC, thereby modulating expression of the protein in the
20 cell. In an embodiment, the PTC comprises UAA, UGA or UAG.

In one aspect, provided herein is a method of modulating expression of a protein in a subject, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a premature termination codon (PTC), comprising contacting the subject with a TREM composition comprising a TREM, a TREM core fragment,
25 or a TREM fragment disclosed herein in an amount and/or for a time sufficient to modulate expression of the encoded protein, wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the PTC, thereby modulating expression of the protein in the subject. In an embodiment, the PTC comprises UAA, UGA or UAG.

In an aspect, provided herein is a method of increasing expression of a protein in a
30 subject wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a premature termination codon (PTC), comprising

contacting the subject, in an amount and/or for a time sufficient to increase expression of the protein, with a TREM composition that (i) has an anticodon that pairs with the PTC, (ii) recognizes an aminoacyl-tRNA synthetase specific for Trp, Tyr, Cys, Glu, Lys, Gln, Ser, Leu, Arg, or Gly, (iii) comprises a sequence of Formula A, or (iv) comprises one or more of a 2'-O-MOE, pseudouridine or 5,6 dihydrouridine modification. In an embodiment, the PTC comprises UAA, UGA or UAG. In an embodiment, the TREM composition comprises (i). In an embodiment, the TREM composition comprises (ii). In an embodiment, the TREM composition comprises (iii). In an embodiment, the TREM composition comprises (iv). In an embodiment, the TREM composition comprises two of (i)-(iv). In an embodiment, the TREM composition comprises three of (i)-(iv). In an embodiment, the TREM composition comprises each of (i)-(iv).

In an aspect, provided herein is a method of increasing expression of a protein in a subject wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a premature termination codon (PTC), comprising:

contacting the subject, in an amount and/or for a time sufficient to increase expression of the protein, with a TREM composition that (i) has an anticodon that pairs with the PTC, (ii) recognizes an aminoacyl-tRNA synthetase specific for Trp, Tyr, Cys, Glu, Lys, Gln, Ser, Leu, Arg, or Gly, (iii) comprises a sequence of Formula B, or (iv) comprises one or more of a 2'-O-MOE, pseudouridine or 5,6 dihydrouridine modification. In an embodiment, the PTC comprises UAA, UGA or UAG. In an embodiment, the TREM composition comprises (i). In an embodiment, the TREM composition comprises (ii). In an embodiment, the TREM composition comprises (iii). In an embodiment, the TREM composition comprises (iv). In an embodiment, the TREM composition comprises two of (i)-(iv). In an embodiment, the TREM composition comprises three of (i)-(iv). In an embodiment, the TREM composition comprises each of (i)-(iv).

In an embodiment of any of the methods disclosed herein, the codon having the first sequence comprises a mutation (e.g., a point mutation, e.g., a nonsense mutation), resulting in a premature termination codon (PTC) chosen from UAA, UGA or UAG. In an embodiment, the codon having the first sequence or the PTC comprises a UAA mutation. In an embodiment, the codon having the first sequence or the PTC comprises a UGA mutation. In an embodiment, the codon having the first sequence or the PTC comprises a UAG mutation.

In an embodiment of any of the methods disclosed herein, the codon having the first sequence or the PTC comprises a UAA, UGA or UAA mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid which preserves, e.g., maintains, a secondary and/or tertiary structure of a polypeptide encoded by the ORF into which the amino acid is incorporated.

In an embodiment of any of the methods disclosed herein, the codon having the first sequence or the PTC comprises a UAA, UGA or UAA mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid which maintains a property, e.g., function, of a polypeptide encoded by the ORF into which the amino acid is incorporated.

In an embodiment of any of the methods disclosed herein, the codon having the first sequence or the PTC comprises a UAA, UGA or UAA mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid which does not alter, e.g., maintains, a production parameter, e.g., an expression parameter and/or a signaling parameter, of an mRNA corresponding to the ORF or a polypeptide encoded by the ORF. In an embodiment, the production parameter is compared to an mRNA corresponding to, or a polypeptide encoded by, an otherwise similar ORF having a pre-mutation, e.g., wildtype, amino acid incorporated at the position corresponding to the first sequence codon or PTC.

In an embodiment of any of the methods disclosed herein, the TREM or TREM fragment comprises a sequence of Formula A. In an embodiment of any of the methods disclosed herein, the TREM core fragment comprises a sequence of Formula B.

In an embodiment of any of the methods disclosed herein, the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for any one of the 20 amino acids. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for Trp, Tyr, Cys, Glu, Lys, Gln, Ser, Leu, Arg, or Gly. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for Trp. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for Tyr. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for Cys. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for Glu. In an embodiment,

the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for Lys. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for Gln. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for Ser. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for Leu. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for Arg. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes an aminoacyl-tRNA synthetase specific for Gly.

10 In an embodiment of any of the methods disclosed herein, the TREM, TREM core fragment or TREM fragment comprises one or more of a 2'-O-MOE, pseudouridine, or a5,6 dihydrouridine modification. In an embodiment of any of the methods disclosed herein, the TREM, TREM core fragment or TREM fragment comprises a 2'-O-MOE modification. In an embodiment of any of the methods disclosed herein, the TREM, TREM core fragment or TREM
15 fragment comprises a pseudouridine modification. In an embodiment of any of the methods disclosed herein, the TREM, TREM core fragment or TREM fragment comprises a 5,6 dihydrouridine modification.

In an aspect, provided herein is a TREM comprising a sequence of Formula A:

[L1]-[ASt Domain1]-[L2]-[DH Domain]-[L3]-[ACH Domain] -[VL Domain]-[TH
20 Domain]-[L4]-[ASt Domain2],

wherein independently, [L1] and [VL Domain], are optional; and one of [L1], [ASt Domain1], [L2]-[DH Domain], [L3], [ACH Domain], [VL Domain], [TH Domain], [L4], and [ASt Domain2] comprises a nucleotide having a non-naturally occurring modification.

In an embodiment, the TREM: (a) retains the ability to: support protein synthesis, be
25 charged by a synthetase, be bound by an elongation factor, introduce an amino acid into a peptide chain, support elongation, or support initiation; (b) comprises at least X contiguous nucleotides without a non-naturally occurring modification, wherein X is greater than 10; (c) comprises at least 3, but less than all of the nucleotides of a type (e.g., A, T, C, G or U) comprise the same non-naturally occurring modification; (d) comprises at least X nucleotides of a type
30 (e.g., A, T, C, G or U) that do not comprise a non-naturally occurring modification, wherein X=1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28,

29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 or 50; (e) comprises no more than 5, 10, or 15 nucleotides of a type (e.g., A, T, C, G or U) that comprise a non-naturally occurring modification; or (f) comprises no more than 5, 10, or 15 nucleotides of a type (e.g., A, T, C, G or U) that do not comprise a non-naturally occurring modification.

5 In an embodiment, the TREM comprises feature (a). In an embodiment, the TREM comprises feature (b). In an embodiment, the TREM comprises feature (c). In an embodiment, the TREM comprises feature (d). In an embodiment, the TREM comprises feature (e). In an embodiment, the TREM comprises feature (f). In an embodiment, the TREM comprises two of features (a)-(f). In an embodiment, the TREM comprises three of features (a)-(f). In an embodiment, the TREM comprises four of features (a)-(f). In an embodiment, the TREM comprises five of features (a)-(f). In an embodiment, the TREM comprises all of features (a)-(f).

In an embodiment, the TREM Domain comprising the non-naturally occurring modification retains a function, e.g., a domain function described herein.

In an aspect, provided herein is a TREM core fragment comprising a sequence of
15 Formula B:

$$[L1]_y-[ASt\ Domain1]_x-[L2]_y-[DH\ Domain]_y-[L3]_y-[ACH\ Domain]_x-[VL\ Domain]_y-[TH\ Domain]_y-[L4]_y-[ASt\ Domain2]_x,$$

wherein $x=1$ and $y=0$ or 1 ; and one of [ASt Domain1], [ACH Domain], and [ASt Domain2] comprises a nucleotide having a non-naturally occurring modification.

20 In an embodiment, the TREM retains the ability to support protein synthesis. In an embodiment, the TREM retains the ability to be able to be charged by a synthetase. In an embodiment, the TREM retains the ability to be bound by an elongation factor. In an embodiment, the TREM retains the ability to introduce an amino acid into a peptide chain. In an embodiment, the TREM retains the ability to support elongation. In an embodiment, the TREM
25 retains the ability to support initiation.

In an embodiment, the [ASt Domain 1] and/or [ASt Domain 2] comprising the non-naturally occurring modification retains the ability to initiate or elongate a polypeptide chain.

In an embodiment, the [ACH Domain] comprising the non-naturally occurring modification retains the ability to mediate pairing with a codon.

30 In an embodiment, $y=1$ for any one, two, three, four, five, six, all or a combination of [L1], [L2], [DH Domain], [L3], [VL Domain], [TH Domain], [L4].

In an embodiment, $y=0$ for any one, two, three, four, five, six, all or a combination of [L1], [L2], [DH Domain], [L3], [VL Domain], [TH Domain], [L4].

In an embodiment, $y=1$ for linker [L1], and L1 comprises a nucleotide having a non-naturally occurring modification.

5 In an embodiment, $y=1$ for linker [L2], and L2 comprises a nucleotide having a non-naturally occurring modification.

In an embodiment, $y=1$ for [DH Domain (DHD)], and DHD comprises a nucleotide having a non-naturally occurring modification. In an embodiment, the DHD comprising the non-naturally occurring modification retains the ability to mediate recognition of aminoacyl-tRNA synthetase.

10 In an embodiment, $y=1$ for linker [L3], and L3 comprises a nucleotide having a non-naturally occurring modification.

In an embodiment, $y=1$ for [VL Domain (VLD)], and VLD comprises a nucleotide having a non-naturally occurring modification.

15 In an embodiment, $y=1$ for [TH Domain (THD)], and THD comprises a nucleotide having a non-naturally occurring modification. In an embodiment, the THD comprising the non-naturally occurring modification retains the ability to mediate recognition of the ribosome.

In an embodiment, $y=1$ for linker [L4], and L4 comprises a nucleotide having a non-naturally occurring modification.

20 In another aspect, the disclosure provides a TREM fragment comprising a portion of a TREM, wherein the TREM comprises a sequence of Formula A:

[L1]-[ASt Domain1]-[L2]-[DH Domain]-[L3]-[ACH Domain]-[VL Domain]-[TH Domain]-[L4]-[ASt Domain2], and wherein the TREM fragment comprises a non-naturally occurring modification.

25 In an embodiment, the TREM fragment comprises one, two, three or all or any combination of the following: (a) a TREM half (*e.g.*, from a cleavage in the ACH Domain, *e.g.*, in the anticodon sequence, *e.g.*, a 5' half or a 3' half); (b) a 5' fragment (*e.g.*, a fragment comprising the 5' end, *e.g.*, from a cleavage in a DH Domain or the ACH Domain); (c) a 3' fragment (*e.g.*, a fragment comprising the 3' end, *e.g.*, from a cleavage in the TH Domain); or (d)
30 an internal fragment (*e.g.*, from a cleavage in any one of the ACH Domain, DH Domain or TH Domain).

In an embodiment, the TREM fragment comprise (a) a TREM half which comprises a nucleotide having a non-naturally occurring modification.

In an embodiment, the TREM fragment comprise (b) a 5' fragment which comprises a nucleotide having a non-naturally occurring modification.

5 In an embodiment, the TREM fragment comprise (c) a 3' fragment which comprises a nucleotide having a non-naturally occurring modification.

In an embodiment, the TREM fragment comprise (d) an internal fragment which comprises a nucleotide having a non-naturally occurring modification.

10 In another aspect, the disclosure provides a pharmaceutical composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein for use in a method disclosed herein.

In another aspect, the disclosure provides a method of making a TREM, a TREM core fragment, or a TREM fragment disclosed herein, comprising linking a first nucleotide to a second nucleotide to form the TREM.

15 In an embodiment, the TREM, TREM core fragment or TREM fragment is synthetic.

In an embodiment, the TREM, TREM core fragment or TREM fragment is made by cell-free solid phase synthesis.

In an embodiment of any of the TREMs, TREM core fragments, or TREM fragments disclosed herein, the TREM Domain comprises a plurality of nucleotides each having a non-naturally occurring modification. In an embodiment, the non-naturally occurring modification comprises a nucleobase modification, a sugar (e.g., ribose) modification, or a backbone modification. In an embodiment, the non-naturally occurring modification is a sugar (e.g., ribose) modification. In an embodiment, the non-naturally occurring modification is 2'-ribose modification, e.g., a 2'-OMe, 2'-halo (e.g., 2'-F), 2'-MOE, or 2'-deoxy modification. In an embodiment, the non-naturally occurring modification is a backbone modification, e.g., a phosphorothioate modification.

20 In an embodiment of any of the TREMs, TREM core fragments, or TREM fragments disclosed herein, the TREM sequence comprises a CCA sequence on a terminus, e.g., the 3' terminus. In an embodiment, the TREM sequence does not comprise a CCA sequence on a terminus, e.g., the 3' terminus.

In an embodiment of any of the TREMs, TREM core fragments, or TREM fragments disclosed herein, the non-naturally occurring modification is a modification in a base or a backbone of a nucleotide, e.g., a modification chosen from any one of Tables 5, 6, 7, 8 or 9.

5 In an embodiment of any of the TREMs, TREM core fragments, or TREM fragments disclosed herein, the non-naturally occurring modification is a base modification chosen from a modification listed in Table 10.

In an embodiment of any of the TREMs, TREM core fragments, or TREM fragments disclosed herein, the non-naturally occurring modification is a base modification chosen from a modification listed in Table 11.

10 In an embodiment of any of the TREMs, TREM core fragments, or TREM fragments disclosed herein, the non-naturally occurring modification is a base modification chosen from a modification listed in Table 12.

In an embodiment of any of the TREMs, TREM core fragments, or TREM fragments disclosed herein, the non-naturally occurring modification is a backbone modification chosen from a modification listed in Table 13.

In an embodiment of any of the TREMs, TREM core fragments, or TREM fragments disclosed herein, the non-naturally occurring modification is a backbone modification chosen from a modification listed in Table 14.

20 Additional features of any of the aforesaid TREMs, TREM core fragments, TREM fragments, TREM compositions, preparations, methods of making TREM compositions and preparations, and methods of using TREM compositions and preparations include one or more of the following enumerated embodiments.

25 Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following enumerated embodiments.

BRIEF DESCRIPTION OF THE DRAWINGS

30 FIGS. 1A-1C are graphs depicting the cell readthrough data of premature termination codons (PTC) in exemplary disease reporters (FIG. 1A - Factor IX at position 298 (*FIX*^{R298X}); FIG. 1B - Tripeptidyl-peptidase 1 at position 208 (*TPPI*^{R208X}); and FIG. 1C - Protocadherin

Related 15 at position 245 (*PCDH15*^{R245X}) after treatment with the unmodified arginine non-cognate TREM and modified arginine non-cognate TREM (TREM-Arg-TGA-Biotin-47), as outlined in Example 15.

5 DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

The present disclosure features methods of modulating a production parameter (e.g., an expression parameter and/or a signaling parameter) of an RNA corresponding to, or polypeptide encoded by, a nucleic acid sequence comprising an endogenous ORF having a premature termination codon (PTC) in a cell or a subject, comprising administering a tRNA-based effector molecule composition (TREM) to the cell or subject. In an embodiment, the TREM composition comprises a TREM, a TREM core fragment, or a TREM fragment comprising a non-naturally occurring modification, e.g., as described herein. Also disclosed herein are methods of modulating expression of a protein in a subject or cell, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF) having a first sequence, e.g., a mutation, e.g., a premature termination codon (PTC), and methods of treating a subject having an endogenous open reading frame (ORF) which comprises a premature termination codon (PTC). Further disclosed herein are TREMs comprising a non-naturally occurring modification, methods of making the same and compositions thereof.

As disclosed herein, TREMs are complex molecules which can mediate a variety of cellular processes. TREM compositions, e.g., pharmaceutical TREM compositions, e.g., TREMs comprising a non-naturally occurring modification, can be administered to a cell, a tissue, or to a subject to modulate these functions. TREMs of the disclosure include TREMs, TREM core fragments and TREM fragments. TREMs, TREM core fragments or TREM fragments can be modified with non-naturally occurring modifications to, e.g., increase the level and/or activity (e.g., stability) of the TREM.

Without wishing to be bound by theory in every case, it is believed that in some embodiments, administration of a TREM composition to a subject or cell having an endogenous ORF having a PTC results in read-through of the PTC, e.g., expression, e.g., increased expression (e.g., increased level and/or activity) of a polypeptide encoded by the ORF having the PTC. In an embodiment, administration of a TREM composition results in modulation of, e.g., increase of, a production parameter of an RNA corresponding to the full length ORF or a

polypeptide encoded by a nucleic acid sequence comprising the full length ORF. In some embodiments, the PTC comprises a UAG, UGA or UAA stop codon. In some embodiments, the TREM comprises an anticodon that pairs with, e.g., recognizes, a stop codon, e.g., a stop codon chosen from UAA, UGA or UAG, and mediates incorporation of an amino acid at the position
5 corresponding to the stop codon. In some embodiments, the production parameter comprises a signaling parameter and/or an expression parameter, e.g., as described herein.

Definitions

“Acquire” or “acquiring” as the terms are used herein, refer to obtaining possession of a
10 value, e.g., a numerical value, by “directly acquiring” or “indirectly acquiring” the physical entity or value. “Directly acquiring” refers to performing a process (e.g., performing an analytical method) to obtain the value. “Indirectly acquiring” refers to receiving the value from another party or source (e.g., a third party laboratory that directly acquired the or value).

An “isoacceptor,” as that term is used herein, refers to a plurality of tRNA molecule or
15 TREMs wherein each molecule of the plurality comprises a different naturally occurring anticodon sequence and each molecule of the plurality mediates the incorporation of the same amino acid and that amino acid is the amino acid that naturally corresponds to the anticodons of the plurality.

A “stop codon” as that term is used herein, refers to a three nucleotide contiguous
20 sequence within messenger RNA that specifies a termination of translation. For example, UAG, UAA, UGA (in RNA) and TAG, TAA or TGA (in DNA) are stop codons. The stop codons are also known as amber (UAG), ochre (UAA), and opal (UGA).

A “premature termination codon” or “PTC” as those terms are used herein, refer to a stop
codon that occurs in an open reading frame (ORF) of a DNA or mRNA. In an embodiment, a
25 PTC occurs at a position upstream of a naturally occurring stop codon in an ORF. In an embodiment, a PTC that occurs upstream of a naturally occurring stop codon, e.g., in an ORF, results in modulation of a production parameter of the corresponding mRNA or polypeptide encoded by the ORF. In an embodiment, a PTC can differ (or arise) from a pre-mutation sequence by a point mutation, e.g., a nonsense mutation. In an embodiment, a PTC can differ (or
30 arise) from a pre-mutation sequence by a genetic change, e.g., abnormality, other than a point mutation, e.g., a frameshift, a deletion, an insertion, a rearrangement, an inversion, a

translocation, a duplication, or a transversion. In an embodiment, a PTC results in the production of a truncated protein which lacks a native activity or which is associated with a mutant, disease, or other unwanted phenotype.

5 A “disease or disorder associated with a PTC” as that term is used herein includes, but is not limited to, a disease or disorder in which cells express, or at one time expressed, a polypeptide encoded by an ORF comprising a PTC. In some embodiments, a disease associated with a PTC is chosen from: a proliferative disorder (e.g., a cancer), a genetic disorder, a metabolic disorder, an immune disorder, an inflammatory disorder or a neurological disorder. Exemplary diseases or disorders associated with a PTC are provided in any one of Tables 15, 16
10 and 17.

An “ORF having a PTC” as that phrase is used herein, refers to an open reading frame (ORF) which comprises a premature termination codon (PTC). In an embodiment, the ORF having the PTC is associated with a disease or disorder associated with a PTC, e.g., as described herein, e.g., a disease or disorder listed in any one of Tables 15, 16 and 17. In an embodiment,
15 the ORF having the PTC is not associated with a disease or disorder associated with a PTC.

A “nucleotide,” as that term is used herein, refers to an entity comprising a sugar, typically a pentameric sugar; a nucleobase; and a phosphate linking group. In an embodiment, a nucleotide comprises a naturally occurring, e.g., naturally occurring in a human cell, nucleotide, e.g., an adenine, thymine, guanine, cytosine, or uracil nucleotide.

20 A “modification,” as that term is used herein with reference to a nucleotide, refers to a modification of the chemical structure, e.g., a covalent modification, of the subject nucleotide. The modification can be naturally occurring or non-naturally occurring. In an embodiment, the modification is non-naturally occurring. In an embodiment, the modification is naturally occurring. In an embodiment, the modification is a synthetic modification. In an embodiment,
25 the modification is a modification provided in Tables 5, 6, 7, 8 or 9.

A “non-naturally occurring modification,” as that term is used herein with reference to a nucleotide, refers to a modification that: (a) a cell, e.g., a human cell, does not make on an endogenous tRNA; or (b) a cell, e.g., a human cell, can make on an endogenous tRNA but wherein such modification is in a location in which it does not occur on a native tRNA, e.g., the
30 modification is in a domain, linker or arm, or on a nucleotide and/or at a position within a domain, linker or arm, which does not have such modification in nature. In either case, the

modification is added synthetically, e.g., in a cell free reaction, e.g., in a solid state or liquid phase synthetic reaction. In an embodiment, the non-naturally occurring modification is a modification that is not present (in identity, location or position) if a sequence of the TREM is expressed in a mammalian cell, e.g., a HEK293 cell line. Exemplary non-naturally occurring
5 modifications are found in Tables 5, 6, 7, 8 or 9.

A “non-naturally modified nucleotide,” as that term is used herein, refers a nucleotide comprising a non-naturally occurring modification on or of a sugar, nucleobase, or phosphate moiety.

A “non-naturally occurring sequence,” as that term is used herein, refers to a sequence
10 wherein an Adenine is replaced by a residue other than an analog of Adenine, a Cytosine is replaced by a residue other than an analog of Cytosine, a Guanine is replaced by a residue other than an analog of Guanine, and a Uracil is replaced by a residue other than an analog of Uracil. An analog refers to any possible derivative of the ribonucleotides, A, G, C or U. In an embodiment, a sequence having a derivative of any one of ribonucleotides A, G, C or U is a non-
15 naturally occurring sequence.

A “naturally occurring nucleotide,” as that term is used herein, refers to a nucleotide that does not comprise a non-naturally occurring modification. In an embodiment, it includes a naturally occurring modification.

A “production parameter,” refers to an expression parameter and/or a signaling
20 parameter. In an embodiment a production parameter is an expression parameter. An expression parameter includes an expression parameter of a polypeptide or protein encoded by the endogenous ORF having a first sequence or PTC; or an expression parameter of an RNA, *e.g.*, messenger RNA, encoded by the endogenous ORF having a first sequence or PTC. In an embodiment, an expression parameter can include:

- 25 (a) protein translation;
(b) expression level (*e.g.*, of polypeptide or protein, or mRNA);
(c) post-translational modification of polypeptide or protein;
(d) folding (*e.g.*, of polypeptide or protein, or mRNA),
(e) structure (*e.g.*, of polypeptide or protein, or mRNA),
30 (f) transduction (*e.g.*, of polypeptide or protein),
(g) compartmentalization (*e.g.*, of polypeptide or protein, or mRNA),

(h) incorporation (*e.g.*, of polypeptide or protein, or mRNA) into a supermolecular structure, *e.g.*, incorporation into a membrane, proteasome, or ribosome,

(i) incorporation into a multimeric polypeptide, *e.g.*, a homo or heterodimer, and/or

(j) stability.

5 In an embodiment, a production parameter is a signaling parameter. A signaling parameter can include:

(1) modulation of a signaling pathway, *e.g.*, a cellular signaling pathway which is downstream or upstream of the protein encoded by the endogenous ORF having a first sequence or PTC;

10 (2) cell fate modulation;

(3) ribosome occupancy modulation;

(4) protein translation modulation;

(5) mRNA stability modulation;

(6) protein folding and structure modulation;

15 (7) protein transduction or compartmentalization modulation; and/or

(8) protein stability modulation.

A “tRNA-based effector molecule” or “TREM,” as that term is used herein, refers to an RNA molecule comprising a structure or property from (a)-(v) below, and which is a recombinant TREM, a synthetic TREM, or a TREM expressed from a heterologous cell. The TREMs described in the present invention are synthetic molecules and are made, *e.g.*, in a cell free reaction, *e.g.*, in a solid state or liquid phase synthetic reaction. TREMs are chemically distinct, *e.g.*, in terms of primary sequence, type or location of modifications from the endogenous tRNA molecules made in cells, *e.g.*, in mammalian cells, *e.g.*, in human cells. A TREM can have a plurality (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9) of the structures and functions of (a)-(v).

25 In an embodiment, a TREM is non-native, as evaluated by structure or the way in which it was made.

In an embodiment, a TREM comprises one or more of the following structures or properties:

(a') an optional linker region of a consensus sequence provided in the “Consensus Sequence” section, *e.g.*, a Linker 1 region;

(a) an amino acid attachment domain that binds an amino acid, *e.g.*, an acceptor stem domain (AStD), wherein an AStD comprises sufficient RNA sequence to mediate, *e.g.*, when present in an otherwise wildtype tRNA, acceptance of an amino acid, *e.g.*, its cognate amino acid or a non-cognate amino acid, and transfer of the amino acid (AA) in the initiation or elongation of a polypeptide chain. Typically, the AStD comprises a 3'-end adenosine (CCA) for acceptor stem charging which is part of synthetase recognition. In an embodiment the AStD has at least 75, 80, 85, 85, 90, 95, or 100% identity with a naturally occurring AStD, *e.g.*, an AStD encoded by a nucleic acid in Table 9. In an embodiment, the TREM can comprise a fragment or analog of an AStD, *e.g.*, an AStD encoded by a nucleic acid in Table 9, which fragment in embodiments has AStD activity and in other embodiments does not have AStD activity. (One of ordinary skill can determine the relevant corresponding sequence for any of the domains, stems, loops, or other sequence features mentioned herein from a sequence encoded by a nucleic acid in Table 9. *E.g.*, one of ordinary skill can determine the sequence which corresponds to an AStD from a tRNA sequence encoded by a nucleic acid in Table 9.)

In an embodiment the AStD falls under the corresponding sequence of a consensus sequence provided in the "Consensus Sequence" section, or differs from the consensus sequence by no more than 1, 2, 5, or 10 positions;

In an embodiment, the AStD comprises residues R₁-R₂-R₃-R₄ -R₅-R₆-R₇ and residues R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁ of Formula I *zzz*, wherein *zzz* indicates any of the twenty amino acids;

In an embodiment, the AStD comprises residues R₁-R₂-R₃-R₄ -R₅-R₆-R₇ and residues R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁ of Formula II *zzz*, wherein *zzz* indicates any of the twenty amino acids;

In an embodiment, the AStD comprises residues R₁-R₂-R₃-R₄ -R₅-R₆-R₇ and residues R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁ of Formula III *zzz*, wherein *zzz* indicates any of the twenty amino acids;

(a'-1) a linker comprising residues R₈-R₉ of a consensus sequence provided in the "Consensus Sequence" section, *e.g.*, a Linker 2 region;

(b) a dihydrouridine hairpin domain (DHD), wherein a DHD comprises sufficient RNA sequence to mediate, *e.g.*, when present in an otherwise wildtype tRNA, recognition of aminoacyl-tRNA synthetase, *e.g.*, acts as a recognition site for aminoacyl-tRNA synthetase for amino acid charging of the TREM. In embodiments, a DHD mediates the stabilization of the TREM's tertiary structure. In an embodiment the DHD has at least 75, 80, 85, 85, 90, 95, or

100% identity with a naturally occurring DHD, *e.g.*, a DHD encoded by a nucleic acid in Table 9. In an embodiment, the TREM can comprise a fragment or analog of a DHD, *e.g.*, a DHD encoded by a nucleic acid in Table 9, which fragment in embodiments has DHD activity and in other embodiments does not have DHD activity.

5 In an embodiment the DHD falls under the corresponding sequence of a consensus sequence provided in the “Consensus Sequence” section, or differs from the consensus sequence by no more than 1, 2, 5, or 10 positions;

In an embodiment, the DHD comprises residues R₁₀-R₁₁-R₁₂-R₁₃-R₁₄ R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈ of Formula I *ZZZ*, wherein *ZZZ* indicates any of the
10 twenty amino acids;

In an embodiment, the DHD comprises residues R₁₀-R₁₁-R₁₂-R₁₃-R₁₄ R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈ of Formula II *ZZZ*, wherein *ZZZ* indicates any of the twenty amino acids;

In an embodiment, the DHD comprises residues R₁₀-R₁₁-R₁₂-R₁₃-R₁₄ R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈ of Formula III *ZZZ*, wherein *ZZZ* indicates any of the
15 twenty amino acids;

(b'-1) a linker comprising residue R₂₉ of a consensus sequence provided in the “Consensus Sequence” section, *e.g.*, a Linker 3 region;

(c) an anticodon that binds a respective codon in an mRNA, *e.g.*, an anticodon hairpin
20 domain (ACHD), wherein an ACHD comprises sufficient sequence, *e.g.*, an anticodon triplet, to mediate, *e.g.*, when present in an otherwise wildtype tRNA, pairing (with or without wobble) with a codon; In an embodiment the ACHD has at least 75, 80, 85, 85, 90, 95, or 100% identity with a naturally occurring ACHD, *e.g.*, an ACHD encoded by a nucleic acid in Table 9. In an embodiment, the TREM can comprise a fragment or analog of an ACHD, *e.g.*, an ACHD
25 encoded by a nucleic acid in Table 9, which fragment in embodiments has ACHD activity and in other embodiments does not have ACHD activity.

In an embodiment the ACHD falls under the corresponding sequence of a consensus sequence provided in the “Consensus Sequence” section, or differs from the consensus sequence by no more than 1, 2, 5, or 10 positions;

In an embodiment, the ACHD comprises residues -R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-R₄₆ of Formula I *ZZZ*, wherein *ZZZ* indicates any of the twenty amino acids;

5 In an embodiment, the ACHD comprises residues -R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-R₄₆ of Formula II *ZZZ*, wherein *ZZZ* indicates any of the twenty amino acids;

In an embodiment, the ACHD comprises residues -R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-R₄₆ of Formula III *ZZZ*, wherein *ZZZ* indicates any of the twenty amino acids;

10 (d) a variable loop domain (VLD), wherein a VLD comprises sufficient RNA sequence to mediate, *e.g.*, when present in an otherwise wildtype tRNA, recognition of aminoacyl-tRNA synthetase, *e.g.*, acts as a recognition site for aminoacyl-tRNA synthetase for amino acid charging of the TREM. In embodiments, a VLD mediates the stabilization of the TREM's tertiary structure. In an embodiment, a VLD modulates, *e.g.*, increases, the specificity of the
15 TREM, *e.g.*, for its cognate amino acid, *e.g.*, the VLD modulates the TREM's cognate adaptor function. In an embodiment the VLD has at least 75, 80, 85, 85, 90, 95, or 100% identity with a naturally occurring VLD, *e.g.*, a VLD encoded by a nucleic acid in Table 9. In an embodiment, the TREM can comprise a fragment or analog of a VLD, *e.g.*, a VLD encoded by a nucleic acid in Table 9, which fragment in embodiments has VLD activity and in other embodiments does not
20 have VLD activity.

In an embodiment the VLD falls under the corresponding sequence of a consensus sequence provided in the "Consensus Sequence" section.

In an embodiment, the VLD comprises residue -[R₄₇]_x of a consensus sequence provided in the "Consensus Sequence" section, wherein x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28,
25
30

x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271);

(e) a thymine hairpin domain (THD), wherein a THD comprises sufficient RNA sequence, to mediate, *e.g.*, when present in an otherwise wildtype tRNA, recognition of the ribosome, *e.g.*, acts as a recognition site for the ribosome to form a TREM-ribosome complex during translation. In an embodiment the THD has at least 75, 80, 85, 85, 90, 95, or 100% identity with a naturally occurring THD, *e.g.*, a THD encoded by a nucleic acid in Table 9. In an embodiment, the TREM can comprise a fragment or analog of a THD, *e.g.*, a THD encoded by a nucleic acid in Table 9, which fragment in embodiments has THD activity and in other embodiments does not have THD activity.

In an embodiment the THD falls under the corresponding sequence of a consensus sequence provided in the “Consensus Sequence” section, or differs from the consensus sequence by no more than 1, 2, 5, or 10 positions;

In an embodiment, the THD comprises residues -R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄ of Formula I ZZZ, wherein ZZZ indicates any of the twenty amino acids;

In an embodiment, the THD comprises residues -R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄ of Formula II ZZZ, wherein ZZZ indicates any of the twenty amino acids;

In an embodiment, the THD comprises residues -R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄ of Formula III ZZZ, wherein ZZZ indicates any of the twenty amino acids;

(e'1) a linker comprising residue R₇₂ of a consensus sequence provided in the “Consensus Sequence” section, *e.g.*, a Linker 4 region;

(f) under physiological conditions, it comprises a stem structure and one or a plurality of loop structures, *e.g.*, 1, 2, or 3 loops. A loop can comprise a domain described herein, *e.g.*, a domain selected from (a)-(e). A loop can comprise one or a plurality of domains. In an embodiment, a stem or loop structure has at least 75, 80, 85, 85, 90, 95, or 100% identity with a naturally occurring stem or loop structure, *e.g.*, a stem or loop structure encoded by a nucleic acid in Table 9. In an embodiment, the TREM can comprise a fragment or analog of a stem or loop structure, *e.g.*, a stem or loop structure encoded by a nucleic acid in Table 9, which

fragment in embodiments has activity of a stem or loop structure, and in other embodiments does not have activity of a stem or loop structure;

(g) a tertiary structure, *e.g.*, an L-shaped tertiary structure;

(h) adaptor function, *i.e.*, the TREM mediates acceptance of an amino acid, *e.g.*, its
5 cognate amino acid and transfer of the AA in the initiation or elongation of a polypeptide chain;

(i) cognate adaptor function wherein the TREM mediates acceptance and incorporation of an amino acid (*e.g.*, cognate amino acid) associated in nature with the anti-codon of the TREM to initiate or elongate a polypeptide chain;

(j) non-cognate adaptor function, wherein the TREM mediates acceptance and
10 incorporation of an amino acid (*e.g.*, non-cognate amino acid) other than the amino acid associated in nature with the anti-codon of the TREM in the initiation or elongation of a polypeptide chain;

(k) a regulatory function, *e.g.*, an epigenetic function (*e.g.*, gene silencing function or signaling pathway modulation function), cell fate modulation function, mRNA stability
15 modulation function, protein stability modulation function, protein transduction modulation function, or protein compartmentalization function;

(l) a structure which allows for ribosome binding;

(m) a post-transcriptional modification, *e.g.*, a naturally occurring post-transcriptional modification;

(n) the ability to inhibit a functional property of a tRNA, *e.g.*, any of properties (h)-(k)
20 possessed by a tRNA;

(o) the ability to modulate cell fate;

(p) the ability to modulate ribosome occupancy;

(q) the ability to modulate protein translation;

(r) the ability to modulate mRNA stability;
25

(s) the ability to modulate protein folding and structure;

(t) the ability to modulate protein transduction or compartmentalization;

(u) the ability to modulate protein stability; or

(v) the ability to modulate a signaling pathway, *e.g.*, a cellular signaling pathway.

30 In an embodiment, a TREM comprises a full-length tRNA molecule or a fragment thereof.

In an embodiment, a TREM comprises the following properties: (a)-(e).

In an embodiment, a TREM comprises the following properties: (a) and (c).

In an embodiment, a TREM comprises the following properties: (a), (c) and (h).

In an embodiment, a TREM comprises the following properties: (a), (c), (h) and (b).

5 In an embodiment, a TREM comprises the following properties: (a), (c), (h) and (e).

In an embodiment, a TREM comprises the following properties: (a), (c), (h), (b) and (e).

In an embodiment, a TREM comprises the following properties: (a), (c), (h), (b), (e) and (g).

In an embodiment, a TREM comprises the following properties: (a), (c), (h) and (m).

10 In an embodiment, a TREM comprises the following properties: (a), (c), (h), (m), and (g).

In an embodiment, a TREM comprises the following properties: (a), (c), (h), (m) and (b).

In an embodiment, a TREM comprises the following properties: (a), (c), (h), (m) and (e).

In an embodiment, a TREM comprises the following properties: (a), (c), (h), (m), (g), (b) and (e).

15 In an embodiment, a TREM comprises the following properties: (a), (c), (h), (m), (g), (b), (e) and (q).

In an embodiment, a TREM comprises:

(i) an amino acid attachment domain that binds an amino acid (*e.g.*, an AStD, as described in (a) herein; and

20 (ii) an anticodon that binds a respective codon in an mRNA (*e.g.*, an ACHD, as described in (c) herein).

In an embodiment the TREM comprises a flexible RNA linker which provides for covalent linkage of (i) to (ii).

In an embodiment, the TREM mediates protein translation.

25 In an embodiment a TREM comprises a linker, *e.g.*, an RNA linker, *e.g.*, a flexible RNA linker, which provides for covalent linkage between a first and a second structure or domain. In an embodiment, an RNA linker comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 ribonucleotides. A TREM can comprise one or a plurality of linkers, *e.g.*, in embodiments a TREM comprising (a), (b), (c), (d) and (e) can have a first linker between a first and second
30 domain, and a second linker between a third domain and another domain.

In an embodiment, the TREM comprises a sequence of Formula A: [L1]-[ASt Domain1]-[L2]-[DH Domain]-[L3]-[ACH Domain]-[VL Domain]-[TH Domain]-[L4]-[ASt Domain2].

In an embodiment, a TREM comprises an RNA sequence at least 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98 or 99% identical with, or which differs by no more than 1, 2, 3, 4, 5, 10, 15, 20, 25, or 30 ribonucleotides from, an RNA sequence encoded by a DNA sequence listed in Table 9, or a fragment or functional fragment thereof. In an embodiment, a TREM comprises an RNA sequence encoded by a DNA sequence listed in Table 9, or a fragment or functional fragment thereof. In an embodiment, a TREM comprises an RNA sequence encoded by a DNA sequence at least 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98 or 99% identical with a DNA sequence listed in Table 9, or a fragment or functional fragment thereof. In an embodiment, a TREM comprises a TREM domain, *e.g.*, a domain described herein, comprising at least 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, or 99% identical with, or which differs by no more than 1, 2, 3, 4, 5, 10, or 15, ribonucleotides from, an RNA encoded by a DNA sequence listed in Table 9, or a fragment or a functional fragment thereof. In an embodiment, a TREM comprises a TREM domain, *e.g.*, a domain described herein, comprising an RNA sequence encoded by DNA sequence listed in Table 9, or a fragment or functional fragment thereof. In an embodiment, a TREM comprises a TREM domain, *e.g.*, a domain described herein, comprising an RNA sequence encoded by DNA sequence at least 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98 or 99% identical with a DNA sequence listed in Table 9, or a fragment or functional fragment thereof.

In an embodiment, a TREM is 76-90 nucleotides in length. In embodiments, a TREM or a fragment or functional fragment thereof is between 10-90 nucleotides, between 10-80 nucleotides, between 10-70 nucleotides, between 10-60 nucleotides, between 10-50 nucleotides, between 10-40 nucleotides, between 10-30 nucleotides, between 10-20 nucleotides, between 20-90 nucleotides, between 20-80 nucleotides, 20-70 nucleotides, between 20-60 nucleotides, between 20-50 nucleotides, between 20-40 nucleotides, between 30-90 nucleotides, between 30-80 nucleotides, between 30-70 nucleotides, between 30-60 nucleotides, or between 30-50 nucleotides.

In an embodiment, a TREM is aminoacylated, *e.g.*, charged, with an amino acid by an aminoacyl tRNA synthetase.

In an embodiment, a TREM is not charged with an amino acid, *e.g.*, an uncharged TREM (uTREM).

In an embodiment, a TREM comprises less than a full length tRNA. In embodiments, a TREM can correspond to a naturally occurring fragment of a tRNA, or to a non-naturally occurring fragment. Exemplary fragments include: TREM halves (*e.g.*, from a cleavage in the ACHD, *e.g.*, in the anticodon sequence, *e.g.*, 5' halves or 3' halves); a 5' fragment (*e.g.*, a fragment comprising the 5' end, *e.g.*, from a cleavage in a DHD or the ACHD); a 3' fragment (*e.g.*, a fragment comprising the 3' end, *e.g.*, from a cleavage in the THD); or an internal fragment (*e.g.*, from a cleavage in one or more of the ACHD, DHD or THD).

A "TREM core fragment," as that term is used herein, refers to a portion of the sequence of Formula B: [L1]_y-[ASt Domain1]_x-[L2]_y-[DH Domain]_y-[L3]_y-[ACH Domain]_x-[VL Domain]_y-[TH Domain]_y-[L4]_y-[ASt Domain2]_x, wherein: x=1 and y=0 or 1.

A "TREM fragment," as used herein, refers to a portion of a TREM, wherein the TREM comprises a sequence of Formula A: [L1]-[ASt Domain1]-[L2]-[DH Domain]-[L3]-[ACH Domain]-[VL Domain]-[TH Domain]-[L4]-[ASt Domain2].

A "cognate adaptor function TREM," as that term is used herein, refers to a TREM which mediates initiation or elongation with the AA (the cognate AA) associated in nature with the anti-codon of the TREM.

"Decreased expression," as that term is used herein, refers to a decrease in comparison to a reference, *e.g.*, in the case where altered control region, or addition of an agent, results in a decreased expression of the subject product, it is decreased relative to an otherwise similar cell without the alteration or addition.

"Increased expression," as that term is used herein, refers to an increase in comparison to a reference, *e.g.*, in the case where altered control region, or addition of an agent, results in an increased expression of the subject product, it is increased relative to an otherwise similar cell without the alteration or addition.

As used herein, the terms "increasing" and "decreasing" refer to modulating that results in, respectively, greater or lesser amounts of function, expression, or activity of a particular metric relative to a reference. For example, subsequent to administration to a cell, tissue or subject of a TREM described herein, the amount of a marker of a metric (*e.g.*, protein translation, mRNA stability, protein folding) as described herein may be increased or decreased by at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98%, 2X, 3X, 5X, 10X or more relative to the amount of the marker prior to

administration or relative to the effect of a negative control agent. The metric may be measured subsequent to administration at a time that the administration has had the recited effect, *e.g.*, at least 12 hours, 24 hours, one week, one month, 3 months, or 6 months, after a treatment has begun.

5 An “exogenous nucleic acid,” as that term is used herein, refers to a nucleic acid sequence that is not present in or differs by at least one nucleotide from the closest sequence in a reference cell, *e.g.*, a cell into which the exogenous nucleic acid is introduced. In an embodiment, an exogenous nucleic acid comprises a nucleic acid that encodes a TREM.

 An “exogenous TREM,” as that term is used herein, refers to a TREM that:

10 (a) differs by at least one nucleotide or one post transcriptional modification from the closest sequence tRNA in a reference cell, *e.g.*, a cell into which the exogenous nucleic acid is introduced;

 (b) has been introduced into a cell other than the cell in which it was transcribed;

 (c) is present in a cell other than one in which it naturally occurs; or

15 (d) has an expression profile, *e.g.*, level or distribution, that is non-wildtype, *e.g.*, it is expressed at a higher level than wildtype. In an embodiment, the expression profile can be mediated by a change introduced into a nucleic acid that modulates expression or by addition of an agent that modulates expression of the RNA molecule. In an embodiment an exogenous TREM comprises 1, 2, 3 or 4 of properties (a)-(d).

20 A “GMP-grade composition,” as that term is used herein, refers to a composition in compliance with current good manufacturing practice (cGMP) guidelines, or other similar requirements. In an embodiment, a GMP-grade composition can be used as a pharmaceutical product.

 A “non-cognate adaptor function TREM,” as that term is used herein, refers to a TREM
25 which mediates initiation or elongation with an AA (a non-cognate AA) other than the AA associated in nature with the anti-codon of the TREM. In an embodiment, a non-cognate adaptor function TREM is also referred to as a mischarged TREM (mTREM).

 A “pharmaceutical TREM composition,” as that term is used herein, refers to a TREM composition that is suitable for pharmaceutical use. Typically, a pharmaceutical TREM
30 composition comprises a pharmaceutical excipient. In an embodiment the TREM will be the only active ingredient in the pharmaceutical TREM composition. In embodiments the

pharmaceutical TREM composition is free, substantially free, or has less than a pharmaceutically acceptable amount, of host cell proteins, DNA, *e.g.*, host cell DNA, endotoxins, and bacteria.

“Post-transcriptional processing,” as that term is used herein, with respect to a subject molecule, *e.g.*, a TREM, RNA or tRNAs, refers to a covalent modification of the subject molecule. In an embodiment, the covalent modification occurs post-transcriptionally. In an embodiment, the covalent modification occurs co-transcriptionally. In an embodiment the modification is made *in vivo*, *e.g.*, in a cell used to produce a TREM. In an embodiment the modification is made *ex vivo*, *e.g.*, it is made on a TREM isolated or obtained from the cell which produced the TREM.

A “synthetic TREM,” as that term is used herein, refers to a TREM which was synthesized other than in or by a cell having an endogenous nucleic acid encoding the TREM, *e.g.*, a synthetic TREM is synthesized by cell-free solid phase synthesis. A synthetic TREM can have the same, or a different, sequence, or tertiary structure, as a native tRNA.

A “recombinant TREM,” as that term is used herein, refers to a TREM that was expressed in a cell modified by human intervention, having a modification that mediates the production of the TREM, *e.g.*, the cell comprises an exogenous sequence encoding the TREM, or a modification that mediates expression, *e.g.*, transcriptional expression or post-transcriptional modification, of the TREM. A recombinant TREM can have the same, or a different, sequence, set of post-transcriptional modifications, or tertiary structure, as a reference tRNA, *e.g.*, a native tRNA.

A “tRNA”, as that term is used herein, refers to a naturally occurring transfer ribonucleic acid in its native state.

A “TREM composition,” as that term is used herein, refers to a composition comprising a plurality of TREMs, a plurality of TREM core fragments and/or a plurality of TREM fragments.

A TREM composition can comprise one or more species of TREMs, TREM core fragments or TREM fragments. In an embodiment, the composition comprises only a single species of TREM, TREM core fragment or TREM fragment. In an embodiment, the TREM composition comprises a first TREM, TREM core fragment or TREM fragment species; and a second TREM, TREM core fragment or TREM fragment species. In an embodiment, the TREM composition comprises X TREM, TREM core fragment or TREM fragment species, wherein X=2, 3, 4, 5, 6, 7, 8, 9, or 10. In an embodiment, the TREM, TREM core fragment or TREM fragment has at

least 70, 75, 80, 85, 90, or 95, or has 100%, identity with a sequence encoded by a nucleic acid in Table 9. A TREM composition can comprise one or more species of TREMs, TREM core fragments or TREM fragments. In an embodiment, the TREM composition is at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 95 or 99% dry weight TREMs (for a liquid composition dry weight refers to the weight after removal of substantially all liquid, *e.g.*, after lyophilization). In an embodiment, the composition is a liquid. In an embodiment, the composition is dry, *e.g.*, a lyophilized material. In an embodiment, the composition is a frozen composition. In an embodiment, the composition is sterile. In an embodiment, the composition comprises at least 0.5 g, 1.0 g, 5.0 g, 10 g, 15 g, 25 g, 50 g, 100 g, 200 g, 400 g, or 500 g (*e.g.*, as determined by dry weight) of TREM.

In an embodiment, at least X% of the TREMs in a TREM composition has a non-naturally occurring modification at a selected position, and X is 80, 90, 95, 96, 97, 98, 99, or 99.5.

In an embodiment, at least X% of the TREMs in a TREM composition has a non-naturally occurring modification at a first position and a non-naturally occurring modification at a second position, and X, independently, is 80, 90, 95, 96, 97, 98, 99, or 99.5. In embodiments, the modification at the first and second position is the same. In embodiments, the modification at the first and second position are different. In embodiments, the nucleotide at the first and second position is the same, *e.g.*, both are adenine. In embodiments, the nucleotide at the first and second position are different, *e.g.*, one is adenine and one is thymine.

In an embodiment, at least X% of the TREMs in a TREM composition has a non-naturally occurring modification at a first position and less than Y% have a non-naturally occurring modification at a second position, wherein X is 80, 90, 95, 96, 97, 98, 99, or 99.5 and Y is 20, 20, 5, 2, 1, .1, or .01. In embodiments, the nucleotide at the first and second position is the same, *e.g.*, both are adenine. In embodiments the nucleotide at the first and second position are different, *e.g.*, one is adenine and one is thymine.

“Pairs with” or “pairing,” as those terms are used herein, refer to the correspondence of a codon with an anticodon and includes fully complementary codon:anticodon pairs as well as “wobble” pairing, in which the third position need not be complementary. Fully complementary pairing refers to pairing of all three positions of the codon with the corresponding anticodon according to Watson-Crick base pairing. Wobble pairing refers to complementary pairing of the

first and second positions of the codon with the corresponding anticodon according to Watson-Crick base pairing, and flexible pairing at the third position of the codon with the corresponding anticodon.

A “subject,” as this term is used herein, includes any organism, such as a human or other animal. In embodiments, the subject is a vertebrate animal (e.g., mammal, bird, fish, reptile, or amphibian). In embodiments, the subject is a mammal, e.g., a human. In embodiments, the method subject is a non-human mammal. In embodiments, the subject is a non-human mammal such as a non-human primate (e.g., monkeys, apes), ungulate (e.g., cattle, buffalo, sheep, goat, pig, camel, llama, alpaca, deer, horses, donkeys), carnivore (e.g., dog, cat), rodent (e.g., rat, mouse), or lagomorph (e.g., rabbit). In embodiments, the subject is a bird, such as a member of the avian taxa Galliformes (e.g., chickens, turkeys, pheasants, quail), Anseriformes (e.g., ducks, geese), Paleognathae (e.g., ostriches, emus), Columbiformes (e.g., pigeons, doves), or Psittaciformes (e.g., parrots). The subject may be a male or female of any age group, e.g., a pediatric subject (e.g., infant, child, adolescent) or adult subject (e.g., young adult, middle-aged adult, or senior adult)). A non-human subject may be a transgenic animal.

The terms modified, replace, derived and similar terms, when used or applied in reference to a product, refer only to the end product or structure of the end product, and are not restricted by any method of making or manufacturing the product, unless expressly provided as such in this disclosure.

Headings, titles, subtitles, numbering or other alpha/numeric hierarchies are included merely for ease of reading and absent explicit language to the contrary do not indicate order of performance, order of importance, magnitude or other value.

Premature termination codons (PTC) and ORFs comprising PTCs

Mutations underlie many diseases. For example, a point mutation in the open reading frame (ORF) of a gene which creates a premature stop codon (PTC) can result in altered expression and/or activity of a polypeptide encoded by the gene. Table 1 provides single mutations in codons encoding amino acids which can result in a stop codon. In an embodiment, a PTC disclosed herein comprises a mutation disclosed in Table 1.

In an embodiment, the codon having the first sequence or the PTC comprises a mutation disclosed in Table 1. In an embodiment, the non-mutated, e.g., wildtype, codon sequence of the

codon having the first sequence or the PTC is an original codon sequence provided in Table 1 and the amino acid corresponding to the non-mutated codon is an original AA provided in Table 1.

In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes a stop codon and mediates incorporation of the original AA provided in Table 1 at the position of the stop codon. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes a stop codon and mediates incorporation of an amino acid belonging to the same group as the original AA, e.g., as provided in Table 2. Other genetic abnormalities, such as insertions and/or deletions can also result in a PTC in an ORF.

10

Table 1. Select amino acids and stop codons

Original AA	Original codon	One mutation to stop codon
TRP	UGG	UGA
TYR	UAU	UAA
	UAC	UAG
CYS	UGU	UGA
	UGC	UGA
GLU	GAA	UAA
	GAG	UAG
LYS	AAA	UAA
	AAG	UAG
GLN	CAA	UAA
	CAG	UAG
SER	UCA	UGA
	UCG	UAG
LEU	UUA	UAA OR UGA
	UUG	UAG
ARG	CGA	UGA
GLY	GGA	UGA

Table 2: Amino acids and amino acid groupings

Group	Amino acid
Nonpolar, aliphatic R group	leucine methionine isoleucine glycine alanine valine
Polar, uncharged R group	serine

	threonine cysteine proline asparagine glutamine
positively charged R group	lysine arginine histidine
Negatively charged R group	aspartate glutamate
Nonpolar, aromatic R group	phenylalanine tyrosine tryptophan

Disclosed herein, inter alia, are endogenous ORFs comprising a codon having a first sequence, e.g., a mutation, e.g., a PTC. An ORF having a PTC, e.g., as described herein, can be present, or part of in any gene. As an example, the ORF can be present or be part of any gene in the human genome.

In an embodiment, a PTC disclosed herein is present in a gene disclosed in any one of Tables 4, 6 or 3. Exemplary genes having ORFs comprising a PTC are provided in Table 3.

Table 3: Exemplary genes with ORFs having a PTC

A2ML1	ARFGEF1	CACNA1G	CNOT1	DLG4
AARS1	ARFGEF2	CACNA1S	COG1	DLL1
AARS2	ARHGAP21	CACNA2D1	COL11A2	DNA2
ABCA13	ARHGEF9	CACNA2D2	COL13A1	DNM1L
ABCB11	ARMC4	CACNB2, NSUN6	COL4A1	DNMT1
ABCG5	ARV1	CAD	COL4A2	DNMT3A
ABHD5	ARX	CAMTA1	COL4A4	DNMT3B
ACAD8	ASCC3	CARS2	COL9A1	DPH1
ACADL	ASH1L	CCDC140	COQ4	DPYD-AS1
ACSF3	ASPH	CCDC8	COQ6	DSEL
ACTA2	ASXL2	CCM2	COX14	DSPP
ACTC1	ATAD3A	CD40LG	CPE	DUOXA1
ACTN2	ATP2A2	CDAN1	CPEB1-AS1	DUOXA2
ACVR1	ATP6V1B1	CDH15	CREBBP	DVL1
ADAR	ATP8A2	CDK11A	CRELD1	EARS2
ADAT3	AUH	CEBPA	CSNK2A1	EBF3
ADCY5	AUTS2	CELF5	CSNK2B	EBP
ADIPOQ	AVPR2	CELSR2	CSTA	EDAR

ADIPOQ-AS1	B3GLCT	CEP135	CTNND2	EFHC1
ADIPOR1	B4GAT1	CEP164	CTSA	EFNB1
AFF2	BCAP31	CEP83	CTSC	EFTUD2
ALG11	BCL11A	CETP	CUL3	EIF2B5
ALG14	BCL11B	CFI	CYLD	ELANE
ALG6	BCORL1	CHAMP1	CYP11A1	EMC1
ALOXE3	BEND2	CHAT	DAG1	EMC1-AS1
AMER1	BGN	CHD1	DARS2	ENO3
AMH	BMP4	CHD4	DCX	ENTPD5
AMMECR1	BRD4	CHD8	DDHD2	EP300
AMN	BRPF1	CHRM2	DDR2	EPM2A
ANK2	BRSK2	CHRNA1	DEAF1	ERLIN2
ANK3	BUB1B	CIC	DENND5A	EVC
ANKS6	BUB1B-PAK6	CLCN7	DGAT1	EZH2
ANOS1	C8G	CLTC	DHFR	FAM111A
AP1S1	CACNA1E	CNGA3	DIAPH3	FAM126A
AP3B2	CACNA1F	CNKS2	DISP1	FAN1
FANCD2	GJB6	ISLR2	LOC110673972	MTR
FANCE	GJC2	ITGA3	LOC112997540	MYCBP2
FASTKD2	GK	ITGA8	LOC113788297	MYCNOS
FAT1	GLI2	ITGB6	LOC349160	MYH7B
FBN2	GNAI1	JMJD1C	LONP1	MYL3
FBP1	GNB1L	KANK1	LORICRIN	MYOM1
FDXR	GNE	KANSL1	LPIN1	MYPN
FGA	GNRH1	KBTBD13	LPIN2	MYT1L
FGD1	GNS	KCNA2	LRP2	NAA15
FGF10	GPAA1	KCNB1	LRRTM4	NAGA
FGFR1	GPD1L	KCND2	MAB21L2	NARS2
FGFR2	GRIN2A	KCNE3	MAF	NAXE
FIBP	GTPBP3	KCNMA1	MARS1	NCAPH2
FLAD1	HACE1	KCNQ5	MARS2	NCF2
FLG-AS1	HADH	KCTD7	MASP1	NDP
FLVCR1	HADHB	KIDINS220	MBOAT7	NDP-AS1
FLVCR2	HDAC4	KIF21B	MCM3AP	NDRG1
FMN2	HERC1	KIF6	MCM3AP-AS1	NDUFA2, TMCO6
FOXA2	HESX1	KIT	MED13	NDUFAF1
FOXC1	HIBCH	KLHL40	MED17	NDUFAF5
FOXC2	HNF4A	KLHL41	MEGF10	NDUFAF6
FOXC2-AS1	HNRNP2	KNL1	MET	NDUFAF7

FOXP2	HPRT1	KRAS	MGAT2	NDUFS1
FREM1	HRG	LAMB1	MIB1	NDUFS3
FRYL	HUWE1	LAMB2	MICU1	NEFH
FTL	IARS1	LAMC3	MIR302CHG	NEK8
FUS	IBA57	LARP7	MIR5004	NEXMIF
GABRG2	IDH2	LARS1	MIR6501	NFIA
GAN	IFNAR1	LCT	MITD1	NFKB1
GATA2	IFT122	LEMD3	MMP13	NHEJ1
GATA4	IFT80	LGI4	MMP21	NICN1
GATAD1	IGF2	LIAS	MNX1	NID1
GDF5	ILDR1	LINS1	MNX1-AS2	NKX2-1
GDF5-AS1	ILK, TAF10	LIPC	MPDU1	NLRP1
GFM1	INF2	LIPT1	MRPS22	NLRP3
GH-LCR	INS-IGF2	LOC101448202	MSL3	NOD2
GHRHR	INSR	LOC106804612, HBA2	MSRB3	NONO
GHSR	IRAK1BP1	LOC107303338	MT-ND2	NOTCH3
GJA1	IRAK3	RARS2	SETD1B	SPTLC1
NPHP4	PLEKHG5	RAX	SETD2	SRD5A3
NPR2	PLEKHM2	RBM10	SETX	SRPX2
NR2F2	PLK4	RELN	SFTA3	SSBP2
NR5A1	PLPBP	RERE	SHANK2	ST3GAL3
NRL, PCK2	PNKD	RFT1	SHH	ST3GAL5
NRXN1	PNPLA1	RMND1	SIN3A	STAMBP
NT5DC1	POC1A	RNASEH1	SIX3	STAT3
NTRK2	POLG2, MILR1	RNF17	SKI	STIL
NUBPL	POMGNT2	ROR2	SLC13A5	STX11
NUS1	PPM1D	RP2	SLC16A1	STX1B
OCRL	PPP3CA	RPL11	SLC16A2	SUCLA2
OPTN	PRDM1	RPS19	SLC17A8	SYN1
P4HA1	PRDM12	RRM2B	SLC18A3	SYN2
PAK6	PREPL	RS1	SLC20A2	SYNJ1
PBX1	PRICKLE1	RUNX2	SLC25A4	TAB2
PCARE	PRKAG2	RXYLT1	SLC25A46	TACR3
PCDH12	PROS1	S100PBP	SLC2A1	TBCD
PDCD10	PRPF31	SALL4	SLC39A8	TBL1XR1
PDE11A-AS1	PRPF8	SAMD9	SLC52A2	TBX1
PDE4D	PRPS1	SAR1B	SLC6A3	TBX18
PDE6A	PRSS1, TRB	SASH3	SLC6A8	TCIRG1
PDHX	PSAT1	SBF2	SLC6A9	TELO2
PDLIM3	PSMD12	SBF2-AS1	SLC7A9	TFAP2A

PDP1	PSTPIP1	SCAMP4	SMAD2	TFG
PDSS1	PTCHD1	SCLT1	SMAD9	TGIF1
PEX5	PTF1A	SCN10A	SMARCA2	THAP1
PHF21A	PTPN23	SCN11A	SMC3	TINF2
PHF8	PTPRQ	SCN1B	SMOC2	TLK2
PHKA2	PTRH2	SCN3A	SNTA1	TMEM43
PHKB	PUS1	SCN4A	SNX14	TMIE
PIEZO1	PYCR2	SCN4B	SNX22	TMPO
PIGG	QARS1	SCN8A	SOCS1	TMPRSS15
PIGL	RAB3GAP1, ZRANB3	SCO2	SOX11	TMTC3
PIGM	RAB3GAP2	SCYL1	SOX17	TNFAIP3
PIGP	RAC1	SEMA4A	SPATA5	TNFRSF11A
PIK3CA	RAF1	SEPTIN9	SPATA7	TOE1
PIN4, ERCC6L	RAG1	SET	SPRED1	TOR1AIP1
PLAT	RARB	SETD1A	SPTBN2	TPK1
TPM1	UTP14C	ZEB1		
TRAPPC9	VPS53	ZFHX4		
TRIM37	WDR19	ZFPM2		
TRIM59-IFT80	WDR26	ZFPM2-AS1		
TRIO	WDR62	ZIC1		
TRIP12	WDR81	ZIC2		
TRIP4	WNT1	ZMYND11		
TRPM1	WNT10A	ZNF335		
TSEN54	WRAP53	ZNF423		
TUBA4A	WWOX	ZNF469		
TUBGCP4	YARS1			
TUBGCP6	YARS2			
TWINK	YY1			
TXNRD2	ZAP70			
UBA2	ZBTB20			
UBA5	ZBTB24			
UMOD	ZC4H2			
UNC5B	ZDHHC9			

Diseases or disorders associated with a PTC

A TREM composition disclosed herein can be used treat a disorder or disease associated with a PTC, e.g., as described herein. Exemplary diseases or disorders associated with a PTC are listed in Tables 4, 5, and 6.

In an embodiment, the subject has a disease or disorder provided in any one of Tables 4-6. In an embodiment, the cell is associated with, e.g., is obtained from a subject who has, a disorder or disease listed in any one of Tables 4-6.

For example, the disorder or disease can be chosen from the left column of Table 4. As another example, the disorder or disease is chosen from the left column of Table 4 and, in embodiments the PTC is in a gene chosen from the right column of Table 4, e.g., any one of the genes provided in the right column of Table 4. In some embodiments, the PTC is in a gene corresponding to the disorder or disease provided in the left column of Table 4. As a further non-limiting example, the PTC can be at a position provided in Table 4.

As another example, the disorder or symptom is chosen from a disorder or disease provided in Table 5.

As yet another example, the disorder or symptom is chosen from a disorder or disease provided in Table 6. In an embodiment, the disorder or symptom is chosen from a disorder or disease provided in Table 6 and, in embodiments, the PTC is in any gene provided in Table 6. In an embodiment, the disorder or symptom is chosen from a disorder or disease provided in Table 6 and the PTC is in a corresponding gene provided in Table 6, e.g., a gene corresponding to the disease or disorder. In an embodiment, the disorder or symptom is chosen from a disorder or disease provided in Table 6 and the PTC is not in a gene provided in Table 6.

In an embodiment of any of the methods disclosed herein, the PTC is at any position within the ORF of the gene, e.g., upstream of the naturally occurring stop codon.

Table 4: Exemplary diseases or disorders

Disease/disorder or protein	Exemplary Point Mutation
G to A point mutations	
Dihydropyrimidine dehydrogenase deficiency	NM 000110.3(DPYD):c.1905+1G>A
Noonan syndrome	NM 005633.3(SOS1):c.2536G>A (p.Glu846Lys)
Lynch syndrome	NM 000251.2(MSH2):c.212-1G>A
Breast-ovarian cancer, familial 1	NM 007294.3(BRCA1):c.963G>A (p.Trp321Ter)
Cystic fibrosis	NM 000492.3(CFTR):c.57G>A (p.Trp19Ter)
Anemia, due to G6PD deficiency	NM 000402.4(G6PD):c.292G>A (p.Val98Met)
AVPR2	NM 000054.4(AVPR2):c.878G>A (p.Trp293Ter)
Nephrogenic diabetes insipidus, X-linked	
FANCC	NM 000054.4(AVPR2):c.878G>A

Fanconi anemia, complementation group C	(p.Trp293Ter)
FANCC Fanconi anemia, complementation group C	NM 000136.2(FANCC):c.1517G>A (p.Trp506Ter)
IL2RG X-linked severe combined immunodeficiency	NM 000206.2(IL2RG):c.710G>A (p.Trp237Ter)
F8 Hereditary factor VIII deficiency disease	NM 000132.3(F8):c.3144G>A (p.Trp1048Ter)
LDLR Familial hypercholesterolemia	NM 000527.4(LDLR):c.1449G>A (p.Trp483Ter)
CBS Homocystinuria due to CBS deficiency	NM 000071.2(CBS):c.162G>A (p.Trp54Ter)
HBB betaThalassemia	NM 000518.4(HBB):c.114G>A (p.Trp38Ter)
ALDOB Hereditary fructosuria	NM 000035.3(ALDOB):c.888G>A (p.Trp296Ter)
DMD Duchenne muscular dystrophy	NM 004006.2(DMD):c.3747G>A (p.Trp1249Ter)
SMAD4 Juvenile polyposis syndrome	NM 005359.5(SMAD4):c.906G>A (p.Trp302Ter)
BRCA2 Familial cancer of breast Breast-ovarian cancer, familial 2	NM 000059.3(BRCA2):c.582G>A (p.Trp194Ter)
GRIN2A Epilepsy, focal, with speech disorder and with or without mental retardation	NM 000833.4(GRIN2A):c.3813G>A (p.Trp1271Ter)
SCN9A Indifference to pain, congenital, autosomal recessive	NM 002977.3(SCN9A):c.2691G>A (p.Trp897Ter)
TARDBP Amyotrophic lateral sclerosis type 10	NM 007375.3(TARDBP):c.943G>A (p.Ala315Thr)
CFTR Cystic fibrosis Hereditary pancreatitis not provided lateral uren response - Efficacy	NM 000492.3(CFTR):c.3846G>A (p.Trp1282Ter)
UBE3A Angelman syndrome	NM 130838.1(UBE3A):c.2304G>A (p.Trp768Ter)
SMPD1 Niemann-Pick disease, type A	NM 000543.4(SMPD1):c.168G>A (p.Trp56Ter)
USH2A Usher syndrome, type 2A	NM 206933.2(USH2A):c.9390G>A (p.Trp3130Ter)
MEN1 Hereditary cancer-predisposing syndrome	NM 130799.2(MEN1):c.1269G>A (p.Trp423Ter)
C8orf37 Retinitis pigmentosa 64	NM 177965.3(C8orf37):c.555G>A (p.Trp185Ter)

MLH1 Lynch syndrome	NM_000249.3(MLH1):c.1998G>A (p.Trp666Ter)
TSC2 Tuberous sclerosis 21Tuberous sclerosis syndrome 46	NM_000548.4(TSC2):c.2108G>A (p.Trp703Ter)
NF1 Neurofibromatosis, type 1	NM_000267.3(NF1):c.7044G>A (p.Trp2348Ter)
MSH6 Lynch syndrome	NM_000179.2(MSH6):c.3020G>A (p.Trp1007Ter)
SMN1 Spinal muscular atrophy, type III Kugelberg- Welander disease	NM_000344.3(SMN1):c.305G>A (p.Trp102Ter)
SH3TC2 Charcot-Marie-Tooth disease, type 4C	NM_024577.3(SH3TC2):c.920G>A (p.Trp307Ter)
DNAH5 Primary ciliary dyskinesia	NM_001369.2(DNAH5):c.8465G>A (p.Trp2822Ter)
MECP2 Rett syndrome	NM_004992.3(MECP2):c.311G>A (p.Trp104Ter)
ADGRV1 Usher syndrome, type 2C	NM_032119.3(ADGRV1):c.7406G>A (p.Trp2469Ter)
AHI1 Joubert syndrome 3	NM_017651.4(AHI1):c.2174G>A (p.Trp725Ter)
PRKN Parkinson disease 2	NM_004562.2(PRKN):c.1358G>A (p.Trp453Ter)
COL3A1 Ehlers-Danlos syndrome, type 4	NM_000090.3(COL3A1):c.3833G>A (p.Trp1278Ter)
BRCA1 Familial cancer ofbreastBreast-ovarian cancer, familial 1	NM_007294.3(BRCA1):c.5511G>A (p.Trp1837Ter)
MYBPC3 Primary familial hypertrophic cardiomyopathy	NM_000256.3(MYBPC3):c.3293G>A (p.Trp1098Ter)
APC Familial adenomatous polyposis 1	NM_000038.5(APC):c.1262G>A (p.Trp421Ter)
BMPR2 Primary pulmonary hypertension	NM_001204.6(BMPR2):c.893G>A (p.W298*)
T to C point mutations	
Wilson disease	NM_000053.3(ATP7B):c.3443T>C (p.Ile 1148Thr)
Leukodystrophy, hypomyelinating, 2	NM_020435.3(GJC2):c.857T>C (p.Met286Thr)
Alport syndrome, X-linked recessive	NM_000495.4(COL4A5):c.438+2T>C
Leigh disease	NC_012920.1:m.9478T>C
Gaucher disease, type 1	NM_001005741.2(GBA):c.751T>C (p.Tyr251His)

Renal dysplasia, retinal pigmentary dystrophy, cerebellar ataxia and skeletal dysplasia	NM_014714.3(IFT140):c.4078T>C (p.Cysl360Arg)
Marfan syndrome	NM_000138.4(FBN1):c.3793T>C (p.Cysl265Arg)
Deficiency of UDPglucose-hexose-1-phosphate uridylyltransferase	NM_000155.3(GALT):c.482T>C (p.Leu161Pro)
Familial hypercholesterolemia	NM_000527.4(LDLR):c.694+2T>C
Episodic pain syndrome, familial, 3	NM_001287223.1(SCN11A):c.1142T>C (p.Ile381Thr)
Navajo neurohepatopathy	NM_002437.4(MPV17):c.186+2T>C
Congenital muscular dystrophy, LMNA-related	NM_170707.3(LMNA):c.1139T>C (p.Leu380Ser)
Hereditary factor VIII deficiency disease	NM_000132.3(F8):c.5372T>C (p.Met1791Thr)
Insulin-dependent diabetes mellitus secretory diarrhea syndrome	NM_014009.3(FOXP3):c.970T>C (p.Phe324Leu)
Hereditary factor IX deficiency disease	NM_000133.3(F9):c.1328T>C (p.Ile443Thr)
Familial cancer of breast, Breast-ovarian cancer, familial 2, Hereditary cancer predisposing syndrome	NM_000059.3(BRCA2):c.316+2T>C
Cardiac arrhythmia	NM_000238.3(KCNH2):c.1945+6T>C
Tangier disease	NM_005502.3(ABCA1):c.4429T>C (p.Cysl477Arg)
Dilated cardiomyopathy IAA	NM_001103.3(ACTN2):c.683T>C (p.Met228Thr)
Mental retardation 3, X-linked	NM_005334.2(HCFC1):c.-970T>C
Limb-girdle muscular dystrophy, type 2B	NM_003494.3(DYSF):c.1284+2T>C
Macular dystrophy, vitelliform, 5	NM_016247.3(IMP2):c.370T>C (p.Phe124Leu)
Retinitis pigmentosa	NM_000322.4(PRPH2):c.736T>C (p.Trp246Arg)

Table 5: Additional exemplary disorders

5q-syndrome	Adams-Oliver syndrome 1
Adams-Oliver syndrome 3	Adams-Oliver syndrome 5
Adams-Oliver syndrome 6	Alagille syndrome 1
Autoimmune lymphoproliferative syndrome type IA	Autoimmune lymphoproliferative syndrome type V

Autosomal dominant deafness-2A	Brain malformations with or without urinary tract defects (BRMUTD)
Carney complex type 1	CHARGE syndrome
Cleidocranial dysplasia	Currarino syndrome
Denys-Drash syndrome / Frasier syndrome	Developmental delay
intellectual disability	obesity
and dysmorphic features (DIDOD)	DiGeorge syndrome (TBX1-associated)
Dravet syndrome	Duane-radial ray syndrome
Ehlers-Danlos syndrome (classic-like)	Ehlers-Danlos syndrome (vascular type)
Feingold syndrome 1	Frontotemporal lobar degeneration with TDP43 inclusions (FTFD-TDP)
GRN-related	GFUT1 deficiency syndrome
Greig cephalopolysyndactyly syndrome	Hereditary hemorrhagic telangiectasia type 1
Holoprosencephaly 3	Holoprosencephaly 4
Holoprosencephaly 5	Holt-Oram syndrome
Hypoparathyroidism	sensorineural deafness
and renal disease (HDR)	Kleefstra syndrome 1
Klippel-Trenaunay syndrome (AAGF-related)	Feri-Weill dyschondrosteosis
Marfan syndrome	Mental retardation and distinctive facial features with or without cardiac defects (MRFACD)
Mental retardation	autosomal dominant 1
Mental retardation	autosomal dominant 19
Mental retardation	autosomal dominant 29
Nail-patella syndrome (NPS)	Phelan-McDermid syndrome
Pitt-Hopkins syndrome	Primary pulmonary hypertension 1

Rett syndrome (congenital variant)	Smith-Magenis syndrome (RAI1-associated)
Sotos syndrome 1	Sotos syndrome 2
Stickler syndrome type I	Supravalvular aortic stenosis
SYNGAP1 -related intellectual disability	Treacher Collins syndrome
Trichorhinophalangeal syndrome type I	Ulnar-mammary syndrome
van der Woude syndrome 1	Waardenburg syndrome type 1
Waardenburg syndrome type 2A	Waardenburg syndrome type 4C.

Table 6: Exemplary genes with ORFs comprising a PTC and exemplary disorders

Gene	Disease/Disorder
AAAS	Glucocorticoid deficiency with achalasia
AAGAB	Keratosis palmoplantaris papulosa
AASS	Hyperlysinemia
ABCA1	Tangier disease
ABCA12, SNHG31	Autosomal recessive congenital ichthyosis 4B
ABCA3	3, Surfactant metabolism dysfunction, pulmonary
ABCA4	Bietti crystalline corneoretinal dystrophy, Cone-rod degeneration, Cone-rod dystrophy 3, Macular dystrophy, Retinal dystrophy, Retinitis pigmentosa, Retinitis pigmentosa 19, Stargardt disease, Stargardt disease 1
ABCB4	Cholestasis, Progressive familial intrahepatic cholestasis 3, intrahepatic, of pregnancy 3
ABCC2	Dubin-Johnson syndrome
ABCC6	Cutis laxa, Generalized arterial calcification of infancy 2, Papule, Pseudoxanthoma elasticum, forme fruste
ABCC8	1, Familial hyperinsulinism, Hyperinsulinemic hypoglycemia, familial
ABCC9	Arrhythmogenic right ventricular cardiomyopathy, Cardiomyopathy, Cardiovascular phenotype, Dilated cardiomyopathy 10, Primary dilated cardiomyopathy
ABCD1	Adrenoleukodystrophy, Spastic gait, Spastic paraplegia
ABHD12	Polyneuropathy, and cataract, ataxia, hearing loss, retinitis pigmentosa
ABRAXAS1	Hereditary breast and ovarian cancer syndrome
ACAD9	Acyl-CoA dehydrogenase family, deficiency of, member 9
ACADM	Medium-chain acyl-coenzyme A dehydrogenase deficiency
ACADS	Deficiency of butyryl-CoA dehydrogenase
ACADVL	Very long chain acyl-CoA dehydrogenase deficiency
ACAN	Osteochondritis dissecans, Spondyloepiphyseal dysplasia, kimberley type
ACAT1	Deficiency of acetyl-CoA acetyltransferase
ACBD5	RETINAL DYSTROPHY WITH LEUKODYSTROPHY
ACBD6, LHX4, LHX4-AS1	Short stature-pituitary and cerebellar defects-small sella turcica syndrome
ACE	Renal dysplasia
ACOX1	Peroxisomal acyl-CoA oxidase deficiency
ACP5	Spondyloenchondrodysplasia with immune dysregulation
ACP5, ZNF627	Spondyloenchondrodysplasia with immune dysregulation
ACTA1	Congenital myopathy with excess of thin filaments
ACTB	Baraitser-Winter syndrome
ACVRL1	Hereditary hemorrhagic telangiectasia type 1, Primary pulmonary hypertension, Pulmonary arterial hypertension related to hereditary hemorrhagic telangiectasia, Telangiectasia, hereditary hemorrhagic, type 2

ACY1	Neurological conditions associated with aminoacylase 1 deficiency
ADA	Severe combined immunodeficiency disease, Severe combined immunodeficiency due to ADA deficiency
ADAM10	Reticulate acropigmentation of Kitamura
ADAMTS17	Weill-Marchesani syndrome 4
ADAMTS2	Ehlers-Danlos syndrome dermatosparaxis type
ADAMTSL4	Ectopia lentis et pupillae
ADAMTSL4	Ectopia lentis, Ectopia lentis 2, Ectopia lentis et pupillae, autosomal recessive, isolated
ADCY3	BODY MASS INDEX QUANTITATIVE TRAIT LOCUS 19
ADCY3, CENPO	BODY MASS INDEX QUANTITATIVE TRAIT LOCUS 19
ADGRG1	Polymicrogyria, bilateral frontoparietal
ADGRG2	Congenital bilateral aplasia of vas deferens from CFTR mutation, Vas deferens, X-linked, congenital bilateral aplasia of
ADGRG6	Arthrogryposis multiplex congenita, Lethal congenital contracture syndrome 9
ADGRV1	4, Febrile seizures, Rare genetic deafness, Retinal dystrophy, Usher syndrome, familial, type 2C
ADNP	Helsmoortel-Van der Aa Syndrome, History of neurodevelopmental disorder, Inborn genetic diseases
AEBP1	2, CLASSIC-LIKE, EHLERS-DANLOS SYNDROME
AGA	Aspartylglucosaminuria
AGK	Sengers syndrome
AGK, DENND11	Cataract, Sengers syndrome, autosomal recessive congenital 5
AGL	Glycogen storage disease, Glycogen storage disease IIIa, Glycogen storage disease IIIb, Glycogen storage disease type III
AGPAT2	Congenital generalized lipodystrophy type 1
AGRN	Congenital myasthenic syndrome
AGT	Renal dysplasia
AGTR1	Renal dysplasia
AGXT	Primary hyperoxaluria, type I
AHDC1	Delayed speech and language development, Global developmental delay, Intellectual disability, Muscular hypotonia, Neonatal hypotonia, Sleep apnea, Xia-Gibbs syndrome
AHI1	Joubert syndrome, Joubert syndrome 3, Retinal dystrophy, Retinitis pigmentosa
AHR	Retinitis pigmentosa 85
AIRE	Autoimmune polyglandular syndrome type 1, Polyglandular autoimmune syndrome, type 1, with reversible metaphyseal dysplasia
ALB	Analbuminemia
ALDH18A1	Cutis laxa-corneal clouding-oligophrenia syndrome
ALDH3A2	Sjögren-Larsson syndrome
ALDH5A1	Succinate-semialdehyde dehydrogenase deficiency
ALDH7A1	Pyridoxine-dependent epilepsy, Seizures
ALDOB	Hereditary fructosuria

ALG1	ALG1-CDG, Congenital disorder of glycosylation
ALG3	ALG3-CDG
ALMS1	Alstrom syndrome
ALOX12B	Autosomal recessive congenital ichthyosis 2
ALPK3	CARDIOMYOPATHY, FAMILIAL HYPERTROPHIC 27, Hypertrophic cardiomyopathy
ALPL	Hypophosphatasia, Infantile hypophosphatasia
ALS2	Amyotrophic lateral sclerosis type 2, Infantile-onset ascending hereditary spastic paralysis, Juvenile primary lateral sclerosis
ALX4	Parietal foramina 2
AMPD2	Pontocerebellar hypoplasia, type 9
AMT	Non-ketotic hyperglycinemia
ANAPC1	Rothmund-Thomson syndrome type 1
ANGPTL3, DOCK7	2, Hypobetalipoproteinemia, familial
ANKRD1	ANKRD1-related dilated cardiomyopathy, Cardiovascular phenotype, Primary dilated cardiomyopathy
ANKRD11	Abnormal facial shape, Clinodactyly of the 5th finger, Conductive hearing impairment, Delayed speech and language development, Global developmental delay, Inborn genetic diseases, Intellectual disability, KBG syndrome, Ptosis, Seizures, Short foot, Short palm, Unilateral cryptorchidism
ANO10	Autosomal recessive cerebellar ataxia, Spinocerebellar ataxia, autosomal recessive 10
ANO5	ANO5-Related Disorders, Achilles tendon contracture, Elevated serum creatine phosphokinase, Gnathodiaphyseal dysplasia, Limb-girdle muscular dystrophy, Lower limb amyotrophy, Lower limb muscle weakness, Miyoshi muscular dystrophy 3, Muscular Diseases, Polycystic kidney dysplasia, type 2L
ANTXR1	Odontotrichomelic syndrome
AP1B1	Autosomal recessive keratitis-ichthyosis-deafness syndrome
AP3B1	Hermansky-Pudlak syndrome 2
AP4B1, AP4B1- AS1	Inborn genetic diseases, Spastic paraplegia 47, autosomal recessive
AP4M1	Spastic paraplegia 50, autosomal recessive
AP5Z1	Spastic paraplegia 48, autosomal recessive
APC	Adenomatous colonic polyposis, Adenomatous polyposis coli with congenital cholesteatoma, Brain tumor-polyposis syndrome 2, Carcinoma of colon, Colon adenocarcinoma, Colorectal cancer, Craniopharyngioma, Desmoid disease, Desmoid tumors, Duodenal polyposis, Familial adenomatous polyposis, Familial adenomatous polyposis 1, Familial multiple polyposis syndrome, Gardner syndrome, Gastric polyposis, Hepatocellular carcinoma, Hereditary cancer-predisposing syndrome, Hyperplastic colonic polyposis, Intestinal polyp, Malignant Colorectal Neoplasm, Neoplasm of stomach, Neoplasm of the large intestine, Periampullary adenoma, hereditary, susceptibility to
APOA1, APOA1- AS	Familial hypoalphalipoproteinemia

APOB	I, Familial hypobetalipoproteinemia, Hypobetalipoproteinemia, familial, normotriglyceridemic
APOC2	APOLIPOPROTEIN C-II (NIJMEGEN), Apolipoprotein C2 deficiency
APOC2, APOC4-APOC2	APOLIPOPROTEIN C-II (PADOVA), Apolipoprotein C2 deficiency
APTIX	Ataxia-oculomotor apraxia type 1
AR	Androgen resistance syndrome, Bulbo-spinal atrophy X-linked, Partial androgen insensitivity syndrome
ARCN1	Short stature, and developmental delay, micrognathia, rhizomelic, with microcephaly
ARG1, MED23	Arginase deficiency
ARHGEF18	Retinitis pigmentosa 78
ARID1A	Mental retardation, autosomal dominant 14
ARID1B	Absent speech, Blepharophimosis, Coffin-Siris syndrome 1, Constipation, Decreased body weight, Failure to thrive, Inborn genetic diseases, Intellectual disability, Long eyelashes, Microcephaly, Recurrent respiratory infections, Seizures, Short stature, Thick lower lip vermillion, Thin upper lip vermillion, moderate
ARID2	COFFIN-SIRIS SYNDROME 6
ARL2BP	Retinitis pigmentosa 82 with or without situs inversus
ARMC2	Male infertility with teratozoospermia due to single gene mutation, SPERMATOGENIC FAILURE 38, Sperm tail anomaly
ARMC2, ARMC2-AS1	Male infertility with teratozoospermia due to single gene mutation, SPERMATOGENIC FAILURE 38
ARMC5	Acth-independent macronodular adrenal hyperplasia 2
ARSA	Metachromatic leukodystrophy, Pseudoarylsulfatase A deficiency, late infantile
ARSB	Metachromatic leukodystrophy, Mucopolysaccharidosis type 6
ART4	Blood group, Dombrock system
ASAH1	Farber disease, Spinal muscular atrophy-progressive myoclonic epilepsy syndrome
ASL	Argininosuccinate lyase deficiency
ASPA, SPATA22	Canavan Disease, Familial Form, Spongy degeneration of central nervous system
ASPM	Microcephaly, Primary autosomal recessive microcephaly, Primary autosomal recessive microcephaly 1, Primary autosomal recessive microcephaly 5
ASS1	Citrullinemia type I
ASXL1	Bohring-Opitz syndrome, Inborn genetic diseases
ASXL3	Bainbridge-Ropers syndrome
ATF6	Achromatopsia, Achromatopsia 7
ATL1	Hereditary spastic paraplegia 3A
ATM	Ataxia-telangiectasia syndrome, Familial cancer of breast, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome, Ovarian Neoplasms
ATM, C11orf65, ATP13A2	Ataxia-telangiectasia syndrome, Ataxia-telangiectasia without immunodeficiency, Breast cancer, Familial cancer of breast, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome, Neoplasm of the breast, susceptibility to Kufor-Rakeb syndrome

ATP1A2	Abnormality of neuronal migration, Arthrogryposis multiplex congenita, Epilepsy, Hydrops fetalis
ATP2A1	Brody myopathy
ATP2C1	Familial benign pemphigus
ATP6V0A2	ALG9 congenital disorder of glycosylation, Cutis laxa with osteodystrophy
ATP6V0A4	Renal tubular acidosis, autosomal recessive, distal
ATP7A	Cutis laxa, Menkes kinky-hair syndrome, X-linked
ATP7B	Inborn genetic diseases, Wilson disease
ATRX	1, Alpha thalassemia-X-linked intellectual disability syndrome, Intellectual disability, Mental retardation-hypotonic facies syndrome, Mental retardation-hypotonic facies syndrome X-linked, X-linked
AXIN2	Oligodontia-colorectal cancer syndrome
B3GALNT1	p phenotype
B3GALNT2	11, Muscular dystrophy-dystroglycanopathy (congenital with brain and eye anomalies), type a
B3GALT6	Spondylo-epi-(meta)-physeal dysplasia
B4GALNT1	Hereditary spastic paraplegia 26, Inborn genetic diseases
B4GALT7	Ehlers-Danlos syndrome progeroid type
B9D1	Joubert syndrome, Meckel syndrome, Meckel-Gruber syndrome, type 9
B9D2	Joubert syndrome
BAG3	BAG3-related, Cardiovascular phenotype, Dilated cardiomyopathy 1HH, Inborn genetic diseases, Myofibrillar myopathy, Primary dilated cardiomyopathy
BAP1	Hereditary cancer-predisposing syndrome, Tumor susceptibility linked to germline BAP1 mutations
BARD1	Breast cancer, Familial cancer of breast, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome, Triple-Negative Breast Cancer Finding, susceptibility to
BBS1	Bardet-Biedl syndrome
BBS1, ZDHHC24	Bardet-Biedl syndrome, Bardet-Biedl syndrome 1
BBS10	Bardet-Biedl syndrome, Bardet-Biedl syndrome 1, Bardet-Biedl syndrome 10, Bardet-biedl syndrome 6/10, Inborn genetic diseases, Retinal dystrophy, Retinitis pigmentosa, digenic
BBS2	Bardet-Biedl syndrome, Bardet-Biedl syndrome 2, Bardet-biedl syndrome 1/2, Bardet-biedl syndrome 2/6, Retinal dystrophy, Retinitis pigmentosa, Retinitis pigmentosa 74, digenic
BBS5	Bardet-Biedl syndrome 5
BBS9	Bardet-Biedl syndrome
BCKDHA	Maple syrup urine disease, Maple syrup urine disease type 1A
BCKDHB	CLASSIC, MAPLE SYRUP URINE DISEASE, Maple syrup urine disease, Maple syrup urine disease type 1B, TYPE 1B
BCOR	Oculofaciocardiodental syndrome
BCS1L	BCS1L-Related Disorders, GRACILE syndrome, Leigh syndrome, Mitochondrial complex III deficiency, Pili torti-deafness syndrome, nuclear type 1
BEST1	Bestrophinopathy, Retinal dystrophy, Vitelliform macular dystrophy type 2, autosomal recessive

BET1	Progressive muscle weakness, Seizures
BFSP1	Cataract 33, multiple types
BLM	Bloom syndrome, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome
BMP1	Osteogenesis imperfecta, type xiii
BMP2	AND SKELETAL ANOMALIES WITH OR WITHOUT CARDIAC ANOMALIES, FACIAL DYSMORPHISM, SHORT STATURE
BMPR1A	Hereditary cancer-predisposing syndrome, Juvenile polyposis syndrome
BMPR2	Primary pulmonary hypertension
BNC1	PREMATURE OVARIAN FAILURE 16
BOLA3	Multiple mitochondrial dysfunctions syndrome 2
BPNT2	Chondrodysplasia with joint dislocations, GPAPP type
BPTF	NEURODEVELOPMENTAL DISORDER WITH DYSMORPHIC FACIES AND DISTAL LIMB ANOMALIES
BRAT1	Inborn genetic diseases, NEURODEVELOPMENTAL DISORDER WITH CEREBELLAR ATROPHY AND WITH OR WITHOUT SEIZURES, Rigidity and multifocal seizure syndrome, lethal neonatal
BRCA1	Breast and/or ovarian cancer, Breast carcinoma, Breast-ovarian cancer, COMPLEMENTATION GROUP S, Dysgerminoma, FANCONI ANEMIA, Familial cancer of breast, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome, Infiltrating duct carcinoma of breast, Neoplasm of ovary, Neoplasm of the breast, Ovarian Neoplasms, Ovarian Serous Surface Papillary Adenocarcinoma, Ovarian cancer, Pancreatic cancer, Pancreatic cancer 4, Porokeratosis punctata palmaris et plantaris, Rhabdomyosarcoma (disease), bilateral breast cancer, breast cancer, familial 1, susceptibility to
BRCA2	Asthma, BRCA2-Related Disorders, Breast and/or ovarian cancer, Breast carcinoma, Breast-ovarian cancer, Cancer of the pancreas, Colorectal cancer, Diffuse intrinsic pontine glioma, Ectopic ossification, Familial cancer of breast, Fanconi anemia, Focal seizures, Genetic non-acquired premature ovarian failure, Glioma susceptibility 3, Headache, Hereditary Cancer Syndrome, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome, Inborn genetic diseases, Malignant tumor of prostate, Medulloblastoma, Migraine, Muscle weakness, Neoplasm of the breast, Nephrolithiasis, Obesity, Ovarian Neoplasms, Ovarian cancer, Pancreatic cancer 2, Polydactyly, Short attention span, Striae distensae, Tracheoesophageal fistula, Tumor susceptibility linked to germline BAP1 mutations, Wilms tumor 1, complementation group D1, familial 1, familial 2
BRIP1	BRIP1-Related Disorders, Breast cancer, Carcinoma of colon, Familial cancer of breast, Fanconi anemia, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome, Neoplasm of ovary, Neoplasm of the breast, Ovarian Cancers, Ovarian Neoplasms, Tracheoesophageal fistula, complementation group J, early-onset
BRWD3	Mental retardation, X-linked 93
BSND	Bartter disease type 4a
BTD	Biotinidase deficiency
BTK	Agammaglobulinemia, X-linked agammaglobulinemia, X-linked agammaglobulinemia with growth hormone deficiency, non-Bruton type

C11orf65, ATM	Ataxia-telangiectasia syndrome, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome
C12orf4	AUTOSOMAL RECESSIVE 66, Attention deficit hyperactivity disorder, Intellectual disability, MENTAL RETARDATION, Muscular hypotonia
C12orf65	Combined oxidative phosphorylation deficiency 7, Spastic paraplegia
C19orf12	Neurodegeneration with brain iron accumulation 4, Spastic paraplegia 43, autosomal recessive
C1QB	C1q deficiency
C1S	Complement component c1s deficiency
C2	Complement component 2 deficiency
C2CD3	Orofaciodigital syndrome xiv
C5	Leiner disease
C6	Complement component 6 deficiency, Immunodeficiency due to a late component of complement deficiency
C7	Complement component 7 deficiency
C8B	Complement component 6 deficiency, Type II complement component 8 deficiency
C8orf37	Cone-rod dystrophy 16
C8orf37	Retinitis pigmentosa 64
CA2	Osteopetrosis with renal tubular acidosis
CABP4	Congenital stationary night blindness, type 2B
CACNA1A	42, Bulbar palsy, Epileptic encephalopathy, Episodic ataxia, Episodic ataxia type 2, Recurrent respiratory infections, and epilepsy, early infantile, type 2
CACNA1C	Long QT syndrome
CACNA2D4	Abnormality of the eye, Retinal cone dystrophy 4
CAPN1	Spastic paraplegia 76, autosomal recessive
CAPN3	Absent Achilles reflex, Absent muscle fiber calpain-3, Arrhythmia, Calf muscle hypertrophy, Congenital muscular dystrophy, Contractures of the joints of the lower limbs, Difficulty walking, EMG: myopathic abnormalities, EMG: neuropathic changes, Elbow flexion contracture, Elevated serum creatine phosphokinase, Limb-Girdle Muscular Dystrophy, Limb-girdle muscle weakness, Limb-girdle muscular dystrophy, Migraine, Muscle weakness, Muscular Diseases, Muscular dystrophy, Myositis, Paresthesia, Positive Romberg sign, Progressive spinal muscular atrophy, Recessive, Shoulder girdle muscle weakness, eosinophilic, type 2A
CASK	Mental retardation and microcephaly with pontine and cerebellar hypoplasia
CASP14	Ichthyosis, autosomal recessive 12, congenital
CASQ2	2, Ventricular tachycardia, catecholaminergic polymorphic
CASR	Hypocalciuric hypercalcemia, Inborn genetic diseases, familial, type 1
CAST	Peeling skin with leukonychia, acral punctate keratoses, and knuckle pads, cheilitis
CAST, ERAP1	Peeling skin with leukonychia, acral punctate keratoses, and knuckle pads, cheilitis
CAT	Acatalasemia, Acatalasia, japanese type
CATSPER1	Spermatogenic failure 7
CAV1	Lipodystrophy, congenital generalized, type 3

CAV3, SSUH2	Long QT syndrome
CBL	Noonan syndrome-like disorder with or without juvenile myelomonocytic leukemia
CBS	CYSTATHIONINE BETA-SYNTHETASE POLYMORPHISM, Classic homocystinuria, Homocystinuria
CC2D1A	Mental Retardation, Mental retardation, Psychosocial, autosomal recessive 3
CC2D2A	Joubert syndrome, Joubert syndrome 9, Meckel syndrome type 6, Meckel-Gruber syndrome
CCBE1	Hennekam lymphangiectasia-lymphedema syndrome 1
CCDC103	Primary ciliary dyskinesia
CCDC28B	Bardet-Biedl syndrome, Bardet-Biedl syndrome 1, modifier of
CCDC39	14, Ciliary dyskinesia, Primary ciliary dyskinesia, primary
CCDC40	15, Ciliary dyskinesia, Primary ciliary dyskinesia, primary
CCDC47	Global developmental delay with dysmorphic features, Trichohepatoneurodevelopmental syndrome, and woolly hair, liver dysfunction, pruritus
CCDC65	27, Ciliary dyskinesia, Kartagener syndrome, Primary ciliary dyskinesia, primary
CCDC78	4, Myopathy, centronuclear
CCDC88C	Congenital hydrocephalus 1
CCN6	Progressive pseudorheumatoid dysplasia
CCNH, RASA1	Capillary malformation-arteriovenous malformation
CCNO	29, Ciliary dyskinesia, Kartagener syndrome, Primary ciliary dyskinesia, primary
CCNQ	Syndactyly-telecanthus-anogenital and renal malformations syndrome
CD19	Common variable immunodeficiency 3
CD247	Immunodeficiency due to defect in cd3-zeta
CD36	Malaria, Platelet glycoprotein IV deficiency, cerebral, susceptibility to
CD46	Atypical hemolytic-uremic syndrome 2
CD55	CROMER BLOOD GROUP SYSTEM, Dr(a-) PHENOTYPE, Protein-losing enteropathy (disease)
CDC14A	Deafness, Rare genetic deafness, autosomal recessive 32
CDC73	Parathyroid adenoma, Parathyroid carcinoma
CDH1	Blepharocheilodontic syndrome 1, Breast cancer, Endometrial carcinoma, Familial cancer of breast, Hereditary cancer-predisposing syndrome, Hereditary diffuse gastric cancer, Malignant tumor of prostate, Neoplasm of ovary, lobular
CDH11	Brachioskeletogenital syndrome
CDH23	Deafness, Inborn genetic diseases, MULTIPLE TYPES, PITUITARY ADENOMA 5, Rare genetic deafness, Usher syndrome type 1D, autosomal recessive 12
CDH23, C10orf105	Rare genetic deafness
CDH23, CDH23-AS1	DIGENIC, TYPE ID/F, USHER SYNDROME, Usher syndrome type 1, Usher syndrome type 1D
CDH3	Congenital hypotrichosis with juvenile macular dystrophy, EEM syndrome, Hypotrichosis with juvenile macular dystrophy, Macular dystrophy

CDHR1	Cone-rod dystrophy 15, Leber congenital amaurosis, Retinal dystrophy, Retinitis pigmentosa 65
CDK10	AL KAISSI SYNDROME
CDK13	Congenital heart defects, and intellectual developmental disorder, dysmorphic facial features
CDK5RAP2	Primary autosomal recessive microcephaly 3
CDKL5	Angelman syndrome-like, Atypical Rett syndrome, Early infantile epileptic encephalopathy 2, Epileptic encephalopathy, Inborn genetic diseases
CDKN2A	Hereditary cancer-predisposing syndrome, Hereditary cutaneous melanoma, Melanoma-pancreatic cancer syndrome, Neoplasm
CDSN, PSORS1C1	Peeling skin syndrome 1
CEL	Maturity-onset diabetes of the young type 8
CELA2A	Coronary artery disease, Diabetes, Familial partial lipodystrophy 6, Hypertensive disorder, Hypertriglyceridemia
CENPF	Stromme syndrome
CENPJ	Congenital microcephaly, Intellectual disability, Perisylvian polymicrogyria, Primary autosomal recessive microcephaly, Primary autosomal recessive microcephaly 1, Primary autosomal recessive microcephaly 6, Seckel syndrome 4, Type III lissencephaly, moderate
CEP120	JOUBERT SYNDROME 31
CEP152	Seckel syndrome
CEP290	Abnormality of the kidney, Bardet-Biedl syndrome 14, Blindness, CEP290-Related Disorders, Cerebellar cyst, Cerebellar vermis hypoplasia, Global developmental delay, Hyperechogenic kidneys, Joubert syndrome, Joubert syndrome 5, Leber congenital amaurosis 10, Meckel syndrome, Meckel-Gruber syndrome, Nephronophthisis, Polycystic kidney dysplasia, Retinal dystrophy, Senior-Loken syndrome 6, type 4
CEP290, C12orf29	Bardet-Biedl syndrome 14, Joubert syndrome, Joubert syndrome 5, Meckel-Gruber syndrome, Nephronophthisis
CEP41	Joubert syndrome 15
CEP78	Cone-rod degeneration, Cone-rod dystrophy and hearing loss 1, Sensorineural hearing loss
CFAP251	Male infertility with teratozoospermia due to single gene mutation, Non-syndromic male infertility due to sperm motility disorder, SPERMATOGENIC FAILURE 18, SPERMATOGENIC FAILURE 33, asthenozoospermia, dysplasia of the mitochondrial sheath, multiple morphologic abnormalities of the sperm flagellum
CFAP410	Axial spondylometaphyseal dysplasia, RETINAL DYSTROPHY WITH OR WITHOUT MACULAR STAPHYLOMA
CFAP43	SPERMATOGENIC FAILURE 19
CFAP44	SPERMATOGENIC FAILURE 20
CFHR5	CFHR5 deficiency
CFTR	Bronchiectasis with or without elevated sweat chloride 1, CFTR-related disorders, Congenital bilateral aplasia of vas deferens from CFTR mutation, Cystic fibrosis, Hereditary pancreatitis, Inborn genetic diseases, ataluren response - Efficacy
CFTR, CFTR-AS1	CFTR-related disorders, Congenital bilateral aplasia of vas deferens from CFTR mutation, Cystic fibrosis

CFTR, LOC111674472	Bronchiectasis with or without elevated sweat chloride 1, CFTR-related disorders, Congenital bilateral aplasia of vas deferens from CFTR mutation, Cystic fibrosis, Hereditary pancreatitis
CFTR, LOC111674475	Bronchiectasis with or without elevated sweat chloride 1, CFTR-related disorders, Congenital bilateral aplasia of vas deferens from CFTR mutation, Cystic fibrosis, Hereditary pancreatitis, Inborn genetic diseases, ataluren response - Efficacy
CFTR, LOC111674477	Cystic fibrosis
CFTR, LOC113633877	Bronchiectasis with or without elevated sweat chloride 1, CFTR-related disorders, Congenital bilateral aplasia of vas deferens from CFTR mutation, Cystic fibrosis, Hereditary pancreatitis
CFTR, LOC113664106	Bronchiectasis with or without elevated sweat chloride 1, Congenital bilateral aplasia of vas deferens from CFTR mutation, Cystic fibrosis, Hereditary pancreatitis
CHD2	CHD2-Related Disorder, Epileptic encephalopathy, History of neurodevelopmental disorder, childhood-onset
CHD7	CHARGE association, Hypogonadism with anosmia, Hypogonadotropic hypogonadism 5 with or without anosmia
CHEK2	3, Astrocytoma, B Lymphoblastic Leukemia/Lymphoma, Breast and colorectal cancer, Breast cancer, CHEK2-Related Cancer Susceptibility, Colitis, Congenital heart defects, Diffuse intrinsic pontine glioma, Familial cancer of breast, Hematochezia, Hereditary breast and ovarian cancer syndrome, Hereditary cancer, Hereditary cancer-predisposing syndrome, Inflammation of the large intestine, Leiomyosarcoma, Li-Fraumeni syndrome, Li-Fraumeni syndrome 2, Malignant tumor of prostate, Neoplasm of the breast, Not Otherwise Specified, Osteosarcoma, Ovarian Neoplasms, Prostate cancer, Thrombocytopenia, multiple types, somatic, susceptibility to
CHM	Retinal dystrophy
CHRD1	Megalocornea
CHRNA1	Congenital myasthenic syndrome
CHRNA2	Autosomal dominant nocturnal frontal lobe epilepsy
CHRNA3	CHRNA3-related condition
CHRND	Lethal multiple pterygium syndrome
CHRNE	4a, Congenital myasthenic syndrome, Congenital myasthenic syndrome 4C, Myasthenic syndrome, congenital, slow-channel
CHRNE, C17orf107	4a, 4b, Congenital myasthenic syndrome, Congenital myasthenic syndrome 4C, Myasthenic syndrome, congenital, fast-channel, slow-channel
CHRNA3	Autosomal recessive multiple pterygium syndrome, CHRNA3-Related Disorders, Inborn genetic diseases, Lethal multiple pterygium syndrome
CHST14	Ehlers-Danlos syndrome, musculocontractural type
CHST3	Spondyloepiphyseal dysplasia with congenital joint dislocations
CHSY1	Temtamy preaxial brachydactyly syndrome
CIB1	3, EPIDERMODYSPLASIA VERRUCIFORMIS, SUSCEPTIBILITY TO
CIITA	Bare lymphocyte syndrome 2
CKAP2L	Filippi syndrome

CLCN1	Autosomal dominant intermediate Charcot-Marie-Tooth disease, Congenital myotonia, EMG: myopathic abnormalities, Muscular Diseases, Myotonia congenita, autosomal dominant form, autosomal recessive form
CLCN2	Epilepsy, Leukoencephalopathy with ataxia, juvenile myoclonic 8
CLCN5	Nephrolithiasis, X-linked recessive, X-linked recessive nephrolithiasis with renal failure
CLDN1, CLDN16	Neonatal ichthyosis-sclerosing cholangitis syndrome
CLIC5	Deafness, autosomal recessive
CLN3	Juvenile neuronal ceroid lipofuscinosis, Neuronal ceroid lipofuscinosis
CLN5, FBXL3	Neuronal ceroid lipofuscinosis, Neuronal ceroid lipofuscinosis 5
CLRN1	Rare genetic deafness, Retinal dystrophy, Retinitis pigmentosa, Usher syndrome, type 3A
CNGA1, LOC101927157	Retinal dystrophy, Retinitis pigmentosa 49
CNGB1	Retinal dystrophy, Retinitis pigmentosa, Retinitis pigmentosa 45
CNGB3	Abnormality of the eye, Achromatopsia, Achromatopsia 3, CNGB3-Related Disorders, Cone-rod dystrophy, Leber congenital amaurosis, Recessive, Retinal dystrophy, Retinitis pigmentosa, Stargardt Disease
CNNM2	Hypomagnesemia 6, renal
CNNM4	Jalili syndrome
CNTNAP1	Lethal congenital contracture syndrome 7
CNTNAP2	Pitt-Hopkins-like syndrome 1
COASY	Neurodegeneration with brain iron accumulation 6
COG4	Congenital disorder of glycosylation type 2J
COG5	Congenital disorder of glycosylation type 2i
COG5, DUS4L, DUS4L-BCAP29	Congenital disorder of glycosylation type 2i
COL10A1	Metaphyseal chondrodysplasia, Schmid type
COL11A1	Fibrochondrogenesis 1
COL12A1	Ullrich congenital muscular dystrophy 2
COL17A1	Epidermolysis bullosa, Junctional epidermolysis bullosa, junctional, localisata variant, non-Herlitz type
COL18A1	GLAUCOMA, Knobloch syndrome 1, PRIMARY CLOSED-ANGLE
COL18A1, SLC19A1	Knobloch syndrome 1, Macular dystrophy, Retinal dystrophy, Retinitis pigmentosa
COL1A1	Ehlers-Danlos syndrome, Infantile cortical hyperostosis, Osteogenesis imperfecta, Osteogenesis imperfecta type I, Osteogenesis imperfecta type III, Osteogenesis imperfecta with normal sclerae, Postmenopausal osteoporosis, dominant form, procollagen proteinase deficient, recessive perinatal lethal
COL1A2	COL1A2-Related Disorder, Ehlers-Danlos syndrome, Inborn genetic diseases, Osteogenesis imperfecta type I, autosomal recessive, cardiac valvular form, classic type
COL2A1	Spondyloperipheral dysplasia-short ulna syndrome, Stickler syndrome type 1
COL3A1	Ehlers-Danlos syndrome, type 4

COL4A3, MFF-DT	Alport syndrome, autosomal recessive
COL4A5	Alport syndrome 1, X-linked recessive
COL5A1	Ehlers-Danlos syndrome, classic type
COL5A2	Ehlers-Danlos syndrome, Ehlers-Danlos syndrome classic type 2, classic type
COL6A1	Bethlem myopathy 1
COL6A2	Bethlem myopathy 1, Ullrich congenital muscular dystrophy 1
COL6A3	Bethlem myopathy 1
COL7A1	Dystrophic epidermolysis bullosa, Epidermolysis bullosa pruriginosa, Recessive dystrophic epidermolysis bullosa, Transient bullous dermolysis of the newborn, autosomal dominant
COL9A2	Stickler syndrome, type 5
COLEC10	3MC syndrome 3
COLEC10, LOC101927513	3MC syndrome 3
COLQ	Congenital myasthenic syndrome, Endplate acetylcholinesterase deficiency
COQ2	Coenzyme Q10 deficiency, primary, primary 1
COQ8A	4, ADCK3-Related Disorders, Coenzyme Q10 deficiency, primary
COQ9	5, Coenzyme Q10 deficiency, primary
COX15	Cardioencephalomyopathy, Leigh syndrome, Leigh syndrome due to mitochondrial complex IV deficiency, due to cytochrome c oxidase deficiency 2, fatal infantile
CP	Ceruloplasmin belfast, Deficiency of ferroxidase, Hemosiderosis, due to aceruloplasminemia, systemic
CPAMD8	Anterior segment dysgenesis 8
CPLANE1	Global developmental delay, Jaundice, Joubert syndrome, Joubert syndrome 1, Joubert syndrome 17, Orofaciodigital syndrome type 6, Typical Joubert syndrome MRI findings
CPOX	Coproporphyrria
CPS1	Congenital hyperammonemia, type I
CPSF1	MYOPIA 27
CPT2	Carnitine palmitoyltransferase II deficiency, infantile, lethal neonatal, myopathic, stress-induced
CRB1	Leber congenital amaurosis 8
CRB2	Focal segmental glomerulosclerosis 9, Steroid-resistant nephrotic syndrome
CRIPT	Ateleiotic dwarfism, Short stature with microcephaly and distinctive facies
CRPPA	7, Congenital muscular dystrophy-dystroglycanopathy with brain and eye anomalies, Muscular dystrophy-dystroglycanopathy (limb-girdle), type A7, type c
CRTAP	Osteogenesis imperfecta type 7
CRX	Leber congenital amaurosis 7
CRYAB	Alpha-B crystallinopathy, Dilated cardiomyopathy III
CRYBA4, CRYBB1	Cataract, autosomal recessive 3, congenital nuclear
CRYBB2	Cataract 3, Congenital cataract, multiple types

CSGALNACT1	MILD, SKELETAL DYSPLASIA, WITH JOINT LAXITY AND ADVANCED BONE AGE
CSPP1	Joubert syndrome 21, Meckel-Gruber syndrome
CSRP3	Cardiovascular phenotype
CSTB	Inborn genetic diseases, Progressive myoclonic epilepsy, Unverricht-Lundborg syndrome
CTC1	Cerebroretinal microangiopathy with calcifications and cysts, Cerebroretinal microangiopathy with calcifications and cysts 1, Dyskeratosis congenita
CTCF	Mental retardation, autosomal dominant 21
CTNNB1	EXUDATIVE VITREORETINOPATHY 7, Exudative vitreoretinopathy 1, Hepatocellular carcinoma, Inborn genetic diseases, Mental retardation, autosomal dominant 19
CTNND1, TMX2-CTNND1	Blepharocheilodontic syndrome 2
CTNS	Cystinosis, Juvenile nephropathic cystinosis, Nephropathic cystinosis, Ocular cystinosis
CTSD	Neuronal ceroid lipofuscinosis 10
CTSH	Variant of unknown significance
CTU2	AND AMBIGUOUS GENITALIA SYNDROME, FACIAL DYSMORPHISM, MICROCEPHALY, RENAL AGENESIS
CUBN	Megaloblastic anemia due to inborn errors of metabolism
CUL4B	Cabezas type, Syndromic X-linked mental retardation
CUL7	Three M syndrome 1
CWC27	Retinitis pigmentosa with or without skeletal anomalies
CWF19L1	Spinocerebellar ataxia, autosomal recessive 17
CYB5R3	Methemoglobinemia type 2
CYBB	Chronic granulomatous disease, X-linked
CYP11B1, LOC106799833	Deficiency of steroid 11-beta-monooxygenase
CYP17A1	20-lyase deficiency, Combined partial 17-alpha-hydroxylase/17, Complete combined 17-alpha-hydroxylase/17, Deficiency of steroid 17-alpha-monooxygenase
CYP1B1	A, Anterior segment dysgenesis 6, CYP1B1-Related Disorders, Congenital glaucoma, Congenital ocular coloboma, Glaucoma, Glaucoma 3, Irido-corneo-trabecular dysgenesis, b, congenital, primary congenital, primary infantile
CYP21A2, LOC106780800	Classic congenital adrenal hyperplasia due to 21-hydroxylase deficiency
CYP21A2, TNXB, LOC106780800	Classic congenital adrenal hyperplasia due to 21-hydroxylase deficiency
CYP24A1	1, Hypercalcemia, infantile
CYP26C1	Optic nerve hypoplasia
CYP27A1	Cholesterol storage disease
CYP27B1	Vitamin D-dependent rickets, type 1

CYP2C19	CYP2C19: no function, Clopidogrel response, Mephenytoin, Proguanil, Toxicity/ADR, amitriptyline response - Efficacy, citalopram response - Efficacy, clomipramine response - Efficacy, clopidogrel response - Efficacy, poor metabolism of
CYP2D6	Debrisoquine, Deutetrabenazine response, Tamoxifen response, Toxicity/ADR, Tramadol response, amitriptyline response - Dosage, antidepressants response - Dosage, clomipramine response - Dosage, desipramine response - Dosage, doxepin response - Dosage, imipramine response - Dosage, nortriptyline response - Dosage, poor metabolism of, tamoxifen response - Efficacy, trimipramine response - Dosage
CYP2U1	Spastic paraplegia 56, autosomal recessive
CYP4F22	Autosomal recessive congenital ichthyosis 5
CZ1P-ASNS, ASNS	Asparagine synthetase deficiency
DBH	Orthostatic hypotension 1
DBT	Maple syrup urine disease, Maple syrup urine disease type 2
DCAF17	Hypogonadism, alopecia, diabetes mellitus, mental retardation and electrocardiographic abnormalities
DCLRE1C	Severe combined immunodeficiency, Severe combined immunodeficiency due to DCLRE1C deficiency, partial
DCN	Congenital Stromal Corneal Dystrophy
DDHD1	Spastic paraplegia 28, autosomal recessive
DDRKG1	Shohat type, Spondyloepimetaphyseal dysplasia
DDX3X	Delayed speech and language development, Global developmental delay, History of neurodevelopmental disorder, Mental retardation, Microcephaly, X-linked 102
DDX41	Acute myeloid leukemia, Myeloproliferative/lymphoproliferative neoplasms, familial (multiple types), susceptibility to
DEPDC5	DEPDC5-Related Disorder, Familial focal epilepsy with variable foci
DES	Muscular dystrophy, Myofibrillar myopathy 1, Neuromuscular disease, Primary dilated cardiomyopathy, limb-girdle, type 2R
DGKE	Nephrotic syndrome, type 7
DGUOK	Mitochondrial DNA depletion syndrome, Mitochondrial DNA-depletion syndrome 3, Progressive external ophthalmoplegia with mitochondrial DNA deletions, autosomal recessive 4, hepatocerebral, hepatocerebral form due to DGUOK deficiency
DHCR7	2-3 toe syndactyly, Congenital microcephaly, Elevated 7-dehydrocholesterol, History of neurodevelopmental disorder, Inborn genetic diseases, Small for gestational age, Smith-Lemli-Opitz syndrome
DHH	46, XY sex reversal, type 7
DHTKD1	2-aminoadipic 2-oxoadipic aciduria
DIAPH1	Seizures, and microcephaly syndrome, cortical blindness
DICER1	DICER1-related pleuropulmonary blastoma cancer predisposition syndrome, Hereditary cancer-predisposing syndrome
DIPK1A, RPL5	Diamond-Blackfan anemia 6
DLD	Maple syrup urine disease, type 3

DLG3	X-Linked mental retardation 90
DLL3, PLEKHG2	Leukodystrophy and acquired microcephaly with or without dystonia, Spondylocostal dysostosis 1, autosomal recessive
DLX3	Amelogenesis imperfecta, Tricho-dento-osseous syndrome, type IV
DLX4	Orofacial cleft 15
DMD	Becker muscular dystrophy, Duchenne muscular dystrophy
DMP1	Autosomal recessive hypophosphatemic vitamin D refractory rickets
DNAAF2	Primary ciliary dyskinesia
DNAAF4, DNAAF4-CCPG1	Primary ciliary dyskinesia
DNAH1	Non-syndromic male infertility due to sperm motility disorder, SPERMATOGENIC FAILURE 18
DNAH11	7, Ciliary dyskinesia, Primary ciliary dyskinesia, primary
DNAH17	SPERMATOGENIC FAILURE 39
DNAH5	3, Ciliary dyskinesia, Primary ciliary dyskinesia, primary
DNAI1	Kartagener syndrome, Primary ciliary dyskinesia
DNAI2	9, Ciliary dyskinesia, Primary ciliary dyskinesia, primary
DNAJB2	5, Charcot-Marie-Tooth disease, Spinal muscular atrophy, autosomal recessive, distal
DNAJC12	Hyperphenylalaninemia, mild, non-bh4-deficient
DNAL1	16, Ciliary dyskinesia, Primary ciliary dyskinesia, primary
DNM2	Charcot-Marie-Tooth disease, dominant intermediate B
DNMBP	CATARACT 48
DOCK6	Adams-Oliver syndrome 2
DOCK6, LOC105372273	Adams-Oliver syndrome, Adams-Oliver syndrome 2
DOCK8	Hyperimmunoglobulin E recurrent infection syndrome, Inborn genetic diseases, autosomal recessive
DOK7	Congenital myasthenic syndrome, Inborn genetic diseases, Myasthenia, Pena-Shokeir syndrome type I, familial, limb-girdle
DOLK	Congenital disorder of glycosylation type 1M
DONSON	AND LIMB ABNORMALITIES, MICROCEPHALY, Microcephaly-micromelia syndrome, SHORT STATURE
DPY19L2	Spermatogenic failure 9
DPYD	Dihydropyrimidine dehydrogenase deficiency, fluorouracil response - Other
DRAM2	Cone-rod dystrophy 21, Retinal dystrophy
DRC1	21, Ciliary dyskinesia, Kartagener syndrome, Primary ciliary dyskinesia, primary
DSC2	11, Arrhythmogenic right ventricular cardiomyopathy, Arrhythmogenic right ventricular dysplasia, familial, type 11, with mild palmoplantar keratoderma and woolly hair
DSC2, DSCAS	Arrhythmogenic right ventricular cardiomyopathy, type 11
DSG1	Palmoplantar keratoderma i, focal, or diffuse, striate
DSG1, DSG1- AS1	Erythroderma, and hyper-ige, congenital, hypotrichosis, with palmoplantar keratoderma

DSG2	Arrhythmogenic right ventricular cardiomyopathy, Cardiac arrest, Cardiomyopathy, Cardiovascular phenotype, Dilated Cardiomyopathy, Dominant, Hypertrophic cardiomyopathy, type 10
DSG2, DSG2-AS1	Dilated cardiomyopathy 1BB
DSG4, DSG1-AS1	Hypotrichosis 6
DSP	Arrhythmogenic right ventricular cardiomyopathy, Arrhythmogenic right ventricular dysplasia/cardiomyopathy, Cardiac arrest, Cardiomyopathy, Cardiovascular phenotype, DSP-Related Disorders, Dilated cardiomyopathy with woolly hair and keratoderma, Keratosis palmoplantaris striata II, Left ventricular noncompaction cardiomyopathy, Lethal acantholytic epidermolysis bullosa, Long QT syndrome 1, Primary dilated cardiomyopathy, Skin fragility-woolly hair-palmoplantar keratoderma syndrome, Ventricular tachycardia, and tooth agenesis, dilated, keratoderma, type 8, with woolly hair
DST	Epidermolysis bullosa simplex, Neuropathy, autosomal recessive 2, hereditary sensory and autonomic, type VI
DUOX2	Congenital hypothyroidism, Familial thyroid dysmorphogenesis, Inborn genetic diseases, Nongoitrous Euthyroid Hyperthyrotropinemia, Thyroid dysmorphogenesis 6
DVL3	Robinow syndrome, autosomal dominant 1, autosomal dominant 3
DYNC2H1	Jeune thoracic dystrophy, Short Rib Polydactyly Syndrome, Short-rib polydactyly syndrome type III, Short-rib thoracic dysplasia 3 with or without polydactyly
DYNC2I1	Short-rib thoracic dysplasia 8 with or without polydactyly
DYNC2I2	Jeune thoracic dystrophy, Short-rib thoracic dysplasia 11 with or without polydactyly
DYNC2LI1	Short-rib thoracic dysplasia 15 with polydactyly
DYRK1A	Mental retardation, autosomal dominant 7
DYSF	Autosomal recessive limb-girdle muscular dystrophy type 2B, Miyoshi muscular dystrophy 1, Myopathy, Qualitative or quantitative defects of dysferlin, distal, with anterior tibial onset
ECEL1	Distal arthrogryposis type 5D, Inborn genetic diseases
ECHS1	Inborn genetic diseases, Mitochondrial short-chain enoyl-coa hydratase 1 deficiency
ECM1	Lipid proteinosis
EDA	Hypohidrotic X-linked ectodermal dysplasia
EDARADD	Ectodermal dysplasia 11b, autosomal recessive, hypohidrotic/hair/tooth type
EDN3	Congenital central hypoventilation, Dominant, Hirschsprung Disease, Hirschsprung disease, Waardenburg syndrome, Waardenburg syndrome type 4B
EDNRB, EDNRB-AS1	Rare genetic deafness
EFEMP2	Autosomal recessive cutis laxa type 1B, Autosomal recessive cutis laxa type 1A
EHMT1	Kleefstra syndrome 1
EIF2AK3	Wolcott-Rallison dysplasia
EIF2AK4	Pulmonary venoocclusive disease 2, autosomal recessive
EIF2B2	Leukoencephalopathy with vanishing white matter, Ovarioleukodystrophy

EIF2S3	MEHMO syndrome
ELN	Inborn genetic diseases, Supravalvar aortic stenosis
ELOVL4	Retinal dystrophy, Stargardt Disease 3
ELP1	Familial dysautonomia
ELP2	ELP2-Related Disorders, Mental retardation, autosomal recessive 58
EMD	Cardiovascular phenotype, Emery-Dreifuss muscular dystrophy 1, Neuromuscular disease, X-linked
ENAM	Amelogenesis imperfecta, Amelogenesis imperfecta - hypoplastic autosomal dominant - local, type IC
ENG	Hereditary hemorrhagic telangiectasia, Hereditary hemorrhagic telangiectasia type 1
ENG, LOC102723566	Hereditary hemorrhagic telangiectasia type 1
EOGT	Adams-Oliver syndrome, Adams-Oliver syndrome 4
EPB42	Spherocytosis type 5
EPCAM	Diarrhea 5, congenital, with tufting enteropathy
EPG5	Vici syndrome
EPHB4	Capillary malformation-arteriovenous malformation 2
EPHB4, SLC12A9	Capillary malformation-arteriovenous malformation 2
EPOR	Primary familial polycythemia due to EPO receptor mutation
ERCC2	Metachromatic leukodystrophy variant, Trichothiodystrophy 1, Xeroderma pigmentosum, group D, photosensitive
ERCC3	Xeroderma pigmentosum, complementation group b
ERCC4	Cockayne syndrome, Fanconi anemia, Hutchinson-Gilford syndrome, Pre-B-cell acute lymphoblastic leukemia, XFE progeroid syndrome, Xeroderma pigmentosum, complementation group Q, group F
ERCC5, BIVM-ERCC5	Xeroderma pigmentosum, group G
ERCC6	Cerebrooculofacioskeletal syndrome 1, Cockayne syndrome B, DE SANCTIS-CACCHIONE SYNDROME
ERCC8	Cockayne syndrome type A
ERCC8, ERCC8-AS1	Cockayne syndrome type A
ERCC8, NDUFAF2	Cockayne syndrome type A, MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 10
ERF	Craniosynostosis 1, Craniosynostosis 4
ERI1	Abnormality of finger, Coarse facial features, Global developmental delay, Unilateral renal agenesis
ESCO2	Roberts-SC phocomelia syndrome
ESRP1	AUTOSOMAL RECESSIVE 109, DEAFNESS
ESRRB	Rare genetic deafness
ETFDH	Multiple acyl-CoA dehydrogenase deficiency
ETHE1	Ethylmalonic encephalopathy
EVC2	Curry-Hall syndrome, Ellis-van Creveld syndrome

EXOSC3	Pontocerebellar hypoplasia, type 1b
EXPH5	Epidermolysis bullosa, autosomal recessive, nonspecific
EXT1	Chondrosarcoma, Multiple congenital exostosis, Multiple exostoses type 1, sporadic
EXT2	Multiple exostoses type 2
EYA1	Branchiootic syndrome, Melnick-Fraser syndrome, Rare genetic deafness
EYA4	Deafness, Dilated cardiomyopathy 1J, Rare genetic deafness, autosomal dominant 10
EYA4, TARID	EYA4-Related Disorders
EYS	Retinal dystrophy, Retinitis pigmentosa, Retinitis pigmentosa 25
F13A1	Factor XIII subunit A deficiency
F13B	Factor XIII, b subunit, deficiency of
F2	Prothrombin deficiency, congenital
F5	Factor V deficiency
F8	Hereditary factor VIII deficiency disease
F9	Hereditary factor IX deficiency disease, Thrombophilia, X-linked, due to factor IX defect
FA2H	Spastic paraplegia 35
FAH	Tyrosinemia type I
FAM161A	Retinal dystrophy, Retinitis pigmentosa, Retinitis pigmentosa 28
FAM20A	Amelogenesis imperfecta type 1G
FANCA	Fanconi anemia, complementation group A
FANCB	Fanconi anemia, complementation group B
FANCC	Fanconi anemia, Hereditary cancer-predisposing syndrome, complementation group C
FANCC, AOPEP	Fanconi anemia, Hereditary cancer-predisposing syndrome, Tracheoesophageal fistula, complementation group C
FANCF	Fanconi anemia, complementation group F
FANCM	Fanconi anemia, Malignant germ cell tumor of ovary, SPERMATOGENIC FAILURE 28
FARS2	Combined oxidative phosphorylation deficiency 14
FARSB	Interstitial lung and liver disease, Rajab interstitial lung disease with brain calcifications
FAS	Autoimmune lymphoproliferative syndrome
FAT4	Van Maldergem syndrome
FBN1	Acromicric dysplasia, Acute aortic dissection, Cardiovascular phenotype, Ectopia lentis, Familial thoracic aortic aneurysm, Familial thoracic aortic aneurysm and aortic dissection, Geleophysic dysplasia 2, Inborn genetic diseases, MASS syndrome, Marfan Syndrome/Loeys-Dietz Syndrome/Familial Thoracic Aortic Aneurysms and Dissections, Marfan lipodystrophy syndrome, Marfan syndrome, Stiff skin syndrome, Weill-Marchesani syndrome 2, autosomal dominant, isolated
FBN1, LOC113939944	Marfan Syndrome/Loeys-Dietz Syndrome/Familial Thoracic Aortic Aneurysms and Dissections, Marfan syndrome
FBXL4	Inborn genetic diseases, Mitochondrial DNA depletion syndrome, Mitochondrial DNA depletion syndrome 13 (encephalomyopathic type)

FERMT1	Kindler's syndrome
FEZF1-AS1, FEZF1	Hypogonadotropic hypogonadism 22 with anosmia
FGD4	Charcot-Marie-Tooth disease, Charcot-Marie-Tooth disease type 4
FGF16	Metacarpal 4-5 fusion
FGF3	Deafness with labyrinthine aplasia microtia and microdontia (LAMM)
FGG	Afibrinogenemia, Hypofibrinogenemia, congenital
FH	Fumarase deficiency, Hereditary cancer-predisposing syndrome, Hereditary leiomyomatosis and renal cell cancer
FIG4	Charcot-Marie-Tooth disease, Charcot-Marie-Tooth disease type 4, Yunis-Varon syndrome, type 4J
FKBP10	Bruck syndrome 1, Osteogenesis imperfecta type 12
FKBP14, FKBP14-AS1	Congenital muscular dystrophy, Ehlers-Danlos syndrome with progressive kyphoscoliosis, Inborn genetic diseases, Joint hypermobility, Muscular hypotonia, Pes valgus, Thoracolumbar scoliosis, and hearing loss, myopathy
FKRP	Limb-girdle muscular dystrophy-dystroglycanopathy, type C5
FKTN	Congenital muscular dystrophy-dystroglycanopathy with brain and eye anomalies, Congenital muscular dystrophy-dystroglycanopathy without mental retardation, FKTN-Related Disorders, Fukuyama congenital muscular dystrophy, Limb-girdle muscular dystrophy-dystroglycanopathy, Walker-Warburg congenital muscular dystrophy, type A4, type B4, type C4
FLCN	Hereditary cancer-predisposing syndrome, Multiple fibrofolliculomas, Pneumothorax, primary spontaneous
FLG	2, Dermatitis, FLG-Related Disorder, Ichthyosis vulgaris, atopic, susceptibility to
FLNA	Periventricular nodular heterotopia 1
FLNB	Spondylocarpotarsal synostosis syndrome
FLNC	26, 4, Cardiomyopathy, Dilated Cardiomyopathy, Dominant, Myofibrillar myopathy, Myopathy, distal, familial hypertrophic, filamin C-related
FLNC, FLNC- AS1	26, 4, Cardiomyopathy, Dilated Cardiomyopathy, Dominant, Myofibrillar myopathy, Myopathy, distal, familial hypertrophic, filamin C-related
FLT4	7, CONGENITAL HEART DEFECTS, MULTIPLE TYPES
FMR1	Intellectual disability
FOXF1	Persistent fetal circulation syndrome
FOXG1	History of neurodevelopmental disorder, Rett syndrome, congenital variant
FOXL2	Blepharophimosis, and epicanthus inversus, and epicanthus inversus syndrome type 1, ptosis
FOXN1	AUTOSOMAL DOMINANT, INFANTILE, T-CELL LYMPHOPENIA, T-cell immunodeficiency, WITH OR WITHOUT NAIL DYSTROPHY, and nail dystrophy, congenital alopecia
FOXP1	Mental retardation with language impairment and with or without autistic features
FOXRED1	Leigh syndrome, Mitochondrial complex I deficiency, nuclear type 1
FRAS1	Fraser syndrome 1
FREM2	Cryptophthalmos, FRASER SYNDROME 2, Fraser syndrome 1, isolated, unilateral or bilateral
FSHB	Hypogonadotropic hypogonadism 24 without anosmia

FSIP2	SPERMATOGENIC FAILURE 34
FSIP2, FSIP2-AS1	SPERMATOGENIC FAILURE 34
FTCD	GLUTAMATE FORMIMINOTRANSFERASE DEFICIENCY
FTSJ1	Mental retardation 9, X-linked
FUCA1	Fucosidosis
FYCO1	Cataract 18
FZD4, PRSS23	Exudative retinopathy, Familial exudative vitreoretinopathy
G6PC	Glycogen storage disease, Glycogen storage disease due to glucose-6-phosphatase deficiency type IA
GAA	Glycogen storage disease, type II
GABRA1	19, Epilepsy, Epileptic encephalopathy, early infantile, juvenile myoclonic 5
GABRA6	GABRA6-Related Disorder
GALC	Galactosylceramide beta-galactosidase deficiency
GALM	GALACTOSEMIA IV
GALNS	MPS-IV-A, Morquio syndrome, Mucopolysaccharidosis
GALT	Deficiency of UDPglucose-hexose-1-phosphate uridylyltransferase
GAMT	Cerebral creatine deficiency syndrome, Deficiency of guanidinoacetate methyltransferase
GAREM2, HADHA	Long-chain 3-hydroxyacyl-CoA dehydrogenase deficiency, Mitochondrial trifunctional protein deficiency
GATA1	Acute megakaryoblastic leukemia
GATA3	Hypoparathyroidism-deafness-renal disease syndrome
GATA6	Abnormality of cardiovascular system morphology, Congenital diaphragmatic hernia, Pancreatic agenesis and congenital heart disease, Persistent truncus arteriosus
GATAD1, PEX1	Deafness enamel hypoplasia nail defects, Peroxisome biogenesis disorder 1A (Zellweger)
GATAD2B	GATAD2B-Related Disorder, Mental retardation, autosomal dominant 18
GBA	Acute neuronopathic Gaucher's disease, Gaucher disease, Gaucher disease type 3C, Gaucher's disease, Subacute neuronopathic Gaucher's disease, type 1
GBA, LOC106627981	Gaucher disease, Gaucher's disease, perinatal lethal, type 1
GBE1	Glycogen storage disease, Glycogen storage disease IV, classic hepatic, fatal perinatal neuromuscular, type IV
GCDH	Glutaric aciduria, type 1
GCH1	Dystonia 5
GCK	Maturity onset diabetes mellitus in young, Maturity-onset diabetes of the young, type 2
GDAP1	Charcot-Marie-Tooth disease, recessive intermediate A, type 4A
GDF1, CERS1	Heterotaxia
GDF9	PREMATURE OVARIAN FAILURE 14
GFER	Mitochondrial diseases

GHR	Laron syndrome with elevated serum GH-binding protein, Laron-type isolated somatotropin defect
GJB1	Charcot-Marie-Tooth Neuropathy X, Charcot-Marie-Tooth disease
GJB2	Bilateral conductive hearing impairment, Bilateral sensorineural hearing impairment, Deafness, Dominant, GJB2-Related Disorders, GJB2/GJB3, GJB2/GJB6, Hearing impairment, Hearing loss, Hystrix-like ichthyosis with deafness, Inborn genetic diseases, Keratitis ichthyosis and deafness syndrome, Keratitis-ichthyosis-deafness syndrome, Knuckle pads, Mutilating keratoderma, Nonsyndromic Hearing Loss, Nonsyndromic hearing loss and deafness, Palmoplantar keratoderma-deafness syndrome, Rare genetic deafness, Recessive, Severe sensorineural hearing impairment, X-linked 2, autosomal dominant, autosomal dominant 3a, autosomal recessive 1A, autosomal recessive 1b, deafness AND leukonychia syndrome, digenic
GJB3	Deafness, autosomal dominant 2b
GLA, RPL36A-HNRNPH2	Fabry disease
GLB1	GLB1-Related Disorders, GM1 gangliosidosis, GM1 gangliosidosis type 2, GM1 gangliosidosis type 3, GM1-gangliosidosis, Infantile GM1 gangliosidosis, MPS-IV-B, Mucopolysaccharidosis, type I, with cardiac involvement
GLDC	Non-ketotic hyperglycinemia
GLDN	Lethal congenital contracture syndrome 11
GLI3	Greig cephalopolysyndactyly syndrome, Pallister-Hall syndrome, Postaxial polydactyly, Preaxial polydactyly 4, type A1/B
GLIS3	Diabetes mellitus, neonatal, with congenital hypothyroidism
GLMN	Glomuvenous malformations
GLRA1	Hyperekplexia 1
GNAS	Progressive osseous heteroplasia, Pseudohypoparathyroidism, Pseudopseudohypoparathyroidism
GNAT2	Achromatopsia 4
GNB5	Intellectual developmental disorder with cardiac arrhythmia, Language delay and attention deficit-hyperactivity disorder/cognitive impairment with or without cardiac arrhythmia
GNPAT	Rhizomelic chondrodysplasia punctata type 2
GNPTAB	GNPTAB-Related Disorders, Inborn genetic diseases, MUCOLIPIDOSIS III ALPHA/BETA, Mucopolipidosis, Mucopolipidosis type II, Pseudo-Hurler polydystrophy
GNPTG	Mucopolipidosis, Mucopolipidosis type III gamma
GORAB	Geroderma osteodysplastica
GOSR2, LRRC37A2	Progressive myoclonic epilepsy
GPC3	Simpson-Golabi-Behmel syndrome, Wilms tumor 1
GPC4	Keipert syndrome
GPC6	Autosomal recessive omodysplasia
GPC6, GPC6-AS2	Autosomal recessive omodysplasia
GPI	Hemolytic anemia, due to glucose phosphate isomerase deficiency, nonspherocytic
GPMB	3, AMYLOIDOSIS, PRIMARY LOCALIZED CUTANEOUS

GPR143	Ocular albinism, type I
GPR179	Congenital stationary night blindness, Retinal dystrophy, type 1E
GPSM2	Chudley-McCullough syndrome, GPSM2-Related Disorders, Rare genetic deafness
GRHL2	Deafness, autosomal dominant 28
GRHL3	Van der Woude syndrome 2
GRHPR	Nephrocalcinosis, Nephrolithiasis, Primary hyperoxaluria, type II
GRIN2B	Mental retardation, autosomal dominant 6
GRIP1	FRASER SYNDROME 3
GRN	Frontotemporal dementia
GRXCR1	Deafness, Rare genetic deafness, autosomal recessive 25
GSDME	Deafness, autosomal dominant 5
GUCY2C, C12orf60	Meconium ileus
GUSB	Mucopolysaccharidosis type 6, Mucopolysaccharidosis type 7
GYG1	Glycogen storage disease XV, Polyglucosan body myopathy 2
GYS1	Glycogen storage disease 0, muscle
GYS2	Glycogen storage disease, Glycogen storage disease due to hepatic glycogen synthase deficiency
GZF1	AND MYOPIA, JOINT LAXITY, SHORT STATURE
H1-4	Inborn genetic diseases, RAHMAN SYNDROME
H6PD	Cortisone reductase deficiency 1
HADHA	HADHA-Related Disorders, Long-chain 3-hydroxyacyl-CoA dehydrogenase deficiency, Mitochondrial trifunctional protein deficiency
HADHA, GAREM2	HADHA-Related Disorders, Inborn genetic diseases, LCHAD Deficiency, Lchad deficiency with maternal acute fatty liver of pregnancy, Long-chain 3-hydroxyacyl-CoA dehydrogenase deficiency, Mitochondrial trifunctional protein deficiency
HAX1	Severe congenital neutropenia 3, autosomal recessive
HBA2, LOC106804612	Alpha plus thalassemia
HBB, LOC106099062, LOC107133510	Anemia, Beta thalassemia major, Beta-plus-thalassemia, Beta-thalassemia, Erythrocytosis 6, Fetal hemoglobin quantitative trait locus 1, HBB-Related Disorders, Hb SS disease, Heinz body anemia, Hemoglobin E, Hemoglobin E disease, Hemoglobin E/beta thalassemia disease, Hemoglobin M disease, Hemoglobinopathy, Malaria, Susceptibility to malaria, alpha Thalassemia, beta Thalassemia, beta ⁰ Thalassemia, dominant inclusion body type, familial, resistance to
HBB, LOC107133510, LOC110006319	beta Thalassemia
HCN4	Brugada syndrome 8, Sick sinus syndrome 2, autosomal dominant
HEXA	Inborn genetic diseases, Tay-Sachs disease
HEXB	Sandhoff disease, infantile
HFM1	Premature ovarian failure 9

HGD	Alkaptonuria
HGSNAT	MPS-III-C, Mucopolysaccharidosis, Retinitis pigmentosa 73, Sanfilippo syndrome
HIVEP2	Angelman syndrome-like, Mental retardation, autosomal dominant 43
HJV	Hemochromatosis type 2A
HLCS	Holocarboxylase synthetase deficiency
HMCN1	Age-related macular degeneration 1
HMGB3	Microphthalmia, syndromic 13
HMGCL	Deficiency of hydroxymethylglutaryl-CoA lyase
HNF1A	20, Clear cell carcinoma of kidney, Diabetes mellitus, Diabetes mellitus type 1, Hepatic adenomas, Maturity onset diabetes mellitus in young, Maturity-onset diabetes of the young, familial, insulin-dependent, type 3
HNF1B	Familial hypoplastic, Renal cysts and diabetes syndrome, glomerulocystic kidney
HNRNPK	AU-KLINE SYNDROME
HNRNPU	Epileptic encephalopathy
HOXA1	Athabaskan brainstem dysgenesis syndrome, Bosley-Salih-Alorainy syndrome
HOXA11	Radioulnar synostosis with amegakaryocytic thrombocytopenia 1
HOXD13	Synpolydactyly 1
HPGD	1, HPGD-Related Disorders, Hypertrophic osteoarthropathy, autosomal recessive, primary
HPS1	Hermansky-Pudlak syndrome, Hermansky-Pudlak syndrome 1
HPS5	Hermansky-Pudlak syndrome, Hermansky-Pudlak syndrome 5
HPS6	Hermansky-Pudlak syndrome, Hermansky-Pudlak syndrome 6
HPSE2	Urofacial syndrome 1
HR	Atrichia with papular lesions
HSD17B10	HSD10 disease
HSD17B4	Bifunctional peroxisomal enzyme deficiency, Perrault syndrome
HSPA9	4, Anemia, Even-plus syndrome, sideroblastic
HSPB1	Charcot-Marie-Tooth disease, Charcot-Marie-Tooth disease axonal type 2F, Distal hereditary motor neuronopathy type 2B
HSPG2	Lethal Kniest-like syndrome, Schwartz-Jampel syndrome
HYAL1	Deficiency of hyaluronoglucosaminidase
HYDIN	5, Ciliary dyskinesia, primary
ICAM4	Landsteiner-Wiener phenotype
IDS	MPS-II, Mucopolysaccharidosis
IDS, LOC106050102	MPS-II, Mucopolysaccharidosis
IDUA	Hurler syndrome, MPS-I-H/S, MPS-I-S, Mucopolysaccharidosis, Mucopolysaccharidosis type 1
IDUA, SLC26A1	Hurler syndrome, MPS-I-H/S, MPS-I-S, Mucopolysaccharidosis, Mucopolysaccharidosis type 1
IFIH1	Aicardi-Goutieres syndrome 7, Singleton-Merten syndrome 1
IFNGR1	Disseminated atypical mycobacterial infection, IFN-gamma receptor 1 deficiency, Immunodeficiency 27b, Inherited Immunodeficiency Diseases
IFNGR2	Immunodeficiency 28

IFT140	Retinitis pigmentosa 80
IFT140, LOC105371046	Jeune thoracic dystrophy, Joubert syndrome with Jeune asphyxiating thoracic dystrophy, Renal dysplasia, cerebellar ataxia and skeletal dysplasia, retinal pigmentary dystrophy
IFT172	Short-rib thoracic dysplasia 10 with or without polydactyly
IFT52	Short Rib Polydactyly Syndrome, Short-rib thoracic dysplasia 16 with or without polydactyly
IGF1	Growth delay due to insulin-like growth factor type 1 deficiency
IGF1R	Inborn genetic diseases
IGFALS	Acid-labile subunit deficiency
IGHM	Agammaglobulinemia, non-Bruton type
IGHMBP2	1, Autosomal dominant distal hereditary motor neuropathy, Charcot-Marie-Tooth disease, Distal spinal muscular atrophy, Inborn genetic diseases, Spinal muscular atrophy, autosomal recessive, axonal, distal, type 2S
IGLL1	Agammaglobulinemia 2, autosomal recessive
IGSF1	Hypothyroidism, and testicular enlargement, central
IGSF3	Lacrimal duct defect
IKBKG	Ectodermal dysplasia and immunodeficiency 1, Immunodeficiency without anhidrotic ectodermal dysplasia, Incontinentia pigmenti, atypical
IL12B	Immunodeficiency 29
IL12RB1	Immunodeficiency 30
IL2RB	Ichthyosis (disease)
IL2RG	Combined immunodeficiency, X-linked, X-linked severe combined immunodeficiency
IL36RN	Pustular psoriasis, generalized
IL7R	B cell-positive, NK cell-positive, Severe combined immunodeficiency, T cell-negative, autosomal recessive
INPP5E	Retinal dystrophy
INPPL1	Opsismodysplasia
INTU	Mohr syndrome, Orofaciodigital syndrome 17
IQCB1	Renal dysplasia and retinal aplasia
IQCE	POLYDACTYLY, POSTAXIAL, TYPE A7
IQSEC2	Mental retardation, Severe intellectual deficiency, X-linked 1
IRAK4	Immunodeficiency due to interleukin-1 receptor-associated kinase-4 deficiency
IRF2BPL	ABNORMAL MOVEMENTS, AND SEIZURES, LOSS OF SPEECH, NEURODEVELOPMENTAL DISORDER WITH REGRESSION, Neurodevelopmental disorder
IRF6	Van der Woude syndrome
IRS4	9, CONGENITAL, HYPOTHYROIDISM, NONGOITROUS
ISCA2	Multiple mitochondrial dysfunctions syndrome 4
ISG15	Immunodeficiency 38 with basal ganglia calcification
ITGA7	Muscular dystrophy, congenital, due to integrin alpha-7 deficiency
ITGB2	Leukocyte adhesion deficiency
ITGB4	Epidermolysis bullosa junctionalis with pyloric atresia
ITPA	35, Epileptic encephalopathy, Inosine triphosphatase deficiency, early infantile

ITPR1	Gillespie syndrome
IVD	Isovaleric acidemia, Isovaleryl-CoA dehydrogenase deficiency, type III
JAG1	Alagille syndrome 1, Arteriohepatic dysplasia, Heart, malformation of
JAK3	B cell-positive, NK cell-negative, Severe combined immunodeficiency, Severe combined immunodeficiency disease, T cell-negative, autosomal recessive
KAT6A	History of neurodevelopmental disorder, Mental retardation, autosomal dominant 32
KAT6B	Blepharophimosis - intellectual disability syndrome, SBBYS type
KAT6B, DUPD1	Blepharophimosis - intellectual disability syndrome, Genitopatellar syndrome, Inborn genetic diseases, SBBYS type
KATNIP	Joubert syndrome 26
KCNA1	Episodic ataxia type 1
KCNA5	7, Atrial fibrillation, familial
KCNC1	Epilepsy, progressive myoclonic 7
KCNE1	Long QT syndrome
KCNH2	Cardiac arrhythmia, Cardiovascular phenotype, Congenital long QT syndrome, Long QT syndrome, Long QT syndrome 1/2, Long QT syndrome 2, digenic
KCNK18	Migraine, with or without aura 13
KCNQ1	Cardiac arrhythmia, Cardiovascular phenotype, Congenital long QT syndrome, Jervell and Lange-Nielsen syndrome, Jervell and Lange-Nielsen syndrome 1, KCNQ1-Related Disorders, LQT1 subtype, Long QT syndrome, Long QT syndrome 1, Rare genetic deafness, Romano-Ward syndrome, recessive
KCNQ1-AS1, KCNQ1	Jervell and Lange-Nielsen syndrome 1
KCNQ1, KCNQ1-AS1	Cardiovascular phenotype, Long QT syndrome, Long QT syndrome 1
KCNQ1, KCNQ1OT1	Congenital long QT syndrome, LQT1 subtype, Long QT syndrome
KCNQ2	Benign familial neonatal seizures 1, Early infantile epileptic encephalopathy, Early infantile epileptic encephalopathy 7, Epileptic encephalopathy, Inborn genetic diseases, Seizures
KCNQ3	Intellectual disability, Seizures
KCNQ4	Autosomal dominant nonsyndromic deafness 2A
KCNT1	5, Early infantile epileptic encephalopathy 14, Epilepsy, nocturnal frontal lobe
KCNV2	Cone dystrophy with supernormal rod response, Progressive cone dystrophy (without rod involvement), Retinal dystrophy, Stargardt disease
KDM5B	Intellectual disability, autosomal recessive 65
KDM5C	Claes-Jensen type, Mental retardation, X-linked, syndromic
KDM6A	Kabuki syndrome 2
KERA	Cornea plana 2
KHDC3L	2, Hydatidiform mole, recurrent
KIAA0586	Congenital cerebellar hypoplasia, Intellectual disability, Joubert syndrome, Joubert syndrome 23, Retinal dystrophy, Rod-cone dystrophy, Short-rib thoracic dysplasia 14 with polydactyly
KIAA0753	Orofaciodigital syndrome XV
KIAA0825	POLYDACTYLY, POSTAXIAL, Postaxial polydactyly type A1, TYPE A10

KIAA1549	RETINITIS PIGMENTOSA 86
KIF11	Microcephaly with or without chorioretinopathy, lymphedema, or mental retardation
KIF7	Acrocallosal syndrome, Joubert syndrome 12
KIFBP	Goldberg-Shprintzen megacolon syndrome
KISS1R	Hypogonadotropic hypogonadism 8 without anosmia
KIZ	Retinitis pigmentosa 69
KMT2A	Wiedemann-Steiner syndrome
KMT2B	Dystonia 28, childhood-onset
KMT2C	Kleefstra syndrome due to a point mutation
KMT2D	CHARGE association, Kabuki syndrome, Kabuki syndrome 1
KMT2E	Epilepsy, Leukoencephalopathy, Macrocephalus, O'DONNELL-LURIA-RODAN SYNDROME, See cases, intellectual deficiency
KPTN	Mental retardation, autosomal recessive 41
KRIT1	Cavernous malformations of CNS and retina, Cerebral cavernous malformation, Cerebral cavernous malformations 1
KRT1	Ichthyosis histrix, curth-macklin type
KRT10	Bullous ichthyosiform erythroderma
KRT10, TMEM99	Bullous ichthyosiform erythroderma
KRT14	Epidermolysis bullosa simplex, autosomal recessive
KRT5	Dowling-Degos disease 1
KRT6A	Pachyonychia congenita 3
KRT85	pure hair-nail type, Ectodermal dysplasia
KYNU	AND LIMB DEFECTS SYNDROME 2, CARDIAC, Congenital NAD deficiency disorder, RENAL, VERTEBRAL
L1CAM	MASA syndrome, Spastic paraplegia
L2HGDH	L-2-hydroxyglutaric aciduria
LACC1	JUVENILE ARTHRITIS
LAMA2	Inborn genetic diseases, Laminin alpha 2-related dystrophy, Merosin deficient congenital muscular dystrophy
LAMA3	Junctional epidermolysis bullosa gravis of Herlitz
LAMA4	Dilated cardiomyopathy 1JJ
LAMB3	Amelogenesis imperfecta, Junctional epidermolysis bullosa, Junctional epidermolysis bullosa gravis of Herlitz, non-Herlitz type, type IA
LAMC2	Junctional epidermolysis bullosa, Junctional epidermolysis bullosa gravis of Herlitz, non-Herlitz type
LAMP2	Cardiomyopathy, Danon disease, Hypertrophic cardiomyopathy, Primary dilated cardiomyopathy
LARGE1	Congenital muscular dystrophy-dystroglycanopathy with mental retardation, type B6
LBR	Disproportionate short stature, Femoral bowing, Pelger-Huët anomaly, Regressive spondylometaphyseal dysplasia, Retrognathia, Rhizomelic arm shortening, Rhizomelic leg shortening, Short long bone
LDB3	Cardiomyopathy, Myofibrillar myopathy, ZASP-related

LDLR	Familial hypercholesterolemia, Familial hypercholesterolemia 1, Homozygous familial hypercholesterolemia
LDLRAP1	Familial hypercholesterolemia 4
LEP	Leptin deficiency or dysfunction
LFNG	Spondylocostal dysostosis 3, autosomal recessive
LGII	Familial temporal lobe epilepsy 1
LHFPL5	Rare genetic deafness
LHX3	Non-acquired combined pituitary hormone deficiency with spine abnormalities
LIFR	StÅ½ve-Wiedemann syndrome
LIG4	LIG4-Related Disorders, Lig4 syndrome
LIPA	Lysosomal acid lipase deficiency
LIPE	Familial partial lipodystrophy 6
LIPE, LIPE-AS1, LOC101930071	Familial partial lipodystrophy 6
LIPH	Hypotrichosis 7, Woolly hair, autosomal recessive 2, with or without hypotrichosis
LIPN	Autosomal recessive congenital ichthyosis 8
LMBR1	Acheiropodia
LMBRD1	Inborn genetic diseases, Methylmalonic aciduria and homocystinuria type cblF
LMNA	Cardiovascular phenotype, Charcot-Marie-Tooth disease, Primary dilated cardiomyopathy, type 2
LMOD3	Nemaline myopathy 10
LMX1B	Nail-patella syndrome
LOC100507346, PTCH1	Gorlin syndrome, Medulloblastoma
LOC101927055, TTN	Dilated cardiomyopathy 1G, Limb-girdle muscular dystrophy, Primary dilated cardiomyopathy, type 2J
LOC101927157, CNGA1	Retinitis pigmentosa, Retinitis pigmentosa 49
LOC101927188, LAMA1	Poretti-Boltshauser syndrome
LOC102723566, ENG	Hereditary hemorrhagic telangiectasia type 1
LOC106694316, MPO	Myeloperoxidase deficiency
LOC110006319, HBB, LOC107133510	beta Thalassemia
LOXHD1	Deafness, Rare genetic deafness, autosomal recessive 77
LPL	Hyperlipoproteinemia, Lpl-arita, type I
LRAT	EARLY-ONSET SEVERE, JUVENILE, LRAT-RELATED, Leber congenital amaurosis, Leber congenital amaurosis 14, RETINAL DYSTROPHY, RETINITIS PIGMENTOSA
LRBA	Common variable immunodeficiency 8, with autoimmunity
LRIT3	Congenital stationary night blindness, type 1F
LRP4	Cenani-Lenz syndactyly syndrome

LRP5	Exudative vitreoretinopathy 4, Familial exudative vitreoretinopathy, autosomal dominant
LRP6	7, Tooth agenesis, selective
LRPAP1	Myopia 23, Rare isolated myopia, autosomal recessive
LRPPRC	Congenital lactic acidosis, Saguenay-Lac-Saint-Jean type
LRSAM1	Charcot-Marie-Tooth disease type 2P
LRTOMT	Deafness, Rare genetic deafness, autosomal recessive 63
LTBP2	Congenital glaucoma, Microspherophakia
LTBP3	Dental anomalies and short stature
LTBP4	Cutis laxa with severe pulmonary, and urinary abnormalities, gastrointestinal
LYRM7	Mitochondrial complex III deficiency, nuclear type 8
LZTFL1	Bardet-Biedl syndrome 17
LZTR1	Noonan syndrome 2, Schwannomatosis 2
MAB21L1, NBEA	AND GENITAL SYNDROME, CEREBELLAR, CRANIOFACIAL, OCULAR
MAFB	Duane retraction syndrome 2, Duane retraction syndrome 3 with or without deafness, Duane syndrome type 1, Duane syndrome type 3
MAGED2	Barter syndrome, antenatal, transient, type 5
MAGEL2	Inborn genetic diseases, Schaaf-Yang syndrome
MAGT1	Epstein-Barr virus infection, Immunodeficiency, X-Linked, and neoplasia, with magnesium defect
MAK	Retinal dystrophy
MAN2B1	Deficiency of alpha-mannosidase
MANBA	Beta-D-mannosidosis
MAP2K2	Rasopathy
MAPRE2	2, Skin creases, congenital symmetric circumferential
MARVELD2	Deafness, Rare genetic deafness, autosomal recessive 49, neurosensory
MAX	Hereditary cancer-predisposing syndrome
MBD5	Mental retardation, autosomal dominant 1
MC2R	ACTH resistance
MC4R	Monogenic diabetes, Obesity, Schizophrenia
MCCC1	3 Methylcrotonyl-CoA carboxylase 1 deficiency
MCCC2	3-methylcrotonyl CoA carboxylase 2 deficiency
MCM5	MEIER-GORLIN SYNDROME 8
MCM8	Premature ovarian failure 10
MCOLN1	Mucopolipidosis type IV
MCPH1	Abnormality of brain morphology, Primary autosomal recessive microcephaly 1
MECP2	Angelman syndrome, Atypical Rett syndrome, Autism, Delayed gross motor development, Delayed speech and language development, Developmental regression, Encephalopathy, Global developmental delay, History of neurodevelopmental disorder, Inborn genetic diseases, Intellectual disability, Loss of ability to walk, Mental retardation, Rett syndrome, Severe neonatal-onset encephalopathy with microcephaly, Smith-Magenis Syndrome-like, Syndromic X-linked intellectual disability Lubs type, X-linked, X-linked 3, neonatal

	severeMental retardation, susceptibility to, syndromic 13, syndromic 13Rett syndrome
MED12	Cardiovascular phenotype, FG syndrome 1, History of neurodevelopmental disorder
MED13L	Mental retardation and distinctive facial features with or without cardiac defects
MED25	Broad-based gait, Charcot-Marie-Tooth disease, Decreased body weight, Failure to thrive, Generalized hypotonia, Impaired distal proprioception, Sensory ataxia, Sensory ataxic neuropathy, Sensory neuropathy, type 2
MEF2C	MEF2C-Related Disorder, Mental retardation, and/or cerebral malformations, epilepsy, stereotypic movements
MEFV	Familial Mediterranean fever
MEN1	Hereditary cancer-predisposing syndrome, Lipoma, Multiple endocrine neoplasia, somatic, type 1
MERTK	Retinitis pigmentosa 38
MESD	OSTEOGENESIS IMPERFECTA, TYPE XX
METTL23	Inborn genetic diseases, Mental retardation, autosomal recessive 44
MFN2	Charcot-Marie-Tooth disease, type 2
MFRP, C1QTNF5	Microphthalmia, Nanophthalmos 2, isolated 5
MFSD8	Neuronal ceroid lipofuscinosis 7
MIP	Cataract 15, multiple types
MIR6886, LDLR	Familial hypercholesterolemia, Familial hypercholesterolemia 1, Homozygous familial hypercholesterolemia
MITF	Coloboma, Rare genetic deafness, Waardenburg syndrome type 2A, albinism, and deafness, macrocephaly, microphthalmia, osteopetrosis
MKRN3	2, Precocious puberty, central
MKS1	Joubert syndrome, Joubert syndrome 28, Meckel syndrome type 1, Meckel-Gruber syndrome
MLC1	Megalencephalic leukoencephalopathy with subcortical cysts 1
MLH1	Carcinoma of colon, Colon cancer, Hereditary cancer-predisposing syndrome, Hereditary nonpolyposis colon cancer, Lynch syndrome, Lynch syndrome I, Lynch syndrome II, Muir-Torr� syndrome, Turcot syndrome
MLH3	Hereditary nonpolyposis colorectal cancer type 7
MLYCD	Deficiency of malonyl-CoA decarboxylase
MMAA	Methylmalonic acidemia, Vitamin B12-responsive methylmalonic acidemia type cblA
MMAB	Methylmalonic acidemia, Vitamin B12-responsive methylmalonic acidemia type cblB
MMACHC	DIGENIC, Disorders of Intracellular Cobalamin Metabolism, METHYLMALONIC ACIDURIA AND HOMOCYSTINURIA, Methylmalonic acidemia with homocystinuria, Methylmalonic aciduria due to methylmalonyl-CoA mutase deficiency, cblC TYPE

MME	Charcot-Marie-Tooth disease, Congenital membranous nephropathy due to fetomaternal anti-neutral endopeptidase alloimmunization, axonal, type 2T
MMUT	Methylmalonic acidemia, Methylmalonic aciduria due to methylmalonyl-CoA mutase deficiency
MOCS2	Molybdenum cofactor deficiency, complementation group B
MPDZ	2, Congenital hydrocephalus, Hydrocephalus, congenital, with or without brain or eye anomalies
MPL	Congenital amegakaryocytic thrombocytopenia, essential thrombocytemia
MPLKIP	Trichothiodystrophy, nonphotosensitive 1
MPO	Myeloperoxidase deficiency
MPV17	Navajo neurohepatopathy
MPZ	Charcot-Marie-Tooth disease
MPZL2	AUTOSOMAL RECESSIVE 111, DEAFNESS
MRE11	Hereditary cancer-predisposing syndrome
MSH2	Carcinoma of colon, Colon cancer, Glioblastoma, Hereditary cancer-predisposing syndrome, Hereditary nonpolyposis colon cancer, Lynch syndrome, Lynch syndrome I, Malignant tumor of ascending colon, Malignant tumor of sigmoid colon, Muir-TorrÃ© syndrome, Ovarian Neoplasms, Turcot syndrome
MSH6	Endometrial carcinoma, Hereditary cancer-predisposing syndrome, Hereditary nonpolyposis colon cancer, Hereditary nonpolyposis colorectal cancer type 5, Hereditary nonpolyposis colorectal carcinoma, Lynch syndrome, Lynch syndrome I, Turcot syndrome
MSTO1	Mitochondrial myopathy-cerebellar ataxia-pigmentary retinopathy syndrome
MSX2	Parietal foramina 1
MTFMT	Abnormal facial shape, Combined oxidative phosphorylation deficiency 15, Cytochrome C oxidase-negative muscle fibers, Decreased activity of mitochondrial complex I, Inability to walk by childhood/adolescence, Leigh syndrome, MITOCHONDRIAL COMPLEX I DEFICIENCY, Mitochondrial oxidative phosphorylation disorder, NUCLEAR TYPE 27, Poor speech, Short stature
MTHFD1	COMBINED IMMUNODEFICIENCY AND MEGALOBlastic ANEMIA WITH OR WITHOUT HYPERHOMOCYSTEINEMIA
MTM1	Severe X-linked myotubular myopathy
MTRR	Disorders of Intracellular Cobalamin Metabolism, Homocystinuria without methylmalonic aciduria, Homocystinuria-Megaloblastic anemia due to defect in cobalamin metabolism, cblE complementation type
MTTP	Abetalipoproteinaemia
MUTYH	Carcinoma of colon, Colon cancer, Familial colorectal cancer, Hereditary cancer-predisposing syndrome, MUTYH-associated polyposis, MYH-associated polyposis, Neoplasm of stomach, Pilomatrixoma
MVK	Hyperimmunoglobulin D with periodic fever, Mevalonic aciduria, Porokeratosis 3, disseminated superficial actinic type
MYBPC3	Asymmetric septal hypertrophy, Cardiomyopathy, Cardiovascular phenotype, Dyspnea, Familial dilated cardiomyopathy, Familial hypertrophic cardiomyopathy 1, Familial hypertrophic cardiomyopathy 4, Heart block, Hypertrophic cardiomyopathy, Inborn genetic diseases, Left ventricular hypertrophy, Left ventricular noncompaction, Left ventricular noncompaction 10, Long QT syndrome, MYBPC3-Related Disorders, Noncompaction cardiomyopathy, Primary

	dilated cardiomyopathy, Primary familial hypertrophic cardiomyopathy, Tachycardia, Ventricular extrasystoles
MYCN	Inborn genetic diseases
MYEF2, SLC24A5	Albinism, oculocutaneous, type VI
MYF5	Abnormality of the ribs, EXTERNAL, External ophthalmoplegia, OPTHALMOPLÉGIA, Scoliosis, WITH RIB AND VERTEBRAL ANOMALIES
MYH11, NDE1	Familial aortopathy
MYH2, MYHAS	Myopathy, and ophthalmoplegia, proximal
MYH3	Contractures, Spondylarcarpotarsal synostosis syndrome, and variable skeletal fusions syndrome 1A, pterygia
MYH6	Familial hypertrophic cardiomyopathy 1
MYH7	Hypertrophic cardiomyopathy, Primary dilated cardiomyopathy
MYH7, MHRT	Cardiomyopathy, Cardiovascular phenotype, Hypertrophic cardiomyopathy, MYH7-Related Disorders
MYL2, LOC114827850	Familial hypertrophic cardiomyopathy 10
MYLK	Visceral myopathy
MYO15A	Congenital sensorineural hearing impairment, Deafness, Nonsyndromic hearing loss and deafness, Rare genetic deafness, autosomal recessive 3
MYO3A	Deafness, autosomal recessive 30
MYO5B	Congenital microvillous atrophy
MYO6	Deafness, Nonsyndromic hearing loss and deafness, Rare genetic deafness, autosomal dominant 22
MYO7A	Deafness, MYO7A-Related Disorders, Rare genetic deafness, Retinal dystrophy, Retinitis pigmentosa, Usher syndrome, Usher syndrome type 1, autosomal dominant 11, autosomal recessive 2, type 1B
MYOCD	CONGENITAL, MEGABLADDER, Prune belly syndrome
MYRF	CARDIAC-UROGENITAL SYNDROME
NADSYN1	AND LIMB DEFECTS SYNDROME 3, CARDIAC, Congenital NAD deficiency disorder, RENAL, VERTEBRAL
NAGLU	Charcot-Marie-Tooth disease, MPS-III-B, Mucopolysaccharidosis, Sanfilippo syndrome, axonal type 2V
NALCN	Hypotonia, infantile, with psychomotor retardation and characteristic facies 1
NBAS	Fever-associated acute infantile liver failure syndrome, Infantile liver failure syndrome 2
NBN	Acute lymphoid leukemia, Aplastic anemia, Breast-ovarian cancer, Familial cancer of breast, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome, Lissencephaly, Microcephaly, Ovarian Neoplasms, familial 1, normal intelligence and immunodeficiency

NCF1, LOC106029312	Chronic granulomatous disease, Chronic granulomatous disease due to deficiency of NCF-1, Granulomatous disease, autosomal recessive, autosomal recessive cytochrome b-positive, chronic, cytochrome b-positive, type 1, type III
NCR1, NLRP7	1, Hydatidiform mole, recurrent
NCSTN	Familial acne inversa 1
NDE1	Lissencephaly 4
NDNF	HYPOGONADOTROPIC HYPOGONADISM 25 WITH ANOSMIA
NDUFA12	Leigh syndrome
NDUFAF2	Inborn genetic diseases, Leigh syndrome, MITOCHONDRIAL COMPLEX I DEFICIENCY, Mitochondrial complex I deficiency, NDUFAF2-Related Disorders, NUCLEAR TYPE 10, nuclear type 1
NDUFAF3	Mitochondrial complex I deficiency
NDUFB11	Linear skin defects with multiple congenital anomalies 3
NDUFS4	Leigh syndrome, Mitochondrial complex I deficiency, nuclear type 1
NDUFS6	MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 9
NDUFV1	MITOCHONDRIAL COMPLEX I DEFICIENCY, Mitochondrial complex I deficiency, NUCLEAR TYPE 4, nuclear type 1
NEB	Inborn genetic diseases, Nemaline myopathy, Nemaline myopathy 2, Non-immune hydrops fetalis
NEB, RIF1	Nemaline myopathy, Nemaline myopathy 2
NEBL	Hypertrophic cardiomyopathy, Long QT syndrome, Primary dilated cardiomyopathy, Primary familial hypertrophic cardiomyopathy, Sudden unexplained death
NEFL	Charcot-Marie-Tooth disease type 2E
NEK1	24, AMYOTROPHIC LATERAL SCLEROSIS, Majewski type, SUSCEPTIBILITY TO, Short rib-polydactyly syndrome, Short-rib thoracic dysplasia 3 with or without polydactyly
NEUROD1	Maturity-onset diabetes of the young type 6
NEXN	Dilated cardiomyopathy 1CC, Familial hypertrophic cardiomyopathy 20
NF1	Axillary freckling, Café-au-lait macules with pulmonary stenosis, Focal T2 hyperintense basal ganglia lesion, Ganglioglioma, Hereditary cancer-predisposing syndrome, Inborn genetic diseases, Juvenile myelomonocytic leukemia, Multiple café-au-lait spots, Neurofibroma, Neurofibromas, Neurofibromatosis, Neurofibromatosis-Noonan syndrome, Optic nerve glioma, Pilocytic astrocytoma, Tibial pseudoarthrosis, familial spinal, type 1
NF1, LOC111811965	Hereditary cancer-predisposing syndrome, Neurofibromatosis, type 1
NF2	Meningioma, Neurofibromatosis, type 2
NFIB	ACQUIRED, Intellectual disability, MACROCEPHALY, Macrocephalus, WITH IMPAIRED INTELLECTUAL DEVELOPMENT
NFIX	Marshall-Smith syndrome
NGLY1	Congenital disorder of deglycosylation, Intellectual disability, Neuromotor delay, Peripheral neuropathy
NHLRC1	Epilepsy, Lafora disease, progressive myoclonic 2b
NHLRC2	AND CEREBRAL ANGIOMATOSIS, FIBROSIS, NEURODEGENERATION
NHS	Nance-Horan syndrome

NIPAL4	Autosomal recessive congenital ichthyosis 6
NIPBL	Cornelia de Lange syndrome 1
NKX2-5	Abnormality of cardiovascular system morphology, Atrial septal defect 7 with or without atrioventricular conduction defects
NKX3-2	Spondylo-megaepiphyseal-metaphyseal dysplasia
NKX6-2	AUTOSOMAL RECESSIVE, SPASTIC ATAXIA 8, WITH HYPOMYELINATING LEUKODYSTROPHY
NLGN4X	Autism, Non-syndromic X-linked intellectual disability, X-linked 2, susceptibility to
NLRP7	1, Hydatidiform mole, recurrent
NOTCH1	Adams-Oliver syndrome 5, Aortic valve disorder, congenital heart defect
NPC1	Niemann-Pick disease, Niemann-Pick disease type C1, type C
NPHP1	Nephronophthisis, Nephronophthisis 1
NPHP3, NPHP3-ACAD11	Meckel syndrome type 7
NPHS1	Finnish congenital nephrotic syndrome
NPHS2	Idiopathic nephrotic syndrome, Nephrotic syndrome, idiopathic, steroid-resistant
NPHS2, AXDND1	Idiopathic nephrotic syndrome, Nephrotic range proteinuria, Nephrotic syndrome, idiopathic, steroid-resistant
NPRL3, HBA-LCR	Epilepsy, familial focal, with variable foci 3
NR0B1	Congenital adrenal hypoplasia, X-linked
NR2E3	Abnormality of color vision, Cone-rod dystrophy, Enhanced s-cone syndrome, Horizontal nystagmus, NR2E3-Related Disorders, Retinal dystrophy, Retinitis pigmentosa, Retinitis pigmentosa 37, Visual impairment
NR3C2	Autosomal dominant pseudohypoaldosteronism type 1
NSD1	Beckwith-Wiedemann syndrome, Inborn genetic diseases, Sotos syndrome 1
NSD2	4p partial monosomy syndrome, Wolf-Hirschhorn like syndrome
NSMCE2	Seckel syndrome 10
NSMF	Hypogonadotropic hypogonadism 9 with or without anosmia
NSUN2	Mental retardation, autosomal recessive 5
NT5E	Calcification of joints and arteries
NTHL1	Familial adenomatous polyposis 3, Hereditary cancer-predisposing syndrome
NTRK1	Hereditary insensitivity to pain with anhidrosis
OAT	Ornithine aminotransferase deficiency
OBSL1	Three M syndrome 2
OCA2	1, Skin/hair/eye pigmentation, Tyrosinase-positive oculocutaneous albinism, variation in
OCLN	Pseudo-TORCH syndrome 1
OFD1	Joubert syndrome, Orofaciodigital syndrome I, Simpson-Golabi-Behmel syndrome, type 2
OPA1	Abortive cerebellar ataxia, Dominant hereditary optic atrophy, Inborn genetic diseases, Mitochondrial diseases, Retinal dystrophy
OPHN1	Mental retardation X-linked with cerebellar hypoplasia and distinctive facial appearance

OPN1LW	Cone monochromatism
ORC6	Meier-Gorlin syndrome 3
OSGIN2, NBN	Hereditary cancer-predisposing syndrome, Microcephaly, normal intelligence and immunodeficiency
OTC	Abnormality of ornithine metabolism, Hyperammonemia, Ornithine carbamoyltransferase deficiency, Protein avoidance
OTOA	Deafness, Rare genetic deafness, autosomal recessive 22
OTOF	Deafness, Rare genetic deafness, autosomal recessive 9
OTOG	Deafness, Intellectual disability, Rare genetic deafness, Seizures, autosomal recessive 18b
OTOGL	Rare genetic deafness
OTUD6B	Dysmorphic features, Epilepsy, Intellectual developmental disorder with dysmorphic facies, Intellectual disability, and distal limb anomalies, seizures
OTX2	Syndromic microphthalmia type 5
P2RY12, MED12L	Platelet-type bleeding disorder 8
P3H1	Osteogenesis imperfecta type 8
P3H2	Myopia, high, with cataract and vitreoretinal degeneration
P4HA2	Myopia 25, autosomal dominant
PAFAH1B1	Inborn genetic diseases, Lissencephaly due to LIS1 mutation
PAH	Phenylketonuria
PALB2	Basal cell carcinoma, Breast cancer, Cancer of the pancreas, Familial cancer of breast, Fanconi anemia, Generalized hypopigmentation, Hereditary breast and ovarian cancer syndrome, Hereditary cancer, Hereditary cancer-predisposing syndrome, Neoplasm of the breast, Ovarian Neoplasms, PALB2-Related Disorders, Pancreatic cancer 3, Pre-B-cell acute lymphoblastic leukemia, Tracheoesophageal fistula, Tumor susceptibility linked to germline BAP1 mutations, complementation group N, susceptibility to
PANK2	Pigmentary pallidal degeneration
PAPSS2	Spondyloepimetaphyseal dysplasia, pakistani type
PARN	Dyskeratosis congenita, autosomal recessive 6
PATL2	OOCYTE MATURATION DEFECT 4
PAX2	Focal segmental glomerulosclerosis 7, Renal coloboma syndrome
PAX3	Rare genetic deafness, Waardenburg syndrome, Waardenburg syndrome type 1
PAX6	Aniridia 1, Keratitis, autosomal dominant
PAX9	3, Tooth agenesis, selective
PC	Pyruvate carboxylase deficiency
PCCA	Propionic acidemia
PCCB	Propionic acidemia
PCDH15	DIGENIC, Deafness, Nonsyndromic Deafness, Rare genetic deafness, Retinal dystrophy, TYPE ID/F, USHER SYNDROME, Usher syndrome, Usher syndrome type 1, Usher syndrome type 1D, Usher syndrome type 1F, autosomal recessive 23, type 1G

PCDH19	Absence seizures, Delayed speech and language development, Early infantile epileptic encephalopathy 9, Frontal cortical atrophy, Generalized seizures, Generalized tonic-clonic seizures, Global developmental delay, Hand tremor, Long palpebral fissure, Prominent fingertip pads, Strabismus, Temporal cortical atrophy
PCLO	Pontocerebellar hypoplasia type 3
PCNT	Microcephalic osteodysplastic primordial dwarfism type II
PCSK1, LOC101929710	Proprotein convertase 1/3 deficiency
PCSK9	Familial hypercholesterolemia, Familial hypercholesterolemia 1, Low density lipoprotein cholesterol level quantitative trait locus 1
PCYT1A	Spondylometaphyseal dysplasia-cone-rod dystrophy syndrome
PDE11A	2, Pigmented nodular adrenocortical disease, primary
PDE6B	Retinal dystrophy, Retinitis pigmentosa, Retinitis pigmentosa 40
PDE6C	Achromatopsia 5
PDE8B	Striatal degeneration, autosomal dominant 1
PDHA1	Inborn genetic diseases, Pyruvate dehydrogenase E1-alpha deficiency
PDX1	DIABETES MELLITUS, Maturity-onset diabetes of the young type 4, PERMANENT NEONATAL 1, Pancreatic agenesis 1
PDZD7	AUTOSOMAL RECESSIVE 57, DEAFNESS, Rare genetic deafness, Usher syndrome, type 2A
PEPD	Prolidase deficiency
PEX1	Deafness enamel hypoplasia nail defects, Peroxisome biogenesis disorder 1A (Zellweger), Peroxisome biogenesis disorder 1B, Peroxisome biogenesis disorders, Retinal dystrophy, Zellweger syndrome spectrum
PEX1, GATAD1	Deafness enamel hypoplasia nail defects, Peroxisome biogenesis disorder 1A (Zellweger), Peroxisome biogenesis disorder 1B, Peroxisome biogenesis disorders, Zellweger syndrome spectrum
PEX10	Peroxisome biogenesis disorder, Peroxisome biogenesis disorder 6A, Peroxisome biogenesis disorder 6B, Peroxisome biogenesis disorders, Zellweger syndrome spectrum, complementation group 7
PEX10, PLCH2	Peroxisome biogenesis disorder 6B
PEX12	Infantile Refsum disease, Peroxisome biogenesis disorder 3A, Peroxisome biogenesis disorders, Zellweger syndrome spectrum
PEX2	Peroxisome biogenesis disorder 5B, Peroxisome biogenesis disorder 5a (zellweger), Peroxisome biogenesis disorders, Zellweger syndrome spectrum
PEX26	Peroxisome biogenesis disorder 7A, Peroxisome biogenesis disorder 7B, Peroxisome biogenesis disorders, Zellweger syndrome spectrum
PEX6	Heimler syndrome 2, Peroxisome biogenesis disorder 4B, Peroxisome biogenesis disorder 4a (zellweger), Peroxisome biogenesis disorders, Retinal dystrophy, Zellweger syndrome spectrum
PEX7	PEX7-Related Disorders, Peroxisome biogenesis disorder 9B, Phytanic acid storage disease, Rhizomelic chondrodysplasia punctata type 1
PGAM2, DBNL	Glycogen storage disease type X
PGAP1	Mental retardation, autosomal recessive 42
PGAP3	Hyperphosphatasia with mental retardation syndrome 4
PGM3, DOP1A	Immunodeficiency 23

PHEX	Familial X-linked hypophosphatemic vitamin D refractory rickets
PHEX, PTCHD1-AS	Familial X-linked hypophosphatemic vitamin D refractory rickets
PHF3, EYS	Retinal dystrophy, Retinitis pigmentosa 25
PHF6	Borjeson-Forssman-Lehmann syndrome
PHGDH	Phosphoglycerate dehydrogenase deficiency
PHIP	Developmental delay, and dysmorphic features, intellectual disability, obesity
PHYH	1, Phytanic acid storage disease, Refsum disease, adult
PI4KA	Polymicrogyria, perisylvian, with cerebellar hypoplasia and arthrogryposis
PIGA	Paroxysmal nocturnal hemoglobinuria 1
PIGN	Multiple congenital anomalies-hypotonia-seizures syndrome 1
PIGO	Hyperphosphatasia with mental retardation syndrome 2, Hyperphosphatasia-intellectual disability syndrome
PIGT	Multiple congenital anomalies-hypotonia-seizures syndrome 3, PIGT-related disorder
PIK3R1	SHORT syndrome
PINK1	Parkinson disease 6, autosomal recessive early-onset
PIRC66, MIR4713HG, CYP19A1	Aromatase deficiency
PITX3	Anterior segment mesenchymal dysgenesis, Cataract 11
PJVK	Deafness, Rare genetic deafness, autosomal recessive 59
PKD1	Autosomal recessive polycystic kidney disease, Polycystic kidney disease, adult type
PKD1, LOC105371049	Polycystic kidney disease, adult type
PKHD1	Autosomal recessive polycystic kidney disease, Polycystic kidney dysplasia, Polycystic liver disease
PKP1	Epidermolysis bullosa simplex due to plakophilin deficiency
PKP2	Arrhythmogenic right ventricular cardiomyopathy, Arrhythmogenic right ventricular dysplasia/cardiomyopathy, Arrhythmogenic ventricular cardiomyopathy, Cardiac arrhythmia, Cardiomyopathy, Cardiovascular phenotype, Sudden unexplained death, type 9
PLA2G5	Fleck retina, familial benign
PLA2G6	Infantile neuroaxonal dystrophy, Iron accumulation in brain, Neurodegeneration with brain iron accumulation 2b, PLA2G6-associated neurodegeneration
PLCB1	Early infantile epileptic encephalopathy 12
PLCB4	Auriculocondylar syndrome 2
PLCD1	Leukonychia totalis
PLD1	Cardiac valvular defect, developmental
PLD3, PRX	Charcot-Marie-Tooth disease, SPINOCEREBELLAR ATAXIA 46
PLEC	Epidermolysis bullosa simplex with muscular dystrophy
PLN, CEP85L	Cardiac arrest, Cardiomyopathy, Cardiovascular phenotype, Dilated cardiomyopathy 1P, Familial hypertrophic cardiomyopathy 18, Hypertrophic cardiomyopathy, Primary dilated cardiomyopathy, Sudden cardiac death
PLOD1	Cardiovascular phenotype, Ehlers-Danlos syndrome, hydroxylysine-deficient

PLOD2	Bruck syndrome 2
PLP1, RAB9B	Hereditary spastic paraplegia 2
PLS3	Bone mineral density quantitative trait locus 18
PMFBP1	SPERMATOGENIC FAILURE 31
PMM2	Congenital disorder of glycosylation, type Ia
PMP22	Charcot-Marie-Tooth disease
PMS2	Acute lymphoid leukemia, Burkitt lymphoma, Colorectal cancer, Glioblastoma, Hereditary cancer, Hereditary cancer-predisposing syndrome, Hereditary nonpolyposis colon cancer, Hereditary nonpolyposis colorectal cancer type 4, Lymphoma, Lynch syndrome, Lynch syndrome I, Pulmonary arterial hypertension, Pulmonary insufficiency, Respiratory insufficiency, Tumor susceptibility linked to germline BAP1 mutations, Turcot syndrome, non-polyposis
PNKD, CATIP-AS2	Paroxysmal nonkinesigenic dyskinesia 1
PNKP	Ataxia-oculomotor apraxia 4, Early infantile epileptic encephalopathy 10, Early infantile epileptic encephalopathy 12, History of neurodevelopmental disorder
PNPLA2	Neutral lipid storage myopathy
PNPLA6	Hereditary spastic paraplegia 39, Laurence-Moon syndrome, PNPLA6-related disorders, Trichomegaly-retina pigmentary degeneration-dwarfism syndrome
PNPLA8	Mitochondrial myopathy-lactic acidosis-deafness syndrome
PNPO	Pyridoxal phosphate-responsive seizures
POC5	Retinitis pigmentosa, Syndromic retinitis pigmentosa
POGLUT1	Dowling-degos disease 4
POGZ	Global developmental delay, Speech apraxia, White-sutton syndrome, dysmorphism, intellectual deficiency
POLA1	VAN ESCH-ODRISCOLL SYNDROME, Van Esch type, X-linked intellectual disability
POLD1	Colorectal cancer 10, Hereditary cancer-predisposing syndrome, Mandibular hypoplasia, and lipodystrophy syndrome, deafness, progeroid features
POLE	12, ADRENAL HYPOPLASIA CONGENITA, AND IMMUNODEFICIENCY, Colorectal cancer, GENITAL ANOMALIES, Hereditary cancer-predisposing syndrome, INTRAUTERINE GROWTH RETARDATION, METAPHYSEAL DYSPLASIA, susceptibility to
POLG	Generalized epilepsy, Global developmental delay, Obesity, Progressive sclerosing poliodystrophy, Seizures
POLH	Xeroderma pigmentosum variant type
POLR1A	Acrofacial dysostosis, Cincinnati type
POLR1C	Treacher Collins syndrome 3
POLR1D	Treacher Collins syndrome 2
POLR2F, SOX10	Rare genetic deafness, Waardenburg syndrome type 4C
POLR3A	Hypomyelinating leukodystrophy 7, Neonatal pseudo-hydrocephalic progeroid syndrome
POLR3B	Cerebellar hypoplasia with endosteal sclerosis, Hypogonadotropic hypogonadism 7 with or without anosmia, Hypomyelinating leukodystrophy 7, Hypomyelinating leukodystrophy 8, with or without oligodontia and/or hypogonadotropic hypogonadism

POMK	12, Muscular dystrophy-dystroglycanopathy (limb-girdle), type c
POMT1	1, Congenital muscular dystrophy-dystroglycanopathy with mental retardation, Limb-girdle muscular dystrophy-dystroglycanopathy, Muscular dystrophy-dystroglycanopathy (congenital with brain and eye anomalies), POMT1-Related Disorders, Walker-Warburg congenital muscular dystrophy, type A, type B1, type C1
POMT2	Congenital muscular dystrophy-dystroglycanopathy with brain and eye anomalies, Limb-girdle muscular dystrophy-dystroglycanopathy, type A2, type C2
POP1	Anauxetic dysplasia 2
POR	Antley-Bixler syndrome with genital anomalies and disordered steroidogenesis, Disordered steroidogenesis due to cytochrome p450 oxidoreductase deficiency
PORCN	Focal dermal hypoplasia
POT1	10, Hereditary cancer-predisposing syndrome, Melanoma, cutaneous malignant, susceptibility to
POU3F4	Deafness, Rare genetic deafness, X-linked 2
POU4F3	Rare genetic deafness
PPARG	Diabetes Mellitus, Diabetes mellitus, Noninsulin-Dependent, digenic, type II, with Acanthosis Nigricans and Hypertension
PPIB	Osteogenesis imperfecta type 9
PPOX	Variegate porphyria
PPP1R12A	GENITOURINARY AND/OR BRAIN MALFORMATION SYNDROME
PPT1	History of neurodevelopmental disorder, Neuronal Ceroid-Lipofuscinosis, Neuronal ceroid lipofuscinosis, Neuronal ceroid lipofuscinosis 1, Recessive
PQBP1	Delayed speech and language development, Hyperactivity, Inborn genetic diseases, Intellectual disability, Microcephaly, Renpenning syndrome 1
PRB3	PRB3M(NULL)
PRDM16	Left ventricular noncompaction 8
PRDM5	Brittle cornea syndrome 2
PRDX1, MMACHC	DIGENIC, METHYLMALONIC ACIDURIA AND HOMOCYSTINURIA, cb1C TYPE
PRF1	Familial hemophagocytic lymphohistiocytosis, Familial hemophagocytic lymphohistiocytosis 2
PRKAR1A	Carney complex, type 1
PRKAR1A, FAM20A	Amelogenesis imperfecta type 1G
PRKAR1B, DNAAF5	18, Ciliary dyskinesia, primary
PRKCSH	Polycystic liver disease 1
PRKN	Parkinson disease 2
PRMT7	Short stature, and seizures, brachydactyly, intellectual developmental disability
PROK2	Hypogonadotropic hypogonadism 4 with or without anosmia
PROKR2	Inborn genetic diseases, Kallmann syndrome 3
PROM1	Cone-rod dystrophy 12, PROM1-Related Disorders, Retinal dystrophy, Retinitis pigmentosa, Retinitis pigmentosa 41
PROP1	Pituitary hormone deficiency, combined, combined 2

PRPH2	Macular dystrophy, Retinal dystrophy, Retinitis pigmentosa 7, Retinitis punctata albescens, adult-onset, autosomal dominant, vitelliform
PRRT2	2, Episodic kinesigenic dyskinesia 1, History of neurodevelopmental disorder, Infantile convulsions and choreoathetosis, Paroxysmal kinesigenic dyskinesia, Paroxysmal nonkinesigenic dyskinesia 1, Seizures, benign familial infantile
PRSS12	Mental retardation, autosomal recessive 1
PRSS56	Microphthalmia, isolated 6
PRX	Charcot-Marie-Tooth disease, demyelinating, type 4F
PSAP	Combined saposin deficiency
PSEN1	3, Acne inversa, familial
PSENEN	2, Acne inversa, familial
PTCH1	Gorlin syndrome, Hereditary cancer-predisposing syndrome
PTCH2	Gorlin syndrome, Medulloblastoma
PTEN	Cowden syndrome, Cowden syndrome 1, Glioblastoma, Glioma susceptibility 2, Hemangioma, Hereditary cancer-predisposing syndrome, Inborn genetic diseases, Macrocephaly/autism syndrome, Malignant tumor of prostate, Meningioma, Neoplasm of brain, Neoplasm of the breast, Neoplasm of the large intestine, Non-small cell lung cancer, Ovarian Neoplasms, PTEN hamartoma tumor syndrome, PTEN-related disorder, Proteus-like syndrome, VACTERL association with hydrocephalus, familial
PTH1R	Chondrodysplasia Blomstrand type
PTPN11	Metachondromatosis
PTPRF	2, Breasts and/or nipples, aplasia or hypoplasia of
PTPRO	Nephrotic syndrome, type 6
PTS	BH4-deficient hyperphenylalaninemia A, Hyperphenylalaninemia, a, bh4-deficient, due to partial pts deficiency
PUF60	Verheij syndrome
PURA	Apnea, Generalized hypotonia, Intellectual disability, Limb dystonia, Mental retardation, PURA Syndrome, PURA-related severe neonatal hypotonia-seizures-encephalopathy syndrome due to a point mutation, autosomal dominant 31
PUS7	AND SHORT STATURE, INTELLECTUAL DEVELOPMENTAL DISORDER WITH ABNORMAL BEHAVIOR, MICROCEPHALY
PXDN	Anterior segment dysgenesis 7
PYCR1	Autosomal recessive cutis laxa type 2B
PYGL	Glycogen storage disease, type VI
PYGM	Glycogen storage disease, type V
RAB23	Carpenter syndrome, Carpenter syndrome 1
RAB27A	Griscelli syndrome, Griscelli syndrome type 2
RAB33B	Smith-McCort dysplasia 2
RAB3GAP1	Warburg micro syndrome 1
RABL3	5, PANCREATIC CANCER, SUSCEPTIBILITY TO
RAD50	Hereditary cancer-predisposing syndrome, Nijmegen breakage syndrome-like disorder

RAD51C	Breast-ovarian cancer, Fanconi anemia, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome, Ovarian Neoplasms, RAD51C-Related Disorders, complementation group O, familial 3
RAD51D, RAD51L3-RFFL	Breast-ovarian cancer, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome, Ovarian Neoplasms, familial 4
RAD51L3-RFFL, RAD51D	Breast-ovarian cancer, Hereditary cancer-predisposing syndrome, familial 4
RAI1	Smith-Magenis syndrome
RAPSN	11, Myasthenic syndrome, associated with acetylcholine receptor deficiency, congenital
RARS1	9, Leukodystrophy, hypomyelinating
RASA1	Capillary malformation-arteriovenous malformation, Capillary malformation-arteriovenous malformation 1
RB1	Hereditary cancer-predisposing syndrome, Neoplasm, Osteosarcoma, Retinoblastoma, Small cell lung cancer, Urinary bladder cancer, trilateral
RBBP8	Microcephaly with mental retardation and digital anomalies
RBM20	Cardiovascular phenotype, Dilated cardiomyopathy 1DD, Primary dilated cardiomyopathy
RBP3	Retinitis pigmentosa 66
RD3	Leber congenital amaurosis 12
RDH12	Retinitis pigmentosa 53
RDH5, BLOC1S1-RDH5	Fundus albipunctatus, Pigmentary retinal dystrophy, autosomal recessive
RECQL	Hereditary cancer-predisposing syndrome
RECQL, PYROXD1	Hereditary cancer-predisposing syndrome
RECQL4	B lymphoblastic leukemia lymphoma with t(12;21)(p13;q22); TEL-AML1 (ETV6-RUNX1), Baller-Gerold syndrome, High Grade Surface Osteosarcoma, Rapadilino syndrome, Rothmund-Thomson syndrome, Rothmund-Thomson syndrome type 2
REEP6	Retinitis pigmentosa 77
RELT	AMELOGENESIS IMPERFECTA, TYPE IIIC
REN	Hyperproreninemia, familial
RET	Hirschsprung disease 1, Sensorineural hearing loss
RFX5	Bare lymphocyte syndrome, complementation group c, type II
RFXANK	Bare lymphocyte syndrome, complementation group B, type II
RFXAP	Bare Lymphocyte Syndrome, Bare lymphocyte syndrome 2, Complementation Group D, Type II
RHAG	Rh-null, regulator type
RHCE	AMORPH TYPE, RH-NUL
RHO	Autosomal dominant retinitis pigmentosa
RIF1, NEB	Nemaline myopathy, Nemaline myopathy 2
RIN2	Macrocephaly, alopecia, and scoliosis, cutis laxa
RIPK1	IMMUNODEFICIENCY 57 WITH AUTOINFLAMMATION
RIPK4	Bartsocas-Papas syndrome
RNASEH2A	Aicardi Goutieres syndrome 4

RNASEH2B	Aicardi Goutieres syndrome 2
RNF113A	Trichothiodystrophy 5, nonphotosensitive
RNF216	Gordon Holmes syndrome
ROBO3	Gaze palsy, familial horizontal, with progressive scoliosis 1
RORA, RORA-AS1	INTELLECTUAL DEVELOPMENTAL DISORDER WITH OR WITHOUT EPILEPSY OR CEREBELLAR ATAXIA
RP1	Retinal dystrophy, Retinitis pigmentosa, Retinitis pigmentosa 1
RP1L1	RETINITIS PIGMENTOSA 88
RPE65	Leber congenital amaurosis 2, RETINITIS PIGMENTOSA 87 WITH CHOROIDAL INVOLVEMENT, RPE65-Related Disorders, Retinal dystrophy, Retinitis pigmentosa 20
RPGR	Inborn genetic diseases, Retinal dystrophy, Retinitis pigmentosa, Retinitis pigmentosa 15, X-linked, and sinorespiratory infections, with deafness
RPGRIP1	Leber congenital amaurosis 6
RPGRIP1L	Joubert syndrome, Joubert syndrome 7
RPL36A-HNRNP2, GLA	Fabry disease
RPL5, DIPK1A	Diamond-Blackfan anemia, Diamond-Blackfan anemia 1
RPS10, RPS10-NUDT3	Diamond-Blackfan anemia 9
RPS27	Diamond-Blackfan anemia 17
RPS6KA3	Coffin-Lowry syndrome, Mental retardation, X-linked 19
RSPH1	Kartagener syndrome, Primary ciliary dyskinesia, Primary ciliary dyskinesia 24
RSPH4A	11, Ciliary dyskinesia, Kartagener syndrome, Primary ciliary dyskinesia, primary
RSPO2	TETRAAMELIA SYNDROME 2
RTKL1, RTKL1-TNFRSF6B	3, 4, 5, Dyskeratosis congenita, Idiopathic fibrosing alveolitis, Pulmonary fibrosis and/or bone marrow failure, autosomal dominant, autosomal recessive, chronic form, telomere-related
RTN2	Hereditary spastic paraplegia 12
RTTN	Congenital microcephaly, Microcephaly, and polymicrogyria with or without seizures, short stature
RUNX1	Acute myeloid leukemia, Familial platelet disorder with associated myeloid malignancy
RYSR1	1, Central core myopathy, Malignant hyperthermia, Minicore myopathy, Multi-minicore disease and atypical periodic paralysis, Neuromuscular disease, RYSR1-Related Disorders, susceptibility to
SACS	Autosomal recessive spastic ataxia, Charlevoix-Saguenay spastic ataxia, Spastic paraplegia
SAG	Oguchi disease, Retinitis pigmentosa 47, SAG-Related Disorders
SALL1	Townes syndrome
SAMD9L	Ataxia-pancytopenia syndrome
SAMHD1	Aicardi Goutieres syndrome 5
SASH1	Dyschromatosis universalis hereditaria 1
SATB2	SATB2-Related Disorder
SBDS	Inborn genetic diseases, Shwachman-Diamond syndrome 1
SBF1	Charcot-Marie-Tooth disease type 4

SCAPER	Attention deficit hyperactivity disorder, INTELLECTUAL DEVELOPMENTAL DISORDER AND RETINITIS PIGMENTOSA, Intellectual disability, Rod-cone dystrophy, moderate
SCARB2	Epilepsy, progressive myoclonic 4, with or without renal failure
SCARF2	Van den Ende-Gupta syndrome
SCN1A	Autosomal dominant epilepsy, Early infantile epileptic encephalopathy, Familial hemiplegic migraine type 3, Generalized epilepsy with febrile seizures plus, History of neurodevelopmental disorder, Severe myoclonic epilepsy in infancy, type 2, Dravet
SCN1A, LOC102724058	Autosomal dominant epilepsy, Early infantile epileptic encephalopathy, Epileptic encephalopathy, Generalized epilepsy with febrile seizures plus, Seizures, Severe myoclonic epilepsy in infancy, type 2
SCN2A	SCN2A-related disorder
SCN5A	Brugada syndrome, Brugada syndrome (shorter-than-normal QT interval), Brugada syndrome 1, Cardiovascular phenotype, Dilated cardiomyopathy 1E, Heart block, Long QT syndrome 1, nonprogressive
SCN5A, LOC110121269	Brugada syndrome, Brugada syndrome (shorter-than-normal QT interval)
SCN9A, SCN1A-AS1	Generalized epilepsy with febrile seizures plus, Hereditary sensory and autonomic neuropathy type IIA, Indifference to pain, autosomal recessive, congenital, type 7
SCNN1A	Autosomal recessive pseudohypoaldosteronism type 1, Idiopathic bronchiectasis
SCNN1B	Liddle syndrome 1
SCNN1G	Autosomal recessive pseudohypoaldosteronism type 1, LIDDLE SYNDROME 2
SCO1	Mitochondrial complex IV deficiency
SCP2	Leukoencephalopathy with dystonia and motor neuropathy
SDCCAG8	Bardet-Biedl syndrome, Bardet-Biedl syndrome 16, Senior-Loken syndrome 7
SDHA	Carney triad, Dilated cardiomyopathy 1GG, Hereditary cancer-predisposing syndrome, Leigh syndrome, Mitochondrial complex II deficiency, Paragangliomas 5, Pilocytic astrocytoma
SDHAF2	Hereditary Paraganglioma-Pheochromocytoma Syndromes
SDHB	Carney-Stratakis syndrome, Gastrointestinal stromal tumor, Hereditary Paraganglioma-Pheochromocytoma Syndromes, Hereditary cancer-predisposing syndrome, Paragangliomas 4, Pheochromocytoma
SDHC	Gastrointestinal stromal tumor, Hereditary Paraganglioma-Pheochromocytoma Syndromes, Hereditary cancer-predisposing syndrome, Paragangliomas 3
SDHD	Carney-Stratakis syndrome, Cowden syndrome 3, Hereditary Paraganglioma-Pheochromocytoma Syndromes, Hereditary cancer-predisposing syndrome, Paragangliomas 1, Paragangliomas 1 with sensorineural hearing loss, Pheochromocytoma
SDR9C7	AUTOSOMAL RECESSIVE 13, CONGENITAL, ICHTHYOSIS
SEC23B	Congenital dyserythropoietic anemia
SEC24D	Cole-Carpenter syndrome 2
SECISBP2	Thyroid hormone metabolism, abnormal
SELENBP1	EXTRAORAL HALITOSIS DUE TO METHANETHIOL OXIDASE DEFICIENCY, Extra oral halitosis
SELENON	Eichsfeld type congenital muscular dystrophy
SEMA3A	Hypogonadotropic hypogonadism 16 with or without anosmia

SEPSECS	Pontocerebellar hypoplasia type 2D
SEPTIN12	Spermatogenic failure 10
SERAC1	3-methylglutaconic aciduria with deafness, Mitochondrial oxidative phosphorylation disorder, and Leigh-like syndrome, encephalopathy
SERPINA6	Corticosteroid-binding globulin deficiency
SERPINA7	Thyroxine-binding globulin quantitative trait locus
SERPINB6	Rare genetic deafness
SERPINB7	Palmoplantar keratoderma, nagashima type
SERPINC1	Antithrombin III deficiency
SERPINF1	Osteogenesis imperfecta, type VI
SERPING1	Hereditary angioedema type 1
SERPINH1	Osteogenesis imperfecta type 10
SETBP1	SETBP1-Related Disorder
SETD5	Inborn genetic diseases, Mental retardation, autosomal dominant 23
SF3B4	Hereditary hearing loss and deafness, Inborn genetic diseases, Nager syndrome
SFRP4	Pyle metaphyseal dysplasia
SFTPA1	Respiratory distress associated with prematurity
SFTPB	1, Surfactant metabolism dysfunction, pulmonary
SGCA	Autosomal recessive limb-girdle muscular dystrophy type 2D
SGCD	Neuromuscular disease
SGCE, CASD1	Myoclonic dystonia
SGCG	Severe autosomal recessive muscular dystrophy of childhood - North African type
SGSH	Developmental regression, Diarrhea, Gastrointestinal dysmotility, Global developmental delay, MPS-III-A, Mucopolysaccharidosis, Nystagmus, Retinal dystrophy, Sanfilippo syndrome, Severe visual impairment
SH2D1A	Lymphoproliferative syndrome 1, X-Linked Lymphoproliferative Syndrome, X-linked
SH3PXD2B	Frank-Ter Haar syndrome
SH3TC2	Charcot-Marie-Tooth disease, Charcot-Marie-Tooth disease type 4, Inborn genetic diseases, Mononeuropathy of the median nerve, SH3TC2-Related Disorders, mild, type 4C
SHANK3	22q13.3 deletion syndrome, Autism spectrum disorder, History of neurodevelopmental disorder, Inborn genetic diseases, SHANK3-Related Disorder
SHOX	Leri-Weill dyschondrosteosis
SI	Sucrase-isomaltase deficiency
SIX6	Colobomatous optic disc-macular atrophy-chorioretinopathy syndrome
SKIV2L	Trichohepatoenteric syndrome 2
SLC10A7	AMELOGENESIS IMPERFECTA, AND SKELETAL DYSPLASIA WITH SCOLIOSIS, SHORT STATURE
SLC12A1	Bartter syndrome, antenatal, type 1
SLC12A3	Familial hypokalemia-hypomagnesemia
SLC12A6	Agenesis of the corpus callosum with peripheral neuropathy, Charcot-Marie-Tooth disease
SLC17A5	Salla disease, Sialic acid storage disease, severe infantile type

SLC19A1, COL18A1	Knobloch syndrome 1
SLC19A2	Megaloblastic anemia, thiamine-responsive, with diabetes mellitus and sensorineural deafness
SLC19A3	Biotin-responsive basal ganglia disease
SLC22A5	Renal carnitine transport defect
SLC25A20	Carnitine acylcarnitine translocase deficiency
SLC26A2	3MC syndrome 2, Achondrogenesis, Atelosteogenesis type II, Diastrophic dysplasia, Multiple epiphyseal dysplasia type 4, Osteochondrodysplasia, SLC26A2-Related Disorders, type IB
SLC26A3	Congenital secretory diarrhea, chloride type
SLC26A4	Enlarged vestibular aqueduct, Pendred syndrome, Rare genetic deafness
SLC2A10	Arterial tortuosity syndrome, Cardiovascular phenotype
SLC2A2	Fanconi-Bickel syndrome
SLC30A8	Diabetes mellitus type 2
SLC33A1	Spastic paraplegia, Spastic paraplegia 42, autosomal dominant
SLC34A3	Autosomal recessive hypophosphatemic bone disease
SLC35A2	SLC35A2-CDG
SLC35D1	Schneckenbecken dysplasia
SLC37A4	Glucose-6-phosphate transport defect, Glycogen storage disease, Inborn genetic diseases, Phosphate transport defect
SLC38A8	FOVEAL HYPOPLASIA 2 WITH OPTIC NERVE MISROUTING AND ANTERIOR SEGMENT DYSGENESIS
SLC39A4	Hereditary acrodermatitis enteropathica
SLC45A2	Oculocutaneous albinism type 4
SLC4A1	Autosomal dominant distal renal tubular acidosis
SLC4A11	4, Corneal dystrophy, Corneal endothelial dystrophy, Fuchs endothelial
SLC52A3	Brown-Vialetto-Van Laere syndrome 1
SLC6A1	Myoclonic-atonic epilepsy, SLC6A1-Related Disorder
SLC9A3	Diarrhea 8, congenital, secretory sodium
SLC9A3, SLC9A3-AS1	Diarrhea 8, congenital, secretory sodium
SLC9A6	Gastrostomy tube feeding in infancy, Global developmental delay, Recurrent respiratory infections, Scoliosis, Seizures, Sleep disturbance
SLCO2A1	Primary hypertrophic osteoarthropathy, autosomal recessive 2
SLITRK1	Tourette Syndrome, Trichotillomania
SLURP1	Acroerythrokeratoderma
SMAD3	Familial thoracic aortic aneurysm and aortic dissection
SMAD4	Carcinoma of pancreas, Hereditary cancer-predisposing syndrome, Juvenile polyposis syndrome, Juvenile polyposis/hereditary hemorrhagic telangiectasia syndrome, Myhre syndrome
SMAD6	Aortic valve disease 2, Aortic valve disorder, CRANIOSYNOSTOSIS 7, SUSCEPTIBILITY TO
SMARCA4	Neuroblastoma
SMARCAL1	Schimke immuno-osseous dysplasia

SMARCB1	Teratoid tumor, atypical
SMARCE1	Meningioma, familial
SMC1A	85, Congenital muscular hypertrophy-cerebral syndrome, EARLY INFANTILE, EPILEPTIC ENCEPHALOPATHY, WITH OR WITHOUT MIDLINE BRAIN DEFECTS
SMN1	Werdnig-Hoffmann disease
SMPD1	Niemann-Pick disease, Sphingomyelin/cholesterol lipidosis, type A, type B
SNAP29	2, CEDNIK syndrome, Leukodystrophy, hypomyelinating
SNRNPB	Cerebro-costo-mandibular syndrome
SOHLH1	Nonsyndromic hypergonadotropic hypogonadism, OVARIAN DYSGENESIS 5
SON	Inborn genetic diseases, ZTTK syndrome
SOS1	Gingival fibromatosis 1
SOX2, SOX2-OT	Anophthalmia/microphthalmia-esophageal atresia syndrome
SOX9	Campomelic dysplasia with autosomal sex reversal, Camptomelic dysplasia
SOX9, LOC108021846	Campomelic dysplasia with autosomal sex reversal
SP110, SP140	Hepatic veno-occlusive disease-immunodeficiency syndrome
SP7	Osteogenesis imperfecta type 12
SPART	Troyer syndrome
SPAST	Spastic paraplegia 4, autosomal dominant
SPEF2	Primary ciliary dyskinesia, SPERMATOGENIC FAILURE 43
SPEG	5, Myopathy, centronuclear
SPEG, ASIC4- AS1	5, Myopathy, centronuclear
SPG11	Amyotrophic lateral sclerosis type 5, Hereditary spastic paraplegia, Spastic paraplegia 11, autosomal recessive
SPG7	Hereditary spastic paraplegia, Hereditary spastic paraplegia 7, Mitochondrial diseases
SPINK2	Spermatogenic failure 29
SPINK5	Netherton syndrome
SPNS2	AUTOSOMAL RECESSIVE 115, DEAFNESS, Inborn genetic diseases
SPRTN	Ruijs-Aalfs syndrome
SPTA1	Elliptocytosis 2, Hereditary pyropoikilocytosis
SPTB	Hereditary spherocytosis, Spherocytosis type 2
SQSTM1	Amyotrophic lateral sclerosis and/or frontotemporal dementia 1, Paget disease of bone 2, SQSTM1-related disorder, early-onset
SRCAP	Floating-Harbor syndrome
SRPK2, KMT2E	See cases
SRY	46, XY sex reversal, type 1
ST14	Ichthyosis, autosomal recessive 11, congenital
STAG1	AUTOSOMAL DOMINANT 47, MENTAL RETARDATION
STAG3	Abnormality of the ovary, Female infertility, Premature ovarian failure 8, Premature ovarian insufficiency
STAT1	Mycobacterial and viral infections, autosomal recessive, susceptibility to

STIM1	1, Combined immunodeficiency due to STIM1 deficiency, Myopathy, Stormorken syndrome, tubular aggregate
STK11	Hereditary cancer-predisposing syndrome, Peutz-Jeghers syndrome
STRA6	Microphthalmia syndromic 9
STRC	Deafness, Rare genetic deafness, autosomal recessive 16
STXBP1	Early infantile epileptic encephalopathy, Early infantile epileptic encephalopathy 4, Epileptic encephalopathy
STXBP2	5, Hemophagocytic lymphohistiocytosis, familial
SUCLG1	Mitochondrial DNA depletion syndrome 9 (encephalomyopathic with methylmalonic aciduria)
SUFU	Gorlin syndrome, Medulloblastoma, Medulloblastoma with extensive nodularity, desmoplastic
SULT2B1	AUTOSOMAL RECESSIVE 14, Autosomal recessive congenital ichthyosis 2, CONGENITAL, ICHTHYOSIS
SUMF1	Multiple sulfatase deficiency
SUN5	Spermatogenic failure 16
SURF1	Abnormal pyramidal signs, Cerebellar ataxia, Charcot-Marie-Tooth disease, Dysarthria, Inborn genetic diseases, Leigh syndrome, Leigh syndrome due to COX IV deficiency, Leigh syndrome due to mitochondrial complex IV deficiency, Mitochondrial complex IV deficiency, Muscle weakness, type 4k
SUZ12	IMAGAWA-MATSUMOTO SYNDROME
SYCP2	Cryptozoospermia, Early spermatogenesis maturation arrest, Oligosynaptic infertility
SYCP3	Spermatogenic failure 4
SYNE1	ARTHROGRYPOSIS MULTIPLEX CONGENITA, Cerebellar ataxia, Emery-Dreifuss muscular dystrophy 4, MYOGENIC TYPE, Spinocerebellar ataxia, autosomal dominant, autosomal recessive 8
SYNE4	Rare genetic deafness
SYNGAP1	Inborn genetic diseases, Mental retardation, autosomal dominant 5
SZT2	Early infantile epileptic encephalopathy 18
TAC3	Hypogonadotropic hypogonadism 10 with or without anosmia
TACO1	Mitochondrial complex IV deficiency
TALDO1	Deficiency of transaldolase
TANGO2	AND NEURODEGENERATION, Acute rhabdomyolysis, CARDIAC ARRHYTHMIAS, Cardiac arrhythmia, Episodic flaccid weakness, Intellectual functioning disability, METABOLIC CRISES, RECURRENT, Seizures, WITH RHABDOMYOLYSIS
TAP1	Bare lymphocyte syndrome type 1
TAP2	Bare lymphocyte syndrome type 1, PEPTIDE TRANSPORTER PSF2 POLYMORPHISM
TAZ	3-Methylglutaconic aciduria type 2
TBC1D20	Warburg micro syndrome 4
TBC1D24	1, Caused by mutation in the TBC1 domain family, DOORS syndrome, Deafness, Epileptic encephalopathy, Inborn genetic diseases, autosomal dominant 65, early infantile, member 24
TBCK	Hypotonia, Inborn genetic diseases, Syndromic Infantile Encephalopathy, infantile, with psychomotor retardation and characteristic facies 3

TBR1	Autism 5, Autistic behavior, Intellectual disability, Moderate global developmental delay, Neurodevelopmental disorder, Severe global developmental delay
TBX19	Adrenocorticotrophic hormone deficiency
TBX22	Cleft palate with ankyloglossia
TBX3	Ulnar-mammary syndrome
TBX4	Coxopodopatellar syndrome
TBX5	Congenital heart disease (variable), Holt-Oram syndrome
TBXAS1	Ghosal hematodiaphyseal dysplasia, Thromboxane synthetase deficiency
TCAP	Autosomal recessive limb-girdle muscular dystrophy type 2G, Dilated cardiomyopathy 1N, Primary familial hypertrophic cardiomyopathy
TCF12	Craniosynostosis 3
TCF20	Neurodevelopmental abnormality
TCF4	Intellectual disability, Pitt-Hopkins syndrome
TCN2	Inborn genetic diseases, Transcobalamin II deficiency
TCOF1	Treacher Collins syndrome 1
TCTEX1D2	Short-rib thoracic dysplasia 17 with or without polydactyly
TCTEX1D2, TM4SF19- TCTEX1D2	Short-rib thoracic dysplasia 17 with or without polydactyly
TCTN2	Joubert syndrome, Meckel syndrome type 8
TCTN3	Orofacial-digital syndrome IV
TDO2	Hypertryptophanemia, familial
TDRD7	Cataract, autosomal recessive congenital 4
TDRD9	SPERMATOGENIC FAILURE 30
TECPR2	Spastic paraplegia 49, autosomal recessive
TECTA	Deafness, Nonsyndromic hearing loss and deafness, Rare genetic deafness, autosomal dominant 12, autosomal recessive 21, neurosensory autosomal recessive 21
TENM3	MICROPTHALMIA, SYNDROMIC 15
TENT5A	Osteogenesis imperfecta, type 18
TEX14	SPERMATOGENIC FAILURE 23
TEX15	SPERMATOGENIC FAILURE 25
TFAP2B	Patent ductus arteriosus 2
TFR2	Hemochromatosis type 3
TG	Iodotyrosyl coupling defect
TGFB2	Cardiovascular phenotype, Holt-Oram syndrome, Loeys-Dietz syndrome 4
TGFB3	Cardiovascular phenotype, Loeys-Dietz syndrome 5
TGFBR1	Familial thoracic aortic aneurysm and aortic dissection
TGFBR2	Familial thoracic aortic aneurysm and aortic dissection, Hereditary nonpolyposis colorectal cancer type 6, Loeys-Dietz syndrome, Loeys-Dietz syndrome 2, Malignant tumor of esophagus
TGM1	Autosomal recessive congenital ichthyosis 1, Ichthyosis (disease)
TGM5	Peeling skin syndrome 2
TH	Segawa syndrome, autosomal recessive

THRB	Thyroid hormone resistance, autosomal dominant, generalized
TICAM1	4, Herpes simplex encephalitis, susceptibility to
TIMM8A	Deafness dystonia syndrome
TIMMDC1	Leigh syndrome
TJP2	Progressive familial intrahepatic cholestasis 4
TK2	Mitochondrial DNA depletion syndrome 2
TLR5	1, Legionellosis, Melioidosis, Systemic lupus erythematosus, resistance to
TM4SF20	Specific language impairment 5
TMC1	Deafness, Dominant, Nonsyndromic Hearing Loss, Rare genetic deafness, autosomal recessive 7
TMCO1	Craniofacial dysmorphism, and mental retardation syndrome, skeletal anomalies
TMCO6, NDUFA2	Cystic Leukoencephalopathy
TMEM127	Hereditary Paraganglioma-Pheochromocytoma Syndromes, Hereditary cancer-predisposing syndrome, Pheochromocytoma
TMEM216	Joubert syndrome, Joubert syndrome 2, Meckel syndrome, TMEM216-Related Disorders, type 2
TMEM237	Joubert syndrome
TMEM260	Structural heart defects and renal anomalies syndrome
TMEM67	Cerebellar vermis hypoplasia, Generalized hypotonia, Iris coloboma, Joubert syndrome, Joubert syndrome 6, Meckel syndrome, Meckel-Gruber syndrome, Nystagmus, TMEM67-Related Disorders, type 3
TMEM70	Mitochondrial proton-transporting ATP synthase complex deficiency, Nuclearly-encoded mitochondrial complex V (ATP synthase) deficiency 2
TMEM94	Intellectual developmental disorder with cardiac defects and dysmorphic facies
TMEM99, KRT10	Bullous ichthyosiform erythroderma
TMPRSS3	Deafness, Inborn genetic diseases, Rare genetic deafness, autosomal recessive 8
TNFRSF10B	Squamous cell carcinoma of the head and neck
TNFRSF11B	Hyperphosphatasemia with bone disease
TNFRSF13B	Absent epiphyses, Chronic lung disease, Cleft palate, Clubfoot, Coat hanger sign of ribs, Common Variable Immune Deficiency, Common variable immunodeficiency 2, Dominant, Hemivertebrae, Immunoglobulin A deficiency 2, Interstitial pulmonary abnormality, Micrognathia, Patent ductus arteriosus, Preaxial foot polydactyly, Pseudoarthrosis, Respiratory failure, Short femur, Skeletal dysplasia, Vertebral hypoplasia, Vertebral segmentation defect
TNFRSF1A	5, Familial Periodic Fever, Multiple sclerosis, susceptibility to
TNFSF11	Autosomal recessive osteopetrosis 2
TNNI3	Cardiovascular phenotype
TNNI3K, FPGT-TNNI3K	Cardiac conduction disease with or without dilated cardiomyopathy
TNNT2	Cardiomyopathy, Cardiovascular phenotype, Familial hypertrophic cardiomyopathy 2, Familial restrictive cardiomyopathy 3, Hypertrophic cardiomyopathy, Left ventricular noncompaction 6, Primary familial hypertrophic cardiomyopathy
TNPO3	Limb-girdle muscular dystrophy, type 1F

TNXB	1, Ehlers-Danlos syndrome, Ehlers-Danlos syndrome due to tenascin-X deficiency, classic-like
TONSL	Sponastrime dysplasia
TONSL, TONSL-AS1	Sponastrime dysplasia
TOP3A	AND INCREASED SISTER CHROMATID EXCHANGE 2, GROWTH RESTRICTION, MICROCEPHALY
TOPORS	Retinal dystrophy, Retinitis pigmentosa
TP53	Head and Neck Neoplasms, Hereditary cancer-predisposing syndrome, Li-Fraumeni syndrome, Li-Fraumeni syndrome 1, Li-Fraumeni-like syndrome, Multiple myeloma, Neoplasm of the large intestine, Ovarian Neoplasms
TP63	Ectrodactyly, Orofacial cleft 8, and cleft lip/palate syndrome 3, ectodermal dysplasia
TPI1	Triosephosphate isomerase deficiency
TPM2	ARTHROGRYPOSIS, DISTAL, TYPE 2B4
TPO	Deficiency of iodide peroxidase
TPP1	Ceroid lipofuscinosis neuronal 2, Childhood-onset autosomal recessive slowly progressive spinocerebellar ataxia, Inborn genetic diseases, Neuronal ceroid lipofuscinosis
TPRN	Deafness, autosomal recessive 79
TRAPPC11	Limb-girdle muscular dystrophy, type 2S
TRAPPC2	Spondyloepiphyseal dysplasia tarda
TRDN	5, Catecholaminergic polymorphic ventricular tachycardia, Ventricular tachycardia, catecholaminergic polymorphic, with or without muscle weakness
TREX1, ATRIP, ATRIP-TREX1	Aicardi Goutieres syndrome 1, Chilblain Lupus, Retinal vasculopathy with cerebral leukoencephalopathy and systemic manifestations, TREX1-Related Disorders
TRIM14, NANS	Genevieve type, Spondyloepimetaphyseal dysplasia
TRIM32, ASTN2	Limb-girdle muscular dystrophy
TRIOBP	Nonsyndromic hearing loss and deafness
TRIP11	Achondrogenesis, Goldblatt hypertension, Osteochondrodysplasia, type IA
TRMU	Acute infantile liver failure due to synthesis defect of mtDNA-encoded proteins
TRNT1	Retinitis pigmentosa and erythrocytic microcytosis, Sideroblastic anemia with B-cell immunodeficiency, and developmental delay, periodic fevers
TRPM4	Cardiomyopathy, Progressive familial heart block type IB, TRPM4-Related Disorders
TRPS1	Trichorhinophalangeal dysplasia type I
TRPV4	Charcot-Marie-Tooth disease axonal type 2C
TRPV6	HYPERPARATHYROIDISM, TRANSIENT NEONATAL
TSC1	Cortical dysplasia, Cortical tubers, Focal cortical dysplasia type II, Hereditary cancer-predisposing syndrome, Lymphangiomyomatosis, Multiple renal cysts, Renal cortical cysts, Renal insufficiency, Seizures, Tuberous sclerosis 1, Tuberous sclerosis syndrome, Urinary bladder cancer
TSC2	Focal cortical dysplasia type II, Lymphangiomyomatosis, Tuberous sclerosis 2, Tuberous sclerosis syndrome
TSMF	Combined oxidative phosphorylation deficiency 3, Primary dilated cardiomyopathy

TSHB	Secondary hypothyroidism
TSHR	1, Hypothyroidism, congenital, nongoitrous
TSHZ1	Aural atresia, congenital
TSPAN1, POMGNT1	Congenital muscular alpha-dystroglycanopathy with brain and eye anomalies, Congenital muscular dystrophy-dystroglycanopathy with mental retardation, Limb-girdle muscular dystrophy-dystroglycanopathy, Muscle eye brain disease, POMGNT1-Related Disorders, Retinitis pigmentosa 76, type B3, type C3
TSPAN12	Exudative vitreoretinopathy 5
TSPAN7	Mental retardation 58, X-linked
TSPEAR	ECTODERMAL DYSPLASIA 14, HAIR/TOOTH TYPE WITH HYPOHIDROSIS
TSPEAR-AS1, TSPEAR	Deafness, ECTODERMAL DYSPLASIA 14, HAIR/TOOTH TYPE WITH HYPOHIDROSIS, autosomal recessive 98
TTC19	Mitochondrial complex III deficiency, nuclear type 2
TTC21A	SPERMATOGENIC FAILURE 37
TTC21B, TTC21B-AS1	SHORT-RIB THORACIC DYSPLASIA 4 WITH POLYDACTYLY
TTC29	SPERMATOGENIC FAILURE 42
TTC37	Trichohepatoenteric syndrome, Trichohepatoenteric syndrome 1
TTC7A	Multiple gastrointestinal atresias
TTLL5	Cone-rod dystrophy 19
TTN	Cardiomyopathy, Cardiovascular phenotype, Dilated cardiomyopathy 1G, Limb-girdle muscular dystrophy, Myotubular myopathy, Primary dilated cardiomyopathy, Tibial muscular dystrophy, type 2J
TTN-AS1, TTN	Cardiovascular phenotype, Dilated cardiomyopathy 1G, Limb-girdle muscular dystrophy, Primary dilated cardiomyopathy, TTN-Related Disorders, type 2J
TTN, LOC101927055	Primary dilated cardiomyopathy
TTN, TTN-AS1	9, Broad-based gait, Cardiomyopathy, Cardiovascular phenotype, Congenital muscular dystrophy, Decreased patellar reflex, Delayed gross motor development, Dilated cardiomyopathy 1G, Dilated cardiomyopathy 1S, Distal muscle weakness, Familial dilated cardiomyopathy, Familial hypertrophic cardiomyopathy 9, Gowers sign, Heart murmur, Limb-girdle muscular dystrophy, Muscular dystrophy, Myopathy, Primary dilated cardiomyopathy, Proximal lower limb amyotrophy, Scoliosis, Severe muscular hypotonia, TTN-Related disorder, Tibial muscular dystrophy, Waddling gait, early-onset, myofibrillar, type 2J, with early respiratory failure, with fatal cardiomyopathy
TTPA	Ataxia, Familial isolated deficiency of vitamin E, Friedreich-like, with isolated vitamin E deficiency
TUB, RIC3	Retinal dystrophy and obesity
TUBA3D, MZT2A	KERATOCONUS 9
TUBB8	Oocyte maturation defect 2
TULP1	Leber congenital amaurosis, Retinitis pigmentosa
TWIST1	Craniosynostosis 1, Robinow-Sorauf syndrome, Saethre-Chotzen syndrome
TXNL4A	Burn-McKeown syndrome
TYK2	Tyrosine kinase 2 deficiency

TYR	3, Albinism, Inborn genetic diseases, Myopia (disease), Nonsyndromic Oculocutaneous Albinism, Nystagmus, Oculocutaneous albinism, Oculocutaneous albinism type 1B, Skin/hair/eye pigmentation, Tyrosinase-negative oculocutaneous albinism, ocular, variation in, with sensorineural deafness
TYRP1, LURAP1L-AS1	Oculocutaneous albinism type 3
UBAP1	AUTOSOMAL DOMINANT, SPASTIC PARAPLEGIA 80
UBE3A, SNHG14	Angelman syndrome, History of neurodevelopmental disorder, Inborn genetic diseases
UBE3B	Kaufman oculocerebrofacial syndrome
UBR1	Johanson-Blizzard syndrome
UCP3	Obesity, and type II diabetes, severe
UGT1A, UGT1A10, UGT1A8, UGT1A7, UGT1A6, UGT1A5, UGT1A9, UGT1A4, UGT1A1, UGT1A3	Crigler-Najjar syndrome, Crigler-Najjar syndrome type 1, type II
UNC13D	Familial hemophagocytic lymphohistiocytosis 3
UNC80	Hypotonia, Hypotonia-speech impairment-severe cognitive delay syndrome, infantile, with psychomotor retardation and characteristic facies 2
UNG	Hyper-IgM syndrome type 5
UPF3B	Mental retardation, X-linked, syndromic 14
USH1C	Deafness, Rare genetic deafness, Retinal dystrophy, Retinitis pigmentosa, Usher syndrome, Usher syndrome type 1, autosomal recessive 18, type 1C
USH2A	Abnormality of the upper limb, Abnormality of upper limb bone, Abnormality of upper limb joint, Anxiety, Brisk reflexes, Chronic pain, Cognitive impairment, Cone-rod dystrophy, Congenital sensorineural hearing impairment, Congenital stationary night blindness, Dislocated radial head, Distal arthrogyposis, Dysautonomia, Hearing impairment, High palate, Inborn genetic diseases, Macular dystrophy, Multiple joint contractures, Rare genetic deafness, Retinal dystrophy, Retinitis pigmentosa, Retinitis pigmentosa 39, Short stature, USH2A-Related Disorders, Usher syndrome, Usher syndrome type 2, type 2A
USH2A, USH2A-AS1	Rare genetic deafness, Retinal dystrophy, Retinitis pigmentosa 39, USH2A-Related Disorders, Usher syndrome, type 2A
USH2A, USH2A-AS2	Rare genetic deafness, Retinitis pigmentosa 39, Usher syndrome, type 2A
USP18	Pseudo-TORCH syndrome 2
USP27X	Mental retardation, X-linked 105
USP9X	Mental retardation, USP9X related disorders, X-linked 99, female-restricted, syndromic
VCL	Dilated cardiomyopathy 1W, Familial hypertrophic cardiomyopathy 15, Primary dilated cardiomyopathy

VHL	2, Erythrocytosis, Hereditary cancer-predisposing syndrome, Von Hippel-Lindau syndrome, familial
VHL, LOC107303340	1, 2, Erythrocytosis, Hereditary cancer-predisposing syndrome, Renal cell carcinoma, Von Hippel-Lindau syndrome, familial, papillary
VIM, VIM-AS1	Cataract 30, Congenital cataract
VIPAS39	Arthrogryposis, and cholestasis 2, renal dysfunction
VPS13A	Choreoacanthocytosis
VPS13B	Abnormality of the eye, Cohen syndrome, Inborn genetic diseases, Intellectual disability, Microcephaly, Neutropenia, Progressive visual loss, Recurrent aphthous stomatitis, Retinal dystrophy, Short foot, Short stature, Small hand
VPS33B	Arthrogryposis, Inborn genetic diseases, and cholestasis 1, renal dysfunction
VRK2, FANCL	Fanconi anemia, complementation group A, complementation group L
VWF	von Willebrand disorder
WAC	Desanto-shinawi syndrome
WAS	Wiskott-Aldrich syndrome, X-linked severe congenital neutropenia, X-linked thrombocytopenia with normal platelets
WDR35	Cranioectodermal dysplasia, Cranioectodermal dysplasia 2, Jeune thoracic dystrophy, SHORT-RIB THORACIC DYSPLASIA 7 WITHOUT POLYDACTYLY, Short Rib Polydactyly Syndrome, Short rib polydactyly syndrome 5, Short-rib thoracic dysplasia 7/20 with polydactyly, WDR35-Related Disorders, digenic
WDR45	Neurodegeneration with brain iron accumulation, Neurodegeneration with brain iron accumulation 5
WDR72	Amelogenesis imperfecta
WDR73	Galloway-Mowat syndrome 1
WEE2-AS1, WEE2	OOCYTE MATURATION DEFECT 5
WFS1	Autosomal dominant nonsyndromic deafness 6, Diabetes mellitus AND insipidus with optic atrophy AND deafness, WFS1-Related Spectrum Disorders, Wolfram-like syndrome, autosomal dominant
WHRN	Deafness, Rare genetic deafness, Usher syndrome, autosomal recessive 31, type 2D
WRN	Medulloblastoma, Werner syndrome
WT1	Drash syndrome, Frasier syndrome, Wilms tumor, Wilms tumor 1, and mental retardation syndrome, aniridia, genitourinary anomalies
WT1, LOC107982234	Drash syndrome, Frasier syndrome, Pre-B-cell acute lymphoblastic leukemia, Wilms tumor, Wilms tumor 1, and mental retardation syndrome, aniridia, genitourinary anomalies
XDH	Deficiency of xanthine oxidase
XIAP	Lymphoproliferative syndrome 2, X-linked
XK	McLeod neuroacanthocytosis syndrome
XPA	Xeroderma pigmentosum, Xeroderma pigmentosum group A
XPC	Xeroderma pigmentosum, group C
XRCC2	Fanconi anemia, Hereditary Cancer Syndrome, Hereditary breast and ovarian cancer syndrome, Hereditary cancer-predisposing syndrome, Ovarian Neoplasms, complementation group U

XRCC4	Short stature, and endocrine dysfunction, microcephaly
XYLT1	Desbuquois dysplasia 2
XYLT1, LOC102723692	Desbuquois dysplasia 2
XYLT2	Inborn genetic diseases, Spondyloocular syndrome, autosomal recessive
YY1AP1	Grange syndrome
ZBTB18	Mental retardation, autosomal dominant 22
ZDBF2	Nasopalpebral lipoma-coloboma syndrome
ZEB2	Mowat-Wilson syndrome
ZFYVE26	Hereditary spastic paraplegia 15, Spastic paraplegia
ZFYVE26, RDH12	Abnormality of the eye, Leber congenital amaurosis 13, RDH12-Related Disorders, Retinal dystrophy, Retinitis pigmentosa
ZMPSTE24	Lethal tight skin contracture syndrome, Mandibuloacral dysplasia with type B lipodystrophy, ZMPSTE24-Related Disorders
ZNF408	Retinitis pigmentosa 72
ZNF462	Craniosynostosis, Mental retardation, WEISS-KRUSZKA SYNDROME, autosomal dominant
ZNF711	ZNF711-Related X-linked Mental Retardation
ZP1	Oocyte maturation defect 1
ZP2	OOCYTE MATURATION DEFECT 6

Use of TREMs

A TREM composition (*e.g.*, a pharmaceutical TREM composition described herein) can modulate a function in a cell, tissue or subject having an endogenous ORF having a codon comprising a first sequence, *e.g.*, a mutation, *e.g.*, a premature termination codon.

In embodiments, a TREM composition (*e.g.*, a pharmaceutical TREM composition) described herein is contacted with a cell or tissue, or administered to a subject in need thereof, in an amount and for a time sufficient to modulate a production parameter of an RNA corresponding to, or a protein encoded by an endogenous ORF having a first sequence, *e.g.*, a mutation, *e.g.*, a premature termination codon.

In embodiments, a TREM composition (*e.g.*, a pharmaceutical TREM composition) described herein is contacted with a cell or tissue, or administered to a subject in need thereof, in an amount and for a time sufficient to modulate expression of a protein encoded by an endogenous ORF having a first sequence, *e.g.*, a mutation, *e.g.*, a premature termination codon.

In embodiments, a TREM composition (*e.g.*, a pharmaceutical TREM composition described

herein is contacted with a cell or tissue, or administered to a subject in need thereof, in an amount and for a time sufficient to treat a disease or disorder associated with a PTC, e.g., as described herein.

5 *Methods of modulating a production parameter of an RNA corresponding to, or a protein encoded by an endogenous ORF having a PTC with a TREM composition*

A production parameter of an RNA corresponding to, or a protein encoded by a nucleic acid sequence comprising an endogenous ORF having a codon having a first sequence, e.g., a mutation, e.g., a premature termination codon, can be modulated by administration of a TREM
10 composition comprising a TREM which pairs with, e.g., recognizes the codon having the first sequence.

In an aspect, provided herein is a method of modulating a production parameter of an RNA corresponding to, or a protein encoded by, a nucleic acid sequence comprising an endogenous ORF having a codon having a first sequence, e.g., a mutation, e.g., a premature
15 termination codon, in a target cell or tissue, comprising:

providing, e.g., administering, to the target cell or tissue, or contacting the target cell or tissue with, an effective amount of a TREM composition, e.g., comprising a TREM, TREM fragment or TREM core fragment,

thereby modulating the production parameter of the RNA, or protein in the target cell or
20 tissue.

The TREM composition can be administered to the subject or the target cell or tissue can be contacted ex vivo with the TREM composition.

Modulation of a production parameter of an RNA corresponding to, or a protein encoded by a nucleic acid sequence comprising an endogenous ORF having a codon having a first
25 sequence, e.g., a mutation, e.g., a premature termination codon, by administration of a TREM composition, e.g., comprising a TREM, TREM fragment or TREM core fragment, comprises modulation of an expression parameter and/or a signaling parameter, e.g., as described herein.

For example, administration of a TREM composition to a target cell or tissue can result in an increase or decrease in any one or more of the following expression parameters for the RNA
30 corresponding to, or protein encoded by a nucleic acid sequence comprising the endogenous ORF having the first sequence, e.g., mutation, e.g., PTC:

- (a) protein translation;
- (b) expression level (*e.g.*, of polypeptide or protein, or mRNA);
- (c) post-translational modification of polypeptide or protein;
- (d) folding (*e.g.*, of polypeptide or protein, or mRNA),
- 5 (e) structure (*e.g.*, of polypeptide or protein, or mRNA),
- (f) transduction (*e.g.*, of polypeptide or protein),
- (g) compartmentalization (*e.g.*, of polypeptide or protein, or mRNA),
- (h) incorporation (*e.g.*, of polypeptide or protein, or mRNA) into a supermolecular structure, *e.g.*, incorporation into a membrane, proteasome, or ribosome,
- 10 (i) incorporation into a multimeric polypeptide, *e.g.*, a homo or heterodimer, and/or
- (j) stability.

As another example, administration of a TREM composition to a target cell or tissue can result in an increase or decrease in any one or more of the following signaling parameters for the RNA corresponding to, or protein encoded by a nucleic acid sequence comprising the

15 endogenous ORF having the first sequence, *e.g.*, mutation, *e.g.*, PTC:

- (1) modulation of a signaling pathway, *e.g.*, a cellular signaling pathway which is downstream or upstream of the protein encoded by the endogenous ORF comprising the PTC;
- (2) cell fate modulation;
- (3) ribosome occupancy modulation;
- 20 (4) protein translation modulation;
- (5) mRNA stability modulation;
- (6) protein folding and structure modulation;
- (7) protein transduction or compartmentalization modulation; and/or
- (8) protein stability modulation.

25 A production parameter (*e.g.*, an expression parameter and/or a signaling parameter) may be modulated, *e.g.*, increased, *e.g.*, by at least 5% (*e.g.*, at least 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 150%, 200% or more) compared to a reference, *e.g.*, an RNA corresponding to or a polypeptide encoded by a nucleic acid sequence comprising an endogenous ORF having a non-mutated codon, *e.g.*, wildtype codon. In some embodiments, the

30 reference polypeptide encoded by the endogenous ORF having a non-mutated codon comprises a

pre-mutation amino acid, e.g., wildtype amino acid, at the position corresponding to the non-mutated codon.

In some embodiments, the production parameter (*e.g.*, an expression parameter and/or a signaling parameter) is increased by at least about 10%, about 20%, about 30%, about 40%,
5 about 50%, about 60%, about 70%, about 80%, about 90%, about 100%, about 110%, about 120%, about 130%, about 140%, about 150%, about 160%, about 170%, about 180%, about 190%, about 200%, about 250%, about 300%, about 350%, about 400%, about 450%, about 500%, about 600%, about 700%, about 800%, about 900%, about 1000% , or more compared to a reference, e.g., as described herein.

10 In some embodiments, the production parameter (*e.g.*, an expression parameter and/or a signaling parameter) is increased from about 100% to about 1000%, about 100% to about 900%, about 100% to about 800%, about 100% to about 700%, about 100% to about 600%, about 100% to about 500%, about 100% to about 400%, about 100% to about 300%, about 100% to about 200%, about 200% to about 1000%, about 200% to about 900%, about 200% to about 800%,
15 about 200% to about 700%, about 200% to about 600%, about 200% to about 500%, about 200% to about 400%, about 200% to about 300%, about 300% to about 1000%, about 300% to about 900%, about 300% to about 800%, about 300% to about 700%, about 300% to about 600%, about 300% to about 500%, about 300% to about 400%, about 400% to about 1000%, about 400% to about 900%, about 400% to about 800%, about 400% to about 700%, about 400% to
20 about 600%, about 400% to about 500%, about 500% to about 1000%, about 500% to about 900%, about 500% to about 800%, about 500% to about 700%, about 500% to about 600%, about 600% to about 1000%, about 600% to about 900%, about 600% to about 800%, about 600% to about 700%, about 700% to about 1000%, about 700% to about 900%, about 700% to about 800%, about 800% to about 1000%, about 800% to about 900%, or about 900% to about
25 1000% compared to a reference, e.g., as described herein.

In some embodiments, the production parameter (*e.g.*, an expression parameter and/or a signaling parameter) is decreased by at least about 10%, about 20%, about 30%, about 40%,
about 50%, about 60%, about 70%, about 80%, about 90%, about 100%, about 110%, about 120%, about 130%, about 140%, about 150%, about 160%, about 170%, about 180%, about
30 190%, about 200%, about 250%, about 300%, about 350%, about 400%, about 450%, about

500%, about 600%, about 700%, about 800%, about 900%, about 1000% , or more compared to a reference, e.g., as described herein.

In some embodiments, the production parameter (*e.g.*, an expression parameter and/or a signaling parameter) is decreased from about 100% to about 1000%, about 100% to about 900%, about 100% to about 800%, about 100% to about 700%, about 100% to about 600%, about 100% to about 500%, about 100% to about 400%, about 100% to about 300%, about 100% to about 200%, about 200% to about 1000%, about 200% to about 900%, about 200% to about 800%, about 200% to about 700%, about 200% to about 600%, about 200% to about 500%, about 200% to about 400%, about 200% to about 300%, about 300% to about 1000%, about 300% to about 900%, about 300% to about 800%, about 300% to about 700%, about 300% to about 600%, about 300% to about 500%, about 300% to about 400%, about 400% to about 1000%, about 400% to about 900%, about 400% to about 800%, about 400% to about 700%, about 400% to about 600%, about 400% to about 500%, about 500% to about 1000%, about 500% to about 900%, about 500% to about 800%, about 500% to about 700%, about 500% to about 600%, about 600% to about 1000%, about 600% to about 900%, about 600% to about 800%, about 600% to about 700%, about 700% to about 1000%, about 700% to about 900%, about 700% to about 800%, about 800% to about 1000%, about 800% to about 900%, or about 900% to about 1000% compared to a reference, e.g., as described herein.

A production parameter described herein may be measured by any method known in the art. For example Western blotting can be used to measure protein levels and quantitative RT-PCR or Northern blotting can be used to measure RNA levels.

Methods of modulating expression of a protein encoded by an endogenous ORF having a PTC with a TREM composition

Expression and/or activity of a protein encoded by a nucleic acid sequence comprising an endogenous ORF having a codon having a first sequence, e.g., a mutation, e.g., a premature termination codon, can be modulated by administration of a TREM composition comprising a TREM which pairs with, e.g., recognizes the codon having the first sequence.

In an aspect, provided herein is a method of modulating the expression and/or activity of a protein encoded by a nucleic acid sequence comprising an endogenous ORF having a codon having a first sequence, e.g., a mutation, e.g., a premature termination codon, in a target cell or tissue, comprising:

providing, *e.g.*, administering, to the target cell or tissue, or contacting the target cell or tissue with, an effective amount of a TREM composition, *e.g.*, comprising a TREM, TREM fragment or TREM core fragment,

5 thereby modulating the expression and/or activity of the protein in the target cell or tissue.

In some embodiments, the expression and/or activity of a polypeptide encoded by an endogenous ORF having a codon comprising a first sequence, *e.g.*, a mutation, *e.g.*, a PTC, is increased by at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 100%, about 110%, about 120%, about 130%, about 10
140%, about 150%, about 160%, about 170%, about 180%, about 190%, about 200%, about 250%, about 300%, about 350%, about 400%, about 450%, about 500%, about 600%, about 700%, about 800%, about 900%, about 1000% , or more compared to a reference, *e.g.*, as described herein.

In some embodiments, the expression and/or activity of a polypeptide encoded by the
15 endogenous ORF having a codon comprising a first sequence, *e.g.*, a mutation, *e.g.*, a PTC, is increased from about 100% to about 1000%, about 100% to about 900%, about 100% to about 800%, about 100% to about 700%, about 100% to about 600%, about 100% to about 500%, about 100% to about 400%, about 100% to about 300%, about 100% to about 200%, about 200% to about 1000%, about 200% to about 900%, about 200% to about 800%, about 200% to about
20 700%, about 200% to about 600%, about 200% to about 500%, about 200% to about 400%, about 200% to about 300%, about 300% to about 1000%, about 300% to about 900%, about 300% to about 800%, about 300% to about 700%, about 300% to about 600%, about 300% to about 500%, about 300% to about 400%, about 400% to about 1000%, about 400% to about 900%, about 400% to about 800%, about 400% to about 700%, about 400% to about 600%,
25 about 400% to about 500%, about 500% to about 1000%, about 500% to about 900%, about 500% to about 800%, about 500% to about 700%, about 500% to about 600%, about 600% to about 1000%, about 600% to about 900%, about 600% to about 800%, about 600% to about 700%, about 700% to about 1000%, about 700% to about 900%, about 700% to about 800%, about 800% to about 1000%, about 800% to about 900%, or about 900% to about 1000%
30 compared to a reference, *e.g.*, as described herein.

In some embodiments, the expression and/or activity of a polypeptide encoded by the endogenous ORF having a codon comprising a first sequence, e.g., a mutation, e.g., a PTC, is decreased by at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 100%, about 110%, about 120%, about 130%, about 140%, about 150%, about 160%, about 170%, about 180%, about 190%, about 200%, about 250%, about 300%, about 350%, about 400%, about 450%, about 500%, about 600%, about 700%, about 800%, about 900%, about 1000% , or more compared to a reference, e.g., as described herein.

In some embodiments, the expression and/or activity of a polypeptide encoded by the endogenous ORF having a codon comprising a first sequence, e.g., a mutation, e.g., a PTC, is decreased from about 100% to about 1000%, about 100% to about 900%, about 100% to about 800%, about 100% to about 700%, about 100% to about 600%, about 100% to about 500%, about 100% to about 400%, about 100% to about 300%, about 100% to about 200%, about 200% to about 1000%, about 200% to about 900%, about 200% to about 800%, about 200% to about 700%, about 200% to about 600%, about 200% to about 500%, about 200% to about 400%, about 300% to about 1000%, about 300% to about 900%, about 300% to about 800%, about 300% to about 700%, about 300% to about 600%, about 300% to about 500%, about 300% to about 400%, about 400% to about 1000%, about 400% to about 900%, about 400% to about 800%, about 400% to about 700%, about 400% to about 600%, about 400% to about 500%, about 500% to about 1000%, about 500% to about 900%, about 500% to about 800%, about 500% to about 700%, about 500% to about 600%, about 600% to about 1000%, about 600% to about 900%, about 600% to about 800%, about 600% to about 700%, about 700% to about 1000%, about 700% to about 900%, about 700% to about 800%, about 800% to about 1000%, about 800% to about 900%, or about 900% to about 1000% compared to a reference, e.g., as described herein.

In some embodiments, the reference comprises a polypeptide encoded by an endogenous ORF having a non-mutated codon, e.g., wildtype codon. In some embodiments, the reference polypeptide encoded by the endogenous ORF having a non-mutated codon comprises a pre-mutation amino acid, e.g., wildtype amino acid, at the position corresponding to the non-mutated codon.

Methods of treating a subject having an endogenous ORF having a PTC with a TREM composition

In an aspect, provided herein is a method of treating a subject having an endogenous open reading frame (ORF) which comprises a codon having a first sequence, comprising:

5 providing a TREM composition comprising a TREM disclosed herein, wherein the TREM comprises a tRNA moiety having an anticodon that pairs with the codon of the ORF having the first sequence;

contacting the subject with the TREM composition in an amount and/or for a time sufficient to treat the subject,

10 thereby treating the subject.

In an embodiment, the subject has a disease or disorder associated with a PTC, e.g., as provided in any one of Tables 15-17.

In an embodiment, the subject has an ORF comprising a PTC in a gene disclosed in any one of Tables 15, 16 or 18.

15

TREM, TREM core fragment and TREM fragment

A “tRNA-based effector molecule” or “TREM” refers to an RNA molecule comprising one or more of the properties described herein. A TREM can comprise a non-naturally occurring modification, e.g., as provided in Tables 4, 5, 6 or 7.

20 In an embodiment, a TREM includes a TREM comprising a sequence of Formula A; a TREM core fragment comprising a sequence of Formula B; or a TREM fragment comprising a portion of a TREM which TREM comprises a sequence of Formula A.

In an embodiment, a TREM comprises a sequence of Formula A: [L1]-[ASt Domain1]-[L2]-[DH Domain]-[L3]-[ACH Domain]-[VL Domain]-[TH Domain]-[L4]-[ASt Domain2]. In an embodiment, [VL Domain] is optional. In an embodiment, [L1] is optional.

25 In an embodiment, a TREM core fragment comprises a sequence of Formula B: [L1]_y-[ASt Domain1]_x-[L2]_y-[DH Domain]_y-[L3]_y-[ACH Domain]_x-[VL Domain]_y-[TH Domain]_y-[L4]_y-[ASt Domain2]_x, wherein: x=1 and y=0 or 1. In an embodiment, y=0. In an embodiment, y=1.;

30 In an embodiment, a TREM fragment comprises a portion of a TREM, wherein the TREM comprises a sequence of Formula A: [L1]-[ASt Domain1]-[L2]-[DH Domain]-[L3]-[ACH Domain]-[VL Domain]-[TH Domain]-[L4]-[ASt Domain2], and wherein the TREM

fragment comprises: one, two, three or all or any combination of the following: a TREM half (*e.g.*, from a cleavage in the ACH Domain, *e.g.*, in the anticodon sequence, *e.g.*, a 5' half or a 3' half); a 5' fragment (*e.g.*, a fragment comprising the 5' end, *e.g.*, from a cleavage in a DH Domain or the ACH Domain); a 3' fragment (*e.g.*, a fragment comprising the 3' end, *e.g.*, from a cleavage in the TH Domain); or an internal fragment (*e.g.*, from a cleavage in any one of the ACH Domain, DH Domain or TH Domain). Exemplary TREM fragments include TREM halves (*e.g.*, from a cleavage in the ACHD, *e.g.*, 5' TREM halves or 3' TREM halves), a 5' fragment (*e.g.*, a fragment comprising the 5' end, *e.g.*, from a cleavage in a DHD or the ACHD), a 3' fragment (*e.g.*, a fragment comprising the 3' end of a TREM, *e.g.*, from a cleavage in the THD), or an internal fragment (*e.g.*, from a cleavage in one or more of the ACHD, DHD or THD).

In an embodiment, a TREM, a TREM core fragment or a TREM fragment can be charged with an amino acid (*e.g.*, a cognate amino acid); charged with a non-cognate amino acid (*e.g.*, a mischarged TREM (mTREM)); or not charged with an amino acid (*e.g.*, an uncharged TREM (uTREM)). In an embodiment, a TREM, a TREM core fragment or a TREM fragment can be charged with an amino acid selected from alanine, arginine, asparagine, aspartate, cysteine, glutamine, glutamate, glycine, histidine, isoleucine, methionine, leucine, lysine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, or valine.

In an embodiment, the TREM, TREM core fragment or TREM fragment is a cognate TREM. In an embodiment, the TREM, TREM core fragment or TREM fragment is a non-cognate TREM. In an embodiment, the TREM, TREM core fragment or TREM fragment recognizes a codon provided in Table 7 or Table 8.

Table 7: List of codons

AAA	AGG
AAC	AGU
AAG	AUA
AAU	AUC
ACA	AUG
ACC	AUU
ACG	CAA
ACU	CAC
AGA	CAG
AGC	CAU

CCA	GGG
CCC	GGU
CCG	GUA
CCU	GUC
CGA	GUG
CGC	GUU
CGG	UAA
CGU	UAC
CUA	UAG
CUC	UAU
CUG	UCA
CUU	UCC
GAA	UCG
GAC	UCU
GAG	UGA
GAU	UGC
GCA	UGG
GCC	UGU
GCG	UUA
GCU	UUC
GGA	UUG
GGC	UUU

Table 8: Amino acids and corresponding codons

Amino Acid	mRNA codons
Alanine	GCU, GCC, GCA, GCG
Arginine	CGU, CGC, CGA, CGG, AGA, AGG
Asparagine	AAU, AAC
Aspartate	GAU, GAC
Cysteine	UGU, UGC
Glutamate	GAA, GAG
Glutamine	CAA, CAG
Glycine	GGU, GGC, GGA, GGG
Histidine	CAU, CAC
Isoleucine	AUU, AUC, AUA
Leucine	UUA, UUG, CUU, CUC, CUA, CUG
Lysine	AAA, AAG
Methionine	AUG
Phenylalanine	UUU, UUC
Proline	CCU, CCC, CCA, CCG

Serine	UCU, UCC, UCA, UCG, AGU, AGC
Stop	UAA, UAG, UGA
Threonine	ACU, ACC, ACA, ACG
Tryptophan	UGG
Tyrosine	UAU, UAC
Valine	GUU, GUC, GUA, GUG

In an embodiment, a TREM comprises a ribonucleic acid (RNA) sequence encoded by a deoxyribonucleic acid (DNA) sequence disclosed in Table 9, *e.g.*, any one of SEQ ID NOs: 1-451 disclosed in Table 9. In an embodiment, a TREM comprises an RNA sequence at least 60%, 65%, 70%, 75%, 80%, 82%, 85%, 87%, 88%, 90%, 92%, 95%, 96%, 97%, 98%, or 99% identical to an RNA sequence encoded by a DNA sequence provided in Table 9, *e.g.*, any one of SEQ ID NOs: 1-451 disclosed in Table 9. In an embodiment, a TREM comprises an RNA sequence encoded by a DNA sequence at least 60%, 65%, 70%, 75%, 80%, 82%, 85%, 87%, 88%, 90%, 92%, 95%, 96%, 97%, 98%, or 99% identical to a DNA sequence provided in Table 9, *e.g.*, any one of SEQ ID NOs: 1-451 disclosed in Table 9.

In an embodiment, a TREM, a TREM core fragment, or TREM fragment comprises at least 5, 10, 15, 20, 25, or 30 consecutive nucleotides of an RNA sequence encoded by a DNA sequence disclosed in Table 9, *e.g.*, at least 5, 10, 15, 20, 25, or 30 consecutive nucleotides of an RNA sequence encoded by any one of SEQ ID NOs: 1-451 disclosed in Table 9. In an embodiment, a TREM, a TREM core fragment, or TREM fragment comprises at least 5, 10, 15, 20, 25, or 30 consecutive nucleotides of an RNA sequence at least 60%, 65%, 70%, 75%, 80%, 82%, 85%, 87%, 88%, 90%, 92%, 95%, 96%, 97%, 98%, or 99% identical to an RNA sequence encoded by a DNA sequence provided in Table 9, *e.g.*, any one of SEQ ID NOs: 1-451 disclosed in Table 9. In an embodiment, a TREM, a TREM core fragment, or TREM fragment comprises at least 5, 10, 15, 20, 25, or 30 consecutive nucleotides of an RNA sequence encoded by a DNA sequence at least 60%, 65%, 70%, 75%, 80%, 82%, 85%, 87%, 88%, 90%, 92%, 95%, 96%, 97%, 98%, or 99% identical to a DNA sequence provided in Table 9, *e.g.*, any one of SEQ ID NOs: 1-451 disclosed in Table 9.

In an embodiment, a TREM core fragment or a TREM fragment comprises at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% of an RNA sequence encoded by a DNA sequence provided in Table 9, *e.g.*, any one of SEQ ID NOs: 1-451 disclosed in Table 9. In an embodiment, a TREM

core fragment or a TREM fragment comprises at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% of an RNA sequence at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to an RNA sequence encoded by a DNA sequence provided in Table 9, *e.g.*, any one of SEQ ID NOs: 1-451 disclosed in Table 9. In an embodiment, a TREM core fragment or a TREM fragment comprises at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% of an RNA sequence encoded by a DNA sequence at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to a DNA sequence provided in Table 9, *e.g.*, any one of SEQ ID NOs: 1-451 disclosed in Table 9.

In an embodiment, a TREM core fragment or a TREM fragment comprises at least 5 ribonucleotides (nt), 10 nt, 15 nt, 20 nt, 25 nt, 30 nt, 35 nt, 40 nt, 45 nt, 50 nt, 55 nt or 60 nt (but less than the full length) of an RNA sequence encoded by a DNA sequence disclosed in Table 9, *e.g.*, any one of SEQ ID NOs: 1-451 disclosed in Table 9. In an embodiment, a TREM core fragment or a TREM fragment comprises at least 5 ribonucleotides (nt), 10 nt, 15 nt, 20 nt, 25 nt, 30 nt, 35 nt, 40 nt, 45 nt, 50 nt, 55 nt or 60 nt (but less than the full length) of an RNA sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to an RNA sequence encoded by a DNA sequence provided in Table 9, *e.g.*, any one of SEQ ID NOs: 1-451 disclosed in Table 9. In an embodiment, a TREM core fragment or a TREM fragment comprises at least 5 ribonucleotides (nt), 10 nt, 15 nt, 20 nt, 25 nt, 30 nt, 35 nt, 40 nt, 45 nt, 50 nt, 55 nt or 60 nt (but less than the full length) of an RNA sequence encoded by a DNA sequence with at least 80%, 82%, 85%, 87%, 88%, 90%, 92%, 95%, 96%, 97%, 98%, 99% or 100% identity to a DNA sequence provided in Table 9, *e.g.*, any one of SEQ ID NOs: 1-451 disclosed in Table 9.

In an embodiment, a TREM core fragment or a TREM fragment comprises a sequence of a length of between 10-90 ribonucleotides (rnt), between 10-80 rnt, between 10-70 rnt, between 10-60 rnt, between 10-50 rnt, between 10-40 rnt, between 10-30 rnt, between 10-20 rnt, between 20-90 rnt, between 20-80 rnt, 20-70 rnt, between 20-60 rnt, between 20-50 rnt, between 20-40 rnt, between 30-90 rnt, between 30-80 rnt, between 30-70 rnt, between 30-60 rnt, or between 30-50 rnt.

Table 9. List of tRNA sequences

SEQ ID NO	tRNA name	tRNA sequence
1	Ala_AGC_chr6:28763741-28763812 (-)	GGGGGTATAGCTCAGTGTAGAGCGCGTGTGCTTAGCATGCACGAGGTCC TGGGTTGATCCCCAGTACCTCCA
2	Ala_AGC_chr6:26687485-26687557 (+)	GGGGAATTAGCTCAAGTGGTAGAGCGCTTGCTTAGCACGCAAGAGGTA GTGGGATCGATGCCACATCTCCA
3	Ala_AGC_chr6:26572092-26572164 (-)	GGGGAATTAGCTCAATGGTAGAGCGCTCGCTTAGCATGCGAGAGGTA GCGGGATCGATGCCCGCATCTCCA
4	Ala_AGC_chr6:26682715-26682787 (+)	GGGGAATTAGCTCAAGTGGTAGAGCGCTTGCTTAGCATGCAAGAGGTA GTGGGATCGATGCCACATCTCCA
5	Ala_AGC_chr6:26705606-26705678 (+)	GGGGAATTAGCTCAAGCGGTAGAGCGCTTGCTTAGCATGCAAGAGGTA GTGGGATCGATGCCACATCTCCA
6	Ala_AGC_chr6:26673590-26673662 (+)	GGGGAATTAGCTCAAGTGGTAGAGCGCTTGCTTAGCATGCAAGAGGTA GTGGGATCAATGCCACATCTCCA
7	Ala_AGC_chr14:89445442-89445514 (+)	GGGGAATTAGCTCAAGTGGTAGAGCGCTCGCTTAGCATGCGAGAGGTA GTGGGATCGATGCCCGCATCTCCA
8	Ala_AGC_chr6:58196623-58196695 (-)	GGGGAATTAGCTCAAGTGGTAGAGCGCTTGCTTAGCATGCAAGAGGTA GTGGGATCGATGCCACATCTCCA
9	Ala_AGC_chr6:28806221-28806292 (-)	GGGGGTGTAGCTCAGTGGTAGAGCGCGTGTGCTTAGCATGCACGAGGCC CGGGTTCAATCCCCGGCACCTCCA
10	Ala_AGC_chr6:28574933-28575004 (+)	GGGGGTGTAGCTCAGTGGTAGAGCGCGTGTGCTTAGCATGTACGAGGTCC CGGGTTCAATCCCCGGCACCTCCA
11	Ala_AGC_chr6:28626014-28626085 (-)	GGGGATGTAGCTCAGTGGTAGAGCGCATGTGCTTAGCATGCATGAGGTCC CGGGTTGATCCCCAGCATCTCCA
12	Ala_AGC_chr6:28678366-28678437 (+)	GGGGGTGTAGCTCAGTGGTAGAGCGCGTGTGCTTAGCATGCACGAGGCC TGGGTTCAATCCCCAGCACCTCCA
13	Ala_AGC_chr6:28779849-28779920 (-)	GGGGGTATAGCTCAGCGGTAGAGCGCGTGTGCTTAGCATGCACGAGGTCC TGGGTTCAATCCCCAATACCTCCA

14	Ala_AGC_chr6:28687481-28687552 (+)	GGGGGTGTAGCTCAGTGGTAGAGCGCGTGTGCTTAGCATGCACGAGGCC CGGGTTCAATCCCTGGCACCTCCA
15	Ala_AGC_chr2:27274082-27274154 (+)	GGGGATTAGCTCAAATGGTAGAGCGCTCGCTTAGCATGCGAGAGGTA GCGGGATCGATGCCCGCATCTCCA
16	Ala_AGC_chr6:26730737-26730809 (+)	GGGGAATTAGCTCAGGCGGTAGAGCGCTCGCTTAGCATGCGAGAGGTA GCGGGATCGACGCCCGCATCTCCA
17	Ala_CGC_chr6:26553731-26553802 (+)	GGGGATGTAGCTCAGTGGTAGAGCGCATGTCTTCGCATGTATGAGGTCC CGGGTTCGATCCCCGGCATCTCCA
18	Ala_CGC_chr6:28641613-28641684 (-)	GGGGATGTAGCTCAGTGGTAGAGCGCATGTCTTCGCATGTATGAGGCC CGGGTTCGATCCCCGGCATCTCCA
19	Ala_CGC_chr2:157257281-157257352 (+)	GGGGATGTAGCTCAGTGGTAGAGCGCGCTTCGCATGTGTGAGGTCC CGGGTTCAATCCCCGGCATCTCCA
20	Ala_CGC_chr6:28697092-28697163 (+)	GGGGGTGTAGCTCAGTGGTAGAGCGCGTGTCTTCGCATGTACGAGGCC CGGGTTCGACCCCGGCTCCTCCA
21	Ala_TGC_chr6:28757547-28757618 (-)	GGGGGTGTAGCTCAGTGGTAGAGCGCATGTCTTTGCATGTATGAGGTCC CGGGTTCGATCCCCGGCACCTCCA
22	Ala_TGC_chr6:28611222-28611293 (+)	GGGGATGTAGCTCAGTGGTAGAGCGCATGTCTTTGCATGTATGAGGTCC CGGGTTCGATCCCCGGCATCTCCA
23	Ala_TGC_chr5:180633868-180633939 (+)	GGGGATGTAGCTCAGTGGTAGAGCGCATGTCTTTGCATGTATGAGGCC CGGGTTCGATCCCCGGCATCTCCA
24	Ala_TGC_chr12:125424512-125424583 (+)	GGGGATGTAGCTCAGTGGTAGAGCGCATGTCTTTGCACGTATGAGGCC CGGGTTCAATCCCCGGCATCTCCA
25	Ala_TGC_chr6:28785012-28785083 (-)	GGGGGTGTAGCTCAGTGGTAGAGCGCATGTCTTTGCATGTATGAGGCCT CGGGTTCGATCCCCGACACCTCCA
26	Ala_TGC_chr6:28726141-28726212 (-)	GGGGGTGTAGCTCAGTGGTAGAGCACATGTCTTTGCATGTGTGAGGCC CGGGTTCGATCCCCGGCACCTCCA
27	Ala_TGC_chr6:28770577-28770647 (-)	GGGGGTGTAGCTCAGTGGTAGAGCGCATGTCTTTGCATGTATGAGGCCT CGGGTTCGATCCCCGACACCTCCA
28	Arg_ACG_chr6:26328368-26328440 (+)	GGGCCAGTGGCGCAATGGATAACGCGTCTGACTACGGATCAGAAGATT CCAGGTTCCGACTCCTGGCTGGCTCG
29	Arg_ACG_chr3:45730491-45730563 (-)	GGGCCAGTGGCGCAATGGATAACGCGTCTGACTACGGATCAGAAGATT CTAGGTTCCGACTCCTGGCTGGCTCG

30	Arg_CCG_chr6:28710729-28710801 (-)	GGCCGCGTGGCCTAATGGATAAGGCGTCTGATTCCGGATCAGAAAGATT GAGGGTTTCGAGTCCCTTCGTGGTCG
31	Arg_CCG_chr17:66016013-66016085 (-)	GACCCAGTGGCCTAATGGATAAGGCATCAGCCCTCCGGAGCTGGGGATT GTGGGTTTCGAGTCCCATCTGGGTCG
32	Arg_CCT_chr17:73030001-73030073 (+)	GCCCCAGTGGCCTAATGGATAAGGCATCTGGCCTCCTAAGCCAGGGATT GTGGGTTTCGAGTCCACCTGGGGTA
33	Arg_CCT_chr17:73030526-73030598 (-)	GCCCCAGTGGCCTAATGGATAAGGCATCTGGCCTCCTAAGCCAGGGATT GTGGGTTTCGAGTCCACCTGGGGTG
34	Arg_CCT_chr16:3202901-3202973 (+)	GCCCCGTGGCCTAATGGATAAGGCATCTGGCCTCCTAAGCCAGGGATT GTGGGTTTCGAGTCCACCCGGGTA
35	Arg_CCT_chr7:139025446-139025518 (+)	GCCCCAGTGGCCTAATGGATAAGGCATCTGGCCTCCTAAGCCAGGGATT GTGGGTTTCGAGTCCCATCTGGGGTG
36	Arg_CCT_chr16:3243918-3243990 (+)	GCCCCAGTGGCCTGATGGATAAGGTACTGGCCTCCTAAGCCAGGGATT GTGGGTTTCGAGTCCACCTGGGGTA
37	Arg_TCG_chr15:89878304-89878376 (+)	GGCCGCGTGGCCTAATGGATAAGGCGTCTGACTTCGGATCAGAAAGATT GCAGGTTTCGAGTCCCTGCCCGGGTCG
38	Arg_TCG_chr6:26323046-26323118 (+)	GACCACGTGGCCTAATGGATAAGGCGTCTGACTTCGGATCAGAAAGATT GAGGGTTTCGAATCCCTCCGTGGTTA
39	Arg_TCG_chr17:73031208-73031280 (+)	GACC CGTGGCCTAATGGATAAGGCGTCTGACTTCGGATCAGAAAGATT GAGGGTTTCGAGTCCCTTCGTGGTCG
40	Arg_TCG_chr6:26299905-26299977 (+)	GACCACGTGGCCTAATGGATAAGGCGTCTGACTTCGGATCAGAAAGATT GAGGGTTTCGAATCCCTTCGTGGTTA
41	Arg_TCG_chr6:28510891-28510963 (-)	GACCACGTGGCCTAATGGATAAGGCGTCTGACTTCGGATCAGAAAGATT GAGGGTTTCGAATCCCTTCGTGGTTG
42	Arg_TCG_chr9:112960803-112960875 (+)	GGCCGTGGCCTAATGGATAAGGCGTCTGACTTCGGATCAAAAAGATT GCAGGTTTCGAGTCTGCCACGGTCG
43	Arg_TCT_chr1:94313129-94313213 (+)	GGTCCGTGGCGCAATGGATAGCGCATTTGGACTTCTAGAGGCTGAAGG CATTCAAAGGTTCCGGGTCGAGTCCCGGGGAGTCG
44	Arg_TCT_chr17:8024243-8024330 (+)	GGCTCTGTGGCGCAATGGATAGCGCATTTGGACTTCTAGTGACGAATAG AGCAATTCAAAGGTTGTGGGTTCCGAATCCACCAAGATCG
45	Arg_TCT_chr9:131102355-131102445 (-)	GGCTCTGTGGCGCAATGGATAGCGCATTTGGACTTCTAGCTGAGCCTAG TGTGGTCAATCAAAGGTTGTGGGTTTCGAGTCCCAAGATCG

46	Arg_TCT_chr11:59318767-59318852 (+)	GGCTCTGTGGCGCAATGGATAGCGCATTTGGACTTCTAGATAGTTAGAG AAATTCAAAGGTTGTGGGTTTCGAGTCCACCCAGAGTCG
47	Arg_TCT_chr1:159111401-159111474 (-)	GTCTCTGTGGCGCAATGGACGAGCGGCTGGACTTCTAATCCAGAGGT TCCGGGTTTCGAGTCCCGCAGAGATG
48	Arg_TCT_chr6:27529963-27530049 (+)	GGCTCTGTGGCGCAATGGATAGCGCATTTGGACTTCTAGCCTAAATCAA GAGATTCAAAGGTTGCGGTTTCGAGTCCCTCCAGAGTCG
49	Asn_GTT_chr1:161510031-161510104 (+)	GTCTCTGTGGCGCAATCGGTTAGCGGTTTCGGCTGTTAACCCGAAAGGT TGGTGGTTTCGATCCCAACCCAGGGACG
50	Asn_GTT_chr1:143879832-143879905 (-)	GTCTCTGTGGCGCAATCGGCTAGCGGTTTGGCTGTTAACTAAAGGTT GGCGGTTTCGAACCCACCCAGAGGCG
51	Asn_GTT_chr1:144301611-144301684 (+)	GTCTCTGTGGCGCAATCGGTTAGCGGTTCCGCTGTTAACCCGAAAGCTT GGTGGTTTCGAGCCCAACCCAGGGATG
52	Asn_GTT_chr1:149326272-149326345 (-)	GTCTCTGTGGCGCAATCGGCTAGCGGTTTGGCTGTTAACTAAAGGTT GGTGGTTTCGAACACACCCAGAGGCG
53	Asn_GTT_chr1:148248115-148248188 (+)	GTCTCTGTGGCGCAATCGGTTAGCGGTTTCGGCTGTTAACCCGAAAGGT TGGTGGTTTCGAGCCCAACCCAGGGACG
54	Asn_GTT_chr1:148598314-148598387 (-)	GTCTCTGTGGCGCAATCGGTTAGCGCATTCGGCTGTTAACCCGAAAGGT TGGTGGTTTCGAGCCCAACCCAGGGACG
55	Asn_GTT_chr1:17216172-17216245 (+)	GTCTCTGTGGCGCAATCGGTTAGCGGTTTCGGCTGTTAACCCGAAAGAT TGGTGGTTTCGAGCCCAACCCAGGGACG
56	Asn_GTT_chr1:16847080-16847153 (-)	GTCTCTGTGGCGCAATCGGTTAGCGGTTTCGGCTGTTAACTGAAAGGTT GGTGGTTTCGAGCCCAACCCAGGGACG
57	Asn_GTT_chr1:149230570-149230643 (-)	GTCTCTGTGGCGCAATGGGTTAGCGGTTTCGGCTGTTAACCCGAAAGGT TGGTGGTTTCGAGCCCAACCCAGGGACG
58	Asn_GTT_chr1:148000805-148000878 (+)	GTCTCTGTGGCGTAGTCGGTTAGCGGTTTCGGCTGTTAACCCGAAAGTT GGTGGTTTCGAGCCCAACCCAGGAACG
59	Asn_GTT_chr1:149711798-149711871 (-)	GTCTCTGTGGCGCAATCGGTTAGCGGTTTGGCTGTTAACTAAAGGTT GGTGGTTTCGAACCCCAACCCAGAGGCG
60	Asn_GTT_chr1:145979034-145979107 (-)	GTCTCTGTGGCGCAATCGGTTAGCGGTTTCGGCTGTTAACTGAAAGGTT AGTGGTTTCGAGCCCAACCCGGGACG
61	Asp_GTC_chr12:98897281-98897352 (+)	TCCTCGTTAGTATAGTGGTTAGTATCCCCGCCCTGTCACGCGGGAGACCG GGTTCAATTCCCCGACGGGAG

62	Asp_GTC_chr1:161410615-161410686 (-)	TCCTCGTTAGTATAGTGGTGAAGTATCCCCCGCTGTACGCGGGAGAGACC GGGGTTCGATTCCCCGACGGGGAG
63	Asp_GTC_chr6:27551236-27551307 (-)	TCCTCGTTAGTATAGTGGTGAAGTGTCCCCCGTCTGTACGCGGGAGAGACC GGGGTTCGATTCCCCGACGGGGAG
64	Cys_GCA_chr7:149007281-149007352 (+)	GGGGGCATAGCTCAGTGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CTGGTTCAAATCCAGGTGCCCCCT
65	Cys_GCA_chr7:149074601-149074672 (-)	GGGGGTATAGCTCAGGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CTGGTTCAAATCCAGGTGCCCCCT
66	Cys_GCA_chr7:149112229-149112300 (-)	GGGGGTATAGCTTAGCGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCGGTTCAAATCCGGGTGCCCCCT
67	Cys_GCA_chr7:149344046-149344117 (-)	GGGGGTATAGCTTAGGGTAGAGCATTTGACTGCAGATCAAAAAGGTCC CTGGTTCAAATCCAGGTGCCCCCT
68	Cys_GCA_chr7:149052766-149052837 (-)	GGGGGTATAGCTCAGGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCAGTTCAAATCTGGGTGCCCCCT
69	Cys_GCA_chr17:37017937-37018008 (-)	GGGGGTATAGCTCAGGGTAGAGCATTTGACTGCAGATCAAGAAGTCC CCGGTTCAAATCCGGGTGCCCCCT
70	Cys_GCA_chr7:149281816-149281887 (+)	GGGGGTATAGCTCAGGGTAGAGCATTTGACTGCAGATCAAGAGGTCT CTGGTTCAAATCCAGGTGCCCCCT
71	Cys_GCA_chr7:149243631-149243702 (+)	GGGGGTATAGCTCAGGGTAGAGCATTTGACTGCAGATCAAGAAGTCC TTGGTTCAAATCCAGGTGCCCCCT
72	Cys_GCA_chr7:149388272-149388343 (-)	GGGGATATAGCTCAGGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCGGTTCAAATCCGGGTGCCCCCT
73	Cys_GCA_chr7:149072850-149072921 (-)	GGGGGTATAGTTCAAGGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CTGGTTCAAATCCAGGTGCCCCCT
74	Cys_GCA_chr7:149310156-149310227 (-)	GGGGGTATAGCTCAGGGTAGAGCATTTGACTGCAAAATCAAGAGGTCC CTGATTCAAATCCAGGTGCCCCCT
75	Cys_GCA_chr4:124430005-124430076 (-)	GGGGGTATAGCTCAGTGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCGGTTCAAATCCGGGTGCCCCCT
76	Cys_GCA_chr7:149295046-149295117 (+)	GGGGGTATAGCTCAGGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCAGTTCAAATCTGGGTGCCCCCT
77	Cys_GCA_chr7:149361915-149361986 (+)	GGGGGTATAGCTCACAGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCGGTTCAAATCTGGGTGCCCCCT

78	Cys_GCA_chr7:149253802-149253871 (+)	GGGCGTATAGCTCAGGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCAGTTCAAATCTGGGTGCCCA
79	Cys_GCA_chr7:149292305-149292376 (-)	GGGGGTATAGCTCACAGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCGGTTCAAATCCGGTTACTCCCT
80	Cys_GCA_chr7:149286164-149286235 (-)	GGGGGTATAGCTCAGGGTAGAGCACTTGACTGCAGATCAAGAGGTCC CTGGTTCAAATCCAGGTGCCCCCT
81	Cys_GCA_chr17:37025545-37025616 (-)	GGGGGTATAGCTCAGTGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CTGGTTCAAATCCGGGTGCCCCCT
82	Cys_GCA_chr15:80036997-80037069 (+)	GGGGGTATAGCTCAGTGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCCGGTTCAAATCCGGGTGCCCCCT
83	Cys_GCA_chr3:131947944-131948015 (-)	GGGGGTATAGCTCAGTGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CTGGTTCAAATCCAGGTGCCCCCT
84	Cys_GCA_chr1:93981834-93981906 (-)	GGGGGTATAGCTCAGTGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCCGGTTCAAATCCGGGTGCCCCCT
85	Cys_GCA_chr14:73429679-73429750 (+)	GGGGGTATAGCTCAGGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCGGTTCAAATCCGGGTGCCCCCT
86	Cys_GCA_chr3:131950642-131950713 (-)	GGGGGTATAGCTCAGGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CTGGTTCAAATCCAGGTGCCCCCT
87	Gln_CTG_chr6:18836402-18836473 (+)	GGTTCCATGGTGTAATGGTAGCAGCTCTGGACTCTGAATCCAGCGATCC GAGTTCAAATCTCGGTGGAACCT
88	Gln_CTG_chr6:27515531-27515602 (-)	GGTTCCATGGTGTAATGGTAGCAGCTCTGGACTCTGAATCCAGCGATCC GAGTTCAAAGTCTCGGTGGAACCT
89	Gln_CTG_chr1:145963304-145963375 (+)	GGTTCCATGGTGTAATGGTAGCAGCTCTGGACTCTGAATCCAGCGATC CGAGTTCCGAGTCTCGGTGGAACCT
90	Gln_CTG_chr1:147737382-147737453 (-)	GGTTCCATGGTGTAATGGTAGCAGCTCTGGACTCTGAATCCAGCGATC CGAGTTCCGAGTCTCGGTGGAACCT
91	Gln_CTG_chr6:27263212-27263283 (+)	GGTTCCATGGTGTAATGGTAGCAGCTCTGGACTCTGAATCCCGGTAATCC GAGTTCAAATCTCGGTGGAACCT
92	Gln_CTG_chr6:27759135-27759206 (-)	GGCCCCATGGTGTAATGGTAGCAGCTCTGGACTCTGAATCCAGCGATC CGAGTTCAAATCTCGGTGGAACCT
93	Gln_CTG_chr1:147800937-147801008 (+)	GGTTCCATGGTGTAATGGTAGCAGCTCTGGACTCTGAATCCAGCGATCT GAGTTCCGAGTCTCTGTGGAACCT

94	Gln_TTG_chr17:47269890-47269961 (+)	GGTCCCATGGTGTAAATGGTTAGCACTCTGGACTTTGAATCCAGCGATCC GAGTTCAAATCTCGTGGGACCT
95	Gln_TTG_chr6:28557156-28557227 (+)	GGTCCCATGGTGTAAATGGTTAGCACTCTGGACTTTGAATCCAGCAATCC GAGTTCGAATCTCGTGGGACCT
96	Gln_TTG_chr6:26311424-26311495 (-)	GGCCCCATGGTGTAAATGGTTAGCACTCTGGACTTTGAATCCAGCGATC CGAGTTCAAATCTCGTGGGACCT
97	Gln_TTG_chr6:145503859-145503930 (+)	GGTCCCATGGTGTAAATGGTTAGCACTCTGGGCTTTGAATCCAGCAAATCC GAGTTCGAATCTTGTGGGACCT
98	Glu_CTC_chr1:145399233-145399304 (-)	TCCCTGGTGGTCTAGTGGTTAGGATTCGGCGCTCTCACCCGCCGCGGCC GGTTTCGATTCCCGGTCAGGAA
99	Glu_CTC_chr1:249168447-249168518 (+)	TCCCTGGTGGTCTAGTGGTTAGGATTCGGCGCTCTCACCCGCCGCGGCC GGTTTCGATTCCCGGTCAGGAA
100	Glu_TTC_chr2:131094701-131094772 (-)	TCCCATATGGTCTAGCGGTTAGGATTCCTGGTTTTCACCCAGGTGGCCC GGTTTCGACTCCCGGTATGGAA
101	Glu_TTC_chr13:45492062-45492133 (-)	TCCACATGGTCTAGCGGTTAGGATTCCTGGTTTTCACCCAGCGGCC GGTTTCGACTCCCGGTGTGGAA
102	Glu_TTC_chr1:17199078-17199149 (+)	TCCCTGGTGGTCTAGTGGTAGGATTCGGCGCTTTCACCCGCCGCGGCC GGTTTCGATTCCCGGCCAGGAA
103	Glu_TTC_chr1:16861774-16861845 (-)	TCCCTGGTGGTCTAGTGGTAGGATTCGGCGCTTTCACCCGCCGCGGCC GGTTTCGATTCCCGGTCAGGAA
104	Gly_CCC_chr1:16872434-16872504 (-)	GCATTGGTGGTTCAGTGGTAGAATTCTCGCCTCCCACGCGGGAGACCC GGTTCAAATTCGCGCCAAATGCA
105	Gly_CCC_chr2:70476123-70476193 (-)	GCGCCGCTGGTGTAGTGGTATCATGCAAGATTCCCATTCTTGCGACCCG GGTTCGATTCCCGGGCGCGCA
106	Gly_CCC_chr17:19764175-19764245 (+)	GCATTGGTGGTTCAGTGGTAGAATTCTCGCCTCCCACGCGAGAGACCC AGGTTTCGATTCTCGGCCAATGCA
107	Gly_GCC_chr1:161413094-161413164 (+)	GCATGGTGGTTCAGTGGTAGAATTCTCGCCTGCCACGCGGGAGGCC GGTTTCGATTCCCGGCCCATGCA
108	Gly_GCC_chr1:161493637-161493707 (-)	GCATTGGTGGTTCAGTGGTAGAATTCTCGCCTGCCACGCGGGAGGCC GGTTTCGATTCCCGGCCAATGCA
109	Gly_GCC_chr16:70812114-70812184 (-)	GCATTGGTGGTTCAGTGGTAGAATTCTCGCCTGCCACGCGGGAGGCC GGTTTCGATTCCCGGCCAGTCA

110	Gly_GCC_chr1:161450356-161450426 (+)	GCATAGGTGGTTCAAGTGGTAGAATTCTTGCCCTGCCACGCAGGAGGCC AGGTTGATTCCTGGCCCATGCA
111	Gly_GCC_chr16:70822597-70822667 (+)	GCATTGGTGGTTCAAGTGGTAGAATTCTCGCCCTGCCATGCGGGCGGCCG GGCTTCGATTCTCTGGCCCAATGCA
112	Gly_TCC_chr19:4724082-4724153 (+)	GCGTTGGTGGTATAGTGGTAGCATAGCTGCCCTTCCAAGCAGTTGACC CGGTTTCGATTCCCGGCCAACGCA
113	Gly_TCC_chr1:145397864-145397935 (-)	GCGTTGGTGGTATAGTGGTAGCATAGCTGCCCTTCCAAGCAGTTGACC CGGTTTCGATTCCCGGCCAACGCA
114	Gly_TCC_chr17:8124866-8124937 (+)	GCGTTGGTGGTATAGTGGTAGCATAGCTGCCCTTCCAAGCAGTTGACC CGGTTTCGATTCCCGGCCAACGCA
115	Gly_TCC_chr1:161409961-161410032 (-)	GCGTTGGTGGTATAGTGGTAGCATAGCTGCCCTTCCAAGCAGTTGACC CGGCTCGATTCCCGGCCAACGCA
116	His_GTG_chr1:145396881-145396952 (-)	GCCGTGATCGTATAGTGGTAGTACTCTGCGTTGTGGCCGCAAGCAACCT CGGTTTCGAAATCCGAGTACGGCA
117	His_GTG_chr1:149155828-149155899 (-)	GCCATGATCGTATAGTGGTAGTACTCTGCGCTGTGGCCGCAAGCAACC TCGGTTTCGAATCCGAGTACGGCA
118	Ile_AAT_chr6:58149254-58149327 (+)	GGCCGGTTAGCTCAGTTGGTTAGAGCGTGGCGCTAATAACGCCCAAGGT CGCGGGTTCGATCCCCGTACGGGCCA
119	Ile_AAT_chr6:27655967-27656040 (+)	GGCCGGTTAGCTCAGTTGGTTAGAGCGTGGTGCTAATAACGCCCAAGGT CGCGGGTTCGATCCCCGTACTGGCCA
120	Ile_AAT_chr6:27242990-27243063 (-)	GGCTGGTTAGCTCAGTTGGTTAGAGCGTGGTGCTAATAACGCCCAAGGT CGCGGGTTCGATCCCCGTACTGGCCA
121	Ile_AAT_chr17:8130309-8130382 (-)	GGCCGGTTAGCTCAGTTGGTTAGAGCGTGGTGCTAATAACGCCCAAGGT CGCGGGTTCGAACCCCGTACGGGCCA
122	Ile_AAT_chr6:26554350-26554423 (+)	GGCCGGTTAGCTCAGTTGGTTAGAGCGTGGTGCTAATAACGCCCAAGGT CGCGGGTTCGATCCCCGTACGGGCCA
123	Ile_AAT_chr6:26745255-26745328 (-)	GGCCGGTTAGCTCAGTTGGTTAGAGCGTGGTGCTAATAACGCCCAAGGT CGCGGGTTCGATCCCCGTACTGGCCA
124	Ile_AAT_chr6:26721221-26721294 (-)	GGCCGGTTAGCTCAGTTGGTTAGAGCGTGGTGCTAATAACGCCCAAGGT CGCGGGTTCGATCCCCGTACGGGCCA
125	Ile_AAT_chr6:27636362-27636435 (+)	GGCCGGTTAGCTCAGTCCGGCTAGAGCGTGGTGCTAATAACGCCCAAGGT CGCGGGTTCGATCCCCGTACGGGCCA

126	Ile_AAT_chr6:27241739-27241812 (+)	GGCTGGTTAGTTCAGTTGGTTAGAGCGTGGTGCTAATAACGCCAAGGT CGTGGGTTTCGATCCCATATCGGCCA
127	Ile_GAT_chrX:3756418-3756491 (-)	GGCCGGTTAGCTCAGTTGGTAAGAGCGGTGGTGCTGATAACACCAAGGT CGCGGGCTCGACTCCCGCACCGGCCA
128	Ile_TAT_chr19:39902808-39902900 (-)	GCTCCAGTGGCGCAATCGGTTAGCGCGCGGTACTTATATGACAGTGCG AGCGAGCAATGCCGAGGTTGTGAGTTCGATCCTCACCTGGAGCA
129	Ile_TAT_chr2:43037676-43037768 (+)	GCTCCAGTGGCGCAATCGGTTAGCGCGCGGTACTTATACAGCAGTACA TGCAAGCAATGCCGAGGTTGTGAGTTCGAGCCTCACCTGGAGCA
130	Ile_TAT_chr6:26988125-26988218 (+)	GCTCCAGTGGCGCAATCGGTTAGCGCGCGGTACTTATATGCGAGTATG TGTGCGAGTGTATGCCGAGGTTGTGAGTTCGAGCCTCACCTGGAGCA
131	Ile_TAT_chr6:27599200-27599293 (+)	GCTCCAGTGGCGCAATCGGTTAGCGCGCGGTACTTATACAGCAGTATA TGTGCGGTTGATGCCGAGGTTGTGAGTTCGAGCCTCACCTGGAGCA
132	Ile_TAT_chr6:28505367-28505460 (+)	GCTCCAGTGGCGCAATCGGTTAGCGCGCGGTACTTATACAGCAGTATA CCTGTGAGCAATGCCGAGGTTGTGAGTTCGAGCCTCACCTGGAGCA
133	Leu_AAG_chr5:180524474-180524555 (-)	GGTAGCGTGGCCGAGCGGTCTAAGGCGCTGGATTAAAGGCTCCAGTCTC TTCGGAGGCGTGGGTTTCGAATCCCAACCGCTGCCA
134	Leu_AAG_chr5:180614701-180614782 (+)	GGTAGCGTGGCCGAGCGGTCTAAGGCGCTGGATTAAAGGCTCCAGTCTC TTCGGGGGCGTGGGTTTCGAATCCCAACCGCTGCCA
135	Leu_AAG_chr6:28956779-28956860 (+)	GGTAGCGTGGCCGAGCGGTCTAAGGCGCTGGATTAAAGGCTCCAGTCTC TTCGGGGGCGTGGGTTTCGAATCCCAACCGCTGCCA
136	Leu_AAG_chr6:28446400-28446481 (-)	GGTAGCGTGGCCGAGTGGTCTAAGACGCTGGATTAAAGGCTCCAGTCTC TTCGGGGGCGTGGGTTTGAATCCCAACCGCTGCCA
137	Leu_CAA_chr6:28864000-28864105 (-)	GTCAAGGATGGCCGAGTGGTCTAAGGCGCCAGACTCAAAGCTAAGCTTCC TCCGCGGTGGGGATTCTGGTCTCCAATGGAGGCGTGGGTTTCGAATCCC
138	Leu_CAA_chr6:28908830-28908934 (+)	GTCAAGGATGGCCGAGTGGTCTAAGGCGCCAGACTCAAAGCTTGGCTTCC TCGTGTTGAGGATTCTGGTCTCCAATGGAGGCGTGGGTTTCGAATCCA
139	Leu_CAA_chr6:27573417-27573524 (-)	GTCAAGGATGGCCGAGTGGTCTAAGGCGCCAGACTCAAAGCTTACTGCTT CCTGTGTTCCGGTCTTCTGGTCTCCGATGGAGGCGTGGGTTTCGAATCC
140	Leu_CAA_chr6:27570348-27570454 (-)	GTCAAGGATGGCCGAGTGGTCTAAGGCGCCAGACTCAAAGCTTACTTCC CCAGGTTTGGGGCTTCTGGTCTCCGCAATGGAGGCGTGGGTTTCGAATCC
141	Leu_CAA_chr1:249168054-249168159 (+)	GTCAAGGATGGCCGAGTGGTCTAAGGCGCCAGACTCAAAGGTAAGCACCT TGCCTGCGGGCTTCTGGTCTCCGGATGGAGGCGTGGGTTTCGAATCCC

142	Leu_CAA_chr11:9296790-9296863 (+)	GCCTCCTTAGTGCAGTAGGTAGCGCATCAGTCTCAAAATCTGAATGGT CCTGAGTTCAAGCCTCAGAGGGGCA
143	Leu_CAA_chr1:161581736-161581819 (-)	GTCAGGATGGCCGAGCAGTCTTAAGGCGTGGTTCAAATCGCACCCCT CCGCTGGAGGCGTGGTTCGAAATCCCACTTTTGACA
144	Leu_CAG_chr1:161411323-161411405 (+)	GTCAGGATGGCCGAGCGGTCTAAGGCGTGGTTCAGGTCGCAAGTCTC CCCTGGAGGCGTGGTTCGAAATCCCACTCCTGACA
145	Leu_CAG_chr16:57333863-57333945 (+)	GTCAGGATGGCCGAGCGGTCTAAGGCGTGGTTCAGGTCGCAAGTCTC CCCTGGAGGCGTGGTTCGAAATCCCACTTCTGACA
146	Leu_TAA_chr6:144537684-144537766 (+)	ACCAAGATGGCCGAGTGGTTAAGGCGTTGGACTTAAGATCCAATGGAC ATATGTCGCGTGGTTCGAAACCCCACTCCTGGTA
147	Leu_TAA_chr6:27688898-27688980 (-)	ACCGGATGGCCGAGTGGTTAAGGCGTTGGACTTAAGATCCAATGGGC TGGTGCCCGCTGGTTCGAAACCCCACTCCTGGTA
148	Leu_TAA_chr1:59319228-59319310 (+)	ACCAAGATGGCCGAGTGGTTAAGGCGTTGGACTTAAGATCCAATGGAT TCATATCCGCGTGGTTCGAAACCCCACTTCTGGTA
149	Leu_TAA_chr6:27198334-27198416 (-)	ACCGGATGGCTGAGTGGTTAAGGCGTTGGACTTAAGATCCAATGGAC AGGTGTCGCGTGGTTCGAGCCCCACTCCCGGTA
150	Leu_TAG_chr17:8023632-8023713 (-)	GGTAGCGTGGCCGAGCGGTCTAAGGCGTGGATTAGGCTCCAGTCTC TTCGGAGGCGTGGTTCGAAATCCCACTCCGCTGCCA
151	Leu_TAG_chr14:21093529-21093610 (+)	GGTAGTGTGGCCGAGCGGTCTAAGGCGTGGATTAGGCTCCAGTCTC TTCGGGGCGTGGTTCGAAATCCCACTCCGCTGCCA
152	Leu_TAG_chr16:22207032-22207113 (-)	GGTAGCGTGGCCGAGTGGTCTAAGGCGTGGATTAGGCTCCAGTCTC TTCGATGGCGTGGTTCGAAATCCCACTCCGCTGCCA
153	Lys_CTT_chr14:58706613-58706685 (-)	GCCCGGCTAGCTCAGTCGGTAGAGCATGGGACTCTTAATCCCAGGGTC GTGGGTTTCGAGCCCCACGTTGGGCG
154	Lys_CTT_chr19:36066750-36066822 (+)	GCCAGCTAGCTCAGTCGGTAGAGCATAGACTCTTAATCTCAGGGTT GTGGATTCTGTGCCCCATGCTGGGTG
155	Lys_CTT_chr19:52425393-52425466 (-)	GCAGCTAGCTCAGTCGGTAGAGCATGAGACTCTTAATCTCAGGGTCTC GGGTTCTGTGCCCCATGTTGGGTGCCA
156	Lys_CTT_chr1:145395522-145395594 (-)	GCCCGGCTAGCTCAGTCGGTAGAGCATGAGACTCTTAATCTCAGGGTC GTGGGTTTCGAGCCCCACGTTGGGCG
157	Lys_CTT_chr16:3207406-3207478 (-)	GCCCGGCTAGCTCAGTCGGTAGAGCATGAGACCTTAATCTCAGGGTC GTGGGTTTCGAGCCCCACGTTGGGCG

158	Lys_CTT_chr16:3241501-3241573 (+)	GCCCGGCTAGCTCAGTCGGTAGAGCATGGGACTCTTAATCTCAGGGTC GTGGGTTTCGAGCCCCACGTTGGGCG
159	Lys_CTT_chr16:3230555-3230627 (-)	GCCCGGCTAGCTCAGTCGATAGAGCATGAGACTCTTAATCTCAGGGTC GTGGGTTTCGAGCCGACGTTGGGCG
160	Lys_CTT_chr1:55423542-55423614 (-)	GCCCAGCTAGCTCAGTCGGTAGAGCATGAGACTCTTAATCTCAGGGTC ATGGGTTTGAGCCCCACGTTTGGTG
161	Lys_CTT_chr16:3214939-3215011 (+)	GCCTGGCTAGCTCAGTCGGCAAGCATGAGACTCTTAATCTCAGGGTC GTGGGCTCGAGTCCATGTTGGGCG
162	Lys_CTT_chr5:26198539-26198611 (-)	GCCCGACTACCTCAGTCGGTGGAGCATGGGACTCTTCATCCACAGGTT GTGGGTTTCGAGCCCCACATTGGGCA
163	Lys_TTT_chr16:73512216-73512288 (-)	GCCTGGATAGCTCAGTTGGTAGAGCATCAGACTTTTAATCTGAGGGTC CAGGGTTCAAGTCCCTGTTCAAGGCA
164	Lys_TTT_chr12:27843306-27843378 (+)	ACCCAGATAGCTCAGTCAGTAGAGCATCAGACTTTTAATCTGAGGGTC CAAGGTTCAATGTCCTTTTGGGTG
165	Lys_TTT_chr11:122430655-122430727 (+)	GCCTGGATAGCTCAGTTGGTAGAGCATCAGACTTTTAATCTGAGGGTC CAGGGTTCAAGTCCCTGTTCAAGGCG
166	Lys_TTT_chr1:204475655-204475727 (+)	GCCCGGATAGCTCAGTCGGTAGAGCATCAGACTTTTAATCTGAGGGTC CAGGGTTCAAGTCCCTGTTCCGGGCG
167	Lys_TTT_chr6:27559593-27559665 (-)	GCCTGGATAGCTCAGTCGGTAGAGCATCAGACTTTTAATCTGAGGGTC CAGGGTTCAAGTCCCTGTTCAAGGCG
168	Lys_TTT_chr11:59323902-59323974 (+)	GCCCGGATAGCTCAGTCGGTAGAGCATCAGACTTTTAATCTGAGGGTC CGGGGTTCAAGTCCCTGTTCCGGGCG
169	Lys_TTT_chr6:27302769-27302841 (-)	GCCTGGGTAAGCTCAGTCGGTAGAGCATCAGACTTTTAATCTGAGGGTC CAGGGTTCAAGTCCCTGTCCAGGCG
170	Lys_TTT_chr6:28715521-28715593 (+)	GCCTGGATAGCTCAGTTGGTAGAACATCAGACTTTTAATCTGACGGTG CAGGGTTCAAGTCCCTGTTCAAGGCG
171	Met_CAT_chr8:124169470-124169542 (-)	GCCTCGTTAGCGCAGTAGGTCGCGTCAGTCTCATAATCTGAAAGGTC GTGAGTTCGATCCTCACACGGGGCA
172	Met_CAT_chr16:71460396-71460468 (+)	GCCCTCTTAGCGCAGTGGGCAGCGCGTCAGTCTCATAATCTGAAAGGTC CTGAGTTCGAGCCTCAGAGAGGGCA
173	Met_CAT_chr6:28912352-28912424 (+)	GCCTCCTTAGCGCAGTAGGTCGCGTCAGTCTCATAATCTGAAAGGTC CTGAGTTCGAACCTCAGAGGGGGCA

174	Met_CAT_chr6:26735574-26735646 (-)	GCCCTCTTAGCGCAGCGGCGAGCGCTCAGTCTCATAATCTGAAAGGTC CTGAGTTCGAGCCTCAGAGAGGGCA
175	Met_CAT_chr6:26701712-26701784 (+)	GCCCTCTTAGCGCAGCTGGCAGCGCTCAGTCTCATAATCTGAAAGGTC CTGAGTTCAAGCCTCAGAGAGGGCA
176	Met_CAT_chr16:87417628-87417700 (-)	GCCTCGTTAGCGCAGTAGGCAGCGCTCAGTCTCATAATCTGAAAGGTC GTGAGTTCGAGCCTCACAGGGGCA
177	Met_CAT_chr6:58168492-58168564 (-)	GCCCTCTTAGTGCAGCTGGCAGCGCTCAGTTTTCATAATCTGAAAGTCC TGAGTTCAAAGCCTCAGAGAGGGCA
178	Phe_GAA_chr6:28758499-28758571 (-)	GCCGAAATAGCTCAGTTGGGAGAGCGTTAGACTGAAAGATCTAAAGGTC CCTGGTTCGATCCCGGGTTTCGGCA
179	Phe_GAA_chr11:59333853-59333925 (-)	GCCGAAATAGCTCAGTTGGGAGAGCGTTAGACTGAAAGATCTAAAGGTC CCTGGTTCAAATCCCGGGTTTCGGCA
180	Phe_GAA_chr6:28775610-28775682 (-)	GCCGAGATAGCTCAGTTGGGAGAGCGTTAGACTGAAAGATCTAAAGGTC CCTGGTTCAAATCCCGGGTTTCGGCA
181	Phe_GAA_chr6:28791093-28791166 (-)	GCCGAAATAGCTCAGTTGGGAGAGCGTTAGACCAGAAAGATCTTAAAGGT CCCTGGTTCAAATCCCGGGTTTCGGCA
182	Phe_GAA_chr6:28731374-28731447 (-)	GCTGAAATAGCTCAGTTGGGAGAGCGTTAGACTGAAAGATCTTAAAGTT CCCTGGTTCAAACCCCTGGGTTTCAGCC
183	Pro_AGG_chr16:3241989-3242060 (+)	GGCTCGTTGGTCTAGGGGTATGATTCGCTTAGGATGCGAGAGGTCC CGGGTTCAAATCCCGGACGAGCCC
184	Pro_AGG_chr1:167684725-167684796 (-)	GGCTCGTTGGTCTAGGGGTATGATTCGCTTAGGGTGCAGAGGTCC CGGGTTCAAATCCCGGACGAGCCC
185	Pro_CGG_chr1:167683962-167684033 (+)	GGCTCGTTGGTCTAGGGGTATGATTCGCTTCGGGTGCAGAGAGGTCC CGGGTTCAAATCCCGGACGAGCCC
186	Pro_CGG_chr6:27059521-27059592 (+)	GGCTCGTTGGTCTAGGGGTATGATTCGCTTCGGGTGTGAGAGGTCCC GGGTTCAAATCCCGGACGAGCCC
187	Pro_TGG_chr14:21101165-21101236 (+)	GGCTCGTTGGTCTAGGGGTATGATTCGCTTTGGGTGCGAGAGGTCCC GGGTTCAAATCCCGGACGAGCCC
188	Pro_TGG_chr11:75946869-75946940 (-)	GGCTCGTTGGTCTAGGGGTATGATTCGCTTTGGGTCCGAGAGGTCCC GGGTTCAAATCCCGGACGAGCCC
189	Pro_TGG_chr5:180615854-180615925 (-)	GGCTCGTTGGTCTAGGGGTATGATTCGCTTTGGGTGCGAGAGGTCCC GGGTTCAAATCCCGGACGAGCCC

190	SeC_TCA_chr19:45981859-45981945 (-)	GCCCGGATGATCCTCAGTGGTCTGGGGTGCAGGCTTCAAACCTGTAGC TGCTAGCGACAGAGTGGTTCAATTCCACCTTTCGGGCG
191	SeC_TCA_chr22:44546537-44546620 (+)	GCTCGGATGATCCTCAGTGGTCTGGGGTGCAGGCTTCAAACCTGTAGC TGCTAGTGACAGAGTGGTTCAATTCCACCTTTGTA
192	Ser_AGA_chr6:27509554-27509635 (-)	GTAGTCGTGGCCGAGTGGTTAAGGCGATGGACTAGAAATCCATTGGGG TTTCCCGCGCAGGTTTCGAATCCTGCCGACTACG
193	Ser_AGA_chr6:26327817-26327898 (+)	GTAGTCGTGGCCGAGTGGTTAAGGCGATGGACTAGAAATCCATTGGGG TCTCCCGCGCAGGTTTCGAATCCTGCCGACTACG
194	Ser_AGA_chr6:27499987-27500068 (+)	GTAGTCGTGGCCGAGTGGTTAAGGCGATGGACTAGAAATCCATTGGGG TTTCCCGCGCAGGTTTCGAATCCTGCCGACTACG
195	Ser_AGA_chr6:27521192-27521273 (-)	GTAGTCGTGGCCGAGTGGTTAAGTGTGACTAGAAACCCATTGGGG TCTCCCGCGCAGGTTTCGAATCCTGCCGACTACG
196	Ser_CGA_chr17:8042199-8042280 (-)	GCTGTGATGGCCGAGTGGTTAAGGCGTTGGACTCGAAATCCAAATGGGG TCTCCCGCGCAGGTTTCGAATCCTGCTCACAGCG
197	Ser_CGA_chr6:27177628-27177709 (+)	GCTGTGATGGCCGAGTGGTTAAGGCGTTGGACTCGAAATCCAAATGGGG TCTCCCGCGCAGGTTCAAATCCTGCTCACAGCG
198	Ser_CGA_chr6:27640229-27640310 (-)	GCTGTGATGGCCGAGTGGTTAAGTGTGACTCGAAATCCAAATGGGG GTTCCCGCGCAGGTTCAAATCCTGCTCACAGCG
199	Ser_CGA_chr12:56584148-56584229 (+)	GTCACGGTGGCCGAGTGGTTAAGGCGTTGGACTCGAAATCCAAATGGGG TTTCCCGCGCAGGTTTCGAATCCTGTTCTGTGACG
200	Ser_GCT_chr6:27065085-27065166 (+)	GACGAGGTGGCCGAGTGGTTAAGGCGATGGACTGCTAATCCATTGTGC TCTGCACGCGTGGGTTTCGAATCCACCCCTCGTCG
201	Ser_GCT_chr6:27265775-27265856 (+)	GACGAGGTGGCCGAGTGGTTAAGGCGATGGACTGCTAATCCATTGTGC TCTGCACGCGTGGGTTTCGAATCCACCCCTCGTCG
202	Ser_GCT_chr11:66115591-66115672 (+)	GACGAGGTGGCCGAGTGGTTAAGGCGATGGACTGCTAATCCATTGTGC TTTGCACGCGTGGGTTTCGAATCCCATCCTCGTCG
203	Ser_GCT_chr6:28565117-28565198 (-)	GACGAGGTGGCCGAGTGGTTAAGGCGATGGACTGCTAATCCATTGTGC TCTGCACGCGTGGGTTTCGAATCCCATCCTCGTCG
204	Ser_GCT_chr6:28180815-28180896 (+)	GACGAGGTGGCCGAGTGGTTAAGGCGATGGACTGCTAATCCATTGTGC TCTGCACACGTGGGTTTCGAATCCCATCCTCGTCG
205	Ser_GCT_chr6:26305718-26305801 (-)	GGAGAGGCCCTGGCCGAGTGGTTAAGGCGATGGACTGCTAATCCATTGT GCTCTGCACGCGTGGGTTTCGAATCCCATCCTCGTCG

206	Ser_TGA_chr10:69524261-69524342 (+)	GCAGCGATGGCCGAGTGTTAAGGCGTTGGACTTGAAATCCAATGGGG TCTCCCCGCGCAGGTTTCGAACCCCTGCTCGCTGCG
207	Ser_TGA_chr6:27513468-27513549 (+)	GTAGTCGTGGCCGAGTGGTTAAGGCGATGGACTTGAAATCCATTGGGG TTTCCCCGCGCAGGTTTCGAATCCTGCCGACTACG
208	Ser_TGA_chr6:26312824-26312905 (-)	GTAGTCGTGGCCGAGTGGTTAAGGCGATGGACTTGAAATCCATTGGGG TCTCCCCGCGCAGGTTTCGAATCCTGCCGACTACG
209	Ser_TGA_chr6:27473607-27473688 (-)	GTAGTCGTGGCCGAGTGGTTAAGGCGATGGACTTGAAATCCATTGGGG TTTCCCCGCGCAGGTTTCGAATCCTGTGCGCTACG
210	Thr_AGT_chr17:8090478-8090551 (+)	GGCGCCGTGGCTTAGTTGGTTAAAGCGCCTGTCTAGTAAACAGGAGAT CCTGGGTTTCGAATCCACGCGGTGCCT
211	Thr_AGT_chr6:26533145-26533218 (-)	GGTCCGTGGCTTAGCTGGTTAAAGCGCCTGTCTAGTAAACAGGAGAT CCTGGGTTTCGAATCCACGCGGGGCCT
212	Thr_AGT_chr6:28693795-28693868 (+)	GGTCCGTAGCTTAGTTGGTTAAAGCGCCTGTCTAGTAAACAGGAGAT CCTGGGTTTCGACTCCACGCGGGGCCT
213	Thr_AGT_chr6:27694473-27694546 (+)	GGCTTCGTGGCTTAGCTGGTTAAAGCGCCTGTCTAGTAAACAGGAGAT CCTGGGTTTCGAATCCACGCGAGGCCT
214	Thr_AGT_chr17:8042770-8042843 (-)	GGCGCCGTGGCTTAGCTGGTTAAAGCGCCTGTCTAGTAAACAGGAGAT CCTGGGTTTCGAATCCACGCGGTGCCT
215	Thr_AGT_chr6:27130050-27130123 (+)	GGCCCTGTGGCTTAGCTGGTCAAGCGCCTGTCTAGTAAACAGGAGAT CCTGGGTTTCGAATCCACGCGGGGCCT
216	Thr_CGT_chr6:28456770-28456843 (-)	GGCTCTATGGCTTAGTTGGTTAAAGCGCCTGTCTCGTAAACAGGAGAT CCTGGGTTTCGACTCCACGCGGGGCCT
217	Thr_CGT_chr16:14379750-14379821 (+)	GGCGCGGTGGCCAAAGTGTAAAGCGCGTCGGTCTCGTAAACCGAAGATCA CGGGTTCGAACCCCGTCCGTGCCT
218	Thr_CGT_chr6:28615984-28616057 (-)	GGCTCTGTGGCTTAGTTGGCTAAAGCGCCTGTCTCGTAAACAGGAGAT CCTGGGTTTCGAATCCACGCGGGGCCT
219	Thr_CGT_chr17:29877093-29877164 (+)	GGCGCGGTGGCCAAAGTGTAAAGCGCGTCGGTCTCGTAAACCGAAGATCG CGGGTTCGAACCCCGTCCGTGCCT
220	Thr_CGT_chr6:27586135-27586208 (+)	GGCCCTGTAGCTACGCGGTGGAGCGCTGGTCTCGTAAACCTAGGGGT CGTAGGTTCAAAATCTCACCAGGGCCT
221	Thr_TGT_chr6:28442329-28442402 (-)	GGCTCTATGGCTTAGTTGGTTAAAGCGCCTGTCTTGTAAACAGGAGAT CCTGGGTTTCGAATCCACGAGTAGAGCCT

222	Thr_TGT_chr1:222638347-222638419 (+)	GGCTCCATAGCTCAGTGGTTAGAGCACTGGTCTTGTAAACACAGGGGTC GCGAGTTCGATCCTCGCTGGGGCCT
223	Thr_TGT_chr14:21081949-21082021 (-)	GGCTCCATAGCTCAGGGGTTAGAGCGCTGGTCTTGTAAACACAGGGGTC GCGAGTTC AATTCTCGCTGGGGCCT
224	Thr_TGT_chr14:21099319-21099391 (-)	GGCTCCATAGCTCAGGGGTTAGAGCACTGGTCTTGTAAACACAGGGGTC GCGAGTTC AATCTCGCTGGGGCCT
225	Thr_TGT_chr14:21149849-21149921 (+)	GGCCCTATAGCTCAGGGGTTAGAGCACTGGTCTTGTAAACACAGGGGTC GCGAGTTC AATCTCGCTGGGGCCT
226	Thr_TGT_chr5:180618687-180618758 (-)	GGCTCCATAGCTCAGGGGTTAGAGCACTGGTCTTGTAAACACAGGGTCTG CGAGTTC AATCTCGCTGGGGCCT
227	Trp_CCA_chr17:8124187-8124258 (-)	GGCTCGTGGCGCAACGGTAGCGGCTCTGACTCCAGATCAGAAAGGTTG CGTGTTC A AATCACGTCGGGGTCA
228	Trp_CCA_chr17:19411494-19411565 (+)	GACCTCGTGGCGCAATGGTAGCGGCTCTGACTCCAGATCAGAAAGGTTG CGTGTTC A AGTCACGTCGGGGTCA
229	Trp_CCA_chr6:26319330-26319401 (-)	GACCTCGTGGCGCAACGGTAGCGGCTCTGACTCCAGATCAGAAAGGTTG CGTGTTC A AATCACGTCGGGGTCA
230	Trp_CCA_chr12:98898030-98898101 (+)	GACCTCGTGGCGCAACGGTAGCGGCTCTGACTCCAGATCAGAAAGGCTG CGTGTTC G AATCACGTCGGGGTCA
231	Trp_CCA_chr7:99067307-99067378 (+)	GACCTCGTGGCGCAACGGCAGCGGCTCTGACTCCAGATCAGAAAGGTTG CGTGTTC A AATCACGTCGGGGTCA
232	Tyr_ATA_chr2:219110549-219110641 (+)	CCTTCAATAGTTCAGCTGGTAGAGCAGAGGACTATAGCTACTTCCCTCA GTAGGAGACGTCCTTAGGTTGCTGGTTCGATTCCAGCTTGAAGGA
233	Tyr_GTA_chr6:26569086-26569176 (+)	CCTTCGATAGCTCAGTTGGTAGAGCGGAGGACTGTAGTTGGCTGTGTC CTTAGACATCCTTAGGTCGCTGGTTCGAAATCCGGCTCGAAGGA
234	Tyr_GTA_chr2:27273650-27273738 (+)	CCTTCGATAGCTCAGTTGGTAGAGCGGAGGACTGTAGTGATAGGGCG TGGCAATCCTTAGGTCGCTGGTTCGATTCCGGCTCGAAGGA
235	Tyr_GTA_chr6:26577332-26577420 (+)	CCTTCGATAGCTCAGTTGGTAGAGCGGAGGACTGTAGGCTCATTAAGC AAGGTATCCTTAGGTCGCTGGTTCGAAATCCGGCTCGGAGGA
236	Tyr_GTA_chr14:21125623-21125716 (-)	CCTTCGATAGCTCAGCTGGTAGAGCGGAGGACTGTAGATTGTATAGAC ATTTCGGGACATCCTTAGGTCGCTGGTTCGATTCCAGCTCGAAGGA
237	Tyr_GTA_chr8:67025602-67025694 (+)	CCTTCGATAGCTCAGCTGGTAGAGCGGAGGACTGTAGCTACTTCCCTCA GCAGGAGACATCCTTAGGTCGCTGGTTCGATTCCGGCTCGAAGGA

238	Tyr_GTA_chr8:67026223-67026311 (+)	CCTTCGATAGCTAGCTGGTAGAGCGGAGGACTGTAGGCGCGGCCCG TGGCCATCCTTAGGTCGCTGGTTCGATTCCGGCTCGAAGGA
239	Tyr_GTA_chr14:21121258-21121351 (-)	CCTTCGATAGCTAGCTGGTAGAGCGGAGGACTGTAGCCTGTAGAAAC ATTTGTGGACATCCTTAGGTCGCTGGTTCGATTCCGGCTCGAAGGA
240	Tyr_GTA_chr14:21131351-21131444 (-)	CCTTCGATAGCTAGCTGGTAGAGCGGAGGACTGTAGATTGTACAGAC ATTTGCGGACATCCTTAGGTCGCTGGTTCGATTCCGGCTCGAAGGA
241	Tyr_GTA_chr14:21151432-21151520 (+)	CCTTCGATAGCTAGCTGGTAGAGCGGAGGACTGTAGTACTTAAATGTG TGGTCATCCTTAGGTCGCTGGTTCGATTCCGGCTCGAAGGA
242	Tyr_GTA_chr6:26595102-26595190 (+)	CCTTCGATAGCTAGCTGGTAGAGCGGAGGACTGTAGGGGTTTGAATG TGGTCATCCTTAGGTCGCTGGTTCGAAATCCGGCTCGGAGGA
243	Tyr_GTA_chr14:21128117-21128210 (-)	CCTTCGATAGCTAGCTGGTAGAGCGGAGGACTGTAGACTGCGGAAAC GTTTGTGGACATCCTTAGGTCGCTGGTTCGAAATCCGGCTCGAAGGA
244	Tyr_GTA_chr6:26575798-26575887 (+)	CTTTCGATAGCTAGCTGGTAGAGCGGAGGACTGTAGGTTCATTAAAC TAAGGCATCCTTAGGTCGCTGGTTCGAAATCCGGCTCGAAGGA
245	Tyr_GTA_chr8:66609532-66609619 (-)	TCTTCAATAGCTAGCTGGTAGAGCGGAGGACTGTAGTGCACGCCCG TGGCCATCCTTAGGTCGCTGGTTCGATTCCGACTTGGAGAG
246	Val_AAC_chr3:169490018-169490090 (+)	GTTTCCGATAGTGTGGTATCACGTTTCGCTAACACGCGAAAGGTCC CCGGTTCGAAACCGGGCGGAAACA
247	Val_AAC_chr5:180615416-180615488 (-)	GTTTCCGATAGTGTGGTATCACGTTTCGCTAACACGCGAAAGGTCC CCCGTTTCGAAACCGGGCGGAAACA
248	Val_AAC_chr6:27618707-27618779 (-)	GTTTCCGATAGTGTGGTATCACGTTTCGCTAACACGCGAAAGGTCC CTGGATCAAAACCAAGCGGAAACA
249	Val_AAC_chr6:27648885-27648957 (-)	GTTTCCGATAGTGTGGTATCACGTTTCGCTAACACGCGAAAGGTCC GCGGTTTCGAAACCGGGCGGAAACA
250	Val_AAC_chr6:27203288-27203360 (+)	GTTTCCGATAGTGTGGTATCACGTTTTCGCTAACACGCGAAAGGTCC CCGGTTCGAAACCGGGCGGAAACA
251	Val_AAC_chr6:28703206-28703277 (-)	GGGGTGTAGCTCAGTGTAGAGCGGTATGCTTAAACATTCATGAGGCTC TGGGTTTCGATCCCCAGCACTTCCA
252	Val_CAC_chr1:161369490-161369562 (-)	GTTTCCGATAGTGTGGTATCACGTTTCGCTCACACGCGAAAGGTCC CCGGTTCGAAACCGGGCGGAAACA
253	Val_CAC_chr6:27248049-27248121 (-)	GCTTCTGTAGTGTGGTATCACGTTTCGCTCACACGCGAAAGGTCC CCGGTTCGAAACCGGGCGGAAAGCA

254	Val_CAC_chrl9:4724647-4724719 (-)	GTTTCCGTAGTGTAGCGGTTATCACATTTCGCCTCACACGCGAAAGGTCC CCGGTTCGATCCCGGCGGAAACA
255	Val_CAC_chrl1:149298555-149298627 (-)	GTTTCCGTAGTGTAGTGGTTATCACGTTTCGCCTCACACGCGAAAGGTCC CCGGTTCGAAACTGGGCGGAAACA
256	Val_CAC_chrl1:149684088-149684161 (-)	GTTTCCGTAGTGTAGTGGTTATCACGTTTCGCCTCACACGCGTAAAGGTC CCCGGTTCGAAACCGGGCGGAAACA
257	Val_CAC_chrl6:27173867-27173939 (-)	GTTTCCGTAGTGGAGTGGTTATCACGTTTCGCCTCACACGCGAAAGGTC CCCGGTTTGAAACCAAGCGGAAACA
258	Val_TAC_chrl1:59318102-59318174 (-)	GGTTCCATAGTGTAGTGGTTATCACGTTCTGCTTTACACGCAGAAAGGTCC TGGGTTTCGAGCCCCAGTGGAACCA
259	Val_TAC_chrl1:59318460-59318532 (-)	GGTTCCATAGTGTAGCGGTTATCACGTTCTGCTTTACACGCAGAAAGGTCC TGGGTTTCGAGCCCCAGTGGAACCA
260	Val_TAC_chrl10:5895674-5895746 (-)	GGTTCCATAGTGTAGTGGTTATCACATCTGCTTTACACGCAGAAAGGTCC TGGGTTCAAAGCCCCAGTGGAACCA
261	Val_TAC_chrl6:27258405-27258477 (+)	GTTTCCGTGGTGTAGTGGTTATCACATTTCGCCTTACACGCGAAAGGTCC TCGGGTCGAAACCGAGCGGAAACA
262	iMet_CAT_chrl1:153643726-153643797 (+)	AGCAGAGTGGCGCAGCGGAAAGCGTGTGGGCCCATAAACCCAGAGGTC GATGGATCGAAACCATCCTCTGCTA
263	iMet_CAT_chrl6:27745664-27745735 (+)	AGCAGAGTGGCGCAGCGGAAAGCGTGTGGGCCCATAAACCCAGAGGTC GATGGATCTAAACCATCCTCTGCTA
264	Glu_TTC_chrl1:16861773-16861845 (-)	TCCCTGGTGGTCTAGTGGCTAGGATTTCGGCGCTTTCACCGCCGCGGCC GGGTTTCGATTCCCGGTCAGGGAAT
265	Gly_CCC_chrl1:17004765-17004836 (-)	GCGTTGGTGGTTTAGTGGTAGAATTTCGCCTCCCATGCGGGAGACCC GGGTTCAATTCCCGGCCACTGCAC
266	Gly_CCC_chrl1:17053779-17053850 (+)	GGCCTTGGTGGTGCAGTGGTAGAATTTCGCCTCCCATGCGGGAGACC CGGGTTCAATTCCCGGCCAATGCA
267	Glu_TTC_chrl1:17199077-17199149 (+)	GTCCCTGGTGGTCTAGTGGCTAGGATTTCGGCGCTTTCACCGCCGCGGCC CGGGTTCGATTCCCGGCCAGGGA
268	Asn_GTT_chrl1:17216171-17216245 (+)	TGTCTCTGTGGCGCAATCGGTTAGCGGTTTCGGCTGTTAACCAGAAAGA TTGGTGGTTCGAGCCACCCAGGACG
269	Arg_TCT_chrl1:94313128-94313213 (+)	TGGCTCCGTGGCGCAATGGATAGCGCATTTGGACTTCTAGAGGCTGAAG GCATTCAAAGGTTCCGGGTTTCGAGTCCCGCGGAGTCG

270	Lys_CTT_chrl:145395521-145395594 (-)	GCCCGGCTAGCTCAGTCGGTAGAGCATGAGACTCTTAATCTCAGGGTC GTGGGTTTCGAGCCCCACGTTGGGCGC
271	His_GTG_chrl:145396880-145396952 (-)	GCCGTGATCGTATAGTGGTACTCTGCGTTGTGGCCGACGAAACCT CGGTTTCAAATCCGAGTCACGGCAG
272	Gly_TCC_chrl:145397863-145397935 (-)	GCGTTGGTGGTATAGTGGTGAGCATAGCTGCCTTCCAAGCAGTTGACC CGGGTTCGATTCCCGGCCAACGCAG
273	Glu_CTC_chrl:145399232-145399304 (-)	TCCCTGGTGGTCTAGTGGTTAGGATTGGGCGCTCTCACCGCCGCGGGCCC GGGTTTCGATTCCCGGTCAGGAAA
274	Gln_CTG_chrl:145963303-145963375 (+)	AGGTTCCATGGTGTAAATGGTGAGCACTCTGGACTCTGAATCCAGCGAT CCGAGTTCGAGTCTCGGTGGAACCT
275	Asn_GTT_chrl:148000804-148000878 (+)	TGTCCTGTGGCGTAGTCGGTTAGCGCGTTCGGCTGTTAACCCGAAAAAGT TGGTGGTTCGAGCCCCACCCAGGAACG
276	Asn_GTT_chrl:148248114-148248188 (+)	TGTCCTGTGGCGCAATCGGTTAGCGCGTTCGGCTGTTAACCCGAAAAAGG TTGGTGGTTCGAGCCCCACCCAGGGACG
277	Asn_GTT_chrl:148598313-148598387 (-)	GTCTCTGTGGCGCAATCGGTTAGCGCATTCGGCTGTTAACCCGAAAAAGGT TGGTGGTTCGAGCCCCACCCAGGGACGC
278	Asn_GTT_chrl:149230569-149230643 (-)	GTCTCTGTGGCGCAATGGGTTAGCGCGTTCGGCTGTTAACCCGAAAAAGGT TGGTGGTTCGAGCCCCATCCAGGGACGC
279	Val_CAC_chrl:149294665-149294736 (-)	GCACCTGGTGGTTCAGTGGTAGAATTCCTCGCCTCACACGCGGGGACACCC GGGTTCAAATCCCGGTCAAAGCAA
280	Val_CAC_chrl:149298554-149298627 (-)	GTTTCCGTAGTGTAGTGGTTATCACGTTTCGCCCTCACACGCGAAAGGTCC CCGGTTCGAAACTGGCGGGAACAG
281	Gly_CCC_chrl:149680209-149680280 (-)	GCACTGGTGGTTCAGTGGTAGAATTCCTCGCCTCCACGCGGGGAGACCC GGGTTTAATTCCCGGTCAAGATAA
282	Val_CAC_chrl:149684087-149684161 (-)	GTTTCCGTAGTGTAGTGGTTATCACGTTTCGCCCTCACACGCGTAAAGGTC CCCGGTTTCGAAACCGGGCGGAAACAT
283	Met_CAT_chrl:153643725-153643797 (+)	TAGCAGAGTGGCGCAGCGGAAGCGTGTGGGCCCATAACCCAGAGGT CGATGGATCGAAACCATCCTCTGCTA
284	Val_CAC_chrl:161369489-161369562 (-)	GTTTCCGTAGTGTAGTGGTTATCACGTTTCGCCCTCACACGCGAAAGGTCC CCGGTTCGAAACCGGGCGGAAACAA
285	Asp_GTC_chrl:161410614-161410686 (-)	TCCTCGTTAGTATAGTGGTAGTATCCCCGCCCTGTACACGCGGGAGACC GGGTTTCGATTCCCGGACGGGGAGG

286	Gly_GCC_chr1:161413093-161413164 (+)	TGCATGGGTGGTTCAAGTGGTAGAAATTCTCGCCTGCCACGCGGGAGGCC CGGGTTCGATTCCCGGCCCATGCA
287	Glu_CTC_chr1:161417017-161417089 (-)	TCCCTGGTGGTCTAGTGGTAGGATTGGGCGCTCTCACCGCCGCGGCC GGGTTTCGATTCCCGGTCAGGGAAG
288	Asp_GTC_chr1:161492934-161493006 (+)	ATCCTTGTACTATAGTGGTAGTATCTCTGCCTGTCTCATGCGTGAGAGA GGGGTCGATTCCCGACGGGGAG
289	Gly_GCC_chr1:161493636-161493707 (-)	GCAATTGGTGGTTTCAGTGGTAGAAATTCTCGCCTGCCACGCGGGAGGCC GGGTTTCGATTCCCGGCCAATGCAC
290	Leu_CAG_chr1:161500131-161500214 (-)	GTCAGGATGGCCGAGCGGTCTAAGGCGCTGCGTTCAGTTCGCAAGTCTC CCCTGGAGGCGTGGGTTCTGAATCCCACTCCTGACAA
291	Gly_TCC_chr1:161500902-161500974 (+)	CGCGTTGGTGGTATAGTGGTAGCATAGCTGCCTTCCCAAGCAGTTGAC CCGGGTTTCGATTCCCGGCCAACGCA
292	Asn_GTT_chr1:161510030-161510104 (+)	CGTCTCTGTGGCGCAATCGGTTAGCGGTTCTGGCTGTTAACCGAAAGG TTGGTGGTTCGATCCCAACCGGACG
293	Glu_TTC_chr1:161582507-161582579 (+)	CGCGTTGGTGGTGTAGTGGTAGACACAGCTGCCTTTCAAGCAGTTAAC GCGGGTTCGATTCCCGGGTAACGAA
294	Pro_CGG_chr1:167683961-167684033 (+)	CGGCTCGTTGGTCTAGGGGTATGATTCTCGCTTCGGGTGCGAGAGGTC CCGGGTTCAAATCCCGGACGAGCCC
295	Pro_AGG_chr1:167684724-167684796 (-)	GGCTCGTTGGTCTAGGGGTATGATTCTCGCTTAGGGTGCGAGAGGTCC CGGGTTCAAATCCCGGACGAGCCCT
296	Lys_TTT_chr1:204475654-204475727 (+)	CGCCCGGATAGCTCAGTCGGTAGAGCATCAGACTTTTAACTCTGAGGGT CCAGGGTTCAAGTCCCTGTTCTGGGCG
297	Lys_TTT_chr1:204476157-204476230 (-)	GCCCGGATAGCTCAGTCGGTAGAGCATCAGACTTTTAACTCTGAGGGTC CAGGGTTCAAAGTCCCTGTTCTGGGCGT
298	Leu_CAA_chr1:249168053-249168159 (+)	TGTCAGGATGGCCGAGTGGTCTAAGGCGCCAGACTCAAGGTAAGCACCC TTGCCCTCGGGGCTTCTGGTCTCCGGATGGAGCGGTGGGTTTGAATCCC
299	Glu_CTC_chr1:249168446-249168518 (+)	TTCCCTGGTGGTCTAGTGGTAGGATTCCGGCGCTCTCACCCCGCGGCC CGGGTTCGATTCCCGGTCAGGAAA
300	Tyr_GTA_chr2:27273649-27273738 (+)	GCCTTCGATAGCTCAGTTGGTAGAGCGGAGGACTGTAGTGGATAGGGC GTGGCAATCCTTAGGTCGCTGGTTCGATTCCGGCTCGAAGGA
301	Ala_AGC_chr2:27274081-27274154 (+)	CGGGGGATTAGCTCAAAATGGTAGAGCGCTCGCTTAGCATGCGAGAGGT AGCGGGATCGATGCCCCGCATCCTCCA

302	Ile_TAT_chr2:43037675-43037768 (+)	AGCTCCAGTGGCGCAATCGGTTAGCGCGCGGTACTTATACAGCAGTAC ATGCAGAGCAATGCCGAGGTTGTGAGTTCGAGCCTCACCTGGAGCA
303	Gly_CCC_chr2:70476122-70476193 (-)	GCGCCGCTGGTGTAGTGGTATCATGCAAGATTCCCATTTCTTGGACCCG GGTTCGATTCCCGGGCGGCGCAT
304	Glu_TTC_chr2:131094700-131094772 (-)	TCCCATATGGTCTAGCGGTAGGATTCTCTGGTTTTCACCCAGGTGGCCC GGTTCGACTCCCGGTATGGGAAAC
305	Ala_CGC_chr2:157257280-157257352 (+)	GGGGATGTAGCTCAGTGTAGAGCGCGCTTCGCAATGTGTGAGGTC CCGGGTTCAATCCCCGGCATCTCCA
306	Gly_GCC_chr2:157257658-157257729 (-)	GCATTGGTGGTTCAGTGGTAGAATTCTCGCCTGCCACGCGGAGGCC GGTTCGATTCCCGGCCAATGCAA
307	Arg_ACG_chr3:45730490-45730563 (-)	GGCCAGTGGCGCAATGGATAACCGCTCTGACTACGGATCAGAAAGATT CTAGGTTCCGACTCCTGGCTGGCTCGC
308	Val_AAC_chr3:169490017-169490090 (+)	GGTTCCGTAGTGTAGTGGTTATCACGTTTCGCCCTAACACGCGAAAGGT CCCCGGTTCGAAACCGGGCGGAAACA
309	Val_AAC_chr5:180596609-180596682 (+)	AGTTCCGTAGTGTAGTGGTTATCACGTTTCGCCCTAACACGCGAAAGGT CCCCGGTTCGAAACCGGGCGGAAACA
310	Leu_AAG_chr5:180614700-180614782 (+)	AGGTAGCGTGGCGAGCGGTCTAAGCGCTGGATTAAGGCTCCAGTCT CTTCGGGGCGGTGGGTTCCGAATCCACCGCTGCCA
311	Val_AAC_chr5:180615415-180615488 (-)	GTTCCTAGTGTAGTGGTATCATCAGTTTCGCCCTAACACGCGAAAGGTC CCCCGGTTCGAAACCGGGCGGAAACA
312	Pro_TGG_chr5:180615853-180615925 (-)	GGCTCGTTGGTCTAGGGGTATGATTCTCGCTTTGGGTGCGAGAGGTCCC GGGTTCAAATCCCGGACGAGCCCA
313	Thr_TGT_chr5:180618686-180618758 (-)	GGCTCCATAGCTCAGGGGTTAGAGCACTGGTCTTGTAAACAGGGTCTG CGAGTTCAAATCTCGCTGGGGCTG
314	Ala_TGC_chr5:180633867-180633939 (+)	TGGGGATGTAGCTCAGTGGTAGAGCGCATGCTTTGCATGTATGAGGCC CCGGGTTTCGATCCCCGGCATCTCCA
315	Lys_CTT_chr5:180634754-180634827 (+)	CGCCCGGTAGCTCAGTCGCGTAGAGCATGAGACTCTTAATCTCAGGGT CGTGGGTTTCGAGCCCCACGTTGGGCG
316	Val_AAC_chr5:180645269-180645342 (-)	GTTCCTAGTGTAGTGGTTATCACGTTTCGCCCTAACACGCGAAAGGTCC CCGGTTCGAAACCGGGCGGAAACA
317	Lys_CTT_chr5:180648978-180649051 (-)	GCCCGGCTAGCTCAGTCGCGTAGAGCATGAGACTCTTAATCTCAGGGTC GTGGGTTTCGAGCCCCACGTTGGGCGT

318	Val_CAC_chr5:180649394-180649467 (-)	GTTTCCGTAGTGTAGTGGTTATCACGTTTCGCCTCACACGCGAAAGGTCC CCGTTTCGAAACCGGGCGGAAACAC
319	Met_CAT_chr6:26286753-26286825 (+)	CAGCAGAGTGGCGCAGCGGAAGCGTGTCTGGGCCCATAAACCCAGAGGT CGATGGATCGAAACCATCCTCTGCTA
320	Ser_GCT_chr6:26305717-26305801 (-)	GGAGAGCCTGGCCGAGTGGTTAAGCGCATGGACTGCTAATCCATTGT GCTCTGCACGCGTGGGTTCGAATCCCATCCTCGTCGC
321	Gln_TTG_chr6:26311423-26311495 (-)	GGCCCCATGGTGTAAATGGTTAGCACTCTGGACTTTGAATCCAGCGATC CGAGTTCAAATCTCGGTGGGACCTG
322	Gln_TTG_chr6:26311974-26312046 (-)	GGCCCCATGGTGTAAATGGTTAGCACTCTGGACTTTGAATCCAGCGATC CGAGTTCAAATCTCGGTGGGACCTA
323	Ser_TGA_chr6:26312823-26312905 (-)	GTAGTCGTGGCCGAGTGGTTAAGCGCATGGACTTGAAATCCATTGGGG TCTCCCCGCGCAGGTTCGAATCCTGCCGACTACGG
324	Met_CAT_chr6:26313351-26313423 (-)	AGCAGAGTGGCGCAGCGGAAGCGTGTGGGCCCATAAACCCAGAGGTC GATGGATCGAAACCATCCTCTGCTAT
325	Arg_TCG_chr6:26323045-26323118 (+)	GGACCACGTGGCCTAATGGATAAGGCGTCTGACTTCGGATCAGAAGAT TGAGGGTTCGAATCCCTCCGTGGTTA
326	Ser_AGA_chr6:26327816-26327898 (+)	TGTAGTCGTGGCCGAGTGGTTAAGCGCATGGACTAGAAATCCATTGGG GTCTCCCCGCGCAGGTTCGAATCCTGCCGACTACG
327	Met_CAT_chr6:26330528-26330600 (-)	AGCAGAGTGGCGCAGCGGAAGCGTGTGGGCCCATAAACCCAGAGGTC GATGGATCGAAACCATCCTCTGCTAG
328	Leu_CAG_chr6:26521435-26521518 (+)	CGTCAGGATGGCCGAGCGGTCTAAGGCGTGCCTTCAGGTCGCAGTCT CCCCTGGAGCGTGGGTTCGAATCCCACTCCTGACA
329	Thr_AGT_chr6:26533144-26533218 (-)	GGCTCCGTGGCTTAGCTGGTTAAAGCGCCTGTCTAGTAAACAGGAGAT CCTGGGTTCGAATCCAGCGGGGCTG
330	Arg_ACG_chr6:26537725-26537798 (+)	AGGGCCAGTGGCGCAATGGATAACGCGTCTGACTACGGATCAGAAGA TTCCAGGTTCGACTCCTGGCTGGCTCG
331	Val_CAC_chr6:26538281-26538354 (+)	GGTTCCGTAGTGTAGTGGTTATCACGTTTCGCCCTCACACGCGAAAGGTC CCCGGTTCGAAACCGGGGGGAAACA
332	Ala_CGC_chr6:26553730-26553802 (+)	AGGGGATGTAGTCACTAGTGGTAGAGCGCATGCTTCGCATGTATGAGGTC CCGGGTTTCGATCCCCGGCATCTCCA
333	Ile_AAT_chr6:26554349-26554423 (+)	TGGCCGGTTAGCTCAGTTGGTTAGAGCGGTGTGTCTAATAACGCCAAGG TCGCGGGTTCGATCCCCCGTACGGGCCA

334	Pro_AGG_chr6:26555497-26555569 (+)	CGGCTCGTTGGTCTAGGGGTATGATTCCTCGCTTAGGGTGCGAGAGGTC CCGGGTTCAAATCCCGGACGAGCCC
335	Lys_CTT_chr6:26556773-26556846 (+)	AGCCCCGCTAGCTCAGTCGGTAGAGCATGAGACTCTTAATCTCAGGGT CGTGGGTTTCGAGCCCCACGTTGGCG
336	Tyr_GTA_chr6:26569085-26569176 (+)	TCCTTCGATAGCTCAGTTGGTAGAGCGGAGGACTGTAGTTGGCTGTGT CCTTAGACATCCTTAGGTCGCTGGTTCGAATCCGGCTCGAAGGA
337	Ala_AGC_chr6:26572091-26572164 (-)	GGGGAATTAGCTCAAATGTAGAGCGCTCGCTTAGCATGCGAGAGGTA GCGGGATCGATGCCCGCATTTCTCCAG
338	Met_CAT_chr6:26766443-26766516 (+)	CGCCCTCTTAGCGCAGCGGGCAGCGCTCAGTCTCATAATCTGAAGGT CCTGAGTTCGAGCCTCAGAGAGGGCA
339	Ile_TAT_chr6:26988124-26988218 (+)	TGCTCCAGTGGCGCAATCGGTTAGCGCGGGTACTTATATGGCAGTAT GTGTGCGAGTGTATGCCGAGGTTGTGAGTTCGAGCCTCACCTGGAGCA
340	His_GTG_chr6:27125905-27125977 (+)	TGCCGTGATCGTATAGTGGTGTAGTACTCTGCGTTGTGGCCGCAAGCAACC TCGGTTCGAATCCGAGTCACGGCA
341	Ile_AAT_chr6:27144993-27145067 (-)	GGCCGGTTAGCTCAGTTGGTTAGAGCGGTGTGCTAATAACGCCAAGGT CGCGGGTTCGATCCCGCTACGGGCCAC
342	Val_AAC_chr6:27203287-27203360 (+)	AGTTCCGTAAGTGTAGTGGTTATCACGTTTGCCCTAACACGCGAAAGGTC CCC GGTTTCGAAACCGGGCAGAAACA
343	Val_CAC_chr6:27248048-27248121 (-)	GCTTCTGTAGTGTAGTGGTTATCACGTTTCGCCCTCACACGCGAAAGGTCC CCGGTTCGAAACCGGGCAGAAAGCAA
344	Asp_GTC_chr6:27447452-27447524 (+)	TTCCCTCGTTAGTATAGTGGTGTAGTATCCCCGCCCTGTACGCGGGAGACC GGGGTTCGATTCCCCGACGGGGAG
345	Ser_TGA_chr6:27473606-27473688 (-)	GTAGTCGTGGCCGAGTGGTTAAGGCGGATGGACTTGAAATCCAATTGGGG TTTCCCCCGCGCAGGTTTCGAATCCGTGCGGCTACGG
346	Gln_CTG_chr6:27487307-27487379 (+)	AGGTTCCATGGTGTAAATGGTTAGCACTCTGGACTCTGAATCCAGCGAT CCGAGTTCAAATCTCGGTGGAACCT
347	Asp_GTC_chr6:27551235-27551307 (-)	TCCTCGTTAGTATAGTGGTGTAGTGTCCCCGCTGTGTACGCGGGGAGACC GGGGTTCGATTCCCCGACGGGGAGA
348	Val_AAC_chr6:27618706-27618779 (-)	GTTTCCGTAGTGTAGTGGTTATCACGTTTCGCCCTAACACGCGAAAGGTCC CTGGATCAAACCCAGGCGGAACAA
349	Ile_AAT_chr6:27655966-27656040 (+)	CGGCCGGTTAGCTCAGTTGGTTAGAGCGGTGTGCTAATAACGCCAAGG TCGCGGGTTCGATCCCCGTA CTGGCCA

350	Gln_CTG_ch6:27759134-27759206 (-)	GGCCCCATGGTGAATGGTCAGCACTCTGGACTCTCTGAATCCAGCGATC CGAGTTCAAATCTCGGTGGACCCA
351	Gln_TTG_ch6:27763639-27763711 (-)	GGCCCCATGGTGAATGGTTAGCACTCTGGACTTTGAATCCAGCGATC CGAGTTCAAATCTCGGTGGACCTT
352	Ala_AGC_ch6:28574932-28575004 (+)	TGGGGTGTAGCTCAGTGTAGAGCGCGTGTCTTAGCATGTACGAGGTC CCGGTTCAATCCCCGGCACCTCCA
353	Ala_AGC_ch6:28626013-28626085 (-)	GGGATGTAGCTCAGTGTAGAGCGCATGTCTTAGCATGCATGAGGTCC CGGGTTCGATCCCCAGCATCTCCAG
354	Ala_CGC_ch6:28697091-28697163 (+)	AGGGGTGTAGCTCAGTGTAGAGCGCGTGTCTTCGCATGTACGAGGCC CCGGTTCGACCCCCGGCTCCTCCA
355	Ala_AGC_ch6:28806220-28806292 (-)	GGGGTGTAGCTCAGTGTAGAGCGCGTGTCTTAGCATGCACGAGGCC CGGGTTCAAATCCCCGGCACCTCCAT
356	Ala_AGC_ch6:28831461-28831533 (-)	GGGGTGTAGCTCAGTGTAGAGCGCGTGTCTTAGCATGCACGAGGCC CGGGTTCAAATCCCCGGCACCTCCAG
357	Leu_CAA_ch6:28863999-28864105 (-)	GTCAGGATGGCCGAGTGTCTAAGGCGCCAGACTCAAAGCTAAAGCTTCC TCCGCGTGGGATTCTGGTCTCCAATGAGGCGTGGGTTCGAAATCCC
358	Leu_CAA_ch6:28908829-28908934 (+)	TGTCAGGATGGCCGAGTGTCTAAGGCGCCAGACTCAAAGCTTGGCTTC CTCGTGTGAGGATTCTGGTCTCCAATGGAGCGGTGGGTTCGAATCCC
359	Gln_CTG_ch6:28909377-28909449 (-)	GGTTCATGGTGAATGGTGTAGCACTCTGGACTCTGAATCCAGCGATCC GAGTTCAAATCTCGGTGGAACTT
360	Leu_AAG_ch6:28911398-28911480 (-)	GGTAGCGTGGCCGAGCGGTCTAAGGCGTGGATTAAAGGCTCCAGTCTC TTCGGGGCGTGGGTTCGAATCCCACCGCTGCCAG
361	Met_CAT_ch6:28912351-28912424 (+)	TGCTTCCTTAGCGCAGTAGGCAGCGGTCTAGTCTCATAATCTGAAGGT CCTGAGTTCGAACCTCAGAGGGGCA
362	Lys_TTT_ch6:28918805-28918878 (+)	AGCCCCGATAGCTCAGTCGGTAGAGCATCAGACTTTTAAATCTGAGGT CCAGGGTTCAGTCCCTGTTCGGGCG
363	Met_CAT_ch6:28921041-28921114 (-)	GCCTCCTTAGCGCAGTAGGCAGCGGTCTAGTCTCATAATCTGAAGGT CTGAGTTCGAACCTCAGAGGGGCGAG
364	Glu_CTC_ch6:28949975-28950047 (+)	TTCCCTGGTGTCTAGTGTAGGATTCGGCGCTCTCACCCGCCGGGCC CGGGTTCGATTCCCCGGTCAGGGAA
365	Leu_TAA_ch6:144537683-144537766 (+)	CACCAGGATGGCCGAGTGGTTAAGGCGTTGGACTTAAGATCCAATGGA CATATGTCGCGGTGGGTTCGAACCCCACTCCTGGTA

366	Pro_AGG_chr7:128423503-128423575 (+)	TGGCTCGTTGGTCTAGGGGTATGATTCTCGCTTAGGGTGCGAGAGGTC CCGGTTCAATATCCCGGACGAGCCC
367	Arg_CCT_chr7:139025445-139025518 (+)	AGCCCCAGTGGCTAATGGATAAGGCATTGGCCCTCTTAAGCCAGGGAT TGTGGTTTCGAGTCCCATCTGGGGTG
368	Cys_GCA_chr7:149388271-149388343 (-)	GGGGATATAGCTCAGGGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCGGTTCAAATCCGGGTGCCCCCCC
369	Tyr_GTA_chr8:67025601-67025694 (+)	CCCTTCGATAGCTCAGCTGGTAGAGCGGAGGACTGTAGCTACTTCTC AGCAGGAGACATCCTTAGGTCTGGTTCGATTCCGGCTCGAAGGA
370	Tyr_GTA_chr8:67026222-67026311 (+)	CCCTTCGATAGCTCAGCTGGTAGAGCGGAGGACTGTAGCGCGCGCCC GTGGCCATCCTTAGGTCTGGTTCGATTCCGGCTCGAAGGA
371	Ala_AGC_chr8:67026423-67026496 (+)	TGGGGATTAGCTCAAAATGGTAGAGCGCTCGCTTAGCATGCGAGAGGT AGCGGGATCGATGCCCCCATCTCTCA
372	Ser_AGA_chr8:96281884-96281966 (-)	GTAGTCGTGGCCGAGTGGTTAAGGCGATGGACTAGAAATCCATTGGGG TCTCCCCGCGCAGGTTCGAAATCTGCCGACTACGG
373	Met_CAT_chr8:124169469-124169542 (-)	GCCTCGTTAGCGCAGTAGGTAGCGCTCAGTCTCATAATCTGAAAGTCT GTGAGTTCGATCCTCACACGGGGCAC
374	Arg_TCT_chr9:131102354-131102445 (-)	GGCTCTGTGGCGCAATGGATAGCGCATTTGGACTTCTAGCTGAGCCTAG TGTGTCATTCAAAAGGTGTGGGTTCGAGTCCACCAGAGTCGA
375	Asn_GTT_chr10:22518437-22518511 (-)	GTCTCTGTGGCGCAATCGGTAGCGCGTTCGGCTGTAAACCGAAAAGGT TGGTGGTTTCGAGCCCCACCCAGGACGC
376	Ser_TGA_chr10:69524260-69524342 (+)	GGCAGCGATGGCCGAGTGGTTAAGCGTTGGACTTGAAATCCAATGGG GTCTCCCCGCGCAGGTTCGAACCTGCTCGCTGCG
377	Val_TAC_chr11:59318101-59318174 (-)	GGTTCCATAGTGTAGTGGTTATCACGCTCTGCTTTACACGCGAGAAGGTCC TGGGTTTCGAGCCCCAGTGAACCAT
378	Val_TAC_chr11:59318459-59318532 (-)	GGTTCCATAGTGTAGCGGTTATCACGCTCTGCTTTACACGCGAGAAGGTCC TGGGTTTCGAGCCCCAGTGAACCAT
379	Arg_TCT_chr11:59318766-59318852 (+)	TGGCTCTGTGGCGCAATGGATAGCGCATTTGGACTTCTAGATAGTTAGA GAAATTCAAAGGTGTGGGTTCGAGTCCACCAGAGTCG
380	Leu_TAA_chr11:59319227-59319310 (+)	TACCAGAAATGGCCGAGTGGTTAAGGCGTTGGACTTAAGATCCAATGGA TTCATATCCGCGTGGGTTCGAACCCCACTTCTGGTA
381	Lys_TTT_chr11:59323901-59323974 (+)	GGCCCCGATAGCTCAGTCGGTAGAGCATCAGACTTTTAAATCTGAGGGT CCGGGGTTCAAGTCCCTGTTCGGGCG

382	Phe_GAA_chr11:59324969-59325042 (-)	GCCGAAATAGCTCAGTTGGGAGAGCGTTAGACTGAAAGATCTAAAGGTC CCTGTTTCGATCCCGGGTTTCGGCAG
383	Lys_TTT_chr11:59327807-59327880 (-)	GCCCGGATAGCTCAGTCGCTAGAGCATCAGACTTTTAAATCTGAGGGTC CAGGGTTCAAGTCCCTGTTCCGGCGG
384	Phe_GAA_chr11:59333852-59333925 (-)	GCCGAAATAGCTCAGTTGGGAGAGCGTTAGACTGAAAGATCTAAAGGTC CCTGTTCAAATCCCGGGTTTCGGCAG
385	Ser_GCT_chr11:66115590-66115672 (+)	GGACGAGTGGCCGAGTGGTTAAGGCGATGGACTGCTAATCCATTGTG CTTTGCACGCTGGGTTTCGAATCCCATCCTCGTCG
386	Pro_TGG_chr11:75946868-75946940 (-)	GGCTCGTTGGTCTAGGGGTATGATTCTCGGTTTGGGTCCGAGAGGTCCCC GGGTTCAAATCCCGGACGAGCCCC
387	Ser_CGA_chr12:56584147-56584229 (+)	AGTCACGGTGGCCGAGTGGTTAAGGCGTTGGACTCGAAATCCAATGGG GTTTCCCCGCACAGGTTTCGAATCCTGTTCTGTCGACG
388	Asp_GTC_chr12:98897280-98897352 (+)	CTCCTCGTTAGTATAGTGGTTAGTATCCCCGCCCTGTCACGCGGGAGACC GGGTTCAAATCCCCGACGGGGAG
389	Trp_CCA_chr12:98898029-98898101 (+)	GGACCTCGTGGCGCAACGTTAGCGCGTCTGACTCCAGATCAGAAAGGCT GCGTGTTCGAATCACGTCGGGGTCA
390	Ala_TGC_chr12:125406300-125406372 (-)	GGGGATGTAGCTCAGTGGTAGAGCGCATGTCTTGCATGTATGAGGCCC CGGGTTCGATCCCCGGCATCTCCAT
391	Phe_GAA_chr12:125412388-125412461 (-)	GCCGAAATAGCTCAGTTGGGAGAGCGTTAGACTGAAAGATCTAAAGGTC CCTGTTTCGATCCCCGGGTTTCGGCAC
392	Ala_TGC_chr12:125424511-125424583 (+)	AGGGGATGTAGCTCAGTGGTAGAGCGCATGCTTTGCACGTATGAGGCC CCGGGTTCAAATCCCCGGCATCTCCA
393	Asn_GTT_chr13:31248100-31248174 (-)	GTCTCTGTGGCGCAATCGGTTAGCGCGTTCGGCTGTAAACCGAAAAGGT TGGTGGTTCGAGCCCCACCCAGGGACGG
394	Glu_TTC_chr13:45492061-45492133 (-)	TCCCACATGGTCTAGCGGTTAGGATTCCTGTGTTTTCACCCAGCGCGGCC GGGTTTCGACTCCCGGTGTGGGAAC
395	Thr_TGT_chr14:21081948-21082021 (-)	GGCTCCATAGCTCAGGGGTTAGAGCGCTGGTCTTGTAAACCAAGGGGTC GCGAGTTCAAATTCGCTGGGGCCCTG
396	Leu_TAG_chr14:21093528-21093610 (+)	TGGTAGTGTGGCCGAGCGGTCTAAGGCGCTGGATTAGGCTCCAGTCT CTTCGGGGCGTGGGTTTCGAATCCCACTGCCA
397	Thr_TGT_chr14:21099318-21099391 (-)	GGCTCCATAGCTCAGGGGTTAGAGCACTGGTCTTGTAAACCAAGGGGTC GCGAGTTCAAATCTCGCTGGGGCCCTC

398	Pro_TGG_chr14:21101164-21101236 (+)	TGGCTCGTTGGTCTAGTGGTATGATTCTCGCTTTGGGTGCGAGAGGTCC CGGGTTCAAATCCCGACGAGCCC
399	Tyr_GTA_chr14:21131350-21131444 (-)	CCTTCGATAGCTCAGCTGGTAGAGCGGAGGACTGTAGATTGTACAGAC ATTTCGGACATCCTTAGTCTGCTGGTTCGATTCGGCTCGAAGGAA
400	Thr_TGT_chr14:21149848-21149921 (+)	AGGCCCTATAGCTCAGGGGTAGAGCACTGGTCTTGTAAACCAAGGGGT CGCGAGTTCAAATCTCGCTGGGGCT
401	Tyr_GTA_chr14:21151431-21151520 (+)	TCCTTCGATAGCTCAGCTGGTAGAGCGGAGGACTGTAGTACTTAATGT GTGTCATCCTTAGGTCGCTGGTTCGATTCCGGCTCGAAGGA
402	Pro_TGG_chr14:21152174-21152246 (+)	TGGCTCGTTGGTCTAGGGGTATGATTCTCGCTTTGGGTGCGAGAGGTCC CGGGTTCAAATCCCGACGAGCCC
403	Lys_CTT_chr14:58706612-58706685 (-)	GCCCGGTAGCTCAGTCGGTAGAGCATGGGACTCTTAATCCCAGGGTC GTGGGTTTCGAGCCCCACGTTGGGCGC
404	Ile_AAT_chr14:102783428-102783502 (+)	CGGCCGGTTAGCTCAGTTGGTTAGAGCGGTGGTGTATAACGCCAAGG TCGCGGGTTCGATCCCCGTACGGGCCA
405	Glu_TTC_chr15:26327380-26327452 (-)	TCCACATGGTCTAGCGGTTAGGATTCTCTGTTTTCACCCAGCGGGCCC GGGTTTCGACTCCCGGTGTGGGAAT
406	Ser_GCT_chr15:40886022-40886104 (-)	GACGAGGTGGCCGAGTGGTTAAGGCGATGGACTGCTAATCCATTGTGC TCTGCACGCGTGGGTTTCGAATCCCATCCTCGTCGA
407	His_GTG_chr15:45490803-45490875 (-)	GCCGTGATCGTATAGTGGTTAGTACTCTGCTTGTGGCCGACGCAACCT CGGTTCGAATCCGAGTCACGGCAT
408	His_GTG_chr15:45493348-45493420 (+)	CGCCGTGATCGTATAGTGGTTAGTACTCTGCTGCGTTGTGGCCGACGCAAC CTCGGTTTCGAATCCGAGTACGGCA
409	Gln_CTG_chr15:66161399-66161471 (-)	GGTTCCATGGTGTAATGGTTAGCACTCTGGACTCTGAATCCAGCGATCC GAGTTCAAATCTCGGTGGAACCTG
410	Lys_CTT_chr15:79152903-79152976 (+)	TGCCCGGCTAGCTCAGTCGGTAGAGCATGGGACTCTTAATCCCAGGGT CGTGGGTTTCGAGCCCCACGTTGGCG
411	Arg_TCG_chr15:89878303-89878376 (+)	GGGCCGCTGGCCTAATGGATAAGGCGTCTGACTTCGGATCAGAAGAT TGCAGGTTTCGAGTCTCGCCGCGGTG
412	Gly_CCC_chr16:686735-686806 (-)	GCGCCGCTGGTGTAGTGGTATCATGCAAGATTCCCATTTCTTCCGACCCG GGTTCGATTCCCGGGCGGGCAC
413	Arg_CCG_chr16:3200674-3200747 (+)	GGGCCGCTGGCCTAATGGATAAGGCGTCTGATTCGGGATCAGAAGAT TGAGGGTTCGAGTCCCTTCGTGGTTCG

414	Arg_CCT_chr16:3202900-3202973 (+)	CGCCCCGGTGGCCTAATGGATAAGGCATTGGCCTCCTAAGCCAGGGAT TGTGGGTTTCGAGTCCACCCGGGTA
415	Lys_CTT_chr16:3207405-3207478 (-)	GCCCGGCTAGCTCAGTCGGTAGACATGAGACCCCTTAATCTCAGGGTC GTGGGTTTCGAGCCCCACGTTGGGCGT
416	Thr_CGT_chr16:14379749-14379821 (+)	AGGCGCGTGGCCAAAGTGGTAAGGCGTCGGTCTCGTAACCCGAAGATC ACGGGTTTCGAACCCCGTCCGTGCCCT
417	Leu_TAG_chr16:22207031-22207113 (-)	GGTAGCGTGGCCGAGTGGTCTAAGGCGCTGGAATTTAGGCTCCAGTTCAT TTCGATGGCGTGGGTTTCGAATCCACCCGCTGCCAC
418	Leu_AAG_chr16:22308460-22308542 (+)	GGTAGCGTGGCCGAGCGGTCTAAGGCGCTGGATTAAAGGCTCCAGTCT CTTCGGGGCGTGGGTTTCGAATCCACCCGCTGCCA
419	Leu_CAG_chr16:57333862-57333945 (+)	AGTCAGGATGGCCGAGCGGTCTAAGGCGCTGCGTTTCAGGTCGCAGTCT CCCCTGAGGCGTGGGTTTCGAATCCCACTTCTGACA
420	Leu_CAG_chr16:57334391-57334474 (-)	GTCAGGATGGCCGAGCGGTCTAAGGCGCTGCGTTTCAGGTCGCAGTCTC CCCTGGAGCGTGGGTTTCGAATCCCACTTCTGACAG
421	Met_CAT_chr16:87417627-87417700 (-)	GCCTGTTAGCGCAGTAGGCGAGCGGTCTAGTCTCATAATCTGAAGGTC GTGAGTTCGAGCCTCACACGGGGCAG
422	Leu_TAG_chr17:8023631-8023713 (-)	GGTAGCGTGGCCGAGCGGTCTAAGGCGCTGGATTAGGCTCCAGTCTC TTCGGAGGCGTGGGTTTCGAATCCACCCGCTGCCAG
423	Arg_TCT_chr17:8024242-8024330 (+)	TGGCTCTGTGGCGCAATGGATAGCGCATTTGGACTTCTAGTGACGAATA GAGCAATTCAAAAGTTGTGGGTTTCGAATCCCAACCAGAGTCG
424	Gly_GCC_chr17:8029063-8029134 (+)	CGCATTGGTGGTTCAGTGGTAGAAATTCCTCGCCTGCCACGCGGGAGGCC CGGGTTCGATTCCCGGCCAATGCA
425	Ser_CGA_chr17:8042198-8042280 (-)	GCTGTGATGGCCGAGTGGTTAAGGCGTTGGACTCGAAATCCAATGGGG TCTCCCCCGCAGGTTTCGAATCCTGCTCACAGCGT
426	Thr_AGT_chr17:8042769-8042843 (-)	GGCGCCGTGGCTTAGCTGGTTAAAGCGCCTGTCTAGTAAACAGGAGAT CCTGGGTTTCGAATCCACCGGTGCGCTG
427	Trp_CCA_chr17:8089675-8089747 (+)	CGACCTCGTGGCGCAACGGTAGCGGCTGTGACTCCAGATCAGAAAGGTT GCGTGTTCAAATCACGTCGGGGTCA
428	Ser_GCT_chr17:8090183-8090265 (+)	AGACGAGTGGCCGAGTGGTTAAGGCGATGGACTGCTAATCCATTGTG CTCTGCACGCGTGGGTTTCGAATCCCACTCTCGTCG
429	Thr_AGT_chr17:8090477-8090551 (+)	CGGCGCCGTGGCTTAGTTGGTTAAAGCGCCTGTCTAGTAAACAGGAGA TCCTGGGTTTCGAATCCACGCGGTGCCT

430	Trp_CCA_chr17:8124186-8124258 (-)	GGCTCGTGGCGCAACGGTAGCGCGTCTGTGACTCCAGATCAGAAAGGTTG CGTGTTCAAATCACGTCGGGGTCAA
431	Gly_TCC_chr17:8124865-8124937 (+)	AGCGTTGGTGGTATAGTGGTAAGCATAGCTGCCTTCCAAAGCAGTTGAC CCGGGTTTCGATTCCCGGCCAACGCA
432	Asp_GTC_chr17:8125555-8125627 (-)	TCCTCGTTAGTATAGTGGTGAATATCCCCGCCCTGTACGCGGGAGACC GGGTTTCGATTCCCCGACGGGGAGA
433	Pro_CGG_chr17:8126150-8126222 (-)	GGCTCGTTGGTCTAGGGGTATGATTCCTCGCTTCGGGTGCGAGAGGTCC CGGGTTCAAATCCCGGACGAGCCCT
434	Thr_AGT_chr17:8129552-8129626 (-)	GGCGCGTGGCTTAGTTGGTTAAAGCGCCTGTCTAGTAAACAGGAGAT CCTGGGTTTCGAATCCACGCGGTGCTT
435	Ser_AGA_chr17:8129927-8130009 (-)	GTAGTCGTGGCCGAGTGGTTAAGCGGATGGACTAGAAATCCATTGGGG TCTCCCCGCGAGGTTTCGAATCCTGCCGACTACGT
436	Trp_CCA_chr17:19411493-19411565 (+)	TGACCTCGTGGCGCAATGGTAGCGCTGTGACTCCAGATCAGAAAGGTT GCGTGTTCAAAGTCAAGTCAGTCGGGGTCA
437	Thr_CGT_chr17:29877092-29877164 (+)	AGCGCGGTGGCCAAAGTGGTAAAGCGTCTCGTAAACCGAAGATC GCGGGTTCGAACCCCGTCCGTGCCT
438	Cys_GCA_chr17:37023897-37023969 (+)	AGGGGTATAGCTCAGTGGTAGAGCATTTGACTGCAGATCAAGAGGTC CCCGGTTCAAATCCGGGTGCCCCCT
439	Cys_GCA_chr17:37025544-37025616 (-)	GGGGTATAGCTCAGTGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CTGGTTCAAATCCGGGTGCCCCCTC
440	Cys_GCA_chr17:37309986-37310058 (-)	GGGGTATAGCTCAGTGGTAGAGCATTTGACTGCAGATCAAGAGGTCC CCGGTTCAAATCCGGGTGCCCCCTC
441	Gln_TTG_chr17:47269889-47269961 (+)	AGTCCCATGGTGTAAATGGTTAGCACTCTGGACTTTTGAATCCAGCGAT CCGAGTTCAAATCTCGGTGGACCT
442	Arg_CCG_chr17:66016012-66016085 (-)	GACCCAGTGGCCTAATGGATAAGGCATCAGCCTCCGGAGCTGGGGATT GTGGGTTTCGAGTCCCATCTGGGTCGC
443	Arg_CCT_chr17:73030000-73030073 (+)	AGCCCCAGTGGCCTAATGGATAAGGCACCTGGCCTCCTAAGCCAGGGAT TGTGGGTTTCGAGTCCCACTGGGGTA
444	Arg_CCT_chr17:73030525-73030598 (-)	GCCCCAGTGGCCTAATGGATAAGGCACCTGGCCTCCTAAGCCAGGGATT GTGGGTTTCGAGTCCCACTGGGGTGT
445	Arg_TCG_chr17:73031207-73031280 (+)	AGACCGCGTGGCCTAATGGATAAGCGGTCTGACTTCGGATCAGAAAGAT TGAGGGTTCGAGTCCCTTCGTGGTCCG

446	Asn_GTT_chr19:1383561-1383635 (+)	CGTCTCTGTGGCGCAATCGGTTAGCGCGTTTCGGCTGTTAACCCGAAAGG TTGGTGGTTCGAGCCACCCAGGGACG
447	Gly_TCC_chr19:4724081-4724153 (+)	GGCGTTGGTGGTATAGTGGTTAGCATAGCTGCCTTCCAAGCAGTTGAC CCGGGTTTCGATTCCCGGCCAACGCA
448	Val_CAC_chr19:4724646-4724719 (-)	GTTTCCGTAGTAGCGGTTATCACATTTCGCCCTCACACGCGAAAGGTCC CCGGTTCGATCCCGGGCGGAAACAG
449	Thr_AGT_chr19:33667962-33668036 (+)	TGGCGCCGTGGCTTAGTTGGTTAAAGCGCCCTGTCTAGTAAACAGGAGA TCCTGGGTTTCGAAATCCAGCGGTGCCT
450	Ile_TAT_chr19:39902807-39902900 (-)	GC'TCCAGTGGCGCAATCGGTTAGCGCGCGGTACTTATATGACAGTGCG AGCGGAGCAATGCCGAGGTTGTGAGTTCGATCCTCACCTGGAGCAC
451	Gly_GCC_chr21:18827106-18827177 (-)	GCATGGGTGGTTCAGTGGTAGAATTCTCGCCTGCCACGCGGGAGGCC GGGTTTCGATTCCCGGCCCATGCAG

Non-naturally occurring modification

A TREM, a TREM core fragment or a TREM fragment described herein may comprise a non-naturally occurring modification, e.g., a modification described in any one of Tables 10-14.

A non-naturally occurring modification can be made according to methods known in the art.

- 5 Methods of making non-naturally occurring modifications are known in the art; for example, several methods are provided in the Examples described herein.

In an embodiment, a non-naturally occurring modification is a modification that a cell, e.g., a human cell, does not make on an endogenous tRNA.

- 10 In an embodiment, a non-naturally occurring modification is a modification that a cell, e.g., a human cell, can make on an endogenous tRNA, but wherein such modification is in a location in which it does not occur on a native tRNA. In an embodiment, the non-naturally occurring modification is in a domain, linker or arm which does not have such modification in nature. In an embodiment, the non-naturally occurring modification is at a position within a domain, linker or arm, which does not have such modification in nature. In an embodiment, the non-naturally occurring modification is on a nucleotide which does not have such modification in nature. In an embodiment, the non-naturally occurring modification is on a nucleotide at a position within a domain, linker or arm, which does not have such modification in nature.

- 15 In an embodiment, a TREM, a TREM core fragment or a TREM fragment described herein comprises a non-naturally occurring modification provided in Table 10 or a combination thereof.

Table 10: Exemplary non-naturally occurring modifications

Modification	Modification
7-deaza-adenosine	2-(aminopropyl)adenine
N1-methyl-adenosine	2-(halo)adenine
N6, N6 (dimethyl)adenine	2-(propyl)adenine
N6-cis-hydroxy-isopentenyl-adenosine	2'-azido-2'-deoxy-adenosine
thio-adenosine	2'-Deoxy-2'-alpha-aminoadenosine
2-(amino)adenine	2'-Deoxy-2'-alpha-azidoadenosine
2-(aminopropyl)adenine	6-(alkyl)adenine
2-(methylthio) N6 (isopentenyl)adenine	6-(methyl)adenine
2-(alkyl)adenine	6-(alkyl)adenine
2-(aminoalkyl)adenine	

Modification
6-(methyl)adenine
7-(deaza)adenine
8-(alkenyl)adenine
8-(alkynyl)adenine
8-(amino)adenine
8-(thioalkyl)adenine
8-(alkenyl)adenine
8-(alkyl)adenine
8-(alkynyl)adenine
8-(amino)adenine
8-(halo)adenine
8-(hydroxyl)adenine
8-(thioalkyl)adenine
8-(thiol)adenine
8-azido-adenosine
azaadenine
deazaadenine
N6-(methyl)adenine
N6-(isopentyl)adenine
7-deaza-8-aza-adenosine
7-methyladenine
1-deazaadenosine
2'-Fluoro-N6-Bz-deoxyadenosine
2'-OMe-2-Amino-adenosine
2'-O-methyl-N6-Bz-deoxyadenosine
2'-alpha-ethynyladenosine
2-aminoadenine
2-Aminoadenosine
2-Amino-adenosine
2'-alpha-Trifluoromethyladenosine
2-Azidoadenosine

Modification
2'-beta-Ethynyladenosine
2-Bromoadenosine
2'-beta-Trifluoromethyladenosine
2-Chloroadenosine
2'-Deoxy-2',2'-difluoroadenosine
2'-Deoxy-2'-alpha-mercaptoadenosine
2'-Deoxy-2'-alpha-thiomethoxyadenosine
2'-Deoxy-2'-beta-aminoadenosine
2'-Deoxy-2'-beta-azidoadenosine
2'-Deoxy-2'-beta-bromoadenosine
2'-Deoxy-2'-beta-chloroadenosine
2'-Deoxy-2'-beta-fluoroadenosine
2'-Deoxy-2'-beta-iodoadenosine
2'-Deoxy-2'-beta-mercaptoadenosine
2'-Deoxy-2'-beta-thiomethoxyadenosine
2-Fluoroadenosine
2-Iodoadenosine
2-Mercaptoadenosine
2-methoxy-adenine
2-methylthio-adenine
2-Trifluoromethyladenosine
3-Deaza-3-bromoadenosine
3-Deaza-3-chloroadenosine
3-Deaza-3-fluoroadenosine
3-Deaza-3-iodoadenosine
3-Deazaadenosine
4'-Azidoadenosine
4'-Carbocyclic adenosine
4'-Ethynyladenosine
5'-Homo-adenosine

Modification
8-Aza-adenosine
8-bromo-adenosine
8-Trifluoromethyladenosine
9-Deazaadenosine
2-aminopurine
7-deaza-2,6-diaminopurine
7-deaza-8-aza-2,6-diaminopurine
7-deaza-8-aza-2-aminopurine
2,6-diaminopurine
7-deaza-8-aza-adenine, 7-deaza-2-aminopurine
4-methylcytidine
5-aza-cytidine
Pseudo-iso-cytidine
pyrrolo-cytidine
alpha-thio-cytidine
2-(thio)cytosine
2' -Amino-2' -deoxy-cytosine
2' -Azido-2' -deoxy-cytosine
2' -Deoxy-2' -alpha-aminocytidine
2' -Deoxy-2' -alpha-azidocytidine
3 (deaza) 5 (aza)cytosine
3 (methyl)cytosine
3-(alkyl)cytosine
3-(deaza) 5 (aza)cytosine
3-(methyl)cytidine
4,2'-O-dimethylcytidine
5 (halo)cytosine
5 (methyl)cytosine
5 (propynyl)cytosine
5 (trifluoromethyl)cytosine

Modification
5-(alkyl)cytosine
5-(alkynyl)cytosine
5-(halo)cytosine
5-(propynyl)cytosine
5-(trifluoromethyl)cytosine
5-bromo-cytidine
5-iodo-cytidine
5-propynyl cytosine
6-(azo)cytosine
6-aza-cytidine
aza cytosine
deaza cytosine
N4 (acetyl)cytosine
1-methyl-1-deaza-pseudoisocytidine
1-methyl-pseudoisocytidine
2-methoxy-5-methyl-cytidine
2-methoxy-cytidine
2-thio-5-methyl-cytidine
4-methoxy-1-methyl-pseudoisocytidine
4-methoxy-pseudoisocytidine
4-thio- 1-methyl-1-deaza-pseudoisocytidine
4-thio-1-methyl-pseudoisocytidine
4-thio-pseudoisocytidine
5-aza-zebularine
5-methyl-zebularine
pyrrolo-pseudoisocytidine
zebularine
(E)-5-(2-Bromo-vinyl)cytidine
2,2'-anhydro-cytidine
2'-Fluor-N4-Bz-cytidine

Modification
2'-Fluoro-N4-Acetyl-cytidine
2'-O-Methyl-N4-Acetyl-cytidine
2'-O-methyl-N4-Bz-cytidine
2'-a-Ethynylcytidine
2'-a-Trifluoromethylcytidine
2'-b-Ethynylcytidine
2'-b-Trifluoromethylcytidine
2'-Deoxy-2',2'-difluorocytidine
2'-Deoxy-2'-alpha-mercaptopcytidine
2'-Deoxy-2'-alpha-thiomethoxycytidine
2'-Deoxy-2'-betab-aminocytidine
2'-Deoxy-2'-beta-azidocytidine
2'-Deoxy-2'-beta-bromocytidine
2'-Deoxy-2'-beta-chlorocytidine
2'-Deoxy-2'-beta-fluorocytidine
2'-Deoxy-2'-beta-iodocytidine
2'-Deoxy-2'-beta-mercaptopcytidine
2'-Deoxy-2'-beta-thiomethoxycytidine TP
2'-O-Methyl-5-(1-propynyl)cytidine
3'-Ethynylcytidine
4'-Azidocytidine
4'-Carbocyclic cytidine
4'-Ethynylcytidine
5-(1-Propynyl)ara-cytidine
5-(2-Chloro-phenyl)-2-thiocytidine
5-(4-Amino-phenyl)-2-thiocytidine
5-Aminoallyl-cytosine
5-Cyanocytidine
5-Ethynylara-cytidine
5-Ethynylcytidine

Modification
5'-Homo-cytidine
5-Methoxycytidine
5-Trifluoromethyl-Cytidine
N4-Amino-cytidine
N4-Benzoyl-cytidine
pseudoisocytidine
6-thio-guanosine
7-deaza-guanosine
8-oxo-guanosine
N1-methyl-guanosine
alpha-thio-guanosine
2-(propyl)guanine
2-(alkyl)guanine
2'-Amino-2'-deoxy-guanosine
2'-Azido-2'-deoxy-guanosine
2'-Deoxy-2'-alpha-aminoguanosine
2'-Deoxy-2'-alpha-azidoguanosine
6-(methyl)guanine
6-(alkyl)guanine
6-(methyl)guanine
6-methyl-guanosine
7-(alkyl)guanine
7-(deaza)guanine
7-(methyl)guanine
7-(alkyl)guanine
7-(deaza)guanine
7-(methyl)guanine
8-(alkyl)guanine
8-(alkynyl)guanine
8-(halo)guanine
8-(thioalkyl)guanine

Modification
8-(alkenyl)guanine
8-(alkyl)guanine
8-(alkynyl)guanine
8-(amino)guanine
8-(halo)guanine
8-(hydroxyl)guanine
8-(thioalkyl)guanine
8-(thiol)guanine
azaguanine
deaza guanine
N (methyl)guanine
N-(methyl)guanine
1-methyl-6-thio-guanosine
6-methoxy-guanosine
6-thio-7-deaza-8-aza-guanosine
6-thio-7-deaza-guanosine
6-thio-7-methyl-guanosine
7-deaza-8-aza-guanosine
7-methyl-8-oxo-guanosine
N2,N2-dimethyl-6-thio-guanosine
N2-methyl-6-thio-guanosine
1-Me-guanosine
2' Fluoro-N2-isobutyl-guanosine
2' O-methyl-N2-isobutyl-guanosine
2' -alpha-Ethynylguanosine
2' -alpha-Trifluoromethylguanosine
2' -beta-Ethynylguanosine
2' -beta-Trifluoromethylguanosine
2' -Deoxy-2',2'-difluoroguanosine
2' -Deoxy-2' -alpha-mercaptoguanosine
2' -Deoxy-2' -alpha-

Modification
thiomethoxyguanosine
2' -Deoxy-2' -beta-aminoguanosine
2' -Deoxy-2' -beta-azidoguanosine
2' -Deoxy-2' -beta-bromoguanosine
2' -Deoxy-2' -beta-chloroguanosine
2' -Deoxy-2' -beta-fluoroguanosine
2' -Deoxy-2' -beta-iodoguanosine
2' -Deoxy-2' -beta-mercaptoguanosine
2' -Deoxy-2' -beta-thiomethoxyguanosine
4' -Azidoguanosine
4' -Carbocyclic guanosine
4' -Ethynylguanosine
5' -Homo-guanosine
8-bromo-guanosine
9-Deazaguanosine
N2-isobutyl-guanosine
7-methylinosine
allyamino- thymidine
aza thymidine
deaza thymidine
deoxy-thymidine
5-propynyl uracil
alpha-thio-uridine
1-(aminoalkylamino-carbonylethynyl)-2(thio)-pseudouracil
1-(aminoalkylaminocarbonylethynyl)-2,4-(dithio)pseudouracil
1-(aminoalkylaminocarbonylethynyl)-4(thio)pseudouracil
1-(aminoalkylaminocarbonylethynyl)-pseudouracil
1-(aminocarbonylethynyl)-2(thio)-pseudouracil
1-(aminocarbonylethynyl)-2,4-(dithio)pseudouracil

Modification
1-(aminocarbonylethyl-4 (thio)pseudouracil
1-(aminocarbonylethyl)-pseudouracil
1-substituted 2-(thio)-pseudouracil
1-substituted 2,4-(dithio)pseudouracil
1-substituted 4 (thio)pseudouracil
1-substituted pseudouracil
1-(aminoalkylamino-carbonylethyl)-2- (thio)-pseudouracil
1-Methyl-3-(3-amino-3-carboxypropyl) pseudouridine
1-Methyl-3-(3-amino-3-carboxypropyl)pseudo-Uridine
1-Methyl-pseudo-UTP
2 (thio)pseudouracil
2' deoxy uridine
2' fluorouridine
2-(thio)uracil
2,4-(dithio)pseudouracil
2'-methyl, 2'-amino, 2'-azido, 2'-fluoro-guanosine
2'-Amino-2'-deoxy-uridine
2'-Azido-2'-deoxy-uridine
2' -Azido-deoxyuridine
2'-O-methylpseudouridine
2' deoxyuridine
2' fluorouridine
2' -Deoxy-2'-alpha-aminouridine TP
2' -Deoxy-2'-alpha-azidouridine TP
2-methylpseudouridine
3-(3-amino-3-carboxypropyl)uracil
4-(thio)pseudouracil
4-(thio)pseudouracil
4-(thio)uracil

Modification
4-thiouracil
5-(1,3-diazole-1-alkyl)uracil
5-(2-aminopropyl)uracil
5-(aminoalkyl)uracil
5-(dimethylaminoalkyl)uracil
5-(guanidiniumalkyl)uracil
5-(methoxycarbonylmethyl)-2-(thio)uracil
5-(methoxycarbonyl-methyl)uracil
5-(methyl)-2-(thio)uracil
5-(methyl)-2,4-(dithio)uracil
5 (methyl) 4 (thio)uracil
5 (methylaminomethyl)-2 (thio)uracil
5 (methylaminomethyl)-2,4 (dithio)uracil
5 (methylaminomethyl)-4 (thio)uracil
5 (propynyl)uracil
5 (trifluoromethyl)uracil
5-(2-aminopropyl)uracil
5-(alkyl)-2-(thio)pseudouracil
5-(alkyl)-2,4 (dithio)pseudouracil
5-(alkyl)-4 (thio)pseudouracil
5-(alkyl)pseudouracil
5-(alkyl)uracil
5-(alkynyl)uracil
5-(allylamino)uracil
5-(cyanoalkyl)uracil
5-(dialkylaminoalkyl)uracil
5-(dimethylaminoalkyl)uracil
5-(guanidiniumalkyl)uracil
5-(halo)uracil
5-(1,3-diazole-1-alkyl)uracil

Modification
5-(methoxy)uracil
5-(methoxycarbonylmethyl)-2-(thio)uracil
5-(methoxycarbonyl-methyl)uracil
5-(methyl) 2(thio)uracil
5-(methyl) 2,4 (dithio)uracil
5-(methyl) 4 (thio)uracil
5-(methyl)-2-(thio)pseudouracil
5-(methyl)-2,4 (dithio)pseudouracil
5-(methyl)-4 (thio)pseudouracil
5-(methyl)pseudouracil
5-(methylaminomethyl)-2 (thio)uracil
5-(methylaminomethyl)-2,4(dithio)uracil
5-(methylaminomethyl)-4-(thio)uracil
5-(propynyl)uracil
5-(trifluoromethyl)uracil
5-aminoallyl-uridine
5-bromo-uridine
5-iodo-uridine
5-uracil
6 (azo)uracil
6-(azo)uracil
6-aza-uridine
allylamino-uracil
aza uracil
deaza uracil
N3 (methyl)uracil
Pseudo-uridine-1-2-ethanoic acid
pseudouracil
4-Thio-pseudouridine
1-carboxymethyl-pseudouridine

Modification
1-methyl-1-deaza-pseudouridine
1-propynyl-uridine
1-taurinomethyl-1-methyl-uridine
1-taurinomethyl-4-thio-uridine
1-taurinomethyl-pseudouridine
2-methoxy-4-thio-pseudouridine
2-thio- 1-methyl-1-deaza-pseudouridine
2-thio-1-methyl-pseudouridine
2-thio-5-aza-uridine
2-thio-dihydropseudouridine
2-thio-dihydrouridine
2-thio-pseudouridine
4-methoxy-2-thio-pseudouridine
4-methoxy-pseudouridine
4-thio-1-methyl-pseudouridine
4-thio-pseudouridine
5-aza-uridine
dihydropseudouridine
(±)1-(2-Hydroxypropyl)pseudouridine
(2R)- 1 -(2-Hydroxypropyl)pseudouridine
(2S)- 1 -(2-Hydroxypropyl)pseudouridine
(E)-5-(2-Bromo-vinyl)ara-uridine
(E)-5-(2-Bromo-vinyl)uridine
(Z)-5-(2-Bromo-vinyl)ara-uridine
(Z)-5-(2-Bromo-vinyl)uridine
1-(2,2,2-Trifluoroethyl)-pseudouridine
1-(2,2,3,3,3-Pentafluoropropyl)pseudouridine
1-(2,2-Diethoxyethyl)pseudouridine
1-(2,4,6-Trimethylbenzyl)pseudouridine
1-(2,4,6-Trimethyl-benzyl)pseudo-uridine

Modification
1-(2,4,6-Trimethyl-phenyl)pseudo-uridine
1-(2-Amino-2-carboxyethyl)pseudo-uridine
1-(2-Amino-ethyl)pseudouridine
1-(2-Hydroxyethyl)pseudouridine
1-(2-Methoxyethyl)pseudouridine
1-(3,4-Bis-trifluoromethoxybenzyl)pseudouridine
1-(3,4-Dimethoxybenzyl)pseudouridine
1-(3-Amino-3-carboxypropyl)pseudo-uridine
1-(3-Amino-propyl)pseudouridine
1-(3-Cyclopropyl-prop-2-ynyl)pseudouridine TP
1-(4-Amino-4-carboxybutyl)pseudouridine
1-(4-Amino-benzyl)pseudouridine
1-(4-Amino-buty l)pseudouridine
1-(4-Amino-phenyl)pseudouridine
1-(4-Azidobenzyl)pseudouridine
1-(4-Bromobenzyl)pseudouridine
1-(4-Chlorobenzyl)pseudouridine
1-(4-Fluorobenzyl)pseudouridin
1-(4-Iodobenzyl)pseudouridine
1-(4-Methanesulfonylbenzyl)pseudouridine
1-(4-Methoxybenzy l)pseudouridine
1-(4-Methoxy-benzyl)pseudouridine
1-(4-Methoxy-phenyl)pseudouridine
1-(4-Methylbenzyl)pseudouridine
1-(4-Methyl-benzyl)pseudouridine
1-(4-Nitrobenzyl)pseudouridine
1-(4-Nitro-benzy l)pseudouridine
1(4-Nitro-phenyl)pseudouridine

Modification
1-(4-Thiomethoxybenzyl)pseudouridine
1-(4-Trifluoromethoxybenzyl)pseudouridine
1-(4-Trifluoromethylbenzyl)pseudouridine
1-(5-Amino-pentyl)pseudouridine
1-(6-Amino-hexyl)pseudouridine
1,6-Dimethyl-pseudouridine
1-[3-(2-{2-[2-(2-Aminoethoxy)-ethoxy]-ethoxy}-ethoxy)-propionyl]pseudouridine
1-{3-[2-(2-Aminoethoxy)-ethoxy]-propionyl } pseudouridine
1-Acetylpsudouridine
1-Alkyl-6-(1-propynyl)-pseudo-uridine
1-Alkyl-6-(2-propynyl)-pseudo-uridine
1-Alkyl-6-allyl-pseudo-uridine
1-Alkyl-6-ethynyl-pseudo-uridine
1 -Alkyl-6-homoallyl-pseudo-uridine
1-Alkyl-6-vinyl-pseudo-uridine
1-Allylpsudouridine
1-Aminomethyl-pseudo-uridine
1-Benzoylpsudouridine
1-Benzyloxymethylpsudouridine
1-Benzyl-pseudo-uridine
1-Biotinyl-PEG2-pseudouridine
1-Biotinylpsudouridine
1-Butyl-pseudo-uridine
1-Cyanomethylpsudouridine
1-Cyclobutylmethyl-pseudo-uridine
1-Cyclobutyl-pseudo-uridine
1-Cycloheptylmethyl-pseudo-uridine
1-Cycloheptyl-pseudo-uridine
1-Cyclohexylmethyl-pseudo-uridine

Modification
1-Cyclohexyl-pseudo-uridine
1-Cyclooctylmethyl-pseudo-uridine
1-Cyclooctyl-pseudo-uridine
1-Cyclopentylmethyl-pseudo-uridine
1-Cyclopentyl-pseudo-uridine
1-Cyclopropylmethyl-pseudo-uridine
1-Cyclopropyl-pseudo-uridine
1-Ethyl-pseudo-uridine
1-Hexyl-pseudo-uridine
1-Homoallylpseudouridine
1-Hydroxymethylpseudouridine
1-iso-propyl-pseudo-uridine
1-Me-2-thio-pseudo-uridine
1-Me-4-thio-pseudo-uridine
1-Me-alpha-thio-pseudo-uridine
1-Methanesulfonylmethylpseudouridine
1-Methoxymethylpseudouridine uridine
1-Methyl-6-(2,2,2-Trifluoroethyl)pseudo-uridine
1-Methyl-6-(4-morpholino)-pseudo-uridine
1-Methyl-6-(4-thiomorpholino)-pseudo-uridine
1-Methyl-6-(substituted phenyl)pseudo-uridine
1-Methyl-6-amino-pseudo-uridine
1-Methyl-6-azido-pseudo-uridine
1-Methyl-6-bromo-pseudo-uridine
1-Methyl-6-butyl-pseudo-uridine
1-Methyl-6-chloro-pseudo-uridine
1-Methyl-6-cyano-pseudo-uridine
1-Methyl-6-dimethylamino-pseudo-uridine
1-Methyl-6-ethoxy-pseudo-uridine

Modification
1-Methyl-6-ethylcarboxylate-pseudo-uridine
1-Methyl-6-ethyl-pseudo-uridine
1-Methyl-6-fluoro-pseudo-uridine
1-Methyl-6-formyl-pseudo-uridine
1-Methyl-6-hydroxyamino-pseudo-uridine
1-Methyl-6-hydroxy-pseudo-uridine
1-Methyl-6-iodo-pseudo-uridine
1-Methyl-6-iso-propyl-pseudo-uridine
1-Methyl-6-methoxy-pseudo-uridine
1-Methyl-6-methylamino-pseudo-uridine
1-Methyl-6-phenyl-pseudo-uridine
1-Methyl-6-propyl-pseudo-uridine
1-Methyl-6-tert-butyl-pseudo-uridine
1-Methyl-6-trifluoromethoxy-pseudo-uridine
1-Methyl-6-trifluoromethyl-pseudo-uridine
1-Morpholinomethylpseudouridine
1-Pentyl-pseudo-uridineuridine
1-Phenyl-pseudo-uridine
1-Pivaloylpseudouridine
1-Propargylpseudouridine
1-Propyl-pseudo-uridine
1-propynyl-pseudouridine
1-p-tolyl-pseudo-uridine
1-tert-Butyl-pseudo-uridine
1-Thiomethoxymethylpseudouridine
1-Thiomorpholinomethylpseudouridine
1-Trifluoroacetylpsudouridine
1-Trifluoromethyl-pseudouridine
1-Vinylpseudouridine

Modification
2,2'-anhydro-uridine
2' -bromo-deoxyuridine
2' -F-5-Methyl-2'-deoxy-uridine
2' -OMe-5-Me-uridine
2' -OMe-pseudouridine
2' -alpha-Ethynyluridine
2' -alpha-Trifluoromethyluridine
2' -beta-Ethynyluridine
2' -beta-Trifluoromethyluridine
2' -Deoxy-2',2'-difluorouridine
2' -Deoxy-2'-a-mercaptopuridine
2' -Deoxy-2'-alpha-thiomethoxyuridine
2' -Deoxy-2'-beta-aminouridine
2' -Deoxy-2'-beta-azidouridine
2' -Deoxy-2'-beta-bromouridine
2' -Deoxy-2'-beta-chlorouridine
2' -Deoxy-2'-beta-fluorouridine
2' -Deoxy-2'-beta-iodouridine
2' -Deoxy-2'-beta-mercaptopuridine
2' -Deoxy-2'-beta-thiomethoxyuridine
2-methoxy-4-thio-uridine
2-methoxyuridine
2' -O-Methyl-5-(1-propynyl)uridine
3-Alkyl-pseudo-uridine
4' -Azidouridine
4'-Carbocyclic uridine
4' -Ethynyluridine
5-(1-Propynyl)ara-uridine
5-(2-Furanyl)uridine
5-Cyanouridine
5-Dimethylaminouridine

Modification
5'-Homo-uridine
5-iodo-2'-fluoro-deoxyuridine
5-Phenylethynyluridine
5-Trideuteromethyl-6-deuterouridine
5-Trifluoromethyl-Uridine
5-Vinylarauridine
6-(2,2,2-Trifluoroethyl)-pseudo-uridine
6-(4-Morpholino)-pseudo-uridine
6-(4-Thiomorpholino)-pseudo-uridine
6-(Substituted-Phenyl)-pseudo-uridine
6-Amino-pseudo-uridine
6-Azido-pseudo-uridine
6-Bromo-pseudo-uridine
6-Butyl-pseudo-uridine
6-Chloro-pseudo-uridine
6-Cyano-pseudo-uridine
6-Dimethylamino-pseudo-uridine
6-Ethoxy-pseudo-uridine
6-Ethylcarboxylate-pseudo-uridine
6-Ethyl-pseudo-uridine
6-Fluoro-pseudo-uridine
6-Formyl-pseudo-uridine
6-Hydroxyamino-pseudo-uridine
6-Hydroxy-pseudo-uridine
6-Iodo-pseudo-uridine
6-iso-Propyl-pseudo-uridine
6-Methoxy-pseudo-uridine
6-Methylamino-pseudo-uridine
6-Methyl-pseudo-uridine
6-Phenyl-pseudo-uridine
6-Phenyl-pseudo-uridine

Modification
6-Propyl-pseudo-uridine
6-tert-Butyl-pseudo- uridine
6-Trifluoromethoxy-pseudo-uridine
6-Trifluoromethyl-pseudo-uridine
alpha-thio-pseudo-uridine
Pseudouridine 1-(4-methylbenzenesulfonic acid)
Pseudouridine 1-(4-methylbenzoic acid) TP
Pseudouridine 1-[3-(2-ethoxy)]propionic acid
Pseudouridine 1-[3-{2-(2-[2-(2-ethoxy)-ethoxy]-ethoxy)-ethoxy}]propionic acid
Pseudouridine 1-[3-{2-(2-[2-{2(2-ethoxy)-ethoxy}-ethoxy]-ethoxy)-ethoxy}]propionic acid

Modification
Pseudouridine 1-[3-{2-(2-[2-ethoxy]-ethoxy)-ethoxy}]propionic acid
Pseudouridine 1-[3-{2-(2-ethoxy)-ethoxy}]propionic acid
Pseudouridine 1-methylphosphonic acid
Pseudouridine TP 1-methylphosphonic acid diethyl ester
Pseudo-uridine-N1-3-propionic acid
Pseudo-uridine-N1-4-butanoic acid
Pseudo-uridine-N 1-5-pentanoic acid
Pseudo-uridine-N1-6-hexanoic acid
Pseudo-uridine-N1-7-heptanoic acid
Pseudo-uridine-N1-methyl-p-benzoic acid
Pseudo-uridine-N1-p-benzoic acid

In an embodiment, a TREM, a TREM core fragment or a TREM fragment described herein comprises a modification provided in Table 11, or a combination thereof. The
5 modifications provided in Table 6 occur naturally in RNAs, and are used herein on a synthetic TREM, a TREM core fragment or a TREM fragment at a position that does not occur in nature.

Table 11: Additional exemplary modifications

Modification
2-methylthio-N6-(cis-hydroxyisopentenyl)adenosine
2-methylthio-N6-methyladenosine
2-methylthio-N6-threonyl carbamoyl adenosine
N6-glycyl carbamoyl adenosine
N6-isopentenyladenosine
N6-methyladenosine
N6-threonyl carbamoyl adenosine
1,2'-O-dimethyladenosine

Modification
1-methyladenosine
2'-O-methyladenosine
2'-O-ribosyladenosine (phosphate)
2-methyladenosine
2-methylthio-N6 isopentenyladenosine
2-methylthio-N6-hydroxynorvalyl carbamoyl adenosine
2'-O-methyladenosine
2'-O-ribosyladenosine (phosphate)
isopentenyl adenosine

Modification
N6-(cis-hydroxyisopentenyl)adenosine
N6,2'-O-dimethyladenosine
N6,2'-O-dimethyladenosine
N6,N6,2'-O-trimethyladenosine
N6,N6-dimethyladenosine
N6-acetyladenosine
N6-hydroxynorvalylcarbamoyl-adenosine
N6-methyl-N6-threonylcarbamoyl-adenosine
2-methyladenosine
2-methylthio-N ⁶ -isopentenyladenosine
2-thiocytidine
3-methylcytidine
5-formylcytidine
5-hydroxymethylcytidine
5-methylcytidine
N4-acetylcytidine
2'-O-methylcytidine
2'-O-methylcytidine
5,2'-O-dimethylcytidine
5-formyl-2'-O-methylcytidine
lysine
N4,2'-O-dimethylcytidine
N4-acetyl-2'-O-methylcytidine
N4-methylcytidine
N4,N4-Dimethyl-2'-OMe-Cytidine
7-methylguanosine
N2,2'-O-dimethylguanosine
N2-methylguanosine
wyosme
1,2'-O-dimethylguanosine
1-methylguanosine
2'-O-methylguanosine

Modification
2'-O-ribosylguanosine (phosphate)
2'-O-methylguanosine
2'-O-ribosylguanosine (phosphate)
7-aminomethyl-7-deazaguanosine
7-cyano-7-deazaguanosine
archaeosine
methylwyosine
N2,7-dimethylguanosine
N2,N2,2'-O-trimethylguanosine
N2,N2,7-trimethylguanosine
N2,N2-dimethylguanosine
N2, 7,2'-O-trimethylguanosine
1-methylinosine
mosme
1,2'-O-dimethylinosine
2'-O-methylinosine
2'-O-methylinosine
epoxyqueuosine
galactosyl-queuosine
mannosyl-queuosine
2'-O-methyluridine
2-thiouridine
3-methyluridine
5-carboxymethyluridine
5-hydroxyuridine
5-methyluridine
5-taurinomethyl-2-thiouridine
5-taurinomethyluridine
dihydrouridine
pseudouridine
(3-(3-amino-3-carboxypropyl)uridine

Modification
1-methyl-3-(3-amino-5-carboxypropyl)pseudouridine
1-methylpseudouridine
1-methyl-pseudouridine
2'-O-methyluridine
2'-O-methylpseudouridine
2' -O-methyluridine
2-thio-2'-O-methyluridine
3-(3-amino-3-carboxypropyl)uridine
3,2'-O-dimethyluridine
3-Methyl-pseudo-Uridine
4-thiouridine
5-(carboxyhydroxymethyl)uridine
5-(carboxyhydroxymethyl)uridine methyl ester
5,2'-O-dimethyluridine
5,6-dihydro-uridine
5-aminomethyl-2-thiouridine
5-carbamoylmethyl-2'-O-methyluridine
5-carbamoylmethyluridine
5-carboxyhydroxymethyluridine
5-carboxyhydroxymethyluridine methyl ester
5-carboxymethylaminomethyl-2'-O-methyluridine
5-carboxymethylaminomethyl-2-thiouridine
5-carboxymethylaminomethyl-2-thiouridine
5-carboxymethylaminomethyluridine
5-carboxymethylaminomethyluridine

Modification
5-Carbamoylmethyluridine
5-methoxycarbonylmethyl-2'-O-methyluridine
5-methoxycarbonylmethyl-2-thiouridine
5-methoxycarbonylmethyluridine
5-methoxyuridine
5-methyl-2-thiouridine
5-methylaminomethyl-2-selenouridine
5-methylaminomethyl-2-thiouridine
5-methylaminomethyluridine
5-Methyldihydrouridine
5-Oxyacetic acid- Uridine
5-Oxyacetic acid-methyl ester-Uridine
5-methyl-pseudo-uridine
uridine 5-oxyacetic acid
uridine 5-oxyacetic acid methyl ester
3-(3-Amino-3-carboxypropyl)-Uridine
5-(iso-Pentenylaminomethyl)- 2-thiouridine
5-(iso-Pentenylaminomethyl)-2'-O-methyluridine
5-(iso-Pentenylaminomethyl)uridine
wybutosine
hydroxywybutosine
isowyosme
peroxywybutosine
undermodified hydroxywybutosine
4-demethylwyosine
altriol

In an embodiment, a TREM, a TREM core fragment or a TREM fragment described herein comprises a non-naturally occurring modification provided in Table 12, or a combination thereof.

Table 12: Additional exemplary non-naturally occurring modifications

Modification	Modification
2,6-(diamino)purine	6-(azo)thymine
1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl	6-(methyl)-7-(aza)indolyl
1,3-(diaza)-2-(oxo)-phenthiazin-1-yl	6-chloro-purine
1,3-(diazia)-2-(oxo)-phenoxazin-1-yl	6-phenyl-pyrrolo-pyrimidin-2-on-3-yl
1,3,5-(triazia)-2,6-(dioxo)-naphthalene	7-(aminoalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)- phenthiazin-1-yl
2 (amino)purine	7-(aminoalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl
2,4,5-(trimethyl)phenyl	7-(aminoalkylhydroxy)-1,3-(diazia)-2-(oxo)-phenoxazin-1-yl
2' methyl, 2' amino, 2' azido, 2' fluoro-cytidine	7-(aminoalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenthiazin- 1-yl
2' methyl, 2' amino, 2' azido, 2' fluoro-adenine	7-(aminoalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenoxazin-1- yl
2' methyl, 2' amino, 2' azido, 2' fluoro-uridine	7-(aminoalkylhydroxy)-1,3-(diaza)-2-(oxo)-phenoxazin-1- yl
2' -amino-2' -deoxyribose	7-(aza)indolyl
2-amino-6-Chloro-purine	7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)- phenoxazinl-yl
2-aza-inosinyl	7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)- phenthiazin-1-yl
2' -azido-2' -deoxyribose	7-(guanidiniumalkylhydroxy)-1-(aza)-2-(thio)-3-(aza)- phenoxazin-1-yl
2' fluoro-2' -deoxyribose	7-(guanidiniumalkylhydroxy)-1,3-(diaza)-2-(oxo)- phenoxazin-1-yl
2' -fluoro-modified bases	7-(guanidiniumalkyl-hydroxy)-1,3-(diaza)-2-(oxo)- phenthiazin-1-yl
2' -O-methyl-ribose	7-(guanidiniumalkylhydroxy)-1,3-(diaza)-2-(oxo)- phenoxazin-1-yl
2-oxo-7-aminopyridopyrimidin-3-yl	7-(propynyl)isocarbostyryl
2-oxo-pyridopyrimidine-3-yl	7-(propynyl)isocarbostyryl, propynyl-7-(aza)indolyl
2-pyridinone	7-deaza-inosinyl
3 nitropyrrole	7-substituted 1-(aza)-2-(thio)-3-(aza)-phenoxazin-1-yl
3-(methyl)-7-(propynyl)isocarbostyryl	7-substituted 1,3-(diazia)-2-(oxo)-phenoxazin-1-yl
3-(methyl)isocarbostyryl	9-(methyl)-imidizopyridinyl
4-(fluoro)-6-(methyl)benzimidazole	
4-(methyl)benzimidazole	
4-(methyl)indolyl	
4,6-(dimethyl)indolyl	
5 nitroindole	
5 substituted pyrimidines	
5-(methyl)isocarbostyryl	
5-nitroindole	
6-(aza)pyrimidine	

Modification
aminoindolyl
anthracenyl
bis-ortho-(aminoalkylhydroxy)-6-phenyl-pyrrolo- nvrimidin-2-on-3-yl
bis-ortho-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl
difluorotolyl
hypoxanthine
imidizopyridinyl
inosinyl
isocarbostyrilyl
isoguanosine
N2-substituted purines
N6-methyl-2-amino-purine
N6-substituted purines
N-alkylated derivative
naphthalenyl
nitrobenzimidazolyl
nitroimidazolyl
nitroindazolyl
nitropyrazolyl
nubularine
O6-substituted purines
O-alkylated derivative
ortho-(aminoalkylhydroxy)-6-phenyl-pyrrolo-pyrimidin-2- on-3-yl
ortho-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl
Oxoformycin TP
para-(aminoalkylhydroxy)-6-phenyl-pyrrolo-pyrimidin-2- on-3-yl
para-substituted-6-phenyl-pyrrolo-pyrimidin-2-on-3-yl

Modification
pentacenyl
phenanthracenyl
phenyl
propynyl-7-(aza)indolyl
pyrenyl
pyridopyrimidin-3-yl
pyridopyrimidin-3-yl, 2-oxo-7-amino-pyridopyrimidin-3- yl
pyrrolo-pyrimidin-2-on-3-yl
pyrrolopyrimidinyl
pyrrolopyrizinyl
stilbenzyl
substituted 1,2,4-triazoles
tetracenyl
tubercidine
xanthine
Xanthosine
2-thio-zebularine
5-aza-2-thio-zebularine
7-deaza-2-amino-purine
pyridin-4-one ribonucleoside
2-Amino-riboside
Formycin A
Formycin B
Pyrrolosine
2' -OH-ara-adenosine
2' -OH-ara-cytidine
2' -OH-ara-uridine
2' -OH-ara-guanosine
5-(2-carbomethoxyvinyl)uridine
N6-(19-Amino-pentaoxonadecyl)adenosine

In an embodiment, a TREM, a TREM core fragment or a TREM fragment described herein comprises a non-naturally occurring modification provided in Table 13, or a combination thereof.

Table 13: Exemplary backbone modifications

Modification
3'-alkylene phosphonates
3'-amino phosphoramidate
alkene containing backbones
aminoalkylphosphoramidates
aminoalkylphosphotriesters
boranophosphates
-CH ₂ -O-N(CH ₃)-CH ₂ -
-CH ₂ -N(CH ₃)-N(CH ₃)-CH ₂ -
-CH ₂ -NH-CH ₂ -
chiral phosphonates
chiral phosphorothioates
formacetyl and thioformacetyl backbones
methylene (methylimino)
methylene formacetyl and thioformacetyl backbones
methyleneimino and methylenehydrazino backbones
morpholino linkages
-N(CH ₃)-CH ₂ -CH ₂ -
oligonucleosides with heteroatom internucleoside linkage
phosphinates
phosphoramidates
phosphorodithioates
phosphorothioate internucleoside linkages
phosphorothioates
phosphotriesters
Peptide nucleic acid (PNA)
siloxane backbones
sulfamate backbones
Sulfide, sulfoxide, and sulfone backbones
sulfonate and sulfonamide backbones
thionoalkylphosphonates
thionoalkylphosphotriesters
thionophosphoramidates

Modification
methylphosphonates
phosphonoacetates
Phosphorothioate
Constrained nucleic acid (CNA)
2'-O-methyl
2'-O-methoxyethyl (MOE)
2' Fluoro
Locked nucleic acid (LNA)
(S)-constrained ethyl (<i>cEt</i>)
Fluoro hexitol nucleic acid (FHNA)
5'-phosphorothioate
Phosphorodiamidate Morpholino Oligomer (PMO)
Tricyclo-DNA (tcDNA)
(S) 5'-C-methyl
(E)-vinylphosphonate
Methyl phosphonate
(S) 5'-C-methyl with phosphate
(R) 5'-C-methyl with phosphate
DNA
(R) 5'-C-methyl
GNA (glycol nucleic acid)
alkyl phosphonates
Phosphorothioate
Constrained nucleic acid (CNA)
2'-O-methyl
2'-O-methoxyethyl (MOE)
2' Fluoro
Locked nucleic acid (LNA)
(S)-constrained ethyl (<i>cEt</i>)
Fluoro hexitol nucleic acid (FHNA)
5'-phosphorothioate
Phosphorodiamidate Morpholino Oligomer (PMO)
Tricyclo-DNA (tcDNA)
(S) 5'-C-methyl
(E)-vinylphosphonate
Methyl phosphonate

Modification
(S) 5'-C-methyl with phosphate
(R) 5'-C-methyl with phosphate
DNA
(R) 5'-C-methyl

Modification
GNA (glycol nucleic acid)
alkyl phosphonates

- In an embodiment, a TREM, a TREM core fragment or a TREM fragment described
 5 herein comprises a non-naturally occurring modification provided in Table 14, or a combination thereof.

Table 14: Exemplary non-naturally occurring backbone modifications

Name of synthetic backbone modifications
Phosphorothioate
Constrained nucleic acid (CNA)
2' O'-methylation
2'-O-methoxyethylribose (MOE)
2'-Fluoro
Locked nucleic acid (LNA)
(S)-constrained ethyl (cEt)
Fluoro hexitol nucleic acid (FHNA)
5'-phosphorothioate
Phosphorodiamidate Morpholino Oligomer (PMO)
Tricyclo-DNA (tcDNA)
(S) 5'-C-methyl
(E)-vinylphosphonate
Methyl phosphonate
(S) 5'-C-methyl with phosphate

TREM, TREM core fragment and TREM fragment fusions

In an embodiment, a TREM, a TREM core fragment or a TREM fragment disclosed herein comprises an additional moiety, *e.g.*, a fusion moiety. In an embodiment, the fusion moiety can be used for purification, to alter folding of the TREM, TREM core fragment or TREM fragment, or as a targeting moiety. In an embodiment, the fusion moiety can comprise a tag, a linker, can be cleavable or can include a binding site for an enzyme. In an embodiment, the fusion moiety can be disposed at the N terminal of the TREM or at the C terminal of the TREM, TREM core fragment or TREM fragment. In an embodiment, the fusion moiety can be encoded by the same or different nucleic acid molecule that encodes the TREM, TREM core fragment or TREM fragment.

TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises a consensus sequence provided herein.

In an embodiment, a TREM disclosed herein comprises a consensus sequence of Formula I *zzz*, wherein *zzz* indicates any of the twenty amino acids and Formula I corresponds to all species.

In an embodiment, a TREM disclosed herein comprises a consensus sequence of Formula II *zzz*, wherein *zzz* indicates any of the twenty amino acids and Formula II corresponds to mammals.

In an embodiment, a TREM disclosed herein comprises a consensus sequence of Formula III *zzz*, wherein *zzz* indicates any of the twenty amino acids and Formula III corresponds to humans.

In an embodiment, *zzz* indicates any of the twenty amino acids: alanine, arginine, asparagine, aspartate, cysteine, glutamine, glutamate, glycine, histidine, isoleucine, methionine, leucine, lysine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, or valine.

In an embodiment, a TREM disclosed herein comprises a property selected from the following:

- a) under physiological conditions residue R₀ forms a linker region, *e.g.*, a Linker 1 region;
- b) under physiological conditions residues R₁-R₂-R₃-R₄-R₅-R₆-R₇ and residues R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁ form a stem region, *e.g.*, an AStD stem region;

c) under physiological conditions residues R₈-R₉ forms a linker region, *e.g.*, a Linker 2 region;

d) under physiological conditions residues -R₁₀-R₁₁-R₁₂-R₁₃-R₁₄ R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈ form a stem-loop region, *e.g.*, a D arm Region;

5 e) under physiological conditions residue -R₂₉ forms a linker region, *e.g.*, a Linker 3 Region;

f) under physiological conditions residues -R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-R₄₆ form a stem-loop region, *e.g.*, an AC arm region;

10 g) under physiological conditions residue -[R₄₇]_x comprises a variable region, *e.g.*, as described herein;

h) under physiological conditions residues -R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄ form a stem-loop region, *e.g.*, a T arm Region; or

i) under physiological conditions residue R₇₂ forms a linker region, *e.g.*, a Linker 4 region.

15

Alanine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{ALA} (SEQ ID NO: 562),

20 R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Ala is:

25 R₀= absent;

R₁₄, R₅₇=are independently A or absent;

R₂₆= A, C, G or absent;

R₅, R₆, R₁₅, R₁₆, R₂₁, R₃₀, R₃₁, R₃₂, R₃₄, R₃₇, R₄₁, R₄₂, R₄₃, R₄₄, R₄₅, R₄₈, R₄₉, R₅₀, R₅₈, R₅₉, R₆₃, R₆₄, R₆₆, R₆₇= are independently N or absent;

30 R₁₁, R₃₅, R₆₅= are independently A, C, U or absent;

R₁, R₉, R₂₀, R₃₈, R₄₀, R₅₁, R₅₂, R₅₆= are independently A, G or absent;

- R₇, R₂₂, R₂₅, R₂₇, R₂₉, R₄₆, R₅₃, R₇₂= are independently A, G, U or absent;
 R₂₄, R₆₉= are independently A, U or absent;
 R₇₀, R₇₁= are independently C or absent;
 R₃, R₄= are independently C, G or absent;
 5 R₁₂, R₃₃, R₃₆, R₆₂, R₆₈= are independently C, G, U or absent;
 R₁₃, R₁₇, R₂₈, R₃₉, R₅₅, R₆₀, R₆₁= are independently C, U or absent;
 R₁₀, R₁₉, R₂₃= are independently G or absent;
 R₂= G, U or absent;
 R₈, R₁₈, R₅₄= are independently U or absent;
 10 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 15 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 20 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{ALA} (SEQ ID NO: 563),

- R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 25 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Ala is:

- R₀, R₁₈= are absent;
 30 R₁₄, R₂₄, R₅₇= are independently A or absent;

- R₁₅, R₂₆, R₆₄= are independently A, C, G or absent;
 R₁₆, R₃₁, R₅₀, R₅₉= are independently N or absent;
 R₁₁, R₃₂, R₃₇, R₄₁, R₄₃, R₄₅, R₄₉, R₆₅, R₆₆= are independently A, C, U or absent;
 R₁, R₅, R₉, R₂₅, R₂₇, R₃₈, R₄₀, R₄₆, R₅₁, R₅₆= are independently A, G or absent;
 5 R₇, R₂₂, R₂₉, R₄₂, R₄₄, R₅₃, R₆₃, R₇₂= are independently A, G, U or absent;
 R₆, R₃₅, R₆₉= are independently A, U or absent;
 R₅₅, R₆₀, R₇₀, R₇₁= are independently C or absent;
 R₃= C, G or absent;
 R₁₂, R₃₆, R₄₈= are independently C, G, U or absent;
 10 R₁₃, R₁₇, R₂₈, R₃₀, R₃₄, R₃₉, R₅₈, R₆₁, R₆₂, R₆₇, R₆₈= are independently C, U or absent;
 R₄, R₁₀, R₁₉, R₂₀, R₂₃, R₅₂= are independently G or absent;
 R₂, R₈, R₃₃= are independently G, U or absent;
 R₂₁, R₅₄= are independently U or absent;
 [R₄₇]_x = N or absent;
 15 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 20 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 residues are N; or no more than 20 residues are absent.

25

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{ALA} (SEQ ID NO: 564),

R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-

R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Ala is:

- R₀, R₁₈= are absent;
- 5 R₁₄, R₂₄, R₅₇, R₇₂=are independently A or absent;
- R₁₅, R₂₆, R₆₄= are independently A, C, G or absent;
- R₁₆, R₃₁, R₅₀= are independently N or absent;
- R₁₁, R₃₂, R₃₇, R₄₁, R₄₃, R₄₅, R₄₉, R₆₅, R₆₆= are independently A, C, U or absent;
- R₅, R₉, R₂₅, R₂₇, R₃₈, R₄₀, R₄₆, R₅₁, R₅₆= are independently A, G or absent;
- 10 R₇, R₂₂, R₂₉, R₄₂, R₄₄, R₅₃, R₆₃= are independently A, G, U or absent;
- R₆, R₃₅= are independently A, U or absent;
- R₅₅, R₆₀, R₆₁, R₇₀, R₇₁= are independently C or absent;
- R₁₂, R₄₈, R₅₉= are independently C, G, U or absent;
- R₁₃, R₁₇, R₂₈, R₃₀, R₃₄, R₃₉, R₅₈, R₆₂, R₆₇, R₆₈= are independently C, U or absent;
- 15 R₁, R₂, R₃, R₄, R₁₀, R₁₉, R₂₀, R₂₃, R₅₂= are independently G or absent;
- R₃₃, R₃₆= are independently G, U or absent;
- R₈, R₂₁, R₅₄, R₆₉= are independently U or absent;
- [R₄₇]_x = N or absent;
- wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
- 20 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
- x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
- x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
- x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
- 25 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
- x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
- provided that the TREM has one or both of the following properties: no more than 15% of the
residues are N; or no more than 20 residues are absent.

Arginine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{ARG} (SEQ ID NO: 565),

5 R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Arg is:

10 R₅₇=A or absent;
R₉,R₂₇=are independently A,C,G or absent;
R₁,R₂,R₃,R₄,R₅,R₆,R₇,R₁₁,R₁₂,R₁₆,R₂₁,R₂₂,R₂₃,R₂₅,R₂₆,R₂₉,R₃₀,R₃₁,R₃₂,R₃₃,R₃₄,R₃₇,R₄₂,R₄₄,R₄₅,
R₄₆,R₄₈,R₄₉,R₅₀,R₅₁,R₅₈,R₆₂,R₆₃,R₆₄,R₆₅,R₆₆,R₆₇,R₆₈,R₆₉,R₇₀,R₇₁=are independently N or
absent;
15 R₁₃,R₁₇,R₄₁=are independently A,C,U or absent;
R₁₉,R₂₀,R₂₄,R₄₀,R₅₆=are independently A,G or absent;
R₁₄,R₁₅,R₇₂=are independently A,G,U or absent;
R₁₈= A,U or absent;
R₃₈= C or absent;
20 R₃₅,R₄₃,R₆₁=are independently C,G,U or absent;
R₂₈,R₅₅,R₅₉,R₆₀=are independently C,U or absent;
R₀,R₁₀,R₅₂=are independently G or absent;
R₈,R₃₉=are independently G,U or absent;
R₃₆,R₅₃,R₅₄=are independently U or absent;
25 [R₄₇]_x = N or absent;
wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
30 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,

x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

5

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{ARG} (SEQ ID NO: 566),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
10 R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Arg is:

R₁₈= absent;

R₂₄,R₅₇=are independently A or absent;

15 R₄₁= A,C or absent;

R₃,R₇,R₃₄,R₅₀=are independently A,C,G or absent;

R₂,R₅,R₆,R₁₂,R₂₆,R₃₂,R₃₇,R₄₄,R₅₈,R₆₆,R₆₇,R₆₈,R₇₀=are independently N or absent;

R₄₉,R₇₁=are independently A,C,U or absent;

R₁,R₁₅,R₁₉,R₂₅,R₂₇,R₄₀,R₄₅,R₄₆,R₅₆,R₇₂=are independently A,G or absent;

20 R₁₄,R₂₉,R₆₃=are independently A,G,U or absent;

R₁₆,R₂₁=are independently A,U or absent;

R₃₈,R₆₁=are independently C or absent;

R₃₃,R₄₈=are independently C,G or absent;

R₄,R₉,R₁₁,R₄₃,R₆₂,R₆₄,R₆₉=are independently C,G,U or absent;

25 R₁₃,R₂₂,R₂₈,R₃₀,R₃₁,R₃₅,R₅₅,R₆₀,R₆₅=are independently C,U or absent;

R₀,R₁₀,R₂₀,R₂₃,R₅₁,R₅₂=are independently G or absent;

R₈,R₃₉,R₄₂=are independently G,U or absent;

R₁₇,R₃₆,R₅₃,R₅₄,R₅₉=are independently U or absent;

[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 5 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the
 10 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III ARG (SEQ ID NO: 567),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 15 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
 R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Arg is:

R₁₈=is absent;
 20 R₁₅,R₂₁,R₂₄,R₄₁,R₅₇=are independently A or absent;
 R₃₄,R₄₄=are independently A,C or absent;
 R₃,R₅,R₅₈=are independently A,C,G or absent;
 R₂,R₆,R₆₆,R₇₀=are independently N or absent;
 R₃₇,R₄₉=are independently A,C,U or absent;
 25 R₁,R₂₅,R₂₉,R₄₀,R₄₅,R₄₆,R₅₀=are independently A,G or absent;
 R₁₄,R₆₃,R₆₈=are independently A,G,U or absent;
 R₁₆= A,U or absent;
 R₃₈,R₆₁=are independently C or absent;
 R₇,R₁₁,R₁₂,R₂₆,R₄₈=are independently C,G or absent;
 30 R₆₄,R₆₇,R₆₉=are independently C,G,U or absent;

R₄,R₁₃,R₂₂,R₂₈,R₃₀,R₃₁,R₃₅,R₄₃,R₅₅,R₆₀,R₆₂,R₆₅,R₇₁=are independently C,U or absent;

R₀,R₁₀,R₁₉,R₂₀,R₂₃,R₂₇,R₃₃,R₅₁,R₅₂,R₅₆,R₇₂=are independently G or absent;

R₈,R₉,R₃₂,R₃₉,R₄₂=are independently G,U or absent;

R₁₇,R₃₆,R₅₃,R₅₄,R₅₉=are independently U or absent;

5 [R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the
15 residues are N; or no more than 20 residues are absent.

Asparagine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{ASN} (SEQ ID NO: 568),

20 R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Asn is:

R₀,R₁₈=are absent;

25 R₄₁= A or absent;

R₁₄,R₄₈,R₅₆=are independently A,C,G or absent;

R₂,R₄,R₅,R₆,R₁₂,R₁₇,R₂₆,R₂₉,R₃₀,R₃₁,R₄₄,R₄₅,R₄₆,R₄₉,R₅₀,R₅₈,R₆₂,R₆₃,R₆₅,R₆₆,R₆₇,R₆₈,R₇₀,R₇₁=are independently N or absent;

R₁₁,R₁₃,R₂₂,R₄₂,R₅₅,R₅₉=are independently A,C,U or absent;

30 R₉,R₁₅,R₂₄,R₂₇,R₃₄,R₃₇,R₅₁,R₇₂=are independently A,G or absent;

R₁,R₇,R₂₅,R₆₉=are independently A,G,U or absent;

- R_{40}, R_{57} are independently A, U or absent;
 R_{60} = C or absent;
 R_{33} = C, G or absent;
 $R_{21}, R_{32}, R_{43}, R_{64}$ are independently C, G, U or absent;
5 $R_3, R_{16}, R_{28}, R_{35}, R_{36}, R_{61}$ are independently C, U or absent;
 $R_{10}, R_{19}, R_{20}, R_{52}$ are independently G or absent;
 R_{54} = G, U or absent;
 $R_8, R_{23}, R_{38}, R_{39}, R_{53}$ are independently U or absent;
 $[R_{47}]_x$ = N or absent;
10 wherein, *e.g.*, $x=1-271$ (*e.g.*, $x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,$
 $x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,$
 $x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,$
 $x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-$
 $271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,$
15 $x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,$
 $x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,$
 $x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, \text{ or } x=271$),
provided that the TREM has one or both of the following properties: no more than 15% of the
residues are N; or no more than 20 residues are absent.
20 In an embodiment, a TREM disclosed herein comprises the sequence of Formula II _{ASN}
(SEQ ID NO: 569),
 $R_0-R_1-R_2-R_3-R_4-R_5-R_6-R_7-R_8-R_9-R_{10}-R_{11}-R_{12}-R_{13}-R_{14}-R_{15}-R_{16}-R_{17}-R_{18}-R_{19}-R_{20}-R_{21}-R_{22}-$
 $R_{23}-R_{24}-R_{25}-R_{26}-R_{27}-R_{28}-R_{29}-R_{30}-R_{31}-R_{32}-R_{33}-R_{34}-R_{35}-R_{36}-R_{37}-R_{38}-R_{39}-R_{40}-R_{41}-R_{42}-R_{43}-R_{44}-R_{45}-$
 $R_{46}-[R_{47}]_x-R_{48}-R_{49}-R_{50}-R_{51}-R_{52}-R_{53}-R_{54}-R_{55}-R_{56}-R_{57}-R_{58}-R_{59}-R_{60}-R_{61}-R_{62}-R_{63}-R_{64}-R_{65}-R_{66}-R_{67}-$
25 $R_{68}-R_{69}-R_{70}-R_{71}-R_{72}$
wherein R is a ribonucleotide residue and the consensus for Asn is:
 R_0, R_{18} are absent
 $R_{24}, R_{41}, R_{46}, R_{62}$ are independently A or absent;
 R_{59} = A, C or absent;
30 R_{14}, R_{56}, R_{66} are independently A, C, G or absent;

R_{17}, R_{29} are independently N or absent;
 $R_{11}, R_{26}, R_{42}, R_{55}$ are independently A, C, U or absent;
 $R_1, R_9, R_{12}, R_{15}, R_{25}, R_{34}, R_{37}, R_{48}, R_{51}, R_{67}, R_{68}, R_{69}, R_{70}, R_{72}$ are independently A, G or absent;
 R_{44}, R_{45}, R_{58} are independently A, G, U or absent;
5 R_{40}, R_{57} are independently A, U or absent;
 R_5, R_{28}, R_{60} are independently C or absent;
 R_{33}, R_{65} are independently C, G or absent;
 R_{21}, R_{43}, R_{71} are independently C, G, U or absent;
 $R_3, R_6, R_{13}, R_{22}, R_{32}, R_{35}, R_{36}, R_{61}, R_{63}, R_{64}$ are independently C, U or absent;
10 $R_7, R_{10}, R_{19}, R_{20}, R_{27}, R_{49}, R_{52}$ are independently G or absent;
 R_{54} is G, U or absent;
 $R_2, R_4, R_8, R_{16}, R_{23}, R_{30}, R_{31}, R_{38}, R_{39}, R_{50}, R_{53}$ are independently U or absent;
 $[R_{47}]_x$ is N or absent;
wherein, *e.g.*, $x=1-271$ (*e.g.*, $x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,$
15 $x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,$
 $x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,$
 $x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-$
 $271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,$
 $x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,$
20 $x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,$
 $x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, \text{ or } x=271$),
provided that the TREM has one or both of the following properties: no more than 15% of the
residues are N; or no more than 20 residues are absent.

25 In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{ASN}
(SEQ ID NO: 570),

$R_0 - R_1 - R_2 - R_3 - R_4 - R_5 - R_6 - R_7 - R_8 - R_9 - R_{10} - R_{11} - R_{12} - R_{13} - R_{14} - R_{15} - R_{16} - R_{17} - R_{18} - R_{19} - R_{20} - R_{21} - R_{22} -$
 $R_{23} - R_{24} - R_{25} - R_{26} - R_{27} - R_{28} - R_{29} - R_{30} - R_{31} - R_{32} - R_{33} - R_{34} - R_{35} - R_{36} - R_{37} - R_{38} - R_{39} - R_{40} - R_{41} - R_{42} - R_{43} - R_{44} - R_{45} -$
 $R_{46} - [R_{47}]_x - R_{48} - R_{49} - R_{50} - R_{51} - R_{52} - R_{53} - R_{54} - R_{55} - R_{56} - R_{57} - R_{58} - R_{59} - R_{60} - R_{61} - R_{62} - R_{63} - R_{64} - R_{65} - R_{66} - R_{67} -$
 $R_{68} - R_{69} - R_{70} - R_{71} - R_{72}$

30 wherein R is a ribonucleotide residue and the consensus for Asn is:

- R₀,R₁₈=are absent
 R₂₄,R₄₀,R₄₁,R₄₆,R₆₂=are independently A or absent;
 R₅₉= A,C or absent;
 R₁₄,R₅₆,R₆₆=are independently A,C,G or absent;
 5 R₁₁,R₂₆,R₄₂,R₅₅=are independently A,C,U or absent;
 R₁,R₉,R₁₂,R₁₅,R₃₄,R₃₇,R₄₈,R₅₁,R₆₇,R₆₈,R₆₉,R₇₀=are independently A,G or absent;
 R₄₄,R₄₅,R₅₈=are independently A,G,U or absent;
 R₅₇= A,U or absent;
 R₅,R₂₈,R₆₀=are independently C or absent;
 10 R₃₃,R₆₅=are independently C,G or absent;
 R₁₇,R₂₁,R₂₉=are independently C,G,U or absent;
 R₃,R₆,R₁₃,R₂₂,R₃₂,R₃₅,R₃₆,R₄₃,R₆₁,R₆₃,R₆₄,R₇₁=are independently C,U or absent;
 R₇,R₁₀,R₁₉,R₂₀,R₂₅,R₂₇,R₄₉,R₅₂,R₇₂=are independently G or absent;
 R₅₄= G,U or absent;
 15 R₂,R₄,R₈,R₁₆,R₂₃,R₃₀,R₃₁,R₃₈,R₃₉,R₅₀,R₅₃=are independently U or absent;
 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 20 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 25 provided that the TREM has one or both of the following properties: no more than 15% of the
 residues are N; or no more than 20 residues are absent.

Aspartate TREM Consensus sequence

- In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{ASP}
 30 (SEQ ID NO: 571),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

5 wherein R is a ribonucleotide residue and the consensus for Asp is:

R₀=absent

R₂₄,R₇₁=are independently A,C or absent;

R₃₃,R₄₆=are independently A,C,G or absent;

10 R₂,R₃,R₄,R₅,R₆,R₁₂,R₁₆,R₂₂,R₂₆,R₂₉,R₃₁,R₃₂,R₄₄,R₄₈,R₄₉,R₅₈,R₆₃,R₆₄,R₆₆,R₆₇,R₆₈,R₆₉=are
independently N or absent;

R₁₃,R₂₁,R₃₄,R₄₁,R₅₇,R₆₅=are independently A,C,U or absent;

R₉,R₁₀,R₁₄,R₁₅,R₂₀,R₂₇,R₃₇,R₄₀,R₅₁,R₅₆,R₇₂=are independently A,G or absent;

R₇,R₂₅,R₄₂=are independently A,G,U or absent;

R₃₉= C or absent;

15 R₅₀,R₆₂=are independently C,G or absent;

R₃₀,R₄₃,R₄₅,R₅₅,R₇₀=are independently C,G,U or absent;

R₈,R₁₁,R₁₇,R₁₈,R₂₈,R₃₅,R₅₃,R₅₉,R₆₀,R₆₁=are independently C,U or absent;

R₁₉,R₅₂=are independently G or absent;

R_i= G,U or absent;

20 R₂₃,R₃₆,R₃₈,R₅₄=are independently U or absent;

[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
25 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),

provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{ASP} (SEQ ID NO: 572),

5 R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Asp is:

10 R₀,R₁₇,R₁₈,R₂₃=are independently absent;
R₉,R₄₀=are independently A or absent;
R₂₄,R₇₁=are independently A,C or absent;
R₆₇,R₆₈=are independently A,C,G or absent;
R₂,R₆,R₆₆=are independently N or absent;
15 R₅₇,R₆₃=are independently A,C,U or absent;
R₁₀,R₁₄,R₂₇,R₃₃,R₃₇,R₄₄,R₄₆,R₅₁,R₅₆,R₆₄,R₇₂=are independently A,G or absent;
R₇,R₁₂,R₂₆,R₆₅=are independently A,U or absent;
R₃₉,R₆₁,R₆₂=are independently C or absent;
R₃,R₃₁,R₄₅,R₇₀=are independently C,G or absent;
20 R₄,R₅,R₂₉,R₄₃,R₅₅=are independently C,G,U or absent;
R₈,R₁₁,R₁₃,R₃₀,R₃₂,R₃₄,R₃₅,R₄₁,R₄₈,R₅₃,R₅₉,R₆₀=are independently C,U or absent;
R₁₅,R₁₉,R₂₀,R₂₅,R₄₂,R₅₀,R₅₂=are independently G or absent;
R₁,R₂₂,R₄₉,R₅₈,R₆₉=are independently G,U or absent;
R₁₆,R₂₁,R₂₈,R₃₆,R₃₈,R₅₄=are independently U or absent;
25 [R₄₇]_x = N or absent;
wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
30 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,

x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{ASP} (SEQ ID NO: 573),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Asp is:

R₀,R₁₇,R₁₈,R₂₃=are absent
 R₉,R₁₂,R₄₀,R₆₅,R₇₁=are independently A or absent;
 R₂,R₂₄,R₅₇=are independently A,C or absent;
 R₆,R₁₄,R₂₇,R₄₆,R₅₁,R₅₆,R₆₄,R₆₇,R₆₈=are independently A,G or absent;
 R₃,R₃₁,R₃₅,R₃₉,R₆₁,R₆₂=are independently C or absent;
 R₆₆= C,G or absent;
 R₅,R₈,R₂₉,R₃₀,R₃₂,R₃₄,R₄₁,R₄₃,R₄₈,R₅₅,R₅₉,R₆₀,R₆₃=are independently C,U or absent;
 R₁₀,R₁₅,R₁₉,R₂₀,R₂₅,R₃₃,R₃₇,R₄₂,R₄₄,R₄₅,R₄₉,R₅₀,R₅₂,R₆₉,R₇₀,R₇₂=are independently G or absent;
 R₂₂,R₅₈=are independently G,U or absent;
 R₁,R₄,R₇,R₁₁,R₁₃,R₁₆,R₂₁,R₂₆,R₂₈,R₃₆,R₃₈,R₅₃,R₅₄=are independently U or absent;
 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,

x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

5 Cysteine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{CYS} (SEQ ID NO: 574),

10 R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
10 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Cys is:

R₀ =absent
R₁₄,R₃₉,R₅₇=are independently A or absent;
15 R₄₁= A,C or absent;
R₁₀,R₁₅,R₂₇,R₃₃,R₆₂=are independently A,C,G or absent;
R₃,R₄,R₅,R₆,R₁₂,R₁₃,R₁₆,R₂₄,R₂₆,R₂₉,R₃₀,R₃₁,R₃₂,R₃₄,R₄₂,R₄₄,R₄₅,R₄₆,R₄₈,R₄₉,R₅₈,R₆₃,R₆₄,R₆₆,
R₆₇,R₆₈,R₆₉,R₇₀=are independently N or absent;
R₆₅= A,C,U or absent;
20 R₉,R₂₅,R₃₇,R₄₀,R₅₂,R₅₆=are independently A,G or absent;
R₇,R₂₀,R₅₁=are independently A,G,U or absent;
R₁₈,R₃₈,R₅₅=are independently C or absent;
R₂= C, G or absent;
R₂₁,R₂₈,R₄₃,R₅₀=are independently C,G,U or absent;
25 R₁₁,R₂₂,R₂₃,R₃₅,R₃₆,R₅₉,R₆₀,R₆₁,R₇₁,R₇₂=are independently C,U or absent;
R₁,R₁₉=are independently G or absent;
R₁₇= G,U or absent;
R₈,R₅₃,R₅₄=are independently U or absent;
[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 5 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the
 10 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{CYS} (SEQ ID NO: 575),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
 15 R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Cys is:

R₀,R₁₈,R₂₃=are absent;
 R₁₄,R₂₄,R₂₆,R₂₉,R₃₉,R₄₁,R₄₅,R₅₇=are independently A or absent;
 20 R₄₄= A,C or absent;
 R₂₇,R₆₂=are independently A,C,G or absent;
 R₁₆= A,C,G,U or absent;
 R₃₀,R₇₀=are independently A,C,U or absent;
 R₅,R₇,R₉,R₂₅,R₃₄,R₃₇,R₄₀,R₄₆,R₅₂,R₅₆,R₅₈,R₆₆=are independently A,G or absent;
 25 R₂₀,R₅₁=are independently A,G,U or absent;
 R₃₅,R₃₈,R₄₃,R₅₅,R₆₉=are independently C or absent;
 R₂,R₄,R₁₅=are independently C,G or absent;
 R₁₃= C,G,U or absent;
 R₆,R₁₁,R₂₈,R₃₆,R₄₈,R₄₉,R₅₀,R₆₀,R₆₁,R₆₇,R₆₈,R₇₁,R₇₂=are independently C,U or absent;
 30 R₁,R₃,R₁₀,R₁₉,R₃₃,R₆₃=are independently G or absent;

R₈,R₁₇,R₂₁,R₆₄=are independently G,U or absent;

R₁₂,R₂₂,R₃₁,R₃₂,R₄₂,R₅₃,R₅₄,R₆₅=are independently U or absent;

R₅₉= U, or absent;

[R₄₇]_x = N or absent;

5 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

15 In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{CYS} (SEQ ID NO: 576),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Cys is:

R₀,R₁₈,R₂₃=are absent

R₁₄,R₂₄,R₂₆,R₂₉,R₃₄,R₃₉,R₄₁,R₄₅,R₅₇,R₅₈=are independently A or absent;

R₄₄,R₇₀=are independently A,C or absent;

25 R₆₂= A,C,G or absent;

R₁₆= N or absent;

R₅,R₇,R₉,R₂₀,R₄₀,R₄₆,R₅₁,R₅₂,R₅₆,R₆₆=are independently A,G or absent;

R₂₈,R₃₅,R₃₈,R₄₃,R₅₅,R₆₇,R₆₉=are independently C or absent;

R₄,R₁₅=are independently C,G or absent;

30 R₆,R₁₁,R₁₃,R₃₀,R₄₈,R₄₉,R₅₀,R₆₀,R₆₁,R₆₈,R₇₁,R₇₂=are independently C,U or absent;

R₁,R₂,R₃,R₁₀,R₁₉,R₂₅,R₂₇,R₃₃,R₃₇,R₆₃=are independently G or absent;

R₈,R₂₁,R₆₄=are independently G,U or absent;

R₁₂,R₁₇,R₂₂,R₃₁,R₃₂,R₃₆,R₄₂,R₅₃,R₅₄, R₅₉,R₆₅=are independently U or absent;

[R₄₇]_x = N or absent;

- 5 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

15 Glutamine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{GLN} (SEQ ID NO: 577),

- R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
20 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Gln is:

R₀,R₁₈=are absent;

R₁₄,R₂₄,R₅₇=are independently A or absent;

- 25 R₉,R₂₆,R₂₇,R₃₃,R₅₆=are independently A,C,G or absent;

R₂,R₄,R₅,R₆,R₁₂,R₁₃,R₁₆,R₂₁,R₂₂,R₂₅,R₂₉,R₃₀,R₃₁,R₃₂,R₃₄,R₄₁,R₄₂,R₄₄,R₄₅,R₄₆,R₄₈,R₄₉,R₅₀,R₅₈,R₆₂,R₆₃,R₆₆,R₆₇,R₆₈,R₆₉,R₇₀=are independently N or absent;

R₁₇,R₂₃,R₄₃,R₆₅,R₇₁=are independently A,C,U or absent;

R₁₅,R₄₀,R₅₁,R₅₂=are independently A,G or absent;

- 30 R₁,R₇,R₇₂=are independently A,G,U or absent;

R₃,R₁₁,R₃₇,R₆₀,R₆₄=are independently C,G,U or absent;

R₂₈,R₃₅,R₅₅,R₅₉,R₆₁=are independently C,U or absent;

R₁₀,R₁₉,R₂₀=are independently G or absent;

R₃₉= G,U or absent;

R₈,R₃₆,R₃₈,R₅₃,R₅₄=are independently U or absent;

5 [R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the
15 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{GLN} (SEQ ID NO: 578),

R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
20 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Gln is:

R₀,R₁₈,R₂₃=are absent

R₁₄,R₂₄,R₅₇=are independently A or absent;

25 R₁₇,R₇₁=are independently A,C or absent;

R₂₅,R₂₆,R₃₃,R₄₄,R₄₆,R₅₆,R₆₉=are independently A,C,G or absent;

R₄,R₅,R₁₂,R₂₂,R₂₉,R₃₀,R₄₈,R₄₉,R₆₃,R₆₇,R₆₈=are independently N or absent;

R₃₁,R₄₃,R₆₂,R₆₅,R₇₀=are independently A,C,U or absent;

R₁₅,R₂₇,R₃₄,R₄₀,R₄₁,R₅₁,R₅₂=are independently A,G or absent;

30 R₂,R₇,R₂₁,R₄₅,R₅₀,R₅₈,R₆₆,R₇₂=are independently A,G,U or absent;

- $R_3, R_{13}, R_{32}, R_{37}, R_{42}, R_{60}, R_{64}$ are independently C, G, U or absent;
 $R_6, R_{11}, R_{28}, R_{35}, R_{55}, R_{59}, R_{61}$ are independently C, U or absent;
 $R_9, R_{10}, R_{19}, R_{20}$ are independently G or absent;
 R_1, R_{16}, R_{39} are independently G, U or absent;
5 $R_8, R_{36}, R_{38}, R_{53}, R_{54}$ are independently U or absent;
 $[R_{47}]_x = N$ or absent;
wherein, *e.g.*, $x=1-271$ (*e.g.*, $x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,$
 $x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,$
 $x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,$
10 $x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-$
 $271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,$
 $x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,$
 $x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,$
 $x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, \text{ or } x=271$),
15 provided that the TREM has one or both of the following properties: no more than 15% of the
residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{GLN} (SEQ ID NO: 579),

- $R_0-R_1-R_2-R_3-R_4-R_5-R_6-R_7-R_8-R_9-R_{10}-R_{11}-R_{12}-R_{13}-R_{14}-R_{15}-R_{16}-R_{17}-R_{18}-R_{19}-R_{20}-R_{21}-R_{22}-$
20 $R_{23}-R_{24}-R_{25}-R_{26}-R_{27}-R_{28}-R_{29}-R_{30}-R_{31}-R_{32}-R_{33}-R_{34}-R_{35}-R_{36}-R_{37}-R_{38}-R_{39}-R_{40}-R_{41}-R_{42}-R_{43}-R_{44}-R_{45}-$
 $R_{46}-[R_{47}]_x-R_{48}-R_{49}-R_{50}-R_{51}-R_{52}-R_{53}-R_{54}-R_{55}-R_{56}-R_{57}-R_{58}-R_{59}-R_{60}-R_{61}-R_{62}-R_{63}-R_{64}-R_{65}-R_{66}-R_{67}-$
 $R_{68}-R_{69}-R_{70}-R_{71}-R_{72}$

wherein R is a ribonucleotide residue and the consensus for Gln is:

- R_0, R_{18}, R_{23} are absent
25 $R_{14}, R_{24}, R_{41}, R_{57}$ are independently A or absent;
 R_{17}, R_{71} are independently A, C or absent;
 $R_5, R_{25}, R_{26}, R_{46}, R_{56}, R_{69}$ are independently A, C, G or absent;
 $R_4, R_{22}, R_{29}, R_{30}, R_{48}, R_{49}, R_{63}, R_{68}$ are independently N or absent;
 $R_{43}, R_{62}, R_{65}, R_{70}$ are independently A, C, U or absent;
30 $R_{15}, R_{27}, R_{33}, R_{34}, R_{40}, R_{51}, R_{52}$ are independently A, G or absent;

- R₂,R₇,R₁₂,R₄₅,R₅₀,R₅₈,R₆₆=are independently A,G,U or absent;
 R₃₁= A,U or absent;
 R₃₂,R₄₄,R₆₀=are independently C,G or absent;
 R₃,R₁₃,R₃₇,R₄₂,R₆₄,R₆₇=are independently C,G,U or absent;
 5 R₆,R₁₁,R₂₈,R₃₅,R₅₅,R₅₉,R₆₁=are independently C,U or absent;
 R₉,R₁₀,R₁₉,R₂₀=are independently G or absent;
 R₁,R₂₁,R₃₉,R₇₂=are independently G,U or absent;
 R₈,R₁₆,R₃₆,R₃₈,R₅₃,R₅₄=are independently U or absent;
 [R₄₇]_x = N or absent;
 10 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 15 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.
- 20 Glutamate TREM Consensus sequence
 In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{GLU} (SEQ ID NO: 580),
- R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
 25 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂
 wherein R is a ribonucleotide residue and the consensus for Glu is:
- R₀=absent;
 R₃₄,R₄₃,R₆₈,R₆₉=are independently A,C,G or absent;
 30 R₁,R₂,R₅,R₆,R₉,R₁₂,R₁₆,R₂₀,R₂₁,R₂₆,R₂₇,R₂₉,R₃₀,R₃₁,R₃₂,R₃₃,R₄₁,R₄₄,R₄₅,R₄₆,R₄₈,R₅₀,R₅₁,R₅₈,R₆₃,R₆₄,R₆₅,R₆₆,R₇₀,R₇₁=are independently N or absent;

- R₁₃,R₁₇,R₂₃,R₆₁=are independently A,C,U or absent;
 R₁₀,R₁₄,R₂₄,R₄₀,R₅₂,R₅₆=are independently A,G or absent;
 R₇,R₁₅,R₂₅,R₆₇,R₇₂=are independently A,G,U or absent;
 R₁₁,R₅₇=are independently A,U or absent;
 5 R₃₉= C,G or absent;
 R₃,R₄,R₂₂,R₄₂,R₄₉,R₅₅,R₆₂=are independently C,G,U or absent;
 R₁₈,R₂₈,R₃₅,R₃₇,R₅₃,R₅₉,R₆₀=are independently C,U or absent;
 R₁₉= G or absent;
 R₈,R₃₆,R₃₈,R₅₄=are independently U or absent;
 10 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 15 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 20 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{GLU} (SEQ ID NO: 581),

- R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
 25 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Glu is:

- R₀,R₁₈,R₂₃=are absent
 R₁₇,R₄₀=are independently A or absent;
 30 R₂₆,R₂₇,R₃₄,R₄₃,R₆₈,R₆₉,R₇₁=are independently A,C,G or absent;

R₁,R₂,R₅,R₁₂,R₂₁,R₃₁,R₃₃,R₄₁,R₄₅,R₄₈,R₅₁,R₅₈,R₆₆,R₇₀=are independently N or absent;
 R₄₄,R₆₁=are independently A,C,U or absent;
 R₉,R₁₄,R₂₄,R₂₅,R₅₂,R₅₆,R₆₃=are independently A,G or absent;
 R₇,R₁₅,R₄₆,R₅₀,R₆₇,R₇₂=are independently A,G,U or absent;
 5 R₂₉,R₅₇=are independently A,U or absent;
 R₆₀= C or absent;
 R₃₉= C,G or absent;
 R₃,R₆,R₂₀,R₃₀,R₃₂,R₄₂,R₅₅,R₆₂,R₆₅=are independently C,G,U or absent;
 R₄,R₈,R₁₆,R₂₈,R₃₅,R₃₇,R₄₉,R₅₃,R₅₉=are independently C,U or absent;
 10 R₁₀,R₁₉=are independently G or absent;
 R₂₂,R₆₄=are independently G,U or absent;
 R₁₁,R₁₃,R₃₆,R₃₈,R₅₄=are independently U or absent;
 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 15 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 20 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{GLU}
 25 (SEQ ID NO: 582),

R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

30 wherein R is a ribonucleotide residue and the consensus for Glu is:

- R₀,R₁₇,R₁₈,R₂₃=are absent
 R₁₄,R₂₇,R₄₀,R₇₁=are independently A or absent;
 R₄₄= A,C or absent;
 R₄₃= A,C,G or absent;
 5 R₁,R₃₁,R₃₃,R₄₅,R₅₁,R₆₆=are independently N or absent;
 R₂₁,R₄₁=are independently A,C,U or absent;
 R₇,R₂₄,R₂₅,R₅₀,R₅₂,R₅₆,R₆₃,R₆₈,R₇₀=are independently A,G or absent;
 R₅,R₄₆=are independently A,G,U or absent;
 R₂₉,R₅₇,R₆₇,R₇₂=are independently A,U or absent;
 10 R₂,R₃₉,R₆₀=are independently C or absent;
 R₃,R₁₂,R₂₀,R₂₆,R₃₄,R₆₉=are independently C,G or absent;
 R₆,R₃₀,R₄₂,R₄₈,R₆₅=are independently C,G,U or absent;
 R₄,R₁₆,R₂₈,R₃₅,R₃₇,R₄₉,R₅₃,R₅₅,R₅₈,R₆₁,R₆₂=are independently C,U or absent;
 R₉,R₁₀,R₁₉,R₆₄=are independently G or absent;
 15 R₁₅,R₂₂,R₃₂=are independently G,U or absent;
 R₈,R₁₁,R₁₃,R₃₆,R₃₈,R₅₄,R₅₉=are independently U or absent;
 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 20 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 25 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 residues are N; or no more than 20 residues are absent.

Glycine TREM Consensus sequence

- In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{GLY}
 30 (SEQ ID NO: 583),

R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

5 wherein R is a ribonucleotide residue and the consensus for Gly is:

R₀=absent;

R₂₄= A or absent;

R₃,R₉,R₄₀,R₅₀,R₅₁=are independently A,C,G or absent;

10 R₄,R₅,R₆,R₇,R₁₂,R₁₆,R₂₁,R₂₂,R₂₆,R₂₉,R₃₀,R₃₁,R₃₂,R₃₃,R₃₄,R₄₁,R₄₂,R₄₃,R₄₄,R₄₅,R₄₆,R₄₈,R₄₉,R₅₈,R₆₃,R₆₄,R₆₅,R₆₆,R₆₇,R₆₈=are independently N or absent;

R₅₉= A,C,U or absent;

R₁,R₁₀,R₁₄,R₁₅,R₂₇,R₅₆=are independently A,G or absent;

R₂₀,R₂₅=are independently A,G,U or absent;

R₅₇,R₇₂=are independently A,U or absent;

15 R₃₈,R₃₉,R₆₀=are independently C or absent;

R₅₂= C,G or absent;

R₂,R₁₉,R₃₇,R₅₄,R₅₅,R₆₁,R₆₂,R₆₉,R₇₀=are independently C,G,U or absent;

R₁₁,R₁₃,R₁₇,R₂₈,R₃₅,R₃₆,R₇₁=are independently C,U or absent;

R₈,R₁₈,R₂₃,R₅₃=are independently U or absent;

20 [R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
25 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
provided that the TREM has one or both of the following properties: no more than 15% of the
30 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{GLY} (SEQ ID NO: 584),

5 R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Gly is:

R₀,R₁₈,R₂₃=are absent
10 R₂₄,R₂₇,R₄₀,R₇₂=are independently A or absent;
R₂₆= A,C or absent;
R₃,R₇,R₆₈=are independently A,C,G or absent;
R₅,R₃₀,R₄₁,R₄₂,R₄₄,R₄₉,R₆₇=are independently A,C,G,U or absent;
R₃₁,R₃₂,R₃₄=are independently A,C,U or absent;
15 R₉,R₁₀,R₁₄,R₁₅,R₃₃,R₅₀,R₅₆=are independently A,G or absent;
R₁₂,R₁₆,R₂₂,R₂₅,R₂₉,R₄₆=are independently A,G,U or absent;
R₅₇= A,U or absent;
R₁₇,R₃₈,R₃₉,R₆₀,R₆₁,R₇₁=are independently C or absent;
R₆,R₅₂,R₆₄,R₆₆=are independently C,G or absent;
20 R₂,R₄,R₃₇,R₄₈,R₅₅,R₆₅=are independently C,G,U or absent;
R₁₃,R₃₅,R₄₃,R₆₂,R₆₉=are independently C,U or absent;
R₁,R₁₉,R₂₀,R₅₁,R₇₀=are independently G or absent;
R₂₁,R₄₅,R₆₃=are independently G,U or absent;
R₈,R₁₁,R₂₈,R₃₆,R₅₃,R₅₄,R₅₈,R₅₉=are independently U or absent;
25 [R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-

271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),

- 5 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{GLY} (SEQ ID NO: 585),

10 R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Gly is:

- 15 R₀,R₁₈,R₂₃=are absent
 R₂₄,R₂₇,R₄₀,R₇₂=are independently A or absent;
 R₂₆= A,C or absent;
 R₃,R₇,R₄₉,R₆₈=are independently A,C,G or absent;
 R₅,R₃₀,R₄₁,R₄₄,R₆₇=are independently N or absent;
 20 R₃₁,R₃₂,R₃₄=are independently A,C,U or absent;
 R₉,R₁₀,R₁₄,R₁₅,R₃₃,R₅₀,R₅₆=are independently A,G or absent;
 R₁₂,R₂₅,R₂₉,R₄₂,R₄₆=are independently A,G,U or absent;
 R₁₆,R₅₇=are independently A,U or absent;
 R₁₇,R₃₈,R₃₉,R₆₀,R₆₁,R₇₁=are independently C or absent;
 25 R₆,R₅₂,R₆₄,R₆₆=are independently C,G or absent;
 R₃₇,R₄₈,R₆₅=are independently C,G,U or absent;
 R₂,R₄,R₁₃,R₃₅,R₄₃,R₅₅,R₆₂,R₆₉=are independently C,U or absent;
 R₁,R₁₉,R₂₀,R₅₁,R₇₀=are independently G or absent;
 R₂₁,R₂₂,R₄₅,R₆₃=are independently G,U or absent;
 30 R₈,R₁₁,R₂₈,R₃₆,R₅₃,R₅₄,R₅₈,R₅₉=are independently U or absent;

[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 10 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

Histidine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{HIS} (SEQ ID NO: 586),

15 R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
 R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for His is:

20 R₂₃=absent;
 R₁₄,R₂₄,R₅₇=are independently A or absent;
 R₇₂= A,C or absent;
 R₉,R₂₇,R₄₃,R₄₈,R₆₉=are independently A,C,G or absent;
 25 R₃,R₄,R₅,R₆,R₁₂,R₂₅,R₂₆,R₂₉,R₃₀,R₃₁,R₃₄,R₄₂,R₄₅,R₄₆,R₄₉,R₅₀,R₅₈,R₆₂,R₆₃,R₆₆,R₆₇,R₆₈=are
 independently N or absent;
 R₁₃,R₂₁,R₄₁,R₄₄,R₆₅=are independently A,C,U or absent;
 R₄₀,R₅₁,R₅₆,R₇₀=are independently A,G or absent;
 R₇,R₃₂=are independently A,G,U or absent;
 R₅₅,R₆₀=are independently C or absent;
 30 R₁₁,R₁₆,R₃₃,R₆₄=are independently C,G,U or absent;
 R₂,R₁₇,R₂₂,R₂₈,R₃₅,R₅₃,R₅₉,R₆₁,R₇₁=are independently C,U or absent;

$R_1, R_{10}, R_{15}, R_{19}, R_{20}, R_{37}, R_{39}, R_{52}$ are independently G or absent;

R_0 = G, U or absent;

$R_8, R_{18}, R_{36}, R_{38}, R_{54}$ are independently U or absent;

$[R_{47}]_x$ = N or absent;

5 wherein, *e.g.*, $x=1-271$ (*e.g.*, $x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or $x=271$), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.$

15

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{HIS} (SEQ ID NO: 587),

$R_0-R_1-R_2-R_3-R_4-R_5-R_6-R_7-R_8-R_9-R_{10}-R_{11}-R_{12}-R_{13}-R_{14}-R_{15}-R_{16}-R_{17}-R_{18}-R_{19}-R_{20}-R_{21}-R_{22}-R_{23}-R_{24}-R_{25}-R_{26}-R_{27}-R_{28}-R_{29}-R_{30}-R_{31}-R_{32}-R_{33}-R_{34}-R_{35}-R_{36}-R_{37}-R_{38}-R_{39}-R_{40}-R_{41}-R_{42}-R_{43}-R_{44}-R_{45}-R_{46}-[R_{47}]_x-R_{48}-R_{49}-R_{50}-R_{51}-R_{52}-R_{53}-R_{54}-R_{55}-R_{56}-R_{57}-R_{58}-R_{59}-R_{60}-R_{61}-R_{62}-R_{63}-R_{64}-R_{65}-R_{66}-R_{67}-R_{68}-R_{69}-R_{70}-R_{71}-R_{72}$

20 wherein R is a ribonucleotide residue and the consensus for His is:

$R_0, R_{17}, R_{18}, R_{23}$ are absent;

$R_7, R_{12}, R_{14}, R_{24}, R_{27}, R_{45}, R_{57}, R_{58}, R_{63}, R_{67}, R_{72}$ are independently A or absent;

25 R_3 = A, C, U or absent;

$R_4, R_{43}, R_{56}, R_{70}$ are independently A, G or absent;

R_{49} = A, U or absent;

$R_2, R_{28}, R_{30}, R_{41}, R_{42}, R_{44}, R_{48}, R_{55}, R_{60}, R_{66}, R_{71}$ are independently C or absent;

R_{25} = C, G or absent;

30 R_9 = C, G, U or absent;

R₈,R₁₃,R₂₆,R₃₃,R₃₅,R₅₀,R₅₃,R₆₁,R₆₈=are independently C,U or absent;

R₁,R₆,R₁₀,R₁₅,R₁₉,R₂₀,R₃₂,R₃₄,R₃₇,R₃₉,R₄₀,R₄₆,R₅₁,R₅₂,R₆₂,R₆₄,R₆₉=are independently G or absent;

R₁₆= G,U or absent;

5 R₅,R₁₁,R₂₁,R₂₂,R₂₉,R₃₁,R₃₆,R₃₈,R₅₄,R₅₉,R₆₅=are independently U or absent;

[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 10 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{HIS} (SEQ ID NO: 588),

R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 20 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for His is:

R₀,R₁₇,R₁₈,R₂₃=are absent

25 R₇,R₁₂,R₁₄,R₂₄,R₂₇,R₄₅,R₅₇,R₅₈,R₆₃,R₆₇,R₇₂=are independently A or absent;

R₃= A,C or absent;

R₄,R₄₃,R₅₆,R₇₀=are independently A,G or absent;

R₄₉= A,U or absent;

R₂,R₂₈,R₃₀,R₄₁,R₄₂,R₄₄,R₄₈,R₅₅,R₆₀,R₆₆,R₇₁=are independently C or absent;

30 R₈,R₉,R₂₆,R₃₃,R₃₅,R₅₀,R₆₁,R₆₈=are independently C,U or absent;

R₁,R₆,R₁₀,R₁₅,R₁₉,R₂₀,R₂₅,R₃₂,R₃₄,R₃₇,R₃₉,R₄₀,R₄₆,R₅₁,R₅₂,R₆₂,R₆₄,R₆₉=are independently G or absent;

R₅,R₁₁,R₁₃,R₁₆,R₂₁,R₂₂,R₂₉,R₃₁,R₃₆,R₃₈,R₅₃,R₅₄,R₅₉,R₆₅=are independently U or absent;

[R₄₇]_x = N or absent;

5 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

15 Isoleucine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{ILE} (SEQ ID NO: 589),

R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
20 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Ile is:

R₂₃=absent;

R₃₈,R₄₁,R₅₇,R₇₂=are independently A or absent;

25 R₁,R₂₆=are independently A,C,G or absent;

R₀,R₃,R₄,R₆,R₁₆,R₃₁,R₃₂,R₃₄,R₃₇,R₄₂,R₄₃,R₄₄,R₄₅,R₄₆,R₄₈,R₄₉,R₅₀,R₅₈,R₅₉,R₆₂,R₆₃,R₆₄,R₆₆,R₆₇,R₆₈,R₆₉=are independently N or absent;

R₂₂,R₆₁,R₆₅=are independently A,C,U or absent;

R₉,R₁₄,R₁₅,R₂₄,R₂₇,R₄₀=are independently A,G or absent;

30 R₇,R₂₅,R₂₉,R₅₁,R₅₆=are independently A,G,U or absent;

R₁₈,R₅₄=are independently A,U or absent;

R₆₀= C or absent;

R₂,R₅₂,R₇₀=are independently C,G or absent;

R₅,R₁₂,R₂₁,R₃₀,R₃₃,R₇₁=are independently C,G,U or absent;

R₁₁,R₁₃,R₁₇,R₂₈,R₃₅,R₅₃,R₅₅=are independently C,U or absent;

5 R₁₀,R₁₉,R₂₀=are independently G or absent;

R₈,R₃₆,R₃₉=are independently U or absent;

[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{Ile} (SEQ ID NO: 590),

20 R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Ile is:

25 R₀,R₁₈,R₂₃=are absent

R₂₄,R₃₈,R₄₀,R₄₁,R₅₇,R₇₂=are independently A or absent;

R₂₆,R₆₅=are independently A,C or absent;

R₅₈,R₅₉,R₆₇=are independently N or absent;

R₂₂= A,C,U or absent;

30 R₆,R₉,R₁₄,R₁₅,R₂₉,R₃₄,R₄₃,R₄₆,R₄₈,R₅₀,R₅₁,R₆₃,R₆₉=are independently A,G or absent;

- R₃₇,R₅₆=are independently A,G,U or absent;
 R₅₄= A,U or absent;
 R₂₈,R₃₅,R₆₀,R₆₂,R₇₁=are independently C or absent;
 R₂,R₅₂,R₇₀=are independently C,G or absent;
 5 R₅= C,G,U or absent;
 R₃,R₄,R₁₁,R₁₃,R₁₇,R₂₁,R₃₀,R₄₂,R₄₄,R₄₅,R₄₉,R₅₃,R₅₅,R₆₁,R₆₄,R₆₆=are independently C,U or absent;
 R₁,R₁₀,R₁₉,R₂₀,R₂₅,R₂₇,R₃₁,R₆₈=are independently G or absent;
 R₇,R₁₂,R₃₂=are independently G,U or absent;
 10 R₈,R₁₆,R₃₃,R₃₆,R₃₉=are independently U or absent;
 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 15 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{Ile} (SEQ ID NO: 591),

- R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 25 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Ile is:

- R₀,R₁₈,R₂₃=are absent
 30 R₁₄,R₂₄,R₃₈,R₄₀,R₄₁,R₅₇,R₇₂=are independently A or absent;

- R₂₆,R₆₅=are independently A,C or absent;
 R₂₂,R₅₉=are independently A,C,U or absent;
 R₆,R₉,R₁₅,R₃₄,R₄₃,R₄₆,R₅₁,R₅₆,R₆₃,R₆₉=are independently A,G or absent;
 R₃₇= A,G,U or absent;
 5 R₁₃,R₂₈,R₃₅,R₄₄,R₅₅,R₆₀,R₆₂,R₇₁=are independently C or absent;
 R₂,R₅,R₇₀=are independently C,G or absent;
 R₅₈,R₆₇=are independently C,G,U or absent;
 R₃,R₄,R₁₁,R₁₇,R₂₁,R₃₀,R₄₂,R₄₅,R₄₉,R₅₃,R₆₁,R₆₄,R₆₆=are independently C,U or absent;
 R₁,R₁₀,R₁₉,R₂₀,R₂₅,R₂₇,R₂₉,R₃₁,R₃₂,R₄₈,R₅₀,R₅₂,R₆₈=are independently G or absent;
 10 R₇,R₁₂=are independently G,U or absent;
 R₈,R₁₆,R₃₃,R₃₆,R₃₉,R₅₄=are independently U or absent;
 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 15 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 20 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 residues are N; or no more than 20 residues are absent.

Methionine TREM Consensus sequence

- 25 In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{MET} (SEQ ID
 NO: 592),

R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

- 30 wherein R is a ribonucleotide residue and the consensus for Met is:

- R_0, R_{23} = are absent;
 $R_{14}, R_{38}, R_{40}, R_{57}$ = are independently A or absent;
 R_{60} = A, C or absent;
 R_{33}, R_{48}, R_{70} = are independently A, C, G or absent;
5 $R_1, R_3, R_4, R_5, R_6, R_{11}, R_{12}, R_{16}, R_{17}, R_{21}, R_{22}, R_{26}, R_{27}, R_{29}, R_{30}, R_{31}, R_{32}, R_{42}, R_{44}, R_{45}, R_{46}, R_{49}, R_{50}, R_{58}, R_{62}, R_{63}, R_{66}, R_{67}, R_{68}, R_{69}, R_{71}$ = are independently N or absent;
 $R_{18}, R_{35}, R_{41}, R_{59}, R_{65}$ = are independently A, C, U or absent;
 R_9, R_{15}, R_{51} = are independently A, G or absent;
 $R_7, R_{24}, R_{25}, R_{34}, R_{53}, R_{56}$ = are independently A, G, U or absent;
10 R_{72} = A, U or absent;
 R_{37} = C or absent;
 R_{10}, R_{55} = are independently C, G or absent;
 $R_2, R_{13}, R_{28}, R_{43}, R_{64}$ = are independently C, G, U or absent;
 R_{36}, R_{61} = are independently C, U or absent;
15 R_{19}, R_{20}, R_{52} = are independently G or absent;
 R_8, R_{39}, R_{54} = are independently U or absent;
 $[R_{47}]_x$ = N or absent;
wherein, *e.g.*, $x=1-271$ (*e.g.*, $x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or $x=271$),
20 $x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271$),
25 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.$

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{MET} (SEQ ID NO: 593),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

5 wherein R is a ribonucleotide residue and the consensus for Met is:

R₀,R₁₈,R₂₂,R₂₃=are absent

R₁₄,R₂₄,R₃₈,R₄₀,R₄₁,R₅₇,R₇₂=are independently A or absent;

R₅₉,R₆₀,R₆₂,R₆₅=are independently A,C or absent;

R₆,R₄₅,R₆₇=are independently A,C,G or absent;

10 R₄= N or absent;

R₂₁,R₄₂=are independently A,C,U or absent;

R₁,R₉,R₂₇,R₂₉,R₃₂,R₄₆,R₅₁=are independently A,G or absent;

R₁₇,R₄₉,R₅₃,R₅₆,R₅₈=are independently A,G,U or absent;

R₆₃=A,U or absent;

15 R₃,R₁₃,R₃₇=are independently C or absent;

R₄₈,R₅₅,R₆₄,R₇₀=are independently C,G or absent;

R₂,R₅,R₆₆,R₆₈=are independently C,G,U or absent;

R₁₁,R₁₆,R₂₆,R₂₈,R₃₀,R₃₁,R₃₅,R₃₆,R₄₃,R₄₄,R₆₁,R₇₁=are independently C,U or absent;

R₁₀,R₁₂,R₁₅,R₁₉,R₂₀,R₂₅,R₃₃,R₅₂,R₆₉=are independently G or absent;

20 R₇,R₃₄,R₅₀=are independently G,U or absent;

R₈,R₃₉,R₅₄=are independently U or absent;

[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
25 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,

x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

5 In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{MET} (SEQ ID NO: 594),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
10 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Met is:

R₀,R₁₈,R₂₂,R₂₃=are absent
R₁₄,R₂₄,R₃₈,R₄₀,R₄₁,R₅₇,R₇₂=are independently A or absent;
R₅₉,R₆₂,R₆₅=are independently A,C or absent;
15 R₆,R₆₇=are independently A,C,G or absent;
R₄,R₂₁=are independently A,C,U or absent;
R₁,R₉,R₂₇,R₂₉,R₃₂,R₄₅,R₄₆,R₅₁=are independently A,G or absent;
R₁₇,R₅₆,R₅₈=are independently A,G,U or absent;
R₄₉,R₅₃,R₆₃=are independently A,U or absent;
20 R₃,R₁₃,R₂₆,R₃₇,R₄₃,R₆₀=are independently C or absent;
R₂,R₄₈,R₅₅,R₆₄,R₇₀=are independently C,G or absent;
R₅,R₆₆=are independently C,G,U or absent;
R₁₁,R₁₆,R₂₈,R₃₀,R₃₁,R₃₅,R₃₆,R₄₂,R₄₄,R₆₁,R₇₁=are independently C,U or absent;
R₁₀,R₁₂,R₁₅,R₁₉,R₂₀,R₂₅,R₃₃,R₅₂,R₆₉=are independently G or absent;
25 R₇,R₃₄,R₅₀,R₆₈=are independently G,U or absent;
R₈,R₃₉,R₅₄=are independently U or absent;
[R₄₇]_x = N or absent;

wherein, e.g., x=1-271 (e.g., x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,

x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, 5 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

Leucine TREM Consensus sequence

10 In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{LEU} (SEQ ID NO: 595),

R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

15 wherein R is a ribonucleotide residue and the consensus for Leu is:

R₀=absent;

R₃₈,R₅₇=are independently A or absent;

R₆₀= A,C or absent;

R₁,R₁₃,R₂₇,R₄₈,R₅₁,R₅₆=are independently A,C,G or absent;

20 R₂,R₃,R₄,R₅,R₆,R₇,R₉,R₁₀,R₁₁,R₁₂,R₁₆,R₂₃,R₂₆,R₂₈,R₂₉,R₃₀,R₃₁,R₃₂,R₃₃,R₃₄,R₃₇,R₄₁,R₄₂,R₄₃,R₄₄,
R₄₅,R₄₆,R₄₉,R₅₀,R₅₈,R₆₂,R₆₃,R₆₅,R₆₆,R₆₇,R₆₈,R₆₉,R₇₀=are independently N or absent;

R₁₇,R₁₈,R₂₁,R₂₂,R₂₅,R₃₅,R₅₅=are independently A,C,U or absent;

R₁₄,R₁₅,R₃₉,R₇₂=are independently A,G or absent;

R₂₄,R₄₀=are independently A,G,U or absent;

25 R₅₂,R₆₁,R₆₄,R₇₁=are independently C,G,U or absent;

R₃₆,R₅₃,R₅₉=are independently C,U or absent;

R₁₉= G or absent;

R₂₀= G,U or absent;

R₈,R₅₄=are independently U or absent;

30 [R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 5 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the
 10 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{LEU} (SEQ ID NO: 596),

R₀- R₁- R₂- R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
 15 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Leu is:

R₀=absent
 R₃₈,R₅₇,R₇₂=are independently A or absent;
 20 R₆₀= A,C or absent;
 R₄,R₅,R₄₈,R₅₀,R₅₆,R₆₉=are independently A,C,G or absent;
 R₆,R₃₃,R₄₁,R₄₃,R₄₆,R₄₉,R₅₈,R₆₃,R₆₆,R₇₀=are independently N or absent;
 R₁₁,R₁₂,R₁₇,R₂₁,R₂₂,R₂₈,R₃₁,R₃₇,R₄₄,R₅₅=are independently A,C,U or absent;
 R₁,R₉,R₁₄,R₁₅,R₂₄,R₂₇,R₃₄,R₃₉=are independently A,G or absent;
 25 R₇,R₂₉,R₃₂,R₄₀,R₄₅=are independently A,G,U or absent;
 R₂₅= A,U or absent;
 R₁₃= C,G or absent;
 R₂,R₃,R₁₆,R₂₆,R₃₀,R₅₂,R₆₂,R₆₄,R₆₅,R₆₇,R₆₈=are independently C,G,U or absent;
 R₁₈,R₃₅,R₄₂,R₅₃,R₅₉,R₆₁,R₇₁=are independently C,U or absent;
 30 R₁₉,R₅₁=are independently G or absent;

R_{10}, R_{20} are independently G, U or absent;

$R_8, R_{23}, R_{36}, R_{54}$ are independently U or absent;

$[R_{47}]_x = N$ or absent;

wherein, *e.g.*, $x=1-271$ (*e.g.*, $x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,$
 5 $x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,$
 $x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,$
 $x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-$
 $271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,$
 $x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,$
 10 $x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,$
 $x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, \text{ or } x=271$),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{LEU}
 15 (SEQ ID NO: 597),

$R_0-R_1-R_2-R_3-R_4-R_5-R_6-R_7-R_8-R_9-R_{10}-R_{11}-R_{12}-R_{13}-R_{14}-R_{15}-R_{16}-R_{17}-R_{18}-R_{19}-R_{20}-R_{21}-R_{22}-$
 $R_{23}-R_{24}-R_{25}-R_{26}-R_{27}-R_{28}-R_{29}-R_{30}-R_{31}-R_{32}-R_{33}-R_{34}-R_{35}-R_{36}-R_{37}-R_{38}-R_{39}-R_{40}-R_{41}-R_{42}-R_{43}-R_{44}-R_{45}-$
 $R_{46}-[R_{47}]_x-R_{48}-R_{49}-R_{50}-R_{51}-R_{52}-R_{53}-R_{54}-R_{55}-R_{56}-R_{57}-R_{58}-R_{59}-R_{60}-R_{61}-R_{62}-R_{63}-R_{64}-R_{65}-R_{66}-R_{67}-$
 $R_{68}-R_{69}-R_{70}-R_{71}-R_{72}$

20 wherein R is a ribonucleotide residue and the consensus for Leu is:

R_0 = absent

R_{38}, R_{57}, R_{72} are independently A or absent;

R_{60} = A, C or absent;

$R_4, R_5, R_{48}, R_{50}, R_{56}, R_{58}, R_{69}$ are independently A, C, G or absent;

25 $R_6, R_{33}, R_{43}, R_{46}, R_{49}, R_{63}, R_{66}, R_{70}$ are independently N or absent;

$R_{11}, R_{12}, R_{17}, R_{21}, R_{22}, R_{28}, R_{31}, R_{37}, R_{41}, R_{44}, R_{55}$ are independently A, C, U or absent;

$R_1, R_9, R_{14}, R_{15}, R_{24}, R_{27}, R_{34}, R_{39}$ are independently A, G or absent;

$R_7, R_{29}, R_{32}, R_{40}, R_{45}$ are independently A, G, U or absent;

R_{25} = A, U or absent;

30 R_{13} = C, G or absent;

- R₂,R₃,R₁₆,R₃₀,R₅₂,R₆₂,R₆₄,R₆₇,R₆₈=are independently C,G,U or absent;
 R₁₈,R₃₅,R₄₂,R₅₃,R₅₉,R₆₁,R₆₅,R₇₁=are independently C,U or absent;
 R₁₉,R₅₁=are independently G or absent;
 R₁₀,R₂₀,R₂₆=are independently G,U or absent;
 5 R₈,R₂₃,R₃₆,R₅₄=are independently U or absent;
 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 10 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

Lysine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{LYS} (SEQ ID NO: 598),

- 20 R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Lys is:

- 25 R₀ =absent
 R₁₄= A or absent;
 R₄₀,R₄₁=are independently A,C or absent;
 R₃₄,R₄₃,R₅₁=are independently A,C,G or absent;
 30 R₁,R₂,R₃,R₄,R₅,R₆,R₇,R₁₁,R₁₂,R₁₆,R₂₁,R₂₆,R₃₀,R₃₁,R₃₂,R₄₄,R₄₅,R₄₆,R₄₈,R₄₉,R₅₀,R₅₈,R₆₂,R₆₃,R₆₅,
 R₆₆,R₆₇,R₆₈,R₆₉,R₇₀=are independently N or absent;
 R₁₃,R₁₇,R₅₉,R₇₁=are independently A,C,U or absent;

R₉,R₁₅,R₁₉,R₂₀,R₂₅,R₂₇,R₅₂,R₅₆=are independently A,G or absent;
 R₂₄,R₂₉,R₇₂=are independently A,G,U or absent;
 R₁₈,R₅₇=are independently A,U or absent;
 R₁₀,R₃₃=are independently C,G or absent;
 5 R₄₂,R₆₁,R₆₄=are independently C,G,U or absent;
 R₂₈,R₃₅,R₃₆,R₃₇,R₅₃,R₅₅,R₆₀=are independently C,U or absent;
 R₈,R₂₂,R₂₃,R₃₈,R₃₉,R₅₄=are independently U or absent;
 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 10 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 15 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{LYS}
 20 (SEQ ID NO: 599),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
 R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

25 wherein R is a ribonucleotide residue and the consensus for Lys is:

R₀,R₁₈,R₂₃=are absent
 R₁₄= A or absent;
 R₄₀,R₄₁,R₄₃=are independently A,C or absent;
 R₃,R₇=are independently A,C,G or absent;
 30 R₁,R₆,R₁₁,R₃₁,R₄₅,R₄₈,R₄₉,R₆₃,R₆₅,R₆₆,R₆₈=are independently N or absent;

- R₂,R₁₂,R₁₃,R₁₇,R₄₄,R₆₇,R₇₁=are independently A,C,U or absent;
- R₉,R₁₅,R₁₉,R₂₀,R₂₅,R₂₇,R₃₄,R₅₀,R₅₂,R₅₆,R₇₀,R₇₂=are independently A,G or absent;
- R₅,R₂₄,R₂₆,R₂₉,R₃₂,R₄₆,R₆₉=are independently A,G,U or absent;
- R₅₇= A,U or absent;
- 5 R₁₀,R₆₁=are independently C,G or absent;
- R₄,R₁₆,R₂₁,R₃₀,R₅₈,R₆₄=are independently C,G,U or absent;
- R₂₈,R₃₅,R₃₆,R₃₇,R₄₂,R₅₃,R₅₅,R₅₉,R₆₀,R₆₂=are independently C,U or absent;
- R₃₃,R₅₁=are independently G or absent;
- R₈=G,U or absent;
- 10 R₂₂,R₃₈,R₃₉,R₅₄=are independently U or absent;
- [R₄₇]_x = N or absent;
- wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
- 15 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
- 20 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{Lys} (SEQ ID NO: 600),

- R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
- 25 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
- R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
- R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Lys is:

- R₀,R₁₈,R₂₃=absent
- 30 R₉,R₁₄,R₃₄,R₄₁=are independently A or absent;

- R₄₀= A,C or absent;
 R₁,R₃,R₇,R₃₁=are independently A,C,G or absent;
 R₄₈,R₆₅,R₆₈=are independently N or absent;
 R₂,R₁₃,R₁₇,R₄₄,R₆₃,R₆₆=are independently A,C,U or absent;
 5 R₅,R₁₅,R₁₉,R₂₀,R₂₅,R₂₇,R₂₉,R₅₀,R₅₂,R₅₆,R₇₀,R₇₂=are independently A,G or absent;
 R₆,R₂₄,R₃₂,R₄₉=are independently A,G,U or absent;
 R₁₂,R₂₆,R₄₆,R₅₇=are independently A,U or absent;
 R₁₁,R₂₈,R₃₅,R₄₃=are independently C or absent;
 R₁₀,R₄₅,R₆₁=are independently C,G or absent;
 10 R₄,R₂₁,R₆₄=are independently C,G,U or absent;
 R₃₇,R₅₃,R₅₅,R₅₉,R₆₀,R₆₂,R₆₇,R₇₁=are independently C,U or absent;
 R₃₃,R₅₁=are independently G or absent;
 R₈,R₃₀,R₅₈,R₆₉=are independently G,U or absent;
 R₁₆,R₂₂,R₃₆,R₃₈,R₃₉,R₄₂,R₅₄=are independently U or absent;
 15 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 20 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 25 residues are N; or no more than 20 residues are absent.

Phenylalanine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{PHE} (SEQ ID NO: 601),

R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 30 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-

R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Phe is:

- R₀,R₂₃=are absent
- 5 R₉,R₁₄,R₃₈,R₃₉,R₅₇,R₇₂=are independently A or absent;
R₇₁= A,C or absent;
R₄₁,R₇₀=are independently A,C,G or absent;
R₄,R₅,R₆,R₃₀,R₃₁,R₃₂,R₃₄,R₄₂,R₄₄,R₄₅,R₄₆,R₄₈,R₄₉,R₅₈,R₆₂,R₆₃,R₆₆,R₆₇,R₆₈,R₆₉=are
independently N or absent;
- 10 R₁₆,R₆₁,R₆₅=are independently A,C,U or absent;
R₁₅,R₂₆,R₂₇,R₂₉,R₄₀,R₅₆=are independently A,G or absent;
R₇,R₅₁=are independently A,G,U or absent;
R₂₂,R₂₄=are independently A,U or absent;
R₅₅,R₆₀=are independently C or absent;
- 15 R₂,R₃,R₂₁,R₃₃,R₄₃,R₅₀,R₆₄=are independently C,G,U or absent;
R₁₁,R₁₂,R₁₃,R₁₇,R₂₈,R₃₅,R₃₆,R₅₉=are independently C,U or absent;
R₁₀,R₁₉,R₂₀,R₂₅,R₃₇,R₅₂=are independently G or absent;
R₁= G,U or absent;
R₈,R₁₈,R₅₃,R₅₄=are independently U or absent;
- 20 [R₄₇]_x = N or absent;
wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
25 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
provided that the TREM has one or both of the following properties: no more than 15% of the
30 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{PHE} (SEQ ID NO: 602),

R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
 5 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Phe is:

R₀,R₁₈,R₂₃=absent
 R₁₄,R₂₄,R₃₈,R₃₉,R₅₇,R₇₂=are independently A or absent;
 10 R₄₆,R₇₁=are independently A,C or absent;
 R₄,R₇₀=are independently A,C,G or absent;
 R₄₅= A,C,U or absent;
 R₆,R₇,R₁₅,R₂₆,R₂₇,R₃₂,R₃₄,R₄₀,R₄₁,R₅₆,R₆₉=are independently A,G or absent;
 R₂₉= A,G,U or absent;
 15 R₅,R₉,R₆₇=are independently A,U or absent;
 R₃₅,R₄₉,R₅₅,R₆₀=are independently C or absent;
 R₂₁,R₄₃,R₆₂=are independently C,G or absent;
 R₂,R₃₃,R₆₈=are independently C,G,U or absent;
 R₃,R₁₁,R₁₂,R₁₃,R₂₈,R₃₀,R₃₆,R₄₂,R₄₄,R₄₈,R₅₈,R₅₉,R₆₁,R₆₆=are independently C,U or absent;
 20 R₁₀,R₁₉,R₂₀,R₂₅,R₃₇,R₅₁,R₅₂,R₆₃,R₆₄=are independently G or absent;
 R₁,R₃₁,R₅₀=are independently G,U or absent;
 R₈,R₁₆,R₁₇,R₂₂,R₅₃,R₅₄,R₆₅=are independently U or absent;
 [R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 25 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,

x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

5 In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{PHE} (SEQ ID NO: 603),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
10 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Phe is:

R₀,R₁₈,R₂₂,R₂₃=absent
R₅,R₇,R₁₄,R₂₄,R₂₆,R₃₂,R₃₄,R₃₈,R₃₉,R₄₁,R₅₇,R₇₂=are independently A or absent;
R₄₆= A,C or absent;
15 R₇₀= A,C,G or absent;
R₄,R₆,R₁₅,R₅₆,R₆₉=are independently A,G or absent;
R₉,R₄₅=are independently A,U or absent;
R₂,R₁₁,R₁₃,R₃₅,R₄₃,R₄₉,R₅₅,R₆₀,R₆₈,R₇₁=are independently C or absent;
R₃₃= C,G or absent;
20 R₃,R₂₈,R₃₆,R₄₈,R₅₈,R₅₉,R₆₁=are independently C,U or absent;
R₁,R₁₀,R₁₉,R₂₀,R₂₁,R₂₅,R₂₇,R₂₉,R₃₇,R₄₀,R₅₁,R₅₂,R₆₂,R₆₃,R₆₄=are independently G or absent;
R₈,R₁₂,R₁₆,R₁₇,R₃₀,R₃₁,R₄₂,R₄₄,R₅₀,R₅₃,R₅₄,R₆₅,R₆₆,R₆₇=are independently U or absent;
[R₄₇]_x = N or absent;
wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
25 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,

x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

5 Proline TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{PRO} (SEQ ID NO: 604),

10 R₀- R₁- R₂- R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
10 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Pro is:

R₀ =absent
R₁₄,R₅₇=are independently A or absent;
15 R₇₀,R₇₂=are independently A,C or absent;
R₉,R₂₆,R₂₇=are independently A,C,G or absent;
R₄,R₅,R₆,R₁₆,R₂₁,R₂₉,R₃₀,R₃₁,R₃₂,R₃₃,R₃₄,R₃₇,R₄₁,R₄₂,R₄₃,R₄₄,R₄₅,R₄₆,R₄₈,R₄₉,R₅₀,R₅₈,R₆₁,R₆₂,
R₆₃,R₆₄,R₆₆,R₆₇,R₆₈=are independently N or absent;
R₃₅,R₆₅=are independently A,C,U or absent;
20 R₂₄,R₄₀,R₅₆=are independently A,G or absent;
R₇,R₂₅,R₅₁=are independently A,G,U or absent;
R₅₅,R₆₀=are independently C or absent;
R₁,R₃,R₇₁=are independently C,G or absent;
R₁₁,R₁₂,R₂₀,R₆₉=are independently C,G,U or absent;
25 R₁₃,R₁₇,R₁₈,R₂₂,R₂₃,R₂₈,R₅₉=are independently C,U or absent;
R₁₀,R₁₅,R₁₉,R₃₈,R₃₉,R₅₂=are independently G or absent;
R₂= are independently G,U or absent;
R₈,R₃₆,R₅₃,R₅₄=are independently U or absent;
[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{PRO} (SEQ ID NO: 605),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Pro is:

R₀,R₁₇,R₁₈,R₂₂,R₂₃=absent;
 R₁₄,R₄₅,R₅₆,R₅₇,R₅₈,R₆₅,R₆₈=are independently A or absent;
 R₆₁= A,C,G or absent;
 R₄₃=N or absent;
 R₃₇= A, C,U or absent;
 R₂₄,R₂₇,R₃₃,R₄₀,R₄₄,R₆₃=are independently A,G or absent;
 R₃,R₁₂,R₃₀,R₃₂,R₄₈,R₅₅,R₆₀,R₇₀,R₇₁,R₇₂=are independently C or absent;
 R₅,R₃₄,R₄₂,R₆₆=are independently C,G or absent;
 R₂₀= C,G,U or absent;
 R₃₅,R₄₁,R₄₉,R₆₂=are independently C,U or absent;
 R₁,R₂,R₆,R₉,R₁₀,R₁₅,R₁₉,R₂₆,R₃₈,R₃₉,R₄₆,R₅₀,R₅₁,R₅₂,R₆₄,R₆₇,R₆₉=are independently G or absent;
 R₁₁,R₁₆=are independently G,U or absent;

R₄,R₇,R₈,R₁₃,R₂₁,R₂₅,R₂₈,R₂₉,R₃₁,R₃₆,R₅₃,R₅₄,R₅₉=are independently U or absent;

[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{PRO} (SEQ ID NO: 606),

R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Pro is:

R₀,R₁₇,R₁₈,R₂₂,R₂₃=absent

R₁₄,R₄₅,R₅₆,R₅₇,R₅₈,R₆₅,R₆₈=are independently A or absent;

R₃₇= A,C,U or absent;

R₂₄,R₂₇,R₄₀=are independently A,G or absent;

R₃,R₅,R₁₂,R₃₀,R₃₂,R₄₈,R₄₉,R₅₅,R₆₀,R₆₁,R₆₂,R₆₆,R₇₀,R₇₁,R₇₂=are independently C or absent;

R₃₄,R₄₂=are independently C,G or absent;

R₄₃= C,G,U or absent;

R₄₁= C,U or absent;

R₁,R₂,R₆,R₉,R₁₀,R₁₅,R₁₉,R₂₀,R₂₆,R₃₃,R₃₈,R₃₉,R₄₄,R₄₆,R₅₀,R₅₁,R₅₂,R₆₃,R₆₄,R₆₇,R₆₉=are independently G or absent;

R₁₆= G,U or absent;

R₄,R₇,R₈,R₁₁,R₁₃,R₂₁,R₂₅,R₂₈,R₂₉,R₃₁,R₃₅,R₃₆,R₅₃,R₅₄,R₅₉=are independently U or absent;

[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

Serine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{SER} (SEQ ID NO: 607),

R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Ser is:

R₀=absent;

R₁₄,R₂₄,R₅₇=are independently A or absent;

R₄₁= A,C or absent;

R₂,R₃,R₄,R₅,R₆,R₇,R₉,R₁₀,R₁₁,R₁₂,R₁₃,R₁₆,R₂₁,R₂₅,R₂₆,R₂₇,R₂₈,R₃₀,R₃₁,R₃₂,R₃₃,R₃₄,R₃₇,R₄₂,R₄₃,R₄₄,R₄₅,R₄₆,R₄₈,R₄₉,R₅₀,R₆₂,R₆₃,R₆₄,R₆₅,R₆₆,R₆₇,R₆₈,R₆₉,R₇₀=are independently N or absent;

R₁₈= A,C,U or absent;

R₁₅,R₄₀,R₅₁,R₅₆=are independently A,G or absent;

R₁,R₂₉,R₅₈,R₇₂=are independently A,G,U or absent;

R₃₉= A,U or absent;

R₆₀= C or absent;

R₃₈= C,G or absent;

R₁₇,R₂₂,R₂₃,R₇₁=are independently C,G,U or absent;

R₈,R₃₅,R₃₆,R₅₅,R₅₉,R₆₁=are independently C,U or absent;

R₁₉,R₂₀=are independently G or absent;

R₅₂= G,U or absent;

5 R₅₃,R₅₄=are independently U or absent;

[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
10 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{SER} (SEQ ID NO: 608),

R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
20 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Ser is:

R₀,R₂₃=absent

25 R₁₄,R₂₄,R₄₁,R₅₇=are independently A or absent;

R₄₄= A,C or absent;

R₂₅,R₄₅,R₄₈=are independently A,C,G or absent;

R₂,R₃,R₄,R₅,R₃₇,R₅₀,R₆₂,R₆₆,R₆₇,R₆₉,R₇₀=are independently N or absent;

R₁₂,R₂₈,R₆₅=are independently A,C,U or absent;

30 R₉,R₁₅,R₂₉,R₃₄,R₄₀,R₅₆,R₆₃=are independently A,G or absent;

- R₇,R₂₆,R₃₀,R₃₃,R₄₆,R₅₈,R₇₂=are independently A,G,U or absent;
 R₃₉= A,U or absent;
 R₁₁,R₃₅,R₆₀,R₆₁=are independently C or absent;
 R₁₃,R₃₈=are independently C,G or absent;
 5 R₆,R₁₇,R₃₁,R₄₃,R₆₄,R₆₈=are independently C,G,U or absent;
 R₃₆,R₄₂,R₄₉,R₅₅,R₅₉,R₇₁=are independently C,U or absent;
 R₁₀,R₁₉,R₂₀,R₂₇,R₅₁=are independently G or absent;
 R₁,R₁₆,R₃₂,R₅₂=are independently G,U or absent;
 R₈,R₁₈,R₂₁,R₂₂,R₅₃,R₅₄=are independently U or absent;
 10 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 15 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 20 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{SER}
 (SEQ ID NO: 609),

- R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-
 25 R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Ser is:

- R₀,R₂₃=absent
 R₁₄,R₂₄,R₄₁,R₅₇,R₅₈=are independently A or absent;
 30 R₄₄= A,C or absent;

- R₂₅,R₄₈=are independently A,C,G or absent;
 R₂,R₃,R₅,R₃₇,R₆₆,R₆₇,R₆₉,R₇₀=are independently N or absent;
 R₁₂,R₂₈,R₆₂=are independently A,C,U or absent;
 R₇,R₉,R₁₅,R₂₉,R₃₃,R₃₄,R₄₀,R₄₅,R₅₆,R₆₃=are independently A,G or absent;
 5 R₄,R₂₆,R₄₆,R₅₀=are independently A,G,U or absent;
 R₃₀,R₃₉=are independently A,U or absent;
 R₁₁,R₁₇,R₃₅,R₆₀,R₆₁=are independently C or absent;
 R₁₃,R₃₈=are independently C,G or absent;
 R₆,R₆₄=are independently C,G,U or absent;
 10 R₃₁,R₄₂,R₄₃,R₄₉,R₅₅,R₅₉,R₆₅,R₆₈,R₇₁=are independently C,U or absent;
 R₁₀,R₁₉,R₂₀,R₂₇,R₅₁,R₅₂=are independently G or absent;
 R₁,R₁₆,R₃₂,R₇₂=are independently G,U or absent;
 R₈,R₁₈,R₂₁,R₂₂,R₃₆,R₅₃,R₅₄=are independently U or absent;
 [R₄₇]_x = N or absent;
 15 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 20 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 residues are N; or no more than 20 residues are absent.

25 Threonine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{THR} (SEQ ID NO: 610),

R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-

R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Thr is:

- R₀,R₂₃=absent
- 5 R₁₄,R₄₁,R₅₇=are independently A or absent;
R₅₆,R₇₀=are independently A,C,G or absent;
R₄,R₅,R₆,R₇,R₁₂,R₁₆,R₂₆,R₃₀,R₃₁,R₃₂,R₃₄,R₃₇,R₄₂,R₄₄,R₄₅,R₄₆,R₄₈,R₄₉,R₅₀,R₅₈,R₆₂,R₆₃,R₆₄,R₆₅,R₆₆,R₆₇,R₆₈,R₇₂=are independently N or absent;
R₁₃,R₁₇,R₂₁,R₃₅,R₆₁=are independently A,C,U or absent;
- 10 R₁,R₉,R₂₄,R₂₇,R₂₉,R₆₉=are independently A,G or absent;
R₁₅,R₂₅,R₅₁=are independently A,G,U or absent;
R₄₀,R₅₃=are independently A,U or absent;
R₃₃,R₄₃=are independently C,G or absent;
R₂,R₃,R₅₉=are independently C,G,U or absent;
- 15 R₁₁,R₁₈,R₂₂,R₂₈,R₃₆,R₅₄,R₅₅,R₆₀,R₇₁=are independently C,U or absent;
R₁₀,R₂₀,R₃₈,R₅₂=are independently G or absent;
R₁₉= G,U or absent;
R₈,R₃₉=are independently U or absent;
[R₄₇]_x = N or absent;
- 20 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
25 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{THR} (SEQ ID NO: 611),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
5 R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Thr is:

R₀,R₁₈,R₂₃=absent
R₁₄,R₄₁,R₅₇=are independently A or absent;
10 R₉,R₄₂,R₄₄,R₄₈,R₅₆,R₇₀=are independently A,C,G or absent;
R₄,R₆,R₁₂,R₂₆,R₄₉,R₅₈,R₆₃,R₆₄,R₆₆,R₆₈=are independently N or absent;
R₁₃,R₂₁,R₃₁,R₃₇,R₆₂=are independently A,C,U or absent;
R₁,R₁₅,R₂₄,R₂₇,R₂₉,R₄₆,R₅₁,R₆₉=are independently A,G or absent;
R₇,R₂₅,R₄₅,R₅₀,R₆₇=are independently A,G,U or absent;
15 R₄₀,R₅₃=are independently A,U or absent;
R₃₅= C or absent;
R₃₃,R₄₃=are independently C,G or absent;
R₂,R₃,R₅,R₁₆,R₃₂,R₃₄,R₅₉,R₆₅,R₇₂=are independently C,G,U or absent;
R₁₁,R₁₇,R₂₂,R₂₈,R₃₀,R₃₆,R₅₅,R₆₀,R₆₁,R₇₁=are independently C,U or absent;
20 R₁₀,R₁₉,R₂₀,R₃₈,R₅₂=are independently G or absent;
R₈,R₃₉,R₅₄=are independently U or absent;
[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
25 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
30 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),

provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{THR} (SEQ ID NO: 612),

5 R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Thr is:

10 R₀,R₁₈,R₂₃=absent
R₁₄,R₄₀,R₄₁,R₅₇=are independently A or absent;
R₄₄= A,C or absent;
R₉,R₄₂,R₄₈,R₅₆=are independently A,C,G or absent;
R₄,R₆,R₁₂,R₂₆,R₅₈,R₆₄,R₆₆,R₆₈=are independently N or absent;
15 R₁₃,R₂₁,R₃₁,R₃₇,R₄₉,R₆₂=are independently A,C,U or absent;
R₁,R₁₅,R₂₄,R₂₇,R₂₉,R₄₆,R₅₁,R₆₉=are independently A,G or absent;
R₇,R₂₅,R₄₅,R₅₀,R₆₃,R₆₇=are independently A,G,U or absent;
R₅₃= A,U or absent;
R₃₅= C or absent;
20 R₂,R₃₃,R₄₃,R₇₀=are independently C,G or absent;
R₅,R₁₆,R₃₄,R₅₉,R₆₅=are independently C,G,U or absent;
R₃,R₁₁,R₂₂,R₂₈,R₃₀,R₃₆,R₅₅,R₆₀,R₆₁,R₇₁=are independently C,U or absent;
R₁₀,R₁₉,R₂₀,R₃₈,R₅₂=are independently G or absent;
R₃₂= G,U or absent;
25 R₈,R₁₇,R₃₉,R₅₄,R₇₂=are independently U or absent;
[R₄₇]_x = N or absent;
wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
30 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-

271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),

- 5 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

Tryptophan TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{TRP} (SEQ ID NO: 613),

10 R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Trp is:

- 15 R₀=absent;
R₂₄,R₃₉,R₄₁,R₅₇=are independently A or absent;
R₂,R₃,R₂₆,R₂₇,R₄₀,R₄₈=are independently A,C,G or absent;
R₄,R₅,R₆,R₂₉,R₃₀,R₃₁,R₃₂,R₃₄,R₄₂,R₄₄,R₄₅,R₄₆,R₄₉,R₅₁,R₅₈,R₆₃,R₆₆,R₆₇,R₆₈=are independently N or absent;
20 R₁₃,R₁₄,R₁₆,R₁₈,R₂₁,R₆₁,R₆₅,R₇₁=are independently A,C,U or absent;
R₁,R₉,R₁₀,R₁₅,R₃₃,R₅₀,R₅₆=are independently A,G or absent;
R₇,R₂₅,R₇₂=are independently A,G,U or absent;
R₃₇,R₃₈,R₅₅,R₆₀=are independently C or absent;
R₁₂,R₃₅,R₄₃,R₆₄,R₆₉,R₇₀=are independently C,G,U or absent;
25 R₁₁,R₁₇,R₂₂,R₂₈,R₅₉,R₆₂=are independently C,U or absent;
R₁₉,R₂₀,R₅₂=are independently G or absent;
R₈,R₂₃,R₃₆,R₅₃,R₅₄=are independently U or absent;
[R₄₇]_x=N or absent;

wherein, e.g., x=1-271 (e.g., x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
30 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,

x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{TRP} (SEQ ID NO: 614),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Trp is:

R₀,R₁₈,R₂₂,R₂₃=absent

R₁₄,R₂₄,R₃₉,R₄₁,R₅₇,R₇₂=are independently A or absent;

R₃,R₄,R₁₃,R₆₁,R₇₁=are independently A,C or absent;

R₆,R₄₄=are independently A,C,G or absent;

R₂₁= A,C,U or absent;

R₂,R₇,R₁₅,R₂₅,R₃₃,R₃₄,R₄₅,R₅₆,R₆₃=are independently A,G or absent;

R₅₈= A,G,U or absent;

R₄₆= A,U or absent;

R₃₇,R₃₈,R₅₅,R₆₀,R₆₂=are independently C or absent;

R₁₂,R₂₆,R₂₇,R₃₅,R₄₀,R₄₈,R₆₇=are independently C,G or absent;

R₃₂,R₄₃,R₆₈=are independently C,G,U or absent;

R₁₁,R₁₆,R₂₈,R₃₁,R₄₉,R₅₉,R₆₅,R₇₀=are independently C,U or absent;

R₁,R₉,R₁₀,R₁₉,R₂₀,R₅₀,R₅₂,R₆₉=are independently G or absent;

R₅,R₈,R₂₉,R₃₀,R₄₂,R₅₁,R₆₄,R₆₆=are independently G,U or absent;

R₁₇,R₃₆,R₅₃,R₅₄=are independently U or absent;

[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 10 provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{TRP} (SEQ ID NO: 615),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 15 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
 R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Trp is:

R₀,R₁₈,R₂₂,R₂₃=absent
 20 R₁₄,R₂₄,R₃₉,R₄₁,R₅₇,R₇₂=are independently A or absent;
 R₃,R₄,R₁₃,R₆₁,R₇₁=are independently A,C or absent;
 R₆,R₄₄=are independently A,C,G or absent;
 R₂₁= A,C,U or absent;
 R₂,R₇,R₁₅,R₂₅,R₃₃,R₃₄,R₄₅,R₅₆,R₆₃=are independently A,G or absent;
 25 R₅₈= A,G,U or absent;
 R₄₆= A,U or absent;
 R₃₇,R₃₈,R₅₅,R₆₀,R₆₂=are independently C or absent;
 R₁₂,R₂₆,R₂₇,R₃₅,R₄₀,R₄₈,R₆₇=are independently C,G or absent;
 R₃₂,R₄₃,R₆₈=are independently C,G,U or absent;
 30 R₁₁,R₁₆,R₂₈,R₃₁,R₄₉,R₅₉,R₆₅,R₇₀=are independently C,U or absent;

$R_1, R_9, R_{10}, R_{19}, R_{20}, R_{50}, R_{52}, R_{69}$ are independently G or absent;

$R_5, R_8, R_{29}, R_{30}, R_{42}, R_{51}, R_{64}, R_{66}$ are independently G, U or absent;

$R_{17}, R_{36}, R_{53}, R_{54}$ are independently U or absent;

$[R_{47}]_x = N$ or absent;

- 5 wherein, *e.g.*, $x=1-271$ (*e.g.*, $x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or $x=271$), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.$

15 Tyrosine TREM Consensus sequence

In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{TYR} (SEQ ID NO: 616),

- $R_0 - R_1 - R_2 - R_3 - R_4 - R_5 - R_6 - R_7 - R_8 - R_9 - R_{10} - R_{11} - R_{12} - R_{13} - R_{14} - R_{15} - R_{16} - R_{17} - R_{18} - R_{19} - R_{20} - R_{21} - R_{22} - R_{23} - R_{24} - R_{25} - R_{26} - R_{27} - R_{28} - R_{29} - R_{30} - R_{31} - R_{32} - R_{33} - R_{34} - R_{35} - R_{36} - R_{37} - R_{38} - R_{39} - R_{40} - R_{41} - R_{42} - R_{43} - R_{44} - R_{45} - R_{46} - [R_{47}]_x - R_{48} - R_{49} - R_{50} - R_{51} - R_{52} - R_{53} - R_{54} - R_{55} - R_{56} - R_{57} - R_{58} - R_{59} - R_{60} - R_{61} - R_{62} - R_{63} - R_{64} - R_{65} - R_{66} - R_{67} - R_{68} - R_{69} - R_{70} - R_{71} - R_{72}$
- 20

wherein R is a ribonucleotide residue and the consensus for Tyr is:

R_0 = absent

R_{14}, R_{39}, R_{57} are independently A or absent;

- 25 $R_{41}, R_{48}, R_{51}, R_{71}$ are independently A, C, G or absent;

$R_3, R_4, R_5, R_6, R_9, R_{10}, R_{12}, R_{13}, R_{16}, R_{25}, R_{26}, R_{30}, R_{31}, R_{32}, R_{42}, R_{44}, R_{45}, R_{46}, R_{49}, R_{50}, R_{58}, R_{62}, R_{63}, R_{66}, R_{67}, R_{68}, R_{69}, R_{70}$ are independently N or absent;

R_{22}, R_{65} are independently A, C, U or absent;

$R_{15}, R_{24}, R_{27}, R_{33}, R_{37}, R_{40}, R_{56}$ are independently A, G or absent;

- 30 $R_7, R_{29}, R_{34}, R_{72}$ are independently A, G, U or absent;

R_{23}, R_{53} are independently A, U or absent;

R₃₅,R₆₀=are independently C or absent;

R₂₀= C,G or absent;

R₁,R₂,R₂₈,R₆₁,R₆₄=are independently C,G,U or absent;

R₁₁,R₁₇,R₂₁,R₄₃,R₅₅=are independently C,U or absent;

5 R₁₉,R₅₂=are independently G or absent;

R₈,R₁₈,R₃₆,R₃₈,R₅₄,R₅₉=are independently U or absent;

[R₄₇]_x = N or absent;

wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13, x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{TYR} (SEQ ID NO: 617),

20 R₀- R₁-R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Tyr is:

25 R₀,R₁₈,R₂₃=absent

R₇,R₉,R₁₄,R₂₄,R₂₆,R₃₄,R₃₉,R₅₇=are independently A or absent;

R₄₄,R₆₉=are independently A,C or absent;

R₇₁= A,C,G or absent;

R₆₈= N or absent;

30 R₅₈= A,C,U or absent;

- R₃₃,R₃₇,R₄₁,R₅₆,R₆₂,R₆₃=are independently A,G or absent;
 R₆,R₂₉,R₇₂=are independently A,G,U or absent;
 R₃₁,R₄₅,R₅₃=are independently A,U or absent;
 R₁₃,R₃₅,R₄₉,R₆₀=are independently C or absent;
 5 R₂₀,R₄₈,R₆₄,R₆₇,R₇₀=are independently C,G or absent;
 R₁,R₂,R₅,R₁₆,R₆₆=are independently C,G,U or absent;
 R₁₁,R₂₁,R₂₈,R₄₃,R₅₅,R₆₁=are independently C,U or absent;
 R₁₀,R₁₅,R₁₉,R₂₅,R₂₇,R₄₀,R₅₁,R₅₂=are independently G or absent;
 R₃,R₄,R₃₀,R₃₂,R₄₂,R₄₆=are independently G,U or absent;
 10 R₈,R₁₂,R₁₇,R₂₂,R₃₆,R₃₈,R₅₀,R₅₄,R₅₉,R₆₅=are independently U or absent;
 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 15 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 20 provided that the TREM has one or both of the following properties: no more than 15% of the
 residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{TYR} (SEQ ID NO: 618),

- 25 R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
 R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Tyr is:

- 30 R₀,R₁₈,R₂₃=absent

- R₇,R₉,R₁₄,R₂₄,R₂₆,R₃₄,R₃₉,R₅₇,R₇₂=are independently A or absent;
 R₄₄,R₆₉=are independently A,C or absent;
 R₇₁= A,C,G or absent;
 R₃₇,R₄₁,R₅₆,R₆₂,R₆₃=are independently A,G or absent;
 5 R₆,R₂₉,R₆₈=are independently A,G,U or absent;
 R₃₁,R₄₅,R₅₈=are independently A,U or absent;
 R₁₃,R₂₈,R₃₅,R₄₉,R₆₀,R₆₁=are independently C or absent;
 R₅,R₄₈,R₆₄,R₆₇,R₇₀=are independently C,G or absent;
 R₁,R₂=are independently C,G,U or absent;
 10 R₁₁,R₁₆,R₂₁,R₄₃,R₅₅,R₆₆=are independently C,U or absent;
 R₁₀,R₁₅,R₁₉,R₂₀,R₂₅,R₂₇,R₃₃,R₄₀,R₅₁,R₅₂=are independently G or absent;
 R₃,R₄,R₃₀,R₃₂,R₄₂,R₄₆=are independently G,U or absent;
 R₈,R₁₂,R₁₇,R₂₂,R₃₆,R₃₈,R₅₀,R₅₃,R₅₄,R₅₉,R₆₅=are independently U or absent;
 [R₄₇]_x = N or absent;
 15 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 20 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
 provided that the TREM has one or both of the following properties: no more than 15% of the
 residues are N; or no more than 20 residues are absent.
 25 Valine TREM Consensus sequence
 In an embodiment, a TREM disclosed herein comprises the sequence of Formula I_{VAL} (SEQ ID
 NO: 619),
 R₀- R₁- R₂- R₃-R₄ -R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂- R₄₃- R₄₄-R₄₅-

R₄₆- [R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Val is:

- R₀,R₂₃=absent;
- 5 R₂₄,R₃₈,R₅₇=are independently A or absent;
- R₉,R₇₂=are independently A,C,G or absent;
- R₂,R₄,R₅,R₆,R₇,R₁₂,R₁₅,R₁₆,R₂₁,R₂₅,R₂₆,R₂₉,R₃₁,R₃₂,R₃₃,R₃₄,R₃₇,R₄₁,R₄₂,R₄₃,R₄₄,R₄₅,R₄₆,R₄₈,R₄₉,R₅₀,R₅₈,R₆₁,R₆₂,R₆₃,R₆₄,R₆₅,R₆₆,R₆₇,R₆₈,R₆₉,R₇₀=are independently N or absent;
- R₁₇,R₃₅,R₅₉=are independently A,C,U or absent;
- 10 R₁₀,R₁₄,R₂₇,R₄₀,R₅₂,R₅₆=are independently A,G or absent;
- R₁,R₃,R₅₁,R₅₃=are independently A,G,U or absent;
- R₃₉= C or absent;
- R₁₃,R₃₀,R₅₅=are independently C,G,U or absent;
- R₁₁,R₂₂,R₂₈,R₆₀,R₇₁=are independently C,U or absent;
- 15 R₁₉= G or absent;
- R₂₀= G ,U or absent;
- R₈,R₁₈,R₃₆,R₅₄=are independently U or absent;
- [R₄₇]_x = N or absent;
- wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
20 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,
25 x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50,
x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271),
provided that the TREM has one or both of the following properties: no more than 15% of the
residues are N; or no more than 20 residues are absent.

In an embodiment, a TREM disclosed herein comprises the sequence of Formula II_{VAL} (SEQ ID NO: 620),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
 R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
 5 R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Val is:

R₀,R₁₈,R₂₃=absent;
 R₂₄,R₃₈,R₅₇=are independently A or absent;
 10 R₆₄,R₇₀,R₇₂=are independently A,C,G or absent;
 R₁₅,R₁₆,R₂₆,R₂₉,R₃₁,R₃₂,R₄₃,R₄₄,R₄₅,R₄₉,R₅₀,R₅₈,R₆₂,R₆₅=are independently N or absent;
 R₆,R₁₇,R₃₄,R₃₇,R₄₁,R₅₉=are independently A,C,U or absent;
 R₉,R₁₀,R₁₄,R₂₇,R₄₀,R₄₆,R₅₁,R₅₂,R₅₆=are independently A,G or absent;
 R₇,R₁₂,R₂₅,R₃₃,R₅₃,R₆₃,R₆₆,R₆₈=are independently A,G,U or absent;
 15 R₆₉= A,U or absent;
 R₃₉= C or absent;
 R₅,R₆₇=are independently C,G or absent;
 R₂,R₄,R₁₃,R₄₈,R₅₅,R₆₁=are independently C,G,U or absent;
 R₁₁,R₂₂,R₂₈,R₃₀,R₃₅,R₆₀,R₇₁=are independently C,U or absent;
 20 R₁₉= G or absent;
 R₁,R₃,R₂₀,R₄₂=are independently G,U or absent;
 R₈,R₂₁,R₃₆,R₅₄=are independently U or absent;
 [R₄₇]_x = N or absent;
 wherein, *e.g.*, x=1-271 (*e.g.*, x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125,
 25 x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24,
 x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,
 x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-
 271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1,
 x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17,

x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

5 In an embodiment, a TREM disclosed herein comprises the sequence of Formula III_{VAL} (SEQ ID NO: 621),

R₀-R₁-R₂-R₃-R₄-R₅-R₆-R₇-R₈-R₉-R₁₀-R₁₁-R₁₂-R₁₃-R₁₄-R₁₅-R₁₆-R₁₇-R₁₈-R₁₉-R₂₀-R₂₁-R₂₂-
R₂₃-R₂₄-R₂₅-R₂₆-R₂₇-R₂₈-R₂₉-R₃₀-R₃₁-R₃₂-R₃₃-R₃₄-R₃₅-R₃₆-R₃₇-R₃₈-R₃₉-R₄₀-R₄₁-R₄₂-R₄₃-R₄₄-R₄₅-
R₄₆-[R₄₇]_x-R₄₈-R₄₉-R₅₀-R₅₁-R₅₂-R₅₃-R₅₄-R₅₅-R₅₆-R₅₇-R₅₈-R₅₉-R₆₀-R₆₁-R₆₂-R₆₃-R₆₄-R₆₅-R₆₆-R₆₇-
10 R₆₈-R₆₉-R₇₀-R₇₁-R₇₂

wherein R is a ribonucleotide residue and the consensus for Val is:

R₀,R₁₈,R₂₃=absent
R₂₄,R₃₈,R₄₀,R₅₇,R₇₂=are independently A or absent;
R₂₉,R₆₄,R₇₀=are independently A,C,G or absent;
15 R₄₉,R₅₀,R₆₂=are independently N or absent;
R₁₆,R₂₆,R₃₁,R₃₂,R₃₇,R₄₁,R₄₃,R₅₉,R₆₅=are independently A,C,U or absent;
R₉,R₁₄,R₂₇,R₄₆,R₅₂,R₅₆,R₆₆=are independently A,G or absent;
R₇,R₁₂,R₂₅,R₃₃,R₄₄,R₄₅,R₅₃,R₅₈,R₆₃,R₆₈=are independently A,G,U or absent;
R₆₉= A,U or absent;
20 R₃₉= C or absent;
R₅,R₆₇=are independently C,G or absent;
R₂,R₄,R₁₃,R₁₅,R₄₈,R₅₅=are independently C,G,U or absent;
R₆,R₁₁,R₂₂,R₂₈,R₃₀,R₃₄,R₃₅,R₆₀,R₆₁,R₇₁=are independently C,U or absent;
R₁₀,R₁₉,R₅₁=are independently G or absent;
25 R₁,R₃,R₂₀,R₄₂=are independently G,U or absent;
R₈,R₁₇,R₂₁,R₃₆,R₅₄=are independently U or absent;
[R₄₇]_x = N or absent;

wherein, e.g., x=1-271 (e.g., x=1-250, x=1-225, x=1-200, x=1-175, x=1-150, x=1-125, x=1-100, x=1-75, x=1-50, x=1-40, x=1-30, x=1-29, x=1-28, x=1-27, x=1-26, x=1-25, x=1-24, x=1-23, x=1-22, x=1-21, x=1-20, x=1-19, x=1-18, x=1-17, x=1-16, x=1-15, x=1-14, x=1-13,

x=1-12, x=1-11, x=1-10, x=10-271, x=20-271, x=30-271, x=40-271, x=50-271, x=60-271, x=70-271, x=80-271, x=100-271, x=125-271, x=150-271, x=175-271, x=200-271, x=225-271, x=1, x=2, x=3, x=4, x=5, x=6, x=7, x=8, x=9, x=10, x=11, x=12, x=13, x=14, x=15, x=16, x=17, x=18, x=19, x=20, x=21, x=22, x=23, x=24, x=25, x=26, x=27, x=28, x=29, x=30, x=40, x=50, 5 x=60, x=70, x=80, x=90, x=100, x=110, x=125, x=150, x=175, x=200, x=225, x=250, or x=271), provided that the TREM has one or both of the following properties: no more than 15% of the residues are N; or no more than 20 residues are absent.

Variable region consensus sequence

In an embodiment, a TREM disclosed herein comprises a variable region at position R₄₇.
 10 In an embodiment, the variable region is 1-271 ribonucleotides in length (*e.g.* 1-250, 1-225, 1-200, 1-175, 1-150, 1-125, 1-100, 1-75, 1-50, 1-40, 1-30, 1-29, 1-28, 1-27, 1-26, 1-25, 1-24, 1-23, 1-22, 1-21, 1-20, 1-19, 1-18, 1-17, 1-16, 1-15, 1-14, 1-13, 1-12, 1-11, 1-10, 10-271, 20-271, 30-271, 40-271, 50-271, 60-271, 70-271, 80-271, 100-271, 125-271, 150-271, 175-271, 200-271, 225-271, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 15 27, 28, 29, 30, 40, 50, 60, 70, 80, 90, 100, 110, 125, 150, 175, 200, 225, 250, or 271 ribonucleotides). In an embodiment, the variable region comprises any one, all or a combination of Adenine, Cytosine, Guanine or Uracil.

In an embodiment, the variable region comprises a ribonucleic acid (RNA) sequence encoded by a deoxyribonucleic acid (DNA) sequence disclosed in Table 15, *e.g.*, any one of SEQ
 20 ID NOs: 452-561 disclosed in Table 15.

Table 15: Exemplary variable region sequences.

	SEQ ID NO	SEQUENCE
1	452	AAAATATAAATATATTTC
2	453	AAGCT
3	454	AAGTT
4	455	AATTCTTCGGAATGT
5	456	AGA
6	457	AGTCC
7	458	CAACC
8	459	CAATC
9	460	CAGC
10	461	CAGGCGGGTTCTGCCC GCGC
11	462	CATACCTGCAAGGGTATC
12	463	CGACCGCAAGGTTGT
13	464	CGACCTTGCGGTCAT

14	465	CGATGCTAATCACATCGT
15	466	CGATGGTGACATCAT
16	467	CGATGGTTTACATCGT
17	468	CGCCGTAAGGTGT
18	469	CGCCTTAGGTGT
19	470	CGCCTTTCGACGCGT
20	471	CGCTTCACGGCGT
21	472	CGGCAGCAATGCTGT
22	473	CGGCTCCGCCTTC
23	474	CGGGTATCACAGGGTC
24	475	CGGTGCGCAAGCGCTGT
25	476	CGTACGGGTGACCGTACC
26	477	CGTCAAAGACTTC
27	478	CGTCGTAAGACTT
28	479	CGTTGAATAAACGT
29	480	CTGTC
30	481	GGCC
31	482	GGGGATT
32	483	GGTC
33	484	GGTTT
34	485	GTAG
35	486	TAACTAGATACTTTCAGAT
36	487	TACTCGTATGGGTGC
37	488	TACTTTGCGGTGT
38	489	TAGGCGAGTAACATCGTGC
39	490	TAGGCGTGAATAGCGCCTC
40	491	TAGGTGCGGAGAGCGGCGC
41	492	TAGGTGCGGTAAGCGGCGC
42	493	TAGGTGGTTATCCACGC
43	494	TAGTC
44	495	TAGTT
45	496	TATACGTGAAAGCGTATC
46	497	TATAGGGTCAAAAACCTCTATC
47	498	TATGCAGAAATACCTGCATC
48	499	TCCCCATACGGGGGC
49	500	TCCCGAAGGGGTTC
50	501	TCTACGTATGTGGGC
51	502	TCTCATAGGAGTTC
52	503	TCTCCTCTGGAGGC
53	504	TCTTAGCAATAAGGT
54	505	TCTTGTAGGAGTTC
55	506	TGAACGTAAGTTCGC
56	507	TGAACTGCGAGGTTCC

57	508	TGAC
58	509	TGACCGAAAGGTCGT
59	510	TGACCGCAAGGTCGT
60	511	TGAGCTCTGCTCTC
61	512	TGAGGCCTCACGGCCTAC
62	513	TGAGGGCAACTTCGT
63	514	TGAGGGTCATACCTCC
64	515	TGAGGGTGCAAATCCTCC
65	516	TGCCGAAAGGCGT
66	517	TGCCGTAAGGCGT
67	518	TGCGGTCTCCGCGC
68	519	TGCTAGAGCAT
69	520	TGCTCGTATAGAGCTC
70	521	TGGACAATTGTCTGC
71	522	TGGACAGATGTCCGT
72	523	TGGACAGGTGTCCGC
73	524	TGGACGGTTGTCCGC
74	525	TGGACTTGTGGTC
75	526	TGGAGATTCTCTCCGC
76	527	TGGCATAGGCCTGC
77	528	TGGCTTATGTCTAC
78	529	TGGGAGTTAATCCCGT
79	530	TGGGATCTTCCCGC
80	531	TGGGCAGAAATGTCTC
81	532	TGGGCGTTCGCCCCG
82	533	TGGGCTTCGCCCCG
83	534	TGGGGGATAACCCCGT
84	535	TGGGGGTTTCCCCGT
85	536	TGGT
86	537	TGGTGGCAACACCGT
87	538	TGGTTTATAGCCGT
88	539	TGTACGGTAATACCGTACC
89	540	TGTCCGCAAGGACGT
90	541	TGTCCTAACGGACGT
91	542	TGTCCTATTAAACGGACGT
92	543	TGTCCTTCACGGGCGT
93	544	TGTCTTAGGACGT
94	545	TGTGCGTTAACGCGTACC
95	546	TGTGTCGCAAGGCACC
96	547	TGTTTCGTAAGGACTT
97	548	TTCACAGAAATGTGTC
98	549	TTCCCTCGTGGAGT
99	550	TTCCCTCTGGGAGC

100	551	TTCCCTTGTGGATC
101	552	TTCCCTTCGGGAGC
102	553	TTCTAGCAATAGAGT
103	554	TTCTCCACTGGGGAGC
104	555	TTCTCGAGAGGGAGC
105	556	TTCTCGTATGAGAGC
106	557	TTTAAGGTTTCCCTTAAC
107	558	TTTCATTGTGGAGT
108	559	TTTCGAAGGAATCC
109	560	TTTCTTCGGAAGC
110	561	TTTGGGGCAACTCAAC

Method of making TREMs, TREM core fragments, and TREM fragments

In vitro methods for synthesizing oligonucleotides are known in the art and can be used to make a TREM, a TREM core fragment or a TREM fragment disclosed herein. For example, a TREM, TREM core fragment or TREM fragment can be synthesized using a synthetic method, e.g., solid state synthesis or liquid phase synthesis. In an embodiment, a synthetic method of making a TREM, TREM core fragment or TREM fragment comprises linking a first nucleotide to a second nucleotide to form the TREM TREM core fragment or TREM fragment.

In an embodiment, a TREM, a TREM core fragment or a TREM fragment made according to an in vitro synthesis method disclosed herein has a different modification profile compared to a TREM expressed and isolated from a cell, or compared to a naturally occurring tRNA.

An exemplary method for making a synthetic TREM via 5'-Silyl-2'-Orthoester (2'-ACE) Chemistry is provided in Example 3. The method provided in Example 3 can also be used to make a synthetic TREM core fragment or synthetic TREM fragment. Additional synthetic methods are disclosed in Hartsel SA et al., (2005) *Oligonucleotide Synthesis*, 033–050, the entire contents of which are hereby incorporated by reference.

TREM composition

In an embodiment, a TREM composition, e.g., a TREM pharmaceutical composition, comprises a pharmaceutically acceptable excipient. Exemplary excipients include those provided in the FDA Inactive Ingredient Database (<https://www.accessdata.fda.gov/scripts/cder/iig/index.Cfm>).

In an embodiment, a TREM composition, *e.g.*, a TREM pharmaceutical composition, comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100 or 150 grams of TREM, TREM core fragment or TREM fragment. In an embodiment, a TREM composition, *e.g.*, a TREM pharmaceutical composition, comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50 or 100 milligrams of TREM, TREM core fragment or TREM fragment.

In an embodiment, a TREM composition, *e.g.*, a TREM pharmaceutical composition, is at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 95 or 99% dry weight TREMs, TREM core fragments or TREM fragments.

In an embodiment, a TREM composition comprises at least 1×10^6 TREM molecules, at least 1×10^7 TREM molecules, at least 1×10^8 TREM molecules or at least 1×10^9 TREM molecules.

In an embodiment, a TREM composition comprises at least 1×10^6 TREM core fragment molecules, at least 1×10^7 TREM core fragment molecules, at least 1×10^8 TREM core fragment molecules or at least 1×10^9 TREM core fragment molecules.

In an embodiment, a TREM composition comprises at least 1×10^6 TREM fragment molecules, at least 1×10^7 TREM fragment molecules, at least 1×10^8 TREM fragment molecules or at least 1×10^9 TREM fragment molecules.

In an embodiment, a TREM composition produced by any of the methods of making disclosed herein can be charged with an amino acid using an in vitro charging reaction as known in the art.

In an embodiment, a TREM composition comprise one or more species of TREMs, TREM core fragments, or TREM fragments. In an embodiment, a TREM composition comprises a single species of TREM, TREM core fragment, or TREM fragment. In an embodiment, a TREM composition comprises a first TREM, TREM core fragment, or TREM fragment species and a second TREM, TREM core fragment, or TREM fragment species. In an embodiment, the TREM composition comprises X TREM, TREM core fragment, or TREM fragment species, wherein X=2, 3, 4, 5, 6, 7, 8, 9, or 10.

In an embodiment, the TREM, TREM core fragment, or TREM fragment has at least 70, 75, 80, 85, 90, or 95, or has 100%, identity with a sequence encoded by a nucleic acid in Table 9.

In an embodiment, the TREM comprises a consensus sequence provided herein.

A TREM composition can be formulated as a liquid composition, as a lyophilized composition or as a frozen composition.

In some embodiments, a TREM composition can be formulated to be suitable for pharmaceutical use, e.g., a pharmaceutical TREM composition. In an embodiment, a
5 pharmaceutical TREM composition is substantially free of materials and/or reagents used to separate and/or purify a TREM, TREM core fragment, or TREM fragment.

In some embodiments, a TREM composition can be formulated with water for injection. In some embodiments, a TREM composition formulated with water for injection is suitable for pharmaceutical use, e.g., comprises a pharmaceutical TREM composition.

10

TREM characterization

A TREM, TREM core fragment, or TREM fragment, or a TREM composition, e.g., a pharmaceutical TREM composition, produced by any of the methods disclosed herein can be assessed for a characteristic associated with the TREM, TREM core fragment, or TREM
15 fragment or the TREM composition, such as purity, sterility, concentration, structure, or functional activity of the TREM, TREM core fragment, or TREM fragment. Any of the above-mentioned characteristics can be evaluated by providing a value for the characteristic, e.g., by evaluating or testing the TREM, TREM core fragment, or TREM fragment, or the TREM composition, or an intermediate in the production of the TREM composition. The value can also
20 be compared with a standard or a reference value. Responsive to the evaluation, the TREM composition can be classified, e.g., as ready for release, meets production standard for human trials, complies with ISO standards, complies with cGMP standards, or complies with other pharmaceutical standards. Responsive to the evaluation, the TREM composition can be subjected to further processing, e.g., it can be divided into aliquots, e.g., into single or multi-
25 dosage amounts, disposed in a container, e.g., an end-use vial, packaged, shipped, or put into commerce. In embodiments, in response to the evaluation, one or more of the characteristics can be modulated, processed or re-processed to optimize the TREM composition. For example, the TREM composition can be modulated, processed or re-processed to (i) increase the purity of the TREM composition; (ii) decrease the amount of fragments in the composition; (iii) decrease the
30 amount of endotoxins in the composition; (iv) increase the in vitro translation activity of the composition; (v) increase the TREM concentration of the composition; or (vi) inactivate or

remove any viral contaminants present in the composition, *e.g.*, by reducing the pH of the composition or by filtration.

In an embodiment, the TREM, TREM core fragment, or TREM fragment (*e.g.*, TREM composition or an intermediate in the production of the TREM composition) has a purity of at least 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%, *i.e.*, by mass.

In an embodiment, the TREM (*e.g.*, TREM composition or an intermediate in the production of the TREM composition) has less than 0.1%, 0.5%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25% TREM fragments relative to full length TREMs.

In an embodiment, the TREM, TREM core fragment, or TREM fragment (*e.g.*, TREM composition or an intermediate in the production of the TREM composition) has low levels or absence of endotoxins, *e.g.*, a negative result as measured by the Limulus amoebocyte lysate (LAL) test.

In an embodiment, the TREM, TREM core fragment, or TREM fragment (*e.g.*, TREM composition or an intermediate in the production of the TREM composition) has in-vitro translation activity, *e.g.*, as measured by an assay described in Examples 12-13.

In an embodiment, the TREM, TREM core fragment, or TREM fragment (*e.g.*, TREM composition or an intermediate in the production of the TREM composition) has a TREM concentration of at least 0.1 ng/mL, 0.5 ng/mL, 1 ng/mL, 5 ng/mL, 10 ng/mL, 50 ng/mL, 0.1 ug/mL, 0.5 ug/mL, 1 ug/mL, 2 ug/mL, 5 ug/mL, 10 ug/mL, 20 ug/mL, 30 ug/mL, 40 ug/mL, 50 ug/mL, 60 ug/mL, 70 ug/mL, 80 ug/mL, 100 ug/mL, 200 ug/mL, 300 ug/mL, 500 ug/mL, 1000 ug/mL, 5000 ug/mL, 10,000 ug/mL, or 100,000 ug/mL.

In an embodiment, the TREM, TREM core fragment, or TREM fragment (*e.g.*, TREM composition or an intermediate in the production of the TREM composition) is sterile, *e.g.*, the composition or preparation supports the growth of fewer than 100 viable microorganisms as tested under aseptic conditions, the composition or preparation meets the standard of USP <71>, and/or the composition or preparation meets the standard of USP <85>.

In an embodiment, the TREM, TREM core fragment, or TREM fragment (*e.g.*, TREM composition or an intermediate in the production of the TREM composition) has an undetectable level of viral contaminants, *e.g.*, no viral contaminants. In an embodiment, any viral contaminant, *e.g.*, residual virus, present in the composition is inactivated or removed. In an

embodiment, any viral contaminant, *e.g.*, residual virus, is inactivated, *e.g.*, by reducing the pH of the composition. In an embodiment, any viral contaminant, *e.g.*, residual virus, is removed, *e.g.*, by filtration or other methods known in the field.

5 **TREM administration**

Any TREM composition or pharmaceutical composition described herein can be administered to a cell, tissue or subject, *e.g.*, by direct administration to a cell, tissue and/or an organ in vitro, ex-vivo or in vivo. In-vivo administration may be via, *e.g.*, by local, systemic and/or parenteral routes, for example intravenous, subcutaneous, intraperitoneal, intrathecal, 10 intramuscular, ocular, nasal, urogenital, intradermal, dermal, enteral, intravitreal, intracerebral, intrathecal, or epidural.

Vectors and Carriers

In some embodiments the TREM, TREM core fragment, or TREM fragment or TREM 15 composition described herein, is delivered to cells, *e.g.* mammalian cells or human cells, using a vector. The vector may be, *e.g.*, a plasmid or a viral vector. In some embodiments, delivery is *in vivo*, *in vitro*, *ex vivo*, or *in situ*. In some embodiments, the viral vector is an adeno associated virus (AAV) vector, a lentivirus vector, an adenovirus or an anellovector. In some embodiments, the system or components of the system are delivered to cells with a viral-like particle or a 20 virosome. In some embodiments, the delivery uses more than one virus, viral-like particle or virosome.

A TREM, a TREM composition or a pharmaceutical TREM composition described herein may comprise, may be formulated with, or may be delivered in, a carrier.

Viral vectors

25 The carrier may be a viral vector (*e.g.*, a viral vector comprising a sequence encoding a TREM, a TREM core fragment or a TREM fragment). The viral vector may be administered to a cell or to a subject (*e.g.*, a human subject or animal model) to deliver a TREM, a TREM core fragment or a TREM fragment, a TREM composition or a pharmaceutical TREM composition.

A viral vector may be systemically or locally administered (*e.g.*, injected). Viral genomes 30 provide a rich source of vectors that can be used for the efficient delivery of exogenous genes into a mammalian cell. Viral genomes are known in the art as useful vectors for delivery

because the polynucleotides contained within such genomes are typically incorporated into the nuclear genome of a mammalian cell by generalized or specialized transduction. These processes occur as part of the natural viral replication cycle, and do not require added proteins or reagents in order to induce gene integration. Examples of viral vectors include a retrovirus (*e.g.*,
5 Retroviridae family viral vector), adenovirus (*e.g.*, Ad5, Ad26, Ad34, Ad35, and Ad48), parvovirus (*e.g.*, adeno-associated viruses), coronavirus, negative strand RNA viruses such as orthomyxovirus (*e.g.*, influenza virus), rhabdovirus (*e.g.*, rabies and vesicular stomatitis virus), paramyxovirus (*e.g.*, measles and Sendai), positive strand RNA viruses, such as picornavirus and alphavirus, and double stranded DNA viruses including adenovirus, herpesvirus (*e.g.*, Herpes
10 Simplex virus types 1 and 2, Epstein-Barr virus, cytomegalovirus, replication deficient herpes virus), and poxvirus (*e.g.*, vaccinia, modified vaccinia Ankara (MVA), fowlpox and canarypox). Other viruses include Norwalk virus, togavirus, flavivirus, reoviruses, papovavirus, hepadnavirus, human papilloma virus, human foamy virus, and hepatitis virus, for example. Examples of retroviruses include: avian leukemia-sarcoma, avian C-type viruses, mammalian C-
15 type, B-type viruses, D-type viruses, oncoretroviruses, HTLV-BLV group, lentivirus, alpharetrovirus, gammaretrovirus, spumavirus (Coffin, J. M., *Retroviridae: The viruses and their replication*, Virology (Third Edition) Lippincott-Raven, Philadelphia, 1996). Other examples include murine leukemia viruses, murine sarcoma viruses, mouse mammary tumor virus, bovine leukemia virus, feline leukemia virus, feline sarcoma virus, avian leukemia virus, human T-cell
20 leukemia virus, baboon endogenous virus, Gibbon ape leukemia virus, Mason Pfizer monkey virus, simian immunodeficiency virus, simian sarcoma virus, Rous sarcoma virus and lentiviruses. In some embodiments, a viral vector is used which does not integrate into the genome, *e.g.*, an anellovector (see, *e.g.*, US20200188456). Other examples of vectors are described, for example, in US Patent No. 5,801,030, the teachings of which are incorporated
25 herein by reference. In some embodiments the system or components of the system are delivered to cells with a viral-like particle or a virosome.

Cell and vesicle-based carriers

A TREM, a TREM core fragment or a TREM fragment, a TREM composition or a
30 pharmaceutical TREM composition described herein can be administered to a cell in a vesicle or other membrane-based carrier.

In embodiments, a TREM, a TREM core fragment or a TREM fragment, or TREM composition, or pharmaceutical TREM composition described herein is administered in or via a cell, vesicle or other membrane-based carrier. In one embodiment, the TREM, TREM core fragment, TREM fragment, or TREM composition or pharmaceutical TREM composition can be formulated in liposomes or other similar vesicles. Liposomes are spherical vesicle structures composed of a uni- or multilamellar lipid bilayer surrounding internal aqueous compartments and a relatively impermeable outer lipophilic phospholipid bilayer. Liposomes may be anionic, neutral or cationic. Liposomes are biocompatible, nontoxic, can deliver both hydrophilic and lipophilic drug molecules, protect their cargo from degradation by plasma enzymes, and transport their load across biological membranes and the blood brain barrier (BBB) (see, *e.g.*, Spuch and Navarro, *Journal of Drug Delivery*, vol. 2011, Article ID 469679, 12 pages, 2011. doi:10.1155/2011/469679 for review).

Vesicles can be made from several different types of lipids; however, phospholipids are most commonly used to generate liposomes as drug carriers. Methods for preparation of multilamellar vesicle lipids are known in the art (see for example U.S. Pat. No. 6,693,086, the teachings of which relating to multilamellar vesicle lipid preparation are incorporated herein by reference). Although vesicle formation can be spontaneous when a lipid film is mixed with an aqueous solution, it can also be expedited by applying force in the form of shaking by using a homogenizer, sonicator, or an extrusion apparatus (see, *e.g.*, Spuch and Navarro, *Journal of Drug Delivery*, vol. 2011, Article ID 469679, 12 pages, 2011. doi:10.1155/2011/469679 for review). Extruded lipids can be prepared by extruding through filters of decreasing size, as described in Templeton et al., *Nature Biotech*, 15:647-652, 1997, the teachings of which relating to extruded lipid preparation are incorporated herein by reference.

Lipid nanoparticles are another example of a carrier that provides a biocompatible and biodegradable delivery system for the TREM, TREM core fragment, TREM fragment, or TREM composition or pharmaceutical TREM composition described herein. Nanostructured lipid carriers (NLCs) are modified solid lipid nanoparticles (SLNs) that retain the characteristics of the SLN, improve drug stability and loading capacity, and prevent drug leakage. Polymer nanoparticles (PNPs) are an important component of drug delivery. These nanoparticles can effectively direct drug delivery to specific targets and improve drug stability and controlled drug release. Lipid-polymer nanoparticles (PLNs), a new type of carrier that combines liposomes and

polymers, may also be employed. These nanoparticles possess the complementary advantages of PNPs and liposomes. A PLN is composed of a core-shell structure; the polymer core provides a stable structure, and the phospholipid shell offers good biocompatibility. As such, the two components increase the drug encapsulation efficiency rate, facilitate surface modification, and prevent leakage of water-soluble drugs. For a review, see, *e.g.*, Li et al. 2017, *Nanomaterials* 7, 122; doi:10.3390/nano7060122.

Exemplary lipid nanoparticles are disclosed in International Application PCT/US2014/053907, the entire contents of which are hereby incorporated by reference. For example, an LNP described in paragraphs [403-406] or [410-413] of PCT/US2014/053907 can be used as a carrier for the TREM, TREM core fragment, TREM fragment, or TREM composition or pharmaceutical TREM composition described herein.

Additional exemplary lipid nanoparticles are disclosed in U.S. Patent 10,562,849 the entire contents of which are hereby incorporated by reference. For example, an LNP of formula (I) as described in columns 1-3 of U.S. Patent 10,562,849 can be used as a carrier for the TREM, TREM core fragment, TREM fragment, or TREM composition or pharmaceutical TREM composition described herein.

Lipids that can be used in nanoparticle formations (*e.g.*, lipid nanoparticles) include, for example those described in Table 4 of WO2019217941, which is incorporated by reference, *e.g.*, a lipid-containing nanoparticle can comprise one or more of the lipids in Table 4 of WO2019217941. Lipid nanoparticles can include additional elements, such as polymers, such as the polymers described in Table 5 of WO2019217941, incorporated by reference.

In some embodiments, conjugated lipids, when present, can include one or more of PEG-diacylglycerol (DAG) (such as 1-(monomethoxy-polyethyleneglycol)-2,3- dimyristoylglycerol (PEG-DMG)), PEG-dialkoxypentyl (DAA), PEG-phospholipid, PEG- ceramide (Cer), a pegylated phosphatidylethanolamine (PEG-PE), PEG succinate diacylglycerol (PEGS-DAG) (such as 4-0-(2,3-di(tetradecanoyloxy)propyl)-1-0-(w- methoxy(polyethoxy)ethyl) butanedioate (PEG-S-DMG)), PEG dialkoxypentylcarbam, N- (carbonyl-methoxypoly ethylene glycol 2000)-1,2-distearoyl-sn-glycero-3-phosphoethanolamine sodium salt, and those described in Table 2 of WO2019051289 (incorporated by reference), and combinations of the foregoing.


In some embodiments, sterols that can be incorporated into lipid nanoparticles include one or more of cholesterol or cholesterol derivatives, such as those in W02009/127060 or

US2010/0130588, which are incorporated by reference. Additional exemplary sterols include phytosterols, including those described in Eygeris et al (2020), incorporated herein by reference.


In some embodiments, the lipid particle comprises an ionizable lipid, a non-cationic lipid, a conjugated lipid that inhibits aggregation of particles, and a sterol. The amounts of these components can be varied independently and to achieve desired properties. For example, in some embodiments, the lipid nanoparticle comprises an ionizable lipid is in an amount from about 20 mol % to about 90 mol % of the total lipids (in other embodiments it may be 20-70% (mol), 30-60% (mol) or 40-50% (mol); about 50 mol % to about 90 mol % of the total lipid present in the lipid nanoparticle), a non-cationic lipid in an amount from about 5 mol % to about 30 mol % of the total lipids, a conjugated lipid in an amount from about 0.5 mol % to about 20 mol % of the total lipids, and a sterol in an amount from about 20 mol % to about 50 mol % of the total lipids. The ratio of total lipid to nucleic acid can be varied as desired. For example, the total lipid to nucleic acid (mass or weight) ratio can be from about 10: 1 to about 30: 1.

In some embodiments, the lipid to nucleic acid ratio (mass/mass ratio; w/w ratio) can be in the range of from about 1 : 1 to about 25: 1, from about 10: 1 to about 14: 1, from about 3 : 1 to about 15: 1, from about 4: 1 to about 10: 1, from about 5: 1 to about 9: 1, or about 6: 1 to about 9: 1. The amounts of lipids and nucleic acid can be adjusted to provide a desired N/P ratio, for example, N/P ratio of 3, 4, 5, 6, 7, 8, 9, 10 or higher. Generally, the lipid nanoparticle formulation's overall lipid content can range from about 5 mg/ml to about 30 mg/mL.


Some non-limiting example of lipid compounds that may be used (e.g., in combination with other lipid components) to form lipid nanoparticles for the delivery of compositions described herein, e.g., nucleic acid (e.g., RNA) described herein includes,



In some embodiments an LNP comprising Formula (i) is used to deliver a TREM composition described herein to the liver and/or hepatocyte cells.




In some embodiments an LNP comprising Formula (ii) is used to deliver a TREM composition described herein to the liver and/or hepatocyte cells.




(iii)

In some embodiments an LNP comprising Formula (iii) is used to deliver a TREM composition described herein to the liver and/or hepatocyte cells.




(iv)





(v)

In some embodiments an LNP comprising Formula (v) is used to deliver a TREM composition described herein to the liver and/or hepatocyte cells.




(vi)


In some embodiments an LNP comprising Formula (vi) is used to deliver a TREM composition described herein to the liver and/or hepatocyte cells.




In some embodiments an LNP comprising Formula (viii) is used to deliver a TREM composition described herein to the liver and/or hepatocyte cells.




In some embodiments an LNP comprising Formula (ix) is used to deliver a TREM composition described herein to the liver and/or hepatocyte cells.




- wherein X^1 is O, NR^1 , or a direct bond, X^2 is C2-5 alkylene, X^3 is $C(=O)$ or a direct bond, R^1 is H or Me, R^3 is C1-3 alkyl, R^2 is C1-3 alkyl, or R^2 taken together with the nitrogen atom to which it is attached and 1-3 carbon atoms of X^2 form a 4-, 5-, or 6-membered ring, or X^1 is NR^1 , R^1 and R^2 taken together with the nitrogen atoms to which they are attached form a 5- or 6-membered ring, or R^2 taken together with R^3 and the nitrogen atom to which they are attached form a 5-, 6-, or 7-membered ring, Y^1 is C2-12 alkylene, Y^2 is selected from





(in either orientation),
orientation),




(in either orientation),




(in either


- 5 n is 0 to 3, R⁴ is C1-15 alkyl, Z¹ is C1-6 alkylene or a direct bond, Z² is , (in either orientation) or absent, provided that if Z¹ is a direct bond, Z² is absent; R⁵ is C5-9 alkyl or C6-10 alkoxy, R⁶ is C5-9 alkyl or C6-10 alkoxy, W is methylene or a direct bond, and R⁷ is H or Me, or a salt thereof, provided that if R³ and R² are C2 alkyls, X¹ is O, X² is linear C3 alkylene, X³ is C(=O), Y¹ is linear C6 alkylene, (Y²)_n-R⁴ is , R⁴ is linear C5 alkyl, Z¹ is C2 alkylene, Z² is absent, W is methylene, and R⁷ is H, then R⁵ and R⁶ are not C_x alkoxy.


- 10 In some embodiments an LNP comprising Formula (xii) is used to deliver a TREM composition described herein to the liver and/or hepatocyte cells.




In some embodiments an LNP comprising Formula (xi) is used to deliver a TREM composition described herein to the liver and/or hepatocyte cells.

- 15  where R =  (xii)








In some embodiments an LNP comprises a compound of Formula (xiii) and a compound of Formula (xiv).





- 5 In some embodiments, an LNP comprising Formula (xv) is used to deliver a TREM composition described herein to the liver and/or hepatocyte cells.




In some embodiments an LNP comprising a formulation of Formula (xvi) is used to deliver a TREM composition described herein to the lung endothelial cells.





In some embodiments, a lipid compound used to form lipid nanoparticles for the delivery of compositions described herein, e.g., a TREM described herein is made by one of the following reactions:



In some embodiments, a composition described herein (e.g., TREM composition) is provided in an LNP that comprises an ionizable lipid. In some embodiments, the ionizable lipid is heptadecan-9-yl 8-((2-hydroxyethyl)(6-oxo-6-(undecyloxy)hexyl)amino)octanoate (SM-102); e.g., as described in Example 1 of US9,867,888 (incorporated by reference herein in its entirety). In some embodiments, the ionizable lipid is 9Z,12Z)-3-((4,4-bis(octyloxy)butanoyl)oxy)-2-(((3-(diethylamino)propoxy)carbonyl)oxy)methyl)propyl octadeca-9,12-dienoate (LP01), e.g., as synthesized in Example 13 of WO2015/095340 (incorporated by reference herein in its entirety). In some embodiments, the ionizable lipid is Di((Z)-non-2-en-1-yl) 9-((4-dimethylamino)-butanoyl)oxy)heptadecanedioate (L319), e.g. as synthesized in Example 7, 8, or 9 of US2012/0027803 (incorporated by reference herein in its entirety). In some embodiments, the ionizable lipid is 1,1-((2-(4-(2-((2-(Bis(2-hydroxydodecyl)amino)ethyl)(2-hydroxydodecyl)

amino)ethyl)piperazin-1-yl)ethyl)azanediyl)bis(dodecan-2-ol) (C12-200), e.g., as synthesized in Examples 14 and 16 of WO2010/053572 (incorporated by reference herein in its entirety). In some embodiments, the ionizable lipid is Imidazole cholesterol ester (ICE) lipid (3S, 10R, 13R, 17R)-10, 13-dimethyl-17- ((R)-6-methylheptan-2-yl)-2, 3, 4, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17-tetradecahydro-1H- cyclopenta[a]phenanthren-3-yl 3-(1H-imidazol-4-yl)propanoate, e.g., Structure (I) from WO2020/106946 (incorporated by reference herein in its entirety).

In some embodiments, an ionizable lipid may be a cationic lipid, an ionizable cationic lipid, e.g., a cationic lipid that can exist in a positively charged or neutral form depending on pH, or an amine-containing lipid that can be readily protonated. In some embodiments, the cationic lipid is a lipid capable of being positively charged, e.g., under physiological conditions. Exemplary cationic lipids include one or more amine group(s) which bear the positive charge. In some embodiments, the lipid particle comprises a cationic lipid in formulation with one or more of neutral lipids, ionizable amine-containing lipids, biodegradable alkyne lipids, steroids, phospholipids including polyunsaturated lipids, structural lipids (e.g., sterols), PEG, cholesterol and polymer conjugated lipids. In some embodiments, the cationic lipid may be an ionizable cationic lipid. An exemplary cationic lipid as disclosed herein may have an effective pKa over 6.0. In embodiments, a lipid nanoparticle may comprise a second cationic lipid having a different effective pKa (e.g., greater than the first effective pKa), than the first cationic lipid. A lipid nanoparticle may comprise between 40 and 60 mol percent of a cationic lipid, a neutral lipid, a steroid, a polymer conjugated lipid, and a therapeutic agent, e.g., a TREM described herein, encapsulated within or associated with the lipid nanoparticle. In some embodiments, the TREM is co-formulated with the cationic lipid. The TREM may be adsorbed to the surface of an LNP, e.g., an LNP comprising a cationic lipid. In some embodiments, the TREM may be encapsulated in an LNP, e.g., an LNP comprising a cationic lipid. In some embodiments, the lipid nanoparticle may comprise a targeting moiety, e.g., coated with a targeting agent. In embodiments, the LNP formulation is biodegradable. In some embodiments, a lipid nanoparticle comprising one or more lipid described herein, e.g., Formula (i), (ii), (iii), (vii) and/or (ix) encapsulates at least 1%, at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 92%, at least 95%, at least 97%, at least 98% or 100% of a TREM.

Exemplary ionizable lipids that can be used in lipid nanoparticle formulations include, without limitation, those listed in Table 1 of WO2019051289, incorporated herein by reference. Additional exemplary lipids include, without limitation, one or more of the following formulae: X of US2016/0311759; I of US20150376115 or in US2016/0376224; I, II or III of

5 US20160151284; I, IA, II, or IIA of US20170210967; I-c of US20150140070; A of US2013/0178541; I of US2013/0303587 or US2013/0123338; I of US2015/0141678; II, III, IV, or V of US2015/0239926; I of US2017/0119904; I or II of WO2017/117528; A of US2012/0149894; A of US2015/0057373; A of WO2013/116126; A of US2013/0090372; A of US2013/0274523; A of US2013/0274504; A of US2013/0053572; A of WO2013/016058; A of

10 WO2012/162210; I of US2008/042973; I, II, III, or IV of US2012/01287670; I or II of US2014/0200257; I, II, or III of US2015/0203446; I or III of US2015/0005363; I, IA, IB, IC, ID, II, IIA, IIB, IIC, IID, or III-XXIV of US2014/0308304; of US2013/0338210; I, II, III, or IV of WO2009/132131; A of US2012/01011478; I or XXXV of US2012/0027796; XIV or XVII of US2012/0058144; of US2013/0323269; I of US2011/0117125; I, II, or III of US2011/0256175;

15 I, II, III, IV, V, VI, VII, VIII, IX, X, XI, XII of US2012/0202871; I, II, III, IV, V, VI, VII, VIII, X, XII, XIII, XIV, XV, or XVI of US2011/0076335; I or II of US2006/008378; I of US2013/0123338; I or X-A-Y-Z of US2015/0064242; XVI, XVII, or XVIII of US2013/0022649; I, II, or III of US2013/0116307; I, II, or III of US2013/0116307; I or II of US2010/0062967; I-X of US2013/0189351; I of US2014/0039032; V of US2018/0028664; I of

20 US2016/0317458; I of US2013/0195920; 5, 6, or 10 of US10,221,127; III-3 of WO2018/081480; I-5 or I-8 of WO2020/081938; 18 or 25 of US9,867,888; A of US2019/0136231; II of WO2020/219876; 1 of US2012/0027803; OF-02 of US2019/0240349; 23 of US10,086,013; cKK-E12/A6 of Miao et al (2020); C12-200 of WO2010/053572; 7C1 of Dahlman et al (2017); 304-O13 or 503-O13 of Whitehead et al; TS-P4C2 of US9,708,628; I of WO2020/106946; I of

25 WO2020/106946.

In some embodiments, the ionizable lipid is MC3 (6Z,9Z,28Z,3 lZ)-heptatriaconta-6,9,28,3 l-tetraen-19-yl-4-(dimethylamino) butanoate (DLin-MC3-DMA or MC3), e.g., as described in Example 9 of WO2019051289A9 (incorporated by reference herein in its entirety). In some embodiments, the ionizable lipid is the lipid ATX-002, e.g., as described in Example 10

30 of WO2019051289A9 (incorporated by reference herein in its entirety). In some embodiments, the ionizable lipid is (13Z,16Z)-A,A-dimethyl-3- nonyldocosa-13, 16-dien-1-amine (Compound

32), e.g., as described in Example 11 of WO2019051289A9 (incorporated by reference herein in its entirety). In some embodiments, the ionizable lipid is Compound 6 or Compound 22, e.g., as described in Example 12 of WO2019051289A9 (incorporated by reference herein in its entirety).

Exemplary non-cationic lipids include, but are not limited to, distearoyl-sn-glycero-
 5 phosphoethanolamine, distearoylphosphatidylcholine (DSPC), dioleoylphosphatidylcholine (DOPC), dipalmitoylphosphatidylcholine (DPPC), dioleoylphosphatidylglycerol (DOPG), dipalmitoylphosphatidylglycerol (DPPG), dioleoyl-phosphatidylethanolamine (DOPE), palmitoyloleoylphosphatidylcholine (POPC), palmitoyloleoylphosphatidylethanolamine (POPE), dioleoyl-phosphatidylethanolamine 4-(N-maleimidomethyl)-cyclohexane- 1 - carboxylate
 10 (DOPE-mal), dipalmitoyl phosphatidyl ethanolamine (DPPE), dimyristoylphosphoethanolamine (DMPE), distearoyl-phosphatidyl-ethanolamine (DSPE), monomethyl-phosphatidylethanolamine (such as 16-O-monomethyl PE), dimethyl- phosphatidylethanolamine (such as 16-O-dimethyl PE), 18-l-trans PE, 1-stearoyl-2-oleoyl- phosphatidylethanolamine (SOPE), hydrogenated soy phosphatidylcholine (HSPC), egg phosphatidylcholine (EPC), dioleoylphosphatidylserine
 15 (DOPS), sphingomyelin (SM), dimyristoyl phosphatidylcholine (DMPC), dimyristoyl phosphatidylglycerol (DMPG), distearoylphosphatidylglycerol (DSPG), dierycoylphosphatidylcholine (DEPC), palmitoyloleoylphosphatidylglycerol (POPG), dielaidoyl-phosphatidylethanolamine (DEPE), lecithin, phosphatidylethanolamine, lysolecithin, lysophosphatidylethanolamine, phosphatidylserine, phosphatidylinositol, sphingomyelin, egg
 20 sphingomyelin (ESM), cephalin, cardiolipin, phosphatidicacid, cerebrosides, dicetylphosphate, lysophosphatidylcholine, dilinoleoylphosphatidylcholine, or mixtures thereof. It is understood that other diacylphosphatidylcholine and diacylphosphatidylethanolamine phospholipids can also be used. The acyl groups in these lipids are preferably acyl groups derived from fatty acids having C10-C24 carbon chains, e.g., lauroyl, myristoyl, palmitoyl, stearoyl, or oleoyl. Additional
 25 exemplary lipids, in certain embodiments, include, without limitation, those described in Kim et al. (2020) [dx.doi.org/10.1021/acs.nanolett.0c01386](https://doi.org/10.1021/acs.nanolett.0c01386), incorporated herein by reference. Such lipids include, in some embodiments, plant lipids found to improve liver transfection with mRNA (e.g., DGTS).

Other examples of non-cationic lipids suitable for use in the lipid nanoparticles include,
 30 without limitation, nonphosphorous lipids such as, e.g., stearylamine, dodecylamine, hexadecylamine, acetyl palmitate, glycerol ricinoleate, hexadecyl stearate, isopropyl myristate,

amphoteric acrylic polymers, triethanolamine-lauryl sulfate, alkyl-aryl sulfate polyethyloxylated fatty acid amides, dioctadecyl dimethyl ammonium bromide, ceramide, sphingomyelin, and the like. Other non-cationic lipids are described in WO2017/099823 or US patent publication US2018/0028664, the contents of which is incorporated herein by reference in their entirety.

5 In some embodiments, the non-cationic lipid is oleic acid or a compound of Formula I, II, or IV of US2018/0028664, incorporated herein by reference in its entirety. The non-cationic lipid can comprise, for example, 0-30% (mol) of the total lipid present in the lipid nanoparticle. In some embodiments, the non-cationic lipid content is 5-20% (mol) or 10-15% (mol) of the total lipid present in the lipid nanoparticle. In embodiments, the molar ratio of ionizable lipid to the
10 neutral lipid ranges from about 2:1 to about 8:1 (e.g., about 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, or 8:1).

In some embodiments, the lipid nanoparticles do not comprise any phospholipids.


In some aspects, the lipid nanoparticle can further comprise a component, such as a sterol, to provide membrane integrity. One exemplary sterol that can be used in the lipid nanoparticle is cholesterol and derivatives thereof. Non-limiting examples of cholesterol
15 derivatives include polar analogues such as 5 α -cholestanol, 5 β -coprostanol, choiesterolyl-(2-hydroxy)-ethyl ether, choiesterolyl-(4-hydroxy)-butyl ether, and 6-ketocholestanol; non-polar analogues such as 5 α -cholestane, cholestenone, 5 α -cholestanone, 5 β -cholestanone, and choiesterolyl decanoate; and mixtures thereof. In some embodiments, the cholesterol derivative is a polar analogue, e.g., choiesterolyl-(4-hydroxy)-butyl ether. Exemplary cholesterol derivatives
20 are described in PCT publication W02009/127060 and US patent publication US2010/0130588, each of which is incorporated herein by reference in its entirety.

In some embodiments, the component providing membrane integrity, such as a sterol, can comprise 0-50% (mol) (e.g., 0-10%, 10-20%, 20-30%, 30-40%, or 40-50%) of the total lipid present in the lipid nanoparticle. In some embodiments, such a component is 20-50% (mol) 30-
25 40% (mol) of the total lipid content of the lipid nanoparticle.

In some embodiments, the lipid nanoparticle can comprise a polyethylene glycol (PEG) or a conjugated lipid molecule. Generally, these are used to inhibit aggregation of lipid nanoparticles and/or provide steric stabilization. Exemplary conjugated lipids include, but are not limited to, PEG-lipid conjugates, polyoxazoline (POZ)-lipid conjugates, polyamide-lipid
30 conjugates (such as ATTA-lipid conjugates), cationic-polymer lipid (CPL) conjugates, and

mixtures thereof. In some embodiments, the conjugated lipid molecule is a PEG-lipid conjugate, for example, a (methoxy polyethylene glycol)-conjugated lipid.

Exemplary PEG-lipid conjugates include, but are not limited to, PEG-diacylglycerol (DAG) (such as 1-(monomethoxy-polyethyleneglycol)-2,3-dimyristoylglycerol (PEG-DMG)),
 5 PEG-dialkylxypropyl (DAA), PEG-phospholipid, PEG-ceramide (Cer), a pegylated phosphatidylethanolamine (PEG-PE), PEG succinate diacylglycerol (PEGS-DAG) (such as 4-0-(2,3-di(tetradecanoyloxy)propyl-1-0-(w-methoxy(polyethoxy)ethyl) butanedioate (PEG-S-DMG)), PEG dialkoxypopylcarbam, N-(carbonyl-methoxypolyethylene glycol 2000)-1,2-distearoyl-sn-glycero-3-phosphoethanolamine sodium salt, or a mixture thereof. Additional
 10 exemplary PEG-lipid conjugates are described, for example, in US5,885,613, US6,287,591, US2003/0077829, US2003/0077829, US2005/0175682, US2008/0020058, US2011/0117125, US2010/0130588, US2016/0376224, US2017/0119904, and US/099823, the contents of all of which are incorporated herein by reference in their entirety. In some embodiments, a PEG-lipid is a compound of Formula III, III-a-1, III-a-2, III-b-1, III-b-2, or V of US2018/0028664, the
 15 content of which is incorporated herein by reference in its entirety. In some embodiments, a PEG-lipid is of Formula II of US20150376115 or US2016/0376224, the content of both of which is incorporated herein by reference in its entirety. In some embodiments, the PEG-DAA conjugate can be, for example, PEG-dilauryloxypropyl, PEG- dimyristyloxypropyl, PEG-dipalmitoyloxypropyl, or PEG-distearoyloxypropyl. The PEG-lipid can be one or more of PEG-
 20 DMG, PEG-dilaurylglycerol, PEG-dipalmitoylglycerol, PEG- disteryllycerol, PEG-dilaurylglycamide, PEG-dimyristylglycamide, PEG- dipalmitoylglycamide, PEG-disterylglycamide, PEG-cholesterol (1-[8-(Cholest-5-en-3[beta]-oxy)carboxamido-3,6-dioxaoctanyl] carbamoyl-[omega]-methyl-poly(ethylene glycol), PEG- DMB (3,4-Ditetradecoxybenzyl- [omega]-methyl-poly(ethylene glycol) ether), and 1,2- dimyristoyl-sn-glycero-3-phosphoethanolamine-N-[methoxy(polyethylene glycol)-2000]. In some embodiments,
 25 the PEG-lipid comprises PEG-DMG, 1,2- dimyristoyl-sn-glycero-3-phosphoethanolamine-N-[methoxy(polyethylene glycol)-2000]. In some embodiments, the PEG-lipid comprises a structure selected from:



In some embodiments, lipids conjugated with a molecule other than a PEG can also be used in place of PEG-lipid. For example, polyoxazoline (POZ)-lipid conjugates, polyamide-lipid conjugates (such as ATTA-lipid conjugates), and cationic-polymer lipid (GPL) conjugates can be used in place of or in addition to the PEG-lipid.

Exemplary conjugated lipids, i.e., PEG-lipids, (POZ)-lipid conjugates, ATTA-lipid conjugates and cationic polymer-lipids are described in the PCT and LIS patent applications listed in Table 2 of WO2019051289A9, the contents of all of which are incorporated herein by reference in their entirety.

In some embodiments, the PEG or the conjugated lipid can comprise 0-20% (mol) of the total lipid present in the lipid nanoparticle. In some embodiments, PEG or the conjugated lipid content is 0.5- 10% or 2-5% (mol) of the total lipid present in the lipid nanoparticle. Molar ratios of the ionizable lipid, non-cationic-lipid, sterol, and PEG/conjugated lipid can be varied as needed. For example, the lipid particle can comprise 30-70% ionizable lipid by mole or by total weight of the composition, 0-60% cholesterol by mole or by total weight of the composition, 0-30% non-cationic-lipid by mole or by total weight of the composition and 1-10% conjugated lipid by mole or by total weight of the composition. Preferably, the composition comprises 30-40% ionizable lipid by mole or by total weight of the composition, 40-50% cholesterol by mole or by total weight of the composition, and 10- 20% non-cationic-lipid by mole or by total weight of the composition. In some other embodiments, the composition is 50-75% ionizable lipid by mole or by total weight of the composition, 20-40% cholesterol by mole or by total weight of the

composition, and 5 to 10% non-cationic-lipid, by mole or by total weight of the composition and 1-10% conjugated lipid by mole or by total weight of the composition. The composition may contain 60-70% ionizable lipid by mole or by total weight of the composition, 25-35% cholesterol by mole or by total weight of the composition, and 5-10% non-cationic-lipid by mole or by total weight of the composition. The composition may also contain up to 90% ionizable lipid by mole or by total weight of the composition and 2 to 15% non-cationic lipid by mole or by total weight of the composition. The formulation may also be a lipid nanoparticle formulation, for example comprising 8-30% ionizable lipid by mole or by total weight of the composition, 5-30% non-cationic lipid by mole or by total weight of the composition, and 0-20% cholesterol by mole or by total weight of the composition; 4-25% ionizable lipid by mole or by total weight of the composition, 4-25% non-cationic lipid by mole or by total weight of the composition, 2 to 25% cholesterol by mole or by total weight of the composition, 10 to 35% conjugate lipid by mole or by total weight of the composition, and 5% cholesterol by mole or by total weight of the composition; or 2-30% ionizable lipid by mole or by total weight of the composition, 2-30% non-cationic lipid by mole or by total weight of the composition, 1 to 15% cholesterol by mole or by total weight of the composition, 2 to 35% conjugate lipid by mole or by total weight of the composition, and 1-20% cholesterol by mole or by total weight of the composition; or even up to 90% ionizable lipid by mole or by total weight of the composition and 2-10% non-cationic lipids by mole or by total weight of the composition, or even 100% cationic lipid by mole or by total weight of the composition. In some embodiments, the lipid particle formulation comprises ionizable lipid, phospholipid, cholesterol and a PEG-ylated lipid in a molar ratio of 50: 10:38.5: 1.5. In some other embodiments, the lipid particle formulation comprises ionizable lipid, cholesterol and a PEG-ylated lipid in a molar ratio of 60:38.5: 1.5.

In some embodiments, the lipid particle comprises ionizable lipid, non-cationic lipid (e.g. phospholipid), a sterol (e.g., cholesterol) and a PEG-ylated lipid, where the molar ratio of lipids ranges from 20 to 70 mole percent for the ionizable lipid, with a target of 40-60, the mole percent of non-cationic lipid ranges from 0 to 30, with a target of 0 to 15, the mole percent of sterol ranges from 20 to 70, with a target of 30 to 50, and the mole percent of PEG-ylated lipid ranges from 1 to 6, with a target of 2 to 5.

In some embodiments, the lipid particle comprises ionizable lipid / non-cationic- lipid / sterol / conjugated lipid at a molar ratio of 50: 10:38.5: 1.5.

In an aspect, the disclosure provides a lipid nanoparticle formulation comprising phospholipids, lecithin, phosphatidylcholine and phosphatidylethanolamine.

In some embodiments, one or more additional compounds can also be included. Those compounds can be administered separately, or the additional compounds can be included in the lipid nanoparticles of the invention. In other words, the lipid nanoparticles can contain other compounds in addition to the nucleic acid or at least a second nucleic acid, different than the first. Without limitations, other additional compounds can be selected from the group consisting of small or large organic or inorganic molecules, monosaccharides, disaccharides, trisaccharides, oligosaccharides, polysaccharides, peptides, proteins, peptide analogs and derivatives thereof, peptidomimetics, nucleic acids, nucleic acid analogs and derivatives, an extract made from biological materials, or any combinations thereof.

In some embodiments, LNPs are directed to specific tissues by the addition of targeting domains. For example, biological ligands may be displayed on the surface of LNPs to enhance interaction with cells displaying cognate receptors, thus driving association with and cargo delivery to tissues wherein cells express the receptor. In some embodiments, the biological ligand may be a ligand that drives delivery to the liver, e.g., LNPs that display GalNAc result in delivery of nucleic acid cargo to hepatocytes that display asialoglycoprotein receptor (ASGPR). The work of Akinc et al. *Mol Ther* 18(7):1357-1364 (2010) teaches the conjugation of a trivalent GalNAc ligand to a PEG-lipid (GalNAc-PEG-DSG) to yield LNPs dependent on ASGPR for observable LNP cargo effect (see, e.g., FIG. 6 of Akinc et al. 2010, *supra*). Other ligand-displaying LNP formulations, e.g., incorporating folate, transferrin, or antibodies, are discussed in WO2017223135, which is incorporated herein by reference in its entirety, in addition to the references used therein, namely Kolhatkar et al., *Curr Drug Discov Technol.* 2011 8:197-206; Musacchio and Torchilin, *Front Biosci.* 2011 16:1388-1412; Yu et al., *Mol Membr Biol.* 2010 27:286-298; Patil et al., *Crit Rev Ther Drug Carrier Syst.* 2008 25:1-61 ; Benoit et al., *Biomacromolecules.* 2011 12:2708-2714; Zhao et al., *Expert Opin Drug Deliv.* 2008 5:309-319; Akinc et al., *Mol Ther.* 2010 18:1357-1364; Srinivasan et al., *Methods Mol Biol.* 2012 820:105-116; Ben-Arie et al., *Methods Mol Biol.* 2012 757:497-507; Peer 2010 *J Control Release.* 20:63-68; Peer et al., *Proc Natl Acad Sci U S A.* 2007 104:4095-4100; Kim et al., *Methods Mol Biol.* 2011 721:339-353; Subramanya et al., *Mol Ther.* 2010 18:2028-2037; Song et al., *Nat*

Biotechnol. 2005 23:709-717; Peer et al., Science. 2008 319:627-630; and Peer and Lieberman, Gene Ther. 2011 18:1127-1133.

In some embodiments, LNPs are selected for tissue-specific activity by the addition of a Selective ORgan Targeting (SORT) molecule to a formulation comprising traditional
 5 components, such as ionizable cationic lipids, amphipathic phospholipids, cholesterol and poly(ethylene glycol) (PEG) lipids. The teachings of Cheng et al. Nat Nanotechnol 15(4):313-320 (2020) demonstrate that the addition of a supplemental "SORT" component precisely alters the in vivo RNA delivery profile and mediates tissue-specific (e.g., lungs, liver, spleen) gene delivery and editing as a function of the percentage and biophysical property of the SORT
 10 molecule.

In some embodiments, the LNPs comprise biodegradable, ionizable lipids. In some embodiments, the LNPs comprise (9Z,12Z)-3-((4,4-bis(octyloxy)butanoyl)oxy)-2-(((3-(diethylamino)propoxy)carbonyl)oxy)methyl)propyl octadeca-9,12-dienoate, also called 3-((4,4-bis(octyloxy)butanoyl)oxy)-2-(((3-(diethylamino)propoxy)carbonyl)oxy)methyl)propyl
 15 (9Z,12Z)-octadeca-9,12-dienoate) or another ionizable lipid. See, e.g, lipids of WO2019/067992, WO/2017/173054, WO2015/095340, and WO2014/136086, as well as references provided therein. In some embodiments, the term cationic and ionizable in the context of LNP lipids is interchangeable, e.g., wherein ionizable lipids are cationic depending on the pH.

In some embodiments, the average LNP diameter of the LNP formulation may be
 20 between 10s of nm and 100s of nm, e.g., measured by dynamic light scattering (DLS). In some embodiments, the average LNP diameter of the LNP formulation may be from about 40 nm to about 150 nm, such as about 40 nm, 45 nm, 50 nm, 55 nm, 60 nm, 65 nm, 70 nm, 75 nm, 80 nm, 85 nm, 90 nm, 95 nm, 100 nm, 105 nm, 110 nm, 115 nm, 120 nm, 125 nm, 130 nm, 135 nm, 140 nm, 145 nm, or 150 nm. In some embodiments, the average LNP diameter of the LNP
 25 formulation may be from about 50 nm to about 100 nm, from about 50 nm to about 90 nm, from about 50 nm to about 80 nm, from about 50 nm to about 70 nm, from about 50 nm to about 60 nm, from about 60 nm to about 100 nm, from about 60 nm to about 90 nm, from about 60 nm to about 80 nm, from about 60 nm to about 70 nm, from about 70 nm to about 100 nm, from about 70 nm to about 90 nm, from about 70 nm to about 80 nm, from about 80 nm to about 100 nm,
 30 from about 80 nm to about 90 nm, or from about 90 nm to about 100 nm. In some embodiments, the average LNP diameter of the LNP formulation may be from about 70 nm to about 100 nm. In

a particular embodiment, the average LNP diameter of the LNP formulation may be about 80 nm. In some embodiments, the average LNP diameter of the LNP formulation may be about 100 nm. In some embodiments, the average LNP diameter of the LNP formulation ranges from about 1 mm to about 500 mm, from about 5 mm to about 200 mm, from about 10 mm to about 100 mm, from about 20 mm to about 80 mm, from about 25 mm to about 60 mm, from about 30 mm to about 55 mm, from about 35 mm to about 50 mm, or from about 38 mm to about 42 mm.

A LNP may, in some instances, be relatively homogenous. A polydispersity index may be used to indicate the homogeneity of a LNP, e.g., the particle size distribution of the lipid nanoparticles. A small (e.g., less than 0.3) polydispersity index generally indicates a narrow particle size distribution. A LNP may have a polydispersity index from about 0 to about 0.25, such as 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.10, 0.11, 0.12, 0.13, 0.14, 0.15, 0.16, 0.17, 0.18, 0.19, 0.20, 0.21, 0.22, 0.23, 0.24, or 0.25. In some embodiments, the polydispersity index of a LNP may be from about 0.10 to about 0.20.

The zeta potential of a LNP may be used to indicate the electrokinetic potential of the composition. In some embodiments, the zeta potential may describe the surface charge of an LNP. Lipid nanoparticles with relatively low charges, positive or negative, are generally desirable, as more highly charged species may interact undesirably with cells, tissues, and other elements in the body. In some embodiments, the zeta potential of a LNP may be from about -10 mV to about +20 mV, from about -10 mV to about +15 mV, from about -10 mV to about +10 mV, from about -10 mV to about +5 mV, from about -10 mV to about 0 mV, from about -10 mV to about -5 mV, from about -5 mV to about +20 mV, from about -5 mV to about +15 mV, from about -5 mV to about +10 mV, from about -5 mV to about +5 mV, from about -5 mV to about 0 mV, from about 0 mV to about +20 mV, from about 0 mV to about +15 mV, from about 0 mV to about +10 mV, from about 0 mV to about +5 mV, from about +5 mV to about +20 mV, from about +5 mV to about +15 mV, or from about +5 mV to about +10 mV.

The efficiency of encapsulation of a TREM describes the amount of TREM that is encapsulated or otherwise associated with a LNP after preparation, relative to the initial amount provided. The encapsulation efficiency is desirably high (e.g., close to 100%). The encapsulation efficiency may be measured, for example, by comparing the amount of TREM in a solution containing the lipid nanoparticle before and after breaking up the lipid nanoparticle with one or more organic solvents or detergents. An anion exchange resin may be used to measure the

amount of free protein or nucleic acid (e.g., RNA) in a solution. Fluorescence may be used to measure the amount of free TREM in a solution. For the lipid nanoparticles described herein, the encapsulation efficiency of a TREM may be at least 50%, for example 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%. In
5 some embodiments, the encapsulation efficiency may be at least 80%. In some embodiments, the encapsulation efficiency may be at least 90%. In some embodiments, the encapsulation efficiency may be at least 95%.

A LNP may optionally comprise one or more coatings. In some embodiments, a LNP may be formulated in a capsule, film, or tablet having a coating. A capsule, film, or tablet
10 including a composition described herein may have any useful size, tensile strength, hardness or density.

Additional exemplary lipids, formulations, methods, and characterization of LNPs are taught by WO2020061457, which is incorporated herein by reference in its entirety.

In some embodiments, in vitro or ex vivo cell lipofections are performed using
15 Lipofectamine MessengerMax (Thermo Fisher) or TransIT-mRNA Transfection Reagent (Mirus Bio). In certain embodiments, LNPs are formulated using the GenVoy_ILM ionizable lipid mix (Precision NanoSystems). In certain embodiments, LNPs are formulated using 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA) or dilinoleylmethyl-4-dimethylaminobutyrate (DLin-MC3-DMA or MC3), the formulation and in vivo use of which
20 are taught in Jayaraman et al. Angew Chem Int Ed Engl 51(34):8529-8533 (2012), incorporated herein by reference in its entirety.

LNP formulations optimized for the delivery of CRISPR-Cas systems, e.g., Cas9-gRNA RNP, gRNA, Cas9 mRNA, are described in WO2019067992 and WO2019067910, both incorporated by reference.

25 Additional specific LNP formulations useful for delivery of nucleic acids are described in US8158601 and US8168775, both incorporated by reference, which include formulations used in patisiran, sold under the name ONPATPRO.

Exosomes can also be used as drug delivery vehicles for the TREM, TREM core fragment, TREM fragment, or TREM compositions or pharmaceutical TREM composition
30 described herein. For a review, see Ha et al. July 2016. Acta Pharmaceutica Sinica B. Volume 6, Issue 4, Pages 287-296; <https://doi.org/10.1016/j.apsb.2016.02.001>.

Ex vivo differentiated red blood cells can also be used as a carrier for a TREM, TREM core fragment, TREM fragment, or TREM composition, or pharmaceutical TREM composition described herein. See, *e.g.*, WO2015073587; WO2017123646; WO2017123644; WO2018102740; WO2016183482; WO2015153102; WO2018151829; WO2018009838; Shi et al. 2014. *Proc Natl Acad Sci USA*. 111(28): 10131–10136; US Patent 9,644,180; Huang et al. 2017. *Nature Communications* 8: 423; Shi et al. 2014. *Proc Natl Acad Sci USA*. 111(28): 10131–10136.

Fusosome compositions, *e.g.*, as described in WO2018208728, can also be used as carriers to deliver the TREM, TREM core fragment, TREM fragment, or TREM composition, or pharmaceutical TREM composition described herein.

Virosomes and virus-like particles (VLPs) can also be used as carriers to deliver a TREM, TREM core fragment, TREM fragment, or TREM composition, or pharmaceutical TREM composition described herein to targeted cells.

Plant nanovesicles, *e.g.*, as described in WO2011097480A1, WO2013070324A1, or WO2017004526A1 can also be used as carriers to deliver the TREM, TREM core fragment, TREM fragment, or TREM composition, or pharmaceutical TREM composition described herein.

Delivery without a carrier

A TREM, a TREM core fragment or a TREM fragment, a TREM composition or a pharmaceutical TREM composition described herein can be administered to a cell without a carrier, *e.g.*, via naked delivery of the TREM, a TREM core fragment or a TREM fragment, a TREM composition or a pharmaceutical TREM composition.

In some embodiments, naked delivery as used herein refers to delivery without a carrier. In some embodiments, delivery without a carrier, *e.g.*, naked delivery, comprises delivery with a moiety, *e.g.*, a targeting peptide.

In some embodiments, a TREM, a TREM core fragment or a TREM fragment, or TREM composition, or pharmaceutical TREM composition described herein is delivered to a cell without a carrier, *e.g.*, via naked delivery. In some embodiments, the delivery without a carrier, *e.g.*, naked delivery, comprises delivery with a moiety, *e.g.*, a targeting peptide.

Enumerated Embodiments

1. A TREM comprising a sequence of Formula A:

- 5 [L1]-[ASt Domain1]-[L2]-[DH Domain]-[L3]-[ACH Domain] -[VL Domain]-[TH Domain]-[L4]-[ASt Domain2],
 wherein:
 independently, [L1] and [VL Domain], are optional;
 one of [L1], [ASt Domain1], [L2]-[DH Domain], [L3], [ACH Domain], [VL Domain],
 10 [TH Domain], [L4], and [ASt Domain2] comprises a nucleotide having a non-naturally occurring modification; and
 wherein:
 (a) the TREM retains the ability to: support protein synthesis, be charged by a synthetase, be bound by an elongation factor, introduce an amino acid into a peptide chain,
 15 support elongation, or support initiation;
 (b) the TREM comprises at least X contiguous nucleotides without a non-naturally occurring modification, wherein X is greater than 10;
 (c) at least 3, but less than all of the nucleotides of a type (e.g., A, T, C, G or U) comprise the same non-naturally occurring modification;
 20 (d) at least X nucleotides of a type (e.g., A, T, C, G or U) do not comprise a non-naturally occurring modification, wherein X=1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 or 50;
 (e) no more than 5, 10, or 15 nucleotides of a type (e.g., A, T, C, G or U) comprise a non-naturally occurring modification; and/or
 25 (f) no more than 5, 10, or 15 nucleotides of a type (e.g., A, T, C, G or U) do not comprise a non-naturally occurring modification.

2. The TREM of embodiment 1, comprising the feature provided in embodiment 1(a).

30

3. The TREM of embodiment 1, comprising the feature provided in embodiment 1(b).

4. The TREM of embodiment 1, comprising the feature provided in embodiment 1(c).
5. The TREM of embodiment 1, comprising the feature provided in embodiment 1(d).
- 5 6. The TREM of embodiment 1, comprising the feature provided in embodiment 1(e).
7. The TREM of embodiment 1, comprising the feature provided in embodiment 1(f).
8. The TREM of embodiment 1, comprising all of the features provided in embodiments 1(a)-(f).
- 10 9. The TREM of any one of embodiments 1-8, wherein the Domain comprising the non-naturally occurring modification retains a function, e.g., a domain function described herein.
10. The TREM of any one of embodiments 1-8, comprising an [L1].
- 15 11. The TREM of any one of embodiments 1-8, comprising a [VL Domain].
12. The TREM of any one of embodiments 1-8, wherein: [L1] is a linker comprising a nucleotide having a non-naturally occurring modification.
- 20 13. The TREM of any one of embodiments 1-8, wherein [ASt Domain1 (AstD1)] comprises a nucleotide having a non-naturally occurring modification.
14. The TREM of any one of embodiments 1-8, wherein [L2] is a linker comprising a nucleotide having a non-naturally occurring modification.
- 25 15. The TREM of any one of embodiments 1-8, wherein [DH Domain (DHD)] comprises a nucleotide having a non-naturally occurring modification.
- 30 16. The TREM of any one of embodiments 1-8, wherein [L3] is a linker comprising a nucleotide having a non-naturally occurring modification.

17. The TREM of any one of embodiments 1-8, wherein [ACH Domain (ACHD)] comprises a nucleotide having a non-naturally occurring modification.

5 18. The TREM of any one of embodiments 1-8, wherein [VL Domain (VLD)] comprises a nucleotide having a non-naturally occurring modification.

19. The TREM of any one of embodiments 1-8, wherein [TH Domain (THD)] comprises a nucleotide having a non-naturally occurring modification.

10

20. The TREM of any one of embodiments 1-8, wherein [L4] is a linker comprises a nucleotide having a non-naturally occurring modification.

21. The TREM of any one of embodiments 1-8, wherein: [ASt Domain2 (AStD2)] comprises a
15 nucleotide having a non-naturally occurring modification.

22. A TREM core fragment comprising a sequence of Formula B:

$[L1]_y-[ASt\ Domain1]_x-[L2]_y-[DH\ Domain]_y-[L3]_y-[ACH\ Domain]_x-[VL\ Domain]_y-[TH\ Domain]_y-[L4]_y-[ASt\ Domain2]_x,$

20

wherein:

$x=1$ and $y=0$ or 1 ;

one of [ASt Domain1], [ACH Domain], and [ASt Domain2] comprises a nucleotide having a non-naturally occurring modification; and

the TREM retains the ability to: support protein synthesis; be able to be charged by a
25 synthetase, be bound by an elongation factor, introduce an amino acid into a peptide chain, support elongation, or support initiation.

23. The TREM core fragment of embodiment 22, wherein AStD1 and AStD2 comprise an ASt Domain (AStD).

30

24. The TREM core fragment of embodiment 22, wherein the [ASt Domain 1], and/or [ASt Domain 2] comprising the non-naturally occurring modification retains the ability to initiate or elongate a polypeptide chain.
- 5 25. The TREM core fragment of embodiment 22, wherein the [ACH Domain] comprising the non-naturally occurring modification retains the ability to mediate pairing with a codon.
26. The TREM core fragment of embodiment 22, wherein y=1 for any one, two, three, four, five, six, all or a combination of [L1], [L2], [DH Domain], [L3], [VL Domain], [TH Domain], [L4].
- 10 27. The TREM core fragment of embodiment 22, wherein y=0 for any one, two, three, four, five, six, all or a combination of [L1], [L2], [DH Domain], [L3], [VL Domain], [TH Domain], [L4].
28. The TREM core fragment of embodiment 22, wherein y=1 for linker [L1], and L1 comprises
- 15 a nucleotide having a non-naturally occurring modification.
29. The TREM core fragment of embodiment 22, wherein y=1 for linker [L2], and L2 comprises a nucleotide having a non-naturally occurring modification.
- 20 30. The TREM core fragment of embodiment 22, wherein y=1 for [DH Domain (DHD)], and DHD comprises a nucleotide having a non-naturally occurring modification.
31. The TREM core fragment of embodiment 30, wherein the DHD comprising the non-naturally occurring modification retains the ability to mediate recognition of aminoacyl-tRNA synthetase.
- 25 32. The TREM core fragment of embodiment 22, wherein y=1 for linker [L3], and L3 comprises a nucleotide having a non-naturally occurring modification.
33. The TREM core fragment of embodiment 22, wherein y=1 for [VL Domain (VLD)], and
- 30 VLD comprises a nucleotide having a non-naturally occurring modification.

34. The TREM core fragment of embodiment 22, wherein $y=1$ for [TH Domain (THD)], and THD comprises a nucleotide having a non-naturally occurring modification.
35. The TREM core fragment of embodiment 34, wherein the THD comprising the non-naturally occurring modification retains the ability to mediate recognition of the ribosome.
36. The TREM core fragment of embodiment 22, wherein $y=1$ for linker [L4], and L4 comprises a nucleotide having a non-naturally occurring modification.
37. A TREM fragment comprising a portion of a TREM, wherein the TREM comprises a sequence of Formula A:
[L1]-[ASt Domain1]-[L2]-[DH Domain]-[L3]-[ACH Domain] -[VL Domain]-[TH Domain]-[L4]-[ASt Domain2], and wherein:
the TREM fragment comprises:
a non-naturally occurring modification; and
one, two, three or all or any combination of the following:
(a) a TREM half (*e.g.*, from a cleavage in the ACH Domain, *e.g.*, in the anticodon sequence, *e.g.*, a 5' half or a 3' half);
(b) a 5' fragment (*e.g.*, a fragment comprising the 5' end, *e.g.*, from a cleavage in a DH Domain or the ACH Domain);
(c) a 3' fragment (*e.g.*, a fragment comprising the 3' end, *e.g.*, from a cleavage in the TH Domain); or
(d) an internal fragment (*e.g.*, from a cleavage in any one of the ACH Domain, DH Domain or TH Domain).
38. The TREM of embodiment 37, wherein the TREM fragment comprise (a) a TREM half which comprises a nucleotide having a non-naturally occurring modification.
39. The TREM of embodiment 37, wherein the TREM fragment comprise (b) a 5' fragment which comprises a nucleotide having a non-naturally occurring modification.

40. The TREM of embodiment 37, wherein the TREM fragment comprise (c) a 3' fragment which comprises a nucleotide having a non-naturally occurring modification.
41. The TREM of embodiment 37, wherein the TREM fragment comprise (d) an internal
5 fragment which comprises a nucleotide having a non-naturally occurring modification.
42. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM Domain comprises a plurality of nucleotides each having a non-naturally occurring modification.
- 10 43. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of AStD1 have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6 or 7.
- 15 44. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein no more than X of the nucleotides of AStD1 have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6 or 7.
- 20 45. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of AStD2 have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6 or 7.
- 25 46. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein no more than X of the nucleotides of AStD2 have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6 or 7.
- 30 47. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of ACHD have a

non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.

5 48. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of ACHD have a non-naturally occurring modification, wherein X is equal to or greater than 11, 12, 13, 14, 15, 16, or 17.

10 49. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein no more than X of the nucleotides of ACHD have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.

15 50. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein no more than X of the nucleotides of ACHD have a non-naturally occurring modification, wherein X is equal to or greater than 11, 12, 13, 14, 15, or 16.

20 51. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of THD have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.

25 52. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of THD have a non-naturally occurring modification, wherein X is equal to or greater than 11, 12, 13, 14, 15, 16, or 17.

30 53. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein no more than X of the nucleotides of THD have

a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.

54. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or
5 the TREM fragment of embodiment 37, wherein no more than X of the nucleotides of THD have a non-naturally occurring modification, wherein X is equal to or greater than 11, 12, 13, 14, 15, or 16.

55. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or
10 the TREM fragment of embodiment 37, wherein at least X of the nucleotides of DHD have a non-naturally occurring modification, wherein X is equal to or greater than 2, 3, 4, 5, 6, 7, 8, 9 or 10.

56. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or
15 the TREM fragment of embodiment 37, wherein at least X of the nucleotides of DHD have a non-naturally occurring modification, wherein X is equal to or greater than 11, 12, 13, 14, 15, 16, 17, 18 or 19.

57. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or
20 the TREM fragment of embodiment 37, wherein no more than X of the nucleotides of DHD have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.

58. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or
25 the TREM fragment of embodiment 37, wherein no more than X of the nucleotides of DHD have a non-naturally occurring modification, wherein X is equal to or greater than 11, 12, 13, 14, 15, 16, 17, or 18.

59. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or
30 the TREM fragment of embodiment 37, wherein at least X of the nucleotides of the VLD have a

non-naturally occurring modification, wherein X is equal to or greater than 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 50, 100, 150, 200 or 271.

5 60. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein all of the nucleotides of the AStD1, AStD2, ACHD, DHD, and/or THD have a non-naturally occurring modification.

61. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of AStD1 and/or
10 AStD2 do not have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6 or 7.

62. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of ACHD do not
15 have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17.

63. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of THD do not
20 have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17.

64. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of DHD do not
25 have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18 or 19.

65. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of VLD do not
30 have a non-naturally occurring modification, wherein X is equal to or greater than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 50, 100, 150, 200 or 271.

66. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM Linker L2 comprises two nucleotides each having a non-naturally occurring modification.

5

67. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X of the nucleotides of the TREM Linker do not have a non-naturally occurring modification, wherein X is equal to 1 or 2.

10 68. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein:

each of a plurality of TREM Domains and Linkers comprises a nucleotide having a non-naturally occurring modification.

15 69. The TREM, TREM core fragment or TREM fragment of embodiment 68, wherein one of the TREM Domains and Linkers of the plurality comprises a plurality of nucleotides each having a non-naturally occurring modification.

70. The TREM, TREM core fragment or TREM fragment of any of the preceding embodiments, wherein the non-naturally occurring modification is a modification in a base or a backbone of a nucleotide, e.g., a modification chosen from any one of Tables 5-9.

71. The TREM, TREM core fragment or TREM fragment of any of the preceding embodiments, wherein the non-naturally occurring modification is a base modification chosen from a modification listed in Table 10.

72. The TREM, TREM core fragment or TREM fragment of any of the preceding embodiments, wherein the non-naturally occurring modification is a base modification chosen from a modification listed in Table 11.

30

73. The TREM, TREM core fragment or TREM fragment of any of the preceding embodiments, wherein the non-naturally occurring modification is a base modification chosen from a modification listed in Table 12.

5 74. The TREM, TREM core fragment or TREM fragment of any of the preceding embodiments, wherein the non-naturally occurring modification is a backbone base modification chosen from a modification listed in Table 13.

75. The TREM, TREM core fragment or TREM fragment of any of the preceding embodiments,
10 wherein the non-naturally occurring modification is a backbone modification chosen from a modification listed in Table 14.

76. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, comprising a nucleotide of a first type comprising a non-
15 naturally occurring modification.

77. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, comprising a nucleotide of a first type and a nucleotide of a second type comprising a non-naturally occurring modification.

20

78. The TREM, TREM core fragment or TREM fragment of embodiment 77, wherein the non-naturally occurring modification on the nucleotide of the first type and the non-naturally occurring modification on the nucleotide of the second type are the same non-naturally occurring modification.

25

79. The TREM, TREM core fragment or TREM fragment of embodiment 77, wherein the non-naturally occurring modification on the nucleotide of the first type and the non-naturally occurring modification on the nucleotide of the second type are different non-naturally occurring modifications.

30

80. The TREM, TREM core fragment or TREM fragment of embodiments 76 or 77, wherein the nucleotide of the first type is chosen from: A, T, C, G or U.
81. The TREM, TREM core fragment or TREM fragment of embodiments 76 or 77, wherein the nucleotide of the second type is chosen from: A, T, C, G or U.
82. The TREM, TREM core fragment or TREM fragment of embodiments 76 or 77, wherein the nucleotide of the first type is an A.
83. The TREM, TREM core fragment or TREM fragment of embodiments 76 or 77, wherein the nucleotide of the first type is a G.
84. The TREM, TREM core fragment or TREM fragment of embodiments 76 or 77, wherein the nucleotide of the first type is a C.
85. The TREM core fragment or TREM fragment of embodiments 76 or 77, wherein the nucleotide of the first type is a T.
86. The TREM, TREM core fragment or TREM fragment of embodiments 76 or 77, wherein the nucleotide of the first type is a U.
87. The TREM, TREM core fragment or TREM fragment of embodiment 77, wherein when the nucleotide of the first type is an A, the nucleotide of the second type is chosen from: T, C, G or U.
88. The TREM, TREM core fragment or TREM fragment of embodiment 77, wherein when the nucleotide of the first type is a G, the nucleotide of the second type is chosen from: T, C, A or U.
89. The TREM, TREM core fragment or TREM fragment of embodiment 77, wherein when the nucleotide of the first type is a C, the nucleotide of the second type is chosen from: T, A, G or U.

90. The TREM, TREM core fragment or TREM fragment of embodiment 77, wherein when the nucleotide of the first type is a T, the nucleotide of the second type is chosen from: A, C, G or U.

5 91. The TREM, TREM core fragment or TREM fragment of embodiment 77, wherein when the nucleotide of the first type is a U, the nucleotide of the second type is chosen from: T, C, G or A.

92. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the non-naturally modification is in a purine (A or G).

10

93. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the non-naturally modification is not in a purine (A or G).

15 94. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the non-naturally modification is in a pyrimidine (U, T or C).

20 95. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the non-naturally modification is not in a pyrimidine (U, T or C).

96. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the DHD has a first sequence, a second
25 sequence and a third sequence, optionally wherein the first sequence and the third sequence form a stem and the second sequence forms a loop, e.g., under physiological conditions.

97. The TREM, TREM core fragment or TREM fragment of embodiment 96, wherein the DHD comprises a non-naturally occurring modification in the first sequence or the third sequence, e.g.,
30 in the stem.

98. The TREM, TREM core fragment or TREM fragment of embodiment 96, wherein the DHD comprises a non-naturally occurring modification in the second sequence, e.g., in the loop.

5 100. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the ACHD has a first sequence, a second sequence and a third sequence, optionally wherein the first sequence and the third sequence form a stem and the second sequence forms a loop, e.g., under physiological conditions.

10 101. The TREM, TREM core fragment or TREM fragment of embodiment 100, wherein the ACHD comprises a non-naturally occurring modification in the first sequence or the third sequence, e.g., in the stem.

15 102. The TREM, TREM core fragment or TREM fragment of embodiment 100, wherein the ACHD comprises a non-naturally occurring modification in the second sequence, e.g., in the loop.

20 103. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the THD has a first sequence, a second sequence and a third sequence, optionally wherein the first sequence and the third sequence form a stem and the second sequence forms a loop, e.g., under physiological conditions.

25 104. The TREM, TREM core fragment or TREM fragment of embodiment 103, wherein the THD comprises a non-naturally occurring modification in the first sequence or the third sequence, e.g., in the stem.

105. The TREM, TREM core fragment or TREM fragment of embodiment 103, wherein the THD comprises a non-naturally occurring modification in the second sequence, e.g., in the loop.

30 106. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the VLD comprises a variable region having 1-271 nucleotides.

107. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM comprises at least X contiguous nucleotides without a non-naturally occurring modification, wherein X is greater than 10.

5

108. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least 3, but less than all of the nucleotides of a type (e.g., A, T, C, G or U) comprise the same non-naturally occurring modification.

109. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein at least X nucleotides of a type (e.g., A, T, C, G or U) do not comprise a non-naturally occurring modification, wherein X=1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 or 50.

15

110. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein no more than 5, 10, or 15 of a type (e.g., A, T, C, G or U) comprise a non-naturally occurring modification.

111. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein no more than 5, 10, or 15 of a type (e.g., A, T, C, G or U) do not comprise a non-naturally occurring modification.

112. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, which specifies X, wherein X is an amino acid selected from alanine, arginine, asparagine, aspartate, cysteine, glutamine, glutamate, glycine, histidine, isoleucine, methionine, leucine, lysine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, or valine.

113. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, which recognizes a codon provided in Table 7 or Table 8.

5 114. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM is a cognate TREM.

115. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM is a non-cognate TREM.

10

116. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM, TREM core fragment, or TREM fragment is encoded by a sequence provided in Table 9, e.g., any one of SEQ ID NOs 1-451.

15 117. The TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM, TREM core fragment, or TREM fragment is encoded by a consensus sequence chosen from any one of SEQ ID NOs: 562-621.

20 118. A pharmaceutical composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37.

119. The pharmaceutical composition of embodiment 118, comprising a pharmaceutically acceptable component, *e.g.*, an excipient.

25 120. A method of making a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, comprising linking a first nucleotide to a second nucleotide to form the TREM.

30 121. The method of embodiment 120, wherein the TREM, TREM core fragment or TREM fragment is synthetic.

122. The method of embodiment 120 or 121, wherein the synthesis is performed in vitro.

123. The method of embodiment 120, wherein the TREM, TREM core fragment or TREM fragment is made by cell-free solid phase synthesis.

5

124. A cell comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37.

125. A cell comprising a TREM, TREM core fragment or TREM fragment made according to the method of embodiment 120.

10

126. A method of modulating a tRNA pool in a cell comprising an endogenous open reading frame (ORF), which ORF comprises a codon having a first sequence, comprising:

optionally, acquiring knowledge of the abundance of one or both of (i) and (ii), e.g.,
15 acquiring knowledge of the relative amounts of: (i) and (ii) in the cell, wherein (i) is a tRNA moiety having an anticodon that pairs with the codon of the ORF having a first sequence (the first tRNA moiety) and (ii) is an isoacceptor tRNA moiety having an anticodon that pairs with a codon other than the codon having the first sequence (the second tRNA moiety) in the cell;
contacting the cell with a TREM composition comprising a TREM of any one of
20 embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with: (a) the codon having the first sequence; or (b) the codon other than the codon having the first sequence, in an amount and/or for a time sufficient to modulate the relative amounts of the first tRNA moiety and the second tRNA moiety in the cell,
25 thereby modulating the tRNA pool in the cell.

127. A method of modulating a tRNA pool in a subject having an endogenous open reading frame (ORF), which ORF comprises a codon having a first sequence, comprising:

optionally, acquiring knowledge of the abundance of one or both of (i) and (ii), e.g.,
30 acquiring knowledge of the relative amounts of: (i) and (ii) in the subject, wherein (i) is a tRNA moiety having an anticodon that pairs with the codon of the ORF having a first sequence (the

first tRNA moiety) and (ii) is an isoacceptor tRNA moiety having an anticodon that pairs with a codon other than the codon having the first sequence (the second tRNA moiety) in the subject;

contacting the subject with a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with: (a) the codon having the first sequence; or (b) the codon other than the codon having the first sequence, in an amount and/or for a time sufficient to modulate the relative amounts of the first tRNA moiety and the second tRNA moiety in the subject, thereby modulating the tRNA pool in the subject.

10

128. The method of embodiment 126 or 127, wherein the TREM composition comprises a TREM, TREM fragment or TREM core fragment comprising an anticodon that pairs with (a).

15

129. The method of embodiment 126 or 127, wherein the TREM composition comprises a TREM, TREM fragment or TREM core fragment comprising an anticodon that pairs with (b).

130. The method of any one of embodiments 126-129, comprising acquiring knowledge of (i).

20

131. The method of any one of embodiments 126-129, comprising acquiring knowledge of (ii).

132. The method of any one of embodiments 126-129, comprising acquiring knowledge of (i) and (ii).

25

133. The method of any one of embodiments 126-130 or 132, wherein acquiring knowledge of (i) comprises acquiring a value for the abundance, e.g., relative amounts, of (i).

134. The method of any one of embodiments 126-129 or 131-312, wherein acquiring knowledge of (ii) comprises acquiring a value for the abundance, e.g., relative amounts, of (ii).

135. The method of embodiment 133 or 134, wherein responsive to said value, the cell or subject is contacted with the TREM composition comprising a TREM, TREM fragment or TREM core fragment having an anticodon that pairs with (a) or (b).

5 136. A method of evaluating a tRNA pool in a cell or subject, comprising acquiring, e.g., directly or indirectly acquiring, knowledge of the abundance of one or both of (i) and (ii), e.g., acquiring knowledge of the relative amounts of (i) and (ii) in the cell wherein (i) is a tRNA moiety having an anticodon that pairs with the codon of the ORF having a first sequence (the first tRNA moiety) and (ii) is an isoacceptor tRNA moiety having an anticodon that pairs with a codon other
10 than the codon having the first sequence (the second tRNA moiety) in the cell, thereby evaluating the tRNA pool in the cell or subject.

137. A method of modulating a production parameter of an RNA corresponding to, or polypeptide encoded by, a nucleic acid sequence comprising an endogenous open reading frame (ORF) in a cell, which ORF comprises a codon having a first sequence, comprising:
15

contacting the cell with a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37 in an amount and/or for a time sufficient to modulate the production parameter of the mRNA or polypeptide,

20 wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,

thereby modulating the production parameter in the cell.

138. A method of modulating a production parameter of an RNA corresponding to, or
25 polypeptide encoded by, a nucleic acid sequence comprising an endogenous open reading frame (ORF) in a subject, which ORF comprises a codon having a first sequence, comprising:

contacting the subject with a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37 in an amount and/or for a time sufficient to modulate the production parameter
30 of the mRNA or polypeptide,

wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,

thereby modulating the production parameter in the subject.

5 139. The method of embodiment 137 or 138, wherein the production parameter comprises a signaling parameter, e.g., as described herein.

140. The method of embodiment 137 or 138, wherein the production parameter comprises an expression parameter, e.g., as described herein.

10

141. A method of modulating expression of a protein in a cell, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a codon having a first sequence, comprising:

15 contacting the cell with a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37 in an amount and/or for a time sufficient to modulate expression of the encoded protein,

wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,

20 thereby modulating expression of the protein in the cell.

142. A method of modulating expression of a protein in a subject, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a codon having a first sequence, comprising:

25 contacting the subject with a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, in an amount and/or for a time sufficient to modulate expression of the encoded protein,

30 wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,

thereby modulating expression of the protein in the subject.

143. A method of treating a subject having an endogenous open reading frame (ORF) which comprises a codon having a first sequence, comprising:

- 5 providing a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM comprises a tRNA moiety having: an anticodon that pairs with the codon of the ORF having the first sequence;
- 10 contacting the subject with the composition comprising a TREM, TREM core fragment or TREM fragment in an amount and/or for a time sufficient to treat the subject, thereby treating the subject.

144. A method of treating a subject having an endogenous open reading frame (ORF) comprising a codon having a first sequence, comprising:

- 15 (i) acquiring, e.g., directly or indirectly acquiring, a value for the status of the codon having the first sequence in the subject, wherein said value comprises a measure of the presence or absence of the codon having the first sequence in a sample from the subject; and identifying the subject as having the codon having the first sequence; and
- 20 (ii) responsive to said value, administering a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM, TREM core fragment or TREM fragment comprises a tRNA moiety having an anticodon that pairs with the codon having the first sequence, to the subject,
- thereby treating the subject.

- 25 145. A method of evaluating a subject having an endogenous open reading frame (ORF) comprising a codon having a first sequence, comprising:

- acquiring, e.g., directly or indirectly acquiring, a value for the status of the codon having the first sequence in the subject, wherein said value comprises a measure of the presence or absence of the codon having the first sequence in a sample from the subject; and
- 30 identifying the subject as having a codon having the first sequence, thereby evaluating the subject.

146. The method of claim 145, wherein responsive to said value the method further comprises administering a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the
5 TREM, TREM core fragment or TREM fragment comprises a tRNA moiety having an anticodon that pairs with the codon having the first sequence, to the subject.

147. A method of modulating a production parameter of an RNA corresponding to, or polypeptide encoded by, a nucleic acid sequence comprising an endogenous open reading frame
10 (ORF) in a cell, which ORF comprises a premature termination codon (PTC),
contacting the cell with a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37 in an amount and/or for a time sufficient to modulate the production parameter of the mRNA or polypeptide,
15 wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,
thereby modulating the production parameter in the cell.

148. A method of modulating a production parameter of an RNA corresponding to, or
20 polypeptide encoded by, a nucleic acid sequence comprising an endogenous open reading frame (ORF) in a subject, which ORF comprises a premature termination codon (PTC),
contacting the subject with a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37 in an amount and/or for a time sufficient to modulate the production parameter
25 of the mRNA or polypeptide,
wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,
thereby modulating the production parameter in the subject.

30 149. The method of embodiment 147 or 148, wherein the production parameter comprises a signaling parameter and/or an expression parameter, e.g., as described herein.

150. A method of treating a subject having an endogenous open reading frame (ORF) which comprises a premature termination codon (PTC), comprising:

- 5 providing a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37, wherein the TREM comprises a tRNA moiety having an anticodon that pairs with the PTC in the ORF;
- contacting the subject with the composition comprising a TREM, TREM core fragment or TREM fragment in an amount and/or for a time sufficient to treat the subject,
- thereby treating the subject.

10

151. The method of embodiment 150, wherein the PTC comprises UAA, UGA or UAG.

152. A method of modulating expression of a protein in a cell, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a

15 premature termination codon (PTC), comprising:

- contacting the cell with a TREM composition comprising a TREM of any one of embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of embodiment 37 in an amount and/or for a time sufficient to modulate expression of the encoded protein,
- 20 wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the PTC,
- thereby modulating expression of the protein in the cell.

25

153. The method of embodiment 152, wherein the PTC comprises UAA, UGA or UAG.

154. A method of modulating expression of a protein in a subject, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a premature termination codon (PTC), comprising:

- contacting the subject with a TREM composition comprising a TREM of any one of
- 30 embodiments 1-8, the TREM core fragment of embodiment 22, or the TREM fragment of

embodiment 37, in an amount and/or for a time sufficient to modulate expression of the encoded protein,

wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the PTC,

5 thereby modulating expression of the protein in the subject.

155. The method of embodiment 154, wherein the PTC comprises UAA, UGA or UAG.

156. The method of any one of embodiments 126-146, wherein the codon having the first
10 sequence comprises a mutation (e.g., a point mutation, e.g., a nonsense mutation), resulting in a premature termination codon (PTC) chosen from UAA, UGA or UAG.

157. The method of any one of embodiments 126-156, wherein the codon having the first
15 sequence or the PTC comprises a UAA mutation.

158. The method of any one of embodiments 126-156, wherein the codon having the first
sequence or the PTC comprises a UGA mutation.

159. The method of any one of embodiments 126-156, wherein the codon having the first
20 sequence or the PTC comprises a UAG mutation.

160. The method of any one of embodiments 126-159, wherein the TREM comprises an
anticodon that pairs with a stop codon, e.g., a stop codon chosen from UAA, UGA or UAG.

25 161. The method of any one of embodiments 126-160, wherein the codon having the first
sequence or the PTC comprises a UAA mutation and the TREM, TREM core fragment or TREM
fragment mediates incorporation of an amino acid which preserves, e.g., maintains, a secondary
and/or tertiary structure of a polypeptide encoded by the ORF into which the amino acid is
incorporated.

30

162. The method of any one of embodiments 126-160, wherein the codon having the first sequence or the PTC comprises a UAG mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid which preserves, e.g., maintains, a secondary and/or tertiary structure of a polypeptide encoded by the ORF into which the amino acid is
5 incorporated.

163. The method of any one of embodiments 126-160, wherein the codon having the first sequence or the PTC comprises a UGA mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid which preserves, e.g., maintains, a secondary
10 and/or tertiary structure of a polypeptide encoded by the ORF into which the amino acid is incorporated.

164. The method of any one of embodiments 126-160, wherein the codon having the first sequence or the PTC comprises a UAA mutation and the TREM, TREM core fragment or TREM
15 fragment mediates incorporation of an amino acid which maintains a property, e.g., function, of a polypeptide encoded by the ORF into which the amino acid is incorporated.

165. The method of any one of embodiments 126-160, wherein the codon having the first sequence or the PTC comprises a UAG mutation and the TREM, TREM core fragment or TREM
20 fragment mediates incorporation of an amino acid which maintains a property, e.g., function, of a polypeptide encoded by the ORF into which the amino acid is incorporated.

166. The method of any one of embodiments 126-160, wherein the codon having the first sequence or the PTC comprises a UGA mutation and the TREM, TREM core fragment or TREM
25 fragment mediates incorporation of an amino acid which maintains a property, e.g., function, of a polypeptide encoded by the ORF into which the amino acid is incorporated.

167. The method of any one of embodiments 161-166, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of any one of the twenty amino acids listed in Table 2
30 or Table 8.

168. The method of any one of embodiments 161-167, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid corresponding to a non-mutated codon, e.g., a wildtype codon sequence of the codon having the first sequence or the PTC.

5 169. The method of any one of embodiments 161-168, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of a pre-mutation, e.g., wildtype amino acid.

170. The method of embodiment 169, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid having a similar property as the pre-mutation, e.g., wildtype amino acid, e.g., an amino acid that belongs to the same group as the pre-mutation amino acid, e.g., as provided in Table 2.

10

171. The method of embodiment 169 or 170, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid that belongs to the same group as the pre-mutation amino acid, e.g., as provided in Table 2.

15

172. The method of embodiment 171, wherein when the pre-mutation, e.g., wildtype, amino acid is a nonpolar amino acid having an aliphatic R group, the TREM mediates incorporation of any one of the following amino acids: leucine, methionine, isoleucine, glycine, alanine or valine.

20

173. The method of embodiment 171, wherein when the pre-mutation, e.g., wildtype, amino acid is a polar amino acid having an uncharged R group, the TREM mediates incorporation of any one of the following amino acids: serine, threonine, cysteine, proline, asparagine, or glutamine.

25 174. The method of embodiment 171, wherein when the pre-mutation, e.g., wildtype, amino acid has a positively charged R group, the TREM mediates incorporation of any one of the following amino acids: lysine, arginine or histidine.

175. The method of embodiment 171, wherein when the pre-mutation, e.g., wildtype, amino acid has a negatively charged R group, the TREM mediates incorporation of any one of the following amino acids: aspartate or glutamate.

30

176. The method of embodiment 171, wherein when the pre-mutation, e.g., wildtype, amino acid is a nonpolar amino acid having an aromatic R group, the TREM mediates incorporation of any one of the following amino acids: phenylalanine, tyrosine or tryptophan.

5

177. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAA mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid which does not alter, e.g., maintains, a production parameter, e.g., an expression parameter and/or a signaling parameter, of an RNA corresponding to the ORF or a polypeptide encoded by the ORF.

10

178. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UGA mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid which does not alter, e.g., maintains, a production parameter, e.g., an expression parameter and/or a signaling parameter, of an RNA corresponding to the ORF or a polypeptide encoded by the ORF.

15

179. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAG mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid which does not alter, e.g., maintains, a production parameter, e.g., an expression parameter and/or a signaling parameter, of an RNA corresponding to the ORF or a polypeptide encoded by the ORF.

20

180. The method of any one of embodiments 177-179, wherein the production parameter is compared to an RNA corresponding to, or a polypeptide encoded by, an otherwise similar ORF having a pre-mutation, e.g., wildtype, amino acid incorporated at the position corresponding to the first sequence codon or PTC.

25

181. The method of any one of embodiments 177-180, wherein the production parameter comprises an expression parameter.

30

182. The method of embodiment 181, wherein the expression parameter comprises:

- (a) protein translation;
- (b) expression level (*e.g.*, of polypeptide or protein, or mRNA);
- (c) post-translational modification of polypeptide or protein;
- 5 (d) folding (*e.g.*, of polypeptide or protein, or mRNA),
- (e) structure (*e.g.*, of polypeptide or protein, or mRNA),
- (f) transduction (*e.g.*, of polypeptide or protein),
- (g) compartmentalization (*e.g.*, of polypeptide or protein, or mRNA),
- 10 (h) incorporation (*e.g.*, of polypeptide or protein, or mRNA) into a supermolecular structure, *e.g.*, incorporation into a membrane, proteasome, or ribosome,
- (i) incorporation into a multimeric polypeptide, *e.g.*, a homo or heterodimer, and/or
- (j) stability.

183. The method of any one of embodiments 177-180, wherein the production parameter
15 comprises a signaling parameter.

184. The method of embodiment 183, wherein the signaling parameter comprises:

- (1) modulation of a signaling pathway, *e.g.*, a cellular signaling pathway which is downstream or upstream of the protein encoded by the endogenous ORF having a first sequence
20 or PTC;
- (2) cell fate modulation;
- (3) ribosome occupancy modulation;
- (4) protein translation modulation;
- (5) mRNA stability modulation;
- 25 (6) protein folding and structure modulation;
- (7) protein transduction or compartmentalization modulation; and/or
- (8) protein stability modulation.

185. The method of any one of embodiments 177-184, wherein the production parameter (*e.g.*,
30 an expression parameter and/or a signaling parameter) may be modulated (*e.g.*, increased), *e.g.*,

by at least 5% (*e.g.*, at least 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 150%, 200% or more), *e.g.*, compared to a reference sequence.

186. The method of any one of embodiments 177-185, wherein the TREM, TREM core fragment
5 or TREM fragment mediates incorporation of any one of the twenty amino acids listed in Table 2 or Table 8.

187. The method of any one of embodiments 177-186, wherein the TREM, TREM core fragment
or TREM fragment mediates incorporation of an amino acid corresponding to a non-mutated
10 codon, *e.g.*, a wildtype codon sequence of the codon having the first sequence or the PTC.

188. The method of any one of embodiments 177-187, wherein the TREM, TREM core fragment
or TREM fragment mediates incorporation of a pre-mutation, *e.g.*, wildtype amino acid.

15 189. The method of embodiment 188, wherein the TREM, TREM core fragment or TREM
fragment mediates incorporation of an amino acid having a similar property as the pre-mutation,
e.g., wildtype amino acid, *e.g.*, an amino acid that belongs to the same group as the pre-mutation
amino acid, *e.g.*, as provided in Table 2.

20 190. The method of embodiment 188 or 189, wherein the TREM, TREM core fragment or
TREM fragment mediates incorporation of an amino acid that belongs to the same group as the
pre-mutation amino acid, *e.g.*, as provided in Table 2.

191. The method of embodiment 190, wherein when the pre-mutation, *e.g.*, wildtype, amino acid
25 is a nonpolar amino acid having an aliphatic R group, the TREM mediates incorporation of any
one of the following amino acids: leucine, methionine, isoleucine, glycine, alanine or valine.

192. The method of embodiment 190, wherein when the pre-mutation, *e.g.*, wildtype, amino acid
is a polar amino acid having an uncharged R group, the TREM mediates incorporation of any
30 one of the following amino acids: serine, threonine, cysteine, proline, asparagine, or glutamine.

193. The method of embodiment 190, wherein when the pre-mutation, e.g., wildtype, amino acid has a positively charged R group, the TREM mediates incorporation of any one of the following amino acids: lysine, arginine or histidine.
- 5 194. The method of embodiment 190, wherein when the pre-mutation, e.g., wildtype, amino acid has a negatively charged R group, the TREM mediates incorporation of any one of the following amino acids: aspartate or glutamate.
195. The method of embodiment 190, wherein when the pre-mutation, e.g., wildtype, amino acid
10 is a nonpolar amino acid having an aromatic R group, the TREM mediates incorporation of any one of the following amino acids: phenylalanine, tyrosine or tryptophan.
196. The method of any one of embodiments 126-160, wherein the codon having the first
sequence or the PTC comprises a UAA mutation and the TREM, TREM core fragment or TREM
15 fragment mediates incorporation of any one of the 20 amino acids listed in Table 8 at the UAA stop codon.
197. The method of embodiment 196, wherein the TREM, TREM core fragment or TREM
fragment mediates incorporation of the amino acid corresponding to a non-mutated codon, e.g., a
20 wildtype codon sequence of the codon having the first sequence or the PTC.
198. The method of embodiment 196 or 197, wherein the TREM, TREM core fragment or
TREM fragment mediates incorporation of a pre-mutation, e.g., wildtype amino acid.
- 25 199. The method of embodiment 198, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid having similar characteristics as the pre-mutation, e.g., wildtype amino acid, e.g., an amino acid that belongs to the same group as the pre-mutation amino acid, e.g., as provided in Table 2.
- 30 200. The method of embodiment 198 or 199, wherein when the pre-mutation, e.g., wildtype, amino acid is a nonpolar amino acid having an aliphatic R group, the TREM mediates

incorporation of any one of the following amino acids: leucine, methionine, isoleucine, glycine, alanine or valine.

201. The method of embodiment 198 or 199, wherein when the pre-mutation, e.g., wildtype,
5 amino acid is a polar amino acid having an uncharged R group, the TREM mediates
incorporation of any one of the following amino acids: serine, threonine, cysteine, proline,
asparagine, or glutamine.

202. The method of embodiment 198 or 199, wherein when the pre-mutation, e.g., wildtype,
10 amino acid has a positively charged R group, the TREM mediates incorporation of any one of
the following amino acids: lysine, arginine or histidine.

203. The method of embodiment 198 or 199, wherein when the pre-mutation, e.g., wildtype,
amino acid has a negatively charged R group, the TREM mediates incorporation of any one of
15 the following amino acids: aspartate or glutamate.

204. The method of embodiment 198 or 199, wherein when the pre-mutation, e.g., wildtype,
amino acid is a nonpolar amino acid having an aromatic R group, the TREM mediates
incorporation of any one of the following amino acids: phenylalanine, tyrosine or tryptophan.
20

205. The method of any one of embodiments 126-160, wherein the codon having the first
sequence or the PTC comprises a UGA mutation and the TREM, TREM core fragment or TREM
fragment mediates incorporation of any one of the 20 amino acids listed in Table 8 at the UGA
stop codon.
25

206. The method of embodiment 205, wherein the TREM, TREM core fragment or TREM
fragment mediates incorporation of the amino acid corresponding to a non-mutated, e.g., a
wildtype codon sequence of the codon having the first sequence or the PTC.

30 207. The method of embodiment 206, wherein the TREM, TREM core fragment or TREM
fragment mediates incorporation of a pre-mutation, e.g., wildtype, amino acid.

208. The method of embodiment 206 or 207, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid having similar characteristics as the pre-mutation, e.g., wildtype, amino acid, e.g., an amino acid that belongs to the same group as the pre-mutation amino acid as provided in Table 2.

209. The method of embodiment 206 or 207, wherein when the pre-mutation, e.g., wildtype, amino acid is a nonpolar amino acid having an aliphatic R group, the TREM mediates incorporation of any one of the following amino acids: leucine, methionine, isoleucine, glycine, alanine or valine.

210. The method of embodiment 206 or 207, wherein when the pre-mutation, e.g., wildtype, amino acid is a polar amino acid having an uncharged R group, the TREM mediates incorporation of any one of the following amino acids: serine, threonine, cysteine, proline, asparagine, or glutamine.

211. The method of embodiment 206 or 207, wherein when the pre-mutation, e.g., wildtype, amino acid has a positively charged R group, the TREM mediates incorporation of any one of the following amino acids: lysine, arginine or histidine.

212. The method of embodiment 206 or 207, wherein when the pre-mutation, e.g., wildtype, amino acid has a negatively charged R group, the TREM mediates incorporation of any one of the following amino acids: aspartate or glutamate.

213. The method of embodiment 206 or 207, wherein when the pre-mutation, e.g., wildtype, amino acid is a nonpolar amino acid having an aromatic R group, the TREM mediates incorporation of any one of the following amino acids: phenylalanine, tyrosine or tryptophan.

214. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAG mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of any one of the 20 amino acids listed in Table 8 at the UAG stop codon.

215. The method of embodiment 214, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of the amino acid corresponding to a non-mutated, e.g., a wildtype codon sequence of the codon having the first sequence or the PTC.

5

216. The method of embodiment 215, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of a pre-mutation, e.g., wildtype, amino acid.

217. The method of embodiment 216, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid having similar characteristics as the pre-mutation, e.g., wildtype, amino acid, e.g., an amino acid that belongs to the same group as the pre-mutation amino acid, e.g., as provided in Table 2.

218. The method of embodiment 216 or 217, wherein when the pre-mutation, e.g., wildtype, amino acid is a nonpolar amino acid having an aliphatic R group, the TREM mediates incorporation of any one of the following amino acids: leucine, methionine, isoleucine, glycine, alanine or valine.

219. The method of embodiment 216 or 217, wherein when the pre-mutation, e.g., wildtype, amino acid is a polar amino acid having an uncharged R group, the TREM mediates incorporation of any one of the following amino acids: serine, threonine, cysteine, proline, asparagine, or glutamine.

220. The method of embodiment 216 or 217, wherein when the pre-mutation, e.g., wildtype, amino acid has a positively charged R group, the TREM mediates incorporation of any one of the following amino acids: lysine, arginine or histidine.

221. The method of embodiment 216 or 217, wherein when the pre-mutation, e.g., wildtype, amino acid has a negatively charged R group, the TREM mediates incorporation of any one of the following amino acids: aspartate or glutamate.

222. The method of embodiment 216 or 217, wherein when the pre-mutation, e.g., wildtype, amino acid is a nonpolar amino acid having an aromatic R group, the TREM mediates incorporation of any one of the following amino acids: phenylalanine, tyrosine or tryptophan.

5 223. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UGA mutation, e.g., a UGG to UGA mutation.

224. The method of embodiment 223, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is UGA and the amino acid corresponding to
10 the non-mutated codon is a tryptophan.

225. The method of claim 224, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of tryptophan at the position of the UGA stop codon.

15 226. The method of claim 224, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as tryptophan, e.g., as provided in Table 2.

20 227. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAA mutation, e.g., a UAU to UAA mutation.

228. The method of embodiment 227, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is UAU and the amino acid corresponding to
25 the non-mutated codon is a tyrosine.

229. The method of claim 228, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAA stop codon and mediates incorporation of tyrosine at the position of the UAA stop codon.

30

230. The method of claim 228, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAA stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as tyrosine, e.g., as provided in Table 2.

5 231. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAG mutation, e.g., a UAC to UAG mutation.

232. The method of embodiment 231, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is UAC and the amino acid corresponding to
10 the non-mutated codon is a tyrosine.

233. The method of claim 232, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of tyrosine at the position of the UAG stop codon.

15 234. The method of claim 232, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as tyrosine, e.g., as provided in Table 2.

20 235. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UGA mutation, e.g., a UGU to UGA mutation.

236. The method of embodiment 235, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is UGU and the amino acid corresponding to
25 the non-mutated codon is a cysteine.

237. The method of claim 236, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of cysteine at the position of the UGA stop codon.

30

238. The method of claim 236, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as cysteine, e.g., as provided in Table 2.

5 239. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UGA mutation, e.g., a UGC to UGA mutation.

240. The method of embodiment 239, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is UGC and the amino acid corresponding to
10 the non-mutated codon is a cysteine.

241. The method of claim 240, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of cysteine at the position of the UGA stop codon.

15 242. The method of claim 240, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGG stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as cysteine, e.g., as provided in Table 2.

20 243. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAA mutation, e.g., a GAA to UAA mutation.

244. The method of embodiment 243, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is GAA and the amino acid corresponding to
25 the non-mutated codon is a glutamate.

245. The method of claim 244, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAA stop codon and mediates incorporation of glutamate at the position of the UAA stop codon.

30

246. The method of claim 244, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAA stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as glutamate, e.g., as provided in Table 2.

5 247. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAG mutation, e.g., a GAG to UAG mutation.

248. The method of embodiment 247, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is GAG and the amino acid corresponding to
10 the non-mutated codon is a glutamate.

249. The method of claim 248, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of glutamate at the position of the UAG stop codon.

15 250. The method of claim 248, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as glutamate, e.g., as provided in Table 2.

20 251. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAA mutation, e.g., a AAA to UAA mutation.

252. The method of embodiment 251, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is AAA and the amino acid corresponding to
25 the non-mutated codon is a lysine.

253. The method of claim 252, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAA stop codon and mediates incorporation of lysine at the position of the UAA stop codon.

30

254. The method of claim 252, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAA stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as lysine, e.g., as provided in Table 2.

5 255. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAG mutation, e.g., a AAG to UAG mutation.

256. The method of embodiment 255, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is AAG and the amino acid corresponding to
10 the non-mutated codon is a lysine.

257. The method of claim 256, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of lysine at the position of the UAG stop codon.

15 258. The method of claim 256, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as lysine, e.g., as provided in Table 2.

20 259. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAA mutation, e.g., a CAA to UAA mutation.

260. The method of embodiment 259, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is CAA and the amino acid corresponding to
25 the non-mutated codon is a glutamine.

261. The method of claim 260, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAA stop codon and mediates incorporation of glutamine at the position of the UAA stop codon.

30

262. The method of claim 260, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAA stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as glutamine, e.g., as provided in Table 2.

5 263. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAG mutation, e.g., a CAG to UAG mutation.

264. The method of embodiment 263, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is CAG and the amino acid corresponding to
10 the non-mutated codon is a glutamine.

265. The method of claim 264, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of glutamine at the position of the UAG stop codon.

15

265.1. The method of claim 264, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as glutamine, e.g., as provided in Table 2.

20 266. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UGA mutation, e.g., a UCA to UGA mutation.

267. The method of embodiment 266, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is UCA and the amino acid corresponding to
25 the non-mutated codon is a serine.

268. The method of claim 267, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of serine at the position of the UGA stop codon.

30

269. The method of claim 267, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as serine, e.g., as provided in Table 2.

5 270. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAG mutation, e.g., a UCG to UAG mutation.

271. The method of embodiment 270, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is UCG and the amino acid corresponding to
10 the non-mutated codon is a serine.

272. The method of claim 271, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of serine at the position of the UAG stop codon.

15 273. The method of claim 271, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as serine, e.g., as provided in Table 2.

20 274. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAA mutation, e.g., a UUA to UAA mutation.

275. The method of embodiment 274, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is UUA and the amino acid corresponding to
25 the non-mutated codon is a leucine.

276. The method of claim 275, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAA stop codon and mediates incorporation of leucine at the position of the UAA stop codon.

30

277. The method of claim 275, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAA stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as leucine, e.g., as provided in Table 2.

5 278. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UGA mutation, e.g., a UUA to UGA mutation.

279. The method of embodiment 278, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is UUA and the amino acid corresponding to
10 the non-mutated codon is a leucine.

280. The method of claim 279, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of leucine at the position of the UGA stop codon.

15 281. The method of claim 279, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as leucine, e.g., as provided in Table 2.

20 282. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UAG mutation, e.g., a UUG to UAG mutation.

283. The method of embodiment 282, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is UUG and the amino acid corresponding to
25 the non-mutated codon is a leucine.

284. The method of claim 283, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of leucine at the position of the UAG stop codon.

30

285. The method of claim 284, wherein the TREM, TREM core fragment or TREM fragment recognizes the UAG stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as leucine, e.g., as provided in Table 2.

- 5 286. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UGA mutation, e.g., a CGA to UGA mutation.

287. The method of embodiment 286, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is CGA and the amino acid corresponding to
10 the non-mutated codon is an arginine.

288. The method of claim 287, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of arginine at the position of the UGA stop codon.

15

289. The method of claim 287, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as arginine, e.g., as provided in Table 2.

- 20 290. The method of embodiment 126-160, wherein the codon having the first sequence or the PTC comprises a UGA mutation, e.g., a GGA to UGA mutation.

291. The method of embodiment 290, wherein the non-mutated, e.g., wildtype, codon sequence of the codon having the first sequence or the PTC is GGA and the amino acid corresponding to
25 the non-mutated codon is a glycine.

292. The method of claim 291, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of glycine at the position of the UGA stop codon.

30

293. The method of claim 291, wherein the TREM, TREM core fragment or TREM fragment recognizes the UGA stop codon and mediates incorporation of an amino acid which belongs to the same amino acid group as glycine, e.g., as provided in Table 2.

5 294. The method of any of embodiments 126-293, wherein incorporation of the amino acid by the TREM, TREM fragment or TREM core fragment results in modulation, e.g., increase, of a production parameter, e.g., an expression parameter and/or a signaling parameter, of an RNA corresponding to the ORF or a polypeptide encoded by the ORF.

10 295. The method of embodiment 294, wherein the production parameter comprises an expression parameter.

296. The method of embodiment 295, wherein the expression parameter comprises:

- (a) protein translation;
- 15 (b) expression level (e.g., of polypeptide or protein, or mRNA);
- (c) post-translational modification of polypeptide or protein;
- (d) folding (e.g., of polypeptide or protein, or mRNA),
- (e) structure (e.g., of polypeptide or protein, or mRNA),
- (f) transduction (e.g., of polypeptide or protein),
- 20 (g) compartmentalization (e.g., of polypeptide or protein, or mRNA),
- (h) incorporation (e.g., of polypeptide or protein, or mRNA) into a supermolecular structure, e.g., incorporation into a membrane, proteasome, or ribosome,
- (i) incorporation into a multimeric polypeptide, e.g., a homo or heterodimer, and/or
- (j) stability.

25 297. The method of embodiment 294, wherein the production parameter comprises a signaling parameter.

298. The method of embodiment 297, wherein the signaling parameter comprises:

(1) modulation of a signaling pathway, *e.g.*, a cellular signaling pathway which is downstream or upstream of the protein encoded by the endogenous ORF having a first sequence or PTC;

- (2) cell fate modulation;
- 5 (3) ribosome occupancy modulation;
- (4) protein translation modulation;
- (5) mRNA stability modulation;
- (6) protein folding and structure modulation;
- (7) protein transduction or compartmentalization modulation; and/or
- 10 (8) protein stability modulation.

299. The method of any one of embodiments 294-298, wherein the production parameter (*e.g.*, an expression parameter and/or a signaling parameter) may be modulated (*e.g.*, increased), *e.g.*, by at least 5% (*e.g.*, at least 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%,
15 100%, 150%, 200% or more), *e.g.*, compared to a reference sequence.

300. The method of any one of embodiments 126-299, wherein the subject has or has been identified as having a disorder or disease listed in any one of Tables 15, 16, or 17.

20 301. The method of any one of embodiments 126-299, wherein the cell is associated with, *e.g.*, obtained from a subject who has, a disorder or disease listed in any one of Tables 15, 16 or 17.

302. The method of embodiment 300 or 301, wherein the disorder or disease is chosen from the left column of Table 4.

25 303. The method of embodiment 300 or 301, wherein the disorder or disease is chosen from the left column of Table 4 and the codon having the first sequence or PTC is in a gene chosen from the right column of Table 4, optionally wherein the codon having the first sequence or PTC is at a position provided in Table 4.

30

304. The method of any one of embodiments 126-299, wherein the codon having the first sequence or PTC is in a gene chosen from the right column of Table 4, optionally wherein the codon having the first sequence or PTC is at a position provided in Table 4.

5 305. The method of embodiment 300 or 301, wherein the disorder or symptom is chosen from a disorder or disease provided in Table 5.

306. The method of embodiment 300 or 301, wherein the disorder or symptom is chosen from a disorder or disease provided in Table 6.

10

307. The method of embodiment 300 or 301, wherein the disorder or symptom is chosen from a disorder or disease provided in Table 6 and the codon having the first sequence or PTC is in any gene provided in Table 6.

15 308. The method of embodiment 300 or 301, wherein the disorder or symptom is chosen from a disorder or disease provided in Table 6 and the codon having the first sequence or PTC is in a corresponding gene provided in Table 6, e.g., a gene corresponding to the disease or disorder.

20 309. The method of embodiment 300 or 301, wherein the disorder or symptom is chosen from a disorder or disease provided in Table 6 and the codon having the first sequence or PTC is not in a gene provided in Table 6.

310. The method of any one of embodiments 126-299, wherein the codon having the first sequence or PTC is in a gene provided in Table 3.

25

311. The method of any one of embodiments 303, 304, 307, 308, 309 or 310, wherein the codon having the first sequence or PTC is at any position within the ORF of the gene, e.g., upstream of the naturally occurring stop codon.

30 Other features, objects, and advantages of the invention will be apparent from the description and from the claims.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In addition, the materials, methods, and examples are
 5 illustrative only and not intended to be limiting.

EXAMPLES

The following examples are provided to further illustrate some embodiments of the present invention, but are not intended to limit the scope of the invention; it will be understood
 10 by their exemplary nature that other procedures, methodologies, or techniques known to those skilled in the art may alternatively be used.

Table of Contents for Examples

Example 1	Synthesis of guanosine 2'-O-MOE phosphoramidite
Example 2	Synthesis of 5,6 dihydrouridine
Example 3	Synthesis of a TREM via 5-Silyl-2-Orthoester (2-ACE) Chemistry
Example 4	Synthesis of an arginine TREM having a 2'-O-MOE modification
Example 5	Synthesis of an arginine TREM having a pseudouridine and a 2'-O-MOE modification
Example 6	Synthesis of a glutamine TREM having a 5,6 dihydrouridine modification
Example 7	Synthesis of a glutamine TREM having a pseudouridine modification
Example 8	Synthesis of nucleotides comprising an aminonucleobase (AN1)
Example 9	Synthesis of biotin conjugated TREM molecules
Example 10	Quality control of synthesized TREM via Mass Spectrometry Analysis
Example 11	Quality control of synthesized TREM via anion-exchange HPLC
Example 12	Quality control of synthesized TREM via PAGE Purification and Analysis
Example 13	Deprotection of synthesized TREM
Example 14	Readthrough of a premature termination codon (PTC) in a reporter protein with administration of a synthetic arginine non-cognate TREM (1)

Example 15	Readthrough of a premature termination codon (PTC) in a reporter protein with administration of a synthetic arginine non-cognate TREM (2)
Example 16	Readthrough of a premature termination codon (PTC) in the Coagulation Factor IX ORF through administration of a synthetic arginine non-cognate TREM
Example 17	Correction of a missense mutation in an ORF with administration of a TREM

Example 1: Synthesis of guanosine 2'-O-MOE phosphoramidite

This example describes the synthesis of guanosine 2'-O-MOE phosphoramidite. Guanosine 2'-O-MOE phosphoramidite is prepared and purified according to previously published procedures (Wen K. et al. (2002) *The Journal of Organic Chemistry*, 67(22), 7887–7889).

Briefly, guanosine and imidazole are dried by co-evaporation with pyridine, dissolved in dry DMF, and treated with bis(diisopropylchlorosilyl) methane added dropwise at 0 °C. The temperature is gradually increased to 25 °C and then held for 5 h. The reaction mixture is poured into ice water, and the precipitated white solid filtered to afford compound 1. To a solution of compound 1, BrCH₂CH₂OCH₃, and TBAI in DMF at -20 °C is added with sodium bis(trimethylsilyl)amide, and the mixture is stirred for 4 hours under argon. After the reaction is quenched with methanol, the THF is evaporated and the residue is precipitated in ice to furnish compound 2. TBAF is added to a solution of compound 2 at 25 °C and then the mixture is stirred at 35 °C for 5 hours. The solvent is then evaporated under reduced pressure, and the residue is filtered in a short pad of silica gel using 10% methanol in dichloromethane to afford guanosine 2'-O-MOE phosphoramidite.

Example 2: Synthesis of 5,6 dihydrouridine

This example describes the synthesis of 5,6 dihydrouridine. 5,6 dihydrouridine phosphoramidite is prepared and purified according to previously published procedures (Hanze AR et al., (1967) *Journal of the American Chemical Society*, 89(25), 6720–6725). Briefly, oxygen is bubbled through a solution uridine in the presence of platinum black. The reaction is followed by spotting the reaction mixture on silica gel thin layer chromatographic plates and developing in methanol-chloroform (1:1). After 1 hour, the mixture is cooled and centrifuged and the clear liquid lyophilized to yield the 5,6 dihydrouridine product.

Example 3: Synthesis of a TREM via 5'-Silyl-2'-Orthoester (2'-ACE) Chemistry

This example describes the synthesis of a TREM via 5'-Silyl-2'-Orthoester (2'-ACE) Chemistry summarized from (Hartsel SA et al., (2005) *Oligonucleotide Synthesis*, 033–050).

Protected Ribonucleoside Monomers

5 5'-O-silyl-2'-O-ACE protected phosphoramidites are prepared and purified according to previously published procedures (Hartsel SA et al., (2005) *Oligonucleotide Synthesis*, 033–050). Briefly, monomer synthesis begins from standard base-protected ribonucleosides [rA(ibu), rC(acetyl), rG(ibu) and U]. Orthogonal, 5'-silyl-2'-ACE protection and amidite preparation is then accomplished in five general steps:

- 10 1. Simultaneous transient protection of the 5' and 3' hydroxyl groups with 1,1,3,3-tetraisopropylidisiloxane (TIPS).
2. Regiospecific conversion of the 2' hydroxyl to the 2'-O-orthoester using tris(acetoxyethyl)orthoformate (ACE orthoformate).
3. Removal of the 5',3'-TIPS protection.
- 15 4. Introduction of the 5'-O-silyl ether protecting group using benzhydryloxybis-(trimethylsilyloxy)-chlorosilane (BzH-Cl).
5. Phosphitylation of the 3'-OH with bis(N,N-diisopropylamino)methoxyphosphine.

The fully protected, phosphitylated monomer is an oil. For ease of handling and dissolution, the phosphoramidite solution is evaporated to dryness in a tared flask to enable quantitation of yields. The phosphoramidite oil is then dissolved in anhydrous acetonitrile, distributed into synthesis vials in 1.0-mmol aliquots, and evaporated to dryness under vacuum in the presence of

20 potassium hydroxide (KOH) and P₂O₅.

*Synthesis of Oligoribonucleosides***Table 16**

Synthesis Step	Reagent	Delivery Time	Reaction Time
Deblock	3% DCA in DCM	35	
Activator	0.5M S-ethyl-tetrazole	6	
Coupling	0.1M amidite8.0	30	
	0.5M S-ethyl-tetrazole	8	30
<i>Repeat Coupling</i>			

Oxidation	t-Butyl hydroperoxide	20	10
<i>Repeat Oxidation Delivery</i>			
Capping	1-methylimidazole and acetic anhydride	12	10
Desilylation	TEAHF	35	

5'-silyl-2'-ACE oligoribonucleotide synthesis begins with the appropriately modified 3'-terminal nucleoside attached through the 3'-hydroxyl to a polystyrene support. The solid support contained in an appropriate reaction cartridge is then placed on the appropriate column position on the instrument. A synthesis cycle is created using the delivery times and wait steps outlined in Table 16.

1. Initial detritylation: The first step in the synthesis cycle is the removal of the 5'-DMT from the nucleoside-bound polystyrene support using 3% DCA in DCM.
2. Coupling: The 5-ethylthio-1H-tetrazole solution is delivered to the solid support,
10 followed by simultaneous delivery of an equal quantity of activator and phosphoramidite solution. Depending on the desired sequence and synthesis scale, excess activator and activator plus amidite are alternately delivered repeatedly to increase coupling efficiency, which is typically in excess of 99% per coupling reaction. The 5-ethylthio-1H-tetrazole activates coupling by protonating the diisopropyl amine attached to the trivalent
15 phosphorous. Nucleophilic attack of the 5-ethylthio-1H-tetrazole leads to the formation of the tetrazolide intermediate that reacts with the free 5'-OH of the support-bound nucleoside forming the internucleotide phosphite linkage.
3. Oxidation: In the next step of chain elongation, the phosphorous(III) linkage is oxidized for 10-20s to the more stable and ultimately desired P(V) linkage using t-
20 butylhydroperoxide.
4. Capping: Although delivery of excess activator and phosphoramidite increases coupling efficiency, a small percentage of unreacted nucleoside may remain support-bound. To prevent the introduction of mixed sequences, the unreacted 5'-OH are "capped" or blocked by acetylating the primary hydroxyl. This acetylation is achieved through
25 simultaneous delivery of 1-methylimidazole and acetic anhydride.

5. 5'-Desilylation: Before the next nucleoside in the sequence can be added to the growing oligonucleotide chain, the 5'-silyl group is removed with fluoride ion. This requires the delivery of triethylamine trihydrogenfluoride for 45 s. The desilylation is rapid and quantitative and no wait step is required.

- 5 Steps 2-5 are repeated for each subsequent nucleotide until the desired sequence is constructed.

Oligonucleotide Deprotection

- A two-stage rapid deprotection strategy is employed to remove phosphate backbone protection, release the oligonucleotide from the solid support, and remove the exocyclic amine protecting groups on A, G, and C. The treatment also removes the acetyl moiety from the acetoxyethyl orthoester, resulting in the 2'-bis-hydroxyethyl protected intermediate that is now 10 times more labile to final acid deprotection. In the first deprotection step, S₂Na₂ is used to selectively remove the methyl protection from the internucleotide phosphate, leaving the oligoribonucleotide attached to the polystyrene support. This configuration allows any residual reagent to be thoroughly washed away before proceeding. Alternatively, a multicolumn, manifold approach can also be used.
1. A syringe barrel is attached to one of the two luer fittings on the synthesis column. 2 mL of the S₂Na₂ reagent is drawn into a second syringe and attached to the opposite side of the synthesis column. The S₂Na₂ reagent is gently pushed through the column and into the empty syringe barrel continuing back and forth several times. The column, filled with reagent is allowed to sit at room temperature for 10 min.
 2. S₂Na₂ reagent is removed from the column. Using a clean syringe, the column is washed thoroughly with water. In the second deprotection step, 40% 1-methylamine in water is used to free the oligoribonucleotide from the solid support, deprotect the exocyclic base amines, and deacylate the 2'-orthoester leaving the deprotected species.

N-Methylamine Deprotection

1. The solid support resin is transferred from the column into a 4-mL vial
 2. 2 mL 40% methylamine is added and heated for 12 min at 60°C.
 3. The methylamine is removed and is transferred into a fresh vial.
 4. The oligonucleotide solution is evaporated to dryness in a SpeedVac or similar device.
- Oligonucleotide yields are measured using an ultraviolet (UV) spectrophotometer (absorbance at 260 nm).

Example 4: Synthesis of an arginine TREM having a 2'-O-MOE modification

This example describes the synthesis of an Arg TREM having one 2'-O-MOE modification. The 2'-O-MOE modification can be placed on a nucleotide on any domain or linker of the Arg TREM, or at any position in said domain or linker.

- 5 A 2 $\times$ ACE RNA oligoribonucleotide synthesis is performed on a modified Applied Biosystems 394 DNA/RNA synthesizer or similar instrument. 2'-O-MOE amidites are synthesized as in Example 2. An oligonucleotide sequence:
GGCUCCGUGGCGCAAUGGAUAGCGCAUUGGACUUCUAAUUCAAAGGUUCCGGGUU
CG(A-MOE)GUCCCCGGCGGAGUCG is synthesized following the protocol described in
10 example 4. A similar method can be used to add a 2'-O-MOE modification on a TREM specifying any one of the other 19 amino acids.

Example 5: Synthesis of an arginine TREM having a pseudouridine and a 2'-O-MOE modification

- This example describes the synthesis of an Arg TREM having a pseudouridine and 2'-O-MOE modification. The modification can be placed on a nucleotide on any domain or linker of
15 the Arg TREM, or at any position in said domain or linker.

- A 2 $\times$ ACE RNA oligoribonucleotide synthesis is performed on a modified Applied Biosystems 394 DNA/RNA synthesizer or similar instrument. 2'-O-MOE amidites are synthesized as in example 1. Pseudouridine (P) amidites are obtained from Glen Research or
20 similar provider. An oligonucleotide sequence:
GGCUCCGUGGCGCAAUGGAUAGCGCAPUGGACUUCUAAUUCAAAGGUUCCGGGUU
CG(A-MOE)GUCCCCGGCGGAGUCG is synthesized following the protocol described in
example 3. A similar method can be used to add a pseudouridine and 2'-O-MOE modification on
a TREM specifying any one of the other 19 amino acids.

- 25 **Example 6: Synthesis of a glutamine TREM having a dihydrouridine modification**

This example describes the synthesis of a Gln TREM having a dihydrouridine modification. The modification can be placed on a nucleotide on any domain or linker of the Gln TREM, or at any position in said domain or linker.

- A 2 $\times$ ACE RNA oligoribonucleotide synthesis is performed on a modified Applied
30 Biosystems 394 DNA/RNA synthesizer or similar instrument. Dihydrouridine (D) is synthesized as in example 2. An oligonucleotide sequence:

GGUCCAUGGUGUAAUGGDAAGCACUCUGGACUCTGAAUCCAGCGAUCCGAGUUC
GAGUCUCGGUGGAACCUCCA is synthesized following the protocol described in example 3.
A similar method can be used to add a dihydrouridine modification on a TREM specifying any
one of the other 19 amino acids.

5 **Example 7: Synthesis of a glutamine TREM having a pseudouridine modification**


This example describes the synthesis of a Gln TREM having a pseudouridine
modification. The modification can be placed on a nucleotide on any domain or linker of the Gln
TREM, or at any position in said domain or linker.

A 2 $\times$ ACE RNA oligoribonucleotide synthesis is performed on a modified Applied
10 Biosystems 394 DNA/RNA synthesizer or similar instrument. Pseudouridine (P) amidites are
obtained from Glen Research or similar provider. An oligonucleotide sequence:
GGUCCAUGGUGPAAUGGUAAGCACUCUGGACUCTGAAUCCAGCGAUCCGAGUUC
GAGUCUCGGUGGAACCUCCA is synthesized following the protocol described in example 3.

A similar method can be used to add a pseudouridine modification on a TREM specifying
15 any one of the other 19 amino acids.

Example 8: Synthesis of nucleotides comprising an aminonucleobase (AN1)


Modified nucleotides comprising an amine handle at the nucleobase, such as AN1 (C6-U
phosphoramidite (5'-Dimethoxytrityl-5-[N-(trifluoroacetylaminohexyl)-3-acrylimido]-Uridine,
2'-O-triisopropylsilyloxymethyl-3'-[(2-cyanoethyl)-(N,N-diisopropyl)]-phosphoramidite)), may
20 be purchased from Glen Research; catalog # 10-3039. Briefly, Amino-Modifier C6-U
phosphoramidite was purchased with the primary amine protected as trifluoroacetate and
incorporated into a TREM to afford the amino nucleobase AN1.



Example 9: Synthesis of biotin conjugated TREM molecules

This example describes the synthesis of biotin conjugated TREM molecule. These molecules may be utilized as test TREMs (e.g., test chemically modified TREMs) for example, and be useful for investigation of which positions along the TREM sequence are suitable for labeling (+)-Biotin N-hydroxysuccinimide ester may be purchased from Sigma-Aldrich (catalog # H1759). The TREM molecules bearing a free amine may be synthesized as described previously, e.g., Example 8, then coupled with (+)-Biotin N-hydroxysuccinimide ester to form an amide bond, according to the method, e.g., as outlined in Bengstrom M. et al. (1990) Nucleos. Nucleot. Nucl. 9, 123-127. Briefly, a solution of TREM molecules with amino base modification and excess (+)-Biotin N-hydroxysuccinimide ester may be mixed together and vortexed for several hours at 37°C. LCMS analysis is used to determine whether the reaction is complete. The solvent is removed under vacuum, and the resulting residue is diluted with water then subjected to purification using reversed phase column chromatography to afford the final compound.

For example, the biotin moiety was installed on the arginine non-cognate TREM molecules at position 47 named TREM-Arg-TGA-Biotin-47. The arginine non-cognate TREM molecules contain the sequence of ARG-UCU-TREM body but with the anticodon sequence corresponding to UCA instead of UCU.



Example 10: Quality control of synthesized TREM via Mass Spectrometry Analysis

This example describes the quality control of a synthesized TREM via Mass Spectrometry Analysis.

Using the Perseptive Biosystems Voyager-DE BioSpectrometry Workstation, the referenced protocol for mass spectrometry analysis (4- Van Ausdall) is followed. Briefly, a 3-hydroxy picolinic acid matrix is used for sample crystallization. It is prepared by mixing (10:1:1)

3-HPA:picolinic acid:ammonium hydrogen citrate where each component is dissolved in 30% aqueous acetonitrile at a concentration of 50 mg/mL. One optical density unit (ODU) of oligonucleotide is dissolved in the matrix and heated at 55°C for 10 min. The sample is spotted on a MALDI plate, allowed to dry, and analyzed accordingly. This method allows confirmation of oligonucleotide identity and detection of low-level impurities present in synthetic oligonucleotide samples.

Example 11: Quality control of synthesized TREM via anion-exchange HPLC

This example describes the quality control of a synthesized TREM via anion-exchange HPLC. Using the Dionex DNA-Pac-PA-100 column, a gradient is employed using HPLC buffer A and HPLC buffer B. 0.5 ODU of a sample that has been dissolved in H₂O or Tris buffer, pH 7.5 is injected onto the gradient. The gradient employed is based on oligonucleotide length and can be applied according to Table 17. The parameters provided in Table 18 can be used to program a linear gradient on the HPLC analyzer.

Table 17: Oligonucleotide length and gradient percentages

Length (bases)	Gradient (%B)
0–5	0–30
6–10	10–40
11–16	20–50
17–32	30–60
33–50	40–70
>50	50–80

Table 18: Parameters for a linear gradient on HPLC analyzer

Time (min)	Flow (mL/min)	% Buffer A	% Buffer B
0	1.5	100	0
1	1.5	100	0
3	1.5	70a	30a
15	1.5	40a	60a

15.5	2.5	0	100
17	2.5	0	100
17.25	2.5	100	0
23	2.5	100	0
s23.1	1.5	100	0
24	1.5	100	0
25	0.1	100	0

Example 12: Quality control of synthesized TREM via PAGE Purification and Analysis

This example describes the quality control of a synthesized TREM via PAGE Purification and Analysis. Gel purification and analysis of 2'-ACE protected RNA follows standard protocols for denaturing PAGE (Ellington and Pollard (1998) *In Current Protocols in Molecular Biology*, Chanda, V). Briefly, the 2'-ACE protected oligo is resuspended in 200 mL of gel loading buffer. Invitrogen™ NuPAGE™ 4-12% Bis-Tris Gels or similar gel is prepared in gel apparatus. Samples are loaded and gel ran at 50-120 W, maintaining the apparatus at 40°C. When complete, the gel is exposed to ultraviolet (UV) light at 254 nm to visualize the purity of the RNA using UV shadowing. If necessary, the desired gel band is excised with a clean razor blade. The gel slice is crushed and 0.3M NaOAc elution buffer is added to the gel particles, and soaked overnight. The mixture is decanted and filtered through a Sephadex column such as Nap-10 or Nap-25.

Example 13: Deprotection of synthesized TREM

This example describes the deprotection of a TREM made according to an in vitro synthesis method, e.g., as described in Example 3. The 2'-protecting groups are removed using 100 mM acetic acid, pH 3.8. The formic acid and ethylene glycol byproducts are removed by incubating at 60°C for 30 min followed by lyophilization or SpeedVac-ing to dryness. After this final deprotection step, the oligonucleotides are ready for use.

Example 14: Readthrough of a premature termination codon (PTC) in a reporter protein with administration of an arginine non-cognate TREM (1)

This example describes an assay to test the ability of a non-cognate TREM to readthrough a PTC in a cell line expressing a reporter protein having a PTC. This Example

describes an arginine non-cognate TREM. A non-cognate TREM specifying any one of the other 19 amino acids can be used.

Host cell modification

A cell line stably expressing a NanoLuc reporter construct containing a premature termination codon (PTC) is generated using the FlpIn system according to manufacturer's instructions. Briefly, HEK293T (293T ATCC® CRL-3216) cells are co-transfected with an expression vector containing a NanoLuc reporter with a PTC, such as pcDNA5/FRT-NanoLuc-TAA and a pOG44 Flp-Recombinase expression vector using Lipofectamine2000 according to manufacturer's instructions. After 24 hours, the media is replaced with fresh media. The next day, the cells are split 1:2 and selected with 100ug/mL Hygromycin for 5 days. The remaining cells are expanded and tested for reporter construct expression.

Synthesis and preparation of non-cognate TREM

In this example, the arginine non-cognate TREM, is produced such that it contains the sequence of the ARG-UCU-TREM body but with the anticodon sequence corresponding to UCA instead of UCU. The arginine non-cognate TREM is synthesized as described herein and quality control methods as described herein are performed. To ensure proper folding, the TREM is heated at 85°C for 2 minutes and then snap cooled at 4°C for 5 minutes.

Transfection of non-cognate TREM into host cells

To deliver the arginine non-cognate TREM to mammalian cells, 100 nM of TREM is transfected into HEK293T (293T ATCC® CRL-3216), U2OS (U-2 OS (ATCC® HTB-96™)), H1299 (NCI-H1299 (ATCC® CRL-5803™)), or HeLa (HeLa (ATCC® CCL-2™)) cells stably expressing the PTC-containing NanoLuc reporter using lipofectamine 2000 reagents according to the manufacturer's instructions. After 6-18 hours, the transfection media is removed and replaced with fresh complete media (U2OS: McCoy's 5A, 10% FBS, 1%PenStrep; H1299: RPMI1640, 10% FBS, 1%PenStrep; Hek/HeLa: EMEM, 10% FBS, 1%PenStrep).

Translation suppression assay

To monitor the efficacy of the arginine non-cognate TREM to readthrough the PTC in the reporter construct, 24-48 hours after transfection, cell media is replaced and allowed to equilibrate to room temperature. An equal volume to the cell media of ONE-Glo™ EX Reagent is added to the well and mixed on the orbital shaker at 500rpm for 3 min followed by addition of an equal volume of cell media of NanoDLR™ Stop & Glo and mixing on the orbital shaker at

500rpm for 3 min. The reaction is incubated at room temperature for 10min and the NanoLuc activity is detected by reading the luminescence in a plate reader. As a positive control, a host cell expressing the NanoLuc reporter construct without a PTC is used. As a negative control, a host cell expressing the NanoLuc reporter construct with a PTC is used but no TREM is transfected. The TREM efficacy is measured as a ratio of the NanoLuc luminescence in the experimental sample to the NanoLuc luminescence of the positive control. It is expected that if the arginine non-cognate TREM is functional, it can read-through the stop mutation in the NanoLuc reporter and produce a luminescent reading higher than the luminescent reading measured in the negative control. If the arginine non-cognate TREM is not functional, the stop mutation is not rescued, and luminescence less or equal to the negative control is detected.

Example 15: Readthrough of a premature termination codon (PTC) in a reporter protein with administration of an arginine non-cognate TREM (2)

This example describes an assay to test the ability of a non-cognate TREM to readthrough a PTC in a cell line expressing a reporter protein having a PTC. This Example describes an arginine non-cognate TREM. A non-cognate TREM specifying any one of the other 19 amino acids can be used.

Host cell modification

A cell line engineered to stably express a HiBiT-tagged disease reporter construct containing a premature termination codon (PTC), such as Factor IX at position 298 (*FIX*^{R298X}), Tripeptidyl-peptidase 1 at position 208 (*TPPI*^{R208X}), or Protocadherin Related 15 at position 245 (*PCDH15*^{R245X}), was generated using the Jump-In system according to manufacturer's instructions. Briefly, Jump-In GripTite HEK293 (Thermo Scientific A14150) cells were co-transfected with an expression vector containing the disease reporter, such as pJTI-R4-DEST-CMV-FIX-R298X-HiBiT-pA for *FIX*^{R298X} to make the Factor IX disease reporter expressing cell line, and a pJTI-R4-Int PhiC31 integrase expression vector using Lipofectamine2000 according to manufacturer's instructions. After 24 hours, the media was replaced with fresh media. The next day, the cells were re-seeded at 50% confluency and selected with 10ug/mL Blasticidin and 600ug/mL G418 for 7 days with media change every 2 days. The remaining cells were expanded and tested for reporter construct expression.

Synthesis and preparation of non-cognate TREM

In this example, the modified arginine non-cognate TREMs were produced such that they contain the sequence of the ARG-UCU-TREM body but with the anticodon sequence corresponding to UCA instead of UCU and modified as described herein. The resulting TREMs may be modified, for example, to contain a biotin as in Example 8-9. To ensure proper folding, the TREM was heated at 85°C for 2 minutes and then snap cooled at 4°C for 5 minutes.

Transfection of non-cognate TREM into host cells

Forty-eight hours after TREM delivery into cells, conditioned media was collected, fresh media was added to the cells, and allowed to equilibrate to room temperature. To measure the efficacy of arginine non-cognate TREMs in PTC readthrough, full-length HiBiT-tagged disease reporter protein was assayed in both cells, and 48-hour conditioned media. Briefly, reconstituted Nano-Glo® HiBiT Lytic Reagent was added to both cells containing fresh media, and 48-hour conditioned media at a 1:1 v/v ratio, mixed on an orbital shaker at 500rpm for 10 minutes, incubated at room temperature for 10 minutes, and the HiBiT-NanoLuc activity is measured by reading the luminescence in a plate reader.

Translation suppression assay

To monitor the efficacy of the arginine non-cognate TREM to readthrough the PTC in the reporter construct, Forty-eight hours after TREM delivery into cells, conditioned media was collected, fresh media was added to the cells, and allowed to equilibrate to room temperature. To measure the efficacy of arginine non-cognate TREMs in PTC readthrough, full-length HiBiT-tagged disease reporter protein was assayed in both cells, and 48-hour conditioned media. Briefly, reconstituted Nano-Glo® HiBiT Lytic Reagent was added to both cells containing fresh media, and 48-hour conditioned media at a 1:1 v/v ratio, mixed on an orbital shaker at 500rpm for 10 minutes, incubated at room temperature for 10 minutes, and the HiBiT-NanoLuc activity is measured by reading the luminescence in a plate reader. The results of this experiment in the three HiBiT-tagged disease reporter constructs is shown in FIGS. 1A-1C.

Example 16: Readthrough of a premature termination codon (PTC) in the Coagulation Factor IX ORF through administration of a synthetic arginine non-cognate TREM

This example describes an assay to test the ability of a non-cognate arginine TREM to readthrough a PTC, such as R252X or R333X, in the Coagulation Factor IX open reading frame (ORF) in a Hemophilia B patient-derived cell line. This Example describes an arginine non-

cognate TREM. A non-cognate TREM specifying any one of the other 19 amino acids can be used.

Patient-derived cells

Fibroblast cells derived from a patient with Hemophilia B having a PTC in the Coagulation Factor IX open reading frame (ORF), such as R252X or R333X, is obtained from a center or an organization, such as the Coriell Institute. The patient-derived fibroblast cells are reprogrammed into hepatocytes as previously shown (Takahashi, K. & Yamanaka, S. (2006) *Cell* 126, 663–676 (2006); Park I. et al. (2008) *Nature* 451, 141–146); Jia, B. et al. (2014) *Life Sci.* 108, 22–29).

Synthesis and preparation of TREM

In this example, the arginine non-cognate TREM, is produced such that it contains the sequence of the ARG-UCU-TREM body but with the anticodon sequence corresponding to UCA instead of UCU. The arginine TREM is synthesized as described in Examples 3–7 and quality control methods as described in Examples herein are performed. To ensure proper folding, the TREM is heated at 85°C for 2 minutes and then snap cooled at 4°C for 5 minutes.

Transfection of non-cognate TREM into host cells

To deliver the arginine TREM to mammalian cells, 100 nM of TREM is transfected into the reprogrammed hepatocyte cells using lipofectamine 2000 reagents according to the manufacturer's instructions. After 6–18 hours, the transfection media is removed and replaced with fresh complete media.

Translation suppression assay

To monitor the efficacy of the arginine non-cognate TREM to readthrough the PTC in the Coagulation Factor IX ORF, 24–48 hours after transfection, cell media is replaced, and cells are lysed. Using Western blot detection, the non-cognate TREM efficacy is measured as the level of full-length protein expression, in this example of Coagulation Factor IX protein, in the reprogrammed hepatocyte cells administered the Arg non-cognate TREM, in comparison to the Coagulation Factor IX protein expression levels found in control cells. For example, as a control, cells of a person unaffected by the disease (i.e. cells having an ORF with a WT Coagulation Factor IX transcript) can be used. It is expected that if the non-cognate TREM is functional, it can read-through the PTC and the full-length protein level will be detected at higher levels than that found in patient-derived fibroblast cells or reprogrammed hepatocyte cells which have not

been administered the non-cognate TREM. If the non-cognate TREM is not functional, the full-length protein level will be detected at a similar level as detected in patient-derived fibroblast cells or reprogrammed hepatocyte cells which have not been administered the non-cognate TREM.

5 **Example 17: Correction of a missense mutation in an ORF with administration of a TREM**

This example describes the administration of a TREM to correct a missense mutation. In this example, a TREM translates a reporter with a missense mutation into a wild type (WT) protein by incorporation of the WT amino acid (at the missense position) in the protein.

Host cell modification

10 A cell line stably expressing a GFP reporter construct containing a missense mutation, for example T203I or E222G, which prevent GFP excitation at the 470 nm and 390 nm wavelengths, is generated using the FlpIn system according to manufacturer's instructions. Briefly, HEK293T (293T ATCC ® CRL-3216) cells are co-transfected with an expression vector containing a GFP reporter with a missense mutation, such as pcDNA5/FRT-NanoLuc-TAA and a pOG44 Flp-Recombinase expression vector using Lipofectamine2000 according to manufacturer's
15 instructions. After 24 hours, the media is replaced with fresh media. The next day, the cells are split 1:2 and selected with 100ug/mL Hygromycin for 5 days. The remaining cells are expanded and tested for reporter construct expression.

Synthesis and preparation of TREM

20 The TREM is synthesized as described in Examples 3-7 and quality control methods as described in Examples 8-10 are performed. To ensure proper folding, the TREM is heated at 85°C for 2 minutes and then snap cooled at 4°C for 5 minutes.

Transfection of non-cognate TREM into host cells

To deliver the TREM to mammalian cells, 100 nM of TREM is transfected into cells
25 expressing the ORF having a missense mutation using lipofectamine 2000 reagents according to the manufacturer's instructions. After 6-18 hours, the transfection media is removed and replaced with fresh complete media.

Missense mutation correction assay

To monitor the efficacy of the TREM to correct the missense mutation in the reporter
30 construct, 24-48 hours after TREM transfection, cell media is replaced, and cell fluorescence is measured. As a negative control, no TREM is transfected in the cells and as a positive control,

cells expressing WT GFP are used for this assay. If the TREM is functional, it is expected that the GFP protein produced fluoresces when illuminated with a 390 nm excitation wavelength using a fluorimeter, as observed in the positive control. If the TREM is not functional, the GFP protein produced fluoresces only when excited with a 470 nm wavelength, as is observed in the

5 negative control.

What is claimed is:

1 A method of modulating a production parameter of an mRNA corresponding to, or polypeptide encoded by, an endogenous open reading frame (ORF) in a cell, which ORF comprises a codon having a first sequence, comprising:

contacting the cell with a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein in an amount and/or for a time sufficient to modulate the production parameter of the mRNA or polypeptide,

wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,

thereby modulating the production parameter in the cell.

2. A method of modulating a production parameter of an mRNA corresponding to, or polypeptide encoded by, an endogenous open reading frame (ORF) in a subject, which ORF comprises a codon having a first sequence, comprising:

contacting the subject with a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein in an amount and/or for a time sufficient to modulate the production parameter of the mRNA or polypeptide,

wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,

thereby modulating the production parameter in the subject.

3. The method of claim 1 or 2, wherein the production parameter comprises a signaling parameter, e.g., as described herein.

4. The method of claim 1 or 2, wherein the production parameter comprises an expression parameter, e.g., as described herein.

5. A method of modulating expression of a protein in a cell, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a codon having a first sequence, comprising:

contacting the cell with a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein in an amount and/or for a time sufficient to modulate expression of the encoded protein,

wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,

thereby modulating expression of the protein in the cell.

6. A method of modulating expression of a protein in a subject, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a codon having a first sequence, comprising:

contacting the subject with a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein in an amount and/or for a time sufficient to modulate expression of the encoded protein,

wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,

thereby modulating expression of the protein in the subject.

7. A method of treating a subject having an endogenous open reading frame (ORF) which comprises a codon having a first sequence, comprising:

providing a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein wherein the TREM comprises a tRNA moiety having: an anticodon that pairs with the codon of the ORF having the first sequence;

contacting the subject with the composition comprising a TREM, TREM core fragment or TREM fragment in an amount and/or for a time sufficient to treat the subject,

thereby treating the subject.

8. A method of treating a subject having an endogenous open reading frame (ORF) comprising a codon having a first sequence, comprising:

(i) acquiring, e.g., directly or indirectly acquiring, a value for the status of the codon having the first sequence in the subject, wherein said value comprises a measure of the presence

or absence of the codon having the first sequence in a sample from the subject; and identifying the subject as having the codon having the first sequence; and

(ii) responsive to said value, administering a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein wherein the TREM, TREM core fragment or TREM fragment comprises a tRNA moiety having an anticodon that pairs with the codon having the first sequence, to the subject,

thereby treating the subject.

9. A method of evaluating a subject having an endogenous open reading frame (ORF) comprising a codon having a first sequence, comprising:

acquiring, e.g., directly or indirectly acquiring, a value for the status of the codon having the first sequence in the subject, wherein said value comprises a measure of the presence or absence of the codon having the first sequence in a sample from the subject; and

identifying the subject as having a codon having the first sequence,

thereby evaluating the subject.

10. The method of claim 9, wherein responsive to said value the method further comprises administering a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein wherein the TREM, TREM core fragment or TREM fragment comprises a tRNA moiety having an anticodon that pairs with the codon having the first sequence, to the subject.

11. A method of modulating a production parameter of an mRNA corresponding to, or polypeptide encoded by, an endogenous open reading frame (ORF) in a cell, which ORF comprises a premature termination codon (PTC),

contacting the cell with a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein in an amount and/or for a time sufficient to modulate the production parameter of the mRNA or polypeptide,

wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,

thereby modulating the production parameter in the cell.

12. A method of modulating a production parameter of an mRNA corresponding to, or polypeptide encoded by, an endogenous open reading frame (ORF) in a subject, which ORF comprises a premature termination codon (PTC),
- contacting the subject with a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein in an amount and/or for a time sufficient to modulate the production parameter of the mRNA or polypeptide,
- wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the codon having the first sequence,
- thereby modulating the production parameter in the subject.
13. The method of claim 11 or 12, wherein the production parameter comprises a signaling parameter and/or an expression parameter, e.g., as described herein.
14. A composition for use in treating a subject having an endogenous open reading frame (ORF) which comprises a premature termination codon (PTC), wherein the composition comprises a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein, wherein the TREM comprises a tRNA moiety having an anticodon that pairs with the PTC in the ORF.
15. A composition for use in modulating expression of a protein in a cell, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a premature termination codon (PTC) wherein the composition comprises a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed herein, and wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the PTC.
16. A composition for use in modulating expression of a protein in a subject, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a premature termination codon (PTC), wherein the composition comprises a TREM composition comprising a TREM, a TREM core fragment, or a TREM fragment disclosed

herein, and wherein the TREM, TREM core fragment or TREM fragment has an anticodon that pairs with the PTC.

17. The composition for use of any one of claims 14-16, wherein the PTC comprises UAA, UGA or UAG.

18. A TREM composition for use in increasing expression of a protein in a subject wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a premature termination codon (PTC), wherein the TREM composition

- (i) has an anticodon that pairs with the PTC,
- (ii) recognizes an aminoacyl-tRNA synthetase specific for Trp, Tyr, Cys, Glu, Lys, Gln, Ser, Leu, Arg, or Gly,
- (iii) comprises a sequence of Formula A, and
- (iv) comprises one or more of a 2'-O-MOE, pseudouridine or 5,6 dihydrouridine modification.

19. A TREM composition for use in increasing expression of a protein in a cell or subject, wherein the protein is encoded by a nucleic acid comprising an endogenous open reading frame (ORF), which ORF comprises a premature termination codon (PTC), wherein the TREM composition::

- (i) has an anticodon that pairs with the PTC,
- (ii) recognizes an aminoacyl-tRNA synthetase specific for Trp, Tyr, Cys, Glu, Lys, Gln, Ser, Leu, Arg, or Gly,
- (iii) comprises a sequence of Formula B, and
- (iv) comprises one or more of a 2'-O-MOE, pseudouridine or 5,6 dihydrouridine modification.

20. The TREM composition of claim 18 or 19, wherein the PTC comprises UAA, UGA or UAG.

21. The method or composition for use of any one of the preceding claims, wherein the codon having the first sequence or the PTC comprises a UAA mutation.

22. The method or composition for use of any one of the preceding claims, wherein the codon having the first sequence or the PTC comprises a UGA mutation.

23. The method or composition for use of any one of the preceding claims, wherein the codon having the first sequence or the PTC comprises a UAG mutation.

24. The method or composition for use of any one of claims 1-23, wherein the codon having the first sequence or the PTC comprises a UAA, UGA or UAA mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid which preserves, e.g., maintains, a secondary and/or tertiary structure of a polypeptide encoded by the ORF into which the amino acid is incorporated.

25. The method or composition for use of any one of claims 1-23, wherein the codon having the first sequence or the PTC comprises a UAA, UGA or UAA mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid which maintains a property, e.g., function, of a polypeptide encoded by the ORF into which the amino acid is incorporated.

26. The method or composition for use of one of claims 1-23, wherein the codon having the first sequence or the PTC comprises a UAA, UGA or UAA mutation and the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid which does not alter, e.g., maintains, a production parameter, e.g., an expression parameter and/or a signaling parameter, of an mRNA corresponding to the ORF or a polypeptide encoded by the ORF.

27. The method or composition for use of claim 26, wherein the production parameter is compared to an mRNA corresponding to, or a polypeptide encoded by, an otherwise similar ORF having a pre-mutation, e.g., wildtype, amino acid incorporated at the position corresponding to the first sequence codon or PTC.

28. The method or composition for use of claim 26 or 27, wherein the production parameter comprises an expression parameter.

29. The method or composition for use of claim 28, wherein the expression parameter comprises:

- (a) protein translation;
- (b) expression level (*e.g.*, of polypeptide or protein, or mRNA);
- (c) post-translational modification of polypeptide or protein;
- (d) folding (*e.g.*, of polypeptide or protein, or mRNA),
- (e) structure (*e.g.*, of polypeptide or protein, or mRNA),
- (f) transduction (*e.g.*, of polypeptide or protein),
- (g) compartmentalization (*e.g.*, of polypeptide or protein, or mRNA),
- (h) incorporation (*e.g.*, of polypeptide or protein, or mRNA) into a supermolecular structure, *e.g.*, incorporation into a membrane, proteasome, or ribosome,
- (i) incorporation into a multimeric polypeptide, *e.g.*, a homo or heterodimer, and/or
- (j) stability.

30. The method or composition for use of claim 26 or 27, wherein the production parameter comprises a signaling parameter.

31. The method or composition for use of claim 30, wherein the signaling parameter comprises:

- (1) modulation of a signaling pathway, *e.g.*, a cellular signaling pathway which is downstream or upstream of the protein encoded by the endogenous ORF comprising the first sequence or PTC;
- (2) cell fate modulation;
- (3) ribosome occupancy modulation;
- (4) protein translation modulation;
- (5) mRNA stability modulation;
- (6) protein folding and structure modulation;
- (7) protein transduction or compartmentalization modulation; and/or
- (8) protein stability modulation.

32. The method or composition for use of any one of claims 26-31, wherein the production parameter (*e.g.*, an expression parameter and/or a signaling parameter) may be modulated (*e.g.*, increased), *e.g.*, by at least 5% (*e.g.*, at least 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 150%, 200% or more), *e.g.*, compared to a reference sequence.
33. The method or composition for use of any one of the preceding claims, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of any one of the twenty amino acids listed in Table 8.
34. The method or composition for use of any one of the preceding claims, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid corresponding to a non-mutated codon, *e.g.*, a wildtype codon sequence of the codon having the first sequence or the PTC.
35. The method or composition for use of any one of the preceding claims, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of a pre-mutation, *e.g.*, wildtype amino acid.
36. The method or composition for use of claim 35, wherein the TREM, TREM core fragment or TREM fragment mediates incorporation of an amino acid having a similar property as the pre-mutation, *e.g.*, wildtype amino acid, *e.g.*, an amino acid that belongs to the same group as the pre-mutation amino acid, *e.g.*, as provided in Table 2.
37. The method or composition for use of any of the preceding claims, wherein incorporation of the amino acid by the TREM, TREM fragment or TREM core fragment results in modulation, *e.g.*, increase, of a production parameter, *e.g.*, an expression parameter and/or a signaling parameter, of an mRNA corresponding to the ORF or a polypeptide encoded by the ORF.
38. The method or composition for use of claim 37, wherein the production parameter comprises an expression parameter.

39. The method or composition for use of claim 38, wherein the expression parameter comprises:

- (a) protein translation;
- (b) expression level (*e.g.*, of polypeptide or protein, or mRNA);
- (c) post-translational modification of polypeptide or protein;
- (d) folding (*e.g.*, of polypeptide or protein, or mRNA),
- (e) structure (*e.g.*, of polypeptide or protein, or mRNA),
- (f) transduction (*e.g.*, of polypeptide or protein),
- (g) compartmentalization (*e.g.*, of polypeptide or protein, or mRNA),
- (h) incorporation (*e.g.*, of polypeptide or protein, or mRNA) into a supermolecular structure, *e.g.*, incorporation into a membrane, proteasome, or ribosome,
- (i) incorporation into a multimeric polypeptide, *e.g.*, a homo or heterodimer, and/or
- (j) stability.

40. The method or composition for use of claim 37, wherein the production parameter comprises a signaling parameter.

41. The method or composition for use of claim 40, wherein the signaling parameter comprises:

- (1) modulation of a signaling pathway, *e.g.*, a cellular signaling pathway which is downstream or upstream of the protein encoded by the endogenous ORF comprising the first sequence or PTC;
- (2) cell fate modulation;
- (3) ribosome occupancy modulation;
- (4) protein translation modulation;
- (5) mRNA stability modulation;
- (6) protein folding and structure modulation;
- (7) protein transduction or compartmentalization modulation; and/or
- (8) protein stability modulation.

42. The method or composition for use of any one of claims 37-41, wherein the production parameter (*e.g.*, an expression parameter and/or a signaling parameter) may be modulated (*e.g.*,

increased), *e.g.*, by at least 5% (*e.g.*, at least 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 150%, 200% or more), *e.g.*, compared to a reference sequence.

43. The method or composition for use of any one of the preceding claims, wherein the subject has or has been identified as having a disorder or disease listed in any one of Tables 4, 5, and 6.

44. The method or composition for use of any one of the preceding claims, wherein the cell is associated with, *e.g.*, obtained from a subject who has, a disorder or disease listed in any one of Tables 4, 5, and 6.

45. The method or composition for use of claim 43 or 44, wherein the disorder or disease is chosen from the left column of Table 4.

46. The method or composition for use of claim 43 or 44, wherein the disorder or disease is chosen from the left column of Table 4 and the codon having the first sequence or PTC is in a gene chosen from the right column of Table 4, optionally wherein the codon having the first sequence or PTC is at a position provided in Table 4.

47. The method or composition for use of any one of the preceding claims, wherein the codon having the first sequence or PTC is in a gene chosen from the right column of Table 4, optionally wherein the codon having the first sequence or PTC is at a position provided in Table 4.

48. The method or composition for use of claim 43 or 44, wherein the disorder or symptom is chosen from a disorder or disease provided in Table 5.

49. The method or composition for use of claim 43 or 44, wherein the disorder or symptom is chosen from a disorder or disease provided in Table 6, optionally wherein the codon having the first sequence or PTC is in any gene provided in Table 6.

50. The method or composition for use of claim 43 or 44, wherein the disorder or symptom is chosen from a disorder or disease provided in Table 6 and the codon having the first sequence or

PTC is in a corresponding gene provided in Table 6, e.g., a gene corresponding to the disease or disorder.

51. The method or composition for use of claim 43 or 44, wherein the disorder or symptom is chosen from a disorder or disease provided in Table 6 and the codon having the first sequence or PTC is not in a gene provided in Table 6.

52. The method or composition for use of any one of the preceding claims, wherein the codon having the first sequence or PTC is in a gene provided in Table 3.

53. The method or composition for use of any one of the preceding claims, wherein the codon having the first sequence or PTC is at any position within the ORF of the gene, e.g., upstream of the naturally occurring stop codon.

54. The method or composition for use of any one of the preceding claims, wherein the TREM comprises a sequence of Formula A:

[L1]-[ASt Domain1]-[L2]-[DH Domain]-[L3]-[ACH Domain] -[VL Domain]-[TH Domain]-[L4]-[ASt Domain2],

wherein:

independently, [L1] and [VL Domain], are optional;

one of [L1], [ASt Domain1], [L2]-[DH Domain], [L3], [ACH Domain], [VL Domain], [TH Domain], [L4], and [ASt Domain2] comprises a nucleotide having a non-naturally occurring modification; and

wherein:

- (a) the TREM retains the ability to: support protein synthesis, be charged by a synthetase, be bound by an elongation factor, introduce an amino acid into a peptide chain, support elongation, or support initiation;
- (b) the TREM comprises at least X contiguous nucleotides without a non-naturally occurring modification, wherein X is greater than 10;
- (c) at least 3, but less than all of the nucleotides of a type (e.g., A, T, C, G or U) comprise the same non-naturally occurring modification;

- (d) at least X nucleotides of a type (e.g., A, T, C, G or U) do not comprise a non-naturally occurring modification, wherein X=1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 or 50;
- (e) no more than 5, 10, or 15 nucleotides of a type (e.g., A, T, C, G or U) comprise a non-naturally occurring modification; and/or
- (f) no more than 5, 10, or 15 nucleotides of a type (e.g., A, T, C, G or U) do not comprise a non-naturally occurring modification.

55. The method or composition for use of claim 53, wherein the Domain comprising the non-naturally occurring modification retains a function, e.g., a domain function described herein.

56. The method or composition for use of any one of claims 1-53, wherein the TREM core fragment comprises a sequence of Formula B:

[L1]_y-[ASt Domain1]_x-[L2]_y-[DH Domain]_y-[L3]_y-[ACH Domain]_x-[VL Domain]_y-[TH Domain]_y-[L4]_y-[ASt Domain2]_x,

wherein:

x=1 and y=0 or 1;

one of [ASt Domain1], [ACH Domain], and [ASt Domain2] comprises a nucleotide having a non-naturally occurring modification; and

the TREM retains the ability to: support protein synthesis; be able to be charged by a synthetase, be bound by an elongation factor, introduce an amino acid into a peptide chain, support elongation, or support initiation.

57. The method or composition for use of any one of the claims 1-53, wherein the TREM fragment comprises a portion of a TREM, wherein the TREM comprises a sequence of Formula A:

[L1]-[ASt Domain1]-[L2]-[DH Domain]-[L3]-[ACH Domain]-[VL Domain]-[TH Domain]-[L4]-[ASt Domain2], and wherein:

the TREM fragment comprises:

a non-naturally occurring modification; and

one, two, three or all or any combination of the following:

- (a) a TREM half (*e.g.*, from a cleavage in the ACH Domain, *e.g.*, in the anticodon sequence, *e.g.*, a 5' half or a 3' half);
- (b) a 5' fragment (*e.g.*, a fragment comprising the 5' end, *e.g.*, from a cleavage in a DH Domain or the ACH Domain);
- (c) a 3' fragment (*e.g.*, a fragment comprising the 3' end, *e.g.*, from a cleavage in the TH Domain); or
- (d) an internal fragment (*e.g.*, from a cleavage in any one of the ACH Domain, DH Domain or TH Domain).

58. The method or composition for use of any one of claims 54-57, wherein the TREM Domain comprises a plurality of nucleotides each having a non-naturally occurring modification.

59. The method or composition for use of any one of claims 54-58, wherein the non-naturally occurring modification is a modification in a base or a backbone of a nucleotide, *e.g.*, a modification chosen from any one of Tables 5, 6, 7, 8 or 9.

60. The method or composition for use of any one of claims 54-59, wherein the modification comprises one or more of a 2'-O-methyl, 2-deoxy, 2'-fluoro, 2'-O-MOE, pseudouridine or 5,6 dihydrouridine modification.

61. The method or composition for use of any one of claims 54-60, wherein the TREM, TREM core fragment or TREM fragment recognizes a codon provided in Table 7 or Table 8.

62. The method or composition for use of any one of claims 54-61, wherein the TREM, TREM core fragment or TREM fragment is a cognate TREM.

63. The method or composition for use of any one of claims 54-61, wherein the TREM, TREM core fragment or TREM fragment is a non-cognate TREM.

64. The method or composition for use of any one of claims 54-63, wherein the TREM, TREM core fragment or TREM fragment is encoded by a sequence provided in Table 9, e.g., any one of SEQ ID NOs 1-451.

65. The method or composition for use of any one of claims 54-63, wherein the TREM, TREM core fragment or TREM fragment is encoded by a consensus sequence chosen from any one of SEQ ID NOs: 562-621.

66. A pharmaceutical composition comprising a TREM, TREM core fragment or TREM fragment of any one of claims 1-65.

67. A method of making a TREM, TREM core fragment or TREM fragment, comprising linking a first nucleotide to a second nucleotide to form the TREM.

68. The method of claim 67, wherein the TREM, TREM core fragment or TREM fragment is synthetic.

69. The method of claim 68, wherein the TREM, TREM core fragment or TREM fragment is made by cell-free solid phase synthesis.