

Anti-HLA-G antibodies and use thereof

The present invention relates to anti-HLA-G antibodies, their preparation, formulations and methods of using the same.

5 Background of the Invention

The human major histocompatibility complex, class I, 6, also known as human leukocyte antigen G (HLA-G), is a protein that in humans is encoded by the HLA-G gene. HLA-G belongs to the HLA nonclassical class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane but can also be
10 shedded/secreted.

- The heavy chain consists of three domains: alpha 1, alpha 2 and alpha 3. The alpha 1 and alpha 2 domains form a peptide binding groove flanked by two alpha helices. Small peptides (approximately 9-mers) can bind to this groove
15 akin to other MHC I proteins.
- The second chain is beta 2 microglobulin which binds to the heavy chain similar to other MHC I proteins.

For HLA-G there exist 7 isoforms, 3 secreted and 4 membrane bound forms (as schematically shown in Fig.1).

20 HLA-G can form functionally active complex oligomeric structures (Kuroki, K et al. Eur J Immunol. 37 (2007) 1727–1729). Disulfide-linked dimers are formed between Cys 42 of two HLA-G molecules. (Shiroishi M et al., J Biol Chem 281 (2006) 10439-10447. Trimers and Tetrameric complexes have also been described e.g. in Kuroki, K et al. Eur J Immunol. 37 (2007) 1727–1729, Allan D.S., et al. J Immunol Methods.
25 268 (2002) 43-50 and T Gonen-Gross et al., J Immunol 171 (2003)1343-1351).

HLA-G is predominantly expressed on cytotrophoblasts in the placenta. Several tumors (including pancreatic, breast, skin, colorectal, gastric & ovarian) express HLA-G (Lin, A. et al., Mol Med. 21 (2015) 782–791; Amiot, L., et al., Cell Mol Life Sci. 68 (2011) 417–431). The expression has also been reported to be associated with
30 pathological conditions like inflammatory diseases, GvHD and cancer. Expression of HLA-G has been reported to be associated with poor prognosis in cancer. Tumor cells escape host immune surveillance by inducing immune tolerance/suppression via HLA-G expression.

Overview polymorphisms HLA family		
• HLA-A:	2579 seqs	} <u>classical</u> class I MHC
• HLA-B:	3283 seqs	
• HLA-C:	2133 seqs	
• HLA-E:	15 seqs	} <u>non-classical</u> class I MHC
• HLA-F:	22 seqs	
• HLA-G:	50 seqs	

HLA-G shares high homology (>98%) with other MHC I molecules, therefore truly HLA-G specific antibodies with no crossreactivity to other MHC I molecules are difficult to generate.

- 5 Certain antibodies which interact in different ways with HLA-G were described previously: Tissue Antigens, 55 (2000) 510-518 relates to monoclonal antibodies e.g. 87G, and MEM-G/9; Neoplasma 50 (2003) 331-338 relates to certain monoclonal antibodies recognizing both, intact HLA-G oligomeric complex (e.g. 87G and MEM-G9) as well as HLA-G free heavy chain (e.g. 4H84, MEM-G/1 and MEM-G/2); Hum
- 10 Immunol. 64 (2003) 315-326 relates to several antibodies tested on HLA-G expressing JEG3 tumor cells (e.g. MEM-G/09 and -G/13 which react exclusively with native HLA-G1 molecules. MEM-G/01 recognizes (similar to the 4H84 mAb) the denatured HLA-G heavy chain of all isoforms, whereas MEM-G/04 recognizes selectively denatured HLA-G1, -G2, and -G5 isoforms; Wiendl et al Brain 2003 176-
- 15 85 relates to different monoclonal HLA-G antibodies as e.g. 87G, 4H84, MEM-G/9.

The above publications report antibodies, which bind to human HLA-G or the human HLA-G- β 2M MHC complex. However, due to the high polymorphism and high homology of the HLA family most of the antibodies lack either truly specific HLA-G binding properties and often also bind or crossreact with other HLA family

20 members (either as MHC complex with β 2M or in its β 2M-free form) or they simply do not inhibit binding of HLA-G β 2M MHC complex to its receptors ILT2 and/or ILT4 (and are regarded as non-antagonistic antibodies).

WO2019/202040 relates to HLA-G antibodies including antibody HLA-G-0090. WO 2019/202041 relates to multispecific HLA-G antibodies including antibody

25 HLA-G-0090.

Summary of the Invention

The invention described herein provides an antibody that binds to human HLA-G comprising

5 A) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:23; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID
10 NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6, or

B) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:25; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID
15 NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6.

One embodiment of the invention is an antibody that binds to human HLA-G wherein the antibody

A) comprises a VH domain comprising the amino acid sequence of SEQ ID
20 NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24; or

B) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26.

One embodiment of the invention is an antibody that binds to human HLA-G wherein the antibody comprises a VH domain comprising the amino acid sequence of SEQ
25 ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24.

One embodiment of the invention is an antibody that binds to human HLA-G wherein the antibody comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26.

Such antibodies have highly valuable properties like their binding properties, their
30 high specificity towards HLA-G with no crossreactivity to HLA-A and HLA MHC I complexes from other species. They can bind to HLA-G on cells and inhibit ILT2

and/or ILT4 binding to HLA-G expressed on these cells. They have been generated from the HLA-G antibody HLA-G-0090.

As the HLA-G antibody HLA-G-0090 described in WO 2019/202040 comprises a glycosylation site in one of the CDRs (CDR-L1 which comprises a NSS motif at amino acids 31, 32 and 33 of the light chain (LC)), its binding properties are impacted by the N-glycosylation which constitutes a potential developability liability.

A homology model of the variable region of HLA-G-0090 indicated that light chain (LC) positions 31 to 33 are highly solvent accessible. Furthermore, the side chains of N31 and S32 are predicted to point inwards, in the direction of CDR-H3, making them likely candidates for being part of the antibody paratope. In fact, a number of published antibody-antigen X-ray complex structures document these residues to be undergoing chemical interactions with the antigen. Therefore, the risk of worsening the binding affinity of the antibody by introducing mutations at LC positions 31-33 was high. Therefore various variants of antibody HLA-G-0090 with mutations on LC positions 31, 32, and 33 were designed from which however most variants worsened binding properties or expressability. Surprisingly it was found that among these various variants of HLA-G-0090 in which the glycosylation site was removed only the two variants HLA-G-0090-VL-S32P and HLA-G-0090-VL-S33A show even improved binding properties, good expressability and stability, while showing no more N-glycosylation at the CDR-L1 of the LC (so no Fab glycosylation could be detected). As all recently approved pharmaceutical antibody products are produced in mammalian cells, especially CHO cells (see e.g. Walsh G., Nature Biotech (2018) 1136-1145), providing an antibody without glycosylation sites in the binding region (VH and VL and especially the CDRs) represents a valuable advantage, as these antibodies can readily be used for production in mammalian expression systems without the risk of (at least partially) impairing the binding properties by glycosylation.

In a further embodiment the HLA-G antibody of the present invention comprises a Fc domain of human origin. In one embodiment the Fc domain is of the IgG isotype, in one preferred embodiment of the IgG1 isotype.

In a further embodiment the HLA-G antibody of the present invention is a bispecific antibody, in particular an a bispecific antibody that binds to human HLA-G and to human CD3, comprising a first antigen binding moiety that binds to human HLA-G and a second antigen binding moiety that binds to human CD3.

- 5 -

In one embodiment the HLA-G antibody of the present invention is a bispecific antibody that binds to human HLA-G and to human CD3, comprising a first antigen binding moiety that binds to human HLA-G and a second antigen binding moiety that binds to human CD3,

5 wherein the first antigen binding moiety that binds to human HLA-G comprises

A) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:23; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6, or

B) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:25; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6;

and wherein the second antigen binding moiety that binds to a T cell activating antigen binds to human CD3 comprises

C) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:52, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:53, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:54; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:55; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:56 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:57, or

D) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:60, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:61, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:62; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:63; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:64 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:65, or

- 6 -

- 5 E) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:68, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:69, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:70; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:71; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:72 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:73.

One embodiment of the invention is such bispecific antibody,

wherein the first antigen binding moiety

- 10 A) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24; or

B) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26,

and wherein the second antigen binding moiety

- 15 C) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:58 and a VL domain comprising the amino acid sequence of SEQ ID NO:59; or

D) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:66 and a VL domain comprising the amino acid sequence of SEQ ID NO:67; or

- 20 E) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:74 and a VL domain comprising the amino acid sequence of SEQ ID NO:75.

One embodiment of the invention is such bispecific antibody,

wherein the first antigen binding moiety

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24;

- 25 and wherein the second antigen binding moiety

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:58 and a VL domain comprising the amino acid sequence of SEQ ID NO:59.

One embodiment of the invention is such bispecific antibody,

- 7 -

wherein the first antigen binding moiety

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and
a VL domain comprising the amino acid sequence of SEQ ID NO:24;

and wherein the second antigen binding moiety

- 5 comprises a VH domain comprising the amino acid sequence of SEQ ID NO:66 and
 a VL domain comprising the amino acid sequence of SEQ ID NO:67.

One embodiment of the invention is such bispecific antibody,

wherein the first antigen binding moiety

- 10 comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and
 a VL domain comprising the amino acid sequence of SEQ ID NO:24;

and wherein the second antigen binding moiety

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:74 and
a VL domain comprising the amino acid sequence of SEQ ID NO:75.

- 15 Such bispecific antibodies binding to human HLA-G and human CD3 show in
addition to the properties of the HLA-G-antibodies further valuable properties like
the induction of antibody mediated IFN gamma secretion by T cells on HLA-G
expressing cells, T cell activation in the presence of HLA-G expressing tumor cells,
induction of T cell mediated tumor cell killing on HLA-G expressing cells and in
vivo anti-tumor efficacy and even tumor regression in different cancer xenograft
20 mouse models.

The invention provides an isolated nucleic acid encoding the antibody or bispecific
antibody as described herein.

The invention provides a host cell comprising such nucleic acid.

- 25 The invention provides a method of producing an antibody comprising culturing the
host cell so that the antibody is produced.

The invention provides such method of producing an antibody, further comprising
recovering the antibody from the host cell.

The invention provides an antibody produced by such an host cell where the host cell is eukaryotic.

The invention provides a pharmaceutical formulation comprising the antibody described herein and a pharmaceutically acceptable carrier.

5 The invention provides the antibody described herein for use as a medicament.

The invention provides the antibody described herein for use in treating cancer.

The invention provides the use of the antibody described herein in the manufacture of a medicament. In one embodiment the medicament is for treatment of cancer.

10 The invention provides a method of treating an individual having cancer comprising administering to the individual an effective amount of the antibody described herein.

Description of the Figures

Figure 1: Different isoforms of HLA-G

Figure 2: **Fig. 2A:** Schematic representation of the HLA-G molecule in association with β 2M

15 **Fig. 2B:** Structure of HLA-G molecule in association with certain receptors : HLA-G structure in complex with given receptors such as ILT4 and KIR2DL1. ILT4 structure (PDB code: 2DYP). The KIR2DL1 structure is taken from PDB code 1IM9 (KIR2DL1: HLA-Cw4 complex structure) and was positioned on HLA-G by superposition of the HLA-Cw4 and HLA-G structures. Receptors are shown in a ribbon representation, HLA-G is shown in a molecular surface representation. HLA-G residues that are unique or conserved in other HLA paralogs are colored in white and gray, respectively. Unique surface residues were replaced by a HLA consensus sequence in the chimeric counter antigen.

20

25

Figure 3: Schematic antibody-antigen binding assay principle - relative active concentration (RAC) of HLA-G antibodies for binding to HLA-G

Figure 4A: Mass spectrum of N-glycosylation of HLA-G-0090 indicates Fab-glycosylation

- Figure 4B:** Mass spectra of N-glycosylation of HLA-G-0090-VL-S32P and HLA-G-0090-VL-S33A: No Fab-glycosylation detectable
- Figure 5:** exemplary FACS staining for anti-HLA-G antibodies HLA-G-0090, HLA-G-0090-VL-S32P and HLA-G-0090-VL-S33A (2µg/ml) on SKOV3 cells (no HLA-G expression, JEG3 cells (expressing HLA-G) and SKOV3-HLA-G cells (SKOV3 cells transfected with HLA-G)
- Figure 6:** Ability of the HLA-G antibodies HLA-G-0090, HLA-G-0090-VL-S32P and HLA-G-0090-VL-S33A to modify/inhibit the interaction and binding of recombinant soluble ILT2 (ILT2Fc domain fusion) to HLA-G naturally expressed on JEG3 tumor cells. JEG3 cells are preincubated/pretreated with HLA-G antibodies so that ILT2 binding to JEG3 cells is inhibited/blocked. Controls were carried out with JEG3 cells without HLA-G antibody pretreatment (only ILT2-Fc) and isotype antibody.
- Figure 7:** Binding of bispecific anti-HLA-G/anti-CD3 T cell bispecific (TCB) antibody to CD3 expressed on T-cells by antibodies P1AF7977, P1AF7978 and P1AF7979
- Figure 8:** Bispecific anti-HLA-G/anti-CD3 T cell bispecific (TCB) antibodies P1AF7977, P1AF7978 and P1AF7979 showed binding to JEG3 cells and SKOV3 cells, transfected with HLA-G.
- Figure 9:** Bispecific anti-HLA-G/anti-CD3 T cell bispecific (TCB) antibody mediated/induced IFN gamma secretion by T cells, by antibodies P1AF7977, P1AF7978 and P1AF7979
- Figure 10:** Bispecific anti-HLA-G/anti-CD3 T cell bispecific (TCB) antibodies P1AF7977, P1AF7978 and P1AF7979 induced T cell mediated cytotoxicity/tumor cell killing.
- Figure 11:** In vivo anti-tumor efficacy of anti-HLA-G/anti-CD3 T cell bispecific (TCB) antibody P1AF7977 in humanized NSG mice bearing SKOV3 human ovarian carcinoma transfected with recombinant HLA-G (SKOV3 HLA-G), leading to tumor regression

- 5 **Figure 12:** Dose-response study with anti-HLA-G/anti-CD3 T cell bispecific (TCB) antibody P1AF7977 in humanized NSG mice bearing human breast cancer PDX tumors (BC004). Strong tumor growth inhibition until tumor regression is observed in mice treated with different doses.
- 10 **Figure 13:** Exemplary configurations of the bispecific antigen binding molecules of the invention. (A, D) Illustration of the “1+1 CrossMab” molecule. (B, E) Illustration of the “2+1 IgG Crossfab” molecule with alternative order of Crossfab and Fab components (“inverted”). (C, F) Illustration of the “2+1 IgG Crossfab” molecule. (G, K) Illustration of the “1+1 IgG Crossfab” molecule with alternative order of Crossfab and Fab components (“inverted”). (H, L) Illustration of the “1+1 IgG Crossfab” molecule. (I, M) Illustration of the “2+1 IgG Crossfab” molecule with two CrossFabs. (J, N). Black dot: optional modification in the Fc domain promoting heterodimerization. ++, --: amino acids of opposite charges optionally introduced in the CH1 and CL domains. Crossfab molecules are depicted as comprising an exchange of VH and VL regions, but may – in embodiments wherein no charge modifications are introduced in CH1 and CL domains – alternatively comprise an exchange of the CH1 and CL domains.
- 15 20
- Figure 14:** Induction of T cell activation by bispecific anti-HLA-G/anti-CD3 antibody P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035 in the presence of SKOV3 HLAG cells.
- 25 **Figure 15:** Relative binding activity of original and optimized CD3 binders, CD3_{orig} and CD3_{opt} (=P035-093 (P035)), to recombinant CD3 as measured by SPR in unstressed condition, after 14 d at 40°C pH 6, or after 14 d at 37°C pH 7.4 (IgG format).
- 30 **Figure 16:** Binding of original and optimized CD3 binders, CD3_{orig} and CD3_{opt}(=P035-093 (P035)), to Jurkat NFAT cells as measured by flow cytometry (IgG format). Antibodies bound to Jurkat NFAT cells were detected with a fluorescently labeled anti-human Fc specific secondary antibody.
- Figure 17:** Schematic illustration of the CD3 activation assay used in Example 8.

Figure 18: Jurkat NFAT activation with original and optimized CD3 binders, CD3_{orig} and CD3_{opt} (=P035-093 (P035)) (IgG format). Jurkat NFAT reporter cells were co-incubated with anti-PGLALA expressing CHO (CHO-PGLALA) cells in the presence of CD3_{orig} or CD3_{opt} (=P035-093 (P035)) IgG PGLALA, or CD3_{opt} IgG wt as negative control. CD3 activation was quantified by measuring luminescence after 24 h.

Detailed Description of the Invention

When used herein, the term “HLA-G”, “human HLA-G”, “HLAG”, refers to the HLA-G human major histocompatibility complex, class I, G, also known as human leukocyte antigen G (HLA-G) (exemplary SEQ ID NO: 35). Typically, HLA-G forms a MHC class I complex together with $\beta 2$ microglobulin (B2M or $\beta 2m$). In one embodiment HLA-G refers to the MHC class I complex of HLA-G and $\beta 2$ microglobulin. In one preferred embodiment HLA-G refers to the cell surface bound MHC class I complex of HLA-G and $\beta 2$ microglobulin, also known as HLA-G1 (see Figure 1 of this description and e.g. Blaschitz et al., Molecular Human Reproduction, 11(2005) 699-710, inter alia Figure 1)

As used herein, an antibody (either mono-, multi- or bispecific) or antigen binding moiety “binding to human HLA-G”, “specifically binding to human HLA-G”, “that binds to human HLA-G” or “anti-HLA-G” refers to an antibody/ antigen binding moiety specifically binding to the human HLA-G antigen or its extracellular domain (ECD) with a binding affinity of a K_D -value of 5.0×10^{-8} mol/l or lower, in one embodiment of a K_D -value of 1.0×10^{-9} mol/l or lower, in one embodiment of a K_D -value of 5.0×10^{-8} mol/l to 1.0×10^{-13} mol/l. In one embodiment the antibody binds to HLA-G $\beta 2m$ MHC I complex comprising SEQ ID NO: 39)

The binding affinity is determined with a standard binding assay, such as surface plasmon resonance technique (BIAcore®, GE-Healthcare Uppsala, Sweden) e.g. using constructs comprising HLA-G extracellular domain (e.g. in its natural occurring 3 dimensional structure). In one embodiment binding affinity is determined with a standard binding assay using exemplary soluble HLA-G comprising MHC class I complex comprising SEQ ID NO: 39.

HLA-G has the regular MHC I fold and consists of two chains: Chain 1 consists of three domains: alpha 1, alpha 2 and alpha 3. The alpha 1 and alpha 2 domains form

a peptide binding groove flanked by two alpha helices. Small peptides (approximately 9mers) can bind to this groove akin to other MHCI proteins. Chain 2 is beta 2 microglobulin ($\beta 2M$) which is shared with various other MHCI proteins.

5 HLA-G can form functionally active complex oligomeric structures (Kuroki, K et al. Eur J Immunol. 37 (2007) 1727–1729). Disulfide-linked dimers are formed between Cys 42 of two HLA-G molecules. (Shiroishi M et al., J Biol Chem 281 (2006) 10439-10447. Trimers and Tetrameric complexes have also been described e.g. in Kuroki, K et al. Eur J Immunol. 37 (2007) 1727–1729, Allan D.S., et al. J Immunol Methods. 268 (2002) 43-50 and T Gonen-Gross et al., J Immunol 171 (2003)1343-1351).
10 HLA-G has several free cysteine residues, unlike most of the other MHC class I molecules. Boyson et al., Proc Nat Acad Sci USA, 99: 16180 (2002) reported that the recombinant soluble form of HLA-G5 could form a disulfide-linked dimer with the intermolecular Cys42-Cys42 disulfide bond. In addition, the membrane-bound form of HLA-G1 can also form a disulfide-linked dimer on the cell surface of the
15 JEG3 cell line, which endogenously expresses HLA-G. Disulfide-linked dimer forms of HLA-G1 and HLA-G5 have been found on the cell surface of trophoblast cells as well (Apps, R., Tissue Antigens, 68:359 (2006)).

HLA-G is predominantly expressed on cytotrophoblasts in the placenta. Several tumors (including pancreatic, breast, skin, colorectal, gastric & ovarian) express
20 HLA-G (Lin, A. et al., Mol Med. 21 (2015) 782–791; Amiot, L., et al., Cell Mol Life Sci. 68 (2011) 417–431). The expression has also been reported to be associated with pathological conditions like inflammatory diseases, GvHD and cancer. Expression of HLA-G has been reported to be associated with poor prognosis in cancer. Tumor cells escape host immune surveillance by inducing immune tolerance/suppression
25 via HLA-G expression.

For HLA-G there exist 7 isoforms, 3 secreted and 4 membrane bound forms (as schematically shown in Fig.1). The most important functional isoforms of HLA-G include $\beta 2$ -microglobulin ($\beta 2M$)-associated HLA-G1 and HLA-G5. However, the tolerogenic immunological effect of these isoforms is different and is dependent on
30 the form (monomer, dimer) of ligands and the affinity of the ligand-receptor interaction.

HLA-G protein can be produced using standard molecular biology techniques. The nucleic acid sequence for HLA-G isoforms is known in the art. See for example GENBANK Accession No. AY359818.

The HLA-G isomeric forms promote signal transduction through ILTs, in particular ILT2, ILT4, or a combination thereof.

5 ILTs: ILTs represent Ig types of activating and inhibitory receptors that are involved in regulation of immune cell activation and control the function of immune cells (Borges, L., et al., Curr Top Microbial Immunol, 244:123-136 (1999)). ILTs are categorized into three groups: (i) inhibitory, those containing a cytoplasmic immunoreceptor tyrosine-based inhibitory motif (ITIM) and transducing an inhibitory signal (ILT2, ILT3, ILT4, ILT5, and LIR8); (ii) activating, those containing a short cytoplasmic tail and a charged amino acid residue in the transmembrane domain (ILT1, ILT7, ILT8, and LIR6alpha) and delivering an activating signal through the cytoplasmic immunoreceptor tyrosine-based activating motif (ITAM) of the associated common gamma chain of Fc receptor; and (iii) the soluble molecule ILT6 lacking the transmembrane domain. A number of recent studies have highlighted immunoregulatory roles for ILTs on the surface of antigen presenting cells (APC). ILT2, ILT3, and ILT4 receptors, the most characterized immune inhibitory receptors, are expressed on a wide range of immune cells including monocytes, B cells, dendritic cells, plasmacytoid dendritic cells and a subset of NK and T cells. ILT2 is expressed on T cells subsets has been shown to inhibit activation and proliferation of these cells upon ligation (Colonna M. et al., J Immunol. 20011, 66:2514-2521, J Immunol 2000; 165:3742-3755). ILT3 and ILT4 are upregulated by exposing immature DC to known immunosuppressive factors, including IL-10, vitamin D3, or suppressor CD8 T cells (Chang, C. C., et al., Nat Immunol, 3:237-243 (2002)). The expression of ILTs on DC is tightly controlled by inflammatory stimuli, cytokines, and growth factors, and is down-regulated following DC activation (Ju, X. S., et al., Gene, 331:159-164 (2004)). The expression of ILT2 and ILT4 receptors is highly regulated by histone acetylation, which contributes to strictly controlled gene expression exclusively in the myeloid lineage of cells (Nakajima, H., J Immunol, 171:6611-6620 (2003)).

30 Engagement of the inhibitory receptors ILT2 and ILT4 alters the cytokine and chemokine secretion/release profile of monocytes and can inhibit Fc receptor signaling (Colonna, M., et al. J Leukoc Biol, 66:375-381 (1999)). The role and function of ILT3 on DC have been precisely described by the Suci-Foca group (Suci-Foca, N., Int Immunopharmacol, 5:7-11 (2005)). Although the ligand for ILT3 is unknown, ILT4 is known to bind to the third domain of HLA class I molecules (HLA-A, HLA-B, HLA-C, and HLA-G), competing with CD8 for MHC class I binding (Shiroishi, M., Proc Natl Acad Sci USA, 100:8856-8861 (2003)). The

preferential ligand for several inhibitory ILT receptors is HLA-G. HLA-G plays a potential role in maternal-fetal tolerance and in the mechanisms of escape of tumor cells from immune recognition and destruction (Hunt, J. S., et al., *Faseb J*, 19:681-693 (2005)). It is most likely that regulation of DC function by HLA-G-ILT interactions is an important pathway in the biology of DC. It has been determined that human monocyte-derived DC that highly express ILT2 and ILT4 receptors, when treated with HLA-G and stimulated with allogeneic T cells, still maintain a stable tolerogenic-like phenotype (CD80^{low}, CD86^{low}, HLA-DR^{low}) with the potential to induce T cell anergy (Ristich, V., et al., *Eur J Immunol*, 35:1133-1142 (2005)). Moreover, the HLA-G interaction with DC that highly express ILT2 and ILT4 receptors resulted in down-regulation of several genes involved in the MHC class II presentation pathway. A lysosomal thiol reductase, IFN-gamma inducible lysosomal thiol reductase (GILT), abundantly expressed by professional APC, was greatly reduced in HLA-G-modified DC. The repertoire of primed CD4⁺ T cells can be influenced by DC expression of GILT, as in vivo T cell responses to select antigens were reduced in animals lacking GILT after targeted gene disruption (Marie, M., et al., *Science*, 294:1361-1365 (2001)). The HLA-G/ILT interaction on DC interferes with the assembly and transport of MHC class II molecules to the cell surface, which might result in less efficient presentation or expression of structurally abnormal MHC class II molecules. It was determined that HLA-G markedly decreased the transcription of invariant chain (CD74), HLA-DMA, and HLA-DMB genes on human monocyte-derived DC highly expressing ILT inhibitory receptors (Ristich, V., et al; *Eur J Immunol* 35:1133-1142 (2005)).

Another receptor of HLA-G is KIR2DL4 because KIR2DL4 binds to cells expressing HLA-G (US2003232051; Cantoni, C. et al. *Eur J Immunol* 28 (1998) 1980; Rajagopalan, S. and E. O. Long. [published erratum appears in *J Exp Med* 191 (2000) 2027] *J Exp Med* 189 (1999) 1093; Ponte, M. et al. *PNAS USA* 96 (1999) 5674). KIR2DL4 (also referred to as 2DL4) is a KIR family member (also designated CD158d) that shares structural features with both activating and inhibitory receptors (Selvakumar, A. et al. *Tissue Antigens* 48 (1996) 285). 2DL4 has a cytoplasmic ITIM, suggesting inhibitory function, and a positively charged amino acid in the transmembrane region, a feature typical of activating KIR. Unlike other clonally distributed KIRs, 2DL4 is transcribed by all NK cells (Valiante, N. M. et al. *Immunity* 7 (1997) 739; Cantoni, C. et al. *Eur J Immunol* 28 (1998) 1980; Rajagopalan, S. and E. O. Long. [published erratum appears in *J Exp Med* 191 (2000) 2027] *J Exp Med* 189 (1999) 1093).

HLA-G has also been shown to interact with CD8 (Sanders et al, J. Exp. Med., 174 (1991), 737–740) on cytotoxic T cells and induce CD95 mediated apoptosis in activated CD8 positive cytotoxic T cells (Fournel et al, J. Immun., 164 (2000), 6100-6104). This mechanism of elimination of cytotoxic T cells has been reported to one of the mechanisms of immune escape and induction of tolerance in pregnancy, inflammatory diseases and cancer (Amodio G. et al, Tissue Antigens, 84 (2014), 255-263).

As used herein an anti-HLA-G antibody (either mono-, multi- or bispecific) or antigen binding moiety that “does not crossreact with “or that “does not (specifically) bind to “a modified human HLA-G β 2M MHC I complex, wherein the HLA-G specific amino acids have been replaced by HLA-A consensus amino acids, the complex comprising SEQ ID NO:40; a mouse H2Kd β 2M MHC I complex comprising SEQ ID NO:41 rat RT1A β 2M MHC I complex comprising SEQ ID NO:43, human HLA-A2 β 2M MHC I complex comprising SEQ ID NO:35 and SEQ ID NO: 33 refers to an anti-HLA-G antibody (either mono-, multi- or bispecific) or antigen binding moiety that does substantially not bind to any of these counterantigens. In one embodiment an anti-HLA-G antibody(either mono-, multi- or bispecific) or antigen binding moiety that “does not crossreact with “ or that “does not specifically bind to“ a modified human HLA-G β 2M MHC I complex, wherein the HLA-G specific amino acids have been replaced by HLA-A consensus amino acids, the complex comprising SEQ ID NO:40; a mouse H2Kd β 2M MHC I complex comprising SEQ ID NO:41, a rat RT1A β 2M MHC I complex comprising SEQ ID NO:43, and/or a human HLA-A2 β 2M MHC I complex comprising SEQ ID NO:35 and SEQ ID NO: 33 refers to an anti-HLA-G antibody (either mono-, multi- or bispecific) or antigen binding moiety that shows no significant binding/interaction in e.g. a Surface plasmon resonance assay (as described e.g. in Example 2) The binding/ interaction is determined with a standard binding assay, such as surface plasmon resonance technique (BIAcore®, GE-Healthcare Uppsala, Sweden) with the respective antigen: a modified human HLA-G β 2M MHC I complex, wherein the HLA-G specific amino acids have been replaced by HLA-A consensus amino acids, the complex comprising SEQ ID NO:40; a mouse H2Kd β 2M MHC I complex comprising SEQ ID NO:41 rat RT1A β 2M MHC I complex comprising SEQ ID NO:43, and/or a human HLA-A2 β 2M MHC I complex comprising SEQ ID NO:35 and SEQ ID NO: 33 The assay setup as well as the construction/preparation of the antigens is described in the Examples.

The term “inhibits ILT2 binding to HLA-G on JEG-3 cells (ATCC HTB36)” refers to the inhibition of binding interaction of (recombinant) ILT2 e.g in an assay as described in Example 5.

5 An “activating T cell antigen” as used herein refers to an antigenic determinant expressed on the surface of a T lymphocyte, particularly a cytotoxic T lymphocyte, which is capable of inducing T cell activation upon interaction with an antibody. Specifically, interaction of an antibody with an activating T cell antigen may induce T cell activation by triggering the signaling cascade of the T cell receptor complex. In a particular embodiment the activating T cell antigen is CD3, particularly the
10 epsilon subunit of CD3 (see UniProt no. P07766 (version 189), NCBI RefSeq no. NP_000724.1, SEQ ID NO: 88 for the human sequence; or UniProt no. Q95LI5 (version 49), NCBI GenBank no. BAB71849.1, SEQ ID NO: 108 for the cynomolgus [Macaca fascicularis] sequence).

15 “CD3” refers to any native CD3 from any vertebrate source, including mammals such as primates (e.g. humans), non-human primates (e.g. cynomolgus monkeys) and rodents (e.g. mice and rats), unless otherwise indicated. CD3 is an exemplary activated T cell antigen. The term “CD3” encompasses “full-length,” unprocessed CD3 as well as any form of CD3 that results from processing in the cell. The term also encompasses naturally occurring variants of CD3, e.g., splice variants or allelic
20 variants. In one embodiment, CD3 is human CD3, particularly the epsilon subunit of human CD3 (CD3ε). The amino acid sequence of human CD3ε is shown in UniProt (www.uniprot.org) accession no. P07766 (version 189), or NCBI (www.ncbi.nlm.nih.gov/) RefSeq NP_000724.1. See also SEQ ID NO: 88. The amino acid sequence of cynomolgus [Macaca fascicularis] CD3ε is shown in NCBI
25 GenBank no. BAB71849.1. See also SEQ ID NO: 89.

As used herein, an antibody (either mono-, multi- or bispecific) or antigen binding moiety “binding to human CD3”, “specifically binding to human CD3”, “that binds to human CD3” or “anti-CD3” refers to an anti-antibody (either mono-, multi- or
30 bispecific) or antigen binding moiety specifically binding to the human CD3 antigen or its extracellular domain (ECD) which shows significant binding/interaction in a surface plasmon resonance assay. In one embodiment with a binding affinity of a K_D -value of 5.0×10^{-8} mol/l or lower, in one embodiment of a K_D -value of 1.0×10^{-9} mol/l or lower, in one embodiment of a K_D -value of 5.0×10^{-8} mol/l to 1.0×10^{-13} mol/l. In one embodiment the antibody binds to CD3 comprising SEQ ID
35 NO: 88.

The binding affinity is determined with a standard binding assay, such as surface plasmon resonance technique (BIAcore®, GE-Healthcare Uppsala, Sweden) e.g. using constructs comprising HLA-G extracellular domain (e.g. in its natural occurring 3 dimensional structure). In one embodiment binding affinity is
5 determined with a standard binding assay using exemplary CD3 comprising SEQ ID NO: 88.

Accordingly a multispecific or bispecific that binds to human HLA-G and to human CD3, comprising a first antigen binding moiety that binds to human HLA-G and a second antigen binding moiety that binds to human CD3 refers to an antibody that
10 binds with an (first) antigen binding moiety to human HLA-G as described herein and that binds with another (second) antigen binding moiety to human CD3 as described herein.

“T cell activation” as used herein refers to one or more cellular response of a T lymphocyte, particularly a cytotoxic T lymphocyte, selected from: proliferation,
15 differentiation, cytokine secretion, cytotoxic effector molecule release, cytotoxic activity, and expression of activation markers. Suitable assays to measure T cell activation are known in the art and described herein.

An “acceptor human framework” for the purposes herein is a framework comprising the amino acid sequence of a light chain variable domain (VL) framework or a heavy chain variable domain (VH) framework derived from a human immunoglobulin
20 framework or a human consensus framework, as defined below. An acceptor human framework “derived from” a human immunoglobulin framework or a human consensus framework may comprise the same amino acid sequence thereof, or it may contain amino acid sequence changes. In some embodiments, the number of amino
25 acid changes are 10 or less, 9 or less, 8 or less, 7 or less, 6 or less, 5 or less, 4 or less, 3 or less, or 2 or less. In some embodiments, the VL acceptor human framework is identical in sequence to the VL human immunoglobulin framework sequence or human consensus framework sequence.

The term "antibody" herein is used in the broadest sense and encompasses various
30 antibody structures, including but not limited to monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments so long as they exhibit the desired antigen-binding activity.

An "antibody fragment" refers to a molecule other than an intact antibody that comprises a portion of an intact antibody that binds the antigen to which the intact

antibody binds. Examples of antibody fragments include but are not limited to Fv, Fab, Fab', Fab'-SH, F(ab')₂, diabodies; linear antibodies; single-chain antibody molecules (e.g. scFv); and multispecific antibodies formed from antibody fragments.

5 The term "bispecific" means that the antibody is able to specifically bind to at least two distinct antigenic determinants. Typically, a bispecific antibody comprises two antigen binding sites or moieties, each of which is specific for a different antigenic determinant. In certain embodiments the bispecific antibody is capable of simultaneously binding two antigenic determinants, particularly two antigenic determinants expressed on two distinct cells.

10 The term "valent" as used herein denotes the presence of a specified number of antigen binding sites in an antibody. As such, the term "monovalent binding to an antigen" denotes the presence of one (and not more than one) antigen binding site specific for the antigen in the antibody.

15 The terms "antigen binding site" and "antigen binding moiety" as used herein are interchangeable and refer to the site, i.e. one or more amino acid residues, of an antibody which provides interaction with the antigen. For example, the antigen binding site of an antibody comprises amino acid residues from the complement determining regions (CDRs). A native immunoglobulin molecule typically has two antigen binding sites, a Fab molecule typically has a single antigen binding site.

20 "Antigen binding site" and "antigen binding moiety" refers to a polypeptide molecule that specifically binds to an antigenic determinant. In one embodiment, an antigen binding moiety is able to direct the entity to which it is attached (e.g. a second antigen binding moiety) to a target site, for example to a specific type of tumor cell bearing the antigenic determinant. In another embodiment an antigen binding moiety

25 is able to activate signaling through its target antigen, for example a T cell receptor complex antigen. Antigen binding moieties include antibodies and fragments thereof as further defined herein. Particular antigen binding moieties include an antigen binding domain of an antibody, comprising an antibody heavy chain variable region and an antibody light chain variable region. In certain embodiments, the antigen

30 binding moieties may comprise antibody constant regions as further defined herein and known in the art. Useful heavy chain constant regions include any of the five isotypes: α , δ , ϵ , γ , or μ . Useful light chain constant regions include any of the two isotypes: κ and λ . In one preferred embodiment such constant regions are of human origin.

As used herein, the term "antigenic determinant" or "antigen" refers to a site on a polypeptide macromolecule to which an antigen binding moiety binds, forming an antigen binding moiety-antigen complex. Useful antigenic determinants can be found, for example, on the surfaces of tumor cells, on the surfaces of virus-infected cells, on the surfaces of other diseased cells, on the surface of immune cells, free in blood serum, and/or in the extracellular matrix (ECM).

The term "chimeric" antibody refers to an antibody in which a portion of the heavy and/or light chain is derived from a particular source or species, while the remainder of the heavy and/or light chain is derived from a different source or species.

The "class" of an antibody refers to the type of constant domain or constant region possessed by its heavy chain. There are five major classes of antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG₁, IgG₂, IgG₃, IgG₄, IgA₁, and IgA₂. The heavy chain constant domains that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. In one preferred embodiment such subclasses (isotypes) are of human origin. In one preferred embodiment the antibodies of the present invention are of the IgG isotype, in another preferred embodiment of the IgG₁ isotype.

In one aspect, the antibody comprises a constant region of human origin. In one aspect, the antibody is an immunoglobulin molecule comprising a human constant region, particularly of the IgG isotype, more particularly of the IgG₁ isotype, comprising a human CH1, CH2, CH3 and/or CL domain. Exemplary sequences of human constant domains are given in SEQ ID Nos: 47 and 48 (human kappa and lambda CL domains, respectively) and SEQ ID NO: 49 (human IgG₁ heavy chain constant domains CH1-CH2-CH3) or SEQ ID NO: 50 (human IgG₁ heavy chain constant region with mutations L234A, L235A and P329G). An "effective amount" of an agent, e.g., a pharmaceutical formulation, refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic or prophylactic result.

The term "Fc domain" or "Fc region" herein is used to define a C-terminal region of an immunoglobulin heavy chain that contains at least a portion of the constant region. The term includes native sequence Fc regions and variant Fc regions. Although the boundaries of the Fc region of an IgG heavy chain might vary slightly, the human IgG heavy chain Fc region is usually defined to extend from Cys226, or from Pro230, to the carboxyl-terminus of the heavy chain. However, antibodies produced by host cells may undergo post-translational cleavage of one or more, particularly one or

two, amino acids from the C-terminus of the heavy chain. Therefore an antibody produced by a host cell by expression of a specific nucleic acid molecule encoding a full-length heavy chain may include the full-length heavy chain, or it may include a cleaved variant of the full-length heavy chain (also referred to herein as a “cleaved variant heavy chain”). This may be the case where the final two C-terminal amino acids of the heavy chain are glycine (G446) and lysine (K447, numbering according to Kabat EU index). Therefore, the C-terminal lysine (Lys447), or the C-terminal glycine (Gly446) and lysine (K447), of the Fc region may or may not be present. Amino acid sequences of heavy chains including Fc domains (or a subunit of an Fc domain as defined herein) are denoted herein without C-terminal glycine-lysine dipeptide if not indicated otherwise. In one embodiment of the invention, a heavy chain including a subunit of an Fc domain as specified herein, comprised in an antibody or bispecific antibody according to the invention, comprises an additional C-terminal glycine-lysine dipeptide (G446 and K447, numbering according to EU index of Kabat). In one embodiment of the invention, a heavy chain including a subunit of an Fc domain as specified herein, comprised in an antibody or bispecific antibody according to the invention, comprises an additional C-terminal glycine residue (G446, numbering according to EU index of Kabat). Compositions/formulations of the invention, such as the pharmaceutical compositions/formulations described herein, comprise a population of antibodies or bispecific antibodies of the invention. The population of antibodies or bispecific antibodies may comprise molecules having a full-length heavy chain and molecules having a cleaved variant heavy chain. The population of antibodies or bispecific antibodies may consist of a mixture of molecules having a full-length heavy chain and molecules having a cleaved variant heavy chain, wherein at least 50%, at least 60%, at least 70%, at least 80% or at least 90% of the antibodies or bispecific antibodies have a cleaved variant heavy chain. In one embodiment of the invention a composition comprising a population of antibodies or bispecific antibodies of the invention comprises an antibody or bispecific antibody comprising a heavy chain including a subunit of an Fc domain as specified herein with an additional C-terminal glycine-lysine dipeptide (G446 and K447, numbering according to EU index of Kabat). In one embodiment of the invention a composition comprising a population of antibodies or bispecific antibodies of the invention comprises an antibody or bispecific antibody comprising a heavy chain including a subunit of an Fc domain as specified herein with an additional C-terminal glycine residue (G446, numbering according to EU index of Kabat). In one embodiment of the invention such a composition comprises a population of antibodies or bispecific antibodies comprised

of molecules comprising a heavy chain including a subunit of an Fc domain as specified herein; molecules comprising a heavy chain including a subunit of a Fc domain as specified herein with an additional C-terminal glycine residue (G446, numbering according to EU index of Kabat); and molecules comprising a heavy chain including a subunit of an Fc domain as specified herein with an additional C-terminal glycine-lysine dipeptide (G446 and K447, numbering according to EU index of Kabat). Unless otherwise specified herein, numbering of amino acid residues in the Fc region or constant region is according to the EU numbering system, also called the EU index, as described in Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD, 1991 (see also above). A “subunit” of an Fc domain as used herein refers to one of the two polypeptides forming the dimeric Fc domain, i.e. a polypeptide comprising C-terminal constant regions of an immunoglobulin heavy chain, capable of stable self-association. For example, a subunit of an IgG Fc domain comprises an IgG CH2 and an IgG CH3 constant domain. In one preferred embodiment such an Fc domain is of human origin, in one preferred of the IgG isotype, in another preferred embodiment of the IgG1 isotype.

"Framework" or "FR" refers to variable domain residues other than complement determining region (CDR) residues. The FR of a variable domain generally consists of four FR domains: FR1, FR2, FR3, and FR4. Accordingly, the CDR and FR sequences generally appear in the following sequence in VH (or VL): FR1-H1(L1)-FR2-H2(L2)-FR3-H3(L3)-FR4.

The terms “full length antibody”, “intact antibody”, and “whole antibody” are used herein interchangeably to refer to an antibody having a structure substantially similar to a native antibody structure or having heavy chains that contain an Fc region as defined herein.

By “fused” is meant that the components (e.g. a Fab molecule and an Fc domain subunit) are linked by peptide bonds, either directly or via one or more peptide linkers.

A “Fab molecule” refers to a protein consisting of the VH and CH1 domain of the heavy chain (the “Fab heavy chain”) and the VL and CL domain of the light chain (the “Fab light chain”) of an immunoglobulin.

By a “crossover” Fab molecule (also termed “Crossfab”) is meant a Fab molecule wherein the variable domains or the constant domains of the Fab heavy and light

chain are exchanged (i.e. replaced by each other), i.e. the crossover Fab molecule comprises a peptide chain composed of the light chain variable domain VL and the heavy chain constant domain 1 CH1 (VL-CH1, in N- to C-terminal direction), and a peptide chain composed of the heavy chain variable domain VH and the light chain constant domain CL (VH-CL, in N- to C-terminal direction). For clarity, in a crossover Fab molecule wherein the variable domains of the Fab light chain and the Fab heavy chain are exchanged, the peptide chain comprising the heavy chain constant domain 1 CH1 is referred to herein as the “heavy chain” of the (crossover) Fab molecule. Conversely, in a crossover Fab molecule wherein the constant domains of the Fab light chain and the Fab heavy chain are exchanged, the peptide chain comprising the heavy chain variable domain VH is referred to herein as the “heavy chain” of the (crossover) Fab molecule.

In contrast thereto, by a “conventional” Fab molecule is meant a Fab molecule in its natural format, i.e. comprising a heavy chain composed of the heavy chain variable and constant domains (VH-CH1, in N- to C-terminal direction), and a light chain composed of the light chain variable and constant domains (VL-CL, in N- to C-terminal direction). The terms “host cell,” “host cell line,” and “host cell culture” are used interchangeably and refer to cells into which exogenous nucleic acid has been introduced, including the progeny of such cells. Host cells include “transformants” and “transformed cells,” which include the primary transformed cell and progeny derived therefrom without regard to the number of passages. Progeny may not be completely identical in nucleic acid content to a parent cell, but may contain mutations. Mutant progeny that have the same function or biological activity as screened or selected for in the originally transformed cell are included herein.

A “human antibody” is one which possesses an amino acid sequence which corresponds to that of an antibody produced by a human or a human cell or derived from a non-human source that utilizes human antibody repertoires or other human antibody-encoding sequences. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues.

A “humanized” antibody refers to a chimeric antibody comprising amino acid residues from non-human CDRs and amino acid residues from human FRs. In certain embodiments, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDRs (e.g., CDRs) correspond to those of a non-human antibody, and all or substantially all of the FRs correspond to those of a human antibody. A humanized antibody

optionally may comprise at least a portion of an antibody constant region derived from a human antibody. A “humanized form” of an antibody, e.g., a non-human antibody, refers to an antibody that has undergone humanization.

5 The term “complementarity determining regions” or “CDRs” as used herein refers to each of the regions of an antibody variable domain which are hypervariable in sequence and/or form structurally defined loops (“hypervariable loops”) and/or contain the antigen-contacting residues (“antigen contacts”). Generally, antibodies comprise six CDRs: three in the VH (CDR-H1, CDR-H2, CDR-H3), and three in the VL (CDR-L1, CDR-L2, CDR-L3). Exemplary CDRs herein include:

- 10 (a) hypervariable loops occurring at amino acid residues 26-32 (CDR-L1), 50-52 (CDR-L2), 91-96 (CDR-L3), 26-32 (CDR-H1), 53-55 (CDR-H2), and 96-101 (CDR-H3) (Chothia and Lesk, *J. Mol. Biol.* 196:901-917 (1987));
- (b) CDRs occurring at amino acid residues 24-34 (CDR-L1), 50-56 (CDR-L2), 89-97 (CDR-L3), 31-35b (CDR-H1), 50-65 (CDR-H2), and 95-102 (CDR-H3) (Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991));
- 15 (c) antigen contacts occurring at amino acid residues 27c-36 (CDR-L1), 46-55 (CDR-L2), 89-96 (CDR-L3), 30-35b (CDR-H1), 47-58 (CDR-H2), and 93-101 (CDR-H3) (MacCallum et al. *J. Mol. Biol.* 262: 732-745 (1996)); and
- 20 (d) combinations of (a), (b), and/or (c), including CDR amino acid residues 24-34 (CDR-L1), 50-56 (CDR-L2), 89-97 (vL3), 31-35 (CDR-H1), 50-63 (CDR-H2), and 95-102 (CDR-H3).

25 Unless otherwise indicated, CDR-residues and other residues in the variable domain (e.g., FR residues) are numbered herein according to Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991).

30 An “individual” or “subject” is a mammal. Mammals include, but are not limited to, domesticated animals (e.g., cows, sheep, cats, dogs, and horses), primates (e.g., humans and non-human primates such as monkeys), rabbits, and rodents (e.g., mice and rats). In certain embodiments, the individual or subject is a human.

An “isolated” antibody is one which has been separated from a component of its natural environment. In one embodiment the antibody is an isolated antibody. In

some embodiments, an antibody is purified to greater than 95% or 99% purity as determined by, for example, electrophoretic (e.g., SDS-PAGE, isoelectric focusing (IEF), capillary electrophoresis) or chromatographic (e.g., ion exchange or reverse phase HPLC). For review of methods for assessment of antibody purity see, e.g.,
5 Flatman, S. et al., J. Chromatogr. B 848 (2007) 79-87.

An "isolated" nucleic acid refers to a nucleic acid molecule that has been separated from a component of its natural environment. An isolated nucleic acid includes a nucleic acid molecule contained in cells that ordinarily contain the nucleic acid molecule, but the nucleic acid molecule is present extrachromosomally or at a
10 chromosomal location that is different from its natural chromosomal location.

"Isolated nucleic acid encoding an anti-HLA-G antibody" refers to one or more nucleic acid molecules encoding antibody heavy and light chains (or fragments thereof), including such nucleic acid molecule(s) in a single vector or separate vectors, and such nucleic acid molecule(s) present at one or more locations in a host
15 cell.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical and/or bind the same epitope, except for possible variant antibodies, e.g., containing naturally occurring mutations or arising during production of a monoclonal antibody preparation, such variants generally
20 being present in minor amounts. In contrast to polyclonal antibody preparations, which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody of a monoclonal antibody preparation is directed against a single determinant on an antigen. Thus, the modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially
25 homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by a variety of techniques, including but not limited to the hybridoma method,
30 recombinant DNA methods, phage-display methods, and methods utilizing transgenic animals containing all or part of the human immunoglobulin loci, such methods and other exemplary methods for making monoclonal antibodies being described herein.

A "modification promoting the association of the first and the second subunit of the Fc domain" is a manipulation of the peptide backbone or the post-translational
35

modifications of an Fc domain subunit that reduces or prevents the association of a polypeptide comprising the Fc domain subunit with an identical polypeptide to form a homodimer. A modification promoting association as used herein particularly includes separate modifications made to each of the two Fc domain subunits desired to associate (i.e. the first and the second subunit of the Fc domain), wherein the modifications are complementary to each other so as to promote association of the two Fc domain subunits. For example, a modification promoting association may alter the structure or charge of one or both of the Fc domain subunits so as to make their association sterically or electrostatically favorable, respectively. Thus, (hetero)dimerization occurs between a polypeptide comprising the first Fc domain subunit and a polypeptide comprising the second Fc domain subunit, which might be non-identical in the sense that further components fused to each of the subunits (e.g. antigen binding moieties) are not the same. In some embodiments the modification promoting association comprises an amino acid mutation in the Fc domain, specifically an amino acid substitution. In a particular embodiment, the modification promoting association comprises a separate amino acid mutation, specifically an amino acid substitution, in each of the two subunits of the Fc domain. Such modification promoting the association of the first and the second subunit of the Fc domain play an important role in the heterodimerization of multi-or bispecific antibodies (see e.g. also below under A.2 Exemplary multispecific anti-HLA-G /anti-CD3 Antibodies)

"Native antibodies" refer to naturally occurring immunoglobulin molecules with varying structures. For example, native IgG antibodies are heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light chains and two identical heavy chains that are disulfide-bonded. From N- to C-terminus, each heavy chain has a variable region (VH), also called a variable heavy domain or a heavy chain variable domain, followed by three constant domains (CH1, CH2, and CH3). Similarly, from N- to C-terminus, each light chain has a variable region (VL), also called a variable light domain or a light chain variable domain, followed by a constant light (CL) domain. The light chain of an antibody may be assigned to one of two types, called kappa (κ) and lambda (λ), based on the amino acid sequence of its constant domain.

The term "package insert" is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, combination therapy, contraindications and/or warnings concerning the use of such therapeutic products.

“Percent (%) amino acid sequence identity” with respect to a reference polypeptide sequence is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the reference polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for aligning sequences, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc., and the source code has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available from Genentech, Inc., South San Francisco, California, or may be compiled from the source code. The ALIGN-2 program should be compiled for use on a UNIX operating system, including digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

100 times the fraction X/Y

where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program’s alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program.

The term "pharmaceutical formulation" refers to a preparation which is in such form as to permit the biological activity of an active ingredient contained therein to be effective, and which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered.

5 A "pharmaceutically acceptable carrier" refers to an ingredient in a pharmaceutical formulation, other than an active ingredient, which is nontoxic to a subject. A pharmaceutically acceptable carrier includes, but is not limited to, a buffer, excipient, stabilizer, or preservative.

10 As used herein, "treatment" (and grammatical variations thereof such as "treat" or "treating") refers to clinical intervention in an attempt to alter the natural course of the individual being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Desirable effects of treatment include, but are not limited to, preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease,
15 preventing metastasis, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis. In some embodiments, antibodies of the invention are used to delay development of a disease or to slow the progression of a disease.

20 The term "variable region" or "variable domain" refers to the domain of an antibody heavy or light chain that is involved in binding the antibody to antigen. The variable domains of the heavy chain and light chain (VH and VL, respectively) of a native antibody generally have similar structures, with each domain comprising four conserved framework regions (FRs) and three complement determining regions (CDRs). (See, e.g., Kindt, T.J. et al. Kuby Immunology, 6th ed., W.H. Freeman and
25 Co., N.Y. (2007), page 91) A single VH or VL domain may be sufficient to confer antigen-binding specificity. Furthermore, antibodies that bind a particular antigen may be isolated using a VH or VL domain from an antibody that binds the antigen to screen a library of complementary VL or VH domains, respectively. See e.g., Portolano, S. et al., J. Immunol. 150 (1993) 880-887; Clackson, T. et al., Nature 352
30 (1991) 624-628).

The term "vector," as used herein, refers to a nucleic acid molecule capable of propagating another nucleic acid to which it is linked. The term includes the vector as a self-replicating nucleic acid structure as well as the vector incorporated into the genome of a host cell into which it has been introduced. Certain vectors are capable

of directing the expression of nucleic acids to which they are operatively linked. Such vectors are referred to herein as "expression vectors".

I. COMPOSITIONS AND METHODS

5 In one aspect, the invention is based, in part, on the finding that surprisingly among various variants of HLA-G-0090 in which the glycosylation site was removed only the two variants HLA-G-0090-VL-S32P and HLA-G-0090-VL-S33A show even improved binding properties, good expressability and stability, while showing no more glycosylation at the CDR-L1 of the LC (so no Fab glycosylation could be detected). As all recently approved pharmaceutical antibody products are produced
10 in mammalian cells, especially CHO cells (see e.g. Walsh G., Nature Biotech (2018) 1136-1145) providing an antibody without glycosylation sites in the binding region (VH and VL and especially the CDRs) represents a valuable advantage, as these antibodies can then be readily used for production in mammalian expression systems without the risk of (at least partially impairing the binding properties by
15 glycosylation. In particular for Fc domain-comprising antibodies, manufacturing the antibody in a host cell lacking a glycosylation machinery (such as a prokaryotic host cell) would not results in a product of comparable quality, since the N-glycans attached to amino acid residue ASN297 (numbering according to EU index of Kabat) in the Fc region are required to maintain solubility and thermal stability, and to
20 prevent aggregation of the antibody in an aqueous solution (e.g. in pharmaceutical compositions).

A.1 Exemplary anti-HLA-G Antibodies

One embodiment of the invention is an antibody that binds to human HLA-G comprising

5 A) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:23; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID
10 NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6, or

B) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:25; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID
15 NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6.

One embodiment of the invention is an antibody that binds to human HLA-G wherein the antibody

A) comprises a VH domain comprising the amino acid sequence of SEQ ID
20 NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24; or

B) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26.

One embodiment of the invention is an antibody that binds to human HLA-G wherein the antibody comprises a VH domain comprising the amino acid sequence of SEQ
25 ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24.

One embodiment of the invention is an antibody that binds to human HLA-G wherein the antibody comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24.

In embodiment such anti-HLA-G antibody has improved binding properties with
30 respect to maximal binding (R_{max}) and/or binding affinity (K_D) compared to the (parental) antibody that comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID

NO:8 (as shown in Example 2). In a further embodiment the HLA-G antibody of the present invention comprises an Fc domain of human origin, in one embodiment the Fc domain is of the IgG isotype, in one preferred embodiment of the IgG1 isotype. In one embodiment, such an IgG1 isotype Fc domain of human origin comprises the amino acid mutations L234A, L235A and P329G (“P329G LALA”, “PGLALA” or “LALAPG”) (numberings according to Kabat EU index).

In a further embodiment the HLA-G antibody of the present invention comprises a constant region of human origin, particularly of the IgG isotype, more particularly of the IgG1 isotype, comprising a human CH1, CH2, CH3 and/or CL domain.

In one embodiment such such constant region of the IgG1 isotype comprises the amino acid mutations L234A, L235A and P329G (“P329G LALA”, “PGLALA” or “LALAPG”) (numberings according to Kabat EU index).

Such Fc domain-comprising antibodies can be typically N-glycosylated at position ASN-297 (numbering according to Kabat EU index) e.g. when produced in eukaryotic cells, like mammalian cells, in particular CHO cells. N-glycosylation at position ASN-297 (numbering according to EU index (see Kabat) represents a valuable contribution to e.g. the high stability, low aggregation tendency and/or good pharmacokinetic and other critical quality properties of such an antibody (see e.g. Zheng et al, mAbs (2011) 568-576; and Reusch et al, Glycobiology (2015) 1325–133). Therefore such an Fc domain-comprising antibody of the present invention is easily ready for production in in eukaryotic cells, like mammalian cells, in particular CHO cells, without the risk of being glycosylated in the binding region (which would interfere with its binding properties), but at the same time with the benefit and valuable quality attributes of the N-glycosylation at position ASN-297.

One embodiment of the invention is an antibody that binds to human HLA-G wherein the antibody comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24, wherein antibody has improved binding properties with respect to maximal binding (R_{max}) and/or binding affinity (K_D) compared to the (parental) antibody that comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:8 (as shown in Example 2).

In embodiment the anti-HLA-G antibody

- 31 -

- a) does not crossreact with a modified human HLA-G β 2M MHC I complex, wherein the HLA-G specific amino acids have been replaced by HLA-A consensus amino acids, the complex comprising SEQ ID NO:40; and/ or
- 5 b) does not crossreact with a mouse H2Kd β 2M MHC I complex comprising SEQ ID NO:41; and/ or
- c) does not crossreact with rat RT1A β 2M MHC I complex comprising SEQ ID NO:43.

In embodiment the anti-HLA-G antibody

- 10 a) inhibits ILT2 binding to (HLA-G expressed on) JEG3 cells (ATCC No. HTB36); or
- b) binds to (HLA-G expressed on) JEG3 cells (ATCC No. HTB36), and inhibits ILT2 binding to (HLA-G expressed on) JEG-3 cells (ATCC No. HTB36).

15 In another aspect, the invention relates to multispecific antibodies comprising the anti-HLA-G antigen binding moiety. These multispecific antibodies (e.g. the bispecific antibodies) as described herein use the selected, improved anti-HLA-G antibodies as first antigen binding moiety/site. These anti-HLA-G antibodies bind to HLA-G with high specificity and affinity (improved binding properties, no crossreactivity with other species and human HLA-A consensus sequences), and have ability to specifically inhibit ILT2 and or ILT4 binding to HLA-G.

20 In one embodiment the invention relates to a multispecific (preferably bispecific) that binds to human HLA-G and to human CD3. In one embodiment the invention relates to a multispecific (preferably bispecific) anti-HLA-G /anti-CD3 antibody, wherein the multispecific (preferably bispecific) antibody that binds to human HLA-G and to human CD3, comprises a first antigen binding moiety that binds to human

25 HLA-G and a second antigen binding moiety that binds to human CD3. This bispecific antibody as described herein binds with specific, second antigen binding moieties /sites to CD3, especially CD3 ϵ and are therefore able to attract CD3 expressing T-cells to HLA-G expressing tumor cells and at the same time to inhibit the HLA-G induced immune suppression in the tumor environment by blocking

30 ILT2/4 binding to HLA-G. Thus these bispecific anti-HLA-G /anti-CD3 antibodies show strong tumor growth inhibition and tumor regression in vivo.

A.2 Exemplary multispecific anti-HLA-G /anti-CD3 Antibodies

One embodiment of the invention is a bispecific antibody that binds to human HLA-G and to human CD3, comprising a first antigen binding moiety that binds to human HLA-G and a second antigen binding moiety that binds to human CD3,
5 wherein the first antigen binding moiety that binds to human HLA-G comprises

A) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence
10 of SEQ ID NO:23; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6, or

B) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence
15 of SEQ ID NO:25; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6;

and wherein the second antigen binding moiety that binds to a T cell activating antigen binds to human CD3 comprises

C) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:52, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:53, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:54; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:55; (ii) CDR-L2 comprising the amino acid sequence of
20 SEQ ID NO:56 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:57, or

C) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:60, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:61, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:62; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:63; (ii) CDR-L2 comprising the amino acid sequence of
30 SEQ ID NO:64 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:65, or

- 33 -

5 D) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:68, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:69, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:70; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:71; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:72 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:73.

10 Another embodiment of the invention is a bispecific antibody that binds to human HLA-G and to human CD3, comprising a first antigen binding moiety that binds to human HLA-G and a second antigen binding moiety that binds to human CD3,

wherein the first antigen binding moiety

A) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24; or

15 B) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26,

and wherein the second antigen binding moiety

C) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:58 and a VL domain comprising the amino acid sequence of SEQ ID NO:59; or

20 D) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:66 and a VL domain comprising the amino acid sequence of SEQ ID NO:67; or

E) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:74 and a VL domain comprising the amino acid sequence of SEQ ID NO:75.

25 Another embodiment of the invention is a bispecific antibody that binds to human HLA-G and to human CD3, comprising a first antigen binding moiety that binds to human HLA-G and a second antigen binding moiety that binds to human CD3,

wherein the first antigen binding moiety

30 comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24;

- 34 -

and wherein the second antigen binding moiety

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:58 and
a VL domain comprising the amino acid sequence of SEQ ID NO:59.

5 Another embodiment of the invention is a bispecific antibody that binds to human
HLA-G and to human CD3, comprising a first antigen binding moiety that
binds to human HLA-G and a second antigen binding moiety that binds to
human CD3,

wherein the first antigen binding moiety

10 comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and
a VL domain comprising the amino acid sequence of SEQ ID NO:24;

and wherein the second antigen binding moiety

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:66 and
a VL domain comprising the amino acid sequence of SEQ ID NO:67.

15 Another embodiment of the invention is a bispecific antibody that binds to human
HLA-G and to human CD3, comprising a first antigen binding moiety that
binds to human HLA-G and a second antigen binding moiety that binds to
human CD3,

wherein the first antigen binding moiety

20 comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and
a VL domain comprising the amino acid sequence of SEQ ID NO:24;

and wherein the second antigen binding moiety

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:74 and
a VL domain comprising the amino acid sequence of SEQ ID NO:75.

25 In one embodiment these bispecific antibodies are characterized by one or more of
the following properties :

a) induction of T cell mediated cytotoxicity/tumor cell killing in the presence
of HLA-G expressing tumor cells (preferably in the presence of JEG3 cells (ATCC
No. HTB36)); and/or

- 35 -

b) induction IFN gamma secretion by T cells in the presence of HLA-G expressing tumor cells (preferably in the presence of JEG3 cells (ATCC No. HTB36)); and/or

c) inhibition of tumor growth in vivo (in a mouse xenograft tumor model),

5 d) in vivo anti-tumor efficacy/ tumor regression in humanized NSG mice bearing SKOV3 human ovarian carcinoma transfected with recombinant HLA-G (SKOV3 HLA-G) humanized NSG mice (see Example 13); and/or

e) in vivo anti-tumor efficacy /tumor of HLA-G CD3 T cell bi-specific in humanized NSG mice bearing human breast cancer PDX tumors (BC004) (see Example 14).

10 In one embodiment these bispecific antibodies are characterized in addition by one or more of the following properties: the bispecific antibody

a) does not crossreact with a modified human HLA-G β 2M MHC I complex, wherein the HLA-G specific amino acids have been replaced by HLA-A consensus amino acids, the complex comprising SEQ ID NO:44; and/ or

15 b) does not crossreact with a mouse H2Kd β 2M MHC I complex comprising SEQ ID NO:41; and/ or

c) does not crossreact with rat RT1A β 2M MHC I complex comprising SEQ ID NO:43.

In one embodiment these bispecific antibodies are characterized in addition by one or more of the following properties: the bispecific antibody

20 a) inhibits ILT2 binding to (HLA-G expressed on) JEG3 cells (ATCC No. HTB36); or

b) binds to (HLA-G expressed on) JEG3 cells (ATCC No. HTB36), and inhibits ILT2 binding to (HLA-G expressed on) JEG-3 cells (ATCC No. HTB36); and/or

Multispecific antibodies

25 Multispecific antibodies are monoclonal antibodies that have binding specificities for at least two different sites, i.e., different epitopes on different antigens or different epitopes on the same antigen. In certain embodiments, the multispecific antibody has three or more binding specificities. In a preferred embodiment the multispecific antibody provided herein is a bispecific antibody. In certain embodiments, one of the

binding specificities is for HLA-G and the other specificity is for CD3. In certain embodiments, bispecific antibodies may bind to two (or more) different epitopes of HLA-G. Multispecific antibodies can be prepared as full length antibodies or antibody fragments.

5 Techniques for making multispecific and in particular bispecific antibodies include, but are not limited to, recombinant co-expression of two immunoglobulin heavy chain-light chain pairs having different specificities (see Milstein and Cuello, *Nature* 305: 537 (1983)) and “knob-in-hole” engineering (see, e.g., U.S. Patent No. 5,731,168, and Atwell et al., *J. Mol. Biol.* 270:26 (1997)). Multi-specific antibodies
10 may also be made by engineering electrostatic steering effects for making antibody Fc-heterodimeric molecules (see, e.g., WO 2009/089004); cross-linking two or more antibodies or fragments (see, e.g., US Patent No. 4,676,980, and Brennan et al., *Science*, 229: 81 (1985)); using leucine zippers to produce bi-specific antibodies (see, e.g., Kostelny et al., *J. Immunol.*, 148(5):1547-1553 (1992) and WO
15 2011/034605); using the common light chain technology for circumventing the light chain mis-pairing problem (see, e.g., WO 98/50431); using “diabody” technology for making bispecific antibody fragments (see, e.g., Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993)); and using single-chain Fv (sFv) dimers (see, e.g. Gruber et al., *J. Immunol.*, 152:5368 (1994)); and preparing trispecific antibodies as
20 described, e.g., in Tutt et al. *J. Immunol.* 147: 60 (1991).

Engineered antibodies with three or more antigen binding sites, including for example, “Octopus antibodies,” or DVD-Ig are also included herein (see, e.g. WO 2001/77342 and WO 2008/024715). Other examples of multispecific antibodies with
25 three or more antigen binding sites can be found in WO 2010/115589, WO 2010/112193, WO 2010/136172, WO2010/145792, and WO 2013/026831. The bispecific antibody or antigen binding fragment thereof also includes a “Dual Acting FAb” or “DAF” comprising an antigen binding site that binds to HLA-G as well as another different antigen, or two different epitopes of HLA-G (see, e.g., US 2008/0069820 and WO 2015/095539).

30 Multi-specific antibodies may also be provided in an asymmetric form with a domain crossover in one or more binding arms of the same antigen specificity, i.e. by exchanging the VH/VL domains (see e.g., WO 2009/080252 and WO 2015/150447), the CH1/CL domains (see e.g., WO 2009/080253) or the complete Fab arms (see e.g., WO 2009/080251, WO 2016/016299, also see Schaefer et al, PNAS, 108 (2011)
35 1187-1191, and Klein et al., *MAbs* 8 (2016) 1010-20). Asymmetrical Fab arms can

also be engineered by introducing charged or non-charged amino acid mutations into domain interfaces to direct correct Fab pairing. See e.g., WO 2016/172485.

Various further molecular formats for multispecific antibodies are known in the art and are included herein (see e.g., Spiess et al., Mol Immunol 67 (2015) 95-106).

5 A particular type of multispecific antibodies, also included herein, are bispecific antibodies designed to simultaneously bind to a surface antigen on a target cell, e.g., a tumor cell, and to an activating, invariant component of the T cell receptor (TCR) complex, such as CD3, for retargeting of T cells to kill target cells. Hence, in certain
10 embodiments, an antibody provided herein is a multispecific antibody, particularly a bispecific antibody, wherein one of the binding specificities is for HLA-G and the other is for CD3.

Examples of bispecific antibody formats that may be useful for this purpose include, but are not limited to, the so-called “BiTE” (bispecific T cell engager) molecules wherein two scFv molecules are fused by a flexible linker (see, e.g.,
15 WO2004/106381, WO2005/061547, WO2007/042261, and WO2008/119567, Nagorsen and Bäuerle, Exp Cell Res 317, 1255-1260 (2011)); diabodies (Holliger et al., Prot Eng 9, 299-305 (1996)) and derivatives thereof, such as tandem diabodies (“TandAb”; Kipriyanov et al., J Mol Biol 293, 41-56 (1999)); “DART” (dual affinity
20 retargeting) molecules which are based on the diabody format but feature a C-terminal disulfide bridge for additional stabilization (Johnson et al., J Mol Biol 399, 436-449 (2010)), and so-called triomabs, which are whole hybrid mouse/rat IgG molecules (reviewed in Seimetz et al., Cancer Treat Rev 36, 458-467 (2010)). Particular T cell bispecific antibody formats included herein are described in WO
25 2013/026833, WO2013/026839, WO 2016/020309; Bacac et al., Oncoimmunology 5(8) (2016) e1203498.

Bispecific antibodies that bind to HLA-G and to CD3

The invention also provides a bispecific antibody, i.e. an antibody that comprises at least two antigen binding moieties capable of specific binding to two distinct antigenic determinants (a first and a second antigen).

30 Based on the anti-HLA-G antigen binding moieties and anti-CD3 antigen binding moieties they developed, the present inventors have developed bispecific antibodies that bind to HLA-G and to CD3.

As shown in the Examples, these bispecific antibodies have a number of remarkable properties, including good efficacy and low toxicity.

Thus, in certain aspects, the invention provides a bispecific antibody, comprising (a) a first antigen binding moiety that binds to human HLA-G, and (b) a second antigen binding moiety which specifically binds to human CD3, wherein the bispecific antibody has any of the following features:-The bispecific antibody of the invention specifically induces T-cell mediated killing of cells expressing HLA-G. In some embodiments, the bispecific antibody of the invention specifically induces T-cell mediated killing of cells expressing HLA-G. In a more specific embodiment, the bispecific antibody specifically induces T-cell mediated killing of cells expressing HLA-G.

-In one embodiment, induction of T-cell mediated killing by the bispecific antibody is determined using HLA-G -expressing cells.

-In one embodiment, activation of T cells by the bispecific antibody is determined by measuring, particularly by flow cytometry, expression of CD25 and/or CD69 by T cells after incubation with the bispecific antibody in the presence of HLA-G -expressing cells,.

In a specific embodiment, induction of T-cell mediated killing by the bispecific antibody is determined as follows:

Ability of anti HLA-G/anti CD3 TCB to activate T cells in the presence of HLA-G expressing tumor cells is tested on SKOV3 cells transfected with recombinant HLA-G (SKOV3HLA-G). Activation of T cells is assessed by FACS analysis of cell surface activation markers CD25 and early activation marker CD69 on T cells. Briefly, Peripheral Blood Mononuclear Cells (PBMCs) are isolated from human peripheral blood by density gradient centrifugation using Lymphocyte Separating Medium Tubes (PAN #P04-60125). PBMC`s and SKOV3HLA-G cells are seeded at a ratio of 10 : 1 in 96-well U bottom plates. The co-culture is then incubated with HLA-G-TCB at different concentrations as described in the Example 12 and incubated for 24h at 37°C in an incubator with 5% Co2. On the next day, expression of CD25 and CD69 is measured by flow cytometry.

For flow cytometry analysis, cells are stained with with PerCP-Cy5.5 Mouse Anti-Human CD8 (BD Pharmingen # 565310), PE Mouse Anti-Human CD25 (eBioscience # 9012-0257) and APC Mouse Anti-Human CD69 (BD Pharmingen #

555533) at 4°C. Briefly, antibodies are diluted to a 2-fold concentration and 25µl of antibody dilution are added in each well with 25µl of pre-washed co-cultures. Cells are stained for 30 min at 4°C and washed twice with 200µl/well staining buffer and centrifugation at 300g for 5min. Cell pellets are resuspended in 200µl of staining buffer and stained with DAPI for live dead discrimination at a final concentration of 2µg/ml. Samples are then measured using BD LSR flow cytometer. Data analysis is performed using FlowJo V.10.1 software.

The bispecific antibody of the invention specifically activates T cells in the presence of cells expressing HLA-G. In some embodiments, the bispecific antibody of the invention specifically activates T cells in the presence of cells expressing HLA-G. In a more specific embodiment, the bispecific antibody specifically activates T cells in the presence of cells expressing HLA-G.

In one embodiment, the bispecific antibody induces T cell mediated killing of, or activate T cells in the presence of, cells expressing HLA-G,. In one embodiment, the bispecific antibody induces T cell mediated killing of, and/or activates T cells in the presence of, cells expressing HLA-G with an EC50 that is at least 5, at least 10, at least 15, at least 20, at least 25, at least 50, at least 75 or at least 100 times lower than the EC50 for induction of T cell mediated killing of, or activation of T cells in the presence of, cells expressing HLA-G

According to particular embodiments of the invention, the antigen binding moieties comprised in the bispecific antibody are Fab molecules (i.e. antigen binding domains composed of a heavy and a light chain, each comprising a variable and a constant domain). In one embodiment, the first and/or the second antigen binding moiety is a Fab molecule. In one embodiment, said Fab molecule is human. In a particular embodiment, said Fab molecule is humanized. In yet another embodiment, said Fab molecule comprises human heavy and light chain constant domains.

Preferably, at least one of the antigen binding moieties is a crossover Fab molecule. Such modification reduces mispairing of heavy and light chains from different Fab molecules, thereby improving the yield and purity of the bispecific antibody of the invention in recombinant production. In a particular crossover Fab molecule useful for the bispecific antibody of the invention, the variable domains of the Fab light chain and the Fab heavy chain (VL and VH, respectively) are exchanged. Even with this domain exchange, however, the preparation of the bispecific antibody may comprise certain side products due to a so-called Bence Jones-type interaction between mispaired heavy and light chains (see Schaefer et al, PNAS, 108 (2011)

- 40 -

11187-11191). To further reduce mispairing of heavy and light chains from different Fab molecules and thus increase the purity and yield of the desired bispecific antibody, charged amino acids with opposite charges may be introduced at specific amino acid positions in the CH1 and CL domains of either the Fab molecule(s) binding to the first antigen (HLA-G), or the Fab molecule binding to the second antigen an activating T cell antigen such as CD3, as further described herein. Charge modifications are made either in the conventional Fab molecule(s) comprised in the bispecific antibody (such as shown e.g. in Figures 13 A-C, G-J), or in the VH/VL crossover Fab molecule(s) comprised in the bispecific antibody (such as shown e.g. in Figure 13 D-F, K-N) (but not in both). In particular embodiments, the charge modifications are made in the conventional Fab molecule(s) comprised in the bispecific antibody (which in particular embodiments bind(s) to the first antigen, i.e. HLA-G).

In a particular embodiment according to the invention, the bispecific antibody is capable of simultaneous binding to the first antigen (i.e. HLA-G), and the second antigen (e.g. an activating T cell antigen, particularly CD3). In one embodiment, the bispecific antibody is capable of crosslinking a T cell and a target cell by simultaneous binding HLA-G and an activating T cell antigen. In an even more particular embodiment, such simultaneous binding results in lysis of the target cell, particularly a HLA-G expressing tumor cell. In one embodiment, such simultaneous binding results in activation of the T cell. In other embodiments, such simultaneous binding results in a cellular response of a T lymphocyte, particularly a cytotoxic T lymphocyte, selected from the group of: proliferation, differentiation, cytokine secretion, cytotoxic effector molecule release, cytotoxic activity, and expression of activation markers. In one embodiment, binding of the bispecific antibody to the activating T cell antigen, particularly CD3, without simultaneous binding to HLA-G does not result in T cell activation.

In one embodiment, the bispecific antibody is capable of re-directing cytotoxic activity of a T cell to a target cell. In a particular embodiment, said re-direction is independent of MHC-mediated peptide antigen presentation by the target cell and and/or specificity of the T cell.

Particularly, a T cell according to any of the embodiments of the invention is a cytotoxic T cell. In some embodiments the T cell is a CD4⁺ or a CD8⁺ T cell, particularly a CD8⁺ T cell.

35

First antigen binding moiety that binds to human HLA-G

The bispecific antibody of the invention comprises at least one antigen binding moiety, particularly a Fab molecule, that binds to human HLA-G (first antigen). In certain embodiments, the bispecific antibody comprises two antigen binding moieties, particularly Fab molecules, which bind to human HLA-G. In a particular such embodiment, each of these antigen binding moieties binds to the same antigenic determinant. In an even more particular embodiment, all of these antigen binding moieties are identical, i.e. they comprise the same amino acid sequences including the same amino acid substitutions in the CH1 and CL domain as described herein (if any). In one embodiment, the bispecific antibody comprises not more than two antigen binding moieties, particularly Fab molecules, which bind to human HLA-G.

In particular embodiments, the antigen binding moiety(ies) which bind to human HLA-G is/are a conventional Fab molecule. In such embodiments, the antigen binding moiety(ies) that binds to a second antigen is a crossover Fab molecule as described herein, i.e. a Fab molecule wherein the variable domains VH and VL or the constant domains CH1 and CL of the Fab heavy and light chains are exchanged / replaced by each other.

In alternative embodiments, the antigen binding moiety(ies) which bind to human HLA-G is/are a crossover Fab molecule as described herein, i.e. a Fab molecule wherein the variable domains VH and VL or the constant domains CH1 and CL of the Fab heavy and light chains are exchanged / replaced by each other. In such embodiments, the antigen binding moiety(ies) that binds a second antigen is a conventional Fab molecule.

The HLA-G binding moiety is able to direct the bispecific antibody to a target site, for example to a specific type of tumor cell that expresses human HLA-G.

The first antigen binding moiety of the bispecific antibody may incorporate any of the features, singly or in combination, described herein in relation to the antibody that binds HLA-G, unless scientifically clearly unreasonable or impossible.

Thus, in one aspect, the invention provides a bispecific antibody, comprising a first antigen binding moiety that binds to a first antigen, wherein the first antigen is human HLA-G (in one embodiment the antibody binds to HLA-G β 2M MHC I complex comprising SEQ ID NO: 39), and the first antigen binding moiety comprises

(a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising an amino acid sequence of SEQ ID NO:3; and wherein the VH domain comprises an amino acid sequence of at least 95%, 96%, 97%, 98%, 99% or 100% (in one preferred embodiment 98% or 99% or 100%) sequence identity to the amino acid sequence of SEQ ID NO: 7; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:23; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6; and wherein the VL domain comprises an amino acid sequence of at least 95%, 96%, 97%, 98%, 99% or 100% (in one preferred embodiment 98% or 99% or 100%) sequence identity to the amino acid sequence of SEQ ID NO: 24.

The term like “a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising an amino acid sequence of SEQ ID NO:3; and wherein the VH domain comprises an amino acid sequence of at least 95%, 96%, 97%, 98%, 99% or 100% (in one preferred embodiment 98% or 99% or 100%) sequence identity to the amino acid sequence of SEQ ID NO: 7” refers to a VH domain with an amino acid sequence of SEQ ID NO: 7 wherein the 3 CDRs are unchanged (i.e. the same as in SEQ ID NO:7) but e.g. no, one, two, three, four or five amino acid residues in the framework regions of the VH is/are changed/substituted with another amino acid without affecting the binding properties of the VH and the antigen binding site. As the framework residues with a high probability to influence on the binding properties are well known (see e.g. Foote J. and Winter G., J. Mol. Biol. (1992) 224, 487-499), the framework residues with no or minor influence can be chosen for substitution. The same applies to analogous terms used herein relating to another VH or VL.

In one embodiment the first antigen binding moiety comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24.

In one embodiment the first binding moiety that binds to human HLA-G (in one embodiment to HLA-G β 2M MHC I complex comprising SEQ ID NO: 39), comprises

(a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising an amino acid sequence of SEQ ID NO:3; and wherein the VH domain comprises an amino acid sequence of at least 95%, 96%, 97%, 98%, 99% or 100% (in one preferred embodiment 98% or 99% or 100%) sequence identity to the amino acid sequence of SEQ ID NO: 7; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:25; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6; and wherein the VL domain comprises an amino acid sequence of at least 95%, 96%, 97%, 98%, 99% or 100% (in one preferred embodiment 98% or 99% or 100%) sequence identity to the amino acid sequence of SEQ ID NO: 26.

In one embodiment the first antigen binding moiety comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26.

Such anti-HLA-G antibodies show highly valuable properties, as they have no N-glycosylation in the antigen binding site (and the CDR-L1) (as shown in Example 2), have improved binding properties with respect to maximal binding (R_{max}) and/or binding affinity (KD) compared to the (parental) antibody that comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:8 (as shown in Example 2), do not crossreact with a HLA-A MHC I complexes, or murine or rat MHC I complexes and bind to (HLA-G expressed on) JEG3 cells (ATCC No. HTB36), and inhibits ILT2 binding to (HLA-G expressed on) JEG-3 cells (ATCC No. HTB36).

Second antigen binding moiety that binds to human CD3

The bispecific antibody of the invention comprises at least one antigen binding moiety, particularly a Fab molecule, that binds to human CD3.

In particular embodiments, the antigen binding moiety that binds human CD3, is a crossover Fab molecule as described herein, i.e. a Fab molecule wherein the variable domains VH and VL or the constant domains CH1 and CL of the Fab heavy and light chains are exchanged / replaced by each other. In such embodiments, the antigen binding moiety(ies) that binds to human HLA-G is preferably a conventional Fab molecule. In embodiments where there is more than one antigen binding moiety,

particularly Fab molecule, that binds to human CD3 comprised in the bispecific antibody, the antigen binding moiety that binds human CD3 preferably is a crossover Fab molecule and the antigen binding moieties that bind to human HLA-G are conventional Fab molecules.

5 In alternative embodiments, the antigen binding moiety that binds to the second antigen is a conventional Fab molecule. In such embodiments, the antigen binding moiety(ies) that binds to the first antigen (i.e. HLA-G) is a crossover Fab molecule as described herein, i.e. a Fab molecule wherein the variable domains VH and VL or the constant domains CH1 and CL of the Fab heavy and light chains are exchanged
10 / replaced by each other. In embodiments where there is more than one antigen binding moiety, particularly Fab molecule, that binds to a second antigen comprised in the bispecific antibody, the antigen binding moiety that binds to human HLA-G preferably is a crossover Fab molecule and the antigen binding moieties that bind to human CD3 are conventional Fab molecules.

15 In some embodiments, the second antigen is an activating T cell antigen (also referred to herein as an “activating T cell antigen binding moiety, or activating T cell antigen binding Fab molecule”). In a particular embodiment, the bispecific antibody comprises not more than one antigen binding moiety capable of specific binding to an activating T cell antigen. In one embodiment the bispecific antibody provides
20 monovalent binding to the activating T cell antigen.

In particular embodiments, the second antigen is CD3, particularly human CD3 (SEQ ID NO: 88) or cynomolgus CD3 (SEQ ID NO: 89), most particularly human CD3. In one embodiment the second antigen binding moiety is cross-reactive for (i.e. specifically binds to) human and cynomolgus CD3. In some embodiments, the
25 second antigen is the epsilon subunit of CD3 (CD3 epsilon).

In one embodiment, the second antigen binding moiety that binds to human CD3 comprises (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:52, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:53, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID
30 NO:54; and wherein the VH domain comprises an amino acid sequence of at least 95%, 96%, 97%, 98%, 99% or 100% (in one preferred embodiment 98% or 99% or 100%) sequence identity to the amino acid sequence of SEQ ID NO: 58; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:55; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:56 and (iii)
35 CDR-L3 comprising the amino acid sequence of SEQ ID NO:57, and wherein the

VL domain comprises an amino acid sequence of at least 95%, 96%, 97%, 98%, 99% or 100% (in one preferred embodiment 98% or 99% or 100%) sequence identity to the amino acid sequence of SEQ ID NO: 59.

5 In one embodiment, the second antigen binding moiety that binds to human CD3 comprises (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:60, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:61, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:62, and wherein the VH domain comprises an amino acid sequence of at least 95%, 96%, 97%, 98%, 99% or 100% (in one preferred embodiment 98% or 99% or 100%) sequence identity to the amino acid sequence of SEQ ID NO: 66; and (b) a
10 VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:63; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:64 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:65, and wherein the VL domain comprises an amino acid sequence of at least 95%, 96%, 97%, 98%, 99%
15 or 100% (in one preferred embodiment 98% or 99% or 100%) sequence identity to the amino acid sequence of SEQ ID NO: 67.

In one embodiment, the second antigen binding moiety that binds to human CD3 comprises (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:68, (ii) CDR-H2 comprising the amino acid sequence of
20 SEQ ID NO:69, and wherein the VH domain comprises an amino acid sequence of at least 95%, 96%, 97%, 98%, 99% or 100% (in one preferred embodiment 98% or 99% or 100%) sequence identity to the amino acid sequence of SEQ ID NO: 74; and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:70; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID
25 NO:71; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:72 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:73, and wherein the VL domain comprises an amino acid sequence of at least 95%, 96%, 97%, 98%, 99% or 100% (in one preferred embodiment 98% or 99% or 100%) sequence identity to the amino acid sequence of SEQ ID NO: 75.

30 In one embodiment, the second antigen binding moiety that binds to human CD3 comprises a VH domain comprising the amino acid sequence of SEQ ID NO: 58, and a VL domain comprising the amino acid sequence of SEQ ID NO: 59.

In one embodiment, the second antigen binding moiety that binds to human CD3 comprises a VH domain comprising the amino acid sequence of SEQ ID NO: 66,
35 and a VL domain comprising the amino acid sequence of SEQ ID NO: 67.

- 46 -

In one embodiment, the second antigen binding moiety that binds to human CD3 comprises a VH domain comprising the amino acid sequence of SEQ ID NO: 74, and a VL domain comprising the amino acid sequence of SEQ ID NO: 75.

5 Such CD3 antigen binding moieties/sites show highly valuable properties (e.g. when provided as bispecific antibodies binding to CD3 and HLA-G (with the HLA-G antigen binding moieties as described herein). They show[^]

a) good thermal stability

10 b) induction IFN gamma secretion by T cells in the presence of HLA-G expressing tumor cells (preferably in the presence of JEG3 cells (ATCC No. HTB36)) (Example 11); and/or

c) induction of T cell mediated cytotoxicity/tumor cell killing in the presence of HLA-G expressing tumor cells (preferably in the presence of JEG3 cells (ATCC No. HTB36)) (Example 12) ; and/or

d) inhibition of tumor growth in vivo (in a mouse xenograft tumor model),

15 e) in vivo anti-tumor efficacy/ tumor regression in humanized NSG mice bearing SKOV3 human ovarian carcinoma transfected with recombinant HLA-G (SKOV3 HLA-G) humanized NSG mice (see Example 13); and/or

f) in vivo anti-tumor efficacy /tumor of HLA-G CD3 T cell bi-specific in humanized NSG mice bearing human breast cancer PDX tumors (BC004) (see Example 14).

20 In some embodiments, the second antigen binding moiety is a Fab molecule wherein the variable domains VL and VH or the constant domains CL and CH1, particularly the variable domains VL and VH, of the Fab light chain and the Fab heavy chain are replaced by each other (i.e. according to such embodiment, the second antigen binding moiety is a crossover Fab molecule wherein the variable or constant domains
25 of the Fab light chain and the Fab heavy chain are exchanged). In one such embodiment, the first (and the third, if any) antigen binding moiety is a conventional Fab molecule.

30 In one embodiment, not more than one antigen binding moiety that binds to the second antigen (e.g. an activating T cell antigen such as CD3) is present in the bispecific antibody (i.e. the bispecific antibody provides monovalent binding to the second antigen).

Charge modifications

The bispecific antibodies of the invention may comprise amino acid substitutions in Fab molecules comprised therein which are particularly efficient in reducing mispairing of light chains with non-matching heavy chains (Bence-Jones-type side products), which can occur in the production of Fab-based bi-/ antibodies with a VH/VL exchange in one (or more, in case of molecules comprising more than two antigen-binding Fab molecules) of their binding arms (see also PCT publication no. WO 2015/150447, particularly the examples therein, incorporated herein by reference in its entirety). The ratio of a desired bispecific antibody compared to undesired side products, in particular Bence Jones-type side products occurring in bispecific antibodies with a VH/VL domain exchange in one of their binding arms, can be improved by the introduction of charged amino acids with opposite charges at specific amino acid positions in the CH1 and CL domains (sometimes referred to herein as “charge modifications”).

Accordingly, in some embodiments wherein the first and the second antigen binding moiety of the bispecific antibody are both Fab molecules, and in one of the antigen binding moieties (particularly the second antigen binding moiety) the variable domains VL and VH of the Fab light chain and the Fab heavy chain are replaced by each other,

i) in the constant domain CL of the first antigen binding moiety the amino acid at position 124 is substituted by a positively charged amino acid (numbering according to Kabat), and wherein in the constant domain CH1 of the first antigen binding moiety the amino acid at position 147 or the amino acid at position 213 is substituted by a negatively charged amino acid (numbering according to Kabat EU index); or

ii) in the constant domain CL of the second antigen binding moiety the amino acid at position 124 is substituted by a positively charged amino acid (numbering according to Kabat), and wherein in the constant domain CH1 of the second antigen binding moiety the amino acid at position 147 or the amino acid at position 213 is substituted by a negatively charged amino acid (numbering according to Kabat EU index).

The bispecific antibody does not comprise both modifications mentioned under i) and ii). The constant domains CL and CH1 of the antigen binding moiety having the VH/VL exchange are not replaced by each other (i.e. remain unexchanged).

In a more specific embodiment,

- 5 i) in the constant domain CL of the first antigen binding moiety the amino acid at position 124 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat), and in the constant domain CH1 of the first antigen binding moiety the amino acid at position 147 or the amino acid at position 213 is substituted independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index); or
- 10 ii) in the constant domain CL of the second antigen binding moiety the amino acid at position 124 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat), and in the constant domain CH1 of the second antigen binding moiety the amino acid at position 147 or the amino acid at position 213 is substituted independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index).
- 15

In one such embodiment, in the constant domain CL of the first antigen binding moiety the amino acid at position 124 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat), and in the constant domain CH1 of the first antigen binding moiety the amino acid at position 147 or the amino acid at position 213 is substituted independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index).

20

In a further embodiment, in the constant domain CL of the first antigen binding moiety the amino acid at position 124 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat), and in the constant domain CH1 of the first antigen binding moiety the amino acid at position 147 is substituted independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index).

25

In a particular embodiment, in the constant domain CL of the first antigen binding moiety the amino acid at position 124 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat) and the amino acid at position 123 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat), and in the constant domain CH1 of the first antigen binding moiety the amino acid at position 147 is substituted independently by glutamic acid (E), or aspartic acid (D)

30

35

(numbering according to Kabat EU index) and the amino acid at position 213 is substituted independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index).

5 In a more particular embodiment, in the constant domain CL of the first antigen binding moiety the amino acid at position 124 is substituted by lysine (K) (numbering according to Kabat) and the amino acid at position 123 is substituted by lysine (K) (numbering according to Kabat), and in the constant domain CH1 of the first antigen binding moiety the amino acid at position 147 is substituted by glutamic acid (E) (numbering according to Kabat EU index) and the amino acid at position 213 is substituted by glutamic acid (E) (numbering according to Kabat EU index).

15 In an even more particular embodiment, in the constant domain CL of the first antigen binding moiety the amino acid at position 124 is substituted by lysine (K) (numbering according to Kabat) and the amino acid at position 123 is substituted by arginine (R) (numbering according to Kabat), and in the constant domain CH1 of the first antigen binding moiety the amino acid at position 147 is substituted by glutamic acid (E) (numbering according to Kabat EU index) and the amino acid at position 213 is substituted by glutamic acid (E) (numbering according to Kabat EU index).

20 In particular embodiments, if amino acid substitutions according to the above embodiments are made in the constant domain CL and the constant domain CH1 of the first antigen binding moiety, the constant domain CL of the first antigen binding moiety is of kappa isotype.

25 Alternatively, the amino acid substitutions according to the above embodiments may be made in the constant domain CL and the constant domain CH1 of the second antigen binding moiety instead of in the constant domain CL and the constant domain CH1 of the first antigen binding moiety. In particular such embodiments, the constant domain CL of the second antigen binding moiety is of kappa isotype.

30 Accordingly, in one embodiment, in the constant domain CL of the second antigen binding moiety the amino acid at position 124 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat), and in the constant domain CH1 of the second antigen binding moiety the amino acid at position 147 or the amino acid at position 213 is substituted

independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index).

5 In a further embodiment, in the constant domain CL of the second antigen binding moiety the amino acid at position 124 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat), and in the constant domain CH1 of the second antigen binding moiety the amino acid at position 147 is substituted independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index).

10 In still another embodiment, in the constant domain CL of the second antigen binding moiety the amino acid at position 124 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat) and the amino acid at position 123 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat), and in the constant domain CH1 of the second antigen binding moiety the amino acid at position
15 147 is substituted independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index) and the amino acid at position 213 is substituted independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index).

20 In one embodiment, in the constant domain CL of the second antigen binding moiety the amino acid at position 124 is substituted by lysine (K) (numbering according to Kabat) and the amino acid at position 123 is substituted by lysine (K) (numbering according to Kabat), and in the constant domain CH1 of the second antigen binding moiety the amino acid at position 147 is substituted by glutamic acid (E) (numbering according to Kabat EU index) and the amino
25 acid at position 213 is substituted by glutamic acid (E) (numbering according to Kabat EU index).

30 In another embodiment, in the constant domain CL of the second antigen binding moiety the amino acid at position 124 is substituted by lysine (K) (numbering according to Kabat) and the amino acid at position 123 is substituted by arginine (R) (numbering according to Kabat), and in the constant domain CH1 of the second antigen binding moiety the amino acid at position 147 is substituted by glutamic acid (E) (numbering according to Kabat EU index) and the amino acid at position 213 is substituted by glutamic acid (E) (numbering according to Kabat EU index).

In a particular embodiment, the bispecific antibody of the invention comprises

I) a first antigen binding moiety that binds to human HLA-G, and the first antigen binding moiety is a Fab molecule comprising

5 A) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:32;24

B) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26;

and

II) a second antigen binding moiety that binds to human CD3,

10 wherein the second antigen binding moiety is a Fab molecule wherein the variable domains VL and VH of the Fab light chain and the Fab heavy chain are replaced by each other, comprising

C) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:58 and a VL domain comprising the amino acid sequence of SEQ ID NO:59, or

15 D) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:66 and a VL domain comprising the amino acid sequence of SEQ ID NO:67, or

E) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:74 and a VL domain comprising the amino acid sequence of SEQ ID NO:75;

and

20 III) wherein in the constant domain CL of the first antigen binding moiety the amino acid at position 124 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat) (in a particular embodiment independently by lysine (K) or arginine (R)) and the amino acid at position 123 is substituted independently by lysine (K), arginine (R) or
25 histidine (H) (numbering according to Kabat) (in a particular embodiment independently by lysine (K) or arginine (R)), and in the constant domain CH1 of the first antigen binding moiety the amino acid at position 147 is substituted independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index) and the amino acid at position 213 is substituted

independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index).

Bispecific antibody formats

5 The components of the bispecific antibody according to the present invention can be fused to each other in a variety of configurations. Exemplary configurations are depicted in **Figure 13**.

10 In particular embodiments, the antigen binding moieties comprised in the bispecific antibody are Fab molecules. In such embodiments, the first, second, third etc. antigen binding moiety may be referred to herein as first, second, third etc. Fab molecule, respectively.

15 In one embodiment, the first and the second antigen binding moiety of the bispecific antibody are fused to each other, optionally via a peptide linker. In particular embodiments, the first and the second antigen binding moiety are each a Fab molecule. In one such embodiment, the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety. In another such embodiment, the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety. In embodiments wherein either (i) the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety or (ii) the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety, additionally the Fab light chain of the first antigen binding moiety and the Fab light chain of the second antigen binding moiety may be fused to each other, optionally via a peptide linker.

25 A bispecific antibody with a single antigen binding moiety (such as a Fab molecule) capable of specific binding to a target cell antigen such as HLA-G (for example as shown in **Figure 13A, D, G, H, K, L**) is useful, particularly in cases where internalization of the target cell antigen is to be expected following binding of a high affinity antigen binding moiety. In such cases, the presence of more than one antigen binding moiety specific for the target cell antigen may enhance internalization of the target cell antigen, thereby reducing its availability.

30

In other cases, however, it will be advantageous to have a bispecific antibody comprising two or more antigen binding moieties (such as Fab molecules) specific for a target cell antigen (see examples shown in **Figure 13B, 13C, 13E, 13F, 13I, 13J, 13M or 13N**), for example to optimize targeting to the target site or to allow crosslinking of target cell antigens.

Accordingly, in particular embodiments, the bispecific antibody according to the present invention comprises a third antigen binding moiety.

In one embodiment, the third antigen binding moiety binds to the first antigen, i.e. HLA-G. In one embodiment, the third antigen binding moiety is a Fab molecule.

In particular embodiments, the third antigen moiety is identical to the first antigen binding moiety.

The third antigen binding moiety of the bispecific antibody may incorporate any of the features, singly or in combination, described herein in relation to the first antigen binding moiety and/or the antibody that binds HLA-G, unless scientifically clearly unreasonable or impossible.

In one embodiment, the third antigen binding moiety

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24; or

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26; or

In particular embodiments, the third and the first antigen binding moiety are each a Fab molecule and the third antigen binding moiety is identical to the first antigen binding moiety. Thus, in these embodiments the first and the third antigen binding moiety comprise the same heavy and light chain amino acid sequences and have the same arrangement of domains (i.e. conventional or crossover)). Furthermore, in these embodiments, the third antigen binding moiety comprises the same amino acid substitutions, if any, as the first antigen binding moiety. For example, the amino acid substitutions described herein as “charge modifications” will be made in the constant domain CL and the constant domain CH1 of each of the first antigen binding moiety and the third antigen binding moiety. Alternatively, said amino acid substitutions may be made in the constant domain CL and the constant domain CH1 of the second antigen binding moiety (which in particular embodiments is also a Fab molecule),

but not in the constant domain CL and the constant domain CH1 of the first antigen binding moiety and the third antigen binding moiety.

Like the first antigen binding moiety, the third antigen binding moiety particularly is a conventional Fab molecule. Embodiments wherein the first and the third antigen binding moieties are crossover Fab molecules (and the second antigen binding moiety is a conventional Fab molecule) are, however, also contemplated. Thus, in particular embodiments, the first and the third antigen binding moieties are each a conventional Fab molecule, and the second antigen binding moiety is a crossover Fab molecule as described herein, i.e. a Fab molecule wherein the variable domains VH and VL or the constant domains CL and CH1 of the Fab heavy and light chains are exchanged / replaced by each other. In other embodiments, the first and the third antigen binding moieties are each a crossover Fab molecule and the second antigen binding moiety is a conventional Fab molecule.

If a third antigen binding moiety is present, in a particular embodiment the first and the third antigen moiety bind to human HLA-G, and the second antigen binding moiety binds to a second antigen human CD3, most particularly CD3 epsilon.

In particular embodiments, the bispecific antibody comprises an Fc domain composed of a first and a second subunit. The first and the second subunit of the Fc domain are capable of stable association.

The bispecific antibody according to the invention can have different configurations, i.e. the first, second (and optionally third) antigen binding moiety may be fused to each other and to the Fc domain in different ways. The components may be fused to each other directly or, preferably, via one or more suitable peptide linkers. Where fusion of a Fab molecule is to the N-terminus of a subunit of the Fc domain, it is typically via an immunoglobulin hinge region.

In some embodiments, the first and the second antigen binding moiety are each a Fab molecule and the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain. In such embodiments, the first antigen binding moiety may be fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety or to the N-terminus of the other one of the subunits of the Fc domain. In particular such embodiments, said first antigen binding moiety is a conventional Fab molecule, and the second antigen binding moiety is a crossover Fab molecule as described herein, i.e. a Fab molecule wherein the variable domains

VH and VL or the constant domains CL and CH1 of the Fab heavy and light chains are exchanged / replaced by each other. In other such embodiments, said first Fab molecule is a crossover Fab molecule and the second Fab molecule is a conventional Fab molecule.

- 5 In one embodiment, the first and the second antigen binding moiety are each a Fab molecule, the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain, and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety. In a
10 specific embodiment, the bispecific antibody essentially consists of the first and the second Fab molecule, the Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the first Fab molecule is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second Fab molecule, and the second Fab molecule is fused at the C-terminus of the
15 Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain. Such a configuration is schematically depicted in **Figures 13G** and **13K** (with the second antigen binding domain in these examples being a VH/VL crossover Fab molecule). Optionally, the Fab light chain of the first Fab molecule and the Fab light chain of the second Fab molecule may additionally be fused to each other.
- 20 In another embodiment, the first and the second antigen binding moiety are each a Fab molecule and the first and the second antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of one of the subunits of the Fc domain. In a specific embodiment, the bispecific antibody essentially consists of the first and the second Fab molecule, the Fc domain composed of a first and a
25 second subunit, and optionally one or more peptide linkers, wherein the first and the second Fab molecule are each fused at the C-terminus of the Fab heavy chain to the N-terminus of one of the subunits of the Fc domain. Such a configuration is schematically depicted in **Figures 13A** and **13D** (in these examples with the second antigen binding domain being a VH/VL crossover Fab molecule and the first antigen
30 binding moiety being a conventional Fab molecule). The first and the second Fab molecule may be fused to the Fc domain directly or through a peptide linker. In a particular embodiment the first and the second Fab molecule are each fused to the Fc domain through an immunoglobulin hinge region. In a specific embodiment, the immunoglobulin hinge region is a human IgG₁ hinge region, particularly where the
35 Fc domain is an IgG₁ Fc domain.

In some embodiments, the first and the second antigen binding moiety are each a Fab molecule and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain. In such embodiments, the second antigen binding moiety may be fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety or (as described above) to the N-terminus of the other one of the subunits of the Fc domain. In particular such embodiments, said first antigen binding moiety is a conventional Fab molecule, and the second antigen binding moiety is a crossover Fab molecule as described herein, i.e. a Fab molecule wherein the variable domains VH and VL or the constant domains CL and CH1 of the Fab heavy and light chains are exchanged / replaced by each other. In other such embodiments, said first Fab molecule is a crossover Fab molecule and the second Fab molecule is a conventional Fab molecule.

In one embodiment, the first and the second antigen binding moiety are each a Fab molecule, the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain, and the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety. In a specific embodiment, the bispecific antibody essentially consists of the first and the second Fab molecule, the Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the second Fab molecule is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first Fab molecule, and the first Fab molecule is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain. Such a configuration is schematically depicted in **Figures 13H** and **13L** (in these examples with the second antigen binding domain being a VH/VL crossover Fab molecule and the first antigen binding moiety being a conventional Fab molecule). Optionally, the Fab light chain of the first Fab molecule and the Fab light chain of the second Fab molecule may additionally be fused to each other.

In some embodiments, a third antigen binding moiety, particularly a third Fab molecule, is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or second subunit of the Fc domain. In particular such embodiments, said first and third Fab molecules are each a conventional Fab molecule, and the second Fab molecule is a crossover Fab molecule as described herein, i.e. a Fab molecule wherein the variable domains VH and VL or the constant domains CL and CH1 of the Fab heavy and light chains are exchanged / replaced by each other. In other such

embodiments, said first and third Fab molecules are each a crossover Fab molecule and the second Fab molecule is a conventional Fab molecule.

In a particular such embodiment, the second and the third antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of one of the subunits of the Fc domain, and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second Fab molecule. In a specific embodiment, the bispecific antibody essentially consists of the first, the second and the third Fab molecule, the Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the first Fab molecule is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second Fab molecule, and the second Fab molecule is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain, and wherein the third Fab molecule is fused at the C-terminus of the Fab heavy chain to the N-terminus of the second subunit of the Fc domain. Such a configuration is schematically depicted in **Figure 13B** and **13E** (in these examples with the second antigen binding moiety being a VH/VL crossover Fab molecule, and the first and the third antigen binding moiety being a conventional Fab molecule), and **Figure 13J** and **13N** (in these examples with the second antigen binding moiety being a conventional Fab molecule, and the first and the third antigen binding moiety being a VH/VL crossover Fab molecule). The second and the third Fab molecule may be fused to the Fc domain directly or through a peptide linker. In a particular embodiment the second and the third Fab molecule are each fused to the Fc domain through an immunoglobulin hinge region. In a specific embodiment, the immunoglobulin hinge region is a human IgG₁ hinge region, particularly where the Fc domain is an IgG₁ Fc domain. Optionally, the Fab light chain of the first Fab molecule and the Fab light chain of the second Fab molecule may additionally be fused to each other.

In another such embodiment, the first and the third antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of one of the subunits of the Fc domain, and the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety. In a specific embodiment, the bispecific antibody essentially consists of the first, the second and the third Fab molecule, the Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the second Fab molecule is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first Fab molecule, and the first Fab molecule

is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain, and wherein the third Fab molecule is fused at the C-terminus of the Fab heavy chain to the N-terminus of the second subunit of the Fc domain. Such a configuration is schematically depicted in **Figure 13C** and **13F** (in these examples with the second antigen binding moiety being a VH/VL crossover Fab molecule, and the first and the third antigen binding moiety being a conventional Fab molecule) and in **Figure 13I** and **13M** (in these examples with the second antigen binding moiety being a conventional Fab molecule, and the first and the third antigen binding moiety being a VH/VL crossover Fab molecule). The first and the third Fab molecule may be fused to the Fc domain directly or through a peptide linker. In a particular embodiment the first and the third Fab molecule are each fused to the Fc domain through an immunoglobulin hinge region. In a specific embodiment, the immunoglobulin hinge region is a human IgG₁ hinge region, particularly where the Fc domain is an IgG₁ Fc domain. Optionally, the Fab light chain of the first Fab molecule and the Fab light chain of the second Fab molecule may additionally be fused to each other.

In configurations of the bispecific antibody wherein a Fab molecule is fused at the C-terminus of the Fab heavy chain to the N-terminus of each of the subunits of the Fc domain through an immunoglobulin hinge regions, the two Fab molecules, the hinge regions and the Fc domain essentially form an immunoglobulin molecule. In a particular embodiment the immunoglobulin molecule is an IgG class immunoglobulin. In an even more particular embodiment the immunoglobulin is an IgG₁ subclass immunoglobulin. In another embodiment the immunoglobulin is an IgG₄ subclass immunoglobulin. In a further particular embodiment the immunoglobulin is a human immunoglobulin. In other embodiments the immunoglobulin is a chimeric immunoglobulin or a humanized immunoglobulin. In one embodiment, the immunoglobulin comprises a human constant region, particularly a human Fc region.

In some of the bispecific antibody of the invention, the Fab light chain of the first Fab molecule and the Fab light chain of the second Fab molecule are fused to each other, optionally via a peptide linker. Depending on the configuration of the first and the second Fab molecule, the Fab light chain of the first Fab molecule may be fused at its C-terminus to the N-terminus of the Fab light chain of the second Fab molecule, or the Fab light chain of the second Fab molecule may be fused at its C-terminus to the N-terminus of the Fab light chain of the first Fab molecule. Fusion of the Fab light chains of the first and the second Fab molecule further reduces mispairing of

unmatched Fab heavy and light chains, and also reduces the number of plasmids needed for expression of some of the bispecific antibodies of the invention.

The antigen binding moieties may be fused to the Fc domain or to each other directly or through a peptide linker, comprising one or more amino acids, typically about 2-20 amino acids. Peptide linkers are known in the art and are described herein.

5 Suitable, non-immunogenic peptide linkers include, for example, $(G_4S)_n$, $(SG_4)_n$, $(G_4S)_n$ or $G_4(SG_4)_n$ peptide linkers. "n" is generally an integer from 1 to 10, typically from 2 to 4. In one embodiment said peptide linker has a length of at least 5 amino acids, in one embodiment a length of 5 to 100, in a further embodiment of 10 to 50

10 amino acids. In one embodiment said peptide linker is $(G_xS)_n$ or $(G_xS)_nG_m$ with G=glycine, S=serine, and $(x=3, n=3, 4, 5 \text{ or } 6, \text{ and } m=0, 1, 2 \text{ or } 3)$ or $(x=4, n=2, 3, 4 \text{ or } 5 \text{ and } m=0, 1, 2 \text{ or } 3)$, in one embodiment $x=4$ and $n=2$ or 3, in a further embodiment $x=4$ and $n=2$. In one embodiment said peptide linker is $(G_4S)_2$. A particularly suitable peptide linker for fusing the Fab light chains of the first and the

15 second Fab molecule to each other is $(G_4S)_2$. An exemplary peptide linker suitable for connecting the Fab heavy chains of the first and the second Fab fragments comprises the sequence $(D)-(G_4S)_2$. Another suitable such linker comprises the sequence $(G_4S)_4$. Additionally, linkers may comprise (a portion of) an immunoglobulin hinge region. Particularly where a Fab molecule is fused to the N-

20 terminus of an Fc domain subunit, it may be fused via an immunoglobulin hinge region or a portion thereof, with or without an additional peptide linker.

In certain embodiments the bispecific antibody according to the invention comprises a polypeptide wherein the Fab light chain variable region of the second Fab molecule shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of

25 the second Fab molecule (i.e. the second Fab molecule comprises a crossover Fab heavy chain, wherein the heavy chain variable region is replaced by a light chain variable region), which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit ($VL_{(2)}-CH1_{(2)}-CH2-CH3(-CH4)$), and a polypeptide wherein the Fab heavy chain of the first Fab molecule shares a carboxy-terminal peptide bond with

30 an Fc domain subunit ($VH_{(1)}-CH1_{(1)}-CH2-CH3(-CH4)$). In some embodiments the bispecific antibody further comprises a polypeptide wherein the Fab heavy chain variable region of the second Fab molecule shares a carboxy-terminal peptide bond with the Fab light chain constant region of the second Fab molecule ($VH_{(2)}-CL_{(2)}$) and the Fab light chain polypeptide of the first Fab molecule ($VL_{(1)}-CL_{(1)}$). In certain

35 embodiments the polypeptides are covalently linked, e.g., by a disulfide bond.

In certain embodiments the bispecific antibody according to the invention comprises a polypeptide wherein the Fab heavy chain variable region of the second Fab molecule shares a carboxy-terminal peptide bond with the Fab light chain constant region of the second Fab molecule (i.e. the second Fab molecule comprises a crossover Fab heavy chain, wherein the heavy chain constant region is replaced by a light chain constant region), which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₂₎-CL₍₂₎-CH2-CH3(-CH4)), and a polypeptide wherein the Fab heavy chain of the first Fab molecule shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₁₎-CH1₍₁₎-CH2-CH3(-CH4)). In some embodiments the bispecific antibody further comprises a polypeptide wherein the Fab light chain variable region of the second Fab molecule shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of the second Fab molecule (VL₍₂₎-CH1₍₂₎) and the Fab light chain polypeptide of the first Fab molecule (VL₍₁₎-CL₍₁₎). In certain embodiments the polypeptides are covalently linked, e.g., by a disulfide bond.

In some embodiments, the bispecific antibody comprises a polypeptide wherein the Fab light chain variable region of the second Fab molecule shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of the second Fab molecule (i.e. the second Fab molecule comprises a crossover Fab heavy chain, wherein the heavy chain variable region is replaced by a light chain variable region), which in turn shares a carboxy-terminal peptide bond with the Fab heavy chain of the first Fab molecule, which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit (VL₍₂₎-CH1₍₂₎-VH₍₁₎-CH1₍₁₎-CH2-CH3(-CH4)). In other embodiments, the bispecific antibody comprises a polypeptide wherein the Fab heavy chain of the first Fab molecule shares a carboxy-terminal peptide bond with the Fab light chain variable region of the second Fab molecule which in turn shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of the second Fab molecule (i.e. the second Fab molecule comprises a crossover Fab heavy chain, wherein the heavy chain variable region is replaced by a light chain variable region), which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₁₎-CH1₍₁₎-VL₍₂₎-CH1₍₂₎-CH2-CH3(-CH4)).

In some of these embodiments the bispecific antibody further comprises a crossover Fab light chain polypeptide of the second Fab molecule, wherein the Fab heavy chain variable region of the second Fab molecule shares a carboxy-terminal peptide bond with the Fab light chain constant region of the second Fab molecule (VH₍₂₎-CL₍₂₎), and the Fab light chain polypeptide of the first Fab molecule (VL₍₁₎-CL₍₁₎). In others

of these embodiments the bispecific antibody further comprises a polypeptide wherein the Fab heavy chain variable region of the second Fab molecule shares a carboxy-terminal peptide bond with the Fab light chain constant region of the second Fab molecule which in turn shares a carboxy-terminal peptide bond with the Fab light chain polypeptide of the first Fab molecule (VH₍₂₎-CL₍₂₎-VL₍₁₎-CL₍₁₎), or a polypeptide wherein the Fab light chain polypeptide of the first Fab molecule shares a carboxy-terminal peptide bond with the Fab heavy chain variable region of the second Fab molecule which in turn shares a carboxy-terminal peptide bond with the Fab light chain constant region of the second Fab molecule (VL₍₁₎-CL₍₁₎-VH₍₂₎-CL₍₂₎), as appropriate.

The bispecific antibody according to these embodiments may further comprise (i) an Fc domain subunit polypeptide (CH₂-CH₃(-CH₄)), or (ii) a polypeptide wherein the Fab heavy chain of a third Fab molecule shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₃₎-CH₁₍₃₎-CH₂-CH₃(-CH₄)) and the Fab light chain polypeptide of a third Fab molecule (VL₍₃₎-CL₍₃₎). In certain embodiments the polypeptides are covalently linked, e.g., by a disulfide bond.

In some embodiments, the bispecific antibody comprises a polypeptide wherein the Fab heavy chain variable region of the second Fab molecule shares a carboxy-terminal peptide bond with the Fab light chain constant region of the second Fab molecule (i.e. the second Fab molecule comprises a crossover Fab heavy chain, wherein the heavy chain constant region is replaced by a light chain constant region), which in turn shares a carboxy-terminal peptide bond with the Fab heavy chain of the first Fab molecule, which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₂₎-CL₍₂₎-VH₍₁₎-CH₁₍₁₎-CH₂-CH₃(-CH₄)). In other embodiments, the bispecific antibody comprises a polypeptide wherein the Fab heavy chain of the first Fab molecule shares a carboxy-terminal peptide bond with the Fab heavy chain variable region of the second Fab molecule which in turn shares a carboxy-terminal peptide bond with the Fab light chain constant region of the second Fab molecule (i.e. the second Fab molecule comprises a crossover Fab heavy chain, wherein the heavy chain constant region is replaced by a light chain constant region), which in turn shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₁₎-CH₁₍₁₎-VH₍₂₎-CL₍₂₎-CH₂-CH₃(-CH₄)).

In some of these embodiments the bispecific antibody further comprises a crossover Fab light chain polypeptide of the second Fab molecule, wherein the Fab light chain variable region of the second Fab molecule shares a carboxy-terminal peptide bond

with the Fab heavy chain constant region of the second Fab molecule (VL₍₂₎-CH1₍₂₎), and the Fab light chain polypeptide of the first Fab molecule (VL₍₁₎-CL₍₁₎). In others of these embodiments the bispecific antibody further comprises a polypeptide wherein the Fab light chain variable region of the second Fab molecule shares a carboxy-terminal peptide bond with the Fab heavy chain constant region of the second Fab molecule which in turn shares a carboxy-terminal peptide bond with the Fab light chain polypeptide of the first Fab molecule (VL₍₂₎-CH1₍₂₎-VL₍₁₎-CL₍₁₎), or a polypeptide wherein the Fab light chain polypeptide of the first Fab molecule shares a carboxy-terminal peptide bond with the Fab heavy chain variable region of the second Fab molecule which in turn shares a carboxy-terminal peptide bond with the Fab light chain constant region of the second Fab molecule (VL₍₁₎-CL₍₁₎-VH₍₂₎-CL₍₂₎), as appropriate.

The bispecific antibody according to these embodiments may further comprise (i) an Fc domain subunit polypeptide (CH2-CH3(-CH4)), or (ii) a polypeptide wherein the Fab heavy chain of a third Fab molecule shares a carboxy-terminal peptide bond with an Fc domain subunit (VH₍₃₎-CH1₍₃₎-CH2-CH3(-CH4)) and the Fab light chain polypeptide of a third Fab molecule (VL₍₃₎-CL₍₃₎). In certain embodiments the polypeptides are covalently linked, e.g., by a disulfide bond.

In all of the different configurations of the bispecific antibody according to the invention, the amino acid substitutions described herein, if present, may either be in the CH1 and CL domains of the first and (if present) the third antigen binding moiety/Fab molecule, or in the CH1 and CL domains of the second antigen binding moiety/Fab molecule. Preferably, they are in the CH1 and CL domains of the first and (if present) the third antigen binding moiety/Fab molecule. In accordance with the concept of the invention, if amino acid substitutions as described herein are made in the first (and, if present, the third) antigen binding moiety/Fab molecule, no such amino acid substitutions are made in the second antigen binding moiety/Fab molecule. Conversely, if amino acid substitutions as described herein are made in the second antigen binding moiety/Fab molecule, no such amino acid substitutions are made in the first (and, if present, the third) antigen binding moiety/Fab molecule. Amino acid substitutions are particularly made in bispecific antibodies comprising a Fab molecule wherein the variable domains VL and VH1 of the Fab light chain and the Fab heavy chain are replaced by each other.

In particular embodiments of the bispecific antibody according to the invention, particularly wherein amino acid substitutions as described herein are made in the

first (and, if present, the third) antigen binding moiety/Fab molecule, the constant domain CL of the first (and, if present, the third) Fab molecule is of kappa isotype. In other embodiments of the bispecific antibody according to the invention, particularly wherein amino acid substitutions as described herein are made in the
5 second antigen binding moiety/Fab molecule, the constant domain CL of the second antigen binding moiety/Fab molecule is of kappa isotype. In some embodiments, the constant domain CL of the first (and, if present, the third) antigen binding moiety/Fab molecule and the constant domain CL of the second antigen binding moiety/Fab molecule are of kappa isotype.

10 In a particular aspect, the invention provides a bispecific antibody comprising

a) a first and a third antigen binding moiety that binds to a first antigen; wherein the first antigen is HLA-G, and wherein the first and the second antigen binding moiety are each a (conventional) Fab molecule comprising a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 7 and a light chain variable
15 region comprising the amino acid sequence of SEQ ID NO: 24,

b) a second antigen binding moiety that binds to a second antigen; wherein the second antigen is CD3 and wherein the second antigen binding moiety is Fab molecule wherein the variable domains VL and VH of the Fab light chain and the Fab heavy chain are replaced by each other, comprising (i) a heavy chain variable
20 region comprising the amino acid sequence of SEQ ID NO: 58 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 59; or (ii) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 66 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 67; or (iii) a heavy chain variable region comprising the amino acid sequence of SEQ
25 ID NO: 74 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 75; and

c) an Fc domain composed of a first and a second subunit;

wherein

in the constant domain CL of the first and the third antigen binding moiety under a)
30 the amino acid at position 124 is substituted by lysine (K) (numbering according to Kabat) and the amino acid at position 123 is substituted by lysine (K) or arginine (R) (numbering according to Kabat) (most particularly by arginine (R)), and wherein in the constant domain CH1 of the first and the third

antigen binding moiety under a) the amino acid at position 147 is substituted by glutamic acid (E) (numbering according to Kabat EU index) and the amino acid at position 213 is substituted by glutamic acid (E) (numbering according to Kabat EU index);

5 and wherein further

the first antigen binding moiety under a) is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety under b), and the second antigen binding moiety under b) and the third antigen binding moiety under a) are each fused at the C-terminus of the Fab heavy chain to the N-terminus of one of the subunits of the Fc domain under c).

In one embodiment according to these aspects of the invention, in the first subunit of the Fc domain the threonine residue at position 366 is replaced with a tryptophan residue (T366W), and in the second subunit of the Fc domain the tyrosine residue at position 407 is replaced with a valine residue (Y407V) and optionally the threonine residue at position 366 is replaced with a serine residue (T366S) and the leucine residue at position 368 is replaced with an alanine residue (L368A) (numberings according to Kabat EU index).

In a further embodiment according to these aspects of the invention, in the first subunit of the Fc domain additionally the serine residue at position 354 is replaced with a cysteine residue (S354C) or the glutamic acid residue at position 356 is replaced with a cysteine residue (E356C) (particularly the serine residue at position 354 is replaced with a cysteine residue), and in the second subunit of the Fc domain additionally the tyrosine residue at position 349 is replaced by a cysteine residue (Y349C) (numberings according to Kabat EU index).

In still a further embodiment according to these aspects of the invention, in each of the first and the second subunit of the Fc domain the leucine residue at position 234 is replaced with an alanine residue (L234A), the leucine residue at position 235 is replaced with an alanine residue (L235A) and the proline residue at position 329 is replaced by a glycine residue (P329G) (numbering according to Kabat EU index).

In still a further embodiment according to these aspects of the invention, the Fc domain is a human IgG₁ Fc domain.

5 A specific embodiment of the invention is a bispecific antibody that binds to human HLA-G and to human CD3 wherein the antibody comprises a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 76, a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 77, a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 78, and a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 79 (wherein in the VH or VL framework regions or in the constant regions amino acids are substituted without affecting the specific binding properties and the properties of constant regions of such bispecific antibody)

10 A specific embodiment of the invention is a bispecific antibody that binds to human HLA-G and to human CD3 wherein the antibody comprises a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 76, a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 77, a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 78, and a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 79,

15 and wherein the bispecific antibody has one or more of the of the following properties:
the bispecific antibody shows

- 20 a) inhibition of ILT2 and/or ILT4 binding to HLA-G (see Example 10); and/or
- 25 b) antibody mediated IFN gamma secretion by T cells on SKOV3 cells transfected with recombinant HLA-G (SKOV3 HLA-G) and/or on JEG3 cells expressing endogenous HLA-G wherein the IFN gamma secretion was detected (by Luminex technology) (see Example 11); and or
- 30 c) T cell mediated cytotoxicity/tumor cell killing on SKOV3 cells transfected with recombinant HLA-G (SKOV 3HLA-G) and/or JEG3 cells expressing endogenous HLA-G wherein the cytotoxicity was detected by measuring Caspase 8 activation in cells after treatment with bispecific antibody (see Example 12); and/or

- 66 -

d) in vivo anti-tumor efficacy/ tumor regression in humanized NSG mice bearing SKOV3 human ovarian carcinoma transfected with recombinant HLA-G (SKOV3 HLA-G) humanized NSG mice (see Example 13); and/or

5 e) in vivo anti-tumor efficacy /tumor of HLA-G CD3 T cell bi-specific in humanized NSG mice bearing human breast cancer PDX tumors (BC004) (see Example 14).

In a further specific embodiment, the bispecific antibody comprises a polypeptide comprising the amino acid sequence of SEQ ID NO: 76, a polypeptide comprising the amino acid sequence of SEQ ID NO: 77, a polypeptide comprising the amino acid sequence of SEQ ID NO: 78 and a polypeptide comprising the amino acid
10 sequence of SEQ ID NO: 79.

A further specific embodiment of the invention is a bispecific antibody that binds to human HLA-G and to human CD3 wherein the antibody comprises a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 80, a polypeptide comprising an amino acid sequence that
15 is at least 98%, or 99% identical to the sequence of SEQ ID NO: 81, a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 82, and a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 83 (wherein in the VH or VL framework regions or in the constant regions amino acids are substituted without affecting the specific binding properties and the properties of
20 constant regions of such bispecific antibody)

A specific embodiment of the invention is a bispecific antibody that binds to human HLA-G and to human CD3 wherein the antibody comprises a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 80, a polypeptide comprising an amino acid sequence that
25 is at least 98%, or 99% identical to the sequence of SEQ ID NO: 81, a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 82, and a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 83,

30 and wherein the bispecific antibody has one or more of the of the following properties:

the bispecific antibody shows

a) inhibition of ILT2 and/or ILT4 binding to HLA-G (see Example 10); and/or

- 67 -

- b) antibody mediated IFN gamma secretion by T cells on SKOV3 cells transfected with recombinant HLA-G (SKOV3 HLA-G) and/or on JEG3 cells expressing endogenous HLA-G wherein the IFN gamma secretion was detected (by Luminex technology) (see Example 11); and or
- 5 c) T cell mediated cytotoxicity/tumor cell killing on SKOV3 cells transfected with recombinant HLA-G (SKOV3 HLA-G) and/or JEG3 cells expressing endogenous HLA-G wherein the cytotoxicity was detected by measuring Caspase 8 activation in cells after treatment with bispecific antibody (see Example 12); and/or
- 10 d) in vivo anti-tumor efficacy/ tumor regression in humanized NSG mice bearing SKOV3 human ovarian carcinoma transfected with recombinant HLA-G (SKOV3 HLA-G) humanized NSG mice (see Example 13); and/or
- e) in vivo anti-tumor efficacy /tumor of HLA-G CD3 T cell bi-specific in humanized NSG mice bearing human breast cancer PDX tumors (BC004) (see Example 14).
- 15 In a further specific embodiment, the bispecific antibody comprises a polypeptide comprising the amino acid sequence of SEQ ID NO: 80, a polypeptide comprising the amino acid sequence of SEQ ID NO: 81, a polypeptide comprising the amino acid sequence of SEQ ID NO: 82 and a polypeptide comprising the amino acid sequence of SEQ ID NO: 83.
- 20 A further specific embodiment of the invention is a bispecific antibody that binds to human HLA-G and to human CD3 wherein the antibody comprises a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 84, a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 85, a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 86, and a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 87 (wherein in the VH or VL framework regions or in the constant regions amino acids are substituted without affecting the specific binding properties and the properties of constant regions of such bispecific antibody)
- 25
- 30 A specific embodiment of the invention is a bispecific antibody that binds to human HLA-G and to human CD3 wherein the antibody comprises a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 84, a polypeptide comprising an amino acid sequence that

- 68 -

is at least 98%, or 99% identical to the sequence of SEQ ID NO: 85, a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 86, and a polypeptide comprising an amino acid sequence that is at least 98%, or 99% identical to the sequence of SEQ ID NO: 87,

5 and wherein the bispecific antibody has one or more of the of the following properties:

the bispecific antibody shows

a) inhibition of ILT2 and/or ILT4 binding to HLA-G (see Example 10); and/or

10 b) antibody mediated IFN gamma secretion by T cells on SKOV3 cells transfected with recombinant HLA-G (SKOV3 HLA-G) and/or on JEG3 cells expressing endogenous HLA-G wherein the IFN gamma secretion was detected (by Luminex technology) (see Example 11); and or

15 c) T cell mediated cytotoxicity/tumor cell killing on SKOV3 cells transfected with recombinant HLA-G (SKOV 3HLA-G) and/or JEG3 cells expressing endogenous HLA-G wherein the cytotoxicity was detected by measuring Caspase 8 activation in cells after treatment with bispecific antibody (see Example 12); and/or

d) in vivo anti-tumor efficacy/ tumor regression in humanized NSG mice bearing SKOV3 human ovarian carcinoma transfected with recombinant HLA-G (SKOV3 HLA-G) humanized NSG mice (see Example 13); and/or

20 e) in vivo anti-tumor efficacy /tumor of HLA-G CD3 T cell bi-specific in humanized NSG mice bearing human breast cancer PDX tumors (BC004) (see Example 14).

25 In a further specific embodiment, the bispecific antibody comprises a polypeptide comprising the amino acid sequence of SEQ ID NO: 84, a polypeptide comprising the amino acid sequence of SEQ ID NO: 85, a polypeptide comprising the amino acid sequence of SEQ ID NO: 86 and a polypeptide comprising the amino acid sequence of SEQ ID NO: 87.

Fc domain

30 In particular embodiments, the bispecific antibody of the invention comprises an Fc domain composed of a first and a second subunit. It is understood, that the features of the Fc domain described herein in relation to the bispecific antibody can equally

apply to an Fc domain comprised in a monospecific anti-HLAG antibody of the invention except for those modifications relevant for Fc heterodimerization.

5 The Fc domain of the bispecific antibody consists of a pair of polypeptide chains comprising heavy chain domains of an immunoglobulin molecule. For example, the Fc domain of an immunoglobulin G (IgG) molecule is a dimer, each subunit of which comprises the CH2 and CH3 IgG heavy chain constant domains. The two subunits of the Fc domain are capable of stable association with each other. In one embodiment, the bispecific antibody of the invention comprises not more than one Fc domain.

10 In one embodiment, the Fc domain of the bispecific antibody is an IgG Fc domain. In a particular embodiment, the Fc domain is an IgG₁ Fc domain. In another embodiment the Fc domain is an IgG₄ Fc domain. In a more specific embodiment, the Fc domain is an IgG₄ Fc domain comprising an amino acid substitution at position S228 (Kabat EU index numbering), particularly the amino acid substitution
15 S228P. This amino acid substitution reduces *in vivo* Fab arm exchange of IgG₄ antibodies (see Stubenrauch et al., Drug Metabolism and Disposition 38, 84-91 (2010)). In a further particular embodiment, the Fc domain is a human Fc domain. In an even more particular embodiment, the Fc domain is a human IgG₁ Fc domain.

Fc domain modifications promoting heterodimerization

20 Bispecific antibodies according to the invention comprise different antigen binding moieties, which may be fused to one or the other of the two subunits of the Fc domain, thus the two subunits of the Fc domain are typically comprised in two non-identical polypeptide chains. Recombinant co-expression of these polypeptides and subsequent dimerization leads to several possible combinations of the two
25 polypeptides. To improve the yield and purity of bispecific antibodies in recombinant production, it will thus be advantageous to introduce in the Fc domain of the bispecific antibody a modification promoting the association of the desired polypeptides.

30 Accordingly, in particular embodiments, the Fc domain of the bispecific antibody according to the invention comprises a modification promoting the association of the first and the second subunit of the Fc domain. The site of most extensive protein-protein interaction between the two subunits of a human IgG Fc domain is in the CH3 domain of the Fc domain. Thus, in one embodiment said modification is in the CH3 domain of the Fc domain.

There exist several approaches for modifications in the CH3 domain of the Fc domain in order to enforce heterodimerization, which are well described e.g. in WO 96/27011, WO 98/050431, EP 1870459, WO 2007/110205, WO 2007/147901, WO 2009/089004, WO 2010/129304, WO 2011/90754, WO 2011/143545, WO 2012058768, WO 2013157954, WO 2013096291. Typically, in all such approaches the CH3 domain of the first subunit of the Fc domain and the CH3 domain of the second subunit of the Fc domain are both engineered in a complementary manner so that each CH3 domain (or the heavy chain comprising it) can no longer homodimerize with itself but is forced to heterodimerize with the complementarily engineered other CH3 domain (so that the first and second CH3 domain heterodimerize and no homodimers between the two first or the two second CH3 domains are formed). These different approaches for improved heavy chain heterodimerization are contemplated as different alternatives in combination with the heavy-light chain modifications (e.g. VH and VL exchange/replacement in one binding arm and the introduction of substitutions of charged amino acids with opposite charges in the CH1/CL interface) in the bispecific antibody which reduce heavy/light chain mispairing and Bence Jones-type side products.

In a specific embodiment said modification promoting the association of the first and the second subunit of the Fc domain is a so-called “knob-into-hole” modification, comprising a “knob” modification in one of the two subunits of the Fc domain and a “hole” modification in the other one of the two subunits of the Fc domain.

The knob-into-hole technology is described e.g. in US 5,731,168; US 7,695,936; Ridgway et al., Prot Eng 9, 617-621 (1996) and Carter, J Immunol Meth 248, 7-15 (2001). Generally, the method involves introducing a protuberance (“knob”) at the interface of a first polypeptide and a corresponding cavity (“hole”) in the interface of a second polypeptide, such that the protuberance can be positioned in the cavity so as to promote heterodimer formation and hinder homodimer formation. Protuberances are constructed by replacing small amino acid side chains from the interface of the first polypeptide with larger side chains (e.g. tyrosine or tryptophan). Compensatory cavities of identical or similar size to the protuberances are created in the interface of the second polypeptide by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine).

Accordingly, in a particular embodiment, in the CH3 domain of the first subunit of the Fc domain of the bispecific antibody an amino acid residue is replaced with an amino acid residue having a larger side chain volume, thereby generating a

protuberance within the CH3 domain of the first subunit which is positionable in a cavity within the CH3 domain of the second subunit, and in the CH3 domain of the second subunit of the Fc domain an amino acid residue is replaced with an amino acid residue having a smaller side chain volume, thereby generating a cavity within the CH3 domain of the second subunit within which the protuberance within the CH3 domain of the first subunit is positionable.

Preferably said amino acid residue having a larger side chain volume is selected from the group consisting of arginine (R), phenylalanine (F), tyrosine (Y), and tryptophan (W).

Preferably said amino acid residue having a smaller side chain volume is selected from the group consisting of alanine (A), serine (S), threonine (T), and valine (V).

The protuberance and cavity can be made by altering the nucleic acid encoding the polypeptides, e.g. by site-specific mutagenesis, or by peptide synthesis.

In a specific embodiment, in (the CH3 domain of) the first subunit of the Fc domain (the “knobs” subunit) the threonine residue at position 366 is replaced with a tryptophan residue (T366W), and in (the CH3 domain of) the second subunit of the Fc domain (the “hole” subunit) the tyrosine residue at position 407 is replaced with a valine residue (Y407V). In one embodiment, in the second subunit of the Fc domain additionally the threonine residue at position 366 is replaced with a serine residue (T366S) and the leucine residue at position 368 is replaced with an alanine residue (L368A) (numberings according to Kabat EU index).

In yet a further embodiment, in the first subunit of the Fc domain additionally the serine residue at position 354 is replaced with a cysteine residue (S354C) or the glutamic acid residue at position 356 is replaced with a cysteine residue (E356C) (particularly the serine residue at position 354 is replaced with a cysteine residue), and in the second subunit of the Fc domain additionally the tyrosine residue at position 349 is replaced by a cysteine residue (Y349C) (numberings according to Kabat EU index). Introduction of these two cysteine residues results in formation of a disulfide bridge between the two subunits of the Fc domain, further stabilizing the dimer (Carter, J Immunol Methods 248, 7-15 (2001)).

In a particular embodiment, the first subunit of the Fc domain comprises the amino acid substitutions S354C and T366W, and the second subunit of the Fc domain

comprises the amino acid substitutions Y349C, T366S, L368A and Y407V (numbering according to Kabat EU index).

5 In a particular embodiment the antigen binding moiety that binds to the second antigen (e.g. an activating T cell antigen) is fused (optionally via the first antigen binding moiety, which binds to HLA-G, and/or a peptide linker) to the first subunit of the Fc domain (comprising the “knob” modification). Without wishing to be bound by theory, fusion of the antigen binding moiety that binds a second antigen, such as an activating T cell antigen, to the knob-containing subunit of the Fc domain will (further) minimize the generation of antibodies comprising two antigen binding
10 moieties that bind to an activating T cell antigen (steric clash of two knob-containing polypeptides).

Other techniques of CH3-modification for enforcing the heterodimerization are contemplated as alternatives according to the invention and are described e.g. in WO 96/27011, WO 98/050431, EP 1870459, WO 2007/110205, WO 2007/147901,
15 WO 2009/089004, WO 2010/129304, WO 2011/90754, WO 2011/143545, WO 2012/058768, WO 2013/157954, WO 2013/096291.

In one embodiment, the heterodimerization approach described in EP 1870459, is used alternatively. This approach is based on the introduction of charged amino acids with opposite charges at specific amino acid positions in the CH3/CH3 domain interface between the two subunits of the Fc domain. One preferred embodiment for
20 the bispecific antibody of the invention are amino acid mutations R409D; K370E in one of the two CH3 domains (of the Fc domain) and amino acid mutations D399K; E357K in the other one of the CH3 domains of the Fc domain (numbering according to Kabat EU index).

25 In another embodiment, the bispecific antibody of the invention comprises amino acid mutation T366W in the CH3 domain of the first subunit of the Fc domain and amino acid mutations T366S, L368A, Y407V in the CH3 domain of the second subunit of the Fc domain, and additionally amino acid mutations R409D; K370E in the CH3 domain of the first subunit of the Fc domain and amino acid mutations
30 D399K; E357K in the CH3 domain of the second subunit of the Fc domain (numberings according to Kabat EU index).

In another embodiment, the bispecific antibody of the invention comprises amino acid mutations S354C, T366W in the CH3 domain of the first subunit of the Fc domain and amino acid mutations Y349C, T366S, L368A, Y407V in the CH3

domain of the second subunit of the Fc domain, or said bispecific antibody comprises amino acid mutations Y349C, T366W in the CH3 domain of the first subunit of the Fc domain and amino acid mutations S354C, T366S, L368A, Y407V in the CH3 domains of the second subunit of the Fc domain and additionally amino acid mutations R409D; K370E in the CH3 domain of the first subunit of the Fc domain and amino acid mutations D399K; E357K in the CH3 domain of the second subunit of the Fc domain (all numberings according to Kabat EU index).

In one embodiment, the heterodimerization approach described in WO 2013/157953 is used alternatively. In one embodiment, a first CH3 domain comprises amino acid mutation T366K and a second CH3 domain comprises amino acid mutation L351D (numberings according to Kabat EU index). In a further embodiment, the first CH3 domain comprises further amino acid mutation L351K. In a further embodiment, the second CH3 domain comprises further an amino acid mutation selected from Y349E, Y349D and L368E (preferably L368E) (numberings according to Kabat EU index).

In one embodiment, the heterodimerization approach described in WO 2012/058768 is used alternatively. In one embodiment a first CH3 domain comprises amino acid mutations L351Y, Y407A and a second CH3 domain comprises amino acid mutations T366A, K409F. In a further embodiment the second CH3 domain comprises a further amino acid mutation at position T411, D399, S400, F405, N390, or K392, e.g. selected from a) T411N, T411R, T411Q, T411K, T411D, T411E or T411W, b) D399R, D399W, D399Y or D399K, c) S400E, S400D, S400R, or S400K, d) F405I, F405M, F405T, F405S, F405V or F405W, e) N390R, N390K or N390D, f) K392V, K392M, K392R, K392L, K392F or K392E (numberings according to Kabat EU index). In a further embodiment a first CH3 domain comprises amino acid mutations L351Y, Y407A and a second CH3 domain comprises amino acid mutations T366V, K409F. In a further embodiment, a first CH3 domain comprises amino acid mutation Y407A and a second CH3 domain comprises amino acid mutations T366A, K409F. In a further embodiment, the second CH3 domain further comprises amino acid mutations K392E, T411E, D399R and S400R (numberings according to Kabat EU index).

In one embodiment, the heterodimerization approach described in WO 2011/143545 is used alternatively, e.g. with the amino acid modification at a position selected from the group consisting of 368 and 409 (numbering according to Kabat EU index).

In one embodiment, the heterodimerization approach described in WO 2011/090762, which also uses the knobs-into-holes technology described above, is used

alternatively. In one embodiment a first CH3 domain comprises amino acid mutation T366W and a second CH3 domain comprises amino acid mutation Y407A. In one embodiment, a first CH3 domain comprises amino acid mutation T366Y and a second CH3 domain comprises amino acid mutation Y407T (numberings according to Kabat EU index).

In one embodiment, the bispecific antibody or its Fc domain is of IgG₂ subclass and the heterodimerization approach described in WO 2010/129304 is used alternatively.

In an alternative embodiment, a modification promoting association of the first and the second subunit of the Fc domain comprises a modification mediating electrostatic steering effects, e.g. as described in PCT publication WO 2009/089004. Generally, this method involves replacement of one or more amino acid residues at the interface of the two Fc domain subunits by charged amino acid residues so that homodimer formation becomes electrostatically unfavorable but heterodimerization electrostatically favorable. In one such embodiment, a first CH3 domain comprises amino acid substitution of K392 or N392 with a negatively charged amino acid (e.g. glutamic acid (E), or aspartic acid (D), preferably K392D or N392D) and a second CH3 domain comprises amino acid substitution of D399, E356, D356, or E357 with a positively charged amino acid (e.g. lysine (K) or arginine (R), preferably D399K, E356K, D356K, or E357K, and more preferably D399K and E356K). In a further embodiment, the first CH3 domain further comprises amino acid substitution of K409 or R409 with a negatively charged amino acid (e.g. glutamic acid (E), or aspartic acid (D), preferably K409D or R409D). In a further embodiment the first CH3 domain further or alternatively comprises amino acid substitution of K439 and/or K370 with a negatively charged amino acid (e.g. glutamic acid (E), or aspartic acid (D)) (all numberings according to Kabat EU index).

In yet a further embodiment, the heterodimerization approach described in WO 2007/147901 is used alternatively. In one embodiment, a first CH3 domain comprises amino acid mutations K253E, D282K, and K322D and a second CH3 domain comprises amino acid mutations D239K, E240K, and K292D (numberings according to Kabat EU index).

In still another embodiment, the heterodimerization approach described in WO 2007/110205 can be used alternatively.

In one embodiment, the first subunit of the Fc domain comprises amino acid substitutions K392D and K409D, and the second subunit of the Fc domain comprises

amino acid substitutions D356K and D399K (numbering according to Kabat EU index).

Fc domain modifications reducing Fc receptor binding and/or effector function

5 The Fc domain confers to the bispecific antibody (or the antibody) favorable pharmacokinetic properties, including a long serum half-life which contributes to good accumulation in the target tissue and a favorable tissue-blood distribution ratio. At the same time it may, however, lead to undesirable targeting of the bispecific antibody (or the antibody) to cells expressing Fc receptors rather than to the preferred antigen-bearing cells. Moreover, the co-activation of Fc receptor signaling pathways
10 may lead to cytokine secretion/release which, in combination with the T cell activating properties (e.g. in embodiments of the bispecific antibody wherein the second antigen binding moiety binds to an activating T cell antigen) and the long half-life of the bispecific antibody, results in excessive activation of cytokine receptors and severe side effects upon systemic administration. Activation of (Fc
15 receptor-bearing) immune cells other than T cells may even reduce efficacy of the bispecific antibody (particularly a bispecific antibody wherein the second antigen binding moiety binds to an activating T cell antigen) due to the potential destruction of T cells e.g. by NK cells.

Accordingly, in particular embodiments, the Fc domain of the bispecific antibody
20 according to the invention exhibits reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a native IgG₁ Fc domain. In one such embodiment the Fc domain (or the bispecific antibody comprising said Fc domain) exhibits less than 50%, preferably less than 20%, more preferably less than 10% and most preferably less than 5% of the binding affinity to an Fc receptor, as compared
25 to a native IgG₁ Fc domain (or a bispecific antibody comprising a native IgG₁ Fc domain), and/or less than 50%, preferably less than 20%, more preferably less than 10% and most preferably less than 5% of the effector function, as compared to a native IgG₁ Fc domain (or a bispecific antibody comprising a native IgG₁ Fc domain). In one embodiment, the Fc domain (or the bispecific antibody
30 comprising said Fc domain) does not substantially bind to an Fc receptor and/or induce effector function. In a particular embodiment the Fc receptor is an Fcγ receptor. In one embodiment the Fc receptor is a human Fc receptor. In one embodiment the Fc receptor is an activating Fc receptor. In a specific embodiment the Fc receptor is an activating human Fcγ receptor, more specifically human
35 FcγRIIIa, FcγRI or FcγRIIa, most specifically human FcγRIIIa. In one embodiment

the effector function is one or more selected from the group of CDC, ADCC, ADCCP, and cytokine secretion. In a particular embodiment, the effector function is ADCC. In one embodiment, the Fc domain domain exhibits substantially similar binding affinity to neonatal Fc receptor (FcRn), as compared to a native IgG₁ Fc domain domain. Substantially similar binding to FcRn is achieved when the Fc domain (or the bispecific antibody comprising said Fc domain) exhibits greater than about 70%, particularly greater than about 80%, more particularly greater than about 90% of the binding affinity of a native IgG₁ Fc domain (or the bispecific antibody comprising a native IgG₁ Fc domain) to FcRn.

10 In certain embodiments the Fc domain is engineered to have reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a non-engineered Fc domain. In particular embodiments, the Fc domain of the bispecific antibody comprises one or more amino acid mutation that reduces the binding affinity of the Fc domain to an Fc receptor and/or effector function. Typically, the same one or

15 more amino acid mutation is present in each of the two subunits of the Fc domain. In one embodiment, the amino acid mutation reduces the binding affinity of the Fc domain to an Fc receptor. In one embodiment, the amino acid mutation reduces the binding affinity of the Fc domain to an Fc receptor by at least 2-fold, at least 5-fold, or at least 10-fold. In embodiments where there is more than one amino acid mutation

20 that reduces the binding affinity of the Fc domain to the Fc receptor, the combination of these amino acid mutations may reduce the binding affinity of the Fc domain to an Fc receptor by at least 10-fold, at least 20-fold, or even at least 50-fold. In one embodiment the bispecific antibody comprising an engineered Fc domain exhibits less than 20%, particularly less than 10%, more particularly less than 5% of the binding affinity to an Fc receptor as compared to a bispecific antibody comprising a non-engineered Fc domain. In a particular embodiment, the Fc receptor is an Fcγ receptor. In some embodiments, the Fc receptor is a human Fc receptor. In some

25 embodiments, the Fc receptor is an activating Fc receptor. In a specific embodiment, the Fc receptor is an activating human Fcγ receptor, more specifically human FcγRIIIa, FcγRI or FcγRIIa, most specifically human FcγRIIIa. Preferably, binding to each of these receptors is reduced. In some embodiments, binding affinity to a complement component, specifically binding affinity to C1q, is also reduced. In one embodiment, binding affinity to neonatal Fc receptor (FcRn) is not reduced. Substantially similar binding to FcRn, i.e. preservation of the binding affinity of the Fc domain to said receptor, is achieved when the Fc domain (or the bispecific antibody comprising said Fc domain) exhibits greater than about 70% of the binding affinity of a non-engineered form of the Fc domain (or the bispecific antibody

30

35

comprising said non-engineered form of the Fc domain) to FcRn. The Fc domain, or bispecific antibodies of the invention comprising said Fc domain, may exhibit greater than about 80% and even greater than about 90% of such affinity. In certain embodiments, the Fc domain of the bispecific antibody is engineered to have reduced effector function, as compared to a non-engineered Fc domain. The reduced effector function can include, but is not limited to, one or more of the following: reduced complement dependent cytotoxicity (CDC), reduced antibody-dependent cell-mediated cytotoxicity (ADCC), reduced antibody-dependent cellular phagocytosis (ADCP), reduced cytokine secretion, reduced immune complex-mediated antigen uptake by antigen-presenting cells, reduced binding to NK cells, reduced binding to macrophages, reduced binding to monocytes, reduced binding to polymorphonuclear cells, reduced direct signaling inducing apoptosis, reduced crosslinking of target-bound antibodies, reduced dendritic cell maturation, or reduced T cell priming. In one embodiment, the reduced effector function is one or more selected from the group of reduced CDC, reduced ADCC, reduced ADCP, and reduced cytokine secretion. In a particular embodiment, the reduced effector function is reduced ADCC. In one embodiment the reduced ADCC is less than 20% of the ADCC induced by a non-engineered Fc domain (or a bispecific antibody comprising a non-engineered Fc domain).

In one embodiment, the amino acid mutation that reduces the binding affinity of the Fc domain to an Fc receptor and/or effector function is an amino acid substitution. In one embodiment, the Fc domain comprises an amino acid substitution at a position selected from the group of E233, L234, L235, N297, P331 and P329 (numberings according to Kabat EU index). In a more specific embodiment, the Fc domain comprises an amino acid substitution at a position selected from the group of L234, L235 and P329 (numberings according to Kabat EU index). In some embodiments, the Fc domain comprises the amino acid substitutions L234A and L235A (numberings according to Kabat EU index). In one such embodiment, the Fc domain is an IgG₁ Fc domain, particularly a human IgG₁ Fc domain. In one embodiment, the Fc domain comprises an amino acid substitution at position P329. In a more specific embodiment, the amino acid substitution is P329A or P329G, particularly P329G (numberings according to Kabat EU index). In one embodiment, the Fc domain comprises an amino acid substitution at position P329 and a further amino acid substitution at a position selected from E233, L234, L235, N297 and P331 (numberings according to Kabat EU index). In a more specific embodiment, the further amino acid substitution is E233P, L234A, L235A, L235E, N297A, N297D or P331S. In particular embodiments, the Fc domain comprises amino acid

substitutions at positions P329, L234 and L235 (numberings according to Kabat EU index). In more particular embodiments, the Fc domain comprises the amino acid mutations L234A, L235A and P329G ("P329G LALA", "PGLALA" or "LALAPG"). Specifically, in particular embodiments, each subunit of the Fc domain comprises the amino acid substitutions L234A, L235A and P329G (Kabat EU index numbering), i.e. in each of the first and the second subunit of the Fc domain the leucine residue at position 234 is replaced with an alanine residue (L234A), the leucine residue at position 235 is replaced with an alanine residue (L235A) and the proline residue at position 329 is replaced by a glycine residue (P329G) (numbering according to Kabat EU index).

In one such embodiment, the Fc domain is an IgG₁ Fc domain, particularly a human IgG₁ Fc domain. The "P329G LALA" combination of amino acid substitutions almost completely abolishes Fcγ receptor (as well as complement) binding of a human IgG₁ Fc domain, as described in PCT publication no. WO 2012/130831, which is incorporated herein by reference in its entirety. WO 2012/130831 also describes methods of preparing such mutant Fc domains and methods for determining its properties such as Fc receptor binding or effector functions.

IgG₄ antibodies exhibit reduced binding affinity to Fc receptors and reduced effector functions as compared to IgG₁ antibodies. Hence, in some embodiments, the Fc domain of the bispecific antibodies of the invention is an IgG₄ Fc domain, particularly a human IgG₄ Fc domain. In one embodiment, the IgG₄ Fc domain comprises amino acid substitutions at position S228, specifically the amino acid substitution S228P (numberings according to Kabat EU index). To further reduce its binding affinity to an Fc receptor and/or its effector function, in one embodiment, the IgG₄ Fc domain comprises an amino acid substitution at position L235, specifically the amino acid substitution L235E (numberings according to Kabat EU index). In another embodiment, the IgG₄ Fc domain comprises an amino acid substitution at position P329, specifically the amino acid substitution P329G (numberings according to Kabat EU index). In a particular embodiment, the IgG₄ Fc domain comprises amino acid substitutions at positions S228, L235 and P329, specifically amino acid substitutions S228P, L235E and P329G (numberings according to Kabat EU index). Such IgG₄ Fc domain mutants and their Fcγ receptor binding properties are described in PCT publication no. WO 2012/130831, incorporated herein by reference in its entirety.

In a particular embodiment, the Fc domain exhibiting reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a native IgG₁ Fc domain, is a human IgG₁ Fc domain comprising the amino acid substitutions L234A, L235A and optionally P329G, or a human IgG₄ Fc domain comprising the amino acid substitutions S228P, L235E and optionally P329G (numberings according to Kabat EU index).

Mutant Fc domains can be prepared by amino acid deletion, substitution, insertion or modification using genetic or chemical methods well known in the art. Genetic methods may include site-specific mutagenesis of the encoding DNA sequence, PCR, gene synthesis, and the like. The correct nucleotide changes can be verified for example by sequencing.

Binding to Fc receptors can be easily determined e.g. by ELISA, or by Surface Plasmon Resonance (SPR) using standard instrumentation such as a BIAcore instrument (GE Healthcare), and Fc receptors such as may be obtained by recombinant expression. Alternatively, binding affinity of Fc domains or bispecific antibodies comprising an Fc domain for Fc receptors may be evaluated using cell lines known to express particular Fc receptors, such as human NK cells expressing FcγIIIa receptor.

Effector function of an Fc domain, or a bispecific antibody comprising an Fc domain, can be measured by methods known in the art. Examples of *in vitro* assays to assess ADCC activity of a molecule of interest are described in U.S. Patent No. 5,500,362; Hellstrom et al. Proc Natl Acad Sci USA 83, 7059-7063 (1986) and Hellstrom et al., Proc Natl Acad Sci USA 82, 1499-1502 (1985); U.S. Patent No. 5,821,337; Bruggemann et al., J Exp Med 166, 1351-1361 (1987). Alternatively, non-radioactive assays methods may be employed (see, for example, ACTIT[™] non-radioactive cytotoxicity assay for flow cytometry (CellTechnology, Inc. Mountain View, CA); and CytoTox 96[®] non-radioactive cytotoxicity assay (Promega, Madison, WI)). Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, e.g. in a animal model such as that disclosed in Clynes et al., Proc Natl Acad Sci USA 95, 652-656 (1998).

In some embodiments, binding of the Fc domain to a complement component, specifically to C1q, is reduced. Accordingly, in some embodiments wherein the Fc domain is engineered to have reduced effector function, said reduced effector

function includes reduced CDC. C1q binding assays may be carried out to determine whether the Fc domain, or the bispecific antibody comprising the Fc domain, is able to bind C1q and hence has CDC activity. See e.g., C1q and C3c binding ELISA in WO 2006/029879 and WO 2005/100402. To assess complement activation, a CDC assay may be performed (see, for example, Gazzano-Santoro et al., J Immunol Methods 202, 163 (1996); Cragg et al., Blood 101, 1045-1052 (2003); and Cragg and Glennie, Blood 103, 2738-2743 (2004)).

FcRn binding and *in vivo* clearance/half life determinations can also be performed using methods known in the art (see, e.g., Petkova, S.B. et al., *Int'l. Immunol.* 18(12):1759-1769 (2006); WO 2013/120929).

In a further aspect, an anti-HLA-G antibody according to any of the above embodiments may incorporate any of the features, singly or in combination, as described in Sections 1-6 below:

1. Antibody Affinity

In certain embodiments, an antibody provided herein has a dissociation constant KD of $\leq 1 \mu\text{M}$, $\leq 100 \text{ nM}$, $\leq 10 \text{ nM}$, $\leq 1 \text{ nM}$, $\leq 0.1 \text{ nM}$, $\leq 0.01 \text{ nM}$, or $\leq 0.001 \text{ nM}$ (e.g. 10^{-8} M or less, e.g. from 10^{-8} M to 10^{-13} M , e.g., from 10^{-9} M to 10^{-13} M).

In one preferred embodiment, KD is measured using surface plasmon resonance assays using a BIACORE[®] at 25°C with immobilized antigen CM5 chips at ~10 response units (RU). Briefly, carboxymethylated dextran biosensor chips (CM5, BIACORE, Inc.) are activated with *N*-ethyl-*N'*-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) and *N*-hydroxysuccinimide (NHS) according to the supplier's instructions. Antigen is diluted with 10 mM sodium acetate, pH 4.8, to 5 $\mu\text{g/ml}$ (~0.2 μM) before injection at a flow rate of 5 $\mu\text{l/minute}$ to achieve approximately 10 response units (RU) of coupled protein. Following the injection of antigen, 1 M ethanolamine is injected to block unreacted groups. For kinetics measurements, two-fold serial dilutions of Fab (0.78 nM to 500 nM) are injected in PBS with 0.05% polysorbate 20 (TWEEN-20TM) surfactant (PBST) at 25°C at a flow rate of approximately 25 $\mu\text{l/min}$. Association rates (k_{ON} or k_a) and dissociation rates (k_{OFF} or k_d) are calculated using a simple one-to-one Langmuir binding model (BIACORE[®] Evaluation Software version 3.2) by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant KD is calculated as the ratio k_d/k_a ($k_{\text{OFF}}/k_{\text{ON}}$). See, e.g., Chen, Y. et al., J. Mol. Biol. 293 (1999) 865-881. If the on-rate exceeds $10^6 \text{ M}^{-1} \text{ s}^{-1}$ by the surface plasmon resonance

assay above, then the on-rate can be determined by using a fluorescent quenching technique that measures the increase or decrease in fluorescence emission intensity (excitation = 295 nm; emission = 340 nm, 16 nm band-pass) at 25°C of a 20 nM anti-antigen antibody (Fab form) in PBS, pH 7.2, in the presence of increasing concentrations of antigen as measured in a spectrometer, such as a stop-flow equipped spectrophotometer (Aviv Instruments) or a 8000-series SLM-AMINCO™ spectrophotometer (ThermoSpectronic) with a stirred cuvette.

2. Antibody Fragments

In certain embodiments, an antibody provided herein is an antibody fragment. Antibody fragments include, but are not limited to, Fab, Fab', Fab'-SH, F(ab')₂, Fv, and scFv fragments, and other fragments described below. For a review of certain antibody fragments, see Hudson, P.J. et al., Nat. Med. 9 (2003) 129-134. For a review of scFv fragments, see, e.g., Plueckthun, A., In: The Pharmacology of Monoclonal Antibodies, Vol. 113, Rosenberg and Moore (eds.), Springer-Verlag, New York (1994), pp. 269-315; see also WO 93/16185; and U.S. Patent Nos. 5,571,894 and 5,587,458. For discussion of Fab and F(ab')₂ fragments comprising salvage receptor binding epitope residues and having increased in vivo half-life, see U.S. Patent No. 5,869,046.

Diabodies are antibody fragments with two antigen-binding sites that may be bivalent or bispecific. See, for example, EP 0 404 097; WO 1993/01161; Hudson, P.J. et al., Nat. Med. 9 (2003) 129-134; and Holliger, P. et al., Proc. Natl. Acad. Sci. USA 90 (1993) 6444-6448. Triabodies and tetrabodies are also described in Hudson, P.J. et al., Nat. Med. 9 (2003) 129-134.

Single-domain antibodies are antibody fragments comprising all or a portion of the heavy chain variable domain or all or a portion of the light chain variable domain of an antibody. In certain embodiments, a single-domain antibody is a human single-domain antibody (Domantis, Inc., Waltham, MA; see, e.g., U.S. Patent No. 6,248,516 B1).

Antibody fragments can be made by various techniques, including but not limited to proteolytic digestion of an intact antibody as well as production by recombinant host cells (e.g. *E. coli* or phage), as described herein.

3. Chimeric and Humanized Antibodies

In certain embodiments, an antibody provided herein is a chimeric antibody. Certain chimeric antibodies are described, e.g., in U.S. Patent No. 4,816,567; and Morrison, S.L. et al., Proc. Natl. Acad. Sci. USA 81 (1984) 6851-6855). In one example, a
5 chimeric antibody comprises a non-human variable region (e.g., a variable region derived from a mouse, rat, hamster, rabbit, or non-human primate, such as a monkey) and a human constant region. In a further example, a chimeric antibody is a “class switched” antibody in which the class or subclass has been changed from that of the parental antibody. Chimeric antibodies include antigen-binding fragments thereof.

10 In certain embodiments, a chimeric antibody is a humanized antibody. Typically, a non-human antibody is humanized to reduce immunogenicity to humans, while retaining the specificity and affinity of the parental non-human antibody. Generally, a humanized antibody comprises one or more variable domains in which CDRs, e.g., CDRs, (or portions thereof) are derived from a non-human antibody, and FRs (or
15 portions thereof) are derived from human antibody sequences. A humanized antibody optionally will also comprise at least a portion of a human constant region. In some embodiments, some FR residues in a humanized antibody are substituted with corresponding residues from a non-human antibody (e.g., the antibody from
20 which the CDR residues are derived), e.g., to restore or improve antibody specificity or affinity.

Humanized antibodies and methods of making them are reviewed, e.g., in Almagro, J.C. and Fransson, J., Front. Biosci. 13 (2008) 1619-1633, and are further described, e.g., in Riechmann, I. et al., Nature 332 (1988) 323-329; Queen, C. et al., Proc. Natl. Acad. Sci. USA 86 (1989) 10029-10033; US Patent Nos. 5, 821,337, 7,527,791,
25 6,982,321, and 7,087,409; Kashmiri, S.V. et al., Methods 36 (2005) 25-34 (describing SDR (a-CDR) grafting); Padlan, E.A., Mol. Immunol. 28 (1991) 489-498 (describing “resurfacing”); Dall’Acqua, W.F. et al., Methods 36 (2005) 43-60 (describing “FR shuffling”); and Osbourn, J. et al., Methods 36 (2005) 61-68 and Klimka, A. et al., Br. J. Cancer 83 (2000) 252-260 (describing the “guided selection”
30 approach to FR shuffling).

Human framework regions that may be used for humanization include but are not limited to: framework regions selected using the “best-fit” method (see, e.g., Sims, M.J. et al., J. Immunol. 151 (1993) 2296-2308; framework regions derived from the consensus sequence of human antibodies of a particular subgroup of light or heavy
35 chain variable regions (see, e.g., Carter, P. et al., Proc. Natl. Acad. Sci. USA 89

(1992) 4285-4289; and Presta, L.G. et al., J. Immunol. 151 (1993) 2623-2632); human mature (somatically mutated) framework regions or human germline framework regions (see, e.g., Almagro, J.C. and Fransson, J., Front. Biosci. 13 (2008) 1619-1633); and framework regions derived from screening FR libraries (see, 5 e.g., Baca, M. et al., J. Biol. Chem. 272 (1997) 10678-10684 and Rosok, M.J. et al., J. Biol. Chem. 271 (1996) 22611-22618).

4. Human Antibodies

In certain embodiments, an antibody provided herein is a human antibody. Human antibodies can be produced using various techniques known in the art. Human 10 antibodies are described generally in van Dijk, M.A. and van de Winkel, J.G., Curr. Opin. Pharmacol. 5 (2001) 368-374 and Lonberg, N., Curr. Opin. Immunol. 20 (2008) 450-459.

Human antibodies may be prepared by administering an immunogen to a transgenic animal that has been modified to produce intact human antibodies or intact antibodies 15 with human variable regions in response to antigenic challenge. Such animals typically contain all or a portion of the human immunoglobulin loci, which replace the endogenous immunoglobulin loci, or which are present extrachromosomally or integrated randomly into the animal's chromosomes. In such transgenic mice, the endogenous immunoglobulin loci have generally been inactivated. For review of 20 methods for obtaining human antibodies from transgenic animals, see Lonberg, N., Nat. Biotech. 23 (2005) 1117-1125. See also, e.g., U.S. Patent Nos. 6,075,181 and 6,150,584 describing XENOMOUSE™ technology; U.S. Patent No. 5,770,429 describing HUMAB® technology; U.S. Patent No. 7,041,870 describing K-M MOUSE® technology, and U.S. Patent Application Publication No. US 25 2007/0061900, describing VELOCIMOUSE® technology). Human variable regions from intact antibodies generated by such animals may be further modified, e.g., by combining with a different human constant region.

Human antibodies can also be made by hybridoma-based methods. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal 30 antibodies have been described. (See, e.g., Kozbor, D., J. Immunol. 133 (1984) 3001-3005; Brodeur, B.R. et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York (1987), pp. 51-63; and Boerner, P. et al., J. Immunol. 147 (1991) 86-95) Human antibodies generated via human B-cell hybridoma technology are also described in Li, J. et al., Proc. Natl. Acad. Sci. USA 35 103 (2006) 3557-3562. Additional methods include those described, for example, in

U.S. Patent No. 7,189,826 (describing production of monoclonal human IgM antibodies from hybridoma cell lines) and Ni, J., *Xiandai Mianyixue* 26 (2006) 265-268 (describing human-human hybridomas). Human hybridoma technology (Trioma technology) is also described in Vollmers, H.P. and Brandlein, S., *Histology and Histopathology* 20 (2005) 927-937 and Vollmers, H.P. and Brandlein, S., *Methods and Findings in Experimental and Clinical Pharmacology* 27 (2005) 185-191.

Human antibodies may also be generated by isolating Fv clone variable domain sequences selected from human-derived phage display libraries. Such variable domain sequences may then be combined with a desired human constant domain. Techniques for selecting human antibodies from antibody libraries are described below.

5. Library-Derived Antibodies

Antibodies of the invention may be isolated by screening combinatorial libraries for antibodies with the desired activity or activities. For example, a variety of methods are known in the art for generating phage display libraries and screening such libraries for antibodies possessing the desired binding characteristics. Such methods are reviewed, e.g., in Hoogenboom, H.R. et al., *Methods in Molecular Biology* 178 (2001) 1-37 and further described, e.g., in the McCafferty, J. et al., *Nature* 348 (1990) 552-554; Clackson, T. et al., *Nature* 352 (1991) 624-628; Marks, J.D. et al., *J. Mol. Biol.* 222 (1992) 581-597; Marks, J.D. and Bradbury, A., *Methods in Molecular Biology* 248 (2003) 161-175; Sidhu, S.S. et al., *J. Mol. Biol.* 338 (2004) 299-310; Lee, C.V. et al., *J. Mol. Biol.* 340 (2004) 1073-1093; Fellouse, F.A., *Proc. Natl. Acad. Sci. USA* 101 (2004) 12467-12472; and Lee, C.V. et al., *J. Immunol. Methods* 284 (2004) 119-132.

In certain phage display methods, repertoires of VH and VL genes are separately cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be screened for antigen-binding phage as described in Winter, G. et al., *Ann. Rev. Immunol.* 12 (1994) 433-455. Phage typically display antibody fragments, either as single-chain Fv (scFv) fragments or as Fab fragments. Libraries from immunized sources provide high-affinity antibodies to the immunogen without the requirement of constructing hybridomas. Alternatively, the naive repertoire can be cloned (e.g., from human) to provide a single source of antibodies to a wide range of non-self and also self antigens without any immunization as described by Griffiths, A.D. et al., *EMBO J.* 12 (1993) 725-734. Finally, naive libraries can also be made synthetically by cloning non-rearranged V-

gene segments from stem cells, and using PCR primers containing random sequence to encode the highly variable CDR3 regions and to accomplish rearrangement *in vitro*, as described by Hoogenboom, H.R. and Winter, G., J. Mol. Biol. 227 (1992) 381-388. Patent publications describing human antibody phage libraries include, for example: US Patent No. 5,750,373, and US Patent Publication Nos. 2005/0079574, 2005/0119455, 2005/0266000, 2007/0117126, 2007/0160598, 2007/0237764, 2007/0292936, and 2009/0002360.

Antibodies or antibody fragments isolated from human antibody libraries are considered human antibodies or human antibody fragments herein.

6. Antibody Variants

In certain embodiments, amino acid sequence variants of the antibodies provided herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence variants of an antibody may be prepared by introducing appropriate modifications into the nucleotide sequence encoding the antibody, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of residues within the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution can be made to arrive at the final construct, provided that the final construct possesses the desired characteristics, e.g., antigen-binding.

a) Substitution, Insertion, and Deletion Variants

In certain embodiments, antibody variants having one or more amino acid substitutions are provided. Sites of interest for substitutional mutagenesis include the CDRs and FRs (in a preferred embodiment framework residues not relevant for the binding properties of the antibodies (see e.g. Foote J. and Winter G., J. Mol. Biol. (1992) 224, 487-499). Exemplary changes are provided in Table 1 under the heading of "exemplary substitutions", and as further described below in reference to amino acid side chain classes. Conservative substitutions are shown in Table 1 under the heading of "preferred substitutions". Amino acid substitutions may be introduced into an antibody of interest and the products screened for a desired activity, e.g., retained/improved antigen binding, decreased immunogenicity, or improved ADCC or CDC.

Table 1

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp, Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala
His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu
Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Val; Ser	Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe

Original Residue	Exemplary Substitutions	Preferred Substitutions
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

Amino acids may be grouped according to common side-chain properties:

(1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;

(2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;

5 (3) acidic: Asp, Glu;

(4) basic: His, Lys, Arg;

(5) residues that influence chain orientation: Gly, Pro;

(6) aromatic: Trp, Tyr, Phe.

10 Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

One type of substitutional variant involves substituting one or more CDRs of a parent antibody (*e.g.* a humanized or human antibody). Generally, the resulting variant(s) selected for further study will have modifications (*e.g.*, improvements) in certain biological properties (*e.g.*, increased affinity, reduced immunogenicity) relative to the parent antibody and/or will have substantially retained certain biological properties of the parent antibody. An exemplary substitutional variant is an affinity matured antibody, which may be conveniently generated, *e.g.*, using phage display-based affinity maturation techniques such as those described herein. Briefly, one or more CDR residues are mutated and the variant antibodies displayed on phage and screened for a particular biological activity (*e.g.* binding affinity).

Alterations (*e.g.*, substitutions) may be made in CDRs, *e.g.*, to improve antibody affinity. Such alterations may be made in CDR “hotspots,” *i.e.*, residues encoded by codons that undergo mutation at high frequency during the somatic maturation process (see, *e.g.*, Chowdhury, P.S., *Methods Mol. Biol.* 207 (2008) 179-196), and/or SDRs (a-CDRs), with the resulting variant VH or VL being tested for binding

affinity. Affinity maturation by constructing and reselecting from secondary libraries has been described, e.g., in Hoogenboom, H.R. et al. in *Methods in Molecular Biology* 178 (2002) 1-37. In some embodiments of affinity maturation, diversity is introduced into the variable genes chosen for maturation by any of a variety of methods (e.g., error-prone PCR, chain shuffling, or oligonucleotide-directed mutagenesis). A secondary library is then created. The library is then screened to identify any antibody variants with the desired affinity. Another method to introduce diversity involves CDR-directed approaches, in which several CDR residues (e.g., 4-6 residues at a time) are randomized. CDR residues involved in antigen binding may be specifically identified, e.g., using alanine scanning mutagenesis or modeling. CDR-H3 and CDR-L3 in particular are often targeted.

In certain embodiments, substitutions, insertions, or deletions may occur within one or more CDRs so long as such alterations do not substantially reduce the ability of the antibody to bind antigen. For example, conservative alterations (e.g., conservative substitutions as provided herein) that do not substantially reduce binding affinity may be made in CDRs. Such alterations may be outside of CDR "hotspots" or SDRs. In certain embodiments of the variant VH and VL sequences provided above, each CDR either is unaltered, or contains no more than one, two or three amino acid substitutions.

A useful method for identification of residues or regions of an antibody that may be targeted for mutagenesis is called "alanine scanning mutagenesis" as described by Cunningham, B.C. and Wells, J.A., *Science* 244 (1989) 1081-1085. In this method, a residue or group of target residues (e.g., charged residues such as arg, asp, his, lys, and glu) are identified and replaced by a neutral or negatively charged amino acid (e.g., alanine or polyalanine) to determine whether the interaction of the antibody with antigen is affected. Further substitutions may be introduced at the amino acid locations demonstrating functional sensitivity to the initial substitutions. Alternatively, or additionally, a crystal structure of an antigen-antibody complex to identify contact points between the antibody and antigen. Such contact residues and neighboring residues may be targeted or eliminated as candidates for substitution. Variants may be screened to determine whether they contain the desired properties.

Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal

methionyl residue. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody to an enzyme (e.g. for ADEPT) or a polypeptide which increases the serum half-life of the antibody.

b) Fc region variants

5 In certain embodiments, one or more amino acid modifications may be introduced into the Fc region of an antibody provided herein, thereby generating an Fc region variant. The Fc region variant may comprise a human Fc region sequence (*e.g.*, a human IgG1, IgG2, IgG3 or IgG4 Fc region) comprising an amino acid modification (*e.g.* a substitution) at one or more amino acid positions.

10 Antibodies with reduced effector function include those with substitution of one or more of Fc region residues 238, 265, 269, 270, 297, 327 and 329 (U.S. Patent No. 6,737,056). Such Fc mutants include Fc mutants with substitutions at two or more of amino acid positions 265, 269, 270, 297 and 327, including the so-called "DANA" Fc mutant with substitution of residues 265 and 297 to alanine (US Patent No.
15 7,332,581).

Certain antibody variants with improved or diminished binding to FcRs are described. (See, *e.g.*, U.S. Patent No. 6,737,056; WO 2004/056312, and Shields, R.L. et al., J. Biol. Chem. 276 (2001) 6591-6604)

20 In one embodiment the invention such antibody is a IgG1 with mutations L234A and L235A or with mutations L234A, L235A and P329G. In another embodiment or IgG4 with mutations S228P and L235E or S228P, L235E or and P329G (numbering according to EU index of Kabat et al, Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD, 1991)

25 Antibodies with increased half lives and improved binding to the neonatal Fc receptor (FcRn), which is responsible for the transfer of maternal IgGs to the fetus (Guyer, R.L. et al., J. Immunol. 117 (1976) 587-593, and Kim, J.K. et al., J. Immunol. 24 (1994) 2429-2434), are described in US 2005/0014934. Those antibodies comprise an Fc region with one or more substitutions therein which improve binding
30 of the Fc region to FcRn. Such Fc variants include those with substitutions at one or more of Fc region residues: 238, 256, 265, 272, 286, 303, 305, 307, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 413, 424 or 434, *e.g.*, substitution of Fc region residue 434 (US Patent No. 7,371,826).

See also Duncan, A.R. and Winter, G., Nature 322 (1988) 738-740; US 5,648,260; US 5,624,821; and WO 94/29351 concerning other examples of Fc region variants.

c) Cysteine engineered antibody variants

In certain embodiments, it may be desirable to create cysteine engineered antibodies, e.g., "thioMAbs," in which one or more residues of an antibody are substituted with cysteine residues. In particular embodiments, the substituted residues occur at accessible sites of the antibody. By substituting those residues with cysteine, reactive thiol groups are thereby positioned at accessible sites of the antibody and may be used to conjugate the antibody to other moieties, such as drug moieties or linker-drug moieties, to create an immunoconjugate, as described further herein. In certain embodiments, any one or more of the following residues may be substituted with cysteine: V205 (Kabat numbering) of the light chain; A118 (EU numbering) of the heavy chain; and S400 (EU numbering) of the heavy chain Fc region. Cysteine engineered antibodies may be generated as described, e.g., in U.S. Patent No. 7,521,541.

d) Antibody Derivatives

In certain embodiments, an antibody provided herein may be further modified to contain additional non-proteinaceous moieties that are known in the art and readily available. The moieties suitable for derivatization of the antibody include but are not limited to water soluble polymers. Non-limiting examples of water soluble polymers include, but are not limited to, polyethylene glycol (PEG), copolymers of ethylene glycol/propylene glycol, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1, 3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), and dextran or poly(n-vinyl pyrrolidone)polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide copolymers, polyoxyethylated polyols (e.g., glycerol), polyvinyl alcohol, and mixtures thereof. Polyethylene glycol propionaldehyde may have advantages in manufacturing due to its stability in water. The polymer may be of any molecular weight, and may be branched or unbranched. The number of polymers attached to the antibody may vary, and if more than one polymer is attached, they can be the same or different molecules. In general, the number and/or type of polymers used for derivatization can be determined based on considerations including, but not limited to, the particular properties or functions of the antibody to be improved, whether the antibody derivative will be used in a therapy under defined conditions, etc.

In another embodiment, conjugates of an antibody and non-proteinaceous moiety that may be selectively heated by exposure to radiation are provided. In one embodiment, the non-proteinaceous moiety is a carbon nanotube (Kam, N.W. et al., Proc. Natl. Acad. Sci. USA 102 (2005) 11600-11605). The radiation may be of any wavelength, and includes, but is not limited to, wavelengths that do not harm ordinary cells, but which heat the non-proteinaceous moiety to a temperature at which cells proximal to the antibody-non-proteinaceous moiety are killed.

B. Recombinant Methods and Compositions

Antibodies may be produced using recombinant methods and compositions, e.g., as described in U.S. Patent No. 4,816,567. In one embodiment, isolated nucleic acid encoding an anti-HLA-G antibody described herein is provided. Such nucleic acid may encode an amino acid sequence comprising the VL and/or an amino acid sequence comprising the VH of the antibody (e.g., the light and/or heavy chains of the antibody). In a further embodiment, one or more vectors (e.g., expression vectors) comprising such nucleic acid are provided. In a further embodiment, a host cell comprising such nucleic acid is provided. In one such embodiment, a host cell comprises (e.g., has been transformed with): (1) a vector comprising a nucleic acid that encodes an amino acid sequence comprising the VL of the antibody and an amino acid sequence comprising the VH of the antibody, or (2) a first vector comprising a nucleic acid that encodes an amino acid sequence comprising the VL of the antibody and a second vector comprising a nucleic acid that encodes an amino acid sequence comprising the VH of the antibody. In one preferred embodiment, the host cell is eukaryotic, e.g. a Chinese Hamster Ovary (CHO) cell, a HEK293 cell or lymphoid cell (e.g., Y0, NS0, Sp20 cell). In one embodiment, a method of making an anti-HLA-G antibody or bispecific antibody is provided, wherein the method comprises culturing a host cell comprising a nucleic acid encoding the antibody, as provided above, under conditions suitable for expression of the antibody, and optionally recovering the antibody from the host cell (or host cell culture medium).

For recombinant production of an anti-HLA-G antibody, nucleic acid encoding an antibody, e.g., as described above, is isolated and inserted into one or more vectors for further cloning and/or expression in a host cell. Such nucleic acid may be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody).

The most suitable host cells for cloning or expression of antibody-encoding vectors eukaryotic cells, preferably mammalian cells, described herein.

Vertebrate cells may be used as hosts. For example, mammalian cell lines that are adapted to grow in suspension may be useful. Other examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7); human embryonic kidney line (293 or 293 cells as described, e.g., in Graham, F.L. et al., J. Gen Virol. 36 (1977) 59-74); baby hamster kidney cells (BHK); mouse sertoli cells (TM4 cells as described, e.g., in Mather, J.P., Biol. Reprod. 23 (1980) 243-252); monkey kidney cells (CV1); African green monkey kidney cells (VERO-76); human cervical carcinoma cells (HELA); canine kidney cells (MDCK); buffalo rat liver cells (BRL 3A); human lung cells (W138); human liver cells (Hep G2); mouse mammary tumor (MMT 060562); TRI cells, as described, e.g., in Mather, J.P. et al., Annals N.Y. Acad. Sci. 383 (1982) 44-68; MRC 5 cells; and FS4 cells. Most useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including DHFR⁻ CHO cells (Urlaub, G. et al., Proc. Natl. Acad. Sci. USA 77 (1980) 4216-4220); and myeloma cell lines such as Y0, NS0 and Sp2/0. For a review of certain mammalian host cell lines suitable for antibody production, see, e.g., Yazaki, P. and Wu, A.M., Methods in Molecular Biology, Vol. 248, Lo, B.K.C. (ed.), Humana Press, Totowa, NJ (2004), pp. 255-268.

To some extent also prokaryotic cells may be used, however with the disadvantage of sometimes higher efforts and more complex procedures,. For expression of antibody fragments and polypeptides in bacteria, see, e.g., US 5,648,237, US 5,789,199, and US 5,840,523. (See also Charlton, K.A., In: Methods in Molecular Biology, Vol. 248, Lo, B.K.C. (ed.), Humana Press, Totowa, NJ (2003), pp. 245-254, describing expression of antibody fragments in *E. coli*.) After expression, the antibody may be isolated from the bacterial cell paste in a soluble fraction and can be further purified.

Plant cell cultures can also be utilized as hosts. See, e.g., US Patent Nos. 5,959,177, 6,040,498, 6,420,548, 7,125,978, and 6,417,429 (describing PLANTIBODIESTM technology for producing antibodies in transgenic plants).

C. Assays

Anti-HLA-G antibodies provided herein may be identified, screened for, or characterized for their physical/chemical properties and/or biological activities by various assays known in the art.

1. Binding assays and other assays

In one aspect, an antibody of the invention is tested for its antigen binding activity, e.g., by known methods such as ELISA, Western blot, etc. Detailed exemplary methods for mapping an epitope to which an antibody binds are provided in Morris, G.E. (ed.), Epitope Mapping Protocols, In: Methods in Molecular Biology, Vol. 66, Humana Press, Totowa, NJ (1996).

2. Activity assays

In one aspect, assays are provided for identifying anti-HLA-G antibodies thereof having biological activity. Biological activity may include, e.g., the ability to enhance the activation and/or proliferation of different immune cells including T-cells. E.g. they enhance secretion of immunomodulating cytokines (e.g. interferon-gamma (IFN-gamma) and/or tumor necrosis factor alpha (TNF alpha)). Other immunomodulating cytokines which are or can be enhance are e.g. IL1 β , IL6, IL12, Granzyme B etc. binding to different cell types. Antibodies having such biological activity in vivo and/or in vitro are also provided.

In certain embodiments, an antibody of the invention is tested for such biological activity as described e.g. in Examples below.

D. Methods and Compositions for Diagnostics and Detection

In certain embodiments, any of the anti-HLA-G antibodies provided herein is useful for detecting the presence of HLA-G in a biological sample. The term “detecting” as used herein encompasses quantitative or qualitative detection. In certain embodiments, a biological sample comprises a cell or tissue, such as immune cell or T cell infiltrates and or tumor cells.

In one embodiment, an anti-HLA-G antibody for use in a method of diagnosis or detection is provided. In a further aspect, a method of detecting the presence of HLA-G in a biological sample is provided. In certain embodiments, the method comprises contacting the biological sample with an anti-HLA-G antibody as described herein under conditions permissive for binding of the anti-HLA-G antibody to HLA-G, and detecting whether a complex is formed between the anti-HLA-G antibody and HLA-G. Such method may be an *in vitro* or *in vivo* method. In one embodiment, an anti-HLA-G antibody is used to select subjects eligible for therapy with an anti-HLA-G antibody, e.g. where HLA-G is a biomarker for selection of patients.

In certain embodiments, labeled anti-HLA-G antibodies are provided. Labels include, but are not limited to, labels or moieties that are detected directly (such as fluorescent, chromophoric, electron-dense, chemiluminescent, and radioactive labels), as well as moieties, such as enzymes or ligands, that are detected indirectly, e.g., through an enzymatic reaction or molecular interaction. Exemplary labels include, but are not limited to, the radioisotopes ^{32}P , ^{14}C , ^{125}I , ^3H , and ^{131}I , fluorophores such as rare earth chelates or fluorescein and its derivatives, rhodamine and its derivatives, dansyl, umbelliferone, luciferases, e.g., firefly luciferase and bacterial luciferase (U.S. Patent No. 4,737,456), luciferin, 2,3-dihydrophthalazinediones, horseradish peroxidase (HRP), alkaline phosphatase, β -galactosidase, glucoamylase, lysozyme, saccharide oxidases, e.g., glucose oxidase, galactose oxidase, and glucose-6-phosphate dehydrogenase, heterocyclic oxidases such as uricase and xanthine oxidase, coupled with an enzyme that employs hydrogen peroxide to oxidize a dye precursor such as HRP, lactoperoxidase, or microperoxidase, biotin/avidin, spin labels, bacteriophage labels, stable free radicals, and the like.

E. Pharmaceutical Formulations

Pharmaceutical formulations of anti-HLA-G antibodies or the anti-HLA-G/anti-CD3 bispecific antibodies as described herein are prepared by mixing such antibodies having the desired degree of purity with one or more optional pharmaceutically acceptable carriers (Remington's Pharmaceutical Sciences, 16th edition, Osol, A. (ed.) (1980)), in the form of lyophilized formulations or aqueous solutions. Pharmaceutically acceptable carriers are generally nontoxic to recipients at the dosages and concentrations employed, and include, but are not limited to: buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyl dimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride; benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as poly(vinylpyrrolidone); amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as

polyethylene glycol (PEG). Exemplary pharmaceutically acceptable carriers herein further include interstitial drug dispersion agents such as soluble neutral-active hyaluronidase glycoproteins (sHASEGP), for example, human soluble PH-20 hyaluronidase glycoproteins, such as rhuPH20 (HYLENEX[®], Baxter International, Inc.). Certain exemplary sHASEGPs and methods of use, including rhuPH20, are described in US Patent Publication Nos. 2005/0260186 and 2006/0104968. In one aspect, a sHASEGP is combined with one or more additional glycosaminoglycanases such as chondroitinases.

Exemplary lyophilized antibody formulations are described in US Patent No. 6,267,958. Aqueous antibody formulations include those described in US Patent No. 6,171,586 and WO 2006/044908, the latter formulations including a histidine-acetate buffer.

The formulation herein may also contain more than one active ingredients as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to further provide. Such active ingredients are suitably present in combination in amounts that are effective for the purpose intended.

Active ingredients may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly- (methyl methacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Osol, A. (ed.) (1980).

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semi-permeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g. films, or microcapsules.

The formulations to be used for *in vivo* administration are generally sterile. Sterility may be readily accomplished, e.g., by filtration through sterile filtration membranes.

F. Therapeutic Methods and Compositions

Any of the anti-HLA-G antibodies or the anti-HLA-G/anti-CD3 bispecific antibodies provided herein may be used in therapeutic methods.

In one aspect, an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody for use as a medicament is provided. In further aspects, an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody for use in treating cancer is provided. In certain embodiments, an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody for use in a method of treatment is provided. In certain embodiments, the invention provides an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody for use in a method of treating an individual having cancer comprising administering to the individual an effective amount of the anti-HLA-G/anti-CD3 bispecific antibody.

In further embodiments, the invention provides an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody for use as immunomodulatory agent/ to directly or indirectly induce proliferation and/or activation of immune cells (like T cells, B cells and myeloid cells including monocytes, macrophages, dendritic cells, plasmacytoid dendritic cells) e.g. by secretion of immunostimulatory cytokines like TNFalpha (TNFa) and IFNgamma (IFNg) or further recruitment of immune cells. In certain embodiments, the invention provides an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody for use in a method of immunomodulatory agent/ to directly or indirectly induce proliferation, activation of immune cells e.g. by secretion of immunostimulatory cytokines like TNFa and IFNgamma or further recruitment of immune cells in an individual comprising administering to the individual an effective of the anti-HLA-G antibody or anti-HLA-G/anti-CD3 bispecific antibody for immunomodulation/ or directly or indirectly induce proliferation, activation of immune cells e.g. by secretion of immunostimulatory cytokines like TNFa and IFNgamma or further recruitment of immune cells.

In further embodiments, the invention provides an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody for use as immunostimulatory agent/or stimulating tumor necrosis factor alpha (TNF alpha) secretion. In certain embodiments, the invention provides an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody for use in a method of immunomodulation to directly or indirectly induce proliferation, activation e.g. by secretion of immunostimulatory cytokines like TNFa and IFNg or further recruitment of immune cells in an individual comprising administering to the individual an effective of the anti-HLA-G antibody or anti-HLA-G/anti-CD3 bispecific antibody immunomodulation to directly or indirectly induce proliferation, activation e.g. by secretion of immunostimulatory cytokines like TNFa and IFNg or further recruitment of immune cells

The term “cancer” as used herein may be, for example, lung cancer, non small cell lung (NSCL) cancer, bronchioloalviolar cell lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach
5 cancer, gastric cancer, colon cancer, breast cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft
10 tissue, cancer of the urethra, cancer of the penis, prostate cancer, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, mesothelioma, hepatocellular cancer, biliary cancer, neoplasms of the central nervous system (CNS), spinal axis tumors, brain stem glioma, glioblastoma multiforme, astrocytomas, schwannomas, ependymomas, medulloblastomas,
15 meningiomas, squamous cell carcinomas, pituitary adenoma, lymphoma, lymphocytic leukemia, including refractory versions of any of the above cancers, or a combination of one or more of the above cancers.

An “individual” according to any of the above embodiments is preferably a human. In a further aspect, the invention provides for the use of an anti-HLA-G antibody in
20 the manufacture or preparation of a medicament. In one embodiment, the medicament is for treatment of cancer. In a further embodiment, the medicament is for use in a method of treating cancer comprising administering to an individual having cancer an effective amount of the medicament. In a further embodiment, the medicament is for inducing cell mediated lysis of cancer cells In a further
25 embodiment, the medicament is for use in a method of inducing cell mediated lysis of cancer cells in an individual suffering from cancer comprising administering to the individual an amount effective of the medicament to induce apoptosis in a cancer cell/ or to inhibit cancer cell proliferation. An “individual” according to any of the above embodiments may be a human.

30 In a further aspect, the invention provides a method for treating cancer. In one embodiment, the method comprises administering to an individual having cancer an effective amount of an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody. An “individual” according to any of the above embodiments may be a human.

In a further aspect, the invention provides a method for inducing cell mediated lysis of cancer cells in an individual suffering from cancer. In one embodiment, the method comprises administering to the individual an effective amount of an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody to induce cell mediated lysis of cancer cells in the individual suffering from cancer. In one
5 embodiment, an “individual” is a human.

In a further aspect, the invention provides pharmaceutical formulations comprising any of the anti-HLA-G antibodies or anti-HLA-G/anti-CD3 bispecific antibodies provided herein, e.g., for use in any of the above therapeutic methods. In one
10 embodiment, a pharmaceutical formulation comprises any of the anti-HLA-G antibodies or anti-HLA-G/anti-CD3 bispecific antibodies provided herein and a pharmaceutically acceptable carrier.

An antibody of the invention (and any additional therapeutic agent) can be administered by any suitable means, including parenteral, intrapulmonary, and
15 intranasal, and, if desired for local treatment, intralesional administration. Parenteral infusions include intramuscular, intravenous, intra-arterial, intraperitoneal, or subcutaneous administration. Dosing can be by any suitable route, e.g. by injections, such as intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic. Various dosing schedules including but not limited
20 to single or multiple administrations over various time-points, bolus administration, and pulse infusion are contemplated herein.

Antibodies of the invention would be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being
25 treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The antibody need not be, but is optionally formulated with one or more agents currently used to prevent or treat the disorder in question. The effective amount of such other agents depends
30 on the amount of antibody present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as described herein, or about from 1 to 99% of the dosages described herein, or in any dosage and by any route that is empirically/clinically determined to be appropriate.

For the prevention or treatment of disease, the appropriate dosage of an antibody of the invention (when used alone or in combination with one or more other additional therapeutic agents) will depend on the type of disease to be treated, the type of antibody, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments. Depending on the type and severity of the disease, about 1 µg/kg to 15 mg/kg (e.g. 0.5mg/kg - 10 mg/kg) of antibody can be an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. One typical daily dosage might range from about 1 µg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment would generally be sustained until a desired suppression of disease symptoms occurs. One exemplary dosage of the antibody would be in the range from about 0.05 mg/kg to about 10 mg/kg. Thus, one or more doses of about 0.5 mg/kg, 2.0 mg/kg, 4.0 mg/kg or 10 mg/kg (or any combination thereof) may be administered to the patient. Such doses may be administered intermittently, e.g. every week or every three weeks (e.g. such that the patient receives from about two to about twenty, or e.g. about six doses of the antibody). An initial higher loading dose, followed by one or more lower doses may be administered. An exemplary dosing regimen comprises administering an initial loading dose of about 4 mg/kg, followed by a weekly maintenance dose of about 2 mg/kg of the antibody. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

It is understood that any of the above formulations or therapeutic methods may be carried out using an immunoconjugate of the invention in place of or in addition to an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody.

It is understood that any of the above formulations or therapeutic methods may be carried out using an immunoconjugate of the invention in place of or in addition to an anti-HLA-G antibody or an anti-HLA-G/anti-CD3 bispecific antibody.

II. Articles of Manufacture

In another aspect of the invention, an article of manufacture containing materials useful for the treatment, prevention and/or diagnosis of the disorders described above is provided. The article of manufacture comprises a container and a label or package

insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, IV solution bags, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is by itself or combined with another composition effective for treating, preventing and/or diagnosing the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is an antibody of the invention. The label or package insert indicates that the composition is used for treating the condition of choice. Moreover, the article of manufacture may comprise (a) a first container with a composition contained therein, wherein the composition comprises an antibody of the invention; and (b) a second container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. The article of manufacture in this embodiment of the invention may further comprise a package insert indicating that the compositions can be used to treat a particular condition. Alternatively, or additionally, the article of manufacture may further comprise a second (or third) container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

The following examples and figures are provided to aid the understanding of the present invention, the true scope of which is set forth in the appended claims. It is understood that modifications can be made in the procedures set forth without departing from the spirit of the invention.

Description of the amino acid sequences

Anti-HLA-G antibodies/antigen binding moieties (SEQ ID Nos of variable regions and complement determining regions (CDRs)):

SEQ ID NO: 1	heavy chain CDR-H1, HLA-G-0090
SEQ ID NO: 2	heavy chain CDR-H2, HLA-G-0090
SEQ ID NO: 3	heavy chain CDR-H3, HLA-G-0090
SEQ ID NO: 4	light chain CDR-L1, HLA-G-0090

	SEQ ID NO: 5	light chain CDR-L2, HLA-G-0090
	SEQ ID NO: 6	light chain CDR-L3, HLA-G-0090
	SEQ ID NO: 7	heavy chain variable domain VH, HLA-G-0090
	SEQ ID NO: 8	light chain variable domain VL, HLA-G-0090
5	SEQ ID NO: 9	light chain CDR-L1, HLA-G-0090-VL-N31D
	SEQ ID NO: 10	light chain variable domain VL, HLA-G-0090-VL-N31D
	SEQ ID NO: 11	light chain CDR-L1, HLA-G-0090-VL-N31L
	SEQ ID NO: 12	light chain variable domain VL, HLA-G-0090-VL-N31L
	SEQ ID NO: 13	light chain CDR-L1, HLA-G-0090-VL-N31Q
10	SEQ ID NO: 14	light chain variable domain VL, HLA-G-0090-VL-N31Q
	SEQ ID NO: 15	light chain CDR-L1, HLA-G-0090-VL-N31S
	SEQ ID NO: 16	light chain variable domain VL, HLA-G-0090-VL-N31S
	SEQ ID NO: 17	light chain CDR-L1, HLA-G-0090-VL-N31T
	SEQ ID NO: 18	light chain variable domain VL, HLA-G-0090-VL-N31T
15	SEQ ID NO: 19	light chain CDR-L1, HLA-G-0090-VL-N31Y
	SEQ ID NO: 20	light chain variable domain VL, HLA-G-0090-VL-N31Y
	SEQ ID NO: 21	light chain CDR-L1, HLA-G-0090-VL-N31Y-N38Y
	SEQ ID NO: 22	light chain variable domain VL, HLA-G-0090-VL-N31Y-N38Y
20	SEQ ID NO: 23	light chain CDR-L1, HLA-G-0090-VL-S32P
	SEQ ID NO: 24	light chain variable domain VL, HLA-G-0090-VL-S32P
	SEQ ID NO: 25	light chain CDR-L1, HLA-G-0090-VL-S33A
	SEQ ID NO: 26	light chain variable domain VL, HLA-G-0090-VL-S33A

SEQ ID NO: 27 light chain CDR-L1, HLA-G-0090-VL-S33D

SEQ ID NO: 28 light chain variable domain VL, HLA-G-0090-VL-S33D

SEQ ID NO: 29 light chain CDR-L1, HLA-G-0090-VL-S33P

SEQ ID NO: 30 light chain variable domain VL, HLA-G-0090-VL-S33P

5 **Further sequences**

SEQ ID NO: 31 exemplary human HLA-G

SEQ ID NO: 32 exemplary human HLA-G extracellular domain (ECD)

SEQ ID NO: 33 exemplary human β 2M

10 SEQ ID NO: 34 modified human HLA-G (wherein the HLA-G specific amino acids have been replaced by HLA-A consensus amino acids (= degrafted HLA-G see also Figure 1) ECD)

SEQ ID NO: 35 exemplary human HLA-A2

SEQ ID NO: 36 exemplary human HLA-A2 ECD

SEQ ID NO: 37 exemplary mouse H2Kd ECD

15 SEQ ID NO: 38 exemplary rat RT1A ECD

SEQ ID NO: 39 exemplary human HLA-G β 2M MHC class I complex

SEQ ID NO: 40 exemplary modified human HLA-G β 2M MHC class I complex (wherein the HLA-G specific amino acids have been replaced by HLA-A consensus amino acids (= degrafted HLA-G) see also Figure 2)

20 SEQ ID NO: 41 exemplary mouse H2Kd β 2M MHC class I complex

SEQ ID NO: 42 exemplary human HLA-G/ mouse H2Kd β 2M MHC class I complex wherein the positions specific for human HLA-G are grafted onto the mouse H2Kd framework

25 SEQ ID NO: 43 exemplary rat RT1A β 2M MHC class I complex

- 103 -

- SEQ ID NO: 44 exemplary human HLA-G/ rat RT1A β 2M MHC class I complex wherein the positions specific for human HLA-G are grafted onto the rat RT1A framework
- SEQ ID NO: 45 linker and his-Tag
- 5 SEQ ID NO: 46 peptide
- SEQ ID NO: 47 human kappa light chain constant region
- SEQ ID NO: 48 human lambda light chain constant region
- SEQ ID NO: 49 human heavy chain constant region derived from IgG1
- 10 SEQ ID NO: 50 human heavy chain constant region derived from IgG1 with mutations L234A, L235A and P329G
- SEQ ID NO: 51 human heavy chain constant region derived from IgG4

Anti-CD3 antibodies/ antigen binding moieties (variable regions and complementarity determining regions (CDRs)):

- 15 SEQ ID NO: 52 heavy chain CDR-H1, P035-093 (abbreviated as P035)
- SEQ ID NO: 53 heavy chain CDR-H2, P035-093
- SEQ ID NO: 54 heavy chain CDR-H3, P035-093
- SEQ ID NO: 55 light chain CDR-L1, P035-093
- SEQ ID NO: 56 light chain CDR-L2, P035-093
- 20 SEQ ID NO: 57 light chain CDR-L3, P035-093
- SEQ ID NO: 58 heavy chain variable domain VH, P035-093
- SEQ ID NO: 59 light chain variable domain VL, P035-093
- SEQ ID NO: 60 heavy chain CDR-H1, Clone 22 (abbreviated as Cl22)
- 25 SEQ ID NO: 61 heavy chain CDR-H2, Clone 22

- 104 -

	SEQ ID NO: 62	heavy chain CDR-H3, Clone 22
	SEQ ID NO: 63	light chain CDR-L1, Clone 22
	SEQ ID NO: 64	light chain CDR-L2, Clone 22
	SEQ ID NO: 65	light chain CDR-L3, Clone 22
5	SEQ ID NO: 66	heavy chain variable domain VH, Clone 22
	SEQ ID NO: 67	light chain variable domain VL, Clone 22
	SEQ ID NO: 68	heavy chain CDR-H1, V9
	SEQ ID NO: 69	heavy chain CDR-H2, V9
10	SEQ ID NO: 70	heavy chain CDR-H3, V9
	SEQ ID NO: 71	light chain CDR-L1, V9
	SEQ ID NO: 72	light chain CDR-L2, V9
	SEQ ID NO: 73	light chain CDR-L3, V9
	SEQ ID NO: 74	heavy chain variable domain VH, V9
15	SEQ ID NO: 75	light chain variable domain VL, V9

Bispecific anti-HLA-G/anti-CD3 T cell bispecific (TCB) antibodies:**P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035-093 (P035)):**

	SEQ ID NO: 76	light chain 1 P1AF7977
20	SEQ ID NO: 77	light chain 2 P1AF7977
	SEQ ID NO: 78	heavy chain 1 P1AF7977
	SEQ ID NO: 79	heavy chain 2 P1AF7977

P1AF7978 (HLA-G-0090-VL-S32P/ CD3 Clone 22 (Cl22)):

	SEQ ID NO: 80	light chain 1 P1AF7978
--	---------------	------------------------

- 105 -

SEQ ID NO: 81 light chain 2 P1AF7978

SEQ ID NO: 82 heavy chain 1 P1AF7978

SEQ ID NO: 83 heavy chain 2 P1AF7978

P1AF7979 (HLA-G-0090-VL-S32P/ CD3 V9):

5 SEQ ID NO: 84 light chain 1 P1AF7979

SEQ ID NO: 85 light chain 2 P1AF7979

SEQ ID NO: 86 heavy chain 1 P1AF7979

SEQ ID NO: 87 heavy chain 2 P1AF7979

10 **Further sequences**

SEQ ID NO: 88 exemplary human CD3

SEQ ID NO: 89 exemplary cynomolgus CD3

SEQ ID NO: 90 Human CD3 epsilon stalk – Fc(knob) – Avi

SEQ ID NO: 91 Human CD3 delta stalk – Fc (hole) – Avi

15 SEQ ID NO: 92 CD3_{orig} VHSEQ ID NO: 93 CD3_{orig} VLSEQ ID NO: 94 CD3_{orig} IgG HC

SEQ ID NO: 95 P035 IgG HC

SEQ ID NO: 96 CD3_{orig} / P035 IgG LC

20

The amino acid sequences of HLA-G-0090 antibody (variable regions with underlined complementarity determining regions (CDRs) and unmodified N-glycosylation site in CDR-L1 (bold)):

SEQ ID NO: 7: heavy chain variable domain VH, HLA-G-0090:

- 106 -

QVQLQQSGPGLLKPSQTLSTCAISGDSVSSNRAAWNWIRQSPSRGLEWLG
 RTYYRSKQWYNDYAVSVQGRITLIPDTSKNQFSLRLNSVTPEDTAVYYCASV
RAVAPFDYWGQGVLVTVSS

SEQ ID NO: 8: light chain variable domain VL, HLA-G-0090

5 DIVMTQSPDSLAVSLGERATINCKSSQSVLNSSNNKNNLAWYQQQPGQPP
 KLLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTP
WTFGQGTKVEIK

**The amino acid sequences of modified HLA-G-0090 antibody light chain
 variable regions (with underlined complementarity determining regions
 (CDRs) and modified N-glycosylation site in CDR-L1 (bold)):**

10

SEQ ID NO:10: light chain variable domain VL, HLA-G-0090-N31D

DIVMTQSPDSLAVSLGERATINCKSSQSVL**D**SSNNKNNLAWYQQQPGQPP
 KLLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTP
WTFGQGTKVEIK

15

SEQ ID NO: 12: light chain variable domain VL, HLA-G-0090-N31L

DIVMTQSPDSLAVSLGERATINCKSSQSVL**L**SSNNKNNLAWYQQQPGQPPK
 LLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTPW
TFGQGTKVEIK

SEQ ID NO: 14: light chain variable domain VL, HLA-G-0090-N31Q

20

DIVMTQSPDSLAVSLGERATINCKSSQSVL**Q**SSNNKNNLAWYQQQPGQPP
 KLLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTP
WTFGQGTKVEIK

SEQ ID NO: 16: light chain variable domain VL, HLA-G-0090-N31S

25

DIVMTQSPDSLAVSLGERATINCKSSQSVL**S**SSNNKNNLAWYQQQPGQPPK
 LLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTPW
TFGQGTKVEIK

SEQ ID NO: 18: light chain variable domain VL, HLA-G-0090-N31T

- 107 -

DIVMTQSPDSLAVSLGERATINCKSSQSVLTSSNNKNNLAWYQQQPGQPPK
LLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTPW
TFGQGTKVEIK

SEQ ID NO: 20: light chain variable domain VL, HLA-G-0090-N31Y

5 DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNNLAWYQQQPGQPPK
LLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTPW
TFGQGTKVEIK

SEQ ID NO: 22: light chain variable domain VL, HLA-G-0090-N31Y-N38Y

10 DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQQPGQPPK
LLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTPW
TFGQGTKVEIK

SEQ ID NO: 24: light chain variable domain VL, HLA-G-0090-S32P

15 DIVMTQSPDSLAVSLGERATINCKSSQSVLNPSNNKNNLAWYQQQPGQPP
KLLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTP
WTFGQGTKVEIK

SEQ ID NO: 26: light chain variable domain VL, HLA-G-0090-S33A

DIVMTQSPDSLAVSLGERATINCKSSQSVLNSANNKNNLAWYQQQPGQPP
KLLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTP
WTFGQGTKVEIK

20 SEQ ID NO: 28: light chain variable domain VL, HLA-G-0090-S33D

DIVMTQSPDSLAVSLGERATINCKSSQSVLNSDNNKNNLAWYQQQPGQPP
KLLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTP
WTFGQGTKVEIK

SEQ ID NO: 30: light chain variable domain VL, HLA-G-0090-S33P

25 DIVMTQSPDSLAVSLGERATINCKSSQSVLNSPNNKNNLAWYQQQPGQPP
KLLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTP
WTFGQGTKVEIK

The amino acid sequences of anti-CD3 binding moieties (variable regions with underlined complementarity determining regions (CDRs) :

SEQ ID NO: 58 heavy chain variable domain VH, P035-093 (abbreviates as P035)

5 EVQLLES GGGLVQP GGS LRLSCAASGFTFSS YAMNWVRQAPGKGLEWVS
RIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCV
RASNFPASYVSYFAYWGQGTLVTVSS

SEQ ID NO: 59 light chain variable domain VL, P035-093 (P035)

10 QAVVTQEPSLTVSPGGTVTLTC GSSTGAVTTSNYANWVQEKPQAFRGLI
GGTNKRAPGTPARFSGSLLGGKAALTLGAQPEDEAEYYC ALWYSNLWV
FGGGTKLTVL

SEQ ID NO: 66 heavy chain variable domain VH, Clone 22 (abbreviated as Cl22)

15 EVQLLES GGGLVQP GGS LRLSCAASGFQFSS YAMNWVRQAPGKGLEWVS
RIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCV
RHTTFPSSYVSYYGYWGQGTLVTVSS

SEQ ID NO: 67 light chain variable domain VL, Clone 22 (Cl22)

20 QAVVTQEPSLTVSPGGTVTLTC GSSTGAVTTSNYANWVQEKPQAFRGLI
GGTNKRAPGTPARFSGSLLGGKAALTLGAQPEDEAEYYC ALWYSNLWV
FGGGTKLTVL

SEQ ID NO: 74 heavy chain variable domain VH, V9

EVQLVES GGGLVQP GGS LRLSCAASGYST GYTMNWVRQAPGKGLEWVA
LINPYKGVSTYNQKFKDRFTISVDKSKNTAYLQMNSLRAEDTAVYYCARS
GYYGDS DWYFDVWGQGTLVTVSS

25 SEQ ID NO: 75 light chain variable domain VL, V9

DIQMTQSPSSLSASVGDRVTITC RASQDIRNYLNWYQQKPGKAPKLLIYYT
SRLESGVPSRFSGSGSGTDYTLTISSLPEDFATYYC CQQGNTLPWTFGQGTK
VEIK

The amino acid sequences of Bispecific anti-HLA-G/anti-CD3 T cell bispecific (TCB) antibodies:

P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035-093 (P035)):

SEQ ID NO: 76 light chain 1 P1AF7977

5 EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMNWVRQAPGKGLEWVS
RIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCV
RASNFPASYVSFYAWGQGLVTVSSASVAAPSVFIFPPSDEQLKSGTASV
VCLLNNFYPRKAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLS
KADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

10 SEQ ID NO: 77 light chain 2 P1AF7977

DIVMTQSPDSLAVSLGERATINCKSSQSVLNPSNNKNNLAWYQQQPGQPP
KLLIYWASTRESGVDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYRTP
WTFGQGTKVEIKRTVAAPSVFIFPPSDRKLKSGTASVVCLLNNFYPRKAKV
QWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEEKHKVYACEV
15 THQGLSSPVTKSFNRGEC

SEQ ID NO: 78 heavy chain 1 P1AF7977

QVQLQQSGPGLLKPSQTLSTCAISGDSVSSNRAAWNWIQSPSRGLEWLG
RTYYRSKQYNDYAVSVQGRITLIPDTSKNQFSLRLNSVTPEDTAVYYCASV
RAVAPFDYWGGVLTIVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVED
20 YFPEPVTIVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYIC
NVNHKPSNTKVDEKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTL
MISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY
RVVSVLTIVLHQQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQVCT
LPPSRDELTKNQVSLSCAVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD
25 GSFFLVSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSP

SEQ ID NO: 79 heavy chain 2 P1AF7977

QVQLQQSGPGLLKPSQTLSTCAISGDSVSSNRAAWNWIQSPSRGLEWLG
RTYYRSKQYNDYAVSVQGRITLIPDTSKNQFSLRLNSVTPEDTAVYYCASV
RAVAPFDYWGGVLTIVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVED
30 YFPEPVTIVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYIC
NVNHKPSNTKVDEKVEPKSCDGGGGSGGGGGQAVVTQEPSTVSPGGTV
TLTGSSTGAVTTSNYANWVQEKPGQAFRGLIGGTNKRAPGTPARFSGSL
GGKAALTLGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLSSASTKGPS
VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTIVSWNSGALTSGVHTFPAVLQ
35 SSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDEKVEPKSCDKTHTC
PPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNW
YVDGVEVHNAKTKPREEQYNSTYRVVSVLTIVLHQQDWLNGKEYKCKVSN
KALGAPIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPSDI
AVEWESNGQPENNYKTTTPVLDSDGSFFLVSKLTVDKSRWQQGNVFSCSV
40 MHEALHNHYTQKSLSLSP

- 110 -

P1AF7978 (HLA-G-0090-VL-S32P/ CD3 Clone 22 (Cl22)):

SEQ ID NO: 80 light chain 1 P1AF7978

5 EVQLLESQGGGLVQPGGSLRLSCAASGFOFSSYAMNWVRQAPGKGLEWVS
RIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCV
RHTTFPSSYVSYYGYWGQGLTVTVSSASVAAPSVFIFPPSDEQLKSGTASV
VCLLNPFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLS
KADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

SEQ ID NO: 81 light chain 2 P1AF7978

10 DIVMTQSPDSLAVSLGERATINCKSSQSVLNPSNNKNNLAWYQQQPGQPP
KLLIYWASTRESGVDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTP
WTFGQGTKVEIKRTVAAPSVFIFPPSDRKLKSGTASVVCLLNPFYPREAKV
QWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEEKHKVYACEV
THQGLSSPVTKSFNRGEC

SEQ ID NO: 82 heavy chain 1 P1AF7978

15 QVQLQQSGPGLLKPSQTLSTCAISGDSVSSNRAAWNWIRQSPSRGLEWLG
RTYYRSKQWYNDYAVSVQGRITLIPDTSKNQFSLRLNSVTPEDTAVYYCASV
RAVAPFDYWGQGVLTSSASTKGPSVFPLAPSSKSTSGGTAALGCLVED
YFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYIC
NVNHKPSNTKVDEKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTL
20 MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY
RVVSVLTVLHQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQVCT
LPPSRDELTKNQVSLSCAVKGFPYPSDIAVEWESNGQPENNYKTPPVLDSD
GSFFLVSKLTVDKSRWQQGNVFSFCSVMHEALHNHYTQKSLSLSP

SEQ ID NO: 83 heavy chain 2 P1AF7978

25 QVQLQQSGPGLLKPSQTLSTCAISGDSVSSNRAAWNWIRQSPSRGLEWLG
RTYYRSKQWYNDYAVSVQGRITLIPDTSKNQFSLRLNSVTPEDTAVYYCASV
RAVAPFDYWGQGVLTSSASTKGPSVFPLAPSSKSTSGGTAALGCLVED
YFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYIC
NVNHKPSNTKVDEKVEPKSCDGGGGSGGGGGQAVVTQEPSTVSPGGTV
30 TLTCGSSTGAVTTSNYANWVQEKPGQAFRGLIGGTNKRAPGTARFSGSL
GGKAALTLGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLSSASTKGPS
VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ
SSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDEKVEPKSCDKTHTC
PPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW
35 YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN
KALGAPIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPSDI
AVEWESNGQPENNYKTPPVLDSDGSFFLVSKLTVDKSRWQQGNVFSFCSV
MHEALHNHYTQKSLSLSP

P1AF7979 (HLA-G-0090-VL-S32P/ CD3 V9):

40 SEQ ID NO: 84 light chain 1 P1AF7979

- 111 -

EVQLVESGGGLVQPGGSLRLSCAASGYSTGYTMNWVRQAPGKGLEWVA
LINPYKGVSTYNQKFKDRFTISVDKSKNTAYLQMNSLRAEDTAVYYCARS
GYYGDSWDYFDVWGQGTLLTVSSASVAAPSVFIFPPSDEQLKSGTASVVC
LLNFPYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTLSSTLTLKA
5 DYEKHKVYACEVTHQGLSSPVTKSFNRGEC

SEQ ID NO: 85 light chain 2 P1AF7979

DIVMTQSPDSLAVSLGERATINCKSSQSVLNPSNNKNNLAWYQQQPGQPP
KLLIYWASTRESGVDPDRFSGSGSGTDFTLTISSLQAEDVAVYFCQQYYRTP
WTFGQGTKVEIKRTVAAPSVFIFPPSDRKLKSGTASVVCLLNFPYPREAKV
10 QWKVDNALQSGNSQESVTEQDSKDSSTLSSTLTLKADYEKHKVYACEV
THQGLSSPVTKSFNRGEC

SEQ ID NO: 86 heavy chain 1 P1AF7979

QVQLQQSGPGLLKPSQTLSTCAISGDSVSSNRAAWNWRQSPSRGLEWLG
RTYYRSKWYNDYAVSVQGRITLIPDTSKNQFSRLNSVTPEDTAVYYCASV
15 RAVAPFDYWGGQVLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVED
YFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYIC
NVNHKPSNTKVDEKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTL
MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY
RVVSVLTVLHQDWLNGKEYKCKVSNKALGAPIEKTKAKGQPREPQVCT
20 LPPSRDELTKNQVSLSCAVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD
GSFFLVSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSP

SEQ ID NO:87 heavy chain 2 P1AF7979

QVQLQQSGPGLLKPSQTLSTCAISGDSVSSNRAAWNWRQSPSRGLEWLG
RTYYRSKWYNDYAVSVQGRITLIPDTSKNQFSRLNSVTPEDTAVYYCASV
25 RAVAPFDYWGGQVLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVED
YFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYIC
NVNHKPSNTKVDEKVEPKSCDGGGSGGGGGDIQMTQSPSSLSASVGDVR
TITCRASQDIRNYLNWYQQKPKAPKLLIYYTSRLESGVPSRFSGSGSGTDY
TLTISSLQPEDFATYYCQQGNTLPWTFGQGTKVEIKSSASTKGPSVFPLAPS
30 SKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
SSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPE
AAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE
VHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGAPI
EKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWES
35 NGQPENNYKTTPPVLDSDGSFFLVSKLTVDKSRWQQGNVFSCSVMHEALH
NHYTQKSLSLSP

In the following specific embodiments of the invention are listed:

1. An antibody that binds to human HLA-G comprising

- 112 -

5 A) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:23; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6, or

10 B) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:25; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6.

2. The antibody according to embodiment 1, wherein the antibody

15 A) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24; or

B) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26.

20 3. The antibody according to any one of embodiments 1 or 2, wherein the antibody comprises a Fc domain of human origin, particularly of the IgG isotype, more particularly of the IgG1 isotype.

4. The antibody according to any one of embodiments 1 or 2, wherein the antibody comprises a constant region of human origin, particularly of the IgG isotype, more particularly of the IgG1 isotype, comprising a human CH1, CH2, CH3 and/or CL domain..

25 5. The antibody according to any one of embodiment 1 to 4, wherein the antibody

30 a) has improved binding properties with respect to maximal binding (R_{max}) and/or binding affinity (K_D) compared to the (parental) antibody that comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:8 (as shown in Example 2).

- 113 -

- b) does not crossreact with a modified human HLA-G β 2M MHC I complex, wherein the HLA-G specific amino acids have been replaced by HLA-A consensus amino acids, the complex comprising SEQ ID NO:40 (as shown in Example 2); and/or
- 5 c) does not crossreact with a mouse H2Kd β 2M MHC I complex comprising SEQ ID NO:41 (as shown in Example 2); and/or
- d) does not crossreact with rat RT1A β 2M MHC I complex comprising SEQ ID NO:43 (as shown in Example 2).
6. The antibody according to any one of embodiments 1 to 3, wherein the
10 antibody
- a) inhibits ILT2 binding to (HLA-G expressed on) JEG3 cells (ATCC No. HTB36) (as shown in Example 5); or
- b) binds to (HLA-G expressed on) JEG3 cells (ATCC No. HTB36), and inhibits
15 ILT2 binding to (HLA-G expressed on) JEG-3 cells (ATCC No. HTB36) (as shown in Example 5).
7. The antibody according to any one of embodiments 1 to 4, wherein the antibody is a multispecific antibody (preferably a bispecific antibody).
8. The antibody according to embodiment 7, wherein the antibody is a bispecific antibody that binds to human HLA-G and to human CD3.
9. The antibody according to embodiment 7, wherein the antibody is a bispecific
20 antibody that binds to human HLA-G and to human CD3, comprising a first antigen binding moiety that binds to human HLA-G and a second antigen binding moiety that binds to human CD3, wherein the first antigen binding moiety that binds to human HLA-G comprises
- A) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid
25 sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:23; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID
30 NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6, or

- 114 -

5 B) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:25; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6;

and wherein the second antigen binding moiety that binds to a T cell activating antigen binds to human CD3 comprises

10 C) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:52, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:53, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:54; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:55; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:56 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:57, or

20 D) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:60, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:61, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:62; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:63; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:64 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:65, or

25 E) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:68, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:69, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:70; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:71; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:72 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:73.

30 10. The bispecific antibody according to embodiment 9,

wherein the first antigen binding moiety

A) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24; or

- 115 -

B) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26,

and wherein the second antigen binding moiety

5 C) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:58 and a VL domain comprising the amino acid sequence of SEQ ID NO:59; or

D) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:66 and a VL domain comprising the amino acid sequence of SEQ ID NO:67; or

E) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:74 and a VL domain comprising the amino acid sequence of SEQ ID NO:75.

10 11. The bispecific antibody according to embodiment 10,

wherein the first antigen binding moiety

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24;

and wherein the second antigen binding moiety

15 comprises a VH domain comprising the amino acid sequence of SEQ ID NO:58 and a VL domain comprising the amino acid sequence of SEQ ID NO:59.

12. The bispecific antibody according to embodiment 108,

wherein the first antigen binding moiety

20 comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24;

and wherein the second antigen binding moiety

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:66 and a VL domain comprising the amino acid sequence of SEQ ID NO:67.

13. The bispecific antibody according to embodiment 10,

25 wherein the first antigen binding moiety

- 116 -

comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24;

and wherein the second antigen binding moiety

5 comprises a VH domain comprising the amino acid sequence of SEQ ID NO:74 and a VL domain comprising the amino acid sequence of SEQ ID NO:75.

14. The bispecific antibody according to any one of embodiments 8 to 13, wherein the bispecific antibody shows

a) inhibition of ILT2 and/or ILT4 binding to HLA-G (as shown in Example 13); and/or

10 b) antibody mediated IFN gamma secretion by T cells on SKOV3 cells transfected with recombinant HLA-G (SKOV3 HLA-G) and/or on JEG3 cells expressing endogenous HLA-G wherein the IFN gamma secretion was detected (by Luminex technology) (as shown in Example 14); and or

15 c) T cell mediated cytotoxicity/tumor cell killing on SKOV3 cells transfected with recombinant HLA-G (SKOV3 HLA-G) and/or JEG3 cells expressing endogenous HLA-G wherein the cytotoxicity was detected by measuring Caspase 8 activation in cells after treatment with bispecific antibody (as shown in Example 15); and/or

20 d) in vivo anti-tumor efficacy/ tumor regression in humanized NSG mice bearing SKOV3 human ovarian carcinoma transfected with recombinant HLA-G (SKOV3 HLA-G) humanized NSG mice (as shown in Example 16); and/or

e) in vivo anti-tumor efficacy /tumor of HLA-G CD3 T cell bi-specific in humanized NSG mice bearing human breast cancer PDX tumors (BC004) (as shown in Example 17).

25 15. The bispecific antibody of any one of embodiments 9 to 14, wherein the first and the second antigen binding moiety is a Fab molecule.

16. The bispecific antibody of any one of embodiments 9 to 15, wherein the second antigen binding moiety is a Fab molecule wherein the variable domains VL and VH or the constant domains CL and CH1, particularly the variable domains VL and VH, of the Fab light chain and the Fab heavy chain are replaced by each other.

17. The bispecific antibody of any one of embodiments 9 to 16, wherein the first antigen binding moiety is a Fab molecule wherein in the constant domain the amino acid at position 124 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat) and the amino acid at position 123 is substituted independently by lysine (K), arginine (R) or histidine (H) (numbering according to Kabat), and in the constant domain CH1 the amino acid at position 147 is substituted independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index) and the amino acid at position 213 is substituted independently by glutamic acid (E), or aspartic acid (D) (numbering according to Kabat EU index).
18. The bispecific antibody of any one of embodiments 9 to 17, comprising a third antigen binding moiety wherein the third antigen moiety is identical to the first antigen binding moiety.
19. The bispecific antibody of any one of embodiments 9 to 18, comprising an Fc domain composed of a first and a second subunit.
20. The bispecific antibody of embodiment 19 wherein the first, the second and, where present, the third antigen binding moiety are each a Fab molecule; and wherein either (i) the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain, or (ii) the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety and the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain; and wherein the third antigen binding moiety, where present, is fused at the C-terminus of the Fab heavy chain to the N-terminus of the second subunit of the Fc domain.
21. The bispecific antibody of embodiment 19 or 20, wherein the Fc domain is a human IgG Fc domain, particularly of the IgG1 isotype.
22. The bispecific antibody of any one of embodiments 19 or 20, wherein the Fc domain comprises one or more amino acid substitution that reduces binding to an Fc receptor and/or effector function.

23. The bispecific antibody according embodiment 22, wherein the antibody is of the IgG1 isotype with mutations L234A, L235A and P329G (numbering according to the EU index of Kabat).
24. The bispecific antibody of any one of embodiments 19 to 23, wherein an amino acid residue in the CH3 domain of the first subunit of the Fc domain is replaced with an amino acid residue having a larger side chain volume, thereby generating a protuberance within the CH3 domain of the first subunit which is positionable in a cavity within the CH3 domain of the second subunit, and an amino acid residue in the CH3 domain of the second subunit of the Fc domain is replaced with an amino acid residue having a smaller side chain volume, thereby generating a cavity within the CH3 domain of the second subunit within which the protuberance within the CH3 domain of the first subunit is positionable.
25. The bispecific antibody according embodiment 24, wherein the antibody is of IgG1 isotype with mutation T366W in the first subunit of the Fc domain and with mutations Y407V, T366S and L368A in the second subunit of the Fc domain (numberings according to Kabat EU index).
26. The bispecific antibody according embodiment 25, wherein the antibody comprises an additional mutation S354C in the first subunit of the Fc domain and an additional mutation Y349C in the second subunit of the Fc domain (numberings according to Kabat EU index).
27. The bispecific antibody according embodiment 25, wherein the antibody comprises an additional mutation Y349C in the first subunit of the Fc domain and an additional S354C mutation in the second subunit of the Fc domain (numberings according to Kabat EU index).
28. Isolated nucleic acid encoding the antibody according to any one of embodiments 1-4 or the bispecific antibody according to any one of embodiments 9-27.
29. A host cell, preferably an eukaryotic host cell, comprising the nucleic acid of embodiment 28.
30. A method of producing the antibody according to any one of embodiments 1-4 or the bispecific antibody according to any one of embodiments 9-27 comprising culturing the host cell of embodiment 29 so that the antibody or bispecific antibody is produced.

31. The method of embodiment 30, further comprising recovering the antibody or bispecific antibody from the host cell.
32. The antibody according to any one of embodiments 1-4 or the bispecific antibody according to any one of embodiments 9-27, wherein the antibody is produced according to a method of embodiments 30 to 31 and wherein the host cell is an eukaryotic host cell (in one preferred embodiment a mammalian host cell, in another preferred embodiment a CHO cell).
33. The antibody according to any one of embodiments 1-4 or the bispecific antibody according to any one of embodiments 9-27, wherein the antibody is produced in an eukaryotic host cell (in one preferred embodiment a mammalian host cell, in another preferred embodiment a CHO cell).
34. The antibody according to any one of embodiments 1-4 or the bispecific antibody according to any one of embodiments 9-27 for use as a medicament.
35. The antibody according to any one of embodiments 1-4 or the bispecific antibody according to any one of embodiments 9-27 for use in treating cancer.
36. Use of the antibody according to any one of embodiments 1-4 or the bispecific antibody according to any one of embodiments 9-27 in the manufacture of a medicament.
37. The use of embodiment 36, wherein the medicament is for treatment of cancer.
38. A method of treating an individual having cancer comprising administering to the individual an effective amount of the antibody according to any one of embodiments 1-4 or the bispecific antibody according to any one of embodiments 9-27.

Examples**Recombinant DNA techniques**

5 Standard methods were used to manipulate DNA as described in Sambrook, J. et al., Molecular cloning: A laboratory manual; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989. The molecular biological reagents were used according to the manufacturer's instructions.

Gene and oligonucleotide synthesis

10 Desired gene segments were prepared by chemical synthesis at Geneart GmbH (Regensburg, Germany). The synthesized gene fragments were cloned into an E. coli plasmid for propagation/amplification. The DNA sequences of subcloned gene fragments were verified by DNA sequencing. Alternatively, short synthetic DNA fragments were assembled by annealing chemically synthesized oligonucleotides or via PCR. The respective oligonucleotides were prepared by metabion GmbH (Planegg-Martinsried, Germany)

15 **Description of the basic/standard mammalian expression plasmid**

For the expression of a desired gene/protein (e.g. full length antibody heavy chain, full length antibody light chain, or an MHC class I molecule, e.g. HLA-G, or an MHC class I molecule fused to peptide and beta-2 microglobulin, e.g. HLA-G fused to HLA-G binding peptide and or beta-2 microglobulin) a transcription unit
20 comprising the following functional elements is used:

- the immediate early enhancer and promoter from the human cytomegalovirus (P-CMV) including intron A,
- a human heavy chain immunoglobulin 5'-untranslated region (5'UTR),
- a murine immunoglobulin heavy chain signal sequence,
- 25 – a gene/protein to be expressed (e.g. full length antibody heavy chain or MHC class I molecule), and
- the bovine growth hormone polyadenylation sequence (BGH pA).

Beside the expression unit/cassette including the desired gene to be expressed the basic/standard mammalian expression plasmid contains

- 121 -

- an origin of replication from the vector pUC18 which allows replication of this plasmid in *E. coli*, and
- a beta-lactamase gene which confers ampicillin resistance in *E. coli*.

Protein determination

- 5 The protein concentration of purified polypeptides was determined by determining the optical density (OD) at 280 nm, using the molar extinction coefficient calculated on the basis of the amino acid sequence of the polypeptide.

Example 1

Generation of HLA-G chimeric molecules

- 10 Due to high homology (>98%) with other MHC I molecules, immunisation with HLA-G molecules results in generation of polyclonal sera, composed of a mixture of MHC-I crossreactive antibodies as well as truly HLA-G specific antibodies.

So far no tools have been provided to select truly HLA-G specific antibodies without crossreactivity to other human MHC-I (e.g. HLA-A), and to further select those with
15 receptor blocking function.

We identified unique HLA-G positions in combination to positions necessary for structural conformity and receptor interaction (ILT2/4 and KIR2DL4.)

- 20 Unique and proximal positions of human HLA-G were then „grafted“ on MHC class I complex molecules from different rodent species (such as rat RT1A and mouse H2kd) to generate „chimeric“ immunogen/screening antigens.

Antibodies generated were subjected to stringent screening/testing for binding/specificity, (and no crossreactivity/ no specificity to counterantigens, respectively)

antigens for binding testing:

- 25 – rec. HLA-G expressed as human HLA-G β 2M MHC complex comprising SEQ ID NO: 43
- HLA-G specific sequences grafted onto rat RT-1 and mouse H2kd (SEQ ID NO: 46: human HLA-G/ mouse H2Kd β 2M MHC class I complex wherein the positions specific for human HLA-G are grafted onto the mouse H2Kd

- 122 -

framework and SEQ ID NO: 48: human HLA-G/ rat RT1A β 2M MHC class I complex wherein the positions specific for human HLA-G are grafted onto the rat RT1A framework)

- 5 – Natural HLA-G MHC class I complex expressing cells (e.g. Jeg3 cells), or human HLA-G transfected cell lines SKOV3 HLA-G⁺ and PA-TU-8902 HLA-G⁺

counter antigens for crossreactivity testing:

- 10 – Counter antigens (MHC class I complexes) with other HLA-A sequences (HLA-A2 and HLA-G^{degrafted} with HLA-A consensus sequence) combined with different peptides) (see e.g. SEQ ID NO 35 (HLA-A2) and SEQ ID NO: 40 HLA-A consensus sequence on HLA-G framework)
- Counter antigens (MHC class I complexes) from other species such as rat RT-1 and mouse H2kd (SEQ ID NO: 43 and SEQ ID NO: 41)
- 15 – Unmodified tumor cell lines SKOV3 and PA-TU-8902, which are characterized by absence of HLA-G expression.

Design of chimeric HLA-G antigens to determine the specific binding of anti-HLA-G antibodies (see Figure 2):

Design of a chimeric rat MHC I molecule (RT1-A) carrying HLA-G unique positions (SEQ ID NO: 44) for use in for use in binding assays:

- 20 HLA-G unique positions were identified by the alignment of 2579 HLA-A, 3283 HLA-B, 2133 HLA-C, 15 HLA-E, 22 HLA-F, and 50 HLA-G sequences from IMGT (as available on 6. Feb 2014). Those residues of HLA-G that occur in less than 1% (mostly ~0%) of the sequences of any of the 3 sequence sets HLA-A, HLA-B, and a combined set of HLA-C + HLA-E + HLA-F are called HLA-G unique positions.
- 25 The 4 core HLA-G unique positions (2 in alpha-1 and 2 in alpha-3) show no polymorphism in the set of HLA-G sequences and none of the other HLA genes contain the HLA-G specific residues at these positions (except 1x HLA-A for M100, 1x HLA-B for Q103, and 1x HLA-C for Q103).

- 30 The crystal structure of rat RT1-A (Rudolph, M.G. et al. J.Mol.Biol. 324: 975-990 (2002); PDB code: 1KJM) was superimposed on the crystal structure of human HLA-G (Clements, C.S. et al. PROC.NATL.ACAD.SCI.USA 102: 3360-3365

(2005); PDB code: 1YDP). The overall structure of the alpha-chain and the associated beta-2-microglobulin is conserved.

5 HLA-G unique positions were identified in the RT1-A structure by comparison of the sequence and structural alignments. In a first step, unique HLA-G positions were identified that are exposed on the molecular surface of HLA-G and RT1-A and thus accessible for an antibody. Unique positions that are buried within the protein fold were excluded for engineering. In a second step, structurally proximal residues were identified, that also need to be exchanged to make the corresponding region „HLA-G-like“, i.e. to generate real HLA-G epitopes containing the unique positions rather than generating HLA-G/rat RT1-A chimeric epitopes that would be artificial. All the positions that were thus selected for mutation were analyzed for structural fit of the respective residue from HLA-G to avoid possible local disturbances of the molecular structure upon mutation.

15 A chimeric mouse MHC I molecule (H2Kd) carrying HLA-G unique positions (SEQ ID NO: 42) for use in binding assays was generated analogously.

Design of HLA-A based counter antigens by “de-grafting” of HLA-G unique positions towards a HLA-A consensus sequence for crossreactivity testing (SEQ ID NO:40 =modified human HLA-G β 2M MHC class I complex (wherein the HLA-G specific amino acids have been replaced by HLA-A consensus amino acids (= degrafted HLA-G))

20 Unique positions derived from the multiple sequence alignment were analyzed in a crystal structure of human HLA-G (PDB code: 1YDP). First, positions that are not exposed on the HLA-G surface and are thus not accessible for an antibody were excluded for engineering. Second, the surface exposed residues were analyzed for feasibility of amino acid exchange (i.e. exclusion of possible local disturbances of the molecular structure upon mutation of the relevant position). In total, 14 positions were validated for exchange. The amino acids in the validated positions were mutated towards a HLA-A consensus sequence derived from a multiple sequence alignment of 2579 HLA-A sequences downloaded from IMGT (as available on 6. Feb 2014).

Generation of expression plasmids for soluble classical and non-classical MHC class I molecules

5 The recombinant MHC class I genes encode N-terminally extended fusion molecules consisting of a peptide known to be bound by the respective MHC class I molecule, beta-2 microglobulin, and the respective MHC class I molecule.

10 The expression plasmids for the transient expression of soluble MHC class I molecules comprised besides the soluble MHC class I molecule expression cassette an origin of replication from the vector pUC18, which allows replication of this plasmid in *E. coli*, and a beta-lactamase gene which confers ampicillin resistance in *E. coli*.

The transcription unit of the soluble MHC class I molecule comprised the following functional elements:

- the immediate early enhancer and promoter from the human cytomegalovirus (P-CMV) including intron A,
- 15 - a human heavy chain immunoglobulin 5'-untranslated region (5'UTR),
- a murine immunoglobulin heavy chain signal sequence,
- an N-terminally truncated *S. aureus* sortase A encoding nucleic acid, and
- the bovine growth hormone polyadenylation sequence (BGH pA).

20 The amino acid sequences of the mature soluble MHC class I molecules derived from the various species are:

SEQ ID NO: 39: exemplary human HLA-G β 2M MHC class I complex

SEQ ID NO: 40: exemplary modified human HLA-G β 2M MHC class I complex (wherein the HLA-G specific amino acids have been replaced by HLA consensus amino acids (= degrafted HLA-G see also Figure 2)

25 SEQ ID NO: 41: exemplary mouse H2Kd β 2M MHC class I complex

SEQ ID NO: 42: exemplary human HLA-G/ mouse H2Kd β 2M MHC complex wherein the positions specific for human HLA-G are grafted onto the mouse H2Kd framework

SEQ ID NO: 43: exemplary rat RT1A β 2M MHC class I complex

SEQ ID NO: 44: exemplary human HLA-G/ rat RT1A β 2M MHC complex wherein the positions specific for human HLA-G are grafted onto the rat RT1A framework

- 5 For the exemplary HLA-A2 β 2M MHC class I complex the following components were used and the complex was expressed in E.Coli and purified.

MHCI complex HLA-A2 / β 2M (SEQ ID NOs 35 and 33) (both with an additional N-terminal methionine) + VLDFAPPGA peptide (SEQ ID NO: 46) + linker and his-Tag (SEQ ID NO: 45)

10 **Example 2**

Removal of N-glycosylation motif in CDR-L1

- The CDR-L1 of anti-HLA-G antibody HLA-G-0090 contains a classical N-glycosylation motif "NSS" comprising positions 31 to 33 of the light chain (LC). It was decided to remove this motif as it could constitute a potential developability liability. An homology model of the variable region of HLA-G-0090 indicated that LC positions 31 to 33 are highly solvent accessible. Furthermore, the side chains of N31 and S32 are predicted to point inwards, in the direction of CDR-H3, making them likely candidates for being part of the antibody paratope. In fact, a number of published antibody-antigen X-ray complex structures document these residues to be undergoing chemical interactions with the antigen. Therefore, the risk of worsening the binding affinity of the antibody by introducing mutations at LC positions 31-33 was considered high. To ameliorate the risk, 11 different variants of antibody HLA-G-0090 with mutations on LC positions 31, 32, and 33 were designed and produced in HEK293F cells in an IgG1 format.

- 25 Note that mutant variant HLA-G-0090-VL-N31Y-N38Y contained a second mutation (N38Y), apart from the N-glycosylation motif, and not related to its removal, to increase germline identity.

Summary of anti-HLA-G antibody sequences (SEQ ID NOs of variable regions and CDRs):

Anti-HLA-G antibody	CDR -H1	CDR -H2	CDR -H3	CDR -L1	CDR -L2	CDR -L3	VH	VL
	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:
HLA-G-0090	1	2	3	4	5	6	7	8
HLA-G-0090-VL-N31D	1	2	3	9	5	6	7	10
HLA-G-0090-VL-N31L	1	2	3	11	5	6	7	12
HLA-G-0090-VL-N31Q	1	2	3	13	5	6	7	14
HLA-G-0090-VL-N31S	1	2	3	15	5	6	7	16
HLA-G-0090-VL-N31T	1	2	3	17	5	6	7	18
HLA-G-0090-VL-N31Y	1	2	3	19	5	6	7	20
HLA-G-0090-VL-N31Y-N38Y	1	2	3	21	5	6	7	22
HLA-G-0090-VL-S32P	1	2	3	23	5	6	7	24
HLA-G-0090-VL-S33A	1	2	3	25	5	6	7	26

- 127 -

Anti-HLA-G antibody	CDR -H1	CDR -H2	CDR -H3	CDR -L1	CDR -L2	CDR -L3	VH	VL
	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:	SEQ ID NO:
HLA-G-0090-VL-S33D	1	2	3	27	5	6	7	28
HLA-G-0090-VL-S33P	1	2	3	29	5	6	7	30

Binding and other properties of the obtained anti-HLA-G specific antibodies and biological activities were determined as described in the following Examples, and compared to the known reference, HLA-G-0090.

Expression and purification

The expression yields from a 0.5 L expression in HEK293F cells after purification by affinity chromatography (MabSelect Sure) and dialysis are shown in the following table.

	mg	Monomer Content [%] (analytical SEC)	Purity [%] (Caliper)
HLA-G-0090	3.2	98	99
HLA-G-0090-VL-N31D	0.4	97	98
HLA-G-0090-VL-N31L	0.4	98	99
HLA-G-0090-VL-N31Q	0.4	91	95
HLA-G-0090-VL-N31S	0.3	93	98
HLA-G-0090-VL-N31T	0.4	85	89
HLA-G-0090-VL-N31Y	0.3	89	94
HLA-G-0090-VL-N31Y-N38Y	0.7	94	98
HLA-G-0090-VL-S32P	2.4	98	99
HLA-G-0090-VL-S33A	2.1	98	99
HLA-G-0090-VL-S33D	2.4	98	99
HLA-G-0090-VL-S33P	1.4	98	99

Variants with mutations involving position LC 31 (N31X) showed strongly decreased expression titers and, often, impaired material quality, while LC 32 and

LC33 positions variant showed good to acceptable expression titers and good material quality.

HLA-G Binding

wt-HLA-G Affinity/Kinetic

- 5 Affinity to wt-HLA-G complex (SEQ ID NO: 39) of individual 5nM anti-HLA-G antibodies was determined by capturing with anti-hFc (GE Healthcare BR-1008-39) on a CM3 sensor chip and the injection of wt-HLA-G antigen at a concentration of 11 nM to 300nM diluted in HBS-P+ (GE Healthcare) running buffer and a flow rate of 60 µl/min with 120 s association time and 600 s dissociation time. After each cycle
10 the surface was regenerated by washing with 3M MgCl₂. The kinetics binding curves were evaluated using T200 evaluation software and for the calculation of binding properties 1:1 Langmuir binding model was used.

RAC of HLA-G antibodies (relative active concentration) (assay scheme is shown in Fig. 3)

- 15 RAC of HLA-G binders of individual anti-HLA-G antibodies (10 nM solutions in HBS-P+) was determined by capturing with anti-hFc (GE Healthcare BR-1008-39) on a CM3 sensor chip and the injection of wt-HLA-G antigen at a concentration of 300nM diluted in HBS-P+ (GE Healthcare) running buffer and a flow rate of 10 µl/min with 60 s association time and 600 s dissociation time. After each cycle the
20 surface was regenerated by washing with 3M MgCl₂. Final RAC and R_{max} values are calculated from “binding” report points and the capturing levels. $R_{max} = (MW \text{ analyte} / MW \text{ ligand}) * RU \text{ capturing ligand} * \text{stoichiometry of interaction}$.

- The 11 variants were evaluated in an SPR binding experiment in which the antibodies were immobilized on a CM3 chip via the Fc part and recombinant single-chain HLA-G monomer/ human HLA-G β2M MHC class I complex (SEQ ID NO: 39) was used
25 as the analyte. The kinetic parameters were determined by a single cycle kinetic measurement on a Biacore T200 device at 25 °C.

	ka [1/ms]	kd [1/s]	t ½ [s]	Kd [nM]	Rmax [%]
HLA-G-0090	1.19E+06	1.38E-03	501	1.2	86
HLA-G-0090-VL-N31D	1.94E+05	3.91E-03	177.3	20.2	66
HLA-G-0090-VL-N31L	5.28E+06	8.71E-03	79.6	1.7	91
HLA-G-0090-VL-N31Q	1.91E+06	3.73E-03	185.9	2.0	91
HLA-G-0090-VL-N31S	3.83E+05	9.22E-04	751.8	2.4	77
HLA-G-0090-VL-N31T	3.42E+05	7.69E-04	901.9	2.3	75
HLA-G-0090-VL-N31Y	5.49E+05	1.07E-03	646.2	2.0	78
HLA-G-0090-VL-N31Y-N38Y-GL	1.82E+09	4.30E+01	0	23.6	57
HLA-G-0090-VL-S32P	1.19E+06	1.24E-03	557.7	1.0	97
HLA-G-0090-VL-S33A	1.24E+06	1.24E-03	558.2	1.0	98
HLA-G-0090-VL-S33D	7.17E+05	3.44E-03	201.7	4.8	95

- 131 -

	ka [1/ms]	kd [1/s]	t ½ [s]	Kd [nM]	Rmax [%]
HLA-G-0090-VL-S33P	1.34E+06	2.36E-03	294.3	1.8	94

Among those, the four variants with acceptable expression titers (HLA-G-0090-VL-S32P, HLA-G-0090-VL-S33A, HLA-G-0090-VL-S33D, HLA-G-0090-VL-S33P) were evaluated further in a more accurate multi cycle kinetics measurement using the same SPR device and experimental setup.

5

	ka [1/ms]	kd [1/s]	t ½ [s]	Kd [nM]	Rmax [%]
HLA-G-0090	1.30E+06	1.78E-03	389.9	1.4	87
HLA-G-0090-VL-S32P	1.27E+06	1.57E-03	440.6	1.2	99
HLA-G-0090-VL-S33A	1.24E+06	1.49E-03	465.9	1.2	99
HLA-G-0090-VL-S33D	8.97E+05	6.67E-03	103.9	7.4	98
HLA-G-0090-VL-S33P	1.36E+06	3.68E-03	188.1	2.7	97

The two variants HLA-G-0090-VL-S32P and HLA-G-0090-VL-S33A have an improved binding affinity compared to the parental antibody HLA-G-0090 while the two variants HLA-G-0090-VL-S33D and HLA-G-0090-VL-S33P are losing binding affinity by a factor of 5 and 2, respectively. Surprisingly, for the tested variants, the removal of the N-glycosylation motif leads to a higher Rmax value in the kinetics measurement, indicating a higher fraction of successful complex formation than for the N-glycosylated antibody.

10

Crossreactivity of anti HLA-G antibodies and variants to soluble human HLA-G, soluble degrafted human HLA-G with HLA-A consensus specific sequence, and rat/ mouse homologues

15

To further investigate the binding properties of the two binding-affinity improved variants HLA-G-0090-VL-S32P and HLA-G-0090-VL-S33A, SPR binding

5 experiments to four counter-screening constructs were performed. These recombinant single-chain peptide-MHC complex constructs were murine H2-K1 (SEQ ID NO: 41), rat RT1 (SEQ ID NO: 43), and human HLA-G β 2M MHC class I complex, wherein the HLA-G specific amino acids have been replaced by HLA-A consensus amino acids (SEQ ID NO:40). The latter construct constitutes a version of HLA-G in which all HLA-G specific residues have been replaced by their HLA-A consensus counterparts. Again, the antibodies were immobilized on CM3 chip and the single-chain peptide-MHC constructs were used as analyte on a Biacore T200 device at 25° C.

Antigen	Antibody	Interaction
murine H2-K1 (SEQ ID NO:41)	HLA-G-0090	no binding interaction
murine H2-K1 (SEQ ID NO:41)	HLA-G-0090-VL-S32P	no binding interaction
murine H2-K1 (SEQ ID NO:41)	HLA-G-0090-VL-S33A	no binding interaction
rat RT1 (SEQ ID NO:43)	HLA-G-0090	no binding interaction
rat RT1 (SEQ ID NO:43)	HLA-G-0090-VL-S32P	no binding interaction
rat RT1 (SEQ ID NO:43)	HLA-G-0090-VL-S33A	no binding interaction
HLA-A consensus on HLA-G frame (SEQ ID NO:40)	HLA-G-0090	no binding interaction
HLA-A consensus on HLA-G frame (SEQ ID NO:40)	HLA-G-0090-VL-S32P	no binding interaction
HLA-A consensus on HLA-G frame (SEQ ID NO:40)	HLA-G-0090-VL-S33A	no binding interaction
HLA-G (SEQ ID NO:39)	HLA-G-0090	very strong binding interaction
HLA-G (SEQ ID NO:39)	HLA-G-0090-VL-S32P	very strong binding interaction

- 133 -

Antigen	Antibody	Interaction
HLA-G (SEQ ID NO:39)	HLA-G-0090-VL-S33A	very strong binding interaction

Stability under stress

5 The parental antibody as well as the two derived variants HLA-G-0090-VL-S32P and HLA-G-0090-VL-S33A were stressed for 13 days under two different conditions:

- pH 6.0 20 mM His/HisCl, 140 mM NaCl; at 40 °C (His 40 °C)
- pH 7.4 PBS; at 37 °C (PBS 37 °C).

10 Afterwards, the material was analysed using SEC, and SPR to investigate chemical degradation and possible effects on target binding. For reference, the stressed material was compared with material kept under storage conditions:

- pH 6.0 20 mM His/HisCl, 140 mM NaCl; frozen at -80°C (Ref.)

The results are listed in the following table (relative in % compared to Ref):

Parameter		HLA-G-0090_VL-S32P	HLA-G-0090_VL-S33A	HLA-G-0090
SEC monomer [%]	Ref.	100	99	99
	His 40 °C	98	98	98
	PBS 37 °C	98	98	98
relative HLA-G binding signal compared to Ref (as 100%) ± stress, by SPR RAC	Ref.	100	100	100
	His 40 °C	99	99	99
	PBS 37 °C	98	96	96

While all three antibodies are showing a very similar stability profile, variant HLAG-0090_VL-S32P is retaining more HLA-G binding (SPR relative active concentration (RAC)) after stress in PBS at 37 °C than the other two, including the parental antibody.

Thermal stability testing

For thermal stability testing of the purified proteins the Uncle device was used (UNCHAINED LABS, Boston, MA, USA). Static light scattering at 266 nm and 473 nm and in parallel intrinsic fluorescence is hereby used to determine aggregation temperature (Tagg) and melting temperature (Tm) of the purified proteins. A temperature ramp from 30°C to 90°C in 0.1°C /min steps was run. Glass cuvettes with 9 µl volume per samples were used and the concentration was 1 mg/ mL in 20 mM Histidin, 140 mM NaCl, pH 6.0 buffer. For analysis, the software UNcle analysis (UNCHAINED LABS) was used.

Mass spectrometry analysis and N-Glycosylation

The deconvoluted mass spectra of the intact samples are documenting the impact on the N-Glycosylation of the removal of the N-glycosylation site/NSS motif in HLA-G-0090_VL-S32P and HLA-G-0090_VL-S33A. The samples were prepared with PNGase F to remove all N-linked glycans and obtain the molecular mass of the antibody only. While PNGase F is fully specific for cleavage of N-linked Fc-glycans, it shows much less efficacy when cleaving N-linked Fab-glycans. HLA-G-0090 is showing a clear N-glycosylation pattern coming from incomplete deglycosylation and therefore indicating Fab-glycosylation ((see Figures 4A). No signs of residual N-glycosylation can be detected for HLA-G-0090_VL-S32P and HLA-G-0090_VL-S33A (see Figure 4B).

Example 3

ILT2 and -4 binding inhibition of anti-HLA-G antibodies

The ELISA is set up by coating the Fc tagged ILT2 and ILT4 respectively to Maxisorp microtiter plates. After incubation and washing steps, the respective antibodies are added at a concentration of 100nM. Soluble His tagged monomeric, dimeric or trimeric HLA-G was added to the wells. After incubation and washing steps, detection of bound receptor is carried out by anti-His-antibody-POD conjugates. Percentage inhibition (%) is calculated in comparison to values obtained from wells with ILT2/4 + HLA-G (mono-, di-, or Trimer) without anti HLA-G or ILT2/4 antibodies (100% binding = 0% inhibition) and shown in a table

Example 4

Binding of HLA-G antibodies to natural or recombinant HLA-G expressed on cells (as assessed by FACS analysis)

For flow cytometry analysis, cells were stained with anti HLA-G mAbs at 4°C. Briefly, each cell suspension was transferred into a polypropylene tube (2×10^5 cells/tube) and prechilled at 5°C for 10 minutes. Cells were then washed with 2ml FACS Buffer (4°C) and centrifuged at 300g for 5 minutes. Anti-HLA-G antibodies HLAG-0090-VL-S32P, HLAG-0090-VL-S33A, HLAG-0090 were diluted in staining buffer to a starting concentration of 50µg/ml. A 5-fold serial dilution of the antibodies was performed to get the final concentrations (10µg/ml, 2µg/ml, 0,4µg/ml, 0,08µg/ml, 0,016µg/ml, 0,0032µg/ml). FACS buffer was then aspirated from the tubes and the cell pellets were resuspended in 100µl of the antibody solution

and incubated for 1h at 5°C. Cells were then washed once with 2ml staining buffer and centrifuged at 300g for 5 minutes.

For detection fluorescent labeled anti-species antibody (goat anti-human IgG (H+L) conjugated to Alexa 488, Life technologies # A11013) was diluted to 10µg/ml in a staining buffer and cell pellets were resuspended in 100µl of detection antibody. After a 1 hour incubation at 5°C cells were again washed once with 2ml of staining buffer, resuspended in 500µl of staining buffer and measured on a FACS CELESTA

An exemplary FACS staining for anti-HLA-G antibodies HLA-G-0090, HLA-G-0090-VL-S32P and HLA-G-0090-VL-S32P (10µg/ml) is shown in the FACS overlays of Figure 5: Both deglycosylated variants of the HLA-G 0090, HLA-G-0090-VL-S32P and HLA-G-0090-VL-S32P show good binding to HLA-G expressing SKOV3 cells and JEG cells but not to parental SKOV3 cells. The MFI values of the respective HLA-G antibodies are indicated in the histograms.

Example 5

Anti HLA-G antibodies inhibit/modulate the binding of ILT2 to HLA-G expressed on JEG3 cells

For analysis, JEG3 cells (ATCC HTB36) were stained with ILT2-c-Myc-Fc fusion protein (control = no inhibition) with or without pre-incubation with different anti-HLA-G antibodies.

Binding/ Inhibition of binding was determined as follows: Recombinant ILT2-c-Myc-Fc protein was added to JEG3 cells either pre-incubated anti HLA-G mAbs as described or to untreated JEG3 cells as reference. For the pre-incubation with anti-HLA-G antibodies, 2×10^5 cells were transferred into a polypropylene tubes. Anti HLA-G antibodies HLAG-0090-VL-S32P, HLAG-0090-VL-S33A and HLA-G-0090 were diluted in staining buffer to a concentration of 80µg/ml and 25µl of the antibody solution was added to the prepared cells and incubated for 1h at 5°C. The ILT2-c-Myc-Fc or (control human IgG (Jackson-Immuno-Research # 009-000-003)) were diluted in staining buffer to a 2-fold concentration of 20µg/ml and added to the prepared cells at a final concentration of 10µg/ml and incubated for 2h at 5°C. Cells were washed twice with 200µl of staining buffer. Human ILT2-c-Myc-Fc protein was detected with fluorescent labeled Anti-Myc tag (9E10) Alexa Fluor 647 (abcam; # ab223895) at a dilution of 10µg/ml in staining buffer. Cells were resuspended in 50µl detection antibody dilution and incubated for 1 hour at 5°C. Cells were then

- 137 -

washed once with 2ml staining buffer and resuspended in 500µl of staining buffer before measuring at a FACS CELESTA

As control, the anti-HLA-G antibodies bound to JEG-3 pre-incubated cells were detected by using anti-species antibody (goat anti-human IgG (H+L) conjugated to Alexa 488, Life technologies # A11013), was diluted to 10µg/ml in staining buffer and cell pellets were resuspended in 100µl/well detection antibody. After a 1 hour incubation at 5°C cells were again washed once with staining buffer, resuspended in 500µl of staining buffer and measured at a FACS CELESTA

The histograms in Figure 6 show the respective ability of the HLA-G antibodies to modify/inhibit the interaction and binding of recombinant ILT2 to HLA-G naturally expressed on JEG3 tumor cells.

The following table summarizes the results from the experiments. The binding of the anti-HLA-G antibodies to JEG3 cells is depicted as + = weak binding - +++=strong binding. The ability of the anti-HLA-G antibodies either to inhibit/block or increase the binding of ILT2 to the HLA-G expressing JEG3 cells. In the last column, the binding of the recombinant ILT2 to the cells or the inhibition/blockade thereof is shown/quantified (staining of ILT2-c-Myc-Fc in the absence of an anti-HLA-G antibody was set to 100% binding = 0% inhibition):

Antibody	Binding to JEG-3 cells	HLA-G:ILT2 interaction	Inhibition of ILT2 binding to Jeg3 cells
ILT2-Fc w/o Antibody	-	-	0% inhibition
HLA-G-0090	+++	inhibits binding of ILT2	72 % inhibition
HLAG-0090-VL-S32P	+++	inhibits binding of ILT2	72 % inhibition
HLAG-0090-VL-S33A	+++	inhibits binding of ILT2	72 % inhibition

Example 6**Generation of optimized CD3 binder**

Starting from a previously described CD3 binder, termed “CD3_{orig}” herein (see for details e.g. WO2014/131712 incorporated herein by reference) comprising the VH and VL sequences of SEQ ID NOs 92 and 93 and we aimed at optimizing properties of this binder by removal of two asparagine deamidation sequence motifs at Kabat positions 97 and 100 of the heavy chain CDR3.

To this aim, we generated an antibody library, suitable for phage display, of the heavy chain with both asparagines at Kabat position 97 and 100 removed, and in addition the CDRs H1, H2, and H3 randomized in order to compensate for loss of affinity caused by replacing Asn97 and Asn100 through an affinity-maturation process.

This library was put on a filamentous phage via fusion to minor coat protein p3 (Marks et al. (1991) *J Mol Biol* **222**, 581-597) and selected for binding to recombinant CD3ε.

10 candidate clones were identified in the initial screening, showing acceptable binding on recombinant antigen as measured by SPR as Fab fragments (produced in *E. coli*).

Only one of these clones, however, showed acceptable binding activity to CD3 expressing cells as measured by flow cytometry after conversion to IgG format.

The selected clone, termed P035-093 (P035) (=“CD3_{opt}”) herein and comprising the VH and VL sequences of SEQ ID NOs 58 and 59, respectively, was further evaluated and converted into bispecific format as described in the following.

Example 7**Binding of optimized CD3 binder to CD3****Binding to recombinant CD3**

Binding to recombinant CD3 was determined by surface plasmon resonance (SPR) for the optimized CD3 binder P035-093 (P035) (=“CD3_{opt}”) and the original CD3

- 139 -

binder “CD3_{orig}”, both in human IgG1 format with P329G L234A L235A (“PGLALA”, EU numbering) mutations in the Fc region (SEQ ID NOs 94 and 96 (CD3_{orig}) and SEQ ID NOs 95 and 96 (P035=CD3_{opt})).

5 In order to assess the effect of the deamidation site removal and its effect on the stability of the antibodies, binding of the original and the optimized CD3 binder to recombinant CD3 was tested after temperature stress for 14 days at 37°C or 40°C. Samples stored at -80°C were used as reference. The reference samples and the samples stressed at 40°C were in 20 mM His, 140 mM NaCl, pH 6.0, and the samples stressed at 37°C in PBS, pH 7.4, all at a concentration of 1.2-1.3 mg/ml. After the
10 stress period (14 days) samples in PBS were dialyzed back to 20 mM His, 140 mM NaCl, pH 6.0 for further analysis.

Relative Active Concentration (RAC) of the samples was determined by SPR as follows.

SPR was performed on a Biacore T200 instrument (GE Healthcare). Anti-Fab
15 capturing antibody (GE Healthcare, #28958325) was immobilized on a Series S Sensor Chip CM5 (GE Healthcare) using standard amine coupling chemistry, resulting in a surface density of 4000 – 6000 resonance units (RU). As running and dilution buffer, HBS-P+ (10 mM HEPES, 150 mM NaCl pH 7.4, 0.05% Surfactant P20) was used. CD3 antibodies with a concentration of 2 µg/ml were injected for 60
20 s at a flow rate of 5 µl/min. CD3 antigen (see below) was injected at a concentration of 10 µg/ml for 120 s and dissociation was monitored at a flow rate of 5 µl/min for 120 s. The chip surface was regenerated by two consecutive injections of 10 mM glycine pH 2.1 for 60 s each. Bulk refractive index differences were corrected by subtracting blank injections and by subtracting the response obtained from the blank
25 control flow cell. For evaluation, the binding response was taken 5 seconds after injection end. To normalize the binding signal, the CD3 binding was divided by the anti-Fab response (the signal (RU) obtained upon capture of the CD3 antibody on the immobilized anti-Fab antibody). The relative active concentration was calculated by referencing each temperature stressed sample to the corresponding, non-stressed
30 sample.

- 140 -

The antigen used was a heterodimer of CD3 delta and CD3 epsilon ectodomains fused to a human Fc domain with knob-into-hole modifications and a C-terminal Avi-tag (see SEQ ID NOs 90 and 91).

5 The results of this experiment are shown in Figure 15. As can be seen, the optimized CD3 binder CD3_{opt} P035-093 (P035) (=CD3_{opt}) showed strongly improved binding to CD3 after temperature stress (2 weeks at 37°C, pH 7.4) as compared to the original CD3 binder CD3_{orig}. This result demonstrates that the deamidation site removal was successful, and has yielded an antibody with superior stability properties, relevant for in vivo half-life, as well as formulation of the antibody at neutral pH.

10 Binding to CD3 on Jurkat cells

Binding to CD3 on the human reporter T-cell line Jurkat NFAT was determined by FACS for the optimized CD3 binder P035-093 (P035) (=CD3_{opt}) and the original CD3 binder "CD3_{orig}", both in human IgG1 format with P329G L234A L235A ("PGLALA", EU numbering) mutations in the Fc region (SEQ ID NOs 94 and 96 (CD3_{orig}) and SEQ ID NOs 95 and 96 (P035=CD3_{opt})).

15 Jurkat-NFAT reporter cells (GloResponse Jurkat NFAT-RE-luc2P; Promega #CS176501) are a human acute lymphatic leukemia reporter cell line with a NFAT promoter, expressing human CD3. The cells were cultured in RPMI1640, 2g/l glucose, 2 g/l NaHCO₃, 10% FCS, 25 mM HEPES, 2 mM L-glutamine, 1 x NEAA, 20 1 x sodium-pyruvate at 0.1-0.5 mio cells per ml. A final concentration of 200 µg per ml hygromycin B was added whenever cells were passaged.

For the binding assay, Jurkat NFAT cells were harvested, washed with PBS and resuspended in FACS buffer. The antibody staining was performed in a 96-well round bottom plate. Therefore 100'000 to 200'000 cells were seeded per well. The plate was centrifuged for 4 min at 400 x g and the supernatant was removed. The test antibodies were diluted in FACS buffer and 20 µl of the antibody solution were added to the cells for 30 min at 4°C. To remove unbound antibody, the cells were washed twice with FACS buffer before addition of the diluted secondary antibody (PE-conjugated AffiniPure F(ab')₂ Fragment goat anti-human IgG Fcg Fragment Specific; Jackson ImmunoResearch #109-116-170). After 30 min incubation at 4°C

- 141 -

unbound secondary antibody was washed away. Before measurement the cells were resuspended in 200 μ l FACS buffer and then analyzed by flow cytometry using a BD Canto II device.

As shown in Figure 16, the optimized CD3 binder P035-093 (P035) (=CD3_{opt}) and the original CD3 binder "CD3_{orig}" bound comparably well to CD3 on Jurkat cells.

Example 8

Functional activity of optimized CD3 binder

The functional activity of the optimized CD3 binder "CD3_{opt}" was tested in a Jurkat reporter cell assay and compared to the activity of the original CD3 binder "CD3_{orig}". To test the functional activity of the IgGs, anti-PGLALA expressing CHO cells were co-incubated with Jurkat NFAT reporter cells in the presence of increasing concentrations of CD3_{opt} human IgG1 PGLALA or CD3_{orig} human IgG1 PGLALA. Activation of CD3 on the Jurkat NFAT reporter cells upon T cell cross-linking induces the production of luciferase and luminescence can be measured as an activation marker. CD3_{orig} human IgG1 wt was included as negative control which cannot bind to anti-PGLALA expressing CHO cells and therefore cannot be crosslinked on Jurkat NFAT cells. A schematic illustration of the assay is provided in Figure 17.

Anti-PGLALA expressing CHO cells are CHO-K1 cells engineered to express on their surface an antibody that specifically binds human IgG₁ Fc(PGLALA) (see WO 2017/072210, incorporated herein by reference). These cells were cultured in DMEM/F12 medium containing 5% FCS + 1% GluMax. The Jurkat NFAT reporter cells are as described in Example 7.

Upon simultaneous binding of the CD3 huIgG1 PGLALA to anti-PGLALA expressed on CHO and CD3 expressed on Jurkat-NFAT reporter cells, the NFAT promoter is activated and leads to expression of active firefly luciferase. The intensity of luminescence signal (obtained upon addition of luciferase substrate) is proportional to the intensity of CD3 activation and signaling. Jurkat-NFAT reporter cells grow in suspension and were cultured in RPMI1640, 2g/l glucose, 2 g/l NaHCO₃, 10 % FCS, 25 mM HEPES, 2 mM L-glutamin, 1 x NEAA, 1 x sodium-

- 142 -

pyruvate at 0.1-0.5 mio cells per ml, 200 µg per ml hygromycin. For the assay, CHO cells were harvested and viability determined using ViCell. 30 000 target cells/well were plated in a flat-bottom, white-walled 96-well-plate (Greiner bio-one #655098) in 100 µl medium and 50 µl/well of diluted antibodies or medium (for controls) were
5 added to the CHO cells. Subsequently, Jurkat-NFAT reporter cells were harvested and viability assessed using ViCell. Cells were resuspended at 1.2 mio cells/ml in cell culture medium without hygromycin B and added to CHO cells at 60 000 cells/well (50 µl/well) to obtain a final effector-to-target (E:T) ratio of 2:1 and a final volume of 200 µl per well. Then, 4 µl of GloSensor (Promega #E1291) was added
10 to each well (2% of final volume). Cells were incubated for 24 h at 37°C in a humidified incubator. At the end of incubation time, luminescence was detected using TECAN Spark 10M.

As shown in Figure 18, the optimized CD3 binder P035-093 (P035) (= CD3_{opt}) had a similar activity on Jurkat NFAT cells upon crosslinking as CD3_{orig}.

15 **Example 9**

Generation of bispecific antibodies that bind to human HLA-G and to human CD3 (Anti-HLA-G/anti-CD3 antibodies)

Recombinant DNA techniques

Standard methods were used to manipulate DNA as described in Sambrook, J. et al.,
20 Molecular cloning: A laboratory manual; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989. The molecular biological reagents were used according to the manufacturer's instructions.

Gene and oligonucleotide synthesis

Desired gene segments were prepared by chemical synthesis at Geneart GmbH
25 (Regensburg, Germany). The synthesized gene fragments were cloned into an E. coli plasmid for propagation/amplification. The DNA sequences of subcloned gene fragments were verified by DNA sequencing. Alternatively, short synthetic DNA fragments were assembled by annealing chemically synthesized oligonucleotides or via PCR. The respective oligonucleotides were prepared by metabion GmbH
30 (Planegg-Martinsried, Germany)

Description of the basic/standard mammalian expression plasmid

For the expression of a desired gene/protein (e.g. antibody heavy chain or antibody light chain) a transcription unit comprising the following functional elements is used:

- 5 - the immediate early enhancer and promoter from the human cytomegalovirus (P-CMV) including intron A,
- a human heavy chain immunoglobulin 5'-untranslated region (5'UTR),
- a murine immunoglobulin heavy chain signal sequence,
- a gene/protein to be expressed (e.g. full length antibody heavy chain or MHC class I molecule), and
- 10 - the bovine growth hormone polyadenylation sequence (BGH pA).

Beside the expression unit/cassette including the desired gene to be expressed the basic/standard mammalian expression plasmid contains

- an origin of replication from the vector pUC18 which allows replication of this plasmid in E. coli, and
- 15 - a beta-lactamase gene which confers ampicillin resistance in E. coli.

Protein determination

The protein concentration of purified polypeptides was determined by determining the optical density (OD) at 280 nm, using the molar extinction coefficient calculated on the basis of the amino acid sequence of the polypeptide.

20 **Generation of expression plasmids for recombinant monoclonal bispecific antibodies**

The recombinant monoclonal antibody genes encode the respective immunoglobulin heavy and light chains.

- 25 The expression plasmids for the transient expression monoclonal antibody molecules comprised besides the immunoglobulin heavy or light chain expression cassette an origin of replication from the vector pUC18, which allows replication of this plasmid in E. coli, and a beta-lactamase gene which confers ampicillin resistance in E. coli.

- 144 -

The transcription unit of a respective antibody heavy or light chain comprised the following functional elements:

- the immediate early enhancer and promoter from the human cytomegalovirus (P-CMV) including intron A,
- 5 - a human heavy chain immunoglobulin 5'-untranslated region (5'UTR),
- a murine immunoglobulin heavy chain signal sequence, and
- the bovine growth hormone polyadenylation sequence (BGH pA).

Transient expression and analytical characterization

10 The recombinant production was performed by transient transfection of HEK293 cells (human embryonic kidney cell line 293-derived) cultivated in F17 Medium (Invitrogen Corp.). For the production of monoclonal antibodies, cells were co-transfected with plasmids containing the respective immunoglobulin heavy- and light chain. For transfection "293-Fectin" Transfection Reagent (Invitrogen) was used. Transfection was performed as specified in the manufacturer's instructions.

15 Cell culture supernatants were harvested three to seven (3-7) days after transfection. Supernatants were stored at reduced temperature (e.g. -80°C).

General information regarding the recombinant expression of human immunoglobulins in e.g. HEK293 cells is given in: Meissner, P. et al., Biotechnol. Bioeng. 75 (2001) 197-203.

20 Using the above described methods for recombinant DNA techniques, the generation of expression plasmids for recombinant monoclonal antibodies and transient expression and analytical characterization, the following bispecific antibodies that bind to human HLA-G and to human CD3 were produced and analyzed:

25 **Bispecific antibodies that bind to human HLA-G and to human CD3 (Anti-HLA-G/anti-CD3 antibodies) (SEQ ID Nos of variable regions VH/VL and hypervariable regions (HVRs) of antigen binding moieties/sites binding human HLA-G and of antigen binding moieties/sites binding human CD3):**

Anti-HLA-G antigen binding site	HVR-H1	HVR-H2	HVR-H3	HVR-L1	HVR-L2	HVR-L3	VH	VL
HLA-G-0090-S32P	SEQ ID NO: 1	SEQ ID NO: 2	SEQ ID NO: 3	SEQ ID NO: 23	SEQ ID NO: 5	SEQ ID NO: 6	SEQ ID NO: 7	SEQ ID NO: 24
Anti-CD3 antigen binding site	HVR-H1	HVR-H2	HVR-H3	HVR-L1	HVR-L2	HVR-L3	VH	VL
P035-093 (P035)	SEQ ID NO: 52	SEQ ID NO: 53	SEQ ID NO: 54	SEQ ID NO: 55	SEQ ID NO: 56	SEQ ID NO: 57	SEQ ID NO: 58	SEQ ID NO: 59
Clone 22 (Cl22)	SEQ ID NO: 60	SEQ ID NO: 61	SEQ ID NO: 62	SEQ ID NO: 63	SEQ ID NO: 64	SEQ ID NO: 65	SEQ ID NO: 66	SEQ ID NO: 67
V9	SEQ ID NO: 68	SEQ ID NO: 69	SEQ ID NO: 70	SEQ ID NO: 71	SEQ ID NO: 72	SEQ ID NO: 73	SEQ ID NO: 74	SEQ ID NO: 75

“Clone 22 (abbreviated as “Cl22”)” is an optimized CD3 binder (see WO 2020/127619); “P035-093 (abbreviated as “P035”)” is another optimized variant CD3 binder; V9 is another CD3 binder described e.g. in Rodrigues et al., Int J Cancer Suppl (1992) 7, 45-50, and WO 1992/22653 (SEQ ID NOs 20 and 17 of WO 1992/22653 are the VH and VL sequences)

Bispecific anti-HLA-G/anti-CD3 T cell bispecific (TCB) antibodies:

P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035-093 (P035)):

- 146 -

SEQ ID NO: 76 light chain 1 P1AF7977

SEQ ID NO: 77 light chain 2 P1AF7977

SEQ ID NO: 78 heavy chain 1 P1AF7977

SEQ ID NO: 79 heavy chain 2 P1AF7977

5 P1AF7978 (HLA-G-0090-VL-S32P/ CD3 Clone 22 (CI22)):

SEQ ID NO: 80 light chain 1 P1AF7978

SEQ ID NO: 81 light chain 2 P1AF7978

SEQ ID NO: 82 heavy chain 1 P1AF7978

SEQ ID NO: 83 heavy chain 2 P1AF7978

10 P1AF7979 (HLA-G-0090-VL-S32P/ CD3 V9):

SEQ ID NO: 84 light chain 1 P1AF7979

SEQ ID NO: 85 light chain 2 P1AF7979

SEQ ID NO: 86 heavy chain 1 P1AF7979

SEQ ID NO: 87 heavy chain 2 P1AF7979

15

Example 10**Binding and Stability of bispecific anti-HLA-G/anti-CD3 antibody (T cell bispecific (TCB) antibody) to HLA-G****Stability under stress**

20 The three TCB molecules featuring the same HLA-G targeting binder HLAG-0090-VL-S32P and three different CD3e binders were stressed for 14 days under two different conditions:

- pH 6.0 20 mM His/HisCl, 140 mM NaCl; at 40 °C (His 40 °C)
- pH 7.4 PBS; at 37 °C (PBS 37 °C).

25 Afterwards, the material was analysed using CD-SDS, SEC, and Surface plasmon resonance) SPR to investigate chemical degradation and possible effects on target binding. For reference, the stressed material was compared with material kept under storage conditions:

- 147 -

- pH 6.0 20 mM His/HisCl, 140 mM NaCl; frozen at -80°C (Ref.)

The results are listed in the following tables:

Parameter		P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035)	P1AF7978 (HLA-G-0090-VL-S32P/ CD3 CI22)	P1AF7979 (HLA-G-0090-VL-S32P/ CD3 V9)
CE-SDS [%] (Caliper, non-reducing)	Ref.	95	95	96
	His 40 °C	93	96	95
	PBS 37 °C	93	94	95
CE-SDS [%] (Caliper, reducing)	Ref.	100	100	100
	His 40 °C	100	100	100
	PBS 37 °C	100	100	100
SEC monomer [%]	Ref.	99	99	99
	His 40 °C	97	98	98
	PBS 37 °C	96	96	97
Thermal stability (DLS T _{agg})		64	64	68

		P1AF7977 (HLA-G- 0090-VL- S32P/ CD3 P035)		P1AF7978 (HLA-G- 0090-VL- S32P/ CD3 CI22)		P1AF7979 (HLA-G- 0090-VL- S32P/ CD3 V9)	
		HLA-G	CD3	HLA-G	CD3	HLA-G	CD3
HLA-G or CD3 specific binding (respectively) $\pm$ stress, by Biacore RAC	Ref.	100	100	100	100	100	100
	His 40 °C	100	99	99	97	99	96
	PBS 37 °C	98	96	98	92	98	91

All three TCBs are showing an acceptable stability profile with moderate loss of binding after stress. Furthermore, Thermal stability was measured by DLS (T_{agg}) and it is in the normal range known for human IgG. RAC Binding was determined by Surface plasmon resonance (Biacore) as described in Example 2.

Example 11

Binding of bispecific anti-HLA-G/anti-CD3 antibody (T cell bispecific (TCB) antibody) to CD3 expressed on T-cells (as assessed by Flow cytometry)

Briefly, 100ml fresh blood was collected in Erlenmeyer flasks and mixed with 100ml of isolation buffer (PBS with 2% FBS and 2mM EDTA). 25 ml of the suspension was then transferred carefully over 15ml of ficoll in a 50ml tube and centrifuged for 15 min at 800g without brakes. The PBMC layer in the ficoll gradient was then transferred to a fresh 50ml tube with isolation buffer and centrifuged at 300g for 10min at 4°C. The PBMCs were then washed twice and the cells were pooled in 10ml of isolation buffer. PBMCs were frozen at -80°C until further use. T cells were isolated from the PBMCs using EasySep negative selection human T cell Isolation kit (Stem cell, # 17951) as per manufacturer's instructions. Binding of HLA-G TCBs to T cells was then measured by flow cytometry. Briefly, 500 μ l of T cells (5x10⁵ cells) were added to each FACS tube. T cells were washed in 2 ml of staining buffer (PBS with 2% FBS) and centrifuged at 300g at 4°C for 5 minutes. The HLA-G TCBs

were diluted at different concentrations ranging from 5 – 0.05µg/ml in medium. T cells were then resuspended in 100µl of HLA-G TCB dilution and incubated for 30 min in the dark at 4°C. After washing once with 2ml staining buffer, cells were centrifuged at 300g for 5 min and then resuspended in 100µl of secondary antibody dilution (Alexa Fluor 488 labeled anti-human IgG, 1:200) for 30 min at 4°C in the dark. T cells were washed twice with 2ml staining buffer and centrifuged at 300g for 5 min at 4°C. Finally cells were resuspended in 500µl medium and binding of HLA-G TCBs to T Cells was detected on BD LSR. The binding of HLA-G TCBs P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035); P1AF7978 (HLA-G-0090-VL-S32P/ CD3 C122) and P1AF7979 (HLA-G-0090-VL-S32P/ CD3 V9) to CD3 on T cells at different concentrations is illustrated in Figure 7.

Example 12

Binding of bispecific anti-HLA-G/anti-CD3 antibody (T cell bispecific (TCB) antibody) to natural or recombinant HLA-G expressed on cells (as assessed by Flow cytometry)

Binding ability of anti HLA-G TCB mAb to HLA-G expressed on different cells and cell lines was assessed by FACS analysis. Either the binding to naturally HLA-G expressing JEG3 tumor cells or Skov3 transfectants and respective parental, untransfected cells is described.

For flow cytometry analysis, cells were stained with anti HLA-G TCB mAb at 4°C. Briefly, 25µl/well of each cell suspension (5x10⁴ cells/well) was transferred into a polypropylene 96-Well V-bottom plate and prechilled in the fridge at 5°C for 10 min. Anti-HLA-G samples were diluted in staining buffer to a 2-fold starting concentration of 80µg/ml. A 4-fold serial dilution of the antibodies was performed and 25µl/well of the antibody solution was added to the prepared cells and incubated for 1h at 5°C. Cells were washed twice with 200µl/well staining buffer and centrifugation at 300g for 5min. Cell pellets were resuspended in 25µl of staining buffer afterwards. For detection fluorescent labeled anti-species antibody (donkey anti human IgG (H+L) conjugated to PE, Jackson Immuno Research # 709-116-149) was diluted 1:100 in staining buffer and 25µl/well detection antibody was added to the cell suspension. After a 1 hour incubation at 5°C cells were again washed twice with staining buffer, resuspended in 70µl of staining buffer and measured at a FACS Canto II. Bispecific anti-HLA-G/anti-CD3 antibodies (T cell bispecific (TCB) antibodies) P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035); P1AF7978 (HLA-G-0090-VL-S32P/ CD3 C122) and P1AF7979 (HLA-G-0090-VL-S32P/ CD3 V9)

- 150 -

showed binding to JEG3 cells and SKOV3 cells, transfected with HLAG (see Fig. 8). The EC₅₀ values for FACS binding are listed in the table below.

Cell binding EC ₅₀ (nM)	JEG3	SKOV3 HLA-G
P1AF7977	0.42	1.5
P1AF7978	0.12	0.15
P1AF7979	0.36	0.58

Example 13

5 ILT2 and -4 binding inhibition of bispecific anti-HLA-G/anti-CD3 antibody (T cell bispecific (TCB) antibody)

The ELISA was set up by coating the Fc tagged ILT2 and ILT4 respectively to Maxisorp microtiter plates. After incubation and washing steps, the respective antibodies are added at a concentration of 100nM. Soluble His tagged monomeric,
 10 dimeric or trimeric HLA-G was added to the wells. After incubation and washing steps, detection of bound receptor was carried out by anti-His-antibody-POD conjugates. Percentage inhibition (%) is calculated in comparison to values obtained from wells with ILT2/4 + HLA-G (mono-, di-, or Trimer) without anti HLA-G or ILT2/4 antibodies (100% binding = 0% inhibition) and shown in the following table.

15

% Binding inhibition (133nM)	ILT2	ILT4
P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035)	100	92
P1AF7978 (HLA-G-0090-VL-S32P/ CD3 C122)	100	91
P1AF7979 (HLA-G-0090-VL-S32P/ CD3 V9)	100	95

Example 14**Bispecific anti-HLA-G/anti-CD3 antibody (T cell bispecific (TCB) antibody) mediated IFN gamma secretion by T cells**

Ability of anti HLA-G TCB to induce IFN gamma secretion by T cells in the presence of HLA-G expressing tumor cells was tested using SKOV3 cells transfected with recombinant HLA-G (SKOV3 HLA-G) and JEG3 cells expressing endogenous HLA-G. IFN gamma secretion was detected by Luminex technology. For measurement of IFN gamma secretion by T cells after TCB treatment, co-cultures of PBMCs and SKOV3HLA-G cells or JEG3 cells were incubated with anti-HLA-G TCB. Briefly, PBMCs were isolated from human peripheral blood by density gradient centrifugation using Lymphocyte Separating Medium Tubes (PAN #P04-60125). PBMC's and SKOV 3 HLA-G cells were seeded at a ratio of 10 : 1 in 96-well U bottom plates. The co-culture was then incubated with HLA-G-TCB at different concentrations as shown in the figure (Fig. 9) and incubated for 24h at 37°C in an incubator with 5% Co2. On the next day, supernatants were collected and IFN gamma secretion was measured using Milliplex MAP kit (Luminex technology) according to the manufacturer's instructions. Bispecific anti-HLA-G/anti-CD3 (T cell bispecific (TCB)) antibodies P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035); P1AF7978 (HLA-G-0090-VL-S32P/ CD3 C122) and P1AF7979 (HLA-G-0090-VL-S32P/ CD3 V9) induced IFN gamma secretion by T cells (Figure 9). The EC50 values are listed in the table below.

IFNgamma induction EC ₅₀ (nM)	JEG3	SKOV3 HLA-G
P1AF7977	20	30
P1AF7978	2.2	2.6
P1AF7979	29	16

Example 15**Induction of T cell mediated cytotoxicity/tumor cell killing by bispecific anti-HLA-G/anti-CD3 antibody (T cell bispecific (TCB) antibody)**

Ability of anti HLA-G TCB to induce T cell mediated cytotoxicity in the presence of HLA-G expressing tumor cells was tested on SKOV3 cells transfected with

recombinant HLA-G (SKOV 3 HLA-G) and JEG3 cells expressing endogenous HLA-G. Cytotoxicity was detected by measuring Caspase 8 activation in cells after treatment with HLA-G TCB. For measurement of Caspase 8 activation after HLA-G/anti-CD3 antibody (TCB) treatment, co-cultures of PBMCs and SKOV3 HLA-G cells or JEG3 cells were incubated with anti-HLA-G TCB for 24 hours and Caspase8 activation was measured using the Caspase8Glo kit (Promega, #G8200). Briefly, PBMCs were isolated from human peripheral blood by density gradient centrifugation using Lymphocyte Separating Medium Tubes (PAN #P04-60125). PBMC's and SKOV3 HLA-G or JEG3 cells were seeded at a ratio of 10: 1 (100µl per well) in black clear bottom 96-well plates. The co-culture was then incubated with HLA-G-TCB at different concentrations as shown in the figure (Fig. 10) and incubated for 24h or 48h at 37°C in an incubator with 5% Co₂. On the next day, 100µl of Caspase8 Glo substrate was added to each well and placed on a shaker for 1 hour at room temperature. The luminescence was measured on a BioTek Synergy 2 machine. The relative luminescence units (RLUs) correspond to the Caspase8 activation /cytotoxicity are plotted in the graph (Fig. 10). Bispecific anti-HLA-G/anti-CD3 (T cell bispecific (TCB)) antibodies P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035); P1AF7978 (HLA-G-0090-VL-S32P/ CD3 C122) and P1AF7979 (HLA-G-0090-VL-S32P/ CD3 V9) induced T cell mediated cytotoxicity/tumor cell killing. The EC₅₀ values of the tumor cell killing are listed in the table below.

T cell mediated cytotoxicity induction EC₅₀ (nM)	JEG3	SKOV3 HLA-G
P1AF7977	12	1.4
P1AF7978	2.6	1.36
P1AF7979	4.6	1

Example 16**In vivo anti-tumor efficacy of bispecific anti-HLA-G/anti-CD3 (T cell bispecific (TCB)) antibody in humanized NSG mice bearing SKOV3 human ovarian carcinoma transfected with recombinant HLA-G (SKOV3 HLA-G)**

5 Humanized NSG (NOD/scid/IL-2R γ null humanized with CD34⁺ cord blood cells by Jackson Laboratories, US) mice (n=15) were injected subcutaneously with 5 $\times$ 10⁶ SKOV3 HLA-G cells in a total volume of 100 μ l. Once the tumors reached an average volume of 200 mm³, mice were randomized and treated weekly with bispecific anti-HLA-G/anti-CD3 (T cell bispecific (TCB)) antibody (P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035)) (5 mg/kg) weekly. As a control, one group of mice received weekly i.v. injections of histidine buffer (vehicle). Tumor volume was measured twice weekly until study termination. The results of the experiment are shown in Fig. 11. Results show tumor volume data (Median and Inter quartile range (IQR)) measured by caliper in the two study groups. The anti-HLA-G/anti-CD3 T cell bispecific (TCB) antibody P1AF7977 showed strong tumor growth inhibition/ tumor regression in the SKOV3-HLA-G tumor model.

Example 17**Dose-response study bispecific anti-HLA-G/anti-CD3 (T cell bispecific (TCB)) antibody in humanized NSG mice bearing human breast cancer PDX tumors (BC004)**

20 Humanized NSG (NOD/scid/IL-2R γ null humanized by intravenous injection of 1 $\times$ 10⁵ CD34⁺ cord blood cells per mouse) mice were injected with 2 $\times$ 10⁶ BC004 breast cancer cells in total volume of 50 μ L PBS into the intra-mammary fat pad. Once the tumors reached an average volume of approximately 200 mm³, mice were randomized (n=15 animals per group) and treated weekly with bispecific anti-HLA-G/anti-CD3 (T cell bispecific (TCB)) antibody (P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035)) with three different doses (5 mg/kg, 2.5 mg/kg, 0.5mg/kg). As a control, one group of mice received weekly i.v. injections of histidine buffer (vehicle). Tumor volume was determined twice weekly via caliper measurement.

30 The results of the experiment shown in Fig. 12 demonstrates tumor volume data (Median and Inter quartile range (IQR)). All three doses of anti-HLA-G/anti-CD3 T cell bispecific (TCB) antibody showed strong tumor growth inhibition/ tumor regression in the BC004 tumor model. The highest dose (5 mg/kg) shows slightly higher efficacy compared to the 2.5 mg/kg and 0.5 mg/kg treatment groups.

Example 18

Induction of T cell activation in the presence of HLA-G expressing tumor cells was tested on SKOV3 cells transfected with recombinant HLA-G (SKOV3 HLA-G) by bispecific anti-HLA-G/anti-CD3 antibody (T cell bispecific (TCB) antibody)

Ability of anti HLA-G/anti CD3 TCB to activate T cells in the presence of HLA-G expressing tumor cells was tested on SKOV3 cells transfected with recombinant HLA-G (SKOV3 HLA-G). Activation of T cells was assessed by FACS analysis of cell surface activation markers CD25 and early activation marker CD69 on T cells. Briefly, Peripheral Blood Mononuclear Cells (PBMCs) are isolated from human peripheral blood by density gradient centrifugation using Lymphocyte Separating Medium Tubes (PAN #P04-60125). PBMC's and SKOV3 HLA-G cells are seeded at a ratio of 10 : 1 in 96-well U bottom plates. The co-culture was then incubated with anti-HLA-G/anti-CD3 (T cell bispecific (TCB)) antibody (P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035)) (0.01nM) and incubated for 24h at 37°C in an incubator with 5% Co2. On the next day, expression of CD25 and CD69 was measured by flow cytometry.

For flow cytometry analysis, cells are stained with with PerCP-Cy5.5 Mouse Anti-Human CD8 (BD Pharmingen # 565310), PE-Cy7 Mouse Anti-Human CD4 (Biologend #317414), FITC Mouse Anti-Human CD25 (Biolegend # 356106) and APC Mouse Anti-Human CD69 (BD Pharmingen # 555533) at 4°C. Briefly, antibodies are diluted to a 2-fold concentration and 25µl of antibody dilution are added in each well with 25µl of pre-washed co-cultures. Cells are stained for 30 min at 4°C and washed twice with 200µl/well staining buffer and centrifugation at 300g for 5min. Cell pellets are resuspended in 200µl of staining buffer and stained with DAPI for live dead discrimination at a final concentration of 2µg/ml. Samples are then measured using BD LSR flow cytometer. Data analysis was performed using FlowJo V.10.1 software. Figure 14 shows the induction of T cell activation by bispecific anti-HLA-G/anti-CD3 antibody P1AF7977 (HLA-G-0090-VL-S32P/ CD3 P035 in the presence of SKOV3 HLAG cells.

Patent Claims

1. An antibody that binds to human HLA-G comprising
 - 5 A) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:23; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6, or
 - 10 B) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:25; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6.
2. The antibody according to claim 1, wherein the antibody
 - 20 A) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24; or
 - B) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26.
3. The antibody of claims 1 or 2, wherein the antibody comprises a Fc domain of human origin, particularly of the IgG isotype, more particularly of the IgG1 isotype.
- 25 4. The antibody according to any one of claims 1 to 3, wherein the antibody comprises a constant region of human origin, particularly of the IgG isotype, more particularly of the IgG1 isotype, comprising a human CH1, CH2, CH3 and/or CL domain..
5. The antibody according to any one of claim 1 to 4, wherein the antibody

- 156 -

- a) does not crossreact with a modified human HLA-G β 2M MHC I complex, wherein the HLA-G specific amino acids have been replaced by HLA-A consensus amino acids, the complex comprising SEQ ID NO:40; and/ or
 - b) does not crossreact with a mouse H2Kd β 2M MHC I complex comprising SEQ ID NO:41; and/ or
 - c) does not crossreact with rat RT1A β 2M MHC I complex comprising SEQ ID NO:43.
6. The antibody according to any one of claims 1 to 5, wherein the antibody
- a) inhibits ILT2 binding to (HLA-G expressed on) JEG3 cells (ATCC No. HTB36); or
 - b) binds to (HLA-G expressed on) JEG3 cells (ATCC No. HTB36), and inhibits ILT2 binding to (HLA-G expressed on) JEG-3 cells (ATCC No. HTB36).
7. The antibody according to any one of claims 1 to 4, wherein the antibody is a multispecific antibody .
8. The antibody according to any one of claims 1 to 4, wherein the antibody is a bispecific antibody that binds to human HLA-G and to human CD3.
9. The antibody according to claim 5, wherein the antibody is a bispecific antibody that binds to human HLA-G and to human CD3, comprising a first antigen binding moiety that binds to human HLA-G and a second antigen binding moiety that binds to human CD3, wherein the first antigen binding moiety that binds to human HLA-G comprises
- A) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:23; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6, or
 - B) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:1, (ii) CDR-H2 comprising the amino acid sequence of SEQ

- 157 -

5 ID NO:2, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:3; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:25; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:5 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:6;

and wherein the second antigen binding moiety that binds to a T cell activating antigen binds to human CD3 comprises

10 C) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:52, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:53, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:54; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:55; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:56 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:57, or

15 D) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:60, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:61, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:62; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:63; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:64 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:65, or

20 E) (a) a VH domain comprising (i) CDR-H1 comprising the amino acid sequence of SEQ ID NO:68, (ii) CDR-H2 comprising the amino acid sequence of SEQ ID NO:69, and (iii) CDR-H3 comprising the amino acid sequence of SEQ ID NO:70; and (b) a VL domain comprising (i) CDR-L1 comprising the amino acid sequence of SEQ ID NO:71; (ii) CDR-L2 comprising the amino acid sequence of SEQ ID NO:72 and (iii) CDR-L3 comprising the amino acid sequence of SEQ ID NO:73.

25 10. The bispecific antibody according to claim 9,
30 wherein the first antigen binding moiety

A) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24; or

- 158 -

- B) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:26, and wherein the second antigen binding moiety
- 5 C) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:58 and a VL domain comprising the amino acid sequence of SEQ ID NO:59; or
- D) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:66 and a VL domain comprising the amino acid sequence of SEQ ID NO:67; or
- 10 E) comprises a VH domain comprising the amino acid sequence of SEQ ID NO:74 and a VL domain comprising the amino acid sequence of SEQ ID NO:75.
11. The bispecific antibody according to claim 10, wherein the first antigen binding moiety
- 15 comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24; and wherein the second antigen binding moiety
- comprises a VH domain comprising the amino acid sequence of SEQ ID NO:58 and a VL domain comprising the amino acid sequence of SEQ ID
- 20 NO:59.
12. The bispecific antibody according to claim 10, wherein the first antigen binding moiety
- comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7 and a VL domain comprising the amino acid sequence of SEQ ID NO:24;
- 25 and wherein the second antigen binding moiety
- comprises a VH domain comprising the amino acid sequence of SEQ ID NO:66 and a VL domain comprising the amino acid sequence of SEQ ID NO:67.

- 159 -

13. The bispecific antibody according to claim 10,
wherein the first antigen binding moiety
comprises a VH domain comprising the amino acid sequence of SEQ ID NO:7
and a VL domain comprising the amino acid sequence of SEQ ID NO:24;
5 and wherein the second antigen binding moiety
comprises a VH domain comprising the amino acid sequence of SEQ ID
NO:74 and a VL domain comprising the amino acid sequence of SEQ ID
NO:75.
- 10 14. The bispecific antibody according to any one of claims 7 to 13, wherein the
bispecific antibody shows
- a) inhibition of ILT2 and/or ILT4 binding to HLA-G; and/or
- b) antibody mediated IFN gamma secretion by T cells on SKOV3 cells
transfected with recombinant HLA-G (SKOV3 HLA-G) and/or on JEG3 cells
expressing endogenous HLA-G wherein the IFN gamma secretion was
15 detected; and or
- c) T cell mediated cytotoxicity/tumor cell killing on SKOV3 cells transfected
with recombinant HLA-G (SKOV 3HLA-G) and/or JEG3 cells expressing
endogenous HLA-G wherein the cytotoxicity was detected by measuring
Caspase 8 activation in cells after treatment with bispecific antibody; and/or
- 20 d) in vivo anti-tumor efficacy/ tumor regression in humanized NSG mice bearing
SKOV3 human ovarian carcinoma transfected with recombinant HLA-G
(SKOV3 HLA-G) humanized NSG mice; and/or
- e) in vivo anti-tumor efficacy /tumor of HLA-G CD3 T cell bi-specific in
humanized NSG mice bearing human breast cancer PDX tumors (BC004).
- 25 15. Isolated nucleic acid encoding the antibody according to any one of claims 1-
4 or the bispecific antibody according to any one of claims 7-13.
16. A host cell, preferably an eukaryotic host cell, comprising the nucleic acid of
embodiment 15.

- 160 -

17. A method of producing the antibody according to any one of claims 1-4 or the bispecific antibody according to any one of claims 7-13 comprising culturing the host cell of claim 16 so that the antibody or bispecific antibody is produced.
- 5 18. The method of claim 30, further comprising recovering the antibody or bispecific antibody from the host cell.
19. The antibody according to any one of claims 1-4 or the bispecific antibody according to any one of claims 7-13, wherein the antibody is produced in an eukaryotic host cell.
- 10 20. A pharmaceutical formulation comprising the antibody according to any one of claims 1-4 or the bispecific antibody according to any one of claims 7-13 and a pharmaceutically acceptable carrier.
21. The antibody according to any one of claims 1-4 or the bispecific antibody according to any one of claims 7-13 for use as a medicament.
- 15 22. The antibody according to any one of claims 1-4 or the bispecific antibody according to any one of claims 7-13 for use in treating cancer.
23. Use of the antibody according to any one of claims 1-4 or the bispecific antibody according to any one of claims 7-13 in the manufacture of a medicament.
24. The use of claim 23, wherein the medicament is for treatment of cancer.
- 20 25. A method of treating an individual having cancer comprising administering to the individual an effective amount of the antibody according to any one of claims 1-4 or the bispecific antibody according to any one of claims 7-13.