

ADAMANTANE DERIVATIVES

The present invention relates to adamantane derivatives, a process for their preparation, pharmaceutical compositions containing them, a process for preparing the pharmaceutical compositions, and their use in therapy.

Adamantane derivatives are known in the art, e.g. from WO 95/04720 for use as gastrin and cholecystokinin receptor ligands, from Chem. Abs. (1977), Volume 86, No. 13 (86: 89560d) for use as analgesics, and from US-A-3 464 998 as antibiotics.

10


The P2X₇ receptor (previously known as P2Z receptor), which is a ligand-gated ion channel, is present on a variety of cell types, largely those known to be involved in the inflammatory/immune process, specifically, macrophages, mast cells and lymphocytes (T and B). Activation of the P2X₇ receptor by extracellular nucleotides, in particular adenosine triphosphate, leads to the release of interleukin-1 β (IL-1 β) and giant cell formation (macrophages/microglial cells), degranulation (mast cells) and proliferation (T cells), apoptosis and L-selectin shedding (lymphocytes). P2X₇ receptors are also located on antigen-presenting cells (APC), keratinocytes, salivary acinar cells (parotid cells), hepatocytes and mesangial cells.

20

It would be desirable to make compounds effective as P2X₇ receptor antagonists for use in the treatment of inflammatory, immune or cardiovascular diseases, in the aetiologies of which the P2X₇ receptor may play a role.

25

In accordance with the present invention, there is therefore provided a compound of general formula




wherein m represents 1, 2 or 3, preferably 1 or 2;

each R¹ independently represents a hydrogen or halogen (e.g. fluorine, chlorine, bromine or iodine) atom, preferably a hydrogen atom;

5 A represents C(O)NH or, preferably, NHC(O);

Ar represents a group



X represents a bond, an oxygen atom or a group CO, (CH₂)₁₋₆, CH=, (CH₂)₁₋₆O, O(CH₂)₁₋₆, O(CH₂)₂₋₆O, O(CH₂)₂₋₃O(CH₂)₁₋₃, CR⁵(OH), (CH₂)₁₋₃O(CH₂)₁₋₃,
 10 (CH₂)₁₋₃O(CH₂)₂₋₃O, NR⁵, (CH₂)₁₋₆NR⁵, NR⁵(CH₂)₁₋₆, (CH₂)₁₋₃NR⁵(CH₂)₁₋₃,
 O(CH₂)₂₋₆NR⁵, O(CH₂)₂₋₃NR⁵(CH₂)₁₋₃, (CH₂)₁₋₃NR⁵(CH₂)₂₋₃O, NR⁵(CH₂)₂₋₆O,
 NR⁵(CH₂)₂₋₃O(CH₂)₁₋₃, CONR⁵, NR⁵CO, S(O)_n, S(O)_nCH₂, CH₂S(O)_n, SO₂NR⁵
 or NR⁵SO₂;

n is 0, 1 or 2;

15 R' represents a hydrogen atom or a C₁-C₆ alkyl, preferably methyl, group;

one of R² and R³ represents a halogen, cyano, nitro, amino, hydroxyl, or a group

selected from (i) C₁-C₆ alkyl optionally substituted by at least one C₃-C₆ cycloalkyl,

(ii) C₃-C₈ cycloalkyl, (iii) C₁-C₆ alkyloxy optionally substituted by at least one

C₃-C₆ cycloalkyl, and (iv) C₃-C₈ cycloalkyloxy, each of these groups being optionally

20 substituted by one or more fluorine atoms, and the other of R² and R³ represents a hydrogen or halogen atom;

- either R^4 represents a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom, the heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl, carboxyl, cyano, C_1 - C_6 alkyl, C_1 - C_6 hydroxyalkyl, $-NR^6R^7$, $-(CH_2)_rNR^6R^7$ and $-CONR^6R^7$,
or R^4 represents a 3- to 8-membered saturated carbocyclic ring system substituted by one or more substituents independently selected from $-NR^6R^7$, $-(CH_2)_rNR^6R^7$ and $-CONR^6R^7$, the ring system being optionally further substituted by one or more substituents independently selected from fluorine atoms, hydroxyl and C_1 - C_6 alkyl;
- r is 1, 2, 3, 4, 5 or 6;
- R^5 represents a hydrogen atom or a C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl group;
- R^6 and R^7 each independently represent a hydrogen atom or a C_1 - C_6 alkyl, C_2 - C_6 hydroxyalkyl or C_3 - C_8 cycloalkyl group, or R^6 and R^7 together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring;
- with the provisos that,
- (a) when A represents $C(O)NH$ and R^4 represents an unsubstituted 3- to 8-membered saturated aliphatic heterocyclic ring system containing one nitrogen atom, then X is other than a bond, and
- (b) when A represents $C(O)NH$ and X represents a group $(CH_2)_{1-6}$ or $O(CH_2)_{1-6}$, then R^4 does not represent an unsubstituted imidazolyl, unsubstituted morpholinyl, unsubstituted piperidinyl or unsubstituted pyrrolidinyl group, and
- (c) when A represents $NHC(O)$ and R^4 represents an unsubstituted 3- to 8-membered saturated aliphatic heterocyclic ring system containing one nitrogen atom, then X is other than a bond, and
- (d) when A represents $NHC(O)$ and X represents $O(CH_2)_{1-6}$, $NH(CH_2)_{1-6}$ or SCH_2 , then R^4 does not represent an unsubstituted 1-piperidinyl or unsubstituted 1-pyrrolidinyl group, and
- (e) when A represents $NHC(O)$ and X represents $O(CH_2)_{2-3}NH(CH_2)_2$, then R^4 does not represent an imidazolyl group;
- or a pharmaceutically acceptable salt or solvate thereof.

In the context of the present specification, unless otherwise indicated, an alkyl substituent or alkyl moiety in a substituent group may be linear or branched. Examples of alkyl groups/moieties containing up to 6 carbon atoms include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl and hexyl. When one of R^2 and R^3 represents a C_1 - C_6 alkyl/ C_1 - C_6 alkyloxy optionally substituted by at least one C_3 - C_6 cycloalkyl, it should be understood that one or both of the alkyl and cycloalkyl moieties may be optionally substituted by fluorine atoms. In relation to R^4 , a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom may be a monocyclic or bicyclic ring system. Further in relation to R^4 , a 3- to 8-membered saturated carbocyclic ring system may be a monocyclic or bicyclic ring system. When R^6 or R^7 represents a C_2 - C_6 hydroxyalkyl in the substituent NR^6R^7 , $-(CH_2)_rNR^6R^7$ or $-CONR^6R^7$, it will be appreciated that the hydroxyl group will not be bonded to the same carbon atom as the nitrogen atom. When R^6 and R^7 together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring, the ring obtained is monocyclic.

Preferably X represents a bond, an oxygen atom or a group CO, $(CH_2)_{1-6}$, $CH=$, $O(CH_2)_{1-6}$, $O(CH_2)_{2-6}O$, $O(CH_2)_{2-3}O(CH_2)_{1-3}$, $CR^1(OH)$, NR^5 , $(CH_2)_{1-6}NR^5$, $CONR^5$, $S(O)_n$ or $S(O)_nCH_2$.


One of R^2 and R^3 represents a halogen (e.g. fluorine, chlorine, bromine or iodine), cyano, nitro, amino, hydroxyl, or a group selected from (i) C_1 - C_6 alkyl, preferably C_1 - C_4 alkyl, optionally substituted by at least one (e.g. 1, 2 or 3) C_3 - C_6 cycloalkyl (i.e. cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl), (ii) C_3 - C_8 cycloalkyl (e.g. cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl), (iii) C_1 - C_6 alkyloxy, preferably C_1 - C_4 alkyloxy, optionally substituted by at least one (e.g. 1, 2 or 3) C_3 - C_6 cycloalkyl (i.e. cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl), and (iv) C_3 - C_8 cycloalkyloxy (e.g. cyclopropyloxy, cyclobutyloxy, cyclopentyloxy or cyclohexyloxy), each of these groups being optionally substituted by one or more (e.g. 1, 2, 3 or 4) fluorine atoms, and the other

of R^2 and R^3 represents a hydrogen or halogen (e.g. fluorine, chlorine, bromine or iodine) atom.


Preferably, one of R^2 and R^3 represents a halogen (especially chlorine or bromine) atom or a nitro, amino or C_1 - C_6 alkyl (especially methyl or ethyl) group and the other of R^2 and R^3 represents a hydrogen atom.

R^4 may represent a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom, the heterocyclic ring system being optionally substituted by one or more (e.g. 1, 2, 3 or 4) substituents independently selected from fluorine atoms, hydroxyl, carboxyl, cyano, C_1 - C_6 alkyl, preferably C_1 - C_4 alkyl, C_1 - C_6 hydroxyalkyl, preferably C_1 - C_4 hydroxyalkyl, $-NR^6R^7$, $-(CH_2)_rNR^6R^7$ and $-CONR^6R^7$.

The 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system in the group R^4 may be a monocyclic ring system such as pyrrolidinyl (e.g. 1-pyrrolidinyl, 2-pyrrolidinyl or 3-pyrrolidinyl), piperidinyl (e.g. 1-piperidinyl, 2-piperidinyl, 3-piperidinyl or 4-piperidinyl), 4-piperiden-3-yl, piperazinyl (e.g. 1-piperazinyl), homopiperazinyl,




or a bicyclic ring system such as





Alternatively, R^4 may represent a 3- to 8-membered saturated carbocyclic ring system substituted by one or more (e.g. 1, 2 or 3) substituents independently selected from NR^6R^7 , $-(CH_2)_rNR^6R^7$ and $-CONR^6R^7$, the ring system being optionally further substituted by one or more (e.g. 1, 2, 3 or 4) substituents independently selected from
 5 fluorine atoms, hydroxyl and C_1 - C_6 alkyl, preferably C_1 - C_4 alkyl.

The 3- to 8-membered saturated carbocyclic ring in the group R^4 is preferably a monocyclic ring system such as a cyclopentyl or cyclohexyl ring.


10 Specific examples of groups R^4 include:









When X represents a bond or a group CO , $(\text{CH}_2)_{1-6}$, $\text{O}(\text{CH}_2)_{2-6}$, $\text{O}(\text{CH}_2)_{2-3}\text{O}(\text{CH}_2)_{2-3}$, $(\text{CH}_2)_{1-3}\text{O}(\text{CH}_2)_{2-3}$, $\text{NR}^5(\text{CH}_2)_{2-6}$, $(\text{CH}_2)_{1-3}\text{NR}^5(\text{CH}_2)_{2-3}$, $\text{O}(\text{CH}_2)_{2-3}\text{NR}^5(\text{CH}_2)_{2-3}$, $\text{NR}^5(\text{CH}_2)_{2-3}\text{O}(\text{CH}_2)_{2-3}$, NR^5CO , SO_2 or NR^5SO_2 , R^4 preferably represents a group:





When X represents an oxygen or sulphur atom or a group CH= , $(\text{CH}_2)_{1-6}\text{O}$, OCH_2 , $\text{O}(\text{CH}_2)_{2-6}\text{O}$, $\text{O}(\text{CH}_2)_{2-3}\text{OCH}_2$, CR^4OH , $(\text{CH}_2)_{1-3}\text{OCH}_2$, $(\text{CH}_2)_{1-3}\text{O}(\text{CH}_2)_{2-3}\text{O}$, NR^5 , $(\text{CH}_2)_{1-6}\text{NR}^5$, $\text{O}(\text{CH}_2)_{2-6}\text{NR}^5$, NR^5CH_2 , $(\text{CH}_2)_{1-3}\text{NR}^5\text{CH}_2$, $\text{O}(\text{CH}_2)_{2-3}\text{NR}^5\text{CH}_2$, $(\text{CH}_2)_{1-3}\text{NR}^5(\text{CH}_2)_{2-3}\text{O}$, $\text{NR}^5(\text{CH}_2)_{2-6}\text{O}$, $\text{NR}^5(\text{CH}_2)_{2-3}\text{OCH}_2$, CONR^5 , SO , $\text{S}(\text{O})_n\text{CH}_2$, $\text{CH}_2\text{S}(\text{O})_n$ or SO_2NR^5 , R^4 preferably represents a group:



R^5 represents a hydrogen atom, or a C_1 - C_6 , preferably C_1 - C_4 , alkyl (e.g. methyl, ethyl, propyl, butyl, pentyl or hexyl) or C_3 - C_8 , preferably C_3 - C_6 , cycloalkyl (e.g. cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl) group.

R^6 and R^7 each independently represent a hydrogen atom, or a C_1 - C_6 , preferably C_1 - C_4 , alkyl (e.g. methyl, ethyl, propyl, butyl, pentyl or hexyl), C_2 - C_6 hydroxyalkyl or C_3 - C_8 , preferably C_3 - C_6 , cycloalkyl (e.g. cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl) group, or R^6 and R^7 together with the nitrogen atom to which they are attached form a 3- to 8-membered, preferably 3- to 6-membered, saturated heterocyclic ring such as a pyrrolidinyl or piperidinyl ring.

Preferred compounds of the invention include:

2-Nitro-3-piperazin-1-yl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Amino-3-piperazin-1-yl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
dihydrochloride salt,

2-Chloro-3-piperazin-1-yl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-piperazin-1-yl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5 2-Chloro-5-(hexahydro-1H-1,4-diazepin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

5-(4-Amino-1-piperidinyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

(+/-)-5-(3-Amino-1-pyrrolidinyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
10 benzamide, hydrochloride salt,

2-Chloro-5-piperazin-1-ylmethyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
hydrochloride salt,

2-Chloro-5-[(hexahydro-1H-1,4-diazepin-1-yl)methyl] -N-(tricyclo[3.3.1.1^{3,7}]dec-1-
ylmethyl)-benzamide, hydrochloride salt,

15 5-[(4-Amino-1-piperidinyl)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

5-[(3-Amino-1-pyrrolidinyl)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

2-Chloro-5-(4-piperidinyloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
20 hydrochloride salt,

(R)-2-Chloro-5-(2-pyrrolidinylmethoxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

(S)-2-Chloro-5-(2-pyrrolidinylmethoxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

25 2-Chloro-5-(3-piperidinylmethoxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

cis-5-[(4-Aminocyclohexyl)oxy]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

2-Methyl-5-(1-piperazinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
30 hydrochloride salt,

2-Chloro-5-(1-piperazinylmethyl)-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-benzamide,
hydrochloride salt,

(+/-)-2-Chloro-5-(3-pyrrolidinylloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

5 (+/-)-2-Chloro-5-(3-piperidinylloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

trans-5-[(4-Aminocyclohexyl)oxy]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide,

10 cis-(+/-)-5-[(3-Aminocyclopentyl)oxy]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
ylmethyl)-benzamide,

(S,S)-2-Chloro-5-(2,5-diazabicyclo[2.2.1]hept-2-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(2-methyl-1-piperazinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

15 (+/-)-2-Chloro-5-(3-pyrrolidinylamino)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

(+/-)-5-(3-Amino-1-piperidinyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide,

20 (+/-)-2-Chloro-5-(3-piperidinylamino)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide,

2-Chloro-5-[hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
ylmethyl)-benzamide,

N-[2-methyl-5-(4-piperidinylloxy)phenyl]-tricyclo[3.3.1.1^{3,7}]decane-1-acetamide,
hydrochloride salt,

25 N-[2-chloro-5-(4-piperidinylloxy)phenyl]-tricyclo[3.3.1.1^{3,7}]decane-1-acetamide,
hydrochloride salt,

2-Chloro-5-[(4-piperidinylamino)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, dihydrochloride salt,

30 5-[[[4-(Aminomethyl)cyclohexyl]amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-
1-ylmethyl)-benzamide, dihydrochloride salt,

5-[[[(4-Aminocyclohexyl)amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt,

5-[(1-Azabicyclo[2.2.2]oct-3-ylamino)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5 N-[4-(3-Aminopyrrolidin-1-yl)-2-methylphenyl]-2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetamide, dihydrochloride salt,

N-(2-Methyl-4-piperazin-1-ylphenyl)-2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetamide, dihydrochloride salt,

10 cis-4-(3-Amino-cyclopentyloxy)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-4-(4-piperidinyloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

(+/-)-2-Chloro-4-(pyrrolidin-3-yloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

15 2-Chloro-4-(piperidin-3-yloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-4-(4-piperazin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

20 2-Chloro-4-(3-pyrrolidinylamino)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-4-(hexahydro-1H-1,4-diazepin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

(±)-5-[(3-Amino-1-piperidinyl)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

25 2-Chloro-5-(2,5-diazabicyclo[2.2.1]hept-2-ylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(9-oxa-3,7-diazabicyclo[3.3.1]non-3-ylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

30 2-Chloro-5-(3,7-diazabicyclo[3.3.1]non-3-ylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

trans-2-Chloro-5-[[8-(methylamino)-3-azabicyclo[3.2.1]oct-3-yl]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

cis-2-Chloro-5-[(hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

5 2-Chloro-5-(4-piperidinylidenemethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(4-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

10 2-Chloro-5-(4-hydroxy-piperidin-4-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(1,2,3,6-tetrahydro-pyridin-4-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Ethyl-5-piperazin-1-ylmethyl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

15 2-Chloro-5-(piperidin-4-ylsulfanyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(piperidin-4-ylsulfinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

20 2-Chloro-5-(piperidin-4-ylsulfonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(piperidin-4-ylmethylsulfanyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(piperidin-4-ylmethanesulfonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

25 2-Chloro-5-(piperazine-1-carbonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-([1,4]diazepane-1-carbonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

30 4-Chloro-N¹-(piperidin-4-yl)-N²-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-isophthalamide, hydrochloride salt,

2-Chloro-5-(hydroxy-4-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

(±)-2-Chloro-5-(hydroxy-3-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
ylmethyl)-benzamide, hydrochloride salt,

5 2-Bromo-5-piperazin-1-ylmethyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
hydrochloride salt,

2-Chloro-5-[2-(1-piperazinyl)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

10 2-Chloro-5-[2-(2,5-diazabicyclo[2.2.1]hept-2-yl)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
ylmethyl)-benzamide, hydrochloride salt,

5-[2-(4-Amino-1-piperidinyl)ethyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

2-Chloro-5-[2-(3-piperidinylamino)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, dihydrochloride salt,

15 5-[2-(3-Amino-1-piperidinyl)ethyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

2-Chloro-5-[2-(3-pyrrolidinylamino)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, dihydrochloride salt,

20 5-[2-[(3R)-3-Aminopyrrolidinyl]ethyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[2-[2-(hydroxymethyl)-1-piperazinyl]ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(hexahydro-1H-1,4-diazepin-1-yl)-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-
benzamide, hydrochloride salt,

25 (+/-)-5-(3-Amino-1-pyrrolidinyl)-2-chloro-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-
benzamide, hydrochloride salt,

2-Chloro-5-(4-piperidinylcarbonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

30 2-Chloro-5-[1-hydroxy-1-(4-piperidinyl)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

2-Chloro-5-[2-(1-piperazinyl)ethoxy]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[2-(4-piperidinyloxy)ethoxy]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

5 2-Chloro-5-[2-(4-piperidinyloxy)ethoxy]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[2-[2-(1-piperazinyl)ethoxy]ethoxy]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

10 2-Chloro-5-[(5,6-dihydro-1(4*H*)-pyrimidinyl)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[[4-[(2-hydroxyethyl)amino]-1-piperidinyl)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[[4-hydroxy-4-[(1-methylethyl)amino]methyl]-1-piperidinyl)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

15 2-Chloro-5-[(1,2,3,6-tetrahydro-3-pyridinyl)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(3-piperidinylmethyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetate salt,


20 2-bromo-5-[[4-[(2-hydroxyethyl)amino]-1-piperidinyl)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, and

2-Chloro-5-[(*E*)-3-piperidinylidenemethyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide.

25 The present invention further provides a process for the preparation of a compound of formula (I) as defined above which comprises:

(i) when X represents a CH₂ group, R⁴ represents a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom, the heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl, carboxyl,

cyano, C₁-C₆ alkyl, C₁-C₆ hydroxyalkyl, -NR⁶R⁷, -(CH₂)_rNR⁶R⁷ and -CONR⁶R⁷ and R⁴ is linked to X through a nitrogen atom, reacting a compound of general formula




wherein one of R¹⁰ and R¹¹ represents a hydrogen atom and the other of R¹⁰ and R¹¹ represents a group -CH₂L¹ in which L¹ represents a leaving group (e.g. a halogen atom) and m, A, R¹, R² and R³ are as defined in formula (I), with a compound of general formula




in the presence of a base (e.g. diisopropylethylamine), wherein R^{4'} represents a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom, the heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl, carboxyl, cyano, C₁-C₆ alkyl, C₁-C₆ hydroxyalkyl, -NR⁶R⁷, -(CH₂)_rNR⁶R⁷ and -CONR⁶R⁷ and wherein R⁶ and R⁷ are as defined in formula (I); or

(ii) when X represents an oxygen atom or a group O(CH₂)₁₋₆, O(CH₂)₂₋₆O, O(CH₂)₂₋₃O(CH₂)₁₋₃, O(CH₂)₂₋₆NR⁵ or O(CH₂)₂₋₃NR⁵(CH₂)₁₋₃, reacting a compound of general formula




wherein one of R^{12} and R^{13} represents a hydrogen atom and the other of R^{12} and R^{13} represents a hydroxyl group and m , A , R^1 , R^2 and R^3 are as defined in formula (I), with a compound of general formula




wherein Y represents a bond or a group $(CH_2)_{1-6}$, $O(CH_2)_{2-6}$, $(CH_2)_{1-3}O(CH_2)_{2-3}$, $NR^5(CH_2)_{2-6}$ or $(CH_2)_{1-3}NR^5(CH_2)_{2-3}$ and R^4 is as defined in formula (I), in the presence of 1,1-(azodicarbonyl)dipiperidine and tributylphosphine (under conditions of the Mitsunobu reaction: Tetrahedron Lett. (1993), 34, 1639); or

(iii) when X represents a bond, an oxygen atom or a group $O(CH_2)_{1-6}$, $O(CH_2)_{2-6}O$, $O(CH_2)_{2-3}O(CH_2)_{1-3}$, NR^5 , $NR^5(CH_2)_{1-6}$, $NR^5(CH_2)_{2-6}O$ or $NR^5(CH_2)_{2-3}O(CH_2)_{1-3}$ and A is $NHC(O)$, reacting a compound of general formula




wherein one of R^{14} and R^{15} represents a group $-X'-R^4$ and the other of R^{14} and R^{15} represents a hydrogen atom, X' represents a bond, an oxygen atom or a group $O(CH_2)_{1-6}$, $O(CH_2)_{2-6}O$, $O(CH_2)_{2-3}O(CH_2)_{1-3}$, NR^5 , $NR^5(CH_2)_{1-6}$, $NR^5(CH_2)_{2-6}O$ or $NR^5(CH_2)_{2-3}O(CH_2)_{1-3}$, L^2 represents a leaving group (e.g. a hydroxyl or chloride leaving

group) and R^2 , R^3 , R^4 and R^5 are as defined in formula (I), with a compound of general formula




5 wherein m and R^1 are as defined in formula (I), optionally in the presence of a coupling agent (e.g. 1,1'-carbonyldiimidazole); or

(iv) when X represents a bond, an oxygen atom or a group $O(CH_2)_{1-6}$, $O(CH_2)_{2-6}O$, $O(CH_2)_{2-3}O(CH_2)_{1-3}$, NR^5 , $NR^5(CH_2)_{1-6}$, $NR^5(CH_2)_{2-6}O$ or
 10 $NR^5(CH_2)_{2-3}O(CH_2)_{1-3}$ and A is $C(O)NH$, reacting a compound of general formula




wherein R^2 and R^3 are as defined in formula (I) and R^{14} and R^{15} are as defined in formula
 15 (VI) in (iii) above, with a compound of general formula



wherein m and R^1 are as defined in formula (I), in the presence of a base (e.g. diisopropylamine); or

(v) when X represents a bond or a group NR^5 , $\text{NR}^5(\text{CH}_2)_{1-6}$, $\text{NR}^5(\text{CH}_2)_{2-6}\text{O}$ or $\text{NR}^5(\text{CH}_2)_{2-3}\text{O}(\text{CH}_2)_{1-3}$, reacting a compound of general formula



wherein one of R^{16} and R^{17} represents a leaving group, L^3 , such as a halogen atom and the other of R^{16} and R^{17} represents a hydrogen atom and m , A , R^1 , R^2 and R^3 are as defined in formula (I), with a compound of general formula



wherein Z represents a hydrogen atom or a group NHR^5 , $(\text{CH}_2)_{1-6}\text{NHR}^5$, $\text{O}(\text{CH}_2)_{2-6}\text{NHR}^5$ or a group $(\text{CH}_2)_{1-3}\text{O}(\text{CH}_2)_{2-3}\text{NHR}^5$ and R^4 and R^5 are as defined in formula (I), optionally in the presence of a palladium catalyst (e.g. palladium acetate), a phosphine ligand (e.g. BINAP) and a base (e.g. cesium carbonate); or

(vi) when X represents a group CH_2O , reacting a compound of formula (II) as defined in (i) above with a compound of formula (V) as defined in (ii) above wherein Y represents a bond, in the presence of a base (e.g. sodium hydride) or in the presence of a metal salt (e.g. silver trifluoromethanesulfonate); or


(vii) when X represents a group CH_2NR^5 , reacting a compound of formula (II) as defined in (i) above with a compound of formula (XI) as defined in (v) above wherein Z represents a group NHR^5 ; or

(viii) when X represents a group $\text{CH}_2\text{O}(\text{CH}_2)_{1-3}$ or $\text{CH}_2\text{O}(\text{CH}_2)_{2-3}\text{O}$, reacting a compound of formula (II) as defined in (i) above with a compound of formula (V) as defined in (ii) above wherein Y represents a group $(\text{CH}_2)_{1-3}$ or $\text{O}(\text{CH}_2)_{2-3}$, in the presence

of a base (e.g. sodium hydride) or in the presence of a metal salt (e.g. silver trifluoromethanesulfonate); or


(ix) when X represents a group $\text{CH}_2\text{NR}^5\text{CH}_2$ or $\text{CH}_2\text{NR}^5(\text{CH}_2)_{2-3}\text{O}$ reacting a compound of formula (II) as defined in (i) above with a compound of formula (XI) as defined in (v) above wherein Z represents a group CH_2NHR^5 or $\text{O}(\text{CH}_2)_{2-3}\text{NHR}^5$; or

(x) when X represents a group CH_2 and R^4 represents an unsubstituted 4- to 6-membered saturated aliphatic heterocyclic ring system containing one nitrogen atom, reacting a compound of formula (II) as defined in (i) above, with a compound of general formula




wherein s and t independently represent 1 or 2; or

(xi) when X represents a group CO , CONR^5 , NR^5CO , SO_2 , NR^5SO_2 or SO_2NR^5 and A is NHC(O) , reacting a compound of general formula



wherein one of R^{18} and R^{19} represents a group $-\text{X}''-\text{R}^4$ and the other of R^{18} and R^{19} represents a hydrogen atom, X'' represents a group CO , CONR^5 , NR^5CO , SO_2 , NR^5SO_2 or SO_2NR^5 , L^4 represents a leaving group (e.g. a hydroxyl or chloride leaving group) and R^2 , R^3 , R^4 and R^5 are as defined in formula (I), with a compound of formula (VII) as defined in (iii) above, optionally in the presence of a coupling agent (e.g. 1,1'-carbonyldiimidazole); or

(xii) when X represents a group CO, CONR^5 , NR^5CO , SO_2 , NR^5SO_2 or SO_2NR^5 and A is C(O)NH , reacting a compound of general formula



wherein R^2 and R^3 are as defined in formula (I) and R^{18} and R^{19} are as defined in formula (XIII) in (xi) above, with a compound of formula (IX) as defined in (iv) above, in the presence of a base (e.g. diisopropylamine); or

(xiii) when X represents a sulfur atom, reacting a compound of formula (X) as defined in (v) above, with an organolithium reagent such as n-butyllithium (e.g. at -70°C) and then with a compound of general formula



wherein Tol represents a tolyl group (4-methylphenyl) and R^4 is as defined in formula (I); or

(xiv) when X represents a CHOH or CH_2 group, reacting a compound of formula (X) as defined in (v) above, with an organolithium reagent (e.g. methylolithium/t-butyllithium or n-butyllithium at -70°C) and then with a compound of general formula



wherein R^4 is as defined in formula (I), optionally followed by a reduction reaction, e.g. with methyloxalylchloride and triethylamine followed by tributyltin hydride in the presence of azobisisobutyronitrile; or


(xv) when X represents a bond, reacting a compound of formula (X) as defined in (v) above, with an organolithium reagent such as n-butyllithium (e.g. at -70°C) and then with a compound of general formula



wherein R^4 is as defined in formula (I), optionally followed by a reduction reaction, e.g. with methyloxalylchloride and triethylamine followed by tributyltin hydride in the presence of azobisisobutyronitrile;

5 (xvi) when X represents a group SO, oxidising a corresponding compound of formula (I) in which X represents a sulphur atom (e.g. using, as oxidising agent, 3-chloroperoxybenzoic acid or potassium peroxymonosulphate (commercially sold under the trade mark "OXONE")); or

10 (xvii) when X represents a group SCH_2 , reacting a compound of formula (X) as defined in (v) above, with an organolithium reagent (e.g. methyllithium and/or t-butyllithium at $-70^\circ C$) and then with a compound of general formula



15 wherein R^4 is as defined in formula (I); or


(xviii) when X represents a group $SOCH_2$ or SO_2CH_2 , oxidising a corresponding compound of formula (I) in which X represents a group SCH_2 (e.g. using, as oxidising agent, 3-chloroperoxybenzoic acid or potassium peroxymonosulphate (commercially sold under the trade mark "OXONE")); or

20

(xix) when X represents a group $CH=$, reacting a compound of formula (II) as defined in (i) above with trimethyl phosphite and then with a compound of formula (XVII) as defined in (xv) above in the presence of a base (e.g. lithium diisopropylamide); or

25

(xx) when X represents a group $(CH_2)_{1-6}$, reacting a compound of general formula




wherein one of R^{20} and R^{21} represents a group CHO or a group $(CH_2)_{1-5}CHO$ and the other of R^{20} and R^{21} represents a hydrogen atom, and m , A , R^1 , R^2 and R^3 are as defined in formula (I), with a compound of general formula (XX), R^4-H , wherein R^4 is as defined in formula (I), in the presence of a reducing agent (e.g. sodium triacetoxyborohydride, in a suitable solvent such as dichloroethane); or

(xxi) when X represents a group $(CH_2)_{1-6}NR^5$, $(CH_2)_{1-3}NR^5(CH_2)_{1-3}$ or $(CH_2)_{1-3}NR^5(CH_2)_{2-3}O$, reacting a compound of formula (XIX) as defined in (xx) above, with a compound of general formula (XXI), R^4-Z' , wherein Z' represents a group NHR^5 , $(CH_2)_{1-3}NHR^5$, $O(CH_2)_{2-3}NHR^5$ and R^4 and R^5 are as defined in formula (I), in the presence of a reducing agent (e.g. sodium triacetoxyborohydride, in a suitable solvent such as dichloroethane); or

(xxii) when X represents a group $(CH_2)_{1-3}O(CH_2)_{1-3}$ or $(CH_2)_{1-3}O(CH_2)_{2-3}O$, reacting a compound of formula (XIX) as defined in (xx) above in which one of R^{20} and R^{21} represents a group CHO or a group $(CH_2)_{1-2}CHO$ and the other of R^{20} and R^{21} represents a hydrogen atom, with a reducing agent (such as sodium borohydride), followed by reaction with a compound of general formula (XXII), R^4-E , wherein E represents a group $(CH_2)_{1-3}L^5$ or $O(CH_2)_{2-3}L^5$, L^5 is a leaving group (such as a halogen atom or a sulphonate ester group, e.g. p-toluenesulphonate) and R^4 is as defined in formula (I), in the presence of a base (such as sodium hydride); or

(xxiii) when X represents a group $(\text{CH}_2)_{1-6}$, reacting a compound of formula (II) as defined in (i) above with trimethylphosphite and then with a compound of formula (XVI) as defined in (xiv) above, a compound of formula (XVII) as defined in (xv) above or a compound of general formula (XVIA), $\text{R}^4(\text{CH}_2)_{1-4}\text{CHO}$ in which R^4 is as defined in formula (I), in the presence of a base (e.g. lithium diisopropylamide), followed by a reduction reaction (for example, with hydrogen and a platinum oxide catalyst); or

(xxiv) when X represents a group $(\text{CH}_2)_{2-6}\text{O}$, reacting a compound of general formula



wherein one of R^{22} and R^{23} represents a group $(\text{CH}_2)_{2-6}\text{L}^6$ and the other of R^{20} and R^{21} represents a hydrogen atom, L^6 represents a leaving group (e.g. a halogen atom or a sulphonate ester group such as p-toluenesulphonate) and m, A, R^1 , R^2 and R^3 are as defined in formula (I), with a compound of formula (V) as defined in (ii) above in which Y represents a bond; or


(xxv) when X represents a group $\text{CR}'(\text{OH})$ in which R' is a $\text{C}_1\text{-C}_6$ alkyl group, oxidising a corresponding compound of formula (I) in which X represents $\text{CH}(\text{OH})$ (e.g. using the oxidant dimethylsulphoxide/oxalyl chloride), followed by reaction with a $\text{C}_1\text{-C}_6$ alkyllithium reagent; or

(xxvi) when X represents a group CH_2S , reacting a compound of formula (II) as defined in (i) above with a compound of general formula (XXIV), $\text{R}^4\text{-SH}$, wherein R^4 is as defined in formula (I), in the presence of a base (e.g. sodium hydride); or

(xxvii) when X represents a group CH_2SO or CH_2SO_2 , oxidising a corresponding compound of formula (I) in which X represents a group CH_2S (e.g. using, as oxidising agent, 3-chloroperoxybenzoic acid or potassium peroxymonosulphate (commercially sold under the trade mark "OXONE")); or


(xxviii) when X represents a group CH_2 and R^4 represents a 3-piperidiny1 or 2-piperazinyl group, reacting a compound of formula (II) as defined in (i) above with a reagent formed by combining pyridine or pyrazine with an aluminium hydride reagent (e.g. lithium aluminium hydride), followed by a reduction reaction (e.g. with hydrogen and a platinum catalyst); or

(xxix) when X represents a group CH= and R^4 represents a 3-piperidiny1 group, reacting a compound of general formula




wherein one of R^{24} and R^{25} represents an aldehyde group -CHO , and the other of R^{24} and R^{25} represents a hydrogen atom and m , A , R^1 , R^2 and R^3 are as defined in formula (I), with 2,3,4,5-tetrahydropyridine (*Bull.Chem.Soc.Jpn.* 1983, 56, 3199), followed by a reduction reaction (e.g. with sodium borohydride in a protic solvent such as methanol); or

(xxx) when X represents a bond, NR^5 or $\text{NR}^5(\text{CH}_2)_{1-6}$ and R^4 represents a carbon-linked piperidyl or piperazinyl group, reducing a compound of general formula




wherein one of R^{26} and R^{27} represents a pyridyl, pyrazinyl, NR^5 -pyridyl, NR^5 -pyrazinyl, $\text{NR}^5(\text{CH}_2)_{1-6}$ -pyridyl or $\text{NR}^5(\text{CH}_2)_{1-6}$ -pyrazinyl group and the other of R^{26} and R^{27} represents a hydrogen atom, and m, A, R^1 , R^2 and R^3 are as defined in formula (I), with a source of hydrogen and a hydrogenation catalyst (such as platinum oxide); or

(xxxi) when X represents a group $\text{CH}_2\text{O}(\text{CH}_2)_{1-3}$ or $\text{CH}_2\text{O}(\text{CH}_2)_{2-3}\text{O}$ and A is NHC(O) , reacting a compound of general formula



wherein one of R^{28} and R^{29} represents a group $-\text{X}^m-\text{R}^4$ and the other of R^{28} and R^{29} represents a hydrogen atom, X^m represents a group $\text{CH}_2\text{O}(\text{CH}_2)_{1-3}$ or $\text{CH}_2\text{O}(\text{CH}_2)_{2-3}\text{O}$, L^7 represents a leaving group (e.g. a hydroxyl or chloride leaving group) and R^2 , R^3 and R^4 are as defined in formula (I), with a compound of formula (VII) as defined in (iii) above, optionally in the presence of a coupling agent (e.g. 1,1'-carbonyldiimidazole); or

(xxxii) when X represents a group $\text{CH}_2\text{O}(\text{CH}_2)_{1-3}$ or $\text{CH}_2\text{O}(\text{CH}_2)_{2-3}\text{O}$ and A is C(O)NH , reacting a compound of general formula




wherein R^2 and R^3 are as defined in formula (I) and R^{28} and R^{29} are as defined in formula (XXVII) in (xxxi) above, with a compound of formula (IX) as defined in (iv) above, in the presence of a base (e.g. diisopropylamine);

and optionally after (i), (ii), (iii), (iv), (v), (vi), (vii), (viii), (ix), (x), (xi), (xii), (xiii), (xiv), (xv), (xvi), (xvii), (xviii), (xix), (xx), (xxi), (xxii), (xxiii), (xxiv), (xxv), (xxvi), (xxvii), (xxviii), (xxix), (xxx), (xxxi) or (xxxii) converting the compound of formula (I) to a further compound of formula (I) and, if desired, forming a pharmaceutically acceptable salt or solvate of the compound of formula (I).

The processes of the invention may conveniently be carried out in a solvent, e.g. an organic solvent such as dichloromethane, dichloroethane, tetrahydrofuran, dioxane, xylene or dimethylformamide, at a temperature, e.g. in the range from 0 to 200 °C, preferably in the range from 0 to 150 °C.


Compounds of formula (II) in which A is NHC(O) may be prepared by reacting a compound of general formula



wherein L^{10} represents a leaving group (e.g. a hydroxyl or chloride leaving group) and R^2 , R^3 , R^{10} and R^{11} are as defined in formula (II), with a compound of formula (VII) as

defined above, optionally in the presence of a coupling agent (e.g. 1,1'-carbonyldiimidazole).


Compounds of formula (XXX) in which one of R^{10} and R^{11} represents a hydrogen atom and the other of R^{10} and R^{11} represents a group $-CH_2L^1$ and L^1 represents a bromine atom can be prepared by reacting a compound of general formula



wherein one of R^{30} and R^{31} represents a hydrogen atom and the other of R^{30} and R^{31} represents a methyl group and R^2 and R^3 are as defined in formula (I), with


N-bromosuccinimide and catalytic azobisisobutyronitrile or dibenzoylperoxide, optionally followed by chlorination with oxalyl chloride and catalytic dimethylformamide or with thionyl chloride.

Compounds of formula (II) in which A is $C(O)NH$ and L^1 represents, for example, a bromine atom may be prepared by reacting a compound of general formula




wherein R^2 and R^3 are as defined in formula (I) and R^{30} and R^{31} are as defined in formula (XXXI) above, with a compound of formula (IX) as defined above, in the presence of a base (e.g. diisopropylethylamine), followed by reaction with N-bromosuccinimide and catalytic azobisisobutyronitrile or dibenzoylperoxide.

Compounds of formula (IV) in which A is NHC(O) may be prepared in an analogous manner to compounds of formula (II) in which A is NHC(O) , using instead of the intermediate compound of formula (XXX), an intermediate compound of general formula




5 wherein L^{11} represents a leaving group (e.g. a hydroxyl or chloride leaving group) and R^2 , R^3 , R^{12} and R^{13} are as defined in formula (IV).

Compounds of formula (IV) in which A is C(O)NH may be prepared by reacting a compound of general formula




10 wherein R^2 , R^3 , R^{12} and R^{13} are as defined in formula (IV), with a compound of formula (IX) as defined above, optionally in the presence of a base (e.g. diisopropylethylamine).

Compounds of formula (VI) can be prepared by reacting a compound of general
15 formula




20 wherein R^{32} represents a hydrogen atom or a $\text{C}_1\text{-C}_6$ alkyl group, one of R^{33} and R^{34} represents a leaving group, L^{12} , such as a halogen atom (e.g. bromine or iodine) or a trifluoromethanesulfonate group and the other of R^{33} and R^{34} represents a hydrogen atom, and R^2 and R^3 are as defined in formula (VI), with a compound of general formula



wherein X' and R^4 are as defined in formula (VI), in the presence of a palladium catalyst (e.g. palladium acetate), a phosphine ligand (e.g. BINAP) and a base (e.g. cesium carbonate) (1996 *J. Am. Chem. Soc.*, 7215-6; 1997 *J. Am. Chem. Soc.*, 3395), followed by
 5 a hydrolysis reaction (e.g. with sodium hydroxide) and optionally a chlorination reaction (e.g. with oxalyl chloride and catalytic dimethylformamide or with thionyl chloride).


Compounds of formula (VIII) may conveniently be prepared by reacting a compound of formula (VI) in which L^2 represents a hydroxyl group with diphenylphosphoryl azide in
 10 the presence of a base such as triethylamine.

Compounds of formula (X) in which A is NHC(O) may be prepared in an analogous manner to compounds of formula (II) in which A is NHC(O) , using instead of the intermediate compound of formula (XXX), an intermediate compound of general formula



15 wherein L^{13} represents a leaving group (e.g. a hydroxyl or chloride leaving group) and R^2 , R^3 , R^{16} and R^{17} are as defined in formula (X).


Compounds of formula (X) in which A is C(O)NH may be prepared in an analogous
 20 manner to compounds of formula (IV) in which A is C(O)NH , using instead of the intermediate compound of formula (XXXIV), an intermediate compound of general formula



wherein R^2 , R^3 , R^{16} and R^{17} are as defined in formula (X).


Compounds of formula (XII) can be prepared as described in Syn. Lett. (1998) 379-380.

Compounds of formula (XIII) in which X'' represents a group CO, $CONR^5$, SO_2 or SO_2NR^5 can be prepared by reacting a compound of general formula



wherein one of R^{35} and R^{36} represents a group COL^{14} or SO_2L^{14} and the other of R^{25} and R^{26} represents a hydrogen atom, L^{14} represents a leaving group (e.g. a halogen atom), R^{37} represents a hydrogen atom or a C_1 - C_6 alkyl group, and R^2 and R^3 are as defined in formula (XIII), with a compound of formula (XXXVI) in which X' represents a bond or a group NR^5 , in the presence of a base such as diisopropylethylamine and catalytic dimethylaminopyridine, followed by a hydrolysis reaction (e.g. sodium hydroxide) and, optionally, a chlorination reaction (e.g. with oxalyl chloride and catalytic dimethylformamide or with thionyl chloride).

Compounds of formula (XIII) in which X'' represents a group represents a group NR^5CO or NR^5SO_2 can be prepared by reacting a compound of general formula




wherein one of R^{38} and R^{39} represents a group NHR^5 and the other of R^{38} and R^{39} represents a hydrogen atom, R^{37} is as defined for compound (XXXIX), and R^2 and R^3 are as defined in formula (XIII), with a compound general formula (XLI), R^4 -J, wherein

J represents a group COCl or SO₂Cl and R⁴ is as defined in formula (I), in the presence of a base such as diisopropylethylamine.

Compounds of formula (XIV) may conveniently be prepared by reacting a compound of formula (XIII) in which L⁴ represents a hydroxyl group with diphenylphosphoryl azide in the presence of a base such as triethylamine.

Compounds of formula (XIX) in which one of R²⁰ and R²¹ represents a group (CH₂)₁₋₅CHO and the other of R²⁰ and R²¹ represents a hydrogen atom can be prepared by oxidising a compound of general formula




wherein one of R⁴⁰ and R⁴¹ represents a group (CH₂)₂₋₆OH and the other of R⁴⁰ and R⁴¹ represents a hydrogen atom, and m, A, R¹, R² and R³ are as defined in formula (I), using as the oxidising agent, for example, Dess Martin Periodinane reagent.

Compounds of formula (XLII) in which one of R⁴⁰ and R⁴¹ represents a group (CH₂)₂OH and the other of R⁴⁰ and R⁴¹ represents a hydrogen atom can be prepared from a compound of general formula (X) as defined above, an organolithium reagent such as methyllithium (at -70 °C) followed by n-butyllithium (at -70 °C), and then treatment with ethylene oxide.

Compounds of formula (XLII) in which one of R⁴⁰ and R⁴¹ represents a group (CH₂)₃₋₆OH and the other of R⁴⁰ and R⁴¹ represents a hydrogen atom can be prepared by

reacting a compound of general formula (X) as defined above with a compound of general formula



- 5 in the presence of a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0), followed by reduction with, for example, hydrogen and a platinum oxide catalyst.

Compounds of formula (XIX) in which one of R^{20} and R^{21} represents a group CHO and the other of R^{20} and R^{21} represents a hydrogen atom (which are equivalent to
10 compounds of formula (XXV)) can be prepared from a compound of general formula (X) as defined above, with an organolithium reagent such as methyllithium (at -70°C) followed by n-butyllithium (at -70°C) and then with dimethylformamide.

Compounds of formula (XXIII) in which L^6 represents an iodine atom or
15 p-toluenesulphonyloxy group may be prepared by reacting a compound of formula (XLII) as defined above with iodine/triphenylphosphine/imidazole or with a sulphonyl chloride such as p-toluenesulphonyl chloride, in the presence of a base such as diisopropylethylamine.

20 Compounds of formula (XXVI) in which one of R^{26} and R^{27} represents a pyridyl or pyrazinyl group and the other of R^{26} and R^{27} represents a hydrogen atom can be prepared from a compound of formula (X) as defined above by reaction with a pyridyl or pyrazinyl boronic acid in the presence of a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0).

25

Compounds of formula (XXVI) in which one of R^{26} and R^{27} represents a NR^5 -pyridyl, NR^5 -pyrazinyl, $\text{NR}^5(\text{CH}_2)_{1-6}$ -pyridyl or $\text{NR}^5(\text{CH}_2)_{1-6}$ -pyrazinyl group and the other of R^{26} and R^{27} represents a hydrogen atom can be prepared from a compound of formula (X) as defined above by reaction with a compound NHR^5 pyridyl,

NHR⁵pyrazinyl, NHR⁵(CH₂)₁₋₆-pyridyl or NHR⁵(CH₂)₁₋₆-pyrazinyl, in the presence of a palladium catalyst (e.g. palladium acetate), a phosphine ligand (e.g. BINAP) and a base (e.g. cesium carbonate).

5 Compounds of formulae (III), (V), (VII), (IX), (XI), (XV), (XVI), (XVIA), (XVII), (XVIII), (XX), (XXI), (XXII), (XXIV), (XXVII), (XXVIII), (XXXI), (XXXII), (XXXIII), (XXXIV), (XXXV), (XXXVI), (XXXVII), (XXXVIII), (XXXIX), (XL), (XLI), (XLII) and (XLIII) are either commercially available, are well known in the literature or may be prepared easily using known techniques.

10 Compounds of formula (I) can be converted into further compounds of formula (I) using standard procedures. For example, compounds of formula (I) in which one of R² and R³ represents a nitro group can be converted to compounds of formula (I) in which one of R² and R³ represents an amino group by reduction using iron powder and ammonium
15 chloride in ethanol/water under reflux conditions. The latter compounds can in turn be converted into compounds of formula (I) in which one of R² and R³ represents a halogen atom, e.g. chlorine, by diazotization (e.g. with sodium nitrite) and reaction with copper chloride. Compounds of formula (I) in which R⁶ or R⁷ represents a hydrogen atom can be converted to compounds of formula (I) in which R⁶ or R⁷ represents a C₁-C₆ alkyl,
20 C₂-C₆ hydroxyalkyl, C₃-C₈ cycloalkyl or a 3- to 8-membered saturated heterocyclic ring by standard chemical procedures.

It will be appreciated by those skilled in the art that in the processes of the present invention certain functional groups such as hydroxyl or amino groups in the starting
25 reagents or intermediate compounds may need to be protected by protecting groups. Thus, the preparation of the compounds of formula (I) may involve, at an appropriate stage, the removal of one or more protecting groups.

The protection and deprotection of functional groups is described in 'Protective
30 Groups in Organic Chemistry', edited by J.W.F. McOmie, Plenum Press (1973) and

'Protective Groups in Organic Synthesis', 2nd edition, T.W. Greene and P.G.M. Wuts, Wiley-Interscience (1991).

The compounds of formula (I) above may be converted to a pharmaceutically acceptable salt or solvate thereof, preferably an acid addition salt such as a hydrochloride, hydrobromide, phosphate, acetate, fumarate, maleate, tartrate, citrate, oxalate, methanesulphonate or *p*-toluenesulphonate, or an alkali metal salt such as a sodium or potassium salt.

Certain compounds of formula (I) are capable of existing in stereoisomeric forms. It will be understood that the invention encompasses all geometric and optical isomers of the compounds of formula (I) and mixtures thereof including racemates. Tautomers and mixtures thereof also form an aspect of the present invention.

The compounds of the present invention are advantageous in that they possess pharmacological activity. They are therefore indicated as pharmaceuticals for use in the treatment of rheumatoid arthritis, osteoarthritis, psoriasis, allergic dermatitis, asthma, chronic obstructive pulmonary disease (COPD), hyperresponsiveness of the airway, septic shock, glomerulonephritis, irritable bowel disease, Crohn's disease, ulcerative colitis, atherosclerosis, growth and metastases of malignant cells, myoblastic leukaemia, diabetes, Alzheimer's disease, meningitis, osteoporosis, burn injury, ischaemic heart disease, stroke and varicose veins.

Accordingly, the present invention provides a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined for use in therapy.

In another aspect, the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for use in therapy.

In the context of the present specification, the term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The terms "therapeutic" and "therapeutically" should be construed accordingly.

5

The invention further provides a method of effecting immunosuppression (e.g. in the treatment of rheumatoid arthritis, irritable bowel disease, atherosclerosis or psoriasis) which comprises administering a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore
10 defined to a patient.

The invention also provides a method of treating an obstructive airways disease (e.g. asthma or COPD) which comprises administering to a patient a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate
15 thereof, as hereinbefore defined to a patient.

For the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disorder indicated. The daily dosage of the compound of formula (I)/salt/solvate
20 (active ingredient) may be in the range from 0.001 mg/kg to 30 mg/kg.

The compounds of formula (I) and pharmaceutically acceptable salts and solvates thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in which the formula (I) compound/salt/solvate (active
25 ingredient) is in association with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of administration, the pharmaceutical composition will preferably comprise from 0.05 to 99 %w (per cent by weight), more preferably from 0.10 to 70 %w, of active ingredient, and, from 1 to 99.95 %w, more preferably from 30 to 99.90 %w, of a pharmaceutically acceptable adjuvant, diluent or carrier, all percentages by weight being
30 based on total composition.

Thus, the present invention also provides a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in association with a pharmaceutically acceptable adjuvant, diluent or carrier.


The invention further provides a process for the preparation of a pharmaceutical composition of the invention which comprises mixing a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined with a pharmaceutically acceptable adjuvant, diluent or carrier.

The pharmaceutical composition of the invention may be administered topically (e.g. to the lung and/or airways or to the skin) in the form of solutions, suspensions, heptafluoroalkane aerosols and dry powder formulations; or systemically, e.g. by oral administration in the form of tablets, capsules, syrups, powders or granules, or by parenteral administration in the form of solutions or suspensions, or by subcutaneous administration or by rectal administration in the form of suppositories or transdermally.

The present invention will now be further explained by reference to the following illustrative examples.

Example 1

2-Nitro-3-piperazin-1-yl -*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



To a suspension of 3-chloro-2-nitrobenzoic acid (2.68 g) in dichloromethane (10 ml) at 0°C was added oxalyl chloride (3 ml) and dimethylformamide (1 drop). The resulting mixture was stirred at room temperature under a nitrogen atmosphere for 1 hour, then concentrated under reduced pressure to yield a solid. The solid was dissolved in dichloromethane (10 ml) and cooled to 0°C. A solution of 1-adamantanemethylamine (2.19g) and *N,N*-diisopropylethylamine (11 ml) in dichloromethane (10 ml) was added portion-wise and the resulting solution allowed to stir at room temperature under a nitrogen atmosphere for 2h. The reaction mixture was poured into water and the organic phase separated and washed with 2N hydrochloric acid, 10% aqueous sodium hydroxide and saturated brine. The organic phase was then dried over sodium sulfate, filtered and concentrated under reduced pressure and the resulting solid re-crystallized from *iso*-propanol to afford the subtitle compound as a solid (3.52 g).

MS (APCI +ve) 349 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.74 (1H, t); 7.89 (1H, m); 7.75-7.69 (2H, m); 2.91 (2H, d), 1.93 (3H, bs); 1.64 (6H, dd); 1.47 (6H, d)

b) 3-(4-{1,1-dimethylethyl}oxycarbonyl]-piperazin-1-yl)-2-nitro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

A mixture of 3-chloro-2-nitro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (2.80 g, Example 1a) and piperazine-1-carboxylic acid, *tert*-butyl ester (7.47 g) in dry dimethyl sulfoxide (10 ml) was heated at 120°C under a nitrogen atmosphere for 24h. The cooled reaction mixture was diluted with water and extracted thrice with ethyl acetate. The combined extracts were washed with water, dried over sodium sulfate, filtered, and the filtrate concentrated under reduced pressure to give a solid. Purification by chromatography over silica gel, eluting with *iso*-hexane/ethyl acetate (2:1) gave the subtitle compound as a solid (3.8 g).

MS (APCI +ve) 499 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.55 (1H, t); 7.62-7.59 (2H, m); 7.43 (1H, dd); 3.38 (4H, bt); 2.90-2.84 (6H, m), 1.93 (3H, bs); 1.63 (6H, dd); 1.47 (6H, d); 1.41 (9H, s)

c) 2-Nitro-3-piperazin-1-yl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide


5 A solution of 3-(4-{1,1-dimethylethyl}oxycarbonyl]-piperazin-1-yl)-2-nitro-N-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (0.58 g, Example 1b) and hydrochloric acid (6.4 ml, 4N in dioxane) in tetrahydrofuran (20 ml) was stirred at room temperature under a nitrogen atmosphere for 18h. The reaction mixture was concentrated under reduced pressure and the residue dissolved in water, made basic with solid sodium
10 bicarbonate and extracted with dichloromethane three times. The combined organic extracts were dried over magnesium sulfate, filtered and the filtrate concentrated under reduce pressure to give a solid. Purification by chromatography over silica gel, eluting with 10% methanol in dichloromethane afforded the title compound as a solid (0.165 g).

15 MS (APCI +ve) 399 (M+H)+

¹H NMR (DMSO-d₆) δ 8.52 (1H, t); 7.59 (1H, t); 7.51 (1H d); 7.35 (1H, d); 2.88 (2H, d); 2.81 (4H, m); 2.37 (4H, m); 1.93 (3H, bs); 1.67 (3H, d); 1.60 (3H, d); 1.47 (6H, s)

Example 2

20 2-Amino-3-piperazin-1-yl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt



a) 2-Amino-3-(4-{1,1-dimethylethyl}oxycarbonyl]-piperazin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

25 A suspension of 3-(4-{1,1-dimethylethyl}oxycarbonyl]-piperazin-1-yl)-2-nitro-N-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (3.8 g, Example 1b), iron powder (2.13 g)

and ammonium chloride (2.04 g) in 2:1 ethanol/water (90 ml) was heated at reflux, under a nitrogen atmosphere, for 2h. The cooled reaction mixture was filtered and the filtrate partitioned between water and ethyl acetate. The organic layer was separated and washed with water twice further, dried over magnesium sulfate, filtered and the filtrate concentrated under reduced pressure to give a residue. Purification of the residue by chromatography over silica gel, eluting with 20% ethyl acetate in *iso*-hexane, yielded the subtitle compound as a solid (2.27 g).

MS (APCI +ve) 469 (M+H)⁺

b) 2-Amino-3-piperazin-1-yl-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt


Prepared as described in Example 1c) using 2-amino-3-(4-{1,1-dimethyl ethyl} oxycarbonyl)-piperazin-1-yl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.2 g, Example 2a) and hydrochloric acid (5 ml, 4N in dioxane). The reaction mixture was concentrated under reduced pressure to give a solid which when triturated with diethyl ether gave the title compound as a solid (0.2 g).

MS (APCI +ve) 369 (M-2HCl)⁺

¹H NMR (DMSO-d₆) δ 9.16 (2H, bs); 8.14 (1H, t); 7.37 (1H, d); 7.07 (1H, d); 6.64 (1H, t); 3.27 (4H, bs); 2.98 (4H, bs); 2.95 (2H, d); 1.93 (3H, bs); 1.67 (3H, d); 1.59 (3H, d); 1.48 (6H, s).

Example 3

2-Chloro-3-piperazin-1-yl -*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



a) **2-Chloro-3-(4-{1,1-dimethylethyl}oxycarbonyl]-piperazin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

To a solution of 2-amino-3-(4-{1,1-dimethylethyl}oxycarbonyl]-piperazin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (1g, Example 2a) in tetrahydrofuran
5 (23 ml) was added 1M aqueous hydrochloric acid (2.78 ml) and water (10 ml). The solution was cooled to 0°C and sodium nitrite (1.91 g) added portion-wise, whilst maintaining the internal temperature below 5°C. After stirring at 0-5°C for 0.5h, a pre-cooled suspension of copper (I) chloride (10.58 g) and copper (II) chloride in water (20ml) was added portion-wise to the pale yellow suspension. The mixture was stirred at 0°C for
10 0.5h then at room temperature for 0.5h. The reaction mixture was poured into a mixture of water and dichloromethane and 1/1 : 0.88 ammonia/water was added until the aqueous phase was homogeneous. The layers were separated and the aqueous phase extracted twice further with dichloromethane. The combined organic extracts were washed with 1/1 : 0.88 ammonia/water until the aqueous layer was colourless, dried over sodium sulfate, filtered
15 and concentrated under reduced pressure to yield an oil. Purification by chromatography on silica gel, eluting with 20-35% ethyl acetate/*iso*-hexane gave the subtitle compound as a solid (0.45 g).

MS (APCI +ve) 388 (M-BOC)⁺

20 ¹H NMR (DMSO-d₆) δ 8.27 (1H, t); 7.32 (1H, t); 7.19 (1H, d); 7.04 (1H, d); 3.48 (4H, m); 2.93-2.91 (6H, m); 1.94 (3H, bs); 1.64 (3H, d); 1.59 (3H, d); 1.52 (6H, s); 1.43 (9H, s).

b) **2-Chloro-3-piperazin-1-yl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

To a solution of 2-Chloro-3-(4-{1,1-dimethylethyl}oxycarbonyl]-piperazin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.45 g, Example 3a) in dichloromethane
25 (10 ml) was added trifluoroacetic acid (5 ml). After stirring at room temperature under a nitrogen atmosphere the reaction mixture was concentrated under reduced pressure to give a gum. The gum was partitioned between water and dichloromethane and made basic with solid sodium bicarbonate. The layers were separated and the aqueous layer extracted twice
30 further with dichloromethane. The combined organic extracts were washed twice with


water, saturated brine then dried over magnesium sulfate, filtered and the filtrate concentrated under reduced pressure to give a foam. The foam was purified by normal phase HPLC (0-20% ethanol/dichloromethane) and chromatography over silica gel, eluting 10% methanol in dichloromethane, to afford the title compound as a foam (0.05 g).

MS (APCI +ve) 388/90 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.24 (1H, t); 7.31 (1H, t); 7.15 (1H, d); 7.00 (1H, d); 2.96-2.87 (10H, m); 1.93 (3H, bs); 1.67 (3H, d); 1.59 (3H, d); 1.52 (6H, s).

Example 4

2-Chloro-5-piperazin-1-yl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



a) 2-Chloro-5-nitro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

To a solution of 2-chloro-5-nitrobenzoic acid (1.22g) in *N,N*-dimethylformamide (1.5 ml) was added carbonyldiimidazole (1.0 g). The resulting reaction mixture was stirred for 2.5h and then 1-adamantanemethylamine (1.0g) was added. After 14h the reaction mixture was partitioned between ethyl acetate and water and the organic layer was separated, washed with water and brine and then dried over sodium sulphate (Na₂SO₄). The organic layer was concentrated under reduced pressure to give a residue which was purified by silica gel chromatography (eluting with 3-10% methanol in dichloromethane) to yield the subtitle compound as a yellow solid (1.7 g).

MS (APCI +ve) 348/350 (M+H)⁺

¹H NMR (CDCl₃) δ 8.53 (1H, d), 8.2 (1H, dd), 7.6 (1H, d), 6.2 (1H, bs), 3.2 (2H, d), 2.0 (3H,bs), 1.8 (12H, m)

b) 5-Amino-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

A solution of the nitro compound from Example 4a, (0.50 g) and ammonium chloride (0.5g) were dissolved in 50% aqueous ethanol. Iron powder (0.5g) was added and the mixture stirred at reflux temperature for 3 hr before being cooled and solids removed by filtration. The mother liquors were treated with 10% sodium hydroxide solution and the product extracted into ethyl acetate. The organic solution was washed with brine, dried over sodium sulphate (Na₂SO₄) and concentrated to give a residue which was purified by silica gel chromatography to give the title compound as a white solid (0.4g).

MS (APCI +ve) 319/21 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.14 (1H, t); 7.03 (1H, dd); 6.56 (2H, m); 5.36 (2H, s); 2.89 (2H, d); 1.95 (3H, s); 1.7 (12H, m)

c) 2-Chloro-5-Piperazin-1-yl -*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide


To a solution of 5-amino-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (1.00 g, Example 4b) in xylene (20 ml) was added *bis*-(2-chloroethyl)amine hydrochloride salt (0.620g). The mixture was heated at 150°C for 12h (a dark solution is obtained). The cold solution was washed with 2M HCl, the aqueous layer washed with ethyl acetate then basified with sodium bicarbonate and extracted twice with dichloromethane. The organic layer was dried over magnesium sulfate, filtered and the filtrate concentrated under reduced pressure to give a foam. The crude material was purified on silica gel (0-10% ethanol/dichloromethane), to afford the title compound as a white solid (0.90 g).

MS (APCI +ve) 388/90 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.22 (1H, t); 7.22 (1H, d); 6.96 (1H, dd); 6.84 (1H, d); 3.50-3.20 (7H, m); 3.00-2.90 (2H, t); 2.91 (2H, d); 1.94 (3H, bs); 1.67 (3H, d); 1.59 (3H, d); 1.52 (6H, s).

Example 5

2-Chloro-5-(hexahydro-1*H*-1,4-diazepin-1-yl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) **4-[4-Chloro-3-(ethoxycarbonyl)phenyl]hexahydro-1*H*-1,4-diazepine-1-carboxylic acid, 1,1-dimethylethyl ester**

A mixture of 5-bromo-2-chloro-benzoic acid, ethyl ester (0.50 g), hexahydro-1*H*-1,4-diazepine-1-carboxylic acid, 1,1-dimethylethyl ester (0.46 g), cesium carbonate (0.86 g), palladium (II) acetate (8.5 mg) and (R)-BINAP (35 mg) in toluene (3 ml) was heated at 100 °C for 14h in a pressure vessel flushed with nitrogen. The cooled reaction mixture was poured into water and extracted (3 times) with ethyl acetate. The combined organic extracts were washed with saturated sodium chloride solution and then dried over magnesium sulfate. Evaporation under reduced pressure gave an oil which was purified by chromatography over silica gel, eluting with 20% ethyl acetate in *iso*-hexane to yield the subtitle compound as an oil (0.21 g).

MS (APCI +ve) 282/284 (M-BOC)⁺

b) **4-(3-Carboxy-4-chlorophenyl)hexahydro-1*H*-1,4-diazepine-1-carboxylic acid, 1,1-dimethylethyl ester**

A suspension of 4-[4-chloro-3-(ethoxycarbonyl)phenyl]hexahydro-1*H*-1,4-diazepine-1-carboxylic acid, 1,1-dimethylethyl ester (Example 5a, 0.21g), lithium hydroxide monohydrate (1.05ml of 3M solution in water) in 1:1 ethanol/water (7 ml) was stirred at room temperature for 14h. More lithium hydroxide monohydrate (0.55ml of 3M solution

in water) was added followed by tetrahydrofuran (1 ml). The resulting solution was stirred for 4h at room temperature then poured into water and extracted with diethyl ether. The aqueous phase was separated, acidified with 2M hydrochloric acid and then extracted with dichloromethane three times. The combined dichloromethane layers were dried over magnesium sulfate and evaporated under reduced pressure to yield the subtitle compound as a glass.

MS (APCI +ve) 298/300 (M-^tBu)⁺

c) **Hexahydro-4-[4-methyl-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-phenyl]-1H-1,4-diazepine-1-carboxylic acid, 1,1-dimethylethyl ester**

A solution of 4-(3-carboxy-4-chlorophenyl)hexahydro-1H-1,4-diazepine-1-carboxylic acid, 1,1-dimethylethyl ester (Example 5b, 0.10 g) and N,N'-carbonyldiimidazole (0.045 g) in dimethylformamide (3 ml) was stirred at room temperature for 2h. 1-Adamantanemethylamine (0.050 ml) was then added and stirring continued for 14h. The reaction mixture was poured into water and extracted with ethyl acetate three times. The ethyl acetate layers were combined and washed with 2M hydrochloric acid, 10% aqueous sodium hydroxide and brine, then dried over magnesium sulfate and concentrated under reduced pressure. Purification by chromatography on silica gel, eluting with 20-30% ethyl acetate in *iso*-hexane, gave the subtitle product as a gum which crystallised on standing.

MS (APCI +ve) 502/504 (M+H)⁺

d) **2-Chloro-5-(hexahydro-1H-1,4-diazepin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide, hydrochloride salt**

Hexahydro-4-[4-methyl-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-phenyl]-1H-1,4-diazepine-1-carboxylic acid, 1,1-dimethylethyl ester (from Example 5c) was dissolved in methanol (5 ml) and hydrochloric acid (0.5ml of a 4N solution in dioxane) was added. After stirring at room temperature for 14h, the mixture was evaporated to 2/3


original volume under reduced pressure. Diethyl ether was gradually added to the solution and the resulting precipitate collected by filtration, washed with diethyl ether and dried in vacuo to afford the title compound as a solid (0.027 g)

5 MS (APCI +ve) 402/404 (M+H)⁺

¹H NMR (DMSO-d₆) δ 9.11 (2H, bs); 8.18 (1H, t); 7.24 (1H, d); 6.81 (1H, dd); 6.71 (1H, d); 3.71 (2H, t); 3.50 (2H, t); 3.19 (2H, bs); 2.93 (2H, bs); 2.92 (2H, d); 2.08 (2H, m); 1.94 (3H, bs); 1.67 (3H, d); 1.59 (3H, bs); 1.52 (6H, s)

10 **Example 6**

5-(4-Amino-1-piperidinyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



15 **a) 2-Chloro-5-[4-[(1,1-dimethylethoxy)carbonyl]amino]-1-piperidinyl]-benzoic acid, ethyl ester**

Prepared as described in Example 5a) using 5-bromo-2-chloro-benzoic acid, ethyl ester (0.50 g), 4-piperidinyl-carbamic acid, 1,1-dimethylethyl ester (0.46 g), cesium carbonate (0.86 g), palladium (II) acetate (8.5 mg) and (R)-BINAP (35 mg) and toluene (3 ml) to afford the subtitle compound as an oil (0.17 g).

20

MS (APCI +ve) 383/385 (M+H)⁺

b) 2-Chloro-5-[4-[[[(1,1-dimethylethoxy)carbonyl]amino]-1-piperidinyl]- benzoic acid

Prepared as described in example 5b) using 2-chloro-5-[4-[[[(1,1-dimethylethoxy)-carbonyl]amino]-1-piperidinyl]-benzoic acid, ethyl ester (Example 6a, 0.17 g), lithium hydroxide monohydrate (0.88 ml of a 3M solution in water), 1:1 ethanol/water (7 ml) and tetrahydrofuran (1ml) to give the subtitle compound as a solid (0.14 g).

MS (APCI +ve) 354/356 (M+H)⁺

c) [1-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-piperidinyl]-carbamic acid, 1,1-dimethylethyl ester

Prepared as described in Example 5c) using 2-chloro-5-[4-[[[(1,1-dimethylethoxy)carbonyl]amino]-1-piperidinyl]- benzoic acid (Example 6b, 0.065 g), N,N'-carbonyldiimidazole (0.030 g), 1-adamantanemethylamine (0.032 ml) and dimethylformamide (3 ml) to give the subtitle compound as a solid.

MS (APCI +ve) 501/503 (M+H)⁺

d) 5-(4-Amino-1-piperidinyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt


Prepared as described in example 5d) above using [1-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-4-piperidinyl]-carbamic acid, 1,1-dimethylethyl ester (Example 6c), hydrochloric acid (0.5ml of a 4N solution in dioxane) and methanol (10 ml). The mixture was heated at reflux for 15 min. to complete the reaction. After evaporation to two-thirds of the original volume, a solid crystallised on standing which was collected by filtration and dried in vacuo to give the title compound as a solid (0.025 g).

MS (APCI +ve) 402/404 (M-HCl)⁺

¹H NMR (DMSO-d₆) δ 8.23 (1H, t); 8.11 (1H, bs); 7.28 (1H, d); 7.03 (1H, dd); 6.94 (1H, s); 3.74 (2H, d); 3.20 (1H, m); 2.91 (2H, d); 2.83 (2H, t); 1.98 (2H, bs); 1.94 (3H, bs); 1.69-1.58 (8H, m); 1.52 (6H, s)

Example 7

(+/-)-5-(3-Amino-1-pyrrolidiny)-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) (+/-)-2-Chloro-5-[3-[[[(1,1-dimethylethoxy)carbonyl]amino]-1-pyrrolidinyl]-benzoic acid, ethyl ester

Prepared as described in Example 5a) using 5-bromo-2-chloro-benzoic acid, ethyl ester (0.50 g), 3-pyrrolidinyl-carbamic acid 1,1-dimethylethyl ester (0.42 g), cesium carbonate (0.86 g), palladium (II) acetate (21 mg) and (R)-BINAP (88 mg) and toluene (3 ml) to afford the subtitle compound as an oil (0.25 g).

MS (APCI +ve) 311/313 (M-BOC)⁺

b) (+/-)-2-Chloro-5-[3-[[[(1,1-dimethylethoxy)carbonyl]amino]-1-pyrrolidinyl]-benzoic acid

Prepared as described in Example 5b) using (+/-)-2-chloro-5-[3-[[[(1,1-dimethylethoxy)carbonyl]amino]-1-pyrrolidinyl]-benzoic acid, ethyl ester (Example 7a, 0.25g), lithium hydroxide monohydrate (1.36ml of a 3M solution in water), 1:1 ethanol/water (7 ml) and tetrahydrofuran (1ml) to give the subtitle compound as a solid (0.23 g).

MS (APCI +ve) 284/286 (M-BOC)⁺

c) (+/-)-[1-[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-
3-pyrrolidinyl]-carbamic acid, 1,1-dimethylethyl ester

Prepared as described in Example 5c) using (+/-)-2-chloro-5-[3-[[1,1-dimethylethoxy]carbonyl]amino]-1-pyrrolidinyl]- benzoic acid (Example 7b, 0.070g), N,N'-carbonyldiimidazole (0.033 g), 1-adamantanemethylamine (0.036 ml) and dimethylformamide (3 ml) to give the subtitle compound as a gum.

MS (APCI +ve) 487/489 (M+H)⁺

d) (+/-)-5-(3-Amino-1-pyrrolidinyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt


Prepared as described in example 5d) above using (+/-)-[1-[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-3-pyrrolidinyl]-carbamic acid, 1,1-dimethylethyl ester (Example 7c), hydrochloric acid (0.5ml of a 4N solution in dioxane) and methanol (5 ml). Evaporation under reduced pressure gave a solid on trituration with diethyl ether. Recrystallisation from methanol/diethyl ether gave the title compound as a solid (0.030 g).

MS (APCI +ve) 388/390 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.24 (3H, bs); 8.20 (1H, t); 7.25 (1H, d); 6.61 (1H, dd); 6.51 (1H, d); 3.94 (1H, m); 3.55-3.32 (2H, m); 3.29 (2H, m); 2.92 (2H, d); 2.37-2.27 (1H, m); 2.13-2.05 (1H, m); 1.94 (3H, bs); 1.68 (3H, d); 1.59 (3H, d); 1.52 (6H, s)

Example 8

2-Chloro-5-piperazin-1-ylmethyl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



5 **a) 5-Bromomethyl-2-chloro-benzoic acid**

To a stirred solution of 2-chloro-5-methyl-benzoic acid (25g) in chloroform (500ml) at 50°C was added N-bromosuccinimide (27.40g). The flask was purged with nitrogen and azobisisobutyronitrile (0.10g) added in one portion. The solution was heated at reflux for 1h. Further azobisisobutyronitrile (0.10g) was added and the mixture heated a further 3h.

10 The solution was concentrated in vacuo, redissolved in diethyl ether and filtered to remove insoluble succinimide. The ether solution was washed with 2N aqueous hydrochloric acid solution followed by brine then dried over magnesium sulphate. The solution was concentrated to a volume of 150ml then diluted with isohexane. After further partial concentration crystallization started. The mixture was allowed to stand in an ice-bath for

15 1h. The resulting crystals were filtered, washed with isohexane and dried in vacuo to give the subtitle compound (17g).

b) 5-Bromomethyl-2-chloro-N-(tricyclo[3.3.1.1.3,7]dec-1-ylmethyl)-benzamide

To a stirred solution of 5-bromomethyl-2-chloro-benzoic acid (Example 8a, 12.4g) in dichloromethane (250ml) and dimethylformamide (0.12ml) at 0°C was added oxalyl chloride (8.7ml). The cooling bath was removed and the solution allowed to warm to room temperature. Once gas evolution had ceased the solution was concentrated in vacuo. The residue was redissolved in dichloromethane (300ml), cooled to 0°C and treated with diisopropylethylamine (12.4 ml) and adamantylmethylamine (7.54ml). After 15min. at 0°C

20 the solution was poured into diethyl ether (1L) and washed with 1N aqueous hydrochloric

25

acid followed by brine. The organics were dried over magnesium sulphate and concentrated in vacuo to give the title compound as a white powder (19g)

MS (APCI +ve) 396/398 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.39 (1H, t); 7.50-7.40 (2H, m); 4.74 (2H, s); 2.92 (2H, d); 2.50 (3H, s); 1.94 (3H, bs); 1.67 (3H, d); 1.59 (3H, d); 1.52 (6H, s).

c) 2-Chloro-5-(4-[[1,1-dimethylethyl]oxycarbonyl]-piperazin-1-yl)methyl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

A mixture of 5-bromomethyl-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 8b, 0.130g), 1-*tert*butyloxycarbonylpiperazine (0.074g) and diisopropylethylamine (6.3 ml) in dimethylformamide (3 ml) was heated at 60°C for 3h. The mixture was diluted with water (10 ml) and extracted with ethyl acetate (3 x 10 ml). The organic layer was dried over magnesium sulfate, filtered and the filtrate concentrated under reduced pressure. The crude material was purified on a silica gel eluting with dichloromethane/ethanol (0-20% gradient) to afford the title compound as a white foam (0.112g).

MS (APCI +ve) MW 502/504 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.28 (1H, t); 7.40 (1H, d); 7.32 (1H, dd); 7.29 (1H, d); 3.74 (2H, s); 3.28 (4H, t); 2.90 (2H, d); 2.31 (4H, t); 1.92 (3H, bs); 1.70-1.50 (6H, m); 1.59 (6H, d); 1.37 (9H, s).

d) 2-Chloro-5-piperazin-1-ylmethyl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt

2-Chloro-5-(4-[[1,1-dimethylethyl]oxycarbonyl]-piperazin-1-yl)methyl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, (Example 8c, 0.080g) was dissolved in methanol (3ml), 4N HCl in dioxane (1ml) was added and the mixture stirred at room temperature for 1.5h. The solvent was removed under *vacuum* and the resulting solid was triturated with ether to afford the title compound as a white powder (0.062g).


MS (APCI +ve) MW 402/404 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.30 (1H, t); 7.63 (2H, bs); 7.55 (1H, d); 4.33 (1H, bs);
4.05 (4H, m); 3.50-3.00 (4H, m); 3.50-3.40 (1H, m); 2.92 (2H, d); 1.92 (3H, bs);
1.70-1.50 (6H, m); 1.57 (6H, bs).

According to the procedure described in Example 8, the following compounds were prepared.

Example 9

2-Chloro-5-[(hexahydro-1H-1,4-diazepin-1-yl)methyl] -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




MS (APCI +ve) MW 416/418 (M+H)⁺

¹H NMR (DMSO-d₆) δ 11.62 (bs, 1H), 9.57 (bs, 1H); 9.30 (bs, 1H); 8.34 (1H, t); 7.80-7.60
(2H, m); 7.59 (1H, d); 4.50-4.30 (bs, 2H); 3.80-3.00 (m, 8H); 2.94 (2H, d); 2.25-2.10
(m, 2H); 1.94 (3H, bs); 1.66 (3H, d); 1.58 (3H, d); 1.54 (6H, s).

Example 10

5-[(4-Amino-1-piperidiny)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




MS (APCI +ve) MW 416/418 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.35 (1H, t); 8.30 (2H, bs); 7.66 (1H, d); 7.65 (1H, s); 7.59 (1H, d); 4.28 (d, 2H); 3.65-3.18 (m, 4H); 3.10-2.90 (1H, m); 2.95 (2H, d); 2.15-2.05 (2H, m); 2.05-1.90 (1H, m); 1.94 (3H, bs); 1.68 (3H, d); 1.61 (3H, d); 1.54 (6H, s).


Example 11

5-[(3-Amino-1-pyrrolidinyl)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



MS (APCI +ve) MW 402/404 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.56 (1H, bs); 8.42 (2H, bs); 8.35 (1H, t); 7.66 (2H, bs); 7.59 (1H, d); 4.60-4.40 (m, 2H); 4.20-3.00 (m, 5H); 2.94 (2H, d); 2.35-1.95 (m, 2H); 1.95 (3H, bs); 1.68 (3H, d); 1.61 (3H, d); 1.54 (6H, s).

Example 12**2-Chloro-5-(4-piperidinyloxy)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt****a) 2-Chloro-5-hydroxy-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

To a solution of 2-chloro-5-hydroxybenzoic acid (3.12g) in *N,N*-dimethylformamide (50 ml) was added 1,1'-carbonyldiimidazole (3.0 g). The resulting reaction mixture was stirred for 2.5h and then 1-adamantanemethylamine (3.0g) was added. Stirring was continued for 14h . The reaction mixture was partitioned between ethyl acetate and water and the organic layer was separated, washed with water and brine and then dried over sodium sulphate (Na₂SO₄). The organic layer was concentrated under reduced pressure to give a residue which was purified by silica gel chromatography (eluting with 3-10% methanol in dichloromethane) to yield the subtitle compound as a white solid (0.15 g).

MS (APCI+ve) 319/321 (M+H)⁺

¹H NMR (DMSO-d₆) δ 9.85(1H,s), 8.25 (1H, t), 7.24(1H, d), 6.76-6.82(2H, m), 2.90 (2H,d), 1.93(3H, s), 1.67 (3H, d), 1.57 (3H, d), 1.51 (6H, s)

b) 2-Chloro-5-(4-piperidinyloxy)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt

To a solution of 2-chloro-5-hydroxy-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-methyl)-benzamide (0.20 g, Example 12a), 4-hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (0.19 g) and tributylphosphine (0.23 ml) in dry tetrahydrofuran (6 ml) was added 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g). The orange solution was heated at


60°C under a nitrogen atmosphere for 2h. At this point additional 4-hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (0.19 g), tributylphosphine (0.23 ml) and 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g) were added.. Heating was continued and the process described above repeated until reaction was complete as judged by LC/MS. The cooled reaction mixture was diluted with diethyl ether then filtered. The filtrate was concentrated and purified by normal phase HPLC (0-2% methanol/dichloromethane) followed by chromatography on silica gel (0-2% methanol/dichloromethane) to give the t-butyloxycarbonyl (BOC)-protected compound as a colourless foam. The foam was dissolved in methanol (5 ml) and 4N hydrochloric acid in dioxane (0.25 ml) added. The solution was stirred at room temperature under a nitrogen atmosphere until the reaction was complete as judged by LC/MS. Evaporation of solvent followed by trituration with diethyl ether gave the title compound as a colourless solid (0.15 g).

MS (APCI +ve) 417/419 (M+H)+

¹H NMR (DMSO-d₆) δ 8.65 (2H, bs); 8.30 (1H, t); 7.39 (1H, d); 7.07 (1H, dd); 6.99 (1H, d); 4.72-4.67 (1H, m); 3.21 (2H, bm); 3.07 (2H, bm); 2.92 (2H, d); 2.12-2.07 (2H, m); 1.94 (3H, bs); 1.88-1.80 (2H, m); 1.67 (3H, d); 1.59 (3H, d); 1.52 (6H, s)

Example 13

(R)-2-Chloro-5-(2-pyrrolidinylmethoxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



Prepared as described in Example 12b using 2-chloro-5-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-methyl)-benzamide (0.20 g, Example 12a), N-tBOC-D-prolinol

(0.19 g) and tributylphosphine (0.23 ml), dry tetrahydrofuran (6 ml) and 1-[[1-(piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g) to obtain the butyloxycarbonyl (BOC)-protected compound, followed by treatment with 4N hydrochloric acid in dioxane (0.4 ml) and methanol (5 ml) to yield the title compound as colourless solid (0.14 g)

5


MS (APCI +ve) 403/405 (M+H)+

¹H NMR (CD₃OD) δ 8.45 (1H, bt); 7.46 (1H, d); 7.14-7.10 (2H, m); 4.41 (1H, dd); 4.18 (1H, t); 4.10-4.04 (1H, m); 3.41 (2H, t); 3.10 (2H, m); 2.36-2.28 (1H, m); 2.25-2.08 (2H, d); 2.03 (3H, s); 2.00-1.90 (1H, m); 1.83 (3H, m); 1.74 (3H, d); 1.68 (6H, s)

10

Example 14

(S)-2-Chloro-5-(2-pyrrolidinylmethoxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



15

Prepared as described in Example 12b using 2-chloro-5-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-methyl)-benzamide (0.20 g- Example 12a), N-tBOC-L-prolinol (0.19 g) and tributylphosphine (0.23 ml), dry tetrahydrofuran (6 ml) and 1-[[1-(piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g) to obtain the t-butyloxycarbonyl (BOC)-protected compound, followed by treatment with 4N hydrochloric acid in dioxane (0.5 ml) and methanol (5 ml) to yield the title compound as colourless solid (0.07 g)

20


MS (APCI +ve) 403/405 (M+H)+

¹H NMR (CD₃OD) δ 8.45 (1H, bt); 7.46 (1H, d); 7.14-7.10 (2H, m); 4.41 (1H, dd); 4.18 (1H, t); 4.10-4.04 (1H, m); 3.41 (2H, t); 3.10 (2H, m); 2.36-2.28 (1H, m); 2.25-2.08 (2H, d); 2.03 (3H, s); 2.00-1.90 (1H, m); 1.83 (3H, m); 1.74 (3H, d); 1.68 (6H, s)

25

Example 15

2-Chloro-5-(3-piperidinylmethoxy)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




- 5 Prepared as described in Example 12b using 2-chloro-5-hydroxy-*N*-(tricyclo[3.3.1.1.3,7]dec-1-methyl)-benzamide (0.20 g- Example 12a), 3-piperidinemethanol (0.20 g) and tributylphosphine (0.23 ml), dry tetrahydrofuran (6 ml) and 1-[(1-piperidinylcarbonyl)azo]carbonyl]- piperidine (0.24 g) to obtain the t-butyloxycarbonyl (BOC)-protected compound. This compound was treated with 4N
10 hydrochloric acid in dioxane (0.5 ml) and methanol (5 ml) to yield the title compound as colourless solid (0.09 g).

MS (APCI +ve) 417/19 (M+H)+

- ¹H NMR (DMSO-d₆) δ 8.34 (2H, bs); 8.29 (1H, t); 7.38 (1H, d); 7.01 (1H, dd); 6.93
15 (1H, d); 3.99-3.95 (1H, m); 3.91-3.87 (1H, m); 3.34 (1H, m); 3.23 (1H, bd); 2.92 (2H, d); 2.82-2.71 (2H, m); 2.22 (1H, m); 1.94 (3H, s); 1.82 (2H, d); 1.72-1.66 (4H, m); 1.59 (3H, d); 1.52 (6H, s); 1.39-1.32 (1H, m)

Example 16

- 20 **cis-5-[(4-Aminocyclohexyl)oxy]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt**



Prepared as described in Example 12b using 2-chloro-5-hydroxy-N-(tricyclo[3.3.1.^{3,7}]dec-1-methyl)-benzamide (0.20 g, Example 12a), trans-4-amino-
 5 cyclohexanol (0.20 g) and tributylphosphine (0.23 ml), dry tetrahydrofuran (6 ml) and
 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]- piperidine (0.24 g) to obtain the
 t-butyloxycarbonyl (BOC)-protected compound. This compound was treated with 4N
 hydrochloric acid in dioxane (0.5 ml) and methanol (5 ml) to yield the title compound as
 colourless solid (0.065 g).

10


MS (APCI +ve) 417/19 BP 417

¹H NMR (DMSO-d₆) δ 8.30 (1H, t); 7.97 (3H, bs); 7.38 (1H, d); 7.02 (1H, dd); 6.92
 (1H, d); 4.62 (1H, bs); 3.11 (1H, bs); 2.92 (2H, d); 1.94 (5H, s); 1.76-1.58 (12H, m); 1.52
 (6H, s).

15

Example 17

**2-Methyl-5-(1-piperazinylmethyl)-N-(tricyclo[3.3.1.^{3,7}]dec-1-ylmethyl)-benzamide,
 hydrochloride salt**




To a solution of 2-bromo-5-(4-[[1,1-dimethylethyl]oxycarbonyl]-piperazin-1-yl)methyl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.20g, Example 65b) and tetrakis(triphenylphosphine)palladium(0) (2mg) in dry toluene (6 ml) was added tetramethyltin (0.2ml). The solution was heated at 130°C in a sealed tube for 18hrs. The cooled reaction mixture was evaporated and the residue was treated with 10% KF solution in acetone and stirred for 45min. The mixture was concentrated and chromatographed on silica gel (isohexane then 60% ethyl acetate/ 40% isohexane) to give the t-butyloxycarbonyl (BOC)-protected compound as a colourless oil. The oil was dissolved in methanol (2 ml) and 4N hydrochloric acid in dioxane (1 ml) added. The solution was stirred at room temperature under a nitrogen atmosphere until the reaction was complete as judged by LC/MS. Evaporation of solvent followed by trituration with diethyl ether gave the title compound as a colourless solid (0.03 g).

MS (APCI +ve) 382 (M+H)+

¹H NMR (CD₃OD) δ 7.61 (1H, s); 7.55 (1H, d); 7.39 (1H, d); 4.45 (2H, s); 3.67-3.46 (8H, bm); 3.08 (2H, s); 2.45 (3H, s); 1.99 (3H, s); 1.78 (3H, d); 1.71 (3H, d); 1.62 (6H, s)

Example 18

2-Chloro-5-(1-piperazinylmethyl)-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-benzamide, hydrochloride salt



a) 5-(bromomethyl)-2-chloro-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-benzamide

To a solution of 2-chloro-5-(bromomethyl)-benzoic acid (1.0 g) in dichloromethane (25ml) at 0°C was added dimethylformamide (0.05ml) followed by oxalyl chloride (0.52 ml). The reaction was allowed to warm to room temperature and stirred for 30min. The volatiles were removed under vacuum and the residue dried under high vacuum. The acylchloride was dissolved in dichloromethane (20 ml) and added to a solution of 2-adamantanethylaniline hydrochloride salt (0.95g) in dichloromethane (20ml) and diisopropylethylamine (2 ml) at 0°C. The reaction was allowed to warm to room temperature and stirred for 2h. The organics were washed with water (20ml) then saturated aqueous ammonium chloride solution and the organic layer dried over magnesium sulfate then filtered. The filtrate was concentrated under reduced pressure to a solid. The crude material was recrystallised from dichloromethane/hexane to afford the subtitle compound as a white solid (1.3 g).

b) 4-[[4-Chloro-3-[(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]phenyl]methyl]-1-piperazinecarboxylic acid, 1,1-dimethylethyl ester

A mixture 5-(bromomethyl)-2-chloro-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-benzamide (Example 18a, 0.35g), 1-tertbutyloxycarbonylpiperazine (0.213g), potassium carbonate (0.20g) and potassium iodide (10 mg) in acetone (5 ml) was heated at 60°C for 2h. The acetone was removed under vacuum, the residue taken into dichloromethane and the solid removed by filtration. The crude material was purified on a silica gel eluting with dichloromethane/ethanol (0-10% gradient) to afford the subtitle compound as a white foam (0.383g).

MS (APCI +ve) MW 516/518 (M+H)+

¹H NMR (CDCl₃) δ 7.63 (1H, bs); 7.34 (2H, bs); 6.09 (1H, bs); 3.60-3.30 (8H, m); 2.50-2.30 (4H, bs); 1.97 (3H, bs); 1.72 (3H, d); 1.68 (3H, d); 1.56 (6H, bs); 1.44 (9H, s); 1.50-1.35 (2H, m)

c) **2-Chloro-5-(1-piperazinylmethyl)-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-benzamide, hydrochloride salt**


4-[[4-chloro-3-[[2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl]amino]carbonyl]phenyl]methyl]-1-piperazinecarboxylic acid, 1,1-dimethylethyl ester, (Example 18b, 0.270g) was dissolved in methanol (3ml), 4N HCl in dioxane (2ml) was added and the mixture stirred for 14h at room temperature. The solvent was removed under vacuum and the resulting solid was triturated with ether to afford the title compound as a white powder (0.207g).

MS (APCI +ve) MW 416/418 (M+H)+

¹H NMR (CD₃OD) δ 7.69 (1H, s); 7.66 (1H, d); 7.60 (1H, d); 4.86 (2H, s); 3.70-3.50 (8H, m); 3.50-3.35 (2H, m); 1.98 (3H, bs); 1.78 (3H, d); 1.70 (3H, d); 1.62 (6H, bs); 1.50-1.35 (2H, m).

Example 19

(+/-)-2-Chloro-5-(3-pyrrolidinyloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



Prepared as described in Example 12b using 2-chloro-5-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.155 g, Example 12a), tributylphosphine


(0.23 ml), (+/-)-3-hydroxy-1-pyrrolidinecarboxylic acid, 1,1-dimethylethyl ester (0.19 g), 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g) and dry tetrahydrofuran (10 ml) to give the t-butyloxycarbonyl (BOC)-protected compound as a colourless foam. This compound was treated with 4N hydrochloric acid in dioxane (0.5 ml) and methanol (5 ml) to yield the title compound as a colourless solid (0.075 g).

MS (APCI+ve) 389/391 (M+H)+

¹H NMR (CD₃OD) δ 8.42 (1H, bt); 7.42 (1H, d); 7.09-7.03 (2H, m); 5.23 (1H, bm); 3.59-3.41 (4H, m); 3.07 (2H, d); 2.36-2.30 (2H, m); 1.99 (3H, bs); 1.79 (3H, d); 1.70 (3H, d); 1.63 (6H, d)

Example 20

(+/-)-2-Chloro-5-(3-piperidinyloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




Prepared as described in Example 12b using 2-chloro-5-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.15 g, Example 12a), tributylphosphine (2 x 0.18 ml), 3-hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (2 x 0.14 g), 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]-piperidine (2 x 0.18 g) and dry tetrahydrofuran (6 ml) to give the t-butyloxycarbonyl (BOC)-protected compound as a colourless foam. This compound was treated with 4N hydrochloric acid in dioxane (0.25 ml) and methanol (5 ml) to yield the title compound as a colourless foam (0.042 g).

MS (APCI+ve) 403/405 (M+H)+

¹H NMR (CD₃OD) δ 8.42 (1H, t); 7.41 (1H, d); 7.14-7.10 (2H, m); 4.82 (1H, bm); 3.51-3.39 (1H, m); 3.38 (2H, m); 3.20-3.17 (1H, m); 3.06 (2H, d); 2.10-2.04 (2H, m); 2.00 (3H, bs); 1.94-1.89 (1H, m); 1.84-1.68 (7H, d); 1.64 (6H, d)

Example 21

trans-5-[(4-Aminocyclohexyl)oxy]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




Prepared as described in Example 12b using 2-chloro-5-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.15 g, Example 12a), tributylphosphine (3 x 0.18 ml), cis-(4-hydroxycyclohexyl)-carbamic acid, 1,1-dimethylethyl ester (3 x 0.15 g), 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]-piperidine (3 x 0.18 g) and dry tetrahydrofuran (6 ml) to give the t-butyloxycarbonyl (BOC)-protected compound as a colourless foam. This compound was treated with 4N hydrochloric acid in dioxane (0.5 ml) and methanol (3 ml) to yield the title compound as a colourless foam (0.080 g).

MS (APCI+ve) 417/419 (M+H)+

¹H NMR (CD₃OD) δ 8.38 (1H, t); 7.34 (1H, d); 6.98 (1H, dd); 6.96 (1H, d); 4.30 (1H, m); 3.17 (1H, m); 3.04 (2H, d); 2.22 (2H, bm); 2.09 (2H, m); 1.98 (3H, bs); 1.77 (3H, d); 1.68 (3H, d); 1.62 (6H, s); 1.55 (4H, m)

Example 22

cis-(+/-)-5-[(3-Aminocyclopentyl)oxy]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




Prepared as described in Example 12b using 2-chloro-5-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.20 g, Example 12a), tributylphosphine (0.24 ml), trans-(+/-)-(3-hydroxycyclopentyl)-carbamic acid, 1,1-dimethylethyl ester (0.19 g), 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g) and dry tetrahydrofuran (3 ml) to give the t-butyloxycarbonyl (BOC)-protected compound as a colourless foam. This compound was treated with 4N hydrochloric acid in dioxane (0.5 ml) and methanol (5 ml) to yield the title compound as a colourless foam (0.15 g).

MS (APCI+ve) 403/405 (M+H)+

¹H NMR (CD₃OD) δ 7.36 (1H, d); 7.02-6.98 (2H, m); 4.94-4.90 (1H, m); 3.75-3.68 (1H, m); 3.04 (2H, s); 2.55 (1H, m); 2.24-2.17 (1H, m); 2.09-2.03 (2H, m); 1.98-1.86 (5H, m); 1.76 (3H, d); 1.68 (3H, d); 1.62 (6H, d)

Example 23

(S,S)-2-Chloro-5-(2,5-diazabicyclo[2.2.1]hept-2-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) **5-Bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

Prepared as in Example 1a from 5-bromo-2-chlorobenzoic acid (7.17g), oxalyl chloride (5.3ml), dichloromethane (150ml), dimethylformamide (0.05ml), diisopropylethylamine (6ml) and adamantylmethylamine (5ml) to give the subtitle compound as white colourless needles (7.3g).

MS (APCI-ve) 382/384 (M-H)+

b) **(S,S)-2-Chloro-5-(2,5-diazabicyclo[2.2.1]hept-2-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt**

A mixture of 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (1.70 g, Example 23a), 2,5-diazabicyclo[2.2.1]heptane-2-carboxylic acid, 1,1-dimethylethyl ester (1.06 g), cesium carbonate (2.20 g), (R)-(+)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl ((R)-(+)-BINAP, 0.20 g), and palladium (II) acetate (0.050 g) in dry toluene (10 ml) was heated at 100°C under nitrogen for 24h. The cooled reaction mixture was filtered, washing the residue with ethyl acetate. The filtrate was washed with water and brine, dried (MgSO₄) and evaporated under reduced pressure to give an orange oil.

The oil was purified by chromatography on silica gel, eluting with 0.5% methanol/dichloromethane to yield the *t*-butyloxycarbonyl (BOC) protected compound as a colourless foam. The foam was dissolved in methanol (20 ml) and 4N hydrochloric acid in


dioxane (2.5 ml) added. The solution was stirred at room temperature until reaction was complete (LCMS). The solution was then evaporated under reduced pressure and the residue triturated with diethyl ether to afford the title compound as an off-white solid (0.92 g).

MS (APCI+ve) 400/402 (M-HCl)+

^1H NMR (CD_3OD) δ 8.32 (1H, t); 7.30 (1H, d); 6.77-3.70 (2H, m); 4.69 (1H, s); 4.50 (1H, s); 3.73 (1H, dd); 3.67 (2H, s); 3.06 (2H, d); 2.30 (1H, bd); 2.06 (1H, bd); 1.99 (3H, bs); 1.78 (3H, d); 1.70 (3H, d); 1.64 (6H, s); 1.55 (4H, m) Methanol peak masks other ^1H signal.

Example 24

2-Chloro-5-(2-methyl-1-piperazinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



Prepared as described in Example 23 above from 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.30 g, Example 23a), 3-methyl-1-piperazinecarboxylic acid, 1,1-dimethylethyl ester (0.20 g), cesium carbonate (0.36 g), (R)-(+)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl ((R)-(+)-BINAP, 0.036 g), palladium (II) acetate (0.009 g) and dry toluene (10 ml) to yield the t-butyloxycarbonyl (BOC)-protected compound. This compound was treated with 4N hydrochloric acid in dioxane (0.5 ml) and methanol (5 ml) to yield the title compound as a solid (0.025 g).


MS (APCI+ve) 402/404 (M-HCl)+

¹H NMR (CD₃OD) δ 8.40 (1H, t); 7.37 (1H, d); 7.11 (1H, dd); 7.07 (1H, d); 4.00-3.96 (1H, m); 3.43-3.39 (3H, m); 3.28-3.19 (3H, m); 3.06 (2H, d); 1.98 (3H, bs); 1.77 (3H, d); 1.70 (3H, d); 1.63 (6H, s); 1.10 (3H, d).

5

Example 25

(+/-)-2-Chloro-5-(3-pyrrolidinylamino)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



10

Prepared as described in Example 23 above from 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.30 g, Example 23a), 3-amino-1-pyrrolidinecarboxylic acid, 1,1-dimethylethyl ester (0.18 g), cesium carbonate (0.36 g),

15 (R)-BINAP (0.036 g), anhydrous toluene (3 ml) and palladium (II) acetate (0.009 g); the mixture was heated for 14h in a pressure vessel flushed with nitrogen. Additional (R)-BINAP (0.036 g) and palladium (II) acetate (0.009 g) were added and heating continued for a further 24h. The cooled reaction mixture was poured into water and extracted with ethyl acetate three times. The organic fractions were combined and washed with water then

20 brine, and dried (MgSO₄). Evaporation under reduced pressure gave an oil which was purified by normal phase HPLC (0-5% methanol/dichloromethane) to yield the t-butyloxycarbonyl (BOC)- protected compound as a colourless foam. The foam was dissolved in methanol (5 ml) and 4N hydrochloric acid in dioxane (0.5 ml) added. The solution was stirred at room temperature under a nitrogen atmosphere until the reaction was


complete as judged by LCMS. Evaporation followed by trituration with diethyl ether and methanol yielded the title compound as an off-white solid/foam (0.040 g).

MS (APCI +ve) 388/390 (M-HCl)+

5 ^1H NMR (CD_3OD) δ 8.20 (1H, bt); 7.12 (1H, d); 6.63-6.60 (2H, m); 4.76-4.08 (1H, m); 3.43-3.38 (2H, m); 3.35-3.28 (1H, m); 3.25 (1H, m); 2.94 (2H, s); 2.31-2.22 (1H, m); 2.01-1.94 (1H, m); 1.89 (3H, bs), 1.67 (3H, d); 1.60 (3H, d); 1.53 (6H, s)

Example 26

10 (+/-)-5-(3-Amino-1-piperidiny)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



Prepared as described in Example 23 above from 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.20 g, Example 23a), 3-piperidiny-
15 carbamic acid, 1,1-dimethylethyl ester (0.12 g), cesium carbonate (0.24 g), (R)-(+)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl ((R)-(+)-BINAP, 0.024 g), palladium (II) acetate (0.006 g) and dry toluene (3 ml) to yield the t-butoxycarbonyl (BOC)-protected compound. The t-butoxycarbonyl (BOC)-protected compound was dissolved in methanol (5 ml) and
20 hydrochloric acid (0.5 ml of a 4N solution in dioxane). After stirring at room temperature for 24h the mixture was evaporated and the residue partitioned between ethyl acetate and saturated sodium bicarbonate. The layers were separated and the aqueous phase acidified with 2M hydrochloric acid and extracted with ethyl acetate. The organic phase was dried


over magnesium sulfate and then evaporated under reduced pressure to give a gum. Purification by chromatography on silica gel, eluting with 4-10% methanol in dichloromethane/aqueous ammonia afforded the title compound as a solid (0.036 g).

5 MS (APCI+ve) 402/404 (M+H)+

^1H NMR (CD_3OD) δ 7.25 (1H, d); 7.00 (1H, dd); 6.96 (1H, d); 3.59 (1H, dd); 3.48-3.45 (1H, m); 3.04 (2H, d); 2.91-2.85 (1H, m); 2.82-2.75 (1H, m); 2.58 (1H, dd); 1.98-1.93 (4H, m); 1.85-1.75 (3H, d); 1.70-1.62 (10H, d); 1.34-1.25 (1H, d).

10 **Example 27**

(+/-)-2-Chloro-5-(3-piperidinylamino)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



15 Prepared as described in Example 23 above from 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.30 g, Example 23a), 3-amino-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (0.19 g), cesium carbonate (0.36 g), (R)-(+)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl ((R)-(+)-BINAP, 0.036 g), palladium (II) acetate (0.008 g) and dry toluene (3 ml) to yield the t-butyloxycarbonyl (BOC)-protected
20 compound. This compound was treated with methanol (5 ml) and hydrochloric acid (0.5 ml of a 4 M solution in dioxane) followed by an acid/base work-up. Purification by


chromatography on silica gel, eluting with 4-10% methanol in dichloromethane/aqueous ammonia afforded the title compound as a solid (0.008 g).

MS (APCI+ve) 402/404 (M+H)+

5 ^1H NMR (CD_3OD) δ 7.14 (1H, d); 6.68-6.65 (2H, m); 3.47-3.40 (1H, m); 3.25 (1H, m); 3.05-3.02 (3H, m); 2.72-2.65 (1H, m); 2.52-2.47 (1H, m); 2.08-2.04 (1H, m); 1.97 (3H, bs); 1.89-1.82 (1H, m); 1.77 (3H, d); 1.70-1.62 (10H, d); 1.50-1.40 (1H, m).

Example 28

10 **2-Chloro-5-[hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**



Prepared as described in Example 23 above from 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.15 g, Example 23a), hexahydro-
15 pyrrolo[3,4-c]pyrrole-2(1H)-carboxylic acid, 1,1-dimethylethyl ester (0.17 g), cesium carbonate (0.33 g), (R)-(+)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl ((R)-(+)-BINAP, 0.018 g), palladium (II) acetate (0.004 g) and dry toluene (2 ml) to yield the t-butyloxycarbonyl (BOC)-protected compound. This compound was treated with methanol (5 ml) and hydrochloric acid (0.5 ml of a 4 M solution in dioxane) followed by an acid/base
20 work-up. Trituration of the residue with dichloromethane afforded the title compound as a solid (0.020 g).


MS (APCI+ve) 414/416(M+H)+

¹H NMR (CD₃OD) δ 8.17 (1H, t); 7.20 (1H, d); 6.63 (1H, dd); 6.54 (1H, d); 3.36 (2H, m); 3.02 (2H, dd); 2.95-2.90 (4H, m); 2.80 (2H, m); 2.60 (2H, dd); 1.94 (3H, bs); 1.67 (3H, d); 1.59 (3H, d); 1.52 (6H, d).

5

Example 29

N-[2-methyl-5-(4-piperidinyloxy)phenyl]- tricyclo[3.3.1.1^{3,7}]decane-1-acetamide, hydrochloride salt



10

Prepared as described in Example 12b using N-(5-hydroxy-2-methylphenyl)-tricyclo[3.3.1.1^{3,7}]decane-1-acetamide (0.51 g, Example 12, WO 99/29660), tributylphosphine (0.64 ml), 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]-piperidine (0.65 g), 4-hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (0.52 g) and dry tetrahydrofuran (10 ml) to give the t-butyloxycarbonyl (BOC) protected compound as a colourless solid. This compound was treated with 4N hydrochloric acid in dioxane (0.5 ml) and methanol (5 ml) to yield the title compound as a colourless solid (0.13 g).

15


MS (APCI+ve) 383 (M-HCl)+

¹H NMR (CD₃OD) δ 7.19 (1H, d); 7.12 (1H, d); 6.83 (1H, dd); 4.71-4.66 (1H, m); 3.46-3.40 (2H, m); 3.28-3.22 (2H, m); 2.25 (3H, s); 2.21 (1H, s); 2.21-2.14 (2H, m); 2.11-2.04 (5H, m); 1.84-1.73 (12H, m).

20

Example 30

N-[2-chloro-5-(4-piperidinyloxy)phenyl]- tricyclo[3.3.1.1^{3,7}]decane-1-acetamide, hydrochloride salt




Prepared as described in Example 12b using N-(2-chloro-5-hydroxyphenyl)-tricyclo[3.3.1.1^{3,7}]decane-1-acetamide (0.25 g, Example 28, WO 99/29660), tributylphosphine (0.29 ml), 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]-piperidine (0.30 g), 4-hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (0.24 g) and dry tetrahydrofuran (10 ml) to give the t-butyloxycarbonyl (BOC) protected compound as a colourless solid. This compound was treated with 4N hydrochloric acid in dioxane (1 ml) and methanol (20 ml) to yield the title compound as a colourless solid (0.08 g).

MS (APCI+ve) 375/377 (M-HCl)+

¹H NMR (CD₃OD) δ 7.55 (1H, d); 7.41 (1H, d); 6.88 (1H, dd); 4.76-4.70 (1H, m); 3.48-3.39 (2H, m); 3.30-3.22 (2H, m); 2.25 (2H, s); 2.22-2.16 (2H, m); 2.14-2.03 (2H, m); 1.84-1.72 (12H, m).

Example 31

2-Chloro-5-[(4-piperidinylamino)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt



a) **2-Chloro-5-formyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

5 A solution of 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (3.25 g, Example 23a) in anhydrous tetrahydrofuran (150 ml) was cooled to -78°C under a nitrogen atmosphere. A solution of methyllithium (1.4M in diethyl ether, 6.1 ml) was added over 2min. The mixture was stirred at -78°C for 10min, then a solution tert-butyllithium (1.7M in pentane, 10.0 ml) was added dropwise. The mixture was stirred at -78°C for a further 10min, then dimethylformamide (1.0 ml) was added. The resulting solution was stirred at -78°C for 30min., quenched with saturated aqueous ammonium chloride solution (100 ml) and extracted with ethyl acetate. The combined extracts were dried over anhydrous magnesium sulfate, filtered, and the filtrate concentrated under reduced pressure to give the subtitle compound as a solid (2.76 g).

15

MS (APCI +ve) 332 (M+H)+

^1H NMR (DMSO- d_6) δ 10.04 (1H, s); 8.49 (1H, t); 7.96-7.91 (2H, m); 7.74 (1H, d); 2.96 (2H, d), 1.95 (3H, s); 1.64 (6H, AB); 1.53 (6H, d).

20 b) **4-[4-Chloro-3-[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]methyl]amino]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester**

2-Chloro-5-formyl-N-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (0.270 g, Example 31a) and 3-amino-1-pyrrolidinecarboxylic acid, 1,1-dimethylethyl ester (0.325 g, Journal of Medicinal Chemistry, 1998, 41(22), 4273-4278) were dissolved in 1,2-dichloroethane (30 ml), under a nitrogen atmosphere. Sodium triacetoxyborohydride (0.24 g) was added and the mixture was stirred for 14h at room temperature. Water and

25

dichloromethane were added and the layers were partitioned. The organic extracts were dried over anhydrous magnesium sulfate, filtered and concentrated under reduced pressure. The residue was purified by HPLC eluting with a gradient of 0-10% of ethanol in dichloromethane, then by chromatography over silica gel eluting with ethyl acetate : iso-hexane (1:1) then ethyl acetate : ethanol (98:2) to give the subtitle compound as a colourless oil (0.158 g).

MS (APCI +ve) 516 (M+H)+

c) **2-Chloro-5-[(4-piperidinylamino)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt**

Prepared from 4-[[[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]methyl]amino]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (0.158 g,

Example 31b) methanol (3 ml) and 4N hydrochloric acid solution in dioxane (2 ml).


Solvents were removed under reduced pressure and the residue was triturated with ethyl acetate, iso-hexane and diethyl ether to give the title compound as a white solid (0.126 g).

MS (APCI +ve) 416 (M+H-2HCl)+

¹H NMR (CD₃OD) δ 8.47 (1H, t); 7.62-7.56 (3H, m), 4.33 (2H, s); 3.58-3.55 (3H, m); 3.12 (2H, t); 3.07 (2H, d); 2.44 (2H, d); 2.03-1.92 (5H, m); 1.73 (6H, q); 1.63 (6H, d).

Example 32

5-[[[4-(Aminomethyl)cyclohexyl]amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt



a) **[[4-[[[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]methyl]amino]cyclohexyl]methyl]-carbamic acid, 1,1-dimethylethyl ester**

Prepared according to the method described in Example 31b from 2-chloro-5-formyl-N-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (0.30 g, Example 31a),
 5 [(4-aminocyclohexyl)methyl]-carbamic acid, 1,1-dimethylethyl ester (0.207 g, WO 97/32882), sodium triacetoxyborohydride (0.135 g) and 1,2-dichloroethane (10 ml).
 The residue was purified by chromatography over silica gel eluting with ethyl acetate : iso-hexane (1:1) then ethyl acetate : ethanol (9:1) to give the subtitle compound as a colourless
 10 oil (0.26 g).

MS (APCI +ve) 544 (M+H)+

b) **5-[[[4-(Aminomethyl)cyclohexyl]amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt**


Prepared from [[4-[[[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]methyl]amino]cyclohexyl]methyl]-carbamic acid, 1,1-dimethylethyl ester (0.26 g, Example 32a) methanol (5 ml) and 4N hydrochloric acid
 solution in dioxane (2 ml). Solvents were removed under reduced pressure and the residue
 20 was triturated with diethyl ether to give the title compound as a white powder (0.191 g).

MS (APCI +ve) 444 (M+H-2HCl)+

¹H NMR (CD₃OD) δ 7.60-7.58 (3H, m), 4.29 (2H, s); 3.28-3.12 (1H, m); 3.09 (2H, s);
 2.84 (2H, d); 2.31 (2H, bd); 2.00 (5H, bs); 1.75 (6H, q); 1.65 (6H, d); 1.71-1.65 (1H, m);
 25 1.63-1.44 (2H, m); 1.31-1.12 (2H, m).

Example 33

5-[[[4-(Aminocyclohexyl)amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt



a) 4-[[[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]methyl]amino]cyclohexyl]-carbamic acid, 1,1-dimethylethyl ester

Prepared according to the method described in Example 31b from 2-chloro-5-formyl-N-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (0.30 g, Example 31a), (4-aminocyclohexyl)-carbamic acid, 1,1-dimethylethyl ester (0.194 g, Journal of Organic Chemistry, 1996, 61(25), 8811-8818), sodium triacetoxyborohydride (0.135 g) and 1,2-dichloroethane (10 ml). The residue was purified by chromatography over silica gel eluting with ethyl acetate : iso-hexane (1:1) then ethyl acetate : ethanol (95:5) to give the subtitle compound as a colourless oil (0.24 g).

MS (APCI +ve) 530 (M+H)+

b) 5-[[[(4-Aminocyclohexyl)amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt


Prepared from 4-[[[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl] methyl]amino]cyclohexyl]-carbamic acid, 1,1-dimethylethyl ester (0.26 g, Example 33a), methanol (5 ml) and 4N hydrochloric acid solution in dioxane (1 ml). Solvents were removed under reduced pressure and the residue was triturated with diethyl ether to give the title compound as a white powder (0.190 g).

MS (APCI +ve) 430 (M+H-2HCl)+

¹H NMR (CD₃OD) δ 7.61-7.59 (3H, m), 4.30 (2H, s); 3.28-3.11 (2H, m); 3.08 (2H, s); 2.40-2.32 (2H, m); 2.21-2.17 (2H, m); 2.00 (3H, s); 1.74 (6H, q); 1.64 (6H, d); 1.63-1.48 (4H, m).

Example 34

5-[(1-Azabicyclo[2.2.2]oct-3-ylamino)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




Prepared according to the method described in Example 31b from 2-chloro-5-formyl-N-(tricyclo [3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.30 g, Example 31a), 1-azabicyclo[2.2.2]octan-3-amine dihydrochloride salt (0.18 g), sodium triacetoxymethylborohydride (0.135 g) and 1,2-dichloroethane (10 ml). The residue was purified by chromatography over silica gel eluting with ethyl acetate : iso-hexane (1:1) followed by ethyl acetate : ethanol (95:5). Repurification by chromatography over silica gel eluting with dichloromethane : methanol (95:5) then (9:1) gave the title compound as a white gum (0.013 g).

MS (APCI +ve) 442 (M+H)+

¹H NMR (CDCl₃) δ 7.68 (1H, d); 7.39 (1H, d); 7.31 (1H, dd); 6.41 (1H, t); 3.75 (2H, s); 3.42-3.31 (2H, m); 3.25-3.09 (6H, m); 2.94 (1H, d); 2.38-2.23 (2H, m); 2.22-2.14 (1H, m); 2.01 (3H, s); 1.92-1.83 (2H, m); 1.69 (6H, q); 1.59 (6H, d).

Example 35

N-[4-(3-Aminopyrrolidin-1-yl)-2-methylphenyl]-2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetamide, dihydrochloride salt



a) [1-(3-Methyl-4-nitrophenyl)-pyrrolidin-3-yl]-carbamate derivative

4-Fluoro-2-methyl-1-nitrobenzene (1 g), pyrrolidin-3-ylcarbamate derivative (1.2 g), potassium carbonate (1.79 g) and dimethyl sulfoxide (10 ml) were heated together at 80°C under nitrogen for 15h. The mixture was then cooled, diluted with ethyl acetate (200 ml), washed with 2N aqueous hydrochloric acid (200 ml), dried (MgSO₄) then concentrated. Purification of the residue by silica gel chromatography (eluting with 20% ethyl acetate in isohexane) gave the subtitle compound (1.744 g).

¹H NMR (DMSO-d₆) δ 8.03 - 8.00 (1H, d), 7.28 - 7.21 (1H, br d), 6.51 - 6.47 (2H, m), 4.20 - 4.12 (1H, br m), 3.61 - 3.16 (4H, m), 2.56 (3H, s), 2.20 - 2.08 (1H, m), 1.98 - 1.85 (1H, m), 1.39 (9H, s).

b) [1-(4-Amino-3-methylphenyl)-pyrrolidin-3-yl]-carbamate derivative

[1-(3-Methyl-4-nitrophenyl)-pyrrolidin-3-yl]-carbamate derivative (1.744 g, Example 35a), iron powder (1.52 g), ammonium chloride (1.45 g), ethanol (50 ml) and water (50 ml) were refluxed together under nitrogen for 2h. The mixture was cooled and the iron was filtered off. Water (200 ml) was added to the residue and the product extracted into ethyl acetate (3 x 200 ml), dried (MgSO₄), and concentrated to give the subtitle compound (1.56 g).

¹H NMR (CDCl₃) 6.65 (1H, br s), 6.38 (2H, br m), 4.80 (1H, m), 4.33 (2H, br m), 3.60 - 2.80 (5H, m), 2.31 - 2.17 (4H, m), 1.92 - 1.82 (1H, m), 1.45 (9H, br s).

c) {1-[4-(2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetylamino)-3-methylphenyl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester

To a solution of adamantan-1-yl-acetic acid (0.46g) in dichloromethane (10ml) at 0°C was added dimethylformamide (0.1ml) followed by oxalyl chloride (2.50 ml). The reaction was allowed to warm to room temperature and stirred for 30min.. The volatiles were removed under vacuum and the residue dried under high vacuum. The residue was dissolved in dichloromethane (10ml) and added to a solution of [1-(4-amino-3-methylphenyl)-pyrrolidin-3-yl]-carbamic acid tert-butyl ester (0.70g, Example 35b) in dichloromethane (10ml) and triethylamine (0.8 ml) at 0°C. The reaction was allowed to warm to room temperature and stirred for 3h. The solution was washed with 2N aqueous hydrochloric acid (20ml), then brine (20 ml) and the organic layer dried over magnesium sulfate then filtered. The filtrate was concentrated under reduced pressure. The crude material was purified by silica gel chromatography (eluting with 1% methanol in dichloromethane) to yield the subtitle compound(1.1g).

MS (APCI +ve) MW 468 (M+H)+

¹H NMR (DMSO-d₆) δ 8.86 (1H, s); 7.01 - 6.98 (1H, d); 7.18 - 7.14 (1H, br d); 6.33 - 6.27 (3H, m); 4.15 - 4.04 (1H, m); 3.42 - 3.15 (3H, m); 3.00 - 2.97 (1H, m); 2.12 (3H, s); 2.00 (2H, s); 1.99 - 1.80 (5H, m); 1.70 - 1.61 (12H, m); 1.39 (9H, s).

d) N-[4-(3-Aminopyrrolidin-1-yl)-2-methylphenyl]-2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetamide, dihydrochloride salt

{1-[4-(2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetylamino)-3-methylphenyl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester (0.20 g, Example 35c) was dissolved in methanol (5 ml) and hydrochloric acid (0.5ml of a 4N solution in dioxane) was added. After stirring at room temperature for 14h, the mixture was evaporated to 2/3 original volume under reduced pressure. Diethyl ether was gradually added to the solution and the resulting precipitate collected by filtration, washed with diethyl ether and dried in vacuo to afford the title compound as a solid (0.15 g)


MS (APCI +ve) 368 (M+H)+

¹H NMR (DMSO-d₆) δ 8.92 (1H, s); 8.21 (2H, br s); 7.07 - 7.04 (1H, d); 6.41 - 6.35 (2H, m); 3.91 (1H, br m); 3.50 - 3.39 (2H, m); 3.29 - 3.20 (2H, m); 2.37 - 2.27 (2H, m); 2.14 (3H, s); 2.02 (2H, s); 1.94 (3H, s); 1.70 - 1.58 (12H, m).

5

Example 36

N-(2-Methyl-4-piperazin-1-ylphenyl)-2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetamide, dihydrochloride salt



- 10 a) **4-(3-Methyl-4-nitrophenyl)piperazine-1-carboxylic acid tert-butyl ester**
4-Fluoro-2-methyl-1-nitrobenzene (2 g), piperazine-1-carboxylic acid tert-butyl ester (4.8 g), potassium carbonate (3.57 g) and dimethyl sulfoxide (20 ml) were heated together at 80 °C under nitrogen for 15h. The mixture was then cooled, diluted with ethyl acetate (200 ml), washed with 2N aqueous hydrochloric acid (200 ml), dried (MgSO₄), and
15 concentrated to give the subtitle compound (4.10g).

MS (APCI+ve) 321 (M)+

¹H NMR (DMSO-d₆) δ 8.02 - 7.98 (1H, d), 6.89 - 6.86 (2H, m), 3.45 (8H, s), 2.55 (3H, s), 1.42 (9H, s).

20

- b) **4-(4-Amino-3-methylphenyl)piperazine-1-carboxylic acid tert-butyl ester**
4-(3-Methyl-4-nitrophenyl)piperazine-1-carboxylic acid tert-butyl ester (2 g, Example 36a), iron powder (1.74 g), ammonium chloride (1.67 g), ethanol (50 ml) and water (50 ml) were refluxed together under nitrogen for 2h. The mixture was cooled and
25 the iron was filtered off. Water (200 ml) was added to the residue and the product

extracted into ethyl acetate (3 x 200 ml), dried (MgSO₄), and concentrated to give the subtitle compound (1.22 g).

¹H NMR (DMSO-d₆) δ 6.62 - 6.52 (3H, m), 4.38 (2H, s), 3.41 (4H, br s), 2.83 (4H, br s),
5 2.02 (3H, s), 1.41 (9H, s).

c) 4-[4-(2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetylamino)-3-methylphenyl]-piperazine-1-carboxylic acid tert-butyl ester

To a solution of adamantan-1-yl-acetic acid (0.40g) in dichloromethane (10ml) at
10 0°C was added dimethylformamide (0.1ml) followed by oxalyl chloride (2.00 ml). The reaction was allowed to warm to room temperature and stirred for 30min.. The volatiles were removed under vacuum and the residue dried under high vacuum. The residue was dissolved in dichloromethane (10ml) and added to a solution of 4-(4-amino-3-methylphenyl)piperazine-1-carboxylic acid tert-butyl ester (0.60g, Example 36b) in
15 dichloromethane (10ml) and triethylamine (0.7 ml) at 0°C. The reaction was allowed to warm to room temperature and stirred for 3h. The solution was washed with 2N aqueous hydrochloric acid (20ml), then brine (20 ml) and the organic layer dried over magnesium sulfate then filtered. The filtrate was concentrated under reduced pressure. The crude material was purified by silica gel chromatography (eluting with 1% methanol in
20 dichloromethane) to yield the subtitle compound(0.42g).

MS (APCI +ve) MW 468 (M+H)+

¹H NMR (DMSO-d₆) δ 8.96 (1H, s); 7.14 - 7.11 (1H, d); 6.79 - 6.72 (2H, m); 3.47 - 3.40
(4H, m); 3.20 - 3.00 (4H, m); 2.14 (3H, s); 2.03 (2H, s); 1.94 (3H, br s); 1.70 - 1.56
25 (12H, m); 1.42 (9H, s).

d) N-(2-Methyl-4-piperazin-1-ylphenyl)-2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetamide, dihydrochloride salt

4-[4-(2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetylamino)-3-methylphenyl]-piperazine-1-carboxylic acid tert-butyl ester (0.05 g, Example 36c) was dissolved in methanol (2 ml) and
30


hydrochloric acid (0.5ml of a 4N solution in dioxane) was added. After stirring at room temperature for 14h, the mixture was evaporated to 2/3 original volume under reduced pressure. Diethyl ether was gradually added to the solution and the resulting precipitate collected by filtration, washed with diethyl ether and dried in vacuo to afford the title compound as a solid (0.043 g)

MS (APCI +ve) 368 (M+H)+

¹H NMR (DMSO-d₆) δ 9.01 (3H, br s); 7.18 - 7.15 (1H, d); 6.84 - 6.82 (1H, d); 6.79 - 6.76 (1H, dd); 3.31 - 3.29 (4H, m); 3.28 - 3.16 (4H, m); 2.16 (3H, s); 2.04 (2H, s); 1.94 (3H, br s); 1.69 - 1.58 (12H, m).

Example 37

cis-4-(3-Amino-cyclopentyloxy)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) 2-Chloro-4-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

To a solution of 2-chloro-4-hydroxybenzoic acid (3.30 g) in dimethylformamide (20 ml) was added 1,1'-carbonyldiimidazole (3.30 g). The reaction mixture was stirred for 2.5h and then 1-adamantanemethylamine (3.4 ml) was added. After 14h the reaction mixture was partitioned between ethyl acetate and 2N aqueous hydrochloric acid and the organic layer was separated, washed with water then brine and dried (MgSO₄). The organic layer was concentrated under reduced pressure to give a residue which was purified by silica gel chromatography (eluting with 10 - 70% ethyl acetate in dichloromethane) to

yield a white solid which was triturated with ethyl acetate to yield the subtitle compound as a white solid (3.6 g).

MS (APCI +ve) 320/322 (M+H)+

¹H NMR (DMSO-d₆) δ 10.12 (1H, s), 8.10 - 8.06 (1H, t), 7.27 - 7.24 (1H, d), 6.81 (1H, d), 6.77 - 6.73 (1H, dd), 2.91 - 2.88 (2H, d), 1.93 (3H, br s), 1.69 - 1.56 (6H, br q), 1.50 (6H, br s).

b) **cis-4-(3-Amino-cyclopentyloxy)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt**


To a solution of 2-chloro-4-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.20 g, Example 37a), *trans*-(3-hydroxycyclopentyl)-carbamic acid, *tert*-butyl ester (0.19 g) and tributylphosphine (0.23 ml) in dry tetrahydrofuran (6 ml) was added 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g). The orange solution was heated at 60°C under a nitrogen atmosphere for 2h. Additional *trans*-(3-hydroxycyclopentyl)-carbamic acid, *tert*-butyl ester (0.19 g), tributylphosphine (0.23 ml) and 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g) were added. Heating was continued and the process described above repeated until reaction was judged complete as judged by LC/MS. The cooled reaction mixture was diluted with diethyl ether then filtered. The filtrate was concentrated and purified by chromatography on silica gel (25 - 33% ethyl acetate/ hexane) to give the t-butyloxycarbonyl (BOC)-protected compound as a colourless foam. The foam was dissolved in methanol (5 ml) and 4N hydrochloric acid in dioxane (0.25 ml) added. The solution was stirred at room temperature under a nitrogen atmosphere until the reaction was complete as judged by LC/MS. Evaporation of solvent followed by trituration with diethyl ether gave the title compound as a colourless solid (0.24 g).

MS (APCI +ve) 403/405 (M+H)+

^1H NMR (DMSO- d_6) δ 8.18 (1H, t); 7.96 (2H, br s); 7.37 - 7.34 (1H, d); 7.05 (1H, m); 6.97 - 6.94 (1H, m); 4.87 (1H, br m); 3.72 - 3.40 (2H, m); 2.93 - 2.90 (2H, d); 2.04 - 1.51 (19H, m); 1.22 (2H, m).

5 **Example 38**

2-Chloro-4-(4-piperidinyloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




To a solution of 2-chloro-4-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-methyl)-benzamide
10 (0.20 g, Example 37a), 4-hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester
(0.19 g) and tributylphosphine (0.25 ml) in dry tetrahydrofuran (6 ml) was added 1-[(1-
piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g). The orange solution was heated at
50°C under a nitrogen atmosphere for 2h. Additional 4-hydroxy-1-piperidinecarboxylic
acid, 1,1-dimethylethyl ester (0.19 g), tributylphosphine (0.25 ml) and 1-[(1-
15 piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g) were added. Heating was continued
and the process described above repeated until reaction was complete as judged by LC/MS.
The cooled reaction mixture was diluted with diethyl ether then filtered. The filtrate was
concentrated and purified by chromatography on silica gel (3:1 iso-hexane/ethyl acetate) to
give the the t-butyloxycarbonyl (BOC)-protected compound as a colourless foam. The
20 foam was dissolved in methanol (10 ml) and 4N hydrochloric acid in dioxane (10 ml)
added. The solution was stirred at room temperature under a nitrogen atmosphere until the
reaction was complete as judged by LC/MS. Evaporation of solvent followed by trituration
with diethyl ether gave the title compound as a colourless solid (0.165 g).

MS (APCI +ve) 403 (M+H)+

¹H NMR (DMSO-d₆) δ 8.80 (2H, bs); 8.21-8.16 (1H, t); 7.37-7.34 (1H, d); 7.16 (1H, m);
7.03-6.99 (1H, m); 4.80-4.68 (1H, m); 3.25-3.18 (2H, m); 3.17-3.01 (2H, m); 2.93-2.90
(2H, d); 2.17-2.02 (2H, m); 1.93 (3H, bs); 1.87-1.73 (2H, m); 1.69-1.57 (6H, AB); 1.51
(6H, s)

Example 39

(+/-)-2-Chloro-4-(pyrrolidin-3-yloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide




Prepared as described in Example 38 from 2-chloro-4-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-methyl)-benzamide (0.20 g, Example 37a), (+/-)-3-hydroxy-1-pyrrolidinecarboxylic acid, 1,1-dimethylethyl ester (0.18 g), tributylphosphine (0.25 ml), dry tetrahydrofuran (6 ml) and 1-[[[(1-piperidinylcarbonyl)azo]carbonyl]-piperidine
(0.24 g) to obtain the the t-butyloxycarbonyl (BOC)-protected compound. This compound
was treated with 4N hydrochloric acid in dioxane (10 ml) and methanol (10 ml) to yield the
title compound as colourless solid (0.165 g).

MS (APCI +ve) 389 (M+H)+

¹H NMR (DMSO-d₆) δ 8.19-8.15 (1H, t); 7.35-7.32 (1H, d); 6.99 (1H, m); 6.93-6.90
(1H, m); 4.94-4.89 (1H, m); 3.24 (1H, s); 3.08-3.02 (1H, dd); 2.92-2.90 (2H, d); 2.88-2.72
(3H, m); 2.08-1.98 (1H, m); 1.93 (3H, s); 1.76-1.57 (7H, m); 1.51 (6H, s).

Example 40

2-Chloro-4-(piperidin-3-yloxy)- N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



5

To a solution of 2-chloro-4-hydroxy-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.20 g, Example 37a), 3-hydroxy-piperidine-1-carboxylic acid, *tert*-butyl ester (0.189 g) and tributylphosphine (0.23 ml) in dry tetrahydrofuran (6 ml) was added 1-[[1-(piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g). The orange solution was heated at 60°C under a nitrogen atmosphere for 2h. Additional 3-hydroxy-piperidine-1-carboxylic acid, *tert*-butyl ester (0.19 g), tributylphosphine (0.23 ml) and 1-[[1-(piperidinylcarbonyl)azo]carbonyl]-piperidine (0.24 g) were added. Heating was continued and the process described above repeated until reaction was complete as judged by LC/MS. The cooled reaction mixture was diluted with diethyl ether then filtered. The filtrate was concentrated and purified by chromatography on silica gel (25% ethyl acetate : *iso*-hexane) followed by normal phase HPLC (0 – 1% ethanol in dichloromethane) to give the *t*-butyloxycarbonyl (BOC)-protected compound as a colourless foam. The foam was dissolved in methanol (5 ml) and 4N hydrochloric acid in dioxane (0.25 ml) added. The solution was stirred at room temperature under a nitrogen atmosphere until the reaction was complete as judged by LC/MS. Evaporation of solvent followed by trituration with diethyl ether gave the title compound as a colourless solid (0.006 g).

10

15


20

MS (APCI +ve) 403/405 (M+H)+

¹H NMR (DMSO-d₆) δ 8.84 (2H, br s), 8.21 (1H, t); 7.38 (1H, d); 7.18 (1H, s); 7.05 (1H, dd); 4.82 (1H, br s); 3.24 (1H, d); 3.20 (1H, dd); 3.06 (2H, br s); 2.92 (2H, d); 1.94 – 1.51 (19H, m).

Example 41

2-Chloro-4-(4-piperazin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) 4-Bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

To a suspension of 4-bromo-2-chlorobenzoic acid (5.00 g) in dichloromethane (25 ml) at 0°C was added oxalyl chloride (3.7 ml) and dimethylformamide (5 drops). The resulting mixture was stirred at room temperature under a nitrogen atmosphere for 1 hour, then concentrated under reduced pressure to yield a solid. The solid was dissolved in dichloromethane (20 ml) and added dropwise to a solution of 1-adamantanemethylamine (3.36g) and N,N-diisopropylethylamine (5.55 ml) in dichloromethane (20 ml). The resulting solution was allowed to stir at room temperature under a nitrogen atmosphere for 20h. The reaction mixture was diluted with dichloromethane and washed with water, 10% aqueous potassium carbonate, 10% aqueous potassium hydrogen sulfate and saturated brine. The organic phase was then dried over sodium sulfate, filtered and concentrated under reduced pressure to afford the subtitle compound as a solid (4.28 g).

MS (APCI +ve) 382/384 (M+H)+

¹H NMR (DMSO-d₆) δ 8.39-8.34 (1H, t); 7.78 (1H, m); 7.62-7.59 (1H, m); 7.37-7.34 (1H, d), 2.94-2.92 (2H, d); 1.94 (3H, br s); 1.69-1.57 (6H, br AB); 1.52 (6H, s).

5 **b) 2-Chloro-4-(4-piperazin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt**


To a suspension of the 4-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 41a, 0.30 g), piperazine-1-carboxylic acid, 1,1-dimethylethyl ester (0.18 g), cesium carbonate (0.36 g) and (R)-BINAP (0.036 g) in anhydrous toluene (3 ml) was added palladium (II) acetate (0.009 g) and the mixture heated at 100 °C for 14h in a pressure vessel flushed with nitrogen. The cooled reaction mixture was evaporated under reduced pressure to give an oil which was purified by chromatography on silica gel (2:1 / *iso*-hexane : ethyl acetate) to give the t-butyloxycarbonyl (BOC)- protected compound as a colourless foam. The foam was dissolved in methanol (15 ml) and 4N hydrochloric acid in dioxane (15 ml) added. The solution was stirred at room temperature under a nitrogen atmosphere until the reaction was complete as judged by LCMS. Evaporation followed by trituration with diethyl ether and methanol yielded the title compound as an off-white solid/foam (0.161 g).

20 MS (APCI +ve) 388/390 (M+H)+

¹H NMR (DMSO-d₆) δ 8.98 (2H, bs); 8.11-8.07 (1H, t); 7.33-7.31 (1H, d); 7.05 (1H, m); 6.99-6.95 (1H, m); 3.46-3.43 (4H, m); 3.20 (4H, bs); 1.94 (3H, bs); 1.69-1.57 (6H, b AB); 1.51 (6H, bs).

25 **Example 42**

2-Chloro-4-(3-pyrrolidinylamino)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




Prepared according to the method described in Example 41b from 4-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.25 g, Example 41a), 3-amino-1-pyrrolidinecarboxylic acid, 1,1-dimethylethyl ester (0.182 g, Journal of Medicinal Chemistry, 1998, 41 (22), 4273-4278), cesium carbonate (0.347 g), (R)-(+)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (0.036 g), palladium (II) acetate (0.009 g) and anhydrous toluene (3 ml). The residue was purified by HPLC eluting with a gradient of 0-5% ethanol in dichloromethane. The product was dissolved in methanol and stirred at room temperature for 3h in the presence of 4N hydrochloric acid solution in dioxane (2 ml). The solution was concentrated under reduced pressure and triturated with diethyl ether to give the title compound as a white powder (0.057 g).

MS (APCI +ve) 388 (M+H-HCl)+

¹H NMR (CD₃OD) δ 7.33 (1H, d); 6.71 (1H, d); 6.63 (1H, dd); 4.27-4.21 (1H, m); 3.57-3.40 (3H, m); 3.23 (1H, dd); 3.05 (2H, s); 2.43-2.33 (1H, m); 2.33-2.01 (1H, m); 1.99 (3H, bs); 1.73 (6H, q); 1.62 (6H, d).

Example 43

2-Chloro-4-(hexahydro-1H-1,4-diazepin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




Prepared according to the method described in Example 41b from 4-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.25 g, Example 41a), hexahydro-1H-1,4-diazepine-1-carboxylic acid, 1,1-dimethylethyl ester (0.182 g), cesium carbonate (0.347 g), (R)-(+)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (0.036 g), palladium (II) acetate (0.009 g) and anhydrous toluene (3 ml). The residue was purified by HPLC eluting with a gradient of 0-5% ethanol in dichloromethane. The product was dissolved in methanol and stirred at room temperature for 3h in presence of 4N hydrochloric acid solution in dioxane (2 ml). The solution was concentrated under reduced pressure and triturated with diethyl ether to give the title compound as a white powder (0.17 g).

MS (APCI +ve) 402 (M+H-HCl)+
¹H NMR (CD₃OD) δ 7.41 (1H, d); 6.88 (1H, d); 6.81 (1H, dd); 3.83 (2H, t); 3.63 (2H, t); 3.40 (2H, t); 3.30 (2H, t); 3.06 (2H, s); 2.24-2.16 (2H, m); 1.99 (3H, bs); 1.74 (6H, q); 1.63 (6H, d).

According to the procedure described in Example 8, the following compounds were prepared:

Example 44

(±)-5-[(3-Amino-1-piperidinyl)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




MS (APCI +ve) MW 416/418 (M+H)+

¹H NMR (CD₃OD) δ 7.70 (1H, bs); 7.67 (1H, dd); 7.60 (1H, d); 4.49 (1H, d); 4.45 (1H, d);
3.73-3.58 (2H, m); 3.57-3.45 (1H, m); 3.14-2.95 (4H, m); 2.25-2.04 (2H, m); 1.98
5 (4H, bs); 1.76 (3H, d); 1.73-1.58 (1H, m); 1.70 (3H, d); 1.63 (6H, bs).

Example 45

2-Chloro-5-(2,5-diazabicyclo[2.2.1]hept-2-ylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide, hydrochloride salt




MS (APCI +ve) MW 414/416 (M+H)+

¹H NMR (CD₃OD) δ 7.74 (1H, d); 7.72 (1H, dd); 7.60 (1H, d); 4.70-4.55 (3H, m); 4.45
(1H, d); 4.00 (1H, d); 3.73 (1H, d); 3.60-3.50 (2H, m); 3.07 (2H, s); 2.71 (1H, d); 2.27
15 (1H, d); 1.98 (3H, bs); 1.77 (3H, d); 1.69 (3H, d); 1.63 (6H, bs).

Example 46

2-Chloro-5-(9-oxa-3,7-diazabicyclo[3.3.1]non-3-ylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide, hydrochloride salt




MS (APCI +ve) MW 442/446 (M+H)+

¹H NMR (CD₃OD) δ 7.60-7.40 (3H, m); 4.25-4.00 (2H, m); 3.70-3.40 (2H, m); 3.46 (4H, m); 3.07 (2H, s); 3.15-2.90 (2H, m); 2.80-2.50 (2H, m); 2.00 (3H, bs); 1.78 (3H, d);
 5 1.71 (3H, d); 1.63 (6H, bs).

Example 47

2-Chloro-5-(3,7-diazabicyclo[3.3.1]non-3-ylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt

10




MS (APCI +ve) MW 442/444 (M+H)+

¹H NMR (CD₃OD) δ 7.91 (1H, s); 7.78 (1H, d); 7.56 (1H, d); 7.46 (1H, bs); 4.44 (2H, bs);
 3.65-3.28 (8H, m); 3.08 (2H, bs); 2.48 (2H, bs); 2.05-1.90 (5H, m); 1.77 (3H, d); 1.71
 15 (3H, d); 1.64 (6H, bs).

Example 48

trans-2-Chloro-5-[[8-(methylamino)-3-azabicyclo[3.2.1]oct-3-yl]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt

20



MS (APCI +ve) MW 456/458 (M+H)+


¹H NMR (CD₃OD) δ 7.79 (1H, d); 7.77 (1H, dd); 7.58 (1H, d); 4.71 (2H, bs); 3.80 (2H, d);

3.40 (1H, t); 3.25 (2H, dd); 3.07 (2H, s); 2.86 (3H, s); 2.70 (2H, bs); 2.10-1.90 (7H, m);

1.77 (3H, d); 1.70 (3H, d); 1.63 (6H, bs).

Example 49

cis-2-Chloro-5-[(hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



MS (APCI +ve) MW 428/430 (M+H)+


¹H NMR (DMSO-d₆) δ 8.00 (1H, t); 7.65 (1H, s); 7.63 (1H, d); 7.52 (1H, d); 4.34

(2H, bs); 3.60-3.05 (10H, m); 2.97 (2H, d); 1.95 (3H, bs); 1.70 (3H, d); 1.63 (3H, d);

1.57 (6H, s).

Example 50

2-Chloro-5-(4-piperidinyldenemethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) **[[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-methyl] phosphonic acid, dimethyl ester**

5 **5-Bromomethyl-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (5.70 g, Example 8b) in 100ml of trimethylphosphite was heated at reflux for 15h. The solvent was removed by azeotropic distillation with toluene under high vacuum to afford the subtitle compound as a yellow solid.**

10 **MS (APCI +ve) MW 426/428 (M+H)+**

¹H NMR (DMSO-d₆) δ 8.34 (1H, t); 7.42 (1H, d); 7.35-7.27 (2H, m); 3.63 (3H, s); 3.58 (3H, s); 3.42 (2H, d); 2.92 (2H, d); 1.94 (3H, bs); 1.67 (3H, d); 1.59 (3H, d); 1.52 (6H, bs).

b) **4-[[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]methylene]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester**

15 **To a solution of the crude [[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl] methyl] phosphonic acid, dimethyl ester (2.50 g, Example 50a) in tetrahydrofuran (50 ml) at -78 °C was added a solution of lithium diisopropylamide (7.30 ml, 2M in tetrahydrofuran). The reaction was allowed to warm to room temperature and stirred for 15min. N-t-butoxycarbonylpiperidin-4-one (1.52g) in tetrahydrofuran (5 ml) was then added and the mixture stirred for 24h. The reaction was diluted with water and extracted with ethylacetate. The organic layer was washed with brine and dried over magnesium sulfate. The crude material was purified on a silica gel (0 to 5% methanol in dichloromethane) to afford the subtitle compound as a white foam.**

25 **MS (APCI +ve) MW 443/445 (M+H)+**

¹H NMR (DMSO-d₆) δ 7.52 (1H, d); 7.34 (1H, d); 7.16 (1H, dd); 6.30 (1H, s); 6.25 (t, 1H); 3.48 (2H, t); 3.40 (2H, t); 3.40 (2H, t); 3.18 (2H, d); 2.42 (t, 2H); 2.32 (t, 2H); 2.05 (3H, bs); 1.73 (3H, d); 1.64 (d, 3H), 1.59 (6H, s); 1.47 (9H, bs).

5 c) **2-Chloro-5-(4-piperidinylidenemethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt**


A solution of 4-[[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl] methylene]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (0.10g, Example 50b) in methanol (3ml) was treated with a 4N solution
10 of hydrochloric acid in dioxane (1 ml) and stirred for 14h at room temperature. The reaction mixture was concentrated under vacuum and the residue recrystallised from isopropanol/ether to give the title compound as a white solid (0.071g).

MS (APCI +ve) MW 399/401 (M+H)+

15 ¹H NMR (DMSO-d₆) δ 7.52 (1H, d); 7.34 (1H, d); 7.16 (1H, dd); 6.30 (1H, s); 6.25 (t, 1H); 3.48 (2H, t); 3.40 (2H, t); 3.40 (2H, t); 3.18 (2H, d); 2.42 (t, 2H); 2.32 (t, 2H); 2.05 (3H, bs); 1.73 (3H, d); 1.64 (d, 3H), 1.59 (6H, s); 1.47 (9H, bs).

Example 51

20 **2-Chloro-5-(4-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt**



To a solution of 2-chloro-5-(4-piperidinylidenemethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt (0.10g, Example 50c) in ethanol (10 ml) was
25 added platinum oxide (2mg). The vessel was placed under 3 bars hydrogen pressure for


3h. The catalyst was removed by filtration through a pad of Celite, washed with ethanol and the solution concentrated under vacuum. The crude material was recrystallised from iso-propanol to afford a white solid. The t-butoxycarbonyl protected compound was dissolved in methanol (10ml) and treated with a solution of 4N HCl in dioxane (2 ml). The reaction was stirred for 14h at room temperature, the volatiles removed under vacuum and the residue recrystallised from iso-propanol/ether to afford the hydrochloride salt as a white powder (0.065g).

MS (APCI +ve) MW 402/404 (M+H)+

¹H NMR (CD₃OD) δ 8.38 (1H, t); 7.38 (1H, d); 7.30-7.20 (2H, m); 3.35 (2H, d); 3.05 (2H, d); 2.92 (2H, td); 2.63 (2H, d); 1.98 (3H, bs); 1.95-1.80 (1H, m); 1.85 (2H, d); 1.77 (3H, d); 1.68 (3H, d); 1.62 (6H, s); 1.40 (2H, q).

Example 52

2-Chloro-5-(4-hydroxy-piperidin-4-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



To a solution of 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.30g, Example 23a) in anhydrous tetrahydrofuran (10ml) at -78 °C was added dropwise a solution of n-butyllithium in hexanes (2.5M, 0.72ml). After 10min. a solution of t-butoxycarbonyl-4-piperidone (0.21g) in tetrahydrofuran (2ml) was added. The solution was stirred an additional 20min. then treated with saturated aqueous ammonium chloride solution. The mixture was allowed to warm to room temperature then partitioned between ethyl acetate and saturated aqueous ammonium chloride solution. The organic extracts were dried over magnesium sulphate and concentrated in vacuo. The residue was


chromatographed on silica (ethyl acetate: isohexane/1:4 to 1:2 gradient) to give the t-butoxycarbonyl protected product (0.153g). This was redissolved in methanol (4ml) and treated for 14h with 4N HCl in dioxan (1ml). The solution was partially concentrated in vacuo and the product precipitated with diethyl ether. The solution was filtered and the white solid washed with diethyl ether to give the title compound (0.091g)

MS (APCI +ve) MW 403 (M+H)+

¹H NMR (CD₃OD) δ 8.43 (1H, m, br); 7.59 (1H, m); 7.55 (1H, d); 7.49 (1H, d); 3.52-3.30 (4H, m); 3.09 (2H, d); 2.23 (2H, m); 2.04-1.88 (5H, m); 1.95-1.80 (1H, m); 1.84-1.66 (6H, m); 1.65 (6H, d).

Example 53

2-Chloro-5-(1,2,3,6-tetrahydro-pyridin-4-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




A solution of 2-chloro-5-(4-hydroxy-piperidin-4-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt (0.25g, Example 52) in concentrated hydrochloric acid (10ml) was heated at 100°C for 5h. The solution was allowed to cool slowly. Colourless crystals separated. These were removed by filtration, washed with diethyl ether then acetonitrile and dried to afford the title compound (0.031g).

MS (APCI +ve) MW 385 (M+H)+

^1H NMR (CD_3OD) δ 8.44 (1H, m, br); 7.55-7.45 (3H, m); 6.23 (1H, m); 3.85 (2H, m); 3.47 (2H, t); 3.07 (2H, d); 2.79 (2H, m); 1.98 (3H, m); 1.77 (3H, m); 1.69 (3H, m); 1.63 (6H, s, br).

5 **Example 54**

2-Ethyl-5-piperazin-1-ylmethyl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




2-Bromo-5-(4-[[1,1-dimethylethyl]oxycarbonyl]-piperazin-1-yl)methyl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 65b, 0.89g) was dissolved in dry tetrahydrofuran. Sodium hydride (60% dispersion, 0.07g) was added and the mixture stirred at room temperature for 5min. The mixture was cooled to -70°C under a nitrogen atmosphere and t-butyllithium (1.9ml, 1.7M solution) added. After 5min, ethyl iodide (0.5ml) was added and the mixture stirred at -70°C for 30min. Aqueous ammonium chloride solution was added and the product extracted with diethyl ether, dried (MgSO_4) and concentrated in vacuo. Chromatography on silica gave the t-butyloxycarbonyl (BOC) protected compound as a foam. This was redissolved in methanol (5ml) and 4N HCl in dioxane (1ml) added. The mixture was stirred at room temperature for 14h. The solution was partially concentrated under vacuum and the product precipitated with diethyl ether.

20 The resulting solid was filtered and washed with ether to afford the title compound as a white powder (0.040g).

¹H NMR (DMSO) δ 9.59 (2H, s, br); 8.15 (1H, t); 7.58 (2H, s, br); 7.35 (1H, d); 4.37 (2H, s, br); 3.49 (m); 3.25 (2H, m, br); 2.95 (2H, d); 2.72 (2H, q); 1.94 (3H, s, br); 1.69-1.59 (6H, m); 1.52 (6H, s); 1.165 (3H, t).

5 **Example 55**

2-Chloro-5-(piperidin-4-ylsulfanyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) **4-(Toluene-4-sulfonylsulfanyl)-piperidine-1-carboxylic acid tert-butyl ester**

10 4-Iodo-piperidine-1-carboxylic acid, *tert*-butyl ester (1.3g) and potassium toluene-4-thiosulfonate (1.0g) were combined in ethanol (10ml) with *cis*-dicyclohexane 18-crown-6 (10mg) and heated under reflux for 12h. After cooling the reaction mixture was partitioned between ethyl acetate and water, the organic layer was separated, washed with water and brine, dried over magnesium sulfate and concentrated under reduced pressure. The residue
15 was purified by silica gel chromatography (eluting with *iso*-hexane / ethyl acetate 4:1 to 7:3), to yield the subtitle compound as an oil (0.65g).

MS (APCI +ve) 315 (M+H-tBu)+

¹H NMR (CDCl₃) δ 7.80-7.85 (2H, m), 7.30-7.40 (2H, m), 3.95-4.10 (1H, m), 3.75-3.85 (1H, m), 3.40-3.50 (1/2H, m), 3.10-3.20 (1/2H, m), 2.95-3.05 (1H, m), 2.80-2.90 (1H, m),
20 2.46 (3H, s), 1.90-2.10 (2H, m), 1.50-1.70 (2H, m), 1.43 & 1.45 (9H, pair s).

b) 2-Chloro-5-(4-[(1,1-dimethylethyl)oxycarbonyl]piperidine-4-ylsulfanyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

To a solution of 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.30g, Example 23a) in anhydrous tetrahydrofuran (10ml) at -78 °C was added dropwise a solution of n-butyllithium in hexanes (2.5M, 0.72ml). After 10min. a solution of 4-(toluene-4-sulfonylsulfanyl)-piperidine-1-carboxylic acid, *tert*-butyl ester (0.38g, Example 55a) in tetrahydrofuran (7ml) was added. After a further 1 hour at -78°C the reaction mixture was warmed to ambient temperature and quenched by the addition of water (5ml). The reaction mixture was diluted with ethyl acetate and washed twice with saturated aqueous sodium hydrogen carbonate solution, then with brine and dried over magnesium sulfate. The organic layer was concentrated under reduced pressure to give a residue which was purified by silica gel chromatography (eluting with 0-5% ethanol in dichloromethane) to yield the subtitle compound (0.20g).

MS (APCI +ve) 419/21 (M+H-BOC)+

c) 2-Chloro-5-(piperidin-4-ylsulfanyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt


2-Chloro-5-(4-[(1,1-dimethylethyl)oxycarbonyl]-piperidin-4-ylsulfanyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.20g, Example 55b) was dissolved in methanol (15 ml) and hydrochloric acid (1.0ml of a 4N solution in dioxane) was added. After stirring at room temperature for 14h, the reaction mixture was basified with saturated sodium hydrogen carbonate solution and extracted twice with dichloromethane. The organic layer was concentrated under reduced pressure to give a residue which was purified by silica gel chromatography (eluting with 0-100% methanol in dichloromethane). The residue was dissolved in dichloromethane (5ml) and hydrochloric acid (1N in diethyl ether, 2ml) added. Evaporation to dryness gave the title compound as the hydrochloride salt (0.050g).

MS (APCI +ve) 419/21 (M+H)+

^1H NMR (DMSO- d_6) δ 8.75 (2H, brd), 8.38 (1H, t), 7.48 (2H, s), 7.37 (1H, s), 3.50-3.60 (1H, m), 3.30 (2H, brd), 2.92-3.05 (4H, m), 2.06 (2H, brd), 1.94 (3H, s), 1.57-1.75 (8H, m), 1.52 (6H, s).

5 Example 56

2-Chloro-5-(piperidin-4-ylsulfinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




m-Chloroperoxybenzoic acid (166mg) was added to a solution of 2-chloro-5-(piperidin-4-ylsulfinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.35g, Example 55c) in dichloromethane (5ml). After 2h calcium hydroxide (0.20g) was added and 30min. later the salts removed by filtration. The filtrate was concentrated under reduced pressure to give a residue which was purified by silica gel chromatography (eluting with 0-25% ethanol in dichloromethane). The residue was dissolved in methanol (5 ml) and hydrochloric acid (0.5ml of a 4N solution in dioxane) was added. After stirring at room temperature for 14h, the solution was concentrated under reduced pressure and triturated with diethyl ether to yield the title compound as the hydrochloride salt (0.030g).

MS (APCI +ve) 435/37 (M+H)+

^1H NMR (DMSO- d_6) δ 8.88 (1H, brs), 8.47 (2H, brt), 7.76 (1H, d), 7.67 (1H, dd), 7.61 (1H, d), 3.30-3.40 (2H, m), 3.13 (1H, t), 2.98 (2H, d), 2.80-2.90 (2H, m), 2.15 (1H, d), 1.95 (3H, m), 1.50-1.85 (15H, m).

Example 57

2-Chloro-5-(piperidin-4-ylsulfonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




m-Chloroperoxybenzoic acid (0.30g) was added to a solution of 2-chloro-5-(piperidin-4-ylsulfonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.30g) in dichloromethane (10ml). After 2h calcium hydroxide (170mg) was added and 30min. later the salts removed by filtration. The filtrate was concentrated under reduced pressure to give a residue which was purified by silica gel chromatography (eluting with 0-2% ethanol in dichloromethane). The residue was dissolved in methanol (5 ml) and hydrochloric acid (0.25ml of a 4N solution in dioxane) was added. After stirring at room temperature for 14h, the solution was concentrated under reduced pressure and triturated with diethyl ether to yield the title compound (0.03g).

MS (APCI +ve) 451/53 (M+H)+

¹H NMR (DMSO-d₆) δ 8.89 (1H, brs), 8.57 (1H, t), 8.50 (1H, brs), 7.87 (2H, ABq), 7.75 (1H, d), 3.71 (1H, td), 3.40 (2H, d), 3.00 (2H, d), 2.80-2.90 (2H, m), 1.95-2.05 (5H, m), 1.50-1.90 (14H, m).

Example 58

2-Chloro-5-(piperidin-4-ylmethylsulfonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) 2-Chloro-5-(4-[(1,1-dimethylethyl)oxycarbonyl] piperidin-4-ylmethylsulfanyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Trifluoroacetic acid anhydride (2ml) was added to a solution of 2-chloro-5-methylsulphanyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.53g, Example 58a, WO 99/29661) in dichloromethane (10ml) and heated under reflux for 1.5h; cooled and concentrated. The resultant residue was dissolved in methanol (30ml), allowed to stand for 1 hour then concentrated. The resultant residue was dissolved in acetone (10ml) and potassium carbonate (0.60g) and 4-iodomethyl-piperidine-1-carboxylic acid tert-butyl ester (0.94g) was added. The reaction mixture was heated under reflux for 3h, cooled and concentrated. The resultant residue was dissolved in ethyl acetate, washed twice with 10% w/w KHSO₄ solution, twice with saturated sodium hydrogen carbonate solution, once with brine and dried over magnesium sulfate. The organic layer was concentrated under reduced pressure to give a residue which was purified by silica gel chromatography (eluting with 0-2% ethanol in dichloromethane), to yield the subtitle compound (0.46g).

MS (APCI +ve) 433/35 (M+H-BOC)+

¹H NMR (CDCl₃) δ 7.61 (1H, d), 7.25-7.31 (2H, m), 6.27 (1H, brt), 4.09 (2H, brd), 3.17 (2H, d), 2.86 (2H, d), 2.66 (2H, t), 2.01 (3H, s), 1.60-1.90 (15H, m), 1.45 (9H, s), 1.18 (2H, dq).

b) 2-Chloro-5-(piperidin-4-ylmethylsulfanyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt

Hydrochloric acid (4N dioxane, 0.5ml) was added to a solution of 2-chloro-5-(4-
[1,1-dimethylethyl]oxycarbonyl] piperidin-4-ylmethylsulfanyl)-N-
(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.235g, Example 58a) in methanol (10ml).
After 24h the reaction mixture was concentrated, then triturated with ether to give the title
5 compound (0.20g).


MS (APCI +ve) 433/35 (M+H)+

¹H NMR (DMSO-d₆) δ 8.76 (1H, brs), 8.48 (1H, brs), 8.35 (1H, t), 7.40 (2H, ABq), 7.28
(1H, d), 3.24 (2H, d), 3.00 (2H, d), 2.92 (2H, d), 2.84 (2H, q), 1.94 (5H, brs), 1.75-1.82
10 (1H, m), 1.65 (6H, q), 1.52 (6H, s), 1.40 (2H, q).

Example 59

2-Chloro-5-(piperidin-4-ylmethanesulfonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt

15



m-Chloroperoxybenzoic acid (0.19g) was added to a solution 2-chloro-5-(4-[[1,1-
dimethylethyl]oxycarbonyl] piperidin-4-ylmethylsulfanyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
ylmethyl)-benzamide (0.165g, Example 58a) in chloroform (10ml). After 5h calcium
20 hydroxide (120mg) was added and 30min. later the salts removed by filtration. The
reaction mixture was concentrated, then dissolved in methanol (10 ml) and hydrochloric
acid (1.0ml of a 4N solution in dioxane) added. After stirring at room temperature for 14h,
concentration under reduced pressure yielded the title compound (0.075g).


MS (APCI +ve) 465/67 (M+H)+

¹H NMR (DMSO-d₆) δ 8.70 (1H, brs), 8.55 (1H, t), 8.48 (1H, brs), 7.95 (1H, dd), 7.88 (1H, d), 7.82 (1H, d), 3.47 (2H, d), 3.21 (2H, d), 2.97 (2H, d), 2.89 (2H, d), 2.10-2.25 (1H, m), 1.95 (5H, brs), 1.40-1.80 (14H, m).

5

Example 60

2-Chloro-5-(piperazine-1-carbonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



10 **a) 4-Chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-isophthalamic acid**

n-Butyllithium (3ml, 2M hexanes) was added at -78°C to a solution of 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide (1.0g, Example 23a) in tetrahydrofuran (20ml). After 10min. the reaction mixture was decanted onto dry solid carbon dioxide and allowed to warm to ambient temperature. The reaction mixture was
15 acidified with concentrated hydrochloric acid and extracted with ether. The organics were separated, dried over magnesium sulfate and concentrated under reduced pressure to give a residue which was purified by silica gel chromatography (eluting with iso-hexane / ethyl acetate 3:1 to 1:1 + 1%AcOH), to yield the subtitle compound as a solid (0.45g).

20 MS (APCI +ve) 348 / 350 (M+H)+

¹H NMR (DMSO-d₆) δ 13.33 (1H, s), 8.44 (1H, t), 7.94 (1H, dd), 7.87 (1H, d), 7.63 (1H, d), 2.95 (2H, d), 1.95 (3H, s), 1.63 (6H, q), 1.53 (6H, s).

b) **2-Chloro-5-(piperazine-1-carbonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt**

Ethyl-diisopropylamine (0.3ml) was added at ambient temperature to a solution of 4-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-isophthalamide (0.15g, Example 60a),
5 piperazine-1-carboxylic acid tert-butyl ester (0.16g) and PyBrOP (0.40g) in N-methylpyrrolidinone (10ml). After 5h the reaction mixture was diluted with ethyl acetate and washed twice with water, twice with 10% KHSO₄ solution, twice with saturated NaHCO₃ solution and once with brine, then dried over magnesium sulfate and concentrated under reduced pressure. The residue was purified by silica gel
10 chromatography (eluting with ethanol (0-5%) in dichloromethane), then redissolved in methanol (10ml) and treated with hydrochloric acid (4N dioxane, 1ml). After 48h the reaction mixture was concentrated under reduced pressure and recrystallised from iso-hexane/propan-2-ol to yield the title compound as a solid (0.10g).

15 MS (APCI +ve) 416 / 418 (M+H)⁺


¹H NMR (DMSO-d₆) δ 9.18 (1H, t), 8.42 (1H, t), 7.59 (1H, d), 7.47-7.52 (2H, m), 3.50-3.90 (4H, brs), 3.05-3.25 (4H, brs), 2.94 (2H, d), 1.95 (3H, s), 1.63 (6H, q), 1.52 (6H, s).

The following Examples were made in an analogous manner.

20

Example 61

2-Chloro-5-([1,4]diazepane-1-carbonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




MS (APCI +ve) 430 / 432 (M+H)+

¹H NMR (DMSO-d₆) δ 9.10 (2H, brs), 8.06 (1H, brs), 7.53 (1H, d), 7.45-7.48 (2H, m), 3.78 (2H, brs), 3.54 (2H, brs), 3.20-3.25 (4H, m), 2.97 (2H, d), 2.00 (2H, m), 1.95 (3H, s), 1.65 (6H, q), 1.55 (6H, s).

Example 62

4-Chloro-N-1-(piperidin-4-yl)-N2-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-isophthalamide, hydrochloride salt




MS (APCI +ve) 430 / 432 (M+H)+

¹H NMR (DMSO-d₆) δ 8.66-8.76 (3H, m), 8.42 (1H, t), 7.89-7.93 (2H, m), 7.60 (1H, d), 4.01-4.09 (1H, m), 3.29 (2H, d), 2.95-3.05 (4H, m), 1.95 (5H, brs), 1.47-1.82 (14H, m).

Example 63

2-Chloro-5-(hydroxy-4-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



5

a) 2-Chloro-5-[4-[[[1,1-dimethylethyl]oxycarbonyl] piperidinyl]-hydroxymethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

To a solution of 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (1.5 g, Example 23a) in anhydrous tetrahydrofuran (50 ml) under a nitrogen atmosphere at -78°C was added dropwise n-butyllithium solution (2.5M in hexanes, 3.4 ml). The mixture was stirred for 10min. at -78°C, then a solution of 4-formyl-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (1.09 g, Journal of Medicinal Chemistry, 1999, 42(12), 2180-2190) in anhydrous tetrahydrofuran (10 ml) was added dropwise. The reaction mixture was stirred at -78°C for 30min. then quenched with saturated aqueous ammonium chloride solution (100 ml). The product was extracted twice with ethyl acetate (2x100 ml). The organic extracts were combined, dried over anhydrous magnesium sulfate, filtered and concentrated under reduced pressure. The residue was purified by chromatography on silica gel eluting with iso-hexane : ethyl acetate / (2:1) then (1:2), then purified further by HPLC eluting with a gradient of 0-5% ethanol in dichloromethane to give the subtitle compound as a white foam (0.61 g).

20

MS (APCI +ve) 517 (M+H)+

¹H NMR (DMSO-d₆) δ 8.28 (1H, t); 7.40 (1H, d); 7.33-7.27 (2H, m); 5.33 (1H, d); 4.34 (1H, t); 3.93 (3H, bs); 2.93 (2H, d); 2.61 (2H, bs); 1.94 (3H, bs); 1.63 (6H, q); 1.53 (6H, d); 1.37 (9H, s); 1.34-1.23 (2H, m); 1.09 (2H, dt).

25

b) **2-Chloro-5-(hydroxy-4-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt**


A solution of 2-chloro-5-[4-[[[1,1-dimethylethyl]oxycarbonyl]-piperidinyl]-hydroxymethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.10 g, Example 63a) in methanol (3 ml) was treated with 4N hydrochloric acid solution in dioxane (1 ml). After 14h the solvents were removed under reduced pressure and the residue was triturated with diethyl ether to give the title compound as a white powder (0.062 g).

MS (APCI +ve) 417 (M+H-HCl)+

¹H NMR (DMSO-d₆) δ 8.70 (1H, bs); 8.29 (2H, bt); 7.44 (1H, d); 7.33 (2H, dt), 5.53 (1H, d); 4.40 (1H, t); 3.23 (2H, bs); 2.93 (2H, d); 2.76 (2H, bd); 1.94 (3H, bs); 1.77-1.36 (1H, m); 1.53 (6H, s).

Example 64


(±)-2-Chloro-5-(hydroxy-3-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



Prepared by an analogous route to Example 63 employing 3-formyl-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester.

MS (APCI +ve) MW 417/419 (M+H)+

¹H NMR (CD₃OD) δ 8.41 (1H, t); 7.46 (1H, d); 7.41 (1H, d); 7.39 (1H, s); 4.65 (0.5H, d); 4.50 (0.5H, d); 3.74 (0.5H, td); 3.66 (1H, q); 3.57 (0.5H, t); 3.44 (0.5H, bd); 3.18 (0.5H, bd); 3.06 (2H, d); 2.86 (2H, qd); 2.10-1.87 (m, 2H); 1.98 (3H, bs); 1.77 (3H, d); 1.68 (3H, bd); 1.63 (6H, s), 1.60-1.50 (m, 1H); 1.50-1.36 (m, 1H).

Example 65**2-Bromo-5-(1-methyl-1-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)methyl)piperazin-5-ylmethyl)-benzamide, hydrochloride salt****a) 2-Bromo-5-bromomethyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

To a solution of 2-bromo-5-bromomethyl-benzoic acid (6.1g) in dichloromethane (100 ml) at 0°C were added dimethylformamide (0.2 ml) followed by oxalylchloride (3 ml). The reaction was stirred at room temperature for 0.5 hour and concentrated under vacuum. The acylchloride was redissolved in dichloromethane (100 ml) and di-isopropylethylamine (6 ml) followed by adamantanemethylamine (3.5ml) added at 0°C. The mixture was stirred at 0°C for 10min., then partitioned between diethyl ether and 1N aqueous hydrochloric acid. The organic layer was dried over magnesium sulfate and concentrated in vacuo. The crude material was purified by recrystallisation from dichloromethane/ethylacetate/i-hexane to afford the title compound as a white solid (6.5 g) (sample contained some of the corresponding benzyl chloride).

b) 2-Bromo-5-(4-[(1,1-dimethylethyl)oxycarbonyl]piperazin-1-yl)methyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

A mixture of 2-bromo-5-bromomethyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 65a, 5.0g), 1-tertbutyloxycarbonylpiperazine (2.3g), potassium carbonate (3.2g), potassium iodide (0.30g) and acetone (75ml) was refluxed in the dark for 14h. The mixture was concentrated in vacuo, partitioned between ethyl acetate and water, then washed with brine. The organic layer was dried over magnesium sulfate and

concentrated in vacuo. Recrystallization from ethyl acetate:isohexane gave the subtitle compound as colourless solid (4.7g).

MS (APCI +ve) MW 546 (M+H)+

5

c) 2-Bromo-5-piperazin-1-ylmethyl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt


2-Bromo-5-(4-[(1,1-dimethylethyl)oxycarbonyl]-piperazin-1-yl)methyl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 65b, 0.40g) was dissolved in
10 methanol (15ml), and 4N HCl in dioxane (3ml) added. The mixture was stirred at room temperature for 14h. The solvent was removed under vacuum and the resulting solid was triturated with ether to afford the title compound as a white powder (0.23g).

MS (APCI +ve) MW 447(M+H)+

15 ¹H NMR (DMSO-d₆) δ 9.53 (1H, s, br); 8.31 (1H, t); 7.72 (1H, d); 7.60 (1H, m), 4.33 (1H, m); 3.50-3.00 (4H, m); 3.50-3.40 (1H, m); 2.94 (2H, d); 1.94 (3H, bs); 1.71-1.58 (6H, m); 1.54 (6H, bs).

20 **Example 66**

2-Chloro-5-[2-(1-piperazinyl)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) 2-Chloro-5-(2-hydroxyethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

A solution of 5-bromo-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (3.0 g, Example 23a) in anhydrous tetrahydrofuran (100 ml) was cooled to -78°C under a nitrogen atmosphere. A solution of methyllithium (1.4M in diethyl ether, 4.9 ml) was added over 2min. The mixture was stirred at -78°C for 10min., then a solution *tert*-butyllithium (1.7M in pentane, 9.3 ml) was added dropwise. The mixture was stirred at -78°C for a further 10min., then ethylene oxide (1.0 ml) was added. The resulting solution was stirred at -78°C for 30min. then was warmed to 0°C and stirred for another 6h. The reaction mixture was quenched with saturated aqueous ammonium chloride solution (70 ml) and extracted with ethyl acetate (3x100 ml). The combined extracts were dried over anhydrous magnesium sulfate, filtered, and the filtrate concentrated under reduced pressure. The residue was purified by chromatography over silica gel eluting with dichloromethane : ethanol (98:2) to give the subtitle compound as a pale yellow solid (0.89 g).

MS (APCI +ve) 348 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.27 (1H, t), 7.36 (1H, d); 7.28-7.23 (2H, m); 4.65 (1H, t); 3.60 (2H, q); 2.92 (2H, d); 2.73 (2H, t); 1.94 (3H, bs); 1.63 (6H, q); 1.52 (6H, d).

b) 2-Chloro-5-(2-oxoethyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

To a solution of 2-chloro-5-(2-hydroxyethyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (1.07 g, Example 66a) in anhydrous dichloromethane (20 ml) was added Dess-Martin periodinane reagent (1.95 g) and the mixture was stirred at room temperature for 1h. Sodium thiosulfate (3.43 g) was dissolved in aqueous sodium bicarbonate solution (28 ml) and added to the reaction mixture. Diethyl ether (50 ml) was then added and the mixture was stirred for 10min.. The layers were partitioned and the organic layer was washed with water then brine. The organic extracts were then dried over anhydrous magnesium sulfate, filtered and concentrated under reduced pressure to afford the subtitle compound as a white solid (1.06 g).

MS (APCI +ve) 346 (M+H)⁺

¹H NMR (DMSO-d₆) δ 9.69 (1H, s); 8.32 (1H, t); 7.45 (1H, d); 7.30-7.24 (2H, m); 3.84 (2H, s); 2.92 (2H, d); 1.94 (3H, bs); 1.63 (6H, q); 1.52 (6H, d).

c) **2-Chloro-5-[2-(1-piperazinyl)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide hydrochloride salt**


To a solution of 2-chloro-5-(2-oxoethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.101 g, Example 66b) in anhydrous 1,2-dichloroethane (5 ml) was added 1-piperazinecarboxylic acid, 1,1-dimethylethyl ester (0.108 g) then sodium triacetoxyborohydride (0.086 g). The reaction mixture was stirred for 14h at room temperature. Water (10 ml) and dichloromethane (10 ml) were added and the layers were partitioned. The organic extract was dried over anhydrous magnesium sulfate, filtered and concentrated under reduced pressure. The residue was purified by HPLC eluting with a gradient of 0-5% ethanol in dichloromethane then by chromatography over silica gel eluting with ethyl acetate. The white powder obtained was dissolved in methanol (5 ml) and a solution of hydrochloric acid in dioxane (4N, 1 ml) was added. The mixture was stirred for 14h at room temperature. Solvents were then removed under reduced pressure and the product obtained was triturated with diethyl ether to afford the title compound as a white powder (0.047 g).

MS (APCI +ve) 416 (M+H)⁺

¹H NMR (CD₃OD) δ 8.42 (1H, t); 7.46 (1H, d); 7.41-7.38 (2H, m); 3.63-3.49 (8H, m); 3.48-3.45 (2H, m); 3.19-3.14 (2H, m); 3.06 (2H, s); 1.99 (3H, bs); 1.73 (6H, q); 1.63 (6H, d).

Example 67

2-Chloro-5-[2-(2,5-diazabicyclo[2.2.1]hept-2-yl)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




Prepared according to the method described in Example 66c from 2-chloro-5-(2-oxoethyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.094 g, Example 66b), 2,5-diazabicyclo[2.2.1]heptane-2-carboxylic acid, 1,1-dimethylethyl ester (0.108 g), sodium triacetoxyborohydride (0.081 g) and 1,2-dichloroethane (2 ml). After work-up, the residue was purified by HPLC eluting with a gradient of 0-5% ethanol in dichloromethane. The white powder obtained was dissolved in methanol (2 ml) and a solution of hydrochloric acid in dioxane (4N, 1ml) was added. The mixture was stirred for 14h at room temperature. Solvents were then removed under reduced pressure and the product obtained was triturated with diethyl ether to afford the title compound as a white powder (0.067 g).

MS (APCI +ve) 428 (M+H)⁺

¹H NMR (CD₃OD) δ 7.49-7.40 (3H, m); 4.64 (2H, d); 3.92 (2H, d); 3.77-3.48 (4H, m); 3.18 (2H, t); 3.08 (2H, s); 2.61 (1H, bd); 2.28 (1H, bd); 2.00 (3H, bs); 1.75 (6H, q); 1.64 (6H, d).

Example 68

5-[2-(4-Amino-1-piperidinyl)ethyl]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




Prepared according to the method described in Example 66c from 2-chloro-5-(2-oxoethyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.094 g, Example 66b), 4-piperidinyl-carbamic acid, 1,1-dimethylethyl ester (0.109 g), sodium triacetoxyborohydride (0.081 g) and 1,2-dichloroethane (2 ml). After work-up, the residue was purified by HPLC eluting with a gradient of 0-5% ethanol in dichloromethane. The white powder obtained was dissolved in methanol (2 ml) and a solution of hydrochloric acid in dioxane (4N, 1ml) was added. The mixture was stirred for 14h at room temperature. Solvents were then removed under reduced pressure and the product obtained was triturated with diethyl ether to afford the title compound as a white powder (0.065 g).

MS (APCI +ve) 430 (M+H)⁺

¹H NMR (CD₃OD) δ 8.43 (1H, t); 7.49-7.38 (3H, m); 3.78 (2H, bd); 3.64-3.42 (2H, m); 3.41-3.35 (2H, m); 3.23-3.14 (3H, m); 3.08 (2H, s); 2.33-2.29 (2H, m); 2.13-2.04 (2H, m); 2.00 (3H, bs); 1.75 (6H, q); 1.64 (6H, d).

Example 69

2-Chloro-5-[2-(3-piperidinylamino)ethyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt



Prepared according to the method described in Example 66c from 2-chloro-5-(2-oxoethyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.094 g, Example 66b), 3-amino-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (0.109 g), sodium

triacetoxyborohydride (0.081 g) and 1,2-dichloroethane (2 ml). After work-up, the residue was purified by HPLC eluting with a gradient of 0-5% ethanol in dichloromethane. The white powder obtained was dissolved in methanol (2 ml) and a solution of hydrochloric acid in dioxane (4N, 1ml) was added. The mixture was stirred for 14h at room temperature.

5 Solvents were then removed under reduced pressure and the product obtained was triturated with diethyl ether to afford the title compound as a white powder (0.065 g).

MS (APCI +ve) 430 (M+H)⁺


¹H NMR (CD₃OD) δ 7.49-7.46 (1H, m); 7.42-7.38 (2H, m); 3.76 (1H, bd); 3.64-3.54 (1H, m); 3.44-3.34 (3H, m); 3.19-2.98 (4H, m); 3.08 (2H, s); 2.34 (1H, bd); 2.18-2.12 (1H, m); 2.00 (3H, bs); 1.89-1.78 (2H, m); 1.74 (6H, q); 1.64 (6H, d).

10

Example 70

5-[2-(3-Amino-1-piperidinyl)ethyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt

15



Prepared according to the method described in Example 66c from 2-chloro-5-(2-oxoethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.094 g, Example 66b), 3-piperidinyl-carbamic acid, 1,1-dimethylethyl ester (0.109 g), sodium triacetoxyborohydride (0.081 g) and 1,2-dichloroethane (2 ml). After work-up, the residue was purified by HPLC eluting with a gradient of 0-5% ethanol in dichloromethane then by HPLC eluting with a gradient of 0-2% ethanol in dichloromethane. The white powder obtained was dissolved in methanol (2 ml) and a solution of hydrochloric acid in dioxane (4N, 1ml) was added. The

20

25


mixture was stirred for 14h at room temperature. Solvents were then removed under reduced pressure and the product obtained was triturated with diethyl ether to afford the title compound as a white powder (0.032 g).

5 MS (APCI +ve) 430 (M+H)⁺

¹H NMR (CD₃OD) δ 7.49-7.46 (1H, m); 7.42-7.39 (2H, m); 3.80-3.67 (3H, m); 3.46 (2H, t); 3.19 (2H, t); 3.08 (2H, s); 3.09-3.04 (1H, m); 2.14 (1H, bt); 2.00 (3H, bs); 1.74 (6H, q); 1.77-1.68 (2H, m); 1.64 (6H, d).

10 **Example 71**

2-Chloro-5-[2-(3-pyrrolidinylamino)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt



15

Prepared according to the method described in Example 66c from 2-chloro-5-(2-oxoethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.094 g, Example 66b), 3-amino-1-pyrrolidinecarboxylic acid, 1,1-dimethylethyl ester (0.101 g), sodium triacetoxyborohydride (0.081 g) and 1,2-dichloroethane (2 ml). After work-up, the residue
20 was purified by HPLC eluting with a gradient of 0-5% ethanol in dichloromethane. The pale orange powder obtained was dissolved in methanol (2 ml) and a solution of hydrochloric acid in dioxane (4N, 1ml) was added. The mixture was stirred for 14h at room temperature. Solvents were then removed under reduced pressure and the product obtained


was triturated with diethyl ether to afford the title compound as a pale orange powder (0.033 g).

MS (APCI +ve) 416 (M+H)⁺

5 ¹H NMR (CD₃OD) δ 7.46-7.39 (3H, m); 4.12 (1H, bs); 3.78-3.71 (1H, m); 3.69-3.57 (2H, m); 3.43-3.32 (4H, m); 3.13 (2H, bt); 3.06 (2H, s); 2.62-2.51 (1H, m); 2.38-2.29 (1H, m); 1.98 (3H, s); 1.73 (6H, q); 1.63 (6H, s).

Example 72

10 **5-[2-[(3*R*)-3-Aminopyrrolidinyl]ethyl]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt**



15 Prepared according to the method described in Example 66c from 2-chloro-5-(2-oxoethyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.094 g, Example 66b), (3*R*)-pyrrolidinyl-carbamic acid, 1,1-dimethylethyl ester (0.101 g), sodium triacetoxymethylborohydride (0.081 g) and 1,2-dichloroethane (2 ml). After work-up, the residue was purified by HPLC eluting with a gradient of 0-5% ethanol in dichloromethane. The


20 white powder obtained was dissolved in methanol (2 ml) and a solution of hydrochloric acid in dioxane (4N, 1ml) was added. The mixture was stirred for 14h at room temperature. Solvents were then removed under reduced pressure and the product obtained was triturated with diethyl ether to afford the title compound as a white powder (0.060 g).

25 MS (APCI +ve) 416 (M+H)⁺

¹H NMR (CD₃OD) δ 7.49-7.41 (3H, m); 4.86 (1H, bs); 4.05-3.80 (2H, m); 3.58 (4H, bs); 3.17 (2H, t); 3.08 (2H, s); 2.66 (1H, bs); 2.28 (1H, bs); 2.00 (3H, s); 1.74 (6H, q); 1.64 (6H, s).

5 **Example 73**


2-Chloro-5-[2-[2-(hydroxymethyl)-1-piperazinyl]ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



Example 74

2-Chloro-5-(hexahydro-1*H*-1,4-diazepin-1-yl)-*N*-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-benzamide, hydrochloride salt

5



a) 4-[4-Chloro-3-[[2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl]amino]carbonyl]phenyl]hexahydro-1*H*-1,4-diazepine-1-carboxylic acid, 1,1-dimethylethyl ester

A solution of 4-(3-carboxy-4-chlorophenyl)hexahydro-1*H*-1,4-diazepine-1-carboxylic acid, 1,1-dimethylethyl ester (0.075 g, Example 5b) and 1,1'-carbonyldiimidazole (0.034 g) in dimethylformamide (3 ml) was stirred at room temperature of 2.5h. Tricyclo[3.3.1.1^{3,7}]decane-1-ethanamine, hydrochloride salt (0.045 g) and *N,N*-diisopropylethylamine (0.037 ml) were then added and stirring continued for 14h. The reaction mixture was poured into water and extracted with ethyl acetate three times.

15 The ethyl acetate layers were combined and washed with 2M hydrochloric acid, 10% aqueous sodium hydroxide and brine, then dried over magnesium sulfate and concentrated under reduced pressure. Purification by chromatography on silica gel eluting with 20% ethyl acetate in *iso*-hexane gave the subtitle compound as a yellow oil (0.053 g).

20 MS (APCI +ve) 460/462 (*M*-^tBu)⁺

b) 2-Chloro-5-(hexahydro-1*H*-1,4-diazepin-1-yl)-*N*-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-benzamide, hydrochloride salt


4-[4-Chloro-3-[[[(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)amino]carbonyl]phenyl]-hexahydro-1*H*-1,4-diazepine-1-carboxylic acid, 1,1-dimethylethyl ester (0.053g, Example 74a) was dissolved in methanol (5 ml) and hydrochloric acid (0.5 ml from a 4N solution in dioxane) was added. After stirring at room temperature for 14h, the mixture was
5 evaporated to $\frac{3}{4}$ original volume under reduced pressure. Diethyl ether was gradually added to the solution and the resulting precipitate was collected by filtration, washed with diethyl ether and dried in vacuo to afford the title compound as a cream solid (0.017 g).

MS (APCI +ve) 416/418 (M-HCl)+

10 ¹H NMR (DMSO-d₆) δ 9.07 (2H, bs); 8.18 (1H, t); 7.22 (1H, d); 6.80 (1H, dd); 6.71 (1H, d); 3.70 (2H, m); 3.50 (2H, t); 3.25-3.17 (4H, m); 3.07 (2H, m); 2.09-2.06 (2H, m); 1.93 (3H, bs); 1.68 (3H, d); 1.61 (3H, d); 1.51 (6H, s); 1.34-1.28 (2H, m).

Example 75

15 (+/-)-5-(3-Amino-1-pyrrolidinyl)-2-chloro-*N*-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-benzamide, hydrochloride salt




Prepared as described in example 74 above using (+/-)-2-chloro-5-[3-[(1,1-dimethylethoxy)carbonyl]amino]-1-pyrrolidinyl]-benzoic acid (0.090 g), 1,1'-carbonyldiimidazole (0.043 g), tricyclo[3.3.1.1^{3,7}]decane-1-ethanamine, hydrochloride salt (0.057 g), *N,N*-diisopropylethylamine (0.046 ml) and dimethylformamide (3 ml). This
20 compound was treated with 4N hydrochloric acid in dioxane (0.5 ml) and methanol (5 ml) to yield the title compound (0.025 g).

MS (APCI +ve) 402/404 (M-HCl)+

¹H NMR (DMSO-d₆) δ 8.24 (3H, bs); 8.18 (1H, t); 7.24 (1H, d); 6.60 (1H, dd); 6.49 (1H, d); 3.93 (1H, m); 3.54-3.37 (2H, m); 3.31-3.17 (4H, m); 2.37-2.28 (1H, m); 2.07 (1H, m); 1.93 (3H, bs); 1.68 (3H, d); 1.61 (3H, d); 1.51 (6H, s); 1.34-1.28 (2H, m).

Example 76

2-Chloro-5-(4-piperidinylcarbonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) 4-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]benzoyl]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester

To dimethyl sulfoxide (0.155 ml) in anhydrous dichloromethane (11 ml) at -78°C was added oxalyl chloride (0.086 ml) and the mixture was stirred for 5min. at -78°C. A solution of 4-[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-hydroxymethyl]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (0.47 g, Example 64) in anhydrous dichloromethane (3 ml) was added dropwise and the mixture was stirred for 15min. at -78°C. Triethylamine (0.633 ml) was then added and the solution was warmed to room temperature. After 45min. at room temperature, the reaction mixture was poured onto water and the layers were separated. The organic extract was dried over anhydrous magnesium sulfate, filtered and concentrated under reduced pressure. The residue was purified by HPLC eluting a gradient of 0-5% ethanol in dichloromethane to give the subtitle compound as a white foam (0.31 g).

MS (APCI +ve) 15 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.46 (1H, t); 8.06-8.01 (1H, m); 7.95-7.92 (1H, m); 7.70-7.65 (1H, m); 3.97 (2H, bd); 3.72-3.61 (1H, m); 2.97 (2H, t); 2.92 (2H, bs); 1.96 (3H, bs); 1.77 (2H, d); 1.65 (6H, q); 1.55 (6H, s); 1.41 (9H, s); 1.48-1.33 (2H, m).

5

b) 2-Chloro-5-(4-piperidinylcarbonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt


A solution of 4-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]benzoyl]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (0.07 g, Example 76a) in methanol (3 ml) was treated with 4N hydrochloric acid solution in dioxane (1 ml). After 14h the solvents were removed under reduced pressure and the residue was triturated with diethyl ether to give the title compound as a white powder (0.025 g).

MS (APCI +ve) 415 (M+H-HCl)⁺

¹H NMR (DMSO-d₆) δ 8.90 (1H, bs); 8.64 (1H, bs); 8.46 (1H, t); 8.03 (1H, d); 7.95 (1H, s); 7.69 (1H, d); 3.81 (1H, t); 3.24-3.18 (2H, m); 3.09-2.99 (2H, m); 2.96 (2H, d); 2.01 (2H, dd); 1.95 (3H, s); 1.79 (2H, t); 1.64 (6H, q); 1.45 (6H, s).

Example 77

2-Chloro-5-[1-hydroxy-1-(4-piperidinyl)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) 4-[1-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-1-hydroxyethyl]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester

25

To methyl magnesiumbromide (3M solution in diethyl ether, 0.225 ml) in anhydrous diethyl ether (7 ml) under a nitrogen atmosphere was added slowly 4-[4-chloro-3-
[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl] benzoyl]-1-piperidinecarboxylic acid,
1,1-dimethylethyl ester (0.23 g, Example 76a) in anhydrous diethyl ether (7 ml). The
5 reaction mixture was stirred for 14h at room temperature, then poured onto crushed ice. A
solution of 10% aqueous potassium hydrogen sulphate was added keeping the pH of the
solution >4. The layers were separated, and the aqueous layer was extracted with ethyl
acetate (4x25 ml). The organic extracts were combined, dried over anhydrous magnesium
sulfate, filtered and concentrated under reduced pressure. The residue was dissolved in
10 methanol (3 ml) and hydrochloric acid (4N solution in dioxane, 2 ml) and stirred for 14h at
room temperature. Solvents were then removed under reduced pressure and the product
(0.11 g) was redissolved in dichloromethane (3 ml). Triethylamine (0.066 ml) was added
followed by di-*tert*-butyl-dicarbonate (0.055 g) and the reaction mixture was stirred for 1
hour at room temperature. Water was added and the layers were partitioned. The organic
15 extract was dried over anhydrous magnesium sulfate, filtered and concentrated under
reduced pressure. The residue was purified by HPLC eluting with a gradient of 0-5%
ethanol in dichloromethane, then by RPHPLC eluting with a gradient of 75-5% of 0.1%
aqueous ammonium acetate in acetonitrile to give the subtitle compound as a white foam
(0.06 g).

20 MS (APCI +ve) 431 (M+H-BOC)⁺

b) 2-Chloro-5-[1-hydroxy-1-(4-piperidinyl)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt


25 To a solution of 4-[1-[4-chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]
carbonyl] phenyl]-1-hydroxyethyl]-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester
(0.07 g, Example 77a) in methanol (3 ml) was added 4N hydrochloric acid solution in
dioxane (1 ml). After 14h the solvents were removed under reduced pressure and the
residue was triturated with diethyl ether to give the title compound as a white powder
30 (0.038 g).

MS (APCI +ve) 431 (M+H-HCl)⁺

¹H NMR (DMSO-d₆) δ 8.78 (1H, bs); 8.28 (1H, t); 7.44-7.40 (3H, m); 5.21 (1H, s); 3.25 (1H, d); 3.16 (1H, d); 2.98-2.89 (2H, m); 2.79-2.67 (2H, m); 1.94 (3H, bs); 1.84-1.75 (2H, m); 1.63 (6H, q); 1.53 (6H, s); 1.42 (3H, s); 1.53-1.31 (3H, m).

Example 78

2-Chloro-5-[2-(1-piperazinyl) ethoxy]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




Prepared as described in Example 12b using 2-chloro-5-hydroxy-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-methyl)-benzamide (Example 12a) and 4-(2-hydroxyethyl)-1-piperazinecarboxylic acid, 1,1-dimethylethyl ester.

MS (APCI +ve) 432 (M+H)⁺

¹H NMR (CD₃OD) δ 8.40 (1H, t); 7.41 (1H, d); 7.14-7.06 (2H, m); 4.45 (2H, t); 3.76-3.58 (10H, m); 3.07-3.03 (2H, m); 1.98 (3H, s); 1.77 (3H, d); 1.69 (3H, d); 1.62 (6H, s).

Example 79

2-Chloro-5-[2-(4-piperidinyloxy)ethoxy]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




- 5 Prepared as described in Example 12b using 2-chloro-5-hydroxy-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-methyl)-benzamide (Example 12a) and 4-(2-hydroxyethyl)-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester.

MS (APCI +ve) 431 (M+H)+

- 10 **¹H NMR (DMSO-d₆) δ** 8.75 (1H, brs); 8.49 (1H, brs); 8.27 (1H, t); 7.37 (1H, d); 6.99 (1H, dd); 6.91 (1H, d); 4.03 (2H, t); 3.22 (2H, d); 2.92 (2H, d); 2.82 (2H, t); 1.94 (3H, s); 1.82 (2H, d); 1.79-1.70 (1H, m); 1.69-1.64 (5H, m); 1.59 (3H, d); 1.53 (6H, s); 1.43-1.30 (2H, m).

15 **Example 80**

2-Chloro-5-[2-(4-piperidinyloxy)ethoxy]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




Prepared as described in Example 12b using 2-chloro-5-hydroxy-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-methyl)-benzamide (Example 12a) and 4-(2-hydroxyethoxy)-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester.

5 MS (APCI +ve) 447 (M+H)+

¹H NMR (CD₃OD) δ 8.39 (1H, brt); 7.38-7.33 (1H, m); 7.06-6.98 (2H, m); 4.22-4.16 (2H, m); 3.88-3.82 (2H, m); 3.80-3.71 (1H, m); 3.36-3.24 (2H, m); 3.16-3.04 (4H, m); 1.99 (3H, s); 2.08-1.86 (4H, m); 1.78 (3H, d); 1.69 (3H, d); 1.62 (6H, s).

10 **Example 81**

2-Chloro-5-[2-[2-(1-piperazinyl)ethoxy]ethoxy]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



Prepared as described in Example 12b using 2-chloro-5-hydroxy-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-methyl)-benzamide (0.20g, Example 12a) and 4-[2-(2-


15 hydroxyethoxy)ethyl]-1-piperazine carboxylic acid;1,1-dimethylethyl ester (0.26g).

MS (APCI +ve) 476 (M+H)+

20 ¹H NMR (CD₃OD) δ 7.7(1H, dd); 7.04-7.01 (2H, m); 4.25-4.18 (2H, m); 3.94 (2H, t); 3.91-3.87 (2H, m); 3.80-3.43 (10H, m); 3.06 (2H, s); 1.99 (3H, s); 1.75 (3H, d); 1.67 (3H, d); 1.62 (6H, s).

Example 82

2-Chloro-5-[(5,6-dihydro-1(4*H*)-pyrimidinyl)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




5 Prepared according to the method described in Example 8 from 5-bromomethyl-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 8b,) and 1,4,5,6-tetrahydro- pyrimidine.

MS (APCI +ve) 400/402 (M+H)⁺

10 ¹H NMR (CDCl₃) δ 7.84 (1H, s); 7.60 (1H, d); 7.43 (1H, d); 7.29 (1H, dd); 6.51 (1H, t); 4.39 (2H, s); 3.40-3.10 (3H, m); 3.17 (2H, d); 3.14 (1H, t); 2.01 (3H, s); 1.91 (q, 2H); 1.74 (3H, d); 1.64 (3H, d); 1.59 (6H, bs).

Example 83

15 **2-Chloro-5-[[4-[(2-hydroxyethyl)amino]-1-piperidinyl]methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt**



a) 2-Chloro-5-[(4-oxo-1-piperidiny)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Prepared according to the method described in Example 8c from 5-bromomethyl-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 8b) and 4-piperidinone.

MS (APCI +ve) 456/458 (M+H)⁺

b) 2-Chloro-5-[[4-[(2-hydroxyethyl)amino]-1-piperidiny)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt


To a solution of 2-chloro-5-[(4-oxo-1-piperidiny)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.150g, Example 83a) in methanol (3ml) at room temperature were added ethanolamine (0.11ml) and sodium cyanoborohydride (0.068g). The pH was adjusted to 6 by adding a 4N solution of hydrogen chloride in dioxane and the reaction stirred for 48h. The reaction was acidified with concentrated hydrochloric acid until gas evolution ceased. The precipitate was removed by filtration and the filtrate concentrated under vacuum. The residue was partitioned between ethyl acetate and water. The aqueous layer was basified with 5% aqueous sodium hydroxide and extracted with dichloromethane. The organics were washed with brine and dried over magnesium sulfate. The crude material was purified on silica gel (5% 7N ammonia in methanol/95% dichloromethane) to afford a white foam which was dissolved in ether/methanol and treated with a 4N solution of hydrogen chloride in dioxane to give the title compound (0.135 g).

MS (APCI +ve) 460/462 (M+H)⁺

¹H NMR (CD₃OD) δ 8.47 (1H, t); 7.67 (1H, d); 7.64 (1H, dd); 7.60 (1H, d); 4.39 (2H, s); 3.81 (2H, t); 3.62 (2H, bd); 3.52 (1H, t); 3.23-3.10 (4H, m); 3.09 (2H, d); 2.39 (2H, d); 2.07 (2H, q); 1.99 (s, 3H); 1.77 (3H, d); 1.70 (3H, d); 1.64 (6H, d).

Example 84

2-Chloro-5-[[4-hydroxy-4-[[[(1-methylethyl)amino]methyl]-1-piperidinyl]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



- 10 a) **2-Chloro-5-(1-oxa-6-azaspiro[2.5]oct-6-ylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

To dimethylsulfoxide (2ml) was added sodium hydride (0.033g, 60 % in oil) at room temperature. The mixture was stirred for 5min. at this temperature and a solution of trimethylsulfoxonium iodide (0.178g) in dimethylsulfoxide (2ml) was added. After 15 30min., 2-chloro-5-[[4-oxo-1-piperidinyl]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.28g, Example 83a) in dimethylsulfoxide (2ml) was added and the reaction stirred at room temperature for 3h before being quenched with ice/water (20ml). The mixture was extracted three times with ethyl acetate, the combined organic layers washed with brine and dried over magnesium sulfate. The crude material was purified on silica gel, 20 eluting with ethyl acetate to afford the subtitle compound as a white foam (0.25g).

MS (APCI +ve) 429/431 (M+H)⁺

¹H NMR (CDCl₃) δ 7.68 (1H, s); 7.40-7.30 (2H, m); 6.27 (1H, t); 3.54 (2H, s); 3.18 (2H, d); 2.70-2.50 (6H, m); 2.00 (s, 3H); 1.90-1.75 (2H, m); 1.74 (3H, d); 1.66 (3H, d); 25 1.59 (6H, bs); 1.80-1.50 (m, 2H).

- b) **2-Chloro-5-[[4-hydroxy-4-[[[(1-methylethyl)amino]methyl]-1-piperidinyl]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

In a sealed tube 2-chloro-5-(1-oxa-6-azaspiro[2.5]oct-6-ylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 84a, 0.15g) was dissolved in a 30

mixture of ethanol (4ml) and di-isopropylamine (1ml) and heated at 65°C for 14h. The volatiles were removed under vacuum and the residue purified on silica gel (5% 7N ammonia in methanol/95% dichloromethane) to afford the title compound as a white solid (0.115g).

5


MS (APCI +ve) 488/490 (M+H)⁺

¹H NMR (CD₃OD) δ 7.45-7.35 (3H, m); 3.55 (2H, s); 3.06 (2H, s); 2.76 (1H, q); 2.70-2.55 (2H, m); 2.54 (2H, s); 2.50-2.35 (2H, m); 1.99 (s, 3H); 1.77 (3H, d); 1.70 (3H, d); 1.63 (10H, bs); 1.07 (m, 2H).

10

Example 85

2-Chloro-5-[(1,2,3,6-tetrahydro-3-pyridinyl)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide, hydrochloride salt



15

To pyridine (6ml) at 0°C was added portionwise lithium aluminium hydride (0.24g). The mixture was allowed to warm to room temperature and was stirred for 24h. Lithium iodide (0.220g) and pyridine (1ml) were added and the reaction stirred for a further 1h.

5-Bromomethyl-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.30g,

Example 8b) in dry pyridine (2ml) was added to the solution at room temperature. After 2h,

20

the mixture was quenched at 0°C with a cold 15 % aqueous solution of acetic acid, stirred for an hour and concentrated under vacuum. The residue was taken in 1N sodium hydroxide, extracted with dichloromethane and the organic layers dried over magnesium sulfate. The crude material was purified on silicagel (2 to 10% 7N ammonia in


methanol/dichloromethane) then treated with a 4N solution of hydrogen chloride in dioxane and methanol to give the title compound (0.20g).

MS (APCI +ve) 399/401 (M+H)⁺

¹H NMR (CD₃OD) δ 8.42 (1H, t); 7.43 (1H, dd); 7.32 (1H, dd); 7.30 (d; 1H); 5.89 (1H, d); 5.80 (1H, d); 3.64 (2H, s); 3.40-3.30 (1H, m); 3.06 (4H, d); 1.99 (s, 3H); 1.78 (3H, d); 1.68 (3H, d); 1.63 (6H, d).

Example 86

2-Chloro-5-(3-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide, acetate salt



a) 2-Chloro-5-(3-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide, acetate salt

To pyridine (12 ml) at 0 °C was added portionwise lithium aluminium hydride (0.46g). The mixture was allowed to warm to room temperature and was stirred for 24h. Lithium iodide (0.44g) and pyridine (5ml) were added and the reaction stirred for a further 1h. The solution was cooled to -10 °C and 5-bromomethyl-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 8b, 0.5g) in dry pyridine (5ml) added. After 1 hour, the mixture was quenched at -10 degrees with cold water, then 1N sodium hydroxide. The solution was stirred for 1h then concentrated under vacuum. The residue was taken in water, extracted with dichloromethane and the organic layers dried over magnesium sulfate. The crude material was purified on silica gel (ethylacetate : i-hexane / 4 : 1) to afford the subtitle compound as a white foam (0.355g).

b) 2-Chloro-5-(3-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetate salt


2-Chloro-5-(3-pyridinylmethyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.10g, Example 86a) was dissolved in methanol and treated with a 4N solution of hydrogen chloride in dioxane. The hydrochloride salt was isolated and hydrogenated in ethanol over platinum oxide following the procedure described in Example 51 to give the title compound as the acetate salt after purification by reverse phase HPLC (0.1% aqueous ammonium acetate/acetonitrile) (0.053g).

MS (APCI +ve) 401/403 (M+H)⁺

¹H NMR (CD₃OD) δ 7.42 (1H, d); 7.31-7.25 (2H, m); 3.38-3.28 (1H; m); 3.28-3.18 (1H; 2m); 3.07 (2H, s); 2.86 (1H, dt); 2.72-2.61 (1H, m); 2.65 (1H, d); 2.13-2.05 (1H; m); 2.01 (3H, s); 1.94 (3H, s); 1.90-1.85 (1H; m); 1.85-1.60 (2H; m); 1.80 (3H, d); 1.71 (3H, d); 1.64 (6H, d); 1.29 (1H, qd).

Example 87

**2-bromo-5-[[4-[(2-hydroxyethyl)amino]-1-piperidinyl)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]
dec -1-yl)methyl- benzamide**




Prepared according to the procedures described in Example 83a and 83b from 2-bromo-5-bromomethyl-N-(tricyclo[3.3.1.^{3,7}]dec-1-ylmethyl)-benzamide (Example 65a), 4-piperidone and ethanolamine

MS (APCI +ve) MW 505/506 (M+H)⁺

¹H NMR (CDCl₃) δ 7.53 (1H, d); 7.52 (1H, d); 7.26 (1H, dd); 6.05 (1H, t); 3.65 (2H, t);
3.46 (2H, s); 3.17 (2H, d); 2.82 (2H, t); 2.60-2.45 (1H, m); 2.20-1.95 (7H, m); 1.95-1.82
5 (2H, bd); 1.75 (3H, d); 1.66 (3H, d); 1.61 (6H, d); 1.50-1.35 (2H, qd).

Example 88

2-Chloro-5-[(E)-3-piperidinylidenemethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



10 **a) 2-Chloro-5-[(E)-(5,6-dihydro-3(4H)-pyridinylidene)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

2-Chloro-5-formyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.152 g, Example 31a) and 2,3,4,5-tetrahydropyridine trimer (Org. Synth., 1977, Vol.56, 118-122,
15 0.038 g) were dissolved in methanol (3 ml) and heated at reflux for 4h. The mixture was evaporated under reduced pressure then purified by HPLC eluting a gradient of 0-5% ethanol in dichloromethane to give the subtitle compound as a white foam (0.046 g).

MS (APCI +ve) 397 (M+H)⁺

20 ¹H NMR (CD₃OD) δ 7.98 (1H, s); 7.51 (3H, s); 6.83 (1H, s); 3.66-3.62 (2H, m); 3.07 (2H, s); 2.78-2.73 (2H, m); 1.99 (3H, bs); 1.73 (6H, q); 1.77-1.69 (2H, m); 1.64 (6H, s).

b) 2-Chloro-5-[(E)-3-piperidinylidenemethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Sodium borohydride (0.009 g) in methanol (0.5 ml) was added to a solution of 2-chloro-5-[(E)-(5,6-dihydro-3(4H)-pyridinylidene)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.046 g, Example 88a,) in methanol (1.5 ml). The reaction mixture was stirred for 4h under an atmosphere of nitrogen at room temperature. Concentrated hydrochloric acid (0.01 ml) was added and the mixture was evaporated under reduced pressure. An aqueous solution of sodium hydroxide (2M, 2 ml) was added to rebaseify the residue followed by water (10 ml) and dichloromethane (10 ml). The layers were partitioned and the aqueous layer was extracted further with dichloromethane (2x10 ml). The organic extracts were combined, dried over anhydrous magnesium sulfate, filtered and concentrated under reduced pressure to give the title compound as a white powder (0.024 g).

MS (APCI +ve) 399 (M+H)⁺

¹H NMR (CD₃OD) δ 7.43 (1H, d); 7.29-7.26 (2H, m); 6.38 (1H, s); 3.45 (2H, s); 3.07 (2H, s); 2.96 (2H, t); 2.55 (2H, t); 2.00 (3H, bs); 1.75 (6H, q); 1.81-1.64 (2H, m); 1.64 (6H, s).

Pharmacological Analysis

Certain compounds such as benzoylbenzoyl adenosine triphosphate (bbATP) are known to be agonists of the P2X₇ receptor, effecting the formation of pores in the plasma membrane (Drug Development Research (1996), 37(3), p.126). Consequently, when the receptor is activated using bbATP in the presence of ethidium bromide (a fluorescent DNA probe), an increase in the fluorescence of intracellular DNA-bound ethidium bromide is observed. The increase in fluorescence can be used as a measure of P2X₇ receptor activation and therefore to quantify the effect of a compound on the P2X₇ receptor.


In this manner, each of the title compounds of Examples 1 to 88 was tested for antagonist activity at the P2X₇ receptor. Thus, the test was performed in 96-well flat bottomed microtitre plates, the wells being filled with 250 µl of test solution comprising 200 µl of a suspension of THP-1 cells (2.5 x 10⁶ cells/ml) containing 10⁻⁴ M ethidium

bromide, 25 μ l of a high potassium buffer solution containing 10^{-5} M bbATP, and 25 μ l of the high potassium buffer solution containing 3×10^{-5} M test compound. The plate was covered with a plastics sheet and incubated at 37 °C for one hour. The plate was then read in a Perkin-Elmer fluorescent plate reader, excitation 520 nm, emission 595 nm, slit widths: Ex 15 nm, Em 20 nm. For the purposes of comparison, bbATP (a P2X₇ receptor agonist) and pyridoxal 5-phosphate (a P2X₇ receptor antagonist) were used separately in the test as controls. From the readings obtained, a pIC₅₀ figure was calculated for each test compound, this figure being the negative logarithm of the concentration of test compound necessary to reduce the bbATP agonist activity by 50%. Each of the compounds of Examples 1 to 88 demonstrated antagonist activity, having a pIC₅₀ figure > 4.50.

CLAIMS



1. A compound of general formula




5 wherein m represents 1, 2 or 3;

each R¹ independently represents a hydrogen or halogen atom;

A represents C(O)NH or NHC(O);

Ar represents a group



10 X represents a bond, an oxygen atom or a group CO, (CH₂)₁₋₆, CH=, (CH₂)₁₋₆O, O(CH₂)₁₋₆, O(CH₂)₂₋₆O, O(CH₂)₂₋₃O(CH₂)₁₋₃, CR(OH), (CH₂)₁₋₃O(CH₂)₁₋₃, (CH₂)₁₋₃O(CH₂)₂₋₃O, NR⁵, (CH₂)₁₋₆NR⁵, NR⁵(CH₂)₁₋₆, (CH₂)₁₋₃NR⁵(CH₂)₁₋₃, O(CH₂)₂₋₆NR⁵, O(CH₂)₂₋₃NR⁵(CH₂)₁₋₃, (CH₂)₁₋₃NR⁵(CH₂)₂₋₃O, NR⁵(CH₂)₂₋₆O, NR⁵(CH₂)₂₋₃O(CH₂)₁₋₃, CONR⁵, NR⁵CO, S(O)_n, S(O)_nCH₂, CH₂S(O)_n, SO₂NR⁵
 15 or NR⁵SO₂;

n is 0, 1 or 2;

R' represents a hydrogen atom or a C₁-C₆ alkyl group;

one of R² and R³ represents a halogen, cyano, nitro, amino, hydroxyl, or a group selected from (i) C₁-C₆ alkyl optionally substituted by at least one C₃-C₆ cycloalkyl,

20 (ii) C₃-C₈ cycloalkyl, (iii) C₁-C₆ alkyloxy optionally substituted by at least one C₃-C₆ cycloalkyl, and (iv) C₃-C₈ cycloalkyloxy, each of these groups being optionally

substituted by one or more fluorine atoms, and the other of R^2 and R^3 represents a hydrogen or halogen atom;

either R^4 represents a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom, the

5 heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl, carboxyl, cyano, C_1 - C_6 alkyl, C_1 - C_6 hydroxyalkyl, $-NR^6R^7$, $-(CH_2)_rNR^6R^7$ and $-CONR^6R^7$,

or R^4 represents a 3- to 8-membered saturated carbocyclic ring system substituted by one or more substituents independently selected from $-NR^6R^7$, $-(CH_2)_rNR^6R^7$ and

10 $-CONR^6R^7$, the ring system being optionally further substituted by one or more substituents independently selected from fluorine atoms, hydroxyl and C_1 - C_6 alkyl;

r is 1, 2, 3, 4, 5 or 6;

R^5 represents a hydrogen atom or a C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl group;

R^6 and R^7 each independently represent a hydrogen atom or a C_1 - C_6 alkyl,

15 C_2 - C_6 hydroxyalkyl or C_3 - C_8 cycloalkyl group, or R^6 and R^7 together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring; with the provisos that,

(a) when A represents $C(O)NH$ and R^4 represents an unsubstituted 3- to 8-membered saturated aliphatic heterocyclic ring system containing one nitrogen atom, then X is other
20 than a bond, and

(b) when A represents $C(O)NH$ and X represents a group $(CH_2)_{1-6}$ or $O(CH_2)_{1-6}$, then R^4 does not represent an unsubstituted imidazolyl, unsubstituted morpholinyl, unsubstituted piperidinyl or unsubstituted pyrrolidinyl group, and

(c) when A represents $NHC(O)$ and R^4 represents an unsubstituted 3- to 8-membered
25 saturated aliphatic heterocyclic ring system containing one nitrogen atom, then X is other than a bond, and


(d) when A represents $NHC(O)$ and X represents $O(CH_2)_{1-6}$, $NH(CH_2)_{1-6}$ or SCH_2 , then R^4 does not represent an unsubstituted 1-piperidinyl or unsubstituted 1-pyrrolidinyl group, and



(e) when A represents NHC(O) and X represents $\text{O(CH}_2\text{)}_{2-3}\text{NH(CH}_2\text{)}_2$, then R^4 does not represent an imidazolyl group;
or a pharmaceutically acceptable salt or solvate thereof.

2. A compound according to claim 1, wherein A represents NHC(O) .

3. A compound according to claim 1 or claim 2, wherein Ar represents a group




4. A compound according to any one of claims 1 to 3, wherein X represents a bond, an oxygen atom or a group CO , $(\text{CH}_2)_{1-6}$, CH= , $\text{O(CH}_2\text{)}_{1-6}$, $\text{O(CH}_2\text{)}_{2-6}\text{O}$, $\text{O(CH}_2\text{)}_{2-3}\text{O(CH}_2\text{)}_{1-3}$, CR(OH) , NR^5 , $(\text{CH}_2)_{1-6}\text{NR}^5$, CONR^5 , S(O)_n or $\text{S(O)}_n\text{CH}_2$.


5. A compound according to any one of claims 1 to 4, wherein R^4 represents a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom, the heterocyclic ring system being optionally substituted by one or two substituents independently selected from hydroxyl, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ hydroxyalkyl, $\text{-NR}^6\text{R}^7$ and $\text{-(CH}_2\text{)}_r\text{NR}^6\text{R}^7$.

6. A compound according to any one of claims 1 to 4, wherein R^4 represents a group selected from:





* C L M S *



- 5 7. A compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, according to claim 1 being:

2-Nitro-3-piperazin-1-yl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Amino-3-piperazin-1-yl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

dihydrochloride salt,



- 2-Chloro-3-piperazin-1-yl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
 2-Chloro-5-piperazin-1-yl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
 2-Chloro-5-(hexahydro-1H-1,4-diazepin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 benzamide, hydrochloride salt,
 5 (5-(4-Amino-1-piperidinyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 benzamide, hydrochloride salt,
 (+/-)-5-(3-Amino-1-pyrrolidinyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 benzamide, hydrochloride salt,
 2-Chloro-5-piperazin-1-ylmethyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
 10 hydrochloride salt,
 2-Chloro-5-[(hexahydro-1H-1,4-diazepin-1-yl)methyl] -N-(tricyclo[3.3.1.1^{3,7}]dec-1-
 ylmethyl)-benzamide, hydrochloride salt,
 5-[(4-Amino-1-piperidinyl)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 benzamide, hydrochloride salt,
 15 5-[(3-Amino-1-pyrrolidinyl)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 benzamide, hydrochloride salt,
 2-Chloro-5-(4-piperidinylloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
 hydrochloride salt,
 (R)-2-Chloro-5-(2-pyrrolidinylmethoxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 20 benzamide, hydrochloride salt,
 (S)-2-Chloro-5-(2-pyrrolidinylmethoxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 benzamide, hydrochloride salt,
 2-Chloro-5-(3-piperidinylmethoxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 benzamide, hydrochloride salt,
 25 cis-5-[(4-Aminocyclohexyl)oxy]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 benzamide, hydrochloride salt,
 2-Methyl-5-(1-piperazinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
 hydrochloride salt,
 2-Chloro-5-(1-piperazinylmethyl)-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-benzamide,
 30 hydrochloride salt,



(+/-)-2-Chloro-5-(3-pyrrolidinyloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

(+/-)-2-Chloro-5-(3-piperidinyloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

5 trans-5-[(4-Aminocyclohexyl)oxy]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide,

cis-(+/-)-5-[(3-Aminocyclopentyl)oxy]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
ylmethyl)-benzamide,

(S,S)-2-Chloro-5-(2,5-diazabicyclo[2.2.1]hept-2-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
10 ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(2-methyl-1-piperazinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

(+/-)-2-Chloro-5-(3-pyrrolidinylamino)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, hydrochloride salt,

15 (+/-)-5-(3-Amino-1-piperidinyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide,

(+/-)-2-Chloro-5-(3-piperidinylamino)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide,

2-Chloro-5-[hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
20 ylmethyl)-benzamide,

N-[2-methyl-5-(4-piperidinyloxy)phenyl]-tricyclo[3.3.1.1^{3,7}]decane-1-acetamide,
hydrochloride salt,

N-[2-chloro-5-(4-piperidinyloxy)phenyl]-tricyclo[3.3.1.1^{3,7}]decane-1-acetamide,
hydrochloride salt,

25 2-Chloro-5-[(4-piperidinylamino)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
benzamide, dihydrochloride salt,

5-[[[4-(Aminomethyl)cyclohexyl]amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-
1-ylmethyl)-benzamide, dihydrochloride salt,

30 5-[[[4-Aminocyclohexyl]amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-
ylmethyl)-benzamide, dihydrochloride salt,



* C L M S *

5-[(1-Azabicyclo[2.2.2]oct-3-ylamino)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

N-[4-(3-Aminopyrrolidin-1-yl)-2-methylphenyl]-2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetamide, dihydrochloride salt,

5 N-(2-Methyl-4-piperazin-1-ylphenyl)-2-(tricyclo[3.3.1.1^{3,7}]dec-1-yl)acetamide, dihydrochloride salt,

cis-4-(3-Amino-cyclopentyloxy)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-4-(4-piperidinyloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

10 (+/-)-2-Chloro-4-(pyrrolidin-3-yloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-4-(piperidin-3-yloxy)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

15 2-Chloro-4-(4-piperazin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-4-(3-pyrrolidinylamino)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-4-(hexahydro-1H-1,4-diazepin-1-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

20 (±)-5-[(3-Amino-1-piperidinyloxy)methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(2,5-diazabicyclo[2.2.1]hept-2-ylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

25 2-Chloro-5-(9-oxa-3,7-diazabicyclo[3.3.1]non-3-ylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(3,7-diazabicyclo[3.3.1]non-3-ylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

trans-2-Chloro-5-[[8-(methylamino)-3-azabicyclo[3.2.1]oct-3-yl]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,



cis-2-Chloro-5-[(hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(4-piperidinylidenemethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

5 2-Chloro-5-(4-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(4-hydroxy-piperidin-4-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(1,2,3,6-tetrahydro-pyridin-4-yl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Ethyl-5-piperazin-1-ylmethyl -N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(piperidin-4-ylsulfanyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

15 2-Chloro-5-(piperidin-4-ylsulfinyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-(piperidin-4-ylsulfonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(piperidin-4-ylmethylsulfanyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(piperidin-4-ylmethanesulfonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(piperazine-1-carbonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

25 2-Chloro-5-([1,4]diazepane-1-carbonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

4-Chloro-N¹-(piperidin-4-yl)-N²-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-isophthalamide, hydrochloride salt,

2-Chloro-5-(hydroxy-4-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,



(±)-2-Chloro-5-(hydroxy-3-piperidinylmethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Bromo-5-piperazin-1-ylmethyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

5 2-Chloro-5-[2-(1-piperazinyl)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[2-(2,5-diazabicyclo[2.2.1]hept-2-yl)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

10 5-[2-(4-Amino-1-piperidinyl)ethyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[2-(3-piperidinylamino)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt,

5-[2-(3-Amino-1-piperidinyl)ethyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

15 2-Chloro-5-[2-(3-pyrrolidinylamino)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt,

5-[2-[(3R)-3-Aminopyrrolidinyl]ethyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

20 2-Chloro-5-[2-[2-(hydroxymethyl)-1-piperazinyl]ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-(hexahydro-1H-1,4-diazepin-1-yl)-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-benzamide, hydrochloride salt,

(+/-)-5-(3-Amino-1-pyrrolidinyl)-2-chloro-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl)-benzamide, hydrochloride salt,

25 2-Chloro-5-(4-piperidinylcarbonyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[1-hydroxy-1-(4-piperidinyl)ethyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

30 2-Chloro-5-[2-(1-piperazinyl)ethoxy]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,



2-Chloro-5-[2-(4-piperidinyloxy)ethoxy]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[2-(4-piperidinyloxy)ethoxy]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

5 2-Chloro-5-[2-[2-(1-piperazinyl)ethoxy]ethoxy]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[(5,6-dihydro-1(4*H*)-pyrimidinyl)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[[4-[(2-hydroxyethyl)amino]-1-piperidinyloxy]methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide hydrochloride salt,

2-Chloro-5-[[4-hydroxy-4-[(1-methylethyl)amino]methyl]-1-piperidinyloxy]methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[(1,2,3,6-tetrahydro-3-pyridinyl)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

15 2-Chloro-5-(3-piperidinylmethyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetate salt,


2-bromo-5-[[4-[(2-hydroxyethyl)amino]-1-piperidinyloxy]methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, or

2-Chloro-5-[(*E*)-3-piperidinylidenemethyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide.

8. A process for the preparation of a compound of formula (I) as defined in claim 1 which comprises:

25 (i) when X represents a CH₂ group, R⁴ represents a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom, the heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl, carboxyl, cyano, C₁-C₆ alkyl, C₁-C₆ hydroxyalkyl, -NR⁶R⁷, -(CH₂)_rNR⁶R⁷ and -CONR⁶R⁷ and
30 R⁴ is linked to X through a nitrogen atom, reacting a compound of general formula





(II)

wherein one of R^{10} and R^{11} represents a hydrogen atom and the other of R^{10} and R^{11} represents a group $-\text{CH}_2\text{L}^1$ in which L^1 represents a leaving group and m , A , R^1 , R^2 and R^3 are as defined in formula (I), with a compound of general formula




in the presence of a base, wherein $R^{4'}$ represents a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom, the heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl, carboxyl, cyano, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ hydroxyalkyl, $-\text{NR}^6\text{R}^7$, $-(\text{CH}_2)_t\text{NR}^6\text{R}^7$ and $-\text{CONR}^6\text{R}^7$ and wherein R^6 and R^7 are as defined in formula (I); or

10

(ii) when X represents an oxygen atom or a group $\text{O}(\text{CH}_2)_{1-6}$, $\text{O}(\text{CH}_2)_{2-6}\text{O}$, $\text{O}(\text{CH}_2)_{2-3}\text{O}(\text{CH}_2)_{1-3}$, $\text{O}(\text{CH}_2)_{2-6}\text{NR}^5$ or $\text{O}(\text{CH}_2)_{2-3}\text{NR}^5(\text{CH}_2)_{1-3}$, reacting a compound


15 of general formula



(IV)




wherein one of R^{12} and R^{13} represents a hydrogen atom and the other of R^{12} and R^{13} represents a hydroxyl group and m , A , R^1 , R^2 and R^3 are as defined in formula (I), with a compound of general formula




wherein Y represents a bond or a group $(CH_2)_{1-6}$, $O(CH_2)_{2-6}$, $(CH_2)_{1-3}O(CH_2)_{2-3}$, $NR^5(CH_2)_{2-6}$ or $(CH_2)_{1-3}NR^5(CH_2)_{2-3}$ and R^4 is as defined in formula (I), in the presence of 1,1-(azodicarbonyl)dipiperidine and tributylphosphine; or

(iii) when X represents a bond, an oxygen atom or a group $O(CH_2)_{1-6}$, $O(CH_2)_{2-6}O$, $O(CH_2)_{2-3}O(CH_2)_{1-3}$, NR^5 , $NR^5(CH_2)_{1-6}$, $NR^5(CH_2)_{2-6}O$ or $NR^5(CH_2)_{2-3}O(CH_2)_{1-3}$ and A is $NHC(O)$, reacting a compound of general formula




wherein one of R^{14} and R^{15} represents a group $-X'-R^4$ and the other of R^{14} and R^{15} represents a hydrogen atom, X' represents a bond, an oxygen atom or a group $O(CH_2)_{1-6}$, $O(CH_2)_{2-6}O$, $O(CH_2)_{2-3}O(CH_2)_{1-3}$, NR^5 , $NR^5(CH_2)_{1-6}$, $NR^5(CH_2)_{2-6}O$ or $NR^5(CH_2)_{2-3}O(CH_2)_{1-3}$, L^2 represents a leaving group and R^2 , R^3 , R^4 and R^5 are as defined in formula (I), with a compound of general formula




wherein m and R^1 are as defined in formula (I), optionally in the presence of a coupling agent; or



(iv) when X represents a bond, an oxygen atom or a group $\text{O}(\text{CH}_2)_{1-6}$, $\text{O}(\text{CH}_2)_{2-6}\text{O}$, $\text{O}(\text{CH}_2)_{2-3}\text{O}(\text{CH}_2)_{1-3}$, NR^5 , $\text{NR}^5(\text{CH}_2)_{1-6}$, $\text{NR}^5(\text{CH}_2)_{2-6}\text{O}$ or $\text{NR}^5(\text{CH}_2)_{2-3}\text{O}(\text{CH}_2)_{1-3}$ and A is $\text{C}(\text{O})\text{NH}$, reacting a compound of general formula




wherein R^2 and R^3 are as defined in formula (I) and R^{14} and R^{15} are as defined in formula (VI) in (iii) above, with a compound of general formula



wherein m and R^1 are as defined in formula (I), in the presence of a base; or

(v) when X represents a bond or a group NR^5 , $\text{NR}^5(\text{CH}_2)_{1-6}$, $\text{NR}^5(\text{CH}_2)_{2-6}\text{O}$ or $\text{NR}^5(\text{CH}_2)_{2-3}\text{O}(\text{CH}_2)_{1-3}$, reacting a compound of general formula



wherein one of R^{16} and R^{17} represents a leaving group, L^3 , and the other of R^{16} and R^{17} represents a hydrogen atom and m, A, R^1 , R^2 and R^3 are as defined in formula (I), with a compound of general formula



5 wherein Z represents a hydrogen atom or a group NHR^5 , $(CH_2)_{1-6}NHR^5$, $O(CH_2)_{2-6}NHR^5$ or a group $(CH_2)_{1-3}O(CH_2)_{2-3}NHR^5$ and R^4 and R^5 are as defined in formula (I), optionally in the presence of a palladium catalyst, a phosphine ligand and a base; or

10 (vi) when X represents a group CH_2O , reacting a compound of formula (II) as defined in (i) above with a compound of formula (V) as defined in (ii) above wherein Y represents a bond, in the presence of a base or in the presence of a metal salt; or


(vii) when X represents a group CH_2NR^5 , reacting a compound of formula (II) as
15 defined in (i) above with a compound of formula (XI) as defined in (v) above wherein Z represents a group NHR^5 ; or

(viii) when X represents a group $CH_2O(CH_2)_{1-3}$ or $CH_2O(CH_2)_{2-3}O$, reacting a
20 compound of formula (II) as defined in (i) above with a compound of formula (V) as defined in (ii) above wherein Y represents a group $(CH_2)_{1-3}$ or $O(CH_2)_{2-3}$, in the presence of a base or in the presence of a metal salt; or

(ix) when X represents a group $CH_2NR^5CH_2$ or $CH_2NR^5(CH_2)_{2-3}O$ reacting a
25 compound of formula (II) as defined in (i) above with a compound of formula (XI) as defined in (v) above wherein Z represents a group CH_2NHR^5 or $O(CH_2)_{2-3}NHR^5$; or


(x) when X represents a group CH_2 and R^4 represents an unsubstituted 4- to
6-membered saturated aliphatic heterocyclic ring system containing one nitrogen atom,
reacting a compound of formula (II) as defined in (i) above, with a compound of general
30 formula






wherein s and t independently represent 1 or 2; or

- 5 (xi) when X represents a group CO, CONR⁵, NR⁵CO, SO₂, NR⁵SO₂ or SO₂NR⁵ and A is NHC(O), reacting a compound of general formula



- wherein one of R¹⁸ and R¹⁹ represents a group -X"-R⁴ and the other of R¹⁸ and R¹⁹ represents a hydrogen atom, X" represents a group CO, CONR⁵, NR⁵CO, SO₂,
 10 NR⁵SO₂ or SO₂NR⁵, L⁴ represents a leaving group and R², R³, R⁴ and R⁵ are as defined in formula (I), with a compound of formula (VII) as defined in (iii) above, optionally in the presence of a coupling agent; or

- (xii) when X represents a group CO, CONR⁵, NR⁵CO, SO₂, NR⁵SO₂ or
 15 SO₂NR⁵ and A is C(O)NH, reacting a compound of general formula



- wherein R² and R³ are as defined in formula (I) and R¹⁸ and R¹⁹ are as defined in formula (XIII) in (xi) above, with a compound of formula (IX) as defined in (iv) above, in the presence of a base; or



(xiii) when X represents a sulfur atom, reacting a compound of formula (X) as defined in (v) above, with an organolithium reagent and then with a compound of general formula



5 wherein Tol represents a tolyl group and R^4 is as defined in formula (I); or

(xiv) when X represents a CHOH or CH_2 group, reacting a compound of formula (X) as defined in (v) above, with an organolithium reagent and then with a compound of general formula



wherein R^4 is as defined in formula (I), optionally followed by a reduction reaction; or

(xv) when X represents a bond, reacting a compound of formula (X) as defined in (v) above, with an organolithium reagent and then with a compound of general formula




wherein R^4 is as defined in formula (I), optionally followed by a reduction reaction;

(xvi) when X represents a group SO, oxidising a corresponding compound of formula (I) in which X represents a sulphur atom; or

20

(xvii) when X represents a group SCH_2 , reacting a compound of formula (X) as defined in (v) above, with an organolithium reagent and then with a compound of general formula




wherein R^4 is as defined in formula (I); or



(xviii) when X represents a group SOCH_2 or SO_2CH_2 , oxidising a corresponding compound of formula (I) in which X represents a group SCH_2 ; or

(xix) when X represents a group CH= , reacting a compound of formula (II) as
 5 defined in (i) above with trimethyl phosphite and then with a compound of formula (XVII) as defined in (xv) above in the presence of a base; or

(xx) when X represents a group $(\text{CH}_2)_{1-6}$, reacting a compound of general formula



10

wherein one of R^{20} and R^{21} represents a group CHO or a group $(\text{CH}_2)_{1-5}\text{CHO}$ and the other of R^{20} and R^{21} represents a hydrogen atom, and m , A , R^1 , R^2 and R^3 are as defined in formula (I), with a compound of general formula (XX), $\text{R}^4\text{-H}$, wherein R^4 is as defined in formula (I), in the presence of a reducing agent; or

15

(xxi) when X represents a group $(\text{CH}_2)_{1-6}\text{NR}^5$, $(\text{CH}_2)_{1-3}\text{NR}^5(\text{CH}_2)_{1-3}$ or $(\text{CH}_2)_{1-3}\text{NR}^5(\text{CH}_2)_{2-3}\text{O}$, reacting a compound of formula (XIX) as defined in (xx) above, with a compound of general formula (XXI), $\text{R}^4\text{-Z'}$, wherein Z' represents a group NHR^5 , $(\text{CH}_2)_{1-3}\text{NHR}^5$, $\text{O}(\text{CH}_2)_{2-3}\text{NHR}^5$ and R^4 and R^5 are as defined in formula (I), in
 20 the presence of a reducing agent; or


(xxii) when X represents a group $(\text{CH}_2)_{1-3}\text{O}(\text{CH}_2)_{1-3}$ or $(\text{CH}_2)_{1-3}\text{O}(\text{CH}_2)_{2-3}\text{O}$, reacting a compound of formula (XIX) as defined in (xx) above in which one of R^{20} and R^{21} represents a group CHO or a group $(\text{CH}_2)_{1-2}\text{CHO}$ and the other of R^{20} and R^{21}



represents a hydrogen atom, with a reducing agent, followed by reaction with a compound of general formula (XXII), R^4-E , wherein E represents a group $(CH_2)_{1-3}L^5$ or $O(CH_2)_{2-3}L^5$, L^5 is a leaving group and R^4 is as defined in formula (I), in the presence of a base; or

(xxiii) when X represents a group $(CH_2)_{1-6}$, reacting a compound of formula (II) as defined in (i) above with trimethylphosphite, and then with a compound of formula (XVI) as defined in (xiv) above, or with a compound of formula (XVII) as defined in (xv) above or with a compound of general formula (XVIA), $R^4(CH_2)_{1-4}CHO$ in which R^4 is as defined in formula (I), in the presence of a base, followed by a reduction reaction; or

(xxiv) when X represents a group $(CH_2)_{2-6}O$, reacting a compound of general formula



wherein one of R^{22} and R^{23} represents a group $(CH_2)_{2-6}L^6$ and the other of R^{20} and R^{21} represents a hydrogen atom, L^6 represents a leaving group and m, A, R^1 , R^2 and R^3 are as defined in formula (I), with a compound of formula (V) as defined in (ii) above in which Y represents a bond; or

(xxv) when X represents a group $CR'(OH)$ in which R' is a C_1-C_6 alkyl group, oxidising a corresponding compound of formula (I) in which X represents $CH(OH)$, followed by reaction with a C_1-C_6 alkyllithium reagent; or




(xxvi) when X represents a group CH_2S , reacting a compound of formula (II) as defined in (i) above with a compound of general formula (XXIV), $\text{R}^4\text{-SH}$, wherein R^4 is as defined in formula (I), in the presence of a base; or

5 (xxvii) when X represents a group CH_2SO or CH_2SO_2 , oxidising a corresponding compound of formula (I) in which X represents a group CH_2S ; or

(xxviii) when X represents a group CH_2 and R^4 represents a 3-piperidinyl or 2-piperazinyl group, reacting a compound of formula (II) as defined in (i) above with a
10 reagent formed by combining pyridine or pyrazine with an aluminium hydride reagent, followed by a reduction reaction; or

(xxix) when X represents a group CH= and R^4 represents a 3-piperidinyl group, reacting a compound of general formula




15

wherein one of R^{24} and R^{25} represents an aldehyde group -CHO , and the other of R^{24} and R^{25} represents a hydrogen atom and m , A , R^1 , R^2 and R^3 are as defined in formula (I), with 2,3,4,5-tetrahydropyridine, followed by a reduction reaction; or

20 (xxx) when X represents a bond, NR^5 or $\text{NR}^5(\text{CH}_2)_{1-6}$ and R^4 represents a carbon-linked piperidyl or piperazinyl group, reducing a compound of general formula






(XXVI)

wherein one of R^{26} and R^{27} represents a pyridyl, pyrazinyl, NR^5 -pyridyl, NR^5 -pyrazinyl, $NR^5(CH_2)_{1-6}$ -pyridyl or $NR^5(CH_2)_{1-6}$ -pyrazinyl group and the other of R^{26} and R^{27} represents a hydrogen atom, and m , A , R^1 , R^2 and R^3 are as defined in formula (I), with a source of hydrogen and a hydrogenation catalyst; or

(xxxi) when X represents a group $CH_2O(CH_2)_{1-3}$ or $CH_2O(CH_2)_{2-3}O$ and A is $NHC(O)$, reacting a compound of general formula




(XXVII)

wherein one of R^{28} and R^{29} represents a group $-X''-R^4$ and the other of R^{28} and R^{29} represents a hydrogen atom, X'' represents a group $CH_2O(CH_2)_{1-3}$ or $CH_2O(CH_2)_{2-3}O$, L^7 represents a leaving group and R^2 , R^3 and R^4 are as defined in formula (I), with a compound of formula (VII) as defined in (iii) above, optionally in the presence of a coupling agent; or

(xxxii) when X represents a group $CH_2O(CH_2)_{1-3}$ or $CH_2O(CH_2)_{2-3}O$ and A is $C(O)NH$, reacting a compound of general formula





wherein R^2 and R^3 are as defined in formula (I) and R^{28} and R^{29} are as defined in formula (XXVII) in (xxxi) above, with a compound of formula (IX) as defined in (iv) above, in the presence of a base;

5

and optionally after (i), (ii), (iii), (iv), (v), (vi), (vii), (viii), (ix), (x), (xi), (xii), (xiii), (xiv), (xv), (xvi), (xvii), (xviii), (xix), (xx), (xxi), (xxii), (xxiii), (xxiv), (xxv), (xxvi), (xxvii), (xxviii), (xxix), (xxx), (xxxi) or (xxxii) converting the compound of formula (I) to a further compound of formula (I) and, if desired, forming a pharmaceutically acceptable salt or solvate of the compound of formula (I).

10

9. A pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

15

10. A process for the preparation of a pharmaceutical composition as claimed in claim 9 which comprises mixing a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as defined in any one of claims 1 to 7 with a pharmaceutically acceptable adjuvant, diluent or carrier.

20

11. A compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 for use in therapy.

25

12. Use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 in the manufacture of a medicament for use in the treatment of rheumatoid arthritis.



13. Use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7 in the manufacture of a medicament for use in the treatment of an obstructive airways disease.

5 14. Use according to claim 13, wherein the obstructive airways disease is asthma or chronic obstructive pulmonary disease.

15. A method of treating rheumatoid arthritis which comprises administering to a patient a therapeutically effective amount of a compound of formula (I), or a pharmaceutically
10 acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7.

16. A method of treating an obstructive airways disease which comprises administering to a patient a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 7.

15

