


WHAT IS CLAIMED IS:

1. A compound of formula (I):



wherein:

Q is F or H;

P¹ is hydrogen or a labile protecting group;

P² is an amino protecting group or hydrogen; and

L¹ is Br, Cl, I, R¹-SO₂O- or R²C(O)O-; wherein R¹ and R² are each independently optionally substituted aryl or optionally substituted C₁-₆alkyl.

2. The compound of claim 1, wherein P¹ is -H.

3. The compound of claim 1, wherein P¹ is selectively cleaved in the presence of P².

4. The compound of claim 3, wherein P¹ is 9-fluorenylmethoxycarbonyl, t-butoxycarbonyl, trimethylsilyl or t-butyldiphenylsilyl.

5. The compound of any of claims 1-4, wherein P² is an amino protecting group selected from R³-C(O)- or R⁴O-C(O)-, wherein R³ and R⁴ are each independently selected from C₁-₆alkyl, aryl, heteroaryl, aryl-C₁-₂alkyl, heteroaryl-C₁-₂alkyl, C₃-₁₀cycloalkyl, C₃-₁₀cycloalkyl-C₁-₂alkyl, ethynyl or vinyl, each of which is optionally substituted.

6. The compound of claim 5, wherein P² is -H.

7. The compound of claim 5, wherein P² is 2,6-dichlorophenylcarbonyl.

8. The compound of any of claims 5, wherein R³ or R⁴ is optionally substituted with 1-3 Rᵃ groups independently selected from halogen, C₁-₆alkyl, fluoro substituted C₁-₆alkyl, fluoro substituted C₁-₆alkoxy, aryl, heteroaryl, C₁-₆alkoxy, -CN, -NO₂, -OH, C₁-₆alkyl-OC(O)-, C₁-₆alkyl-C(O)O- or -SiMe₃, wherein the aliphatic or aromatic portion of Rᵃ is further optionally substituted with from 1-3 Rᵇ groups independently selected from halogen, C₁-₆alkyl, C₁-₆alkoxy, -CN, -NO₂ or -OH.

9. The compound of claim 8, wherein R^a is F, Cl, Br, I, $-CH_3$, Phenyl, t-butyl, MeO-, $-NO_2$, $-CN$, $-CF_3$, CF_3O- , $-OH$ or $-CH=CH_2$.

10. The compound of any of claims 1-9, wherein R^3 and R^4 are each independently methyl, ethyl, phenyl, 2,2,2-trichloroethyl, $(CH_3)_2CHC\equiv C-$, 2-trimethylsilylethyl, 1-methyl-1-phenylethyl, cyclobutyl, cyclopropyl, allyl, vinyl, 1-adamantyl, benzyl or diphenylmethyl, each of which is optionally substituted with from 1-3 R^a groups.


11. The compound of any of claims 1-10, wherein L^1 is Br, Cl, I, tosyl-O-, mesyl-O-, trifluoromethanesulfonyl-O-, R^1-SO_2O- or $R^2C(O)O-$, wherein R^1 and R^2 are each independently selected from aryl, aryl- C_{1-4} alkyl or C_{1-6} alkyl, each of which is optionally substituted with from 1-3 R^c substituents selected from halogen, $-CH=CH_2$, $-CN$, $-OH$, $-NH_2$, $-NO_2$, $-C(O)OH$, $-C(O)NH_2$, $-S(O)_2NH_2$, $-NHC(O)NH_2$, $-NHC(S)NH_2$, $-NHS(O)_2NH_2$, $-C(NH)NH_2$, $-OR^d$, $-SR^d$, $-OC(O)R^d$, $-C(O)R^d$, $-C(O)OR^d$, $-C(S)OR^d$, $-S(O)R^d$, $-S(O)_2R^d$, $-C(O)NHR^d$, $-C(O)NR^dR^d$, $-S(O)_2NHR^d$, $-S(O)_2NR^dR^d$, $-C(NH)NHR^d$, $-C(NH)NR^dR^d$, $-NHC(O)R^d$, $-NR^dC(O)R^d$, $-NHS(O)_2R^d$, $-NR^dS(O)_2R^d$, $-NHC(O)NHR^d$, $-NHR^d$ or $-NR^dR^d$, wherein each R^d is independently selected from C_{1-6} alkyl or aryl.

12. The compound of claim 11, wherein L^1 is Br, Cl, I, tosyl-O-, mesyl-O-, trifluoromethanesulfonyl-O-, $CF_3C(O)O-$ or $CH_3C(O)O-$.

13. The compound of claim 1, wherein P^1 is H; P^2 is 2,6-dichlorophenylcarbonyl; and L^1 is Br.


14. The compound of claim 1, wherein P^1 is H; P^2 is H; and L^1 is Br.

15. A method for preparing a compound of formula (Ia), said method comprising: contacting a compound of formula (II):



II

with an agent of the formula: P^2-X^1 under conditions sufficient to form the compound of formula (Ia);



(Ia); and

wherein:


X^1 is selected from Br, Cl, I, tosyl-O-, mesyl-O-, trifluoromethanesulfonyl-O-, $CF_3C(O)O$ - or $CH_3C(O)O$ -;

P^2 is an amino protecting group;

Q is H or F; and

L^1 is Br, Cl, I, R^1-SO_2O - or $R^2C(O)O$ -; wherein R^1 and R^2 are each independently optionally substituted aryl or optionally substituted C_{1-6} alkyl.


16. A method for preparing a compound of formula (I)



(I),


said method comprising:

contacting a compound of formula (II):



II

with an agent of the formula: P^2-X^1 under conditions sufficient to form the compound of formula (Ia):



(Ia); and

reacting a compound of formula (Ia) with an agent of formula: P^1-X^3 under conditions sufficient to form the compound of formula (I);

wherein:

X¹ is selected from Br, Cl, I, tosyl-O-, mesyl-O-, trifluoromethanesulfonyl-O-,
CF₃C(O)O- or CH₃C(O)O-;

X³ is a leaving group;

P¹ is a labile protecting group;

P² is an amino protecting group;

Q is H or F;

L¹ is Br, Cl, I, R¹-SO₂O- or R²C(O)O; wherein R¹ and R² are each independently
optionally substituted aryl or optionally substituted C₁₋₆alkyl.

17. The method of claim 15, wherein said contacting is carried out in the presence of triethylamine and 4-dimethylaminopyridine.


18. The method of claim 15 or 16, wherein L¹ is Br.

19. The method of claim 15 or 16, wherein P² is 2,6-dichlorophenyl; and L¹ is Br.

20. The method of claim 15 or 16, wherein Q is F.

What is claimed is:

1. A compound of formula (I):



(I)

wherein:

Q is F or H;

P¹ is H, 9-fluorenylmethoxycarbonyl, t-butoxycarbonyl, trimethylsilyl, or t-butyldiphenylsilyl;

P² is 2,6-dichlorophenylcarbonyl; and

L¹ is Br, Cl, I, R¹—SO₂O— or R²C(O)O—; wherein R¹ and R² are each optionally substituted aryl or optionally substituted C₁-₆alkyl.

2. The compound of claim 1, wherein P¹ is H.


3. The compound of claim 1, wherein L¹ is Br, Cl, I, tosyl-O—, mesyl-O—, trifluoromethanesulfonyl-O—, R¹—SO₂O— or R²C(O)O—, wherein R¹ and R² are each aryl, aryl-C₁-₄alkyl or C₁-₆alkyl, each of which is optionally substituted with from 1-3 Rᶜ substituents selected from halogen, —CH=CH₂, —CN, —OH, —NH₂, —NO₂, —C(O)OH, —C(O)NH₂, —S(O)₂NH₂, —NHC(O)NH₂, —NHC(S)NH₂, —NHS(O)₂NH₂, —C(NH)NH₂, —ORᵈ, —SRᵈ, —OC(O)Rᵈ, —C(O)Rᵈ, —C(O)ORᵈ, —C(S)ORᵈ, —S(O)Rᵈ, —S(O)₂Rᵈ, —C(O)NHRᵈ, —C(O)NRᵈRᵈ, —S(O)₂NHRᵈ, —S(O)₂NRᵈRᵈ, —C(NH)NHRᵈ, —C(NH)NRᵈRᵈ, —NHC(O)Rᵈ, —NRᵈC(O)Rᵈ, —NHS(O)₂Rᵈ, —NRᵈS(O)₂Rᵈ, —NHC(O)NHRᵈ, —NHRᵈ or —NRᵈRᵈ, wherein each Rᵈ is independently selected from C₁-₆alkyl or aryl.

4. The compound of claim 3, wherein L¹ is Br, Cl, I, tosyl-O—, mesyl-O—, trifluoromethanesulfonyl-O—, CF₃C(O)O— or CH₃C(O)O—.


5. The compound of claim 1, wherein P¹ is H and L¹ is Br.

6. A method for preparing a compound of formula (Ia), said method comprising:

contacting a compound of formula (II):



with an agent of formula: P^2-X^1 under conditions sufficient to form the compound of formula (Ia):



and

wherein:

X^1 is Br, Cl, I, tosyl-O—, mesyl-O—, trifluoromethanesulfonyl-O—, $CF_3C(O)O—$ or $CH_3C(O)O—$;


P^2 is 2,6-dichlorophenylcarbonyl;

Q is H or F; and

L^1 is Br, Cl, I, $R^1-SO_2O—$ or $R^2C(O)O—$; wherein R^1 and R^2 are each optionally substituted aryl or optionally substituted C_{1-6} alkyl.

7. A method for preparing a compound of formula (I)


(I)



said method comprising:


contacting a compound of formula (II):

(II)



with an agent of formula: P^2-X^1 under conditions sufficient to form the compound of formula (Ia):

(Ia)



and

reacting a compound of formula (Ia) with an agent of formula: P^1-X^3 under conditions sufficient to form the compound of formula (I);

wherein:

X^1 is Br, Cl, I, tosyl-O—, mesyl-O—, trifluoromethanesulfonyl-O—, $CF_3C(O)O$ — or $CH_3C(O)O$ —;

X^3 is a leaving group;

P^1 is H, 9-fluorenylmethoxycarbonyl, t-butoxycarbonyl, trimethylsilyl, or t-butyldiphenylsilyl;

P^2 is 2,6-dichlorophenylcarbonyl;

Q is H or F;

L^1 is Br, Cl, I, R^1-SO_2O — or $R^2C(O)O$; wherein R^1 and R^2 are each optionally substituted aryl or optionally substituted C_{1-6} alkyl.

8. The method of claim 6, wherein said contacting is carried out in the presence of triethylamine and 4-dimethylaminopyridine.

9. The method of claim 6 or 7, wherein L^1 is Br.

10. The method of claim 6 or 7, wherein Q is F.