cause lysis of host cells. In the polysaccharide capsule, the presence of complex polysaccharides forms the basis for dividing pneumococci into different serotypes. To date, close to 100 serotypes of *S. pneumoniae* have been identified.

[0006] Two vaccines for *S. pneumoniae* are currently available in the U.S.; PCV13 and PPSV23. PCV13 cannot confer protection against most of the known serotypes of *S. pneumoniae*. While PPSV23 includes polysaccharide components of more serotypes of *S. pneumoniae* than PCV13, it induces an immune response that is neither long-lasting nor anamnestic upon subsequent challenge. PPSV23 protects adults and the elderly against invasive pneumococcal disease; however, no consistent effect has been observed in the prevention of pneumonia [Gruber et al, 2008].

[0007] Thus, there is a medical need for a vaccine that provides T-cell dependent immunity against a broad range of serotypes of *S. pneumoniae.*

Summary

[0008] The present disclosure addresses the lack of suitable technologies for the prevention and/or treatment of pneumococcal infection. Among other things, the present disclosure addresses challenges in providing vaccines with sufficient immunogenicity to protect against invasive pneumococcal disease and pneumonia, by inducing a T- and B-cell response providing immunity against a broad range of *S. pneumoniae* serotypes including those serotypes not included in the vaccine.

[0009] Among other things, the present disclosure provides compositions and methods for prevention and/or treatment of pneumococcal infections in patient populations in need thereof.

[0010] In some examples, a vaccine comprises an immunogenic complex, wherein the immunogenic complex comprises: (a) a biotinylated polysaccharide antigen; and (b) a fusion protein comprising: (i) a biotin-binding moiety; and (ii) at least one polypeptide antigen; wherein the biotinylated polysaccharide antigen is non-covalently associated with the biotin-binding moiety of the fusion protein to form an immunogenic complex.

[0011] In some examples, the at least one polypeptide antigen comprises: an SP1500 polypeptide or antigenic fragment thereof; or an SP0785 polypeptide or antigenic fragment thereof. In some examples, the at least one polypeptide antigen comprises: (a) a first polypeptide antigen comprising an SP1500 polypeptide or antigenic fragment thereof; and (b) a second polypeptide antigen comprising an SP0785 polypeptide or antigenic fragment thereof.

[0011a] In one aspect, the present disclosure refers to a multivalent vaccine comprising fifteen or more species of immunogenic complexes, wherein each species of the immunogenic complexes comprises: (a)a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from a distinct *Streptococcus pneumoniae* serotype; and (b)a fusion protein comprising: (i) a biotin-binding moiety; and (ii)at least one polypeptide antigen comprising: a first polypeptide antigen comprising an SP1500 polypeptide or an antigenic fragment thereof; and a second polypeptide antigen comprising an SP0785 polypeptide or an antigenic fragment thereof; wherein the fusion protein comprises an amino acid sequence at least 90% identical to SEQ ID NO:6; and wherein in each species of the immunogenic complexes, the biotinylated polysaccharide antigen is non-covalently associated with the biotin-binding moiety of the fusion protein.

[0012] In some examples, the biotinylated polysaccharide antigen comprises a polysaccharide of *Streptococcus pneumoniae.* In some examples, the biotinylated polysaccharide antigen comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 1, 2, 3, 4, 5, 6A, 6B, 6C, 6D, 6E, 6F, 6G, 6H, 7A, 7B, 7C, 7F, 8, 9A, 9L, 9N, 9V, 10A, 10B, 10C, 10F, 11A, 11B, 11C, 11D, 11E, 11F, 12A, 12B, 12F, 13, 14, 15A, 15B, 15C, 15F, 16A, 16F, 17A, 17F, 18A, 18B, 18C, 18F, 19A, 19B, 19C, 19F, 20A, 20B, 21, 22A, 22F, 23A, 23B, 23F, 24A, 24B, 24F, 25A, 25F, 27, 28A, 28F, 29, 31, 32A, 32F, 33A, 33B, 33C, 33D, 33E, 33F, 34, 35A, 35B, 35C, 35F, 36, 37, 38, 39, 40, 41A, 41F, 42, 43, 44, 45, 46, 47A, 47F, and 48.

[0013] In some examples, the biotinylated polysaccharide antigen comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20B, 22F, 23F, and 33F.

[0014] In some examples, the biotinylated polysaccharide antigen comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 22F, 23F, and 33F.

[0015] In some examples, the biotinylated polysaccharide antigen comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 2, 8, 9N, 10A, 11A, 12F, 15B, 17F, and 20B.

[0016] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising:(a) a plurality of biotinylated polysaccharide antigens, wherein the plurality comprises polysaccharide antigens of one or more of *Streptococcus pneumoniae* serotypes 1, 2, 3, 4, 5, 6A, 6B, 6C, 6D, 6E, 6F, 6G, 6H, 7A, 7B, 7C, 7F, 8, 9A, 9L, 9N, 9V, 10A, 10B, 10C, 10F, 11A, 11B, 11C, 11D, 11E, 11F, 12A, 12B, 12F, 13, 14, 15A, 15B, 15C, 15F,

16A, 16F, 17A, 17F, 18A, 18B, 18C, 18F, 19A, 19B, 19C, 19F, 20A, 20B, 21, 22A, 22F, 23A, 23B, 23F, 24A, 24B, 24F, 25A, 25F, 27, 28A, 28F, 29, 31, 32A, 32F, 33A, 33B, 33C, 33D, 33E, 33F, 34, 35A, 35B, 35C, 35F, 36, 37, 38, 39, 40, 41A, 41F, 42, 43, 44, 45, 46, 47A, 47F, and 48; and (b) a plurality of fusion proteins, each fusion protein comprising (i) a biotin-binding moiety; (ii) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (iii) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof; wherein each of the first plurality of biotinylated polysaccharide antigens is non-covalently associated with the biotin-binding moiety of one or more of the plurality of fusion proteins to form an immunogenic complex.

[0017] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising: (a) a plurality of biotinylated polysaccharide antigens, wherein the plurality comprises polysaccharide antigens of one or more of *Streptococcus pneumoniae* serotypes 1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20B, 22F, 23F, and 33F; and (b) a plurality of fusion proteins, each fusion protein comprising (i) a biotin-binding moiety; (ii) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (iii) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof; wherein each of the first plurality of biotinylated polysaccharide antigens is non-covalently associated with the biotin-binding moiety of one or more of the plurality of fusion proteins to form an immunogenic complex.

[0018] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising: (a) a plurality of biotinylated polysaccharide antigens, wherein the plurality comprises polysaccharide antigens of each of *Streptococcus pneumoniae* serotypes 1, 2,

3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20B, 22F, 23F, and 33F; and (b) a plurality of fusion proteins, each fusion protein comprising (i) a biotin-binding moiety; (ii) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (iii) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof; wherein each of the first plurality of biotinylated polysaccharide antigens is non-covalently associated with the biotin-binding moiety of one or more of the plurality of fusion proteins to form an immunogenic complex.

[0019] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising: (a) a plurality of biotinylated polysaccharide antigens, wherein the plurality comprises polysaccharide antigens of one or more of *Streptococcus pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 22F, 23F, and 33F; and (b) a plurality of fusion proteins, each fusion protein comprising (i) a biotin-binding moiety; (ii) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (iii) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof; wherein each of the first plurality of biotinylated polysaccharide antigens is non-covalently associated with the biotin-binding moiety of one or more of the plurality of fusion proteins to form an immunogenic complex.

[0020] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising: (a) a plurality of biotinylated polysaccharide antigens, wherein the plurality comprises polysaccharide antigens of each of *Streptococcus pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 22F, 23F, and 33F; and (b) a plurality of fusion proteins, each fusion protein comprising (i) a biotin-binding moiety; (ii) a first polypeptide

antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (iii) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof; wherein each of the first plurality of biotinylated polysaccharide antigens is non-covalently associated with the biotin-binding moiety of one or more of the plurality of fusion proteins to form an immunogenic complex.

[0021] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising: (a) a plurality of biotinylated polysaccharide antigens, wherein the plurality comprises polysaccharide antigens of one or more of *Streptococcus pneumoniae* serotypes 2, 8, 9N, 10A, 11A, 12F, 15B, 17F, and 20B; and (b) a plurality of fusion proteins, each fusion protein comprising (i) a biotin-binding moiety; (ii) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (iii) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof; wherein each of the first plurality of biotinylated polysaccharide antigens is non-covalently associated with the biotin-binding moiety of one or more of the plurality of fusion proteins to form an immunogenic complex.

[0022] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising: (a) a plurality of biotinylated polysaccharide antigens, wherein the plurality comprises polysaccharide antigens of each of *Streptococcus pneumoniae* serotypes 2, 8, 9N, 10A, 11A, 12F, 15B, 17F, and 20B; and (b) a plurality of fusion proteins, each fusion protein comprising (i) a biotin-binding moiety; (ii) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5

or an antigenic fragment thereof; and (iii) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof; wherein each of the first plurality of biotinylated polysaccharide antigens is non-covalently associated with the biotin-binding moiety of one or more of the plurality of fusion proteins to form an immunogenic complex.

[0023] In some examples, a vaccine comprises a biotin-binding moiety, wherein the biotin-binding moiety is a polypeptide that has or comprises an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:1 or a biotin-binding fragment thereof; or a polypeptide that has or comprises an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:2 or a biotin-binding fragment thereof. In some examples, a vaccine comprises a fusion protein that is or comprises an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:6. In some examples, a vaccine comprises the fusion protein CP1.

[0024] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising one or more species of immunogenic complexes, wherein a species refers to complexes comprising a polysaccharide antigen of a single *Streptococcus pneumoniae* serotype. In some examples, a vaccine comprises a plurality of immunogenic complexes comprising two or more species of immunogenic complexes, wherein a species refers to complexes comprising a polysaccharide antigen of a single *Streptococcus pneumoniae* serotype. In some examples, a vaccine comprises a plurality of immunogenic complexes comprising nine or more species of immunogenic complexes, wherein a species refers to complexes comprising a polysaccharide antigen of a single *Streptococcus pneumoniae* serotype.

[0025] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising fifteen or more species of immunogenic complexes, wherein a species refers to complexes comprising a polysaccharide antigen of a single *Streptococcus pneumoniae* serotype.

[0026] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising twenty-four or more species of immunogenic complexes, wherein a species refers to complexes comprising a polysaccharide antigen of a single *Streptococcus pneumoniae* serotype.

[0027] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising: a first species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 1 non-covalently complexed with a fusion protein; a second species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 2 non-covalently complexed with a fusion protein; a third species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 3 non-covalently complexed with a fusion protein; a fourth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 4 non-covalently complexed with a fusion protein; a fifth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 5 non-covalently complexed with a fusion protein; a sixth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 6A non-covalently complexed with a fusion protein; a seventh species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 6B non-covalently complexed with a fusion protein; an eighth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 7F non-covalently complexed with a fusion protein; a ninth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 8 non-covalently complexed with a fusion protein; a tenth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 9N non-covalently complexed with a fusion protein; an eleventh species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 9V non-covalently complexed with a fusion protein; a twelfth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 10A non-covalently complexed with a fusion protein; a thirteenth species of immunogenic complex comprising a biotinylated

polysaccharide antigen of *Streptococcus pneumoniae* serotype 11A non-covalently complexed with a fusion protein; a fourteenth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 12F non-covalently complexed with a fusion protein; a fifteenth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 14 non-covalently complexed with a fusion protein; a sixteenth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 15B non-covalently complexed with a fusion protein; a seventeenth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 17F non-covalently complexed with a fusion protein; an eighteenth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 18C non-covalently complexed with a fusion protein; a nineteenth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 19A non-covalently complexed with a fusion protein; a twentieth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 19F non-covalently complexed with a fusion protein; a twenty-first species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 20B non-covalently complexed with a fusion protein; a twenty-second species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 22F non-covalently complexed with a fusion protein; a twenty-third species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 23F non-covalently complexed with a fusion protein; a twenty-fourth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 33F non-covalently complexed with a fusion protein; wherein each fusion protein comprises (a) a biotin-binding moiety;(b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof;

and wherein each of the first through twenty-fourth biotinylated polysaccharide antigens is non-covalently associated with the biotin-binding moiety of at least one fusion protein to form immunogenic complexes.

[0028] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising: a first species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 1 non-covalently complexed with a fusion protein; a second species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 3 non-covalently complexed with a fusion protein; a third species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 4 non-covalently complexed with a fusion protein; a fourth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 5 non-covalently complexed with a fusion protein; a fifth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 6A non-covalently complexed with a fusion protein; a sixth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 6B non-covalently complexed with a fusion protein; a seventh species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 7F non-covalently complexed with a fusion protein; an eighth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 9V non-covalently complexed with a fusion protein; a ninth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 14 non-covalently complexed with a fusion protein; a tenth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 18C non-covalently complexed with a fusion protein; an eleventh species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 19A non-covalently complexed with a fusion protein; a twelfth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 19F non-covalently complexed with a fusion protein; a thirteenth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 22F non-covalently complexed

with a fusion protein; a fourteenth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 23F non-covalently complexed with a fusion protein; a fifteenth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 33F non-covalently complexed with a fusion protein; wherein each fusion protein comprises (a) a biotin-binding moiety;(b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof; and wherein each of the first through fifteenth biotinylated polysaccharide antigens is non-covalently associated with the biotin-binding moiety of at least one fusion protein to form immunogenic complexes.

[0029] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising: a first species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 2 non-covalently complexed with a fusion protein; a second species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 8 non-covalently complexed with a fusion protein; a third species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 9N non-covalently complexed with a fusion protein; a fourth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 10A non-covalently complexed with a fusion protein; a fifth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 11A non-covalently complexed with a fusion protein; a sixth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 12F non-covalently complexed with a biotin-binding fusion protein; a seventh species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 15B non-covalently complexed with a fusion protein; an eighth species of immunogenic complex comprising a

biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 17F non-covalently complexed with a fusion protein; and a ninth species of immunogenic complex comprising a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 20B non-covalently complexed with a fusion protein; wherein each fusion protein comprises (a) a biotin-binding moiety;(b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof; and wherein each of the first through ninth biotinylated polysaccharide antigens is non-covalently associated with the biotin-binding moiety of at least one fusion protein to form immunogenic complexes.

[0030] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising two or more species of immunogenic complexes, wherein each species of immunogenic complexes contributes a stoichiometrically equal ratio, by weight, of the polysaccharide antigen.

[0031] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising two or more species of immunogenic complexes, wherein at least one species of immunogenic complex contributes a stoichiometrically different ratio, by weight, of the polysaccharide antigen.

[0032] In some examples, a vaccine comprises a plurality of immunogenic complexes comprising two or more species of immunogenic complexes, wherein different species of immunogenic complexes contribute a stoichiometrically different ratio, by weight, of the polysaccharide antigen.

[0033] In some examples, a vaccine comprises a fusion protein comprising a biotin-binding moiety, wherein the biotin-binding moiety is a polypeptide comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:1

or a biotin-binding fragment thereof; or a polypeptide comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:2 or a biotin-binding fragment thereof.

[0034] In some examples, an immunogenic complex comprises a fusion protein comprising a biotin-binding moiety, wherein the biotin-binding moiety is a polypeptide comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:1 or a biotin-binding fragment thereof; or a polypeptide comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:2 or a biotin-binding fragment thereof.

[0035] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* non-covalently associated with a fusion protein, wherein the fusion protein comprises a biotin-binding moiety and at least one polypeptide antigen.

[0036] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen, wherein the biotinylated polysaccharide antigen comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 1, 2, 3, 4, 5, 6A, 6B, 6C, 6D, 6E, 6F, 6G, 6H, 7A, 7B, 7C, 7F, 8, 9A, 9L, 9N, 9V, 10A, 10B, 10C, 10F, 11A, 11B, 11C, 11D, 11E, 11F, 12A, 12B, 12F, 13, 14, 15A, 15B, 15C, 15F, 16A, 16F, 17A, 17F, 18A, 18B, 18C, 18F, 19A, 19B, 19C, 19F, 20A, 20B, 21, 22A, 22F, 23A, 23B, 23F, 24A, 24B, 24F, 25A, 25F, 27, 28A, 28F, 29, 31, 32A, 32F, 33A, 33B, 33C, 33D, 33E, 33F, 34, 35A, 35B, 35C, 35F, 36, 37, 38, 39, 40, 41A, 41F, 42, 43, 44, 45, 46, 47A, 47F, and 48.

[0037] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen, wherein the biotinylated polysaccharide antigen comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20B, 22F, 23F, and 33F.

[0038] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen, wherein the biotinylated polysaccharide antigen comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 22F, 23F, and 33F.

[0039] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen, wherein the biotinylated polysaccharide antigen comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 2, 8, 9N, 10A, 11A, 12F, 15B, 17F, and 20B.

[0040] In some examples, an immunogenic complex comprises at least one peptide antigen, wherein the at least one polypeptide antigen comprises an SP1500 polypeptide or antigenic fragment thereof; an SP0785 polypeptide or antigenic fragment thereof; or a combination thereof.

[0041] In some examples, an immunogenic complex comprises at least one peptide antigen, wherein the at least one polypeptide antigen comprises: (a) a first polypeptide antigen comprising an SP1500 polypeptide or antigenic fragment thereof; and (b) a second polypeptide antigen comprising an SP0785 polypeptide or antigenic fragment thereof.

[0042] In some examples, an immunogenic complex comprises at least one peptide antigen, wherein the at least one polypeptide antigen comprises: (a) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (b) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0043] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 1 non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or

at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0044] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 2 non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0045] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 3 non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0046] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 4 non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%,

at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0047] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 5 non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0048] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 6A non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0049] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 6B non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0050] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 7F non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0051] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 8 non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0052] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 9N non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0053] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 9V non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first

polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0054] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 10A non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0055] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 11A non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0056] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 12F non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second

polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0057] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 14 non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0058] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 15B non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0059] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 17F non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0060] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 18C non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0061] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 19A non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0062] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 19F non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0063] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 20B non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first

polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0064] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 22F non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0065] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 23F non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0066] In some examples, an immunogenic complex comprises a biotinylated polysaccharide antigen of *Streptococcus pneumoniae* serotype 33F non-covalently associated with a fusion protein, wherein the fusion protein comprises (a) a biotin-binding moiety; (b) a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and (c) a second

polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

[0067] In some examples, an immunogenic complex comprises a biotin-binding moiety, wherein the biotin-binding moiety is a polypeptide comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:1 or a biotin-binding fragment thereof; or a polypeptide comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:2 or a biotin-binding fragment thereof.

[0068] In some examples, an immunogenic complex comprises a fusion protein and a polysaccharide antigen, wherein the ratio of fusion protein to polysaccharide antigen in the complex is about 1:1, about 2:1, about 3:1, about 4:1, about 5:1, about 6:1, about 7:1, about 8:1, about 9:1, or about 10:1 by weight. In some examples, a vaccine comprises one or more immunogenic complexes disclosed herein.

[0069] In some examples, a pharmaceutical composition comprises a vaccine or vaccine composition disclosed herein, and a pharmaceutically acceptable carrier. In some examples, a pharmaceutical composition comprises an immunogenic complex disclosed herein, and a pharmaceutically acceptable carrier. In some examples, a pharmaceutical composition disclosed herein comprises one or more adjuvants. In some examples, the one or more adjuvants is or comprises a co-stimulation factor.

[0069a] In one aspect, the present disclosure refers to a pharmaceutical composition comprising the vaccine disclosed herein, and a pharmaceutically acceptable carrier.

[0070] In some examples, the one or more adjuvants are selected from the group consisting of aluminum phosphate, aluminum hydroxide, and phosphated aluminum hydroxide. In some examples, wherein the one or more adjuvants is or comprises aluminum phosphate. In some examples, a pharmaceutical composition disclosed herein is formulated for injection.

[0071] In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein induces an immune response. In some examples, the immune

response comprises an innate immune response. In some examples, the immune response comprises an antibody or B cell response. In some examples, the immune response comprises a CD4+ T cell response (*e.g.*, T_H1, T_H2, or T_H17 response), a CD8+ T cell response, a CD4+ and CD8+ T cell response, or a CD4-/CD8- T cell response. In some examples, the immune response comprises (i) an antibody or B cell response, and (ii) a T cell response. In some examples, the immune response comprises (i) an antibody or B cell response, (ii) a T cell response, and (iii) an innate immune response. In some examples, the immune response is to at least one polysaccharide antigen or at least one polypeptide antigen of a fusion protein. In some examples, the immune response comprises (i) an antibody or B cell response to at least one polysaccharide antigen, and (ii) a CD4+ T cell response (*e.g.*, T_H1, T_H2, or T_H17 response), a CD8+ T cell response, a CD4+ and CD8+ T cell response, or a CD4-/CD8- T cell response to at least one polypeptide of a fusion protein. In some examples, the immune response comprises (i) an antibody or B cell response to at least one polysaccharide antigen, and (ii) an antibody or B cell response to at least one polypeptide of a fusion protein. In some examples, the immune response comprises (i) an antibody or B cell response to at least one polysaccharide antigen, and (ii) an antibody or B cell response, and a CD4+ T cell response, including a T_H1, T_H2, or T_H17 response, a CD8+ T cell response, a CD4+ and CD8+ T cell response, or a CD4-/CD8- T cell response, to at least one polypeptide of a fusion protein.

[0072] In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein induces an opsonic/bactericidal response against one or more serotypes of *Streptococcus pneumoniae.* In some examples, such an opsonic/bactericidal response may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such an opsonic/bactericidal response may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such an opsonic/bactericidal response may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus*

pneumoniae, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, an opsonic/bactericidal response can include production of antibodies that opsonize one or more serotypes of *Streptococcus pneumoniae*, which leads to killing of such serotype(s) by one or more immune pathways. In some examples, such an opsonic/bactericidal response can be determined by an opsonophagocytic assay (OPA) or a concentrated opsonophagocytic assay (COPA) known in the art and/or as described herein. In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein inhibits transmission of one or more serotypes of *Streptococcus pneumoniae* from the subject to another subject. In some examples, such inhibition of transmission may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of transmission may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of transmission may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0073] In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein inhibits or reduces the rate of occurrence of Invasive Pneumonoccal Disease (IPD) associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition of IPD or reduction in the rate of occurrence of IPD may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described

herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of IPD or reduction in the rate of occurrence of IPD may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of IPD or reduction in the rate of occurrence of IPD may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein reduces the severity of Invasive Pneumonoccal Disease (IPD) associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such reduction in the severity of IPD may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of IPD may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of IPD may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0074] In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein inhibits or reduces the rate of occurrence of bacteremia, sepsis, and/or meningitis associated with or induced by one or more serotypes of *Streptococcus*

pneumoniae. In some examples, such inhibition, or reduction in the rate of occurrence, of bacteremia, sepsis, and/or meningitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of bacteremia, sepsis, and/or meningitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of bacteremia, sepsis, and/or meningitis may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein reduces the severity of bacteremia, sepsis, and/or meningitis associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such reduction in the severity of bacteremia, sepsis, and/or meningitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of bacteremia, sepsis, and/or meningitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of bacteremia, sepsis, and/or meningitis may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include

polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0075] In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein inhibits or reduces the rate of occurrence of organ damage associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition, or reduction in the rate of occurrence, of organ damage may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of organ damage may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of organ damage may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein reduces, the severity of organ damage associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such reduction in the severity of organ damage may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of organ damage may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of organ damage may be

directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0076] In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein inhibits or reduces the rate of occurrence of pneumonia associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition, or reduction in the rate of occurrence, of pneumonia may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of pneumonia may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of pneumonia may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein reduces the severity of pneumonia associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such reduction in the severity of pneumonia may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of pneumonia may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a

pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of pneumonia may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0077] In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein inhibits or reduces the rate of occurrence of otitis media associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition, or reduction in the rate of occurrence, of otitis media may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of otitis media may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of otitis media may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein reduces the severity of otitis media associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such reduction in the severity of otitis media may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s)

present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of otitis media may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of otitis media may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0078] In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein inhibits or reduces the rate of occurrence of sinusitis associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition, or reduction in the rate of occurrence, of sinusitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of sinusitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of sinusitis may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein reduces the severity of sinusitis associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some

examples, such reduction in the severity of sinusitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of sinusitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of sinusitis may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0079] In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein inhibits colonization of mucosal surface(s) by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition of colonization of mucosal surface(s) may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of colonization of mucosal surface(s) may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of colonization of mucosal surface(s) may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, upon administration to a subject, a pharmaceutical composition disclosed herein inhibits colonization

of the nasopharynx by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition of colonization of nasopharynx may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of colonization of nasopharynx may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of colonization of nasopharynx may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0080] Some examples provide a method of making a vaccine, the method comprising non-covalently complexing a plurality of biotinylated polysaccharide antigens with a plurality of a fusion protein comprising at least one polypeptide antigen selected from: an SP1500 polypeptide or antigenic fragment thereof; an SP0785 polypeptide or antigenic fragment thereof; or a combination thereof. In some examples, the plurality of biotinylated polysaccharide antigens comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 1, 2, 3, 4, 5, 6A, 6B, 6C, 6D, 6E, 6F, 6G, 6H, 7A, 7B, 7C, 7F, 8, 9A, 9L, 9N, 9V, 10A, 10B, 10C, 10F, 11A, 11B, 11C, 11D, 11E, 11F, 12A, 12B, 12F, 13, 14, 15A, 15B, 15C, 15F, 16A, 16F, 17A, 17F, 18A, 18B, 18C, 18F, 19A, 19B, 19C, 19F, 20A, 20B, 21, 22A, 22F, 23A, 23B, 23F, 24A, 24B, 24F, 25A, 25F, 27, 28A, 28F, 29, 31, 32A, 32F, 33A, 33B, 33C, 33D, 33E, 33F, 34, 35A, 35B, 35C, 35F, 36, 37, 38, 39, 40, 41A, 41F, 42, 43, 44, 45, 46, 47A, 47F, and 48. In some examples, the plurality of biotinylated polysaccharide antigens comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20B, 22F, 23F, and 33F. In some examples, the plurality of biotinylated

polysaccharide antigens comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from each of 1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20B, 22F, 23F, and 33F. In some examples, the plurality of biotinylated polysaccharide antigens comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 22F, 23F, and 33F. In some examples, the plurality of biotinylated polysaccharide antigens comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from each of 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 22F, 23F, and 33F. In some examples, the plurality of biotinylated polysaccharide antigens comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from one or more of 2, 8, 9N, 10A, 11A, 12F, 15B, 17F, and 20B. In some examples, the plurality of biotinylated polysaccharide antigens comprises a polysaccharide of *Streptococcus pneumoniae* having a serotype selected from each of 2, 8, 9N, 10A, 11A, 12F, 15B, 17F, and 20B.

[0080a] In one aspect, the present disclosure refers to a method of making a multivalent vaccine, comprising mixing fifteen or more species of immunogenic complexes in a single formulation, wherein each species of the immunogenic complexes comprises: (a)a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from a distinct *Streptococcus pneumoniae* serotype; and (b)a fusion protein comprising: (i) a biotin-binding moiety; and (ii)at least one polypeptide antigen comprising: a first polypeptide antigen comprising an SP1500 polypeptide or an antigenic fragment thereof; and a second polypeptide antigen comprising an SP0785 polypeptide or an antigenic fragment thereof; wherein the fusion protein comprises an amino acid sequence at least 90% identical to SEQ ID NO:6; and wherein in each species of the immunogenic complexes, the biotinylated polysaccharide antigen is non-covalently associated with the biotin-binding moiety of the fusion protein.

[0081] Some examples provide for a method of immunizing a subject against *Streptococcus pneumoniae* infection and/or colonization, the method comprising administering to the subject an immunologically effective amount of a vaccine disclosed herein. In some examples, a method of immunizing a subject against *Streptococcus pneumoniae* infection and/or colonization, comprises administering to a subject an immunologically effective amount of an immunogenic complex disclosed herein. In some examples, a method of immunizing a subject against *Streptococcus pneumoniae* infection and/or colonization, comprises administering to a subject an immunologically effective amount of a pharmaceutical composition disclosed herein.

[0081a] In one aspect, the present disclosure refers to use of the multivalent vaccine disclosed herein or the pharmaceutical composition disclosed herein, in the manufacture of a medicament for immunizing a subject against *Streptococcus pneumoniae* infection and/or colonization.

[0082] In some examples, administration of the vaccine, immunogenic complex, or pharmaceutical composition induces an immune response. In some examples, the immune response comprises an innate immune response. In some examples, the immune response comprises an antibody or B cell response. In some examples, the immune response comprises a CD4+ T cell response (*e.g.*, T_H1, T_H2, or T_H17 response), a CD8+ T cell response, a CD4+ and CD8+ T cell response, or a CD4-/CD8- T cell response. In some examples, the immune response comprises (i) an antibody or B cell response and (ii) a T cell response. In some examples, the immune response comprises (i) an antibody or B cell response, (ii) a T

cell response, and (iii) an innate immune response. In some examples, the immune response is to at least one polysaccharide antigen or at least one polypeptide of a fusion protein. In some examples, the immune response comprises (i) an antibody or B cell response to at least one polysaccharide antigen, and (ii) a CD4+ T cell response (*e.g.*, T_H1, T_H2, or T_H17 response), a CD8+ T cell response, a CD4+ and CD8+ T cell response, or a CD4-/CD8- T cell response to at least one polypeptide of a fusion protein. In some examples, the immune response comprises (i) an antibody or B cell response to at least one polysaccharide antigen, and (ii) an antibody or B cell response to at least one polypeptide of a fusion protein. In some examples, the immune response comprises (i) an antibody or B cell response to at least one polysaccharide antigen, and (ii) an antibody or B cell response, and a CD4+ T cell response, including T_H1, T_H2, or T_H17 response, or a CD8+ T cell response, a CD4+ and CD8+ T cell response, or a CD4-/CD8- T cell response, to at least one polypeptide of a fusion protein.

[0083] In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) induces an opsonic/bactericidal response against one or more serotypes of *Streptococcus pneumoniae.* In some examples, such an opsonic/bactericidal response may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such an opsonic/bactericidal response may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such an opsonic/bactericidal response may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, administration of a vaccine, immunogenic complex,

or pharmaceutical composition (*e.g.*, ones as described and/or utilized herein) inhibits transmission of one or more serotypes of *Streptococcus pneumoniae* from the subject to another subject. In some examples, such inhibition of transmission may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of transmission may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of transmission may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0084] In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) inhibits or reduces the rate of occurrence of Invasive Pneumococcal Disease (IPD) associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition, or reduction in the rate of occurrence, of IPD may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of IPD may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of IPD may be directed against two or more

(*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) reduces the severity of Invasive Pneumococcal Disease (IPD) associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such reduction in the severity of IPD may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of IPD may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of IPD may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0085] In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) inhibits or reduces the rate of occurrence of bacteremia, sepsis, and/or meningitis associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition, or reduction in rate of occurrence, of bacteremia, sepsis, and/or meningitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or

more) of such serotype(s). In some examples, such inhibition, or reduction in rate of occurrence, of bacteremia, sepsis, and/or meningitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in rate of occurrence, of bacteremia, sepsis, and/or meningitis may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) reduces the severity of bacteremia, sepsis, and/or meningitis associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such reduction in the severity of bacteremia, sepsis, and/or meningitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of bacteremia, sepsis, and/or meningitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of bacteremia, sepsis, and/or meningitis may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0086] In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) inhibits or reduces the rate of occurrence of organ damage associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition, or reduction in the rate of occurrence, of organ damage may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of organ damage may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of organ damage may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) reduces the severity of organ damage associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such reduction in the severity of organ damage may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of organ damage may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some

examples, such reduction in the severity of organ damage may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0087] In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) inhibits or reduces the rate of occurrence of pneumonia associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition, or reduction in the rate of occurrence, of pneumonia may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of pneumonia may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of pneumonia may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) reduces the severity of pneumonia associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such reduction in the severity of pneumonia may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition

described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of pneumonia may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of pneumonia may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0088] In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) inhibits or reduces the rate of occurrence of otitis media associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition, or reduction in the rate of occurrence, of otitis media may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of otitis media may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of otitis media may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more)

of such serotype(s). In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) reduces the severity of otitis media associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such reduction in the severity of otitis media may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of otitis media may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of otitis media may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0089] In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) inhibits or reduces the rate of occurrence of sinusitis associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition, or reduction in the rate of occurrence, of sinusitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of sinusitis be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more)

of such serotype(s). In some examples, such inhibition, or reduction in the rate of occurrence, of sinusitis may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) reduces the severity of sinusitis associated with or induced by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such reduction in the severity of sinusitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of sinusitis may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such reduction in the severity of sinusitis may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0090] In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) inhibits colonization of mucosal surface(s) by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition of colonization of mucosal surface(s) may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein

includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of colonization of mucosal surface(s) may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of colonization of mucosal surface(s) may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, administration of a vaccine, immunogenic complex, or pharmaceutical composition (*e.g.*, ones described and/or utilized herein) inhibits colonization of nasopharynx by one or more serotypes of *Streptococcus pneumoniae*. In some examples, such inhibition of colonization of nasopharynx may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of colonization of nasopharynx may be directed against one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s). In some examples, such inhibition of colonization of nasopharynx may be directed against two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) serotypes of *Streptococcus pneumoniae*, wherein a vaccine, immunogenic complex, or pharmaceutical composition described herein (i) includes polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s); and (ii) does not include polysaccharide(s) present in at least one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) of such serotype(s).

[0091] In some examples of the methods disclosed herein, the *Streptococcus pneumoniae* has a serotype selected from one or more of 1, 2, 3, 4, 5, 6A, 6B, 6C, 6D, 6E, 6F, 6G, 6H, 7A, 7B, 7C, 7F, 8, 9A, 9L, 9N, 9V, 10A, 10B, 10C, 10F, 11A, 11B, 11C, 11D, 11E, 11F, 12A, 12B, 12F, 13, 14, 15A, 15B, 15C, 15F, 16A, 16F, 17A, 17F, 18A, 18B, 18C, 18F, 19A, 19B, 19C, 19F, 20A, 20B, 21, 22A, 22F, 23A, 23B, 23F, 24A, 24B, 24F, 25A, 25F, 27, 28A, 28F, 29, 31, 32A, 32F, 33A, 33B, 33C, 33D, 33E, 33F, 34, 35A, 35B, 35C, 35F, 36, 37, 38, 39, 40, 41A, 41F, 42, 43, 44, 45, 46, 47A, 47F, and 48. In some examples, the *Streptococcus pneumoniae* has a serotype selected from one or more of 1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20B, 22F, 23F, and 33F. In some examples, the *Streptococcus pneumoniae* has a serotype selected from one or more of 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 22F, 23F, and 33F. In some examples, the *Streptococcus pneumoniae* has a serotype selected from one or more of 2, 8, 9N, 10A, 11A, 12F, 15B, 17F, and 20B.

[0092] In some examples, the subject is immunized against *Streptococcus pneumoniae* infection and/or colonization with one dose of a vaccine. In some examples, the subject is immunized against *Streptococcus pneumoniae* infection and/or colonization with two doses of a vaccine. In some examples, the subject is immunized against *Streptococcus pneumoniae* infection and/or colonization with three doses of a vaccine.

Brief Description of the Drawings

[0093] The present teachings described herein will be more fully understood from the following description of various illustrative embodiments, when read together with the accompanying drawings. It should be understood that the drawings described below are for illustration purposes only and are not intended to limit the scope of the present teachings in any way.

[0094] Figure 1 is a schematic representation of an exemplary Multiple Antigen Presenting System (MAPS). In the exemplary embodiment shown, MAPS immunogenic complexes comprise one or more polypeptide antigens fused to the biotin-binding protein rhizavidin, or a biotin-binding domain or biotin-binding fragment thereof, and a biotinylated

Claims

We claim:

1. A multivalent vaccine comprising fifteen or more species of immunogenic complexes, wherein each species of the immunogenic complexes comprises:

(a)a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from a distinct *Streptococcus pneumoniae* serotype; and

(b)a fusion protein comprising:

(i) a biotin-binding moiety; and

(ii)at least one polypeptide antigen comprising:

a first polypeptide antigen comprising an SP1500 polypeptide or an antigenic fragment thereof; and

a second polypeptide antigen comprising an SP0785 polypeptide or an antigenic fragment thereof;

wherein the fusion protein comprises an amino acid sequence at least 90% identical to SEQ ID NO:6; and

wherein in each species of the immunogenic complexes, the biotinylated polysaccharide antigen is non-covalently associated with the biotin-binding moiety of the fusion protein.

2. The vaccine of claim 1, wherein the *Streptococcus pneumoniae* serotype for each species of the immunogenic complexes is selected from 1, 2, 3, 4, 5, 6A, 6B, 6C, 6D, 6E, 6F, 6G, 6H, 7A, 7B, 7C, 7F, 8, 9A, 9L, 9N, 9V, 10A, 10B, 10C, 10F, 11A, 11B, 11C, 11D, 11E, 11F, 12A, 12B, 12F, 13, 14, 15A, 15B, 15C, 15F, 16A, 16F, 17A, 17F, 18A, 18B, 18C, 18F, 19A, 19B, 19C, 19F, 20A, 20B, 21, 22A, 22F, 23A, 23B, 23F, 24A, 24B, 24F, 25A, 25F, 27, 28A, 28F, 29, 31, 32A, 32F, 33A, 33B, 33C, 33D, 33E, 33F, 34, 35A, 35B, 35C, 35F, 36, 37, 38, 39, 40, 41A, 41F, 42, 43, 44, 45, 46, 47A, 47F, and 48.

3. The vaccine of claim 1, wherein the *Streptococcus pneumoniae* serotype for each species of the immunogenic complexes is selected from 1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20B, 22F, 23F, and 33F.

4. The vaccine of claim 1, wherein the first polypeptide antigen comprises an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof.

5. The vaccine of claim 1, wherein the second polypeptide antigen comprises an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof.

6. The vaccine of claim 1, wherein the fusion protein comprises an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:6.

7. The vaccine of claim 1, wherein the biotin-binding moiety is a polypeptide comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:2 or a biotin-binding fragment thereof.

8. The vaccine of claim 1, wherein the fusion protein comprises the amino acid sequence of SEQ ID NO: 6.

9. The vaccine of claim 2, comprising twenty four or more species of the immunogenic complexes.

10. The vaccine of claim 9, comprising:

a first immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 1, and (b) a fusion protein;

a second immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 2, and (b) a fusion protein;

a third immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 3, and (b) a fusion protein;

a fourth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 4, and (b) a fusion protein;

a fifth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 5, and (b) a fusion protein;

a sixth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 6A, and (b) a fusion protein;

a seventh immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 6B, and (b) a fusion protein;

an eighth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 7F, and (b) a fusion protein;

a ninth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 8, and (b) a fusion protein;

a tenth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 9N, and (b) a fusion protein;

an eleventh immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 9V, and (b) a fusion protein;

a twelfth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 10A, and (b) a fusion protein;

a thirteenth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 11A, and (b) a fusion protein;

a fourteenth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 12F, and (b) a fusion protein;

a fifteenth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 14, and (b) a fusion protein;

a sixteenth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 15B, and (b) a fusion protein;

a seventeenth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 17F, and (b) a fusion protein;

an eighteenth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 18C, and (b) a fusion protein;

a nineteenth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 19A, and (b) a fusion protein;

a twentieth immunogenic complex comprising (a) a biotinylated polysaccharide antigen, comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 19F, and (b) a fusion protein;

a twenty-first immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 20B, and (b) a fusion protein;

a twenty-second immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 22F, and (b) a fusion protein;

a twenty-third immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 23F and (b) a fusion protein; and

a twenty-fourth immunogenic complex comprising (a) a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from *Streptococcus pneumoniae* serotype 33F, and (b) a fusion protein;

wherein each fusion protein comprises:

(i) a biotin-binding moiety;

(ii) at least one polypeptide antigen comprising: a first polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:5 or an antigenic fragment thereof; and a second polypeptide antigen comprising an amino acid sequence at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to SEQ ID NO:4 or an antigenic fragment thereof;

and wherein in each of the immunogenic complexes, the biotinylated polysaccharide antigen is non-covalently associated with the biotin-binding moiety of the fusion protein.

11. A pharmaceutical composition comprising the vaccine of claim 1, and a pharmaceutically acceptable carrier.

12. The pharmaceutical composition of claim 11, further comprising one or more adjuvants.

13. The pharmaceutical composition of claim 12, wherein the one or more adjuvants is or comprises a co-stimulation factor.

14. The pharmaceutical composition of claim 12, wherein the one or more adjuvants are selected from the group consisting of aluminum phosphate, aluminum hydroxide, and phosphated aluminum hydroxide.

15. A method of making a multivalent vaccine, comprising mixing fifteen or more species of immunogenic complexes in a single formulation, wherein each species of the immunogenic complexes comprises:

(a)a biotinylated polysaccharide antigen comprising biotin and a polysaccharide antigen from a distinct *Streptococcus pneumoniae* serotype; and

(b)a fusion protein comprising:

(i) a biotin-binding moiety; and

(ii)at least one polypeptide antigen comprising:

a first polypeptide antigen comprising an SP1500 polypeptide or an antigenic fragment thereof; and

a second polypeptide antigen comprising an SP0785 polypeptide or an antigenic fragment thereof;

wherein the fusion protein comprises an amino acid sequence at least 90% identical to SEQ ID NO:6; and

wherein in each species of the immunogenic complexes, the biotinylated polysaccharide antigen is non-covalently associated with the biotin-binding moiety of the fusion protein.

16. The method of claim 15, wherein the *Streptococcus pneumoniae* serotype for each species of the immunogenic complexes is selected from 1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20B, 22F, 23F, and 33F.

17. Use of the multivalent vaccine of any one of claims 1-10 or the pharmaceutical composition of any one of claims 11-14, in the manufacture of a medicament for immunizing a subject against *Streptococcus pneumoniae* infection and/or colonization.

18. The use according to claim 17, wherein the vaccine or pharmaceutical composition induces

an immune response.

19. The use according to claim 18, wherein the immune response comprises (i) an antibody or B cell response to at least one polysaccharide antigen, and (ii) a CD4+ T cell response (*e.g.*, T_H1, T_H2, or T_H17 response), a CD8+ T cell response, a CD4+ and CD8+ T cell response, or a CD4-/CD8- T cell response to at least one polypeptide antigen of a fusion protein.

20. The use according to claim 18 or claim 19, wherein the immune response comprises (i) an antibody or B cell response to at least one polysaccharide antigen, and (ii) an antibody or B cell response and a CD4+ T cell response, including T_H1, T_H2, or T_H17 response, or a CD8+ T cell response, a CD4+ and CD8+ T cell response, or a CD4-/CD8- T cell response to at least one polypeptide antigen of a fusion protein.

21. The use according to any one of claims 17-20, wherein the vaccine inhibits or reduces the rate of occurrence of Invasive Pneumococcal Disease (IPD) associated with or induced by one or more serotypes of *Streptococcus pneumoniae*.

22. The use according to any one of claims 17-21, wherein the vaccine reduces the severity of Invasive Pneumococcal Disease (IPD) associated with or induced by one or more serotypes of *Streptococcus pneumoniae*.

23. The use according to any one of claims 17-22, wherein the vaccine reduces the severity of bacteremia, sepsis, and/or meningitis associated with or induced by one or more serotypes of *Streptococcus pneumoniae*.