

Description


The present invention relates to a process for the manufacture of 23-(C₁-C₆alkyloxime)-LL-F28249 compounds. The designation of LL-F28249 is used to describe a series of compounds produced by the fermentation broth of Streptomyces cyaneogriseus subspecies noncyanogenus, deposited in the NRRL collection under deposit accession number 15773.

It is an object of this invention to provide a process for the manufacture of 23-(C₁-C₆alkyloxime)-LL-F28249 compounds and more specifically 23-(methyloxime)-LL-F28249 α (moxidectin), a potent endectocidal agent.

The present invention relates to a process for the preparation of 23-(C₁-C₆alkyloxime)-LL-F28249 compounds which comprises protecting the 5-hydroxy group of LL-F28249 compounds with p-nitrobenzoyl chloride to give the corresponding 5-O(p-nitrobenzoyl)-LL-F28249 compound; oxidizing said compound to give a 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 derivative in a crystalline state; reacting said derivative with a C₁-C₆alkoxylamine or salt thereof to give the 23-(C₁-C₆alkyloxime)-5-O(p-nitrobenzoyl)-LL-F28249 intermediate in a crystalline state; and deprotecting said intermediate in the presence of base to give the desired 23-(C₁-C₆alkyloxime)-LL-F28249 compound. Optionally, the crystalline 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 derivatives are deprotected in the presence of base to give the corresponding 23-oxo-LL-F28249 compounds and said compounds are reacted with a C₁-C₆alkoxylamine, or salt thereof, to give the desired 23-(C₁-C₆alkyloxime)-LL-F28249 compound.

The 23-(C₁-C₆alkyloxime)-LL-F28249 compounds and their use as anthelmintic, insecticidal, nematocidal, ectoparasitocidal and acaricidal agents are described in U. S. Patent No. 4,916,154. The antibiotic compounds designated LL-F28249 are described in co-pending U. S. patent application Serial No. 732,252, filed on May 10, 1985.

The invention herein described relates to a process for the manufacture of 23-(C₁-C₆alkyloxime) derivatives of LL-F28249 compounds. The LL-F28249 compounds are represented by the following structural formula



Component	R ₁	R ₂	R ₃	R ₇
LL-F28249 α	CH(CH ₃) ₂	H	CH ₃	CH ₃
LL-F28249 β	CH ₃	H	CH ₃	CH ₃
LL-F28249c	CH ₃	CH ₃	CH ₃	CH ₃
LL-F28249e	CH(CH ₃) ₂	H	H	CH ₃
LL-F28249f	CH ₂ CH ₃	H	CH ₃	CH ₃
LL-F28249h	CH(CH ₃) ₂	H	CH ₃	CH ₂ CH ₃
LL-F28249i	CH(CH ₃) ₂	H	CH ₂ CH ₃	CH ₃
LL-F28249k	CH(CH ₃) ₂	CH ₃	CH ₃	CH ₃

There is provided a process for the manufacture of 23-(C₁-C₆alkyloxime)-LL-F28249 compounds wherein R₂ is hydrogen which comprises protecting the 5-hydroxy group of said LL-F28249 compounds with p-nitrobenzoyl chloride to yield the corresponding 5-O(p-nitrobenzoyl)-LL-F28249 compound; oxidizing said compound to yield the 5-O(p-nitrobenzoyl)-33-oxo-LL-F28249 derivative in a crystalline state; reacting said derivative with a C₁-C₆alkoxyamine or salt thereof to yield the 23-(C₁-C₆alkyloxime)-5-O(p-nitrobenzoyl)-LL-F28249 intermediate in a crystalline state; and deprotecting said intermediate in the presence of base to yield the desired 23-(C₁-C₆alkyloxime)-LL-F28249 compound. There is further provided a process for the manufacture of 23-(C₁-C₆alkyloxime)-LL-F28249 compounds wherein the crystalline 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 derivatives are deprotected in the presence of base to yield the corresponding 23-oxo-LL-F28249 compounds and said compounds are reacted with a C₁-C₆alkoxyamine or salt thereof to yield the desired 23-(C₁-C₆alkyloxime)-LL-F28249 compound.

Using LL-F28249 α as starting material and methoxyamine hydrochloride as the C₁-C₆alkoxyamine reagent, the processes of the invention can be illustrated as shown in flow diagram I wherein PNB designates the functionality p-nitrobenzoyl.

FLOW DIAGRAM I

5

10

15

20

25


30

35

40

45

50



Protection of the 5-hydroxy group of LL-F28249 α is achieved by the reaction of LL-F28249 α with p-nitrobenzoyl chloride in the presence of an organic solvent such as toluene, methylene chloride, ethyl acetate, acetonitrile and the like, preferably toluene, and an organic base such as pyridine, triethylamine, N-methylpyrrolidone, and the like, preferably triethylamine.

Surprisingly, it has been found that oxidation of 5-O(p-nitrobenzoyl)-LL-F28249 compounds gives a crystalline product, the corresponding 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 compound. A crystalline intermediate allows a simplified, efficient purification of said intermediate via recrystallization from a suitable organic solvent and eliminates complex, time consuming purification by procedures such as column chromatography. Oxidation of the 5-O(p-nitrobenzoyl)-LL-F28249 compound is successfully achieved using an oxidizing system selected from the group consisting of pyridinium dichromate and acetic anhydride; pyridinium dichromate and dimethylformamide; aluminum t-butoxide and o-benzoquinone; phosphorous pentoxide and dimethyl sulfoxide; dicyclohexylcarbodiimide and dimethyl sulfoxide; manganese dioxide; and acetic anhydride and dimethyl sulfoxide.

A preferred oxidizing system for the oxidation of 5-O(p-nitrobenzoyl)-LL-F28249 compounds is manganese dioxide in the presence of a solvent such as methylene chloride, acetonitrile, ethyl acetate or the like, preferably ethyl acetate.

A more preferred oxidizing system for the oxidation of 5-O(p-nitrobenzoyl)-LL-F28249 compounds is acetic anhydride and dimethyl sulfoxide in the presence of pyridine and an acid such as acetic acid, trifluoroacetic acid, dichloroacetic acid, monochloroacetic acid and the like, preferably monochloroacetic acid. Surprisingly, it has been found that the use of acetic anhydride and dimethyl sulfoxide in the presence of pyridine and an acid greatly enhances the yield of the 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 derivatives over the use of acetic anhydride and dimethyl sulfoxide alone. For example, the acetic anhydride and dimethyl sulfoxide oxidation is carried out in the presence of pyridine and monochloroacetic acid, the yield of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α is 76%; whereas when the same oxidation is carried out without the presence of pyridine and monochloroacetic acid, the yield is 2-4%.

Optionally, the crystalline 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 derivative is purified by recrystallization from a suitable solvent (preferably n-propanol) prior to deprotection (removal of the p-nitrobenzoyl group) or reaction with C₁-C₆alkoxylamine hydrochloride.

Preferably, a solution of the crude reaction product, 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 compound, in an organic solvent, such as toluene, is reacted with an aqueous solution of C₁-C₆alkoxylamine hydrochloride and sodium acetate and stirred until oxime formation is complete. The thus-formed 23-(C₁-C₆alkyloxime)-5-O(p-nitrobenzoyl)-LL-F28249 intermediate is isolated and then purified by recrystallization from a suitable solvent, preferably n-butanol.

The recrystallized 23-(C₁-C₆alkyloxime)-5-O(p-nitrobenzoyl)-LL-F28249 compound is deprotected by reaction with sodium hydroxide at 0°-25°C to give the desired 23-(C₁-C₆alkyloxime)-LL-F28249 product. The deprotection is achieved by reacting a solution of a 23-(C₁-C₆alkyloxime)-5-O(p-nitrobenzoyl)-LL-F28249 compound in an organic solvent such as toluene, dioxane, n-butanol or the like, preferably dioxane, with an aqueous solution of sodium hydroxide at 0°-25°C and isolating the product 23-(C₁-C₆alkyloxime)-LL-F28249 compound from the organic phase using standard procedures such as concentration and filtration or removal of the solvent.

In order to facilitate a further understanding of the invention, the following examples are presented primarily for the purpose of illustrating more specific details thereof. The invention is not to be limited thereby except as defined in the claims.

Unless otherwise noted, all parts are by weight, all high pressure liquid chromatographic analyses are designated as HPLC analyses, and all proton nuclear magnetic spectroscopic analyses are designated as ¹HNMR analyses.

EXAMPLE 1

Preparation of 5-O(p-nitrobenzoyl)-LL-F28249 α

A stirred solution of LL-F28249 α (6.36 g, 10.4 mmole) in methylene chloride is treated with pyridine (1.98 g, 25.0 mmole) and p-nitrobenzoyl chloride (2.45 g, 13.2 mmole) at 20°-25°C. After 4 hours at 20°-25°, the reaction mixture is treated with saturated sodium bicarbonate and methylene chloride and stirred until solution is complete. The phases are separated, the organic phase is washed sequentially with saturated sodium bicarbonate, 5% hydrochloric acid, and saturated sodium chloride and concentrated in vacuo to give the title compound as a solid foam, 7.9 g (quantitative yield) identified by liquid chromatography, ¹HNMR and mass spectral analyses.

EXAMPLE 2**Preparation of 5-O(p-nitrobenzoyl)-LL-F28249 α**

5 A stirred solution of LL-F28249 α (6.13 g, 10.0 mmole) in toluene is treated with triethylamine (2.53 g, 25 mmole), cooled to 15°C, treated portionwise with p-nitrobenzoyl chloride (2.60 g, 14 mmole) at a temperature range of 15°C-22°C, and stirred for 6 hours at 20°C-24°C. The reaction mixture is treated with
10 water, stirred for 10 minutes and filtered. The filtrate is separated, the organic phase is washed sequentially with saturated sodium bicarbonate, 3N hydrochloric acid and water, and concentrated in vacuo to give the title product as a solid foam, 7.55 g.

EXAMPLE 3**Preparation of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α
using pyridinium dichromate and dimethylformide**

15 A stirred solution of 5-O(p-nitrobenzoyl)-LL-F28249 α (3.12 g, 4.10 mmole) in dimethylformamide is treated with pyridinium dichromate (18.8 g, 50 mmole) in a single portion, stirred at 20°C-25°C for 6 hours and poured into water. The reaction mixture is stirred for 15 minutes and filtered. The filter cake is washed
25 with water, air-dried and taken up in ethyl acetate. The resulting mixture is heated at reflux temperature for 15 minutes, treated with diatomaceous earth and filtered. The filtrate is concentrated in vacuo to give a red-brown solid which is recrystallized from n-propanol to give the title product as white crystals, 3.33 g (52% overall yield from LL-F28249 α) mp 217-221°C identified by ¹HNMR and mass spectral analyses.

EXAMPLE 4**Preparation of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α
using pyridinium dichromate and acetic anhydride.**

35 A solution of 5-O(p-nitrobenzoyl)-LL-F28249 α (0.38 g, 0.5 mmol) in methylene chloride is added to a freshly prepared mixture of pyridinium dichromate (0.19 g, 0.5 mmol) and acetic anhydride (0.3g, 3.0 mmol) with vigorous stirring. The reaction mixture is stirred at room temperature for 15 minutes, heated at reflux temperature for 6-8 hours, cooled to room temperature, and treated with water. After vigorous stirring, the phases are separated and the organic phase is washed with saturated sodium bicarbonate solution and concentrated in vacuo to give a residue. The residue is taken up in ethyl acetate and chromatographed
45 using silica gel and ethyl acetate as eluent to give a pale yellow/grey solid. This solid is stirred with ethyl acetate; hexanes (55:45 v/v) filtered, and the filtrate is concentrated in vacuo to give the title compound as a pale yellow solid, 0.26 g, identified by ¹HNMR and HPLC analyses.

EXAMPLE 5**Preparation of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α
using phosphorous pentoxide and dimethyl sulfoxide**

50 A stirred mixture of 5-O(p-nitrobenzoyl)-LL-F28249 α (0.38 g, 0.50 mmole), aluminum t-butoxide (0.184 g, 0.75 mmole) and o-benzoquinone (0.216 g, 2.0 mmole) in toluene is heated at reflux temperature for 2 hours, cooled to room temperature, treated with toluene and dilute sulfuric acid (16%), stirred for 5 minutes and filtered. The filtrate is separated and the organic phase is washed with water and concentrated in vacuo
55

to give a glassy solid residue. The residue is taken up in ethyl acetate and filtered through neutral alumina. The filtrate is concentrated in vacuo to give the title product as a white solid, 0.324 g, (71% yield by HPLC analyses).

5 EXAMPLE 6

10 Preparation of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α
using phosphorous pentoxide and dimethyl sulfoxide

A solution of 5-O(p-nitrobenzoyl)-LL-F28249 α (0.38 g, 0.50 mmole) and dimethyl sulfoxide (0.75 g, 9.6 mmole) in methylene chloride is treated with powdered phosphorous pentoxide (0.107 g, 0.75 mmole) in one portion, stirred at 20°-25°C for 19 hours, treated dropwise with triethylamine (0.30 g, 3.0 mmole), stirred for 30 minutes, treated further with methylene chloride and water and stirred for 5 minutes. The phases are separated and the organic phase is washed with dilute hydrochloric acid (7%) and concentrated in vacuo to give the title product as a white solid, 0.28 g (47% purity by HPLC analysis).

20 EXAMPLE 7

25 Preparation of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α
via manganese dioxide using methylene chloride as
solvent

A solution of 5-O(p-nitrobenzoyl)-LL-F28249 α (0.19 g, 0.25 mmole) in methylene chloride is treated with manganese dioxide (8.0 g, 92 mmole), stirred at 20°-25°C for 2 hours, treated further with methylene chloride, stirred for 5 minutes and filtered. The filtrate is concentrated in vacuo to give the title compound as a white solid, 0.08 g, 51% purity by HPLC analysis.

35 EXAMPLE 8

40 Preparation of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α
via manganese dioxide using acetonitrile as solvent

A solution of 5-O(p-nitrobenzoyl)-LL-F28249 α (0.19 g, 0.25 mmole) in acetonitrile is treated with manganese dioxide (8.0 g, 92 mmole), stirred for 3 hours at room temperature, treated further with acetonitrile, stirred for 5 minutes and filtered. The filter cake is slurried in methylene chloride and filtered. The acetonitrile and methylene chloride filtrates were combined, washed with water and concentrated in vacuo to give the title compound as a white solid, 0.14 g, 69% purity by HPLC analysis.

EXAMPLE 9

50 Preparation of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α
via manganese dioxide and using ethyl acetate as
solvent

55

A solution of 5-O(p-nitrobenzoyl)-LL-F28249 α (1.0 g, 1.3 mmole) in ethyl acetate is treated with manganese dioxide (20.0 g, 230 mmole), stirred at 20°-25°C for 3 hours and filtered. The filter cake is

washed with ethyl acetate. The filtrates are combined, treated with manganese dioxide (8.0 g, 92 mmole), stirred at 20°-25°C for 2 hours and filtered. The filter cake is washed with ethyl acetate; the filtrates are combined and concentrated in vacuo to give the title product as a white solid, 0.8 g, (70% purity by HPLC analysis). The solid is recrystallized from n-propanol to give white crystals, mp 218-222°C.

EXAMPLE 10

Preparation of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α via dicyclohexylcarbodiimide and dimethyl sulfoxide

A solution of 5-O(p-nitrobenzoyl)-LL-F28249 α (0.38 g, 0.50 mmole) in benzene is treated sequentially with dimethyl sulfoxide (0.78 g, 10 mmole), pyridine (0.04 g, 0.5 mmole) trifluoroacetic acid (0.03 g, 0.25 mmole) and dicyclohexylcarbodiimide (0.31 g, 1.5 mmole), stirred at 20°-25°C for 21 hours, treated further with benzene and filtered. The filter cake is washed with benzene. The combined filtrates are washed with water and concentrated in vacuo to give the title compound as a light orange-brown solid, 0.34 g, 75% yield by HPLC analysis.

EXAMPLE 11

Preparation of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α via acetic anhydride and dimethyl sulfoxide in the presence of pyridinium trifluoroacetate

A mixture of 5-O(p-nitrobenzoyl)-LL-F28249 α (0.38 g, 0.5 mmole), dimethyl sulfoxide (0.78 g, 10 mmole) and pyridinium trifluoroacetate (0.97 g, 0.5 mmole) in ethyl acetate is treated dropwise with acetic anhydride (0.26 g, 2.5 mmole), stirred for 24 hours at 20°-25°C and treated with ethyl acetate and water. The phases are separated; the organic phase is washed with water and concentrated in vacuo to give a viscous oil residue. The residue is taken up in methylene chloride and concentrated in vacuo to give the title product as a yellow solid, 0.36 g, identified by HPLC analysis.

EXAMPLE 12

Preparation of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α via acetic anhydride and dimethyl sulfoxide in the presence of pyridine and dichloroacetic acid

A mixture of 5-O(p-nitrobenzoyl)-LL-F28249 α (7.26 g, 10 mole) and pyridine (31.6 g, 400 mmole) is treated with dimethyl sulfoxide (15.6 g, 200 mmole) and dichloroacetic acid (1.29 g, 10 mmole), cooled to 2°-3°C, treated dropwise with acetic anhydride (5.1 g, 50 mmole) at 3°-7°C and treated with methylene chloride and water. The reaction mixture is stirred at ambient temperatures for 15-30 minutes and the phases are separated. The organic phase is washed with cold dilute hydrochloric acid (5%), and 5% sodium chloride solution and concentrated in vacuo to give the title product as a yellow solid foam, 7.47 g, 73% purity by HPLC analysis.

Using essentially the same procedure, but varying the acid reagent used, the following yields are obtained and reported in Table I.

Table I

Oxidation of 5-O(p-nitrobenzoyl)-LL-F28249alpha to
5-O(p-nitrobenzoyl)-23-oxo-LL-F28249alpha via
Acetic Anhydride and Dimethyl Sulfoxide
in the Presence of Pyridine and An Acid

Exp.	Molar Equivalents				Acid ^a -Mol. eq	Solvent	Temp (°C)	Time (h)	Yield (%)
	Dimethyl Sulfoxide	Acetic Anhydride	Pyridine	Pyridine					
1	20	5	40	40	DCAA	Pyridine	2-3	4	76
2	20	5	40	40	DCAA	Pyridine	2-3	4	57
3	20	4	20	20	NCAA	Pyridine	2-3	7	75
4	20	5	40	40	NCAA	Pyridine	20-25	4	51
5	20	5	40	40	TERE	Pyridine	20-25	24	55-60
6	20	5	40	40	TCAA	Pyridine	20-25	24	40-50
7	20	5	40	40	H ₃ PO ₄	Pyridine	20-25	24	20-25
8	20	5	40	40	BA	Pyridine	20-25	4	15-20
9	20	5	10	10	DCAA	Ethyl acetate	20-25	6	66
10 ^b	20	5	1	1	TFAA	Ethyl acetate	20-25	24	60
11	20	8	22	22	DCAA	Ethyl acetate	20-25	3	63

Table I (Continued)

Exp.	Molar Equivalents		Pyridine	Acid ^a -Mol eq	Solvent	Temp (°C)	Time (h)	Yield %
	Dimethyl Sulfoxide	Acetic Anhydride						
12	20	3	1	TFAA 1.0	Benzene	20-25	6	42
13	40	5	1	TFAA 1.0	Benzene	20-25	8	54
14	40	5	1	TFAA 1.0	Toluene	20-25	24	45
15	40	3	1	TFAA 1.0	CH ₃ Cl ₂	20-25	4	34
16	40	3	1	TFAA 1.0	CH ₂ CN	20-25	5	20-25
17	40	3	1	TFAA 1.0	Dimethyl- formamide	20-25	4	20-25
18	40	40	0	0	Dimethyl sulfoxide	20-25	16	2-4

- a. DCAA = dichloroacetic acid, NCA = monochloroacetic acid, TCAA = trichloroacetic acid,
 TERE = terephthalic acid, BA = benzoic acid, TFAA = trifluoroacetic acid
- b. Reaction is carried out in the presence of 4A molecular sieves.

EXAMPLE 13

5 **Preparation of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α**
via acetic anhydride and dimethyl sulfoxide in the
presence of pyridine and monochloroacetic acid

10 A mixture of 5-O(p-nitrobenzoyl)-LL-F28249 α (1.52 g, 2.0 mmole) and pyridine (3.16 g, 40 mmole) in toluene is treated with dimethyl sulfoxide (3.12 g, 40 mmole) and monochloroacetic acid (0.19 g, 2.0 mmole), cooled to 2°-3°C, treated dropwise with acetic anhydride (0.82 g, 8.0 mmole) at 3°-5°C, stirred at 2-3°C for 7 hours and treated further with toluene and water. After stirring the reaction mixture at 15°-20°C for 10 minutes, the phases are separated. The organic phase is washed sequentially with cold 2.4N
15 hydrochloric acid and water at 15-20°C and concentrated in vacuo to give the title product as a yellow solid foam, 1.4 g, 71% purity by HPLC analysis.

EXAMPLE 14

20 **Evaluation of the oxidation of 5-O(p-nitrobenzoyl)-LL-**
F28249 α to form 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α
using a variety of oxidizing reagents

25

A variety of oxidizing systems are evaluated for the conversion of 5-O(p-nitrobenzoyl)-LL-F28249 α to 5-O(p-nitrobenzoyl)-23-oxo-LL-28249 α . The reagents, reaction conditions and percent 5-O(p-nitrobenzoyl)-23-oxo-LL-28249 α obtained as determined by HPLC analysis are reported in Table II.

30

35

40

45

50

55

Table II

Attempted Oxidation of 5-O(p-Nitrobenzoyl)-LL-F28249alpha to
5-O(p-Nitrobenzoyl)-23-oxo-LL-F28249alpha
Using A Variety of Oxidizing Reagents

<u>Experiment</u>	<u>Oxidant</u>	<u>Molar Equivalents</u>	<u>Solvent</u>	<u>Temp(°C)</u>	<u>Time(h)</u>	<u>% Product (HPLC area %)</u>
1	MoO ₃ Dimethyl sulfoxide	0.04 0.80	Toluene	Reflux	3.5	14%
2	N-Bromo- succinimide	1.0	CH ₃ CN	20-25	3	10%
3	K ₂ S ₂ O ₈ Bu ₄ NBr RuCl ₃	1.2 0.1 0.2	CH ₂ Cl ₂ / water	20-25	3	6%
4	NaOCl Bu ₄ NBr	6.0 0.5	CH ₂ Cl ₂ / water	20-25	ca 4	0%

Table II (Continued)

<u>Experiment</u>	<u>Oxidant</u>	<u>Molar Equivalents</u>	<u>Solvent</u>	<u>Temp(°C)</u>	<u>Time(h)</u>	<u>% Product (HPLC area %)</u>
13	Pyridinium	3.0	Methylene	20-25	1	No reaction
	Sulfite		Chloride			
	Dimethyl	40.0				
	Sulfoxide					
14	Pyridinium	3.0	Methylene	20-25	22	No reaction
	Sulfite	20.0	Chloride			
	Pyridine	1.0				
	Trifluoro- acetic acid	0.5				
15	H ₂ O ₂	4.0	Methylene	20-25	5	No reaction
	Bu ₄ NBr	0.1	Chloride Water			
16	p-Benzoquinone Alumina	20	Benzene	Reflux	5	No reaction

Table II (Continued)

Experiment	Oxidant	Molar Equivalents	Solvent	Temp(°C)	Time(h)	% Product (HPLC area %)
5	$(\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$	10	Ethyl Acetate	20-25	3	0%
6	KMnO_4	1.5	t-Butanol/water	20-25	1	0%
7	KMnO_4	1.5	Acetic Acid	20-25	0.5	0%
8	KMnO_4	1.0	Acetone	20-25	5	0%
9	KMnO_4 $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$	1.5 6.0	Methylene Chloride	20-25	5	No reaction
10	BaMnO_4	10	Toluene	Reflux	6	0%
11	BaMnO_4 $\text{CH}_3\text{NET}_3\text{Cl}$	15.0 0.5	Methylene Chloride	20-25	6	No reaction
12	BaMnO_4	15	Methylene Chloride	20-25	5	No reaction

EXAMPLE 15**Preparation of 5-O(p-nitrobenzoyl)-23-methyloxime)-LL-F28249 α**

A solution of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α (10.67 g, 14.0 mmole) in *n*-butanol is treated with a solution of methoxylamine hydrochloride (2.34 g, 28.1 mmole) and anhydrous sodium acetate (2.30 g, 28.1 mmole) in water at 20°-22°C, stirred for 2 hours at 20°-22°C and filtered. The filter cake is air-dried and recrystallized from *n*-butanol (filtered hot) to give the title compound as a colorless solid, 3.6 g, 91% purity by HPLC analysis.

EXAMPLE 16**Preparation of 5-O(p-nitrobenzoyl)-23-(methyloxime)-LL-F28249 α**

A solution of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α (1.5 g, 2.0 mmole) in toluene is treated with a solution of methoxylamine hydrochloride (0.25 g, 3.0 mmol) and anhydrous sodium acetate (0.25 g, 3.0 mmol) in water and stirred at 20°-25°C for 10 hours. The toluene phase is separated, washed with water and concentrated *in vacuo* to give a solid residue. The solid is recrystallized from *n*-butanol to give the title product, 0.65 g, identified by HPLC analysis.

EXAMPLE 17**Preparation of 23-(methyloxime)-LL-F28249 α**

A solution of 5-O(p-nitrobenzoyl)-23-(methyloxime)-LL-F28249 α (1.58 g, 2.0 mmole) in dioxane is treated dropwise with 4% sodium hydroxide (3.0 g, 3.0 mmole NaOH) at 8°-12°C, stirred for 3 hours at 8°-12°C, treated with toluene and water, and stirred for 5 minutes at ambient temperatures. The phases are separated and the organic phase is washed with 10% sodium chloride and concentrated *in vacuo* to give the title compound as a white solid foam, 1.15 g, 89% purity by HPLC analysis.

EXAMPLE 18**Preparation of 23-oxo-LL-F28249 α**

A mixture of 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 α (1.52 g, 2.0 mmole) and 4% sodium hydroxide (3.3 g, 3.3 mmole NaOH) in dioxane is stirred at 23°C for 2 hours, treated with toluene and water and shaken. The phases are separated and the organic phase is washed with water and concentrated *in vacuo* to give the title compound as a solid foam, 0.90 g identified by ¹HNMR.

EXAMPLE 19**Preparation of 23-(methyloxime)-LL-F28249 α**

A mixture of 23-oxo-LL-F28249 α (0.90 g, 1.5 mmole), methoxylamine hydrochloride (0.42 g, 5.0 mmole), anhydrous sodium acetate (0.41 g, 5.0 mmole), acetic acid and dioxane is stirred at 20°-25°C for 22 hours, treated with toluene and water and stirred for 5 minutes. The phases are separated, and the organic phase is washed with water and concentrated *in vacuo* to give the title compound as a solid foam, 0.84 g, 71% purity by HPLC analysis.

Claims

1. A process for the preparation of a 23-(C₁-C₆alkyloxime)-LL-F28249 compound characterized by protecting the 5-hydroxy group of a LL-F28249 compound with p-nitrobenzoyl chloride to yield a 5-O(p-nitrobenzoyl)-LL-F28249 compound; oxidizing said compound to yield a 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 derivative in a crystalline state; reacting said derivative with a C₁-C₆alkoxylamine or salt thereof to yield a 23-(C₁-C₆alkyloxime)-5-O(p-nitrobenzoyl)-LL-F28249 intermediate; and deprotecting said intermediate in the presence of base to yield the product 23-(C₁-C₆alkyloxime)-LL-F28249 compound.
2. A process for the preparation of a 23-(C₁-C₆alkyloxime)-LL-F28249 compound characterized by protecting the 5-hydroxy group of a LL-F28249 compound with p-nitrobenzoyl chloride to yield a 5-O(p-nitrobenzoyl)-LL-F28249 compound; oxidizing said compound to yield a 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 derivative in a crystalline state; deprotecting said derivative in the presence of base to yield a 23-oxo-LL-F28249 compound; and reacting said compound with a C₁-C₆alkoxylamine or salt thereof to yield the product 23-(C₁-C₆alkyloxime)-LL-F28249 compound.
3. The process according to claim 1 wherein the oxidation is carried out using an oxidizing system selected from the group consisting of acetic anhydride and dimethyl sulfoxide; manganese dioxide; pyridinium dichromate and acetic anhydride; aluminum t-butoxide and o-benzoquinone; phosphorous pentoxide and dimethyl sulfoxide; and dicyclohexylcarbodiimide and dimethyl sulfoxide.
4. The process according to claim 2, wherein the oxidation is carried out using an oxidizing system selected from the group consisting of acetic anhydride and dimethyl sulfoxide; manganese dioxide; pyridinium dichromate and acetic anhydride; aluminum t-butoxide and o-benzoquinone; phosphorous pentoxide and dimethyl sulfoxide; and dicyclohexylcarbodiimide and dimethyl sulfoxide.
5. The process according to claims 3 or 4, wherein the oxidizing system is acetic anhydride and dimethyl sulfoxide and the oxidation is carried out in the presence of pyridine and an acid.
6. The process according to claims 3 or 4, wherein the oxidizing system is manganese dioxide and the oxidation is carried out in the presence of ethyl acetate.
7. The process according to claims 3 or 4, wherein the oxidizing system is pyridinium dichromate and acetic anhydride.
8. The process according to claims 1 or 2, wherein the 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 derivative is purified by recrystallization from a suitable solvent.
9. The process according to claim 1, wherein the 23-(C₁-C₆alkyloxime)-5-O(p-nitrobenzoyl)-LL-F28249 intermediate is purified by recrystallization from a suitable solvent.
10. The process according to claims 1 or 2, wherein the product compound is 23-(methyloxime)-LL-F28249 α .

Patentansprüche

1. Verfahren zur Herstellung einer 23-(C₁-C₆-Alkyloxim)-LL-F28249-Verbindung, gekennzeichnet durch Schützen der 5-Hydroxygruppe einer LL-F28249-Verbindung mit p-Nitrobenzoylchlorid unter Herstellung einer 5-O-(p-Nitrobenzoyl)-LL-F28249-Verbindung; Oxidieren der Verbindung zu einem 5-O-(p-Nitrobenzoyl)-23-oxo-LL-F28249-Derivat im kristallinen Zustand, Umsetzen des Derivats mit einem C₁-C₆-Alkoxylamin oder einem Salz davon zu einem 23-(C₁-C₆-Alkyloxim)-5-O-(p-nitrobenzoyl)-LL-F28249-Zwischenprodukt; und Entfernen der Schutzgruppen aus dem Zwischenprodukt in Gegenwart einer Base unter Herstellung der Produktverbindung 23-(C₁-C₆-Alkyloxim)-LL-F28249.
2. Verfahren zur Herstellung einer 23-(C₁-C₆-Alkyloxim)-LL-F28249-Verbindung, gekennzeichnet durch Schützen der 5-Hydroxygruppe einer LL-F28249-Verbindung mit p-Nitrobenzoylchlorid unter Herstellung einer 5-O-(p-Nitrobenzoyl)-LL-F28249-Verbindung; Oxidieren der Verbindung zu einem 5-O-(p-Nitrobenzoyl)-23-oxo-LL-F28249-Derivat im kristallinen Zustand; Entfernen der Schutzgruppen aus dem



Derivat in Gegenwart einer Base unter Herstellung einer 23-Oxo-LL-F28249-Verbindung und Umsetzen der Verbindung mit einem C₁-C₆-Alkoxyaminoder einem Salz davon zu der Produktverbindung 23-(C₁-C₆-Alkyloxim)-LL-F28249.

- 5 3. Verfahren nach Anspruch 1, wobei die Oxidation unter Verwendung eines Oxidationssystems, ausgewählt aus der Gruppe, bestehend aus Essigsäureanhydrid und Dimethylsulfoxid; Mangandioxid; Pyridiniumdichromat und Essigsäureanhydrid; Aluminium-tert.-butoxid und o-Benzochinon; Phosphorpentoxid und Dimethylsulfoxid; und Dicyclohexylcarbodiimid und Dimethylsulfoxid ausgeführt wird.
- 10 4. Verfahren nach Anspruch 2, wobei die Oxidation unter Verwendung eines Oxidationssystems, ausgewählt aus der Gruppe, bestehend aus Essigsäureanhydrid und Dimethylsulfoxid; Mangandioxid; Pyridiniumdichromat und Essigsäureanhydrid; Aluminium-tert.-butoxid und o-Benzochinon; Phosphorpentoxid und Dimethylsulfoxid; und Dicyclohexylcarbodiimid und Dimethylsulfoxid ausgeführt wird.
- 15 5. Verfahren nach den Ansprüchen 3 oder 4, wobei das Oxidationssystem Essigsäureanhydrid und Dimethylsulfoxid ist und die Oxidation in Gegenwart von Pyridin und einer Säure ausgeführt wird.
6. Verfahren nach den Ansprüchen 3 oder 4, wobei das Oxidationssystem Mangandioxid ist und die Oxidation in Gegenwart von Essigsäureethylester ausgeführt wird.
- 20 7. Verfahren nach den Ansprüchen 3 oder 4, wobei das Oxidationssystem Pyridiniumdichromat und Essigsäureanhydrid ist.
8. Verfahren nach den Ansprüchen 1 oder 2, wobei das 5-O(p-Nitrobenzoyl)-23-oxo-LL-F28249-Derivat durch Umkristallisieren aus einem geeigneten Lösungsmittel gereinigt wird.
- 25 9. Verfahren nach Anspruch 1, wobei das 23-(C₁-C₆-Alkyloxim)-5-O(p-nitrobenzoyl)-LL-F28249-Zwischenprodukt durch Umkristallisieren aus einem geeigneten Lösungsmittel gereinigt wird.
- 30 10. Verfahren nach den Ansprüchen 1 oder 2, wobei die Produktverbindung 23-(Methyloxim)-LL-F28249 α ist.

Revendications

- 35 1. Procédé pour la préparation d'un composé 23-(alkyloxime en C₁-C₆)-LL-F28249 caractérisé par la protection du groupe 5-hydroxy d'un composé LL-F28249 avec du chlorure de p-nitrobenzoyl pour obtenir un composé 5-O(p-nitrobenzoyl)-LL-F28249 ; l'oxydation dudit composé pour obtenir un dérivé 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 à l'état cristallin ; la réaction dudit dérivé avec une alcoxylamine en C₁-C₆ ou un de ses sels pour obtenir un 23-(alkyloxime en C₁-C₆)-5-O(p-nitrobenzoyl)-LL-F28249 intermédiaire ; et la déprotection dudit intermédiaire en présence d'une base pour obtenir le composé 23-(alkyloxime en C₁-C₆)-LL-F28249 produit.
- 40 2. Procédé pour la préparation d'un composé 23-(alkyloxime en C₁-C₆)-LL-F28249 caractérisé par la protection du groupe 5-hydroxy d'un composé LL-F28249 avec du chlorure de p-nitrobenzoyl pour obtenir un composé 5-O(p-nitrobenzoyl)-LL-F28249 ; l'oxydation dudit composé pour obtenir un dérivé 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 à l'état cristallin ; la déprotection dudit dérivé en présence d'une base pour obtenir un composé 23-oxo-LL-F28249 ; et la réaction dudit composé avec une alcoxylamine en C₁-C₆ ou un de ses sels pour obtenir le composé 23-(alkyloxime en C₁-C₆)-LL-F28249 produit.
- 45 3. Procédé selon la revendication 1, dans lequel l'oxydation est effectuée en utilisant un système oxydant choisi parmi l'ensemble constitué par l'anhydride acétique et le diméthylsulfoxyde ; le dioxyde de manganèse ; le dichromate de pyridinium et l'anhydride acétique ; le tert-butylate d'aluminium et l'o-benzoquinone ; le pentoxyde de phosphore et le diméthylsulfoxyde ; et le dicyclohexylcarbodiimide et le diméthylsulfoxyde.
- 50 4. Procédé selon la revendication 2, dans lequel l'oxydation est effectuée en utilisant un système oxydant choisi parmi l'ensemble constitué par l'anhydride acétique et le diméthylsulfoxyde ; le dioxyde de manganèse ; le dichromate de pyridinium et l'anhydride acétique ; le tert-butylate d'aluminium et l'o-
- 55



benzoquinone ; le pentoxyde de phosphore et le diméthylsulfoxyde ; et le dicyclohexylcarbodiimide et le diméthylsulfoxyde.

- 5 5. Procédé selon les revendications 3 ou 4, dans lequel le système oxydant est l'anhydride acétique et le diméthylsulfoxyde et l'oxydation est effectuée en présence de pyridine et d'un acide.
6. Procédé selon les revendications 3 ou 4, dans lequel le système oxydant est le dioxyde de manganèse et l'oxydation est effectuée en présence d'acétate d'éthyle.
- 10 7. Procédé selon les revendications 3 ou 4, dans lequel le système oxydant est le dichromate de pyridinium et l'anhydride acétique.
8. Procédé selon les revendications 1 ou 2, dans lequel on purifie le dérivé 5-O(p-nitrobenzoyl)-23-oxo-LL-F28249 par recristallisation dans un solvant approprié.
- 15 9. Procédé selon la revendication 1, dans lequel on purifie le 23-(alkyloxime en C₁-C₆)-5-O(p-nitrobenzoyl)-LL-F28249 intermédiaire par recristallisation dans un solvant approprié.
- 20 10. Procédé selon les revendications 1 ou 2, dans lequel le composé produit est le 23-(méthyloxime)-LL-F28249 α .

25

30

35

40

45

50

55

