

METHODS FOR DIAGNOSING AND EVALUATING NON-ALCOHOLIC STEATOHEPATITIS

5 FIELD OF THE INVENTION

The invention relates to a method for the diagnosis of non-alcoholic steatohepatitis (NASH), for determining the activity, the stage, or the severity of NASH or for classifying a subject as a potential receiver or non receiver of a treatment of NASH. It also relates to a kit for implementing the method
10 of the invention, and the compounds for use in a method for the treatment of NASH, wherein the subject to be treated is identified, evaluated or classified according to the method of the invention.

BACKGROUND OF THE INVENTION

15 Non-alcoholic steatohepatitis (NASH) is a progressive disease of the liver characterized histologically by fatty acid accumulation, hepatocyte damage and inflammation resembling alcoholic hepatitis. NASH is a critical stage in the process that spans from hepatic steatosis to cirrhosis and liver failure. A careful history of a lack of significant alcohol intake is essential to establish this diagnostic. NASH is one of the most common causes of elevated aminotransferases in patients referred for evaluation to
20 hepatologists. Obesity and type-2 diabetes are associated to NASH. Since the prevalence of these diseases is increasing, the prevalence of NASH is also expected to increase and therefore, this disease is becoming an emerging public issue in the United States as well as in other countries.

Although several attempts have been made to propose non-invasive methods for diagnosing and
25 determining the activity, the stage, or the severity of NASH, as of today histological analysis of liver biopsies remains the optimal approach for differentiating NASH from early stage steatosis. Steatosis, lobular and portal inflammation, hepatocyte injury in the forms of ballooning and apoptosis, and fibrosis are features of NASH evaluated from the biopsy. However, liver biopsy has a number of obvious drawbacks. First, the material collected in liver biopsy represents only a very small part of the
30 liver of the diagnosed subject, thereby raising doubts on whether the collected sample is representative of the global state of the subject's organ. Moreover, liver biopsy is a very invasive procedure that may be cumbersome, worrisome and painful for the patient, and which raises concerns about morbidity and mortality. At last, in view of the foregoing, liver biopsy cannot be reasonably proposed as a routine procedure for determining whether a person in the general population, or even patients at risk of
35 NASH, suffers from NASH and/or for determining the activity, the stage, or the severity of NASH in said person.

Therefore, a strong need exists in providing accurate, non-invasive, means for diagnosing and determining the activity, the stage, or the severity of NASH in a subject.

SUMMARY OF THE INVENTION

5

The present invention is based on a very fine and complete analysis of a vast number of variables determined from more than 300 high quality samples obtained from NASH patients during a clinical trial conducted by the Applicant (study GFT505-212-7 - NCT01694849), including histological data obtained from liver biopsies of the patients. This study led to the discovery of key circulating factors
10 (or biomarkers) indicative of NASH and of its severity or stage or activity.

Accordingly, a first aspect of the invention relates to a method for the diagnosis of non-alcoholic steatohepatitis (NASH) or for determining the activity, the stage, or the severity of NASH in a subject, comprising the measure of the level of at least one blood, serum or plasma circulating micro-RNA
15 marker of NASH and at least one other blood, serum or plasma circulating marker of liver damage.

Preferably, this first aspect of the invention relates to a method for the diagnosis of non-alcoholic steatohepatitis (NASH) or for determining the activity, the stage, or the severity of NASH in a subject, comprising the step of measuring the level of at least one marker selected from either group (i), (ii) or
20 (iii), and at least one marker selected from either group (iv) or (v) defined as follows:

- (i) alpha 2 macroglobulin
- (ii) glycated haemoglobin (HbA1c), fasting glucose level, fructosamine level
- (iii) N-terminal pro-peptide of collagen type III
- (iv) hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p), hsa-miR-122 (in
25 particular hsa-miR-122-5p) and hsa-miR-192 (in particular hsa-miR-192-5p)
- (v) hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).

In a particular embodiment of the first aspect, the method comprises the implementation of the following measures from a blood, serum or plasma sample of said subject:

- 30 (i) determining the level of alpha 2 macroglobulin;
- (ii) determining the level of at least one marker selected in the group consisting of HbA1c, fasting glucose level, fructosamine level;
- (iii) determining the level of N-terminal pro-peptide of collagen type III;
- (iv) determining the level of at least one marker selected in the group consisting of hsa-miR-34,
35 hsa-miR-122 and hsa-miR-192; and
- (v) determining the level of hsa-miR-200;

wherein an increase in said levels relative to that of a control sample is indicative of NASH and/or of the activity, the stage, or the severity of NASH.

In a further embodiment of the first aspect of the invention:

- 5 - measure (ii) comprises determining the level of at least HbA1c; and/or
- measure (iv) comprises determining the level of at least hsa-miR-34.

In another aspect, the method of the first aspect comprises determining the level of alpha 2 macroglobulin, HbA1c, N-terminal pro-peptide of collagen type III, hsa-miR-34 and hsa-miR-200.

10

In yet another embodiment of the first aspect, a NASH score is calculated from the levels determined in measures (i) to (v) and wherein NASH is diagnosed and/or NASH stage or activity or severity is determined on the basis of said NASH score.

- 15 In another embodiment of the first aspect, the NASH score is calculated according to the following logistic function:

$$S = \frac{e^Y}{1 + e^Y}$$

wherein:

20

$$Y = k + a \cdot A + b \cdot B + c \cdot C + d \cdot D + f \cdot F$$

wherein:

S is the NASH score;

A is the level of alpha 2 macroglobulin in g/L;

B is the level of HbA1c in percent;

- 25 C is the level of N-terminal pro-peptide of collagen type III in ng/mL;

D is the level of hsa-miR-34 in Cq;

F is the level of hsa-miR-200 in Cq;

k is a number comprised between 5.26 and 36.76;

a is a number comprised between 0.11 and 1.07;

- 30 b is a number comprised between 0 and 1.02;

c is a number comprised between 0 and 0.23;

d is a number comprised between -0.77 and 0;

f is a number comprised between -0.78 and 0.

- 35 In a specific variant of this embodiment:

k is a number comprised between 5.26 and 36.76;

- a is a number comprised between 0.11 and 1.07;
b is a number comprised between 0.001 and 1.02;
c is a number comprised between 0.001 and 0.23;
d is a number comprised between -0.77 and -0.001;
5 f is a number comprised between -0.78 and -0.04.

In a further embodiment of the first aspect, a NASH score higher than a threshold value comprised between 0.102 and 0.6465 is indicative of a severe NASH, or of a moderate or high NASH activity.

In a variant of this embodiment:

- 10 k is equal to 25.13;
a is equal to 0.60;
b is equal to 0.48;
c is equal to 0.1;
d is equal to -0.43; and
15 f is equal to -0.45;
and wherein the threshold value is comprised between 0.102 and 0.462, and is more particularly equal to 0.23.

In an alternative variant of this embodiment:

- k is equal to 21.35;
20 a is equal to 0.62;
b is equal to 0.55;
c is equal to 0.14;
d is equal to -0.38; and
f is equal to -0.39;
25 and wherein the threshold value is comprised between 0.2319 and 0.6465, and is more particularly equal to 0.4343

- A second aspect of the invention relates to a method for classifying a subject as a receiver or non receiver of a treatment for non-alcoholic steatohepatitis (NASH), comprising the measure of the level
30 of at least one blood, serum or plasma circulating micro-RNA marker of NASH and at least one other blood, serum or plasma circulating marker of liver damage.

- Preferably, this second aspect of the invention relates to a method for classifying a subject as a receiver or non receiver of a treatment for non-alcoholic steatohepatitis (NASH), comprising the step
35 of measuring the level of at least one marker selected from either group (i), (ii) or (iii), and at least one marker selected from either group (iv) or (v) defined as follows:

(i) alpha 2 macroglobulin

- (ii) glycated haemoglobin (HbA1c), fasting glucose level, fructosamine level
- (iii) N-terminal pro-peptide of collagen type III
- (iv) hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p), hsa-miR-122 (in particular hsa-miR-122-5p) and hsa-miR-192 (in particular hsa-miR-192-5p)
- (v) hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).

In a particular embodiment of the second aspect of the invention, the method comprises the implementation of the following measures from a blood, serum or plasma sample of said subject:

- (i) determining the level of alpha 2 macroglobulin;
- (ii) determining the level of at least one marker selected in the group consisting of HbA1c, fasting glucose level, fructosamine level;
- (iii) determining the level of N-terminal pro-peptide of collagen type III;
- (iv) determining the level of at least one marker selected in the group consisting of hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p), hsa-miR-122 (in particular hsa-miR-122-5p) and hsa-miR-192 (in particular hsa-miR-192-5p); and
- (v) determining the level of hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p);

wherein a NASH score is calculated from the levels determined in measures (i) to (v) and wherein the subject is classified as a receiver or non receiver of the treatment for NASH based on said score.

In a further embodiment of the second aspect:

- measure (ii) comprises determining the level of at least HbA1c; and/or
- measure (iv) comprises determining the level of at least hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p).

In another embodiment of the second aspect, the method comprises determining the level of alpha 2 macroglobulin, HbA1c, N-terminal pro-peptide of collagen type III, hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p) and hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).

In yet another embodiment of the second aspect, the NASH score is calculated according to the following logistic function:

$$S = \frac{e^Y}{1 + e^Y}$$

wherein:

$$Y = k + a * A + b * B + c * C + d * D + f * F$$

wherein:

S is the NASH score;

A is the level of alpha 2 macroglobulin in g/L;

B is the level of HbA1c in percent;

5 C is the level of N-terminal pro-peptide of collagen type III in ng/mL;

D is the level of hsa-miR-34 in Cq;

F is the level of hsa-miR-200 in Cq;

k is a number comprised between 5.26 and 36.76;

a is a number comprised between 0.11 and 1.07;

10 b is a number comprised between 0 and 1.02;

c is a number comprised between 0 and 0.23;

d is a number comprised between -0.77 and 0;

f is a number comprised between -0.78 and 0.

15 In a variant of this embodiment:

k is a number comprised between 5.26 and 36.76;

a is a number comprised between 0.11 and 1.07;

b is a number comprised between 0.001 and 1.02;

c is a number comprised between 0.001 and 0.23;

20 d is a number comprised between -0.77 and -0.001;

f is a number comprised between -0.78 and -0.04.

In yet another embodiment of the second aspect, the subject is classified as a receiver of the treatment if S is higher than a threshold value comprised between 0.102 and 0.6465.

25 In a variant of this embodiment:

k is equal to 25.13;

a is equal to 0.6;

b is equal to 0.48;

c is equal to 0.1;

30 d is equal to -0.43; and

f is equal to -0.45;

and wherein the threshold value is comprised between 0.102 and 0.462, and is more particularly equal to 0.23.

35 In an alternative variant of this embodiment:

k is equal to 21.35;

a is equal to 0.62;

b is equal to 0.55;

c is equal to 0.14;

d is equal to -0.38; and

f is equal to -0.39;

- 5 and wherein the threshold value is comprised between 0.2319 and 0.6465, and is more particularly equal to 0.4343.

A third aspect of the invention relates to a method for calculating a NASH score which is indicative of non-alcoholic steatohepatitis (NASH), or of the activity, the stage, or the severity of NASH in a
10 subject, wherein said score is based on the measure of at least one marker selected from either group (i), (ii) or (iii), and at least one marker selected from either group (iv) or (v) defined as follows:

(i) alpha 2 macroglobulin

(ii) glycated haemoglobin (HbA1c), fasting glucose level, fructosamine level

(iii) N-terminal pro-peptide of collagen type III

15 (iv) hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p), hsa-miR-122 (in particular hsa-miR-122-5p) and hsa-miR-192 (in particular hsa-miR-192-5p)

(v) hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).

In an embodiment of the third aspect, the following (i) to (v) measures are determined from a blood,
20 serum or plasma sample of said subject:

(i) the level of alpha 2 macroglobulin;

(ii) the level of at least one marker selected in the group consisting of HbA1c, fasting glucose level, fructosamine level;

(iii) the level of N-terminal pro-peptide of collagen type III;

25 (iv) the level of at least one marker selected in the group consisting of hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p), hsa-miR-122 (in particular hsa-miR-122-5p) and hsa-miR-192 (in particular hsa-miR-192-5p); and

(v) the level of hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).

30 In a particular embodiment of the third aspect, the method comprises determining the level of alpha 2 macroglobulin, HbA1c, N-terminal pro-peptide of collagen type III, hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p) and hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p), wherein the NASH score is calculated according to the following logistic function:

35

$$S = \frac{e^Y}{1 + e^Y}$$

wherein:

$$Y=k+a*A+b*B+c*C+d*D+f*F$$

wherein:

S is the NASH score;

- 5 A is the level of alpha 2 macroglobulin in g/L;
 B is the level of HbA1c in percent;
 C is the level of N-terminal pro-peptide of collagen type III in ng/mL;
 D is the level of hsa-miR-34 in Cq;
 F is the level of hsa-miR-200 in Cq;
- 10 k is a number comprised between 5.26 and 36.76;
 a is a number comprised between 0.11 and 1.07;
 b is a number comprised between 0 and 1.02;
 c is a number comprised between 0 and 0.23;
 d is a number comprised between -0.77 and 0;
- 15 f is a number comprised between -0.78 and 0.
- In a specific variant of this embodiment:
 k is a number comprised between 5.26 and 36.76;
 a is a number comprised between 0.11 and 1.07;
 b is a number comprised between 0.001 and 1.02;
- 20 c is a number comprised between 0.001 and 0.23;
 d is a number comprised between -0.77 and -0.001;
 f is a number comprised between -0.78 and -0.04.

In another specific variant of this embodiment:

- 25 k is equal to 25.13;
 a is equal to 0.6;
 b is equal to 0.48;
 c is equal to 0.1;
 d is equal to -0.43; and
- 30 f is equal to -0.45.

In yet another specific variant of this embodiment:

- k is equal to 21.35;
 a is equal to 0.62;
- 35 b is equal to 0.55;
 c is equal to 0.14;
 d is equal to -0.38; and

f is equal to -0.39.

In a fourth aspect, the invention relates to a kit comprising means for determining the level of:

- (i) alpha 2 macroglobulin;
- 5 (ii) at least one marker selected in the group consisting of HbA1c, fasting glucose level, fructosamine level;
- (iii) N-terminal pro-peptide of collagen type III;
- (iv) at least one marker selected in the group consisting of has-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p), hsa-miR-122 (in particular hsa-miR-122-5p) and hsa-miR-192 (in particular hsa-miR-192-5p); and
- 10 (v) hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).

In an embodiment, the kit of the fourth aspect comprises means for determining the level of alpha 2 macroglobulin, HbA1c, N-terminal pro-peptide of collagen type III, hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p) and hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).


In a fifth aspect of the invention, the invention relates to a method for treating non-alcoholic steatohepatitis (NASH) in a subject, comprising:

- 20 (a) diagnosing NASH or determining the activity, the stage, or the severity of NASH in a subject according to the method described above;
- (b) administering a treatment for NASH to said subject based on step (a).

In a sixth aspect, the invention relates to a method for treating non-alcoholic steatohepatitis (NASH) in a subject, comprising:

- 25 (a) classifying said subject according to the method described above;
- (b) administering a treatment for NASH to the subject if said subject is classified as receiver of the treatment according to step (a).

30 In an embodiment of the fifth and sixth aspects, treatment comprises administering to the subject a compound of formula (I):



wherein:

X1 represents a halogen, a R1, or G1-R1 group;

A represents a CH=CH or a CH₂-CH₂ group;

X2 represents a G2-R2 group;

G1 and G2, identical or different, represent an atom of oxygen or sulfur;

5 R1 represents a hydrogen atom, an unsubstituted alkyl group, an aryl group or an alkyl group that is substituted by one or more halogen atoms, an alkoxy or an alkylthio group, cycloalkyl groups, cycloalkylthio groups or heterocyclic groups;

R2 represents an alkyl group substituted by at least a -COOR₃ group, wherein R₃ represents a hydrogen atom, or an alkyl group that is substituted or not by one or more halogen atoms, cycloalkyl groups, or heterocyclic groups.

10 R4 and R5, identical or different, representing an alkyl group that is substituted or not by one or more halogen atoms, cycloalkyl groups, heterocyclic groups; or a pharmaceutically acceptable salt thereof.

15 In another particular embodiment of the fifth and sixth aspects, the compound is selected in the group consisting of 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxy phenyl]prop-2-en-1-one, 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-isopropoxy carbonyldimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyldimethylmethoxyphenyl] prop-2-en-1-one, 1-[4-trifluoromethylphenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyl dimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-trifluoromethylphenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-trifluoromethyl oxyphenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyldimethylmethoxy phenyl] prop-2-en-1-one, 1-[4-trifluoromethoxyphenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxyphenyl]prop-2-en-1-one, 2-[2,6-dimethyl-4-[3-[4-(methylthio)phenyl]-3-oxo-propyl] phenoxy]-2-methylpropanoic acid, and 2-[2,6-dimethyl-4-[3-[4-(methylthio) phenyl]-3-oxo-propyl]phenoxy]-2-methyl-propanoic acid isopropyl ester; or a pharmaceutically acceptable salt thereof.

25 In another particular embodiment of the fifth and sixth aspects, the compound is 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxy phenyl]prop-2-en-1-one or a pharmaceutically acceptable salt thereof.

30 In another particular embodiment of the fifth and sixth aspects, the treatment comprises administering to the subject a compound of Table 1 as shown below.

35 Other aspects and embodiments will be apparent from the following detailed description.

LEGENDS OF THE FIGURES

Figures 1a to 1g show comparison of the NASH scoring according to the method of the invention derived from the Bootstrap model (Figure 1a) with existing scores.

5 DETAILED DESCRIPTION OF THE INVENTION

The invention relates to a method for the diagnosis of non-alcoholic steatohepatitis (NASH) or for determining the activity, the stage, or the severity of NASH in a subject. It also relates to a method for classifying a subject as a receiver or non receiver of a treatment for non-alcoholic steatohepatitis
10 (NASH). This classification may also be the base for further determining whether a subject should undergo further confirmation of NASH by methods known in the art such as by undergoing liver biopsy.

The invention stems from the identification of a defined set of circulating markers that, when
15 considered collectively, are indicative of NASH and/or of the activity, the stage, or the severity of NASH in the subject tested for NASH or for its severity or stage or activity. The invention relates in particular to a method for the diagnosis of non-alcoholic steatohepatitis (NASH) or for determining the activity, the stage, or the severity of NASH in a subject, combining the determination of the level of circulating micro-RNA(s) (miRNA(s)) and other circulating markers of liver damage.

20 Furthermore, the data collected by the inventors were processed according to two different biostatistical approaches leading to the determination of a score that is indicative of the disease and/or its severity or stage or activity, thereby providing a powerful, accurate and highly predictive tool for the physician to easily determine whether a subject is suffering from NASH, or to determine the stage
25 or activity or severity of the disease. This score, otherwise termed "NASH score" in the following description, is an invaluable tool for both the practitioner and the subjects in need thereof, as it allows the diagnosis or the determination of the severity or stage or activity of NASH using a non-invasive procedure, i.e. without relying on liver biopsy.

30 The methods of the invention comprise the measure of at least one marker which is a micro-RNA marker and at least one other circulating markers of liver damage. The methods of the invention more particularly comprise the measure of the level of at least one marker selected from groups (i), (ii), and (iii), and at least one marker selected from groups (iv) and (v) below.

35 In a particular embodiment, the methods of the invention comprise the measure of the level of:
- at least one marker selected in groups (i), (ii) and (iii) and at least one marker from each groups (iv) and (v);

- at least one marker selected from each groups (i) and (ii) or (ii) and (iii) or (i) and (iii), and at least one marker selected from groups (iv) and (v);
- at least one marker selected from each groups (i), (ii) and (iii) and at least one marker selected from groups (iv) and (v);
- 5 - at least one marker in each groups of markers (i), (ii), (iii), (iv) and (v) below, i.e.:
 - the marker listed in (i); and
 - at least one marker listed in (ii); and
 - the marker listed in (iii); and
 - at least one marker listed in (iv);
 - 10 - the marker listed in (v).

The groups of markers are defined as follows:

- (i) alpha 2 macroglobulin
- (ii) glycated haemoglobin (HbA1c), fasting glucose level, fructosamine level
- 15 (iii) N-terminal pro-peptide of collagen type III
- (iv) hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p), hsa-miR-122 (in particular hsa-miR-122-5p) and hsa-miR-192 (in particular hsa-miR-192-5p)
- (v) hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).
- 20 According to the present invention "non-alcoholic steatohepatitis" or "NASH" is defined as the presence of liver steatosis, hepatocyte ballooning and liver inflammation. To this basic definition of the disease, NASH may further include liver fibrosis.

According to the present invention, the subject is a mammal, in particular a human subject. The invention, thanks to its non-invasive nature, may be implemented on any subject, such as a subject with no known or suspected predisposition to NASH. However, in a particular embodiment, the subject is a subject at risk of having NASH or of developing NASH in the future, such as a subject having obesity, diabetes, suffering from the metabolic syndrome, or having elevated liver enzymes, or who was previously diagnosed with non-alcoholic fatty liver disease (or liver steatosis). The subject may also be a subject with already identified NASH, the method of the invention thereby allowing determining the risks of evolution of the disease towards hepatocarcinoma or liver transplantation or a cardiovascular disease.

The levels of the markers measured in the present invention are determined from a body fluid of the subject, which may in particular be a blood, more particularly serum or plasma sample.

The level of the markers identified by the inventors may be determined by conventional methods well known in the art, such as immunoassays (e.g. ELISA), or molecular and biochemical assays (quantitative PCR, colorimetric assays), or analytical methods (such as mass spectrometry), depending on the biomarker type (such as a protein, a micro-RNA, glucose level, ...). When the tested marker is a micro-RNA, its measure may be carried out according to a number of methods well-known in the art, such as that presented in the examples below. Briefly, the measures are carried out from total RNA extracted from a plasma sample, in particular a cell-free, platelet-free plasma sample. An appropriate internal control (such as a micro-RNA of known sequence and quantity, e.g. *C. elegans* miR-39) may be added to the sample before RNA extraction. Cq values are determined using quantitative PCR. Commercial kits are available for conducting such assays. For example, the Taqman miRNA qRT-PCR assay: Taqman MicroRNA Reverse transcription Kit, TaqMan MicroRNA Assay 20X, and TaqMan Universal Master Mix II (Applied Biosystems) may be used according to the manufacturer's instructions. Reverse transcription may be performed using readily available PCR systems, such as the GeneAmp® PCR System 9700 thermal cycler (Applied Biosystems), with appropriate cycling parameters such as 16°C for 30 minutes followed by 42°C for 30 minutes and 85°C for 5 minutes before holding at 4°C. The reverse transcription may be implemented in the multiplexed format. Quantitative PCT is then conducted using a quantitative PCR system such as the CFX96™ Real-Time System (C1000 Touch™ Thermal Cycler, BioRad). Cycling conditions may be the following: 95°C for 10 minutes followed by 95°C for 15 sec and 60°C for 60 sec for a total of 50 cycles and 30°C for 30 sec. Cq determination mode may be set of Regression in the quantitative PCR system. In a particular embodiment, the Cq value determined according to the method of the invention is the Cq value which is obtainable using the above specific parameters and material. Cq values of samples may be excluded from the analysis if values are above the maximum Cq of the standard curve of each miRNA. The standard curve may be used to assess the reaction efficiency. Serial dilution may be performed over eight points starting from the most concentrated cDNA sample, to ensure the standard curve covers all potential template concentrations that may be encountered during the study. The standard curve may be constructed by plotting the log of the starting quantity of the template against the Cq values obtained.

According to the present invention, a subject is diagnosed having NASH, or is diagnosed as potentially having NASH or likely to have NASH, when all the levels of markers selected and measured in each of group (i) to (v) are increased when compared to reference values obtained from a control (or reference) sample. The reference sample may correspond to a body fluid sample, such as a blood, serum or plasma sample obtained from one or more subjects, such as two or more, who do not have NAFLD or NASH.

In the present invention, the levels of biomarkers measured as described above may also be used to determine a disease score, otherwise named NASH score. The disease score may be compared with a threshold value that distinguishes between low, moderate, and high NASH activity, or moderate and severe NASH. According to the present invention, this information may be advantageously used to determine whether a subject will receive or not a therapeutic treatment for NASH. Only those subjects with a moderate to high NASH activity or severe NASH will be treated. Accordingly, the present invention relates to a method for classifying a subject as a receiver or non receiver of a treatment for non-alcoholic steatohepatitis (NASH), said method comprising the determination of the levels of the markers as defined above, the calculation of a NASH score, and the classification of the subject as receiver or non receiver of the treatment based on said NASH score. In addition, the classification of the subject may also be used to determine a low NASH activity or a moderate NASH in the subject, and providing to the subject based on this classification diet and lifestyle recommendations to reverse NASH.

According to the present invention, those subjects classified as moderate to high NASH activity, or as severe NASH patients are considered as receivers, or potential receivers, of a treatment for NASH. In the context of the present invention, a moderate to high NASH activity or a severe NASH patient is defined as, or equivalent to, a patient presenting the following liver biopsy-derived grades:

- steatosis grade ≥ 1
- hepatocyte ballooning grade ≥ 1
- lobular inflammation grade ≥ 1
- NAS (NAFLD Activity Score) ≥ 4 (NAS is defined as the sum of the steatosis grade, hepatocyte ballooning grade and lobular inflammation grade)
- fibrosis stage ≥ 2 (such as a fibrosis equal to 2, 3 or 4, in particular 2 or 3)

In a particular embodiment, the NASH score of the invention is calculated from the following marker levels measured as provided above:

- level of alpha 2 macroglobulin (LVBP_A2M_GL); and
- level of HbA1c (LBBC_HBA1C_PERC); and
- level of N-terminal pro-peptide of collagen type III (LVB_P3NP); and
- level of hsa-miR-34 (in particular miR-34a, more particularly miR-34a-5p) (LVMK_HSA_MIR34A_MULTI1_CQ); and
- level of hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p) (LVMK_HSA_MIR200A_MULTI1_CQ).

In a further particular embodiment, the score is defined as a logistic function:

$$S = \frac{e^Y}{1 + e^Y}$$

wherein:

$$Y = k + a \cdot A + b \cdot B + c \cdot C + d \cdot D + f \cdot F$$

5 wherein:

S is the NASH score;

A is the level of alpha 2 macroglobulin in g/L;

B is the level of HbA1c in percent (e.g. B is equal to 10 if measured HbA1c percentage is 10%);

C is the level of N-terminal pro-peptide of collagen type III in ng/mL;

10 D is the level of hsa-miR-34 (in particular hsa-miR-34a , more particularly hsa-miR-34a-5p) in Cq;

F is the level of hsa-miR-200 (in particular c hsa-miR-200a , more particularly hsa-miR-200a-3p) in Cq;

k is a number comprised between 5.26 and 36.76;

a is a number comprised between 0.11 and 1.07;

15 b is a number comprised between 0 and 1.02;

c is a number comprised between 0 and 0.23;

d is a number comprised between -0.77 and 0;

f is a number comprised between -0.78 and 0.

20 In a further particular embodiment, the score is calculated according to the above logistic function, wherein:

k is a number comprised between 5.26 and 36.76;

a is a number comprised between 0.11 and 1.07;

b is a number comprised between 0.001 and 1.02;

25 c is a number comprised between 0.001 and 0.23;

d is a number comprised between -0.77 and -0.001;

f is a number comprised between -0.78 and -0.04.

30 The NASH score thus is the probability of having a moderate to high NASH activity or a severe NASH. If S as determined for a subject is greater or equal to a threshold value, the subject is classified as to be treated for NASH, or to potentially be treated for NASH. If S is lower than a threshold value, the subject is classified as not to be treated, and/or the subject is classified as receiver, or potential receiver, of diet and lifestyle advices for managing its low NASH activity or moderate NASH.

35 According to a particular embodiment, the subject is classified as a receiver, or potential receiver, of a treatment if S is higher than or equal to a threshold value comprised between 0.102 and 0.6465.

In a particular embodiment, derived from the median model as described in the experimental part of this application:

k is equal to 25.13;

5 a is equal to 0.6;

b is equal to 0.48;

c is equal to 0.1;

d is equal to -0.43; and

f is equal to -0.45;

10 and wherein the threshold value is comprised between 0.102 and 0.462, and is more particularly equal to 0.23.

In a particular embodiment, derived from the bootstrap model as described in the experimental part of this application:

15 k is equal to 21.35;

a is equal to 0.62;

b is equal to 0.55;

c is equal to 0.14;

d is equal to -0.38; and

20 f is equal to -0.39;

and wherein the threshold value is comprised between 0.2319 and 0.6465, and is more particularly equal to 0.4343.

The present invention also relates to a kit comprising means for determining the level of:

25 (i) alpha 2 macroglobulin;

(ii) at least one marker selected in the group consisting of HbA1c, fasting glucose level, fructosamine level;

(iii) N-terminal pro-peptide of collagen type III;

30 (iv) at least one marker selected in the group consisting of hsa-miR-34 (in particular hsa-miR-34a, more particularly miR-34a-5p), hsa-miR-122 (in particular hsa-miR-122-5p) and hsa-miR-192 (in particular hsa-miR-192-5p); and

(v) hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).

35 In a particular embodiment of the invention, said kit comprises means for determining the level of alpha 2 macroglobulin, HbA1c, N-terminal pro-peptide of collagen type III, hsa-miR-34 (in particular hsa-miR-34a, more particularly miR-34a-5p) and hsa-miR-200a (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).

The kit of the invention is useful for implementing the methods described above may further optionally include instructions for implementing said methods. The kit may comprise reagents and buffers appropriate for conducting measures of the levels of markers identified above. In particular, the kit may comprise antibodies specific for a protein to be quantified, and/or primers useful for quantifying micro-RNA levels, as well-known in the art.

The present invention further relates to a method for treating NASH in a subject, comprising:

- (a) diagnosing NASH or determining the activity, the stage, or the severity of NASH in a subject according to the methods provided above;
- (b) administering a treatment for NASH to said subject based on step (a).


The present invention further relates to a method for treating NASH in a subject, comprising:

- (a) classifying said subject according to the method of classification as defined above;
- (b) administering a treatment for NASH to the subject if said subject is classified as receiver of the treatment according to step (a).

The present invention further relates to a method for managing low NASH activity, or moderate NASH in a subject, comprising:

- (a) classifying said subject according to the method of classification as defined above;
- (b) providing diet and lifestyle recommendations to the subject if said subject is classified as non receiver of the treatment according to step (a).

The method for treating NASH according to the present invention comprise the administration of one or more compound(s) to the subject in need thereof such as a compound of formula (I):



wherein:

X1 represents a halogen, a R1, or G1-R1 group;

A represents a CH=CH or a CH2-CH2 group;

X2 represents a G2-R2 group;

G1 and G2, identical or different, represent an atom of oxygen or sulfur;

R1 represents a hydrogen atom, an unsubstituted alkyl group, an aryl group or an alkyl group that is substituted by one or more halogen atoms, an alkoxy or an alkylthio group, cycloalkyl groups, cycloalkylthio groups or heterocyclic groups;

R2 represents an alkyl group substituted by at least a –COOR₃ group, wherein R₃ represents a hydrogen atom, or an alkyl group that is substituted or not by one or more halogen atoms, cycloalkyl groups, or heterocyclic groups.

R4 and R5, identical or different, representing an alkyl group that is substituted or not by one or more halogen atoms, cycloalkyl groups, heterocyclic groups;

or a pharmaceutically acceptable salt thereof.

In a particular embodiment of this method for the treatment of NASH, the compound of formula (I) is selected in the group consisting of 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxy phenyl]prop-2-en-1-one, 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-isopropoxy carbonyldimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyldimethylmethoxyphenyl] prop-2-en-1-one, 1-[4-trifluoromethylphenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyl dimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-trifluoromethylphenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-trifluoromethyl oxyphenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyldimethylmethoxy phenyl] prop-2-en-1-one, 1-[4-trifluoromethoxyphenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxyphenyl]prop-2-en-1-one, 2-[2,6-dimethyl-4-[3-[4-(methylthio)phenyl]-3-oxo-propyl] phenoxy]-2-methylpropanoic acid, and 2-[2,6-dimethyl-4-[3-[4-(methylthio) phenyl]-3-oxo-propyl]phenoxy]-2-methyl-propanoic acid isopropyl ester; or a pharmaceutically acceptable salt thereof. In a further particular embodiment of the invention, the compound of formula (I) is 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxy phenyl]prop-2-en-1-one or a pharmaceutically acceptable salt thereof.

Other compounds that might be used in the method for the treatment of NASH of the present invention may include, without limitation, a compound selected from the following classes and specific compounds of table 1.

Table 1

Class	Compound
Alanine transaminase stimulator	HU-002
11-beta hydroxysteroid dehydrogenase (HSD) inhibitor	RO5093151
Aldosterone antagonist, anti-oxydant	Spiro-lactone /Vit E
Amino-acid	SAMe (S-adenosyl-L-methionine)
Analog of human leptin	Metreleptin
Anti-apoptotic caspase inhibitor	Emricasan (IDN-6556)

Antibiotic	Rifaximin
Anti-oxidant	MitoQ Vitamin E
Anti-Oxidant PPAR γ	Vitamin E Pioglitazone
Anti-oxydant	Cysteamine (Cystagon-EC) Siliphos
Anti-oxydant Biguanide	Vitamin E Metformin
Bovine Colostrum Inflammation	IMM 124-E
Caspase Inib	GS-9450
SCD1 inhibitors	Aramchol
DGAT1 inhibitors	LCQ908
Fish oil supplement	Omega 3 PUFA Opti-EPA
Flavonoid	MEDOX (Anthocyanin) Silymarin
FXR agonist	OCA (obeticholic acid)
galectin inhibitor	GR-MD-02
GLP-1 Agonist	Exenatide
GLP1 mimetic	Liraglutide
Immuno-modulator	Oral anti α -CD3 EGS21
Bile acids	UDCA
Iron Chelator	Deferasirox ICL 670 Exjade®
Lipid Lowering agent	Ezetimibe
musculotropic vasodilator	Pentoxifylline
Omega 3	Lovaza EPA-E (MND-21)

EXAMPLES

5

Sample collection and analysis

Clinical study

- 10 The clinical trial (phase 2 GOLDEN-505 trial in NASH (GFT505-212-7 - NCT01694849) is a multicentre, randomized, double blind, placebo-controlled study to evaluate the efficacy and safety of Elafibranor (1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxy phenyl]prop-2-en-1-one) once daily on steatohepatitis in patients with non-alcoholic steatohepatitis (NASH).
- 15 Liver biopsy was performed to confirm the diagnosis of NASH after appropriate exclusion of liver disease of other etiology. NASH was diagnosed as steatohepatitis evaluated by liver biopsy within 6 months before randomization. Steatohepatitis confirmation was based on central reading of liver

biopsies. NASH patients were defined with a NAS ≥ 3 including steatosis grade ≥ 1 and hepatocyte ballooning ≥ 1 and lobular inflammation ≥ 1 .

The study was approved by appropriate regulatory bodies at each participating center and all patients had given consent for participation in medical research.

5

Blood sampling and Laboratory testing

Blood samples were collected according to the Central Laboratory Protocol and Manual– Genfit – GFT505-212-7.

10

According to the study protocol, following analyses were performed.

HEMATOLOGY includes hemoglobin, hematocrit, RBC count, leukocytes, differential leukocyte count (neutrophils, lymphocytes, eosinophils, monocytes, basophils -abs. and % values), platelets and reticulocytes.

15

BIOCHEMISTRY Panel I includes plasma glucose, triglycerides (TG), creatinine, creatinine clearance, gamma-glutamyltransferase (GGT), aspartate aminotransferase (AST), alanine aminotransferase (ALT), creatine phosphokinase (CPK), alkaline phosphatase, thyroid stimulating hormone (TSH) and HbA1c.

20

BIOCHEMISTRY Panel II includes plasma glucose, creatinine, creatinine clearance, total protein, albumin, sodium, potassium, chloride, calcium, uric acid, urea expressed as blood urea nitrogen (BUN), aspartate aminotransferase (AST), alanine aminotransferase (ALT), gamma-glutamyltransferase (GGT), alkaline phosphatase, creatine phosphokinase (CPK), bilirubin total, bilirubin conjugated, C-reactive protein (hsCRP), AST/ALT Ratio and HbA1c.

25

URINALYSIS includes:

- Dipstick analysis (specific gravity, pH, RBC, leukocytes, glucose, protein, ketones, bilirubin, urobilinogen and nitrite)
- Microscopy analysis includes RBC, WBC, casts, crystals, bacteria, epithelial cells and yeasts.
- Chemistry analysis (albumin and creatinine)

30

SEROLOGY includes HIV ab I/ II, HCV ab, HCV RNA (only tested upon receipt of HCV RNA Visit samples and in case of 'reactive' or 'indeterminate' result for HCV Ab) and HbsAg.

35

LIPID PANEL includes triglycerides (TG), total cholesterol, non HDL-C (calculation), high-density lipoprotein cholesterol (HDL-C), low density lipoprotein (LDL-C) (calculation), calculated very low density lipoprotein cholesterol (VLDL-C) (calculation), apolipoprotein AI (ApoAI) and apolipoprotein B (ApoB).

5

URINE CHEMISTRY includes alpha-1-microglobulin, beta-N-acetylglucosaminidase(beta-NAG) and neutrophil-gelatinase associated lipocalin(N-Gal)

10

SAFETY MARKERS includes homocysteine, NT-ProBNP, Troponin T, Cystatin C, and Beta2-microglobulin.

GLYCEMIC AND OTHER LIPIDIC PARAMETERS includes leptin, insulin, homeostatic model assesment (HOMA-IR), serum glucose (for calculation of HOMA-IR), fructosamine, C-peptide and free fatty acids (FFA).

15

INFLAMMATORY MARKERS includes haptoglobin, fibrinogen, tumor necrosis factor alpha (TNF- α), interleukine 6 (IL-6) and plasminogen activator inhibitor 1 (PAI-1) Ag (citrate).

20

LIVER MARKERS includes cytokeratin-18 (CK18)(M65 & M30), adinopectin, ferritin, alpha2 macroglobulin, FGF19 & FGF21, hyaluronic acid (Advia centaur, reagentiaproced by Siemens Belgium and charged to Genfit in pass-through), N-terminal pro-peptide of collagen type III (PIIINP) (Advia centaur, reagentia proced by Siemens Belgium) and tissue inhibitor of matrix metalloprotease-1 (TIMP-1) (Advia centaur, reagentiaproced by Siemens).

25

Sample Collection & Storage

Blood samples used in this biomarker study were drawn from patients of the 505.212.7 study before treatment period. Written, informed consent for collection, storage and use of additional samples was obtained from every patient.

30

Blood collected in citrate containing tubes 2.7mL was processed by separating cell-free plasma from blood cells within 15 minutes of collection by centrifugation at 1,500xg for 15 minutes. The supernatant plasma was transferred to a new tube. Tubes were kept at -70°C.

35

To proceed to RNA extraction, plasma tubes were then centrifuged at 13,000xg for 2 min to pellet and remove the platelets. The supernatant platelet-free plasma was transferred to a new tube, frozen in liquid nitrogen and stored at -80°C.

Total RNA extraction and quantitation of miRNA

Total RNA with preserved miRNAs was extracted from 400 µl of platelet-free plasma by miRNeasy extraction kit (*miRNeasy Serum/Plasma Kit (cat. no. 217184)*) and using a plasma/QIAzol ratio of 1:5 according to the manufacturer's instructions. Synthetic spiked-in *C. elegans* miR-39 was added to the samples [3,125 fmoles] prior to RNA extraction as internal control of RNA extraction process. The elution was performed in 18µl of elution buffer.

Expression of mature miRNAs was detected according to the manufacturer's instructions using the Taqman miRNA qRT-PCR Assay: TaqMan MicroRNA Reverse transcription Kit (Ref: 4366596, Applied Biosystems, Carlsbad, CA), TaqMan MicroRNA Assay 20X (Ref: 4440887, Applied Biosystems) and TaqMan Universal Master Mix II (Ref: 4440040, Applied Biosystems).

Reverse Transcription were performed using a GeneAmp® PCR System 9700 thermal cycler (Ref: 200005, Applied Biosystems) with cycling conditions of 16°C for 30 minutes followed by 42°C for 30 minutes and 85°C for 5 minutes before holding at 4°C. The Reverse transcription was multiplexed with 4 miRNA specific-reverse primers in order to conserve RNA sample.

Quantitative PCR were performed using a CFX96™ Real-Time System (C1000 Touch™ Thermal Cycler, BioRad) with cycling conditions of 95°C for 10 minutes followed by 95°C for 15 sec and 60°C for 60 sec for a total of 50 cycles and 30°C for 30 sec.

The sequences of specific primers for miRNA of potential interest were chosen according to the following table 2.

Table 2: sequences of tested miRNA

Mature miRNA	Assay id (Life Technologies)	Sequence of mature miRNA	mirBase number	Supplier
cel-miR-39-3p	000200	UCACCGGGUGUAAAUCAGCUUG (SEQ ID NO:1)	MIMAT0000010	Life technologies
hsa-miR-122-5p	002245	UGGAGUGUGACAAUGGUGUUU G (SEQ ID NO:2)	MIMAT0000421	Life technologies
hsa-miR-34a-5p	000426	UGGCAGUGUCUUAGCUGGUUGU (SEQ ID NO:3)	MIMAT0000255	Life technologies
hsa-miR-200a-3p	000502	UAACACUGUCUGGUAACGAUGU (SEQ ID NO:4)	MIMAT0000318	Life technologies
hsa-miR-29b-3p	000413	UAGCACCAUUUGAAAUCAGUGU U (SEQ ID NO:5)	MIMAT0000100	Life technologies
hsa-miR-192-5p	000491	CUGACCUAUGAAUUGACAGCC	MIMAT0000222	Life technologies

		(SEQ ID NO:6)		
hsa-miR-221-3p	000524	AGCUACAUUGUCUGCGGGUUU C (SEQ ID NO:7)	MIMAT0000278	Life technologies
hsa-miR-103a-3p	000439	AGCAGCAUUGUACAGGGCUAUG A (SEQ ID NO:8)	MIMAT0000101	Life technologies
hsa-miR-155-5p	002623	UUA AUGCUAAUCGUGAUAGGGG U (SEQ ID NO:9)	MIMAT0000646	Life technologies
hsa-miR-199a -5p	000498	CCCAGUGUUCAGACUACCUGUU C (SEQ ID NO:10)	MIMAT0000231	Life technologies

Data used in the construction of the algorithm were in Cq format. The Cq Determination mode was Regression.

5 Statistical analysis

Objective and definition

The objective of these analyses was to discover biomarkers that can be related to the identification of NASH patients to be treated. Patients to be treated (TBT) were defined as:

- Fibrosis stage ≥ 2 and
- NAS ≥ 4 and
- Lobular inflammation grade ≥ 1 and
- Steatosis grade ≥ 1 and
- Ballooning grade ≥ 1 .

Other patients were stated as not to be treated (NTBT). For the analysis, TBT patients were categorized as 1 and NTBT as 0 in the response variable. As shown above, explicative variables encompassed a wide range of biomarkers measured in blood (hematology, biochemistry, coagulation, liver markers, circulating miRNA) or in urinary (dipstick, sediment) samples, as long as demographic (age, sex, race), region (study centre, country, continent) or medical (diabetes) recordings.

Dataset management

The dataset used in this study was coming from the Golden-505 trial and was initially composed of 274 patients and 121 variables. The dataset management included the following steps:

1. Removal of 28 patients without miRNAs measurement
2. Removal of 9 variables with more than 10 missing values
3. Removal of 7 patients with extreme values and 28 with remaining missing values

It resulted in a dataset with 211 patients and 112 variables that were used to test collinearity between explicative variables.

Test of collinearity

5

Pearson correlation was calculated two by two between quantitative variables. When two variables presented a correlation superior to 0.7, univariate test of difference in their mean in relation to the response variable defining patients TBT were done. The selected variable was the more significant. The collinear and selected variables are presented in Table 2.

10

Thirty-five unique collinear variables were removed and the final dataset used in the analysis included 77 variables (patient id, response variable, 5 histological grades and 70 explicative variables) and 211 patients (95 TBT and 116 NTBT).

Table 3. Selected and dropped variables in collinearity test

<i>Selected variables in collinear variables</i>	<i>Dropped variables in collinear variables</i>		
GLP_HOMA_IR	GLP_INS_PMOLL	GLP_CPEPTID_NMOLL	
INMK_CK18_M65_UL	LVBK_CK18_M30_PMOLL	LBBC_AST_UL	LBBC_ALT_UL
INMK_FGF_21_PGML	SFMK_HOMOCYS_UMOLL		
LBBC_HBA1C_PERC	LBBC_GLU_FAST_MMOLL	GLP_GLU_FAST_MMOLL	GLP_FRUCTO_UMOLL
LBHA_BASO_10_9_L	LBHA_BASO_PERC		
LBHA_EOSINO_10_9_L	LBHA_EOSINO_PERC		
LBHA_HEMATO_VV	LBHA_HEMO_GL		
LBHA_LEUKO_10_9_L	LBHA_NEUTRO_10_9_L	LBHA_RBC_COUNT_10_12_L	
LBHA_LYMPOH_PERC	LBHA_NEUTRO_PERC		
LBHA_PROTHR_TIME_SEC	LBHA_INR	LBHA_PROTHR_TIME_PERC	
LBHA_RETICULO_10_9_L	LBHA_RETICULO_RBC		
LBHP_LIPOPRINT_HDL_MGDL	LBHP_HDL_MMOLL	LBHP_APOA1_MGDL	
LBHP_LIPOPRINT_IDLA_MGDL	LBHP_LIPOPRINT_LDL1_MGDL		
LBHP_LIPOPRINT_IDLB_MGDL	LBHP_NHDL_MMOLL		
LBHP_LIPOPRINT_IDLC_MGDL	LBHP_TC_MMOLL		
LBHP_LIPOPRINT_LDL2_MGDL	LBHP_APOB_MGDL	LBHP_APOB_MGDL	LBHP_NHDL_MMOLL
LBHP_TG_MMOLL	LBHP_VLDL_MMOLL	LBHP_LDL_MMOLL	
LVMK_HSA_MIR103_MULTII_CQ	LVMK_HSA_MIR155_MULTII_CQ	LBHP_LIPOPRINT_VLDL_MGDL	
LVMK_HSA_MIR34A_MULTII_CQ	LVMK_HSA_MIR122_MULTII_CQ	LVMK_HSA_MIR199A_MULTII_CQ	LVMK_HSA_MIR221_MULTII_CQ
URI_SPECGRV	URI_CREAT_GL	LVMK_HSA_MIR192_MULTII_CQ	
WAIST	BMI	WEIGHT	

Variables	Parameter
GLP_HOMA_IR	Homeostatic Model Assesment (HOMA-IR)
INMK_CK18_M65_UL	Cytokeratin-18 (CK18) (M65)
INMK_FGF_21_PGML	FGF21
LBBC_HBA1C_PERC	HbA1c
LBHA_BASO_10_9_L	Basophils (abs.)
LBHA_EOSINO_10_9_L	Eosinophils (abs)
LBHA_HEMATO_VV	Hematocrit
LBHA_LEUKO_10_9_L	Leukocytes
LBHA_LYMPHO_PERC	Lymphocytes (%)
LBHA_PROTHR_TIME_SEC	Prothrombin time (sec)
LBHA_RETICULO_10_9_L	Reticulocytes (abs.)
LBLP_LIPOPRINT_HDL_MGDL	High-density lipoprotein (HDL)
LBLP_LIPOPRINT_IDLA_MGDL	Intermediate-density lipoprotein
LBLP_LIPOPRINT_IDLB_MGDL	Intermediate-density lipoprotein
LBLP_LIPOPRINT_IDLC_MGDL	Intermediate-density lipoprotein
LBLP_LIPOPRINT_LDL2_MGDL	low Density Lipoprotein (LDL)
LBLP_TG_MMOLL	Triglycerides
lvmk_hsa_mir103_multi1_cq	miRNA103
lvmk_hsa_mir34a_multi1_cq	miRNA34a
URI_SPECGRAV	Specific gravity
GLP_INS_PMOLL	insulin
LVBP_CK18_M30_PMOLL	cytokeratin-18 (CK18) (M30)
SFMK_HOMOCYS_UMOLL	homocysteine
LBBC_GLU_FAST_MMOLL	glucose
LBHA_BASO_PERC	basophils (%)
LBHA_EOSINO_PERC	eosinophils (%)
LBHA_HEMO_GL	hemoglobin
LBHA_NEUTRO_10_9_L	neutrophils (abs)
LBHA_NEUTRO_PERC	neutrophils (%)
LBHA_INR	INR
LBHA_RETICULO_RBC	reticulocytes (% of RBC)
LBLP_HDL_MMOLL	high-density lipoprotein cholesterol (HDL-C)
LBLP_LIPOPRINT_LDL1_MGDL	low Density Lipoprotein (LDL)
LBLP_NHDL_MMOLL	non HDL-C
LBLP_TC_MMOLL	total cholesterol
LBLP_APOB_MGDL	apolipoprotein B (Apo B)
LBLP_VLDL_MMOLL	calculated Very Low Density Lipoprotein Cholesterol (VLDL-C)
lvmk_hsa_mir155_multi1_cq	miR155
lvmk_hsa_mir122_multi1_cq	miR122
URI_CREAT_GL	creatinine
bmi	body mass index
GLP_CPEPTID_NMOLL	C-peptide

LBBC_AST_UIL	Aspartate Aminotransferase (AST)
GLP_GLU_FAST_MMOLL	glucose
LBHA_RBC_COUNT_10_12_L	RBC Count
LBHA_PROTHR_TIME_PERC	Prothrombin time
LBLP_APOA1_MGDL	Apolipoprotein AI (Apo AI)
LBLP_LDL_MMOLL	Low Density Lipoprotein (LDL-C)
LBLP_LIPOPRINT_VLDL_MGDL	Very Low Density Lipoprotein Cholesterol
lvmk_hsa_mir199a_multi1_cq	miR199a
lvmk_hsa_mir192_multi1_cq	miR192
LBBC_ALT_UIL	Alanine Aminotransferase (ALT)
GLP_FRUCTO_UMOLL	Fructosamine
lvmk_hsa_mir221_multi1_cq	miR221

Median model

- 5 Dataset was partitioned one million times in 2/3 training set and in 1/3 verification set, stratified on TBT and NTBT patient distribution. In these different possible partitions, 27,477 presented no significant difference between the training and verification sets in all explicative variables and histological grades and were used for analysis.
- 10 A logistic regression was performed between the response variable and 70 explicative variables, followed by a variable selection using the Akaike Information Criterion (AIC), in the 27,477 training sets. It resulted in 27,477 different coefficients for the 70 variables. The median model was created by combining variables with a median coefficient different from zero (Table 4).
- 15 **Table 4.** Results of median model (^{cor}: collinear variable)

<i>Variables included in median model</i>	<i>Coefficients</i>	<i>1st-3rd quartiles</i>
Constant	25.13	18.39; 32.66
LVBP_A2M_GL	0.60	0.43; 0.77
LBBC_HBA1C_PERC ^{cor}	0.48	0.001; 0.69
LVB_P3NP	0.10	0.001; 0.14
lvmk_hsa_mir34a_multi1_cq ^{cor}	-0.43	-0.60; -0.001
lvmk_hsa_mir200a_multi1_cq	-0.45	-0.58; -0.31

LVBP_A2M_GL: Alpha 2 macroglobuline; LBBC_HBA1C_PERC: HbA1c; LVB_P3NP: N-terminal pro-peptide of collagen type III; lvmk_hsa_mir34a_multi1_cq: miR-34a; lvmk_hsa_mir200a_multi1_cq: miR-200a.

- 20 The model was then verified by calculating the Area Under the ROC curve (AUC) for each of the 27,477 sets. The AUC was of 0.82 (95% CI: 0.72-0.91). Using the median AUC, 3 thresholds were defined for predicting patients as TBT or NTBT:

1. The threshold closest to the point with 1-Specificity=0% and Sensitivity=100% equalled 0.23. The corresponding contingency table is presented in Table 5a and produced the following indices:

- Total accuracy: 80.28%
- Sensitivity: 81.25%
- Specificity: 79.49%
- PPV (Positive Predictive Value): 76.47%
- NPV (Negative Predictive Value): 83.78%

2. The threshold giving a sensitivity superior to 90% equalled 0.102. The corresponding contingency table is presented in Table 5b and produced the following indices:

- Total accuracy: 63.38 %
- Sensitivity: 90.63 %
- Specificity: 41.03 %
- PPV (Positive Predictive Value): 55.77 %
- NPV (Negative Predictive Value): 84.21 %

3. The threshold giving a specificity superior to 90% equalled 0.462. The corresponding contingency table is presented in Table 5c and produced the following indices:

- Total accuracy: 70.42 %
- Sensitivity: 43.75 %
- Specificity: 92.31 %
- PPV (Positive Predictive Value): 82.35 %
- NPV (Negative Predictive Value): 66.67 %

Table 5a. Contingency table of TBT/NTBT patients observed and predicted by the median model and a threshold of 0.23.

	<i>Predicted</i>		
		NTBT (0)	TBT (1)
<i>Observed</i>	NTBT (0)	31	8
	TBT (1)	6	26

Table 5b. Contingency table of TBT/NTBT patients observed and predicted by the median model and a threshold of 0.102.

	<i>Predicted</i>		
		NTBT (0)	TBT (1)
<i>Observed</i>	NTBT (0)	16	23
	TBT (1)	3	29

Table 5c. Contingency table of TBT/NTBT patients observed and predicted by the median model and a threshold of 0.462.

	<i>Predicted</i>		
		NTBT (0)	TBT (1)
<i>Observed</i>	NTBT (0)	36	3
	TBT (1)	18	14

Bootstrap model

Using all 211 patients a logistic regression was calculated as full model using all 70 explicative variables and an optimal model is obtained by selecting variables with the AIC (Table 6). The significance of variable coefficients was then tested using 1000 bootstrap samples (Table 6).

Table 6. Results of bootstrap model (^{cor}: collinear variable)

<i>Variables included in median model</i>	<i>Coefficients</i>	<i>95% CI</i>
Constant	21.35	5.26; 36.76
LVBP_A2M_GL	0.62	0.11; 1.07
LBBC_HBA1C_PERC ^{cor}	0.55	0.13; 1.02
LVB_P3NP	0.14	0.03; 0.23
lvmk_hsa_mir34a_multi1_cq ^{cor}	-0.38	-0.77; -0.001
lvmk_hsa_mir200a_multi1_cq	-0.39	-0.78; -0.04

Of outmost importance, it is very satisfying to note that the variables identified with the bootstrap model are the same as those identified with the median model described above. This observation strengthens the statistical relevance and confidence of these variables as predictive of NASH and of its stage or severity.

The optimal model is verified by calculating the AUC in 1000 bootstrap samples. The AUC was of 0.82 (95% CI: 0.76-0.87). Using the AUC of the optimal model, 3 thresholds were defined for predicting patients as TBT or NTBT:

1. The threshold closest to the point with 1-Specificity=0% and Sensitivity=100%, equalled 0.4343. The corresponding contingency table is presented in Table 7a and produced the following indices:

- Total accuracy: 75.36%
- Sensitivity: 74.74%
- Specificity: 75.86%
- PPV: 71.72%

- NPV (Negative Predictive Value): 78.57%
- 2. The threshold giving a sensitivity superior to 90% equalled 0.2319. The corresponding contingency table is presented in Table 7b and produced the following indices:
 - Total accuracy: 65.88%
 - Sensitivity: 90.53%
 - Specificity: 45.69%
 - PPV (Positive Predictive Value): 57.72%
 - NPV (Negative Predictive Value): 85.48%
- 3. The threshold giving a specificity superior to 90% equalled 0.6465. The corresponding contingency table is presented in Table 7c and produced the following indices:
 - Total accuracy: 73.46%
 - Sensitivity: 52.63%
 - Specificity: 90.52%
 - PPV (Positive Predictive Value): 81.97%
 - NPV (Negative Predictive Value): 70.00%

Table 7a. Contingency table of TBT/NTBT patients observed and predicted by the bootstrap model and a threshold of 0.4343.

	<i>Predicted</i>		
		NTBT (0)	TBT (1)
<i>Observed</i>	NTBT (0)	88	28
	TBT (1)	24	71

Table 7b. Contingency table of TBT/NTBT patients observed and predicted by the bootstrap model and a threshold of 0.2319.

	<i>Predicted</i>		
		NTBT (0)	TBT (1)
<i>Observed</i>	NTBT (0)	53	63
	TBT (1)	9	86

Table 7c. Contingency table of TBT/NTBT patients observed and predicted by the bootstrap model and a threshold of 0.6465.

	<i>Predicted</i>		
		NTBT (0)	TBT (1)
<i>Observed</i>	NTBT (0)	105	11
	TBT (1)	45	50

Comparison to other NASH scoring methods

Methods to establish NASH scores were previously described in literature:

- the Angulo Index (Angulo et al, 2007)
- 5 - ELF (Guha et al, 2008) (see also the ELF specification sheet
<http://www.healthcare.siemens.com/clinical-specialities/liver-disease/elf-test-now-avail>)
- Fibrotest (Ratziu et al, 2006)
- Fibrometer (Cales et al, 2009)
- FLI (Bedogni et al, 2006)
- 10 - Steatosis Grade or SteatoTest (Poynard et al, 2005)

The inventors thus evaluated the NASH score of the invention to a number of these previous methods. Figures 1a to 1g show comparison of the NASH scoring according to the method of the invention derived from the Bootstrap model (Figure 1a) with existing scores.

15

The data presented herein show a spectacular improvement of the accuracy of the identification of NASH patients, or potential NASH patients, determination of NASH activity, stage or severity. This finding is of outmost importance and will be an invaluable tool to improve patient management efficiency.

20

FURTHER OBJECTS DISCLOSED HEREIN

The present disclosure relates in particular to the following objects:

25

1. A method for the diagnosis of non-alcoholic steatohepatitis (NASH) or for determining the activity, the stage, or the severity of NASH in a subject, comprising the measure of the level of at least one blood, serum or plasma circulating micro-RNA marker of NASH and at least one other blood, serum or plasma circulating marker of liver damage.

30

2. The method according to object 1, comprising the step of measuring the level of at least one marker selected from either group (i), (ii) or (iii), and at least one marker selected from either group (iv) or (v) defined as follows:

- (i) alpha 2 macroglobulin
- 35 (ii) glycated haemoglobin (HbA1c), fasting glucose level, fructosamine level
- (iii) N-terminal pro-peptide of collagen type III

- (iv) hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p), hsa-miR-122 (in particular hsa-miR-122-5p) and hsa-miR-192 (in particular hsa-miR-192-5p)
- (v) hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).

- 5 3. The method according to object 1 or 2, comprising the implementation of the following measures from a blood, serum or plasma sample of said subject:
- (i) determining the level of alpha 2 macroglobulin;
 - (ii) determining the level of at least one marker selected in the group consisting of HbA1c, fasting glucose level, fructosamine level;
 - 10 (iii) determining the level of N-terminal pro-peptide of collagen type III;
 - (iv) determining the level of at least one marker selected in the group consisting of hsa-miR-34, hsa-miR-122 and hsa-miR-192; and
 - (v) determining the level of hsa-miR-200;
- wherein an increase in said levels relative to that of a control sample is indicative of NASH and/or of
- 15 the activity, the stage, or the severity of NASH.

4. The method according to object 2 to 3, wherein:
- measure (ii) comprises determining the level of at least HbA1c; and/or
 - measure (iv) comprises determining the level of at least hsa-miR-34.

20

5. The method according to any one of objects 1 or 4, comprising determining the level of alpha 2 macroglobulin, HbA1c, N-terminal pro-peptide of collagen type III, hsa-miR-34 and hsa-miR-200.

6. The method according to any one of objects 2 to 5, wherein a NASH score is calculated from the
- 25 levels determined in measures (i) to (v) and wherein NASH is diagnosed and/or NASH stage or activity or severity is determined on the basis of said NASH score.

7. The method according to object 6, wherein a NASH score is calculated according to the following logistic function:

30

$$S = \frac{e^Y}{1 + e^Y}$$

wherein:

$$Y = k + a * A + b * B + c * C + d * D + f * F$$

wherein:

- 35 S is the NASH score;

A is the level of alpha 2 macroglobulin in g/L;

B is the level of HbA1c in percent;

C is the level of N-terminal pro-peptide of collagen type III in ng/mL;

D is the level of hsa-miR-34 in Cq;

F is the level of hsa-miR-200 in Cq;

- 5 k is a number comprised between 5.26 and 36.76;
a is a number comprised between 0.11 and 1.07;
b is a number comprised between 0 and 1.02;
c is a number comprised between 0 and 0.23;
d is a number comprised between -0.77 and 0;
10 f is a number comprised between -0.78 and 0.

8. The method according to object 7, wherein:

k is a number comprised between 5.26 and 36.76;

a is a number comprised between 0.11 and 1.07;

- 15 b is a number comprised between 0.001 and 1.02;
c is a number comprised between 0.001 and 0.23;
d is a number comprised between -0.77 and -0.001;
f is a number comprised between -0.78 and -0.04.

- 20 9. The method according to object 7 or 8, wherein a NASH score higher than a threshold value comprised between 0.102 and 0.6465 is indicative of a severe NASH, or of a moderate or high NASH activity.

10. The method according to object 9, wherein:

- 25 k is equal to 25.13;
a is equal to 0.60;
b is equal to 0.48;
c is equal to 0.1;
d is equal to -0.43; and
30 f is equal to -0.45;
and wherein the threshold value is comprised between 0.102 and 0.462, and is more particularly equal to 0.23.

11. The method according to object 9, wherein:

- 35 k is equal to 21.35;
a is equal to 0.62;
b is equal to 0.55;

c is equal to 0.14;

d is equal to -0.38; and

f is equal to -0.39;

and wherein the threshold value is comprised between 0.2319 and 0.6465, and is more particularly equal to 0.4343

12. A method for classifying a subject as a receiver or non receiver of a treatment for non-alcoholic steatohepatitis (NASH), comprising the measure of the level of at least one blood, serum or plasma circulating micro-RNA marker of NASH and at least one other blood, serum or plasma circulating marker of liver damage.

13. The method according to object 12, comprising the step of measuring the level of at least one marker selected from either group (i), (ii) or (iii), and at least one marker selected from either group (iv) or (v) defined as follows:

(i) alpha 2 macroglobulin

(ii) glycated haemoglobin (HbA1c), fasting glucose level, fructosamine level

(iii) N-terminal pro-peptide of collagen type III

(iv) hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p), hsa-miR-122 (in particular hsa-miR-122-5p) and hsa-miR-192 (in particular hsa-miR-192-5p)

(v) hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).

14. The method according to object 12 or 13, comprising the implementation of the following measures from a blood, serum or plasma sample of said subject:

(i) determining the level of alpha 2 macroglobulin;

(ii) determining the level of at least one marker selected in the group consisting of HbA1c, fasting glucose level, fructosamine level;

(iii) determining the level of N-terminal pro-peptide of collagen type III;

(iv) determining the level of at least one marker selected in the group consisting of hsa-miR-34, hsa-miR-122 and hsa-miR-192; and

(v) determining the level of hsa-miR-200;

wherein a NASH score is calculated from the levels determined in measures (i) to (v) and wherein the subject is classified as a receiver or non receiver of the treatment for NASH based on said score.

15. The method according to object 13 or 14, wherein:

- measure (ii) comprises determining the level of at least HbA1c; and/or

- measure (iv) comprises determining the level of at least hsa-miR-34.

16. The method according to any one of objects 13 to 15, comprising determining the level of alpha 2 macroglobulin, HbA1c, N-terminal pro-peptide of collagen type III, hsa-miR-34 and hsa-miR-200.

5 17. The method according to object 16, wherein the NASH score is calculated according to the following logistic function:

$$S = \frac{e^Y}{1 + e^Y}$$

wherein:

$$Y = k + a \cdot A + b \cdot B + c \cdot C + d \cdot D + f \cdot F$$

10 wherein:

S is the NASH score;

A is the level of alpha 2 macroglobulin in g/L;

B is the level of HbA1c in percent;

C is the level of N-terminal pro-peptide of collagen type III in ng/mL;

15 D is the level of hsa-miR-34 in Cq;

F is the level of hsa-miR-200 in Cq;

k is a number comprised between 5.26 and 36.76;

a is a number comprised between 0.11 and 1.07;

b is a number comprised between 0 and 1.02;

20 c is a number comprised between 0 and 0.23;

d is a number comprised between -0.77 and 0;

f is a number comprised between -0.78 and 0.

18. The method according to object 17, wherein:

25 k is a number comprised between 5.26 and 36.76;

a is a number comprised between 0.11 and 1.07;

b is a number comprised between 0.001 and 1.02;

c is a number comprised between 0.001 and 0.23;

d is a number comprised between -0.77 and -0.001;

30 f is a number comprised between -0.78 and -0.04.

19. The method according to any one of objects 17 or 18, wherein the subject is classified as a receiver of the treatment if S is higher than a threshold value comprised between 0.102 and 0.6465.

35 20. The method according to object 19, wherein:

k is equal to 25.13;

a is equal to 0.6;

b is equal to 0.48;

c is equal to 0.1;

d is equal to -0.43; and

5 f is equal to -0.45;

and wherein the threshold value is comprised between 0.102 and 0.462, and is more particularly equal to 0.23.

21. The method according to object 19, wherein:

10 k is equal to 21.35;

a is equal to 0.62;

b is equal to 0.55;

c is equal to 0.14;

d is equal to -0.38; and

15 f is equal to -0.39;

and wherein the threshold value is comprised between 0.2319 and 0.6465, and is more particularly equal to 0.4343.

22. A method for calculating a NASH score which is indicative of non-alcoholic steatohepatitis (NASH), or of the activity, the stage, or the severity of NASH in a subject, wherein said score is based on the measure of at least one marker selected from either group (i), (ii) or (iii), and at least one marker selected from either group (iv) or (v) defined as follows:

(i) alpha 2 macroglobulin

(ii) glycated haemoglobin (HbA1c), fasting glucose level, fructosamine level

25 (iii) N-terminal pro-peptide of collagen type III

(iv) hsa-miR-34 (in particular hsa-miR-34a, more particularly hsa-miR-34a-5p), hsa-miR-122 (in particular hsa-miR-122-5p) and hsa-miR-192 (in particular hsa-miR-192-5p)

(v) hsa-miR-200 (in particular hsa-miR-200a, more particularly hsa-miR-200a-3p).

30 23. The method according to object 22, wherein the following (i) to (v) measures are determined from a blood, serum or plasma sample of said subject:

(i) the level of alpha 2 macroglobulin;

(ii) the level of at least one marker selected in the group consisting of HbA1c, fasting glucose level, fructosamine level;

35 (iii) the level of N-terminal pro-peptide of collagen type III;

(iv) the level of at least one marker selected in the group consisting of hsa-miR-34, hsa-miR-122 and hsa-miR-192; and

(v) the level of hsa-miR-200.

24. The method of object 23, comprising determining the level of alpha 2 macroglobulin, HbA1c, N-terminal pro-peptide of collagen type III, hsa-miR-34 and hsa-miR-200.

5

25. The method of object 24, wherein the NASH score is calculated according to the following logistic function:

$$S = \frac{e^Y}{1 + e^Y}$$

10 wherein:

$$Y = k + a * A + b * B + c * C + d * D + f * F$$

wherein:

S is the NASH score;

A is the level of alpha 2 macroglobulin in g/L;

15 B is the level of HbA1c in percent;

C is the level of N-terminal pro-peptide of collagen type III in ng/mL;

D is the level of hsa-miR-34 in Cq;

F is the level of hsa-miR-200 in Cq;

k is a number comprised between 5.26 and 36.76;

20 a is a number comprised between 0.11 and 1.07;

b is a number comprised between 0 and 1.02;

c is a number comprised between 0 and 0.23;

d is a number comprised between -0.77 and 0;

f is a number comprised between -0.78 and 0.

25

26. The method according to object 25, wherein:

k is a number comprised between 5.26 and 36.76;

a is a number comprised between 0.11 and 1.07;

b is a number comprised between 0.001 and 1.02;

30 c is a number comprised between 0.001 and 0.23;

d is a number comprised between -0.77 and -0.001;

f is a number comprised between -0.78 and -0.04

27. The method according to object 25, wherein:

35 k is equal to 25.13;

a is equal to 0.6;

b is equal to 0.48;
c is equal to 0.1;
d is equal to -0.43; and
f is equal to -0.45.

5

28. The method according to object 25, wherein:

k is equal to 21.35;
a is equal to 0.62;
b is equal to 0.55;

10

c is equal to 0.14;
d is equal to -0.38; and
f is equal to -0.39.

29. A kit comprising means for determining the level of:

15

- (i) alpha 2 macroglobulin;
- (ii) at least one marker selected in the group consisting of HbA1c, fasting glucose level, fructosamine level;
- (iii) N-terminal pro-peptide of collagen type III;
- (iv) at least one marker selected in the group consisting of has-miR-34, hsa-miR-122 and hsa-miR-192; and
- (v) hsa-miR-200.

20

30. The kit according to object 29, comprising means for determining the level of alpha 2 macroglobulin, HbA1c, N-terminal pro-peptide of collagen type III, hsa-miR-34 and hsa-miR-200.

25

31. A method for treating non-alcoholic steatohepatitis (NASH) in a subject, comprising:

- (a) diagnosing NASH or determining the activity, the stage, or the severity of NASH in a subject according to the method of any one of objects 1 to 11;
- (b) administering a treatment for NASH to said subject based on step (a).


30

32. A method for treating non-alcoholic steatohepatitis (NASH) in a subject, comprising:

- (a) classifying said subject according to the method of any one of objects 12 to 21;
- (b) administering a treatment for NASH to the subject if said subject is classified as receiver of the treatment according to step (a).

35

33. The method according to object 31 or 32, wherein the treatment comprises administering to the subject a compound of formula (I):



wherein:

X1 represents a halogen, a R1, or G1-R1 group;

A represents a CH=CH or a CH2-CH2 group;

5 X2 represents a G2-R2 group;

G1 and G2, identical or different, represent an atom of oxygen or sulfur;

R1 represents a hydrogen atom, an unsubstituted alkyl group, an aryl group or an alkyl group that is substituted by one or more halogen atoms, an alkoxy or an alkylthio group, cycloalkyl groups, cycloalkylthio groups or heterocyclic groups;

10 R2 represents an alkyl group substituted by at least a -COOR3 group, wherein R3 represents a hydrogen atom, or an alkyl group that is substituted or not by one or more halogen atoms, cycloalkyl groups, or heterocyclic groups.

R4 and R5, identical or different, representing an alkyl group that is substituted or not by one or more halogen atoms, cycloalkyl groups, heterocyclic groups;

15 or a pharmaceutically acceptable salt thereof.

34. The method according to object 33, wherein the compound is selected in the group consisting of 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxy phenyl]prop-2-en-1-one, 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-isopropoxy carbonyldimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyldimethylmethoxyphenyl] prop-2-en-1-one, 1-[4-trifluoromethylphenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyl dimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-trifluoromethylphenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-trifluoromethyl oxyphenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyldimethylmethoxy phenyl] prop-2-en-1-one, 1-[4-trifluoromethyloxyphenyl]-3-[3,5-dimethyl-4-carboxydimethylmethyl oxyphenyl]prop-2-en-1-one, 2-[2,6-dimethyl-4-[3-[4-(methylthio)phenyl]-3-oxo-propyl] phenoxy]-2-methylpropanoic acid, and 2-[2,6-dimethyl-4-[3-[4-(methylthio) phenyl]-3-oxo-propyl]phenoxy]-2-methyl-propanoic acid isopropyl ester; or a pharmaceutically acceptable salt thereof.

30 35. The method according to object 33 or 34, wherein the compound is 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxy phenyl]prop-2-en-1-one or a pharmaceutically acceptable salt thereof.

36. The method according to object 31 or 32, wherein the treatment comprises administering to the subject a compound of Table 1.

5 REFERENCES

- Angulo P, Hui JM, Marchesini G, Bugianesi E, George J, Farrell GC, Enders F, Saksena S, Burt AD, Bida JP, Lindor K, Sanderson SO, Lenzi M, Adams LA, Kench J, Therneau TM, Day CP (2007) The NAFLD fibrosis score: a noninvasive system that identifies liver fibrosis
10 in patients with NAFLD. *Hepatology* **45**: 846-854
- Bedogni G, Bellentani S, Miglioli L, Masutti F, Passalacqua M, Castiglione A, Tiribelli C (2006) The Fatty Liver Index: a simple and accurate predictor of hepatic steatosis in the general population. *BMC Gastroenterol* **6**: 33
15
- Cales P, Laine F, Boursier J, Deugnier Y, Moal V, Oberti F, Hunault G, Rousselet MC, Hubert I, Laafi J, Ducluzeaux PH, Lunel F (2009) Comparison of blood tests for liver fibrosis specific or not to NAFLD. *J Hepatol* **50**: 165-173
- 20 Guha IN, Parkes J, Roderick P, Chattopadhyay D, Cross R, Harris S, Kaye P, Burt AD, Ryder SD, Aithal GP, Day CP, Rosenberg WM (2008) Noninvasive markers of fibrosis in nonalcoholic fatty liver disease: Validating the European Liver Fibrosis Panel and exploring simple markers. *Hepatology* **47**: 455-460
- 25 Poynard T, Ratziu V, Naveau S, Thabut D, Charlotte F, Messous D, Capron D, Abella A, Massard J, Ngo Y, Munteanu M, Mercadier A, Manns M, Albrecht J (2005) The diagnostic value of biomarkers (SteatoTest) for the prediction of liver steatosis. *Comp Hepatol* **4**: 10
- 30 Ratziu V, Massard J, Charlotte F, Messous D, Imbert-Bismut F, Bonyhay L, Tahiri M, Munteanu M, Thabut D, Cadranet JF, Le Bail B, de Ledinghen V, Poynard T (2006) Diagnostic value of biochemical markers (FibroTest-FibroSURE) for the prediction of liver fibrosis in patients with non-alcoholic fatty liver disease. *BMC Gastroenterol* **6**: 6

CLAIMS

1. A method for the diagnosis of non-alcoholic steatohepatitis (NASH) or for determining the activity, the stage, or the severity of NASH in a subject, comprising the step of measuring the blood, serum, or plasma level of:

- (i) alpha 2 macroglobulin;
- (ii) glycated haemoglobin (HbA1c);
- (iii) N-terminal pro-peptide of collagen type III;
- (iv) hsa-miR-34a-5p; and
- (v) hsa-miR-200a-3p.

2. The method according to claim 1, comprising the implementation of the following measures from a blood, serum or plasma sample of said subject:

- (i) determining the level of alpha 2 macroglobulin;
- (ii) determining the level of HbA1c, ;
- (iii) determining the level of N-terminal pro-peptide of collagen type III;
- (iv) determining the level of hsa-miR-34a-5p; and
- (v) determining the level of hsa-miR-200a-3p;

wherein an increase in said levels relative to that of a control sample is indicative of NASH and/or of the activity, the stage, or the severity of NASH.

3. The method according to claim 1 or 2, wherein a NASH score is calculated from the levels determined in measures (i) to (v) and wherein NASH is diagnosed and/or NASH stage or activity or severity is determined on the basis of said NASH score.

4. The method according to claim 3, wherein a NASH score is calculated according to the following logistic function:

$$S = \frac{e^Y}{1 + e^Y}$$

wherein:

$$Y = k + a \cdot A + b \cdot B + c \cdot C + d \cdot D + f \cdot F$$

wherein:

S is the NASH score;

A is the level of alpha 2 macroglobulin in g/L;

B is the level of HbA1c in percent;

C is the level of N-terminal pro-peptide of collagen type III in ng/mL;

D is the level of hsa-miR-34a-5p in Cq;

F is the level of hsa-miR-200a-3p in Cq;

k is a number comprised between 5.26 and 36.76;

5 a is a number comprised between 0.11 and 1.07;

b is a number comprised between 0 and 1.02;

c is a number comprised between 0 and 0.23;

d is a number comprised between -0.77 and 0; and

f is a number comprised between -0.78 and 0.

10

5. The method according to claim 4, wherein a NASH score higher than a threshold value comprised between 0.102 and 0.6465 is indicative of a severe NASH, or of a moderate or high NASH activity.

6. A method for classifying a subject as a receiver or non receiver of a treatment for NASH,
15 comprising the step of measuring the blood, serum, or plasma level of:

(i) alpha 2 macroglobulin;

(ii) glycated haemoglobin (HbA1c);

(iii) N-terminal pro-peptide of collagen type III;

(iv) hsa-miR-34a-5p; and

20

(v) hsa-miR-200a-3p.

7. The method according to claim 6, comprising the implementation of the following measures from a blood, serum or plasma sample of said subject:

25

(i) determining the level of alpha 2 macroglobulin;

(ii) determining the level of HbA1c;

(iii) determining the level of N-terminal pro-peptide of collagen type III;

(iv) determining the level of hsa-miR-34a-5p; and

(v) determining the level of hsa-miR-200a-3p;

30

wherein a NASH score is calculated from the levels determined in measures (i) to (v) and wherein the subject is classified as a receiver or non receiver of the treatment for NASH based on said score.

8. The method according to claim 6 or 7, comprising determining the level of alpha 2 macroglobulin, HbA1c, N-terminal pro-peptide of collagen type III, hsa-miR-34a-5p and hsa-miR-200a-3p wherein
35 the NASH score is calculated according to the following logistic function:

$$S = \frac{e^Y}{1 + e^Y}$$

wherein:

$$Y = k + a \cdot A + b \cdot B + c \cdot C + d \cdot D + f \cdot F$$

wherein:

- 5 S is the NASH score;
 A is the level of alpha 2 macroglobulin in g/L;
 B is the level of HbA1c in percent;
 C is the level of N-terminal pro-peptide of collagen type III in ng/mL;
 D is the level of hsa-miR-34a-5p in Cq;
 10 F is the level of hsa-miR-200a-3p in Cq;
 k is a number comprised between 5.26 and 36.76;
 a is a number comprised between 0.11 and 1.07;
 b is a number comprised between 0 and 1.02;
 c is a number comprised between 0 and 0.23;
 15 d is a number comprised between -0.77 and 0;
 f is a number comprised between -0.78 and 0.

9. The method according to claim 8, wherein the subject is classified as a receiver of the treatment if S is higher than a threshold value comprised between 0.102 and 0.6465.

20

10. A method for calculating a NASH score which is indicative of non-alcoholic steatohepatitis (NASH), or of the activity, the stage, or the severity of NASH in a subject, wherein said score is based on the measure of the blood, serum, or plasma level of:

- (i) alpha 2 macroglobulin;
 25 (ii) glycated haemoglobin (HbA1c);
 (iii) N-terminal pro-peptide of collagen type III;
 (iv) hsa-miR-34a-5p; and
 (v) hsa-miR-200a-3p.

- 30 11. The method of claim 10, comprising determining the level of alpha 2 macroglobulin, HbA1c, N-terminal pro-peptide of collagen type III, hsa-miR-34a-5p and hsa-miR-200a-3p, wherein the NASH score is calculated according to the following logistic function:

35
$$S = \frac{e^Y}{1 + e^Y}$$

wherein:

$$Y=k+a*A+b*B+c*C+d*D+f*F$$

wherein:

S is the NASH score;

5 A is the level of alpha 2 macroglobulin in g/L;

B is the level of HbA1c in percent;

C is the level of N-terminal pro-peptide of collagen type III in ng/mL;

D is the level of hsa-miR-34a-5p in Cq;

F is the level of hsa-miR-200a-3p in Cq;

10 k is a number comprised between 5.26 and 36.76;

a is a number comprised between 0.11 and 1.07;

b is a number comprised between 0 and 1.02;

c is a number comprised between 0 and 0.23;

d is a number comprised between -0.77 and 0;

15 f is a number comprised between -0.78 and 0.

12. A kit comprising means for determining the level of:

(i) alpha 2 macroglobulin;

(ii) HbA1c;

20 (iii) N-terminal pro-peptide of collagen type III;

(iv) hsa-miR-34a-5p; and

(v) hsa-miR-200a-3p;


wherein the kit comprises as means antibodies specific for a protein to be quantified and/or primers useful for quantifying micro-RNA levels.

25

13. Use of a compound suitable for the treatment of non-alcoholic steatohepatitis (NASH) in the manufacture of a medicament for treating non-alcoholic steatohepatitis (NASH) in a subject, wherein the subject has been diagnosed according to the method of any one of claims 1 to 5, or classified according to the method of any one of claims 6 to 11.

30

14. The use according to claim 13, wherein the compound is a compound of formula (I):



wherein:

X1 represents a halogen, a R1, or G1-R1 group;

A represents a CH=CH or a CH₂-CH₂ group;

X2 represents a G2-R2 group;

G1 and G2, identical or different, represent an atom of oxygen or sulfur;

- 5 R1 represents a hydrogen atom, an unsubstituted alkyl group, an aryl group or an alkyl group that is substituted by one or more halogen atoms, an alkoxy or an alkylthio group, cycloalkyl groups, cycloalkylthio groups or heterocyclic groups;

R2 represents an alkyl group substituted by at least a –COOR₃ group, wherein R₃ represents a hydrogen atom, or an alkyl group that is substituted or not by one or more halogen atoms, cycloalkyl groups, or heterocyclic groups.

- 10 R4 and R5, identical or different, representing an alkyl group that is substituted or not by one or more halogen atoms, cycloalkyl groups, heterocyclic groups; or a pharmaceutically acceptable salt thereof.

- 15 15. The use according to claim 14, wherein the compound is selected in the group consisting of 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxy phenyl]prop-2-en-1-one, 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-isopropoxy carbonyldimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyldimethylmethoxyphenyl] prop-2-en-1-one, 1-[4-trifluoromethylphenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyl
- 20 dimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-trifluoromethylphenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxyphenyl]prop-2-en-1-one, 1-[4-trifluoromethyl oxyphenyl]-3-[3,5-dimethyl-4-tertbutyloxycarbonyldimethylmethoxy phenyl] prop-2-en-1-one, 1-[4-trifluoromethyloxyphenyl]-3-[3,5-dimethyl-4-carboxydimethylmethyl oxyphenyl]prop-2-en-1-one, 2-[2,6-dimethyl-4-[3-[4-(methylthio)phenyl]-3-oxo-propyl] phenoxy]-2-methylpropanoic acid, and 2-[2,6-dimethyl-4-[3-[4-(methylthio) phenyl]-3-oxo-propyl]phenoxy]-2-methyl-propanoic acid isopropyl ester; or a
- 25 pharmaceutically acceptable salt thereof.

16. The use for according to any one of claims 13 to 15, wherein said compound is 1-[4-methylthiophenyl]-3-[3,5-dimethyl-4-carboxydimethylmethoxyphenyl]prop-2-en-1-one or a
- 30 pharmaceutically acceptable salt thereof.