

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

(19) Weltorganisation für geistiges Eigentum
Internationales Büro



(43) Internationales Veröffentlichungsdatum
10. September 2004 (10.09.2004)

PCT

(10) Internationale Veröffentlichungsnummer
WO 2004/076454 A1

(51) Internationale Patentklassifikation⁷: **C07D 475/04**,
A61K 31/4375, 31/4985

(21) Internationales Aktenzeichen: PCT/EP2003/001935

(22) Internationales Anmeldedatum:
26. Februar 2003 (26.02.2003)

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme von
US): **BOEHRINGER INGELHEIM PHARMA GMBH
& CO KG** [DE/DE]; Binger Strasse 173, 55216 INGEL-
HEIM (DE).

(72) Erfinder; und

(75) Erfinder/Anmelder (nur für US): **HOFFMANN**,
Matthias [DE/DE]; Sonnenweg 17, 88441 MITTEL-
BIBERACH (DE). **GRAUERT**, Matthias [DE/DE];
Osterbergstrasse 10, 88400 BIBERACH (DE). **BRANDL**,
Trixi [DE/DE]; Königbergallee 9/1, 88400 BIBERACH
(DE). **BREITFELDER**, Steffen [DE/DE]; Weiher-
gasse 21, 88433 ASSMANNSHARDT (DE). **EICK-
MEIER**, Christian [DE/DE]; Ayestrasse 10/2, 88441

MITTELBIBERACH (DE). **STEEGMAIER**, Martin
[DE/AT]; Schloeglasse 12/B5, A-1120 WIEN (AT).
SCHNAPP, Gisela [DE/DE]; Esterbuch 5, 88400 BIB-
ERACH-RINDENMOOS (DE). **BAUM**, Anke [DE/AT];
Groisbach 13, A-2534 ALLAND (AT). **QUANT**, Jens,
Jürgen [DE/AT]; Hafergasse 32, A-2353 GUNTRAMS-
DORF (AT). **SOLCA**, Flavio [CH/AT]; Gesslgasse 10/6,
A-1230 WIEN (AT). **COLBATZKY**, Florian [DE/DE];
Beim Wiesental 16, 88441 STAFFLANGEN (DE).

(74) Gemeinsamer Vertreter: **BOEHRINGER INGEL-
HEIM PHARMA GMBH & CO KG**; Binger Strasse
173, 55216 Ingelheim (DE).


(81) Bestimmungsstaaten (national): AE, AG, AL, AM, AT,
AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR,
CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Bestimmungsstaaten (regional): ARIPO Patent (GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ,

[Fortsetzung auf der nächsten Seite]

(54) Title: DIHYDROPTERIDINONES, METHOD FOR THE PRODUCTION AND USE THEREOF IN THE FORM OF DRUGS

(54) Bezeichnung: DIHYDROPTERIDINONE, VERFAHREN ZU DEREN HERSTELLUNG UND DEREN VERWENDUNG
ALS ARZNEIMITTEL



(I)

(57) Abstract: The invention relates to novel dihydropteridi-
nones of general formula (I), wherein rests L, R¹-R⁵ have the
significance indicated in claims and a specification, in the iso-
mers thereof, in a method for producing and using said dihy-
dropteridinones in the form of drugs.


(57) Zusammenfassung: Die vorliegende Erfindung
betrifft neue Dihydropteridinone der allgemeinen Formel
(I), wobei die Reste L, R¹-R⁵ die in den Ansprüchen und der
Beschreibung genannten Bedeutungen haben, deren Isomere,
Verfahren zur Herstellung dieser Dihydropteridinone sowie
deren Verwendung als Arzneimittel.

WO 2004/076454 A1

80560pct

**Dihydropteridinones, method for the production and use thereof
in the form of drugs**

- 5 The present invention relates to new dihydropteridinones of general formula (I)



- 10 wherein the groups L, R¹, R², R³, R⁴ and R⁵ have the meanings given in the claims and specification, the isomers thereof, processes for preparing these dihydropteridinones and the use thereof as pharmaceutical compositions.

Background to the invention

- 15 Pteridinone derivatives are known from the prior art as active substances with an antiproliferative activity. WO 01/019825 describes the use of pteridinone derivatives for the treatment of neoplastic and viral diseases. The resistance of many types of tumours calls for the development of new pharmaceutical compositions for combating tumours.


20

The aim of the present invention is to prepare new compounds with antiinflammatory and antiproliferative activity.

Detailed description of the invention

Surprisingly it has been found that compounds of general formula (I) wherein the groups L and R¹ to R⁵ have the meanings given hereinafter act as inhibitors of specific cell cycle kinases. Thus, the compounds according to the invention may be used for example to treat diseases connected with the activity of specific cell cycle kinases and characterised by excessive or abnormal cell proliferation.

The present invention therefore relates to compounds of general formula (I)



wherein

R¹, R² which may be identical or different, denote hydrogen or optionally substituted C₁-C₆-alkyl,

or

R¹ and R² together denote a 2- to 5-membered alkyl bridge which may contain 1 to 2 heteroatoms,

R³ denotes hydrogen or a group selected from among optionally substituted C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, C₂-C₁₂-alkynyl and C₆-C₁₄-aryl, or a group selected from among optionally substituted and/or bridged C₃-C₁₂-cycloalkyl, C₃-C₁₂-cycloalkenyl, C₇-C₁₂-polycycloalkyl, C₇-C₁₂-polycycloalkenyl, C₅-

C₁₂-spirocycloalkyl, C₃-C₁₂-heterocycloalkyl which contains 1 to 2 heteroatoms, and C₃-C₁₂-heterocycloalkenyl which contains 1 to 2 heteroatoms, or

5 R¹ and R³ or R² and R³ together denote a saturated or unsaturated C₃-C₄-alkyl bridge which may contain 1 heteroatom,

R⁴ denotes a group selected from among hydrogen, -CN, hydroxy, -NR₆R₇ and halogen, or

10 a group selected from among optionally substituted C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₅-alkyloxy, C₂-C₅-alkenyloxy, C₂-C₅-alkynyloxy, C₁-C₆-alkylthio, C₁-C₆-alkylsulphoxo and C₁-C₆-alkylsulphonyl,

15 L denotes a linker selected from among optionally substituted C₂-C₁₀-alkyl, C₂-C₁₀-alkenyl, C₆-C₁₄-aryl, -C₂-C₄-alkyl-C₆-C₁₄-aryl, -C₆-C₁₄-aryl-C₁-C₄-alkyl, optionally bridged C₃-C₁₂-cycloalkyl and heteroaryl which contains 1 or 2 nitrogen atoms,

n denotes 0 or 1

m denotes 1 or 2

20

R⁵ denotes a group selected from among optionally substituted morpholinyl, piperidinyl, piperazinyl, piperazinylcarbonyl, pyrrolidinyl, tropanyl, R⁸-diketomethylpiperazinyl, sulphoxomorpholinyl, sulphonylmorpholinyl, thiomorpholinyl, -NR⁸R⁹ and azacycloheptyl,

25 R⁶, R⁷ which may be identical or different, denote hydrogen or C₁-C₄-alkyl, and

R⁸, R⁹ denote unsubstituted nitrogen substituents at R⁵, which may be identical or different, denote either hydrogen or a group selected from among C₁-C₆-alkyl,

-C₁-C₄-alkyl-C₃-C₁₀-cycloalkyl, C₃-C₁₀-cycloalkyl, C₆-C₁₄-aryl,

30 -C₁-C₄-alkyl-C₆-C₁₄-aryl, pyranyl, pyridinyl, pyrimidinyl, C₁-C₄-alkyloxycarbonyl, C₆-C₁₄-arylcarbonyl, C₁-C₄-alkylcarbonyl, C₆-C₁₄-arylmethyloxycarbonyl, C₆-C₁₄-arylsulphonyl, C₁-C₄-alkylsulphonyl- and C₆-C₁₄-aryl-C₁-C₄-alkylsulphonyl-,

optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.

5 Preferred compounds of formula (I) are those wherein

R^1 to R^4 , R^6 and R^7 are as hereinbefore defined, and

L denotes a linker selected from among optionally substituted C_2 - C_{10} -alkyl, C_2 - C_{10} -alkenyl, C_6 - C_{14} -aryl, $-C_2$ - C_4 -alkyl- C_6 - C_{14} -aryl, $-C_6$ - C_{14} -aryl- C_1 - C_4 -alkyl, optionally bridged C_3 - C_{12} -cycloalkyl and heteroaryl which contains 1 or 2 nitrogen
10 atoms

n denotes 1

m denotes 1 or 2

R^5 denotes a group which is bound to L via a nitrogen atom, selected from among optionally substituted morpholinyl, piperidinyl, R^8 -piperazinyl, pyrrolidinyl, tropanyl,

15 R^8 -diketomethylpiperazinyl, sulphoxomorpholinyl, sulphonylmorpholinyl, thiomorpholinyl, $-NR^8R^9$ and azacycloheptyl,

R^8 , R^9 denote unsubstituted nitrogen substituents at R^5 , which may be identical or different, hydrogen or a group selected from among C_1 - C_6 -alkyl, $-C_1$ - C_4 -alkyl-

C_3 - C_{10} -cycloalkyl, C_3 - C_{10} -cycloalkyl, C_6 - C_{14} -aryl, $-C_1$ - C_4 -alkyl- C_6 - C_{14} -aryl, pyranyl, 20 pyridinyl, pyrimidinyl, C_1 - C_4 -alkyloxycarbonyl, C_6 - C_{14} -arylcarbonyl, C_1 - C_4 -alkylcarbonyl, C_6 - C_{14} -arylmethyloxycarbonyl, C_6 - C_{14} -arylsulphonyl, C_1 - C_4 -alkylsulphonyl and C_6 - C_{14} -aryl- C_1 - C_4 -alkylsulphonyl,

optionally in the form of the tautomers, the racemates, the enantiomers, the

25 diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.

Also preferred are compounds of formula (I), wherein

R^1 to R^4 , R^6 and R^7 are as hereinbefore defined,

30 L denotes a linker selected from among optionally substituted C_2 - C_{10} -alkyl, C_2 - C_{10} -alkenyl, C_6 - C_{14} -aryl, $-C_2$ - C_4 -alkyl- C_6 - C_{14} -aryl, $-C_6$ - C_{14} -aryl- C_1 - C_4 -alkyl,

optionally bridged C₃-C₁₂-cycloalkyl and heteroaryl which contains 1 or 2 nitrogen atoms

n denotes 0 or 1

m denotes 1 or 2

- 5 R⁵ denotes a group which is bound to L via a carbon atom, selected from among R⁸ - piperidinyl, R⁸R⁹-piperazinyl, R⁸-pyrrolidinyl, R⁸- piperazinylcarbonyl, R⁸-tropenyl, R⁸-morpholinyl and R⁸-azacycloheptyl, and

- 10 R⁸, R⁹ denote unsubstituted nitrogen substituents at R⁵, which may be identical or different, hydrogen or a group selected from among C₁-C₆-alkyl, -C₁-C₄-alkyl-C₃-C₁₀-cycloalkyl, C₃-C₁₀-cycloalkyl, C₆-C₁₄-aryl, -C₁-C₄-alkyl-C₆-C₁₄-aryl, pyranyl, pyridinyl, pyrimidinyl, C₁-C₄-alkyloxycarbonyl, C₆-C₁₄-arylcarbonyl, C₁-C₄-alkylcarbonyl, C₆-C₁₄-arylmethyloxycarbonyl, C₆-C₁₄-arylsulphonyl, C₁-C₄-alkylsulphonyl and C₆-C₁₄-aryl-C₁-C₄-alkylsulphonyl,

15

optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.

- 20 Particularly preferred are compounds of formula I wherein L, m, n and R³ to R⁹ are as hereinbefore defined, and R¹, R² which may be identical or different, denote a group selected from among hydrogen, Me, Et, Pr, or R¹ and R² together form a C₂-C₄-alkyl bridge ,

- 25 optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.

Especially preferred are compounds of formula I wherein

- 30 R¹, R², m, n and R⁵ to R⁸ are as hereinbefore defined, and R³ denotes a group selected from among optionally substituted C₁-C₁₀-alkyl, C₃-C₇-cycloalkyl, C₃-C₆-heterocycloalkyl and C₆-C₁₄-aryl or

R¹ and R³ or R² and R³ together denote a saturated or unsaturated C₃-C₄-alkyl bridge which may contain 1 to 2 heteroatoms,

R⁴ denotes a group selected from among hydrogen, OMe, OH, Me, Et, Pr, OEt, NHMe, NH₂, F, Cl, Br, O-propargyl, O-butynyl, CN, SMe, NMe₂, CONH₂, ethynyl, propynyl, butynyl and allyl, and

L denotes a linker selected from among optionally substituted phenyl, phenylmethyl, cyclohexyl and branched C₁-C₆-alkyl, optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.

The invention further relates to compounds of formula I for use as pharmaceutical compositions.


Of particular importance according to the invention are compounds of formula I for use as pharmaceutical compositions with an antiproliferative activity.

The invention also relates to the use of a compound of formula I for preparing a pharmaceutical composition for the treatment and/or prevention of cancer, infections, inflammatory and autoimmune diseases.

The invention also relates to a method of treating and/or preventing cancer, infections, inflammatory and autoimmune diseases, characterised in that a patient is given an effective amount of a compound of formula I.

The invention also relates to pharmaceutical preparations, containing as active substance one or more compounds of general formula (I) or the physiologically acceptable salts thereof, optionally combined with conventional excipients and/or carriers.

The invention also relates to a process for preparing a compound of general formula (I),




5

wherein

R^1 - R^5 , m , n and L are as hereinbefore defined,

characterised in that a compound of general formula (II)




10

wherein

R^1 - R^3 are as hereinbefore defined and A is a leaving group,


is reacted with an optionally substituted compound of general formula (III),



wherein

R^4 is as hereinbefore defined and

- 5 R^{10} denotes OH, $NH-L-R^5$, -O-methyl, -O-ethyl,
and optionally then the product of general formula (IV)



wherein

- 10 R^1 to R^4 is as hereinbefore defined and
 R^{10} denotes OH, $-NH-L-R^5$, -O-methyl or -O-ethyl,

optionally after previous hydrolysis of the ester group $-COR^{10}$, is reacted with an
amine of general formula (V)


15



wherein

R⁵ is as hereinbefore defined.

The invention further relates to a compound of formula (II),



5 wherein

R¹-R³ are as hereinbefore defined and A is a leaving group.

The term alkyl groups, including alkyl groups which are a part of other groups, denotes branched and unbranched alkyl groups with 1 to 12 carbon atoms,
10 preferably 1 – 6, most preferably 1-4 carbon atoms, such as, for example: methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl and dodecyl. Unless otherwise stated, the abovementioned terms propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl and dodecyl include all the possible isomeric forms. For example, the term propyl includes the two isomeric groups n-propyl and iso-propyl,
15 the term butyl includes n-butyl, iso-butyl, sec. butyl and tert.-butyl, the term pentyl includes iso-pentyl, neopentyl, etc.

In the abovementioned alkyl groups one or more hydrogen atoms may optionally be replaced by other groups. For example these alkyl groups may be substituted
20 by fluorine. All the hydrogen atoms of the alkyl group may optionally also be replaced.

The term alkyl bridge, unless otherwise stated, denotes branched and unbranched alkyl groups with 1 to 5 carbon atoms, for example methylene, ethylene,
25 propylene, isopropylene, n-butylene, iso-butyl, sec. butyl and tert.-butyl etc. bridges. Methylene, ethylene, propylene and butylene bridges are particularly preferred. In the alkyl bridges mentioned 1 to 2 C-atoms may optionally be

replaced by one or more heteroatoms selected from among oxygen, nitrogen or sulphur.

5 The term alkenyl groups (including those which are a part of other groups) denotes branched and unbranched alkylene groups with 2 to 10 carbon atoms, preferably 2 - 6 carbon atoms, most preferably 2 - 3 carbon atoms, provided that they have at least one double bond. Examples include: ethenyl, propenyl, butenyl, pentenyl etc. Unless otherwise stated, the abovementioned terms propenyl, butenyl, etc also include all the possible isomeric forms. For example, the term butenyl
10 includes 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-1-propenyl, 2-methyl-2-propenyl and 1-ethyl-1-ethenyl.

In the abovementioned alkenyl groups, unless otherwise stated, one or more hydrogen atoms may optionally be replaced by other groups. For example, these
15 alkyl groups may be substituted by the halogen atom fluorine. All the hydrogen atoms of the alkenyl group may optionally also be replaced.

The term alkynyl groups (including those which are a part of other groups) denotes branched and unbranched alkynyl groups with 2 to 10 carbon atoms, provided that
20 they have at least one triple bond, for example ethynyl, propargyl, butynyl, pentynyl, hexynyl etc., preferably ethynyl or propynyl.

In the abovementioned alkynyl groups, unless otherwise stated, one or more hydrogen atoms may optionally be replaced by other groups. For example, these
25 alkyl groups may be substituted by fluorine. All the hydrogen atoms of the alkynyl group may optionally also be replaced.

The term aryl denotes an aromatic ring system with 6 to 14 carbon atoms, preferably 6 or 10 carbon atoms, preferably phenyl, which, unless otherwise
30 stated, may carry one or more of the following substituents, for example: OH, NO₂, CN, OMe, -OCHF₂, -OCF₃, -NH₂, halogen, for example fluorine or chlorine, C₁-C₁₀-alkyl, preferably C₁-C₅-alkyl, preferably C₁-C₃-alkyl, most preferably methyl or

ethyl, -O-C₁-C₃-alkyl, preferably -O-methyl or -O-ethyl, -COOH, -COO-C₁-C₄-alkyl, preferably -O-methyl or -O-ethyl, -CONH₂.

5 Examples of heteroaryl groups wherein up to two carbon atoms are replaced by one or two nitrogen atoms include pyrrole, pyrazole, imidazole, triazole, pyridine, pyrimidine, while each of the abovementioned heteroaryl rings may optionally also be anellated to a benzene ring, preferably benzimidazole, and unless otherwise stated these heterocycles may carry one or more of the following substituents, for example: F, Cl, Br, OH, OMe, methyl, ethyl, CN, CONH₂, NH₂, optionally
10 substituted phenyl, optionally substituted heteroaryl, preferably optionally substituted pyridyl.

Examples of cycloalkyl groups are cycloalkyl groups with 3 - 12 carbon atoms, for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl,
15 preferably cyclopropyl, cyclopentyl or cyclohexyl, while each of the abovementioned cycloalkyl groups may optionally also carry one or more substituents, for example: OH, NO₂, CN, OMe, -OCHF₂, -OCF₃, -NH₂ or halogen, preferably fluorine or chlorine, C₁-C₁₀-alkyl, preferably C₁-C₅-alkyl, preferably C₁-C₃-alkyl, more preferably methyl or ethyl, -O-C₁-C₃-alkyl, preferably -O-methyl or -
20 O-ethyl, -COOH, -COO-C₁-C₄-alkyl, preferably -COO-methyl or -COO-ethyl or -CONH₂. Particularly preferred substituents of the cycloalkyl groups are =O, OH, NH₂, methyl or F.

Examples of cycloalkenyl groups are cycloalkyl groups with 3 - 12 carbon atoms
25 which have at least one double bond, for example cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl or cycloheptenyl, preferably cyclopropenyl, cyclopentenyl or cyclohexenyl, while each of the abovementioned cycloalkenyl groups may optionally also carry one or more substituents.

30 "=O" denotes an oxygen atom linked via a double bond.

Examples of heterocycloalkyl groups, unless otherwise described in the definitions, include 3- to 12- membered, preferably 5- , 6- or 7-membered, saturated or unsaturated heterocycles which may contain as heteroatoms nitrogen, oxygen or sulphur, for example tetrahydrofuran, tetrahydrofuranone, γ -butyrolactone, α -pyran, γ -pyran, dioxolane, tetrahydropyran, dioxane, 5 dihydrothiophene, thiolan, dithiolan, pyrroline, pyrrolidine, pyrazoline, pyrazolidine, imidazoline, imidazolidine, tetrazole, piperidine, pyridazine, pyrimidine, pyrazine, piperazine, triazine, tetrazine, morpholine, thiomorpholine, diazepan, oxazine, tetrahydro-oxazinyI, isothiazole, pyrazolidine, preferably morpholine, pyrrolidine, 10 piperidine or piperazine, while the heterocyclic group may optionally carry substituents, for example C1-C4-alkyl, preferably methyl, ethyl or propyl.

Examples of polycycloalkyl groups are optionally substituted, bi-, tri-, tetra- or pentacyclic cycloalkyl groups, for example pinane, 2,2,2-octane, 2,2,1-heptane or 15 adamantane. Examples of polycycloalkenyl groups are optionally bridged and/or substituted 8-membered bi-, tri-, tetra- or pentacyclic cycloalkenyl groups, preferably bicycloalkenyl or tricycloalkenyl groups, if they have at least one double bond, for example norbornene.

20 Examples of spiroalkyl groups are optionally substituted spirocyclic C₅-C₁₂ alkyl groups.

Generally, the term halogen denotes fluorine, chlorine, bromine or iodine, preferably fluorine, chlorine or bromine, most preferably chlorine.

25

The leaving group A denotes either identical or different leaving groups such as for example -o-methyl, -SCN, chlorine, bromine, iodine, methanesulphonyl, trifluoromethanesulphonyl or p-toluenesulphonyl, preferably chlorine.

30 The compounds according to the invention may be present in the form of the individual optical isomers, mixtures of the individual enantiomers, diastereomers or racemates, in the form of the tautomers and also in the form of the free bases or the corresponding acid addition salts with pharmacologically acceptable acids - such as for example acid addition salts with hydrohalic acids, for example

hydrochloric or hydrobromic acid, or organic acids, such as for example oxalic, fumaric, diglycolic or methanesulphonic acid.

5 The substituent R^1 may denote hydrogen or a group selected from among optionally substituted and/or branched C_1 - C_6 -alkyl, preferably methyl or ethyl, more preferably methyl or ethyl.

10 The substituent R^2 may denote hydrogen or a group selected from among optionally substituted and/or branched C_1 - C_6 -alkyl, preferably methyl or ethyl.

R^1 and R^2 together may denote a 2- to 5-membered alkyl bridge, preferably an ethylene, propylene or butylene bridge which may contain 1 to 2 heteroatoms, preferably oxygen or nitrogen, more preferably ethylene, propylene.

15 The substituent R^3 may denote hydrogen or a group selected from among optionally substituted and/or branched C_1 - C_{12} -alkyl, preferably ethyl, propyl, butyl, pentyl or hexyl, more preferably propyl, butyl, pentyl or hexyl, C_2 - C_{12} -alkenyl, preferably C_5 - C_7 -alkenyl, C_2 - C_{12} -alkynyl, preferably C_5 - C_7 -alkynyl and C_6 - C_{14} -aryl, preferably phenyl, a group selected from among optionally substituted and/or
20 bridged C_3 - C_{12} -cycloalkyl, preferably cyclopentyl or cyclohexyl, C_3 - C_{12} -cycloalkenyl, preferably C_5 - C_7 -cycloalkenyl, C_7 - C_{12} -polycycloalkyl, C_7 - C_{12} -polycycloalkenyl, C_5 - C_{12} -spirocycloalkyl, C_3 - C_{12} -heterocycloalkyl, preferably pyranyl or piperinyl, pyrrolidinyl, pyrazinyl or morpholinyl which contains 1 to 2 heteroatoms, preferably oxygen or nitrogen, and C_3 - C_{12} -heterocycloalkenyl which
25 contains 1 to 2 heteroatoms, preferably oxygen or nitrogen.

Most preferably, the substituent R^3 denotes isopropyl, isobutyl, isopentyl, cyclopentyl, phenyl or cyclohexyl.

30 R^1 and R^3 or R^2 and R^3 together may denote a saturated or unsaturated C_3 - C_4 -alkyl bridge which may contain 1 heteroatom, preferably oxygen or nitrogen.

The substituent R^4 may denote a group selected from among hydrogen, -CN, hydroxy, $-NR^6R^7$ and Halogen, preferably chlorine or fluorine, more preferably chlorine or a group selected from among optionally substituted C_1 - C_6 -alkyl, preferably methyl, ethyl or propyl, C_2 - C_6 -alkenyl, preferably ethenyl or propenyl, C_2 - C_6 -alkynyl, preferably ethynyl, propynyl or butynyl, C_1 - C_5 -alkyloxy, preferably methoxy, ethoxy or propargyloxy, C_2 - C_5 -alkenyloxy, C_2 - C_5 -alkynyloxy, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulphoxo and C_1 - C_6 -alkylsulphonyl.

Most preferably, the substituent R^4 denotes methoxy, methyl, ethoxy, ethyl, propargyloxy or chlorine.

L may denote a linker selected from among optionally substituted C_2 - C_{10} -alkyl, preferably ethyl, propyl, butyl or pentyl, C_2 - C_{10} -alkenyl, C_6 - C_{14} -aryl, preferably phenyl, $-C_2$ - C_4 -alkyl- C_6 - C_{14} -aryl, $-C_6$ - C_{14} -aryl- C_1 - C_4 -alkyl, preferably -phenyl-methyl, optionally bridged C_3 - C_{12} -cycloalkyl, preferably cyclohexyl, and heteroaryl which contains 1 or 2 nitrogen atoms.

n denotes 0 or 1

m denotes 1 or 2, preferably 1.

R^5 may denote a group selected from among optionally substituted morpholinyl, piperidinyl, piperazinyl, piperazinylcarbonyl, pyrrolidinyl, tropenyl, R^8 -

diketomethylpiperazinyl, sulphoxomorpholinyl, sulphonylmorpholinyl, thiomorpholinyl, $-NR^8R^9$ and azacycloheptyl, preferably piperidinyl, morpholinyl, pyrrolidinyl, sulphoxomorpholinyl, piperazinyl, thiomorpholinyl or tropenyl.

The groups R^6 and R^7 may be identical or different and may denote hydrogen or C_1 - C_4 -alkyl, preferably methyl or ethyl.

The groups R^8 and R^9 may be unsubstituted nitrogen substituents at R^5 , they may be identical or different and denote either hydrogen or a group selected from among C_1 - C_6 -alkyl, preferably methyl, ethyl or propyl, $-C_1$ - C_4 -alkyl- C_3 - C_{10} -cycloalkyl, preferably $-CH_2$ -cyclopropyl, C_3 - C_{10} -cycloalkyl, C_6 - C_{14} -aryl, preferably phenyl, $-C_1$ - C_4 -alkyl- C_6 - C_{14} -aryl, preferably benzyl, pyranyl, pyridinyl, pyrimidinyl, pyranyl, C_1 - C_4 -alkyloxycarbonyl, C_6 - C_{14} -arylcarbonyl, C_1 - C_4 -alkylcarbonyl, C_6 - C_{14} -

arylmethyloxycarbonyl, C₆-C₁₄-arylsulphonyl, C₁-C₄-alkylsulphonyl and C₆-C₁₄-aryl-C₁-C₄-alkylsulphonyl.

Most preferably, the substituent R⁸ denotes methyl, ethyl or propyl.

5 Most preferably, the substituent R⁹ denotes methyl, ethyl or propyl.

R¹⁰ may be a substituent selected from among OH, NH₂-LR₅, -O-methyl and -O-ethyl, preferably OH, LR₅, -O-methyl or -O-ethyl.

10 All the groups mentioned in the definition of R¹ to R¹⁰ may optionally be branched and/or substituted.


The compounds according to the invention may be prepared by synthesis methods A described hereinafter, while the substituents of general formulae (A1) to (A9) have the meanings given hereinbefore. This method is to be understood as an
15 illustration of the invention without restricting it to the subject matter thereof.

Method A

Step 1A

A compound of formula (A1) is reacted with a compound of formula (A2) to obtain
20 a compound of formula (A3) (Diagram 1A). This reaction may be carried out according to WO 0043369 or WO 0043372. Compound (A1) is commercially obtainable, for example, from City Chemical LLC, 139 Allings Crossing Road, West Haven, CT, 06516, USA. Compound (A2) may be prepared by procedures known from the literature: (a) F. Effenberger, U. Burkhart, J. Willfahrt *Liebigs Ann. Chem.* **1986**, 314-333; b) T. Fukuyama, C.-K. Jow, M. Cheung, *Tetrahedron Lett.* **1995**, 36, 6373-6374; c) R. K. Olsen, *J. Org. Chem.* **1970**, 35, 1912-1915; d) F.E. Dutton, B.H. Byung *Tetrahedron Lett.* **1998**, 30, 5313-5316; e) J. M. Ranajuhi, M. M. Joullie *Synth. Commun.* **1996**, 26, 1379-1384.).
25

Diagram 1A



In Step 1A, 1 equivalent of the compound (A1) and 1 to 1.5 equivalents, preferably 1.1 equivalents of a base, preferably potassium carbonate, potassium hydrogen carbonate, sodium carbonate or sodium hydrogen carbonate, calcium carbonate, most preferably potassium carbonate, are stirred in a diluent optionally mixed with water, for example acetone, tetrahydrofuran, diethylether, cyclohexane, petroleum ether or dioxane, preferably cyclohexane or diethylether.

At a temperature of 0 to 15 °C, preferably 5 to 10 °C, 1 equivalent of an amino acid of formula (A2), dissolved in an organic solvent, for example acetone, tetrahydrofuran, diethylether, cyclohexane or dioxane, is added dropwise. The reaction mixture is heated to a temperature of 18°C to 30 °C, preferably about 22°C, with stirring and then stirred for a further 10 to 24 hours, preferably about 12 hours. Then the diluent is distilled off, the residue is combined with water and the mixture is extracted two to three times with an organic solvent, such as diethylether or ethyl acetate, preferably ethyl acetate. The combined organic extracts are dried and the solvent is distilled off. The residue (compound A3) may be used in Step 2 without any prior purification.


Step 2A

The compound obtained in Step 1A (A3) is reduced at the nitro group and cyclised to form the compound of formula (A4) (Diagram 2A).

5

Diagram 2A

Schema 2A



10 [Reduktion = reduction]


In Step 2A, 1 equivalent of the nitro compound (A3) is dissolved in an acid, preferably glacial acetic acid, formic acid or hydrochloric acid, preferably glacial acetic acid, and heated to 50 to 70°C, preferably about 60°C. Then a reducing agent, for example zinc, tin or iron, preferably iron filings, is added to complete the exothermic reaction and the mixture is stirred for 0.2 to 2 hours, preferably 0.5 hours, at 100 to 125 °C, preferably at about 117 °C. After cooling to ambient temperature the iron salt is filtered off and the solvent is distilled off. The residue is taken up in a solvent or mixture of solvents, for example ethyl acetate or dichloromethane/ methanol 9/1 and semisaturated NaCl solution, and filtered through kieselgur, for example. The organic phase is dried and evaporated down. The residue (compound (A4)) may be purified by chromatography or by crystallisation or used as the crude product in Step 3A of the synthesis.

25

Step 3A

The compound obtained in Step 2A (A4) may be reacted by electrophilic substitution as shown in Diagram 3A to obtain the compound of formula (A5).

5 Diagram 3A




- In Step 3A 1 equivalent of the amide of formula (A4) is dissolved in an organic solvent, for example dimethylformamide or dimethylacetamide, preferably dimethylacetamide, and cooled to about -5 to 5 °C, preferably 0°C.
- 10 Then 0.9 to 1.3 equivalents of sodium hydride and 0.9 to 1.3 equivalents of a methylating reagent, e.g. methyl iodide, are added. The reaction mixture is stirred for 0.1 – 3 hours, preferably about 1 hour, at about 0 to 10 °C, preferably at about 5 °C, and may optionally be left to stand for a further 12 hours at this temperature.
- 15 The reaction mixture is poured onto ice water and the precipitate is isolated. The residue (compound (A5)) may be purified by chromatography, preferably over silica gel, or by crystallisation, or used as the crude product in step 4A of the synthesis.

Step 4A

The amination of the compound (A5) obtained in Step 3A to yield the compound of formula (A9) (Diagram 4A) may be carried out using the methods known from the literature of variants 4.1 A (a) M.P.V. Boarland, J.F.W. McOmie *J. Chem. Soc.*

- 5 **1951**, 1218-1221; b) F. H. S. Curd, F. C. Rose *J. Chem. Soc.* **1946**, 343-348., 4.2 A (a) Banks *J. Am. Chem. Soc.* **1944**, 66, 1131 b) Ghosh and Dolly *J. Indian Chem. Soc.* **1981**, 58, 512-513; c) N. P. Reddy and M. Tanaka *Tetrahedron Lett.* **1997**, 38, 4807-4810.

10 Diagram 4A



For example, in variant 4.1 A, 1 equivalent of the compound (A5) and 1 to 3 equivalents, preferably about 2 equivalents of the compound (A6) are heated without a solvent or in an organic solvent such as for example sulpholane, dimethylformamide, dimethylacetamide, toluene, N-methylpyrrolidone, dimethylsulphoxide or dioxane, preferably sulpholane, for 0.1 to 4 hours, preferably 1 hour, at 100 to 220 °C, preferably at about 160 °C. After cooling, the product (A9) is crystallised by the addition of organic solvents or mixtures of solvents, e.g. diethylether/methanol, ethyl acetate, methylene chloride, or diethylether, preferably diethylether/methanol 9/1, or purified by chromatography.

10

For example, in variant 4.2 A, 1 equivalent of the compound (A5) and 1 to 3 equivalents of the compound (A6) are stirred with acid, for example 1-10 equivalents of 10-38% hydrochloric acid and/or an alcohol, for example ethanol, propanol, butanol, preferably ethanol, at reflux temperature for 1 to 48 hours, preferably about 5 hours.

15

The product precipitated (A9) is filtered off and optionally washed with water, dried and crystallised from a suitable organic solvent.

For example, in variant 4.3 A, 1 equivalent of the compound (A5) and 1 to 3 equivalents of the compound (A7) are dissolved in a solvent, for example toluene or dioxane and combined with a phosphine ligand, for example 2,2'-bis-(diphenylphosphino)-1,1'-binaphthyl and a palladium catalyst, for example tris(dibenzylidene-acetone)-dipalladium(0) and a base, for example caesium carbonate, and refluxed for 1-24 h, preferably 17 h. The reaction mixture is purified for example over silica gel and the product (A8) is isolated from the solution or obtained by suitable crystallisation.

20

25


The product (A8) is dissolved in a suitable solvent, for example dioxane and mixed with acid, for example semiconcentrated hydrochloric acid, for example in the ratio of solvent to acid of 3: 1. Then the mixture is refluxed for 1 - 48 h, for example 12 h, and the precipitate formed is isolated. If desired the product (A9) is purified by crystallisation.

30

Step 5A

Diagram 5A

5



Variant 5.1 A:


- 10 For example, 1 equivalent of the compound (A9) is dissolved with 1 equivalent of an activating reagent, e.g. O-benzotriazolyl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU) and a base, for example 1.5 equivalents of diisopropylethylamine (DIPEA) in an organic diluent, for example dichloromethane, tetrahydrofuran, dimethylformamide, N-methylpyrrolidone, dimethylacetamide,
- 15 preferably dichloromethane or dimethylformamide. After the addition of 1 equivalent of the amine (A10) the reaction mixture is stirred for 0.1 to 24 hours, preferably about 2 hours at 20°C to 100°C. The product of formula (A11) is obtained for example by crystallisation or chromatographic purification.
- 20 The new compounds of general formula (I) may be synthesised analogously to the following examples of synthesis. These Examples are, however, intended only as examples of procedures to illustrate the invention further, without restricting the invention to their subject matter.

The preparation of some intermediate compounds used to synthesise the examples is also described hereinafter.

Preparation of the acids

5

To synthesise the compounds of Examples 94 and 95 first an intermediate



compound Z1

is prepared as described hereinafter.

10

50.0 g (0.48 mol) of D-alanine methyl ester x HCl and 49.1 g (0.50 mol) cyclohexanone were placed in 300 mL dichloromethane and then combined with 41.0 g (0.50 mol) sodium acetate and 159.0 g (0.75 mol) sodium triacetoxyborohydride. the mixture was stirred overnight and then 300 mL of 10% sodium hydrogen carbonate solution were added. The aqueous phase was extracted with dichloromethane. The combined organic phases were washed with 10% sodium hydrogen carbonate solution, dried over Na₂SO₄ and evaporated down.

15

Yield: 72.5 g of a compound Z1a (clear liquid)

20

72.5 g of the compound Z1a were placed in 500 mL water and 76.6 g (0.39 mol) of 2,4-dichloro-5-nitropyrimidine in 500 mL diethyl ether were added. At a temperature of -5°C 100 mL 10% potassium hydrogen carbonate solution were added dropwise. The mixture was stirred for 3 h at -5°C and for a further 12 h at

ambient temperature. The organic phase was separated off and dried over Na_2SO_4 . On evaporation the product crystallised out.

Yield: 48.0 g of a compound Z1b (yellow crystals)

- 5 48.0 g of the compound Z1b were dissolved in 350 mL glacial acetic acid and heated to 60°C. 47.5 g of iron powder were added, while the temperature rose to 105°C. The reaction mixture was stirred for three hours at 80°C, then filtered hot through cellulose and evaporated down. The residue was stirred in water and ethyl acetate, suction filtered and the light-grey precipitate was washed with ethyl
- 10 acetate. The filtrate was washed with dilute ammonia and water, the organic phase was dried over Na_2SO_4 , filtered through activated charcoal and evaporated down. Some more light-grey solid was obtained.

Yield: 29.5 g of a compound Z1c (light-grey crystals)

- 15 32.1 g of the compound Z1c were placed in 300 mL dimethylacetamide and combined with 13 mL (0.2 mol) methyl iodide. At -5°C 6.4 g (0.16 mol) sodium hydride as a 60% dispersion in mineral oil was added batchwise. After 2 h the reaction mixture was poured onto 800 mL ice water. The precipitate formed was suction filtered and washed with petroleum ether.


- 20 Yield: 33.0 g of a compound Z1d (beige crystals)

4.0 g of the compound Z1d and 2.3 g (15 mmol) 4-amino-3-methylbenzoic acid were suspended in 50 mL ethanol and 120 mL water, combined with 2 mL conc. hydrochloric acid and refluxed for 48 h. The precipitate formed on cooling was

25 suction filtered and washed with water, ethanol and diethyl ether.

Yield: 2.9 g of a compound Z1 (colourless crystals)

To synthesise the compounds Example 188 and Example 203 first of all an intermediate compound Z2



5 is prepared as described below.

A solution of 128.2 g (0.83 mol) D-alanine ethyl ester x HCl and 71.5 g (0.85 mol) cyclopentanone in 1500 mL dichloromethane was combined with 70.1 (0.85 mol) sodium acetate and 265.6 g (1.25 mol) sodium triacetoxyborohydride. The reaction
10 mixture was stirred for 12 h and then poured into 1.5 L of a 10% sodium hydrogen carbonate solution. The aqueous phase was extracted with dichloromethane. The combined organic phases were dried over Na₂SO₄ and evaporated down.
Yield: 143.4 g of a compound Z2a (colourless oil)

15 66.0 g of the compound Z2a were placed in 500 mL water and combined with 85.0 g (0.44 mol) 2,4-dichloro-5-nitropyrimidine in 500 mL diethyl ether. At -5°C 100 mL 10% potassium hydrogen carbonate solution were added dropwise and the reaction mixture was stirred for 48 h at ambient temperature. The aqueous phase was extracted with diethyl ether, the combined organic phases were dried over
20 Na₂SO₄ and evaporated down. The dark red solid was stirred with petroleum ether and suction filtered.

Yield: 88.0 g of a compound Z2b (yellow crystals)

88.0 g of the compound Z2b were dissolved in 1000 mL glacial acetic acid and at 60°C combined batchwise with 85 g iron powder, while the temperature rose to 110°C. It was stirred for 1 h at 60°C, then suction filtered hot through cellulose and evaporated down. The brown solid was stirred with 700 mL water and suction
5 filtered.

Yield: 53.3 g of a compound Z2c (light brown crystals)

53.3 g of the compound Z2c were dissolved in 300 mL dimethylacetamide and combined with 13 mL (0.21 mol) methyl iodide. At -5°C 5.0 g (0.21 mol) sodium
10 hydride as a 60% dispersion in mineral oil were added batchwise. After 12 h the reaction mixture was poured onto 1000 mL ice water and the precipitate formed was suction filtered.


Yield: 40.0 g of a compound Z2d (colourless crystals)

15 4.0 g of the compound Z2d and 2.8 g (16 mmol) 4-amino-3-chlorbenzoic acid were suspended in 25 mL ethanol and 60 mL water, combined with 3 mL conc. hydrochloric acid and refluxed for 43 h. The precipitate formed on cooling was suction filtered and washed with water, ethanol and diethyl ether.

Yield: 0.9 g of a compound Z2 (colourless crystals)

20

To synthesise the compounds of Examples 19, 21, 22, 23, 45, 55, 58, 116, 128, 131, 133, 134, 136, 138, 177, 217, 231, 239, 46, 184, 166 and 187 first of all an intermediate compound Z3



is prepared as described below.

54.0 g (0.52 mol) D-2-aminobutyric acid were suspended in 540 mL methanol and slowly combined with 132 g (1.1 mol) thionyl chloride while cooling with ice. The mixture was refluxed for 1.5 h and then evaporated down. The oil remaining was combined with 540 mL tert-butylmethylether and the colourless crystals formed were suction filtered.

Yield: 78.8 g of a compound Z3a (colourless crystals)

74.2 g of the compound Z3a and 43.5 mL (0.49 mol) cyclopentanone were dissolved in 800 mL dichloromethane. After the addition of 40.0 g (0.49 mol) sodium acetate and 150.0 g (0.71 mol) sodium triacetoxyborohydride at 0°C the mixture was stirred for 12 h at ambient temperature and then 500 mL of 20% sodium hydrogen carbonate solution were added. The aqueous phase was extracted with dichloromethane. The combined organic phases were washed with water, dried over MgSO₄ and evaporated down.

Yield: 85.8 g of a compound Z3b (light yellow oil)

40.0 g of the compound Z3b and 30.0 g (0.22 mol) potassium carbonate were suspended in 600 mL acetone and combined with 45.0 g (0.23 mol) 2,4-dichloro-5-nitropyrimidin in 200 mL acetone while cooling with ice. After 12 h a further 5.0 g 2,4-dichloro-5-nitropyrimidin were added and stirred for 3 h. The reaction mixture was evaporated down, taken up in 800 mL ethyl acetate and 600 mL water and the aqueous phase was extracted with ethyl acetate. The combined organic phases were washed with water, dried over MgSO₄ and evaporated down.

Yield: 75.0 g of a compound Z3c (brown oil)

100 g of the compound Z3c were dissolved in 650 mL glacial acetic acid and at 70°C 20 g of iron powder were added batchwise. The mixture was stirred for 1 h at 70°C, then for 1.5 h at 100°C and then filtered hot through kieselgur. The reaction mixture was evaporated down, taken up in methanol/dichloromethane, applied to

silica gel and purified with ethyl acetate by Soxhlet extraction. The solvent was removed and the residue stirred with methanol.

Yield: 30.0 g of a compound Z3d (light brown crystals)

- 5 25.0 g of the compound Z3d and 6.5 mL (0.1 mol) methyl iodide were placed in 250 mL dimethylacetamide and at -10°C 3.8 g (0.95 mol) sodium hydride as a 60% dispersion in mineral oil was added. It was stirred for 20 min at 0°C , then for 30 min at ambient temperature and finally ice was added. The reaction mixture was evaporated down and combined with 300 mL water. The precipitate formed
- 10 was suction filtered and washed with petroleum ether.


Yield: 23.0 g of a compound Z3e (colourless solid)

6.0 g of the compound Z3e and 5.1 g (31 mmol) 4-amino-3-methoxybenzoic acid were suspended in 90 mL ethanol and 350 mL water, combined with 3.5 mL conc.

- 15 hydrochloric acid and refluxed for 48 h. The reaction mixture was evaporated down, the residue stirred with methanol/diethyl ether and the precipitate formed was suction filtered.

Yield: 6.3 g of a compound Z3 (light beige crystals)

- 20 To synthesise the compound of Examples 81, 82, 93, 137 first of all an intermediate compound Z4



Z4

is prepared as described below.

25.0 g (0.19 mol) of ethyl 1-aminocyclopropane-1-carboxylate x HCl and 16.8 g (0.20 mol) of cyclopentanone were dissolved in 300 mL of dichloromethane and combined with 16.4 g (0.20 mol) of sodium acetate and 61.7 g (0.29 mol) of sodium triacetoxyborohydride. It was stirred overnight and the reaction mixture was then poured onto 400 mL of 10% sodium hydrogen carbonate solution. The aqueous phase was extracted with dichloromethane. The combined organic phases were dried over Na₂SO₄ and evaporated down.

Yield: 34.5 g of a compound Z4a (colourless oil)

42.5 g (0.22 mol) of 2,4-dichloro-5-nitropyrimidine in 350 mL of diethyl ether were added to a mixture of 34.5 g of the compound Z4a in 350 mL water. At -5°C the mixture was combined with 80 mL 10% potassium hydrogen carbonate solution and stirred overnight at ambient temperature. The aqueous phase was extracted with diethyl ether. The combined organic phases were dried over Na₂SO₄ and evaporated down.

Yield: 53.8 g of a compound Z4b (brown oil)

20.1 g of the compound Z4b were dissolved in 200 mL glacial acetic acid and combined batchwise at 60°C with 19.1 g iron powder, during which time the temperature rose to 100°C. The mixture was stirred for 3 h at 60°C, then suction filtered through cellulose and evaporated down. The residue was stirred in water and ethyl acetate and the yellow precipitate was suction filtered. The filtrate was washed with dilute ammonia and water, the organic phase dried over Na₂SO₄ and evaporated down. After the addition of diethyl ether additional product crystallised out.

Yield: 4.0 g of a compound Z4c (yellow crystals)


7.8 g of the compound Z4c and 2.6 mL (0.04 mol) methyl iodide were dissolved in 100 mL dimethylacetamide and at -5°C 1.5 g (0.04 mol) sodium hydride were

added batchwise as a 60% dispersion in mineral oil. After 2 h the reaction mixture was poured onto ice water and the precipitate formed was suction filtered.

Yield: 7.5 g of a compound Z4d (light brown crystals)

- 5 3.0 g of the compound Z4d and 1.9 g (11 mmol) 4-amino-3-methoxybenzoic acid were suspended in 40 mL ethanol and 80 mL water, combined with 2 mL conc. hydrochloric acid and refluxed for 20 h. A further 0.5 g of 4-amino-3-methoxybenzoic acid were added and refluxed for 48 h. The precipitate formed on cooling was suction filtered and washed with water, ethanol and diethyl ether.
- 10 Yield: 2.1 g of a compound Z4 (colourless crystals) m.p.: 222-223°C

To synthesise the compounds of Examples 162, 43, 53, 161, 202, 211, 215 and 212 first of all an intermediate compound Z5



Z5

15

is prepared as described below.

- A mixture of 73.4 mL (0.5 mol) ethyl 2-bromoisobutyrate, 87.1 mL (0.75 mol) of
- 20 3-methyl-1-butylamine, 82.5 g (0.6 mol) sodium iodide and 76.0 g (0.6 mol) of potassium carbonate in 1000 mL ethyl acetate was refluxed for 3 days. Any salts present were filtered off and the filtrate evaporated down.

Yield: 97.0 g of a compound Z5a (red oil)

49.0 g (0.25 mol) of 2,4-dichloro-5-nitropyrimidine and 38.3 g (0.28 mol) of potassium carbonate were suspended in 500 mL acetone and at 0°C combined with 93.0 g of the compound Z5a in 375 mL acetone. The reaction mixture was stirred overnight at ambient temperature, filtered and evaporated down. The
5 residue dissolved in ethyl acetate was washed with water and the organic phase dried over MgSO₄ and evaporated down.

Yield: 102.7 g of a compound Z5b (brown oil)

22.7 g of the compound Z5b were dissolved in 350 mL glacial acetic acid and at
10 60°C combined batchwise with 17.4 g iron powder. After the addition had ended the mixture was refluxed for 0.5 h, filtered hot and evaporated down. The residue was taken up in 200 mL dichloromethane/methanol (9:1) and washed with sodium chloride solution. The organic phase was suction filtered through kieselguhr, dried
15 over MgSO₄, evaporated down and purified by column chromatography (eluant: ethyl acetate/cyclohexane 1:1).

Yield: 1.9 g of a compound Z5c (colourless crystals)

1.9 g of the compound Z5c were dissolved in 32 mL dimethylacetamide and while cooling with ice combined with 0.3 g (7 mmol) sodium hydride as a 60% dispersion
20 in mineral oil. After 10 min 0.5 mL (7 mmol) methyl iodide were added and stirred for 3 h at ambient temperature. The reaction mixture was evaporated down and combined with water. The precipitate formed was suction filtered and washed with petroleum ether.

Yield: 1.6 g of a compound Z5d (colourless crystals)


25

14.0 g of the compound Z5d and 10.0 g (0.06 mol) 4-amino-3-methoxybenzoic acid were suspended in 200 mL dioxane and 80 mL water, combined with 10 mL conc. hydrochloric acid and refluxed for 40 h. The precipitate formed on cooling was suction filtered and washed with water, dioxane and diethyl ether.

30

Yield: 13.9 g of a compound Z5 (colourless crystals)

To synthesise the compounds of Examples 88, 194, 229 and 89 first of all an intermediate compound Z6



5

Z6

is prepared as described below.

6.0 g (0.06 mol) L-2-aminobutyric acid was placed in 80 mL 0.5 M sulphuric acid
10 and at 0°C combined with 5.5 g (0.08 mol) sodium nitrite in 15 mL water. The
reaction mixture was stirred for 22 h at 0°C, combined with ammonium sulphate
and filtered. The filtrate was extracted with diethyl ether and the combined organic
dried over MgSO₄ and evaporated down.

Yield: 6.0 g of a compound Z6a (yellow oil)

15

200 mL methanol were combined successively with 65.0 mL (0.89 mol) thionyl
chloride and 76.0 g of the compound Z6a in 50 mL methanol while cooling with
ice. The resulting mixture was stirred for 1 h at 0°C and 2 h at ambient
temperature and then the methanol and remaining thionyl chloride were eliminated
20 in vacuo at 0°C.

Yield: 40.0 g of a compound Z6b (yellow oil)

30.0 mL (0.17 mol) of trifluoromethanesulphonic acid anhydride were placed in 150 mL dichloromethane and while cooling with ice a solution of 20.0 g of the compound Z6b and 14.0 mL (0.17 mol) pyridine in 50 mL dichloromethane was added within one hour. The mixture was stirred for 2 h at ambient temperature,
5 any salts formed were suction filtered and then washed with 100 mL water. The organic phase was dried over MgSO_4 and evaporated down.

Yield: 42.0 g of a compound Z6c (light yellow oil)

42.0 g of the compound Z6c in 200 mL dichloromethane was added dropwise
10 within one hour to a solution of 15.5 mL (0.17 mol) of aniline and 24.0 mL (0.17 mol) of triethylamine in 400 mL dichloromethane while cooling with ice. The mixture was stirred for 1 h at ambient temperature and a further 2 h at 35°C. The reaction mixture was washed with water, dried over MgSO_4 and evaporated down. The residue remaining was purified by distillation (95-100°C, $1 \cdot 10^{-3}$ mbar).

15 Yield: 14.0 of a compound Z6d (colourless oil)

14.0 g of the compound Z6d and 16.0 g (0.1 mol) potassium carbonate were suspended in 100 mL acetone and at 10°C combined with 16.0 g (0.08 mol) of 2,4-dichloro-5-nitropyrimidine. The mixture was stirred for 4 h at 40°C, any salts
20 formed were suction filtered and the filtrate evaporated down. The residue was taken up in 300 mL ethyl acetate and washed with water. The organic phase was dried over MgSO_4 and evaporated down.

Yield: 31.0 g of a compound Z6e (brown oil)

25 31.0 g of the compound Z6e were dissolved in 200 mL glacial acetic acid and at 60°C combined batchwise with 10 g iron powder, during which time the temperature rose to 85°C. The mixture was stirred for a further hour at 60°C, filtered through kieselguhr and evaporated down. The residue was stirred with methanol.

30 Yield: 4.5 g of a compound Z6f (brown crystals)

At -20°C 0.6 g (16 mmol) of sodium hydride as a 60% dispersion in mineral oil were added batchwise to a mixture of 4.5 g of the compound Z6f and 1.0 mL (16 mmol) methyl iodide in 100 mL dimethylacetamide. After 1 h the reaction mixture was combined with 50 mL water and evaporated down. The residue was stirred
5 with 200 mL water, the precipitate is suction filtered and washed with petroleum ether.

Yield: 4.5 g of a compound Z6g (colourless crystals)


A suspension of 1.5 g of the compound Z6g and 1.4 g (8 mmol) of methyl 4-amino-
10 3-methoxybenzoate in 30 mL toluene was combined with 0.4 g (0.6 mmol) of 2,2'-bis-(diphenylphosphino)-1,1'-binaphthyl, 0.23 g (0.3 mmol) of tris(dibenzylideneacetone)-dipalladium(0) and 7.0 g (21 mmol) of caesium carbonate and refluxed for 17 h. The reaction mixture was applied to silica gel and purified by column chromatography (eluant: dichloromethane/methanol 9:1).

15 Yield: 1.7 g of a compound Z6h (yellow crystals)

1.7 g of the compound Z6h were dissolved in 50 mL dioxane, combined with 15 mL of semiconc. hydrochloric acid and refluxed for 12 h. After cooling the precipitate formed was suction filtered.

20 Yield: 1.1 g of a compound Z6 (colourless solid)

To synthesise the compound of Examples 26, 20, 32, 56, 101, 112, 209 first of all an intermediate compound Z7



5

is prepared as described below.

50.0 g (0.36 mol) D-alanine methyl ester x HCl was suspended in 500 mL of dichloromethane and 35 mL of acetone and combined with 30.0 g (0.37 mol) of sodium acetate and 80.0 g (0.38 mol) of sodium triacetoxyborohydride. The mixture was stirred for 12 h and then poured onto 400 mL of 10% sodium hydrogen carbonate solution. The organic phase was dried over Na₂SO₄ and evaporated down.

Yield: 51.0 g of a compound Z7a (yellow oil)

15

A suspension of 51.0 g of the compound Z7a in 450 mL water was combined with 80.0 g (0.41 mol) of 2,4-dichloro-5-nitropyridine in 450 mL of diethyl ether. At -5°C 100 mL of 10% potassium hydrogen carbonate solution were added dropwise. The reaction mixture was stirred for 3 h, the organic phase dried over Na₂SO₄ and evaporated down.

20

Yield: 74 g of a compound Z7b (yellow oil)

18.6 g of the compound Z7b were dissolved in 200 mL glacial acetic acid and at 60°C combined batchwise with 20.0 g iron powder. The mixture was stirred for 2 h at 60°C and then suction filtered through cellulose. The residue was dissolved in ethyl acetate and washed with water and conc. ammonia. The organic phase was
5 dried over Na₂SO₄ and evaporated down. The residue was crystallised from diethyl ether.

Yield: 9.8 g of a compound Z7c (colourless crystals)

17.0 g of the compound Z7c and 7 mL (0.1 mol) methyl iodide were dissolved in
10 200 mL dimethylacetamide and at -5°C combined with 4.0 g (0.1 mol) of sodium hydride as a 60% dispersion in mineral oil. The reaction mixture was stirred for 30 min and then poured onto 300 mL ice water. The precipitate formed was suction filtered and stirred with petroleum ether.

Yield: 14.8 g of a compound Z7d (beige crystals)



15

0.9 g of the compound Z7d and 1.5 g (9 mmol) 4-amino-3-methoxybenzoic acid were heated to 210°C for 30 min. After cooling the residue was stirred with ethyl acetate and the precipitate obtained was suction filtered.



Yield: 1.2 g of a compound Z7 (grey crystals)

20

The following acids were prepared, for example, analogously to the methods of synthesis described:





5



Synthesis of the amino components L-R5

10 The following amines were obtained as follows:

1,1-dimethyl-2-dimethylamino-1-yl-ethylamine and 1,1-dimethyl-2-piperidin-1-yl-ethylamine



15


The compounds were prepared according to the following references: a) S. Schuetz et al. *Arzneimittel-Forschung* **1971**, 21, 739-763 b) V. M. Belikov et al. *Tetrahedron* **1970**, 26, 1199-1216. c) E.B. Butler and McMillan *J. Amer. Chem. Soc.* **1950**, 72, 2978.

5

Other amines were prepared as follows, in a modified manner compared with the literature described above.

1,1-dimethyl-2-morpholin-1-yl-ethylamine

10



8.7 mL morpholine and 9.3 mL 2-nitropropane were prepared, while cooling with ice, 7.5 mL formaldehyde (37%) and 4 mL of a 0.5 mol/L NaOH solution were slowly added dropwise ($<10^{\circ}\text{C}$). Then the mixture was stirred for 1h at 25°C and 1h at 50°C . The solution was treated with water and ether and the aqueous phase was extracted 3x with ether. The combined org. phase was dried over NaSO_4 and combined with HCl in dioxane (4mol/l), the precipitate formed was suction filtered. Yield: 21.7 of white powder

15


5 g of the white powder were dissolved in 80 mL methanol and with the addition of 2 g RaNi treated with hydrogen at 35°C and 50 psi for 40 minutes. This yielded 3.6 g of 1,1-dimethyl-2-morpholin-1-yl-ethylamine.

20


The following amines were prepared by this method:

1,1- dimethyl-N-methylpiperazin-1-yl-ethylamine


25



1,1-dimethyl-2-pyrrolidin-1-yl-ethylamine




5 Synthesis of 1,3 -dimorpholin-2-amino-propane




- 5 g of 1,3 Dimorpholine-2-nitropropane obtained from Messrs. Aldrich was dissolved in 80 mL methanol and treated with hydrogen for 5.5 h at 30°C and 50
10 psi with the addition of 2 g RaNi. This yielded 4.2 g of 1,3 dimorpholin-2-amino-propane.

4-Aminobenzylmorpholine



- 15 The preparation of this amine is described in the following reference:
S. Mitsuru et al. *J. Med. Chem.* **2000**, 43, 2049-2063.

4-amino-1-tetrahydro-4H-pyran-4-yl-piperidine



20 g (100 mmol) of 4-tert-butyloxycarbonyl-aminopiperidine were dissolved in 250 mL CH₂Cl₂ and stirred for 12 h at RT with 10 g (100 mmol) tetrahydro-4H-pyran-4-one and 42 g (200 mmol) NaBH(OAc)₃. Then water and potassium carbonate were added, the org. phase was separated off, dried and the solvent was eliminated in vacuo. The residue was dissolved in 200 mL CH₂Cl₂ and stirred for 12 h at RT with 100 mL trifluoroacetic acid. The solvent was eliminated in vacuo, the residue taken up with CHCl₃ and evaporated down again, then taken up in acetone and the hydrochloride was precipitated with ethereal HCl. Yield: 14.3 g (56%).

cis- and trans-4-morpholino-cyclohexylamine



15 Dibenzyl-4-morpholino-cyclohexylamine

3.9 g (30 mmol) of 4-dibenzylcyclohexanone were dissolved in 100 mL of CH₂Cl₂ and stirred for 12 h at RT with 3.9 g (45 mmol) of morpholine and 9.5 g (45 mmol) NaBH(OAc)₃. Then water and potassium carbonate were added, the org. phase was separated off, dried and the solvent was eliminated in vacuo. The residue was purified through a silica gel column (about 20 mL silica gel; about 500 mL of ethyl acetate 90/ methanol 10 + 1% conc. ammonia). The appropriate fractions were evaporated down in vacuo. Yield : 6.6 g (60%) of cis-isomer and 2 g (18%) of trans-isomer.

Alternatively the *trans*-dibenzyl-4-morpholino-cyclohexylamine may be prepared by the following method:

33 g (112 mmol) of 4-dibenzylcyclohexanone were dissolved in 300 mL MeOH,
5 combined with 17.4 g (250 mmol) of hydroxylamine hydrochloride and stirred for 4 h at 60°C. The solvent was evaporated down in vacuo, combined with 500 mL water and 50 g potassium carbonate and extracted twice with 300 mL of dichloromethane. The org. phase was dried, evaporated down in vacuo, the residue was crystallised from petroleum ether, dissolved in 1.5 L of EtOH and
10 heated to 70°C. 166 g of sodium were added batchwise and the mixture was refluxed until the sodium dissolved. The solvent is eliminated in vacuo, the residue combined with 100 mL water and extracted twice with 400 mL of ether. The org. phase is washed with water, dried, evaporated down in vacuo and the *trans* isomer is isolated using a column (about 1.5 L silica gel; about 2 L of ethyl acetate
15 80/ methanol 20 + 2 % conc. ammonia). Yield: 12.6 g (41.2%).
6.8 g (23 mmol) of *trans*-1-amino-4-dibenzylaminocyclohexane was dissolved in 90 mL of DMF and stirred for 8 h at 100°C with 5 mL (42 mmol) of 2,2'-dichloroethyl ether and 5 g of potassium carbonate. After cooling 30 mL of water was added, the precipitated crystals were suction filtered and purified through a
20 short column (about 20 mL silica gel, about 100 mL ethyl acetate). The residue is crystallised from methanol and conc. HCl as the dihydrochloride. Yield: 7.3 g (72.4%).


trans-4-morpholino-cyclohexylamine

25 7.2 g (16.4 mmol) of *trans*-dibenzyl-4-morpholino-cyclohexylamine were dissolved in 100 mL of MeOH and hydrogenated on 1.4 g of Pd/C (10%) at 30-50°C. The solvent was eliminated in vacuo and the residue was crystallised from ethanol and conc. HCl. Yield: 3.9 g (93%); m.p. 312 °C.

30

The *cis* isomer can be prepared analogously.

cis- and trans-4-piperidino-cyclohexylamine



5 trans-dibenzyl-4-piperidino-cyclohexylamine

2.0 g (6.8 mmol) of trans-1-amino-4-dibenzylaminocyclohexane (see Example 2) was dissolved in 50 mL DMF and stirred for 48 h at RT with 1.6 g (7 mmol) of 1,5-dibromopentane and 2 g of potassium carbonate. The mixture was cooled,
10 combined with water, extracted twice with 100 mL of dichloromethane, dried and the solvent was eliminated in vacuo. The residue is purified over a column (about 100 mL silica gel, about 500 mL ethyl acetate 80/methanol 20 +1% conc. ammonia). The desired fractions were evaporated down in vacuo and crystallised from petroleum ether. Yield: 1.2 g (49%).


15

trans-4-piperidino-cyclohexylamine

1.7 g (4.8 mmol) of trans-dibenzyl-4-piperidino-cyclohexylamine were dissolved in 35 mL MeOH and hydrogenated on 350 mg of Pd/C (10%) at 20°C. The solvent was eliminated in vacuo and the residue crystallised from ethanol and conc. HCl.
20 Yield: 1.1 g (78%).

The *cis* isomer may be prepared analogously.

cis- and trans-4-(4-phenyl-piperazin-1-yl)-cyclohexylamine




- 4.1 g (25.3 mmol) of 4-dibenzylcyclohexanone was dissolved in 50 mL of
5 dichloromethane and stirred for 12 h at RT with 7.4 g (25.3 mmol) of
N-phenylpiperazine and 7.4 g (35 mmol) of $\text{NaBH}(\text{OAc})_3$. Then water and
potassium carbonate were added, the org. phase was separated off, dried and the
solvent was eliminated in vacuo. The residue was purified over a silica gel column
(ethyl acetate 80/ methanol 20 + 0.5% conc. ammonia). Yield: 1.7 g (15.8%) of cis-
10 isomer and 0.27 (2.5%) of trans-isomer.

trans-4-(4-phenyl-piperazin-1-yl)-cyclohexylamine

- 15 270 mg (0.61 mmol) of trans-dibenzyl-[4-(4-phenyl-piperazin-1-yl)-cyclohexyl]-
amine were dissolved in 5 mL MeOH and hydrogenated on 40 mg of Pd/C (10%)
at 20-30 °C. The solvent was eliminated in vacuo and the residue crystallised from
ethanol and conc. HCl. Yield: 110 mg (69%).
- 20 The cis isomer may be prepared analogously.

cis- and trans-4-(4-cyclopropylmethyl-piperazin-1-yl)-cyclohexylamine



9.8 g (33.4 mmol) of 4-dibenzylcyclohexanone was dissolved in 100 mL dichloromethane and stirred for 12 h at RT with 5.6 g (40 mmol) of N-cyclopropylmethylpiperazine and 8.5 g (40 mmol) of NaBH(OAc)₃. Then water and potassium carbonate were added, the org. phase was separated off, dried and
5 the solvent was eliminated in vacuo. The residue was purified over a silica gel column (about 50 mL silica gel, about 3 L ethyl acetate 95/ methanol 5 + 0.25% conc. ammonia. The appropriate fractions were evaporated down in vacuo. The faster eluting cis compound crystallised from ethyl acetate. The trans-compound was crystallised from ethanol + conc. HCl. Yield: 8.5 g (61%) cis-isomer and 2.2
10 (13%) trans-isomer.

cis-4-(4-cyclopropylmethyl-piperazin-1-yl)-cyclohexylamine

8.5 g (20 mmol) of cis-dibenzyl-[4-(4-cyclopropylmethyl-piperazin-1-yl)-cyclohexyl]-
15 amine were dissolved in 170 mL MeOH and hydrogenated on 1.7 g Pd/C (10%) at 30-50 °C. The solvent was eliminated in vacuo and the residue was crystallised from ethanol and conc. HCl. Yield: 4.4 g (91%).

The trans-isomer may be prepared analogously.

Synthesis of the Examples

Example 152

- 5 0.15g of the compound Z10, 0.14 g TBTU, 0.13 mL DIPEA were dissolved in dichloromethane and stirred for 20 minutes at 25°C. Then 90 µL 1-(3-aminopropyl)-4-methylpiperazine was added and stirred for a further 2 hours at 25°C. The solution was then diluted with dichloromethane and extracted with water. The product was precipitated by the addition of petroleum ether, ether and
10 ethyl acetate to the organic phase. Yield: 0.16 g of beige solid

Example 164

- 15 0.10g of the compound Z10, 0.1 g TBTU, 0.08 mL DIPEA were dissolved in 4 mL dichloromethane and stirred for 20 minutes at 25°C. Then 44 µL dimethylaminopropylamine were added and stirred for a further 2 hours at 25°C. The solution was then diluted with dichloromethane and extracted with water. The product was precipitated by the addition of petroleum ether, ether and acetone to
20 the organic phase. Yield: 0.08 g yellow solid.

Example 242

- 25 0.15g of the compound Z10, 0.14 g TBTU, 0.13 mL DIPEA were dissolved in 5 mL dichloromethane and stirred for 20 minutes at 25°C. Then 75 µL 1-(2-aminoethyl) piperidine were added and stirred for a further 2 hours at 25°C. The solution was then diluted with dichloromethane and extracted with water. The product was precipitated by the addition of petroleum ether, ether and ethyl acetate to the
30 organic phase. Yield: 0.14 g yellow solid.

Example 188

0.1 g of the compound Z2, 0.09 g TBTU, 0.05 mL DIPEA were dissolved in 15 mL dichloromethane and stirred for 20 minutes at 25°C. Then 33 mg 1-methyl-4-aminopiperidin were added and the mixture was stirred for a further 3 hours at 25°C. The solution was extracted with 20 mL water, then evaporated down in vacuo. The product was crystallised using ether. Yield: 0.047 g of white crystals.

Example 203

10

0.1 g of the compound Z2, 0.09 g TBTU, 0.5 mL DIPEA were dissolved in 15 mL dichloromethane and stirred for 30 minutes at 25°C. Then 50 mg 4-amino-1-benzylpiperidin were added and the mixture was stirred for a further 3 hours at 25°C. The solution was extracted with 20 mL water, then evaporated down in vacuo. Then the residue was chromatographed over silica gel and the isolated product was crystallised with ether. Yield: 0.015 g of white crystals.

Example 94

20 0.17 g of the compound Z1, 0.19 g TBTU, 0.11 mL DIPEA were dissolved in 50 mL dichloromethane and stirred for 30 minutes at 25°C. Then 63 mg of 1-methyl-4-aminopiperidine were added and the mixture was stirred for a further 17 hours at 25°C. 50 mL of water and 1 g of potassium carbonate were added to the solution and the organic phase was separated off using a phase separation cartridge, then 25 evaporated down in vacuo. Then the product was purified by silica gel chromatography and the purified product was crystallised with ether. Yield: 0.1 g of white crystals.

Example 95

30

0.17 g of the compound Z1, 0.19 g TBTU, 0.11 mL DIPEA were dissolved in 50 mL dichloromethane and stirred for 30 minutes at 25°C. Then 77 mg of exo-3-β-

amino-tropane were added and the mixture was stirred for a further 17 hours at 25°C. 50 mL of water and 1 g of potassium carbonate were added to the solution and the organic phase was separated off using a phase separation cartridge, then evaporated down in vacuo. Then the product was purified by silica gel chromatography and the purified product was crystallised with ether. Yield: 0.03 g of white crystals.

Example 46

0.15 g of the compound Z3, 0.12 g TBTU, 0.12 mL DIPEA were dissolved in 5 mL dichloromethane and stirred for 30 minutes at 25°C. Then 50 mg 1-methyl-4-aminopiperidin were added and the mixture was stirred for a further 2.5 hours at 25°C. The solution was then extracted with water and then evaporated down. The residue was dissolved in warm ethyl acetate and crystallised from ether and petroleum ether. Yield: 0.025 g of white crystals. M.p.: 203°C as the base

Example 80

0.2 g of the compound Z8, 0.2 g of TBTU, 0.1 mL of DIPEA were dissolved in 10 mL dichloromethane and stirred for 30 minutes at 25°C. Then 100 mg of 1-methyl-4-aminopiperidine were added and the mixture was stirred for a further 17 hours at 25°C. The solution was then extracted with a dilute potassium carbonate solution and evaporated down. The residue was crystallised using ether. Yield: 0.12 g of white crystals

Example 190

0.2 g of compound Z8, 0.2 g of TBTU, 0.3 mL of DIPEA were dissolved in 5 mL dichloromethane and stirred for 1h at 25°C. Then 0.13 g of 4-amino-1-benzylpiperidine were added and the mixture was stirred for a further hour at 25°C. The solution was then diluted with 10 mL methylene chloride and extracted

with 20 mL water. Then the product was purified over silica gel and crystallised from ethyl acetate and ether. Yield: 0.23 g of the compound Z8.

0.23 g of the benzylamine Z8 are dissolved in 10 mL methanol, combined with 50 mg of Pd/C and hydrogenated under 3 bar for 3h at 25°C. By adding petroleum
5 ether and ethyl acetate white crystals are produced. These are chromatographed over silica gel and crystallised from ethyl acetate and ether. Yield: 0.075 g of white crystals.

Example 196

10

0.1g of compound Z10, 0.09 g of TBTU, 0.3 mL of DIPEA were dissolved in 4 mL of dichloromethane and stirred for 20 minutes at 25°C. Then 67 mg xx amine was added and stirred for a further 2 hours at 25°C. The solution was then diluted with dichloromethane and extracted with water. It was then chromatographed over
15 silica gel and the residue was dissolved in acetone, combined with ethereal HCl and the precipitate formed was isolated. Yield: 0.09 g light yellow solid

Example 166

20 0.1 g of the compound Z10, 0.11 g of TBTU, 0.14 mL of DIPEA were dissolved in 2 mL dimethylformamide and stirred for 3h at 50°C. Then 55 mg of 4-morpholinomethylphenylamine was added. The reaction mixture was then cooled to ambient temperature within 17 h. Then the dimethylformamide was eliminated in vacuo, the residue was taken up in dichloromethane and extracted
25 with water. It was then chromatographed over silica gel and the product crystallised from ethyl acetate and ether. Yield: 0.06 g yellowish crystals

Example 81

30 0.2 g of the compound Z4, 0.2 g of TBTU, 0.1 mL of DIPEA were dissolved in 10 mL dichloromethane and stirred for 30 minutes at 25°C. Then 0.1 g of 1-methyl-4-aminopiperidine were added and the mixture was stirred for a further 17 hours at

25°C. The solution was then extracted with aqueous potassium carbonate solution and then evaporated down. The product was crystallised using ether. Yield: 0.16 g of white crystals.

5 Example 162

0.1 g of the compound Z5, 0.07 g of TBTU, 0.15 mL of DIPEA were dissolved in 5 mL dichloromethane and stirred for 20 minutes at 25°C. Then 0.04 g 1-methyl-4-aminopiperidine were added and the mixture was stirred for a further 2 hours at
10 25°C. The solution was then diluted with 15 mL dichloromethane and extracted with 20 mL water. The residue was dissolved in MeOH and acetone, combined with 1 mL ethereal HCl and evaporated down. A crystalline product was produced using ether, ethyl acetate and a little MeOH. Yield: 0.1 g of white crystals.

15 Example 88

0.1 g of the compound Z6, 0.12 g of TBTU, 0.12 mL of DIPEA were in 10 mL dichloromethane dissolved and stirred for 30 minutes at 25°C. Then 0.04 g of 1-methyl-4-aminopiperidine were added and the mixture was stirred for a further 2
20 hours at 25°C. The solution was then diluted with 10 mL dichloromethane and extracted with 10 mL water. A crystalline product was produced using ether, ethyl acetate and petroleum ether. Yield: 0.6 g of white crystals.

Example 89

25

0.1 g of the compound Z6, 0.08 g of TBTU, 0.08 mL of DIPEA were dissolved in 10 mL dichloromethane and stirred for 30 minutes at 25°C. Then 37 µL g N,N-dimethylneopentanediamine were added and the mixture was stirred for a further 2 hours at 25°C. The solution was then diluted with 10 mL dichloromethane and
30 extracted with 10 mL water. The product was then chromatographed over silica gel and crystallised from ethyl acetate, ether and petroleum ether. Yield: 0.005 g of white crystals.

Example 26

0.15 g of the compound Z7, 0.16 g of TBTU, 1 mL of DIPEA were dissolved in 5
5 mL dichloromethane and stirred for 30 minutes at 25°C. Then 0.1 g
4-morpholinocyclohexylamine were added and the mixture was stirred for a further
17 hours at 25°C. The residue was then combined with 10 mL of 10% potassium
carbonate solution, the precipitate was isolated and washed with water. It was
then dissolved in dichloromethane and evaporated down again. The product was
10 crystallised from ethyl acetate. Yield: 0.1 g of white crystals.

Example 9

150 mg of the compound Z9 and 93 mg of amine were dissolved in 5 mL
15 dichloromethane and stirred with 160 mg of TBTU and 1 mL of DIPEA for 12 h at
RT. The solvent was eliminated in vacuo, the residue was combined with 10 mL
of 10% potassium carbonate solution. The precipitate was suction filtered, washed
with water, taken up in dichloromethane, dried and the solvent eliminated in
vacuo. The residue was crystallised from ethyl acetate. Yield: 82.0 mg; m.p. 253°C
20 (as base).

Example 16

150 mg of the compound Z8 and 73 mg of trans-4-piperidino-cyclohexylamine
25 were dissolved in 5 mL dichloromethane and stirred with 160 mg (0.50 mmol) of
TBTU and 1 mL of DIPEA for 12 h at RT. The solvent was eliminated in vacuo, the
residue was combined with 10 mL of 10% potassium carbonate solution. The
precipitate was suction filtered, washed with water, taken up in dichloromethane,
dried and the solvent eliminated in vacuo. The residue was crystallised from ethyl
30 acetate. Yield: 87.0 mg; m.p. 237 °C (as base).

Example 37

100 mg of the compound Z9 and 42 mg of 3-amino-1-ethyl-pyrrolidine were dissolved in 10 mL dichloromethane and stirred with 90 mg of TBTU and 0.5 mL of
5 DIPEA for 12 h at RT. The solvent was eliminated in vacuo, the residue was combined with 10 mL of 10% potassium carbonate solution. The precipitate was suction filtered, washed with water, taken up in dichloromethane, dried and the solvent was eliminated in vacuo. The residue was crystallised from ethyl acetate/petroleum ether. Yield: 24.0 mg .

10

Example 120

100 mg of the compound Z11 and 73 mg of 4-amino-1-tetrahydro-4H-pyran-4-yl-piperidine were dissolved in 10 mL dichloromethane and stirred with 90 mg of
15 TBTU and 0.5 mL of DIPEA for 1 h at RT. The solvent was eliminated in vacuo, the residue was combined with 10 mL of 10% potassium carbonate solution. The precipitate was suction filtered, washed with water, taken up in dichloromethane, dried and the solvent was eliminated in vacuo. The residue was crystallised from ethyl acetate/petroleum ether. Yield: 89 mg.

20

Example 212

150 mg of the compound Z5 and 150 mg of trans-4-(4-cyclopropylmethyl-piperazin-1-yl)-cyclohexylamine (as the hydrochloride) were dissolved in 5 mL of
25 dichloromethane and stirred with 160 mg of TBTU and 2 mL of DIPEA for 2 h at RT. The solvent was eliminated in vacuo, the residue was combined with 10 mL of 10% potassium carbonate solution. The precipitate was suction filtered, washed with water, taken up in dichloromethane, dried and the solvent eliminated in vacuo. The residue was purified over a column (20 mL silica gel, 300 mL ethyl acetate 90/
30 methanol 10 + 2% conc. ammonia). The appropriate fractions were evaporated down in vacuo and crystallised from ethyl acetate. Yield: 140 mg; m.p. 187 °C (as base).

Example 232

390 mg of the compound Z11 and 240 mg of trans-4-(4-tbutoxycarbonyl-piperazin-1-yl)-cyclohexylamine were dissolved in 2.5 mL of NMP and stirred with
5 482 mg of TBTU and 1 mL triethylamine for 2 h at RT. Then 100 mL of water and 200 mg of potassium carbonate were added, the precipitate was suction filtered, washed with water and purified through a silica gel column. The appropriate fractions were evaporated down in vacuo, dissolved in 2 mL dichloromethane, combined with 2 mL of trifluoroacetic acid and stirred for 2 h at RT, combined with
10 another 100ml of water and 200 mg potassium carbonate and the precipitate was suction filtered and washed with water. Then the precipitate was purified through a silica gel column. The appropriate fractions were evaporated down in vacuo and the residue was crystallised from ethanol and conc. hydrochloric acid. Yield: 95 mg; m.p. 291 °C.

15

Example 213

60 mg of the compound of Example 232 was dissolved in 10 mL ethyl acetate and stirred with 1 mL of acetic anhydride and 1 mL of triethylamine for 30 min. at RT.
20 The solvent was eliminated in vacuo, the residue combined with water and ammonia, the crystals precipitated were suction filtered and washed with water and a little cold acetone. Yield: 40 mg; m.p. 248 °C.

Example 218

25

1.2 g of the compound Z9 and 0.5g of 1,4-dioxaspiro[4.5]dec-8-ylamine were dissolved in 20 mL dichloromethane and stirred with 1.28 g of TBTU and 4 mL of triethylamine for 12 h at RT. Then 50 mL of water and 0.5 g of potassium carbonate were added, the org. phase was separated off, dried and evaporated
30 down in vacuo. The residue was crystallised from ethyl acetate, combined with 25 mL of 1 N hydrochloric acid and 20 mL of methanol and stirred for 30 min at 50°C.

The methanol was eliminated in vacuo, the precipitate was suction filtered, washed with water and dried.

5 The residue was taken up in 20 mL dichloromethane, stirred with 0.5 g of thiomorpholine and 0.5 g of $\text{NaBH}(\text{OAc})_3$ for 12 h at RT. Then water and potassium carbonate were added, the org. phase was separated off, dried and the solvent was eliminated in vacuo. The residue was purified over a silica gel column. The appropriate fractions were evaporated down in vacuo and the hydrochloride was precipitated with ethereal HCl. Yield: 86 mg of trans-isomer; amorphous powder.

10

Example 187

200 mg of the compound Z3 in 5 mL dichloromethane was combined with 0.1 mL of diisopropylethylamine and 180 mg of TBTU and stirred for 30 min. Then 191 mg
15 of 4-(4-methyl-piperazin-1-yl)-phenylamine were added and the mixture was stirred overnight. The reaction mixture was combined with water and the aqueous phase extracted with dichloromethane. The combined organic phases were dried over Na_2SO_4 and Z9 orated down. The residue was purified by column chromatography (eluant: dichloromethane/methanol 100:7).

20 Yield: 128 mg (light yellow crystals)

The compounds of formula (I) listed in Table 1, *inter alia*, are obtained analogously to the procedure described hereinbefore. The abbreviations X_1 , X_2 , X_3 , X_4 and X_5 used in Table 1 in each case denote a link to a position in the general formula
25 shown under Table 1 instead of the corresponding groups R^1 , R^2 , R^3 , R^4 and $L-R^5$.

Table 1

54

















7	0.084	0.865	H	X_2  CH ₃	R		X_4 - O - CH ₃		
8	0.063		H	X_2  CH ₃	R		H		
9	0.057	0.207	H	X_2  CH ₃	R		X_4 - O - CH ₃		253
10	0.091		H	X_2  CH ₃	R		H		
11	0.086	1.300	H	X_2  CH ₃	R		H		
12	0.099	1.069	H	X_2  CH ₃	R		H		
13	0.026	1.320	H	X_2  CH ₃	R		CH_3 - O - X_4		
14	0.055	3.000	H	X_2  CH ₃	R		H		

Table 1

55









15	0.020	1.000	H	X_2  CH ₃	R		X_3  O-CH ₃		231
16	0.017	0.483	H	X_2  CH ₃	R		X_3  O-CH ₃		237
17	0.024	0.730	H	X_2  CH ₃	R		CH ₃  O- X_4		
18	0.040	1.764	H	X_2  CH ₃	R		H		
19	0.013	0.747	H	X_2  CH ₃	R		X_3  O-CH ₃		254
20	0.083	3.000	H	X_2  CH ₃	R		CH ₃  O- X_4		
21	0.009	0.488	H	X_2  CH ₃	R		CH ₃  O- X_4		
22	0.008	0.347	H	X_2  CH ₃	R		CH ₃  O- X_4		

Table 1

56









23	0.013	0.679	H	X_2  CH_3	R		CH_3-O-X_4		
24	0.017	0.277	H	X_2  CH_3	R		X_4-O-CH_3		152
25	0.038	1.000	H	X_2  CH_3	R		X_4-O-CH_3		254
26	0.059	0.424	H	X_2  CH_3	R	X_3  CH_3 H_3C	X_4-O-CH_3		
27	0.014	0.164	H	X_2  CH_3	R	X_3  CH_3 H_3C	X_4-O-CH_3		292
28	0.018	0.160	H	X_2  CH_3	R	X_3  CH_3 H_3C	X_4-O-CH_3		160
29	0.049	1.000	H	X_2  CH_3	R	X_3  CH_3 H_3C	X_4-O-CH_3		251
30	0.012	0.574	H	X_2  CH_3	R		CH_3-O-X_4		

Table 1

57























31	0.023	3.000	H	X_2  CH ₃	R		H		
32	0.049	3.000	H	X_2  CH ₃	R				
33	0.097	3.000	H	X_2  CH ₃	R		H		
34	0.055	1.000	H	X_2  CH ₃	R				242
35	0.088	1.000	H	X_2  CH ₃	R				213
36	0.021	1.000	H	X_2  CH ₃	R				232
37	0.072	0.762	H	X_2  CH ₃	R				160
38	0.087	0.278	H	X_2  CH ₃	R				225

Table 1

58

39	0.075	1.000	H	X_2 -CH ₃	R		H		
40	0.038	0.187	H	X_2 -CH ₃	R		X_4 -O-CH ₃		226
41	0.083	3.000	H	X_2 -CH ₃	R		CH_3 -O- X_4		
42	0.076	2.310	H	X_2 -CH ₃	R		CH_3 -O- X_4		
43	0.094	3.000	X_1 -CH ₃	X_2 -CH ₃			H_3 C-O- X_4		218
44	0.009	0.204	H	X_2 -CH ₃	R		H		
45	0.022	0.792	H	X_2 -CH ₃	R		H_3 C-O- X_4		
46	0.012	0.208	H	X_2 -CH ₃	R		H_3 C-O- X_4		203

Table 1

59



















47	0.028		H	X_2 	R		H		
48	0.083	3.000	H	X_2 			H		
49	0.059	0.529	H	X_2 					
50	0.081	0.436	H	X_2 					
51	0.080	0.150	H	X_2 					
52	0.079	0.153	H	X_2 					
53	0.076	0.245	X_1 	X_2 					
54	0.031	0.226	H	X_2 	R				

Table 1

60

55	0.013	0.086	H	X_2 -CH ₃	R				
56	0.064	0.324	H	X_2 -CH ₃	R				
57	0.038	0.067	H	X_2 -CH ₃	R				
58	0.010	0.023	H	X_2 -CH ₃	R				
59	0.095	1.000	H	X_2 -CH ₃	R				
60	0.041	0.189	H	X_2 -CH ₃	R				
61	0.065	1.040	X_1 -CH ₃	X_2 -CH ₃					217
62	0.058	0.538	H	X_2 -CH ₃	R				259

Table 1

61







63	0.081	0.206	H	X_2  CH ₃	R				184
64	0.064	0.894	H	X_2  CH ₃	R				175
65	0.052	0.216	H	X_2  CH ₃	R				172
66	0.064	0.250	H	X_2  CH ₃	R				164
67	0.052	0.566	H	X_2  CH ₃	R				115
68	0.092	1.242	H	X_2  CH ₃	R		H		
69	0.055	0.633	H	X_2  CH ₃	R		H		
70	0.092	1.000	H	X_2  CH ₃	R		H		

Table 1

62





















71	0.035	0.227	H		R			146
72	0.050		H		R			
73	0.091		H		R	H		
74	0.043		H		R			
75	0.046	0.557	H		R			
76	0.068		H		R			
77	0.055	1.000	H		R	H		
78	0.035	0.610	H		R	H		

Table 1

63


























79	0.043		H		R		H		
80	0.038	0.286	H						
81	0.050	0.237	H		R				194
82	0.083	1.580	H		R				
83	0.069	0.263	H		R				
84	0.015	0.312	H		R				
85	0.025	0.754	H		R		H		
86	0.038	0.024	H		R				

Table 1

64

87	0.029	0.066	H		R				
88	0.020	1.018	H		R				
89	0.047	0.245	H		R				
90	0.032	0.137	H		R				
91	0.041	1.780	H		R				
92	0.043		H		R		H		181
93	0.060		H		R		H		
94	0.018	0.510	H		R				178

Table 1

65













95	0.047	1.000	H	X_2 	R		H_3C-X_4			
96	0.011	0.577	H	X_2 	R		H_3C-O-X_4		203	
97	0.032	0.066	H	X_2 	R		H_3C-O-X_4		179	
98	0.077	1.319	H	X_2 	R		X_4-O-CH_3			
99	0.025	0.209	H	X_2 	R	$H_3C-CH(X_3)-CH_3$	X_4-O-CH_3		280	
100	0.058	0.124	H	X_2 	R	$H_3C-CH(X_3)-CH_3$	X_4-O-CH_3		233	
101	0.031	0.124	H	X_2 	R	$H_3C-CH(X_3)-CH_3$	CH_3-O-X_4			
102	0.006	0.053	H	X_2 	R	$H_3C-CH(X_3)-CH_3$	CH_3-O-X_4			

Table 1

66













103	0.005	0.076	H	X_2 	R				
104	0.041	0.133	H	X_2 	R				
105	0.015	0.055	H	X_2 	R				229
106	0.025	1.000	H	X_2 	R				213
107	0.043	0.363	H	X_2 	R				285
108	0.016	1.000	H	X_2 	R				
109	0.031	0.760	H	X_2 	R				
110	0.027	0.122	H	X_2 	R				278

Table 1

67



















111	0.023	0.503	H	X_2 	R				
112	0.075	0.345	H	X_2 	R				264
113	0.037	0.687	H	X_2 	R				279
114	0.064	0.766	H	X_2 	R				234
115	0.016	0.076	H	X_2 	R				
116	0.006	0.174	H	X_2 	R				
117	0.010	0.370	H	X_2 	R				
118	0.010	0.279	H	X_2 	R				

Table 1

68




























119	0.008	0.438	H		R				
120	0.014	0.374	H		R				154
121	0.011	0.427	H		R				
122	0.009	0.261	H		R				
123	0.043	0.490	H		R				
124	0.006	0.391	H		R				
125	0.049	0.795	H		R				283
126	0.051	0.742	H		R				238

Table 1

69

























127	0.045	0.787	H	X_2 	R				221
128	0.015	0.282	H	X_2 	R				257
129	0.035	0.646	H	X_2 	R	X_3 			284
130	0.015	0.091	H	X_2 	R				236
131	0.023	0.101	H	X_2 	R				141
132	0.062	1.000	H	X_2 	R	X_3 			268
133	0.011	0.075	H	X_2 	R				272
134	0.015	0.063	H	X_2 	R				319

Table 1

70

135	0.012	0.699	H	X_2 -CH ₃	R	$H_3C-CH(X_3)-CH_3$	X_4-O-CH_3		289
136	0.041	0.821	H	X_2 -CH ₃	R		X_4-O-CH_3		201
137	0.039	0.148	H	X_2 -CH ₃	R		X_4-O-CH_3		223
138	0.043	1.137	H	X_2 -CH ₃	R		X_4-O-CH_3		217
139	0.097	3.000	H	X_2 -CH ₃	R	$H_3C-CH(X_3)-CH_3$	H_3C-O-X_4		112
140	0.029	1.970	H	X_2 -CH ₃	R	$H_3C-CH(X_3)-CH_3$	H_3C-O-X_4		215
141	0.057		H	X_2 -CH ₃	R	$H_3C-CH(X_3)-CH_3$	H_3C-O-X_4		198
142	0.030		H	X_2 -CH ₃	R	$H_3C-CH(X_3)-CH_3$	H_3C-O-X_4		192

Table 1

71

143	0.086		H	X_2 -CH ₃	R	$H_3C-CH(X_3)-CH_3$	H		139
144	0.044		H	X_2 -CH ₃	R	$H_3C-CH(X_3)-CH_3$	H		191
145	0.043	3.000	H	H_3C-X_2	R	$H_3C-C(CH_3)(X_3)-CH_3$	CH_3-O-X_4		
146	0.066	0.966	H	X_2 -CH ₃	R	$H_3C-CH(X_3)-CH_3$	X_4-O-CH_3		
147	0.055	1.760	H	H_3C-X_2	R	$H_3C-C(CH_3)(X_3)-CH_3$	H		
148	0.087		H	X_2 -CH ₃	R	$H_3C-CH(X_3)-CH_3$	CH_3-O-X_4		
149	0.085		H	X_2 -CH ₃	R	$H_3C-CH(X_3)-CH_3$	CH_3-O-X_4		
150	0.043		H	X_2 -CH ₃	R	$H_3C-CH(X_3)-CH_3$	H		

Table 1

72








151	0.043		H		R				155
152	0.081		H		R				147
153	0.072	2.640	H		R				128
154	0.097		H		R		H		
155	0.079		H		R		H		
156	0.048	0.481	H		R				173
157	0.051		H		R				134
158	0.043	0.536	H		R				230

Table 1

73

159	0.029	1.000	H	X_2-CH_3	R	$X_3-CH_2-CH_2-CH_3$	CH_3-O-X_4	$X_5-CH_2-CH_2-CH_2-N(CH_3)_2$	260
160	0.037	0.728	H	X_2-CH_3	R	$X_3-CH_2-CH_2-CH_2-CH_3$	CH_3-O-X_4	$X_5-CH_2-CH_2-CH_2-N(CH_3)-CH(CH_3)_2$	193
161	0.081		X_1-CH_3	X_2-CH_3		$X_3-CH_2-CH_2-CH_2-CH_3$	H_3C-O-X_4	$X_5-CH_2-CH_2-CH_2-N(CH_3)-CH_2-CH_3$	199
162	0.071	0.903	X_1-CH_3	X_2-CH_3		$X_3-CH_2-CH_2-CH_2-CH_3$	H_3C-O-X_4	$X_5-CH_2-CH_2-CH_2-N(CH_3)-CH_2-CH_3$	254
163	0.071	1.000	H	X_2-CH_3	R	$X_3-CH_2-CH_2-CH_2-CH_3$	CH_3-O-X_4	$X_5-CH_2-CH_2-CH_2-N(CH_3)-CH_2-CH_3$	249
164	0.030	0.975	H	X_2-CH_3	R	$X_3-CH_2-CH_2-CH_2-CH_3$	CH_3-O-X_4	$X_5-CH_2-CH_2-CH_2-N(CH_3)-CH_2-CH_3$	
165	0.046	0.668	H	X_2-CH_3	R	$X_3-CH_2-CH_2-CH_2-CH_3$	CH_3-O-X_4	$X_5-CH_2-CH_2-CH_2-N(CH_3)-CH_2-CH_3$	
166			H	X_1-CH_3	R	$X_3-CH_2-CH_2-CH_2-CH_3$	CH_3-O-X_4	$X_5-CH_2-CH_2-CH_2-N(CH_3)-CH_2-CH_3$	

Table 1

























167	0.034	0.817	H		R				
168	0.042	0.088	H						
169	0.044	0.503	H						
170	0.045	0.427	H						170
171	0.031	0.790	H						196
172	0.020	0.772	H						166
173	0.070	0.871	H						
174	0.000	0.155	H						

Table 1

75

175	0.000	0.296	H		R				
176	0.000	0.301	H		R				
177	0.000	0.082	H		R				
178	0.000	1.000	H		R				
179	0.000	1.000	H		R				
180	0.000	0.147	H		R				136
181	0.000	0.754	H		R				
182			H		R				

Table 1

76







183			H		R				
184			H		R				
185			H		R				217
186			H		R				287
187			H		R				223
188			H		R				161
189			H		R				267
190			H		R				

Table 1

77

191			H		R				
192			H		R				
193			H		R				
194			H		R				
195			H		R				
196			H		R				
197			H						
198			H		R				301

Table 1

78

199			H		R				291
200			H		R				
201			H		R				
202									
203			H		R				
204			H		R				
205			H		R				255
206			H		R				190

Table 1

79









207			H	X_2-CH_3	R	$X_3-CH(CH_3)_2$	X_4-O-CH_3		236
208			H	X_2-CH_3	R	$X_3-CH(CH_3)_2$	X_4-O-CH_3		192
209			H	X_2-CH_3	R	$X_3-CH(CH_3)_2$	X_4-O-CH_3		211
210			H	X_2-CH_3	R	$X_3-C_5H_9$	X_4-O-CH_3		210
211			X_1-CH_3	X_2-CH_3		$X_3-CH(CH_3)_2$	X_4-O-CH_3		182
212			X_1-CH_3	X_2-CH_3		$X_3-CH(CH_3)_2$	X_4-O-CH_3		187
213			H	X_2-CH_3	R	$X_3-C_6H_{11}$	X_4-O-CH_3		248
214			H	X_2-CH_3	R	$X_3-CH(CH_3)_2$	X_4-O-CH_3		338

Table 1

80
















215			X_1-CH_3	X_2-CH_3			X_3-O-CH_3		232
216			H	X_2-CH_3	R		X_3-O-CH_3		217
217			H	X_2-CH_3	R		X_3-O-CH_3		139
218			H	X_2-CH_3	R		X_3-O-CH_3		
219			H	X_2-CH_3	R		X_3-O-CH_3		298
220			H	X_2-CH_3	R		X_3-O-CH_3		279
221			H	X_2-CH_3	R		X_3-O-CH_3		336
222			H	X_2-CH_3	R		X_3-O-CH_3		241

Table 1

81
























223		1	H	X_2  CH ₃	R				
224	1.056	1	H	X_2  CH ₃	R				
225	1.18	1	H	X_2  CH ₃	R		H		
226	0.295	1	H	X_2  CH ₃	R				
227	0.853	0.946	H	X_2  CH ₃	R				
228		1	H	X_2  CH ₃	R				
229		0.381	H	X_2  CH ₃	R				
230		0.193	H	X_2  CH ₃	R				













Table 1

82

231		0.17	H		R				
232		1	H		R				
233			H		R				
234			H		R				
235			H		R				
236			H		R				
237	0.012	1	H		R				
238	0.015	1	H		R				

Table 1

83

239	0.051	0.07	H		R				
240	0.017	0.054	H		R				
241	0.044	0.093	H		R				
242	0.019	0.524	H		R				
243	0.018	0.975	H		R				
244	0.0115		H		R				

As has been found, the compounds of general formula (I) are characterised by their wide range of applications in the therapeutic field. Particular mention should be made of those applications in which the inhibition of specific cell cycle kinases, particularly the inhibiting effect on the proliferation of cultivated
5 human tumour cells but also the proliferation of other cells, such as endothelial cells, for example, plays a part.

As could be demonstrated by FACS analysis, the inhibition of proliferation brought about by the compounds according to the invention is mediated by
10 the arrest of the cells, particularly at the G2/M phase of the cell cycle. The cells arrest, independently of the cells used, for a specific length of time in this phase of the cell cycle before programmed cell death is initiated. An arrest in the G2/M phase of the cell cycle is triggered, for example, by the inhibition of specific cell cycle kinases. Studies in model organisms such as
15 *Schizosaccharomyces pombe* or *Xenopus*, or investigations in human cells have shown that the transition from the G2 phase to mitosis is regulated by the CDK1/cyclin B kinase (Nurse, 1990). This kinase, which is also known as the "mitosis promoting factor" (MPF), phosphorylates and thereby regulates a number of proteins, such as e.g. nuclear lamins, kinesin-like motor proteins,
20 condensins and Golgi matrix proteins, which play an important part in the breakdown of the nuclear envelope, in centrosome separation, the formation of the mitotic spindle apparatus, chromosome condensation and the breakdown of the Golgi apparatus (Nigg. E., 2001). A murine cell line with a temperature-sensitive CDK1 kinase mutant shows a rapid breakdown of the
25 CDK1 kinase and a subsequent arrest in the G2/M phase after a temperature increase (Th'ng et al., 1990). The treatment of human tumour cells with inhibitors against CDK1/cyclin B such as e.g. butyrolactone also leads to an arrest in the G2/M phase and subsequent apoptosis (Nishio, et al. 1996). Another kinase which is involved in the G2 and mitosis phase is polo-like
30 kinase 1 (Plk1), which is responsible for the maturation of the centrosomes, for the activation of the phosphatase Cdc25C, as well as for the activation of the anaphase promoting complex (Glover et al., 1998, Qian, et al., 2001). The

injection of Plk1 antibodies leads to a G2 arrest in untransformed cells whereas tumour cells arrest in the mitosis phase (Lane and Nigg, 1996). In addition, the protein kinase aurora B has been described as having an essential function during entry into mitosis. Aurora B phosphorylates histone H3 at Ser11 and thereby initiates chromosome condensation (Hsu, J.Y. et al., 2000). A specific cell cycle arrest in the G2/M phase may, however, also be triggered e.g. by the inhibition of specific phosphatases such as e.g. Cdc25C (Russell and Nurse, 1986). Yeasts with a defective cdc25 gene arrest in the G2 phase, while overexpression of cdc25 leads to early entry into the mitosis phase (Russell and Nurse, 1987). However, an arrest in the G2/M phase can also be triggered by the inhibition of certain motor proteins, so-called kinesins such as e.g. Eg5 (Mayer et al., 1999), or by agents which stabilise or destabilise microtubules (e.g. colchicin, taxol, etoposide, vinblastin, vincristin) (Schiff and Horwitz, 1980).

In view of their biological properties the compounds of general formula I according to the invention, their isomers and their physiologically acceptable salts are suitable for the treatment of diseases characterised by excessive or abnormal cell proliferation.

Such diseases include, for example: viral infections (e.g. HIV and Kaposi's sarcoma); inflammatory and autoimmune diseases (e.g. colitis, arthritis, Alzheimer's disease, glomerulonephritis and wound healing); bacterial, fungal and/or parasitic infections; leukaemias, lymphoma and solid tumours; skin diseases (e.g. psoriasis); bone diseases; cardiovascular diseases (e.g. restenosis and hypertrophy). They are also suitable for protecting proliferating cells (e.g. hair, intestinal, blood and progenitor cells) from damage to their DNA caused by radiation, UV treatment and/or cytostatic treatment (Davis et al., 2001).

The new compounds may be used for the prevention, short-term or long-term treatment of the abovementioned diseases, also in combination with other active substances used for the same indications, e.g. cytostatics.

The activity of the compounds according to the invention was determined in the cytotoxicity test on cultivated human tumour cells and/or in a FACS analysis, for example on HeLaS3 cells. In both test methods, the compounds exhibited a good to very good activity, i.e. for example an EC_{50} value in the HeLaS3 cytotoxicity test of less than 5 μmol , generally less than 1 μmol .

Measurement of cytotoxicity on cultivated human tumour cells

To measure the cytotoxicity on cultivated human tumour cells, cells of the cervical cancer tumour cell line HeLaS3 (obtained from American Type Culture Collection (ATCC)) in Ham's F12 Medium (Life Technologies) and 10% foetal calf serum (Life Technologies) were cultivated and harvested in the logarithmic growth phase. Then the HeLaS3 cells were placed in 96-well plates (Costar) at a density of 1000 cells per well and incubated overnight in an incubator (at 37°C and 5 % CO_2), while on each plate 6 wells were filled only with medium (3 wells as a control of the medium, 3 wells for incubation with reduced AlamarBlue). The active substances were added to the cells in various concentrations (dissolved in DMSO; final concentration: 1%) (in each case as a triple measurement). After 72 hours' incubation, 20 μl of AlamarBlue (AccuMed International) were added to each well, and the cells were incubated for a further 7 hours. As a control, 20 μl of reduced Alamar Blue (AlamarBlue reagent which had been autoclaved for 30 min) were added to 3 wells. After 7 h incubation the colour change of the AlamarBlue reagent in the individual wells was determined in a Perkin Elmer fluorescence spectrophotometer (excitation 530 nm, emission 590 nm, slits 15, integrate time 0.1). The amount of AlamarBlue reagent reacted represents the metabolic activity of the cells. The relative cell activity was calculated as a percentage of the control (HeLa S3 cells without inhibitor) and the active substance concentration which inhibits the cell activity by 50% (IC_{50}) was obtained. The values were calculated from the average of three individual measurements, correcting for the control value (medium control).

FACS Analysis

Propidium iodide (PI) binds stoichiometrically to double-stranded DNA, and is thus suitable for determining the percentage of cells in the G1, S and G2/M phase of the cell cycle on the basis of the cell DNA content. Cells in the G0 and G1 phase have a diploid DNA content (2N), whereas cells in G2 or mitosis have a 4N DNA content.

For PI staining, 0.4 million HeLaS3 cells were seeded, for example, on a 75 cm² cell culture flask, and after 24 h either 1 % DMSO was added as control or the substance was added in various concentrations (in 1% DMSO). The cells were incubated for 24 h with the substance or with DMSO, before the cells were washed with 2 x PBS and detached with trypsin /EDTA. The cells were centrifuged (1000 rpm, 5 min, 4°C), and the cell pellet was washed 2 x with PBS, before the cells were resuspended in 0.1 ml of PBS. Then the cells were fixed with 80% ethanol for 16 hours at 4°C or alternatively for 2 hours at -20°C. The fixed cells (10⁶ cells) were centrifuged (1000 rpm, 5 min, 4°C), washed with PBS and then centrifuged again. The cell pellet was resuspended in 2 ml of Triton X-100 in 0.25 % PBS, and incubated for 5 min on ice, before 5 ml of PBS were added and the mixture was centrifuged again. The cell pellet was resuspended in 350 µl of PI stain solution (0.1 mg/ml of Raze A, 10 µg/ml of propidium iodide in 1 x PBS). The cells were incubated for 20 min in the dark with the stain buffer before being transferred into sample measuring vessels for the FACS scan. The DNA measurement was carried out in a Becton Dickinson FACS Analyzer, with an argon laser (500 mW, emission 488 nm), and the DNA Cell Quest Program (BD). The logarithmic PI fluorescence was determined with a band-pass filter (BP 585/42). The cell populations in the individual phases of the cell cycle were quantified with the ModFit LT program of Becton Dickinson.

The compounds of general formula (I) may be used on their own or combined with other active substances according to the invention, optionally also in conjunction with other pharmacologically active substances. Suitable preparations include for example tablets, capsules, suppositories, solutions,

particularly solutions for injection (s.c., i.v., i.m.) and infusion, syrups, emulsions or dispersible powders. The amount of pharmaceutically active compound in each case should be in the range from 0.1 - 90 wt.%, preferably 0.5 - 50 wt.% of the total composition, i.e. in amounts which are sufficient to
5 achieve the dosage range given below. The doses specified may, if necessary, be given several times a day.

Suitable tablets may be obtained, for example, by mixing the active substance(s) with known excipients, for example inert diluents such as
10 calcium carbonate, calcium phosphate or lactose, disintegrants such as corn starch or alginic acid, binders such as starch or gelatine, lubricants such as magnesium stearate or talc and/or agents for delaying release, such as carboxymethyl cellulose, cellulose acetate phthalate, or polyvinyl acetate. The tablets may also comprise several layers.

15 Coated tablets may be prepared accordingly by coating cores produced analogously to the tablets with substances normally used for tablet coatings, for example collidone or shellac, gum arabic, talc, titanium dioxide or sugar. To achieve delayed release or prevent incompatibilities the core may also
20 consist of a number of layers. Similarly the tablet coating may consist of a number of layers to achieve delayed release, possibly using the excipients mentioned above for the tablets.

Syrups or elixirs containing the active substances or combinations thereof
25 according to the invention may additionally contain a sweetener such as saccharin, cyclamate, glycerol or sugar and a flavour enhancer, e.g. a flavouring such as vanillin or orange extract. They may also contain suspension adjuvants or thickeners such as sodium carboxymethyl cellulose, wetting agents such as, for example, condensation products of fatty alcohols
30 with ethylene oxide, or preservatives such as p-hydroxybenzoates.

Solutions for injection and infusion are prepared in the usual way, e.g. with the addition of preservatives such as p-hydroxybenzoates, or stabilisers such as alkali metal salts of ethylenediamine tetraacetic acid, optionally using emulsifiers and/or dispersants, while if water is used as the diluent organic solvents may optionally be used as solubilisers or auxiliary solvents, and transferred into injection vials or ampoules or infusion bottles.

Capsules containing one or more active substances or combinations of active substances may for example be prepared by mixing the active substances with inert carriers such as lactose or sorbitol and packing them into gelatine capsules.

Suitable suppositories may be made for example by mixing with carriers provided for this purpose, such as neutral fats or polyethyleneglycol or the derivatives thereof.

Suitable excipients may be, for example, water, pharmaceutically acceptable organic solvents, such as paraffins (e.g. petroleum fractions), oils of vegetable origin (e.g. groundnut or sesame oil), mono- or polyfunctional alcohols (e.g. ethanol or glycerol), carriers such as e.g. natural mineral powders (e.g. kaolin, clays, talc, chalk), synthetic mineral powders (e.g. highly dispersed silica and silicates), sugar (e.g. glucose, lactose and dextrose), emulsifiers (e.g. lignin, spent sulphite liquors, methylcellulose, starch and polyvinylpyrrolidone) and lubricants (e.g. magnesium stearate, talc, stearic acid and sodium lauryl sulphate).

The preparations are administered in the usual way, preferably by oral or transdermal route, particularly preferably by oral route. When administered orally the tablets may, of course, contain additives, such as e.g. sodium citrate, calcium carbonate and dicalcium phosphate together with various additives, such as starch, preferably potato starch, gelatine and the like, in addition to the abovementioned carriers. Lubricants such as magnesium

stearate, sodium laurylsulphate and talc may also be used to form tablets. In the case of aqueous suspensions the active substances may be combined with various flavour enhancers or colourings in addition to the abovementioned excipients.

- 5 For parenteral use, solutions of the active substances may be prepared using suitable liquid carrier materials.

The dosage for intravenous use is 1 - 1000 mg per hour, preferably between 5 - 500 mg per hour.

- 10 However, it may optionally be necessary to deviate from the amounts specified, depending on the body weight or method of administration, the individual response to the medication, the nature of the formulation used and the time or interval over which it is administered. Thus, in some cases, it may be sufficient to use less than the minimum quantity specified above, while in
15 other cases the upper limit specified will have to be exceeded. When large amounts are administered it may be advisable to spread them over the day in a number of single doses.

- The formulation examples that follow illustrate the present invention without
20 restricting its scope:

Examples of pharmaceutical formulations

25	A)	<u>Tablets</u>	<u>per tablet</u>
		active substance	100 mg
		lactose	140 mg
		corn starch	240 mg
		polyvinylpyrrolidone	15 mg
30		magnesium stearate	5 mg
			500 mg

- 5 The finely ground active substance, lactose and some of the corn starch are mixed together. The mixture is screened, then moistened with a solution of polyvinylpyrrolidone in water, kneaded, wet-granulated and dried. The granules, the remaining corn starch and the magnesium stearate are screened and mixed together. The mixture is compressed to produce tablets of suitable shape and size.

B) <u>Tablets</u>		<u>per tablet</u>
10	active substance	80 mg
	lactose	55 mg
	corn starch	190 mg
	microcrystalline cellulose	35 mg
	polyvinylpyrrolidone	15 mg
15	sodium-carboxymethyl starch	23 mg
	magnesium stearate	2 mg
		400 mg

- 20 The finely ground active substance, some of the corn starch, lactose, microcrystalline cellulose and polyvinylpyrrolidone are mixed together, the mixture is screened and worked with the remaining corn starch and water to form a granulate which is dried and screened. The sodiumcarboxymethyl starch and the magnesium stearate are added and mixed in and the mixture is
- 25 compressed to form tablets of a suitable size.

C) Ampoule solution

30	active substance	50 mg
	sodium chloride	50 mg
	water for inj.	5 ml

- 35 The active substance is dissolved in water at its own pH or optionally at pH 5.5 to 6.5 and sodium chloride is added to make it isotonic. The solution obtained is filtered free from pyrogens and the filtrate is transferred under


- 92 -

aseptic conditions into ampoules which are then sterilised and sealed by fusion. The ampoules contain 5 mg, 25 mg and 50 mg of active substance.

Patent Claims

1) Compounds of general formula (I),

5



wherein

10

R^1 , R^2 which may be identical or different, denote hydrogen or optionally substituted C_1 - C_6 -alkyl,

or

R^1 and R^2 together denote a 2- to 5-membered alkyl bridge which may contain

15

1 to 2 heteroatoms,

R^3 denotes hydrogen or a group selected from among optionally substituted C_1 - C_{12} -alkyl, C_2 - C_{12} -alkenyl, C_2 - C_{12} -alkynyl and C_6 - C_{14} -aryl, or a group selected from among optionally substituted and/or bridged C_3 - C_{12} -cycloalkyl, C_3 - C_{12} -cycloalkenyl, C_7 - C_{12} -polycycloalkyl, C_7 - C_{12} -polycycloalkenyl, C_5 - C_{12} -spirocycloalkyl, C_3 - C_{12} -heterocycloalkyl which contains 1 to 2 heteroatoms, and C_3 - C_{12} -heterocycloalkenyl which contains 1 to 2 heteroatoms, or

20

R^1 and R^3 or R^2 and R^3 together denote a saturated or unsaturated C_3 - C_4 -alkyl bridge which may contain 1 heteroatom,

- 5 R^4 denotes a group selected from among hydrogen, -CN, hydroxy, $-NR_6R_7$ and halogen, or
a group selected from among optionally substituted C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_1 - C_5 -alkyloxy, C_2 - C_5 -alkenyloxy, C_2 - C_5 -alkynyloxy, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulphoxo and C_1 - C_6 -alkylsulphonyl,

10

L denotes a linker selected from among optionally substituted C_2 - C_{10} -alkyl, C_2 - C_{10} -alkenyl, C_6 - C_{14} -aryl, $-C_2$ - C_4 -alkyl- C_6 - C_{14} -aryl, $-C_6$ - C_{14} -aryl- C_1 - C_4 -alkyl, optionally bridged C_3 - C_{12} -cycloalkyl and heteroaryl which contains 1 or 2 nitrogen atoms,

15

n denotes 0 or 1

m denotes 1 or 2

- 20 R^5 denotes a group selected from among optionally substituted morpholinyl, piperidinyl, piperazinyl, piperazinylcarbonyl, pyrrolidinyl, tropenyl, R^8 -diketomethylpiperazinyl, sulphoxomorpholinyl, sulphonylmorpholinyl, thiomorpholinyl, $-NR^8R^9$ and azacycloheptyl,

R^6 , R^7 which may be identical or different, denote hydrogen or C_1 - C_4 -alkyl, and

- 25 R^8 , R^9 denote unsubstituted nitrogen substituents at R^5 , which may be identical or different, denote either hydrogen or a group selected from among C_1 - C_6 -alkyl, $-C_1$ - C_4 -alkyl- C_3 - C_{10} -cycloalkyl, C_3 - C_{10} -cycloalkyl, C_6 - C_{14} -aryl, $-C_1$ - C_4 -alkyl- C_6 - C_{14} -aryl, pyranyl, pyridinyl, pyrimidinyl, C_1 - C_4 -alkyloxycarbonyl, C_6 - C_{14} -arylcarbonyl, C_1 - C_4 -alkylcarbonyl, C_6 - C_{14} -arylmethyloxycarbonyl, C_6 - C_{14} -arylsulphonyl, C_1 - C_4 -alkylsulphonyl and C_6 - C_{14} -aryl- C_1 - C_4 -alkylsulphonyl,
- 30

optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.

5

2) Compounds of general formula (I), wherein

R^1 to R^4 , R^6 and R^7 are as hereinbefore defined, and

10 L denotes a linker selected from among optionally substituted C_2 - C_{10} -alkyl, C_2 - C_{10} -alkenyl, C_6 - C_{14} -aryl, $-C_2$ - C_4 -alkyl- C_6 - C_{14} -aryl, $-C_6$ - C_{14} -aryl- C_1 - C_4 -alkyl, optionally bridged C_3 - C_{12} -cycloalkyl and heteroaryl which contains 1 or 2 nitrogen atoms

n denotes 1

m denotes 1 or 2

15 R^5 denotes a group which is bound to L via a nitrogen atom, selected from among optionally substituted morpholinyl, piperidinyl, R^8 -piperazinyl, pyrrolidinyl, tropenyl, R^8 -diketomethylpiperazinyl, sulphoxomorpholinyl, sulphonylmorpholinyl, thiomorpholinyl, $-NR^8R^9$ and azacycloheptyl,

20 R^8 , R^9 denote unsubstituted nitrogen substituents at R^5 , which may be identical or different, hydrogen or a group selected from among C_1 - C_6 -alkyl, $-C_1$ - C_4 -alkyl- C_3 - C_{10} -cycloalkyl, C_3 - C_{10} -cycloalkyl, C_6 - C_{14} -aryl, $-C_1$ - C_4 -alkyl- C_6 - C_{14} -aryl, pyranyl, pyridinyl, pyrimidinyl, C_1 - C_4 -alkyloxycarbonyl, C_6 - C_{14} -arylcarbonyl, C_1 - C_4 -alkylcarbonyl, C_6 - C_{14} -arylmethyloxycarbonyl, C_6 - C_{14} -arylsulphonyl, C_1 - C_4 -alkylsulphonyl and C_6 - C_{14} -aryl- C_1 - C_4 -alkylsulphonyl,

25

optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.

30

3) Compounds according to claim 1, wherein

R^1 to R^4 , R^6 and R^7 are as hereinbefore defined,

L denotes a linker selected from among optionally substituted C_2 - C_{10} -alkyl, C_2 - C_{10} -alkenyl, C_6 - C_{14} -aryl, $-C_2$ - C_4 -alkyl- C_6 - C_{14} -aryl, $-C_6$ - C_{14} -aryl- C_1 - C_4 -alkyl, optionally bridged C_3 - C_{12} -cycloalkyl and heteroaryl which contains 1 or 2 nitrogen atoms

n denotes 0 or 1

m denotes 1 or 2

R^5 denotes a group which is bound to L via a carbon atom, selected from among R^8 - piperidinyl, R^8R^9 -piperazinyl, R^8 -pyrrolidinyl, R^8 -piperazinylcarbonyl, R^8 -tropenyl, R^8 -morpholinyl and R^8 -azacycloheptyl, and

R^8 , R^9 denote unsubstituted nitrogen substituents at R^5 , which may be identical or different, hydrogen or a group selected from among C_1 - C_6 -alkyl, $-C_1$ - C_4 -alkyl- C_3 - C_{10} -cycloalkyl, C_3 - C_{10} -cycloalkyl, C_6 - C_{14} -aryl, $-C_1$ - C_4 -alkyl- C_6 - C_{14} -aryl, pyranyl, pyridinyl, pyrimidinyl, C_1 - C_4 -alkyloxycarbonyl, C_6 - C_{14} -arylcarbonyl, C_1 - C_4 -alkylcarbonyl, C_6 - C_{14} -arylmethyloxycarbonyl, C_6 - C_{14} -arylsulphonyl, C_1 - C_4 -alkylsulphonyl and C_6 - C_{14} -aryl- C_1 - C_4 -alkylsulphonyl,

optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.

25

4) Compounds according to one of claims 1 to 3, wherein

L, m, n and R^3 to R^9 are as hereinbefore defined, and

R^1 , R^2 which may be identical or different, denote a group selected from among hydrogen, Me, Et, Pr, or


30

R^1 and R^2 together form a C_2 - C_4 -alkyl bridge,

optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.


- 5) Compounds according to one of claims 1 to 4, wherein
R¹, R², m, n and R⁵ to R⁸ are as hereinbefore defined, and
R³ denotes a group selected from among optionally substituted C₁-C₁₀-alkyl, C₃-C₇-cycloalkyl, C₃-C₆-heterocycloalkyl and C₆-C₁₄-aryl or
R¹ and R³ or R² and R³ together denote a saturated or unsaturated C₃-C₄-alkyl
bridge which may contain 1 to 2 heteroatoms,
R⁴ denotes a group selected from among hydrogen, OMe, OH, Me, Et, Pr, OEt, NHMe, NH₂, F, Cl, Br, O-propargyl, O-butynyl, CN, SMe, NMe₂, CONH₂, ethynyl, propynyl, butynyl and allyl,
and
L denotes a linker selected from among optionally substituted phenyl, phenylmethyl, cyclohexyl and branched C₁-C₆-alkyl,
optionally in the form of the tautomers, the racemates, the enantiomers, the diastereomers and the mixtures thereof, and optionally the pharmacologically acceptable acid addition salts thereof.
- 6) Compound of formula I according to one of claims 1 to 5 for use as a pharmaceutical composition.
- 7) Compound of formula I according to one of claims 1 to 5 for use as a pharmaceutical composition with an antiproliferative activity.
- 8) Use of a compound of formula I according to one of claims 1 to 5 for preparing a pharmaceutical composition for the treatment and/or prevention of cancer, infections, inflammatory and autoimmune diseases.

- 9) Method of treating and/or preventing cancer, infections, inflammatory and autoimmune diseases, characterised in that a patient is given an effective quantity of a compound of formula I according to claims 1 to 4.
- 5 10) Pharmaceutical preparations, containing as active substance one or more compounds of general formula (I) according to one of claims 1 to 4 or the physiologically acceptable salts thereof, optionally combined with conventional excipients and/or carriers.
- 10 11) Process for preparing a compound of general formula (I),




wherein

- 15 R¹-R⁵, m, n and L have the meanings given in claims 1 to 5, characterised in that a compound of general formula (II)




wherein

R^1 - R^3 have the meanings given in claims 1 to 4 and A is a leaving group,
is reacted with an optionally substituted compound of general formula (III),



5 wherein

R^4 has the meanings given in claims 1 to 5 and
 R^{10} denotes OH, $NH-L-R^5$, -O-methyl, -O-ethyl,
and then optionally the product of general formula (IV)



10

wherein

R^1 to R^4 has the meanings given in claims 1 to 5 and
 R^{10} denotes OH, $-NH-L-R^5$, -O-methyl or -O-ethyl,

15 optionally after previous hydrolysis of the ester group $-COR^{10}$,
is reacted with an amine of general formula (V)




wherein

R^5 has the meanings given in claims 1 to 5.

5

12) Compound of formula (II),



(II)

10 wherein

$\text{R}^1\text{-R}^3$ have the meanings given in claims 1 to 5 and A is a leaving group.