

**Immunotherapy with B*07 restricted peptides and combination of peptides
against cancers and related methods**

The present invention relates to peptides, proteins, nucleic acids and cells for use in immunotherapeutic methods. In particular, the present invention relates to the immunotherapy of cancer. The present invention furthermore relates to tumor-associated T-cell peptide epitopes, alone or in combination with other tumor-associated peptides that can for example serve as active pharmaceutical ingredients of vaccine compositions that stimulate anti-tumor immune responses, or to stimulate T cells ex vivo and transfer into patients. Peptides bound to molecules of the major histocompatibility complex (MHC), or peptides as such, can also be targets of antibodies, soluble T-cell receptors, and other binding molecules.

The present invention relates to several novel peptide sequences and their variants derived from HLA class I molecules of human tumor cells that can be used in vaccine compositions for eliciting anti-tumor immune responses, or as targets for the development of pharmaceutically / immunologically active compounds and cells.

BACKGROUND OF THE INVENTION

According to the World Health Organization (WHO), cancer ranged among the four major non-communicable deadly diseases worldwide in 2012. For the same year, colorectal cancer, breast cancer and respiratory tract cancers were listed within the top 10 causes of death in high income countries.

In 2012, 14.1 million new cancer cases, 32.6 million patients suffering from cancer (within 5 years of diagnosis) and 8.2 million cancer deaths were estimated worldwide (Bray et al., 2013; Ferlay et al., 2013).

- 2 -

Within the groups of brain cancer, leukemia and lung cancer the current invention specifically focuses on glioblastoma (GBM), chronic lymphocytic leukemia (CLL) and acute myeloid leukemia (AML), non-small cell and small cell lung cancer (NSCLC and SCLC), respectively.

GBM is the most common central nervous system malignancy with an age-adjusted incidence rate of 3.19 per 100,000 inhabitants within the United States. GBM has a very poor prognosis with a 1-year survival rate of 35% and a 5-year survival rate lower than 5%. Male gender, older age and ethnicity appear to be risk factors for GBM (Thakkar et al., 2014).

CLL is the most common leukemia in the Western world where it comprises about one third of all leukemia. Incidence rates are similar in the US and Europe, and estimated new cases are about 16,000 per year. CLL is more common in Caucasians than in Africans, rarer in Hispanics and Native Americans and seldom in Asians. In people of Asian origin, CLL incidence rates are 3-fold lower than in Caucasians (Gunawardana et al., 2008). The five-year overall survival for patients with CLL is about 79%.

AML is the second most common type of leukemia diagnosed in both adults and children. Estimated new cases in the United States are about 21,000 per year. The five-year survival rate of people with AML is approximately 25%.

Lung cancer is the most common type of cancer worldwide and the leading cause of death from cancer in many countries. Lung cancer is subdivided into small cell lung cancer and non-small cell lung cancer. NSCLC includes the histological types adenocarcinoma, squamous cell carcinoma and large cell carcinoma and accounts for 85% of all lung cancers in the United States. The incidence of NSCLC is closely correlated with smoking prevalence, including current and former smokers and the five-year survival rate was reported to be 15% (Molina et al., 2008; World Cancer Report, 2014).

Considering the severe side-effects and expense associated with treating cancer, there is a need to identify factors that can be used in the treatment of cancer in general and acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer in particular. There is also a need to identify factors representing biomarkers for cancer in general and acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer in particular, leading to better diagnosis of cancer, assessment of prognosis, and prediction of treatment success.

Immunotherapy of cancer represents an option of specific targeting of cancer cells while minimizing side effects. Cancer immunotherapy makes use of the existence of tumor associated antigens.

The current classification of tumor associated antigens (TAAs) comprises the following major groups:

a) Cancer-testis antigens: The first TAAs ever identified that can be recognized by T cells belong to this class, which was originally called cancer-testis (CT) antigens because of the expression of its members in histologically different human tumors and, among normal tissues, only in spermatocytes/spermatogonia of testis and, occasionally, in placenta. Since the cells of testis do not express class I and II HLA molecules, these

- 4 -

antigens cannot be recognized by T cells in normal tissues and can therefore be considered as immunologically tumor-specific. Well-known examples for CT antigens are the MAGE family members and NY-ESO-1.

b) Differentiation antigens: These TAAs are shared between tumors and the normal tissue from which the tumor arose. Most of the known differentiation antigens are found in melanomas and normal melanocytes. Many of these melanocyte lineage-related proteins are involved in biosynthesis of melanin and are therefore not tumor specific but nevertheless are widely used for cancer immunotherapy. Examples include, but are not limited to, tyrosinase and Melan-A/MART-1 for melanoma or PSA for prostate cancer.

c) Over-expressed TAAs: Genes encoding widely expressed TAAs have been detected in histologically different types of tumors as well as in many normal tissues, generally with lower expression levels. It is possible that many of the epitopes processed and potentially presented by normal tissues are below the threshold level for T-cell recognition, while their over-expression in tumor cells can trigger an anticancer response by breaking previously established tolerance. Prominent examples for this class of TAAs are Her-2/neu, survivin, telomerase, or WT1.

d) Tumor-specific antigens: These unique TAAs arise from mutations of normal genes (such as β -catenin, CDK4, etc.). Some of these molecular changes are associated with neoplastic transformation and/or progression. Tumor-specific antigens are generally able to induce strong immune responses without bearing the risk for autoimmune reactions against normal tissues. On the other hand, these TAAs are in most cases only relevant to the exact tumor on which they were identified and are usually not shared between many individual tumors. Tumor-specificity (or -association) of a peptide may also arise if the peptide originates from a tumor- (-associated) exon in case of proteins with tumor-specific (-associated) isoforms.

e) TAAs arising from abnormal post-translational modifications: Such TAAs may arise from proteins which are neither specific nor overexpressed in tumors but nevertheless become tumor associated by posttranslational processes primarily active in tumors. Examples for this class arise from altered glycosylation patterns leading to novel epitopes in tumors as for MUC1 or events like protein splicing during degradation which may or may not be tumor specific.

f) Oncoviral proteins: These TAAs are viral proteins that may play a critical role in the oncogenic process and, because they are foreign (not of human origin), they can evoke a T-cell response. Examples of such proteins are the human papilloma type 16 virus proteins, E6 and E7, which are expressed in cervical carcinoma.

T-cell based immunotherapy targets peptide epitopes derived from tumor-associated or tumor-specific proteins, which are presented by molecules of the major histocompatibility complex (MHC). The antigens that are recognized by the tumor specific T lymphocytes, that is, the epitopes thereof, can be molecules derived from all protein classes, such as enzymes, receptors, transcription factors, etc. which are expressed and, as compared to unaltered cells of the same origin, usually up-regulated in cells of the respective tumor.

There are two classes of MHC-molecules, MHC class I and MHC class II. MHC class I molecules are composed of an alpha heavy chain and beta-2-microglobulin, MHC class II molecules of an alpha and a beta chain. Their three-dimensional conformation results in a binding groove, which is used for non-covalent interaction with peptides.

MHC class I molecules can be found on most nucleated cells. They present peptides that result from proteolytic cleavage of predominantly endogenous proteins, defective ribosomal products (DRIPs) and larger peptides. However, peptides derived from endosomal compartments or exogenous sources are also frequently found on MHC class I molecules. This non-classical way of class I presentation is referred to as cross-presentation in the literature (Brossart and Bevan, 1997; Rock et al., 1990). MHC class II molecules can be found predominantly on professional antigen presenting cells (APCs), and primarily present peptides of exogenous or transmembrane proteins that are taken up by APCs e.g. during endocytosis and are subsequently processed.

Complexes of peptide and MHC class I are recognized by CD8-positive T cells bearing the appropriate T-cell receptor (TCR), whereas complexes of peptide and MHC class II molecules are recognized by CD4-positive-helper-T cells bearing the appropriate TCR. It is well known that the TCR, the peptide and the MHC are thereby present in a stoichiometric amount of 1:1:1.

CD4-positive helper T cells play a key role in inducing and sustaining effective responses by CD8-positive cytotoxic T cells. The identification of CD4-positive T-cell epitopes derived from tumor associated antigens (TAA) is of great importance for the development of pharmaceutical products for triggering anti-tumor immune responses (Gnjatic et al., 2003). At the tumor site, T helper cells, support a cytotoxic T cell- (CTL-) friendly cytokine milieu (Mortara et al., 2006) and attract effector cells, e.g. CTLs, natural killer (NK) cells, macrophages, and granulocytes (Hwang et al., 2007).

In the absence of inflammation, expression of MHC class II molecules is mainly restricted to cells of the immune system, especially professional antigen-presenting cells (APC), e.g., monocytes, monocyte-derived cells, macrophages, dendritic cells. In cancer patients, cells of the tumor have been found to express MHC class II molecules (Dengjel et al., 2006).

Longer (elongated) peptides of the invention can act as MHC class II active epitopes.

T-helper cells, activated by MHC class II epitopes, play a key role in orchestrating the effector function of CTLs in anti-tumor immunity. T-helper cell epitopes that trigger a T-helper cell response of the TH1 type support effector functions of CD8-positive killer T cells, which include cytotoxic functions directed against tumor cells displaying tumor-associated peptide/MHC complexes on their cell surfaces. In this way tumor-associated T-helper cell peptide epitopes, alone or in combination with other tumor-associated peptides, can serve as active pharmaceutical ingredients of vaccine compositions that stimulate anti-tumor immune responses.

It was shown in mammalian animal models, e.g., mice, that even in the absence of CD8-positive T lymphocytes, CD4-positive T cells are sufficient for inhibiting manifestation of tumors via inhibition of angiogenesis by secretion of interferon-gamma (IFN γ) (Beatty and Paterson, 2001; Mumberg et al., 1999). There is evidence for CD4 T cells as direct anti-tumor effectors (Braumuller et al., 2013; Tran et al., 2014).

Since the constitutive expression of HLA class II molecules is usually limited to immune cells, the possibility of isolating class II peptides directly from primary tumors was previously not considered possible. However, Dengjel et al. were successful in identifying a number of MHC Class II epitopes directly from tumors (WO 2007/028574, EP 1 760 088 B1).

Since both types of response, CD8 and CD4 dependent, contribute jointly and synergistically to the anti-tumor effect, the identification and characterization of tumor-associated antigens recognized by either CD8+ T cells (ligand: MHC class I molecule + peptide epitope) or by CD4-positive T-helper cells (ligand: MHC class II molecule + peptide epitope) is important in the development of tumor vaccines.

For an MHC class I peptide to trigger (elicit) a cellular immune response, it also must bind to an MHC-molecule. This process is dependent on the allele of the MHC-molecule and specific polymorphisms of the amino acid sequence of the peptide. MHC-class-I-binding peptides are usually 8-12 amino acid residues in length and usually contain two conserved residues ("anchors") in their sequence that interact with the corresponding binding groove of the MHC-molecule. In this way each MHC allele has a "binding motif" determining which peptides can bind specifically to the binding groove.

In the MHC class I dependent immune reaction, peptides not only have to be able to bind to certain MHC class I molecules expressed by tumor cells, they subsequently also have to be recognized by T cells bearing specific T cell receptors (TCR).

For proteins to be recognized by T-lymphocytes as tumor-specific or -associated antigens, and to be used in a therapy, particular prerequisites must be fulfilled. The antigen should be expressed mainly by tumor cells and not, or in comparably small amounts, by normal healthy tissues. In a preferred embodiment, the peptide should be over-presented by tumor cells as compared to normal healthy tissues. It is furthermore desirable that the respective antigen is not only present in a type of tumor, but also in

high concentrations (i.e. copy numbers of the respective peptide per cell). Tumor-specific and tumor-associated antigens are often derived from proteins directly involved in transformation of a normal cell to a tumor cell due to their function, e.g. in cell cycle control or suppression of apoptosis. Additionally, downstream targets of the proteins directly causative for a transformation may be up-regulated and thus may be indirectly tumor-associated. Such indirect tumor-associated antigens may also be targets of a vaccination approach (Singh-Jasuja et al., 2004). It is essential that epitopes are present in the amino acid sequence of the antigen, in order to ensure that such a peptide ("immunogenic peptide") that is derived from a tumor associated antigen leads to an *in vitro* or *in vivo* T-cell-response.

Basically, any peptide able to bind an MHC molecule may function as a T-cell epitope. A prerequisite for the induction of an *in vitro* or *in vivo* T-cell-response is the presence of a T cell having a corresponding TCR and the absence of immunological tolerance for this particular epitope.

Therefore, TAAs are a starting point for the development of a T cell-based therapy including but not limited to tumor vaccines. The methods for identifying and characterizing the TAAs are usually based on the use of T-cells that can be isolated from patients or healthy subjects, or they are based on the generation of differential transcription profiles or differential peptide expression patterns between tumors and normal tissues. However, the identification of genes over-expressed in tumor tissues or human tumor cell lines, or selectively expressed in such tissues or cell lines, does not provide precise information as to the use of the antigens being transcribed from these genes in an immune therapy. This is because only an individual subpopulation of epitopes of these antigens are suitable for such an application since a T cell with a corresponding TCR has to be present and the immunological tolerance for this particular epitope needs to be absent or minimal. In a very preferred embodiment of the invention it is therefore important to select only those over- or selectively presented peptides against which a functional and/or a proliferating T cell can be found. Such a

functional T cell is defined as a T cell, which upon stimulation with a specific antigen can be clonally expanded and is able to execute effector functions ("effector T cell").

In case of targeting peptide-MHC by specific TCRs (e.g. soluble TCRs) and antibodies or other binding molecules (scaffolds) according to the invention, the immunogenicity of the underlying peptides is secondary. In these cases, the presentation is the determining factor.

In a first aspect of the present invention, the present invention relates to a peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499 or a variant sequence thereof which is at least 77%, preferably at least 88%, homologous (preferably at least 77% or at least 88% identical) to SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499, or a variant sequence comprising one conservative amino acid sequence exchange compared with to SEQ ID NO: 1 through SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499, wherein said variants bind to MHC and/or induces T cells cross-reacting with said peptide, or a pharmaceutical acceptable salt thereof, wherein said peptide is not the underlying full-length polypeptide.

The present invention further relates to a peptide of the present invention comprising a sequence that is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499 or a variant thereof, which is at least 77%, preferably at least 88%, homologous (preferably at least 77% or at least 88% identical) to SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499, wherein said peptide or variant thereof has an overall length of between 8 and 100, preferably between 8 and 30, and most preferred of between 8 and 14 amino acids.

The following tables show the peptides according to the present invention, their respective SEQ ID NOs, and the prospective source (underlying) genes for these peptides. In Table 1a, peptides with SEQ ID NO: 1 to SEQ ID NO: 357 bind to HLA-B*07. In Table 1b, peptides with SEQ ID NO: 448 to SEQ ID NO: 497 bind to HLA-B*07.

The peptides in Table 2 have been disclosed before in large listings as results of high-throughput screenings with high error rates or calculated using algorithms but have not been associated with cancer at all before. In Table 2a, peptides with SEQ ID NO: 358 to SEQ ID NO: 383 bind to HLA-B*07. In Table 2b, peptides with SEQ ID NO: 498 to SEQ ID NO 499: bind to HLA-B*07. The peptides in Table 3 are additional peptides that may be useful in combination with the other peptides of the invention. In Table 3, peptides with SEQ ID NO: 384 to SEQ ID NO: 445 bind to HLA-B*07.

Table 1a: Peptides according to the present invention.

Seq ID No	Sequence	Gene(s)	HLA allotype
1	RPRSLQCVSL	KLK2	B*07
2	VPGSDPARYEFL	MAGEA10	B*07
3	MPYVVLTA	SLC6A3	B*07
4	GPKKFIVKL	TGM4	B*07
5	LPSLSHCSQL	PRAME	B*07
6	LPLNSSTSL	MMP12	B*07
7	RPSQLAPATL	LOC100507003	B*07
8	MPKTNLSKM	LOC100507003	B*07
9	RPDSRLLEL	CTAG2	B*07
10	SPMEAELVRRIL	CTAG2	B*07
11	APLPRPGAV	CTAG2	B*07
12	LPNTGRIGQLL	TGM4	B*07
13	LPNTGRIGQL	TGM4	B*07
14	AVHEIGHSL	MMP12	B*07
15	KPGFNISIL	CYP4Z1, CYP4Z2P	B*07
16	FPAPPAHWF	CYP4Z1, CYP4Z2P	B*07
17	SPAAPLSPASSL	CRH	B*07
18	MALSVLRLAL	SPINK2	B*07
19	WPRLPGAGL	FAM178B	B*07
20	MVLGIGPVL	SLC45A3	B*07
21	APSRLEL	LOC645382, LOC645399, PRAMEF10	B*07
22	LPQLKPAL	AMTN	B*07
23	VPRPTSTVGL	TRIM51, TRIM51BP, TRIM51EP, TRIM51HP	B*07
24	VPRPTSTVGLFL	TRIM51, TRIM51BP, TRIM51EP, TRIM51HP	B*07
25	RPQGAVGGL	RHOXF2, RHOXF2B	B*07
26	SPSFSSTLLSL	MAGEC1	B*07
27	RIRVTSEVL	ABCC11	B*07

Seq ID No	Sequence	Gene(s)	HLA allotype
28	LPAPTSVL	COL20A1	B*07
29	APLRVHITSL	PAEP	B*07
30	YPGFTKRL	ACTL8	B*07
31	RPGSPDPA	KISS1R	B*07
32	APMAPGRSPL	PGR	B*07
33	LPRGLSPARQL	PGR	B*07
34	APMAPGRSP	PGR	B*07
35	APLPPPRAL	COL20A1	B*07
36	RPFSREMDL	FLT3	B*07
37	LPPPRALTL	COL20A1	B*07
38	RPSFPNLTSF	FLT3	B*07
39	QPRPSSEKI	POU4F1, POU4F2, POU4F3	B*07
40	FPRTVKHIDAAL	MMP1	B*07
41	MPAGGGPRSL	CKAP4	B*07
42	APLKMLAL	CCL17	B*07
43	APTPRPKVL	BMPR1B	B*07
44	SPSQTVQRAV	IGHE	B*07
45	TRPWSGPYIL	DCT	B*07
46	QPISGNPVT	FCRL5	B*07
47	RPRQTGALM	DNTT	B*07
48	RPRYSIGL	BMPR1B	B*07
49	APEKARAFL	SOX14	B*07
50	PEKARAFL	SOX14	B*07
51	SPVFYVQTL	ABCC11	B*07
52	KEDNPSGHTYTL	MAGEB1	B*07
53	SPRIPFSTF	ROPN1	B*07
54	VPSCGRSVEGL	PTHLH	B*07
55	LPALLRSGL	RAPGEF5	B*07
56	LPALLRSGLTL	RAPGEF5	B*07
57	APLLPIRTL	UPK2	B*07
58	APLLPIRTLPL	UPK2	B*07
59	KPRTIYSSL	DLX1, DLX4, DLX6	B*07
60	RPYSIYPHGVTF	F5	B*07
61	LPRIPFSTF	ROPN1B	B*07
62	KPQSTISGL	F5	B*07
63	FPHMATTAF	HEPHL1	B*07
64	VPRPIFSQLYL	GREB1, GREB1L	B*07
65	FPNVYSTLV	SLC45A2	B*07
66	LPMTVISVL	NMUR2	B*07
67	VPVSRPVL	FCRL2, FCRL3, FCRL5	B*07
68	FPNEVSVVL	HMCN1	B*07
69	RPEDGRPRL	BTBD17	B*07

Seq ID No	Sequence	Gene(s)	HLA allotype
70	VPAQRLGLL	IGKV1D-43	B*07
71	APFAPLHGGGSL	NEFH	B*07
72	APCNGNVAV	KRT81	B*07
73	LPVSSPVTL	AKNAD1	B*07
74	VPVSHPVL	FCRL3, FCRL5	B*07
75	HVRPQTNCI	DCC	B*07
76	KPKVESQAL	ABCC11	B*07
77	QPRLPVGETL	FMN1, LOC101059984	B*07
78	HPSQESPTL	DLX5	B*07
79	GPASDHPSL	GREB1	B*07
80	SALPTTISF	MAGEA4	B*07/B*35
81	IAYPSLREAAL	MAGEA4	B*07/C*03
82	DAAHPGPSV	ALPPL2	B*07/B*51
83	AVSSHNNVI	POTEG	B*07
84	MPMQDIKMI	PRAME	B*07/B*51
85	MPMQDIKMILKM	PRAME	B*07/B*35
86	ALLLRGVTL	SLC6A3	B*07/C*06
87	APVGGNVTSSF	CT45A1, CT45A2, CT45A3, CT45A4, CT45A6, LOC101060208, LOC101060210, LOC101060211	B*07/B*35
88	KPSAVKDSIY	MMP26	B*07/A*26
89	FLIPRAGWLAGL	SLC45A3	B*07/C*06
90	HAIEKAFQL	MMP1	B*07/C*04
91	FPRLVGPdff	MMP11	B*07/B*35
92	TSPLPPTV	ENPP3	B*07
93	SVAIAHGVF	SLC35D3	B*07/B*15
94	LPMSKRQEY	DSCR6	B*07/B*35
95	RTKEEINEL	KRT121P, KRT81, KRT83, KRT85, KRT86	B*07/B*57
96	QPSLVQAIF	ANKRD28	B*07/B*35
97	LPPGTQIQI	ROBO3	B*07
98	FPCSALLACF	FAM107B	B*07/B*35
99	MPVSAFTVI	COL10A1	B*07/B*51
100	TPIPFDKILY	COL10A1	B*07/B*35
101	KPGLPGLK	COL10A1	B*07/C*07
102	MAGPAIHTAPM	DRAXIN	B*07
103	REPIMKADML	MAGEB1	B*07
104	RPLPNSVIHV	TFDP3	B*07
105	RPRYETGVCA	WISP3	B*07
106	APFHIDRLF	NMUR2	B*07
107	GQRLESPGGAA	SOX1	B*07
108	APRGSPPI	FCRL5	B*07
109	LPRALMRST	IGHE	B*07

Seq ID No	Sequence	Gene(s)	HLA allotype
110	YPSSPASAV	MAGEB17	B*07/B*51
111	YPLQQTNVY	SEMA5B	B*07/B*35
112	YPSPLNKHSF	SEMA5B	B*07/B*35
113	KPHLDRRGAVI	TSPY1, TSPY10, TSPY2, TSPY3, TSPY4, TSPY8	B*07
114	FVPPSGPSNPM	PAX3	B*07
115	KTKSLAQQL	LAMC2	B*07/A*30
116	RSYQHSLRL	LAMC2	B*07/C*07
117	IPHQRSSL	GCSAML	B*07
118	NPERHKPSF	C1orf186	B*07
119	KATEYVHSL	MYCN	B*07/C*16
120	TVRPKNAAL	MYCN	B*07
121	GPFQRAAL	MSX1	B*07
122	KPRTPFTTAQLL	MSX1	B*07
123	RPRLRDLPAL	RAPGEF5	B*07
124	KTIDGHINL	FAM111B	B*07/C*04
125	SPAKQFNIY	FAM111B	B*07/B*35
126	MPREDAHF	MLANA	B*07/B*35
127	KSKQVITLL	ENTHD1	B*07/C*02
128	SPPATLFLFLL	LOXL4	B*07
129	MTLPATTLTL	IGHE	B*07
130	GAYDKARYL	ABCC11	B*07/C*06
131	MPFRNIRSIL	LOC101060288, LOC101060295, LOC101060308, LOC645359, PRAMEF11, PRAMEF15, PRAMEF23, PRAMEF4, PRAMEF5, PRAMEF6, PRAMEF9	B*07
132	LPRLPVGSLV	EPYC	B*07/B*51
133	LPELPTTLTF	EPYC	B*07
134	VAAAARLTL	ITIH6	B*07
135	MPRLRLSL	INSL6	B*07
136	QPDHAGIFRVF	HEPHL1	B*07
137	SPQLSYFEY	LAMC2	B*07/B*35
138	VPFNLITEY	SLC45A2	B*07/B*35
139	LVILQHQMAM	NMUR2	B*07/A*26
140	TPFPDIHWF	HMCN1	B*07/B*35
141	YGPYGSWSGW	HRNR	B*07
142	HPFKMKWQY	QRFPR	B*07/B*35
143	YPMIPPAQL	BTBD17	B*07
144	RPVQGIPTY	HMCN1	B*07/B*35
145	VPAQRLGLLL	IGKV1D-43	B*07
146	SPTRGSEF	PTPRZ1	B*07
147	HVAQPAVVL	CTLA4	B*07/C*07

Seq ID No	Sequence	Gene(s)	HLA allotype
148	KPHLIYRQ	ADAMTS20	B*07
149	SPSSVFTSK	NAT1	B*07
150	APLHGGGSL	NEFH	B*07
151	QPTWNTQTRL	CLNK	B*07
152	SSASFSSSELFRLH	PTPRZ1	B*07
153	QPVHSLCSA	DPCR1	B*07/B*54
154	RPPPSRRRAVL	BCAN	B*07
155	APIPRLIEG	CCDC38	B*07
156	GSRPVTGSV	KRT121P, KRT81, KRT83	B*07
157	RPVTGSVC	KRT121P, KRT81, KRT83	B*07
158	SVPVATVEL	ELL3	B*07/C*01
159	TPMTVPRI	KBTBD8	B*07/B*51
160	IPVSHPVLTFL	FCRL3	B*07
161	ASKILETSL	HMCN1	B*07/C*03
162	IPIRVDQNGAF	ADAMTS6	B*07/B*35
163	FRYPNGVSL	RHBG	B*07/C*07
164	RAAGRRLQL	RHBG	B*07
165	RPSKEMQVTI	ONECUT3	B*07
166	YAYTSRVIL	KBTBD8	B*07/C*12
167	NVNPARKGL	DNMT3B	B*07
168	SPSGQRDVSL	ENTHD1	B*07
169	RPFSVSSISQL	HMCN1	B*07
170	APEGKRLGF	FMN1, LOC101059984	B*07
171	LPLGGHKSL	FMN1, LOC101059984	B*07
172	SAQSLHPVL	POU3F4	B*07/C*12
173	VIINNPISL	HMCN1	B*07/C*03
174	IPVTSSVTL	KRTAP24-1	B*07
175	AAFPHKIIF	SPINK13	B*07/C*12/C*02
176	QPLDICHSF	ETV4	B*07/B*35
177	VWEPQGSSRSL	DLX5	B*07
178	VPYYPRANL	STK31	B*07
179	PRVRLLLL	PTH2	B*07
180	RLRDYISSL	RALGPS2	B*07/C*07
181	LPVSPARAL	ADAM12	B*07
182	RPLPVSPARAL	ADAM12	B*07
183	VPRRPARAL	BCAN	B*07
184	KIIASGNHL	ERVV-1, ERVV-2	B*07
185	RPVLTASSF	FCRL2	B*07
186	VPLPAGGGTVLT	GRP	B*07
187	APPPPPPPF	BEND4	B*07
188	HAAASFETL	FCRLA	B*07/B*35
189	QPQCSTHIL	APOB	B*07

Seq ID No	Sequence	Gene(s)	HLA allotype
190	RLAHVRARL	GREB1	B*07/C*07
191	KPKAVKPKAA	HIST1H1B	B*07
192	IPFADVKSF	ROS1	B*07
193	LPALKEEAF	NKX3-1	B*07/B*35
194	HPNKKASL	TRPS1	B*07
195	NPIFGRKEL	CAPN6	B*07
196	RPSGTAGAAL	EGFR	B*07
197	RPSGTAGAALLA	EGFR	B*07/B*56
198	LPSPAHAFRAL	SIGLEC15	B*07
199	SPFWIHQA	MPL	B*07
200	EPFHLIVSY	FCRLA	B*07/B*35
201	LPIARVLTV	LRP1B	B*07/B*51
202	SPSREASAL	MPL	B*07
203	KPRGPIDI	COL9A1	B*07
204	FPYSGDKILV	VCAN	B*07/B*51
205	LPPALLTTV	VCAN	B*07/B*51
206	TPRIGPKVSL	VCAN	B*07
207	VPSDITTAF	VCAN	B*07/B*35
208	RNRQVATAL	SUCNR1	B*07
209	KIEQIRAVL	LAMB3	B*07
210	IPENRVVSY	CCL24	B*07/B*35
211	IPDTIASVL	SLC24A5	B*07
212	VPYAAQAAL	SP5	B*07
213	RPYQDAPVA	SIGLEC8	B*07/B*55
214	LPLKFFPII	TMPRSS3	B*07
215	IPVAIKEL	EGFR	B*07
216	LPWEQNEQV	APOB	B*07/B*51
217	SPGDKRLAAYL	APOB	B*07
218	VQRTPPPI	USH1C	B*07
219	VPHTGRYTCL	HMCN1	B*07
220	RPAPGPAPFV	GRIN2D	B*07
221	LPQRPNARL	LAMB3	B*07
222	MLKTTLTAF	APOB	B*07
223	KAHVRIEL	RNF43, SUPT4H1	B*07/C*16
224	SPIIHSILL	PLEKHG4B	B*07
225	SPIIHSIL	PLEKHG4B	B*07
226	APGGSSRSSL	IGLL1	B*07
227	RPGTGQGGL	IGLL1	B*07
228	RPTAASQSRAL	IGLL1	B*07
229	FPNAGPRHLL	RALGPS2	B*07
230	DVIDDIISL	MITF	B*07/A*25
231	SPITLQAL	APOB	B*07

Seq ID No	Sequence	Gene(s)	HLA allotype
232	TAYPSLRLL	ST8SIA2	B*07/C*06
233	MAYDRFIAI	OR10V1, OR4K1, OR51B2, OR51B5, OR9K2	B*07/C*03
234	HPRAPGEQQRL	RNF43, SUPT4H1	B*07
235	AQNPAGTAL	HMCN1	B*07/B*15
236	TPELKSSIL	APOB	B*07
237	LPRAGGAF	LAMB3	B*07
238	LPRAGGAFLM	LAMB3	B*07
239	VLPRAGGAFLM	LAMB3	B*07
240	RVMLPKAAL	NLRP2	B*07
241	LPKAALLV	NLRP2	B*07/B*51
242	IPETASVVAI	FCRLA	B*07
243	MARTGGMVVI	UPK2	B*07
244	VPAHLVAA	ESM1	B*07/B*55
245	GPVPSPLAL	GDF7	B*07
246	RPILKEQSSSSF	KRT13, KRT16	B*07
247	SPVGVGQRL	SOX3	B*07
248	KPYDGIPAS	EGFR	B*07/B*56
249	SPRSGVLL	LGR5	B*07
250	APAAPAAVPS	FEZF1	B*07
251	MPVDSFNSM	NFE2L3	B*07/C*04
252	QPENSLEGISL	NFE2L3	B*07
253	MPVDSFNSML	NFE2L3	B*07
254	RVIQGTTL	PHEX	B*07/A*32
255	VPSHWMVAL	CD79B	B*07
256	SPVPSHWMVAL	CD79B	B*07
257	APYGTLRKS	SYT12	B*07
258	SVIGPNSRL	BTLA	B*07/A*26
259	VPMPGVQAV	TMEM211	B*07
260	LVQSSRSEV	KRT16	B*07
261	SPSTSRTPL	EGFR	B*07
262	SPSTSRTPLLSSL	EGFR	B*07
263	SQRPPATSQA	HTR3A	B*07
264	APRPGNWIL	LAMA1	B*07
265	FPRKPYEGRV	LOXL3	B*07
266	KPYEGRVEI	LOXL3	B*07
267	MPVPGILL	PMEL	B*07
268	EPLSVTASY		B*07/A*25
269	FTVSSSSAM	LOC100131514, LOC101060622, MUC3A	B*07/C*03
270	SPRGTTSTL		B*07
271	SPTPVFTTL	LOC100131514, LOC101060622, LOC101060740, LOC101060797, MUC3A,	B*07

Seq ID No	Sequence	Gene(s)	HLA allotype
		MUC3B	
272	SSPRGTTSTL		B*07
273	TDTPSTTPTTI	MUC3A	B*07
274	KPIRTGISPLAL	APOB	B*07
275	IPAPQGAVL	KISS1	B*07
276	RPVWDVRSAL	RNASE10	B*07/B*55
277	MPPLLIVAF	HCAR1	B*07/B*35
278	LIAARGSLVL	NTN3	B*07
279	APADEPRTVL	SAPCD2	B*07
280	LPRAFLQSL	SAPCD2	B*07
281	NPRSPPATM	SIGLEC15	B*07
282	AVRLPTPNL	MPL	B*07
283	RPQPGWRESL	FMR1NB	B*07
284	RPSAPRGPP	ASCL2	B*07
285	RITWKGLLL	PSG3, PSG8, PSG9	B*07/C*07
286	APARPAAAF	TDRD9	B*07
287	SPIPRPLFL	SYCP2	B*07
288	LHAMNSLSAM	DMBX1	B*07
289	LPYEKVSRL	CHST4	B*07
290	RPTHPLRSF	DCC	B*07
291	SPSKSLEM	L1TD1	B*07
292	LPMTHRLQL	PSG4, PSG5, PSG6, PSG7	B*07
293	FPYDKPLIM	MET	B*07/B*51
294	VPKPAIPSSSVL	DCC	B*07
295	HPRWIEPTVM	HOXA13	B*07
296	SPLLMQRTL	TFEC	B*07/B*51
297	FPIKYVNDF	MET	B*07/B*35
298	RVLLRWISL	ILDR2	B*07
299	SPFSGGPVSF	POU5F1, POU5F1B	B*07
300	HPYSDLADV	RTL1	B*07
301	FPAFLEAM	RTL1	B*07/B*35
302	IPIDQILNSF	RTL1	B*07/B*35
303	RPPPPCIAL	RTL1	B*07
304	SPLIGDFPAF	RTL1	B*07/B*35
305	AASPVGSAL	SLC16A11	B*07/C*03
306	RPFPLALL	CYP2W1	B*07
307	RPHQKGWLSI	KLB	B*07
308	RPDVVRTLL	ARSI	B*07
309	FAFYGGKSL	SLC44A5	B*07/C*03
310	SPGWAQTQL	MACC1	B*07
311	APRLALDPDAL	AMH	B*07
312	SPSLQSSRESL	KIF26B	B*07

Seq ID No	Sequence	Gene(s)	HLA allotype
313	VPLSSYVSI	PCDHGA11	B*07
314	IALMKLAGPL	TMPRSS3	B*07
315	APVVFPAL	GSC	B*07
316	NPPEPEKSL	LOC100124692	B*07
317	MPYNSAHHCVV	KLHL14	B*07/B*51
318	TPISNTGVL	SIGLEC6	B*07
319	RPLDTFRVL	EVX1	B*07
320	APMHIDRNIL	SLC10A5	B*07
321	QPQQPGFFL	MED12L	B*07
322	RAVPVGSGL	TRIT1	B*07/C*03
323	TPHGITDIL	NKX6-3	B*07
324	LPAPLRISL	GPR45	B*07
325	SPRSNPVRL	FCRL4	B*07
326	IPPFTQRFV	RASSF9	B*07/B*35
327	GPRTTQSSVL	SIGLEC6	B*07
328	LPLHRGDLVI	DNMBP	B*07/B*51
329	QPANFIVLL	LOC100124692	B*07
330	RPFSAIAEL	MORC4	B*07
331	SPDSAILKL	ZNF827	B*07
332	SPYAGSTAF	IRX5	B*07
333	SVLPRALSL	C5orf34	B*07/C*07
334	YPLSLALHF	C5orf34	B*07/B*35
335	VPPQNPRPSL	RUNX2	B*07
336	YPLQGPGLLSV	NUP155	B*07/B*51
337	IPTSRVITL	NUGGC	B*07
338	MPATPSLKV	TRIM15	B*07
339	GPQRTTSV	DTX1	B*07
340	APEPRAALL	UAP1L1	B*07
341	LPRSPPLKVL	RMI2	B*07
342	RPRPPKVL	TXNRD1	B*07
343	RPRPPKVLGL	TXNRD1	B*07
344	VPYPSPTCV	AR	B*07
345	SAAPPGASL	AR	B*07/C*03
346	IPMPRITWL	FSTL4	B*07
347	SPEKINSF	BRIP1	B*07
348	HPAPPVTSa	STON2	B*07/B*54
349	QPRDGWPMML	STON2	B*07
350	RPKSTLMNF	STON2	B*07
351	SPYADIIPSA	GLI3	B*07/B*54
352	LPAFSKIGGIL	IQGAP3	B*07
353	KPRATWTL	RTP4	B*07
354	APAKDARASL	TRIO	B*07

Seq ID No	Sequence	Gene(s)	HLA allotype
355	APKTSFIAA	GALNT5	B*07/B*55
356	RPFLRLPSL	UCHL3	B*07
357	LPPHIFAI	MYO10	B*07/B*51

Table 1b: Peptides according to the present invention.

Seq ID No.	Sequence	Gene(s)
448	TVYGEPRKL	MAGEA4
449	RPKTSVNLISL	MMP12
450	SAAARALLP	MMP11
451	VPILPLNHAL	PGR
452	QPVKKNTL	ENPP3
453	APALPGQVTI	MEX3A
454	SPDPAHLESL	MAGEA8
455	FPVQATIDF	DSCR6
456	VPVSHPVLT	FCRL5
457	RPPLSQRHTF	CLNK
458	VIPTHYFVVL	ENPP3,OR2A4
459	SPAPWRPWI	DSCR6
460	APLMPLGKTL	LAMC2
461	MPHLGPGILL	ITIH6
462	MPLLADVRL	ITIH6
463	LPTDLFNSVM	ROPN1,ROPN1B
464	VPFVPRTSV	NFE2L3
465	VPKSLPYVL	GTSF1
466	SPMEAILVSRL	MAP7D2
467	VPRNQDWLGVSRL	PMEL
468	LPSLHVLVL	DCT
469	LPLDTRTSI	CLNK
470	RPNGEVKSEL	GRM8
471	LPLLAGTLLL	ALPI,ALPP,ALPPL2
472	LPMPAITWY	HMCN1
473	LPGEREAAL	FMN1,LOC101059984
474	VPKADLLTL	FMN1,LOC101059984
475	IPLEIQKL	CDRT1,FBXW10
476	EPNPVEEIF	COL11A1
477	NPVPVITWYKDNRL	HMCN1
478	APKFISPASQL	GRM8
479	APHAGGALL	MIXL1

480	VPAAPTKAL	SPOCD1
481	SPARALLALA	ADAM12
482	QPSLKKIIL	INSL4
483	KPAAAGVKKVA	HIST1H1B
484	LPASAEPKGTM	OFCC1
485	LPLKTKVFA	ZPLD1
486	TPTRPLPSA	USH1C
487	IPWFIKTAF	SLC24A5
488	LPLGGLPLLI	BTLA
489	WPNHIMLVL	GREB1,GREB1L
490	APRGPAQGEAA	RUNX1
491	MPEADLKRIF	SLC24A5
492	IPFSVDEI	NFE2L3
493	LPLQQYKLV	FEZF1
494	LPLRAVNLNL	FAM64A
495	SPSYTQASL	TRPS1
496	MPAVSGPGPLF	LMAN1L
497	YVVKPLHPF	ROS1

Table 2a: Additional peptides according to the present invention with no prior known cancer association.

SEQ ID No.	Sequence	Official Gene Symbol(s)
358	APSMRLRKNQL	DCAF4L2
359	SPRRLVELAGQSL	PRAME
360	SPASRSISLL	CD70
361	SPYGSDRLVQL	ZBTB32
362	KPMLPPAAF	MSX1
363	KPRTPFHTTA	MSX1
364	KPRTPFHTTAQL	MSX1
365	RPKHFLQML	NLRP7
366	SPTLRQLDL	NLRP11
367	APQVHIFSL	OXTR
368	NPASRLTAL	BMPR1A, BMPR1B
369	RPYGCVLRAA	CD70
370	AAHEFGHVL	MMP11
371	APRSPGQVTPRGL	C6orf223
372	SPSSASLTL	FAM227B
373	LPKPDLPQLI	DEFB132
374	KPRNMTGLDL	CLEC17A
375	LPRGVLEGL	SAPCD2
376	FPQVGRTAL	CXCR3
377	SPFSKRIKL	BCL11A

378	SPRQPRLDF	E2F7
379	GPRQVLFPL	PCDHGB6, PCDHGB7
380	SPYPGSAAF	IRX2
381	APRPRLLL	TGFBR1
382	RPGPQRTTSV	DTX1
383	KPRATWTLKL	RTP4

Table 2b: Additional peptides according to the present invention with no prior known cancer association.

Seq ID No	Sequence	Gene(s)
498	RPQPQPRPAL	WNT10A
499	QPVLPSPAC	PMEL

Table 3: Peptides of the invention useful for e.g. personalized cancer therapies.

SEQ ID No.	Sequence	Official Gene Symbol(s)
384	APLLARAA	ACPP
385	FPSLREAAL	MAGEA1
386	YPLRGSSI	ALPP, ALPPL2
387	MPMQDIKM	PRAME
388	SPSVSQLSVL	PRAME
389	APLPRPGAVL	CTAG2
390	SPRMSGLLSQT	DLL3
391	APRPASSL	MMP11
392	GPQPWHAAL	MMP11
393	APAAWLRSA	MMP11
394	APAAWLRSAA	MMP11
395	APAAWLRSAAA	MMP11
396	VPDVAQFVL	MMP1
397	GPALGRSFL	CD70
398	SPASRSISL	CD70
399	NPFYPEVEL	MMP1
400	TIASQRLTPL	CD70
401	RPAPADSAL	KISS1R
402	LPSPVDAAF	MMP11
403	RGVPSEIDAAF	MMP11
404	VPSEIDAAF	MMP11
405	LPFDGPGGIL	MMP11
406	LPDGSRVEL	ACTL8
407	FPRLVGPDF	MMP11
408	YPKDIYSSF	MMP1
409	IPASHPVL	FCRL5
410	SPRSWIQVQI	FCRL5
411	IPNWARQDL	NLRP7

SEQ No.	ID	Sequence	Official Gene Symbol(s)
412		HPSAHDVIL	LAMC2
413		SVHKITSTF	LAMC2
414		KPSESIYSAL	FAM111B
415		LPSDSHFKITF	FAM111B
416		RAKNAGVTI	LAMC2
417		VPRPIFSQL	GREB1, GREB1L
418		RPMTPTQIGPSL	TCL1A
419		SPMWHVQQL	QRFPR
420		SPRWLPVSL	BTBD17
421		FPTEVTPHAF	PTPRZ1
422		IPVSHPVL	FCRL3
423		SPRVYWLGL	CLEC17A
424		YPRGNHWAVGH	GRP
425		YPRGNHWAVGHL	GRP
426		RYLPNPSLNAF	CYP1A1
427		VPSSRILQL	THEG
428		SPADAHRL	NLRP2
429		RPRALRDLQL	NLRP7
430		RPRALRDLQLL	NLRP7
431		VPLPAGGGTV	GRP
432		VPLPAGGGTVL	GRP
433		IPEPSAQQ	APOB
434		LPRIPFADV	ROS1
435		MPLSTIREV	CDK6
436		RPMQQARAQL	KLHDC7B
437		FPYPYAERL	GRIN2D
438		LPRAGGAFL	LAMB3
439		NPFPHLITL	ROS1
440		YPRTITPGM	KLK14
441		SPVPSHWMVA	CD79B
442		SPSTSRTPLL	EGFR
443		VPDGVSKVL	APOB
444		RVEEVRALL	CDKN2A
445		RPAATAVISL	SLC7A11

The present invention furthermore generally relates to the peptides according to the present invention for use in the treatment of proliferative diseases, such as, for example, acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma,

non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

Particularly preferred are the peptides – alone or in combination - according to the present invention selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499. More preferred are the peptides – alone or in combination - selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 79 (see Table 1a), and their uses in the immunotherapy of cancer, and preferably acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

Thus, another aspect of the present invention relates to the use of the peptides according to the present invention for the - preferably combined - treatment of a proliferative disease selected from the group of acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

- 24 -

The present invention furthermore relates to peptides according to the present invention that have the ability to bind to a molecule of the human major histocompatibility complex (MHC) class-I or - in an elongated form, such as a length-variant - MHC class -II.

The present invention further relates to the peptides according to the present invention wherein said peptides (each) consist or consist essentially of an amino acid sequence according to SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499.

The present invention further relates to the peptides according to the present invention, wherein said peptide is modified and/or includes non-peptide bonds.

The present invention further relates to the peptides according to the present invention, wherein said peptide is part of a fusion protein, in particular fused to the N-terminal amino acids of the HLA-DR antigen-associated invariant chain (Ii), or fused to (or into the sequence of) an antibody, such as, for example, an antibody that is specific for dendritic cells.

The present invention further relates to a nucleic acid, encoding the peptides according to the present invention. The present invention further relates to the nucleic acid according to the present invention that is DNA, cDNA, PNA, RNA or combinations thereof.

The present invention further relates to an expression vector capable of expressing and/or expressing a nucleic acid according to the present invention.

The present invention further relates to a peptide according to the present invention, a nucleic acid according to the present invention or an expression vector according to the present invention for use in the treatment of diseases and in medicine, in particular in the treatment of cancer.

The present invention further relates to antibodies that are specific against the peptides according to the present invention or complexes of said peptides according to the present invention with MHC, and methods of making these.

The present invention further relates to T-cell receptors (TCRs), in particular soluble TCR (sTCRs) and cloned TCRs engineered into autologous or allogeneic T cells, and methods of making these, as well as NK cells or other cells bearing said TCR or cross-reacting with said TCRs.

The antibodies and TCRs are additional embodiments of the immunotherapeutic use of the peptides according to the invention at hand.

The present invention further relates to a host cell comprising a nucleic acid according to the present invention or an expression vector as described before. The present invention further relates to the host cell according to the present invention that is an antigen presenting cell, and preferably is a dendritic cell.

The present invention further relates to a method for producing a peptide according to the present invention, said method comprising culturing the host cell according to the present invention, and isolating the peptide from said host cell or its culture medium.

The present invention further relates to said method according to the present invention, wherein the antigen is loaded onto class I or II MHC molecules expressed on the surface of a suitable antigen-presenting cell or artificial antigen-presenting cell by contacting a sufficient amount of the antigen with an antigen-presenting cell.

The present invention further relates to the method according to the present invention, wherein the antigen-presenting cell comprises an expression vector capable of expressing or expressing said peptide containing SEQ ID No. 1 to SEQ ID No.: 383 and SEQ ID NO: 448 to SEQ ID NO: 499, preferably containing SEQ ID No. 1 to SEQ ID No. 79, or a variant amino acid sequence.

The present invention further relates to activated T cells, produced by the method according to the present invention, wherein said T cell selectively recognizes a cell which expresses a polypeptide comprising an amino acid sequence according to the present invention.

In an aspect, the activated T cells may be produced by contacting in vitro T cells with antigen loaded human class I or II MHC molecules presented on the surface of a suitable antigen-presenting cell or an artificial construct mimicking an antigen-presenting cell for a period of time sufficient to activate said T cells.

The present invention further relates to a method of killing target cells in a patient, in which the target cells aberrantly express a polypeptide comprising any amino acid sequence according to the present invention, the method comprising administering to the patient an effective number of T cells as produced according to the present invention.

The present invention further relates to a method of treating a patient who has cancer, in which the cancer cells aberrantly express a polypeptide comprising any amino acid sequence according to the present invention, the method comprising administering to the patient an effective number of T cells as produced according to the present invention.

The present invention further relates to a method of eliciting an immune response in a patient who has cancer, in which the cancer cells aberrantly express a polypeptide comprising any amino acid sequence according to the present invention, the method comprising administering to the patient an effective number of T cells as produced according to the present invention.

In one aspect, the immune response may include cytotoxic T cell response.

The present invention further relates to the use of any peptide as described, the nucleic acid according to the present invention, the expression vector according to the present invention, the cell according to the present invention, the activated T lymphocyte, the T cell receptor or the antibody or other peptide- and/or peptide-MHC-binding molecules according to the present invention as a medicament or in the manufacture of a medicament. Preferably, said medicament is active against cancer.

Preferably, said medicament is a cellular therapy, a vaccine or a protein based on a soluble TCR or antibody.

The present invention further relates to a use according to the present invention, wherein said cancer cells are acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer, and preferably acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer cells.

The present invention further relates to biomarkers based on the peptides according to the present invention, herein called "targets" that can be used in the diagnosis of cancer, preferably acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma,

melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer. The marker can be over-presentation of the peptide(s) themselves, or over-expression of the corresponding gene(s). The markers may also be used to predict the probability of success of a treatment, preferably an immunotherapy, and most preferred an immunotherapy targeting the same target that is identified by the biomarker. For example, an antibody or soluble TCR can be used to stain sections of the tumor to detect the presence of a peptide of interest in complex with MHC.

Optionally the antibody carries a further effector function such as an immune stimulating domain or toxin.

The present invention also relates to the use of these novel targets in the context of cancer treatment.

Stimulation of an immune response is dependent upon the presence of antigens recognized as foreign by the host immune system. The discovery of the existence of tumor associated antigens has raised the possibility of using a host's immune system to intervene in tumor growth. Various mechanisms of harnessing both the humoral and cellular arms of the immune system are currently being explored for cancer immunotherapy.

Specific elements of the cellular immune response are capable of specifically recognizing and destroying tumor cells. The isolation of T-cells from tumor-infiltrating cell populations or from peripheral blood suggests that such cells play a significant role in natural immune defense against cancer. CD8-positive T-cells in particular, which recognize class I molecules of the major histocompatibility complex (MHC)-bearing peptides of usually 8 to 10 amino acid residues derived from proteins or defect ribosomal products (DRIPS) located in the cytosol, play a significant role in this

response. The MHC-molecules of the human are also designated as human leukocyte-antigens (HLA).

As used herein and except as noted otherwise all terms are defined as given below.

The term "T-cell response" means the specific proliferation and activation of effector functions induced by a peptide *in vitro* or *in vivo*. For MHC class I restricted cytotoxic T cells, effector functions may be lysis of peptide-pulsed, peptide-precursor pulsed or naturally peptide-presenting target cells, secretion of cytokines, preferably Interferon-gamma, TNF-alpha, or IL-2 induced by peptide, secretion of effector molecules, preferably granzymes or perforins induced by peptide, or degranulation.

The term "peptide" is used herein to designate a series of amino acid residues, connected one to the other typically by peptide bonds between the alpha-amino and carbonyl groups of the adjacent amino acids. The peptides are preferably 9 amino acids in length but can be as short as 8 amino acids in length, and as long as 10, 11, 12, or 13 or longer, and in case of MHC class II peptides (elongated variants of the peptides of the invention) they can be as long as 14, 15, 16, 17, 18, 19 or 20 or more amino acids in length.

Furthermore, the term "peptide" shall include salts of a series of amino acid residues, connected one to the other typically by peptide bonds between the alpha-amino and carbonyl groups of the adjacent amino acids. Preferably, the salts are pharmaceutical acceptable salts of the peptides, such as, for example, the chloride or acetate (trifluoroacetate) salts. It has to be noted that the salts of the peptides according to the present invention differ substantially from the peptides in their state(s) *in vivo*, as the peptides are not salts *in vivo*.

The term "peptide" shall also include "oligopeptide". The term "oligopeptide" is used herein to designate a series of amino acid residues, connected one to the other typically by peptide bonds between the alpha-amino and carbonyl groups of the adjacent amino

acids. The length of the oligopeptide is not critical to the invention, as long as the correct epitope or epitopes are maintained therein. The oligopeptides are typically less than about 30 amino acid residues in length, and greater than about 15 amino acids in length.

The term "polypeptide" designates a series of amino acid residues, connected one to the other typically by peptide bonds between the alpha-amino and carbonyl groups of the adjacent amino acids. The length of the polypeptide is not critical to the invention as long as the correct epitopes are maintained. In contrast to the terms peptide or oligopeptide, the term polypeptide is meant to refer to molecules containing more than about 30 amino acid residues.

A peptide, oligopeptide, protein or polynucleotide coding for such a molecule is "immunogenic" (and thus is an "immunogen" within the present invention), if it is capable of inducing an immune response. In the case of the present invention, immunogenicity is more specifically defined as the ability to induce a T-cell response. Thus, an "immunogen" would be a molecule that is capable of inducing an immune response, and in the case of the present invention, a molecule capable of inducing a T-cell response. In another aspect, the immunogen can be the peptide, the complex of the peptide with MHC, oligopeptide, and/or protein that is used to raise specific antibodies or TCRs against it.

A class I T cell "epitope" requires a short peptide that is bound to a class I MHC receptor, forming a ternary complex (MHC class I alpha chain, beta-2-microglobulin, and peptide) that can be recognized by a T cell bearing a matching T-cell receptor binding to the MHC/peptide complex with appropriate affinity. Peptides binding to MHC class I molecules are typically 8-14 amino acids in length, and most typically 9 amino acids in length.

In humans there are three different genetic loci that encode MHC class I molecules (the MHC-molecules of the human are also designated human leukocyte antigens (HLA)):

HLA-A, HLA-B, and HLA-C. HLA-A*01, HLA-A*02, and HLA-B*07 are examples of different MHC class I alleles that can be expressed from these loci.

Table 4: Expression frequencies F of HLA-A*02, HLA-A*01, HLA-A*03, HLA-A*24, HLA-B*07, HLA-B*08 and HLA-B*44 serotypes. Haplotype frequencies Gf are derived from a study which used HLA-typing data from a registry of more than 6.5 million volunteer donors in the U.S. (Gragert et al., 2013). The haplotype frequency is the frequency of a distinct allele on an individual chromosome. Due to the diploid set of chromosomes within mammalian cells, the frequency of genotypic occurrence of this allele is higher, and can be calculated employing the Hardy-Weinberg principle ($F = 1 - (1 - Gf)^2$).

Allele	Population	Calculated phenotype from allele frequency (F)
A*02	African (N=28557)	32.3%
	European Caucasian (N=1242890)	49.3%
	Japanese (N=24582)	42.7%
	Hispanic, S + Cent Amer. (N=146714)	46.1%
	Southeast Asian (N=27978)	30.4%
A*01	African (N=28557)	10.2%
	European Caucasian (N=1242890)	30.2%
	Japanese (N=24582)	1.8%
	Hispanic, S + Cent Amer. (N=146714)	14.0%
	Southeast Asian (N=27978)	21.0%
A*03	African (N=28557)	14.8%
	European Caucasian (N=1242890)	26.4%
	Japanese (N=24582)	1.8%
	Hispanic, S + Cent Amer. (N=146714)	14.4%
	Southeast Asian (N=27978)	10.6%
A*24	African (N=28557)	2.0%
	European Caucasian (N=1242890)	8.6%
	Japanese (N=24582)	35.5%
	Hispanic, S + Cent Amer. (N=146714)	13.6%

	Southeast Asian (N=27978)	16.9%
B*07	African (N=28557)	14.7%
	European Caucasian (N=1242890)	25.0%
	Japanese (N=24582)	11.4%
	Hispanic, S + Cent Amer. (N=146714)	12.2%
	Southeast Asian (N=27978)	10.4%
B*08	African (N=28557)	6.0%
	European Caucasian (N=1242890)	21.6%
	Japanese (N=24582)	1.0%
	Hispanic, S + Cent Amer. (N=146714)	7.6%
	Southeast Asian (N=27978)	6.2%
B*44	African (N=28557)	10.6%
	European Caucasian (N=1242890)	26.9%
	Japanese (N=24582)	13.0%
	Hispanic, S + Cent Amer. (N=146714)	18.2%
	Southeast Asian (N=27978)	13.1%

The peptides of the invention, preferably when included into a vaccine of the invention as described herein bind to B*07. A vaccine may also include pan-binding MHC class II peptides. Therefore, the vaccine of the invention can be used to treat cancer in patients that are B*07-positive, whereas no selection for MHC class II allotypes is necessary due to the pan-binding nature of these peptides.

If B*07 peptides of the invention are combined with peptides binding to another allele, for example A*24, a higher percentage of any patient population can be treated compared with addressing either MHC class I allele alone. While in most populations less than 50% of patients could be addressed by either allele alone, a vaccine comprising HLA-A*24 and HLA-A*02 epitopes can treat at least 60% of patients in any relevant population. Specifically, the following percentages of patients will be positive for

at least one of these alleles in various regions: USA 61%, Western Europe 62%, China 75%, South Korea 77%, Japan 86% (calculated from www.allelefreqencies.net).

Table 5: HLA alleles coverage in European Caucasian population (calculated from (Gragert et al., 2013)).

	coverage (at least one A- allele)	combined with B*07	combined with B*44	combined with B*07 and B*44
A*02 / A*01	70%	78%	78%	84%
A*02 / A*03	68%	76%	76%	83%
A*02 / A*24	61%	71%	71%	80%
A*01 / A*03	52%	64%	65%	75%
A*01 / A*24	44%	58%	59%	71%
A*03 / A*24	40%	55%	56%	69%
A*02 / A*01 / A*03	84%	88%	88%	91%
A*02 / A*01 / A*24	79%	84%	84%	89%
A*02 / A*03 / A*24	77%	82%	83%	88%
A*01 / A*03 / A*24	63%	72%	73%	81%
A*02 / A*01 / A*03 / A*24	90%	92%	93%	95%

In a preferred embodiment, the term “nucleotide sequence” refers to a heteropolymer of deoxyribonucleotides.

The nucleotide sequence coding for a particular peptide, oligopeptide, or polypeptide may be naturally occurring or they may be synthetically constructed. Generally, DNA segments encoding the peptides, polypeptides, and proteins of this invention are assembled from cDNA fragments and short oligonucleotide linkers, or from a series of oligonucleotides, to provide a synthetic gene that is capable of being expressed in a recombinant transcriptional unit comprising regulatory elements derived from a microbial or viral operon.

As used herein the term “a nucleotide coding for (or encoding) a peptide” refers to a nucleotide sequence coding for the peptide including artificial (man-made) start and

stop codons compatible for the biological system the sequence is to be expressed by, for example, a dendritic cell or another cell system useful for the production of TCRs.

As used herein, reference to a nucleic acid sequence includes both single stranded and double stranded nucleic acid. Thus, for example for DNA, the specific sequence, unless the context indicates otherwise, refers to the single strand DNA of such sequence, the duplex of such sequence with its complement (double stranded DNA) and the complement of such sequence.

The term “coding region” refers to that portion of a gene which either naturally or normally codes for the expression product of that gene in its natural genomic environment, i.e., the region coding *in vivo* for the native expression product of the gene.

The coding region can be derived from a non-mutated (“normal”), mutated or altered gene, or can even be derived from a DNA sequence, or gene, wholly synthesized in the laboratory using methods well known to those of skill in the art of DNA synthesis.

The term “expression product” means the polypeptide or protein that is the natural translation product of the gene and any nucleic acid sequence coding equivalents resulting from genetic code degeneracy and thus coding for the same amino acid(s).

The term “fragment”, when referring to a coding sequence, means a portion of DNA comprising less than the complete coding region, whose expression product retains essentially the same biological function or activity as the expression product of the complete coding region.

The term “DNA segment” refers to a DNA polymer, in the form of a separate fragment or as a component of a larger DNA construct, which has been derived from DNA isolated at least once in substantially pure form, i.e., free of contaminating endogenous materials and in a quantity or concentration enabling identification, manipulation, and recovery of

the segment and its component nucleotide sequences by standard biochemical methods, for example, by using a cloning vector. Such segments are provided in the form of an open reading frame uninterrupted by internal non-translated sequences, or introns, which are typically present in eukaryotic genes. Sequences of non-translated DNA may be present downstream from the open reading frame, where the same do not interfere with manipulation or expression of the coding regions.

The term “primer” means a short nucleic acid sequence that can be paired with one strand of DNA and provides a free 3'-OH end at which a DNA polymerase starts synthesis of a deoxyribonucleotide chain.

The term “promoter” means a region of DNA involved in binding of RNA polymerase to initiate transcription.

The term “isolated” means that the material is removed from its original environment (e.g., the natural environment, if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

The polynucleotides, and recombinant or immunogenic polypeptides, disclosed in accordance with the present invention may also be in “purified” form. The term “purified” does not require absolute purity; rather, it is intended as a relative definition, and can include preparations that are highly purified or preparations that are only partially purified, as those terms are understood by those of skill in the relevant art. For example, individual clones isolated from a cDNA library have been conventionally purified to electrophoretic homogeneity. Purification of starting material or natural material to at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly contemplated. Furthermore, a claimed

polypeptide which has a purity of preferably 99.999%, or at least 99.99% or 99.9%; and even desirably 99% by weight or greater is expressly encompassed.

The nucleic acids and polypeptide expression products disclosed according to the present invention, as well as expression vectors containing such nucleic acids and/or such polypeptides, may be in "enriched form". As used herein, the term "enriched" means that the concentration of the material is at least about 2, 5, 10, 100, or 1000 times its natural concentration (for example), advantageously 0.01%, by weight, preferably at least about 0.1% by weight. Enriched preparations of about 0.5%, 1%, 5%, 10%, and 20% by weight are also contemplated. The sequences, constructs, vectors, clones, and other materials comprising the present invention can advantageously be in enriched or isolated form. The term "active fragment" means a fragment, usually of a peptide, polypeptide or nucleic acid sequence, that generates an immune response (i.e., has immunogenic activity) when administered, alone or optionally with a suitable adjuvant or in a vector, to an animal, such as a mammal, for example, a rabbit or a mouse, and also including a human, such immune response taking the form of stimulating a T-cell response within the recipient animal, such as a human. Alternatively, the "active fragment" may also be used to induce a T-cell response *in vitro*.

As used herein, the terms "portion", "segment" and "fragment", when used in relation to polypeptides, refer to a continuous sequence of residues, such as amino acid residues, which sequence forms a subset of a larger sequence. For example, if a polypeptide were subjected to treatment with any of the common endopeptidases, such as trypsin or chymotrypsin, the oligopeptides resulting from such treatment would represent portions, segments or fragments of the starting polypeptide. When used in relation to polynucleotides, these terms refer to the products produced by treatment of said polynucleotides with any of the endonucleases.

In accordance with the present invention, the term "percent identity" or "percent identical", when referring to a sequence, means that a sequence is compared to a claimed or described sequence after alignment of the sequence to be compared (the

- 37 -

“Compared Sequence”) with the described or claimed sequence (the “Reference Sequence”). The percent identity is then determined according to the following formula:

$$\text{percent identity} = 100 [1 - (C/R)]$$

wherein C is the number of differences between the Reference Sequence and the Compared Sequence over the length of alignment between the Reference Sequence and the Compared Sequence, wherein

- (i) each base or amino acid in the Reference Sequence that does not have a corresponding aligned base or amino acid in the Compared Sequence and
- (ii) each gap in the Reference Sequence and
- (iii) each aligned base or amino acid in the Reference Sequence that is different from an aligned base or amino acid in the Compared Sequence, constitutes a difference and
- (iv) the alignment has to start at position 1 of the aligned sequences;

and R is the number of bases or amino acids in the Reference Sequence over the length of the alignment with the Compared Sequence with any gap created in the Reference Sequence also being counted as a base or amino acid.

If an alignment exists between the Compared Sequence and the Reference Sequence for which the percent identity as calculated above is about equal to or greater than a specified minimum Percent Identity, then the Compared Sequence has the specified minimum percent identity to the Reference Sequence even though alignments may exist in which the herein above calculated percent identity is less than the specified percent identity.

As mentioned above, the present invention thus provides a peptide comprising a sequence that is selected from the group of consisting of SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499 or a variant thereof which is 88% homologous to SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499, or a variant thereof that will induce T cells cross-reacting with said peptide. The peptides of the invention have the ability to bind to a molecule of the human major histocompatibility complex (MHC) class-I or elongated versions of said peptides to class II.

In the present invention, the term "homologous" refers to the degree of identity (see percent identity above) between sequences of two amino acid sequences, i.e. peptide or polypeptide sequences. The aforementioned "homology" is determined by comparing two sequences aligned under optimal conditions over the sequences to be compared. Such a sequence homology can be calculated by creating an alignment using, for example, the ClustalW algorithm. Commonly available sequence analysis software, more specifically, Vector NTI, GENETYX or other tools are provided by public databases.

A person skilled in the art will be able to assess, whether T cells induced by a variant of a specific peptide will be able to cross-react with the peptide itself (Appay et al., 2006; Colombetti et al., 2006; Fong et al., 2001; Zaremba et al., 1997).

By a "variant" of the given amino acid sequence the inventors mean that the side chains of, for example, one or two of the amino acid residues are altered (for example by replacing them with the side chain of another naturally occurring amino acid residue or some other side chain) such that the peptide is still able to bind to an HLA molecule in substantially the same way as a peptide consisting of the given amino acid sequence in consisting of SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499. For example, a peptide may be modified so that it at least maintains, if not improves, the ability to interact with and bind to the binding groove of a suitable MHC molecule, such as HLA-A*02 or -DR, and in that way, it at least maintains, if not improves, the ability to bind to the TCR of activated T cells.

These T cells can subsequently cross-react with cells and kill cells that express a polypeptide that contains the natural amino acid sequence of the cognate peptide as defined in the aspects of the invention. As can be derived from the scientific literature and databases (Godkin et al., 1997; Rammensee et al., 1999), certain positions of HLA binding peptides are typically anchor residues forming a core sequence fitting to the binding motif of the HLA receptor, which is defined by polar, electrophysical,

hydrophobic and spatial properties of the polypeptide chains constituting the binding groove. Thus, one skilled in the art would be able to modify the amino acid sequences set forth in SEQ ID NO: 1 to SEQ ID NO 383 and SEQ ID NO: 448 to SEQ ID NO: 499, by maintaining the known anchor residues, and would be able to determine whether such variants maintain the ability to bind MHC class I or II molecules. The variants of the present invention retain the ability to bind to the TCR of activated T cells, which can subsequently cross-react with and kill cells that express a polypeptide containing the natural amino acid sequence of the cognate peptide as defined in the aspects of the invention.

The original (unmodified) peptides as disclosed herein can be modified by the substitution of one or more residues at different, possibly selective, sites within the peptide chain, if not otherwise stated. Preferably those substitutions are located at the end of the amino acid chain. Such substitutions may be of a conservative nature, for example, where one amino acid is replaced by an amino acid of similar structure and characteristics, such as where a hydrophobic amino acid is replaced by another hydrophobic amino acid. Even more conservative would be replacement of amino acids of the same or similar size and chemical nature, such as where leucine is replaced by isoleucine. In studies of sequence variations in families of naturally occurring homologous proteins, certain amino acid substitutions are more often tolerated than others, and these often show correlation with similarities in size, charge, polarity, and hydrophobicity between the original amino acid and its replacement, and such is the basis for defining "conservative substitutions."

Conservative substitutions are herein defined as exchanges within one of the following five groups: Group 1-small aliphatic, nonpolar or slightly polar residues (Ala, Ser, Thr, Pro, Gly); Group 2-polar, negatively charged residues and their amides (Asp, Asn, Glu, Gln); Group 3-polar, positively charged residues (His, Arg, Lys); Group 4-large, aliphatic, nonpolar residues (Met, Leu, Ile, Val, Cys); and Group 5-large, aromatic residues (Phe, Tyr, Trp).

In an aspect, conservative substitutions may include those, which are described by Dayhoff in "The Atlas of Protein Sequence and Structure. Vol. 5", *Natl. Biomedical Research*, the contents of which are incorporated by reference in their entirety. For example, in an aspect, amino acids, which belong to one of the following groups, can be exchanged for one another, thus, constituting a conservative exchange: Group 1: alanine (A), proline (P), glycine (G), asparagine (N), serine (S), threonine (T); Group 2: cysteine (C), serine (S), tyrosine (Y), threonine (T); Group 3: valine (V), isoleucine (I), leucine (L), methionine (M), alanine (A), phenylalanine (F); Group 4: lysine (K), arginine (R), histidine (H); Group 5: phenylalanine (F), tyrosine (Y), tryptophan (W), histidine (H); and Group 6: aspartic acid (D), glutamic acid (E). In an aspect, a conservative amino acid substitution may be selected from the following of T→A, G→A, A→I, T→V, A→M, T→I, A→V, T→G, and/or T→S.

In an aspect, a conservative amino acid substitution may include the substitution of an amino acid by another amino acid of the same class, for example, (1) nonpolar: Ala, Val, Leu, Ile, Pro, Met, Phe, Trp; (2) uncharged polar: Gly, Ser, Thr, Cys, Tyr, Asn, Gln; (3) acidic: Asp, Glu; and (4) basic: Lys, Arg, His. Other conservative amino acid substitutions may also be made as follows: (1) aromatic: Phe, Tyr, His; (2) proton donor: Asn, Gln, Lys, Arg, His, Trp; and (3) proton acceptor: Glu, Asp, Thr, Ser, Tyr, Asn, Gln (see, for example, U.S. Patent No. 10,106,805, the contents of which are incorporated by reference in their entirety).

In another aspect, conservative substitutions may be made in accordance with Table A. Methods for predicting tolerance to protein modification may be found in, for example, Guo et al., *Proc. Natl. Acad. Sci., USA*, 101(25):9205-9210 (2004), the contents of which are incorporated by reference in their entirety.

Table A

Conservative Amino Acid Substitutions	
Amino Acid	Substitutions (others are known in the art)
Ala	Ser, Gly, Cys
Arg	Lys, Gln, His
Asn	Gln, His, Glu, Asp
Asp	Glu, Asn, Gln
Cys	Ser, Met, Thr
Gln	Asn, Lys, Glu, Asp, Arg
Glu	Asp, Asn, Gln
Gly	Pro, Ala, Ser
His	Asn, Gln, Lys
Ile	Leu, Val, Met, Ala
Leu	Ile, Val, Met, Ala
Lys	Arg, Gln, His
Met	Leu, Ile, Val, Ala, Phe
Phe	Met, Leu, Tyr, Trp, His
Ser	Thr, Cys, Ala
Thr	Ser, Val, Ala
Trp	Tyr, Phe
Tyr	Trp, Phe, His
Val	Ile, Leu, Met, Ala, Thr

In another aspect, conservative substitutions may be those shown in Table B under the heading of “conservative substitutions.” If such substitutions result in a change in biological activity, then more substantial changes, denominated “exemplary substitutions” in Table B, may be introduced and the products screened if needed.

Table B

Amino Acid Substitutions		
Original Residue (naturally occurring amino acid)	Conservative Substitutions	Exemplary Substitutions
Ala (A)	Val	Val; Leu; Ile
Arg (R)	Lys	Lys; Gln; Asn
Asn (N)	Gln	Gln; His; Asp, Lys; Arg
Asp (D)	Glu	Glu; Asn
Cys (C)	Ser	Ser; Ala
Gln (Q)	Asn	Asn; Glu
Glu (E)	Asp	Asp; Gln
Gly (G)	Ala	Ala
His (H)	Arg	Asn; Gln; Lys; Arg
Ile (I)	Leu	Leu; Val; Met; Ala; Phe; Norleucine
Leu (L)	Ile	Norleucine; Ile; Val; Met; Ala; Phe
Lys (K)	Arg	Arg; Gln; Asn
Met (M)	Leu	Leu; Phe; Ile
Phe (F)	Tyr	Leu; Val; Ile; Ala; Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Ser	Ser
Trp (W)	Tyr	Tyr; Phe
Tyr (Y)	Phe	Trp; Phe; Thr; Ser
Val (V)	Leu	Ile; Leu; Met; Phe; Ala; Norleucine

Less conservative substitutions might involve the replacement of one amino acid by another that has similar characteristics but is somewhat different in size, such as replacement of an alanine by an isoleucine residue. Highly non-conservative replacements might involve substituting an acidic amino acid for one that is polar, or even for one that is basic in character. Such “radical” substitutions cannot, however, be dismissed as potentially ineffective since chemical effects are not totally predictable and

radical substitutions might well give rise to serendipitous effects not otherwise predictable from simple chemical principles.

Of course, such substitutions may involve structures other than the common L-amino acids. Thus, D-amino acids might be substituted for the L-amino acids commonly found in the antigenic peptides of the invention and yet still be encompassed by the disclosure herein. In addition, non-standard amino acids (i.e., other than the common naturally occurring proteinogenic amino acids) may also be used for substitution purposes to produce immunogens and immunogenic polypeptides according to the present invention.

If substitutions at more than one position are found to result in a peptide with substantially equivalent or greater antigenic activity as defined below, then combinations of those substitutions will be tested to determine if the combined substitutions result in additive or synergistic effects on the antigenicity of the peptide. At most, no more than four positions within the peptide would be simultaneously substituted.

A peptide consisting essentially of the amino acid sequence as indicated herein can have one or two non-anchor amino acids (see below regarding the anchor motif) exchanged without that the ability to bind to a molecule of the human major histocompatibility complex (MHC) class-I or –II is substantially changed or is negatively affected, when compared to the non-modified peptide. In another embodiment, in a peptide consisting essentially of the amino acid sequence as indicated herein, one or two amino acids can be exchanged with their conservative exchange partners (see herein below) without that the ability to bind to a molecule of the human major histocompatibility complex (MHC) class-I or –II is substantially changed, or is negatively affected, when compared to the non-modified peptide.

The amino acid residues that do not substantially contribute to interactions with the T-cell receptor can be modified by replacement with other amino acid whose incorporation

Table 6: Variants and motif of the peptides according to SEQ ID NO: 13, 25, 38, 54 and 362

[illegible]

										M	
										A	
										I	
Position	1	2	3	4	5	6	7	8	9	10	11
Seq ID No 54	V	P	S	C	G	R	S	V	E	G	L
Variant											F
											V
											M
											A
											I
Position	1	2	3	4	5	6	7	8	9		
Seq ID No 362	K	P	M	L	P	P	A	A	F		
Variant									L		
									V		
									M		
									A		
									I		

Longer (elongated) peptides may also be suitable. It is possible that MHC class I epitopes, although usually between 8 and 11 amino acids long, are generated by peptide processing from longer peptides or proteins that include the actual epitope. It is preferred that the residues that flank the actual epitope are residues that do not substantially affect proteolytic cleavage necessary to expose the actual epitope during processing.

The peptides of the invention can be elongated by up to four amino acids, that is 1, 2, 3 or 4 amino acids can be added to either end in any combination between 4:0 and 0:4. Combinations of the elongations according to the invention can be found in Table 7.

Table 7: Combinations of the elongations of peptides of the invention

C-terminus	N-terminus
4	0
3	0 or 1
2	0 or 1 or 2
1	0 or 1 or 2 or 3
0	0 or 1 or 2 or 3 or 4
N-terminus	C-terminus
4	0
3	0 or 1
2	0 or 1 or 2
1	0 or 1 or 2 or 3
0	0 or 1 or 2 or 3 or 4

The amino acids for the elongation/extension can be the peptides of the original sequence of the protein or any other amino acid(s). The elongation can be used to enhance the stability or solubility of the peptides.

Thus, the epitopes of the present invention may be identical to naturally occurring tumor-associated or tumor-specific epitopes or may include epitopes that differ by no more than four residues from the reference peptide, as long as they have substantially identical antigenic activity.

In an alternative embodiment, the peptide is elongated on either or both sides by more than 4 amino acids, preferably to a total length of up to 30 amino acids. This may lead to MHC class II binding peptides. Binding to MHC class II can be tested by methods known in the art.

Accordingly, the present invention provides peptides and variants of MHC class I epitopes, wherein the peptide or variant has an overall length of between 8 and 100, preferably between 8 and 30, and most preferred between 8 and 14, namely 8, 9, 10, 11, 12, 13, 14 amino acids, in case of the elongated class II binding peptides the length can also be 15, 16, 17, 18, 19, 20, 21 or 22 amino acids.

Of course, the peptide or variant according to the present invention will have the ability to bind to a molecule of the human major histocompatibility complex (MHC) class I or II. Binding of a peptide or a variant to an MHC complex may be tested by methods known in the art.

Preferably, when the T cells specific for a peptide according to the present invention are tested against the substituted peptides, the peptide concentration at which the substituted peptides achieve half the maximal increase in lysis relative to background is no more than about 1 mM, preferably no more than about 1 μ M, more preferably no more than about 1 nM, and still more preferably no more than about 100 pM, and most preferably no more than about 10 pM. It is also preferred that the substituted peptide be recognized by T cells from more than one individual, at least two, and more preferably three individuals.

In a particularly preferred embodiment of the invention the peptide consists or consists essentially of an amino acid sequence according to SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499.

“Consisting essentially of” shall mean that a peptide according to the present invention, in addition to the sequence according to any of SEQ ID NO: 1 to SEQ ID NO 383 and SEQ ID NO: 448 to SEQ ID NO: 499 or a variant thereof contains additional N- and/or C-terminally located stretches of amino acids that are not necessarily forming part of the peptide that functions as an epitope for MHC molecules epitope.

Nevertheless, these stretches can be important to provide an efficient introduction of the peptide according to the present invention into the cells. In one embodiment of the present invention, the peptide is part of a fusion protein which comprises, for example, the 80 N-terminal amino acids of the HLA-DR antigen-associated invariant chain (p33, in the following “Ii”) as derived from the NCBI, GenBank Accession number X00497. In other fusions, the peptides of the present invention can be fused to an antibody as described herein, or a functional part thereof, in particular into a sequence of an

antibody, so as to be specifically targeted by said antibody, or, for example, to or into an antibody that is specific for dendritic cells as described herein.

In addition, the peptide or variant may be modified further to improve stability and/or binding to MHC molecules in order to elicit a stronger immune response. Methods for such an optimization of a peptide sequence are well known in the art and include, for example, the introduction of reverse peptide bonds or non-peptide bonds.

In a reverse peptide bond amino acid residues are not joined by peptide (-CO-NH-) linkages, but the peptide bond is reversed. Such retro-inverso peptidomimetics may be made using methods known in the art, for example such as those described in Meziere et al (1997) (Meziere et al., 1997), incorporated herein by reference. This approach involves making pseudopeptides containing changes involving the backbone, and not the orientation of side chains. Meziere et al. (Meziere et al., 1997) show that for MHC binding and T helper cell responses, these pseudopeptides are useful. Retro-inverse peptides, which contain NH-CO bonds instead of CO-NH peptide bonds, are much more resistant to proteolysis.

A non-peptide bond is, for example, -CH₂-NH-, -CH₂S-, -CH₂CH₂-, -CH=CH-, -COCH₂-, -CH(OH)CH₂-, and -CH₂SO-. US 4,897,445 provides a method for the solid phase synthesis of non-peptide bonds (-CH₂-NH) in polypeptide chains which involves polypeptides synthesized by standard procedures and the non-peptide bond synthesized by reacting an amino aldehyde and an amino acid in the presence of NaCNBH₃.

Peptides comprising the sequences described above may be synthesized with additional chemical groups present at their amino and/or carboxy termini, to enhance the stability, bioavailability, and/or affinity of the peptides. For example, hydrophobic groups such as carbobenzoxy, dansyl, or t-butyloxycarbonyl groups may be added to the peptides' amino termini. Likewise, an acetyl group or a 9-fluorenylmethoxy-carbonyl group may be placed at the peptides' amino termini. Additionally, the hydrophobic

group, t-butyloxycarbonyl, or an amido group may be added to the peptides' carboxy termini.

Further, the peptides of the invention may be synthesized to alter their steric configuration. For example, the D-isomer of one or more of the amino acid residues of the peptide may be used, rather than the usual L-isomer. Still further, at least one of the amino acid residues of the peptides of the invention may be substituted by one of the well-known non-naturally occurring amino acid residues. Alterations such as these may serve to increase the stability, bioavailability and/or binding action of the peptides of the invention.

Similarly, a peptide or variant of the invention may be modified chemically by reacting specific amino acids either before or after synthesis of the peptide. Examples for such modifications are well known in the art and are summarized e.g. in R. Lundblad, *Chemical Reagents for Protein Modification*, 3rd ed. CRC Press, 2004 (Lundblad, 2004), which is incorporated herein by reference. Chemical modification of amino acids includes but is not limited to, modification by acylation, amidination, pyridoxylation of lysine, reductive alkylation, trinitrobenzylation of amino groups with 2,4,6-trinitrobenzene sulphonic acid (TNBS), amide modification of carboxyl groups and sulphydryl modification by performic acid oxidation of cysteine to cysteic acid, formation of mercurial derivatives, formation of mixed disulphides with other thiol compounds, reaction with maleimide, carboxymethylation with iodoacetic acid or iodoacetamide and carbamoylation with cyanate at alkaline pH, although without limitation thereto. In this regard, the skilled person is referred to Chapter 15 of *Current Protocols In Protein Science*, Eds. Coligan et al. (John Wiley and Sons NY 1995-2000) (Coligan et al., 1995) for more extensive methodology relating to chemical modification of proteins.

Briefly, modification of e.g. arginyl residues in proteins is often based on the reaction of vicinal dicarbonyl compounds such as phenylglyoxal, 2,3-butanedione, and 1,2-cyclohexanedione to form an adduct. Another example is the reaction of methylglyoxal with arginine residues. Cysteine can be modified without concomitant modification of

other nucleophilic sites such as lysine and histidine. As a result, a large number of reagents are available for the modification of cysteine. The websites of companies such as Sigma-Aldrich (<http://www.sigma-aldrich.com>) provide information on specific reagents.

Selective reduction of disulfide bonds in proteins is also common. Disulfide bonds can be formed and oxidized during the heat treatment of biopharmaceuticals. Woodward's Reagent K may be used to modify specific glutamic acid residues. N-(3-(dimethylamino)propyl)-N'-ethylcarbodiimide can be used to form intra-molecular crosslinks between a lysine residue and a glutamic acid residue. For example, diethylpyrocarbonate is a reagent for the modification of histidyl residues in proteins. Histidine can also be modified using 4-hydroxy-2-nonenal. The reaction of lysine residues and other α -amino groups is, for example, useful in binding of peptides to surfaces or the cross-linking of proteins/peptides. Lysine is the site of attachment of poly(ethylene)glycol and the major site of modification in the glycosylation of proteins. Methionine residues in proteins can be modified with e.g. iodoacetamide, bromoethylamine, and chloramine T.

Tetranitromethane and N-acetylimidazole can be used for the modification of tyrosyl residues. Cross-linking via the formation of dityrosine can be accomplished with hydrogen peroxide/copper ions.

Recent studies on the modification of tryptophan have used N-bromosuccinimide, 2-hydroxy-5-nitrobenzyl bromide or 3-bromo-3-methyl-2-(2-nitrophenylmercapto)-3H-indole (BPNS-skatole).

Successful modification of therapeutic proteins and peptides with PEG is often associated with an extension of circulatory half-life while cross-linking of proteins with glutaraldehyde, polyethylene glycol diacrylate and formaldehyde is used for the preparation of hydrogels. Chemical modification of allergens for immunotherapy is often achieved by carbamylation with potassium cyanate.

A peptide or variant, wherein the peptide is modified or includes non-peptide bonds is a preferred embodiment of the invention.

Another embodiment of the present invention relates to a non-naturally occurring peptide wherein said peptide consists or consists essentially of an amino acid sequence according to SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499 and has been synthetically produced (e.g. synthesized) as a pharmaceutically acceptable salt. Methods to synthetically produce peptides are well known in the art. The salts of the peptides according to the present invention differ substantially from the peptides in their state(s) *in vivo*, as the peptides as generated *in vivo* are no salts. The non-natural salt form of the peptide mediates the solubility of the peptide, in particular in the context of pharmaceutical compositions comprising the peptides, e.g. the peptide vaccines as disclosed herein. A sufficient and at least substantial solubility of the peptide(s) is required in order to efficiently provide the peptides to the subject to be treated. Preferably, the salts are pharmaceutically acceptable salts of the peptides. These salts according to the invention include alkaline and earth alkaline salts such as salts of the Hofmeister series comprising as anions PO_4^{3-} , SO_4^{2-} , CH_3COO^- , Cl^- , Br^- , NO_3^- , ClO_4^- , I^- , SCN^- and as cations NH_4^+ , Rb^+ , K^+ , Na^+ , Cs^+ , Li^+ , Zn^{2+} , Mg^{2+} , Ca^{2+} , Mn^{2+} , Cu^{2+} and Ba^{2+} . Particularly salts are selected from $(\text{NH}_4)_3\text{PO}_4$, $(\text{NH}_4)_2\text{HPO}_4$, $(\text{NH}_4)\text{H}_2\text{PO}_4$, $(\text{NH}_4)_2\text{SO}_4$, $\text{NH}_4\text{CH}_3\text{COO}$, NH_4Cl , NH_4Br , NH_4NO_3 , NH_4ClO_4 , NH_4I , NH_4SCN , Rb_3PO_4 , Rb_2HPO_4 , RbH_2PO_4 , Rb_2SO_4 , $\text{Rb}_4\text{CH}_3\text{COO}$, Rb_4Cl , Rb_4Br , Rb_4NO_3 , Rb_4ClO_4 , Rb_4I , Rb_4SCN , K_3PO_4 , K_2HPO_4 , KH_2PO_4 , K_2SO_4 , KCH_3COO , KCl , KBr , KNO_3 , KClO_4 , KI , KSCN , Na_3PO_4 , Na_2HPO_4 , NaH_2PO_4 , Na_2SO_4 , NaCH_3COO , NaCl , NaBr , NaNO_3 , NaClO_4 , NaI , NaSCN , ZnCl_2 , Cs_3PO_4 , Cs_2HPO_4 , CsH_2PO_4 , Cs_2SO_4 , CsCH_3COO , CsCl , CsBr , CsNO_3 , CsClO_4 , CsI , CsSCN , Li_3PO_4 , Li_2HPO_4 , LiH_2PO_4 , Li_2SO_4 , LiCH_3COO , LiCl , LiBr , LiNO_3 , LiClO_4 , LiI , LiSCN , Cu_2SO_4 , $\text{Mg}_3(\text{PO}_4)_2$, Mg_2HPO_4 , $\text{Mg}(\text{H}_2\text{PO}_4)_2$, Mg_2SO_4 , $\text{Mg}(\text{CH}_3\text{COO})_2$, MgCl_2 , MgBr_2 , $\text{Mg}(\text{NO}_3)_2$, $\text{Mg}(\text{ClO}_4)_2$, MgI_2 , $\text{Mg}(\text{SCN})_2$, MnCl_2 , $\text{Ca}_3(\text{PO}_4)_2$, Ca_2HPO_4 , $\text{Ca}(\text{H}_2\text{PO}_4)_2$, CaSO_4 , $\text{Ca}(\text{CH}_3\text{COO})_2$, CaCl_2 , CaBr_2 , $\text{Ca}(\text{NO}_3)_2$, $\text{Ca}(\text{ClO}_4)_2$, CaI_2 , $\text{Ca}(\text{SCN})_2$, $\text{Ba}_3(\text{PO}_4)_2$, Ba_2HPO_4 , $\text{Ba}(\text{H}_2\text{PO}_4)_2$, BaSO_4 , $\text{Ba}(\text{CH}_3\text{COO})_2$, BaCl_2 , BaBr_2 , $\text{Ba}(\text{NO}_3)_2$, $\text{Ba}(\text{ClO}_4)_2$, BaI_2 , and $\text{Ba}(\text{SCN})_2$.

Particularly preferred are NH acetate, MgCl_2 , KH_2PO_4 , Na_2SO_4 , KCl, NaCl, and CaCl_2 , such as, for example, the chloride or acetate (trifluoroacetate) salts.

In a preferred embodiment, the pharmaceutical compositions comprise the peptides as salts of acetic acid (acetates), trifluoro acetates or hydrochloric acid (chlorides).

In an aspect, a polypeptide described herein is in the form of a pharmaceutically acceptable salt.

In another aspect, a polypeptide in the form of a pharmaceutical salt is in crystalline form.

In an aspect, a pharmaceutically acceptable salt described herein refers to salts which possess toxicity profiles within a range that is acceptable for pharmaceutical applications.

In an aspect, pharmaceutically acceptable salts may increase the solubility and/or stability of peptides of described herein. In another aspect, pharmaceutical salts described herein may be prepared by conventional means from the corresponding carrier peptide or complex by reacting, for example, the appropriate acid or base with peptides or complexes as described herein.

In another aspect, the pharmaceutically acceptable salts are in crystalline form or semi-crystalline form.

In yet another aspect, pharmaceutically acceptable salts may include, for example, those described in Handbook of Pharmaceutical Salts: Properties, Selection, and Use By P. H. Stahl and C. G. Wermuth (Wiley-VCH 2002) and L. D. Bighley, S. M. Berge, D. C. Monkhouse, in "Encyclopedia of Pharmaceutical Technology". Eds. J. Swarbrick and J. C. Boylan, Vol. 13, Marcel Dekker, Inc., New York, Basel, Hong Kong 1995, pp. 453-499, each of these references is herein incorporated by reference in their entirety.

Generally, peptides and variants (at least those containing peptide linkages between amino acid residues) may be synthesized by the Fmoc-polyamide mode of solid-phase peptide synthesis as disclosed by Lukas et al. (Lukas et al., 1981) and by references as cited therein. Temporary N-amino group protection is afforded by the 9-fluorenylmethyloxycarbonyl (Fmoc) group. Repetitive cleavage of this highly base-labile protecting group is done using 20% piperidine in N, N-dimethylformamide. Side-chain functionalities may be protected as their butyl ethers (in the case of serine threonine and tyrosine), butyl esters (in the case of glutamic acid and aspartic acid), butyloxycarbonyl derivative (in the case of lysine and histidine), trityl derivative (in the case of cysteine) and 4-methoxy-2,3,6-trimethylbenzenesulphonyl derivative (in the case of arginine). Where glutamine or asparagine are C-terminal residues, use is made of the 4,4'-dimethoxybenzhydryl group for protection of the side chain amido functionalities. The solid-phase support is based on a polydimethyl-acrylamide polymer constituted from the three monomers dimethylacrylamide (backbone-monomer), bisacryloylethylene diamine (cross linker) and acryloylsarcosine methyl ester (functionalizing agent). The peptide-to-resin cleavable linked agent used is the acid-labile 4-hydroxymethyl-phenoxyacetic acid derivative. All amino acid derivatives are added as their preformed symmetrical anhydride derivatives with the exception of asparagine and glutamine, which are added using a reversed N, N-dicyclohexyl-carbodiimide/1hydroxybenzotriazole mediated coupling procedure. All coupling and deprotection reactions are monitored using ninhydrin, trinitrobenzene sulphonic acid or isotin test procedures. Upon completion of synthesis, peptides are cleaved from the resin support with concomitant removal of side-chain protecting groups by treatment with 95% trifluoroacetic acid containing a 50 % scavenger mix. Scavengers commonly used include ethanedithiol, phenol, anisole and water, the exact choice depending on the constituent amino acids of the peptide being synthesized. Also a combination of solid phase and solution phase methodologies for the synthesis of peptides is possible (see, for example, (Bruckdorfer et al., 2004), and the references as cited therein).

Trifluoroacetic acid is removed by evaporation in vacuo, with subsequent trituration with diethyl ether affording the crude peptide. Any scavengers present are removed by a simple extraction procedure which on lyophilization of the aqueous phase affords the crude peptide free of scavengers. Reagents for peptide synthesis are generally available from e.g. Calbiochem-Novabiochem (Nottingham, UK).

Purification may be performed by anyone, or a combination of, techniques such as re-crystallization, size exclusion chromatography, ion-exchange chromatography, hydrophobic interaction chromatography and (usually) reverse-phase high performance liquid chromatography using e.g. acetonitrile/water gradient separation.

Analysis of peptides may be carried out using thin layer chromatography, electrophoresis, in particular capillary electrophoresis, solid phase extraction (CSPE), reverse-phase high performance liquid chromatography, amino-acid analysis after acid hydrolysis and by fast atom bombardment (FAB) mass spectrometric analysis, as well as MALDI and ESI-Q-TOF mass spectrometric analysis.

In order to select over-presented peptides, a presentation profile is calculated showing the median sample presentation as well as replicate variation. The profile juxtaposes samples of the tumor entity of interest to a baseline of normal tissue samples. Each of these profiles can then be consolidated into an over-presentation score by calculating the p-value of a Linear Mixed-Effects Model (Pinheiro et al., 2015) adjusting for multiple testing by False Discovery Rate (Benjamini and Hochberg, 1995) (cf. Example 1, Figures 1A-1I).

For the identification and relative quantitation of HLA ligands by mass spectrometry, HLA molecules from shock-frozen tissue samples were purified and HLA-associated peptides were isolated. The isolated peptides were separated, and sequences were identified by online nano-electrospray-ionization (nanoESI) liquid chromatography-mass spectrometry (LC-MS) experiments. The resulting peptide sequences were verified by comparison of the fragmentation pattern of natural tumor-associated peptides

(TUMAPs) recorded from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer samples (N = 149 samples) with the fragmentation patterns of corresponding synthetic reference peptides of identical sequences. Since the peptides were directly identified as ligands of HLA molecules of primary tumors, these results provide direct evidence for the natural processing and presentation of the identified peptides on primary cancer tissue obtained from 149 acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer patients.

The discovery pipeline XPRESIDENT® v2.1 (see, for example, US 2013-0096016, which is hereby incorporated by reference in its entirety) allows the identification and selection of relevant over-presented peptide vaccine candidates based on direct relative quantitation of HLA-restricted peptide levels on cancer tissues in comparison to several different non-cancerous tissues and organs. This was achieved by the development of label-free differential quantitation using the acquired LC-MS data processed by a proprietary data analysis pipeline, combining algorithms for sequence identification, spectral clustering, ion counting, retention time alignment, charge state deconvolution and normalization.

Presentation levels including error estimates for each peptide and sample were established. Peptides exclusively presented on tumor tissue and peptides over-presented in tumor versus non-cancerous tissues and organs have been identified.

HLA-peptide complexes from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer tissue samples were purified and HLA-associated peptides were isolated and analyzed by LC-MS (see example 1). All TUMAPs contained in the present application were identified with this approach on acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer samples confirming their presentation on acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

TUMAPs identified on multiple acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer,

glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer and normal tissues were quantified using ion-counting of label-free LC-MS data. The method assumes that LC-MS signal areas of a peptide correlate with its abundance in the sample. All quantitative signals of a peptide in various LC-MS experiments were normalized based on central tendency, averaged per sample and merged into a bar plot, called presentation profile. The presentation profile consolidates different analysis methods like protein database search, spectral clustering, charge state deconvolution (decharging) and retention time alignment and normalization.

Besides over-presentation of the peptide, mRNA expression of the underlying gene was tested. mRNA data were obtained via RNASeq analyses of normal tissues and cancer tissues (cf. Example 2, Figures 2A-2T). An additional source of normal tissue data was a database of publicly available RNA expression data from around 3000 normal tissue samples (Lonsdale, 2013). Peptides which are derived from proteins whose coding mRNA is highly expressed in cancer tissue, but very low or absent in vital normal tissues, were preferably included in the present invention.

The present invention provides peptides that are useful in treating cancers/tumors, preferably acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer that over- or exclusively present the peptides of the invention. These peptides were shown by mass spectrometry to be naturally presented by HLA molecules on primary human acute

myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer samples.

Many of the source gene/proteins (also designated “full-length proteins” or “underlying proteins”) from which the peptides are derived were shown to be highly over-expressed in cancer compared with normal tissues – “normal tissues” in relation to this invention shall mean either healthy blood cells, blood vessels, brain, heart, liver, lung, adrenal gland, bile duct, bladder, bone marrow, esophagus, gallbladder, large intestine, small intestine, kidney, lymph node, peripheral nerve, pancreas, pituitary, skin, spinal cord, spleen, stomach, thyroid, trachea cells or other normal tissue cells, demonstrating a high degree of tumor association of the source genes (see Example 2). Moreover, the peptides themselves are strongly over-presented on tumor tissue – “tumor tissue” in relation to this invention shall mean a sample from a patient suffering from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer, but not on normal tissues (see Example 1).

HLA-bound peptides can be recognized by the immune system, specifically T lymphocytes. T cells can destroy the cells presenting the recognized HLA/peptide complex, e.g. acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma,

gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer cells presenting the derived peptides.

The peptides of the present invention have been shown to be capable of stimulating T cell responses and/or are over-presented and thus can be used for the production of antibodies and/or TCRs, such as soluble TCRs, according to the present invention (see Example 3, Example 4). Furthermore, the peptides when complexed with the respective MHC can be used for the production of antibodies and/or TCRs, in particular sTCRs, according to the present invention, as well. Respective methods are well known to the person of skill and can be found in the respective literature as well (see also below). Thus, the peptides of the present invention are useful for generating an immune response in a patient by which tumor cells can be destroyed. An immune response in a patient can be induced by direct administration of the described peptides or suitable precursor substances (e.g. elongated peptides, proteins, or nucleic acids encoding these peptides) to the patient, ideally in combination with an agent enhancing the immunogenicity (i.e. an adjuvant). The immune response originating from such a therapeutic vaccination can be expected to be highly specific against tumor cells because the target peptides of the present invention are not presented on normal tissues in comparable copy numbers, preventing the risk of undesired autoimmune reactions against normal cells in the patient.

The present description further relates to T-cell receptors (TCRs) comprising an alpha chain and a beta chain ("alpha/beta TCRs"). Also provided are peptides according to the invention capable of binding to TCRs and antibodies when presented by an MHC molecule.

The present description also relates to fragments of the TCRs according to the invention that are capable of binding to a peptide antigen according to the present invention when presented by an HLA molecule. The term particularly relates to soluble TCR fragments, for example TCRs missing the transmembrane parts and/or constant regions, single chain TCRs, and fusions thereof to, for example, with Ig.

The present description also relates to nucleic acids, vectors and host cells for expressing TCRs and peptides of the present description; and methods of using the same.

The term "T-cell receptor" (abbreviated TCR) refers to a heterodimeric molecule comprising an alpha polypeptide chain (alpha chain) and a beta polypeptide chain (beta chain), wherein the heterodimeric receptor is capable of binding to a peptide antigen presented by an HLA molecule. The term also includes so-called gamma/delta TCRs.

In one embodiment the description provides a method of producing a TCR as described herein, the method comprising culturing a host cell capable of expressing the TCR under conditions suitable to promote expression of the TCR.

The description in another aspect relates to methods according to the description, wherein the antigen is loaded onto class I or II MHC molecules expressed on the surface of a suitable antigen-presenting cell or artificial antigen-presenting cell by contacting a sufficient amount of the antigen with an antigen-presenting cell or the antigen is loaded onto class I or II MHC tetramers by tetramerizing the antigen/class I or II MHC complex monomers.

The alpha and beta chains of alpha/beta TCR's, and the gamma and delta chains of gamma/delta TCRs, are generally regarded as each having two "domains", namely variable and constant domains. The variable domain consists of a concatenation of variable region (V) and joining region (J). The variable domain may also include a leader region (L). Beta and delta chains may also include a diversity region (D). The

alpha and beta constant domains may also include C-terminal transmembrane (TM) domains that anchor the alpha and beta chains to the cell membrane.

With respect to gamma/delta TCRs, the term "TCR gamma variable domain" as used herein refers to the concatenation of the TCR gamma V (TRGV) region without leader region (L), and the TCR gamma J (TRGJ) region, and the term TCR gamma constant domain refers to the extracellular TRGC region, or to a C-terminal truncated TRGC sequence. Likewise the term "TCR delta variable domain" refers to the concatenation of the TCR delta V (TRDV) region without leader region (L) and the TCR delta D/J (TRDD/TRDJ) region, and the term "TCR delta constant domain" refers to the extracellular TRDC region, or to a C-terminal truncated TRDC sequence.

TCRs of the present description preferably bind to a peptide-HLA molecule complex with a binding affinity (KD) of about 100 μ M or less, about 50 μ M or less, about 25 μ M or less, or about 10 μ M or less. More preferred are high affinity TCRs having binding affinities of about 1 μ M or less, about 100 nM or less, about 50 nM or less, about 25 nM or less. Non-limiting examples of preferred binding affinity ranges for TCRs of the present invention include about 1 nM to about 10 nM; about 10 nM to about 20 nM; about 20 nM to about 30 nM; about 30 nM to about 40 nM; about 40 nM to about 50 nM; about 50 nM to about 60 nM; about 60 nM to about 70 nM; about 70 nM to about 80 nM; about 80 nM to about 90 nM; and about 90 nM to about 100 nM.

As used herein in connect with TCRs of the present description, "specific binding" and grammatical variants thereof are used to mean a TCR having a binding affinity (KD) for a peptide-HLA molecule complex of 100 μ M or less.

Alpha/beta heterodimeric TCRs of the present description may have an introduced disulfide bond between their constant domains. Preferred TCRs of this type include those which have a TRAC constant domain sequence and a TRBC1 or TRBC2 constant domain sequence except that Thr 48 of TRAC and Ser 57 of TRBC1 or TRBC2 are replaced by cysteine residues, the said cysteines forming a disulfide bond between the

TRAC constant domain sequence and the TRBC1 or TRBC2 constant domain sequence of the TCR.

With or without the introduced inter-chain bond mentioned above, alpha/beta heterodimeric TCRs of the present description may have a TRAC constant domain sequence and a TRBC1 or TRBC2 constant domain sequence, and the TRAC constant domain sequence and the TRBC1 or TRBC2 constant domain sequence of the TCR may be linked by the native disulfide bond between Cys4 of exon 2 of TRAC and Cys2 of exon 2 of TRBC1 or TRBC2.

TCRs of the present description may comprise a detectable label selected from the group consisting of a radionuclide, a fluorophore and biotin. TCRs of the present description may be conjugated to a therapeutically active agent, such as a radionuclide, a chemotherapeutic agent, or a toxin.

In an embodiment, a TCR of the present description having at least one mutation in the alpha chain and/or having at least one mutation in the beta chain has modified glycosylation compared to the unmutated TCR.

In an embodiment, a TCR comprising at least one mutation in the TCR alpha chain and/or TCR beta chain has a binding affinity for, and/or a binding half-life for, a peptide-HLA molecule complex, which is at least double that of a TCR comprising the unmutated TCR alpha chain and/or unmutated TCR beta chain. Affinity-enhancement of tumor-specific TCRs, and its exploitation, relies on the existence of a window for optimal TCR affinities. The existence of such a window is based on observations that TCRs specific for HLA-B*07-restricted pathogens have KD values that are generally about 10-fold lower when compared to TCRs specific for HLA-B*07-restricted tumor-associated self-antigens. It is now known, although tumor antigens have the potential to be immunogenic, because tumors arise from the individual's own cells only mutated proteins or proteins with altered translational processing will be seen as foreign by the immune system. Antigens that are upregulated or overexpressed (so called self-

antigens) will not necessarily induce a functional immune response against the tumor: T-cells expressing TCRs that are highly reactive to these antigens will have been negatively selected within the thymus in a process known as central tolerance, meaning that only T-cells with low-affinity TCRs for self-antigens remain. Therefore, affinity of TCRs or variants of the present description to peptides can be enhanced by methods well known in the art.

The present description further relates to a method of identifying and isolating a TCR according to the present description, said method comprising incubating PBMCs from HLA-B*07-negative healthy donors with B*07/peptide monomers, incubating the PBMCs with tetramer-phycoerythrin (PE) and isolating the high avidity T-cells by fluorescence activated cell sorting (FACS)–Calibur analysis.

The present description further relates to a method of identifying and isolating a TCR according to the present description, said method comprising obtaining a transgenic mouse with the entire human TCR $\alpha\beta$ gene loci (1.1 and 0.7 Mb), whose T-cells express a diverse human TCR repertoire that compensates for mouse TCR deficiency, immunizing the mouse with a peptide, incubating PBMCs obtained from the transgenic mice with tetramer-phycoerythrin (PE), and isolating the high avidity T-cells by fluorescence activated cell sorting (FACS)–Calibur analysis.

In one aspect, to obtain T-cells expressing TCRs of the present description, nucleic acids encoding TCR-alpha and/or TCR-beta chains of the present description are cloned into expression vectors, such as gamma retrovirus or lentivirus. The recombinant viruses are generated and then tested for functionality, such as antigen specificity and functional avidity. An aliquot of the final product is then used to transduce the target T-cell population (generally purified from patient PBMCs), which is expanded before infusion into the patient.

In another aspect, to obtain T-cells expressing TCRs of the present description, TCR RNAs are synthesized by techniques known in the art, e.g., in vitro transcription

systems. The in vitro-synthesized TCR RNAs are then introduced into primary CD8+ T-cells obtained from healthy donors by electroporation to re-express tumor specific TCR-alpha and/or TCR-beta chains.

To increase the expression, nucleic acids encoding TCRs of the present description may be operably linked to strong promoters, such as retroviral long terminal repeats (LTRs), cytomegalovirus (CMV), murine stem cell virus (MSCV) U3, phosphoglycerate kinase (PGK), β -actin, ubiquitin, and a simian virus 40 (SV40)/CD43 composite promoter, elongation factor (EF)-1a and the spleen focus-forming virus (SFFV) promoter. In a preferred embodiment, the promoter is heterologous to the nucleic acid being expressed.

In addition to strong promoters, TCR expression cassettes of the present description may contain additional elements that can enhance transgene expression, including a central polypurine tract (cPPT), which promotes the nuclear translocation of lentiviral constructs (Follenzi et al., 2000), and the woodchuck hepatitis virus posttranscriptional regulatory element (wPRE), which increases the level of transgene expression by increasing RNA stability (Zufferey et al., 1999).

The alpha and beta chains of a TCR of the present invention may be encoded by nucleic acids located in separate vectors or may be encoded by polynucleotides located in the same vector.

Achieving high-level TCR surface expression requires that both the TCR-alpha and TCR-beta chains of the introduced TCR be transcribed at high levels. To do so, the TCR-alpha and TCR-beta chains of the present description may be cloned into bicistronic constructs in a single vector, which has been shown to be capable of overcoming this obstacle. The use of a viral intraribosomal entry site (IRES) between the TCR-alpha and TCR-beta chains results in the coordinated expression of both chains, because the TCR-alpha and TCR-beta chains are generated from a single transcript

that is broken into two proteins during translation, ensuring that an equal molar ratio of TCR-alpha and TCR-beta chains are produced (Schmitt et al., 2009).

Nucleic acids encoding TCRs of the present description may be codon optimized to increase expression from a host cell. Redundancy in the genetic code allows some amino acids to be encoded by more than one codon, but certain codons are less "optimal" than others because of the relative availability of matching tRNAs as well as other factors (Gustafsson et al., 2004). Modifying the TCR-alpha and TCR-beta gene sequences such that each amino acid is encoded by the optimal codon for mammalian gene expression, as well as eliminating mRNA instability motifs or cryptic splice sites, has been shown to significantly enhance TCR-alpha and TCR-beta gene expression (Scholten et al., 2006).

Furthermore, mispairing between the introduced and endogenous TCR chains may result in the acquisition of specificities that pose a significant risk for autoimmunity. For example, the formation of mixed TCR dimers may reduce the number of CD3 molecules available to form properly paired TCR complexes, and therefore can significantly decrease the functional avidity of the cells expressing the introduced TCR (Kuball et al., 2007).

To reduce mispairing, the C-terminus domain of the introduced TCR chains of the present description may be modified in order to promote interchain affinity, while decreasing the ability of the introduced chains to pair with the endogenous TCR. These strategies may include replacing the human TCR-alpha and TCR-beta C-terminus domains with their murine counterparts (murinized C-terminus domain); generating a second interchain disulfide bond in the C-terminus domain by introducing a second cysteine residue into both the TCR-alpha and TCR-beta chains of the introduced TCR (cysteine modification); swapping interacting residues in the TCR-alpha and TCR-beta chain C-terminus domains ("knob-in-hole"); and fusing the variable domains of the TCR-alpha and TCR-beta chains directly to CD3 ζ (CD3 ζ fusion) (Schmitt et al., 2009).

In an embodiment, a host cell is engineered to express a TCR of the present description. In preferred embodiments, the host cell is a human T-cell or T-cell progenitor. In some embodiments the T-cell or T-cell progenitor is obtained from a cancer patient. In other embodiments the T-cell or T-cell progenitor is obtained from a healthy donor. Host cells of the present description can be allogeneic or autologous with respect to a patient to be treated. In one embodiment, the host is a gamma/delta T-cell transformed to express an alpha/beta TCR.

A "pharmaceutical composition" is a composition suitable for administration to a human being in a medical setting. Preferably, a pharmaceutical composition is sterile and produced according to GMP guidelines.

The pharmaceutical compositions comprise the peptides either in the free form or in the form of a pharmaceutically acceptable salt (see also above). As used herein, "a pharmaceutically acceptable salt" refers to a derivative of the disclosed peptides wherein the peptide is modified by making acid or base salts of the agent. For example, acid salts are prepared from the free base (typically wherein the neutral form of the drug has a neutral -NH_2 group) involving reaction with a suitable acid. Suitable acids for preparing acid salts include both organic acids, e.g., acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methane sulfonic acid, ethane sulfonic acid, p-toluene sulfonic acid, salicylic acid, and the like, as well as inorganic acids, e.g., hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid phosphoric acid and the like. Conversely, preparation of basic salts of acid moieties which may be present on a peptide are prepared using a pharmaceutically acceptable base such as sodium hydroxide, potassium hydroxide, ammonium hydroxide, calcium hydroxide, trimethylamine or the like.

In an especially preferred embodiment, the pharmaceutical compositions comprise the peptides as salts of acetic acid (acetates), trifluoro acetates or hydrochloric acid (chlorides).

Preferably, the medicament of the present invention is an immunotherapeutic such as a vaccine. It may be administered directly into the patient, into the affected organ or systemically i.d., i.m., s.c., i.p. and i.v., or applied *ex vivo* to cells derived from the patient or a human cell line which are subsequently administered to the patient or used *in vitro* to select a subpopulation of immune cells derived from the patient, which are then re-administered to the patient. If the nucleic acid is administered to cells *in vitro*, it may be useful for the cells to be transfected so as to co-express immune-stimulating cytokines, such as interleukin-2. The peptide may be substantially pure or combined with an immune-stimulating adjuvant (see below) or used in combination with immune-stimulatory cytokines, or be administered with a suitable delivery system, for example liposomes. The peptide may also be conjugated to a suitable carrier such as keyhole limpet haemocyanin (KLH) or mannan (see WO 95/18145 and (Longenecker et al., 1993)). The peptide may also be tagged, may be a fusion protein, or may be a hybrid molecule. The peptides whose sequence is given in the present invention are expected to stimulate CD4 or CD8 T cells. However, stimulation of CD8 T cells is more efficient in the presence of help provided by CD4 T-helper cells. Thus, for MHC Class I epitopes that stimulate CD8 T cells the fusion partner or sections of a hybrid molecule suitably provide epitopes which stimulate CD4-positive T cells. CD4- and CD8-stimulating epitopes are well known in the art and include those identified in the present invention.

In one aspect, the vaccine comprises at least one peptide having the amino acid sequence set forth SEQ ID No. 1 to SEQ ID No. 383 and SEQ ID NO: 448 to SEQ ID NO: 499, and at least one additional peptide, preferably two to 50, more preferably two to 25, even more preferably two to 20 and most preferably two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen or eighteen peptides. The peptide(s) may be derived from one or more specific TAAs and may bind to MHC class I molecules.

A further aspect of the invention provides a nucleic acid (for example a polynucleotide) encoding a peptide or peptide variant of the invention. The polynucleotide may be, for

example, DNA, cDNA, PNA, RNA or combinations thereof, either single- and/or double-stranded, or native or stabilized forms of polynucleotides, such as, for example, polynucleotides with a phosphorothioate backbone and it may or may not contain introns so long as it codes for the peptide. Of course, only peptides that contain naturally occurring amino acid residues joined by naturally occurring peptide bonds are encodable by a polynucleotide. A still further aspect of the invention provides an expression vector capable of expressing a polypeptide according to the invention.

A variety of methods have been developed to link polynucleotides, especially DNA, to vectors for example via complementary cohesive termini. For instance, complementary homopolymer tracts can be added to the DNA segment to be inserted to the vector DNA. The vector and DNA segment are then joined by hydrogen bonding between the complementary homopolymeric tails to form recombinant DNA molecules.

Synthetic linkers containing one or more restriction sites provide an alternative method of joining the DNA segment to vectors. Synthetic linkers containing a variety of restriction endonuclease sites are commercially available from a number of sources including International Biotechnologies Inc. New Haven, CN, USA.

A desirable method of modifying the DNA encoding the polypeptide of the invention employs the polymerase chain reaction as disclosed by Saiki RK, et al. (Saiki et al., 1988). This method may be used for introducing the DNA into a suitable vector, for example by engineering in suitable restriction sites, or it may be used to modify the DNA in other useful ways as is known in the art. If viral vectors are used, pox- or adenovirus vectors are preferred.

The DNA (or in the case of retroviral vectors, RNA) may then be expressed in a suitable host to produce a polypeptide comprising the peptide or variant of the invention. Thus, the DNA encoding the peptide or variant of the invention may be used in accordance with known techniques, appropriately modified in view of the teachings contained herein, to construct an expression vector, which is then used to transform an

appropriate host cell for the expression and production of the polypeptide of the invention. Such techniques include those disclosed, for example, in US 4,440,859, 4,530,901, 4,582,800, 4,677,063, 4,678,751, 4,704,362, 4,710,463, 4,757,006, 4,766,075, and 4,810,648.

The DNA (or in the case of retroviral vectors, RNA) encoding the polypeptide constituting the compound of the invention may be joined to a wide variety of other DNA sequences for introduction into an appropriate host. The companion DNA will depend upon the nature of the host, the manner of the introduction of the DNA into the host, and whether episomal maintenance or integration is desired.

Generally, the DNA is inserted into an expression vector, such as a plasmid, in proper orientation and correct reading frame for expression. If necessary, the DNA may be linked to the appropriate transcriptional and translational regulatory control nucleotide sequences recognized by the desired host, although such controls are generally available in the expression vector. The vector is then introduced into the host through standard techniques. Generally, not all of the hosts will be transformed by the vector. Therefore, it will be necessary to select for transformed host cells. One selection technique involves incorporating into the expression vector a DNA sequence, with any necessary control elements, that codes for a selectable trait in the transformed cell, such as antibiotic resistance.

Alternatively, the gene for such selectable trait can be on another vector, which is used to co-transform the desired host cell.

Host cells that have been transformed by the recombinant DNA of the invention are then cultured for a sufficient time and under appropriate conditions known to those skilled in the art in view of the teachings disclosed herein to permit the expression of the polypeptide, which can then be recovered.

Many expression systems are known, including bacteria (for example *E. coli* and *Bacillus subtilis*), yeasts (for example *Saccharomyces cerevisiae*), filamentous fungi (for example *Aspergillus spec.*), plant cells, animal cells and insect cells. Preferably, the system can be mammalian cells such as CHO cells available from the ATCC Cell Biology Collection.

A typical mammalian cell vector plasmid for constitutive expression comprises the CMV or SV40 promoter with a suitable poly A tail and a resistance marker, such as neomycin. One example is pSVL available from Pharmacia, Piscataway, NJ, USA. An example of an inducible mammalian expression vector is pMSG, also available from Pharmacia. Useful yeast plasmid vectors are pRS403-406 and pRS413-416 and are generally available from Stratagene Cloning Systems, La Jolla, CA 92037, USA. Plasmids pRS403, pRS404, pRS405 and pRS406 are Yeast Integrating plasmids (YIps) and incorporate the yeast selectable markers HIS3, TRP1, LEU2 and URA3. Plasmids pRS413-416 are Yeast Centromere plasmids (Ycps). CMV promoter-based vectors (for example from Sigma-Aldrich) provide transient or stable expression, cytoplasmic expression or secretion, and N-terminal or C-terminal tagging in various combinations of FLAG, 3xFLAG, c-myc or MAT. These fusion proteins allow for detection, purification and analysis of recombinant protein. Dual-tagged fusions provide flexibility in detection.

The strong human cytomegalovirus (CMV) promoter regulatory region drives constitutive protein expression levels as high as 1 mg/L in COS cells. For less potent cell lines, protein levels are typically ~0.1 mg/L. The presence of the SV40 replication origin will result in high levels of DNA replication in SV40 replication permissive COS cells. CMV vectors, for example, can contain the pMB1 (derivative of pBR322) origin for replication in bacterial cells, the b-lactamase gene for ampicillin resistance selection in bacteria, hGH polyA, and the f1 origin. Vectors containing the pre-pro-trypsin leader (PPT) sequence can direct the secretion of FLAG fusion proteins into the culture medium for purification using ANTI-FLAG antibodies, resins, and plates. Other vectors and expression systems are well known in the art for use with a variety of host cells.

- 71 -

In another embodiment two or more peptides or peptide variants of the invention are encoded and thus expressed in a successive order (similar to “beads on a string” constructs). In doing so, the peptides or peptide variants may be linked or fused together by stretches of linker amino acids, such as for example LLLLLL, or may be linked without any additional peptide(s) between them. These constructs can also be used for cancer therapy and may induce immune responses both involving MHC I and MHC II.

The present invention also relates to a host cell transformed with a polynucleotide vector construct of the present invention. The host cell can be either prokaryotic or eukaryotic. Bacterial cells may be preferred prokaryotic host cells in some circumstances and typically are a strain of *E. coli* such as, for example, the *E. coli* strains DH5 available from Bethesda Research Laboratories Inc., Bethesda, MD, USA, and RR1 available from the American Type Culture Collection (ATCC) of Rockville, MD, USA (No ATCC 31343). Preferred eukaryotic host cells include yeast, insect and mammalian cells, preferably vertebrate cells such as those from a mouse, rat, monkey or human fibroblastic and colon cell lines. Yeast host cells include YPH499, YPH500 and YPH501, which are generally available from Stratagene Cloning Systems, La Jolla, CA 92037, USA. Preferred mammalian host cells include Chinese hamster ovary (CHO) cells available from the ATCC as CCL61, NIH Swiss mouse embryo cells NIH/3T3 available from the ATCC as CRL 1658, monkey kidney-derived COS-1 cells available from the ATCC as CRL 1650 and 293 cells which are human embryonic kidney cells. Preferred insect cells are Sf9 cells which can be transfected with baculovirus expression vectors. An overview regarding the choice of suitable host cells for expression can be found in, for example, the textbook of Paulina Balbás and Argelia Lorence “Methods in Molecular Biology Recombinant Gene Expression, Reviews and Protocols,” Part One, Second Edition, ISBN 978-1-58829-262-9, and other literature known to the person of skill.

Transformation of appropriate cell hosts with a DNA construct of the present invention is accomplished by well-known methods that typically depend on the type of vector used.

With regard to transformation of prokaryotic host cells, see, for example, Cohen et al. (Cohen et al., 1972) and (Green and Sambrook, 2012) . Transformation of yeast cells is described in Sherman et al. (Sherman et al., 1986) . The method of Beggs (Beggs, 1978) is also useful. With regard to vertebrate cells, reagents useful in transfecting such cells, for example calcium phosphate and DEAE-dextran or liposome formulations, are available from Stratagene Cloning Systems, or Life Technologies Inc., Gaithersburg, MD 20877, USA. Electroporation is also useful for transforming and/or transfecting cells and is well known in the art for transforming yeast cell, bacterial cells, insect cells and vertebrate cells.

Successfully transformed cells, i.e. cells that contain a DNA construct of the present invention, can be identified by well-known techniques such as PCR. Alternatively, the presence of the protein in the supernatant can be detected using antibodies.

It will be appreciated that certain host cells of the invention are useful in the preparation of the peptides of the invention, for example bacterial, yeast and insect cells. However, other host cells may be useful in certain therapeutic methods. For example, antigen-presenting cells, such as dendritic cells, may usefully be used to express the peptides of the invention such that they may be loaded into appropriate MHC molecules. Thus, the current invention provides a host cell comprising a nucleic acid or an expression vector according to the invention.

In a preferred embodiment the host cell is an antigen presenting cell, in particular a dendritic cell or antigen presenting cell. APCs loaded with a recombinant fusion protein containing prostatic acid phosphatase (PAP) were approved by the U.S. Food and Drug Administration (FDA) on April 29, 2010, to treat asymptomatic or minimally symptomatic metastatic HRPc (Sipuleucel-T) (Rini et al., 2006; Small et al., 2006).

A further aspect of the invention provides a method of producing a peptide or its variant, the method comprising culturing a host cell and isolating the peptide from the host cell or its culture medium.

In another embodiment, the peptide, the nucleic acid or the expression vector of the invention are used in medicine. For example, the peptide or its variant may be prepared for intravenous (i.v.) injection, sub-cutaneous (s.c.) injection, intradermal (i.d.) injection, intraperitoneal (i.p.) injection, intramuscular (i.m.) injection. Preferred methods of peptide injection include s.c., i.d., i.p., i.m., and i.v. Preferred methods of DNA injection include i.d., i.m., s.c., i.p. and i.v. Doses of e.g. between 50 µg and 1.5 mg, preferably 125 µg to 500 µg, of peptide or DNA may be given and will depend on the respective peptide or DNA. Dosages of this range were successfully used in previous trials (Walter et al., 2012).

The polynucleotide used for active vaccination may be substantially pure or contained in a suitable vector or delivery system. The nucleic acid may be DNA, cDNA, PNA, RNA or a combination thereof. Methods for designing and introducing such a nucleic acid are well known in the art. An overview is provided by e.g. Teufel et al. (Teufel et al., 2005). Polynucleotide vaccines are easy to prepare, but the mode of action of these vectors in inducing an immune response is not fully understood. Suitable vectors and delivery systems include viral DNA and/or RNA, such as systems based on adenovirus, vaccinia virus, retroviruses, herpes virus, adeno-associated virus or hybrids containing elements of more than one virus. Non-viral delivery systems include cationic lipids and cationic polymers and are well known in the art of DNA delivery. Physical delivery, such as via a "gene-gun" may also be used. The peptide or peptides encoded by the nucleic acid may be a fusion protein, for example with an epitope that stimulates T cells for the respective opposite CDR as noted above.

The medicament of the invention may also include one or more adjuvants. Adjuvants are substances that non-specifically enhance or potentiate the immune response (e.g., immune responses mediated by CD8-positive T cells and helper-T (TH) cells to an antigen and would thus be considered useful in the medicament of the present invention. Suitable adjuvants include, but are not limited to, 1018 ISS, aluminum salts, AMPLIVAX®, AS15, BCG, CP-870,893, CpG7909, CyaA, dSLIM, flagellin or TLR5

ligands derived from flagellin, FLT3 ligand, GM-CSF, IC30, IC31, Imiquimod (ALDARA®), resiquimod, ImuFact IMP321, Interleukins as IL-2, IL-13, IL-21, Interferon-alpha or -beta, or pegylated derivatives thereof, IS Patch, ISS, ISCOMATRIX, ISCOMs, JuvImmune®, LipoVac, MALP2, MF59, monophosphoryl lipid A, Montanide IMS 1312, Montanide ISA 206, Montanide ISA 50V, Montanide ISA-51, water-in-oil and oil-in-water emulsions, OK-432, OM-174, OM-197-MP-EC, ONTAK, OspA, PepTel® vector system, poly(lactid co-glycolid) [PLG]-based and dextran microparticles, talactoferrin SRL172, Virosomes and other Virus-like particles, YF-17D, VEGF trap, R848, beta-glucan, Pam3Cys, Aquila's QS21 stimulon, which is derived from saponin, mycobacterial extracts and synthetic bacterial cell wall mimics, and other proprietary adjuvants such as Ribi's Detox, Quil, or Superfos. Adjuvants such as Freund's or GM-CSF are preferred. Several immunological adjuvants (e.g., MF59) specific for dendritic cells and their preparation have been described previously (Allison and Krummel, 1995). Also cytokines may be used. Several cytokines have been directly linked to influencing dendritic cell migration to lymphoid tissues (e.g., TNF-), accelerating the maturation of dendritic cells into efficient antigen-presenting cells for T-lymphocytes (e.g., GM-CSF, IL-1 and IL-4) (U.S. Pat. No. 5,849,589, specifically incorporated herein by reference in its entirety) and acting as immunoadjuvants (e.g., IL-12, IL-15, IL-23, IL-7, IFN-alpha, IFN-beta) (Gabrilovich et al., 1996).

CpG immunostimulatory oligonucleotides have also been reported to enhance the effects of adjuvants in a vaccine setting. Without being bound by theory, CpG oligonucleotides act by activating the innate (non-adaptive) immune system via Toll-like receptors (TLR), mainly TLR9. CpG triggered TLR9 activation enhances antigen-specific humoral and cellular responses to a wide variety of antigens, including peptide or protein antigens, live or killed viruses, dendritic cell vaccines, autologous cellular vaccines and polysaccharide conjugates in both prophylactic and therapeutic vaccines. More importantly it enhances dendritic cell maturation and differentiation, resulting in enhanced activation of TH1 cells and strong cytotoxic T-lymphocyte (CTL) generation, even in the absence of CD4 T cell help. The TH1 bias induced by TLR9 stimulation is maintained even in the presence of vaccine adjuvants such as alum or incomplete

Freund's adjuvant (IFA) that normally promote a TH2 bias. CpG oligonucleotides show even greater adjuvant activity when formulated or co-administered with other adjuvants or in formulations such as microparticles, nanoparticles, lipid emulsions or similar formulations, which are especially necessary for inducing a strong response when the antigen is relatively weak. They also accelerate the immune response and enable the antigen doses to be reduced by approximately two orders of magnitude, with comparable antibody responses to the full-dose vaccine without CpG in some experiments (Krieg, 2006). US 6,406,705 B1 describes the combined use of CpG oligonucleotides, non-nucleic acid adjuvants and an antigen to induce an antigen-specific immune response. A CpG TLR9 antagonist is dSLIM (double Stem Loop Immunomodulator) by Mologen (Berlin, Germany) which is a preferred component of the pharmaceutical composition of the present invention. Other TLR binding molecules such as RNA binding TLR 7, TLR 8 and/or TLR 9 may also be used.

Other examples for useful adjuvants include, but are not limited to chemically modified CpGs (e.g. CpR, Idera), dsRNA analogues such as Poly(I:C) and derivatives thereof (e.g. AmpliGen®, Hiltonol®, poly-(ICLC), poly(IC-R), poly(I:C12U), non-CpG bacterial DNA or RNA as well as immunoactive small molecules and antibodies such as cyclophosphamide, sunitinib, immune checkpoint inhibitors including ipilimumab, nivolumab, pembrolizumab, atezolizumab, avelumab, durvalumab, and cemiplimab, Bevacizumab®, celebrex, NCX-4016, sildenafil, tadalafil, vardenafil, sorafenib, temozolomide, temsirolimus, XL-999, CP-547632, pazopanib, VEGF Trap, ZD2171, AZD2171, anti-CTLA4, other antibodies targeting key structures of the immune system (e.g. anti-CD40, anti-TGFbeta, anti-TNFalpha receptor) and SC58175, which may act therapeutically and/or as an adjuvant. The amounts and concentrations of adjuvants and additives useful in the context of the present invention can readily be determined by the skilled artisan without undue experimentation.

Preferred adjuvants are anti-CD40, imiquimod, resiquimod, GM-CSF, cyclophosphamide, sunitinib, bevacizumab, interferon-alpha, interferon-beta, CpG oligonucleotides and derivatives, poly-(I:C) and derivatives, RNA, sildenafil, particulate

formulations with poly(lactide co-glycolide) (PLG), virosomes, atezolizumab, interleukin (IL)-1, IL-2, IL-4, IL-7, IL-12, IL-13, IL-15, IL-21, and IL-23.

In a preferred embodiment, the pharmaceutical composition according to the invention the adjuvant is selected from the group consisting of colony-stimulating factors, such as Granulocyte Macrophage Colony Stimulating Factor (GM-CSF, sargramostim), cyclophosphamide, imiquimod, resiquimod, and interferon-alpha.

In a preferred embodiment, the pharmaceutical composition according to the invention the adjuvant is selected from the group consisting of colony-stimulating factors, such as Granulocyte Macrophage Colony Stimulating Factor (GM-CSF, sargramostim), cyclophosphamide, imiquimod and resiquimod. In a preferred embodiment of the pharmaceutical composition according to the invention, the adjuvant is cyclophosphamide, imiquimod or resiquimod. Even more preferred adjuvants are Montanide IMS 1312, Montanide ISA 206, Montanide ISA 50V, Montanide ISA-51, poly-ICLC (Hiltonol®) and anti-CD40 mAB, or combinations thereof.

This composition is used for parenteral administration, such as subcutaneous, intradermal, intramuscular or oral administration. For this, the peptides and optionally other molecules are dissolved or suspended in a pharmaceutically acceptable, preferably aqueous carrier. In addition, the composition can contain excipients, such as buffers, binding agents, blasting agents, diluents, flavors, lubricants, etc. The peptides can also be administered together with immune stimulating substances, such as cytokines. An extensive listing of excipients that can be used in such a composition, can be, for example, taken from A. Kibbe, Handbook of Pharmaceutical Excipients (Kibbe, 2000). The composition can be used for a prevention, prophylaxis and/or therapy of adenomatous or cancerous diseases. Exemplary formulations can be found in, for example, EP2112253.

It is important to realize that the immune response triggered by the vaccine according to the invention attacks the cancer in different cell-stages and different stages of

development. Furthermore different cancer associated signaling pathways are attacked. This is an advantage over vaccines that address only one or few targets, which may cause the tumor to easily adapt to the attack (tumor escape). Furthermore, not all individual tumors express the same pattern of antigens. Therefore, a combination of several tumor-associated peptides ensures that every single tumor bears at least some of the targets. The composition is designed in such a way that each tumor is expected to express several of the antigens and cover several independent pathways necessary for tumor growth and maintenance. Thus, the vaccine can easily be used "off-the-shelf" for a larger patient population. This means that a pre-selection of patients to be treated with the vaccine can be restricted to HLA typing, does not require any additional biomarker assessments for antigen expression, but it is still ensured that several targets are simultaneously attacked by the induced immune response, which is important for efficacy (Banchereau et al., 2001; Walter et al., 2012).

As used herein, the term "scaffold" refers to a molecule that specifically binds to an (e.g. antigenic) determinant. In one embodiment, a scaffold is able to direct the entity to which it is attached (e.g. a (second) antigen binding moiety) to a target site, for example to a specific type of tumor cell or tumor stroma bearing the antigenic determinant (e.g. the complex of a peptide with MHC, according to the application at hand). In another embodiment a scaffold is able to activate signaling through its target antigen, for example a T cell receptor complex antigen. Scaffolds include but are not limited to antibodies and fragments thereof, antigen binding domains of an antibody, comprising an antibody heavy chain variable region and an antibody light chain variable region, binding proteins comprising at least one ankyrin repeat motif and single domain antigen binding (SDAB) molecules, aptamers, (soluble) TCRs and (modified) cells such as allogenic or autologous T cells. To assess whether a molecule is a scaffold binding to a target, binding assays can be performed.

"Specific" binding means that the scaffold binds the peptide-MHC-complex of interest better than other naturally occurring peptide-MHC-complexes, to an extent that a scaffold armed with an active molecule that is able to kill a cell bearing the specific

- 78 -

target is not able to kill another cell without the specific target but presenting other peptide-MHC complex(es). Binding to other peptide-MHC complexes is irrelevant if the peptide of the cross-reactive peptide-MHC is not naturally occurring, i.e. not derived from the human HLA-peptidome. Tests to assess target cell killing are well known in the art. They should be performed using target cells (primary cells or cell lines) with unaltered peptide-MHC presentation, or cells loaded with peptides such that naturally occurring peptide-MHC levels are reached.

Each scaffold can comprise a labelling which provides that the bound scaffold can be detected by determining the presence or absence of a signal provided by the label. For example, the scaffold can be labelled with a fluorescent dye or any other applicable cellular marker molecule. Such marker molecules are well known in the art. For example a fluorescence-labelling, for example provided by a fluorescence dye, can provide a visualization of the bound aptamer by fluorescence or laser scanning microscopy or flow cytometry.

Each scaffold can be conjugated with a second active molecule such as for example IL-21, anti-CD3, and anti-CD28.

For further information on polypeptide scaffolds see for example the background section of WO 2014/071978A1 and the references cited therein.

The present invention further relates to aptamers. Aptamers (see for example WO 2014/191359 and the literature as cited therein) are short single-stranded nucleic acid molecules, which can fold into defined three-dimensional structures and recognize specific target structures. They have appeared to be suitable alternatives for developing targeted therapies. Aptamers have been shown to selectively bind to a variety of complex targets with high affinity and specificity.

Aptamers recognizing cell surface located molecules have been identified within the past decade and provide means for developing diagnostic and therapeutic approaches.

Since aptamers have been shown to possess almost no toxicity and immunogenicity they are promising candidates for biomedical applications. Indeed aptamers, for example prostate-specific membrane-antigen recognizing aptamers, have been successfully employed for targeted therapies and shown to be functional in xenograft *in vivo* models. Furthermore, aptamers recognizing specific tumor cell lines have been identified.

DNA aptamers can be selected to reveal broad-spectrum recognition properties for various cancer cells, and particularly those derived from solid tumors, while non-tumorigenic and primary healthy cells are not recognized. If the identified aptamers recognize not only a specific tumor sub-type but rather interact with a series of tumors, this renders the aptamers applicable as so-called broad-spectrum diagnostics and therapeutics.

Further, investigation of cell-binding behavior with flow cytometry showed that the aptamers revealed very good apparent affinities that are within the nanomolar range.

Aptamers are useful for diagnostic and therapeutic purposes. Further, it could be shown that some of the aptamers are taken up by tumor cells and thus can function as molecular vehicles for the targeted delivery of anti-cancer agents such as siRNA into tumor cells.

Aptamers can be selected against complex targets such as cells and tissues and complexes of the peptides comprising, preferably consisting of, a sequence according to any of SEQ ID NO 1 to SEQ ID NO 383 and SEQ ID NO: 448 to SEQ ID NO: 499, according to the invention at hand with the MHC molecule, using the cell-SELEX (Systematic Evolution of Ligands by Exponential enrichment) technique.

The peptides of the present invention can be used to generate and develop specific antibodies against MHC/peptide complexes. These can be used for therapy, targeting toxins or radioactive substances to the diseased tissue. Another use of these antibodies

can be targeting radionuclides to the diseased tissue for imaging purposes such as PET. This use can help to detect small metastases or to determine the size and precise localization of diseased tissues.

Therefore, it is a further aspect of the invention to provide a method for producing a recombinant antibody specifically binding to a human major histocompatibility complex (MHC) class I or II being complexed with a HLA-restricted antigen (preferably a peptide according to the present invention), the method comprising: immunizing a genetically engineered non-human mammal comprising cells expressing said human major histocompatibility complex (MHC) class I or II with a soluble form of a MHC class I or II molecule being complexed with said HLA-restricted antigen; isolating mRNA molecules from antibody producing cells of said non-human mammal; producing a phage display library displaying protein molecules encoded by said mRNA molecules; and isolating at least one phage from said phage display library, said at least one phage displaying said antibody specifically binding to said human major histocompatibility complex (MHC) class I or II being complexed with said HLA-restricted antigen.

It is thus a further aspect of the invention to provide an antibody that specifically binds to a human major histocompatibility complex (MHC) class I or II being complexed with an HLA-restricted antigen, wherein the antibody preferably is a polyclonal antibody, monoclonal antibody, bi-specific antibody and/or a chimeric antibody.

Respective methods for producing such antibodies and single chain class I major histocompatibility complexes, as well as other tools for the production of these antibodies are disclosed in WO 03/068201, WO 2004/084798, WO 01/72768, WO 03/070752, and in publications (Cohen et al., 2003a; Cohen et al., 2003b; Denkberg et al., 2003), which for the purposes of the present invention are all explicitly incorporated by reference in their entireties.

- 81 -

Preferably, the antibody is binding with a binding affinity of below 20 nanomolar, preferably of below 10 nanomolar, to the complex, which is also regarded as "specific" in the context of the present invention.

The present invention relates to a peptide comprising a sequence that is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499, or a variant thereof which is at least 88% homologous (preferably identical) to SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499 or a variant thereof that induces T cells cross-reacting with said peptide, wherein said peptide is not the underlying full-length polypeptide.

The present invention further relates to a peptide comprising a sequence that is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499 or a variant thereof which is at least 88% homologous (preferably identical) to SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499, wherein said peptide or variant has an overall length of between 8 and 100, preferably between 8 and 30, and most preferred between 8 and 14 amino acids.

The present invention further relates to the peptides according to the invention that have the ability to bind to a molecule of the human major histocompatibility complex (MHC) class-I or -II.

The present invention further relates to the peptides according to the invention wherein the peptide consists or consists essentially of an amino acid sequence according to SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499.

The present invention further relates to the peptides according to the invention, wherein the peptide is (chemically) modified and/or includes non-peptide bonds.

The present invention further relates to the peptides according to the invention, wherein the peptide is part of a fusion protein, in particular comprising N-terminal amino acids of

- 82 -

the HLA-DR antigen-associated invariant chain (Ii), or wherein the peptide is fused to (or into) an antibody, such as, for example, an antibody that is specific for dendritic cells.

The present invention further relates to a nucleic acid, encoding the peptides according to the invention, provided that the peptide is not the complete (full) human protein.

The present invention further relates to the nucleic acid according to the invention that is DNA, cDNA, PNA, RNA or combinations thereof.

The present invention further relates to an expression vector capable of expressing a nucleic acid according to the present invention.

The present invention further relates to a peptide according to the present invention, a nucleic acid according to the present invention or an expression vector according to the present invention for use in medicine, in particular in the treatment of acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

The present invention further relates to a host cell comprising a nucleic acid according to the invention or an expression vector according to the invention.

The present invention further relates to the host cell according to the present invention that is an antigen presenting cell, and preferably a dendritic cell.

The present invention further relates to a method of producing a peptide according to the present invention, said method comprising culturing the host cell according to the present invention, and isolating the peptide from said host cell or its culture medium.

The present invention further relates to the method according to the present invention, where-in the antigen is loaded onto class I or II MHC molecules expressed on the surface of a suitable antigen-presenting cell by contacting a sufficient amount of the antigen with an antigen-presenting cell.

The present invention further relates to the method according to the invention, wherein the antigen-presenting cell comprises an expression vector capable of expressing said peptide containing SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499 or said variant amino acid sequence.

The present invention further relates to activated T cells, produced by the method according to the present invention, wherein said T cells selectively recognizes a cell which aberrantly expresses a polypeptide comprising an amino acid sequence according to the present invention.

The present invention further relates to a method of killing target cells in a patient which target cells aberrantly express a polypeptide comprising any amino acid sequence according to the present invention, the method comprising administering to the patient an effective number of T cells as according to the present invention.

The present invention further relates to the use of any peptide described, a nucleic acid according to the present invention, an expression vector according to the present invention, a cell according to the present invention, or an activated cytotoxic T lymphocyte according to the present invention as a medicament or in the manufacture of a medicament. The present invention further relates to a use according to the present invention, wherein the medicament is active against cancer.

The present invention further relates to a use according to the invention, wherein the medicament is a vaccine. The present invention further relates to a use according to the invention, wherein the medicament is active against cancer.

The present invention further relates to a use according to the invention, wherein said cancer cells are acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer cells or other solid or hematological tumor cells such as acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

The present invention further relates to particular marker proteins and biomarkers based on the peptides according to the present invention, herein called "targets" that can be used in the diagnosis and/or prognosis of acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and

endometrial cancer. The present invention also relates to the use of these novel targets for cancer treatment.

The term "antibody" or "antibodies" is used herein in a broad sense and includes both polyclonal and monoclonal antibodies. In addition to intact or "full" immunoglobulin molecules, also included in the term "antibodies" are fragments (e.g. CDRs, Fv, Fab and Fc fragments) or polymers of those immunoglobulin molecules and humanized versions of immunoglobulin molecules, as long as they exhibit any of the desired properties (e.g., specific binding of an acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer marker (poly)peptide, delivery of a toxin to an acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer cell expressing a cancer marker gene at an increased level, and/or inhibiting the activity of an acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer marker polypeptide) according to the invention.

Whenever possible, the antibodies of the invention may be purchased from commercial sources. The antibodies of the invention may also be generated using well-known methods. The skilled artisan will understand that either full length acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer marker polypeptides or fragments thereof may be used to generate the antibodies of the invention. A polypeptide to be used for generating an antibody of the invention may be partially or fully purified from a natural source or may be produced using recombinant DNA techniques.

For example, a cDNA encoding a peptide according to the present invention, such as a peptide according to SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499 polypeptide, or a variant or fragment thereof, can be expressed in prokaryotic cells (e.g., bacteria) or eukaryotic cells (e.g., yeast, insect, or mammalian cells), after which the recombinant protein can be purified and used to generate a monoclonal or polyclonal antibody preparation that specifically bind the acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer marker polypeptide used to generate the antibody according to the invention.

One of skill in the art will realize that the generation of two or more different sets of monoclonal or polyclonal antibodies maximizes the likelihood of obtaining an antibody with the specificity and affinity required for its intended use (e.g., ELISA, immunohistochemistry, *in vivo* imaging, immunotoxin therapy). The antibodies are tested for their desired activity by known methods, in accordance with the purpose for which the antibodies are to be used (e.g., ELISA, immunohistochemistry, immunotherapy, etc.; for further guidance on the generation and testing of antibodies, see, e.g., Greenfield, 2014 (Greenfield, 2014)). For example, the antibodies may be tested in ELISA assays or, Western blots, immunohistochemical staining of formalin-fixed cancers or frozen tissue sections. After their initial *in vitro* characterization, antibodies intended for therapeutic or *in vivo* diagnostic use are tested according to known clinical testing methods.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a substantially homogeneous population of antibodies, i.e.; the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. The monoclonal antibodies herein specifically include "chimeric" antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired antagonistic activity (US 4,816,567, which is hereby incorporated in its entirety).

Monoclonal antibodies of the invention may be prepared using hybridoma methods. In a hybridoma method, a mouse or other appropriate host animal is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized *in vitro*.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in US 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies).

In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly Fab fragments, can be accomplished using routine techniques known in the art. For instance, digestion can be performed using papain. Examples of papain digestion are described in WO 94/29348 and US 4,342,566. Papain digestion of antibodies typically produces two identical antigen binding fragments, called Fab fragments, each with a single antigen binding site, and a residual Fc fragment. Pepsin treatment yields a F(ab')₂ fragment and a pFc' fragment.

The antibody fragments, whether attached to other sequences or not, can also include insertions, deletions, substitutions, or other selected modifications of particular regions or specific amino acids residues, provided the activity of the fragment is not significantly altered or impaired compared to the non-modified antibody or antibody fragment. These modifications can provide for some additional property, such as to remove/add amino acids capable of disulfide bonding, to increase its bio-longevity, to alter its secretory characteristics, etc. In any case, the antibody fragment must possess a bioactive property, such as binding activity, regulation of binding at the binding domain, etc. Functional or active regions of the antibody may be identified by mutagenesis of a specific region of the protein, followed by expression and testing of the expressed polypeptide. Such methods are readily apparent to a skilled practitioner in the art and can include site-specific mutagenesis of the nucleic acid encoding the antibody fragment.

The antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric

immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab' or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin.

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (US 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Transgenic animals (e.g., mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin

- 90 -

production can be employed. For example, it has been described that the homozygous deletion of the antibody heavy chain joining region gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. Human antibodies can also be produced in phage display libraries.

Antibodies of the invention are preferably administered to a subject in a pharmaceutically acceptable carrier. Typically, an appropriate amount of a pharmaceutically-acceptable salt is used in the formulation to render the formulation isotonic. Examples of the pharmaceutically-acceptable carrier include saline, Ringer's solution and dextrose solution. The pH of the solution is preferably from about 5 to about 8, and more preferably from about 7 to about 7.5. Further carriers include sustained release preparations such as semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, liposomes or microparticles. It will be apparent to those persons skilled in the art that certain carriers may be more preferable depending upon, for instance, the route of administration and concentration of antibody being administered.

The antibodies can be administered to the subject, patient, or cell by injection (e.g., intravenous, intraperitoneal, subcutaneous, intramuscular), or by other methods such as infusion that ensure its delivery to the bloodstream in an effective form. The antibodies may also be administered by intratumoral or peritumoral routes, to exert local as well as systemic therapeutic effects. Local or intravenous injection is preferred.

Effective dosages and schedules for administering the antibodies may be determined empirically, and making such determinations is within the skill in the art. Those skilled in the art will understand that the dosage of antibodies that must be administered will vary depending on, for example, the subject that will receive the antibody, the route of administration, the particular type of antibody used, and other drugs being administered. A typical daily dosage of the antibody used alone might range from about 1 µg/kg to up

to 100 mg/kg of body weight or more per day, depending on the factors mentioned above. Following administration of an antibody, preferably for treating acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer, the efficacy of the therapeutic antibody can be assessed in various ways well known to the skilled practitioner. For instance, the size, number, and/or distribution of cancer in a subject receiving treatment may be monitored using standard tumor imaging techniques. A therapeutically-administered antibody that arrests tumor growth, results in tumor shrinkage, and/or prevents the development of new tumors, compared to the disease course that would occur in the absence of antibody administration, is an efficacious antibody for treatment of cancer.

It is a further aspect of the invention to provide a method for producing a soluble T-cell receptor (sTCR) recognizing a specific peptide-MHC complex. Such soluble T-cell receptors can be generated from specific T-cell clones, and their affinity can be increased by mutagenesis targeting the complementarity-determining regions. For the purpose of T-cell receptor selection, phage display can be used (US 2010/0113300, (Liddy et al., 2012)). For the purpose of stabilization of T-cell receptors during phage display and in case of practical use as drug, alpha and beta chain can be linked e.g. by non-native disulfide bonds, other covalent bonds (single-chain T-cell receptor), or by dimerization domains (Boulter et al., 2003; Card et al., 2004; Willcox et al., 1999). The T-cell receptor can be linked to toxins, drugs, cytokines (see, for example, US 2013/0115191), and domains recruiting effector cells such as an anti-CD3 domain, etc., in order to execute particular functions on target cells. Moreover, it could be expressed in T cells used for adoptive transfer. Further information can be found in WO 2004/033685A1 and WO 2004/074322A1. A combination of sTCRs is described in WO

2012/056407A1. Further methods for the production are disclosed in WO 2013/057586A1.

In addition, the peptides and/or the TCRs or antibodies or other binding molecules of the present invention can be used to verify a pathologist's diagnosis of a cancer based on a biopsied sample.

The antibodies or TCRs may also be used for *in vivo* diagnostic assays. Generally, the antibody is labeled with a radionucleotide (such as ^{111}In , ^{99}Tc , ^{14}C , ^{131}I , ^3H , ^{32}P or ^{35}S) so that the tumor can be localized using immunoscintigraphy. In one embodiment, antibodies or fragments thereof bind to the extracellular domains of two or more targets of a protein selected from the group consisting of the above-mentioned proteins, and the affinity value (K_d) is less than $1 \times 10\mu\text{M}$.

Antibodies for diagnostic use may be labeled with probes suitable for detection by various imaging methods. Methods for detection of probes include, but are not limited to, fluorescence, light, confocal and electron microscopy; magnetic resonance imaging and spectroscopy; fluoroscopy, computed tomography and positron emission tomography. Suitable probes include, but are not limited to, fluorescein, rhodamine, eosin and other fluorophores, radioisotopes, gold, gadolinium and other lanthanides, paramagnetic iron, fluorine-18 and other positron-emitting radionuclides. Additionally, probes may be bi- or multi-functional and be detectable by more than one of the methods listed. These antibodies may be directly or indirectly labeled with said probes. Attachment of probes to the antibodies includes covalent attachment of the probe, incorporation of the probe into the antibody, and the covalent attachment of a chelating compound for binding of probe, amongst others well recognized in the art. For immunohistochemistry, the disease tissue sample may be fresh or frozen or may be embedded in paraffin and fixed with a preservative such as formalin. The fixed or embedded section contains the sample are contacted with a labeled primary antibody and secondary antibody, wherein the antibody is used to detect the expression of the proteins *in situ*.

Another aspect of the present invention includes an *in vitro* method for producing activated T cells, the method comprising contacting *in vitro* T cells with antigen loaded human MHC molecules expressed on the surface of a suitable antigen-presenting cell for a period of time sufficient to activate the T cell in an antigen specific manner, wherein the antigen is a peptide according to the invention. Preferably a sufficient amount of the antigen is used with an antigen-presenting cell.

Preferably the mammalian cell lacks or has a reduced level or function of the TAP peptide transporter. Suitable cells that lack the TAP peptide transporter include T2, RMA-S and Drosophila cells. TAP is the transporter associated with antigen processing.

The human peptide loading deficient cell line T2 is available from the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, USA under Catalogue No CRL 1992; the Drosophila cell line Schneider line 2 is available from the ATCC under Catalogue No CRL 19863; the mouse RMA-S cell line is described in Ljunggren et al. (Ljunggren and Karre, 1985).

Preferably, before transfection the host cell expresses substantially no MHC class I molecules. It is also preferred that the stimulator cell expresses a molecule important for providing a co-stimulatory signal for T-cells such as any of B7.1, B7.2, ICAM-1 and LFA 3. The nucleic acid sequences of numerous MHC class I molecules and of the co-stimulator molecules are publicly available from the GenBank and EMBL databases.

In case of an MHC class I epitope being used as an antigen; the T cells are CD8-positive T cells.

If an antigen-presenting cell is transfected to express such an epitope, preferably the cell comprises an expression vector capable of expressing a peptide containing SEQ ID NO: 1 to SEQ ID NO: 383 and SEQ ID NO: 448 to SEQ ID NO: 499, or a variant amino acid sequence thereof.

A number of other methods may be used for generating T cells in vitro. For example, autologous tumor-infiltrating lymphocytes can be used in the generation of CTL. Plebanski et al. (Plebanski et al., 1995) made use of autologous peripheral blood lymphocytes (PLBs) in the preparation of T cells. Furthermore, the production of autologous T cells by pulsing dendritic cells with peptide or polypeptide, or via infection with recombinant virus is possible. Also, B cells can be used in the production of autologous T cells. In addition, macrophages pulsed with peptide or polypeptide, or infected with recombinant virus, may be used in the preparation of autologous T cells. S. Walter et al. (Walter et al., 2003) describe the in vitro priming of T cells by using artificial antigen presenting cells (aAPCs), which is also a suitable way for generating T cells against the peptide of choice. In the present invention, aAPCs were generated by the coupling of preformed MHC:peptide complexes to the surface of polystyrene particles (microbeads) by biotin:streptavidin biochemistry. This system permits the exact control of the MHC density on aAPCs, which allows to selectively elicit high- or low-avidity antigen-specific T cell responses with high efficiency from blood samples. Apart from MHC:peptide complexes, aAPCs should carry other proteins with co-stimulatory activity like anti-CD28 antibodies coupled to their surface. Furthermore such aAPC-based systems often require the addition of appropriate soluble factors, e. g. cytokines, like interleukin-12.

Allogeneic cells may also be used in the preparation of T cells and a method is described in detail in WO 97/26328, incorporated herein by reference. For example, in addition to *Drosophila* cells and T2 cells, other cells may be used to present antigens such as CHO cells, baculovirus-infected insect cells, bacteria, yeast, and vaccinia-infected target cells. In addition plant viruses may be used (see, for example, Porta et al. (Porta et al., 1994) which describes the development of cowpea mosaic virus as a high-yielding system for the presentation of foreign peptides.

The activated T cells that are directed against the peptides of the invention are useful in therapy. Thus, a further aspect of the invention provides activated T cells obtainable by the foregoing methods of the invention.

Activated T cells, which are produced by the above method, will selectively recognize a cell that aberrantly expresses a polypeptide that comprises an amino acid sequence of SEQ ID NO: 1 to SEQ ID NO 383 and SEQ ID NO: 448 to SEQ ID NO: 499.

Preferably, the T cell recognizes the cell by interacting through its TCR with the HLA/peptide-complex (for example, binding). The T cells are useful in a method of killing target cells in a patient whose target cells aberrantly express a polypeptide comprising an amino acid sequence of the invention wherein the patient is administered an effective number of the activated T cells. The T cells that are administered to the patient may be derived from the patient and activated as described above (i.e. they are autologous T cells). Alternatively, the T cells are not from the patient but are from another individual. Of course, it is preferred if the individual is a healthy individual. By "healthy individual" the inventors mean that the individual is generally in good health, preferably has a competent immune system and, more preferably, is not suffering from any disease that can be readily tested for and detected.

In vivo, the target cells for the CD8-positive T cells according to the present invention can be cells of the tumor (which sometimes express MHC class II) and/or stromal cells surrounding the tumor (tumor cells) (which sometimes also express MHC class II; (Dengjel et al., 2006)).

The T cells of the present invention may be used as active ingredients of a therapeutic composition. Thus, the invention also provides a method of killing target cells in a patient whose target cells aberrantly express a polypeptide comprising an amino acid sequence of the invention, the method comprising administering to the patient an effective number of T cells as defined above.

By "aberrantly expressed" the inventors also mean that the polypeptide is over-expressed compared to levels of expression in normal tissues or that the gene is silent in the tissue from which the tumor is derived but, in the tumor, it is expressed. By "over-expressed" the inventors mean that the polypeptide is present at a level at least 2-fold of that present in normal tissue; preferably at least 3-fold, and more preferably at least 5-fold or 10-fold the level present in normal tissue.

T cells may be obtained by methods known in the art, e.g. those described above.

Protocols for this so-called adoptive transfer of T cells are well known in the art. Reviews can be found in: Gattioni et al. and Morgan et al. (Gattinoni et al., 2006; Morgan et al., 2006).

Another aspect of the present invention includes the use of the peptides complexed with MHC to generate a T-cell receptor whose nucleic acid is cloned and is introduced into a host cell, preferably a T cell. This engineered T cell can then be transferred to a patient for therapy of cancer.

Any molecule of the invention, i.e. the peptide, nucleic acid, antibody, expression vector, cell, activated T cell, T-cell receptor or the nucleic acid encoding it, is useful for the treatment of disorders, characterized by cells escaping an immune response. Therefore any molecule of the present invention may be used as medicament or in the manufacture of a medicament. The molecule may be used by itself or combined with other molecule(s) of the invention or (a) known molecule(s).

The present invention is further directed at a kit comprising:

- (a) a container containing a pharmaceutical composition as described above, in solution or in lyophilized form;
- (b) optionally a second container containing a diluent or reconstituting solution for the lyophilized formulation; and

(c) optionally, instructions for (i) use of the solution or (ii) reconstitution and/or use of the lyophilized formulation.

The kit may further comprise one or more of (iii) a buffer, (iv) a diluent, (v) a filter, (vi) a needle, or (v) a syringe. The container is preferably a bottle, a vial, a syringe or test tube; and it may be a multi-use container. The pharmaceutical composition is preferably lyophilized.

Kits of the present invention preferably comprise a lyophilized formulation of the present invention in a suitable container and instructions for its reconstitution and/or use. Suitable containers include, for example, bottles, vials (e.g. dual chamber vials), syringes (such as dual chamber syringes) and test tubes. The container may be formed from a variety of materials such as glass or plastic. Preferably the kit and/or container contain/s instructions on or associated with the container that indicates directions for reconstitution and/or use. For example, the label may indicate that the lyophilized formulation is to be reconstituted to peptide concentrations as described above. The label may further indicate that the formulation is useful or intended for subcutaneous administration.

The container holding the formulation may be a multi-use vial, which allows for repeat administrations (e.g., from 2-6 administrations) of the reconstituted formulation. The kit may further comprise a second container comprising a suitable diluent (e.g., sodium bicarbonate solution).

Upon mixing of the diluent and the lyophilized formulation, the final peptide concentration in the reconstituted formulation is preferably at least 0.15 mg/mL/peptide (=75 µg) and preferably not more than 3 mg/mL/peptide (=1500 µg). The kit may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use.

Kits of the present invention may have a single container that contains the formulation of the pharmaceutical compositions according to the present invention with or without other components (e.g., other compounds or pharmaceutical compositions of these other compounds) or may have distinct container for each component.

Preferably, kits of the invention include a formulation of the invention packaged for use in combination with the co-administration of a second compound (such as adjuvants (e.g. GM-CSF), a chemotherapeutic agent, a natural product, a hormone or antagonist, an anti-angiogenesis agent or inhibitor, an apoptosis-inducing agent or a chelator) or a pharmaceutical composition thereof. The components of the kit may be pre-complexed, or each component may be in a separate distinct container prior to administration to a patient. The components of the kit may be provided in one or more liquid solutions, preferably, an aqueous solution, more preferably, a sterile aqueous solution. The components of the kit may also be provided as solids, which may be converted into liquids by addition of suitable solvents, which are preferably provided in another distinct container.

The container of a therapeutic kit may be a vial, test tube, flask, bottle, syringe, or any other means of enclosing a solid or liquid. Usually, when there is more than one component, the kit will contain a second vial or other container, which allows for separate dosing. The kit may also contain another container for a pharmaceutically acceptable liquid. Preferably, a therapeutic kit will contain an apparatus (e.g., one or more needles, syringes, eye droppers, pipette, etc.), which enables administration of the agents of the invention that are components of the present kit.

The present formulation is one that is suitable for administration of the peptides by any acceptable route such as oral (enteral), nasal, ophthal, subcutaneous, intradermal, intramuscular, intravenous or transdermal. Preferably, the administration is s.c., and most preferably i.d. administration may be by infusion pump.

Since the peptides of the invention were isolated from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer, the medicament of the invention is preferably used to treat acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

The present invention further relates to a method for producing a personalized pharmaceutical for an individual patient comprising manufacturing a pharmaceutical composition comprising at least one peptide selected from a warehouse of pre-screened TUMAPs, wherein the at least one peptide used in the pharmaceutical composition is selected for suitability in the individual patient. In one embodiment, the pharmaceutical composition is a vaccine. The method could also be adapted to produce T cell clones for down-stream applications, such as TCR isolations, or soluble antibodies, and other treatment options.

A "personalized pharmaceutical" shall mean specifically tailored therapies for one individual patient that will only be used for therapy in such individual patient, including actively personalized cancer vaccines and adoptive cellular therapies using autologous patient tissue.

As used herein, the term “warehouse” shall refer to a group or set of peptides that have been pre-screened for immunogenicity and/or over-presentation in a particular tumor type. The term “warehouse” is not intended to imply that the particular peptides included in the vaccine have been pre-manufactured and stored in a physical facility, although that possibility is contemplated. It is expressly contemplated that the peptides may be manufactured *de novo* for each individualized vaccine produced or may be pre-manufactured and stored. The warehouse (e.g. in the form of a database) is composed of tumor-associated peptides which were highly overexpressed in the tumor tissue of acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer patients with various HLA-A HLA-B and HLA-C alleles. It may contain MHC class I and MHC class II peptides or elongated MHC class I peptides. In addition to the tumor associated peptides collected from several acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer tissues, the warehouse may contain HLA-A*02, HLA-A*01, HLA-A*03, HLA-A*24, HLA-B*07, HLA-B*08 and HLA-B*44 marker peptides. These peptides allow comparison of the magnitude of T-cell immunity induced by TUMAPS in a quantitative manner and hence allow important conclusion to be drawn on the capacity of the vaccine to elicit anti-tumor responses. Secondly, they function as important positive control peptides derived from a “non-self” antigen in the case that any vaccine-induced T-cell responses to TUMAPs derived from “self” antigens in a patient are not

observed. And thirdly, it may allow conclusions to be drawn, regarding the status of immunocompetence of the patient.

TUMAPs for the warehouse are identified by using an integrated functional genomics approach combining gene expression analysis, mass spectrometry, and T-cell immunology (XPresident®). The approach assures that only TUMAPs truly present on a high percentage of tumors but not or only minimally expressed on normal tissue, are chosen for further analysis. For initial peptide selection, acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer samples from patients and blood from healthy donors were analyzed in a stepwise approach:

1. HLA ligands from the malignant material were identified by mass spectrometry
2. Genome-wide messenger ribonucleic acid (mRNA) expression analysis was used to identify genes over-expressed in the malignant tissue (acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer) compared with a range of normal organs and tissues
3. Identified HLA ligands were compared to gene expression data. Peptides over-presented or selectively presented on tumor tissue, preferably encoded by selectively expressed or over-expressed genes as detected in step 2 were considered suitable TUMAP candidates for a multi-peptide vaccine.

- 102 -

4. Literature research was performed in order to identify additional evidence supporting the relevance of the identified peptides as TUMAPs
5. The relevance of over-expression at the mRNA level was confirmed by redetection of selected TUMAPs from step 3 on tumor tissue and lack of (or infrequent) detection on healthy tissues.
6. In order to assess, whether an induction of *in vivo* T-cell responses by the selected peptides may be feasible, *in vitro* immunogenicity assays were performed using human T cells from healthy donors as well as from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer patients.

In an aspect, the peptides are pre-screened for immunogenicity before being included in the warehouse. By way of example, and not limitation, the immunogenicity of the peptides included in the warehouse is determined by a method comprising *in vitro* T-cell priming through repeated stimulations of CD8⁺ T cells from healthy donors with artificial antigen presenting cells loaded with peptide/MHC complexes and anti-CD28 antibody.

This method is preferred for rare cancers and patients with a rare expression profile. In contrast to multi-peptide cocktails with a fixed composition as currently developed, the warehouse allows a significantly higher matching of the actual expression of antigens in the tumor with the vaccine. Selected single or combinations of several “off-the-shelf” peptides will be used for each patient in a multitarget approach. In theory an approach based on selection of e.g. 5 different antigenic peptides from a library of 50 would already lead to approximately 17 million possible drug product (DP) compositions.

In an aspect, the peptides are selected for inclusion in the vaccine based on their suitability for the individual patient based on the method according to the present invention as described herein, or as below.

The HLA phenotype, transcriptomic and peptidomic data is gathered from the patient's tumor material, and blood samples to identify the most suitable peptides for each patient containing "warehouse" and patient-unique (i.e. mutated) TUMAPs. Those peptides will be chosen, which are selectively or over-expressed in the patients' tumor and, where possible, show strong *in vitro* immunogenicity if tested with the patients' individual PBMCs.

Preferably, the peptides included in the vaccine are identified by a method comprising: (a) identifying tumor-associated peptides (TUMAPs) presented by a tumor sample from the individual patient; (b) comparing the peptides identified in (a) with a warehouse (database) of peptides as described above; and (c) selecting at least one peptide from the warehouse (database) that correlates with a tumor-associated peptide identified in the patient. For example, the TUMAPs presented by the tumor sample are identified by: (a1) comparing expression data from the tumor sample to expression data from a sample of normal tissue corresponding to the tissue type of the tumor sample to identify proteins that are over-expressed or aberrantly expressed in the tumor sample; and (a2) correlating the expression data with sequences of MHC ligands bound to MHC class I and/or class II molecules in the tumor sample to identify MHC ligands derived from proteins over-expressed or aberrantly expressed by the tumor. Preferably, the sequences of MHC ligands are identified by eluting bound peptides from MHC molecules isolated from the tumor sample and sequencing the eluted ligands. Preferably, the tumor sample and the normal tissue are obtained from the same patient.

In addition to, or as an alternative to, selecting peptides using a warehousing (database) model, TUMAPs may be identified in the patient *de novo*, and then included in the vaccine. As one example, candidate TUMAPs may be identified in the patient by (a1) comparing expression data from the tumor sample to expression data from a sample of

normal tissue corresponding to the tissue type of the tumor sample to identify proteins that are over-expressed or aberrantly expressed in the tumor sample; and (a2) correlating the expression data with sequences of MHC ligands bound to MHC class I and/or class II molecules in the tumor sample to identify MHC ligands derived from proteins over-expressed or aberrantly expressed by the tumor. As another example, proteins may be identified containing mutations that are unique to the tumor sample relative to normal corresponding tissue from the individual patient, and TUMAPs can be identified that specifically target the mutation. For example, the genome of the tumor and of corresponding normal tissue can be sequenced by whole genome sequencing: For discovery of non-synonymous mutations in the protein-coding regions of genes, genomic DNA and RNA are extracted from tumor tissues and normal non-mutated genomic germline DNA is extracted from peripheral blood mononuclear cells (PBMCs). The applied NGS approach is confined to the re-sequencing of protein coding regions (exome re-sequencing). For this purpose, exonic DNA from human samples is captured using vendor-supplied target enrichment kits, followed by sequencing with e.g. a HiSeq2000 (Illumina). Additionally, tumor mRNA is sequenced for direct quantification of gene expression and validation that mutated genes are expressed in the patients' tumors. The resultant millions of sequence reads are processed through software algorithms. The output list contains mutations and gene expression. Tumor-specific somatic mutations are determined by comparison with the PBMC-derived germline variations and prioritized. The *de novo* identified peptides can then be tested for immunogenicity as described above for the warehouse, and candidate TUMAPs possessing suitable immunogenicity are selected for inclusion in the vaccine.

In one exemplary embodiment, the peptides included in the vaccine are identified by: (a) identifying tumor-associated peptides (TUMAPs) presented by a tumor sample from the individual patient by the method as described above; (b) comparing the peptides identified in a) with a warehouse of peptides that have been prescreened for immunogenicity and overpresentation in tumors as compared to corresponding normal tissue; (c) selecting at least one peptide from the warehouse that correlates with a

tumor-associated peptide identified in the patient; and (d) optionally, selecting at least one peptide identified de novo in (a) confirming its immunogenicity.

In one exemplary embodiment, the peptides included in the vaccine are identified by: (a) identifying tumor-associated peptides (TUMAPs) presented by a tumor sample from the individual patient; and (b) selecting at least one peptide identified de novo in (a) and confirming its immunogenicity.

Once the peptides for a personalized peptide-based vaccine are selected, the vaccine is produced. The vaccine preferably is a liquid formulation consisting of the individual peptides dissolved in between 20-40% DMSO, preferably about 30-35% DMSO, such as about 33% DMSO.

Each peptide to be included into a product is dissolved in DMSO. The concentration of the single peptide solutions has to be chosen depending on the number of peptides to be included into the product. The single peptide-DMSO solutions are mixed in equal parts to achieve a solution containing all peptides to be included in the product with a concentration of ~2.5 mg/ml per peptide. The mixed solution is then diluted 1:3 with water for injection to achieve a concentration of 0.826 mg/ml per peptide in 33% DMSO. The diluted solution is filtered through a 0.22 µm sterile filter. The final bulk solution is obtained.

Final bulk solution is filled into vials and stored at -20°C until use. One vial contains 700 µL solution, containing 0.578 mg of each peptide. Of this, 500 µL (approx. 400 µg per peptide) will be applied for intradermal injection.

In addition to being useful for treating cancer, the peptides of the present invention are also useful as diagnostics. Since the peptides were generated from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin

lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer cells and since it was determined that these peptides are not or at lower levels present in normal tissues, these peptides can be used to diagnose the presence of a cancer.

The presence of claimed peptides on tissue biopsies in blood samples can assist a pathologist in diagnosis of cancer. Detection of certain peptides by means of antibodies, mass spectrometry or other methods known in the art can tell the pathologist that the tissue sample is malignant or inflamed or generally diseased, or can be used as a biomarker for acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer. Presence of groups of peptides can enable classification or sub-classification of diseased tissues.

The detection of peptides on diseased tissue specimen can enable the decision about the benefit of therapies involving the immune system, especially if T-lymphocytes are known or expected to be involved in the mechanism of action. Loss of MHC expression is a well described mechanism by which infected or malignant cells escape immunosurveillance. Thus, presence of peptides shows that this mechanism is not exploited by the analyzed cells.

The peptides of the present invention might be used to analyze lymphocyte responses against those peptides such as T cell responses or antibody responses against the peptide or the peptide complexed to MHC molecules. These lymphocyte responses can be used as prognostic markers for decision on further therapy steps. These responses

can also be used as surrogate response markers in immunotherapy approaches aiming to induce lymphocyte responses by different means, e.g. vaccination of protein, nucleic acids, autologous materials, adoptive transfer of lymphocytes. In gene therapy settings, lymphocyte responses against peptides can be considered in the assessment of side effects. Monitoring of lymphocyte responses might also be a valuable tool for follow-up examinations of transplantation therapies, e.g. for the detection of graft versus host and host versus graft diseases.

The present invention will now be described in the following examples which describe preferred embodiments thereof, and with reference to the accompanying figures, nevertheless, without being limited thereto. For the purposes of the present invention, all references as cited herein are incorporated by reference in their entireties.

FIGURES

Figures 1A through 1I show the over-presentation of various peptides in different cancer tissues (black dots). Upper part: Median MS signal intensities from technical replicate measurements are plotted as dots for single HLA-B*07 positive normal (grey dots, left part of figure) and tumor samples (black dots, right part of figure) on which the peptide was detected. Boxes display median, 25th and 75th percentile of normalized signal intensities, while whiskers extend to the lowest data point still within 1.5 interquartile range (IQR) of the lower quartile, and the highest data point still within 1.5 IQR of the upper quartile. Normal organs are ordered according to risk categories (blood cells, blood vessels, brain, liver, lung: high risk, grey dots; all other organs: medium risk; grey dots). Lower part: The relative peptide detection frequency in every organ is shown as spine plot. Numbers below the panel indicate number of samples on which the peptide was detected out of the total number of samples analyzed for each organ (N = 87 for normal samples, N = 149 for tumor samples). If the peptide has been detected on a sample but could not be quantified for technical reasons, the sample is included in this representation of detection frequency, but no dot is shown in the upper part of the figure. Tissues (from left to right) Normal samples: *blood cells*; *bloodvess* (blood vessels); *brain*; *heart*; *liver*; *lung*; *adrenal gl* (adrenal gland); *bile duct*; *bladder*; *bone*

marrow; *esoph* (esophagus); *gall bl* (gallbladder); *intest. la* (large intestine); *intest. sm* (small intestine); *kidney*; *lymph node*; *nerve periph* (peripheral nerve); *pancreas*; *pituit* (pituitary); *skin*; *spinal cord*; *spleen*; *stomach*; *thyroid*; *trachea*; Tumor samples: *AML* (acute myeloid leukemia); *BRCA* (breast cancer); *CCC* (cholangiocellular carcinoma); *CLL* (chronic lymphocytic leukemia); *CRC* (colorectal cancer); *GBC* (gallbladder cancer); *GBM* (glioblastoma); *GC* (gastric cancer); *HCC* (hepatocellular carcinoma); *HNSCC* (head and neck squamous cell carcinoma); *MEL* (melanoma); *NHL* (non-hodgkin lymphoma); *NSCLCadeno* (non-small cell lung cancer adenocarcinoma); *NSCLCother* (NSCLC samples that could not unambiguously be assigned to NSCLCadeno or NSCLCsquam); *NSCLCsquam* (squamous cell non-small cell lung cancer); *OC* (ovarian cancer); *OSCAR* (esophageal cancer); *PACA* (pancreatic cancer); *PRCA* (prostate cancer); *RCC* (renal cell carcinoma); *SCLC* (small cell lung cancer); *UBC* (urinary bladder carcinoma); *UEC* (uterine and endometrial cancer).

Figure 1A) Peptide: TPRIGPKVSL (SEQ ID NO.: 206), Figure 1B) Peptide: RPILKEQSSSSF (SEQ ID NO.: 246), Figure 1C) Peptide: RPDVVRTL (SEQ ID NO.: 308), Figure 1D) Peptide: SPSLQSSRESL (SEQ ID NO.: 312), Figure 1E) Peptide: NPPEPEKSL (SEQ ID NO.: 316), Figure 1F) Peptide: VPPQNPRPSL (SEQ ID NO.: 335), Figure 1G) Peptide: IPTSRVITL (SEQ ID NO.: 337), Figure 1H) Peptide: KPRATWTL (SEQ ID NO.: 353), Figure 1I) Peptide: SPRRLVELAGQSL (SEQ ID NO.: 359).

Figures 2A through 2T show exemplary expression profile of source genes of the present invention that are over-expressed in different cancer samples. Tumor (black dots) and normal (grey dots) samples are grouped according to organ of origin. Box-and-whisker plots represent median FPKM value, 25th and 75th percentile (box) plus whiskers that extend to the lowest data point still within 1.5 interquartile range (IQR) of the lower quartile and the highest data point still within 1.5 IQR of the upper quartile. Normal organs are ordered according to risk categories. FPKM: fragments per kilobase per million mapped reads. Normal samples: *blood cells*; *bloodvess* (blood vessels); *brain*; *heart*; *liver*; *lung*; *adipose* (adipose tissue); *adrenal gl* (adrenal gland); *bile duct*; *bladder*; *bone marrow*; *esoph* (esophagus); *eye*; *gall bl* (gallbladder); *head&neck*; *intest.*

la (large intestine); *intest. sm* (small intestine); *kidney*; *lymph node*; *nerve periph* (peripheral nerve); *pancreas*; *parathyr* (parathyroid gland); *perit* (peritoneum); *pituit* (pituitary); *pleura*; *skel. mus* (skeletal muscle); *skin*; *spleen*; *stomach*; *thyroid*; *trachea*; *ureter*; *breast*; *ovary*; *placenta*; *prostate*; *testis*; *thymus*; *uterus*. Tumor samples: *AML* (acute myeloid leukemia); *BRCA* (breast cancer); *CCC* (cholangiocellular carcinoma); *CLL* (chronic lymphocytic leukemia); *CRC* (colorectal cancer); *GBC* (gallbladder cancer); *GBM* (glioblastoma); *GC* (gastric cancer); *HCC* (hepatocellular carcinoma); *HNSCC* (head and neck squamous cell carcinoma); *MEL* (melanoma); *NHL* (non-hodgkin lymphoma); *NSCLCadeno* (non-small cell lung cancer adenocarcinoma); *NSCLCother* (NSCLC samples that could not unambiguously be assigned to NSCLCadeno or NSCLCsquam); *NSCLCsquam* (squamous cell non-small cell lung cancer); *OC* (ovarian cancer); *OSCAR* (esophageal cancer); *other* (other cancers, e.g. multiple myeloma); *PACA* (pancreatic cancer); *PRCA* (prostate cancer); *RCC* (renal cell carcinoma); *SCLC* (small cell lung cancer); *UBC* (urinary bladder carcinoma); *UEC* (uterine and endometrial cancer).

Figure 2A) Gene symbol: *KLK2*, Peptide: RPRSLQCVSL (SEQ ID No.: 1), Figure 2B) Gene symbol: *TGM4*, Peptide: GPKKFIVKL (SEQ ID No.: 4), Figure 2C) Gene symbol: *PRAME*, Peptide: LPSLSHCSQL (SEQ ID No.: 5), Figure 2D) Gene symbol: *MMP12*, Peptide: AVHEIGHSL (SEQ ID No.: 14), Figure 2E) Gene symbol: *FAM178B*, Peptide: WPRLPGAGL (SEQ ID No.: 19), Figure 2F) Gene symbol: *LOC645382/LOC645399/PRAMEF10*, Peptide: APSRLLEL (SEQ ID No.: 21), Figure 2G) Gene symbol: *MAGEC1*, Peptide: SPSFSSTLLSL (SEQ ID No.: 26), Figure 2H) Gene symbol: *CYP4Z1/CYP4Z2P*, Peptide: FPAPPAHWF (SEQ ID No.: 16), Figure 2I) Gene symbol: *UPK2*, Peptide: APLLPIRTLPL (SEQ ID No.: 58), Figure 2J) Gene symbol: *ALPPL2*, Peptide: DAAHPGPSV (SEQ ID No.: 82), Figure 2K) Gene symbol: *MMP1*, Peptide: HAIEKAFQL (SEQ ID No.: 90), Figure 2L) Gene symbol: *COL10A1*, Peptide: TPIPFDKILY (SEQ ID No.: 100), Figure 2M) Gene symbol: *LAMC2*, Peptide: RSYQHSLRL (SEQ ID No.: 116), Figure 2N) Gene symbol: *LAMC2*, Peptide: SPQLSYFEY (SEQ ID No.: 137), Figure 2O) Gene symbol: *SLC45A2*, Peptide: VPFNLITEY (SEQ ID No.: 138), Figure 2P) Gene symbol: *QRFPR*, Peptide: HPFKMKWQY (SEQ ID No.: 142), Figure 2Q) Gene symbol: *KBTBD8*, Peptide:

TPMTVPRI (SEQ ID No.: 159), Figure 2R) Gene symbol: KBTBD8, Peptide: YAYTSRVIL (SEQ ID No.: 166), Figure 2S) Gene symbol: NLRP2, Peptide: LPKAALLV (SEQ ID No.: 241), Figure 2T) Gene symbol: DCAF4L2, Peptide: APSMLRKNQL (SEQ ID No.: 358).

Figures 3A through 3G show exemplary results of peptide-specific in vitro CD8⁺ T cell responses of a healthy HLA-B*07⁺ donor. CD8⁺ T cells were primed using artificial APCs coated with anti-CD28 mAb and HLA-B*07 in complex with SEQ ID No.: 388 peptide (SPSVSQLSVL) (Figure 3A, left panel), SEQ ID No.: 406 peptide (LPDGSRVEL) (Figure 3B, left panel), SEQ ID No.: 23 peptide (VPRPTSTVGL) (Figure 3C, left panel), SEQ ID No.: 33 peptide (LPRGLSPARQL) (Figure 3D, left panel), SEQ ID No.: 61 peptide (LPRIPFSTF) (Figure 3E, left panel), SEQ ID No.: 64 peptide (VPRPIFSQLYL) (Figure 3F, left panel), SEQ ID No.: 68 peptide (FPNEVSVVL) (Figure 3G, left panel). After three cycles of stimulation, the detection of peptide-reactive cells was performed by 2D multimer staining with B*07/SEQ ID No.: 388 (Figure 3A), B*07/SEQ ID No.: 406 (Figure 3B), B*07/SEQ ID No.: 23 (Figure 3C), B*07/SEQ ID No.: 33 (Figure 3D), B*07/SEQ ID No.: 61 (Figure 3E), B*07/SEQ ID No.: 64 (Figure 3F) or B*07/SEQ ID No.: 68 (Figure 3G). Right panel (Figure 3A through 3G) show control staining of cells stimulated with irrelevant B*07/peptide complexes. Viable singlet cells were gated for CD8⁺ lymphocytes. Boolean gates helped excluding false-positive events detected with multimers specific for different peptides. Frequencies of specific multimer⁺ cells among CD8⁺ lymphocytes are indicated.

EXAMPLES

EXAMPLE 1

Identification and quantitation of tumor associated peptides presented on the cell surface

Tissue samples

Patients' tumor tissues were obtained from: Asterand (Detroit, MI, USA & Royston, Herts, UK), Bio-Options Inc. (Brea, CA, USA), BioServe (Beltsville, MD, USA), Conversant Bio (Huntsville, AL, USA), Geneticist Inc. (Glendale, CA, USA), Kyoto Prefectural University of Medicine (KPUM) (Kyoto, Japan), Osaka City University (Osaka, Japan), ProteoGenex Inc. (Culver City, CA, USA), Stanford Cancer Center (Stanford, CA, USA), Tissue Solutions Ltd (Glasgow, UK), University Hospital Bonn (Bonn, Germany), University Hospital Geneva (Geneva, Switzerland), University Hospital Heidelberg (Heidelberg, Germany), and University Hospital Tübingen (Tübingen, Germany)

Normal tissues were obtained from: Asterand (Detroit, MI, USA & Royston, Herts, UK), BioServe (Beltsville, MD, USA), Capital BioScience Inc. (Rockville, MD, USA), Centre for Clinical Transfusion Medicine Tübingen (Tübingen, Germany), Geneticist Inc. (Glendale, CA, USA), Kyoto Prefectural University of Medicine (KPUM) (Kyoto, Japan), ProteoGenex Inc. (Culver City, CA, USA), Tissue Solutions Ltd (Glasgow, UK), and University Hospital Tübingen (Tübingen, Germany)

Written informed consents of all patients had been given before surgery or autopsy. Tissues were shock-frozen immediately after excision and stored until isolation of TUMAPs at -70°C or below.

Isolation of HLA peptides from tissue samples

HLA peptide pools from shock-frozen tissue samples were obtained by immune precipitation from solid tissues according to a slightly modified protocol (Falk et al., 1991; Seeger et al., 1999) using the HLA-A*02-specific antibody BB7.2, the HLA-A, -B, C-specific antibody W6/32, the HLA-DR specific antibody L243 and the HLA DP specific antibody B7/21, CNBr-activated sepharose, acid treatment, and ultrafiltration.

Mass spectrometry analyses

The HLA peptide pools as obtained were separated according to their hydrophobicity by reversed-phase chromatography (nanoAcquity UPLC system, Waters) and the eluting

peptides were analyzed in LTQ-velos and fusion hybrid mass spectrometers (ThermoElectron) equipped with an ESI source. Peptide pools were loaded directly onto the analytical fused-silica micro-capillary column (75 μm i.d. x 250 mm) packed with 1.7 μm C18 reversed-phase material (Waters) applying a flow rate of 400 nL per minute. Subsequently, the peptides were separated using a two-step 180 minute-binary gradient from 10% to 33% B at a flow rate of 300 nL per minute. The gradient was composed of Solvent A (0.1% formic acid in water) and solvent B (0.1% formic acid in acetonitrile). A gold coated glass capillary (PicoTip, New Objective) was used for introduction into the nanoESI source. The LTQ-Orbitrap mass spectrometers were operated in the data-dependent mode using a TOP5 strategy. In brief, a scan cycle was initiated with a full scan of high mass accuracy in the Orbitrap ($R = 30\,000$), which was followed by MS/MS scans also in the Orbitrap ($R = 7500$) on the 5 most abundant precursor ions with dynamic exclusion of previously selected ions. Tandem mass spectra were interpreted by SEQUEST at a fixed false discovery rate ($q \leq 0.05$) and additional manual control. In cases where the identified peptide sequence was uncertain it was additionally validated by comparison of the generated natural peptide fragmentation pattern with the fragmentation pattern of a synthetic sequence-identical reference peptide.

Label-free relative LC-MS quantitation was performed by ion counting i.e. by extraction and analysis of LC-MS features (Mueller et al., 2007). The method assumes that the peptide's LC-MS signal area correlates with its abundance in the sample. Extracted features were further processed by charge state deconvolution and retention time alignment (Mueller et al., 2008; Sturm et al., 2008). Finally, all LC-MS features were cross-referenced with the sequence identification results to combine quantitative data of different samples and tissues to peptide presentation profiles. The quantitative data were normalized in a two-tier fashion according to central tendency to account for variation within technical and biological replicates. Thus each identified peptide can be associated with quantitative data allowing relative quantification between samples and tissues. In addition, all quantitative data acquired for peptide candidates was inspected manually to assure data consistency and to verify the accuracy of the automated analysis. For each peptide a presentation profile was calculated showing the mean

sample presentation as well as replicate variations. The profiles juxtapose acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer samples to a baseline of normal tissue samples. Presentation profiles of exemplary over-presented peptides are shown in Figures 1A-1I.

Table 8a and 8b show the presentation on various cancer entities for selected peptides, and thus the particular relevance of the peptides as mentioned for the diagnosis and/or treatment of the cancers as indicated (e.g. peptide SEQ ID No. 22 for head and neck squamous cell carcinoma and esophageal cancer, peptide SEQ ID No. 34 for breast cancer and uterine and endometrial cancer).

Table 8a: Overview of presentation of selected tumor-associated peptides of the present invention across entities.

AML: acute myeloid leukemia; BRCA: breast cancer; CCC: cholangiocellular carcinoma; CLL: chronic lymphocytic leukemia; CRC: colorectal cancer; GBC: gallbladder cancer; GBM: glioblastoma; GC: gastric cancer; HCC: hepatocellular carcinoma; HNSCC: head and neck squamous cell carcinoma; MEL: melanoma; NHL: non-Hodgkin lymphoma; NSCLCaden: non-small cell lung cancer adenocarcinoma; NSCLCother: NSCLC samples that could not unambiguously be assigned to NSCLCaden or NSCLCsquam; NSCLCsquam: squamous cell non-small cell lung cancer; OC: ovarian cancer; OSCAR: esophageal cancer; PACA: pancreatic cancer; PRCA: prostate cancer; RCC: renal cell carcinoma; SCLC: small cell lung cancer; UBC: urinary bladder carcinoma; UEC: uterine and endometrial cancer.

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
1	RPRSLQCVSL	PRCA

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
2	VPGSDPARYEFL	OSCAR
3	MPYVVLTAL	RCC
4	GPKKFIVKL	PRCA
5	LPSLSHCSQL	MEL
6	LPLNSSTSL	SCLC
8	MPKTNLSKM	HNSCC
12	LPNTGRIGQLL	NSCLCadenoc
13	LPNTGRIGQL	PRCA
14	AVHEIGHSL	NSCLCadenoc
15	KPGFNISIL	BRCA
16	FPAPPAHWF	BRCA
17	SPAAPLSPASSL	UBC
18	MALSVLRAL	GBC, GC, NSCLCsquam, OC, OSCAR, PACA, PRCA, SCLC
20	MVLGIGPVL	CRC, GBC, HNSCC, NSCLCadenoc, NSCLCsquam, OC, SCLC, UEC
21	APSRLEL	HCC
22	LPQLKPAL	HNSCC, OSCAR
23	VPRPTSTVGL	MEL
24	VPRPTSTVGLFL	MEL
25	RPQGAVGGL	HCC
27	RIRVTSEVL	BRCA
29	APLRVHITSL	MEL
31	RPGPSDPAA	NSCLCadenoc
32	APMAPGRSPL	UEC
33	LPRGLSPARQL	UEC
34	APMAPGRSP	BRCA, UEC
35	APLPPPRAL	GBM
36	RPFSREMDL	AML, OC
38	RPSFPNLTSF	AML, OC
39	QPRPSSEKI	AML
40	FPRTVKHIDAAL	CRC
42	APLKMLAL	MEL, OC
43	APTPRPKVL	BRCA
45	TRPWSGPYIL	MEL
46	QPISGNPVTI	NHL, NSCLCsquam, SCLC, UBC
47	RPRQTGALM	AML
48	RPRYSIGL	BRCA
49	APEKARAFL	OSCAR
50	PEKARAFL	OSCAR

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
51	SPVFYVQTL	BRCA
53	SPRIPFSTF	MEL
54	VPSCGRSVEGL	CLL, GBC, NHL, NSCLCadenoc, OC, OSCAR, PACA, RCC, UEC
55	LPALLRSGSL	CCC, GBC, HCC, HNSCC, NSCLCadenoc, NSCLCsquam, OC, OSCAR, RCC, SCLC
56	LPALLRSGSLTL	BRCA, CCC, CRC, GBC, HCC, HNSCC, MEL, NHL, NSCLCadenoc, OC, OSCAR, RCC
57	APLLPIRTL	UBC
58	APLLPIRTLPL	UBC
59	KPRTIYSSL	OSCAR, UEC
60	RPYSIYPHGVTF	CCC, GBC, HCC
61	LPRIPFSTF	MEL
63	FPHMATTAF	HNSCC
64	VPRPIFSQLYL	NHL, UEC
65	FPNVYSTLV	MEL
66	LPMTVISVL	GBC
67	VPVSRPVL	CLL, NHL
68	FPNEVSVVL	BRCA, MEL, NSCLCadenoc, NSCLCsquam, OC, OSCAR
69	RPEDGRPRL	GBM
70	VPAQRLGSL	AML, NSCLCadenoc, NSCLCsquam, OC, OSCAR
71	APFAPLHGGGSL	PRCA
72	APCNGNVAV	BRCA, NSCLCadenoc
73	LPVSSPVT	BRCA, NSCLCsquam
74	VPVSHPV	CLL, NHL
76	KPKVESQAL	BRCA
77	QPRLPGETL	GBC, MEL, NHL, NSCLCadenoc, NSCLCsquam
78	HPSQESPTL	NSCLCsquam, OC, RCC, UBC, UEC
79	GPASDHPSL	OC, PRCA, UEC
80	SALPTTISF	HNSCC, OSCAR, SCLC
82	DAAHPGPSV	UEC
83	AVSSHNV	RCC
84	MPMQDIKMI	UEC
85	MPMQDIKMILKM	OC, SCLC
86	ALLLRGVTL	SCLC
87	APVGGNVTSF	OSCAR
88	KPSAVKDSIY	HNSCC, RCC
89	FLIPRAGWLAGL	CRC
90	HAIEKAFQL	CRC
91	FPRLVGPDDF	OC

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
92	TSPLPPTV	AML, HNSCC
93	SVAIAHGVF	CLL, CRC
94	LPMSKRQEY	OC
95	RTKEEINEL	NSCLCadenos
96	QPSLVQAIF	OC
97	LPPGTQIQI	CRC
98	FPCSALLACF	OSCAR
99	MPVSAFTVI	GC, NSCLCsquam
100	TIPIFDKILY	OSCAR
101	KPGLPGLK	BRCA, CRC, GBM, HCC, MEL, NHL, NSCLCadenos, NSCLCsquam, OC, OSCAR, RCC, SCLC, UBC, UEC
102	MAGPAIHTAPM	HNSCC
105	RPRYETGVCA	CRC
106	APFHIDRLF	OC, RCC
107	GQRLESPGGAA	AML, CCC, CLL, CRC, GBC, HCC, HNSCC, NSCLCsquam, PACA, UBC, UEC
108	APRGSPPI	HCC, OC, OSCAR, PACA, RCC
110	YPSSPASAV	GC
111	YPLQQTNVY	RCC
112	YPSPLNKHSF	RCC
114	FVPPSGPSNPM	HNSCC
115	KTKSLAQQL	CRC, NSCLCadenos, UEC
116	RSYQHSLRL	CRC
117	IPHQRSSL	CLL
118	NPERHKPSF	AML
119	KATEYVHSL	CRC, UEC
120	TVRPKNAAL	AML, HCC, OSCAR, UEC
121	GPFQRAAL	UEC
122	KPRTPFTTAQLL	UEC
123	RPRLRDLPAL	CLL, GBC, NHL, OC, OSCAR, RCC
124	KTIDGHINL	CRC, NSCLCadenos, NSCLCother
125	SPAKQFNIY	SCLC
126	MPREDAHF	MEL
127	KSKQVITLL	NSCLCsquam
128	SPPATLFLFLL	OSCAR
129	MTLPATTLTL	BRCA, CCC, CLL, CRC, GBC, GBM, HCC, HNSCC, MEL, NSCLCadenos, NSCLCsquam, OC, OSCAR, PACA, RCC, SCLC, UBC, UEC
130	GAYDKARYL	BRCA
131	MPFRNIRSIL	HCC

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
132	LPRLPVGSLV	GC
133	LPELPTTLTF	NSCLCadenos
134	VAAAARLTL	MEL
135	MPRLRLSL	UEC
136	QPDHAGIFRVF	HNSCC, OSCAR
137	SPQLSYFEY	OSCAR
138	VPFNLITEY	MEL
139	LVILQHQAM	RCC
140	TPFPDIHWF	NSCLCsquam, RCC, SCLC
141	YGPYGGSGGW	GC
142	HPFKMKWQY	RCC
143	YPMIPPAQL	SCLC
144	RPVQGIPTY	MEL
145	VPAQRLGLLL	BRCA, GBM, HNSCC, NSCLCsquam, OC
146	SPTRGSEF	GBM
147	HVAQPAVVL	BRCA, HCC
148	KPHLIYRQ	AML, BRCA, CRC, HCC, HNSCC, MEL, NSCLCadenos, OSCAR, RCC, SCLC, UEC
149	SPSSVFTSK	BRCA
150	APLHGGGSL	PRCA
151	QPTWNTQTRL	NHL
152	SSASFSSSELFRLH	BRCA, CCC, OSCAR, PACA, SCLC, UEC
153	QPVHSLCSA	GC
154	RPPPSRRAVL	GBM
155	APIPRLIEG	OSCAR
156	GSRPVTGSV	BRCA
157	RPVTGSVC	BRCA, NSCLCadenos
158	SVPVATVEL	CLL
159	TPMTVPRI	CLL, NHL
160	IPVSHPVLTFL	BRCA, CRC, GBM, HCC, NSCLCadenos
161	ASKILETSL	MEL
163	FRYPNGVSL	HCC
164	RAAGRRRLQL	HCC
166	YAYTSRVIL	NHL
167	NVNPARKGL	AML
168	SPSGQRDVSL	MEL
169	RPFSVSSISQL	MEL
170	APEGKRLGF	MEL
171	LPLGGHKSL	MEL
172	SAQSLHPVL	GBM

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
173	VIINNPISL	NSCLCother
174	IPVTSSVTL	NSCLCsquam, OSCAR
175	AAFPHKIIF	OSCAR
176	QPLDICHSF	OC
177	VWEPQGSSRSL	UEC
178	VPYYPRANL	OSCAR
179	PRVRLLLL	GBC, HNSCC
180	RLRDYISSL	CLL, NHL
181	LPVSPARAL	HNSCC, NSCLCadenoc, NSCLCsquam, OSCAR
182	RPLPVSPARAL	BRCA, NSCLCadenoc, NSCLCsquam
183	VPRRPARAL	BRCA, GBM, MEL
184	KIIASGNHL	NSCLCother
185	RPVLTASSF	NHL
188	HAAASFETL	NSCLCadenoc
189	QPQCSTHIL	CCC, HCC, OSCAR
190	RLAHVRARL	UEC
191	KPKAVKPKAA	NSCLCadenoc
192	IPFADVKSF	NSCLCadenoc, NSCLCsquam, SCLC
193	LPALKEEAF	PRCA
194	HPNKKASL	BRCA
195	NPIFGRKEL	UEC
196	RPSGTAGAAL	HNSCC, NSCLCsquam, RCC
197	RPSGTAGAALLA	HCC
198	LPSPAHAFRAL	OC
199	SPFWIHQA	AML
200	EPFHLIVSY	CRC, HNSCC, MEL, NSCLCadenoc, NSCLCsquam, OSCAR, RCC, SCLC
201	LPIARVLTV	UEC
202	SPSREASAL	CLL, GBM, UBC
203	KPRGPIDI	CRC, GBC, NSCLCadenoc
204	FPYSGDKILV	AML, CRC, GC, HCC, MEL, NSCLCsquam, PACA, SCLC, UBC, UEC
205	LPPALLTTV	GC
206	TPRIGPKVSL	AML, BRCA, CCC, CLL, CRC, GBC, GBM, GC, HCC, HNSCC, MEL, NSCLCadenoc, NSCLCsquam, OSCAR, PACA, RCC, SCLC, UBC, UEC
207	VPSDITTAF	NSCLCadenoc, OSCAR, SCLC
208	RNRQVATAL	CCC, CRC, GBC, HNSCC
209	KIEQIRAVL	OSCAR
210	IPENRVVSY	OSCAR

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
211	IPDTIASVL	MEL
212	VPYAAQAAL	BRCA
214	LPLKFFPII	UEC
215	IPVAIKEL	NSCLCother
216	LPWEQNEQV	HCC
217	SPGDKRLAAYL	CCC, HCC
218	VQRTPPPI	GC, NSCLCadenoc, NSCLCsquam, OSCAR
219	VPHTGRYTCL	RCC
220	RPAPGPAPFV	BRCA
221	LPQRPNARL	CCC, CRC, GBC, GC, HCC, HNSCC, MEL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, PACA, PRCA, RCC, SCLC, UBC, UEC
222	MLKTTLTAF	HCC
223	KAHVRIEL	CRC
226	APGGSSRSSL	AML
229	FPNAGPRHLL	AML, CLL, NHL, OC, RCC
230	DVIDDIISL	GBM, HCC, HNSCC, NHL, NSCLCsquam
231	SPITLQAL	HCC
232	TAYPSLRLL	BRCA, HCC, MEL, NHL, NSCLCadenoc, NSCLCsquam, PACA, SCLC, UEC
233	MAYDRFIAI	CRC, GBC, NSCLCadenoc, RCC
234	HPRAPGEQQRL	BRCA, CRC, UEC
235	AQNPAGTAL	MEL
236	TPELKSSIL	HCC
237	LPRAGGAF	GBC, GC, NSCLCadenoc, PACA, RCC
238	LPRAGGAFLM	HNSCC
239	VLPRAGGAFLM	NSCLCadenoc
241	LPKAALLV	CLL
242	IPETASVVAI	AML, CCC, CRC, NSCLCadenoc, NSCLCsquam
243	MARTGGMVVI	CRC, NSCLCadenoc, NSCLCsquam
244	VPAHLVAA	GC
245	GPVPSPLAL	OC
246	RPILKEQSSSF	GBC, HNSCC, NSCLCsquam, OSCAR
247	SPVGVGQRL	SCLC
249	SPRSGVLL	OSCAR, UEC
250	APAAPAAVPS	CRC, GBM, OC
251	MPVDSFNSM	MEL, NSCLCadenoc, NSCLCsquam, RCC, SCLC, UBC
252	QPENSLEGISL	NSCLCsquam
253	MPVDSFNML	CRC, MEL, NSCLCsquam, SCLC, UBC
254	RVIQGTTL	HCC

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
255	VPSHWMVAL	NHL
256	SPVPSHWMVAL	CLL, NHL
257	APYGTLRKS	MEL
258	SVIGPNSRL	SCLC
259	VPMPGVQAV	UEC
260	LVQSSRSEV	HNSCC, OSCAR
261	SPSTSRTPL	GBC, GBM, HCC, HNSCC, NSCLCadenoc
262	SPSTSRTPLLSSL	BRCA, CCC, GBC, GBM, HCC, HNSCC, NHL, OSCAR, RCC, UEC
263	SQRPPATSQA	AML, BRCA, CCC, CLL, CRC, GBM, HCC, HNSCC, NHL, OC, OSCAR, PACA, PRCA, RCC
264	APRPGNWIL	MEL, NSCLCsquam
266	KPYEGRVEI	CLL, GBM
267	MPVPGILL	MEL
268	EPLSVTASY	HNSCC
269	FTVSSSSAM	GBC
270	SPRGTTSTL	CCC, CRC, GBC, HCC, NHL
271	SPTPVFTTL	CRC, GBC
272	SSPRGTTSTL	GBC, OSCAR
273	TDTPSTPTTI	CCC, CRC, GBC, GBM, HCC, MEL, NSCLCadenoc, OSCAR, PACA, PRCA, RCC, SCLC, UEC
274	KPIRTGISPLAL	CCC, HCC
276	RPVWDVRSAL	HNSCC
277	MPPLLIVAF	OC
278	LIAARGSLVL	AML, CRC, MEL, NSCLCadenoc, OSCAR, SCLC, UEC
279	APADEPRTVL	NSCLCsquam
280	LPRAFLQSL	CRC, NHL, NSCLCsquam
281	NPRSPPATM	AML, OC
282	AVRLPTPNL	AML
284	RPSAPRGPP	CRC, OSCAR
285	RITWKGLLL	NSCLCother
286	APARPAAAF	BRCA, MEL, NSCLCadenoc, OC, RCC
287	SPIPRPLFL	BRCA
288	LHAMNSLSAM	NHL
289	LPYEKVSRL	CCC, NHL
291	SPSKSLEM	CRC
292	LPMTHRLQL	MEL
293	FPYDKPLIM	GC, NSCLCadenoc
296	SPLLMQRTL	CLL, NHL
297	FPIKYVNDF	RCC

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
298	RVLLRWISL	GBM
299	SPFSGGPVSF	RCC
300	HPYSDLADV	MEL
301	FPAFLEAM	MEL
302	IPIDQILNSF	MEL
303	RPPPPCIAL	MEL
304	SPLIGDFPAF	MEL
305	AASPVGSAL	HCC
306	RPFPLALL	CRC
307	RPHQKGWLSI	HCC
308	RPDVVRTLL	CRC, GBC, GC, HNSCC, MEL, NSCLCadenoc, NSCLCsquam, OSCAR, PACA, RCC, UBC
311	APRLALDPDAL	BRCA, SCLC
312	SPSLQSSRESL	GBC, HNSCC, NSCLCadenoc, NSCLCsquam, OSCAR, UBC
313	VPLSSYVSI	MEL
314	IALMKLAGPL	AML, CRC, HCC, MEL, NHL, NSCLCadenoc, UEC
316	NPREPEKSL	BRCA, CRC, GBC, GC, HCC, OSCAR, SCLC
317	MPYNSAHHCVV	CLL, NHL
318	TPISNTGVL	AML, GBC, NHL, NSCLCadenoc, NSCLCother, RCC
319	RPLDTRVL	OSCAR
320	APMHIDRNIL	HCC
321	QPQQPGFFL	CLL
322	RAVPVGSGL	MEL, NSCLCadenoc, NSCLCsquam
323	TPHGITDIL	CRC, UEC
324	LPAPLRISL	BRCA, GBM, MEL, NSCLCadenoc, OC, UEC
325	SPRSNPVRL	NHL
326	IPPFTQRVF	OC
327	GPRTTQSSVL	NSCLCadenoc
328	LPLHRGDLVI	CLL
329	QPANFIVLL	GBC
330	RPFSAIAEL	GBC
331	SPDSAILKL	NSCLCsquam, UEC
332	SPYAGSTAF	AML, NSCLCadenoc
333	SVLPRALSL	BRCA
334	YPLSLALHF	OC
335	VPPQNPRPSL	AML, GBM, HNSCC, NHL, NSCLCadenoc, OSCAR
336	YPLQGPGLLSV	NHL
337	IPTSRVITL	BRCA, CLL, HCC, MEL, NHL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, PACA, SCLC

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
338	MPATPSLKV	HNSCC, OC, PACA, UBC
339	GPQRTTSV	CCC
340	APEPRAALL	MEL, NSCLCsquam, OC
343	RPRPPKVLGL	AML, CCC, HCC, MEL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, RCC, SCLC, UEC
344	VPYPSPTCV	BRCA, HCC, UEC
345	SAAPPGASL	HCC
346	IPMPRITWL	BRCA
347	SPLEKINSF	GBC
348	HPAPPV TSA	OC
349	QPRDGWPMML	HNSCC, OSCAR
350	RPKSTLMNF	OC, OSCAR
351	SPYADIIPSA	HNSCC
352	LPAFSKIGGIL	CRC, MEL
353	KPRATWTL	AML, BRCA, CCC, CLL, GBC, GC, HCC, HNSCC, MEL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, PACA, RCC, SCLC, UEC
354	APAKDARASL	CRC, GBC, HNSCC, NSCLCadenoc, NSCLCsquam, OC, OSCAR, UBC
355	APKTSFIAA	GC
356	RPFLRLPSL	CRC, HCC, HNSCC, OC, OSCAR
357	LPPHIFAI	CRC, GC, HCC, UEC
358	APSMRLRKNQL	HCC
359	SPRRLVELAGQSL	MEL, OC, OSCAR, UEC
361	SPYGSDRLVQL	CLL, NHL
362	KPMLPPAAF	UEC
363	KPRTPF TTA	UEC
364	KPRTPF TTAQL	BRCA, CRC, GBM, RCC, UEC
365	RPKHFLQML	OSCAR
366	SPTLRQLDL	CCC, HCC
367	APQVHIFSL	BRCA, CRC, GBM, RCC, UEC
368	NPASRLTAL	BRCA, CCC, CRC, GBC, GBM, GC, HCC, HNSCC, MEL, NHL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, PACA, PRCA, RCC, SCLC, UBC, UEC
370	AAHEFGHVL	BRCA, UEC
371	APRSPGQVTPRGL	NSCLCadenoc, OC
372	SPSSASLTL	GBC, NSCLCsquam, OSCAR
373	LPKPDLPQLI	AML, BRCA, HCC, MEL, NSCLCadenoc, PRCA, RCC, UEC
374	KPRNMTGLDL	CLL
375	LPRGVLEGL	CRC, NSCLCadenoc, UEC

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
376	FPQVGRTAL	AML, BRCA, CCC, CLL, CRC, GBC, GC, HCC, HNSCC, MEL, NHL, NSCLCadenoc, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, PRCA, RCC, SCLC, UBC, UEC
377	SPFSKRIKL	AML, BRCA, CLL, GC, HNSCC, MEL, NHL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, RCC, SCLC, UBC, UEC
379	GPRQVLFPL	AML, BRCA, CCC, CRC, GBC, GBM, GC, HCC, HNSCC, MEL, NHL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, PACA, PRCA, RCC, SCLC, UBC, UEC
380	SPYPGSAAF	BRCA, NSCLCadenoc, OSCAR
381	APRPRLLL	AML, BRCA, CCC, CLL, CRC, GBC, GBM, GC, HCC, HNSCC, MEL, NHL, NSCLCadenoc, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, PRCA, RCC, SCLC, UBC, UEC
382	RPGPQRTTSV	BRCA, CCC, CLL, HCC, MEL, NHL, OC, OSCAR, PRCA, SCLC, UEC
383	KPRATWTLKL	BRCA, CCC, CRC, GBC, GC, HCC, HNSCC, MEL, NHL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, RCC, SCLC, UEC

Table 8b: Overview of presentation of selected tumor-associated peptides of the present invention across entities.

AML: acute myeloid leukemia; BRCA: breast cancer; CCC: cholangiocellular carcinoma; CLL: chronic lymphocytic leukemia; CRC: colorectal cancer; GBC: gallbladder cancer; GBM: glioblastoma; GC: gastric cancer; HCC: hepatocellular carcinoma; HNSCC: head and neck squamous cell carcinoma; MEL: melanoma; NHL: non-Hodgkin lymphoma; NSCLCadenoc: non-small cell lung cancer adenocarcinoma; NSCLCother: NSCLC samples that could not unambiguously be assigned to NSCLCadenoc or NSCLCsquam; NSCLCsquam: squamous cell non-small cell lung cancer; OC: ovarian cancer; OSCAR: esophageal cancer; PACA: pancreatic cancer; PRCA: prostate cancer; RCC: renal cell carcinoma; SCLC: small cell lung cancer; UBC: urinary bladder carcinoma; UEC: uterine and endometrial cancer.

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
448	TVYGEPRKL	OSCAR
449	RPKTSVNLISSL	SCLC
450	SAAARALLP	CCC, CRC, GBM, GC, HCC, NSCLCother,

		NSCLCsquam, PACA, SCLC, UEC
451	VPILPLNHAL	UEC
454	SPDPAHLESL	NSCLCother
455	FPVQATIDF	OC
456	VPVSHPVLT	NSCLCsquam
457	RPPLSQRHTF	NHL
459	SPAPWRPWI	SCLC
460	APLMPLGKTL	PACA
461	MPHLGPGILL	MEL
462	MPLLADVRL	MEL
463	LPTDLFNSVM	MEL
467	VPRNQDWLGVSRL	MEL
468	LPSLHVLVL	MEL
469	LPLDTRTSI	CLL
470	RPNGEVKSEL	SCLC
471	LPLLAGTLLL	NSCLCaden
472	LPMPAITWY	MEL
473	LPGEREAL	MEL
474	VPKADLLTL	MEL
475	IPLEIQKL	MEL
476	EPNPVEEIF	NSCLCsquam
477	NPVPVITWYKDNRL	MEL
478	APKFISPASQL	SCLC
479	APHAGGALL	NSCLCsquam
480	VPAAPTAL	CLL
481	SPARALLALA	HNSCC
482	QPSLKKIIL	PACA
484	LPASAEPKGTM	HNSCC
485	LPLKTKVFA	NSCLCsquam
486	TPTRPLPSA	NSCLCaden
487	IPWFIKTAF	MEL
488	LPLGGLPLLI	CLL
489	WPNHIMLVL	MEL
491	MPEADLKRIF	MEL
492	IPFSVDEI	NSCLCother, UEC
493	LPLQQYKLV	HCC
494	LPLRAVNLNL	SCLC
495	SPSYTQASL	BRCA
496	MPAVSGPGPLF	NSCLCaden
497	YVVKPLHPF	CRC

498	RPQPQPRPAL	HNSCC, OSCAR
499	QPVLPSPAC	MEL

EXAMPLE 2

Expression profiling of genes encoding the peptides of the invention

Over-presentation or specific presentation of a peptide on tumor cells compared to normal cells is sufficient for its usefulness in immunotherapy, and some peptides are tumor-specific despite their source protein occurring also in normal tissues. Still, mRNA expression profiling adds an additional level of safety in selection of peptide targets for immunotherapies. Especially for therapeutic options with high safety risks, such as affinity-matured TCRs, the ideal target peptide will be derived from a protein that is unique to the tumor and not found on normal tissues.

RNA sources and preparation

Surgically removed tissue specimens were provided as indicated above (see Example 1) after written informed consent had been obtained from each patient. Tumor tissue specimens were snap-frozen immediately after surgery and later homogenized with mortar and pestle under liquid nitrogen. Total RNA was prepared from these samples using TRI Reagent (Ambion, Darmstadt, Germany) followed by a cleanup with RNeasy (QIAGEN, Hilden, Germany); both methods were performed according to the manufacturer's protocol.

Total RNA from healthy human tissues for RNASeq experiments was obtained from: Asterand (Detroit, MI, USA & Royston, Herts, UK); Bio-Options Inc. (Brea, CA, USA); Geneticist Inc. (Glendale, CA, USA); ProteoGenex Inc. (Culver City, CA, USA); Tissue Solutions Ltd (Glasgow, UK). Total RNA from tumor tissues for RNASeq experiments was obtained from: Asterand (Detroit, MI, USA & Royston, Herts, UK); BioCat GmbH (Heidelberg, Germany); BioServe (Beltsville, MD, USA); Geneticist Inc. (Glendale, CA, USA); Istituto Nazionale Tumori "Pascale" (Naples, Italy); ProteoGenex Inc. (Culver City, CA, USA); University Hospital Heidelberg (Heidelberg, Germany).

Quality and quantity of all RNA samples were assessed on an Agilent 2100 Bioanalyzer (Agilent, Waldbronn, Germany) using the RNA 6000 Pico LabChip Kit (Agilent).

RNAseq experiments

Gene expression analysis of - tumor and normal tissue RNA samples was performed by next generation sequencing (RNAseq) by CeGaT (Tübingen, Germany). Briefly, sequencing libraries are prepared using the Illumina HiSeq v4 reagent kit according to the provider's protocol (Illumina Inc., San Diego, CA, USA), which includes RNA fragmentation, cDNA conversion and addition of sequencing adaptors. Libraries derived from multiple samples are mixed equimolar and sequenced on the Illumina HiSeq 2500 sequencer according to the manufacturer's instructions, generating 50 bp single end reads. Processed reads are mapped to the human genome (GRCh38) using the STAR software. Expression data are provided on transcript level as RPKM (Reads Per Kilobase per Million mapped reads, generated by the software Cufflinks) and on exon level (total reads, generated by the software Bedtools), based on annotations of the ensembl sequence database (Ensembl77). Exon reads are normalized for exon length and alignment size to obtain RPKM values.

Exemplary expression profiles of source genes of the present invention that are highly over-expressed or exclusively expressed in acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer are shown in Figures 2A-2T. Expression scores for further exemplary genes are shown in Table 9a and 9b.

Table 9a: Expression scores. The table lists peptides from genes that are very highly over-expressed in tumors compared to a panel of normal tissues (+++), highly over-

expressed in tumors compared to a panel of normal tissues (++) or over-expressed in tumors compared to a panel of normal tissues (+). The baseline for this score was calculated from measurements of the following relevant normal tissues: adipose tissue, adrenal gland, bile duct, blood cells, blood vessels, bone marrow, brain, esophagus, eye, gallbladder, heart, head&neck, kidney, large intestine, liver, lung, lymph node, nerve, parathyroid, pancreas, pituitary, pleura, skeletal muscle, skin, small intestine, spleen, stomach, thyroid gland, trachea, ureter, urinary bladder. In case expression data for several samples of the same tissue type were available, the arithmetic mean of all respective samples was used for the calculation. AML: acute myeloid leukemia; BRCA: breast cancer; CCC: cholangiocellular carcinoma; CLL: chronic lymphocytic leukemia; CRC: colorectal cancer; GBC: gallbladder cancer; GBM: glioblastoma; GC: gastric cancer; HCC: hepatocellular carcinoma; HNSCC: head and neck squamous cell carcinoma; MEL: melanoma; NHL: non-Hodgkin lymphoma; NSCLCaden: non-small cell lung cancer adenocarcinoma; NSCLCother: NSCLC samples that could not unambiguously be assigned to NSCLCaden or NSCLCsquam; NSCLCsquam: squamous cell non-small cell lung cancer; OC: ovarian cancer; OSCAR: esophageal cancer; PACA: pancreatic cancer; PRCA: prostate cancer; RCC: renal cell carcinoma; SCLC: small cell lung cancer; UBC: urinary bladder carcinoma; UEC: uterine and endometrial cancer

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
1	RPRSLQCVSL	CLL		PRCA
2	VPGSDPARYEFL	HCC, UEC	GBC, GC, HNSCC, NSCLCaden, NSCLCsquam, OC, OSCAR, SCLC	MEL, UBC
3	MPYVVLTA	NSCLCsquam, SCLC	NSCLCaden	RCC
4	GPKKFIVKL			PRCA
5	LPSLSHCSQL	AML, BRCA, NHL, OSCAR, UBC	GBC, HNSCC, NSCLCaden, NSCLCsquam,	MEL, UEC

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
			OC, RCC, SCLC	
6	LPLNSSTSL	BRCA, HCC, RCC	CCC, CRC, GC, HNSCC, MEL, NHL, NSCLCadenoc, OC, OSCAR, PACA, UBC, UEC	GBC, NSCLCother, NSCLCsquam, SCLC
7	RPSQLAPATL	AML, GC, HCC, HNSCC, MEL, NSCLCadenoc, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, UBC, UEC	GBC	SCLC
8	MPKTNLSKM	AML, GC, HCC, HNSCC, MEL, NSCLCadenoc, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, UBC, UEC	GBC	SCLC
9	RPDSRLLEL	GC, HNSCC, NSCLCsquam, OC, UBC	HCC, MEL, SCLC	GBC
10	SPMEAELVRRIL	GC, HNSCC, NSCLCsquam, OC, UBC	HCC, MEL, SCLC	GBC
11	APLPRPGAV	GC, HNSCC, NSCLCsquam, OC, UBC	HCC, MEL, SCLC	GBC
12	LPNTGRIGQLL			PRCA
13	LPNTGRIGQL			PRCA
14	AVHEIGHSL	HCC, RCC	CCC, GC, MEL, NHL, NSCLCadenoc, OC, PACA, UEC	CRC, GBC, HNSCC, NSCLCother, NSCLCsquam, OSCAR, SCLC, UBC
15	KPGFNISIL	OC		BRCA

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
16	FPAPPAHWF	OC		BRCA
17	SPAAPLSPASSL			UBC
18	MALSVLRLAL			AML
19	WPRLPGAGL	GC, HCC, NSCLCsquam, PACA, SCLC, UBC, UEC	GBC, HNSCC, OSCAR	MEL
20	MVLGIGPVL			PRCA
21	APSRLEL	CCC, CRC, GBC, MEL, SCLC		HCC
22	LPQLKPAL	BRCA, NSCLCsquam	OSCAR	HNSCC
23	VPRPTSTVGL		MEL	
24	VPRPTSTVGLFL		MEL	
25	RPQGAVGGL	GBC	HCC	
26	SPSFSTLLSL	BRCA, GBC, SCLC	HCC, MEL	
27	RIRVTSEVL	HCC, MEL, UEC	BRCA, PRCA	
28	LPAPTSLVL		GBM	
29	APLRVHITSL	HCC	CCC, NSCLCaden, OC	
30	YPGFTKRL	GC, HNSCC, MEL, NSCLCaden, NSCLCsquam, OSCAR, PACA, RCC, SCLC	BRCA, CCC, CRC, GBC, HCC, OC, UBC	
31	RPGPSDPAA	NSCLCaden, UEC	RCC	
32	APMAPGRSPL	BRCA	UEC	
33	LPRGLSPARQL	BRCA	UEC	
34	APMAPGRSP	BRCA	UEC	
35	APLPPPRAL		GBM	
36	RPFSREMDL		AML	
37	LPPPRALTL		GBM	
38	RPSFPNLTSF		AML	
39	QPRPSSEKI	AML, GBM	SCLC	
40	FPRTVKHIDAAL	CRC, GC, NSCLCaden, PACA, UBC	HNSCC, NSCLCsquam, OSCAR	
41	MPAGGGPRSL	CCC, HCC, NHL, UEC	CLL	
42	APLKMLAL		NHL	
43	APTPRPKVL	PRCA, UEC	BRCA	

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
44	SPSQTVQRAV		NHL	
45	TRPWSGPYIL		MEL	
46	QPISGNPVTI	NHL	CLL	
47	RPRQTGALM		AML	
48	RPRYSIGL	PRCA, UEC	BRCA	
49	APEKARAFL	GBC, HCC, HNSCC, NSCLCadenoc, NSCLCother, SCLC, UBC, UEC	CRC, GC, PRCA	
50	PEKARAFL	GBC, HCC, HNSCC, NSCLCadenoc, NSCLCother, SCLC, UBC, UEC	CRC, GC, PRCA	
51	SPVFYVQTL	HCC	BRCA, PRCA	
52	KEDNPSGHTYTL	BRCA, MEL, NSCLCadenoc, NSCLCsquam, OC, SCLC	GBC, HCC, NHL	
53	SPRIPFSTF		MEL	
54	VPSCGRSVEGL	CCC, NSCLCother, UBC	HNSCC, NSCLCsquam, OSCAR, RCC	
55	LPALLRSGI	GBC, NHL, OSCAR	CCC	
56	LPALLRSGITL	GBC, NHL, OSCAR	CCC	
57	APLLPIRTI		UBC	
58	APLLPIRTIPL		UBC	
59	KPRTIYSSL	HNSCC, NSCLCsquam	SCLC, UEC	
60	RPYSIYPHGVT	CCC	HCC	
61	LPRIPIFSTF		MEL	
62	KPQSTISGL	CCC	HCC	
63	FPHMATTAF	NSCLCother, OSCAR	HNSCC	
64	VPRPIFSQLYL	HCC, OC, SCLC, UEC	BRCA, PRCA	
65	FPNVYSTLV	HCC	MEL	
66	LPMTVISV	PACA, UEC	GC	
67	VPVSRPV	NHL	CLL	
68	FPNEVSVV	NSCLCadenoc	MEL	
69	RPEDGRPRL	SCLC	GBM	
70	VPAQRLGLL	NSCLCother	CLL	

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
71	APFAPLHGGGSL		PRCA	
72	APCNGNVAV	BRCA, GBC, MEL, OC	NSCLCadenoc, UBC	
73	LPVSSPVTL		GBM	
74	VPVSHPV		CLL, NHL	
75	HVRPQTNCI	HCC, NSCLCother, NSCLCsquam, SCLC	GBM	
76	KPKVESQAL	PRCA	BRCA	
77	QPRLVPGETL		MEL	
78	HPSQESPTL	SCLC	UEC	
79	GPASDHPSL	BRCA, HCC, OC, UEC	PRCA	
80	SALPTTISF	BRCA, CRC, GC, UEC	HCC, NSCLCadenoc	GBC, HNSCC, MEL, NSCLCsquam, OC, OSCAR, SCLC, UBC
81	IAYPSLREAAL	BRCA, CRC, GC, UEC	HCC, NSCLCadenoc	GBC, HNSCC, MEL, NSCLCsquam, OC, OSCAR, SCLC, UBC
82	DAAHPGPSV		OC	UEC
83	AVSSHNNVI	BRCA, GBC, HCC, NHL, OC, SCLC, UBC		PRCA
84	MPMQDIKMI	AML, BRCA, GC, NHL, OSCAR, UBC	GBC, HNSCC, NSCLCadenoc, NSCLCsquam, RCC	MEL, OC, SCLC, UEC
85	MPMQDIKMILKM	AML, BRCA, GC, NHL, OSCAR, UBC	GBC, HNSCC, NSCLCadenoc, NSCLCsquam, RCC	MEL, OC, SCLC, UEC
86	ALLLRGVTL	NSCLCsquam, SCLC	NSCLCadenoc	RCC
87	APVGGNVTSSF	AML, HCC, HNSCC, MEL, OSCAR	GBC, NHL	NSCLCsquam, OC
88	KPSAVKDSIY	OC	PRCA, SCLC	UEC
89	FLIPRAGWLAGL			PRCA
90	HAIEKAFQL	CRC, MEL, NSCLCadenoc, PACA	GC, NSCLCsquam, OSCAR, UBC	HNSCC

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
91	FPRLVGPDDF	HCC, OC	BRCA, CCC, CRC, GBC, GC, HNSCC, NSCLCadenoc, NSCLCsquam, OSCAR, PACA, UBC, UEC	
92	TSPLPPTV		RCC	
93	SVAIAHGVF	CRC	SCLC	
94	LPMSKRQEY	BRCA, CRC, NSCLCsquam, OC, OSCAR, PACA, UBC	SCLC, UEC	
95	RTKEEINEL	HNSCC, NSCLCother, NSCLCsquam, PACA, SCLC	BRCA, GBC, MEL, NSCLCadenoc, OC, UBC	
96	QPSLVQAIF	MEL	AML	
97	LPPGTQIQI	SCLC	GBM	
98	FPCSALLACF	BRCA	RCC	
99	MPVSAFTVI	CRC, NSCLCsquam, OC, UBC, UEC	BRCA, CCC, GBC, GC, HNSCC, NSCLCadenoc, OSCAR, PACA	
100	TPIPFDKILY	CRC, NSCLCsquam, OC, UBC, UEC	BRCA, CCC, GBC, GC, HNSCC, NSCLCadenoc, OSCAR, PACA	
101	KPGLPGLK	CRC, NSCLCsquam, OC, UBC, UEC	BRCA, CCC, GBC, GC, HNSCC, NSCLCadenoc, OSCAR, PACA	
102	MAGPAIHTAPM	SCLC	GBM	
103	REPIMKADML	BRCA, MEL, NSCLCadenoc, NSCLCsquam, OC, SCLC	GBC, HCC, NHL	
104	RPLPNSVIHV		HCC	

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
105	RPRYETGVCA	CRC, GC, PACA	HNSCC, MEL, OC, OSCAR, UBC	
106	APFHIDRLF	NSCLCsquam, PACA, UEC	GC	
107	GQRLESPGGAA	GBM	SCLC	
108	APRGSPPI	NHL	CLL	
109	LPRALMRST		NHL	
110	YPSSPASAV		HCC	
111	YPLQQTENVY	GBM	RCC	
112	YPSPLNKHSF	GBM	RCC	
113	KPHLDRRGAVI		GBC, HCC	
114	FVPPSGPSNPM		MEL	
115	KTKSLAQQ	GBC, GC, NSCLCadeno, NSCLCother, PACA	CCC, HNSCC, NSCLCsquam, OSCAR, UBC	
116	RSYQHSLRL	GBC, GC, NSCLCadeno, NSCLCother, PACA	CCC, HNSCC, NSCLCsquam, OSCAR, UBC	
117	IPHQRSSL	PRCA	AML	
118	NPERHKPSF		AML, OC, RCC, UEC	
119	KATEYVHSL	GBM, MEL, OC, SCLC, UBC	AML, UEC	
120	TVRPKNAAL	GBM, MEL, OC, SCLC, UBC	AML, UEC	
121	GPFQRAAL		UEC	
122	KPRTPTTAQLL		UEC	
123	RPRLRDLPAL	GBC, NHL, OSCAR	CCC	
124	KTIDGHINL	AML, BRCA, CCC, CRC, GBC, GBM, GC, HCC, HNSCC, MEL, NSCLCadeno, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, UBC, UEC	CLL, NHL, SCLC	
125	SPAKQFNIY	AML, BRCA, CCC, CRC, GBC, GBM, GC,	CLL, SCLC	

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
		HCC, HNSCC, MEL, NHL, NSCLCaden, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, UBC, UEC		
126	MPREDAHF		MEL	
127	KSKQVITLL		MEL	
128	SPPATLFLFLL	CCC	MEL	
129	MTLPATTLTL		NHL	
130	GAYDKARYL	HCC, PRCA	BRCA	
131	MPFRNIRSIL	BRCA, OC, SCLC	CCC, HCC	
132	LPRLPVGSLV	BRCA, CRC, GBC, GC, MEL, NHL, NSCLCother, NSCLCsquam, OC, UEC	HNSCC, NSCLCaden, OSCAR, PACA	
133	LPELPTTLTF	BRCA, CRC, GBC, GC, HNSCC, MEL, NHL, NSCLCsquam, OC, UEC	NSCLCaden, OSCAR, PACA	
134	VAAAARLTL		MEL	
135	MPRLRLSL	GBC, HCC, NHL, OC, SCLC, UEC	BRCA	
136	QPDHAGIFRVF	NSCLCother, OSCAR	HNSCC	
137	SPQLSYFEY	GC, NSCLCaden, NSCLCother, PACA, UBC	CCC, HNSCC, NSCLCsquam, OSCAR	
138	VPFNLITEY	HCC	MEL	
139	LVILQHQAM	PACA, UEC	GC	
140	TPFPDIHWF	NSCLCaden	MEL	
141	YGPYGSWSGW	NHL	HNSCC, MEL	
142	HPFKMKWQY	OSCAR	RCC	
143	YPMIPPAQL	SCLC	GBM	
144	RPVQGIPTY		MEL	
145	VPAQRLGLLL	NSCLCother	CLL	
146	SPTRGSEF		GBM	
147	HVAQPAVVL	NHL	CLL	
148	KPHLIYRQ	BRCA, OC, UEC	SCLC	

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
149	SPSSVFTSK		BRCA	
150	APLHGGGSL		PRCA	
151	QPTWNTQTRL	NHL	CLL	
152	SSASFSSSEFRH		GBM	
153	QPVHSLCSA		GC, PACA	
154	RPPPSRRAVL		GBM	
155	APIPRLIEG	HCC, OSCAR	HNSCC	
156	GSRPVTGSV	BRCA, GBC, MEL, OC	NSCLCadenoc, UBC	
157	RPVTGSVC	BRCA, GBC, MEL, OC	NSCLCadenoc, UBC	
158	SVPVATVEL		NHL	
159	TPMTVPRI	NHL	CLL	
160	IPVSHPLVTF		CLL, NHL	
161	ASKILETSL		MEL	
162	IPIRVDQNGAF	AML, GBM	CLL	
163	FRYPNGVSL		HCC	
164	RAAGRRLQL		HCC	
165	RPSKEMQVTI	CCC, GC	PACA	
166	YAYTSRVIL	AML	CLL, NHL	
167	NVNPARKGL	OC, OSCAR	AML	
168	SPSGQRDVSL		MEL	
169	RPFSVSSISQL		MEL	
170	APEGKRLGF		MEL	
171	LPLGGHKSL		MEL	
172	SAQSLHPVL		GBM	
173	VIINNPISL		MEL	
174	IPVTSSVTL		MEL	
175	AAFPHKIIF		RCC	
176	QPLDICHSF	CCC, GBC, GC, HCC, HNSCC, NSCLCadenoc, NSCLCsquam, OC, OSCAR, PACA, UBC	CRC, MEL	
177	VWEPQGSSRSL		UEC	
178	VPYYPRANL	CRC, GC, HNSCC, NSCLCadenoc, NSCLCsquam, OC, OSCAR, UBC	GBC	

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
179	PRVRLLLL	AML	SCLC	
180	RLRDYISSL	NHL	CLL	
181	LPVSPARAL	CCC, CRC, GBC, GBM, GC, MEL, NHL, NSCLCaden, NSCLCother, NSCLCsquam, OC, PACA, UBC, UEC	BRCA, HNSCC, OSCAR	
182	RPLPVSPARAL	CCC, CRC, GBC, GBM, GC, MEL, NHL, NSCLCaden, NSCLCother, NSCLCsquam, OC, PACA, UBC, UEC	BRCA, HNSCC, OSCAR	
183	VPRRPARAL		GBM	
184	KIIASGNHL	BRCA, CLL, NHL, RCC	UBC	
185	RPVLTASSF	NHL	CLL	
186	VPLPAGGGTVLT		SCLC	
187	APPPPPPPF	AML, CLL, NHL	PRCA	
188	HAAASFETL	MEL	CLL, NHL	
189	QPQCSTHIL		HCC	
190	RLAHVRARL	BRCA, HCC, OC	PRCA	
191	KPKAVKPKAA	CLL	NHL	
192	IPFADVKSF	NSCLCsquam	NSCLCaden, NSCLCother	
193	LPALKEEAF		PRCA	
194	HPNKKIKASL		BRCA	
195	NPIFGRKEL		UEC	
196	RPSGTAGAAL		GBM	
197	RPSGTAGAALLA		GBM	
198	LPSPAHAFRAL	HCC, MEL, NHL, OC, PACA, UEC	UBC	
199	SPFWIHQA		AML	
200	EPFHLIVSY	MEL	CLL, NHL	
201	LPIARVLTV		BRCA	
202	SPSREASAL		AML	
203	KPRGPIDI	MEL	PRCA	
204	FPYSGDKILV		GBM	

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
205	LPPALLTTV		GBM	
206	TPRIGPKVSL		GBM	
207	VPSDITTAF		GBM	
208	RNRQVATAL	AML	CCC	
209	KIEQIRAVL	CCC, GBC, GC, HNSCC, NSCLCaden, NSCLCsquam, PACA, UBC	OSCAR	
210	IPENRVVSY		CRC	
211	IPDTIASVL		MEL	
212	VPYAAQAAL	CRC, GBC, SCLC, UEC	HCC	
213	RPYQDAPVA	NHL, RCC	GBM	
214	LPLKFFPII	CCC, GBC, GC, HCC, NSCLCaden, PACA, SCLC	OC	
215	IPVAIKEL		GBM	
216	LPWEQNEQV		HCC	
217	SPGDKRLAAYL		HCC	
218	VQRTPPPI	CCC, CRC, GBC, GC, PACA	RCC, SCLC	
219	VPHTGRYTCL		MEL	
220	RPAPGPAPFV	GBC, HNSCC, MEL, NSCLCaden, OSCAR, PACA, UBC, UEC	CRC, GC	
221	LPQRPNARL	CCC, GBC, GC, NSCLCaden, NSCLCoher, NSCLCsquam, PACA, UBC	HNSCC, OSCAR	
222	MLKTTLTAF		HCC	
223	KAHVRIEL	HCC	CRC	
224	SPIIHSILL	BRCA, HNSCC, NSCLCaden, NSCLCsquam, OC, UEC	SCLC	
225	SPIIHSIL	BRCA, HNSCC, NSCLCaden, NSCLCsquam, OC, UEC	SCLC	

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
226	APGGSSRSSL		AML	
227	RPGTGQGGL		AML	
228	RPTAASQSRAL		AML	
229	FPNAGPRHLL	NHL	CLL	
230	DVIDDIISL		MEL	
231	SPITLQAL		HCC	
232	TAYPSLRLL	HNSCC, MEL, NSCLCsquam	SCLC	
233	MAYDRFIAI	BRCA, GBC, HCC, HNSCC, MEL, NSCLCaden, NSCLCother, NSCLCsquam, PACA, SCLC	NHL	
234	HPRAPGEQQRL	HCC	CRC	
235	AQNPAGTAL		MEL	
236	TPELKSSIL		HCC	
237	LPRAGGAF	CCC, GC, HNSCC, NSCLCaden, NSCLCsquam, PACA, UBC	OSCAR	
238	LPRAGGAFLM	CCC, GC, HNSCC, NSCLCaden, NSCLCsquam, PACA, UBC	OSCAR	
239	VLPRAGGAFLM	CCC, GC, HNSCC, NSCLCaden, NSCLCsquam, PACA, UBC	OSCAR	
240	RVMLPKAAL	BRCA, CCC, NHL, NSCLCsquam, OC, OSCAR, SCLC, UBC, UEC	GBC	
241	LPKAALLV	BRCA, CCC, NHL, NSCLCsquam, OC, OSCAR, SCLC, UBC, UEC	GBC	
242	IPETASVVAI	MEL	CLL, NHL	
243	MARTGGMVVI		UBC	
244	VPAHLVAA		RCC	

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
245	GPVPSPLAL		CLL	
246	RPILKEQSSSF		HNSCC	
247	SPVGVGQRL	GBM	SCLC	
248	KPYDGIPAS		GBM	
249	SPRSGVLL	OC	UEC	
250	APAAPAAVPS	CRC, GBM, GC, HNSCC, SCLC	NSCLCadenocarcinoma, NSCLCsquamous	
251	MPVDSFNSM	CCC, GBC, NSCLCadenocarcinoma, UEC	CRC	
252	QPENSLEGISL	CCC, GBC, NSCLCadenocarcinoma, NSCLCother, PACA, RCC, UEC	CRC	
253	MPVDSFNML	CCC, GBC, NSCLCadenocarcinoma, UEC	CRC	
254	RVIQGTTL	NHL	CLL	
255	VPSHWMVAL	CLL	NHL	
256	SPVPSHWMVAL	CLL	NHL	
257	APYGTLRKS		CCC	
258	SVIGPNSRL	NHL	CLL	
259	VPMPGVQAV	CRC	UEC	
260	LVQSSRSEV	OSCAR	HNSCC	
261	SPSTSRTPL		GBM	
262	SPSTSRTPLLSSL		GBM	
263	SQRPPATSQA	OC	NHL	
264	APRPGNWIL	MEL, NSCLCsquamous, OC, RCC, SCLC	UEC	
265	FPRKPYEGRV		MEL	
266	KPYEGRVEI		MEL	
267	MPVPGILL		MEL	
268	EPLSVTASY	GBC, SCLC	HCC	
269	FTVSSSSAM		PACA	
270	SPRGTTSTL		PACA	
271	SPTPVFTTL		PACA	
272	SSPRGTTSTL		PACA	
273	TDTPSTPTTI		PACA	
274	KPIRTGISPLAL		HCC	
275	IPAPQGAVL		NSCLCother	

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
276	RPVWDVRS	NSCLCadeno	HNSCC	
277	MPPLLIVAF	UBC	BRCA	
278	LIAARGSLVL	GBM, HCC, MEL, OC	SCLC	
279	APADEPRTVL	BRCA, CCC, CRC, GBC, GC, HNSCC, NHL, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, PRCA, SCLC, UBC, UEC	GBM	
280	LPRAFLQSL	BRCA, CCC, CRC, GBC, GC, HNSCC, NHL, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, PRCA, SCLC, UBC, UEC	GBM	
281	NPRSPATM	HCC, UEC	UBC	
282	AVRLPTPNL	AML		
283	RPQPGWRESL	GBC, SCLC		
284	RPSAPRGPP	CRC		
285	RITWKGLLL	CCC		
286	APARPAAAF	MEL, OC		
287	SPIRPLFL	BRCA, SCLC		
288	LHAMNSLSAM	CCC, OC, SCLC, UBC, UEC		
289	LPYEKVSRL	CCC		
290	RPTHPLRSF	GBM		
291	SPSKSLEM	CLL, CRC		
292	LPMTHRLQL	CCC		
293	FPYDKPLIM	CCC, GBC, NSCLCadeno		
294	VPKPAIPSSSVL	GBM, NSCLCsquam, SCLC		
295	HPRWIEPTVM	PRCA		
296	SPLLMQRTL	AML, CLL		
297	FPIKYVNDF	CCC, GBC, NSCLCadeno		
298	RVLLRWISL	GBM		
299	SPFSGGPVSF	RCC		
300	HPYSDLADV	SCLC		

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
301	FPAFLEAM	SCLC		
302	IPIDQILNSF	SCLC		
303	RPPPPCIAL	SCLC		
304	SPLIGDFPAF	SCLC		
305	AASPVGSAL	HCC		
306	RPFPLALL	CRC		
307	RPHQKGWLSI	HCC		
308	RPDVVRTLL	CCC, HNSCC, OSCAR		
309	FAFYGGKSL	GBM, HCC, HNSCC, NSCLCadenoc, NSCLCsquam, OSCAR, SCLC, UBC		
310	SPGWAQTQL	CRC, GC, NSCLCadenoc, PACA, RCC		
311	APRLALDPDAL	GBM, SCLC, UEC		
312	SPSLQSSRESL	BRCA, CCC, CRC, GBC, GBM, GC, HNSCC, MEL, NHL, NSCLCadenoc, NSCLCsquam, OSCAR, PACA, SCLC, UBC, UEC		
313	VPLSSYVSI	GBM		
314	IALMKLAGPL	CCC, GBC, GC, HCC, OC, PACA, SCLC		
315	APVVFPAL	SCLC		
316	NPREPEKSL	CRC, GBC, GC, PACA		
317	MPYNSAHHCVV	CLL, OC		
318	TPISNTGVL	CLL, NHL		
319	RPLDTFRVL	PRCA, UBC		
320	APMHIDRNIL	HCC		
321	QPQQPGFFL	AML		
322	RAVPVGSGSL	SCLC		
323	TPHGITDIL	GC		
324	LPAPLRISL	GBM		
325	SPRSNPVRL	NHL		
326	IPPFTQRVF	HNSCC, NSCLCsquam		

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
327	GPRTTQSSVL	CLL, NHL		
328	LPLHRGDLVI	CLL		
329	QPANFIVLL	GBC, GC, PACA		
330	RPFSAIAEL	CCC		
331	SPDSAILKL	PRCA		
332	SPYAGSTAF	BRCA		
333	SVLPRALSL	SCLC		
334	YPLSLALHF	SCLC		
335	VPPQNPRPSL	AML		
336	YPLQGPGLLSV	NSCLCother, SCLC		
337	IPTSRVITL	HCC, NHL		
338	MPATPSLKV	CCC, CRC		
339	GPQRTTSV	NHL		
340	APEPRAALL	MEL		
341	LPRSPPLKVL	GBC, NHL		
342	RPRPPKVL	HCC, NSCLCother, NSCLCsquam		
343	RPRPPKVLGL	HCC, NSCLCother, NSCLCsquam		
344	VPYPSPTCV	HCC, PRCA		
345	SAAPPGASL	HCC, PRCA		
346	IPMPRITWL	UBC		
347	SPLEKINSF	MEL, SCLC		
348	HPAPPVTSA	HNSCC, OC, OSCAR, UBC		
349	QPRDGWPMML	HNSCC, OC, OSCAR, UBC		
350	RPKSTLMNF	HNSCC, OC, OSCAR, UBC		
351	SPYADIIPSA	BRCA, MEL		
352	LPAFSKIGGIL	SCLC		
353	KPRATWTL	HNSCC, OC		
354	APAKDARASL	GBM		
355	APKTSFIAA	CRC, GC, NSCLCadeno, PACA		
356	RPFLRLPSL	CRC, HNSCC, OSCAR		
357	LPPHIFAI	MEL		
358	APSMLRKNQL	BRCA, GBM, MEL,	GBC, UBC	HCC

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
		NSCLCadeno, OSCAR, SCLC		
359	SPRRLVELAGQSL	AML, BRCA, GC, HCC, NHL, OSCAR, UBC	GBC, HNSCC, NSCLCadeno, NSCLCsquam, RCC	MEL, OC, SCLC, UEC
360	SPASRSISLL	AML, GC, HNSCC, OC, OSCAR	CLL, NHL	RCC
361	SPYGSDRLVQL	NHL	CLL	
362	KPMLPPAAF		UEC	
363	KPRTPFHTTA		UEC	
364	KPRTPFHTTAQL		UEC	
365	RPKHFLQML	HNSCC, NHL, NSCLCsquam, OSCAR	OC	
366	SPTLRQLDL	HCC	NHL	
367	APQVHIFSL	GBM, OC	CCC	
368	NPASRLTAL	PRCA, UEC	BRCA	
369	RPYGCVLRAA	AML, CCC, GBC, GC, HNSCC, OC, OSCAR	CLL, NHL	RCC
370	AAHEFGHVL	HCC, NSCLCother, OC, RCC, SCLC	BRCA, CCC, CRC, GBC, GC, HNSCC, NSCLCadeno, NSCLCsquam, OSCAR, PACA, UBC, UEC	
371	APRSPGQVTPRGL	HCC, NSCLCadeno, NSCLCsquam, PACA, SCLC	CRC, GBC, GC, RCC	
372	SPSSASLTL	GBC, GC, HCC, MEL, NSCLCadeno, NSCLCother, PACA, UEC	HNSCC, NSCLCsquam, OC, OSCAR, SCLC, UBC	
373	LPKPDLPQLI	BRCA, HCC	PRCA	
374	KPRNMTGLDL	NHL	CLL	
375	LPRGVLEGL	BRCA, CCC, CRC, GBC, GC, HNSCC, NHL, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, PRCA,	GBM	

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
		SCLC, UBC, UEC		
376	FPQVGRTAL	CLL		
377	SPFSKRIKL	AML, CLL, NHL		
378	SPRQPRLDF	NSCLCsquam, SCLC		
379	GPRQVLFPL	GBM		
380	SPYPGSAAF	BRCA, GC		
381	APRPRLLLL	CCC, GBC, MEL		
382	RPGPQRTTSV	NHL		
383	KPRATWTLKL	HNSCC, OC		

Table 9b: Expression scores. The table lists peptides from genes that are very highly over-expressed in tumors compared to a panel of normal tissues (+++), highly over-expressed in tumors compared to a panel of normal tissues (++) or over-expressed in tumors compared to a panel of normal tissues (+). The baseline for this score was calculated from measurements of the following relevant normal tissues: adipose tissue, adrenal gland, bile duct, blood cells, blood vessels, bone marrow, brain, esophagus, eye, gallbladder, heart, head&neck, kidney, large intestine, liver, lung, lymph node, nerve, parathyroid, pancreas, pituitary, pleura, skeletal muscle, skin, small intestine, spleen, stomach, thyroid gland, trachea, ureter, urinary bladder. In case expression data for several samples of the same tissue type were available, the arithmetic mean of all respective samples was used for the calculation. AML: acute myeloid leukemia; BRCA: breast cancer; CCC: cholangiocellular carcinoma; CLL: chronic lymphocytic leukemia; CRC: colorectal cancer; GBC: gallbladder cancer; GBM: glioblastoma; GC: gastric cancer; HCC: hepatocellular carcinoma; HNSCC: head and neck squamous cell carcinoma; MEL: melanoma; NHL: non-Hodgkin lymphoma; NSCLCaden: non-small cell lung cancer adenocarcinoma; NSCLCother: NSCLC samples that could not unambiguously be assigned to NSCLCaden or NSCLCsquam; NSCLCsquam: squamous cell non-small cell lung cancer; OC: ovarian cancer; OSCAR: esophageal cancer; PACA: pancreatic cancer; PRCA: prostate cancer; RCC: renal cell carcinoma;

SCLC: small cell lung cancer; UBC: urinary bladder carcinoma; UEC: uterine and endometrial cancer

Seq ID No	Sequence	Gene Expression in tumor samples		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
448	TVYGEPRKL	BRCA, CRC, GC, UEC	HCC, NSCLCadeno	GBC, HNSCC, MEL, NSCLCsquam, OC, OSCAR, SCLC, UBC
449	RPKTSVNLISL	HCC, NHL, RCC	CCC, GBC, GC, MEL, NSCLCadeno, OC, OSCAR, PACA, UBC, UEC	CRC, HNSCC, NSCLCother, NSCLCsquam, SCLC
450	SAAARALLP	HCC, MEL, RCC, SCLC	BRCA, CRC, GBC, GC, HNSCC, NSCLCadeno, NSCLCsquam, OC, OSCAR, UBC, UEC	CCC, PACA
451	VPILPLNHAL	BRCA	UEC	
452	QPVKKNTL	HCC, UEC	RCC	
453	APALPGQVTI	CCC, CRC, GBC, HCC, MEL, NSCLCadeno, NSCLCother, NSCLCsquam, PACA, PRCA	BRCA, GBM, OC, SCLC, UBC, UEC	
454	SPDPAHLESL	BRCA, GC, NSCLCadeno, OSCAR, PACA, UEC	GBC, HCC, HNSCC, MEL, NSCLCsquam, OC, SCLC, UBC	
455	FPVQATIDF	BRCA, CRC, NSCLCsquam, OC, OSCAR, PACA, UBC	SCLC, UEC	
456	VPVSHPVLTl		CLL, NHL	
457	RPPLSQRHTF	NHL	CLL	
458	VIPIPTHYFVVL		RCC	

459	SPAPWRPWI	BRCA, NSCLCsquam, OC, OSCAR, PRCA, UBC, UEC	SCLC	
460	APLMPLGKTL	GBC, GC, NSCLCadenos, NSCLCothers, PACA, UEC	CCC, HNSCC, NSCLCsquam, OSCAR, UBC	
461	MPHLGPGILL		MEL	
462	MPLLADVRL		MEL	
463	LPTDLFNSVM		MEL	
464	VPFVPRTSV	BRCA, GC, HNSCC, MEL, NHL, NSCLCadenos, NSCLCothers, NSCLCsquam, OC, OSCAR, PACA, RCC, UBC, UEC	CCC, CRC, GBC	
465	VPKSLPYVL	HNSCC, NSCLCsquam, OSCAR	AML, HCC, NHL	
466	SPMEAILVSRL	CRC, NSCLCadenos, NSCLCsquam, SCLC	RCC	
467	VPRNQDWLGVSRL		MEL	
468	LPSLHVLVL		MEL	
469	LPLDTRTSI	NHL	CLL	
470	RPNGEVKSEL	RCC, SCLC	CRC	
471	LPLLAGTLLL		UEC	
472	LPMPAITWY		MEL	
473	LPGEREAAL		MEL	
474	VPKADLLTL		MEL	
475	IPLEIQKL		HCC	
476	EPNPVEEIF	CCC, CRC, GBM, GC, HNSCC, NSCLCothers, NSCLCsquam, OC, OSCAR	BRCA, GBC, MEL, NSCLCadenos, PACA	
477	NPVPVITWYKDNRL		MEL	
478	APKFISPASQL	RCC, SCLC	CRC	
479	APHAGGALL	GBC, HNSCC, MEL, NSCLCadenos, NSCLCsquam, OC, SCLC, UEC	NHL	

480	VPAAPTKAL	UBC	GBM	
481	SPARALLLALA	CCC, CRC, GBC, GBM, GC, MEL, NHL, NSCLCadenos, NSCLCother, NSCLCsquam, OC, PACA, UBC, UEC	BRCA, HNSCC, OSCAR	
482	QPSLKKIIL	CCC, CLL, GBC, GC, HCC, NSCLCsquam, PACA	NSCLCadenos, UEC	
483	KPAAAGVKKVA	CLL	NHL	
484	LPASAEPKGTM	CCC, CRC, GBC, GC, HCC, HNSCC, NSCLCadenos, NSCLCother, NSCLCsquam, OC, PACA	OSCAR, UBC	
485	LPLKTKVFA	CCC, NSCLCadenos, NSCLCother, PACA, RCC	BRCA	
486	TPTRPLPSA	CCC, CRC, GBC, GC, PACA	RCC, SCLC	
487	IPWFIKTAF		MEL	
488	LPLGGLPLLI	NHL	CLL	
489	WPNHIMLVL	BRCA, MEL, OC	PRCA	
490	APRGPAQGEAA		AML	
491	MPEADLKRIF		MEL	
492	IPFSVDEI	CCC, GBC, NSCLCadenos, UEC	CRC	
493	LPLQQYKLV	CRC, GBM, GC, HNSCC, NSCLCsquam, SCLC	NSCLCadenos	
494	LPLRAVNLNL	GBC, GBM, MEL, NSCLCsquam, OC, OSCAR	SCLC	
495	SPSYTQASL		BRCA	
496	MPAVSGPGPLF		PRCA	
497	YVVKPLHPF	NSCLCother, NSCLCsquam	NSCLCadenos	
498	RPQPQPRPAL	CLL, NHL, OC, OSCAR, UBC	CCC, GBC, HNSCC	
499	QPVLPSPAC		MEL	

EXAMPLE 3

In vitro immunogenicity for MHC class I presented peptides

In order to obtain information regarding the immunogenicity of the TUMAPs of the present invention, the inventors performed investigations using an *in vitro* T-cell priming assay based on repeated stimulations of CD8+ T cells with artificial antigen presenting cells (aAPCs) loaded with peptide/MHC complexes and anti-CD28 antibody. This way the inventors could show immunogenicity for HLA-B*07 restricted TUMAPs of the invention, demonstrating that these peptides are T-cell epitopes against which CD8+ precursor T cells exist in humans (Table 10a and 10b).

In vitro priming of CD8+ T cells

In order to perform *in vitro* stimulations by artificial antigen presenting cells loaded with peptide-MHC complex (pMHC) and anti-CD28 antibody, the inventors first isolated CD8+ T cells from fresh HLA-B*07 leukapheresis products via positive selection using CD8 microbeads (Miltenyi Biotec, Bergisch-Gladbach, Germany) of healthy donors obtained from the University clinics Mannheim, Germany, after informed consent.

PBMCs and isolated CD8+ lymphocytes were incubated in T-cell medium (TCM) until use consisting of RPMI-Glutamax (Invitrogen, Karlsruhe, Germany) supplemented with 10% heat inactivated human AB serum (PAN-Biotech, Aidenbach, Germany), 100 U/ml Penicillin/100 µg/ml Streptomycin (Cambrex, Cologne, Germany), 1 mM sodium pyruvate (CC Pro, Oberdorla, Germany), 20 µg/ml Gentamycin (Cambrex). 2.5 ng/ml IL-7 (PromoCell, Heidelberg, Germany) and 10 U/ml IL-2 (Novartis Pharma, Nürnberg, Germany) were also added to the TCM at this step.

Generation of pMHC/anti-CD28 coated beads, T-cell stimulations and readout was performed in a highly defined *in vitro* system using four different pMHC molecules per stimulation condition and 8 different pMHC molecules per readout condition.

The purified co-stimulatory mouse IgG2a anti human CD28 Ab 9.3 (Jung et al., 1987) was chemically biotinylated using Sulfo-N-hydroxysuccinimidobiotin as recommended

by the manufacturer (Perbio, Bonn, Germany). Beads used were 5.6 μm diameter streptavidin coated polystyrene particles (Bangs Laboratories, Illinois, USA).

pMHC used for positive and negative control stimulations were A*0201/MLA-001 (peptide ELAGIGILTV (SEQ ID NO. 446) from modified Melan-A/MART-1) and A*0201/DDX5-001 (YLLPAIVHI from DDX5, SEQ ID NO. 447), respectively.

800.000 beads / 200 μl were coated in 96-well plates in the presence of 4 x 12.5 ng different biotin-pMHC, washed and 600 ng biotin anti-CD28 were added subsequently in a volume of 200 μl . Stimulations were initiated in 96-well plates by co-incubating 1×10^6 CD8⁺ T cells with 2×10^5 washed coated beads in 200 μl TCM supplemented with 5 ng/ml IL-12 (PromoCell) for 3 days at 37°C. Half of the medium was then exchanged by fresh TCM supplemented with 80 U/ml IL-2 and incubating was continued for four days at 37°C. This stimulation cycle was performed for a total of three times. For the pMHC multimer readout using 8 different pMHC molecules per condition, a two-dimensional combinatorial coding approach was used as previously described (Andersen et al., 2012) with minor modifications encompassing coupling to five different fluorochromes. Finally, multimeric analyses were performed by staining the cells with Live/dead near IR dye (Invitrogen, Karlsruhe, Germany), CD8-FITC antibody clone SK1 (BD, Heidelberg, Germany) and fluorescent pMHC multimers. For analysis, a BD LSRII SORP cytometer equipped with appropriate lasers and filters was used. Peptide specific cells were calculated as percentage of total CD8⁺ cells. Evaluation of multimeric analysis was done using the FlowJo software (Tree Star, Oregon, USA). *In vitro* priming of specific multimer⁺ CD8⁺ lymphocytes was detected by comparing to negative control stimulations. Immunogenicity for a given antigen was detected if at least one evaluable *in vitro* stimulated well of one healthy donor was found to contain a specific CD8⁺ T-cell line after *in vitro* stimulation (i.e. this well contained at least 1% of specific multimer⁺ among CD8⁺ T-cells and the percentage of specific multimer⁺ cells was at least 10x the median of the negative control stimulations).

In vitro immunogenicity for acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer peptides.

For tested HLA class I peptides, *in vitro* immunogenicity could be demonstrated by generation of peptide specific T-cell lines. Exemplary flow cytometry results after TUMAP-specific multimer staining for two peptides of the invention are shown in Figures 3A through 3G together with corresponding negative controls. Results for 58 peptides from the invention are summarized in Table 10a and 10b.

Table 10a: *in vitro* immunogenicity of HLA class I peptides of the invention

Exemplary results of *in vitro* immunogenicity experiments conducted by the applicant for the peptides of the invention. <20 % = +; 20 % - 49 % = ++; 50 % - 69 % = +++; >= 70 % = ++++

Seq ID No	Sequence	Wells positive [%]
388	SPSVSQLSVL	++++
389	APLPRPGAVL	++
390	SPRMSGLLSQT	+++
391	APRPASSL	+
392	GPQPWHAAL	++
393	APAAWLRSA	+++
396	VPDVAQFVL	++
398	SPASRSISL	+
399	NPFYPEVEL	+
405	LPFDGPGGIL	++++
406	LPDGSRVEL	++++
407	FPRLVGPDF	+++
408	YPKDIYSSF	++
410	SPRSWIQVQI	+
411	IPNWARQDL	++++

- 151 -

Seq ID No	Sequence	Wells positive [%]
414	KPSESIYSAL	++
415	LPSDSHFKITF	++
418	RPMTPTQIGPSL	+
419	SPMWHVQQL	++++
420	SPRWLPVSL	++++
423	SPRVYWLGL	++++
425	YPRGNHWAVGHL	++
427	VPSSRILQL	+++
429	RPRALRDLQL	++
430	RPRALRDLQLL	+++
433	IPEPSAQQQL	+
437	FPYPYAERL	+++
440	YPRITITPGM	+
443	VPDGVSKVL	+++
445	RPAATAVISL	+

Table 10b: *in vitro* immunogenicity of HLA class I peptides of the invention

Exemplary results of *in vitro* immunogenicity experiments conducted by the applicant for the peptides of the invention. <20 % = +; 20 % - 49 % = ++; 50 % - 69 % = +++; >= 70 % = ++++

Seq ID No	Sequence	Wells positive [%]
1	RPRSLQCVSL	+
3	MPYVVLTAL	+
4	GPKKFIVKL	+
5	LPSLSHCSQL	++
6	LPLNSSTSL	+
9	RPDSRLLEL	+++
12	LPNTGRIGQLL	++
17	SPAAPLSPASSL	+
21	APSRLEL	+
23	VPRPTSTVGL	++
24	VPRPTSTVGLFL	+
26	SPSFSTLLSL	+
33	LPRGLSPARQL	++
38	RPSFPNLTFS	+
45	TRPWSGPYIL	+
48	RPRYSIGL	+

Seq ID No	Sequence	Wells positive [%]
49	APEKARAFL	+
51	SPVFYVQTL	+
53	SPRIPFSTF	+
56	LPALLRSGLTL	+++
59	KPRTIYSSL	+
61	LPRIPIFSTF	++
64	VPRPIFSQLYL	+++
68	FPNEVSVVL	++
361	SPYGSDRLVQL	+
364	KPRTPFHTAQL	+
365	RPKHFLQML	+
366	SPTLRQLDL	++

EXAMPLE 4

Synthesis of peptides

All peptides were synthesized using standard and well-established solid phase peptide synthesis using the Fmoc-strategy. Identity and purity of each individual peptide have been determined by mass spectrometry and analytical RP-HPLC. The peptides were obtained as white to off-white lyophilizes (trifluoro acetate salt) in purities of >50%. All TUMAPs are preferably administered as trifluoro-acetate salts or acetate salts, other salt-forms are also possible.

EXAMPLE 5

MHC Binding Assays

Candidate peptides for T cell based therapies according to the present invention were further tested for their MHC binding capacity (affinity). The individual peptide-MHC complexes were produced by UV-ligand exchange, where a UV-sensitive peptide is cleaved upon UV-irradiation and exchanged with the peptide of interest as analyzed. Only peptide candidates that can effectively bind and stabilize the peptide-receptive MHC molecules prevent dissociation of the MHC complexes. To determine the yield of the exchange reaction, an ELISA was performed based on the detection of the light

chain ($\beta 2m$) of stabilized MHC complexes. The assay was performed as generally described in Rodenko et al. (Rodenko et al., 2006)

96 well MAXISorp plates (NUNC) were coated over night with 2ug/ml streptavidin in PBS at room temperature, washed 4x and blocked for 1h at 37°C in 2% BSA containing blocking buffer. Refolded HLA-A*02:01/MLA-001 monomers served as standards, covering the range of 15-500 ng/ml. Peptide-MHC monomers of the UV-exchange reaction were diluted 100-fold in blocking buffer. Samples were incubated for 1h at 37°C, washed four times, incubated with 2ug/ml HRP conjugated anti- $\beta 2m$ for 1h at 37°C, washed again and detected with TMB solution that is stopped with NH_2SO_4 . Absorption was measured at 450nm. Candidate peptides that show a high exchange yield (preferably higher than 50%, most preferred higher than 75%) are generally preferred for a generation and production of antibodies or fragments thereof, and/or T cell receptors or fragments thereof, as they show sufficient avidity to the MHC molecules and prevent dissociation of the MHC complexes.

Table 11: MHC class I binding scores. Binding of HLA-class I restricted peptides to HLA-B*07 was ranged by peptide exchange yield: >10% = +; >20% = ++; >50 = +++; > 75% = ++++

SEQ ID	Sequence	Peptide exchange
1	RPRSLQCVSL	++++
2	VPGSDPARYEFL	+++
3	MPYVVLTA	++++
4	GPKKFIVKL	+++
5	LPSLSHCSQL	++++
6	LPLNSSTSL	++++
7	RPSQLAPATL	+++
8	MPKTNLSKM	+++
9	RPDSRLLEL	++++
10	SPMEAEVRRIL	++++
11	APLPRPGAV	++++
12	LPNTGRIGQLL	++++
13	LPNTGRIGQL	+++
14	AVHEIGHSL	+++

15	KPGFNISIL	++++
16	FPAPPAHWF	++++
17	SPAAPLSPASSL	++++
18	MALSVLRAL	++
19	WPRLPGAGL	+++
20	MVLGIGPVL	++++
21	APSRLEL	++++
22	LPQLKPAL	++++
23	VPRPTSTVGL	+++
24	VPRPTSTVGLFL	++++
25	RPQGAVGGL	+++
26	SPSFSSTLLSL	++++
27	RIRVTSEVL	+++
28	LPAPTSVL	++++
29	APLRVHITS	++++
30	YPGFTKRL	+++
31	RPGPSDPAA	+++

32	APMAPGRSPL	++++
33	LPRGLSPARQL	+++
34	APMAPGRSP	+++
35	APLPPPRAL	+++
36	RPFSREMDL	+++
37	LPPPRALTL	+++
38	RPSFPNLTSF	++++
39	QPRPSSEKI	+++
40	FPRTVKHIDAAL	++++
41	MPAGGGPRSL	++++
42	APLKMLAL	++++
43	APTPRPKVL	++++
44	SPSQTVQRAV	++++
45	TRPWSGPYIL	+++
46	QPISGNPVT	+++
47	RPRQTGALM	++++
48	RPRYSIGL	+++
49	APEKARAFL	+++
50	PEKARAFL	+++
51	SPVFYVQTL	++++
52	KEDNPSGHTYTL	+++
53	SPRIPFSTF	++++
54	VPSCGRSVEGL	+++
56	LPALLRSGLTL	++++
59	KPRTIYSSL	++++
60	RPYSIYPHGVT	++++
61	LPRIPFSTF	++++
62	KPQSTISGL	+++
63	FPHMATTAF	+++
64	VPRPIFSQLYL	++++
65	FPNVYSTLVL	++++
66	LPMTVISVL	++++
67	VPVSRPVL	++++
68	FPNEVSVVL	++++
69	RPEDGRPRL	+++
70	VPAQRLGLL	++++
71	APFAPLHGGSL	+++
72	APCNGNVAV	+++
73	LPVSSPVT	++++
74	VPVSHPVL	++++
75	HVRPQTNCI	+++
76	KPKVESQAL	++++
77	QPRLVPGETL	++++
78	HPSQESPTL	+++
79	GPASDHPSL	+++

80	SALPTTISF	+++
81	IAYPSLREAAL	++
83	AVSSHNNVI	++
84	MPMQDIKMI	+++
85	MPMQDIKMILKM	+++
86	ALLLRGVTL	++
87	APVGGNVTSF	+++
88	KPSAVKDSIY	++
89	FLIPRAGWLAGL	++
91	FPRLVGPDDF	++++
92	TSPLPPTV	+++
93	SVAIAHGVF	++++
94	LPMSKRQEY	+++
95	RTKEEINEL	++
96	QPSLVQAIF	++++
98	FPCSALLACF	++++
99	MPVSAFTVI	++++
101	KPGLPGLK	+
102	MAGPAHTAPM	++
103	REPIMKADML	++
104	RPLPNSVIHV	+++
105	RPRYETGVCA	++++
106	APFHIDRLF	+++
108	APRGSPPI	+++
109	LPRALMRST	+++
110	YPSSPASAV	++++
111	YPLQQTENVY	+
112	YPSPLNKHSF	+++
113	KPHLDRRGAVI	++++
114	FVPPSGPSNPM	++
115	KTKSLAQQ	+++
117	IPHQRSSL	++++
118	NPERHKPSF	+++
119	KATEYVHSL	+
120	TVRPKNAAL	+++
121	GPFQRAAL	+++
122	KPRTPTTTAQLL	++++
123	RPRLRDLPAL	+++
125	SPAKQFNII	++
126	MPREDAHF	+
127	KSKQVITLL	+
129	MTLPATTLTL	++
130	GAYDKARYL	++
131	MPFRNIRSIL	+++
132	LPRLPVGSLV	++++

133	LPELPTTLTF	++
134	VAAAARLTL	+++
135	MPRLRLSL	+++
136	QPDHAGIFRVF	++++
139	LVILQHQAM	+++
140	TPFPDIHWF	++++
141	YGPYGGSGSW	++
143	YPMIPPAQL	+++
144	RPVQGIPTY	+++
145	VPAQRLGLLL	+++
146	SPTRGSEF	++++
147	HVAQPAVVL	+++
149	SPSSVFTSK	+++
150	APLHGGGSL	+++
151	QPTWNTQTRL	++++
152	SSASFSELFRH	+++
153	QPVHSLCSA	++++
154	RPPPSRRAVL	+++
156	GSRPVTGSV	+++
157	RPVTGSVC	++++
158	SVPVATVEL	++
160	IPVSHPVLTf	++
161	ASKILETSL	++
162	IPIRVDQNGAF	+++
164	RAAGRRLQL	++++
165	RPSKEMQVTI	++++
167	NVNPARKGL	++++
168	SPSGQRDVSL	++++
169	RPFSVSSISQL	++++
170	APEGKRLGF	++++
171	LPLGGHKSL	+++
172	SAQSLHPVL	+++
173	VIINNPISL	++
174	IPVTSSVTL	++++
176	QPLDICHSF	++++
177	VWEPQGSSRSL	+++
178	VPYYPRANL	+++
180	RLRDYISSL	++++
181	LPVSPARAL	+++
182	RPLPVSPARAL	+++
183	VPRRPARAL	+++
184	KIIASGNHL	++
185	RPVLTASSF	++++
186	VPLPAGGGTVLT	+
187	APPPPPPPF	++++

188	HAAASFETL	++
189	QPQCSTHIL	++++
190	RLAHVRARL	+++
191	KPKAVKPKAA	++++
192	IPFADVKSF	++
193	LPALKEEAF	++++
194	HPNKKIKASL	+++
195	NPIFGRKEL	+++
196	RPSGTAGAAL	+++
197	RPSGTAGAALLA	++
198	LPSPAHAFRAL	+++
199	SPFWIHQA	++
201	LPIARVLTV	++++
202	SPSREASAL	+++
203	KPRGPIDI	++
204	FPYSGDKILV	+++
206	TPRIGPKVSL	+++
207	VPSDITTAF	+++
208	RNRQVATAL	+++
209	KIEQIRAVL	+++
210	IPENRVVSY	++
211	IPDTIASVL	+++
212	VPYAAQAAL	++++
213	RPYQDAPVA	++++
214	LPLKFFPII	+
215	IPVAIKEL	++
216	LPWEQNEQV	+++
217	SPGDKRLAAYL	+++
219	VPHTGRYTCL	+++
220	RPAPGPAPFV	+++
221	LPQRPNARL	+++
222	MLKTTLTAF	+
223	KAHVRIEL	++
224	SPIIHSILL	++++
225	SPIIHSIL	+++
226	APGGSSRSSL	+++
227	RPGTGQGGL	+++
228	RPTAASQSRAL	+++
229	FPNAGPRHLL	+++
230	DVIDDIISL	++++
231	SPITLQAL	++++
232	TAYPSLRLL	+++
233	MAYDRFIAI	+++
234	HPRAPGEQQRL	+++
235	AQNPAGTAL	+

236	TPELKSSIL	+++
237	LPRAGGAF	++++
238	LPRAGGAFLM	++++
239	VLPRAGGAFLM	+++
240	RVMLPKAAL	+++
242	IPETASVVAI	+++
243	MARTGGMVVI	++
244	VPAHLVAA	++
245	GPVPSPLAL	++++
246	RPILKEQSSSF	++++
247	SPVGVGQRL	+++
248	KPYDGIPAS	+++
249	SPRSGVLL	+++
250	APAAPAAVPS	++
251	MPVDSFNSM	++++
252	QPENSLEGISL	+++
253	MPVDSFNML	++++
254	RVIQGTTL	++++
255	VPSHWMVAL	++++
256	SPVPSHWMVAL	++++
258	SVIGPNSRL	++
259	VPMPGVQAV	++++
260	LVQSSRSEV	++++
261	SPSTSRTPL	+++
262	SPSTSRTPLLSSL	++++
263	SQRPPATSQA	++
264	APRPGNWIL	+++
265	FPRKPYEGRV	+++
266	KPYEGRVEI	++++
267	MPVPGILL	+++
268	EPLSVTASY	++
269	FTVSSSSAM	++
270	SPRGTTSTL	+++
271	SPTPVFTTL	+++
272	SSPRGTTSTL	+++
273	TDTPSTPTTI	++
274	KPIRTGISPLAL	+++
275	IPAPQGAVL	+++
276	RPVWDVRSAL	++++
277	MPPLLIVAF	++
278	LIAARGSLVL	+++
279	APADEPRTVL	+++
280	LPRAFLQSL	++++
281	NPRSPPATM	++++
282	AVRLTPPNL	++++

283	RPQPGWRESL	++++
284	RPSAPRGPP	++++
285	RITWKGLLL	+++
286	APARPAAAF	++++
287	SPIPRPLFL	++++
288	LHAMNSLSAM	+++
289	LPYEKVSRL	+++
290	RPTHPLRSF	++++
291	SPSKSLEM	+++
292	LPMTHRLQL	+++
293	FPYDKPLIM	++++
294	VPKPAIPSSSVL	++++
295	HPRWIEPTVM	++++
296	SPLLMQRTL	++++
298	RVLLRWISL	+++
299	SPFSGGPVSF	++++
300	HPYSDLADV	+++
301	FPAFLEAM	+++
302	IPIDQILNSF	+++
303	RPPPPCIAL	++++
304	SPLIGDFPAF	++
305	AASPVGSAL	+++
306	RPFPLALL	+++
307	RPHQKGWLSI	+++
308	RPDVVRTLL	++++
309	FAFYGGKSL	+++
310	SPGWAQTQL	++++
311	APRLALDPDAL	++++
312	SPSLQSSRESL	+++
313	VPLSSYVSI	++++
314	IALMKLAGPL	++
315	APVVFPAL	++++
316	NPREPEKSL	+++
317	MPYNSAHHCVV	++++
318	TPISNTGVL	++++
319	RPLDTFRVL	++++
320	APMHIDRNIL	++++
321	QPQQPGFFL	+++
322	RAVPVGSGL	+++
323	TPHGTDIL	++++
324	LPAPLRISL	++++
325	SPRSNPVRL	++++
327	GPRTTQSSVL	++++
328	LPLHRGDLVI	+++
329	QPANFIVLL	+++

330	RPFSAIAEL	++++
331	SPDSAILKL	+++
332	SPYAGSTAF	++++
333	SVLPRALSL	+++
334	YPLSLALHF	++++
335	VPPQNPRPSL	+++
336	YPLQGPGLLSV	+++
337	IPTSRVITL	++++
338	MPATPSLKV	+++
339	GPQRTTSV	+++
340	APEPRAALL	+++
341	LPRSPPLKVL	++++
342	RPRPPKVL	+++
343	RPRPPKVLGL	+++
344	VPYPSPTCV	++++
345	SAAPPGASL	+++
346	IPMPRITWL	++++
347	SPLEKINSF	++++
348	HPAPPV TSA	+++
349	QPRDGWPMML	++++
350	RPKSTLMNF	++++
351	SPYADIIPSA	+++
352	LPAFSKIGGIL	++++
353	KPRATWTL	+++
354	APAKDARASL	++++
355	APKTSFIAA	++++
356	RPFLRLPSL	+++
357	LPPHIFAI	++
358	APSMRLKNQL	+++
359	SPRRLVELAGQSL	++++
360	SPASRSISLL	++++
361	SPYGSDRLVQL	++++
362	KPMLPPAAF	+++
363	KPRTPF TTA	++++
364	KPRTPF TTAQL	++++
365	RPKHFLQML	++++
366	SPTLRQLDL	++++
367	APQVHIFSL	++++
368	NPASRLTAL	++++
369	RPYGCVLRAA	++++
370	AAHEFGHVL	++
371	APRSPGQVTPRG L	+++
372	SPSSASLTL	++++
373	LPKPDLPQLI	+

374	KPRNMTGLDL	++++
375	LPRGVLEGL	++++
376	FPQVGRTAL	++++
377	SPFSKRIKL	+++
378	SPRQPRLDF	++++
379	GPRQVLFPL	++++
380	SPYPGSAAF	++++
381	APRPRLLLL	+++
382	RPGPQRTTSV	++++
383	KPRATWTLKL	+++
448	TVYGEPRKL	++
449	RPKTSVNLISL	++++
450	SAAARALLP	+
451	VPILPLNHAL	++++
452	QPVKKNTL	++++
453	APALPGQVTI	++++
454	SPDPAHLESL	++++
455	FPVQATIDF	++++
456	VPVSHPVLT L	+++
457	RPPLSQRHTF	++++
458	VPIPTHYFVVL	++++
459	SPAPWRPWI	++++
460	APLMPLGKTL	++++
461	MPHLGPGILL	++++
462	MPLLADVRL	++++
463	LPTDLFNSVM	++++
464	VPFVPRTSV	+++
465	VPKSLPYVL	++++
466	SPMEAILVSRL	++++
467	VPRNQDWLGVSR QL	++++
468	LPSLHVLVL	++++
469	LPLDTRTSI	++++
470	RPNGEVKSEL	++++
471	LPLLAGTLLL	++
472	LPMPAITWY	++
473	LPGEREAAL	++++
474	VPKADLLTL	++++
475	IPLEIQKL	+
476	EPNPVEEIF	++
477	NPVPVITWYKDNR L	++++
478	APKFISPASQL	++++
479	APHAGGALL	++++
480	VPAAPT KAL	+++

- 158 -

481	SPARALLLALA	+++
482	QPSLKKIIL	++++
483	KPAAAGVKKVA	+++
484	LPASAEPKGTM	++
486	TPTRPLPSA	++
487	IPWFIKTAF	++++
488	LPLGGLPLLI	+++
489	WPNHIMLVL	++++
490	APRGPAQGEAA	++++

491	MPEADLKRIF	++
492	IPFSVDEI	+
493	LPLQQYKLV	+
494	LPLRAVNLNL	+++
495	SPSYTQASL	++++
496	MPAVSGPGPLF	+++
497	YVVKPLHPF	+++
498	RPQPQPRPAL	++++
499	QPVLPSPAC	++++

Reference List

- Alcoser, S. Y. et al., BMC Biotechnol. **11** (2011): 124
Allison, J. P. et al., Science. **270** (1995): 932-933
Andersen, R. S. et al., Nat Protoc. **7** (2012): 891-902
Anderson, N. L. et al., J Proteome Res. **11** (2012): 1868-1878
Appay, V. et al., Eur J Immunol. **36** (2006): 1805-1814
Banchereau, J. et al., Cell. **106** (2001): 271-274
Beatty, G. et al., J Immunol. **166** (2001): 2276-2282
Beggs, J. D., Nature. **275** (1978): 104-109
Benjamini, Y. et al., Journal of the Royal Statistical Society Series B (Methodological), **Vol.57** (1995): 289-300
Boulter, J. M. et al., Protein Eng. **16** (2003): 707-711
Braumuller, H. et al., Nature. (2013)
Bray, F. et al., Int J Cancer. **132** (2013): 1133-1145
Brossart, P. et al., Blood. **90** (1997): 1594-1599
Bruckdorfer, T. et al., Curr Pharm Biotechnol. **5** (2004): 29-43
Card, K. F. et al., Cancer Immunol Immunother. **53** (2004): 345-357
Cohen, C. J. et al., J Mol Recognit. **16** (2003a): 324-332
Cohen, C. J. et al., J Immunol. **170** (2003b): 4349-4361
Cohen, S. N. et al., Proc Natl Acad Sci U S A. **69** (1972): 2110-2114
Coligan, J. E. et al., Current Protocols in Protein Science. (1995)
Colombetti, S. et al., J Immunol. **176** (2006): 2730-2738
Dengjel, J. et al., Clin Cancer Res. **12** (2006): 4163-4170
Denkberg, G. et al., J Immunol. **171** (2003): 2197-2207
Falk, K. et al., Nature. **351** (1991): 290-296
Ferlay et al., GLOBOCAN 2012 v1 0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No 11 [Internet] (2013), <http://globocan.iarc.fr>
Follenzi, A. et al., Nat Genet. **25** (2000): 217-222
Fong, L. et al., Proc Natl Acad Sci U S A. **98** (2001): 8809-8814
Forsey, R. W. et al., Biotechnol Lett. **31** (2009): 819-823
Gabrilovich, D. I. et al., Nat Med. **2** (1996): 1096-1103
Gattinoni, L. et al., Nat Rev Immunol. **6** (2006): 383-393
Gnjatic, S. et al., Proc Natl Acad Sci U S A. **100** (2003): 8862-8867
Godkin, A. et al., Int Immunol. **9** (1997): 905-911

- Gragert, L. et al., Hum Immunol. **74** (2013): 1313-1320
- Green, M. R. et al., Molecular Cloning, A Laboratory Manual. **4th** (2012)
- Greenfield, E. A., Antibodies: A Laboratory Manual. **2nd** (2014)
- Gunawardana, C. et al., Br J Haematol. **142** (2008): 606-609
- Gustafsson, C. et al., Trends Biotechnol. **22** (2004): 346-353
- Hwang, M. L. et al., J Immunol. **179** (2007): 5829-5838
- Jung, G. et al., Proc Natl Acad Sci U S A. **84** (1987): 4611-4615
- Kibbe, A. H., Handbook of Pharmaceutical Excipients. **rd** (2000)
- Krieg, A. M., Nat Rev Drug Discov. **5** (2006): 471-484
- Kuball, J. et al., Blood. **109** (2007): 2331-2338
- Liddy, N. et al., Nat Med. **18** (2012): 980-987
- Ljunggren, H. G. et al., J Exp Med. **162** (1985): 1745-1759
- Longenecker, B. M. et al., Ann N Y Acad Sci. **690** (1993): 276-291
- Lonsdale, J., Nat Genet. **45** (2013): 580-585
- Lukas, T. J. et al., Proc Natl Acad Sci U S A. **78** (1981): 2791-2795
- Lundblad, R. L., Chemical Reagents for Protein Modification. **3rd** (2004)
- Meziere, C. et al., J Immunol. **159** (1997): 3230-3237
- Molina, J. R. et al., Mayo Clin Proc. **83** (2008): 584-594
- Morgan, R. A. et al., Science. **314** (2006): 126-129
- Mortara, L. et al., Clin Cancer Res. **12** (2006): 3435-3443
- Mueller, L. N. et al., J Proteome Res. **7** (2008): 51-61
- Mueller, L. N. et al., Proteomics. **7** (2007): 3470-3480
- Mumberg, D. et al., Proc Natl Acad Sci U S A. **96** (1999): 8633-8638
- Pinheiro, J. et al., nlme: Linear and Nonlinear Mixed Effects Models (<http://CRANR-project.org/package=nlme>) (2015)
- Plebanski, M. et al., Eur J Immunol. **25** (1995): 1783-1787
- Porta, C. et al., Virology. **202** (1994): 949-955
- Rammensee, H. et al., Immunogenetics. **50** (1999): 213-219
- Rini, B. I. et al., Cancer. **107** (2006): 67-74
- Rock, K. L. et al., Science. **249** (1990): 918-921
- Rodenko, B. et al., Nat Protoc. **1** (2006): 1120-1132
- Saiki, R. K. et al., Science. **239** (1988): 487-491
- Schmitt, T. M. et al., Hum Gene Ther. **20** (2009): 1240-1248
- Scholten, K. B. et al., Clin Immunol. **119** (2006): 135-145
- Seeger, F. H. et al., Immunogenetics. **49** (1999): 571-576
- Sherman, F. et al., Laboratory Course Manual for Methods in Yeast Genetics. (1986)
- Silva, L. P. et al., Anal Chem. **85** (2013): 9536-9542
- Singh-Jasuja, H. et al., Cancer Immunol Immunother. **53** (2004): 187-195
- Small, E. J. et al., J Clin Oncol. **24** (2006): 3089-3094
- Sturm, M. et al., BMC Bioinformatics. **9** (2008): 163
- Teufel, R. et al., Cell Mol Life Sci. **62** (2005): 1755-1762
- Thakkar, J. P. et al., Cancer Epidemiol Biomarkers Prev. **23** (2014): 1985-1996
- Tran, T. T. et al., Photochem Photobiol. **90** (2014): 1136-1143
- Walter, S. et al., J Immunol. **171** (2003): 4974-4978
- Walter, S. et al., Nat Med. **18** (2012): 1254-1261
- Willcox, B. E. et al., Protein Sci. **8** (1999): 2418-2423

World Cancer Report (2014)

Zaremba, S. et al., Cancer Res. **57** (1997): 4570-4577

Zufferey, R. et al., J Virol. **73** (1999): 2886-2892

CLAIMS

1. A peptide comprising an amino acid sequence selected from the group consisting of SEQ ID No. 1 to SEQ ID No. 383 and SEQ ID No. 448 to SEQ ID No. 499, and variant sequences thereof which are at least 88% homologous to SEQ ID No. 1 to SEQ ID No. 383 and SEQ ID No. 448 to SEQ ID No. 499, and wherein said variant binds to molecule(s) of the major histocompatibility complex (MHC) and/or induces T cells cross-reacting with said variant peptide; and a pharmaceutical acceptable salt thereof, wherein said peptide is up to 16 amino acids in length.
2. The peptide according to claim 1, wherein said peptide has the ability to bind to an MHC class-I or –II molecule, and wherein said peptide, when bound to said MHC, is capable of being recognized by CD4 and/or CD8 T cells.
3. The peptide or variant thereof according to claim 1 or 2, wherein the amino acid sequence thereof comprises a continuous stretch of amino acids according to any one of SEQ ID No. 1 to SEQ ID No. 383 and SEQ ID No. 448 to SEQ ID No. 499.
4. The peptide or variant thereof according to any of claims 1 to 3, wherein said peptide or variant thereof has an overall length of from 8 to 16 amino acids, and most preferred wherein the peptide consists or consists essentially of an amino acid sequence according to any of SEQ ID No. 1 to SEQ ID No. 383 and SEQ ID No. 448 to SEQ ID No. 499.
5. The peptide or variant thereof according to any of Claims 1 to 4, wherein said peptide is modified and/or includes non-peptide bonds.
6. The peptide or variant thereof according to any of Claims 1 to 5, wherein said peptide is part of a fusion protein, in particular comprising N-terminal amino acids of the HLA-DR antigen-associated invariant chain (Ii).

7. An antibody, in particular a soluble or membrane-bound antibody, preferably a monoclonal antibody or fragment thereof, that specifically recognizes the peptide or variant thereof according to any of claims 1 to 5, preferably the peptide or variant thereof according to any of claims 1 to 5 when bound to an MHC molecule.
8. A T-cell receptor, preferably soluble or membrane-bound, or a fragment thereof, that is reactive with an HLA ligand, wherein said ligand is the peptide or variant thereof according to any of claims 1 to 5, preferably the peptide or variant thereof according to any of claims 1 to 5 when bound to an MHC molecule.
9. The T-cell receptor according to claim 8, wherein said ligand amino acid sequence is at least 88% identical to any one of SEQ ID No. 1 to SEQ ID No. 383 and SEQ ID No. 448 to SEQ ID No. 499, or wherein said ligand amino acid sequence consists of any one of SEQ ID No. 1 to SEQ ID No. 383 and SEQ ID No. 448 to SEQ ID No. 499.
10. The T-cell receptor according to claim 8 or 9, wherein said T-cell receptor is provided as a soluble molecule and optionally carries a further effector function such as an immune stimulating domain or toxin.
11. An aptamer that specifically recognizes the peptide or variant thereof according to any of claims 1 to 5, preferably the peptide or variant thereof according to any of claims 1 to 5 that is bound to an MHC molecule.
12. A nucleic acid, encoding for a peptide or variant thereof according to any one of claims 1 to 5, an antibody or fragment thereof according to claim 7, a T-cell receptor or fragment thereof according to claim 8 or 9, optionally linked to a heterologous promoter sequence, or an expression vector expressing said nucleic acid.
13. A recombinant host cell comprising the peptide according to any one of claims 1 to 6, the antibody or fragment thereof according to claim 7, the T-cell receptor or fragment thereof according to claim 8 or 9 or the nucleic acid or the expression

vector according to claim 12, wherein said host cell preferably is selected from an antigen presenting cell, such as a dendritic cell, a T cell or an NK cell.

14. An *in vitro* method for producing activated T lymphocytes, the method comprising contacting *in vitro* T cells with antigen loaded human class I or II MHC molecules expressed on the surface of a suitable antigen-presenting cell or an artificial construct mimicking an antigen-presenting cell for a period of time sufficient to activate said T cells in an antigen specific manner, wherein said antigen is a peptide according to any one of claims 1 to 4.
15. An activated T lymphocyte, produced by the method according to claim 14, that selectively recognizes a cell which presents a polypeptide comprising an amino acid sequence given in any one of claims 1 to 4.
16. A pharmaceutical composition comprising at least one active ingredient selected from the group consisting of the peptide according to any one of claims 1 to 6, the antibody or fragment thereof according to claim 7, the T-cell receptor or fragment thereof according to claim 8 or 9, the aptamer according to claim 11, the nucleic acid or the expression vector according to claim 12, the host cell according to claim 13, or the activated T lymphocyte according to claim 15, or a conjugated or labelled active ingredient, and a pharmaceutically acceptable carrier, and optionally, pharmaceutically acceptable excipients and/or stabilizers.
17. A method for producing the peptide or variant thereof according to any of claims 1 to 6, the antibody or fragment thereof according to claim 7, or the T-cell receptor or fragment thereof according to claim 8 or 9, the method comprising culturing the host cell according to claim 13, and isolating the peptide or variant thereof, the antibody or fragment thereof or the T cell receptor or fragment thereof from said host cell and/or its culture medium.
18. The peptide according to any one of claims 1 to 6, the antibody or fragment thereof according to claim 7, the T-cell receptor or fragment thereof according to claim 8 or 9, the aptamer according to claim 11, the nucleic acid or the expression vector

- 164 -

according to claim 12, the host cell according to claim 13, or the activated T lymphocyte according to claim 15 for use in medicine.

19. A method for killing target cells in a patient which target cells present a polypeptide comprising an amino acid sequence given in any one of claims 1 to 4, the method comprising administering to the patient an effective number of activated T cells as defined in claim 15.
20. The peptide according to any one of claims 1 to 6, the antibody or fragment thereof according to claim 7, the T-cell receptor or fragment thereof according to claim 8 or 9, the aptamer according to claim 11, the nucleic acid or the expression vector according to claim 12, the host cell according to claim 13, or the activated T lymphocyte according to claim 15 for use in diagnosis and/or treatment of cancer, or for use in the manufacture of a medicament against cancer.
21. The use according to claim 20, wherein said cancer is selected from the group of acute myeloid leukaemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukaemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer and other tumors that show an overexpression of a protein from which a peptide SEQ ID No. 1 to SEQ ID No. 383 and SEQ ID No. 448 to SEQ ID No. 499 is derived from.
22. A kit comprising:
 - (a) a container comprising a pharmaceutical composition containing the peptide(s) or the variant according to any one of claims 1 to 6, the antibody or fragment thereof according to claim 7, the T-cell receptor or fragment thereof according to claim 8 or 9, the aptamer according to claim 11, the nucleic acid or

- 165 -

- the expression vector according to claim 12, the host cell according to claim 13, or the activated T lymphocyte according to claim 15, in solution or in lyophilized form;
- (b) optionally, a second container containing a diluent or reconstituting solution for the lyophilized formulation;
 - (c) optionally, at least one more peptide selected from the group consisting of SEQ ID No. 1 to SEQ ID No. 445 and SEQ ID No. 448 to SEQ ID No. 499, and
 - (d) optionally, instructions for (i) use of the solution or (ii) reconstitution and/or use of the lyophilized formulation.
23. The kit according to claim 22, further comprising one or more of (iii) a buffer, (iv) a diluent, (v) a filter, (vi) a needle, or (v) a syringe.
24. A method for producing a personalized anti-cancer vaccine or a compound-based and/or cellular therapy for an individual patient, said method comprising:
- a) identifying tumor-associated peptides (TUMAPs) presented by a tumor sample from said individual patient;
 - b) comparing the peptides as identified in a) with a warehouse of peptides that have been pre-screened for immunogenicity and/or over-presentation in tumors as compared to normal tissues;
 - c) selecting at least one peptide from the warehouse that matches a TUMAP identified in the patient; and
 - d) manufacturing and/or formulating the personalized vaccine or compound-based or cellular therapy based on step c).
25. The method according to claim 24, wherein said TUMAPs are identified by:
- a1) comparing expression data from the tumor sample to expression data from a sample of normal tissue corresponding to the tissue type of the tumor sample to identify proteins that are over-expressed or aberrantly expressed in the tumor sample; and

- 166 -

- a2) correlating the expression data with sequences of MHC ligands bound to MHC class I and/or class II molecules in the tumor sample to identify MHC ligands derived from proteins over-expressed or aberrantly expressed by the tumor.
26. The method according to claim 24 or 25, wherein the sequences of MHC ligands are identified by eluting bound peptides from MHC molecules isolated from the tumor sample, and sequencing the eluted ligands.
27. The method according to any of claims 24 to 26, wherein the normal tissue corresponding to the tissue type of the tumor sample is obtained from the same patient.
28. The method according to any of claims 24 to 27, wherein the peptides included in the warehouse are identified based on the following steps:
- aa. Performing genome-wide messenger ribonucleic acid (mRNA) expression analysis by highly parallel methods, such as microarrays or sequencing-based expression profiling, comprising identify genes that over-expressed in a malignant tissue, compared with a normal tissue or tissues;
 - ab. Selecting peptides encoded by selectively expressed or over-expressed genes as detected in step aa, and
 - ac. Determining an induction of *in vivo* T-cell responses by the peptides as selected comprising *in vitro* immunogenicity assays using human T cells from healthy donors or said patient; or
 - ba. Identifying HLA ligands from said tumor sample using mass spectrometry;
 - bb. Performing genome-wide messenger ribonucleic acid (mRNA) expression analysis by highly parallel methods, such as microarrays or sequencing-based expression profiling, comprising identify genes that over-expressed in a malignant tissue, compared with a normal tissue or tissues;
 - bc. Comparing the identified HLA ligands to said gene expression data;
 - bd. Selecting peptides encoded by selectively expressed or over-expressed genes as detected in step bc;

- 167 -

- be. Re-detecting of selected TUMAPs from step bd on tumor tissue and lack of or infrequent detection on healthy tissues and confirming the relevance of over-expression at the mRNA level; and
- bf. Determining an induction of *in vivo* T-cell responses by the peptides as selected comprising *in vitro* immunogenicity assays using human T cells from healthy donors or said patient.
29. The method according to any of claims 24 to 28, wherein the immunogenicity of the peptides included in the warehouse is determined by a method comprising *in vitro* immunogenicity assays, patient immunomonitoring for individual HLA binding, MHC multimer staining, ELISPOT assays and/or intracellular cytokine staining.
30. The method according to any of claims 24 to 29, wherein said warehouse comprises a plurality of peptides selected from the group consisting of SEQ ID No. 1 to SEQ ID No. 445 and SEQ ID No. 448 to SEQ ID No. 499.
31. The method according to any of claims 24 to 30, further comprising identifying at least one mutation that is unique to the tumor sample relative to normal corresponding tissue from the individual patient, and selecting a peptide that correlates with the mutation for inclusion in the vaccine or for the generation of cellular therapies.
32. The method according to claim 31, wherein said at least one mutation is identified by whole genome sequencing.