

DESCRIPTION

METHOD FOR PROMOTING EFFICIENCY OF PURIFICATION OF FC
REGION-CONTAINING POLYPEPTIDETechnical Field

The present invention relates to methods for increasing the dynamic binding capacity of an Fc region-containing polypeptide for a Protein A resin in Protein A column chromatography, purification methods that use such methods, and the like.

Background Art

In the production of antibody pharmaceuticals, purification steps that use Protein A columns, ion exchange columns, and such, greatly affect the production efficiency (yield) of antibodies; therefore, it is desired that the efficiency of such steps be increased. Means for accomplishing the efficiency include the following two methods: (1) increasing the binding capacity per unit volume of resin; and (2) reducing the time required for purification by high-flow treatment.

Recently, the improvement of Protein A resins has advanced, and more antibodies can bind to Protein A resins, and accordingly, efficient antibody purification is being achieved. For example, for rProtein A sepharose Fast Flow manufactured by GE Healthcare, which is a typical first-generation Protein A resin, the ordinary antibody-binding capacity is 15 to 20 g/L resin; whereas, for the second-generation Protein A resin manufactured by the same company, Mab Select SuRe, which is currently most commonly used worldwide, the generally observed binding capacity is approximately 30 g/L resin. In addition, compared to the former resin, the latter can accommodate a linear flow rate approximately 1.5- to 2-times higher, and a more efficient Protein A purification of antibody molecules has been possible.

Bispecific antibodies have properties of recognizing two different types of antigens, and accordingly, they carry two types of H chains. Therefore, culture supernatants containing bispecific antibodies contain not only the bispecific antibodies comprising the two types of H chains, but also antibodies comprising only one type of H chain. To separate these antibodies from the bispecific antibodies, Fc region variants with modified binding activities to Protein A resins have been used (Patent Documents 1 and 2). It was concerned that such molecular modification could have an effect of decreasing the efficiency of Protein A purification of bispecific antibodies.

Under such circumstances, new means for performing more efficient bispecific antibody purification using Protein A resin columns have been desired.

Citation List

[Patent Documents]

[Patent Document 1] US20100331527

5 [Patent Document 2] US20130018174

Summary

[Problems to be Solved by the Invention]

10 An objective of the present invention is to provide methods for efficiently purifying an Fc region-containing polypeptide, in particular a bispecific antibody, using a Protein A resin column.

[Means for Solving the Problems]

15 As a result of dedicated research to solve the above-mentioned problems, the present inventors discovered that, by preparing an Fc region whose first polypeptide chain and second polypeptide chain have binding activities to a Protein A resin that are different from each other, the dynamic binding capacity of the antibody is increased, and the antibody purification efficiency is increased, and completed the present disclosure.

20 In one aspect, the present disclosure refers to a method for producing and purifying a heterologous Fc region-containing polypeptide, which comprises the steps of: (a) preparing a first polypeptide chain and a second polypeptide chain of an Fc region having binding activities to a Protein A resin that are different from each other; (b) comparing the dynamic binding capacity of a heterologous Fc region-containing polypeptide comprising the first polypeptide chain and the second polypeptide chain prepared in step (a) for the Protein A resin in Protein A column chromatography with the dynamic binding capacity of a homologous Fc region-containing polypeptide comprising two polypeptide chains having substantially the same binding activity to said resin for the Protein A resin in Protein A column chromatography to confirm the elevated dynamic binding capacity of the former Fc region-containing polypeptide compared with that of the latter Fc region-containing polypeptide; (c) contacting a sample comprising the heterologous Fc region-containing polypeptide with said resin; and (d) collecting the
30 heterologous Fc region-containing polypeptide bound to the resin.

Specifically, the present disclosure provides the following:

[1] a method for increasing the dynamic binding capacity of an Fc region-containing polypeptide for a Protein A resin in Protein A column chromatography;

- [2] The method of [1], comprising the step of preparing a first polypeptide chain and a second polypeptide chain having binding activities to the resin that are different from each other, as the first polypeptide chain and the second polypeptide chain of the Fc region;
- [3] The method of [1] or [2], comprising the steps of preparing a first polypeptide chain that
5 binds to the resin as the first polypeptide chain of the Fc region, and preparing a second polypeptide chain that does not bind to the resin or shows weaker binding to the resin compared to the first polypeptide chain, as the second polypeptide chain of the Fc region;
- [4] The method of any one of [1] to [3], comprising the step of modifying the Fc region of the Fc region-containing polypeptide so that the first polypeptide chain of the Fc region binds to the
10 resin, and the second polypeptide chain of the Fc region does not bind to the resin or shows weaker binding to the resin compared to the first polypeptide chain;
- [5] The method of any one of [1] to [4], wherein the first polypeptide chain of the Fc region comprises a CH3 of IgG1, IgG2, or IgG4, and the second polypeptide chain of the Fc region comprises a CH3 of IgG3;
- 15 [6] The method of any one of [1] to [5], wherein the amino acid at position 435 according to EU

numbering in the first polypeptide chain of the Fc region is His, and the amino acid at position 435 according to EU numbering in the second polypeptide chain of the Fc region is Arg;

[7] The method of any one of [1] to [6], wherein the increase in the binding capacity is 5 g/L resin or more;

5 [8] The method of any one of [1] to [7], wherein the dynamic binding capacity after the increase is 45 g/L resin or more;

[9] The method of any one of [1] to [8], wherein the Fc region-containing polypeptide is an antibody;

[10] The method of [9], wherein the antibody is a bispecific antibody;

10 [11] A method for purifying an Fc region-containing polypeptide using the method of any one of [1] to [10];

[12] An Fc region-containing polypeptide purified by the method of [11];

[13] A Protein A resin bound by an Fc region-containing polypeptide, wherein the dynamic binding capacity of the Fc region-containing polypeptide for the Protein A resin in Protein A column chromatography is 45 g/L resin or more;

15 [14] An Fc region-containing polypeptide in which the dynamic binding capacity for a Protein A resin in Protein A column chromatography has been increased;

[15] A method for producing an Fc region-containing polypeptide using a Protein A resin, which comprises the steps of:

20 (a) preparing a first polypeptide chain and a second polypeptide chain of an Fc region having binding activities to said resin that are different from each other;

(b) comparing the dynamic binding capacity of the Fc region-containing polypeptide of step (a) for the Protein A resin in Protein A column chromatography with the dynamic binding capacity of an Fc region-containing polypeptide comprising two polypeptide chains having substantially the same binding activity to said resin for the Protein A resin in Protein A column chromatography;

25 (c) contacting a sample comprising a polypeptide comprising the first polypeptide chain of the Fc region and a polypeptide comprising the second polypeptide chain of the Fc region with said resin; and

30 (d) collecting an Fc region-containing polypeptide binding to the resin and comprising a heterologous polypeptide which comprises the polypeptide comprising the first polypeptide chain of the Fc region and the polypeptide comprising the second polypeptide chain of the Fc region;

[16] The method of [15], wherein said step (a) is preparing a first polypeptide chain that binds to the resin as the first polypeptide chain of the Fc region, and preparing a second polypeptide chain that does not bind to the resin or shows weaker binding to the resin compared to the first

polypeptide chain, as the second polypeptide chain of the Fc region;

[17] The method of [15] or [16], wherein said step (a) is modifying the Fc region of the Fc region-containing polypeptide for purification, so that the first polypeptide chain of the Fc region binds to the resin, and the second polypeptide chain of the Fc region does not bind to the resin or shows weaker binding to the resin compared to the first polypeptide chain;

[18] the method of any one of [15] to [17], wherein the sample in step (c) comprises a common L chain polypeptide that can provide binding ability for both a polypeptide comprising the first polypeptide chain of the Fc region and a polypeptide comprising the second polypeptide chain of the Fc region;

[19] the purification method of [11], wherein the Fc region-containing polypeptide is an antibody;

[20] the purification method of [19], wherein the antibody is a bispecific antibody;

[21] an antibody purified by the method of [19];

[22] a bispecific antibody purified by the method of [20]; and

[23] a column containing the resin of [13].

[Effects of the Invention]

The present invention provides methods for more efficient purification of Fc region-containing polypeptides, in particular bispecific antibodies, using Protein A resins.

Brief Description of the Drawings

Fig. 1 shows a breakthrough curve chromatogram detecting proteins discharged from the column when a BiAb solution was continuously loaded onto a Protein A resin column.

Mode for Carrying Out the Invention

Herein below, the present invention will be described in detail.

The Fc region-containing polypeptides to be used in the present invention may contain an antibody Fc region, and they include polypeptides formed by fusing an Fc region with another polypeptide, for example, antibodies.

“Polypeptides” of the present invention generally refers to peptides and proteins approximately ten amino acids or more in length. Furthermore, they are generally polypeptides derived from organisms, but are not particularly limited, and for example, they may be polypeptides comprising an artificially designed sequence. Furthermore, they may be any of naturally-occurring polypeptides, synthetic polypeptides, recombinant polypeptides, or such.

“Fc region” generally refers to the region comprising two polypeptide chains which consist of a hinge portion or a portion thereof, CH2 domain, and CH3 domain in an antibody

molecule, but is not particularly limited thereto, and there are also cases where the hinge portion or portion thereof is not included. According to EU numbering by Kabat, a human IgG-class Fc region refers to, for example, the region from cysteine at position 226 to the C terminus, or from proline at position 230 to the C terminus, but not limited thereto. Furthermore, the human CH2 domain refers to positions 231 to 340 according to EU numbering by Kabat, and the human CH3 domain refers to positions 341 to 447 according to EU numbering by Kabat, but not limited thereto.

The Fc region may be obtained preferably by partially digesting IgG1, IgG2, IgG3, Fc region-containing monoclonal antibodies or such using a protease such as pepsin, and then re-eluting the fraction adsorbed onto protein A resins. The protease is not particularly limited as long as it can digest a full-length antibody so that Fab and F(ab')₂ will be produced in a restrictive manner by appropriately setting the enzyme reaction conditions such as pH, and examples include pepsin and papain.

Examples of the Fc region include human IgG-type Fc, and for example, they may be any of the IgG1, IgG2, IgG3, and IgG4 isotypes.

The Fc region of the present invention comprises the first polypeptide chain and the second polypeptide chain mentioned above.

An embodiment of the present invention is a method for increasing the dynamic binding capacity of an Fc region-containing polypeptide for a Protein A resin in Protein A column chromatography. The first polypeptide chain and the second polypeptide chain comprised in the Fc region preferably have binding activities to the Protein A resin that are different from each other. For example, when using a polypeptide chain that binds to a Protein A resin as the first polypeptide chain, a polypeptide chain that does not bind to or shows weaker binding to the Protein A resin compared to the first polypeptide chain can be used as the second polypeptide chain. As the first polypeptide chain, a polypeptide chain comprising the CH3 of IgG1, IgG2, or IgG4 may be used, and as the second polypeptide chain, a polypeptide chain comprising the CH3 of IgG3 may be used. In this case, IgG1, IgG2, IgG3, and IgG4 may be naturally-occurring, or they may include mutations within a range that allows the objective of the present invention to be accomplished. Furthermore, as the first polypeptide chain, a polypeptide chain in which position 435 according to EU numbering is His (H) can be used. As the second polypeptide chain, a polypeptide chain in which position 435 according to EU numbering is Arg (R) can be used. Furthermore, a polypeptide chain in which positions 435 and 436 according to EU numbering are His (H) and Tyr (Y), respectively, can be used as the first polypeptide chain. A polypeptide chain in which positions 435 and 436 according to EU numbering are Arg (R) and Phe (F), respectively, can be used as the second polypeptide chain. The positions other than position 435 or 436 according to EU numbering may be the same as or

different from those of the naturally-occurring IgGs.

In this embodiment, increasing the dynamic binding capacity of an Fc region-containing polypeptide for the resin in Protein A column chromatography can be accomplished by modifying the Fc region of the Fc region-containing polypeptide that binds to the resin, so that the binding activities of the first polypeptide chain of the Fc region and the second polypeptide chain of the Fc region to the resin will be different from each other.

In another embodiment of the present invention, increasing the dynamic binding capacity of an Fc region-containing polypeptide for the resin in Protein A column chromatography can be accomplished by modifying the Fc region of the Fc region-containing polypeptide that binds to the resin, so that the first polypeptide chain of the Fc region binds to the resin, but the second polypeptide chain of the Fc region does not bind to the resin or shows weaker binding to the resin compared to the first polypeptide chain.

Examples of the modification include, but are not limited to, modifications performed so that the first and second polypeptide chains of the Fc region will contain CH3 regions such as those mentioned above, for example, modifications performed so that the above-mentioned specific amino acid residues at specified positions are contained.

On the other hand, the regions other than the Fc region in the Fc region-containing polypeptides used in the present invention may be in a homologous or heterologous form.

The homologous form has one or two or more, uniform or the same antigen-binding activities (i.e., when the Fc region-containing polypeptide is an antigen-binding molecule, it refers to an antigen-binding molecule having one or two or more, uniform or the same antigen-binding activity, which is, for example, an IgG-type antibody having two identical antigen-binding sites).

The heterologous form preferably has different antigen-binding activities (i.e., the Fc region-containing polypeptide is a bispecific antigen-binding molecule, for example, a bispecific antibody). When the Fc region-containing polypeptide used in the present invention is a bispecific antibody, while the H chains may be heterologous, the L chains may be common L chains, and the common L chains preferably provide binding abilities for both H chains. When the bispecific antibody is an IgG-type antibody, it is composed of two heterologous H chains and two identical common L chains.

Binding capacities include static binding capacity (SBC) and dynamic binding capacity (DBC). Static binding capacity refers to the upper limit of the amount of polypeptides that a resin can adsorb, and dynamic binding capacity refers to the degree to which polypeptides can be collected when a polypeptide-containing solution is flowing through the column. A resin having a large dynamic binding capacity allows efficient polypeptide adsorption even under high linear flow rate, and polypeptide purification can be accomplished in a short time.

For example, dynamic binding capacity (DBC) can be determined by the following method. First, a column loaded with a resin is placed in a chromatography apparatus, and a polypeptide-containing sample solution is allowed to flow through the column at a specified linear flow rate. Then, the absorbance of the eluate is measured, and DBC is determined by identifying the mass of the added polypeptide when breakthrough (BT) for a specified proportion (for example, 5%) of absorbance of the added sample solution is measured.

The following apparatus and such can be used for the DBC calculation:

- LC apparatus: AKTA AVANT25 manufactured by GE Healthcare
- Software: Unicorn version 6.1 manufactured by GE Healthcare
- Protein A resin: Mab Select SuRe (Cat No.17-5438-05) or Hitrap Mab Select SuRe (Cat No. 11-0034-93) manufactured by GE Healthcare
- Buffers:
 - equilibration/preliminary washing – 20 mmol/L Na-phosphate, pH 7.5
 - elution – 50 mmol/L Acetic acid
 - regeneration – 0.1 mol/L NaOH

The method for calculating the DBC can be carried out as follows.

The above-mentioned apparatus, software, and resins are used, and by performing the chromatography operation by the following procedure, the DBC is calculated. A calculation method when using 5% BT as the indicator is shown below.

- (1) The load fraction (IgG concentration: P g/L) is allowed to flow through the LC apparatus without passing it through the column, and the value of OD_{280 nm} for 100% leakage (= 100% BT) was confirmed. This value is denoted as a.

(2) The value obtained by multiplying 0.05 to a is defined as the OD_{280 nm} at 5% BT. This value is denoted as b_{5%}.

- (3) The load fraction is allowed to flow continuously through a set amount of equilibrated resin (r L), and when the OD_{280 nm} value reaches b_{5%}, the volume of the load fraction is read from the chromatogram. This value is denoted as c_{5%} L.

(4) The value obtained by the equation $(P \times c_{5\%})/r$ is calculated as DBC_{5%} which is the dynamic binding capacity at 5% BT.

$$DBC_{5\%} = (P \times c_{5\%})/r \text{ (unit: g/L resin)}$$

When determining DBC_{10%}, the calculation is possible by determining c_{10%} in a similar manner.

In an embodiment of the present invention, the increase in the dynamic binding capacity of an Fc region-containing polypeptide for a Protein A resin in Protein A column chromatography is at least 5 g/L resin, preferably 10 g/L resin or more, 15 g/L resin or more, 20 g/L resin or more, and 25 g/L resin or more, when taking 5% BT as the standard.

In a specific embodiment of the present invention, the increase in the dynamic binding capacity of an Fc region-containing polypeptide for a Protein A resin in Protein A column chromatography is at least 5 g/L resin, preferably 10 g/L resin or more, 15 g/L resin or more, 20 g/L resin or more, and 25 g/L resin or more, at contact time of 3.4 minutes, when taking 5% BT as the standard.

In an embodiment of the present invention, according to the method of the present invention, the dynamic binding capacity of an Fc region-containing polypeptide for a Protein A resin in Protein A column chromatography is at least 45 g/L resin or more, preferably 50 g/L resin or more, 55 g/L resin or more, and 60 g/L resin or more, when taking 5% BT as the standard.

In a specific embodiment of the present invention, according to the method of the present invention, the dynamic binding capacity of an Fc region-containing polypeptide for a Protein A resin in Protein A column chromatography is at least 50 g/L resin or more, preferably 51 g/L resin or more, 52 g/L resin or more, 53 g/L resin or more, 54 g/L resin or more, and 55 g/L resin or more, at contact time of 3.4 minutes, when taking 5% BT as the standard.

In one embodiment of the present invention, the Fc region-containing polypeptide may be a polypeptide in which an Fc region is linked to another protein, bioactive peptide, or such. Examples of other proteins and bioactive peptides include receptors, adhesion molecules, ligands (cytokines, chemokines, and such), and enzymes, but are not limited thereto. They may be blood coagulation factors, and examples include FIX, FIXa, and FX.

In one embodiment of the present invention, the Fc region-containing polypeptide may be an immunoadhesin.

In another embodiment of the present invention, the Fc region-containing polypeptides may be antibodies. Antibodies of the present invention are not particularly limited, as long as they bind to antigens of interest, and they may be polyclonal or monoclonal antibodies. Monoclonal antibodies are preferred because they can be stably produced as homogeneous antibodies.

The monoclonal antibodies used in the present invention include not only those derived from animals such as humans, mice, rats, hamsters, rabbits, sheep, camels, and monkeys, but also artificially modified, gene recombinant antibodies such as chimeric antibodies, humanized antibodies (also referred to as reshaped human antibodies), and bispecific antibodies. Furthermore, they also include gene recombinant antibodies produced by artificially modifying the antibody constant region and such to alter the physical properties of the antibody molecule, specifically, alteration of the isoelectric point (pI), modification of affinity for Fc receptor, etc., for the purpose of improving retention in blood and in vivo kinetics.

The immunoglobulin class of the antibodies used in the present invention is not

particularly limited; and the class may be any class, including IgG such as IgG1, IgG2, IgG3, and IgG4, IgA, IgD, IgE, and IgM. However, IgG is preferred.

The antibodies used in the present invention also include not only whole antibodies but also antibody fragments such as Fv, Fab, and F(ab)₂, and minibodies (low molecular weight
5 antibodies) such as monovalent or bivalent or higher valency single-chain Fv formed by linking antibody variable regions *via* a linker such as a peptide linker (scFv, sc(Fv)₂, diabodies such as scFv dimer, etc).

The above-described antibodies used in the present invention can be prepared by methods well known to those skilled in the art.

10 Basically, monoclonal antibody-producing hybridomas can be prepared using known techniques as described below. Specifically, immunization is carried out according to a conventional immunization method using a desired antigen or cells expressing the desired antigen as a sensitizing antigen. The yielded immunocytes are fused with known parental cells by a conventional cell fusion method. The fused cells are screened for monoclonal
15 antibody-producing cells (hybridomas) by conventional screening methods to produce the hybridomas. Hybridomas can be produced, for example, according to the method by Milstein *et al.* (Kohler, G. and Milstein, C., Methods Enzymol. (1981) 73:3-46).

Amino acid residues can be modified by modifying one or more bases in a DNA encoding a polypeptide and expressing the DNA in a host cell, as described below. Those
20 skilled in the art can easily determine the number, positions, and types of nucleotides that should be modified depending on the types of amino acid residues after the modification.

In the present invention, “modification” refers to substitution, deletion, addition, insertion, or a combination thereof.

Antibodies used in the present invention can also include additional alterations, besides
25 the above-mentioned amino acid sequence modifications. The additional modifications can be selected from any of amino acid substitution, deletion, and modification, or a combination thereof. Specifically, polypeptides containing the following modifications in their amino acid sequences are all included in the present invention:

- amino acid modifications for increasing the rate of heterologous association of two types of H
30 chains of a bispecific antibody;
- amino acid modifications for stabilizing a disulfide bond formed between a first polypeptide having antigen-binding activity and a second polypeptide with or without antigen-binding activity;
- amino acid modifications for improving antibody retention in plasma;
- 35 · modifications for improving the stability under acidic conditions;
- modifications for decreasing the heterogeneity;

- modifications for suppressing deamidation reactions;
- modifications for introducing a difference in the isoelectric points between two types of polypeptides; and
- modifications for changing the affinity towards an Fc γ receptor.

5 Methods for obtaining human antibodies are also known. For example, desired human antibodies having antigen-binding activity can be obtained by sensitizing human lymphocytes with an antigen of interest or cells expressing an antigen of interest *in vitro*; and fusing the sensitized lymphocytes with human myeloma cells. Alternatively, desired human antibodies can also be obtained by immunizing transgenic animals having the entire repertoire of human
10 antibody genes with an antigen. Furthermore, techniques for obtaining human antibodies by panning using a human antibody library are known. For example, the variable regions of human antibodies can be expressed as single-chain antibodies (scFvs) on the surface of phages using a phage display method, and then phages that bind to the antigen can be selected to obtain human antibodies. The antibodies used in the present invention also include such human
15 antibodies.

When the antibody genes are isolated and introduced into appropriate hosts to produce antibodies, hosts and expression vectors can be used in appropriate combinations. When eukaryotic cells are used as a host, animal cells, plant cells, and fungal cells can be used. The animal cells include mammalian cells such as CHO, COS, myeloma, baby hamster kidney
20 (BHK), HeLa, and Vero cells. Antibodies can be obtained by introducing the antibody genes of interest into these cells by transformation and then culturing the transformed cells *in vitro*.

The antigen of the antibody used in the present invention is not particularly limited, and it may be any antigens. Examples of antigens preferably include ligands (cytokines, chemokines, and such), receptors, cancer antigens, MHC antigens, differentiation antigens,
25 immunoglobulins, and immune complexes partly containing immunoglobulins. Examples include blood coagulation factors such as FIX, FIXa, and FX.

For collection of expression products, the medium is collected when the polypeptides are secreted into the medium. When the polypeptides are produced within cells, the cells are dissolved, and then the polypeptides are collected.

30 The polypeptides can be collected and purified from recombinant cell cultures by using known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxyapatite chromatography and lectin chromatography. In the present invention, Protein A affinity chromatography is preferred.
35 Herein, purification methods using a column, separation methods using a column, and chromatography may be used synonymously. Examples of columns that use Protein A resins

include POROS A (manufactured by Applied Biosystems), rProtein A Sepharose F. F. (manufactured by GE), ProSep vA (manufactured by Millipore), but are not limited thereto. Furthermore, resins to which ligands produced by modifying the amino acid sequence of intact Protein A, and such, can be used for Protein A affinity chromatography. When such modified Protein A resins are used, amino acid modifications of the present invention produce differences in the binding activities, and polypeptide multimers of interest can be separated and purified. Examples of resins to which modified Protein A are bound include mabSelect SuRE (manufactured by GE Healthcare) and Hitrap MabSelect Sure (manufactured by GE Healthcare), but are not limited thereto. Herein, a column packed with a Protein A resin, a column that uses a Protein A resin, a Protein A resin column, and a Protein A column are synonymous. Furthermore, a purification method that uses a Protein A resin and a purification method that uses a Protein A column may also be used synonymously.

An embodiment of the present invention is a method for purifying an Fc region-containing polypeptide which uses the method for increasing dynamic binding capacity of an Fc region-containing polypeptide for a Protein A resin in Protein A column chromatography. More specifically, an embodiment is a method for purifying an antibody, and another embodiment is a method for purifying a bispecific antibody.

Another embodiment of the present invention is an Fc region-containing polypeptide purified by the above-mentioned purification method. More specifically, an embodiment is an antibody purified by the above-mentioned purification method, and another embodiment is a bispecific antibody purified by the above-mentioned purification method.

Furthermore, another embodiment of the present invention is a Protein A resin to which an Fc region-containing polypeptide is bound at 45 g/L resin or more, when taking 5% BT as the standard; and a column containing the resin. The dynamic binding capacity of the Fc region-containing polypeptide for the resin and the column containing the resin is preferably 50 g/L resin or more, 55 g/L resin or more, 60 g/L resin or more, and 65 g/L resin or more, when taking 5% BT as the standard.

Furthermore, a specific embodiment of the present invention is a Protein A resin to which an Fc region-containing polypeptide is bound, where the dynamic binding capacity of the Fc region-containing polypeptide for a Protein A resin in Protein A column chromatography is 50 g/L resin or more at contact time of 3.4 minutes, when taking 5% BT as the standard; and a column containing the resin. The dynamic binding capacity of the Fc region-containing polypeptide for the resin and the column containing the resin is preferably 51 g/L resin or more, 52 g/L resin or more, 53 g/L resin or more, 54 g/L resin or more, and 55 g/L resin or more, at contact time of 3.4 minutes, when taking 5% BT as the standard.

An embodiment of the present invention is an Fc region-containing polypeptide having

an increased dynamic binding capacity for a Protein A resin in Protein A column chromatography. More specifically, in one embodiment, the Fc region-containing polypeptide is an antibody, and in another embodiment, the Fc region-containing polypeptide is a bispecific antibody.

5 In an embodiment of the present invention, the increase in the dynamic binding capacity of the Fc region-containing polypeptide for a Protein A resin in Protein A column chromatography is, when taking 5% BT as the standard, at least 5 g/L resin, preferably 10 g/L resin or more, 15 g/L resin or more, 20 g/L resin or more, and 25 g/L resin or more.

10 In a specific embodiment of the present invention, the increase in the dynamic binding capacity of the Fc region-containing polypeptide for a Protein A resin in Protein A column chromatography is, when taking 5% BT as the standard, at least 5 g/L resin, preferably 10 g/L resin or more, 15 g/L resin or more, 20 g/L resin or more, and 25 g/L resin or more, at contact time of 3.4 minutes.

In the Fc region-containing polypeptide having an increased dynamic binding capacity for a Protein A resin in Protein A column chromatography, it is preferred that a first polypeptide chain and a second polypeptide chain contained in the Fc region have different binding activities for the Protein A resin. For example, when using a polypeptide chain that binds to the Protein A resin as the first polypeptide chain, a polypeptide chain that does not bind to the Protein A resin or binds weakly to the Protein A resin compared to the first polypeptide may be used as the second polypeptide chain. As the first polypeptide chain, a polypeptide chain comprising the CH3 of IgG1, IgG2, or IgG4 may be used. As the second polypeptide chain, a polypeptide chain comprising the CH3 of IgG3 may be used. In this case, naturally-occurring IgG1, IgG2, IgG3, and IgG4 may be used, or they may contain mutations within a range that allows the objectives of the present invention to be accomplished. Furthermore, as the first polypeptide chain, a polypeptide chain in which position 435 according to EU numbering is His (H) can be used. As the second polypeptide chain, a polypeptide chain in which position 435 according to EU numbering is Arg (R) can be used. Furthermore, a polypeptide chain in which positions 435 and 436 according to EU numbering are His (H) and Tyr (Y), respectively, can be used as the first polypeptide chain. A polypeptide chain in which positions 435 and 436 according to EU numbering are Arg (R) and Phe (F), respectively, can be used as the second polypeptide chain. The positions other than position 435 or 436 according to EU numbering may be the same as those of the naturally-occurring IgG, or different from those of the naturally-occurring IgGs.

An embodiment of the present invention is a method for producing an Fc region-containing polypeptide using a Protein A resin, which comprises the steps of:

- 35 (a) preparing a first polypeptide chain and a second polypeptide chain of an Fc region, which have binding activities to the resin that are different from each other;
- (b) comparing the dynamic binding capacity of the Fc region-containing polypeptide of step (a)

for the Protein A resin in Protein A column chromatography with the dynamic binding capacity of an Fc region-containing polypeptide comprising two polypeptide chains having substantially the same binding activities to the resin for the Protein A resin in Protein A column chromatography;

5 (c) contacting a sample comprising a polypeptide comprising the first polypeptide chain of the Fc region and a polypeptide comprising the second polypeptide of the Fc region with the resin; and

(d) collecting an Fc region-containing polypeptide comprising a heterologous polypeptide which comprises the polypeptide comprising the first polypeptide chain of the Fc region and the
10 polypeptide comprising the second polypeptide of the Fc region.

The above-mentioned step (a) may be a step of preparing a first polypeptide chain that binds to the resin as the first polypeptide chain of the Fc region, and preparing a second polypeptide chain that does not bind to or shows weaker binding to the resin (compared to binding of the aforementioned first polypeptide chain to the resin) as the second polypeptide
15 chain of the Fc region. Furthermore, the above-mentioned step (a) may be a step of modifying the Fc region of the Fc region-containing polypeptide, which is the target of purification, so that the first polypeptide chain of the Fc region binds to the resin, but the second polypeptide chain of the Fc region does not bind to or shows weaker binding to the resin (compared to binding of the aforementioned first polypeptide chain to the resin). The modification is not particularly
20 limited as long as it is a modification for obtaining an Fc region having the above-described features, and examples include modifying the first polypeptide chain to be a polypeptide chain comprising the CH3 of IgG1, IgG2, or IgG4, and modifying the second polypeptide chain to be a polypeptide chain comprising the CH3 of IgG3. Examples of such modifications include modifying position 435 according to EU numbering in the first polypeptide chain to be His, and
25 position 435 according to EU numbering in the second polypeptide chain to be Arg. Examples of other modifications include modifying positions 435 and 436 according to EU numbering of the first polypeptide chain to be His (H) and Tyr (Y), respectively, and positions 435 and 436 according to EU numbering of the second polypeptide chain to be Arg (R) and Phe (F), respectively. The positions other than position 435 or 436 according to EU numbering may be
30 the same as or different from those of the naturally-occurring IgGs.

In the above-mentioned step (b), the two polypeptide chains may be any polypeptide chains as long as their binding activities to the resin are substantially the same, and the homology between the two polypeptide chains may be high or low. For example, “an Fc region-containing polypeptide comprising two polypeptide chains having substantially the same
35 binding activities to the resin” is an Fc region-containing polypeptide comprising two of the first polypeptide chains, or an Fc region-containing polypeptide comprising two of the second

polypeptide chains. Furthermore, examples of two polypeptide chains having substantially the same binding activities to a Protein A resin include: two polypeptides chains which are polypeptide chains each comprising any of the CH3 of IgG1, IgG2, or IgG4; two polypeptides chains which are polypeptide chains each comprising the CH3 of IgG3; two polypeptide chains in which position 435 according to EU numbering in the polypeptide chains are both His (H) or both Arg (R); two polypeptide chains both of which are polypeptide chains in which positions 435 and 436 according to EU numbering are His (H) and Tyr (Y), respectively; two polypeptide chains both of which are polypeptide chains in which positions 435 and 436 according to EU numbering are Arg (R) and Phe (F), respectively.

“Substantially the same” means being not necessarily completely identical as long as this is within a range that can accomplish the objectives of the present invention, and the meaning includes being the “same”.

In one embodiment of the present invention, “comparing” in the above-mentioned step (b) may be a step of “confirming the elevated” dynamic binding capacity of the Fc region-containing polypeptide of step (a) for a Protein A resin in Protein A column chromatography compared with the dynamic binding capacity of an Fc region-containing polypeptide comprising two polypeptide chains having substantially the same binding activities to the resin for the Protein A resin in Protein A column chromatography.

By comparing or confirming the dynamic binding capacity, one can know the maximum amount of antibodies that can be loaded onto the Protein A resin column when producing the antibodies, and this enables efficient antibody production.

The sample described in the above-mentioned step (c) may comprise two different L chain polypeptides, or common L chain polypeptides that can provide binding ability to both the H chain of the polypeptide comprising the first polypeptide chain of the Fc region and the H chain of the polypeptide comprising the second polypeptide chain of the Fc region.

In an embodiment of the present invention, in the above-mentioned method for purifying an Fc region-containing polypeptide, the Fc region-containing polypeptide is an antibody, and in another embodiment the Fc region-containing polypeptide is a bispecific antibody.

The aforementioned steps (a) to (d) do not have to be performed in this order, and each step may be included multiple times.

An embodiment of the present invention is a method for purifying an Fc region-containing polypeptide using a Protein A resin, which comprises the steps of (a) to (d) above.

All patents and reference documents explicitly cited herein are incorporated by reference into this specification in their entirety.

The present invention will be further illustrated by the following Examples, but the technical scope of the present invention is not to be construed as being limited thereto.

Examples

5 [Example 1] Preparation of antibody gene expression vectors and expression of each antibody

In the Examples, the anti-FIXa/FX bispecific antibody (H1 chain / H2 chain / L chain: SEQ ID NOs: 1/2/3) having activity of substituting for FVIII function, which is described in WO2012/067176, was used (herein below, this is referred to as “BiAb”, which is a so-called heterologous antibody). The BiAb comprises four chains consisting of three types of chains. The four chains consist of an H1 chain and an H2 chain which are two types of H chains, and two common L chains which are one type of L chain. This antibody was obtained by the method described in WO2012/067176. An antibody gene was inserted into an animal cell expression vector. By transfecting CHO cells with the vector, the bispecific antibody was expressed. Furthermore, “Q homo” comprising the two L chains and the two H1 chains, and “J

10 homo” comprising the two L chains and the two H2 chains were obtained by the above method.

This antibody is IgG4-type, and His at position 435 according to EU numbering in the Fc region of the H1 chain has been substituted with Arg. This substitution weakens or eliminates the binding activity of the Fc region for the Protein A resin.

20 [Example 2] Method for evaluating the dynamic binding capacity (DBC)

Generally, DBC is determined by depicting the behavior in which the continuously loaded protein is discharged from the column as a breakthrough curve (hereinafter, “BTC”) in a chromatogram by UV monitoring using a purification device connected to a UV detector. The BTC chromatogram when using BiAb is shown in Fig. 1 as an example.

25 The DBC was evaluated by comparing the amount of load at the 5% breakthrough point (BT point) among the antibody molecules and their mixtures.

The following apparatus and such were used for the DBC calculations:

- LC apparatus: AKTA AVANT25 manufactured by GE Healthcare
- Software: Unicorn version 6.1 manufactured by GE Healthcare
- 30 · Protein A resin: Mab Select SuRe (Cat No.17-5438-05) or Hitrap Mab Select SuRe (Cat No. 11-0034-93) manufactured by GE Healthcare
- Buffers:
 - equilibration/preliminary washing – 20 mmol/L Na-phosphate, pH 7.5
 - elution – 50 mmol/L Acetic acid
 - 35 regeneration – 0.1 mol/L NaOH

The method for calculating the DBC was carried out as follows.

The above-mentioned apparatus, software, and resins were used, and the DBC was calculated by performing the chromatography operation as follows.

(1) The load fraction (IgG concentration: P g/L) was once allowed to flow through the LC apparatus without passing it through the column, and the value of OD_{280 nm} for 100% leakage (= 100% BT) was confirmed. This value was denoted as a.

(2) The value obtained by multiplying 0.05 and a was defined as the OD_{280 nm} at 5% BT. This value was denoted as b_{5%}.

(3) The load fraction was allowed to flow continuously through a set amount equilibrated resin (r L), and when the OD_{280 nm} value reached b_{5%}, the volume of the load fraction was read from the chromatogram. This value was denoted as c_{5%} L.

(4) The value obtained by the equation $(P \times c_{5\%})/r$ was calculated as DBC_{5%} which is the dynamic binding capacity at 5% BT.

$$DBC_{5\%} = (P \times c_{5\%})/r \text{ (unit: g/L resin)}$$

When determining DBC_{10%}, calculations were carried out by determining c_{10%} in the same manner.

[Example 3] DBC for each antibody molecule alone

The DBC of each of Q homo, J homo, and BiAb was determined under the following conditions:

Column: Hitrap MabSelect Sure (hereinafter, referred to as MSS) (GE Healthcare), 0.7 x 2.5 cm

Load material: material mimicking the IgG concentration, pH, and conductivity of actual load CM, using each purified antibody standard. IgG concentration: approximately 2 g/L; pH7.5; conductivity: 1.2 S/m. J homo with a purity of approximately 80%, Q homo with a purity of approximately 85%, and BiAb with a purity of approximately 95% were used.

Contact time: 3.4 min (43.75 cm/h)

The results are shown in Table 1.

[Table 1]

IgG	5% BT g/L resin
BiAb	58.0
J Homo	32.8
Q Homo	31.2

The results showed that the DBC of BiAb is significantly higher than the DBC of J

homo and Q homo.

[Example 4] Verification of pH and contact time for the DBC of BiAb

Next, the DBC of BiAb alone when changing pH of the load solution and the contact time on the resin was confirmed, and the effects of both of the parameters were verified. The conditions are shown below:

- Column: MabSelect Sure (GE Healthcare), 1.0 x 20 cm
- Load material: a diluted preparation of a purified BiAb standard mimicking CM (BiAb: 95%): 2 g/L; pH 6.5-8.0 (verification); Conductivity: 1.2 S/m
- Contact time: 3-8 min (verification)

The results are shown in Table 2.

[Table 2]

5% BT g/L resin	150 cm/h 8 min	350 cm/h 3.4 min	400 cm/h 3 min
pH 6.5	Not tested	55.5	Not tested
pH 7.0	Not tested	54.2	Not tested
pH 7.5	63.2	52.3	49.2
pH 8.0	Not tested	51.7	Not tested

[Example 5] The DBC in a mixture of BiAb and Homo

In the culture supernatant actually loaded onto the Protein A resin (hereinafter, “HCCF”), BiAb, J homo, and Q homo are present as a mixture. More specifically, from the viewpoint of recovering BiAb which is the substance of interest, J homo and Q homo can be regarded as substances that compete with BiAb. Therefore, when considering actual production, the verification of the DBC of BiAb under conditions where certain amounts of J homo and Q homo are present in HCCF is meaningful. For this verification, experiments were carried out under the following conditions:

- Column: Hitrap MabSelect Sure (MSS) (GE Healthcare), 0.7 x 2.5cm
- Load Material: a mixture of purified BiAb and Homo standards mimicking CM: 2 g/L; pH 7.5; Cond: 1.2 S/m

Control BiAb (95%) J homo:BiAb:Q homo = 5:95:0

Mimic A J homo:BiAb:Q homo = 10:83:7

Mimic B J homo:BiAb:Q homo = 10:68:22

- Contact time: 3.4 min (43.75 cm/h)

- The load of BTC was fractioned, and the BiAb/Homo ratio at each of the BT points was

confirmed by Analytical CIEC.

Conditions of Analytical CIEC were as follows:

- HPLC apparatus: Alliance 2695/2487 manufactured by Waters
- 5 · Software: Empower3 manufactured by Waters
- CIEC column: ProPac WCX-10, Product No.054993 manufactured by Thermo scientific
- Column temperature: 30°C
- Amount injected: 30 µg/shot
- Buffers:
- 10 Mobile phase A – 9.6 mmol/L Tris, 6.0 mmol/L piperazine, 11.0 mmol/L imidazole, pH 6.0
- Mobile phase B – 9.6 mmol/L Tris, 6.0 mmol/L piperazine, 11.0 mmol/L imidazole, 150 mmol/L NaCl, pH 10.1
- Gradient conditions:

Time (min)	Flow rate (mL/min)	%A	%B
0.0	1.0	100	0
1.0	1.0	100	0
20.0	1.0	0	100
35.0	1.0	0	100

The results are shown in Table 3

15

[Table 3]

Sample		5% BT	10% BT	15% BT	20% BT
Control	g/L resin	58.0	-	-	-
BiAb 95%	J : BiAb : Q	4 : 96 : 0			
Mimic A	g/L resin	51.0	58.2	63.1	65.5
	J : BiAb : Q	5 : 55 : 40	5 : 55 : 40	5 : 57 : 38	6 : 59 : 36
Mimic B	g/L resin	38.8	42.8	45.9	-
	J : BiAb : Q	0 : 3 : 97	0 : 3 : 97	0 : 2 : 98	

The following was found from the above results.

- DBC: J homo $\approx$ Q homo < BiAb
- 20 · Affinity to MSS: Q homo < BiAb < J homo
- Effects of the parameters on BiAb DBC:

pH: In the range of pH 6.5 to 8.0, while lower pH tended to yield higher DBC, the impact was small.

Contact time: In the range of three to eight minutes, while longer contact time tended to yield higher DBC, the DBC was not less than those of Q homo and J homo even at three minutes.

5 Regarding the affinity to MSS, the results reflected the features of the present invention, and it is manifested by the order of leakage in Example 5

10 On the other hand, regarding DBC, it is presumed that the differences in the affinity to the MSS resin and the availability of the ligand produced the results. More specifically, since J homo has two sequences where it binds strongly to the MSS ligand, it is bound to the MSS resin at the two sites. That is, the MSS ligands present in the area that is spatially occupied by J
15 homo cannot be used. On the other hand, since BiAb has only one site with a sequence that strongly binds to MSS, it has spatial freedom higher than J homo, and high DBC could be realized by effective utilization of more MSS ligands. The reason why Q homo has a low DBC would be simply that the binding activity of the molecule as the whole is low. Furthermore, it is thought that Q homo and J homo competitively inhibit the binding of BiAb to MSS.

CLAIMS

1. A method for producing and purifying a heterologous Fc region-containing polypeptide, which comprises the steps of:
 - 5 (a) preparing a first polypeptide chain and a second polypeptide chain of an Fc region having binding activities to a Protein A resin that are different from each other;
 - (b) comparing the dynamic binding capacity of a heterologous Fc region-containing polypeptide comprising the first polypeptide chain and the second polypeptide chain prepared in step (a) for the Protein A resin in Protein A column chromatography with the dynamic binding capacity of a homologous Fc region-containing polypeptide comprising two polypeptide chains having substantially the same binding activity to said resin for the Protein A resin in Protein A column chromatography to confirm the elevated dynamic binding capacity of the former Fc region-containing polypeptide compared with that of the latter Fc region-containing polypeptide;
 - 10 (c) contacting a sample comprising the heterologous Fc region-containing polypeptide with said resin; and
 - 15 (d) collecting the heterologous Fc region-containing polypeptide bound to the resin.
2. The method of claim 1, wherein said step (a) is preparing a first polypeptide chain that binds to the resin as the first polypeptide chain of the Fc region, and preparing a second polypeptide chain that does not bind to the resin or shows weaker binding to the resin compared to the first polypeptide chain, as the second polypeptide chain of the Fc region.
- 20 3. The method of claim 1 or 2, wherein said step (a) is modifying the Fc region of the Fc region-containing polypeptide for purification, so that the first polypeptide chain of the Fc region binds to the resin, and the second polypeptide chain of the Fc region does not bind to the resin or shows weaker binding to the resin compared to the first polypeptide chain.
- 25