

A METHOD FOR THE PREPARATION OF A HIGH-TEMPERATURE STABLE OXYGEN-CARRIER-CONTAINING PHARMACEUTICAL COMPOSITION AND THE USE THEREOF

Cross-Reference to Related Application:

[0001] This application claims priority of U.S. provisional patent application number 61/348,764 filed on May 27, 2010 and U.S. patent application serial no. 13/013,847 filed on 26 January, 2011, the disclosures of which are incorporated by reference.

Technical Field

[0002] The present invention relates to a method for the preparation of a high-temperature stable oxygen-carrier-containing pharmaceutical composition and the composition made by the process. The present invention also relates to the use of the high-temperature stable oxygen carrier-containing pharmaceutical composition for cancer treatment, oxygen-deprivation disorder and organ preservation for humans and other animals.

Background of Invention

[0003] Hemoglobin plays an important role in most vertebrates for gaseous exchange between the vascular system and tissue. It is responsible for carrying oxygen from the respiratory system to the body cells *via* blood circulation and also carrying the metabolic waste product carbon dioxide away from body cells to the respiratory system, where the carbon dioxide is exhaled. Since hemoglobin has this oxygen transport feature, it can be used as a potent oxygen supplier if it can be stabilized *ex vivo* and used *in vivo*.

[0004] Naturally-occurring hemoglobin is a tetramer which is generally stable when present within red blood cells. However, when naturally-occurring hemoglobin is removed from red blood cells, it becomes unstable in plasma and splits into two α - β dimers. Each of these dimers is approximately 32 kDa in molecular weight. These dimers may cause substantial renal injury when filtered through the kidneys and excreted. The breakdown of the tetramer linkage also negatively impacts the sustainability of the functional hemoglobin in circulation.

[0005] In order to prevent breakdown of the tetramer, recent developments in hemoglobin processing have incorporated various cross-linking techniques to create intramolecular bonds within the tetramer as well as intermolecular bonds between the tetramers to form polymeric hemoglobin. The prior art teaches that polymeric hemoglobin is the preferred form in order to increase circulatory half-life of the hemoglobin. However, as determined by the present inventors, polymeric hemoglobin more readily converts to met-hemoglobin in blood circulation. Met-hemoglobin cannot bind oxygen and therefore cannot oxygenate tissue. Therefore, the cross-linking taught by the prior art that causes the formation of polymeric hemoglobin is a problem. There is a need in the art for a technique that permits intramolecular crosslinking to create stable tetramers without the simultaneous formation of polymeric hemoglobin.

[0006] Further problems with the prior art attempts to stabilize hemoglobin include production of tetrameric hemoglobin that includes an unacceptably high percentage of dimer units; the presence of dimers makes the hemoglobin composition unsatisfactory for administration to mammals. The dimeric form of the hemoglobin can cause severe renal injury in a mammalian body; this renal injury can be severe enough to cause death.

Therefore, there is a need in the art to create stable tetrameric hemoglobin with low unwanted dimeric form in the final product.

[0007] Further problems with prior art attempts to create stable hemoglobin include the presence of protein impurities such as immunoglobulin G that can cause allergic effects in mammals. Therefore, there is a need in the art for a process which can produce stable tetrameric hemoglobin without protein impurities.

[0008] Other problems with prior art hemoglobin preparations include vasoconstriction following transfusion in mammals. It has been indicated that this vasoconstriction is due to endothelium-derived relaxing factor binding to reactive sulfhydryl groups of a hemoglobin molecule. Thus, there is a need in the art to create a stabilized tetrameric hemoglobin that does not cause vasoconstriction following transfusion.

[0009] In addition to the above problems, there is a need in the art for a stabilized tetrameric hemoglobin that is phospholipid free and capable of production on an industrial scale.

Summary of Invention

[00010] The present invention provides a method for producing high-temperature stable, purified, cross-linked tetrameric hemoglobin suitable for use in mammals without causing severe renal injury, vascular detrimental effects, or other severe adverse effects (including death). The invention also includes the high-temperature stable, purified, cross-linked tetrameric hemoglobin and the use of the hemoglobin for oxygenation of *in vivo* and *ex vivo* tissue.

[00011] The method includes a starting material of mammalian whole blood including at least red blood cells and plasma. Red blood cells are separated from the plasma in the mammalian whole blood followed by filtering to obtain a filtered red blood cell fraction. The filtered red blood cell fraction is washed to remove plasma protein impurities. The washed red blood cells are disrupted by a controlled hypotonic lysis for a time sufficient to lyse red blood cells without lysing white blood cells in an instant cytolysis apparatus at a flow rate of at 50-1000 liter/hr. Filtration is performed to remove at least a portion of the waste retentate from the lysate. A first hemoglobin solution is extracted from the lysate.

[00012] A first ultrafiltration process is performed using an ultrafiltration filter configured to remove impurities having a higher molecular weight than tetrameric hemoglobin and to further remove any viruses and residual waste retentate from the first hemoglobin solution to obtain a second hemoglobin solution. Flowthrough column chromatography is performed on the second hemoglobin solution to remove protein impurities, dimeric hemoglobin and phospholipids to form a phospholipid-free and low dimer hemoglobin solution. A second ultrafiltration process is performed on the phospholipid-free and low dimer hemoglobin solution using a filter configured to remove impurities resulting in a concentrated purified phospholipid-free and low dimer hemoglobin solution.

[00013] Sulfhydryl groups of hemoglobin molecules are blocked in the concentrated, purified phospholipid-free and low dimer hemoglobin solution by a sulfhydryl reagent in a fully oxygenated environment. The resultant hemoglobin molecules each have at least one cysteine moiety including a thiol-protecting group such that the hemoglobin

molecules are incapable of binding endothelium-derived relaxing factor at the cysteine site.

[00014] At least the α - α and/or β - β subunits of the thiol-protected hemoglobin are cross-linked by bis-3,5-dibromosalicyl fumarate to form high-temperature stable cross-linked tetrameric hemoglobin without the formation of polymeric hemoglobin such that the molecular weight of the resultant cross-linked tetrameric hemoglobin is 60 – 70 kDa. A suitable physiological buffer is exchanged for the cross-linked tetrameric hemoglobin. Any residual non-cross-linked tetrameric hemoglobin and any residual chemicals are removed by using tangential-flow ultrafiltration. N-acetyl cysteine is added at a concentration of 0.2-0.4% to the cross-linked tetrameric hemoglobin to maintain a level of met-hemoglobin below 5%. The phospholipid-free, low dimer, thiol-protected high-temperature stable cross-linked tetrameric hemoglobin is then added to a pharmaceutically acceptable carrier; the pharmaceutically acceptable carrier can be a physiological buffer or water.

[00015] Following this procedure, the resultant hemoglobin is optionally packaged in air-tight polyethylene, ethylene-vinyl-acetate, ethylene-vinyl alcohol (PE, EVA, EVOH) infusion packages. The packaging prevents oxygen contamination which results in the formation of inactive met-hemoglobin.

[00016] The high-temperature stable cross-linked hemoglobin produced by the above method is used for the treatment of various cancers such as leukemia, colorectal cancer, lung cancer, breast cancer, nasopharyngeal cancer and esophageal cancer. The mechanism for destroying cancer cells is to improve oxygenation in tumor cells, thereby enhancing the sensitivity towards radiation and chemotherapeutic agents. The high-

temperature stable cross-linked tetrameric hemoglobin is also used for preservation of organ tissue during transplant or for preservation of the heart in situations where there is a lack of oxygen supply *in vivo*, such as in an oxygen-deprived heart.

Brief description of the drawings

[00017] FIG. 1 depicts the amino acid sequence alignment of different hemoglobins.

[00018] FIG. 2 is a flow-chart depicting an overview of the process of the present invention.

[00019] FIG. 3 schematically depicts an instant cytolysis apparatus used in the process of the present invention.

[00020] FIG. 4 is a graph showing the reaction of hemoglobin with sulfhydryl reagent in oxygenated and in deoxygenated environments.

[00021] FIG. 5 depicts high performance liquid chromatography analysis for cross-linked tetrameric hemoglobin.

[00022] FIG. 6 depicts electrospray ionization mass spectrometry (ESI-MS) analysis for the cross-linked tetrameric hemoglobin.

[00023] FIG. 7 shows a circular dichroism (CD) spectroscopy analysis for (a) purified hemoglobin solution and (b) cross-linked tetrameric hemoglobin.

[00024] FIG. 8 shows the chemo-sensitization effects of cross-linked tetrameric hemoglobin *in vitro*.

[00025] FIG. 9 depicts the improvement of oxygenation in normal tissue using the cross-linked tetrameric hemoglobin of the present invention.

[00026] FIG. 10 shows the improvement of oxygenation in an extremely hypoxic tumor area using the cross-linked tetrameric hemoglobin of the present invention.

[00027] FIG. 11 shows the mean arterial pressure changes in a rat model of severe hemorrhagic shock after the treatment with the stabilized cross-linked tetrameric hemoglobin of the present invention.

[00028] FIG. 12 is an elution profile for flowthrough column chromatography; the hemoglobin solution is in the flowthrough fraction.

[00029] FIG. 13 schematically depicts a flowthrough CM column chromatography system with ultrafiltration for an industrial scale operation.

[00030] FIG. 14 is a comparison between sulfhydryl reaction in an oxygenated environment and reaction in a deoxygenated environment.

[00031] FIG. 15 is a graph demonstrating the heat stability of the cross-linked tetrameric hemoglobin of the present invention compared to prior art hemoglobin.

[00032] FIG. 16 is a schematic depiction of an infusion bag for the cross-linked tetrameric hemoglobin of the present invention.

[00033] FIG. 17 is a schematic depiction of an apparatus used to test the formation of met-hemoglobin *in vitro*.

[00034] FIG. 18 depicts the rate of met-hemoglobin formation for polymeric hemoglobin and the inventive hemoglobin in the apparatus of FIG. 17.

Detailed Description of Invention

[00035] Hemoglobin is an iron-containing oxygen-transport protein in red blood cells of the blood of mammals and other animals. Hemoglobin exhibits characteristics of both

the tertiary and quaternary structures of proteins. Most of the amino acids in hemoglobin form alpha helices connected by short non-helical segments. Hydrogen bonds stabilize the helical sections inside the hemoglobin causing attractions within the molecule thereto folding each polypeptide chain into a specific shape. A hemoglobin molecule is an assembly of four globular protein subunits. Each subunit is composed of a polypeptide chain arranged into a set of α -helix structural segments connected in a “myoglobin fold” arrangement with an embedded heme group.

[00036] The heme group consists of an iron atom held in a heterocyclic ring, known as a porphyrin. The iron atom binds equally to all four nitrogen atoms in the center of the ring which lie in one plane. Oxygen is then able to bind to the iron center perpendicular to the plane of the porphyrin ring. Thus a single hemoglobin molecule has the capacity to combine with four molecules of oxygen.

[00037] In adult humans, the most common type of hemoglobin is a tetramer called hemoglobin A consisting of two α and two β non-covalently bound subunits designated as $\alpha_2\beta_2$, each made of 141 and 146 amino acid residues respectively. The size and structure of α and β subunits are very similar to each other. Each of the subunits has a molecular weight of about 16 kDa for a total molecular weight of the tetramer of about 65 kDa. The four polypeptide chains are bound to each other by salt bridges, hydrogen bonds and hydrophobic interaction. The structure of bovine hemoglobin is similar to human hemoglobin (90.14% identity in α chain; 84.35% identity in β chain). The difference is that the two sulfhydryl groups in the bovine hemoglobin are positioned at β Cys 93, while the sulfhydryls in human hemoglobin are at positioned at a Cys 104, β Cys 93 and β Cys 112 respectively. FIG. 1 shows the amino acid sequences alignment of

bovine, human, canine, porcine and equine hemoglobin, respectively labeled B, H, C, P, and E. The unlike amino acids from various sources are shaded. FIG. 1 indicates that human hemoglobin shares high similarity with bovine, canine, porcine and equine hemoglobin when comparing their amino acid sequences.

[00038] In naturally-occurring hemoglobin inside red blood cells, the association of an α chain with its corresponding β chain is very strong and does not disassociate under physiological conditions. However, the association of one $\alpha\beta$ dimer with another $\alpha\beta$ dimer is fairly weak outside red blood cells. The bond has a tendency to split into two $\alpha\beta$ dimers each approximately 32 kDa. These undesired dimers are small enough to be filtered by the kidneys and be excreted, with the result being potential renal injury and substantially decreased intravascular retention time.

[00039] Therefore, it is necessary to stabilize any hemoglobin that is used outside of red blood cells both for efficacy and safety. The process for producing the stabilized hemoglobin is outlined below; an overview of the process of the present invention is presented in the flow chart of FIG. 2.

[00040] Initially, a whole blood source is selected as a source of hemoglobin from red blood cells. Mammalian whole blood is selected including, but not limited to, human, bovine, porcine, equine, and canine whole blood. The red blood cells are separated from the plasma, filtered, and washed to remove plasma protein impurities.

[00041] In order to release the hemoglobin from the red blood cells, the cell membrane is lysed. Although various techniques can be used to lyse red blood cells, the present invention uses lysis under hypotonic conditions in a manner which can be precisely controlled at volumes suitable for industrial-scale production. To this end, an

instant cytolysis apparatus as seen in FIG. 3 is used to lyse the red blood cells. Hypotonic lysis creates a solution of lysate including hemoglobin and a waste retentate. To enable industrial-scale production, the lysis is carefully controlled such that only red blood cells are lysed without lysing white blood cells or other cells. In one embodiment, the size of the instant cytolysis apparatus is selected such that the red blood cells traverse the apparatus in approximately 30 seconds and the instant cytolysis apparatus includes a static mixer. Deionized and distilled water is used as a hypotonic solution. Of course it is understood that the use of other hypotonic solutions having different saline concentrations would result in different time periods for red blood cell lysis. Because the controlled lysis procedure breaks the red blood cells only, not white blood cells or cellular matter, it minimizes the release of toxic proteins, phospholipids or DNA from white blood cells and other cellular matter. A hypertonic solution is added immediately after 30 seconds, that is, after the red blood-cell containing solution has traversed the static mixer portion of the instant cytolysis apparatus. The resultant hemoglobin has a higher purity and lower levels of contaminants such as undesired DNA and phospholipids than hemoglobin resulted from using other lysis techniques. Undesired nucleic acid from white blood cells and phospholipids impurities are not detected in the hemoglobin solution by polymerase chain reaction (detection limit = 64 pg) and HPLC (detection limit = 1 µg/mL) method respectively.

[00042] Two ultrafiltration processes are performed; one which removes impurities having molecular weights greater than hemoglobin before flowthrough column chromatography, and another which removes impurities having molecular weights less than hemoglobin after flowthrough column chromatography. The latter ultrafiltration

process concentrates the hemoglobin. In some embodiments, a 100 kDa filter is used for the first ultrafiltration, while a 30 kDa filter is used for the second ultrafiltration.

[00043] Flowthrough column chromatography is used to remove protein impurities in the purified hemoglobin solution such as immunoglobulin-G, albumin and carbonic anhydrase. In some embodiments, column chromatography is carried out by using one or a combination of commercially available ion exchange columns such as a DEAE column, CM column, hydroxyapatite column, etc. The pH for column chromatography is typically from 6 to 8.5. In one embodiment, a flowthrough CM column chromatography step is used to remove protein impurities at pH 8.0. Enzyme-linked immunosorbent assay (ELISA) is performed to detect the protein impurities and phospholipids remaining in the sample after elution from the column chromatography. This unique flowthrough column chromatography separation enables a continuous separation scheme enabling industrial-scale production. The ELISA result shows that the amount of these impurities are substantially low in the eluted cross-linked tetrameric hemoglobin (immunoglobulin-G: 44.3 ng/mL; albumin: 20.37 ng/mL; carbonic anhydrase: 81.2 µg/mL). The protein impurities removal results using different kinds of column with different pH values are shown in Table 1 below.

[00044] Table 1: Removal of different protein impurities using different ion-exchange columns

Column (pH condition)	Removal percentage (%)		
	Carbonic anhydrase	Albumin	Immunoglobulin-G
DEAE (at pH 7.5)	---	68	29.8
DEAE (at pH 7.8)	---	60	50.9
CM (at pH 6.2)	---	32	21.8
CM (at pH 8.0)	5.6	53.2	66.4
Hydroxyapatite (at pH 7.5)	4.5	23.5	22.8

[00045] In order to prevent the binding of endothelium-derived relaxing factor to a cysteine site of the hemoglobin causing undesirable vasoconstriction, the hemoglobin is subjected to a reaction with a sulfhydryl reagent under oxygenated conditions. This is in direct contrast to the teachings of the prior art which emphasize reaction between hemoglobin and a sulfhydryl reagent under deoxygenated conditions. The present invention demonstrates that the sulfhydryl reagent reacts faster and in completion with the reactive sulfhydryl groups of hemoglobin under an oxygenated condition. The reaction of sulfhydryl reagent with sulfhydryl groups of hemoglobin produces iodide. Therefore, the completeness of the alkylation reaction can be monitored by measuring the iodide release. During a time-course experiment, the alkylation reaction is performed faster and more efficiently in an oxygenated environment when compared to a deoxygenated environment, as indicated in FIG. 4 and FIG. 14. The reaction time required to reach completion is reduced to under 5 hours in an oxygenated environment when compared to a deoxygenated environment. A shorter reaction time is very

important for an industrial-scale process. It also reduces reactions caused by unwanted impurities that lead to adverse effects in the final product.

[00046] Following the sulfhydryl reaction process, the hemoglobin is subjected to α - α and/or β - β cross-linking by bis-3,5-dibromosalicyl fumarate (DBSF). In order to prevent formation of polymeric hemoglobin, the reaction is carefully controlled in a deoxygenated environment with a molar ratio of hemoglobin to DBSF between 1:2.5 to 1:4.0 such that the resultant α - α and/or β - β cross-linked hemoglobin is tetrameric hemoglobin having a molecular weight of 60-70 kDa, demonstrating that polymeric hemoglobin is not present. The yield of the DBSF reaction is high, > 99% and the dimer concentration in the final product is low; in the context of this invention, low dimer content means less than 5% and, more preferably, less than 2% dimer. In one embodiment, the cross-linking is selected such that the β - β cross-linked hemoglobin is greater than 50% of the total cross-linked tetrameric hemoglobin. In another embodiment, the cross-linking is selected such that the β - β cross-linked hemoglobin is 60% or greater of the total. Further, conditions can be selected such that the β - β cross-linked hemoglobin is greater than 70% of the total tetrameric hemoglobin. There is some evidence that β - β cross-linked hemoglobin may have a lower oxygen affinity than α - α cross-linked tetrameric hemoglobin. Therefore, there may be situations in which a higher percentage of β - β cross-linked hemoglobin is preferred to α - α cross-linked hemoglobin to enhance the efficiency of oxygen transport. Further, β - β cross-linked hemoglobin may result in fewer α - β dimers which would assist in reducing renal toxicity.

[00047] N-acetyl cysteine is added at a concentration of 0.2-0.4% to the α - α and/or β - β cross-linked tetrameric hemoglobin to maintain a level of met-hemoglobin below 5%.

[00048] Depending upon the final application of the hemoglobin, the purified, cross-linked tetrameric hemoglobin of the present invention is optionally packaged in an air-tight package in a deoxygenated environment. The packaging used in the present invention results in the α - α and/or β - β cross-linked tetrameric hemoglobin being stable for more than two years. In contrast, the hemoglobin of the present invention converts into inactive met-hemoglobin rapidly within few days under oxygenated conditions. Prior art hemoglobin solutions have been packaged in PVC or Stericon blood bags which have high oxygen permeability, thus shortening the life span of the product.

[00049] In many embodiments of the present invention, the oxygen-carrier-containing cross-linked hemoglobin-containing pharmaceutical composition is delivered by intravenous injection. Therefore, the packaging design and material choice are directed towards intravenous injection applications. A multi-layer package of EVA/EVOH material is used to minimize the gas permeability and to avoid the formation of inactive met-hemoglobin. A 100 mL infusion bag designed for use with the purified and cross-linked hemoglobin of the present invention is made from a five layers EVA/EVOH laminated material with a thickness of 0.4 mm that has an oxygen permeability of 0.006-0.132 cm³ per 100 square inches per 24 hours per atmosphere at room temperature. This material is a Class VI plastic (as defined in USP<88>), which meets the *in-vivo* Biological Reactivity Tests and the Physico-Chemical Test and is suitable for fabricating an infusion bag for intravenous injection purpose. This primary bag is particularly useful to protect the α - α and/or β - β cross-linked tetrameric hemoglobin solution from long term oxygen exposure that cause its instability and eventually affects its therapeutic properties.

[00050] For secondary protection of blood products, it has been known to use aluminum overwrap to protect against potential air leakage and to maintain the product in a deoxygenated state. However, there is a potential of pin holes in the aluminum overwrap that compromise its air tightness and make the product unstable. Therefore the present invention uses as secondary packaging an aluminum overwrap pouch which prevents oxygenation and also prevents light exposure. The composition of the overwrap pouch includes 0.012mm of polyethylene terephthalate (PET), 0.007mm of aluminum (AL), 0.015mm of nylon (NY) and 0.1mm of polyethylene (PE). The overwrap film has a thickness of 0.14mm and an oxygen transmission rate of 0.006 cm³ per 100 square inches per 24 hours per atmosphere at room temperature. This secondary packaging lengthens the stability time for the hemoglobin, extending the product shelf-life.

[00051] High performance liquid chromatography (HPLC), electrospray ionization mass spectrometry (ESI-MS), and circular dichroism (CD) spectroscopy are used to analyze and characterize the α - α and/or β - β cross-linked tetrameric hemoglobin. For a bovine blood source, FIG. 5 shows the composition of the product in terms of molecular weight distribution by HPLC analysis. An HPLC analytical method is used to detect the amount of tetramer and dimer respectively. The mobile phase for HPLC analysis contains magnesium chloride (0.75M) which can separate dimer, uncross-linked tetramer, and stabilized α - α and/or β - β cross-linked tetramer. For promoting hemoglobin dissociation into dimers, magnesium chloride is approximately 30 times more effective than sodium chloride at the same ionic strength.

[00052] ESI-MS allows the analysis of very large molecules. It is an ionization technique that analyzes the high molecular weight compound by ionizing the protein, and

then separating the ionized protein based on mass/charge ratio. Therefore, the molecular weight and the protein interactions can be determined accurately. In FIG. 6, ESI-MS analysis result indicates that the size of stabilized tetramer is 65 kDa. The far UV CD spectra from 190 to 240 nm reveal the secondary structures of globin portion of the hemoglobin. In FIG. 7, the consistency of the spectra of the purified and cross-linked hemoglobin reveals that the hemoglobin chains are properly folded even after the cross-linking by DBSF. The CD result shows that the cross-linked hemoglobin has around 42 % of alpha-helix, 38 % of beta-sheet, 2.5 % of beta-turn and 16 % of random coil. It further confirms that cross-linking step with DBSF to form cross-linked tetrameric hemoglobin does not affect the secondary structure of hemoglobin.

[00053] The purified cross-linked tetrameric hemoglobin produced by the process of the present invention has a molecular weight of 60-70 kDa and has at least one cysteine moiety wherein said cysteine moiety includes a thiol-protecting group such that the hemoglobin is incapable of binding endothelium-derived relaxing factor at the cysteine site. Further, the cross-linked tetrameric hemoglobin is non-pyrogenic, endotoxin free (< 0.05 EU/mL), and stroma free (< 1%).

[00054] The process of the present invention is applicable to large-scale industrial production of cross-linked tetrameric hemoglobin. In addition, the cross-linked tetrameric hemoglobin in combination with a pharmaceutical carrier (e.g. water, physiological buffer, in capsule) is suitable for mammalian use.

[00055] The cross-linked tetrameric hemoglobin of the present invention is used for tissue oxygenation, for cancer treatment, for the treatment of an oxygen-deprivation disorder such as hemorrhagic shock, and in heart preservation under a low oxygen

content environment (e.g. heart transplant). The dosage of cross-linked tetrameric hemoglobin is selected at a concentration range of approximately 0.3-1.3 g/kg.

[00056] For use in cancer treatment, the oxygen-carrier-containing pharmaceutical composition of the present invention serves as a tissue oxygenation agent to improve the oxygenation in tumor tissues, thereby enhancing chemo-sensitivity (e.g., sensitivity to chemotherapy) and radiation sensitivity.

[00057] FIG. 8 demonstrates the enhanced chemo-sensitivity of cancer tumor cells after applying a composition including cross-linked tetrameric hemoglobin *in vitro*. Five different cancer cell lines (A) Jurkat (leukemia), (B) HKESC1 (Esophageal cancer), (C) COLO205 (Colon cancer), (D) A549/Cisp (Lung cancer) and (E) MCF-7/ADM (Breast cancer) are treated with various chemotherapeutic agents alone, or in combination with the cross-linked tetrameric hemoglobin of the present invention. Inhibition of tumor cell growth is determined by the ATP tumor chemo-sensitivity assay (ATP-TCA) (for Jurkat, COLO205, A549/Cisp, and MCF-7/ADM cell lines) or MTT cell proliferation assay (for HKESC1 cell line). The results show that the chemo-sensitivity is highly enhanced by addition of the inventive cross-linked tetrameric hemoglobin in all cancer cell lines including two cell lines, A549/Cisp and MCF-7/ADM, which are highly resistant to chemotherapy. The results in FIG. 8 show that as a result of the enhanced chemo-sensitivity by addition of the cross-linked tetrameric hemoglobin, the treatment efficacy for leukemia cells, esophageal cancer cells, lung cancer cells, colon cancer cells, and breast cancer cells is greatly increased.

[00058] In addition, the ability of the inventive cross-linked tetrameric hemoglobin to improve oxygenation in normal tissues (FIG. 9) and in extremely hypoxic tumor tissue

(FIG. 10) (human nasopharyngeal carcinoma (CNE2)), is demonstrated in this invention. The representative oxygen profile along the tissue track of a human CNE2 xenograft is showed in FIG. 10. Oxygen partial pressure within the tumor mass is directly monitored by a fiber-optic oxygen sensor (Oxford Optronix Limited) coupled with a micro-positioning system (DTI Limited). After intravenous injection of 1.2 g/kg of the cross-linked tetrameric hemoglobin, the median pO₂ value rises from 0.2 mmHg to 3.9 mmHg in 3 hours and 10.6 mmHg in 6 hours respectively. Even in the most hypoxic regions, the oxygen carrier-containing pharmaceutical composition of the present invention significantly increased oxygen tension. No other similar commercial products or any existing technologies show as high an efficacy when compared to the oxygen-carrier-containing pharmaceutical composition prepared in this invention.

[00059] For the use in the treatment of oxygen-deprivation disorder and in heart preservation, the oxygen-carrier-containing pharmaceutical composition of the present invention serves as a blood substitute providing oxygen to a target organ. In some embodiments, the composition is used as a cardioplegic solution for heart preservation.

[00060] The mean arterial pressure changes in a rat model of severe hemorrhagic shock (see Example 12b) after the treatment with 0.5 g/kg of the cross-linked tetrameric hemoglobin of the present invention are showed in FIG. 11. In a rat model of severe hemorrhagic shock, the mean arterial pressure is returned back to a safe and stable level and maintained below the baseline after treatment with the stabilized cross-linked tetrameric hemoglobin. Following treatment with the hemoglobin of the present invention, the reaction time for the mean arterial pressure to return to normal is even shorter than that following administration of rat whole blood which serves as a positive

control. The above results indicate that the vaso-active event after the transfusion of the inventive hemoglobin is beneficial to maintain a stable and hemodynamic state. Prior hemoglobin-based oxygen carriers caused many vasoconstriction events. For example, the Hemopure® product (Biopure Co., USA) resulted in higher mean arterial pressure (124 ± 9 mmHg) when compared to the baseline (96 ± 10 mmHg) as disclosed by Katz et al., 2010.

[00061] Use of the inventive hemoglobin for infusion increases the survival rate in a shock model in mouse (from 18% to 75%) and beagle dogs (from 46% to 97%). The survival rate of a severe hemorrhagic shock animal model is significantly increased by infusing different amounts of the cross-linked tetrameric hemoglobin (see Example 12). Thus the inventive hemoglobin is used as a treatment for hemorrhagic shock.

Examples

[00062] The following examples are provided by way of describing specific embodiments of the invention without intending to limit the scope of the invention in any way.

[00063] **Example 1**

[00064] **Overall process**

[00065] A schematic flow diagram of the process of the present invention is illustrated in FIG. 2. Bovine whole blood is collected into an enclosed sterile container/bag containing 3.8% (w/v) tri-sodium citrate solution as anti-coagulant. Blood is then immediately mixed well with tri-sodium citrate solution to inhibit blood clots. Red blood cells (RBC) are isolated and collected from plasma and other smaller blood cells by an

apheresis mechanism. A “cell washer” is used for this procedure with gamma sterilized disposable centrifuge bowl. RBC are washed with an equal volume of 0.9% (w/v sodium chloride) saline.

[00066] Washed RBC are lysed to release hemoglobin content by manipulating hypotonic shock to the RBC cell membrane. A specialized instant cytolysis apparatus for RBC lysis device depicted in FIG. 3 is used for this purpose. Following RBC lysis, hemoglobin molecules are isolated from other proteins by tangential-flow ultrafiltration using 100 kDa membrane. Hemoglobin in the filtrate is collected for flowthrough column chromatography and further concentrated to 12-14 g/dL by a 30 kDa membrane. Column chromatography is carried out to remove the protein impurities.

[00067] The concentrated hemoglobin solution is first modified by a sulfhydryl reagent (alkylation reaction) under natural oxygenated condition, then the components are reacted with DBSF to form stable α - α and/or β - β cross-linked tetrameric hemoglobin molecules.

[00068] **Example 2**

[00069] **Time & Controlled Hypotonic lysis and filtration**

[00070] Bovine whole blood is freshly collected and transported under a cool condition. The red blood cells are separated from the plasma *via* a cell washer and subsequently with a 0.65 μ m filtration. After washing the red blood cells (RBC) filtrate with 0.9% saline, the filtrate is disrupted by hypotonic lysis. The hypotonic lysis is performed by using the instant cytolysis apparatus depicted in FIG. 3. The instant cytolysis apparatus includes a static mixer to assist in cell lysis. A RBC suspension with

controlled hemoglobin concentration (12-14 g/dL) is mixed with 4 volumes of purified water to generate a hypotonic shock to RBC cell membranes. The period of hypotonic shock is controlled to avoid unwanted lysis of white blood cells and platelets. The hypotonic solution passes through the static mixer portion of the instant cytolysis apparatus for approximately 30 seconds. The shock is terminated after 30 seconds by mixing the lysate with 1/10 volume of hypertonic buffer as it exits the static mixer. The hypertonic solution used is 0.1M phosphate buffer, 7.4% NaCl, pH 7.4. The instant cytolysis apparatus of FIG. 3 can process at 50 to 1000 liters of lysate per hour and, preferably at least 300 liters per hour in a continuous manner.

[00071] Following the RBC lysis, the lysate of red blood cells is filtered by a 0.22 μ m filter to obtain a hemoglobin solution. Nucleic acid from white blood cells and phospholipids impurities are not detected in the hemoglobin solution by polymerase chain reaction (detection limit = 64 pg) and HPLC (detection limit = 1 μ g/mL) method respectively. A first 100 kDa ultrafiltration is performed to remove impurities having a higher molecular weight than hemoglobin. A flowthrough column chromatography is followed to further purify the hemoglobin solution. A second 30 kDa ultrafiltration is then performed to remove impurities having a lower molecular weight than hemoglobin and for concentration.

[00072] **Example 3**

[00073] **Viral clearance study on stroma-free hemoglobin solution**

[00074] In order to demonstrate the safety of the product from this invention, the virus removal abilities of (1) 0.65 μ m diafiltration step and (2) 100 kDa ultrafiltration step are

demonstrated by virus validation study. This is done by the deliberate spiking of a down-scaled version of these two processes with different model viruses (encephalomyocarditis virus, pseudorabies virus, bovine viral diarrhoea virus and bovine parvovirus). In this study, four types of viruses (see the following table 2) were used. These viruses vary in their biophysical and structural features and they display a variation in resistance to physical and chemical agents or treatments.

[00075] Table 2

Target Virus	Model Virus	Taxonomy	Genome	Structure	Size [nm]	Stability*
Hepatitis C virus (HCV)	Bovine viral diarrhea virus (BVDV)	Flaviviridae	ssRNA	enveloped	40-60	low
-	Encephalomyocarditis virus (EMCV)	Picornavirus	ssRNA	non-enveloped	25-30	medium
Parvovirus B19	Bovine parvovirus (BPV)	Parvoviridae	ssDNA	non-enveloped	18-26	very high
Hepatitis B virus (HBV)	Pseudorabies virus (PRV)	Herpesviridae	dsDNA	enveloped	120-200	Low to medium

[00076] The virus validation scheme is briefly shown in the following table 3.

[00077] Table 3

Diafiltration	Ultrafiltration
Cell Washing ↓ Virus spiking ↓ Diafiltration ↓ Virus tests	Virus spiking ↓ Ultrafiltration ↓ Virus tests

[00078] The summary of the log reduction results of the 4 viruses in (1) 0.65 μm diafiltration and (2) 100 kDa ultrafiltration is shown below in Table 4. All four viruses, BVDV, BPV, EMCV and PRV, were effectively removed by 0.65 μm diafiltration and 100 kDa ultrafiltration.

[00079] Table 4

Viruses	BVDV		BPV		EMCV		PRV	
Run	1	2	1	2	1	2	1	2
0.65 μ m Diafiltration	2.69	3.20	3.73	3.53	3.25	= 3.90	2.67	2.63
100 kDa Ultrafiltration	= 4.68	= 4.38	5.87	5.92	3.60	3.43	= 6.05	3.27
Cumulative maximum	= 7.88		9.65		= 7.50		= 8.72	
Cumulative minimum	= 7.07		9.40		6.68		5.90	

Annotation:

= no residual infectivity determined

[00080] **Example 4**

[00081] **Flowthrough column chromatography**

[00082] A CM column (commercially available from GE healthcare) is used to further remove any protein impurities. The starting buffer is 20mM sodium acetate (pH 8.0), and the elution buffer is 20mM sodium acetate, 2M NaCl (pH 8.0). After the equilibration of the CM column with starting buffer, the protein sample is loaded into the column. The unbound protein impurities are washed with at least 5 column volume of starting buffer. The elution is performed using 25% elution buffer (0-0.5M NaCl) in 8 column volume. The elution profile is shown in FIG. 12; the hemoglobin solution is in the flowthrough fraction.

[00083] The purity of flowthrough fraction is analyzed by ELISA. The results are indicated in the following table 5.

[00084] Table 5

	Protein impurities		
	Immunoglobulin-G	Carbonic anhydrase	Albumin
Before CM column	1320 ng/mL	860.3 µg/mL	435.2 ng/mL
Flowthrough (containing hemoglobin)	44.3 ng/mL	81.2 µg/mL	20.4 ng/mL

[00085] As the hemoglobin solution is in the flowthrough from the CM column chromatography at pH 8 (not in the eluate), it is a good approach for continuous industrial scale operation. The first ultrafiltration set-up is connected directly to the flowthrough CM column chromatography system, and the flowthrough tubing can connect to the second ultrafiltration set-up for industrial scale operation. The schematic industrial process configuration is shown in FIG. 13.

[00086] **Example 5**

[00087] **Sulfhydryl reaction and cross-linking**

[00088] (5a) Sulfhydryl reaction

[00089] In the present invention, reaction between hemoglobin and sulfhydryl is performed in an oxygenated environment in contrast to the teachings of the prior art, in which the reaction typically takes place in an inert atmosphere such as nitrogen. An alkylating sulfhydryl reagent is added to alkylate the free sulfhydryl groups of hemoglobin. In this embodiment, the molar ratio of hemoglobin to sulfhydryl reagent is

1:2 to 1:4. This reaction can eliminate the binding of endothelium-derived relaxing factor which reacts with hemoglobin sulfhydryl groups. Endothelium-derived relaxing factor has been demonstrated to bind to the reactive sulfhydryl groups of hemoglobin, which may account for the increase of blood pressure observed after the infusion of earlier generations of hemoglobin-based oxygen carriers. The completion of the sulfhydryl reaction can be monitored by measuring the release of product or the residual sulfhydryl reagent (by UV spectrometry, at 265nm). FIG. 14 is a comparison between reaction in an oxygenated environment and reaction in a deoxygenated environment. FIG. 14 shows that the residual sulfhydryl reagent levels off at 3 hours in an oxygenated environment during a time-course experiment. In contrast, a lot of unreacted sulfhydryl reagent remains in a deoxygenated environment. The yield of the sulfhydryl reaction in an oxygenated environment is high (92.7%) after 4 hours of reaction.

[00090] (5b) Cross-linking reaction to form stable tetramers

[00091] The α - α and/or β - β cross-linking reaction is carried out in a deoxygenated environment. Bis-3,5-dibromosalicyl fumarate (DBSF) is added to the hemoglobin solution to form cross-linked tetrameric hemoglobin that includes α - α cross-linking and/or β - β cross-linking within the tetramer. Optionally, some α - β cross-linking is included within the tetramer.

[00092] In this embodiment, the molar ratio of hemoglobin and DBSF is between 1:2.5 to 1:4.0. The DBSF stabilization procedure stabilizes the tetrameric form of hemoglobin (65 kDa) and prevents dissociation into dimers (32 kDa) which are excreted through the kidneys. In this cross-linking process, only tetrameric hemoglobin is formed;

no polymeric hemoglobin is formed. This process is carried out in an inert atmosphere of nitrogen to prevent oxidation of the hemoglobin to form ferric met-hemoglobin which is physiologically inactive. The completeness of DBSF reaction is monitored by measuring the residual DBSF using HPLC. The yield of DBSF reaction is high, > 99%.

[00093] In one embodiment, the cross-linking reaction is carried out in a deoxygenated environment (dissolved oxygen < 0.1mg/L) at room temperature (15 – 25°C). DBSF is added to the hemoglobin solution to form cross-linked tetrameric hemoglobin; this cross-linked tetrameric hemoglobin is characterized as set forth below. Note that this characterization is based on one embodiment only. The various proportions and types of cross-links within the tetramer can be controlled by the amount of DBSF used, the time and temperature of the reaction, and other process conditions.

[00094] When changing the molar ratio of hemoglobin and DBSF to 1:5.0, the percentage of octamer (130 kDa) is sharply increased to 18%.

[00095] (5b-1) Electrophoretic analysis on cross-linked hemoglobin using 15% SDS-PAGE

[00096] Cross-linked hemoglobin solution was mixed with an equal volume of reducing sample buffer (62 mM Tris-HCl (pH 6.8), 10% (v/v) glycerol, 5% (v/v) mercaptoethanol and 2.3% (w/v) SDS), and heated at 95°C for 10 min. The sample mixtures were resolved using a 15% acrylamide slab gel with a 4% stacking gel. The electrophoresis was run with a constant current of 60 mA. After electrophoresis, the SDS-PAGE gel was stained with 0.1% (w/v) Coomassie blue R350, 20% (v/v) methanol, and 10% (v/v) acetic acid. The intensities of protein bands expressed in Black Light Unit (BLU) were quantified using Bio-Rad Quantity One Software.

[00097] (5b-2) Trypsin Digestion of protein bands excised from 15% SDS-PAGE

[00098] Protein bands were excised from the SDS-PAGE gel, cut into cubes (1 X 1 mm), destained with 10% methanol/10% acetic acid. The destained gel were reduced with 10mM DTT in 25mM NH_4CO_3 and alkylated with 55mM iodoacetamide in 25mM NH_4CO_3 for 45 min in dark, and then in-gel digested with 20 ng/ μL modified trypsin in 25mM NH_4CO_3 at 37°C overnight. After trypsin digestion, the trypsin-digested peptides were extracted by diffusion into 50% (v/v) acetonitrile and 1% (v/v) trifluoroacetic acid (TFA).

[00099] (5b-3) Matrix-assisted laser dissociation/ ionization – time of flight mass spectrometry (MALDI-TOF MS) analysis of trypsin-digested protein bands

[000100] The trypsin digested peptides extracted from the gel cubes were spotted onto Anchorchip plate, which was pre-spotted with 1 μL of matrix solution (2 mg/mL HCCA, saturated in 50% acetonitrile/ 0.1% TFA, and allowed to air-dry. After drying, the sample spot was washed with 10mM monophosphate buffer and recrystallized using a solution of EtOH: acetone: 0.1% TFA (6:3:1 ratio). MALDI-TOF MS analysis was performed with a Bruker Autoflex III (Bruker Daltonic GmbH, Bremen, Germany) operated in the reflectron mode over the m/z range of 800–3500 Da and the parameters were set as follows: ion source 25kV for peptide mass fingerprint (PMF), and reflector 26.3 kV for PMF. External calibration was performed using Bruker Peptide Mix Calibration Standard. The peaks with ratio of S/N over 4 were automatically labelled by Flex-Analysis (Bruker Daltonic GmbH, Bremen, Germany). The MS data was further analyzed through MASCOT 2.2.04 and Biotools 2.1 softwares (Bruker Daltonic GmbH, Bremen, Germany), and these data were searched against Mammalian proteins in NCBI

nonredundant (NCBI nr) database. The following parameters were used for database searches: monoisotopic mass accuracy < 250 ppm, parent charge +1, missed cleavages 1, carbamidomethylation of cysteine as fixed modification, oxidation of methionine as variable modification. The results with protein score more than 67 ($p = 0.05$) based on PMF was considered as positive identification. Other criteria for confident identification were that the protein match should have at least 15% sequence coverage and match at least 4 peptides.

[000101] After combination the analysis results from (a) 15% SDS-PAGE and (b) MALDI-TOF MS, the composition of cross-linked hemoglobin is determined. From the results, $\geq 70\%$ of cross-linked hemoglobin is β - β linked.

[000102] **Example 6**

[000103] **Maintaining a low level of inactive met-hemoglobin in the product after formulation**

[000104] When compared to other oxygen carrier pharmaceutical products or a product formed in accordance with the methods disclosed in US patent 7494974 B2 and US patent 7504377 B2, the product from this invention contains low levels of inactive met-hemoglobin molecules. In this embodiment, an anti-oxidant such as N-acetyl cysteine is added at 0.2% to the cross-linked tetrameric hemoglobin. Without adding the anti-oxidant, inactive met-hemoglobin is found at a high level (12-20%). The product from this invention has a low level of inactive met-hemoglobin (< 5%) and it is high-temperature stable resulting in greater efficacy when used in treatment.

[000105] To demonstrate the stability of the product from this invention, a heat stability test is performed at 80°C. The results are depicted in FIG. 15. The product

made according to US patent 7494974 and US patent 7504377, shows a high met-hemoglobin content (22-28%). However, the product of the present invention shows a low level of met-hemoglobin (< 5%).

[000106] **Example 7**

[000107] **Packaging & Product stability**

[000108] Because the product of the present invention is stable under de-oxygenated conditions, the packaging for the product is important to minimize gas permeability. For intravenous application a custom designed, 100 mL infusion bag is made from a five layers EVA/EVOH laminated material with a thickness of 0.4 mm that has an oxygen permeability of 0.006 to 0.132 cm³ per 100 square inches per 24 hours per atmosphere at room temperature. This specific material is a Class VI plastic (as defined in USP<88>), which meets the *in-vivo* biological reactivity tests and the physico-chemical test and are suitable for fabricating containers for intravenous injection purpose (note that other forms of packaging can be made from this material as well depending upon the desired application). A secondary packaging aluminum overwrap pouch is also applied to the primary packaging infusion bag that provides additional barrier in minimizing light exposure and oxygen diffusion. The composition of the pouch consists of: 0.012mm of Polyethylene terephthalate (PET), 0.007mm of Aluminum (AL), 0.015mm of Nylon (NY) and 0.1mm of Polyethylene (PE). The overwrap film has a thickness of 0.14mm and oxygen transmission rate of 0.006 cm³ per 100 square inches per 24 hours per atmosphere at room temperature. A schematic depiction of the infusion bag is depicted in FIG. 16.

The overall oxygen permeability for each infusion bag from this present invention is 0.0025 cm³ per 24 hours per atmosphere at room temperature.

[000109] A stability study is performed on the above packaging material at 40 °C and 75% relative humidity. The results are depicted in the following table 6. The results demonstrate that the packaging material maintains a low level of met-hemoglobin over an extended period.

[000110] **Table 6**

Stability Study

Conditions: Temperature 40 +/- 2 °C, Relative humidity 75 +/- 5%

Package Material	Time Points (month)	Test Items			
		Total Hb	Met-Hb	Oxy-Hb	Endotoxin
		(g/dL)	(%)	(%)	(EU/mL)
Infusion Bag & Aluminum Overwrap	0	6.3	3.2	7.3	<0.02
	1	6.3	1.0	6.8	<0.02
	2	6.4	0.8	6.4	<0.02
	3	6.3	0.8	6.6	<0.02
	6	6.3	0.6	5.2	<0.02

[000111] **Example 8**

[000112] **Toxicity Study**

[000113] Experiments are carried out to evaluate the potential toxicity of the purified cross-linked tetrameric hemoglobin of the present invention. 10 male Sprague-Dawley rats are assigned to 3 groups, 3 in control group, 3 in low dose group and 4 in high dose group. Normal saline (control), purified cross-linked tetrameric hemoglobin at a low dose (5.52 g/kg), purified cross-linked tetrameric hemoglobin at high dose (6.90 g/kg) are administered respectively to the rats *via* continuous intravenous infusion *via* the jugular vein at an infusion rate of 3 mL/kg/hr. The drug treatment schedule is shown in the

following table 7. The cross-linked tetrameric hemoglobin is infused to the rats for up to 33.3 hours. The animals are closely observed for the duration of 9 days.

[000114] **Table 7**

Study Groups	Control (n=3)	Low Dose (n=3)	High Dose (n=4)
Day -7 to Day -6	Surgery on animals and then NS infusion for 24 hours		
Day -5 to Day -1	Daily injection of heparin		
Day 0	Randomization and Acclimatization, NS infusion for 24 hours		
Day 1	Metabolic Analysis (Water and Urine) collection. NS Infusion	Metabolic Analysis (Water and Urine) collection. 26.7 hours Hb infusion and then 21.3 hours NS infusion	Metabolic Analysis (Water and Urine) collection. 33.3 hours Hb infusion and then 14.7 hours NS infusion
Day 2	Metabolic Analysis (Water and Urine) collection		
Day 3	Metabolic Analysis (Water and Urine) collection		
Day 4	Stop NS infusion		
Day 5	Remove jacket and seal the tube, return animals to normal cages		
Day 8	Move Animal to metabolic cages for urine collection, and water consumed record		
Day 9	Metabolic Analysis (Water and Urine) collection Euthanasia of remaining animals Collection of blood (for both serum and plasma) Major organs preserved in formalin		

* NS: normal saline, Hb: cross-linked tetrameric hemoglobin

[000115] Blood samples are collected from all study animals for clinical pathology evaluations on Day 9. Urine is collected on Day 0 to Day 3 and Day 9 for urinalysis. Terminal necropsy is conducted on Day 9. Parameters evaluated include clinical observations, body weights, body weight changes, water consumption, clinical pathology (hematology, clinical chemistry and urinalysis), organ weights, organ-to-body weight ratio, organ-to-brain weight ratio, gross pathology and histopathology. Treatment with the cross-linked tetrameric hemoglobin of the present invention does not result in mortality or adverse clinical findings and no remarkable effects on body weight changes, and water consumption of rats.

[000116] There are no obvious changes observed on clinical chemistry and hematology parameters related to the treatment. Urine examination shows that the concentration of chloride and potassium is lower in treatment group than that of the control group on Day 2. Blood staining, higher protein concentration and more red blood cells in urine are found in treatment groups than control group on Day 2 to 3. Those observations all soon recover to normal. No significant difference is observed between lung-to-body weight ratio. During the study, no unexpected deaths and no obvious clinical signs of toxicity are observed. Additionally, gross pathological and histological analysis reveals no major abnormalities in all organs including lung, heart, liver, spleen and kidneys (such abnormalities are attributed to toxic side effects of infusion).

[000117] **Example 9**

[000118] **Heart preservation**

[000119] Cross-linked tetrameric hemoglobin can be used as a cardioplegic solution in a model of heart preservation during cardiopulmonary bypass in Beagle dogs. 18 Beagle dogs are randomly divided into 3 groups: sham, St. Thomas' solution (STS), and 0.1% of cross-linked tetrameric hemoglobin groups. Cardiopulmonary bypass is established in a standard fashion with cannulation of the ascending aorta, superior and inferior vena cava, and left ventricle for venting. Except for the sham group, STS without (STS group) or with 0.1 g/dL cross-linked tetrameric hemoglobin (0.1% Hb group) are infused into the aortic root after aortic clamping to achieve cardiac arrest that is maintained for 120 minutes. The cardiac function, including cardiac output, pulmonary artery pressure, pulmonary artery wedge pressure, mean artery pressure, central venous pressure and heart rate, and the blood gas are measured during reperfusion and compared to baseline. The release of cardiac enzymes, including creatine kinase MB, lactate dehydrogenase and Troponin-1 are also measured as surrogate markers of cardiac injury. Hematoxylin and eosin staining and cardiac water content detection are performed to determine the morphological and pathological changes of myocardium 120 minutes after reperfusion.

[000120] Under the basal condition, the measures of the cardiac function, the oxygen consumption and the release of cardiac enzymes are similar among the 3 groups. During reperfusion, the heart rate, cardiac output and central venous pressure are greatly decreased in the STS group when compared to the sham group. However, the heart rate, cardiac output, central venous pressure and cardiac oxygen consumption are greatly preserved in 0.1% Hb group, which is similar to those of the sham group. The cross-linked tetrameric hemoglobin in STS also greatly reduces the appearance of lactate dehydrogenase, creatine kinase MB and troponin-I when compared to the STS group.

Moreover, the cellular swelling, fatty changes and hyaline changes are significantly lessened in the 0.1% Hb group when compared to the STS group. The other measures, including pulmonary artery pressure, pulmonary artery wedge pressure, mean artery pressure and the cardiac water content, do not have significant differences among the 3 groups. All the measures in this study of the 0.1% Hb group have no significant difference as compared to the sham group. The 0.1 g/dL cross-linked tetrameric hemoglobin in STS during cardiopulmonary bypass exhibit a better cardiac protective effect than that of STS (a current standard cardioplegic solution) and the outcome is also comparable to the sham group.

[000121] **Example 10**

[000122] **Studies on Tissue Oxygenation, normal and cancerous tissues**

[000123] (10a) Improvement of oxygenation in normal tissue

[000124] Some studies for the normal tissue oxygenation by cross-linked tetrameric hemoglobin are carried out (as shown in FIG. 9). A comparative pharmacokinetic and pharmacodynamic study is conducted in buffalo rats. Male inbred buffalo rats are individually administered with 0.2 g/kg cross-linked tetrameric hemoglobin solution or ringer's acetate buffer (control group), through intravenous injection. The concentration-time profile of plasma hemoglobin is determined by HemocueTM photometer at 1, 6, 24, 48 hours and compared with the baseline reading. The methods are based on photometric measurement of hemoglobin where the concentration of hemoglobin is directly read out as g/dL. Oxygen partial pressure (pO₂) in the hind leg muscle of buffalo rats is directly measured by the OxylabTM tissue oxygenation and temperature monitor (Oxford Optronix

Limited). Rats are anesthetized by intra-peritoneal injection of 30-50 mg/kg pentobarbitone solution followed by insertion of the oxygen sensor into the muscle. All pO₂ readings are recorded by Datatrax2 data acquisition system (World Precision Instrument) in a real-time manner.

[000125] As seen in FIG. 9, injection of 0.2 g/kg cross-linked tetrameric hemoglobin solution demonstrate a correlation between the pharmacokinetic (plasma hemoglobin concentration) and pharmacodynamic (delivery of oxygen to muscular tissue) properties of the inventive cross-linked tetrameric hemoglobin solution. Importantly, a significant increase in oxygenation is observed for a longer period of time compared with the plasma hemoglobin level. Plasma hemoglobin concentration is seen in graph (A) and oxygen delivery to muscle is seen in graph (B).

[000126] (10b) Improvement of oxygenation in extremely hypoxic tumor area

[000127] Improvement of oxygenation in an extremely hypoxic tumor area is evaluated by a human nasopharyngeal carcinoma (CNE2) xenograft model. The CNE2 cell line is obtained from the Laboratory of Cancer Genetics, University of Hong Kong. Approximately 1×10^6 cancer cells are injected subcutaneously into 4- to 6-week-old inbred BALB/c AnN-nu (nude) mice. When the tumor xenograft reaches a diameter of 8-10 mm, oxygen partial pressure within the tumor mass is directly monitored by the Oxylab™ tissue oxygenation and temperature monitor (Oxford Optronix Limited). Oxygen partial pressure is measured along a tissue track with a fully computerized PTS30 Micro-positioning system (Discovery Technology International). All pO₂ readings are recorded by the Datatrax2 data acquisition system (World Precision Instrument) in a real-time manner. When the pO₂ reading is stabilized, 1.2 g/kg the

inventive cross-linked tetrameric hemoglobin solution is injected intravenously through the tail vein of the mice and the tissue oxygenation is measured. Results demonstrate a significant increase of oxygenation in the most hypoxic tumor area. After intravenous injection of 1.2 g/kg of cross-linked tetrameric hemoglobin, the median pO₂ value rise from 0.2 mmHg to 3.9 mmHg (in 3 hours) and 10.6 mmHg (in 6 hours), respectively (shown in FIG. 10).

[000128] **Example 11**

[000129] **Cancer treatment study (Chemo-sensitization effects of cross-linked tetrameric hemoglobin)**

[000130] The chemo-sensitization effects of the inventive cross-linked tetrameric hemoglobin are evaluated in different cancer cell lines. Leukemia cell line (Jurkat), colon cancer cell line (COLO205), cisplatin-resistant lung cancer cell line (A549/Cisp) and adriamycin-resistant breast cancer cell line (MCF-7/ADM) are obtained from Chinese Academy of Medical Sciences Cancer Institute. 8000 Jurkat cells, 4000 COLO205 cells, 3000 A549/Cisp cells and 3000 MCF-7/ADM cells are seeded individually in triplicate, onto a 96-well plate. After attachment, cells are incubated at 37°C with various chemotherapeutic agents alone, or in conjunction with 0.5 mg/mL cross-linked tetrameric hemoglobin solution. Jurkat cells are treated with vincristine sulfate at 0.31, 0.63, 1.25, 2.5, 5 and 10 µg/mL; COLO205 cells are treated with 5-fluorouracil at 0.78, 1.56, 3.13, 6.25 and 12.5 µg/mL; A549/Cisp cells are treated with cisplatin at 0.39, 0.78, 1.56, 3.13, 6.25 and 12.5 µg/mL and MCF-7/ADM cells are treated with adriamycin at 0.39, 0.78, 1.56, 3.13, 6.25 and 12.5 µg/mL. After treatment,

inhibition of cancer cell growth is determined by the ATP tumor chemo-sensitivity assay (ATP-TCA).

[000131] Esophageal cancer cell line HKESC-1 is obtained from Laboratory of Cancer Genetics, the University of Hong Kong. 2000 cancer cells are seeded onto a 96-well plate. After attachment, cells are incubated at 37°C with cisplatin alone, at 0.08, 0.4, 2, 10 and 50 µg/mL, or in conjunction with 3 mg/mL cross-linked tetrameric hemoglobin solution. After incubation, cytotoxicity is evaluated by MTT cell proliferation assay. Results show that addition of the inventive cross-linked tetrameric hemoglobin significantly enhances chemo-sensitivity in various cancer cell lines including A549/Cisp and MCF-7/ADM cells that are highly resistant to chemotherapy (shown in FIG. 8).

[000132] **Example 12**

[000133] **Treatment of Acute Severe Hemorrhagic Shock**

[000134] (12a) Treatment of Acute Severe Hemorrhagic Shock in Beagle dogs

[000135] The inventive cross-linked tetrameric hemoglobin is used as a resuscitation agent in a model of Acute Severe Hemorrhagic Shock in Beagle dogs. 60 Beagle dogs are randomly divided into 4 groups according to resuscitation agents, 15 dogs in each group.

[000136] Group 1: Dextran (Negative Control)

[000137] Group 2: Animal autologous blood (Positive Control)

[000138] Group 3: Low Dose Treatment (0.35 g cross-linked tetrameric hemoglobin/kg of body weight)

[000139] Group 4: Mid Dose Treatment (1.05 g cross-linked tetrameric hemoglobin/kg of body weight)

[000140] Acute severe hemorrhagic shock is established by withdrawing animal whole blood at a volume of 50 mL/kg body weight. Ten minutes after hemorrhagic shock established, Dextran (50 mL/kg), animal autologous blood (50 mL/kg), different doses of cross-linked tetrameric hemoglobin (5 mL/kg, 15 mL/kg) are infused into the animals. The infusion rate of cross-linked tetrameric hemoglobin is set at 10 mL/kg/h, hereafter, all experimental animals are observed for 7 days. A panel of parameters is observed and analyzed during the study period including survival, body weight, electro-cardiography (ECG), blood pressure, heart rate, respiration rate, body temperature, plasma hemoglobin concentration, hematology, arterial blood gas, urinalysis, clinical chemistry, coagulation, physical conditions and adverse events. Among all, survival is the primary end point. After seven days of treatment, the cross-linked tetrameric hemoglobin treatment group has a much higher survival rate compared with the normal group and the autologous blood group (as shown in the following table 8).

[000141] **Table 8**

[000142] **Canine study in hemorrhagic shock model**

Group	Survival no. after 7 days (n=15)	Survival rate after 7 days (%)
Dextran Negative Control	7	46
Dog's Autologous Blood	12	80
Low Dose Treatment	14	97
Mid Dose Treatment	15	100

[000143] (12b) Treatment of Acute Severe Hemorrhagic Shock in Rats

[000144] The inventive cross-linked tetrameric hemoglobin is also used as a resuscitation agent in a model of Acute Severe Hemorrhagic Shock in rats. 80 Sprague-Dawley rats are randomly divided into 5 groups according to resuscitation agents, 16 rats in each group.

[000145] Group 1: Lactate Ringer's solution (Negative Control)

[000146] Group 2: Animal autologous blood (Positive Control)

[000147] Group 3: Low Dose Treatment (0.1g cross-linked tetrameric hemoglobin/kg of body weight)

[000148] Group 4: Mid Dose Treatment (0.3g cross-linked tetrameric hemoglobin/kg of body weight)

[000149] Group 5: High Dose Treatment (0.5 g cross-linked tetrameric hemoglobin/kg of body weight)

[000150] Acute severe hemorrhagic shock is established by withdrawing 50% of animal whole blood, which is estimated as 35 mL/kg of body weight. Ten minutes after hemorrhagic shock established, Lactate Ringer's solution, animal autologous blood, different doses of cross-linked tetrameric hemoglobin (0.1 g Hb/kg, 0.3 g Hb/kg, 0.5 g Hb/kg) are infused into the animals. The infusion rate of cross-linked tetrameric hemoglobin is set at 5 mL/h, hereafter, all experimental animals are observed for 24 hours. A panel of parameters is observed and analyzed during the study period including survival, hemodynamics, myocardial mechanics, cardiac output, cardiac function, blood gas, tissue oxygen delivery & consumption, tissue perfusion & oxygen tension (liver, kidney and brain), liver & renal function, hemorheology (blood viscosity), mitochondrial respiratory control rate (liver, kidney and brain). Among all, survival is the primary end point. After 24 hours of observation, the inventive cross-linked tetrameric hemoglobin treatment group has much higher survival rate compared with the normal group and the autologous blood group (shown in the following table 9).

[000151] **Table 9**

Groups	Survival no. after 24-hour (n=16)	24-hour survival rate (%)
Negative control	3	18.75
Low Dose Treatment (0.1 g Hb/kg)	6	37.5
Mid Dose Treatment (0.3 g Hb/kg)	8	50
High Dose Treatment (0.5 g Hb/kg)	12	75
Rat's Autologous Blood	10	62.5

*Hb: cross-linked tetrameric hemoglobin

[000152] **Example 13: An *In vitro* met-hemoglobin formation study**

[000153] The prior art has stated that high molecular weight polymerized hemoglobin solution is preferable because it results in longer persistence in circulation. Therefore, the stabilized cross-linked tetrameric hemoglobin of the present invention is compared to high molecular weight polymerized hemoglobin *in vitro* to analyze stability in conditions that simulate circulation conditions. The test circuit is depicted in FIG. 17. In the met-hemoglobin formation by-pass circuit, A is the sample reservoir, B is the sample outlet port, C is the pump, D is Liqui-Cel contactor, E is a sample collection point and F is a sample inlet port. Time course met-hemoglobin formation of the cross-linked tetrameric hemoglobin solution of the present invention is compared with a commercially available product (Oxyglobin®), which contains approximately 68% of polymeric fraction ($\geq 128,000$ MW). Before the experiment, all samples are diluted to 5 g/dL and 50 mL of diluted sample is introduced into the sample reservoir from the sample inlet port. The

liquid pump rate is set at 30 mL/min, and the sample is allowed to fill the circuit. Throughout the experiment, the temperature of the sample reservoir is maintained at 37°C. To measure the met-hemoglobin level, 0.2 mL of test sample is collected from the sample collection point for co-oximetry (IL-682, Instrumentation Laboratory). Compressed air is then passed through the liqui-cel membrane contactor at flow rate 2.0 mL/ min to start oxygenation of the samples. A 0.2 mL sample is collected at 30-minute intervals. After treatment for 5 hours, the met-hemoglobin fractions of the inventive cross-linked tetrameric hemoglobin solution and the commercial available product (Oxyglobin[®]) increase to 8.7% and 16.4%, respectively. This demonstrates that the inactive met-hemoglobin formation rate for polymeric hemoglobin is substantially greater than that for the inventive cross-linked tetrameric hemoglobin (FIG. 18).

[000154] While the foregoing invention has been described with respect to various embodiments, such embodiments are not limiting. Numerous variations and modifications would be understood by those of ordinary skill in the art. Such variations and modifications are considered to be included within the scope of the following claims.

What is claimed:

1. A method for the preparation of a highly purified and high-temperature stable oxygen-carrier-containing pharmaceutical composition, the oxygen-carrier-containing pharmaceutical composition including hemoglobin, the method comprising:

a) providing mammalian whole blood including at least red blood cells and plasma;

b) separating the red blood cells from the plasma in the mammalian whole blood;

c) filtering the red blood cells that were separated from the plasma to obtain a filtered red blood cell fraction;

d) washing the filtered red blood cell fraction to remove plasma protein impurities, resulting in washed red blood cells;

e) disrupting the washed red blood cells by a controlled hypotonic lysis for a time sufficient to lyse red blood cells without lysing white blood cells in an instant cytolysis apparatus at a flow rate of at 50-1000 liter/hr to create a solution comprising a lysate of disrupted red blood cells;

f) performing filtration to remove at least a portion of the waste retentate from the lysate;

g) extracting a first hemoglobin solution from the lysate;

h) performing a first ultrafiltration process using an ultrafiltration filter configured to remove impurities having a higher molecular weight than tetrameric hemoglobin to further remove any viruses and residual waste retentate from the first hemoglobin solution to obtain a second hemoglobin solution;

i) performing flowthrough column chromatography on the second hemoglobin solution to remove phospholipids, protein impurities and dimeric hemoglobin and to form a phospholipid-free, low protein impurities and low dimer hemoglobin solution;

j) performing a second ultrafiltration process on the phospholipid-free, low protein impurities and low dimer hemoglobin solution using a filter configured to remove impurities resulting in a concentrated purified phospholipid-free, low protein impurities and low dimer hemoglobin solution;

k) blocking the sulfhydryl groups of hemoglobin molecules in the concentrated purified phospholipid-free, low protein impurities and low dimer hemoglobin solution by a sulfhydryl reagent in a fully oxygenated environment such that the hemoglobin molecules each have at least one cysteine moiety including a thiol-protecting group such that the hemoglobin molecules are incapable of binding endothelium-derived relaxing factor at the cysteine site;

l) cross-linking the thiol-protected hemoglobin by bis-3,5-dibromosalicyl fumarate to form high-temperature stable cross-linked tetrameric hemoglobin without the formation of polymeric hemoglobin such that the molecular weight of the resultant cross-linked tetrameric hemoglobin is 60 – 70 kDa and consists essentially of nonpolymeric cross-linked tetrameric hemoglobin;

m) exchanging a suitable physiological buffer for the cross-linked tetrameric hemoglobin;

n) removing any residual non-cross-linked tetrameric hemoglobin and any residual chemicals by washing;

o) adding N-acetyl cysteine at a concentration of 0.2-0.4% to the cross-linked tetrameric hemoglobin to maintain a level of met-hemoglobin below 5%; and

p) adding the low dimer, phospholipid-free, thiol-protected high-temperature stable up to a temperature of 80° C cross-linked tetrameric hemoglobin free from polymeric hemoglobin to a pharmaceutically acceptable carrier.

2. The method for the preparation of the oxygen-carrier-containing pharmaceutical composition of claim 1 wherein the instant cytolysis apparatus includes a static mixer.

3. The method for the preparation of the oxygen-carrier-containing pharmaceutical composition of claim 1 wherein said cross-linked tetrameric hemoglobin is derived from bovine, porcine, canine or equine hemoglobin.

4. The method for the preparation of the oxygen-carrier-containing pharmaceutical composition of claim 1 wherein the column chromatography comprises one or more cation-exchange columns and anion-exchange columns.

5. The method for the preparation of the oxygen-carrier-containing pharmaceutical composition of claim 4 wherein the ion exchange columns are one or more DEAE column, CM column and/or hydroxyapatite column.

6. The method for the preparation of the oxygen-carrier-containing pharmaceutical composition of claim 1 wherein the hemoglobin cross-linking by bis-3,5-dibromosalicy fumarate includes at least α - α and/or β - β cross-linking.
7. The method for the preparation of an oxygen-carrier-containing pharmaceutical composition of claim 6 wherein the cross-linking conditions are selected such that β - β cross-linking is greater than 50% of the total cross-linking.
8. The method for the preparation of an oxygen-carrier-containing pharmaceutical composition of claim 6 wherein the cross-linking conditions are selected such that β - β cross-linking is greater than 60% of the total cross-linking.
9. The method for the preparation of an oxygen-carrier-containing pharmaceutical composition of claim 6 wherein the cross-linking conditions are selected such that β - β cross-linking is greater than 70% of the total cross-linking.
10. The method for the preparation of an oxygen-carrier-containing pharmaceutical composition of claim 1 wherein the pharmaceutically acceptable carrier is a physiological buffer or water.
11. The method for the preparation of an oxygen-carrier-containing pharmaceutical composition of claim 1 further comprising packaging the prepared composition in a

multilayer, flexible infusion package having an oxygen permeability of less than 0.0025 cm³ per 24 hours at ambient conditions.

12. A highly purified and high temperature stable oxygen carrier-containing pharmaceutical composition comprising hemoglobin, the hemoglobin consisting essentially of nonpolymeric cross-linked tetrameric hemoglobin formed by the process of claim 1.

13. A method of oxygenating tissue comprising providing the composition of claim 12 to the tissue either *in vivo* or *ex vivo*.

P2739PC01_SequenceListing.txt
SEQUENCE LISTING

<110> WONG, Bing Lou
KWOK, Sui Yi

<120> A METHOD FOR THE PREPARATION OF HIGH-TEMPERATURE STABLE
OXYGEN-CARRIER-CONTAINING PHARMACEUTICAL COMPOSITION AND THE USE

<130> P2739PC01

<140> PCT/US2011/032594

<141> 2011-04-15

<160> 9

<170> PatentIn version 3.5

<210> 1

<211> 141

<212> PRT

<213> Bovine

<400> 1

Val Leu Ser Ala Ala Asp Lys Gly Asn Val Lys Ala Ala Trp Gly Lys
1 5 10 15

Val Gly Gly His Ala Ala Glu Tyr Gly Ala Glu Ala Leu Glu Arg Met
20 25 30

Phe Leu Ser Phe Pro Thr Thr Lys Thr Tyr Phe Pro His Phe Asp Leu
35 40 45

Ser His Gly Ser Ala Gln Val Lys Gly His Gly Ala Lys Val Ala Ala
50 55 60

Ala Leu Thr Lys Ala Val Glu His Leu Asp Asp Leu Pro Gly Ala Leu
65 70 75 80

Ser Glu Leu Ser Asp Leu His Ala His Lys Leu Arg Val Asp Pro Val
85 90 95

Asn Phe Lys Leu Leu Ser His Ser Leu Leu Val Thr Leu Ala Ser His
100 105 110

Leu Pro Ser Asp Phe Thr Pro Ala Val His Ala Ser Leu Asp Lys Phe
115 120 125

Leu Ala Asn Val Ser Thr Val Leu Thr Ser Lys Tyr Arg
130 135 140

<210> 2

<211> 142

<212> PRT

<213> Homo sapiens

<400> 2

Met Val Leu Ser Pro Ala Asp Lys Thr Asn Val Lys Ala Ala Trp Gly

P2739PC01_SequenceListing.txt

```

1              5              10              15
Lys Val Gly Ala His Ala Gly Glu Tyr Gly Ala Glu Ala Leu Glu Arg
20              25              30
Met Phe Leu Ser Phe Pro Thr Thr Lys Thr Tyr Phe Pro His Phe Asp
35              40              45
Leu Ser His Gly Ser Ala Gln Val Lys Gly His Gly Lys Lys Val Ala
50              55              60
Asp Ala Leu Thr Asn Ala Val Ala His Val Asp Asp Met Pro Asn Ala
65              70              75              80
Leu Ser Ala Leu Ser Asp Leu His Ala His Lys Leu Arg Val Asp Pro
85              90              95
Val Asn Phe Lys Leu Leu Ser His Cys Leu Leu Val Thr Leu Ala Ala
100             105             110
His Leu Pro Ala Glu Phe Thr Pro Ala Val His Ala Ser Leu Asp Lys
115             120             125
Phe Leu Ala Ser Val Ser Thr Val Leu Thr Ser Lys Tyr Arg
130             135             140

<210> 3
<211> 141
<212> PRT
<213> Canine

<400> 3
Val Leu Ser Pro Ala Asp Lys Thr Asn Ile Lys Ser Thr Trp Asp Lys
1      5      10      15
Ile Gly Gly His Ala Gly Asp Tyr Gly Gly Glu Ala Leu Asp Arg Thr
20      25      30
Phe Gln Ser Phe Pro Thr Thr Lys Thr Tyr Phe Pro His Phe Asp Leu
35      40      45
Ser Pro Gly Ser Ala Gln Val Lys Ala His Gly Lys Lys Val Ala Asp
50      55      60
Ala Leu Thr Thr Ala Val Ala His Leu Asp Asp Leu Pro Gly Ala Leu
65      70      75      80
Ser Ala Leu Ser Asp Leu His Ala Tyr Lys Leu Arg Val Asp Pro Val
85      90      95
Asn Phe Lys Leu Leu Ser His Cys Leu Leu Val Thr Leu Ala Cys His
100     105     110

```

P2739PC01_SequenceListing.txt

His Pro Thr Glu Phe Thr Pro Ala Val His Ala Ser Leu Asp Lys Phe
115 120 125

Phe Ala Ala Val Ser Thr Val Leu Thr Ser Lys Tyr Arg
130 135 140

<210> 4
<211> 141
<212> PRT
<213> Porcine

<400> 4

Val Leu Ser Ala Ala Asp Lys Ala Asp Val Lys Ala Ala Tyr Gly Lys
1 5 10 15

Val Gly Ala His Ala Gly Glu Ala Gly Ala Glu Ala Leu Glu Arg Met
20 25 30

Phe Leu Gly Phe Thr Thr Thr Lys Thr Tyr Phe Pro His Phe Asp Leu
35 40 45

Ser His Gly Ser Asp Glu Val Lys Ala His Gly Glu Lys Val Ala Asp
50 55 60

Ala Leu Thr Lys Ala Val Gly His Leu Asp Asp Met Pro Gly Ala Leu
65 70 75 80

Ser Ala Leu Ser Asp Leu His Ala His Lys Leu Arg Val Asp Pro Val
85 90 95

Asp Phe Lys Leu Leu Ser His Cys Leu Leu Ser Thr Leu Ala Val His
100 105 110

Leu Pro Asp Asp Phe Thr Pro Ala Val His Ala Asp Leu Asp Lys Phe
115 120 125

Leu Ala Asp Val Ser Thr Val Leu Asp Ser Lys Tyr Arg
130 135 140

<210> 5
<211> 142
<212> PRT
<213> Equine

<400> 5

Met Val Leu Ser Ala Ala Asp Lys Thr Asn Val Lys Ala Ala Trp Ser
1 5 10 15

Lys Val Gly Gly His Ala Gly Glu Phe Gly Ala Glu Ala Leu Glu Arg
20 25 30

P2739PC01_SequenceListing.txt

Met Phe Leu Gly Phe Pro Thr Thr Lys Thr Tyr Phe Pro His Phe Asp
35 40 45

Leu Ser His Gly Ser Ala Gln Val Lys Ala His Gly Lys Lys Val Gly
50 55 60

Asp Ala Leu Thr Leu Ala Val Gly His Leu Asp Asp Leu Pro Gly Ala
65 70 75 80

Leu Ser Asn Leu Ser Asp Leu His Ala His Lys Leu Arg Val Asp Pro
85 90 95

Val Asn Phe Lys Leu Leu Ser His Cys Leu Leu Ser Thr Leu Ala Val
100 105 110

His Leu Pro Asn Asp Phe Thr Pro Ala Val His Ala Ser Leu Asp Lys
115 120 125

Phe Leu Ser Ser Val Ser Thr Val Leu Thr Ser Lys Tyr Arg
130 135 140

<210> 6
<211> 145
<212> PRT
<213> Bovine

<400> 6

Met Leu Thr Ala Glu Glu Lys Ala Ala Val Thr Ala Phe Trp Gly Lys
1 5 10 15

Val Lys Val Asp Glu Val Gly Gly Glu Ala Leu Gly Arg Leu Leu Val
20 25 30

Val Tyr Pro Trp Thr Gln Arg Phe Phe Glu Ser Phe Gly Asp Leu Ser
35 40 45

Thr Ala Asp Ala Val Met Asn Asn Pro Lys Val Lys Ala His Gly Lys
50 55 60

Lys Val Leu Asp Ser Phe Ser Asn Gly Met Lys His Leu Asp Asp Leu
65 70 75 80

Lys Gly Thr Phe Ala Ala Leu Ser Glu Leu His Cys Asp Lys Leu His
85 90 95

Val Asp Pro Glu Asn Phe Lys Leu Leu Gly Asn Val Leu Val Val Val
100 105 110

Leu Ala Arg Asn Phe Gly Lys Glu Phe Thr Pro Val Leu Gln Ala Asp
115 120 125

Phe Gln Lys Val Val Ala Gly Val Ala Asn Ala Leu Ala His Arg Tyr

130

His
145

<210> 7
<211> 147
<212> PRT
<213> Homo sapiens

<400> 7

Met Val His Leu Thr Pro Glu Glu Lys Ser Ala Val Thr Ala Leu Trp
1 5 10 15

Gly Lys Val Asn Val Asp Glu Val Gly Gly Glu Ala Leu Gly Arg Leu
20 25 30

Leu Val Val Tyr Pro Trp Thr Gln Arg Phe Phe Glu Ser Phe Gly Asp
35 40 45

Leu Ser Thr Pro Asp Ala Val Met Gly Asn Pro Lys Val Lys Ala His
50 55 60

Gly Lys Lys Val Leu Gly Ala Phe Ser Asp Gly Leu Ala His Leu Asp
65 70 75 80

Asn Leu Lys Gly Thr Phe Ala Thr Leu Ser Glu Leu His Cys Asp Lys
85 90 95

Leu His Val Asp Pro Glu Asn Phe Arg Leu Leu Gly Asn Val Leu Val
100 105 110

Cys Val Leu Ala His His Phe Gly Lys Glu Phe Thr Pro Pro Val Gln
115 120 125

Ala Ala Tyr Gln Lys Val Val Ala Gly Val Ala Asn Ala Leu Ala His
130 135 140

Lys Tyr His
145

<210> 8
<211> 147
<212> PRT
<213> Porcine

<400> 8

Met Val His Leu Ser Ala Glu Glu Lys Glu Ala Val Leu Gly Leu Trp
1 5 10 15

Gly Lys Val Asn Val Asp Glu Val Gly Gly Glu Ala Leu Gly Arg Leu
20 25 30

P2739PC01_SequenceListing.txt

Leu Val Val Tyr Pro Trp Thr Gln Arg Phe Phe Glu Ser Phe Gly Asp
35 40 45

Leu Ser Asn Ala Asp Ala Val Met Gly Asn Pro Lys Val Lys Ala His
50 55 60

Gly Lys Lys Val Leu Gln Ser Phe Ser Asp Gly Leu Lys His Leu Asp
65 70 75 80

Asn Leu Lys Gly Thr Phe Ala Lys Leu Ser Glu Leu His Cys Asp Gln
85 90 95

Leu His Val Asp Pro Glu Asn Phe Arg Leu Leu Gly Asn Val Ile Val
100 105 110

Val Val Leu Ala Arg Arg Leu Gly His Asp Phe Asn Pro Asn Val Gln
115 120 125

Ala Ala Phe Gln Lys Val Val Ala Gly Val Ala Asn Ala Leu Ala His
130 135 140

Lys Tyr His
145

<210> 9
<211> 147
<212> PRT
<213> Equine

<400> 9

Met Val Gln Leu Ser Gly Glu Glu Lys Ala Ala Val Leu Ala Leu Trp
1 5 10 15

Asp Lys Val Asn Glu Glu Glu Val Gly Gly Glu Ala Leu Gly Arg Leu
20 25 30

Leu Val Val Tyr Pro Trp Thr Gln Arg Phe Phe Asp Ser Phe Gly Asp
35 40 45

Leu Ser Asn Pro Gly Ala Val Met Gly Asn Pro Lys Val Lys Ala His
50 55 60

Gly Lys Lys Val Leu His Ser Phe Gly Glu Gly Val His His Leu Asp
65 70 75 80

Asn Leu Lys Gly Thr Phe Ala Ala Leu Ser Glu Leu His Cys Asp Lys
85 90 95

Leu His Val Asp Pro Glu Asn Phe Arg Leu Leu Gly Asn Val Leu Val
100 105 110

P2739PC01_SequenceListing.txt

Val Val Leu Ala Arg His Phe Gly Lys Asp Phe Thr Pro Glu Leu Gln
115 120 125

Ala Ser Tyr Gln Lys Val Val Ala Gly Val Ala Asn Ala Leu Ala His
130 135 140

Lys Tyr His
145