

MACROCYCLIZATION REACTIONS AND INTERMEDIATES AND OTHER FRAGMENTS USEFUL IN THE SYNTHESIS OF HALICHONDRIN MACROLIDES


Background

5 The invention relates to intermediates useful in the synthesis of pharmaceutically active macrolide compounds and methods of synthesizing macrolide compounds. Halichondrin B is a potent anticancer agent originally isolated from the marine sponge *Halichondria okadai*, and subsequently found in *Axinella* sp., *Phakellia carteri*, and *Lissodendoryx* sp. A total synthesis of halichondrin B was published in 1992 (Aicher, T. D. et al., J. Am. Chem. Soc. 114:3162-3164). Further synthetic and structure-activity relationship studies
10 have been disclosed in U.S. Patent Nos. 5,338,865 and 5,436,238 and in Towle et al., *Annual Meeting of the American Association for Cancer Research*, April 6-10, 2002, 5721 and Wang et al., *Bioorg. Med. Chem. Lett.*, 10:1029-1032, 2000. Methods and intermediates for the synthesis of certain halichondrin B analogs and intermediates are described in International Publication Nos. WO 2005/118565, WO 2009/046308, WO 2009/064029, and WO 2009/124237; U.S. Patent No. 6,214,865; Austad et al., *Synlett* 24(3):333-337, 2013;
15 Austad et al., *Synlett*, 24(3):327-332, 2013; and Chase et al., *Synlett* 24(3):323-326, 2013. New methods for the synthesis of halichondrin B and its analogs (e.g., macrolide analogs) are desirable.

Summary

20 In general, the present disclosure provides methods for macrocyclization of intermediates in the synthesis of a halichondrin macrolide. The disclosure also provides intermediates that can be employed in the macrocyclization reactions described herein.


25 In one aspect, the disclosure provides a method of preparing a macrocyclic intermediate in the synthesis of a halichondrin macrolide, said method comprising performing a macrocyclization reaction on a non-macrocyclic intermediate, wherein (1) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.2-C.3 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with an organic base and a Lewis acid, wherein said non-macrocyclic intermediate is (1) a compound of formula (IA):



30 or a salt or a tautomer thereof,
wherein

each R is independently optionally substituted alkyl or optionally substituted aryl;

35 each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;


or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_V))-$, wherein P_V is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and


P_5 is H or a hydroxyl protecting group; and

wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IB):




or a salt or a tautomer thereof;

(2) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.2-C.3 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with an olefin metathesis catalyst, wherein said non-macrocyclic intermediate is a compound of formula (IIA):



or a salt or a tautomer thereof,
wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;


or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_V))- , wherein P_V is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

P₅ is H or a hydroxyl protecting group; and


wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IB):



(IB)


or a salt or a tautomer thereof;

5 (3) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.3-C.4 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with an olefin metathesis catalyst, wherein said non macrocyclic intermediate is a compound of formula (IIIA):



10 or a salt or a tautomer thereof,
wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

20 L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:

25



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;


or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_V))- , wherein P_V is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and


P₅ is H or a hydroxyl protecting group; and

wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IIIB):




or a salt or a tautomer thereof;

(4) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.12-C.13 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with an olefin metathesis catalyst, wherein said non-macrocyclic intermediate is a compound of formula (IVA):



or a salt thereof,
wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:





wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(i) R₃ is H or a hydroxyl protecting group, R₄ is alkyl ether, and R₅ is H;

(ii) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;


or

(iii) R₃ and R₄ combine to form a bond, and R₅ is H;

each P₄ and P₅ is independently H or a hydroxyl protecting group;


R₆ is H, and P₆ is H or a hydroxyl protecting group; or R₆ and P₆ combine to form a double bond; and wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of

formula (IVB):




or a salt thereof;

(5) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.15-C.16 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with an olefin metathesis catalyst, wherein said non-macrocyclic intermediate is a compound of formula (VA):



or a salt or a tautomer thereof,
wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:





wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:





, wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(a1) R₃ is H or a hydroxyl protecting group, R₄ is alkyl ether, and R₅ is H;

(a2) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

(a3) R₃ and R₄ combine to form a bond, and R₅ is H;

(b1) A₁ and R₇ combine to form oxo, R₆ is H or a hydroxyl protecting group, and R₈ is H;

or

(b2) A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₆ is H or a hydroxyl protecting group, and R₇ and R₈ combine to form a double bond;

or

(ii) R₆ and R₇ combine to form a bond, and R₈ is H or OP'';

(c1) R₉ is H, and P₄ is H or a hydroxyl protecting group;


or

(c2) R₉ and P₄ combine to form a double bond;

R₁₀ is H or -CH₂X₁CH₂CH=CH₂, wherein X₁ is O, -C(R₁₁)₂-, or NP₆, and wherein each R₁₁ is independently H or -COOR₁₂, P₆ is an N-protecting group, and R₁₂ is alkyl;


each P₅ is independently H or a hydroxyl protecting group; and

wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (VB):



or a salt or a tautomer thereof;

(6) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.19-C.20 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with a Cr(II) salt and a Ni(II) salt, wherein said non-macrocyclic intermediate is a compound of formula (VIA):




or a salt or a tautomer thereof,
wherein

Y is iodide, bromide, or trifluoromethanesulfonate;

b designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or *b* designates (*S*)-stereogenic center, and Z is OR₆, wherein R₆ is a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,

5 NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-

alkyl;

k is 0 or 1;

10 n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OPⁿ, where Pⁿ is H or a hydroxyl protecting group, and:


15 (i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OPⁿ;

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

20 wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IVB):



or a salt or a tautomer thereof,


25 wherein

b designates (R)-stereogenic center, c designates (S)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide;

or

30 b designates (S)-stereogenic center, c designates (R)-stereogenic center, and Z is OR₆, wherein R₆ is a hydroxyl protecting group; or

(7) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.26-C.27 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with a Cr(II) salt and a Ni(II) salt, wherein said non-macrocyclic intermediate is a compound of formula (VIIA):




(VIIA)

or a salt thereof,
wherein

Y is iodide, bromide, or trifluoromethanesulfonate;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(a1) R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$, wherein R₈ is H or a hydroxyl protecting group;

or

(a2) R₃ and R₄ combine to form a bond, R₅ is H, and

each P_4 is independently H or a hydroxyl protecting group, and X_1 , together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$;

or

both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal;


(b1) Z is chloride, bromide, or iodide, and R_6 and R_7 combine to form a bond;

or

(b2) Z and R_7 combine to form a double bond, and R_6 is a hydroxyl protecting group;

and


wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (VIIB):



or a salt thereof,

wherein each of P_5 and R_6 is independently H or a hydroxyl protecting group.

In another aspect, the disclosure provides a method of preparing:




halichondrin macrolide

or a salt thereof,

wherein

each of A_1 and A_2 is independently H or OP'' , where each P'' is independently H or a hydroxyl protecting group;


each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:



wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;
 R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo;
 or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$,

$NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, wherein each of R_A and


R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(1) said method comprising:

(A) producing a compound of formula (IB) or a salt or a tautomer thereof from a compound of formula (IA) or a salt or a tautomer thereof, the compound of formula (IA) having the following structure:



wherein

each R is independently optionally substituted alkyl or optionally substituted aryl;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;


or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and

P_5 is H or a hydroxyl protecting group; and

the compound of formula (IB) having the following structure:




and

(B) producing the halichondrin macrolide or the salt thereof from compound (IB) or the salt or the tautomer thereof;

(2) said method comprising:

(A) producing a compound of formula (IB) or the salt or the tautomer thereof from a compound of formula (IIA), the compound of formula (IIA) or the salt or the tautomer thereof having the following structure:



wherein

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or


A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_V))-$, wherein P_V is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and P_5 is H or a hydroxyl protecting group; and the compound of formula (IB) having the following structure:




and

(B) producing the halichondrin macrolide or the salt thereof from compound (IB) or the salt or the tautomer thereof;

(3) said method comprising:

(A) producing a compound of formula (IIIB) or the salt or the tautomer thereof from a compound of formula (IIIA) or the salt or the tautomer thereof, the compound of formula (IIIA) having the following structure:



wherein

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:


(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_V))-$, wherein P_V is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and P_5 is H or a hydroxyl protecting group; and

the compound of formula (IIIB) having the following structure:




and

(B) producing the halichondrin macrolide or the salt thereof from the compound of formula (IIIB) or the salt or the tautomer thereof;

(4) said method comprising:

(A) producing a compound of formula (IVB) or a salt thereof from a compound of formula (IVA) or a salt thereof, the compound of formula (IVA) having the following structure:



wherein

(i) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;

(ii) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;


or

(iii) R_3 and R_4 combine to form a bond, and R_5 is H;

each P_4 and P_5 is independently H or a hydroxyl protecting group;

R_6 is H, and P_6 is H or a hydroxyl protecting group; or R_6 and P_6 combine to form a double bond; and

the compound of formula (IVB) having the following structure:




and

(B) producing the halichondrin macrolide or the salt thereof from the compound of formula (IVB) or the salt thereof;


(5) said method comprising:

(A) producing a compound of formula (VB) or a salt or a tautomer thereof from a compound of formula (VA) or a salt or a tautomer thereof, the compound of formula (VA) having the following structure:




wherein

- (a1) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;
 (a2) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;
 or
 (a3) R_3 and R_4 combine to form a bond, and R_5 is H;
 (b1) A_1 and R_7 combine to form oxo, R_6 is H or a hydroxyl protecting group, and R_8 is H;
 or
 (b2) A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:
 (i) R_6 is H or a hydroxyl protecting group, and R_7 and R_8 combine to form a double bond;
 or
 (ii) R_6 and R_7 combine to form a bond, and R_8 is H or OP'' ;
 (c1) R_9 is H, and P_4 is H or a hydroxyl protecting group;
 or
 (c2) R_9 and P_4 combine to form a double bond;
 R_{10} is H or $-CH_2X_1CH_2CH=CH_2$, wherein X_1 is O, $-C(R_{11})_2-$, or NP_6 , and wherein each R_{11} is independently H or $-COOR_{12}$, P_6 is an *N*-protecting group, and R_{12} is alkyl;
 each P_5 is independently H or a hydroxyl protecting group; and
 the compound of formula (VB) having the following structure:




(B) producing a compound of formula (VC) or a salt thereof from the compound of formula (VB) or the salt or the tautomer thereof, the compound of formula (VC) having the following structure:



(VC),

wherein each of A_1 and A_2 is independently H or OP'' ;

- (C) producing a compound of formula (IVC) or a salt thereof from the compound of formula (VC) or a salt thereof, the compound of formula (IVC) having the following structure:




(IVC),

wherein each of A_1 and A_2 is independently H or OP'' ; and

- (D) producing the halichondrin macrolide or a salt thereof from the compound of formula (IVC) or a salt thereof;

(6) said method comprising:

- (A) producing a compound of formula (VIB) or a salt or a tautomer thereof from a compound of formula (VIA) or a salt or a tautomer thereof, the compound of formula (VIA) having the following structure:



(VIA),

wherein

Y is iodide, bromide, or trifluoromethanesulfonate;

- b designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or b designates (*S*)-stereogenic center, and Z is OR_6 , wherein R_6 is a hydroxyl protecting group;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H; or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

14g


- (i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

- (ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP*;

5

and each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; the compound of formula (VIB) having the following structure:



(VIB).

10

wherein

b designates (*R*)-stereogenic center, *c* designates (*S*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide;

or

b designates (S)-stereogenic center, *c* designates (R)-stereogenic center, and Z is OR₆, wherein R₆ is a hydroxyl protecting group;


15

(B) producing the halichondrin macrolide or the salt thereof from the compound of formula (VIB) or the salt or the tautomer thereof; or

(7) said method comprising:

20

(A) producing a compound of formula (VIIB) or the salt thereof from a compound of formula (VIIA) or the salt thereof, the compound of formula (VIIA) having the following structure:



(VIIA),

wherein

Y is iodide, bromide, or trifluoromethanesulfonate;

25

(a1) R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))_n-$;

or

(a2) R_3 and R_4 combine to form a bond, R_5 is H, and

30

each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))_n-$;

or

both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal;

35

14h

(b1) Z is chloride, bromide, or iodide, and R₆ and R₇ combine to form a bond;


or

(b2) Z and R₇ combine to form a double bond, and R₆ is a hydroxyl protecting group;

and


wherein R₈ is H or a hydroxyl protecting group

the compound of formula (VIIB) having the following structure:



wherein each of P₅ and R₆ is H or a hydroxyl protecting group;

(B) producing a compound of formula (VIIC) or a salt or a tautomer thereof from the compound of formula (VIIB) or the salt thereof, the compound of formula (VIIC) having the following structure:



wherein A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP^{''}, where P^{''} is H or a hydroxyl protecting group, and:


(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP”;

(C) producing a compound of formula (VIID) or a salt thereof from the compound of formula (VIIC) or the salt or the tautomer thereof, the compound of formula (VIID) having the following structure:

25



wherein each of A_1 and A_2 is independently H or OP'' ;


and

(D) producing the halichondrin macrolide or the salt thereof from the compound of formula (VIID) or the salt thereof.

In another aspect, the disclosure provides a method of preparing an intermediate in the synthesis of a halichondrin macrolide, said method comprising performing an allene-Prins reaction by contacting a compound of formula (VIIIA) or a salt thereof with a compound of formula (VIIIB) and R_4OH ,


wherein R_4 is an optionally substituted acyl;

wherein the compound of formula (VIIIA) has the following structure:



wherein

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;


k is 0 or 1;

n is 0, 1, or 2;


X is O, or X, together with the carbon atom to which it is attached, forms -(C(OP_Z)₂)-, wherein each P_Z is independently optionally substituted alkyl or optionally substituted aryl, or both P_Z combine to form optionally substituted alkylene or optionally substituted arylene; and

P₄ is H or a hydroxyl protecting group;

wherein the compound of formula (VIIB) has the following structure:




wherein



wherein P₅ is H or a hydroxyl protecting group; each P₆ is independently a hydroxyl protecting group, or both P₆ groups, together with the atoms to which each is attached, combine to form a cyclic protected diol; and R₅ is H or -

CH₂X₁CH₂CH=CH₂, wherein X₁ is O, -CH₂-, or NP₇, wherein P₇ is a sulfonyl; and wherein said intermediate is a compound of formula (VIIC):


14k



or a salt thereof.

In yet another aspect, the disclosure provides a compound of

5 (1) formula (IA) or formula (IB),




or a salt or a tautomer thereof,

wherein

each R is independently optionally substituted alkyl or optionally substituted aryl;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and

R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

5 n is 0, 1, or 2;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

10

or


(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both

15 P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and

P_5 is H or a hydroxyl protecting group;-

(2) formula (IIA), formula (IIIA), or formula (IIIB):




20

or a salt or a tautomer thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(\text{CH}(\text{OP}_2))-$, $-(\text{C}(\text{OH})(\text{OP}_2))-$, or $-\text{C}(\text{O})-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(\text{CH}_2)_n\text{OP}_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl,

arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

5 A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;


10 or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and

15 P_5 is H or a hydroxyl protecting group;

(3) formula (IVA), formula (IVB), or formula (IVC):




20

or a salt thereof,

wherein

each of A_1 and A_2 is H or OP'' , where P'' is H or a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl,

arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(i) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;

(ii) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;



or

(iii) R_3 and R_4 combine to form a bond, and R_5 is H;

each P_4 and P_5 is independently H or a hydroxyl protecting group; and

R_6 is H, and P_6 is H or a hydroxyl protecting group; or R_6 and P_6 combine to form a double bond;


(4) formula (VA), formula (VB), or formula (VC):



or a salt or a tautomer thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, wherein each P' is independently H or a hydroxyl protecting group;

A₂ is H or OP'';

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(a1) R₃ is H or a hydroxyl protecting group, R₄ is alkyl ether, and R₅ is H;

(a2) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;
or

(a3) R_3 and R_4 combine to form a bond, and R_5 is H;

(b1) A_1 and R_7 combine to form oxo, R_6 is H or a hydroxyl protecting group, and R_8 is H;
or

(b2) A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_6 is H or a hydroxyl protecting group, and R_7 and R_8 combine to form a double bond;

or

(ii) R_6 and R_7 combine to form a bond, and R_8 is H or OP'' ;

(c1) R_9 is H, and P_4 is H or a hydroxyl protecting group;

or


(c2) R_9 and P_4 combine to form a double bond;

R_{10} is H or $-CH_2X_1CH_2CH=CH_2$, wherein X_1 is O, $-C(R_{11})_2-$, or NP_6 , and wherein each R_{11} is

independently H or $-COOR_{12}$, P_6 is an *N*-protecting group, and R_{12} is alkyl;

each P_5 is independently H or a hydroxyl protecting group;

(5) formula (VIA):




(VIA)

or a salt or a tautomer thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



10 E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

Y is iodide, bromide, or trifluoromethanesulfonate;

20 b designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or b designates (*S*)-stereogenic center, and Z is OR_6 , wherein R_6 is a hydroxyl protecting group;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:


25 (i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal;

(6) formula (VIB):



or a salt or a tautomer thereof,


wherein

b designates (*R*)-stereogenic center, c designates (*S*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide;

or

b designates (*S*)-stereogenic center, c designates (*R*)-stereogenic center, and Z is OR_6 , wherein R_6 is a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:





wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo;

or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, wherein each P' is independently H or a hydroxyl protecting

group;

E is optionally substituted alkyl or optionally substituted alkoxy;

5 G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

10 A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:


(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

15 each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))-, wherein P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal;

(7) formula (VIIA) or formula (VIIB):




or a salt thereof,

wherein

- 5 each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:

10



wherein


L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;


R₁ is H, or R₁ and P₁ combine to form a bond;

15

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:

20





, wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

5 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

10 k is 0 or 1;

n is 0, 1, or 2;

(a1) R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or -(CH(OR₈))-;

15 or

(a2) R₃ and R₄ combine to form a bond, R₅ is H, and

each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or -(CH(OR₈))-;

or

20 both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal;

(b1) Z is chloride, bromide, or iodide, and R₆ and R₇ combine to form a bond;

(b2) Z and R₇ combine to form a double bond, and R₆ is a hydroxyl protecting group;


or

25 (b3) when Z and R₇ are absent, R₆ is H or a hydroxyl protecting group;

P₅ is H or a hydroxyl protecting group;

wherein R₈ is H or a hydroxyl protecting group; or


(8) formula (VIIC):



or a salt or a tautomer thereof,
wherein

- 5 each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:

10



wherein


L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;


R₁ is H, or R₁ and P₁ combine to form a bond;

15

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:

20





, wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

5 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

10 k is 0 or 1;

n is 0, 1, or 2;

each P₄ is H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))-, wherein P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form a ketal; and

15 A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or


A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

20 or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP''.


In yet another aspect, the disclosure provides a compound of formula (VIIC):



25

or a salt thereof,
wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo;

or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

20


each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl,

arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

5 P_4 is H or a hydroxyl protecting group; and



wherein P_5 is H or a hydroxyl protecting group; each P_6 is independently a hydroxyl protecting group, or both P_6 groups, together with the atoms to which each is attached, combine to form a cyclic protected diol; and R_5 is H or $-\text{CH}_2\text{X}_1\text{CH}_2\text{CH}=\text{CH}_2$, wherein X_1 is O, $-\text{CH}_2-$, or NP_7 , wherein P_7 is a sulfonyl.


10

As described herein, the disclosure provides a method of preparing a macrocyclic intermediate in the synthesis of a halichondrin macrolide, the method involving performing a macrocyclization reaction on a non-macrocyclic intermediate, the macrocyclization reaction producing the macrocyclic intermediate by forming C.2-C.3, C.3-C.4, C.12-C.13, C.15-C.16, C.19-C.20, or C.26-C.27 bond in the structure of the halichondrin macrolide.

15

In some examples, performing the macrocyclization reaction involves contacting the non-macrocyclic intermediate (e.g., a compound of formula (IA)) with an organic base (e.g., DBU or triethylamine) and a Lewis acid (e.g., a salt of Li or Zn). The non-macrocyclic intermediate can be a compound of formula (IA):

20



or a salt or a tautomer thereof,

where


each R is independently optionally substituted alkyl or optionally substituted aryl;

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6}

saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10

25

substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:





where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

- (i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;


or

- (ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and

P_5 is H or a hydroxyl protecting group; and

the macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IB):




or a salt or a tautomer thereof.

In formula (IA) or (IB), each R can be optionally substituted C_{1-6} alkyl. In formula (IA) or (IB), both P_4 groups and X_1 , together with the atoms to which each is attached, can combine to form ketal. In formula (IA) or (IB), X_1 , together with the carbon atom to which it is attached, can be $-(CH(OP_Y))-$, where P_Y is H or a hydroxyl protecting group. In formula (IA) or (IB), P_5 can be a hydroxyl protecting group. In formula (IA) or (IB), R_3 and R_4 can combine to form a bond, and R_5 can be H.

In particular examples, performing the macrocyclization reaction involves contacting the non-macrocyclic intermediate (e.g., a compound of formula (IIA), formula (IIIA), formula (IVA), or formula (VA)) with an olefin metathesis catalyst (e.g., a ruthenium-carbene complex).

The non-macrocyclic intermediate can be a compound of formula (IIA):




or a salt or a tautomer thereof,

where

- 5 each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:

10





where

L is -(CH(OP₂))-, -(C(OH)(OP₂))-, or -C(O)-;

R₁ is H, or R₁ and P₁ combine to form a bond;

15

R₂ is H or -(CH₂)_nOP₃, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

10 k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

15 (i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;


or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

20 each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))- , wherein P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and


P₅ is H or a hydroxyl protecting group; and

where the macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IB):




5 or a salt or a tautomer thereof.

The non-macrocyclic intermediate can be a compound of formula (IIIA):



10 or a salt or a tautomer thereof,
where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:



where


L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;


R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is

20

attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



5



, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$,

10 $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

15

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

20

or


(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and

25

P_5 is H or a hydroxyl protecting group; and


where the macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IIIB):



5 or a salt or a tautomer thereof.


In formula (IIA), (IIB), (IIIA), or (IIIB), both P_4 groups and X_1 , together with the atoms to which each is attached, can combine to form ketal. In formula (IIA), (IIB), (IIIA), or (IIIB), P_5 can be a hydroxyl protecting group. In formula (IIA), (IIB), (IIIA), or (IIIB), X_1 , together with the carbon atom to which it is attached, can be
 10 $-(CH(OP_Y))-$, where P_Y is H. In formula (IIA), (IIB), (IIIA), or (IIIB), X_1 can be oxo.

the non-macrocyclic intermediate is a compound of formula (IVA):



15 or a salt thereof,
 where

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10
 20 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




where

L is $-(\text{CH}(\text{OP}_2))-$, $-(\text{C}(\text{OH})(\text{OP}_2))-$, or $-\text{C}(\text{O})-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(\text{CH}_2)_n\text{OP}_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $\text{NR}_B(\text{CO})\text{R}_A$, $\text{NR}_B(\text{CO})(\text{CO})\text{R}_A$,

$\text{NR}_A(\text{CO})\text{NR}_B\text{R}_A$, $\text{NR}_B(\text{CO})\text{OR}_A$, $(\text{CO})\text{OR}_A$, $\text{O}(\text{CO})\text{R}_A$, $(\text{CO})\text{NR}_B\text{R}_A$, or $\text{O}(\text{CO})\text{NR}_B\text{R}_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(i) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;

(ii) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;


or

(iii) R_3 and R_4 combine to form a bond, and R_5 is H;

each P_4 and P_5 is independently H or a hydroxyl protecting group;

R_6 is H, and P_6 is H or a hydroxyl protecting group; or R_6 and P_6 combine to form a double bond; and


where the macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IVB):



5 or a salt thereof.

In formula (IVA) or (IVB), at least one of P_4 and P_5 can be H. In formula (IVA) or (IVB), R_6 and P_6 can combine to form a double bond.


10 The non-macrocyclic intermediate can be a compound of formula (VA):



or a salt or a tautomer thereof,

where

15 each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




20

where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

- 10 G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;
 each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;
- 15 k is 0 or 1;
 n is 0, 1, or 2;
- (a1) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;
 (a2) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;
 or
- 20 (a3) R_3 and R_4 combine to form a bond, and R_5 is H;
 (b1) A_1 and R_7 combine to form oxo, R_6 is H or a hydroxyl protecting group, and R_8 is H;
 or
- (b2) A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:
- (i) R_6 is H or a hydroxyl protecting group, and R_7 and R_8 combine to form a double bond;
- 25 or


(ii) R_6 and R_7 combine to form a bond, and R_8 is H or OP'' ;

(c1) R_9 is H, and P_4 is H or a hydroxyl protecting group;

or

(c2) R_9 and P_4 combine to form a double bond;


- 5 R_{10} is H or $-CH_2X_1CH_2CH=CH_2$, where X_1 is O, $-C(R_{11})_2-$, or NP_6 , and where each R_{11} is independently H or $-COOR_{12}$, P_6 is an *N*-protecting group, and R_{12} is alkyl;
- each P_5 is independently H or a hydroxyl protecting group; and
- where the macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (VB):



or a salt or a tautomer thereof.

- 15 In formula (VA) or (VB), R_9 can be H, and P_4 can be H. In formula (VA) or (VB), R_{10} can be $-CH_2X_1CH_2CH=CH_2$, and X_1 can be O. In formula (VA) or (VB), R_6 and R_7 can combine to form a bond, and R_8 can be H. In formula (VA) or (VB), each P_5 can be independently a hydroxyl protecting group. In formula (VA) or (VB), at least one P_5 can be H. In formula (VA) or (VB), R_3 and R_4 can combine to form a bond, and R_5 can be H.
- 20 In certain examples, performing the macrocyclization reaction involves contacting the non-macrocyclic intermediate (e.g., a compound of formula (VIA) or (VIIA)) with a Cr(II) salt and a Ni(II) salt.

The non-macrocyclic intermediate can be a compound of formula (VIA):




or a salt or a tautomer thereof,

where

5 Y is iodide, bromide, or trifluoromethanesulfonate;

b designates (*R*)-stereogenic center, and *Z* is a sulfonate, chloride, bromide, or iodide; or *b* designates (*S*)-stereogenic center, and *Z* is OR₆, where R₆ is a hydroxyl protecting group;

each of *D* and *D'* is independently H, optionally substituted alkyl, or OP₁, provided that only one of *D* and *D'* is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; *A* is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:





15 where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:





, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

5 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

10 n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

15


or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

20

where the macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IVB):



(IVB)

or a salt or a tautomer thereof,

where

25

b designates (*R*)-stereogenic center, *c* designates (*S*)-stereogenic center, and *Z* is a sulfonate, chloride, bromide, or iodide;

or


b designates (*S*)-stereogenic center, *c* designates (*R*)-stereogenic center, and *Z* is OR₆, where R₆ is a hydroxyl protecting group.

5

10

In formula (VIA) or (VIB), *Y* can be bromide. In formula (VIA) or (VIB), R₃ and R₄ can combine to form a bond, and R₅ can be H. In formula (VIA) or (VIB), both P₄ groups and X₁, together with the atoms to which each is attached, can combine to form ketal. In formula (VIA) or (VIB), *Z* can be a sulfonate. In formula (VIA) or (VIB), *Z* can be OR₆, where R₆ is a hydroxyl protecting group. In formula (VIA) or (VIB), *Z* can be an ester, carbonate, or carbamate.

The non-macrocyclic intermediate can be a compound of formula (VIIA):



15


or a salt thereof,

where

Y is iodide, bromide, or trifluoromethanesulfonate;

20

each of *D* and *D'* is independently H, optionally substituted alkyl, or OP₁, provided that only one of *D* and *D'* is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; *A* is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:



25

where



L is -(CH(OP₂))-, -(C(OH)(OP₂))-, or -C(O)-;

R₁ is H, or R₁ and P₁ combine to form a bond;

30

R₂ is H or -(CH₂)_nOP₃, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo;

or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting

group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(a1) R_3 is H or a hydroxyl protecting group, R_4 and R_5 combine to form a double bond, each P_4 is independently H or a hydroxyl protecting group, and X_1 , together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$, where R_8 is H or a hydroxyl protecting group;

or

(a2) R_3 and R_4 combine to form a bond, R_5 is H, and

each P_4 is independently H or a hydroxyl protecting group, and X_1 , together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$;

or

both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal;


(b1) Z is chloride, bromide, or iodide, and R_6 and R_7 combine to form a bond;

or

(b2) Z and R_7 combine to form a double bond, and R_6 is a hydroxyl protecting group;

and

where the macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (VIIB):




(VIIB)

or a salt thereof,

where each of P_5 and R_6 is independently H or a hydroxyl protecting group.

In formula (VIA), Z can be iodide. In formula (VIA) or (VIIB), Y can be trifluoromethanesulfonate. In formula (VIA) or (VIIB), R_3 can be H or a hydroxyl protecting group, R_4 and R_5 can combine to form a double bond, each P_4 can independently be H or a hydroxyl protecting group, and X_1 , together with the carbon to which it is attached, can form $-(CH(OR_8))-$, where R_8 is H or a hydroxyl protecting group. In formula (VIA) or (VIIB), at least one of P_3 , P_4 , and R_6 can be a hydroxyl protecting group. In formula (VIA), Z can be chloride, bromide, or iodide, and R_6 and R_7 combine to form a bond.

Further described herein is a method of preparing:



halichondrin macrolide


or a salt thereof,

where

each of A_1 and A_2 is independently H or OP'' , where each P'' is independently H or a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10

substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$,


$NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

the method involving:

(A) producing a compound of formula (IB) from a compound of formula (IA), the compound of formula (IA) having the following structure:



5 or a salt or a tautomer thereof,

wherein

each R is independently optionally substituted alkyl or optionally substituted aryl;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

10 A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;


or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

15 each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))-, wherein P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

P₅ is H or a hydroxyl protecting group; and

the compound of formula (IB) having the following structure:



20


or a salt or a tautomer thereof;

and

(B) producing the halichondrin macrolide from compound (IB).

In some examples, the producing the compound of formula (IB) involves reacting the compound of formula (IA) with with an organic base and a Lewis acid. In certain examples, the producing the halichondrin macrolide involves reacting the compound of formula (IB) with a hydroxyl protecting group removing agent. In particular examples, each R is optionally substituted alkyl.

Further described herein is a method of preparing:




halichondrin macrolide

or a salt thereof,
where

each of A_1 and A_2 is independently H or OP'' , where each P'' is independently H or a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:





where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;


each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

the method involving:

(A) producing a compound of formula (IB) from a compound of formula (IIA), the compound of formula (IIA) having the following structure:



or a salt or a tautomer thereof,

wherein

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;
or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or


(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both

P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and

P_5 is H or a hydroxyl protecting group; and

the compound of formula (IB) having the following structure:



(IB)


or a salt or a tautomer thereof;

and

(B) producing the halichondrin macrolide from compound (IB).

In some examples, the producing the compound of formula (IB) involves reacting the compound of formula (IIA) with an olefin metathesis catalyst (e.g., a ruthenium-carbene complex). In some examples, the producing the halichondrin macrolide involves reacting the compound of formula (IB) with a hydroxyl protecting group removing agent.

Further described herein is a method of preparing:




halichondrin macrolide

or a salt thereof,

wherein

each of A_1 and A_2 is independently H or OP'' , where each P'' is independently H or a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;


each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

the method involving:

(A) producing a compound of formula (IIIB) from a compound of formula (IIIA), the compound of formula (IIIA) having the following structure:



or a salt or a tautomer thereof,

wherein

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;


or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and

P_5 is H or a hydroxyl protecting group; and

the compound of formula (IIIB) having the following structure:




or a salt or a tautomer thereof;

and

(B) producing the halichondrin macrolide from the compound of formula (IIIB).

In some examples, the producing the compound of formula (IIIB) involves reacting the compound of formula (IIIA) with an olefin metathesis catalyst. In certain examples, the producing the halichondrin macrolide involves reacting the compound of formula (IIIB) with a hydroxyl protecting group removing agent. In particular examples, both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal. In further examples, P₅ is a hydroxyl protecting group.

Further described herein is a method of preparing:



halichondrin macrolide


or a salt thereof,

where

each of A₁ and A₂ is independently H or OPⁿ, where each Pⁿ is independently H or a hydroxyl

protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;


each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

the method involving:

(A) producing a compound of formula (IVB) from a compound of formula (IVA), the compound of formula (IVA) having the following structure:




or a salt thereof,

where

- (i) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;
 (ii) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;
 or
 (iii) R_3 and R_4 combine to form a bond, and R_5 is H;

each P_4 and P_5 is independently H or a hydroxyl protecting group;

R_6 is H, and P_6 is H or a hydroxyl protecting group; or R_6 and P_6 combine to form a double bond; and
 the compound of formula (IVB) having the following structure:




or a salt thereof;

and

(B) producing the halichondrin macrolide from the compound of formula (IVB).

In some examples, the producing the compound of formula (IVB) involves reacting the compound of formula (IVA) with an olefin metathesis catalyst. In certain examples, the producing the halichondrin macrolide from the compound of formula (IVB) involves reacting the compound of formula (IVB) with a Brønsted acid. In other examples, the producing the halichondrin macrolide involves producing the compound of formula (IVC), the compound of formula (IVC) having the following structure:

57a



or a salt thereof;


and producing the halichondrin macrolide from the compound of formula (IVC).

5

In yet other examples, the producing the compound of formula (IVC) involves reacting the compound of formula (IVB) with a hydroxyl protecting group removing agent. In still other examples, the producing the halichondrin macrolide from the compound of formula (IVC) involves reacting the compound of formula (IVC) with a Brønsted acid. In particular examples, at least one of P₄ and P₅ is H.

10

Further described herein is a method of preparing:



halichondrin macrolide

or a salt thereof,


15

where

each of A₁ and A₂ is independently H or OPⁿ, where each Pⁿ is independently H or a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:

20





25

where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$,


15 $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

20 the method involving:

(A) producing a compound of formula (VB) from a compound of formula (VA), the compound of formula (VA) having the following structure:



or a salt or a tautomer thereof,

where

(a1) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;
 (a2) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

(a3) R_3 and R_4 combine to form a bond, and R_5 is H;
 (b1) A_1 and R_7 combine to form oxo, R_6 is H or a hydroxyl protecting group, and R_8 is H;

or

(b2) A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:
 (i) R_6 is H or a hydroxyl protecting group, and R_7 and R_8 combine to form a double bond;

or

(ii) R_6 and R_7 combine to form a bond, and R_8 is H or OP'' ;

(c1) R_9 is H, and P_4 is H or a hydroxyl protecting group;

or


(c2) R_9 and P_4 combine to form a double bond;

R_{10} is H or $-CH_2X_1CH_2CH=CH_2$, where X_1 is O, $-C(R_{11})_2-$, or NP_6 , and where each R_{11} is

independently H or $-COOR_{12}$, P_6 is an *N*-protecting group, and R_{12} is alkyl;


each P_5 is independently H or a hydroxyl protecting group; and

where the macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (VB):



or a salt or a tautomer thereof;


(B) producing a compound of formula (VC) from the compound of formula (VB), the compound of formula (VC) having the following structure:



or a salt thereof,

where each of A_1 and A_2 is independently H or OP'' ;

(C) producing a compound of formula (IVC) from the compound of formula (VC), the compound of formula (IVC) having the following structure:




or a salt thereof,

where each of A_1 and A_2 is independently H or OP'' ; and

(D) producing the halichondrin macrolide from the compound of formula (IVC).

In some examples, the producing the compound of formula (VB) involves contacting the compound of formula (VA) with an olefin metathesis catalyst. In certain examples, P_4 is H, R_9 is H, and the producing the compound of formula (VC) involves oxidizing the compound of formula (VB) to produce a compound of formula (VBa):



or a salt thereof;


where each of A₁ and A₂ is independently H or OP''

5 and reacting the compound of formula (VBa) with a 1,4-reducing agent to produce the compound of formula (VC).

In particular examples, the producing the compound of formula (IVC) involves contacting the compound of formula (VC) with a hydroxyl protecting group removing agent. In further examples, the producing the

10 halichondrin macrolide involves contacting the compound of formula (IVC) with a Brønsted acid.

Further described herein is a method of preparing:



halichondrin macrolide


15 or a salt thereof,

where

each of A₁ and A₂ is independently H or OP'', where each P'' is independently H or a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:

20




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$,

$NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;


k is 0 or 1;

n is 0, 1, or 2;

the method involving:

(A) producing a compound of formula (VIB) from a compound of formula (VIA), the compound of formula (VIA) having the following structure:

57g



or a salt or a tautomer thereof,

where

Y is iodide, bromide, or trifluoromethanesulfonate;

b designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or *b* designates (*S*)-stereogenic center, and Z is OR₆, where R₆ is a hydroxyl protecting group;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OPⁿ, where Pⁿ is H or a hydroxyl protecting group, and:


(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OPⁿ;

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal;

the compound of formula (VIB) having the following structure:



or a salt or a tautomer thereof,

where

b designates (*R*)-stereogenic center, *c* designates (*S*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide;

or


b designates (*S*)-stereogenic center, *c* designates (*R*)-stereogenic center, and *Z* is OR₆, where R₆ is a hydroxyl protecting group;

(B) producing the halichondrin macrolide from the compound of formula (VIB).

- 5 In some examples, the producing the compound of formula (VIB) involves reacting the compound of formula (VIA) with a Cr(II) salt and a Ni(II) salt. In particular examples, the producing the halichondrin macrolide involves the step of nucleophilic ring-closing of the compound of formula (VIB). In certain examples, both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal. In further examples, R₃ and R₄ combine to form a bond, and R₅ is H.

10

Further described herein is a method of preparing:



halichondrin macrolide

or a salt thereof,


15

where

each of A₁ and A₂ is independently H or OP'', where each P'' is independently H or a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:

20



25

where



L is -(CH(OP₂))- , -(C(OH)(OP₂))- , or -C(O)-;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or -(CH₂)_nOP₃, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is

30

attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,


NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

10 k is 0 or 1;

n is 0, 1, or 2;

the method involving:

(A) producing a compound of formula (VIIB) from a compound of formula (VIIA), the compound of formula (VIIA) having the following structure:



15

or a salt thereof,

where

Y is iodide, bromide, or trifluoromethanesulfonate;

(a1) R_3 is H or a hydroxyl protecting group, R_4 and R_5 combine to form a double bond, each P_4 is independently H or a hydroxyl protecting group, and X_1 , together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$;

or

(a2) R_3 and R_4 combine to form a bond, R_5 is H, and each P_4 is independently H or a hydroxyl protecting group, and X_1 , together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$;

or

both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal;

(b1) Z is chloride, bromide, or iodide, and R_6 and R_7 combine to form a bond;


or

(b2) Z and R_7 combine to form a double bond, and R_6 is a hydroxyl protecting group;

where R_8 is H or a hydroxyl protecting group;

and


the compound of formula (VIIB) having the following structure:



or a salt thereof,

where each of P_5 and R_6 is independently H or a hydroxyl protecting group;

(B) producing a compound of formula (VIIC) from the compound of formula (VIIB), the compound of formula (VIIC) having the following structure:



or a salt or a tautomer thereof,

where

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or


A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

(C) producing a compound of formula (VIID) from the compound of formula (VIIC), the compound of formula (VIID) having the following structure:



or a salt thereof,

where each of A_1 and A_2 is independently H or OP'' ;

and

(D) producing the halichondrin macrolide from the compound of formula (VIID).

In some examples, the producing the compound of formula (VIIB) involves reacting the compound of formula (VIIA) with a Cr(II) salt and a Ni(II) salt. In certain examples, the producing the compound of formula (VIIC) involves the step of nucleophilic ring-closing of the compound of formula (VIIB). In particular examples, the producing the compound of formula (VIID) involves reacting the compound of formula (VIIC) with a hydroxyl protecting group removing agent. In further examples, the producing the halichondrin macrolide involves reacting the compound of formula (VIID) with a Brønsted acid. In other examples, Z is iodide. In yet other examples, Y is trifluoromethanesulfonate. In other examples, R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond. In some examples, each P_4 is independently H or a hydroxyl protecting group, and X_1 , together with the carbon to which it is attached, forms $-(CH(OR_8))-$, where R_8 is H or a hydroxyl protecting group. In particular examples, P_5 is H. In certain examples, Z is chloride, bromide, or iodide, and R_6 and R_7 combine to form a bond.

In some examples, a designates (S)-stereogenic center. In certain examples, one and only one of D and D' is optionally substituted alkyl or OP_1 . In particular examples, one and only one of D and D' is OP_1 , where P_1


is H, alkyl, or a hydroxyl protecting group. In further examples, A is a group of formula (1). In certain examples, L is $-(CH(OP_2))-$. In other examples, R_1 and P_1 combine to form a bond. In yet other examples, G is O. In still other examples, E is optionally substituted alkyl. In some examples, k is 1. In certain examples, R_2 is $-(CH_2)_nOP_3$. In particular examples, at least one of P_2 and P_3 is a hydroxyl protecting group.

5 In certain examples, A_1 is H. In some examples, A_2 is H.

Further described herein is a method of preparing an intermediate in the synthesis of a halichondrin macrolide. The method involves performing an allene-Prins reaction by contacting a compound of formula (VIII A) with a compound of formula (VIII B) and R_4OH ,

10 where R_4 is an optionally substituted acyl;

where the compound of formula (VIII A) has the following structure:




or a salt thereof,

15 where

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group

20 of formula (1) having the structure:





where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

25 R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from

30 the group consisting of:



, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;


10 k is 0 or 1;

n is 0, 1, or 2;


X is O, or X, together with the carbon atom to which it is attached, forms -(C(OP_Z)₂)-, wherein each P_Z is independently optionally substituted alkyl or optionally substituted aryl, or both P_Z combine to form optionally substituted alkylene or optionally substituted arylene; and

15 P₄ is H or a hydroxyl protecting group;

where the compound of formula (VIIIB) has the following structure:




where



where P_5 is H or a hydroxyl protecting group; each P_6 is independently a hydroxyl protecting group, or both P_6 groups, together with the atoms to which each is attached, combine to form a cyclic protected diol; and R_5 is H or $-\text{CH}_2\text{X}_1\text{CH}_2\text{CH}=\text{CH}_2$, where X_1 is O, $-\text{CH}_2-$, or NP_7 , where P_7 is a sulfonyl;

and where the intermediate is a compound of formula (VIIC):




or a salt thereof,

where a designates (R)-stereogenic center or (S)-stereogenic center.

In some examples, the performing a Prins reaction involves reacting the compound of formula (VIII A) with a Lewis acid. In certain examples, k is 1.

Further described herein is a compound of formula (IA) or formula (IB),




or a salt or a tautomer thereof,

where

each R is independently optionally substituted alkyl or optionally substituted aryl;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




where

L is $-(\text{CH}(\text{OP}_2))-$, $-(\text{C}(\text{OH})(\text{OP}_2))-$, or $-\text{C}(\text{O})-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(\text{CH}_2)_n\text{OP}_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;
or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:


(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and P_5 is H or a hydroxyl protecting group.

Further described herein is a compound of formula (IIA), formula (IIIA), or formula (IIIB):




or a salt or a tautomer thereof,

where

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10

substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$,

$NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;


or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and

P_5 is H or a hydroxyl protecting group.

Further described herein is a compound of formula (IVA), formula (IVB), or formula (IVC):




or a salt thereof,

where

each of A_1 and A_2 is H or OP'' , where P'' is H or a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$,

$NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(i) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;

(ii) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;


or

(iii) R_3 and R_4 combine to form a bond, and R_5 is H;

each P_4 and P_5 is independently H or a hydroxyl protecting group; and

R_6 is H, and P_6 is H or a hydroxyl protecting group; or R_6 and P_6 combine to form a double bond.


Further described herein is a compound of formula (VA), formula (VB), or formula (VC):



or a salt or a tautomer thereof,

where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,
 NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B
 is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl,
 alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

10 k is 0 or 1;

n is 0, 1, or 2;

(a1) R₃ is H or a hydroxyl protecting group, R₄ is alkyl ether, and R₅ is H;

(a2) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

15 (a3) R₃ and R₄ combine to form a bond, and R₅ is H;

(b1) A₁ and R₇ combine to form oxo, R₆ is H or a hydroxyl protecting group, and R₈ is H;

or

(b2) A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₆ is H or a hydroxyl protecting group, and R₇ and R₈ combine to form a
 20 double bond;

or

(ii) R₆ and R₇ combine to form a bond, and R₈ is H or OP'';


(c1) R₉ is H, and P₄ is H or a hydroxyl protecting group;

or

25 (c2) R₉ and P₄ combine to form a double bond;

R_{10} is H or $-\text{CH}_2\text{X}_1\text{CH}_2\text{CH}=\text{CH}_2$, where X_1 is O, $-\text{C}(\text{R}_{11})_2-$, or NP_6 , and where each R_{11} is independently H or $-\text{COOR}_{12}$, P_6 is an *N*-protecting group, and R_{12} is alkyl;
each P_5 is independently H or a hydroxyl protecting group; and


5 Further described herein is a compound of formula (VIA):



or a salt or a tautomer thereof,

where

10 each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:



15


where

L is $-(\text{CH}(\text{OP}_2))-$, $-(\text{C}(\text{OH})(\text{OP}_2))-$, or $-\text{C}(\text{O})-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

20

R_2 is H or $-(\text{CH}_2)_n\text{OP}_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

Y is iodide, bromide, or trifluoromethanesulfonate;

b designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or *b* designates (*S*)-stereogenic center, and Z is OR₆, where R₆ is a hydroxyl protecting group;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;
or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:


(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))-, wherein P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal.

Further described herein is a compound of formula (VIB):



(VIB)

or a salt or a tautomer thereof,

where


b designates (*R*)-stereogenic center, *c* designates (*S*)-stereogenic center, and *Z* is a sulfonate, chloride, bromide, or iodide;

or

b designates (*S*)-stereogenic center, *c* designates (*R*)-stereogenic center, and *Z* is *OR*₆,

where *R*₆ is a hydroxyl protecting group;

each of *D* and *D'* is independently H, optionally substituted alkyl, or *OP*₁, provided that only one of *D* and *D'* is *OP*₁, where *P*₁ is H, alkyl, or a hydroxyl protecting group; *A* is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and *Q*₁, the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

*R*₁ is H, or *R*₁ and *P*₁ combine to form a bond;

*R*₂ is H or $-(CH_2)_nOP_3$, and each of *P*₂ and *P*₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or *P*₂ and *P*₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or *R*₂ and *P*₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



group;

E is optionally substituted alkyl or optionally substituted alkoxy;

5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

10 A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:


(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

15 each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))-, wherein P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal.


Further described herein is a compound of formula (VIIA) or formula (VIIB):



or a salt thereof,

where

each of D and D' is independently H , optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H , alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:





where

L is $-(CH(OP_2))_n-$, $-(C(OH)(OP_2))_n-$, or $-C(O)-$;

R_1 is H , or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H , optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:





, where each P' is independently H or a hydroxyl protecting

group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

5 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

10 n is 0, 1, or 2;

(a1) R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or -(CH(OR₈))-;

or

15 (a2) R₃ and R₄ combine to form a bond, R₅ is H, and

each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or -(CH(OR₈))-;

or

20 both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal;

(b1) Z is chloride, bromide, or iodide, and R₆ and R₇ combine to form a bond;

(b2) Z and R₇ combine to form a double bond, and R₆ is a hydroxyl protecting group;


or

(b3) when Z and R₇ are absent, R₆ is H or a hydroxyl protecting group;

25 P₅ is H or a hydroxyl protecting group;

where R₈ is H or a hydroxyl protecting group.


Further described herein is a compound of formula (VIIC):



or a salt or a tautomer thereof,

where

each of D and D' is independently H , optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H , alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:





where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H , or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H , optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:





, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

5 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

10 n is 0, 1, or 2;

each P₄ is H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))-, wherein P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form a ketal; and

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

15 or


A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

20 (ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP''.

Further described herein is a compound of formula (VIII C):




25 or a salt thereof,

where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆

saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




where

L is $-(\text{CH}(\text{OP}_2))-$, $-(\text{C}(\text{OH})(\text{OP}_2))-$, or $-\text{C}(\text{O})-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(\text{CH}_2)_n\text{OP}_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;


each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,

NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

P₄ is H or a hydroxyl protecting group; and



where P_5 is H or a hydroxyl protecting group; each P_6 is independently a hydroxyl protecting group, or both P_6 groups, together with the atoms to which each is attached, combine to form a cyclic protected diol; and R_5 is H or $-\text{CH}_2\text{X}_1\text{CH}_2\text{CH}=\text{CH}_2$, where X_1 is O, $-\text{CH}_2-$, or NP_7 , where P_7 is a sulfonyl.

The disclosure also features compounds of formula (IA), (IB), (IC), (IE), (IF), (IIA), (IIB), (IIC), (IICa), (IIIA), (IIIB), (IVA), (IVB), (IVC), (IVD), (IVE), (VA), (VB), (VBa), (VC), (VIA), (VIB), (VIC), (VIIA), (VIIB), (VIIC), (VIID), (VIIE), (VIIF), (VIIG), (VIIIA), (VIIIB), (IIIC), or (IIID).

In certain examples of formula (IA), (IB), (IC), (IIA), (IIB), (IIIA), (IIIB), (VIA), (VIB), (VIC), (VIIA), (VIIB), and/or (VIIC) as described above, each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal (e.g., each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo).

In particular examples of formula (VIIIA) as described above, X is O or X , together with the carbon to which it is attached, forms an acetal (e.g., cyclic acetal).

Definitions

Compounds useful in the invention may be isotopically labeled compounds. Useful isotopes include hydrogen, carbon, nitrogen, and oxygen (e.g., ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , and ^{17}O). Isotopically-labeled compounds can be prepared by synthesizing a compound using a readily available isotopically-labeled reagent in place of a non-isotopically-labeled reagent.

For any of the following chemical definitions, a number following an atomic symbol indicates that total number of atoms of that element that are present in a particular chemical moiety. As will be understood, other atoms, such as hydrogen atoms, or substituent groups, as described herein, may be present, as necessary, to satisfy the valences of the atoms. For example, an unsubstituted C_2 alkyl group has the formula $-\text{CH}_2\text{CH}_3$. When used with the groups defined herein, a reference to the number of carbon atoms includes the divalent carbon in acetal and ketal groups but does not include the carbonyl carbon in acyl, ester, carbonate, or carbamate groups. A reference to the number of oxygen, nitrogen, or sulfur atoms in a heteroaryl group only includes those atoms that form a part of a heterocyclic ring.

By "acetal" is meant $-\text{O}-(\text{CHR})-\text{O}-$, where R is H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, optionally substituted arylalkyl, or a bond to the chain of carbon atoms within an intermediate in the synthesis of the halichondrin macrolide.

By "acyl" is meant $-C(O)R$, where R is H, alkyl, alkenyl, aryl, or arylalkyl. In exemplary acyl groups, R is H, C_{1-12} alkyl (e.g., C_{1-8} , C_{1-6} , C_{1-4} , C_{2-7} , C_{3-12} , or C_{3-6} alkyl), C_{2-12} alkenyl (e.g., C_{2-8} , C_{2-6} , C_{2-4} , C_{3-12} , or C_{3-6} alkenyl), C_{6-20} aryl (e.g., C_{6-15} , C_{6-10} , C_{8-20} , or C_{8-15} aryl), monocyclic C_{1-6} heteroaryl (e.g., monocyclic C_{1-4} or C_{2-6} heteroaryl), C_{4-19} heteroaryl (e.g., C_{4-10} heteroaryl), $(C_{6-15})\text{aryl}(C_{1-6})\text{alkyl}$, $(C_{1-6})\text{heteroaryl}(C_{1-6})\text{alkyl}$, or $(C_{4-19})\text{heteroaryl}(C_{1-6})\text{alkyl}$. As defined herein, any heteroaryl group present in an acyl group has from 1 to 4 heteroatoms selected independently from O, N, and S. An acyl group can be unsubstituted or substituted (e.g., optionally substituted acyl). In the optionally substituted acyl group, the substituent R is H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl. In some embodiments, acyl is C_{2-10} acyl.

By "acylating agent" is meant a compound that reacts with an amine or a hydroxyl group to produce an amide or an ester, respectively. An acylating agent has a formula $R-LG$, where R is acyl, and LG is halogen, carbonate, or $-OR'$, where R' is acyl.

By "alkoxide" is meant an anionic compound RO^- , where R is alkyl. A counterion for alkoxide can be an alkali metal cation, an alkali earth metal cation, or a tetraalkylammonium cation. Alkoxide can be optionally substituted in the same manner as alkyl.

By "alkoxy" is meant $-OR$, where R is alkyl. Alkoxy can be optionally substituted in the same manner as alkyl.

By "alkoxyalkyl" is meant $-OR$, where R is alkyl substituted by alkoxy. Each portion of the alkoxyalkyl can be optionally substituted in the same manner as alkyl.

By "alkoxyaryl" is meant $-R'(R'')_n$, where n is 1 or 2, R' is arylene and R'' is alkoxy, as defined herein. R' can be further optionally substituted in the same manner as aryl. R'' can be optionally substituted in the same manner as alkyl.

By "alkoxyarylalkyl" is meant $-R'(R''(R'''))_n$, where n is an integer from 1 to 3, R' is arylene, R'' is arylene, and R''' is alkoxy, as defined herein. R' can be optionally substituted in the same manner as alkyl. R'' can be further optionally substituted in the same manner as aryl. R''' can be optionally substituted in the same manner as alkyl.

By "alkyl" is meant a straight or branched chain saturated cyclic (i.e., cycloalkyl) or acyclic hydrocarbon group of from 1 to 12 carbons, unless otherwise specified. In some embodiments, alkyl is C_{1-6} alkyl. Exemplary alkyl groups include C_{1-8} , C_{1-6} , C_{1-4} , C_{2-7} , C_{3-12} , and C_{3-6} alkyl. Specific examples include methyl, ethyl, 1-propyl, 2-propyl, 2-methyl-1-propyl, 1-butyl, 2-butyl, and the like. Alkyl group can be optionally substituted with 1, 2, 3, or 4 substituents selected from the group consisting of halogen, hydroxy, alkoxy, aryloxy, arylalkyloxy, amino, oxo, alkylthio, alkylenedithio, alkylamino,

[alkenyl]alkylamino, [aryl]alkylamino, [arylalkyl]alkylamino, dialkylamino, silyl, sulfonyl, cyano, nitro, carboxyl, and azido.

By "alkylamino" is meant -NHR , where R is alkyl. By "[alkenyl]alkylamino" is meant $\text{-NRR}'$, where R is alkyl, and R' is alkenyl. By "[aryl]alkylamino" is meant $\text{-NRR}'$, where R is alkyl, and R' is aryl. By "[arylalkyl]alkylamino" is meant $\text{-NRR}'$, where R is alkyl, and R' is arylalkyl. By "dialkylamino" is meant -NR_2 , where each R is alkyl, selected independently.

By "alkylaryl" is meant $\text{-R}'(\text{R})_n$, where n is an integer from 1 to 3, R' is arylene, and R is alkyl. Alkylaryl can be optionally substituted in the same manner as defined for each R' and R group.

By "alkylene" is meant a multivalent alkyl group. Alkylene groups can be optionally substituted in the same manner as alkyl groups. For example, a C_1 alkylene group is $\text{-CH}_2\text{-}$.

By "alkylenedithio" is meant -S-alkylene-S- . Alkylenedithio can be optionally substituted in the same manner as an alkylene group.

By "alkylhaloaryl" is meant $\text{-R}'(\text{R})_n\text{-R}''$, where n is an integer from 1 to 5 and R' is arylene, R is halogen, and R'' is alkylene, as defined herein. R' can be further optionally substituted in the same manner as aryl. R'' can be further optionally substituted in the same manner as alkyl.

By "alkylthio" is meant -SR , where R is alkyl. Alkylthio can be optionally substituted in the same manner as an alkyl group.

By "alkenyl" is meant a straight or branched chain cyclic or acyclic hydrocarbon group of, unless otherwise specified, from 2 to 12 carbons and containing one or more carbon-carbon double bonds. In some embodiments, alkenyl is C_{2-6} alkenyl. Exemplary alkenyl groups include C_{2-8} , C_{2-7} , C_{2-6} , C_{2-4} , C_{3-12} , and C_{3-6} alkenyl. Specific examples include ethenyl (i.e., vinyl), 1-propenyl, 2-propenyl (i.e., allyl), 2-methyl-1-propenyl, 1-butenyl, 2-butenyl (i.e., crotyl), and the like. Alkenyl group can be optionally substituted in the same manner as alkyl groups. Alkenyl groups, used in any context herein, may also be substituted with an aryl group.

By "amido" is meant -NHR , where R is acyl. Amido can be optionally substituted in the same manner as acyl.

By "aminal" is meant $\text{-O-CR}_2\text{-NR}'$, where each R is independently H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl, or both R groups are together optionally substituted alkylene, and R' is H or an N-protecting group. In particular, R' can be an N-protecting group (e.g., Boc).

By "amino" is meant $-NR_2$, where N and R_2 combine to form azido, or each R is independently H or an *N*-protecting group, or both R combine to form an *N*-protecting group. Amino can be unmasked, when each R is H, or masked, when at least one R is not H. Thus, optionally masked amino can be masked or unmasked amino.

5

By "aminoalkyl" is meant $-R'(R)_n$, where n is 1 or 2, R' is alkylene, and R" is amino, as defined herein. R' can be optionally substituted in the same manner as an alkyl group.

By "aryl" is meant a monocyclic or multicyclic ring system having one or more aromatic rings, where the ring system is carbocyclic. Exemplary aryl groups include C_{6-20} , C_{6-15} , C_{6-10} , C_{8-20} , and C_{8-15} aryl. A preferred aryl group is a C_{6-10} aryl group. Specific examples of carbocyclic aryl groups include phenyl, indanyl, indenyl, naphthyl, phenanthryl, anthracyl, and fluorenyl. Aryl group can be optionally substituted with 1, 2, 3, 4, or 5 substituents selected from the group consisting of alkyl, alkenyl, aryl, arylalkyl, halogen, alkoxy, aryloxy, arylalkyloxy, alkylthio, alkylenedithio, alkylamino, [alkenyl]alkylamino, [aryl]alkylamino, [arylalkyl]alkylamino, dialkylamino, silyl, sulfonyl, cyano, nitro, carboxyl, and azido.

15

By "arylalkyl" is meant $-R'R$, where R' is alkylene, and R" is aryl. Arylalkyl can be optionally substituted in the same manner as defined for each R' and R" group.

By "arylalkyloxy" is meant $-OR$, where R is arylalkyl. Arylalkyloxy can be optionally substituted in the same manner as defined for arylalkyl.

20

By "arylene" is meant a multivalent aryl group. Arylene groups can be optionally substituted in the same manner as aryl groups. For example, a C_6 arylene group is phenylene.

By "aryloxy" is meant $-OR$, where R is aryl. Aryloxy can be optionally substituted in the same manner as aryl.

By "azido" is meant $-N_3$.

25

By "boronate" is meant $-OBRO-$, where R is alkyl, alkenyl, aryl, arylalkyl, alkoxy, or 2,6-diacetamidophenyl. Boronate can be substituted, when R is a substituted alkyl, substituted alkenyl, substituted aryl, substituted arylalkyl, or substituted alkoxy. Alternatively, boronate can be unsubstituted, when R is unsubstituted alkyl, unsubstituted alkenyl, aryl, unsubstituted arylalkyl, unsubstituted alkoxy, or 2,6-diacetamidophenyl.

30

By "carbamate" is meant a group, when a hydroxyl protecting group, having the formula $-OC(O)NR_2$, or, when an amine protecting group, having the formula $-NR'-C(O)OR$, where each R and R' is independently H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl.

35

40

By "carbonate" is meant --OC(O)OR , where R is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl.

By "carbonyl" is meant --C(O)-- .

By "carboxyl" is meant --C(O)OH , in free acid, ionized, or salt form.

By "carboxylic acid" is meant R-OH , where R is optionally substituted acyl.

By "carboxylic acid anhydride" is meant R-O-R , where each R is independently optionally substituted acyl.

By "cyclic carbonate" is meant --OC(O)O-- that is part of a ring.

By "dicarbonyl" is meant --C(O)--C(O)-- . Dicarbonyl-dioxo is --OC(O)--COO-- .

By "ester" is meant --OC(O)R , where --C(O)R is an optionally substituted acyl group.

By "ether" is meant --OR , where R is alkyl, alkenyl, arylalkyl, silyl, or 2-tetrahydropyranyl. Ether can be optionally substituted as defined for each R group.

By "halichondrin macrolide" is meant a lactone including the structure of carbons 1-30 as shown in Chart 1, wherein carbons 29 and 30 form part of a five- or six-membered ring.

By "haloalkyl" is meant --R'(R)_n , where n is an integer from 1 to 5 and R' is alkylene and R is halogen, as defined herein. R' can be further optionally substituted in the same manner as alkyl.


By "haloaryl" is meant --R'(R)_n , where n is an integer from 1 to 5 and R' is arylene and R is halogen, as defined herein. R' can be further optionally substituted in the same manner as aryl.

By "haloarylalkyl" is meant --R'(R''(R'''))_n , where n is an integer from 1 to 5 and R' is alkylene, R'' is arylene, and R''' is halogen, as defined herein. R' can be further optionally substituted in the same manner as alkyl. R'' can be further optionally substituted in the same manner as aryl.

By "halogen" is meant fluoro, chloro, bromo, or iodo.

By "heterocyclic radical" is meant a 5-, 6- or 7-membered ring, unless otherwise specified, containing one, two, three, or four heteroatoms independently selected from the group containing nitrogen, oxygen, and sulfur. The 5-membered ring has zero to one double bonds, and the 6- and 7-membered rings have zero to two double bonds. Certain heterocyclyl groups include from 1 to 9 carbon atoms. Other such groups may include up to 12 carbon atoms. The term "heterocyclyl" also represents a heterocyclic compound having a bridged multicyclic structure in which one or more carbons and/or heteroatoms bridges two non-

adjacent members of a monocyclic ring, *e.g.*, a quinuclidinyl group. The term "heterocyclyl" includes bicyclic, tricyclic, and tetracyclic groups in which any of the above heterocyclic rings is fused to one, two, or three carbocyclic rings, *e.g.*, an aryl ring, a cyclohexane ring, a cyclohexene ring, a cyclopentane ring, a cyclopentene ring, or another monocyclic heterocyclic ring, such as indolyl, quinolyl, isoquinolyl, tetrahydroquinolyl, benzofuryl, benzothienyl and the like. Examples of fused heterocyclyls include tropanes and 1,2,3,5,8,8a-hexahydroindolizine. Heterocyclics include pyrrolyl, pyrrolinyl, pyrrolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyridyl, piperidinyl, homopiperidinyl, pyrazinyl, piperazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiomorpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, benzothiazolyl, benzoxazolyl, furyl, thienyl, thiazolidinyl, isothiazolyl, isoindazolyl, triazolyl, tetrazolyl, oxadiazolyl, purinyl, thiadiazolyl (*e.g.*, 1,3,4-thiadiazole), tetrahydrofuryl, dihydrofuryl, tetrahydrothienyl, dihydrothienyl, dihydroindolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, pyranyl, dihydropyranyl, dithiazolyl, benzofuryl, benzothienyl and the like. Still other exemplary heterocyclyls include: 2,3,4,5-tetrahydro-2-oxo-oxazolyl; 2,3-dihydro-2-oxo-1H-imidazolyl; 2,3,4,5-tetrahydro-5-oxo-1H-pyrazolyl (*e.g.*, 2,3,4,5-tetrahydro-2-phenyl-5-oxo-1H-pyrazolyl); 2,3,4,5-tetrahydro-2,4-dioxo-1H-imidazolyl (*e.g.*, 2,3,4,5-tetrahydro-2,4-dioxo-5-methyl-5-phenyl-1H-imidazolyl); 2,3-dihydro-2-thioxo-1,3,4-oxadiazolyl (*e.g.*, 2,3-dihydro-2-thioxo-5-phenyl-1,3,4-oxadiazolyl); 4,5-dihydro-5-oxo-1H-triazolyl (*e.g.*, 4,5-dihydro-3-methyl-4-amino 5-oxo-1H-triazolyl); 1,2,3,4-tetrahydro-2,4-dioxypyridinyl (*e.g.*, 1,2,3,4-tetrahydro-2,4-dioxo-3,3-diethylpyridinyl); 2,6-dioxo-piperidinyl (*e.g.*, 2,6-dioxo-3-ethyl-3-phenylpiperidinyl); 1,6-dihydro-6-oxopyridiminy; 1,6-dihydro-4-oxopyrimidinyl (*e.g.*, 2-(methylthio)-1,6-dihydro-4-oxo-5-methylpyrimidin-1-yl); 1,2,3,4-tetrahydro-2,4-dioxopyrimidinyl (*e.g.*, 1,2,3,4-tetrahydro-2,4-dioxo-3-ethylpyrimidinyl); 1,6-dihydro-6-oxo-pyridazinyl (*e.g.*, 1,6-dihydro-6-oxo-3-ethylpyridazinyl); 1,6-dihydro-6-oxo-1,2,4-triazinyl (*e.g.*, 1,6-dihydro-5-isopropyl-6-oxo-1,2,4-triazinyl); 2,3-dihydro-2-oxo-1H-indolyl (*e.g.*, 3,3-dimethyl-2,3-dihydro-2-oxo-1H-indolyl and 2,3-dihydro-2-oxo-3,3'-spiropropane-1H-indol-1-yl); 1,3-dihydro-1-oxo-2H-iso-indolyl; 1,3-dihydro-1,3-dioxo-2H-iso-indolyl; 1H-benzopyrazolyl (*e.g.*, 1-(ethoxycarbonyl)-1H-benzopyrazolyl); 2,3-dihydro-2-oxo-1H-benzimidazolyl (*e.g.*, 3-ethyl-2,3-dihydro-2-oxo-1H-benzimidazolyl); 2,3-dihydro-2-oxo-benzoxazolyl (*e.g.*, 5-chloro-2,3-dihydro-2-oxo-benzoxazolyl); 2,3-dihydro-2-oxo-benzoxazolyl; 2-oxo-2H-benzopyranyl; 1,4-benzodioxanyl; 1,3-benzodioxanyl; 2,3-dihydro-3-oxo,4H-1,3-benzothiazinyl; 3,4-dihydro-4-oxo-3H-quinazolinyl (*e.g.*, 2-methyl-3,4-dihydro-4-oxo-3H-quinazolinyl); 1,2,3,4-tetrahydro-2,4-dioxo-3H-quinazolyl (*e.g.*, 1-ethyl-1,2,3,4-tetrahydro-2,4-dioxo-3H-quinazolyl); 1,2,3,6-tetrahydro-2,6-dioxo-7H-purinyl (*e.g.*, 1,2,3,6-tetrahydro-1,3-dimethyl-2,6-dioxo-7H-purinyl); 1,2,3,6-tetrahydro-2,6-dioxo-1H-purinyl (*e.g.*, 1,2,3,6-tetrahydro-3,7-dimethyl-2,6-dioxo-1H-purinyl); 2-oxobenz[*c,d*]indolyl; 1,1-dioxo-2H-naphth[1,8-*c,d*]isothiazolyl; and 1,8-naphthylenedicarboxamido. Heterocyclic groups also include groups of the formula



F' is selected from the group consisting of -CH₂-, -CH₂O- and -O-, and G' is selected from the group consisting of -C(O)- and -(C(R')(R''))_v-, where each of R' and R'' is, independently, selected from the group consisting of hydrogen or alkyl of one to four carbon atoms, and v is one to three and includes groups, such as 1,3-benzodioxolyl, 1,4-benzodioxanyl, and the like. Any of the heterocyclyl groups

mentioned herein may be optionally substituted with one, two, three, four or five substituents independently selected from the group consisting of: (1) alkanoyl (*e.g.*, formyl, acetyl, and the like); (2) alkyl (*e.g.*, alkoxyalkylene, alkylsulfanylalkylene, aminoalkylene, azidoalkylene, acylalkylene, haloalkylene (*e.g.*, perfluoroalkyl), hydroxyalkylene, nitroalkylene, or thioalkoxyalkylene); (3) alkenyl; (4) alkynyl; (5) alkoxy (*e.g.*, perfluoroalkoxy); (6) alkylsulfanyl; (7) aryl; (8) amino; (9) aryl-alkylene; (10) azido; (11) cycloalkyl; (12) cycloalkyl-alkylene; (13) cycloalkenyl; (14) cycloalkenyl-alkylene; (15) halo; (16) heterocyclyl (*e.g.*, heteroaryl); (17) (heterocyclyl)oxy; (18) (heterocyclyl)aza; (19) hydroxy; (20) oxo; (21) nitro; (22) sulfide; (23) thioalkoxy; (24) $-(CH_2)_qCO_2R^A$, where q is an integer from zero to four, and R^A is selected from the group consisting of (a) alkyl, (b) aryl, (c) hydrogen, and (d) aryl-alkylene; (25) $-(CH_2)_qCONR^BR^C$, where q is an integer from zero to four and where R^B and R^C are independently selected from the group consisting of (a) hydrogen, (b) alkyl, (c) aryl, and (d) aryl-alkylene; (26) $-(CH_2)_qSO_2R^D$, where q is an integer from zero to four and where R^D is selected from the group consisting of (a) alkyl, (b) aryl, and (c) aryl-alkylene; (27) $-(CH_2)_qSO_2NR^ER^F$, where q is an integer from zero to four and where each of R^E and R^F is, independently, selected from the group consisting of (a) hydrogen, (b) alkyl, (c) aryl, and (d) aryl-alkylene; (28) thiol; (29) aryloxy; (30) cycloalkoxy; (31) arylalkoxy; (31) heterocyclyl-alkylene (*e.g.*, heteroaryl-alkylene); (32) silyl; (33) cyano; and (34) $-S(O)R^H$ where R^H is selected from the group consisting of (a) hydrogen, (b) alkyl, (c) aryl, and (d) aryl-alkylene. In some embodiments, each of these groups can be further substituted as described herein. For example, the alkylene group of an aryl- C_1 -alkylene or a heterocyclyl- C_1 -alkylene can be further substituted with an oxo group to afford the respective aryloyl and (heterocyclyl)oyl substituent group. In addition, when a heterocyclyl group is present in a bioreversible group of the invention it may be substituted with an ester, thioester, or disulfide group that is bound to a conjugating moiety, a hydrophilic functional group, or an auxiliary moiety as defined herein.

By "heterocyclic radical alkyl," as used herein, represents an alkyl group substituted with a heterocyclic radical. The heterocyclic radical and alkyl portions may be substituted as the individual groups as described herein.

By "hydroxyalkyl" is meant $-R'(R'')_n$, where n 1 or 2, R' is alkylene and R'' is hydroxyl, as defined herein. R' can be further optionally substituted in the same manner as alkyl.

By "hydroxyaryl" is meant $-R'(R'')_n$, where n is 1 or 2, R' is arylene and R'' is hydroxyl, as defined herein. R' can be further optionally substituted in the same manner as aryl.

By "hydroxyl" is meant $-OH$.

By "hydroxyl protecting group" is meant any group capable of protecting the oxygen atom to which it is attached from reacting or bonding. Hydroxyl protecting groups are known in the art, *e.g.*, as described in Wuts, Greene's Protective Groups in Organic Synthesis, Wiley-Interscience, 4th Edition, 2006.

Exemplary protecting groups (with the oxygen atom to which they are attached) are independently selected from the group consisting of esters, carbonates, carbamates, sulfonates, and ethers. In

exemplary ester hydroxyl protecting groups, R of the acyl group is C₁₋₁₂ alkyl (e.g., C₁₋₈, C₁₋₆, C₁₋₄, C₂₋₇, C₃₋₁₂, and C₃₋₆ alkyl), C₂₋₁₂ alkenyl (e.g., C₂₋₈, C₂₋₆, C₂₋₄, C₃₋₁₂, and C₃₋₆ alkenyl), carbocyclic C₆₋₂₀ aryl (e.g., C₆₋₁₅, C₆₋₁₀, C₈₋₂₀, and C₈₋₁₅ aryl), monocyclic C₁₋₆ heteroaryl (e.g., C₁₋₄ and C₂₋₆ heteroaryl), C₄₋₁₉ heteroaryl (e.g., C₄₋₁₀ heteroaryl), (C₆₋₁₅)aryl(C₁₋₆)alkyl, (C₄₋₁₉)heteroaryl(C₁₋₆)alkyl, or (C₁₋₆)heteroaryl(C₁₋₆)alkyl.

5 Specific examples of acyl groups for use in esters include formyl, benzoylformyl, acetyl (e.g., unsubstituted or chloroacetyl, trifluoroacetyl, methoxyacetyl, triphenylmethoxyacetyl, and p-chlorophenoxyacetyl), 3-phenylpropionyl, 4-oxopentanoyl, 4,4-(ethylenedithio)pentanoyl, pivaloyl (Piv), vinylpivaloyl, crotonoyl, 4-methoxy-crotonoyl, naphthoyl (e.g., 1- or 2-naphthoyl), and benzoyl (e.g., unsubstituted or substituted, e.g., p-methoxybenzoyl, phthaloyl (including salts, such as triethylamine and potassium), p-bromobenzoyl, and 2,4,6-trimethylbenzoyl). As defined herein, any heteroaryl group
10 present in an ester group has from 1 to 4 heteroatoms selected independently from O, N, and S. In exemplary carbonate hydroxyl protecting groups, R is C₁₋₁₂ alkyl (e.g., C₁₋₈, C₁₋₆, C₁₋₄, C₂₋₇, C₃₋₁₂, and C₃₋₆ alkyl), C₂₋₁₂ alkenyl (e.g., C₂₋₈, C₂₋₆, C₂₋₄, C₃₋₁₂, and C₃₋₆ alkenyl), carbocyclic C₆₋₂₀ aryl (e.g., C₆₋₁₅, C₆₋₁₀, C₈₋₂₀, and C₈₋₁₅ aryl), monocyclic C₁₋₆ heteroaryl (e.g., C₁₋₄ and C₂₋₆ heteroaryl), C₄₋₁₉ heteroaryl (e.g., C₄₋₁₀ heteroaryl), (C₆₋₁₅)aryl(C₁₋₆)alkyl, (C₄₋₁₉)heteroaryl(C₁₋₆)alkyl, or (C₁₋₆)heteroaryl(C₁₋₆)alkyl. Specific
15 examples include methyl, 9-fluorenylmethyl, ethyl, 2,2,2-trichloroethyl, 2-(trimethylsilyl)ethyl, 2-(phenylsulfonyl)ethyl, vinyl, allyl, t-butyl, p-nitrobenzyl, and benzyl carbonates. As defined herein, any heteroaryl group present in a carbonate group has from 1 to 4 heteroatoms selected independently from O, N, and S. In exemplary carbamate hydroxyl protecting groups, each R is independently H, C₁₋₁₂ alkyl (e.g., C₁₋₈, C₁₋₆, C₁₋₄, C₂₋₇, C₃₋₁₂, and C₃₋₆ alkyl), C₂₋₁₂ alkenyl (e.g., C₂₋₈, C₂₋₆, C₂₋₄, C₃₋₁₂, and C₃₋₆ alkenyl), carbocyclic C₆₋₂₀ aryl (e.g., C₆₋₁₅, C₆₋₁₀, C₈₋₂₀, and C₈₋₁₅ aryl), monocyclic C₁₋₆ heteroaryl (e.g., C₁₋₄ and C₂₋₆ heteroaryl), C₄₋₁₉ heteroaryl (e.g., C₄₋₁₀ heteroaryl), (C₆₋₁₅)aryl(C₁₋₆)alkyl, (C₄₋₁₉)heteroaryl(C₁₋₆)alkyl, or (C₁₋₆)heteroaryl(C₁₋₆)alkyl. Specific examples include N-phenyl and N-methyl-N-(o-nitrophenyl) carbamates.

As defined herein, any heteroaryl group present in a carbamate group has from 1 to 4 heteroatoms
25 selected independently from O, N, and S. Exemplary ether hydroxyl protecting groups include C₁₋₁₂ alkyl (e.g., C₁₋₈, C₁₋₆, C₁₋₄, C₂₋₇, C₃₋₁₂, and C₃₋₆ alkyl), C₂₋₁₂ alkenyl (e.g., C₂₋₈, C₂₋₆, C₂₋₄, C₃₋₁₂, and C₃₋₆ alkenyl), (C₆₋₁₅)aryl(C₁₋₆)alkyl, (C₄₋₁₉)heteroaryl(C₁₋₆)alkyl, (C₁₋₆)heteroaryl(C₁₋₆)alkyl, (C₁₋₆)alkoxy(C₁₋₆)alkyl, (C₁₋₆)alkylthio(C₁₋₆)alkyl, (C₆₋₁₀)aryl(C₁₋₆)alkoxy(C₁₋₆)alkyl, and silyl (e.g., tri(C₁₋₆ alkyl)silyl, tri(C₆₋₁₀ aryl or C₁₋₆ heteroaryl)silyl, di(C₆₋₁₀ aryl or C₁₋₆ heteroaryl)(C₁₋₆ alkyl)silyl, and (C₆₋₁₀ aryl or C₁₋₆ heteroaryl)di(C₁₋₆ alkyl)silyl). Specific examples of alkylethers include methyl and t-butyl, and an example of an alkenyl
30 ether is allyl. Ether hydroxyl protecting groups can be used to protect a carboxyl group (e.g., with a C₁₋₁₂ alkyl (e.g., C₁₋₈, C₁₋₆, C₁₋₄, C₂₋₇, C₃₋₁₂, and C₃₋₆ alkyl), (C₆₋₁₅)aryl(C₁₋₆)alkyl, (C₁₋₆)alkoxy(C₁₋₆)alkyl, (C₁₋₆)alkylthio(C₁₋₆)alkyl, or (C₆₋₁₀)aryl(C₁₋₆)alkoxy(C₁₋₆)alkyl). Examples of alkoxyalkyls and alkylthioalkyls that can be used as ether hydroxyl protecting groups include methoxymethyl, methylthiomethyl, (2-methoxyethoxy)methyl, and β-(trimethylsilyl)ethoxymethyl. Examples of arylalkyl groups that can be used
35 as ether hydroxyl protecting groups include benzyl, p-methoxybenzyl (MPM), 3,4-dimethoxybenzyl, triphenylmethyl (trityl), o-nitrobenzyl, p-nitrobenzyl, p-halobenzyl, 2,6-dichlorobenzyl, p-cyanobenzyl, naphthylmethyl, and 2- and 4-picolyl ethers. Specific examples of silylethers include trimethylsilyl (TMS), triethylsilyl (TES), t-butyl dimethylsilyl (TBS), t-butyl diphenylsilyl (TBDPS), triisopropylsilyl (TIPS), and triphenylsilyl (TPS) ethers. An example of an arylalkyloxyalkylether is benzyloxymethyl ether. As defined
40 herein, any heteroaryl group present in an ether group has from 1 to 4 heteroatoms selected

independently from O, N, and S. Vicinal or 1,3-diols may be protected with a diol protecting group (e.g., to produce a "cyclic protected diol"), such as acetal (e.g., containing C₁₋₆ alkylene), ketal (e.g., containing C₃₋₆ alkylene or C₃₋₆ cycloalkyl), cyclic silylene, cyclic carbonate, and cyclic boronate. Examples of acetal and ketal groups include methylene-dioxo, ethylidene-dioxo, benzylidene-dioxo, isopropylidene-dioxo, cyclohexylidene-dioxo, and cyclopentylidene-dioxo. An example of a cyclic silylene is di-*t*-butylsilylene. Another diol protecting group is 1,1,3,3-tetraisopropylsiloxanediyl. Examples of cyclic boronates include methyl, ethyl, phenyl, and 2,6-diacetamidophenyl boronates. Protecting groups may be substituted as is known in the art; for example, aryl and arylalkyl groups, such as phenyl, benzyl, naphthyl, or pyridinyl, can be substituted with C₁₋₆ alkyl, C₁₋₆ alkoxy, nitro, cyano, carboxyl, or halogen. Alkyl groups, such as methyl, ethyl, isopropyl, *n*-propyl, *t*-butyl, *n*-butyl, and *sec*-butyl, and alkenyl groups, such as vinyl and allyl, can also be substituted with oxo, arylsulfonyl, halogen, and trialkylsilyl groups. Preferred protecting groups are TBS and Piv. Protecting groups that are orthogonal are removed under different conditions, as is known in the art.

By "imido" is meant NR_2 , where each R is independently optionally substituted acyl.

By "ketal" is meant $\text{O-CR}_2\text{-O-}$, where each R is independently optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, optionally substituted arylalkyl, or a bond to the chain of carbon atoms within an intermediate in the synthesis of the halichondrin macrolide; or both R groups are together optionally substituted alkylene.

By "macrocyclization" is meant a reaction converting a non-macrocyclic compound into a compound containing at least one *n*-membered ring, where *n* is equal to or greater than 16.

By "MNBA" is meant 2-methyl-6-nitrobenzoic anhydride.

By "non-enolizable" is meant a group that, either alone or in combination with a group to which it is attached, cannot form an enol through a deprotonation/reprotonation sequence. For example, a "non-enolizable alkyl" can be bonded to a sulfone group or to a carbonyl group through a quaternary carbon atom (i.e., the carbon atom that is not bonded to a hydrogen atom).

By "non-macrocyclic" is meant a compound not containing rings or containing one or more *m*-membered rings, where *m* is less than or equal to 15.

By "*N*-protecting group" is meant a group protecting a nitrogen atom in a molecule from participating in one or more undesirable reactions during chemical synthesis (e.g., oxidation reactions, or certain nucleophilic and electrophilic substitutions). Commonly used *N*-protecting groups are disclosed in Wuts, Greene's Protective Groups in Organic Synthesis, Wiley-Interscience, 4th Edition, 2006. Exemplary *N*-protecting groups include acyl (e.g., formyl, acetyl, trifluoroacetyl, propionyl, pivaloyl, *t*-butylacetyl, 2-chloroacetyl, 2-bromoacetyl, trifluoroacetyl, trichloroacetyl, phthalyl, *o*-nitrophenoxyacetyl, α -chlorobutyryl, benzoyl, 4-chlorobenzoyl, and 4-bromobenzoyl); sulfonyl-containing groups (e.g., benzenesulfonyl, *p*-

toluenesulfonyl, o-nitrobenzenesulfonyl, and p-nitrobenzenesulfonyl); carbamate forming groups (e.g., benzyloxycarbonyl, p-chlorobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, 3,4-dimethoxybenzyloxycarbonyl, 3,5-dimethoxybenzyloxycarbonyl, 2,4-dimethoxybenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitro-4,5-dimethoxybenzyloxycarbonyl, 3,4,5-trimethoxybenzyloxycarbonyl, 1-(p-biphenyl)-1-methylethoxycarbonyl, α,α -dimethyl-3,5-dimethoxybenzyloxycarbonyl, benzhydryloxy carbonyl, t-butylloxycarbonyl, diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, methoxycarbonyl, allyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, phenoxycarbonyl, 4-nitrophenoxy carbonyl, fluorenyl-9-methoxycarbonyl, cyclopentylloxycarbonyl, adamantylloxycarbonyl, cyclohexylloxycarbonyl, and phenylthiocarbonyl), arylalkyl (e.g., triphenylmethyl); silyl groups (e.g., trimethylsilyl); and imine-forming groups (e.g., diphenylmethylene). Preferred *N*-protecting groups are acetyl, benzoyl, phenylsulfonyl, p-toluenesulfonyl, p-nitrobenzenesulfonyl, o-nitrobenzenesulfonyl, t-butylloxycarbonyl (Boc), and benzyloxycarbonyl (Cbz).

By "oxo" or (O) is meant =O.

By "pharmaceutically acceptable salt" is meant a salt within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and animals without undue toxicity, irritation, allergic response and the like and commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, pharmaceutically acceptable salts are described in: Berge et al., J. Pharmaceutical Sciences 66:1-19, 1977 and in Pharmaceutical Salts: Properties, Selection, and Use, (Eds. P.H. Stahl and C.G. Wermuth), Wiley-VCH, 2008. Representative acid addition salts include acetate, adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, hydrobromide, hydrochloride, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate, undecanoate, valerate salts and the like. A preferred salt is the mesylate salt.

By "silyl" is meant $-\text{SiR}_3$, where each R is independently alkyl, alkenyl, aryl, or arylalkyl. Examples of silyl groups include $\text{tri}(\text{C}_{1-6} \text{ alkyl})\text{silyl}$, $\text{tri}(\text{C}_{6-10} \text{ aryl or C}_{1-6} \text{ heteroaryl})\text{silyl}$, $\text{di}(\text{C}_{6-10} \text{ aryl or C}_{1-6} \text{ heteroaryl})(\text{C}_{1-6} \text{ alkyl})\text{silyl}$, and $(\text{C}_{6-10} \text{ aryl or C}_{1-6} \text{ heteroaryl})\text{di}(\text{C}_{1-6} \text{ alkyl})\text{silyl}$. It will be understood that, when a silyl group includes two or more alkyl, alkenyl, aryl, heteroaryl, or arylalkyl groups, these groups are independently selected. As defined herein, any heteroaryl group present in a silyl group has from 1 to 4 heteroatoms selected independently from O, N, and S. Silyl can be optionally substituted in the same manner as defined for each R group.

By “silylene” is meant $-\text{SiR}_2-$, where each R is independently alkyl, alkenyl, aryl, arylalkyl, or alkoxy. By “dialkylsilylene” is meant a silylene, where each R is alkyl. Silylene can be optionally substituted in the same manner as defined for each R group. Silylene-dioxo is a group having the formula $-\text{O}-\text{SiR}_2-\text{O}-$.

- 5 By “strong base” is meant a Brønsted base, the conjugate acid of which has pKa that is greater than or equal to 13. Non-limiting examples of strong bases include alkyl alkali metals (e.g., butyl lithium or Schlosser’s base), Grignard reagents (e.g., alkyl magnesium halide), alkoxides (e.g., tertiary alkoxides, such as *t*-butoxide), amides (e.g., diisopropylamide, tetramethylpiperidide, or bis(trimethylsilyl)amide), and phosphazene bases (e.g., Schwesinger base).

10

By “sulfonamide” is meant $-\text{NR}$, where R is sulfonyl.

- By “sulfonate” is meant $-\text{OS}(\text{O})_2\text{R}$, where R is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl. In exemplary sulfonates, R is C₁₋₁₂ alkyl (e.g., C₁₋₈, C₁₋₆, C₁₋₄, C₂₋₇, C₃₋₁₂, or C₃₋₆ alkyl), C₂₋₁₂ alkenyl (e.g., C₂₋₈, C₂₋₆, C₂₋₄, C₃₋₁₂, or C₃₋₆ alkenyl), carbocyclic C₆₋₂₀ aryl (e.g., C₆₋₁₅, C₆₋₁₀, C₈₋₂₀, or C₈₋₁₅ aryl), monocyclic C₁₋₆ heteroaryl (e.g., C₁₋₄ and C₂₋₆ heteroaryl), C₄₋₁₉ heteroaryl (e.g., C₄₋₁₀ heteroaryl), (C₆₋₁₅)aryl(C₁₋₆)alkyl, (C₄₋₁₉)heteroaryl(C₁₋₆)alkyl, or (C₁₋₆)heteroaryl(C₁₋₆)alkyl. As defined herein, any heteroaryl group present in a sulfonate group has from 1 to 4 heteroatoms selected independently from O, N, and S.

20

By “sulfonyl” is meant $-\text{S}(\text{O})_2\text{R}$, where R is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, optionally substituted arylalkyl, or silyl. Preferred R groups for sulfonyl are the same as those described above for sulfonates.

- 25 By “thioacetal” is meant $-\text{S}-(\text{CHR})-\text{S}-$, where R is H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl.

By “thioketal” is meant $-\text{S}-(\text{CR}_2)-\text{S}-$, where each R is independently optionally substituted alkyl, optionally substituted alkenyl, optionally substituted aryl, or optionally substituted arylalkyl.

30

By “triflate” is meant trifluoromethanesulfonate.

The pKa values recited herein refer to the pKa values of a conjugate Brønsted acid in water at room temperature, unless stated otherwise.

35

Brief Description of the Drawings

Figure 1 is a scheme showing preparation of a halichondrin macrolide involving a C.2-C.3 bond-forming macrocyclization through Horner-Wadsworth-Emmons olefination.

- 40 Figure 2 is a scheme showing preparation of a halichondrin macrolide involving a C.2-C.3 bond-forming macrocyclization through ring-closing olefin metathesis.

Figure 3 is a scheme showing preparation of a halichondrin macrolide involving a C.3-C.4 bond-forming macrocyclization through ring-closing olefin metathesis.

- 5 Figure 4 is a scheme showing preparation of a halichondrin macrolide involving a C.12-C.13 bond-forming macrocyclization through ring-closing olefin metathesis.

Figure 5 is a scheme showing preparation of a halichondrin macrolide involving a C.15-C.16 bond-forming macrocyclization through ring-closing olefin metathesis.

10

Figure 6 is a scheme showing preparation of a halichondrin macrolide involving a C.19-C.20 bond-forming macrocyclization through Nozaki-Hiyama-Kishi reaction.


15

Figure 7 is a scheme showing preparation of a halichondrin macrolide involving a C.26-C.27 bond-forming macrocyclization through Nozaki-Hiyama-Kishi reaction.

Detailed Description

The present invention provides methods for the synthesis of a halichondrin macrolide (see Chart 1) through a macrocyclization. Preferably, the halichondrin macrolide is a halichondrin B macrolide. The macrocyclizations of the present invention involve subjecting a non-macrocyclic intermediate to a carbon-carbon bond-forming reaction (e.g., an olefination reaction (e.g., Horner-Wadsworth-Emmons olefination), catalytic Ring-Closing Olefin Metathesis, or Nozaki-Hiyama-Kishi reaction) to afford a macrocyclic intermediate (e.g., a compound of formula (IB), (IIIB), (IVB), (VB), (VIB), or (VIIB)). The carbon-carbon bond forming reaction provides C.2-C.3, C.3-C.4, C.12-C.13, C.15-C.16, C.19-C.20, or C.26-C.27 bond in the structure of a halichondrin macrolide. The carbon-atom numbering scheme for a halichondrin macrolide is shown in Chart 1.


Chart 1



30 in which

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to

10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




where

L is $-(\text{CH}(\text{OP}_2))-$, $-(\text{C}(\text{OH})(\text{OP}_2))-$, or $-\text{C}(\text{O})-$;

R₁ is H, or R₁ and P₁ combine to form a bond;;

R₂ is H or $-(\text{CH}_2)_n\text{OP}_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



where each P' is independently H or a hydroxyl protecting group;

each of A₁ and A₂ is independently H or OP'', where each P'' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;


n is 0, 1, or 2.

In some embodiments of the halichondrin macrolide, C and D' are not the same, and a designates (*S*)-stereogenic center. In certain embodiments of the halichondrin macrolide, one of D and D' is optionally substituted alkyl or OP₁, the other of D and D' is H, and G is O. In particular embodiments of the halichondrin macrolide, one and only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group. In one embodiment of the halichondrin macrolide, k is 1, E is optionally substituted alkyl (e.g., Me); and D is a group of formula (1); in further embodiments, L is -(CH(OP₂))-, R₁ and P₁ combine to form a bond, R₂ is -(CH₂)_nOP₃, each of P₂ and P₃ is H, and n is 1 or 2 (e.g., n is 2).

The invention also provides intermediates in the synthesis of halichondrin and analogs of the macrolide in the structure of halichondrin and methods of preparing the same, as described herein.

C.2-C.3 Bond-forming Macrocyclization

The macrocyclization reaction of the invention can be a carbon-carbon bond-forming reaction (e.g., Horner-Wadsworth-Emmons olefination) that provides a C.2-C.3 bond in a halichondrin macrolide. The general synthetic sequence including Horner-Wadsworth-Emmons olefination that can be used to prepare a halichondrin macrolide is shown in Figure 1. As shown in Figure 1, the non-macrocyclic intermediate in the synthesis of the halichondrin macrolide can be a compound of formula (IA):




or a salt or a tautomer thereof,

in which

each R is independently optionally substituted alkyl or optionally substituted aryl;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , $NR_B R_A$, $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_B R_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_B R_A$, or $O(CO)NR_B R_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H; or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;


each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, where P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

P₅ is H or a hydroxyl protecting group.

5

In certain embodiments of the compound of formula (IA), R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo. In other embodiments of the compound of formula (IA), R₃ and R₄ combine to form a bond, and R₅ is H.

10 A macrocyclic intermediate in the synthesis of a halichondrin macrolide can be a compound of formula (IB), which can be produced by reacting the compound of formula (IA) with an organic base and a Lewis acid. The compound of formula (IB) has the following structure:



15 or a salt or a tautomer thereof,
where all variables are as defined for compound of formula (IA).

The reaction conditions that can be used to convert the compound of formula (IA) into the compound of formula (IB) include those known in the art for Horner-Wadsworth-Emmons reaction, e.g., Masamune-Roush conditions or Helquist protocol. In particular, the compound of formula (IA) can be reacted with an organic base (e.g., an organic base, the conjugate acid of which has a pKa of from 11 ± 2) and a Lewis acid (e.g., a salt of Li, Mg, or Zn). Non-limiting examples of an organic base that can be used in the Horner-Wadsworth-Emmons reaction include trialkylamines (e.g., triethylamine or Hünig's base), DBU, and DBN. Non-limiting examples of Lewis acids that can be used in the Horner-Wadsworth-Emmons reaction include LiCl, $\text{Zn}(\text{OTf})_2$, and MgCl_2 .

The compound of formula (IB), in which, e.g., P₅ is a hydroxyl protecting group, can be converted into a halichondrin macrolide by reacting the compound of formula (IB) with a hydroxyl protecting group removing agent (e.g., a fluoride source, if the hydroxyl protecting group is a silyl group).

30

If, in the compound of formula (IB), each P₄ is H, and X₁ is oxo, the synthesis may further involve a reaction with a Brønsted acid (e.g., a Brønsted acid having a pK_a of 5 ± 3) after the reaction of the

compound of formula (IB) with a hydroxyl protecting group removing agent (e.g., to convert P₄ from a hydroxyl protecting group into H).

If, in the compound of formula (IB), X₁ is oxo, R₃ is a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond, treatment with a hydroxyl protecting group removing agent can provide the compound of formula (IB), in which R₃ and R₄ combine to form a bond, and R₅ is H.


The compound of formula (IA) or formula (IB), in which X₁, together with the carbon atom to which it is attached, is $-(CH(OP_V))-$, where P_V is H, may be treated with an oxidizing agent capable of oxidizing hydroxyl to a carbonyl group (e.g., Dess-Martin periodinane or a dimethylsulfonium compound) to give the compound of formula (IA) or formula (IB), in which X₁ is oxo.

Preparation of certain compounds of formula (IB) may further involve conversion of the compound of formula (IB), in which A₁ is H, and R₄ and R₅ combine to form a double bond, into the compound of formula (IB) in which R₄ and A₁ combine to form oxo. In a non-limiting example, the enone in the compound of formula (IB), in which R₄ and R₅ combine to form a double bond can be converted into a C.12-C.13 epoxide using a nucleophilic peroxide agent, e.g., *t*-butyl hydroperoxide, which can then be converted into the compound of formula (IB), in which A₁ and R₄ combine to form oxo, using methods known in the art, e.g., by reacting with a bidentate phosphine ligand and a source of Pd(0) (see, e.g., Muzart, J., *Eur. J. Org. Chem.*, 4717-4741, 2011). Thus, the compound of formula (IB), in which A₁ is OP'', can be prepared. Other transformations may involve α -oxygenation to produce the compound of formula (VIB), in which R₅ is OP''.

As described herein, one of skill in the art can identify the sequence of reactions involving hydroxyl protecting group removing agents and Brønsted acids to convert the compound of formula (IB) into a halichondrin macrolide.

C.2-C.3 Bond-forming Macrocyclization


The macrocyclization reaction of the invention can be a carbon-carbon bond-forming reaction (e.g., catalytic Ring-Closing olefin Metathesis (RCM)) that provides a C.2-C.3 bond in a halichondrin macrolide. The general synthetic sequence including RCM that can be used to prepare the halichondrin macrolide is shown in Figure 2. As shown in Figure 2, the non-macrocyclic intermediate in the synthesis of the halichondrin macrolide can be a compound of formula (IIA):



or a salt or a tautomer thereof,

in which

- 5 each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:





where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

- 15 R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:





, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

5 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

10 k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

15 (i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or


(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

20 each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))- , where P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

P₅ is H or a hydroxyl protecting group.

25 In certain embodiments of the compound of formula (IIA), R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo. In particular embodiments of formula (IIA), X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))- , where P_Y is H. In some embodiments of formula (IIA), X₁ is oxo. In other embodiments of the compound of formula (IIA), R₃ and R₄ combine to form a bond, and R₅ is H.

30 A macrocyclic intermediate in the synthesis of a halichondrin macrolide can be a compound of formula (IB), which can be produced by reacting the compound of formula (IIA) with an olefin metathesis catalyst. The compound of formula (IB) has the following structure:



or a salt or a tautomer thereof,

where all variables are as defined for compound of formula (IIA).

5

The catalysts that can be used to convert the compound of formula (IIA) to the compound of formula (IB) can be those known in the art. Olefin metathesis catalysts include Ru-carbene complexes (e.g., Grubbs and Hoveyda-Grubbs catalysts). Olefin metathesis-competent catalysts that may be used in this reaction are known in the art (e.g., second generation Hoveyda-Grubbs-type catalysts, e.g., those in which the Ru-benzylidene moiety is modified to include electron-withdrawing and/or electron-donating groups).

10

The compound of formula (IIA) or formula (IB), in which X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_V))-$, where P_V is H, may be treated with an oxidizing agent capable of oxidizing hydroxyl to a carbonyl group (e.g., Dess-Martin periodinane or a dimethylsulfonium compound) to give the compound of formula (IIA) or formula (IB), in which X_1 is oxo.


15

The halichondrin macrolide can be prepared from the compound of formula (IB) as described herein.

C.3-C.4 Bond-forming Macrocyclization

20


The macrocyclization reaction of the invention can be a carbon-carbon bond-forming reaction (e.g., catalytic Ring-Closing olefin Metathesis (RCM)) that provides a C.3-C.4 bond in a halichondrin macrolide. The general synthetic sequence including RCM that can be used to prepare the halichondrin macrolide is shown in Figure 3. As shown in Figure 3, the non-macrocyclic intermediate in the synthesis of the halichondrin macrolide can be a compound of formula (IIIA):



or a salt or a tautomer thereof,

where

- 5 each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:





where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

- 15 R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:





, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

5 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

10 k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

15 (i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;


or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

20 each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_V))- , wherein P_V is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and P₅ is H or a hydroxyl protecting group.

25 In certain embodiments of the compound of formula (IIIA), R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo. In other embodiments of the compound of formula (IIIA), R₃ and R₄ combine to form a bond, and R₅ is H. In particular embodiments of formula (IIIA), X₁, together with the carbon atom to which it is attached, is -(CH(OP_V))- , where P_V is H. In some embodiments of formula (IIIA), X₁ is oxo.

30 A macrocyclic intermediate in the synthesis of a halichondrin macrolide can be a compound of formula (IIIB), which can be produced by reacting the compound of formula (IIIA) with an olefin metathesis catalyst. The compound of formula (IIIB) has the following structure:



or a salt or a tautomer thereof,

where all variables are as defined for the compound of formula (IIIA).

5

The catalysts that can be used to convert the compound of formula (IIIA) to the compound of formula (IIIB) can be those known in the art. Olefin metathesis catalysts include Ru-carbene complexes (e.g., Grubbs and Hoveyda-Grubbs catalysts). Olefin metathesis-competent catalysts that may be used in this reaction are known in the art (e.g., second generation Hoveyda-Grubbs-type catalysts, e.g., those in which the Ru-benzylidene moiety is modified to include electron-withdrawing and/or electron-donating groups).


The halichondrin macrolide can be prepared from the compound of formula (IIIB), as described for the synthesis involving the compound of formula (IB), as the C.3-C.4 double bond can undergo isomerization to give the compound of formula (IB) described herein upon exposure to basic (e.g., isomerization mediated by a hydroxyl protecting group removing agent, such as a fluoride source) or acidic (e.g., isomerization mediated by a Brønsted acid) conditions.

The compound of formula (IIIA) or formula (IIIB), in which X₁, together with the carbon atom to which it is attached, is $-(CH(OP_V))_n-$, where P_V is H, may be treated with an oxidizing agent capable of oxidizing hydroxyl to a carbonyl group to give the compound of formula (IIIA) or formula (IIIB), in which X₁ is oxo.

Preparation of the halichondrin macrolide may further involve conversion of the compound of formula (IIIB), in which A₁ is H, and R₄ and R₅ combine to form a double bond, into the compound of formula (VIB) in which R₄ and A₁ combine to form oxo. In a non-limiting example, the enone in the compound of formula (IIIB), in which R₄ and R₅ combine to form a double bond can be converted into a C.12-C.13 epoxide using a nucleophilic peroxide agent, e.g., *t*-butyl hydroperoxide, which can then be converted into the compound of formula (IIIB), in which A₁ and R₄ combine to form oxo, using methods known in the art, e.g., by reacting with a bidentate phosphine ligand and a source of Pd(0) (see, e.g., Muzart, J., *Eur. J. Org. Chem.*, 4717-4741, 2011). Thus, the halichondrin macrolide, in which A₁ is OP'', can be prepared. Other transformations may involve α-oxygenation to produce the halichondrin macrolide, in which R₅ is OP''.

C.12-C.13 Bond-forming Macrocyclization


The macrocyclization reaction of the invention can be a carbon-carbon bond-forming reaction (e.g., catalytic Ring-Closing olefin Metathesis (RCM)) that provides a C.12-C.13 bond in a halichondrin macrolide. The general synthetic sequence including RCM that can be used to prepare the halichondrin macrolide is shown in Figure 4. As shown in Figure 4, the non-macrocyclic intermediate in the synthesis of the halichondrin macrolide can be a compound of formula (IVA):



or a salt thereof,

where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

- 5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;
 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,
 NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and
 R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl,
 arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic
 10 radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

- (i) R₃ is H or a hydroxyl protecting group, R₄ is alkyl ether, and R₅ is H;
 (ii) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

15 or


- (iii) R₃ and R₄ combine to form a bond, and R₅ is H;

each P₄ and P₅ is independently H or a hydroxyl protecting group; and

R₆ is H, and P₆ is H or a hydroxyl protecting group; or R₆ and P₆ combine to form a double bond.

- 20 In certain embodiments of the compound of formula (IVA), P₅ is H. In particular embodiments of the
 compound of formula (IVA), R₃ and R₄ combine to form a bond, and R₅ is H. In some embodiments, R₆ is
 H, and P₆ is H or a hydroxyl protecting group. In other embodiments, R₆ and P₆ combine to form a double
 bond.

- 25 A macrocyclic intermediate in the synthesis of a halichondrin macrolide can be a compound of formula
 (IVB), which can be produced by reacting the compound of formula (IVA) with an olefin metathesis
 catalyst. The compound of formula (IVB) has the following structure:



or a salt thereof,

where all variables are as defined for the compound of formula (IVA).


5

The catalysts that can be used to convert the compound of formula (IVA) to the compound of formula (IVB) can be those known in the art. Olefin metathesis catalysts include Ru-carbene complexes (e.g., Grubbs and Hoveyda-Grubbs catalysts). Olefin metathesis-competent catalysts that may be used in this reaction are known in the art (e.g., second generation Hoveyda-Grubbs-type catalysts, e.g., those in which the Ru-benzylidene moiety is modified to include electron-withdrawing and/or electron-donating groups).

10

The compound of formula (IVB) (e.g., the compound of formula (IVB), in which R₆ and P₆ combine to form a double bond) can be converted to the halichondrin macrolide by a reaction with a Brønsted acid (if each of P₄ and P₅ is independently H or a hydroxyl protecting group capable of being deprotected by the Brønsted acid). Alternatively, if at least one P₄ or P₅ is a hydroxyl protecting group that is not capable of being deprotected by the Brønsted acid, the compound of formula (IVB) can be reacted with a hydroxyl protecting group removing agent to effect the removal of at least one hydroxyl protecting group, provided that the product of the reaction is the compound of formula (IVC):

15



20

or a salt thereof,

where each of A₁ and A₂ is H or OP'', where P'' is H or a hydroxyl protecting group; and the remaining variables are as defined for the compound of formula (IVA).

25

In a non-limiting example of the transformation of the compound of formula (IVB) to provide the compound of formula (IVC), at least one of P₄ and P₅ is a silyl, and the hydroxyl protecting group removing agent is a fluoride source.

- 5 The compound of formula (IVC) can then be converted to the halichondrin macrolide by a reaction with a Brønsted acid.


If, in the compound of formula (IVB) or (IVC), R₃ is a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond, treatment with a hydroxyl protecting group removing agent can provide the
10 compound of formula (IVB) or (IVC), in which R₃ and R₄ combine to form a bond, and R₅ is H.

Preparation of the compound of formula (IVC) from the compound of formula (IVB) may further involve reaction with a hydroxyl protecting group removing agent to convert P₆ in the compound of formula (IVB) to H and subsequent reaction with an oxidizing agent capable of converting a hydroxyl group into a
15 carbonyl (e.g., capable of oxidizing allylic alcohol into an enone). In a non-limiting example, the enone in the compound of formula (IVB) formed through the oxidation can then be converted into a C.12-C.13 epoxide using a nucleophilic peroxide agent, e.g., *t*-butyl hydroperoxide, which can then be converted into the compound of formula (IVB), in which A₁ and R₄ combine to form oxo, using methods known in the art, e.g., by reacting with a bidentate phosphine ligand and a source of Pd(0) (see, e.g., Muzart, J., *Eur. J. Org. Chem.*, 4717-4741, 2011). Thus, the compound of formula (IVC), in which A₁ is OP⁺, can be
20 prepared. Other transformations may involve α -oxygenation to produce the compound of formula (IVC), in which A₂ is OP⁺.

As described herein, one of skill in the art can identify the sequence of reactions involving the reactions
25 described above to convert the compound of formula (IVB) into a halichondrin macrolide with or without the isolation of the compound of formula (IVC).

C.15-C.16 Bond-forming Macrocyclization


The macrocyclization reaction of the invention can be a carbon-carbon bond-forming reaction (e.g.,
30 catalytic Ring-Closing olefin Metathesis (RCM)) that provides a C.15-C.16 bond in a halichondrin macrolide. The general synthetic sequence including RCM that can be used to prepare the halichondrin macrolide is shown in Figure 5. As shown in Figure 5, the non-macrocyclic intermediate in the synthesis of the halichondrin macrolide can be a compound of formula (VA):



or a salt or a tautomer thereof,

where

- 5 each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:





where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

- 15 R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:





, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

5 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

10 k is 0 or 1;

n is 0, 1, or 2;

(a1) R₃ is H or a hydroxyl protecting group, R₄ is alkyl ether, and R₅ is H;

(a2) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

15 or

(a3) R₃ and R₄ combine to form a bond, and R₅ is H;

(b1) A₁ and R₇ combine to form oxo, R₆ is H or a hydroxyl protecting group, and R₈ is H;

or

20 (b2) A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₆ is H or a hydroxyl protecting group, and R₇ and R₈ combine to form a double bond;

or

(ii) R₆ and R₇ combine to form a bond, and R₈ is H or OP'';

25 (c1) R₉ is H, and P₄ is H or a hydroxyl protecting group;

or

(c2) R₉ and P₄ combine to form a double bond;


R₁₀ is H or -CH₂X₁CH₂CH=CH₂, where X₁ is O, -C(R₁₁)₂-, or NP₆, and where each R₁₁ is independently H or -COOR₁₂, P₆ is an N-protecting group, and R₁₂ is alkyl; and

30 each P₅ is independently H or a hydroxyl protecting group.

In certain embodiments of the compound of formula (VA), R₆ and R₇ combine to form a bond, and R₈ is H. In particular embodiments of the compound of formula (VA), R₃ and R₄ combine to form a bond, and R₅ is H. In some embodiments of the compound of formula (VA), each P₅ is independently a hydroxyl

protecting group. In further embodiments of the compound of formula (VA), R₉ is H, and P₄ is a hydroxyl protecting group. In other embodiments, A₁ is H.

A macrocyclic intermediate in the synthesis of a halichondrin macrolide can be a compound of formula (VB), which can be produced by reacting the compound of formula (VA) with an olefin metathesis catalyst. The compound of formula (VB) has the following structure:




or a salt or a tautomer thereof,

where all variables are as defined for the compound of formula (VA).

The catalysts that can be used to convert the compound of formula (VA) to the compound of formula (VB) can be those known in the art. Olefin metathesis catalysts include Ru-carbene complexes (e.g., Grubbs and Hoveyda-Grubbs catalysts). Olefin metathesis-competent catalysts that may be used in this reaction are known in the art (e.g., second generation Hoveyda-Grubbs-type catalysts, e.g., those in which the Ru-benzylidene moiety is modified to include electron-withdrawing and/or electron-donating groups).


The compound of formula (VB) can be reacted with a hydroxyl protecting group removing agent and, optionally, an oxidizing agent (e.g., when R₉ is H in the compound of formula (VB)) capable of converting an alcohol to a carbonyl group (e.g., capable of converting an allylic alcohol to an enone) to afford the compound of formula (VBa):



or a salt thereof,

where A_2 is H or OP'' , and the remaining variables are as defined for the compound of formula (VA).

The compound of formula (VBa) can be reacted with a 1,4-reducing agent to furnish the compound of formula (VC):



or a salt thereof,


where all variables are as defined for the compound of formula (VBa).

The 1,4-reducing agents include copper (I) hydrides, which can be isolated (e.g., Stryker's reagent) or prepared *in situ* (e.g., from a copper (I) or copper (II) salt and a hydride source). Catalytic quantities of a copper salt (either copper (I) or copper (II) salt) in combination with stoichiometric or superstoichiometric quantities of a hydride source (e.g., a borohydride salt, borane, PMHS, or a hydrosilane (e.g., Ph_2SiH_2)).

A non-limiting example of the reaction conditions that can be used in the reaction sequence from the compound of formula (VB) to the compound of formula (VC) are described, e.g., in Baker et al., *Org. Lett.*, 10:289-292, 2008, the disclosure of which is incorporated herein by reference. Other metals can be used to catalyze 1,4-reduction, e.g., Ru, Pd, and Ir compounds.

Alternatively, if R_9 and P_4 combine to form a double bond in the compound of formula (VB), the reaction of the compound of formula (VB) with a hydroxyl protecting group removing agent can furnish the compound of formula (VC) directly.

If the compound of formula (VC) includes hydroxyl protecting groups as P_5 , these hydroxyl protecting groups can be removed with a hydroxyl protecting group removing agent to give the compound of formula (IVC):



(IVC)

or a salt thereof,

where all variables are as defined for the compound of formula (VBa).


5

The halichondrin macrolide can be prepared from the compound of formula (IVC) as described herein.

C.19-C.20 Bond-forming Macrocyclization

The macrocyclization reaction of the invention can be a carbon-carbon bond-forming reaction (e.g., Nozaki-Hiyama-Kishi reaction (NHK)) that provides a C.19-C.20 bond in a halichondrin macrolide. The general synthesis sequence including NHK that can be used to prepare the halichondrin macrolide is shown in Figure 6. As shown in Figure 6, the non-macrocyclic intermediate in the synthesis of the halichondrin macrolide can be a compound of formula (VIA):

10



(VIA),

15


or a salt or a tautomer thereof,

where

b designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or *b* designates (*S*)-stereogenic center, and Z is OR₆, where R₆ is a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:

20




(1),

where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

Y is iodide, bromide, or trifluoromethanesulfonate;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or


(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

5 and

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal.

10 In particular embodiments of the compound of formula (VIA), R_3 and R_4 combine to form a bond, and R_5 is H. In some embodiments of the compound of formula (VIA), both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal.

15 An intermediate in the synthesis of the halichondrin macrolide can be a compound of formula (VIB), which can be produced by subjecting the compound of formula (VIA) to Nozaki-Hiyama-Kishi reaction conditions (e.g., by reacting with a Cr(II) salt and a Ni(II) salt). The compound of formula (VIB) has the following structure:



or a salt or a tautomer thereof,

20 where

b designates (*R*)-stereogenic center, c designates (*S*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide;

or

b designates (*S*)-stereogenic center, c designates (*R*)-stereogenic center, and Z is OR_6 ,
25 where R_6 is a hydroxyl protecting group;

and the remaining variables are as defined for the compound of formula (VIA).

Nozaki-Hiyama-Kishi reaction conditions that may be used to prepare the compound of formula (VIB) from the compound of formula (VIA) can be those known in the art. Nozaki-Hiyama-Kishi reaction on the
30 compound of formula (VIA) can include reacting the compound of formula (VIA) with a Cr(II) salt and a Ni(II) salt. Ancillary ligands can be used in combination with the metal salts. In a non-limiting example, a substituted 1,10-phenanthroline can be used in combination with a Ni(II) salt. Chiral ancillary ligands can be used to render the reaction stereoselective. In a non-limiting example, chiral N-(dihydrooxazolyl-

phenyl)-sulfonamides can be used with a Cr(II) salt to control the stereochemistry of the carbonyl carbon, to which a vinyl nucleophile is added in the course of Nozaki-Hiyama-Kishi reaction.

The compound of formula (VIB) can be converted to the halichondrin macrolide using the step of nucleophilic ring-closing of the compound of formula (VIB).

In particular, the compound of formula (VIB) (in which Z is a sulfonate, chloride, bromide, or iodide; *b* designates (*R*)-stereogenic center; and *c* designates (*S*)-stereogenic center) can be converted to the halichondrin macrolide directly, e.g., upon isolation from the Nozaki-Hiyama-Kishi reaction mixture (e.g., by treatment with a base) or by contacting a mixture containing the product of the Nozaki-Hiyama-Kishi reaction with silica gel. Additionally, the compound of formula (VIB) can be reacted with a hydroxyl protecting group removing agent (before or after the step of nucleophilic ring-closing) if, e.g., each of P₄ is a hydroxyl protecting group, optionally, with a subsequent reaction with a Brønsted acid to form a compound, in which both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal.


Alternatively, if, in the compound of formula (VIB), *b* designates (*S*)-stereogenic center, *c* designates (*R*)-stereogenic center, and Z is OR₆, where R₆ is a hydroxyl protecting group, the step of nucleophilic ring-closing of the compound of formula (VIB) can be effected after the hydroxyl group attached to the carbon atom designated with *c* is converted into a leaving group (e.g., a sulfonate by a reaction with a sulfonyl electrophile, such as a sulfonyl chloride or a sulfonyl anhydride). In these embodiments, the step of nucleophilic ring-closing can be carried out by, e.g., first, reacting the compound of formula (VIB), after the conversion of the hydroxyl into the leaving group, with a hydroxyl protecting group removing agent to convert R₆ into H and then reacting the product with a base.

Preparation of certain compounds of formula (VIB) may further involve conversion of the compound of formula (VIB), in which A₁ is H, and R₄ and R₅ combine to form a double bond, into the compound of formula (VIB) in which R₄ and A₁ combine to form oxo. In a non-limiting example, the enone in the compound of formula (VIB), in which R₄ and R₅ combine to form a double bond can be converted into a C.12-C.13 epoxide using a nucleophilic peroxide agent, e.g., *t*-butyl hydroperoxide, which can then be converted into the compound of formula (VIB), in which A₁ and R₄ combine to form oxo, using methods known in the art, e.g., by reacting with a bidentate phosphine ligand and a source of Pd(0) (see, e.g., Muzart, J., *Eur. J. Org. Chem.*, 4717-4741, 2011). Thus, the compound of formula (VIB), in which A₁ is OPⁿ, can be prepared. Other transformations may involve α-oxygenation to produce the compound of formula (VIB), in which R₅ is OPⁿ.

One of skill in the art will recognize that any unprotected hydroxyl groups can be protected with a hydroxyl protecting group, if the unprotected hydroxyl groups are not intended for reacting in the step of nucleophilic ring-closing.


C.26-C.27 Bond-forming Macrocyclization

The macrocyclization reaction of the invention can be a carbon-carbon bond-forming reaction (e.g., Nozaki-Hiyama-Kishi reaction (NHK)) that provides a C.26-C.27 bond in a halichondrin macrolide. The general synthesis sequence including NHK that can be used to prepare the halichondrin macrolide is shown in Figure 7. As shown in Figure 7, the non-macrocyclic intermediate in the synthesis of the halichondrin macrolide can be a compound of formula (VIIA):



where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,
 NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and
 R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl,
 arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic
 10 radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

Y is iodide, bromide, or trifluoromethanesulfonate; and

(a1) R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, each

15 P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is
 attached, forms a carbonyl or -(CH(OR₈))-; where R₈ is H or a hydroxyl protecting group;

or

(a2) R₃ and R₄ combine to form a bond, R₅ is H, and

20 each P₄ is independently H or a hydroxyl protecting group, and X₁, together with
 the carbon to which it is attached, forms a carbonyl or -(CH(OR₈))-;

or

both P₄ groups and X₁, together with the atoms to which each is attached,
 combine to form ketal;


(b1) Z is chloride, bromide, or iodide, and R₆ and R₇ combine to form a bond;

25 or

(b2) Z and R₇ combine to form a double bond, and R₆ is H or a hydroxyl protecting group.

In particular embodiments of the compound of formula (VIIA), Z and R₇ combine to form a double bond, and R₆ is a hydroxyl protecting group. In some embodiments of the compound of formula (VIIA), each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$. In certain embodiments of the compound of formula (VIIA), R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond.

An intermediate in the synthesis of the halichondrin macrolide can be a compound of formula (VIIB), which can be produced by subjecting the compound of formula (VIIA) to Nozaki-Hiyama-Kishi reaction conditions (e.g., by reacting with a Cr(II) salt and a Ni(II) salt). The compound of formula (VIIB) has the following structure:




or a salt thereof,

where each of P₅ and R₆ is independently H or a hydroxyl protecting group, and the remaining variables are as defined for the compound of formula (VIIA).

Nozaki-Hiyama-Kishi reaction conditions that may be used to prepare the compound of formula (VIIB) from the compound of formula (VIIA) can be those known in the art. Nozaki-Hiyama-Kishi reaction on the compound of formula (VIIA) can include reacting the compound of formula (VIIA) with a Cr(II) salt and a Ni(II) salt. Ancillary ligands can be used in combination with the metal salts. In a non-limiting example, a substituted 1,10-phenanthroline can be used in combination with a Ni(II) salt. Chiral ancillary ligands can be used to render the reaction stereoselective. In a non-limiting example, chiral N-(dihydrooxazolyl-phenyl)-sulfonamides can be used with a Cr(II) salt to control the stereochemistry of the carbonyl carbon, to which a vinyl nucleophile is added in the course of Nozaki-Hiyama-Kishi reaction.

The compound of formula (VIIB) can be converted to a compound of formula (VIIC) through the step of nucleophilic ring-closing of the compound of formula (VIIB), where the compound of formula (VIIC) has the following structure:



or a salt or a tautomer thereof,

where each P_4 is H or a hydroxyl protecting group, and X_1 is oxo; or both P_4 groups and X_1 ,
 5 together with the atoms to which each is attached, combine to form a ketal;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a
 10 double bond;

or


(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

and the remaining variables are as defined for the compound of formula (VIIA).

15 The step of nucleophilic ring-closing of the compound of formula (VIIB) can be effected after the hydroxyl group produced by the NHK reaction is converted into a leaving group (e.g., a sulfonate by a reaction with a sulfonyl electrophile, such as a sulfonyl chloride or a sulfonyl anhydride). In these embodiments, the step of nucleophilic ring-closing can be carried out by reacting the compound of formula (VIB), in which R_6 is H, after the conversion of the hydroxyl into the leaving group, with a strong base (e.g., alkoxide).

20 Preparation of the compound of formula (VIIC) can involve Vasella fragmentation prior to the step of nucleophilic ring-closing, if Z and R_7 combine to form a double bond, and R_6 is a hydroxyl protecting group in the compound of formula (VIIB).

25 The compound of formula (VIIC) can be reacted with a hydroxyl protecting group removing agent to give a compound of formula (VIID):



or a salt thereof,

where each of A_1 and A_2 is independently H or OP'' , and the remaining variables are as defined

5 for the compound of formula (VIIC).

The compound of formula (VIID) can be converted to the halichondrin macrolide by reacting with a Brønsted acid (e.g., a pyridinium salt, such as pyridinium *p*-toluene sulfonate).

10 In some embodiments, the preparation of the halichondrin macrolide from the compound of formula (VIIA) can further involve an oxidation reaction (e.g., contacting the compound of formula (VIIA), (VIIB), or (VIIC), in which X_1 is $-(CH(OR_8))-$ and R_8 is H, with an oxidizing agent capable of oxidizing an alcohol to a carbonyl group).

15 Preparation of the compound of formula (VIID) from the compound of formula (VIIB) may further involve conversion of the compound of formula (VIIB), in which R_4 and R_5 combine to form a double bond into the compound of formula (VIIC) in which R_4 and A_1 combine to form oxo. In a non-limiting example, the enone in the compound of formula (VIIB), in which R_4 and R_5 combine to form a double bond can be converted into a C.12-C.13 epoxide using a nucleophilic peroxide agent, e.g., *t*-butyl hydroperoxide,

20 which can then be converted into the compound of formula (VIIC), in which A_1 and R_4 combine to form oxo, using methods known in the art, e.g., by reacting with a bidentate phosphine ligand and a source of Pd(0) (see, e.g., Muzart, J., *Eur. J. Org. Chem.*, 4717-4741, 2011). Thus, the compound of formula (VIID), in which A_1 is OP'' , can be prepared. Other transformations may involve α -oxygenation to produce the compound of formula (VIID), in which A_2 is OP'' .

25 One of skill in the art will recognize that any unprotected hydroxyl groups can be protected with a hydroxyl protecting group, if the unprotected hydroxyl groups are not intended for reacting in the step of nucleophilic ring-closing.

30 **Synthesis of Non-macrocyclic intermediates**

The compounds of formula (IA), (IIA), (IIIA), (IVA), (VA), (VIA), and (VIIA) can be prepared using methods and intermediates disclosed, e.g., in U.S. Patent Nos. 5,338,865 and 5,436,238; in International Patent application No. PCT/US2014/063960; and in Towle et al., *Annual Meeting of the American Association for*

Cancer Research, April 6-10, 2002, 5721; Wang et al., *Bioorg. Med. Chem. Lett.*, 10:1029-1032, 2000; Aicher et al., *J. Am. Chem. Soc.*, 114:3162-3164, 1992; Ueda et al., *J. Am. Chem. Soc.*, 136:5171-5176; and Yamamoto et al., *J. Am. Chem. Soc.*, 134:893-896, 2012; each of which is incorporated herein by reference in its entirety.


5

Compound (IA)

The compound of formula (IA) can be prepared using methods described in the references incorporated by reference above. In particular, the compound of formula (IA) can be prepared from compound (IC), in which P_x is H, and compound (ID), e.g., using esterification reaction conditions (e.g., Shiina esterification reaction conditions (e.g., using MNBA) or Yamaguchi esterification reaction conditions, e.g., the reaction conditions described in Aicher et al., *J. Am. Chem. Soc.*, 114:3162-3164, 1992).

10

The compound of formula (IC) has the following structure




(IC),

15

where

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:

20



(1),

where


L is $-(CH(OP_2))_n-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently optionally substituted alkyl or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:

25

30



E is optionally substituted alkyl or optionally substituted alkoxy;

- 5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;
 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,
 NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and
 R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl,
 arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic
 10 radical-alkyl;

P_X is H or a hydroxyl protecting group;

k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

- 15 or

A₁ is H or OP'', and:

- (i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a
 double bond;

or

- 20 (ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

R₆ is OP'' and R₇ is H, or R₆ and R₇ combine to form a double bond;

each P₄ is independently a hydroxyl protecting group, and X₁ is oxo or X₁, together with the
 carbon atom to which it is attached, is -(CH(OP_Y))- , wherein P_Y is H or a hydroxyl protecting group; or
 both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

- 25 P₅ is a hydroxyl protecting group; and

where each P'' is independently H or a hydroxyl protecting group.

The compound of formula (IC), in which X_1 is oxo, can be prepared by reacting an oxidizing agent capable of oxidizing hydroxyl to a carbonyl group (e.g., Dess-Martin periodinane) with the compound of formula (IC), in which X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, where P_Y is H.


5

The compound of formula (ID) has the following structure: $(RO)_2P(O)CH_2COOH$, where each R is independently optionally substituted alkyl or optionally substituted aryl.

The preparation of the compound of formula (IA) from the compound of formula (IC) and the compound of formula (ID) may involve a hydroboration/oxidation reaction to furnish the aldehyde group in the compound of formula (IA) after the esterification reaction, if R_6 and R_7 combine to form a double bond in the compound of formula (IC). Alternatively, the preparation of the compound of formula (IA) from the compound of formula (IC) and the compound of formula (ID) may involve reacting the esterification product with a hydroxyl protecting group removing agent and then reacting the product with an oxidizing agent capable of converting a hydroxyl group into a carbonyl to furnish the aldehyde group in the compound of formula (IA), if, in the compound of formula (IC) R_6 is OP and R_7 is H.


The compound of formula (IC) can be prepared from a compound of formula (IE) and the compound of formula (IF), e.g., using Nozaki-Hiyama-Kishi reaction conditions, as described herein. The hydroxyl group used in the subsequent esterification reaction can be deprotected using a hydroxyl protecting group removing agent. Specifically, the preparation of the compound of formula (IC), in which P_X is H, may further involve treating the compound of formula (IC), in which P_X is a hydroxyl protecting group, with a hydroxyl protecting group removing agent.

The compound of formula (IE) has the following structure:



where

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




(1),

where

L is $-(CH(OP_2))_n-$, or $-C(O)-$; R_1 is H, or R_1 and P_1 combine to form a bond;


R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently optionally substituted alkyl or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:

each P' is independently a hydroxyl protecting group; E is optionally substituted alkyl or optionally substituted alkoxy; G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

 P_X is a hydroxyl protecting group; k is 0 or 1; n is 0, 1, or 2; and X_1 is oxo.

The compound of formula (IF) has the following structure:



where

- 5 each R₃, P₄, and P₅ is independently a hydroxyl protecting group;
 A₁ is H or OP^{''}, where P^{''} is a hydroxyl protecting group;
 R₆ is OP₅, and R₇ is H, or R₆ and R₇ combine to form a double bond;
 and
 Y is chloro, bromo, iodo, trifluoromethanesulfonate, or trialkylsilane.

10

The compound of formula (IE), in which X₁ is oxo, can be reacted with the compound of formula (IF) under the Nozaki-Hiyama-Kishi reaction conditions, as described herein.

The compound of formula (IE) can be accessed, e.g., from the compound of formula (VIII D).

15

The compound of formula (IF) can be prepared using methods known in the art, e.g., those described in WO 2015/066729.

20


The preparation of compound (IA) may further involve reactions deprotecting protected hydroxyl groups or protecting unprotected hydroxyl groups using methods described herein.

Compound (IIA)

25


The compound of formula (IIA) can be prepared using methods described in the references incorporated by reference above. In particular, the compound of formula (IIIA) can be prepared from acrylic acid a compound of formula (IIB), e.g., using esterification reaction conditions (e.g., Shiina esterification reaction conditions (e.g., using MNBA) or Yamaguchi esterification reaction conditions, e.g., the reaction conditions described in Aicher et al., *J. Am. Chem. Soc.*, 114:3162-3164, 1992).

The compound of formula (IIB) has the following structure:



where

- 5 each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

- 15 R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently optionally substituted alkyl or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

- 5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;
 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,
 NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and
 R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl,
 arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic
 10 radical-alkyl;

P_X is H or a hydroxyl protecting group;

k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

- 15 or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

- (i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a
 double bond;

or

- 20 (ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

each P₄ is independently a hydroxyl protecting group, and X₁ is oxo or X₁, together with the
 carbon atom to which it is attached, is -(CH(OP_Y))- , wherein P_Y is H or a hydroxyl protecting group; or
 both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

P₅ is a hydroxyl protecting group.

25

The compound of formula (IIB), in which X₁ is oxo, can be prepared by reacting an oxidizing agent
 capable of oxidizing hydroxyl to a carbonyl group (e.g., Dess-Martin periodinane or a dimethylsulfonium


compound) with the compound of formula (IIB), in which X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_V))-$, where P_V is H.

The compound of formula (IIB), in which P_X is H, can be prepared by reacting a hydroxyl protecting group removing agent with the compound of formula (IIB), in which P_X is a hydroxyl protecting group.

In non-limiting examples, the compound of formula (IIB) can be prepared from a compound of formula (IE) and a compound of formula (IIC) (e.g., via the compound of formula (IICa)).

The preparation of the compound of formula (IIB) may further involve hydroboration/oxidation reaction with subsequent olefination (e.g., Wittig reaction (e.g., with Ph_3PCH_2) to provide a terminal alkene). For example, compound of formula (IIC) may be subjected to hydroboration/oxidation sequence to produce an aldehyde, which upon treatment with Ph_3PCH_2 , produces the compound of formula (IICa).

The compound of formula (IIC) has the following structure:




where

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group;

each R_3 , P_4 , and P_5 is independently a hydroxyl protecting group; and

Y is chloro, bromo, iodo, or trifluoromethanesulfonate.

The compound of formula (IICa) has the following structure:



where all variables are as defined for formula (IIC).

The compound of formula (IE), in which X_1 is oxo, can be reacted with the compound of formula (IIC) under the Nozaki-Hiyama-Kishi reaction conditions, as described herein.

The compound of formula (IIC) can be prepared using methods known in the art, e.g., those described in WO 2015/066729.


The preparation of compound (IIA) may also involve reactions deprotecting protected hydroxyl groups or protecting unprotected hydroxyl groups using methods described herein.

Compound (IIIA)

- 5 The compound of formula (IIIA) can be prepared using methods described in the references incorporated by reference above. In particular, the compound of formula (IIIA) can be prepared from but-3-enoic acid a compound of formula (IC), in which R₆ and R₇ combine to form a double bond, e.g., using esterification reaction conditions (e.g., Shiina esterification reaction conditions (e.g., using MNBA) or Yamaguchi esterification reaction conditions, e.g., the reaction conditions described in Aicher et al., *J. Am. Chem.*
- 10 *Soc.*, 114:3162-3164, 1992). The preparation of compound (IIIA) may also involve reactions deprotecting protected hydroxyl groups or protecting unprotected hydroxyl groups using methods described herein.

Compound (IVA)


- The compound of formula (IVA) can be prepared using methods described in the references incorporated
- 15 by reference above. In particular, the compound of formula (IVA) can be prepared from a compound of formula (IVD) and a compound of formula (IVE), e.g., using esterification reaction conditions (e.g., Shiina esterification reaction conditions (e.g., using MNBA) or Yamaguchi esterification reaction conditions, e.g., the reaction conditions described in Aicher et al., *J. Am. Chem. Soc.*, 114:3162-3164, 1992).
- 20 The compound of formula (IVD) has the following structure:



where


- (i) R₃ is a hydroxyl protecting group, R₄ is alkyl ether, and R₅ is H;
- 25 (ii) R₃ is a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;
- or
- (iii) R₃ and R₄ combine to form a bond, and R₅ is H;
- each P₄ and P₅ is independently a hydroxyl protecting group.

The compound of formula (IVE) has the following structure:



where

each of D and D' is independently H, optionally substituted alkyl, or OP₁; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:





where

L is $-(CH(OP_2))-$ or $-C(O)-$;

R₁ and P₁ combine to form a bond; or R₁ is H, and P₁ is alkyl;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently optionally substituted alkyl or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:





, where each P' is independently a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;


5 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

10 k is 0 or 1;

n is 0, 1, or 2; and

R₆ is H, and P₆ is H or a hydroxyl protecting group; or R₆ and P₆ combine to form a double bond.

15 The compound (IVE) can be prepared using methods known in the art or using the allene-Prins method described herein. In particular, after the allene-Prins reaction and allylic reduction, the substrate, in which




R₃ is can be subjected to a reaction sequence involving reactions with oxidizing agents capable of oxidizing an alcohol to a carbonyl and vinyl addition to carbonyl.

20 The preparation of compound (IVA) may also involve reactions deprotecting protected hydroxyl groups or protecting unprotected hydroxyl groups using methods described herein.

Compound (VA)

25 The compound of formula (VA) can be prepared using methods described in the references incorporated by reference above. In particular, the compound of formula (VA) can be prepared from a compound of formula (VE) and a compound of formula (VD), in which P_x is H, e.g., using esterification reaction conditions (e.g., Shiina esterification reaction conditions (e.g., using MNBA) or Yamaguchi esterification reaction conditions, e.g., the reaction conditions described in Aicher et al., *J. Am. Chem. Soc.*, 114:3162-3164, 1992).

The compound of formula (VE) has the following structure:




(VE),

where

- (a1) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;
 - (a2) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;
 - or
 - (a3) R_3 and R_4 combine to form a bond, and R_5 is H;
 - (b1) R_6 is a hydroxyl protecting group, and R_7 and R_8 combine to form a double bond;
 - or
 - (b2) R_6 and R_7 combine to form a bond, and R_8 is H;
 - (c1) R_9 is H, and P_4 is a hydroxyl protecting group;
 - or
 - (c2) R_9 and P_4 combine to form a double bond;
- each P_5 is independently a hydroxyl protecting group; and
 A_1 is H or OP'' , where P'' is a hydroxyl protecting group;.


The compound of formula (VD) has the following structure:



(VD),

where

- each of D and D' is independently H, optionally substituted alkyl, or OP_1 ; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




(1),

where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 and P_1 combine to form a bond; or R_1 is H, and P_1 is a hydroxyl protecting group or alkyl;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently optionally substituted alkyl or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N -protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

P_X is H or a hydroxyl protecting group;


k is 0 or 1;

n is 0, 1, or 2; and

R_{10} is H or $-CH_2X_1CH_2CH=CH_2$, where X_1 is O, $-C(R_{11})_2-$, or NP_6 , and where each R_{11} is independently H or $-COOR_{12}$, P_6 is an N -protecting group, and R_{12} is alkyl.

The compound of formula (VD), in which P_X is H, can be prepared by reacting a hydroxyl protecting group removing agent with the compound of formula (VD), in which P_X is a hydroxyl protecting group.

The compound of formula (VD) can be prepared using methods known in the art or using the allene-Prins method described herein. For example, the compound of formula (VD), in which R_{10} is $-CH_2X_1CH_2CH=CH_2$, can be prepared from the compound of formula (VF):



where R_G is $-\text{CHO}$, $-\text{CH}=\text{CH}-\text{COOR}_H$, where R_H is H or a hydroxyl protecting group, and all remaining variables are as defined for formula (VD).

5

Thus, in a non-limiting example, the compound of formula (VF), in which R_G is $-\text{CHO}$, may be subjected to Horner-Wadsworth-Emmons reaction conditions (e.g., with $(\text{EtO})_2\text{P}(\text{O})-\text{CH}_2-\text{COOR}_H$) to produce the compound of formula (VF), in which R_G is $-\text{CH}=\text{CH}-\text{COOR}_H$. The latter compound may be treated with a 1,2-reducing agent (e.g., DIBAL) followed by a quench with allyl halide (e.g., allyl bromide) to give the compound of formula (VD).

10

The compound of formula (VF) may be prepared from the compound of formula (VIID). In a non-limiting example, the compound of formula (VIID), in which R_3 is $-\text{CH}_2-\text{OP}_5$, and P_5 is a hydroxyl protecting group, may be treated with a hydroxyl protecting group removing agent (e.g., if P_5 is Piv, the hydroxyl protecting group removing agent may be DIBAL), followed by an oxidizing agent capable of oxidizing hydroxyl to a carbonyl group (e.g., Dess-Martin periodinane), to give the compound of formula (VF), in which R_G is $-\text{CHO}$.

15

The preparation of compound (VA) may also involve reactions deprotecting protected hydroxyl groups or protecting unprotected hydroxyl groups using methods described herein.


20

Compound (VIA)

The compound of formula (VIA) can be prepared using methods described in the references incorporated by reference above. In particular, the compound of formula (VIA) can be prepared from a compound of formula (VIC) and a compound of formula (VID), e.g., using esterification reaction conditions (e.g., Shiina esterification reaction conditions (e.g., using MNBA) or Yamaguchi esterification reaction conditions, e.g., the reaction conditions described in Aicher et al., *J. Am. Chem. Soc.*, 114:3162-3164, 1992).

25

The compound of formula (VIC) has the following structure:



(VIC),

or a tautomer thereof

where

b designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or *b* designates (*S*)-stereogenic center, and Z is OR₆, where R₆ is a hydroxyl protecting group;

Y is iodide, bromide, or trifluoromethanesulfonate;


(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H;

each P₄ is independently a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))-, wherein P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal.


The compound of formula (VID) has the following structure:



(VID),

where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




where

L is -(CH(OP₂))-, -(C(OH)(OP₂))-, or -C(O)-;

R₁ and P₁ combine to form a bond; or R₁ is absent or H, and P₁ is absent, a hydroxyl protecting group, or alkyl;

R₂ is H or -(CH₂)_nOP₃, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each

is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$,

10 $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

15 n is 0, 1, or 2; and


P_5 is a hydroxyl protecting group

The compound of formula (VIC) can be prepared using methods described in WO 2015/066729.

20 The preparation of compound (VIA) may also involve reactions deprotecting protected hydroxyl groups or protecting unprotected hydroxyl groups using methods described herein.

Compound (VIIA)


The compound of formula (VIIA) can be prepared using methods described in the references incorporated
 25 by reference above. In particular, the compound of formula (VIIA) can be prepared from a compound of formula (VIIIE):



or a salt or a tautomer thereof,

where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:





where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:





, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

Y is iodide, bromide, or trifluoromethanesulfonate;

P₅ is H or a hydroxyl protecting group; and

(a1) R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, each

P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or -(CH(OR₈))- , where R₈ is H or a hydroxyl protecting group;

or

(a2) R₃ and R₄ combine to form a bond, R₅ is H, and

each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or -(CH(OR₈))-;

or

both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal;


(b1) Z is chloride, bromide, or iodide, and R₆ and R₇ combine to form a bond;

or

(b2) Z and R₇ combine to form a double bond, and R₆ is H or a hydroxyl protecting group.

The preparation of the compound of formula (VIIA) can involve oxidizing the compound of formula (VIIE) (e.g., the compound of formula (VIIE), in which P₅ is H or a hydroxyl protecting group), e.g., by reacting with an oxidizing agent capable of oxidizing hydroxyl to a carbonyl group. The mixture containing the oxidizing agent can also act as a hydroxyl protecting group removing agent.

The compound of formula (VIIE) can be formed by reacting a compound of formula (VIIF) with the compound of formula (VIIG). The compound of formula (VIIF) has the following structure:



or a salt or a tautomer thereof,

where

5 Y is iodide, bromide, or trifluoromethanesulfonate; and

(a1) R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$, where R₈ is H or a hydroxyl protecting group;

or

10 (a2) R₃ and R₄ combine to form a bond, R₅ is H, and

each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$;

or


15 both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal;

(b1) Z is chloride, bromide, or iodide, and R₆ and R₇ combine to form a bond;

or


(b2) Z and R₇ combine to form a double bond, and R₆ is H or a hydroxyl protecting group.

20 The compound of formula (VIIG) has the following structure:



or a salt thereof,

25 each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2; and


P₅ is H or a hydroxyl protecting group.

Allene-Prins Reaction in the Preparation of Fragments (e.g., C.14-C.38 Fragment) of Halichondrin

The invention further features a method of preparing a fragment of halichondrin macrolide (e.g., C.20-C.35, C.20-C.38, C.16-C.35, C.16-C.38, C.14-C.35, and C.14-C.38 fragments), which can be an intermediate in the synthesis of halichondrin macrolide or a salt thereof. The advantages of the process relative to the current synthesis of C.14-C.35, C.14-C.38, C.16-C.35, C.16-C.38, C.20-C.35, and C.20-C.38 fragments of halichondrin include non-metal mediated assembly that does not employ a C.23 leaving group. Additionally, the process obviates the need for a chiral ligand. The method involves performing an allene-Prins reaction on a compound of formula (VIII A), a compound of formula (VIII B), and R_4OH to afford the intermediate in the synthesis of halichondrin macrolide,

where R_4 is an optionally substituted acyl;


where the compound of formula (VIII A) has the following structure:



or a salt thereof,

where

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , where P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:





where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, where each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

- 5 G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;
 each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A,
 NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and
 R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl,
 arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic
 10 radical-alkyl;


k is 0 or 1;

n is 0, 1, or 2;


X is O, or X, together with the carbon atom to which it is attached, forms -(C(OPz)₂)-, wherein
 each Pz is independently optionally substituted alkyl or optionally substituted aryl, or both Pz combine to
 15 form optionally substituted alkylene or optionally substituted arylene; and


P₄ is H or a hydroxyl protecting group;

where the compound of formula (VIII B) has the following structure:



20 where






where P₅ is H or a hydroxyl protecting group; each P₆ is independently a hydroxyl protecting group, or both P₆ groups, together with the atoms to which each is attached, combine to form a cyclic protected diol; and R₅ is H or -

CH₂X₁CH₂CH=CH₂, where X₁ is O, -CH₂-, or NP₇, where P₇ is a sulfonyl;

and where the intermediate is a compound of formula (VIIIc):




or a salt thereof,

where *a* designates (*R*)-stereogenic center or (*S*)-stereogenic center.

In some embodiments, *a* designates (*S*)-stereogenic center. In certain embodiments, one and only one of *D* and *D'* is optionally substituted alkyl or OP₁. In other embodiments, one and only one of *D* and *D'* is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group. In yet other embodiments, *A* is a group of formula (1). In still other embodiments, *L* is $-(CH(OP_2))-$. In particular embodiments, *R*₁ and P₁ combine to form a bond. In certain embodiments, *G* is O. In further embodiments, *E* is optionally substituted alkyl (e.g., methyl). In some embodiments, *k* is 1. In particular embodiments, *R*₂ is $-(CH_2)_nOP_3$ (e.g., *n* is 1 or 2). In further embodiments, at least one of P₂ and P₃ is a hydroxyl protecting group.

The reaction conditions that can be used to prepare the compound of formula (VIII C) using allene-Prins reaction are those known in the art for Prins reaction and can include reacting the compound of formula (VIII A), the compound of formula (VIII B), and R_4OH with a Lewis acid (e.g., boron trifluoride or a solvate thereof).


The compound of formula (VIIC) can be subjected to allylic reducing conditions to afford a compound of formula (VIID):



or a salt thereof,

where

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, where P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




where

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, where R_N is H, an N-protecting group, or optionally substituted alkyl;


each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, where each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

and

P₄ is H or a hydroxyl protecting group; and



where P₅ is H or a hydroxyl protecting group; each P₆ is independently a hydroxyl protecting group, or both P₆ groups, together with the atoms to which each is attached, combine to form a cyclic protected diol; and R₅ is is H or $-\text{CH}_2\text{X}_1\text{CH}_2\text{CH}=\text{CH}_2$, where X₁ is

10 O, $-\text{CH}_2-$, or NP₇, where P₇ is a sulfonyl.

Preparation of a compound of formula (VIIIB) is described in detail in International Patent Application No. PCT/US2014/063960, and the disclosure of the preparation is incorporated herein by reference in its entirety.

Compounds

The present invention also provides compounds that can be used in the synthesis of a halichondrin macrolide, e.g., the compounds of formula (IA), (IB), (IC), (IE), (IF), (IIA), (IIB), (IIC), (IICa), (IIIA), (IIIB), (IVA), (IVB), (IVC), (IVD), (IVE), (VA), (VB), (VBa), (VC), (VIA), (VIB), (VIC), (VIIA), (VIIB), (VIIC), (VIID), (VIIIE), (VIIF), (VIIG), (VIIIA), (VIIIB), (VIIC), or (VIID).

20

Oxidizing Agents Capable of Converting an Alcohol to a Carbonyl Group

Oxidizing agents capable of converting an alcohol to a carbonyl group are known in the art. Non-limiting examples of these oxidizing agents include Dess-Martin periodinane, TEMPO (in the presence of bleach or BAIB), a dimethylsulfonium compound (e.g., dimethylchlorosulfonium chloride), aluminum trialkoxide with an excess of a ketone (e.g., acetone), and catalytic tetrapropylammonium perruthenate (TPAP) (in the presence of *N*-methylmorpholine oxide). The dimethylsulfonium compound can be prepared *in situ* under the conditions known for Parikh-Doering oxidation, Swern oxidation, Corey-Kim oxidation, or Pfitzner-Moffatt oxidation. An oxidation reaction of an alcohol to a carbonyl group (e.g., a ketone) can be performed using aluminum trialkoxide and an excess of a ketone (e.g., acetone) under the conditions known in the art for Oppenauer oxidation. Allylic and benzylic alcohols can also be oxidized with MnO₂.

25

30

Reducing Agents

Reducing agents that can be used in the methods of the invention are those known in the art. A reducing agent can be an electron-transfer reducing agent, a metal hydride, or a metalloid hydride. Non-limiting examples of electron-transfer reducing agent include alkali metals in oxidation state (0), alkali earth metals in oxidation state (0), alkali arenides, lanthanide (II) salts (e.g., SmI₂), Zn(0), Fe(0), and Mn(0).

35

Non-limiting examples of metal hydrides and metalloid hydrides include boron hydride compounds (e.g., NaBH_4 , LiBH_4 , $\text{LiB}(\text{Et})_3\text{H}$, selectrides (e.g., L-selectride), and boranes (e.g., 9-BBN and alpine borane)), aluminum hydride compounds (e.g., LiAlH_4 , Red-Al®, and alanes (e.g., DIBAL)), hydrosilanes (e.g., PMHS and Ph_2SiH_2), hydrostannanes (e.g., Bu_3SnH), copper hydride complexes (e.g., Stryker's reagent),
5 palladium hydride complexes, platinum hydride complexes, iridium hydride complexes, rhodium hydride complexes, and ruthenium hydride complexes. Reducing agents can be formed *in situ*, e.g., a copper hydride complex can be formed by a reaction of a copper salt with, e.g., a boron hydride compound or a hydrosilane. Thus, some reducing reagents (e.g., boron hydride compounds, hydrosilanes, and hydrostannanes) can be used in combination with a catalytic quantity of a metal salt (e.g., Cu, Pd, Pt, Ir,
10 Rh, or Ru salt). Alternatively, catalytic reducing agents can be metal salts (e.g., aluminum isopropoxide or a ruthenium complex) in combination with an alcohol, which undergo transfer hydrogenation of carbonyl-containing compounds without intermediacy of a metal hydride. Non-limiting examples of transfer hydrogenation reactions include Meerwein-Ponndorf-Verley reduction (e.g., using aluminum isopropoxide/isopropanol) and Ru-catalyzed transfer hydrogenation (e.g., Hashiguchi et al., *J. Am. Chem.*
15 *Soc.*, 117:7562-7563, 1995).

When a substrate is an α,β -unsaturated carbonyl compound (e.g., an α,β -enone), a reducing agent can be a 1,2-reducing agent or a 1,4-reducing agent. For example, a reaction between an α,β -unsaturated carbonyl compound and a 1,2-reducing agent can afford, e.g., an allylic alcohol (or an allylic amine, if the
20 starting compound is an enamide), whereas a reaction between an α,β -unsaturated carbonyl compound and a 1,4-reducing agent can afford an α,β -saturated compound and can leave the carbonyl group intact after work up of the reaction mixture. Non-limiting examples of 1,2-reducing agents include metal hydrides and metalloid hydrides, e.g., aluminum hydride compounds, boron hydride compounds (e.g., CeCl_3 with NaBH_4), and ruthenium hydride complexes. Non-limiting examples of 1,4-reducing agents
25 include boron hydride compounds, hydrostannanes, copper hydride complexes (e.g., Stryker's reagent), palladium hydride complexes, platinum hydride complexes, iridium hydride complexes, rhodium hydride complexes, and ruthenium hydride complexes.

A compound having an allylic leaving group (e.g., a carboxylate, a halide, or a sulfonate) can be treated
30 with an allylic reducing agent to replace the leaving group with a hydrogen atom. A non-limiting example of allylic reducing agent is a palladium salt in combination with a formic acid salt (e.g., trialkylammonium formate).

Hydroxyl Protecting Groups and Hydroxyl Protecting Group Removing Agents

Hydroxyl protecting groups can be as defined herein. In particular, a hydroxyl protecting group can be an acyl, a sulfonyl, an arylalkyl (e.g., benzyl or *p*-methoxybenzyl), an aryl (e.g., *p*-methoxyphenyl), or an optionally substituted silyl (e.g., TMS, TES, TBS, TIPS, TBDPS, DMPS, or TPS). Hydroxyl protecting groups, hydroxyl protecting agents, and hydroxyl protecting reaction conditions can be selected to protect selectively certain hydroxyl groups in a compound, while leaving other hydroxyl groups unprotected. The
40 choice of hydroxyl protecting groups for a compound can facilitate subsequent deprotection strategies, as some hydroxyl protecting groups can be removed in the presence of others using appropriate hydroxyl

protecting group removing agents. Some of these strategies involving the choice of silyl hydroxyl protecting groups are discussed in, e.g., *Silicon-Based Blocking Agents*, Gelest, Inc., 2011.

Hydroxyl protecting group removing agents are those agents that can react with a compound having a protected hydroxyl group to afford the compound with a deprotected hydroxyl group. Hydroxyl protecting group removing agents and deprotection reaction conditions can be those known in the art. In a non-limiting example, hydroxyl masked as silyl ether can be unmasked by a reaction with a fluoride source (e.g., a fluoride salt, such as KF or TBAF). Alternatively, hydroxyl protected as TMS or TES ether can be deprotected by a reaction with a Brønsted acid (e.g., a carboxylic acid). In another non-limiting example, hydroxyl protected as an ester can be deprotected by a reaction with a C₁₋₆ alkoxide (e.g., alkali C₁₋₆ alkoxide or alkali earth C₁₋₆ alkoxide). In yet another non-limiting example, hydroxyl protected as an arylalkyl ether (e.g., 1-arylalk-1-yl ether) can be deprotected using a reduction reaction, e.g., with Pd/C and H₂ or with Na/NH₃. Alternatively, hydroxyl protected as an alkoxy-arylalkyl ether (e.g., MPM ether) can be deprotected by a reaction with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ). In still another non-limiting example, hydroxyl protected as alkoxyalkyl ether (e.g., 1-alkoxyalk-1-yl) or THP ether can be deprotected by a reaction with a Brønsted acid. Cyclic protected diols, such as acetals or ketals (e.g., as 2-alkyl-1,3-dioxolane, 2,2-dialkyl-1,3-dioxolane, 2-alkyl-1,3-dioxane, or 2,2-dialkyl-1,3-dioxane), can be deprotected by a reaction with a Brønsted acid (e.g., a carboxylic acid).

Dihydroxylation and Oxidative Carbon-Carbon Bond Cleavage

Dihydroxylation reaction produces diols from olefins. The reaction conditions can be those known in the art. Dihydroxylation reaction may be direct (e.g., Sharpless dihydroxylation, Upjohn dihydroxylation, or Milas dihydroxylation) or indirect (e.g., through epoxidation/hydrolysis, diborylation/oxidation, or carboxylation/hydrolysis (e.g., Woodward reaction or Prévost reaction)). A direct dihydroxylation reaction can be performed using an osmium complex (e.g., OsO₄ or osmate salt). A catalytic quantity of the osmium complex can be used in the dihydroxylation reactions in combination with an additional oxidizing agent (e.g., *N*-methylmorpholine *N*-oxide or hydrogen peroxide). The dihydroxylation reaction can be rendered stereoselective by including a chiral ligand (e.g., in AD-mix, which is a mixture of osmate and a chiral ligand).

When a dihydroxylation reaction is performed with an osmium complex and a periodate salt as oxidizing agent, the carbon-carbon double bond of an olefin can be cleaved to afford two carbonyl groups. The transformation of an olefin to two carbonyl groups is referred to herein as a cleavage reaction. Cleavage reaction conditions can be those known in the art. Non-limiting examples of cleavage reaction conditions can be ozonolysis reaction conditions or a reaction with an osmium complex (e.g., OsO₄ or osmate salt) and a periodate salt. A vicinal diol can be cleaved by sodium periodate or potassium permanganate.

Epimerizations

Epimerization reactions can be used to invert a stereogenic center having an undesired stereochemical identity. For example, through epimerization, *R* stereogenic center can be converted to *S* stereogenic center and *vice versa*. Epimerization of a stereogenic *sp*³-carbon bonded to one hydrogen atom and to

one hydroxyl group can be achieved through a reaction sequence involving oxidation of the hydroxyl group to a carbonyl group followed by a 1,2-reduction reaction. The 1,2-reduction reaction can provide the desired stereochemical identity diastereoselectively, or the reaction can be carried out using a chiral catalyst, chiral auxiliary, or a chiral reducing agent. Non-limiting examples of chiral reducing agents include alpine borane and prapine borane. Non-limiting examples of 1,2-reduction reactions involving chiral catalysts are Corey-Bakshi-Shibata reduction, Noyori hydrogenation, and Noyori transfer hydrogenation. The oxidation/reduction reaction sequence can be carried out *in situ* using dynamic kinetic resolution. A dynamic kinetic resolution can further involve a reaction with a hydroxyl protecting agent, which removes the desired stereoisomer from the reduction/oxidation equilibrium. In a non-limiting example, a dynamic kinetic resolution of chiral secondary alcohols can involve reduction/oxidation equilibration using $\eta^5\text{-Ph}_5\text{CpRu(CO)}_2\text{H}$ in combination with enantioselective esterification using isopropenyl acetate catalyzed by a lipase enzyme (e.g., lipase B from *Candida Antarctica*, see, e.g., Martin-Matute et al., *J. Am. Chem. Soc.*, 127:8817-8825, 2005).


Epimerization can also be carried out on a compound containing a tetrahydropyran-2-yl-acetaldehyde moiety, in which carbon 2 of the pyran ring exhibits an undesired stereochemical identity. Contacting this compound with L-proline can provide equilibrium between two stereoisomers. If other, non-equilibrating stereogenic centers are present in the compound, the most stable stereoisomer will be present in a larger quantity relative to other stereoisomer(s) in equilibrium with the most stable stereoisomer.

The following examples are meant to illustrate the invention. They are not meant to limit the invention in any way.

Examples

Example 1 – C.2-C.3 Macrocyclization through Horner-Wadsworth-Emmons Olefination



A halichondrin macrolide **3** can be prepared according to the below sequence.



- 5 Compound **1** can be converted to compound **2** through the Horner-Wadsworth-Emmons olefination (e.g., under Masamune conditions or according to Helquist protocol). Compound **2** can be converted to compound **3** after global deprotection (e.g., removal of silyl groups with a fluoride source, such as TBAF).

Example 2 – C.2-C.3 Macrocyclization through Ring-Closing Olefin Metathesis

A halichondrin macrolide **3** was prepared according to the below sequence.


**5 Ring-closing Olefin Metathesis**

Compound **4** (0.047 g, 0.027 mmol) was dissolved in toluene (57.0 mL) and 1,4-benzoquinone (1 mg, 10 μ mol), and Hoveyda-Grubbs 2nd generation catalyst (3 mg, 5.3 μ mol) was added. The reaction mixture was warmed to 70 °C. Upon completion (2h), DMSO (9.51 μ L, 0.134 mmol) was added, and the solution was allowed to stir at room temperature for 16 h. After concentration, the resulting green residue was purified by column chromatography (Biotage Ultra cartridge, 10g, heptane/EtOAc) to provide two RCM adducts, **5-cis** and **5-trans** with cis impurity (19 mg total, 0.011 mmol, 41%, 3:1 ratio cis/trans).

5-cis: ^1H NMR (400MHz, BENZENE- d_6) δ = 7.97 - 7.88 (m, 2H), 7.80 (dd, J =2.9, 6.4 Hz, 2H), 7.76 - 7.66 (m, 4H), 7.28 - 7.17 (m, 12H), 6.66 - 6.55 (m, 1H), 6.04 (d, J =15.6 Hz, 1H), 5.30 (dd, J =3.7, 6.8 Hz, 1H), 5.02 (s, 1H), 4.92 (t, J =1.0 Hz, 1H), 4.89 (br. s., 1H), 4.87 (s, 1H), 4.77 - 4.73 (m, 1H), 4.77 - 4.73 (m, 1H), 4.75 (s, 1H), 4.65 (br. s., 1H), 4.47 (br. s., 1H), 4.20 (dd, J =3.7, 6.8 Hz, 1H), 4.16 - 4.09 (m, 2H), 4.03 - 3.89 (m, 3H), 3.80 - 3.66 (m, 3H), 3.43 (d, J =9.4 Hz, 4H), 3.33 (d, J =3.5 Hz, 1H), 3.22 (d, J =9.4 Hz, 1H), 2.96 - 2.90 (m, 1H), 2.85 - 2.75 (m, 1H), 2.63 - 2.44 (m, 1H), 2.31 (d, J =5.5 Hz, 1H), 2.25 - 1.97 (m, 2H), 1.96 - 1.57 (m, 1H), 1.52 - 1.23 (m, 1H), 1.21 (s, 1H), 1.15 - 1.13 (m, 1H), 1.08 (s, 1H), 1.15 (s, 1H), 0.99 - 0.97 (m, 1H), 0.93 (s, 1H), 0.85 - 0.85 (m, 1H), 0.85 (s, 1H), 0.92 (s, 1H), 0.81 (br. s., 1H), 0.39 (s, 1H),

0.32 - 0.30 (m, 1H), 0.30 (s, 1H), 0.30 - 0.29 (m, 1H), 0.30 (s, 1H), 1.21 (s, 9H), 1.14 (s, 9H), 1.13 - 1.12 (m, 3H), 1.08 (s, 9H), 0.98 (s, 9H), 0.97 - 0.95 (m, $J=2.0$ Hz, 3H), 0.93 (s, 9H), 0.85 (s, 9H), 0.30 (s, 3H), 0.30 (s, 3H), 0.18 (s, 3H), 0.14 (s, 3H), 0.13 (s, 3H), 0.11 (s, 3H), 0.01 (s, 3H), -0.11 (s, 3H)

5 Compound 3



The mixture of compounds **5-cis** and **5-trans** (0.019 g, 0.011 mmol) was dissolved in N,N-dimethylacetamide (0.304 mL, 3.27 mmol). Imidazole hydrochloride (5.76 mg, 0.055 mmol) and TBAF (0.110 mL, 0.11 mmol) were added, and the reaction monitored by LCMS. Upon completion, water (5 mL) was added. Extraction with heptane removed lipophilic impurities. The aqueous layer was then extracted with toluene (5 mL) followed by toluene/THF (3:1 v/v, 4 x 8 mL). Brine was added to the aqueous layer to assist the recovery after the third extraction. The organic layers were combined and washed with saturated NaCl (2 x 6 mL). The organic layers were then dried over MgSO₄, filtered, and concentrated. The colorless residue was azeotroped with acetonitrile/water (3:1, v/v) at 30 °C to provide compound **6** (27 mg, 310% mass recovery) as a pale tan solid that was used directly in the next step. LCMS (CI): calcd. 790, found, Na ion 813 ($M + 1 + 22(\text{Na})$). Crude **6** was dissolved in DCM (0.087 mL, 1.352 mmol), and PPTS (0.028 g, 0.11 mmol) was added. The reaction was monitored by LCMS. At 2 h, product forms as determined by $M+1$ of 773.


The residue (8.70 mg, 11.00 μmol) was dissolved in dichloromethane (0.5 mL) at ambient temperature and PPTS (18 mg, 72 μmol) was added. The reaction mixture was filtered through a SiO₂ plug (0.75 cm x 2 cm) using EtOAc/MeOH to provide 5.5 mg of residue after concentration. The structure was confirmed by LCMS comparison with an authentic standard.

UPLC column:	Waters Acquity UPLC HSS T3 1.8 μ m, 2.1*50 mm, Part no. 186003538, Serial no. 01753602015820		
Temperature:	50 °C		
Flow rate:	0.8 mL/min		
Gradient:	Time, min	%-Solvent A	%-Solvent B
	Initial	97	3
	0.1	97	3
	2.0	1	99
	2.5	1	99
	2.51	97	3
	3.0	97	3
Injection volume:	1 μ L		
UV Detection:	PDA 210-400 nm		
Mass Detection(SQD):	ES(+), mass scan 100.00 to 2000.00		
Run time:	3 min		

Compound 3 standard, RT 1.90 min, ES+(M+1) 773


Found, RT 1.90 min, ES+ (M+1) 773

Compound **4** was prepared according to the below sequence:



(6S,E)-6-((2S,3R,4S,5S,6S)-6-(but-3-en-1-yl)-3,4,5-tris((tert-butyldimethylsilyl)oxy)tetrahydro-2H-pyran-2-yl)-6-((tert-butyldimethylsilyl)oxy)-1-((2S,5S)-5-(2-((2S,4R,6R)-6-

5 **(((2S,3R,4R,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-3-((4-methoxybenzyl)oxy)-4-methyloctahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)hex-4-en-3-ol**




Compound **8** (0.200 g, 0.242 mmol) and compound **7** (0.175 g, 0.156 mmol) were dissolved in N_2 sparged THF (0.5 mL). In a separate flask, the ligand (0.255 g, 0.86 mmol) and chromium(II) chloride (0.106 g, 0.862 mmol) were suspended in N_2 sparged THF (1.5 mL) at 30 °C. Triethylamine (0.12 mL, 0.861 mmol) was added, and the green solution was stirred for 30 minutes before cooling to 0 °C, at which time, Ni-DMP complex (8.00 mg, 0.024 mmol) was added. The solution of compounds **7** and **8** was added, and the reaction warmed to room temperature. Upon completion, the reaction was cooled to 0 °C, and ethylenediamine (0.158 mL, 2.345 mmol) was added and stirred for 1 h. Water (2 mL) was

added followed by heptane (10 mL) after 10 min. The phases were separated, and the aqueous phase was extracted with MTBE twice. The combined extracts were sequentially washed with HCl (1 N), water, and brine. After drying with MgSO_4 , the filtrates were concentrated to provide a pale green solid.

Heptane trituration provided a solution that was loaded directly on a Redisep cartridge (4 g, heptane/EtOAc) and provided the compound **9** (217 mg, 0.119 mmol, 76%, (est. 3:1 ratio of diastereomers)). The product was used without further purification in the next step.

(S,E)-6-((2S,3R,4S,5S,6S)-6-(but-3-en-1-yl)-3,4,5-tris((tert-butyldimethylsilyl)oxy)tetrahydro-2H-pyran-2-yl)-6-((tert-butyldimethylsilyl)oxy)-1-((2S,5S)-5-(2-((2S,4R,6R)-6-

(((2S,3R,4R,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-3-((4-methoxybenzyl)oxy)-4-methyloctahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)hex-4-en-3-one




Compound **9** (0.217 g, 0.119 mmol) was dissolved in DCM (2.170 mL), and water (0.107 μL) was added followed by Dess-Martin periodinane (0.061 g, 0.143 mmol). Upon completion, aqueous sodium bicarbonate (2 mL) was added, and the mixture was stirred for 15 minutes before extracting with DCM (3 x 2 mL). The combined extracts were washed with water and dried (Na_2SO_4) to provide 246 mg of crude product as a white foam. The material was purified using a Redisep cartridge (4 g, conditioned with 9:1::heptane/EtOAc, loaded with 1:1 heptane/DCM, and eluted as a single 30 mL fraction of heptane/EtOAc (9:1, (v/v)). After concentration, **15** (211 mg, 0.116 mmol, 97%) was provided as a white crunchy foam. ^1H NMR (400MHz, CDCl_3) δ = d = 7.92 - 7.84 (m, 2H), 7.79 (dd, J =2.9, 6.4 Hz, 2H), 7.77 - 7.70 (m, 4H), 7.34 - 7.16 (m, 14H), 6.78 (d, J =1.0 Hz, 2H), 6.62 (d, J =1.0 Hz, 1H), 5.96 - 5.81 (m, 1H), 5.88 (d, J =5.9 Hz, 1H), 5.39 - 5.30 (m, 1H), 5.35 (dd, J =4.1, 6.8 Hz, 1H), 5.11 - 5.03 (m, 1H), 5.12 - 5.02 (m, 1H), 4.96 (d, J =10.9 Hz, 1H), 5.00 - 4.92 (m, 1H), 4.85 (s, 1H), 4.82 - 4.79 (m, 1H), 4.79 - 4.76 (m, 1H), 4.79 (d, J =8.2 Hz, 1H), 4.50 (t, J =5.5 Hz, 1H), 4.40 (d, J =7.8 Hz, 1H), 4.25 - 4.19 (m, 1H), 4.18 - 4.11 (m, 1H), 4.11 - 4.00 (m, 1H), 3.96 (dd, J =5.1, 9.4 Hz, 1H), 3.85 - 3.67 (m, 1H), 3.61 - 3.41 (m, 2H), 3.34 - 3.25 (m, 3H), 3.20 - 3.10 (m, 2H), 2.85 - 2.57 (m, 1H), 2.55 - 2.30 (m, 2H), 2.28 - 1.89 (m, 7H), 1.89 - 1.76 (m, 4H), 1.76 - 1.28 (m, 3H), 1.18 (s, 7H), 1.24 - 1.16 (m, 6H), 1.15 (s, 10H), 1.09 (s, 9H), 0.97 (s, 11H), 0.96 (br. s., 4H), 0.93 (s, 9H), 0.91 (s, 9H), 0.85 (t, J =6.8 Hz, 4H), 0.33 (br. s., 3H), 0.32 (br. s., 3H), 0.19 - 0.17 (m, 3H), 0.14 (br. s., 3H), 0.14 - 0.13 (m, 3H), 0.12 (s, 6H), 0.07 - 0.05 (m, 3H), 0.07 - 0.05 (m, 3H), 0.01 - -0.03 (m, 1H), -0.01 (s, 3H).

(S,E)-6-((2R,3R,4S,5S,6S)-6-(but-3-en-1-yl)-3,4,5-tris((tert-butyldimethylsilyl)oxy)tetrahydro-2H-pyran-2-yl)-6-((tert-butyldimethylsilyl)oxy)-1-((2S,5S)-5-(2-((2S,4R,6R)-6-


(((2S,3R,4R,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-3-((4-methoxybenzyl)oxy)-4-methyloctahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)hex-4-en-3-one

3-hydroxy-4-methyloctahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)hex-4-en-3-one



- 5 Compound **10** (0.211 g, 0.116 mmol) was dissolved in DCM (4.22 mL), *t*-BuOH (0.042 mL) and pH 7 phosphate buffer were added (0.422 mL). DDQ (53 mg, 0.23 mmol) was added. Upon completion, aqueous sodium bicarbonate (10 mL) was added, and the solution stirred for 10 minutes before it was extracted with heptane (3 x 5 mL). The combined extracts were washed with aqueous sodium bicarbonate and water before drying with Na₂SO₄. The filtrate was concentrated, and residue was
- 10 purified by column chromatography (Biotage Ultra cartridge, 10 g, heptane/EtOAc) to provide **16** (139 mg, 0.083 mmol, 70%). ¹H NMR (400MHz, BENZENE-*d*₆) δ = 7.93 - 7.84 (m, 2H), 7.84 - 7.67 (m, 7H), 7.34 - 7.14 (m, 11H), 6.62 (d, *J*=1.0 Hz, 1H), 5.96 - 5.82 (m, 1H), 5.35 (dd, *J*=4.3, 7.4 Hz, 1H), 5.08 (d, *J*=15.6 Hz, 1H), 4.97 (d, *J*=10.2 Hz, 2H), 4.92 (s, 2H), 4.89 (s, 3H), 4.86 (s, 3H), 4.75 (s, 3H), 4.48 - 4.34 (m, 2H), 4.25 - 4.18 (m, 1H), 4.16 (s, 1H), 4.13 - 4.07 (m, 1H), 4.07 - 3.95 (m, 2H), 3.95 - 3.85 (m, 1H), 3.85 - 3.75
- 15 (m, 1H), 3.75 - 3.60 (m, 3H), 3.52 - 3.42 (m, 4H), 3.33 - 3.26 (m, 1H), 3.12 - 3.08 (m, 1H), 2.86 - 2.72 (m, 2H), 2.72 - 2.58 (m, 3H), 2.54 - 2.00 (m, 11H), 1.98-1.20 (m, 11H), 1.12 (s, 9H), 1.28 - 1.11 (m, 60H), 1.10 (s, 9H), 0.98 (s, 9H), 0.93- 0.91 (m, 3H), 0.87 (s, 9H), 0.85 (s, 9H), 0.83 (s, 9H), 0.33 (br. s., 3H), 0.32 (br. s., 3H), 0.18 (s, 3H), 0.15 (s, 3H), 0.14 (br. s., 3H), 0.12 (s, 3H), 0.06 (s, 3H), -0.01 (s, 3H)

- 20 **(2S,3R,4S,4aS,6R,7R,8aS)-2-(((2R,4R,6S)-6-(2-((2S,5S)-5-((S,E)-6-((2R,3R,4S,5S,6S)-6-(but-3-en-1-yl)-3,4,5-tris((tert-butyldimethylsilyl)oxy)tetrahydro-2H-pyran-2-yl)-6-((tert-butyldimethylsilyl)oxy)-3-oxohex-4-en-1-yl)-3-methylenetetrahydrofuran-2-yl)ethyl)-4-methyl-3-methylenetetrahydro-2H-pyran-2-yl)methyl)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-4-methyloctahydropyrano[3,2-b]pyran-3-yl acrylate**




25


Compound **11** (0.082 g, 0.048 mmol) was dissolved in DCM (820 μ L), and the resulting solution was cooled to 0 °C. Triethylamine (0.05 mL, 0.359 mmol) was added, followed by acryloyl chloride (4 μ L,

0.049 mmol). Additional 0.05 mL TEA and 6 μ L acryoyl chloride were added. Upon completion of the reaction, saturated sodium bicarbonate (5 mL) was added, and the reaction mixture was extracted with heptane (3 x 3 mL). The combined extracts were washed with water and dried (Na_2SO_4). After concentration, the residue was purified by column chromatography (Biotage Ultra cartridge, 10 g, heptane/EtOAc) to provide compound **4** (48 mg, 0.027 mmol, 57%) and a mixture of compounds **16** and **17** (35 mg). ^1H NMR (400MHz, BENZENE- d_6) δ = 7.95 - 7.86 (m, 3H), 7.79 (dd, J =3.1, 6.6 Hz, 3H), 7.73 - 7.66 (m, 7H), 7.34 - 7.27 (m, 1H), 7.26 - 7.15 (m, 7H), 6.66 - 6.57 (m, 1H), 6.34 (dd, J =1.6, 17.6 Hz, 1H), 5.99 (dd, J =10.2, 17.2 Hz, 1H), 5.94 - 5.80 (m, 1H), 5.35 (dd, J =4.1, 7.2 Hz, 1H), 5.31 - 5.25 (m, 1H), 5.15 - 4.92 (m, 1H), 4.91 - 4.83 (m, 3H), 4.77 (s, 1H), 4.51 - 4.41 (m, 1H), 4.34 (t, J =1.0 Hz, 1H), 4.25 - 4.19 (m, 1H), 4.16 (s, 1H), 4.09 - 3.95 (m, 4H), 3.89 (dt, J =4.7, 9.4 Hz, 1H), 3.80 (t, J =8.8 Hz, 1H), 3.75 - 3.68 (m, 1H), 3.66 (br. s., 2H), 3.60 - 3.49 (m, 1H), 3.42 - 3.35 (m, 2H), 3.29 (d, J =1.0 Hz, 1H), 2.95 (t, J =1.6 Hz, 1H), 2.86 - 2.61 (m, 2H), 2.56 - 2.33 (m, 3H), 2.31 - 1.34 (m, 15H), 2.30-2.00 (m, 2H), 1.12 (s, 9H), 1.09 (s, 9H), 1.06 (d, J =1.0 Hz, 3H), 0.97 (s, 9H), 0.95 (d, J =1.0 Hz, 3H), 0.93 (s, 9H), 0.91 (s, 9H), 0.85 (s, 9H), 0.32 (br. s., 3H), 0.18 (s, 3H), 0.14 (s, 3H), 0.13 (br. s., 3H), 0.12 (s, 3H), 0.07 - 0.05 (m, 3H), 0.06 (s, 3H), -0.01 (s, 3H)

Compound **8** was prepared as shown in the following sequence:




(2S,3S,4S,5R,6S)-4,5-bis((tert-butyldimethylsilyl)oxy)-6-((S,E)-1-((tert-butyldimethylsilyl)oxy)-3-(trimethylsilyl)allyl)-2-(3-hydroxypropyl)tetrahydro-2H-pyran-3-yl benzoate




Compound **15** (1.160 g, 1.548 mmol) was dissolved in THF (10 mL), and 9-borabicyclo[3.3.1]nonane (5.0 mL, 2.50 mmol) was added. Upon completion, the solution was cooled to 0 °C, and water (10 mL) was added followed by sodium perborate tetrahydrate (1.429 g, 9.288 mmol). The mixture was stirred for 12 h. Aqueous sodium bicarbonate was added, and the mixture was extracted twice with heptane/MTBE (1:1 (v/v)). The combined organic layers were washed with aqueous sodium bicarbonate followed by water and then dried over sodium sulfate. The resulting solution was concentrated, and the resulting residue was purified by column chromatography to provide 0.820 g (1.13 mmol, 73%) of compound **8A**. ¹H NMR (400MHz, CDCl₃) δ = 8.09 (br. s., 1H), 7.44 - 7.36 (m, 1H), 7.27 - 7.22 (m, 2H), 7.18 - 7.13 (m, 2H), 5.96 (br. s., 1H), 5.21 (br. s., 1H), 4.47 (br. s., 1H), 4.11 (br. s., 2H), 4.01 - 3.91 (m, 3H), 3.65 (br. s., 3H), 3.30 (br. s., 1H), 0.96 (s, 9H), 0.89 (s, 9H), 0.87 (s, 9H), 0.16 - 0.08 (m, 27 H).

(2S,3S,4S,5R,6S)-2-(3-acetoxypropyl)-4,5-bis((tert-butyldimethylsilyl)oxy)-6-((S,E)-1-((tert-butyldimethylsilyl)oxy)-3-(trimethylsilyl)allyl)tetrahydro-2H-pyran-3-yl benzoate




Compound **8A** (0.870 g, 1.134 mmol) was dissolved in DCM (8.70 mL) and cooled to 0 °C. Triethylamine (1 mL, 7.175 mmol) and DMAP (0.014 g, 0.113 mmol) were added followed by Ac₂O (0.214 mL, 2.268 mmol). Upon completion, saturated aqueous sodium bicarbonate (30 mL) was added. The mixture was extracted with MTBE (3 x 20 mL). The combined extracts were washed with saturated aqueous sodium bicarbonate, then water, and then brine. After drying over Na₂SO₄, the solution was concentrated to provide compound **8B** (0.895 g, 1.106 mmol, 98%) as a colorless oil. ¹H NMR (400MHz, CDCl₃) δ = 8.12 - 8.04 (m, 1H), 7.55 - 7.48 (m, 1H), 7.43 - 7.34 (m, 2H), 6.49 - 6.44 (m, 1H), 5.97 - 5.89 (m, 1H), 5.18 - 5.09 (m, 1H), 4.46 - 4.33 (m, 1H), 4.14 - 4.01 (m, 3H), 3.96 - 3.89 (m, 2H), 3.20 (s, 3H), 3.00 - 2.97 (m, 1H), 2.35 - 2.33 (m, 1H), 2.06 - 1.99 (m, 2H), 1.81 - 1.74 (m, 1H), 1.64 - 1.54 (m, 2H), 0.96 (s, 9H), 0.88 (s, 9H), 0.86 (s, 9H), 0.18 - 0.08 (m, 9H), 0.07 (s, 6H), 0.04 (s, 6H), -0.05 - -0.11 (m, 23).

(2S,3S,4S,5R,6S)-2-(3-acetoxypropyl)-4,5-bis((tert-butyldimethylsilyl)oxy)-6-((S,E)-1-((tert-butyldimethylsilyl)oxy)-3-iodoallyl)tetrahydro-2H-pyran-3-yl benzoate




Compound **8B** (0.895 g, 1.106 mmol) was dissolved in toluene (5.37 ml) and acetonitrile (10.74 mL, 205.633 mmol). *t*-Butyldimethylchlorosilane (0.017 g, 0.111 mmol) was added, and the solution was warmed to 30 °C. NIS (1.990 g, 8.846 mmol) was added. Upon completion, the reaction mixture was poured into a solution of sodium thiosulfate (2 g) in saturated aqueous sodium bicarbonate (40 mL). MTBE (20 mL) was added, and the mixture stirred for 45 minutes. The MTBE layer was removed *in vacuo*, and the aqueous layer was extracted with MTBE (2 x 20 mL). The combined MTBE extracts were washed with 10% (w/v) aqueous sodium thiosulfate, followed by water and then by brine. The solution was dried over MgSO₄ and concentrated to provide crude **8C** (857 g, 0.993 mmol, 90%) as pale yellow oil. ¹H NMR (400MHz, CDCl₃) δ = 8.06 (br. s., 2H), 7.53 (br. s., 1H), 7.40 (br. s., 2H), 6.73 (br. s., 1H), 6.38 (d, *J*=14.8 Hz, 1H), 5.20 (br. s., 1H), 5.01 (br. s., 1H), 4.67 - 4.56 (m, 1H), 4.54 - 4.46 (m, 1H), 4.44 - 4.36 (m, 1H), 4.17 - 3.98 (m, 3H), 3.98 - 3.82 (m, 2H), 3.34 - 3.19 (m, 1H), 2.72 (s, 1H), 2.33 (s, 1H), 2.14 - 1.88 (m, 3H), 1.88 - 1.50 (m, 4H), 1.37 - 1.17 (m, 2H), 0.94 (s, 9H), 0.88 (s, 9H), 0.87 (s, 10H), 0.05 (s, 18H)

(2S,3S,4S,5R,6S)-4,5-bis((tert-butyldimethylsilyl)oxy)-6-((S,E)-1-((tert-butyldimethylsilyl)oxy)-3-iodoallyl)-2-(3-hydroxypropyl)tetrahydro-2H-pyran-3-ol



Compound **8C** (0.814 g, 0.943 mmol) was dissolved in toluene (8.14 mL) and cooled to -10 °C. DIBAL (4.72 ml, 4.716 mmol) was added. After 15 minutes, methanol (0.191 mL, 4.716 mmol) was added followed by aqueous HCl (0.1 N, 20 mL). The mixture was stirred for 15 minutes and the extracted with MTBE three times. The combined extracts were washed with water and brine before drying over Na₂SO₄. The resulting solution was concentrated *in vacuo*, and the residue was purified by column chromatography (Biotage Ultra cartridge, 25 g; heptane/EtOAc) to provide compound **8D** (601 g, 0.838 mmol, 89%).


(2S,3S,4S,5R,6S)-4,5-bis((tert-butyldimethylsilyl)oxy)-6-((S,E)-1-((tert-butyldimethylsilyl)oxy)-3-iodoallyl)-2-(3-((triethylsilyl)oxy)propyl)tetrahydro-2H-pyran-3-ol



Step 1. Compound **8D** (0.601 g, 0.838 mmol) in DCM (6.01 mL) was cooled to 0 °C. 2,4-Lutidine (0.291 mL, 2.515 mmol) was added followed by chlorotriethylsilane (0.148 mL, 0.88 mmol). Upon completion, saturated aqueous sodium bicarbonate (20 mL) was added, and the mixture was extracted with DCM (3 x 20 mL). The combined extracts were washed with water. After drying over Na₂SO₄, the organic layers were concentrated to provide crude TES-ether (0.697 g).

Step 2. The TES-ether (0.697 g) was dissolved in DCM (6.97 mL) and cooled to 0 °C. 2,4-Dimethylpyridine (0.388 mL, 3.354 mmol) was added followed by *t*-butyldimethylsilyl trifluoromethanesulfonate (0.231 mL, 1.006 mmol). Upon completion, saturated aqueous sodium bicarbonate (20 mL) was added, and the mixture was extracted with MTBE (3 x 20 mL). The combined extracts were washed with water, HCl (0.1 N), water, and then aqueous sodium bicarbonate. After drying, the filtrate was concentrated to a colorless oil and purified by column chromatography (Biotage Snap 25 cartridge, heptane/EtOAc) to provide compound **8E** (0.650 g, 0.687 mmol, 82%). ¹H NMR (400 MHz, CDCl₃) δ = 6.80 - 6.57 (m, 1H), 6.26 (dd, J=0.8, 14.5 Hz, 1H), 4.58 - 4.25 (m, 1H), 3.92 - 3.66 (m, 3H), 3.64-3.55 (m, 3H), 3.05 (d, J=8.6 Hz, 1H), 1.68 - 1.55 (m, 1H), 1.51 - 1.38 (m, 1H), 1.49 - 1.33 (m, 2H), 1.00 - 0.83 (m, 36H), 0.60 (q, J=8.2 Hz, 6H), 0.13 - -0.03 (m, 33H)

((2S,3R,4S,5S,6S)-2-((S,E)-1-((tert-butyldimethylsilyl)oxy)-3-iodoallyl)-6-(3-((triethylsilyl)oxy)propyl)tetrahydro-2H-pyran-3,4,5-triyl)tris(oxy))tris(tert-butyldimethylsilane)




Step 1 (Swern oxidation): DMSO (0.19 mL, 1.685 mmol) in DCM (0.4 mL) was added to oxalyl chloride (2 M in DCM, 0.357 mL, 0.715 mmol) in DCM (1 mL) at -78 °C. After 10 minutes, a solution of compound **8E** (0.169 g, 0.179 mmol) in DCM (1 mL with a 1 mL rinse) was added dropwise at -78 °C. The solution was warmed to -40 °C and stirred for 30 min. The solution was cooled to -78 °C and triethylamine (0.299 mL, 2.145 mmol) was added. The reaction mixture was warmed to 0 °C and HCl (0.1 N) was added. The mixture was extracted with DCM three times. The organic layers were combined, washed with water and then dried with Na₂SO₄. Concentration provided the aldehyde (171 mg, 134% mass recovery) as a pale yellow oil that was used directly in step 2.

Step 2. (Wittig reaction): Methyltriphenylphosphonium bromide (0.255 g, 0.715 mmol) was suspended in THF (1.6 mL) and cooled to 0 °C. Butyllithium (2M in cyclohexane, 0.349 mL, 0.697 mmol) was added,

and the resulting solution was stirred for 30 min. The aldehyde was dissolved in THF (1.6 mL) and added to the ylide mixture. Upon completion, HCl (0.1 N) was added and the mixture extracted with heptane (3 x 20 mL). The combined organics were washed with water followed by aqueous sodium bicarbonate. After drying with Na₂SO₄, the filtrate was concentrated to provide crude compound **8F** (0.155 g, 0.217 mmol, 122% mass recovery, contaminated with PPh₃). ¹H NMR (400 MHz, CDCl₃) δ = 6.70 (q, *J* = 6.3 Hz, 1H), 6.38 - 6.22 (m, 1H), 5.88 - 5.71 (m, 1H), 5.13 - 4.96 (m, 2H), 4.39 - 4.28 (m, 1H), 3.78 (s, 3H), 3.69 - 3.62 (m, 2H), 3.06 - 3.01 (m, 1H), 2.21 - 2.01 (m, 1H), 2.20 - 2.00 (m, 1H), 1.82 - 1.63 (m, 1H), 1.45 - 1.20 (m, 1H), 0.92 (s, 9H), 0.90 (s, 9H), 0.88 (s, 9H), 0.11 (s, 3H), 0.10 (s, 3H), 0.09 (s, 3H), 0.07 (s, 3H), 0.05 (s, 3H), 0.04 (s, 3H)


(((2S,3S,4S,5R,6S)-2-(but-3-en-1-yl)-6-((S,E)-1-((tert-butyldimethylsilyl)oxy)-3-iodoallyl)tetrahydro-2H-pyran-3,4,5-triyl)tris(oxy))tris(tert-butyldimethylsilane)




Compound **8F** (0.127 g, 0.178 mmol) was dissolved in dichloromethane (1.270 mL) and cooled to 0 °C. 2,4-Dimethylpyridine (0.057 g, 0.534 mmol) was added followed by *t*-butyldimethylsilyl trifluoromethanesulfonate (0.049 mL, 0.214 mmol). After 3 h, additional 2,4-dimethylpyridine (0.4 mL) and *t*-butyldimethylsilyl trifluoromethanesulfonate (0.1 mL) were added. Upon completion, aqueous sodium bicarbonate was added, and the reaction was stirred for 15 min. The product was extracted with heptane (3 x 10 mL). The combined extracts were washed with water, HCl (1N, 2 x 10 mL), and water. After drying with Na₂SO₄/MgSO₄, the filtrate was concentrated to provide a pale yellow oil that was purified by column chromatography (Biotage Ultra cartridge, 10 g, H/EtOAc) to provide compound **8** (141 mg, 0.17 mmol, 95% over 3-steps (Swern, Wittig, TBS-protection)).

Example 3 – C.3-C.4 Macrocyclization through Ring-Closing Olefin Metathesis

A halichondrin macrolide **3** can be prepared according to the below sequence.




Compound **12** can be prepared according to the following sequence:



Example 4 – C.12-C.13 Macrocyclization through Ring-Closing Olefin Metathesis


A halichondrin macrolide **3** can be prepared according to the below sequence.




- Compound **21** can be converted to compound **22** through the catalytic ring-closing olefin metathesis (e.g.,
5 with catalytic quantities of Hoveyda-Grubbs 2nd generation catalyst). Compound **22** can be converted to
compound **3** after global deprotection (e.g., removal of silyl groups with a fluoride source, such as TBAF).

Example 5 – C. 15-C. 16 Macrocyclization through Ring-Closing Olefin Metathesis

A halichondrin macrolide **3** can be prepared according to the below sequence.




5 2-((2R,3S,3aR,4aS,7R,8aS,9S,9aR)-3,9-bis((tert-butyldiphenylsilyl)oxy)-2-((triethylsilyl)oxy)but-3-en-1-yl)decahydrofuro[3,2-b]pyrano[2,3-e]pyran-7-yl)acetic acid



2-((2R,3S,3aR,4aS,7R,8aS,9S,9aR)-3,9-bis((tert-butyldiphenylsilyl)oxy)-2-((triethylsilyl)oxy)but-3-en-1-yl)decahydrofuro[3,2-b]pyrano[2,3-e]pyran-7-yl)ethanol (compound **23**, 70 mg, 0.076 mmol) and TEMPO (1.187 mg, 7.597 μmol) were dissolved in toluene (2 mL) at ambient temperature. To the resulting solution were added pH 7 phosphate buffer (1 mL), a solution of sodium chlorite (42.9 mg, 0.38 mmol) in water (1 mL), and 4% bleach (a solution of sodium hypochlorite in water) (0.012 mL, 7.597 μmol) slowly. After being stirred overnight, the reaction mixture was treated with sodium bisulfite (63.2 mg, 0.608 mmol) and water (5 mL). The resulting mixture was extracted twice with MTBE (10 mL each time). The combined organic layers were washed with 30% aqueous NaCl (2.0 mL) and dried over MgSO₄.

Filtration, concentration *in vacuo*, and purification by silica gel column chromatography using a 25 - 66% gradient of ethyl acetate in *n*-heptane as eluent provided 54 mg of the target product as a white foam solid. ¹H NMR (1:1 diastereomeric mixture, 400 MHz, CDCl₃) δ ppm 0.50 - 0.66 (m, 6 H) 0.86 - 1.01 (m, 9 H) 1.05 - 1.17 (m, 18 H) 1.30 - 1.45 (m, 3 H) 1.55- 1.65 (m, 1.5H) 1.68 - 1.79 (m, 1.5H) 1.87 - 1.99 (m, 1 H) 2.09 - 2.25 (m, 1.5 H) 2.38 (d, *J*=5.5 Hz, 1 H) 2.94 (dd, *J*=9.4, 4.3 Hz, 0.5 H) 3.11 (dd, *J*=9.6, 6.4 Hz, 0.5 H) 3.34 (t, *J*=4.9 Hz, 0.5 H) 3.46-3.54 (m, 1 H) 3.58 (t, *J*=4.7 Hz, 0.5 H) 3.62 - 3.78 (m, 2 H) 3.88 (t, *J*=6.3 Hz, 0.5 H) 3.95 - 4.09 (m, 1 H) 4.16 - 4.25 (m, 2 H) 4.26-4.42 (m, 1.5 H) 4.95 - 5.18 (m, 2 H) 5.71 (m, *J*=6.3 Hz, 1 H) 7.27 - 7.45 (m, 12 H) 7.64 - 7.77 (m, 8 H)


10 ((2R,5S)-5-(2-((2S,4R,6R)-6-(((2S,3R,4R,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-3-hydroxy-4-methyloctahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)methyl pivalate



15 A solution of 2-((2S,3R,4R,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-3-((4-methoxybenzyl)oxy)-4-methyloctahydropyrano[3,2-b]pyran-2-yl)acetaldehyde (compound **25**; 340 mg, 0.347 mmol) and ((2R,5S)-5-((S)-5-methyl-3-((triethylsilyl)oxy)hepta-5,6-dien-1-yl)-4-methylenetetrahydrofuran-2-yl)methyl pivalate (compound **26**; 197 mg, 0.452 mmol) in dichloromethane (10 mL) was cooled to -37 °C. Methoxyacetic acid (0.533 mL, 6.946 mmol) and BF₃.OEt₂ (0.132 mL, 1.042 mmol) were added, and the resulting solution was stirred for 3 h between -20 to -35 °C. The reaction mixture was then treated with sat. NaHCO₃ (15 mL) and extracted twice with MTBE (30.3 mL each). The organic layers were combined, washed with 30% aqueous NaCl (15 mL), and dried over MgSO₄. Filtration, concentration *in vacuo*, and purification by silica gel column chromatography using a 20 - 50% gradient of ethyl acetate in *n*-heptane as eluent provided 0.367 g of a mixture of the target product and byproducts, identified as compound **25** that underwent MPM deprotection and compound **26** that underwent TES deprotection. Without further purification, the mixture was dissolved in THF (7.5 mL). In another reactor, Pd(PPh₃)₄ (0.037 g, 0.032 mmol) and triphenylphosphine (0.034 g, 0.128 mmol) were dissolved in THF (2 mL) and treated with formic acid (0.123 mL, 3.204 mmol) and triethylamine (0.447 mL, 3.204 mmol). The resulting solution was added to the substrate solution prepared above. After being stirred for 20 h at 60 to 65 °C, the reaction mixture was diluted with MTBE (20 mL) and washed with sat. aq. NaHCO₃ (5 mL) and 30% aqueous NaCl (5 mL). The organic layer was dried over MgSO₄, filtered, and concentrated *in vacuo*. The residue was purified by silica gel column chromatography using a 20 - 100% gradient of ethyl acetate in *n*-heptane as eluent to give 98 mg of the target product. ¹H NMR (400 MHz, CDCl₃) δ ppm 1.01 (s, 9 H) 1.05 (s, 9 H) 1.10 (d,


$J=6.3$ Hz, 3 H) 1.13 (d, $J=7.8$ Hz, 3 H) 1.20 (s, 9 H) 1.28 - 1.37 (m, 1 H) 1.45 - 1.73 (m, 8 H) 1.77 - 1.85 (m, 1 H) 1.91 - 2.15 (m, 5 H) 2.24 - 2.33 (m, 1 H) 2.37 - 2.47 (m, 1 H) 2.63 - 2.74 (m, 1 H) 3.22 (t, $J=2.5$ Hz, 1 H) 3.29 - 3.38 (m, 1 H) 3.49 (d, $J=8.2$ Hz, 1 H) 3.55 - 3.64 (m, 2 H) 3.68 (d, $J=2.0$ Hz, 1 H) 3.72 - 3.82 (m, 3 H) 3.91 (t, $J=5.9$ Hz, 1 H) 4.01 (dd, $J=11.7$, 4.7 Hz, 1 H) 4.05 - 4.10 (m, 1 H) 4.13 (dd, $J=11.7$, 4.7 Hz, 1 H) 4.22 - 4.34 (m, 1 H) 4.41 - 4.52 (m, 1 H) 4.77 - 4.84 (m, 2 H) 4.90 (d, $J=2.0$ Hz, 1 H) 4.99 (d, $J=2.3$ Hz, 1 H) 7.28 - 7.44 (m, 12 H) 7.60 (dq, $J=8.1$, 1.5 Hz, 4 H) 7.66 - 7.69 (m, 2 H) 7.75 - 7.78 (m, 2 H)

((2R,5S)-5-(2-((2S,4R,6R)-6-(((2S,3R,4S,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-4-methyl-3-((triethylsilyl)oxy)octahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)methyl pivalate



((2R,5S)-5-(2-((2S,4R,6R)-6-(((2S,3R,4R,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-3-hydroxy-4-methyloctahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)methyl pivalate (compound **27**; 98 mg, 0.093 mmol) was dissolved in dichloromethane (5 mL) and treated with imidazole (31.5 mg, 0.463 mmol) and chlorotriethylsilane (0.047 mL, 0.278 mmol) at ambient temperature. After 15 h, the reaction was quenched with saturated aqueous NH_4Cl (27% (w/v)) (10 mL). The resulting mixture was extracted twice with MTBE (20 mL). The combined organic layers were washed with 30% aqueous NaCl (5.0 mL) and dried over MgSO_4 . Filtration, concentration *in vacuo*, and purification by silica gel column chromatography using a 10 - 50% gradient of ethyl acetate in *n*-heptane as eluent provided 94 mg of the target product. ^1H NMR (400 MHz, CDCl_3) δ ppm 0.65 (q, $J=7.8$ Hz, 6 H) 0.97 (t, $J=7.8$ Hz, 9 H) 1.02 (s, 9 H) 1.04 (s, 9 H) 1.06 - 1.10 (m, 6 H) 1.20 (s, 9 H) 1.44 - 1.52 (m, 2 H) 1.58 - 1.68 (m, 4 H) 1.71 - 1.84 (m, 3 H) 1.85 - 2.01 (m, 2 H) 2.07 - 2.23 (m, 2 H) 2.35 - 2.45 (m, 1 H) 2.63 - 2.75 (m, 1 H) 3.18 - 3.28 (m, 2 H) 3.43 - 3.51 (m, 2 H) 3.57 - 3.61 (m, 2 H) 3.64 - 3.72 (m, 1 H) 3.74 - 3.85 (m, 3 H) 4.01 (dd, $J=11.7$, 4.7 Hz, 1 H) 4.12 (dd, $J=11.3$, 5.1 Hz, 1 H) 4.23 - 4.31 (m, 1 H) 4.39 - 4.47 (m, 1 H) 4.78 (s, 1 H) 4.84 (br. s., 1 H) 4.86 (br. s., 1 H) 4.97 (s, 1 H) 7.29 - 7.44 (m, 12 H) 7.61 - 7.71 (m, 8 H)

((2R,5S)-5-(2-((2S,4R,6R)-6-(((2S,3R,4S,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-4-methyl-3-((triethylsilyl)oxy)octahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)methanol




((2R,5S)-5-(2-((2S,4R,6R)-6-(((2S,3R,4S,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-4-methyl-3-((triethylsilyl)oxy)octahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)methyl pivalate (compound **28**; 94 mg, 0.08 mmol) was dissolved in dichloromethane (3.0 mL) and cooled to -78 °C.

DIBAL in toluene (1.0 M, 0.24 mL) was charged at this temperature. After being stirred for 1 h between -65 to -78 °C, the reaction mixture was treated with MeOH (0.5 mL) and a Rochelle's salt solution (5.0 mL). The resulting mixture was warmed to ambient temperature and extracted with MTBE (15 mL). The organic layers were washed 30% aqueous NaCl (5.0 mL) and dried over MgSO₄. Filtration and concentration *in vacuo* provided the target product (86mg), which was used in next step without further


purification. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.65 (q, *J*=8.1 Hz, 6 H) 0.97 (t, *J*=7.8 Hz, 9 H) 1.02 (s, 9 H) 1.04 (s, 9 H) 1.06 - 1.10 (m, 6 H) 1.45 - 1.52 (m, 2 H) 1.58 - 1.69 (m, 4 H) 1.72 - 1.83 (m, 3 H) 1.83 - 1.98 (m, 3 H) 2.09 - 2.25 (m, 2 H) 2.37 - 2.46 (m, 1 H) 2.56 - 2.65 (m, 1 H) 3.18 - 3.27 (m, 2 H) 3.43 - 3.53 (m, 3 H) 3.55 - 3.64 (m, 3 H) 3.65 - 3.72 (m, 1 H) 3.74 - 3.85 (m, 1 H) 4.10 - 4.21 (m, 1 H) 4.37 - 4.44 (m, 1 H) 4.78 (s, 1 H) 4.84 (d, *J*=2.0 Hz, 1 H) 4.86 (s, 1 H) 4.97 (d, *J*=2.3 Hz, 1 H) 7.30 - 7.42 (m, 12 H) 7.61 - 7.71 (m, 8 H)

(E)-ethyl 3-((2R,5S)-5-(2-((2S,4R,6R)-6-(((2S,3R,4S,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-4-methyl-3-((triethylsilyl)oxy)octahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)acrylate




To a solution of ((2R,5S)-5-(2-((2S,4R,6R)-6-(((2S,3R,4S,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-4-methyl-3-((triethylsilyl)oxy)octahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)methanol (compound **35**; 86 mg, 0.079 mmol) in dichloromethane (3.0 mL) at ambient temperature was added sodium bicarbonate (26.6 mg, 0.316 mmol) and Dess-Martin periodinane (50.3 mg, 0.119 mmol). After being stirred for 2 h, the reaction mixture was diluted with MTBE (20 ml) and treated with a saturated aqueous Na₂S₂O₃ solution (10 mL) and water (5.0 mL). The resulting mixture was stirred for 0.5 h at ambient temperature, and the layers were separated. The organic layer was washed with a saturated aqueous NaHCO₃ solution (10 mL) twice and 30% aqueous NaCl (5.0 mL) and dried over MgSO₄. Filtration followed by concentration *in vacuo* gave the target product ((2R,5S)-5-(2-((2S,4R,6R)-6-(((2S,3R,4S,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-4-methyl-3-((triethylsilyl)oxy)octahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-carbaldehyde, 80 mg) as an oil. Without further purification, the crude product was dissolved in THF (4.0 mL) and cooled to 0 °C. In an another reaction flask, ethyl 2-(diethoxyphosphoryl)acetate (73.7 μL, 0.368 mmol) was dissolved in THF (2.7 mL), cooled to 0 °C, and treated with 60% sodium hydride (13.26 mg, 0.332 mmol). After 20 min at 0 °C, the resulting solution was added to the starting material reactor. After 30 min, saturated aqueous NH₄Cl (27% (w/v)) (8.0 mL), water (2.8 mL), and MTBE (20 mL) were added, and the resulting mixture was warmed to ambient temperature. The layers were separated, and the aqueous layer was extracted with MTBE (12 mL). The combined organic layers were washed with 30% aqueous NaCl (10 mL) and dried over MgSO₄. Filtration, concentration *in vacuo*, and purification by silica gel column chromatography using a 20 - 50% gradient of ethyl acetate in *n*-heptane as eluent provided 71 mg of the target product (compound **29**). ¹H NMR (400 MHz, CDCl₃) δ ppm 0.65 (q, *J*=8.1 Hz, 6 H) 0.97 (t, *J*=8.2 Hz, 9 H) 1.02 (s, 9 H) 1.04 (s, 9 H) 1.06 - 1.10 (m, 6 H) 1.45 - 1.52 (m, 2 H) 1.57 - 1.70 (m, 5 H) 1.72 - 1.83 (m, 3 H) 1.83 - 1.98 (m, 2 H) 2.08 - 2.23 (m, 2 H) 2.38 - 2.47 (m, 1 H) 2.76 - 2.86 (m, 1 H) 3.18 - 3.28 (m, 2 H) 3.43 - 3.51 (m, 2 H) 3.56 - 3.62 (m, 2 H) 3.64 - 3.71 (m, 1 H) 3.73 - 3.85 (m, 3 H) 4.19 (q, *J*=7.3 Hz, 2 H) 4.40 - 4.47 (m, 1 H) 4.60 - 4.69 (m, 1 H) 4.78 (s, 1 H) 4.86 (s, 2 H) 4.98 (d, *J*=2.0 Hz, 1 H) 6.00 (dd, *J*=15.6, 1.6 Hz, 1 H) 6.87 (dd, *J*=15.6, 5.1 Hz, 1 H) 7.30 - 7.42 (m, 12 H) 7.61 - 7.71 (m, 8 H)

(E)-3-((2R,5S)-5-(2-((2S,4R,6R)-6-(((2S,3R,4S,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-4-methyl-3-((triethylsilyl)oxy)octahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)prop-2-en-1-ol



To a solution of (E)-ethyl 3-((2R,5S)-5-(2-((2S,4R,6R)-6-(((2S,3R,4S,4aS,6R,7R,8aS)-7-((tert-butyl)diphenylsilyl)oxy)-6-(2-((tert-butyl)diphenylsilyl)oxy)ethyl)-4-methyl-3-((triethylsilyl)oxy)octahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)acrylate (compound **29**; 71 mg, 0.061 mmol) in dichloromethane (4mL), at -78 °C, was added a solution of DIBAL-H in toluene (1.0 M, 0.246 mL, 0.246 mmol). After being stirred for 1 h between -65 to -78 °C, the reaction mixture was treated with MeOH (0.5 mL) and a Rochelle's salt solution (5.0 mL). The resulting mixture was warmed to ambient temperature and extracted twice with MTBE (15 mL). The combined organic layers were washed 30% aqueous NaCl (5.0 mL) and dried over MgSO₄. Filtration and concentration *in vacuo* provided the target product (compound **36**, 68mg), which was used in the next step without further purification. ¹H NMR (400 MHz, CDCl₃) δ ppm 0.65 (q, *J*=8.1 Hz, 6 H) 0.97 (t, *J*=8.2 Hz, 9 H) 1.02 (s, 9 H) 1.04 (s, 9 H) 1.06 - 1.10 (m, 6 H) 1.45 - 1.52 (m, 2 H) 1.57 - 1.70 (m, 4 H) 1.72 - 1.83 (m, 3 H) 1.84 - 1.98 (m, 2 H) 2.08 - 2.22 (m, 2 H) 2.34 - 2.42 (m, 1 H) 2.68 - 2.76 (m, 1 H) 3.20 - 3.28 (m, 2 H) 3.43 - 3.50 (m, 2 H) 3.56 - 3.62 (m, 2 H) 3.64 - 3.71 (m, 1 H) 3.73 - 3.85 (m, 3 H) 4.14 (br s, 2 H) 4.40 - 4.43 (m, 1 H) 4.49 (q, *J*=6.6 Hz, 1 H) 4.78 (s, 1 H) 4.83 (d, *J*=1.9 Hz, 1 H) 4.86 (s, 1 H) 4.97 (d, *J*=1.9 Hz, 1 H) 5.70-5.76 (m, 1 H) 5.83-5.89 (m, 1 H) 7.30 - 7.43 (m, 12 H) 7.61 - 7.72 (m, 8 H)

(((2R,3R,4aS,6S,7R,8S,8aS)-6-(((2R,4R,6S)-6-(2-((2S,5R)-5-((E)-3-(allyloxy)prop-1-en-1-yl)-3-methylenetetrahydrofuran-2-yl)ethyl)-4-methyl-3-methylenetetrahydro-2H-pyran-2-yl)methyl)-2-(2-((tert-butyl)diphenylsilyl)oxy)ethyl)-8-methyl-7-((triethylsilyl)oxy)octahydropyrano[3,2-b]pyran-3-yl)oxy)(tert-butyl)diphenylsilane




To a solution of (E)-3-((2R,5S)-5-(2-((2S,4R,6R)-6-(((2S,3R,4S,4aS,6R,7R,8aS)-7-((tert-butyl)diphenylsilyl)oxy)-6-(2-((tert-butyl)diphenylsilyl)oxy)ethyl)-4-methyl-3-((triethylsilyl)oxy)octahydropyrano[3,2-b]pyran-2-yl)methyl)-4-methyl-5-methylenetetrahydro-2H-pyran-2-yl)ethyl)-4-methylenetetrahydrofuran-2-yl)prop-2-en-1-ol (compound **36**, 68 mg, 0.061 mmol) in THF (2.0 mL) at 0 °C was added sodium hydride (60% in oil, 6.1 mg, 0.15 mmol). The resulting slurry was stirred at ambient temperature for 20 min. Allyl bromide (0.032 mL, 0.37 mmol) and DMF (0.5 mL, 6.5 mmol) were then added. The resulting mixture was stirred for 16 h at ambient temperature. Saturated aqueous NH₄Cl (27% (w/v)) (5.0 mL), water (1.0 mL), and MTBE (10 mL) were added. The layers were separated, and the aqueous layer was extracted with MTBE (10 mL). The combined organic layers were washed with 30% aqueous NaCl (3.0 mL) and dried over MgSO₄. Filtration and concentration *in vacuo* provided 71 mg of the target product. Without further purification, the crude product (((2R,3R,4aS,6S,7R,8S,8aS)-6-(((2R,4R,6S)-6-(2-((2S,5R)-5-((E)-3-(allyloxy)prop-1-en-1-yl)-3-methylenetetrahydrofuran-2-yl)ethyl)-4-

methyl-3-methylenetetrahydro-2H-pyran-2-yl)methyl)-2-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-8-methyl-7-((triethylsilyl)oxy)octahydropyrano[3,2-b]pyran-3-yl)oxy)(tert-butyl)diphenylsilane (72 mg) was dissolved in acetic acid (4.0 mL) and treated with water (1.0 mL) at ambient temperature. After 5 h, the reaction mixture was concentrated *in vacuo*. The resulting residue was purified by silica gel column

chromatography using a 20 - 33% gradient of ethyl acetate in *n*-heptane as eluent to give 54 mg of the target product (compound **30**). ¹H NMR (400 MHz, CDCl₃) δ ppm 1.00 (s, 9 H) 1.04 - 1.06 (m, 9 H) 1.08 - 1.15 (m, 6 H) 1.45 - 1.76 (m, 11 H) 1.80 (br. s., 1 H) 1.91 - 2.15 (m, 5 H) 2.24 - 2.34 (m, 1 H) 2.36 - 2.45 (m, 1 H) 2.67 - 2.76 (m, 1 H) 3.22 (br. s., 1 H) 3.33 (br. s., 1 H) 3.49 (d, *J*=7.8 Hz, 1 H) 3.55 - 3.65 (m, 2 H) 3.68 (br. s., 1 H) 3.72 - 3.83 (m, 3 H) 3.91 (t, *J*=5.9 Hz, 1 H) 3.95 - 3.99 (m, 4 H) 4.07 (q, *J*=5.6 Hz, 1 H) 4.41 - 4.53 (m, 2 H) 4.81 (br. s., 1 H) 4.81 - 4.84 (m, 1 H) 4.88 (d, *J*=2.0 Hz, 1 H) 4.98 (d, *J*=2.0 Hz, 1 H) 5.18 (d, *J*=10.2 Hz, 1 H) 5.27 (dd, *J*=17.2, 1.6 Hz, 1 H) 5.73 (dd, *J*=16.0, 5.1 Hz, 1 H) 5.79 (dd, *J*=15.2, 5.1 Hz, 1 H) 5.85 - 5.97 (m, 1 H) 7.30 - 7.43 (m, 12 H) 7.57 - 7.79 (m, 8 H)


(2S,3R,4S,4aS,6R,7R,8aS)-2-(((2R,4R,6S)-6-(2-((2S,5R)-5-((E)-3-(allyloxy)prop-1-en-1-yl)-3-methylenetetrahydrofuran-2-yl)ethyl)-4-methyl-3-methylenetetrahydro-2H-pyran-2-yl)methyl)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-4-methyloctahydropyrano[3,2-b]pyran-3-yl 2-((2R,3S,3aR,4aS,7R,8aS,9S,9aR)-3,9-bis((tert-butyldiphenylsilyl)oxy)-2-(2-((triethylsilyl)oxy)but-3-en-1-yl)decahydrofuro[3,2-b]pyrano[2,3-e]pyran-7-yl)acetate




To a solution of 2-((2R,3S,3aR,4aS,7R,8aS,9S,9aR)-3,9-bis((tert-butyldiphenylsilyl)oxy)-2-(2-((triethylsilyl)oxy)but-3-en-1-yl)decahydrofuro[3,2-b]pyrano[2,3-e]pyran-7-yl)acetic acid (compound **24**, 66 mg, 0.071 mmol) in dichloromethane (2 mL) at ambient temperature were added TEA (0.022 mL, 0.156 mmol), 2-methyl-6-nitrobenzoic anhydride (26.8 mg, 0.078 mmol) and 4-dimethylaminopyridine (3.17 mg, 0.026 mmol). After 10 min, the resulting solution was added to a solution of (2S,3R,4R,4aS,6R,7R,8aS)-2-(((2R,4R,6S)-6-(2-((2S,5R)-5-((E)-3-(allyloxy)prop-1-en-1-yl)-3-methylenetetrahydrofuran-2-yl)ethyl)-4-methyl-3-methylenetetrahydro-2H-pyran-2-yl)methyl)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-4-methyloctahydropyrano[3,2-b]pyran-3-ol (compound **30**, 54 mg, 0.052 mmol) in dichloromethane (1.0 mL). After being stirred overnight, the reaction mixture was treated with a saturated aqueous NH₄Cl solution (5.0 mL). The resulting mixture was stirred for 30 min and extracted twice with MTBE (10 mL each). The combined organic layers were washed with 30% aqueous NaCl (5.0 mL) and dried over MgSO₄. Filtration, concentration *in vacuo*, and purification by silica gel column chromatography using a 10 - 33% gradient of ethyl acetate in *n*-heptane as eluent provided 87 mg of the target product (compound **31**). ¹H NMR (1:1 diastereomeric mixture, 400 MHz, CDCl₃) δ ppm 0.46 - 0.62 (m, 6 H) 0.83 - 0.98 (m, 12 H) 1.00 (s, 9 H) 1.06 (s, 9 H) 1.07 - 1.15 (m, 6 H) 1.25 - 2.04 (m, 19 H) 2.08-

2.24 (m., 2.5 H) 2.32 - 2.44 (m, 1.5 H) 2.66 - 2.76 (m, 1 H) 2.86 - 2.96 (m, 1 H) 3.22 - 3.28 (m, 1 H) 3.37 - 3.42 (m, 0.5 H) 3.46 - 3.71 (m, 6.5 H) 3.73 - 3.86 (m, 4 H) 3.93 - 4.00 (m, 4 H) 4.05 - 4.23 (m., 3 H) 4.27 - 4.34 (m, 1 H) 4.40 - 4.46 (m, 1 H) 4.49 (q, $J=6.4$ Hz, 1 H) 4.58 - 4.68 (m, 1 H) 4.70 (s, 1 H) 4.76 (s, 1 H) 4.84 (br. s., 1 H) 4.90 - 5.14 (m, 2 H) 4.95 - 4.97 (m, 1 H) 5.17 (d, $J=10.6$ Hz, 1 H) 5.27 (dd, $J=17.4$, 1.8 Hz, 1 H) 5.59 - 5.70 (m, 1 H) 5.73 (dd, $J=15.6$, 5.1 Hz, 1 H) 5.80 (dd, $J=15.6$, 5.1 Hz, 1 H) 5.84 - 5.97 (m, 1 H) 7.27 - 7.43 (m, 24 H) 7.55 - 7.79 (m, 16H)

Compound **31** can be transformed into compound **3** according to the following sequence:



Ring-Closing Olefin Metathesis




- To a solution of (2S,3R,4S,4aS,6R,7R,8aS)-2-(((2R,4R,6S)-6-(2-(((2S,5R)-5-((E)-3-(allyloxy)prop-1-en-1-yl)-3-methylenetetrahydrofuran-2-yl)ethyl)-4-methyl-3-methylenetetrahydro-2H-pyran-2-yl)methyl)-7-((tert-butylidiphenylsilyl)oxy)-6-(2-((tert-butylidiphenylsilyl)oxy)ethyl)-4-methyloctahydropyrano[3,2-b]pyran-3-yl 2-((2R,3S,3aR,4aS,7R,8aS,9S,9aR)-3,9-bis((tert-butylidiphenylsilyl)oxy)-2-(2-((triethylsilyl)oxy)but-3-en-1-yl)decahydrofuro[3,2-b]pyrano[2,3-e]pyran-7-yl)acetate (compound **31**, 87 mg, 0.044 mmol) in toluene (61 mL) at 80 °C was added quinone (4.8 mg, 0.044 mmol) and a solution of Hoveyda-Grubbs 2nd generation catalyst (5.59 mg, 8.891 μmol) in toluene (10 mL). After being stirred at 80 °C for 8h, the reaction mixture was cooled to ambient temperature and treated with DMSO (0.063 mL, 0.889 mmol). The resulting solution was stirred overnight and concentrated *in vacuo*. Silica gel column chromatography of the residue using a 20 - 33% gradient of ethyl acetate in *n*-heptane as eluent afforded 71 mg of a mixture of the target product (compound **33a**) and compound **32a**: MS *m/z* 1908.0 and 1936.0 [M+Na]⁺. The mixture was dissolved in THF (2.0 mL) at ambient temperature. To the solution was added acetic acid (3.2 mL) and water (0.8 mL). After being stirred for 3 h at ambient temperature, the resulting mixture was concentrated *in vacuo*. The residue was purified by silica gel column chromatography using a 10 - 33% gradient of ethyl acetate in *n*-heptane as eluent to give 46 mg of a mixture of compound **33** and compound **32**. The mixture was dissolved in dichloromethane (45 mL) and treated with Hoveyda-Grubbs 2nd generation catalyst (2.91 mg, 4.63 μmol) at reflux (45°C oil bath) over 15 h. The reaction mixture was cooled to ambient temperature and concentrated *in vacuo*. Purification by silica gel column chromatography of the residue using a 15 - 33% gradient of ethyl acetate in *n*-heptane as eluent afforded 15 mg of the target product (compound **33**). ¹H NMR (major isomer, 400 MHz, CDCl₃) δ ppm 0.99 - 1.10

(m, 42 H) 1.25-2.27 (m, 19 H) 2.32-2.42 (m, $J=7.4$ Hz, 3 H) 2.45 - 2.51 (m, 1 H) 2.55 (dd, $J=14.9$, 3.1 Hz, 1 H) 2.69 - 2.79 (m, 1 H) 2.85 (t, $J=9.0$ Hz, 1 H) 3.17 - 3.27 (m, 1 H) 3.33 - 3.55 (m, 4 H) 3.59 - 3.81 (m, 11 H) 3.87 - 3.95 (m, 1 H) 4.15 (dd, $J=8.6$, 3.9 Hz, 1 H) 4.29 - 4.39 (m, 2 H) 4.41 - 4.48 (m, 1 H) 4.49 - 4.55 (m, 1 H) 4.58 - 4.67 (m, 2 H) 4.76 (s, 1 H) 4.83 (s, 1 H) 4.92 (br. s., 2 H) 5.77 (ddd, $J=15.6$, 7.6, 1.4 Hz, 1 H) 6.31 (dd, $J=15.4$, 4.5 Hz, 1 H) 7.30 - 7.43 (m, 24 H) 7.58 - 7.77 (m, 16 H)

(2S,3R,4S,4aS,6R,7R,8aS)-7-((tert-butyldiphenylsilyl)oxy)-6-(2-((tert-butyldiphenylsilyl)oxy)ethyl)-4-methyl-2-(((2R,4R,6S)-4-methyl-3-methylene-6-(2-((2S,5R)-3-methylene-5-vinyltetrahydrofuran-2-yl)ethyl)tetrahydro-2H-pyran-2-yl)methyl)octahydropyrano[3,2-b]pyran-3-yl 2-(((2R,3S,3aR,4aS,7R,8aS,9S,9aR)-3,9-bis((tert-butyldiphenylsilyl)oxy)-2-(2-hydroxybut-3-en-1-yl)decahydrofuro[3,2-b]pyrano[2,3-e]pyran-7-yl)acetate

^1H NMR (compound **32**, 400 MHz, CDCl_3) δ ppm 1.00-1.11 (m, 42 H) 1.25 - 2.03 (m, 19 H) 2.12 - 2.22 (m, 2 H) 2.33 - 2.42 (m, 2 H) 2.72 (ddd, $J=15.3$, 6.7, 1.8 Hz, 1 H) 2.95 (dd, $J=9.8$, 4.3 Hz, 1 H) 3.24 (dd, $J=5.7$, 3.7 Hz, 1 H) 3.47 - 3.56 (m, 3 H) 3.59 - 3.69 (m, 5 H) 3.72 - 3.80 (m, 3 H) 3.81 - 3.88 (m, 1 H) 3.99 - 4.08 (m, 1 H) 4.10 - 4.20 (m, 2 H) 4.25 (dd, $J=6.3$, 4.3 Hz, 1 H) 4.41 - 4.50 (m, 2 H) 4.67 (t, $J=8.0$ Hz, 1 H) 4.72 (s, 1 H) 4.78 (s, 1 H) 4.85 (d, $J=2.0$ Hz, 1 H) 4.96 (d, $J=2.0$ Hz, 1 H) 5.04 (d, $J=10.2$ Hz, 1 H) 5.10 (d, $J=10.2$ Hz, 1 H) 5.16 (d, $J=16.8$ Hz, 1 H) 5.23 (d, $J=17.2$ Hz, 1 H) 5.72 (ddd, $J=16.9$, 10.6, 6.1 Hz, 1 H) 5.84 (ddd, $J=17.0$, 10.4, 6.3 Hz, 1 H) 7.27 - 7.45 (m, 24 H) 7.56 - 7.78 (m, 16 H)


Compound **37**



Compound **33** (25.5 mg, 0.015 mmol) was dissolved in dichloromethane (1.0 ml) at ambient temperature. Sodium bicarbonate (12.3 mg, 0.146 mmol) and Dess-Martin periodinane (18.6 mg, 0.044 mmol) were added, and the resulting mixture was stirred at ambient temperature overnight. MTBE (10.0 ml), water (2.0 mL) and a saturated aqueous sodium thiosulfate solution (5 mL) were added. The resulting mixture was stirred at ambient temperature over 30 min. The organic layer was separated out, washed with 30% aqueous NaCl (3.0 mL), and dried over MgSO_4 . Filtration, concentration *in vacuo*, and purification by silica gel column chromatography using a 10 - 33% gradient of ethyl acetate in *n*-heptane as eluent provided 17 mg of the target product (compound **37**). ^1H NMR (400 MHz, CDCl_3) δ ppm 0.97 - 1.17 (m, 42 H) 1.28 - 2.25 (m, 20 H) 2.49 (dd, $J=16.0$, 7.0 Hz, 1 H) 2.56 (dd, $J=15.0$, 10.0 Hz, 1 H) 2.72 (dd, $J=15.2$, 3.5 Hz, 1 H) 2.75 - 2.82 (m, 1 H) 2.98 (t, $J=8.8$ Hz, 1 H) 3.19 - 3.25 (m, 1 H) 3.36 (d, $J=12.5$ Hz, 1 H) 3.42 (t, $J=4.3$ Hz, 1 H) 3.48 - 3.57 (m, 4 H) 3.60 - 3.70 (m, 5 H) 3.71 - 3.82 (m, 3 H) 3.90 (td, $J=10.4$, 3.9 Hz, 1 H) 4.01 (ddd, $J=10.6$, 7.4, 3.1 Hz, 1 H) 4.08 (dd, $J=8.0$, 4.5 Hz, 1 H) 4.24 - 4.35 (m, 2 H) 4.55 -


4.59 (m, 1 H) 4.62 (t, $J=7.4$ Hz, 1 H) 4.66 - 4.73 (m, 1 H) 4.76 (s, 1 H) 4.85 (s, 1 H) 4.91 (s, 1 H) 5.00 (d, $J=1.6$ Hz, 1 H) 6.41 (d, $J=16.0$ Hz, 1 H) 6.86 (dd, $J=16.2, 5.7$ Hz, 1 H) 7.28 - 7.43 (m, 24 H) 7.57 - 7.77 (m, 16 H)

5 Compound 34



Compound **37** (4.0 mg, 2.3 μ mol) was dissolved in deoxygenated toluene (0.6 mL) at ambient temperature. Deoxygenated (purged with nitrogen for 40 min) water (2.4 μ L, 0.13 mmol) was added followed by hydrido(triphenylphosphine)copper(I) hexamer (9.0 mg, 4.6 μ mol). After being stirred for 1 h, the reaction mixture was treated with air. Copper-containing decomposition products precipitated. Concentration *in vacuo* followed by purification by column chromatography using a 10 - 33% gradient of ethyl acetate in *n*-heptane as eluent afforded 3.0 mg of the target product (compound **34**). ^1H NMR (400 MHz, CDCl_3) δ ppm 0.98 - 1.14 (m, 42 H) 1.28 - 2.40 (m, 22 H) 2.55 - 2.77 (m, 4 H) 2.81 - 2.90 (m, 1 H) 2.91 (t, $J=9.2$ Hz, 1 H) 3.06 (t, $J=11.7$ Hz, 1 H) 3.20 (dd, $J=6.3, 3.9$ Hz, 1 H) 3.39 - 3.47 (m, 2 H) 3.50 - 3.57 (m, 2 H) 3.62 - 3.69 (m, 3 H) 3.72 (dd, $J=6.1, 4.5$ Hz, 1 H) 3.75 - 3.85 (m, 3 H) 3.94 - 4.02 (m, 1 H) 4.09 (dd, $J=8.4, 3.7$ Hz, 2 H) 4.26 - 4.32 (m, 1 H) 4.33 - 4.39 (m, 1 H) 4.43 - 4.48 (m, 1 H) 4.64 (t, $J=7.0$ Hz, 1 H) 4.77 (s, 1 H) 4.84 (s, 1 H) 4.86 (s, 1 H) 4.93 (s, 1 H) 7.28 - 7.44 (m, 24 H) 7.57 - 7.77 (m, 16 H)

20 Compound 3




To compound **34** (3.0 mg, 1.72 μ mol) in a vial was added THF (0.6 mL) and *N,N*-dimethylacetamide (0.21 mL) at ambient temperature. A mixture of TBAF (1.0 M in THF, 52 μ L, 0.052 mmol) and imidazole hydrochloride (2.7 mg, 0.026 mmol) was added, and the resulting mixture was stirred over 3 days at

ambient temperature. 30% aqueous NaCl (2.0 mL) was added, and the resulting mixture was extracted twice with a mixture of THF (5.0 mL) and toluene (5.0 mL). The combined organic layers were concentrated with a stream of nitrogen. The residue was dissolved in dichloromethane (1.0 mL) at ambient temperature, and PPTS (18 mg, 72 μ mol) was added. Once all starting material was consumed (2 h), the reaction mixture was purified by silica gel column chromatography using a 0 - 10% gradient of methanol in ethyl acetate as eluent to give 0.9 mg of the target product (compound **3**). The structure was confirmed by ^1H NMR comparison with the reported spectrum. ^1H NMR (400 MHz, CDCl_3) δ ppm 1.10 (d, $J=6.3$ Hz, 3 H) 1.18 (d, $J=7.8$ Hz, 3 H) 1.27 - 2.40 (m, 25 H) 2.41 (dd, $J=16.8, 2.7$ Hz, 1 H) 2.53 (dd, $J=17.2, 10.2$ Hz, 1 H) 2.76 - 2.86 (m, 1 H) 2.90 (dd, $J=9.8, 2.3$ Hz, 1 H) 3.29 (s, 1 H) 3.52 - 3.57 (m, 2 H) 3.64 (d, $J=10.9$ Hz, 1 H) 3.69 - 3.89 (m, 5 H) 4.04 (dd, $J=6.3, 4.3$ Hz, 1 H) 4.10 - 4.15 (m, 1 H) 4.18 (dd, $J=6.4, 4.5$ Hz, 1 H) 4.21 - 4.28 (m, 1 H) 4.36 (d, $J=11.7$ Hz, 1 H) 4.38 - 4.44 (m, 2 H) 4.50 (t, $J=2.3$ Hz, 1 H) 4.58 - 4.62 (m, 1 H) 4.68 (t, $J=4.5$ Hz, 1 H) 4.80 (br. s., 2 H) 4.98 (br. s., 1 H) 5.08 (s, 1 H)

Example 6 – C.19-C.20 Macrocyclization through Nozaki-Hiyama-Kishi Reaction


A halichondrin macrolide **3** can be prepared according to the below sequence.



Compound **38** can be converted to compound **39** through the Nozaki-Hiyama-Kishi reaction (e.g., with Cr(II) and Ni(II) salts). Compound **39** can give compound **40** through nucleophilic ring-closing substitution reaction. Compound **40** can be converted to compound **3** after global deprotection (e.g., removal of silyl groups with a fluoride source, such as TBAF).

Example 7 – C.26-C.27 Macrocyclization through Nozaki-Hiyama-Kishi Reaction


A halichondrin macrolide **3** can be prepared according to the below sequence.



- 5 Compound **41** can be converted to compound **42** through the Nozaki-Hiyama-Kishi reaction (e.g., with Cr(II) and Ni(II) salts) with subsequent silylation of the formed hydroxyl by chlorotriethylsilane. Compound **42** can be subjected to Vasella fragmentation conditions to provide **43**. DDQ deprotection of the MPM-group and selective oxidation of the resulting allylic alcohol by manganese (II) oxide can provide **44**. The remaining hydroxyl group can be reacted with methanesulfonyl chloride to give compound **45**.
- 10 Global desilylation of **45** with tetrabutylammonium fluoride (TBAF) can form the compound **46** by nucleophilic displacement of the mesylate and conjugate addition of a deprotected alcohol to an enone. Compound **46** can be reacted with a Brønsted acid (e.g., PPTs) to afford compound **3**.

Example 8 – C.14-C.38 Halichondrin Intermediate through Allene-Prins reaction.

A halichondrin C.14-C.38 intermediate **53** can be prepared according to the below sequence.



- 5 Compound **47** (Aicher et al. *Tetrahedron Letters*, 33:1549-1552, 1992) can be converted to **48** in a three step process [(1) diisobutylaluminum hydride reduction to the aldehyde, (2) Acid catalyzed acetonide deprotection and dimethylacetal formation and (3) benzoylation]. The allene-Prins reaction between **48** and **49** can be achieved by treatment with boron trifluoride diethyl etherate and methoxyacetic acid to provide **50**. Allylic reduction of the methoxyacetate can be accomplished using
- 10 tetrakis(triphenylphosphine)palladium(0) with formic acid and triethylamine resulting in **51**. Deprotection of the benzoate groups can be achieved by treatment with magnesium methoxide to provide **52**. Finally, treatment of **52** with tert-butyldimethylsilyl chloride can provide the known C.14-C.38 halichondrin intermediate **53** (Aicher et al. *J. Am. Chem. Soc.*, 114:3162-3164, 1992).

15 **Other Embodiments**


Various modifications and variations of the described compositions and methods of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various

20 modifications of the described modes for carrying out the invention that are obvious to those skilled in the art are intended to be within the scope of the invention.

Other embodiments are in the claims.

Claims

1. A method of preparing a macrocyclic intermediate in the synthesis of a halichondrin macrolide, said method comprising performing a macrocyclization reaction on a non-macrocyclic intermediate, wherein (1) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.2-C.3 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with an organic base and a Lewis acid, wherein said non-macrocyclic intermediate is (1) a compound of formula (IA):




or a salt or a tautomer thereof,

wherein

each R is independently optionally substituted alkyl or optionally substituted aryl;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-

dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;


or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_V))-$, wherein P_V is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and


P_5 is H or a hydroxyl protecting group; and

wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IB):



or a salt or a tautomer thereof;

(2) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.2-C.3 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with an olefin metathesis catalyst, wherein said non-macrocyclic intermediate is a compound of formula (IIA):




or a salt or a tautomer thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to

10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OPⁿ, where Pⁿ is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;


or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OPⁿ;

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and


P₅ is H or a hydroxyl protecting group; and

wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IB):



or a salt or a tautomer thereof;


(3) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.3-C.4 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with an olefin metathesis catalyst, wherein said non macrocyclic intermediate is a compound of formula (IIIA):



or a salt or a tautomer thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;


or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_V))- , wherein P_V is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and


P₅ is H or a hydroxyl protecting group; and

wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IIIB):



or a salt or a tautomer thereof;


(4) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.12-C.13 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with an olefin metathesis catalyst, wherein said non-macrocyclic intermediate is a compound of formula (IVA):



or a salt thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(i) R₃ is H or a hydroxyl protecting group, R₄ is alkyl ether, and R₅ is H;


(ii) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

- (iii) R_3 and R_4 combine to form a bond, and R_5 is H;
 each P_4 and P_5 is independently H or a hydroxyl protecting group;
 R_6 is H, and P_6 is H or a hydroxyl protecting group; or R_6 and P_6 combine to form a double bond;


and

wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IVB):



or a salt thereof;


- (5) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.15-C.16 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with an olefin metathesis catalyst, wherein said non-macrocyclic intermediate is a compound of formula (VA):



or a salt or a tautomer thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;


G is O, S, CH_2 , or NR_N , wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;


n is 0, 1, or 2;

- (a1) R₃ is H or a hydroxyl protecting group, R₄ is alkyl ether, and R₅ is H;
(a2) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;
or
(a3) R₃ and R₄ combine to form a bond, and R₅ is H;
(b1) A₁ and R₇ combine to form oxo, R₆ is H or a hydroxyl protecting group, and R₈ is H;
or
(b2) A₁ is H or OP^{''}, where P^{''} is H or a hydroxyl protecting group, and:
(i) R₆ is H or a hydroxyl protecting group, and R₇ and R₈ combine to form a double bond;
or
(ii) R₆ and R₇ combine to form a bond, and R₈ is H or OP^{''};
(c1) R₉ is H, and P₄ is H or a hydroxyl protecting group;
or
(c2) R₉ and P₄ combine to form a double bond;
R₁₀ is H or -CH₂X₁CH₂CH=CH₂, wherein X₁ is O, -C(R₁₁)₂-, or NP₆, and wherein each R₁₁ is independently H or -COOR₁₂, P₆ is an *N*-protecting group, and R₁₂ is alkyl;
each P₅ is independently H or a hydroxyl protecting group; and
wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (VB):



or a salt or a tautomer thereof;

(6) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.19-C.20 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with a Cr(II) salt and a Ni(II) salt, wherein said non-macrocyclic intermediate is a compound of formula (VIA):




or a salt or a tautomer thereof,

wherein

Y is iodide, bromide, or trifluoromethanesulfonate;

b designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or *b* designates (*S*)-stereogenic center, and Z is OR₆, wherein R₆ is a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:


(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (IVB):



(VIB)

or a salt or a tautomer thereof,


wherein

b designates (*R*)-stereogenic center, *c* designates (*S*)-stereogenic center, and *Z* is a sulfonate, chloride, bromide, or iodide;

or

b designates (*S*)-stereogenic center, *c* designates (*R*)-stereogenic center, and *Z* is OR₆, wherein R₆ is a hydroxyl protecting group; or

(7) said macrocyclization reaction produces said macrocyclic intermediate by forming a C.26-C.27 bond in the structure of the halichondrin macrolide, said performing said macrocyclization reaction comprising contacting said non-macrocyclic intermediate with a Cr(II) salt and a Ni(II) salt, wherein said non-macrocyclic intermediate is a compound of formula (VIIA):



(VIIA)


or a salt thereof,

wherein

Y is iodide, bromide, or trifluoromethanesulfonate;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to

10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(a1) R_3 is H or a hydroxyl protecting group, R_4 and R_5 combine to form a double bond, each P_4 is independently H or a hydroxyl protecting group, and X_1 , together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$, wherein R_8 is H or a hydroxyl protecting group;

or

(a2) R_3 and R_4 combine to form a bond, R_5 is H, and each P_4 is independently H or a hydroxyl protecting group, and X_1 , together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$;

or

both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal;


(b1) Z is chloride, bromide, or iodide, and R_6 and R_7 combine to form a bond;

or

(b2) Z and R_7 combine to form a double bond, and R_6 is a hydroxyl protecting group;

and

wherein said macrocyclic intermediate in the synthesis of a halichondrin macrolide is a compound of formula (VIIB):



(VIIB)

or a salt thereof,

wherein each of P_5 and R_6 is independently H or a hydroxyl protecting group.

2. The method of claim 1, wherein for (1),

(a) each R is optionally substituted alkyl;

(b) both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal;

(c) P_5 is a hydroxyl protecting group;

(d) R_3 and R_4 combine to form a bond, and R_5 is H;

(e) said organic base is DBU or trimethylamine; and/or

(f) said Lewis acid is a salt of Li or Zn.

3. The method of claim 1, wherein for (2) or (3),

(a) both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and/or

(b) P₅ is a hydroxyl protecting group.

4. The method of claim 1, wherein for (4), at least one of P₄ and P₅ is H.

5. The method of claim 1, wherein for (5),

(a) R₉ is H, and P₄ is H;

(b) R₁₀ is -CH₂X₁CH₂CH=CH₂, and X₁ is O;

(c) R₆ and R₇ combine to form a bond, and R₈ is H;

(d) each P₅ is independently a hydroxyl protecting group

(e) at least one P₅ is H; and/or

(f) R₃ and R₄ combine to form a bond, and R₅ is H.

6. The method of claim 1, wherein for (2), (3), (4), or (5), said olefin metathesis catalyst is a ruthenium-carbene complex.

7. The method of claim 1, wherein for (6),

(a) Y is bromide;

(b) R₃ and R₄ combine to form a bond, and R₅ is H;

(c) both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal;

(d) Z is a sulfonate;

(e) Z is OR₆, wherein R₆ is a hydroxyl protecting group; and/or

(f) Z is an ester, carbonate, or carbamate.

8. The method of claim 1, wherein for (7),

(a) Z is iodide;


(b) Y is trifluoromethanesulfonate;

(c) R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms -(CH(OR₈))- , wherein R₈ is H or a hydroxyl protecting group;

(d) at least one of P₃, P₄, and R₆ is a hydroxyl protecting group; and/or

(e) Z is chloride, bromide, or iodide, and R₆ and R₇ combine to form a bond.

9. A method of preparing:




halichondrin macrolide

or a salt thereof,

wherein

each of A_1 and A_2 is independently H or OP'' , where each P'' is independently H or a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




(1),

wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;


each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(1) said method comprising:

(A) producing a compound of formula (IB) or a salt or a tautomer thereof from a compound of formula (IA) or a salt or a tautomer thereof, the compound of formula (IA) having the following structure:



wherein

each R is independently optionally substituted alkyl or optionally substituted aryl;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP^{''}, where P^{''} is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or


(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP^{''};

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))-, wherein P_Y is H or a hydroxyl protecting group; or

both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

P₅ is H or a hydroxyl protecting group; and

the compound of formula (IB) having the following structure:




and

(B) producing the halichondrin macrolide or the salt thereof from compound (IB) or the salt or the tautomer thereof;

(2) said method comprising:

(A) producing a compound of formula (IB) or the salt or the tautomer thereof from a compound of formula (IIA), the compound of formula (IIA) or the salt or the tautomer thereof having the following structure:



wherein

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;


or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and

P_5 is H or a hydroxyl protecting group; and

the compound of formula (IB) having the following structure:




(IB);

and

(B) producing the halichondrin macrolide or the salt thereof from compound (IB) or the salt or the tautomer thereof;

(3) said method comprising:

(A) producing a compound of formula (IIIB) or the salt or the tautomer thereof from a compound of formula (IIIA) or the salt or the tautomer thereof, the compound of formula (IIIA) having the following structure:



wherein

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OPⁿ, where Pⁿ is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;


or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OPⁿ;

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_γ))-, wherein P_γ is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

P₅ is H or a hydroxyl protecting group; and

the compound of formula (IIIB) having the following structure:




and

(B) producing the halichondrin macrolide or the salt thereof from the compound of formula (IIIB) or the salt or the tautomer thereof;

(4) said method comprising:

(A) producing a compound of formula (IVB) or a salt thereof from a compound of formula (IVA) or a salt thereof, the compound of formula (IVA) having the following structure:



wherein

- (i) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;
 - (ii) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;
- or


- (iii) R_3 and R_4 combine to form a bond, and R_5 is H;

each P_4 and P_5 is independently H or a hydroxyl protecting group;

R_6 is H, and P_6 is H or a hydroxyl protecting group; or R_6 and P_6 combine to form a double bond;

and

the compound of formula (IVB) having the following structure:




and

(B) producing the halichondrin macrolide or the salt thereof from the compound of formula (IVB) or the salt thereof;

(5) said method comprising:

(A) producing a compound of formula (VB) or a salt or a tautomer thereof from a compound of formula (VA) or a salt or a tautomer thereof, the compound of formula (VA) having the following structure:



wherein

- (a1) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;
- (a2) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;
- or
- (a3) R_3 and R_4 combine to form a bond, and R_5 is H;
- (b1) A_1 and R_7 combine to form oxo, R_6 is H or a hydroxyl protecting group, and R_8 is H;
- or
- (b2) A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:
 - (i) R_6 is H or a hydroxyl protecting group, and R_7 and R_8 combine to form a double bond;
- or


(ii) R_6 and R_7 combine to form a bond, and R_8 is H or OP^r ;

(c1) R_9 is H, and P_4 is H or a hydroxyl protecting group;

or


(c2) R_9 and P_4 combine to form a double bond;

R_{10} is H or $-CH_2X_1CH_2CH=CH_2$, wherein X_1 is O, $-C(R_{11})_2-$, or NP_6 , and wherein each R_{11} is independently H or $-COOR_{12}$, P_6 is an *N*-protecting group, and R_{12} is alkyl; each P_5 is independently H or a hydroxyl protecting group; and the compound of formula (VB) having the following structure:



(VB);


(B) producing a compound of formula (VC) or a salt thereof from the compound of formula (VB) or the salt or the tautomer thereof, the compound of formula (VC) having the following structure:



(VC),

wherein each of A_1 and A_2 is independently H or OP^r ;

(C) producing a compound of formula (IVC) or a salt thereof from the compound of formula (VC) or a salt thereof, the compound of formula (IVC) having the following structure:




(IVC),

wherein each of A₁ and A₂ is independently H or OP^{''}; and

(D) producing the halichondrin macrolide or a salt thereof from the compound of formula (IVC) or a salt thereof;

(6) said method comprising:

(A) producing a compound of formula (VIB) or a salt or a tautomer thereof from a compound of formula (VIA) or a salt or a tautomer thereof, the compound of formula (VIA) having the following structure:



(VIA),

wherein

Y is iodide, bromide, or trifluoromethanesulfonate;

b designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or *b* designates (*S*)-stereogenic center, and Z is OR₆, wherein R₆ is a hydroxyl protecting group;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP^{''}, where P^{''} is H or a hydroxyl protecting group, and:


- (i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

- (ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

and each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal;

the compound of formula (VIB) having the following structure:



(VIB),

wherein

b designates (*R*)-stereogenic center, c designates (*S*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide;


or

b designates (*S*)-stereogenic center, c designates (*R*)-stereogenic center, and Z is OR_6 , wherein R_6 is a hydroxyl protecting group;

(B) producing the halichondrin macrolide or the salt thereof from the compound of formula (VIB) or the salt or the tautomer thereof; or

(7) said method comprising:

(A) producing a compound of formula (VIIB) or the salt thereof from a compound of formula (VIIA) or the salt thereof, the compound of formula (VIIA) having the following structure:



(VIIA),

wherein

Y is iodide, bromide, or trifluoromethanesulfonate;

(a1) R_3 is H or a hydroxyl protecting group, R_4 and R_5 combine to form a double bond, each P_4 is independently H or a hydroxyl protecting group, and X_1 , together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$;

or

(a2) R_3 and R_4 combine to form a bond, R_5 is H, and each P_4 is independently H or a hydroxyl protecting group, and X_1 , together with the carbon to which it is attached, forms a carbonyl or $-(CH(OR_8))-$;

or

both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal;

(b1) Z is chloride, bromide, or iodide, and R_6 and R_7 combine to form a bond;


or

(b2) Z and R_7 combine to form a double bond, and R_6 is a hydroxyl protecting group;

and

wherein R_8 is H or a hydroxyl protecting group


the compound of formula (VIIB) having the following structure:



(VIIB),

wherein each of P_5 and R_6 is H or a hydroxyl protecting group;

(B) producing a compound of formula (VIIC) or a salt or a tautomer thereof from the compound of formula (VIIB) or the salt thereof, the compound of formula (VIIC) having the following structure:



(VIIC),

wherein A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or


A₁ is H or OP^{''}, where P^{''} is H or a hydroxyl protecting group, and:

- (i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

- (ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP^{''};

(C) producing a compound of formula (VIID) or a salt thereof from the compound of formula (VIIC) or the salt or the tautomer thereof, the compound of formula (VIID) having the following structure:



(VIID),

wherein each of A₁ and A₂ is independently H or OP^{''};

and

(D) producing the halichondrin macrolide or the salt thereof from the compound of formula (VIID) or the salt thereof.

10. The method of claim 9, wherein for (1),

(a) said producing the compound of formula (IB) or the salt or the tautomer thereof comprises reacting the compound of formula (IA) or the salt or the tautomer thereof with an organic base and a Lewis acid;

(b) said producing the halichondrin macrolide or the salt thereof comprises reacting the compound of formula (IB) or the salt or the tautomer thereof with a hydroxyl protecting group removing agent; and/or

(c) each R is optionally substituted alkyl.

11. The method of claim 9, wherein for (2)

(a) said producing the compound of formula (IB) or the salt or the tautomer thereof comprises reacting the compound of formula (IIA) or the salt or the tautomer thereof with an olefin metathesis catalyst; and/or

(b) said producing the halichondrin macrolide or the salt thereof comprises reacting the compound of formula (IB) or the salt or the tautomer thereof with a hydroxyl protecting group removing agent.

12. The method of claim 9, wherein for (3)

(a) said producing the compound of formula (IIIB) or the salt or the tautomer thereof comprises reacting the compound of formula (IIIA) or the salt or the tautomer thereof with an olefin metathesis catalyst;

(b) said producing the halichondrin macrolide or the salt thereof comprises reacting the compound of formula (IIIB) or the salt or the tautomer thereof with a hydroxyl protecting group removing agent;

(c) both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and/or

(d) P₅ is a hydroxyl protecting group.


13. The method of claim 9, wherein for (4)

(a) said producing the compound of formula (IVB) or the salt thereof comprises reacting the compound of formula (IVA) or the salt thereof with an olefin metathesis catalyst;

(b) said producing the halichondrin macrolide or the salt thereof from the compound of formula (IVB) or the salt thereof comprises reacting the compound of formula (IVB) or the salt thereof with a Brønsted acid; and/or

(c) at least one of P₄ and P₅ is H.

14. The method of claim 9, wherein said producing the halichondrin macrolide or the salt thereof comprises producing the compound of formula (IVC) or the salt thereof, the compound of formula (IVC) having the following structure:




wherein each of A_1 and A_2 is H or OP'' , and wherein P'' is H or a hydroxyl protecting group; and producing the halichondrin macrolide or the salt thereof from the compound of formula (IVC).

15. The method of claim 14, wherein

- (a) said producing the compound of formula (IVC) or the salt thereof comprises reacting the compound of formula (IVB) or the salt thereof with a hydroxyl protecting group removing agent; and/or
- (b) said producing the halichondrin macrolide from the compound of formula (IVC) or the salt thereof comprises reacting the compound of formula (IVC) or the salt thereof with a Brønsted acid.

16. The method of claim 9, wherein for (5),

- (a) said producing the compound of formula (VB) or the salt or the tautomer thereof comprises contacting the compound of formula (VA) or the salt or the tautomer thereof with an olefin metathesis catalyst;
- (b) P_4 is H, R_9 is H, and said producing the compound of formula (VC) or the salt thereof comprises oxidizing the compound of formula (VB) or the salt or the tautomer thereof to produce a compound of formula (VBa):



(VBa)

or a salt thereof;

and reacting the compound of formula (VBa) or the salt thereof with a 1,4-reducing agent to produce the compound of formula (VC) or the salt thereof;

(c) said producing the compound of formula (IVC) or the salt thereof comprises contacting the compound of formula (VC) or the salt thereof with a hydroxyl protecting group removing agent; and/or

(d) said producing the halichondrin macrolide or the salt thereof comprises contacting the compound of formula (IVC) or the salt thereof with a Brønsted acid.

17. The method of claim 9, wherein for (6),

(a) said producing the compound of formula (VIB) or the salt or the tautomer thereof comprises reacting the compound of formula (VIA) with a Cr(II) salt and a Ni(II) salt;

(b) said producing the halichondrin macrolide or the salt thereof comprises the step of nucleophilic ring-closing of the compound of formula (VIB) or the salt or the tautomer thereof;

(c) wherein both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and/or

(d) R₃ and R₄ combine to form a bond, and R₅ is H.

18. The method of claim 9, wherein for (7),

(a) said producing the compound of formula (VIIB) or the salt thereof comprises reacting the compound of formula (VIIA) or the salt thereof with a Cr(II) salt and a Ni(II) salt;

(b) said producing the compound of formula (VIIC) or the salt or the tautomer thereof comprises the step of nucleophilic ring-closing of the compound of formula (VIIB) or the salt thereof;

(c) said producing the compound of formula (VIID) or the salt thereof comprises reacting the compound of formula (VIIC) or the salt or the tautomer thereof with a hydroxyl protecting group removing agent;

(d) said producing the halichondrin macrolide comprises reacting the compound of formula (VIID) or the salt thereof with a Brønsted acid;

- (e) Z is iodide;
- (f) Y is trifluoromethanesulfonate;
- (g) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;
- (h) each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms $-(CH(OR_8))-$, wherein R₈ is H or a hydroxyl protecting group;
- (i) P₅ is H; and/or
- (j) Z is chloride, bromide, or iodide, and R₆ and R₇ combine to form a bond.

19. The method of claim 1, wherein


- (a) a designates (S)-stereogenic center;
- (b) one and only one of D and D' is optionally substituted alkyl or OP₁;
- (c) A is a group of formula (1);
- (d) L is $-(CH(OP_2))-$;
- (e) R₁ and P₁ combine to form a bond;
- (f) G is O;
- (g) E is optionally substituted alkyl;
- (h) k is 1;
- (i) R₂ is $-(CH_2)_nOP_3$;
- (j) at least one of P₂ and P₃ is a hydroxyl protecting group; and/or
- (k) A₁ is H.

20. The method of claim 19, wherein one and only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group.

21. A method of preparing an intermediate in the synthesis of a halichondrin macrolide, said method comprising performing an allene-Prins reaction by contacting a compound of formula (VIII A) or a salt thereof with a compound of formula (VIII B) and R₄OH,

wherein R₄ is an optionally substituted acyl;


wherein the compound of formula (VIII A) has the following structure:



wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆

saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl,

alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;


k is 0 or 1;

n is 0, 1, or 2;


X is O, or X, together with the carbon atom to which it is attached, forms $-(C(OP_z)_2)-$, wherein each P_z is independently optionally substituted alkyl or optionally substituted aryl, or both P_z combine to form optionally substituted alkylene or optionally substituted arylene; and

P₄ is H or a hydroxyl protecting group;


wherein the compound of formula (VIIB) has the following structure:



wherein




wherein P₅ is H or a hydroxyl protecting group; each P₆ is independently a hydroxyl protecting group, or both P₆ groups, together with the atoms to which each is attached, combine to form a cyclic protected diol; and R₅ is H or -CH₂X₁CH₂CH=CH₂, wherein X₁ is O, -CH₂-, or NP₇, wherein P₇ is a sulfonyl; and wherein said intermediate is a compound of formula (VIIC):



or a salt thereof.

22. The method of claim 21, wherein said performing a Prins reaction comprises reacting the compound of formula (VIIIa) or a salt thereof with a Lewis acid; and/or k is 1.

23. A compound of
(1) formula (IA) or formula (IB),




or a salt or a tautomer thereof,

wherein

each R is independently optionally substituted alkyl or optionally substituted aryl;

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or

A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;


or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP'';

each P₄ is independently H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_V))-, wherein P_V is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal; and

P₅ is H or a hydroxyl protecting group;-


(2) formula (IIA), formula (IIIA), or formula (IIIB):



or a salt or a tautomer thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each

is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:


(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_V))-$, wherein P_V is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal; and P_5 is H or a hydroxyl protecting group;

(3) formula (IVA), formula (IVB), or formula (IVC):




or a salt thereof,

wherein

each of A_1 and A_2 is H or OP'' , where P'' is H or a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



, wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(i) R₃ is H or a hydroxyl protecting group, R₄ is alkyl ether, and R₅ is H;

(ii) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;


or

(iii) R₃ and R₄ combine to form a bond, and R₅ is H;

each P_4 and P_5 is independently H or a hydroxyl protecting group; and

R_6 is H, and P_6 is H or a hydroxyl protecting group; or R_6 and P_6 combine to form a double bond;


(4) formula (VA), formula (VB), or formula (VC):



or a salt or a tautomer thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



A_2 is H or OP'' ;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(a1) R_3 is H or a hydroxyl protecting group, R_4 is alkyl ether, and R_5 is H;

(a2) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

(a3) R_3 and R_4 combine to form a bond, and R_5 is H;

(b1) A_1 and R_7 combine to form oxo, R_6 is H or a hydroxyl protecting group, and R_8 is H;

or

(b2) A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:

(i) R_6 is H or a hydroxyl protecting group, and R_7 and R_8 combine to form a double bond;

or

(ii) R_6 and R_7 combine to form a bond, and R_8 is H or OP'' ;


(c1) R_9 is H, and P_4 is H or a hydroxyl protecting group;

or

(c2) R_9 and P_4 combine to form a double bond;

R_{10} is H or $-CH_2X_1CH_2CH=CH_2$, wherein X_1 is O, $-(R_{11})_2-$, or NP_6 , and wherein each R_{11} is independently H or $-COOR_{12}$, P_6 is an *N*-protecting group, and R_{12} is alkyl;
each P_5 is independently H or a hydroxyl protecting group;


(5) formula (VIA):



or a salt or a tautomer thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

Y is iodide, bromide, or trifluoromethanesulfonate;

b designates (*R*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide; or b designates (*S*)-stereogenic center, and Z is OR_6 , wherein R_6 is a hydroxyl protecting group;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:


- (i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

- (ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_Y))-$, wherein P_Y is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal;

(6) formula (VIB):



(VIB)

or a salt or a tautomer thereof,


wherein

b designates (*R*)-stereogenic center, c designates (*S*)-stereogenic center, and Z is a sulfonate, chloride, bromide, or iodide;

or

b designates (*S*)-stereogenic center, c designates (*R*)-stereogenic center, and Z is OR_6 , wherein R_6 is a hydroxyl protecting group;

each of D and D' is independently H, optionally substituted alkyl, or OP_1 , provided that only one of D and D' is OP_1 , wherein P_1 is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C_{1-6} saturated or C_{2-6} unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q_1 , the group of formula (1) having the structure:




(1),

wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R_1 is H, or R_1 and P_1 combine to form a bond;

R_2 is H or $-(CH_2)_nOP_3$, and each of P_2 and P_3 is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P_2 and P_3 , together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R_2 and P_2 combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH_2 , or NR_N , wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

k is 0 or 1;

n is 0, 1, or 2;

A_1 and R_4 combine to form oxo, R_3 is H or a hydroxyl protecting group, and R_5 is H;

or

A_1 is H or OP'' , where P'' is H or a hydroxyl protecting group, and:


(i) R_3 is H or a hydroxyl protecting group, and R_4 and R_5 combine to form a double bond;

or

(ii) R_3 and R_4 combine to form a bond, and R_5 is H or OP'' ;

each P_4 is independently H or a hydroxyl protecting group, and X_1 is oxo or X_1 , together with the carbon atom to which it is attached, is $-(CH(OP_V))-$, wherein P_V is H or a hydroxyl protecting group; or both P_4 groups and X_1 , together with the atoms to which each is attached, combine to form ketal;


(7) formula (VIIA) or formula (VIIB):



or a salt thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:





wherein

L is $\text{---(CH(OP}_2\text{))---}$, $\text{---(C(OH)(OP}_2\text{))---}$, or ---C(O)--- ;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $\text{---(CH}_2\text{)}_n\text{OP}_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:





, wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

(a1) R₃ is H or a hydroxyl protecting group, R₄ and R₅ combine to form a double bond, each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or -(CH(OR₈))-;

or

(a2) R₃ and R₄ combine to form a bond, R₅ is H, and

each P₄ is independently H or a hydroxyl protecting group, and X₁, together with the carbon to which it is attached, forms a carbonyl or -(CH(OR₈))-;

or

both P₄ groups and X₁, together with the atoms to which each is attached, combine to form ketal;

(b1) Z is chloride, bromide, or iodide, and R₆ and R₇ combine to form a bond;

(b2) Z and R₇ combine to form a double bond, and R₆ is a hydroxyl protecting group;


or

(b3) when Z and R₇ are absent, R₆ is H or a hydroxyl protecting group;

P₅ is H or a hydroxyl protecting group;

wherein R₈ is H or a hydroxyl protecting group; or


(8) formula (VIIC):



or a salt or a tautomer thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:





wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:





, wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;

G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q₁ is independently OR_A, SR_A, SO₂R_A, OSO₂R_A, NR_BR_A, NR_B(CO)R_A, NR_B(CO)(CO)R_A, NR_A(CO)NR_BR_A, NR_B(CO)OR_A, (CO)OR_A, O(CO)R_A, (CO)NR_BR_A, or O(CO)NR_BR_A, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

each P₄ is H or a hydroxyl protecting group, and X₁ is oxo or X₁, together with the carbon atom to which it is attached, is -(CH(OP_Y))-, wherein P_Y is H or a hydroxyl protecting group; or both P₄ groups and X₁, together with the atoms to which each is attached, combine to form a ketal; and

A₁ and R₄ combine to form oxo, R₃ is H or a hydroxyl protecting group, and R₅ is H;

or


A₁ is H or OP'', where P'' is H or a hydroxyl protecting group, and:

(i) R₃ is H or a hydroxyl protecting group, and R₄ and R₅ combine to form a double bond;

or

(ii) R₃ and R₄ combine to form a bond, and R₅ is H or OP''.


24. A compound of formula (VIIC):



or a salt thereof,

wherein

each of D and D' is independently H, optionally substituted alkyl, or OP₁, provided that only one of D and D' is OP₁, wherein P₁ is H, alkyl, or a hydroxyl protecting group; A is a group of formula (1) or a C₁₋₆ saturated or C₂₋₆ unsaturated hydrocarbon skeleton, the skeleton being unsubstituted or having from 1 to 10 substituents independently selected from the group consisting of cyano, halo, azido, oxo, and Q₁, the group of formula (1) having the structure:




wherein

L is $-(CH(OP_2))-$, $-(C(OH)(OP_2))-$, or $-C(O)-$;

R₁ is H, or R₁ and P₁ combine to form a bond;

R₂ is H or $-(CH_2)_nOP_3$, and each of P₂ and P₃ is independently H, optionally substituted alkyl, or a hydroxyl protecting group, or P₂ and P₃, together with the atoms to which each is attached, combine to form a ketal, a cyclic carbonate, a dicarbonyl-dioxo, or silylene-dioxo; or R₂ and P₂ combine to form an optionally substituted ethylene or a structure selected from the group consisting of:



wherein each P' is independently H or a hydroxyl protecting group;

E is optionally substituted alkyl or optionally substituted alkoxy;


G is O, S, CH₂, or NR_N, wherein R_N is H, an N-protecting group, or optionally substituted alkyl;

each Q_1 is independently OR_A , SR_A , SO_2R_A , OSO_2R_A , NR_BR_A , $NR_B(CO)R_A$, $NR_B(CO)(CO)R_A$, $NR_A(CO)NR_BR_A$, $NR_B(CO)OR_A$, $(CO)OR_A$, $O(CO)R_A$, $(CO)NR_BR_A$, or $O(CO)NR_BR_A$, wherein each of R_A and R_B is independently H, alkyl, haloalkyl, hydroxyalkyl, aminoalkyl, aryl, haloaryl, hydroxyaryl, alkoxyaryl, arylalkyl, alkylaryl, haloarylalkyl, alkylhaloaryl, (alkoxyaryl)alkyl, heterocyclic radical, or heterocyclic radical-alkyl;

k is 0 or 1;

n is 0, 1, or 2;

P_4 is H or a hydroxyl protecting group; and



wherein P_5 is H or a hydroxyl protecting group; each P_6 is independently a hydroxyl protecting group, or both P_6 groups, together with the atoms to which each is attached, combine to form a cyclic protected diol; and R_5 is H or $-\text{CH}_2\text{X}_1\text{CH}_2\text{CH}=\text{CH}_2$, wherein X_1 is O, $-\text{CH}_2-$, or NP_7 , wherein P_7 is a sulfonyl.