

Quinoline derivatives as PDE10A enzyme inhibitors**Field of the Invention**

The present invention provides quinoline derivatives that are PDE10A enzyme inhibitors, and as such are useful to treat neurodegenerative and psychiatric disorders. Especially, the invention provides compounds that are highly selective for PDE10 over other PDE subtypes. The present invention also provides pharmaceutical compositions comprising compounds of the invention and methods of treating disorders using the compounds of the invention.

Background of the Invention

The cyclic nucleotides cyclic-adenosine monophosphate (cAMP) and cyclic-guanosine monophosphate (cGMP) function as intracellular second messengers regulating a vast array of processes in neurons. Intracellular cAMP and cGMP are generated by adenylyl and guanylyl cyclases, and are degraded by cyclic nucleotide phosphodiesterases (PDEs) via hydrolysis of the cyclic nucleotides into their respective nucleotide monophosphates.

Phosphodiesterase 10A (PDE10A) is a dual-specificity phosphodiesterase that can convert both cAMP to AMP and cGMP to GMP (Soderling, S. *et al. Proc. Natl. Acad. Sci.* **1999**, 96, 7071-7076). PDE10A is primarily expressed in the neurons in the striatum, n. accumbens and in the olfactory tubercle (Kotera, J. *et al. Biochem. Biophys. Res. Comm.* **1999**, 261, 551-557 and Seeger, T.F. *et al. Brain Research*, **2003**, 985, 113-126).

Studies indicate that within the brain, PDE10 expression is expressed at high levels by the medium spiny neurons (MSN) of the caudate nucleus, the accumbens nucleus and the corresponding neurons of the olfactory tubercle. MSN express two functional classes of neurons: the D₁ class expressing D₁ dopamine receptors and the D₂ class expressing D₂ dopamine receptors. The D₁ class of neurons is part of the 'direct' striatal output pathway, which broadly functions to facilitate behavioral responses. The D₂ class of neurons is part of the 'indirect' striatal output pathway,

which functions to suppress behavioral responses that compete with those being facilitated by the 'direct' pathway.

Dopamine D₂ receptor antagonism is well established in the treatment of schizophrenia. Since the 1950's, dopamine D₂ receptor antagonism has been the mainstay in psychosis treatment and all effective antipsychotic drugs antagonise D₂ receptors. The effects of D₂ are likely to be mediated primarily through neurons in the striatum, nucleus accumbens and olfactory tubercle, since these areas receive the densest dopaminergic projections and have the strongest expression of D₂ receptors (Konradi, C. and Heckers, S. *Society of Biological Psychiatry*, **2001**, 50, 729-742).

Because PDE10A, in this context, has the desired expression profile with high and relatively specific expression in neurons in striatum, nucleus accumbens and olfactory tubercle, PDE10A inhibition is likely to have effects similar to D₂ receptor antagonism and therefore have antipsychotic effects.

While PDE10A inhibition is expected to mimic D₂ receptor antagonism in part, it might be expected to have a different profile. The D₂ receptor has signaling components besides cAMP (Neve, K. A. *et al. Journal of Receptors and Signal Transduction* **2004**, 24, 165-205), wherefore interference with cAMP through PDE10A inhibition may reduce the risk of the extrapyramidal side effects that are seen with strong D₂ antagonism. Conversely, PDE10A inhibition may have some effects not seen with D₂ receptor antagonism. PDE10A is also expressed in D₁ receptors expressing striatal neurons (Seeger, T. F. *et al. Brain Research*, **2003**, 985, 113-126).

Further, since D₁ receptor agonism leads to stimulation of adenylate cyclase and resulting increase in cAMP levels, PDE10A inhibition is likely to also have effects that mimic D₁ receptor agonism.

Finally, PDE10A inhibition will not only increase cAMP in cells, but might also be expected to increase cGMP levels, since PDE10A is a dual specificity

phosphodiesterase. cGMP activates a number of target protein in cells like cAMP and also interacts with the cAMP signaling pathways.

In conclusion, PDE10A inhibition is likely to mimic D₂ receptor antagonism in part and therefore has antipsychotic effect, but the profile might differ from that observed with classical D₂ receptor antagonists.

The PDE10A inhibitor papaverine is shown to be active in several antipsychotic models. Papaverine potentiated the cataleptic effect of the D₂ receptor antagonist haloperidol in rats, but did not cause catalepsy on its own (WO 03/093499). Papaverine reduced hyperactivity in rats induced by PCP, while reduction of amphetamine induced hyperactivity was insignificant (WO 03/093499). These models suggest that PDE10A inhibition has the classic antipsychotic potential that would be expected from the theoretical considerations outlined above. WO 03/093499 further discloses the use of selective PDE10 inhibitors for the treatment of associated neurologic and psychiatric disorders. Furthermore, PDE10A inhibition reverses subchronic PCP-induced deficits in attentional set-shifting in rats (Rodefer *et al. Eur. J. Neurosci.* **2005**, 4, 1070-1076). This model suggests that PDE10A inhibition might alleviate cognitive deficits associated with schizophrenia.

The tissue distribution of PDE10A indicates that PDE10A inhibitors can be used to raise levels of cAMP and/or cGMP within cells that express the PDE10A enzyme, especially neurons that comprise the basal ganglia, and the PDE10A inhibitors of the present invention would therefore be useful in treating a variety of associated neuropsychiatric conditions involving the basal ganglia such as neurological and psychiatric disorders, schizophrenia, bipolar disorder, psychosis, obsessive compulsive disorder and addiction, and may have the benefit of not possessing unwanted side effects, which are associated with the current therapies on the market.

Furthermore, recent publications (WO 2005/120514, WO 2005012485, Cantin *et al*, Bioorganic & Medicinal Chemistry Letters 17 (2007) 2869–2873) suggest that

PDE10A inhibitors may be useful for treatment of obesity and non-insulin dependent diabetes.

Furthermore, recent publications suggest that PDE10A inhibitors may be useful for the treatment of Huntingtons Disease (Giampa et al. PLoS One 2010, 5(10), Giampa et al. Neurobiology of Disease (2009), 34(3), 450-456, Hebb et al. Current Opinion in Pharmacology 2007, 7(1), 86-92.)

Pyrrolodihydroisoquinolines and variants thereof are disclosed as inhibitors of PDE10 in WO 05/03129 and WO 05/02579. Piperidinyl-substituted quinazolines and isoquinolines that serve as PDE10 inhibitors are disclosed in WO 05/82883. WO 06/11040 discloses substituted quinazoline and isoquinoline compounds that serve as inhibitors of PDE10. US 20050182079 discloses substituted tetrahydro-isoquinoliny derivatives of quinazoline and isoquinoline that serve as effective phosphodiesterase (PDE) inhibitors. In particular, US 20050182079 relates to said compounds, which are selective inhibitors of PDE10. Analogously, US 20060019975 discloses piperidine derivatives of quinazoline and isoquinoline that serve as effective phosphodiesterase (PDE) inhibitors. US 20060019975 also relates to compounds that are selective inhibitors of PDE10. WO 06/028957 discloses cinnoline derivatives as inhibitors of PDE10 for the treatment of psychiatric and neurological syndromes. WO09/152825 discloses phenylimidazole derivatives as compounds that serve as inhibitors of PDE10.

However, these disclosures do not pertain to the compounds of the invention, which are structurally unrelated to any of the known PDE10 inhibitors (Kehler, J. *et al. Expert Opin. Ther. Patents* **2007**, 17, 147-158), and which have now been found by the inventors to be highly active and selective PDE10A enzyme inhibitors.

The present invention provides compounds that are PDE10A enzyme inhibitors and thus useful for treatment for neurodegenerative and/or psychiatric disorders, which are not efficacious in all patients. Hence, there remains a need for alternative methods of treatment.

Summary of the Invention

The objective of the present invention is to provide compounds that are selective PDE10A enzyme inhibitors.

- 5 A further objective of the present invention is to provide compounds which have such activity, and which have improved solubility, metabolic stability and/or bioavailability compared to prior art compounds.

- 10 Another objective of the invention is to provide an effective treatment, in particular long-term treatment, of a human patient, without causing the side effects typically associated with current therapies for neurological and psychiatric disorders.


Further objectives of the invention will become apparent upon reading the present specification.

15

Detailed description of the invention

Embodiments of the invention


- 20 In a first embodiment (E1) the present invention relates to compounds of formula I:



- 25 wherein R1, R2, R3, R4, R5 and R6 are individually selected from the group consisting of hydrogen, hydroxyl, cyano, C₁-C₆ alkyl; C₁-C₆ alkoxy, halogen, methylenedioxy, difluoromethylenedioxy and ethylenedioxy

wherein -L- is a linker selected from -CH₂-CH₂- and -CH=CH-

wherein HET is selected from the group consisting of



5

wherein one or more of the carbon-bound hydrogen in the HET optionally may be substituted with up to three substituents R7, R8 and R9 individually selected from C₁-C₆ alkyl; halogen; cyano, halo(C₁-C₆)alkyl; aryl, alkoxy and C₁-C₆ hydroxyalkyl

10 and wherein * denotes the attachment point,

and tautomers and pharmaceutically acceptable salts thereof, and polymorphic forms thereof.

15 In an embodiment (E2) of embodiment (E1) one or more of R1-R6 is selected from the group consisting of C₁-C₃ alkyl substituted with one or more F and unsubstituted C₁-C₃ alkyl.

In an embodiment (E3) of embodiment (E1) or (E2) one or more of R1-R6 is
20 selected from the group consisting of methyl, ethyl, propyl, isopropyl, monofluoromethyl, difluoromethyl and trifluoromethyl.

In an embodiment (E4) of embodiment (E1) one or more of R1-R6 is selected from the group consisting of methoxy, difluoromethoxy and trifluoromethoxy.

25


In an embodiment (E5) of embodiment (E1) one or more of R1-R6 is selected from the group consisting of fluorine and chlorine.

In an embodiment (E6) of any of embodiments (E1) to (E5) -L- is -CH₂-CH₂-

In an embodiment (E7) of any of embodiments (E1) to (E5) -L- is -CH=CH-

5

In an embodiment (E8) of any of embodiments (E1) to (E7) HET is selected from the group consisting of



10

wherein HET optionally is substituted with one or more of R⁷-R⁹, and wherein * denotes the attachment point.

15 In an embodiment (E9) of any of embodiments (E1) to (E8) HET is substituted with one substituent R⁷ selected from the group consisting of C₁-C₆ alkyl, such as methyl; halogen, such as chlorine or bromine; cyano; halo(C₁-C₆)alkyl, such as trifluoromethyl; aryl such as phenyl; and C₁-C₆ hydroxyalkyl such as CH₂CH₂OH.

20 In an embodiment (E10) of any of embodiments (E1) to (E8), HET is substituted with two substituents R⁷ and R⁸ individually selected from C₁-C₆ alkyl, such as methyl; halogen, such as chlorine or bromine; cyano; halo(C₁-C₆)alkyl, such as trifluoromethyl; aryl, such as phenyl; and C₁-C₆ hydroxyalkyl, such as CH₂CH₂OH.

25 In an embodiment (E11) of any of embodiment (E1) to (E8), HET is substituted with three substituents R⁷, R⁸ and R⁹ individually selected from C₁-C₆ alkyl, such as methyl; halogen, such as chlorine or bromine; cyano; halo(C₁-C₆)alkyl, such as trifluoromethyl; aryl, such as phenyl; and C₁-C₆ hydroxyalkyl, such as CH₂CH₂OH.

In an embodiment (E12) of any of embodiments (E1) to (E8) HET is unsubstituted.

30

In an embodiment (E13) of any of embodiments (E1), (E9), (E10) and (E11) HET is substituted with at least one C₁-C₆ alkyl, such as methyl

In an embodiment (E14) of any of embodiments (E1) to (E11) HET is selected from the group consisting of (5,7-Dimethyl-imidazo[1,2-a]pyrimidin-2-yl), 5,7-Dimethyl-
5 [1,2,4]triazolo[1,5-a]pyrimidin-2-yl), (5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl), (8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl) and 5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl).

10 In a specific embodiment (E15) of embodiment (E1) the compound is selected from the group of compounds listed in Table 1.

In an embodiment (E16) of any of the embodiments (E1) to (E15) the invention provides a compound of formula I, or a pharmaceutically acceptable salt thereof, for
15 use as a medicament.

In an embodiment (E17) of any of the embodiments (E1) to (E15) the invention provides a pharmaceutical composition comprising a therapeutically effective amount of a compound of formula I and a pharmaceutically acceptable carrier, diluent or excipient.
20

In an embodiment (E18) of any of the embodiments (E1) to (E15) the present invention provides the use of a compound of formula I, or a pharmaceutically acceptable salt thereof, for the preparation of a medicament for the treatment of a neurodegenerative or psychiatric disorder.
25


Furthermore, in an embodiment (E19) of any of the embodiments (E1) to (E15) the present invention provides a method of treating a subject suffering from a neurodegenerative disorder, comprising administering to the subject a therapeutically effective amount of a compound of formula I.
30

In an embodiment (E20) of any of the embodiments (E1) to (E15) the present invention provides a method of treating a subject suffering from a psychiatric

disorder, comprising administering to the subject a therapeutically effective amount of a compound of formula I.

In an embodiment (E21) of any of the embodiments (E1) to (E15) the present invention provides a method of treating a subject suffering from a drug addiction, such as an alcohol, amphetamine, cocaine, or opiate addiction.

In an embodiment (E22) the present invention relates to compounds of formula I:



wherein R1-R6 and HET are as described in any of the previous embodiments (E1) to (E14) and -L- is a linker selected from -S-CH₂-, -CH₂-S- and -C≡C-.

15 Definition of substituents

As used in the context of the present invention, the terms “halo” and “halogen” are used interchangeably and refer to fluorine, chlorine, bromine or iodine.

The term “C₁-C₆ alkyl” refers to a straight-chain or branched saturated hydrocarbon having from one to six carbon atoms, inclusive. Examples of such groups include, but are not limited to, methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl, 2-methyl-1-butyl, and n-hexyl. The expression “C₁-C₆ hydroxyalkyl” refers to a C₁-C₆ alkyl group as defined above which is substituted with one hydroxy group. The term “halo(C₁-C₆)alkyl” refers to a C₁-C₆ alkyl group as defined above which is substituted with up to three halogen atoms, such as trifluoromethyl.

The expression “C₁-C₆ alkoxy” refers to a straight-chain or branched saturated alkoxy group having from one to six carbon atoms, inclusive, with the open valency on the oxygen. Examples of such groups include, but are not limited to, methoxy,

ethoxy, n-butoxy, 2-methyl-pentoxo and n-hexyloxy. The alkoxy may optionally be substituted with up to three halogen atoms, such as trifluoromethoxy.

The term "C₃-C₈ cycloalkyl" typically refers to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl. The expression "C₁-C₆ alkyl(C₃-C₈)cycloalkyl" refers to a C₃-C₈ cycloalkyl as defined above which is substituted with a straight-chain or branched C₁-C₆ alkyl. Examples of such groups include, but are not limited to, cyclopropylmethyl.

The term "heterocycloalkyl" refers to a four to eight membered ring containing carbon atoms and up to three N, O or S atoms, provided that the four to eight membered ring does not contain adjacent O or adjacent S atoms. The open valency is on either the heteroatom or carbon atom. Examples of such groups include, but are not limited to, azetidiny, oxetanyl, piperaziny, morpholinyl, thiomorpholinyl and [1,4]diazepanyl. The term "hydroxyheterocycloalkyl" refers to a heterocycloalkyl as defined above which is substituted with one hydroxy group. The term "C₁-C₆ alkyl-heterocycloalkyl" refers to a heterocycloalkyl as defined above which is substituted with a C₁-C₆ alkyl group. Examples of such groups include, but are not limited to, tetrahydropyran-4-yl-methyl and 2-morpholin-4-yl-ethyl.

The term "aryl" refers to a phenyl ring, optionally substituted with halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy or halo(C₁-C₆)alkyl as defined above. Examples of such groups include, but are not limited to, phenyl and 4-chlorophenyl.

The term "C₁-C₆arylalkyl" refers to an aryl as defined above which is substituted with a straight-chain or branched C₁-C₆ alkyl. Examples of such groups include, but are not limited to, benzyl and 4-chlorobenzyl.

In a further embodiment one or more of the hydrogen atoms of the compound of formula I have been substituted by deuterium.

In the context of this application it should be understood that the meaning of "R₁-R₆", "R₁ to R₆" and "R₁, R₂, R₃, R₄, R₅ and R₆" is the same.

Additionally, the present invention further provides certain embodiments of the invention, that are described below.

In separate embodiments of the invention, the compound of formula I is selected among the following specific compounds listed in Table 1, in the form of the free base, one or more tautomers thereof or a pharmaceutically acceptable salt thereof. Table 1 lists compounds of the invention and the corresponding IC₅₀ values determined as described in the section "PDE10A inhibition assay". Each of the compounds constitutes an individual embodiment, of the present invention.

It should be understood that the various aspects, embodiments, implementations and features of the invention mentioned herein may be claimed separately, or in any combination, as illustrated by the following non-limiting examples.

Table 1: Compounds of the invention and IC₅₀ values

Compound	IC ₅₀ (nM)
2-[(E)-2-(5,7-Dimethyl-imidazo[1,2-a]pyrimidin-2-yl)-vinyl]-quinoline	320
2-[(Z)-2-(5,7-Dimethyl-imidazo[1,2-a]pyrimidin-2-yl)-vinyl]-quinoline	240
2-[2-(5,7-Dimethyl-imidazo[1,2-a]pyrimidin-2-yl)-ethyl]-quinoline	310
2-[2-(5,7-Dimethyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl)-ethyl]-quinoline	72
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-quinoline	19
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-quinoline	7,4
2-[2-(5,7-Dimethyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl)-ethyl]-6-methoxy-quinoline	170
2-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline	12
2-[2-(8-Ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-6-fluoro-quinoline	25
6-Fluoro-2-[2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline	34
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline	24
2-[2-(6-Fluoro-quinolin-2-yl)-ethyl]-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-8-ol	85

Compound	IC50 (nM)
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-4-methyl-quinoline	18
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-4-methoxy-quinoline	12
4-Methoxy-2-[2-(5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline	190
4-Methoxy-2-[2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline	35
4-Methyl-2-[2-(5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline	140
2-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-4-methyl-quinoline	42
4-Chloro-8-fluoro-2-[(E)-2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-vinyl]-quinoline	270
8-Fluoro-2-[2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline	150
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-7-fluoro-quinoline	15
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-fluoro-quinoline	24
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-fluoro-quinoline	26
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-4-fluoro-quinoline	28
2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-fluoro-quinoline	190
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-7-fluoro-4-methoxy-quinoline	26
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-7-fluoro-quinolin-4-ol	500
2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-7-trifluoromethyl-quinoline	550
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-fluoro-4-methoxy-quinoline	14
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-7-trifluoromethyl-quinoline	240
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-fluoro-quinolin-4-ol	410
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-5-fluoro-quinoline	120
7-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-quinoline	52
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-c]pyrimidin-2-yl)-ethyl]-6-isopropyl-quinoline	61
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-5,7-	110

Compound	IC ₅₀ (nM)
difluoro-quinoline	
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-5,6,8-trifluoro-quinoline	360
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6,8-difluoro-quinoline	300
6-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-[1,3]dioxolo[4,5-g]quinoline	13
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-fluoro-8-methyl-quinoline	2000
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-fluoro-7-methyl-quinoline	8,8
6-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-2,2-difluoro-[1,3]dioxolo[4,5-g]quinoline	520
7-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-quinoline-6-carbonitrile	59
7-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-2,3-dihydro-[1,4]dioxino[2,3-g]quinoline	4
6-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-quinoline	42
6-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-8-fluoro-quinoline	240
8-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-methyl-quinoline	290
5,7-Dichloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-quinoline	39
2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-trifluoromethoxy-quinoline	2800
2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-trifluoromethyl-quinoline	490
2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-cyano-quinoline	2100
2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-7-methoxy-quinoline	13
2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-5-methoxy-quinoline	19

In a particular embodiment of the present invention the compounds of the present invention have an IC₅₀ value of less than 50 nM, such as in the range of 0.2 – 20 nM, particularly in the range of 0.2 – 10 nM, such as in the range of 0.2 – 5 nM.

Pharmaceutically Acceptable Salts

The present invention also comprises salts of the compounds, typically, pharmaceutically acceptable salts. Such salts include pharmaceutically acceptable acid addition salts. Acid addition salts include salts of inorganic acids as well as organic acids.

Representative examples of suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, phosphoric, sulfuric, sulfamic, nitric acids and the like. Representative examples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, itaconic, lactic, methanesulfonic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, methane sulfonic, ethanesulfonic, tartaric, ascorbic, pamoic, bismethylene salicylic, ethanedisulfonic, gluconic, citraconic, aspartic, stearic, palmitic, EDTA, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, p-toluenesulfonic acids, theophylline acetic acids, as well as the 8-halotheophyllines, for example 8-bromotheophylline and the like. Further examples of pharmaceutically acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in Berge, S.M. et al., *J. Pharm. Sci.* 1977, 66, 2, the contents of which are hereby incorporated by reference.

Furthermore, the compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of this invention.

Pharmaceutical compositions

The present invention further provides a pharmaceutical composition comprising a therapeutically effective amount of a compound of formula I and a pharmaceutically acceptable carrier or diluent. The present invention also provides a pharmaceutical composition comprising a therapeutically effective amount of one of the specific compounds disclosed in the Experimental Section herein and a pharmaceutically acceptable carrier or diluent.

The compounds of the invention may be administered alone or in combination with pharmaceutically acceptable carriers, diluents or excipients, in either single or multiple doses. The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), transdermal, intracisternal, intraperitoneal, vaginal and parenteral (including subcutaneous, intramuscular, intrathecal, intravenous and intradermal) routes. It will be appreciated that the route will depend on the general condition and age of the subject to be treated, the nature of the condition to be treated and the active ingredient.

Pharmaceutical compositions for oral administration include solid dosage forms such as capsules, tablets, dragees, pills, lozenges, powders and granules. Where appropriate, the compositions may be prepared with coatings such as enteric coatings or they may be formulated so as to provide controlled release of the active ingredient such as sustained or prolonged release according to methods well known in the art. Liquid dosage forms for oral administration include solutions, emulsions, suspensions, syrups and elixirs.

Pharmaceutical compositions for parenteral administration include sterile aqueous and nonaqueous injectable solutions, dispersions, suspensions or emulsions as well as sterile powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Other suitable administration forms include, but are not limited to, suppositories, sprays, ointments, creams, gels, inhalants, dermal patches and implants.

Typical oral dosages range from about 0.001 to about 100 mg/kg body weight per day. Typical oral dosages also range from about 0.01 to about 50 mg/kg body weight per day. Typical oral dosages further range from about 0.05 to about 10 mg/kg body weight per day. Oral dosages are usually administered in one or more dosages, typically, one to three dosages per day. The exact dosage will depend upon the frequency and mode of administration, the sex, age, weight and general condition of the subject treated, the nature and severity of the condition treated and any concomitant diseases to be treated and other factors evident to those skilled in the art.

The formulations may also be presented in a unit dosage form by methods known to those skilled in the art. For illustrative purposes, a typical unit dosage form for oral administration may contain from about 0.01 to about 1000 mg, from about 0.05 to about 500 mg, or from about 0.5 mg to about 200 mg.

For parenteral routes such as intravenous, intrathecal, intramuscular and similar administration, typical doses are in the order of half the dose employed for oral administration.

The present invention also provides a process for making a pharmaceutical composition comprising admixing a therapeutically effective amount of a compound of formula I and at least one pharmaceutically acceptable carrier or diluent. In an embodiment, of the present invention, the compound utilized in the aforementioned process is one of the specific compounds disclosed in the Experimental Section herein.

The compounds of this invention are generally utilized as the free substance or as a pharmaceutically acceptable salt thereof. One example is an acid addition salt of a compound having the utility of a free base. When a compound of formula I contains a free base such salts are prepared in a conventional manner by treating a solution or suspension of a free base of formula I with a molar equivalent of a pharmaceutically acceptable acid. Representative examples of suitable organic and inorganic acids are described above.

For parenteral administration, solutions of the compounds of formula I in sterile aqueous solution, aqueous propylene glycol, aqueous vitamin E or sesame or peanut oil may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The compounds of formula I may be readily incorporated into known sterile aqueous media using standard techniques known to those skilled in the art.

Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solutions and various organic solvents. Examples of solid carriers include lactose, terra alba, sucrose, cyclodextrin, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and lower alkyl ethers of cellulose. Examples of liquid carriers include, but are not limited to, syrup, peanut oil, olive oil, phospholipids, fatty acids, fatty acid amines, polyoxyethylene and water. Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax. The pharmaceutical compositions formed by combining the compounds of formula I and a pharmaceutically acceptable carrier are then readily administered in a variety of dosage forms suitable for the disclosed routes of administration. The formulations may conveniently be presented in unit dosage form by methods known in the art of pharmacy.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules or tablets, each containing a

predetermined amount of the active ingredient, and optionally a suitable excipient. Furthermore, the orally available formulations may be in the form of a powder or granules, a solution or suspension in an aqueous or non-aqueous liquid, or an oil-in-water or water-in-oil liquid emulsion.

5

If a solid carrier is used for oral administration, the preparation may be tableted, placed in a hard gelatin capsule in powder or pellet form or it may be in the form of a troche or lozenge. The amount of solid carrier will vary widely but will range from about 25 mg to about 1 g per dosage unit. If a liquid carrier is used, the preparation
10 may be in the form of a syrup, emulsion, soft gelatin capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

The pharmaceutical compositions of the invention may be prepared by conventional methods in the art. For example, tablets may be prepared by mixing the active
15 ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine prepare tablets. Examples of adjuvants or diluents comprise: corn starch, potato starch, talcum, magnesium stearate, gelatin, lactose, gums, and the like. Any other adjuvants or additives usually used for such purposes such as colorings, flavorings, preservatives etc. may
20 be used provided that they are compatible with the active ingredients.

Therapeutically effective amount

In the present context, the term "therapeutically effective amount" of a compound means an amount sufficient to cure, alleviate or partially arrest the clinical
25 manifestations of a given disease and its complications in a therapeutic intervention comprising the administration of said compound. An amount adequate to accomplish this is defined as "therapeutically effective amount". Effective amounts for each purpose will depend on the severity of the disease or injury as well as the weight and general state of the subject. It will be understood that determining an
30 appropriate dosage may be achieved using routine experimentation, by constructing a matrix of values and testing different points in the matrix, which is all within the ordinary skills of a trained physician.

In the present context, the term "treatment" and "treating" means the management and care of a patient for the purpose of combating a condition, such as a disease or a disorder. The term is intended to include the full spectrum of treatments for a given condition from which the patient is suffering, such as administration of the active compound to alleviate the symptoms or complications, to delay the progression of the disease, disorder or condition, to alleviate or relieve the symptoms and complications, and/or to cure or eliminate the disease, disorder or condition as well as to prevent the condition, wherein prevention is to be understood as the management and care of a patient for the purpose of combating the disease, condition, or disorder and includes the administration of the active compounds to prevent the onset of the symptoms or complications. Nonetheless, prophylactic (preventive) and therapeutic (curative) treatments are two separate aspects of the invention. The patient to be treated is preferably a mammal, in particular a human being.

Treatment of Disorders

As mentioned above, the compounds of formula I are PDE10A enzyme inhibitors and as such are useful to treat associated neurological and psychiatric disorders.

The invention thus provides a compound of formula I or a pharmaceutically acceptable acid addition salt thereof, as well as a pharmaceutical composition containing such a compound, for use in the treatment of a neurodegenerative disorder, psychiatric disorder or drug addiction in humans.

In one embodiment of the present invention, the neurodegenerative disorder or condition involves neurodegeneration of striatal medium spiny neurons in a human. In a specific embodiment of the present invention, the neurodegenerative disorder or condition is Huntington's disease. In a further embodiment the disorder is dyskinesia associated with dopamine agonist therapy.

In an embodiment the psychiatric disorder is selected from the group consisting of schizophrenia, for example of the paranoid, disorganized, catatonic,

undifferentiated, or residual type; schizophreniform disorder; schizoaffective disorder, for example of the delusional type or the depressive type; delusional disorder; substance-induced psychotic disorder, for example psychosis induced by alcohol, amphetamine, cannabis, cocaine, hallucinogens, inhalants, opioids, or phencyclidine; personality disorder of the paranoid type; and personality disorder of the schizoid type.

This invention further provides a method of treating a drug addiction, for example an alcohol, amphetamine, cocaine, or opiate addiction, in a human, which method comprises administering to said human an amount of a compound of formula I effective in treating addiction, such as drug addiction.

The term "drug addiction", as used herein, means an abnormal desire for a drug and is generally characterized by motivational disturbances such a compulsion to take the desired drug and episodes of intense drug craving.

Other disorders that can be treated according to the present invention are obsessive/compulsive disorders, non-insuline demanding diabetes mellitus (NIDDM), and Tourette's syndrome and other tic disorders as well as Attention Deficit/Hyperactivity Disorder (ADHD).

The compounds of formula I or pharmaceutically acceptable salts thereof may be used in combination with one or more other drugs (including typical and atypical antipsychotic agent) in the treatment of diseases or conditions for which the compounds of the present invention have utility, where the combination of the drugs together are safer or more effective than either drug alone. Additionally, the compounds of the present invention may be used in combination with one or more other drugs that treat, prevent, control, ameliorate, or reduce the risk of side effects or toxicity of the compounds of the present invention. The combinations, uses and methods of treatment of the invention may also provide advantages in treatment of patients who fail to respond adequately or who are resistant to other known treatments.

Such other drugs may be administered, by a route and in an amount commonly used therefore, contemporaneously or sequentially with the compounds of the present invention. Accordingly, the pharmaceutical compositions of the present invention include those that contain one or more other active ingredients, in addition
5 to the compounds of the present invention. The combinations may be administered as part of a unit dosage form combination product, or as a kit or treatment protocol wherein one or more additional drugs are administered in separate dosage forms as part of a treatment regimen.

10 The term "neuroleptic agent" as used herein refers to drugs, which have the effect on cognition and behaviour of antipsychotic agent drugs that reduce confusion, delusions, hallucinations, and psychomotor agitation in patients with psychoses. Also known as major tranquilizers and antipsychotic drugs, neuroleptic agents include, but are not limited to: typical antipsychotic drugs, including phenothiazines,
15 further divided into the aliphatics, piperidines, and piperazines, thioxanthenes (e.g., cisordinol), butyrophenones (e.g., haloperidol), dibenzoxazepines (e.g., loxapine), dihydroindolones (e.g., molindone), diphenylbutylpiperidines (e.g., pimozide), and atypical antipsychotic drugs, including benzisoxazoles (e.g., risperidone), sertindole, olanzapine, quetiapine, osanetant and ziprasidone.

20 Particularly preferred neuroleptic agents for use in the invention are sertindole, olanzapine, risperidone, quetiapine, aripiprazole, haloperidol, clozapine, ziprasidone and osanetant.

25 As used herein, and unless otherwise indicated, a "neurodegenerative disorder or condition" refers to a disorder or condition that is caused by the dysfunction and/or death of neurons in the central nervous system. The treatment of these disorders and conditions can be facilitated by administration of an agent which prevents the dysfunction or death of neurons at risk in these disorders or conditions and/or
30 enhances the function of damaged or healthy neurons in such a way as to compensate for the loss of function caused by the dysfunction or death of at-risk neurons. The term "neurotrophic agent" as used herein refers to a substance or agent that has some or all of these properties.

All references, including publications, patent applications and patents, cited herein are hereby incorporated by reference in their entirety and to the same extent as if each reference were individually and specifically indicated to be incorporated by reference and were set forth in its entirety (to the maximum extent permitted by law).

Headings and sub-headings are used herein for convenience only, and should not be construed as limiting the invention in any way.

The use of any and all examples, or exemplary language (including “for instance”, “for example”, “e.g.”, and “as such”) in the present specification is intended merely to better illuminate the invention, and does not pose a limitation on the scope of invention unless otherwise indicated.

The citation and incorporation of patent documents herein is done for convenience only, and does not reflect any view of the validity, patentability and/or enforceability of such patent documents.

The present invention includes all modifications and equivalents of the subject-matter recited in the claims appended hereto, as permitted by applicable law.

The invention disclosed herein is further illustrated by the following non-limiting examples.


Experimental Section

Preparation of the compounds of the invention

Compounds of the general formula I of the invention may be prepared as described in the following reaction schemes.

Compounds of formula I, wherein L is $-\text{CH}=\text{CH}-$ or $-\text{CH}_2-\text{CH}_2-$ can be prepared by the reaction sequence shown in scheme 1.

23



Scheme 1.

- 5 Specifically, compounds of formula I, wherein L is -CH₂-CH₂- can be prepared by reduction of an alkene of formula I, wherein L is -CH=CH-, by hydrogenation using a transition metal catalyst, such as palladium metal, together with a hydrogen source, such as hydrogen gas, ammonium hydrogen carbonate, or cyclohexadiene. Said alkenes of formula I, wherein L is -CH=CH- can be prepared by the Wittig
- 10 reaction between a phosphonium salt of formula IV and an aldehyde of formula V in a suitable solvent, such as tetrahydrofuran, in the presence of a suitable base, such as 1,8-diazabicyclo[5.4.0]undec-7-ene. Phosphonium salts of formula IV are readily available by reaction of compounds of formula IV (see scheme 1 above) with triphenylphosphine by methods known to chemists skilled in the art and as
- 15 described in e.g. WO-2011072696, WO-2011072694 and WO-2009152825. Aldehydes of formula V are commercially available or available by methods described in the literature see e.g. *Organometallics* (2011), 30(5), 1008-1012, *Journal of Medicinal Chemistry* (2010), 53(24), 8663-8678. *Chemical Communications* (2010), 46(35), 6554-6556, *Journal of Medicinal Chemistry* (2010),

53(5), Science of Synthesis (2005), 15 389-549. Journal of the Chemical Society (1932), Journal of the American Chemical Society (1941), 63 2654-5.

General Methods

5

Analytical LC-MS data were obtained using the following method:

Method 111:

LC-MS were run on a Sciex API150EX equipped with APPI-source operating in positive ion mode. The HPLC consisted of Shimadzu LC10-ADvp LC pumps. SPD-M20A PDA UV-detector (operating at 254 nm) and shimadzu CBM-20A system controller. Autosampler was Gilson 215. Column oven was a Metalox model 200-C and column temperature: 60°C. injector: Gilson model 841 (1 microliter loop). ELS detector was a Sedere Sedex 85.

LC-conditions: The column was a Waters Symmetry C-18. 4.6 x 30 mm. 3.5µm operating at 60 °C with 3.3 ml/min of a binary gradient consisting of Solvent A: 100% H₂O 0.05% TFA and Solvent B: 95% ACN 5% H₂O 0.035% TFA

Injection vol: 10µl (1µl injected on the column)

Gradient: 10% B to 100% B in 2.4 min

10% B in 0.4 min.

Total run time: 2.8 min

Method 131:

LC-MS were run on a Sciex API150EX equipped with APPI-source operating in positive ion mode. The HPLC consisted of Shimadzu LC10-ADvp LC pumps. SPD-M20A PDA UV-detector (operating at 254 nm) and shimadzu CBM-20A system controller. Autosampler was Gilson 215. Column oven was a Jones Chromatography 7990R and column temperature: 60°C. ELS detector was a Sedere Sedex 85.

LC-conditions: The column was a Waters Symmetry C-18. 4.6 x 30 mm. 3.5µm operating at 60 °C with 3.0 ml/min of a binary gradient consisting of Solvent A: H₂O with 0.05% v/v TFA and Solvent B: Methanol with 0.05% TFA

Injection vol: 10µl (1µl injected on the column)

Gradient:

	0.01 min	17% B (v/v)
	0.27 min	28% B (v/v)
5	0.53 min	39% B (v/v)
	0.80 min	50% B (v/v)
	1.07 min	59% B (v/v)
	1.34 min	68% B (v/v)
	1.60 min	78% B (v/v)
10	1.87 min	86% B (v/v)
	2.14 min	93% B (v/v)
	2.38 min	100% B (v/v)
	2.40 min	17% B (v/v)
	2.80 min	17% B (v/v)
15	Total run time: 2.8 min	

Method 132

LC-MS were run on a Sciex API150EX equipped with APPI-source operating in positive ion mode. The HPLC consisted of Shimadzu LC10-ADvp LC pumps. SPD-
20 M20A PDA detector (operating at 254 nm) and SCL-10A system controller. Autosampler was Gilson 215. Column oven was a Jones Chromatography 7990R and ELS detector was a Sedere Sedex 85.

LC-conditions: The column was a Waters Symmetry C-18. 4.6 x 30 mm. 3.5µ
operating at 60 °C with 2.5 ml/min of a binary gradient consisting of water + 0.05 %
25 TFA (A) and methanol + 0.05 % TFA (B).

Gradient:	0.01 min.	5% B
	2.38 min.	100% B
	2.40 min.	5% B
	2.80 min.	5% B
30	Total run time: 2.8 minutes	

Method 350

LC-MS were run on a Sciex API300 equipped with APPI source operating in positive ion mode. The UPLC consisted of Waters Aquity including column manager, binary solvent manager, sample organizer, PDA detector (operating at 254 nm) and ELS
5 detector.

LC-conditions: The column was a Waters Aquity UPLC BEH C-18, 2.1 x 50 mm, 1.7 μ m operating at 60 °C with 1.2 ml/min of a binary gradient consisting of water + 0.05 % TFA (A) and 95 % acetonitrile containing 5 % water + 0.03 % TFA (B).

Gradient:	Time (min.)	%B
10	0.00	10.0
	1.00	100.0
	1.01	10.0
	1.15	10.0

Total run time 1.15 min

Preparative LC-MS-purification was performed on a PE Sciex API 150EX instrument with atmospheric pressure chemical ionization. Column: 50 X 20 mm YMC ODS-A with 5 μ m particle size; Method: Linear gradient elution with A:B = 80:20 to 0:100 in 7 minutes and with a flow rate of 22.7 mL/minute. Fraction collection was
20 performed by split-flow MS detection.


¹H NMR spectra were recorded at 500.13 MHz on a Bruker Avance AV500 instrument or at 600.16 MHz on a Bruker Avance Ultrashield plus instrument. TMS was used as internal reference standard. Chemical shift values are expressed in
25 ppm. The following abbreviations are used for multiplicity of NMR signals: s = singlet, d = doublet, t = triplet, q = quartet, qui = quintet, h = heptet, dd = double doublet, dt = double triplet, dq = double quartet, td = triplet of doublets, tt = triplet of triplets, m = multiplet, br s = broad singlet and br = broad signal.

Abbreviations are in accordance with to the ACS Style Guide: "The ACS Styleguide – A manual for authors and editors" Janet S. Dodd, Ed. 1997, ISBN: 0841234620
30

Preparation of intermediates

Phosphonium salts of formula **IV** shown in **scheme 1** are readily available by reaction of compounds of formula **IV** (see scheme 1 above) with triphenylphosphine by methods known to chemists skilled in the art and as described in e.g. WO-2011072696, WO-2011072694 and WO-2009152825. Aldehydes of formula **V** are commercially available or available by methods described in the literature see e.g. Organometallics (2011), 30(5), 1008-1012, Journal of Medicinal Chemistry (2010), 53(24), 8663-8678. Chemical Communications (2010), 46(35), 6554-6556, Journal of Medicinal Chemistry (2010), 53(5), Science of Synthesis (2005), 15 389-549. Journal of the Chemical Society (1932), Journal of the American Chemical Society (1941), 63 2654-5.

2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-quinoline-6-carbonitrile



To a suspension of (5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)methyl-triphenylphosphonium; chloride (0.222 g, 0.483 mmol) and 2-Formyl-quinoline-6-carbonitrile (80 mg, 0.4 mmol) in dry Tetrahydrofuran (6 mL, 80 mmol) was added 1,8-Diazabicyclo[5.4.0]undec-7-ene (66 μ L, 0.44 mmol) (reaction mixture turns slightly yellow for a while and precipitation changes character) and the mixture was stirred at room temperature under an atmosphere of Argon overnight. The mixture was rotovaped and the THF was evaporated off. The solid was dissolved in DCM and was chromatographed on silicagel (0-30% MeOH in EtOAc). Pure product 2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-quinoline-6-carbonitrile precipitates in one of the fractions (10): isolated by filtration. Yield: 6 mg white solid.

The following intermediates were made in a similar way:


2-[2-(5,7-Dimethyl-imidazo[1,2-a]pyrimidin-2-yl)-vinyl]-quinoline

- 2-[2-(5,7-Dimethyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl)-vinyl]-quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-quinoline
2-[2-(5,7-Dimethyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl)-vinyl]-6-methoxy-quinoline
5 2-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-vinyl]-quinoline
2-[2-(8-Ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-vinyl]-6-fluoro-quinoline
6-Fluoro-2-[2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-vinyl]-quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-vinyl]-quinoline
2-[2-(6-Fluoro-quinolin-2-yl)-vinyl]-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-8-ol
10 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-4-methyl-quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-4-methoxy-quinoline
4-Methoxy-2-[2-(5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-vinyl]-quinoline
4-Methoxy-2-[2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-vinyl]-
quinoline
15 4-Methyl-2-[2-(5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-vinyl]-quinoline
2-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-vinyl]-4-methyl-quinoline
4-Chloro-8-fluoro-2-[(E)-2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-
vinyl]-quinoline
8-Fluoro-2-[2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-vinyl]-quinoline
20 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-7-fluoro-quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-fluoro-quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-fluoro-quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-4-fluoro-quinoline
2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-fluoro-quinoline
25 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-7-fluoro-4-methoxy-
quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-7-fluoro-quinolin-4-ol
2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-7-trifluoromethyl-
quinoline
30 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-fluoro-4-methoxy-
quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-7-trifluoromethyl-quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-fluoro-quinolin-4-ol

- 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-5-fluoro-quinoline
7-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-c]pyrimidin-2-yl)-vinyl]-6-isopropyl-quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-5,7-difluoro-quinoline
5 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-5,6,8-trifluoro-quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6,8-difluoro-quinoline
6-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-[1,3]dioxolo[4,5-
g]quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-fluoro-8-methyl-
10 quinoline
2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-fluoro-7-methyl-
quinoline
6-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-2,2-difluoro-
[1,3]dioxolo[4,5-g]quinoline
15 7-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-quinoline-6-
carbonitrile
7-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-2,3-dihydro-
[1,4]dioxino[2,3-g]quinoline
6-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-quinoline
20 6-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-8-fluoro-
quinoline
8-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-methyl-
quinoline
5,7-Dichloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-quinoline

Preparation of compounds of the invention

Example 1: 2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-trifluoromethyl-quinoline



- 10 To a solution of (5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-ylmethyl)-triphenylphosphonium; chloride (0.48 g, 1.0 mmol) and 6-Trifluoromethyl-quinoline-2-carbaldehyde (0.24 g, 1.0 mmol) in dry N,N-Dimethylformamide (25 mL, 320 mmol) was added 1,8-Diazabicyclo[5.4.0]undec-7-ene (0.16 mL, 1.0 mmol) (reaction mixture turns more dark) and the mixture was stirred at room
- 15 temperature under an atmosphere of Argon overnight. The reaction mixture shows precipitation the day after.

- Precipitation filtered off. Washed with water and diethyl ether. Dried on filter by vacuum, then in vacuo for 2 hours at 60 °C. Filtercake: Giving a white solid containing the final product 2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-trifluoromethyl-quinoline. LC-MS: m/z = 369.7 (MH⁺). Rt = 1.96 min., method = 131.
- 20

The following compounds were made in a similar way:


- 2-[(E)-2-(5,7-Dimethyl-imidazo[1,2-a]pyrimidin-2-yl)-vinyl]-quinoline LC-MS: m/z = 301.1 (MH⁺). Rt = 0.55 min., method = 111.
- 25 2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-trifluoromethoxy-quinoline LC-MS: m/z = 386.1, Rt = 1.97 min, method = 131
- 2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-6-cyano-quinoline LC-MS: m/z = 327.3, Rt = 1.97 min, method = 131

2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-7-methoxy-quinoline LC-MS: m/z = 332.1, R_t = 1,22 min, method = 131

2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-5-methoxy-quinoline LC-MS: m/z = 332.2, R_t = 1,41 min, method = 131

5

Example 2: 6-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-1,3-dioxolo[4,5-g]quinoline



10

6-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-1,3-dioxolo[4,5-g]quinoline (0.183 g, 0.530 mmol) was dissolved in N,N-Dimethylformamide (11 mL, 140 mmol). [B] p-Toluenesulfonylhydrazide (0.296 g, 1.59 mmol; Supplier =

15 Avocado) was added and the reaction was stirred at 130 °C under an atmosphere of Argon ON. LCMS was done and showed almost complete conversion. 0.100 g [B] was added to the mixture was stirred 2 days at 130 C.

DMF was evaporated. The solid was dissolved in 50 mL EtOAc and extracted with 2 x 25 mL sat. NaHCO₃ and washed with 50 mL brine. The organic phase was
20 rotovaped and chromatographed on silicagel using EtOAc: heptane (1:1) and then 0-30% MeOH in EtOAc. Yield: 40 mg solid. LC-MS: m/z = 348.4 (MH⁺). R_t = 0.34 min., method = 350.

The following compounds were prepared in a similar way:

25

2-[2-(5,7-Dimethyl-imidazo[1,2-a]pyrimidin-2-yl)-ethyl]-quinoline LC-MS: m/z = 303,4 (MH⁺). R_t = 0,34 min., method = 111

2-[2-(5,7-Dimethyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl)-ethyl]-quinoline LC-MS: m/z = 304,3 (MH⁺). R_t = 0,46 min., method = 111

30 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-quinoline LC-MS: m/z = 304,3 (MH⁺). R_t = 0,61 min., method = 131

2-[2-(5,7-Dimethyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl)-ethyl]-6-methoxy-quinoline

LC-MS: m/z = 334,5 (MH⁺). Rt = 0,62 min., method = 131

2-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline LC-MS:

m/z = 319,1 (MH⁺). Rt = 0,71 min., method = 131

5 2-[2-(8-Ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-6-fluoro-quinoline LC-

MS: m/z = 335,2 (MH⁺). Rt = 1,12 min., method = 131

6-Fluoro-2-[2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline

LC-MS: m/z = 337,5 (MH⁺). Rt = 0,96 min., method = 131

2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline LC-MS: m/z =

10 303,5 (MH⁺). Rt = 0,76 min., method = 131

2-[2-(6-Fluoro-quinolin-2-yl)-ethyl]-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-8-ol LC-MS:

m/z = 323,1 (MH⁺). Rt = 0,41 min., method = 350

2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-4-methyl-quinoline LC-

MS: m/z = 318,2 (MH⁺). Rt = 0,83 min., method = 131

15 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-4-methoxy-quinoline LC-

MS: m/z = 334,5 (MH⁺). Rt = 0,9 min., method = 131

4-Methoxy-2-[2-(5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline LC-MS:

m/z = 319,2 (MH⁺). Rt = 0,93 min., method = 131

4-Methoxy-2-[2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-

20 quinoline LC-MS: m/z = 349,1 (MH⁺). Rt = 1,01 min., method = 131

4-Methyl-2-[2-(5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline LC-MS:

m/z = 303,4 (MH⁺). Rt = 0,85 min., method = 131

2-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-4-methyl-quinoline

LC-MS: m/z = 333,2 (MH⁺). Rt = 0,95 min., method = 131

25 4-Chloro-8-fluoro-2-[(E)-2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-

vinyl]-quinoline LC-MS: m/z = 369,2 (MH⁺). Rt = 1,98 min., method = 131

8-Fluoro-2-[2-(8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-quinoline

LC-MS: m/z = 337,5 (MH⁺). Rt = 1,32 min., method = 131

2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-7-fluoro-quinoline LC-MS:

30 m/z = 321,8 (MH⁺). Rt = 0,43 min., method = 350

2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-fluoro-quinoline LC-MS:

m/z = 322,1 (MH⁺). Rt = 0,44 min., method = 350

2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-4-fluoro-quinoline LC-MS:
m/z = 321,9 (MH⁺). Rt = 0,44 min., method = 350

2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-7-fluoro-4-methoxy-
quinoline LC-MS: m/z = 352,3 (MH⁺). Rt = 0,87 min., method = 131

5 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-7-fluoro-quinolin-4-ol LC-
MS: m/z = 338,4 (MH⁺). Rt = 1,08 min., method = 131

2-[(E)-2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-vinyl]-7-trifluoromethyl-
quinoline LC-MS: m/z = 370,2 (MH⁺). Rt = 0,79 min., method = 350

10 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-fluoro-4-methoxy-
quinoline LC-MS: m/z = 352,3 (MH⁺). Rt = 0,9 min., method = 131

2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-7-trifluoromethyl-quinoline
LC-MS: m/z = 372,3 (MH⁺). Rt = 0,61 min., method = 350

2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-fluoro-quinolin-4-ol LC-
MS: m/z = 338,1 (MH⁺). Rt = 1,05 min., method = 131

15 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-5-fluoro-quinoline LC-MS:
m/z = 322,1 (MH⁺). Rt = 1,25 min., method = 131

7-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-quinoline LC-
MS: m/z = 338,3 (MH⁺). Rt = 1,82 min., method = 132

20 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-c]pyrimidin-2-yl)-ethyl]-6-isopropyl-quinoline
LC-MS: m/z = 346,2 (MH⁺). Rt = 0,47 min., method = 350

2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-5,7-difluoro-quinoline LC-
MS: m/z = 339,8 (MH⁺). Rt = 0,57 min., method = 350

2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-5,6,8-trifluoro-quinoline
LC-MS: m/z = 358,4 (MH⁺). Rt = 0,67 min., method = 350

25 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6,8-difluoro-quinoline LC-
MS: m/z = 339,7 (MH⁺). Rt = 0,6 min., method = 350

6-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-[1,3]dioxolo[4,5-
g]quinoline LC-MS: m/z = 348,4 (MH⁺). Rt = 0,34 min., method = 350

30 2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-fluoro-8-methyl-
quinoline LC-MS: m/z = 336,3 (MH⁺). Rt = 0,57 min., method = 350

2-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-fluoro-7-methyl-
quinoline LC-MS: m/z = 336,3 (MH⁺). Rt = 0,41 min., method = 350

6-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-2,2-difluoro-
[1,3]dioxolo[4,5-g]quinoline LC-MS: m/z = 384,2 (MH⁺). Rt = 1,6 min., method =
131

7-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-quinoline-6-
5 carbonitrile LC-MS: m/z = 363,2 (MH⁺). Rt = 1,61 min., method = 131

7-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-2,3-dihydro-
[1,4]dioxino[2,3-g]quinoline LC-MS: m/z = 362,3 (MH⁺). Rt = 0,82 min., method =
131

6-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-quinoline LC-
10 MS: m/z = 338,3 (MH⁺). Rt = 1,31 min., method = 131

6-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-8-fluoro-
quinoline LC-MS: m/z = 356,2 (MH⁺). Rt = 1,72 min., method = 131

8-Chloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-6-methyl-
quinoline LC-MS: m/z = 352,4 (MH⁺). Rt = 1,7 min., method = 131

15 5,7-Dichloro-2-[2-(5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-quinoline LC-
MS: m/z = 372,0 (MH⁺). Rt = 2,01 min., method = 131

Pharmacological Testing

20 PDE10A enzyme

Active PDE10A enzyme is prepared in a number of ways for use in PDE assays
(Loughney, K. *et al. Gene* **1999**, 234, 109-117; Fujishige, K. *et al. Eur J Biochem.*
1999, 266, 1118-1127 and Soderling, S. *et al. Proc. Natl. Acad. Sci.* **1999**, 96, 7071-
7076). PDE10A can be expressed as full-length proteins or as truncated proteins, as
25 long as they express the catalytic domain. PDE10A can be prepared in different cell
types, for example insect cells or E. coli. An example of a method to obtain
catalytically active PDE10A is as follows: The catalytic domain of human PDE10A
(amino acids 440-779 from the sequence with accession number NP 006652) is
amplified from total human brain total RNA by standard RT-PCR and is cloned into
30 the BamH1 and Xho1 sites of the pET28a vector (Novagen). Expression in coli is
performed according to standard protocols. Briefly, the expression plasmids are
transformed into the BL21(DE3) E. coli strain, and 50 mL cultures inoculated with

the cells allowed to grow to an OD600 of 0.4-0.6 before protein expression is induced with 0.5mM IPTG. Following induction, the cells are incubated overnight at room temperature, after which the cells are collected by centrifugation. Cells expressing PDE10A are resuspended in 12 mL (50 mM TRIS-HCl-pH8.0, 1 mM MgCl₂ and protease inhibitors). The cells are lysed by sonication, and after all cells are lysed, TritonX100 is added according to Novagen protocols. PDE10A is partially purified on Q sepharose and the most active fractions were pooled.

PDE10A inhibition assay

A PDE10A assay may for example, be performed as follows: The assay is performed in 60 uL samples containing a fixed amount of the relevant PDE enzyme (sufficient to convert 20-25% of the cyclic nucleotide substrate), a buffer (50 mM HEPES7.6; 10mM MgCl₂; 0.02% Tween20), 0.1mg/ml BSA, 225 pCi of ³H-labelled cyclic nucleotide substrate, tritium labeled cAMP to a final concentration of 5 nM and varying amounts of inhibitors. Reactions are initiated by addition of the cyclic nucleotide substrate, and reactions are allowed to proceed for one hr at room temperature before being terminated through mixing with 15 uL 8 mg/mL yttrium silicate SPA beads (Amersham). The beads are allowed to settle for one hr in the dark before the plates are counted in a Wallac 1450 Microbeta counter. The measured signal can be converted to activity relative to an uninhibited control (100 %) and IC₅₀ values can be calculated using the Xlfit extension to EXCEL.

Phencyclidine (PCP) induced hyperactivity

Male mice (NMRI, Charles River) weighing 20-25g are used. Eight mice are used in each group receiving the test compound (5 mg/kg) plus PCP (2.3 mg/kg) including the parallel control groups receiving the vehicle of the test compound plus PCP or vehicle injections only. The injection volumen is 10 ml/kg. The experiment is made in normal light conditions in an undisturbed room. The test substance is injected per oss 60 min before injection of PCP, which is administered subcutaneous.

Immediately after injection of PCP the mice are placed individually in special designed test cage (20 cm x 32 cm). The activity is measured by 5X8 infrared light


sources and photocells spaced by 4 cm. The light beams cross the cage 1.8 cm above the bottom of the cage. Recording of a motility count requires interruption of adjacent light beams, thus avoiding counts induced by stationary movements of the mice.

- 5 Motility is recorded in 5 min intervals for a period of 1 hour. The drug effect is calculated on the total counts during the 1 hour behavioral test period in the following manner:

The mean motility induced by vehicle treatment in the absence of PCP is used as baseline. The 100 per cent effect of PCP is accordingly calculated to be total motility
10 counts minus baseline. The response in groups receiving test compound is thus determined by the total motility counts minus baseline, expressed in per cent of the similar result recorded in the parallel PCP control group. The per cent responses are converted to per cent inhibition.

What is claimed:


1. A compound having the structure of formula I:



wherein R_1 , R_2 , R_3 , R_4 , R_5 and R_6 are individually selected from the group consisting of hydrogen, hydroxyl, cyano, C_1 - C_6 alkyl; C_1 - C_6 alkoxy, halogen, methylenedioxy, difluoromethylenedioxy and ethylenedioxy

wherein -L- is a linker selected from $-CH_2-CH_2-$ and $-CH=CH-$

wherein HET is selected from the group consisting of



wherein one or more of the carbon-bound hydrogen in the HET optionally may be substituted with up to three substituents R_7 , R_8 and R_9 individually selected from C_1 - C_6 alkyl; halogen; cyano, halo(C_1 - C_6)alkyl; aryl, alkoxy and C_1 - C_6 hydroxyalkyl

and wherein * denotes the attachment point,

and tautomers and pharmaceutically acceptable salts thereof, and polymorphic forms thereof.

- 5 2. The compound of claim 1 wherein one or more of R1 to R6 is selected from the group consisting of C₁-C₃ alkyl substituted with one or more F and unsubstituted C₁-C₃ alkyl.
 - 10 3. The compound of claim 1 wherein one or more of R1-R6 is selected from the group consisting of methoxy, difluoromethoxy and trifluoromethoxy.
 4. The compound of claim 1 wherein one or more of R1-R6 is selected from the group consisting of fluorine and chlorine.
 - 15 5. The compound of claim 1 wherein -L- is -CH₂-CH₂-
 6. The compound of claim 1 wherein -L- is -CH=CH-
 - 20 7. The compound of any of claims 1 to 6 wherein HET is selected from the group consisting of
- wherein * denotes the attachment point.

- 25 8. The compound of claim 1 wherein HET is substituted with one substituent R7 selected from the group consisting of C₁-C₆ alkyl, such as methyl; halogen, such as chlorine or bromine; cyano; halo(C₁-C₆)alkyl, such as trifluoromethyl; aryl such as phenyl; and C₁-C₆ hydroxyalkyl such as CH₂CH₂OH.
- 30 9. The compound of claim 1 or 8 wherein HET is substituted with two substituents R7 and R8 individually selected from the group consisting of C₁-C₆ alkyl, such as methyl; halogen, such as chlorine or bromine; cyano;

halo(C₁-C₆)alkyl, such as trifluoromethyl; aryl such as phenyl; and C₁-C₆ hydroxyalkyl such as CH₂CH₂OH.

- 5 10. The compound of claim 1 or 8 wherein HET is substituted with three substituents R₇, R₈ and R₉ individually selected from the group consisting of C₁-C₆ alkyl, such as methyl; halogen, such as chlorine or bromine; cyano; halo(C₁-C₆)alkyl, such as trifluoromethyl; aryl such as phenyl; and C₁-C₆ hydroxyalkyl such as CH₂CH₂OH.
- 10 11. The compound of claim 1 or 8 wherein HET is unsubstituted.
12. The compound of claim 1 wherein HET is selected from the group consisting of (5,7-Dimethyl-imidazo[1,2-a]pyrimidin-2-yl), 5,7-Dimethyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl), (5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl), (8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl) and 5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl).
- 15 13. The compound of any of claim 1 to 12, or a pharmaceutically acceptable salt thereof, for use as a medicament.
- 20 14. A pharmaceutical composition comprising a therapeutically effective amount of the compound of any of claims 1-12 and a pharmaceutically acceptable carrier, diluent or excipient.
- 25 15. The use of the compound of any of claims 1-12, or a pharmaceutically acceptable salt thereof, for the preparation of a medicament for the treatment of a neurodegenerative or psychiatric disorder.