

Bacteria-based protein delivery

The field of the invention

The present invention relates to recombinant Gram-negative bacterial strains and the use thereof for delivery of repeated domains of a heterologous protein or two or more domains of different heterologous proteins into eukaryotic cells.

Background of the invention

Transient transfection techniques have been applied in cell biological research over many years to address protein functions. These methods generally result in a massive overrepresentation of the protein under study, which might lead to oversimplified models of signalling. For proteins controlling short-lived signalling processes, the protein of interest is present for far longer as the signalling event it controls. Even more, DNA transfection based transient over-expression leads to a heterogenous and unsynchronized cell population, which complicates functional studies and hampers – omics approaches. Besides this, the upscaling of such assays to a larger scale is very expensive. Some of the above mentioned points are covered by existing techniques as microinjection or proteo-fection of purified proteins, the inducible translocation strategy to rapidly target plasmid born small GTPases to the cell membrane or the addition of purified proteins fused to cell-permeable bacterial toxins. But these techniques are all time-consuming and cumbersome and to our knowledge none fulfils all mentioned criteria.

Bacteria have evolved different mechanisms to directly inject proteins into target cells [1]. The type III secretion system (T3SS) used by bacteria like *Yersinia*, *Shigella* and *Salmonella* [2] functions like a nano-syringe that injects so-called bacterial effector proteins into host cells. Bacterial proteins to be secreted via the T3SS, called effectors, harbour a short N-terminal secretion signal [3]. Inside bacteria, some effectors are bound by chaperones. Chaperones might mask toxic domains, they contribute to exposition of the secretion signal and keep the substrates in a secretion-competent conformation, therefore facilitating secretion. Upon induction of secretion, an ATPase adjacent to the T3SS removes the chaperones and the effectors travel

unfolded or only partially folded through the needle, and refold once in the host cytoplasm.

T3S has been exploited to deliver hybrid peptides and proteins into target cells. Heterologous bacterial T3SS effectors have been delivered in case the bacterium under study is hardly accessible by genetics (like *Chlamydia trachomatis*). Often reporter proteins were fused to possible T3SS secretion signals as to study requirements for T3SS dependent protein delivery, such as the *Bordetella pertussis* adenylate cyclase, murine DHFR or a phosphorylatable tag. Peptide delivery was mainly conducted with the aim of vaccination. This includes viral epitopes, bacterial epitopes (listeriolysin O) as well as peptides representing epitopes of human cancer cells. In few cases functional eukaryotic proteins have been delivered to modulate the host cell, as done with nanobodies [4], nuclear proteins (Cre-recombinase, MyoD) [5,6] or IL10 and IL1ra [7]. None of the above-mentioned systems allows single-protein delivery as in each case one or multiple endogenous effector proteins are still encoded. Furthermore, the vectors used have not been designed in a way allowing simple cloning of other DNA fragments encoding proteins of choice, hindering broad application of the system. Surprisingly it has been found that delivery of repeated domains of heterologous proteins or combinations of domains of different heterologous proteins to eukaryotic cells enlarges the impact on a desired cellular pathway.

Summary of the invention

The present invention relates generally to recombinant Gram-negative bacterial strains and the use thereof for delivery of repeated domains of a heterologous protein or two or more domains of different heterologous proteins into eukaryotic cells. The present invention provides Gram-negative bacterial strains and the use thereof, which allows the translocation of repeated domains of a heterologous protein or two or more domains of different heterologous proteins such as various type III effectors, but also of type IV effectors, viral proteins and most importantly functional eukaryotic proteins. Means for fluorescent tracking of delivery, for relocalization to the nucleus and notably for removal of the bacterial appendage after delivery to the host cell are provided. The presented T3SS based system results in scalable, rapid, synchronized,

homogenous and tunable delivery of repeated domains of a heterologous protein or two or more domains of different heterologous proteins of interest. The delivery system of the present invention is suitable to inject repeated domains of a eukaryotic protein or two or more domains of different eukaryotic proteins in living animals and can be used for therapeutic purposes.

In a first aspect the present invention relates to a recombinant Gram-negative bacterial strain transformed with a vector which comprises in the 5' to 3' direction:

a promoter;

a first DNA sequence encoding a delivery signal from a bacterial T3SS effector protein, operably linked to said promoter; and

a second DNA sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins fused in frame to the 3' end of said first DNA sequence, wherein the heterologous proteins are selected from the group consisting of proteins involved in apoptosis or apoptosis regulation, cell cycle regulators, ankyrin repeat proteins, cell signaling proteins, reporter proteins, transcription factors, proteases, small GTPases, GPCR related proteins, nanobody fusion constructs and nanobodies, bacterial T3SS effectors, bacterial T4SS effectors and viral proteins.

In a further aspect the present invention relates to a recombinant Gram-negative bacterial strain transformed with a vector which comprises in the 5' to 3' direction:

a first DNA sequence encoding a delivery signal or a fragment thereof from a bacterial effector protein; and

a second DNA sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins fused in frame to the 3' end of said first DNA sequence, wherein the heterologous proteins are selected from the group consisting of proteins involved in apoptosis or apoptosis regulation, cell cycle regulators, ankyrin repeat proteins, cell signaling proteins, reporter proteins, transcription factors, proteases, small GTPases, GPCR related proteins, nanobody fusion constructs and nanobodies, bacterial T3SS effectors, bacterial T4SS effectors and viral proteins.

In a further aspect the present invention relates to a vector which comprises in the 5' to 3' direction:

a first DNA sequence encoding a delivery signal or a fragment thereof from a bacterial effector protein; and
a second DNA sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins fused in frame to the 3' end of said first DNA sequence, wherein the heterologous proteins are selected from the group consisting of proteins involved in apoptosis or apoptosis regulation, cell cycle regulators, ankyrin repeat proteins, cell signaling proteins, reporter proteins, transcription factors, proteases, small GTPases, GPCR related proteins, nanobody fusion constructs and nanobodies, bacterial T3SS effectors, bacterial T4SS effectors and viral proteins.

In a further aspect the present invention relates to a vector which comprises in the 5' to 3' direction:

a promoter;
a first DNA sequence encoding a delivery signal from a bacterial T3SS effector protein, operably linked to said promoter;
a second DNA sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins fused in frame to the 3' end of said first DNA sequence,
wherein the heterologous proteins are selected from the group consisting of proteins involved in apoptosis or apoptosis regulation, cell cycle regulators, ankyrin repeat proteins, cell signaling proteins, reporter proteins, transcription factors, proteases, small GTPases, GPCR related proteins, nanobody fusion constructs and nanobodies, bacterial T3SS effectors, bacterial T4SS effectors and viral proteins.

The present invention further relates to a method for delivering repeated domains of a heterologous protein or two or more domains of different heterologous proteins into a eukaryotic cell comprising the following steps:

- i) culturing a Gram-negative bacterial strain; and
- ii) contacting a eukaryotic cell with the Gram-negative bacterial strain of i) wherein a fusion protein which comprises a delivery signal from a bacterial T3SS effector protein and the repeated domains of a heterologous protein or two or more domains of

different heterologous proteins is expressed by the Gram-negative bacterial strain and is translocated into the eukaryotic cell.

The present invention further relates to a method for delivering repeated domains of a heterologous protein or two or more domains of different heterologous proteins into a eukaryotic cell comprising the following steps:

- i) culturing a Gram-negative bacterial strain;
- ii) contacting a eukaryotic cell with the Gram-negative bacterial strain of i) wherein a fusion protein which comprises a delivery signal from a bacterial T3SS effector protein and the repeated domains of a heterologous protein or two or more domains of different heterologous proteins is expressed by the Gram-negative bacterial strain and is translocated into the eukaryotic cell; and
- iii) cleaving the fusion protein so that the repeated domains of a heterologous protein or the two or more domains of different heterologous proteins are cleaved from the delivery signal from the bacterial T3SS effector protein.

In a further aspect the present invention relates to a library of Gram-negative bacterial strains, wherein the repeated domains of a heterologous protein or the two or more domains of different heterologous proteins encoded by the second DNA sequence of the expression vector of the Gram-negative bacterial strains are domains of a human or murine protein and, wherein each domain of a human or murine protein expressed by a Gram-negative bacterial strain is different in amino acid sequence.

Brief description of the figures

Fig. 1: Characterization of T3SS protein delivery. (A) Schematic representation of T3SS dependent protein secretion into the surrounding medium (in-vitro secretion)(left side) or into eukaryotic cells (right side). I: shows the type 3 secretion system. II indicates proteins secreted into the surrounding medium, III proteins translocated through the membrane into the cytosol of eukaryotic cells (VII). VI shows a stretch of the two bacterial membranes in which the T3SS is inserted and the bacterial cytosol underneath. IV is a fusion protein attached to the YopE₁₋₁₃₈ N-terminal fragment (V) (B) In-vitro secretion of I: *Y. enterocolitica* E40 wild type, II: *Y. enterocolitica* Δ HOPEMT *asd* or III: *Y. enterocolitica* Δ HOPEMT *asd* + pBadSi_2

as revealed by Western blotting on total bacterial lysates (IV) and precipitated culture supernatants (V) using an anti-YopE antibody.

Fig. 2: Characterization of T3SS protein delivery into epithelial cells. (A) Anti-Myc immunofluorescence staining on HeLa cells infected at an MOI of 100 for 1 h with I: *Y. enterocolitica* Δ HOPEMT *asd* or II: *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si2. **(B)** Quantification of anti-Myc immunofluorescence staining intensity from (A) within HeLa cells. Data were combined from n = 20 sites, error bars indicated are standard error of the mean. I: uninfected, II: *Y. enterocolitica* Δ HOPEMT *asd* or III: *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si2. Y-axis indicates anti-Myc staining intensity [arbitrary unit], x-axis indicates time of infection in minutes **(C)** Quantification of Anti-Myc immunofluorescence staining intensity within cells. HeLa cells were infected for 1h with *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si2 at an MOI indicated on the x-axis. Data were combined from n = 20 sites, error bars indicated are standard error of the mean. Y-axis indicates anti-Myc staining intensity [a.u.].

Fig. 3: Modifications of the T3SS based protein delivery allow nuclear localization of a YopE₁₋₁₃₈ fusion protein (EGFP). EGFP signal in HeLa cells infected with I: *Y. enterocolitica* Δ HOPEMT *asd* or II: *Y. enterocolitica* Δ HOPEMT *asd* Δ yopB carrying the plasmids III: +YopE₁₋₁₃₈-EGFP or IV: +YopE₁₋₁₃₈-EGFP-NLS at an MOI of 100. EGFP signal is shown in “a”, for localization comparison nuclei were stained in “b”.

Fig. 4: Modifications of the T3SS based protein delivery allow removal of the YopE₁₋₁₃₈ appendage. HeLa cells are infected with two different *Y. enterocolitica* strains at the same time, which is reached by simple mixing of the two bacterial suspensions. One strain is delivering the TEV protease fused to YopE₁₋₁₃₈, while the other strain delivers a protein of interest fused to YopE₁₋₁₃₈ with a linker containing a double TEV protease cleavage site. After protein delivery into the eukaryotic cell, the TEV protease will cleave the YopE₁₋₁₃₈ appendage from the protein of interest **(A)** Digitonin lysed HeLa cells uninfected (II) or after infection (MOI of 100) for 2h with I: *Y. enterocolitica* Δ HOPEMT *asd* and III: + pBadSi_2, IV: + YopE₁₋₁₃₈-2x TEV cleavage site-Flag-INK4C, V: + YopE₁₋₁₃₈-2x TEV cleavage site-Flag-INK4C and further overnight treatment with purified TEV protease and VI: + YopE₁₋₁₃₈-2x TEV cleavage site-Flag-INK4C and a second strain + YopE₁₋₁₃₈-TEV were analyzed by

Western blotting anti-INK4C (shown in “a”) for the presence of YopE₁₋₁₃₈ – 2x TEV cleavage site –Flag-INK4C or its cleaved form Flag-INK4C. As a loading control western blotting anti-Actin was performed (shown in “b”). In one case (V) the lysed cells were incubated overnight with purified TEV protease. **(B)** Actin normalized quantification of anti-INK4C staining intensity (shown as [a.u.] on the y-axis) from (A) at the size of full length YopE₁₋₁₃₈-2x TEV cleavage site-Flag-INK4C, where sample IV is set to 100%. I: *Y. enterocolitica* ΔHOPEMT *asd* and IV: + YopE₁₋₁₃₈-2x TEV cleavage site-Flag-INK4C, V: + YopE₁₋₁₃₈-2x TEV cleavage site-Flag-INK4C and further overnight treatment with purified TEV protease and VI: + YopE₁₋₁₃₈-2x TEV cleavage site-Flag-INK4C and a second strain + YopE₁₋₁₃₈-TEV. Data were combined from n = 2 independent experiments, error bars indicated are standard error of the mean **(C)** Digitonin lysed HeLa cells uninfected (II) or after infection (MOI of 100) for 2h with I: *Y. enterocolitica* ΔHOPEMT *asd* and III: + pBadSi₂, IV: + YopE₁₋₁₃₈-2x TEV cleavage site-ET1-Myc, V: + YopE₁₋₁₃₈-2x TEV cleavage site-ET1-Myc and further overnight treatment with purified TEV protease and VI: + YopE₁₋₁₃₈-2x TEV cleavage site-ET1-Myc and a second strain + YopE₁₋₁₃₈-TEV were analyzed by Western blotting anti-Myc (shown in “a”) for the presence of YopE₁₋₁₃₈ – 2x TEV cleavage site - ET1-Myc or its cleaved form ET1-Myc. As a loading control western blotting anti-Actin was performed (shown in “b”) In one case (V) the lysed cells were incubated overnight with purified TEV protease.

Fig. 5: Delivery of bacterial effector proteins into eukaryotic cells (A) HeLa cells were infected with I: *Y. enterocolitica* ΔHOPEMT *asd* carrying II: pBad_Si2 or III: YopE₁₋₁₃₈-SopE at an MOI of 100 for the time indicated above the images (2, 10 or 60 minutes). After fixation cells were stained for the actin cytoskeleton **(B)** HeLa cells were left uninfected (II) or infected with I: *Y. enterocolitica* ΔHOPEMT *asd* carrying III: YopE₁₋₁₃₈-SopE-Myc and in some cases coinfecting with IV: YopE₁₋₁₃₈-SptP at the MOI indicated below the strain (MOI 50; MOI50:MOI50 or MOI50:MOI100) for 1h. After fixation cells were stained for the actin cytoskeleton (shown in “a”) and the presence of the YopE₁₋₁₃₈-SopE-Myc fusion protein was followed via staining anti-Myc (shown in “b”).

Fig. 6: Delivery of bacterial effector proteins into eukaryotic cells (A) Phospho-p38 (“a”), total p38 (“b”) and actin (“c”) western blot analysis on HeLa cells left untreated (II) or infected for 75 min with I: *Y. enterocolitica* ΔHOPEMT *asd* carrying

III: pBad_Si2 or IV: YopE₁₋₁₃₈-OspF at an MOI of 100. Cells were stimulated with TNF α for the last 30 min of the infection as indicated (+ stands for addition of TNF α , - represent no treatment with TNF α) **(B)** Phospho-Akt T308 (“a”) and S473 (“b”) and actin (“c”) western blot analysis on HeLa cells left untreated (II) or infected for 22.5 or 45 min (indicated below the blots) with I: *Y. enterocolitica* Δ HOPEMT *asd* carrying III: pBad_Si2, IV: YopE₁₋₁₃₈-SopE or V: YopE₁₋₁₃₈-SopB at an MOI of 100 **(C)** cAMP levels (in fmol/well shown on y-axis) in HeLa cells left untreated (I) or infected for 2.5h with V: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-BepA, VI: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-BepA_{E305-end}, VII: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-BepG_{Bid} or VIII: *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si2 at an MOI of 100. Cholera toxin (CT) was added for 1h as positive control to samples II (1 μ g/ml), III (25 μ g/ml) or IV (50 μ g/ml). Data were combined from n = 3 independent experiments, error bars indicated are standard error of the mean. Statistical analysis was performed using an unpaired two-tailed t-test (ns indicates a non significant change, ** indicates a p value < 0.01, *** indicates a p value < 0.001).

Fig. 7: Delivery of human tBid into eukaryotic cells induces massive apoptosis.

(A) Cleaved Caspase 3 p17 (“a”) and actin (“b”) western blot analysis on HeLa cells left untreated (II) or infected for 60 min with I: *Y. enterocolitica* Δ HOPEMT *asd* carrying III: pBad_Si2, IV: YopE₁₋₁₃₈-Bid or V: YopE₁₋₁₃₈-t-Bid at an MOI of 100. In some cases, cells were treated with VI: 0.5 μ M Staurosporine or VII: 1 μ M Staurosporine **(B)** Digitonin lysed HeLa cells left untreated (II) or after infection for 1h with I: *Y. enterocolitica* Δ HOPEMT *asd* carrying III: pBad_Si2, IV: YopE₁₋₁₃₈-Bid or V: YopE₁₋₁₃₈-t-Bid at an MOI of 100 were analyzed by Western blotting anti-Bid (“a”) allowing comparison of endogenous Bid levels (marked Z) to translocated YopE₁₋₁₃₈-Bid (marked X) or YopE₁₋₁₃₈-tBid (marked Y) levels. As a loading control western blotting anti-Actin was performed (shown in “b”). In some cases, cells were treated with VI: 0.5 μ M Staurosporine or VII: 1 μ M Staurosporine **(C)** HeLa cells were left untreated (I) or infected at an MOI of 100 for 1h with II: *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si2, III: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-Bid, IV: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-tBid. In some cases, cells were treated with V: 0.5 μ M Staurosporine or VI: 1 μ M Staurosporine. After fixation cells were stained for the actin cytoskeleton (gray).

Fig. 8: T3SS dependent delivery of zebrafish BIM induces apoptosis in zebrafish embryos. (A) 2 dpf zebrafish embryos were infected with the EGFP expressing *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si1 control strain (I) or zBIM translocating strain (II: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-zBIM) by injection of about 400 bacteria into the hindbrain region. After 5.5 h the embryos were fixed, stained for activated Caspase 3 (cleaved Caspase 3, p17; shown in “c”) and analyzed for presence of bacteria (EGFP signal, shown in “b”). Maximum intensity z projections are shown for fluorescent images. Bright-field z projection are shown in “a“ (B) Automated image analysis on maximum intensity z projections of recorded z-stack images of (A). Briefly, bacteria were detected via the GFP channel. Around each area of a bacterial spot a circle with a radius of 10 pixels was created. Overlapping regions were separated equally among the connecting members. In those areas closely surrounding bacteria, the Caspase 3 p17 staining intensity was measured and is plotted on the y-axis (as [a.u.]). Statistical analysis was performed using a Mann-Whitney test (***) indicates a p value < 0.001). Data were combined from n=14 for *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si1 control strain (I) or n=19 for II: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-zBIM infected animals, error bars indicated are standard error of the mean.

Fig. 9: tBid dependent phosphoproteome: HeLa cells were infected for 30 min with *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-t-Bid at an MOI of 100 and as a control with *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si2. (A) Graphical representation of the tBID phosphoproteome. Proteins containing phosphopeptides that were significantly regulated in a tBid dependent manner (gray) (q-value <0.01) as well as known apoptosis related proteins (dark gray) are represented in a STRING network of known and predicted protein-protein interactions (high-confidence, score 0.7). Only proteins with at least one connection in STRING are represented. (B) Confocal images of HeLa cells infected with either *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si2 (I) or *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-t-Bid (II) reveal the induction of an apoptotic phenotype upon tBid delivery. Cells were stained for the nuclei with Hoechst (“a“), for F-actin with phalloidin (“b“), for tubulin with an anti-tubulin antibody (“c“) and for mitochondria with mitotracker (“d“). Scale bar represents 40µm.

Fig. 10: Description of the type III secretion-based delivery toolbox. (A) Vector maps of the cloning plasmids pBad_Si1 and pBad_Si2 used to generate fusion constructs with YopE₁₋₁₃₈. The chaperone SycE and the YopE₁₋₁₃₈-fusion are under the native *Y. enterocolitica* promoter. The two plasmids only differ in presence of an arabinose inducible EGFP present on pBad_Si1 (B) Multiple cloning site directly following the yopE₁₋₁₃₈ fragment on pBad_Si1 and pBad_Si2 plasmids.

Fig. 11: Characterization of T3SS protein delivery into various cell lines. Anti-Myc immunofluorescence staining on Swiss 3T3 fibroblasts (“a”), Jurkat cells (“b”) and HUVEC cells (“c”) left untreated (II) or infected with *Y. enterocolitica* ΔHOPEMT *asd* + pBad_Si2 (I) at the MOI indicated above the images (MOI 25, 50, 100, 200 and 400 for HUVECs) for 1 h.

Fig. 12: T3SS dependency of delivery of bacterial effector proteins into eukaryotic cell. Digitonin lysed HeLa cells after infection at an MOI of 100 for time indicated above the blots (0, 5, 15, 10, 60 and 120 minutes) with *Y. enterocolitica* ΔHOPEMT *asd* Δ*yopB* + YopE₁₋₁₃₈-SopE-Myc (I) or *Y. enterocolitica* ΔHOPEMT *asd* + YopE₁₋₁₃₈-SopE-Myc (II) were analyzed by Western blotting anti-Myc. The size corresponding to YopE₁₋₁₃₈-SopE-Myc is marked with “a”, while the size of the endogenous c-Myc protein is marked with “b”.

Fig. 13 and 14: T3SS dependent secretion of various other proteins into the culture supernatant. In-vitro secretion experiment of I: *Y. enterocolitica* ΔHOPEMT *asd* + YopE₁₋₁₃₈ fused to the protein as indicated. Protein content of total bacterial lysates (“A”) and precipitated culture supernatants (“B”) was analyzed by Western blotting using an anti-YopE antibody. Numbers written indicate molecular weight in kDa at the corresponding height.

Figs. 15A to N: *Y. enterocolitica* and *S. enterica* strains used in this study.

List of *Y. enterocolitica* and *S. enterica* strains used in this study providing information on background strains, plasmids and proteins for T3SS dependent delivery encoded on corresponding plasmids. Further, information on oligonucleotides used for construction of the corresponding plasmid, the backbone plasmid and antibiotic resistances is provided.

Fig. 16: Delivery of murine tBid, murine Bid BH3 and murine Bax BH3 into B16F10 cells induces massive apoptosis. B16F10 cells uninfected (I) or after infection (MOI of 50) for 2.5h with *Y. enterocolitica* ΔHOPEMT *asd* and II: +

pBadSi_2, III: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine tBid, IV: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine Bid BH3 or V: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine Bax BH3. After fixation cells were stained for the actin cytoskeleton and nuclei (both in gray).

Fig. 17: Delivery of murine tBid, murine Bid BH3 and murine Bax BH3 into D2A1 cells induces massive apoptosis. D2A1 cells uninfected (I) or after infection (MOI of 50) for 2.5h with *Y. enterocolitica* ΔHOPEMT *asd* and II: + pBadSi_2, III: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine tBid, IV: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine Bid BH3 or V: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine Bax BH3. After fixation cells were stained for the actin cytoskeleton and nuclei (both in gray).

Fig. 18: Delivery of murine tBid, murine Bid BH3 and murine Bax BH3 into HeLa cells induces massive apoptosis. HeLa cells uninfected (I) or after infection (MOI of 50) for 2.5h with *Y. enterocolitica* ΔHOPEMT *asd* and II: + pBadSi_2, III: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine tBid, IV: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine Bid BH3 or V: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine Bax BH3. After fixation cells were stained for the actin cytoskeleton and nuclei (both in gray).

Fig. 19: Delivery of murine tBid, murine Bid BH3 and murine Bax BH3 into 4T1 cells induces massive apoptosis. 4T1 cells uninfected (I) or after infection (MOI of 50) for 2.5h with *Y. enterocolitica* ΔHOPEMT *asd* and II: + pBadSi_2, III: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine tBid, IV: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine Bid BH3 or V: + YopE₁₋₁₃₈-*Y. enterocolitica* codon optimized murine Bax BH3. After fixation cells were stained for the actin cytoskeleton and nuclei (both in gray).

Fig. 20: Delivery of murine tBid by *S. enterica* grown under SPI-1 T3SS inducing conditions into eukaryotic cells induces apoptosis. Cleaved Caspase 3 p17 western blot analysis on HeLa cells left untreated (I) or infected for 4h with III: *S. enterica* *aroA* carrying IV: SteA₁₋₂₀-t-Bid, V: SteA_{FL}-Bid, VI: SopE₁₋₈₁-t-Bid or VII: SopE₁₋₁₀₅-t-Bid at an MOI of 100. For this experiment, all *S. enterica* *aroA* strains were grown under SPI-1 T3SS inducing conditions. In some cases, cells were treated with II: 1 μM Staurosporine. Numbers written indicate molecular weight in kDa at the corresponding height.

Fig. 21: Delivery of murine tBid by *S. enterica* grown under SPI-2 T3SS inducing conditions into eukaryotic cells induces apoptosis. Cleaved Caspase 3 p17 western blot analysis on HeLa cells left untreated (I) or infected for 4h with III: *S. enterica aroA* carrying IV: SteA₁₋₂₀-t-Bid, V: SteA_{FL}-Bid, VI: SopE₁₋₈₁-t-Bid or VII: SopE₁₋₁₀₅-t-Bid at an MOI of 100. For this experiment, all *S. enterica aroA* strains were grown under SPI-2 T3SS inducing conditions. In some cases, cells were treated with II: 1 μ M Staurosporine. Numbers written indicate molecular weight in kDa at the corresponding height.

Fig. 22: *S. enterica* T3SS dependent secretion of various cell cycle proteins into the culture supernatant. In-vitro secretion experiment of *S. enterica aroA* + either SteA_{FL} (I, III, V, VII) or SopE₁₋₁₀₅ (II, IV, VI, VIII) fused to proteins as listed following. I and II: Ink4a-MycHis; III and IV: Ink4c-MycHis; V and VI: Mad2-MycHis; VII and VIII: Cdk1-MycHis. Protein content of precipitated culture supernatants (“A”) and total bacterial lysates (“B”) was analyzed by Western blotting using an anti-myc antibody. Numbers written indicate molecular weight in kDa at the corresponding height.

Fig. 23: T3SS dependent secretion of various known cell cycle interfering peptides into the culture supernatant. In-vitro secretion experiment of I: *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si2. II-VII: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈ fused to peptides as listed following: II: Ink4A₈₄₋₁₀₃; III: p107/RBL1₆₅₇₋₆₆₂; IV: p21₁₄₁₋₁₆₀D149A; V: p21₁₄₅₋₁₆₀D149A; VI: p21₁₇₋₃₃; VII: cyclin D2₁₃₉₋₁₄₇. Protein content of precipitated culture supernatants (“A”) and total bacterial lysates (“B”) was analyzed by Western blotting using an anti-YopE antibody. Numbers written indicate molecular weight in kDa at the corresponding height.

Fig. 24: Fusion of the T3SS delivered protein to Ubiquitin allows removal of the YopE₁₋₁₃₈ appendage. HeLa cells are infected with a strain delivering a protein of interest fused to YopE₁₋₁₃₈ with a directly fused Ubiquitin (YopE₁₋₁₃₈-Ubi). After protein delivery into the eukaryotic cell, endogenous Ubiquitin specific proteases will cleave the YopE₁₋₁₃₈-Ubi appendage from the protein of interest. Digitonin lysed HeLa cells uninfected (I) or after infection (MOI of 100) for 1h with II: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-Flag-INK4C-MycHis or III: + YopE₁₋₁₃₈-Flag-Ubiquitin-INK4C-MycHis were analyzed by Western blotting anti-INK4C for the presence of IV: YopE₁₋₁₃₈-Flag-Ubiquitin-INK4C-MycHis or V: YopE₁₋₁₃₈-Flag-

INK4C-MycHis, the cleaved form VI: INK4C-MycHis and VII: the endogenous INK4C.

Fig. 25: Schematic representation of heterologous proteins and domains thereof to be delivered via the bacteria T3SS. I: Human/ murine full-length protein with domains colored in various grayscale, II: Truncated human/ murine protein with domains colored in various grayscale, III: Motif/domain of human/ murine full-length protein only, IV: Motif/domain only repeated (left) or combination of two different motifs/domains of human/murine full-length protein (right).

Fig. 26: Delivery of BH3 domains of murine tBid and murine Bax into eukaryotic cells and fused repeats thereof induce apoptosis in cancerous cells.

B16F10 murine melanoma cells were infected at a MOI of 5, 10, 25 and 50 (left to right in each condition) of corresponding bacteria as indicated for 4h. Effect on cell viability was assessed by counting cell numbers via nuclear counting. I: nuclear count. II: *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si2, III: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-Bid-BH3, IV: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-(Bid-BH3)₂ and V: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-(Bid-BH3)-(Bax-BH3). Nuclei were stained with Hoechst. Images were acquired using an automated microscope and cell number was automatically determined using CellProfiler.

Fig. 27: Delivery of BH3 domains of murine tBid and murine Bax into eukaryotic cells and fused repeats thereof induce apoptosis in cancerous cells.

4T1 murine breast cancer cells were infected at a MOI of 5, 10, 25 and 50 (left to right in each condition) of corresponding bacteria as indicated for 4h. Effect on cell viability was assessed by counting cell numbers via nuclear counting. I: nuclear count. II: *Y. enterocolitica* Δ HOPEMT *asd* + pBad_Si2, III: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-Bid-BH3, IV: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-(Bid-BH3)₂ and V: *Y. enterocolitica* Δ HOPEMT *asd* + YopE₁₋₁₃₈-(Bid-BH3)-(Bax-BH3). Nuclei were stained with Hoechst. Images were acquired using an automated microscope and cell number was automatically determined using CellProfiler.

Fig. 28: Delivery of synthetic increased pro-apoptotic proteins. Delivery of single synthetic proteins consisting of single or tandem repeats of BH3 domains originating from pro-apoptotic proteins t-BID or BAX leads to enhanced apoptosis induction in

4T1 and B16F10 cancerous cells. 4T1 (I) or B16F10 (II) cells were infected with *Y. enterocolitica* $\Delta yopHOPEMT$ encoding on pBad-MycHisA IV: YopE₁₋₁₃₈-tBID BH3 extended domain, V: YopE₁₋₁₃₈-linker-tBID BH3, VI: YopE₁₋₁₃₈-tBID BH3, VII: YopE₁₋₁₃₈-(tBID BH3)₂, VIII: YopE₁₋₁₃₈-tBID BH3 - BAX BH3 or IX: YopE₁₋₁₃₈-BAX BH3 - tBID BH3. A titration of the bacteria added to the cells (MOI) was performed for each strain, cell counts determined and IC50 calculated using non-linear regression. IC50 MOI is indicated (III).

Fig. 29: Induction of apoptosis by pYV-encoded synthetic pro-apoptotic proteins.

Delivery of a single or a tandem repeat of BID BH3 domain encoded on the pYV leads to apoptosis induction in 4T1 and B16F10 cancerous cells. 4T1 (I) or B16F10 (II) cells were infected with *Y. enterocolitica* $\Delta HOPEMT$ + IV: pYV-YopE₁₋₁₃₈-BH3-Bid, or V: + pYV-YopE₁₋₁₃₈-(BH3-Bid)₂ or VI: with *Y. enterocolitica* $\Delta HOPEMT$ pBad-MycHisA-YopE₁₋₁₃₈-(BH3-Bid)₂ for 3 hours. A titration of the bacteria added to the cells (MOI) was performed for each strain, cell counts determined and IC50 (III) calculated using non-linear regression.

Fig. 30: Tumor colonization of i.v. injected *Y. enterocolitica* $\Delta yopH,O,P,E,M,T$ in the 4T1 breast cancer allograft model. Bacterial counts in tumors are indicated as colony forming units (CFU) per gram of tissue (III). Counts were assessed in tumors at day 8 (I) and 14 (II) post infection. Each dot represents an individual mouse. The horizontal dashed line indicates the detection limit.

Fig. 31: Biodistribution of i.v. injected *Y. enterocolitica* $\Delta yopH,O,P,E,M,T$ in the 4T1 breast cancer allograft model. Bacterial counts in blood (I), spleen (II), liver (III), lung (IV) and tumor (V) are indicated as colony forming units (CFU) per gram of tissue or per ml of blood (VI). Counts were assessed at day 14 post infection. Each dot represents an individual mouse. The horizontal dashed line indicates the detection limit. * indicates a mouse with large metastases found on lung.

Fig. 32: Delay of tumor progression in wildtype Balb/C mice allografted s.c. with 4T1 breast cancer cells. Wildtype Balb/C mice allografted s.c. with 4T1 breast cancer cells were i.v. injected with I: PBS or II: 1×10^7 *Y. enterocolitica* $\Delta HOPEMT$ + pYV-YopE₁₋₁₃₈-(BH3-Bid)₂, once the tumor had reached a size of 150-250 mm³. The day of the i.v. injection of bacteria was defined as day 0. Tumor volume was measured over the following days (III; day 0 to day 9 post i.v. injection of bacteria) with calipers. The relative tumor volume, normalized to the tumor volume at day 0, is indicated (IV) as mm³. The mean is indicated with symbols, error bars depicted show

the standard error of the mean. Statistical significance is measured with a 2way ANOVA, * indicates p value <0.05, ** a p value < 0.005.

Fig. 33: Tumor progression in wildtype Balb/C mice allografted s.c. with 4T1 breast cancer cells. Wildtype Balb/C mice allografted s.c. with 4T1 breast cancer cells were i.v. injected with I: PBS or II: 1×10^7 Y. enterocolitica dHOPEMT, once the tumor had reached a size of 150-250 mm³. The day of the i.v. injection of bacteria was defined as day 0. Tumor volume was measured over the following days (III; day 0 to day 9 post i.v. injection of bacteria) with calipers. The relative tumor volume, normalized to the tumor volume at day 0, is indicated (IV) as mm³. The mean is indicated with symbols, error bars depicted show the standard error of the mean.

Detailed description of the invention

The present invention provides recombinant Gram-negative bacterial strains and the use thereof for delivery of repeated domains of a heterologous protein or two or more domains of different heterologous proteins into eukaryotic cells.

For the purposes of interpreting this specification, the following definitions will apply and whenever appropriate, terms used in the singular will also include the plural and vice versa. It is to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

The term “Gram-negative bacterial strain“ as used herein includes the following bacteria: *Aeromonas salmonicida*, *Aeromonas hydrophila*, *Aeromonas veronii*, *Anaeromyxobacter dehalogenans*, *Bordetella bronchiseptica*, *Bordetella parapertussis*, *Bordetella pertussis*, *Bradyrhizobium japonicum*, *Burkholderia cenocepacia*, *Burkholderia cepacia*, *Burkholderia mallei*, *Burkholderia pseudomallei*, *Chlamydia muridarum*, *Chlamydia trachomatis*, *Chlamydophila abortus*, *Chlamydophila pneumoniae*, *Chromobacterium violaceum*, *Citrobacter rodentium*, *Desulfovibrio vulgaris*, *Edwardsiella tarda*, *Endozoicomonas elysicola*, *Erwinia amylovora*, *Escherichia albertii*, *Escherichia coli*, *Lawsonia intracellularis*, *Mesorhizobium loti*, *Myxococcus xanthus*, *Pantoea agglomerans*, *Photobacterium damsela*, *Photobacterium luminescens*, *Photobacterium temperate*, *Pseudoalteromonas spongiae*, *Pseudomonas aeruginosa*, *Pseudomonas plecoglossicida*, *Pseudomonas*

syringae, *Ralstonia solanacearum*, *Rhizobium* sp, *Salmonella enterica* and other *Salmonella* sp, *Shigella flexneri* and other *Shigella* sp, *Sodalis glossinidius*, *Vibrio alginolyticus*, *Vibrio azureus*, *Vibrio campellii*, *Vibrio caribbenthicus*, *Vibrio harvey*, *Vibrio parahaemolyticus*, *Vibrio tasmaniensis*, *Vibrio tubiashii*, *Xanthomonas axonopodis*, *Xanthomonas campestris*, *Xanthomonas oryzae*, *Yersinia enterocolitica*, *Yersinia pestis*, *Yersinia pseudotuberculosis*. Preferred Gram-negative bacterial strains of the invention are Gram-negative bacterial strains comprised by the family of *Enterobacteriaceae* and *Pseudomonadaceae*. The Gram-negative bacterial strain of the present invention is normally used for delivery of heterologous proteins by the bacterial T3SS into eukaryotic cells in vitro and/or in vivo, preferably in vivo.

The term "recombinant Gram-negative bacterial strain" used herein refers to a Gram-negative bacterial strain genetically transformed with a vector. A useful vector of the present invention is e.g. an expression vector, a vector for chromosomal or virulence plasmid insertion or a DNA or RNA fragment for chromosomal or virulence plasmid insertion or modification.

The term "recombinant Gram-negative bacterial strains which are deficient in producing at least one T3SS functional effector protein" used herein refers to a recombinant Gram-negative bacterial strain in which at least one T3SS effector protein is mutated such that the resulting recombinant Gram-negative bacterial strain no longer produces a functional form of at least one T3SS effector protein i.e. that the expression of such effector gene is abolished so that the resulting recombinant Gram-negative bacterial strains does not produce any of the at least one T3SS effector protein or that the catalytic activity of the encoded effector protein is abolished so that the at least one T3SS effector protein produced does not have its catalytic activity. e.g. does not exercise its effector functions. For the purpose of delivering proteins, the secretion and translocation system of the recombinant Gram-negative bacterial strains which are deficient in producing at least one T3SS functional effector protein needs to be intact. The term "T3SS effector protein" or "bacterial T3SS effector protein" as used herein refers to proteins which are naturally injected by T3S systems into the cytosol of eukaryotic cells and to proteins which are naturally secreted by T3S systems that might e.g. form the translocation pore into the eukaryotic membrane (including pore-forming translocators (as *Yersinia* YopB and YopD) and tip-proteins

like *Yersinia* LcrV). Preferably proteins which are naturally injected by T3S systems into the cytosol of eukaryotic cells are used. These virulence factors will paralyze or reprogram the eukaryotic cell to the benefit of the pathogen. T3S effectors display a large repertoire of biochemical activities and modulate the function of crucial host regulatory molecules and include AvrA, AvrB, AvrBs2, AvrBS3, AvrBsT, AvrD, AvrD1, AvrPphB, AvrPphC, AvrPphEpto, AvrPpiBPto, AvrPto, AvrPtoB, AvrRpm1, AvrRpt2, AvrXv3, CigR, EspF, EspG, EspH, EspZ, ExoS, ExoT, GogB, GtgA, GtgE, GALA family of proteins, HopAB2, HopAO1, HopI1, HopM1, HopN1, HopPtoD2, HopPtoE, HopPtoF, HopPtoN, HopU1, HsvB, IcsB, IpaA, IpaB, IpaC, IpaH, IpaH7.8, IpaH9.8, IpgB1, IpgB2, IpgD, LcrV, Map, OspC1, OspE2, OspF, OspG, OspI, PipB, PipB2, PopB, PopP2, PthXo1, PthXo6, PthXo7, SifA, SifB, SipA/SspA, SipB, SipC/SspC, SipD/SspD, SlrP, SopA, SopB/SigD, SopD, SopE, SopE2, SpiC/SsaB, SptP, SpvB, SpvC, SrfH, SrfJ, Sse, SseB, SseC, SseD, SseF, SseG, SseI/SrfH, SseJ, SseK1, SseK2, SseK3, SseL, SspH1, SspH2, SteA, SteB, SteC, SteD, SteE, TccP2, Tir, VirA, VirPphA, VopF, XopD, YopB, YopD YopE, YopH, YopJ, YopM, YopO, YopP, YopT, YpkA.

The terms “Gram-negative bacterial strain deficient to produce an amino acid essential for growth” and “auxotroph mutant” are used herein interchangeably and refer to Gram-negative bacterial strains which can not grow in the absence of at least one exogenously provided essential amino acid or a precursor thereof. The amino acid the strain is deficient to produce is e.g. aspartate, *meso*-2,6-diaminopimelic acid, aromatic amino acids or leucine-arginine [8]. Such a strain can be generated by e.g. deletion of the aspartate-beta-semialdehyde dehydrogenase gene (Δ asd). Such an auxotroph mutant cannot grow in absence of exogenous *meso*-2,6-diaminopimelic acid [9]. The mutation, e.g. deletion of the aspartate-beta-semialdehyde dehydrogenase gene is preferred herein for a Gram-negative bacterial strain deficient to produce an amino acid essential for growth of the present invention.

The term “Gram-negative bacterial strain deficient to produce adhesion proteins binding to the eukaryotic cell surface or extracellular matrix” refers to mutant Gram-negative bacterial strains which do not express at least one adhesion protein compared to the adhesion proteins expressed by the corresponding wild type strain. Adhesion proteins may include e.g. extended polymeric adhesion molecules like pili/fimbriae or

non-fimbrial adhesins. Fimbrial adhesins include type-1 pili (such as *E. coli* Fim-pili with the FimH adhesin), P-pili (such as Pap-pili with the PapG adhesin from *E. coli*), type 4 pili (as pilin protein from e.g. *P. aeruginosa*) or curli (Csg proteins with the CsgA adhesin from *S. enterica*). Non-fimbrial adhesions include trimeric autotransporter adhesins such as YadA from *Y. enterocolitica*, BpaA (*B. pseudomallei*), Hia (*H. influenzae*), BadA (*B. henselae*), NadA (*N. meningitidis*) or UspA1 (*M. catarrhalis*) as well as other autotransporter adhesins such as AIDA-1 (*E. coli*) as well as other adhesins/invasins such as InvA from *Y. enterocolitica* or Intimin (*E. coli*) or members of the Dr-family or Afa-family (*E. coli*). The terms YadA and InvA as used herein refer to proteins from *Y. enterocolitica*. The autotransporter YadA [10,11] binds to different forms of collagen as well as fibronectin, while the invasin InvA [12-14] binds to β -integrins in the eukaryotic cell membrane. If the Gram-negative bacterial strain is a *Y. enterocolitica* strain the strain is preferably deficient in InvA and/or YadA.

As used herein, the term “family of *Enterobacteriaceae*” comprises a family of gram-negative, rod-shaped, facultatively anaerobic bacteria found in soil, water, plants, and animals, which frequently occur as pathogens in vertebrates. The bacteria of this family share a similar physiology and demonstrate a conservation within functional elements and genes of the respective genomes. As well as being oxidase negative, all members of this family are glucose fermenters and most are nitrate reducers.

Enterobacteriaceae bacteria of the invention may be any bacteria from that family, and specifically includes, but is not limited to, bacteria of the following genera:

Escherichia, *Shigella*, *Edwardsiella*, *Salmonella*, *Citrobacter*, *Klebsiella*, *Enterobacter*, *Serratia*, *Proteus*, *Erwinia*, *Morganella*, *Providencia*, or *Yersinia*. In more specific embodiments, the bacterium is of the *Escherichia coli*, *Escherichia blattae*, *Escherichia fergusonii*, *Escherichia hermannii*, *Escherichia vulneris*, *Salmonella enterica*, *Salmonella bongori*, *Shigella dysenteriae*, *Shigella flexneri*, *Shigella boydii*, *Shigella sonnei*, *Enterobacter aerogenes*, *Enterobacter gergoviae*, *Enterobacter sakazakii*, *Enterobacter cloacae*, *Enterobacter agglomerans*, *Klebsiella pneumoniae*, *Klebsiella oxytoca*, *Serratia marcescens*, *Yersinia pseudotuberculosis*, *Yersinia pestis*, *Yersinia enterocolitica*, *Erwinia amylovora*, *Proteus mirabilis*, *Proteus vulgaris*, *Proteus penneri*, *Proteus hauseri*, *Providencia alcalifaciens*, or

Morganella morganii species.

Preferably the Gram-negative bacterial strain is selected from the group consisting of the genera *Yersinia*, *Escherichia*, *Salmonella*, *Shigella*, *Pseudomonas*, *Chlamydia*, *Erwinia*, *Pantoea*, *Vibrio*, *Burkholderia*, *Ralstonia*, *Xanthomonas*, *Chromobacterium*, *Sodalis*, *Citrobacter*, *Edwardsiella*, *Rhizobiae*, *Aeromonas*, *Photorhabdus*, *Bordetella* and *Desulfovibrio*, more preferably from the group consisting of the genera *Yersinia*, *Escherichia*, *Salmonella*, and *Pseudomonas*, most preferably from the group consisting of the genera *Yersinia* and *Salmonella*.

The term "*Yersinia*" as used herein includes all species of *Yersinia*, including *Yersinia enterocolitica*, *Yersinia pseudotuberculosis* and *Yersinia pestis*. Preferred is *Yersinia enterocolitica*.

The term "*Salmonella*" as used herein includes all species of *Salmonella*, including *Salmonella enterica* and *S. bongori*. Preferred is *Salmonella enterica*.

"Promoter" as used herein refers to a nucleic acid sequence that regulates expression of a transcriptional unit. A "promoter region" is a regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. Within the promoter region will be found a transcription initiation site (conveniently defined by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase such as the putative -35 region and the Pribnow box. The term "operably linked" when describing the relationship between two DNA regions simply means that they are functionally related to each other and they are located on the same nucleic acid fragment. A promoter is operably linked to a structural gene if it controls the transcription of the gene and it is located on the same nucleic acid fragment as the gene. Usually the promoter is functional in said Gram-negative bacterial strain, i.e. the promoter is capable of expressing the fusion protein of the present invention, i.e. the promoter is capable of expressing the fusion protein of the present invention without further genetic engineering or expression of further proteins. Furthermore, a functional promoter must not be naturally counter-regulated to the bacterial T3SS.

The term "delivery" used herein refers to the transportation of a protein from a

recombinant Gram-negative bacterial strain to a eukaryotic cell, including the steps of expressing the heterologous protein in the recombinant Gram-negative bacterial strain, secreting the expressed protein(s) from such Gram-negative bacterial strain and translocating the secreted protein(s) by such Gram-negative bacterial strain into the cytosol of the eukaryotic cell. Accordingly, the terms "delivery signal" or "secretion signal" which are used interchangeably herein refer to a polypeptide sequence which can be recognized by the secretion and translocation system of the Gram-negative bacterial strain and directs the delivery of a protein from the Gram-negative bacterial strain to eukaryotic cells.

The term "delivery signal from a bacterial effector protein" used herein refers to a delivery signal from a bacterial effector protein functional in the recombinant Gram-negative bacterial strain, i.e. which allows an expressed heterologous protein in the recombinant Gram-negative bacterial strain to be secreted from such recombinant Gram-negative bacterial strain by a secretion system such as e.g. the type III secretion system or to be translocated by such recombinant Gram-negative bacterial strain into the cytosol of a eukaryotic cell by a secretion system such as e.g. the type III secretion system. The term "delivery signal from a bacterial effector protein" used herein also comprises a fragment of a delivery signal from a bacterial effector protein i.e. shorter versions of a delivery signal e.g. a delivery signal comprising up to 10, preferably up to 20, more preferably up to 50, even more preferably up to 100, in particular up to 140 amino acids of a delivery signal e.g. of a naturally occurring delivery signal. Thus a nucleotide sequence such as e.g. a DNA sequence encoding a delivery signal from a bacterial effector protein may encode a full length delivery signal or a fragment thereof wherein the fragment usually comprises usually up to 30, preferably up to 60, more preferably up to 150, even more preferably up to 300, in particular up to 420 nucleic acids.

As used herein, the "secretion" of a protein refers to the transportation of a heterologous protein outward across the cell membrane of a recombinant Gram-negative bacterial strain. The "translocation" of a protein refers to the transportation of a heterologous protein from a recombinant Gram-negative bacterial strain across the plasma membrane of a eukaryotic cell into the cytosol of such eukaryotic cell.

The term "eukaryotic cells" as used herein includes e.g. the following eukaryotic cells: Hi-5, HeLa, Hek, HUVECs, 3T3, CHO, Jurkat, Sf-9, HepG2, Vero, MDCK, Mefs, THP-1, J774, RAW, Caco2, NCI60, DU145, Lncap, MCF-7, MDA-MB-438, PC3, T47D, A549, U87, SHSY5Y, Ea.Hy926, Saos-2, 4T1, D2A1, B16F10, and primary human hepatocytes. "Eukaryotic cells" as used herein, are also referred to as "target cells" or "target eukaryotic cells".

The term "T3SS effector protein" as used herein refers to proteins which are naturally injected by T3S systems into the cytosol of eukaryotic cells and to proteins which are naturally secreted by T3S systems that might e.g. form the translocation pore into the eukaryotic membrane (including pore-forming translocators (as *Yersinia* YopB and YopD) and tip-proteins like *Yersinia* LcrV). Preferably proteins which are naturally injected by T3S systems into the cytosol of eukaryotic cells are used. These virulence factors will paralyze or reprogram the eukaryotic cell to the benefit of the pathogen. T3S effectors display a large repertoire of biochemical activities and modulate the function of crucial host regulatory molecules [2,15] and include AvrA, AvrB, AvrBs2, AvrBS3, AvrBsT, AvrD, AvrD1, AvrPphB, AvrPphC, AvrPphE_{Pto}, AvrPpiB_{Pto}, AvrPto, AvrPtoB, AvrRpm1, AvrRpt2, AvrXv3, CigR, EspF, EspG, EspH, EspZ, ExoS, ExoT, GogB, GtgA, GtgE, GALA family of proteins, HopAB2, HopAO1, HopI1, HopM1, HopN1, HopPtoD2, HopPtoE, HopPtoF, HopPtoN, HopU1, HsvB, IcsB, IpaA, IpaB, IpaC, IpaH, IpaH7.8, IpaH9.8, IpgB1, IpgB2, IpgD, LcrV, Map, OspC1, OspE2, OspF, OspG, OspI, PipB, PipB2, PopB, PopP2, PthXo1, PthXo6, PthXo7, SifA, SifB, SipA/SspA, SipB, SipC/SspC, SipD/SspD, SlrP, SopA, SopB/SigD, SopD, SopE, SopE2, SpiC/SsaB, SptP, SpvB, SpvC, SrfH, SrfJ, Sse, SseB, SseC, SseD, SseF, SseG, SseI/SrfH, SseJ, SseK1, SseK2, SseK3, SseL, SspH1, SspH2, SteA, SteB, SteC, SteD, SteE, TccP2, Tir, VirA, VirPphA, VopF, XopD, YopB, YopD, YopE, YopH, YopJ, YopM, YopO, YopP, YopT, YpkA.

T3SS effector genes of *Yersinia* have been cloned from e.g. *Y. enterocolitica* which are YopE, YopH, YopM, YopO, YopP/YopJ, and YopT [16]. The respective effector genes can be cloned from *Shigella flexneri* (e.g. OspF, IpgD, IpgB1), *Salmonella enterica* (e.g. SopE, SopB, SptP), *P. aeruginosa* (e.g. ExoS, ExoT, ExoU, ExoY) or *E. coli* (e.g. Tir, Map, EspF, EspG, EspH, EspZ). The nucleic acid sequences of these genes are available to those skilled in the art, e.g., in the Genebank Database (yopH,

yopO, yopE, yopP, yopM, yopT from NC_002120 GI:10955536; *S. flexneri* effector proteins from AF386526.1 GI:18462515; *S. enterica* effectors from NC_016810.1 GI:378697983 or FQ312003.1 GI:301156631; *P. aeruginosa* effectors from AE004091.2 GI:110227054 or CP000438.1 GI:115583796 and *E. coli* effector proteins from NC_011601.1 GI:215485161).

For the purpose of the present invention, genes are denoted by letters of lower case and italicised to be distinguished from proteins. In case the genes (denoted by letters of lower case and italicised) are following a bacterial species name (like *E. coli*), they refer to a mutation of the corresponding gene in the corresponding bacterial species. For example, YopE refers to the effector protein encoded by the *yopE* gene. *Y. enterocolitica yopE* represents a *Y. enterocolitica* having a mutation in the *yopE* gene.

As used herein, the terms “polypeptide”, “peptide”, “protein”, “polypeptidic” and “peptidic” are used interchangeably to designate a series of amino acid residues connected to each other by peptide bonds between the alpha-amino and carboxy groups of adjacent residues. Preferred are proteins which have an amino acid sequence comprising at least 10 amino acids, more preferably at least 20 amino acids.

According to the present invention, “a domain of a heterologous protein” includes domains of naturally occurring proteins and also includes domains of artificially engineered proteins. As used herein, the term “domain of a heterologous protein” refers to a domain of a heterologous protein other than a domain of a T3SS effector protein or a domain other than a domain comprising the N-terminal fragment thereof to which it can be fused to achieve a fusion protein. In particular the domain of a heterologous protein as used herein refers to a domain of a heterologous protein, which do not belong to the proteome, i.e. the entire natural protein complement of the specific recombinant Gram-negative bacterial strain provided and used by the invention, e.g. which do not belong to the proteome, i.e. the entire natural protein complement of a specific bacterial strain of the genera *Yersinia*, *Escherichia*, *Salmonella* or *Pseudomonas*. Usually the domain of the heterologous protein is of animal origin including human origin. Preferably the domain of the heterologous protein is a domain of a human protein. More preferably the domain of the heterologous protein is a domain of a protein selected from the group consisting of

proteins involved in apoptosis or apoptosis regulation, cell cycle regulators, ankyrin repeat proteins, cell signaling proteins, reporter proteins, transcription factors, proteases, small GTPases, GPCR related proteins, nanobody fusion constructs and nanobodies, bacterial T3SS effectors, bacterial T4SS effectors and viral proteins. Particular preferably the domain of the heterologous protein is a domain of a protein selected from the group consisting of proteins involved in apoptosis or apoptosis regulation, cell cycle regulators, ankyrin repeat proteins, reporter proteins, small GTPases, GPCR related proteins, nanobody fusion constructs, bacterial T3SS effectors, bacterial T4SS effectors and viral proteins. Even more particular preferred are domains of heterologous proteins selected from the group consisting of proteins involved in apoptosis or apoptosis regulation, cell cycle regulators, and ankyrin repeat proteins. Most preferred are domains of proteins involved in apoptosis or apoptosis regulation, like animal proteins involved in apoptosis or apoptosis regulation, preferably domains of human heterologous proteins involved in apoptosis or apoptosis regulation.

The term "repeated domains of a heterologous protein" as used herein refers to a fusion protein consisting of several repetitions of a domain of a heterologous protein, where these domains might either be directly fused to each other or where a variable linker e.g. a linker between 1 and 30, preferably between 2 and 15, more preferably between 3 and 10 amino acids might be introduced in between the domains. Preferably repeated identical domains or repeated domains which have an amino acid sequence identity of more than 80 %, usually more than 85 %, preferably more than 90 %, even more preferably more than 95 %, in particular more than 96%, more particular more than 97%, even more particular more than 98 %, most particular more than 99% are used. Also preferred are identical domains which have an amino acid identity of 100 %. Preferably two repeated domains, more preferably two repeated identical domains or two repeated domains having an amino acid sequence identity of more than 90 %, preferably more than 95% most preferably 100 % are comprised by the fusion protein as referred herein. More than two, e.g. three, four, five or six repeated domains are also contemplated by the present invention.

The term "two or more domains of different heterologous proteins" as used herein refers to a fusion protein consisting of one or several repetitions of at least two

domains of different heterologous proteins e.g. at least two domains of heterologous proteins having an amino acid sequence identity of 80% or less, where these different domains might either be directly fused to each other or where a variable linker e.g. a linker between 1 and 30, preferably between 2 and 15, more preferably between 3 and 10 amino acids might be introduced in between the domains.

Preferably two domains of different heterologous proteins are comprised by the fusion protein as referred herein. More than two, e.g. three, four, five or six domains of different heterologous proteins are also contemplated by the present invention.

The term “heterologous proteins which belong to the same functional class of proteins” as used herein refers to heterologous proteins which have the same function e.g. heterologous proteins having enzymatic activity, heterologous proteins which act in the same pathway such as e.g. cell cycle regulation, or share a common specific feature as e.g. belonging to the same class of bacterial effector proteins. Functional classes of proteins are e.g. proteins involved in apoptosis or apoptosis regulation, proteins which act as cell cycle regulators, ankyrin repeat proteins, cell signaling proteins, reporter proteins, transcription factors, proteases, small GTPases, GPCR related proteins, nanobody fusion constructs and nanobodies, bacterial T3SS effectors, bacterial T4SS effectors or viral proteins which act jointly in the biological process of establishing virulence to eukaryotic cells.

The domain of a heterologous protein expressed by the recombinant Gram-negative bacterial strain has usually a molecular weight of between 1-50 kDa, preferably between 1-30 kDa, more preferably between 1-20 kDa, most preferably between 1-10 kDa.

According to the present invention " proteins involved in apoptosis or apoptosis regulation" or “human heterologous proteins involved in apoptosis or apoptosis regulation” include, but are not limited to, Bad, Bcl2, Bak, Bmt, Bax, Puma, Noxa, Bim, Bcl-xL, Apaf1, Caspase 9, Caspase 3, Caspase 6, Caspase 7, Caspase 10, DFFA, DFFB, ROCK1, APP, CAD, ICAD, CAD, EndoG, AIF, HtrA2, Smac/Diablo, Arts, ATM, ATR, Bok/Mtd, Bmf, Mcl-1(S), IAP family, LC8, PP2B, 14-3-3 proteins, PKA, PKC, PI3K, Erk1/2, p90RSK, TRAF2, TRADD, FADD, Daxx, Caspase8, Caspase2, RIP, RAIDD, MKK7, JNK, FLIPs, FKHR, GSK3, CDKs and their

inhibitors like the INK4-family (p16(Ink4a), p15(Ink4b), p18(Ink4c), p19(Ink4d)), and the Cip1/Waf1/Kip1-2-family (p21(Cip1/Waf1), p27(Kip1), p57(Kip2)). Preferably Bad, Bmt, Bcl2, Bak, Bax, Puma, Noxa, Bim, Bcl-xL, Caspase9, Caspase3, Caspase6, Caspase7, Smac/Diablo, Bok/Mtd, Bmf, Mcl-1(S), LC8, PP2B, TRADD, Daxx, Caspase8, Caspase2, RIP, RAIDD, FKHR, CDKs and their inhibitors like the INK4-family (p16(Ink4a), p15(Ink4b), p18(Ink4c), p19(Ink4d)), most preferably BIM, Bid, truncated Bid, FADD, Caspase 3 (and subunits thereof), Bax, Bad, Akt, CDKs and their inhibitors like the INK4-family (p16(Ink4a), p15(Ink4b), p18(Ink4c), p19(Ink4d)) are used [17-19]. Additionally proteins involved in apoptosis or apoptosis regulation include DIVA, Bcl-Xs, Nbk/Bik, Hrk/Dp5, Bid and tBid, Egl-1, Bcl-Gs, Cytochrome C, Beclin, CED-13, BNIP1, BNIP3, Bcl-B, Bcl-W, Ced-9, A1, NR13, Bfl-1, Caspase 1, Caspase 2, Caspase 4, Caspase 5, Caspase 8.

Proteins involved in apoptosis or apoptosis regulation are selected from the group consisting of pro-apoptotic proteins, anti-apoptotic proteins, inhibitors of apoptosis-prevention pathways and inhibitors of pro-survival signalling or pathways. Pro-apoptotic proteins comprise proteins selected from the group consisting of Bax, Bak, Diva, Bcl-Xs, Nbk/Bik, Hrk/Dp5, Bmf, Noxa, Puma, Bim, Bad, Bid and tBid, Bok, Apaf1, Smac/Diablo, BNIP1, BNIP3, Bcl-Gs, Beclin 1, Egl-1 and CED-13, Cytochrome C, FADD, the Caspase family, and CDKs and their inhibitors like the INK4-family (p16(Ink4a), p15(Ink4b), p18(Ink4c), p19(Ink4d)) or selected from the group consisting of Bax, Bak, Diva, Bcl-Xs, Nbk/Bik, Hrk/Dp5, Bmf, Noxa, Puma, Bim, Bad, Bid and tBid, Bok, Egl-1, Apaf1, Smac/Diablo, BNIP1, BNIP3, Bcl-Gs, Beclin 1, Egl-1 and CED-13, Cytochrome C, FADD, and the Caspase family.

Preferred are Bax, Bak, Diva, Bcl-Xs, Nbk/Bik, Hrk/Dp5, Bmf, Noxa, Puma, Bim, Bad, Bid and tBid, Bok, Egl-1, Apaf1, BNIP1, BNIP3, Bcl-Gs, Beclin 1, Egl-1 and CED-13, Smac/Diablo, FADD, the Caspase family, CDKs and their inhibitors like the INK4-family (p16(Ink4a), p15(Ink4b), p18(Ink4c), p19(Ink4d)). Equally preferred are Bax, Bak, Diva, Bcl-Xs, Nbk/Bik, Hrk/Dp5, Bmf, Noxa, Puma, Bim, Bad, Bid and tBid, Bok, Apaf1, BNIP1, BNIP3, Bcl-Gs, Beclin 1, Egl-1 and CED-13, Smac/Diablo, FADD, the Caspase family.

Anti-apoptotic proteins comprise proteins selected from the group consisting of Bcl-2, Bcl-Xl, Bcl-B, Bcl-W, Mcl-1, Ced-9, A1, NR13, IAP family and Bfl-1. Preferred are Bcl-2, Bcl-Xl, Bcl-B, Bcl-W, Mcl-1, Ced-9, A1, NR13 and Bfl-1.

Inhibitors of apoptosis-prevention pathways comprise proteins selected from the group consisting of Bad, Noxa and Cdc25A. Preferred are Bad and Noxa.

Inhibitors of pro-survival signalling or pathways comprise proteins selected from the group consisting of PTEN, ROCK, PP2A, PHLPP, JNK, p38. Preferred are PTEN, ROCK, PP2A and PHLPP.

In some embodiments the heterologous proteins involved in apoptosis or apoptosis regulation are selected from the group consisting of BH3-only proteins, caspases and intracellular signalling proteins of death receptor control of apoptosis.

BH3-only proteins comprise proteins selected from the group consisting of Bad, BIM, Bid and tBid, Puma, Bik/Nbk, Bod, Hrk/Dp5, BNIP1, BNIP3, Bmf, Noxa, Mcl-1, Bcl-Gs, Beclin 1, Egl-1 and CED-13. Preferred are Bad, BIM, Bid and tBid.

Caspases comprise proteins selected from the group consisting of Caspase 1, Caspase 2, Caspase 3, Caspase 4, Caspase 5, Caspase 6, Caspase 7, Caspase 8, Caspase 9, Caspase 10. Preferred are Caspase 3, Caspase 8 and Caspase 9.

Intracellular signalling proteins of death receptor control of apoptosis comprise proteins selected from the group consisting of FADD, TRADD, ASC, BAP31, GULP1/CED-6, CIDEA, MFG-E8, CIDEA, RIPK1/RIP1, CRADD, RIPK3/RIP3, Crk, SHB, CrkL, DAXX, the 14-3-3 family, FLIP, DFF40 and 45, PEA-15, SODD. Preferred are FADD and TRADD.

In some embodiments two domains of a heterologous proteins involved in apoptosis or apoptosis regulation are comprised by the Gram-negative bacterial strain and/or the vector of the present invention, preferably two repeated, more preferably two identical repeated domains of a protein involved in apoptosis or apoptosis regulation or two domains of different proteins involved in apoptosis or apoptosis regulation. In some embodiments two domains of a heterologous proteins involved in apoptosis or apoptosis regulation are comprised by the Gram-negative bacterial strain and/or the vector of the present invention, wherein one is a domain of a pro-apoptotic protein and the other is a domain of a protein which is an inhibitor of apoptosis-prevention pathways or wherein one is a domain of a pro-apoptotic protein and the other domain is a domain of a protein which is an inhibitor of pro-survival signalling or pathways.

Pro-apoptotic proteins encompassed by the present invention have usually an alpha helical structure, preferably a hydrophobic helix surrounded by amphipathic helices and usually comprise at least one of BH1, BH2, BH3 or BH4 domains, preferably comprise at least one BH3 domain. Usually pro-apoptotic proteins encompassed by the present invention have no enzymatic activity.

Anti-apoptotic proteins encompassed by the present invention have usually an alpha helical structure, preferably a hydrophobic helix surrounded by amphipathic helices and comprises a combination of different BH1, BH2, BH3 and BH4 domains, preferably a combination of different BH1, BH2, BH3 and BH4 domains wherein a BH1 and a BH2 domain is present, more preferably BH4-BH3-BH1-BH2, BH1-BH2, BH4-BH1-BH2 or BH3-BH1-BH2 (from N- to the C-terminus). Additionally, proteins containing at least one BIR domain are also encompassed.

Inhibitors of apoptosis-prevention pathways encompassed by the present invention have usually an alpha helical structure, preferably a hydrophobic helix surrounded by amphipathic helices and usually comprise one BH3 domain.

BH1, BH2, BH3 or BH4 domains are each usually between about 5 to about 50 amino acids in length. Thus in some embodiments the domains of heterologous proteins are selected from the group consisting of domains of heterologous proteins which are about 5 to about 200, preferably about 5 to about 150, more preferably about 5 to about 100, most preferably about 5 to about 50, in particular about 5 to about 25 amino acids in length.

A particular preferred domain is the BH3 domain of apoptosis inducer tBID, more particular the BH3 domain comprising a sequence selected from the group consisting of SEQ ID NOs: 209, 210, 211 and 212, preferably SEQ ID NO: 211 or SEQ ID NO: 212. Equally preferred is the BH3 domain of apoptosis regulator BAX, more particular the BAX domain comprising a sequence selected from the group consisting of SEQ ID NOs: 213, 214, 215 and 216, preferably SEQ ID NO: 215 or SEQ ID NO: 216. The human and murine sequences are given in SEQ ID NOs 209-216, but tBID and BAX BH3 domains of all other species are equally included.

In some embodiments the repeated domains of the heterologous proteins are the BH3 domain, in particular repeated BH3 domains of apoptosis inducer tBID, more particular two repeated BH3 domains of apoptosis inducer tBID, most particular two repeated BH3 domains of apoptosis inducer tBID comprised by the sequence of SEQ

ID NO: 202. Thus in a preferred embodiment the vector of the Gram-negative bacterial strain of the present invention comprises a second DNA sequence encoding two repeated domains of a BH3 domain, more preferably two repeated BH3 domains of apoptosis inducer tBID. The two repeated domains are preferably connected by a linker of 1-30 amino acid length, preferably 2-15 amino acids, more preferred 3-10 amino acids long.

In some embodiments the two or more domains of different heterologous proteins are domains of heterologous proteins which belong to the same functional class of proteins, preferably the different heterologous proteins of the two or more domains are different heterologous proteins from the class of proteins involved in apoptosis or apoptosis regulation. In a preferred embodiment the two or more domains of different heterologous proteins are the BH3 domain of apoptosis inducer tBID and the BH3 domain of apoptosis regulator BAX, in particular the fused BH3 domains comprised by the sequence of SEQ ID NO: 203. The two domains of different heterologous proteins are preferably connected by a linker of 1-30 amino acid length, preferably 2-15 amino acids, more preferred 3-10 amino acids long.

In some embodiments the heterologous proteins is a pro-drug converting enzyme. In these embodiments the recombinant virulence attenuated Gram-negative bacterial strain expresses, preferably expresses and secretes a pro-drug converting enzyme. A prodrug converting enzyme as referred herein comprises enzymes converting non-toxic prodrugs into a toxic drug, preferably enzymes selected from the group consisting of cytosine deaminase, purine nucleoside phosphorylase, thymidine kinase, beta-galactosidase, carboxylesterases, nitroreductase, carboxypeptidases and beta-glucuronidases., more preferably enzymes selected from the group consisting of cytosine deaminase, purine nucleoside phosphorylase, thymidine kinase, and beta-galactosidase.

The term "protease cleavage site" as used herein refers to a specific amino acid motif within an amino acid sequence e.g. within an amino acid sequence of a protein or a fusion protein, which is cleaved by a specific protease, which recognizes the amino acid motif. For review see [20]. Examples of protease cleavage sites are amino acid motifs, which are cleaved by a protease selected from the group consisting of

enterokinase (light chain), enteropeptidase, prescission protease, human rhinovirus protease (HRV 3C), TEV protease, TVMV protease, FactorXa protease and thrombin.

The following amino acid motif is recognized by the respective protease:

- Asp-Asp-Asp-Asp-Lys: Enterokinase (light chain) / Enteropeptidase
- Leu-Glu-Val-Leu-Phe-Gln/Gly-Pro: PreScission Protease/human Rhinovirus protease (HRV 3C)
- Glu-Asn-Leu-Tyr-Phe-Gln-Ser and modified motifs based on the Glu-X-X-Tyr-X-Gln-Gly/Ser (where X is any amino acid) recognized by TEV protease (tobacco etch virus)
- Glu-Thr-Val-Arg-Phe-Gln-Ser: TVMV protease
- Ile-(Glu or Asp)-Gly-Arg: FactorXa protease
- Leu-Val-Pro-Arg/Gly-Ser: Thrombin.

Encompassed by the protease cleavage sites as used herein is ubiquitin. Thus in some preferred embodiments ubiquitin is used as protease cleavage site, i.e. the third DNA sequence encodes ubiquitin as protease cleavage site, which can be cleaved by a specific ubiquitin processing proteases at the N-terminal site, e.g. which can be cleaved by a specific ubiquitin processing proteases called Deubiquitinating enzymes at the N-terminal site endogeneously in the cell where the fusion protein has been delivered to. Ubiquitin is processed at its C-terminus by a group of endogenous Ubiquitin-specific C-terminal proteases (Deubiquitinating enzymes, DUBs). The cleavage of Ubiquitin by DUBs is supposed to happen at the very C-terminus of Ubiquitin (after G76).

An "individual," "subject" or "patient" is a vertebrate. In certain embodiments, the vertebrate is a mammal. Mammals include, but are not limited to, primates (including human and non-human primates) and rodents (e.g., mice and rats). In certain embodiments, a mammal is a human.

The term "mutation" is used herein as a general term and includes changes of both single base pair and multiple base pairs. Such mutations may include substitutions, frame-shift mutations, deletions, insertions and truncations.

The term "labelling molecule or an acceptor site for a labelling molecule" as used herein refers to a small chemical compound binding to a specific amino acid sequence

resulting in fluorescence of the bound chemical compound, preferably coumarine ligase/coumarine acceptor site (and derivates thereof), resorufin ligase/ resorufin acceptor site (and derivates thereof) and the tetra-Cysteine motif (as Cys-Cys-Pro-Gly-Cys-Cys and derivates thereof) in use with FlAsH/ReAsH dye (life technologies) or a fluorescent protein as Enhanced Green Fluorescent Protein (EGFP).

The term “nuclear localization signal” as used herein refers to an amino acid sequence that marks a protein for import into the nucleus of a eukaryotic cell and includes preferably a viral nuclear localization signal such as the SV40 large T-antigen derived NLS (PPKKKRKV).

The term “multiple cloning site” as used herein refers to a short DNA sequence containing several restriction sites for cleavage by restriction endonucleases such as AclI, HindIII, SspI, MluCI, Tsp509I, PciI, AgeI, BspMI, BfuAI, SexAI, MluI, BceAI, HpyCH4IV, HpyCH4III, BaeI, BsaXI, AflIII, SpeI, BsrI, BmrI, BglII, AfeI, AluI, StuI, ScaI, ClaI, BspDI, PI-SceI, NsiI, AseI, SwaI, CspCI, MfeI, BssSI, BmgBI, PmlI, DraIII, AleI, EcoP15I, PvuII, AlwNI, BtsIMutI, TspRI, NdeI, NlaIII, CviAII, FatI, MslI, FspEI, XcmI, BstXI, PflMI, BccI, NcoI, BseYI, FauI, SmaI, XmaI, TspMI, Nt.CviPII, LpnPI, AciI, SacII, BsrBI, MspI, HpaII, ScrFI, BssKI, StyD4I, BsaJI, BslI, BtgI, NciI, AvrII, MnlI, BbvCI, Nb.BbvCI, Nt.BbvCI, SbfI, Bpu10I, Bsu36I, EcoNI, HpyAV, BstNI, PspGI, StyI, BcgI, PvuI, BstUI, EagI, RsrII, BsiEI, BsiWI, BsmBI, Hpy99I, MspAII, MspJI, SgrAI, BfaI, BspCNI, XhoI, EarI, AcuI, PstI, BpmI, DdeI, SfcI, AflII, BpuEI, SmlI, AvaI, BsoBI, MboII, BbsI, XmnI, BsmI, Nb.BsmI, EcoRI, HgaI, AatII, ZraI, Tth111I PflFI, PshAI, AhdI, DrdI, Eco53kI, SacI, BseRI, PleI, Nt.BstNBI, MlyI, HinfI, EcoRV, MboI, Sau3AI, DpnII BfuCI, DpnI, BsaBI, TfiI, BsrDI, Nb.BsrDI, BbvI, BtsI, Nb.BtsI, BstAPI, SfaNI, SphI, NmeAIII, NaeI, NgoMIV, BglI, AsiSI, BtgZI, HinPII, HhaI, BssHII, NotI, Fnu4HI, Cac8I, MwoI, NheI, BmtI, SapI, BspQI, Nt.BspQI, BlpI, TseI, ApeKI, Bsp1286I, AlwI, Nt.AlwI, BamHI, FokI, BtsCI, HaeIII, PhoI, FseI, SfiI, NarI, KasI, SfoI, PluTI, AscI, EciI, BsmFI, ApaI, PspOMI, Sau96I, NlaIV, KpnI, Acc65I, BsaI, HphI, BstEII, AvaII, BanI, BaeGI, BsaHI, BanII, RsaI, CviQI, BstZ17I, BciVI, SalI, Nt.BsmAI, BsmAI, BcoDI, ApaLI, BsgI, AccI, Hpy166II, Tsp45I, HpaI, PmeI, HincII, BsiHKAII, ApoI, NspI, BsrFI, BstYI, HaeII, CviKI-1, EcoO109I, PpuMI, I-CeuI, SnaBI, I-SceI, BspHI, BspEI, MmeI, TaqαI, NruI, Hpy188I, Hpy188III, XbaI, BclI, HpyCH4V, FspI, PI-

PspI, MscI, BsrGI, MseI, PacI, PstI, BstBI, DraI, PspXI, BsaWI, BsaAI, EaeI, preferably XhoI, XbaI, HindIII, NcoI, NotI, EcoRI, EcoRV, BamHI, NheI, SacI, SalI, BstBI. The term "multiple cloning site" as used herein further refers to a short DNA sequence used for recombination events as e.g. in Gateway cloning strategy or for methods such as Gibson assembly or topo cloning.

The term "*Yersinia* wild type strain" as used herein refers to a naturally occurring variant (as *Y. enterocolitica* E40) or a naturally occurring variant containing genetic modifications allowing the use of vectors, such as deletion mutations in restriction endonucleases or antibiotic resistance genes (as *Y. enterocolitica* MRS40, the Ampicillin sensitive derivative of *Y. enterocolitica* E40). These strains contain chromosomal DNA as well as an unmodified virulence plasmid (called pYV).

The term "comprise" is generally used in the sense of include, that is to say permitting the presence of one or more features or components.

The term "about" refers to a range of values $\pm 10\%$ of a specified value. For example, the phrase "about 200" includes $\pm 10\%$ of 200, or from 180 to 220.

In one embodiment of the present invention the recombinant Gram-negative bacterial strain is transformed with a vector which comprises in the 5' to 3' direction:

- a first DNA sequence encoding a delivery signal or a fragment thereof from a bacterial effector protein;
- a second DNA sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins fused in frame to the 3' end of said first DNA sequence, wherein the heterologous proteins are selected from the group consisting of proteins involved in apoptosis or apoptosis regulation, cell cycle regulators, ankyrin repeat proteins, cell signaling proteins, reporter proteins, transcription factors, proteases, small GTPases, GPCR related proteins, nanobody fusion constructs and nanobodies, bacterial T3SS effectors, bacterial T4SS effectors and viral proteins. Preferably the DNA sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins is flanked on its 3' end by a DNA sequence homologous to the DNA sequence of the chromosome or of the endogenous virulence plasmid at the 3' end of the delivery

signal from a bacterial effector protein or to a fragment thereof. More preferably, this DNA sequence flanking the homologous protein on its 3' end is homologous to the DNA sequence and is lying within 10kbp on the chromosome or on an endogenous virulence plasmid at the 3' end of the delivery signal from a bacterial effector protein or to a fragment thereof. In particular, this nucleotide sequence flanking the homologous protein on its 3' end is homologous to the DNA sequence and is within the same operon on the chromosome or on an endogenous virulence plasmid as the delivery signal from a bacterial effector protein or a fragment thereof. In this embodiment, transformation is usually performed so that the fused first and the second DNA sequence are inserted by homologous recombination on an endogenous virulence plasmid or a chromosome, preferably on an endogenous virulence plasmid, of the recombinant virulence attenuated Gram-negative bacterial strain, and the fused first and the second DNA sequence is operably linked to a promoter of an endogenous virulence plasmid or of a chromosome e.g. of a chromosomal pathogenicity island. Preferably the fused first and the second DNA sequence is operably linked to a promoter of an endogenous virulence plasmid. In this embodiment the first DNA sequence comprises a delivery signal or fragment thereof from a bacterial effector protein, preferably a fragment thereof, which provides for homologous recombination at the homologous site at the chromosome or at an endogenous virulence plasmid to result in the second DNA sequence be placed in frame to the 3' end of the chromosomal or endogenous virulence plasmid delivery signal which is operatively linked to the endogenous promoter.

In a further embodiment of the present invention the recombinant Gram-negative bacterial strain, is transformed with a nucleotide molecule, preferably a DNA nucleotide molecule, comprising a nucleotide sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins and a nucleotide sequence which is homologous or identical to a nucleotide sequence encoding a delivery signal from a bacterial effector protein or which is homologous or identical to a nucleotide sequence encoding a fragment of a delivery signal from a bacterial effector protein, wherein the delivery signal from a bacterial effector protein is encoded on the chromosome or on an endogenous virulence plasmid of the recombinant virulence attenuated Gram-negative bacterial strain. Preferably the nucleotide sequence which is homologous or identical to a nucleotide sequence of a

delivery signal from a bacterial effector protein or to a fragment thereof is located on the 5' end of the nucleotide sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins. More preferably the nucleotide sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins is flanked on its 3' end by a nucleotide sequence homologous to the DNA sequence of the chromosome or of the endogenous virulence plasmid at the 3' end of the delivery signal from a bacterial effector protein or to a fragment thereof. Even more preferably, this nucleotide sequence flanking the homologous protein on its 3' end is homologous to the DNA sequence lying within 10kbp on the chromosome or on an endogenous virulence plasmid at the 3' end of the delivery signal from a bacterial effector protein or to a fragment thereof. In particular, this nucleotide sequence flanking the homologous protein on its 3' end is homologous to the DNA sequence is within the same operon on the chromosome or on an endogenous virulence plasmid as the delivery signal from a bacterial effector protein or a fragment thereof. In this embodiment, transformation is usually performed so that the nucleotide sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins is inserted on an endogenous virulence plasmid or a chromosome of the recombinant virulence attenuated Gram-negative bacterial strain at the 3' end of a delivery signal from a bacterial effector protein encoded by the chromosome or the endogenous virulence plasmid, wherein the repeated domains of a heterologous protein or two or more domains of different heterologous proteins fused to the delivery signal are expressed and secreted.

In case the recombinant virulence attenuated Gram-negative bacterial strain is a *Yersinia* strain the endogenous virulence plasmid for insertion is pYV (plasmid of *Yersinia* Virulence). In case the recombinant virulence attenuated Gram-negative bacterial strain is a *Salmonella* strain, the endogenous location for insertion is one of the gene clusters called SpiI or SpiII (for *Salmonella* pathogenicity island), a position where an effector protein is elsewhere encoded or alternatively one of the *Salmonella* virulence plasmids (SVPs)..

Preferably the first and the second DNA sequence or the nucleotide molecule are inserted on an endogenous virulence plasmid at the native site of a bacterial effector

protein e.g. at the native site of a virulence factor, preferably in case the recombinant virulence attenuated Gram-negative bacterial strain is a *Yersinia* strain, at the native site of YopE or another Yop (YopH, YopO, YopP, YopM, YopT), preferably at the native site of YopE or in case the recombinant virulence attenuated Gram-negative bacterial strain is a *Salmonella* strain at the native site of an effector protein encoded within SpiI, SpiII or encoded elsewhere, preferably at the native site of an effector protein encoded within SpiI or SpiII, more preferably at the native site of SopE or SteA. Preferably the first and the second DNA sequence or the nucleotide molecule are operably linked to a native promoter of a bacterial effector protein present on an endogenous virulence plasmid e.g. in case the recombinant virulence attenuated Gram-negative bacterial strain is a *Yersinia* strain to a native promoter from a *Yersinia* virulon gene as outlined below, more preferably to the native YopE promoter or another Yop (YopH, YopO, YopP, YopM, YopT) promoter, preferably to the native YopE promoter or in case the recombinant virulence attenuated Gram-negative bacterial strain is a *Salmonella* strain to a native promoter from SpiI or SpiII pathogenicity island or from an effector protein elsewhere encoded as outlined below, more preferably to the native SopE, InvB or SteA promoter.

In one embodiment the present invention provides a recombinant Gram-negative bacterial strain, wherein the Gram-negative bacterial strain is selected from the group consisting of the genera *Yersinia*, *Escherichia*, *Salmonella* and *Pseudomonas*. In one embodiment the present invention provides a recombinant Gram-negative bacterial strain, wherein the Gram-negative bacterial strain is selected from the group consisting of the genera *Yersinia* and *Salmonella*. Preferably the Gram-negative bacterial strain is a *Yersinia* strain, more preferably a *Yersinia enterocolitica* strain. Most preferred is *Yersinia enterocolitica* E40 (O:9, biotype 2) [21] or Ampicilline sensitive derivatives thereof as *Y. enterocolitica* MRS40 (also named *Y. enterocolitica* subsp. *paleartica* MRS40) as described in [22]. *Y. enterocolitica* E40 and its derivative *Y. enterocolitica* MRS40 as described in [22] is identical to *Y. enterocolitica* subsp. *paleartica* E40 and its derivative *Y. enterocolitica* subsp. *paleartica* MRS40 as described in [23-25]. Also preferably the Gram-negative bacterial strain is a *Salmonella* strain, more preferably a *Salmonella enterica* strain. Most preferred is *Salmonella enterica* Serovar Typhimurium SL1344 as described by the Public health England culture collection (NCTC 13347).

In one embodiment of the present invention the delivery signal from a bacterial T3SS effector protein comprises a bacterial T3SS effector protein or a N-terminal fragment thereof wherein the T3SS effector protein or a N-terminal fragment thereof may comprise a chaperone binding site. A T3SS effector protein or a N-terminal fragment thereof which comprises a chaperone binding site is particular useful as delivery signal in the present invention. Preferred T3SS effector proteins or N-terminal fragments thereof are selected from the group consisting of SopE, SopE2, SptP, YopE, ExoS, SipA, SipB, SipD, SopA, SopB, SopD, IpgB1, IpgD, SipC, SifA, SseJ, Sse, SrfH, YopJ, AvrA, AvrBsT, YopT, YopH, YpkA, Tir, EspF, TccP2, IpgB2, OspF, Map, OspG, OspI, IpaH, SspH1, VopF, ExoS, ExoT, HopAB2, XopD, AvrRpt2, HopAO1, HopPtoD2, HopU1, GALA family of proteins, AvrBs2, AvrD1, AvrBS3, YopO, YopP, YopE, YopM, YopT, EspG, EspH, EspZ, IpaA, IpaB, IpaC, VirA, IcsB, OspC1, OspE2, IpaH9.8, IpaH7.8, AvrB, AvrD, AvrPphB, AvrPphC, AvrPphEpto, AvrPpiBpto, AvrPto, AvrPtoB, VirPphA, AvrRpm1, HopPtoE, HopPtoF, HopPtoN, PopB, PopP2, AvrBs3, XopD, and AvrXv3. More preferred T3SS effector proteins or N-terminal fragments thereof are selected from the group consisting of SopE, SptP, YopE, ExoS, SopB, IpgB1, IpgD, YopJ, YopH, EspF, OspF, ExoS, YopO, YopP, YopE, YopM, YopT, whereof most preferred T3SS effector proteins or N-terminal fragments thereof are selected from the group consisting of IpgB1, SopE, SopB, SptP, OspF, IpgD, YopH, YopO, YopP, YopE, YopM, YopT, in particular YopE or an N-terminal fragment thereof.

Equally preferred T3SS effector proteins or N-terminal fragments thereof are selected from the group consisting of SopE, SopE2, SptP, SteA, SipA, SipB, SipD, SopA, SopB, SopD, IpgB1, IpgD, SipC, SifA, SifB, SseJ, Sse, SrfH, YopJ, AvrA, AvrBsT, YopH, YpkA, Tir, EspF, TccP2, IpgB2, OspF, Map, OspG, OspI, IpaH, VopF, ExoS, ExoT, HopAB2, AvrRpt2, HopAO1, HopU1, GALA family of proteins, AvrBs2, AvrD1, YopO, YopP, YopE, YopT, EspG, EspH, EspZ, IpaA, IpaB, IpaC, VirA, IcsB, OspC1, OspE2, IpaH9.8, IpaH7.8, AvrB, AvrD, AvrPphB, AvrPphC, AvrPphEpto, AvrPpiBpto, AvrPto, AvrPtoB, VirPphA, AvrRpm1, HopPtoD2, HopPtoE, HopPtoF, HopPtoN, PopB, PopP2, AvrBs3, XopD, and AvrXv3. Equally more preferred T3SS effector proteins or N-terminal fragments thereof are selected from the group consisting of SopE, SptP, SteA, SifB, SopB, IpgB1, IpgD, YopJ, YopH, EspF, OspF, ExoS, YopO, YopP, YopE, YopT, whereof equally most

preferred T3SS effector proteins or N-terminal fragments thereof are selected from the group consisting of IpgB1, SopE, SopB, SptP, SteA, SifB, OspF, IpgD, YopH, YopO, YopP, YopE, and YopT, in particular SopE, SteA, or YopE or an N-terminal fragment thereof, more particular SteA or YopE or an N-terminal fragment thereof, most particular YopE or an N-terminal fragment thereof.

In some embodiments the delivery signal from the bacterial T3SS effector protein encoded by the first DNA sequence comprises the bacterial T3SS effector protein or an N-terminal fragment thereof, wherein the N-terminal fragment thereof includes at least the first 10, preferably at least the first 20, more preferably at least the first 100 amino acids of the bacterial T3SS effector protein.

In some embodiments the delivery signal from the bacterial T3SS effector protein encoded by the first DNA sequence comprises the bacterial T3SS effector protein or an N-terminal fragment thereof, wherein the bacterial T3SS effector protein or the N-terminal fragment thereof comprises a chaperone binding site.

Preferred T3SS effector proteins or a N-terminal fragment thereof, which comprise a chaperone binding site comprise the following combinations of chaperone binding site and T3SS effector protein or N-terminal fragment thereof: SycE-YopE, InvB-SopE, SicP-SptP, SycT-YopT, SycO-YopO, SycN/YscB-YopN, SycH-YopH, SpsS-ExoS, CesF-EspF, SycD-YopB, SycD-YopD. More preferred are SycE-YopE, InvB-SopE, SycT-YopT, SycO-YopO, SycN/YscB-YopN, SycH-YopH, SpsS-ExoS, CesF-EspF. Most preferred is a YopE or an N-terminal fragment thereof comprising the SycE chaperone binding site such as an N-terminal fragment of a YopE effector protein containing the N-terminal 138 amino acids of the YopE effector protein designated herein as YopE₁₋₁₃₈ and as shown in SEQ ID NO. 2 or a SopE or an N-terminal fragment thereof comprising the InvB chaperone binding site such as an N-terminal fragment of a SopE effector protein containing the N-terminal 81 or 105 amino acids of the SopE effector protein designated herein as SopE₁₋₈₁ or SopE₁₋₁₀₅ respectively, and as shown in SEQ ID NO.: 142 or 143.

In one embodiment of the present invention the recombinant Gram-negative bacterial strain is a *Yersinia* strain and the delivery signal from the bacterial T3SS effector protein encoded by the first DNA sequence comprises a YopE effector protein or an N-terminal part, preferably the *Y. enterocolitica* YopE effector protein or an N-terminal part thereof. Preferably the SycE binding site is comprised within the N-

terminal part of the YopE effector protein. In this connection an N-terminal fragment of a YopE effector protein may comprise the N-terminal 12, 16, 18, 52, 53, 80 or 138 amino acids [26-28]. Most preferred is an N-terminal fragment of a YopE effector protein containing the N-terminal 138 amino acids of the YopE effector protein e.g. as described in Forsberg and Wolf-Watz [29] designated herein as YopE₁₋₁₃₈ and as shown in SEQ ID NO.: 2.

In one embodiment of the present invention the recombinant Gram-negative bacterial strain is a *Salmonella* strain and the delivery signal from the bacterial T3SS effector protein encoded by the first DNA sequence comprises a SopE or SteA effector protein or an N-terminal part thereof, preferably the *Salmonella enterica* SopE or SteA effector protein or an N-terminal part thereof. Preferably the chaperon binding site is comprised within the N-terminal part of the SopE effector protein. In this connection an N-terminal fragment of a SopE effector protein may comprise the N-terminal 81 or 105 amino acids. Most preferred is the full length SteA and an N-terminal fragment of a SopE effector protein containing the N-terminal 105 amino acids of the effector protein e.g. as described in SEQ ID NO. 142 or 143.

One skilled in the art is familiar with methods for identifying the polypeptide sequences of an effector protein that are capable of delivering a protein. For example, one such method is described by Sory et al. [21]. Briefly, polypeptide sequences from e.g. various portions of the Yop proteins can be fused in-frame to a reporter enzyme such as the calmodulin-activated adenylate cyclase domain (or Cya) of the *Bordetella pertussis* cyclolysin. Delivery of a Yop-Cya hybrid protein into the cytosol of eukaryotic cells is indicated by the appearance of cyclase activity in the infected eukaryotic cells that leads to the accumulation of cAMP. By employing such an approach, one skilled in the art can determine, if desired, the minimal sequence requirement, i.e., a contiguous amino acid sequence of the shortest length, that is capable of delivering a protein, see, e.g. [21]. Accordingly, preferred delivery signals of the present invention consists of at least the minimal sequence of amino acids of a T3SS effector protein that is capable of delivering a protein.

In one embodiment the present invention provides mutant recombinant Gram-negative bacterial strains in particular recombinant Gram-negative bacterial strains

which are deficient in producing at least one T3SS functional effector protein. According to the present invention, such a mutant Gram-negative bacterial strain e.g. such a mutant *Yersinia* strain can be generated by introducing at least one mutation into at least one effector-encoding gene. Preferably, such effector-encoding genes include YopE, YopH, YopO/YpkA, YopM, YopP/YopJ and YopT as far as a *Yersinia* strain is concerned. Preferably, such effector-encoding genes include AvrA, CigR, GogB, GtgA, GtgE, PipB, SifB, SipA/SspA, SipB, SipC/SspC, SipD/SspD, SlrP, SopB/SigD, SopA, SpiC/SsaB, SseB, SseC, SseD, SseF, SseG, SseI/SrfH, SopD, SopE, SopE2, SspH1, SspH2, PipB2, SifA, SopD2, SseJ, SseK1, SseK2, SseK3, SseL, SteC, SteA, SteB, SteD, SteE, SpvB, SpvC, SpvD, SrfJ, SptP, as far as a *Salmonella* strain is concerned. Most preferably, all effector-encoding genes are deleted. The skilled artisan may employ any number of standard techniques to generate mutations in these T3SS effector genes. Sambrook et al. describe in general such techniques. See Sambrook et al. [30].

In accordance with the present invention, the mutation can be generated in the promoter region of an effector-encoding gene so that the expression of such effector gene is abolished.

The mutation can also be generated in the coding region of an effector-encoding gene such that the catalytic activity of the encoded effector protein is abolished. The "catalytic activity" of an effector protein refers normally to the anti-target cell function of an effector protein, i.e., toxicity. Such activity is governed by the catalytic motifs in the catalytic domain of an effector protein. The approaches for identifying the catalytic domain and/or the catalytic motifs of an effector protein are well known by those skilled in the art. See, for example, [31,32].

Accordingly, one preferred mutation of the present invention is a deletion of the entire catalytic domain. Another preferred mutation is a frameshift mutation in an effector-encoding gene such that the catalytic domain is not present in the protein product expressed from such "frameshifted" gene. A most preferred mutation is a mutation with the deletion of the entire coding region of the effector protein. Other mutations are also contemplated by the present invention, such as small deletions or base pair substitutions, which are generated in the catalytic motifs of an effector protein leading to destruction of the catalytic activity of a given effector protein.

The mutations that are generated in the genes of the T3SS functional effector proteins may be introduced into the particular strain by a number of methods. One such

method involves cloning a mutated gene into a "suicide" vector which is capable of introducing the mutated sequence into the strain via allelic exchange. An example of such a "suicide" vector is described by [33].

In this manner, mutations generated in multiple genes may be introduced successively into a Gram-negative bacterial strain giving rise to polymutant, e.g. a sextuple mutant recombinant strain. The order in which these mutated sequences are introduced is not important. Under some circumstances, it may be desired to mutate only some but not all of the effector genes. Accordingly, the present invention further contemplates polymutant *Yersinia* other than sextuple-mutant *Yersinia*, e.g., double-mutant, triple-mutant, quadruple-mutant and quintuple-mutant strains. For the purpose of delivering proteins, the secretion and translocation system of the instant mutant strain needs to be intact.

A preferred recombinant Gram-negative bacterial strain of the present invention is a recombinant Gram-negative bacterial strain which is deficient in producing at least one, preferably at least two, more preferably at least three, even more preferably at least four, in particular at least five, more particular at least six, most particular all T3SS effector proteins e.g. a sextuple-mutant Gram-negative bacterial strain in which all the effector-encoding genes are mutated such that the resulting Gram-negative bacterial strain no longer produce any functional effector proteins.

A more preferred recombinant Gram-negative bacterial strain of the present invention is a sextuple-mutant *Yersinia* strain in which all the effector-encoding genes are mutated such that the resulting *Yersinia* no longer produce any functional effector proteins. Such sextuple-mutant *Yersinia* strain is designated as $\Delta yopH,O,P,E,M,T$ for *Y. enterocolitica*. As an example such a sextuple-mutant can be produced from the *Y. enterocolitica* MRS40 strain giving rise to *Y. enterocolitica* MRS40 $\Delta yopH,O,P,E,M,T$, which is preferred.

A further aspect of the present invention is directed to a vector for use in combination with the recombinant Gram-negative bacterial strains to deliver a desired protein into eukaryotic cells, wherein the vector comprises in the 5' to 3' direction:

a promoter;

a first DNA sequence encoding a delivery signal from a bacterial T3SS effector protein, operably linked to said promoter;

a second DNA sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins fused in frame to the 3' end of said first DNA sequence,

wherein the heterologous proteins are selected from the group consisting of proteins involved in apoptosis or apoptosis regulation, cell cycle regulators, ankyrin repeat proteins, cell signaling proteins, reporter proteins, transcription factors, proteases, small GTPases, GPCR related proteins, nanobody fusion constructs and nanobodies, bacterial T3SS effectors, bacterial T4SS effectors and viral proteins.

Promoter, heterologous protein and protease cleavage site as described supra can be used for the vector of the Gram-negative bacterial strain.

A further aspect of the present invention is directed to a vector for use in combination with the recombinant Gram-negative bacterial strains to deliver a desired protein into eukaryotic cells, wherein the vector comprises in the 5' to 3' direction:

a first DNA sequence encoding a delivery signal or a fragment thereof from a bacterial effector protein;

a second DNA sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins fused in frame to the 3' end of said first DNA sequence,

wherein the heterologous proteins are selected from the group consisting of proteins involved in apoptosis or apoptosis regulation, cell cycle regulators, ankyrin repeat proteins, cell signaling proteins, reporter proteins, transcription factors, proteases, small GTPases, GPCR related proteins, nanobody fusion constructs and nanobodies, bacterial T3SS effectors, bacterial T4SS effectors and viral proteins.

Preferably the DNA sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins of the vector is flanked on its 3' end by a DNA sequence homologous to the DNA sequence of the chromosome or of the endogenous virulence plasmid at the 3' end of the delivery signal from a bacterial effector protein or to a fragment thereof. More preferably, this DNA sequence flanking the homologous protein on its 3' end is homologous to the DNA sequence and is lying within 10kbp on the chromosome or on an endogenous virulence plasmid at the 3' end of the delivery signal from a bacterial effector protein or to a fragment thereof. In particular, this nucleotide sequence flanking the homologous protein on its 3' end is homologous to the DNA sequence and is within the same operon on the

chromosome or on an endogenous virulence plasmid as the delivery signal from a bacterial effector protein or a fragment thereof. Heterologous protein and protease cleavage site as described supra can be used for the vector of the Gram-negative bacterial strain.

Vectors which can be used according to the invention depend on the Gram-negative bacterial strains used as known to the skilled person. Vectors which can be used according to the invention include expression vectors (including synthetic or otherwise generated modified versions of endogenous virulence plasmids), vectors for chromosomal or virulence plasmid insertion and DNA fragments for chromosomal or virulence plasmid insertion. Expression vectors which are useful in e.g. *Yersinia*, *Escherichia*, *Salmonella* or *Pseudomonas* strain are e.g. pUC, pBad, pACYC, pUCP20 and pET plasmids. Vectors for chromosomal or virulence plasmid insertion which are useful in e.g. *Yersinia*, *Escherichia*, *Salmonella* or *Pseudomonas* strain are e.g. pKNG101. DNA fragments for chromosomal or virulence plasmid insertion refer to methods used in e.g. *Yersinia*, *Escherichia*, *Salmonella* or *Pseudomonas* strain as e.g. lambda-red genetic engineering. Vectors for chromosomal or virulence plasmid insertion or DNA fragments for chromosomal or virulence plasmid insertion may insert the first, second and/or third DNA sequence of the present invention so that the first, second and/or third DNA sequence is operably linked to an endogenous promoter of the recombinant Gram-negative bacterial strain. Thus if a vector for chromosomal or virulence plasmid insertion or a DNA fragment for chromosomal or virulence plasmid insertion is used, an endogenous promoter can be encoded on the endogenous bacterial DNA (chromosomal or plasmid DNA) and only the first and second DNA sequence will be provided by the engineered vector for chromosomal or virulence plasmid insertion or DNA fragment for chromosomal or virulence plasmid insertion. Alternatively, if a vector for chromosomal or virulence plasmid insertion or a nucleotide molecule such as e.g. a DNA sequence for chromosomal or virulence plasmid insertion is used, an endogenous promoter and the delivery signal from a bacterial effector protein can be encoded on the endogenous bacterial DNA (chromosomal or plasmid DNA) and only the nucleotide molecule such as e.g. a DNA sequence encoding the heterologous protein will be provided by a vector for chromosomal or virulence plasmid insertion or by a nucleotide molecule such as e.g. a DNA sequence for chromosomal or virulence plasmid insertion. Thus a promoter is

not necessarily needed to be comprised by the vector used for transformation of the recombinant Gram-negative bacterial strains i.e. the recombinant Gram-negative bacterial strains of the present invention may be transformed with a vector which does not comprise a promoter. The vector of the present invention is normally used for delivery of the heterologous proteins by the bacterial T3SS into eukaryotic cells in vitro and in vivo.

A preferred vector e.g. a preferred expression vector for *Yersinia* is selected from the group consisting of pBad_Si_1 and pBad_Si_2. pBad_Si2 was constructed by cloning of the SycE-YopE₁₋₁₃₈ fragment containing endogenous promoters for YopE and SycE from purified pYV40 into KpnI/HindIII site of pBad-MycHisA (Invitrogen).

Additional modifications include removal of the NcoI/BglII fragment of pBad-MycHisA by digest, Klenow fragment treatment and religation. Further at the 3' end of YopE₁₋₁₃₈ the following cleavage sites were added: XbaI-XhoI-BstBI-(HindIII).

pBad_Si1 is equal to pBad_Si2 but encodes EGFP amplified from pEGFP-C1 (Clontech) in the NcoI/BglII site under the Arabinose inducible promoter. Equally preferred is the use of modified versions of the endogenous *Yersinia* virulence plasmid pYV encoding heterologous proteins as fusions to a T3SS signal sequence.

A preferred vector e.g. a preferred expression vector for *Salmonella* is selected from the group consisting of pSi_266, pSi_267, pSi_268 and pSi_269. Plasmids pSi_266, pSi_267, pSi_268 and pSi_269 containing the corresponding endogenous promoter and the SteA₁₋₂₀ fragment (pSi_266), the full length SteA sequence (pSi_267), the SopE₁₋₈₁ fragment (pSi_268) or the SopE₁₋₁₀₅ fragment (pSi_269) were amplified from *S. enterica* SL1344 genomic DNA and cloned into NcoI/KpnI site of pBad-MycHisA (Invitrogen).

The vectors of the instant invention may include other sequence elements such as a 3' termination sequence (including a stop codon and a poly A sequence), or a gene conferring a drug resistance which allows the selection of transformants having received the instant vector.

The vectors of the present invention may be transformed by a number of known methods into the recombinant Gram-negative bacterial strains. For the purpose of the present invention, the methods of transformation for introducing a vector include, but are not limited to, electroporation, calcium phosphate mediated transformation, conjugation, or combinations thereof. For example, a vector can be transformed into a

first bacteria strain by a standard electroporation procedure. Subsequently, such a vector can be transferred from the first bacteria strain into the desired strain by conjugation, a process also called "mobilization". Transformant (i.e., Gram-negative bacterial strains having taken up the vector) may be selected, e.g., with antibiotics. These techniques are well known in the art. See, for example, [21].

In accordance with the present invention, the promoter of the expression vector of the recombinant Gram-negative bacterial strain of the invention can be a native promoter of a T3SS effector protein of the respective strain or a compatible bacterial strain or a promoter used in expression vectors which are useful in e.g. *Yersinia*, *Escherichia*, *Salmonella* or *Pseudomonas* strain e.g. pUC and pBad. Such promoters are the T7 promoter, Plac promoter or Ara-bad promoter.

If the recombinant Gram-negative bacterial strain is a *Yersinia* strain the promoter can be from a *Yersinia* virulon gene. A "Yersinia virulon gene" refers to genes on the *Yersinia* pYV plasmid, the expression of which is controlled both by temperature and by contact with a target cell. Such genes include genes coding for elements of the secretion machinery (the Ysc genes), genes coding for translocators (YopB, YopD, and LcrV), genes coding for the control elements (YopN, TyeA and LcrG), genes coding for T3SS effector chaperones (SycD, SycE, SycH, SycN, SycO and SycT), and genes coding for effectors (YopE, YopH, YopO/YpkA, YopM, YopT and YopP/YopJ) as well as other pYV encoded proteins as VirF and YadA.

In a preferred embodiment of the present invention, the promoter is the native promoter of a T3SS functional effector encoding gene. If the recombinant Gram-negative bacterial strain is a *Yersinia* strain the promoter is selected from any one of YopE, YopH, YopO/YpkA, YopM and YopP/YopJ. More preferably, the promoter is from YopE or SycE.

If the recombinant Gram-negative bacterial strain is a *Salmonella* strain the promoter can be from SpiII or SpiIII pathogenicity island or from an effector protein elsewhere encoded. Such genes include genes coding for elements of the secretion machinery, genes coding for translocators, genes coding for the control elements, genes coding for T3SS effector chaperones, and genes coding for effectors as well as other proteins encoded by SPI-1 or SPI-2. In a preferred embodiment of the present invention, the promoter is the native promoter of a T3SS functional effector encoding gene. If the recombinant Gram-negative bacterial strain is a *Salmonella* strain the promoter is

selected from any one of the effector proteins. More preferably, the promoter is from SopE, InvB or SteA.

In a preferred embodiment the vector e.g. the expression vector comprises a DNA sequence encoding a protease cleavage site. Generation of a functional and generally applicable cleavage site allows cleaving off the delivery signal after translocation. As the delivery signal can interfere with correct localization and/or function of the translocated protein within the target cells the introduction of a protease cleavage site between the delivery signal and the protein of interest provides for the first time delivery of almost native proteins into eukaryotic cells. Preferably the protease cleavage site is an amino acid motif which is cleaved by a protease or the catalytic domains thereof selected from the group consisting of enterokinase (light chain), enteropeptidase, prescission protease, human rhinovirus protease 3C, TEV protease, TVMV protease, FactorXa protease and thrombin, more preferably an amino acid motif which is cleaved by TEV protease. Equally preferable the protease cleavage site is an amino acid motif which is cleaved by a protease or the catalytic domains thereof selected from the group consisting of enterokinase (light chain), enteropeptidase, prescission protease, human rhinovirus protease 3C, TEV protease, TVMV protease, FactorXa protease, ubiquitin processing protease, called Deubiquitinating enzymes, and thrombin. Most preferred is an amino acid motif which is cleaved by TEV protease or by an ubiquitin processing protease.

Thus in a further embodiment of the present invention, the heterologous protein is cleaved from the delivery signal from a bacterial T3SS effector protein by a protease. Preferred methods of cleavage are methods wherein:

a) the protease is translocated into the eukaryotic cell by a recombinant Gram-negative bacterial strain as described herein which expresses a fusion protein which comprises the delivery signal from the bacterial T3SS effector protein and the protease as heterologous protein; or

b) the protease is expressed constitutively or transiently in the eukaryotic cell.

Usually the recombinant Gram-negative bacterial strain used to deliver a desired protein into a eukaryotic cell and the recombinant Gram-negative bacterial strain translocating the protease into the eukaryotic cell are different.

In one embodiment of the present invention the vector comprises a further DNA sequence encoding a labelling molecule or an acceptor site for a labelling molecule. The further DNA sequence encoding a labelling molecule or an acceptor site for a labelling molecule is usually fused to the 5' end or to the 3' end of the second DNA sequence. A preferred labelling molecule or an acceptor site for a labelling molecule is selected from the group consisting of enhanced green fluorescent protein (EGFP), coumarin, coumarin ligase acceptor site, resorufin, resorufin ligase acceptor site, the tetra-Cysteine motif in use with FLAsH/ReAsH dye (life technologies). Most preferred is resorufin and a resorufin ligase acceptor site or EGFP. The use of a labelling molecule or an acceptor site for a labelling molecule will lead to the attachment of a labelling molecule to the heterologous protein of interest, which will then be delivered as such into the eukaryotic cell and enables tracking of the protein by e.g. live cell microscopy.

In one embodiment of the present invention the vector comprises a further DNA sequence encoding a peptide tag. The further DNA sequence encoding a peptide tag is usually fused to the 5' end or to the 3' end of the second DNA sequence. A preferred peptide tag is selected from the group consisting of Myc-tag, His-tag, Flag-tag, HA tag, Strep tag or V5 tag or a combination of two or more tags out of these groups. Most preferred is Myc-tag, Flag-tag, His-tag and combined Myc- and His-tags. The use of a peptide tag will lead to traceability of the tagged protein e.g. by immunofluorescence or Western blotting using anti-tag antibodies. Further, the use of a peptide tag allows affinity purification of the desired protein either after secretion into the culture supernatant or after translocation into eukaryotic cells, in both cases using a purification method suiting the corresponding tag (e.g. metal-chelate affinity purification in use with a His-tag or anti-Flag antibody based purification in use with the Flag-tag).

In one embodiment of the present invention the vector comprises a further DNA sequence encoding a nuclear localization signal (NLS). The further DNA sequence encoding a nuclear localization signal (NLS) is usually fused to the 5' end or to the 3' end of the second DNA sequence wherein said further DNA sequence encodes a nuclear localization signal (NLS). A preferred NLS is selected from the group consisting of SV40 large T-antigen NLS and derivatives thereof [34] as well as other

viral NLS. Most preferred is SV40 large T-antigen NLS and derivatives thereof.

In one embodiment of the present invention the vector comprises a multiple cloning site. The multiple cloning site is usually located at the 3' end of the first DNA sequence and/or at the 5' end or 3' end of the second DNA sequence. One or more than one multiple cloning sites can be comprised by the vector. A preferred multiple cloning site is selected from the group of restriction enzymes consisting of XhoI, XbaI, HindIII, NcoI, NotI, EcoRI, EcoRV, BamHI, NheI, SacI, SalI, BstBI. Most preferred is XbaI, XhoI, BstBI and HindIII.

The fused protein expressed from the first and second and optional third DNA sequences of the vector is also termed as a "fusion protein" or a "hybrid protein", i.e., a fused protein or hybrid of delivery signal and repeated domains of a heterologous protein or two or more domains of different heterologous proteins.

The present invention contemplates a method for delivering repeated domains of a heterologous protein or two or more domains of different heterologous proteins as hereinabove described into eukaryotic cells in cell culture as well as in-vivo.

Thus in one embodiment the method for delivering repeated domains of a heterologous protein or two or more domains of different heterologous proteins comprises

- i) culturing the Gram-negative bacterial strain as described herein;
- ii) contacting a eukaryotic cell with the Gram-negative bacterial strain of i) wherein a fusion protein which comprises a delivery signal from a bacterial T3SS effector protein and the repeated domains of a heterologous protein or the two or more domains of different heterologous proteins is expressed by the Gram-negative bacterial strain and is translocated into the eukaryotic cell; and optionally
- iii) cleaving the fusion protein so that the repeated domains of a heterologous protein or the two or more domains of different heterologous proteins are cleaved from the delivery signal from the bacterial T3SS effector protein.

In some embodiments at least two fusion proteins which comprises each a delivery signal from a bacterial effector protein and repeated domains of a heterologous

protein or the two or more domains of different heterologous proteins are expressed by the recombinant virulence attenuated Gram-negative bacterial strain and are translocated into the eukaryotic cell by the methods of the present inventions.

The recombinant Gram-negative bacterial strain can be cultured so that a fusion protein is expressed which comprises the delivery signal from the bacterial T3SS effector protein and the repeated domains of a heterologous protein or the two or more domains of different heterologous proteins according to methods known in the art (e.g. FDA, Bacteriological Analytical Manual (BAM), chapter 8: *Yersinia enterocolitica*). Preferably the recombinant Gram-negative bacterial strain can be cultured in Brain Heart infusion broth e.g. at 28°C. For induction of expression of T3SS and e.g. YopE/SycE promoter dependent genes, bacteria can be grown at 37°C.

In a preferred embodiment, the eukaryotic cell is contacted with two Gram-negative bacterial strains of i), wherein the first Gram-negative bacterial strain expresses a first fusion protein which comprises the delivery signal from the bacterial T3SS effector protein and repeated domains of a heterologous protein or two or more domains of different heterologous proteins and the second Gram-negative bacterial strain expresses a second fusion protein which comprises the delivery signal from the bacterial T3SS effector protein and repeated domains of a second heterologous protein or two or more domains of a different second heterologous proteins of a second heterologous protein, so that the first and the second fusion protein are translocated into the eukaryotic cell. This embodiment provided for co-infection of e.g. eukaryotic cells with two bacterial strains as a valid method to deliver e.g. two different hybrid proteins into single cells to address their functional interaction.

Those skilled in the art can also use a number of assays to determine whether the delivery of a fusion protein is successful. For example, the fusion protein may be detected via immunofluorescence using antibodies recognizing a fused tag (like Myc-tag). The determination can also be based on the enzymatic activity of the protein being delivered.

The present invention contemplates a wide range of eukaryotic cells that may be targeted by the instant recombinant Gram-negative bacterial strain e.g. Hi-5 (BTI-TN-

5B1-4; life technologies B855-02), HeLa cells, e.g. HeLa Ccl2 (as ATCC No. CCL-2), fibroblast cells, e.g. 3T3 fibroblast cells (as ATCC No. CCL-92) or Mef (as ATCC No. SCRC-1040), Hek (as ATCC No. CRL-1573), HUVECs (as ATCC No. PCS-100-013), CHO (as ATCC No. CCL-61), Jurkat (as ATCC No. TIB-152), Sf-9 (as ATCC No. CRL-1711), HepG2 (as ATCC No. HB-8065), Vero (as ATCC No. CCL-81), MDCK (as ATCC No. CCL-34), THP-1 (as ATCC No. TIB-202), J774 (as ATCC No. TIB-67), RAW (as ATCC No. TIB-71), Caco2 (as ATCC No. HTB-37), NCI cell lines (as ATCC No. HTB-182), DU145 (as ATCC No. HTB-81), Lncap (as ATCC No. CRL-1740), MCF-7 (as ATCC No. HTB-22), MDA-MB cell lines (as ATCC No. HTB-128), PC3 (as ATCC No. CRL-1435), T47D (as ATCC No. CRL-2865), A549 (as ATCC No. CCL-185), U87 (as ATCC No. HTB-14), SHSY5Y (as ATCC No. CRL-2266s), Ea.Hy926 (as ATCC No. CRL-2922), Saos-2 (as ATCC No. HTBH-85), 4T1 (as ATCC No. CRL-2539), B16F10 (as ATCC No. CRL-6475), or primary human hepatocytes (as life technologies HMCPIs), preferably HeLa, Hek, HUVECs, 3T3, CHO, Jurkat, Sf-9, HepG2 Vero, THP-1, Caco2, Mef, A549, 4T1, B16F10 and primary human hepatocytes and most preferably HeLa, Hek, HUVECs, 3T3, CHO, Jurkat, THP-1, A549 and Mef. By "target", is meant the extracellular adhesion of the recombinant Gram-negative bacterial strain to a eukaryotic cell.

In accordance with the present invention, the delivery of a protein can be achieved by contacting a eukaryotic cell with a recombinant Gram-negative bacterial strain under appropriate conditions. Various references and techniques are conventionally available for those skilled in the art regarding the conditions for inducing the expression and translocation of virulon genes, including the desired temperature, Ca^{++} concentration, addition of inducers as Congo Red, manners in which the recombinant Gram-negative bacterial strain and target cells are mixed, and the like. See, for example, [35]. The conditions may vary depending on the type of eukaryotic cells to be targeted and the recombinant bacterial strain to be used. Such variations can be addressed by those skilled in the art using conventional techniques.

Those skilled in the art can also use a number of assays to determine whether the delivery of a fusion protein is successful. For example, the fusion protein may be detected via immunofluorescence using antibodies recognizing a fused tag (like Myc-tag). The determination can also be based on the enzymatic activity of the protein being delivered, e.g., the assay described by [21].

In one embodiment the present invention provides the recombinant Gram-negative bacterial strain as described herein for use in medicine.

In one embodiment the present invention provides the recombinant Gram-negative bacterial strain as described herein for use in the delivery of repeated domains of a heterologous protein or two or more domains of different heterologous proteins as a medicament or as a vaccine to a subject. The repeated domains of a heterologous protein or the two or more domains of different heterologous proteins can be delivered to a subject as a vaccine by contacting the Gram-negative bacterial strain with eukaryotic cells, e.g. with a living animal in vivo so that the repeated domains of a heterologous protein or the two or more domains of different heterologous proteins are translocated into the living animal which then produces antibodies against the repeated domains of a heterologous protein or the two or more domains of different heterologous proteins. The antibodies produced can be directly used or be isolated and purified and used in diagnosis, in research use as well as in therapy. The B-cells producing the antibodies or the therein contained DNA sequence can be used for further production of specific antibodies for use in diagnosis, in research use as well as in therapy.

In one embodiment the present invention provides a method for delivering repeated domains of a heterologous protein or the two or more domains of different heterologous proteins, wherein the repeated domains of a heterologous protein or two or more domains of different heterologous proteins are delivered in vitro into a eukaryotic cell.

In a further embodiment the present invention provides a method for delivering repeated domains of a heterologous protein or two or more domains of different heterologous proteins, wherein the eukaryotic cell is a living animal wherein the living animal is contacted with the Gram-negative bacterial strain in vivo so that a fusion protein is translocated into the living animal. The preferred animal is a mammal, more preferably a human being.

In a further embodiment the present invention provides the use of the recombinant Gram-negative bacterial strain as described supra for High Throughput Screenings of inhibitors for a cellular pathway or event triggered by the translocated heterologous protein(s).

In a further aspect the present invention provides a library of Gram-negative bacterial strains, wherein the the repeated domains of a heterologous protein or the two or more domains of different heterologous proteins encoded by the second DNA sequence of the expression vector of the Gram-negative bacterial strains are domains of a human or murine protein, preferably a domain of a human protein and, wherein each domain of a human or murine protein expressed by a Gram-negative bacterial strain is different in amino acid sequence. As cloning vector for expression the above described expression vectors can be used.

In a further aspect the present invention provides a kit comprising a vector as described herein and a bacterial strain expressing and secreting a protease capable of cleaving the protease cleavage site comprised by the vector. A particular useful vector is a vector for use in combination with the bacterial strain to deliver a desired protein into eukaryotic cells as described above, wherein the vector comprises in the 5' to 3' direction:

a promoter;

a first DNA sequence encoding a delivery signal from a bacterial T3SS effector protein, operably linked to said promoter;

a second DNA sequence encoding repeated domains of a heterologous protein or two or more domains of different heterologous proteins fused in frame to the 3'end of said first DNA sequence,

wherein the heterologous proteins are selected from the group consisting of proteins involved in apoptosis or apoptosis regulation, cell cycle regulators, ankyrin repeat proteins, cell signaling proteins, reporter proteins, transcription factors, proteases, small GTPases, GPCR related proteins, nanobody fusion constructs and nanobodies, bacterial T3SS effectors, bacterial T4SS effectors and viral proteins.

Examples

Example 1:

A) Materials and Methods

Bacterial strains and growth conditions. The strains used in this study are listed in Figs. 15A to N. *E. coli* Top10, used for plasmid purification and cloning, and *E. coli* Sm10 λ pir, used for conjugation, as well as *E. coli* BW19610 [36], used to propagate pKNG101, were routinely grown on LB agar plates and in LB broth at 37°C. Ampicillin was used at a concentration of 200 μ g/ml (*Yersinia*) or 100 μ g/ml (*E. coli*) to select for expression vectors. Streptomycin was used at a concentration of 100 μ g/ml to select for suicide vectors. *Y. enterocolitica* MRS40 [22] a non Ampicillin resistant E40-derivate [21] and strains derived thereof were routinely grown on Brain Heart Infusion (BHI; Difco) at RT. To all *Y. enterocolitica* strains Nalidixic acid was added (35 μ g/ml) and all *Y. enterocolitica* *asd* strains were additionally supplemented with 100 μ g/ml *meso*-2,6-Diaminopimelic acid (*m*DAP, Sigma Aldrich). *S. enterica* SL1344 were routinely grown on LB agar plates and in LB broth at 37°C. Ampicillin was used at a concentration of 100 μ g/ml to select for expression vectors in *S. enterica*.

Genetic manipulations of *Y. enterocolitica*. Genetic manipulations of *Y. enterocolitica* has been described [37,38]. Briefly, mutators for modification or deletion of genes in the pYV plasmids or on the chromosome were constructed by 2-fragment overlapping PCR using purified pYV40 plasmid or genomic DNA as template, leading to 200–250 bp of flanking sequences on both sides of the deleted or modified part of the respective gene. Resulting fragments were cloned in pKNG101 [33] in *E. coli* BW19610 [36]. Sequence verified plasmids were transformed into *E. coli* Sm10 λ pir, from where plasmids were mobilized into the corresponding *Y. enterocolitica* strain. Mutants carrying the integrated vector were propagated for several generations without selection pressure. Then sucrose was used to select for clones that have lost the vector. Finally mutants were identified by colony PCR. Specific mutators (pSi_408, pSi_419) are listed in Table III.

Construction of plasmids. Plasmid pBad_Si2 or pBad_Si1 (Fig. 10) were used for

cloning of fusion proteins with the N-terminal 138 amino acids of YopE (SEQ ID No. 2). pBad_Si2 was constructed by cloning of the SycE-YopE₁₋₁₃₈ fragment containing endogenous promoters for YopE and SycE from purified pYV40 into KpnI/HindIII site of pBad-MycHisA (Invitrogen). Additional modifications include removal of the NcoI/BglII fragment of pBad-MycHisA by digestion, Klenow fragment treatment and religation. A bidirectional transcriptional terminator (BBa_B1006; iGEM foundation) was cloned into KpnI cut and Klenow treated (pBad_Si2) or BglII cut site (pBad_Si1). Further at the 3' end of YopE₁₋₁₃₈ the following cleavage sites were added: XbaI-XhoI-BstBI-(HindIII) (Fig. 10 B). pBad_Si1 is equal to pBad_Si2 but encodes EGFP amplified from pEGFP-C1 (Clontech) in the NcoI/BglII site under the Arabinose inducible promoter. Plasmids pSi_266, pSi_267, pSi_268 and pSi_269 containing the corresponding endogenous promoter and the SteA₁₋₂₀ fragment (pSi_266), the full length SteA sequence (pSi_267), the SopE₁₋₈₁ fragment (pSi_268) or the SopE₁₋₁₀₅ fragment (pSi_269) were amplified from *S. enterica* SL1344 genomic DNA and cloned into NcoI/KpnI site of pBad-MycHisA (Invitrogen).

Full length genes or fragments thereof were amplified with the specific primers listed in Table I below and cloned as fusions to YopE₁₋₁₃₈ into plasmid pBad_Si2 or in case of z-BIM (SEQ ID No. 21) into pBad_Si1 (see Table II below). For fusion to SteA or SopE, synthetic DNA constructs were cleaved by KpnI/HindII and cloned into pSi_266, pSi_267, pSi_268 or pSi_269 respectively. In case of genes of bacterial species, purified genomic DNA was used as template (*S. flexneri* M90T, *Salmonella enterica* subsp. *enterica* serovar *Typhimurium* SL1344, *Bartonella henselae* ATCC 49882). For human genes a universal cDNA library (Clontech) was used if not otherwise stated (Figs. 15A to N), zebrafish genes were amplified from a cDNA library (a kind gift of M. Affolter). Ligated plasmids were cloned in *E. coli* Top10. Sequenced plasmids were electroporated into the desired *Y. enterocolitica* or *S. enterica* strain using settings as for standard *E. coli* electroporation.

Table I (Primer Nr. Si_:Sequence)

285: CATAACCATGGGAGTGAGCAAGGGCGAG
 286: GGAAGATCTtACTTGTAACAGCTCGTCCAT
 287: CGGGGTACCTCAACTAAATGACCGTG

288: GTTAAAGCTTTtcgaatctagactcgagCGTGGCGAACTGGTC
292: CAGTctcgagCAAATTCTAAACAAAATACTTCCAC
293: cagtTTCGAATTAATTTGTATTGCTTTGACGG
296: CAGTctcgagACTAACATAACACTATCCACCCAG
297: GTTAAAGCTTTTCAGGAGGCATTCTGAAG
299: CAGTctcgagCAGGCCATCAAGTGTGTG
300: cagtTTCGAATCATTCTCTTCCTCTTCTTCA
301: CAGTctcgagGCTGCCATCCGGAA
302: cagtTTCGAATCACAAGACAAGGCACCC
306: GTTAAAGCTTTGGAGGCATTCTGAAGatacttatt
307: CAGTctcgagCAAATACAGAGCTTCTATCACTCAG
308: GTTAAAGCTTTCAAGATGTGATTAATGAAGAAATG
317: cagtTTCGAACCCATAAAAAAGCCCTGTC
318: GTTAAAGCTTCTACTCTATCATCAAACGATAAAATGg
324: CAGTctcgagTTCACTCAAGAAACGCAAA
339: cagtTTCGAATTTTCTCTTCCTCTTCTTCAcg
341: cgtaTCTAGAAAAATGATGAAAATGGAGACTG
342: GTTAAAGCTTTtaGCTGGAGACGGTGAC
346: CAGTctcgagTTCCAGATCCCAGAGTTTG
347: GTTAAAGCTTTCACTGGGAGGGGG
351: CAGTctcgagctcgagTTATCTACTCATAGAACTACTTTTGCAG
352: cgcGGATCCtcagtgctctgcggcatta
353: CATTTATTCTCCTAGTTAGTCAcagcaactgctgctccttc
354: gaaaggagcagcagttgctgTGAATACTAGGAGGAATAAATG
355: cgattcacggattgctttctCATTATTCCCTCCAGGTACTA
356: TAGTACCTGGAGGGAATAATGagaaagcaatccgtgaatcg
357: cgtaTCTAGAcggctttaagtgcgacattc
364: cgtaTCTAGACTAAAGTATGAGGAGAGAAAATTGAA
365: GTTAAAGCTTTTCAGCTTGCCGTCGT
367: CGTAtctagaGACCCGTTCTGGTGC
369: cgtaTCTAGAccccccaagaagaagc
373: GTTAAAGCTTGCTGGAGACGGTGACC
386: CGTAtctagaTCAGGACGCTTCGGAGGTAG

387: CGTAAtctagaATGGACTGTGAGGTCAACAA
389: CGTAAtctagaGGCAACCGCAGCA
391: GTTAAAGCTTTCAGTCCATCCCATTCTg
403: CGTAAtctagatctggaatatccctggaca
406: GTTAAAGCTTgtctgtctcaatgccacagt
410: CAGTctcgagATGTCCGGGGTGGTg
413: cagtTTCGAATCACTGCAGCATGATGTC
417: CAGTctcgagAGTGGTGTGATGATGACATG
420: cagtTTCGAATTAGTGATAAAAATAGAGTTCTTTTGTGAG
423: CAGTctcgagATGCACATAACTAATTTGGGATT
424: cagtTTCGAATTATACAAATGACGAATACCCTTT
425: GTTAAAGCTTttacacctgctgtcttcttgggcggGCTGGAGACGGTGAC
428: CGTAAtctagaATGGACTTCAACAGGAACTTT
429: CGTAAtctagaGGACATAGTCCACCAGCG
430: GTTAAAGCTTTCAGTTGGATCCGAAAAAC
433: CGTAAtctagaGAATTAAAAAAAACACTCATCCCA
434: CGTAAtctagaCCAAAGGCAAAAGCAAAAA
435: GTTAAAGCTTTTAGCTAGCCATGGCAAGC
436: CGTAAtctagaATGCCCCGCCCC
437: GTTAAAGCTTCTACCCACCGTACTCGTCAAT
438: CGTAAtctagaATGTCTGACACGTCCAGAGAG
439: GTTAAAGCTTTCATCTTCTTCGCAGGAAAAAG
445: cgcGGATCCttatgggttctcacagcaaaa
446: CATTTATTCTCTCTAGTTAGTCAaggcaacagccaatcaagag
447: ctcttgattggctgttgccTACTAACTAGGAGGAATAAATG
448: ttgattgcagtgacatggtgCATTATTCCCTCCAGGTACTA
449: TAGTACCTGGAGGGAATAATGcaccatgtcactgcaatcaa
450: cgtaTCTAGAtagccgcagatgttggtatg
451: CGTAAtctagaGATCAAGTCCAACCTGGTGG
463: CAGTctcgaggaaagcttggttaaggggc
464: cagtTTCGAAttagcgacggcgacg
476: GTTAAAGCTTttACTTGTACAGCTCGTCCAT
477: CGTAAtctagaGTGAGCAAGGGCGAG

478: CAGTctcgagATGGAAGATTATACCAAAATAGAGAAA
479: GTTAAAGCTTCTACATCTTCTTAATCTGATTGTCCa
482: CGTAtctagaATGGCGCTGCAGCt
483: GTTAAAGCTTTCAGTCATTGACAGGAATTTTg
486: CGTAtctagaATGGAGCCGGCGGCG
487: GTTAAAGCTTTCATCGGGGATGTCTg
492: CGTAtctagaATGCGCGAGGAGAACAAGGG
493: GTTAAAGCTTTCAGTCCCCTGTGGCTGTGc
494: CGTAtctagaATGGCCGAGCCTTG
495: GTTAAAGCTTttaTTGAAGATTTGTGGCTCC
504:CGTAtctagaGAAAATCTGTATTTTCAAAGTGAAAATCTGTATTTTCAAA
GTATGCCCCGCCCC
505: GTTAAAGCTTCCCACCGTACTCGTCAATtc
508:CGTAtctagaGAAAATCTGTATTTTCAAAGTGAAAATCTGTATTTTCAAA
GTATGGCCGAGCCTTG
509: GTTAAAGCTTTTGAAGATTTGTGGCTCCc
511:CGTAtctagaGAAAATCTGTATTTTCAAAGTGAAAATCTGTATTTTCAAA
GTGTGAGCAAGGGCGAG
512:CGTAtctagaGAAAATCTGTATTTTCAAAGTGAAAATCTGTATTTTCAAA
GTCCGCCGAAAAAAAAACGTAAAGTTGTGAGCAAGGGCGAG
513:GTTAAAGCTTtAAACTTTACGTTTTTTTTTCGGCGGCTTGTACAGCTCG
TCCAT
515:CGTAtctagaGAAAATCTGTATTTTCAAAGTGAAAATCTGTATTTTCAAA
GTGATTATAAAGATGATGATGATAAAATGGCCGAGCCTTG
558: CGTATCTAGAATGACCAGTTTTGAAGATGC
559:GTTAAAGCTTTCATGACTCATTTTCATCCAT
561:CGTATCTAGAATGAGTCTCTTAAACTGTGAGAACAG
562:GTTAAAGCTTCTACACCCCCGCATCA
580: catgccatgATTTATGGTCATAGATATGACCTC
585: CAGTctcgagATGCAGATCTTCGTCAAGAC
586: GTTAAAGCTTgctagcttcgaaACCACCACGTAGACGTAAGAC
588: cagtTTCGAAGATTATAAAGATGATGATGATAAAATGGCCGAGCCTTG
612: CGGGGTACCatgaggtagcttatttctgataaag

613: CGGGGTACCataattgtccaaatagttatggtagc
 614: catgcatggCGGCAAGGCTCCTC
 615: cggggtaccTTTATTTGTCAACACTGCCC
 616: cggggtaccTGCGGGGTCTTTACTCG
 677:TTACTATTTCGAAGAAATTATTCATAATATTGCCCCGCCATCTGGCCCCAA
 ATTGGTGATGAAATGGATCATTAAGCTTGGAGTA
 678:TACTCCAAGCTTAATGATCCATTTTCATCACCAATTTGGGCCAGATGGC
 GGGCAATATTATGAATAATTTCTTCGAATAGTAA
 682:TTACTACTCGAGAAAAAACTGAGCGAATGTCTGCGCCGCATTGGTGAT
 GAACTGGATAGCTAAGCTTGGAGTA
 683:TACTCCAAGCTTAGCTATCCAGTTCATCACCAATGCGGGCGCAGACATT
 CGCTCAGTTTTTTTCTCGAGTAGTAA
 725: TTACTATTTCGAAGAAATTATTCATAATATTGCC
 726:
 TACTCCAAGCTTACGGTTGAATATTATGATCCATTTTCATCACCAATTTGG
 727:
 TTACTATTTCGAAGCCGGTGGTGCCGAAGAAATTATTCATAATATTGCC
 728: TACTCCAAGCTTAATGATCCATTTTCATCA
 733:TTACTACTCGAGGGTGCCATCGATGCCGAAGAAATTATTCATAATATT
 GCCCCG
 734:TACTCCTTCGAAGGCACCATGATCCATTTTCATCACCAATTTGG
 735:TACTCCTTCGAATTAATGATCCATTTTCATCACCAATTTG
 736:TTACTACTCGAGGGTGCCATCGATGCCAAAAAACTGAGCGAATGTCT
 GCG
 737: TACTCCTTCGAAGGCACCGCTATCCAGTTCATCACCAATG
 738:TACTCCTTCGAATTAGCTATCCAGTTCATCACCAATG

Table II: Cloned fusion proteins

Protein to be delivered by T3SS	Protein Seq. ID. No.	Backbone plasmid	Resulting plasmid name	Primers. Si_Nr.:	Primer Seq. ID No.
YopE1-138-MycHis	3	pBad-MycHisA (Invitrogen)	pBad_Si_1	285/286 (EGFP), 287/288 (sycE-YopE1-138)	44/45 and 46/47

YopE1-138-MycHis	3	pBad-MycHisA (Invitrogen)	pBad_Si_2	287/288 (sycE-YopE1-138)	46/47
YopE1-138-IpgB1	4	pBad_Si_2	pSi_16	292/293	48/49
YopE1-138-SopE	5	pBad_Si_2	pSi_20	296/297	50/51
YopE1-138-Rac1 Q61L	26	pBad_Si_2	pSi_22	299/300	52/53
YopE1-138-RhoA Q61E	27	pBad_Si_2	pSi_24	301/302	54/55
YopE1-138-SopE-MycHis	135	pBad_Si_2	pSi_28	296/306	50/56
YopE1-138-SopB	6	pBad_Si_2	pSi_30	307/308	57/58
YopE1-138-FADD	28	pBad_Si_2	pSi_37	367/386	76/79
YopE1-138-OspF	7	pBad_Si_2	pSi_38	317/318	59/60
YopE1-138-BepG 715-end	136	pBad_Si_2	pSi_43	324/351	61/67
YopE1-138-Rac1 Q61L-MycHis	137	pBad_Si_2	pSi_51	299/339	52/62
YopE1-138-Slmb1-VhH4	32	pBad_Si_2	pSi_53	341/342	63/64
YopE1-138-Bad	29	pBad_Si_2	pSi_57	346/347	65/66
YopE1-138-SptP	8	pBad_Si_2	pSi_64	364/365	74/75
YopE1-138-NLS-Slmb1-VhH4	33	pBad_Si_2	pSi_70	369/342	77/64
YopE1-138-Bid	24	pBad_Si_2	pSi_85	387/391	80/82
YopE1-138-t-Bid	25	pBad_Si_2	pSi_87	389/391	81/82
YopE1-138-Caspase3 p17	22	pBad_Si_2	pSi_97	403/406	83/84
YopE1-138-GPCR GNA12	30	pBad_Si_2	pSi_103	410/413	85/86
YopE1-138-Caspase3 p10/12	23	pBad_Si_2	pSi_106	417/420	87/88
YopE1-138-IpgD	9	pBad_Si_2	pSi_111	423/424	89/90
YopE1-138-Slmb1-VhH4-NLS	34	pBad_Si_2	pSi_112	341/425	63/91
YopE1-138-z-Bid	19	pBad_Si_2	pSi_116	428/430	92/94

YopE1-138-z-t-Bid	20	pBad_Si_2	pSi_117	429/430	93/94
YopE1-138-BepA E305-end	11	pBad_Si_2	pSi_118	433/435	95/97
YopE1-138-BepA	10	pBad_Si_2	pSi_119	434/435	96/97
YopE1-138-ET1	36	pBad_Si_2	pSi_120	436/437	98/99
YopE1-138-z-BIM	21	pbad_Si_1	pSi_121	438/439	100/101
YopE1-138-VhH4 nanobody recognizing EGFP	31	pBad_Si_2	pSi_124	451/373	108/78
YopE1-138-TEV protease S219V	42	pBad_Si_2	pSi_132	463/464	109/110
YopE1-138-EGFP	37	pBad_Si_2	pSi_140	477/476	112/111
YopE1-138-Cdk1	14	pBad_Si_2	pSi_143	478/479	113/114
YopE1-138-Mad2	15	pBad_Si_2	pSi_145	482/483	115/116
YopE1-138-Ink4A	16	pBad_Si_2	pSi_147	486/487	117/118
YopE1-138-Ink4B	17	pBad_Si_2	pSi_150	492/493	119/120
YopE1-138-Ink4C	18	pBad_Si_2	pSi_151	494/495	121/122
YopE1-138-TIFA	13	pBad_Si_2	pSi_153	558/559	131/132
YopE1-138-2x TEVsite - ET1	41	pBad_Si_2	pSi_156	504/505	123/124
YopE1-138-2xTEVsite - EGFP - NLS	39	pBad_Si_2	pSi_159	511/513	127/129
YopE1-138-2xTEVsite - NLS - EGFP	38	pBad_Si_2	pSi_160	512/476	128/111
YopE1-138-2x TEVsite - INK4C	40	pBad_Si_2	pSi_161	508/509	125/126
YopE1-138-2x TEVsite - Flag - INK4C	43	pBad_Si_2	pSi_164	515/509	130/126
YopE1-138-murine Traf6	12	pBad_Si_2	pSi_166	561/562	133/134
YopE1-138-Y. enterocolitica codon optimized	138	pBad_Si_2	pSi_318	677/678	148/149

murine tBid BH3 part					
YopE1-138-Y. enterocolitica codon optimized murine Bax BH3 part	139	pBad_Si_2	pSi_322	682/683	150/151
SteA1-20	140	pBad- MycHisA (Invitrogen)	pSi_266	580/612	152/153
SteA	141	pBad- MycHisA (Invitrogen)	pSi_267	580/613	152/154
SopE1-81	142	pBad- MycHisA (Invitrogen)	pSi_268	614/615	155/156
SopE1-105	143	pBad- MycHisA (Invitrogen)	pSi_269	614/616	155/157
SteA1-20-S. enterica codon optimized murine tBid	144	pSi_266	pSi_270	synthetic construct	/
SteA-S. enterica codon optimized murine tBid	145	pSi_267	pSi_271	synthetic construct	/
SopE1-81-S. enterica codon optimized murine tBid	146	pSi_268	pSi_272	synthetic construct	/
SopE1-105-S. enterica codon optimized murine tBid	147	pSi_269	pSi_273	synthetic construct	/
YopE1-138-Y. enterocolitica codon optimized Ink4A 84-103	158	pBad_Si_2	pSi_362	745/746	172/173
YopE1-138-Y. enterocolitica codon optimized p107/RBL1 657-662 (AAA02489.1)	159	pBad_Si_2	pSi_363	747/748	174/175
YopE1-138-Y. enterocolitica codon optimized p21 141-160 (AAH13967.1)	160	pBad_Si_2	pSi_364	749/750	176/177

YopE1-138-Y. enterocolitica codon optimized p21 145-160 (AAH13967.1)	161	pBad_Si_2	pSi_366	753/754	178/179
YopE1-138-Y. enterocolitica codon optimized p21 17-33 (AAH13967.1)	162	pBad_Si_2	pSi_367	755/756	180/181
YopE1-138-Y. enterocolitica codon optimized cyclin D2 139- 147 (CAA48493.1)	163	pBad_Si_2	pSi_368	757/758	182/183
SteA-Ink4a- MycHis	164	pSi_267	pSi_333	703/704	184/185
SopE1-105- Ink4a-MycHis	165	pSi_269	pSi_334	703/704	184/185
SteA-Ink4c- MycHis	166	pSi_267	pSi_335	PCR1: 705/706; PCR2: 707/708; overlapping PCR: 705/708	186/187, 188/189
SopE1-105- Ink4c-MycHis	167	pSi_269	pSi_336	PCR1: 705/706; PCR2: 707/708; overlapping PCR: 705/708	186/187, 188/189
SteA-Mad2- MycHis	168	pSi_267	pSi_337	709/710	190/191
SopE1-105- Mad2-MycHis	169	pSi_269	pSi_338	709/710	190/191
SteA-Cdk1- MycHis	170	pSi_267	pSi_339	711/712	192/193
SopE1-105- Cdk1-MycHis	171	pSi_269	pSi_340	711/712	192/193
YopE1-138-Y. enterocolitica codon optimized murine tBid	194	pBad_Si_2	pSi_315	synthetic construct	/
YopE1-138- Ubiquitin	195	pBad_Si_2	pSi_236	585/586	197/198
YopE1-138- Ubiquitin-Flag-	196	pSi_236	pSi_237_II	588/509	199/126

INK4C-MycHis					
YopE1-138-(Y. enterocolitica codon optimized murine tBid BH3 part) ready for insertion of further domains	200	pBad_Si_2	pSi_357	733/735	204/205
YopE1-138-(Y. enterocolitica codon optimized murine BAX BH3 part) ready for insertion of further domains	201	pBad_Si_2	pSi_358	736/738	206/207
YopE1-138-(Y. enterocolitica codon optimized murine tBid BH3 part) ₂	202	pSi_357	pSi_371	733/734	204/208
YopE1-(138-Y. enterocolitica codon optimized murine tBid BH3 part- Y. enterocolitica codon optimized murine BAX BH3 part	203	pSi_358	pSi_373	733/734	204/208
YopE ₁₋₁₃₈ -codon optimized murine tBid BH3 extended part	209	pBad_Si_2	pSi_353	725/726	212/213
YopE ₁₋₁₃₈ -10 Aa linker - Y. enterocolitica codon optimized murine tBid BH3 part	210	pBad_Si_2	pSi_354	727/728	214/215
YopE1-(138-Y. enterocolitica codon optimized murine Bax BH3 part- Y. enterocolitica codon optimized murine tBid BH3 part	211	pSI_357	pSi_374	736/737	206/216

Table III: Mutators for genetic modification

Mutator/ Construct	To be inserted onto:	Backbone plasmid	Resulting plasmid name	Primers Si_Nr.:	Primers Seq.Id No.	used with parent strain
YopE ₁₋₁₃₈ - tBID BH3	pYV	pKNG101	pSi_408	Synthetic gene	/	/
YopE ₁₋₁₃₈ - (tBID BH3) ₂	pYV	pKNG101	pSi_419	Synthetic gene	/	Strain mutated with pSi_408

Yop secretion. Induction of the *yop* regulon was performed by shifting the culture to 37°C in BHI-Ox (secretion-permissive conditions) [39]. As carbon source glucose was added (4 mg/ml).

Total cell and supernatant fractions were separated by centrifugation at 20 800 g for 10 min at 4°C. The cell pellet was taken as total cell fraction. Proteins in the supernatant were precipitated with trichloroacetic acid 10% (w/v) final for 1 h at 4°C. After centrifugation (20 800 g for 15 min) and removal of the supernatant, the resulting pellet was washed in ice-cold Acetone over-night. The samples were centrifuged again, the supernatant was discarded and the pellet was air-dried and resuspended in 1x SDS loading dye.

Secreted proteins were analysed by SDS–PAGE; in each case, proteins secreted by 3×10^8 bacteria were loaded per lane. Detection of specific secreted proteins by immunoblotting was performed using 12.5% SDS–PAGE gels. For detection of proteins in total cells, 2×10^8 bacteria were loaded per lane, if not stated otherwise, and proteins were separated on 12.5% SDS–PAGE gels before detection by immunoblotting.

Immunoblotting was carried out using rat monoclonal antibodies against YopE (MIPA193 – 13A9 ; 1:1000, [40]). The antiserum was preabsorbed twice overnight against *Y. enterocolitica* Δ HOPEMT *asd* to reduce background staining. Detection was performed with secondary antibodies directed against rat antibodies and

conjugated to horseradish peroxidase (1:5000; Southern biotech), before development with ECL chemiluminescent substrate (LumiGlo, KPM).

Cell culture and infections. HeLa Ccl2, swiss 3T3 fibroblast cells, 4T1, B16F10 and D2A1 were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% FCS and 2mM L-Glutamine (cDMEM). HUVECs were isolated and cultivated as described [41]. Jurkat and 4T1 cells were cultured in RPMI 1640 supplemented with 10% FCS and 2mM L-Glutamine. *Y. enterocolitica* were grown in BHI with additives overnight at RT, diluted in fresh BHI to an OD₆₀₀ of 0.2 and grown for 2h at RT before a temperature shift to a 37°C waterbath shaker for further 30 min or for 1h in case of delivery of EGFP. Finally, the bacteria were collected by centrifugation (6000 rcf, 30 sec) and washed once with DMEM supplemented with 10 mM HEPES and 2 mM L-glutamine. *S. enterica* were grown in LB with additives overnight at 37°C and either diluted 1:40 in fresh LB and grown for 2.5h at 37°C (SpiI T3SS inducing conditions) or the overnight culture was further incubated at 37°C (SpiII T3SS inducing conditions). Finally, the bacteria were collected by centrifugation (6000 rcf, 30 sec) and washed once with DMEM supplemented with 10 mM HEPES and 2 mM L-glutamine. Cells seeded in 96-well (for Immunofluorescence) or 6-well (for Western blotting) plates were infected at indicated MOIs in DMEM supplemented with 10 mM HEPES and 2 mM L-glutamine. After adding bacteria, plates were centrifuged for 1 min at 1750 rpm and placed at 37°C for indicated time periods. Extracellular bacteria were killed by gentamicin (100 mg/ml) if indicated. In case of immunofluorescence analysis, infection assays were stopped by 4% PFA fixation. For Western blot analysis cells were washed twice with ice-cold PBS and Phospho-safe lysis buffer (Novagen) was added to lyse the cells. After incubation on ice, the cells were centrifuged (16 000 rcf, 25 min, 4°C). Supernatants were collected and analyzed for total protein content by Bradford BCA assay (Pierce) before SDS PAGE and Western blotting using anti-Phospho-Akt (Ser473 and T308, both Cell Signaling), anti-Actin (Millipore), Anti-Bid (Cell Signaling), anti-Myc (Santa Cruz), anti-p38 (Cell Signaling), anti-phospho-p-38 (Thr180/Tyr182; Cell Signaling), anti-Caspase-3 p17 (Cell Signaling) and anti-Ink4C (Cell Signaling) antibody.

Secretion analysis with *S. enterica*. For induction of protein secretion by *S. enterica*,

S. enterica were cultivated overnight in LB containing 0.3 M NaCl on an orbital shaker (set to 150rpm). *S. enterica* were then diluted 1:50 in fresh LB containing 0.3 M NaCl and grown for 4h at 37°C without shaking.

Total cell and supernatant fractions were separated by centrifugation at 20 800 g for 20 min at 4°C. The cell pellet was taken as total cell fraction. Proteins in the supernatant were precipitated with trichloroacetic acid 10% (w/v) final for 1 h at 4°C. After centrifugation (20 800 g for 15 min) and removal of the supernatant, the resulting pellet was washed in ice-cold Acetone over-night. The samples were centrifuged again, the supernatant was discarded and the pellet was air-dried and resuspended in 1x SDS loading dye.

Secreted proteins were analysed by SDS–PAGE; in each case, proteins secreted by 3×10^8 bacteria were loaded per lane. Detection of specific secreted proteins by immunoblotting was performed using 12.5% SDS–PAGE gels. For detection of proteins in total cells, 2×10^8 bacteria were loaded per lane, if not stated otherwise, and proteins were separated on 12.5% SDS–PAGE gels before detection by immunoblotting. Immunoblotting was carried out using anti-Myc (Santa Cruz) antibody.

Western blotting of T3SS translocated proteins from infected cells. HeLa cells in 6-well plates were infected at an MOI of 100 as described above. In case of coinfection with the TEV protease translocating *Y. enterocolitica* strain, the OD₆₀₀ of the strains was set and the two bacterial suspensions were mixed in a tube at a ratio of 1:1 (if not otherwise indicated) before addition to the cells. At the end of the infection, the cells were washed twice with ice-cold PBS and collected by scraping in a small volume of ice-cold PBS. After centrifugation (16 000 rcf, 5 min, 4°C) the pellet was dissolved in 0.002% digitonin supplemented with a protease inhibitor cocktail (Roche complete, Roche). The dissolved pellets were incubated for 5 minutes on ice and then centrifuged (16 000 rcf, 25 min, 4°C). Supernatants were collected and analyzed for total protein content by Bradford BCA assay (Pierce) before SDS PAGE and Western blotting using an anti-Myc (Santa Cruz, 9E11) or anti-Ink4C (Cell Signaling) antibody.

Immunofluorescence. Cell seeded in 96-well plates (Corning) were infected as described above and after fixation with 4% PFA the cells were washed three times

with PBS. The wells were then blocked using 5% goat serum in PBS 0.3% Triton X-100 for 1h at RT. The primary antibody (anti-Myc, Santa Cruz, 1:100) was diluted in PBS with 1% BSA and 0.3% Triton X-100 and cells were incubated overnight at 4°C. Cells were washed 4 times with PBS before the secondary antibody (AF 488 anti-mouse, life technologies, 1:250) diluted in PBS with 1% BSA and 0.3% Triton X-100 was added. If needed Hoechst DNA staining (life technologies, 1:2500) and/or actin staining (Dy647-Phalloidin, DyeOmics) were included. In some cases only the DNA and/or actin stain was applied directly after washing the PFA off. Cells were incubated for 1h at RT, washed three times with PBS and analyzed by automated image analysis as described below.

Automated Microscopy and Image Analysis. Images were automatically acquired with an ImageXpress Micro (Molecular devices, Sunnyvale, USA). Quantification of anti-Myc staining intensities was performed using MetaXpress (Molecular devices, Sunnyvale, USA). Regions within cells excluding nuclear regions and regions containing bacteria were manually chosen (circles with an area of 40 pixels) and average intensity was recorded.

TNF α stimulation and Western blotting of phospho-p38. HeLa cells seeded in 6-well plates were infected with an MOI of 100 as described above. 30min p.i. Gentamicin was added and 45 min p.i. TNF α was added (10 ng/ml). 1 h 15min p.i. cells were washed twice with ice-cold PBS and Phospho-safe lysis buffer (Novagen) was added to lyse the cells. After incubation on ice, the cells were centrifuged (16 000 rcf, 25 min, 4°C). Supernatants were collected and analyzed for total protein content by Bradford BCA assay (Pierce) before SDS PAGE and Western blotting using an anti-Phospho-p38, total p38 antibodies (Cell Signaling) and anti-Actin antibody (Millipore).

cAMP level determination of infected HeLa cells. HeLa cells seeded in 96-well plates were infected as described above. 30 min before the infection cDMEM was changed to DMEM supplemented with 10 mM HEPES and 2 mM L-glutamine and 100 μ M 3-Isobutyl-1-methylxanthin (IBMX, Sigma Aldrich). 60 min p.i. Gentamicin was added and cells were further incubated at 37°C for another 90 min. Determination

of cAMP was performed using a competitive ELISA according to the manufacturers instructions (Amersham, cAMP Biotrak, RPN225). As a positive control indicated amount of cholera toxin (C8052, Sigma Aldrich) was added for 1 h to cells in DMEM supplemented with 10 mM HEPES and 2 mM L-glutamine and 100uM IBMX.

Zebrafish embryo infections, imaging and automated image quantification. All animal experiments were performed according to approved guidelines. Zebrafish were maintained at standard conditions [42]. Embryos were staged by hours postfertilization (hpf) at 28.5°C [43]. The following zebrafish lines were used in this study: wild type fish (AB/EK and EK/TL). Infection protocol followed guidelines given in [44]. 12 hpf embryos were maintained in E3 medium containing 0.2 mM N-phenylthiourea (PTU) to prevent pigment formation. 2 days postfertilization (dpf) embryos were anesthetized by 0.2 mg/ml Tricaine and aligned on 1% agar plates in E3 using a hair loop tool [44]. *Y. enterocolitica* were grown in BHI supplemented with 0.4% Arabinose and antibiotics and mDap overnight at RT, diluted in fresh BHI with 0.5% Arabinose and other additives to an OD₆₀₀ of 0.2 and grown for 2h at RT before a temperature shift to a 37°C waterbath shaker for further 45 min. Finally, the bacteria were collected by centrifugation (6000 rcf, 30 sec) and washed once with PBS. The OD₆₀₀ was set to 2 in PBS containing mDAP. 1-2 nL of this suspension were injected into the hindbrain of aligned zebrafish embryos using an Femtojet Microinjector (Eppendorf) using Femtotips II (Eppendorf), where the tip of the needle had been broken off with fine tweezers. The injection time was set to 0.2 s and the compensation pressure to 15 hPa (Eppendorf, Femtojet) and the injection pressure was adjusted between 600 and 800 hPa. Drop size and thus the inoculum was checked by microscopy and by control plating. Following microinjection the fish were collected in E3 containing Tricaine and PTU and incubated for 30 min at 37°C and incubated for further 5 h at 28°C. A fluorescence binocular (Leica) was used to observe bacterial EGFP fluorescence 1h post infection in zebrafish hindbrains, and embryos that are not properly injected were discarded. At the end of the infection, fish were fixed with 2% ice-cold PFA for 1h on ice and further with fresh ice-cold PFA overnight at 4°C. Antibody staining was performed as described previously [45,46]. Briefly, embryos were washed 4 times with PBS 0.1%Tween for 5 min each wash and permeabilized with PBS-T + 0.5% Triton X-100 for 30 min at RT. Embryos were blocked in blocking solution (PBS 0.1% Tween 0.1% TritonX-100 5% goat serum

and 1% BSA) at 4°C overnight. Antibody (Cleaved Caspase-3 (Asp175), Cell Signaling) was diluted 1:100 in blocking solution and incubated under shaking at 4°C in the dark. Fish were washed 7 times with PBS 0.1% Tween for 30 min before the secondary antibody (goat anti-rabbit AF647, Invitrogen, 1:500) diluted in blocking solution was added and incubated at 4°C overnight. Larvae were washed with PBS 0.1% Tween four times 30 min at 4°C and once overnight and further washed 3-4 times. Images were taken with Leica TCS SP5 confocal microscope using a 40× water immersion objective. Images were analyzed using Imaris (Bitplane) and Image J software (<http://imagej.nih.gov/ij/>).

Image analysis (on n=14 for pBad_Si2 or n=19 for z-BIM) was performed via CellProfiler [47] on maximum intensity z projections of recorded z-stack images. Briefly, bacteria were detected via the GFP channel. Around each area of a bacterial spot a circle with a radius of 10 pixels was created. Overlapping regions were separated equally among the connecting members. In those areas closely surrounding bacteria, the Caspase 3 p17 staining intensity was measured.

Sample Preparation for Phosphoproteomics. For each condition, two 6-well plates of HeLa CCL-2 cells were grown to confluency. Cells were infected for 30 min as described above. At the indicated time-points, the plates were put on ice and washed twice with ice-cold PBS. Samples were then collected in urea solution [8 M Urea (AppliChem), 0.1 M Ammoniumbicarbonate (Sigma), 0.1% RapiGest (Waters), 1× PhosSTOP (Roche)]. The samples were briefly vortexed, sonicated at 4 °C (Hielscher), shaken for 5 min on a thermomixer (Eppendorf) and centrifuged for 20 min at 4 °C and 16'000g. Supernatants were collected and stored at −80 °C for further processing. BCA Protein Assay (Pierce) was used to measure protein concentration.

Phosphopeptide Enrichment. Disulfide bonds were reduced with tris(2-carboxyethyl)phosphine at a final concentration of 10 mM at 37 °C for 1 h. Free thiols were alkylated with 20 mM iodoacetamide (Sigma) at room temperature for 30 min in the dark. The excess of iodoacetamide was quenched with N-acetyl cysteine at a final concentration of 25 mM for 10 min at room temperature. Lys-C endopeptidase (Wako) was added to a final enzyme/protein ratio of 1:200 (w/w) and incubated for 4 h at 37 °C. The solution was subsequently diluted with 0.1 M ammoniumbicarbonate (Sigma) to a

final concentration below 2 M urea and digested overnight at 37 °C with sequencing-grade modified trypsin (Promega) at a protein-to-enzyme ratio of 50:1. Peptides were desalted on a C18 Sep-Pak cartridge (Waters) and dried under vacuum. Phosphopeptides were isolated from 2 mg of total peptide mass with TiO₂ as described previously [48]. Briefly, dried peptides were dissolved in an 80% acetonitrile (ACN)–2.5% trifluoroacetic acid (TFA) solution saturated with phthalic acid. Peptides were added to the same amount of equilibrated TiO₂ (5-µm bead size, GL Sciences) in a blocked Mobicol spin column (MoBiTec) that was incubated for 30 min with end-over-end rotation. The column was washed twice with the saturated phthalic acid solution, twice with 80% ACN and 0.1% TFA, and finally twice with 0.1% TFA. The peptides were eluted with a 0.3 M NH₄OH solution. The pH of the eluates was adjusted to be below 2.5 with 5% TFA solution and 2 M HCl. Phosphopeptides were again desalted with microspin C18 cartridges (Harvard Apparatus).

LC-MS/MS analysis. Chromatographic separation of peptides was carried out using an EASY nano-LC system (Thermo Fisher Scientific), equipped with a heated RP-HPLC column (75 µm x 45 cm) packed in-house with 1.9 µm C18 resin (Reprosil-AQ Pur, Dr. Maisch). Aliquots of 1 µg total phosphopeptide sample were analyzed per LC-MS/MS run using a linear gradient ranging from 98% solvent A (0.15% formic acid) and 2% solvent B (98% acetonitrile, 2% water, 0.15% formic acid) to 30% solvent B over 120 minutes at a flow rate of 200 nl/min. Mass spectrometry analysis was performed on a dual pressure LTQ-Orbitrap mass spectrometer equipped with a nanoelectrospray ion source (both Thermo Fisher Scientific). Each MS1 scan (acquired in the Orbitrap) was followed by collision-induced dissociation (CID, acquired in the LTQ) of the 20 most abundant precursor ions with dynamic exclusion for 30 seconds. For phosphopeptide analysis the 10 most abundant precursor ions were subjected to CID with enabled multistage activation. Total cycle time was approximately 2 s. For MS1, 10⁶ ions were accumulated in the Orbitrap cell over a maximum time of 300 ms and scanned at a resolution of 60,000 FWHM (at 400 m/z). MS2 scans were acquired using the normal scan mode, a target setting of 10⁴ ions, and accumulation time of 25 ms. Singly charged ions and ions with unassigned charge state were excluded from triggering MS2 events. The normalized collision energy was set to 32%, and one microscan was acquired for each spectrum.

Label-free quantification and database searching. The acquired raw-files were imported into the Progenesis software tool (Nonlinear Dynamics, Version 4.0) for label-free quantification using the default parameters. MS2 spectra were exported directly from Progenesis in mgf format and searched using the MASCOT algorithm (Matrix Science, Version 2.4) against a decoy database [49] containing normal and reverse sequences of the predicted SwissProt entries of *Homo sapiens* (www.ebi.ac.uk, release date 16/05/2012) and commonly observed contaminants (in total 41,250 sequences) generated using the SequenceReverser tool from the MaxQuant software (Version 1.0.13.13). To identify proteins originating from *Y. enterocolitica*, non phosphopeptide enriched samples were searched against the same database above including predicted SwissProt entries of *Y. enterocolitica* (www.ebi.ac.uk, release date 15/08/2013). The precursor ion tolerance was set to 10 ppm and fragment ion tolerance was set to 0.6 Da. The search criteria were set as follows: full tryptic specificity was required (cleavage after lysine or arginine residues unless followed by proline), 2 missed cleavages were allowed, carbamidomethylation (C) was set as fixed modification and phosphorylation (S,T,Y) or oxidation (M) as a variable modification for TiO2 enriched or not enriched samples, respectively. Finally, the database search results were exported as an xml-file and imported back to the Progenesis software for MS1 feature assignment. For phosphopeptide quantification, a csv-file containing the MS1 peak abundances of all detected features was exported and for not enriched samples, a csv-file containing all protein measurements based on the summed feature intensities of all identified peptides per protein was created. Importantly, the Progenesis software was set that proteins identified by similar sets of peptides are grouped together and that only non-conflicting peptides with specific sequences for single proteins in the database were employed for protein quantification. Both files were further processed using the in-house developed SafeQuant v1.0 R script (unpublished data, available at <https://github.com/cahrne/SafeQuant/>). In brief, the software sets the identification level False Discovery Rate to 1% (based on the number of decoy protein sequence database hits) and normalizes the identified MS1 peak abundances (Extracted Ion Chromatogram, XIC) across all samples, *i.e.* the summed XIC of all confidently identified peptide features is scaled to be equal for all LC-MS runs. Next, all quantified phosphopeptides/proteins are assigned an abundance ratio for each time point, based on the median XIC per time point. The statistical significance of each

ratio is given by its q-value (False Discovery Rate adjusted p-values), obtained by calculating modified t-statistic p-values [50] and adjusting for multiple testing [51]. The location of the phosphorylated residues was automatically assigned by MASCOT (score >10). All annotated spectra together with the MS raw files and search parameters employed, will be deposited to the ProteomeXchange Consortium (<http://proteomecentral.proteomexchange.org>) via the PRIDE partner repository [52]. **Sequence alignment** was performed using EMBL-EBI web based ClustalW2 multiple sequence alignment tool at <http://www.ebi.ac.uk/Tools/msa/clustalw2/>.

Biodistribution in 4T1 tumor allograft mouse models

All animal experiments were approved (license 1908; Kantonales Veterinäramt Basel-Stadt) and performed according to local guidelines (Tierschutz-Verordnung, Basel-Stadt) and the Swiss animal protection law (Tierschutz-Gesetz). 6 week old BALB/c mice were ordered from Janvier Labs. After at least one week of accommodation, mice were anesthetized using isoflurane and 100 μ l 4T1 cells (1×10^5 - 1×10^6 cells) were subcutaneously injected into the flank of BALB/c mice. Throughout the experiment, mice were scored for behavior and physical appearance, and surface temperature, as well as body weight was measured.

Once tumors had developed, mice were administered an 8 mg/ml desferal solution (10 ml/kg) through i.p. injection. On the following day, mice were infected with *Y. enterocolitica* MRS40 or *Y. enterocolitica* MRS40 Δ HOPEMT (2×10^5 , 1×10^6 or 1×10^7 bacteria) by injection into the tail vein. The inoculum i.v. administered to the mice was validated by dilution plating. In some experiments, tumor progression was followed by daily measurements of tumor length and width with digital calipers. Tumor volume was determined as $0.523 \times \text{length} \times \text{width}^2$. On respective days postinfection, mice were sacrificed by CO₂ inhalation. A blood sample was immediately isolated through aspiration from the heart. Liver, spleen, lung and the tumor were isolated and their weight determined. The organs and the tumor were homogenized. CFU in each sample was determined by spotting of serial dilutions onto LB agar plates containing nalidixic acid (35 μ g/ml).

B) RESULTS

A protein delivery system based on type 3 secretion of YopE fusion proteins

While the very N-terminus of the *Y. enterocolitica* T3SS effector YopE (SEQ ID No. 1) contains the secretion signal sufficient to translocate heterologous proteins [26], the chaperone-binding site (CBS) for its chaperone (SycE) is not included [53]. We selected the N-terminal 138 amino acids of YopE (SEQ ID No. 2) to be fused to proteins to be delivered, as this had been shown to give best results for translocation of other heterologous T3S substrates [28]. As these N-terminal 138 amino acids of YopE contain the CBS, we further decided to coexpress SycE. The SycE-YopE₁₋₁₃₈ fragment cloned from purified *Y. enterocolitica* pYV40 virulence plasmid contains the endogenous promoters of YopE and of its chaperone SycE (Fig. 10). Therefore, SycE and any YopE₁₋₁₃₈ fusion protein are induced by a rapid temperature shift from growth at RT to 37°C. Culture time at 37°C will affect fusion protein amount present in bacteria. A multiple cloning site (MCS) was added at the 3' end of YopE₁₋₁₃₈ (Fig. 10 B) followed by a Myc and a 6xHis tag and a Stop codon.

The background strain was carefully selected. First, to limit the translocation of endogenous effectors, we used a *Y. enterocolitica* strain that was deleted for all known effectors, Yop H, O, P, E, M and T (named Δ HOPEMT) [54]. In addition, we occasionally used an auxotroph mutant that cannot grow in absence of exogenous *meso*-2,6-diaminopimelic acid [55]. This strain was deleted for the aspartate-beta-semialdehyde dehydrogenase gene (Δ asd), and classified as biosafety level 1 by the Swiss safety agency (amendment to A010088/2). In addition, we deleted the adhesion proteins YadA and/or InvA to offer a larger choice of background strains. While the use of the *yadA* or *yadA/invA* strains reduce the background signalling induced [56], the delivered protein amount is affected as well [57].

Characterization of YopE fusion protein delivery into eukaryotic cells

In an *in-vitro* secretion assay (see Fig. 1 A), protein secretion into the surrounding liquid is artificially induced. After TCA based protein precipitation, Western blot analysis with anti-YopE antibody was used to determine protein amounts secreted (Fig. 1 B). While a wt strain secreted full length YopE, the Δ HOPEMT *asd* strains did not. Upon presence of YopE₁₋₁₃₈-Myc-His (further termed YopE₁₋₁₃₈-Myc; SEQ_ID_No._3) a smaller YopE band became visible (Fig. 1 B). Hence, the YopE₁₋

138 fragment is well secreted in the set up described here. To analyze homogeneity of protein translocation into eukaryotic cells, we infected HeLa cells with the YopE₁₋₁₃₈-Myc encoding strain and stained the Myc tag by IF (Fig. 2 A and B). While in the beginning only the bacteria were stained, at 30 min post infection (p.i.) cell outlines start to be visible, which is enhanced upon increased infection time (Fig. 2 B). This trend is well reflected by the Myc tag staining intensity inside HeLa cells (Fig. 2 A and B). The YopE₁₋₁₃₈-Myc can be detected everywhere in the cells (Fig. 2 A), except in the nuclei [58]. Remarkably, most if not all cells were reached by this approach in a comparable way. As *Y. enterocolitica* is known to infect many different cell types [59], we followed YopE₁₋₁₃₈-Myc delivery into various cell lines. The same homogenous anti-Myc IF staining was observed in infected murine fibroblasts, Jurkat cells and HUVECs (Fig. 11). Even more, tuning the MOI up or down allows modulating the protein amount delivered (Fig. 2 C), while still most of the cells remain targeted. A low bacterial number will not result in few cells with lots of delivered protein but rather with most cells containing a low amount of delivered protein (Fig. 2 C).

Redirection of T3SS delivered proteins to the nucleus

As YopE itself localized to the cytoplasm (Fig. 2 A), it is of special interest to test if the YopE₁₋₁₃₈ fragment hampers localization of nuclear fusion proteins. We therefore added the SV40 NLS to the C-terminus (and N-terminus, similar results) of YopE₁₋₁₃₈-EGFP (SEQ ID No. 39 and SEQ ID No. 38, respectively). While YopE₁₋₁₃₈-EGFP (SEQ ID No. 37) led to a weak cytoplasmic staining, YopE₁₋₁₃₈-EGFP-NLS gave rise to a stronger nuclear EGFP signal in HeLa cells infected (Fig. 3). This indicates that the YopE₁₋₁₃₈ fragment is compatible with the use of an NLS. While mCherry had already been used in plant pathogens [60], this represents a successful delivery of a GFP-like protein via human or animal pathogenic bacteria encoding a T3SS. This validates the SycE and YopE₁₋₁₃₈ dependent strategy to be very promising for delivery of many proteins of choice.

Removal of the YopE₁₋₁₃₈ appendage after translocation of the fusion protein to the eukaryotic cell

While for bacterial delivery the YopE₁₋₁₃₈ fragment is of great benefit, it might hamper the fusion proteins function and/or localization. Therefore, its removal after

protein delivery would be optimal. To this end, we introduced two TEV cleavage sites (ENLYFQS) [61-63] in between YopE₁₋₁₃₈ and a fusion partner (the transcriptional regulator ET1-Myc (SEQ ID No. 36 and 41) [64] and human INK4C (SEQ ID No. 40 and SEQ ID No. 43)). To keep the advantages of the presented method, we further fused the TEV protease (S219V variant; [65]) to YopE₁₋₁₃₈ (SEQ ID No. 42) in another *Y. enterocolitica* strain. HeLa cells were infected with both strains at once. To allow analysis of the translocated fraction of proteins only, infected HeLa cells were lysed at 2 h p.i. (Fig. 4) with Digitonin, which is known not to lyse the bacteria ([66]; see Fig. 12 for control). Western blot analysis revealed the presence of the YopE₁₋₁₃₈-2xTEV-cleavage-site-ET1-Myc or YopE₁₋₁₃₈-2xTEV-cleavage-site-Flag-INK4C-Myc only when cells had been infected with the corresponding strain (Fig. 4 A and C). Upon overnight digestion of this cell-lysate with purified TEV protease, a shifted band could be observed (Fig. 4 A and C). This band corresponds to ET1-Myc (Fig. 4 C) or Flag-INK4C (Fig. 4 A) with the N-terminal remnants of the TEV cleavage site, most likely only one Serine. Upon coinfection of cells with the strain delivering the TEV protease, the same cleaved ET1-Myc or Flag-INK4C fragment became visible, indicating that the TEV protease delivered via T3SS is functional and that single cells had been infected by both bacterial strains (Fig. 4 A and C). While cleavage is not complete, the majority of translocated protein is cleaved already 2h post infection and even over-night digestion with purified TEV protease did not yield better cleavage rates (Fig. 4 B). As reported, TEV protease dependent cleavage might need optimization dependent on the fusion protein [67,68]. TEV protease dependent removal of the YopE₁₋₁₃₈ appendage after translocation hence provides for the first time a T3SS protein delivery of almost native heterologous proteins, changing the amino acid composition by only one N-terminal amino acid.

An alternative approach to the TEV protease dependent cleavage of the YopE fragment consisted in incorporating Ubiquitin into the fusion protein of interest. Indeed, Ubiquitin is processed at its C-terminus by a group of endogenous Ubiquitin-specific C-terminal proteases (Deubiquitinating enzymes, DUBs). As the cleavage is supposed to happen at the very C-terminus of Ubiquitin (after G76), the protein of interest should be free of additional amino acid sequence. This method was tested on the YopE₁₋₁₃₈-Ubiquitin-Flag-INK4C-MycHis fusion protein. In control cells infected by YopE₁₋₁₃₈-Flag-INK4C-MycHis-expressing bacteria, a band corresponding to YopE₁₋₁₃₈-Flag-INK4C-MycHis was found, indicative of efficient

translocation of the fusion protein (Figure 24). When cells were infected for 1h with YopE1-138-Ubiquitin-Flag-INK4C-MycHis-expressing bacteria, an additional band corresponding to the size of Flag-INK4C-MycHis was visible, indicating that part of the fusion protein was cleaved. This result shows that the introduction of Ubiquitin into the fusion protein enables to cleave off the YopE1-138 fragment without a need for an exogenous protease.

Translocation of type III and type IV bacterial effectors

SopE from *Salmonella enterica* is a well-characterized guanine nucleotide exchange factor (GEF) that interacts with Cdc42, promoting actin cytoskeletal remodeling [69]. Whereas the translocation of YopE₁₋₁₃₈-Myc into HeLa cells has no effect, translocated YopE₁₋₁₃₈-SopE (SEQ ID No. 5 and 135) induced dramatic changes in the actin network (Fig. 5 A). Similar results were obtained with another GEF effector protein, IpgB1 from *Shigella flexneri* (SEQ ID No. 4). Remarkably, first changes in the actin cytoskeleton were observed as fast as 2 min p.i. (Fig. 5 A). Therefore, one can conclude that T3SS dependent protein delivery happens immediately after infection is initiated by centrifugation. To proof strict T3SS dependent transport, one of the T3SS proteins forming the translocation pore into the eukaryotic cell membrane was deleted (YopB, see [70]) (Fig. 12).

During *Salmonella* infection, SopE translocation is followed by translocation of SptP, which functions as a GTPase activating protein (GAP) for Cdc42 [71]. Whereas the translocation of YopE₁₋₁₃₈-SopE-Myc (SEQ ID No. 135) alone triggered massive F-actin rearrangements, the co-infection with YopE₁₋₁₃₈-SptP (SEQ ID No. 8) expressing bacteria abolished this effect in a dose dependent manner (Fig. 5 B). An anti-Myc staining indicated that this inhibition was not due to a reduced level of YopE₁₋₁₃₈-SopE-Myc translocation (Fig. 5 B). Together these results showed that the co-infection of cells with two bacterial strains is a valid method to deliver two different effectors into single cells to address their functional interaction.

The *S. flexneri* type III effector OspF functions as a phosphothreonine lyase that dephosphorylates MAP kinases p38 and ERK [72]. To test the functionality of translocated YopE₁₋₁₃₈-OspF (SEQ ID No. 7), we monitored the phosphorylation of p38 after stimulation with TNF α . In uninfected cells or in cells infected with YopE₁₋₁₃₈-Myc expressing bacteria, TNF α induced p38 phosphorylation. In contrast, after

translocation of YopE₁₋₁₃₈-OspF, TNF α -induced phosphorylation was abolished, showing that the delivered OspF is active towards p38 (Fig. 6 A).

During *Salmonella* infection, the type III effector SopB protects epithelial cells from apoptosis by sustained activation of Akt [73]. Whereas the translocation of YopE₁₋₁₃₈-Myc or YopE₁₋₁₃₈-SopE had no effect on Akt, the translocation of YopE₁₋₁₃₈-SopB (SEQ ID No. 6) induced a strong phosphorylation of Akt at T308 and S473, reflecting the active form (Fig. 6 B). Similar results were obtained with the SopB-homolog from *S. flexneri* (IpgD, SEQ ID No. 9). Altogether, our results show that the YopE₁₋₁₃₈-based delivery system functions for all T3S effectors tested so far, and that it allows investigating proteins involved in the control of central cellular functions including the cytoskeleton, inflammation and cell survival.

A number of bacteria, including *Agrobacterium tumefaciens*, *Legionella pneumophila* and *Bartonella henselae*, use type IV secretion to inject effectors into cells. We tested whether the type IV effector BepA from *B. henselae* could be translocated into HeLa cells using our tool. Full length BepA (SEQ ID No. 10) and BepA_{E305-end} (SEQ ID No. 11) containing the C-terminal Bid domain, were cloned and cells were infected with the respective strains. As BepA was shown to induce the production of cyclic AMP (cAMP) [74], the level of cAMP in HeLa cells was measured after infection. Whereas the translocation of the Bid domain of the *B. henselae* effector BepG (SEQ ID No. 136) failed to induce cAMP, full length BepA and BepA_{E305-end} triggered cAMP production in expected amounts [74] (Fig. 6 C). This result shows, that type IV effectors can also be effectively delivered by the YopE₁₋₁₃₈-based delivery system into host cell targets and that they are functional.

Translocation of eukaryotic proteins into epithelial cells

To show that human proteins can translocate via type III secretion we fused human apoptosis inducers for delivery by *Y. enterocolitica* to YopE₁₋₁₃₈ or for delivery by *S. enterica* to SteA₁₋₂₀, SteA, SopE₁₋₈₁ or SopE₁₋₁₀₅. We then monitored the translocation of the human BH3 interacting-domain death agonist (BID, SEQ ID No. 24), which is a pro-apoptotic member of the Bcl-2 protein family. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8). CASP8 cleaves BID, and the truncated BID (tBID, SEQ ID No. 25) translocates to mitochondria where it triggers cytochrome c release. The latter leads to the intrinsic mode of caspase 3 (CASP3) activation during

which it is cleaved into 17 and 12 kDa subunits [75]. Whereas infection for 1 h with YopE₁₋₁₃₈-Myc or YopE₁₋₁₃₈-tBID expressing *Y. enterocolitica* failed to induce apoptosis, the translocation of human tBID triggered cell death in larger extend than the well-characterized apoptosis inducer staurosporin (Fig. 7 A and C). As expected, the translocation of tBID lead to the production of CASP3 p17 subunit, even in larger amounts as with staurosporin (Fig. 7 A). To be able to compare translocated protein amounts to endogenous Bid, HeLa cells were lysed with Digitonin and analyzed by Western blotting using an anti Bid antibody (Fig. 7 B). T3SS delivered YopE₁₋₁₃₈-tBID reached about endogenous Bid levels in HeLa cells, while delivered YopE₁₋₁₃₈-BID was present in even higher quantities (2.5 fold) (Fig. 7 B). A deep proteome and transcriptome mapping of HeLa cells estimated 4.4 fold 10⁵ copies of BID per single cell [76]. Therefore, one can conclude that T3SS dependent human protein delivery reaches 10⁵ to 10⁶ proteins per cell. These numbers fit the copies per cell of nanobodies translocated via *E. coli* T3SS [4]. Assuming a levelling of a factor of 10 for the MOI and for the duration of the infection, a factor of 3.2 for the time-point of antibiotic addition and for the culture time at 37°C before infection, the delivered protein copies/cell can be tuned from some 1000 copies/cell up to some 10⁶ copies/cell Altogether, these results indicated that translocated tBID was functional and delivered at relevant levels. This validated the translocation tool to study the role of proteins in the regulation of apoptosis, a central aspect of cell biology.

We further fused murine tBID (codon optimized for *Y. enterocolitica*; SEQ ID No. 194) or the BH3 domains of murine tBID or murine BAX (in both cases codon optimized for *Y. enterocolitica*; SEQ ID No. 138 and 139) to YopE₁₋₁₃₈ for delivery by *Y. enterocolitica*. Whereas infection for 2.5 h with *Y. enterocolitica* ΔHOPEMT *asd* delivering no protein or YopE₁₋₁₃₈-Myc failed to induce apoptosis, the translocation of murine tBID (codon optimized to *Y. enterocolitica*, SEQ ID No. 194) triggered cell death in B16F10 (Fig. 16), D2A1 (Fig. 17), HeLa (Fig. 18) and 4T1 (Fig. 19) cells. The translocation of the BH3 domain of murine BID codon optimized for *Y. enterocolitica* (SEQ ID 138) or murine BAX codon optimized for *Y. enterocolitica* (SEQ ID 139) were as well found to induce massive cell death in B16F10 (Fig. 16), D2A1 (Fig. 17), HeLa (Fig. 18) and 4T1 (Fig. 19) cells.

Whereas infection for 4 h with *S. enterica aroA* bacteria failed to induce apoptosis, the translocation of murine tBID triggered apoptosis, as the translocation of murine

tBID lead to the production of CASP3 p17 subunit (Figs. 20 and 21). The extent of apoptosis induction for SopE fusion proteins was larger when using Spil T3SS inducing conditions (Fig. 20), which reflects the transport of SopE exclusively by Spil T3SS. SteA₁₋₂₀ fused murine tBID failed to induce apoptosis, very likely because the secretion signal within the 20 N-terminal amino acids of SteA is not sufficient to allow delivery of a fusion protein (Figs. 20 and 21). Murine tBID fused to full length SteA lead to apoptosis induction in HeLa cells (Figs. 20 and 21), both in Spil and SpilI T3SS inducing conditions, reflecting the ability of SteA to be transported by both T3SS. It has to be noted that even under SpilI T3SS inducing conditions, a partial activity of the Spil T3SS is expected as seen by the activity of SopE fusion proteins in SpilI T3SS inducing conditions (Fig. 21).

Besides the here functionally elaborated translocated eukaryotic proteins, several other eukaryotic proteins have been secreted using the here-described tool. This includes for delivery by *Y. enterocolitica* (Figs. 13, 14 and 23) proteins from cell cycle regulation (Mad2 (SEQ ID No. 15), CDK1 (SEQ ID No. 14), INK4A (SEQ ID No. 16), INK4B (SEQ ID No. 17) and INK4C (SEQ ID No. 18)) as well as parts thereof (INK4A 84-103 (SEQ ID No. 158), p107 657-662 (SEQ ID No. 159), p21 141-160 (SEQ ID No. 160), p21 145-160 (SEQ ID No. 161), p21 17-33 (SEQ ID No. 162) and cyclin D2 139-147 (SEQ ID No. 163)), apoptosis related proteins (Bad (SEQ ID No. 29), FADD (SEQ ID No. 28), and Caspase 3 p17 (SEQ ID No. 22) and p12 (SEQ ID No. 23), zebrafish Bid (SEQ ID No. 19) and t-Bid (SEQ ID No. 20)) as well as parts thereof (tBid BH3 (SEQ ID No. 138), Bax BH3 (SEQ ID No. 139)), signalling proteins (murine TRAF6 (SEQ ID No. 12), TIFA (SEQ ID No. 13)), GPCR α subunit (GNA12, shortest isoform, (SEQ ID No. 30)), nanobody (vhhGFP4, (SEQ ID No. 31)) and nanobody fusion constructs for targeted protein degradation (Slmb-vhhGFP4; (SEQ ID Nos. 32, 33, 34) [77]) (Fig. 13 and 14) as well as small GTPases (Rac1 Q61E (SEQ ID No. 26 and 137) and RhoA Q63L (SEQ ID No. 27) and Pleckstrin homology domain from human Akt (SEQ ID No. 35). Besides the functionally elaborated apoptosis related proteins (murine tBid, SEQ ID No. 144-147), this further includes for delivery by *S. enterica* (Fig. 22) proteins from cell cycle regulation (Mad2 (SEQ ID No. 168-169), CDK1 (SEQ ID No. 170-171), INK4A (SEQ ID No. 164-165) and INK4C (SEQ ID No. 166-167)). While those proteins

have not been functionally validated, the possibility of T3SS dependent secretion of diverse eukaryotic proteins in combination with the possible removal of the YopE appendage opens up new vistas on the broad applicability of T3SS in cell biology and therapeutic applications.

In vivo translocation of truncated Bid in zebrafish embryos induces apoptosis

An interesting feature of this bacterial tool is the potential use in living animals. Zebrafish in their embryonic state can be kept transparent allowing fluorescent staining and microscopy [44,78,79]. Few zebrafish apoptosis inducers have been described in detail, whereof z-BIM is the most potent [80]. Therefore, we decided to clone z-BIM into our system. Even if weakly homologous to human BIM, we assayed the potency of apoptosis induction of YopE₁₋₁₃₈-z-BIM (SEQ ID No. 21) in human epithelial cells. HeLa cells infected for 1h with the strain translocating YopE₁₋₁₃₈-z-BIM showed clear signs of cell death. We then performed *in-vivo* experiments with 2 days post fertilization (dpf) zebrafish embryos, using a localized infection model via microinjection of bacteria into the hindbrain [44]. After infection for 5.5 h the fish were fixed, permeabilized and stained for presence of CASP3 p17. Upon infection with the YopE₁₋₁₃₈-Myc expressing strain, bacteria were visible in the hindbrain region (staining “b”, Fig. 8 A I) but no induction of apoptosis around the bacteria was detected (staining “c”, Fig. 8 A I). In contrast, upon infection with the strain delivering YopE₁₋₁₃₈-z-BIM a strong increase in presence of cleaved CASP3 was observed in regions surrounding the bacteria (Fig. 8 A II). Automated image analysis on maximum intensity z projections confirms that YopE₁₋₁₃₈-z-BIM translocating bacteria induce apoptosis in nearby cells by far more than control bacteria do (Fig. 8 B). This indicates that z-BIM is functional in zebrafish upon bacterial translocation. These results further validate the use of T3SS for eukaryotic protein delivery in living animals.

Phosphoproteomics reveal the global impact of translocated proteins on protein phosphorylation

Phosphorylation is a wide-spread post-translational modification which can either activate or inactivate biological processes and is therefore a suitable target to study signaling events [81,82]. Despite this, no systems-level analysis of phosphorylation in apoptosis is available today. To analyze the impact of human tBid delivered into

HeLa cells, we used a label-free phosphoproteomic approach by LC-MS/MS. In three independent experiments, cells were either left untreated, infected with $\Delta HOPEMT$ *asd* + YopE₁₋₁₃₈-Myc or with $\Delta HOPEMT$ *asd* + YopE₁₋₁₃₈-tBid for 30 minutes. Cells were lysed, followed by enzymatic digestion, phosphopeptide enrichment and quantification and identification of individual phosphopeptides. We compared cells infected with $\Delta HOPEMT$ *asd* + YopE₁₋₁₃₈-Myc to cells infected with $\Delta HOPEMT$ *asd* + YopE₁₋₁₃₈-tBid, allowing us to identify 363 tBid dependent phosphorylation events. 286 phosphopeptides showed an increase in phosphorylation whereas 77 were less phosphorylated upon tBid delivery, corresponding to 243 different proteins, which we defined as the tBid phosphoproteome. The STRING database was used to create a protein-protein interaction network of the tBid phosphoproteome [83] (Fig. 9 A). Additionally 27 proteins known to be related to mitochondrial apoptosis were added to the network, building a central cluster. Interestingly, only few proteins from the tBid phosphoproteome are connected to this central cluster indicating that many proteins undergo a change in phosphorylation that were so far not directly linked to apoptotic proteins. To characterize the biological functions covered by the tBid phosphoproteome, we performed a gene ontology analysis using the functional annotation tool of the Database for Annotation, Visualization, and Integrated Discovery (DAVID, <http://david.abcc.ncifcrf.gov/>) [84,85]. Identified biological functions show that diverse cellular processes are affected by tBid. Many proteins involved in chromatin rearrangement and the regulation of transcription undergo a change in phosphorylation (i.e. CBX3, CBX5, TRIM28, HDAC1). HDAC1 for example is a histone deacetylase playing a role in regulation of transcription. It has been shown that HDAC1 can modulate transcriptional activity of NF- κ B, a protein also participating in apoptosis. We additionally identified a cluster of proteins involved in RNA processing which has previously been shown to play an important role in the regulation of apoptosis [86]. HNRPK for instance mediates a p53/TP53 response to DNA damage and is necessary for the induction of apoptosis [87]. Furthermore, the phosphorylation of proteins involved in protein translation is also affected. Several eukaryotic initiation factors (i.e. EIF4E2, EIF4B, EIF3A, EIF4G2) undergo a change in phosphorylation, which is in line with the observation that overall protein synthesis is decreased in apoptotic cells. Interestingly, the phosphorylation of many proteins involved in cytoskeleton remodeling (e.g. PXN, MAP1B9) are altered upon tBid delivery. This is in concordance with the observation

that the morphology of cells changes dramatically upon tBid delivery (Figure 9 B). Cells shrinkage and loss of contact is reflected by the fact that we observe phosphorylation of adhesion related proteins like ZO2 and Paxillin. Similarly, shrinkage of the nuclei is accompanied by phosphorylation of laminar proteins like LaminA/C and Lamin B1. Altogether, tBID delivery induces a rapid apoptotic response also indicated by rupture of the mitochondrial integrity (Fig. 9 B). We showed that tBid induced apoptosis affects hundreds of phosphorylation events participating in diverse cellular processes. While many identified proteins have been related to apoptosis, only few were known to be phosphorylated upon apoptosis induction. The phosphoproteomic approach thus provides a useful resource for further studies on apoptosis.

Translocation of eukaryotic heterologous fusion proteins consisting of repeated identical or variable protein domains into epithelial cells

To show that heterologous fusion proteins consisting of repeated identical or variable protein domains can translocate via type III secretion we fused murine apoptosis inducers for delivery by *Y. enterocolitica* to YopE₁₋₁₃₈. As control, we fused murine tBID (codon optimized for *Y. enterocolitica*; SEQ ID No. 194) or the BH3 domains of murine tBID or murine BAX (in both cases codon optimized for *Y. enterocolitica*; SEQ ID No. 200 and 201) to YopE₁₋₁₃₈ for delivery by *Y. enterocolitica*. The heterologous fusion protein consisted in one case of murine BH3 domain of tBID fused to itself, resulting in YopE₁₋₁₃₈-(tBID-BH3)₂ (SEQ ID No. 202). In a second case, the heterologous fusion proteins consisted of murine BH3 domain of tBID fused to murine BH3 domain of BAX, resulting in YopE₁₋₁₃₈-(tBID-BH3)-(BAX-BH3) (SEQ ID No. 203). In the case of murine tBID and murine BAX the codon was optimized for *Y. enterocolitica*. Schematic representation of repeated identical domains or combination of different protein domains is shown in Fig. 25.

Whereas infection for 4 h with *Y. enterocolitica* ΔHOPEMT *asd* delivering YopE₁₋₁₃₈-Myc failed to induce apoptosis, the translocation of murine BH3 domain tBID (codon optimized to *Y. enterocolitica*, SEQ ID No. 194) triggered cell death in B16F10 and 4T1 cells (Figs. 26 and 27), with a clear dose-response effect upon increasing multiplicity of infection (MOI). Surprisingly, delivered YopE₁₋₁₃₈-(tBID-BH3)-(BAX-

BH3) or YopE₁₋₁₃₈-(tBID-BH3)₂ were found more active than YopE₁₋₁₃₈-(tBID-BH3) at lower MOI (Figs. 26 and 27). This indicates that upon delivery of repeated identical domains or combination of different protein domains, the impact on a desired cellular pathway as apoptosis can be enlarged.

Generation of enhanced pro-apoptotic bacteria

In above mentioned experiments it is shown that the T3SS-based delivery of pro-apoptotic proteins (e.g. t-BID (SEQ ID No. 25) or BIM (SEQ ID No. 21)) efficiently induces cell death in both murine and human cells, including cancerous cells, and that this effect could be increased when using murine t-BID optimized to the bacterial codon usage (SEQ ID No. 138). This increased cell killing very likely reflects increased amount of protein production and following delivery via T3SS due to optimal codons used.

In order to optimize the delivery of pro-apoptotic proteins, strains transformed with different pro-apoptotic proteins have been generated according to Table IV.

Table IV: Strains transformed with different pro-apoptotic proteins

Strain Name	Background strain	Protein to be delivered by T3SS	Backbone plasmid	Resulting plasmid name	Primers. Si_Nr.:	resistances
YopE1-138- (Y. enterocolitica codon optimized murine tBid BH3 extended part)	Y. enterocolitica Δ yopH,O,P,E,M,T Δ asd	YopE1-138- Y. enterocolitica codon optimized murine tBid BH3 extended (by 4 Aa)	pBad_Si_2	pSi_353	725/726	Nal Amp
YopE1-138- 10 Aa linker- (Y. enterocolitica)	Y. enterocolitica Δ yopH,O,P	YopE1-138- Y. enterocolitica 10 Aa linker-	pBad_Si_2	pSi_354	727/728	Nal Amp

codon optimized murine tBid BH3 part)	E,M,T Δasd	a codon optimized murine tBid BH3				
YopE1-(138- Y. enterocolitica codon optimized murine Bax BH3 part- Y. enterocolitica codon optimized murine tBid BH3 part	Y. enterocolitic a ΔyopH,O,P, E,M,T Δasd	YopE1-138- Y. enterocolitic a codon optimized murine Bax BH3-. enterocolitic a codon optimized murine tBid BH3	pSi_357			
				pSi_374	736/737	Nal Amp

Shortening the delivered proteins to the essential domains required for signaling (SEQ ID No. 138 or 200)) could increase the efficiency of cell killing (**Fig. 28**). Without being bound by theory, this increase in efficacy is likely to be related to increased amount of protein production and following delivery via T3SS due to smaller size of the delivered protein. Introduction of a linker between the YopE part and the BH3 domain of tBID (SEQ ID No. 218) decreased efficacy, as well as extending the BH3 domain by 4 further amino acids (SEQ ID No. 217) (**Fig. 28**).

Additionally, synthetic cargos with repeats of such essential domains (e.g. the BH3 domain of t-BID (SEQ ID No. 202))) or combinations of these essential domains (e.g. the BH3 domain of t-BID and the BH3 domain of BAX (SEQ ID No. 203 and 219)) were generated. Surprisingly, tandem repeats of the same or different BH3 domains were found to result in enhanced apoptosis induction on cancerous cell lines (including 4T1 and B16F10 cells, **Fig. 28**). The IC₅₀ (half maximal inhibitory concentration), referring to the number of bacteria per eukaryotic cell (MOI) needed in order to kill 50% of such cells, was found to be decreased upon delivery of tandem repeats of tBID BH3 domain as compared to a single tBID BH3 domain (**Fig. 28**).

This finding was surprising, as the protein size is increased by fusing as second BH3 domain of t-BID. Due to this, decreased expression and delivery levels of YopE₁₋₁₃₈-(tBID BH3)₂ (SEQ ID No. 202) as compared to YopE₁₋₁₃₈-tBID BH3 (SEQ ID No. 138 or 200) would be expected, and might maximally reach equivalent levels. In order to reach an increase in cell killing activity, the fused tBID BH3 domains must simultaneously act side by side upon delivery by the T3SS into eukaryotic cells. In case only one tBID BH3 domain in the YopE₁₋₁₃₈-(tBID BH3)₂ construct would be functional, at best the same efficiency as with YopE₁₋₁₃₈-tBID BH3 might be expected.

In order to increase the genetic stability of YopE₁₋₁₃₈-(tBID BH3)₂ (SEQ ID No. 202) for in vivo studies, we cloned YopE₁₋₁₃₈-(tBID BH3)₂ (SEQ ID No. 202) by homologous recombination on the Yersinia virulence plasmid pYV at the native site of YopE and under the native YopE promoter (using mutator plasmids pSI_408 and pSI_419). Such mutators contain the DNA sequence coding for the desired protein, flanked by 200–250 bp of sequences on both sides corresponding to the site of the respective gene, where the integration shall take place. These plasmids are transformed into *E. coli* Sm10 λ pir, from where plasmids were mobilized into the corresponding *Y. enterocolitica* strain. Mutants carrying the integrated vector were propagated for several generations without selection pressure. Then sucrose was used to select for clones that have lost the vector. Finally mutants were identified by colony PCR. The endogenous proteins for the transport by the T3SS (called “*Yersinia* outer proteins”, Yops) are encoded by *Y. enterocolitica* on this 70kb plasmid, named plasmid of *Yersinia* Virulence (pYV), which further encodes the T3SS apparatus.

Yersinia strains encoding YopE₁₋₁₃₈-(tBID BH3) (SEQ ID No. 138 or 200) or YopE₁₋₁₃₈-(tBID BH3)₂ (SEQ ID No. 202) on the Yersinia virulence plasmid pYV at the native site of YopE and under the native YopE promoter were assessed for their capacity of inducing apoptosis in cancerous cells (including 4T1 and B16F10 cells, **Fig. 29**). The IC₅₀ (half maximal inhibitory concentration), referring to the number of bacteria per eukaryotic cell (MOI) needed in order to kill 50% of such cells, was found to be decreased upon delivery of tandem repeats of tBID BH3 domain as compared to a single tBID BH3 domain, when both proteins are encoded on the Yersinia virulence plasmid pYV at the native site of YopE and under the native YopE promoter (**Fig. 29**). This is in agreement with findings from expression plasmid borne delivery of these proteins (**Fig. 28**). Again, this finding was surprising, as the protein

size is increased by fusing a second BH3 domain of t-BID. Due to this, decreased expression and delivery levels of YopE₁₋₁₃₈-(tBID BH3)₂ (SEQ ID No. 202) as compared to YopE₁₋₁₃₈-tBID BH3 (SEQ ID No. 138 or 200) would be expected, and might maximally reach equivalent levels. In order to reach an increase in cell killing activity, the fused tBID BH3 domains must simultaneously act side by side upon delivery by the T3SS into eukaryotic cells. In case only one tBID BH3 domain in the YopE₁₋₁₃₈-(tBID BH3)₂ construct would be functional, at best the same efficiency as with YopE₁₋₁₃₈-tBID BH3 might be expected. Furthermore, *Yersinia* strains encoding YopE₁₋₁₃₈-(tBID BH3)₂ (SEQ ID No. 202) on the *Yersinia* virulence plasmid pYV at the native site of YopE and under the native YopE promoter were compared for their capacity of inducing apoptosis in cancerous cells to expression plasmid (pBad-MycHisA based) derived delivery of YopE₁₋₁₃₈-(tBID BH3)₂. In agreement with the higher copy number of pBad-MycHisA (20-25 copies) as compared to the pYV (1-6 copies are reported), pBad-MycHisA based delivery of YopE₁₋₁₃₈-(tBID BH3)₂ (SEQ ID No. 202) resulted in a slightly decreased IC50 value on 4T1 and B16F10 cells (Fig. 29).

Validation of tumor specific growth in vivo up to day 14 post bacterial administration

The experiment of tumor colonization by genetically modified *Y. enterocolitica* was repeated in a syngeneic murine allograft model (4T1 breast cancer model) and bacterial colonization was followed over two weeks. This time, mice were infected with 1×10^6 colony forming units (CFU) of *Y. enterocolitica* $\Delta yopH, O, P, E, M, T$. While obtaining similar results to the B16F10 model at early days post infection, we could further show that the tumor colonization is consistently found at day 8 and up to day 14 after infection (Fig. 30). Furthermore, the colonization remains highly specific with only low counts of bacteria detected in all other organs assessed (Fig. 31). These findings indicate that *Y. enterocolitica* $\Delta yopH, O, P, E, M, T$ is able to establish a persistent colonization of the tumor thereby preventing clearance by the immune system.

Efficacy of *Y. enterocolitica* $\Delta HOPEMT$ in delaying tumor progression

In order to assess the impact of YopE₁₋₁₃₈-(tBID BH3)₂ (SEQ ID No. 202) delivered to tumor cells in vivo, we performed studies in wildtype Balb/C mice allografted s.c.

with 4T1 breast cancer cells. We aimed at assessing the *Y. enterocolitica* $\Delta HOPEMT$ strain encoding YopE₁₋₁₃₈-(tBID BH3)₂ (SEQ ID No. 202) on the Yersinia virulence plasmid pYV at the native site of YopE and under the native YopE promoter. Mice were i.v. injected with PBS or 1×10^7 *Y. enterocolitica* $\Delta HOPEMT$ pYV-YopE₁₋₁₃₈-(tBID BH3)₂, once the tumor had reached a size of 150-250 mm³. The day of the i.v. injection of bacteria was defined as day 0. Tumor volume was measured over the following days (day 0 to day 9 post i.v. injection of bacteria) with calipers. The tumor volume was normalized to the tumor volume at day 0 to compensate for any initial heterogeneity in tumor size. Treatment with *Y. enterocolitica* $\Delta HOPEMT$ pYV-YopE₁₋₁₃₈-(tBID BH3)₂ showed an impact on tumor volume progression, with statistically significant tumor reduction at day 8, 9 and 10 post bacterial administration (**Fig. 32**). Importantly, *Y. enterocolitica* $\Delta HOPEMT$ alone was found not to impact tumor progression in the 4T1 murine cancer model (**Fig. 33**). These findings highlight that such bacteria and their T3SS can be employed for interference with tumor progression.

List of references

- 1 Hayes, C. S., Aoki, S. K. & Low, D. A. Bacterial contact-dependent delivery systems. *Annu Rev Genet* 44, 71-90, doi:10.1146/annurev.genet.42.110807.091449 (2010).
- 2 Cornelis, G. R. The type III secretion injectisome. *Nat Rev Microbiol* 4, 811-825, doi:nrmicro1526 [pii]10.1038/nrmicro1526 (2006).
- 3 Michiels, T., Wattiau, P., Brasseur, R., Ruyschaert, J. M. & Cornelis, G. Secretion of Yop proteins by *Yersinia*. *Infect Immun* 58, 2840-2849 (1990).
- 4 Blanco-Toribio, A., Muyldermans, S., Frankel, G. & Fernandez, L. A. Direct injection of functional single-domain antibodies from *E. coli* into human cells. *PLoS One* 5, e15227, doi:10.1371/journal.pone.0015227 (2010).
- 5 Bichsel, C. et al. Direct reprogramming of fibroblasts to myocytes via bacterial injection of MyoD protein. *Cell Reprogram* 15, 117-125, doi:10.1089/cell.2012.0058 (2013).
- 6 Bichsel, C. et al. Bacterial delivery of nuclear proteins into pluripotent and differentiated cells. *PLoS One* 6, e16465, doi:10.1371/journal.pone.0016465 (2011).
- 7 Chamekh, M. et al. Delivery of biologically active anti-inflammatory cytokines IL-10 and IL-1ra in vivo by the *Shigella* type III secretion apparatus. *J Immunol* 180, 4292-4298, doi:180/6/4292 [pii] (2008).
- 8 Hoffman, R. M. Tumor-seeking *Salmonella* amino acid auxotrophs. *Curr Opin Biotechnol* 22, 917-923, doi:S0958-1669(11)00061-9 [pii]10.1016/j.copbio.2011.03.009 (2011).
- 9 Hoang, T. T., Williams, S., Schweizer, H. P. & Lam, J. S. Molecular genetic analysis of the region containing the essential *Pseudomonas aeruginosa* *asd* gene encoding aspartate-beta-semialdehyde dehydrogenase. *Microbiology* 143 (Pt 3), 899-907 (1997).
- 10 Skurnik, M. & Wolf-Watz, H. Analysis of the *yopA* gene encoding the YopI virulence determinants of *Yersinia* spp. *Mol Microbiol* 3, 517-529 (1989).
- 11 Terti, R., Skurnik, M., Vartio, T. & Kuusela, P. Adhesion protein YadA of *Yersinia* species mediates binding of bacteria to fibronectin. *Infect Immun* 60, 3021-3024 (1992).
- 12 Isberg, R. R. & Leong, J. M. Multiple beta 1 chain integrins are receptors for invasins, a protein that promotes bacterial penetration into mammalian cells. *Cell* 60, 861-871, doi:0092-8674(90)90099-Z [pii] (1990).
- 13 Isberg, R. R., Voorhis, D. L. & Falkow, S. Identification of invasins: a protein that allows enteric bacteria to penetrate cultured mammalian cells. *Cell* 50, 769-778, doi:0092-8674(87)90335-7 [pii] (1987).
- 14 Leong, J. M., Fournier, R. S. & Isberg, R. R. Identification of the integrin binding domain of the *Yersinia pseudotuberculosis* invasins protein. *EMBO J* 9, 1979-1989 (1990).
- 15 Mota, L. J. & Cornelis, G. R. The bacterial injection kit: type III secretion systems. *Ann Med* 37, 234-249, doi:R673752030212825 [pii]10.1080/07853890510037329 (2005).
- 16 Trosky, J. E., Liverman, A. D. & Orth, K. *Yersinia* outer proteins: Yops. *Cell Microbiol* 10, 557-565, doi:CM1109 [pii]10.1111/j.1462-5822.2007.01109.x (2008).

- 17 Brenner, D. & Mak, T. W. Mitochondrial cell death effectors. *Curr Opin Cell Biol* 21, 871-877, doi:S0955-0674(09)00160-4 [pii]10.1016/j.ceb.2009.09.004 (2009).
- 18 Chalah, A. & Khosravi-Far, R. The mitochondrial death pathway. *Adv Exp Med Biol* 615, 25-45, doi:10.1007/978-1-4020-6554-5_3 (2008).
- 19 Fuchs, Y. & Steller, H. Programmed cell death in animal development and disease. *Cell* 147, 742-758, doi:S0092-8674(11)01283-9 [pii]10.1016/j.cell.2011.10.033 (2011).
- 20 Waugh, D. S. An overview of enzymatic reagents for the removal of affinity tags. *Protein Expr Purif* 80, 283-293, doi:S1046-5928(11)00203-8 [pii]10.1016/j.pep.2011.08.005 (2011).
- 21 Sory, M. P. & Cornelis, G. R. Translocation of a hybrid YopE-adenylate cyclase from *Yersinia enterocolitica* into HeLa cells. *Mol Microbiol* 14, 583-594 (1994).
- 22 Sarker, M. R., Neyt, C., Stainier, I. & Cornelis, G. R. The *Yersinia* Yop virulon: LcrV is required for extrusion of the translocators YopB and YopD. *J Bacteriol* 180, 1207-1214 (1998).
- 23 Howard, S. L. et al. Application of comparative phylogenomics to study the evolution of *Yersinia enterocolitica* and to identify genetic differences relating to pathogenicity. *Journal of bacteriology* 188, 3645-3653, doi:10.1128/JB.188.10.3645-3653.2006 (2006).
- 24 Neubauer, H., Aleksic, S., Hensel, A., Finke, E. J. & Meyer, H. *Yersinia enterocolitica* 16S rRNA gene types belong to the same genospecies but form three homology groups. *Int J Med Microbiol* 290, 61-64, doi:10.1016/S1438-4221(00)80107-1 (2000).
- 25 Pelludat, C., Hogardt, M. & Heesemann, J. Transfer of the core region genes of the *Yersinia enterocolitica* WA-C serotype O:8 high-pathogenicity island to *Y. enterocolitica* MRS40, a strain with low levels of pathogenicity, confers a yersiniabactin biosynthesis phenotype and enhanced mouse virulence. *Infect Immun* 70, 1832-1841 (2002).
- 26 Feldman, M. F., Muller, S., Wuest, E. & Cornelis, G. R. SycE allows secretion of YopE-DHFR hybrids by the *Yersinia enterocolitica* type III Ysc system. *Mol Microbiol* 46, 1183-1197, doi:3241 [pii] (2002).
- 27 Ramamurthi, K. S. & Schneewind, O. A synonymous mutation in *Yersinia enterocolitica* yopE affects the function of the YopE type III secretion signal. *J Bacteriol* 187, 707-715, doi:187/2/707 [pii]10.1128/JB.187.2.707-715.2005 (2005).
- 28 Wolke, S., Ackermann, N. & Heesemann, J. The *Yersinia enterocolitica* type 3 secretion system (T3SS) as toolbox for studying the cell biological effects of bacterial Rho GTPase modulating T3SS effector proteins. *Cell Microbiol* 13, 1339-1357, doi:10.1111/j.1462-5822.2011.01623.x (2011).
- 29 Forsberg, A. & Wolf-Watz, H. Genetic analysis of the yopE region of *Yersiniaspp.*: identification of a novel conserved locus, yerA, regulating yopE expression. *J Bacteriol* 172, 1547-1555 (1990).
- 30 Sambrook, J. (ed David W. Russell) (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. :, 2001).
- 31 Alto, N. M. & Dixon, J. E. Analysis of Rho-GTPase mimicry by a family of bacterial type III effector proteins. *Methods Enzymol* 439, 131-143, doi:S0076-6879(07)00410-7 [pii]10.1016/S0076-6879(07)00410-7 (2008).

- 32 Alto, N. M. et al. Identification of a bacterial type III effector family with G protein mimicry functions. *Cell* 124, 133-145, doi:S0092-8674(05)01229-8 [pii]10.1016/j.cell.2005.10.031 (2006).
- 33 Kaniga, K., Delor, I. & Cornelis, G. R. A wide-host-range suicide vector for improving reverse genetics in gram-negative bacteria: inactivation of the *blaA* gene of *Yersinia enterocolitica*. *Gene* 109, 137-141, doi:0378-1119(91)90599-7 [pii] (1991).
- 34 Yoneda, Y. et al. A long synthetic peptide containing a nuclear localization signal and its flanking sequences of SV40 T-antigen directs the transport of IgM into the nucleus efficiently. *Exp Cell Res* 201, 313-320 (1992).
- 35 Cornelis, G. R. in *Molecular aspects of host-pathogen interactions* (ed SAUNDERS MoCRAE, SMYTH, STOW) (Cambridge University Press, 1997).
- 36 Metcalf, W. W., Jiang, W. & Wanner, B. L. Use of the *rep* technique for allele replacement to construct new *Escherichia coli* hosts for maintenance of R6K gamma origin plasmids at different copy numbers. *Gene* 138, 1-7 (1994).
- 37 Diepold, A. et al. Deciphering the assembly of the *Yersinia* type III secretion injectisome. *EMBO J* 29, 1928-1940, doi:emboj201084 [pii]10.1038/emboj.2010.84 (2010).
- 38 Iriarte, M., Stainier, I. & Cornelis, G. R. The *rpoS* gene from *Yersinia enterocolitica* and its influence on expression of virulence factors. *Infect Immun* 63, 1840-1847 (1995).
- 39 Cornelis, G., Vanoortegem, J. C. & Sluiter, C. Transcription of the *yop* regulon from *Y. enterocolitica* requires trans acting pYV and chromosomal genes. *Microb Pathog* 2, 367-379, doi:0882-4010(87)90078-7 [pii] (1987).
- 40 Grosdent, N., Maridonneau-Parini, I., Sory, M. P. & Cornelis, G. R. Role of Yops and adhesins in resistance of *Yersinia enterocolitica* to phagocytosis. *Infect Immun* 70, 4165-4176 (2002).
- 41 Dehio, C., Meyer, M., Berger, J., Schwarz, H. & Lanz, C. Interaction of *Bartonella henselae* with endothelial cells results in bacterial aggregation on the cell surface and the subsequent engulfment and internalisation of the bacterial aggregate by a unique structure, the invasome. *J Cell Sci* 110 (Pt 18), 2141-2154 (1997).
- 42 Westerfield, M. *The Zebrafish Book: A Guide for the Laboratory Use of Zebrafish Danio rerio* (University of Oregon Press, Eugene, OR, 2000).
- 43 Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. & Schilling, T. F. Stages of embryonic development of the zebrafish. *Dev Dyn* 203, 253-310, doi:10.1002/aja.1002030302 (1995).
- 44 Benard, E. L. et al. Infection of zebrafish embryos with intracellular bacterial pathogens. *J Vis Exp*, doi:3781 [pii]10.3791/3781 (2012).
- 45 Blum, Y. et al. Complex cell rearrangements during intersegmental vessel sprouting and vessel fusion in the zebrafish embryo. *Dev Biol* 316, 312-322, doi:S0012-1606(08)00079-1 [pii]10.1016/j.ydbio.2008.01.038 (2008).
- 46 Herwig, L. et al. Distinct cellular mechanisms of blood vessel fusion in the zebrafish embryo. *Curr Biol* 21, 1942-1948, doi:S0960-9822(11)01140-7 [pii]10.1016/j.cub.2011.10.016 (2011).
- 47 Carpenter, A. E. et al. CellProfiler: image analysis software for identifying and quantifying cell phenotypes. *Genome Biol* 7, R100, doi:gb-2006-7-10-r100 [pii]10.1186/gb-2006-7-10-r100 (2006).

- 48 Bensimon, A. et al. ATM-dependent and -independent dynamics of the nuclear phosphoproteome after DNA damage. *Sci Signal* 3, rs3, doi:10.1126/scisignal.20010343/151/rs3 [pii] (2010).
- 49 Perkins, D. N., Pappin, D. J., Creasy, D. M. & Cottrell, J. S. Probability-based protein identification by searching sequence databases using mass spectrometry data. *Electrophoresis* 20, 3551-3567, doi:10.1002/(SICI)1522-2683(19991201)20:18<3551::AID-ELPS3551>3.0.CO;2-2 [pii]10.1002/(SICI)1522-2683(19991201)20:18<3551::AID-ELPS3551>3.0.CO;2-2 (1999).
- 50 Smyth, G. K. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* 3, Article3, doi:10.2202/1544-6115.1027 (2004).
- 51 Ting, L. et al. Normalization and statistical analysis of quantitative proteomics data generated by metabolic labeling. *Mol Cell Proteomics* 8, 2227-2242, doi:10.1074/mcp.M800462-MCP200M800462-MCP200 [pii] (2009).
- 52 Vizcaino, J. A. et al. The PRoteomics IDentifications (PRIDE) database and associated tools: status in 2013. *Nucleic Acids Res* 41, D1063-1069, doi:gks1262 [pii]10.1093/nar/gks1262 (2013).
- 53 Boyd, A. P., Lambermont, I. & Cornelis, G. R. Competition between the Yops of *Yersinia enterocolitica* for delivery into eukaryotic cells: role of the SycE chaperone binding domain of YopE. *J Bacteriol* 182, 4811-4821 (2000).
- 54 Iriarte, M. & Cornelis, G. R. YopT, a new *Yersinia* Yop effector protein, affects the cytoskeleton of host cells. *Mol Microbiol* 29, 915-929 (1998).
- 55 Kudryashev, M. et al. In situ structural analysis of the *Yersinia enterocolitica* injectisome. *Elife* 2, e00792, doi:10.7554/eLife.0079200792 [pii] (2013).
- 56 Schulte, R. et al. *Yersinia enterocolitica* invasin protein triggers IL-8 production in epithelial cells via activation of Rel p65-p65 homodimers. *FASEB J* 14, 1471-1484 (2000).
- 57 Mota, L. J., Journet, L., Sorg, I., Agrain, C. & Cornelis, G. R. Bacterial injectisomes: needle length does matter. *Science* 307, 1278, doi:307/5713/1278 [pii]10.1126/science.1107679 (2005).
- 58 Isaksson, E. L. et al. The membrane localization domain is required for intracellular localization and autoregulation of YopE in *Yersinia pseudotuberculosis*. *Infect Immun* 77, 4740-4749, doi:IAI.00333-09 [pii]10.1128/IAI.00333-09 (2009).
- 59 Denecker, G. et al. Effect of low- and high-virulence *Yersinia enterocolitica* strains on the inflammatory response of human umbilical vein endothelial cells. *Infect Immun* 70, 3510-3520 (2002).
- 60 Sharma, S. et al. Deployment of the *Burkholderia glumae* type III secretion system as an efficient tool for translocating pathogen effectors to monocot cells. *Plant J* 74, 701-712, doi:10.1111/tpj.12148 (2013).
- 61 Carrington, J. C. & Dougherty, W. G. A viral cleavage site cassette: identification of amino acid sequences required for tobacco etch virus polyprotein processing. *Proc Natl Acad Sci U S A* 85, 3391-3395 (1988).
- 62 Kapust, R. B., Tozser, J., Copeland, T. D. & Waugh, D. S. The P1' specificity of tobacco etch virus protease. *Biochem Biophys Res Commun* 294, 949-955, doi:10.1016/S0006-291X(02)00574-0S0006-291X(02)00574-0 [pii] (2002).

- 63 Liang, H., Gao, H., Maynard, C. A. & Powell, W. A. Expression of a self-processing, pathogen resistance-enhancing gene construct in *Arabidopsis*. *Biotechnol Lett* 27, 435-442, doi:10.1007/s10529-005-1884-9 (2005).
- 64 Weber, W. et al. Macrolide-based transgene control in mammalian cells and mice. *Nat Biotechnol* 20, 901-907, doi:10.1038/nbt731nbt731 [pii] (2002).
- 65 Kapust, R. B. et al. Tobacco etch virus protease: mechanism of autolysis and rational design of stable mutants with wild-type catalytic proficiency. *Protein Eng* 14, 993-1000 (2001).
- 66 Lee, V. T., Anderson, D. M. & Schneewind, O. Targeting of *Yersinia* Yop proteins into the cytosol of HeLa cells: one-step translocation of YopE across bacterial and eukaryotic membranes is dependent on SycE chaperone. *Mol Microbiol* 28, 593-601 (1998).
- 67 Gray, D. C., Mahrus, S. & Wells, J. A. Activation of specific apoptotic caspases with an engineered small-molecule-activated protease. *Cell* 142, 637-646, doi:S0092-8674(10)00783-X [pii]10.1016/j.cell.2010.07.014 (2010).
- 68 Henrichs, T. et al. Target-directed proteolysis at the ribosome. *Proc Natl Acad Sci U S A* 102, 4246-4251, doi:102/12/4246 [pii]10.1073/pnas.0408520102 (2005).
- 69 Hardt, W. D., Chen, L. M., Schuebel, K. E., Bustelo, X. R. & Galan, J. E. *S. typhimurium* encodes an activator of Rho GTPases that induces membrane ruffling and nuclear responses in host cells. *Cell* 93, 815-826, doi:S0092-8674(00)81442-7 [pii] (1998).
- 70 Hakansson, S. et al. The YopB protein of *Yersinia pseudotuberculosis* is essential for the translocation of Yop effector proteins across the target cell plasma membrane and displays a contact-dependent membrane disrupting activity. *EMBO J* 15, 5812-5823 (1996).
- 71 Stebbins, C. E. & Galan, J. E. Structural mimicry in bacterial virulence. *Nature* 412, 701-705, doi:10.1038/3508900035089000 [pii] (2001).
- 72 Li, H. et al. The phosphothreonine lyase activity of a bacterial type III effector family. *Science* 315, 1000-1003, doi:315/5814/1000 [pii]10.1126/science.1138960 (2007).
- 73 Norris, F. A., Wilson, M. P., Wallis, T. S., Galyov, E. E. & Majerus, P. W. SopB, a protein required for virulence of *Salmonella dublin*, is an inositol phosphate phosphatase. *Proc Natl Acad Sci U S A* 95, 14057-14059 (1998).
- 74 Pulliainen, A. T. et al. Bacterial effector binds host cell adenylyl cyclase to potentiate Galphas-dependent cAMP production. *Proc Natl Acad Sci U S A* 109, 9581-9586, doi:1117651109 [pii]10.1073/pnas.1117651109 (2012).
- 75 Li, H., Zhu, H., Xu, C. J. & Yuan, J. Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis. *Cell* 94, 491-501, doi:S0092-8674(00)81590-1 [pii] (1998).
- 76 Nagaraj, N. et al. Deep proteome and transcriptome mapping of a human cancer cell line. *Mol Syst Biol* 7, 548, doi:msb201181 [pii]10.1038/msb.2011.81 (2011).
- 77 Caussinus, E., Kanca, O. & Affolter, M. Fluorescent fusion protein knockout mediated by anti-GFP nanobody. *Nat Struct Mol Biol* 19, 117-121, doi:nsmb.2180 [pii]10.1038/nsmb.2180 (2011).
- 78 Cosma, C. L., Swaim, L. E., Volkman, H., Ramakrishnan, L. & Davis, J. M. Zebrafish and frog models of *Mycobacterium marinum* infection. *Curr Protoc Microbiol* Chapter 10, Unit 10B 12, doi:10.1002/0471729256.mc10b02s3 (2006).

- 79 Mathias, J. R. et al. Characterization of zebrafish larval inflammatory macrophages. *Dev Comp Immunol* 33, 1212-1217, doi:S0145-305X(09)00149-9 [pii]10.1016/j.dci.2009.07.003 (2009).
- 80 Jette, C. A. et al. BIM and other BCL-2 family proteins exhibit cross-species conservation of function between zebrafish and mammals. *Cell Death Differ* 15, 1063-1072, doi:cdd200842 [pii]10.1038/cdd.2008.42 (2008).
- 81 Olsen, J. V. et al. Global, in vivo, and site-specific phosphorylation dynamics in signaling networks. *Cell* 127, 635-648, doi:S0092-8674(06)01274-8 [pii]10.1016/j.cell.2006.09.026 (2006).
- 82 Schmutz, C. et al. Systems-Level Overview of Host Protein Phosphorylation During *Shigella flexneri* Infection Revealed by Phosphoproteomics. *Mol Cell Proteomics* 12, 2952-2968, doi:M113.029918 [pii]10.1074/mcp.M113.029918 (2013).
- 83 Szklarczyk, D. et al. The STRING database in 2011: functional interaction networks of proteins, globally integrated and scored. *Nucleic Acids Res* 39, D561-568, doi:gkq973 [pii]10.1093/nar/gkq973 (2011).
- 84 Huang da, W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 37, 1-13, doi:gkn923 [pii]10.1093/nar/gkn923 (2009).
- 85 Huang da, W. et al. DAVID gene ID conversion tool. *Bioinformation* 2, 428-430 (2008).
- 86 Schwerk, C. & Schulze-Osthoff, K. Regulation of apoptosis by alternative pre-mRNA splicing. *Mol Cell* 19, 1-13, doi:S1097-2765(05)01375-4 [pii]10.1016/j.molcel.2005.05.026 (2005).
- 87 Papagiannakopoulos, T., Shapiro, A. & Kosik, K. S. MicroRNA-21 targets a network of key tumor-suppressive pathways in glioblastoma cells. *Cancer Res* 68, 8164-8172, doi:68/19/8164 [pii]10.1158/0008-5472.CAN-08-1305 (2008).

Claims

1. A recombinant Gram-negative bacterial strain transformed with a vector which comprises in the 5' to 3' direction:
a promoter;
a first DNA sequence encoding a delivery signal from a bacterial T3SS effector protein, operably linked to said promoter; and
a second DNA sequence encoding repeated domains of a heterologous protein, wherein the repeated domains of a heterologous protein comprise a fusion protein consisting of several repetitions of a domain of a heterologous protein, or two or more domains of different heterologous proteins, wherein the two or more domains of different heterologous proteins comprise a fusion protein consisting of one or several repetitions of at least two domains of different heterologous proteins fused in frame to the 3' end of said first DNA sequence, wherein the heterologous proteins are proteins involved in apoptosis or apoptosis regulation, and
wherein the delivery signal from the bacterial T3SS effector protein encoded by the first DNA sequence comprises the YopE effector protein or an N-terminal fragment thereof or the SopE or the SteA effector protein or an N-terminal fragment thereof.
2. A recombinant Gram-negative bacterial strain transformed with a vector which comprises in the 5' to 3' direction:
a first DNA sequence encoding a delivery signal or a fragment thereof from a bacterial T3SS effector protein; and
a second DNA sequence encoding repeated domains of a heterologous protein, wherein the repeated domains of a heterologous protein comprise a fusion protein consisting of several repetitions of a domain of a heterologous protein, or two or more domains of different heterologous proteins, wherein the two or more domains of different heterologous proteins comprise a fusion protein consisting of one or several repetitions of at least two domains of different heterologous proteins fused in frame to the 3' end of said first DNA sequence, wherein the heterologous proteins are proteins involved in apoptosis or apoptosis regulation, and
wherein the delivery signal from the bacterial T3SS effector protein encoded by the first DNA sequence comprises the YopE effector protein or an N-terminal fragment thereof

or the SopE or the SteA effector protein or an N-terminal fragment thereof.

3. The recombinant Gram-negative bacterial strain of claim 1 or 2, wherein said recombinant Gram-negative bacterial strain is deficient in producing at least one T3SS effector protein.
4. The recombinant Gram-negative bacterial strain of anyone of claims 1-3 , wherein the recombinant Gram-negative bacterial strain is selected from the group consisting of the genera *Yersinia*, *Escherichia*, *Salmonella* and *Pseudomonas*.
5. The recombinant Gram-negative bacterial strain of anyone of claims 1-3, wherein the recombinant Gram-negative bacterial strain is a *Yersinia* strain or wherein the recombinant Gram-negative bacterial strain is a *Salmonella* strain.
6. The recombinant Gram-negative bacterial strain of claim 1 or 2, wherein the recombinant Gram-negative bacterial strain is a *Yersinia* strain and wherein said *Yersinia* strain is wild type or deficient in the production of at least one T3SS effector protein and wherein the delivery signal from the bacterial T3SS effector protein comprises the N-terminal 138 amino acids of the *Y. enterocolitica* YopE effector protein or, wherein the recombinant Gram-negative bacterial strain is a *Salmonella* strain and wherein said *Salmonella* strain is wild type or deficient in the production of at least one T3SS effector protein and wherein the delivery signal from the bacterial T3SS effector protein comprises the *S. enterica* SteA effector protein or the N-terminal 81 or 105 amino acids of the *S. enterica* SopE effector protein.
7. A vector which comprises in the 5' to 3' direction:
 - a promoter;
 - a first DNA sequence encoding a delivery signal from a bacterial T3SS effector protein, operably linked to said promoter, wherein the delivery signal from the bacterial T3SS effector protein comprises the YopE effector protein or an N-terminal fragment thereof or the SopE or the SteA effector protein or an N-terminal fragment thereof;
 - a second DNA sequence encoding repeated domains of a heterologous protein, wherein the repeated domains of a heterologous protein comprise a fusion protein consisting of several repetitions of a domain of a heterologous protein, or two or more domains of

different heterologous proteins, wherein the two or more domains of different heterologous proteins comprise a fusion protein consisting of one or several repetitions of at least two domains of different heterologous proteins fused in frame to the 3' end of said first DNA sequence,
wherein the heterologous proteins are proteins involved in apoptosis or apoptosis regulation.

8. A vector which comprises in the 5' to 3' direction:
a first DNA sequence encoding a delivery signal or a fragment thereof from a bacterial T3SS effector protein, wherein the delivery signal from the bacterial T3SS effector protein comprises the YopE effector protein or an N-terminal fragment thereof or the SopE or the SteA effector protein or an N-terminal fragment thereof; and
a second DNA sequence encoding repeated domains of a heterologous protein, wherein the repeated domains of a heterologous protein comprise a fusion protein consisting of several repetitions of a domain of a heterologous protein, or two or more domains of different heterologous proteins, wherein the two or more domains of different heterologous proteins comprise a fusion protein consisting of one or several repetitions of at least two domains of different heterologous proteins, fused in frame to the 3' end of said first DNA sequence, wherein the heterologous proteins are proteins involved in apoptosis or apoptosis regulation.
9. The recombinant Gram-negative bacterial strain of any one of claims 1-6 or the vector of claims 7 or 8, wherein the proteins involved in apoptosis or apoptosis regulation are selected from the group consisting of BH3-only proteins selected from the group consisting of Bad, BIM, Bid and tBid, Puma, Bik/Nbk, Bod, Hrk/Dp5, BNIP1, BNIP3, Bmf, Noxa, Mcl-1, Bcl-Gs, Beclin 1, Egl-1 and CED-13; caspases and intracellular signalling proteins of death receptor control of apoptosis.
10. The recombinant Gram-negative bacterial strain of any one of claims 1-6 or the vector of claim 7 or 8, wherein the repeated domain is the BH3 domain of apoptosis inducer tBID.
11. The recombinant Gram-negative bacterial strain of any one of claims 1-6 or the vector of claim 7 or 8, wherein the two or more domains of different heterologous proteins are

domains of heterologous proteins which belong to the same functional class of proteins.

12. The recombinant Gram-negative bacterial strain of any one of claims 1-6 or the vector of claim 7 or 8, wherein the two or more domains of different heterologous proteins are the BH3 domain of apoptosis inducer tBID and the BH3 domain of apoptosis regulator BAX.

13. The recombinant Gram-negative bacterial strain of any one of claims 1-6 or the vector of claim 7 or 8, wherein the vector comprises a third DNA sequence encoding a protease cleavage site, wherein the third DNA sequence is located between the 3'end of said first DNA sequence and the 5'end of said second DNA sequence.