

Claims
1. CLAIMS

1. A peptide comprising an amino acid sequence of SEQ ID No. 72 or an amino acid sequence selected from the group consisting of SEQ ID No. 72; NO: 1 to SEQ ID No. 71, and SEQ ID No. 73 to SEQ ID NO: 126, wherein said peptide has ~~an overall~~a length of ~~from 9 up to 16 amino acids;~~up to 16 amino acids, or a ~~pharmaceutically~~pharmaceutical acceptable salt thereof, ~~for use in medicine.~~
- ~~2.~~
2. The peptide ~~for use~~or salt thereof according to claim 1, wherein said peptide ~~consists of the amino acid sequence according to SEQ ID No. 72~~or salt thereof has the ability to bind to an MHC class-I or -II molecule, and wherein said peptide, when bound to said MHC, is capable of being recognized by CD4 and/or CD8 T cells.
- ~~3.~~
3. The peptide ~~for use~~or salt thereof according to claim 1 or 2, wherein the amino acid sequence thereof comprises a continuous stretch of amino acids according to any one of SEQ ID NO: 72, SEQ ID No. 1 to SEQ ID No. 71, and SEQ ID No. 73 to SEQ ID NO: 126.
4. The peptide or salt thereof according to any of claims 1 to 3, wherein said peptide or salt thereof consists of an amino acid sequence according to any of SEQ ID NO: 72, SEQ ID No. 1 to SEQ ID No. 71, and SEQ ID No. 73 to SEQ ID NO: 126.
- ~~3.5.~~ The peptide or salt thereof according to any of Claims 1 to 4, wherein said peptide is modified and/or includes non-peptide bonds.
- ~~4.~~

~~4.6.~~ The peptide for use or salt thereof according to any of claims 1 to 35,
wherein said peptide is part of a fusion protein, in particular comprising N-
terminal amino acids of the HLA-DR antigen-associated invariant chain (Ii).

~~5. A T-cell receptor, preferably a recombinant, soluble or membrane-bound T-cell
receptor that is reactive with an HLA ligand, wherein said ligand has at least 88%
identity to an amino acid sequence according to SEQ ID No. 72, and preferably
consists of the amino acid sequence of SEQ ID No. 72.~~

~~6. An antibody, in particular a soluble or membrane-bound antibody, that specifically
recognizes the peptide according to any of claims 1 to 3, preferably the peptide
according to any of claims 1 to 3 when bound to an MHC molecule.~~

~~7.~~

~~7.~~ A nucleic acid, encoding for a peptide according to any one of claims 1 to 4 or
for a TCR according to claim 5, or an antibody according to claim 6,
optionally linked to a heterologous promoter sequence, or an

~~5.8.~~ An expression vector capable of expressing said the nucleic acid according to
claim 7.

8.

~~6-9.~~ A recombinant host cell comprising the peptide according to claim 1 to 4, ~~or 6,~~
the nucleic acid according to claim 7 or the expression vector according to claim
78, wherein said host cell preferably is an antigen presenting cell such as a
dendritic cell, ~~or wherein said host cell preferably is a T cell or NK cell.~~

9.

10. The peptide or salt thereof according to any one of claims 1 to 6, the nucleic
acid according to claim 7, the expression vector according to claim 8, or the host
cell according to claim 9 for use in medicine.

~~7-11.~~ A method for producing the peptide according to any one of claims 1 to 4, ~~or for
producing the T cell receptor according to claim 5, or an antibody according
to claim 6,~~ the method comprising culturing the host cell according to claim 89
that presents the peptide according to claim 1 to 46, or expresses the nucleic
acid ~~or according to claim 7~~ or comprises the expression vector according to
claim 78, and isolating the peptide ~~or the TCR or the antibody~~ from the host cell
or its culture medium.

~~10.~~

~~8-12.~~ An *in vitro* method for producing activated T lymphocytes, the method
comprising contacting *in vitro* T cells with antigen loaded human class I or II
MHC molecules expressed on the surface of a suitable antigen-presenting cell
or an artificial construct mimicking an antigen-presenting cell for a period of time
sufficient to activate said T cells in an antigen specific manner, wherein said
antigen is a peptide according to any one of claims 1 ~~or 2~~ to 4.

~~11.~~

~~9-13.~~ An activated T lymphocyte, produced by the method according to claim 12, that
selectively recognizes a cell which presents a polypeptide comprising an amino
acid sequence given in any one of claims 1 ~~or 2~~ to 4.

~~12. A pharmaceutical composition comprising at least one active ingredient selected
from the group consisting of~~

14. A method for killing target cells in a patient which target cells present a
polypeptide comprising an amino acid sequence given in any one of claims 1 to

4, the method comprising administering to the patient an effective number of activated T cells as defined in claim 13.

15. An antibody, in particular a soluble or membrane-bound antibody, that specifically recognizes the peptide according to any one of claims 1 to 4, the nucleic acid or 5, preferably the peptide according to any of claims 1 to 5 when bound to an MHC molecule.

Use of a peptide according to any one of claims 1 to 6, the nucleic acid according to claim 7, the expression vector according to claim 78, the cell according to claim 89, the activated T lymphocyte according to claim 11 or the antibody according to claim 6 or T-cell receptor according to claim 5, and a pharmaceutically acceptable carrier, and optionally additional pharmaceutically acceptable excipients and/or stabilizers.

~~13. The peptide for use according to any one of claims 1 to 4, the nucleic acid or the expression vector according to claim 7, the cell according to claim 8, the activated T lymphocyte according to claim 11 or the antibody according to claim 6 or T-cell receptor according to claim 5 for use in medicine, preferably for use in the diagnosis and/or treatment of cancer.~~

~~16. 14. The peptide for use according to any one of claims 1 to 4, the nucleic acid or the expression vector according to claim 7, the cell according to claim 8, the activated T lymphocyte according to claim 11~~13~~ or the antibody according to claim 6 or T-cell receptor according to claim 5~~15~~ for use in the diagnosis and/or treatment of cancer, or for use in the manufacture of a medicament against cancer;~~

~~10.17. The use according to claim 4~~3~~16, wherein said cancer is selected from the group of ovariansmall cell lung cancer, non-small cell lung cancer, small cell lung cancer, kidneyrenal cell cancer, brain cancer, colon or rectumgastric cancer, stomachcolorectal cancer, liverhepatocellular cancer, pancreatic cancer, prostate cancer, leukemia, breast cancer, Merkel cell carcinoma, melanoma, esophagealovarian cancer, urinary bladder cancer, uterine cancer, gallbladder cancer, and bile duct cancer and esophageal cancer and other tumors that show an overexpression of thea protein from which a peptide according toSEQ ID NO: 72, SEQ ID No. 72~~1~~ to SEQ ID No. 71, and SEQ ID No. 73 to SEQ ID NO: 126 is derived from, wherein said protein is ECT2.~~

~~15.~~

~~11.18. A kit comprising:~~

- (a) a container comprising a pharmaceutical composition containing the peptide(s) or the salt according to any one of claims 1 to 4~~6~~, the nucleic acid or the expression vector(s) according to claim 7, the cell~~expression vector(s)~~ according to claim 8, the cell(s) according to claim 10, the activated T lymphocyte(s) according to claim 11~~13~~ or the antibody according to claim 6 or T-cell receptor according to claim 5~~15~~, in solution or in lyophilized form;

- (b) optionally, a second container containing a diluent or reconstituting solution for the lyophilized formulation;
- (c) optionally, at least one more peptide selected from the group consisting of SEQ ID No. ~~NO: 1 to 71, and 73 to 126~~ SEQ ID NO: 141, and
- (d) optionally, instructions for (i) use of the solution or (ii) reconstitution and/or use of the lyophilized formulation, ~~and~~.

e) optionally

19. The kit according to claim 18, further comprising one or more of (iii) a buffer, (iv) a diluent, (v) a filter, (vi) a needle, ~~or~~ (vii) a syringe, or (viii) an adjuvant.

20. The kit according to claim 18 or 19, wherein said peptide is selected from the group consisting of SEQ ID NO: 72, SEQ ID No. 1 to SEQ ID No. 71, and SEQ ID No. 73 to SEQ ID NO: 126.

21. A method for producing a personalized anti-cancer vaccine or a compound-based and/or cellular therapy for an individual patient, said method comprising:

a) identifying tumor-associated peptides (TUMAPs) presented by a tumor sample from said individual patient;

b) comparing the peptides as identified in a) with a warehouse of peptides that have been pre-screened for immunogenicity and/or over-presentation in tumors as compared to normal tissues

c) selecting at least one peptide from the warehouse that matches a TUMAP identified in the patient; and

d) formulating the personalized vaccine or compound-based or cellular therapy based on step c).

22. The method according to claim 21, wherein said TUMAPs are identified by:

a1) comparing expression data from the tumor sample to expression data from a sample of normal tissue corresponding to the tissue type of the tumor sample to identify proteins that are over-expressed or aberrantly expressed in the tumor sample; and

a2) correlating the expression data with sequences of MHC ligands bound to MHC class I and/or class II molecules in the tumor sample to identify MHC ligands derived from proteins over-expressed or aberrantly expressed by the tumor.

23. The method according to claim 21 or 22, wherein the sequences of MHC ligands are identified by eluting bound peptides from MHC molecules isolated from the tumor sample, and sequencing the eluted ligands.
24. The method according to any of claims 21 to 23, wherein the normal tissue corresponding to the tissue type of the tumor sample is obtained from the same patient.
25. The method according to any of claims 21 to 24, wherein the peptides included in the warehouse are identified based on the following steps:
- aa. Performing genome-wide messenger ribonucleic acid (mRNA) expression analysis by highly parallel methods, such as microarrays or sequencing-based expression profiling, comprising identify genes that over-expressed in a malignant tissue, compared with a normal tissue or tissues;
 - ab. Selecting peptides encoded by selectively expressed or over-expressed genes as detected in step aa, and
 - ac. Determining an induction of in vivo T-cell responses by the peptides as selected comprising *in vitro* immunogenicity assays using human T cells from healthy donors or said patient; or
 - ba. Identifying HLA ligands from said tumor sample using mass spectrometry;
 - bb. Performing genome-wide messenger ribonucleic acid (mRNA) expression analysis by highly parallel methods, such as microarrays or sequencing-based expression profiling, comprising identify genes that over-expressed in a malignant tissue, compared with a normal tissue or tissues;
 - bc. Comparing the identified HLA ligands to said gene expression data;
 - bd. Selecting peptides encoded by selectively expressed or over-expressed genes as detected in step bc;
 - be. Re-detecting of selected TUMAPs from step bd on tumor tissue and lack of or infrequent detection on healthy tissues and confirming the relevance of over-expression at the mRNA level; and

- bf. Determining an induction of in vivo T-cell responses by the peptides as selected comprising *in vitro* immunogenicity assays using human T cells from healthy donors or said patient.
26. The method according to any of claims 21 to 25, wherein the immunogenicity of the peptides included in the warehouse is determined by a method comprising in vitro immunogenicity assays, patient immunomonitoring for individual HLA binding, MHC multimer staining, ELISPOT assays and/or intracellular cytokine staining.
27. The method according to any of claims 21 to 26, wherein said warehouse comprises a plurality of peptides selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 141.
28. The method according to any of claims 21 to 27, further comprising identifying at least one mutation that is unique to the tumor sample relative to normal corresponding tissue from the individual patient, and selecting a peptide that correlates with the mutation for inclusion in the vaccine or for the generation of cellular therapies.
29. The method according to claim 28, wherein said at least one mutation is identified by whole genome sequencing.
30. A T-cell receptor, preferably soluble or membrane-bound, that is reactive with an HLA ligand, wherein said ligand has at least 75% identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 72, SEQ ID No. 1 to SEQ ID No. 71, and SEQ ID No. 73 to SEQ ID NO: 126.
31. The T-cell receptor according to claim 30, wherein said amino acid sequence is at least 88% identical to SEQ ID NO: 72, SEQ ID No. 1 to SEQ ID No. 71, and SEQ ID No. 73 to SEQ ID NO: 126.

32. The T-cell receptor according to claim 30 or 31, wherein said amino acid sequence consists any of SEQ ID NO: 72, SEQ ID No. 1 to SEQ ID No. 71, and SEQ ID No. 73 to SEQ ID NO: 126.
33. The T-cell receptor according to any of claims 30 to 32, wherein said T-cell receptor is provided as a soluble molecule and optionally carries a further effector function such as an immune stimulating domain or toxin.
34. A nucleic acid, encoding for a TCR according to any one of claims 30 to 33, optionally linked to a heterologous promoter sequence.
35. An expression vector capable of expressing the nucleic acid according to claim 34.
36. A host cell comprising the nucleic acid according to claim 34 or the nucleic acid encoding an antibody according to claim 15 or the expression vector according to claim 35, wherein said host cell preferably is a T cell or NK cell.
37. A method for producing the T cell receptor according to any claims 30 to 33, said method comprising culturing a host cell according to Claim 36, and isolating said T cell receptor from said host cell and/or its culture medium.
38. A pharmaceutical composition comprising at least one active ingredient selected from the group consisting of-
- a) a peptide selected from the group consisting of SEQ ID NO: 72, SEQ ID No. 1 to SEQ ID No. 71, and SEQ ID No. 73 to SEQ ID NO: 126;
 - b) a T-cell receptor reactive with a peptide and/or the peptide-MHC complex according to a);
 - c) a fusion protein comprising a peptide according to a), and the N-terminal amino acids 1 to 80 of the HLA-DR antigen-associated invariant chain (Ii);
 - d) a nucleic acid encoding for any of a) to c) or an expression vector comprising said nucleic acid,
 - e) a host cell comprising the expression vector of d,

f) an activated T-lymphocyte, obtained by a method comprising contacting in vitro T cells with a peptide according to a) expressed on the surface of a suitable antigen presenting cell for a period of time sufficient to activate said T cell in an antigen specific manner, as well as a method to transfer these activated T cells into the autologous or other patients;

g) an antibody, or soluble T-cell receptor, reactive to a peptide and/or the peptide – MHC complex according to a) and/or a cell presenting a peptide according to a), and potentially modified by fusion with for example immune-activating domains or toxins.

h) an aptamer recognizing a peptide selected from the group consisting of SEQ ID NO: 72, SEQ ID No. 1 to SEQ ID No. 71, and SEQ ID No. 73 to SEQ ID NO: 126 and/or a complex of a peptide selected from the group consisting of SEQ ID NO: 72, SEQ ID No. 1 to SEQ ID No. 71, and SEQ ID No. 73 to SEQ ID NO: 126 with a MHC molecule.

i) a conjugated or labelled peptide or scaffold according to any of a) to h) and a pharmaceutically acceptable carrier, and optionally, pharmaceutically acceptable excipients and/or stabilizers.

39. The pharmaceutical composition according to claim 38, wherein (i) the pharmaceutical composition further comprises an adjuvant, and/or (ii) wherein the pharmaceutical composition is a vaccine or a cellular therapeutic agent.

40. The pharmaceutical composition according to claim 39, wherein the adjuvant is an interleukin, including IL-2, IL-15 or a combination thereof.

41. An aptamer that specifically recognizes the peptide according to any of claims 1 to 5, preferably the peptide according to any of claims 1 to 5 that is bound to an MHC molecule.