

9608698-8 -1-
IP05
copyTHIS SHEET
REPLACEDNOVEL TREATMENTTHIS SHEET HAS BEEN
REPLACED

5 This invention relates to a novel method of treatment and to novel compounds for use in such a method.

10 European Published Patent Application No. 0126311 discloses substituted benzopyran compounds having blood pressure lowering activity, including 6-acetyl-trans-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol.

15 Also EP-A-0 376 524, EP-A-0 205 292, EP-A-0 250 077, EP-A-0 093 535, EP-A-0 150 202, EP-A-0 076 075 and WO/89/05808 (Beecham Group plc) describe certain benzopyran derivatives which possess anti-hypertensive activity.

20 EP-A-0 350 805 (Biersdorf), EP-A-0 277 611, EP-A-0 277612, EP-A-0 337 179 and EP-A-0 355 565 (Hoechst Aktiengesellschaft); and EP-A-0 466 131 (Nissan Chemical Industries Ltd) also describe certain benzopyran derivatives which are believed to possess anti-hypertensive activity.

25 EP-A-0 430 621 and EP-A-0 385 584 (Beecham Group plc) describe the resolution of certain intermediates useful in the preparation of the compounds described in the above mentioned patent applications.

EP-A-0 194 885 (E. Lilly) describes certain amino substituted benzopyran derivatives possessing anti-convulsant activity.

30 It has now been surprisingly found that certain compounds of formula (I) possess anxiolytic and anti-convulsant activity, and are also believed to have utility in the treatment or prevention of mania, depression and the effects associated with withdrawal from substances of abuse.


35 Accordingly, the present invention provides a method of treatment and/or prophylaxis of anxiety and/or mania, and/or depression and/or the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, and/or disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy, comprising

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

administering to the sufferer in need thereof an effective or prophylactic amount of a compound of formula (I) or pharmaceutically acceptable salt thereof:



5

wherein:

either Y is N and R₂ is hydrogen, or Y is C-R₁

10

where:

- either one of R₁ and R₂ is hydrogen and the other is selected from the class of hydrogen, C₃₋₈ cycloalkyl, C₁₋₆ alkyl optionally interrupted by oxygen or substituted by hydroxy, C₁₋₆ alkoxy or substituted aminocarbonyl, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy carbonyl, C₁₋₆ alkylcarbonyloxy, C₁₋₆ alkoxy, nitro, cyano, halo, trifluoromethyl, CF₃S, or a group CF₃-A-, where A is -CF₂-, -CO-, -CH₂- or CH(OH), trifluoromethoxy, C₁₋₆ alkylsulphinyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkoxy sulphinyl, C₁₋₆ alkoxy sulphonyl, aryl, heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphinyl, heteroarylsulphinyl, arylsulphonyl, heteroarylsulphonyl in which any aromatic moiety is optionally substituted, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxy carbonylamino, C₁₋₆ alkyl-thiocarbonyl, C₁₋₆ alkoxy-thiocarbonyl, C₁₋₆ alkyl-thiocarbonyloxy, 1-mercapto C₂₋₇ alkyl, formyl, or aminosulphinyl, aminosulphonyl or aminocarbonyl, any amino moiety being optionally substituted by one or two C₁₋₆ alkyl groups, or C₁₋₆ alkylsulphinylamino, C₁₋₆ alkylsulphonylamino, C₁₋₆ alkoxy sulphinylamino or C₁₋₆ alkoxy sulphonylamino, or ethylenyl terminally substituted by C₁₋₆ alkylcarbonyl, nitro or cyano, or -C(C₁₋₆ alkyl)NOH or -C(C₁₋₆ alkyl)NNH₂, or one of R₁ and R₂ is nitro, cyano or C₁₋₃ alkylcarbonyl and the other is methoxy or amino optionally substituted by one or two C₁₋₆ alkyl or by C₂₋₇ alkanoyl;

This Sheet Has Been Replaced



one of R_3 and R_4 is hydrogen or C_{1-4} alkyl and the other is C_{1-4} alkyl or R_3 and R_4 together are C_{2-5} polymethylene;

- R_5 is C_{1-6} alkylcarbonyloxy, benzoyloxy, ONO_2 , benzyloxy, phenyloxy or
- 5 C_{1-6} alkoxy and R_6 and R_9 are hydrogen or R_5 is hydroxy and R_6 is hydrogen or C_{1-2} alkyl and R_9 is hydrogen;

R_7 is fluorophenyl;

- 10 R_8 is hydrogen or C_{1-6} alkyl;

the $R_8-N-CO-R_7$ group being trans to the R_5 group;

and X is oxygen or NR_{10} where R_{10} is hydrogen or C_{1-6} alkyl.

15

Compounds of formula (I) include those in which Y is $C-R_1$, where;

- either one of R_1 and R_2 is hydrogen and the other is selected from the class of hydrogen, C_{1-6} alkylcarbonyl, C_{1-6} alkoxy carbonyl, C_{1-6}
- 20 alkylcarbonyloxy, C_{1-6} alkylhydroxymethyl, nitro, cyano, chloro, trifluoromethyl, trifluoromethoxy, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, C_{1-6} alkoxy sulphinyl, C_{1-6} alkoxy sulphonyl, C_{1-6} alkylcarbonylamino, C_{1-6} alkoxy carbonylamino, C_{1-6} alkyl-thiocarbonyl, C_{1-6} alkoxy-thiocarbonyl, C_{1-6} alkyl-thiocarbonyloxy, 1-mercapto C_{2-7}
- 25 alkyl, formyl, or aminosulphinyl, aminosulphonyl or aminocarbonyl, the amino moiety being optionally substituted by one or two C_{1-6} alkyl groups, or C_{1-6} alkylsulphinylamino, C_{1-6} alkylsulphonylamino, C_{1-6} alkoxy sulphinylamino or C_{1-6} alkoxy sulphonylamino, or ethylenyl terminally substituted by C_{1-6} alkylcarbonyl, nitro or cyano, or $-C(C_{1-6}$
- 30 alkyl)NOH or $-C(C_{1-6} \text{ alkyl})NNH_2$, or one of R_1 and R_2 is nitro, cyano or C_{1-3} alkylcarbonyl and the other is methoxy or amino optionally substituted by one or two C_{1-6} alkyl or by C_{2-7} alkanoyl;

- one of R_3 and R_4 is hydrogen or C_{1-4} alkyl and the other is C_{1-4} alkyl or
- 35 R_3 and R_4 together are C_{2-5} polymethylene;

R_5 is hydroxy and R_6 is hydrogen;

This Sheet Has Been Replaced



R₇ is fluorophenyl;

THIS SHEET HAS BEEN
REPLACED

R₈ is hydrogen or C₁₋₆ alkyl; and

5 R₉ is hydrogen;

the R₈-N-CO-R₇ group being trans to the R₅ group.

10 All C₁₋₆ alkyl or alkyl containing groups in formula (I) are preferably selected from methyl, ethyl, *n*- and *iso*-propyl, *n*-, *iso*-, *sec*- and *tert*-butyl.


Suitable C₃₋₈ cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

15 Aryl includes but is not limited to phenyl and naphthyl.

Heteroaryl includes a 5- or 6- membered monocyclic or 9- or 10- membered bicyclic of which 5- or 6- membered monocyclic heteroaryl is preferred. In addition, 5- or 6-membered monocyclic or 9- or 10-membered bicyclic
20 heteroaryl preferably contains one, two or three heteroatoms which are selected from the class of oxygen, nitrogen and sulphur and which, in the case of there being more than one heteroatom, are the same or different. Examples of 5- or 6-membered monocyclic heteroaryl containing one, two or three heteroatoms which are selected from the class of oxygen, nitrogen
25 and sulphur include furyl, thienyl, pyrrol, oxazolyl, thiazolyl, imidazolyl and thiadiazolyl, and pyridyl, pyridazyl, pyrimidyl, pyrazolyl and triazolyl. Preferred examples of such groups include furanyl, thienyl, pyrrol and pyridyl, in particular 2- and 3-furyl, 2- and 3-pyrrol, 2- and 3-thienyl, and 2-, 3- and 4-pyridyl. Examples of 9- or 10-membered
30 bicyclic heteroaryl containing one, two or three heteroatoms which are selected from the class of oxygen, nitrogen and sulphur include benzofuranyl, benzothienyl, indolyl and indazolyl, quinolyl and isoquinolyl, and quinazolyl. Preferred examples of such groups include 2- and 3-benzofuryl, 2- and 3-benzothienyl, and 2- and 3-indolyl, and 2- and
35 3-quinolyl.

Suitable examples of groups or atoms for optional substitution of aryl and heteroaryl include one, two or three substituents independently selected

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

from C₁₋₄ alkyl, C₁₋₄ alkoxy, halo (such as fluoro, chloro, bromo), hydroxy, nitro, cyano and SO_nH, where n=0 to 2.

Preferably R₂ is hydrogen.

5

Examples of suitable R₁ substituents include cyano, methoxy, trifluoromethoxy, chloro, trifluoromethyl, ethylcarbonyl, acetyl, hydrogen, methyl, ethyl, ~~iso-propyl, tertiary-butyl~~, nitro, C₂F₅, methoxycarbonyl, phenylsulphonyl, phenyl, fluoro, iodo, cyclopentyl, aminocarbonylmethyl and 1-hydroxyethyl. Preferably R₁ is cyano, acetyl or ethyl.

10

Preferably R₃ and R₄ are both methyl.

Preferably R₅ is hydroxy and R₆ and R₉ are hydrogen or R₅ is hydroxy, R₆ is C₁₋₂ alkyl and R₉ is hydrogen, more preferably R₅ is hydroxy and R₆ and R₉ are hydrogen.

15

It should be appreciated that the term fluorophenyl relating to R₇ encompasses phenyl which has 1,2,3,4 or 5 fluoro groups attached to the phenyl ring. Preferably there are 1 or 2 fluoro groups attached to the phenyl ring and most preferably there is 1 fluoro group attached to the ring.

20

The fluoro group or groups may be in any position around the phenyl ring, preferably in the case of mono fluorophenyl the fluoro is in the 3 or 4 position.

25

In the case of difluorophenyl, preferably the fluoro substituents are at positions 2,4 or 3,4.

30

Preferably R₈ is hydrogen or C₁₋₄ alkyl, more preferably R₈ is hydrogen, methyl or ethyl.

Preferably X is oxygen.

35

It should be appreciated that the compounds of formula (I) may have chiral carbon atoms at positions 2, 3 and 4 and therefore may exist as enantiomers. The present invention extends to each enantiomer and to

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

mixtures thereof including racemates. Preferably the compounds of formula (I) exist as 4S, 3R enantiomers.

5 It should also be appreciated that certain R_1 substituents also have chiral centres and therefore may exist as enantiomers. The present invention extends to each enantiomer and to mixtures thereof including racemates

10 An example of a compound of formula (I) is trans-6-acetyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (compound X).

The following novel compounds of formula (I) also form an aspect of the present invention, (herein referred to as compounds of formula (I')):

15 trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 5),

trans-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 6),

20 trans-6-Trifluoromethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 7),

25 trans-6-Cyano-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 8),

trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 9),

30 trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylmethylamino)2H-1-benzopyran-3-ol, (Example 10),

trans-6-Acetyl-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 11),

35 trans-6-Cyano-3,4-dihydro-4-(4-fluorobenzoylethylamino)-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 12),

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-7-nitro-2H-1-benzopyran-3-ol (Example 13),

5 trans-6-Acetyl-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 18),

trans-6-Acetyl-4R-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 19),

10 trans-6-Acetyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 20),

15 trans-6-Cyano-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 25),

trans-6-Cyano-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2,3-trimethyl-2H-1-benzopyran-3-ol (Example 26),

20 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-nitro-2H-1-benzopyran-3-ol (Example 27),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxycarbonyl-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 30),

25 trans-6-Fluoro-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 34),

trans-4-(4-fluorobenzoyl-methylamino)-6-trifluoromethyl-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 42),

30 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-iodo-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 47) and

35 trans-6-Aminocarbonylmethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 49).

The administration to the mammal may be by way of oral or parenteral administration.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

An amount effective to treat the disorders hereinbefore described depends on the usual factors such as the nature and severity of the disorders being treated and the weight of the mammal. However, a unit dose will
5 normally contain 1 to 1000 mg, suitably 1 to 500 mg, for example an amount in the range of from 2 to 100 mg such as 2, 5, 10, 20, 30, 40, 50 and 100 mg of the active compound. Unit doses will normally be administered once or more than once per day, for example 2, 3, 4, 5 or 6 times a day, more usually 2 to 4 times a day, such that the total daily dose
10 is normally in the range, for a 70 kg adult of 1 to 1000 mg, for example 50 to 500 mg, that is in the range of approximately 0.01 to 50 mg/kg/day, more usually 0.1 to 50 mg/kg/day, for example 1 to 50 mg/kg/day.

It is greatly preferred that the compound of formula (I) is administered in
15 the form of a unit-dose composition, such as a unit dose oral, rectal, topical or parenteral (especially intravenous) composition.

Such compositions are prepared by admixture and are suitably adapted for oral or parenteral administration, and as such may be in the form of
20 tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutible powders, injectable and infusible solutions or suspensions or suppositories. Orally administrable compositions are preferred, in particular shaped oral compositions, since they are more convenient for general use.

25 Tablets and capsules for oral administration are usually presented in a unit dose, and contain conventional excipients such as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, colourants, flavourings, and wetting agents. The tablets may be coated according to
30 well known methods in the art.

Suitable fillers for use include cellulose, mannitol, lactose and other similar agents. Suitable disintegrants include starch,
polyvinylpyrrolidone and starch derivatives such as sodium starch
35 glycollate. Suitable lubricants include, for example, magnesium stearate. Suitable pharmaceutically acceptable wetting agents include sodium lauryl sulphate.

This Sheet Has Been Replaced



These solid oral compositions may be prepared by conventional methods of blending, filling, tableting or the like. Repeated blending operations may be used to distribute the active agent throughout those compositions employing large quantities of fillers. Such operations are, of course,
5 conventional in the art.

Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable
10 vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethyl cellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles
15 (which may include edible oils), for example, almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

20 Oral formulations also include conventional sustained release formulations, such as tablets or granules having an enteric coating.

For parenteral administration, fluid unit dose forms are prepared
25 containing the compound and a sterile vehicle. The compound, depending on the vehicle and the concentration, can be either suspended or dissolved. Parenteral solutions are normally prepared by dissolving the compound in a vehicle and filter sterilising before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic,
30 preservatives and buffering agents are also dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum.

Parenteral suspensions are prepared in substantially the same manner
35 except that the compound is suspended in the vehicle instead of being dissolved and sterilised by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is



THIS SHEET HAS BEEN
REPLACED

included in the composition to facilitate uniform distribution of the compound of the invention.

5 As is common practice, the compositions will usually be accompanied by written or printed directions for use in the medical treatment concerned.

10 The present invention further provides a pharmaceutical composition for use in the treatment and/or prophylaxis of anxiety and/or mania and/or depression and/or the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines and/or disorders treatable or preventable with anti-convulsive agents, such as epilepsy, which comprises a compound of formula (I), or a pharmaceutically acceptable salt thereof, in particular, compounds of examples 1 to 50, and a pharmaceutically acceptable carrier.

15 In a further aspect the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, in particular compounds of examples 1 to 50 inclusive, for the manufacture of a medicament for the treatment and/or prophylaxis of anxiety and/or mania and/or depression and/or the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines and/or disorders treatable or preventable with anti-convulsive agents, such as epilepsy.

25 Such compositions may be prepared in the manner as hereinbefore described.


The invention also provides novel compounds of formula (I) and pharmaceutically acceptable salts thereof wherein R₇ is 2- or 30 3-fluorophenyl, 2, 4- or 3,4- difluorophenyl. Such compounds will hereinafter be referred to as compounds of formula (Ia).

Examples of compounds of formula (Ia) include:

35 trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(3-fluorobenzoylamino)2H-1-benzopyran-3-ol (Example 1),

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(2-fluoro-

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

benzoylamino)2H-1-benzopyran-3-ol (Example 2),

trans-6-Trifluoromethoxy-3,4-dihydro-2,2-dimethyl-4-(3-fluorobenzoylamino)2H-1-benzopyran-3-ol (Example 3),

5

trans-6-Cyano-4S-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 21),

10

trans-6-Cyano-4R-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 22),

trans-6-Cyano-4S-(2-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 24),

15

trans-6-Cyano-4S-(2,4-difluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 28),

trans-6-Acetyl-4-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 31),

20

trans-4-(2-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 38),

25

trans-4-(3-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 39),

trans-6-Cyano-4-(3,4-difluorobenzoyl-methylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 41) and

30

trans-6-cyano-4S-(3,4-difluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 44).

35 The invention further provides novel compounds of formula (I) and pharmaceutically acceptable salts thereof wherein Y is CR₁ where R₁ and R₂ are both hydrogen or one of R₁ and R₂ is trifluoromethoxy, C₁₋₆ alkyl optionally interrupted with oxygen or substituted with hydroxy, C₁₋₆ alkoxy or substituted amino-carbonyl, CF₃A-(where A is -CF₂-, -CO-, -CH₂) or CH(OH)), aryl sulphonyl, aryl C₃₋₈ cycloalkyl, C₁₋₆ alkoxy,

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphinyl, heteroarylsulphinyl, heteroarylsulphonyl, in which any aromatic moiety is optionally substituted and the other is hydrogen. Such compounds will hereinafter be referred to as compounds of formula (Ib).

5

Examples of compounds of formula (Ib) include:

trans-6-Trifluoromethoxy-3,4-dihydro-2,2-dimethyl-4-(3-fluorobenzoylamino)-2H-1-benzopyran-3-ol (Example 3),

10

trans-3,4-Dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)-2H-1-benzopyran-3-ol (Example 4),

15

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2,6-trimethyl-2H-1-benzopyran-3-ol (Example 14),

trans-6-Ethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 15),

20

trans-6-Ethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 16),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-isopropyl-2H-1-benzopyran-3-ol (Example 17),

25

trans-6-Ethyl-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 23),

30

trans-4-(4-Fluorobenzoylamino)-6-pentafluoroethyl-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 29),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-phenylsulphonyl-2H-1-benzopyran-3-ol (Example 32),

35

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-phenyl-2H-1-benzopyran-3-ol (Example 33),

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

trans-6-Ethyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 35),

5 trans-4R-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 36),

trans-4S-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 37),

10 trans-4-(4-fluorobenzoyl-methylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 43),

trans-6-t-Butyl-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 46),

15 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-cyclopentyl-2H-1-benzopyran-3-ol (Example 48) and

20 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxy-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 50).

The present invention also provides novel compounds of formula (I) and pharmaceutically acceptable salts thereof, where Y is N and R₂ is hydrogen. Such compounds will hereinafter be referred to as compounds of formula (Ic):

Examples of compounds of formula (Ic) include:

30 trans-4-(2-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 38),

trans-4-(3-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 39) and

35 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 40).

This Sheet Has Been Replaced



The present invention also provides novel compounds of formula (I) and pharmaceutically acceptable salts thereof, where X is NR₁₀ where R₁₀ is hydrogen or C₁₋₆ alkyl. Such compounds will hereinafter be referred to as compounds of formula (Id).

5

An example of a compound of formula (Id) is:

trans-6-Cyano-4-(4-fluorobenzoylamino)-2,2-dimethyl-1,2,3,4-tetrahydroquinolin-3-ol (Example 45).


10

Generally, compounds of formula (I) may be prepared by procedures generally described or analogous to those described in EP-0126311, EP-0376524, EP-205292, EP-0250077, EP-0093535, EP-0150202, EP-0076075, WO/89/05808, EP-0350805, EP-0277611, EP-0277612, EP-0337179, EP-0355565 and EP-0466131.

15

The invention also provides a process for the preparation of novel compounds of formula (I) or pharmaceutically acceptable salts thereof, which comprises acylating a compound of formula (II'):

20



25


wherein Y', R₂' and R₅' are the required variables Y, R₂ or R₅ as defined in formula (I) or a group convertible thereto and R₃, R₄, R₆, R₈, R₉ and X are the required variables as defined in formula (I), the R₈NH group being *trans* to the R₅ group, with and acylating agent of formula (IIIb):



30

where R₇ is as required and as defined in formula (I) and L₁ is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any R₁', R₂' and R₅' groups to R₁, R₂ and R₅ respectively, interconverting R₈ when hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.


35



THIS SHEET HAS BEEN
REPLACED

In particular, the present invention also provides a process for the preparation of compounds of formula (Ia), or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIa):

5



wherein Y, R₂' and R₅' are Y, R₂ or R₅ as defined in formula (I) or a group convertible thereto and R₃, R₄, R₆, R₈, R₉ and X are as defined in formula (I), the R₈NH group being trans to the R₅' group, with an acylating agent of formula (IIIa):

10




where R₇^a is 2- or 3-fluorophenyl or 2,4- or 3,4- difluorophenyl and L₁ is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any Y, R₂' or R₅' group to Y, R₂ or R₅ respectively, interconverting R₈ when hydrogen to C₁₋₆ alkyl, separating any entantioners and forming a pharmaceutically acceptable salt or solvate.

20

The present invention also provides a process for the preparation of compounds of formula (Ib), or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIb):

25



where R₁' and R₂' are both hydrogen or one of R₁ and R₂ is trifluoromethoxy, C₁₋₆ alkyl optionally interrupted with oxygen or substituted with hydroxy, C₁₋₆ alkoxy or substituted aminocarbonyl, CF₃A- (where A is CF₂-, CO-, -CH₂- or CH(OH)), aryl, aryl sulphonyl, aryl

30

This Sheet Has Been Replaced

C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphiny, heteroarylsulphiny, heteroarylsulphonyl, in which any aromatic moiety is optionally substituted and the other is hydrogen, or groups convertible to any of these; R₅' is R₅ as defined in formula (I) or a group convertible thereto and R₃, R₄, R₆, R₈, R₉ and X are as defined in formula (I) the R₈NH group being trans to the R₅' group, with a compound of formula (IIIb):




10

where R₇ is as defined in formula (I) and L₁ is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any R₁', R₂' and R₅' groups to R₁, R₂ and R₅ respectively, interconverting R₈ when hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

15

The present invention also provides a process for the preparation of compounds of formula (Ic), or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIc):

20




in which Y is N, R₅' is R₅, as defined in relation to formula (I) or a group convertible to R₅ and R₂' is hydrogen or a group convertible thereto, R₆, R₃, R₄, R₈, R₉ and X are as defined in formula (I), the R₈NH group being trans to the R₅' group, with an acylating agent of formula (IIIb):

25



30


where R₇ is as defined in formula (I) and L₁ is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any R₂' or R₅' group to hydrogen or R₅ respectively, interconverting R₈ when



THIS SHEET HAS BEEN
REPLACED

hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

- 5 The present invention also provides a process for the preparation of compounds of formula (Id), or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIId):



- 10 in which X' is NR_{10'} where R_{10'} is hydrogen or C₁₋₆ alkyl or a group convertible thereto, Y', R_{2'} and R_{5'} are Y, R₂ and R₅ respectively as defined in formula (I) or groups convertible thereto and R₃, R₄, R₆, R₈ and R₉ are as defined in formula (I), the R₈NH group being trans to the R_{5'} group, with an acylating agent of formula (IIIb):

15




- where R₇ is as defined in formula (I) and L₁ is a leaving group, thereafter optionally or as necessary and in any appropriate order converting any Y',
20 R_{10'}, R_{2'} or R_{5'} group to Y, R₁₀, R₂ or R₅ respectively, interconverting R₈ when hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

- 25 Examples of suitable leaving groups L₁ include those mentioned in the above-mentioned patents, in particular EP-A-0 126 311 or are conventional in the art.

- 30 The reaction conditions which may be used to carry out the above reactions are as outlined or analogous to those described in the above-mentioned patents, in particular EP-A-0 126 311.

- In particular, the leaving group (L₁) is a group that is displaceable by a primary or secondary amino nucleophile. Examples of such a group include C₁₋₄ alkylcarbonyloxy and halogen, such as chloro and bromo.
35 When the leaving group (L₁) is any of these examples, the acylating agent

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

of formula (IIIa) or (IIIb) is either an acid anhydride or an acid halide.

When it is an acid anhydride, it is, preferably, a mixed anhydride, which may be prepared in situ from an aromatic or heteroaromatic carboxylic acid and an alkyl chlorocarbonate, such as ethyl chloroformate.

5

When the acylating agent of formula (IIIa) or (IIIb) is an acid anhydride, the acylation of the compound formula (IIa), (IIb), (IIc) or (IId) is, preferably carried out using the anhydride as the solvent in the presence of an acid acceptor, such as sodium acetate.

10

When the acylating agent of formula (IIIa) or (IIIb) is an acid halide, the acylation of the compound of formula (IIa), (IIb), (IIc) or (IId), is, preferably, carried out in a non-aqueous medium, such as methylene chloride, in the presence of an acid acceptor, such as triethylamine, trimethylamine or pyridine.

15

Examples of suitable groups convertible to Y (or R₁), R₂ and R₅ include those described in the above-mentioned patents or are conventional in the art.

20

Interconversions of R₈ when hydrogen to C₁₋₆ alkyl may be carried out using conventional alkylation procedures for example using an alkylhalide in the presence of a base.

25

It should be appreciated that racemates for formula (I) may be resolved or enantiomerically purified compounds of formula (I) may be prepared using procedures conventional in the art and in particular using the procedures outlined in EP-0430631 and EP-0355584.

30

Suitably, the procedures outlined in or analogous to those described in Example 35 of the present specification may be used to prepare specific enantiomers of any compounds of formulae (I), (I'), (Ia), (Ib), (Ic) or (Id).

35

Compounds of formulae (I'), (IIa), (IIb), (IIc) and (IId) may be prepared from readily available starting materials using the procedures outlined or analogous to those described in the above-mentioned patents.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN

Compounds of formulae (IIIa) and (IIIb) are either commercially available or may be prepared according to conventional procedures known in the art of organic chemistry.

- 5 Novel compounds of formulae (I'), (IIa), (IIb), (IIc) and (IId) form an aspect of the present invention.

- Compounds of formula (I) in which R₅ is hydroxy, R₆ is C₁₋₂ alkyl and R₉ is hydrogen may be prepared according to the procedures outlined in
10 R. Gericke *et al.* J. Med. Chem. Vol.34, p3074(1991).


- The invention also provides a pharmaceutical composition comprising a compound of formula (I'), (Ia), (Ib), (Ic) or (d), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

- 15 The invention also provides the use of a compound of formula (I'), (Ia), (Ib), (Ic) or (Id), or a pharmaceutically acceptable salt thereof, in the treatment and/or prophylaxis of anxiety and/or mania and/or depression and/or the effects associated with withdrawal from substances of abuse
20 such as cocaine, nicotine, alcohol and benzodiazepines and/or disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy.

- The following examples and pharmacological test results illustrate the
25 present invention:

The following compounds were prepared by methods analogous to those described in the abovementioned patents publications.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

Example 1

5 trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(3-fluoro-
benzoylamino)2H-1-benzopyran-3-ol.

Mpt. 193-194°C.

Example 2

10

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(2-fluoro-
benzoylamino)2H-1-benzopyran-3-ol

Mpt. 163-5°

15

Example 3

20 trans-6-Trifluoromethoxy-3,4-dihydro-2,2-dimethyl-4-(3-
fluorobenzoylamino)2H-1-benzopyran-3-ol
Mpt. 127-30°C

Example 4

25 trans-3,4-Dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)-
2H-1-benzopyran-3-ol

Mpt. 174-5°C

Example 5

30

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(4-fluoro-
benzoylamino)2H-1-benzopyran-3-ol,

Mpt. 229-230°C.

35 3R, 4S isomer (compound B) Mpt. 224-5°C

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

Example 6

5 **trans-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(4-fluoro-
benzoylamino)2H-1-benzopyran-3-ol,**

Mpt. 177-9°C

Example 7

10

**trans-6-Trifluoromethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-
dimethyl-2H-1-benzopyran-3-ol,**

Mpt. 185-7°C

15

Example 8

20

**trans-6-Cyano-4-(4-fluorobenzoylmethylamino)-3,4-dihydro- 2,2-dimethyl-
2H-1-benzopyran-3-ol,**

Mpt. 230-4°C

Example 9

25

**trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-4-(4-
fluorobenzoylamino)2H-1-benzopyran-3-ol,**

Mpt. 205-207°C.

30

Example 10

**trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-
4-(4-fluorobenzoylmethylamino)2H-1-benzopyran-3-ol,**

35

Mpt. 210-212°C.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

Example 11

5 trans-6-Acetyl-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-
2H-1-benzopyran-3-ol,

Mpt. 207-8°C

Example 12

10

trans-6-Cyano-3,4-dihydro-4-(4-fluorobenzoylethylamino)-2,2-dimethyl-
2H-1-benzopyran-3-ol
mp 172-175 °C

15 Example 13

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-7-nitro-2H-1-
benzopyran-3-ol
mp 231-233 °C

20

Example 14

25 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2,6-trimethyl-2H-1-
benzopyran-3-ol
mp 185-186 °C

Example 15

30 trans-6-Ethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-
benzopyran-3-ol
mp 235-237 °C

Example 16

35 trans-6-Ethyl-4-(4-fluorobenzoylethylamino)-3,4-dihydro-2,2-dimethyl-2H-
1-benzopyran-3-ol
mp 175 °C

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

Example 17

- 5 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-isopropyl-2H-1-benzopyran-3-ol
mp 235-236 °C

Example 18

- 10 trans-6-Acetyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 203 °C

Example 19

- 15 trans-6-Acetyl-4R-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol
mp 162-163 °C

20 Example 20

- trans-6-Cyano-4S-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol
mp 164 °C

25

Example 21

- trans-6-Cyano-4S-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol
30 mp 221 °C

Example 22

- trans-6-Cyano-4R-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol
35 mp 221 °C

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

Example 23

- 5 **trans-6-Ethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol**
mp 162-164 °C

Example 24

- 10 **trans-6-Cyano-4S-(2-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol**
mp 163-165 °C

Example 25

- 15 **trans-6-Cyano-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol**
mp 224-225 °C

20 Example 26

trans-6-Cyano-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2,3-trimethyl-2H-1-benzopyran-3-ol
mp 218-220 °C

25

Example 27

- 30 **trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-nitro-2H-1-benzopyran-3-ol**
mp 197-198 °C

Example 28

- 35 **trans-6-Cyano-4S-(2,4-difluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol**
mp 175-176 °C

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

Example 29

- 5 trans-4-(4-Fluorobenzoylamino)-6-pentafluoroethyl-3,4-dihydro-2,2-
dimethyl-2H-1-benzopyran-3-ol
mp 181 °C

Example 30

- 10 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxycarbonyl-2,2-
dimethyl-2H-1-benzopyran-3-ol
mp 198 °C

Example 31

- 15 trans-6-Acetyl-4-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-
benzopyran-3-ol
mp 212 °C

20 Example 32

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-
phenylsulphonyl-2H-1-benzopyran-3-ol
mp 239-240 °C

25

Example 33

- 30 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-phenyl-2H-1-
benzopyran-3-ol
mp 164-165 °C

Example 34

- 35 trans-6-Fluoro-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-
benzopyran-3-ol
mp 164-165 °C

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED**Example 35**

trans-6-Ethyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol

5

(-)-Mandelic acid was used to resolve the residue from resolution of trans-4-amino-6-ethyl-2,2-dimethyl-chroman-3-ol (to give the 3S, 4R enantiomer - as described in EPA 412 760) under standard resolving conditions using acetone as recrystallisation solvent. This furnished trans-4S-amino-6-ethyl-2,2-dimethylchroman-3R-ol D-mandellate.

10

The 4S, 3R-aminoalcohol mandelate (4.0 g) was dissolved in dichloromethane (250 ml) and triethylamine (3.34 ml) and cooled in an ice bath. To this stirred solution was added 4-fluorobenzoyl chloride (1.7 g) dropwise. On completion of addition the reaction mixture was allowed to attain room temperature and was stirred overnight. The solvent was evaporated and the residue was taken up in ethyl acetate (150 ml). This solution was washed with 5% sodium bicarbonate solution, and brine, and dried over anhydrous magnesium sulphate. Filtration and evaporation and recrystallisation of the residue from Ethyl acetate-hexane gave the title compound of example 35. mp 132-136°C. $\alpha_D/20 + 69.1^\circ$ (methanol, $c=1.0$).

15

20

25

30

35

NMR(CDCL₃) δ

1.21 (3H, t, J = 8 Hz)
1.28 (3H, s)
1.49 (3H, s)
2.88 (2H, q, J = 8 Hz)
3.76 (1H, d, J = 9 Hz)
4.46 (1H, broad s)
5.21 (1H, t, J = 9,8 Hz)
6.41 (1H, d, J = 8 Hz)
6.80 (1H, d, J = 9 Hz)
7.07 (2H, irregular m)
7.17 (2H, t, J = 9 Hz)
7.74 (2H, m)

This Sheet Has Been Replaced


THIS SHEET HAS BEEN
REPLACED

Example 36

- 5 **trans-4R-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-**
dimethyl-2H-1-benzopyran-3S-ol
mp 132-133 °C

Example 37

- 10 **trans-4S-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-**
dimethyl-2H-1-benzopyran-3R-ol
mp 133-134 °C

Example 38

- 15 **trans-4-(2-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-**
clpyridin-3-ol
mp 254 °C

20 **Example 39**

trans-4-(3-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-
clpyridin-3-ol
mp 259-261 °C

25

Example 40

- 30 **trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-**
clpyridin-3-ol
mp 254-255 °C

Example 41

- 35 **trans-6-Cyano-4-(3,4-difluorobenzoyl-methylamino)-3,4-dihydro-2,2-**
dimethyl-2H-1-benzopyran-3-ol
mp 199-200°C

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

Example 42

- 5 trans-4-(4-fluorobenzoyl-methylamino)-6-trifluoromethyl-3,4-dihydro-2,2-
dimethyl-2H-1-benzopyran-3-ol
mp 196-197°C

Example 43

- 10 trans-4-(4-fluorobenzoyl-methylamino)-3,4-dihydro-2,2-dimethyl-2H-1-
benzopyran-3-ol
mp 160-161°C

Example 44

- 15 trans-6-cyano-4S-(3,4-difluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-
2H-1-benzopyran-3R-ol
mp 227°C

20 Example 45

trans-6-Cyano-4-(4-fluorobenzoylamino)-2,2-dimethyl-1,2,3,4-
tetrahydroquinolin-3-ol
mp 244-248 °C

25

Example 46

- 30 trans-6-t-Butyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-
benzopyran-3-ol
mp 163-166 °C

Example 47

- 35 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-iodo-2,2-dimethyl-2H-1-
benzopyran-3-ol
mp 204-205 °C

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

Example 48

- 5 **trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-cyclopentyl-
2H-1-benzopyran-3-ol**
mp 173-174 °C

Example 49

- 10 **trans- 6-Aminocarbonylmethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-
dimethyl-2H-1-benzopyran-3-ol**
mp 190 °C

Example 50

- 15 **trans- 4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxy-2,2-dimethyl-2H-
1-benzopyran-3-ol**
mp 155-156 °C

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

PHARMACOLOGICAL DATA

1. Geller-Seifter Procedure

5

Potential anxiolytic properties are evaluated using the Geller-Seifter procedure based on that originally described by Geller and Seifter, (1960) Psychopharmacologia, 1, 482-492. This procedure has been shown to be selective for drugs with anxiolytic properties (Cook and Sepinwall, (1975) "Mechanism of Action of Benzodiazepines" ed. Costa, E. and Greengard, P., Raven Press, New York, pp. 1-28).

10

Rats are trained on a variable interval 30 sec schedule (VI30) to press a lever in order to obtain food reward. The 5 min sessions of the VI30 schedule alternate with 2-5 min of a schedule (FR5) in which every 5th lever press is followed by presentation of a food pellet paired with a 0.5 sec mild footshock. The total study lasts approximately 30 mins. Rats typically respond with high rates of lever pressing under the VI30 schedule and low response rates under the FR5 'conflict' session.

15

20

25

Drugs are administered intraperitoneally or orally to groups of 3-8 rats 30 to 60 mins before testing.

The results are expressed as the percentage increase in the square root of the total number of lever presses in the FR5 'conflict' session. Square root transformation is necessary to normalise the data for statistical analysis using parametric methods.

30

The compound of Example 4 showed a significant increase in responding in the 'conflict' session at a dose of 10mg/kg p.o.

2. MES TEST

35

The maximal electroshock seizure (MES) test in rodents is one of the most widely used models of human grand mal epilepsy¹. In this model,

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

anticonvulsant agents elevate the threshold to electrically-induced seizures whilst proconvulsants lower the seizure threshold.

Method

5 Mice (male, Charles River, U.K. CD-1 strain, 25-30g) are randomly assigned to groups of 10-20 and dosed orally or intraperitoneally at a dose volume of 10ml/kg with various doses of compound (1-100 mg/kg) or vehicle. Mice are then subjected at 30 or 60 min post dose to a variable
10 voltage electroshock (0.1 sec., 50 Hz, sine wave form) via a buccal and a subcutaneous electrode. The mean voltage and standard error required to induce a tonic seizure in 50% (CV₅₀) of the mice in the group is determined by the 'up and down' method of Dixon and Mood (1948)². Statistical comparisons between vehicle- and drug-treated groups are
15 made using the method of Litchfield and Wilcoxon (1949)³.

In control animals the CV₅₀ is usually 40 - 50V. Hence the first animal in the control group is subjected to a voltage of 45V. If a tonic seizure does not ensue, the voltage is increased for a subsequent mouse. If a tonic
20 convulsion does occur, then the voltage is decreased, and so on until all the animals in the group have been tested.

The percentage increase or decrease in CV₅₀ for each group compared to the control is calculated.

25 Studies are carried out using a Heathkit shock generator with totally variable control of shock level from 0 to 200V and voltage steps of 5V are used.

30 Drugs are suspended in 1% methyl cellulose.

Reference

1. Swinyard, E.A. (1972). Electrically-induced convulsions. In:
35 Experimental Models of Epilepsy ed. Purpura, D.P. et al., 433-458, Raven Press, New York.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

2. Dixon, W.J. and Mood, A.M. (1948). J. Amer. Stat. Assn., **43**, 109-126.

5 3. Litchfield, J.T. and Wilcoxon, F. (1949). J. Pharmacol. Exp. Ther. **96**, 99-113.

Results

10 Compounds of Examples 1-3, 5, 7, 8, 18, 20, 21, 25, 30, 31, 35 and 37 and compound X showed a significant increase in CV₅₀ at a dose of 10 mg/kg p.o.

3. X-Maze

15 Introduction


The X-maze test of anxiety (Handley and Mithani, 1984) examines the exploratory response of naive rats in an environment which offers both anxiogenic (open arms) and relatively non-anxiogenic (closed arms) areas.

20 A selective increase in exploration of the open arms following drug pretreatment is therefore postulated to indicate anxiolytic effects.

Method

25 The X-maze was raised 70cm above the floor and consisted of two enclosed arms 45cm (long) x 15cm (wide) x 10cm (high) and two open arms 45 x 10 x 1cm arranged such that the two arms of each type were opposite each other. Both arm types were marked into two equal sections. Rats were placed onto the centre of the X-maze and observed for a period of 10
30 minutes during which time the following parameters were recorded: 1) the number of entries onto, and the time spent on, (a) open arms, (b) closed arms, (c) end of open arms and (d) end of closed arms. 2) the number of sections crossed. The fear-drive evoked in the open arms exceeds that in the enclosed arms and rats typically show a clear
35 preference for the enclosed arms. Anxiolytic drugs increase the number of entries made onto, and the time spent on, the outer half of the open arms, and also the percentage of entries made onto, and the time spent on, the whole of the open arms. These four measures of anxiety, and also the

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

total number of sections traversed, were calculated for each animal.

Drugs are administered intraperitoneally or orally to groups of 6 to 12 rats 30 to 60 mins before testing. Statistical comparisons between vehicle- and drug-treated groups were made using a Mann-Whitney 'U' test (two tailed).

5


S.L. Handley and S. Mithani, Arch. Pharmacol., 1984 327 1-5

RESULTS

10

The compound of Example 21, caused a significant increase in open arm entries at a dose of 30 mg/kg p.o.


This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

Claims

1. A method of treatment and/or prophylaxis of anxiety and/or mania, and/or depression and/or the effects associated with withdrawal from substances of abuse, and/or disorders treatable and/or preventable with anti-convulsive agents, comprising administering to the sufferer in need thereof an effective or prophylactic amount of a compound of formula (I) or pharmaceutically acceptable salt thereof:




wherein:

either Y is N and R₂ is hydrogen, or Y is C-R₁

where:

either one of R₁ and R₂ is hydrogen and the other is selected from the class of hydrogen, C₃₋₈ cycloalkyl, C₁₋₆ alkyl optionally interrupted by oxygen or substituted by hydroxy, C₁₋₆ alkoxy or substituted aminocarbonyl, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy carbonyl, C₁₋₆ alkylcarbonyloxy, C₁₋₆ alkoxy, nitro, cyano, halo, trifluoromethyl, CF₃S, or a group CF₃-A-, where A is -CF₂-, -CO-, -CH₂- or CH(OH), trifluoromethoxy, C₁₋₆ alkylsulphanyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkoxy sulphanyl, C₁₋₆ alkoxy sulphonyl, aryl, heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphanyl, heteroarylsulphanyl, arylsulphonyl, heteroarylsulphonyl in which any aromatic moiety is optionally substituted, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxy carbonylamino, C₁₋₆ alkyl-thiocarbonyl, C₁₋₆ alkoxy-thiocarbonyl, C₁₋₆ alkyl-thiocarbonyloxy, 1-mercapto C₂₋₇ alkyl, formyl, or aminosulphanyl, aminosulphonyl or aminocarbonyl, any amino moiety being optionally substituted by one or two C₁₋₆ alkyl groups, or C₁₋₆ alkylsulphanylamino, C₁₋₆ alkylsulphonylamino, C₁₋₆ alkoxy sulphanylamino or C₁₋₆ alkoxy sulphonylamino, or ethylenyl



This Sheet Has Been Replaced



terminally substituted by C₁₋₆ alkylcarbonyl, nitro or cyano, or
-C(C₁₋₆ alkyl)NOH or -C(C₁₋₆ alkyl)NNH₂, or one of R₁ and R₂ is
nitro, cyano or C₁₋₃ alkylcarbonyl and the other is methoxy or
amino optionally substituted by one or two C₁₋₆ alkyl or by C₂₋₇
alkanoyl;

one of R₃ and R₄ is hydrogen or C₁₋₄ alkyl and the other is C₁₋₄
alkyl or R₃ and R₄ together are C₂₋₅ polymethylene;

R₅ is C₁₋₆ alkylcarbonyloxy, benzoyloxy, ONO₂, benzyloxy,
phenyloxy or C₁₋₆ alkoxy and R₆ and R₉ are hydrogen or R₅ is
hydroxy and R₆ is hydrogen or C₁₋₂ alkyl and R₉ is hydrogen;

R₇ is fluorophenyl;

R₈ is hydrogen or C₁₋₆ alkyl;

the R₈-N-CO-R₇ group being trans to the R₅ group;


and X is oxygen or NR₁₀ where R₁₀ is hydrogen or C₁₋₆ alkyl.

2. A pharmaceutical composition for use in the treatment and/or
prophylaxis of anxiety and/or mania and/or depression and/or the
effects associated with withdrawal from substances of abuse and/or
disorders treatable or preventable with anti-convulsive agents,
which comprises an effective amount of a compound of formula (I),
or a pharmaceutically acceptable salt thereof, as defined in claim 1,
and a pharmaceutically acceptable carrier.

3. The use of a compound of formula (I), as defined in claim 1, or a
pharmaceutically acceptable salt thereof, for the manufacture of a
medicament for the treatment and/or prophylaxis of anxiety and/or
mania and/or depression and/or the effects associated with
withdrawal from substances of abuse and/or disorders treatable or
preventable with anti-convulsive agents.



This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

4. A method according to claim 1 in which the compound of formula (I) has the following variables wherein;

Y is C-R₁, where:

5

10

15

20

25

30

35

either one of R₁ and R₂ is hydrogen and the other is selected from the class of hydrogen, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy carbonyl, C₁₋₆ alkylcarbonyloxy, C₁₋₆ alkylhydroxymethyl, nitro, cyano, chloro, trifluoromethyl, trifluoromethoxy, C₁₋₆ alkylsulphinyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkoxy sulphinyl, C₁₋₆ alkoxy sulphonyl, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxy carbonylamino, C₁₋₆ alkyl-thiocarbonyl, C₁₋₆ alkoxy-thiocarbonyl, C₁₋₆ alkyl-thiocarbonyloxy, 1-mercapto C₂₋₇ alkyl, formyl, or aminosulphinyl, aminosulphonyl or aminocarbonyl, the amino moiety being optionally substituted by one or two C₁₋₆ alkyl groups, or C₁₋₆ alkylsulphinylamino, C₁₋₆ alkylsulphonylamino, C₁₋₆ alkoxy sulphinylamino or C₁₋₆ alkoxy sulphonylamino, or ethylenyl terminally substituted by C₁₋₆ alkylcarbonyl, nitro or cyano, or -C(C₁₋₆ alkyl)NOH or -C(C₁₋₆ alkyl)NNH₂, or one of R₁ and R₂ is nitro, cyano or C₁₋₃ alkylcarbonyl and the other is methoxy or amino optionally substituted by one or two C₁₋₆ alkyl or by C₂₋₇ alkanoyl;

one of R₃ and R₄ is hydrogen or C₁₋₄ alkyl and the other is C₁₋₄ alkyl or R₃ and R₄ together are C₂₋₅ polymethylene;

R₅ is hydroxy and R₆ is hydrogen;

R₇ is fluorophenyl;

R₈ is hydrogen or C₁₋₆ alkyl; and

R₉ is hydrogen;

the R₈-N-CO-R₇ group being trans to the R₅ group.

5. A pharmaceutical composition according to claim 2 wherein the compound of formula (I) is as defined in claim 4.




This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

6. A use according to claim 3 wherein the compound of formula (I) is as defined in claim 4.
- 5 7. A compound of formula (Ia) or a pharmaceutically acceptable salt thereof



- 10 wherein variables Y, R₂, R₃, R₄, R₅, R₆, R₈, R₉ and X are as defined in relation to formula (I) in claim 1 and R₇ is 2- or 3-fluorophenyl or where R₇ is 2-, 4- or 3,4- difluorophenyl.
8. A compound of formula (Ia) as defined in claim 7 selected from the group consisting of;
- 15

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(3-fluorobenzoylamino)2H-1-benzopyran-3-ol (Example 1),

- 20 trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(2-fluorobenzoylamino)2H-1-benzopyran-3-ol (Example 2),

trans-6-Trifluoromethoxy-3,4-dihydro-2,2-dimethyl-4-(3-fluorobenzoylamino)2H-1-benzopyran-3-ol (Example 3),

- 25 trans-6-Cyano-4S-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 21),

- 30 trans-6-Cyano-4R-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 22),

trans-6-Cyano-4S-(2-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 24),

This Sheet Has Been Replaced



trans-6-Cyano-4S-(2,4-difluorobenzoylamino)-3,4-dihydro-2,2-
dimethyl-2H-1-benzopyran-3R-ol (Example 28)


THIS SHEET HAS BEEN
 REPLACED

5 trans-6-Acetyl-4-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-
2H-1-benzopyran-3-ol (Example 31),

trans-6-Cyano-4-(3,4-difluorobenzoyl-methylamino)-3,4-dihydro-2,2-
dimethyl-2H-1-benzopyran-3-ol (Example 41) and

10 trans-6-cyano-4S-(3,4-difluorobenzoylamino)-3,4-dihydro-2,2-
dimethyl-2H-1-benzopyran-3R-ol (Example 44) or a
pharmaceutically acceptable salt thereof.

15 9. A compound of formula (Ib) or pharmaceutically acceptable salts
 thereof:




20 wherein variables R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 and X are as defined
 in relation to formula (I) in claim 1 and Y is CR_1 where R_1 and R_2
 are both hydrogen or one of R_1 and R_2 is trifluoromethoxy, C_{1-6}
 alkyl optionally interrupted with oxygen or substituted with
 hydroxy, C_{1-6} alkoxy or substituted amino-carbonyl, CF_3A -(where
 25 A is $-CF_2-$, $-CO-$, $-CH_2$) or $CH(OH)$, aryl sulphonyl, aryl C_{3-8}
 cycloalkyl, C_{1-6} alkoxy, heteroaryl, arylcarbonyl,
 heteroarylcarbonyl, arylsulphinyl, heteroarylsulphinyl,
 heteroarylsulphonyl, in which any aromatic moiety is optionally
 substituted and the other is hydrogen.

30 10. A compound of formula (Ib) as defined in claim 9 selected from the
 group consisting of:

trans-3,4-Dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)-
2H-1-benzopyran-3-ol (Example 4),

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2,6-trimethyl-2H-1-benzopyran-3-ol (Example 14),

5 trans-6-Ethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 15),

trans-6-Ethyl-4-(4-fluorobenzoylethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 16),

10 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-isopropyl-2H-1-benzopyran-3-ol (Example 17),

15 trans-6-Ethyl-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 23),

trans-4-(4-Fluorobenzoylamino)-6-pentafluoroethyl-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 29),

20 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-phenylsulphonyl-2H-1-benzopyran-3-ol (Example 32),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-phenyl-2H-1-benzopyran-3-ol (Example 33),

25 trans-6-Ethyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 35),

30 trans-4R-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 36),

trans-4S-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 37),

35 trans-4-(4-fluorobenzoyl-methylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 43),

trans-6-t-Butyl-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 46),



This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-cyclopentyl-2H-1-benzopyran-3-ol (Example 48) and

5 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxy-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 50), or a pharmaceutically acceptable salt thereof.

10 11. A compound of formula (Ic), or a pharmaceutically acceptable salt thereof:



15 where R₃, R₄, R₅, R₆, R₇, R₈, R₉ and X are as defined in relation to formula (I) in claim 1 and Y is N and R₂ is hydrogen.

12. A compound of formula (Ic) as defined in claim 11, selected from the group consisting of:

20 trans-4-(2-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 38),

25 trans-4-(3-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 39) and


30 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 40), or a pharmaceutically acceptable salt thereof.

13. A compound of formula (Id), or a pharmaceutically acceptable salt thereof:



This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

where X is NR₁₀ where R₁₀ is hydrogen or C₁₋₆ alkyl and Y, R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₉ are as defined in relation to formula (I) in claim 1.

14. A compound of formula (Id) as defined in claim 13, selected from the group consisting of:
- 10 trans-6-Cyano-4-(4-fluorobenzoylamino)-2,2-dimethyl-1,2,3,4-tetrahydroquinolin-3-ol (Example 45), or a pharmaceutically acceptable salt thereof.
15. A compound of formula (I') selected from the group consisting of:
- 15 trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 5),
- 20 trans-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 6),
- trans-6-Trifluoromethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 7),
- 25 trans-6-Cyano-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 8),
- trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 9),
- 30 trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylmethylamino)2H-1-benzopyran-3-ol, (Example 10),



This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

trans-6-Acetyl-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 11),

5 trans-6-Cyano-3,4-dihydro-4-(4-fluorobenzoylethylamino)-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 12),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-7-nitro-2H-1-benzopyran-3-ol (Example 13),

10 trans-6-Acetyl-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 18),

trans-6-Acetyl-4R-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 19),

15 trans-6-Acetyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 20),

20 trans-6-Cyano-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 25),

trans-6-Cyano-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2,3-trimethyl-2H-1-benzopyran-3-ol (Example 26),

25 trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-nitro-2H-1-benzopyran-3-ol (Example 27),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxycarbonyl-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 30),

30 trans-6-Fluoro-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 34),

35 trans-4-(4-fluorobenzoyl-methylamino)-6-trifluoromethyl-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 42),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-iodo-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 47) and



This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED

trans- 6-Aminocarbonylmethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 49) or a pharmaceutically acceptable salt thereof.

- 5
16. A compound of formula (Ia), (Ib), (Ic) or (Id), according to any one of claims 7, 9, 11 and 13, or a pharmaceutically acceptable salt thereof wherein R₃ and R₄ are both methyl.
- 10 17. A compound of formula (Ia), (Ib), (Ic) or (Id), according to any one of claims 7, 9, 11, 13 and 16, or a pharmaceutically acceptable salt thereof wherein R₅ is hydroxy and R₆ and R₉ are hydrogen.
- 15 18. A compound of formula (Ia), (Ib) or (Ic) according to any one of claims 7, 9, 11, 16 and 17, or a pharmaceutically acceptable salt thereof wherein X is oxygen.
- 20 19. A compound of formula (Ia), (Ib) or (Id) according to any one of claims 7, 9, 13 and 16 to 19, or a pharmaceutically acceptable salt thereof wherein Y is R₁ and R₁ is cyano, methoxy, trifluoromethoxy, chloro, trifluoromethyl, ethylcarbonyl, acetyl, hydrogen, methyl, ethyl, ~~iso-propyl~~, ~~tertiary-butyl~~, nitro, C₂F₅, methoxycarbonyl, phenylsulphonyl, phenyl, fluoro, iodo, cyclopentyl, aminocarbonylmethyl and 1-hydroxyethyl and R₂ is
- 25 hydrogen
20. A compound according to claim 19 wherein R₁ is cyano, ethyl or acetyl and R₂ is hydrogen.
- 30 21. A compound of formula (Ib), (Ic) or (Id) according to any one of claims 9, 11, 13 and 16 to 20, or a pharmaceutically acceptable salt thereof wherein R₇ is 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2,4-difluorophenyl or 3,4-difluorophenyl.
- 35 22. A compound of formula (Ia), (Ib), (Ic) or (Id), according to any one of claims 7, 9, 11, 13 and 16 to 21, or a pharmaceutically acceptable salt thereof wherein R₈ is hydrogen, methyl or ethyl.



This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED

23. A compound of formula (Ia), (Ib), (Ic) or (Id), according to any one of claims 7, 9, 11, 13 and 16 to 22, or a pharmaceutically acceptable salt thereof wherein the compound has the 4S,3R configuration.

5 24. A process for the preparation of compounds of formula (I') as listed in claim 15, or pharmaceutically acceptable salts thereof:

which comprises acylating a compound of formula (II'):




10 wherein Y, R2' and R5' are the required variables Y, R2 or R5 as defined in formula (I) or a group convertible thereto and R3, R4, R6, R8, R9 and X are the required variables as defined in formula (I), the R8NH group being trans to the R5 group, with and acylating agent of formula (IIIb):



20 where R7 is as required and as defined in formula (I) and L1 is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any R1', R2' and R5' groups to R1, R2 and R5 respectively, interconverting R8 when hydrogen to C1-6 alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

25 25. A process for the preparation of compounds of formula (Ia) as defined in claim 7, or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIa):



This Sheet Has Been Replaced




THIS SHEET HAS BEEN
REPLACED

wherein Y' , R_2' and R_5' are Y , R_2 or R_5 as defined in formula (I) or a group convertible thereto and R_3 , R_4 , R_6 , R_8 , R_9 and X are as defined in formula (I), the R_8NH group being trans to the R_5' group, with an acylating agent of formula (IIIa):



where R_7^a is 2- or 3-fluorophenyl or 2,4- or 3,4- difluorophenyl and L_1 is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any Y' , R_2' or R_5' group to Y , R_2 or R_5 respectively, interconverting R_8 when hydrogen to C_{1-6} alkyl, separating any entantioners and forming a pharmaceutically acceptable salt or solvate.


26. A process for the preparation of compounds of formula (Ib) as defined in claim 9, or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIb):



where R_1' and R_2' are both hydrogen or one of R_1 and R_2 is trifluoromethoxy, C_{1-6} alkyl optionally interrupted with oxygen or substituted with hydroxy, C_{1-6} alkoxy or substituted aminocarbonyl, CF_3A- (where A is CF_2 , $-CO-$, $-CH_2-$ or $CH(OH)$), aryl, aryl sulphonyl, aryl C_{3-8} cycloalkyl, C_{1-6} alkoxy, heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphonyl, heteroarylsulphonyl, heteroarylsulphonyl, in which any aromatic moiety is optionally substituted and the other is hydrogen, or groups convertible to any of these; R_5' is R_5 as defined in formula (I) or a group convertible thereto and R_3 , R_4 , R_6 , R_8 , R_9 and X are as defined in formula (I) the R_8NH group being trans to the R_5' group, with a compound of formula (IIIb):




This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

where R_7 is as defined in formula (I) and L_1 is a leaving group;
thereafter optionally or as necessary and in any appropriate order
converting any R_1' , R_2' and R_5' groups to R_1 , R_2 and R_5
respectively, interconverting R_8 when hydrogen to C_{1-6} alkyl,
separating any enantiomers and forming a pharmaceutically
acceptable salt or solvate.

27. A process for the preparation of compounds of formula (Ic) as
defined in claim 11, or a pharmaceutically acceptable salt thereof,
or a pharmaceutically acceptable salt thereof, which comprises
acylating a compound of formula (IIc):




in which Y is N, R_5' is R_5 , as defined in relation to formula (I) or a
group convertible to R_5 and R_2' is hydrogen or a group convertible
thereto, R_6 , R_3 , R_4 , R_8 , R_9 and X are as defined in formula (I), the
 R_8NH group being trans to the R_5' group, with an acylating agent
of formula (IIIb):



where R_7 is as defined in formula (I) and L_1 is a leaving group;
thereafter optionally or as necessary and in any appropriate order
converting any R_2' or R_5' group to hydrogen or R_5 respectively,
interconverting R_8 when hydrogen to C_{1-6} alkyl, separating any
enantiomers and forming a pharmaceutically acceptable salt or
solvate.




This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

28. A process for the preparation of compounds of formula (I) as defined in claim 11, or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIId):



- in which X' is NR₁₀' where R₁₀' is hydrogen or C₁₋₆ alkyl or a group convertible thereto, Y', R₂' and R₅' respectively as defined in formula (I) or groups convertible thereto and R₃, R₄, R₆, R₈ and R₉ are as defined in formula (I), the R₈NH group being trans to the R₅' group, with an acylating agent of formula (IIIb):



- where R₇ is as defined in formula (I) and L₁ is a leaving group, thereafter optionally or as necessary and in any appropriate order converting any Y', R₁₀', R₂' or R₅' group to Y, R₁₀, R₂ or R₅ respectively, interconverting R₈ when hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

29. A pharmaceutical composition comprising a compound of formula (I'), (Ia), (Ib), (Ic) or (Id), as defined in claims 15, 7, 9, 11 and 13 respectively, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
30. The use of a compound of formula (I'), (Ia), (Ib), (Ic) or (Id), as defined in claims 15, 7, 9, 11 and 13 respectively, or a pharmaceutically acceptable salt thereof, in the treatment and/or prophylaxis of anxiety and/or mania and/or depression and/or the effects associated with withdrawal from substances of abuse and/or disorders treatable and/or preventable with anti-convulsive agents.




This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED**Abstract**

A method of treatment and/or prophylaxis of anxiety and/or mania, and/or depression and/or the effects associated with withdrawal from substances of abuse, and/or disorders treatable and/or preventable with anti-convulsive agents, comprising administering to the sufferer in need thereof an effective or prophylactic amount of a compound of formula (I) or pharmaceutically acceptable salt thereof:



10 wherein:

either Y is N and R₂ is hydrogen, or Y is C-R₁

where:

15

either one of R₁ and R₂ is hydrogen and the other is selected from the class of hydrogen, C₃₋₈ cycloalkyl, C₁₋₆ alkyl optionally interrupted by oxygen or substituted by hydroxy, C₁₋₆ alkoxy or substituted aminocarbonyl, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy carbonyl, C₁₋₆ alkylcarbonyloxy, C₁₋₆ alkoxy, nitro, cyano, halo, trifluoromethyl, CF₃S, or a group CF₃-A-, where A is -CF₂-, -CO-, -CH₂- or CH(OH), trifluoromethoxy, C₁₋₆ alkylsulphinyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkoxy sulphinyl, C₁₋₆ alkoxy sulphonyl, aryl, heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphinyl, heteroarylsulphinyl, arylsulphonyl, heteroarylsulphonyl in which any aromatic moiety is optionally substituted, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxy carbonylamino, C₁₋₆ alkyl-thiocarbonyl, C₁₋₆ alkoxy-thiocarbonyl, C₁₋₆ alkyl-thiocarbonyloxy, 1-mercapto C₂₋₇ alkyl, formyl, or aminosulphinyl, aminosulphonyl or aminocarbonyl, any amino moiety being optionally substituted by one or two C₁₋₆ alkyl groups, or C₁₋₆ alkylsulphinylamino, C₁₋₆ alkylsulphonylamino, C₁₋₆ alkoxy sulphinylamino or C₁₋₆ alkoxy sulphonylamino, or ethylenyl terminally substituted by C₁₋₆ alkylcarbonyl, nitro or cyano, or -C(C₁₋₆ alkyl)NOH or -C(C₁₋₆ alkyl)NNH₂, or one of R₁ and R₂ is nitro, cyano or C₁₋₃ alkylcarbonyl and

This Sheet Has Been Replaced



the other is methoxy or amino optionally substituted by one or two C₁₋₆ alkyl or by C₂₋₇ alkanoyl;

- 5 one of R₃ and R₄ is hydrogen or C₁₋₄ alkyl and the other is C₁₋₄ alkyl or R₃ and R₄ together are C₂₋₅ polymethylene;

- 10 R₅ is C₁₋₆ alkylcarbonyloxy, benzoyloxy, ONO₂, benzyloxy, phenyloxy or C₁₋₆ alkoxy and R₆ and R₉ are hydrogen or R₅ is hydroxy and R₆ is hydrogen or C₁₋₂ alkyl and R₉ is hydrogen;

R₇ is fluorophenyl;

R₈ is hydrogen or C₁₋₆ alkyl;

- 15 the R₈-N-CO-R₇ group being trans to the R₅ group;

and X is oxygen or NR₁₀ where R₁₀ is hydrogen or C₁₋₆ alkyl.

This Sheet Has Been Replaced



9608698-8

3/05 copy

REPLACEMENT SHEET

06 MAR 1997

-1-

BENZO-AND PYRIDOPYRAN DERIVATIVES HAVING ANXIOLYTIC
AND ANTI-CONVULSANT ACTIVITY

This invention relates to a novel method of treatment and to novel compounds for use in such a method.

European Published Patent Application No. 0126311 discloses substituted benzopyran compounds having blood pressure lowering activity, including 6-acetyl-trans-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol.

Also EP-A-0 376 524, EP-A-0 205 292, EP-A-0 250 077, EP-A-0 093 535, EP-A-0 150 202, EP-A-0 076 075 and WO/89/05808 (Beecham Group plc) describe certain benzopyran derivatives which possess anti-hypertensive activity.

EP-A-0 350 805 (Biersdorf), EP-A-0 277 611, EP-A-0 277612, EP-A-0 337 179 and EP-A-0 355 565 (Hoechst Aktiengesellschaft); and EP-A-0 466 131 (Nissan Chemical Industries Ltd) also describe certain benzopyran derivatives which are believed to possess anti-hypertensive activity.

EP-A-0 430 621 and EP-A-0 385 584 (Beecham Group plc) describe the resolution of certain intermediates useful in the preparation of the compounds described in the above mentioned patent applications.


EP-A-0 194 885 (E. Lilly) describes certain amino substituted benzopyran derivatives possessing anti-convulsant activity.

It has now been surprisingly found that certain compounds of formula (I) possess anxiolytic and anti-convulsant activity, and are also believed to have utility in the treatment or prevention of mania, depression and the effects associated with withdrawal from substances of abuse.

Accordingly, the present invention provides a method of treatment and/or prophylaxis of anxiety and/or mania, and/or depression and/or the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, and/or disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy, comprising

06 MAR 1997

administering to the sufferer in need thereof an effective or prophylactic amount of a compound of formula (I) or pharmaceutically acceptable salt thereof:



wherein:

either Y is N and R₂ is hydrogen, or Y is C-R₁

where:

either one of R₁ and R₂ is hydrogen and the other is selected from the class of hydrogen, C₃₋₈ cycloalkyl, C₁₋₆ alkyl optionally interrupted by oxygen or substituted by hydroxy, C₁₋₆ alkoxy or substituted aminocarbonyl, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylcarbonyloxy, C₁₋₆ alkoxy, nitro, cyano, halo, trifluoromethyl, CF₃S, or a group CF₃-A-, where A is -CF₂-, -CO-, -CH₂- or CH(OH), trifluoromethoxy, C₁₋₆ alkylsulphinyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkoxysulphinyl, C₁₋₆ alkoxysulphonyl, aryl, heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphinyl, heteroarylsulphinyl, arylsulphonyl, heteroarylsulphonyl in which any aromatic moiety is optionally substituted, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxycarbonylamino, C₁₋₆ alkyl-thiocarbonyl, C₁₋₆ alkoxy-thiocarbonyl, C₁₋₆ alkyl-thiocarbonyloxy, 1-mercapto C₂₋₇ alkyl, formyl, or aminosulphinyl, aminosulphonyl or aminocarbonyl, any amino moiety being optionally substituted by one or two C₁₋₆ alkyl groups, or C₁₋₆ alkylsulphinylamino, C₁₋₆ alkylsulphonylamino, C₁₋₆ alkoxysulphinylamino or C₁₋₆ alkoxysulphonylamino, or ethylenyl terminally substituted by C₁₋₆ alkylcarbonyl, nitro or cyano, or -C(C₁₋₆ alkyl)NOH or -C(C₁₋₆ alkyl)NNH₂, or one of R₁ and R₂ is nitro, cyano or C₁₋₃ alkylcarbonyl and the other is methoxy or amino optionally substituted by one or two C₁₋₆ alkyl or by C₂₋₇ alkanoyl;

06 MAR 1997

one of R_3 and R_4 is hydrogen or C_{1-4} alkyl and the other is C_{1-4} alkyl or R_3 and R_4 together are C_{2-5} polymethylene;

R_5 is C_{1-6} alkylcarbonyloxy, benzoyloxy, ONO_2 , benzyloxy, phenyloxy or C_{1-6} alkoxy and R_6 and R_9 are hydrogen or R_5 is hydroxy and R_6 is hydrogen or C_{1-2} alkyl and R_9 is hydrogen;

R_7 is fluorophenyl;

R_8 is hydrogen or C_{1-6} alkyl;

the $R_8-N-CO-R_7$ group being trans to the R_5 group;


and X is oxygen or NR_{10} where R_{10} is hydrogen or C_{1-6} alkyl.

Compounds of formula (I) include those in which Y is $C-R_1$, where;

either one of R_1 and R_2 is hydrogen and the other is selected from the class of hydrogen, C_{1-6} alkylcarbonyl, C_{1-6} alkoxycarbonyl, C_{1-6} alkylcarbonyloxy, C_{1-6} alkylhydroxymethyl, nitro, cyano, chloro, trifluoromethyl, trifluoromethoxy, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, C_{1-6} alkoxysulphinyl, C_{1-6} alkoxysulphonyl, C_{1-6} alkylcarbonylamino, C_{1-6} alkoxycarbonylamino, C_{1-6} alkyl-thiocarbonyl, C_{1-6} alkoxy-thiocarbonyl, C_{1-6} alkyl-thiocarbonyloxy, 1-mercapto C_{2-7} alkyl, formyl, or aminosulphinyl, aminosulphonyl or aminocarbonyl, the amino moiety being optionally substituted by one or two C_{1-6} alkyl groups, or C_{1-6} alkylsulphinylamino, C_{1-6} alkylsulphonylamino, C_{1-6} alkoxysulphinylamino or C_{1-6} alkoxysulphonylamino, or ethylenyl terminally substituted by C_{1-6} alkylcarbonyl, nitro or cyano, or $-C(C_{1-6} \text{ alkyl})NOH$ or $-C(C_{1-6} \text{ alkyl})NNH_2$, or one of R_1 and R_2 is nitro, cyano or C_{1-3} alkylcarbonyl and the other is methoxy or amino optionally substituted by one or two C_{1-6} alkyl or by C_{2-7} alkanoyl;

one of R_3 and R_4 is hydrogen or C_{1-4} alkyl and the other is C_{1-4} alkyl or R_3 and R_4 together are C_{2-5} polymethylene;

R_5 is hydroxy and R_6 is hydrogen;



R₇ is fluorophenyl;

R₈ is hydrogen or C₁₋₆ alkyl; and

R₉ is hydrogen;

the R₈-N-CO-R₇ group being trans to the R₅ group.


All C₁₋₆ alkyl or alkyl containing groups in formula (I) are preferably selected from methyl, ethyl, *n*- and *iso*-propyl, *n*-, *iso*-, *sec*- and *tert*-butyl.

Suitable C₃₋₈ cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

Aryl includes but is not limited to phenyl and naphthyl.

Heteroaryl includes a 5- or 6- membered monocyclic or 9- or 10- membered bicyclic of which 5- or 6- membered monocyclic heteroaryl is preferred. In addition, 5- or 6-membered monocyclic or 9- or 10-membered bicyclic heteroaryl preferably contains one, two or three heteroatoms which are selected from the class of oxygen, nitrogen and sulphur and which, in the case of there being more than one heteroatom, are the same or different. Examples of 5- or 6-membered monocyclic heteroaryl containing one, two or three heteroatoms which are selected from the class of oxygen, nitrogen and sulphur include furyl, thienyl, pyrrol, oxazolyl, thiazolyl, imidazolyl and thiadiazolyl, and pyridyl, pyridazyl, pyrimidyl, pyrazolyl and triazolyl. Preferred examples of such groups include furanyl, thienyl, pyrrol and pyridyl, in particular 2- and 3-furyl, 2- and 3-pyrrol, 2- and 3-thienyl, and 2-, 3- and 4-pyridyl. Examples of 9- or 10-membered bicyclic heteroaryl containing one, two or three heteroatoms which are selected from the class of oxygen, nitrogen and sulphur include benzofuranyl, benzothienyl, indolyl and indazolyl, quinolyl and isoquinolyl, and quinazolyl. Preferred examples of such groups include 2- and 3-benzofuryl, 2- and 3-benzothienyl, and 2- and 3-indolyl, and 2- and 3-quinolyl.

Suitable examples of groups or atoms for optional substitution of aryl and heteroaryl include one, two or three substituents independently selected



from C₁₋₄ alkyl, C₁₋₄ alkoxy, halo (such as fluoro, chloro, bromo), hydroxy, nitro, cyano and SO_nH, where n=0 to 2.

Preferably R₂ is hydrogen.

Examples of suitable R₁ substituents include cyano, methoxy, trifluoromethoxy, chloro, trifluoromethyl, ethylcarbonyl, acetyl, hydrogen, methyl, ethyl, iso-propyl, tertiary-butyl, nitro, C₂F₅, methoxycarbonyl, phenylsulphonyl, phenyl, fluoro, iodo, cyclopentyl, aminocarbonylmethyl and 1-hydroxyethyl. Preferably R₁ is cyano, acetyl or ethyl.

Preferably R₃ and R₄ are both methyl.

Preferably R₅ is hydroxy and R₆ and R₉ are hydrogen or R₅ is hydroxy, R₆ is C₁₋₂ alkyl and R₉ is hydrogen, more preferably R₅ is hydroxy and R₆ and R₉ are hydrogen.

It should be appreciated that the term fluorophenyl relating to R₇ encompasses phenyl which has 1,2,3,4 or 5 fluoro groups attached to the phenyl ring. Preferably there are 1 or 2 fluoro groups attached to the phenyl ring and most preferably there is 1 fluoro group attached to the ring.

The fluoro group or groups may be in any position around the phenyl ring, preferably in the case of mono fluorophenyl the fluoro is in the 3 or 4 position.

In the case of difluorophenyl, preferably the fluoro substituents are at positions 2,4 or 3,4.

Preferably R₈ is hydrogen or C₁₋₄ alkyl, more preferably R₈ is hydrogen, methyl or ethyl.

Preferably X is oxygen.

It should be appreciated that the compounds of formula (I) may have chiral carbon atoms at positions 2, 3 and 4 and therefore may exist as enantiomers. The present invention extends to each enantiomer and to

06 MAR 1997

-6-

mixtures thereof including racemates. Preferably the compounds of formula (I) exist as 4S, 3R enantiomers.

It should also be appreciated that certain R_1 substituents also have chiral centres and therefore may exist as enantiomers. The present invention extends to each enantiomer and to mixtures thereof including racemates

An example of a compound of formula (I) is trans-6-acetyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (compound X).

The following novel compounds of formula (I) also form an aspect of the present invention, (herein referred to as compounds of formula (I')):

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 5),

trans-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 6),

trans-6-Trifluoromethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 7),

trans-6-Cyano-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 8),

trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 9),

trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylmethylamino)2H-1-benzopyran-3-ol, (Example 10),

trans-6-Acetyl-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 11),

trans-6-Cyano-3,4-dihydro-4-(4-fluorobenzoylethylamino)-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 12),

06 MAR 1997

-7-

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-7-nitro-2H-1-benzopyran-3-ol (Example 13),

trans-6-Acetyl-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 18),

trans-6-Acetyl-4R-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 19),

trans-6-Acetyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 20),

trans-6-Cyano-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 25),

trans-6-Cyano-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2,3-trimethyl-2H-1-benzopyran-3-ol (Example 26),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-nitro-2H-1-benzopyran-3-ol (Example 27),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxycarbonyl-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 30),


trans-6-Fluoro-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 34),

trans-4-(4-fluorobenzoyl-methylamino)-6-trifluoromethyl-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 42),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-iodo-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 47) and

trans-6-Aminocarbonylmethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 49).

The administration to the mammal may be by way of oral or parenteral administration.



An amount effective to treat the disorders hereinbefore described depends on the usual factors such as the nature and severity of the disorders being treated and the weight of the mammal. However, a unit dose will normally contain 1 to 1000 mg, suitably 1 to 500 mg, for example an amount in the range of from 2 to 100 mg such as 2, 5, 10, 20, 30, 40, 50 and 100 mg of the active compound. Unit doses will normally be administered once or more than once per day, for example 2, 3, 4, 5 or 6 times a day, more usually 2 to 4 times a day, such that the total daily dose is normally in the range, for a 70 kg adult of 1 to 1000 mg, for example 50 to 500 mg, that is in the range of approximately 0.01 to 50 mg/kg/day, more usually 0.1 to 50 mg/kg/day, for example 1 to 50 mg/kg/day.

It is greatly preferred that the compound of formula (I) is administered in the form of a unit-dose composition, such as a unit dose oral, rectal, topical or parenteral (especially intravenous) composition.

Such compositions are prepared by admixture and are suitably adapted for oral or parenteral administration, and as such may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable and infusible solutions or suspensions or suppositories. Orally administrable compositions are preferred, in particular shaped oral compositions, since they are more convenient for general use.

Tablets and capsules for oral administration are usually presented in a unit dose, and contain conventional excipients such as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, colourants, flavourings, and wetting agents. The tablets may be coated according to well known methods in the art.

Suitable fillers for use include cellulose, mannitol, lactose and other similar agents. Suitable disintegrants include starch, polyvinylpyrrolidone and starch derivatives such as sodium starch glycollate. Suitable lubricants include, for example, magnesium stearate. Suitable pharmaceutically acceptable wetting agents include sodium lauryl sulphate.

06 MAR 1997

These solid oral compositions may be prepared by conventional methods of blending, filling, tableting or the like. Repeated blending operations may be used to distribute the active agent throughout those compositions employing large quantities of fillers. Such operations are, of course, conventional in the art.

Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethyl cellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example, almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl *p*-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

Oral formulations also include conventional sustained release formulations, such as tablets or granules having an enteric coating.

For parenteral administration, fluid unit dose forms are prepared containing the compound and a sterile vehicle. The compound, depending on the vehicle and the concentration, can be either suspended or dissolved. Parenteral solutions are normally prepared by dissolving the compound in a vehicle and filter sterilising before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are also dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum.

Parenteral suspensions are prepared in substantially the same manner except that the compound is suspended in the vehicle instead of being dissolved and sterilised by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is

06 MAR 1997

included in the composition to facilitate uniform distribution of the compound of the invention.

As is common practice, the compositions will usually be accompanied by written or printed directions for use in the medical treatment concerned.

The present invention further provides a pharmaceutical composition for use in the treatment and/or prophylaxis of anxiety and/or mania and/or depression and/or the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines and/or disorders treatable or preventable with anti-convulsive agents, such as epilepsy, which comprises a compound of formula (I), or a pharmaceutically acceptable salt thereof, in particular, compounds of examples 1 to 50, and a pharmaceutically acceptable carrier.

In a further aspect the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, in particular compounds of examples 1 to 50 inclusive, for the manufacture of a medicament for the treatment and/or prophylaxis of anxiety and/or mania and/or depression and/or the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines and/or disorders treatable or preventable with anti-convulsive agents, such as epilepsy.

Such compositions may be prepared in the manner as hereinbefore described.

The invention also provides novel compounds of formula (I) and pharmaceutically acceptable salts thereof wherein R₇ is 2- or 3-fluorophenyl, 2, 4- or 3,4- difluorophenyl. Such compounds will hereinafter be referred to as compounds of formula (Ia).

Examples of compounds of formula (Ia) include:

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(3-fluorobenzoylamino)2H-1-benzopyran-3-ol (Example 1),

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(2-fluoro-

06 MAR 1997

benzoylamino)2H-1-benzopyran-3-ol (Example 2),

trans-6-Trifluoromethoxy-3,4-dihydro-2,2-dimethyl-4-(3-fluorobenzoylamino)2H-1-benzopyran-3-ol (Example 3),

trans-6-Cyano-4S-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 21),

trans-6-Cyano-4R-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 22),

trans-6-Cyano-4S-(2-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 24),

trans-6-Cyano-4S-(2,4-difluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 28),

trans-6-Acetyl-4-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 31),

trans-4-(2-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 38),

trans-4-(3-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 39),

trans-6-Cyano-4-(3,4-difluorobenzoyl-methylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 41) and

trans-6-cyano-4S-(3,4-difluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 44).

The invention further provides novel compounds of formula (I) and pharmaceutically acceptable salts thereof wherein Y is CR₁ where R₁ and R₂ are both hydrogen or one of R₁ and R₂ is trifluoromethoxy, C₁₋₆ alkyl optionally interrupted with oxygen or substituted with hydroxy, C₁₋₆ alkoxy or substituted amino-carbonyl, CF₃A-(where A is -CF₂-, -CO-, -CH₂) or CH(OH)), aryl sulphonyl, aryl C₃₋₈ cycloalkyl, C₁₋₆ alkoxy,

06 MAR 1997

heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphanyl, heteroarylsulphanyl, heteroarylsulphonyl, in which any aromatic moiety is optionally substituted and the other is hydrogen. Such compounds will hereinafter be referred to as compounds of formula (Ib).

Examples of compounds of formula (Ib) include:

trans-6-Trifluoromethoxy-3,4-dihydro-2,2-dimethyl-4-(3-fluorobenzoylamino)-2H-1-benzopyran-3-ol (Example 3),

trans-3,4-Dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)-2H-1-benzopyran-3-ol (Example 4),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2,6-trimethyl-2H-1-benzopyran-3-ol (Example 14),

trans-6-Ethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 15),

trans-6-Ethyl-4-(4-fluorobenzoyl-ethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 16),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-isopropyl-2H-1-benzopyran-3-ol (Example 17),

trans-6-Ethyl-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 23),

trans-4-(4-Fluorobenzoylamino)-6-pentafluoroethyl-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 29),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-phenylsulphonyl-2H-1-benzopyran-3-ol (Example 32),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-phenyl-2H-1-benzopyran-3-ol (Example 33),

06 MAR 1997

trans-6-Ethyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 35),

trans-4R-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 36),

trans-4S-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 37),

trans-4-(4-fluorobenzoyl-methylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 43),

trans-6-t-Butyl-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 46),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-cyclopentyl-2H-1-benzopyran-3-ol (Example 48) and

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxy-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 50).

The present invention also provides novel compounds of formula (I) and pharmaceutically acceptable salts thereof, where Y is N and R₂ is hydrogen. Such compounds will hereinafter be referred to as compounds of formula (Ic).

Examples of compounds of formula (Ic) include:

trans-4-(2-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 38),

trans-4-(3-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 39) and

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 40).

06 MAR 1997

-14-


The present invention also provides novel compounds of formula (I) and pharmaceutically acceptable salts thereof, where X is NR₁₀ where R₁₀ is hydrogen or C₁₋₆ alkyl. Such compounds will hereinafter be referred to as compounds of formula (Id).

An example of a compound of formula (Id) is:

trans-6-Cyano-4-(4-fluorobenzoylamino)-2,2-dimethyl-1,2,3,4-tetrahydroquinolin-3-ol (Example 45).

Generally, compounds of formula (I) may be prepared by procedures generally described or analogous to those described in EP-0126311, EP-0376524, EP-205292, EP-0250077, EP-0093535, EP-0150202, EP-0076075, WO/89/05808, EP-0350805, EP-0277611, EP-0277612, EP-0337179, EP-0355565 and EP-0466131.

The invention also provides a process for the preparation of novel compounds of formula (I') or pharmaceutically acceptable salts thereof, which comprises acylating a compound of formula (II'):



wherein Y, R₂' and R₅' are the required variables Y, R₂ or R₅ as defined in formula (I) or a group convertible thereto and R₃, R₄, R₆, R₈, R₉ and X are the required variables as defined in formula (I), the R₈NH group being trans to the R₅ group, with and acylating agent of formula (IIIb):




where R₇ is as required and as defined in formula (I) and L₁ is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any R₁', R₂' and R₅' groups to R₁, R₂ and R₅ respectively, interconverting R₈ when hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

06 MAR 1997

- 15 -

In particular, the present invention also provides a process for the preparation of compounds of formula (Ia), or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIa):




wherein Y', R2' and R5' are Y, R2 or R5 as defined in formula (I) or a group convertible thereto and R3, R4, R6, R8, R9 and X are as defined in formula (I), the R8NH group being trans to the R5' group, with an acylating agent of formula (IIIa):



where R7^a is 2- or 3-fluorophenyl or 2,4- or 3,4- difluorophenyl and L₁ is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any Y', R2' or R5' group to Y, R2 or R5 respectively, interconverting R8 when hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

The present invention also provides a process for the preparation of compounds of formula (Ib), or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIb):



where R1' and R2' are both hydrogen or one of R1 and R2 is trifluoromethoxy, C₁₋₆ alkyl optionally interrupted with oxygen or substituted with hydroxy, C₁₋₆ alkoxy or substituted aminocarbonyl, CF₃A- (where A is CF₂-, CO-, -CH₂- or CH(OH)), aryl, aryl sulphonyl, aryl

06 MAR 1997


-16-

C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphinyl, heteroarylsulphinyl, heteroarylsulphonyl, in which any aromatic moiety is optionally substituted and the other is hydrogen, or groups convertible to any of these; R_{5'} is R₅ as defined in formula (I) or a group convertible thereto and R₃, R₄, R₆, R₈, R₉ and X are as defined in formula (I) the R₈NH group being trans to the R_{5'} group, with a compound of formula (IIIb):



where R₇ is as defined in formula (I) and L₁ is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any R_{1'}, R_{2'} and R_{5'} groups to R₁, R₂ and R₅ respectively, interconverting R₈ when hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

The present invention also provides a process for the preparation of compounds of formula (Ic), or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIc):




in which Y is N, R_{5'} is R₅, as defined in relation to formula (I) or a group convertible to R₅ and R_{2'} is hydrogen or a group convertible thereto, R₆, R₃, R₄, R₈, R₉ and X are as defined in formula (I), the R₈NH group being trans to the R_{5'} group, with an acylating agent of formula (IIIb):



where R₇ is as defined in formula (I) and L₁ is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any R_{2'} or R_{5'} group to hydrogen or R₅ respectively, interconverting R₈ when

hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

The present invention also provides a process for the preparation of compounds of formula (Id), or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIId):



in which X' is NR_{10'} where R_{10'} is hydrogen or C₁₋₆ alkyl or a group convertible thereto, Y', R_{2'} and R₅ are Y, R₂ and R₅ respectively as defined in formula (I) or groups convertible thereto and R₃, R₄, R₆, R₈ and R₉ are as defined in formula (I), the R₈NH group being trans to the R_{5'} group, with an acylating agent of formula (IIIb):



where R₇ is as defined in formula (I) and L₁ is a leaving group, thereafter optionally or as necessary and in any appropriate order converting any Y', R_{10'}, R_{2'} or R_{5'} group to Y, R₁₀, R₂ or R₅ respectively, interconverting R₈ when hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

Examples of suitable leaving groups L₁ include those mentioned in the above-mentioned patents, in particular EP-A-0 126 311 or are conventional in the art.

The reaction conditions which may be used to carry out the above reactions are as outlined or analogous to those described in the above-mentioned patents, in particular EP-A-0 126 311.

In particular, the leaving group (L₁) is a group that is displaceable by a primary or secondary amino nucleophile. Examples of such a group include C₁₋₄ alkylcarbonyloxy and halogen, such as chloro and bromo. When the leaving group (L₁) is any of these examples, the acylating agent

06 MAR 1997

of formula (IIIa) or (IIIb) is either an acid anhydride or an acid halide. When it is an acid anhydride, it is, preferably, a mixed anhydride, which may be prepared in situ from an aromatic or heteroaromatic carboxylic acid and an alkyl chlorocarbonate, such as ethyl chloroformate.

When the acylating agent of formula (IIIa) or (IIIb) is an acid anhydride, the acylation of the compound formula (IIa), (IIb), (IIc) or (IId) is, preferably carried out using the anhydride as the solvent in the presence of an acid acceptor, such as sodium acetate.

When the acylating agent of formula (IIIa) or (IIIb) is an acid halide, the acylation of the compound of formula (IIa), (IIb), (IIc) or (IId), is, preferably, carried out in a non-aqueous medium, such as methylene chloride, in the presence of an acid acceptor, such as triethylamine, trimethylamine or pyridine.

Examples of suitable groups convertible to Y (or R₁), R₂ and R₅ include those described in the above-mentioned patents or are conventional in the art.

Interconversions of R₈ when hydrogen to C₁₋₆ alkyl may be carried out using conventional alkylation procedures for example using an alkylhalide in the presence of a base.

It should be appreciated that racemates for formula (I) may be resolved or enantiomerically purified compounds of formula (I) may be prepared using procedures conventional in the art and in particular using the procedures outlined in EP-0430631 and EP-0355584.

Suitably, the procedures outlined in or analogous to those described in Example 35 of the present specification may be used to prepare specific enantiomers of any compounds of formulae (I), (I'), (Ia), (Ib), (Ic) or (Id).

Compounds of formulae (I'), (IIa), (IIb), (IIc) and (IId) may be prepared from readily available starting materials using the procedures outlined or analogous to those described in the above-mentioned patents.

06 MAR 1987

-19-

Compounds of formulae (IIIa) and (IIIb) are either commercially available or may be prepared according to conventional procedures known in the art of organic chemistry.

Novel compounds of formulae (I'), (IIa), (IIb), (IIc) and (IId) form an aspect of the present invention.

Compounds of formula (I) in which R₅ is hydroxy, R₆ is C₁₋₂ alkyl and R₉ is hydrogen may be prepared according to the procedures outlined in R. Gericke *et al.* J. Med. Chem. Vol.34, p3074(1991).

The invention also provides a pharmaceutical composition comprising a compound of formula (I'), (Ia), (Ib), (Ic) or (Id), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The invention also provides the use of a compound of formula (I'), (Ia), (Ib), (Ic) or (Id), or a pharmaceutically acceptable salt thereof, in the treatment and/or prophylaxis of anxiety and/or mania and/or depression and/or the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines and/or disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy.

The following examples and pharmacological test results illustrate the present invention:

The following compounds were prepared by methods analogous to those described in the abovementioned patents publications.

05 MAR 1997

Example 1

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(3-fluoro-benzoylamino)2H-1-benzopyran-3-ol.

Mpt. 193-194°C.

Example 2

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(2-fluoro-benzoylamino)2H-1-benzopyran-3-ol

Mpt. 163-5°

Example 3

trans-6-Trifluoromethoxy-3,4-dihydro-2,2-dimethyl-4-(3-fluorobenzoylamino)2H-1-benzopyran-3-ol

Mpt. 127-30°C

Example 4

trans-3,4-Dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)-2H-1-benzopyran-3-ol


Mpt. 174-5°C

Example 5

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(4-fluoro-benzoylamino)2H-1-benzopyran-3-ol,

Mpt. 229-230°C.

3R, 4S isomer (compound B) Mpt. 224-5°C



Example 6

trans-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol,

Mpt. 177-9°C

Example 7

trans-6-Trifluoromethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol,

Mpt. 185-7°C

Example 8

trans-6-Cyano-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol,

Mpt. 230-4°C

Example 9



trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol,

Mpt. 205-207°C.

Example 10

trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylmethylamino)2H-1-benzopyran-3-ol,

Mpt. 210-212°C.



Example 11

trans-6-Acetyl-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol,

Mpt. 207-8°C

Example 12

trans-6-Cyano-3,4-dihydro-4-(4-fluorobenzoylethylamino)-2,2-dimethyl-2H-1-benzopyran-3-ol

mp 172-175 °C

Example 13

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-7-nitro-2H-1-benzopyran-3-ol

mp 231-233 °C

Example 14

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2,6-trimethyl-2H-1-benzopyran-3-ol

mp 185-186 °C

Example 15

trans-6-Ethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol

mp 235-237 °C

Example 16

trans-6-Ethyl-4-(4-fluorobenzoylethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol

mp 175 °C

06 MAR 1997

Example 17

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-isopropyl-2H-1-benzopyran-3-ol
mp 235-236 °C

Example 18

trans-6-Acetyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 203 °C

Example 19

trans-6-Acetyl-4R-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol
mp 162-163 °C

Example 20

trans-6-Cyano-4S-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol
mp 164 °C

Example 21

trans-6-Cyano-4S-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol
mp 221 °C

Example 22

trans-6-Cyano-4R-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol
mp 221 °C

06 MAR 1987

Example 23

trans-6-Ethyl-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 162-164 °C

Example 24

trans-6-Cyano-4S-(2-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol
mp 163-165 °C

Example 25

trans-6-Cyano-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol
mp 224-225 °C

Example 26

trans-6-Cyano-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2,3-trimethyl-2H-1-benzopyran-3-ol
mp 218-220 °C

Example 27

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-nitro-2H-1-benzopyran-3-ol
mp 197-198 °C

Example 28

trans-6-Cyano-4S-(2,4-difluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol
mp 175-176 °C

06 MAR 1997

Example 29

trans-4-(4-Fluorobenzoylamino)-6-pentafluoroethyl-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 181 °C

Example 30

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxycarbonyl-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 198 °C

Example 31

trans-6-Acetyl-4-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 212 °C

Example 32


trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-phenylsulphonyl-2H-1-benzopyran-3-ol
mp 239-240 °C

Example 33

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-phenyl-2H-1-benzopyran-3-ol
mp 164-165 °C

Example 34

trans-6-Fluoro-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 164-165 °C



- 26 -


Example 35

trans-6-Ethyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol

(-)-Mandelic acid was used to resolve the residue from resolution of trans-4-amino-6-ethyl-2,2-dimethyl-chroman-3-ol (to give the 3S, 4R enantiomer - as described in EPA 412 760) under standard resolving conditions using acetone as recrystallisation solvent. This furnished trans-4S-amino-6-ethyl-2,2-dimethylchroman-3R-ol D-mandelate.

The 4S, 3R-aminoalcohol mandelate (4.0 g) was dissolved in dichloromethane (250 ml) and triethylamine (3.34 ml) and cooled in an ice bath. To this stirred solution was added 4-fluorobenzoyl chloride (1.7 g) dropwise. On completion of addition the reaction mixture was allowed to attain room temperature and was stirred overnight. The solvent was evaporated and the residue was taken up in ethyl acetate (150 ml). This solution was washed with 5% sodium bicarbonate solution, and brine, and dried over anhydrous magnesium sulphate. Filtration and evaporation and recrystallisation of the residue from Ethyl acetate-hexane gave the title compound of example 35. mp 132-136°C. $\alpha_D/20 + 69.1^\circ$ (methanol, $c=1.0$).

NMR(CDCL ₃) δ	1.21 (3H, t, $J = 8$ Hz)
	1.28 (3H, s)
	1.49 (3H, s)
	2.88 (2H, q, $J = 8$ Hz)
	3.76 (1H, d, $J = 9$ Hz)
	4.46 (1H, broad s)
	5.21 (1H, t, $J = 9,8$ Hz)
	6.41 (1H, d, $J = 8$ Hz)
	6.80 (1H, d, $J = 9$ Hz)
	7.07 (2H, irregular m)
	7.17 (2H, t, $J = 9$ Hz)
	7.74 (2H, m)



Example 36

trans-4R-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-dimethyl-2H-1-benzopyran-3S-ol
mp 132-133 °C

Example 37

trans-4S-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-dimethyl-2H-1-benzopyran-3R-ol
mp 133-134 °C

Example 38

trans-4-(2-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol
mp 254 °C

Example 39

trans-4-(3-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol
mp 259-261 °C

Example 40

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol
mp 254-255 °C

Example 41

trans-6-Cyano-4-(3,4-difluorobenzoyl-methylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 199-200°C

06 MAR 1997

-28-

Example 42

trans-4-(4-fluorobenzoyl-methylamino)-6-trifluoromethyl-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 196-197°C

Example 43

trans-4-(4-fluorobenzoyl-methylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 160-161°C

Example 44

trans-6-cyano-4S-(3,4-difluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol
mp 227°C

Example 45

trans-6-Cyano-4-(4-fluorobenzoylamino)-2,2-dimethyl-1,2,3,4-tetrahydroquinolin-3-ol
mp 244-248 °C

Example 46

trans-6-t-Butyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 163-166 °C

Example 47

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-iodo-2,2-dimethyl-2H-1-benzopyran-3-ol
mp 204-205 °C

REPLACEMENT SHEET

06 MAR 1957

-29-

Example 48


trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-cyclopentyl-
2H-1-benzopyran-3-ol
mp 173-174 °C

Example 49

trans- 6-Aminocarbonylmethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-
dimethyl-2H-1-benzopyran-3-ol
mp 190 °C

Example 50

trans- 4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxy-2,2-dimethyl-2H-
1-benzopyran-3-ol
mp 155-156 °C



PHARMACOLOGICAL DATA

1. Geller-Seifter Procedure

Potential anxiolytic properties are evaluated using the Geller-Seifter procedure based on that originally described by Geller and Seifter, (1960) *Psychopharmacologia*, 1, 482-492. This procedure has been shown to be selective for drugs with anxiolytic properties (Cook and Sepinwall, (1975) "Mechanism of Action of Benzodiazepines" ed. Costa, E. and Greengard, P., Raven Press, New York, pp. 1-28).

Rats are trained on a variable interval 30 sec schedule (VI30) to press a lever in order to obtain food reward. The 5 min sessions of the VI30 schedule alternate with 2-5 min of a schedule (FR5) in which every 5th lever press is followed by presentation of a food pellet paired with a 0.5 sec mild footshock. The total study lasts approximately 30 mins. Rats typically respond with high rates of lever pressing under the VI30 schedule and low response rates under the FR5 'conflict' session. Anxiolytic drugs increase the suppressed response rates of rats in a 'conflict' session.

Drugs are administered intraperitoneally or orally to groups of 3-8 rats 30 to 60 mins before testing.

The results are expressed as the percentage increase in the square root of the total number of lever presses in the FR5 'conflict' session. Square root transformation is necessary to normalise the data for statistical analysis using parametric methods.

The compound of Example 4 showed a significant increase in responding in the 'conflict' session at a dose of 10mg/kg p.o.

2. MES TEST

The maximal electroshock seizure (MES) test in rodents is one of the most widely used models of human grand mal epilepsy¹. In this model,

06 MAR 1997

anticonvulsant agents elevate the threshold to electrically-induced seizures whilst proconvulsants lower the seizure threshold.

Method

Mice (male, Charles River, U.K. CD-1 strain, 25-30g) are randomly assigned to groups of 10-20 and dosed orally or intraperitoneally at a dose volume of 10ml/kg with various doses of compound (1-100 mg/kg) or vehicle. Mice are then subjected at 30 or 60 min post dose to a variable voltage electroshock (0.1 sec., 50 Hz, sine wave form) via a buccal and a subcutaneous electrode. The mean voltage and standard error required to induce a tonic seizure in 50% (CV₅₀) of the mice in the group is determined by the 'up and down' method of Dixon and Mood (1948)². Statistical comparisons between vehicle- and drug-treated groups are made using the method of Litchfield and Wilcoxon (1949)³.

In control animals the CV₅₀ is usually 40 - 50V. Hence the first animal in the control group is subjected to a voltage of 45V. If a tonic seizure does not ensue, the voltage is increased for a subsequent mouse. If a tonic convulsion does occur, then the voltage is decreased, and so on until all the animals in the group have been tested.

The percentage increase or decrease in CV₅₀ for each group compared to the control is calculated.

Studies are carried out using a Heathkit shock generator with totally variable control of shock level from 0 to 200V and voltage steps of 5V are used.

Drugs are suspended in 1% methyl cellulose.

Reference

1. Swinyard, E.A. (1972). Electrically-induced convulsions. In: Experimental Models of Epilepsy ed. Purpura, D.P. et al., 433-458, Raven Press, New York.

06 MAR 1997

2. Dixon, W.J. and Mood, A.M. (1948). J. Amer. Stat. Assn., 43, 109-126.

3. Litchfield, J.T. and Wilcoxon, F. (1949). J. Pharmacol. Exp. Ther. 96, 99-113.

Results

Compounds of Examples 1-3, 5, 7, 8, 18, 20, 21, 25, 30, 31, 35 and 37 and compound X showed a significant increase in CV₅₀ at a dose of 10 mg/kg p.o.

3. X-Maze

Introduction

The X-maze test of anxiety (Handley and Mithani, 1984) examines the exploratory response of naive rats in an environment which offers both anxiogenic (open arms) and relatively non-anxiogenic (closed arms) areas. A selective increase in exploration of the open arms following drug pretreatment is therefore postulated to indicate anxiolytic effects.

Method

The X-maze was raised 70cm above the floor and consisted of two enclosed arms 45cm (long) x 15cm (wide) x 10cm (high) and two open arms 45 x 10 x 1cm arranged such that the two arms of each type were opposite each other. Both arm types were marked into two equal sections. Rats were placed onto the centre of the X-maze and observed for a period of 10 minutes during which time the following parameters were recorded: 1) the number of entries onto, and the time spent on, (a) open arms, (b) closed arms, (c) end of open arms and (d) end of closed arms. 2) the number of sections crossed. The fear-drive evoked in the open arms exceeds that in the enclosed arms and rats typically show a clear preference for the enclosed arms. Anxiolytic drugs increase the number of entries made onto, and the time spent on, the outer half of the open arms, and also the percentage of entries made onto, and the time spent on, the whole of the open arms. These four measures of anxiety, and also the

REPLACEMENT SHEET

06 MAR 1997

total number of sections traversed, were calculated for each animal. Drugs are administered intraperitoneally or orally to groups of 6 to 12 rats 30 to 60 mins before testing. Statistical comparisons between vehicle- and drug-treated groups were made using a Mann-Whitney 'U' test (two tailed).

S.L. Handley and S. Mithani, Arch. Pharmacol., 1984 327 1-5


RESULTS

The compound of Example 21, caused a significant increase in open arm entries at a dose of 30 mg/kg p.o.

06 MAR 1997

Claims

1. A method of treatment and/or prophylaxis of anxiety and/or mania, and/or depression and/or the effects associated with withdrawal from substances of abuse, and/or disorders treatable and/or preventable with anti-convulsive agents, comprising administering to the sufferer in need thereof an effective or prophylactic amount of a compound of formula (I) or pharmaceutically acceptable salt thereof:



wherein:

either Y is N and R₂ is hydrogen, or Y is C-R₁

where:

either one of R₁ and R₂ is hydrogen and the other is selected from the class of hydrogen, C₃₋₈ cycloalkyl, C₁₋₆ alkyl optionally interrupted by oxygen or substituted by hydroxy, C₁₋₆ alkoxy or substituted aminocarbonyl, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy carbonyl, C₁₋₆ alkylcarbonyloxy, C₁₋₆ alkoxy, nitro, cyano, halo, trifluoromethyl, CF₃S, or a group CF₃-A-, where A is -CF₂-, -CO-, -CH₂- or CH(OH), trifluoromethoxy, C₁₋₆ alkylsulphinyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkoxy sulphinyl, C₁₋₆ alkoxy sulphonyl, aryl, heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphinyl, heteroarylsulphinyl, arylsulphonyl, heteroarylsulphonyl in which any aromatic moiety is optionally substituted, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxy carbonylamino, C₁₋₆ alkyl-thiocarbonyl, C₁₋₆ alkoxy-thiocarbonyl, C₁₋₆ alkyl-thiocarbonyloxy, 1-mercapto C₂₋₇ alkyl, formyl, or aminosulphinyl, aminosulphonyl or aminocarbonyl, any amino moiety being optionally substituted by one or two C₁₋₆ alkyl groups, or C₁₋₆ alkylsulphinylamino, C₁₋₆ alkylsulphonylamino, C₁₋₆ alkoxy sulphinylamino or C₁₋₆ alkoxy sulphonylamino, or ethylenyl

06 MAR 1997

terminally substituted by C₁₋₆ alkylcarbonyl, nitro or cyano, or -C(C₁₋₆ alkyl)NOH or -C(C₁₋₆ alkyl)NNH₂, or one of R₁ and R₂ is nitro, cyano or C₁₋₃ alkylcarbonyl and the other is methoxy or amino optionally substituted by one or two C₁₋₆ alkyl or by C₂₋₇ alkanoyl;

one of R₃ and R₄ is hydrogen or C₁₋₄ alkyl and the other is C₁₋₄ alkyl or R₃ and R₄ together are C₂₋₅ polymethylene;

R₅ is C₁₋₆ alkylcarbonyloxy, benzoyloxy, ONO₂, benzyloxy, phenyloxy or C₁₋₆ alkoxy and R₆ and R₉ are hydrogen or R₅ is hydroxy and R₆ is hydrogen or C₁₋₂ alkyl and R₉ is hydrogen;

R₇ is fluorophenyl;



R₈ is hydrogen or C₁₋₆ alkyl;

the R₈-N-CO-R₇ group being trans to the R₅ group;

and X is oxygen or NR₁₀ where R₁₀ is hydrogen or C₁₋₆ alkyl.

- 2.. A pharmaceutical composition for use in the treatment and/or prophylaxis of anxiety and/or mania and/or depression and/or the effects associated with withdrawal from substances of abuse and/or disorders treatable or preventable with anti-convulsive agents, which comprises an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof, as defined in claim 1, and a pharmaceutically acceptable carrier.
3. The use of a compound of formula (I), as defined in claim 1, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment and/or prophylaxis of anxiety and/or mania and/or depression and/or the effects associated with withdrawal from substances of abuse and/or disorders treatable or preventable with anti-convulsive agents.

06 MAR 1957

-36-

4. A method according to claim 1 in which the compound of formula (I) has the following variables wherein;

Y is C-R₁, where:

either one of R₁ and R₂ is hydrogen and the other is selected from the class of hydrogen, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy carbonyl, C₁₋₆ alkylcarbonyloxy, C₁₋₆ alkylhydroxymethyl, nitro, cyano, chloro, trifluoromethyl, trifluoromethoxy, C₁₋₆ alkylsulphinyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkoxy sulphinyl, C₁₋₆ alkoxy sulphonyl, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxy carbonylamino, C₁₋₆ alkyl-thiocarbonyl, C₁₋₆ alkoxy-thiocarbonyl, C₁₋₆ alkyl-thiocarbonyloxy, 1-mercapto C₂₋₇ alkyl, formyl, or aminosulphinyl, aminosulphonyl or aminocarbonyl, the amino moiety being optionally substituted by one or two C₁₋₆ alkyl groups, or C₁₋₆ alkylsulphinylamino, C₁₋₆ alkylsulphonylamino, C₁₋₆ alkoxy sulphinylamino or C₁₋₆ alkoxy sulphonylamino, or ethylenyl terminally substituted by C₁₋₆ alkylcarbonyl, nitro or cyano, or -C(C₁₋₆ alkyl)NOH or -C(C₁₋₆ alkyl)NNH₂, or one of R₁ and R₂ is nitro, cyano or C₁₋₃ alkylcarbonyl and the other is methoxy or amino optionally substituted by one or two C₁₋₆ alkyl or by C₂₋₇ alkanoyl;

one of R₃ and R₄ is hydrogen or C₁₋₄ alkyl and the other is C₁₋₄ alkyl or R₃ and R₄ together are C₂₋₅ polymethylene;

R₅ is hydroxy and R₆ is hydrogen;

R₇ is fluorophenyl;

R₈ is hydrogen or C₁₋₆ alkyl; and

R₉ is hydrogen;

the R₈-N-CO-R₇ group being trans to the R₅ group.


5. A pharmaceutical composition according to claim 2 wherein the compound of formula (I) is as defined in claim 4.



* C L H S *

06 MAR 1997

6. A use according to claim 3 wherein the compound of formula (I) is as defined in claim 4.
7. A compound of formula (Ia) or a pharmaceutically acceptable salt thereof



wherein variables Y, R₂, R₃, R₄, R₅, R₆, R₈, R₉ and X are as defined in relation to formula (I) in claim 1 and R₇ is 2- or 3-fluorophenyl or where R₇ is 2, 4- or 3,4- difluorophenyl.

8. A compound of formula (Ia) as defined in claim 7 selected from the group consisting of;

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(3-fluorobenzoylamino)2H-1-benzopyran-3-ol (Example 1),

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(2-fluorobenzoylamino)2H-1-benzopyran-3-ol (Example 2),

trans-6-Trifluoromethoxy-3,4-dihydro-2,2-dimethyl-4-(3-fluorobenzoylamino)2H-1-benzopyran-3-ol (Example 3),

trans-6-Cyano-4S-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 21),

trans-6-Cyano-4R-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 22),

trans-6-Cyano-4S-(2-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 24),

06 MAR 1997


trans-6-Cyano-4S-(2,4-difluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 28),

trans-6-Acetyl-4-(3-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 31),

trans-6-Cyano-4-(3,4-difluorobenzoyl-methylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 41) and

trans-6-cyano-4S-(3,4-difluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 44) or a pharmaceutically acceptable salt thereof.

9. A compound of formula (Ib) or pharmaceutically acceptable salts thereof:




wherein variables R₃, R₄, R₅, R₆, R₇, R₈, R₉ and X are as defined in relation to formula (I) in claim 1 and Y is CR₁ where R₁ and R₂ are both hydrogen or one of R₁ and R₂ is trifluoromethoxy, C₁₋₆ alkyl optionally interrupted with oxygen or substituted with hydroxy, C₁₋₆ alkoxy or substituted amino-carbonyl, CF₃A-(where A is -CF₂-, -CO-, -CH₂) or CH(OH), aryl sulphonyl, aryl C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphinyl, heteroarylsulphinyl, heteroarylsulphonyl, in which any aromatic moiety is optionally substituted and the other is hydrogen.

10. A compound of formula (Ib) as defined in claim 9 selected from the group consisting of:

trans-3,4-Dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)-2H-1-benzopyran-3-ol (Example 4),





trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2,6-trimethyl-2H-1-benzopyran-3-ol (Example 14),

trans-6-Ethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 15),

trans-6-Ethyl-4-(4-fluorobenzoylethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 16),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-isopropyl-2H-1-benzopyran-3-ol (Example 17),

trans-6-Ethyl-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 23),

trans-4-(4-Fluorobenzoylamino)-6-pentafluoroethyl-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 29),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-phenylsulphonyl-2H-1-benzopyran-3-ol (Example 32),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-phenyl-2H-1-benzopyran-3-ol (Example 33),

trans-6-Ethyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 35),

trans-4R-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 36),

trans-4S-(4-Fluorobenzoylamino)-3,4-dihydro-6-(1-hydroxyethyl)-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 37),

trans-4-(4-fluorobenzoyl-methylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 43),

trans-6-t-Butyl-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 46),




06 MAR 1997

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-cyclopentyl-2H-1-benzopyran-3-ol (Example 48) and

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxy-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 50), or a pharmaceutically acceptable salt thereof.

11. A compound of formula (Ic), or a pharmaceutically acceptable salt thereof:



where R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 and X are as defined in relation to formula (I) in claim 1 and Y is N and R_2 is hydrogen.

12. A compound of formula (Ic) as defined in claim 11, selected from the group consisting of:

trans-4-(2-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 38),


trans-4-(3-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 39) and

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-pyrano[3,2-c]pyridin-3-ol (Example 40), or a pharmaceutically acceptable salt thereof.

13. A compound of formula (Id), or a pharmaceutically acceptable salt thereof:



06 MAR 1997



where X is NR₁₀ where R₁₀ is hydrogen or C₁₋₆ alkyl and Y, R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₉ are as defined in relation to formula (I) in claim 1.

14. A compound of formula (Id) as defined in claim 13, selected from the group consisting of:

trans-6-Cyano-4-(4-fluorobenzoylamino)-2,2-dimethyl-1,2,3,4-tetrahydroquinolin-3-ol (Example 45), or a pharmaceutically acceptable salt thereof.

15. A compound of formula (I') selected from the group consisting of:

trans-6-Cyano-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 5),

trans-6-Chloro-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 6),

trans-6-Trifluoromethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 7),

trans-6-Cyano-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 8),

trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylamino)2H-1-benzopyran-3-ol, (Example 9),

trans-6-Ethylcarbonyl-3,4-dihydro-2,2-dimethyl-4-(4-fluorobenzoylmethylamino)2H-1-benzopyran-3-ol, (Example 10),



06 MAR 1997

trans-6-Acetyl-4-(4-fluorobenzoylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol, (Example 11),

trans-6-Cyano-3,4-dihydro-4-(4-fluorobenzoylethylamino)-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 12),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-7-nitro-2H-1-benzopyran-3-ol (Example 13),

trans-6-Acetyl-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 18),

trans-6-Acetyl-4R-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3S-ol (Example 19),

trans-6-Acetyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 20),

trans-6-Cyano-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (Example 25),

trans-6-Cyano-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2,3-trimethyl-2H-1-benzopyran-3-ol (Example 26),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-6-nitro-2H-1-benzopyran-3-ol (Example 27),


trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-methoxycarbonyl-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 30),

trans-6-Fluoro-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 34),

trans-4-(4-fluorobenzoyl-methylamino)-6-trifluoromethyl-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 42),

trans-4-(4-Fluorobenzoylamino)-3,4-dihydro-6-iodo-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 47) and





trans- 6-Aminocarbonylmethyl-4-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3-ol (Example 49) or a pharmaceutically acceptable salt thereof.

16. A compound of formula (Ia), (Ib), (Ic) or (Id), according to any one of claims 7, 9, 11 and 13, or a pharmaceutically acceptable salt thereof wherein R_3 and R_4 are both methyl.
17. A compound of formula (Ia), (Ib), (Ic) or (Id), according to any one of claims 7, 9, 11, 13 and 16, or a pharmaceutically acceptable salt thereof wherein R_5 is hydroxy and R_6 and R_9 are hydrogen.
18. A compound of formula (Ia), (Ib) or (Ic) according to any one of claims 7, 9, 11, 16 and 17, or a pharmaceutically acceptable salt thereof wherein X is oxygen.
19. A compound of formula (Ia), (Ib) or (Id) according to any one of claims 7, 9, 13 and 16 to 19, or a pharmaceutically acceptable salt thereof wherein Y is R_1 and R_1 is cyano, methoxy, trifluoromethoxy, chloro, trifluoromethyl, ethylcarbonyl, acetyl, hydrogen, methyl, ethyl, iso-propyl, tertiary-butyl, nitro, C_2F_5 , methoxycarbonyl, phenylsulphonyl, phenyl, fluoro, iodo, cyclopentyl, aminocarbonylmethyl and 1-hydroxyethyl and R_2 is hydrogen
20. A compound according to claim 19 wherein R_1 is cyano, ethyl or acetyl and R_2 is hydrogen.
21. A compound of formula (Ib), (Ic) or (Id) according to any one of claims 9, 11, 13 and 16 to 20, or a pharmaceutically acceptable salt thereof wherein R_7 is 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2,4-difluorophenyl or 3,4-difluorophenyl.
22. A compound of formula (Ia), (Ib), (Ic) or (Id), according to any one of claims 7, 9, 11, 13 and 16 to 21, or a pharmaceutically acceptable salt thereof wherein R_8 is hydrogen, methyl or ethyl.




* C L M S *

06 MAR 1997

-44-

23. A compound of formula (Ia), (Ib), (Ic) or (Id), according to ~~any one of~~ claims 7, 9, 11, 13 and 16 to 22, or a pharmaceutically acceptable salt thereof wherein the compound has the 4S,3R configuration.
24. A process for the preparation of compounds of formula (I') as listed in claim 15, or pharmaceutically acceptable salts thereof:

which comprises acylating a compound of formula (II'):





wherein Y, R₂' and R₅' are the required variables Y, R₂ or R₅ as defined in formula (I) or a group convertible thereto and R₃, R₄, R₆, R₈, R₉ and X are the required variables as defined in formula (I), the R₈NH group being trans to the R₅ group, with and acylating agent of formula (IIIb):



where R₇ is as required and as defined in formula (I) and L₁ is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any R₁', R₂' and R₅' groups to R₁, R₂ and R₅ respectively, interconverting R₈ when hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

25. A process for the preparation of compounds of formula (Ia) as defined in claim 7, or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIa):






wherein Y' , R_2' and R_5' are Y , R_2 or R_5 as defined in formula (I) or a group convertible thereto and R_3 , R_4 , R_6 , R_8 , R_9 and X are as defined in formula (I), the R_8NH group being trans to the R_5' group, with an acylating agent of formula (IIIa):




where R_7^a is 2- or 3-fluorophenyl or 2,4- or 3,4- difluorophenyl and L_1 is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any Y' , R_2' or R_5' group to Y , R_2 or R_5 respectively, interconverting R_8 when hydrogen to C_{1-6} alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

26. A process for the preparation of compounds of formula (IIb) as defined in claim 9, or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIb):




where R_1' and R_2' are both hydrogen or one of R_1 and R_2 is trifluoromethoxy, C_{1-6} alkyl optionally interrupted with oxygen or substituted with hydroxy, C_{1-6} alkoxy or substituted aminocarbonyl, CF_3A - (where A is CF_2 -, $-CO$ -, $-CH_2$ - or $CH(OH)$), aryl, aryl sulphonyl, aryl C_{3-8} cycloalkyl, C_{1-6} alkoxy, heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphinyl, heteroarylsulphinyl, heteroarylsulphonyl, in which any aromatic moiety is optionally substituted and the other is hydrogen, or groups convertible to any of these; R_5' is R_5 as defined in formula (I) or a group convertible thereto and R_3 , R_4 , R_6 , R_8 , R_9 and X are as defined in formula (I) the R_8NH group being trans to the R_5' group, with a compound of formula (IIIb):



where R_7 is as defined in formula (I) and L_1 is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any R_1' , R_2' and R_5' groups to R_1 , R_2 and R_5 respectively, interconverting R_8 when hydrogen to C_{1-6} alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

27. A process for the preparation of compounds of formula (Ic) as defined in claim 11, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIc):




in which Y is N , R_5' is R_5 , as defined in relation to formula (I) or a group convertible to R_5 and R_2' is hydrogen or a group convertible thereto, R_6 , R_3 , R_4 , R_8 , R_9 and X are as defined in formula (I), the R_8NH group being trans to the R_5' group, with an acylating agent of formula (IIIb):




where R_7 is as defined in formula (I) and L_1 is a leaving group; thereafter optionally or as necessary and in any appropriate order converting any R_2' or R_5' group to hydrogen or R_5 respectively, interconverting R_8 when hydrogen to C_{1-6} alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.





28. A process for the preparation of compounds of formula (I) as defined in claim 11, or a pharmaceutically acceptable salt thereof, which comprises acylating a compound of formula (IIId):



in which X' is NR_{10'} where R_{10'} is hydrogen or C₁₋₆ alkyl or a group convertible thereto, Y', R_{2'} and R₅ are Y, R₂ and R₅ respectively as defined in formula (I) or groups convertible thereto and R₃, R₄, R₆, R₈ and R₉ are as defined in formula (I), the R₈NH group being trans to the R_{5'} group, with an acylating agent of formula (IIIb):



where R₇ is as defined in formula (I) and L₁ is a leaving group, thereafter optionally or as necessary and in any appropriate order converting any Y', R_{10'}, R_{2'} or R_{5'} group to Y, R₁₀, R₂ or R₅ respectively, interconverting R₈ when hydrogen to C₁₋₆ alkyl, separating any enantiomers and forming a pharmaceutically acceptable salt or solvate.

29. A pharmaceutical composition comprising a compound of formula (I'), (Ia), (Ib), (Ic) or (Id), as defined in claims 15, 7, 9, 11 and 13 respectively, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
30. The use of a compound of formula (I'), (Ia), (Ib), (Ic) or (Id), as defined in claims 15, 7, 9, 11 and 13 respectively, or a pharmaceutically acceptable salt thereof, in the treatment and/or prophylaxis of anxiety and/or mania and/or depression and/or the effects associated with withdrawal from substances of abuse and/or disorders treatable and/or preventable with anti-convulsive agents.




06 MAR 1997

-48-

ABSTRACT

BENZO-AND PYRIDOPYRAN DERIVATIVES HAVING ANXIOLYTIC AND ANTI-
CONVULSANT ACTIVITY

A method of treatment and/or prophylaxis of anxiety and/or mania, and/or depression and/or the effects associated with withdrawal from substances of abuse, and/or disorders treatable and/or preventable with anti-convulsive agents, comprising administering to the sufferer in need thereof an effective or prophylactic amount of a compound of formula (I) or pharmaceutically acceptable salt thereof:




wherein:

either Y is N and R₂ is hydrogen, or Y is C-R₁

where:

either one of R₁ and R₂ is hydrogen and the other is selected from the class of hydrogen, C₃₋₈ cycloalkyl, C₁₋₆ alkyl optionally interrupted by oxygen or substituted by hydroxy, C₁₋₆ alkoxy or substituted aminocarbonyl, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylcarbonyloxy, C₁₋₆ alkoxy, nitro, cyano, halo, trifluoromethyl, CF₃S, or a group CF₃-A-, where A is -CF₂-, -CO-, -CH₂- or CH(OH), trifluoromethoxy, C₁₋₆ alkylsulphinyl, C₁₋₆ alkylsulphonyl, C₁₋₆ alkoxysulphinyl, C₁₋₆ alkoxysulphonyl, aryl, heteroaryl, arylcarbonyl, heteroarylcarbonyl, arylsulphinyl, heteroarylsulphinyl, arylsulphonyl, heteroarylsulphonyl in which any aromatic moiety is optionally substituted, C₁₋₆ alkylcarbonylamino, C₁₋₆ alkoxycarbonylamino, C₁₋₆ alkyl-thiocarbonyl, C₁₋₆ alkoxy-thiocarbonyl, C₁₋₆ alkyl-thiocarbonyloxy, 1-mercapto C₂₋₇ alkyl, formyl, or aminosulphinyl, aminosulphonyl or aminocarbonyl, any amino moiety being optionally substituted by one or two C₁₋₆ alkyl groups, or C₁₋₆ alkylsulphinylamino, C₁₋₆ alkylsulphonylamino, C₁₋₆ alkoxysulphinylamino or C₁₋₆ alkoxysulphonylamino, or ethylenyl terminally substituted by C₁₋₆ alkylcarbonyl, nitro or cyano, or -C(C₁₋₆ alkyl)NOH or -C(C₁₋₆ alkyl)NNH₂, or one of R₁ and R₂ is nitro, cyano or C₁₋₃ alkylcarbonyl and



the other is methoxy or amino optionally substituted by one or two C₁₋₆ alkyl or by C₂₋₇ alkanoyl;

one of R₃ and R₄ is hydrogen or C₁₋₄ alkyl and the other is C₁₋₄ alkyl or R₃ and R₄ together are C₂₋₅ polymethylene;

R₅ is C₁₋₆ alkylcarbonyloxy, benzoyloxy, ONO₂, benzyloxy, phenyloxy or C₁₋₆ alkoxy and R₆ and R₉ are hydrogen or R₅ is hydroxy and R₆ is hydrogen or C₁₋₂ alkyl and R₉ is hydrogen;

R₇ is fluorophenyl;

R₈ is hydrogen or C₁₋₆ alkyl;

the R₈-N-CO-R₇ group being trans to the R₅ group;

and X is oxygen or NR₁₀ where R₁₀ is hydrogen or C₁₋₆ alkyl.