


N-OXIDES AS NK1 RECEPTOR ANTAGONIST PRODRUGS OF 4-PHENYL-PYRIDINE DERIVATIVES

According to a first aspect, the present invention provides compounds of the general formula



wherein

R is hydrogen, lower alkyl, lower alkoxy, halogen or trifluoromethyl;

5 R¹ is hydrogen or halogen; or

R and R¹ may be together with the ring carbon atoms to which they are attached
-CH=CH-CH=CH-;

R² and R^{2'} are independently from each other hydrogen, halogen, trifluoromethyl, lower alkoxy or cyano; or


10 R² and R^{2'} may be together -CH=CH-CH=CH-, optionally substituted by one or two substituents selected from lower alkyl or lower alkoxy;

R³, R^{3'} are independently from each other hydrogen, lower alkyl or cycloalkyl;

R⁴, R^{4'} are independently from each other -(CH₂)_mOR⁶ or lower alkyl;

or

15 R⁴ and R^{4'} form together with the N-atom to which they are attached a cyclic tertiary amine of the group



R^5 is hydrogen, hydroxy, lower alkyl, -lower alkoxy, $-(CH_2)_mOH$, $-COOR^3$, $-CON(R^3)_2$, $-N(R^3)CO$ -lower alkyl or $-C(O)R^3$;

R^6 is hydrogen, lower alkyl or phenyl;

5

X is $-C(O)N(R^6)-$, $-N(R^6)C(O)-$, $-(CH_2)_mO-$ or $-O(CH_2)_m-$;

n is 0, 1, 2, 3 or 4; and

10 m is 1, 2 or 3;

and to pharmaceutically acceptable acid addition salts thereof.

According to a second aspect, the present invention provides a medicament containing one or more compounds of formula I and pharmaceutically acceptable excipients.


15

According to a third aspect, the present invention provides a medicament according to the second aspect for the treatment of diseases related to NK-1 receptor antagonists.


According to a fourth aspect, the present invention provides a process for preparing a compound of formula I as defined in the first aspect of the invention, which process comprises

20

oxidizing a compound of formula



with a suitable oxidizing agent to give a compound of formula




wherein the substituents have the significances given in accordance to the first aspect of the invention and if desired, converting the compound obtained into a pharmaceutically acceptable acid addition salt.

- 5 According to a fifth aspect, the present invention provides a compound of formula I, whenever prepared by a process according to the fourth aspect of the invention or by an equivalent method.

- 10 According to a sixth aspect, the present invention provides the use of a compound of formula I for the manufacture of medicaments containing one or more compounds of formula I for the treatment of diseases related to NK-1 receptor antagonists.


According to a seventh aspect, the present invention provides a prodrug for a parent compound of formula II



- 15 for use in medicine,
 wherein the prodrug is a compound of formula I according to the first aspect of the present invention.

-2b-

It has been found that these N-oxides of the present invention have an in vitro activity on the NK1 receptor and/or may be used as prodrugs of compounds of formula



which are antagonists of the Neurokinin 1 (NK-1, substance P) receptor.

However, the advantage of a prodrug lies in its physical properties, such as enhanced water solubility for parenteral administration compared to the parent drug, or it enhances absorption from the digestive tract, or it may enhance drug stability for long-term storage. Compounds of formula II have limited water solubility, not allowing bolus injections. It was therefore useful to find derivatives of the compound of formula II to render these compounds suitable for parenteral and intramuscular application. It has been shown that N-oxides of compounds of formula I fulfil all requirements of a good prodrug.

A prodrug is in most cases a pharmacologically inactive derivative of a parent drug molecule that requires spontaneous or enzymatic transformation within the body in order to release the active drug, and that has improved delivery properties over the parent drug molecule. It has been shown that a molecule with optimal structural configuration and physicochemical properties for eliciting the desired therapeutic response at its target site does not necessarily possess the best molecular form and properties for its delivery to its point of ultimate action. Usually, only a minor fraction of doses administered reach the target area and since most agents interact with non-target sites as well, an inefficient delivery may result in undesirable side effects. This fact of differences in transport and in

- 3 -

situ effect characteristics for many drug molecules is the basic reason why bioreversible chemical derivatization of drugs, i.e., prodrug formation is a means by which a substantial improvement in the overall efficacy of drugs can often be achieved. Prodrugs are designed to overcome pharmaceutically and/or pharmacokinetically based problems associated with
5 the parent drug molecule that would otherwise limit the clinical usefulness of the drug.

In recent years several types of bioreversible derivatives have been exploited for utilization in designing prodrugs. Using esters as a prodrug type for drugs containing carboxyl or hydroxyl function is most popular. Further well-known are prodrug derivatives of peptides, 4-imidazolidinones and the like, described in *Drugs of the Future*, 1991, 16(5),
10 443-458 or N-oxides, described for example in US 5.691.336.

As mentioned earlier, the compounds of formula II are antagonists of the neurokinin receptor. The central and peripheral actions of the mammalian tachykinin substance P have been associated with numerous inflammatory conditions including migraine, rheumatoid arthritis, asthma, and inflammatory bowel disease as well as mediation of the
15 emetic reflex and the modulation of central nervous system (CNS) disorders such as Parkinson's disease (Neurosci. Res., 1996, 7, 187-214), anxiety (Can. J. Phys., 1997, 75, 612-621) and depression (Science, 1998, 281, 1640-1645).

Evidence for the usefulness of tachykinin receptor antagonists in pain, headache, especially migraine, Alzheimer's disease, multiple sclerosis, attenuation of morphine
20 withdrawal, cardiovascular changes; oedema, such as oedema caused by thermal injury, chronic inflammatory diseases such as rheumatoid arthritis, asthma/bronchial hyperreactivity and other respiratory diseases including allergic rhinitis, inflammatory diseases of the gut including ulcerative colitis and Crohn's disease, ocular injury and ocular inflammatory diseases reviewed in "Tachykinin Receptor and Tachykinin Receptor
25 Antagonists", J. Auton. Pharmacol., 13, 23-93, 1993.

Furthermore, Neurokinin 1 receptor antagonists are being developed for the treatment of a number of physiological disorders associated with an excess or imbalance of tachykinin, in particular substance P. Examples of conditions in which substance P has been implicated include disorders of the central nervous system such as anxiety, depression
30 and psychosis (WO 95/16679, WO 95/18124 and WO 95/23798).

The neurokinin-1 receptor antagonists are further useful for the treatment of motion sickness and for treatment induced vomiting.

In addition, in The New England Journal of Medicine, Vol. 340, No. 3 190-195, 1999 has been described the reduction of cisplatin-induced emesis by a selective neurokinin-1-
35 receptor antagonist.

The usefulness of neurokinin 1 receptor antagonists for the treatment of certain forms of urinary incontinence is further described in *Neuropeptides*, 32(1), 1-49, (1998) and *Eur. J. Pharmacol.*, 383(3), 297-303, (1999).

Furthermore, US 5,972,938 describes a method for treating a psychoimmunologic or a psychosomatic disorder by administration of a tachykinin receptor, such as NK-1 receptor antagonist.

Objects of the present invention are the compounds of formula I and pharmaceutically acceptable salts thereof, the preparation of the above-mentioned compounds, medicaments containing them and their manufacture as well as the use of the above-mentioned compounds in the control or prevention of illnesses, especially of illnesses and disorders of the kind referred to earlier or in the manufacture of corresponding medicaments.

The most preferred indications in accordance with the present invention are those, which include disorders of the central nervous system, for example the treatment or prevention of certain depressive disorders or emesis by the administration of NK-1 receptor antagonists. A major depressive episode has been defined as being a period of at least two weeks during which, for most of the day and nearly every day, there is either depressed mood or the loss of interest or pleasure in all, or nearly all activities.

The following definitions of the general terms used in the present description apply irrespective of whether the terms in question appear alone or in combination. As used herein, the term "lower alkyl" denotes a straight- or branched-chain alkyl group containing from 1-7 carbon atoms, for example, methyl, ethyl, propyl, isopropyl, n-butyl, i-butyl, t-butyl and the like.

Preferred lower alkyl groups are groups with 1-4 carbon atoms.

The term "lower alkoxy" denotes a group wherein the alkyl residues are as defined above, and which is attached via an oxygen atom.

The term "halogen" denotes chlorine, iodine, fluorine and bromine.

The term "cycloalkyl" denotes a saturated carbocyclic group, containing 3-6 carbon atoms.

The term "cyclic tertiary amine" denotes, for example, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, morpholin-4-yl or 1,1-dioxo-thiomorpholin-4-yl.

Exemplary preferred are compounds, in which X is $-\text{C}(\text{O})\text{N}(\text{R}^6)-$, wherein R^6 is methyl, for example the following compounds:

- 4-{5-[(3,5-bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4-o-tolyl-pyridin-2-yl}-4-oxy-piperazine-1-carboxylic acid tert-butyl ester,
 5'-[(3,5-bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4'-o-tolyl-1-oxy-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-carboxylic acid ethyl ester,
 5 (RS)-6-[3-(acetyl-methyl-amino)-1-oxo-pyrrolidin-1-yl]-N-(3,5-bis-trifluoromethyl-benzyl)-N-methyl-4-o-tolyl-nicotinamide,
 N-(3,5-bis-trifluoromethyl-benzyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide,
 N-(3,5-bis-trifluoromethyl-benzyl)-6-(1,1-dioxo-1 λ ⁶-4-oxy-thiomorpholin-4-yl)-N-
 10 methyl-4-o-tolyl-nicotinamide,
 N-(3,5-bis-trifluoromethyl-benzyl)-6-(4-formyl-1-oxy-piperazin-1-yl)-N-methyl-4-o-tolyl-nicotinamide,
 N-methyl-N-(2-methyl-naphthalen-1-ylmethyl)-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide,
 15 N-methyl-6-(4-oxy-morpholin-4-yl)-N-naphthalen-1-ylmethyl-4-o-tolyl-nicotinamide,
 N-(2-methoxy-naphthalen-1-ylmethyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide,
 N-(2-methoxy-benzyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide,
 N-(5-chloro-2-methoxy-benzyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-
 20 nicotinamide,
 N-(2-chloro-5-methoxy-benzyl)-N-methyl-6-morpholin-4-yl-4-o-tolyl-nicotinamide,
 N-methyl-6-(4-oxy-morpholin-4-yl)-N-pentafluorophenylmethyl-4-o-tolyl-nicotinamide,
 N-methyl-6-(4-oxy-morpholin-4-yl)-N-naphthalen-2-ylmethyl-4-o-tolyl-nicotinamide,
 N-[2-methoxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-N-methyl-6-(4-oxy-
 25 morpholin-4-yl)-4-o-tolyl-nicotinamide,
 N-(1,4-dimethoxy-naphthalen-2-ylmethyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide or
 5'-[(3,5-bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4'-o-tolyl-1-oxy-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-carboxylic acid.

- 30 Further preferred are compounds, in which X is -N(R⁶)-CO-, wherein R⁶ is hydrogen or methyl.

Examples of such compounds are:

2-(3,5-bis-trifluoromethyl-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-isobutyramide,


- 6 -

- 2-(3,5-bis-trifluoromethyl-phenyl)-N-[4-(2-chloro-phenyl)-6-(4-oxy-morpholin-4-yl)-pyridin-3-yl]-N-methyl-isobutyramide,
- 2-(3,5-bis-trifluoromethyl-phenyl)-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-isobutyramide,
- 5 2-(3,5-bis-trifluoromethyl-phenyl)-N-[4'-(2-chloro-phenyl)-1-oxy-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl]-N-methyl-isobutyramide,
- 2-(3,5-bis-trifluoromethyl-phenyl)-N-(6-oxy-dimethylamino-4-o-tolyl-pyridin-3-yl)-N-methyl-isobutyramide,
- 2-(3,5-bis-trifluoromethyl-phenyl)-N-[4-(2-chloro-phenyl)-6-oxy-dimethylamino-pyridin-3-yl]-isobutyramide,
- 10 2-(3,5-bis-trifluoromethyl-phenyl)-N-1-(4-hydroxy-1-oxy-4'-o-tolyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-N-methyl-isobutyramide,
- 2-(3,5-bis-trifluoromethyl-phenyl)-N-[6-[(2-hydroxy-ethyl)-1-oxy-methyl-amino]-4-o-tolyl-pyridin-3-yl]-N-methyl-isobutyramide,
- 15 (R)-2-(3,5-bis-trifluoromethyl-phenyl)-N-[6-(3-hydroxy-1-oxy-pyrrolidin-1-yl)-4-o-tolyl-pyridin-3-yl]-N-methyl-isobutyramide,
- 2-(3,5-bis-trifluoromethyl-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-acetamide,
- 2-(3,5-dimethoxy-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-acetamide or
- 20 2-(3-fluoro-5-trifluoromethyl-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-acetamide.

The present compounds of formula I and their pharmaceutically acceptable salts can be prepared by methods known in the art, for example, by processes described below,


25 which process comprises

- a) oxidizing a compound of formula



with a suitable oxidizing agent to give a compound of formula

- 7 -



wherein the substituents have the significances given above, and if desired, converting the compound obtained into a pharmaceutically acceptable acid addition salt.

- 5 In accordance with this procedure, a compound of formula I may be prepared, for example, as follows: To a solution of 10 mmol of a compound of general formula II in 50 ml of a suitable solvent such as dichloromethane is added under ice cooling a solution of 10 mmol of a suitable oxidizing reagent such as 3-chloroperbenzoic acid in 50 ml of a suitable solvent such as dichloromethane. Stirring is continued for an appropriate time
- 10 (typically 1 h to 24 h) at 0 °C and reaction progress may be followed by thin-layer-chromatography. In cases in which product formation is too slow, the reaction mixture may be stirred at room temperature. After evaporation of the solvent products of general formula I can be isolated by flash-chromatography in 15 % to 85 % yield. Further purification of crystalline products may be achieved by recrystallization from a suitable
- 15 solvent.

For this transformation other oxidizing reagents may be used instead of 3-chloroperbenzoic acid. Those oxidizing reagents are familiar to any person skilled in the art such as dimethyldioxirane in acetone, hydrogenperoxide in acetic acid or potassium peroxymonosulfate in a suitable solvent such as water.


- 20 The salt formation is effected at room temperature in accordance with methods which are known per se and which are familiar to any person skilled in the art. Not only salts with inorganic acids, but also salts with organic acids come into consideration. Hydrochlorides, hydrobromides, sulphates, nitrates, citrates, acetates, maleates, succinates, methan-sulphonates, p-toluenesulphonates and the like are examples of such salts.

- 25 The following schemes 1-8 describe the processes for preparation of compounds of formula I in more detail. The starting materials are known compounds and may be prepared according to methods known in the art, for example in accordance with methods, described in EP 99103504.9.

In the schemes the following abbreviations have been used:


PivCl	pivaloyl chloride
THF	tetrahydrofuran
TMEDA	N,N,N',N'-tetramethylethylene diamine
5 DIPEA	N-ethyl-diisopropyl-amine
KHMDS	potassium hexamethyldisilazide

Scheme 1



The definition of substituents is given above.


Scheme 2



The definition of substituents is given above.

- 10 -

Scheme 3




$\text{Z} = \text{Cl, Br, I or OS(O)}_2\text{C}_6\text{H}_4\text{CH}_3$

and the definition of the other substituents is given above.


- 11 -

Scheme 4



The definition of substituents is given above.


Scheme 5



- 12 -

Z is Cl, Br, I or $-\text{OS}(\text{O})_2\text{C}_6\text{H}_4\text{CH}_3$ and the definition of the other substituents is described above.


Scheme 6



- 5 Z is Cl, Br, I or $-\text{OS}(\text{O})_2\text{C}_6\text{H}_4\text{CH}_3$ and the definition of the other substituents is given above.

- 13 -

Scheme 7



R^5 is the group $-C(O)R^3$ and the definition of the remaining substituents is given above.


5

10

15

- 14 -

Scheme 8



The definition of substituents is given above.

As mentioned earlier, the compounds of formula I and their pharmaceutically usable addition salts may be used as prodrugs of the parent compounds of formula II, which possess valuable pharmacological properties. These compounds are antagonists of the Neurokinin 1 (NK-1, substance P) receptor.

Furthermore, in addition, some N-oxides of formula I have a good affinity to the NK1 receptor. For some preferred compounds the pK_i value is in the range of 8.3 to 8.7.

These compounds were investigated in accordance with the tests given hereinafter.

Binding assay (in vitro)

- 15 -

The affinity of test compounds for the NK₁ receptor was evaluated at human NK₁ receptors in CHO cells infected with the human NK₁ receptor (using the Semliki virus expression system) and radiolabelled with [³H]substance P (final concentration 0.6 nM). Binding assays were performed in HEPES buffer (50 mM, pH 7.4) containing BSA (0.04 %) 5 leupeptin (8 µg / ml), MnCl₂ (3 mM) and phosphoramidon (2 µM). Binding assays consisted of 250 µl of membrane suspension (1.25 x 10⁵ cells / assay tube), 0.125 µl of buffer of displacing agent and 125 µl of [³H]substance P. Displacement curves were determined with at least seven concentrations of the compound. The assay tubes were 10 filtered under vacuum through GF/C filters presoaked for 60 min with PEI (0.3 %) with 2 x 2 ml washes of HEPES buffer (50 mM, pH 7.4). The radioactivity retained on the filters was measured by scintillation counting. All assays were performed in triplicate in at least 2 separate experiments.

The evidence, that the compounds of formula I may be used as prodrugs of their 15 parent compounds of formula II is shown in accordance with the description given hereinafter.

The conversion of N-oxide prodrugs to the corresponding parent compounds is due to a reduction mechanism and there is some evidence from the literature that similar reactions occur in vivo and are probably catalysed by hemoglobin, hence the decision to 20 study both the stability in plasma and blood was taken. The presence of an oxidant in the work-up solution should help preventing the reduction of the N-oxides.

Conversion in plasma: 10 µLs of a 100 µg/mL DMSO solution of the pro-drug were added to 1 mL plasma to reach a final concentration of 1 µg/mL. The incubation was performed at 37 °C and 8 aliquots were taken at different time points over 30 min. These aliquots were 25 treated with 3 volumes of cold MeOH containing H₂O₂ (final concentration 10 %v/v) and centrifuged at 3500 g for 20 min at 10 °C. The supernatant was directly used to determine the drug levels by LC-MS-MS (HPLC chromatography on reversed phase column X-Terra MS C18 3.5 µM 2.1 x 30 mm Waters at 40 °C, using a polarity gradient MeOH/Form.Ac. 1 % 20/80 /MeOH; run time; 3.0 min; inj. Vol.: 10 µL; Flow: 0.2 µL/min and MS/MS 30 detection on a PE Sciex API-2000 MS/MS spectrometer; ion source: Turbospray; ionisation mode: ESP+).

Conversion in fresh blood: The same procedure was used for the stability studies in blood, even if much more care had to be taken after treatment with the H₂O₂.

Sample stability (plasma and blood): The final matrix were first prepared (plasma or blood 35 treated with 3 volumes of cold MeOH containing H₂O₂ - 10%v/v - and centrifuged at 3500

g for 20 min at 10 °C) and then incubated at 37 °C into two tubes; the pro-drug or the drug were then incubated and finally their concentration determined by LC-MS-MS as described above.

5 The method used to stop the reaction in both plasma and blood was found to be enough reliable to perform the studies at least when the analysis was performed immediately after the incubations.

The half-life obtained for the conversion prodrug to drug in plasma are reported in the following table (Plasma sample preparation was found to be critical for exact determination of values for $t_{1/2}$):

$t_{1/2}$ (hours)			
Example No.	Dog plasma	Human plasma	Rat plasma
1	18	8	4
2	18	12	5
16	16	6	2

10 The stability in blood is much lower ($t_{1/2} < 30$ min) and it was impossible to determine a precise value of $t_{1/2}$. However we can conclude that there are no major species difference with respect with stability in blood and that the prodrugs are converted to the desired drugs in high yield. (> 90 %)

15 In accordance with the tests the compounds of formula I can function as prodrugs of their parent compounds of formula II.

The compounds of formula I as well as their pharmaceutically usable acid addition salts can be used as medicaments, e.g. in the form of pharmaceutical preparations. The pharmaceutical preparations can be administered orally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions. The administration can, however, also be effected rectally, e.g. in the form of suppositories, or parenterally, e.g. in the form of injection solutions.

25 The compounds of formula I and their pharmaceutically usable acid addition salts can be processed with pharmaceutically inert, inorganic or organic excipients for the production of tablets, coated tablets, dragees and hard gelatine capsules. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts etc can be used as such excipients e.g. for tablets, dragées and hard gelatine capsules.

Suitable excipients for soft gelatine capsules are e.g. vegetable oils, waxes, fats, semi-solid and liquid polyols etc.

Suitable excipients for the manufacture of solutions and syrups are e.g. water, polyols, saccharose, invert sugar, glucose etc.

- 5 Suitable excipients for injection solutions are e.g. water, alcohols, polyols, glycerol, vegetable oils etc.

Suitable excipients for suppositories are e.g. natural or hardened oils, waxes, fats, semi-liquid or liquid polyols etc.

- 10 Moreover, the pharmaceutical preparations can contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

- 15 The dosage can vary within wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, in the case of oral administration a daily dosage of about 10 to 1000 mg per person of a compound of general formula I should be appropriate, although the above upper limit can also be exceeded when necessary.

The following Examples illustrate the present invention without limiting it. All temperatures are given in degrees Celsius.

- 20 The preparation of compounds of formula I, starting with compounds of formula II, is described generically in the description. This oxidation procedure is always the last step to obtain the N-oxides of compounds of formula I. A detailed description of this last step is specifically described in the following examples 1, 2 and 13. The N-oxidation of the remaining compounds 3 to 12 and 14 to 29 is generically described in accordance with the above mentioned description.

25

Example 1

2-(3,5-Bis-trifluoromethyl-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-isobutyramide

a) 4-(5-Nitro-2-pyridyl)-morpholine

- 30 To a solution of 20 g (126 mmol) of 2-chloro-5-nitropyridine in 150 ml tetrahydrofuran were added dropwise 27 ml (315 mmol) morpholine within 10 min. The reaction mixture

was refluxed for additional 2 h. After cooling to room temperature, the solvent was removed *in vacuo* and the residue was re-dissolved in 200 ml ethyl acetate. The organic phase was washed with 200 ml 1 N sodium bicarbonate solution, dried (magnesium sulfate) and evaporated to give 27.3 g (quantitative) of the title compound as a yellow solid.
5 M.p. 142-143 °C.

b) 2,2-Dimethyl-N-(6-morpholin-4-yl-pyridin-3-yl)-propionamide

To a solution of 27.3 g (126 mmol) of 4-(5-nitro-2-pyridyl)-morpholine in 600 ml methanol were added 2.5 g of 10 % of palladium on activated charcoal. The reaction mixture was hydrogenated (room temperature to *ca.* 45 °C, 1 bar) until the theoretical
10 amount of hydrogen was taken up (about 3 h). The catalyst was filtered off and was washed twice with 100 ml portions of methanol. The filtrate was evaporated *in vacuo* to give 22.6 g of a purple oil which consisted to *ca.* 95 % of the desired aniline derivative according to analysis by thin layer chromatography.

This crude product was dissolved in a mixture of 240 ml tetrahydrofuran and 60 ml diethyl
15 ether. After cooling to 0 °C, 26 ml (189 mmol) of triethylamine were added in one portion. Stirring was continued while 23 g (189 mmol) of pivaloyl chloride were added dropwise within a period of 10 min. The ice bath was removed and the reaction mixture was stirred for 1h at room temperature. Then, the solvent was removed *in vacuo* and the residue was suspended in 200 ml 1 N sodium bicarbonate solution. The product was extracted three
20 times with 200 ml portions of dichloromethane, dried (sodium sulfate) and evaporated. Recrystallization of the solid residue from ethyl acetate/hexane 1:8 gave 28.6 g (86 %) of the title compound as white crystals.

MS m/e (%): 264 (M+H⁺, 100).

c) N-(4-Iodo-6-morpholin-4-yl-pyridin-3-yl)-2,2-dimethyl-propionamide

25 A solution of 28.4 g (108 mmol) 2,2-dimethyl-N-(6-morpholin-4-yl-pyridin-3-yl)-propionamide and 49 ml (324 mmol) N,N,N',N'-tetramethylethylenediamine under argon in 600 ml tetrahydrofuran was cooled in a dry ice bath to -78 °C. Within 1h, 202 ml (324 mmol) of a 1.6 N n-butyllithium solution in hexane were added dropwise. The reaction mixture was allowed to warm up to -35 °C overnight. After cooling again to -78 °C, 37 g
30 (146 mmol) iodine dissolved in 60 ml tetrahydrofuran were added dropwise during 15 min. The dry ice bath was replaced by an ice bath and a solution of 90 g (363 mmol) sodium thiosulfate pentahydrate in 250 ml water were added within 10 min when the temperature of the reaction mixture had reached 0 °C. Then, 1000 ml diethyl ether were

added and the organic layer was separated. The aqueous layer was extracted twice with 500 ml dichloromethane and the combined organic layers were dried (magnesium sulfate) and evaporated. Flash chromatography gave 15.6 g (37 %) of the title compound as a light brown oil which crystallized upon standing at room temperature.

- 5 MS m/e (%): 389 (M^+ , 71), 358 (25), 304 (43), 57 (100).

d) 2,2-Dimethyl-N-(6-morpholin-4-yl-4-o-tolyl-pyridin-3-yl)-propionamide

- A mixture of 3.50 g (9.0 mmol) N-(4-iodo-6-morpholin-4-yl-pyridin-3-yl)-2,2-dimethyl-propionamide, 35 ml toluene, 18 ml 2 N sodium carbonate solution, 312 mg (0.27 mmol) tetrakis(triphenylphosphine)palladium(0) and 1.34 g (9.9 mmol) o-tolylboronic acid was
10 heated under argon at 80 °C for 12 h. After cooling to room temperature, the aqueous phase was separated and washed twice with ethyl acetate. The combined organic layers were washed with 50 ml brine, dried (sodium sulfate) and evaporated. Purification by flash-chromatography gave 3.23 g (quantitative) of the title compound as a white foam.

MS m/e (%): 354 ($M+H^+$, 100).

- 15 e) 6-Morpholin-4-yl-4-o-tolyl-pyridin-3-ylamine

- A suspension of 2.93 g (8.28 mmol) 2,2-dimethyl-N-(6-morpholin-4-yl-4-o-tolyl-pyridin-3-yl)-propionamide in 80 ml 3 N hydrochloric acid solution and 5 ml 1-propanol was heated to 90-95 °C overnight. The reaction mixture was cooled to room temperature, washed with three 20 ml portions diethyl ether and filtered over celite. The filtrate was
20 diluted with 20 ml water and was adjusted to pH 7-8 by addition of 28 % sodium hydroxide solution under ice cooling. The product was extracted with four 100 ml portions of dichloromethane. The combined organic layers were washed with 50 ml brine, dried (magnesium sulfate) and evaporated to give 2.31 g (quantitative) of the title compound as a white foam.

- 25 MS m/e (%): 269 (M^+ , 100).

f) Methyl-(6-morpholin-4-yl-4-o-tolyl-pyridin-3-yl)-amine

- A solution of 2.24 g (8.3 mmol) 6-morpholin-4-yl-4-o-tolyl-pyridin-3-ylamine in 17 ml trimethyl orthoformate and 3 drops trifluoroacetic acid was heated for 2 h at 130 °C. The reaction mixture was evaporated and dried *in vacuo* for 30 min. The residual oil was
30 dissolved in 5 ml tetrahydrofuran and was added dropwise under ice cooling to 630 mg (16.6 mmol) lithium aluminum hydride in 20 ml tetrahydrofuran. The reaction mixture was stirred for 1h at room temperature, cooled to 0 °C again and acidified (pH 1-2) by

- 20 -

addition of 28 % hydrochloric acid solution. After stirring for 5 min, 28 % sodium hydroxide solution was added to reach pH 10. The solution was filtered over celite, evaporated and purified by flash chromatography to give 1.56 g (66 %) of the title compound as a white foam.

5 MS m/e (%): 283 (M^+ , 100).

g) 2-(3,5-Bis-trifluoromethyl-phenyl)-N-methyl-N-(6-morpholin-4-yl-4-o-tolyl-pyridin-3-yl)-isobutyramide

A solution of 1.46 g (5.15 mmol) methyl-(6-morpholin-4-yl-4-o-tolyl-pyridin-3-yl)-amine and 1.32 ml (7.73 mmol) N-ethyldiisopropylamine in 15 ml dichloromethane was cooled
10 in an ice bath and 1.8 g (5.67 mmol) 2-(3,5-bis-trifluoromethyl-phenyl)-2-methyl-propionyl chloride were added dropwise. The reaction mixture was warmed to 35-40 °C for 3h, cooled to room temperature again and was stirred with 25 ml saturated sodium bicarbonate solution. The organic layer was separated and the aqueous phase was extracted with dichloromethane. The combined organic layers were dried (magnesium sulfate) and
15 evaporated. The residue was purified by flash chromatography to give 2.9 g (quantitative) of the title compound as white crystals. M.p. 131-132 °C.

h) 2-(3,5-Bis-trifluoromethyl-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-isobutyramide

To a solution of 5.0 g (8.84 mmol) 2-(3,5-bis-trifluoromethyl-phenyl)-N-methyl-N-(6-morpholin-4-yl-4-o-tolyl-pyridin-3-yl)-isobutyramide in 50 ml dichloromethane was
20 added under ice cooling a solution of 2.18 g (8.84 mmol) of 3-chloroperbenzoic acid (ca. 70 %) in 35 ml dichloromethane. After stirring for 1 h at 0 °C, 2.6 g (25.7 mmol) triethylamine were added slowly. The reaction mixture was concentrated to a total volume of 10 mL and the residue was purified by flash-chromatography. The crude material was
25 suspended in 20 ml diethyl ether, filtered and dried *in vacuo* to give 4.2 g (82 %) of the title compound as white crystals. M.p. 149-151 °C (partial decomposition).

MS m/e (%): 582 ($M+H^+$, 100).

Example 2

2-(3,5-Bis-trifluoromethyl-phenyl)-N-[4-(2-chloro-phenyl)-6-(4-oxy-morpholin-4-yl)-pyridin-3-yl]-N-methyl-isobutyramide
30

The title compound was obtained as white crystals in comparable yields according to the procedures described above for Example 1 using 2-chlorophenylboronic acid instead of o-

- 21 -

tolylboronic acid in step d). M.p. 141-143 °C (partial decomposition),
MS m/e (%): 602 (M+H⁺, 100), 624 (M+Na⁺, 10).

Example 3

2-(3,5-Bis-trifluoromethyl-phenyl)-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-
5 isobutyramide

The parent compound was obtained as white powder in comparable yields according to the procedures described above for the preparation of Example 1 in steps a) to g). Step f) was omitted.

MS m/e (%): 552 (M+H⁺, 100).

10 The N-oxide was obtained in accordance with step h) in Example 1.

Example 4

2-(3,5-Bis-trifluoromethyl-phenyl)-N-[4'-(2-chloro-phenyl)-1-oxy-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl]-N-methyl-isobutyramide

The parent compound was obtained as a white powder in comparable yields according to
15 the procedures described above for the preparation of Example 1 in steps a) to g) using piperidine instead of morpholine in step a) and using 2-chlorophenylboronic acid instead of o-tolylboronic acid in step d).

MS m/e (%): 583 (M⁺, 20), 296 (78), 255 (100).

The N-oxide was obtained in accordance with step h) in Example 1.

20

Example 5

2-(3,5-Bis-trifluoromethyl-phenyl)-N-(6-oxy-dimethylamino-4-o-tolyl-pyridin-3-yl)-N-methyl-isobutyramide

The parent compound was obtained as white solid in comparable yields according to the procedures described above for the preparation of Example 1, steps a) to g) using
25 dimethylamine hydrochloride instead of morpholine in step a). M.p. 174-175 °C,
MS m/e (%): 524 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

Example 6

2-(3,5-Bis-trifluoromethyl-phenyl)-N-[4-(2-chloro-phenyl)-6-oxy-dimethylamino-
30 pyridin-3-yl]-isobutyramide

- 22 -

The parent compound was obtained as white solid in comparable yields according to the procedures described above for the preparation of Example 1 steps a) to g) using dimethylamine hydrochloride instead of morpholine in step a) and using 2-chlorophenylboronic acid instead of o-tolylboronic acid in step d). M.p. 162-163 °C.

5 MS m/e (%): 544 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

Example 7

2-(3,5-Bis-trifluoromethyl-phenyl)-N-1-(4-hydroxy-1-oxy-4'-o-tolyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl)-N-methyl-isobutyramide

10 The parent compound was obtained as white foam in comparable yields according to the procedures described above for the preparation of Example 1 steps a) to g) using 4-hydroxypiperidine instead of morpholine in step a).

MS m/e (%): 580 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

15

Example 8

2-(3,5-Bis-trifluoromethyl-phenyl)-N-[6-[(2-hydroxy-ethyl)-1-oxy-methyl-amino]-4-o-tolyl-pyridin-3-yl]-N-methyl-isobutyramide

The parent compound was obtained as white foam in comparable yields according to the procedures described above for the preparation of Example 1 in steps a) to g) using N-methylethanolamine instead of morpholine in step a).

20 MS m/e (%): 554 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

Example 9

25 (R)-2-(3,5-Bis-trifluoromethyl-phenyl)-N-[6-(3-hydroxy-1-oxy-pyrrolidin-1-yl)-4-o-tolyl-pyridin-3-yl]-N-methyl-isobutyramide

The parent compound was obtained as white foam in comparable yields according to the procedures described above for the preparation of Example 1 in steps a) to g) using (R)-3-hydroxypyrrolidine instead of morpholine in step a).

MS m/e (%): 566 (M+H⁺, 100).

30 The N-oxide was obtained in accordance with step h) in Example 1.

Example 10

2-(3,5-Bis-trifluoromethyl-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-acetamide

To a solution of 300 mg (1.1 mmol) 3,5-bis(trifluoromethyl)-phenylacetic acid in 7 ml
5 N,N-dimethylformamide were added 185 mg (1.14 mmol) 1,1'-carbonyl-diimidazole and
the solution was stirred for 30 min at room temperature. After addition of 283 mg (1
mmol) of methyl-(6-morpholin-4-yl-4-tolyl-pyridin-3-yl)-amine (as described in step f)
for the preparation of Example 1), the reaction mixture was heated over night at 90°C.
After cooling to room temperature, the solvent was removed *in vacuo* and the residue was
10 re-dissolved in 30 ml ethyl acetate. The organic phase was washed with water (2 x 30 ml),
brine, dried (magnesium sulfate) and evaporated. Flash chromatography gave 506 mg (94
%) of the parent compound as a light brown foam.

MS m/e (%): 538 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

15

Example 11

2-(3,5-Dimethoxy-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-acetamide

To a solution of 226 mg (1.15 mmol) 3,5-dimethoxy-phenylacetic acid in 7 ml N,N-
dimethylformamide were added 244 mg (1.5 mmol) 1,1'-carbonyl-diimidazole and the
20 solution was stirred for 30 min at room temperature. After addition of 283 mg (1 mmol) of
methyl-(6-morpholin-4-yl-4-tolyl-pyridin-3-yl)-amine (as described in step f) for the
preparation of Example 1), the reaction mixture was heated at 70 °C for 7h. After cooling
to room temperature, the solvent was removed *in vacuo* and the residue was re-dissolved in
30 ml ethyl acetate. The organic phase was washed with water (2 x 30 ml), brine, dried
25 (magnesium sulfate) and evaporated. Flash chromatography gave 347 mg (75 %) of the
parent compound as a white foam.

MS m/e (%): 462 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

- 24 -

Example 12

2-(3-Fluoro-5-trifluoromethyl-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-acetamide

To a solution of 266 mg (1.2 mmol) 3-fluoro-5-trifluoromethyl-phenylacetic acid in 7 ml
5 N,N-dimethylformamide were added 195 mg (1.2 mmol) 1,1'-carbonyl-diimidazole and
the solution was stirred for 30 min at room temperature. After addition of 283 mg (1
mmol) of methyl-(6-morpholin-4-yl-4-tolyl-pyridin-3-yl)-amine (as described in step f)
for the preparation of Example 1), the reaction mixture was heated at 90 °C for 6h. After
cooling to room temperature, the solvent was removed *in vacuo* and the residue was re-
10 dissolved in 30 ml ethyl acetate. The organic phase was washed with water (2 x 30 ml),
brine, dried (magnesium sulfate) and evaporated. Flash chromatography gave 432 mg (88
%) of the parent compound as a light yellow foam.

MS m/e (%): 488 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

15

Example 13

4-{5-[(3,5-Bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4-o-tolyl-pyridin-2-yl}-4-oxy-
piperazine-1-carboxylic acid tert-butyl ester

a) 6-Chloro-N-methyl-4-o-tolyl-nicotinamide

To a solution of 3.41 g (20.0 mmol) 6-chloro-N-methyl-nicotinamide in 80 ml
20 tetrahydrofuran 50 ml (50 mmol) of a 1 M solution of o-tolyl magnesium chloride in
tetrahydrofuran was added dropwise at 0 °C. After completed addition the reaction
mixture was allowed to warm to room temperature and stirred for 1.5h. The mixture was
again cooled to 0 °C, followed by the dropwise addition of 5.7 ml (100 mmol) acetic acid
and a solution of 5.1 g (22 mmol) 2,3-dichloro-5,6-dicyano-1,4-benzoquinone in 18 ml
25 tetrahydrofuran. After completed addition the reaction mixture was allowed to warm to
room temperature and stirred for 15 min. Addition of 30 ml 2 N aqueous sodium
hydroxide solution was followed by dilution with 1 l ethyl acetate and 200 ml water. The
layers were separated and the organic layer was washed with 4 250-ml portions of 2 N
aqueous sodium hydroxide solution. The combined aqueous layers were extracted with 3
30 500-ml portions of ethyl acetate. The combined organic extracts were washed with
saturated aqueous sodium chloride solution and dried with sodium sulfate. Concentration
gave 5.44 g of a brown-red oil. Flash column chromatography afforded 2.15 g (41.3 %) of
the title compound as a light yellow solid.

- 25 -

MS m/e (%): 260 (M^+ , 11), M.p. 91 – 93°C.

b) 4-(5-Methylcarbamoyl-4-o-tolyl-pyridin-2-yl)-piperazine-1-carboxylic acid tert-butyl ester

A mixture of 8.31 g (31.9 mmol) 6-chloro-N-methyl-4-o-tolyl-nicotinamide, 6.53 g (35.0 mmol) 1-tert-butoxycarbonyl piperazine, 16.7 ml (95.6 mmol) N-ethyl-diisopropylamine and a catalytic amount of 4-(N,N-dimethylamino)-pyridine was heated at reflux over night. After cooling to room temperature the mixture was dissolved in dichloromethane and washed with two portions of 0.1 N aqueous hydrochloric acid solution. Drying with sodium sulfate and concentration gave 10.7 g of the crude product. Flash column chromatography afforded 6.28 g (48.0 %) of the title compound as an off-white solid.

MS m/e (%): 411 ($M+H^+$, 100).

c) 4-{5-[(3,5-Bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4-o-tolyl-pyridin-2-yl}-piperazine-1-carboxylic acid tert-butyl ester

To a solution of 6.28 g (15.3 mmol) 4-(5-methylcarbamoyl-4-o-tolyl-pyridin-2-yl)-piperazine-1-carboxylic acid tert-butyl ester in 250 ml tetrahydrofuran 20 ml of a 1 M solution (20 mmol) of potassium hexamethyldisilazide in tetrahydrofuran were added at 0 °C. After 30 min, 2.81 ml (15.3 mmol) 3,5-bis(trifluoromethyl)benzyl bromide were added dropwise. The reaction mixture was allowed to warm to room temperature over night. Addition of water and 1 M aqueous sodium hydroxide solution was followed by extraction with three portions of ethyl acetate. The combined organic extracts were dried with sodium sulfate and concentrated. Flash column chromatography gave 6.89 g (70.8 %) of the parent compound as a white solid.

MS m/e (%): 637 ($M+H^+$, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

25

Example 14

5'-[(3,5-Bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4'-o-tolyl-1-oxy-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-carboxylic acid ethyl ester

The parent compound was obtained as a white solid in comparable yields according to the procedures described above for the preparation of 4-{5-[(3,5-bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4-o-tolyl-pyridin-2-yl}-piperazine-1-carboxylic acid tert-butyl ester (Example 13) using ethyl isonipecotat instead of 1-tert-butoxycarbonyl piperazine in step b) and using 5'-methylcarbamoyl-4'-o-tolyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-

- 26 -

carboxylic acid ethyl ester instead of 4-(5-methylcarbamoyl-4-o-tolyl-pyridin-2-yl)-piperazine-1-carboxylic acid tert-butyl ester in step c).

MS m/e (%): 608 ($M+H^+$, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

5

Example 15

(RS)-6-[3-(Acetyl-methyl-amino)-1-oxo-pyrrolidin-1-yl]-N-(3,5-bis-trifluoromethyl-benzyl)-N-methyl-4-o-tolyl-nicotinamide

The parent compound was obtained as a light-yellow solid in comparable yields according to the procedures described above for the preparation of 4-{5-[(3,5-bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4-o-tolyl-pyridin-2-yl}-piperazine-1-carboxylic acid tert-butyl ester (Example 13) using (RS)-3-(acetyl-methyl-amino)-pyrrolidine instead of 1-tert-butoxycarbonyl piperazine in step b) and using (RS)-6-[3-(acetyl-methyl-amino)-pyrrolidin-1-yl]-N-methyl-4-o-tolyl-nicotinamide instead of 4-(5-methylcarbamoyl-4-o-tolyl-pyridin-2-yl)-piperazine-1-carboxylic acid tert-butyl ester in step c).

10 MS m/e (%): 593 ($M+H^+$, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

Example 16

N-(3,5-Bis-trifluoromethyl-benzyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide monohydrate

20 a) 6-Chloro-N-methyl-nicotinamide

To 50 g (317 mmol) of 2-chloronicotinic acid were added 230 ml (3.16 mol) thionyl chloride at 0 °C. After heating the mixture at reflux for 2 h excess thionyl chloride was removed by distillation. The oily brown residue was dissolved in 250 ml dichloromethane. The solution was treated with methylamine gas at 0 °C until no exothermic reaction was observed any longer. The resulting suspension was diluted with 1000 ml dichloromethane/water. The layers were separated and the aqueous layer extracted with three 300-ml portions of dichloromethane. Drying of the combined organic layers with sodium sulfate and concentration gave 53.2 g (98 %) of the title compound as a light yellow solid.

30 MS m/e (%): 171 ($M+H^+$, 15).

b) 6-Chloro-N-methyl-4-o-tolyl-nicotinamide

- 27 -

To a solution of 3.41 g (20.0 mmol) 6-chloro-N-methyl-nicotinamide in 80 ml tetrahydrofuran 50 ml (50 mmol) of a 1 M solution of o-tolyl magnesium chloride in tetrahydrofuran was added dropwise at 0 °C. After completed addition the reaction mixture was allowed to warm to room temperature and stirred for 1.5 h. The mixture was again cooled to 0 °C, followed by the dropwise addition of 5.7 ml (100 mmol) acetic acid and a solution of 5.1 g (22 mmol) 2,3-dichloro-5,6-dicyano-1,4-benzoquinone in 18 ml tetrahydrofuran. After completed addition the reaction mixture was allowed to warm to room temperature and stirred for 15 min. Addition of 30 ml 2 N aqueous sodium hydroxide solution was followed by dilution with 1 l ethyl acetate and 200 ml water. The layers were separated and the organic layer was washed with 4 250-ml portions of 2 N aqueous sodium hydroxide solution. The combined aqueous layers were extracted with 3 500-ml portions of ethyl acetate. The combined organic extracts were washed with saturated aqueous sodium chloride solution and dried with sodium sulfate. Concentration gave 5.44 g of a brown-red oil. Flash column chromatography afforded 2.15 g (41.3 %) of the title compound as a light yellow solid. M.p. 91–93 °C.

MS m/e (%): 260 (M^+ , 11).

c) N-Methyl-6-morpholin-4-yl-4-o-tolyl-nicotinamide

A mixture of 1.00 g (3.84 mmol) 6-chloro-N-methyl-4-o-tolyl-nicotinamide, 0.37 ml (4.22 mmol) morpholine, 2.0 ml (12 mmol) N-ethyl-diisopropylamine and a catalytic amount of 4-(N,N-dimethylamino)-pyridine was heated at 100 °C over night. After cooling to room temperature the mixture was dissolved in ethyl acetate and washed with two portions of water. The combined aqueous layers were extracted with 3 portions of dichloromethane. Drying with sodium sulfate and concentration gave 1.23 g of the crude product. Flash column chromatography afforded 1.11 g (92.9 %) of the title compound as an off-white solid. M.p. 156–158 °C.

MS m/e (%): 311 (M^+ , 64).

d) N-(3,5-Bis-trifluoromethyl-benzyl)-N-methyl-6-morpholin-4-yl-4-o-tolyl-nicotinamide

To a solution of 0.27 g (0.87 mmol) N-methyl-6-morpholin-4-yl-4-o-tolyl-nicotinamide in 15 ml tetrahydrofuran, 1.12 ml of a 1 M solution (1.12 mmol) of potassium hexamethyldisilazide in tetrahydrofuran was added at 0 °C. After 30 min, 0.16 ml (0.87 mmol) 3,5-bis(trifluoromethyl)benzyl bromide were added dropwise and the reaction mixture was allowed to warm to room temperature over night. Quenching with water was

- 28 -

followed by extraction with ethyl acetate. The combined organic extracts were dried with sodium sulfate and concentrated. Column chromatography gave 0.20 g (44 %) of the title compound as a white solid.

MS m/e (%): 538 (M+H⁺, 100).

5 e) N-(3,5-Bis-trifluoromethyl-benzyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide monohydrate

To a solution of 0.40 g (0.74 mmol) N-(3,5-bis-trifluoromethyl-benzyl)-N-methyl-6-morpholin-4-yl-4-o-tolyl-nicotinamide in 4 ml dichloromethane 0.17 g 3-chloroperbenzoic acid (70 %; 0.71 mmol) were added at 0 °C. After 4h the reaction
10 mixture was diluted with dichloromethane and washed with 3 portions of saturated sodium carbonate solution. The combined aqueous layers were extracted with dichloromethane. The combined organic extracts were washed with saturated sodium chloride solution, dried with sodium sulfate and concentrated. Column chromatography gave 0.31 g (73 %) of the title compound as a white solid.

15 MS m/e (%): 534 (M+H⁺, 100).

Crystallisation of a portion of 100 mg from a mixture of t-butylmethyl ether and cyclohexane afforded 90 mg of the title compound as white crystals. M.p. 116-117 °C.

Example 17

N-(3,5-Bis-trifluoromethyl-benzyl)-6-(1,1-dioxo-1λ⁶-4-oxy-thiomorpholin-4-yl)-N-methyl-4-o-tolyl-nicotinamide
20

a) N-(3,5-Bis-trifluoromethyl-benzyl)-N-methyl-6-thiomorpholin-4-yl-4-o-tolyl-nicotinamide

The title compound was obtained as a white solid in comparable yields according to the procedures described above for the preparation of 4-[5-[(3,5-bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4-o-tolyl-pyridin-2-yl]-piperazine-1-carboxylic acid tert-butyl ester
25 using thiomorpholine instead of 1-tert-butoxycarbonyl piperazine in step b) and using N-methyl-6-thiomorpholin-4-yl-4-o-tolyl-nicotinamide instead of 4-(5-methylcarbamoyl-4-o-tolyl-pyridin-2-yl)-piperazine-1-carboxylic acid tert-butyl ester in step c).

MS m/e (%): 554 (M+H⁺, 100).

30 b) N-(3,5-Bis-trifluoromethyl-benzyl)-N-methyl-6-(1-oxo-1λ⁴-thiomorpholin-4-yl)-4-o-tolyl-nicotinamide

- 29 -

To a solution of 1.24 g (2.24 mmol) N-(3,5-bis-trifluoromethyl-benzyl)-N-methyl-6-thiomorpholin-4-yl-4-o-tolyl-nicotinamide (step a)) in 25 ml methanol were added 689 mg (1.12 mmol) Oxone® at 0 °C. After completed addition the reaction mixture was allowed to warm to room temperature and stirred for 1.5 h. Quenching with 5 ml 40% aqueous sodium hydrogen sulfite solution was followed by addition of 6 ml 1N sodium hydroxide solution to adjust the pH to 7-8. The mixture was diluted with 50 ml water and extracted with 3 150-ml portions of dichloromethane. The combined extracts were dried with sodium sulfate and concentrated to give 1.20 g of crude product. Flash chromatography afforded 1.02 g (79.9 %) of the title compound as a white solid.

10 MS m/e (%): 570 (M+H⁺, 100).

c) N-(3,5-Bis-trifluoromethyl-benzyl)-6-(1,1-dioxo-1λ⁶-thiomorpholin-4-yl)-N-methyl-4-o-tolyl-nicotinamide

The parent compound was obtained as a white solid in comparable yield according to the procedure described above (step b)) using N-(3,5-bis-trifluoromethyl-benzyl)-N-methyl-6-(1-oxo-1λ⁴-thiomorpholin-4-yl)-4-o-tolyl-nicotinamide instead of N-(3,5-bis-trifluoromethyl-benzyl)-N-methyl-6-thiomorpholin-4-yl-4-o-tolyl-nicotinamide.

15 MS m/e (%): 586 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

Example 18

20 N-(3,5-Bis-trifluoromethyl-benzyl)-6-(4-formyl-1-oxy-piperazin-1-yl)-N-methyl-4-o-tolyl-nicotinamide

To a mixture of 0.089 ml (1.1 mmol) N,N-dimethylformamide and 38 mg (0.56 mmol) imidazole 0.071 ml (0.56 mmol) trimethylchlorosilane were added dropwise at room temperature. The reaction mixture was cooled to 0 °C, and 0.10 g (0.19 mmol) N-(3,5-bis-trifluoromethyl-benzyl)-N-methyl-6-piperazin-1-yl-4-o-tolyl-nicotinamide were added.

25 The ice-water bath was removed and the mixture stirred over night. The reaction was quenched with a mixture of 2 ml 1 N aqueous hydrochloric acid solution and 4 ml water, and the mixture was extracted with ethyl acetate. The combined extracts were dried with sodium sulfate and concentrated. Flash column chromatography afforded 81 mg (82 %) of the parent compound as a white solid.

30 MS m/e (%): 565 (M+H⁺, 100).

The title compound was obtained in accordance with step h) in Example 1.

Example 19

N-Methyl-N-(2-methyl-naphthalen-1-ylmethyl)-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide

a) N-Methyl-6-morpholin-4-yl-4-o-tolyl-nicotinamide

- 5 The title compound was obtained as an off-white solid in comparable yield according to the procedure described above for the preparation of 4-{5-[(3,5-bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4-o-tolyl-pyridin-2-yl}-piperazine-1-carboxylic acid tert-butyl ester (Example 13, step b) using morpholine instead of 1-tert-butoxycarbonyl piperazine.

MS m/e (%): 311 (M^+ , 63).

10 b) N-Methyl-N-(2-methyl-naphthalen-1-ylmethyl)-6-morpholin-4-yl-4-o-tolyl-nicotinamide

The parent compound was obtained as a white solid in comparable yield according to the procedure described above for the preparation of N-(3,5-bis-trifluoromethyl-benzyl)-N-methyl-6-morpholin-4-yl-4-o-tolyl-nicotinamide (Example 16, step d) using 1-

- 15 chloromethyl-2-methylnaphthalene instead of 3,5-bis-trifluoromethyl-benzyl bromide.
MS m/e (%): 466 ($M+H^+$, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

Example 20

N-Methyl-6-(4-oxy-morpholin-4-yl)-N-naphthalen-1-ylmethyl-4-o-tolyl-nicotinamide

- 20 The parent compound was obtained as a colorless viscous oil in comparable yields according to the procedures described above for the preparation of N-methyl-N-(2-methyl-naphthalen-1-ylmethyl)-6-morpholin-4-yl-4-o-tolyl-nicotinamide (Example 19) using 1-chloromethylnaphthalene instead of 1-chloromethyl-2-methylnaphthalene in step b).

- 25 MS m/e (%): 452 ($M+H^+$, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

Example 21

N-(2-Methoxy-naphthalen-1-ylmethyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide

- 30 The parent compound was obtained as a colorless viscous oil in comparable yields according to the procedures described above for the preparation of N-methyl-N-(2-

- 31 -

methyl-naphthalen-1-yl-methyl)-6-morpholin-4-yl-4-o-tolyl-nicotinamide (Example 19) using toluene-4-sulfonic acid 2-methoxy-naphthalen-1-yl-methyl ester instead of 1-chloromethyl-2-methylnaphthalene in step b).

MS m/e (%): 482 (M+H⁺, 100).

- 5 The N-oxide was obtained in accordance with step h) in Example 1.

Example 22

N-(2-Methoxy-benzyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide

- The parent compound was obtained as a colorless viscous oil in comparable yields according to the procedures described above for the preparation of N-methyl-N-(2-methyl-naphthalen-1-yl-methyl)-6-morpholin-4-yl-4-o-tolyl-nicotinamide (Example 19) using 2-methoxy-benzyl chloride instead of 1-chloromethyl-2-methylnaphthalene in step b).

MS m/e (%): 432 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

15

Example 23

N-(5-Chloro-2-methoxy-benzyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide

- The parent compound was obtained as a white solid in comparable yields according to the procedures described above for the preparation of N-methyl-N-(2-methyl-naphthalen-1-ylmethyl)-6-morpholin-4-yl-4-o-tolyl-nicotinamide (Example 19) using 5-chloro-2-methoxy-benzyl chloride instead of 1-chloromethyl-2-methylnaphthalene in step b).

MS m/e (%): 466 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

Example 24

- 25 N-(2-Chloro-5-methoxy-benzyl)-N-methyl-6-morpholin-4-yl-4-o-tolyl-nicotinamide

The parent compound was obtained as a white solid in comparable yields according to the procedures described above for the preparation of N-methyl-N-(2-methyl-naphthalen-1-ylmethyl)-6-morpholin-4-yl-4-o-tolyl-nicotinamide (Example 19) using 2-chloro-5-methoxy-benzyl bromide instead of 1-chloromethyl-2-methylnaphthalene in step b).

- 30 MS m/e (%): 466 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

- 32 -

Example 25

N-Methyl-6-(4-oxy-morpholin-4-yl)-N-pentafluorophenylmethyl-4-o-tolyl-nicotinamide

The parent compound was obtained as a white solid in comparable yields according to the procedures described above for the preparation of N-methyl-N-(2-methyl-naphthalen-1-ylmethyl)-6-morpholin-4-yl-4-o-tolyl-nicotinamide (Example 19) using 2,3,4,5,6-pentafluoro-benzyl bromide instead of 1-chloromethyl-2-methylnaphthalene in step b).

MS m/e (%): 492 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

Example 26

10 N-Methyl-6-(4-oxy-morpholin-4-yl)-N-naphthalen-2-ylmethyl-4-o-tolyl-nicotinamide

The parent compound was obtained as a white solid in comparable yields according to the procedures described above for the preparation of N-methyl-N-(2-methyl-naphthalen-1-ylmethyl)-6-morpholin-4-yl-4-o-tolyl-nicotinamide (Example 19) using 2-chloromethyl-naphthalene instead of 1-chloromethyl-2-methylnaphthalene in step b).

15 MS m/e (%): 452 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

Example 27

N-[2-Methoxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide

20 The parent compound was obtained as a white solid in comparable yields according to the procedures described above for the preparation of N-methyl-N-(2-methyl-naphthalen-1-ylmethyl)-6-morpholin-4-yl-4-o-tolyl-nicotinamide (Example 19) using toluene-4-sulfonic acid [2-methoxy-5-(5-trifluoromethyl-tetrazol-1-yl)-phenyl]-methyl ester instead of 1-chloromethyl-2-methylnaphthalene in step b).

25 MS m/e (%): 568 (M+H⁺, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

Example 28

N-(1,4-Dimethoxy-naphthalen-2-ylmethyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide

30 The parent compound was obtained as a colorless viscous oil in comparable yields according to the procedures described above for the preparation of N-methyl-N-(2-methyl-naphthalen-1-ylmethyl)-6-morpholin-4-yl-4-o-tolyl-nicotinamide (Example 19)

MS m/e (%): 512 ($M+H^+$, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

5. Example 29

A mixture of 200 mg (0.33 mmol) 5'-[(3,5-bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4'-o-tolyl-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-carboxylic acid ethyl ester

Extraction with dichloromethane, drying with sodium sulfate and flash column chromatography afforded 81 mg (42 %) of the parent compound as a white solid.

15 MS m/e (%): 580 ($M+H^+$, 100).

The N-oxide was obtained in accordance with step h) in Example 1.

- 34 -

Example A

Tablets of the following composition are manufactured in the usual manner:

	<u>mg/tablet</u>
Prodrug	5
5 Lactose	45
Corn starch	15
Microcrystalline cellulose	34
Magnesium stearate	1
Tablet weight	100

10

Example B

Capsules of the following composition are manufactured:

	<u>mg/capsule</u>
Prodrug	10
Lactose	155
15 Corn starch	30
Talc	5
Capsule fill weight	200

The active substance, lactose and corn starch are firstly mixed in a mixer and then in a comminuting machine. The mixture is returned to the mixer, the talc is added thereto
 20 and mixed thoroughly. The mixture is filled by machine into hard gelatine capsules.

Example C

Suppositories of the following composition are manufactured:

	<u>mg/supp.</u>
Prodrug	15
25 Suppository mass	1285
Total	1300

The suppository mass is melted in a glass or steel vessel, mixed thoroughly and cooled to 45°C. Thereupon, the finely powdered active substance is added thereto and


- 35 -

stirred until it has dispersed completely. The mixture is poured into suppository moulds of suitable size, left to cool, the suppositories are then removed from the moulds and packed individually in wax paper or metal foil.

- 36 -

Claims

1. Compounds of the general formula




5 wherein

R is hydrogen, lower alkyl, lower alkoxy, halogen or trifluoromethyl;

R¹ is hydrogen or halogen; orR and R¹ may be together with the ring carbon atoms to which they are attached
-CH=CH-CH=CH-;10 R² and R^{2'} are independently from each other hydrogen, halogen, trifluoromethyl, lower alkoxy or cyano; orR² and R^{2'} may be together -CH=CH-CH=CH-, optionally substituted by one or two substituents selected from lower alkyl or lower alkoxy;R³, R^{3'} are independently from each other hydrogen, lower alkyl or cycloalkyl;15 R⁴, R^{4'} are independently from each other -(CH₂)_mOR⁶ or lower alkyl;

or

R⁴ and R^{4'} form together with the N-atom to which they are attached a cyclic tertiary amine of the group20 R⁵ is hydrogen, hydroxy, lower alkyl, -lower alkoxy, -(CH₂)_mOH, -COOR³,
-CON(R³)₂, -N(R³)CO-lower alkyl or -C(O)R³;

- 37 -

R^6 is hydrogen, lower alkyl or phenyl;

X is $-C(O)N(R^6)-$, $-N(R^6)C(O)-$, $-(CH_2)_mO-$ or $-O(CH_2)_m-$;

n is 0, 1, 2, 3 or 4; and

m is 1, 2 or 3;

5 and pharmaceutically acceptable acid addition salts thereof.

2. A compound according to claim 1, wherein X is $-C(O)N(R^6)-$ and R^6 is methyl.

3. A compound according to claim 2, which is

4-{5-[(3,5-bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4-o-tolyl-pyridin-2-yl}-4-oxy-piperazine-1-carboxylic acid tert-butyl ester,

10 5'-[(3,5-bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4'-o-tolyl-1-oxy-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-carboxylic acid ethyl ester,

(RS)-6-[3-(acetyl-methyl-amino)-1-oxo-pyrrolidin-1-yl]-N-(3,5-bis-trifluoromethyl-benzyl)-N-methyl-4-o-tolyl-nicotinamide,

N-(3,5-bis-trifluoromethyl-benzyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-

15 nicotinamide monohydrate,

N-(3,5-bis-trifluoromethyl-benzyl)-6-(1,1-dioxo-1 λ^6 -4-oxy-thiomorpholin-4-yl)-N-methyl-4-o-tolyl-nicotinamide,

N-(3,5-bis-trifluoromethyl-benzyl)-6-(4-formyl-1-oxy-piperazin-1-yl)-N-methyl-4-o-tolyl-nicotinamide,

20 N-methyl-N-(2-methyl-naphthalen-1-ylmethyl)-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide,

N-methyl-6-(4-oxy-morpholin-4-yl)-N-naphthalen-1-ylmethyl-4-o-tolyl-nicotinamide,

N-(2-methoxy-naphthalen-1-ylmethyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide,

25 N-(2-methoxy-benzyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide,
N-(5-chloro-2-methoxy-benzyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide,

N-(2-chloro-5-methoxy-benzyl)-N-methyl-6-morpholin-4-yl-4-o-tolyl-nicotinamide,

N-methyl-6-(4-oxy-morpholin-4-yl)-N-pentafluorophenylmethyl-4-o-tolyl-nicotinamide,

30 N-methyl-6-(4-oxy-morpholin-4-yl)-N-naphthalen-2-ylmethyl-4-o-tolyl-nicotinamide,
N-[2-methoxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl]-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-tolyl-nicotinamide,

N-(1,4-dimethoxy-naphthalen-2-ylmethyl)-N-methyl-6-(4-oxy-morpholin-4-yl)-4-o-

tolyl-nicotinamide or

5'-[(3,5-bis-trifluoromethyl-benzyl)-methyl-carbamoyl]-4'-o-tolyl-1-oxy-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-carboxylic acid.

4. A compound according to claim 1, wherein X is $-N(R^6)C(O)-$ and R^6 is hydrogen
5 or methyl.

5. A compound according to claim 4, which is


- 2-(3,5-bis-trifluoromethyl-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-isobutyramide,
2-(3,5-bis-trifluoromethyl-phenyl)-N-[4-(2-chloro-phenyl)-6-(4-oxy-morpholin-4-yl)-
10 pyridin-3-yl]-N-methyl-isobutyramide,
2-(3,5-bis-trifluoromethyl-phenyl)-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-isobutyramide,
2-(3,5-bis-trifluoromethyl-phenyl)-N-[4'-(2-chloro-phenyl)-1-oxy-3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-5'-yl]-N-methyl-isobutyramide,
15 2-(3,5-bis-trifluoromethyl-phenyl)-N-(6-oxy-dimethylamino-4-o-tolyl-pyridin-3-yl)-N-methyl-isobutyramide,
2-(3,5-bis-trifluoromethyl-phenyl)-N-[4-(2-chloro-phenyl)-6-oxy-dimethylamino-pyridin-3-yl]-isobutyramide,
2-(3,5-bis-trifluoromethyl-phenyl)-N-1-(4-hydroxy-1-oxy-4'-o-tolyl-3,4,5,6-tetrahydro-
20 2H-[1,2']bipyridinyl-5'-yl)-N-methyl-isobutyramide,
2-(3,5-bis-trifluoromethyl-phenyl)-N-[6-[(2-hydroxy-ethyl)-1-oxy-methyl-amino]-4-o-tolyl-pyridin-3-yl]-N-methyl-isobutyramide,
(R)-2-(3,5-bis-trifluoromethyl-phenyl)-N-[6-(3-hydroxy-1-oxy-pyrrolidin-1-yl)-4-o-tolyl-pyridin-3-yl]-N-methyl-isobutyramide,
25 2-(3,5-bis-trifluoromethyl-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-acetamide,
2-(3,5-dimethoxy-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-acetamide or
2-(3-fluoro-5-trifluoromethyl-phenyl)-N-methyl-N-[6-(4-oxy-morpholin-4-yl)-4-o-tolyl-pyridin-3-yl]-acetamide.
30

6. A medicament containing one or more compounds of formula I as claimed in any one of claims 1-5 and pharmaceutically acceptable excipients.


7. A medicament according to claim 6 for the treatment of diseases related to NK-1 receptor antagonists.

8. A process for preparing a compound of formula I as defined in claim 1, which process comprises

5 oxidizing a compound of formula



with a suitable oxidizing agent to give a compound of formula




wherein the substituents have the significances given in claim I and
 10 if desired, converting the compound obtained into a pharmaceutically acceptable acid addition salt.

9. A compound of formula I according to any one of claims 1-5, whenever prepared by a process as claimed in claim 8 or by an equivalent method.

15 10. The use of a compound of formula I in any one of claims 1-5 for the manufacture of medicaments containing one or more compounds of formula I for the treatment of diseases related to NK-1 receptor antagonists.

11. A prodrug for a parent compound of formula II

- 40 -



for use in medicine,

wherein the prodrug is a compound of formula I according to any one of claims

1-5.

12. The invention as hereinbefore described.