

FACTOR Xa INHIBITORS

CROSS-REFERENCES TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Patent Application No. 60/580,899, filed
5 June 18, 2004, the content of which is incorporated herein by reference.

STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

[0002] NOT APPLICABLE

REFERENCE TO A "SEQUENCE LISTING," A TABLE, OR A COMPUTER PROGRAM LISTING APPENDIX SUBMITTED ON A COMPACT DISK.

[0003] NOT APPLICABLE

BACKGROUND OF THE INVENTION

[0004] Hemostasis, the control of bleeding, occurs by surgical means, or by the
physiological properties of vasoconstriction and coagulation. This invention is particularly
concerned with blood coagulation and ways in which it assists in maintaining the integrity of
mammalian circulation after injury, inflammation, disease, congenital defect, dysfunction or
other disruption. Although platelets and blood coagulation are both involved in thrombus
formation, certain components of the coagulation cascade are primarily responsible for the
amplification or acceleration of the processes involved in platelet aggregation and fibrin
deposition.

[0005] Thrombin is a key enzyme in the coagulation cascade as well as in hemostasis.
Thrombin plays a central role in thrombosis through its ability to catalyze the conversion of
fibrinogen into fibrin and through its potent platelet activation activity. Direct or indirect
inhibition of thrombin activity has been the focus of a variety of recent anticoagulant
strategies as reviewed by Claeson, G., "Synthetic Peptides and Peptidomimetics as Substrates
and Inhibitors of Thrombin and Other Proteases in the Blood Coagulation System", *Blood
Coag. Fibrinol.*, 5:411-436 (1994). Several classes of anticoagulants currently used in the

clinic directly or indirectly affect thrombin (i.e. heparins, low-molecular weight heparins, heparin-like compounds and coumarins).

- [0006] A prothrombinase complex, including Factor Xa (a serine protease, the activated form of its Factor X precursor and a member of the calcium ion binding, gamma carboxyglutamyl (Gla)-containing, vitamin K dependent, blood coagulation glycoprotein family), converts the zymogen prothrombin into the active procoagulant thrombin. Unlike thrombin, which acts on a variety of protein substrates as well as at a specific receptor, factor Xa appears to have a single physiologic substrate, namely prothrombin. Since one molecule of factor Xa may be able to generate up to 138 molecules of thrombin (Elodi *et al.*, *Thromb. Res.* 15:617-619 (1979)), direct inhibition of factor Xa as a way of indirectly inhibiting the formation of thrombin may be an efficient anticoagulant strategy. Therefore, it has been suggested that compounds which selectively inhibit factor Xa may be useful as *in vitro* diagnostic agents, or for therapeutic administration in certain thrombotic disorders, *see e.g.*, WO 94/13693.
- [0007] Polypeptides derived from hematophagous organisms have been reported which are highly potent and specific inhibitors of factor Xa. U.S. Patent No. 4,588,587 describes anticoagulant activity in the saliva of the Mexican leech, *Haementeria officinalis*. A principal component of this saliva was shown to be the polypeptide factor Xa inhibitor, antistasin (ATS), by Nutt, E. *et al.*, "The Amino Acid Sequence of Antistasin, a Potent Inhibitor of Factor Xa Reveals a Repeated Internal Structure", *J. Biol. Chem.*, 263:10162-10167 (1988). Another potent and highly specific inhibitor of Factor Xa, called tick anticoagulant peptide (TAP), has been isolated from the whole body extract of the soft tick *Ornithodoros moubata*, as reported by Waxman, L., *et al.*, "Tick Anticoagulant Peptide (TAP) is a Novel Inhibitor of Blood Coagulation Factor Xa", *Science*, 248:593-596 (1990).
- [0008] Factor Xa inhibitory compounds which are not large polypeptide-type inhibitors have also been reported (*see e.g.* Tidwell, R.R. *et al.*, "Strategies for Anticoagulation With Synthetic Protease Inhibitors. Xa Inhibitors Versus Thrombin Inhibitors", *Thromb. Res.*, 19:339-349 (1980); Turner, A.D. *et al.*, "*p*-Amidino Esters as Irreversible Inhibitors of Factor IXa and Xa and Thrombin", *Biochemistry*, 25:4929-4935 (1986); Hitomi, Y. *et al.*, "Inhibitory Effect of New Synthetic Protease Inhibitor (FUT-175) on the Coagulation System", *Haemostasis*, 15:164-168 (1985); Sturzebecher, J. *et al.*, "Synthetic Inhibitors of Bovine Factor Xa and Thrombin. Comparison of Their Anticoagulant Efficiency", *Thromb.*


Res., 54:245-252 (1989); Kam, C.M. *et al.*, "Mechanism Based Isocoumarin Inhibitors for Trypsin and Blood Coagulation Serine Proteases: New Anticoagulants", *Biochemistry*, 27:2547-2557 (1988); Hauptmann, J. *et al.*, "Comparison of the Anticoagulant and Antithrombotic Effects of Synthetic Thrombin and Factor Xa Inhibitors", *Thromb. Haemost.*, 63:220-223 (1990)).

[0009] Others have reported Factor Xa inhibitors which are small molecule organic compounds, such as nitrogen containing heterocyclic compounds which have amidino substituent groups, wherein two functional groups of the compounds can bind to Factor Xa at two of its active sites. For example, WO 98/28269 describes pyrazole compounds having a terminal C(=NH)-NH₂ group; WO 97/21437 describes benzimidazole compounds substituted by a basic radical which are connected to a naphthyl group via a straight or branched chain alkylene, C(=O) or S(=O)₂ bridging group; WO 99/10316 describes compounds having a 4-phenyl-N-alkylamidino-piperidine and 4-phenoxy-N-alkylamidino-piperidine group connected to a 3-amidinophenyl group via a carboxamidealkyleneamino bridge; and EP 798295 describes compounds having a 4-phenoxy-N-alkylamidino-piperidine group connected to an amidinonaphthyl group via a substituted or unsubstituted sulfonamide or carboxamide bridging group.

[0010] There exists a need for effective therapeutic agents for the regulation of hemostasis, and for the prevention and treatment of thrombus formation and other pathological processes in the vasculature induced by thrombin such as restenosis and inflammation. In particular, there continues to be a need for compounds which selectively inhibit factor Xa or its precursors. Compounds that have different combinations of bridging groups and functional groups than compounds previously discovered are needed, particularly compounds which selectively or preferentially bind to Factor Xa. Compounds with a higher degree of binding to Factor Xa than to thrombin are desired, especially those compounds having good bioavailability and/or solubility.

BRIEF SUMMARY OF THE INVENTION


[0011] The present invention provides in one aspect, compounds having the formula:



(I)

and pharmaceutically acceptable salts thereof. In formula (I), A represents benzene, a six-membered heterocycle containing 1-3 nitrogen atoms or a five-membered heterocycle containing 1-4 nitrogen, oxygen or sulfur atoms; with 0-4 substituents selected from halogen, oxo, R^A , $-OR^A$, $N(R^A)_2$, $CO_2 R^A$, $CON(R^A)_2$, $S(O)_2 R^A$, $S(O)_2 N(R^A)_2$, $X^1 OR^A$, $X^1 CO_2 R^A$, $X^1 CON(R^A)_2$, $X^1 N(R^A)_2$, wherein each R^A is independently selected from H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, or optionally two R^A attached to the same atom are combined to form a 3-, 4-, 5-, 6- or 7-membered ring; R represents a member selected from the group consisting of H and C_{1-4} alkyl; R^1 represents a member selected from the group consisting of halogen, C_{1-8} alkyl, C_{2-8} alkenyl and C_{2-8} alkynyl; R^2 represents a member selected from the group consisting of H, C_{1-8} alkyl, $-X-OR^{2a}$, $-X-SR^{2a}$, $-X-COR^{2a}$, $-X-CO_2 R^{2a}$ and $-X-N(R^{2a})_2$, wherein X is C_{1-8} alkylene and each R^{2a} is independently selected from H and C_{1-8} alkyl, or optionally two R^{2a} groups attached to the same nitrogen atom are combined to form a 4-, 5-, 6- or 7-membered ring.

[0012] The letter W represent a moiety having a formula selected from (a), (b), and (c):



[0013] With reference to formula (b), the symbols Z^1 , Z^2 , and Z^3 each independently represent N, C or CH with the proviso that two of Z^1 , Z^2 , and Z^3 are other than N.

[0014] With reference to formulas (a) and (b), the subscript n is an integer of from 0 to 3 and the subscript m is an integer of from 0 to 1. The symbol R^4 represents, in each occurrence, a member selected from the group consisting of halogen, $-OR^{4a}$, $-OC(O)R^{4a}$, $-NR^{4a}R^{4b}$, $-SR^{4a}$, $S(O)R^{4a}$, $S(O)_2 R^{4a}$, $S(O)_2 NR^{4a}R^{4b}$, $-R^{4c}$, $-CN$, $-NO_2$, $-CO_2 R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-OC(O)NR^{4a}R^{4b}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)_2 R^{4c}$, $-NR^{4a}-C(O)NR^{4a}R^{4b}$, $-X^1 OR^{4a}$, $-X^1 OC(O)R^{4a}$, $-X^1 NR^{4a}R^{4b}$, $-X^1 SR^{4a}$, $-X^1 S(O)R^{4a}$, $-X^1 S(O)_2 R^{4a}$, $-X^1 CN$, $-X^1 NO_2$, $-X^1 CO_2 R^{4a}$, $-O-X^1 CO_2 R^{4a}$, $-X^1 CONR^{4a}R^{4b}$, $-O-X^1 CONR^{4a}R^{4b}$, $-X^1 C(O)R^{4a}$, $-O-X^1 NR^{4a}R^{4b}$, $-S(O)X^1 NR^{4a}R^{4b}$, $-S(O)_2 X^1 NR^{4a}R^{4b}$, $-X^1 OC(O)NR^{4a}R^{4b}$, $-X^1 NR^{4a}C(O)R^{4a}$, $-X^1 NR^{4b}C(O)_2 R^{4c}$, $-X^1 NR^{4a}C(O)NR^{4a}R^{4b}$, $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2 Y$. Similarly, $R^{4'}$ represents a member selected from the group consisting of $-C(=NH)R^{4a}$, $-C(=NR^{4c})R^{4a}$, $-C(=NH)NR^{4a}R^{4b}$, $-C(=NR^{4c})NR^{4a}R^{4b}$, $-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$, $-X^2-C(=NH)NR^{4a}R^{4b}$,

- $-X^2-C(=NR^{4c})NR^{4a}R^{4b}$, $-X^2-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$ and $-C(=NR^{4a})NR^{4a}-Y$, wherein Y is a five or six-membered aryl, heterocyclyl or aryl- C_{1-2} alkyl, optionally substituted with from one to three substituents selected from the group consisting of halogen, oxo, $-OR^{4a}$, $-OC(O)R^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2NR^{4a}R^{4b}$, $-CN$, $-NO_2$, $-CO_2R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)_2R^{4c}$, $-X^1R^{4a}$, $-X^1OR^{4a}$, $-X^1SR^{4a}$, $-X^1S(O)R^{4a}$, $-X^1S(O)_2R^{4a}$, $-X^1CN$, $-X^1NO_2$, $-X^1CO_2R^{4a}$, $-X^1CONR^{4a}R^{4b}$, $-X^1C(O)R^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-S(O)X^1NR^{4a}R^{4b}$, $-S(O)_2X^1NR^{4a}R^{4b}$, $-X^1OC(O)NR^{4a}R^{4b}$, $-X^1NR^{4b}C(O)R^{4a}$, $-X^1NR^{4b}C(O)_2R^{4c}$, $-X^1NR^{4a}-C(O)NR^{4a}R^{4b}$, $-X^1OC(O)R^{4a}$, $-X^1NR^{4a}R^{4b}$, $-O-X^1OR^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-O-X^1CO_2R^{4a}$, $-O-X^1CONR^{4a}R^{4b}$, $-NR^{4b}-X^1OR^{4a}$, $-NR^{4b}-X^1NR^{4a}R^{4b}$, $-NR^{4b}-X^1CO_2R^{4a}$, and $-NR^{4b}-X^1CONR^{4a}R^{4b}$; each X^1 and X^2 are members independently selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene; each R^{4a} and R^{4b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; each R^{4c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl, and optionally any two of R^{4a} , R^{4b} and R^{4c} when part of a common R^4 or R^4 substituent can be combined with the atoms to which each is attached to form a saturated or unsaturated, four to nine-membered mono- or bicyclic ring having from 0 to 2 additional heteroatoms as ring members. Additionally, each of R^{4a} , R^{4b} and R^{4c} is optionally further substituted with from one to three members selected from the group consisting of $-OH$, oxo, $-R^m$, $-OR^m$, $-OC(O)NHR^m$, $-OC(O)N(R^m)_2$, $-SH$, $-SR^m$, $-S(O)R^m$, $-S(O)_2R^m$, $-SO_2NH_2$, $-S(O)_2NHR^m$, $-S(O)_2N(R^m)_2$, $-NHS(O)_2R^m$, $-NR^mS(O)_2R^m$, $-C(O)NH_2$, $-C(O)NHR^m$, $-C(O)N(R^m)_2$, $-C(O)R^m$, $-NHC(O)R^m$, $-NR^mC(O)R^m$, $-NHC(O)NH_2$, $-NR^mC(O)NH_2$, $-NR^mC(O)NHR^m$, $-NHC(O)NHR^m$, $-NR^mC(O)N(R^m)_2$, $-NHC(O)N(R^m)_2$, $-CO_2H$, $-CO_2R^m$, $-NHCO_2R^m$, $-NR^mCO_2R^m$, $-CN$, $-NO_2$, $-NH_2$, $-NHR^m$, $-N(R^m)_2$, $-NR^mS(O)NH_2$ and $-NR^mS(O)_2NHR^m$, wherein each R^m is independently an unsubstituted C_{1-6} alkyl, benzyl or combined with the atoms to which each is attached to form a saturated four-, five-, six- or seven-membered ring having from 0 to 2 additional heteroatoms as ring members.
- [0015]** Turning next to formula (c), the symbol R^5 represents a member selected from the group consisting of $-R^{5a}$, $-CO_2R^{5c}$, $-CONR^{5a}R^{5b}$, $-C(O)R^{5a}$, $-C(=NR^{5a})NR^{5a}R^{5b}$, $-C(=N^+R^{5c}R^{5c})NR^{5a}R^{5b}$, $-C(=NR^{5a})R^{5a}$, $-C(=NR^{5a})Y^1$, $-X^3OR^{5a}$, $-X^3OC(O)R^{5a}$, $-X^3NR^{5a}R^{5b}$, $-X^3SR^{5a}$, $-X^3CN$, $-X^3NO_2$, $-X^3CO_2R^{5a}$, $-X^3CONR^{5a}R^{5b}$, $-X^3C(O)R^{5a}$, $-X^3OC(O)NR^{5a}R^{5b}$, -

$X^3NR^{5a}C(O)R^{5a}$, $-X^3NR^{5b}C(O)_2R^{5c}$, $-X^3NR^{5a}C(O)NR^{5a}R^{5b}$, $-X^3-C(=NR^{5a})NR^{5a}R^{5b}$, $-Y^1$, and $-X^3-Y^1$, wherein Y^1 is a five or six-membered aryl or heteroaryl ring or aryl- C_{1-2} alkyl, optionally substituted with from one to three substituents selected from the group consisting of halogen, $-OR^{5a}$, $-OC(O)R^{5a}$, $-NR^{5a}R^{5b}$, $-R^{5c}$, $-SR^{5a}$, $-CN$, $-NO_2$, $-CO_2R^{5a}$, $-CONR^{5a}R^{5b}$, $-C(O)R^{5a}$, $-NR^{5b}C(O)R^{5a}$, $-NR^{5b}C(O)_2R^{5c}$, $-X^3OR^{5a}$, $-X^3SR^{5a}$, $-X^3CN$, $-X^3NO_2$, $-X^3CO_2R^{5a}$, $-X^3CONR^{5a}R^{5b}$, $-X^3C(O)R^{5a}$, $-X^3OC(O)NR^{5a}R^{5b}$, $-X^3NR^{5b}C(O)R^{5a}$, $-X^3NR^{5b}C(O)_2R^{5c}$, $-X^3NR^{5a}-C(O)NR^{5a}R^{5b}$, $-X^3OC(O)R^{5a}$, $-X^3NR^{5a}R^{5b}$, $-O-X^3OR^{5a}$, $-O-X^3NR^{5a}R^{5b}$, $-O-X^3CO_2R^{5a}$, $-O-X^3CONR^{5a}R^{5b}$, $-NR^{5b}-X^3OR^{5a}$, $-NR^{5b}-X^3NR^{5a}R^{5b}$, $-NR^{5b}-X^3CO_2R^{5a}$, and $-NR^{5b}-X^3CONR^{5a}R^{5b}$, wherein each X^3 is a member independently selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene; each R^{5a} and R^{5b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; each R^{5c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; and optionally any two of R^{5a} , R^{5b} and R^{5c} when part of a common R^5 substituent can be combined with the atoms to which each is attached to form a saturated four-, five-, six- or seven-membered ring having from 0 to 1 additional heteroatoms as ring members. Additionally, each of R^{5a} , R^{5b} and R^{5c} is optionally further substituted with from one to three members selected from the group consisting of $-OH$, $-OR^n$, $-OC(O)NHR^n$, $-OC(O)N(R^n)_2$, $-SH$, $-SR^n$, $-S(O)R^n$, $-S(O)_2R^n$, $-SO_2NH_2$, $-S(O)_2NHR^n$, $-S(O)_2N(R^n)_2$, $-NHS(O)_2R^n$, $-NR^nS(O)_2R^n$, $-C(O)NH_2$, $-C(O)NHR^n$, $-C(O)N(R^n)_2$, $-C(O)R^n$, $-NHC(O)R^n$, $-NR^nC(O)R^n$, $-NHC(O)NH_2$, $-NR^nC(O)NH_2$, $-NR^nC(O)NHR^n$, $-NHC(O)NHR^n$, $-NR^nC(O)N(R^n)_2$, $-NHC(O)N(R^n)_2$, $-CO_2H$, $-CO_2R^n$, $-NHCO_2R^n$, $-NR^nCO_2R^n$, $-CN$, $-NO_2$, $-NH_2$, $-NHR^n$, $-N(R^n)_2$, $-NR^nS(O)NH_2$ and $-NR^nS(O)_2NHR^n$, wherein each R^n is independently an unsubstituted C_{1-6} alkyl or benzyl.

[0016] The present invention further provides chemical intermediates, pharmaceutical compositions and methods for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising the step of administering to said mammal a therapeutically effective amount of a compound of the present invention. Such conditions include but are not limited to acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous

thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, conditions requiring the fitting of prosthetic devices, and the like.

[0017] The present invention further provides methods for inhibiting the coagulation of a blood sample comprising contacting said sample with a compound of the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0018] Not applicable

DETAILED DESCRIPTION OF THE INVENTION

Abbreviations and Definitions

[0019] The term "alkyl", by itself or as part of another substituent, means, unless otherwise stated, a straight or branched chain hydrocarbon radical, having the number of carbon atoms designated (*i.e.* C₁₋₈ means one to eight carbons). Examples of alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. The term "alkenyl" refers to an unsaturated alkyl group is one having one or more double bonds. Similarly, the term "alkynyl" refers to an unsaturated alkyl group having one or more triple bonds. Examples of such unsaturated alkyl groups include vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butylnyl, and the higher homologs and isomers. The term "cycloalkyl" refers to hydrocarbon rings having the indicated number of ring atoms (e.g., C₃₋₆cycloalkyl) and being fully saturated or having no more than one double bond between ring vertices. When "cycloalkyl" is used in combination with "alkyl", as in C₃₋₅cycloalkyl-alkyl, the cycloalkyl portion is meant to have from three to five carbon atoms, while the alkyl portion is an alkylene moiety having from one to three carbon atoms (e.g., -CH₂-, -CH₂CH₂- or -CH₂CH₂CH₂-).

[0020] The term "alkylene" by itself or as part of another substituent means a divalent radical derived from an alkane, as exemplified by -CH₂CH₂CH₂CH₂-. Typically, an alkyl (or

alkylene) group will have from 1 to 24 carbon atoms, with those groups having 10 or fewer carbon atoms being preferred in the present invention. A "lower alkyl" or "lower alkylene" is a shorter chain alkyl or alkylene group, generally having four or fewer carbon atoms.

5 [0021] The terms "alkoxy," "alkylamino" and "alkylthio" (or thioalkoxy) are used in their conventional sense, and refer to those alkyl groups attached to the remainder of the molecule via an oxygen atom, an amino group, or a sulfur atom, respectively. Additionally, for dialkylamino groups (typically provided as $-NR^aR^b$ or a variant thereof), the alkyl portions can be the same or different and can also be combined to form a 3-7 membered ring with the nitrogen atom to which each is attached. Accordingly, a group represented as $-NR^aR^b$ is
10 meant to include piperidinyl, pyrrolidinyl, morpholinyl, azetidiny and the like.

[0022] The terms "halo" or "halogen," by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as "haloalkyl," are meant to include monohaloalkyl and polyhaloalkyl. For example, the term "C₁₋₄ haloalkyl" is mean to include trifluoromethyl, 2,2,2-trifluoroethyl, 4-
15 chlorobutyl, 3-bromopropyl, and the like.

[0023] The term "aryl" means, unless otherwise stated, a polyunsaturated, typically aromatic, hydrocarbon group which can be a single ring or multiple rings (up to three rings) which are fused together or linked covalently. The term "heteroaryl" refers to aryl groups (or rings) that contain from one to five heteroatoms selected from N, O, and S, wherein the
20 nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. A heteroaryl group can be attached to the remainder of the molecule through a heteroatom or through a carbon atom. Non-limiting examples of aryl groups include phenyl, naphthyl and biphenyl, while non-limiting examples of heteroaryl groups include 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 1-pyrazolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-
25 thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl, 2-benzimidazolyl, benzopyrazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxaliny, 5-quinoxaliny, 3-quinolyl, and 6-quinolyl. If not specifically stated, substituents for each of the above noted aryl and
30 heteroaryl ring systems are selected from the group of acceptable substituents described below.

[0024] For brevity, the term "aryl" when used in combination with other terms (*e.g.*, aryloxy, arylthioxy, arylalkyl) includes both aryl and heteroaryl rings as defined above. Thus, the term "arylalkyl" is meant to include those radicals in which an aryl group is attached to an alkyl group (*e.g.*, benzyl, phenethyl, pyridylmethyl and the like).

5 [0025] The term "heterocycle" or "heterocycle" refers to a saturated or unsaturated non-aromatic cyclic group containing at least one sulfur, nitrogen or oxygen heteroatom. Each heterocycle can be attached at any available ring carbon or heteroatom. Each heterocycle may have one or more rings. When multiple rings are present, they can be fused together or linked covalently. Each heterocycle must contain at least one heteroatom (typically 1 to 5
10 heteroatoms) selected from nitrogen, oxygen or sulfur. Preferably, these groups contain 0-5 nitrogen atoms, 0-2 sulfur atoms and 0-2 oxygen atoms. More preferably, these groups contain 0-3 nitrogen atoms, 0-1 sulfur atoms and 0-1 oxygen atoms. Non-limiting examples of heterocycle groups include benzene, pyridine, pyridimidine, pyrazine, morpholin-3-one, piperazine-2-one, pyridine-2-one, piperidine, morpholine, piperazine, isoxazole, isothiazole,
15 pyrazole, imidazole, oxazole, thiazole, isoxazoline, pyrazoline, imidazoline, 1,2,3-triazole, 1,2,4-triazole, 1,3,4-oxadiazole, 1,3,4-thiadiazole, 1,2,4-oxadiazole, 1,2,4-thiadiazole, pyrazol-5-one, pyrrolidine-2,5-dione, imidazolidine-2,4-dione, pyrrolidine, pyrrole, furan, thiophene, and the like.

[0026] The above terms (*e.g.*, "aryl" and "heteroaryl"), in some embodiments, will include
20 both substituted and unsubstituted forms of the indicated radical. Preferred substituents for each type of radical are provided below. For brevity, the terms aryl and heteroaryl will refer to substituted or unsubstituted versions as provided below.

[0027] Substituents for the aryl and heteroaryl groups are varied and are generally selected from: -halogen, -oxo, -OR', -OC(O)R', -NR'R'', -SR', -R', -CN, -NO₂, -CO₂R', -CONR'R'',
25 -C(O)R', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)₂R', -NR'-C(O)NR''R'', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NR'S(O)₂R'', -N₃, perfluoro(C₁-C₄)alkoxy, and perfluoro(C₁-C₄)alkyl, in a number ranging from zero to the total number of open valences on the aromatic ring system; and where R', R'' and R''' are independently selected from hydrogen, C₁₋₈alkyl, C₃₋₆cycloalkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, unsubstituted aryl and heteroaryl, (unsubstituted aryl)-C₁₋₄alkyl, and unsubstituted
30 aryloxy-C₁₋₄alkyl. Other suitable substituents include each of the above aryl substituents attached to a ring atom by an alkylene tether of from 1-4 carbon atoms.

[0028] Two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula $-T-C(O)-(CH_2)_q-U-$, wherein T and U are independently $-NH-$, $-O-$, $-CH_2-$ or a single bond, and q is an integer of from 0 to 2.

Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may

optionally be replaced with a substituent of the formula $-A-(CH_2)_r-B-$, wherein A and B are independently $-CH_2-$, $-O-$, $-NH-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-S(O)_2NR'-$ or a single bond, and r is an integer of from 1 to 3. One of the single bonds of the new ring so formed may optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula $-(CH_2)_s-$
X-(CH₂)_t-, where s and t are independently integers of from 0 to 3, and X is $-O-$, $-NR'-$, $-S-$, $-S(O)-$, $-S(O)_2-$, or $-S(O)_2NR'-$. The substituent R' in $-NR'-$ and $-S(O)_2NR'-$ is selected from hydrogen or unsubstituted C₁₋₆alkyl.

[0029] As used herein, the term "heteroatom" is meant to include oxygen (O), nitrogen (N), sulfur (S) and silicon (Si).

[0030] The term "pharmaceutically acceptable salts" is meant to include salts of the active compounds which are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of salts derived from pharmaceutically-acceptable inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic, manganous, potassium, sodium, zinc and the like. Salts derived from pharmaceutically-acceptable organic bases include salts of primary, secondary and tertiary amines, including substituted amines, cyclic amines, naturally-
occurring amines and the like, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid

addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, malonic, benzoic, succinic, suberic, fumaric, mandelic, phthalic, benzenesulfonic, *p*-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (*see, e.g., Berge, S.M., et al, "Pharmaceutical Salts", Journal of Pharmaceutical Science, 1977, 66, 1-19*). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

[0031] The neutral forms of the compounds may be regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form of the compound for the purposes of the present invention.

[0032] In addition to salt forms, the present invention provides compounds which are in a prodrug form. Prodrugs of the compounds described herein are those compounds that readily undergo chemical changes under physiological conditions to provide the compounds of the present invention. Additionally, prodrugs can be converted to the compounds of the present invention by chemical or biochemical methods in an *ex vivo* environment. For example, prodrugs can be slowly converted to the compounds of the present invention when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent.


[0033] Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention. Certain compounds of the present invention may exist in multiple crystalline or amorphous forms. In general, all physical forms are equivalent for the uses contemplated by the present invention and are intended to be within the scope of the present invention.

[0034] Certain compounds of the present invention possess asymmetric carbon atoms (optical centers) or double bonds; the racemates, diastereomers, geometric isomers, regioisomers and individual isomers (e.g., separate enantiomers) are all intended to be

encompassed within the scope of the present invention. The compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I) or carbon-14 (^{14}C). All isotopic variations of the compounds of the present invention, whether radioactive or not, are intended to be encompassed within the scope of the present invention.


General**Embodiments of the Invention****Compounds**

[0035] In one aspect, the present invention provides compounds having the formula:



and pharmaceutically acceptable salts thereof. In formula (I), A represents benzene, a six-membered heterocycle containing 1-3 nitrogen atoms or a five-membered heterocycle containing 1-4 nitrogen, oxygen or sulfur atoms; with 0-4 substituents selected from halogen, oxo, R^A , $-OR^A$, $N(R^A)_2$, CO_2R^A , $CON(R^A)_2$, $S(O)_2R^A$, $S(O)_2N(R^A)_2$, X^1OR^A , $X^1CO_2R^A$, $X^1CON(R^A)_2$, $X^1N(R^A)_2$, wherein each R^A is independently selected from H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, or optionally two R^A attached to the same atom are combined to form a 3-, 4-, 5-, 6- or 7-membered ring; R represents a member selected from the group consisting of H and C_{1-4} alkyl; R^1 represents a member selected from the group consisting of halogen, C_{1-8} alkyl, C_{2-8} alkenyl and C_{2-8} alkynyl; R^2 represents a member selected from the group consisting of H, C_{1-8} alkyl, $-X-OR^{2a}$, $-X-SR^{2a}$, $-X-COR^{2a}$, $-X-CO_2R^{2a}$ and $-X-N(R^{2a})_2$; wherein X is C_{1-8} alkylene and each R^{2a} is independently selected from the group consisting of H and C_{1-8} alkyl, or optionally two R^{2a} groups attached to the same nitrogen atom are combined to form a 4-, 5-, 6- or 7-membered ring. For the embodiments in which R^2 is present and other than H, a preferred group of embodiments are those in which the carbon to which R^2 is attached has the (R)- stereochemical configuration.

[0036] The letter W represent a moiety having a formula selected from (a), (b), and (c):



wherein the wavy line indicates the point of attachment to the rest of the molecule.

[0037] With reference to formula (b), the symbols Z^1 , Z^2 , and Z^3 each independently represent N, C or CH with the proviso that two of Z^1 , Z^2 , and Z^3 are other than N. The pyridyl

nitrogen can be in a position *para* to the attachment of the A ring (Z^1 (4-pyridyl)), meta (Z^2 (3-pyridyl)) or ortho (Z^3 (2-pyridyl)).

[0038] With reference to formulas (a) and (b), the subscript n is an integer of from 0 to 3 and the subscript m is an integer of from 0 to 1. The symbol R^4 represents, in each


- 5 occurrence, a member selected from the group consisting of halogen, $-OR^{4a}$, $-OC(O)R^{4a}$, $-NR^{4a}R^{4b}$, $-SR^{4a}$, $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2NR^{4a}R^{4b}$, $-R^{4c}$, $-CN$, $-NO_2$, $-CO_2R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-OC(O)NR^{4a}R^{4b}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)_2R^{4c}$, $-NR^{4a}-C(O)NR^{4a}R^{4b}$, $-X^1OR^{4a}$, $-X^1OC(O)R^{4a}$, $-X^1NR^{4a}R^{4b}$, $-X^1SR^{4a}$, $-X^1S(O)R^{4a}$, $-X^1S(O)_2R^{4a}$, $-X^1CN$, $-X^1NO_2$, $-X^1CO_2R^{4a}$, $-O-X^1CO_2R^{4a}$, $-X^1CONR^{4a}R^{4b}$, $-O-X^1CONR^{4a}R^{4b}$, $-X^1C(O)R^{4a}$, $-O-X^1NR^{4a}R^{4b}$, -
- 10 $S(O)X^1NR^{4a}R^{4b}$, $-S(O)_2X^1NR^{4a}R^{4b}$, $-X^1OC(O)NR^{4a}R^{4b}$, $-X^1NR^{4a}C(O)R^{4a}$, $-X^1NR^{4b}C(O)_2R^{4c}$, $-X^1NR^{4a}C(O)NR^{4a}R^{4b}$, $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2Y$. Similarly, $R^{4'}$ represents a member selected from the group consisting of $-C(=NH)R^{4a}$, $-C(=NR^{4c})R^{4a}$, $-C(=NH)NR^{4a}R^{4b}$, $-C(=NR^{4c})NR^{4a}R^{4b}$, $-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$, $-X^2-C(=NH)NR^{4a}R^{4b}$, $-X^2-C(=NR^{4c})NR^{4a}R^{4b}$, $-X^2-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$ and $-C(=NR^{4a})NR^{4a}-Y$; wherein Y is a
- 15 five or six-membered aryl, heterocyclyl or aryl- C_{1-2} alkyl, optionally substituted with from one to three substituents selected from the group consisting of halogen, oxo, $-OR^{4a}$, $-OC(O)R^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2NR^{4a}R^{4b}$, $-CN$, $-NO_2$, $-CO_2R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)_2R^{4c}$, $-X^1R^{4a}$, $-X^1OR^{4a}$, $-X^1SR^{4a}$, -
- 20 $X^1S(O)R^{4a}$, $-X^1S(O)_2R^{4a}$, $-X^1CN$, $-X^1NO_2$, $-X^1CO_2R^{4a}$, $-X^1CONR^{4a}R^{4b}$, $-X^1C(O)R^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-S(O)X^1NR^{4a}R^{4b}$, $-S(O)_2X^1NR^{4a}R^{4b}$, $-X^1OC(O)NR^{4a}R^{4b}$, $-X^1NR^{4b}C(O)R^{4a}$, $-X^1NR^{4b}C(O)_2R^{4c}$, $-X^1NR^{4a}-C(O)NR^{4a}R^{4b}$, $-X^1OC(O)R^{4a}$, $-X^1NR^{4a}R^{4b}$, $-O-X^1OR^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-O-X^1CO_2R^{4a}$, $-O-X^1CONR^{4a}R^{4b}$, $-NR^{4b}-X^1OR^{4a}$, $-NR^{4b}-X^1NR^{4a}R^{4b}$, $-NR^{4b}-X^1CO_2R^{4a}$, and $-NR^{4b}-X^1CONR^{4a}R^{4b}$; each X^1 and X^2 are members independently selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene; each
- 25 R^{4a} and R^{4b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; each R^{4c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl, and optionally any two of
- 30 R^{4a} , R^{4b} and R^{4c} when part of a common R^4 or $R^{4'}$ substituent can be combined with the atoms to which each is attached to form a saturated or unsaturated, four to nine-membered mono- or bicyclic ring having from 0 to 2 additional heteroatoms as ring members. Additionally, each of R^{4a} , R^{4b} and R^{4c} is optionally further substituted with from one to three members selected

from the group consisting of -OH, oxo, -R^m, -OR^m, -OC(O)NHR^m, -OC(O)N(R^m)₂, -SH, -SR^m, -S(O)R^m, -S(O)₂R^m, -SO₂NH₂, -S(O)₂NHR^m, -S(O)₂N(R^m)₂, -NHS(O)₂R^m, -NR^mS(O)₂R^m, -C(O)NH₂, -C(O)NHR^m, -C(O)N(R^m)₂, -C(O)R^m, -NHC(O)R^m, -NR^mC(O)R^m, -NHC(O)NH₂, -NR^mC(O)NH₂, -NR^mC(O)NHR^m, -NHC(O)NHR^m, -NR^mC(O)N(R^m)₂, -NHC(O)N(R^m)₂, -CO₂H, -CO₂R^m, -NHCO₂R^m, -NR^mCO₂R^m, -CN, -NO₂, -NH₂, -NHR^m, -N(R^m)₂, -NR^mS(O)NH₂ and -NR^mS(O)₂NHR^m, wherein each R^m is independently an unsubstituted C₁₋₆alkyl, benzyl or combined with the atoms to which each is attached to form a saturated four-, five-, six- or seven-membered ring having from 0 to 2 additional heteroatoms as ring members.


- 10 **[0039]** Turning next to formula (c), the symbol R⁵ represents a member selected from the group consisting of -R^{5a}, -CO₂R^{5c}, -CONR^{5a}R^{5b}, -C(O)R^{5a}, -C(=NR^{5a})NR^{5a}R^{5b}, -C(=N⁺R^{5c}R^{5c})NR^{5a}R^{5b}, -C(=NR^{5a})R^{5a}, -C(=NR^{5a})Y¹, -X³OR^{5a}, -X³OC(O)R^{5a}, -X³NR^{5a}R^{5b}, -X³SR^{5a}, -X³CN, -X³NO₂, -X³CO₂R^{5a}, -X³CONR^{5a}R^{5b}, -X³C(O)R^{5a}, -X³OC(O)NR^{5a}R^{5b}, -X³NR^{5a}C(O)R^{5a}, -X³NR^{5b}C(O)₂R^{5c}, -X³NR^{5a}C(O)NR^{5a}R^{5b}, -X³-C(=NR^{5a})NR^{5a}R^{5b}, -Y¹, and
- 15 -X³-Y¹, wherein Y¹ is a five or six-membered aryl or heteroaryl ring or aryl-C₁₋₂ alkyl, optionally substituted with from one to three substituents selected from the group consisting of halogen, -OR^{5a}, -OC(O)R^{5a}, -NR^{5a}R^{5b}, -R^{5c}, -SR^{5a}, -CN, -NO₂, -CO₂R^{5a}, -CONR^{5a}R^{5b}, -C(O)R^{5a}, -NR^{5b}C(O)R^{5a}, -NR^{5b}C(O)₂R^{5c}, -X³OR^{5a}, -X³SR^{5a}, -X³CN, -X³NO₂, -X³CO₂R^{5a}, -X³CONR^{5a}R^{5b}, -X³C(O)R^{5a}, -X³OC(O)NR^{5a}R^{5b}, -X³NR^{5b}C(O)R^{5a}, -X³NR^{5b}C(O)₂R^{5c}, -X³NR^{5a}-C(O)NR^{5a}R^{5b}, -X³OC(O)R^{5a}, -X³NR^{5a}R^{5b}, -O-X³OR^{5a}, -O-X³NR^{5a}R^{5b}, -O-X³CO₂R^{5a}, -O-X³CONR^{5a}R^{5b}, -NR^{5b}-X³OR^{5a}, -NR^{5b}-X³NR^{5a}R^{5b}, -NR^{5b}-X³CO₂R^{5a}, and -NR^{5b}-X³CONR^{5a}R^{5b}, wherein each X³ is a member independently selected from the group consisting of C₁₋₄ alkylene, C₂₋₄ alkenylene and C₂₋₄ alkynylene; each R^{5a} and R^{5b} are independently selected from the group consisting of hydrogen, C₁₋₈ alkyl, C₁₋₈ haloalkyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocycloalkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl, heteroaryl, aryl-C₁₋₄ alkyl, and aryloxy-C₁₋₄ alkyl; each R^{5c} is independently selected from the group consisting of C₁₋₈ alkyl, C₁₋₈ haloalkyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocycloalkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl, heteroaryl, aryl-C₁₋₄ alkyl, and aryloxy-C₁₋₄ alkyl; and optionally any two of R^{5a}, R^{5b} and R^{5c} when part of a common R⁵ substituent can be combined with the atoms to which each is
- 25 attached to form a saturated four-, five-, six- or seven-membered ring having from 0 to 1 additional heteroatoms as ring members. Additionally, each of R^{5a}, R^{5b} and R^{5c} is optionally further substituted with from one to three members selected from the group consisting of -OH, -ORⁿ, -OC(O)NHRⁿ, -OC(O)N(Rⁿ)₂, -SH, -SRⁿ, -S(O)Rⁿ, -S(O)₂Rⁿ, -SO₂NH₂,
- 30

- S(O)₂NHRⁿ, -S(O)₂N(Rⁿ)₂, -NHS(O)₂Rⁿ, -NRⁿS(O)₂Rⁿ, -C(O)NH₂, -C(O)NHRⁿ,
 -C(O)N(Rⁿ)₂, -C(O)Rⁿ, -NHC(O)Rⁿ, -NRⁿC(O)Rⁿ, -NHC(O)NH₂, -NRⁿC(O)NH₂,
 -NRⁿC(O)NHRⁿ, -NHC(O)NHRⁿ, -NRⁿC(O)N(Rⁿ)₂, -NHC(O)N(Rⁿ)₂, -CO₂H, -CO₂Rⁿ,
 -NHCO₂Rⁿ, -NRⁿCO₂Rⁿ, -CN, -NO₂, -NH₂, -NHRⁿ, -N(Rⁿ)₂, -NRⁿS(O)NH₂ and
 5 -NRⁿS(O)₂NHRⁿ, wherein each Rⁿ is independently an unsubstituted C₁₋₆ alkyl or benzyl.


[0040] With the above formula are a number of specific embodiments of the invention. In one group of embodiments, W has a formula selected from the group consisting of (a), (d), (e) and (c):



In another group of embodiments, W has the formula (a), (d) or (e). In another group of embodiments, W has the formula (c). In a specific group of embodiments, at least one of n and m is 1 and each R⁴ and R^{4'} can be any of the substituents provided with reference to the full scope of the invention. In one group of embodiments, R^{4'} is selected from




[0041] In another group of embodiments, at least one of n and m is 1 and each R⁴ and R^{4'} is independently selected from the group consisting of H, halogen, OR^{4a}, C₁₋₄ alkyl, SR^{4a}, S(O)R^{4a}, S(O)₂R^{4a}, S(O)₂N(R^{4a}R^{4b}), NR^{4a}R^{4b}, C₁₋₄alkylOR^{4a}, C₁₋₄alkylNR^{4a}R^{4b}, C₁₋₄alkylCO₂R^{4a}, OC₁₋₄alkylOR^{4a}, OC₁₋₄alkylN(R^{4a}N^{4b}), N(R^{4a})C₁₋₄alkylOR^{4b}, N(R^{4a})C₁₋

$$_4\text{alkylN(R}^{4a}\text{R}^{4b}), \text{S(O)}_2\text{C}_{1-4}\text{alkylN(R}^{4a}\text{R}^{4b}),$$


; each R^{4a} is independently H or C₁₋₄alkyl; each R^{4d} is independently selected from the group consisting of H, OH, NH₂, N(CH₃)₂, CONH₂ and CON(CH₃)₂; and the wavy line indicates the

5 point of attachment to the rest of the molecule.

[0042] In another group of embodiments, at least one of n and m is 1 and each R^4 and $R^{4'}$ is selected from the group consisting of:




; and the wavy line indicates the point of attachment to the rest of the molecule.

[0043] In other embodiments, compounds of formula I are provided in which $R^{4'}$ is attached to the ring at a position para to A. In other embodiments, compounds of formula I are provided in which R^4 is attached to the ring at a position ortho to A. Within these

5 embodiments, specific embodiments are provided in which Y is an optionally substituted member selected from the group consisting of phenyl, benzyl, pyridyl, pyridylmethyl, pyrimidinyl, pyrazolyl, triazolyl and imidazolyl. Preferably, A represents benzene or a six-membered heterocycle containing 1-3 nitrogen atoms. One group of embodiments are those in which A is selected from the group consisting of: benzene, pyridine, pyridimidine, pyrazine, morpholin-3-one, piperazine-2-one, pyridine-2-one, piperidine, morpholine and piperazine.


10 Yet another group of embodiments are those in which A is selected from the group consisting of:



15 n is an integer from 0 to 3; each R^3 is independently selected from the group consisting of halogen, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, OR^{3a} , $N(R^{3a})_2$, $X^0CO_2R^{3a}$, $X^0CON(R^{3a})_2$, SO_2C_{1-4} , $SO_2N(R^{3a})_2$; each R^{3a} is independently selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl and C_{2-8} alkynyl; and X^0 is a bond or C_{1-8} alkylene; and the dashed line indicates the point of attachment to A and the wavy line indicates the point of attachment to the rest of


20 the molecule.

[0044] In one aspect, the present invention provides compounds having the formula:



and pharmaceutically acceptable salts thereof. In formula (I), R represents a member selected from the group consisting of H and C₁₋₄ alkyl; R¹ represents a member selected from the group consisting of halogen, C₁₋₈alkyl, C₂₋₈alkenyl and C₂₋₈alkynyl; R² represents a member selected from the group consisting of H, C₁₋₈ alkyl, -X-OR^{2a}, -X-SR^{2a}, -X-COR^{2a}, -X-CO₂R^{2a} and -X-N(R^{2a})₂; wherein X is C₁₋₈alkylene and each R^{2a} is independently selected from the group consisting of H and C₁₋₈alkyl, or optionally two R^{2a} groups attached to the same nitrogen atom are combined to form a 4-, 5-, 6- or 7-membered ring. The symbol R³ represents a member selected from the group consisting of H and halogen.

10 [0045] The letter W represent a moiety having a formula selected from (a), (b), and (c):



[0046] With reference to formula (b), the symbols Z¹, Z², and Z³ each independently represent N, C or CH with the proviso that two of Z¹, Z², and Z³ are other than N.


15 [0047] With reference to formulas (a) and (b), the subscript n is an integer of from 0 to 3 and the subscript m is an integer of from 0 to 1. The symbol R⁴ represents, in each occurrence, a member selected from the group consisting of halogen, -OR^{4a}, -OC(O)R^{4a}, -NR^{4a}R^{4b}, -SR^{4a}, S(O)R^{4a}, S(O)₂R^{4a}, S(O)₂NR^{4a}R^{4b}, -R^{4c}, -CN, -NO₂, -CO₂R^{4a}, -CONR^{4a}R^{4b}, -C(O)R^{4a}, -OC(O)NR^{4a}R^{4b}, -NR^{4b}C(O)R^{4a}, -NR^{4b}C(O)₂R^{4c}, -NR^{4a}-C(O)NR^{4a}R^{4b}, -X¹OR^{4a}, -X¹OC(O)R^{4a}, -X¹NR^{4a}R^{4b}, -X¹SR^{4a}, -X¹S(O)R^{4a}, -X¹S(O)₂R^{4a}, -X¹CN, -X¹NO₂, -X¹CO₂R^{4a}, -O-X¹CO₂R^{4a}, -X¹CONR^{4a}R^{4b}, -O-X¹CONR^{4a}R^{4b}, -X¹C(O)R^{4a}, -O-X¹NR^{4a}R^{4b}, -S(O)X¹NR^{4a}R^{4b}, -S(O)₂X¹NR^{4a}R^{4b}, -X¹OC(O)NR^{4a}R^{4b}, -X¹NR^{4a}C(O)R^{4a}, -X¹NR^{4b}C(O)₂R^{4c}, -X¹NR^{4a}C(O)NR^{4a}R^{4b}, -Y, -X¹-Y, -O-Y, -NR^{4a}Y, -SY, -S(O)Y and -S(O)₂Y. Similarly, R^{4'} represents a member selected from the group consisting of -C(=NH)R^{4a}, -C(=NR^{4c})R^{4a}, -C(=NH)NR^{4a}R^{4b}, -C(=NR^{4c})NR^{4a}R^{4b}, -C(=N⁺R^{4c}R^{4c})NR^{4a}R^{4b}, -X²-C(=NH)NR^{4a}R^{4b}, -X²-C(=NR^{4c})NR^{4a}R^{4b}, -X²-C(=N⁺R^{4c}R^{4c})NR^{4a}R^{4b} and -C(=NR^{4a})NR^{4a}-Y, wherein Y is a five or six-membered aryl, heterocyclyl or aryl-C₁₋₂alkyl, optionally substituted with from one to three substituents selected from the group consisting of halogen, oxo, -OR^{4a},

$-\text{OC(O)}\text{R}^{4a}$, $-\text{NR}^{4a}\text{R}^{4b}$, $-\text{R}^{4c}$, $-\text{SR}^{4a}$, $\text{S(O)}\text{R}^{4a}$, $\text{S(O)}_2\text{R}^{4a}$, $\text{S(O)}_2\text{NR}^{4a}\text{R}^{4b}$, $-\text{CN}$, $-\text{NO}_2$, $-\text{CO}_2\text{R}^{4a}$,
 $-\text{CONR}^{4a}\text{R}^{4b}$, $-\text{C(O)}\text{R}^{4a}$, $-\text{NR}^{4b}\text{C(O)}\text{R}^{4a}$, $-\text{NR}^{4b}\text{C(O)}_2\text{R}^{4c}$, $-\text{X}^1\text{R}^{4a}$, $-\text{X}^1\text{OR}^{4a}$, $-\text{X}^1\text{SR}^{4a}$, $-\text{X}^1\text{S(O)}\text{R}^{4a}$,
 $-\text{X}^1\text{S(O)}_2\text{R}^{4a}$, $-\text{X}^1\text{CN}$, $-\text{X}^1\text{NO}_2$, $-\text{X}^1\text{CO}_2\text{R}^{4a}$, $-\text{X}^1\text{CONR}^{4a}\text{R}^{4b}$, $-\text{X}^1\text{C(O)}\text{R}^{4a}$, $-\text{O}-\text{X}^1\text{NR}^{4a}\text{R}^{4b}$,
 $-\text{S(O)}\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{S(O)}_2\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{X}^1\text{OC(O)}\text{NR}^{4a}\text{R}^{4b}$, $-\text{X}^1\text{NR}^{4b}\text{C(O)}\text{R}^{4a}$, $-\text{X}^1\text{NR}^{4b}\text{C(O)}_2\text{R}^{4c}$,
 $-\text{X}^1\text{NR}^{4a}\text{C(O)}\text{NR}^{4a}\text{R}^{4b}$, $-\text{X}^1\text{OC(O)}\text{R}^{4a}$, $-\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{O}-\text{X}^1\text{OR}^{4a}$, $-\text{O}-\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{O}-\text{X}^1\text{CO}_2\text{R}^{4a}$,
 $-\text{O}-\text{X}^1\text{CONR}^{4a}\text{R}^{4b}$, $-\text{NR}^{4b}-\text{X}^1\text{OR}^{4a}$, $-\text{NR}^{4b}-\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{NR}^{4b}-\text{X}^1\text{CO}_2\text{R}^{4a}$, and $-\text{NR}^{4b}-\text{X}^1\text{CONR}^{4a}\text{R}^{4b}$; each X^1 and X^2 are members independently
 selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene; each R^{4a} and R^{4b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8}
 haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; each R^{4c} is independently selected from the group
 consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl, and optionally any two of
 R^{4a} , R^{4b} and R^{4c} when part of a common R^4 or $\text{R}^{4'}$ substituent can be combined with the atoms
 to which each is attached to form a saturated or unsaturated, four to nine-membered mono- or
 bicyclic ring having from 0 to 2 additional heteroatoms as ring members. Additionally, each
 of R^{4a} , R^{4b} and R^{4c} is optionally further substituted with from one to three members selected
 from the group consisting of $-\text{OH}$, oxo, $-\text{R}^m$, $-\text{OR}^m$, $-\text{OC(O)}\text{NHR}^m$, $-\text{OC(O)}\text{N(R}^m)_2$, $-\text{SH}$,
 $-\text{SR}^m$, $-\text{S(O)}\text{R}^m$, $-\text{S(O)}_2\text{R}^m$, $-\text{SO}_2\text{NH}_2$, $-\text{S(O)}_2\text{NHR}^m$, $-\text{S(O)}_2\text{N(R}^m)_2$, $-\text{NHS(O)}_2\text{R}^m$,
 $-\text{NR}^m\text{S(O)}_2\text{R}^m$, $-\text{C(O)}\text{NH}_2$, $-\text{C(O)}\text{NHR}^m$, $-\text{C(O)}\text{N(R}^m)_2$, $-\text{C(O)}\text{R}^m$, $-\text{NHC(O)}\text{R}^m$, $-\text{NR}^m\text{C(O)}\text{R}^m$,
 $-\text{NHC(O)}\text{NH}_2$, $-\text{NR}^m\text{C(O)}\text{NH}_2$, $-\text{NR}^m\text{C(O)}\text{NHR}^m$, $-\text{NHC(O)}\text{NHR}^m$, $-\text{NR}^m\text{C(O)}\text{N(R}^m)_2$,
 $-\text{NHC(O)}\text{N(R}^m)_2$, $-\text{CO}_2\text{H}$, $-\text{CO}_2\text{R}^m$, $-\text{NHCO}_2\text{R}^m$, $-\text{NR}^m\text{CO}_2\text{R}^m$, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{NHR}^m$,
 $-\text{N(R}^m)_2$, $-\text{NR}^m\text{S(O)}\text{NH}_2$ and $-\text{NR}^m\text{S(O)}_2\text{NHR}^m$, wherein each R^m is independently an
 unsubstituted C_{1-6} alkyl, benzyl or combined with the atoms to which each is attached to form
 a saturated four-, five-, six- or seven-membered ring having from 0 to 2 additional
 heteroatoms as ring members.

[0048] Turning next to formula (c), the symbol R^5 represents a member selected from the
 group consisting of $-\text{R}^{5a}$, $-\text{CO}_2\text{R}^{5c}$, $-\text{CONR}^{5a}\text{R}^{5b}$, $-\text{C(O)}\text{R}^{5a}$, $-\text{C(=NH)}\text{NR}^{5a}\text{R}^{5b}$, $-\text{C(=NR}^{5c})\text{NR}^{5a}\text{R}^{5b}$,
 $-\text{C(=N}^+\text{R}^{5c}\text{R}^{5c})\text{NR}^{5a}\text{R}^{5b}$, $-\text{C(=NR}^{5a})\text{R}^{5a}$, $-\text{C(=NR}^{5a})\text{Y}^1$, $-\text{X}^3\text{OR}^{5a}$,
 $-\text{X}^3\text{OC(O)}\text{R}^{5a}$, $-\text{X}^3\text{NR}^{5a}\text{R}^{5b}$, $-\text{X}^3\text{SR}^{5a}$, $-\text{X}^3\text{CN}$, $-\text{X}^3\text{NO}_2$, $-\text{X}^3\text{CO}_2\text{R}^{5a}$, $-\text{X}^3\text{CONR}^{5a}\text{R}^{5b}$,
 $-\text{X}^3\text{C(O)}\text{R}^{5a}$, $-\text{X}^3\text{OC(O)}\text{NR}^{5a}\text{R}^{5b}$, $-\text{X}^3\text{NR}^{5a}\text{C(O)}\text{R}^{5a}$, $-\text{X}^3\text{NR}^{5b}\text{C(O)}_2\text{R}^{5c}$, $-\text{X}^3\text{NR}^{5a}\text{C(O)}\text{NR}^{5a}\text{R}^{5b}$,
 $-\text{X}^3-\text{C(=NR}^{5a})\text{NR}^{5a}\text{R}^{5b}$, $-\text{Y}^1$, and $-\text{X}^3-\text{Y}^1$, wherein Y^1 is a five or six-membered aryl or
 heteroaryl ring or aryl- C_{1-2} alkyl, optionally substituted with from one to three substituents

selected from the group consisting of halogen, $-OR^{5a}$, $-OC(O)R^{5a}$, $-NR^{5a}R^{5b}$, $-R^{5c}$, $-SR^{5a}$, $-CN$, $-NO_2$, $-CO_2R^{5a}$, $-CONR^{5a}R^{5b}$, $-C(O)R^{5a}$, $-NR^{5b}C(O)R^{5a}$, $-NR^{5b}C(O)_2R^{5c}$, $-X^3OR^{5a}$, $-X^3SR^{5a}$, $-X^3CN$, $-X^3NO_2$, $-X^3CO_2R^{5a}$, $-X^3CONR^{5a}R^{5b}$, $-X^3C(O)R^{5a}$, $-X^3OC(O)NR^{5a}R^{5b}$, $-X^3NR^{5b}C(O)R^{5a}$, $-X^3NR^{5b}C(O)_2R^{5c}$, $-X^3NR^{5a}-C(O)NR^{5a}R^{5b}$, $-X^3OC(O)R^{5a}$, $-X^3NR^{5a}R^{5b}$, $-O-X^3OR^{5a}$, $-O-X^3NR^{5a}R^{5b}$, $-O-X^3CO_2R^{5a}$, $-O-X^3CONR^{5a}R^{5b}$, $-NR^{5b}-X^3OR^{5a}$, $-NR^{5b}-X^3NR^{5a}R^{5b}$, $-NR^{5b}-X^3CO_2R^{5a}$, and $-NR^{5b}-X^3CONR^{5a}R^{5b}$, wherein each X^3 is a member independently selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene; each R^{5a} and R^{5b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; each R^{5c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; and optionally any two of R^{5a} , R^{5b} and R^{5c} when part of a common R^5 substituent can be combined with the atoms to which each is attached to form a saturated four-, five-, six- or seven-membered ring having from 0 to 1 additional heteroatoms as ring members. Additionally, each of R^{5a} , R^{5b} and R^{5c} is optionally further substituted with from one to three members selected from the group consisting of $-OH$, $-OR^n$, $-OC(O)NHR^n$, $-OC(O)N(R^n)_2$, $-SH$, $-SR^n$, $-S(O)R^n$, $-S(O)_2R^n$, $-SO_2NH_2$, $-S(O)_2NHR^n$, $-S(O)_2N(R^n)_2$, $-NHS(O)_2R^n$, $-NR^nS(O)_2R^n$, $-C(O)NH_2$, $-C(O)NHR^n$, $-C(O)N(R^n)_2$, $-C(O)R^n$, $-NHC(O)R^n$, $-NR^nC(O)R^n$, $-NHC(O)NH_2$, $-NR^nC(O)NH_2$, $-NR^nC(O)NHR^n$, $-NHC(O)NHR^n$, $-NR^nC(O)N(R^n)_2$, $-NHC(O)N(R^n)_2$, $-CO_2H$, $-CO_2R^n$, $-NHCO_2R^n$, $-NR^nCO_2R^n$, $-CN$, $-NO_2$, $-NH_2$, $-NHR^n$, $-N(R^n)_2$, $-NR^nS(O)NH_2$ and $-NR^nS(O)_2NHR^n$, wherein each R^n is independently an unsubstituted C_{1-6} alkyl or benzyl.


[0049] With the above formulas are a number of other embodiments of the invention. In one group of embodiments, W has formula (a). In another group of embodiments, the subscript m is 1. In another group of embodiments $R^{4'}$ can be any of the substituents provided with reference to the full scope of the invention. In one group of embodiments, $R^{4'}$ is selected from



wherein the wavy line indicates the point of attachment to the remainder of the compound.

More preferably, $R^{4'}$ is attached to the phenyl ring at a position *para* to the attachment of the ring A.


- 5 [0050] In one aspect, the present invention provides compounds having the formula:



- and pharmaceutically acceptable salts thereof. In formula (I), A represents a five-membered heterocycle containing 1-4 nitrogen, oxygen or sulfur atoms; with 0-4 substituents selected from halogen, oxo, R^A , $-OR^A$, $N(R^A)_2$, $CO_2 R^A$, $CON(R^A)_2$, $S(O)_2 R^A$, $S(O)_2 N(R^A)_2$, $X^1 OR^A$, $X^1 CO_2 R^A$, $X^1 CON(R^A)_2$, $X^1 N(R^A)_2$, wherein each R^A is independently selected from H, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, or optionally two R^A attached to the same atom are combined to form a 3-, 4-, 5-, 6- or 7-membered ring. In one group of embodiments A is selected from the group consisting of: isoxazole, isothiazole, pyrazole, imidazole, oxazole, thiazole, isoxazoline, pyrazoline, imidazoline, 1,2,3-triazole, 1,2,4-triazole, 1,3,4-oxadiazole, 1,3,4-thiadiazole, 1,2,4-oxadiazole, 1,2,4-thiadiazole, pyrazol-5-one, pyrrolidine-2,5-dione, imidazolidine-2,4-dione, pyrrolidine, pyrrole, furan and thiophene. R represents a member

selected from the group consisting of H and C₁₋₄ alkyl; R¹ represents a member selected from the group consisting of halogen, C₁₋₈alkyl, C₂₋₈alkenyl and C₂₋₈alkynyl; R² represents a member selected from the group consisting of H, C₁₋₈ alkyl, -X-OR^{2a}, -X-SR^{2a}, -X-COR^{2a}, -X-CO₂R^{2a} and -X-N(R^{2a})₂; wherein X is C₁₋₈alkylene and each R^{2a} is independently selected from the group consisting of H and C₁₋₈alkyl, or optionally two R^{2a} groups attached to the same nitrogen atom are combined to form a 4-, 5-, 6- or 7-membered ring.

[0051] The letter W represent a moiety having a formula selected from (a), (b), and (c):



[0052] With reference to formula (b), the symbols Z¹, Z², and Z³ each independently represent N, C or CH with the proviso that two of Z¹, Z², and Z³ are other than N.

[0053] With reference to formulas (a) and (b), the subscript n is an integer of from 0 to 3 and the subscript m is an integer of from 0 to 1. The symbol R⁴ represents, in each occurrence, a member selected from the group consisting of halogen, -OR^{4a}, -OC(O)R^{4a}, -NR^{4a}R^{4b}, -SR^{4a}, S(O)R^{4a}, S(O)₂R^{4a}, S(O)₂NR^{4a}R^{4b}, -R^{4c}, -CN, -NO₂, -CO₂R^{4a}, -CONR^{4a}R^{4b}, -C(O)R^{4a}, -OC(O)NR^{4a}R^{4b}, -NR^{4b}C(O)R^{4a}, -NR^{4b}C(O)₂R^{4c}, -NR^{4a}-C(O)NR^{4a}R^{4b}, -X¹OR^{4a}, -X¹OC(O)R^{4a}, -X¹NR^{4a}R^{4b}, -X¹SR^{4a}, -X¹S(O)R^{4a}, -X¹S(O)₂R^{4a}, -X¹CN, -X¹NO₂, -X¹CO₂R^{4a}, -O-X¹CO₂R^{4a}, -X¹CONR^{4a}R^{4b}, -O-X¹CONR^{4a}R^{4b}, -X¹C(O)R^{4a}, -O-X¹NR^{4a}R^{4b}, -S(O)X¹NR^{4a}R^{4b}, -S(O)₂X¹NR^{4a}R^{4b}, -X¹OC(O)NR^{4a}R^{4b}, -X¹NR^{4a}C(O)R^{4a}, -X¹NR^{4b}C(O)₂R^{4c}, -X¹NR^{4a}C(O)NR^{4a}R^{4b}, -Y, -X¹-Y, -O-Y, -NR^{4a}Y, -SY, -S(O)Y and -S(O)₂Y. Similarly, R^{4'} represents a member selected from the group consisting of -C(=NH)R^{4a}, -C(=NR^{4c})R^{4a}, -C(=NH)NR^{4a}R^{4b}, -C(=NR^{4c})NR^{4a}R^{4b}, -C(=N⁺R^{4c}R^{4c})NR^{4a}R^{4b}, -X²-C(=NH)NR^{4a}R^{4b}, -X²-C(=NR^{4c})NR^{4a}R^{4b}, -X²-C(=N⁺R^{4c}R^{4c})NR^{4a}R^{4b} and -C(=NR^{4a})NR^{4a}-Y, wherein Y is a five or six-membered aryl, heterocyclyl or aryl-C₁₋₂alkyl, optionally substituted with from one to three substituents selected from the group consisting of halogen, oxo, -OR^{4a}, -OC(O)R^{4a}, -NR^{4a}R^{4b}, -R^{4c}, -SR^{4a}, S(O)R^{4a}, S(O)₂R^{4a}, S(O)₂NR^{4a}R^{4b}, -CN, -NO₂, -CO₂R^{4a}, -CONR^{4a}R^{4b}, -C(O)R^{4a}, -NR^{4b}C(O)R^{4a}, -NR^{4b}C(O)₂R^{4c}, -X¹R^{4a}, -X¹OR^{4a}, -X¹SR^{4a}, -X¹S(O)R^{4a}, -X¹S(O)₂R^{4a}, -X¹CN, -X¹NO₂, -X¹CO₂R^{4a}, -X¹CONR^{4a}R^{4b}, -X¹C(O)R^{4a}, -O-X¹NR^{4a}R^{4b}, -S(O)X¹NR^{4a}R^{4b}, -S(O)₂X¹NR^{4a}R^{4b}, -X¹OC(O)NR^{4a}R^{4b}, -X¹NR^{4b}C(O)R^{4a}, -X¹NR^{4b}C(O)₂R^{4c}, -X¹NR^{4a}-C(O)NR^{4a}R^{4b}, -X¹OC(O)R^{4a}, -X¹NR^{4a}R^{4b}, -O-X¹OR^{4a},


$-O-X^1NR^{4a}R^{4b}$, $-O-X^1CO_2R^{4a}$, $-O-X^1CONR^{4a}R^{4b}$, $-NR^{4b}-X^1OR^{4a}$, $-NR^{4b}-X^1NR^{4a}R^{4b}$,
 $-NR^{4b}-X^1CO_2R^{4a}$, and $-NR^{4b}-X^1CONR^{4a}R^{4b}$; each X^1 and X^2 are members independently
 selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene; each
 R^{4a} and R^{4b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8}
 5 haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl,
 aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; each R^{4c} is independently selected from the group
 consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8}
 alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl, and optionally any two of
 R^{4a} , R^{4b} and R^{4c} when part of a common R^4 or R^4' substituent can be combined with the atoms
 10 to which each is attached to form a saturated or unsaturated, four to nine-membered mono- or
 bicyclic ring having from 0 to 2 additional heteroatoms as ring members. Additionally, each
 of R^{4a} , R^{4b} and R^{4c} is optionally further substituted with from one to three members selected
 from the group consisting of -OH, oxo, $-R^m$, $-OR^m$, $-OC(O)NHR^m$, $-OC(O)N(R^m)_2$, -SH,
 $-SR^m$, $-S(O)R^m$, $-S(O)_2R^m$, $-SO_2NH_2$, $-S(O)_2NHR^m$, $-S(O)_2N(R^m)_2$, $-NHS(O)_2R^m$,
 15 $-NR^mS(O)_2R^m$, $-C(O)NH_2$, $-C(O)NHR^m$, $-C(O)N(R^m)_2$, $-C(O)R^m$, $-NHC(O)R^m$, $-NR^mC(O)R^m$,
 $-NHC(O)NH_2$, $-NR^mC(O)NH_2$, $-NR^mC(O)NHR^m$, $-NHC(O)NHR^m$, $-NR^mC(O)N(R^m)_2$,
 $-NHC(O)N(R^m)_2$, $-CO_2H$, $-CO_2R^m$, $-NHCO_2R^m$, $-NR^mCO_2R^m$, -CN, $-NO_2$, $-NH_2$, $-NHR^m$,
 $-N(R^m)_2$, $-NR^mS(O)NH_2$ and $-NR^mS(O)_2NHR^m$, wherein each R^m is independently an
 unsubstituted C_{1-6} alkyl, benzyl or combined with the atoms to which each is attached to form
 20 a saturated four-, five-, six- or seven-membered ring having from 0 to 2 additional
 heteroatoms as ring members.

[0054] Turning next to formula (c), the symbol R^5 represents a member selected from the
 group consisting of $-R^{5a}$, $-CO_2R^{5c}$, $-CONR^{5a}R^{5b}$, $-C(O)R^{5a}$, $-C(=NH)NR^{5a}R^{5b}$, -
 $C(=NR^{5c})NR^{5a}R^{5b}$, $-C(=N^+R^{5c}R^{5c})NR^{5a}R^{5b}$, $-C(=NR^{5a})R^{5a}$, $-C(=NR^{5a})Y^1$, $-X^3OR^{5a}$,
 25 $-X^3OC(O)R^{5a}$, $-X^3NR^{5a}R^{5b}$, $-X^3SR^{5a}$, $-X^3CN$, $-X^3NO_2$, $-X^3CO_2R^{5a}$, $-X^3CONR^{5a}R^{5b}$,
 $-X^3C(O)R^{5a}$, $-X^3OC(O)NR^{5a}R^{5b}$, $-X^3NR^{5a}C(O)R^{5a}$, $-X^3NR^{5b}C(O)_2R^{5c}$, $-X^3NR^{5a}C(O)NR^{5a}R^{5b}$,
 $-X^3-C(=NR^{5a})NR^{5a}R^{5b}$, $-Y^1$, and $-X^3-Y^1$, wherein Y^1 is a five or six-membered aryl or
 heteroaryl ring or aryl- C_{1-2} alkyl, optionally substituted with from one to three substituents
 selected from the group consisting of halogen, $-OR^{5a}$, $-OC(O)R^{5a}$, $-NR^{5a}R^{5b}$, $-R^{5c}$, $-SR^{5a}$, -CN,
 30 $-NO_2$, $-CO_2R^{5a}$, $-CONR^{5a}R^{5b}$, $-C(O)R^{5a}$, $-NR^{5b}C(O)R^{5a}$, $-NR^{5b}C(O)_2R^{5c}$, $-X^3OR^{5a}$, $-X^3SR^{5a}$,
 $-X^3CN$, $-X^3NO_2$, $-X^3CO_2R^{5a}$, $-X^3CONR^{5a}R^{5b}$, $-X^3C(O)R^{5a}$, $-X^3OC(O)NR^{5a}R^{5b}$,
 $-X^3NR^{5b}C(O)R^{5a}$, $-X^3NR^{5b}C(O)_2R^{5c}$, $-X^3NR^{5a}-C(O)NR^{5a}R^{5b}$, $-X^3OC(O)R^{5a}$, $-X^3NR^{5a}R^{5b}$,
 $-O-X^3OR^{5a}$, $-O-X^3NR^{5a}R^{5b}$, $-O-X^3CO_2R^{5a}$, $-O-X^3CONR^{5a}R^{5b}$, $-NR^{5b}-X^3OR^{5a}$,

$-\text{NR}^{\text{5a}}-\text{X}^3\text{NR}^{\text{5b}}\text{R}^{\text{5c}}$, $-\text{NR}^{\text{5b}}-\text{X}^3\text{CO}_2\text{R}^{\text{5a}}$, and $-\text{NR}^{\text{5b}}-\text{X}^3\text{CONR}^{\text{5a}}\text{R}^{\text{5c}}$, wherein each X^3 is a member independently selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene; each R^{5a} and R^{5b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; each R^{5c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; and optionally any two of R^{5a} , R^{5b} and R^{5c} when part of a common R^{5} substituent can be combined with the atoms to which each is attached to form a saturated four-, five-, six- or seven-membered ring having from 0 to 1 additional heteroatoms as ring members.


Additionally, each of R^{5a} , R^{5b} and R^{5c} is optionally further substituted with from one to three members selected from the group consisting of $-\text{OH}$, $-\text{OR}^{\text{n}}$, $-\text{OC}(\text{O})\text{NHR}^{\text{n}}$, $-\text{OC}(\text{O})\text{N}(\text{R}^{\text{n}})_2$, $-\text{SH}$, $-\text{SR}^{\text{n}}$, $-\text{S}(\text{O})\text{R}^{\text{n}}$, $-\text{S}(\text{O})_2\text{R}^{\text{n}}$, $-\text{SO}_2\text{NH}_2$, $-\text{S}(\text{O})_2\text{NHR}^{\text{n}}$, $-\text{S}(\text{O})_2\text{N}(\text{R}^{\text{n}})_2$, $-\text{NHS}(\text{O})_2\text{R}^{\text{n}}$, $-\text{NR}^{\text{n}}\text{S}(\text{O})_2\text{R}^{\text{n}}$, $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NHR}^{\text{n}}$, $-\text{C}(\text{O})\text{N}(\text{R}^{\text{n}})_2$, $-\text{C}(\text{O})\text{R}^{\text{n}}$, $-\text{NHC}(\text{O})\text{R}^{\text{n}}$, $-\text{NR}^{\text{n}}\text{C}(\text{O})\text{R}^{\text{n}}$, $-\text{NHC}(\text{O})\text{NH}_2$, $-\text{NR}^{\text{n}}\text{C}(\text{O})\text{NH}_2$, $-\text{NR}^{\text{n}}\text{C}(\text{O})\text{NHR}^{\text{n}}$, $-\text{NHC}(\text{O})\text{NHR}^{\text{n}}$, $-\text{NR}^{\text{n}}\text{C}(\text{O})\text{N}(\text{R}^{\text{n}})_2$, $-\text{NHC}(\text{O})\text{N}(\text{R}^{\text{n}})_2$, $-\text{CO}_2\text{H}$, $-\text{CO}_2\text{R}^{\text{n}}$, $-\text{NHCO}_2\text{R}^{\text{n}}$, $-\text{NR}^{\text{n}}\text{CO}_2\text{R}^{\text{n}}$, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{NHR}^{\text{n}}$, $-\text{N}(\text{R}^{\text{n}})_2$, $-\text{NR}^{\text{n}}\text{S}(\text{O})\text{NH}_2$ and $-\text{NR}^{\text{n}}\text{S}(\text{O})_2\text{NHR}^{\text{n}}$, wherein each R^{n} is independently an unsubstituted C_{1-6} alkyl or benzyl.

[0055] With the above formulas are a number of other embodiments of the invention. In one group of embodiments, A is selected from the group consisting of:



; each R^3 is independently selected from the group consisting of halogen, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, OR^{3a} , $N(R^{3a})_2$, $X^0CO_2R^{3a}$, $X^0CON(R^{3a})_2$, SO_2C_{1-4} , $SO_2N(R^{3a})_2$; each R^{3a} is independently selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl and C_{2-8} alkynyl; and X^0 is a bond or C_{1-8} alkylene; and the dashed line indicates the point of attachment to A and the wavy line indicates the point of attachment to the rest of the molecule. In another group of embodiments, W has formula (a). In another group of embodiments, the subscript m is 1. In another group of embodiments R^4 can be any of the


substituents provided with reference to the full scope of the invention. In one group of embodiments, $R^{4'}$ is selected from



wherein the wavy line indicates the point of attachment to the remainder of the compound.

- 5 More preferably, $R^{4'}$ is attached to the phenyl ring at a position *para* to the attachment of the ring A. In another group of embodiments, the subscript n is an integer of from 0 to 1. In another group of embodiments n is 1; $R^{4'}$ is attached to the phenyl ring at a position *ortho* to A and $R^{4'}$ is attached to the phenyl ring at a position *para* to A. In one group of embodiments, $R^{4'}$ is selected from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-SR^{4a}$,
 10 $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$ and $-SY$.

[0056] In another group of embodiments, m is 0 or 1, n is 0, 1 or 2 and at least one $R^{4'}$ is selected from the group consisting of



wherein the wavy line indicates the point of attachment to the remainder of the compound.

[0057] In another group of embodiments, m is 0 or 1; n is 0, 1 or 2; and at least one R^4 is selected from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2Y$,

wherein Y is an optionally substituted member selected from the group consisting of phenyl, benzyl, pyridyl, pyridylmethyl, pyrimidinyl and pyrazolyl. More preferably, R^4 is selected from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$ and $-NR^{4a}Y$, wherein Y is an optionally substituted member selected from the group consisting of phenyl, benzyl, pyridyl and pyridylmethyl. Still further preferred are those embodiments in which the optional


substituents are selected from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $-CO_2R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-X^1OR^{4a}$ and $-X^1NR^{4a}R^{4b}$.

[0058] In other embodiments, compounds of formula I are provided in which R^2 is present the carbon attached to R^2 has the R- configuration. Other embodiments are provided in which R^3 is other than H the carbon attached to R^3 has the R- configuration.

[0059] In other embodiments, compounds of formula I are provided in which W has the formula (b). Within this group, specific embodiments are provided in which Z^1 is N. Other

embodiments are provided in which Z^2 is N. Other embodiments are provided in which Z^3 is N.

[0060] In another group of embodiments, the subscript m is 1. In another group of embodiments $R^{4'}$ can be any of the substituents provided with reference to the full scope of the invention. In one group of embodiments, $R^{4'}$ is selected from




wherein the wavy line indicates the point of attachment to the remainder of the compound.

More preferably, $R^{4'}$ is attached to the pyridyl ring at a position *para* to the attachment of the

ring A. In another group of embodiments, the subscript n is an integer of from 0 to 1. In

10 another group of embodiments n is 1; R^4 is attached to the pyridyl ring at a position *ortho* to A and $R^{4'}$ is attached to the pyridyl ring at a position *para* to A. In one group of embodiments, R^4 is selected from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-SR^{4a}$, $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$ and $-SY$.

[0061] In another group of embodiments, m is 0 or 1, n is 0, 1 or 2 and at least one R^4 is
15 selected from the group consisting of



wherein the wavy line indicates the point of attachment to the remainder of the compound.


[0062] In another group of embodiments, m is 0 or 1; n is 0, 1 or 2; and at least one R^4 is selected from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2Y$,

wherein Y is an optionally substituted member selected from the group consisting of phenyl, benzyl, pyridyl, pyridylmethyl, pyrimidinyl and pyrazolyl. More preferably, R^4 is selected from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$ and $-NR^{4a}Y$, wherein Y is an optionally substituted member selected from the group consisting of phenyl, benzyl, pyridyl and pyridylmethyl. Still further preferred are those embodiments in which the optional

substituents are selected from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $-CO_2R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-X^1OR^{4a}$ and $-X^1NR^{4a}R^{4b}$.


[0063] In other embodiments, compounds of formula I are provided in which W has the formula (c). Other embodiments are provided in which Y^1 is selected from the group consisting of optionally substituted phenyl and optionally substituted pyridyl. Other embodiments are provided in which R^5 is selected from the group consisting of $-R^{5a}$, $-C(=NH)NR^{5a}R^{5b}$, $-C(=NH)R^{5a}$, $-C(=NH)Y^1$ and $-X^3-Y^1$.

[0064] In one aspect, the present invention provides compounds having the formula:



and pharmaceutically acceptable salts thereof. In formula (I), R represents a member selected from the group consisting of H and C₁₋₄ alkyl; R¹ represents a member selected from the group consisting of halogen, C₁₋₈alkyl, C₂₋₈alkenyl and C₂₋₈alkynyl; R² represents a member selected from the group consisting of H, C₁₋₈ alkyl, -X-OR^{2a}, -X-SR^{2a}, -X-COR^{2a}, -X-CO₂R^{2a} and -X-N(R^{2a})₂; wherein X is C₁₋₈alkylene and each R^{2a} is independently selected from the group consisting of H and C₁₋₈alkyl, or optionally two R^{2a} groups attached to the same nitrogen atom are combined to form a 4-, 5-, 6- or 7-membered ring. Each dashed line in formula (I) independently represents an optional bond, with the understanding that only one optional bond is present. The symbol R³ represents a member selected from the group consisting of H and C₁₋₄ alkyl; or is absent when a carbon-carbon double bond is present in the ring to which R³ would otherwise be attached. For the embodiments in which R² is present or R³ is present and other than H, a preferred group of embodiments are those in which the carbon to which R² or R³ is attached has the (R)- stereochemical configuration.

[0065] The letter W represent a moiety having a formula selected from (a), (b), and (c):



[0066] With reference to formula (b), the symbols Z¹, Z², and Z³ each independently represent N, C or CH with the proviso that two of Z¹, Z², and Z³ are other than N. The pyridyl nitrogen can be in a position *para* to the attachment of the isoxazole or isoxazoline ring (Z¹ (4-pyridyl)), meta (Z² (3-pyridyl)) or ortho (Z³ (2-pyridyl)).


[0067] With reference to formulas (a) and (b), the subscript n is an integer of from 0 to 3 and the subscript m is an integer of from 0 to 1. The symbol R⁴ represents, in each

occurrence, a member selected from the group consisting of halogen, -OR^{4a}, -OC(O)R^{4a}, -NR^{4a}R^{4b}, -SR^{4a}, S(O)R^{4a}, S(O)₂R^{4a}, S(O)₂NR^{4a}R^{4b}, -R^{4c}, -CN, -NO₂, -CO₂R^{4a}, -CONR^{4a}R^{4b}, -C(O)R^{4a}, -OC(O)NR^{4a}R^{4b}, -NR^{4b}C(O)R^{4a}, -NR^{4b}C(O)₂R^{4c}, -NR^{4a}-C(O)NR^{4a}R^{4b}, -X¹OR^{4a}, -X¹OC(O)R^{4a}, -X¹NR^{4a}R^{4b}, -X¹SR^{4a}, -X¹S(O)R^{4a}, -X¹S(O)₂R^{4a}, -X¹CN, -X¹NO₂, -X¹CO₂R^{4a}, -O-X¹CO₂R^{4a}, -X¹CONR^{4a}R^{4b}, -O-X¹CONR^{4a}R^{4b}, -X¹C(O)R^{4a}, -O-X¹NR^{4a}R^{4b}, -

- $S(O)X^1NR^{4a}R^{4b}$, $-S(O)_2X^1NR^{4a}R^{4b}$, $-X^1OC(O)NR^{4a}R^{4b}$, $-X^1NR^{4a}C(O)R^{4a}$, $-X^1NR^{4b}C(O)_2R^{4c}$,
 $-X^1NR^{4a}C(O)NR^{4a}R^{4b}$, $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2Y$. Similarly, $R^{4'}$
represents a member selected from the group consisting of $-C(=NH)R^{4a}$, $-C(=NR^{4c})R^{4a}$,
 $-C(=NH)NR^{4a}R^{4b}$, $-C(=NR^{4c})NR^{4a}R^{4b}$, $-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$, $-X^2-C(=NH)NR^{4a}R^{4b}$,
5 $-X^2-C(=NR^{4c})NR^{4a}R^{4b}$, $-X^2-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$ and $-C(=NR^{4a})NR^{4a}-Y$, wherein Y is a
five or six-membered aryl, heterocyclyl or aryl- C_{1-2} alkyl, optionally substituted with from
one to three substituents selected from the group consisting of halogen, oxo, $-OR^{4a}$,
 $-OC(O)R^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2NR^{4a}R^{4b}$, $-CN$, $-NO_2$, $-CO_2R^{4a}$,
 $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)_2R^{4c}$, $-X^1R^{4a}$, $-X^1OR^{4a}$, $-X^1SR^{4a}$, -
10 $X^1S(O)R^{4a}$, $-X^1S(O)_2R^{4a}$, $-X^1CN$, $-X^1NO_2$, $-X^1CO_2R^{4a}$, $-X^1CONR^{4a}R^{4b}$, $-X^1C(O)R^{4a}$, $-O-$
 $X^1NR^{4a}R^{4b}$, $-S(O)X^1NR^{4a}R^{4b}$, $-S(O)_2X^1NR^{4a}R^{4b}$, $-X^1OC(O)NR^{4a}R^{4b}$, $-X^1NR^{4b}C(O)R^{4a}$, -
 $X^1NR^{4b}C(O)_2R^{4c}$, $-X^1NR^{4a}-C(O)NR^{4a}R^{4b}$, $-X^1OC(O)R^{4a}$, $-X^1NR^{4a}R^{4b}$, $-O-X^1OR^{4a}$,
 $-O-X^1NR^{4a}R^{4b}$, $-O-X^1CO_2R^{4a}$, $-O-X^1CONR^{4a}R^{4b}$, $-NR^{4b}-X^1OR^{4a}$, $-NR^{4b}-X^1NR^{4a}R^{4b}$,
 $-NR^{4b}-X^1CO_2R^{4a}$, and $-NR^{4b}-X^1CONR^{4a}R^{4b}$; each X^1 and X^2 are members independently
15 selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene; each
 R^{4a} and R^{4b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8}
haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl,
aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; each R^{4c} is independently selected from the group
consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-}
20 8 alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl, and optionally any two of
 R^{4a} , R^{4b} and R^{4c} when part of a common R^4 or $R^{4'}$ substituent can be combined with the atoms
to which each is attached to form a saturated or unsaturated, four to nine-membered mono- or
bicyclic ring having from 0 to 2 additional heteroatoms as ring members. Additionally, each
of R^{4a} , R^{4b} and R^{4c} is optionally further substituted with from one to three members selected
25 from the group consisting of $-OH$, oxo, $-R^m$, $-OR^m$, $-OC(O)NHR^m$, $-OC(O)N(R^m)_2$, $-SH$,
 $-SR^m$, $-S(O)R^m$, $-S(O)_2R^m$, $-SO_2NH_2$, $-S(O)_2NHR^m$, $-S(O)_2N(R^m)_2$, $-NHS(O)_2R^m$,
 $-NR^mS(O)_2R^m$, $-C(O)NH_2$, $-C(O)NHR^m$, $-C(O)N(R^m)_2$, $-C(O)R^m$, $-NHC(O)R^m$, $-NR^mC(O)R^m$,
 $-NHC(O)NH_2$, $-NR^mC(O)NH_2$, $-NR^mC(O)NHR^m$, $-NHC(O)NHR^m$, $-NR^mC(O)N(R^m)_2$,
 $-NHC(O)N(R^m)_2$, $-CO_2H$, $-CO_2R^m$, $-NHCO_2R^m$, $-NR^mCO_2R^m$, $-CN$, $-NO_2$, $-NH_2$, $-NHR^m$,
30 $-N(R^m)_2$, $-NR^mS(O)NH_2$ and $-NR^mS(O)_2NHR^m$, wherein each R^m is independently an
unsubstituted C_{1-6} alkyl, benzyl or combined with the atoms to which each is attached to form
a saturated four-, five-, six- or seven-membered ring having from 0 to 2 additional
heteroatoms as ring members.

- [0068] Turning next to formula (c), the symbol R^5 represents a member selected from the group consisting of $-R^{5a}$, $-\text{CO}_2R^{5c}$, $-\text{CONR}^{5a}R^{5b}$, $-\text{C}(\text{O})R^{5a}$, $\text{C}(=\text{NR}^{5a})\text{NR}^{5a}R^{5b}$, $-\text{C}(=\text{N}^+R^{5c}R^{5c})\text{NR}^{5a}R^{5b}$, $-\text{C}(=\text{NR}^{5a})R^{5a}$, $-\text{C}(=\text{NR}^{5a})Y^1$, $-\text{X}^3\text{OR}^{5a}$, $-\text{X}^3\text{OC}(\text{O})R^{5a}$, $-\text{X}^3\text{NR}^{5a}R^{5b}$, $-\text{X}^3\text{SR}^{5a}$, $-\text{X}^3\text{CN}$, $-\text{X}^3\text{NO}_2$, $-\text{X}^3\text{CO}_2R^{5a}$, $-\text{X}^3\text{CONR}^{5a}R^{5b}$, $-\text{X}^3\text{C}(\text{O})R^{5a}$, $-\text{X}^3\text{OC}(\text{O})\text{NR}^{5a}R^{5b}$, $-\text{X}^3\text{NR}^{5a}\text{C}(\text{O})R^{5a}$, $-\text{X}^3\text{NR}^{5b}\text{C}(\text{O})_2R^{5c}$, $-\text{X}^3\text{NR}^{5a}\text{C}(\text{O})\text{NR}^{5a}R^{5b}$, $-\text{X}^3-\text{C}(=\text{NR}^{5a})\text{NR}^{5a}R^{5b}$, $-Y^1$, and $-\text{X}^3-Y^1$, wherein Y^1 is a five or six-membered aryl or heteroaryl ring or aryl- C_{1-2} alkyl, optionally substituted with from one to three substituents selected from the group consisting of halogen, $-\text{OR}^{5a}$, $-\text{OC}(\text{O})R^{5a}$, $-\text{NR}^{5a}R^{5b}$, $-R^{5c}$, $-\text{SR}^{5a}$, $-\text{CN}$, $-\text{NO}_2$, $-\text{CO}_2R^{5a}$, $-\text{CONR}^{5a}R^{5b}$, $-\text{C}(\text{O})R^{5a}$, $-\text{NR}^{5b}\text{C}(\text{O})R^{5a}$, $-\text{NR}^{5b}\text{C}(\text{O})_2R^{5c}$, $-\text{X}^3\text{OR}^{5a}$, $-\text{X}^3\text{SR}^{5a}$, $-\text{X}^3\text{CN}$, $-\text{X}^3\text{NO}_2$, $-\text{X}^3\text{CO}_2R^{5a}$, $-\text{X}^3\text{CONR}^{5a}R^{5b}$, $-\text{X}^3\text{C}(\text{O})R^{5a}$, $-\text{X}^3\text{OC}(\text{O})\text{NR}^{5a}R^{5b}$, $-\text{X}^3\text{NR}^{5b}\text{C}(\text{O})R^{5a}$, $-\text{X}^3\text{NR}^{5b}\text{C}(\text{O})_2R^{5c}$, $-\text{X}^3\text{NR}^{5a}-\text{C}(\text{O})\text{NR}^{5a}R^{5b}$, $-\text{X}^3\text{OC}(\text{O})R^{5a}$, $-\text{X}^3\text{NR}^{5a}R^{5b}$, $-\text{O}-\text{X}^3\text{OR}^{5a}$, $-\text{O}-\text{X}^3\text{NR}^{5a}R^{5b}$, $-\text{O}-\text{X}^3\text{CO}_2R^{5a}$, $-\text{O}-\text{X}^3\text{CONR}^{5a}R^{5b}$, $-\text{NR}^{5b}-\text{X}^3\text{OR}^{5a}$, $-\text{NR}^{5b}-\text{X}^3\text{NR}^{5a}R^{5b}$, $-\text{NR}^{5b}-\text{X}^3\text{CO}_2R^{5a}$, and $-\text{NR}^{5b}-\text{X}^3\text{CONR}^{5a}R^{5b}$, wherein each X^3 is a member independently selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene; each R^{5a} and R^{5b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; each R^{5c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl; and optionally any two of R^{5a} , R^{5b} and R^{5c} when part of a common R^5 substituent can be combined with the atoms to which each is attached to form a saturated four-, five-, six- or seven-membered ring having from 0 to 1 additional heteroatoms as ring members. Additionally, each of R^{5a} , R^{5b} and R^{5c} is optionally further substituted with from one to three members selected from the group consisting of $-\text{OH}$, $-\text{OR}^n$, $-\text{OC}(\text{O})\text{NHR}^n$, $-\text{OC}(\text{O})\text{N}(\text{R}^n)_2$, $-\text{SH}$, $-\text{SR}^n$, $-\text{S}(\text{O})\text{R}^n$, $-\text{S}(\text{O})_2\text{R}^n$, $-\text{SO}_2\text{NH}_2$, $-\text{S}(\text{O})_2\text{NHR}^n$, $-\text{S}(\text{O})_2\text{N}(\text{R}^n)_2$, $-\text{NHS}(\text{O})_2\text{R}^n$, $-\text{NR}^n\text{S}(\text{O})_2\text{R}^n$, $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NHR}^n$, $-\text{C}(\text{O})\text{N}(\text{R}^n)_2$, $-\text{C}(\text{O})\text{R}^n$, $-\text{NHC}(\text{O})\text{R}^n$, $-\text{NR}^n\text{C}(\text{O})\text{R}^n$, $-\text{NHC}(\text{O})\text{NH}_2$, $-\text{NR}^n\text{C}(\text{O})\text{NH}_2$, $-\text{NR}^n\text{C}(\text{O})\text{NHR}^n$, $-\text{NHC}(\text{O})\text{NHR}^n$, $-\text{NR}^n\text{C}(\text{O})\text{N}(\text{R}^n)_2$, $-\text{NHC}(\text{O})\text{N}(\text{R}^n)_2$, $-\text{CO}_2\text{H}$, $-\text{CO}_2\text{R}^n$, $-\text{NHCO}_2\text{R}^n$, $-\text{NR}^n\text{CO}_2\text{R}^n$, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{NHR}^n$, $-\text{N}(\text{R}^n)_2$, $-\text{NR}^n\text{S}(\text{O})\text{NH}_2$ and $-\text{NR}^n\text{S}(\text{O})_2\text{NHR}^n$, wherein each R^n is independently an unsubstituted C_{1-6} alkyl or benzyl.
- [0069] With the above formula are a number of specific embodiments of the invention. In one group of embodiments, W has formula (a). In another group of embodiments, the subscript m is 1. In another group of embodiments $\text{R}^{4'}$ can be any of the substituents


provided with reference to the full scope of the invention. In one group of embodiments, R^4 is selected from



wherein the wavy line indicates the point of attachment to the remainder of the compound.

- 5 More preferably, R^4 is attached to the phenyl ring at a position *para* to the attachment of the isoxazole or isoxazoline ring. In another group of embodiments, the subscript n is an integer of from 0 to 1. In another group of embodiments n is 1; R^4 is attached to the phenyl ring at a position *ortho* to the isoxazole or isoxazoline ring and R^4 is attached to the phenyl ring at a position *para* to the isoxazole or isoxazoline ring. In one group of embodiments, R^4
- 10 is selected from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-SR^{4a}$, $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$ and $-SY$.

[0070] In another group of embodiments, m is 0 or 1, n is 0, 1 or 2 and at least one R^4 is selected from the group consisting of



wherein the wavy line indicates the point of attachment to the remainder of the compound.

[0071] In another group of embodiments, m is 0 or 1; n is 0, 1, or 2; and at least one R^4 is selected from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2Y$,

wherein Y is an optionally substituted member selected from the group consisting of phenyl, benzyl, pyridyl, pyridylmethyl, pyrimidinyl and pyrazolyl. More preferably, R^4 is selected from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$ and $-NR^{4a}Y$, wherein Y is an optionally substituted member selected from the group consisting of phenyl, benzyl, pyridyl and pyridylmethyl. Still further preferred are those embodiments in which the optional


substituents are selected from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $-CO_2R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-X^1OR^{4a}$ and $-X^1NR^{4a}R^{4b}$.

[0072] In other embodiments, compounds of formula I are provided in which R^2 is present the carbon attached to R^2 has the R- configuration. Other embodiments are provided in which R^3 is other than H the carbon attached to R^3 has the R- configuration.

[0073] In other embodiments, compounds of formula I are provided in which W has the formula (b). Within this group, specific embodiments are provided in which Z^1 is N. Other

embodiments are provided in which Z^2 is N. Other embodiments are provided in which Z^3 is N.


[0074] In another group of embodiments, the subscript m is 1. In another group of embodiments $R^{4'}$ can be any of the substituents provided with reference to the full scope of the invention. In one group of embodiments, $R^{4'}$ is selected from



wherein the wavy line indicates the point of attachment to the remainder of the compound.

More preferably, $R^{4'}$ is attached to the pyridyl ring at a position *para* to the attachment of the isoxazole or isoxazoline ring. In another group of embodiments, the subscript n is an integer of from 0 to 1. In another group of embodiments n is 1; R^4 is attached to the pyridyl ring at a position *ortho* to the isoxazole or isoxazoline ring and $R^{4'}$ is attached to the pyridyl ring at a position *para* to the isoxazole or isoxazoline ring. In one group of embodiments, R^4 is selected from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-SR^{4a}$, $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$ and $-SY$.

[0075] In another group of embodiments, m is 0 or 1, n is 0, 1 or 2 and at least one R^4 is selected from the group consisting of



wherein the wavy line indicates the point of attachment to the remainder of the compound.

[0076] In another group of embodiments, m is 0 or 1; n is 1, 2, or 3; and at least one R^4 is selected from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2Y$,

wherein Y is an optionally substituted member selected from the group consisting of phenyl, benzyl, pyridyl, pyridylmethyl, pyrimidinyl and pyrazolyl. More preferably, R^4 is selected from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$ and $-NR^{4a}Y$, wherein Y is an optionally substituted member selected from the group consisting of phenyl, benzyl, pyridyl and pyridylmethyl. Still further preferred are those embodiments in which the optional

substituents are selected from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $-CO_2R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-X^1OR^{4a}$ and $-X^1NR^{4a}R^{4b}$.

[0077] In other embodiments, compounds of formula I are provided in which W has the formula (c). Other embodiments are provided in which Y^1 is selected from the group consisting of optionally substituted phenyl and optionally substituted pyridyl. Other embodiments are provided in which R^5 is selected from the group consisting of $-R^{5a}$, $-C(=NH)NR^{5a}R^{5b}$, $-C(=NH)R^{5a}$, $-C(=NH)Y^1$ and $-X^3-Y^1$.

[0078] Within the present invention, the compounds provided in the examples below are each preferred embodiments, along with their pharmaceutically acceptable salts. Preferred examples of compounds of formula (I) include:

- 5 [0079] 5-Chloro-thiophene-2-carboxylic acid [3-(4-bromo-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0080] 5-Chloro-thiophene-2-carboxylic acid [3-(3-bromo-phenyl)-isoxazol-5-ylmethyl]-amide;
- [0081] 5-Chloro-thiophene-2-carboxylic acid [3-(4-phenoxy-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 10 [0082] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methylcarbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0083] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide;
- [0084] 5-Chloro-thiophene-2-carboxylic acid methyl-{3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- 15 [0085] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide;
- [0086] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide;
- 20 [0087] 1-{[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-methyl-amino]-methyl}-isoxazol-3-yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid ethyl ester;
- [0088] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-imino-methyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide;
- [0089] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-azetidin-1-ylidene-methyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide;
- 25 [0090] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-ethyl-N-methyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide;
- [0091] 5-Chloro-thiophene-2-carboxylic acid methyl-{3-[4-(N-methyl-N-propyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide;

- [0092] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-dimethylamino-ethyl)-N-methyl-carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-methyl-amide;
- [0093] 1-{[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-methyl-amino]-methyl}-isoxazol-3-yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid amide;
- 5 [0094] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-methoxy-ethyl)-N-methyl-carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-methyl-amide;
- [0095] 5-Chloro-thiophene-2-carboxylic acid methyl-{3-[4-(N-methyl-N-prop-2-ynyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0096] (R)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide;
- 10 [0097] (R)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide;
- [0098] (S)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-yl}-ethyl)-amide;
- 15 [0099] (S)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide;
- [0100] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-pyrrolidin-1-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0101] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-methylsulfanyl-phenyl]-isoxazol-5-ylmethyl}-amide;
- 20 [0102] 5-Chloro-thiophene-2-carboxylic acid (3-phenyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
- [0103] 5-Chloro-thiophene-2-carboxylic acid (5-methyl-3-phenyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
- 25 [0104] 5-Chloro-thiophene-2-carboxylic acid (3-phenyl-isoxazol-5-ylmethyl)-amide;
- [0105] 5-Chloro-thiophene-2-carboxylic acid (3-p-tolyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide;

- [0106] 5-Chloro-thiophene-2-carboxylic acid (5-methyl-3-p-tolyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
- [0107] 5-Chloro-thiophene-2-carboxylic acid (3-p-tolyl-isoxazol-5-ylmethyl)-amide;
- [0108] 5-Chloro-thiophene-2-carboxylic acid [3-(4-methoxy-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 5 [0109] 5-Chloro-thiophene-2-carboxylic acid [3-(4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0110] 5-Chloro-thiophene-2-carboxylic acid [3-(4-methoxy-phenyl)-isoxazol-5-ylmethyl]-amide;
- 10 [0111] 5-Chloro-thiophene-2-carboxylic acid [3-(4-chloro-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0112] 5-Chloro-thiophene-2-carboxylic acid [3-(4-chloro-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0113] 5-Chloro-thiophene-2-carboxylic acid [3-(4-chloro-phenyl)-isoxazol-5-ylmethyl]-amide;
- 15 [0114] 4-(5-{{(5-Chloro-thiophene-2-carbonyl)-amino}-methyl})-4,5-dihydro-isoxazol-3-yl)-benzoic acid methyl ester;
- [0115] 4-(5-{{(5-Chloro-thiophene-2-carbonyl)-amino}-methyl})-isoxazol-3-yl)-benzoic acid methyl ester;
- 20 [0116] 4-(5-{{(5-Chloro-thiophene-2-carbonyl)-amino}-methyl})-4,5-dihydro-isoxazol-3-yl)-benzoic acid;
- [0117] 4-(5-{{(5-Chloro-thiophene-2-carbonyl)-amino}-methyl})-isoxazol-3-yl)-benzoic acid;
- [0118] 5-Chloro-thiophene-2-carboxylic acid [3-(4-amino-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 25 [0119] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[(2-diethylamino-ethyl)-methyl-carbamoyl]-phenyl}-isoxazol-5-ylmethyl)-amide;

- [0120] 5-Chloro-thiophene-2-carboxylic acid [3-(1-ethyl-piperidin-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0121] 5-Chloro-thiophene-2-carboxylic acid [3-(1-benzyl-piperidin-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 5 [0122] 5-Chloro-thiophene-2-carboxylic acid [3-(1-isopropyl-piperidin-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0123] 5-Chloro-thiophene-2-carboxylic acid [3-(1-carbamimidoyl-piperidin-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0124] 5-Chloro-thiophene-2-carboxylic acid {3-[1-(1-imino-ethyl)-piperidin-4-yl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 10 [0125] 5-Chloro-thiophene-2-carboxylic acid {3-[1-(imino-phenyl-methyl)-piperidin-4-yl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0126] 5-Chloro-thiophene-2-carboxylic acid {3-[1-(imino-pyridin-2-yl-methyl)-piperidin-4-yl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 15 [0127] 5-Chloro-thiophene-2-carboxylic acid (3-piperidin-4-yl-isoxazol-5-ylmethyl)-amide;
- [0128] 5-Chloro-thiophene-2-carboxylic acid [3-(1-ethyl-piperidin-4-yl)-isoxazol-5-ylmethyl]-amide;
- [0129] 5-Chloro-thiophene-2-carboxylic acid [3-(1-benzyl-piperidin-4-yl)-isoxazol-5-ylmethyl]-amide;
- 20 [0130] 5-Chloro-thiophene-2-carboxylic acid [3-(1-carbamimidoyl-piperidin-4-yl)-isoxazol-5-ylmethyl]-amide;
- [0131] 5-Chloro-thiophene-2-carboxylic acid {3-[1-(1-imino-ethyl)-piperidin-4-yl]-isoxazol-5-ylmethyl}-amide;
- 25 [0132] 5-Chloro-thiophene-2-carboxylic acid {3-[1-(imino-phenyl-methyl)-piperidin-4-yl]-isoxazol-5-ylmethyl}-amide;
- [0133] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-formyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;

- [0134] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-chloro-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0135] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-methylsulfanyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 5 [0136] 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-4-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0137] 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-3-yl-phenyl)-isoxazol-5-ylmethyl]-amide;
- [0138] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-azepan-1-ylmethyl-biphenyl-4-yl)-4,5-
10 dihydro-isoxazol-5-ylmethyl]-amide;
- [0139] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-5'-methoxy-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0140] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-hydroxymethyl-4'-methoxy-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 15 [0141] 5-Chloro-thiophene-2-carboxylic acid [3-(3'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide;
- [0142] 5-Chloro-thiophene-2-carboxylic acid [3-(4'-dimethylaminomethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide;
- [0143] 5-Chloro-thiophene-2-carboxylic acid [3-(4'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-
20 isoxazol-5-ylmethyl]-amide;
- [0144] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-5'-methoxy-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide;
- [0145] 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-2-yl-phenyl)-isoxazol-5-ylmethyl]-amide;
- 25 [0146] 5-Chloro-thiophene-2-carboxylic acid [3-(4-benzylamino-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0147] 5-Chloro-thiophene-2-carboxylic acid [3-(4-phenylamino-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;

- [0148] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(4-fluoro-phenylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0149] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-2-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 5 [0150] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-methyl-pyridin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0151] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[(2-amino-pyrimidin-4-yl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
- [0152] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-ethyl-2H-pyrazol-3-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 10 [0153] 5-Chloro-thiophene-2-carboxylic acid [3-(4-morpholin-4-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0154] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-methoxymethyl-pyrrolidin-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 15 [0155] 5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(5-oxo-[1,4]oxazepan-4-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0156] 5-Chloro-thiophene-2-carboxylic acid [3-(4-phenylamino-phenyl)-isoxazol-5-ylmethyl]-amide;
- [0157] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(4-methoxy-phenylamino)-phenyl]-isoxazol-5-ylmethyl}-amide;
- 20 [0158] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-2-ylamino)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0159] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(methyl-pyridin-4-yl-amino)-phenyl]-isoxazol-5-ylmethyl}-amide;
- 25 [0160] 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyrrolidin-1-yl-phenyl)-isoxazol-5-ylmethyl]-amide;
- [0161] 5-Chloro-thiophene-2-carboxylic acid [3-(4-morpholin-4-yl-phenyl)-isoxazol-5-ylmethyl]-amide;

- [0162] 5-Chloro-thiophene-2-carboxylic acid [3-(4-benzylamino-phenyl)-isoxazol-5-ylmethyl]-amide;
- [0163] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-piperidin-4-ylamino)-phenyl]-isoxazol-5-ylmethyl}-amide;
- 5 [0164] 2-[4-(5-{{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-phenylamino]-benzoic acid methyl ester;
- [0165] 3-[4-(5-{{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-phenylamino]-benzoic acid methyl ester;
- [0166] 4-[4-(5-{{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-phenylamino]-benzoic acid methyl ester;
- 10 [0167] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0168] 5-Chloro-thiophene-2-carboxylic acid [3-(3'-dimethylaminomethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 15 [0169] 5-Chloro-thiophene-2-carboxylic acid [3-(3'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0170] 5-Chloro-thiophene-2-carboxylic acid [3-(4'-dimethylaminomethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0171] 5-Chloro-thiophene-2-carboxylic acid [3-(4'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 20 [0172] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide;
- [0173] 5-Chloro-thiophene-2-carboxylic acid [3-(3'-dimethylaminomethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide;
- 25 [0174] 5-Chloro-thiophene-2-carboxylic acid [3-(3'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide;
- [0175] 5-Chloro-thiophene-2-carboxylic acid [3-(4'-dimethylaminomethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide;

- [0176] 5-Chloro-thiophene-2-carboxylic acid [3-(4'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide;
- [0177] 5-Chloro-thiophene-2-carboxylic acid [3-(3-pyridin-3-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 5 [0178] 5-Chloro-thiophene-2-carboxylic acid [3-(3-pyridin-4-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; and
- [0179] 5-Chloro-thiophene-2-carboxylic acid [3-(3-pyridin-3-yl-phenyl)-isoxazol-5-ylmethyl]-amide.
- [0180] More preferred examples of the compounds of formula (I) include:
- 10 [0181] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0182] 1-{[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid ethyl ester;
- [0183] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methyl-N-prop-2-ynyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 15 [0184] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-methoxy-ethyl)-N-methyl-carbamimidoyl]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
- [0185] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-furan-2-ylmethyl-N-methyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 20 [0186] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methyl-N-prop-2-ynyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0187] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-methoxy-ethyl)-N-methyl-carbamimidoyl]-phenyl}-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
- [0188] 1-{[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-5-methyl-4,5-dihydro-isoxazol-3-yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid amide;
- 25 [0189] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-fluoro-phenyl]-dihydro-isoxazol-5-ylmethyl}-amide;

- [0190] 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(N,N-dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0191] 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 5 [0192] 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0193] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-pyrrolidin-1-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0194] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-pyrrolidin-1-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 10 [0195] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-pyrrolidin-1-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0196] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-morpholin-4-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 15 [0197] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-morpholin-4-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0198] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-morpholin-4-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0199] 5-Chloro-thiophene-2-carboxylic acid {3-[2-[(2-methoxy-ethyl)-methyl-amino]-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 20 [0200] 5-Chloro-thiophene-2-carboxylic acid (3-{4-(imino-pyrrolidin-1-yl-methyl)-2-[(2-methoxy-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
- [0201] 5-Chloro-thiophene-2-carboxylic acid (3-{4-(imino-piperidin-1-yl-methyl)-2-[(2-methoxy-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
- 25 [0202] 5-Chloro-thiophene-2-carboxylic acid {3-[2-azepan-1-yl-4-(N,N-dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0203] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-methyl-pyrrolidine-1-carbonyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;

- [0204] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyrrolidine-1-carbonyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0205] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-amino-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 5 [0206] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-trifluoromethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0207] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-methoxy-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0208] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-methoxymethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide;
- 10 [0209] 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-3-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0210] 5-Chloro-thiophene-2-carboxylic acid [3-(5'-chloro-2'-{[(2-methoxy-ethyl)-methyl-amino]-methyl}-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 15 [0211] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide;
- [0212] 5-Chloro-thiophene-2-carboxylic acid [3-(3'-dimethylaminomethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide;
- [0213] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-chloro-pyridin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 20 [0214] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyrimidin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0215] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-1H-pyrazol-3-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 25 [0216] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(4-methyl-piperazin-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0217] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-piperidin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;

[0218] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[(2-dimethylamino-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide;

[0219] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-oxo-piperidin-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; and

5 [0220] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(3-oxo-thiomorpholin-4-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide.

[0221] Particularly preferred examples of the compounds of formula (I) include:

[0222] 5-Chloro-thiophene-2-carboxylic acid [3-(4-cyano-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;

10 [0223] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;

[0224] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;

15 [0225] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;

[0226] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-dimethylamino-ethyl)-N-methyl-carbamimidoyl]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide;

[0227] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methyl-N-propyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;

20 [0228] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-ethyl-N-methyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;

[0229] 5-Chloro-thiophene-2-carboxylic acid [3-(4-cyano-phenyl)-isoxazol-5-ylmethyl]-amide;

25 [0230] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide;

[0231] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide;

- [0232] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0233] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- 5 [0234] 1-{[4-(5-{{(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-isoxazol-3-yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid ethyl ester;
- [0235] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-imino-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0236] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-azetidin-1-ylidene-10 methyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0237] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-ethyl-N-methyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0238] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methyl-N-propyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- 15 [0239] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-dimethylamino-ethyl)-N-methyl-carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-amide;
- [0240] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-furan-2-ylmethyl-N-methyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0241] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-methoxy-ethyl)-N-methyl-20 carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-amide;
- [0242] 5-Chloro-thiophene-2-carboxylic acid methyl-{3-[4-(N-methylcarbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0243] (R)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide;
- 25 [0244] (R)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-yl}-ethyl)-amide;
- [0245] (R)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide;

- [0246] (S)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide;
- [0247] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 5 [0248] 5-Chloro-thiophene-2-carboxylic acid {5-methyl-3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0249] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0250] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-
10 5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0251] 1-{[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-5-methyl-4,5-dihydro-isoxazol-3-yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid ethyl ester;
- [0252] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-imino-methyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 15 [0253] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-azetidin-1-ylidene-methyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0254] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-ethyl-N-methyl-carbamimidoyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0255] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-dimethylamino-ethyl)-N-methyl-carbamimidoyl]-phenyl}-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
20
- [0256] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-furan-2-ylmethyl-N-methyl-carbamimidoyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0257] 5-Chloro-thiophene-2-carboxylic acid {5-methyl-3-[4-(N-methylcarbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 25 [0258] 5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0259] 5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide;

- [0260] 5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0261] 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(imino-piperidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 5 [0262] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-morpholin-4-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0263] 5-Chloro-thiophene-2-carboxylic acid (3-{4-(N,N-dimethyl-carbamimidoyl)-2-[(2-methoxy-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
- [0264] 5-Chloro-thiophene-2-carboxylic acid {3-[2-azepan-1-yl-4-(1-methyl-4,5-dihydro-10 1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0265] 5-Chloro-thiophene-2-carboxylic acid {3-[2-azepan-1-yl-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0266] 5-Chloro-thiophene-2-carboxylic acid {3-[2-azepan-1-yl-4-(imino-piperidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 15 [0267] 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(N,N-dimethyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0268] 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0269] 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(imino-pyrrolidin-1-20 yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0270] 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0271] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-morpholin-4-yl-phenyl]-isoxazol-5-ylmethyl}-amide;
- 25 [0272] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-morpholin-4-yl-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0273] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-morpholin-4-yl-phenyl]-isoxazol-5-ylmethyl}-amide;

- [0274] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-morpholin-4-yl-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0275] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-methoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 5 [0276] 5-Chloro-thiophene-2-carboxylic acid {3-[2-methoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0277] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-methoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0278] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-methoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 10 [0279] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-ethoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0280] 5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 15 [0281] 5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0282] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-isopropoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0283] 5-Chloro-thiophene-2-carboxylic acid {3-[2-isopropoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 20 [0284] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-isopropoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0285] 5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclopentyloxy-4-(N,N-dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 25 [0286] 5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclopentyloxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0287] 5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclopentyloxy-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;

- [0288] 5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclohexyloxy-4-(N,N-dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0289] 5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclohexyloxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 5 [0290] 5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclohexyloxy-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0291] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-methoxy-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0292] 5-Chloro-thiophene-2-carboxylic acid {3-[2-methoxy-4-(1-methyl-4,5-dihydro-1H-
10 imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0293] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-methoxy-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0294] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-methoxy-phenyl]-isoxazol-5-ylmethyl}-amide;
- 15 [0295] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-ethoxy-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0296] 5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0297] 5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(imino-pyrrolidin-1-yl-
20 methyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0298] 5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0299] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-isopropoxy-phenyl]-isoxazol-5-ylmethyl}-amide;
- 25 [0300] 5-Chloro-thiophene-2-carboxylic acid {3-[2-isopropoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0301] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-isopropoxy-phenyl]-isoxazol-5-ylmethyl}-amide;

- [0302] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-isopropoxy-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0303] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-phenoxy-phenyl]-isoxazol-5-ylmethyl}-amide;
- 5 [0304] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-phenoxy-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0305] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-phenoxy-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0306] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-(4-methoxy-phenoxy)-phenyl]-isoxazol-5-ylmethyl}-amide;
- 10 [0307] 5-Chloro-thiophene-2-carboxylic acid {3-[2-(4-methoxy-phenoxy)-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0308] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-(4-methoxy-phenoxy)-phenyl]-isoxazol-5-ylmethyl}-amide;
- 15 [0309] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-methylsulfanyl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0310] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-methylsulfanyl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0311] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-methylsulfanyl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 20 [0312] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-methylsulfanyl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0313] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-methylsulfanyl-phenyl]-isoxazol-5-ylmethyl}-amide;
- 25 [0314] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-methylsulfanyl-phenyl]-isoxazol-5-ylmethyl}-amide;
- [0315] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-methylsulfanyl-phenyl]-isoxazol-5-ylmethyl}-amide;

- [0316] 5-Chloro-thiophene-2-carboxylic acid [3-(4-amino-phenyl)-isoxazol-5-ylmethyl]-amide;
- [0317] 5-Chloro-thiophene-2-carboxylic acid [3-(4-dimethylcarbamoyl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 5 [0318] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidine-1-carbonyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0319] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyrrolidine-1-carbonyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0320] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[(2-diethylamino-ethyl)-methyl-carbamoyl]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
- 10 [0321] 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-4-yl-phenyl)-isoxazol-5-ylmethyl]-amide;
- [0322] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 15 [0323] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0324] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-piperidin-1-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0325] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-morpholin-4-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 20 [0326] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-{[(2-methoxy-ethyl)-methyl-amino]-methyl}-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0327] 5-Chloro-thiophene-2-carboxylic acid {3-[2'-(4-methyl-piperazin-1-ylmethyl)-biphenyl-4-yl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 25 [0328] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-{[(2-dimethylamino-ethyl)-methyl-amino]-methyl}-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0329] 5-Chloro-thiophene-2-carboxylic acid [3-(5'-methoxy-2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;

- [0330] 5-Chloro-thiophene-2-carboxylic acid [3-(5'-chloro-2'-dimethylaminomethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- [0331] 5-Chloro-thiophene-2-carboxylic acid [3-(5'-chloro-2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 5 [0332] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-formyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide;
- [0333] 5-Chloro-thiophene-2-carboxylic acid [3-(5'-methoxy-2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide;
- 10 [0334] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-4'-methoxy-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide;
- [0335] 5-Chloro-thiophene-2-carboxylic acid [3-(4'-methoxy-2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide;
- [0336] 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-2-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
- 15 [0337] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-3-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0338] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 20 [0339] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(methyl-pyridin-4-yl-amino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0340] 5-Chloro-thiophene-2-carboxylic acid (3-{4-[(3-dimethylamino-propyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
- [0341] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-oxo-pyrrolidin-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 25 [0342] 4-[4-(5-[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl)-4,5-dihydro-isoxazol-3-yl)-phenyl]-3-oxo-piperazine-1-carboxylic acid benzyl ester;
- [0343] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(3-oxo-morpholin-4-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;

- [0344] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(3-methyl-2-oxo-imidazolidin-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0345] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(5-oxo-[1,4]oxazepan-4-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- 5 [0346] 5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(pyridin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0347] 5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(4-methyl-[1,4]diazepan-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide;
- [0348] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-3-ylamino)-phenyl]-isoxazol-10 5-ylmethyl}-amide;
- [0349] 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-4-ylamino)-phenyl]-isoxazol-5-ylmethyl}-amide; and
- [0350] 5-Chloro-thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide.
- 15 [0351] 5-Chloro-N-((3-(4-(2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;
- [0352] 5-Chloro-N-((3-(4-(3-fluoro-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;
- [0353] 5-Chloro-N-((3-(4-(5-fluoro-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-20 yl)methyl)thiophene-2-carboxamide;
- [0354] 5-Chloro-N-((3-(4-(3-methyl-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;
- [0355] 5-Chloro-N-((3-(4-(3-methoxy-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;
- 25 [0356] 5-Chloro-N-((3-(4-(3-chloro-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;
- [0357] 5-Chloro-N-((3-(4-(4-methyl-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;

- [0358] 5-Chloro-N-((3-(4-(5-methyl-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;
- [0359] 5-Chloro-N-((3-(4-(5-methoxy-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;
- 5 [0360] 5-Chloro-N-((3-(4-(5-chloro-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;
- [0361] 5-Chloro-N-((3-(4-iodophenyl)isoxazol-5-yl)methyl)thiophene-2-carboxamide;
- [0362] 5-Chloro-N-((3-(4-(2-oxopyridin-1(2H)-yl)phenyl)isoxazol-5-yl)methyl)thiophene-2-carboxamide;
- 10 [0363] 5-Chloro-N-((1-(4-(2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0364] 5-Chloro-N-((1-(4-(3-fluoro-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0365] 5-Chloro-N-((1-(4-(3-chloro-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 15 [0366] 5-Chloro-N-((1-(4-(3-methyl-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0367] 5-Chloro-N-((1-(4-(3-methoxy-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 20 [0368] 5-Chloro-N-((1-(4-(4-methyl-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0369] 5-Chloro-N-((1-(4-(5-methyl-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0370] 5-Chloro-N-((1-(4-(5-methoxy-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 25 [0371] 5-Chloro-N-((1-(4-(5-fluoro-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;

- [0372] 5-Chloro-N-((1-(4-(5-chloro-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0373] 5-Chloro-N-((1-(4-(2-oxopiperidin-1-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 5 [0374] 5-Chloro-N-((1-(4-(3-oxomorpholino)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0375] 5-Chloro-N-((1-(4-(3-oxothiomorpholino)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0376] 5-Chloro-N-((1-(4-(5-oxo-1,4-oxazepan-4-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 10 [0377] 5-Chloro-N-((1-(2-fluoro-4-iodophenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0378] 5-Chloro-N-((1-(2-fluoro-4-(2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 15 [0379] 5-Chloro-N-((1-(2-fluoro-4-(3-oxomorpholino)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0380] 5-Chloro-N-((1-(2-hydroxy-4-(2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0381] 5-Chloro-N-((1-(2-(4-methylpiperazin-1-yl)-4-(2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 20 [0382] 5-Chloro-N-((1-(5-(2-oxopyridin-1(2H)-yl)pyridin-2-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0383] 5-Chloro-N-((3-(6-(2-oxopyridin-1(2H)-yl)pyridin-3-yl)isoxazol-5-yl)methyl)thiophene-2-carboxamide;
- 25 [0384] 5-Chloro-N-((3-(4-(2-oxopyridin-1(2H)-yl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)thiophene-2-carboxamide;
- [0385] 5-Chloro-N-((1-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;

- [0386] N-((1-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0387] N-((1-(4-(N-Ethyl-N-methylcarbamimidoyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- 5 [0388] N-((1-(4-(N-Methyl-N-propylcarbamimidoyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0389] N-((1-(4-(N-(2-Methoxyethyl)-N-methylcarbamimidoyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0390] N-((1-(4-(Azetidin-1-yl(imino)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- 10 [0391] 5-Chloro-N-((1-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0392] 5-Chloro-N-((1-(4-(imino(piperidin-1-yl)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 15 [0393] N-((1-(4-(N,N-Dimethylcarbamimidoyl)-2-fluorophenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0394] N-((1-(4-(Azetidin-1-yl(imino)methyl)-2-fluorophenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0395] 5-Chloro-N-((1-(2-fluoro-4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 20 [0396] 5-Chloro-N-((1-(2-fluoro-4-(imino(piperidin-1-yl)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0397] 5-Chloro-N-((1-(2-fluoro-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 25 [0398] N-((1-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-1H-pyrazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0399] 5-Chloro-N-((1-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1H-pyrazol-4-yl)methyl)thiophene-2-carboxamide;

- [0400] N-((1-(4-(Azetidin-1-yl(imino)methyl)phenyl)-1H-pyrazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0401] 5-Chloro-N-((1-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1H-pyrazol-4-yl)methyl)thiophene-2-carboxamide;
- 5 [0402] 5-Chloro-N-((1-(4-(imino(piperidin-1-yl)methyl)phenyl)-1H-pyrazol-4-yl)methyl)thiophene-2-carboxamide;
- [0403] N-((5-(4-(N,N-Dimethylcarbamidoyl)phenyl)isoxazol-3-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0404] 5-Chloro-N-((5-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)isoxazol-3-yl)methyl)thiophene-2-carboxamide;
- 10 [0405] 5-Chloro-N-((5-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)isoxazol-3-yl)methyl)thiophene-2-carboxamide;
- [0406] 5-Chloro-N-((5-(4-(imino(piperidin-1-yl)methyl)phenyl)isoxazol-3-yl)methyl)thiophene-2-carboxamide;
- 15 [0407] N-((3-(4-(N,N-Dimethylcarbamidoyl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0408] 5-Chloro-N-((3-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)thiophene-2-carboxamide;
- [0409] 5-Chloro-N-((3-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)thiophene-2-carboxamide;
- 20 [0410] 5-Chloro-N-((3-(4-(imino(piperidin-1-yl)methyl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)thiophene-2-carboxamide;
- [0411] N-((2-(4-(N,N-Dimethylcarbamidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- 25 [0412] 5-Chloro-N-((2-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)oxazol-4-yl)methyl)thiophene-2-carboxamide;
- [0413] 5-Chloro-N-((2-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)oxazol-4-yl)methyl)thiophene-2-carboxamide;

- [0414] 5-Chloro-N-((2-(4-(imino(piperidin-1-yl)methyl)phenyl)oxazol-4-yl)methyl)thiophene-2-carboxamide;
- [0415] Ethyl 1-((4-(4-((2-chlorothiophene-5-carboxamido)methyl)oxazol-2-yl)phenyl)(imino)methyl)piperidine-4-carboxylate;
- 5 [0416] N-((2-(4-(N-Ethyl-N-methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0417] N-((2-(4-(N-Methyl-N-propylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0418] N-((2-(4-(N-(2-(Dimethylamino)ethyl)-N-methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- 10 [0419] N-((2-(4-(N-(2-Methoxyethyl)-N-methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0420] 1-((4-(4-((2-Chlorothiophene-5-carboxamido)methyl)oxazol-2-yl)phenyl)(imino)methyl)piperidine-4-carboxamide;
- 15 [0421] N-((2-(4-(N-Methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0422] N-((2-(4-(N-(Furan-2-ylmethyl)-N-methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0423] N-((2-(4-(N,N-Dimethylcarbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- 20 [0424] 5-Chloro-N-((2-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)thiazol-4-yl)methyl)thiophene-2-carboxamide;
- [0425] 5-Chloro-N-((2-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)thiazol-4-yl)methyl)thiophene-2-carboxamide;
- 25 [0426] 5-Chloro-N-((2-(4-(imino(piperidin-1-yl)methyl)phenyl)thiazol-4-yl)methyl)thiophene-2-carboxamide;
- [0427] Ethyl 1-((4-(4-((2-chlorothiophene-5-carboxamido)methyl)thiazol-2-yl)phenyl)(imino)methyl)piperidine-4-carboxylate;

- [0428] N-((2-(4-(N-(2-(Dimethylamino)ethyl)-N-methylcarbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0429] N-((2-(4-(N-(Furan-2-ylmethyl)-N-methylcarbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- 5 [0430] N-((2-(4-(N-Methyl-N-(prop-2-ynyl)carbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0431] N-((2-(4-(N-Methylcarbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0432] 5-Chloro-N-((4-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)oxazol-2-yl)methyl)thiophene-2-carboxamide;
- 10 [0433] N-((4-(4-(N,N-Dimethylcarbamimidoyl)phenyl)thiazol-2-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0434] 5-Chloro-N-((4-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)thiazol-2-yl)methyl)thiophene-2-carboxamide;
- 15 [0435] 5-Chloro-N-((4-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)thiazol-2-yl)methyl)thiophene-2-carboxamide;
- [0436] 5-Chloro-N-((4-(4-(imino(piperidin-1-yl)methyl)phenyl)thiazol-2-yl)methyl)thiophene-2-carboxamide;
- [0437] N-((1-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-1H-imidazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- 20 [0438] 5-Chloro-N-((1-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1H-imidazol-4-yl)methyl)thiophene-2-carboxamide;
- [0439] N-((1-(4-(Azetidin-1-yl(imino)methyl)phenyl)-1H-imidazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- 25 [0440] 5-Chloro-N-((1-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1H-imidazol-4-yl)methyl)thiophene-2-carboxamide;
- [0441] 5-Chloro-N-((1-(4-(imino(piperidin-1-yl)methyl)phenyl)-1H-imidazol-4-yl)methyl)thiophene-2-carboxamide;

- [0442] N-((5-(4-(N,N-Dimethylcarbamimidoyl)phenyl)pyridin-3-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0443] 5-Chloro-N-((5-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)pyridin-3-yl)methyl)thiophene-2-carboxamide;
- 5 [0444] N-((5-(4-(Azetidin-1-yl(imino)methyl)phenyl)pyridin-3-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0445] 5-Chloro-N-((5-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)pyridin-3-yl)methyl)thiophene-2-carboxamide;
- [0446] 5-Chloro-N-((5-(4-(imino(piperidin-1-yl)methyl)phenyl)pyridin-3-yl)methyl)thiophene-2-carboxamide;
- 10 [0447] N-((5-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-2-fluorophenyl)methyl)-5-chlorothiophene-2-carboxamide;
- [0448] 5-Chloro-N-((5-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-2-fluorophenyl)methyl)thiophene-2-carboxamide;
- 15 [0449] 5-Chloro-N-((5-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-2-fluorophenyl)methyl)thiophene-2-carboxamide;
- [0450] N-((3-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-4-fluorophenyl)methyl)-5-chlorothiophene-2-carboxamide;
- [0451] 5-Chloro-N-((3-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-4-fluorophenyl)methyl)thiophene-2-carboxamide;
- 20 [0452] 5-Chloro-N-((3-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-4-fluorophenyl)methyl)thiophene-2-carboxamide;
- [0453] N-((3-(4-(N,N-Dimethylcarbamimidoyl)phenyl)phenyl)methyl)-5-chlorothiophene-2-carboxamide;
- 25 [0454] 5-Chloro-N-((3-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)phenyl)methyl)thiophene-2-carboxamide;
- [0455] 5-Chloro-N-((3-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)phenyl)methyl)thiophene-2-carboxamide;

- [0456] 5-Chloro-N-((1-(6-(4-methyl-1,4-diazepan-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0457] N-((1-(6-(1,4-Diazepan-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- 5 [0458] N-((1-(6-(1,4-Oxazepan-4-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
- [0459] 5-Chloro-N-((1-(6-morpholinopyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0460] 5-Chloro-N-((1-(6-(3-oxomorpholino)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 10 [0461] 5-Chloro-N-((1-(6-(2-oxopiperidin-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0462] 5-Chloro-N-((1-(6-(2-oxopyridin-1(2H)-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 15 [0463] 5-Chloro-N-((1-(6-(piperazin-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0464] 5-Chloro-N-((1-(6-(4-methylpiperazin-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- [0465] 5-Chloro-N-((1-(6-(piperidin-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide;
- 20 [0466] 5-Chloro-N-((3-(4-(8-(trifluoromethyl)quinolin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;
- [0467] 5-Chloro-N-((3-(4-(7-(trifluoromethyl)quinolin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;
- 25 [0468] 5-Chloro-N-((3-(4-(6-(trifluoromethyl)quinolin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;
- [0469] 5-Chloro-N-((3-(4-(quinolin-6-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;

[0470] 5-Chloro-N-((3-(4-(2,5-dihydro-1H-pyrrole-1-carbonyl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;

[0471] 5-Chloro-N-((3-(4-(2,5-dihydro-1H-pyrrole-1-carbonyl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;

5 [0472] N-((3-(4-((2-Hydroxypropyl)carbamoyl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)-5-chlorothiophene-2-carboxamide;

[0473] 5-Chloro-N-((3-(4-(3-methylpyridin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;

10 [0474] 5-Chloro-N-((3-(4-(2-ethoxypyridin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide;

[0475] 5-Chloro-N-((3-(4-(3,5-dichloropyridin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide; and

[0476] 5-Chloro-N-((3-(4-(3-(trifluoromethyl)pyridin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide

15 [0477] All the preferred, more preferred, and most preferred compounds listed above are selective inhibitors of Factor Xa.

Compositions

20 [0478] The present invention further provides compositions comprising one or more compounds of formula (I) or a tautomer or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier. It will be appreciated that the compounds of formula (I) in this invention may be derivatized at functional groups to provide prodrug derivatives which are capable of conversion back to the parent compounds in vivo. Examples of such
25 prodrugs include the physiologically acceptable and metabolically labile ester derivatives, such as methoxymethyl esters, methylthiomethyl esters, or pivaloyloxymethyl esters derived from a hydroxyl group of the compound or a carbamoyl moiety derived from an amino group of the compound. Additionally, any physiologically acceptable equivalents of the compounds of formula (I), similar to metabolically labile esters or carbamates, which are

capable of producing the parent compounds of formula (I) *in vivo*, are within the scope of this invention.

[0479] If pharmaceutically acceptable salts of the compounds of this invention are utilized in these compositions, those salts are preferably derived from inorganic or organic acids and bases. Included among such acid salts are the following: acetate, adipate, alginate, aspartate, benzoate, benzene sulfonate, bisulfate, butyrate, citrate, camphorate, camphor sulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, lucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate and undecanoate. Base salts include ammonium salts, alkali metal salts, such as sodium and potassium salts, alkaline earth metal salts, such as calcium and magnesium salts, salts with organic bases, such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth.

[0480] Furthermore, the basic nitrogen-containing groups may be quaternized with agents like lower alkyl halides, such as methyl, ethyl, propyl and butyl chlorides, bromides and iodides; dialkyl sulfates, such as dimethyl, diethyl, dibutyl and diamyl sulfates, long chain halides, such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; aralkyl halides, such as benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products are thereby obtained.

[0481] The compounds utilized in the compositions and methods of this invention may also be modified by appending appropriate functionalities to enhance selective biological properties. Such modifications are known in the art and include those which increase biological penetration into a given biological system (*e.g.*, blood, lymphatic system, central nervous system, etc.), increase oral availability, increase solubility to allow administration by injection, alter metabolism and alter rate of excretion.

[0482] The pharmaceutical compositions of the invention can be manufactured by methods well known in the art such as conventional granulating, mixing, dissolving, encapsulating, lyophilizing, or emulsifying processes, among others. Compositions may be produced in various forms, including granules, precipitates, or particulates, powders, including freeze dried, rotary dried or spray dried powders, amorphous powders, tablets, capsules, syrup,

suppositories, injections, emulsions, elixirs, suspensions or solutions. Formulations may optionally contain stabilizers, pH modifiers, surfactants, bioavailability modifiers and combinations of these.

[0483] Pharmaceutical formulations may be prepared as liquid suspensions or solutions

5 using a sterile liquid, such as oil, water, alcohol, and combinations thereof. Pharmaceutically suitable surfactants, suspending agents or emulsifying agents, may be added for oral or parenteral administration. Suspensions may include oils, such as peanut oil, sesame oil, cottonseed oil, corn oil and olive oil. Suspension preparation may also contain esters of fatty acids, such as ethyl oleate, isopropyl myristate, fatty acid glycerides and acetylated fatty acid
10 glycerides. Suspension formulations may include alcohols, such as ethanol, isopropyl alcohol, hexadecyl alcohol, glycerol and propylene glycol. Ethers, such as poly(ethyleneglycol), petroleum hydrocarbons, such as mineral oil and petrolatum, and water may also be used in suspension formulations.

[0484] Pharmaceutically acceptable carriers that may be used in these compositions include

15 ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances, such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-
20 based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[0485] According to a preferred embodiment, the compositions of this invention are

formulated for pharmaceutical administration to a mammal, preferably a human being. Such pharmaceutical compositions of the invention may be administered orally, parenterally, by
25 inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally or intravenously. The formulations of the invention may be designed as short-acting,
30 fast-releasing, or long-acting. Still further, compounds can be administered in a local rather than systemic means, such as administration (*e.g.*, injection) as a sustained release formulation.

[0486] Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation. Compounds may be formulated for parenteral administration by injection such as by bolus injection or continuous infusion. A unit dosage form for injection may be in ampoules or in multi-dose containers.

[0487] The pharmaceutical compositions of this invention may be in any orally acceptable dosage form, including capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers that are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

[0488] Alternatively, the pharmaceutical compositions of this invention may be in the form of suppositories for rectal administration. These may be prepared by mixing the agent with a suitable non-irritating excipient which is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

[0489] The pharmaceutical compositions of this invention may also be in a topical form, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

5 [0490] Topical application for the lower intestinal tract may be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used. For topical applications, the pharmaceutical compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the
10 compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical compositions may be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include
15 mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters, wax, cetyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[0491] For ophthalmic use, the pharmaceutical compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative, such as
20 benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutical compositions may be formulated in an ointment, such as petrolatum.

[0492] The pharmaceutical compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques known in the art of pharmaceutical formulation and may be prepared as solutions in saline,
25 employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons and/or other conventional solubilizing or dispersing agents.

[0493] Any of the above dosage forms containing effective amounts are within the bounds of routine experimentation and within the scope of the invention. A therapeutically effective dose may vary depending upon the route of administration and dosage form. The preferred
30 compound or compounds of the invention is a formulation that exhibits a high therapeutic index. The therapeutic index is the dose ratio between toxic and therapeutic effects which can be expressed as the ratio between LD₅₀ and ED₅₀. The LD₅₀ is the dose lethal to 50% of

the population and the ED₅₀ is the dose therapeutically effective in 50% of the population. The LD₅₀ and ED₅₀ are determined by standard pharmaceutical procedures in animal cell cultures or experimental animals.

[0494] Besides those representative dosage forms described above, pharmaceutically acceptable excipients and carriers and dosage forms are generally known to those skilled in the art and are included in the invention. It should be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex and diet of the patient, and the time of administration, rate of excretion, drug combination, judgment of the treating physician and severity of the particular disease being treated. The amount of active ingredient(s) will also depend upon the particular compound and other therapeutic agent, if present, in the composition.

Methods of Use

15

[0495] The invention provides methods of inhibiting or decreasing Factor Xa activity as well as treating or ameliorating a Factor Xa associated state, symptom, disorder or disease in a patient in need thereof (*e.g.*, human or non-human). "Treating" within the context of the invention means an alleviation of symptoms associated with a disorder or disease, or halt of further progression or worsening of those symptoms, or prevention or prophylaxis of the disease or disorder.

20

[0496] The term "mammal" includes organisms which express Factor Xa. Examples of mammals include mice, rats, cows, sheep, pigs, goats, horses, bears, monkeys, dogs, cats and, preferably, humans. Transgenic organisms which express Factor Xa are also included in this definition.

25

[0497] The inventive methods comprise administering an effective amount of a compound or composition described herein to a mammal or non-human animal. As used herein, "effective amount" of a compound or composition of the invention includes those amounts that antagonize or inhibit Factor Xa. An amount which antagonizes or inhibits Factor Xa is detectable, for example, by any assay capable of determining Factor Xa activity, including the one described below as an illustrative testing method. Effective amounts may also include those amounts which alleviate symptoms of a Factor Xa associated disorder treatable

30

by inhibiting Factor Xa. Accordingly, “antagonists of Factor Xa” include compounds which interact with the Factor Xa and modulate, *e.g.*, inhibit or decrease, the ability of a second compound, *e.g.*, another Factor Xa ligand, to interact with the Factor Xa. The Factor Xa binding compounds are preferably antagonists of Factor Xa. The language “Factor Xa binding compound” (*e.g.*, exhibits binding affinity to the receptor) includes those compounds which interact with Factor Xa resulting in modulation of the activity of Factor Xa. Factor Xa binding compounds may be identified using an *in vitro* (*e.g.*, cell and non-cell based) or *in vivo* method. A description of an *in vitro* method is provided below.

[0498] The amount of compound present in the methods and compositions described herein should be sufficient to cause a detectable decrease in the severity of the disorder, as measured by any of the assays described in the examples. The amount of Factor Xa modulator needed will depend on the effectiveness of the modulator for the given cell type and the length of time required to treat the disorder. In certain embodiments, the compositions of this invention may further comprise another therapeutic agent. When a second agent is used, the second agent may be administered either as a separate dosage form or as part of a single dosage form with the compounds or compositions of this invention. While one or more of the inventive compounds can be used in an application of monotherapy to treat a disorder, disease or symptom, they also may be used in combination therapy, in which the use of an inventive compound or composition (therapeutic agent) is combined with the use of one or more other therapeutic agents for treating the same and/or other types of disorders, symptoms and diseases. Combination therapy includes administration of the two or more therapeutic agents concurrently or sequentially. The agents may be administered in any order. Alternatively, the multiple therapeutic agents can be combined into a single composition that can be administered to the patient. For instance, a single pharmaceutical composition could comprise the compound or pharmaceutically acceptable salt or solvate according to the formula I, another therapeutic agent (*e.g.*, methotrexate) or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable excipient or carrier.

[0499] The invention comprises a compound having the formula I, a method for making an inventive compound, a method for making a pharmaceutical composition from at least one inventive compound and at least one pharmaceutically acceptable carrier or excipient, and a method of using one or more inventive compounds to treat a variety of disorders, symptoms and diseases (*e.g.*, inflammatory, autoimmune, neurological, neurodegenerative, oncology and cardiovascular), such as RA, osteoarthritis, irritable bowel disease IBD, asthma, chronic

obstructive pulmonary disease COPD and MS. The inventive compounds and their pharmaceutically acceptable salts and/or neutral compositions may be formulated together with a pharmaceutically acceptable excipient or carrier and the resulting composition may be administered in vivo to mammals, such as men, women and animals, to treat a variety of disorders, symptoms and diseases. Furthermore, the inventive compounds can be used to prepare a medicament that is useful for treating a variety of disorders, symptoms and diseases.

Kits


[0500] Still another aspect of this invention is to provide a kit comprising separate containers in a single package, wherein the inventive pharmaceutical compounds, compositions and/or salts thereof are used in combination with pharmaceutically acceptable carriers to treat states, disorders, symptoms and diseases where Factor Xa plays a role.

EXAMPLES

EXAMPLE 1

5-Chloro-thiophene-2-carboxylic acid [3-(4-bromo-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide (7)

SCHEME 1



Step 1:

[0501] Aldehyde **1** was dissolved in 200 mL ethanol, then treated with pyridine (24 mL, 297 mmol) and hydroxylamine hydrochloride (19.58 g, 284 mmol) affording a clear colorless solution which was stirred overnight. The following day the reaction was determined to be complete by HPLC and TLC and was diluted with water (1 L) and filtered affording **2** as a white solid (69 g, quantitative) which contained a small amount of ethanol. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 11.38 (s, 1H), δ 8.09 (s, 1H), δ 7.79 (d, 2H), δ 7.50 (d, 2H).

Step 2:

[0502] The undried product from the previous step (270 mmol, theoretical) was dissolved in DMF (140 mL) and treated with N-chlorosuccinimide (38 g, 284 mmol) affording a yellow suspension which was stirred at room temperature overnight. The following day the reaction was diluted with water (1 L), filtered, then aspirated to dryness affording desired product **3** as a slightly yellow solid (75.6 g, quantitative) which contained a small amount of DMF. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 7.71 (d, 2H), δ 7.68 (d, 2H).

Step 3:

[0503] Carboxylic acid **4** (40 g, 246 mmol) was suspended in dichloromethane (200 mL) then treated with oxalyl chloride (32 mL, 368 mmol) and 15 drops of DMF. The reaction generated a steady flow of gas and was stirred at room temperature overnight. The following day all solids had dissolved and the reaction was pale brown. A small amount was removed, diluted with methanol and a small amount of triethyl amine, concentrated and examined by NMR which showed only the methyl ester. It was concentrated to a brown oil, the flask was then stoppered and sealed with Teflon tape. No yield calculated. Methyl ester: ¹H NMR (DMSO-*d*₆, 400 MHz): δ 7.58 (d, 1H), δ 6.90 (d, 1H), δ 3.82 (s, 3H).

Step 4:

[0504] Allyl amine (56.8 mL, 91 mmol) was dissolved in 100 mL dichloromethane, cooled to 0°C, then treated with triethylamine (21 mL, 155 mmol) followed by slow addition of acid chloride **5** (15 g, 83 mmol) in a small amount of dichloromethane. After the addition was determined to be complete by TLC it was then partitioned with aqueous NaHCO₃ separated, and the organic layer extracted once again with dichloromethane. The combined organic layers were dried over Na₂SO₄, concentrated then purified by filtration through a short plug of silica gel affording the desired product (**6**) as a light beige crystalline solid (24 g, quant.).


¹H NMR (DMSO-*d*₆, 400 MHz): δ 8.79 (m, 1H), 7.64 (d, 1H), 7.15 (d, 1H), 5.81 (m, 1H), 5.11 (d, 1H), 5.05 (d, 1H), 3.80 (m, 2H).

Step 5:

[0505] Alkene **6** (4.00g, 20 mmol) and triethylamine (3.62 mL, 26 mmol) were heated to 90°C until all solids dissolved. The mixture was then treated with **3** (5.14 g, 22 mmol) slowly over 5 min. After all **3** was added the reaction was checked by TLC and more **3** was added if any starting alkene was present. Upon complete consumption of **6** the reaction mixture was cooled to room temperature, concentrated *in vacuo*, then suspended in a small amount of methanol (ca. 5 mL) and triturated with water affording a white solid. The solid was filtered and dried affording the desired isoxazolidinone (**7**) as a white solid (6.02 g, 75%). ¹H NMR (DMSO-*d*₆, 400 MHz): δ 8.88 (m, 1H), δ 7.89 (d, 2H), δ 7.80 (d, 2H), δ 7.61 (s, 1H), δ 7.14 (s, 1H), δ 4.90 (m, 1H), δ 3.52 (m, 1H), δ 3.40 (t, 1H), δ 3.21 (dd, 1H).

EXAMPLE 2

15 5-Chloro-thiophene-2-carboxylic acid [3-(4-bromo-phenyl)-isoxazol-5-ylmethyl]-amide




[0506] The titled compound was made by the procedure similar to that described in Example 1 using propargyl amine in place of allyl amine. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 9.50 (m, 1H), δ 7.79 (d, 2H), 7.62 (m, 3H), δ 7.18 (d, 1H), δ 6.94 (s, 1H), δ 4.78 (d, 2H).

20

EXAMPLE 3


5-Chloro-thiophene-2-carboxylic acid [3-(3-bromo-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



25 [0507] The titled compound was made by the procedure similar to that described in Example 1. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 8.87 (t, 1H), δ 7.78 (s, 1H), δ 7.63 (m, 3H), δ 7.37 (t, 1H), δ 7.15 (s, 1H), 4.81 δ (m, 1H), 3.47 (dd, 1H), δ 3.39 (t, 1H), δ 3.19 (dd, 1H).

EXAMPLE 4


5-Chloro-thiophene-2-carboxylic acid [3-(3-bromo-phenyl)-isoxazol-5-ylmethyl]-amide



- 5 [0508] The titled compound was made by the procedure similar to that described in Example 2. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 9.30 (t, 1H), δ 8.02 (s, 1H), δ 7.87 (d, 1H), δ 7.62 (m, 3H), δ 7.42 (t, 1H), δ 7.18 (m, 2H), 7.02 (s, 1H), δ 4.59 (d, 2H).

EXAMPLE 5

- 10 5-Chloro-thiophene-2-carboxylic acid [3-(4-phenoxy-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




- [0509] The titled compound was made by the procedure similar to that described in Example 1. MS found for C₂₁H₁₇ClN₂O₃S as (M+H)⁺ 413.5.

15

EXAMPLE 6

5-Chloro-thiophene-2-carboxylic acid [3-(4-phenoxy-phenyl)-isoxazol-5-ylmethyl]-amide




- [0510] The titled compound was made by the procedure similar to that described in Example 2. MS found for C₂₁H₁₅ClN₂O₃S as (M+H)⁺ 411.4, 413.4.

20

EXAMPLE 7

5-Chloro-thiophene-2-carboxylic acid [3-(4-cyano-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0511] The titled compound was made by the procedure similar to that described in Example 1. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz): δ 8.89 (t, 1H), δ 7.88 (d, 2H), δ 7.78 (d, 2H), δ 7.61 (d, 1H), δ 7.12 (d, 1H), δ 7.90 (m, 1H), δ 3.51 (dd, 1H), 3.41 (t, 1H), 3.20 (dd, 1H).

5

EXAMPLE 8


5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



10 [0512] The nitrile from example 7 (1.00 g, 2.9 mmol) and triethylamine (1.62 mL, 11.6 mmol) were combined with 30 mL of 1, 4-dioxane and placed under an atmosphere of H_2S overnight. The next day the crude mixture was evaporated *in vacuo* then diluted with 30 mL of acetone and methyl iodide (1.44 mL, 23.2 mmol), then refluxed until all of the starting thioamide was converted to the imidate (two hours). The reaction mixture was then cooled
15 and concentrated by rotary evaporation affording a thick orange foam. The orange foam was dissolved in 30 mL of THF and a 7.5 mL aliquot was removed and treated with 7.5 mL of a THF solution containing dimethyl amine (0.5 M) and acetic acid (7.5 M). The resulting light yellow mixture was stirred at rt overnight giving clean conversion to the amidine. The reaction mixture was concentrated and purified by prep-HPLC affording the titled compound
20 as a white powder containing TFA. MS found for $\text{C}_{18}\text{H}_{19}\text{ClN}_4\text{O}_2\text{S}$ as $(\text{M}+\text{H})^+$ 391.0, 393.0.

EXAMPLE 9

5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0513] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{19}H_{19}ClN_4O_2S$ as $(M+H)^+$ 403.1, 405.1.

5

EXAMPLE 10

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0514] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{20}H_{21}ClN_4O_2S$ as $(M+H)^+$ 417.1, 419.1.

10

EXAMPLE 11

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0515] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{21}H_{23}ClN_4O_2S$ as $(M+H)^+$ 431.1, 433.1.

EXAMPLE 12

20 1-[[4-(5-[[[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl]-4,5-dihydro-isoxazol-3-yl]-phenyl]-imino-methyl]-piperidine-4-carboxylic acid ethyl ester




[0516] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{24}H_{27}ClN_4O_4S$ as $(M+H)^+$ 503.1, 505.1.

5

EXAMPLE 13

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methyl-N-prop-2-ynyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0517] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{20}H_{19}ClN_4O_2S$ as $(M+H)^+$ 415.1, 417.1.

10

EXAMPLE 14

5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-methoxy-ethyl)-N-methyl-carbamimidoyl]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide




15

[0518] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{20}H_{23}ClN_4O_3S$ as $(M+H)^+$ 435.1, 437.1.

EXAMPLE 15

20 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-furan-2-ylmethyl-N-methyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0519] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{22}H_{21}ClN_4O_3S$ as $(M+H)^+$ 457.1, 459.1.

5

EXAMPLE 16

5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-dimethylamino-ethyl)-N-methyl-carbamimidoyl]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide




[0520] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{21}H_{26}ClN_5O_2S$ as $(M+H)^+$ 448.1, 450.1.

10

EXAMPLE 17

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methyl-N-propyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0521] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{20}H_{23}ClN_4O_2S$ as $(M+H)^+$ 419.1, 421.1.

20

EXAMPLE 18

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-ethyl-N-methyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0522] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{19}H_{21}ClN_4O_2S$ as $(M+H)^+$ 405.1, 407.0

5

EXAMPLE 19

5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-imino-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0523] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{19}H_{19}ClN_4O_2S$ as $(M+H)^+$ 403.1, 405.1.

10

EXAMPLE 20

5-Chloro-thiophene-2-carboxylic acid [3-(4-cyano-phenyl)-isoxazol-5-ylmethyl]-amide




[0524] The titled compound was made by the procedure similar to that described in Example 2. ^1H NMR ($DMSO-d_6$, 400 MHz): δ 9.32 (t, 1H), δ 8.03 (d, 2H), δ 7.91 (d, 2H), δ 7.68 (s, 1H), δ 7.13 (s, 1H), δ 7.05 (s, 1H), δ 4.60 (d, 2H).

15

EXAMPLE 21

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide

20




[0525] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{18}H_{17}ClN_4O_2S$ as $(M+H)^+$ 389.1, 391.1.

5

EXAMPLE 22


5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide



10 [0526] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{19}H_{17}ClN_4O_2S$ as $(M+H)^+$ 401.1, 403.1.

EXAMPLE 23

15 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide




[0527] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{20}H_{19}ClN_4O_2S$ as $(M+H)^+$ 415.1, 417.1.

20

EXAMPLE 24


5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide



- 5 [0528] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{21}H_{21}ClN_4O_2S$ as $(M+H)^+$ 429.1, 431.1.

EXAMPLE 25


- 10 1-{[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-isoxazol-3-yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid ethyl ester



- [0529] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{24}H_{25}ClN_4O_4S$ as $(M+H)^+$ 501.0, 503.1.

EXAMPLE 26


5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-imino-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide



- 20 [0530] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{19}H_{17}ClN_4O_2S$ as $(M+H)^+$ 401.1, 403.1.

EXAMPLE 27


5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-azetidin-1-ylidene-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide



[0531] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{22}H_{22}ClN_4O_2S$ as $(M)^+$ 441.1, 443.1.

EXAMPLE 28


5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-ethyl-N-methyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide



[0532] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{19}H_{19}ClN_4O_2S$ as $(M+H)^+$ 403.1, 405.1.

EXAMPLE 29


5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methyl-N-propyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide



[0533] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{20}H_{21}ClN_4O_2S$ as $(M+H)^+$ 417.1, 419.1.

EXAMPLE 30


5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-dimethylamino-ethyl)-N-methyl-carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-amide



[0534] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{21}H_{24}ClN_5O_2S$ as (M+H)⁺ 446.1, 448.1.

EXAMPLE 31


5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-furan-2-ylmethyl-N-methyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide



[0535] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{22}H_{19}ClN_4O_3S$ as (M+H)⁺ 455.1, 457.1.

EXAMPLE 32


5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-methoxy-ethyl)-N-methyl-carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-amide



[0536] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{20}H_{21}ClN_4O_3S$ as (M+H)⁺ 433.1, 435.1.

EXAMPLE 33


5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methyl-N-prop-2-ynyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide



[0537] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{20}H_{17}ClN_4O_2S$ as (M+H)⁺ 413.1, 415.1.

EXAMPLE 34


5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methylcarbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide



[0538] The titled compound was made by the procedure similar to that described in Example 8, using the nitrile from Example 20. MS found for $C_{17}H_{15}ClN_4O_2S$ as (M+H)⁺ 375.1, 377.1.

EXAMPLE 35


5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide



[0539] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{19}H_{19}ClN_4O_2S$ as (M+H)⁺ 403.1, 405.1.

EXAMPLE 36


5-Chloro-thiophene-2-carboxylic acid methyl-{3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide



[0540] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{20}H_{19}ClN_4O_2S$ as $(M+H)^+$ 415.1, 417.1.

EXAMPLE 37


5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide



[0541] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{21}H_{21}ClN_4O_2S$ as $(M+H)^+$ 429.1, 431.1.

EXAMPLE 38


5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide



[0542] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{22}H_{23}ClN_4O_2S$ as $(M+H)^+$ 443.1, 445.1.

EXAMPLE 39


1-[[4-(5-{{[(5-Chloro-thiophene-2-carbonyl)-methyl-amino]-methyl}-isoxazol-3-yl)-phenyl]-imino-methyl]-piperidine-4-carboxylic acid ethyl ester



5 [0543] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{25}H_{27}ClN_4O_4S$ as (M+H)⁺ 515.1, 517.1.

EXAMPLE 40


5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-imino-methyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide



[0544] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{20}H_{19}ClN_4O_2S$ as (M+H)⁺ 415.1, 417.0.

EXAMPLE 41


5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-azetidin-1-ylidene-methyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide



[0545] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{23}H_{24}ClN_4O_2S$ as (M)⁺ 455.1, 457.1.

EXAMPLE 42


5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-ethyl-N-methyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide



- 5 [0546] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{20}H_{21}ClN_4O_2S$ as $(M+H)^+$ 417.1, 419.1.

EXAMPLE 43


5-Chloro-thiophene-2-carboxylic acid methyl-{3-[4-(N-methyl-N-propyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide



- [0547] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{21}H_{23}ClN_4O_2S$ as $(M+H)^+$ 431.1, 433.1.

EXAMPLE 44


5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-dimethylamino-ethyl)-N-methyl-carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-methyl-amide



- [0548] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{22}H_{26}ClN_5O_2S$ as $(M+H)^+$ 460.1, 462.1.

EXAMPLE 45


1-{{4-(5-{{[(5-Chloro-thiophene-2-carbonyl)-methyl-amino]-methyl}-isoxazol-3-yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid amide



5 [0549] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{23}H_{24}ClN_5O_3S$ as $(M+H)^+$ 486.1, 488.1.

EXAMPLE 46


5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-methoxy-ethyl)-N-methyl-carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-methyl-amide



[0550] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{21}H_{23}ClN_4O_3S$ as $(M+H)^+$ 447.1, 449.1.

EXAMPLE 47


5-Chloro-thiophene-2-carboxylic acid methyl-{3-[4-(N-methyl-N-prop-2-ynyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide



[0551] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{21}H_{19}ClN_4O_2S$ as $(M+H)^+$ 427.0, 429.1.


EXAMPLE 48

5-Chloro-thiophene-2-carboxylic acid methyl- $\{3-[4-(N\text{-methylcarbamimidoyl})\text{-phenyl}]\text{-isoxazol-5-ylmethyl}\}$ -amide



- 5 [0552] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{18}H_{17}ClN_4O_2S$ as $(M+H)^+$ 389.1, 391.1.


SCHEME 2



10

EXAMPLE 49

(R)-5-Chloro-thiophene-2-carboxylic acid (1- $\{3-[4-(N,N\text{-dimethyl-carbamimidoyl})\text{-phenyl}]\text{-isoxazol-5-yl}\}$ -ethyl)-amide



Step 1:

[0553] Methyl ester **8** (4.03 g, 19.9 mmol) was diluted with 40 mL of toluene and cooled to -78°C. Diisobutylaluminum hydride in toluene (26 mL, 40 mmol) was added slowly to the precooled mixture down the side of the flask. The reaction was stirred at -78°C for two hours at which time it was treated slowly with ca. 5 mL of methanol precooled to -78°C (vigorous reaction). The mixture was then warmed to rt and partitioned with saturated sodium potassium tartrate and ethyl acetate. The layers were separated, the organic phase was extracted with ethyl acetate and the combined organic layers were dried over MgSO₄. After filtration and concentration *in vacuo* the crude material was purified by filtration through a short plug of silica gel eluted with dichloromethane affording the aldehyde **9** as a colorless oil (2.23g, 65%). ¹H NMR (DMSO-*d*₆, 400 MHz): δ 9.38 (s, 1H), δ 7.30 (d, 1H), δ 3.79 (m, 1H), δ 1.31 (s, 9H), 1.06 (d, 3H).

Step 2:

[0554] Triphenylphosphine (6.0 g, 23.1 mmol) was dissolved in 30 mL of dichloromethane and cooled to 0°C. To this was added CBr₄ (3.84 g, 11.6 mmol), slowly, resulting in a clear orange solution. After stirring 10 min aldehyde **9** (1.0 g, 5.78 mmol) was added slowly resulting in a dark red, cloudy solution. This was then checked by TLC which showed complete consumption of the starting material and a less polar spot. The reaction was concentrated *in vacuo* then filtered through a short pad of silica gel (eluted with dichloromethane) affording the desired olefin **10** as an off-white crystalline solid (1.25 g, 66%). ¹H NMR (DMSO-*d*₆, 400 MHz): δ 7.36 (d, 1H), δ 5.18 (s, 1H), δ 4.23 (m, 1H), δ 1.35 (s, 9H), δ 1.28 (d, 3H).

Step 3:

[0555] Dibromoolefin **10** (1.25 g, 3.8 mmol) was dissolved with THF (10 mL), cooled to 0°C, then treated with freshly prepared lithium diisopropyl amide (1.7 M, 20 mL) during which time the reaction changed from colorless to dark yellow-orange. The ice bath was removed and the reaction stirred for 10 min. The reaction was then partitioned with 1 M HCl and ethyl acetate, the layers separated and the aqueous phase extracted again with ethyl acetate. The combined organic layers were dried over MgSO₄, filtered, concentrated and purified by silica gel chromatography (dichloromethane) affording the desired alkyne (**11**) as a white solid (0.3652 g, 57%). ¹H NMR (CDCl₃, 400 MHz): δ 4.71 (br s, 1H), δ 4.42 (br s, 1H), δ 2.20 (s, 1H), δ 1.38 (s, 9H), δ 1.32 (d, 3H).

Step 4:

[0556] Performed in the same manner as in step 5, Example 1. NMR (DMSO-*d*₆, 400 MHz): δ 8.01 (d, 2H), δ 7.95 (d, 2H), δ 7.54 (d, 1H), δ 6.91 (s, 1H), δ 4.81 (m, 1H), δ 1.36 (m, 12H).

5 Step 5:

[0557] Boc protected amine **12** was diluted with 4 M HCl/dioxane (10 mL) and stirred for 2 hrs at which time it was determined to be complete by TLC. The reaction was concentrated affording the desired amine as a white solid which was used immediately for the next step.

10 NMR (DMSO-*d*₆, 400 MHz): δ 8.72 (s, 3H), δ 8.06 (d, 2H), δ 8.02 (d, 2H), δ 7.29 (s, 1H), δ 4.73 (m, 1H), δ 1.59 (d, 3H).

Step 6:


[0558] Performed in the same manner as in step 5, Example 1. NMR (DMSO-*d*₆, 400 MHz): δ 9.11 (d, 1H), δ 8.03 (d, 2H), δ 7.95 (d, 2H), δ 7.70 (d, 2H), δ 7.18 (d, 1H), δ 7.51 (s, 1H), δ 5.29 (m, 1H), δ 1.50 (d, 3H).

15 Step 7:

[0559] Performed in the same manner as in Example 8. MS found for C₁₉H₁₉ClN₄O₂S as (M+H)⁺ 403.1, 405.1.

EXAMPLE 50

20 (R)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-yl}-ethyl)-amide




[0560] The titled compound was made by the procedure similar to that described in Example 49. MS found for C₂₀H₁₉ClN₄O₂S as (M+H)⁺ 415.1, 417.1.

25

EXAMPLE 51

(R)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide




[0561] The titled compound was made by the procedure similar to that described in Example 49. MS found for $C_{21}H_{21}ClN_4O_2S$ as $(M+H)^+$ 429.1, 430.0.

5

EXAMPLE 52

(R)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide




[0562] The titled compound was made by the procedure similar to that described in Example 49. MS found $C_{22}H_{23}ClN_4O_2S$ as $(M+H)^+$ 443.1, 445.1.

10

EXAMPLE 53

(R)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide




15

[0563] The titled compound was made by the procedure similar to that described in Example 49. MS found for $C_{19}H_{19}ClN_4O_2S$ as $(M+H)^+$ 403.0, 405.0

EXAMPLE 54

(S)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-yl}-ethyl)-amide

20




[0564] The titled compound was made by the procedure similar to that described in Example 49. MS found for $C_{20}H_{19}ClN_4O_2S$ as $(M+H)^+$ 415.0, 417.0.

5

EXAMPLE 55

(S)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide




[0565] The titled compound was made by the procedure similar to that described in Example 49. MS found for $C_{21}H_{21}ClN_4O_2S$ as $(M+H)^+$ 429.0, 430.0.

10

EXAMPLE 56

(S)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide




15

[0566] The titled compound was made by the procedure similar to that described in Example 49. MS found $C_{22}H_{23}ClN_4O_2S$ as $(M+H)^+$ 443.0, 445.0.

EXAMPLE 57

20

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0567] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{19}H_{21}ClN_4O_2S$ as $(M+H)^+$ 405.1, 407.1

5

EXAMPLE 58

5-Chloro-thiophene-2-carboxylic acid {5-methyl-3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0568] The titled compound was made by the procedure similar to that described in

10 Example 8. MS found for $C_{20}H_{21}ClN_4O_2S$ as $(M+H)^+$ 417.1, 419.1

EXAMPLE 59

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0569] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{21}H_{23}ClN_4O_2S$ as $(M+H)^+$ 431.1, 433.1

EXAMPLE 60

20 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0570] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{22}H_{25}ClN_4O_2S$ as $(M+H)^+$ 445.1, 447.1

5

EXAMPLE 61

1 - {[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-5-methyl-4,5-dihydro-isoxazol-3-yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid ethyl ester




[0571] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{25}H_{29}ClN_4O_4S$ as $(M+H)^+$ 517.1, 519.1

10

EXAMPLE 62

5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-imino-methyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0572] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{20}H_{21}ClN_4O_2S$ as $(M+H)^+$ 418.0, 420.0

EXAMPLE 63

20

5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-azetidin-1-ylidene-methyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0573] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{23}H_{26}ClN_4O_2S$ as $(M+H)^+$ 457.1, 459.1

5

EXAMPLE 64

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-ethyl-N-methyl-carbamimidoyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0574] The titled compound was made by the procedure similar to that described in

10 Example 8. MS found for $C_{20}H_{23}ClN_4O_2S$ as $(M+H)^+$ 419.1, 421.1

EXAMPLE 65

5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-dimethylamino-ethyl)-N-methyl-carbamimidoyl]-phenyl}-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide




15

[0575] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{22}H_{28}ClN_5O_2S$ as $(M+H)^+$ 462.1, 464.1

EXAMPLE 66

20 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-methoxy-ethyl)-N-methyl-carbamimidoyl]-phenyl}-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide




[0576] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{21}H_{25}ClN_4O_3S$ as (M+H)⁺ 449.1, 451.1

5

EXAMPLE 67

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-furan-2-ylmethyl-N-methyl-carbamimidoyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0577] The titled compound was made by the procedure similar to that described in

10 Example 8. MS found for $C_{21}H_{25}ClN_4O_3S$ as (M+H)⁺ 444.1, 446.1

EXAMPLE 68

1-{{4-(5-{{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}}-5-methyl-4,5-dihydro-isoxazol-3-yl)-phenyl}-imino-methyl}-piperidine-4-carboxylic acid amide




15

[0578] The titled compound was made by the procedure similar to that described in

Example 8. MS found for $C_{23}H_{26}ClN_5O_3S$ as (M+H)⁺ 488.1, 490.1

EXAMPLE 69


20 5-Chloro-thiophene-2-carboxylic acid {5-methyl-3-[4-(N-methylcarbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



[0579] The titled compound was made by the procedure similar to that described in Example 8. MS found for $C_{18}H_{19}ClN_4O_2S$ as $(M+H)^+$ 391.1, 393.1

5


SCHEME 3



EXAMPLE 70

5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide

10



Step 1:

[0580] Carboxylic acid **17** (20.18 g, 138 mmol) was diluted with 150 mL of THF and cooled to 0°C. To this stirring solution was added $BH_3 \cdot S(CH_3)_2$ complex (26 mL, 276 mmol)

slowly which resulted in a vigorous reaction and significant gas evolution. The bath was removed and the reaction stirred at rt for 2 hrs at which time TLC and HPLC showed consumption of the starting material. The reaction was quenched by slow addition of water (vigorous exotherm), then the crude mixture was concentrated, diluted with ethyl acetate and
5 filtered. The filtrate was concentrated to dryness affording the desired alcohol (**18**) as a white solid (17 g, 93%). NMR (DMSO-*d*₆, 400 MHz): δ 7.76 (d, 1H), δ 7.67 (m, 3H), δ 5.50 (t, 1H), δ 4.55 (d, 2H).

Step 2:

[0581] Benzyl alcohol **18** (17 g, 129 mmol) was dissolved in 300 mL of dichloromethane
10 then treated with 33 g of celite followed by PCC (33 g, 155 mmol) resulting in a dark brown suspension which was stirred vigorously. After stirring 3 hrs the reaction was complete by TLC and was filtered through a short pad (2 in. x 5 in.) of silica gel and eluted slowly with dichloromethane. The filtrate was concentrated affording the desired aldehyde (**19**) as a white solid (15.62 g, 93%). NMR (DMSO-*d*₆, 400 MHz): δ 10.38 (s, 1H), δ 7.95 (t, 1H), δ
15 7.59 (d, 1H), δ 7.48 (d, 1H).

Step 3:

[0582] Performed in the same manner as in step 1, Example 1. NMR (DMSO-*d*₆, 400 MHz): δ 12.03 (s, 1H), δ 8.20 (s, 1H), δ 7.90 (m, 2H), δ 7.64 (d, 1H).

Step 4:

20 [0583] Performed in the same manner as in step 2, Example 1. NMR (DMSO-*d*₆, 400 MHz): δ 8.00 (d, 1H), 7.82 (m, 3H).

Step 5:


[0584] Performed in the same manner as in step 5, Example 1. NMR (DMSO-*d*₆, 400 MHz): δ 8.85 (distorter t, 1H), δ 7.93 (d, 1H), δ 7.85 (t, 1H), δ 7.70 (d, 1H), δ 7.60 (s, 1H), δ
25 7.11 (s, 1H), 4.89 (m, 1H), δ 3.20 – 3.55 (m, 4H).

Step 6:

[0585] Performed in the same manner as in Example 8. MS found C₁₉H₁₆ClFN₄O₂S as (M+H)⁺ 419.1, 421.1.

EXAMPLE 71


5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide



- 5 [0586] The titled compound was made by the procedure similar to that described in Example 70. MS found for $C_{20}H_{18}ClFN_4O_2S$ as (M+H)⁺ 433.0, 435.0

EXAMPLE 72


5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide



- [0587] The titled compound was made by the procedure similar to that described in Example 70. MS found for $C_{21}H_{20}ClFN_4O_2S$ as (M+H)⁺ 447.0, 449.0

EXAMPLE 73


5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-fluoro-phenyl]-dihydro-isoxazol-5-ylmethyl}-amide



- [0588] The titled compound was made by the procedure similar to that described in Example 70. MS found for $C_{18}H_{16}ClFN_4O_2S$ as (M+H)⁺ 407.0, 409.0

EXAMPLE 74

5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(N,N-dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




- 5 [0589] A solution of nitrile (Example 70, step 5) (0.090 g, 0.248 mmol) and dimethylamine (0.28 ml, 2.0M solution in tetrahydrofuran) in dimethylsulfoxide (5 mL) was treated with diisopropylethylamine (0.08ml, 0.58 mmol) in a flask equipped with a reflux condenser. The mixture was heated to reflux overnight. The following day the reaction was checked by TLC which showed complete consumption of the starting material and a new polar spot. The
- 10 crude reaction mixture was loaded onto a silica gel plug and eluted with dichloromethane affording the substituted dimethylamino compound as the major product. The product was treated with the procedure from Example 8 to obtain the titled compound. MS found for $C_{20}H_{24}ClN_5O_2S$ as (M+H)⁺ 434.1, 436.0

15

EXAMPLE 75


5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- [0590] The titled compound was made by the procedure similar to that described in
- 20 Example 74. MS found for $C_{21}H_{24}ClN_5O_2S$ as (M+H)⁺ 446.1, 448.1

EXAMPLE 76

5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0591] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{22}H_{26}ClN_5O_2S$ as (M+H)⁺ 460.1, 462.1

5

EXAMPLE 77

5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(imino-piperidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0592] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{23}H_{28}ClN_5O_2S$ as (M+H)⁺ 474.1, 476.1

10

EXAMPLE 78

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-pyrrolidin-1-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0593] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{22}H_{26}ClN_5O_2S$ as (M+H)⁺ 460.1, 462.1

EXAMPLE 79

20 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-pyrrolidin-1-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0594] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{23}H_{26}ClN_5O_2S$ as (M+H)⁺ 472.1, 474.1

5

EXAMPLE 80

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-pyrrolidin-1-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0595] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{24}H_{28}ClN_5O_2S$ as (M+H)⁺ 486.1, 488.1

10

EXAMPLE 81

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-pyrrolidin-1-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0596] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{25}H_{30}ClN_5O_2S$ as (M+H)⁺ 500.1, 502.1

EXAMPLE 82

20 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-morpholin-4-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0597] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{22}H_{26}ClN_5O_3S$ as (M+H)⁺ 474.1, 476.1

5

EXAMPLE 83

5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-morpholin-4-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0598] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{23}H_{26}ClN_5O_3S$ as (M+H)⁺ 486.1, 488.1

10

EXAMPLE 84

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-morpholin-4-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0599] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{24}H_{28}ClN_5O_3S$ as (M+H)⁺ 500.1, 502.1

EXAMPLE 85

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-morpholin-4-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide

20




[0600] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{25}H_{30}ClN_5O_3S$ as (M+H)⁺ 514.1, 516.1

5

EXAMPLE 86

5-Chloro-thiophene-2-carboxylic acid (3-{4-(N,N-dimethyl-carbamimidoyl)-2-[(2-methoxy-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide




[0601] The titled compound was made by the procedure similar to that described in

10 Example 74. MS found for $C_{22}H_{28}ClN_5O_3S$ as (M+H)⁺ 478.1, 480.1

EXAMPLE 87

5-Chloro-thiophene-2-carboxylic acid {3-[2-[(2-methoxy-ethyl)-methyl-amino]-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0602] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{23}H_{28}ClN_5O_3S$ as (M+H)⁺ 490.1, 492.1.


EXAMPLE 88

20 5-Chloro-thiophene-2-carboxylic acid (3-{4-(imino-pyrrolidin-1-yl-methyl)-2-[(2-methoxy-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide



[0603] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{24}H_{30}ClN_5O_3S$ as (M+H)⁺ 504.1, 506.1


5
EXAMPLE 89
5-Chloro-thiophene-2-carboxylic acid (3-{4-(imino-piperidin-1-yl-methyl)-2-[(2-methoxy-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide



[0604] The titled compound was made by the procedure similar to that described in
10 Example 74. MS found for $C_{25}H_{32}ClN_5O_3S$ as (M+H)⁺ 518.1, 520.2

EXAMPLE 90


5-Chloro-thiophene-2-carboxylic acid {3-[2-azepan-1-yl-4-(N,N-dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



15
[0605] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{24}H_{30}ClN_5O_2S$ as (M+H)⁺ 489.2, 491.2

EXAMPLE 91

20
5-Chloro-thiophene-2-carboxylic acid {3-[2-azepan-1-yl-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0606] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{25}H_{30}ClN_5O_2S$ as (M+H)⁺ 501.1, 503.1

5

EXAMPLE 92

5-Chloro-thiophene-2-carboxylic acid {3-[2-azepan-1-yl-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




10 [0607] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{26}H_{32}ClN_5O_2S$ as (M+H)⁺ 515.1, 517.1

EXAMPLE 93

15


5-Chloro-thiophene-2-carboxylic acid {3-[2-azepan-1-yl-4-(imino-piperidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



[0608] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{27}H_{34}ClN_5O_2S$ as (M+H)⁺ 529.2, 531.2

EXAMPLE 94


5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(N,N-dimethyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide



- 5 [0609] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{20}H_{22}ClN_5O_2S$ as (M+H)⁺ 432.1, 434.1

EXAMPLE 95


5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide



- [0610] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{21}H_{22}ClN_5O_2S$ as (M+H)⁺ 444.1, 446.1

EXAMPLE 96


5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide



- [0611] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{22}H_{24}ClN_5O_2S$ as (M+H)⁺ 458.1, 460.1

EXAMPLE 97


5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide



- 5 [0612] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{23}H_{26}ClN_5O_2S$ as $(M+H)^+$ 472.1, 474.1

EXAMPLE 98


5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-morpholin-4-yl-phenyl]-isoxazol-5-ylmethyl}-amide



- [0613] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{22}H_{24}ClN_5O_3S$ as $(M+H)^+$ 474.1, 476.1

EXAMPLE 99


5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-morpholin-4-yl-phenyl]-isoxazol-5-ylmethyl}-amide



- [0614] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{23}H_{24}ClN_5O_3S$ as $(M+H)^+$ 486.1, 488.1

EXAMPLE 100


5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-morpholin-4-yl-phenyl]-isoxazol-5-ylmethyl}-amide



- 5 [0615] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{24}H_{26}ClN_5O_3S$ as $(M+H)^+$ 500.1, 502.1

EXAMPLE 101


5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-morpholin-4-yl-phenyl]-isoxazol-5-ylmethyl}-amide



- [0616] The titled compound was made by the procedure similar to that described in Example 74. MS found for $C_{25}H_{28}ClN_5O_3S$ as $(M+H)^+$ 514.1, 516.1

EXAMPLE 102

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-methoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




- [0617] A solution of the nitrile compound (Example 70, step 5) (0.264 g, 0.727 mmol) in dimethylsulfoxide was treated with sodium methoxide (0.086 g, 1.60 mmol). The mixture was stirred at room temperature for 2 hours and was checked by TLC which showed complete consumption of the starting material and a new polar spot. The crude reaction

mixture was loaded onto a silica gel plug and eluted with dichloromethane affording the substituted methoxy compound as the major product. The product was treated with the procedure similar to that described in Example 8 to obtain the titled compound. MS found for C₁₉H₂₁ClN₄O₃S as (M+H)⁺ 421.0, 423.0.

5

EXAMPLE 103


5-Chloro-thiophene-2-carboxylic acid {3-[2-methoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



10 [0618] The titled compound was made by the procedure similar to that described in Example 102. MS found for C₂₀H₂₁ClN₄O₃S as (M+H)⁺ 433.0, 435.0

EXAMPLE 104

15 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-methoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0619] The titled compound was made by the procedure similar to that described in Example 102. MS found for C₂₁H₂₃ClN₄O₃S as (M+H)⁺ 447.1, 449.1

20

EXAMPLE 105

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-methoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0620] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{22}H_{25}ClN_4O_3S$ as (M+H)⁺ 461.1, 463.1

5

EXAMPLE 106

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-ethoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0621] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{20}H_{23}ClN_4O_3S$ as (M+H)⁺ 435.1, 437.1

10

EXAMPLE 107

5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0622] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{21}H_{23}ClN_4O_3S$ as (M+H)⁺ 447.1, 449.1

EXAMPLE 108

5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide

20




[0623] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{22}H_{25}ClN_4O_3S$ as $(M+H)^+$ 461.1, 463.1

5

EXAMPLE 109

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-isopropoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0624] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{21}H_{25}ClN_4O_3S$ as $(M+H)^+$ 449.1, 451.1

10

EXAMPLE 110

5-Chloro-thiophene-2-carboxylic acid {3-[2-isopropoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0625] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{22}H_{25}ClN_4O_3S$ as $(M+H)^+$ 461.1, 463.1

EXAMPLE 111

20 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-isopropoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0626] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{23}H_{27}ClN_4O_3S$ as (M+H)⁺ 475.1, 477.1

5

EXAMPLE 112


5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclopentyloxy-4-(N,N-dimethylcarbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



[0627] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{23}H_{27}ClN_4O_3S$ as (M+H)⁺ 475.5, 477.5

EXAMPLE 113

5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclopentyloxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0628] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{24}H_{27}ClN_4O_3S$ as (M+H)⁺ 487.5, 489.5

EXAMPLE 114

20 5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclopentyloxy-4-(imino-pyrrolidin-1-ylmethyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0629] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{25}H_{29}ClN_4O_3S$ as $(M+H)^+$ 501.6, 503.6

5

EXAMPLE 115

5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclohexyloxy-4-(N,N-dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0630] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{24}H_{29}ClN_4O_3S$ as $(M+H)^+$ 489.6, 491.6

10

EXAMPLE 116

5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclohexyloxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0631] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{25}H_{29}ClN_4O_3S$ as $(M+H)^+$ 501.6, 503.6

EXAMPLE 117

5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclohexyloxy-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide

20




[0632] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{26}H_{31}ClN_4O_3S$ as $(M+H)^+$ 515.6, 517.6

5

EXAMPLE 118

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-methoxy-phenyl]-isoxazol-5-ylmethyl}-amide




[0633] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{19}H_{19}ClN_4O_3S$ as $(M+H)^+$ 419.0, 421.0

10

EXAMPLE 119

5-Chloro-thiophene-2-carboxylic acid {3-[2-methoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide




15

[0634] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{20}H_{19}ClN_4O_3S$ as $(M+H)^+$ 431.0, 433.0

EXAMPLE 120

20 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-methoxy-phenyl]-isoxazol-5-ylmethyl}-amide




[0635] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{21}H_{21}ClN_4O_3S$ as $(M+H)^+$ 445.0, 447.1

5

EXAMPLE 121

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-methoxy-phenyl]-isoxazol-5-ylmethyl}-amide




[0636] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{22}H_{23}ClN_4O_3S$ as $(M+H)^+$ 459.1, 461.1

10

EXAMPLE 122

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-ethoxy-phenyl]-isoxazol-5-ylmethyl}-amide




15

[0637] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{20}H_{21}ClN_4O_3S$ as $(M+H)^+$ 433.1, 435.1

EXAMPLE 123

5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide

20




[0638] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{21}H_{21}ClN_4O_3S$ as $(M+H)^+$ 445.1, 447.1

5

EXAMPLE 124

5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide




[0639] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{22}H_{23}ClN_4O_3S$ as $(M+H)^+$ 459.1, 461.1

10

EXAMPLE 125

5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide




15

[0640] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{23}H_{25}ClN_4O_3S$ as $(M+H)^+$ 473.1, 475.1

EXAMPLE 126

20 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-isopropoxy-phenyl]-isoxazol-5-ylmethyl}-amide




[0641] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{23}H_{25}ClN_4O_3S$ as $(M+H)^+$ 447.1, 449.1

5

EXAMPLE 127

5-Chloro-thiophene-2-carboxylic acid {3-[2-isopropoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide




[0642] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{22}H_{23}ClN_4O_3S$ as $(M+H)^+$ 459.1, 461.1

10

EXAMPLE 128

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-isopropoxy-phenyl]-isoxazol-5-ylmethyl}-amide




15

[0643] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{23}H_{25}ClN_4O_3S$ as $(M+H)^+$ 473.1, 475.1

EXAMPLE 129

20 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-isopropoxy-phenyl]-isoxazol-5-ylmethyl}-amide




[0644] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{24}H_{27}ClN_4O_3S$ as $(M+H)^+$ 487.1, 489.1

5

EXAMPLE 130

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-phenoxy-phenyl]-isoxazol-5-ylmethyl}-amide




[0645] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{24}H_{21}ClN_4O_3S$ as $(M+H)^+$ 511.1, 513.1

10

EXAMPLE 131

5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-phenoxy-phenyl]-isoxazol-5-ylmethyl}-amide




15

[0646] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{24}H_{21}ClN_4O_3S$ as $(M+H)^+$ 523.1, 525.1

EXAMPLE 132

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-phenoxy-phenyl]-isoxazol-5-ylmethyl}-amide

20




[0647] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{26}H_{23}ClN_4O_3S$ as $(M+H)^+$ 537.1, 539.1

5

EXAMPLE 133

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-(4-methoxy-phenoxy)-phenyl]-isoxazol-5-ylmethyl}-amide




[0648] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{25}H_{23}ClN_4O_4S$ as $(M+H)^+$ 481.1, 483.1

10

EXAMPLE 134

5-Chloro-thiophene-2-carboxylic acid {3-[2-(4-methoxy-phenoxy)-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide




15

[0649] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{26}H_{23}ClN_4O_4S$ as $(M+H)^+$ 493.1, 495.1

EXAMPLE 135

20 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-(4-methoxy-phenoxy)-phenyl]-isoxazol-5-ylmethyl}-amide




[0650] The titled compound was made by the procedure similar to that described in Example 102. MS found for $C_{27}H_{25}ClN_4O_4S$ as $(M+H)^+$ 507.1, 509.1

5

EXAMPLE 136

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-methylsulfanyl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0651] The titled compound was made by the procedure similar to that described in Example 102 using sodium thiomethoxide in place of sodium methoxide. MS found for $C_{19}H_{21}ClN_4O_2S_2$ as $(M+H)^+$ 437.0, 439.0

15

EXAMPLE 137

5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-methylsulfanyl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0652] The titled compound was made by the procedure similar to that described in Example 136. MS found for $C_{20}H_{21}ClN_4O_2S_2$ as $(M+H)^+$ 449.0, 451.0

20

EXAMPLE 138

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-methylsulfanyl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0653] The titled compound was made by the procedure similar to that described in Example 136. MS found for $C_{21}H_{23}ClN_4O_2S_2$ as (M+H)⁺ 463.0, 465.1

5

EXAMPLE 139

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-methylsulfonyl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0654] The titled compound was made by the procedure similar to that described in Example 136. MS found for $C_{22}H_{25}ClN_4O_2S_2$ as (M+H)⁺ 477.0, 479.1

10

EXAMPLE 140

5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-methylsulfonyl-phenyl]-isoxazol-5-ylmethyl}-amide




15

[0655] The titled compound was made by the procedure similar to that described in Example 136. MS found for $C_{19}H_{19}ClN_4O_2S_2$ as (M+H)⁺ 435.0, 437.0

EXAMPLE 141

20 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-methylsulfonyl-phenyl]-isoxazol-5-ylmethyl}-amide




[0656] The titled compound was made by the procedure similar to that described in Example 136. MS found for $C_{20}H_{19}ClN_4O_2S_2$ as (M+H)⁺ 447.0, 449.0

5

EXAMPLE 142

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-methylsulfanyl-phenyl]-isoxazol-5-ylmethyl}-amide




[0657] The titled compound was made by the procedure similar to that described in Example 136. MS found for $C_{21}H_{21}ClN_4O_2S_2$ as (M+H)⁺ 461.0, 463.0

10

EXAMPLE 143

5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-methylsulfanyl-phenyl]-isoxazol-5-ylmethyl}-amide

15




[0658] The titled compound was made by the procedure similar to that described in Example 136. MS found for $C_{22}H_{23}ClN_4O_2S_2$ as (M+H)⁺ 475.1, 477.0

20

EXAMPLE 144

5-Chloro-thiophene-2-carboxylic acid (3-phenyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide




[0659] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{15}H_{13}ClN_2O_2S$ as $(M+H)^+$ 321.3, 323.3

5

EXAMPLE 145

5-Chloro-thiophene-2-carboxylic acid (5-methyl-3-phenyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide




[0660] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{16}H_{15}ClN_2O_2S$ as $(M+H)^+$ 335.4, 337.3

10

EXAMPLE 146

5-Chloro-thiophene-2-carboxylic acid (3-phenyl-isoxazol-5-ylmethyl)-amide




[0661] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{15}H_{11}ClN_2O_2S$ as $(M+H)^+$ 319.2, 321.1

15

EXAMPLE 147

5-Chloro-thiophene-2-carboxylic acid (3-p-tolyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide




20

[0662] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{16}H_{15}ClN_2O_2S$ as $(M+H)^+$ 335.2, 337.2

EXAMPLE 148


5-Chloro-thiophene-2-carboxylic acid (5-methyl-3-p-tolyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide



[0663] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{17}H_{17}ClN_2O_2S$ as $(M+H)^+$ 349.2, 351.2

EXAMPLE 149


5-Chloro-thiophene-2-carboxylic acid (3-p-tolyl-isoxazol-5-ylmethyl)-amide



[0664] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{16}H_{13}ClN_2O_2S$ as $(M+H)^+$ 333.2, 335.1

EXAMPLE 150


5-Chloro-thiophene-2-carboxylic acid [3-(4-methoxy-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



[0665] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{16}H_{15}ClN_2O_3S$ as $(M+H)^+$ 379.2, 381.2

EXAMPLE 151

5-Chloro-thiophene-2-carboxylic acid [3-(4-methoxy-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0666] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{17}H_{17}ClN_2O_3S$ as $(M+H)^+$ 393.2, 395.1

5

EXAMPLE 152

5-Chloro-thiophene-2-carboxylic acid [3-(4-methoxy-phenyl)-isoxazol-5-ylmethyl]-amide




[0667] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{16}H_{13}ClN_2O_3S$ as $(M+H)^+$ 377.3, 379.3

10

EXAMPLE 153

5-Chloro-thiophene-2-carboxylic acid [3-(4-chloro-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




15

[0668] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{15}H_{12}Cl_2N_2O_2S$ as $(M+H)^+$ 356.7, 358.7

EXAMPLE 154

5-Chloro-thiophene-2-carboxylic acid [3-(4-chloro-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl]-amide


20



[0669] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{16}H_{14}Cl_2N_2O_2S$ as (M+H)⁺ 370.7, 372.8

EXAMPLE 155

5 5-Chloro-thiophene-2-carboxylic acid [3-(4-chloro-phenyl)-isoxazol-5-ylmethyl]-amide




[0670] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{15}H_{10}Cl_2N_2O_2S$ as (M+H)⁺ 354.8, 356.7

10

EXAMPLE 156

4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-benzoic acid methyl ester




[0671] The titled compound was made by the procedure similar to that described in

15 Example 1. MS found for $C_{17}H_{15}ClN_2O_4S$ as (M+H)⁺ 379.4, 381.4

EXAMPLE 157

4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-isoxazol-3-yl)-benzoic acid methyl ester




20

[0672] The titled compound was made by the procedure similar to that described in Example 1. MS found for $C_{17}H_{13}ClN_2O_4S$ as (M+H)⁺ 377.3, 379.3

EXAMPLE 158


4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-benzoic acid



- 5 [0673] A solution of the compound 156 (0.360 g, 0.952 mmol) in methanol and water was treated with lithium hydroxide (2.86 ml, 1M in methanol). The mixture was stirred at room temperature overnight. The following day the reaction was checked by TLC which showed complete consumption of the starting material and a new, more polar spot. The crude reaction mixture was acidified to pH=3, concentrated *in vacuo*, then suspended in a small amount of methanol (ca. 5 mL) and triturated with water affording a white solid. The solid was filtered and dried affording the desired product. MS found for C₁₆H₁₃ClN₂O₄S as (M+H)⁺ 365.3, 367.3

EXAMPLE 159

15 4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-isoxazol-3-yl)-benzoic acid




[0674] The titled compound was made by the procedure similar to that described in Example 158. MS found for C₁₆H₁₁ClN₂O₄S as (M+H)⁺ 363.4, 365.4

20

EXAMPLE 160


5-Chloro-thiophene-2-carboxylic acid [3-(4-amino-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



[0675] A solution of the nitroisoxazoline (prepared in a similar manner to Example 1 from 4-nitrobenzaldehyde) (0.078 g, 0.214 mmol) in ethanol was treated with tin chloride dihydrate (0.155 g, 0.684 mmol). The mixture was heated to reflux for 2 hour. The reaction was checked by TLC which showed complete consumption of the starting material and a new, less polar spot. The crude reaction mixture was diluted with saturated potassium carbonate, stirred ca. 30 min, then filtered through a pad of Celite. The filtrate was concentrated affording the desired product which was purified by preparative HPLC. MS found for $C_{15}H_{14}ClN_3O_2S$ as $(M+H)^+$ 336.3, 338.3

EXAMPLE 161


5-Chloro-thiophene-2-carboxylic acid [3-(4-amino-phenyl)-isoxazol-5-ylmethyl]-amide



[0676] The titled compound was made by the procedure similar to that described in Example 160. MS found for $C_{15}H_{12}ClN_3O_2S$ as $(M+H)^+$ 334.3, 336.3

EXAMPLE 162


5-Chloro-thiophene-2-carboxylic acid [3-(4-dimethylcarbamoyl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



[0677] A solution of the acid compound 158 (0.088 g, 0.243 mmol), dimethylamine (0.032ml, 0.316 mmol) and triethylamine (0.051 ml, 0.365 mmol) in dimethylformide was treated with BOP chloride (0.074 g, 0.292 mmol). The mixture was stirred at room temperature overnight. The following day the reaction was checked by TLC which showed complete consumption of the starting material and a new, less polar spot. The crude reaction mixture was purified by preparative HPLC to give the desired product. MS found for $C_{18}H_{18}ClN_3O_3S$ as $(M+H)^+$ 392.5, 394.5

EXAMPLE 163


5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidine-1-carbonyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



[0678] The titled compound was made by the procedure similar to that described in Example 162. MS found for $C_{19}H_{18}ClN_3O_3S$ as $(M+H)^+$ 404.3, 406.3

EXAMPLE 164


5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyrrolidine-1-carbonyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



[0679] The titled compound was made by the procedure similar to that described in Example 162. MS found for $C_{20}H_{20}ClN_3O_3S$ as $(M+H)^+$ 418.0, 420.0

EXAMPLE 165


5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-methyl-pyrrolidine-1-carbonyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



[0680] The titled compound was made by the procedure similar to that described in Example 162. MS found for $C_{21}H_{22}ClN_3O_3S$ as $(M+H)^+$ 432.3, 434.3

EXAMPLE 166


5-Chloro-thiophene-2-carboxylic acid (3-{4-[(2-diethylamino-ethyl)-methyl-carbamoyl]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide



- 5 [0681] The titled compound was made by the procedure similar to that described in Example 162. MS found for $C_{23}H_{29}ClN_4O_3S$ as $(M+H)^+$ 477.2, 479.1

EXAMPLE 167


10 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyrrolidine-1-carbonyl)-phenyl]-isoxazol-5-ylmethyl}-amide



- [0682] The titled compound was made by the procedure similar to that described in Example 162. MS found for $C_{20}H_{18}ClN_3O_3S$ as $(M+H)^+$ 416.1, 418.0


EXAMPLE 168

15 5-Chloro-thiophene-2-carboxylic acid (3-{4-[(2-diethylamino-ethyl)-methyl-carbamoyl]-phenyl}-isoxazol-5-ylmethyl)-amide




- 20 [0683] The titled compound was made by the procedure similar to that described in Example 162. MS found for $C_{20}H_{18}ClN_3O_3S$ as $(M+H)^+$ 475.1, 477.1

SCHEME 4



EXAMPLE 169

- 5 5-Chloro-thiophene-2-carboxylic acid [3-(1-ethyl-piperidin-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide (31).



Step 1:

- [0684] Ethyl isonepecotate, **24** (9.8 mL, 64 mmol), was diluted with dichloromethane (100 mL) and treated with *t*-butylpyrocarbonate (16 mL, 70 mmol) which resulted in significant gas evolution. After 20 min the clear solution was checked by TLC (dichloromethane) which showed complete consumption of the starting amine. The reaction mixture was concentrated affording the protected amine (**25**) as a colorless oil containing a small amount of *t*-butanol. NMR (DMSO-*d*₆, 400 MHz): δ 4.01 (m, 1H), δ 3.75 (m, 2H), δ 3.28 (m, 1H), δ 2.75 (m, 2H), δ 2.45 (m, 2H), δ 1.69 (m, 2H), δ 1.31 (m, 11H), 1.09 (m, 3H).

Step 2:

[0685] Performed in the same manner as in step 1, Example 49. (DMSO-*d*₆, 400 MHz): δ 9.52 (s, 1H), δ 3.72 (m, 2H), 2.81 (m, 2H), δ 1.73 (m, 3H), δ 1.31 (m, 13H).

5 Step 3:

[0686] Performed in the same manner as in step 1, Example 1. (DMSO-*d*₆, 400 MHz): δ 8.88 (d, 0.66H), δ 8.50 (t, 0.33H), δ 8.00 (t, 0.66H), δ 7.20 (d, 0.33H), δ 3.83 (m, 2H), 2.73 (m, 2H), δ 2.26 (m, 1H), δ 1.61 (m, 2H), δ 1.32 (s, 9H), δ 1.10 (m, 2H).

Step 4:

10 [0687] Performed in the same manner as in step 2, Example 1. The material was used immediately for step 5 without purification or characterization.

Step 5:

15 [0688] Performed in the same manner as in step 5, Example 1. (DMSO-*d*₆, 400 MHz): δ 8.78 (m, 1H), δ 7.62 (s, 1H), δ 7.18 (s, 1H), δ 4.58 (m, 1H), 3.82 (m, 2H), δ 3.26 (m, 2H), δ 2.99 (m, 1H), δ 2.72 (m, 2H), δ 2.46 (m, 2H), δ 1.69 (m, 2H), δ 1.36 (m, 11H).

Step 6:


20 [0689] Performed in the same manner as in step 5, Example 49. MS found for C₁₄H₁₈ClN₃O₂S as (M+H)⁺ 328.1, 330.1.

Step 7:

[0690] Amine hydrochloride **30** (40 mg, 0.11 mmol) was diluted with 4 mL dichloroethane then treated with 5 drops of acetaldehyde, triethylamine (18 μ L, 0.13 mmol) and
25 Na(OAc)₃BH (47 mg, 0.22 mmol). The resulting slightly cloudy mixture was stirred overnight. The next day the reaction was checked by HPLC and determined to be complete. It was then concentrated and purified by preparative HPLC affording the desired tertiary amine salt as a white solid. MS found for C₁₆H₂₂ClN₃O₂S as (M+H)⁺ 356.4, 358.4.

EXAMPLE 170


5-Chloro-thiophene-2-carboxylic acid [3-(1-benzyl-piperidin-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



5 [0691] The titled compound was made by the procedure similar to that described in Example 169. MS found for $C_{21}H_{24}ClN_3O_2S$ as $(M+H)^+$ 418.6, 420.6.

EXAMPLE 171


5-Chloro-thiophene-2-carboxylic acid [3-(1-isopropyl-piperidin-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



[0692] The titled compound was made by the procedure similar to that described in Example 169. MS found for $C_{17}H_{24}ClN_3O_2S$ as $(M+H)^+$ 370.6, 372.5.

EXAMPLE 172


5-Chloro-thiophene-2-carboxylic acid [3-(1-carbamimidoyl-piperidin-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



[0693] Amine 30 (40 mg, 0.11 mmol) was dissolved in methanol (5 mL) then treated with triethyl amine (46 μ L, 0.33 mmol) and 1H-pyrazole-1-carboxamide (32 mg, 0.22 mmol) overnight. The next day the reaction was concentrated and the crude reaction mixture purified by reverse phase HPLC affording the desired guanidine as a white powder. MS found for $C_{15}H_{20}ClN_5O_2S$ as $(M+H)^+$ 370.6, 372.4.

EXAMPLE 173


5-Chloro-thiophene-2-carboxylic acid {3-[1-(1-imino-ethyl)-piperidin-4-yl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



[0694] Combined amine **30** (40 mg, 0.11 mmol), triethylamine (46 μ L, 0.33 mmol) ethyl acetimidate – hydrochloride (27 mg, 0.22 mmol) and ethanol (5 mL) then stirred the resulting cloudy white solution overnight. The following day the reaction was found to be done by HPLC and was concentrated and purified by preparative HPLC affording the desired amidine as a TFA salt. MS found for $C_{16}H_{21}ClN_4O_2S$ as $(M+H)^+$ 369.5, 371.5.

EXAMPLE 174


5-Chloro-thiophene-2-carboxylic acid {3-[1-(imino-phenyl-methyl)-piperidin-4-yl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



[0695] The titled compound was made by the procedure similar to that described in Example 173. MS found for $C_{21}H_{23}ClN_4O_2S$ as $(M+H)^+$ 431.5, 433.5.

EXAMPLE 175


5-Chloro-thiophene-2-carboxylic acid {3-[1-(imino-pyridin-2-yl-methyl)-piperidin-4-yl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



[0696] The titled compound was made by the procedure similar to that described in Example 173. MS found for $C_{20}H_{22}ClN_5O_2S$ as $(M+H)^+$ 432.5, 434.5.

EXAMPLE 176


5-Chloro-thiophene-2-carboxylic acid (3-piperidin-4-yl-isoxazol-5-ylmethyl)-amide



[0697] The titled compound was made by the procedure similar to that described in Example 169, step 6. MS found for $C_{14}H_{16}ClN_3O_2S$ as $(M+H)^+$ 326.0, 328.0.

EXAMPLE 177


5-Chloro-thiophene-2-carboxylic acid [3-(1-ethyl-piperidin-4-yl)-isoxazol-5-ylmethyl]-amide



[0698] The titled compound was made by the procedure similar to that described in Example 169. MS found for $C_{16}H_{20}ClN_3O_2S$ as $(M+H)^+$ 354.1, 356.1.

EXAMPLE 178


5-Chloro-thiophene-2-carboxylic acid [3-(1-benzyl-piperidin-4-yl)-isoxazol-5-ylmethyl]-amide



[0699] The titled compound was made by the procedure similar to that described in Example 169. MS found for $C_{21}H_{22}ClN_3O_2S$ as $(M+H)^+$ 416.1, 418.1.

EXAMPLE 179

5-Chloro-thiophene-2-carboxylic acid [3-(1-carbamimidoyl-piperidin-4-yl)-isoxazol-5-ylmethyl]-amide




[0700] The titled compound was made by the procedure similar to that described in Example 172. MS found for $C_{15}H_{18}ClN_5O_2S$ as $(M+H)^+$ 368.0, 370.1.

5

EXAMPLE 180

5-Chloro-thiophene-2-carboxylic acid {3-[1-(1-imino-ethyl)-piperidin-4-yl]-isoxazol-5-ylmethyl}-amide




[0701] The titled compound was made by the procedure similar to that described in Example 173. MS found for $C_{16}H_{19}ClN_4O_2S$ as $(M+H)^+$ 367.1, 369.1.

10

EXAMPLE 181

5-Chloro-thiophene-2-carboxylic acid {3-[1-(imino-phenyl-methyl)-piperidin-4-yl]-isoxazol-5-ylmethyl}-amide




15

[0702] The titled compound was made by the procedure similar to that described in Example 173. MS found for $C_{21}H_{21}ClN_4O_2S$ as $(M+H)^+$ 429.1, 431.1

EXAMPLE 182

20 5-Chloro-thiophene-2-carboxylic acid [3-(1-carbamimidoyl-piperidin-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0703] The titled compound was made by the procedure similar to that described in Example 173. MS found for $C_{15}H_{18}ClN_5O_2S$ as $(M+H)^+$ 368.1, 370.1.

5

EXAMPLE 183

5-Chloro-thiophene-2-carboxylic acid [3-(2'-formyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0704] Aryl bromide (Example 1, 200 mg, 0.50 mmol), 2-formylphenylboronic acid (90 mg, 0.60 mmol) and K_2CO_3 (106 mg, 1.0 mmol) were diluted with diglyme (5 mL) and water (1 mL). The mixture was degassed for 3 – 5 min, then treated with $Pd(PPh_3)_4$ (58 mg, 0.05 mmol) and heated to 120 °C until all starting bromide was consumed as determined by HPLC. The resulting mixture was diluted with water, filtered and the remaining solids purified by silica gel chromatography (0 – 10% Ethyl acetate / dichloromethane) affording the desired biphenyl aldehyde as a white solid. MS found for $C_{22}H_{17}ClN_2O_3S$ as $(M+H)^+$ 425.4, 427.4.

15

EXAMPLE 184

5-Chloro-thiophene-2-carboxylic acid [3-(2'-chloro-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide


20



[0705] The titled compound was made by the procedure similar to that described in Example 183. MS found for $C_{21}H_{16}Cl_2N_2O_2S$ as (M+H)⁺ 431.3, 433.0

EXAMPLE 185


- 5 5-Chloro-thiophene-2-carboxylic acid [3-(2'-methylsulfanyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- [0706] The titled compound was made by the procedure similar to that described in
10 Example 183. MS found for $C_{22}H_{19}ClN_2O_2S_2$ as (M+H)⁺ 429.0, 431.0

EXAMPLE 186


- 5-Chloro-thiophene-2-carboxylic acid [3-(2'-amino-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- 15 [0707] The titled compound was made by the procedure similar to that described in Example 183. MS found for $C_{21}H_{18}ClN_3O_2S$ as (M+H)⁺ 412.3, 414.3

EXAMPLE 187

- 20 5-Chloro-thiophene-2-carboxylic acid [3-(2'-trifluoromethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0708] The titled compound was made by the procedure similar to that described in Example 183. MS found for $C_{22}H_{16}ClF_3N_2O_2S$ as (M+H)⁺ 465.5, 467.5

5

EXAMPLE 188

5-Chloro-thiophene-2-carboxylic acid [3-(2'-methoxy-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0709] The titled compound was made by the procedure similar to that described in

10 Example 183. MS found for $C_{22}H_{19}ClN_2O_3S$ as (M+H)⁺ 505.4, 507.4

EXAMPLE 189

5-Chloro-thiophene-2-carboxylic acid [3-(2'-methoxymethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide




15

[0710] The titled compound was made by the procedure similar to that described in Example 183. MS found for $C_{23}H_{19}ClN_2O_3S$ as (M+H)⁺ 519.4, 521.4

EXAMPLE 190


5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-3-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- 5 [0711] The titled compound was made by the procedure similar to that described in Example 183. MS found for $C_{20}H_{16}ClN_3O_2S$ as $(M+H)^+$ 398.4, 400.4.

EXAMPLE 191


5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-4-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- [0712] The titled compound was made by the procedure similar to that described in Example 183. MS found for $C_{20}H_{16}ClN_3O_2S$ as $(M+H)^+$ as 398.4, 400.4.

EXAMPLE 192


5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-3-yl-phenyl)-isoxazol-5-ylmethyl]-amide



- [0713] The titled compound was made by the procedure similar to that described in Example 183. MS found for $C_{20}H_{14}ClN_3O_2S$ as $(M+H)^+$ 396.4, 398.4.

EXAMPLE 193

5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-4-yl-phenyl)-isoxazol-5-ylmethyl]-amide




[0714] The titled compound was made by the procedure similar to that described in

5 Example 183. MS found for $C_{20}H_{14}ClN_3O_2S$ as $(M+H)^+$ 396.4, 398.4.

EXAMPLE 194

5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




10 [0715] The titled compound was made by the procedure similar to that described in

Example 169, step 7 using the material from Example 183 as starting material. MS found for $C_{24}H_{24}ClN_3O_2S$ as $(M+H)^+$ 454.0, 456.0.

EXAMPLE 195

15 5-Chloro-thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0716] The titled compound was made by the procedure similar to that described in

20 Example 194. MS found for $C_{26}H_{26}ClN_3O_2S$ as $(M+H)^+$ 481.2, 482.2.

EXAMPLE 196


5-Chloro-thiophene-2-carboxylic acid [3-(2'-piperidin-1-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- 5 [0717] The titled compound was made by the procedure similar to that described in Example 194. MS found for $C_{27}H_{28}ClN_3O_2S$ as (M+H)⁺ 494.3, 496.3

EXAMPLE 197


5-Chloro-thiophene-2-carboxylic acid [3-(2'-morpholin-4-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- [0718] The titled compound was made by the procedure similar to that described in Example 194. MS found for $C_{26}H_{26}ClN_3O_3S$ as (M+H)⁺ 496.3, 498.3

EXAMPLE 198


5-Chloro-thiophene-2-carboxylic acid [3-(2'-{[(2-methoxy-ethyl)-methyl-amino]-methyl}-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- [0719] The titled compound was made by the procedure similar to that described in Example 194. MS found for $C_{26}H_{28}ClN_3O_3S$ as (M+H)⁺ 498.3, 500.3

EXAMPLE 199


5-Chloro-thiophene-2-carboxylic acid {3-[2'-(4-methyl-piperazin-1-ylmethyl)-biphenyl-4-yl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- 5 [0720] The titled compound was made by the procedure similar to that described in Example 194. MS found for $C_{27}H_{29}ClN_4O_2S$ as $(M+H)^+$ 509.7, 511.7

EXAMPLE 200

- 10 5-Chloro-thiophene-2-carboxylic acid [3-(2'-{[(2-dimethylamino-ethyl)-methyl-amino]-methyl}-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




- [0721] The titled compound was made by the procedure similar to that described in Example 194. MS found for $C_{27}H_{31}ClN_4O_2S$ as $(M+H)^+$ 511.3, 513.3

15

EXAMPLE 201


5-Chloro-thiophene-2-carboxylic acid [3-(2'-azepan-1-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- [0722] The titled compound was made by the procedure similar to that described in
20 Example 194. MS found for $C_{28}H_{30}ClN_3O_2S$ as $(M+H)^+$ 508.3, 510.4

EXAMPLE 202


5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-5'-methoxy-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- 5 [0723] The titled compound was made by the procedure similar to that described in Example 194. MS found for $C_{25}H_{26}ClN_3O_3S$ as $(M+H)^+$ 484.1, 486.1

EXAMPLE 203

10 5-Chloro-thiophene-2-carboxylic acid [3-(5'-methoxy-2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




- [0724] The titled compound was made by the procedure similar to that described in Example 194. MS found for $C_{27}H_{28}ClN_3O_3S$ as $(M+H)^+$ 510.1, 512.1

15

EXAMPLE 204


5-Chloro-thiophene-2-carboxylic acid [3-(5'-chloro-2'-dimethylaminomethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



[0725] The titled compound was made by the procedure similar to that described in Example 194. MS found for $C_{24}H_{23}Cl_2N_3O_2S$ as $(M+H)^+$ 488.3, 490.3

EXAMPLE 205

- 5 5-Chloro-thiophene-2-carboxylic acid [3-(5'-chloro-2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0726] The titled compound was made by the procedure similar to that described in Example 194. MS found for $C_{26}H_{25}Cl_2N_3O_2S$ as $(M+H)^+$ 514.3, 516.3

10

EXAMPLE 206


- 5-Chloro-thiophene-2-carboxylic acid [3-(5'-chloro-2'-{[(2-methoxy-ethyl)-methyl-amino]-methyl}-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- 15 [0727] The titled compound was made by the procedure similar to that described in Example 194. MS found for $C_{26}H_{27}Cl_2N_3O_3S$ as $(M+H)^+$ 532.4, 534.4

EXAMPLE 207

- 20 5-Chloro-thiophene-2-carboxylic acid [3-(2'-hydroxymethyl-4'-methoxy-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0728] The titled compound was made by the procedure similar to that described in Example 183. MS found for $C_{23}H_{21}ClN_2O_4S$ as $(M+H)^+$ 457.5, 459.5

5

EXAMPLE 208

5-Chloro-thiophene-2-carboxylic acid [3-(2'-formyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide




[0729] The titled compound was made by the procedure similar to that described in Example 183. MS found for $C_{22}H_{15}ClN_2O_3S$ as $(M+H)^+$ 423.4, 425.4

10

EXAMPLE 209

5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide




15

[0730] The titled compound was made by the procedure similar to that described in Example 169 using material from Example 208. MS found for $C_{24}H_{22}ClN_3O_2S$ as $(M+H)^+$ 452.1, 454.1

EXAMPLE 210


5-Chloro-thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide



- 5 [0731] The titled compound was made by the procedure similar to that described in Example 209. MS found for $C_{26}H_{24}ClN_3O_2S$ as $(M+H)^+$ 478.1, 480.1

EXAMPLE 211


5-Chloro-thiophene-2-carboxylic acid [3-(3'-dimethylaminomethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide



- [0732] The titled compound was made by the procedure similar to that described in Example 209. MS found for $C_{24}H_{22}ClN_3O_2S$ as $(M+H)^+$ 452.5, 454.5.

EXAMPLE 212


5-Chloro-thiophene-2-carboxylic acid [3-(3'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide



- [0733] The titled compound was made by the procedure similar to that described in Example 209. MS found for $C_{26}H_{24}ClN_3O_2S$ as $(M+H)^+$ 478.5, 480.5

EXAMPLE 213


5-Chloro-thiophene-2-carboxylic acid [3-(4'-dimethylaminomethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide



- 5 [0734] The titled compound was made by the procedure similar to that described in Example 209. MS found for $C_{24}H_{22}ClN_3O_2S$ as $(M+H)^+$ 452.5, 454.5.

EXAMPLE 214


5-Chloro-thiophene-2-carboxylic acid [3-(4'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide



- [0735] The titled compound was made by the procedure similar to that described in Example 209. MS found for $C_{26}H_{24}ClN_3O_2S$ as $(M+H)^+$ 478.5, 480.5

EXAMPLE 215


5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-5'-methoxy-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide



- [0736] The titled compound was made by the procedure similar to that described in Example 209. MS found for $C_{25}H_{24}ClN_3O_3S$ as $(M+H)^+$ 482.1, 484.1.

EXAMPLE 216


5-Chloro-thiophene-2-carboxylic acid [3-(5'-methoxy-2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide



[0737] The titled compound was made by the procedure similar to that described in Example 209. MS found for $C_{27}H_{26}ClN_3O_3S$ as $(M+H)^+$ 508.1, 510.1

EXAMPLE 217


5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-4'-methoxy-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide



[0738] The titled compound was made by the procedure similar to that described in Example 209. MS found for $C_{25}H_{24}ClN_3O_3S$ as $(M+H)^+$

EXAMPLE 218

5-Chloro-thiophene-2-carboxylic acid [3-(4'-methoxy-2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide




[0739] The titled compound was made by the procedure similar to that described in Example 209. MS found for $C_{27}H_{26}ClN_3O_3S$ as $(M+H)^+$

5

EXAMPLE 219

5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-2-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0740] Combined the bromide from Example 1 (50 mg, 0.13 mmol) with 2-pyridyltributyl stannane (55 mg, 0.15 mmol) and 5 mL of toluene. The mixture was degassed for three minutes then treated with tetrakis(triphenylphosphine) palladium (14 mg, 0.013 mmol) and stirred at reflux overnight. The following day the reaction was checked by TLC which showed consumption of the starting bromide and a more polar spot. The reaction was partitioned with aqueous sodium fluoride and ethyl acetate, separated and dried over magnesium sulfate. After filtration and concentration the crude residue was purified by preparative HPLC affording the desired product as a white solid. MS found for $C_{20}H_{16}ClN_3O_2S$ as $(M+H)^+$ 398.1, 400.1

20

EXAMPLE 220

5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-2-yl-phenyl)-isoxazol-5-ylmethyl]-amide




[0741] The titled compound was made by the procedure similar to that described in Example 219. MS found for $C_{20}H_{14}ClN_3O_2S$ as $(M+H)^+$ 396.1, 398.1.

5

EXAMPLE 221

5-Chloro-thiophene-2-carboxylic acid [3-(4-benzylamino-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0742] The titled compound was made by the procedure similar to that described in Example 169, step 7. MS found for $C_{22}H_{20}ClN_3O_2S$ as $(M+H)^+$ 426.1, 428.1

10

EXAMPLE 222

5-Chloro-thiophene-2-carboxylic acid [3-(4-phenylamino-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



15

[0743] Aryl Bromide (Example 1, 100 mg, 0.25 mmol), aniline (27 μ L, 0.30 mmol), Cs_2CO_3 (163 mg, 0.50 mmol) and Xantphos (43 mg, 0.075 mmol) were diluted with 4 mL of degassed THF. The resulting suspension was degassed for an additional 3 min then $Pd_2(dba)_3$ (23 mg, 0.25 mmol) was added giving a dark violet suspension which became dark yellow-orange within a few minutes. The mixture was then refluxed overnight, changing to cloudy orange as the aryl bromide was consumed. The next day the mixture was partitioned with water and ethyl acetate, separated, and the organic layer extracted twice with ethyl acetate.


20

The combined organic layers were dried over MgSO_4 , filtered, concentrated and purified by preparative HPLC affording the desired product as a white powder. MS found for $\text{C}_{21}\text{H}_{18}\text{ClN}_3\text{O}_2\text{S}$ as $(\text{M}+\text{H})^+$ 412.0, 414.0.

5

EXAMPLE 223

5-Chloro-thiophene-2-carboxylic acid {3-[4-(4-fluoro-phenylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0744] The titled compound was made by the procedure similar to that described in Example 222. MS found for $\text{C}_{21}\text{H}_{17}\text{ClFN}_3\text{O}_2\text{S}$ as $(\text{M}+\text{H})^+$ 430.1, 433.1.

10

EXAMPLE 224

5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-2-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




15

[0745] The titled compound was made by the procedure similar to that described in Example 222. MS found for $\text{C}_{20}\text{H}_{17}\text{ClN}_4\text{O}_2\text{S}$ as $(\text{M}+\text{H})^+$ 413.0, 415.0.

EXAMPLE 225

5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-3-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide


20



[0746] The titled compound was made by the procedure similar to that described in Example 222. found for $C_{20}H_{17}ClN_4O_2S$ as $(M+H)^+$ 413.0, 415.0.

EXAMPLE 226

5 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide




[0747] The titled compound was made by the procedure similar to that described in Example 222. found for $C_{20}H_{17}ClN_4O_2S$ as $(M+H)^+$ 413.0, 415.0.

10

EXAMPLE 227


5-Chloro-thiophene-2-carboxylic acid {3-[4-(methyl-pyridin-4-yl-amino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



15 [0748] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{21}H_{19}ClN_4O_2S$ as $(M+H)^+$ 427.4, 429.4.

EXAMPLE 228


20 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-methyl-pyridin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



[0749] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{21}H_{19}ClN_4O_2S$ as $(M+H)^+$ 427.1, 429.1.

EXAMPLE 229


5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-chloro-pyridin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- 5 [0750] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{16}Cl_2N_4O_2S$ as $(M+H)^+$ 447.2, 449.2.

EXAMPLE 230


- 10 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyrimidin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- [0751] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{19}H_{16}ClN_5O_2S$ as $(M+H)^+$ 414.1, 416.1, $(M+Na)^+$ 436.0, 438.0

- 15 EXAMPLE 231


5-Chloro-thiophene-2-carboxylic acid (3-{4-[(2-amino-pyrimidin-4-yl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide



- 20 [0752] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{19}ClN_6O_2S$ as $(M+H)^+$ 433.3, 435.3.

EXAMPLE 232


5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-ethyl-2H-pyrazol-3-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- 5 [0753] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{20}ClN_5O_2S$ as $(M+H)^+$ 430.2, 432.3.

EXAMPLE 233


5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-1H-pyrazol-3-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- [0754] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{19}H_{18}ClN_5O_2S$ as $(M+H)^+$ 416.5, 418.5.

EXAMPLE 234

5-Chloro-thiophene-2-carboxylic acid [3-(4-morpholin-4-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




- [0755] The titled compound was made by the procedure similar to that described in

20 Example 222. MS found for $C_{19}H_{20}ClN_3O_3S$ as $(M+H)^+$ 406.3, 408.3; $(M+Na)^+$ 428.2, 430.2.

EXAMPLE 235


5-Chloro-thiophene-2-carboxylic acid {3-[4-(4-methyl-piperazin-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- 5 [0756] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{23}ClN_4O_2S$ as $(M+H)^+$ 419.3, 421.3.

EXAMPLE 236


5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-methoxymethyl-pyrrolidin-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- [0757] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{21}H_{24}ClN_3O_3S$ as $(M+H)^+$ 434.0, 436.0.

EXAMPLE 237


5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-piperidin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- [0758] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{21}H_{25}ClN_4O_2S$ as $(M+H)^+$ 433.3, 435.3.

EXAMPLE 238


5-Chloro-thiophene-2-carboxylic acid (3-{4-[(2-dimethylamino-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide



- 5 [0759] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{25}ClN_4O_2S$ as $(M+H)^+$ 421.3, 423.3.

EXAMPLE 239


5-Chloro-thiophene-2-carboxylic acid (3-{4-[(3-dimethylamino-propyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide



- [0760] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{21}H_{27}ClN_4O_2S$ as $(M+H)^+$ 421.0, 423.0.

EXAMPLE 240


5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-oxo-pyrrolidin-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- 15 [0761] The titled compound was made by the procedure similar to that described in
20 Example 222. MS found for $C_{19}H_{18}ClN_3O_3S$ as $(M+H)^+$ 404.2, 406.2.

EXAMPLE 241


5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-oxo-piperidin-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- 5 [0762] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{20}ClN_3O_3S$ as $(M+H)^+$ 418.2, 420.2

EXAMPLE 242


4-[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-phenyl]-3-oxo-piperazine-1-carboxylic acid benzyl ester



- [0763] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{27}H_{25}ClN_4O_5S$ as $(M+H)^+$ 553.3, 555.3

EXAMPLE 243


5-Chloro-thiophene-2-carboxylic acid {3-[4-(3-oxo-morpholin-4-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- [0764] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{19}H_{18}ClN_3O_4S$ as $(M-H)^+$ 418.1, 420.1.

EXAMPLE 244


5-Chloro-thiophene-2-carboxylic acid {3-[4-(3-oxo-thiomorpholin-4-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- 5 [0765] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{19}H_{18}ClN_3O_3S_2$ as $(M+H)^+$ 436.2, 438.2.

EXAMPLE 245


5-Chloro-thiophene-2-carboxylic acid {3-[4-(3-methyl-2-oxo-imidazolidin-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- [0766] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{19}H_{19}ClN_4O_3S$ as $(M+H)^+$ 419.2, 421.2.

EXAMPLE 246


5-Chloro-thiophene-2-carboxylic acid {3-[4-(5-oxo-[1,4]oxazepan-4-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- [0767] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{20}ClN_3O_4S$ as $(M+H)^+$ 434.5, 436.5.

EXAMPLE 247


5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(pyridin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- 5 [0768] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{16}ClFN_4O_2S$ as $(M+H)^+$ 431.1, 433.1

EXAMPLE 248


5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(4-methyl-[1,4]diazepan-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- [0769] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{21}H_{24}ClFN_4O_2S$ as $(M+H)^+$ 451.2, 453.3

EXAMPLE 249


5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(5-oxo-[1,4]oxazepan-4-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide



- [0770] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{19}ClFN_3O_4S$ as $(M+H)^+$ 452.0, 454.0

EXAMPLE 250


5-Chloro-thiophene-2-carboxylic acid [3-(4-phenylamino-phenyl)-isoxazol-5-ylmethyl]-amide



- 5 [0771] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{21}H_{16}ClN_3O_2S$ as $(M+H)^+$ 410.4, 412.3.

EXAMPLE 251


5-Chloro-thiophene-2-carboxylic acid {3-[4-(4-methoxy-phenylamino)-phenyl]-isoxazol-5-ylmethyl}-amide



- [0772] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{22}H_{18}ClN_3O_3S$ as $(M+H)^+$ 440.0, 442.0.

EXAMPLE 252


5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-2-ylamino)-phenyl]-isoxazol-5-ylmethyl}-amide



- [0773] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{15}ClN_4O_2S$ as $(M+H)^+$ 411.0, 413.0.

EXAMPLE 253


5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-3-ylamino)-phenyl]-isoxazol-5-ylmethyl}-amide



[0774] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{15}ClN_4O_2S$ as $(M+H)^+$ 411.0, 413.1.

EXAMPLE 254


5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-4-ylamino)-phenyl]-isoxazol-5-ylmethyl}-amide



[0775] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{15}ClN_4O_2S$ as $(M+H)^+$ 411.0, 413.0.

EXAMPLE 255


5-Chloro-thiophene-2-carboxylic acid {3-[4-(methyl-pyridin-4-yl-amino)-phenyl]-isoxazol-5-ylmethyl}-amide



[0776] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{21}H_{17}ClN_4O_2S$ as $(M+H)^+$ 425.0, 427.0.

EXAMPLE 256


5-Chloro-thiophene-2-carboxylic acid [3-(4-pyrrolidin-1-yl-phenyl)-isoxazol-5-ylmethyl]-amide



- 5 [0777] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{19}H_{18}ClN_3O_2S$ as $(M+H)^+$ 388.4, 390.4

EXAMPLE 257


5-Chloro-thiophene-2-carboxylic acid [3-(4-morpholin-4-yl-phenyl)-isoxazol-5-ylmethyl]-amide



- [0778] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{19}H_{18}ClN_3O_3S$ as $(M+H)^+$ 404.3, 406.3.

EXAMPLE 258


5-Chloro-thiophene-2-carboxylic acid [3-(4-benzylamino-phenyl)-isoxazol-5-ylmethyl]-amide



- [0779] The titled compound was made by the procedure similar to that described in Example 169, step 7. MS found for $C_{22}H_{18}ClN_3O_2S$ as $(M+H)^+$ 424.0, 426.0

EXAMPLE 259


5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-piperidin-4-ylamino)-phenyl]-isoxazol-5-ylmethyl}-amide



[0780] The titled compound was made by the procedure similar to that described in Example 169, step 7. MS found for $C_{21}H_{23}ClN_4O_2S$ as $(M+H)^+$ 431.4, 433.4.

EXAMPLE 260


2-[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-phenylamino]-benzoic acid methyl ester



[0781] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{23}H_{20}ClN_3O_4S$ as $(M+H)^+$ 470.0, 472.0.

EXAMPLE 261


3-[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-phenylamino]-benzoic acid methyl ester



[0782] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{23}H_{20}ClN_3O_4S$ as $(M+H)^+$ 470.0, 472.0.

EXAMPLE 262


4-[4-(5-{{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-phenylamino]-benzoic acid methyl ester



- 5 [0783] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{23}H_{20}ClN_3O_4S$ as $(M+H)^+$ 470.0, 472.0.

EXAMPLE 263


- 10 5-Chloro-thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- [0784] The titled compound was made using the starting bromide from Example 3 by the procedure similar to that described in Example 183, followed by the procedure of Example 15 169, step 7. MS found for $C_{26}H_{26}ClN_3O_2S$ as $(M+H)^+$ 480.5, 482.5.

EXAMPLE 264


5-Chloro-thiophene-2-carboxylic acid [3-(3'-dimethylaminomethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



- 20 [0785] The titled compound was made using the starting bromide from Example 3 by the procedure similar to that described in Example 183, followed by the procedure of Example 169, step 7. MS found for $C_{24}H_{24}ClN_3O_2S$ as $(M+H)^+$ 454.1, 456.1.

EXAMPLE 265


5-Chloro-thiophene-2-carboxylic acid [3-(3'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



[0786] The titled compound was made using the starting bromide from Example 3 by the procedure similar to that described in Example 183, followed by the procedure of Example 169, step 7. MS found for $C_{26}H_{26}ClN_3O_2S$ as $(M+H)^+$ 480.1, 482.1.

EXAMPLE 266


5-Chloro-thiophene-2-carboxylic acid [3-(4'-dimethylaminomethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



[0787] The titled compound was made using the starting bromide from Example 3 by the procedure similar to that described in Example 183, followed by the procedure of Example 169, step 7. MS found for $C_{24}H_{24}ClN_3O_2S$ as $(M+H)^+$ 454.1, 456.1.

EXAMPLE 267


5-Chloro-thiophene-2-carboxylic acid [3-(4'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide



[0788] The titled compound was made using the starting bromide from Example 3 by the procedure similar to that described in Example 183, followed by the procedure of Example 169, step 7. MS found for $C_{26}H_{26}ClN_3O_2S$ as $(M+H)^+$ 480.1, 482.1.

EXAMPLE 268


5-Chloro-thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide



[0789] The titled compound was made using the starting bromide from Example 4 by the procedure similar to that described in Example 183, followed by the procedure of Example 169, step 7. MS found for $C_{24}H_{22}ClN_3O_2S$ as $(M+H)^+$ 452.5, 454.5.

EXAMPLE 269


5-Chloro-thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide



[0790] The titled compound was made using the starting bromide from Example 4 by the procedure similar to that described in Example 183, followed by the procedure of Example 169, step 7. MS found for $C_{26}H_{24}ClN_3O_2S$ as $(M+H)^+$ 478.5, 480.5.

EXAMPLE 270

5-Chloro-thiophene-2-carboxylic acid [3-(3'-dimethylaminomethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide




[0791] The titled compound was made using the starting bromide from Example 4 by the procedure similar to that described in Example 183, followed by the procedure of Example 169, step 7. MS found for $C_{24}H_{22}ClN_3O_2S$ as (M+H)⁺ 452.1, 454.1.

5

EXAMPLE 271

5-Chloro-thiophene-2-carboxylic acid [3-(3'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide




10

[0792] The titled compound was made using the starting bromide from Example 4 by the procedure similar to that described in Example 183, followed by the procedure of Example 169, step 7. MS found for $C_{26}H_{24}ClN_3O_2S$ as (M+H)⁺ 478.1, 480.1.

EXAMPLE 272

5-Chloro-thiophene-2-carboxylic acid [3-(4'-dimethylaminomethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide

15




[0793] The titled compound was made using the starting bromide from Example 4 by the procedure similar to that described in Example 183, followed by the procedure of Example 169, step 7. MS found for $C_{24}H_{22}ClN_3O_2S$ as (M+H)⁺ 452.1, 454.1.

20

EXAMPLE 273

5-Chloro-thiophene-2-carboxylic acid [3-(4'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide




[0794] The titled compound was made using the starting bromide from Example 4 by the procedure similar to that described in Example 183, followed by the procedure of Example 169, step 7. MS found for $C_{26}H_{24}ClN_3O_2S$ as $(M+H)^+$ 478.1, 480.1.

5

EXAMPLE 274

5-Chloro-thiophene-2-carboxylic acid [3-(3-pyridin-3-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0795] The titled compound was made by the procedure similar to that described in Example 183. ^1H NMR ($DMSO-d_6$, 400 MHz): δ 9.02 (s, 1H), δ 8.89 (t, 1H), 8.68 (d, 1H), δ 8.38 (d, 1H), 7.92 (s, 1H), 7.82 (d, 1H), δ 7.71 (m, 2H), δ 7.63 (d, 1H), δ 7.58 (t, 1H), δ 7.12 (d, 1H), δ 4.88 (m, 1H), δ 3.53 (dd, 1H), δ 3.41 (m, 2H), δ 3.258 (dd, 1H).

10

EXAMPLE 275

15 5-Chloro-thiophene-2-carboxylic acid [3-(3-pyridin-4-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide




[0796] The titled compound was made by the procedure similar to that described in Example 183. ^1H NMR ($DMSO-d_6$, 400 MHz): δ 8.89 (t, 1H), δ 8.79 (d, 2H), δ 8.04 (m, 3H), δ 7.94 (d, 1H), δ 7.82 (d, 1H), δ 7.62 (m, 2H), δ 7.13 (d, 1H), δ 4.90 (m, 1H), δ 3.59 (dd, 1H), δ 3.41 (m, 2H), δ 3.28 (dd, 1H).

20

EXAMPLE 276

5-Chloro-thiophene-2-carboxylic acid [3-(3-pyridin-3-yl-phenyl)-isoxazol-5-ylmethyl]-amide




25

[0797] The titled compound was made by the procedure similar to that described in Example 183. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 9.31 (t, 1H), δ 9.07 (s, 1H), δ 8.65 (s, 1H), δ 8.38 (d, 1H), δ 8.19 (s, 1H), δ 7.91 (d, 1H), δ 7.87 (d, 1H), δ 7.63 (m, 3H), δ 7.19 (d, 1H), δ 7.09 (s, 1H), 4.60 (d, 2H).

EXAMPLE 277


5-Chloro-thiophene-2-carboxylic acid [3-(3-pyridin-4-yl-phenyl)-isoxazol-5-ylmethyl]-amide



[0798] The titled compound was made by the procedure similar to that described in Example 183. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 9.35 (t, 1H), δ 8.77 (d, 1H), δ 8.30 (s, 1H), δ 8.08 (d, 1H), δ 8.02 (d, 1H), δ 7.98 (d, 1H), δ 7.68 (m, 2H), δ 7.19 (d, 1H), δ 7.10 (s, 1H), 4.61 (d, 2H).

EXAMPLE 278


5-Chloro-N-((3-(4-(2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



[0799] Aryl bromide (Example 1, 100 mg, 0.25 mmol) and 2-hydroxypyridine (95 mg, 1.0 mmol) were dissolved in 1.5 mL dry dioxane and 0.5 mL dry DMSO in a sealed tube. To it were added N,N'-dimethylethylenediamine (27 μL, 0.25 mmol), CuI (95 mg, 0.125 mmol) and K₃PO₄ (106 mg, 0.5 mmol) in order. The mixture was stirred and heated in 120°C bath till all the bromide starting material had been consumed (3 hrs, monitored by analytical HPLC). The mixture was filtered and directly subjected to reverse phase preparative HPLC to isolate the title compound as a white powder after lyophilization. MS found for C₂₀H₁₆ClN₃O₃S as (M+H)⁺ 414.1, 416.1

EXAMPLE 279


5-Chloro-N-((3-(4-(3-fluoro-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



[0800] The title compound was prepared by the same procedure described in Example 278 using 3-fluoro-2-hydroxypyridine in place of 2-hydroxypyridine. MS found for $C_{20}H_{15}ClFN_3O_3S$ as (M+H)⁺ 432.1, 434.1

EXAMPLE 280


5-Chloro-N-((3-(4-(5-fluoro-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



[0801] The title compound was prepared by the same procedure described in Example 278 using 5-fluoro-2-hydroxypyridine in place of 2-hydroxypyridine. MS found for $C_{20}H_{15}ClFN_3O_3S$ as (M+H)⁺ 432.1, 434.1

EXAMPLE 281

5-Chloro-N-((3-(4-(3-methyl-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide




[0802] The title compound was prepared by the same procedure described in Example 278 using 3-methyl-2-hydroxypyridine in place of 2-hydroxypyridine. MS found for $C_{21}H_{18}ClN_3O_3S$ as (M+H)⁺ 428.1, 430.1

5

EXAMPLE 282


5-Chloro-N-((3-(4-(3-methoxy-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



10 [0803] The title compound was prepared by the same procedure described in Example 278 using 3-methoxy-2-hydroxypyridine in place of 2-hydroxypyridine. MS found for $C_{21}H_{18}ClN_3O_4S$ as (M+H)⁺ 444.1, 446.1

EXAMPLE 283

15 5-Chloro-N-((3-(4-(3-chloro-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide




[0804] The title compound was prepared by the same procedure described in Example 278 using 3-chloro-2-hydroxypyridine in place of 2-hydroxypyridine. MS found for

20 $C_{20}H_{15}Cl_2N_3O_3S$ as (M+H)⁺ 448.0, 450.0

EXAMPLE 284


5-Chloro-N-((3-(4-(4-methyl-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



- 5 [0805] The title compound was prepared by the same procedure described in Example 278 using 4-methyl-2-hydroxypyridine in place of 2-hydroxypyridine. MS found for $C_{21}H_{18}ClN_3O_3S$ as (M+H)⁺ 428.1, 430.1

EXAMPLE 285


- 10 5-Chloro-N-((3-(4-(5-methyl-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



- [0806] The title compound was prepared by the same procedure described in Example 278 using 5-methyl-2-hydroxypyridine in place of 2-hydroxypyridine. MS found for $C_{21}H_{18}ClN_3O_3S$ as (M+H)⁺ 428.1, 430.1
- 15

EXAMPLE 286

5-Chloro-N-((3-(4-(5-methoxy-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide




[0807] The title compound was prepared by the same procedure described in Example 278 using 5-methoxy-2-hydroxypyridine in place of 2-hydroxypyridine. MS found for $C_{21}H_{18}ClN_3O_4S$ as (M+H)⁺ 444.1, 446.1

5

EXAMPLE 287

5-Chloro-N-((3-(4-(5-chloro-2-oxypyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide




10

[0808] The title compound was prepared by the same procedure described in Example 278 using 5-chloro-2-hydroxypyridine in place of 2-hydroxypyridine. MS found for $C_{20}H_{15}Cl_2N_3O_3S$ as (M+H)⁺ 448.0, 450.0

EXAMPLE 288

5-Chloro-N-((3-(4-iodophenyl)isoxazol-5-yl)methyl)thiophene-2-carboxamide




15

[0809] The title compound was prepared by the same procedure described in Example 2 using 4-iodobenzaldehyde in place of 4-bromobenzaldehyde. MS found for $C_{15}H_{10}ClIN_2O_2S$ as (M+H)⁺ 445.0, 447.0

20

EXAMPLE 289

5-Chloro-N-((3-(4-(2-oxypyridin-1(2H)-yl)phenyl)isoxazol-5-yl)methyl)thiophene-2-carboxamide




[0810] Aryl iodide (Example 288, 43 mg, 0.1 mmol) and 2-hydroxypyridine (38 mg, 0.4 mmol) were dissolved in 1.2 mL dry dioxane and 0.5 mL dry DMSO in a sealed tube. To it were added N,N'-dimethylethylenediamine (11 μ L, 0.1 mmol), CuI (10 mg, 0.05 mmol) and K₃PO₄ (42 mg, 0.2 mmol) in order. The mixture was stirred and heated in 120°C bath for 6 hrs. The mixture was filtered and directly subjected to reverse phase preparative HPLC to isolate the title compound as a white powder after lyophilization. MS found for C₂₀H₁₄ClN₃O₃S as (M+H)⁺ 412.0, 414.0

EXAMPLE 290

5-Chloro-N-((1-(4-(2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide (36)

SCHEME 5



Step 1:

[0811] 4-Iodoaniline (32, 6.00 g, 27.4 mmol) was dissolved in 25 mL TFA and stirred in ice bath. Solid NaNO₂ (2.07 g, 30.1 mmol) was added in small portions. The resulting mixture was stirred for 30 min in the ice bath. Sodium azide (1.87 g, 28.8 mmol) was

dissolved in 10 mL water and chilled in ice bath. This cold solution was then added to the TFA solution in three portions. The mixture was stirred in ice bath for 1 hr and concentrated in vacuo to remove TFA. The residue was taken into 600 mL DCM and washed with water three times. The organic phase was dried using MgSO_4 and concentrated in vacuo to afford 1-azido-4-iodobenzene **33** as a brownish waxy solid in >99% yield. In the mean time, 5-chlorothiophene-2-carboxylic acid (**4**, 9.13 g, 56 mmol) was dissolved in 200 mL dry DCM along with 0.5 mL dry DMF. To the vigorously stirred solution was carefully added oxalyl chloride (14.7 mL, 169 mmol) dropwise. The resulting solution was stirred for 3 hrs at RT and concentrated in vacuo. The residue was pumped to dryness and then dissolved in 300 mL dry DCM. To this solution was added propargylamine (5.8 mL, 84 mmol) dropwise. The mixture was stirred at RT for overnight and a lot of solid precipitated out. To this mixture was added 600 mL hexane. The mixture was vigorously stirred for a few hours and filtered for the solid (product **34**). The solid (9.47 g, 85%) was washed with hexane and was pure enough for direct use without further purification. MS found for $\text{C}_8\text{H}_6\text{ClNOS}$ as $(\text{M}+\text{H})^+$ 200.0, 202.0.

Step 2:


[0812] Aryl azide **33** (27 mmol) and alkyne **34** (5.37 g, 27 mmol) were dissolved in 40 mL dry DMF. To it was added 200 mL toluene. The mixture was refluxed in 135°C bath for 1 day. The mixture was concentrated in vacuo and subjected to flash column chromatography to isolate 1,4-disubstituted triazole **35** (major product: $\text{Rf}=0.30$ in 1:1 hexane/EtOAc) and its 1,5-disubstituted triazole isomer (minor product: $\text{Rf}=0.17$ in 1:1 hexane/EtOAc). MS found for **35** $\text{C}_{14}\text{H}_{10}\text{ClIN}_4\text{OS}$ as $(\text{M}+\text{H})^+$ 444.9, 446.9.

Step 3:

[0813] Aryl iodide **35** (100 mg, 0.22 mmol) and 2-hydroxypyridine (42 mg, 0.44 mmol) were dissolved in 5 mL dry DMSO in a sealed tube. To it were added 8-hydroxyquinoline (10 mg, 0.007 mmol), CuI (13 mg, 0.07 mmol) and Cs_2CO_3 (145 mg, 0.44 mmol). The mixture was stirred in 120°C bath for overnight. Then it was filtered and the filtrate was directly subjected to reverse phase preparative HPLC to isolate the title compound **36** as a white powder in 60-75% yield after lyophilization. MS found for $\text{C}_{19}\text{H}_{14}\text{ClN}_5\text{O}_2\text{S}$ as $(\text{M}+\text{H})^+$ 412.1, 414.1.

EXAMPLE 291


5-Chloro-N-((1-(4-(3-fluoro-2-oxypyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



[0814] Aryl iodide **35** (Example 290, 50 mg, 0.11 mmol) and 3-fluoro-2-hydroxypyridine (38 mg, 0.33 mmol) were dissolved in 1.0 mL dry dioxane and 0.3 mL dry DMSO in a sealed tube. To it were added N,N'-dimethylethylenediamine (8 μ L, 0.066 mmol), CuI (11 mg, 0.055 mmol) and K₃PO₄ (48 mg, 0.22 mmol) in order. The mixture was stirred and heated in 120°C bath for overnight. The mixture was filtered and directly subjected to reverse phase preparative HPLC to isolate the title compound as a white powder after lyophilization. MS found for C₁₉H₁₃ClFN₅O₂S as (M+H)⁺ 430.0, 432.0.

EXAMPLE 292


5-Chloro-N-((1-(4-(3-chloro-2-oxypyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



[0815] The title compound was prepared by the same procedure described in Example 290 using 3-chloro-2-hydroxypyridine in place of 3-fluoro-2-hydroxypyridine. MS found for C₁₉H₁₃Cl₂N₅O₂S as (M+H)⁺ 446.0, 448.0

EXAMPLE 293

5-Chloro-N-((1-(4-(3-methyl-2-oxypyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




[0816] The title compound was prepared by the same procedure described in Example 290 using 3-methyl-2-hydroxypyridine in place of 3-fluoro-2-hydroxypyridine. MS found for $C_{20}H_{16}ClN_5O_2S$ as (M+H)⁺ 426.1, 428.1

5

EXAMPLE 294


5-Chloro-N-((1-(4-(3-methoxy-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



10 [0817] The title compound was prepared by the same procedure described in Example 290 using 3-methoxy-2-hydroxypyridine in place of 3-fluoro-2-hydroxypyridine. MS found for $C_{20}H_{16}ClN_5O_3S$ as (M+H)⁺ 442.1, 444.1

EXAMPLE 295

15 5-Chloro-N-((1-(4-(4-methyl-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




[0818] The title compound was prepared by the same procedure described in Example 290 using 4-methyl-2-hydroxypyridine in place of 3-fluoro-2-hydroxypyridine. MS found for $C_{20}H_{16}ClN_5O_2S$ as (M+H)⁺ 426.1, 428.1

20

EXAMPLE 296


5-Chloro-N-((1-(4-(5-methyl-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



- 5 [0819] The title compound was prepared by the same procedure described in Example 290 using 5-methyl-2-hydroxypyridine in place of 3-fluoro-2-hydroxypyridine. MS found for $C_{20}H_{16}ClN_5O_2S$ as (M+H)⁺ 426.1, 428.1

EXAMPLE 297


- 10 5-Chloro-N-((1-(4-(5-methoxy-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



- [0820] The title compound was prepared by the same procedure described in Example 290 using 5-methoxy-2-hydroxypyridine in place of 3-fluoro-2-hydroxypyridine. MS found for $C_{20}H_{16}ClN_5O_3S$ as (M+H)⁺ 442.1, 444.1
- 15

EXAMPLE 298

5-Chloro-N-((1-(4-(5-fluoro-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




[0821] The title compound was prepared by the same procedure described in Example 290 using 5-fluoro-2-hydroxypyridine in place of 3-fluoro-2-hydroxypyridine. MS found for $C_{19}H_{13}ClFN_5O_2S$ as (M+H)⁺ 430.0, 432.0

5

EXAMPLE 299


5-Chloro-N-((1-(4-(5-chloro-2-oxypyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



10 [0822] The title compound was prepared by the same procedure described in Example 290 using 5-chloro-2-hydroxypyridine in place of 3-fluoro-2-hydroxypyridine. MS found for $C_{19}H_{13}Cl_2N_5O_2S$ as (M+H)⁺ 446.0, 448.0

EXAMPLE 300

15 5-Chloro-N-((1-(4-(2-oxopiperidin-1-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




[0823] The title compound was prepared by the same procedure described in Example 290 using δ -valerolactam (2-piperidone) in place of 3-fluoro-2-hydroxypyridine. MS found for $C_{19}H_{18}ClN_5O_2S$ as (M+H)⁺ 416.1, 418.1

20

EXAMPLE 301


5-Chloro-N-((1-(4-(3-oxomorpholino)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



[0824] The title compound was prepared by the same procedure described in Example 290 using 3-morpholinone in place of 3-fluoro-2-hydroxypyridine. MS found for $C_{18}H_{16}ClN_5O_3S$ as (M+H)⁺ 418.1, 420.1

EXAMPLE 302


5-Chloro-N-((1-(4-(3-oxothiomorpholino)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



[0825] The title compound was prepared by the same procedure described in Example 290 using 3-thiomorpholinone in place of 3-fluoro-2-hydroxypyridine. MS found for $C_{18}H_{16}ClN_5O_2S_2$ as (M+H)⁺ 434.0, 436.0

EXAMPLE 303

5-Chloro-N-((1-(4-(5-oxo-1,4-oxazepan-4-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




[0826] The title compound was prepared by the same procedure described in Example 290 using 1,4-oxazepan-5-one in place of 3-fluoro-2-hydroxypyridine. MS found for $C_{19}H_{18}ClN_5O_3S$ as (M+H)⁺ 432.1, 434.1

5

EXAMPLE 304

5-Chloro-N-((1-(2-fluoro-4-iodophenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




10

[0827] The title compound was prepared by the same procedure for compound 35 described in Example 290 in Scheme 5, using 4-iodo-2-fluoroaniline in place of 4-iodoaniline. MS found for $C_{14}H_9ClFIN_4OS$ as (M+H)⁺ 462.0, 464.0

EXAMPLE 305

5-Chloro-N-((1-(2-fluoro-4-(2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide

15




20

[0828] Example 304 (50 mg, 0.11 mmol) and 2-hydroxypyridine (42 mg, 0.44 mmol) were dissolved in 1.2 mL dry dioxane and 0.5 mL dry DMSO in a sealed tube. To it were added N,N'-dimethylethylenediamine (12 μ L, 0.11 mmol), CuI (11 mg, 0.055 mmol) and K_3PO_4 (47 mg, 0.22 mmol). The mixture was stirred in 120°C bath for 2.5 hr. The mixture was filtered and the filtrate was directly subjected to reverse phase preparative HPLC to isolate the title compound in 50-65% yield. MS found for $C_{19}H_{13}ClFN_5O_2S$ as (M+H)⁺ 430.0, 432.0

EXAMPLE 306

25

5-Chloro-N-((1-(2-fluoro-4-(3-oxomorpholino)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




[0829] The title compound was prepared by the same procedure described in Example 305 using 3-morpholinone in place of 2-hydroxypyridine. MS found for $C_{18}H_{15}ClFN_5O_3S$ as $(M+H)^+$ 436.1, 438.1

5

EXAMPLE 307

5-Chloro-N-((1-(2-hydroxy-4-(2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




- 10 [0830] Example 305 (58 mg, 0.13 mmol) was dissolved in 2 mL DMSO and 1 mL water in a sealed tube. To it was added Cs_2CO_3 (88 mg, 0.26 mmol). The mixture was stirred in $120^\circ C$ bath for overnight. It was cooled to RT and directly subjected to reverse phase HPLC to isolate the title compound as a white powder after lyophilization. MS found for $C_{19}H_{14}ClN_5O_3S$ as $(M+H)^+$ 428.1, 430.1

15

EXAMPLE 308

5-Chloro-N-((1-(2-(4-methylpiperazin-1-yl)-4-(2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




[0831] Example 305 (58 mg, 0.13 mmol) was dissolved in 2 mL anhydrous DMSO water in a sealed tube. To it were added N-methylpiperazine (44 μ L, 0.40 mmol) and Cs₂CO₃ (88 mg, 0.26 mmol). The mixture was stirred in 120°C bath for overnight. It was cooled to RT and directly subjected to reverse phase HPLC to isolate the title compound as a white powder after lyophilization. MS found for C₂₄H₂₄ClN₇O₂S (M+H)⁺ 510.1, 512.1

EXAMPLE 309

5-Chloro-N-((1-(5-(2-oxopyridin-1(2H)-yl)pyridin-2-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide (**40**)

SCHEME 6



Step 1:

[0832] 2-Fluoro-5-bromopyridine (2.18 g, 12.3 mmol) was dissolved in 25 mL anhydrous DMSO. To it was added NaN₃ (1.61 g, 24.6 mmol) and the mixture was stirred in 120°C bath for overnight. It was diluted with 300 mL ETOAc and washed with brine three times. The organic phase was dried and concentrated in vacuo to afford compound **38** (1.00 g, 41%) in high purity. MS found for C₅H₃BrN₄ as (M+H)⁺ 199.0, 201.0.

Step 2:

[0833] Azidopyridine **38** (65 mg, 0.33 mmol) and alkyne **34** (see Example 290 and Scheme 5; 66 mg, 0.33 mmol) were dissolved in 4 mL dry MeOH in a sealed tube. To it were added DIPEA (0.29 μ L, 1.64 mmol) and CuI (312 mg, 1.64 mmol). The mixture was stirred in 100 °C bath for 3 hr and cooled to RT. It was diluted with 200 mL acetone, stirred well, and filtered to remove the insoluble copper salt. The filtrate was concentrated in vacuo and

purified by flash column to yield 1,4-disubstituted triazole **39** (52 mg, 40%). MS found for $C_{13}H_9BrClN_5OS$ as (M+H)⁺ 398.0, 400.0.


Step 3:

[0834] Aryl bromide **39** (52 mg, 0.13 mmol) and 2-hydroxypyridine (37 mg, 0.39 mmol) were dissolved in 2 mL anhydrous dioxane and 2 mL anhydrous DMSO in a sealed tube. To it were added N,N'-dimethylethylenediamine (14 μ L, 0.13 mmol), CuI (13 mg, 0.07 mmol) and K_3PO_4 (55 mg, 0.26 mmol). The mixture was stirred in 120°C bath for overnight. Then it was filtered and the filtrate was directly subjected to reverse phase preparative HPLC to isolate the title compound **40** as a white powder in 50% yield after lyophilization. MS found for $C_{18}H_{13}ClN_6O_2S$ as (M+H)⁺ 413.1, 415.1.

EXAMPLE 310

5-Chloro-N-((3-(6-(2-oxopyridin-1(2H)-yl)pyridin-3-yl)isoxazol-5-yl)methyl)thiophene-2-carboxamide (**44**)

SCHEME 7



Step 1:

[0835] 2-Chloropyridine-5-carbaldehyde (3.00 g, 21.2 mmol) was dissolved in 20 mL pyridine and 20 mL ethanol. To it was added hydroxylamine hydrochloride (1.48 g, 21.1 mmol). The mixture was stirred at RT for 2 hr and then was concentrated in vacuo. The dry residue was dissolved in 50 mL anhydrous DMF and stirred in ice bath. To it was added NCS (3.25 g, 24.4 mmol). The mixture was stirred for overnight and diluted with 600 mL EtOAc. It was washed with brine four times, dried and concentrated in vacuo to give compound **42** in

quantitative yield in high purity as a white solid. MS found for $C_6H_4Cl_2N_2O$ as (M+H)⁺ 191.0, 193.0.

Step 2:

- 5 [0836] Compound **42** (3.80 g, 20 mmol) and alkyne **34** (see Example 290 and Scheme 5; 3.98 g, 20 mmol) were dissolved in 200 mL dry toluene. To it was added triethylamine (3.4 mL, 24 mmol). The mixture was refluxed for 3 hr and cooled to RT. White solid (compound **43**) precipitated out and was collected by filtration. It was washed by cold toluene and dried in vacuo. MS found for $C_{14}H_9Cl_2N_3O_2S$ as (M+H)⁺ 354.0, 356.0.

Step 3:


- 10 [0837] Compound **43** (53 mg, 0.15 mmol) and 2-hydroxypyridine (43 mg, 0.45 mmol) were dissolved in 2 mL anhydrous DMSO in a sealed tube. To it was added Cs_2CO_3 (73 mg, 0.22 mmol). The mixture was stirred in 100°C bath for overnight. Then it was filtered and the filtrate was directly subjected to reverse phase preparative HPLC to isolate the title compound **44** as a white powder in 40% yield after lyophilization. MS found for
15 $C_{19}H_{13}ClN_4O_3S$ as (M+H)⁺ 413.0, 415.0.

EXAMPLE 311

5-Chloro-N-((3-(4-(2-oxopyridin-1(2H)-yl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)thiophene-2-carboxamide (**51**)

20

SCHEME 8



Step 1:

[0838] 4-Iodobenzonitrile (3.00 g, 13 mmol) and 2-hydroxypyridine (2.47 g, 26 mmol) were dissolved in 30 mL anhydrous DMSO in a sealed tube. To it were added 8-hydroxyquinoline (0.57 g, 4 mmol), CuI (0.74 g, 4 mmol) and Cs₂CO₃ (8.5 g, 26 mmol). The mixture was stirred in 120°C bath for overnight. The mixture was partitioned in 500 mL EtOAc and 250 mL water. The organic phase was separated, and the aqueous was extracted with EtOAc three times. All the organic phases were combined, dried, concentrated and purified using flash column to give compound 46 (1.67 g, 66%). MS found for C₁₂H₈N₂O as (M+H)⁺ 197.1.

Step 2:

[0839] Compound 46 (1.67 g, 8.5 mmol) was dissolved in 40 mL DMSO. To it were added DIPEA (7.5 mL, 42.6 mmol) and hydroxylamine hydrochloride (3.0 g, 42.6 mmol). The mixture was stirred for 3 hr and to it was added 250 mL brine. White solid (compound 47) crashed out and was collected. It was washed with brine and cold water, and was dried in vacuo for overnight. MS found for C₁₂H₁₁N₃O₂ as (M+H)⁺ 230.1.

Step 3:

[0840] Glycine methyl ester hydrochloride (48, 2.00 g, 16 mmol) and 5-chlorothiophene-2-carboxylic acid (2.60 g, 16 mmol) were dissolved in 50 mL pyridine and stirred in ice bath.

To it was added POCl₃ (4.5 mL, 48 mmol) dropwise. After 5 minutes, ice chips were added into the mixture, followed by addition of 500 mL water. The mixture was well stirred and filtered. The aqueous filtrate was extracted with EtOAc four times. All the organic phases were combined, dried and concentrated in vacuo to give compound **49** was a light yellow solid (2.27 g, 61%). MS found for C₈H₈ClNO₃S as (M+H)⁺ 234.0, 236.0.

Step 4:

[0841] Methyl ester **49** (2.27 g, 9.7 mmol) was dissolved in 20 mL MeOH, 20 mL dioxane and 20 mL water. To it was added lithium hydroxide monohydrate (1.22 g, 29 mmol). The mixture was stirred for 2 hr at RT and acidified using 2M HCl till pH=7. The neutral mixture was concentrated in vacuo to remove organic solvent. The aqueous residue was extracted with EtOAc four times. The organic phases were combined, dried and concentrated in vacuo to afford acid **50** in quantitative yield. MS found for C₇H₆ClNO₃S as (M+H)⁺ 220.0, 222.0.


Step 5:

[0842] Acid **50** (219 mg, 1.0 mmol) was dissolved in 10 mL dry DMF. To it was added carbonyldiimidazole (180 mg, 1.1 mmol). The mixture was stirred for 1 hr at RT. To it was then added compound **47** (229 mg, 1.0 mmol). The mixture was stirred at RT for 30 min and slowly heated to 120°C via an oil bath. It was stirred then for 2 hr. After cooling down to RT, to the mixture was added 50 mL brine. The mixture was extracted with EtOAc four times. All the organic phases were combined, dried and concentrated in vacuo. The resulting mixture was purified with reverse phase preparative HPLC to give the title compound **51**. MS found for C₁₉H₁₃ClN₄O₃S as (M+H)⁺ 413.0, 415.0.

EXAMPLE 312

5-Chloro-N-((1-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide (**55**)

SCHEME 9



Step 1:

[0843] 4-Aminobenzonitrile (**52**, 2.0 g, 17 mmol) was dissolved in 8 mL TFA and stirred in
 5 ice bath. To it was added NaNO_2 (1.3 g, 18.7 mmol) in small portions. The mixture was
 stirred for 30 min in ice bath. Then NaN_3 (1.1 g, 17 mmol) was dissolved in 6 mL water and
 chilled in ice. This solution was added into the reaction mixture. After stirring for 1 hr in ice
 bath, the mixture was concentrated in vacuo to remove TFA, and diluted with water. The
 aqueous mixture was extracted with DCM three times. The organic phases were combined,
 10 dried and concentrated in vacuo to give compound **53** (2.35 g, 96%). MS found for $\text{C}_7\text{H}_4\text{N}_4$
 as $(\text{M}+\text{H})^+$ 145.0.

Step 2:

[0844] Azidobenzene **53** (1.77 g, 12.3 mmol) and alkyne **34** (see Scheme 5 in Example
 290; 2.45 g, 12.3 mmol) were refluxed in 200 mL toluene for overnight. The mixture was
 15 then concentrated in vacuo to half of the volume and chilled in ice bath. The solid
 precipitated out was pure major product 1,4-disubstituted triazole **54**, which was collected by
 filtration and washed with cold toluene. The minor product (the corresponding 1,5-triazole
 isomer) completely stayed in the filtrate along with some 1,4-disubstituted triazole **54**, which
 might be further recovered by multiple recrystallization process using DCM or EtOAc as
 20 solvent. 1,4-Disubstituted triazole **54** was dried in vacuo. MS found for $\text{C}_{15}\text{H}_{10}\text{ClN}_5\text{OS}$ as
 $(\text{M}+\text{H})^+$ 344.0, 346.0.


Step 3:

[0845] 1,4-Disubstituted triazole **54** (100 mg, 0.29 mmol) was dissolved in 50 mL
 anhydrous dioxane and 20 mL triethylamine. To this solution was bubbled in dry H_2S gas till
 25 saturation. The mixture was stirred for overnight at RT and was concentrated in vacuo. The

residue was placed in 50 mL dry acetone. To it was added 1.0 mL iodomethane. The mixture was refluxed for 1.5 hr and concentrated in vacuo. The residue was then dissolved in 20 mL anhydrous methanol. To it were added 0.5 mL acetic acid and 0.2 mL N-methylethylenediamine. The mixture was refluxed for 1 hr, concentrated in vacuo and subjected to reverse phase preparative HPLC to isolate the title compound **55** as a white powder after lyophilization. MS found for $C_{18}H_{17}ClN_6OS$ as (M+H)⁺ 401.1, 403.1.

EXAMPLE 313


N-((1-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



[0846] The title compound was prepared by the same procedure described in Scheme 9 in Example 312, using dimethylamine in place of N-methylethylenediamine. MS found for $C_{17}H_{17}ClN_6OS$ as (M+H)⁺ 389.1, 391.1.

EXAMPLE 314


N-((1-(4-(N-Ethyl-N-methylcarbamimidoyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



[0847] The title compound was prepared by the same procedure described in Scheme 9 in Example 312, using N-ethyl-N-methylamine in place of N-methylethylenediamine. MS found for $C_{18}H_{19}ClN_6OS$ as (M+H)⁺ 403.1, 405.1.

EXAMPLE 315


N-((1-(4-(N-Methyl-N-propylcarbamimidoyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



- 5 [0848] The title compound was prepared by the same procedure described in Scheme 9 in Example 312, using N-methyl-N-propylamine in place of N-methylethylenediamine. MS found for $C_{19}H_{21}ClN_6OS$ as $(M+H)^+$ 417.1, 419.1.

EXAMPLE 316


- 10 N-((1-(4-(N-(2-Methoxyethyl)-N-methylcarbamimidoyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



- [0849] The title compound was prepared by the same procedure described in Scheme 9 in Example 312, using N-(2-methoxyethyl)-N-methylamine in place of N-methylethylenediamine. MS found for $C_{19}H_{21}ClN_6O_2S$ as $(M+H)^+$ 433.1, 435.1.
- 15

EXAMPLE 317

N-((1-(4-(Azetidin-1-yl(imino)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide




[0850] The title compound was prepared by the same procedure described in Scheme 9 in Example 312, using azetidine in place of N-methylethylenediamine. MS found for $C_{18}H_{17}ClN_6OS$ as (M+H)⁺ 401.1, 403.1.

5

EXAMPLE 318

5-Chloro-N-((1-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




[0851] The title compound was prepared by the same procedure described in Scheme 9 in Example 312, using pyrrolidine in place of N-methylethylenediamine. MS found for $C_{19}H_{19}ClN_6OS$ as (M+H)⁺ 415.1, 417.1.

10

EXAMPLE 319

5-Chloro-N-((1-(4-(imino(piperidin-1-yl)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide

15




[0852] The title compound was prepared by the same procedure described in Scheme 9 in Example 312, using piperidine in place of N-methylethylenediamine. MS found for $C_{20}H_{21}ClN_6OS$ as (M+H)⁺ 429.1, 431.1.

20

EXAMPLE 320

N-((1-(4-(N,N-Dimethylcarbamimidoyl)-2-fluorophenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide (**59**)

SCHEME 10



Step 1:

[0853] To a solution of 4-amino-3-fluorobenzonitrile (**56**, 1.0 g, 7.35 mmol) in 5 mL TFA at 0°C was added NaNO₂ (558 mg, 8.09 mmol) in small portions. The reaction mixture was stirred at 0°C for 0.5 hr. Then a solution of NaN₃ (478 mg, 7.35 mmol) in 4 mL water was added. The resulting mixture was stirred at 0°C for 2 hr. The reaction solution was concentrated in vacuo and then neutralized to pH = 7 with the aqueous saturated NaHCO₃ solution and extracted with DCM three times. The combined organic phase was washed with water and brine, dried over Na₂SO₄ and evaporated in vacuo to give compound **57** (0.94g, 79%).

Step 2:

[0854] A solution of azidobenzene **57** (0.94 g, 5.80 mmol) and alkyne **34** (see Scheme 5 in Example 290; 1.3 g, 6.53 mmol) in 30 mL toluene was refluxed for 3 hr and concentrated in vacuo. The residue was dissolved into chloroform and was washed with water and brine, dried, evaporated and subjected to flash column chromatography to give compound **58**. MS found for C₁₅H₉ClFN₅OS as (M+Na)⁺: 384.0, 386.0.


Step 3:

[0855] To a solution of compound **58** (111 mg, 0.31 mmol) in the 1.2 mL triethylamine and 12 mL dioxane was bubbled dry H₂S gas to saturation. The mixture was stirred at RT for overnight and concentrated in vacuo. The residue was placed with 10 mL dry acetone and followed by the addition of 0.4 mL iodomethane. The reaction mixture was refluxed for 2 hr

and was then concentrated in vacuo. The residue was dissolved with 9 mL dry methanol. To it were added 0.2 mL acetic acid and dimethylamine (0.8 mL, 2.0 M in THF). The resulting mixture was refluxed for 1 hr, concentrated in vacuo and subjected to reverse phase preparative HPLC to give the title compound **59** as a white powder after lyophilization. MS found for $C_{17}H_{16}ClFN_6OS$ as $(M+H)^+$: 407.1, 409.1.

EXAMPLE 321


N-((1-(4-(Azetidin-1-yl(imino)methyl)-2-fluorophenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



[0856] The title compound was prepared by the same procedure described in Scheme 10 in Example 320, using azetidine in place of dimethylamine. MS found for $C_{18}H_{16}ClFN_6OS$ as $(M+H)^+$ 419.1, 421.1.

EXAMPLE 322


5-Chloro-N-((1-(2-fluoro-4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



[0857] The title compound was prepared by the same procedure described in Scheme 10 in Example 320, using pyrrolidine in place of dimethylamine. MS found for $C_{19}H_{18}ClFN_6OS$ as $(M+H)^+$ 433.1, 435.1.

EXAMPLE 323

5-Chloro-N-((1-(2-fluoro-4-(imino(piperidin-1-yl)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




[0858] The title compound was prepared by the same procedure described in Scheme 10 in Example 320, using piperidine in place of dimethylamine. MS found for $C_{20}H_{20}ClFN_6OS$ as (M+H)⁺ 447.1, 449.1.

5

EXAMPLE 324

5-Chloro-N-((1-(2-fluoro-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




10 [0859] The title compound was prepared by the same procedure described in Scheme 10 in Example 320, using N-methylethylenediamine in place of dimethylamine. MS found for $C_{18}H_{16}ClFN_6OS$ as (M+H)⁺ 419.1, 421.1.

EXAMPLE 325

15 N-((1-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-1H-pyrazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide (66)

SCHEME 11



Step 1:

[0860] The mixture of 4-cyanobenzeneboronic acid (**60**, 3.6 g, 24 mmol), 4-methylpyrazole (2.0 mL, 24 mmol), pyridine (5.8 mL, 72 mmol), Cu(OAc)₂ (8.7 g, 48 mmol) and activated molecular sieve powder in 100 mL chloroform was refluxed for 24 hr. The reaction was not complete. The mixture was filtered, and the solid cake was washed with plenty amount of methanol. The filtrate was concentrated in vacuo and purified by flash column to give compound **61** (0.93 g, 21%) as a white powder. MS found for C₁₁H₉N₃ as (M+H)⁺ 184.1.

Step 2:

[0861] Compound **61** (0.89 g, 4.8 mmol) was dissolved in 100 mL CCl₄. To it were added NBS (1.3 g, 7.3 mmol) and benzoyl peroxide (1.18 g, 4.8 mmol). The mixture was refluxed for 1 hr and concentrated in vacuo. The residue was subjected to flash column chromatography to give compound **62** (0.33 g, 26%). MS found for C₁₁H₈BrN₃ as (M+H)⁺ 262.0, 264.0.

Step 3:

[0862] Compound **62** (0.33 g, 1.26 mmol) was dissolved in 6 mL dry DMF. To it was added NaN₃ (164 mg, 2.52 mmol). The mixture was stirred for 2 hr and taken into 200 mL EtOAc. It was washed with water and brine, dried, concentrated and purified by flash column to give compound **63** (0.27 g, 96%). MS found for C₁₁H₈N₆ as (M+H)⁺ 225.1.

Step 4:

[0863] Compound **63** (0.27 g, 1.2 mmol) was dissolved in 10 mL THF and 20 mL MeOH. To it was added tin(II) chloride dihydrate (0.54 g, 2.4 mmol). The mixture was refluxed for 2 hr and was diluted with 500 mL chloroform. It was washed with 0.1N NaOH and water, dried, concentrated and purified by reverse phase HPLC to give compound **64** (0.13 g, 55%).

5 MS found for $C_{11}H_{10}N_4$ as (M+H)⁺ 199.1.

Step 5:

[0864] Compound **64** (0.13 g, 0.66 mmol) was dissolved in 12 mL dry pyridine. To it was added 5-chlorothiophene-2-carbonylchloride (0.24 g, 1.32 mmol). The mixture was stirred for 2 hr at RT, concentrated in vacuo and purified using flash column to give compound **65** (0.22 g, 99%). MS found for $C_{16}H_{11}ClN_4OS$ as (M+H)⁺ 343.0, 345.0.

10

Step 6:


[0865] Compound **65** (100 mg, 0.30 mmol) was placed in 50 mL anhydrous methanol in ice bath. To it was bubbled dry HCl gas till saturation. The mixture was stirred for overnight at RT and concentrated in vacuo. The dry residue was then dissolved in 50 mL dry methanol.

15 To it was added dimethylamine (1.2 mmol, 0.6 mL of 2.0M dimethylamine/THF solution). The mixture was refluxed for 1 hr, concentrated in vacuo, and purified with reverse phase HPLC to yield the title compound **66** as a white powder after lyophilization. MS found for $C_{18}H_{18}ClN_5OS$ as (M+H)⁺ 388.1, 390.1.

20

EXAMPLE 326

5-Chloro-N-((1-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1H-pyrazol-4-yl)methyl)thiophene-2-carboxamide




[0866] The title compound was prepared by the same procedure described in Scheme 11 in Example 325, using N-methylethylenediamine in place of dimethylamine. MS found for $C_{19}H_{18}ClN_5OS$ as (M+H)⁺ 400.1, 402.1.

25

EXAMPLE 327


N-((1-(4-(Azetidin-1-yl(imino)methyl)phenyl)-1H-pyrazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



- 5 [0867] The title compound was prepared by the same procedure described in Scheme 11 in Example 325, using azetidine in place of dimethylamine. MS found for $C_{19}H_{18}ClN_5OS$ as $(M+H)^+$ 400.1, 402.1.

EXAMPLE 328


10 5-Chloro-N-((1-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1H-pyrazol-4-yl)methyl)thiophene-2-carboxamide



- [0868] The title compound was prepared by the same procedure described in Scheme 11 in Example 325, using pyrrolidine in place of dimethylamine. MS found for $C_{20}H_{20}ClN_5OS$ as $(M+H)^+$ 414.1, 416.1.
- 15

EXAMPLE 329

5-Chloro-N-((1-(4-(imino(piperidin-1-yl)methyl)phenyl)-1H-pyrazol-4-yl)methyl)thiophene-2-carboxamide



20


[0869] The title compound was prepared by the same procedure described in Scheme 11 in Example 325, using piperidine in place of dimethylamine. MS found for $C_{21}H_{22}ClN_5OS$ as $(M+H)^+$ 428.1, 430.1.

5

EXAMPLE 330

N-((5-(4-(N,N-Dimethylcarbamimidoyl)phenyl)isoxazol-3-yl)methyl)-5-chlorothiophene-2-carboxamide (74).

SCHEME 12



10

Step 1:

[0870] Performed in the same manner as in Step 1, Example 1. Compound 68 was used immediately without purification.

Step 2:

15 [0871] Performed in the same manner as in Step 2, Example 1. Crude compound 69 was used for Step 5.

Step 3:

[0872] Combined aryl bromide 70 (5.00 g, 27 mmol) with CuI (0.10 g, 0.55 mmol), tri *t*-butyl phosphine (0.41 mL, 1.6 mmol) and $Pd(PhCN)_2Cl_2$ (0.32 g, 0.82 mmol). Degassed 3 min then added TMS acetylene (4.6 mL, 33 mmol) and diisopropylamine (9.5 mL, 68 mmol), the reaction changing colors from pale beige to black as the amine was added. After stirring

20

three hours all starting bromide was consumed as shown by HPLC. The reaction was filtered through a short pad of silica gel (eluted with dichloromethane) and concentrated. The resulting semisolid was triturated with 20 mL of hexane affording the desired product (**71**) as a brown solid (2.51 g, 47%). ¹H NMR (DMSO-*d*₆, 400 MHz): δ 7.79 (d, 2H), δ 7.56 (d, 2H),
5 0.19 (s, 9H).

Step 4:

[0873] TMS protected alkyne **71** (1.35 g, 1.01 mmol) was diluted with 10 mL of THF and treated with 1M tetrabutylammonium fluoride in THF (1.1 mL, 1.1 mmol). After 30 min HPLC showed complete consumption of the starting material and a more polar peak. The
10 reaction was diluted with 90 mL of water and 10 mL of 1 M aqueous HCl. The resulting solid was filtered and washed with water affording **72** as a brown solid (0.6237 g, 49%). ¹H NMR (DMSO-*d*₆, 400 MHz): δ 7.81 (d, 2H), δ 7.62 (d, 2H), δ 4.53 (s, 1H).

Step 5:


[0874] Alkyne **72** (0.36 g, 2.87 mmol) and triethylamine (0.60 mL, 4.31 mmol) were
15 diluted with 10 mL of toluene and heated to 100°C. To this solution was added **69** as the crude reaction mixture from Step 3, slowly. After the addition was complete the reaction was checked by TLC which showed ca. 50% reaction. The reaction mixture was cooled, partitioned with 1 M HCl and ethyl acetate and separated. The aqueous phase was extracted again with ethyl acetate and the combined organic layers were filtered through a short pad of
20 silica gel and concentrated. The resulting brown oil was then purified by silica gel chromatography (10% ethyl acetate / dichloromethane) affording isoxazole **73** as an off white solid (0.37 g, 41%). ¹H NMR (DMSO-*d*₆, 400 MHz): δ 8.03 (d, 2H), δ 7.94 (d, 2H), δ 7.47 (t, 1H), δ 7.08 (s, 1H), δ 4.18 (d, 2H), δ 1.38 (s, 9H).

Step 6:

25 [0875] Performed in the same manner as in Example 8. MS found for C₁₈H₁₇ClN₄O₂S as (M+H)⁺ 389.1, 391.1.

EXAMPLE 331

5-Chloro-N-((5-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)isoxazol-3-
30 yl)methyl)thiophene-2-carboxamide




[0876] The titled compound was made by the procedure similar to that described in Scheme 12 in Example 330. MS found for $C_{19}H_{17}ClN_4O_2S$ as $(M+H)^+$ 401.0, 403.0.

5

EXAMPLE 332

5-Chloro-N-((5-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)isoxazol-3-yl)methyl)thiophene-2-carboxamide




10

[0877] The titled compound was made by the procedure similar to that described in Scheme 12 in Example 330. MS found for $C_{20}H_{19}ClN_4O_2S$ as $(M+H)^+$ 415.0, 417.0.

EXAMPLE 333

15 5-Chloro-N-((5-(4-(imino(piperidin-1-yl)methyl)phenyl)isoxazol-3-yl)methyl)thiophene-2-carboxamide




[0878] The titled compound was made by the procedure similar to that described in Scheme 20 12 in Example 330. MS found for $C_{21}H_{21}ClN_4O_2S$ as $(M+H)^+$ 429.0, 431.1.

EXAMPLE 334

N-((3-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)-5-chlorothiophene-2-carboxamide (**80**)

5

SCHEME 13



Step 1:

[0879] Combined **3** (4.13 g, 23 mmol) and dioxane (40 mL). To this was added gaseous NH_3 as a steady stream for 2 min during which time a white precipitate formed. The reaction was checked after 5 min by HPLC which showed complete consumption of the starting chloride. The reaction mixture was diluted with a small amount of ether and filtered, the filtrate was then concentrated affording desired product **75** as a light yellow powder (3.27 g, 89%). ^1H NMR ($\text{DMSO}-d_6$, 400 MHz): δ 10.00 (s, 1H), δ 7.80 (m, 4H), 5.95 (s, 2H).

Step 2:

[0880] Boc protected glycine (**76**) (0.54 g, 3.11 mmol) was combined with **75** (0.50 g, 3.11 mmol) and 5 mL of DMF followed by carbonyldiimidazole (0.604 g, 3.73 mmol) which resulted in the evolution of gas. The mixture was stirred at RT for 1 hr, then heated to 110°C . After 30 min, another portion of CDI was added to the heated mixture (0.600 g, 3.73 mmol) again resulting in gas evolution. The reaction was checked by HPLC which showed complete consumption of **75** and a new peak. The reaction mixture was dilution with water and filtered, the solid washed with water and a small amount of methanol (which appeared to dissolve the product) affording the desired product (**77**) as a white solid (0.335 g, 36 %). ^1H

NMR (DMSO-*d*₆, 400 MHz): δ 8.10 (d, 1H), δ 7.99 (d, 2H), 7.72 (t, 1H), δ 4.44 (d, 2H), 1.32 (s, 9H).

Step 3:

[0881] Performed in the same manner as in Step 5, Example 49. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 9.03 (broad s, 3H), δ 8.13 (d, 2H), δ 8.02 (d, 2H), δ 4.54 (s, 2H).

Step 4:


[0882] Performed in the same manner as in Step 5, Example 325. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 9.57 (t, 1H), δ 8.11 (d, 2H), δ 7.99 (d, 2H), δ 7.70 (d, 1H), 7.20 (d, 1H), δ 4.78 (d, 2H).

Step 5:

[0883] Performed in the same manner as in Step 6, Example 325. MS found for C₁₇H₁₆ClN₅O₂S as (M+H)⁺ 390.8, 392.5.

EXAMPLE 335


5-Chloro-N-((3-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)thiophene-2-carboxamide



[0884] The titled compound was made by the procedure similar to that described in Example 334. MS found for C₁₈H₁₆ClN₅O₂S as (M+H)⁺ 402.1, 404.1.

EXAMPLE 336

5-Chloro-N-((3-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)thiophene-2-carboxamide




[0885] The titled compound was made by the procedure similar to that described in Example 334. MS found for $C_{19}H_{18}ClN_5O_2S$ as (M+H)⁺ 416.1, 418.1.

5

EXAMPLE 337

5-Chloro-N-((3-(4-(imino(piperidin-1-yl)methyl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)thiophene-2-carboxamide



10


[0886] The titled compound was made by the procedure similar to that described in Example 334. MS found for $C_{20}H_{20}ClN_5O_2S$ as (M+H)⁺ 430.1, 432.1.

EXAMPLE 338

N-((2-(4-(N,N-Dimethylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide (87).

15

SCHEME 14



Step 1:

[0887] Combined amide **81** (5.00 g, 31 mmol), dibromoacetone **82** (13 g, 62 mmol) and 60 mL of toluene in a flask equipped with a condenser. The mixture was refluxed overnight, then was checked by HPLC the following day which showed complete consumption of the starting material. The reaction was concentrated and purified by silica gel chromatography (dichloromethane as eluent) affording desired product **83** as a beige solid (5.21 g, 60%). ¹H NMR (DMSO-*d*₆, 400 MHz): δ 8.12 (d, 2H), δ 7.71 (m, 3H), δ 4.40 (s, 2H).

Step 2:

[0888] Combined bromide **83** (3.94 g, 15 mmol) and 30 mL DMF. To this was added sodium azide (1.46 g, 23 mmol) and approximately 10 mg of tetrabutylammonium iodide. The reaction was stirred 1 hr, then was checked by mass spectroscopy which showed only the desired azide. The reaction was diluted with 100 mL of water and filtered affording the desired product as a beige solid. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 8.31 (s, 1H), δ 8.08 (d, 2H), δ 7.93 (d, 2H), δ 4.41 (s, 1H).

Step 3:

[0889] Diluted azide **84** (1.19 g, 5.3 mmol) with 20 mL of ethyl acetate. The solution was degassed with argon, treated with ca. 100 mg of 10% Pd/C then purged with hydrogen gas (balloon). After two hours all starting material had been consumed as determined by TLC. The mixture was filtered through a short pad of celite, concentrated to a beige semisolid and used immediately for the next step.

Step 4:


[0890] The titled compound was made by the procedure similar to that described in Step 5, Example 325. MS ¹H NMR (DMSO-*d*₆, 400 MHz): δ 9.11 (t, 1H), δ 8.14 (s, 1H), δ 8.08 (d, 2H), δ 7.93 (d, 2H), δ 7.64 (d, 1H), δ 7.13 (d, 1H), δ 4.35 (d, 2H).

Step 5:

[0891] The titled compound was made by the procedure similar to that described in Step 6, Example 325. MS found for C₁₈H₁₇ClN₄O₂S as (M+H)⁺ 389.1, 391.1.

EXAMPLE 339

5-Chloro-N-((2-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)oxazol-4-yl)methyl)thiophene-2-carboxamide




[0892] The titled compound was made by the procedure similar to that described in Example 338. MS found for $C_{19}H_{17}ClN_4O_2S$ as $(M+H)^+$ 401.1, 403.1.

5

EXAMPLE 340

5-Chloro-N-((2-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)oxazol-4-yl)methyl)thiophene-2-carboxamide




10

[0893] The titled compound was made by the procedure similar to that described in Example 338. MS found for $C_{20}H_{19}ClN_4O_2S$ as $(M+H)^+$ 415.1, 417.1.

EXAMPLE 341

15 5-Chloro-N-((2-(4-(imino(piperidin-1-yl)methyl)phenyl)oxazol-4-yl)methyl)thiophene-2-carboxamide




[0894] The titled compound was made by the procedure similar to that described in Example 338. MS found for $C_{21}H_{21}ClN_4O_2S$ as $(M+H)^+$ 429.1, 431.1.

20

EXAMPLE 342

Ethyl 1-((4-(4-((2-chlorothiophene-5-carboxamido)methyl)oxazol-2-yl)phenyl)(imino)methyl)piperidine-4-carboxylate




5

[0895] The titled compound was made by the procedure similar to that described in Example 338. MS found for $C_{24}H_{25}ClN_4O_4S$ as $(M+H)^+$ 501.1, 503.1.

EXAMPLE 343


10 N-((2-(4-(N-Ethyl-N-methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



15 [0896] The titled compound was made by the procedure similar to that described in Example 338. MS found for $C_{19}H_{19}ClN_4O_2S$ as $(M+H)^+$ 403.1, 405.1.

EXAMPLE 344

N-((2-(4-(N-Methyl-N-propylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide




20

[0897] The titled compound was made by the procedure similar to that described in Example 338. MS found for $C_{20}H_{21}ClN_4O_2S$ as $(M+H)^+$ 417.1, 419.1.

EXAMPLE 345


N-((2-(4-(N-(2-(Dimethylamino)ethyl)-N-methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



[0898] The titled compound was made by the procedure similar to that described in Example 338. MS found for $C_{21}H_{24}ClN_5O_2S$ as $(M+H)^+$ 446.1, 448.1.

EXAMPLE 346


N-((2-(4-(N-(2-Methoxyethyl)-N-methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



[0899] The titled compound was made by the procedure similar to that described in Example 338. MS found for $C_{20}H_{21}ClN_4O_3S$ as $(M+H)^+$ 433.1, 435.1.

EXAMPLE 347


1-((4-(4-((2-Chlorothiophene-5-carboxamido)methyl)oxazol-2-yl)phenyl)(imino)methyl)piperidine-4-carboxamide



[0900] The titled compound was made by the procedure similar to that described in Example 338. MS found for $C_{22}H_{22}ClN_5O_3S$ as $(M+H)^+$ 472.0, 474.0.

EXAMPLE 348


5 N-((2-(4-(N-Methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



[0901] The titled compound was made by the procedure similar to that described in Example 338. MS found for $C_{17}H_{15}ClN_4O_2S$ as $(M+H)^+$ 375.0, 377.0.

EXAMPLE 349

N-((2-(4-(N-(Furan-2-ylmethyl)-N-methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide




[0902] The titled compound was made by the procedure similar to that described in Example 338. MS found for $C_{22}H_{19}ClN_4O_3S$ as $(M+H)^+$ 455.1, 457.1.

EXAMPLE 350

N-((2-(4-(N,N-Dimethylcarbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide (94)

SCHEME 15



Step 1:

[0903] Combined amide **81** (0.64 g, 4.1 mmol), Lawesson's reagent (1.00 g, 2.4 mmol) and 15 mL of toluene in a flask equipped with a condenser. The resulting beige suspension was refluxed for two hrs at which time all starting amide was consumed as determined by HPLC. The reaction was cooled, filtered through silica gel and concentrated, then purified by eluting through a short plug of silica gel (eluted with dichloromethane) affording the desired thioamide (**88**) as an orange solid (0.260 g, 39%). ¹H NMR (DMSO-*d*₆, 400 MHz): δ 10.12 (s, 1H), δ 9.71 (s, 1H), δ 7.90 (d, 2H), δ 7.84 (d, 2H).

Step 2:

[0904] Thioamide **88** (0.48 g, 30 mmol), dichloroacetone **89** (0.45 g, 36 mmol) and 20 mL of ethanol were combined in a flask equipped with a reflux condenser. The mixture was refluxed for three hours at which time all starting thioamide was consumed as determined by TLC. The mixture was cooled, partitioned with water and ethyl acetate and separated. The aqueous phase was extracted again with ethyl acetate and the combined organic layers were dried over Na₂SO₄, filtered through a short pad of silica gel and concentrated affording a quantitative amount of thiazole **90** as an off-white solid. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 8.06 (d, 2H), δ 7.65 (d, 2H), δ 7.39 (s, 1H), δ 4.30 (s, 2H).

Step 3:

[0905] Performed in the same manner as in Step 2, Example 338. $C_{11}H_7N_5S$ as $(M+H)^+$ 242.3.

Step 4:

5 [0906] Performed in the same manner as in Step 3, Example 338. ^1H NMR ($DMSO-d_6$, 400 MHz): δ 8.09 (d, 2H), δ 7.78 (d, 2H), δ 7.44 (s, 1H), 4.88 (s, 2H), δ 3.90 (s, 2H).

Step 5:

10 [0907] Performed in the same manner as in Step 4, Example 338. ^1H NMR ($CDCl_3$, 400 MHz): δ 8.00 (d, 2H), δ 7.68 (d, 2H), δ 7.27 (m, 2H), δ 6.89 (s, 1H), δ 6.65 (broad s, 1H), δ 4.70 (m, 2H).


Step 6:

[0908] Performed in the same manner as in Step 5, Example 338. MS found for $C_{18}H_{17}ClN_4OS_2$ as $(M+H)^+$ 405.0, 407.0.

15

EXAMPLE 351

5-Chloro-N-((2-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)thiazol-4-yl)methyl)thiophene-2-carboxamide




20

[0909] The titled compound was made by the procedure similar to that described in Example 350. MS found for $C_{19}H_{17}ClN_4OS_2$ as $(M+H)^+$ 417.0, 419.0.

EXAMPLE 352

5-Chloro-N-((2-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)thiazol-4-yl)methyl)thiophene-2-carboxamide


25



[0910] The titled compound was made by the procedure similar to that described in Example 350. MS found for $C_{20}H_{19}ClN_4OS_2$ as $(M+H)^+$ 431.0, 433.0.

EXAMPLE 353


- 5 5-Chloro-N-((2-(4-(imino(piperidin-1-yl)methyl)phenyl)thiazol-4-yl)methyl)thiophene-2-carboxamide



- 10 [0911] The titled compound was made by the procedure similar to that described in Example 350. MS found for $C_{21}H_{21}ClN_4OS_2$ as $(M+H)^+$ 445.1, 447.0.

EXAMPLE 354

Ethyl 1-((4-(4-((2-chlorothiophene-5-carboxamido)methyl)thiazol-2-yl)phenyl)(imino)methyl)piperidine-4-carboxylate




15

[0912] The titled compound was made by the procedure similar to that described in Example 350. MS found for $C_{24}H_{25}ClN_4O_3S_2$ as $(M+H)^+$ 517.1, 519.1.

20

EXAMPLE 355


N-((2-(4-(N-(2-(Dimethylamino)ethyl)-N-methylcarbamidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



[0913] The titled compound was made by the procedure similar to that described in Example 350. MS found for $C_{21}H_{24}ClN_5OS_2$ as $(M+H)^+$ 462.1, 464.1.

EXAMPLE 356


5 N-((2-(4-(N-(Furan-2-ylmethyl)-N-methylcarbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



[0914] The titled compound was made by the procedure similar to that described in Example 350. MS found for $C_{22}H_{19}ClN_4O_2S_2$ as $(M+H)^+$ 471.0, 473.0.

EXAMPLE 357


N-((2-(4-(N-Methyl-N-(prop-2-ynyl)carbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



[0915] The titled compound was made by the procedure similar to that described in Example 350. MS found for $C_{20}H_{17}ClN_4OS_2$ as $(M+H)^+$ 429.0, 431.0.

EXAMPLE 358

20 N-((2-(4-(N-Methylcarbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide




[0916] The titled compound was made by the procedure similar to that described in Example 350. MS found for $C_{17}H_{15}ClN_4OS_2$ as $(M+H)^+$ 391.0, 393.0.

EXAMPLE 359

- 5 5-Chloro-N-((4-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)oxazol-2-yl)methyl)thiophene-2-carboxamide (103)

SCHEME 16



Step 1:

[0917] Compound **95** (2.24 g, 10 mmol) and compound **96** (0.59 g, 10 mmol) were refluxed in ethanol for overnight. The mixture was concentrated in vacuo and subjected to flash column chromatography to give compound **97**. ^1H NMR (DMSO- d_6 , 400 MHz): δ 8.63 (s, 1H), δ 7.89 (d, 2H), δ 7.83 (d, 2H), δ 2.42 (s, 3H).

Step 2:

[0918] Combined **97** (0.115 g, 0.62 mmol), N-bromosuccinimide (0.11 g, 0.62 mmol) and carbontetrachloride in a flask equipped with a reflux condenser. To this was added a small amount (ca. 10 mg) of benzoyl peroxide and the mixture heated to reflux overnight. The

following day the reaction was checked by TLC which showed complete consumption of the starting material and a new, less polar spot. The crude reaction mixture was loaded onto a silica gel plug and eluted with dichloromethane affording the 5-substituted bromide **98** as the major product as determined by NMR. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 7.99 (d, 2H), δ

7.90 (d, 2H), δ 2.44 (s, 3H).

Step 3:

[0919] Performed in the same manner as in Step 2 to get compound **99**. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 8.03 (d, 2H), δ 7.93 (d, 2H), δ 4.80 (s, 2H).

Step 4:

[0920] Performed in the same manner as in Step 2, Example 338. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 8.08 (d, 2H), δ 7.71 (d, 2H), δ 4.47 (s, 2H).

Step 5:

[0921] Performed in the same manner as in Step 3, Example 338. ¹H NMR (DMSO-*d*₆, 400 MHz): δ 8.91 (s, 1H), δ 8.50 (broad s, 3H), δ 7.91 (m, 4H), δ 4.32 (s, 2H).

Step 6:


[0922] Performed in the same manner as in Step 4, Example 338. ¹H NMR (CD₃OD, 400 MHz): δ 8.71 (s, 1H), δ 8.05 (d, 2H), δ 7.90 (d, 2H), δ 7.77 (d, 1H), δ 7.21 (d, 1H), δ 7.76 (d, 2H).

Step 7:

[0923] Performed in the same manner as in Step 5, Example 338. MS found for C₂₀H₁₉ClN₄O₂S as (M+H)⁺ 415.0, 417.0.

EXAMPLE 360


N-((4-(4-(N,N-Dimethylcarbamimidoyl)phenyl)thiazol-2-yl)methyl)-5-chlorothiophene-2-carboxamide



[0924] The titled compound was made by the procedure similar to that described in Example 359. MS found for C₁₈H₁₇ClN₄OS₂ as (M+H)⁺ 405.0, 407.0.

EXAMPLE 361

5-Chloro-N-((4-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)thiazol-2-yl)methyl)thiophene-2-carboxamide




5

[0925] The titled compound was made by the procedure similar to that described in Example 359. MS found for $C_{19}H_{17}ClN_4OS_2$ as $(M+H)^+$ 417.0, 419.0.

EXAMPLE 362


10 5-Chloro-N-((4-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)thiazol-2-yl)methyl)thiophene-2-carboxamide



15 [0926] The titled compound was made by the procedure similar to that described in Example 359. MS found for $C_{20}H_{19}ClN_4OS_2$ as $(M+H)^+$ 431.0, 433.0.

EXAMPLE 363

5-Chloro-N-((4-(4-(imino(piperidin-1-yl)methyl)phenyl)thiazol-2-yl)methyl)thiophene-2-carboxamide



20


[0927] The titled compound was made by the procedure similar to that described in Example 359. MS found for $C_{21}H_{21}ClN_4OS_2$ as $(M+H)^+$ 445.0, 447.0.

EXAMPLE 364

N-((1-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-1H-imidazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide (**110**)

5

SCHEME 17



Step 1:

[0928] 4-Iodobenzonitrile (**45**, 11.9 g, 52 mmol) and 4-methylimidazole (**104**, 5.16 g, 62.8 mmol) were dissolved in 100 mL dry dioxane. To it were added trans-1,2-cyclohexanediamine (3.8 mL, 31 mmol), CuI (1.18 mg, 6.2 mmol) and K₂CO₃ (14.7 g, 106 mmol). The mixture was refluxed for overnight. It was taken into 1000 mL EtOAc, washed with water and brine, dried, concentrated in vacuo and subjected to flash column chromatography to give compound **105** (8.21 g, 86%). MS found for C₁₁H₉N₃ as (M+H)⁺ 184.1.

Step 2:

[0929] Compound **105** (1.88 g, 10.3 mmol) was dissolved in 50 mL CCl₄. To it were added NBS (7.05 g, 39.6 mmol) and benzoyl peroxide (480 mg, 1.98 mmol). The mixture was refluxed for 3 hr. It was concentrated in vacuo and subjected to silica flash column chromatography to give compound **106** (60%) as the major product. MS found for C₁₁H₆Br₃N₃ as (M+H)⁺ 417.8, 419.8, 421.8.

Step 3:

[0930] Compound **106** (820 mg, 1.95 mmol) was dissolved in 10 mL DMF. To it was added NaN₃ (240 mg, 3.7 mmol). The mixture was stirred for 3 hr and concentrated in vacuo. The residue was taken into 300 mL EtOAc and washed with water and brine. The organic phase

Step 4:

[0931] Compound **107** (390 mg, 1.02 mmol) was dissolved in 20 mL methanol. To it were added triethylamine (0.6 mL, 4.1 mmol) and 10% Pd on carbon (337 mg). The mixture was shaken under 10 PSI hydrogen for 30 min. It was then filtered through celite and concentrated in vacuo to give compound **108** as a yellow solid. MS found for C₁₁H₁₀N₄ as (M+H)⁺ 199.1.

Step 5:

[0932] The above crude compound **108** was dissolved in 10 mL DMF. To it were added 5-chlorothiophene-2-carboxylic acid (231 mg, 1.42 mmol), EDC (277 mg, 1.44 mmol), 1-hydroxy-7-azabenzotriazole (HOAt, 17 mg, 0.12 mol) and triethylamine (0.8 mL, 5.7 mmol).


[0933] The mixture was stirred for overnight at RT and was diluted with brine. The mixture was extracted with DCM three times. The organic phase was combined, dried, concentrated and subjected to flash column to give compound **109** (180 mg, 52% for 2 steps). MS found for C₁₆H₁₁ClN₄OS as (M+H)⁺ 343.0, 345.0.

Step 6:

[0934] Performed in the same manner as in Step 6, Example 325. MS found for C₁₈H₁₈ClN₅OS as (M+H)⁺ 388.1, 390.1.

EXAMPLE 365


5-Chloro-N-((1-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1H-imidazol-4-yl)methyl)thiophene-2-carboxamide



[0935] The titled compound was made by the procedure similar to that described in Example 364. MS found for C₁₉H₁₈ClN₅OS as (M+H)⁺ 400.1, 402.1.

EXAMPLE 366


N-((1-(4-(Azetidin-1-yl(imino)methyl)phenyl)-1H-imidazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



- 5 [0936] The titled compound was made by the procedure similar to that described in Example 364. MS found for $C_{19}H_{18}ClN_5OS$ as (M+H)⁺ 400.1, 402.1.

EXAMPLE 367


5-Chloro-N-((1-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1H-imidazol-4-yl)methyl)thiophene-2-carboxamide



- 10 [0937] The titled compound was made by the procedure similar to that described in Example 364. MS found for $C_{20}H_{20}ClN_5OS$ as (M+H)⁺ 414.1, 416.1.

EXAMPLE 368

15 5-Chloro-N-((1-(4-(imino(piperidin-1-yl)methyl)phenyl)-1H-imidazol-4-yl)methyl)thiophene-2-carboxamide




- 20 [0938] The titled compound was made by the procedure similar to that described in Example 364. MS found for $C_{21}H_{22}ClN_5OS$ as (M+H)⁺ 428.1, 430.1.

EXAMPLE 369

N-((5-(4-(N,N-Dimethylcarbamimidoyl)phenyl)pyridin-3-yl)methyl)-5-chlorothiophene-2-carboxamide (**115**)

5

SCHEME 18



Step 1:

[0939] Compound **111** (2.00 g, 10.6 mmol) and triphenylphosphine (3.31 g, 12.6 mmol) were dissolved in 100 mL THF and 50 mL DCM. At 0°C, to it was added DEAD (3.2 mL, 20 mmol) dropwise. After stirring for 2 min, DPPA (4.3 mL, 20 mmol) was added. The mixture was stirred for overnight (from 0°C to RT). The mixture was diluted with water, and it was extracted with DCM three times. The organic phase was combined, dried, concentrated and purified using flash column to give compound **112** (690 mg, 30%). MS found for C₆H₅BrN₄ as (M+H)⁺ 213.0, 215.0.

Step 2:

[0940] Compound **112** (675 mg, 3.17 mmol) was dissolved in 10 mL THF and 1 mL water. To it was added triphenylphosphine (0.96 g, 3.67 mmol). The mixture was stirred for 24 hrs at RT. Then to it were added EDC (617 mg, 3.22 mmol), 5-chlorothiophene-2-carboxylic acid (502 mg, 3.1 mmol) and HOAt (70 mg, 0.51 mmol). The mixture was stirred for overnight at RT. The mixture was diluted with 2N NaOH, and was extracted with DCM three times. The organic phase was combined, dried, concentrated and purified using flash column to give compound **113** (450 mg, 43%). MS found for C₁₁H₈BrClN₂OS as (M+H)⁺ 331.0, 333.0.

25 Step 3:


[0941] A mixture of compound **113** (450 mg, 1.35 mmol), 4-cyanobenzeneboronic acid (282 mg, 1.92 mmol), Pd(Ph₃P)₄ (130 mg, 0.11 mmol), K₂CO₃ (596 mg, 4.31 mmol) in 5 mL Diglyme and 5 mL water was degasses using argon stream for 5 min. The mixture was then stirred in 80°C bath for overnight. The mixture was poured into water and was extracted with DCM three times. The organic phase was combined, dried, concentrated and purified using flash column to give compound **114** (350 mg, 73%). MS found for C₁₈H₁₂ClN₃OS as (M+H)⁺ 354.0, 356.0.

Step 4:

[0942] Performed in the same manner as in Step 6, Example 325. MS found for C₂₀H₁₉ClN₄OS as (M+H)⁺ 399.1, 401.1.

EXAMPLE 370


5-Chloro-N-((5-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)pyridin-3-yl)methyl)thiophene-2-carboxamide



[0943] The titled compound was made by the procedure similar to that described in Example 369. MS found for C₂₁H₁₉ClN₄OS as (M+H)⁺ 411.1, 413.1.

EXAMPLE 371


N-((5-(4-(Azetidin-1-yl(imino)methyl)phenyl)pyridin-3-yl)methyl)-5-chlorothiophene-2-carboxamide



[0944] The titled compound was made by the procedure similar to that described in Example 369. MS found for C₂₁H₁₉ClN₄OS as (M+H)⁺ 411.1, 413.1.

EXAMPLE 372


5-Chloro-N-((5-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)pyridin-3-yl)methyl)thiophene-2-carboxamide



- 5 [0945] The titled compound was made by the procedure similar to that described in Example 369. MS found for $C_{22}H_{21}ClN_4OS$ as $(M+H)^+$ 425.1, 427.1.

EXAMPLE 373

5-Chloro-N-((5-(4-(imino(piperidin-1-yl)methyl)phenyl)pyridin-3-yl)methyl)thiophene-2-carboxamide




- 10 [0946] The titled compound was made by the procedure similar to that described in Example 369. MS found for $C_{23}H_{23}ClN_4OS$ as $(M+H)^+$ 439.1, 441.1.

EXAMPLE 374

N-((5-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-2-fluorophenyl)methyl)-5-chlorothiophene-2-carboxamide (119)

SCHEME 19



Step 1:

[0947] Compound **116** hydrochloride (490 mmol, 2.04 mmol), 5-chlorothiophene-2-carboxylic acid (390 mg, 2.40 mmol) and triethylamine (0.35 mL, 2.51 mmol) were dissolved in 20 mL DCM. To it were added DMAP (25 mg) and EDC (450 mg, 2.35 mmol). The mixture was stirred for overnight. It was diluted with DCM, washed with water, dried, concentrated and purified using flash column to give compound **117** (90%). MS found for $C_{12}H_8BrClFNOS$ as (M+H)⁺ 348.0, 350.0.

Step 2:


[0948] Performed in the same manner as in Step 3, Example 369. MS found for $C_{19}H_{12}ClFN_2OS$ as (M+H)⁺ 371.0, 373.0

Step 3:

[0949] Performed in the same manner as in Step 4, Example 369. MS found for $C_{21}H_{19}ClFN_3OS$ as (M+H)⁺ 416.1, 418.1.

EXAMPLE 375


5-Chloro-N-((5-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-2-fluorophenyl)methyl)thiophene-2-carboxamide



[0950] The titled compound was made by the procedure similar to that described in Example 374. MS found for $C_{22}H_{19}ClFN_3OS$ as (M+H)⁺ 428.1, 430.1.

EXAMPLE 376


5-Chloro-N-((5-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-2-fluorophenyl)methyl)thiophene-2-carboxamide



[0951] The titled compound was made by the procedure similar to that described in Example 374. MS found for $C_{23}H_{21}ClFN_3OS$ as (M+H)⁺ 442.1, 444.1.

EXAMPLE 377

- 5 N-((3-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-4-fluorophenyl)methyl)-5-chlorothiophene-2-carboxamide




[0952] The titled compound was made by the procedure similar to that described in Example 374. MS found for $C_{21}H_{19}ClFN_3OS$ as (M+H)⁺ 416.1, 418.1.

10

EXAMPLE 378

- 5-Chloro-N-((3-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-4-fluorophenyl)methyl)thiophene-2-carboxamide




- 15 [0953] The titled compound was made by the procedure similar to that described in Example 374. MS found for $C_{22}H_{19}ClFN_3OS$ as (M+H)⁺ 428.1, 430.1.

EXAMPLE 379

- 5-Chloro-N-((3-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-4-fluorophenyl)methyl)thiophene-2-carboxamide


20



[0954] The titled compound was made by the procedure similar to that described in Example 374. MS found for $C_{23}H_{21}ClFN_3OS$ as (M+H)⁺ 442.1, 444.1.

EXAMPLE 380

N-((3-(4-(N,N-Dimethylcarbamimidoyl)phenyl)phenyl)methyl)-5-chlorothiophene-2-carboxamide




5

[0955] The titled compound was made by the procedure similar to that described in Example 374. MS found for $C_{21}H_{20}ClN_3OS$ as (M+H)⁺ 398.1, 400.1.

EXAMPLE 381

5-Chloro-N-((3-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)phenyl)methyl)thiophene-2-carboxamide




10

[0956] The titled compound was made by the procedure similar to that described in Example 374. MS found for $C_{22}H_{20}ClN_3OS$ as (M+H)⁺ 410.1, 412.1.

15

EXAMPLE 382

5-Chloro-N-((3-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)phenyl)methyl)thiophene-2-carboxamide



20


[0957] The titled compound was made by the procedure similar to that described in Example 374. MS found for $C_{23}H_{22}ClN_3OS$ as (M+H)⁺ 424.1, 426.1.

EXAMPLE 383

5-Chloro-N-((1-(6-(4-methyl-1,4-diazepan-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide (**123**)

5

SCHEME 20



Step 1:

[0958] 5-Amino-2-chloropyridine (**120**, 4.91 g, 38.3 mmol) was dissolved in 15 mL TFA. In ice bath to it was added NaNO_2 (2.91 g, 42.2 mmol) in small portions. The mixture was stirred in the cold bath for 30 min. To it was then added an ice-cold solution of NaN_3 (2.50 g, 38.3 mmol, in 10 mL water). The mixture was stirred for 1 hr. The mixture was concentrated in vacuo and diluted with 400 mL chloroform. It was washed with water three times, dried, concentrated and purified by flash column to yield compound **121** (3.46 g, 59%). MS found for $\text{C}_5\text{H}_3\text{ClN}_4$ as $(\text{M}+\text{H})^+$ 155.0, 157.0.

Step 2:

[0959] Azidopyridine **121** (3.46 g, 22.5 mmol) and alkyne **34** (see Scheme 5 in Example 291; 4.38 g, 22.0 mmol) were refluxed in 10 mL toluene and 15 mL DMF for 7 hr. The reaction was still in complete. The mixture was concentrated and subjected to flash column to isolate major 1,4-disubstituted triazole product **122** (3.19 g, 40%) as a white solid. MS found for $\text{C}_{13}\text{H}_9\text{Cl}_2\text{N}_5\text{OS}$ as $(\text{M}+\text{H})^+$ 354.0, 356.0. The minor 1,5-triazole isomer was isolated in 7% yield (0.56 g).


Step 3:

[0960] The mixture of compound **122** (50 mg, 0.14 mmol), N-methylhomopiperazine (53 μL , 0.42 mmol) and Cs_2CO_3 (91 mg, 0.28 mmol) in 1 mL DMSO in a sealed tube was stirred in 120°C bath for 2 hr. The mixture was directly subjected to reverse phase preparative HPLC to

give the title compound as a white powder after lyophilization. MS found for $C_{19}H_{22}ClN_7OS$ as $(M+H)^+$ 432.1, 434.1.

EXAMPLE 384

- 5 N-((1-(6-(1,4-Diazepan-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide




[0961] The titled compound was made by the procedure similar to that described in Example 383. MS found for $C_{18}H_{20}ClN_7OS$ as $(M+H)^+$ 418.1, 420.1.

10

EXAMPLE 385


- N-((1-(6-(1,4-Oxazepan-4-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide



- 15 [0962] The titled compound was made by the procedure similar to that described in Example 383. MS found for $C_{18}H_{19}ClN_6O_2S$ as $(M+H)^+$ 419.1, 421.1.

EXAMPLE 386


- 20 5-Chloro-N-((1-(6-morpholinopyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



[0963] The titled compound was made by the procedure similar to that described in Example 383. MS found for $C_{17}H_{17}ClN_6O_2S$ as (M+H)⁺ 405.1, 407.1.

EXAMPLE 387

- 5 5-Chloro-N-((1-(6-(3-oxomorpholino)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




[0964] The titled compound was made by the procedure similar to that described in Example 383. MS found for $C_{17}H_{15}ClN_6O_3S$ as (M+H)⁺ 419.1, 421.1.

10

EXAMPLE 388

- 5-Chloro-N-((1-(6-(2-oxopiperidin-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide




- 15 [0965] The titled compound was made by the procedure similar to that described in Example 383. MS found for $C_{18}H_{17}ClN_6O_2S$ as (M+H)⁺ 417.1, 419.1.

EXAMPLE 389

- 5-Chloro-N-((1-(6-(2-oxopyridin-1(2H)-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide


20



[0966] The titled compound was made by the procedure similar to that described in Example 383. MS found for $C_{18}H_{13}ClN_6O_2S$ as (M+H)⁺ 413.0, 415.0.

EXAMPLE 390


5-Chloro-N-((1-(6-(piperazin-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



[0967] The titled compound was made by the procedure similar to that described in Example 383. MS found for $C_{17}H_{18}ClN_7OS$ as (M+H)⁺ 404.1, 406.1.

EXAMPLE 391


5-Chloro-N-((1-(6-(4-methylpiperazin-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



[0968] The titled compound was made by the procedure similar to that described in Example 383. MS found for $C_{18}H_{20}ClN_7OS$ as (M+H)⁺ 418.1, 420.1.

EXAMPLE 392


5-Chloro-N-((1-(6-(piperidin-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide



[0969] The titled compound was made by the procedure similar to that described in Example 383. MS found for $C_{18}H_{19}ClN_6OS$ as (M+H)⁺ 403.1, 405.1.

EXAMPLE 393


5-Chloro-N-((3-(4-(8-(trifluoromethyl)quinolin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



- 5 [0970] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{25}H_{18}ClF_3N_4O_2S$ as $(M+H)^+$ 531.0, 533.0.

EXAMPLE 394


5-Chloro-N-((3-(4-(7-(trifluoromethyl)quinolin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



- [0971] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{25}H_{18}ClF_3N_4O_2S$ as $(M+H)^+$ 531.0, 533.0.

EXAMPLE 395


5-Chloro-N-((3-(4-(6-(trifluoromethyl)quinolin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



[0972] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{25}H_{18}ClF_3N_4O_2S$ as $(M+H)^+$ 531.0, 533.0.

EXAMPLE 396

- 5 5-Chloro-N-((3-(4-(quinolin-6-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide




[0973] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{24}H_{19}ClN_4O_2S$ as $(M+H)^+$ 463.1, 465.1.

10

EXAMPLE 397


- 5-Chloro-N-((3-(4-(2,5-dihydro-1H-pyrrole-1-carbonyl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



- 15 [0974] The titled compound was made by the procedure similar to that described in Example 162. MS found for $C_{20}H_{18}ClN_3O_3S$ as $(M+H)^+$ 416.0, 418.0.

EXAMPLE 398


- 20 5-Chloro-N-((3-(4-(2,5-dihydro-1H-pyrrole-1-carbonyl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



[0975] The titled compound was made by the procedure similar to that described in Example 162. MS found for $C_{22}H_{24}ClN_3O_4S$ as $(M+H)^+$ 462.0, 464.0.

EXAMPLE 399

5 N-((3-(4-((2-Hydroxypropyl)carbamoyl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)-5-chlorothiophene-2-carboxamide




[0976] The titled compound was made by the procedure similar to that described in Example 162. MS found for $C_{19}H_{20}ClN_3O_4S$ as $(M+H)^+$ 422.0, 424.0.

10

EXAMPLE 400

5-Chloro-N-((3-(4-(3-methylpyridin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide




15 [0977] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{21}H_{19}ClN_4O_2S$ as $(M+H)^+$ 427.0, 429.0.

EXAMPLE 401

5-Chloro-N-((3-(4-(2-ethoxypyridin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide


20



[0978] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{22}H_{21}ClN_4O_3S$ as $(M+H)^+$ 457.1, 459.1.

EXAMPLE 402

5 5-Chloro-N-((3-(4-(3,5-dichloropyridin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide




[0979] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{20}H_{15}Cl_3N_4O_2S$ as $(M+H)^+$ 481.0, 483.0.

10

EXAMPLE 403

5-Chloro-N-((3-(4-(3-(trifluoromethyl)pyridin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide



15 [0980] The titled compound was made by the procedure similar to that described in Example 222. MS found for $C_{21}H_{16}ClF_3N_4O_2S$ as $(M+H)^+$ 481.1, 483.1.

EXAMPLE 404

[0981] This example illustrates methods for evaluating the compounds of the invention, along with results obtained for such assays. The *in vitro* and *in vivo* Factor Xa isoform activities of the inventive compounds can be determined by various procedures known in the art, such as a test for their ability to inhibit the activity of the Factor Xa isoform. The potent affinities for Factor Xa isoform exhibited by the inventive compounds can be measured by an IC_{50} value (in nM). The IC_{50} value is the concentration (in nM) of the compound required to

provide 50% inhibition of Factor Xa isoform. The smaller the IC_{50} value, the more active (potent) is a compound for inhibiting Factor Xa isoform.

[0982] An *in vitro* assay for detecting and measuring inhibition activity against Factor Xa is as follows:

5 *IC₅₀ and Ki Determinations:*

Substrate:

The substrate S-2765 (Z-D-Arg-Gly-Arg-pNA·HCl) was obtained from Diapharma (West Chester, OH).

10 Enzyme:

The human plasma protein factor Xa was purchased from Haematologic Technologies (Essex Junction, VT).

Methods

15 *IC₅₀ determinations*

[0983] All assays, which are performed in 96-well microtiter plates, measure proteolytic activity of the enzyme (factor Xa) by following cleavage of paranitroanilide substrate. The assay buffer used for proteolytic assays was Tris buffered saline (20mM Tris, 150mM NaCl, 5mM $CaCl_2$, 0.1% Bovine serum albumin (BSA), 5% Dimethyl Sulfoxide (DMSO) pH 7.4).

20 In a 96-well microtiter plate, inhibitor was serially diluted to give a range of concentrations from 0.01nM to 10 μ M (final). Duplicate sets of wells were assayed and control wells without inhibitor were included. Enzyme was added to each well, (fXa concentration = 1nM), the plate was shaken for 5 seconds and then incubated for 5 minutes at room temperature. S2765 was added (100 μ M final) and the plate was shaken for 5 seconds (final liquid volume
25 in each well was 200 μ l). The degree of substrate hydrolysis was measured at 405 nm on a Thermomax plate reader (Molecular Devices, Sunnyvale, CA) for 2 minutes. The initial velocities (mOD/min), for each range of inhibitor concentrations, were fitted to a four parameter equation using Softmax data analysis software. The parameter C, derived from the resulting curve-fit, corresponded to the concentration for half maximal inhibition (IC_{50}).

30

K_i determination

[0984] The assay buffer for this series of assays was Hepes buffered saline (20mM Hepes, 150mM NaCl, 5mM $CaCl_2$, 0.1% PEG-8000, pH 7.4). In a 96-well microtiter plate, inhibitor

was serially diluted in a duplicate set of wells to give a range of final concentrations from 5pM to 3µM final. Controls without inhibitor (8 wells) were included. The enzyme, fXa (1nM final) was added to the wells. The substrate S-2765 (200µM final) was added and the degree of substrate hydrolysis was measured at 405nm on a Thermomax plate reader for 5 minutes, using Softmax software. Initial velocities (mOD/min) were analyzed by non-linear least squares regression in the Plate Ki software (BioKin Ltd, Pullman, WA)(Literature reference: Kusmic P, Sideris S, Cregar LM, Elrod KC, Rice KD, Janc J. High-throughput screening of enzyme inhibitors: Automatic determination of tight-binding inhibition constants. *Anal. Biochemistry* 2000, **281**:62-67). The model used for fitting the inhibitor dose-response curves was the Morrison equation. An apparent K_i (K_i^*) was determined. The overall K_i was calculated using the following equation:

$$K_i = \frac{K_i^*}{1 + \frac{[S]}{K_m}}$$

where [S] is substrate concentration (200µM) and K_m , the Michaelis constant for S2765.

The hERG (human ether-a-go-go related gene protein) Membrane Binding Assay

[0985] Human embryonic kidney (HEK293) cells stably transfected with hERG cDNA were used for preparation of membranes (Literature reference: Zhou, Z., Gong, Q., Ye, B., Fan, Z., Makielski, C., Robertson, G., January, CT., Properties of HERG stably expressed in HEK293 cells studied at physiological temperature. *Biophys. J*, 1998, **74**:230-241). The assay buffer was comprised of 50 mM Tris, 10 mM KCl, 1 mM MgCl₂, pH 7.4. Competition assays for hERG binding were performed, in a 96 well plate, with 50 µL ³H- dofetilide, at a concentration of 3.5 nM (final concentration of 0.01% ethanol). Test compound was added at final concentrations of 100 µM, 33.33 µM, 11.11 µM, 3.70 µM, 1.23 µM, 0.41 µM, 0.14 µM, 0.046 µM, 0.015 µM, and 0.005 µM (1.0% DMSO). Each compound was run in duplicate on each of two plates. Total binding was determined by addition of 50 µL of assay buffer in place of compound. Non-specific binding was determined by addition of 50 µL of 50 µM terfenadine in place of test compound. All assays were initiated by addition of 150 µL of membrane homogenates (15 ug protein/well as final concentration) to the wells (total volume = 250 µL per well), and the plates were incubated at room temperature for 80 minutes on a shaking platform. All assays were terminated by vacuum filtration on to glass fiber filters, followed by two washes with cold assay buffer. The filter plates were dried at 55°C for 90

minutes, after which, Microscint 0 (50 μ L) was added to each well of the dried filter plate. The plates were counted on a Packard Topcount (Perkin Elmer, Boston, MA) using a one minute protocol. Scintillation reading (counts per minute, CPM) data generated by the Packard TopCount was used to calculate the percent inhibition of ^3H -dofetilide binding, for each compound at each concentration, using the total binding control value corrected for non-specific binding. The IC_{50} value was calculated from the percent inhibition curve generated using Excel XL Fit software (Microsoft). The equilibrium dissociation constant (K_i) was calculated using the equation of Cheng and Prusoff (see "Relationship between the inhibition constant (K_i) and the concentration of inhibitor which causes 50 per cent inhibition (I_{50}) of an enzymatic reaction," *Biochem Pharmacol.*, 1973, **22(23)**:3099-108.

$$K_i = \text{IC}_{50} / [1 + ([L] / K_D)].$$

[0986] A compound can be run through this assay and its corresponding IC_{50} inhibition value can be calculated from the assay data.

[0987] The following examples exhibited Factor Xa IC_{50} values less than or equal to 100 nM:

7 – 10, 16 – 18, 20 – 32, 48 – 51, 56 – 65, 67, 69 – 72, 77, 82, 86, 91 – 142, 161 – 164, 166, 193 – 200, 203 – 205, 208, 209, 216 – 219, 225 – 227, 239, 240, 242, 243, 245 – 248, 253, 254, 269, 278-286, 289-303, 305-341, 352, 364-377, 379-384, 387-389, 394, 395, 397, 400, 401, 403.

[0988] The following examples exhibited Factor Xa IC_{50} values greater than 100 nM and less than 500 nM:

11 – 15, 33, 66, 68, 73 – 76, 78 – 80, 83 – 85, 87 – 90, 165, 167, 186 – 190, 206, 210, 211, 229, 230, 233, 235, 237, 238, 241, 244, 342-347, 349-351, 353, 354, 359, 378, 385, 386, 390-392, 398.

[0989] The following examples exhibited Factor Xa IC_{50} values greater than or equal to 500 nM:


1, 4, 5, 34 – 47, 52 – 55, 81, 143 – 160, 168 – 181, 183 – 185, 191, 192, 201, 202, 207, 212 – 215, 220 – 224, 228, 231, 232, 234, 236, 249 – 252, 255 – 268, 270 – 276, 288, 304, 348, 355-358, 360-363, 393, 396, 399, 402.

[0990] The present invention provides a number of embodiments. It is apparent that the examples may be altered to provide other embodiments of this invention. Therefore, it will

be appreciated that the scope of this invention is to be defined by the appended claims rather than by the specific embodiments, which have been represented by way of example.

WHAT IS CLAIMED IS:

1. A compound having the formula:



and pharmaceutically acceptable salts thereof, wherein


A represents benzene, a six-membered heterocycle containing 1-3 nitrogen atoms or a five-membered heterocycle containing 1-4 nitrogen, oxygen or sulfur atoms; with 0-4 substituents selected from halogen, oxo, R^A , $-OR^A$, $N(R^A)_2$, $CO_2 R^A$, $CON(R^A)_2$, $S(O)_2 R^A$, $S(O)_2 N(R^A)_2$, $X^1 OR^A$, $X^1 CO_2 R^A$, $X^1 CON(R^A)_2$, $X^1 N(R^A)_2$, wherein each R^A is independently selected from H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, or optionally two R^A attached to the same atom are combined to form a 3-, 4-, 5-, 6- or 7-membered ring;

R is a member selected from the group consisting of H and C_{1-4} alkyl;

R^1 is a member selected from the group consisting of halogen, C_{1-8} alkyl, C_{2-8} alkenyl and C_{2-8} alkynyl;

R^2 is a member selected from the group consisting of H, C_{1-8} alkyl, $-X-OR^{2a}$, $-X-SR^{2a}$, $-X-COR^{2a}$, $-X-CO_2 R^{2a}$ and $-X-N(R^{2a})_2$; wherein X is C_{1-8} alkylene and each R^{2a} is independently selected from the group consisting of H and C_{1-8} alkyl, or optionally two R^{2a} groups attached to the same nitrogen atom are combined to form a 4-, 5-, 6- or 7-membered ring;

W is a member selected from the group consisting of



wherein

each of Z^1 , Z^2 , and Z^3 is independently N, C, or CH and two of Z^1 , Z^2 , and Z^3 are other than N;

the subscript n is an integer of from 0 to 3 and the subscript m is an integer of from 0 to 1;

R^4 is a member selected from the group consisting of halogen, $-OR^{4a}$, $-OC(O)R^{4a}$,

$-NR^{4a}R^{4b}$, $-SR^{4a}$, $S(O)R^{4a}$, $S(O)_2 R^{4a}$, $S(O)_2 NR^{4a}R^{4b}$, $-R^{4c}$, $-CN$, $-NO_2$, $-CO_2 R^{4a}$,

$-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-OC(O)NR^{4a}R^{4b}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)_2 R^{4c}$, $-NR^{4a}-$

$C(O)NR^{4a}R^{4b}$, $-X^1OR^{4a}$, $-X^1OC(O)R^{4a}$, $-X^1NR^{4a}R^{4b}$, $-X^1SR^{4a}$, $-X^1S(O)R^{4a}$, $-X^1S(O)_2R^{4a}$, $-X^1CN$, $-X^1NO_2$, $-X^1CO_2R^{4a}$, $-O-X^1CO_2R^{4a}$, $-X^1CONR^{4a}R^{4b}$, $-O-X^1CONR^{4a}R^{4b}$, $-X^1C(O)R^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-S(O)X^1NR^{4a}R^{4b}$, $-S(O)_2X^1NR^{4a}R^{4b}$, $-X^1OC(O)NR^{4a}R^{4b}$, $-X^1NR^{4a}C(O)R^{4a}$, $-X^1NR^{4b}C(O)_2R^{4c}$, $-X^1NR^{4a}C(O)NR^{4a}R^{4b}$, $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2Y$;

$R^{4'}$ is a member selected from the group consisting of $-C(=NH)R^{4a}$, $-C(=NR^{4c})R^{4a}$, $-C(=NH)NR^{4a}R^{4b}$, $-C(=NR^{4c})NR^{4a}R^{4b}$, $-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$, $-X^2-C(=NH)NR^{4a}R^{4b}$, $-X^2-C(=NR^{4c})NR^{4a}R^{4b}$, $-X^2-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$ and $-C(=NR^{4a})NR^{4a}-Y$,

wherein

Y is a five or six-membered aryl, heterocyclyl or aryl- C_{1-2} alkyl, optionally substituted with from one to three substituents selected from the group consisting of halogen, oxo, $-OR^{4a}$, $-OC(O)R^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2NR^{4a}R^{4b}$, $-CN$, $-NO_2$, $-CO_2R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)_2R^{4c}$, $-X^1R^{4a}$, $-X^1OR^{4a}$, $-X^1SR^{4a}$, $-X^1S(O)R^{4a}$, $-X^1S(O)_2R^{4a}$, $-X^1CN$, $-X^1NO_2$, $-X^1CO_2R^{4a}$, $-X^1CONR^{4a}R^{4b}$, $-X^1C(O)R^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-S(O)X^1NR^{4a}R^{4b}$, $-S(O)_2X^1NR^{4a}R^{4b}$, $-X^1OC(O)NR^{4a}R^{4b}$, $-X^1NR^{4b}C(O)R^{4a}$, $-X^1NR^{4b}C(O)_2R^{4c}$, $-X^1NR^{4a}-C(O)NR^{4a}R^{4b}$, $-X^1OC(O)R^{4a}$, $-X^1NR^{4a}R^{4b}$, $-O-X^1OR^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-O-X^1CO_2R^{4a}$, $-O-X^1CONR^{4a}R^{4b}$, $-NR^{4b}-X^1OR^{4a}$, $-NR^{4b}-X^1NR^{4a}R^{4b}$, $-NR^{4b}-X^1CO_2R^{4a}$, and $-NR^{4b}-X^1CONR^{4a}R^{4b}$,

each X^1 and X^2 are members independently selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene;

each R^{4a} and R^{4b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,

each R^{4c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,

optionally any two of R^{4a} , R^{4b} and R^{4c} when part of a common R^4 or $R^{4'}$ substituent can be combined with the atoms to which each is attached to form a saturated or unsaturated, four to nine-membered mono- or bicyclic ring having from 0 to 2 additional heteroatoms as ring members;


each of R^{4a} , R^{4b} and R^{4c} is optionally further substituted with from one to three members selected from the group consisting of $-OH$, oxo, $-R^m$, $-OR^m$, $-OC(O)NHR^m$, $-OC(O)N(R^m)_2$, $-SH$, $-SR^m$, $-S(O)R^m$, $-S(O)_2R^m$, $-SO_2NH_2$,

62 $-S(O)_2NHR^m$, $-S(O)_2N(R^m)_2$, $-NHS(O)_2R^m$, $-NR^mS(O)_2R^m$, $-C(O)NH_2$,
 63 $-C(O)NHR^m$, $-C(O)N(R^m)_2$, $-C(O)R^m$, $-NHC(O)R^m$, $-NR^mC(O)R^m$, $-NHC(O)NH_2$,
 64 $-NR^mC(O)NH_2$, $-NR^mC(O)NHR^m$, $-NHC(O)NHR^m$, $-NR^mC(O)N(R^m)_2$,
 65 $-NHC(O)N(R^m)_2$, $-CO_2H$, $-CO_2R^m$, $-NHCO_2R^m$, $-NR^mCO_2R^m$, $-CN$, $-NO_2$, $-NH_2$,
 66 $-NHR^m$, $-N(R^m)_2$, $-NR^mS(O)NH_2$ and $-NR^mS(O)_2NHR^m$, wherein each R^m is
 67 independently an unsubstituted C_{1-6} alkyl, benzyl or combined with the atoms to
 68 which each is attached to form a saturated four-, five-, six- or seven-membered
 69 ring having from 0 to 2 additional heteroatoms as ring members;
 70 R^5 is a member selected from the group consisting of $-R^{5a}$, $-CO_2R^{5c}$, $-CONR^{5a}R^{5b}$,
 71 $-C(O)R^{5a}$, $-C(=NR^{5a})NR^{5a}R^{5b}$, $-C(=N^+R^{5c}R^{5c})NR^{5a}R^{5b}$, $-C(=NR^{5a})R^{5a}$, $-C(=NR^{5a})Y^1$,
 72 $-X^3OR^{5a}$, $-X^3OC(O)R^{5a}$, $-X^3NR^{5a}R^{5b}$, $-X^3SR^{5a}$, $-X^3CN$, $-X^3NO_2$, $-X^3CO_2R^{5a}$,
 73 $-X^3CONR^{5a}R^{5b}$, $-X^3C(O)R^{5a}$, $-X^3OC(O)NR^{5a}R^{5b}$, $-X^3NR^{5a}C(O)R^{5a}$, $-X^3NR^{5b}C(O)_2R^{5c}$,
 74 $-X^3NR^{5a}C(O)NR^{5a}R^{5b}$, $-X^3-C(=NR^{5a})NR^{5a}R^{5b}$, $-Y^1$, and $-X^3-Y^1$,
 75 wherein
 76 Y^1 is a five or six-membered aryl or heteroaryl ring or aryl- C_{1-2} alkyl, optionally
 77 substituted with from one to three substituents selected from the group consisting
 78 of halogen, $-OR^{5a}$, $-OC(O)R^{5a}$, $-NR^{5a}R^{5b}$, $-R^{5c}$, $-SR^{5a}$, $-CN$, $-NO_2$, $-CO_2R^{5a}$,
 79 $-CONR^{5a}R^{5b}$, $-C(O)R^{5a}$, $-NR^{5b}C(O)R^{5a}$, $-NR^{5b}C(O)_2R^{5c}$, $-X^3OR^{5a}$, $-X^3SR^{5a}$,
 80 $-X^3CN$, $-X^3NO_2$, $-X^3CO_2R^{5a}$, $-X^3CONR^{5a}R^{5b}$, $-X^3C(O)R^{5a}$, $-X^3OC(O)NR^{5a}R^{5b}$,
 81 $-X^3NR^{5b}C(O)R^{5a}$, $-X^3NR^{5b}C(O)_2R^{5c}$, $-X^3NR^{5a}-C(O)NR^{5a}R^{5b}$, $-X^3OC(O)R^{5a}$,
 82 $-X^3NR^{5a}R^{5b}$, $-O-X^3OR^{5a}$, $-O-X^3NR^{5a}R^{5b}$, $-O-X^3CO_2R^{5a}$, $-O-X^3CONR^{5a}R^{5b}$,
 83 $-NR^{5b}-X^3OR^{5a}$, $-NR^{5b}-X^3NR^{5a}R^{5b}$, $-NR^{5b}-X^3CO_2R^{5a}$, and $-NR^{5b}-X^3CONR^{5a}R^{5b}$,
 84 each X^3 is a member independently selected from the group consisting of C_{1-4} alkylene,
 85 C_{2-4} alkenylene and C_{2-4} alkynylene;
 86 each R^{5a} and R^{5b} are independently selected from the group consisting of hydrogen,
 87 C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8}
 88 alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,
 89 each R^{5c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8}
 90 haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl,
 91 heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,
 92 optionally any two of R^{5a} , R^{5b} and R^{5c} when part of a common R^5 substituent can be
 93 combined with the atoms to which each is attached to form a saturated four-, five-,
 94 six- or seven-membered ring having from 0 to 1 additional heteroatoms as ring
 95 members;

each of R^{5a} , R^{5b} and R^{5c} is optionally further substituted with from one to three members selected from the group consisting of -OH, -ORⁿ, -OC(O)NHRⁿ, -OC(O)N(Rⁿ)₂, -SH, -SRⁿ, -S(O)Rⁿ, -S(O)₂Rⁿ, -SO₂NH₂, -S(O)₂NHRⁿ, -S(O)₂N(Rⁿ)₂, -NHS(O)₂Rⁿ, -NRⁿS(O)₂Rⁿ, -C(O)NH₂, -C(O)NHRⁿ, -C(O)N(Rⁿ)₂, -C(O)Rⁿ, -NHC(O)Rⁿ, -NRⁿC(O)Rⁿ, -NHC(O)NH₂, -NRⁿC(O)NH₂, -NRⁿC(O)NHRⁿ, -NHC(O)NHRⁿ, -NRⁿC(O)N(Rⁿ)₂, -NHC(O)N(Rⁿ)₂, -CO₂H, -CO₂Rⁿ, -NHCO₂Rⁿ, -NRⁿCO₂Rⁿ, -CN, -NO₂, -NH₂, -NHRⁿ, -N(Rⁿ)₂, -NRⁿS(O)NH₂ and -NRⁿS(O)₂NHRⁿ, wherein each Rⁿ is independently an unsubstituted C₁₋₆ alkyl or benzyl; and

the wavy line indicates the point of attachment to the rest of the molecule.


2. The compound of claim 1, wherein W has a formula selected from the group consisting of (a), (d), (e) and (c):



3. The compound of claims 1 or 2, wherein W has a formula selected from the group consisting of (a), (d) and (e).


4. The compound of any of the preceding claims, wherein W has a formula (c).

5. The compound of any of the preceding claims, wherein at least one of n and m is 1 and each R⁴ and R^{4'} is selected from the group consisting of:




3 ; and the wavy line indicates the point of attachment to the rest of the molecule.

1 6. The compound of any one of claims 1 to 4, wherein at least one of n
 2 and m is 1 and each R⁴ and R^{4'} is independently selected from the group consisting of H,
 3 halogen, OR^{4a}, C₁₋₄ alkyl, SR^{4a}, S(O)R^{4a}, S(O)₂R^{4a}, S(O)₂N(R^{4a}R^{4b}), NR^{4a}R^{4b}, C₁₋₄alkylOR^{4a},
 4 C₁₋₄alkylNR^{4a}R^{4b}, C₁₋₄alkylCO₂R^{4a}, OC₁₋₄alkylOR^{4a}, OC₁₋₄alkylN(R^{4a}N^{4b}), N(R^{4a})C₁₋₄
 5 alkylOR^{4b}, N(R^{4a})C₁₋₄alkylN(R^{4a}R^{4b}), S(O)₂C₁₋₄alkylN(R^{4a}R^{4b}),


$$\text{--}\zeta\text{--CO}_2\text{H}, \text{--}\zeta\text{--SO}_2\text{NH}_2, \text{--}\zeta\text{--SO}_2\text{CH}_3 \text{ and } \text{--}\zeta\text{--SOCH}_3$$

; each R^{4a} is independently H or C₁₋₄alkyl; each R^{4d} is independently selected from the group consisting of H, OH, NH₂, N(CH₃)₂, CONH₂ and CON(CH₃)₂; and the wavy line indicates the point of attachment to the rest of the molecule.

- 1 7. The compound of any one of claims 1 to 4, wherein at least one of n
2 and m is 1 and each R⁴ and R^{4'} is selected from the group consisting of:



4 ; and the wavy line indicates the point of attachment to the rest of the molecule.

1 8. The compound of any of the preceding claims, wherein R^4 is attached
2 to the ring at a position para to A.


1 9. The compound of any of the preceding claims, wherein R^4 is attached
2 to the ring at a position ortho to A.

1 10. The compound of any of the preceding claims, wherein Y is an
2 optionally substituted member selected from the group consisting of phenyl, benzyl, pyridyl,
3 pyridylmethyl, pyrimidinyl, pyrazolyl, triazolyl and imidazolyl.

1 11. The compound of any of the preceding claims, wherein A is benzene
2 or a six-membered heterocycle containing 1-3 nitrogen atoms.

1 12. The compound of any of the preceding claims, wherein A is member
2 selected from the group consisting of: benzene, pyridine, pyridimidine, pyrazine, morpholin-
3 3-one, piperazine-2-one, pyridine-2-one, piperidine, morpholine and piperazine.

1 13. The compound of any of the preceding claims, wherein A is selected
2 from the group consisting of:




3
4 ;

5 n is an integer from 0 to 3; each R^3 is independently selected from the group
6 consisting of halogen, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, OR^{3a} , $N(R^{3a})_2$, $X^0CO_2R^{3a}$,
7 $X^0CON(R^{3a})_2$, SO_2C_{1-4} , $SO_2N(R^{3a})_2$; each R^{3a} is independently selected from the group
8 consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl and C_{2-8} alkynyl; and X^0 is a bond or C_{1-8} alkylene; and

9 the dashed line indicates the point of attachment to A and the wavy line indicates the point of
10 attachment to the rest of the molecule.

1 14. The compound of any of the preceding claims having the formula:



2
3
4 and pharmaceutically acceptable salts thereof, wherein


5 R is a member selected from the group consisting of H and C₁₋₄ alkyl

6 R¹ is a member selected from the group consisting of halogen, C₁₋₈alkyl, C₂₋₈alkenyl and
7 C₂₋₈alkynyl;

8 R² is a member selected from the group consisting of H, C₁₋₈ alkyl, -X-OR^{2a}, -X-SR^{2a},
9 -X-COR^{2a}, -X-CO₂R^{2a} and -X-N(R^{2a})₂; wherein X is C₁₋₈alkylene and each R^{2a} is
10 independently selected from the group consisting of H and C₁₋₈alkyl, or optionally
11 two R^{2a} groups attached to the same nitrogen atom are combined to form a 4-, 5-, 6-
12 or 7-membered ring;

13 R³ is a member selected from the group consisting of H and halogen;

14 W is a member selected from the group consisting of



15
16
17 wherein

18 each of Z¹, Z², Z³ and Z⁴ is independently N, C, or CH and two of Z¹, Z², and Z³ are other
19 than N;

20 the subscript n is an integer of from 0 to 3 and the subscript m is an integer of from 0 to 1;

21 R⁴ is a member selected from the group consisting of halogen, -OR^{4a}, -OC(O)R^{4a},
22 -NR^{4a}R^{4b}, -SR^{4a}, S(O)R^{4a}, S(O)₂R^{4a}, S(O)₂NR^{4a}R^{4b}, -R^{4c}, -CN, -NO₂, -CO₂R^{4a},
23 -CONR^{4a}R^{4b}, -C(O)R^{4a}, -OC(O)NR^{4a}R^{4b}, -NR^{4b}C(O)R^{4a}, -NR^{4b}C(O)₂R^{4c}, -NR^{4a}-
24 C(O)NR^{4a}R^{4b}, -X¹OR^{4a}, -X¹OC(O)R^{4a}, -X¹NR^{4a}R^{4b}, -X¹SR^{4a}, -X¹S(O)R^{4a}, -
25 X¹S(O)₂R^{4a}, -X¹CN, -X¹NO₂, -X¹CO₂R^{4a}, -O-X¹CO₂R^{4a}, -X¹CONR^{4a}R^{4b},
26 -O-X¹CONR^{4a}R^{4b}, -X¹C(O)R^{4a}, -O-X¹NR^{4a}R^{4b}, -S(O)X¹NR^{4a}R^{4b}, -S(O)₂X¹NR^{4a}R^{4b},

$-X^1OC(O)NR^{4a}R^{4b}$, $-X^1NR^{4a}C(O)R^{4a}$, $-X^1NR^{4b}C(O)_2R^{4c}$, $-X^1NR^{4a}C(O)NR^{4a}R^{4b}$, $-Y$,
 $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2Y$;

$R^{4'}$ is a member selected from the group consisting of $-C(=NH)R^{4a}$, $-C(=NR^{4c})R^{4a}$,
 $-C(=NH)NR^{4a}R^{4b}$, $-C(=NR^{4c})NR^{4a}R^{4b}$, $-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$, $-X^2-C(=NH)NR^{4a}R^{4b}$,
 $-X^2-C(=NR^{4c})NR^{4a}R^{4b}$, $-X^2-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$ and $-C(=NR^{4a})NR^{4a}-Y$,

wherein

Y is a five or six-membered aryl, heterocyclyl or aryl- C_{1-2} alkyl, optionally substituted
 with from one to three substituents selected from the group consisting of halogen,
 oxo, $-OR^{4a}$, $-OC(O)R^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2NR^{4a}R^{4b}$,
 $-CN$, $-NO_2$, $-CO_2R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)_2R^{4c}$,
 $-X^1R^{4a}$, $-X^1OR^{4a}$, $-X^1SR^{4a}$, $-X^1S(O)R^{4a}$, $-X^1S(O)_2R^{4a}$, $-X^1CN$, $-X^1NO_2$, $-X^1CO_2R^{4a}$,
 $-X^1CONR^{4a}R^{4b}$, $-X^1C(O)R^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-S(O)X^1NR^{4a}R^{4b}$, $-S(O)_2X^1NR^{4a}R^{4b}$,
 $-X^1OC(O)NR^{4a}R^{4b}$, $-X^1NR^{4b}C(O)R^{4a}$, $-X^1NR^{4b}C(O)_2R^{4c}$, $-X^1NR^{4a}-C(O)NR^{4a}R^{4b}$,
 $-X^1OC(O)R^{4a}$, $-X^1NR^{4a}R^{4b}$, $-O-X^1OR^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-O-X^1CO_2R^{4a}$,
 $-O-X^1CONR^{4a}R^{4b}$, $-NR^{4b}-X^1OR^{4a}$, $-NR^{4b}-X^1NR^{4a}R^{4b}$, $-NR^{4b}-X^1CO_2R^{4a}$, and
 $-NR^{4b}-X^1CONR^{4a}R^{4b}$,

each X^1 and X^2 are members independently selected from the group consisting of C_{1-4}
 alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene;

each R^{4a} and R^{4b} are independently selected from the group consisting of hydrogen,
 C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8}
 alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,

each R^{4c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8}
 haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl,
 heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,

optionally any two of R^{4a} , R^{4b} and R^{4c} when part of a common R^4 or $R^{4'}$ substituent can
 be combined with the atoms to which each is attached to form a saturated or
 unsaturated, four to nine-membered mono- or bicyclic ring having from 0 to 2
 additional heteroatoms as ring members;

each of R^{4a} , R^{4b} and R^{4c} is optionally further substituted with from one to three
 members selected from the group consisting of $-OH$, oxo, $-R^m$, $-OR^m$,
 $-OC(O)NHR^m$, $-OC(O)N(R^m)_2$, $-SH$, $-SR^m$, $-S(O)R^m$, $-S(O)_2R^m$, $-SO_2NH_2$,
 $-S(O)_2NHR^m$, $-S(O)_2N(R^m)_2$, $-NHS(O)_2R^m$, $-NR^mS(O)_2R^m$, $-C(O)NH_2$,
 $-C(O)NHR^m$, $-C(O)N(R^m)_2$, $-C(O)R^m$, $-NHC(O)R^m$, $-NR^mC(O)R^m$, $-NHC(O)NH_2$,
 $-NR^mC(O)NH_2$, $-NR^mC(O)NHR^m$, $-NHC(O)NHR^m$, $-NR^mC(O)N(R^m)_2$,


61 $-\text{NHC(O)N(R}^m)_2$, $-\text{CO}_2\text{H}$, $-\text{CO}_2\text{R}^m$, $-\text{NHCO}_2\text{R}^m$, $-\text{NR}^m\text{CO}_2\text{R}^m$, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$,
 62 $-\text{NHR}^m$, $-\text{N(R}^m)_2$, $-\text{NR}^m\text{S(O)NH}_2$ and $-\text{NR}^m\text{S(O)}_2\text{NHR}^m$, wherein each R^m is
 63 independently an unsubstituted C_{1-6} alkyl, benzyl or combined with the atoms to
 64 which each is attached to form a saturated four-, five-, six- or seven-membered
 65 ring having from 0 to 2 additional heteroatoms as ring members;
 66 R^5 is a member selected from the group consisting of $-\text{R}^{5a}$, $-\text{CO}_2\text{R}^{5c}$, $-\text{CONR}^{5a}\text{R}^{5b}$,
 67 $-\text{C(O)R}^{5a}$, $-\text{C(=NH)NR}^{5a}\text{R}^{5b}$, $-\text{C(=NR}^{5c})\text{NR}^{5a}\text{R}^{5b}$, $-\text{C(=N}^+\text{R}^{5c}\text{R}^{5c})\text{NR}^{5a}\text{R}^{5b}$,
 68 $-\text{C(=NR}^{5a})\text{R}^{5a}$, $-\text{C(=NR}^{5a})\text{Y}^1$, $-\text{X}^3\text{OR}^{5a}$, $-\text{X}^3\text{OC(O)R}^{5a}$, $-\text{X}^3\text{NR}^{5a}\text{R}^{5b}$, $-\text{X}^3\text{SR}^{5a}$, $-\text{X}^3\text{CN}$, $-\text{X}^3\text{NO}_2$,
 69 $-\text{X}^3\text{CO}_2\text{R}^{5a}$, $-\text{X}^3\text{CONR}^{5a}\text{R}^{5b}$, $-\text{X}^3\text{C(O)R}^{5a}$, $-\text{X}^3\text{OC(O)NR}^{5a}\text{R}^{5b}$, $-\text{X}^3\text{NR}^{5a}\text{C(O)R}^{5a}$,
 70 $-\text{X}^3\text{NR}^{5b}\text{C(O)}_2\text{R}^{5c}$, $-\text{X}^3\text{NR}^{5a}\text{C(O)NR}^{5a}\text{R}^{5b}$, $-\text{X}^3\text{-C(=NR}^{5a})\text{NR}^{5a}\text{R}^{5b}$,
 71 $-\text{Y}^1$, and $-\text{X}^3\text{-Y}^1$,
 72 wherein
 73 Y^1 is a five or six-membered aryl or heteroaryl ring or aryl- C_{1-2} alkyl, optionally
 74 substituted with from one to three substituents selected from the group consisting
 75 of halogen, $-\text{OR}^{5a}$, $-\text{OC(O)R}^{5a}$, $-\text{NR}^{5a}\text{R}^{5b}$, $-\text{R}^{5c}$, $-\text{SR}^{5a}$, $-\text{CN}$, $-\text{NO}_2$, $-\text{CO}_2\text{R}^{5a}$,
 76 $-\text{CONR}^{5a}\text{R}^{5b}$, $-\text{C(O)R}^{5a}$, $-\text{NR}^{5b}\text{C(O)R}^{5a}$, $-\text{NR}^{5b}\text{C(O)}_2\text{R}^{5c}$, $-\text{X}^3\text{OR}^{5a}$, $-\text{X}^3\text{SR}^{5a}$,
 77 $-\text{X}^3\text{CN}$, $-\text{X}^3\text{NO}_2$, $-\text{X}^3\text{CO}_2\text{R}^{5a}$, $-\text{X}^3\text{CONR}^{5a}\text{R}^{5b}$, $-\text{X}^3\text{C(O)R}^{5a}$, $-\text{X}^3\text{OC(O)NR}^{5a}\text{R}^{5b}$,
 78 $-\text{X}^3\text{NR}^{5b}\text{C(O)R}^{5a}$, $-\text{X}^3\text{NR}^{5b}\text{C(O)}_2\text{R}^{5c}$, $-\text{X}^3\text{NR}^{5a}\text{-C(O)NR}^{5a}\text{R}^{5b}$, $-\text{X}^3\text{OC(O)R}^{5a}$,
 79 $-\text{X}^3\text{NR}^{5a}\text{R}^{5b}$, $-\text{O-X}^3\text{OR}^{5a}$, $-\text{O-X}^3\text{NR}^{5a}\text{R}^{5b}$, $-\text{O-X}^3\text{CO}_2\text{R}^{5a}$, $-\text{O-X}^3\text{CONR}^{5a}\text{R}^{5b}$,
 80 $-\text{NR}^{5b}\text{-X}^3\text{OR}^{5a}$, $-\text{NR}^{5b}\text{-X}^3\text{NR}^{5a}\text{R}^{5b}$, $-\text{NR}^{5b}\text{-X}^3\text{CO}_2\text{R}^{5a}$, and $-\text{NR}^{5b}\text{-X}^3\text{CONR}^{5a}\text{R}^{5b}$,
 81 each X^3 is a member independently selected from the group consisting of C_{1-4} alkylene,
 82 C_{2-4} alkenylene and C_{2-4} alkynylene;
 83 each R^{5a} and R^{5b} are independently selected from the group consisting of hydrogen,
 84 C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8}
 85 alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,
 86 each R^{5c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8}
 87 haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl,
 88 heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,
 89 optionally any two of R^{5a} , R^{5b} and R^{5c} when part of a common R^5 substituent can be
 90 combined with the atoms to which each is attached to form a saturated four-, five-,
 91 six- or seven-membered ring having from 0 to 1 additional heteroatoms as ring
 92 members;
 93 each of R^{5a} , R^{5b} and R^{5c} is optionally further substituted with from one to three
 94 members selected from the group consisting of $-\text{OH}$, $-\text{OR}^n$, $-\text{OC(O)NHR}^n$,

95 -OC(O)N(Rⁿ)₂, -SH, -SRⁿ, -S(O)Rⁿ, -S(O)₂Rⁿ, -SO₂NH₂, -S(O)₂NHRⁿ,
 96 -S(O)₂N(Rⁿ)₂, -NHS(O)₂Rⁿ, -NRⁿS(O)₂Rⁿ, -C(O)NH₂, -C(O)NHRⁿ, -C(O)N(Rⁿ)₂,
 97 -C(O)Rⁿ, -NHC(O)Rⁿ, -NRⁿC(O)Rⁿ, -NHC(O)NH₂, -NRⁿC(O)NH₂,
 98 -NRⁿC(O)NHRⁿ, -NHC(O)NHRⁿ, -NRⁿC(O)N(Rⁿ)₂, -NHC(O)N(Rⁿ)₂, -CO₂H,
 99 -CO₂Rⁿ, -NHCO₂Rⁿ, -NRⁿCO₂Rⁿ, -CN, -NO₂, -NH₂, -NHRⁿ, -N(Rⁿ)₂,
 100 -NRⁿS(O)NH₂ and -NRⁿS(O)₂NHRⁿ, wherein each Rⁿ is independently an
 101 unsubstituted C₁₋₆ alkyl or benzyl.

1 15. The compound of any one of claims 1 to 3 and 5 to 14, wherein W has
 2 formula (a).


1 16. The compound of any of the preceding claims, wherein m is 1.

1 17. The compound of any of the preceding claims, wherein R^{4'} is selected
 2 from the group consisting of:



1 18. The compound of any one of claims 1 to 8 and 10 to 17, wherein R^{4'} is
 2 attached to the phenyl ring at a position *para* to the ring A.

19. The compound of any one of claims 1 to 10 having the formula:



and pharmaceutically acceptable salts thereof, wherein

A represents a five-membered heterocycle containing 1-4 nitrogen, oxygen or sulfur atoms; with 0-4 substituents selected from halogen, oxo, R^A , $-OR^A$, $N(R^A)_2$, CO_2R^A , $CON(R^A)_2$, $S(O)_2R^A$, $S(O)_2N(R^A)_2$, X^1OR^A , $X^1CO_2R^A$, $X^1CON(R^A)_2$, $X^1N(R^A)_2$, wherein each R^A is independently selected from H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, or optionally two R^A attached to the same atom are combined to form a 3-, 4-, 5-, 6- or 7-membered ring;


R is a member selected from the group consisting of H and C_{1-4} alkyl

R^1 is a member selected from the group consisting of halogen, C_{1-8} alkyl, C_{2-8} alkenyl and C_{2-8} alkynyl;

R^2 is a member selected from the group consisting of H, C_{1-8} alkyl, $-X-OR^{2a}$, $-X-SR^{2a}$, $-X-COR^{2a}$, $-X-CO_2R^{2a}$ and $-X-N(R^{2a})_2$; wherein X is C_{1-8} alkylene and each R^{2a} is independently selected from the group consisting of H and C_{1-8} alkyl, or optionally two R^{2a} groups attached to the same nitrogen atom are combined to form a 4-, 5-, 6- or 7-membered ring;

R^3 is a member selected from the group consisting of H and C_{1-4} alkyl; or is absent when a carbon-carbon double bond is present in the ring to which R^3 would otherwise be attached;

W is a member selected from the group consisting of



wherein

each of Z^1 , Z^2 , and Z^3 is independently N, C, or CH and two of Z^1 , Z^2 , and Z^3 are other than N;

the subscript n is an integer of from 0 to 3 and the subscript m is an integer of from 0 to 1;

R^4 is a member selected from the group consisting of halogen, $-OR^{4a}$, $-OC(O)R^{4a}$, $-NR^{4a}R^{4b}$, $-SR^{4a}$, $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2NR^{4a}R^{4b}$, $-R^{4c}$, $-CN$, $-NO_2$, $-CO_2R^{4a}$,

30 $-\text{CONR}^{4a}\text{R}^{4b}$, $-\text{C}(\text{O})\text{R}^{4a}$, $-\text{OC}(\text{O})\text{NR}^{4a}\text{R}^{4b}$, $-\text{NR}^{4b}\text{C}(\text{O})\text{R}^{4a}$, $-\text{NR}^{4b}\text{C}(\text{O})_2\text{R}^{4c}$, $-\text{NR}^{4a}-$
 31 $\text{C}(\text{O})\text{NR}^{4a}\text{R}^{4b}$, $-\text{X}^1\text{OR}^{4a}$, $-\text{X}^1\text{OC}(\text{O})\text{R}^{4a}$, $-\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{X}^1\text{SR}^{4a}$, $-\text{X}^1\text{S}(\text{O})\text{R}^{4a}$, $-$
 32 $\text{X}^1\text{S}(\text{O})_2\text{R}^{4a}$, $-\text{X}^1\text{CN}$, $-\text{X}^1\text{NO}_2$, $-\text{X}^1\text{CO}_2\text{R}^{4a}$, $-\text{O}-\text{X}^1\text{CO}_2\text{R}^{4a}$, $-\text{X}^1\text{CONR}^{4a}\text{R}^{4b}$,
 33 $-\text{O}-\text{X}^1\text{CONR}^{4a}\text{R}^{4b}$, $-\text{X}^1\text{C}(\text{O})\text{R}^{4a}$, $-\text{O}-\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{S}(\text{O})\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{S}(\text{O})_2\text{X}^1\text{NR}^{4a}\text{R}^{4b}$,
 34 $-\text{X}^1\text{OC}(\text{O})\text{NR}^{4a}\text{R}^{4b}$, $-\text{X}^1\text{NR}^{4a}\text{C}(\text{O})\text{R}^{4a}$, $-\text{X}^1\text{NR}^{4b}\text{C}(\text{O})_2\text{R}^{4c}$, $-\text{X}^1\text{NR}^{4a}\text{C}(\text{O})\text{NR}^{4a}\text{R}^{4b}$, $-\text{Y}$,
 35 $-\text{X}^1-\text{Y}$, $-\text{O}-\text{Y}$, $-\text{NR}^{4a}\text{Y}$, $-\text{SY}$, $-\text{S}(\text{O})\text{Y}$ and $-\text{S}(\text{O})_2\text{Y}$;
 36 $\text{R}^{4'}$ is a member selected from the group consisting of $-\text{C}(=\text{NH})\text{R}^{4a}$, $-\text{C}(=\text{NR}^{4c})\text{R}^{4a}$,
 37 $-\text{C}(=\text{NH})\text{NR}^{4a}\text{R}^{4b}$, $-\text{C}(=\text{NR}^{4c})\text{NR}^{4a}\text{R}^{4b}$, $-\text{C}(=\text{N}^+\text{R}^{4c}\text{R}^{4c})\text{NR}^{4a}\text{R}^{4b}$, $-\text{X}^2-\text{C}(=\text{NH})\text{NR}^{4a}\text{R}^{4b}$,
 38 $-\text{X}^2-\text{C}(=\text{NR}^{4c})\text{NR}^{4a}\text{R}^{4b}$, $-\text{X}^2-\text{C}(=\text{N}^+\text{R}^{4c}\text{R}^{4c})\text{NR}^{4a}\text{R}^{4b}$ and $-\text{C}(=\text{NR}^{4a})\text{NR}^{4a}-\text{Y}$,
 39 wherein
 40 Y is a five or six-membered aryl, heterocyclyl or aryl- C_{1-2} alkyl, optionally substituted
 41 with from one to three substituents selected from the group consisting of halogen,
 42 oxo, $-\text{OR}^{4a}$, $-\text{OC}(\text{O})\text{R}^{4a}$, $-\text{NR}^{4a}\text{R}^{4b}$, $-\text{R}^{4c}$, $-\text{SR}^{4a}$, $\text{S}(\text{O})\text{R}^{4a}$, $\text{S}(\text{O})_2\text{R}^{4a}$, $\text{S}(\text{O})_2\text{NR}^{4a}\text{R}^{4b}$,
 43 $-\text{CN}$, $-\text{NO}_2$, $-\text{CO}_2\text{R}^{4a}$, $-\text{CONR}^{4a}\text{R}^{4b}$, $-\text{C}(\text{O})\text{R}^{4a}$, $-\text{NR}^{4b}\text{C}(\text{O})\text{R}^{4a}$, $-\text{NR}^{4b}\text{C}(\text{O})_2\text{R}^{4c}$,
 44 $-\text{X}^1\text{R}^{4a}$, $-\text{X}^1\text{OR}^{4a}$, $-\text{X}^1\text{SR}^{4a}$, $-\text{X}^1\text{S}(\text{O})\text{R}^{4a}$, $-\text{X}^1\text{S}(\text{O})_2\text{R}^{4a}$, $-\text{X}^1\text{CN}$, $-\text{X}^1\text{NO}_2$, $-\text{X}^1\text{CO}_2\text{R}^{4a}$,
 45 $-\text{X}^1\text{CONR}^{4a}\text{R}^{4b}$, $-\text{X}^1\text{C}(\text{O})\text{R}^{4a}$, $-\text{O}-\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{S}(\text{O})\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{S}(\text{O})_2\text{X}^1\text{NR}^{4a}\text{R}^{4b}$,
 46 $-\text{X}^1\text{OC}(\text{O})\text{NR}^{4a}\text{R}^{4b}$, $-\text{X}^1\text{NR}^{4b}\text{C}(\text{O})\text{R}^{4a}$, $-\text{X}^1\text{NR}^{4b}\text{C}(\text{O})_2\text{R}^{4c}$, $-\text{X}^1\text{NR}^{4a}-\text{C}(\text{O})\text{NR}^{4a}\text{R}^{4b}$,
 47 $-\text{X}^1\text{OC}(\text{O})\text{R}^{4a}$, $-\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{O}-\text{X}^1\text{OR}^{4a}$, $-\text{O}-\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{O}-\text{X}^1\text{CO}_2\text{R}^{4a}$,
 48 $-\text{O}-\text{X}^1\text{CONR}^{4a}\text{R}^{4b}$, $-\text{NR}^{4b}-\text{X}^1\text{OR}^{4a}$, $-\text{NR}^{4b}-\text{X}^1\text{NR}^{4a}\text{R}^{4b}$, $-\text{NR}^{4b}-\text{X}^1\text{CO}_2\text{R}^{4a}$, and
 49 $-\text{NR}^{4b}-\text{X}^1\text{CONR}^{4a}\text{R}^{4b}$,
 50 each X^1 and X^2 are members independently selected from the group consisting of C_{1-4}
 51 alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene;
 52 each R^{4a} and R^{4b} are independently selected from the group consisting of hydrogen,
 53 C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8}
 54 alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,
 55 each R^{4c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8}
 56 haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl,
 57 heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,
 58 optionally any two of R^{4a} , R^{4b} and R^{4c} when part of a common R^4 or $\text{R}^{4'}$ substituent can
 59 be combined with the atoms to which each is attached to form a saturated or
 60 unsaturated, four to nine-membered mono- or bicyclic ring having from 0 to 2
 61 additional heteroatoms as ring members;
 62 each of R^{4a} , R^{4b} and R^{4c} is optionally further substituted with from one to three
 63 members selected from the group consisting of $-\text{OH}$, oxo, $-\text{R}^m$, $-\text{OR}^m$,

64 -OC(O)NHR^m, -OC(O)N(R^m)₂, -SH, -SR^m, -S(O)R^m, -S(O)₂R^m, -SO₂NH₂,
 65 -S(O)₂NHR^m, -S(O)₂N(R^m)₂, -NHS(O)₂R^m, -NR^mS(O)₂R^m, -C(O)NH₂,
 66 -C(O)NHR^m, -C(O)N(R^m)₂, -C(O)R^m, -NHC(O)R^m, -NR^mC(O)R^m, -NHC(O)NH₂,
 67 -NR^mC(O)NH₂, -NR^mC(O)NHR^m, -NHC(O)NHR^m, -NR^mC(O)N(R^m)₂,
 68 -NHC(O)N(R^m)₂, -CO₂H, -CO₂R^m, -NHCO₂R^m, -NR^mCO₂R^m, -CN, -NO₂, -NH₂,
 69 -NHR^m, -N(R^m)₂, -NR^mS(O)NH₂ and -NR^mS(O)₂NHR^m, wherein each R^m is
 70 independently an unsubstituted C₁₋₆alkyl, benzyl or combined with the atoms to
 71 which each is attached to form a saturated four-, five-, six- or seven-membered
 72 ring having from 0 to 2 additional heteroatoms as ring members;

73 R⁵ is a member selected from the group consisting of -R^{5a}, -CO₂R^{5c}, -CONR^{5a}R^{5b},
 74 -C(O)R^{5a}, -C(=NH)NR^{5a}R^{5b}, -C(=NR^{5c})NR^{5a}R^{5b}, -C(=N⁺R^{5c}R^{5c})NR^{5a}R^{5b},
 75 -C(=NR^{5a})R^{5a}, -C(=NR^{5a})Y¹, -X³OR^{5a}, -X³OC(O)R^{5a}, -X³NR^{5a}R^{5b}, -X³SR^{5a}, -X³CN, -
 76 X³NO₂, -X³CO₂R^{5a}, -X³CONR^{5a}R^{5b}, -X³C(O)R^{5a}, -X³OC(O)NR^{5a}R^{5b}, -
 77 X³NR^{5a}C(O)R^{5a}, -X³NR^{5b}C(O)₂R^{5c}, -X³NR^{5a}C(O)NR^{5a}R^{5b}, -X³-C(=NR^{5a})NR^{5a}R^{5b},
 78 -Y¹, and -X³-Y¹,

79 wherein

80 Y¹ is a five or six-membered aryl or heteroaryl ring or aryl-C₁₋₂ alkyl, optionally
 81 substituted with from one to three substituents selected from the group consisting
 82 of halogen, -OR^{5a}, -OC(O)R^{5a}, -NR^{5a}R^{5b}, -R^{5c}, -SR^{5a}, -CN, -NO₂, -CO₂R^{5a},
 83 -CONR^{5a}R^{5b}, -C(O)R^{5a}, -NR^{5b}C(O)R^{5a}, -NR^{5b}C(O)₂R^{5c}, -X³OR^{5a}, -X³SR^{5a},
 84 -X³CN, -X³NO₂, -X³CO₂R^{5a}, -X³CONR^{5a}R^{5b}, -X³C(O)R^{5a}, -X³OC(O)NR^{5a}R^{5b},
 85 -X³NR^{5b}C(O)R^{5a}, -X³NR^{5b}C(O)₂R^{5c}, -X³NR^{5a}-C(O)NR^{5a}R^{5b}, -X³OC(O)R^{5a},
 86 -X³NR^{5a}R^{5b}, -O-X³OR^{5a}, -O-X³NR^{5a}R^{5b}, -O-X³CO₂R^{5a}, -O-X³CONR^{5a}R^{5b},
 87 -NR^{5b}-X³OR^{5a}, -NR^{5b}-X³NR^{5a}R^{5b}, -NR^{5b}-X³CO₂R^{5a}, and -NR^{5b}-X³CONR^{5a}R^{5b},

88 each X³ is a member independently selected from the group consisting of C₁₋₄ alkylene,
 89 C₂₋₄ alkenylene and C₂₋₄ alkynylene;

90 each R^{5a} and R^{5b} are independently selected from the group consisting of hydrogen,
 91 C₁₋₈ alkyl, C₁₋₈ haloalkyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocycloalkyl, C₂₋₈ alkenyl, C₂₋₈
 92 alkynyl, aryl, heteroaryl, aryl-C₁₋₄ alkyl, and aryloxy-C₁₋₄ alkyl,


93 each R^{5c} is independently selected from the group consisting of C₁₋₈ alkyl, C₁₋₈
 94 haloalkyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocycloalkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl,
 95 heteroaryl, aryl-C₁₋₄ alkyl, and aryloxy-C₁₋₄ alkyl,

96 optionally any two of R^{5a}, R^{5b} and R^{5c} when part of a common R⁵ substituent can be
 97 combined with the atoms to which each is attached to form a saturated four-, five-,

98 six- or seven-membered ring having from 0 to 1 additional heteroatoms as ring
 99 members;
 100 each of R^{5a} , R^{5b} and R^{5c} is optionally further substituted with from one to three
 101 members selected from the group consisting of -OH, -ORⁿ, -OC(O)NHRⁿ,
 102 -OC(O)N(Rⁿ)₂, -SH, -SRⁿ, -S(O)Rⁿ, -S(O)₂Rⁿ, -SO₂NH₂, -S(O)₂NHRⁿ,
 103 -S(O)₂N(Rⁿ)₂, -NHS(O)₂Rⁿ, -NRⁿS(O)₂Rⁿ, -C(O)NH₂, -C(O)NHRⁿ, -C(O)N(Rⁿ)₂,
 104 -C(O)Rⁿ, -NHC(O)Rⁿ, -NRⁿC(O)Rⁿ, -NHC(O)NH₂, -NRⁿC(O)NH₂,
 105 -NRⁿC(O)NHRⁿ, -NHC(O)NHRⁿ, -NRⁿC(O)N(Rⁿ)₂, -NHC(O)N(Rⁿ)₂, -CO₂H,
 106 -CO₂Rⁿ, -NHCO₂Rⁿ, -NRⁿCO₂Rⁿ, -CN, -NO₂, -NH₂, -NHRⁿ, -N(Rⁿ)₂,
 107 -NRⁿS(O)NH₂ and -NRⁿS(O)₂NHRⁿ, wherein each Rⁿ is independently an
 108 unsubstituted C₁₋₆ alkyl or benzyl.

1 **20.** The compound of any one of claims **1** to **10** and **19**, wherein A is
 2 member selected from the group consisting of: isoxazole, isothiazole, pyrazole, imidazole,
 3 oxazole, thiazole, isoxazoline, pyrazoline, imidazoline, 1,2,3-triazole, 1,2,4-triazole, 1,3,4-
 4 oxadiazole, 1,3,4-thiadiazole, 1,2,4-oxadiazole, 1,2,4-thiadiazole, pyrazol-5-one, pyrrolidine-
 5 2,5-dione, imidazolidine-2,4-dione, pyrrolidine, pyrrole, furan and thiophene.

1 **21.** The compound of any one of claims **1** to **10** and **20**, wherein A is
 2 selected from the group consisting of:




3
4 ; each R^3 is independently selected from the group consisting of halogen, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, OR^{3a} , $N(R^{3a})_2$, $X^0CO_2R^{3a}$, $X^0CON(R^{3a})_2$, SO_2C_{1-4} , $SO_2N(R^{3a})_2$; each R^{3a}
5
6 is independently selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl and C_{2-8} alkynyl; and X^0 is a bond or C_{1-8} alkylene; and the dashed line indicates the point of
7
8 attachment to A and the wavy line indicates the point of attachment to the rest of the
9 molecule.

22. The compound of any one of claims 19 to 21, wherein W has formula (a)..

23. The compound of claim 22, wherein m is 1.

24. The compound of any one of claims 22 to 23, wherein R^{4'} is selected from the group consisting of:




25. The compound of any one of claims 22 to 24, wherein R^{4'} is attached to the phenyl ring at a position *para* to A.

26. The compound of any one of claims 22 to, wherein the subscript n is an integer of from 0 to 1.


27. The compound of any one of claims 22 to 26, wherein the subscript n is 1; R⁴ is attached to the phenyl ring at a position *ortho* to A and R^{4'} is attached to the phenyl ring at a position *para* to A.

1 **28.** The compound of any one of claims **22** to **27**, wherein R^4 is selected
2 from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-SR^{4a}$, $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$ and
3 $-SY$.

1 **29.** The compound of any one of claims **22** to **28**, wherein n is 0, 1 or 2,
2 and R^4 is independently selected from the group consisting of H, halogen, OR^{4a} , C_{1-4} alkyl,
3 SR^{4a} , $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2N(R^{4a}R^{4b})$, $NR^{4a}R^{4b}$, $C_{1-4}alkylOR^{4a}$, $C_{1-4}alkylNR^{4a}R^{4b}$, C_{1-}
4 $4alkylCO_2R^{4a}$, $OC_{1-4}alkylOR^{4a}$, $OC_{1-4}alkylN(R^{4a}R^{4b})$, $N(R^{4a})C_{1-4}alkylOR^{4b}$, $N(R^{4a})C_{1-}$
5 $4alkylN(R^{4a}R^{4b})$, $S(O)_2C_{1-4}alkylN(R^{4a}R^{4b})$,



265

11
12

1 **30.** The compound of any one of claims **22** to **29**, wherein m is 0 or 1; n is 0,
2 1 or 2; and at least one R^4 is selected from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$,
3 $-SY$, $-S(O)Y$ and $-S(O)_2Y$, wherein Y is an optionally substituted member selected from the
4 group consisting of phenyl, benzyl, pyridyl, pyridylmethyl, pyrimidinyl and pyrazolyl.

1 **31.** The compound of any one of claims **22** to **30**, wherein R^4 is selected
2 from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$ and $-NR^{4a}Y$, wherein Y is an optionally
3 substituted member selected from the group consisting of phenyl, benzyl, pyridyl and
4 pyridylmethyl.

1 **32.** The compound of claims **22** to **31**, wherein the optional substituents
2 are selected from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $-CO_2R^{4a}$,
3 $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-X^1OR^{4a}$ and $-X^1NR^{4a}R^{4b}$.

1 **33.** The compound of any one of claims **22** to **32**, wherein when R^2 is
2 present the carbon attached to R^2 has the R- configuration.

34. The compound of any one of claims 22 to 33, wherein when R³ is other than H the carbon attached to R³ has the R- configuration.

35. The compound of any one of claims **19** to **21**, wherein W has formula (b).


36. The compound of claim **35**, wherein Z^1 is N.

37. The compound of claim **36**, wherein Z^2 is N.

38. The compound of claim 37, wherein Z³ is N.

39. The compound of any one of claims 33 to 38, wherein m is 1.

40. The compound of any one of claims **33** to **39**, wherein R⁴ is selected from the group consisting of:




41. The compound of any one of claims **33** to **40**, wherein R^{4'} is attached to the pyridyl ring at a position *para* to A.

1 **42.** The compound of any one of claims **33** to **41**, wherein the subscript n
2 is an integer of from 0 to 1.


1 **43.** The compound of any one of claims **33** to **42**, wherein the subscript n
2 is 1; R^4 is attached to the pyridyl ring at a position *ortho* to A and $R^{4'}$ is attached to the
3 pyridyl ring at a position *para* to A.

1 **44.** The compound of any one of claims **33** to **43**, wherein R^4 is selected
2 from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-SR^{4a}$, $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$ and
3 $-SY$.

1 **45.** The compound of any one of claims **33** to **44**, wherein n is 0, 1 or 2,
2 and R^4 is independently selected from the group consisting of H, halogen, OR^{4a} , C_{1-4} alkyl,
3 SR^{4a} , $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2N(R^{4a}R^{4b})$, $NR^{4a}R^{4b}$, $C_{1-4}alkylOR^{4a}$, $C_{1-4}alkylNR^{4a}R^{4b}$, $C_{1-4}alkylCO_2R^{4a}$,
4 $OC_{1-4}alkylOR^{4a}$, $OC_{1-4}alkylN(R^{4a}N^{4b})$, $N(R^{4a})C_{1-4}alkylOR^{4b}$, $N(R^{4a})C_{1-4}alkylN(R^{4a}R^{4b})$, $S(O)_2C_{1-4}alkylN(R^{4a}R^{4b})$,
5 $S(O)_2C_{1-4}alkylN(R^{4a}R^{4b})$,



; each R^{4a} is independently H or C₁₋₄alkyl; each R^{4d} is independently selected from the group consisting of H, OH, NH₂, N(CH₃)₂, CONH₂ and CON(CH₃)₂; and the wavy line indicates the point of attachment to the rest of the molecule, and wherein m is 0 or 1, and R^{4'} is selected from the group consisting of



11
12

1 **46.** The compound of any one of claims **33** to **45**, wherein m is 0 or 1; n is
2 1, 2, or 3; and at least one R^4 is selected from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$,
3 $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2Y$, wherein Y is an optionally substituted member selected
4 from the group consisting of phenyl, benzyl, pyridyl, pyridylmethyl, pyrimidinyl and
5 pyrazolyl.

1 **47.** The compound of any one of claims **33** to **46**, wherein R^4 is selected
2 from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$ and $-NR^{4a}Y$, wherein Y is an optionally
3 substituted member selected from the group consisting of phenyl, benzyl, pyridyl and
4 pyridylmethyl.

1 **48.** The compound of any one of claims **33** to **47**, wherein the optional
2 substituents are selected from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $-$
3 CO_2R^{4a} , $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-X^1OR^{4a}$ and $-X^1NR^{4a}R^{4b}$.


1 **49.** The compound of any one of claims **19** to **21**, wherein W has formula
2 (c).

50. The compound of claim 49, wherein R^5 is selected from the group consisting of $-R^{5a}$, $-\text{CONR}^{5a}R^{5b}$, $-\text{C}(=\text{NR}^{5a})\text{NR}^{5a}R^{5b}$, $-\text{C}(=\text{N}^+R^{5c}R^{5c})\text{NR}^{5a}R^{5b}$, $-\text{C}(=\text{NR}^{5a})R^{5a}$, $-\text{C}(=\text{NH})Y^1$, $-\text{C}(=\text{NR}^{5c})Y^1$ and $-X^3-Y^1$.

51. The compound of any one of claims 49 to 50, wherein Y^1 is selected from the group consisting of optionally substituted phenyl and optionally substituted pyridyl.

52. The compound of any one of claims 49 to 51, wherein R^5 is selected from the group consisting of $-R^{5a}$, $-\text{C}(=\text{NH})\text{NR}^{5a}R^{5b}$, $-\text{C}(=\text{NH})R^{5a}$, $-\text{C}(=\text{NH})Y^1$ and $-X^3-Y^1$.

53. The compound of any one of claims 19 to 21 having the formula:



and pharmaceutically acceptable salts thereof, wherein

R is a member selected from the group consisting of H and C_{1-4} alkyl


R^1 is a member selected from the group consisting of halogen, C_{1-8} alkyl, C_{2-8} alkenyl and C_{2-8} alkynyl;

R^2 is a member selected from the group consisting of H, C_{1-8} alkyl, $-\text{X}-\text{OR}^{2a}$, $-\text{X}-\text{SR}^{2a}$, $-\text{X}-\text{COR}^{2a}$, $-\text{X}-\text{CO}_2R^{2a}$ and $-\text{X}-\text{N}(\text{R}^{2a})_2$; wherein X is C_{1-8} alkylene and each R^{2a} is independently selected from the group consisting of H and C_{1-8} alkyl, or optionally two R^{2a} groups attached to the same nitrogen atom are combined to form a 4-, 5-, 6- or 7-membered ring;

each dashed line independently represents an optional bond, wherein only one optional bond is present;

R^3 is a member selected from the group consisting of H and C_{1-4} alkyl; or is absent when a carbon-carbon double bond is present in the ring to which R^3 would otherwise be attached;

W is a member selected from the group consisting of



wherein

each of Z^1 , Z^2 , and Z^3 is independently N, C, or CH and two of Z^1 , Z^2 , and Z^3 are other than N;

the subscript n is an integer of from 0 to 3 and the subscript m is an integer of from 0 to 1;

R^4 is a member selected from the group consisting of halogen, $-OR^{4a}$, $-OC(O)R^{4a}$, $-NR^{4a}R^{4b}$, $-SR^{4a}$, $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2NR^{4a}R^{4b}$, $-R^{4c}$, $-CN$, $-NO_2$, $-CO_2R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-OC(O)NR^{4a}R^{4b}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)_2R^{4c}$, $-NR^{4a}-C(O)NR^{4a}R^{4b}$, $-X^1OR^{4a}$, $-X^1OC(O)R^{4a}$, $-X^1NR^{4a}R^{4b}$, $-X^1SR^{4a}$, $-X^1S(O)R^{4a}$, $-X^1S(O)_2R^{4a}$, $-X^1CN$, $-X^1NO_2$, $-X^1CO_2R^{4a}$, $-O-X^1CO_2R^{4a}$, $-X^1CONR^{4a}R^{4b}$, $-O-X^1CONR^{4a}R^{4b}$, $-X^1C(O)R^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-S(O)X^1NR^{4a}R^{4b}$, $-S(O)_2X^1NR^{4a}R^{4b}$, $-X^1OC(O)NR^{4a}R^{4b}$, $-X^1NR^{4a}C(O)R^{4a}$, $-X^1NR^{4b}C(O)_2R^{4c}$, $-X^1NR^{4a}C(O)NR^{4a}R^{4b}$, $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2Y$;

$R^{4'}$ is a member selected from the group consisting of $-C(=NH)R^{4a}$, $-C(=NR^{4c})R^{4a}$, $-C(=NH)NR^{4a}R^{4b}$, $-C(=NR^{4c})NR^{4a}R^{4b}$, $-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$, $-X^2-C(=NH)NR^{4a}R^{4b}$, $-X^2-C(=NR^{4c})NR^{4a}R^{4b}$, $-X^2-C(=N^+R^{4c}R^{4c})NR^{4a}R^{4b}$ and $-C(=NR^{4a})NR^{4a}-Y$,

wherein

Y is a five or six-membered aryl, heterocyclyl or aryl- C_{1-2} alkyl, optionally substituted with from one to three substituents selected from the group consisting of halogen, oxo, $-OR^{4a}$, $-OC(O)R^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2NR^{4a}R^{4b}$, $-CN$, $-NO_2$, $-CO_2R^{4a}$, $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-NR^{4b}C(O)R^{4a}$, $-NR^{4b}C(O)_2R^{4c}$, $-X^1R^{4a}$, $-X^1OR^{4a}$, $-X^1SR^{4a}$, $-X^1S(O)R^{4a}$, $-X^1S(O)_2R^{4a}$, $-X^1CN$, $-X^1NO_2$, $-X^1CO_2R^{4a}$, $-X^1CONR^{4a}R^{4b}$, $-X^1C(O)R^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-S(O)X^1NR^{4a}R^{4b}$, $-S(O)_2X^1NR^{4a}R^{4b}$, $-X^1OC(O)NR^{4a}R^{4b}$, $-X^1NR^{4b}C(O)R^{4a}$, $-X^1NR^{4b}C(O)_2R^{4c}$, $-X^1NR^{4a}-C(O)NR^{4a}R^{4b}$, $-X^1OC(O)R^{4a}$, $-X^1NR^{4a}R^{4b}$, $-O-X^1OR^{4a}$, $-O-X^1NR^{4a}R^{4b}$, $-O-X^1CO_2R^{4a}$, $-O-X^1CONR^{4a}R^{4b}$, $-NR^{4b}-X^1OR^{4a}$, $-NR^{4b}-X^1NR^{4a}R^{4b}$, $-NR^{4b}-X^1CO_2R^{4a}$, and $-NR^{4b}-X^1CONR^{4a}R^{4b}$

each X^1 and X^2 are members independently selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene;

each R^{4a} and R^{4b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,

each R^{4c} is independently selected from the group consisting of C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,

optionally any two of R^{4a} , R^{4b} and R^{4c} when part of a common R^4 or $R^{4'}$ substituent can be combined with the atoms to which each is attached to form a saturated or unsaturated, four to nine-membered mono- or bicyclic ring having from 0 to 2 additional heteroatoms as ring members;

each of R^{4a} , R^{4b} and R^{4c} is optionally further substituted with from one to three members selected from the group consisting of -OH, oxo, $-R^m$, $-OR^m$, $-OC(O)NHR^m$, $-OC(O)N(R^m)_2$, -SH, $-SR^m$, $-S(O)R^m$, $-S(O)_2R^m$, $-SO_2NH_2$, $-S(O)_2NHR^m$, $-S(O)_2N(R^m)_2$, $-NHS(O)_2R^m$, $-NR^mS(O)_2R^m$, $-C(O)NH_2$, $-C(O)NHR^m$, $-C(O)N(R^m)_2$, $-C(O)R^m$, $-NHC(O)R^m$, $-NR^mC(O)R^m$, $-NHC(O)NH_2$, $-NR^mC(O)NH_2$, $-NR^mC(O)NHR^m$, $-NHC(O)NHR^m$, $-NR^mC(O)N(R^m)_2$, $-NHC(O)N(R^m)_2$, $-CO_2H$, $-CO_2R^m$, $-NHCO_2R^m$, $-NR^mCO_2R^m$, -CN, $-NO_2$, $-NH_2$, $-NHR^m$, $-N(R^m)_2$, $-NR^mS(O)NH_2$ and $-NR^mS(O)_2NHR^m$, wherein each R^m is independently an unsubstituted C_{1-6} alkyl, benzyl or combined with the atoms to which each is attached to form a saturated four-, five-, six- or seven-membered ring having from 0 to 2 additional heteroatoms as ring members;

R^5 is a member selected from the group consisting of $-R^{5a}$, $-CO_2R^{5c}$, $-CONR^{5a}R^{5b}$, $-C(O)R^{5a}$, $C(=NR^{5a})NR^{5a}R^{5b}$, $-C(=N^+R^{5c}R^{5c})NR^{5a}R^{5b}$, $-C(=NR^{5a})R^{5a}$, $-C(=NR^{5a})Y^1$, $-X^3OR^{5a}$, $-X^3OC(O)R^{5a}$, $-X^3NR^{5a}R^{5b}$, $-X^3SR^{5a}$, $-X^3CN$, $-X^3NO_2$, $-X^3CO_2R^{5a}$, $-X^3CONR^{5a}R^{5b}$, $-X^3C(O)R^{5a}$, $-X^3OC(O)NR^{5a}R^{5b}$, $-X^3NR^{5a}C(O)R^{5a}$, $-X^3NR^{5b}C(O)_2R^{5c}$, $-X^3NR^{5a}C(O)NR^{5a}R^{5b}$, $-X^3-C(=NR^{5a})NR^{5a}R^{5b}$, $-Y^1$, and $-X^3-Y^1$,

wherein

Y^1 is a five or six-membered aryl or heteroaryl ring or aryl- C_{1-2} alkyl, optionally substituted with from one to three substituents selected from the group consisting of halogen, $-OR^{5a}$, $-OC(O)R^{5a}$, $-NR^{5a}R^{5b}$, $-R^{5c}$, $-SR^{5a}$, -CN, $-NO_2$, $-CO_2R^{5a}$, $-CONR^{5a}R^{5b}$, $-C(O)R^{5a}$, $-NR^{5b}C(O)R^{5a}$, $-NR^{5b}C(O)_2R^{5c}$, $-X^3OR^{5a}$, $-X^3SR^{5a}$, $-X^3CN$, $-X^3NO_2$, $-X^3CO_2R^{5a}$, $-X^3CONR^{5a}R^{5b}$, $-X^3C(O)R^{5a}$, $-X^3OC(O)NR^{5a}R^{5b}$, $-X^3NR^{5b}C(O)R^{5a}$, $-X^3NR^{5b}C(O)_2R^{5c}$, $-X^3NR^{5a}-C(O)NR^{5a}R^{5b}$, $-X^3OC(O)R^{5a}$, $-X^3NR^{5a}R^{5b}$, $-O-X^3OR^{5a}$, $-O-X^3NR^{5a}R^{5b}$, $-O-X^3CO_2R^{5a}$, $-O-X^3CONR^{5a}R^{5b}$, $-NR^{5b}-X^3OR^{5a}$, $-NR^{5b}-X^3NR^{5a}R^{5b}$, $-NR^{5b}-X^3CO_2R^{5a}$, and $-NR^{5b}-X^3CONR^{5a}R^{5b}$,

each X^3 is a member independently selected from the group consisting of C_{1-4} alkylene, C_{2-4} alkenylene and C_{2-4} alkynylene;

each R^{5a} and R^{5b} are independently selected from the group consisting of hydrogen, C_{1-8} alkyl, C_{1-8} haloalkyl, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, aryl, heteroaryl, aryl- C_{1-4} alkyl, and aryloxy- C_{1-4} alkyl,


each R^{5c} is independently selected from the group consisting of C₁₋₈ alkyl, C₁₋₈ haloalkyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocycloalkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl, heteroaryl, aryl-C₁₋₄ alkyl, and aryloxy-C₁₋₄ alkyl, optionally any two of R^{5a}, R^{5b} and R^{5c} when part of a common R⁵ substituent can be combined with the atoms to which each is attached to form a saturated four-, five-, six- or seven-membered ring having from 0 to 1 additional heteroatoms as ring members;

each of R^{5a}, R^{5b} and R^{5c} is optionally further substituted with from one to three members selected from the group consisting of -OH, -ORⁿ, -OC(O)NHRⁿ, -OC(O)N(Rⁿ)₂, -SH, -SRⁿ, -S(O)Rⁿ, -S(O)₂Rⁿ, -SO₂NH₂, -S(O)₂NHRⁿ, -S(O)₂N(Rⁿ)₂, -NHS(O)₂Rⁿ, -NRⁿS(O)₂Rⁿ, -C(O)NH₂, -C(O)NHRⁿ, -C(O)N(Rⁿ)₂, -C(O)Rⁿ, -NHC(O)Rⁿ, -NRⁿC(O)Rⁿ, -NHC(O)NH₂, -NRⁿC(O)NH₂, -NRⁿC(O)NHRⁿ, -NHC(O)NHRⁿ, -NRⁿC(O)N(Rⁿ)₂, -NHC(O)N(Rⁿ)₂, -CO₂H, -CO₂Rⁿ, -NHCO₂Rⁿ, -NRⁿCO₂Rⁿ, -CN, -NO₂, -NH₂, -NHRⁿ, -N(Rⁿ)₂, -NRⁿS(O)NH₂ and -NRⁿS(O)₂NHRⁿ, wherein each Rⁿ is independently an unsubstituted C₁₋₆ alkyl or benzyl.

54. The compound of claim **53**, wherein W has formula (a).

55. The compound of any one of claims **53** to **54**, wherein m is 1.

56. The compound of any one of claims **53** to **55**, wherein R^{4'} is selected from the group consisting of:



57. The compound of any one of claims 53 to 56, wherein R⁴ is attached to the phenyl ring at a position *para* to the isoxazole or isoxazoline ring.


58. The compound of any one of claims 53 to 57, wherein the subscript n is an integer of from 0 to 1.

59. The compound of any one of claims 53 to 58, wherein the subscript n is 1; R⁴ is attached to the phenyl ring at a position *ortho* to the isoxazole or isoxazoline ring and R^{4'} is attached to the phenyl ring at a position *para* to the isoxazole or isoxazoline ring.

60. The compound of any one of claims 53 to 59, wherein R⁴ is selected from the group consisting of halogen, -OR^{4a}, -NR^{4a}R^{4b}, -SR^{4a}, -Y, -X¹-Y, -O-Y, -NR^{4a}Y and -SY.


61. The compound of any one of claims 53 to 60, wherein n is 0, 1 or 2, and R⁴ is independently selected from the group consisting of H, halogen, OR^{4a}, C₁₋₄ alkyl, SR^{4a}, S(O)R^{4a}, S(O)₂R^{4a}, S(O)₂N(R^{4a}R^{4b}), NR^{4a}R^{4b}, C₁₋₄alkylOR^{4a}, C₁₋₄alkylNR^{4a}R^{4b}, C₁₋

- 4 $_{4}\text{alkylCO}_2\text{R}^{4a}$, $\text{OC}_{1-4}\text{alkylOR}^{4a}$, $\text{OC}_{1-4}\text{alkylN}(\text{R}^{4a}\text{N}^{4b})$, $\text{N}(\text{R}^{4a})\text{C}_{1-4}\text{alkylOR}^{4b}$, $\text{N}(\text{R}^{4a})\text{C}_{1-}$
5 $_{4}\text{alkylN}(\text{R}^{4a}\text{R}^{4b})$, $\text{S}(\text{O})_2\text{C}_{1-4}\text{alkylN}(\text{R}^{4a}\text{R}^{4b})$,



- 6 $\text{Z-CO}_2\text{H}$, $\text{Z-SO}_2\text{NH}_2$, $\text{Z-SO}_2\text{CH}_3$ and Z-SOCH_3
7 ; each R^{4a} is independently H or C_{1-4} alkyl; each R^{4d} is independently selected from the group
8 consisting of H, OH, NH_2 , $\text{N}(\text{CH}_3)_2$, CONH_2 and $\text{CON}(\text{CH}_3)_2$; and the wavy line indicates the

9 point of attachment to the rest of the molecule, and wherein m is 0 or 1, and R^{4'} is selected
 10 from the group consisting of



11
 12 .

1 **62.** The compound of any one of claims **53** to **59**, wherein m is 0 or 1; n is
 2 1, 2, or 3; and at least one R⁴ is selected from the group consisting of -Y, -X¹-Y, -O-Y,
 3 -NR^{4a}Y, -SY, -S(O)Y and -S(O)₂Y, wherein Y is an optionally substituted member selected
 4 from the group consisting of phenyl, benzyl, pyridyl, pyridylmethyl, pyrimidinyl and
 5 pyrazolyl.

1 **63.** The compound of any one of claims **53** to **62**, wherein R⁴ is selected
 2 from the group consisting of -Y, -X¹-Y, -O-Y and -NR^{4a}Y, wherein Y is an optionally
 3 substituted member selected from the group consisting of phenyl, benzyl, pyridyl and
 4 pyridylmethyl.

1 **64.** The compound of any one of claims **53** to **63**, wherein the optional
 2 substituents are selected from the group consisting of halogen, -OR^{4a}, -NR^{4a}R^{4b}, -R^{4c}, -SR^{4a}, -
 3 CO₂R^{4a}, -CONR^{4a}R^{4b}, -C(O)R^{4a}, -X¹OR^{4a} and -X¹NR^{4a}R^{4b}.

65. The compound of any one of claims 53 to 64, wherein when R^2 is present the carbon attached to R^2 has the R- configuration.

66. The compound of any one of claims 53 to 65, wherein when R^3 is other than H the carbon attached to R^3 has the R- configuration.

67. The compound of claim 53, wherein W has formula (b).


68. The compound of claim 67, wherein Z^1 is N.

69. The compound of claim 67, wherein Z^2 is N.

70. The compound of claim 67, wherein Z^3 is N.

71. The compound of any one of claims 53 to 70, wherein m is 1.

72. The compound of any one of claims 53 to 71, wherein $R^{4'}$ is selected from the group consisting of:




1 **73.** The compound of any one of claims **53** to **72**, wherein $R^{4'}$ is attached to
2 the pyridyl ring at a position *para* to the isoxazole or isoxazoline ring.

1 **74.** The compound of any one of claims **53** to **73**, wherein the subscript n
2 is an integer of from 0 to 1.

1 **75.** The compound of any one of claims **53** to **74**, wherein the subscript n
2 is 1; R^4 is attached to the pyridyl ring at a position *ortho* to the isoxazole or isoxazoline ring
3 and $R^{4'}$ is attached to the pyridyl ring at a position *para* to the isoxazole or isoxazoline ring.


1 **76.** The compound of any one of claims **53** to **75**, wherein R^4 is selected
2 from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-SR^{4a}$, $-Y$, $-X^1-Y$, $-O-Y$, $-NR^{4a}Y$ and
3 $-SY$.

1 **77.** The compound of any one of claims **53** to **76**, wherein n is 0, 1 or 2,
2 and R^4 is independently selected from the group consisting of H, halogen, OR^{4a} , C_{1-4} alkyl,
3 SR^{4a} , $S(O)R^{4a}$, $S(O)_2R^{4a}$, $S(O)_2N(R^{4a}R^{4b})$, $NR^{4a}R^{4b}$, $C_{1-4}alkylOR^{4a}$, $C_{1-4}alkylNR^{4a}R^{4b}$, C_{1-}
4 $4alkylCO_2R^{4a}$, $OC_{1-4}alkylOR^{4a}$, $OC_{1-4}alkylN(R^{4a}R^{4b})$, $N(R^{4a})C_{1-4}alkylOR^{4b}$, $N(R^{4a})C_{1-}$
5 $4alkylN(R^{4a}R^{4b})$, $S(O)_2C_{1-4}alkylN(R^{4a}R^{4b})$,



6 ζ -CO₂H, ζ -SO₂NH₂, ζ -SO₂CH₃ and ζ -SOCH₃

7 ; each R^{4a} is independently H or C₁₋₄alkyl; each R^{4d} is independently selected from the group
 8 consisting of H, OH, NH₂, N(CH₃)₂, CONH₂ and CON(CH₃)₂; and the wavy line indicates the
 9 point of attachment to the rest of the molecule, and wherein m is 0 or 1, and R^{4'} is selected
 10 from the group consisting of



11
12

1 **78.** The compound of any one of claims **53** to **77**, wherein m is 0 or 1; n is
2 1, 2, or 3; and at least one R^4 is selected from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$,
3 $-NR^{4a}Y$, $-SY$, $-S(O)Y$ and $-S(O)_2Y$, wherein Y is an optionally substituted member selected
4 from the group consisting of phenyl, benzyl, pyridyl, pyridylmethyl, pyrimidinyl and
5 pyrazolyl.

1 **79.** The compound of any one of claims **53** to **78**, wherein R^4 is selected
2 from the group consisting of $-Y$, $-X^1-Y$, $-O-Y$ and $-NR^{4a}Y$, wherein Y is an optionally
3 substituted member selected from the group consisting of phenyl, benzyl, pyridyl and
4 pyridylmethyl.

1 **80.** The compound of any one of claims **53** to **79**, wherein the optional
2 substituents are selected from the group consisting of halogen, $-OR^{4a}$, $-NR^{4a}R^{4b}$, $-R^{4c}$, $-SR^{4a}$, $-$
3 CO_2R^{4a} , $-CONR^{4a}R^{4b}$, $-C(O)R^{4a}$, $-X^1OR^{4a}$ and $-X^1NR^{4a}R^{4b}$.

1 **81.** The compound of claim **53**, wherein W has formula (c).

1 **82.** The compound of claim **81**, wherein R^5 is selected from the group
 2 consisting of $-R^{5a}$, $-\text{CONR}^{5a}R^{5b}$, $-\text{C}(=\text{NR}^{5a})\text{NR}^{5a}R^{5b}$, $-\text{C}(=\text{N}^+R^{5c}R^{5c})\text{NR}^{5a}R^{5b}$, $-\text{C}(=\text{NR}^{5a})R^{5a}$,
 3 $-\text{C}(=\text{NR}^{5a})Y^1$ and $-X^3-Y^1$.

1 **83.** The compound of any of claims **81** to **82**, wherein Y^1 is selected from
 2 the group consisting of optionally substituted phenyl and optionally substituted pyridyl.

1 **84.** The compound of claim **83**, wherein R^5 is selected from the group
 2 consisting of $-R^{5a}$, $-\text{C}(=\text{NH})\text{NR}^{5a}R^{5b}$, $-\text{C}(=\text{NH})R^{5a}$, $-\text{C}(=\text{NH})Y^1$ and $-X^3-Y^1$.

1 **85.** A compound selected from the group consisting of: 5-Chloro-
 2 thiophene-2-carboxylic acid [3-(4-bromo-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
 3 5-Chloro-thiophene-2-carboxylic acid [3-(3-bromo-phenyl)-isoxazol-5-ylmethyl]-amide; 5-
 4 Chloro-thiophene-2-carboxylic acid [3-(4-phenoxy-phenyl)-4,5-dihydro-isoxazol-5-
 5 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methylcarbamimidoyl)-
 6 phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-
 7 dimethyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide; 5-Chloro-thiophene-
 8 2-carboxylic acid methyl-{3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-
 9 5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-
 10 methyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide; 5-Chloro-thiophene-2-carboxylic acid
 11 {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide; 1-{[4-(5-
 12 {[(5-Chloro-thiophene-2-carbonyl)-methyl-amino]-methyl}-isoxazol-3-yl)-phenyl]-imino-
 13 methyl}-piperidine-4-carboxylic acid ethyl ester; 5-Chloro-thiophene-2-carboxylic acid {3-
 14 [4-(azetidin-1-yl-imino-methyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide; 5-Chloro-
 15 thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-azetidin-1-ylidene-methyl)-phenyl]-
 16 isoxazol-5-ylmethyl}-methyl-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-ethyl-
 17 N-methyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-methyl-amide; 5-Chloro-thiophene-
 18 2-carboxylic acid methyl-{3-[4-(N-methyl-N-propyl-carbamimidoyl)-phenyl]-isoxazol-5-
 19 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-dimethylamino-ethyl)-
 20 N-methyl-carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-methyl-amide; 1-{[4-(5-
 21 Chloro-thiophene-2-carbonyl)-methyl-amino]-methyl}-isoxazol-3-yl)-phenyl]-imino-
 22 methyl}-piperidine-4-carboxylic acid amide; 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-
 23 (2-methoxy-ethyl)-N-methyl-carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-methyl-amide;
 24 5-Chloro-thiophene-2-carboxylic acid methyl-{3-[4-(N-methyl-N-prop-2-ynyl-

25 carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide; (R)-5-Chloro-thiophene-2-carboxylic
26 acid (1-{3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide; (R)-5-
27 Chloro-thiophene-2-carboxylic acid (1-{3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-
28 isoxazol-5-yl}-ethyl)-amide; (S)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(1-methyl-
29 4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-yl}-ethyl)-amide; (S)-5-Chloro-thiophene-
30 2-carboxylic acid (1-{3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-yl}-ethyl)-
31 amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-
32 pyrrolidin-1-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
33 carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-methylsulfanyl-phenyl]-isoxazol-5-
34 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid (3-phenyl-4,5-dihydro-isoxazol-5-
35 ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid (5-methyl-3-phenyl-4,5-dihydro-
36 isoxazol-5-ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid (3-phenyl-isoxazol-5-
37 ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid (3-p-tolyl-4,5-dihydro-isoxazol-5-
38 ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid (5-methyl-3-p-tolyl-4,5-dihydro-
39 isoxazol-5-ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid (3-p-tolyl-isoxazol-5-
40 ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-methoxy-phenyl)-4,5-dihydro-
41 isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-methoxy-phenyl)-5-
42 methyl-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-
43 methoxy-phenyl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-
44 chloro-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic
45 acid [3-(4-chloro-phenyl)-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-
46 thiophene-2-carboxylic acid [3-(4-chloro-phenyl)-isoxazol-5-ylmethyl]-amide; 4-(5-{{(5-
47 Chloro-thiophene-2-carbonyl)-amino]-methyl}}-4,5-dihydro-isoxazol-3-yl)-benzoic acid
48 methyl ester; 4-(5-{{(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}}-isoxazol-3-yl)-benzoic
49 acid methyl ester; 4-(5-{{(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}}-4,5-dihydro-
50 isoxazol-3-yl)-benzoic acid; 4-(5-{{(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}}-
51 isoxazol-3-yl)-benzoic acid; 5-Chloro-thiophene-2-carboxylic acid [3-(4-amino-phenyl)-4,5-
52 dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid (3-{4-[(2-
53 diethylamino-ethyl)-methyl-carbamoyl]-phenyl}-isoxazol-5-ylmethyl)-amide; 5-Chloro-
54 thiophene-2-carboxylic acid [3-(1-ethyl-piperidin-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-
55 amide; 5-Chloro-thiophene-2-carboxylic acid [3-(1-benzyl-piperidin-4-yl)-4,5-dihydro-
56 isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(1-isopropyl-piperidin-
57 4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(1-
58 carbamimidoyl-piperidin-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-

59 2-carboxylic acid {3-[1-(1-imino-ethyl)-piperidin-4-yl]-4,5-dihydro-isoxazol-5-ylmethyl}-
60 amide; 5-Chloro-thiophene-2-carboxylic acid {3-[1-(imino-phenyl-methyl)-piperidin-4-yl]-
61 4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[1-
62 (imino-pyridin-2-yl-methyl)-piperidin-4-yl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-
63 Chloro-thiophene-2-carboxylic acid (3-piperidin-4-yl-isoxazol-5-ylmethyl)-amide; 5-Chloro-
64 thiophene-2-carboxylic acid [3-(1-ethyl-piperidin-4-yl)-isoxazol-5-ylmethyl]-amide; 5-
65 Chloro-thiophene-2-carboxylic acid [3-(1-benzyl-piperidin-4-yl)-isoxazol-5-ylmethyl]-amide;
66 5-Chloro-thiophene-2-carboxylic acid [3-(1-carbamimidoyl-piperidin-4-yl)-isoxazol-5-
67 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[1-(1-imino-ethyl)-piperidin-4-
68 yl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[1-(imino-phenyl-
69 methyl)-piperidin-4-yl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid
70 [3-(2'-formyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-
71 carboxylic acid [3-(2'-chloro-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-
72 Chloro-thiophene-2-carboxylic acid [3-(2'-methylsulfanyl-biphenyl-4-yl)-4,5-dihydro-
73 isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-4-yl-
74 phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-
75 (4-pyridin-3-yl-phenyl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid
76 [3-(2'-azepan-1-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-
77 thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-5'-methoxy-biphenyl-4-yl)-4,5-
78 dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(2'-
79 hydroxymethyl-4'-methoxy-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-
80 Chloro-thiophene-2-carboxylic acid [3-(3'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-isoxazol-5-
81 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4'-dimethylaminomethyl-
82 biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4'-
83 pyrrolidin-1-ylmethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-
84 carboxylic acid [3-(2'-dimethylaminomethyl-5'-methoxy-biphenyl-4-yl)-isoxazol-5-
85 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-2-yl-phenyl)-isoxazol-
86 5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-benzylamino-phenyl)-4,5-
87 dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-
88 phenylamino-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-
89 carboxylic acid {3-[4-(4-fluoro-phenylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-
90 amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-2-ylamino)-phenyl]-4,5-
91 dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-methyl-
92 pyridin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-

93 carboxylic acid (3-{4-[(2-amino-pyrimidin-4-yl)-methyl-amino]-phenyl}-4,5-dihydro-
94 isoxazol-5-ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-ethyl-2H-
95 pyrazol-3-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
96 carboxylic acid [3-(4-morpholin-4-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-
97 Chloro-thiophene-2-carboxylic acid {3-[4-(2-methoxymethyl-pyrrolidin-1-yl)-phenyl]-4,5-
98 dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-
99 (5-oxo-[1,4]oxazepan-4-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-
100 thiophene-2-carboxylic acid [3-(4-phenylamino-phenyl)-isoxazol-5-ylmethyl]-amide; 5-
101 Chloro-thiophene-2-carboxylic acid {3-[4-(4-methoxy-phenylamino)-phenyl]-isoxazol-5-
102 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-2-ylamino)-phenyl]-
103 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(methyl-pyridin-4-
104 yl-amino)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-
105 pyrrolidin-1-yl-phenyl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid
106 [3-(4-morpholin-4-yl-phenyl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic
107 acid [3-(4-benzylamino-phenyl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-
108 carboxylic acid {3-[4-(1-methyl-piperidin-4-ylamino)-phenyl]-isoxazol-5-ylmethyl}-amide;
109 2-[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-
110 phenylamino]-benzoic acid methyl ester; 3-[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-
111 methyl}-4,5-dihydro-isoxazol-3-yl)-phenylamino]-benzoic acid methyl ester; 4-[4-(5-{[(5-
112 Chloro-thiophene-2-carbonyl)-amino]-methyl}-4,5-dihydro-isoxazol-3-yl)-phenylamino]-
113 benzoic acid methyl ester; 5-Chloro-thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-
114 ylmethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-
115 carboxylic acid [3-(3'-dimethylaminomethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-
116 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(3'-pyrrolidin-1-ylmethyl-
117 biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic
118 acid [3-(4'-dimethylaminomethyl-biphenyl-3-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-
119 Chloro-thiophene-2-carboxylic acid [3-(4'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-4,5-dihydro-
120 isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(2'-
121 dimethylaminomethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-
122 carboxylic acid [3-(3'-dimethylaminomethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide; 5-
123 Chloro-thiophene-2-carboxylic acid [3-(3'-pyrrolidin-1-ylmethyl-biphenyl-3-yl)-isoxazol-5-
124 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4'-dimethylaminomethyl-
125 biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4'-
126 pyrrolidin-1-ylmethyl-biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-

127 carboxylic acid [3-(3-pyridin-3-yl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-
 128 Chloro-thiophene-2-carboxylic acid [3-(3-pyridin-4-yl-phenyl)-4,5-dihydro-isoxazol-5-
 129 ylmethyl]-amide; and 5-Chloro-thiophene-2-carboxylic acid [3-(3-pyridin-3-yl-phenyl)-
 130 isoxazol-5-ylmethyl]-amide.

1 **86.** A compound selected from the group consisting of: 5-Chloro-
 2 thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-4,5-dihydro-
 3 isoxazol-5-ylmethyl}-amide; 1-{[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-
 4 4,5-dihydro-isoxazol-3-yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid ethyl ester;
 5 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methyl-N-prop-2-ynyl-carbamimidoyl)-
 6 phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid (3-
 7 {4-[N-(2-methoxy-ethyl)-N-methyl-carbamimidoyl]-phenyl}-4,5-dihydro-isoxazol-5-
 8 ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-furan-2-ylmethyl-N-
 9 methyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-
 10 thiophene-2-carboxylic acid {3-[4-(N-methyl-N-prop-2-ynyl-carbamimidoyl)-phenyl]-
 11 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-methoxy-
 12 ethyl)-N-methyl-carbamimidoyl]-phenyl}-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide;
 13 1-{[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-5-methyl-4,5-dihydro-isoxazol-
 14 3-yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid amide; 5-Chloro-thiophene-2-
 15 carboxylic acid {3-[4-(N,N-dimethyl- carbamidoyl)-2-fluoro-phenyl]-dihydro-isoxazol-5-
 16 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(N,N-
 17 dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-
 18 thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(1-methyl-4,5-dihydro-1H-imidazol-2-
 19 yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid
 20 {3-[2-dimethylamino-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-
 21 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-
 22 carbamidoyl)-2-pyrrolidin-1-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-
 23 Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-
 24 pyrrolidin-1-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
 25 carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-pyrrolidin-1-yl-phenyl]-4,5-dihydro-
 26 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-
 27 dihydro-1H-imidazol-2-yl)-2-morpholin-4-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-
 28 amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-
 29 morpholin-4-yl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-

30 carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-morpholin-4-yl-phenyl]-4,5-dihydro-
31 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-[(2-methoxy-
32 ethyl)-methyl-amino]-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-
33 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid (3-{4-(imino-pyrrolidin-
34 1-yl-methyl)-2-[(2-methoxy-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-
35 ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid (3-{4-(imino-piperidin-1-yl-methyl)-
36 2-[(2-methoxy-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-amide; 5-
37 Chloro-thiophene-2-carboxylic acid {3-[2-azepan-1-yl-4-(N,N-dimethyl-carbamimidoyl)-
38 phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-
39 [4-(2-methyl-pyrrolidine-1-carbonyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-
40 Chloro-thiophene-2-carboxylic acid {3-[4-(pyrrolidine-1-carbonyl)-phenyl]-isoxazol-5-
41 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(2'-amino-biphenyl-4-yl)-4,5-
42 dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(2'-
43 trifluoromethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-
44 2-carboxylic acid [3-(2'-methoxy-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-
45 Chloro-thiophene-2-carboxylic acid [3-(2'-methoxymethyl-biphenyl-4-yl)-isoxazol-5-
46 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-3-yl-phenyl)-4,5-
47 dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(5'-chloro-2'-
48 {[(2-methoxy-ethyl)-methyl-amino]-methyl}-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-
49 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-ylmethyl-
50 biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(3'-
51 dimethylaminomethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-
52 carboxylic acid {3-[4-(2-chloro-pyridin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-
53 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyrimidin-4-ylamino)-
54 phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-
55 [4-(1-methyl-1H-pyrazol-3-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-
56 Chloro-thiophene-2-carboxylic acid {3-[4-(4-methyl-piperazin-1-yl)-phenyl]-4,5-dihydro-
57 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-
58 piperidin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-
59 2-carboxylic acid (3-{4-[(2-dimethylamino-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-
60 isoxazol-5-ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-oxo-piperidin-1-
61 yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; and 5-Chloro-thiophene-2-carboxylic
62 acid {3-[4-(3-oxo-thiomorpholin-4-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide.

1 **87.** A compound selected from the group consisting of: 5-Chloro-
2 thiophene-2-carboxylic acid [3-(4-cyano-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-
3 Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-4,5-
4 dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-
5 4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-
6 thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-
7 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-
8 dimethylamino-ethyl)-N-methyl-carbamimidoyl]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-
9 amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-methyl-N-propyl-carbamimidoyl)-
10 phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-
11 [4-(N-ethyl-N-methyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-
12 Chloro-thiophene-2-carboxylic acid [3-(4-cyano-phenyl)-isoxazol-5-ylmethyl]-amide; 5-
13 Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-isoxazol-
14 5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-
15 imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid
16 {3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-
17 thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-
18 ylmethyl}-amide; 1-{[4-(5-{[(5-Chloro-thiophene-2-carbonyl)-amino]-methyl}-isoxazol-3-
19 yl)-phenyl]-imino-methyl}-piperidine-4-carboxylic acid ethyl ester; 5-Chloro-thiophene-2-
20 carboxylic acid {3-[4-(azetidin-1-yl-imino-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-
21 Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-azetidin-1-ylidene-methyl)-phenyl]-
22 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N-ethyl-N-
23 methyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
24 carboxylic acid {3-[4-(N-methyl-N-propyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-
25 amide; 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-dimethylamino-ethyl)-N-methyl-
26 carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid
27 {3-[4-(N-furan-2-ylmethyl-N-methyl-carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide;
28 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-methoxy-ethyl)-N-methyl-
29 carbamimidoyl]-phenyl}-isoxazol-5-ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid
30 methyl-{3-[4-(N-methylcarbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide; (R)-5-Chloro-
31 thiophene-2-carboxylic acid (1-{3-[4-(N,N-dimethyl-carbamimidoyl)-phenyl]-isoxazol-5-yl}-
32 ethyl)-amide; (R)-5-Chloro-thiophene-2-carboxylic acid (1-{3-[4-(1-methyl-4,5-dihydro-1H-
33 imidazol-2-yl)-phenyl]-isoxazol-5-yl}-ethyl)-amide; (R)-5-Chloro-thiophene-2-carboxylic
34 acid (1-{3-[4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-yl}-ethyl)-amide; (S)-5-

35 Chloro-thiophene-2-carboxylic acid (1-{3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-
36 isoxazol-5-yl}-ethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-
37 carbamimidoyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-
38 thiophene-2-carboxylic acid {5-methyl-3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-
39 phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-
40 [4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-
41 amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-phenyl]-5-
42 methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 1-{[4-(5-{[(5-Chloro-thiophene-2-
43 carbonyl)-amino]-methyl}-5-methyl-4,5-dihydro-isoxazol-3-yl)-phenyl]-imino-methyl}-
44 piperidine-4-carboxylic acid ethyl ester; 5-Chloro-thiophene-2-carboxylic acid {3-[4-
45 (azetidin-1-yl-imino-methyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-
46 Chloro-thiophene-2-carboxylic acid {3-[4-(azetidin-1-yl-azetidin-1-ylidene-methyl)-phenyl]-
47 5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-
48 [4-(N-ethyl-N-methyl-carbamimidoyl)-phenyl]-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-
49 amide; 5-Chloro-thiophene-2-carboxylic acid (3-{4-[N-(2-dimethylamino-ethyl)-N-methyl-
50 carbamimidoyl]-phenyl}-5-methyl-4,5-dihydro-isoxazol-5-ylmethyl)-amide; 5-Chloro-
51 thiophene-2-carboxylic acid {3-[4-(N-furan-2-ylmethyl-N-methyl-carbamimidoyl)-phenyl]-
52 5-methyl-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {5-
53 methyl-3-[4-(N-methylcarbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-
54 Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(1-methyl-4,5-dihydro-1H-imidazol-2-
55 yl)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-
56 4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-
57 2-carboxylic acid {3-[2-fluoro-4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-
58 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(imino-
59 piperidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-
60 thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-morpholin-4-yl-phenyl]-
61 4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid (3-{4-(N,N-
62 dimethyl-carbamimidoyl)-2-[(2-methoxy-ethyl)-methyl-amino]-phenyl}-4,5-dihydro-
63 isoxazol-5-ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-azepan-1-yl-4-(1-
64 methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-
65 Chloro-thiophene-2-carboxylic acid {3-[2-azepan-1-yl-4-(imino-pyrrolidin-1-yl-methyl)-
66 phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-
67 [2-azepan-1-yl-4-(imino-piperidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-
68 amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(N,N-dimethyl-

69 carbamimidoyl)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid
70 {3-[2-dimethylamino-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-
71 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-dimethylamino-4-(imino-
72 pyrrolidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
73 carboxylic acid {3-[2-dimethylamino-4-(imino-piperidin-1-yl-methyl)-phenyl]-isoxazol-5-
74 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-
75 carbamimidoyl)-2-morpholin-4-yl-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-
76 2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-morpholin-4-yl-phenyl]-
77 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-
78 1-yl-methyl)-2-morpholin-4-yl-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
79 carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-morpholin-4-yl-phenyl]-isoxazol-5-
80 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-
81 carbamimidoyl)-2-methoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-
82 thiophene-2-carboxylic acid {3-[2-methoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-
83 phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-
84 [4-(imino-pyrrolidin-1-yl-methyl)-2-methoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-
85 amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-
86 methoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic
87 acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-ethoxy-phenyl]-4,5-dihydro-isoxazol-5-
88 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(1-methyl-4,5-
89 dihydro-1H-imidazol-2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-
90 thiophene-2-carboxylic acid {3-[2-ethoxy-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-
91 dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-
92 dimethyl-carbamimidoyl)-2-isopropoxy-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-
93 Chloro-thiophene-2-carboxylic acid {3-[2-isopropoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-
94 2-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic
95 acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-isopropoxy-phenyl]-4,5-dihydro-isoxazol-5-
96 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclopentyloxy-4-(N,N-
97 dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-
98 thiophene-2-carboxylic acid {3-[2-cyclopentyloxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-
99 yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid
100 {3-[2-cyclopentyloxy-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-
101 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-cyclohexyloxy-4-(N,N-
102 dimethyl-carbamimidoyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-

103 thiophene-2-carboxylic acid {3-[2-cyclohexyloxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-
104 yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid
105 {3-[2-cyclohexyloxy-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-4,5-dihydro-isoxazol-5-
106 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-
107 carbamimidoyl)-2-methoxy-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
108 carboxylic acid {3-[2-methoxy-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-
109 5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-
110 methyl)-2-methoxy-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic
111 acid {3-[4-(imino-piperidin-1-yl-methyl)-2-methoxy-phenyl]-isoxazol-5-ylmethyl}-amide; 5-
112 Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-ethoxy-phenyl]-
113 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(1-
114 methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-
115 thiophene-2-carboxylic acid {3-[2-ethoxy-4-(imino-pyrrolidin-1-yl-methyl)-phenyl]-
116 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-ethoxy-4-(imino-
117 piperidin-1-yl-methyl)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
118 carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-isopropoxy-phenyl]-isoxazol-5-
119 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-isopropoxy-4-(1-methyl-4,5-
120 dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
121 carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-isopropoxy-phenyl]-isoxazol-5-
122 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-
123 2-isopropoxy-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid
124 {3-[4-(N,N-dimethyl-carbamimidoyl)-2-phenoxy-phenyl]-isoxazol-5-ylmethyl}-amide; 5-
125 Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-
126 phenoxy-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-
127 (imino-pyrrolidin-1-yl-methyl)-2-phenoxy-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-
128 thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-(4-methoxy-phenoxy)-
129 phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-(4-
130 methoxy-phenoxy)-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-phenyl]-isoxazol-5-
131 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-
132 methyl)-2-(4-methoxy-phenoxy)-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-
133 2-carboxylic acid {3-[4-(N,N-dimethyl-carbamimidoyl)-2-methylsulfanyl-phenyl]-4,5-
134 dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(1-methyl-
135 4,5-dihydro-1H-imidazol-2-yl)-2-methylsulfanyl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-
136 amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-1-yl-methyl)-2-

137 methylsulfanyl-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
138 carboxylic acid {3-[4-(imino-piperidin-1-yl-methyl)-2-methylsulfanyl-phenyl]-4,5-dihydro-
139 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(N,N-dimethyl-
140 carbamimidoyl)-2-methylsulfanyl-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-
141 2-carboxylic acid {3-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-2-methylsulfanyl-phenyl]-
142 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(imino-pyrrolidin-
143 1-yl-methyl)-2-methylsulfanyl-phenyl]-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
144 carboxylic acid [3-(4-amino-phenyl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-
145 carboxylic acid [3-(4-dimethylcarbamoyl-phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide;
146 5-Chloro-thiophene-2-carboxylic acid {3-[4-(azetidine-1-carbonyl)-phenyl]-4,5-dihydro-
147 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyrrolidine-1-
148 carbonyl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
149 carboxylic acid (3-{4-[(2-diethylamino-ethyl)-methyl-carbamoyl]-phenyl}-4,5-dihydro-
150 isoxazol-5-ylmethyl)-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-4-yl-
151 phenyl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(2'-
152 dimethylaminomethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-
153 thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-4,5-dihydro-
154 isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(2'-piperidin-1-
155 ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-
156 carboxylic acid [3-(2'-morpholin-4-ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-
157 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(2'-{[(2-methoxy-ethyl)-methyl-
158 amino]-methyl}-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-
159 thiophene-2-carboxylic acid {3-[2'-(4-methyl-piperazin-1-ylmethyl)-biphenyl-4-yl]-4,5-
160 dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(2'-{[(2-
161 dimethylamino-ethyl)-methyl-amino]-methyl}-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-
162 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(5'-methoxy-2'-pyrrolidin-1-
163 ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-
164 carboxylic acid [3-(5'-chloro-2'-dimethylaminomethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-
165 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(5'-chloro-2'-pyrrolidin-1-
166 ylmethyl-biphenyl-4-yl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-
167 carboxylic acid [3-(2'-formyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-
168 thiophene-2-carboxylic acid [3-(2'-dimethylaminomethyl-biphenyl-4-yl)-isoxazol-5-
169 ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(5'-methoxy-2'-pyrrolidin-1-
170 ylmethyl-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid

171 [3-(2'-dimethylaminomethyl-4'-methoxy-biphenyl-4-yl)-isoxazol-5-ylmethyl]-amide; 5-
 172 Chloro-thiophene-2-carboxylic acid [3-(4'-methoxy-2'-pyrrolidin-1-ylmethyl-biphenyl-4-yl)-
 173 isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid [3-(4-pyridin-2-yl-
 174 phenyl)-4,5-dihydro-isoxazol-5-ylmethyl]-amide; 5-Chloro-thiophene-2-carboxylic acid {3-
 175 [4-(pyridin-3-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-
 176 thiophene-2-carboxylic acid {3-[4-(pyridin-4-ylamino)-phenyl]-4,5-dihydro-isoxazol-5-
 177 ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(methyl-pyridin-4-yl-amino)-
 178 phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid (3-
 179 {4-[(3-dimethylamino-propyl)-methyl-amino]-phenyl}-4,5-dihydro-isoxazol-5-ylmethyl)-
 180 amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(2-oxo-pyrrolidin-1-yl)-phenyl]-4,5-
 181 dihydro-isoxazol-5-ylmethyl}-amide; 4-[4-(5-[(5-Chloro-thiophene-2-carbonyl)-amino]-
 182 methyl)-4,5-dihydro-isoxazol-3-yl]-phenyl]-3-oxo-piperazine-1-carboxylic acid benzyl ester;
 183 5-Chloro-thiophene-2-carboxylic acid {3-[4-(3-oxo-morpholin-4-yl)-phenyl]-4,5-dihydro-
 184 isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[4-(3-methyl-2-oxo-
 185 imidazolidin-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-
 186 carboxylic acid {3-[4-(5-oxo-[1,4]oxazepan-4-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-
 187 amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-4-(pyridin-4-ylamino)-phenyl]-
 188 4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-thiophene-2-carboxylic acid {3-[2-fluoro-
 189 4-(4-methyl-[1,4]diazepan-1-yl)-phenyl]-4,5-dihydro-isoxazol-5-ylmethyl}-amide; 5-Chloro-
 190 thiophene-2-carboxylic acid {3-[4-(pyridin-3-ylamino)-phenyl]-isoxazol-5-ylmethyl}-amide;
 191 5-Chloro-thiophene-2-carboxylic acid {3-[4-(pyridin-4-ylamino)-phenyl]-isoxazol-5-
 192 ylmethyl}-amide; and 5-Chloro-thiophene-2-carboxylic acid [3-(2'-pyrrolidin-1-ylmethyl-
 193 biphenyl-3-yl)-isoxazol-5-ylmethyl]-amide.

1 **88.** A compound selected from the group consisting of: 5-Chloro-N-((3-
 2 (4-(2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-
 3 carboxamide; 5-Chloro-N-((3-(4-(3-fluoro-2-oxopyridin-1(2H)-yl)phenyl)-4,5-
 4 dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(5-fluoro-2-
 5 oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-
 6 Chloro-N-((3-(4-(3-methyl-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-
 7 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(3-methoxy-2-oxopyridin-1(2H)-
 8 yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(3-
 9 chloro-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-
 10 carboxamide; 5-Chloro-N-((3-(4-(4-methyl-2-oxopyridin-1(2H)-yl)phenyl)-4,5-

11 dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(5-methyl-2-
12 oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-
13 Chloro-N-((3-(4-(5-methoxy-2-oxopyridin-1(2H)-yl)phenyl)-4,5-dihydroisoxazol-5-
14 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(5-chloro-2-oxopyridin-1(2H)-
15 yl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-
16 iodophenyl)isoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(2-
17 oxopyridin-1(2H)-yl)phenyl)isoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-
18 ((1-(4-(2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-
19 carboxamide; 5-Chloro-N-((1-(4-(3-fluoro-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-
20 4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(4-(3-chloro-2-oxopyridin-1(2H)-
21 yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(4-(3-
22 methyl-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-
23 carboxamide; 5-Chloro-N-((1-(4-(3-methoxy-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-
24 triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(4-(4-methyl-2-oxopyridin-
25 1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(4-
26 (5-methyl-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-
27 carboxamide; 5-Chloro-N-((1-(4-(5-methoxy-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-
28 triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(4-(5-fluoro-2-oxopyridin-
29 1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(4-
30 (5-chloro-2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-
31 carboxamide; 5-Chloro-N-((1-(4-(2-oxopiperidin-1-yl)phenyl)-1H-1,2,3-triazol-4-
32 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(4-(3-oxomorpholino)phenyl)-1H-
33 1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(4-(3-
34 oxothiomorpholino)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-
35 Chloro-N-((1-(4-(5-oxo-1,4-oxazepan-4-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-
36 2-carboxamide; 5-Chloro-N-((1-(2-fluoro-4-iodophenyl)-1H-1,2,3-triazol-4-
37 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(2-fluoro-4-(2-oxopyridin-1(2H)-
38 yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(2-fluoro-
39 4-(3-oxomorpholino)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-
40 Chloro-N-((1-(2-hydroxy-4-(2-oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-
41 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(2-(4-methylpiperazin-1-yl)-4-(2-
42 oxopyridin-1(2H)-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-
43 Chloro-N-((1-(5-(2-oxopyridin-1(2H)-yl)pyridin-2-yl)-1H-1,2,3-triazol-4-
44 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(6-(2-oxopyridin-1(2H)-yl)pyridin-3-

45 yl)isoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(2-oxopyridin-1(2H)-
46 yl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(4-(1-
47 methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-
48 carboxamide; N-((1-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-
49 5-chlorothiophene-2-carboxamide; N-((1-(4-(N-Ethyl-N-methylcarbamimidoyl)phenyl)-1H-
50 1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide; N-((1-(4-(N-Methyl-N-
51 propylcarbamimidoyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-
52 carboxamide; N-((1-(4-(N-(2-Methoxyethyl)-N-methylcarbamimidoyl)phenyl)-1H-1,2,3-
53 triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide; N-((1-(4-(Azetidin-1-
54 yl(imino)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide; 5-
55 Chloro-N-((1-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1H-1,2,3-triazol-4-
56 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(4-(imino(piperidin-1-
57 yl)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; N-((1-(4-(N,N-
58 Dimethylcarbamimidoyl)-2-fluorophenyl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-
59 2-carboxamide; N-((1-(4-(Azetidin-1-yl(imino)methyl)-2-fluorophenyl)-1H-1,2,3-triazol-4-
60 yl)methyl)-5-chlorothiophene-2-carboxamide; 5-Chloro-N-((1-(2-fluoro-4-(imino(pyrrolidin-
61 1-yl)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-
62 (2-fluoro-4-(imino(piperidin-1-yl)methyl)phenyl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-
63 carboxamide; 5-Chloro-N-((1-(2-fluoro-4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-
64 1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; N-((1-(4-(N,N-
65 Dimethylcarbamimidoyl)phenyl)-1H-pyrazol-4-yl)methyl)-5-chlorothiophene-2-
66 carboxamide; 5-Chloro-N-((1-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1H-
67 pyrazol-4-yl)methyl)thiophene-2-carboxamide; N-((1-(4-(Azetidin-1-
68 yl(imino)methyl)phenyl)-1H-pyrazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide; 5-
69 Chloro-N-((1-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1H-pyrazol-4-yl)methyl)thiophene-
70 2-carboxamide; 5-Chloro-N-((1-(4-(imino(piperidin-1-yl)methyl)phenyl)-1H-pyrazol-4-
71 yl)methyl)thiophene-2-carboxamide; N-((5-(4-(N,N-
72 Dimethylcarbamimidoyl)phenyl)isoxazol-3-yl)methyl)-5-chlorothiophene-2-carboxamide; 5-
73 Chloro-N-((5-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)isoxazol-3-
74 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((5-(4-(imino(pyrrolidin-1-
75 yl)methyl)phenyl)isoxazol-3-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((5-(4-
76 (imino(piperidin-1-yl)methyl)phenyl)isoxazol-3-yl)methyl)thiophene-2-carboxamide; N-((3-
77 (4-(N,N-Dimethylcarbamimidoyl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)-5-chlorothiophene-2-
78 carboxamide; 5-Chloro-N-((3-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)phenyl)-1,2,4-

79 oxadiazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(imino(pyrrolidin-1-
80 yl)methyl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-
81 (4-(imino(piperidin-1-yl)methyl)phenyl)-1,2,4-oxadiazol-5-yl)methyl)thiophene-2-
82 carboxamide; N-((2-(4-(N,N-Dimethylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-
83 chlorothiophene-2-carboxamide; 5-Chloro-N-((2-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-
84 yl)phenyl)oxazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((2-(4-
85 (imino(pyrrolidin-1-yl)methyl)phenyl)oxazol-4-yl)methyl)thiophene-2-carboxamide; 5-
86 Chloro-N-((2-(4-(imino(piperidin-1-yl)methyl)phenyl)oxazol-4-yl)methyl)thiophene-2-
87 carboxamide; Ethyl 1-((4-(4-((2-chlorothiophene-5-carboxamido)methyl)oxazol-2-
88 yl)phenyl)(imino)methyl)piperidine-4-carboxylate; N-((2-(4-(N-Ethyl-N-
89 methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide; N-((2-
90 (4-(N-Methyl-N-propylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-
91 carboxamide; N-((2-(4-(N-(2-(Dimethylamino)ethyl)-N-
92 methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide; N-((2-
93 (4-(N-(2-Methoxyethyl)-N-methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-
94 chlorothiophene-2-carboxamide; 1-((4-(4-((2-Chlorothiophene-5-
95 carboxamido)methyl)oxazol-2-yl)phenyl)(imino)methyl)piperidine-4-carboxamide; N-((2-(4-
96 (N-Methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide; N-
97 ((2-(4-(N-(Furan-2-ylmethyl)-N-methylcarbamimidoyl)phenyl)oxazol-4-yl)methyl)-5-
98 chlorothiophene-2-carboxamide; N-((2-(4-(N,N-Dimethylcarbamimidoyl)phenyl)thiazol-4-
99 yl)methyl)-5-chlorothiophene-2-carboxamide; 5-Chloro-N-((2-(4-(1-methyl-4,5-dihydro-1H-
100 imidazol-2-yl)phenyl)thiazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((2-(4-
101 (imino(pyrrolidin-1-yl)methyl)phenyl)thiazol-4-yl)methyl)thiophene-2-carboxamide; 5-
102 Chloro-N-((2-(4-(imino(piperidin-1-yl)methyl)phenyl)thiazol-4-yl)methyl)thiophene-2-
103 carboxamide; Ethyl 1-((4-(4-((2-chlorothiophene-5-carboxamido)methyl)thiazol-2-
104 yl)phenyl)(imino)methyl)piperidine-4-carboxylate; N-((2-(4-(N-(2-(Dimethylamino)ethyl)-
105 N-methylcarbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide; N-
106 ((2-(4-(N-(Furan-2-ylmethyl)-N-methylcarbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-
107 chlorothiophene-2-carboxamide; N-((2-(4-(N-Methyl-N-(prop-2-
108 ynyl)carbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide; N-((2-
109 (4-(N-Methylcarbamimidoyl)phenyl)thiazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide;
110 5-Chloro-N-((4-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)oxazol-2-yl)methyl)thiophene-2-
111 carboxamide; N-((4-(4-(N,N-Dimethylcarbamimidoyl)phenyl)thiazol-2-yl)methyl)-5-
112 chlorothiophene-2-carboxamide; 5-Chloro-N-((4-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-

113 yl)phenyl)thiazol-2-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((4-(4-
114 (imino(pyrrolidin-1-yl)methyl)phenyl)thiazol-2-yl)methyl)thiophene-2-carboxamide; 5-
115 Chloro-N-((4-(4-(imino(piperidin-1-yl)methyl)phenyl)thiazol-2-yl)methyl)thiophene-2-
116 carboxamide; N-((1-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-1H-imidazol-4-yl)methyl)-5-
117 chlorothiophene-2-carboxamide; 5-Chloro-N-((1-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-
118 yl)phenyl)-1H-imidazol-4-yl)methyl)thiophene-2-carboxamide; N-((1-(4-(Azetidin-1-
119 yl(imino)methyl)phenyl)-1H-imidazol-4-yl)methyl)-5-chlorothiophene-2-carboxamide; 5-
120 Chloro-N-((1-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-1H-imidazol-4-yl)methyl)thiophene-
121 2-carboxamide; 5-Chloro-N-((1-(4-(imino(piperidin-1-yl)methyl)phenyl)-1H-imidazol-4-
122 yl)methyl)thiophene-2-carboxamide; N-((5-(4-(N,N-Dimethylcarbamimidoyl)phenyl)pyridin-
123 3-yl)methyl)-5-chlorothiophene-2-carboxamide; 5-Chloro-N-((5-(4-(1-methyl-4,5-dihydro-
124 1H-imidazol-2-yl)phenyl)pyridin-3-yl)methyl)thiophene-2-carboxamide; N-((5-(4-(Azetidin-
125 1-yl(imino)methyl)phenyl)pyridin-3-yl)methyl)-5-chlorothiophene-2-carboxamide; 5-Chloro-
126 N-((5-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)pyridin-3-yl)methyl)thiophene-2-
127 carboxamide; 5-Chloro-N-((5-(4-(imino(piperidin-1-yl)methyl)phenyl)pyridin-3-
128 yl)methyl)thiophene-2-carboxamide; N-((5-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-2-
129 fluorophenyl)methyl)-5-chlorothiophene-2-carboxamide; 5-Chloro-N-((5-(4-(1-methyl-4,5-
130 dihydro-1H-imidazol-2-yl)phenyl)-2-fluorophenyl)methyl)thiophene-2-carboxamide; 5-
131 Chloro-N-((5-(4-(imino(pyrrolidin-1-yl)methyl)phenyl)-2-fluorophenyl)methyl)thiophene-2-
132 carboxamide; N-((3-(4-(N,N-Dimethylcarbamimidoyl)phenyl)-4-fluorophenyl)methyl)-5-
133 chlorothiophene-2-carboxamide; 5-Chloro-N-((3-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-
134 yl)phenyl)-4-fluorophenyl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-
135 (imino(pyrrolidin-1-yl)methyl)phenyl)-4-fluorophenyl)methyl)thiophene-2-carboxamide; N-
136 ((3-(4-(N,N-Dimethylcarbamimidoyl)phenyl)phenyl)methyl)-5-chlorothiophene-2-
137 carboxamide; 5-Chloro-N-((3-(4-(1-methyl-4,5-dihydro-1H-imidazol-2-
138 yl)phenyl)phenyl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(imino(pyrrolidin-1-
139 yl)methyl)phenyl)phenyl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(6-(4-methyl-
140 1,4-diazepan-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; N-
141 ((1-(6-(1,4-Diazepan-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)-5-chlorothiophene-2-
142 carboxamide; N-((1-(6-(1,4-Oxazepan-4-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)-5-
143 chlorothiophene-2-carboxamide; 5-Chloro-N-((1-(6-morpholinopyridin-3-yl)-1H-1,2,3-
144 triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(6-(3-oxomorpholino)pyridin-
145 3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(6-(2-
146 oxopiperidin-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-

147 5-Chloro-N-((1-(6-(2-oxopyridin-1(2H)-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-
 148 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(6-(piperazin-1-yl)pyridin-3-yl)-1H-
 149 1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(6-(4-methylpiperazin-1-
 150 yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((1-(6-
 151 (piperidin-1-yl)pyridin-3-yl)-1H-1,2,3-triazol-4-yl)methyl)thiophene-2-carboxamide; 5-
 152 Chloro-N-((3-(4-(8-(trifluoromethyl)quinolin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-
 153 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(7-(trifluoromethyl)quinolin-4-
 154 ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-
 155 ((3-(4-(6-(trifluoromethyl)quinolin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-
 156 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(quinolin-6-ylamino)phenyl)-4,5-
 157 dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(2,5-dihydro-1H-
 158 pyrrole-1-carbonyl)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide; 5-
 159 Chloro-N-((3-(4-(2,5-dihydro-1H-pyrrole-1-carbonyl)phenyl)-4,5-dihydroisoxazol-5-
 160 yl)methyl)thiophene-2-carboxamide; N-((3-(4-((2-Hydroxypropyl)carbamoyl)phenyl)-4,5-
 161 dihydroisoxazol-5-yl)methyl)-5-chlorothiophene-2-carboxamide; 5-Chloro-N-((3-(4-(3-
 162 methylpyridin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-
 163 carboxamide; 5-Chloro-N-((3-(4-(2-ethoxypyridin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-
 164 yl)methyl)thiophene-2-carboxamide; 5-Chloro-N-((3-(4-(3,5-dichloropyridin-4-
 165 ylamino)phenyl)-4,5-dihydroisoxazol-5-yl)methyl)thiophene-2-carboxamide; and 5-Chloro-
 166 N-((3-(4-(3-(trifluoromethyl)pyridin-4-ylamino)phenyl)-4,5-dihydroisoxazol-5-
 167 yl)methyl)thiophene-2-carboxamide.

1 **89.** A composition comprising a pharmaceutically acceptable excipient and
 2 a compound of claim 1 to 88.

1 **90.** A method for preventing or treating a condition in a mammal
 2 characterized by undesired thrombosis comprising the step of administering to said mammal
 3 a therapeutically effective amount of a compound of claims 1 to 89.

1 **91.** The method in accordance with claim 90, wherein the condition is
 2 selected from the group consisting of acute coronary syndrome, myocardial infarction,
 3 unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic
 4 therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome,
 5 embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep

6 venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular
7 coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic
8 disease associated with heparin-induced thrombocytopenia, thrombotic complications
9 associated with extracorporeal circulation, thrombotic complications associated with
10 instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon
11 pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic
12 devices.

1 **92.** The method for inhibiting the coagulation of a blood sample
2 comprising contacting said sample with a compound of claim **1** to **91**.