Claims

1. A process for manufacturing an aqueous composition comprising ibuprofen and paracetamol in combination, comprising the steps of:
    a) providing an aqueous solvent of pH 6.0-8.0, preferably 6.2-7.5, more preferably 6.3-7.3,
    b) dissolving in said aqueous solvent 2.8-3.2 mg ibuprofen and 9.8-10.2 mg paracetamol per ml of said composition,
    c) ensuring that the final composition comprising ibuprofen and paracetamol has a pH of 6.3-7.3.

2. Process according to claim 1, wherein said aqueous solvent has a dissolved oxygen content below 2 ppm.

3. Process according to any of the above claims 1-2, wherein the pH of the aqueous solvent in step a) is 6.3 to 6.5.

4. Process according to any of the above claims 1-3, wherein the composition comprises an anti-oxidant; preferably said anti-oxidant is cysteine or acetyl cysteine.

5. Process according to any one of claims 1 to 4, wherein the aqueous solvent comprises an isotonic agent, preferably mannitol.

6. An aqueous composition, comprising 2.8 to 3.2 mg ibuprofen and 9.8 to 10.2 mg paracetamol expressed per ml of said composition, characterized in that, the pH of said composition is 6.3 to 7.3.

7. Composition according to claim 6, comprising 3 mg ibuprofen and 10 mg paracetamol expressed per ml of said composition.

8. Composition according to any one of claims 6-7, comprising an isotonic agent, preferably said isotonic agent is mannitol.

9. Composition according to any one of claims 6-8, wherein said composition has a dissolved oxygen content below 2 ppm.

10. Composition according to any one of claims 6-9, for use as a medicament.

11. Use of a composition according to any one of claims 6-10 in the manufacture of a medicament, for the treatment of pain and/or inflammation; wherein the composition is to be administered preferably by intravenous injection or infusion.

12. A device (1) comprising a composition according to any of claims 6-11 for delivering said aqueous composition by means of injection or infusion.

13. Device (1) according to claim 12, in the form of a pre-filled syringe.

14. Device (1) according to claim 12, in the form of an infusion bag.