

SINGLE DOMAIN SERUM ALBUMIN BINDING PROTEIN

CROSS-REFERENCE

[0001] This application claims the benefit of U.S. Provisional Application No 62/339,682 filed May 20, 2016 which is incorporated by reference herein in its entirety.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on May 18, 2017, is named 47517-703_601_SEQ.txt and is 22,218 bytes in size.

INCORPORATION BY REFERENCE

[0003] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference, and as if set forth in their entireties.

BACKGROUND OF THE INVENTION

[0004] Albumin is the most abundant plasma protein, is highly soluble, very stable and has an extraordinarily long circulatory half-life. Albumin can be used in a variety of ways to increase the circulatory half-life of therapeutic molecules. The present disclosure provides a single domain albumin binding protein which can be used for extending the half-life of therapeutic molecules.

SUMMARY OF THE INVENTION

[0005] Provided herein in one embodiment is a single domain serum albumin binding protein, comprising complementarity determining regions CDR1, CDR2, and CDR3, wherein (a) the amino acid sequence of CDR1 is as set forth in GFX₁X₂X₃X₄FGMS (SEQ ID NO. 1), X₁ is threonine, arginine, lysine, serine, or proline, X₂ is phenylalanine or tyrosine, X₃ is serine, arginine or lysine, X₄ is serine, lysine, arginine, or alanine; (b) the amino acid sequence of CDR2 is as set forth in SISGSGX₅X₆TLYAX₇SX₈K (SEQ ID NO. 2), X₅ is serine, arginine, threonine, or alanine, X₆ is aspartic acid, histidine, valine, or threonine, X₇ is aspartic acid, histidine, arginine, or serine, X₈ is valine or leucine; and (c) the amino acid sequence of CDR3 is as set forth in GGSLX₉X₁₀ (SEQ ID NO. 3), X₉ is serine, arginine, threonine, or lysine, and X₁₀ is arginine, lysine, valine, proline, or asparagine, wherein X₁,

X₂, X₃, X₄, X₅, X₆, X₇, X₈, X₉, and X₁₀ are not simultaneously threonine, phenylalanine, serine, serine, serine, aspartic acid, aspartic acid, valine, serine, and arginine, respectively. In some embodiments, the single domain serum albumin binding protein comprises the following formula: f₁-r₁-f₂-r₂-f₃-r₃-f₄, wherein, r₁ is SEQ ID NO. 1; r₂ is SEQ ID NO. 2; and r₃ is SEQ ID NO. 3; and wherein f₁, f₂, f₃ and f₄ are framework residues selected so that said protein is at least eighty percent identical to the amino acid sequence set forth in SEQ ID NO. 10. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₁ comprises SEQ ID NO. 14, SEQ ID NO. 15 or SEQ ID NO. 16. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₂ comprises SEQ ID NO. 17, SEQ ID NO. 18, SEQ ID NO. 19, SEQ ID NO. 20, SEQ ID NO. 21, or SEQ ID NO. 22. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₃ comprises SEQ ID NO. 23 or SEQ ID NO. 24. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₁ comprises SEQ ID NO. 14. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₁ comprises SEQ ID NO. 15, r₂ comprises SEQ ID NO. 17, and r₃ comprises SEQ ID NO. 23. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₁ comprises SEQ ID NO. 16, and r₃ comprises SEQ ID NO. 23. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₁ comprises SEQ ID NO. 15, and r₂ comprises SEQ ID NO. 18. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₁ comprises SEQ ID NO. 14, and r₃ comprises SEQ ID NO. 23. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₁ comprises SEQ ID NO. 15, r₂ comprises SEQ ID NO. 19 and r₃ comprises SEQ ID NO. 24. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₁ comprises SEQ ID NO. 14, and r₂ comprises SEQ ID NO. 20. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₁ comprises SEQ ID NO. 15, and r₂ comprises SEQ ID NO. 21. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein r₁ comprises SEQ ID NO. 15, r₂ comprises SEQ ID NO. 22, and r₃ comprises SEQ ID NO. 24. In some embodiments, the single domain serum albumin binding protein has an amino acid sequence selected from SEQ ID NO. 4, SEQ ID NO. 5, SEQ ID NO. 6, SEQ ID NO. 7, SEQ ID NO. 8, SEQ ID NO. 9, SEQ ID NO. 25, SEQ ID NO. 26, and

SEQ ID NO. 27. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 4. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 7. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 9. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 26. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 27. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 5. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 6. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 8. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 25.

[0006] In some embodiments, the single domain serum albumin binding protein binds to serum albumin selected from human serum albumin, cynomolgus serum albumin, and mouse serum albumin. In some embodiments, the single domain serum albumin binding protein binds to human serum albumin and cynomolgus serum albumin with comparable binding affinity (Kd). In some embodiments, the single domain serum albumin binding protein binds to mouse serum albumin with a binding affinity (Kd) that is about 1.5 fold to about 20 fold weaker than the binding affinity (Kd) of said protein towards human and cynomolgus serum albumin. In some embodiments, the single domain serum albumin binding protein binds to human serum albumin with a human Kd (hKd) between about 1 nM and about 100 nM and to cynomolgus serum albumin with a cynomolgus Kd (cKd) between 1 nM and 100 nM. In some embodiments, the hKd and the cKd of the single domain serum albumin binding protein are between 1 nM and about 5 nM, or about 5 nM and about 10 nM. In some embodiments, the hKd and cKd of the single domain serum albumin binding protein are between about 1 nM and about 2 nM, about 2 nM and about 3 nM, about 3 nM and about 4 nM, about 4 nM and about 5 nM, about 5 nM and about 6 nM, about 6 nM and about 7 nM, about 7 nM and about 8 nM, about 8 nM and about 9 nM, or about 9 nM and about 10 nM. In some embodiments, the ratio between the hKd and cKd (hKd: cKd) of the single domain serum albumin binding protein ranges from about 20:1 to about 1: 2.

[0007] In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 4, and wherein the hKd and the cKd are

between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 5, and wherein the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 6, and wherein the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 7, and wherein the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 8, and wherein the hKd and the cKd are between about 5 nM and about 10 nM. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 9, and wherein the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 22, and wherein the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 23, and wherein the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises the amino acid sequence set forth as SEQ ID NO. 24, and wherein the hKd and the cKd are between about 1 nM and about 5 nM.

[0008] In some embodiments, the single domain serum albumin binding protein comprises elimination half-time of at least 12 hours, at least 20 hours, at least 25 hours, at least 30 hours, at least 35 hours, at least 40 hours, at least 45 hours, at least 50 hours, or at least 100 hours.

[0009] In another embodiment is provided a single domain serum albumin binding protein comprising CDR1, CDR2, and CDR3, comprising the sequence set forth as SEQ ID NO. 10 wherein one or more amino acid residues selected from amino acid positions 28, 29, 30, or 31 of CDR1; positions 56, 57, 62, or 64 of CDR2; and positions 103, and 104 of CDR3 are substituted, wherein amino acid position 28 is substituted with arginine, lysine, serine, or proline, amino acid position 29 is substituted with tyrosine, amino acid position 30 is substituted with arginine or lysine, amino acid position 31 is substituted with lysine, arginine, or alanine, amino acid position 56 is substituted with arginine, threonine, or alanine, amino acid position 57 is substituted with histidine, valine, or threonine, amino acid position 62 is substituted with histidine, arginine, glutamic acid, or serine, amino acid position 64 is

substituted with leucine, amino acid position 103 is substituted with arginine, threonine, or lysine, amino acid position 104 is substituted with lysine, valine, proline, or asparagine. In some embodiments, the single domain serum albumin binding comprises one or more additional substitutions in amino acid positions other than positions 28, 29, 30, 31, 56, 57, 62, 64, 103, and 104. In some embodiments, the single domain serum albumin binding protein comprises a substitution in position 29. In some embodiments, the single domain serum albumin binding protein comprises a substitution in position 31. In some embodiments, the single domain serum albumin binding protein comprises a substitution in position 56. In some embodiments, the single domain serum albumin binding protein comprises a substitution in position 62. In some embodiments, the single domain serum albumin binding protein comprises a substitution in position 64. In some embodiments, the single domain serum albumin binding protein comprises a substitution in position 104. In some embodiments, the single domain serum albumin binding protein comprises substitutions in amino acid positions 31 and 62. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein position 31 is substituted with arginine. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein position 31 is substituted with arginine and amino acid position 62 is substituted with glutamic acid. In some embodiments, the single domain serum albumin binding protein comprises substitutions in amino acid positions 31, 56, 64, and 104. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein position 31 is substituted with lysine, amino acid position 56 is substituted with alanine, amino acid position 64 is substituted with leucine, and amino acid position 104 is substituted with lysine. In some embodiments, the single domain serum albumin binding protein comprises substitutions in amino acid positions 29 and 104. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 29 is substituted with tyrosine, and amino acid position 104 is substituted with lysine. In some embodiments, the single domain serum albumin binding protein comprises substitutions in amino acid positions 31 and 56. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with lysine, and amino acid position 56 is substituted with threonine. In some embodiments, the single domain serum albumin binding protein comprises substitutions in amino acid positions 31, 56, and 62. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with lysine, amino acid position 56 is

substituted with threonine, and amino acid position 62 is substituted with glutamic acid. In some embodiments, the single domain serum albumin binding protein comprises substitutions in amino acid positions 31 and 104. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with arginine, and amino acid position 104 is substituted with lysine. In some embodiments, the single domain serum albumin binding protein comprises substitutions in amino acid positions 31, 56, and 104. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with lysine, amino acid position 56 is substituted with arginine, and amino acid position 104 is substituted with valine. In some embodiments, the single domain serum albumin binding protein comprises substitutions in amino acid positions 31, 56, 62, and 104. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with lysine, amino acid position 56 is substituted with arginine, amino acid position 62 is substituted with glutamic acid, and amino acid position 104 is substituted with valine. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with arginine, and the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with arginine, amino acid position 62 is substituted with glutamic acid, and the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with lysine, amino acid position 56 is substituted with alanine, amino acid position 64 is substituted with leucine, amino acid position 104 is substituted with lysine, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 29 is substituted with tyrosine, amino acid position 104 is substituted with lysine, and wherein the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with lysine, amino acid position 56 is substituted with threonine, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain

serum albumin comprises an amino acid sequence wherein amino acid position 31 is substituted with lysine, amino acid position 56 is substituted with threonine, and amino acid position 62 is substituted with glutamic acid, and wherein the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with arginine, amino acid position 104 is substituted with lysine, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 5 nM and about 10 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with lysine, amino acid position 56 is substituted with arginine, amino acid position 104 is substituted with valine, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 31 is substituted with lysine, amino acid position 56 is substituted with arginine, amino acid position 62 is substituted with glutamic acid, and amino acid position 104 is substituted with valine, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM.

[0010] Provided herein in another embodiment is a single domain serum albumin binding protein comprising at least one mutation in CDR1, CDR2 or CDR3, wherein CDR1 comprises the sequence as set forth is SEQ ID NO:11, CDR2 comprises the sequence as set forth in SEQ ID NO:12, CDR3 comprises the sequence as set forth in SEQ ID NO. 13, and wherein the at least one mutation is not in amino acid positions 1, 2, 7, 8, 9, or 10 of SEQ ID NO: 11, positions 1, 3, 6, 10, or 11 of SEQ ID NO: 12, or positions 1 or 2 of SEQ ID NO: 13. In some embodiments, the single domain serum albumin binding protein comprises at least one mutation in amino acid positions selected from positions 3, 4, 5, and 6 of CDR1 (SEQ ID NO: 11), amino acid positions 7, 8, 13, and 15 of CDR2 (SEQ ID NO: 12), and amino acid positions 5 and 6 of CDR3 (SEQ ID NO: 13). In some embodiments, the single domain serum albumin binding protein comprises one or more additional substitutions in amino acid positions other than 3, 4, 5, and 6 of CDR1 (SEQ ID NO: 11), amino acid positions 7, 8, 13, and 15 of CDR2 (SEQ ID NO: 12), and amino acid positions 5 and 6 of CDR3 (SEQ ID NO: 13). In some embodiments, the single domain serum albumin binding protein comprises a mutation in amino acid position 6 of CDR 1 (SEQ ID NO:11). In some embodiments, the single domain serum albumin binding protein comprises a mutation in amino acid position 6 of CDR 1 (SEQ ID NO:11), and amino acid position 13 of CDR2 (SEQ ID NO: 12). In some

embodiments, the single domain serum albumin binding protein comprises mutations in amino acid position 6 of (SEQ ID NO:11), amino acid positions 7 and 15 of CDR2 (SEQ ID NO: 12), and amino acid position 6 of CDR3 (SEQ ID NO: 13). In some embodiments, the single domain serum albumin binding protein comprises mutations in amino acid position 4 of CDR1 (SEQ ID NO:11), and amino acid position 6 of CDR3 (SEQ ID NO: 13). In some embodiments, the single domain serum albumin binding protein comprises mutations in amino acid position 6 of CDR 1 (SEQ ID NO:11), and amino acid position 7 of CDR2 (SEQ ID NO: 12). In some embodiments, the single domain serum albumin binding protein comprises mutations in amino acid position 6 of CDR 1 (SEQ ID NO:11), and amino acid positions 7 and 13 of CDR2 (SEQ ID NO: 12). In some embodiments, the single domain serum albumin binding protein comprises mutations in amino acid position 6 of CDR1 (SEQ ID NO:11), and amino acid position 6 of CDR3 (SEQ ID NO: 13). In some embodiments, the single domain serum albumin binding protein comprises mutations in amino acid position 6 of CDR1 (SEQ ID NO:11), amino acid position 7 of CDR2 (SEQ ID NO: 12), and amino acid position 6 of CDR3 (SEQ ID NO: 13). In some embodiments, the single domain serum albumin binding protein comprises mutations in amino acid position 6 of CDR1 (SEQ ID NO:11), amino acid positions 7 and 13 of CDR2 (SEQ ID NO: 12), and amino acid position 6 of CDR3 (SEQ ID NO: 13). In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 6 of CDR1 (SEQ ID NO:11) is mutated to arginine, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 6 of CDR1 (SEQ ID NO:11) is mutated to arginine, and amino acid position 13 of CDR2 (SEQ ID NO: 12) is mutated to glutamic acid, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 6 is of CDR 1 (SEQ ID NO:11) is mutated to lysine, amino acid positions 7 and 15 of CDR2 (SEQ ID NO: 12) are mutated to alanine and leucine, respectively, amino acid position 6 of CDR3 (SEQ ID NO: 13) is mutated to lysine, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 4 of CDR1 (SEQ ID NO:11) is mutated to tyrosine, and amino acid position 6 of CDR3 (SEQ ID NO: 13) is mutated to lysine, and wherein the

hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 6 of CDR 1 (SEQ ID NO:11) is mutated to lysine, amino acid position 7 of CDR2 (SEQ ID NO: 12) is mutated to threonine, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 6 of CDR 1 (SEQ ID NO:11) is mutated to lysine, amino acid positions 7 and 13 of CDR2 (SEQ ID NO: 12) are mutated to threonine and glutamic acid, respectively, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 6 of CDR1 (SEQ ID NO:11) is mutated to arginine, amino acid position 6 of CDR3 (SEQ ID NO: 12) is mutated to lysine, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 5 nM and about 10 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 6 of CDR 1 (SEQ ID NO:11) is mutated to lysine, amino acid position 7 of CDR2 (SEQ ID NO: 12) is mutated to arginine, amino acid position 6 of CDR3 (SEQ ID NO: 13) is mutated to valine, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein comprises an amino acid sequence wherein amino acid position 6 of CDR 1 (SEQ ID NO:11) is mutated to lysine, amino acid positions 7 and 13 of CDR2 (SEQ ID NO: 12) is mutated to arginine and glutamic acid, respectively, and amino acid position 6 of CDR3 (SEQ ID NO: 13) is mutated to valine, and wherein the hKd and the cKd of the single domain serum albumin binding protein are between about 1 nM and about 5 nM.

[0010a] Provided herein in another embodiment is a single domain antibody that binds to a serum albumin, wherein said single domain antibody comprises complementarity determining regions CDR1, CDR2, and CDR3, and wherein

- (a) the amino acid sequence of CDR1 is GFTFSRFGMS (SEQ ID NO. 14);
- (b) the amino acid sequence of CDR2 is SISGSGSDTLYADSVK (SEQ ID NO. 12);
- and
- (c) the amino acid sequence of CDR3 is GGSLSR (SEQ ID NO. 13).

[0010b] Provided herein in another embodiment is a polynucleotide encoding the antibody as described herein or the multispecific binding protein as described herein.

[0011] Provided herein in another embodiment, a polynucleotide encoding a single domain serum albumin binding protein according to the present disclosure. A further embodiment describes a vector comprising the polynucleotide as disclosed herein. Another embodiment describes a host cell transformed with the vector according to the present disclosure. In one embodiment is provided a pharmaceutical composition comprising (i) a single domain serum albumin binding protein according to the present disclosure, a polynucleotide according to the present disclosure, a vector according to the present disclosure or a host cell according to the present disclosure, and (ii) a pharmaceutically acceptable carrier.

[0011a] Provided herein in another embodiment, pharmaceutical composition that comprises (i) a single domain antibody as described herein, the multispecific binding protein as described herein, the polynucleotide as described herein, the vector as described herein, or the host cell as described herein, and (ii) a pharmaceutically acceptable carrier.

[0011b] Provided herein in another embodiment is a process for the production of the single domain antibody as described herein or the multispecific binding protein as described herein, said process comprising culturing a host transformed or transfected with a vector comprising a polynucleotide as described herein under conditions allowing the expression of the single domain antibody or the multispecific binding protein, and recovering and purifying the produced single domain antibody or the produced multispecific binding protein from the culture.

[0012] Described herein in another embodiment, is a process for the production of a single domain serum albumin binding protein according to the present disclosure, said process comprising culturing a host transformed or transfected with a vector comprising a nucleic acid sequence encoding a single domain serum albumin binding protein as described herein under conditions allowing the expression of the serum albumin binding protein and recovering and purifying the produced protein from the culture.

[0013] Further described is a method for the treatment or amelioration of a proliferative disease, a tumorous disease, an inflammatory disease, an immunological disorder, an autoimmune disease, an infectious disease, a viral disease, an allergic reaction, a parasitic reaction, a graft-versus-host disease or a host-versus-graft disease comprising the administration of the single domain serum albumin binding protein according to the present disclosure, to a subject in need thereof. In some embodiments, the subject is human. In some embodiments, the method further comprises administration of an agent in combination with the single domain serum albumin binding protein according to the present disclosure.

[0014] In another embodiment is described a multispecific binding protein comprising the single domain serum albumin binding protein according to the present disclosure. In another embodiment is described an antibody comprising the single domain serum albumin binding protein according to the present disclosure.

[0015] A further embodiment describes a multispecific antibody, a bispecific antibody, an sdAb, a variable heavy domain, a peptide, or a ligand, comprising the single domain serum albumin binding protein according to the present disclosure. In one embodiment is provided an antibody comprising the single domain serum albumin binding protein according to the present disclosure, wherein said antibody is a single domain antibody. In some embodiments, the single domain antibody is derived from a heavy chain variable region of IgG.

BRIEF DESCRIPTION OF THE DRAWINGS

[0017] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

[0018] **FIGURE 1** illustrates the specific binding of parental anti-HSA phage as determined by ELISA titration to an HSA antigen and a CD3 antigen.

[0019] **FIGURE 2** illustrates the cross reactivity of anti-HSA phage to human, cynomolgus monkey, and mouse serum albumin as determined by ELISA titration.

[0020] **FIGURE 3** provides the binding affinity profiles of nine clones selected for more precise K_d determinations using purified sdAbs.

[0021] **FIGURE 4** illustrates the temperature of hydrophobic exposure (T_h °C) for several anti-HSA sdAb variants.

[0022] **FIGURE 5** illustrates the propensity of several anti-HSA sdAb variants to form dimer versus monomer at low pH.

DETAILED DESCRIPTION OF THE INVENTION

[0023] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby

Certain definitions

[0024] The terminology used herein is for the purpose of describing particular cases only and is not intended to be limiting. As used herein, the singular forms “a”, “an” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise. Furthermore, to the extent that the terms “including”, “includes”, “having”, “has”, “with”, or variants thereof are used in either the detailed description and/or the claims, such terms are intended to be inclusive in a manner similar to the term “comprising.”

[0025] The term “about” or “approximately” means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, *e.g.*, the limitations of the measurement system. For example, “about” can mean within 1 or more than 1 standard deviation, per the practice in the given value. Where particular values are described in the application and claims, unless otherwise stated the term “about” should be assumed to mean an acceptable error range for the particular value.

[0026] The terms “individual,” “patient,” or “subject” are used interchangeably. None of the terms require or are limited to situation characterized by the supervision (*e.g.* constant or intermittent) of a health care worker (*e.g.* a doctor, a registered nurse, a nurse practitioner, a physician’s assistant, an orderly, or a hospice worker).

[0027] The term “Framework” or “FR” residues (or regions) refer to variable domain residues other than the CDR or hypervariable region residues as herein defined. A “human consensus framework” is a framework which represents the most commonly occurring amino acid residue in a selection of human immunoglobulin VL or VH framework sequences.

[0028] As used herein, “Variable region” or “variable domain” refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called complementarity-determining regions (CDRs) or hypervariable regions both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat et al., Sequences of Proteins of Immunological Interest, Fifth Edition, National Institute of Health, Bethesda, Md. (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity. “Variable domain residue numbering as in Kabat” or “amino acid position numbering as in Kabat,” and variations thereof, refers to the numbering system used for heavy chain variable domains or light chain variable domains of the compilation of antibodies in Kabat et al., Sequences of Proteins of Immunological

Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991). Using this numbering system, the actual linear amino acid sequence may contain fewer or additional amino acids corresponding to a shortening of, or insertion into, a FR or CDR of the variable domain. For example, a heavy chain variable domain may include a single amino acid insert (residue 52a according to Kabat) after residue 52 of H2 and inserted residues (e.g. residues 82a, 82b, and 82c, etc according to Kabat) after heavy chain FR residue 82. The Kabat numbering of residues may be determined for a given antibody by alignment at regions of homology of the sequence of the antibody with a “standard” Kabat numbered sequence. It is not intended that CDRs of the present disclosure necessarily correspond to the Kabat numbering convention.

[0029] As used herein, the term “Percent (%) amino acid sequence identity” with respect to a sequence is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared.

[0030] As used herein, “elimination half-time” is used in its ordinary sense, as is described in *Goodman and Gillman's The Pharmaceutical Basis of Therapeutics* 21-25 (Alfred Goodman Gilman, Louis S. Goodman, and Alfred Gilman, eds., 6th ed. 1980). Briefly, the term is meant to encompass a quantitative measure of the time course of drug elimination. The elimination of most drugs is exponential (i.e., follows first-order kinetics), since drug concentrations usually do not approach those required for saturation of the elimination process. The rate of an exponential process may be expressed by its rate constant, k , which expresses the fractional change per unit of time, or by its half-time, $t_{1/2}$ the time required for 50% completion of the process. The units of these two constants are time^{-1} and time, respectively. A first-order rate constant and the half-time of the reaction are simply related ($k \times t_{1/2} = 0.693$) and may be interchanged accordingly. Since first-order elimination kinetics dictates that a constant fraction of drug is lost per unit time, a plot of the log of drug concentration versus time is linear at all times following the initial distribution phase (i.e.

after drug absorption and distribution are complete). The half-time for drug elimination can be accurately determined from such a graph.

[0031] As used herein, the term “binding affinity” refers to the affinity of the proteins described in the disclosure to their binding targets, and is expressed numerically using “Kd” values. If two or more proteins are indicated to have comparable binding affinities towards their binding targets, then the Kd values for binding of the respective proteins towards their binding targets, are within ± 2 -fold of each other. If two or more proteins are indicated to have comparable binding affinities towards single binding target, then the Kd values for binding of the respective proteins towards said single binding target, are within ± 2 -fold of each other. If a protein is indicated to bind two or more targets with comparable binding affinities, then the Kd values for binding of said protein to the two or more targets are within ± 2 -fold of each other. In general, a higher Kd value corresponds to a weaker binding. In some embodiments, the “Kd” is measured by a radiolabeled antigen binding assay (RIA) or surface plasmon resonance assays using a BIAcore™-2000 or a BIAcore™-3000 (BIAcore, Inc., Piscataway, N.J.). In certain embodiments, an “on-rate” or “rate of association” or “association rate” or “kon” and an “off-rate” or “rate of dissociation” or “dissociation rate” or “koff” are also determined with the surface plasmon resonance technique using a BIAcore™-2000 or a BIAcore™-3000 (BIAcore, Inc., Piscataway, N.J.). In additional embodiments, the “Kd”, “kon”, and “koff” are measured using the Octet® Systems (Pall Life Sciences).

[0032] Described herein are single domain serum albumin binding proteins, pharmaceutical compositions as well as nucleic acids, recombinant expression vectors, and host cells for making such single domain serum albumin binding proteins. Also provided are methods of using the disclosed single domain serum albumin binding proteins in the prevention, and/or treatment of diseases, conditions and disorders. The single domain serum albumin binding proteins are capable specifically binding to serum albumin. In some embodiments, the single domain serum albumin binding proteins include additional domains, such as a CD3 binding domain, as well as binding domains for other target antigens.

Single domain Serum albumin binding protein

[0033] Contemplated herein are single domain serum albumin binding proteins. Serum albumin is produced by the liver, occurs dissolved in blood plasma and is the most abundant blood protein in mammals. Albumin is essential for maintaining the oncotic pressure needed for proper distribution of body fluids between blood vessels and body tissues; without albumin, the high pressure in the blood vessels would force more fluids out into the tissues. It also acts as a plasma carrier by non-specifically binding several hydrophobic steroid

hormones and as a transport protein for heme and fatty acids. Human serum albumin (HSA) (molecular mass ~67 kDa) is the most abundant protein in plasma, present at about 50 mg/ml (600 μ M), and has a half-life of around 20 days in humans. HSA serves to maintain plasma pH, contributes to colloidal blood pressure, functions as carrier of many metabolites and fatty acids, and serves as a major drug transport protein in plasma. In some embodiments, the single domain serum albumin binding proteins bind to HSA. In some embodiments, the single domain serum albumin binding proteins bind to serum albumin protein from cynomolgus monkeys. In some embodiments, the single domain serum albumin binding proteins bind to HSA and serum albumin protein from cynomolgus monkeys. In some embodiments, the single domain serum albumin binding proteins also bind to mouse serum albumin protein. In some embodiments, the binding affinity towards mouse serum albumin is about 1.5- fold to about 20-fold weaker than that towards human or cynomolgus serum albumin.

[0034] Noncovalent association with albumin extends the elimination half-time of short lived proteins. For example, a recombinant fusion of an albumin binding domain to a Fab fragment resulted in a decrease in *in vivo* clearance by 25- and 58-fold and a half-life extension of 26- and 37-fold when administered intravenously to mice and rabbits respectively as compared to the administration of the Fab fragment alone. In another example, when insulin is acylated with fatty acids to promote association with albumin, a protracted effect was observed when injected subcutaneously in rabbits or pigs. Together, these studies demonstrate a linkage between albumin binding and prolonged action/serum half-life.

[0035] In some embodiments, the single-domain serum albumin binding proteins described herein is a single domain antibody such as a heavy chain variable domain (VH), a variable domain (VHH) of camelid derived sdAb, peptide, ligand or small molecule entity specific for serum albumin. In some embodiments, the single-domain serum albumin binding proteins described herein is a single domain antibody such as a heavy chain variable domain (VH), a variable domain (VHH) of camelid derived sdAb, peptide, ligand or small molecule entity specific for HSA. In some embodiments, the serum albumin binding domain of a single domain serum albumin binding protein described herein is any domain that binds to serum albumin including but not limited to domains from a monoclonal antibody, a polyclonal antibody, a recombinant antibody, a human antibody, a humanized antibody. In certain embodiments, the serum albumin binding domain is a single-domain antibody. In other embodiments, the serum albumin binding domain is a peptide. In further embodiments, the serum albumin binding domain is a small molecule. It is contemplated that the single domain

serum albumin binding protein is fairly small and no more than 25 kD, no more than 20 kD, no more than 15 kD, or no more than 10 kD in some embodiments. In certain instances, the single domain serum albumin binding protein binding is 5 kD or less if it is a peptide or small molecule entity.

[0036] In some embodiments, the single domain serum albumin binding protein described herein is a half-life extension domain which provides for altered pharmacodynamics and pharmacokinetics of the single domain serum albumin binding protein itself. As above, the half-life extension domain extends the elimination half-time. The half-life extension domain also alters pharmacodynamic properties including alteration of tissue distribution, penetration, and diffusion of the single domain serum albumin binding protein. In some embodiments, the half-life extension domain provides for improved tissue (including tumor) targeting, tissue distribution, tissue penetration, diffusion within the tissue, and enhanced efficacy as compared with a protein without a half-life extension domain. In one embodiment, therapeutic methods effectively and efficiently utilize a reduced amount of the single domain serum albumin binding protein, resulting in reduced side effects, such as reduced non-tumor cell cytotoxicity.

[0037] Further, the binding affinity of the single domain serum albumin binding protein towards its binding target can be selected so as to target a specific elimination half-time in a particular single domain serum albumin binding protein. Thus, in some embodiments, the single domain serum albumin binding protein has a high binding affinity towards its binding target. In other embodiments, the single domain serum albumin binding protein has a medium binding affinity towards its binding target. In yet other embodiments, the single domain serum albumin binding protein has a low or marginal binding affinity towards its binding target. Exemplary binding affinities include K_D of 10 nM or less (high), between 10 nM and 100 nM (medium), and greater than 100 nM (low). As above, binding affinities of the single domain serum albumin binding proteins towards binding targets are determined by known methods such as Surface Plasmon Resonance (SPR).

[0038] In certain embodiments, the single domain serum albumin binding protein disclosed herein binds to HSA with a human K_d (hKd). In certain embodiments, the single domain serum albumin binding protein disclosed herein binds to cynomolgus monkey serum albumin with a cyno K_d (cKd). In certain embodiments, the single domain serum albumin binding protein disclosed herein binds to cynomolgus monkey serum albumin with a cyno K_d (cKd) and to HSA with a human K_d (hKd). In some embodiments, the hKd ranges between 1 nM and 100 nM. In some embodiments, the hKd ranges between 1 nM and 10 nM. In some

embodiments, the cKd ranges between 1 nM and 100 nM. In some embodiments, the cKd ranges between 1 nM and 10 nM. In some embodiments, the hKd and the cKd range between about 1 nM and about 5 nM or between about 5 nM and 10 nM. In some embodiments, the single domain serum albumin binding protein binds to serum albumin selected from human serum albumin, cynomolgus serum albumin, and mouse serum albumin. In some embodiments, the single domain serum albumin binding protein binds to human serum albumin, cynomolgus serum albumin, and mouse serum albumin with comparable binding affinity (Kd). In some embodiments, the single domain serum albumin binding protein binds to human serum albumin with a human Kd (hKd) between about 1 nM and about 10 nM and to cynomolgus serum albumin with a cynomolgus Kd (cKd) between 1 nM and 10 nM. In some embodiments, the single domain serum albumin binding protein binds to mouse serum albumin with a mouse Kd (mKd) between about 10 nM and about 50 nM.

[0039] In some embodiments, the hKd is about 1.5 nM, about 1.6 nM, about 1.7 nM, about 1.8 nM, about 1.9 nM, about 2 nM, about 2.1 nM, about 2.2 nM, about 2.3 nM, about 2.4 nM, about 2.5 nM, about 2.6 nM, about 2.7 nM, about 2.8 nM, about 2.9 nM, about 3 nM, 3.1 nM, about 3.2 nM, about 3.3. nM, about 3.4 nM, about 3.5 nM, about 3.6 nM, about 3.7 nM, about 3.8 nM, about 3.9 nM, about 4 nM, about 4.5 nM, about 5 nM, about 6, about 6.5 nM, about 7 nM, about 7.5 nM, about 8 nM, about 8.5 nM, about 9.0 nM, about 9.5 nM, or about 10 nM.

[0040] In some embodiments, the cKd is about 1.5 nM, about 1.6 nM, about 1.7 nM, about 1.8 nM, about 1.9 nM, about 2 nM, about 2.1 nM, about 2.2 nM, about 2.3 nM, about 2.4 nM, about 2.5 nM, about 2.6 nM, about 2.7 nM, about 2.8 nM, about 2.9 nM, about 3 nM, 3.1 nM, about 3.2 nM, about 3.3. nM, about 3.4 nM, about 3.5 nM, about 3.6 nM, about 3.7 nM, about 3.8 nM, about 3.9 nM, about 4 nM, about 4.5 nM, about 5 nM, about 6, about 6.5 nM, about 7 nM, about 7.5 nM, about 8 nM, about 8.5 nM, about 9.0 nM, about 9.5 nM, or about 10 nM.

[0041] In some embodiments, the mKd is about 10 nM, about 11 nM, about 12 nM, about 13 nM, about 14 nM, about 15 nM, about 16 nM, about 17 nM, about 18 nM, about 19 nM, about 20 nM, about 21 nM, about 22 nM, about 23 nM, about 24 nM, about 25 nM, about 26 nM, about 27. nM, about 28 nM, about 29 nM, about 30 nM, about 31 nM, about 32 nM, about 33 nM, about 34 nM, about 35 nM, about 36 nM, about 37, about 38 nM, about 39 nM, about 40 nM, about 41 nM, about 42 nM, about 43 nM, about 44 nM, about 45 nM, about 46 nM, about 47 nM, about 48 nM, or about 50 nM.

[0042] In some embodiments, the single domain serum albumin binding protein has an amino acid sequence selected from SEQ ID NO. 4, SEQ ID NO. 5, SEQ ID NO. 6, SEQ ID NO. 7, SEQ ID NO. 8, SEQ ID NO. 9, SEQ ID NO. 25, SEQ ID NO. 26, and SEQ ID NO. 27.

[0043] In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 4, and the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 4, and the hKd is about 2.3 nM and the cKd is about 2.4 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 25, and the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 25, and the hKd is about 2.1 nM and the cKd is about 2.2 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 5, and the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 5, and the hKd is about 1.9 nM and the cKd is about 1.7 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 6, and the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 6, and the hKd is about 3.2 nM and the cKd is about 3.6 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 7, and the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 7, and the hKd is about 2.7 nM and the cKd is about 2.6 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 26, and the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 26, and the hKd is about 2.1 nM and the cKd is about 2 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 8, and the hKd and the cKd are between about 5 nM and about 10 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 8, and the hKd is about 6 nM and the cKd is about 7.5 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence

set forth as SEQ ID NO. 9, and wherein the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 9, and wherein the hKd is about 2.2 nM and the cKd is about 2.3 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 27 and wherein the hKd and the cKd are between about 1 nM and about 5 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 27 and wherein the hKd is about 1.6 nM and the cKd is about 1.6 nM.

[0044] In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 4 and has a mKd of about 17 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 5 and has a mKd of about 12 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 6 and has a mKd of about 33 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 7 and has a mKd of about 14 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 9 and has a mKd of about 16 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 25 and has a mKd of about 17 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 26 and has a mKd of about 17 nM. In some embodiments, the single domain serum albumin binding protein has the amino acid sequence set forth as SEQ ID NO. 27 and has a mKd of about 16 nM.

[0045] In some embodiments, the ratio between the hKd and cKd (hKd: cKd) ranges from about 20:1 to about 1:2.

[0046] In some embodiments, the single domain serum albumin binding protein has an elimination half-time of at least 1 hour, at least 2 hours, at least 4 hours, at least 6 hours, at least 12 hours, at least 20 hours, at least 25 hours, at least 30 hours, at least 35 hours, at least 40 hours, at least 45 hours, at least 50 hours, or at least 100 hours..

CD3 Binding Domain

[0047] The specificity of the response of T cells is mediated by the recognition of antigen (displayed in context of a major histocompatibility complex, MHC) by the T cell receptor complex. As part of the T cell receptor complex, CD3 is a protein complex that includes a CD3 γ (gamma) chain, a CD3 δ (delta) chain, and two CD3 ϵ (epsilon) chains which are present

on the cell surface. CD3 associates with the α (alpha) and β (beta) chains of the T cell receptor (TCR) as well as and CD3 ζ (zeta) altogether to comprise the T cell receptor complex. Clustering of CD3 on T cells, such as by immobilized anti-CD3 antibodies leads to T cell activation similar to the engagement of the T cell receptor but independent of its clone-typical specificity.

[0048] In one aspect is described herein a multispecific protein comprising a single domain serum albumin binding protein according to the present disclosure. In some embodiments, the multispecific protein further comprises a domain which specifically binds to CD3. In some embodiments, the multispecific protein further comprises a domain which specifically binds to human CD3. In some embodiments, the multispecific protein further comprises a domain which specifically binds to CD3 γ . In some embodiments, the multispecific protein further comprises a domain which specifically binds to CD3 δ . In some embodiments, the multispecific protein further comprises a domain which specifically binds to CD3 ϵ .

[0049] In additional embodiments, the multispecific protein further comprises a domain which specifically binds to the T cell receptor (TCR). In some embodiments, the multispecific protein further comprises a domain which specifically binds the α chain of the TCR. In some embodiments, the multispecific protein further comprises a domain which specifically binds the β chain of the TCR.

[0050] In certain embodiments, the CD3 binding domain of the multispecific protein comprising a single domain serum albumin binding protein described herein exhibit not only potent CD3 binding affinities with human CD3, but show also excellent crossreactivity with the respective cynomolgus monkey CD3 proteins. In some instances, the CD3 binding domain of the multispecific proteins are cross-reactive with CD3 from cynomolgus monkey. In certain instances, human:cynomolgus K_D (hKd: cKd) ratios for CD3 binding are between 20:1 and 1:2.

[0051] In some embodiments, the CD3 binding domain of the multispecific protein comprising a single domain serum albumin binding protein described herein can be any domain that binds to CD3 including but not limited to domains from a monoclonal antibody, a polyclonal antibody, a recombinant antibody, a human antibody, a humanized antibody, or antigen binding fragments of the CD3 binding antibodies, such as single domain antibodies (sdAb), Fab, Fab', F(ab)2, and Fv fragments, fragments comprised of one or more CDRs, single-chain antibodies (e.g., single chain Fv fragments (scFv)), disulfide stabilized (dsFv) Fv fragments, heteroconjugate antibodies (e.g., bispecific antibodies), pFv fragments, heavy chain monomers or dimers, light chain monomers or dimers, and dimers consisting of one

heavy chain and one light chain. In some instances, it is beneficial for the CD3 binding domain to be derived from the same species in which the multispecific protein comprising a single domain serum albumin binding protein described herein will ultimately be used in. For example, for use in humans, it may be beneficial for the CD3 binding domain of the multispecific protein comprising a single domain serum albumin binding protein described herein to comprise human or humanized residues from the antigen binding domain of an antibody or antibody fragment.

[0052] Thus, in one aspect, the antigen-binding domain comprises a humanized or human antibody or an antibody fragment, or a murine antibody or antibody fragment. In one embodiment, the humanized or human anti-CD3 binding domain comprises one or more (e.g., all three) light chain complementary determining region 1 (LC CDR1), light chain complementary determining region 2 (LC CDR2), and light chain complementary determining region 3 (LC CDR3) of a humanized or human anti-CD3 binding domain described herein, and/or one or more (e.g., all three) heavy chain complementary determining region 1 (HC CDR1), heavy chain complementary determining region 2 (HC CDR2), and heavy chain complementary determining region 3 (HC CDR3) of a humanized or human anti-CD3 binding domain described herein, e.g., a humanized or human anti-CD3 binding domain comprising one or more, e.g., all three, LC CDRs and one or more, e.g., all three, HC CDRs.

[0053] In some embodiments, the humanized or human anti-CD3 binding domain comprises a humanized or human light chain variable region specific to CD3 where the light chain variable region specific to CD3 comprises human or non-human light chain CDRs in a human light chain framework region. In certain instances, the light chain framework region is a λ (lambda) light chain framework. In other instances, the light chain framework region is a κ (kappa) light chain framework.

[0054] In some embodiments, the humanized or human anti-CD3 binding domain comprises a humanized or human heavy chain variable region specific to CD3 where the heavy chain variable region specific to CD3 comprises human or non-human heavy chain CDRs in a human heavy chain framework region.

[0055] In certain instances, the complementary determining regions of the heavy chain and/or the light chain are derived from known anti-CD3 antibodies, such as, for example, muromonab-CD3 (OKT3), oteelixizumab (TRX4), teplizumab (MGA031), visilizumab (Nuvion), SP34, TR-66 or X35-3, VIT3, BMA030 (BW264/56), CLB-T3/3, CRIS7, YTH12.5, F111-409, CLB-T3.4.2, TR-66, WT32, SPv-T3b, 11D8, XIII-141, XIII-46, XIII-

87, 12F6, T3/RW2-8C8, T3/RW2-4B6, OKT3D, M-T301, SMC2, F101.01, UCHT-1 and WT-31.

[0056] The affinity to bind to CD3 can be determined, for example, by the ability of the multispecific protein comprising a single domain serum albumin binding protein itself or its CD3 binding domain to bind to CD3 coated on an assay plate; displayed on a microbial cell surface; in solution; etc. The binding activity of multispecific protein comprising a single domain serum albumin binding protein itself or its CD3 binding domain according to the present disclosure to CD3 can be assayed by immobilizing the ligand (e.g., CD3) or said multispecific protein itself or its CD3 binding domain, to a bead, substrate, cell, etc. Agents can be added in an appropriate buffer and the binding partners incubated for a period of time at a given temperature. After washes to remove unbound material, the bound protein can be released with, for example, SDS, buffers with a high pH, and the like and analyzed, for example, by Surface Plasmon Resonance (SPR).

Target Antigen Binding Domain

[0057] In addition to the described serum albumin binding and CD3 domains, the multispecific binding protein comprising a single domain serum albumin binding proteins described herein, in certain embodiments, also comprise a domain that binds to a target antigen. A target antigen is involved in and/or associated with a disease, disorder or condition. In particular, a target antigen associated with a proliferative disease, a tumorous disease, an inflammatory disease, an immunological disorder, an autoimmune disease, an infectious disease, a viral disease, an allergic reaction, a parasitic reaction, a graft-versus-host disease or a host-versus-graft disease. In some embodiments, the target antigen is a tumor antigen expressed on a tumor cell. Alternatively in some embodiments, the target antigen is associated with a pathogen such as a virus or bacterium.

[0058] In some embodiments, the target antigen is a cell surface molecule such as a protein, lipid or polysaccharide. In some embodiments, the target antigen is a on a tumor cell, virally infected cell, bacterially infected cell, damaged red blood cell, arterial plaque cell, or fibrotic tissue cell.

[0059] The design of the multispecific binding proteins comprising a single domain serum albumin binding protein according to the present disclosure allows the binding domain to a target antigen to be flexible in that the binding domain to a target antigen can be any type of binding domain, including but not limited to, domains from a monoclonal antibody, a polyclonal antibody, a recombinant antibody, a human antibody, a humanized antibody. In some embodiments, the binding domain to a target antigen is a single chain variable

fragments (scFv), single-domain antibody such as a heavy chain variable domain (VH), a light chain variable domain (VL) and a variable domain (VHH) of camelid derived sdAb. In other embodiments, the binding domain to a target antigen is a non-Ig binding domain, i.e., antibody mimetic, such as anticalins, affilins, affibody molecules, affimers, affitins, alphabodies, avimers, DARPins, fynomers, kunitz domain peptides, and monobodies. In further embodiments, the binding domain to a target antigen is a ligand or peptide that binds to or associates with a target antigen. In yet further embodiments, the binding domain to a target antigen is a knottin. In yet further embodiments, the binding domain to a target antigen is a small molecular entity.

Single Domain Serum Albumin Binding Protein Modifications

[0060] The single domain serum albumin binding proteins described herein encompass derivatives or analogs in which (i) an amino acid is substituted with an amino acid residue that is not one encoded by the genetic code, (ii) the mature polypeptide is fused with another compound such as polyethylene glycol, or (iii) additional amino acids are fused to the protein, such as a leader or secretory sequence or a sequence to block an immunogenic domain and/or for purification of the protein.

[0061] Typical modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

[0062] Modifications are made anywhere in single domain serum albumin binding proteins described herein, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Certain common peptide modifications that are useful for modification of single domain serum albumin binding proteins include glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation, blockage of the amino or carboxyl group in a polypeptide, or both, by a covalent modification, and ADP-ribosylation.

Polynucleotides Encoding Single Domain Serum Albumin Binding Proteins

[0063] Also provided, in some embodiments, are polynucleotide molecules encoding a single domain serum albumin binding protein described herein. In some embodiments, the polynucleotide molecules are provided as a DNA construct. In other embodiments, the polynucleotide molecules are provided as a messenger RNA transcript.

[0064] The polynucleotide molecules are constructed by known methods such as by combining the genes encoding the three binding domains either separated by peptide linkers or, in other embodiments, directly linked by a peptide bond, into a single genetic construct operably linked to a suitable promoter, and optionally a suitable transcription terminator, and expressing it in bacteria or other appropriate expression system such as, for example CHO cells.

[0065] Also provided, in some embodiments, are polynucleotide molecules encoding a multispecific binding protein comprising a single domain serum albumin binding protein according to the present disclosure. In some embodiments, the polynucleotide encoding said multispecific binding protein also includes coding sequence for a CD3 binding domain. In some embodiments, the polynucleotide encoding said multispecific binding protein also includes coding sequence for a target antigen binding domain. In some embodiments, the polynucleotide encoding said multispecific binding protein also includes coding sequences for a CD3 binding domain and a target antigen binding domain. In some embodiments, the polynucleotide molecules are provided as a DNA construct. In other embodiments, the polynucleotide molecules are provided as a messenger RNA transcript. In the embodiments where the target antigen binding domain is a small molecule, the polynucleotides contain genes encoding the serum albumin binding domain and the CD3 binding domain. In the embodiments where the half-life extension domain is a small molecule, the polynucleotides contain genes encoding the domains that bind to CD3 and the target antigen. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. The promoter is selected such that it drives the expression of the polynucleotide in the respective host cell.

[0066] In some embodiments, the polynucleotide is inserted into a vector, preferably an expression vector, which represents a further embodiment. This recombinant vector can be constructed according to known methods. Vectors of particular interest include plasmids, phagemids, phage derivatives, virii (e.g., retroviruses, adenoviruses, adeno-associated viruses, herpes viruses, lentiviruses, and the like), and cosmids.

[0067] A variety of expression vector/host systems may be utilized to contain and express the polynucleotide encoding the polypeptide of the described single domain serum albumin binding protein. Examples of expression vectors for expression in *E.coli* are pSKK (Le Gall et al., J Immunol Methods. (2004) 285(1):111-27), pcDNA5 (Invitrogen) for expression in mammalian cells, PICHIAPINK™ Yeast Expression Systems (Invitrogen), BACUVANCE™ Baculovirus Expression System (GenScript).

[0068] Thus, the single domain serum albumin binding proteins as described herein, in some embodiments, are produced by introducing a vector encoding the protein as described above into a host cell and culturing said host cell under conditions whereby the protein domains are expressed, may be isolated and, optionally, further purified.

Production of Single Domain Serum Albumin Binding Proteins

[0069] Disclosed herein, in some embodiments, is a process for the production of a single domain serum albumin binding protein. In some embodiments, the process comprises culturing a host transformed or transfected with a vector comprising a nucleic acid sequence encoding a single domain serum albumin binding protein under conditions allowing the expression of the serum albumin binding protein and recovering and purifying the produced protein from the culture.

[0070] In an additional embodiment is provided a process directed to improving one or more properties, e.g. affinity, stability, heat tolerance, cross-reactivity, etc., of the single domain serum albumin binding proteins and/or the multispecific binding proteins comprising a single domain serum albumin binding protein described herein, compared to a reference binding compound. In some embodiments, a plurality of single-substitution libraries is provided each corresponding to a different domain, or amino acid segment of the single domain serum albumin binding protein or reference binding compound such that each member of the single-substitution library encodes only a single amino acid change in its corresponding domain, or amino acid segment. (This allows all of the potential substitutions in a large protein or protein binding site to be probed with a few small libraries.) In some embodiments, the plurality of domains forms or covers a contiguous sequence of amino acids of the single domain serum albumin binding protein or a reference binding compound. Nucleotide sequences of different single-substitution libraries overlap with the nucleotide sequences of at least one other single-substitution library. In some embodiments, a plurality of single-substitution libraries are designed so that every member overlaps every member of each single-substitution library encoding an adjacent domain.

[0071] Binding compounds expressed from such single-substitution libraries are separately selected to obtain a subset of variants in each library which has properties at least as good as those of the reference binding compound and whose resultant library is reduced in size. (That is, the number of nucleic acids encoding the selected set of binding compounds is smaller than the number of nucleic acids encoding members of the original single-substitution library) Such properties include, but are not limited to, affinity to a target compound, stability with respect to various conditions such as heat, high or low pH, enzymatic degradation, cross-reactivity to other proteins and the like. The selected compounds from each single-substitution library are referred to herein interchangeably as “pre-candidate compounds,” or “pre-candidate proteins.” Nucleic acid sequences encoding the pre-candidate compounds from the separate single-substitution libraries are then shuffled in a PCR to generate a shuffled library, using PCR-based gene shuffling techniques.

[0072] An exemplary work flow of the screening process is described herein. Libraries of pre-candidate compounds are generated from single substitution libraries and selected for binding to the target protein(s), after which the pre-candidate libraries are shuffled to produce a library of nucleic acids encoding candidate compounds which, in turn, are cloned into a convenient expression vector, such as a phagemid expression system. Phage expressing candidate compounds then undergo one or more rounds of selection for improvements in desired properties, such as binding affinity to a target molecule. Target molecules may be adsorbed or otherwise attached to a surface of a well or other reaction container, or target molecules may be derivatized with a binding moiety, such as biotin, which after incubation with candidate binding compounds may be captured with a complementary moiety, such as streptavidin, bound to beads, such as magnetic beads, for washing. In exemplary selection regimens, the candidate binding compounds undergo a prolonged wash step so that only candidate compounds with very low dissociation rates from a target molecule are selected. Exemplary wash times for such embodiments are at least 8 hours; or in other embodiments, at least 24 hours; or in other embodiments, at least 48 hours; or in other embodiments, at least 72 hours. Isolated clones after selection are amplified and subjected to an additional cycle of selection or analyzed, for example by sequencing and by making comparative measurements of binding affinity, for example, by ELISA, surface plasmon resonance binding, bio-layer interferometry (e.g. Octet system, ForteBio, Menlo Park, CA) or the like. In some embodiments, the process is implemented to identify one or more a single domain serum albumin binding proteins and/or a multispecific binding protein comprising a single domain serum albumin binding protein with improved thermal stability, improved cross reactivity to a

selected set of binding targets compared to that of a reference serum albumin binding protein, such as a protein having the amino acid sequence of SEQ ID NO. 10. Single substitution libraries are prepared by varying codons in the VH region of the reference serum albumin binding protein, including both codons in framework regions and in CDRs; in another embodiment, the locations where codons are varied comprise the CDRs of the heavy chain of the reference serum albumin binding protein, or a subset of such CDRs, such as solely CDR1, solely CDR2, solely CDR3, or pairs thereof. In another embodiment, locations where codons are varied occur solely in framework regions. In some embodiments, a library comprises single codon changes solely from a reference serum albumin binding protein solely in framework regions of VH numbering in the range of from 10 to 250. In another embodiment, the locations where codons are varied comprise the CDR3s of the heavy chain of the reference serum albumin binding protein, or a subset of such CDR3s. In another embodiment, the number of locations where codons of VH encoding regions are varied are in the range of from 10 to 250, such that up to 100 locations are in framework region. After preparation of the single substitution library, as outlined above, the following steps are carried out: (a) expressing separately each member of each single substitution library as a pre-candidate protein; (b) selecting members of each single substitution library which encode pre-candidate proteins which bind to a binding partner that may or may not differ from the original binding target [e.g. a desired cross-reaction target(s)]; (c) shuffling members of the selected libraries in a PCR to produce a combinatorial shuffled library; (d) expressing members of the shuffled library as candidate serum albumin binding proteins; and (e) selecting members of the shuffled library one or more times for candidate serum albumin binding proteins which bind the original binding partner and potentially (f) further selecting the candidate proteins for binding to the desired cross-reactive target(s) thereby providing a nucleic acid encoded serum albumin binding protein with increased cross reactivity for the one or more substances with respect to the reference serum albumin binding protein without loss of affinity for the original ligand. In additional embodiments, the method may be implemented for obtaining a serum albumin binding protein with decreased reactivity to a selected cross-reactive substance(s) or compound(s) or epitope(s) by substituting step (f) with the following step: depleting candidate binding compounds one or more times from the subset of candidate serum albumin binding protein which bind to the undesired cross-reactive compound.

[0073] Recent studies have reported that during manufacturing, storage and *in vivo* use, therapeutic antibodies are at risk for degradation via a number of pathways. Amongst the most frequently occurring chemical degradation reactions in proteins are deamidation of

asparagine (N) and isomerization of aspartic acid (D) residues. In particular, it has been hypothesized that if N and D residues are involved in antigen recognition, their chemical alteration can lead to severe loss of potency. Asparagine and aspartic acid residues are known to share a degradation pathway that proceeds via the formation of a cyclic succinimide intermediate. The formation of succinimide intermediates and their hydrolysis products (aspartate and isoaspartate), at the aspartic acid sites of an antibody represents a stability problem. When isomerization occurs, the chemical structure of the antibody alters, which could lead to poor stability, manifested for example by aggregation, shorter shelf-life. Accordingly, in some embodiments of the present disclosure are provided single domain serum albumin binding proteins wherein one or more aspartic acid residue is mutated, thereby reducing the isomerization potential of the single domain serum albumin binding protein. In some embodiments, the aspartic acid residue is in CDR2 of the single domain serum albumin binding protein and the aspartic acid residue is mutated to glutamic acid. In certain embodiments, the aspartic acid residue in position 62 of the protein defined by SEQ ID NO. 10 is mutated to glutamic acid (D62E). In some embodiments, the serum albumin binding affinity of the single domain serum albumin proteins containing the D62E mutation is not affected by the mutation. In some embodiments, single domain serum albumin binding proteins with and without the D62E mutation have comparable binding affinity towards serum albumin.

Pharmaceutical Compositions

[0074] Also provided, in some embodiments, are pharmaceutical compositions comprising a single domain serum albumin binding protein described herein, a vector comprising the polynucleotide encoding the polypeptide of the single domain serum albumin binding proteins or a host cell transformed by this vector and at least one pharmaceutically acceptable carrier. The term "pharmaceutically acceptable carrier" includes, but is not limited to, any carrier that does not interfere with the effectiveness of the biological activity of the ingredients and that is not toxic to the patient to whom it is administered. Examples of suitable pharmaceutical carriers are well known in the art and include phosphate buffered saline solutions, water, emulsions, such as oil/water emulsions, various types of wetting agents, sterile solutions etc. Such carriers can be formulated by conventional methods and can be administered to the subject at a suitable dose. Preferably, the compositions are sterile. These compositions may also contain adjuvants such as preservative, emulsifying agents and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents.

[0075] In some embodiments of the pharmaceutical compositions, the single domain serum albumin binding protein described herein is encapsulated in nanoparticles. In some embodiments, the nanoparticles are fullerenes, liquid crystals, liposome, quantum dots, superparamagnetic nanoparticles, dendrimers, or nanorods. In other embodiments of the pharmaceutical compositions, the single domain serum albumin binding protein is attached to liposomes. In some instances, the single domain serum albumin binding protein is conjugated to the surface of liposomes. In some instances, the single domain serum albumin binding protein is encapsulated within the shell of a liposome. In some instances, the liposome is a cationic liposome.

[0076] The single domain serum albumin binding proteins described herein are contemplated for use as a medicament. Administration is effected by different ways, e.g. by intravenous, intraperitoneal, subcutaneous, intramuscular, topical or intradermal administration. In some embodiments, the route of administration depends on the kind of therapy and the kind of compound contained in the pharmaceutical composition. The dosage regimen will be determined by the attending physician and other clinical factors. Dosages for any one patient depends on many factors, including the patient's size, body surface area, age, sex, the particular compound to be administered, time and route of administration, the kind of therapy, general health and other drugs being administered concurrently. An "effective dose" refers to amounts of the active ingredient that are sufficient to affect the course and the severity of the disease, leading to the reduction or remission of such pathology and may be determined using known methods.

Methods of Treatment

[0077] Also provided herein, in some embodiments, are methods and uses for stimulating the immune system of an individual in need thereof comprising administration of a single domain serum albumin binding protein or a multispecific binding protein comprising the a single domain serum albumin binding protein described herein. In some instances, the administration of a single domain serum albumin binding protein described herein induces and/or sustains cytotoxicity towards a cell expressing a target antigen. In some instances, the cell expressing a target antigen is a cancer or tumor cell, a virally infected cell, a bacterially infected cell, an autoreactive T or B cell, damaged red blood cells, arterial plaques, or fibrotic tissue.

[0078] Also provided herein are methods and uses for a treatment of a disease, disorder or condition associated with a target antigen comprising administering to an individual in need thereof a single domain serum albumin binding protein or a multispecific binding protein

comprising the a single domain serum albumin binding protein described herein. Diseases, disorders or conditions associated with a target antigen include, but are not limited to, viral infection, bacterial infection, auto-immune disease, transplant rejection, atherosclerosis, or fibrosis. In other embodiments, the disease, disorder or condition associated with a target antigen is a proliferative disease, a tumorous disease, an inflammatory disease, an immunological disorder, an autoimmune disease, an infectious disease, a viral disease, an allergic reaction, a parasitic reaction, a graft-versus-host disease or a host-versus-graft disease. In one embodiment, the disease, disorder or condition associated with a target antigen is cancer. In one instance, the cancer is a hematological cancer. In another instance, the cancer is a solid tumor cancer.

[0079] As used herein, in some embodiments, “treatment” or “treating” or “treated” refers to therapeutic treatment wherein the object is to slow (lessen) an undesired physiological condition, disorder or disease, or to obtain beneficial or desired clinical results. For the purposes described herein, beneficial or desired clinical results include, but are not limited to, alleviation of symptoms; diminishment of the extent of the condition, disorder or disease; stabilization (*i.e.*, not worsening) of the state of the condition, disorder or disease; delay in onset or slowing of the progression of the condition, disorder or disease; amelioration of the condition, disorder or disease state; and remission (whether partial or total), whether detectable or undetectable, or enhancement or improvement of the condition, disorder or disease. Treatment includes eliciting a clinically significant response without excessive levels of side effects. Treatment also includes prolonging survival as compared to expected survival if not receiving treatment. In other embodiments, “treatment” or “treating” or “treated” refers to prophylactic measures, wherein the object is to delay onset of or reduce severity of an undesired physiological condition, disorder or disease, such as, for example is a person who is predisposed to a disease (e.g., an individual who carries a genetic marker for a disease such as breast cancer).

[0080] In some embodiments of the methods described herein, the single domain serum albumin binding proteins or a multispecific binding protein comprising the a single domain serum albumin binding protein described herein are administered in combination with an agent for treatment of the particular disease, disorder or condition. Agents include but are not limited to, therapies involving antibodies, small molecules (e.g., chemotherapeutics), hormones (steroidal, peptide, and the like), radiotherapies (γ -rays, X-rays, and/or the directed delivery of radioisotopes, microwaves, UV radiation and the like), gene therapies (e.g., antisense, retroviral therapy and the like) and other immunotherapies. In some embodiments,

the single domain serum albumin binding proteins or a multispecific binding protein comprising the a single domain serum albumin binding protein described herein are administered in combination with anti-diarrheal agents, anti-emetic agents, analgesics, opioids and/or non-steroidal anti-inflammatory agents. In some embodiments, the single domain serum albumin binding proteins or a multispecific binding protein comprising a single domain serum albumin binding protein as described herein are administered before, during, or after surgery.

EXAMPLES

Example 1: Generation of anti-HSA single domain antibody variants with equivalent binding properties to a parental anti-HSA single domain antibody

Characterization of Parental anti-HSA Phage

[0081] Specific binding of the parental anti-HSA phage to an HSA antigen was determined, using CD3 as a negative control (**FIG. 1**) and cross reactivity of the anti-HSA phage to human, cynomolgus monkey, and mouse serum albumin was determined (**FIG. 2**).

Single Substitution HSA sdAb Phage Libraries

[0082] A single substitution library was provided for each of the three CDR domains. Single substitution libraries were bound to HSA and then washed in buffer containing various levels of HSA. Phages bound at 0 and 24 hours were rescued and counted. Phages selected using a 24 hour wash with 2.5 mg/ml HSA in the buffer were used to create two independent combinatorial phage libraries.

Combinatorial Anti-HSA Libraries

[0083] MSA was used as the selection target in the first Round. Wells were washed for 24 hours after combinatorial phage binding from two independent libraries. HSA was used as the selection target in the second round. Wells were washed in 1mg/ml HSA for 24 hours after binding of both libraries. Inserts PCR'd from the second round of selection were subcloned into the ME10 His6 expression vector (6XHis sequence disclosed as SEQ ID NO: 38). 96 clones were picked, DNA was purified, sequenced, and transfected into Expi293 cells.

Binding affinity measurement

[0084] Supernatants were used to estimate K_d to HSA and CSA using the Octet platform. Nine clones were selected for further characterization (**FIG. 3**), based on binding affinities compared to the parental sdAb as well as robust production, aggregation and stability profiles.

Example 2: Pharmacokinetics of a trispecific antibody comprising the anti-HSA single domain antibody

[0085] The anti-HSA single domain antibody of Example 1 is used to prepare a trispecific antibody which is evaluated for half-time elimination in animal studies.

[0086] The trispecific antibody is administered to cynomolgus monkeys as a 0.5 mg/kg bolus injection intramuscularly. Another cynomolgus monkey group receives a comparable protein in size with binding domains to CD3 and CD20, but lacking HSA binding. A third and fourth group receive an antibody with CD3 and HSA binding domains and a protein with CD20 and HSA binding domains respectively, and both comparable in size to the trispecific antibody. Each test group consists of 5 monkeys. Serum samples are taken at indicated time points, serially diluted, and the concentration of the proteins is determined using a binding ELISA to CD3 and/or CD20.

[0087] Pharmacokinetic analysis is performed using the test article plasma concentrations. Group mean plasma data for each test article conforms to a multi-exponential profile when plotted against the time post-dosing. The data are fit by a standard two-compartment model with bolus input and first-order rate constants for distribution and elimination phases. The general equation for the best fit of the data for i.v. administration is: $c(t) = Ae^{-\alpha t} + Be^{-\beta t}$, where $c(t)$ is the plasma concentration at time t , A and B are intercepts on the Y-axis, and α and β are the apparent first-order rate constants for the distribution and elimination phases, respectively. The α -phase is the initial phase of the clearance and reflects distribution of the protein into all extracellular fluid of the animal, whereas the second or β -phase portion of the decay curve represents true plasma clearance. Methods for fitting such equations are well known in the art. For example, $A = D/V(\alpha - k_{21})/(\alpha - \beta)$, $B = D/V(\beta - k_{21})/(\alpha - \beta)$, and α and β (for $\alpha > \beta$) are roots of the quadratic equation: $r^2 + (k_{12} + k_{21} + k_{10})r + k_{21}k_{10} = 0$ using estimated parameters of V =volume of distribution, k_{10} =elimination rate, k_{12} =transfer rate from compartment 1 to compartment 2 and k_{21} =transfer rate from compartment 2 to compartment 1, and D =the administered dose.

[0088] Data analysis: Graphs of concentration versus time profiles are made using KaleidaGraph (KaleidaGraph™ V. 3.09 Copyright 1986-1997. Synergy Software. Reading, Pa.). Values reported as less than reportable (LTR) are not included in the PK analysis and are not represented graphically. Pharmacokinetic parameters are determined by compartmental analysis using WinNonlin software (WinNonlin® Professional V. 3.1 WinNonlin™ Copyright 1998-1999. Pharsight Corporation. Mountain View, Calif.). Pharmacokinetic parameters are computed as described in Ritschel W A and Kearns G L,

1999, IN: *Handbook Of Basic Pharmacokinetics Including Clinical Applications*, 5th edition, American Pharmaceutical Assoc., Washington, D.C.

[0089] It is expected that the trispecific antibody comprising the anti-HSA single domain antibody of Example 1 has improved pharmacokinetic parameters such as an increase in elimination half-time as compared to proteins lacking an HSA binding domain.

Example 3: Thermal stability of anti-HSA single domain antibody variants

[0090] The temperature of hydrophobic exposure (T_h) of a protein corresponds to the derivative of the inflection point of peak dye fluorescence and is known to correlate with melting temperature (T_m), which is a measure of protein stability. The goal of this study was to assess the T_h for several anti-HAS single domain antibody variants.

Protein Production

[0091] Sequences of anti-huALB single domain antibodies were cloned into pcDNA3.4 (Invitrogen) preceded by a leader sequence and followed by a 6x Histidine tag (SEQ ID NO: 38). Expi293F cells (Life Technologies A14527) were maintained in suspension in Optimum Growth Flasks (Thomson) between 0.2 to 8 x 1e6 cells/ml in Expi 293 media. Purified plasmid DNA was transfected into Expi293F cells in accordance with Expi293 Expression System Kit (Life Technologies, A14635) protocols, and maintained for 4-6 days post transfection. Conditioned media was partially purified by affinity and desalting chromatography. Anti-huCD3e scFv proteins were concentrated with Amicon Ultra centrifugal filtration units (EMD Millipore), applied to Superdex 200 size exclusion media (GE Healthcare) and resolved in a neutral buffer containing excipients. Fraction pooling and final purity were assessed by SDS-PAGE and analytical size exclusion chromatography (SEC). The absorbance of purified protein solutions were determined at 280nm using a SpectraMax M2 (Molecular Devices) and UV-transparent 96-well plates (Corning 3635) and their concentrations were calculated from molar extinction coefficients.

Differential Scanning Fluorimetry

[0092] Purified anti-HSA single domain antibody proteins were diluted from 0.2 to 0.25 mg/ml together with 5x SYPRO orange dye (Life Technologies S6651) in 0.15% DMSO final concentration in a neutral buffer containing excipients into MicroAmp EnduraPlate optical microplates and adhesive film (Applied Biosystems 4483485 and 4311971). A plate containing diluted protein and dye mixtures was loaded into an ABI 7500 Fast real-time PCR instrument (Applied Biosystems) and subjected to a multi-step thermal gradient from 25 °C to 95 °C. The thermal gradient comprised of a two minute hold at each one degree Celsius with

excitation at 500 nm and emission collected with a ROX filter. T_h in degrees celsius is presented for purified anti-HSA single domain antibody proteins in **Fig. 4**.

Example 4: Relative propensity of dimerization of anti-HSA single domain antibody variants when exposed to low pH

[0093] Anti-HSA single domain antibody proteins were expressed in Expi293-F cells as above. Conditioned medium for each variant was applied to protein A agarose (GE Healthcare, 17519901) packed in a column, washed extensively with TRIS buffered saline, eluted with 0.05% (vol/vol) acetic acid at pH 3, held at room temperature for up to ten minutes prior to partial neutralization to pH 5, and subsequently desalted into a neutral buffer containing excipients using Sephadex G25 columns (GE Healthcare 17058401).

[0094] Concentrations of purified anti-HSA single domain antibody variants were determined by absorbance at 280 nm as described in Example 3. Purified proteins were evaluated by SDS-PAGE and analytical SEC using a Yarra 2000 SEC column (Phenomenex 00H-4512-E0) resolved in phosphate buffer containing solvent on a 1200 LC with Chemstation software (Agilent). Peaks corresponding to dimer and monomer were manually integrated and values are presented in **Fig. 5**.

<u>SEQ ID NO:</u>	<u>Description</u>	<u>AA Sequence</u>
1	CDR1 with variant positions	GFX ₁ X ₂ X ₃ X ₄ FGMS
2	CDR2 with variant positions	SISGSGX ₅ X ₆ TLYAX ₇ SX ₈ K
3	CDR3 with variant positions	GGSLX ₉ X ₁₀
4	Anti-HSA sdAb clone 6C	EVQLVESGGGLVQPGNSLRRLSCAASGFTFSRFGMSWVRQAPGKGL EWVSSISGSGSDTLYADSVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSRSSQGTLVTVSS
5	Anti-HSA sdAb clone 7A	EVQLVESGGGLVQPGNSLRRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGADTLYADSLKGRFTISRDNAAKTTLYLQMNSLRPEDT AVYYCTIGGSLSKSSQGTLVTVSS
6	Anti-HSA sdAb clone 7G	EVQLVESGGGLVQPGNSLRRLSCAASGFTYSSFGMSWVRQAPGKGL EWVSSISGSGSDTLYADSVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSKSSQGTLVTVSS
7	Anti-HSA sdAb clone 8H	EVQLVESGGGLVQPGNSLRRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGTDTLYADSVKGRFTISRDNAAKTTLYLQMNSLRPEDT AVYYCTIGGSLSRSSQGTLVTVSS
8	Anti-HSA sdAb clone 9A	EVQLVESGGGLVQPGNSLRRLSCAASGFTFSRFGMSWVRQAPGKGL EWVSSISGSGSDTLYADSVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSKSSQGTLVTVSS
9	Anti-HSA sdAb clone 10G	EVQLVESGGGLVQPGNSLRRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYADSVKGRFTISRDNAAKTTLYLQMNSLRPEDT AVYYCTIGGSLSVSSQGTLVTVSS
10	wt anti-HSA	EVQLVESGGGLVQPGNSLRRLSCAASGFTFSFGMSWVRQAPGKGLE WVSSISGSGSDTLYADSVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSRSSQGTLVTVSS
11	wt anti-HSA CDR1	GFTFSFGMS
12	wt anti-HSA CDR2	SISGSGSDTLYADSVK
13	wt anti-HSACDR3	GGSLSR
14	CDR1 variant 1	GFTFSRFGMS
15	CDR1 variant 2	GFTFSKFGMS
16	CDR1 variant 3	GFTYSSFGMS
17	CDR2 variant 1	SISGSGADTLYADSLK
18	CDR2 variant 2	SISGSGTDTLYADSVK
19	CDR2 variant 3	SISGSGRDTLYADSVK
20	CDR2 variant 4	SISGSGSDTLYAESVK
21	CDR2 variant 5	SISGSGTDTLYAESVK
22	CDR2 variant 6	SISGSGRDTLYAESVK

23	CDR3 variant 1	GGSLSK
24	CDR3 variant 2	GGSLSV
25	Anti-HSA sdAb clone 6CE	EVQLVESGGGLVQPGNSLRRLSCAASGFTFSRFGMSWVRQAPGKGL EWVSSISGSGSDTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTA VYYCTIGGSLSRSSQGTLVTVSS
26	Anti-HSA sdAb clone 8HE	EVQLVESGGGLVQPGNSLRRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGTDTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTA VYYCTIGGSLSRSSQGTLVTVSS
27	Anti-HSA sdAb clone 10GE	EVQLVESGGGLVQPGNSLRRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTA VYYCTIGGSLSVSSQGTLVTVSS
28	Exemplary linker sequence	(GS)n
29	Exemplary linker sequence	(GGS)n
30	Exemplary linker sequence	(GGGS)n
31	Exemplary linker sequence	(GGSG)n
32	Exemplary linker sequence	(GGSGG)n
33	Exemplary linker sequence	(GGGGS)n
34	Exemplary linker sequence	(GGGGG)n
35	Exemplary linker sequence	(GGG)n
36	Exemplary linker sequence	(GGGGS)3
37	Exemplary linker sequence	(GGGGS)4
38	6X Histidine	HHHHHH

CLAIMS

WHAT IS CLAIMED IS:

1. A single domain antibody that binds to a serum albumin, wherein said single domain antibody comprises complementarity determining regions CDR1, CDR2, and CDR3, and wherein

(a) the amino acid sequence of CDR1 is GFTFSRFGMS (SEQ ID NO. 14);

(b) the amino acid sequence of CDR2 is SISGSGSDTLYADSVK (SEQ ID NO. 12);

and

(c) the amino acid sequence of CDR3 is GGSLSR (SEQ ID NO. 13).

2. The single domain antibody according to claim 1, that comprises the following formula:

$$f1-r1-f2-r2-f3-r3-f4$$

wherein, r1 is SEQ ID NO. 14; r2 is SEQ ID NO. 12; and r3 is SEQ ID NO. 13; and wherein f₁, f₂, f₃ and f₄ are framework residues selected so that said single domain antibody is at least eighty percent identical to the amino acid sequence set forth in SEQ ID NO. 10.

3. The single domain antibody according to claim 1, that has an amino acid sequence set forth as SEQ ID NO. 4.

4. The single domain antibody according to any one of claims 1-3, that binds to a serum albumin selected from the group consisting of human serum albumin, cynomolgus serum albumin, and mouse serum albumin.

5. The single domain antibody according to any one of claims 1-4, that binds to human serum albumin and cynomolgus serum albumin with comparable binding affinity (K_d).

6. The single domain antibody according to any one of claims 1-5, that binds to mouse serum albumin with a binding affinity (K_d) of from 1.5 fold to 20 fold weaker than the binding affinity (K_d) of said single domain antibody towards human serum albumin and cynomolgus serum albumin.

7. The single domain antibody according to any one of claims 1-6, that binds to human serum albumin with a human K_d (hK_d) between 1 nM and 100 nM and to cynomolgus serum albumin with a cynomolgus K_d (cK_d) between 1 nM and 100 nM.

8. The single domain antibody according to claim 7, wherein the hK_d and cK_d are between 1 nM and 5 nM, or 5 nM and 10 nM.

9. The single domain antibody according to claim 7, wherein the hKd and the cKd are between 1 nM and 2 nM, 2 nM and 3 nM, 3 nM and 4 nM, 4 nM and 5 nM, 5 nM and 6 nM, 6 nM and 7 nM, 7 nM and 8 nM, 8 nM and 9 nM, or 9 nM and 10 nM.
10. The single domain antibody according to any one of claims 4 and 6-9, wherein a ratio between the hKd and the cKd (hKd:cKd) ranges from 20:1 to 1:2.
11. The single domain antibody according to any one of claims 1-10, that comprises the amino acid sequence set forth as SEQ ID NO. 4, wherein the hKd and the cKd are between 1 nM and 5 nM.
12. The single domain antibody according to any one of claims 1-11, that has an elimination half-time of at least 12 hours, at least 20 hours, at least 25 hours, at least 30 hours, at least 35 hours, at least 40 hours, at least 45 hours, at least 50 hours, or at least 100 hours.
13. The single domain antibody according to any one of claims 1-12, that is derived from a heavy chain variable region of IgG.
14. A multispecific binding protein that comprises (i) the single domain antibody according to claim 13, and (ii) a CD3-binding domain.
15. A polynucleotide encoding the antibody according to any one of claims 1-13 or the multispecific binding protein of claim 14.
16. A vector that comprises the polynucleotide according to claim 15.
17. A host cell transformed with the vector according to claim 16.
18. A pharmaceutical composition that comprises (i) a single domain antibody according to any one of claims 1-13, the multispecific binding protein according to claim 14, the polynucleotide according to claim 15, the vector according to claim 16, or the host cell according to claim 17, and (ii) a pharmaceutically acceptable carrier.
19. A process for the production of the single domain antibody according to any one of claims 1-13 or the multispecific binding protein according to claim 14, said process comprising culturing a host transformed or transfected with a vector comprising a polynucleotide according to claim 15 under conditions allowing the expression of the single domain antibody or the multispecific binding protein, and recovering and purifying the produced single domain antibody or the produced multispecific binding protein from the culture.
20. The multispecific binding protein according to claim 14 for use in therapy.
21. Use of the multispecific binding protein according to claim 14 in the manufacture of a medicament for the treatment or amelioration of a proliferative disease, a

tumorous disease, an inflammatory disease, an immunological disorder, an autoimmune disease, an infectious disease, a viral disease, an allergic reaction, a parasitic reaction, a graft-versus-host disease or a host-versus-graft disease, wherein the medicament is to be administered to a subject in need thereof.

22. The use of claim 21, wherein the subject is human.

23. The use of claim 21 or 22, wherein an agent is to be administered in combination with the multispecific binding protein.