

-1-

5 **USE OF SULFAMIC ACID DERIVATIVES, ACYL SULFONAMIDES OR SULFONYL
CARBAMATES FOR THE MANUFACTURE OF A MEDICAMENT FOR LOWERING
LIPOPROTEIN LEVELS**

10 **BACKGROUND OF INVENTION**

10

 The compounds of the instant invention are fully described in copending United States Application Serial Number 08/223,932 filed April 13, 1994. The use of the compounds taught is hypercholesterolemia and atherosclerosis. This application is hereby incorporated by reference.

15

 The compounds of the copending application show increased chemical stability over those of United States Patent Number 5,245,068.

20

 The present invention relates generally to lipoprotein(a), Lp(a), and more particularly to methods and agents to lower its plasma concentrations to achieve therapeutic benefit.

25

 The macromolecule known as lipoprotein(a), or Lp(a), is a complex of low-density lipoproteins (LDL), and a hydrophilic glycoprotein that has been given the name apolipoprotein(a), or apo(a). The principal protein of LDL is apo B-100, and apo(a) is attached to the apo B moiety of LDL by a disulfide bond. LDL is the major transporter of cholesterol in human plasma. The physiological function of Lp(a) is unknown.

30

 Apo(a) is not similar in structure to other apolipoproteins but exhibits similarity to another plasma protein called plasminogen. The structure of plasminogen includes five tandemly repeated homologous domains called kringles (Kringles I-V), which are pretzel-like structures stabilized by three internal disulfide bridges followed by a protease domain. Kringle structures have been identified in various

35

-2-

other proteins such as prothrombin, tissue-type plasminogen activator (t-PA), urokinase and coagulation Factor XII (Utermann, Science, 1989;246:904-910).

5 Apo(a) lacks kringles similar to I to III of plasminogen but has multiple copies of the kringle domain similar to the fourth one of plasminogen, and a single copy of a kringle domain similar to the fifth one of plasminogen (kringle-5). Apo(a) also contains a protease domain.

10 Lp(a) was first identified by Berg in 1963 (Berg, Acta Pathol. Microbiol. Scand., 1963;59:369) as an antigenic activity associated with the LDL fraction in the plasma of some individuals. Plasma Lp(a) levels vary in different individuals from less than 2 mg/dL to
15 greater than 200 mg/dL. Increased plasma Lp(a) levels are considered to be a risk factor for atherosclerosis, either alone or in conjunction with elevated LDL levels (Kostner, et al., Circulation, 1989;80(5):1313-1319 citing previous investigators). The plasma
20 concentration of Lp(a) and the size of apo(a) are genetically determined (Gavish, et al., J. Clin. Invest., 1989;84:2021-2027).

The discovery of the homology of apo(a) to plasminogen has prompted further investigation as to
25 the role played by Lp(a). Hajjar, et al., Nature, 1989;339:303-305 considered the similarity between the apo(a) component of Lp(a) and plasminogen and investigated the effect that Lp(a) might have on the interaction between plasminogen and the endothelial
30 cell, and found that Lp(a) competed for plasminogen binding sites and appeared to be capable of inhibiting the activation of plasminogen on the surface of endothelial cells by t-PA. This suggests that elevated levels of Lp(a) might impair and inhibit cell surface
35 fibrinolysis, thus interfering with the fibrinolytic system. In Kostner, et al. (vide supra), HMG-CoA

-3-

reductase inhibitors such as simvastatin and lovastatin, as well as other known cholesterol-lowering agents were administered to a test group of patients, and plasma samples thereafter were taken and examined.

5 Most of the tested agents failed to lower Lp(a) levels, and, in fact, in some instances, Lp(a) levels appeared to rise, possibly due to stimulation of Lp(a) production. The authors identified only two agents, namely neomycin and niacin, that decreased both LDL and

10 Lp(a) levels. These agents only lower Lp(a) to a limited extent and because of this, as well as toxic side effects, do not appear to present a viable therapeutic avenue.

A need, therefore, exists for an effective method and associated agents for decreasing plasma Lp(a) levels.

15

The role of lipoprotein(a) has been studied in patients suffering from ischemic cerebrovascular disease and it has been determined they have

20 significantly higher levels of lipoprotein(a), lipids carried by intermediate-density proteins, low-density lipoprotein cholesterol, and lower levels of high-density lipoproteins than control subjects. These then are major risk factors for ischemic cerebrovascular

25 disease (Pedro-Botck, Stroke, 1992;23(11):1556-1562).

Lipoprotein(a) is a genetic, independent, and critical risk factor for ischemic stroke, especially in young adults (Nagayama, Stroke, 1994;25(1):74-78).

High serum lipoprotein(a) levels are an

30 independent risk factor in the development of cerebral infarction (Shintani, Stroke, 1993;24(7):965-969.

Elevated plasma levels of Lp(a) lipoprotein have been linked to the development of premature atherosclerosis in coronary circulation (Valentine, Arch. Intern. Med., 1994;154:801-806).

35

-4-


The prevention of restenosis after percutaneous transluminal coronary angioplasty by reducing lipoprotein(a) levels with low-density lipoprotein apheresis is reported in Daida, Am. J. Card., 1994;73(15):1037-1040.

Serum Lp(a) is an independent factor associated with stenosis of saphenous vein grafts (Hoff, Circulation, 1988;77(6):1238-1243.

SUMMARY OF THE INVENTION

The present invention is directed to new uses of compounds of Formula I below.

The compounds are those of formula



or a pharmaceutically acceptable salt thereof wherein:
X and Y are selected from oxygen, sulfur and $(\text{CR}'\text{R}'')_n$,
wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, alkyl, alkoxy, halogen, hydroxy, acyloxy, cycloalkyl, phenyl optionally substituted or R' and R'' together form a spirocycloalkyl or a carbonyl;
with the proviso at least one of X and Y is $-(\text{CR}'\text{R}'')_n-$ and with the further proviso when X and Y are both $(\text{CR}'\text{R}'')_n$ and R' and R'' are hydrogen and n is one, R₁ and R₂ are aryl;
R is hydrogen, a straight or branched alkyl of from 1 to 8 carbon atoms or benzyl;
R₁ and R₂ are each independently selected from

-5-

(a) phenyl or phenoxy each of which is unsubstituted or is substituted with 1 to 5 substituents selected from

- phenyl,
5 an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,
an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched;
phenoxy,
10 hydroxy,
fluorine,
chlorine,
bromine,
nitro,
15 trifluoromethyl,
-COOH,
-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,
20 $-(CH_2)_pNR_3R_4$ wherein p is zero or one, and each of R_3 and R_4 is selected from hydrogen or a straight or branched alkyl group having 1 to 4 carbon atoms;

(b) 1- or 2-naphthyl unsubstituted or substituted with from 1 to 3 substituents selected from

- 25 phenyl,
an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,
an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched;
30 hydroxy,
phenoxy,
fluorine,
chlorine,
bromine,
35 nitro,
trifluoromethyl,

-6-

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

-(CH₂)_pNR₃R₄ wherein p, R₃ and R₄ have the meanings defined above;


(c) arylalkyl;

(d) a straight or branched alkyl chain having from 1 to 20 carbon atoms and which is saturated or contains from 1 to 3 double bonds; or

(e) adamantyl or a cycloalkyl group wherein the cycloalkyl moiety has from 3 to 6 carbon atoms; with the provisos:

(i) where X is (CH₂)_n, Y is oxygen, and R₁ is a substituted phenyl, then R₂ is a substituted phenyl;(ii) where Y is oxygen, X is (CH₂)_n, R₂ is phenyl or naphthyl, then R₁ is not a straight or branched alkyl chain; and

(iii) the following compounds are excluded:

X	Y	R	R ₁	R ₂
CH ₂	O	H	(CH ₂) ₂ CH ₃	Ph
CH ₂	O	H	CH ₃	Ph
CH ₂	O	H		i-Pr

Preferred compounds of the instant invention are those of Formula I:

wherein R₁ is phenyl or is phenyl disubstituted in the 2,6-positions,wherein R₂ is phenyl or is phenyl disubstituted in the 2,6-positions,wherein each of R₁ and R₂ is phenyl,


-7-

wherein each phenyl is disubstituted in the 2,6-position,

wherein R_1 is phenyl disubstituted in the 2,6-positions and R_2 is phenyl trisubstituted in the 2,4,6-positions,

wherein R_1 is 2,6-bis(1-methylethyl)phenyl and R_2 is 2,6-bis(1-methylethyl)phenyl or 2,4,6-tris(1-methylethyl)phenyl,

wherein one of R_1 and R_2 is the group



wherein t is zero or 1 to 4; w is zero or 1 to 4 with the proviso that the sum of t and w is not greater than 5; R_5 and R_6 are each independently selected from hydrogen or alkyl having from 1 to 6 carbon atoms, or when R_5 is hydrogen, R_6 can be selected from the groups defined for R_7 ; and R_7 is phenyl or phenyl substituted with from 1 to 3 substituents selected from a straight or branched alkyl group having from 1 to 6 carbon atoms, straight or branched alkoxy group having from 1 to 6 carbon atoms, phenoxy, hydroxy, fluorine, chlorine, bromine, nitro, trifluoromethyl, $-COOH$, $COOalkyl$ wherein alkyl has from 1 to 4 carbon atoms, or $-(CH_2)_pNR_3R_4$ wherein P , R_3 and R_4 have the meanings defined above.

Also preferred compounds of the instant invention are those of Formula I wherein

X is oxygen, sulfur or $(CR'R'')_n$;

Y is oxygen, sulfur or $(CR'R'')_n$, with the proviso that at least one of X or Y is $(CR'R'')_n$ wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, straight or branched alkyl of from 1 to 6 carbons, optionally substituted phenyl, halogen, hydroxy, alkoxy, acyloxy,

-8-

cycloalkyl, or R' and R" taken together form a carbonyl or a spirocycloalkyl group of from 3 to 10 carbons;

R is hydrogen;

- 5 R₁ is phenyl optionally substituted, straight or branched alkyl of from 1 to 10 carbon atoms, cycloalkyl of from 3 to 10 carbon atoms;
- R₂ is phenyl optionally substituted, straight or branched alkyl of from 1 to 10 carbon atoms, cycloalkyl of from 3 to 8 carbon atoms, phenoxy optionally substituted with the proviso that only if X is (CR'R")_n can R₁ be optionally substituted phenoxy and only if Y is (CR'R")_n can R₂ be optionally substituted phenoxy, and with the
- 10 further proviso that at least one of R₁ and R₂ is optionally substituted phenyl or phenoxy.
- 15

More preferred compounds of the instant invention are those of Formula I wherein

- 20 X is oxygen;
- Y is (CR'R")_n wherein n is an integer of from 1 to 2;
- R is hydrogen;
- R₁ is optionally substituted phenyl;
- R₂ is optionally substituted phenyl or phenoxy,
- 25 straight or branched alkyl of from 1 to 10 carbons, or cycloalkyl of from 3 to 10 carbons; and
- R' and R" are each independently hydrogen, straight or branched alkyl of from 1 to 6 carbons, optionally
- 30 substituted phenyl, halogen, hydroxy, alkoxy, acyloxy, cycloalkyl, or R' and R" taken together form a carbonyl or a spirocycloalkyl.

-9-

DETAILED DESCRIPTION OF THE INVENTION

The compounds useful in the present invention provide a class of N-acyl sulfamic acid esters (or thioesters), N-acyl sulfonamides, and N-sulfonyl carbamic acid esters (or thioesters) which are ACAT inhibitors, now found to be useful in treating cerebrovascular disease such as stroke, peripheral vascular disease, and restenosis. The compounds of the present invention have been found to be effective in lowering Lp(a).

In Formula I above, illustrative examples of straight or branched saturated hydrocarbon chains having from 1 to 20 carbon atoms include methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-butyl, tert-butyl, n-pentyl, isopentyl, n-hexyl, n-heptyl, n-octyl, n-undecyl, n-dodecyl, n-hexadecyl, 2,2-dimethyldodecyl, 2-tetradecyl, and n-octadecyl groups.

Illustrative examples of straight or branched hydrocarbon chains having from 1 to 20 carbon atoms and having from 1 to 3 double bonds include ethenyl, 2-propenyl, 2-butenyl, 3-pentenyl, 2-octenyl, 5-nonenyl, 4-undecenyl, 5-heptadecenyl, 3-octadecenyl, 9-octadecenyl, 2,2-dimethyl-11-eicosenyl, 9,12-octadecadienyl, and hexadecenyl.

Straight or branched alkoxy groups having from 1 to 6 carbon atoms include, for example, methoxy, ethoxy, n-propoxy, t-butoxy, and pentyloxy.

Illustrative examples of straight or branched alkyl groups having from 1 to 6 carbon atoms as used in Formula I include methyl, ethyl, n-propyl, isopropyl, n-pentyl, n-butyl, and tert-butyl.

Illustrative examples of cycloalkyl groups, as used in Formula I, include cyclopentyl, cyclohexyl, cyclooctyl, tetrahydronaphthyl, and 1- or 2-adamantyl.

-10-

Spirocycloalkyl groups are, for example, spirocyclopropyl, spirocyclobutyl, spirocyclopentyl, and spirocyclohexyl.

Illustrative examples of arylalkyl groups are:

5 benzyl, phenethyl, 3-phenylpropyl, 2-phenylpropyl, 4-phenylbutyl, 2-phenylbutyl, 3-phenylbutyl, benzhydryl, 2,2-diphenylethyl, and 3,3-diphenylpropyl.

10 Pharmaceutically acceptable salts of the compounds of Formula I are also included as a part of the present invention.

15 The base salts may be generated from compounds of Formula I by reaction of the latter with one equivalent of a suitable nontoxic, pharmaceutically acceptable base followed by evaporation of the solvent employed for the reaction and recrystallization of the salt, if required. The compounds of Formula I may be recovered from the base salt by reaction of the salt with an aqueous solution of a suitable acid such as hydrobromic, hydrochloric, or acetic acid.

20 Suitable bases for forming base salts of the compounds of this invention include amines such as triethylamine or dibutylamine, or alkali metal bases and alkaline earth metal bases. Preferred alkali metal hydroxides and alkaline earth metal hydroxides as salt
25 formers are the hydroxides of lithium, sodium, potassium, magnesium, or calcium. The class of bases suitable for the formation of nontoxic, pharmaceutically acceptable salts is well known to practitioners of the pharmaceutical formulation arts.
30 See, for example, Berge SN, et al, J. Pharm. Sci., 1977;66:1-19.

35 Suitable acids for forming acid salts of the compounds of this invention containing a basic group include, but are not necessarily limited to acetic, benzoic, benzenesulfonic, tartaric, hydrobromic, hydrochloric, citric, fumaric, gluconic, glucuronic,

-11-

glutamic, lactic, malic, maleic, methanesulfonic, pamoic, salicylic, stearic, succinic, sulfuric, and tartaric acids. The acid addition salts are formed by procedures well known in the art.

5 The compounds of the present invention may also exist in different stereoisomeric forms by virtue of the presence of asymmetric centers in the compound. The present invention contemplates all stereoisomeric forms of the compounds as well as mixtures thereof,
10 including racemic mixtures.

 Further, the compounds of this invention may exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms
15 are considered equivalent to the unsolvated forms for the purposes of this invention.

 Preferred compounds of the present invention are those wherein one of R_1 and R_2 is phenyl, and more preferably wherein one of R_1 and R_2 is substituted
20 phenyl, and still more preferably wherein one of R_1 and R_2 is phenyl disubstituted in the 2,6-positions.

 In one preferred embodiment both R_1 and R_2 are phenyl disubstituted in the 2,6-positions. In another preferred embodiment R_1 is phenyl disubstituted in the
25 2,6-position and R_2 is trisubstituted in the 2,4,6-positions.

 In another preferred embodiment of the present invention, R_1 is 2,6-bis(1-methylethyl)phenyl; and R_2 is 2,6-bis(1-methylethyl)phenyl or 2,4,6-tris(1-methylethyl)phenyl.
30

 Preferred compounds of Formula I include, but are not limited to the following:

 Sulfamic acid (phenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

35 Sulfamic acid[[2,4,6-tris(1-methylethyl)phenyl]-acetyl-2,6-bis(1-methylethyl)phenyl ester,

-12-

- Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-
acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]-
acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,
5 Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-
acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,
Sulfamic acid[adamantaneacetyl]-2,6-bis[1-methyl-
ethyl)phenyl ester,
Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-
10 acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,
Sulfamic acid[[2,4,6-tris(1-methylethyl)phenyl]-
acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,
Sulfamic acid (decanoyl)-2,6-bis-(1-methylethyl)-
phenyl ester,
15 Sulfamic acid (dodecanoyl)-2,6-bis-(1-methyl-
ethyl)phenyl ester,
2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methyl-
ethyl)phenyl]methyl]sulfonyl]benzeneacetamide,
2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris-
20 (1-methylethyl)phenyl]methyl]sulfonyl]benzeneacetamide-
sodium salt,
2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris-
(1-methylethyl)phenyl]methyl]sulfonyl]carbamate,
2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris-
25 (1-methylethyl)phenyl]methyl]sulfonyl]carbamate-sodium
salt,
Sulfamic acid (1-oxo-3,3-diphenylpropyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,6-dichlorophenyl(acetyl)]-
30 2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,6-dichlorophenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid trans-[(2-phenylcyclopropyl)-
carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
35 Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,

-13-

- Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,4,6-trimethylphenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,
5 Sulfamic acid [2-thiophenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [3-thiophenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-methoxyphenyl(acetyl)]-
10 2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (oxophenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-trifluoromethylphenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,
15 Sulfamic acid (1-oxo-2-phenylpropyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclopentylphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylacetyl)-
20 2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (diphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (triphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
25 Sulfamic acid [(1-phenylcyclopentyl)carbonyl]-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2-phenylbutyl)-
30 2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2,2-diphenylpropyl)-
2,6-bis(1-methylethyl)phenyl ester,
35 Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-
2,6-bis(1-methylethyl)phenyl ester,

-14-

Sulfamic acid (1-oxo-3-phenylpropyl)-
2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [1-oxo-3-[2,4,6-tris(1-methylethyl)-
phenyl]-2-propenyl]-2,6-bis(1-methylethyl)phenyl ester,

5 Sulfamic acid [1-oxo-3-[2,4,6-tris(1-methylethyl)-
phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(acetyloxy)[2,4,6-tris(1-methyl-
ethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl
ester,

10 Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)-
phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)-
phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

15 Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-
2,6-bis(1-methylethyl)phenyl ester sodium salt,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenoxy]-
acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenoxy]-
acetyl]-2,6-bis(1-methylethyl)phenyl ester, and

20 Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]-
acetyl]-2,6-bis(phenyl)phenyl ester.

The ability of the compounds of the present
invention to lower $Lp(a)$ is summarized in Table I
below. The procedure was: nine male cynomolgus
25 monkeys (*Macaca fascicularis*, 4-5 kg) were maintained
on a standard monkey chow diet (containing less than 5%
fat and only trace amounts of cholesterol). The diet
was available daily from 9 AM until 2 PM. These
animals transport approximately equal amounts of
30 cholesterol in HDL (47%) and LDL (51%) and have low
triglycerides compared to humans (approximately
50 mg/dL). Five weekly blood samples were taken from
anesthetized, restrained animals, and then the animals
were dosed with sulfamic acid [[2,4,6-tris-(1-methyl-
ethyl)phenyl]acetyl-2,6-bis(1-methylethyl)phenyl ester
35 (hereinafter the compound) daily before meals (for

-15-

3 weeks at 30 mg/kg) by incorporating it into oatmeal cream pies (Little Debbie Snack Cakes, McKee Foods, Collegedale, Tennessee). Tang breakfast beverage crystals (Kraft General Foods, Inc., White Plains, New York), and additional cream filling was also added to individual servings. Most animals consumed the drug-containing treat immediately since they were without food during the night. They were not given their daily meal until they consumed the treat. Mean plasma cholesterol (top line) and Lp(a) (bottom line) values are shown below (all values in mg/dL). The drug treatment was in Weeks 6 to 8 and are bolded in Table I.

15

TABLE I

Week											
1	2	3	4	5	6	7	8	9	10	11	12
147	147	155	155	167	112	112	111	117	128	127	160
17.1	18.9	15.1	13.8	18.2	13.6	12.5	11.4	13.6	17.7	18.7	21.3

20

The average baseline values for cholesterol and Lp(a) were 154 and 16.6 mg/dL, respectively. Using these values, the percentage decreases for cholesterol and Lp(a) are 28% and 31%, respectively. It is important to note that every animal demonstrated a decrease in cholesterol and Lp(a), i.e., there were no nonresponders to the compound. The decrease in total cholesterol was due primarily to a decrease in LDL-cholesterol.

25

30

The compounds of the present invention are thus useful in pharmaceutical formulations for the treatment of stroke, peripheral vascular disease, and restenosis.

35

In therapeutic use as agents for treating stroke, peripheral vascular disease, and restenosis, the compounds of Formulas I or II or pharmaceutically acceptable salts thereof are administered to the

-16-


patient at dosage levels of from 250 to 3000 mg per day. For a normal human adult of approximately 70 kg of body weight, this translates into a dosage of from 5 to 40 mg/kg of body weight per day. The specific dosages employed, however, may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the activity of the compound being employed. The determination of optimum dosages for a particular situation is within the skill of the art.



-17-

CLAIMS

1. A method for lowering the serum or plasma level of Lp(a) in a mammal in need of said treatment, comprising administering to said mammal an amount effective for lowering the serum or plasma level of said Lp(a) of a compound of formula



I

or a pharmaceutically acceptable salt thereof wherein:

- X and Y are selected from oxygen, sulfur and (CR'R'')_n wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, alkyl, alkoxy, halogen, hydroxy, acyloxy, cycloalkyl, phenyl optionally substituted or R' and R'' together form a spirocycloalkyl or a carbonyl; with the proviso at least one of X and Y is (CR'R'')_n and with the further proviso when X and Y are both (CR'R'')_n and R' and R'' are hydrogen and n is one, R₁ and R₂ are aryl; R is hydrogen, a straight or branched alkyl of from 1 to 8 carbon atoms or benzyl; R₁ and R₂ are each independently selected from (a) phenyl or phenoxy each of which is unsubstituted or is substituted with 1 to 5 substituents selected from phenyl, an alkyl group having from 1 to 6 carbon atoms and which is straight or branched, an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched;



-18-

- 40 phenoxy,
 hydroxy,
 fluorine,
 chlorine,
 bromine,
 nitro,
 trifluoromethyl,
 -COOH,
45 -COOalkyl wherein alkyl has from 1 to
 4 carbon atoms and is straight or
 branched,
 -(CH₂)_pNR₃R₄ wherein p is zero or one, and
50 each of R₃ and R₄ is selected from
 hydrogen or a straight or branched alkyl
 group having 1 to 4 carbon atoms;
 (b) 1- or 2-naphthyl unsubstituted or
substituted with from 1 to 3 substituents selected
from
55 phenyl,
 an alkyl group having from 1 to 6 carbon
 atoms and which is straight or branched,
 an alkoxy group having from 1 to 6 carbon
 atoms and which is straight or branched;
60 hydroxy,
 phenoxy,
 fluorine,
 chlorine,
 bromine,
65 nitro,
 trifluoromethyl,
 -COOH,
 -COOalkyl wherein alkyl has from 1 to
 4 carbon atoms and is straight or
70 branched,
 -(CH₂)_pNR₃R₄ wherein p, R₃ and R₄ have the
 meanings defined above;



-19-

(c) arylalkyl;

75 (d) a straight or branched alkyl chain having from 1 to 20 carbon atoms and which is saturated or contains from 1 to 3 double bonds; or

(e) adamantyl or a cycloalkyl group wherein the cycloalkyl moiety has from 3 to 6 carbon atoms;


80 with the provisos:

(i) where X is $(CH_2)_n$, Y is oxygen, and R_1 is a substituted phenyl, then R_2 is a substituted phenyl;

85 (ii) where Y is oxygen, X is $(CH_2)_n$, R_2 is phenyl or naphthyl, then R_1 is not a straight or branched alkyl chain; and

(iii) the following compounds are excluded:

90


X	Y	R	R_1	R_2
CH_2	O	H	$(CH_2)_2CH_3$	Ph
CH_2	O	H	CH_3	Ph
CH_2	O	H		i-Pr

2. A method according to Claim 1 wherein R_1 is phenyl.
3. A method according to Claim 2 wherein R_1 is phenyl disubstituted in the 2,6-positions.
4. A method according to Claim 1 wherein R_2 is phenyl.
5. A method according to Claim 4 wherein R_2 is phenyl disubstituted in the 2,6-positions.



-20-

6. A method according to Claim 1 wherein each of R_1 and R_2 is phenyl.
7. A method according to Claim 6 wherein each phenyl is disubstituted in the 2,6-positions.
8. A method according to Claim 1 wherein R_1 is phenyl disubstituted in the 2,6-positions and R_2 is phenyl trisubstituted in the 2,4,6-positions.
9. A method according to Claim 1 wherein R_1 is 2,6-bis(1-methylethyl)phenyl and R_2 is 2,6-bis(1-methylethyl)phenyl or 2,4,6-tris-(1-methylethyl)phenyl.
10. A method according to Claim 1 wherein R_1 is phenyl or phenyl disubstituted in the 2,6-positions, wherein R_2 is phenyl or is phenyl disubstituted in the 2,6-positions, wherein each of R_1 and R_2 is phenyl, wherein each phenyl is disubstituted in the 2,6-position, wherein R_1 is phenyl disubstituted in the 2,6-positions and R_2 is phenyl trisubstituted in the 2,4,6-positions, wherein R_1 is 2,6-bis(1-methylethyl)phenyl and R_2 is 2,6-bis(1-methylethyl)phenyl or 2,4,6-tris-(1-methylethyl)phenyl, wherein one of R_1 and R_2 is the group



20 wherein t is zero or 1 to 4; w is zero or 1 to 4 with the proviso that the sum of t and w is not greater than 5; R_5 and R_6 are each independently selected from hydrogen or alkyl having from 1 to



-21-

25 6 carbon atoms, or when R_5 is hydrogen, R_6 can be
selected from the groups defined for R_7 ; and R_7 is
phenyl or phenyl substituted with from 1 to
3 substituents selected from a straight or
branched alkyl group having from 1 to 6 carbon
atoms, straight or branched alkoxy group having
30 from 1 to 6 carbon atoms, phenoxy, hydroxy,
fluorine, chlorine, bromine, nitro,
trifluoromethyl, $-COOH$, $COOalkyl$ wherein alkyl has
from 1 to 4 carbon atoms, or $-(CH_2)_pNR_3R_4$ wherein
 P , R_3 and R_4 have the meanings defined above.

11. A method according to Claim 1 wherein
 X is oxygen, sulfur or $(CR'R'')_n$;
 Y is oxygen, sulfur or $(CR'R'')_n$ with the proviso
that at least one of X or Y is $(CR'R'')_n$
wherein n is an integer of from 1 to 4 and R'
5 and R'' are each independently hydrogen,
straight or branched alkyl of from 1 to
6 carbons, optionally substituted phenyl,
halogen, hydroxy, alkoxy, acyloxy,
cycloalkyl, or R' and R'' taken together form
10 a carbonyl or a spirocycloalkyl group of from
3 to 10 carbons;
 R is hydrogen;
 R_1 is phenyl optionally substituted, straight or
branched alkyl of from 1 to 10 carbon atoms,
15 cycloalkyl of from 3 to 10 carbon atoms;
 R_2 is phenyl optionally substituted, straight or
branched alkyl of from 1 to 10 carbon atoms,
cycloalkyl of from 3 to 8 carbon atoms,
phenoxy optionally substituted with the
20 proviso that only if X is $(CR'R'')_n$ can R_1 be
optionally substituted phenoxy and only if Y
is $(CR'R'')_n$ can R_2 be optionally substituted
phenoxy, and with the further proviso that at



-22-

- 25 least one of R_1 and R_2 is optionally substituted phenyl or phenoxy.
12. A method according to Claim 1 wherein
X is oxygen;
Y is $(CR'R'')_n$ wherein n is an integer of from 1 to 2;
5 R is hydrogen;
 R_1 is optionally substituted phenyl;
 R_2 is optionally substituted phenyl or phenoxy, straight or branched alkyl of from 1 to 10 carbons, or cycloalkyl of from 3 to 10 carbons;
10 R' and R'' are each independently hydrogen, straight or branched alkyl of from 1 to 6 carbons, optionally substituted phenyl, halogen, hydroxy, alkoxy, acyloxy, cycloalkyl, or R' and R'' taken together form a carbonyl or a spirocycloalkyl.
15
13. A method according to Claim 1 wherein the compound used is selected from:
Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester,
5 Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl-2,4,6-tris(1-methylethyl)phenyl ester,
10 Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,
Sulfamic acid[adamantaneacetyl]-2,6-bis-[1-methylethyl)phenyl ester,
Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,
15



-23-

- Sulfamic acid[[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,
- 20 Sulfamic acid (decanoyl)-2,6-bis-(1-methylethyl)phenyl ester,
- Sulfamic acid (dodecanoyl)-2,6-bis-(1-methylethyl)phenyl ester,
- 25 2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]benzene-acetamide,
- 2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]benzene-acetamide-sodium salt,
- 30 2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]carbamate,
- 2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]carbamate-sodium salt,
- 35 Sulfamic acid (1-oxo-3,3-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- 40 Sulfamic acid trans-[(2-phenylcyclopropyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- 45 Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2,4,6-trimethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- 50 Sulfamic acid [3-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,



-24-

- Sulfamic acid [2-methoxyphenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,
- 55 Sulfamic acid (oxophenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2-trifluoromethylphenyl-
(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (1-oxo-2-phenylpropyl)-
- 60 2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (cyclopentylphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (cyclohexylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
- 65 Sulfamic acid (diphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (triphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [(1-phenylcyclopentyl)-
carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
- 70 Sulfamic acid (3-methyl-1-oxo-
2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl
ester,
- Sulfamic acid (1-oxo-2-phenylbutyl)-
- 75 2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (cyclohexylphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (1-oxo-2,2-diphenylpropyl)-
2,6-bis(1-methylethyl)phenyl ester,
- 80 Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-
2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (1-oxo-3-phenylpropyl)-
2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [1-oxo-3-[2,4,6-tris-
(1-methylethyl)phenyl]-2-propenyl]-2,6-bis-
- 85 (1-methylethyl)phenyl ester,




-25-

- Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,
- 90 Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- 95 Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- 100 Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,
- Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- 105 Sulfamic acid [[2,6-bis(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and
- Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(phenyl)phenyl ester.
14. A method according to Claim 1 wherein sulfamic acid[[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester is administered.
15. A method of treating peripheral vascular disease comprising administering to a mammal in need of said treatment a therapeutically effective amount of a compound of formula



-26-



I

or a pharmaceutically acceptable salt thereof
wherein:

X and Y are selected from oxygen, sulfur and
(CR'R")_n wherein n is an integer of from
1 to 4 and R' and R" are each independently
hydrogen, alkyl, alkoxy, halogen, hydroxy,
acyloxy, cycloalkyl, phenyl optionally
substituted or R' and R" together form a
spirocycloalkyl or a carbonyl;

with the proviso at least one of X and Y is
(CR'R")_n and with the further proviso when
X and Y are both (CR'R")_n and R' and R" are
hydrogen and n is one, R₁ and R₂ are aryl;
R is hydrogen, a straight or branched alkyl of
from 1 to 8 carbon atoms or benzyl;

R₁ and R₂ are each independently selected from
(a) phenyl or phenoxy each of which is
unsubstituted or is substituted with 1 to
5 substituents selected from

phenyl,
an alkyl group having from 1 to 6 carbon
atoms and which is straight or branched,
an alkoxy group having from 1 to 6 carbon
atoms and which is straight or branched;
phenoxy,
hydroxy,
fluorine,
chlorine,
bromine,
nitro,
trifluoromethyl,
-COOH,




-27-

- 45 -COOalkyl wherein alkyl has from 1 to
4 carbon atoms and is straight or
branched,
- $(\text{CH}_2)_p\text{NR}_3\text{R}_4$ wherein p is zero or one, and
each of R_3 and R_4 is selected from
hydrogen or a straight or branched alkyl
group having 1 to 4 carbon atoms;
- 50 (b) 1- or 2-naphthyl unsubstituted or
substituted with from 1 to 3 substituents selected
from
phenyl,
an alkyl group having from 1 to 6 carbon
55 atoms and which is straight or branched,
an alkoxy group having from 1 to 6 carbon
atoms and which is straight or branched;
hydroxy,
phenoxy,
60 fluorine,
chlorine,
bromine,
nitro,
trifluoromethyl,
- 65 -COOH,
-COOalkyl wherein alkyl has from 1 to
4 carbon atoms and is straight or
branched,
- 70 - $(\text{CH}_2)_p\text{NR}_3\text{R}_4$ wherein p, R_3 and R_4 have the
meanings defined above;
- (c) arylalkyl;
- (d) a straight or branched alkyl chain
having from 1 to 20 carbon atoms and which is
saturated or contains from 1 to 3 double bonds; or
- 75 (e) adamantyl or a cycloalkyl group wherein
the cycloalkyl moiety has from 3 to 6 carbon
atoms;
with the provisos:



-28-

- 80 (i) where X is $(CH_2)_n$, Y is oxygen, and R_1 is a substituted phenyl, then R_2 is a substituted phenyl;
- (ii) where Y is oxygen, X is $(CH_2)_n$, R_2 is phenyl or naphthyl, then R_1 is not a straight or branched alkyl chain; and
- 85 (iii) the following compounds are excluded:

X	Y	R	R_1	R_2
CH_2	O	H	$(CH_2)_2CH_3$	Ph
CH_2	O	H	CH_3	Ph
90 CH_2	O	H		i-Pr

16. A method according to Claim 15 wherein the compound administered is selected from:

- Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester,
- 5 Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl-2,4,6-tris(1-methylethyl)phenyl ester,
- 10 Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,
- Sulfamic acid[adamantaneacetyl]-2,6-bis-[1-methylethyl)phenyl ester,
- Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,
- 15 Sulfamic acid[[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,



-29-

- 20 Sulfamic acid (decanoyl)-2,6-bis-(1-methylethyl)phenyl ester,
Sulfamic acid (dodecanoyl)-2,6-bis-(1-methylethyl)phenyl ester,
2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris-
- 25 (1-methylethyl)phenyl]methyl]sulfonyl]benzene-
acetamide,
2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris-
(1-methylethyl)phenyl]methyl]sulfonyl]benzene-
acetamide-sodium salt,
- 30 2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris-
(1-methylethyl)phenyl]methyl]sulfonyl]carbamate,
2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris-
(1-methylethyl)phenyl]methyl]sulfonyl]carbamate-
sodium salt,
- 35 Sulfamic acid (1-oxo-3,3-diphenylpropyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,6-dichlorophenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,6-dichlorophenyl(acetyl)]-
- 40 2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid trans-[(2-phenylcyclopropyl)-
carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,
- 45 Sulfamic acid [2,4,6-trimethoxyphenyl-
(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,4,6-trimethylphenyl-
(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-thiophenyl(acetyl)]-
- 50 2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [3-thiophenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-methoxyphenyl(acetyl)]-
2,6-bis(1-methylethyl)phenyl ester,



-30-

- 55 Sulfamic acid (oxophenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-trifluoromethylphenyl-
(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2-phenylpropyl)-
60 2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclopentylphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
65 Sulfamic acid (diphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (triphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [(1-phenylcyclopentyl)-
70 carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (3-methyl-1-oxo-
2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl
ester,
Sulfamic acid (1-oxo-2-phenylbutyl)-
75 2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylphenylacetyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2,2-diphenylpropyl)-
2,6-bis(1-methylethyl)phenyl ester,
80 Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-3-phenylpropyl)-
2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [1-oxo-3-[2,4,6-tris-
85 (1-methylethyl)phenyl]-2-propenyl]-2,6-bis-
(1-methylethyl)phenyl ester,
Sulfamic acid [1-oxo-3-[2,4,6-tris-
(1-methylethyl)phenyl]propyl]-2,6-bis(1-methyl-
ethyl)phenyl ester,




-31-

- 90 Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- 95 Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- 100 Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,
- Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- 105 Sulfamic acid [[2,6-bis(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and
- Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(phenyl)phenyl ester.

17. A method of treating peripheral vascular disease comprising administering to a patient in need of said treatment a compound named sulfamic acid
- 5 [[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester.
18. A method of treating restenosis comprising administering to a mammal in need of said treatment a therapeutically effective amount of a compound of formula

5



I



-32-

- 10 or a pharmaceutically acceptable salt thereof
wherein:
X and Y are selected from oxygen, sulfur and
(CR'R")_n wherein n is an integer of from
1 to 4 and R' and R" are each independently
15 hydrogen, alkyl, alkoxy, halogen, hydroxy,
acyloxy, cycloalkyl, phenyl optionally
substituted or R' and R" together form a
spirocycloalkyl or a carbonyl;
with the proviso at least one of X and Y is
20 (CR'R")_n and with the further proviso when
X and Y are both (CR'R")_n and R' and R" are
hydrogen and n is one, R₁ and R₂ are aryl;
R is hydrogen, a straight or branched alkyl of
from 1 to 8 carbon atoms or benzyl;
25 R₁ and R₂ are each independently selected from
(a) phenyl or phenoxy each of which is
unsubstituted or is substituted with 1 to
5 substituents selected from
phenyl,
30 an alkyl group having from 1 to 6 carbon
atoms and which is straight or branched,
an alkoxy group having from 1 to 6 carbon
atoms and which is straight or branched;
phenoxy,
35 hydroxy,
fluorine,
chlorine,
bromine,
nitro,
40 trifluoromethyl,
-COOH,
-COOalkyl wherein alkyl has from 1 to
4 carbon atoms and is straight or
branched,



-33-


- 45 $-(CH_2)_pNR_3R_4$ wherein p is zero or one, and
each of R_3 and R_4 is selected from
hydrogen or a straight or branched alkyl
group having 1 to 4 carbon atoms;
- 50 (b) 1- or 2-naphthyl unsubstituted or
substituted with from 1 to 3 substituents selected
from
phenyl,
an alkyl group having from 1 to 6 carbon
atoms and which is straight or branched,
55 an alkoxy group having from 1 to 6 carbon
atoms and which is straight or branched;
hydroxy,
phenoxy,
fluorine,
60 chlorine,
bromine,
nitro,
trifluoromethyl,
-COOH,
65 -COOalkyl wherein alkyl has from 1 to
4 carbon atoms and is straight or
branched,
 $-(CH_2)_pNR_3R_4$ wherein p , R_3 and R_4 have the
meanings defined above;
- 70 (c) arylalkyl;
(d) a straight or branched alkyl chain
having from 1 to 20 carbon atoms and which is
saturated or contains from 1 to 3 double bonds; or
(e) adamantyl or a cycloalkyl group wherein
75 the cycloalkyl moiety has from 3 to 6 carbon
atoms;
with the provisos:
(i) where X is $(CH_2)_n$, Y is oxygen, and R_1
is a substituted phenyl, then R_2 is a
80 substituted phenyl;



-34-

- (ii) where Y is oxygen, X is $(CH_2)_n$, R_2 is phenyl or naphthyl, then R_1 is not a straight or branched alkyl chain; and
 (iii) the following compounds are excluded:

85

X	Y	R	R_1	R_2
CH_2	O	H	$(CH_2)_2CH_3$	Ph
CH_2	O	H	CH_3	Ph
CH_2	O	H		i-Pr

90

19. A method of treating restenosis according to Claim 18 wherein the compound administered is selected from:

Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester,

5

Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

10

Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid[adamantaneacetyl]-2,6-bis-[1-methylethyl)phenyl ester,

15

Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,

Sulfamic acid[[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,

20

Sulfamic acid (decanoyl)-2,6-bis-(1-methylethyl)phenyl ester,



-35-

- Sulfamic acid (dodecanoyl)-2,6-bis-(1-methylethyl)phenyl ester,
- 25 2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]benzene-acetamide,
- 2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]benzene-acetamide-sodium salt,
- 30 2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]carbamate,
- 2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]carbamate-sodium salt,
- 35 Sulfamic acid (1-oxo-3,3-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- 40 Sulfamic acid trans-[(2-phenylcyclopropyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- 45 Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2,4,6-trimethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- 50 Sulfamic acid [3-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [2-methoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- 55 Sulfamic acid (oxophenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,



-36-

- Sulfamic acid [2-trifluoromethylphenyl-(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- 60 Sulfamic acid (1-oxo-2-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (cyclopentylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (cyclohexylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
- 65 Sulfamic acid (diphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (triphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
- 70 Sulfamic acid [(1-phenylcyclopentyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (1-oxo-2-phenylbutyl)-2,6-bis(1-methylethyl)phenyl ester,
- 75 Sulfamic acid (cyclohexylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (1-oxo-2,2-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
- 80 Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (1-oxo-3-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]-2-propenyl]-2,6-bis(1-methylethyl)phenyl ester,
- 85 Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,
- 90 Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,



-37-

95 Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

100 Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

105 Sulfamic acid [[2,6-bis(1-methylethyl)phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and


Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(phenyl)phenyl ester.

20. A method of treating restenosis comprising administering to a patient in need of said treatment a compound named sulfamic acid [[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester.

5

21. A method of treating cerebrovascular disease comprising administering to a mammal in need of said treatment a therapeutically effective amount of a compound of formula

5



I

10

or a pharmaceutically acceptable salt thereof wherein:



-38-

- X and Y are selected from oxygen, sulfur and
15 (CR'R")_n wherein n is an integer of from
1 to 4 and R' and R" are each independently
hydrogen, alkyl, alkoxy, halogen, hydroxy,
acyloxy, cycloalkyl, phenyl optionally
substituted or R' and R" together form a
20 spirocycloalkyl or a carbonyl;
with the proviso at least one of X and Y is
(CR'R")_n and with the further proviso when
X and Y are both (CR'R")_n and R' and R" are
hydrogen and n is one, R₁ and R₂ are aryl;
25 R is hydrogen, a straight or branched alkyl of
from 1 to 8 carbon atoms or benzyl;
R₁ and R₂ are each independently selected from
(a) phenyl or phenoxy each of which is
unsubstituted or is substituted with 1 to
30 5 substituents selected from
phenyl,
an alkyl group having from 1 to 6 carbon
atoms and which is straight or branched,
an alkoxy group having from 1 to 6 carbon
35 atoms and which is straight or branched;
phenoxy,
hydroxy,
fluorine,
chlorine,
40 bromine,
nitro,
trifluoromethyl,
-COOH,
-COOalkyl wherein alkyl has from 1 to
45 4 carbon atoms and is straight or
branched,
-(CH₂)_pNR₃R₄ wherein p is zero or one, and
each of R₃ and R₄ is selected from




-39-

- hydrogen or a straight or branched alkyl
50 group having 1 to 4 carbon atoms;
- (b) 1- or 2-naphthyl unsubstituted or
substituted with from 1 to 3 substituents selected
from
- phenyl,
55 an alkyl group having from 1 to 6 carbon
atoms and which is straight or branched,
an alkoxy group having from 1 to 6 carbon
atoms and which is straight or branched;
hydroxy,
60 phenoxy,
fluorine,
chlorine,
bromine,
nitro,
65 trifluoromethyl,
-COOH,
-COOalkyl wherein alkyl has from 1 to
4 carbon atoms and is straight or
branched,
70 $-(CH_2)_pNR_3R_4$ wherein p, R_3 and R_4 have the
meanings defined above;
- (c) arylalkyl;
(d) a straight or branched alkyl chain
having from 1 to 20 carbon atoms and which is
75 saturated or contains from 1 to 3 double bonds; or
- (e) adamantyl or a cycloalkyl group wherein
the cycloalkyl moiety has from 3 to 6 carbon
atoms;
with the provisos:
- 80 (i) where X is $(CH_2)_n$, Y is oxygen, and R_1
is a substituted phenyl, then R_2 is a
substituted phenyl;



-40-

- 85 (ii) where Y is oxygen, X is $(CH_2)_n$, R_2 is phenyl or naphthyl, then R_1 is not a straight or branched alkyl chain; and
(iii) the following compounds are excluded:

X	Y	R	R_1	R_2
CH_2	O	H	$(CH_2)_2CH_3$	Ph
90 CH_2	O	H	CH_3	Ph
CH_2	O	H		i-Pr

22. A method of treating cerebrovascular disease according to Claim 21 wherein the compound administered is selected from:

- 5 Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester,
Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,4,6-tris(1-methylethyl)phenyl
10 ester,
Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,
Sulfamic acid[adamantaneacetyl]-2,6-bis-[1-methylethyl)phenyl ester,
15 Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,
Sulfamic acid[[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,
20 Sulfamic acid (decanoyl)-2,6-bis-(1-methylethyl)phenyl ester,



-41-

Sulfamic acid (dodecanoyl)-2,6-bis-(1-methylethyl)phenyl ester,

25 2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]benzene-acetamide,

 2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]benzene-acetamide-sodium salt,

30 2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]carbamate,

 2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris-(1-methylethyl)phenyl]methyl]sulfonyl]carbamate-sodium salt,

35 Sulfamic acid (1-oxo-3,3-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

 Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

40 Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

 Sulfamic acid trans-[(2-phenylcyclopropyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,

 Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

45 Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

 Sulfamic acid [2,4,6-trimethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

50 Sulfamic acid [2-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

 Sulfamic acid [3-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

 Sulfamic acid [2-methoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

55 Sulfamic acid (oxophenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,



-42-

- 60 Sulfamic acid [2-trifluoromethylphenyl-(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclopentylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
65 Sulfamic acid (diphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (triphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
70 Sulfamic acid [(1-phenylcyclopentyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester,
75 Sulfamic acid (1-oxo-2-phenylbutyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2,2-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
80 Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-3-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
85 Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]-2-propenyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,
90 Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,



-43-

95 Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

100 Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,

105 Sulfamic acid [[2,4,6-tris(1-methylethyl)phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and

110 Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(phenyl)phenyl ester.

23. A method of treating cerebrovascular disease comprising administering to a patient in need of said treatment a compound named sulfamic acid [[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester.

5

19 SEP 2000


44

REPLACEMENT SHEETS SG

- ☒ Description : paged to paged
☒ Claim(s) : paged 44 to paged 80
☐ Drawing(s) : paged to paged
☐ Abstract : paged to paged

DATE OF RECEIPT 19/9/00

24. Use of a compound of formula



(I)



or a pharmaceutically acceptable salt thereof wherein:

X and Y are selected from oxygen, sulfur and (CR'R'')_n wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, alkyl, alkoxy, halogen, hydroxy, acyloxy, cycloalkyl, phenyl optionally substituted or R' and R'' together form a spirocycloalkyl or a carbonyl;

with the proviso at least one of X and Y is (CR'R'')_n and with the further proviso when X and Y are both (CR'R'')_n and R' and R'' are hydrogen and n is one, R₁ and R₂ are aryl;

R is hydrogen, a straight or branched alkyl of from 1 to 8 carbon atoms or benzyl;

R₁ and R₂ are each independently selected from

(a) phenyl or phenoxy each of which is unsubstituted or is substituted with 1 to 5 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

phenoxy,

hydroxy,

fluorine,

chlorine,

bromine,

nitro,

trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

-(CH₂)_pNR₃R₄ wherein p is zero or one, and each of R₃ and R₄ is selected from hydrogen or a straight or branched alkyl group having 1 to 4 carbon atoms;

(b) 1- or 2-naphthyl unsubstituted or substituted with from 1 to 3 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

19 SEP 2000



an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,
 hydroxy,
 phenoxy,
 fluorine,
 chlorine,
 bromine,
 nitro,
 trifluoromethyl,
 -COOH,
 -COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,
 -(CH₂)_pNR₃R₄ wherein p, R₃ and R₄ have the meanings defined above;

(c) arylalkyl;

(d) a straight or branched alkyl chain having from 1 to 20 carbon atoms and which is saturated or contains from 1 to 3 double bonds; or


(e) adamantyl or a cycloalkyl group wherein the cycloalkyl moiety has from 3 to 6 carbon atoms;

with the provisos

(i) where X is (CH₂)_n, Y is oxygen, and R₁ is a substituted phenyl, then R₂ is a substituted phenyl;

(ii) where Y is oxygen, X is (CH₂)_n, R₂ is phenyl or naphthyl, then R₁ is not a straight or branched alkyl chain; and

(iii) the following compounds are excluded:


X	Y	R	R ₁	R ₂
CH ₂	O	H	(CH ₂) ₂ CH ₃	Ph
CH ₂	O	H	CH ₃	Ph
CH ₂	O	H		i-Pr

for the manufacture of a medicament for lowering the serum or plasma level of Lp(a) in a mammal.


25. Use according to Claim 24 wherein in the compound of formula I R₁ is phenyl.

26. Use according to Claim 25 wherein in the compound of formula I R₁ is phenyl disubstituted in the 2,6-positions.

27. Use according to Claim 24 wherein in the compound of formula I R₂ is phenyl.



28. Use according to Claim 27 wherein in the compound of formula I R_2 is phenyl disubstituted in the 2,6-positions.
29. Use according to Claim 24 wherein in the compound of formula I R_1 and R_2 is phenyl.
30. Use according to Claim 29 wherein each phenyl is disubstituted in the 2,6-positions.
31. Use according to Claim 24 wherein in the compound of formula I R_1 is phenyl disubstituted in the 2,6-positions and R_2 is phenyl trisubstituted in the 2,4,6-positions.
32. Use according to Claim 24 wherein in the compound of formula I R_1 is 2,6-bis(1-methylethyl)phenyl and R_2 is 2,6-bis(1-methylethyl)phenyl or 2,4,6-tris(1-methylethyl)phenyl.
33. Use according to Claim 24 wherein R_1 is phenyl or phenyl disubstituted in the 2,6-positions, wherein R_2 is phenyl or is phenyl disubstituted in the 2,6-positions, wherein each of R_1 and R_2 is phenyl, wherein each phenyl is disubstituted in the 2,6-position, wherein R_1 is phenyl disubstituted in the 2,6-positions and R_2 is phenyl trisubstituted in the 2,4,6-positions, wherein R_1 is 2,6-bis(1-methylethyl)phenyl and is 2,6-bis(1-methylethyl)phenyl or 2,4,6-tris(1-methylethyl)phenyl, wherein one of R_1 and R_2 is the group



wherein t is zero or 1 to 4; w is zero or 1 to 4 with the proviso that the sum of t and w is not greater than 5; R_5 and R_6 are each independently selected from hydrogen or alkyl having from 1 to 6 carbon atoms, or when R_5 is hydrogen, R_6 can be selected from the groups defined for R_7 ; and R_7 is phenyl or phenyl substituted with from 1 to 3 substituents selected from a straight or branched alkyl group having from 1 to 6 carbon atoms, straight or branched alkoxy group having from 1 to 6 carbon atoms, phenoxy, hydroxy, fluorine, chlorine, bromine, nitro, trifluoromethyl, -COOH, COOalkyl wherein alkyl has from 1 to 4 carbon atoms, or $-(CH_2)_pNR_3R_4$ wherein P, R_3 and R_4 have the meanings defined above.

34. Use according to Claim 24 wherein in the compound of formula I

X is oxygen, sulfur or $(CR'R'')_n$;

Y is oxygen, sulfur or $(CR'R'')_n$ with the proviso that at least one of X or Y is $(CR'R'')_n$ wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, straight or branched alkyl of from 1 to 6 carbons, optionally substituted phenyl, halogen, hydroxy, alkoxy, acyloxy, cycloalkyl,

19 SEP 2000



or R' and R" taken together form a carbonyl or a spirocycloalkyl group of from 3 to 10 carbons;

R is hydrogen;

R₁ is phenyl optionally substituted, straight or branched alkyl of from 1 to 10 carbon atoms, cycloalkyl of from 3 to 10 carbon atoms;

R₂ is phenyl optionally substituted, straight or branched alkyl of from 1 to 10 carbon atoms, cycloalkyl of from 3 to 8 carbon atoms, phenoxy optionally substituted with the proviso that only if X is (CR'R")_n can R₁ be optionally substituted phenoxy and only if Y is (CR'R")_n can be optionally substituted phenoxy, and with the further proviso that at least one of R₁ and R₂ is optionally substituted phenyl or phenoxy.

35. Use according to Claim 24 wherein in the compound of formula I

X is oxygen;

Y is (CR'R")_n wherein n is an integer of from 1 to 2;

R is hydrogen;

R₁ is optionally substituted phenyl;

R₂ is optionally substituted phenyl or phenoxy, straight or branched alkyl of from 1 to 10 carbons, or cycloalkyl of from 3 to 10 carbons;

R' and R" are each independently hydrogen, straight or branched alkyl of from 1 to 6 carbons, optionally substituted phenyl, halogen, hydroxy, alkoxy, acyloxy, cycloalkyl, or R' and R" taken together form a carbonyl or a spirocycloalkyl.

36. Use according to Claim 24 wherein in the compound of formula I is selected from:

Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester,


Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,


Sulfamic acid [adamantaneacetyl]-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester-sodium salt,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,



- Sulfamic acid (decanoyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (dodecanoyl)-2,6-bis(1-methylethyl)phenyl ester,
2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]sulfonyl]-benzene-acetamide,
2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]benzene-acetamide-sodium salt,
2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate,
2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate-sodium salt,
Sulfamic acid (1-oxo-3,3-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid trans-[(2-phenylcyclopropyl)carbonyl]-2,6-bis(1-methylethyl)-phenyl ester,
Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,4,6-trimethylphenyl-(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [3-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-methoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (oxophenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-trifluoromethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclopentylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (diphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (triphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,



Sulfamic acid [(1-phenylcyclopentyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2-phenylbutyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (cyclohexylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2,2-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-3-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]-2-propenyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,


Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and

Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(phenyl)-phenyl ester.

37. Use according to Claim 24 wherein in the compound of formula I is sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester.

38. Use of a compound of formula



or a pharmaceutically acceptable salt thereof wherein:

19 SEP 2000



X and Y are selected from oxygen, sulfur and $(CR'R'')_n$, wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, alkyl, alkoxy, halogen, hydroxy, acyloxy, cycloalkyl, phenyl optionally substituted or R' and R'' together form a spirocycloalkyl or a carbonyl;

with the proviso at least one of X and Y is $(CR'R'')_n$ and with the further proviso when X and Y are both $(CR'R'')_n$ and R' and R'' are hydrogen and n is one, R₁ and R₂ are aryl;

R is hydrogen, a straight or branched alkyl of from 1 to 8 carbon atoms or benzyl;

R₁ and R₂ are each independently selected from

(a) phenyl or phenoxy each of which is unsubstituted or is substituted with 1 to 5 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

phenoxy,

hydroxy,

fluorine,

chlorine,

bromine,

nitro,

trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

$-(CH_2)_pNR_3R_4$ wherein p is zero or one, and each of R₃ and R₄ is selected from hydrogen or a straight or branched alkyl group having 1 to 4 carbon atoms;

(b) 1- or 2-naphthyl unsubstituted or substituted with from 1 to 3 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,


an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

hydroxy,

phenoxy,

fluorine,

chlorine,



bromine,

nitro,

trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

-(CH₂)_pNR₃R₄ wherein p, R₃ and R₄ have the meanings defined above;

(c) arylalkyl;

(d) a straight or branched alkyl chain having from 1 to 20 carbon atoms and which is saturated or contains from 1 to 3 double bonds; or


(e) adamantyl or a cycloalkyl group wherein the cycloalkyl moiety has from 3 to 6 carbon atoms;

with the provisos

(i) where X is (CH₂)_n, Y is oxygen, and R₁ is a substituted phenyl, then R₂ is a substituted phenyl;

(ii) where Y is oxygen, X is (CH₂)_n, R₂ is phenyl or naphthyl, then R₁ is not a straight or branched alkyl chain; and

(iii) the following compounds are excluded:

X	Y	R	R ₁	R ₂
CH ₂	O	H	(CH ₂) ₂ CH ₃	Ph
CH ₂	O	H	CH ₃	Ph
CH ₂	O	H		i-Pr

for the manufacture of a medicament for treating peripheral vascular disease.

39. Use according to Claim 38 wherein the compound of formula I is selected from:

Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester,


Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [adamantaneacetyl]-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester-sodium salt,



Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,

Sulfamic acid (decanoyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (dodecanoyl)-2,6-bis(1-methylethyl)phenyl ester,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]sulfonyl]-benzene-acetamide,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]benzene-acetamide-sodium salt,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate-sodium salt,

Sulfamic acid (1-oxo-3,3-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,6-dichlorophenyl(acetyl)]2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid trans-[(2-phenylcyclopropyl)carbonyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,4,6-trimethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [3-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-methoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (oxophenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-trifluoromethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (cyclopentylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,


Sulfamic acid (cyclohexylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (diphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,



* C L M S *

53



Sulfamic acid (triphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(1-phenylcyclopentyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2-phenylbutyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (cyclohexylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2,2-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-3-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]-2-propenyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,


Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,

Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and

Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(phenyl)-phenyl ester.

40. Use according to Claim 39 wherein the compound of formula I is sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester.
41. Use of a compound of formula



(I)



or a pharmaceutically acceptable salt thereof wherein:

X and Y are selected from oxygen, sulfur and $(CR'R'')_n$ wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, alkyl, alkoxy, halogen, hydroxy, acyloxy, cycloalkyl, phenyl optionally substituted or R' and R'' together form a spirocycloalkyl or a carbonyl;
with the proviso at least one of X and Y is $(CR'R'')_n$ and with the further proviso when X and Y are both $(CR'R'')_n$ and R' and R'' are hydrogen and n is one, R₁ and R₂ are aryl;

R is hydrogen, a straight or branched alkyl of from 1 to 8 carbon atoms or benzyl;

R₁ and R₂ are each independently selected from

(a) phenyl or phenoxy each of which is unsubstituted or is substituted with 1 to 5 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

phenoxy,

hydroxy,

fluorine,

chlorine,

bromine,

nitro,

trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

$-(CH_2)_pNR_3R_4$ wherein p is zero or one, and each of R₃ and R₄ is selected from hydrogen or a straight or branched alkyl group having 1 to 4 carbon atoms;

(b) 1- or 2-naphthyl unsubstituted or substituted with from 1 to 3 substituents selected from

phenyl,


an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

hydroxy,

phenoxy,

fluorine,



chlorine,

bromine,

nitro,

trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

-(CH₂)_pNR₃R₄ wherein p, R₃ and R₄ have the meanings defined above;


(c) arylalkyl;

(d) a straight or branched alkyl chain having from 1 to 20 carbon atoms and which is saturated or contains from 1 to 3 double bonds; or

(e) adamantyl or a cycloalkyl group wherein the cycloalkyl moiety has from 3 to 6 carbon atoms;

with the provisos

- (i) where X is (CH₂)_n, Y is oxygen, and R₁ is a substituted phenyl, then R₂ is a substituted phenyl;
- (ii) where Y is oxygen, X is (CH₂)_n, R₂ is phenyl or naphthyl, then R₁ is not a straight or branched alkyl chain; and
- (iii) the following compounds are excluded:

X	Y	R	R ₁	R ₂
CH ₂	O	H	(CH ₂) ₂ CH ₃	Ph
CH ₂	O	H	CH ₃	Ph
CH ₂	O	H		i-Pr

for the manufacture of a medicament for treating restenosis.

42. Use according to Claim 41 wherein the compound of formula I is selected from:

Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [adamantaneacetyl]-2,6-bis-(1-methylethyl)phenyl ester,



Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester-sodium salt,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,

Sulfamic acid (decanoyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (dodecanoyl)-2,6-bis(1-methylethyl)phenyl ester,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]sulfonyl]-benzene-acetamide,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]benzene-acetamide-sodium salt,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate-sodium salt,

Sulfamic acid (1-oxo-3,3-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,6-dichlorophenyl(acetyl)]2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid trans-[(2-phenylcyclopropyl)carbonyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,4,6-trimethylphenyl-(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [3-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-methoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (oxophenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-trifluoromethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (cyclopentylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,



19 SEP 2000


Sulfamic acid (cyclohexylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (diphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (triphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [(1-phenylcyclopentyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2-phenylbutyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2,2-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-3-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]-2-propenyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,
Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [[2,6-bis(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and
Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(phenyl)-phenyl ester.

43. Use according to Claim 41 wherein the compound of formula I is sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester.
44. Use of a compound of formula



19 SEP 2000

58



or a pharmaceutically acceptable salt thereof wherein:

X and Y are selected from oxygen, sulfur and $(\text{CR}'\text{R}'')_n$ wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, alkyl, alkoxy, halogen, hydroxy, acyloxy, cycloalkyl, phenyl optionally substituted or R' and R'' together form a spirocycloalkyl or a carbonyl;

with the proviso at least one of X and Y is $(\text{CR}'\text{R}'')_n$ and with the further proviso when X and Y are both $(\text{CR}'\text{R}'')_n$ and R' and R'' are hydrogen and n is one, R₁ and R₂ are aryl;

R is hydrogen, a straight or branched alkyl of from 1 to 8 carbon atoms or benzyl;

R₁ and R₂ are each independently selected from

(a) phenyl or phenoxy each of which is unsubstituted or is substituted with 1 to 5 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

phenoxy,

hydroxy,

fluorine,

chlorine,

bromine,

nitro,

trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

$-(\text{CH}_2)_p\text{NR}_3\text{R}_4$ wherein p is zero or one, and each of R₃ and R₄ is selected from hydrogen or a straight or branched alkyl group having 1 to 4 carbon atoms;

(b) 1- or 2-naphthyl unsubstituted or substituted with from 1 to 3 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,



19 SEP 2000

hydroxy,
phenoxy,
fluorine,
chlorine,
bromine,
nitro,
trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

-(CH₂)_pNR₃R₄ wherein p, R₃ and R₄ have the meanings defined above;

(c) arylalkyl;

(d) a straight or branched alkyl chain having from 1 to 20 carbon atoms and which is saturated or contains from 1 to 3 double bonds; or


(e) adamantyl or a cycloalkyl group wherein the cycloalkyl moiety has from 3 to 6 carbon atoms;

with the provisos

(i) where X is (CH₂)_n, Y is oxygen, and R₁ is a substituted phenyl, then R₂ is a substituted phenyl;

(ii) where Y is oxygen, X is (CH₂)_n, R₂ is phenyl or naphthyl, then R₁ is not a straight or branched alkyl chain; and

(iii) the following compounds are excluded:

X	Y	R	R ₁	R ₂
CH ₂	O	H	(CH ₂) ₂ CH ₃	Ph
CH ₂	O	H	CH ₃	Ph
CH ₂	O	H		i-Pr


for the manufacture of a medicament for treating cerebrovascular disease.

45. Use according to Claim 44 wherein the compound of formula I is selected from:

Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,



Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [adamantaneacetyl]-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester-sodium salt,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,

Sulfamic acid (decanoyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (dodecanoyl)-2,6-bis(1-methylethyl)phenyl ester,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]sulfonyl]-benzene-acetamide,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]benzene-acetamide-sodium salt,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate-sodium salt,

Sulfamic acid (1-oxo-3,3-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,6-dichlorophenyl(acetyl)]2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid trans-[(2-phenylcyclopropyl)carbonyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,4,6-trimethylphenyl-(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [3-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-methoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (oxophenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,




- Sulfamic acid [2-trifluoromethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (1-oxo-2-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (cyclopentylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (cyclohexylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (diphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (triphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [(1-phenylcyclopentyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (1-oxo-2-phenylbutyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (cyclohexylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (1-oxo-2,2-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (1-oxo-3-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]-2-propenyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,
- Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
- Sulfamic acid [[2,6-bis(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and
- Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(phenyl)-phenyl ester.



19 SEP 2000

46. Use according to Claim 45 wherein the compound of formula I is sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl-2,6-bis(1-methylethyl)phenyl ester.
47. A compound of formula



or a pharmaceutically acceptable salt thereof wherein:

X and Y are selected from oxygen, sulfur and $(\text{CR}'\text{R}'')_n$, wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, alkyl, alkoxy, halogen, hydroxy, acyloxy, cycloalkyl, phenyl optionally substituted or R' and R'' together form a spirocycloalkyl or a carbonyl;

with the proviso at least one of X and Y is $(\text{CR}'\text{R}'')_n$ and with the further proviso when X and Y are both $(\text{CR}'\text{R}'')_n$ and R' and R'' are hydrogen and n is one, R₁ and R₂ are aryl;

R is hydrogen, a straight or branched alkyl of from 1 to 8 carbon atoms or benzyl;

R₁ and R₂ are each independently selected from

(a) phenyl or phenoxy each of which is unsubstituted or is substituted with 1 to 5 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

phenoxy,

hydroxy,

fluorine,

chlorine,

bromine,

nitro,


trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

$-(\text{CH}_2)_p\text{NR}_3\text{R}_4$ wherein p is zero or one, and each of R₃ and R₄ is selected from hydrogen or a straight or branched alkyl group having 1 to 4 carbon atoms;

(b) 1- or 2-naphthyl unsubstituted or substituted with from 1 to 3 substituents selected from



phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

hydroxy,

phenoxy,

fluorine,

chlorine,

bromine,

nitro,

trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

-(CH₂)_pNR₃R₄ wherein p, R₃ and R₄ have the meanings defined above;

(c) arylalkyl;

(d) a straight or branched alkyl chain having from 1 to 20 carbon atoms and which is saturated or contains from 1 to 3 double bonds; or


(e) adamantyl or a cycloalkyl group wherein the cycloalkyl moiety has from 3 to 6 carbon atoms;

with the provisos

(i) where X is (CH₂)_n, Y is oxygen, and R₁ is a substituted phenyl, then R₂ is a substituted phenyl;

(ii) where Y is oxygen, X is (CH₂)_n, R₂ is phenyl or naphthyl, then R₁ is not a straight or branched alkyl chain; and

(iii) the following compounds are excluded:

X	Y	R	R ₁	R ₂
CH ₂	O	H	(CH ₂) ₂ CH ₃	Ph
CH ₂	O	H	CH ₃	Ph
CH ₂	O	H		i-Pr

when used in an effective amount for lowering the serum or plasma level of Lp(a) in a mammal in need of said treatment.


48. A compound when used according to Claim 47 wherein R₁ is phenyl.

49. A compound when used according to Claim 48 wherein R₁ is phenyl disubstituted in the 2,6-positions.



19 SEP 2000

50. A compound when used according to Claim 47 wherein R_2 is phenyl.
51. A compound when used according to Claim 50 wherein R_2 is phenyl disubstituted in the 2,6-positions.
52. A compound when used according to Claim 47 wherein each of R_1 and R_2 is phenyl.
53. A compound when used according to Claim 52 wherein each phenyl is disubstituted in the 2,6-positions.
54. A compound when used according to Claim 47 wherein R_1 is phenyl disubstituted in the 2,6-positions and R_2 is phenyl trisubstituted in the 2,4,6-positions.
55. A compound when used according to Claim 47 wherein R_1 is 2,6-bis(1-methylethyl)phenyl and R_2 is 2,6-bis(1-methylethyl)phenyl or 2,4,6-tris-(1-methylethyl)phenyl.
56. A compound when used according to Claim 47 wherein R_1 is phenyl or phenyl disubstituted in the 2,6-positions, wherein R_2 is phenyl or is phenyl disubstituted in the 2,6-positions, wherein each of R_1 and R_2 is phenyl, wherein each phenyl is disubstituted in the 2,6-position, wherein R_1 is phenyl disubstituted in the 2,6-positions and R_2 is phenyl trisubstituted in the 2,4,6-positions, wherein R_1 is 2,6-bis(1-methylethyl)phenyl and is 2,6-bis(1-methylethyl)phenyl or 2,4,6-tris-(1-methylethyl)phenyl, wherein one of R_1 and R_2 is the group



wherein t is zero or 1 to 4; w is zero or 1 to 4 with the proviso that the sum of t and w is not greater than 5; R_5 and R_6 are each independently selected from hydrogen or alkyl having from 1 to 6 carbon atoms, or when R_5 is hydrogen, R_6 can be selected from the groups defined for R_7 ; and R_7 is phenyl or phenyl substituted with from 1 to 3 substituents selected from a straight or branched alkyl group having from 1 to 6 carbon atoms, straight or branched alkoxy group having from 1 to 6 carbon atoms, phenoxy, hydroxy, fluorine, chlorine, bromine, nitro, trifluoromethyl, $-COOH$, $COOalkyl$ wherein alkyl has from 1 to 4 carbon atoms, or $-(CH_2)_pNR_3R_4$ wherein P , R_3 and R_4 have the meanings defined above.

57. A compound when used according to Claim 47 wherein

X is oxygen, sulfur or $(CR'R'')_n$;

Y is oxygen, sulfur or $(CR'R'')_n$ with the proviso that at least one of X or Y is $(CR'R'')_n$ wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, straight or branched alkyl of from 1 to 6 carbons,



19 SEP 2000

optionally substituted phenyl, halogen, hydroxy, alkoxy, acyloxy, cycloalkyl, or R' and R'' taken together form a carbonyl or a spirocycloalkyl group of from 3 to 10 carbons;

R is hydrogen;

R₁ is phenyl optionally substituted, straight or branched alkyl of from 1 to 10 carbon atoms, cycloalkyl of from 3 to 10 carbon atoms;

R₂ is phenyl optionally substituted, straight or branched alkyl of from 1 to 10 carbon atoms, cycloalkyl of from 3 to 8 carbon atoms, phenoxy optionally substituted with the proviso that only if X is (CR'R'')_n can R₁ be optionally substituted phenoxy and only if Y is (CR'R'')_n can be optionally substituted phenoxy, and with the further proviso that at least one of R₁ and R₂ is optionally substituted phenyl or phenoxy.

58. A compound when used according to Claim 47 wherein

X is oxygen;

Y is (CR'R'')_n wherein n is an integer of from 1 to 2;

R is hydrogen;

R₁ is optionally substituted phenyl;

R₂ is optionally substituted phenyl or phenoxy, straight or branched alkyl of from 1 to 10 carbons, or cycloalkyl of from 3 to 10 carbons;

R' and R'' are each independently hydrogen, straight or branched alkyl of from 1 to 6 carbons, optionally substituted phenyl, halogen, hydroxy, alkoxy, acyloxy, cycloalkyl, or R' and R'' taken together form a carbonyl or a spirocycloalkyl.

59. A compound selected from:

Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester,


Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,


Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [adamantaneacetyl]-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester-sodium salt,



Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,
Sulfamic acid (decanoyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (dodecanoyl)-2,6-bis(1-methylethyl)phenyl ester,
2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]sulfonyl]-benzene-acetamide,
2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]benzene-acetamide-sodium salt,
2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate,
2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate-sodium salt,
Sulfamic acid (1-oxo-3,3-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid trans-[(2-phenylcyclopropyl)carbonyl]-2,6-bis(1-methylethyl)-phenyl ester,
Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2,4,6-trimethylphenyl-(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [3-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-methoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (oxophenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-trifluoromethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclopentylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (diphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,




- Sulfamic acid (triphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [(1-phenylcyclopentyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2-phenylbutyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2,2-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-3-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]-2-propenyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,
Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [[2,6-bis(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and
Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(phenyl)-phenyl ester;


when used in an effective amount for lowering the serum or plasma level of Lp(a) in a mammal.

60. Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)-phenyl ester when used in an effective amount for lowering the serum or plasma level of Lp(a) in a mammal.

61. A compound of formula



68



or a pharmaceutically acceptable salt thereof wherein:

X and Y are selected from oxygen, sulfur and $(\text{CR}'\text{R}'')_n$ wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, alkyl, alkoxy, halogen, hydroxy, acyloxy, cycloalkyl, phenyl optionally substituted or R' and R'' together form a spirocycloalkyl or a carbonyl;

with the proviso at least one of X and Y is $(\text{CR}'\text{R}'')_n$ and with the further proviso when X and Y are both $(\text{CR}'\text{R}'')_n$ and R' and R'' are hydrogen and n is one, R₁ and R₂ are aryl;

R is hydrogen, a straight or branched alkyl of from 1 to 8 carbon atoms or benzyl;

R₁ and R₂ are each independently selected from

(a) phenyl or phenoxy each of which is unsubstituted or is substituted with 1 to 5 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

phenoxy,

hydroxy,

fluorine,

chlorine,

bromine,

nitro,

trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

$-(\text{CH}_2)_p\text{NR}_3\text{R}_4$ wherein p is zero or one, and each of R₃ and R₄ is selected from hydrogen or a straight or branched alkyl group having 1 to 4 carbon atoms;

(b) 1- or 2-naphthyl unsubstituted or substituted with from 1 to 3 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,



19 SEP 2000

hydroxy,

phenoxy,

fluorine,

chlorine,

bromine,

nitro,

trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

-(CH₂)_pNR₃R₄ wherein p, R₃ and R₄ have the meanings defined above;

(c) arylalkyl;

(d) a straight or branched alkyl chain having from 1 to 20 carbon atoms and which is saturated or contains from 1 to 3 double bonds; or


(e) adamantyl or a cycloalkyl group wherein the cycloalkyl moiety has from 3 to 6 carbon atoms;

with the provisos

(i) where X is (CH₂)_n, Y is oxygen, and R₁ is a substituted phenyl, then R₂ is a substituted phenyl;

(ii) where Y is oxygen, X is (CH₂)_n, R₂ is phenyl or naphthyl, then R₁ is not a straight or branched alkyl chain; and

(iii) the following compounds are excluded:

X	Y	R	R ₁	R ₂
CH ₂	O	H	(CH ₂) ₂ CH ₃	Ph
CH ₂	O	H	CH ₃	Ph
CH ₂	O	H		i-Pr


when used in an effective amount for treating peripheral vascular disease.

62. A compound selected from:

Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,



Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [adamantaneacetyl]-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester-sodium salt,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,

Sulfamic acid (decanoyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (dodecanoyl)-2,6-bis(1-methylethyl)phenyl ester,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]sulfonyl]-benzene-acetamide,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]benzene-acetamide-sodium salt,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate-sodium salt,

Sulfamic acid (1-oxo-3,3-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester.

Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,6-dichlorophenyl(acetyl)]2,6-bis(1-methylethyl)phenyl ester.

Sulfamic acid trans-[(2-phenylcyclopropyl)carbonyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,4,6-trimethylphenyl-(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [3-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-methoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (oxophenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,



19 SEP 2000

Sulfamic acid [2-trifluoromethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (cyclopentylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (cyclohexylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (diphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (triphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(1-phenylcyclopentyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2-phenylbutyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (cyclohexylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2,2-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-3-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]-2-propenyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,

Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and

Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(phenyl)-phenyl ester;




19 SEP 2000

when used in an effective amount for treating peripheral vascular disease.

63. Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl-2,6-bis(1-methylethyl)-phenyl ester when used in an effective amount for treating peripheral vascular disease.

64. A compound of formula



or a pharmaceutically acceptable salt thereof wherein:

X and Y are selected from oxygen, sulfur and $(\text{CR}'\text{R}'')_n$ wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, alkyl, alkoxy, halogen, hydroxy, acyloxy, cycloalkyl, phenyl optionally substituted or R' and R'' together form a spirocycloalkyl or a carbonyl;
with the proviso at least one of X and Y is $(\text{CR}'\text{R}'')_n$ and with the further proviso when X and Y are both $(\text{CR}'\text{R}'')_n$ and R' and R'' are hydrogen and n is one, R₁ and R₂ are aryl;

R is hydrogen, a straight or branched alkyl of from 1 to 8 carbon atoms or benzyl;

R₁ and R₂ are each independently selected from

(a) phenyl or phenoxy each of which is unsubstituted or is substituted with 1 to 5 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

phenoxy,

hydroxy,

fluorine,

chlorine,

bromine,

nitro,

trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

-(CH₂)_pNR₃R₄ wherein p is zero or one, and each of R₃ and R₄ is selected from hydrogen or a straight or branched alkyl group having 1 to 4 carbon atoms;



(b) 1- or 2-naphthyl unsubstituted or substituted with from 1 to 3 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

hydroxy,

phenoxy,

fluorine,

chlorine,

bromine,

nitro,

trifluoromethyl,

-COOH,

-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,

-(CH₂)_pNR₃R₄ wherein p, R₃ and R₄ have the meanings defined above;

(c) arylalkyl;

(d) a straight or branched alkyl chain having from 1 to 20 carbon atoms and which is saturated or contains from 1 to 3 double bonds; or


(e) adamantyl or a cycloalkyl group wherein the cycloalkyl moiety has from 3 to 6 carbon atoms;

with the provisos

(i) where X is (CH₂)_n, Y is oxygen, and R₁ is a substituted phenyl. then R₂ is a substituted phenyl;

(ii) where Y is oxygen, X is (CH₂)_n, R₂ is phenyl or naphthyl, then R₁ is not a straight or branched alkyl chain; and

(iii) the following compounds are excluded:

X	Y	R	R ₁	R ₂
CH ₂	O	H	(CH ₂) ₂ CH ₃	Ph
CH ₂	O	H	CH ₃	Ph
CH ₂	O	H		i-Pr

when used in a therapeutically effective amount for treating restenosis.

65. A compound selected from:

Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester.



Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [adamantaneacetyl]-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester-sodium salt,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,

Sulfamic acid (decanoyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (dodecanoyl)-2,6-bis(1-methylethyl)phenyl ester,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]sulfonyl]-benzene-acetamide,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]benzene-acetamide-sodium salt,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate-sodium salt,

Sulfamic acid (1-oxo-3,3-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,6-dichlorophenyl(acetyl)]2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid trans-[(2-phenylcyclopropyl)carbonyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,4,6-trimethylphenyl-(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,



- Sulfamic acid [3-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-methoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (oxophenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [2-trifluoromethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclopentylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (diphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (triphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [(1-phenylcyclopentyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2-phenylbutyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (cyclohexylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-2,2-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (1-oxo-3-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]-2-propenyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,
Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,
Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,



* C L M S *

76

REPLACEMENT SHEET

19 SEP 2000


Sulfamic acid [[2,6-bis(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and

Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyllacetyl]-2,6-bis(phenyl)-phenyl ester;

when used in a therapeutically effective amount for treating restenosis

66. Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)-phenyl ester when used in a therapeutically effective amount for treating restenosis

67. A compound of formula



or a pharmaceutically acceptable salt thereof wherein:

X and Y are selected from oxygen, sulfur and $(\text{CR}'\text{R}'')_n$ wherein n is an integer of from 1 to 4 and R' and R'' are each independently hydrogen, alkyl, alkoxy, halogen, hydroxy, acyloxy, cycloalkyl, phenyl optionally substituted or R' and R'' together form a spirocycloalkyl or a carbonyl;

with the proviso at least one of X and Y is $(\text{CR}'\text{R}'')_n$ and with the further proviso when X and Y are both $(\text{CR}'\text{R}'')_n$ and R' and R'' are hydrogen and n is one, R₁ and R₂ are aryl;

R is hydrogen, a straight or branched alkyl of from 1 to 8 carbon atoms or benzyl;

R₁ and R₂ are each independently selected from

(a) phenyl or phenoxy each of which is unsubstituted or is substituted with 1 to 5 substituents selected from

phenyl,

an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,

an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,

phenoxy,

hydroxy,

fluorine,

chlorine,

bromine,

nitro,


trifluoromethyl,

-COOH,




* C L M S *

.7



- COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,
-(CH₂)_pNR₃R₄ wherein p is zero or one, and each of R₃ and R₄ is selected from hydrogen or a straight or branched alkyl group having 1 to 4 carbon atoms;
(b) 1- or 2-naphthyl unsubstituted or substituted with from 1 to 3 substituents selected from
phenyl,
an alkyl group having from 1 to 6 carbon atoms and which is straight or branched,
an alkoxy group having from 1 to 6 carbon atoms and which is straight or branched,
hydroxy,
phenoxy,
fluorine,
chlorine,
bromine,
nitro,
trifluoromethyl,
-COOH,
-COOalkyl wherein alkyl has from 1 to 4 carbon atoms and is straight or branched,
-(CH₂)_pNR₃R₄ wherein p, R₃ and R₄ have the meanings defined above;
(c) arylalkyl;
(d) a straight or branched alkyl chain having from 1 to 20 carbon atoms and which is saturated or contains from 1 to 3 double bonds; or
(e) adamantyl or a cycloalkyl group wherein the cycloalkyl moiety has from 3 to 6 carbon atoms;
with the provisos
(i) where X is (CH₂)_n, Y is oxygen, and R₁ is a substituted phenyl, then R₂ is a substituted phenyl;
(ii) where Y is oxygen, X is (CH₂)_n, R₂ is phenyl or naphthyl, then R₁ is not a straight or branched alkyl chain; and
(iii) the following compounds are excluded:

X	Y	R	R ₁	R ₂
CH ₂	O	H	(CH ₂) ₂ CH ₃	Ph
CH ₂	O	H	CH ₃	Ph
CH ₂	O	H		i-Pr



* C L M S *

78

REPLACEMENT SHEET

19 SEP 2000

when used in a therapeutically effective amount for treating cerebrovascular disease.

68. A compound selected from:

Sulfamic acid (phenylacetyl)-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid[[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,4,6-tris(1-methylethyl)phenyl ester,

Sulfamic acid [adamantaneacetyl]-2,6-bis-(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)phenyl]-acetyl]-2,6-bis(1-methylethyl)-phenyl ester-sodium salt,

Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester-sodium salt,

Sulfamic acid (decanoyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (dodecanoyl)-2,6-bis(1-methylethyl)phenyl ester,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]sulfonyl]-benzene-acetamide,

2,6-Bis(1-methylethyl)-N-[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]benzene-acetamide-sodium salt,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate,

2,6-Bis(1-methylethyl)phenyl[[[2,4,6-tris(1-methylethyl)phenyl]methyl]-sulfonyl]carbamate-sodium salt,

Sulfamic acid (1-oxo-3,3-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,6-dichlorophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,6-dichlorophenyl(acetyl)]2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid trans-[(2-phenylcyclopropyl)carbonyl]-2,6-bis(1-methylethyl)-phenyl ester,

Sulfamic acid [2,5-dimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2,4,6-trimethoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

19 SEP 2000



79

Sulfamic acid [2,4,6-trimethylphenyl-(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [3-thiophenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-methoxyphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (oxophenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [2-trifluoromethylphenyl(acetyl)]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (cyclopentylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (cyclohexylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (diphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (triphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(1-phenylcyclopentyl)-carbonyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2-phenylbutyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (cyclohexylphenylacetyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-2,2-diphenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(9H-fluoren-9-yl)carbonyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (1-oxo-3-phenylpropyl)-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]-2-propenyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [1-oxo-3-[2,4,6-tris-(1-methylethyl)phenyl]propyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [(acetyloxy)[2,4,6-tris-(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,


Sulfamic acid [hydroxy[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [fluoro[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid (3-methyl-1-oxo-2-phenylpentyl)-2,6-bis(1-methylethyl)phenyl ester sodium salt,



80



Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester,

Sulfamic acid [[2,6-bis(1-methylethyl)-phenoxy]acetyl]-2,6-bis(1-methylethyl)phenyl ester, and

Sulfamic acid [[2,4,6-tris(1-methylethyl)-phenyl]acetyl]-2,6-bis(phenyl)-phenyl ester;

when used in a therapeutically effective amount for treating cerebrovascular disease

69. Sulfamic acid [[2,4,6-tris(1-methylethyl)phenyl]acetyl]-2,6-bis(1-methylethyl)-phenyl ester when used in a therapeutically effective amount for treating cerebrovascular disease.

9801859-1

1P053 copy.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 31/325, 31/18	A1	(11) International Publication Number: ✓ WO 97/05868 (43) International Publication Date: 20 February 1997 (20.02.97)
(21) International Application Number: ✓ PCT/US96/11366 (22) International Filing Date: ✓ 8 July 1996 (08.07.96) (30) Priority Data: ✓ 60/003,031 <i>see PD ...</i> ✓ 3/8/95 ✓ 4 August 1995 (04.08.95) ✓ US (71) Applicant (for all designated States except US): WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US). ✓ (72) Inventor; and (75) Inventor/Applicant (for US only): KRAUSE, Brian Robert [US/US]; 1984 Encore Court, Ann Arbor, MI 48103 (US). ✓ (74) Agents: RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al.		(81) Designated States: AU, BG, CA, CN, CZ, EE , GE, HU, IL, JP, KR, LT, LV, MX, NO, NZ, PL, RO, SG , SI, SK, UA, US, UZ, Eurasian patent (AM, AZ, BY, KG , KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: USE OF SULFAMIC ACID DERIVATIVES, ACYL SULFONAMIDES OR SULFONYL CARBAMATES FOR THE MANUFACTURE OF A MEDICAMENT FOR LOWERING LIPOPROTEIN LEVELS ✓ (57) Abstract The present invention is directed to new therapeutic uses of compounds of formula (I) wherein X and Y are oxygen, sulfur, or (CR'R'')_n wherein n is 1 to 4; R is hydrogen, alkyl, or benzyl; R₁ and R₂ are phenyl, substituted phenyl, naphthyl, substituted naphthyl, an aralkyl group, an alkyl chain, adamantyl, or a cycloalkyl group. The uses are for cerebrovascular disease such as stroke, peripheral vascular diseases, and restenosis. $\begin{array}{c} \text{O} \qquad \text{O} \\ \qquad \\ \text{R}_1 - \text{X} - \text{S} - \text{N} - \text{C} - \text{Y} - \text{R}_2 \\ \qquad \\ \text{O} \quad \text{R} \end{array} \quad (I)$		