

DESCRIPTION

BENZO OXYGEN-CONTAINING HETEROCYCLIC COMPOUNDS AND MEDICAL APPLICATION THEREOF

5

TECHNOLOGY FIELD OF THE INVENTION

The present invention relates to a new benzo oxygen-containing heterocyclic compounds, especially for a benzo oxygen-containing five-membered heterocyclic compounds. The described compounds can be used as agonists of the GPR40 target to
10 lower the blood glucose levels by stimulating the release of insulin from pancreatic β cells for the safe and effective treatment of diseases such as type II diabetes.


BACKGROUND OF THE INVENTION

Diabetes is a chronic endocrine and metabolic disease characterised by
15 hyperglycaemia due to defective insulin secretion, insulin resistance or both. Diabetes can severely damage all major organ systems in the body, causing heart disease, stroke, nerve damage, kidney failure, blindness and infections that may lead to amputation, and the disease is associated with numerous complications and high rates of disability and mortality.

20 Type II diabetes accounts for about 90% of all cases of diabetes and patients are usually able to produce their own insulin, but are unable to produce enough or to utilise it properly. The medications commonly used clinically to treat diabetes include insulin stimulating agents which are currently the first line hypoglycemic agents, such as sulphonylureas such as glipizide and non-sulphonylurea drugs such as metformin. The
25 main side effects common to glucose-lowering drugs are gastrointestinal discomfort, oedema and hypoglycaemia. Some patients may suffer from a strong feeling of fasting, cold sweats, general weakness, palpitations, trembling hands and feet, blurred eyes, headaches, dizziness and, in severe cases, coma due to the side effects of hypoglycaemia. Therefore, the development of new drugs against type II diabetes that are safe and effective,
30 free of hypoglycaemic side effects, easy to take orally is still the direction scientists are trying to explore.

Recent studies have identified free fatty acid receptor 1 (FFAR1), or G

protein-coupled receptor 40 (GPR40), as playing a key role in the stimulation and regulation of insulin production. When glucose and fatty acids are elevated in the blood after a meal, GPR40 agonists lower blood glucose levels by stimulating insulin release from pancreatic beta cells. Drugs that activate GPR40 are therefore effective in controlling blood glucose levels by helping diabetics to release more insulin. These drugs have the characteristic of promoting insulin secretion only at high blood glucose levels, thus greatly reducing the risk of hypoglycaemia. A representative of this group is TAK-875, a selective GPR40 agonist developed by Takeda, Japan, which has entered phase III clinical trials and has shown no effect on insulin secretion when blood glucose levels are normal. Clinically, the dose of 50 mg administered once daily was more effective and well tolerated by patients and its risk of causing hypoglycaemia was significantly lower than that of the control sulphonylureas, verifying that TAK-875-induced insulin secretion is glucose-dependent. Although TAK-875 has shown good efficacy, Takeda Japan terminated its clinical studies in 2013 due to problems with hepatotoxicity from the drug. It is worth mentioning that studies on the hepatotoxicity of TAK-875 have shown that the problem is not related to the mechanism of action of the drug, but is mainly a problem caused by the lack of safety in the design of its molecular structure. Therefore, this GPR40 target is still a good development idea with very important application prospects.



Structure of the compound TAK-875

A series of patent applications for GPR40 agonists are currently disclosed, including W02005087710, W02007106469A1, W02004106276A1, W02010143733A1, CN103030646A1, WO2013104257A1, WO2015062486A1 and others.


Thus, despite the advances that have been made in the field, there remains a critical need in the field for GPR40 agonists that specifically activate GPR40 targets for the treatment of diabetes and related metabolic diseases, as well as associated compositions and methods of disease treatment. The present invention is dedicated to meeting that need and providing other related benefits.

SUMMARY OF THE INVENTION

The present invention relates to compounds that activate the GPR40 target, as well as cis-trans isomers, enantiomers, diastereoisomers, racemates, tautomer, solvate, hydrates, or pharmaceutically acceptable salts of said compounds or mixtures thereof. The invention
5 also relates to pharmaceutically acceptable compositions comprising said compounds, and related methods for the treatment of conditions that derive beneficial effects from activation of the GPR40 target, such as diabetes and related metabolic diseases.

The present invention provides a benzo oxygen-containing heterocyclic compound with a different structure from the existing ones. The benzo oxygen-containing
10 five-membered heterocyclic compound of the present invention has better inhibitory activity against type II diabetes and can be used more effectively in the treatment of patients with type II diabetes and has a better safety profile.

In the first aspect, the present invention provides a compound represented by formula **Ib**, its cis-trans isomer, enantiomer, diastereoisomer, racemate, tautomer, solvate, hydrate,
15 or pharmaceutically acceptable salt, or mixtures thereof;



wherein,

$n = 0, 1$ or 2 ; $n = 0, 1$ or 2 ;

When $n = 0$, Y does not exist and Y^1 is directly single bonded to Z^1 adjacent (ortho-) to Y to form a five-membered heterocyclic group;

When E is not directly linked to G^1 to form a cyclic compound, E is selected from: hydrogen, deuterium(D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, carboxyl, amino(NH_2), aminocarbonyl (H_2NCO), alkyl, alkoxy, alkoxycarbonyl,
25 alkylcarbonylamino, alkylaminocarbonyl, aryl, aryloxy, or heterocyclic aryl; G^1 is CH, or C(Rb), wherein Rb is hydrogen, deuterium(D), alkyl, cycloalkyl, alkenyl, alkynyl alkoxy, cycloalkoxy, alkoxycarbonyl, alkylaminyl, cycloalkylaminyl, alkylaminylcarbonyl, cycloalkylaminylcarbonyl, aryl, aryloxy, heterocyclic, heterocyclic aryl, or heterocyclic aryloxy, or Rb and R^4 may be interconnected to form a cycloalkyl, or heterocyclic group;

When E is linked to G1 to form a cyclic compound, G¹ is -C-; E is -O-, -C(RcRd)-, -OC(RcRd)-, -C(RcRd)O-, or -NRe-, wherein Rc and Rd are hydrogen, deuterium(D), alkyl, cycloalkyl, alkenyl, alkyl, alkoxy, cycloalkoxy, alkoxycarbonyl, alkylamino, cycloalkylamino, alkylaminocarbonyl, cycloalkylaminocarbonyl, aryl, aryloxy heterocyclic, heterocyclic aryl, or heterocyclic aryloxy, Rc and Rd may be interconnected to form a cycloalkyl, or heterocyclic group; Re is hydrogen, deuterium(D), alkyl, cycloalkyl, alkylcarbonyl, alkoxycarbonyl, cycloalkoxycarbonyl, alkylaminocarbonyl, cycloalkylaminocarbonyl, alkylsulfonyl, or aryl sulfonyl group.

L¹ and L² are each independently -O-, -S-, -C(O)-, -S(O)₂-, -CH₂-, -C(RfRg)-, -OC(RfRg)-, -C(RfRg)O-, -N(Re)-, -N(Rc)C(RfRg)-, or -C(RfRg)N(Re)-; wherein Rf and Rg are hydrogen, deuterium(D), alkyl, cycloalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, alkoxycarbonyl, alkylaminyl, cycloalkylaminyl, alkylaminylcarbonyl, cycloalkylaminylcarbonyl, aryl, aryloxy, heterocyclic, heterocyclic aryl, or heterocyclic aryloxy, respectively, and Re is defined as the same as that described for Re in E above;

R¹, R² and R³ are each independently hydrogen, deuterium(D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, aminocarbonyl (H₂NCO), alkyl, alkoxy, alkoxycarbonyl, alkylaminocarbonyl, alkylcarbonylamino, aryl, aryloxy, or heterocyclic aryl group; wherein R² and L¹ in formula **1b** may be interconnected to form a 4~8-membered heterocyclic compound, and L¹ is -CH-.

R⁴, R⁵ and R^{5b} are each independently hydrogen, deuterium(D), halogen, hydroxyl, amino, nitrile, alkyl, alkoxy, cycloalkyl, heterocycloalkyl, cycloalkoxy, optionally substituted alkenyl, optionally substituted alkynyl, alkoxycarbonyl, alkylaminocarbonyl, alkylcarbonylamino, alkoxycarbonylamino, aryl, aryloxy, or heterocycloaryl; wherein R⁵ and R^{5b} may be interconnected as cycloalkyl, heterocycloalkyl, or heterocycloaryl group;

R⁶ is a carboxyl group, an alkoxycarbonyl group, an aryloxycarbonyl group, an alkylaminocarbonyl group, a cycloalkylaminocarbonyl group, an alkylsulfonamido carbonyl group, a cycloalkylsulfonamido carbonyl group, a heterocyclic group, or a heterocyclic aryl group; or R⁶ and the adjacent (ortho-) substituent R⁵ may be interconnected to become a heterocyclic group, or a heterocyclic aryl group.


X⁵, X⁶ and X⁷ are each independently hydrogen, deuterium(D), halogen, nitrile, amino, trifluoromethyl, trifluoromethoxy, aminocarbonyl (H₂NCO), alkyl, heterocycloalkyl, alkoxy, heteroatomically substituted alkoxy, alkylaminyl (NR_iR_j), heteroatomically substituted alkylaminyl, alkoxycarbonyl, alkylaminocarbonyl,

alkylcarbonylaminy, alkoxy carbonylaminy, cyclo alkoxy carbonylamino group, alkyl sulfonamido group, cycloalkyl sulfonamido group, aryl group, aryloxy group, aryl amido carbonyl group, aryl carbonyl amido group, aryloxy carbonyl amido group, heterocyclic aryl group, heterocyclic aryloxy group, or heterocyclic aryl amido group, wherein R_i and R_j are each independently hydrogen, deuterium(D), alkyl, heterocycloalkyl, alkylcarbonyl, alkoxy carbonyl, cycloalkoxy carbonyl, alkylaminocarbonyl, alkylsulfonyl, cycloalkylsulfonyl, aryl, aryloxy carbonyl, arylaminocarbonyl, heterocycloaryl group, or R_i and R_j are interconnected to form a 3-8-membered heterocyclic group containing 1-3 heteroatoms;

Y and Y^1 are each independently -O-, -S-, -CH₂-, -CHF-, -CF₂-, -CCl₂-, -C(R_f R_g)-, or -N(Re)-; wherein R_f and R_g are defined as described above for R_f and R_g in L¹, respectively, and Re is defined as described above for Re in E.

Z and Z¹ are each independently selected from: -O-, -S-, -CF₂-, -CHF-, -CF₂-, -C(R_f R_g)-, -N(Re)-, or -C(O)-; wherein R_f and R_g are defined as described above for R_f and R_g in L¹, respectively, and Re is defined as described above for Re in E.

In the second aspect, the present invention provides a compound of formula **IIb**:



wherein;

n, E, G¹, L¹, R¹, R², R³, R⁴, R⁵, R^{5b}, X⁵, X⁶, X⁷, Y, Y¹, Z have the same definitions as n, E, G¹, L¹, R¹, R², R³, R⁴, R⁵, R^{5b}, X⁵, X⁶, X⁷, Y, Y¹, Z in claim 1.

R⁷ is hydroxy, alkoxy, alkylamino, cycloalkylamino, heterocyclicamino, alkylsulfonamido, cycloalkylsulfonamido, aryloxy, heterocyclic aryloxy, arylamido, or heterocyclic arylamido; or R⁷ and the adjacent (ortho-) substituent R⁵ may be interconnected to form a heterocyclic group.

In some preferred embodiments, n = 0 or 1;

when n = 0, Y does not exist and Y¹ is directly single bonded to the oxygen adjacent (ortho-) to Y to form a five-membered heterocyclic group;

When n = 1, Y is -CH₂-.

When E is not directly linked to G¹ to form a cyclic compound, E is hydrogen, halogen, trifluoromethyl, trifluoromethoxy, or alkoxy; G¹ is -CH-, or -C(Rb)-, wherein Rb is hydrogen, an alkyl group, an optionally substituted alkenyl group, an optionally substituted alkynyl group, an alkoxy group, or Rb and R⁴ are interconnected to form an oxygen-containing heterocyclic group; R⁴ is hydrogen, an alkyl group, an alkoxy group, an optionally substituted alkynyl group, or R⁴ and Rb are interconnected to form an oxygen-containing heterocyclic group.

When E is directly linked to G¹ to form a cyclic compound, E is -OC(RcRd)-, G¹ is -C-, and R⁴ is hydrogen, wherein Rc and Rd are each independently hydrogen;

L¹ is -CH₂-, or when R² and L¹ in formula **I**ib**** are interconnected to form a 5-6-membered heterocyclic compound, L¹ is -OCH-;

R¹, R² and R³ are each independently hydrogen, halogen, alkyl or alkoxy;

R⁵ is hydrogen, halogen, hydroxyl, amino, alkylamino, alkyl or alkoxy;

R^{5b} is hydrogen, halogen, alkyl or alkoxy;

R⁷ is an alkoxy, hydroxy, alkylsulfonamido, or cycloalkylsulfonamido group;

X⁵ is hydrogen, deuterium(D), halogen, nitrile, amino, trifluoromethyl, trifluoromethoxy, alkyl, alkoxy, alkylamino (NR_iR_j), alkylcarbonylamino, alkylaminocarbonylamino, alkoxy carbonylamino, cycloalkoxy carbonylamino, alkylsulfonamido, cycloalkylsulfonamido, aryl, or aryloxy carbonylamino; wherein R_i and R_j are each independently hydrogen, alkyl, alkylcarbonyl, alkoxy carbonyl, or cycloalkoxy carbonyl group;

X⁶ and X⁷ are each independently hydrogen, deuterium(D), halogen, C₁-C₈ alkylamino group or C₁-C₈ alkoxy carbonylamino group;

Y¹ is -CH₂-, -CHF-, -CH₂-, or -C(CH₃)₂-;

Z is -O-, or -CH₂-.

In some more preferred embodiments, n = 0 and Y does not exist;

When E is not directly connected to G¹ to form a cyclic compound, E is hydrogen, or halogen; G¹ is -CH-, or -C(Rb)-, wherein Rb is alkyl, alkenyl, alkynyl or alkoxy; and R⁴ is hydrogen, alkyl or alkoxy group;

E is directly linked to G¹ in a cyclic compound, E is -OC(RcRd)-, G¹ is -C- and R⁴ is hydrogen, where Rc and Rd are each independently hydrogen;

L¹ is -CH₂-, or L¹ is -OCH- when R² and L¹ in formula **I**ib**** are interconnected to form a 5-6-membered heterocyclic compound;

R^1 , R^2 and R^3 are each independently hydrogen, halogen, or alkoxy group;

R^5 is hydrogen, halogen, hydroxyl, amino, alkyl or alkoxy group;

R^{5b} is hydrogen, halogen, alkyl or alkoxy group;

R^7 is selected from: alkoxy, hydroxy, alkylsulfonamido, or cycloalkylsulfonamido

5 group;

X^5 is hydrogen, deuterium(D), halogen, nitrile, amino, trifluoromethyl, trifluoromethoxy, alkyl, alkoxy, alkylamino (NR_iR_j), alkylcarbonylamino, alkylaminocarbonylamino, alkoxy carbonylamino, cycloalkoxy carbonylamino, alkylsulfonamido, cycloalkylsulfonamido, aryl, or aryloxy carbonylamino; wherein R_i and

10 R_j are each independently hydrogen, alkyl, alkylcarbonyl, alkoxy carbonyl, or cycloalkoxy carbonyl group;


X^6 is hydrogen, deuterium(D), halogen, C_1 - C_8 alkylamino group or C_1 - C_8 alkoxy carbonylamino group;

X^7 is hydrogen;


15 Y^1 is $-CH_2-$, $-CHF-$, $-CF_2-$, or $-C(CH_3)_2-$;

Z is $-O-$.


In some specific embodiments, the compound is represented by one of the following structures:




20 **IIb-18**



IIb-25



IIb-43



IIb-46.

Another aspect of the invention provides a composition comprising (i) a therapeutically effective amount of at least one compound of formula Ib or IIb, its cis-trans isomer, enantiomer, diastereoisomer, racemate, tautomer, solvate, hydrate, or pharmaceutically acceptable salt or a mixture thereof; and (ii) a pharmaceutically acceptable diluent and/or excipient.

The invention also relates to use of the compounds or compositions as GPR40 agonists.

The invention also provides methods for treating or preventing diabetes or related metabolic syndromes, characterized by comprising administering to a patient a therapeutically effective amount of said compound or composition.

The invention also relates to the use of said compounds or compositions in the preparation of pharmaceuticals for the treatment or prevention of diabetes mellitus or related metabolic syndromes.

In preferred embodiments, the compounds or compositions of the present invention are particularly suitable for the treatment of type II diabetes.

The compounds of the invention promote the secretion of insulin only at high blood glucose concentrations in patients with type II diabetes, thus effectively reduce the risk of hypoglycaemia in patients. At the same time, the compounds of the present invention have a better GPR40 target selectivity and safety profile relative to commercially available drugs.

The foregoing embodiments and other aspects of the present invention will be apparent in the following detailed description. Wherefore, various references are set forth

herein which describe in greater detail certain background information, processes, compounds and/or compositions and are each incorporated herein by reference in their entirety.

DETAILED DESCRIPTION OF THE INVENTIONS

In the following description, a number of specific details are stated in order to provide a thorough understanding of the various embodiments of the invention. However, it should be understood by those skilled in the art that the invention can be implemented without requiring these details. Unless the context indicates otherwise, throughout the specification and claims of the invention, the word "comprising" and variations thereof. For example, the words "comprising" and "including" are understood to be in an open, combinable form (i.e., "including (comprising) but not limited to").

References in the specification to "an embodiment" or "an embodiment" indicate that the specific features, structures or characteristics described together with the embodiment are included in at least one embodiment of the invention. Thus, the phrases "in an embodiment" or "in an embodiment" appearing throughout the specification do not necessarily all relate to the same embodiment. Furthermore, specific features, structures, or properties may be combined in any suitable manner in one or more embodiments.

I. DEFINITIONS

In the present invention, "alkyl", when not specifically designated, means a branched and straight chain saturated aliphatic hydrocarbon group comprising from one to twenty carbon atoms and consisting only of carbon and hydrogen atoms. In preferred embodiments, said alkyl groups have one to twelve carbon atoms (C₁-C₁₂ alkyl), one to eight carbon atoms (C₁-C₈ alkyl) or one to six carbon atoms (C₁-C₆ alkyl) and are attached to the remainder of the molecule by a single bond. Exemplary alkyl groups include: methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl, isobutyl, pentyl, hexyl, heptyl, octyl, and their various isomers, etc. The alkyl groups are optionally substituted with groups selected from the group consisting of: deuterium (D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, carboxyl, amino (NH₂), aminocarbonyl (H₂NCO), alkyl, alkoxy, alkoxycarbonyl, alkylcarbonylamino, alkylaminocarbonyl, cycloalkyl, cycloalkoxy, cycloalkoxycarbonyl, cycloalkylamino, cycloalkylaminocarbonyl, cycloalkenyl, cycloetheryl, heterocyclic, alkylureido, aryl, aryloxy, heterocyclic aryl, heterocyclic aryloxy, fused-ring aryl, fused

hetero-ring aryl, fused-ring oxy, fused-ring aryloxy, fused-ring heterocyclic aryloxy, aryl ureido, or heterocyclic aryl ureido.

In the present invention, when not specifically designated, the term "alkylidene", where "alkylidene" means a second-order group derived from an alkyl group comprising 1 to 20 carbon atoms with one hydrogen atom removed, such as methylene, ethylene, propylidene, etc., is used when not specifically specified.

In the present invention, when not specifically designated, the term "aryl" is meant any stable monocyclic, bicyclic, tricyclic or tetracyclic hydrocarbon ring group comprising up to seven carbon atoms per ring, at least one of which is an aromatic ring. Exemplary aryl groups are hydrocarbon ring system groups comprising hydrogen and 6-9 carbon atoms and at least one aromatic ring; hydrocarbon ring system groups comprising hydrogen and 9-12 carbon atoms and at least one aromatic ring; hydrocarbon ring system groups comprising hydrogen and 12-15 carbon atoms and at least one aromatic ring; or hydrocarbon ring system groups comprising hydrogen and 15-18 carbon atoms and at least one aromatic ring. For the purposes of the present invention, aryl groups may be monocyclic, bicyclic, tricyclic or tetracyclic ring systems, which may include dense or bridged ring systems. Aryl groups include, but are not limited to, aryl groups derived from the following compositions: benzene, biphenyl, anthracene, azulene, fluorene, indene, indene, naphthalene, phenanthrene, pyrene, etc. "optionally substituted aryl" means: an aryl group or a substituted aryl group. The term aryl group may optionally be substituted with a group selected from the group consisting of: deuterium(D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, carboxyl, amino(NH₂), aminocarbonyl (H₂NCO), alkyl, alkoxy, alkoxycarbonyl, alkylcarbonylamino, alkylaminocarbonyl, alkyl-sulfonyl-alkoxy, cycloalkyl, cycloalkoxy, cycloalkoxycarbonyl, cycloalkylamino, cycloalkyl aminocarbonyl, cycloalkenyl, cycloetheryl, heterocyclic, aryl, aryloxy, heterocyclic aryl, fused-ring aryl, fused-ring alkyl, fused-ring oxy, unsubstituted or benzene or biphenyl groups containing 1 to 4 of the above optional substituents.

In the present invention, when not specifically designated, the term "heterocyclic aryl" means a stable monocyclic, bicyclic or tricyclic ring comprising up to 7 atoms each, at least one of which is an aromatic ring, and at least one of which contains 1-4 heteroatoms selected from O, N, and/or S. Heterocyclic aryl groups within this definition include, but are not limited to: acridinyl, carbazolyl, cinnolinyl, quinoxalinyl, pyrazolyl, indolyl, benzotriazolyl, furanyl, thiophenyl, benzothiazolyl, benzothiophenyl, benzofuranyl,

quinolinyl, isoquinolinyl, oxazolyl, isoxazolyl, pyrazinyl, pyridazinyl, pyridinyl, pyrimidinyl, pyrrole, tetrahydroquinoline. In addition, "heterocyclic aryl" should also be understood to include any quaternary ammonium salt or N-oxide derivative of a nitrogen-containing heterocyclic aryl group. Said heterocyclic aryl group may optionally
5 be substituted with a group selected from the group consisting of: deuterium (D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, carboxyl, amino (NH₂), aminocarbonyl (H₂NCO), alkyl, alkoxy, alkoxycarbonyl, alkylcarbonylamino, alkylaminocarbonyl, cycloalkyl, cycloalkoxy, cycloalkoxycarbonyl, cycloalkylamino, cycloalkylaminocarbonyl, cycloalkenyl, cycloalkenyl, cycloetheryl, heterocyclic, heterocyclic, heterocyclic. cyclic
10 ether group, heterocyclic group, aryl group, aryloxy group, heterocyclic aryl group, fused-ring aryl group, fused-ring alkyl group, fused-ring oxy group, unsubstituted or benzene or biphenyl group containing 1 to 4 of the above optional substituents.

In the present invention, when not specifically designated, the term "fused-ring aryl" means a stable bicyclic or tricyclic ring comprising up to 7 atoms each, at least one of
15 which is an aromatic ring. Said fused-ring aryl groups may optionally be substituted with a group selected from the group consisting of: Deuterium (D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, carboxyl, amino (NH₂), aminocarbonyl (H₂NCO), alkyl, alkoxy, alkoxycarbonyl, alkylcarbonylamino, alkylaminocarbonyl, cycloalkyl, cycloalkoxy, cycloalkoxycarbonyl, cycloalkylamino, cycloalkylaminocarbonyl, cycloalkenyl, cycloether,
20 heterocyclic, aryl, aryloxy, heterocyclic aryl, fused-ring aryl, fused-ring alkyl, fused-ring oxy, unsubstituted or benzene or biphenyl groups containing 1 to 4 of the above optional substituents.

In the present invention, when not specifically designated, the term "dense ring heterocyclic aryl" means a stable bicyclic or tricyclic ring comprising up to 7 atoms each,
25 wherein at least one ring is an aromatic ring and contains 1-4 heteroatoms selected from O, N, and/or S. The fused-ring heterocyclic aryl groups may optionally be substituted with a group selected from the group consisting of: Deuterium (D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, carboxyl, amino (NH₂), aminocarbonyl (H₂NCO), alkyl, alkoxy, alkoxycarbonyl, alkylcarbonylamino, alkylaminocarbonyl, cycloalkyl, cycloalkoxy, cycloalkoxycarbonyl, cycloalkylamino, cycloalkylaminocarbonyl, cycloalkenyl, cycloether,
30 heterocyclic, aryl, aryloxy, heterocyclic aryl, fused-ring aryl, fused-ring alkyl, fused-ring oxy, unsubstituted or benzene or biphenyl groups containing 1 to 4 of the above optional substituents.

In the present invention, when not specifically designated, the term "alkoxy" refers to the group formed when an alkyl group is attached to an oxygen atom, i.e. " $R-O-\overset{\text{}}{\underset{\text{}}{\text{Z}}}-$ ", and R is an alkyl group, which is defined as described above for alkyl groups. Examples of alkoxy groups include, but are not limited to: -O-methyl (methoxy), -O-ethyl (ethoxy),
5 -O-propyl (propoxy), -O-isopropyl (isopropoxy), -O-tert-butyl (tert-butoxy), etc.

In the present invention, when not specifically designated, the term "Alkenyl" means unsaturated aliphatic alkenyl containing 1 to 3 "carbon-carbon double bonds" in branched and straight chains of 2 to 20 carbon atoms, preferably 2 to 10 carbon atoms (C_2-C_{10} alkenyl), more preferably 2 to 8 carbon atoms (C_2-C_8 alkenyl) or 2 to 6 carbon atoms
10 (C_2-C_6 alkenyl), and their various isomers, etc., which are attached to the remaining part of the molecule by single bonds. Examples of alkenyl groups include, but are not limited to: vinyl, propylenyl, butenyl, pentenyl, hexenyl, etc. Said alkenyl may optionally be substituted with a group selected from the group consisting of: deuterium (D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, carboxyl, amino (NH_2), aminocarbonyl
15 (H_2NCO), alkyl, alkoxycarbonyl, alkylcarbonylamino, alkylaminocarbonyl, cycloalkyl, cycloalkoxy, cycloalkoxycarbonyl, cycloalkylamino, cycloalkylaminocarbonyl, cycloalkenyl, cycloetheryl, heterocyclic alkyl, alkyl ureido, aryl, heterocyclic aryl, fused-ring aryl, fused-ring heterocyclic aryl, aryl ureido, or heterocyclic aryl ureido.

In the present invention, when not specifically designated, the term "alkynyl" means
20 unsaturated aliphatic alkynyl containing 1 to 2 "carbon-carbon triple bonds" in branched and straight chains including 2 to 20 carbon atoms, preferably 2 to 10 carbon atoms (C_2-C_{10} alkynyl), more preferably 2 to 8 carbon atoms (C_2-C_8 alkynyl) or 2 to 6 carbon atoms (C_2-C_6 alkynyl), and their various isomers, etc., which are connected to the remaining part of the molecule by single bonds. Examples of alkynyl groups include, but
25 are not limited to: ethynyl groups, propargyl groups, butynyl groups, pentynyl groups, hexynyl groups, etc. The alkynyl group may optionally be substituted with a group selected from the group consisting of: deuterium(D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, carboxyl, amino(NH_2), aminocarbonyl (H_2NCO), alkyl, alkoxycarbonyl, alkylcarbonylamino, alkylaminocarbonyl, cycloalkyl, cycloalkoxy, cycloalkoxycarbonyl,
30 cycloalkylamino, cycloalkylaminocarbonyl, cycloalkenyl, cycloether, heterocyclic, alkylureido, aryl, heterocyclic aryl, fused-ring aryl, fused-ring heterocyclic aryl, arylureido, or heterocyclic arylureido.

In the present invention, unless otherwise specified, "alkylthio" denotes the group formed when an alkyl group is attached to a sulfur atom, i.e. " $R-S-\text{---}$ ", and R is an alkyl group, which is defined in the same way as described above for alkyl groups.

In the present invention, when not specifically designated, the term "aryloxy" means
5 the group formed by the attachment of an aryl group to an oxygen atom, i.e. " $Ar-O-$ ", and Ar is an aryl group, which is defined in the same way as described above for aryl groups.

In the present invention, unless otherwise specified, the term "arylamino group" refers to an amino group in which one of the hydrogens in "NH₃" is replaced by an aryl
10 group, where aryl is defined in the same way as described above for aryl.

In the present invention, unless otherwise specified, "heterocyclic aromatic amino group" means an amino group in which one of the hydrogens in "NH₃" is replaced by a heterocyclic aromatic group, where the definition of heterocyclic aromatic group is the same as described above for heterocyclic aromatic groups.

15 In the present invention, when not specifically designated, the term "dense-ring aryl" means an all-carbon monocyclic or polycyclic group, wherein each ring does not contain a double or triple bond. Preferably, the ring has 3 to 20 carbon atoms, has 3 to 15 carbon atoms, preferably has 3 to 10 carbon atoms, 3 to 8 carbon atoms, 3 to 6 carbon atoms, 3 to 5 carbon atoms, a ring with 4 carbon atoms, or a ring with 3 carbon atoms.
20 Examples of cycloalkyl include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, and the like. Said cycloalkyl group may optionally be substituted with a group selected from the group consisting of: deuterium (D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, carboxyl, amino (NH₂), aminocarbonyl (H₂NCO), alkyl, alkoxy, alkoxycarbonyl, alkylcarbonylamino, alkylaminocarbonyl,
25 cycloalkyl, cycloalkoxy, cycloalkoxycarbonyl, cycloalkylamino, cycloalkylaminocarbonyl, cycloalkenyl, cycloalkenyl cyclic ether group, heterocyclic group, alkyl ureido group, aryl group, aryloxy group, heterocyclic aryl group, heterocyclic aryl oxy group, fused-ring aryl group, fused-ring heterocyclic aryl group, fused-ring oxy group, fused-ring aryl oxy group, fused-ring heterocyclic aryl oxy group, aryl ureido group, or heterocyclic aryl ureido group,
30 wherein the definition of aryl group is the same as described above for aryl group.

In the present invention, when not specifically designated, the term "cycloalkenyl" means an all-carbon monocyclic or polycyclic group in which one or each ring may


contain one or more "carbon-carbon double bonds". Preferably, the ring has 3 to 20 carbon atoms, has 3 to 15 carbon atoms, preferably has 3 to 10 carbon atoms, 3 to 8 carbon atoms, 3 to 6 carbon atoms, 3 to 5 carbon atoms, a ring with 4 carbon atoms, or a ring with 3 carbon atoms. Examples of cycloalkenyl groups include, for example: cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl. Said cycloalkenyl may optionally be substituted with a group selected from the group consisting of: deuterium (D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, carboxyl, amino (NH₂), aminocarbonyl (H₂NCO), alkyl, alkoxy, alkoxycarbonyl, alkylcarbonyl, alkylaminocarbonyl, cycloalkyl, cycloalkoxy, cycloalkoxycarbonyl, cycloalkylaminocarbonyl, cycloalkylaminocarbonyl, cycloalkenyl, cycloalkenyl cyclic ether group, heterocyclic group, alkyl ureido group, aryl group, aryloxy group, heterocyclic aryl group, heterocyclic aryl oxy group, fused-ring aryl group, fused-ring heterocyclic aryl group, fused-ring oxy group, fused-ring aryl-oxy group, fused-ring heterocyclic aryl-oxy group, aryl ureido group, or heterocyclic aryl ureido group. Cycloalkyl can be formed when the substituent of the cycloalkenyl group is substituted on the carbon-carbon double bond and saturates the double bond.

In the present invention, when not specifically designated, the term "cycloether group" means to a cycloalkyl group with an ether group on the ring.

In the present invention, when not specifically designated, the term "heterocyclic group" means an aromatic or non-aromatic heterocyclic group containing one or more heteroatoms selected from O, N and S, and includes a bicyclic group. Thus, "heterocyclic group" includes the above-mentioned heterocyclic aromatic groups and their di- or tetrahydro analogs, and includes, but is not limited to, the following "heterocyclic groups": Benzimidazolyl, benzofuranyl, benzopyrazolyl, benzotriazolyl, benzothiazolyl, benzothiophenyl, benzoxazolyl, isobenzofuranyl, pyridinopyridinyl, and heterocyclic groups can be linked to other organic small molecule groups by carbon atoms or heteroatoms to form new compounds with medicinal effects.

In the present invention, when not specifically designated, the term "dense ring aryl" means two or more aryl and/or heterocyclic aryl groups, formed by fused-ring of polycyclic organic compounds, said dense ring aryl groups may also be substituted in a reasonable manner by alkyl, alkoxy, alkylthio, aryloxy, arylamino, heterocyclic, cycloalkyl, cyclic alkoxy, cycloalkoxycarbonyl, cycloalkylaminyl, cycloalkylaminylcarbonyl, cycloalkenyl, cycloetheryl, aryl, halogen, carbonyl, hydroxyl, heterocyclic aryl groups as

defined in the present invention in a reasonable manner-; wherein- includes, but is not limited to, naphthalene, anthracene, quinone, phenanthrene, fluorene, benzimidazolyl,




In the present invention, when not specifically designated, the term "fused cycloalkyl" refers a non-aromatic polycyclic system resulting from the reduction of one or more double bonds in a fused-ring aryl group, where the carbon number is "C₁₀₋₂₀".

In the present invention, when not specifically designated, the term "fused cycloalkyl aryl" refers the group resulting from the substitution of hydrogen on the carbon of the aryl group by a fused cycloalkyl group, where the carbon number is "C₁₅₋₂₀".

In the present invention, when not specifically designated, the term "fused-ring oxy group" refers a group formed by a fused-ring aryl or fused-ring alkyl group attached to oxygen, i.e. "Ar-O-", Ar is C₁₀₋₂₀ fused-ring aryl or fused-ring alkyl group.

In the present invention, when not specifically designated, the term "alkoxycarbonyl" means the group formed when an alkoxy group is attached to a carbonyl group, i.e.



In the present invention, when not specifically designated, the term "aryloxycarbonyl" means the group formed by the attachment of an aryloxy group to a carbonyl group, i.e., "Ar-O-C(=O)-", and Ar is a C₆₋₂₀ aryl group.

In the present invention, when not specifically designated, the term "heterocyclic oxy" indicates the group formed by the heterocyclic group attached to oxygen, i.e. "R-O-", and R is a C₂₋₂₀ heterocyclic group.

In the present invention, when not specifically designated, the term "alkylamino group" means to the group formed when an alkyl group is attached to an amino group, i.e.,

" $\begin{array}{c} \text{R} \\ | \\ \text{N} \\ | \\ \text{H} \end{array}$ ", and R is a C₁₋₂₀ alkyl group.

In the present invention, when not specifically designated, the term "alkylaminocarbonyl" refers to the group formed when an alkylamino group is attached to

a carbonyl group, i.e., " $\begin{array}{c} \text{O} \\ || \\ \text{R}-\text{N} \\ | \\ \text{H} \end{array}$ ", and R is a C₁₋₂₀ alkyl group.

5 In the present invention, when not specifically designated, the term "arylamino group" refers to the group formed when an aryl group is attached to an amino group, i.e.,

" $\begin{array}{c} \text{Ar} \\ | \\ \text{N} \\ | \\ \text{H} \end{array}$ ", and Ar is a C₆₋₂₀ aryl group.

In the present invention, when not specifically designated, the term "heterocyclic amino group" refers to the group formed when the heterocyclic group is attached to the

10 amino group, i.e., " $\begin{array}{c} \text{R} \\ | \\ \text{N} \\ | \\ \text{H} \end{array}$ ", and R is a C₂₋₂₀ heterocyclic group.

In the present invention, when not specifically designated, the term "aryl sulfonyl" indicates the group formed by the attachment of an arylamino group to a sulfonyl group,

i.e., " $\begin{array}{c} \text{O} \\ || \\ \text{Ar}-\text{N} \\ | \\ \text{H} \end{array}$ ", and Ar is a C₆₋₂₀ aryl group.

15 In the present invention, when not specifically designated, the term "alkylaminosulfonyl" refers to the group formed when an alkylamino group is attached to a

sulfonyl group, i.e., " $\begin{array}{c} \text{O} \\ || \\ \text{R}-\text{N} \\ | \\ \text{H} \end{array}$ ", and R is a C₁₋₂₀ alkyl group.

In the present invention, when not specifically designated, the term "heterocyclic amino sulfonyl" refers the group formed by the attachment of a heterocyclic amino group

to a sulfonyl group, i.e., " $\begin{array}{c} \text{O} \\ || \\ \text{R}-\text{N} \\ | \\ \text{H} \end{array}$ ", and R is a C₂₋₂₀ heterocyclic group.

20 In the present invention, when not specifically designated, the term "alkyl sulfonamido group" refers the group formed by the attachment of an alkyl group to a

sulfonamido group, i.e., " $\begin{array}{c} \text{O} \\ || \\ \text{R}-\text{S} \\ | \\ \text{O} \end{array}$ -NH-", and R is a C₁₋₂₀ alkyl group.

In the present invention, when not specifically designated, the term "heterocyclic sulfonamido group" refers the group formed by the attachment of the heterocyclic group to

the sulfonamido group, i.e. " $\text{R}-\text{S}(=\text{O})_2-\text{NH}-$ ", and R is a C₂₋₂₀ heterocyclic group.

In the present invention, when not specifically designated, the term "aryl sulfonamido group" refers to the group formed by the attachment of an aryl group to a sulfonamido

group, i.e., " $\text{Ar}-\text{S}(=\text{O})_2-\text{NH}-$ ", and Ar is a C₆₋₂₀ aryl group.

5 In the present invention, when not specifically designated, the term "alkylamino sulfonamido group" refers to the group formed when an alkylamino group is attached to a

sulfonamido group, i.e., " $\text{R}-\text{NH}-\text{S}(=\text{O})_2-\text{NH}-$ ", and R is a C₁₋₂₀ alkyl group.

In the present invention, when not specifically designated, the term "alkyl carbonyl amino group" refers to a group formed by linking an alkyl group to a carbonyl group and

10 then to an amino group, i.e. " $\text{R}-\text{C}(=\text{O})-\text{NH}-$ ", with R being a C₁₋₂₀ alkyl group.

In the present invention, when not specifically designated, the term "alkyl ureido" refers to the group formed by the attachment of an alkyl group to a ureido group, i.e. "

$\text{R}-\text{NH}-\text{C}(=\text{O})-\text{NH}-$ ", with R being a C₁₋₂₀ alkyl group.

15 In the present invention, when not specifically designated, the term "aryl ureido" refers to the group formed by joining an aryl group with a ureido group, i.e., "

$\text{Ar}-\text{NH}-\text{C}(=\text{O})-\text{NH}-$ ", with Ar being a C₆₋₂₀ aryl group.

In the present invention, when not specifically designated, the term "alkylthiourea group" refers to the group formed by joining an alkyl group with a thiourea group, i.e.,

" $\text{R}-\text{NH}-\text{C}(=\text{S})-\text{NH}-$ ", with R being a C₁₋₂₀ alkyl group.

20 In the present invention, the term "halogen" refers to "fluorine, chlorine, bromine, or iodine atom".

In the present invention, the term "hydroxyl" refers to $-\text{OH}$.

In the present invention, the term "amino group" refers to $-\text{NH}_2$.

In the present invention, the term "cyano" refers to $-\text{CN}$.

In the present invention, the term "carboxyl" refers to $-\overset{\sim}{\text{C}}\text{OOH}$.

In the present invention, the term "sulfonyl" refers to $-\overset{\sim}{\text{S}}(\text{O})_2-\overset{\sim}{\text{C}}-$.

In the present invention, the term "Sulfonamido" refers to $-\overset{\sim}{\text{S}}(\text{O})_2\text{NH}-\overset{\sim}{\text{C}}-$.

In the present invention, the term "Carbonyl" refers to $-\overset{\sim}{\text{C}}(\text{O})-\overset{\sim}{\text{C}}-$.


5 In the present invention, the term "Urea-based" refers to $-\overset{\sim}{\text{N}}(\text{H})-\overset{\sim}{\text{C}}(\text{O})-\overset{\sim}{\text{N}}(\text{H})-\overset{\sim}{\text{C}}-$.

In the present invention, the term "Thiourea-based" refers to $-\overset{\sim}{\text{N}}(\text{H})-\overset{\sim}{\text{C}}(\text{S})=\overset{\sim}{\text{N}}(\text{H})-\overset{\sim}{\text{C}}-$.

The term "halogen" refers to fluorine, chlorine, bromine and iodine.


II. COMPOUNDS OF THE PRESENT INVENTION

10 The present invention was innovatively designed to introduce a heterocyclic functional group containing the following novel benzo-oxo-containing heterocyclic



(where R^8 is a halogen, hydroxyl, amino, carboxyl, alkyl sulfonyloxy, aryl sulfonyloxy, or a leaving group that can be substituted), and to synthesize a class of benzo oxygen-containing five-membered heterocyclic compounds that are effective as novel GPR40 target agonists for the treatment of type II diabetes.

The present invention relates generally to compounds covered by formula Ib, their cis-trans isomers, enantiomers, diastereoisomers, racemates, tautomer, solvate, hydrates, or pharmaceutically acceptable salts, or mixtures thereof:



Ib

For compounds of formula Ib, n, E, G^1 , L^1 , L^2 , R^1 , R^2 , R^3 , R^4 , R^5 , R^{5b} , R^6 , R^a , R^b , R^c ,

Rd, Re, Rf, Rg, Ri, Rj, X⁵, X⁶, X⁷, Y, Y¹, Z and Z¹ are as defined in the specification.

Specific embodiments of compounds of Formula Ib are also described below.

In one embodiment, n = 0, Y does not exist, and Y¹ is directly connected to Z¹ with a single bond.

5 In one embodiment, E is linked to G¹ to form a cyclic compound. In a preferred embodiment, E is linked to G¹ as a five-membered cyclic compound. In a preferred embodiment, G¹ is -C- and E is -O-, -C(RcRd)-, -OC(RcRd)-, or -C(RcRd)O-. In one embodiment, Rc and Rd are independently selected from: Hydrogen, deuterium(D), alkyl, cycloalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, alkoxycarbonyl, alkylaminyl, cycloalkylaminyl, alkylaminylcarbonyl, cycloalkylaminylcarbonyl, aryl, aryloxy, heterocyclic, heterocyclic aryl, or heterocyclic aryloxy. In another embodiment, Rc and Rd may be connected to each other as a cycloalkyl or heterocyclic group. In a preferred embodiment, G¹ is -C- and E is -OC(RcRd)- or -C(RcRd)O-, wherein Rc and Rd are independently selected from: hydrogen, deuterium (D), C₁-C₈ alkyl, C₃-C₈ cycloalkyl, C₂-C₈ alkenyl, C₂-C₈alkynyl, C₁-C₈ alkoxy, C₃-C₈cycloalkoxy, C₁-C₈ alkoxycarbonyl, C₁-C₈ alkylamino, C₃-C₈ cycloalkylamino, C₁-C₈ alkylamino carbonyl, C₃-C₈ cycloalkylamino carbonyl, aryl, aryloxy, heterocyclic, heterocyclic aryl, heterocyclic, heterocyclic aryl, or heterocyclic aryloxy. In a more preferred embodiment, G¹ is -C- and E is -OC(RcRd)- or -C(RcRd)O-, wherein Rc and Rd are hydrogen, respectively.

20 In one embodiment, L¹ and L² are each independently selected from: -O-, -S-, -C(O)-, -SO₂-, -CH₂-, -C(RfRg)-, -OC(RfRg)-, -C(RfRg)O-, -N(Re)-, -N(Rc)C(RfRg)-, or -C(RfRg)N(Re)-. In a preferred embodiment, one of L¹ and L² is -CH₂- and the other is -O-. In another preferred embodiment, R₂ and L¹ in Formula Ib are interconnected into a 5-6-membered heterocyclic compound, with at least one of L¹ and L² being -OCH-.

25 In one embodiment, R¹, R², and R³ are each independently hydrogen, deuterium(D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxy, aminocarbonyl (H₂NCO), C₁-C₈ alkyl, C₁-C₈ alkoxy, C₁-C₈ alkoxycarbonyl, C₁-C₈alkylaminocarbonyl, C₁-C₈ alkylcarbonylamino, aryl, aryloxy, or heterocyclic aryl. In preferred embodiments, R¹, R², and R³ are each independently hydrogen, halogen, or C₁-C₈ alkoxy group.

30 In one embodiment, R⁴ and R⁵ are each independently selected from: hydrogen, halogen, hydroxyl, amino, or alkoxy. In preferred embodiments, R⁴ and R⁵ are each independently hydrogen, halogen, or C₁-C₈ alkoxy group.

In one embodiment, R⁶ is -COR⁷. In a preferred embodiment, R⁷ is -OH, an alkoxy,

an alkyl amino, a cycloalkyl amino, a heterocyclic amino, an alkyl sulfonamido, a cycloalkyl sulfonamido, an aryloxy, a heterocyclic aryloxy, an aryl amino, or a heterocyclic aryl amino group; or R^7 and a adjacent (ortho-) substituent R^5 may be interconnected to become a heterocyclic group. In a preferred embodiment, R^7 is a C_1 - C_8 alkoxy, hydroxy,
5 C_1 - C_8 alkyl sulfonamido, or C_3 - C_8 cycloalkyl sulfonamido group.

In one embodiment, Y^1 is selected from: -oxygen-, -sulfur-, $-CH_2-$, $-CHF-$, $-CF_2-$, $-CCl_2-$, , $-C(R_fR_g)-$, or $-N(Re)-$. In a preferred embodiment, Y^1 is selected from: CH_2 , $-CHF-$, CF_2 , or $C(CH_3)_2$.

In one embodiment, Z and Z^1 are independently selected from: $-O-$, $-S-$, $-CH_2-$,
10 $-CHF-$, $-CF_2-$, $-C(R_fR_g)-$, $-N(Re)-$, or $-C(O)-$; wherein R_f and R_g are defined as described above for R_f and R_g in L1, respectively, and Re is defined is the same as that described for Re in E above. In a more preferred embodiment, at least one of Z and Z^1 is $-O-$. In a more preferred embodiment, both Z and Z^1 are $-O-$.

In one embodiment, Z is $-O-$.

15 In one embodiment, Z^1 is $-O-$.

In one embodiment, X^5 , X^6 and X^7 are each independently hydrogen, deuterium(D), halogen, nitrile, amino, trifluoromethyl, trifluoromethoxy, aminocarbonyl (H_2NCO), alkyl, heterocycloalkyl, alkoxy, heteroatomically substituted alkoxy, alkylamino (NR_iR_j), heteroatomically substituted alkylamino, alkoxycarbonyl, alkylaminocarbonyl,
20 alkylcarbonylamino, alkoxycarbonyl amino group, cycloalkoxycarbonyl amino group, alkyl sulfonamido group, cycloalkyl sulfonamido group, aryl, aryloxy, aryl amino carbonyl, aryl carbonyl amino group, aryloxy carbonyl amino group, heterocyclic aryl, heterocyclic aryl oxy, or heterocyclic aryl amino group; wherein R_i and R_j are each independently hydrogen, deuterium(D), alkyl, heterocycloalkyl, alkyl carbonyl, alkoxy carbonyl,
25 cycloalkoxy carbonyl, alkyl amino carbonyl, alkyl sulfonyl, cycloalkyl sulfonyl, aryl, aryloxy carbonyl, aryl amino carbonyl, heterocyclic aryl, or R_i and R_j are interconnected to form a 3-8-membered heterocyclic group containing 1-3 heteroatoms.

In one embodiment, X^5 is selected from: hydrogen, deuterium(D), halogen, nitrile, amino(NH_2), trifluoromethyl, trifluoromethoxy, C_1 - C_8 alkyl, C_1 - C_8 alkoxy, C_1 - C_8
30 alkylamino, C_1 - C_8 alkylcarbonylamino, C_1 - C_8 alkylamino carbonylamino, C_1 - C_8 alkoxycarbonylamino, C_3 - C_8 cycloalkoxycarbonylamino, C_1 - C_8 alkyl sulfonamido, C_3 - C_8 cycloalkyl sulfonamido, aryl, or aryloxycarbonyl amino groups. In a preferred embodiment, X^5 is selected from: hydrogen, deuterium(D), halogen, nitrile, trifluoromethyl,

trifluoromethoxy, C₁-C₈ alkoxy carbonyl amino group.

In one embodiment, X⁶ and X⁷ are each independently selected from: hydrogen, deuterium(D), halogen, C₁-C₈ alkylamino group, or C₁-C₈ alkoxy carbonylamino group. In a preferred embodiment, X⁶ is selected from: hydrogen, deuterium(D), halogen, C₁-C₈ alkylamino group or C₁-C₈ alkoxy carbonylamino group, and X⁷ is selected from: hydrogen.

The compounds of the present invention may exist in multiple isomeric forms, as well as one or more mutually variable isomeric forms, including two single mutually variable isomers, and mixtures of mutually variable isomers. The term "isomer" is intended to cover the entire range of isomeric forms of the compounds of the present invention, including the mutually variable isomeric forms of said compounds.

Some of the compounds described herein may have asymmetric centers and thus exist in different enantiomeric and diastereoisomeric forms. The compounds of the present invention may be in the form of optical isomers or diastereoisomers. Thus, the present invention covers the compounds of the present invention and their application in their optical isomeric, diastereoisomeric and mixtures thereof, including racemic mixtures, as described herein. The optical isomers of the compounds of the present invention can be obtained by known techniques, such as asymmetric synthesis, chiral chromatography, or by chemical separation of the stereoisomers using optically active splitting agents.

Unless otherwise indicated, "stereoisomer" means a stereoisomer of a compound that is substantially free of other stereoisomers of the compound. Thus, a stereoisomerically pure compound having a chiral center is substantially free of the opposite enantiomer of the compound. A stereoisomerically pure compound having two chiral centers is substantially free of other diastereoisomers of the compound. A typical stereoisomerically pure compound comprises greater than about 80 wt% of one stereoisomer of the compound and less than about 20 wt% of the other stereoisomers of the compound. For example, greater than about 90 wt% of one stereoisomer of the compound and less than about 10 wt% of the other stereoisomers of the compound, or greater than about 95 wt% of one stereoisomer of the compound and less than about 5 wt% of the other stereoisomers of the compound, or greater than about 97 wt% of one stereoisomer of the compound and less than about 3 wt% of the other stereoisomers of the compound.

If there is a discrepancy between the described structure and the name given to that structure, the described structure prevails. In addition, if the stereochemistry of a structure

or part of a structure is not indicated by, for example, a bold or dashed line, the structure or part of a structure should be understood to cover its entire stereoisomerism. However, in some cases where more than one chiral center is present, the structure and name may be indicated in the form of a single enantiomer to aid in describing the relevant stereochemistry. Those skilled in the field of organic synthesis should be aware of their preparation by means of the preparation of single enantiomers of said compounds.

In this specification, the "pharmaceutically acceptable salt" is a pharmaceutically acceptable, organic or inorganic acid or base salt of a compound of the present invention. Typical pharmaceutically acceptable salts include, for example, alkali metal salts, alkaline earth metal salts, ammonium salts, water-soluble salts, and water-insoluble salts. For example, acetate, aminostilbene sulfonate (4,4-diaminostilbene 2,2-disulfonate), benzenesulfonate, benzoate, bicarbonate, hydrogen sulfate, hydrogen tartrate, borate, bromide, butyrate, calcium, calcium edetate, camphor sulfonate, carbonate, chloride, citrate, clavulanate, dihydrochloride, edetate, ethyl disulfonate, propionate lauryl sulfate, ethanesulfonate fumarate, glucosinolate, glucosinolate, glutamate, ethanoic acid asanate, hexafluorophosphate, hexylresorcinol, haramino, hydrobromide, hydrochloride, hydroxynaphthoic acid salt, iodate, isothiosulfate, lactate, lactobionate, laurate, malate, maleate, lentilate, methanesulfonate, methyl bromide, methyl nitrate, methanesulfonate, mucilage Naphthalene sulfonate, nitrate, N-methyl glucosamino ammonium salt, 3-hydroxy-2-naphthoic acid salt, oleate, oxalate, palmitate, bis-hydroxynaphthalate (1,1-methylene-bis-2-hydroxy- 3-naphthoic acid salt, bis-hydroxynaphthalate, pantothenate, phosphate/diphosphate, picric acid salt, polygalacturonate, propionate, p-toluenesulfonate, salicylate, stearate, alkali acetate salicylate, alkali acetate, succinate, sulfate, sulfosalicylate, sulforaphane, ellagic acid, tartarate, 8-chlorophyllide, toluene sulfonate, triethyl iodide and valerate. A pharmaceutically acceptable salt may have more than one charged atom in its structure. In this embodiment, the pharmaceutically acceptable salt may have a plurality of counterbalance ions. Thus, the pharmaceutically acceptable salt may have one or more charged atoms and/or one or more counterbalancing ions.

The compounds of the present invention may be isotopically labeled in which one or more atoms are replaced by atoms having a different atomic mass or mass number. Examples of isotopes that may be incorporated into a compound of Formula Ib or IIb include: isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine, chlorine, or iodine. Examples of such isotopes are, respectively: ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O ,

¹⁸O, ³¹P, ³²P, ³⁵S, ¹⁸F, ³⁶Cl, ¹²³I and ¹²⁵I. These radiolabeled compounds can be used to detect biodistribution, tissue concentrations, and kinetics of transport and excretion from biological tissues, including the subject to which the labeled compound is administered. The labeled compounds are also used to determine the therapeutic effect, site or mode of action, and binding affinity of the therapeutic candidate for a pharmacologically important target. Thus, certain radiolabeled compounds of formula Ib or IIb can be used for drug and/or tissue distribution studies. The radioisotopes tritium, i.e., ³H, and carbon-14, i.e., ¹⁴C, are particularly useful for this purpose because they are easy to incorporate and detection means are readily available.

Substitution with a heavy isotope such as deuterium, i.e., ²H, can provide certain therapeutic advantages due to higher metabolic stability (e.g., prolonged in vivo half-life of deuterium-containing compounds). Substitution of hydrogen with deuterium reduces the dose required to achieve therapeutic efficacy and can therefore be preferably used in discovery or clinical settings.

Substitution of positron-emitting isotopes (e.g., ¹¹C, ¹⁸F, ¹⁵O and ¹³N) can provide labeled analogs of compounds of the present invention that have been used in positron imaging (PET) studies, for example, for the detection of substance receptor occupancy. For example, for the detection of substance receptor occupancy. Isotopically labeled compounds of Formula Ib or IIb can be prepared generally by conventional techniques known to those skilled in the art or by employing suitable isotopically labeled reagents in a manner similar to those described in the preparation and embodiments section below.

Embodiments of the present invention described herein are also intended to cover in vivo metabolites of compounds of Formula Ib or IIb. These products may be derived, for example, from processes such as oxidation, reduction, hydrolysis, amidation, esterification, etc. that are primarily attributable to the enzymatic activity of the compounds of the present invention after administration. Thus, the present invention includes such compounds which are produced in the form of by-products based on the enzymatic or non-enzymatic activity exerted on the compounds of the present invention after a period of time sufficient to produce metabolites in mammals given the compounds of the present invention. Metabolites, particularly those having pharmaceutical activity, are typically identified by administering a detectable dose of a radiolabeled compound of the present invention to a subject, such as a rat, mouse, guinea pig, monkey or human, for a sufficient period of time during which metabolism occurs, and by isolating said metabolite from a

urine, blood or other biological sample obtained from the subject receiving said radiolabeled compound.

The present invention also provides pharmaceutically acceptable salt forms of compounds of Formula Ib or IIb. The scope of the present invention covers acid addition salts and base addition salts, which are formed by bringing a pharmaceutically suitable acid or a pharmaceutically suitable base into contact with a compound of the present invention.

"Pharmacologically acceptable acid addition salts" means those salts that retain biological validity and free base properties, which are not undesirable biologically or otherwise, and which are formed using inorganic acids, such as, but not limited to hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, etc., and organic acids such as, but not limited to, acetic acid, 2,2-dichloroacetic acid, adipic acid, alginic acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetylaminobenzoic acid, camphoric acid, camphor-10-sulfonic acid, capric acid, capric acid, octanoic acid, carbonic acid, cinnamic acid, citric acid, cyclohexane sulfamic acid, dodecyl sulfate, ethane-1, 2-disulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, formic acid, fumaric acid, galactosuccinic acid, gentianic acid, glucuronic acid, glucuronic acid, glutamic acid, glutaric acid, 2-oxoglutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, isobutyric acid, lactic acid, lactobionic acid, lauric acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, muconic acid, naphthalene-1,5-disulfonic acid, naphthalene 2-sulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, dihydroxynaphthalic acid, propionic acid, pyroglutamic acid, pyruvic acid, salicylic acid, 4-aminosalicylic acid, sebacic acid, stearic acid, succinic acid, tartaric acid, thiocyanic acid, p-toluenesulfonic acid, trifluoroacetic acid, undecylenic acid, etc.

"Pharmacologically acceptable base addition salts" are those salts that retain the biological validity and properties of the free acid and are not biologically or otherwise undesirable. These salts are prepared by adding inorganic or organic bases to the free acid. Salts derived from inorganic bases include, but are not limited to, sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts, etc. Preferred inorganic salts are ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of: primary, secondary and tertiary amines, substituted amines, including naturally occurring substituted amines, cyclic amines and basic ion exchange resins such as ammonia,

isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, diethanolamine, ethanolamine, dimethylethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydantoin, choline, betaine, phenethylbenzylamine, benzylpenicillin, ethylenediamine, glucosamine, methylglucosamine, theobromine, triethanolamine, aminobutriol, purine, piperazine, piperidine, N ethyl piperidine, polyamine resin, etc. Particularly preferred organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline, and caffeine.

Crystallization typically produces solvent complexes of the compounds of the present invention. The term "solvent complex" as used herein refers to an aggregate comprising one or more molecules of the compound of the invention and one or more molecules of the solvent. The solvent may be water, in which case the solvent complex may be a hydrate. Alternatively, the solvent may be an organic solvent. Thus, the compounds of the present invention may exist in hydrate form, including monohydrate, dihydrate, hemihydrate, sesquihydrate, trihydrate, tetrahydrate, etc., and the corresponding solvent complex forms. The compounds of the invention may be true solvent compounds, while in other cases, the compounds of the invention may retain only indefinite water or a mixture of water plus some indefinite solvent.

"Stereoisomer" means a compound which is not interchangeable by being composed of identical atoms linked by the same bonds, but having different three-dimensional structures. The invention envisages various stereoisomers and mixtures thereof, and includes "enantiomeric", which means: two stereoisomers whose molecules are non-overlapping mirror images of each other.

The compounds of the invention, or pharmaceutically acceptable salts thereof, may contain one or more asymmetric centers, resulting in enantiomers, diastereoisomers, and other stereoisomeric forms that can be identified in terms of absolute stereochemistry, such as (R)- or (S)-, or, in the case of amino acids, such as (D) or (L). The present invention is intended to include all these possible isomers, as well as their racemic and optically pure forms. The optically active (+) and (), (R)- and (S)- or (D)- and (L)-isomers can be prepared using chiral synthons or chiral reagents, or split using conventional techniques, such as chromatography and graded crystallization. Conventional techniques used to prepare/isolate individual enantiomers include chiral synthesis from suitable optically pure precursors or the use of, for example, chiral high-pressure liquid chromatography (HPLC)

to split racemates (or racemates of salts or derivatives). When a compound described herein contains an alkene double bond or other geometric asymmetry center, it is intended to indicate that the compound includes the E and Z geometric isomers unless otherwise stated. Likewise, it is intended to include the full range of tautomerforms.

5

III. DRUG COMPOSITIONS

In one embodiment, a compound of Formula Ib or IIb, formulated in the form of a pharmaceutically acceptable composition, said composition comprising an amount of a compound of Formula Ib or IIb that is effective in treating a specific disease or condition of interest after administration of said pharmaceutical composition to a mammal. The pharmaceutical composition of the present invention may comprise a compound of Formula Ib or IIb in combination with a pharmaceutically acceptable carrier, diluent or excipient.

For this purpose, "pharmaceutically acceptable carrier, diluent, or excipient" includes, but is not limited to, any adjuvant, carrier, excipient, flow aid, sweetener, diluent, preservative, dye/colorant, flavor enhancer, surfactant, wetting agent, dispersant, suspending agent, stabilizer, isotonic agent, solvent, or emulsifier that is approved by the U.S. Food and Drug Administration as acceptable to humans or domestic animals.

In addition, "mammal" includes human and domestic animals, e.g., laboratory animals and domestic pets (e.g., cats, dogs, pigs, cattle, sheep, goats, horses, rabbits), and non-domestic animals, e.g., wild animals, etc.

Pharmaceutical compositions of the present invention can be prepared by combining compounds of the present invention with suitable pharmaceutically acceptable carriers, diluents, or excipients, and can be formulated in solid, semi-solid, liquid, or gaseous forms, such as, for example, tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalations, gels, microspheres, and aerosols. Typical routes of administration of such pharmaceutical compositions include, but are not limited to, oral, topical, transdermal, inhalation, parenteral, sublingual, buccal, rectal, vaginal, and intranasal. The term parenteral as used herein includes, subcutaneous, intravenous, intramuscular, intrathoracic, or infusion techniques. The pharmaceutical compositions of the present invention are formulated to allow the active ingredient contained therein to be bioavailable after administration of the composition to a patient. The composition to be given to the subject or patient may be in the form of one or more dosage units, for example,

the tablet may be a single dosage unit and the container of the compound of the invention in aerosol form may contain a plurality of dosage units. Practical methods for preparing such dosage forms are known or understood by those skilled in the art; see, for example, Remington: The Science and Practice of Pharmacy, 20th edition (Philadelphia College of
5 Pharmacy and Science), 2000). In any case the composition to be given contains a therapeutically effective amount of a compound of the present invention or a pharmaceutically acceptable salt thereof for the treatment of a disease or condition of interest according to the teachings of the present invention.

The pharmaceutical compositions of the present invention can be in solid or liquid
10 form. On the one hand, the carrier is a particle, whereby the composition is, for example, in the form of a tablet or a powder. The carrier may be a liquid, wherein said composition is, for example, an oral type syrup, an injectable liquid, or an aerosol, which may be used, for example, for inhalation administration. When intended for oral administration, said pharmaceutical composition is preferably in solid or liquid form, wherein, semi-solid,
15 semi-liquid, suspension, and gel forms are included in the solid or liquid forms contemplated herein.

As solid compositions for oral administration, the pharmaceutical compositions can be formulated in the form of powders, granules, compressed tablets, pills, capsules, chewable gums, flakes, and the like. Such solid compositions will typically contain one or
20 more inert diluents or edible carriers. In addition, one or more of the following substances may be present: binders, such as carboxymethyl cellulose, ethyl cellulose, microcrystalline cellulose, yarrow gum or gelatin; excipients, such as starch, lactose or dextrin, disintegrants, such as alginate, sodium alginate, Primogel, corn starch, etc.; lubricants, such as magnesium stearate or Sterotex; flow aids, such as colloidal silica; sweeteners, such as
25 sucrose or saccharin; flavoring agents, such as menthol, methyl salicylate, or orange flavoring agents; and coloring agents.

If the pharmaceutical composition is in the form of a capsule (e.g., a gelatin capsule), it may contain a liquid carrier, e.g., a polyethylene glycol or an oil, in addition to the types of substances described above.

30 The pharmaceutical composition may be in the form of a liquid, such as an elixir, syrup, solution, emulsion or suspension. As two examples, the liquid may be for oral administration or for delivery by injection. If intended for oral administration, the preferred composition may contain one or more sweeteners, preservatives, dyes/colors, and flavor

"Effective amount" or "therapeutically effective amount" means an amount of a compound of the present invention which (when administered to a mammal, preferably a human) is sufficient to provide effective treatment of an Mnk-related condition or disease in that mammal (preferably a human), as described below. The amount of compound of the present invention that constitutes a "therapeutically effective amount" will vary depending on the compound, the condition and its severity, the manner of administration, and the age of the mammal to be treated, but can be routinely determined by those skilled in the art based on their knowledge and the content of the present invention.

The compounds of the present invention, or pharmaceutically acceptable salts thereof, may also be given prior to, concurrently with, or after the administration of one or more other therapeutic agents. Such combination therapy includes the administration of a single drug-dose formulation comprising the compound of the present invention and one or more other active substances, and the administration of the compound of the present invention and each active substance in separate drug-dose formulations. For example, the compounds of the present invention and other active substances may be given to the patient together in a single oral dose combination (e.g., tablets or capsules), or each substance may be given in separate oral dose formulations. When using separate dose formulations, the compounds of the present invention and one or more other active agents may be given at substantially the same time (i.e., simultaneously) or at separate staggered times (i.e., sequentially); it should be understood that combination therapy includes all of these regimens.

The present invention also optimizes the structure of GPR40 target agonist compound formula **Ib-IIb** by further study of structure activity relationship (SAR) and can effectively reduce the risk of hypoglycemia production for safer and more effective treatment of type II diabetes.

The benzo oxygen-containing heterocyclic compounds of the present invention can lower blood glucose levels by activating GPR40 targets to stimulate insulin release from pancreatic beta cells, and such compounds of formula Ib-IIb are characterized by promoting insulin secretion only when blood glucose concentrations are high in patients with type II diabetes, thus effectively reducing the risk of hypoglycemia in patients and providing better GPR40 target selectivity and safety.

DETAILED DESCRIPTION OF THE INVENTION


Details of the present invention are further illustrated below by way of embodiments,

but the invention is not thereby limited to the scope of the described embodiments. Experimental methods for which specific conditions are not indicated in the following embodiments are completed according to conventionally known methods and conditions, or according to the commodity specification of choice for implementation.


5 The abbreviated notes of the chemical reagents and solvents used in the synthesis of the novel benzo oxygen-containing heterocyclic compounds of the present invention are all summarized in the description of the apparatus and raw materials section of the Examples.

10 The key innovation of the present invention is to follow synthetic reaction routes 1 or 2 as shown in the following schemes 1 and 2, first preparation of the target products Ib-IIb or intermediate RM-2b by reaction of raw material RM-1b with raw material SM-1 (e.g. reagents of the following SM-1a or RM-1b structures), respectively; when R^7 is an alkoxy group (e.g., $R^7 = \text{OCH}_3$ or OEt), the individual target products of the present invention in which R^7 is a hydroxyl group in Formula **IIb** can then be synthesized by the LiOH hydrolysis reaction.

15



The synthetic reaction route 1b described above is as follows:




Scheme 1: Synthesis reactions of compounds of formula Ib-IIb

In the above synthesis reaction route 1b was operated as follows:

- 5 1. Synthesis of RM-2b: RM-Ib (1.0eq), SM-1b (1.0eq), and DIAD and PPh₃ (1.2eq) were added to THF (5x) in a round bottom reaction flask, nitrogen was replaced and protected, and the reaction was followed by TLC and/or HPLC until the end of the reaction, and RM-2b (or RM-IIb) was obtained after post-treatment and other routine operations.
- 10 2. Synthesis of target products Ib-IIb: RM-Ib (or RM-IIb, 1eq) was hydrolyzed in a mixture of MeOH-H₂O with LiOH (1:1, 10X) in a round-bottom reaction flask, detected by TLC and/or HPLC until the end of the reaction, and then the target products Ib-IIb were obtained after post-treatment and purification and other routine operations.

The synthetic reaction route 2b is as follows:




































Scheme 2: Synthesis reactions of the formula Ib-IIb compounds

In the above synthesis reaction route 2 was operated as follows:

- 5 1. Synthesis of RM-2b: RM-Ib (1.0eq), SM-1b (1.0eq), and basic reagents (e.g. potassium phosphate or potassium carbonate, 2-3eq) were added to DMF (5x) in a round bottom reaction flask, nitrogen was replaced and the reaction was protected. The reaction solution was slowly poured into 40X ice water with stirring, and RM-Ib (or RM-IIb) was obtained after post-processing and purification as usual.
- 10 2. Synthesis of target product IIb: RM-Ib (or RM-IIb) was hydrolyzed in a mixture of MeOH-H₂O with LiOH (1:1, 10X) in a round-bottom reaction flask, respectively, and detected by TLC and/or HPLC until the end of the reaction, and then the target products Ib-IIb were obtained after post-treatment and purification and other routine operations.

The above synthesized compounds of formula Ib-IIb, where n, n, L¹, R¹, R², R³, R⁵, R^{5b}, R⁷, Ra, Rb, Rc, Rd, Re, X⁵, X⁶, Y, Y¹ and Z are defined the same as n, L¹, R¹, R², R³, R⁵, R^{5b}, R⁷, Ra, Rb, Rc, Rd, Re, X⁵, X⁶, Y, Y¹, and Z of claims 1-4 of the present invention, respectively. Examples of synthetic reactions for the compounds in Formulas Ib-IIb of the present invention are shown in Table 3 below, respectively, for the preparation of each specific compound of different structures, which were successfully synthesized using the benzo-oxanthene raw material RM-1b in Table 1 and the raw material SM-1 in Table 2 below, respectively:

 RM-Ib-1	 RM-Ib-2	 RM-Ib-3
 RM-Ib-4	 RM-Ib-5	 RM-Ib-6
 RM-Ib-7	 RM-Ib-8	 RM-Ib-9
 RM-Ib-10	 RM-Ib-11	 RM-Ib-12
 RM-Ib-13	 RM-Ib-14	 RM-Ib-15
 RM-Ib-16	 RM-Ib-17	 RM-Ib-18

 RM-Ib-19	 RM-Ib-20	 RM-Ib-21
 RM-Ib-22	 RM-Ib-23	 RM-Ib-24
 RM-Ib-25	 RM-Ib-26	 RM-Ib-27
 RM-Ib-28	 RM-Ib-29	 RM-Ib-30
 RM-Ib-31	 RM-Ib-32	 RM-Ib-33




 RM-Ib-34	 RM-Ib-35	 RM-Ib-36
---	---	---

Table 1: The raw materials SM-Ib-01 to SM-Ib-30 used in the present invention and their structures














 SM1-1	 SM1-2	 SM1-3
 SM1-4	 SM1-5	 SM1-6
 SM1-7	 SM1-8	 SM1-9
 SM1-10	 SM1-11	 SM1-12

Table 2: Raw materials SM1-01 to SM1-15 used in the present invention and their

5 structures

The above-mentioned raw materials RM-1b and SM1 were used, and then intermediate RM-2b was prepared by the synthetic reaction route in Scheme 3 below, and finally each specific compound of formula Ib-IIb was synthesized by hydrolysis reaction, respectively. Examples of specific reactions are as follows, respectively:

10



Scheme 3: Examples of synthetic reactions of the formula Ib-IIb compounds



5 In the above example of synthesis reaction of Scheme 3:













Step 1 reaction: The key intermediate compound (RM-Ib-01) is first obtained by first passing raw material RM-Ib-01 with another raw material SM1-01 in the presence of an inorganic base (e.g., K₃PO₄) in a solvent (e.g., DMSO or DMF);













10 The second reaction step: the intermediate (RM-2b-01) is then hydrolyzed in the presence of an inorganic base (e.g., LiOH) in a solvent (e.g., MeOH or a mixture of methanol and water) to obtain the target product Ib-01.











The intermediate structures of the intermediate compound RM-2b obtained by the first step reactions of the synthetic reaction routes in options 1, 2 and 3 above, respectively, are shown in the following structural formula series of RM-2b; and the target products of formula IIb obtained by the second step hydrolysis reactions, respectively, are shown in the following structural formula series of formula IIb:













RM-2b structural series:













1	 RM-Iib-1	2	 RM-Iib-2
---	--	---	---



3	 RM-IIb-3	4	 RM-IIb-4
5	 RM-IIb-5	6	 RM-IIb-6
7	 RM-IIb-7	8	 RM-IIb-8
9	 RM-IIb-9	10	 RM-IIb-10
11	 RM-IIb-11	12	 RM-IIb-12
13	 RM-IIb-13	14	 RM-IIb-14

15	 RM-IIb-15	16	 RM-IIb-16
17	 RM-IIb-17	18	 RM-IIb-18
19	 RM-IIb-19	20	 RM-IIb-20
21	 RM-IIb-21	22	 RM-IIb-22
23	 RM-IIb-23	24	 RM-IIb-24
25	 RM-IIb-25	26	 RM-IIb-26









			RM-IIb-26
27	 RM-IIb-27	28	 RM-IIb-28
29	 RM-IIb-29	30	 RM-IIb-30
31	 RM-IIb-31	32	 RM-IIb-32
33	 RM-IIb-33	34	 RM-IIb-34
35	 RM-IIb-35	36	 RM-IIb-36











37		38	
	RM-IIb-37		RM-IIb-38
39		40	
	RM-IIb-39		RM-IIb-40
41		42	
	RM-IIb-41		RM-IIb-42
43		44	
	RM-IIb-43		RM-IIb-44
45		46	
	RM-IIb-45		RM-IIb-46
47		48	
	RM-IIb-47		RM-IIb-48













49	 RM-IIb-49	50	 RM-IIb-50
51	 RM-IIb-51	52	 RM-IIb-52
53	 RM-IIb-53	54	 RM-IIb-54
55	 RM-IIb-55	56	 RM-IIb-56
57	 RM-IIb-57	58	 RM-IIb-58
59	 RM-IIb-59	60	 RM-IIb-60











61	 RM-IIb-61	62	 RM-IIb-62
----	--	----	---











Formula IIb structural series











1	 IIb-1	2	 IIb-2
3	 IIb-3	4	 IIb-4
5	 IIb-5	6	 IIb-6
7	 IIb-7	8	 IIb-8





9	 IIb-9	10	 IIb-10
11	 IIb-11	12	 IIb-12
13	 IIb-13	14	 IIb-14
15	 IIb-15	16	 IIb-16
17	 IIb-17	18	 IIb-18

19	 IIb-19	20	 IIb-20
21	 IIb-21	22	 IIb-22
23	 IIb-23	24	 IIb-24
25	 IIb-25	26	 IIb-26
27	 IIb-27	28	 IIb-28
29		30	

	IIb-29		IIb-30
31	 IIb-31	32	 IIb-32
33	 IIb-33	34	 IIb-34
35	 IIb-35	36	 IIb-36
37	 IIb-37	38	 IIb-38
39		40	

	IIb-39		IIb-40
41	 IIb-41	42	 IIb-42
43	 IIb-43	44	 IIb-44
45	 IIb-45	46	 IIb-46
47	 IIb-47	48	 IIb-48
49	 IIb-49	50	 IIb-50

51	 IIb-51	52	 IIb-52
53	 IIb-53	54	 IIb-54
55	 IIb-55	56	 IIb-56
57	 IIb-57	58	 IIb-58
59	 IIb-59	60	 IIb-60

61	 Iib-61	62	 Iib-62
63	 Iib-63	64	 Iib-64

The experimental conditions and product analysis results for each specific reaction step are listed separately in the Examples.

Specific results concerning the synthesis and analysis of each of the novel structural compounds of formula **Ib-Iib** described above are detailed in the final embodiment of the invention, with the chemistry of each compound and its chiral asymmetric purity determined by HPLC on a chiral column and the corresponding chemical structure characterization determined by LC-MS and/or hydrogen spectrum nuclear magnetic resonance (¹H-NMR) analysis, respectively.

The synthesis and effects of the various types of intermediates and compounds of the present invention are illustrated below by means of embodiments.

The apparatus and raw materials involved in the embodiment are described as follows:

The infrared spectral data were analyzed using a Thermo Nicolet Fourier Transform AVATAR™ 360 E.S.P™ infrared instrument and are expressed in cm⁻¹ units.

The NMR hydrogen spectra were obtained from the analysis of a Varian Mercury Plus 400 (400 MHz) NMR instrument. The chemical shifts were recorded using tetramethylsilane as an internal standard and expressed in ppm (CHCl₃: δ= 7.26 ppm). The following data information was recorded: chemical shifts and their cleavage and coupling constants (s: single peak; d : double peak; t: triple peak; q: quadruple peak; br: broad peak;

m: multiple peak).

The mass spectrometry data were analyzed using a liquid mass spectrometer from Finnigan Advanced LCQ (Finnigan LCQ Advantage), among other needs. The molecular weights of the organic carboxylic acid (-COOH) compounds of formulas Ib-IIb in this invention are mainly in the anionic mode ESI-MS [(M-H)+], but the ester intermediate compounds of formulas RM-IIb and some compounds containing amine groups in **Ib-IIb** have molecular weights in the cationic mode ESI-MS [(M+H)+].

The special raw materials and intermediates involved in this invention were provided by Zannan Technology Co., Ltd. and other custom processing, and all other chemical reagents were purchased from reagent suppliers such as Shanghai Reagent Company, Aldrich Company (Aldrich), Acros Company (Acros), etc. If the intermediates or products required for the reaction during the synthesis were not enough for the next step and other tests, the synthesis was repeated several times until sufficient quantities were available. The GPR40 activity tests of the compounds prepared by the invention as well as pharmacological and toxicological tests were done by CRO service units in Shanghai and Beijing.

The abbreviations of the relevant chemical raw materials, reagents and solvents involved in the present invention and its embodiments are annotated as follows:

AIBN: Azo diisobutyronitrile

Boc: tert-Butoxycarbonyl

(Boc)₂O: Di-tert-butyl dicarbonate

CDI: N,N'-carbonyl diimidazole

DBU: 1,8-Diazabicyclo[5.4.0]undec-7-ene

Et: Ethyl

EDCI: N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide hydrochloride

HATU: 2-(7-azobenzotriazole)-N,N,N',N'-tetramethyluronium hexafluorophosphate

NBS: N-bromosuccinimide

SOCl₂: Sulfoxide chloride

Pd/C: Palladium carbon

DMAP: 4-Dimethylaminopyridine

HMTA: Hexamethylenetetramine

DIEA: N,N-diisopropylethylamine

Py: Pyridine

HBr: Hydrobromic acid

HCl: Hydrochloric acid

HOAc: Glacial acetic acid

TFA: Trifluoroacetic acid

5 TsOH: p-Toluenesulfonic acid

NaOH: Sodium hydroxide

LiOH: Lithium hydroxide

ACN: Acetonitrile

DCM: Dichloromethane

10 DCE: Dichloroethane

DMF: N,N-dimethylformamide

DMSO: Dimethyl sulfoxide

Et₂O: Diethyl ether

EA: Ethyl acetate

15 PE: Petroleum ether


THF: Tetrahydrofuran

TBME: Methyl tert-butyl ether

Example 1

20 Synthesis of compound **IIb-1**

The synthesis reaction is shown in the following general formula I:



Step 1:

The raw materials **RM-Ib-1** (0.051g, 0.24mmol) and SM1-1 (0.050g, 0.24mmol, 1.0eq.) were dissolved in 2mL of DMF, and potassium carbonate (0.050g, 0.36mmol, 1.5eq.) was added with stirring, and the reaction was completed by heating to 90°C under nitrogen protection overnight. HPLC showed that the reaction solution was cooled to room temperature after the reaction was finished, water was added, ethyl acetate was added for extraction, and then the organic phases were combined, washed with saline and dried, and the intermediate RM-IIb-1 (0.1g) was concentrated.

ESI-MS [(M+H)⁺]: m/z calculated: 343.1, founded: 343.0.

Step 2:

Added MeOH (2 mL), THF (1 mL) and aqueous LiOH (1N, 1 mL) were added sequentially to intermediate RM-IIb-1, stirred at room temperature for 1 hr. After the reaction was completed, the reaction solution was neutralized with hydrochloric acid (1N) to pH about 4, added water, added ethyl acetate for extraction, and then the organic phases were combined, washed with saline and dried, and finally purified by column chromatography, obtained the light yellow solid product IIb-1 (0.040 g) in two-step yield: 50.7%.


Upon examination, ¹H NMR (400 MHz, CDCl₃) of product IIb-1: δ 7.07-7.05 (m, 1H), 6.95-6.93 (m, 1H), 6.86-6.79 (m, 2H), 6.52-6.49 (m, 2H), 5.99 (s, 2H), 5.01 (s, 2H), 4.78-4.74 (m, 1H), 4.31-4.27 (m, 1H), 3.83-3.80 (m, 1H), 2.83-2.78 (m, 1H), 2.65-2.59 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of IIb-1 calculated: 327.1, founded: 327.0.

Example 2

Synthesis of compound **IIb-2**

The synthesis reaction is shown in the following general formula II:



Step 1:

The raw materials RM-Ib-2 (0.043 g, 0.19 mmol), SM1-1 (0.039 g, 0.19 mmol, 1.0 eq.) and PPh3 (0.146 g, 0.56 mmol, 3 eq.) were dissolved in 4 mL of THF, protected by nitrogen, under cooling in an ice-water bath, and DIAD (0.113 g, 0.56 mmol, 3eq.), stirred for 1h in an ice water bath after addition, and the reaction was completed at 20-30 degrees C overnight. HPLC showed that the reaction solution was cooled to room temperature after the reaction was finished, water was added, DCM was added for extraction, and then the organic phases were combined, washed with saline and dried, and the crude product was concentrated; the crude product was separated by TLC scraper to obtain intermediate **RM-IIb-2**.

ESI-MS [(M+H)⁺]: m/z of RM-IIb-2 calculated: 421.0, founded: 421.1.

Step 2:

MeOH (2 mL), THF (1 mL) and aqueous LiOH (1N, 1 mL) were added to intermediate RM-IIb-2 and stirred at room temperature for 1h. After the reaction was completed as indicated by HPLC, the reaction solution was neutralized with hydrochloric acid (1N) to pH about 4, water was added, ethyl acetate was added for extraction, and then the organic phases were combined, washed with saline and dried, and finally purified by column chromatography to give The pale yellow solid product IIb-2 (0.025 g) was obtained in two-step yield: 33%.

¹H-NMR for the **IIb-2** hydrochloride (400 MHz, CDCl₃) δ: 7.09-7.05 (m, 2H), 6.92 (m, 1H), 6.50-6.46 (m, 2H), 6.01 (s, 2H), 4.95 (s, 2H), 4.79-4.75 (m, 1H), 4.31-4.28 (m, 1H), 3.84 -3.79 (m, 1H), 2.84-2.79 (m, 1H), 2.66-2.59 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of IIb-2 calculated: 405.0, founded: 405.2.

Example 3

Synthesis of compound **IIb-3**

The synthesis of compound IIb-3 was prepared in the same way as in Example 1, and the product IIb-3 was obtained by etherification and hydrolysis reaction, wherein compound RM-Ib-3 (0.48 mmol) was used in the reaction instead of compound RM-Ib-1 to obtain intermediate RM-IIb-3, and the white solid product IIb-3 (0.039 g) was obtained from the intermediate by hydrolysis in two-step yield :20%.

ESI-MS [(M-H)⁺]: m/z of IIb-3 calculated 405.0, founded: 404.8.

Example 4

Synthesis of compound **IIb-4**

The synthesis method for the preparation of compound IIb-4 was the same as that of Example 1, and the product IIb-4 was obtained by etherification and hydrolysis reaction, wherein compound RM-Ib-4 (0.24 mmol) was used in the reaction instead of compound RM-Ib-1 to obtain intermediate RM-IIb-4, and the light yellow solid product IIb-4 (0.060 g) was obtained from the intermediate by hydrolysis in two-step yield: 76%.

¹H-NMR for the **IIb-4**hydrochloride (400 MHz, CDCl₃)δ: 7.07-7.05 (m, 1H), 6.91-6.86 (m, 2H), 6.81-6.79 (m, 1H), 6.49-6.45 (m, 2H), 5.96 (s, 2H), 4.91 (s, 2H), 4.78-4.74 (m, 1H) , 4.31-4.27 (m, 1H), 3.83-3.79 (m, 1H), 2.84-2.78 (m, 1H), 2.65-2.59 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of IIb-4 calculated:327.1, founded: 327.0.

Example 5

Synthesis of compound **IIb-5**

The synthesis of compound IIb-5 was prepared in the same way as in Example 1, and the product IIb-5 was obtained by etherification and hydrolysis reaction, wherein compound RM-Ib-5 (0.54 mmol) was used in the reaction instead of compound RM-Ib-1 to obtain intermediate RM-IIb-5, and the white solid product IIb-5 (0.068 g) was obtained from the intermediate by hydrolysis, in two-step yield : 34.8%.

¹H-NMR for the **IIb-5**(400 MHz, CDCl₃) δ: 7.08-7.06 (m, 1H), 6.95 (s, 1H), 6.78 (s, 1H), 6.50-6.46 (m, 2H), 6.02 (s, 2H), 4.95 (s, 2H), 4.79-4.75 (m, 1H), 4.32-4.28 (m, 1H), 3.83-3.80 (m, 1H), 2.84-2.78 (m, 1H), 2.66-2.59 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of Iib-5 calculated:361.1,founded:360.9.

Example 6

Synthesis of compound **Iib-6**

The synthesis of compound **Iib-6** was prepared in the same way as in Example 1, and the product **Iib-6** was obtained by etherification and hydrolysis reaction, wherein compound **RM-Ib-6** (0.41 mmol) was used in the reaction instead of compound **RM-Ib-1** to obtain intermediate **RM-Iib-6**, and the white solid product **Iib-6** (0.090 g) was obtained from the intermediate by hydrolysis in two-step yield: 63.6%.

¹H-NMR for the **Iib-6** hydrochloride (400 MHz, DMSO) δ: 7.10-7.09 (m, 1H), 6.92-6.89 (m, 1H), 6.74-6.71 (m, 1H), 6.46-6.44 (m, 2H), 6.08 (s, 2H), 4.93 (s, 2H), 4.69-4.64 (m, 1H), 4.19-4.15 (m, 1H), 3.69-3.64 (m, 1H), 2.71-2.65 (m, 1H), 2.48-2.43 (m, 1H). ¹⁹F NMR (376 MHz, DMSO) δ -120.47.

ESI-MS [(M-H)⁺]: m/z of Iib-6 calculated:345.1, founded:345.0.

Example 7

Synthesis of compound **Iib-7**

The synthesis method for preparing compound **Iib-7** was the same as that of Example 1, and the product **Iib-7** was obtained by etherification and hydrolysis reaction, in which compound **RM-Ib-6** (0.35 mmol) was used in the reaction instead of compound **RM-Ib-1** and compound **SM1-2** (0.35 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-Iib-7**, and the intermediate was hydrolyzed to give the white solid product **Iib-7** (0.085 g) in two-step yield: 70.8%. The white solid product **Iib-7** (0.085 g) was obtained in two-step yield: 70.8%.

¹H-NMR for the **Iib-7**hydrochloride (400 MHz, CDCl₃) δ: 7.07-7.05 (m, 1H), 6.66-6.63 (m, 1H), 6.56-6.54 (m, 1H), 6.49-6.46 (m, 2H), 6.01 (s, 2H), 4.96 (s, 2H), 4.79-4.74 (m, 1H) , 4.31-4.27 (m, 1H), 3.84-3.77 (m, 1H), 2.84-2.79 (m, 1H), 2.66-2.59 (m, 1H); ¹⁹F NMR (376 MHz, CDCl₃): δ -119.95.

The ESI-MS [(M-H)⁺]: m/z of Iib-7 calculated: 345.1, founded: 345.2.

Example 8

Synthesis of Compound **Iib-8**

The synthesis of compound **Iib-8** was prepared in the same way as in Example 1, and

the product **Ib-8** was obtained by etherification and hydrolysis reaction, wherein compound **RM-Ib-7** (0.24 mmol) was used in the reaction instead of compound **RM-Ib-1** to obtain intermediate **RM-Ib-8**, and the white solid product **Ib-8** (0.059 g) was obtained from the intermediate by hydrolysis, in two-step yield : 65.5%.

¹H-NMR for the **Ib-8** hydrochloride (400 MHz, CDCl₃) δ : 7.09-7.07 (m, 2H), 6.79-6.76 (m, 1H), 6.57-6.54 (m, 2H), 5.35 (s, 2H), 4.79-4.75 (m, 1H), 4.32-4.25 (m, 5H), 3.84-3.80 (m, 1H), 2.84-2.79 (m, 1H), 2.65-2.59 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of **Ib-8** calculated: 419.0;founded: 418.9.

Example 9

Synthesis of Compound **Ib-9**

The synthesis of compound **Ib-9** was prepared in the same way as in Example 2, and the product **Ib-9** was obtained by etherification and hydrolysis reaction, wherein compound **RM-Ib-8** (0.24 mmol) was used in the reaction instead of compound **RM-Ib-2** to obtain intermediate **RM-Ib-9**, and the white solid product **Ib-9** (0.035 g) was obtained from the intermediate by hydrolysis in two-step yield : 45%.

ESI-MS [(M-H)⁺]: m/z of **Ib-9** calculated: 419.0,founded: 418.8.

Example 10

Synthesis of compound **Ib-10**

The synthesis of compound **Ib-10** was carried out in the same way as in Example 1, and the product **Ib-10** was obtained by etherification and hydrolysis reaction, in which compound **RM-Ib-9** (0.65 mmol) was used in the reaction instead of compound **RM-Ib-1** to obtain intermediate **RM-Ib-10**, and the white solid product **Ib-10** (0.196 g) was obtained from the intermediate by hydrolysis in two steps. Yield: 53.2%.

¹H-NMR for the **Ib-10**hydrochloride (400 MHz, CDCl₃) δ: 7.34 (s, 1H), 7.07-7.05 (m, 1H), 7.00 (m, 1H), 6.48 (m, 1H), 6.51-6.46 (m, 2H), 4.98 (s, 2H), 4.79-4.74 (m, 1H), 4.31- 4.27 (m, 5H), 3.85-3.78 (m, 1H), 2.84-2.79 (m, 1H), 2.66-2.60 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of **Ib-10** calculated: 375.1,founded: 374.9.

Example 11

Synthesis of compound **Ib-11**

The synthesis method for the preparation of compound **Ib-11** was the same as that of

Example 2, and the product **I Ib-11** was obtained by etherification and hydrolysis reaction, wherein compound **RM-Ib-10** (0.24 mmol) was used in the reaction instead of compound **RM-Ib-2** to obtain intermediate **RM-I Ib-11**, and the white solid product **I Ib-11** (0.033 g) was obtained from the intermediate by hydrolysis in two steps Yield:37%.

5 ESI-MS [(M-H)⁺]: m/z of I Ib-11calculated: 359.1, founded: 359.0.

Example 12

Synthesis of compound **I Ib-12**

The synthesis of compound **I Ib-12** was carried out in the same way as in Example 2,
10 and the product **I Ib-12** was obtained by etherification and hydrolysis, in which compound **RM-Ib-11** (0.63 mmol) was used in the reaction instead of **RM-Ib-2** to obtain intermediate **RM-I Ib-12**, and the light yellow solid product **I Ib-12** (0.028 g) was obtained from the intermediate by hydrolysis. Two-step yield: 12.8%.

ESI-MS [(M-H)⁺]: m/z of I Ib-12calculated: 345.1, founded: 344.9.

15

Example 13

Synthesis of Compound **I Ib-13**

The synthesis of compound **I Ib-13** was carried out in the same way as in Example 1,
and the product **I Ib-13** was obtained by etherification and hydrolysis, in which compound
20 **RM-Ib-12** (0.43 mmol) was used in the reaction instead of **RM-Ib-1** to obtain intermediate **RM-I Ib-13**, and the light yellow solid product **I Ib-13** (0.060 g) was obtained from the intermediate by hydrolysis. Two-step yield: 40%.

¹H-NMR for the **I Ib-13**hydrochloride (400 MHz, CDCl₃)δ: 6.92-6.90 (m, 1H),
6.65-6.62 (m, 1H), 6.49-6.39 (m, 3H), 5.90 (s, 2H), 4.90 (s, 2H), 4.62-4.58 (m, 1H),
25 4.20-4.16 (m, 1H), 3.64-3.61 (m, 1H), 2.87-2.57 (m, 1H), 2.41-2.19 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of I Ib-13 calculated:345.0,founded: 345.0.

Example 14

Synthesis of Compound **I Ib-14**

30 The synthesis method for the preparation of compound **I Ib-14** was the same as that of Example 1, and the product **I Ib-14** was obtained by etherification and hydrolysis reaction, in which compound **RM-Ib-13** (0.22 mmol) was used in the reaction instead of compound **RM-Ib-1** to obtain intermediate **RM-I Ib-14**, and the white solid product **I Ib-14**

(0.045 g) was obtained from the intermediate by hydrolysis in two steps. Yield: 51.3%.

¹H-NMR for the **I Ib-14**hydrochloride (400 MHz, CDCl₃)δ: 7.28 (s, 1H), 7.08-7.06 (m, 1H), 7.00 (s, 1H), 6.51-6.47 (m, 2H), 6.09 (s, 2H), 5.00 (s, 2H), 4.79-4.75 (m, 1H), 4.32- 4.28 (m, 1H), 3.84-3.79 (m, 1H), 2.84-2.78 (m, 1H), 2.66-2.59 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of I Ib-14 calculated: 395.1, founded: 395.0.

Example 15

Synthesis of compound **I Ib-15**

The synthesis of compound **I Ib-15** was carried out in the same way as in Example 1, and the product **I Ib-15** was obtained by etherification and hydrolysis reaction, in which compound **RM-Ib-14** (0.42 mmol) was used in the reaction instead of compound **RM-Ib-1** to obtain intermediate RM-I Ib-15, and the white solid product **I Ib-15** (0.072 g) was obtained from the intermediate by hydrolysis in two steps. Yield: 54%.

¹H-NMR for the **I Ib-15**hydrochloride (400 MHz, CDCl₃) δ:7.34 (s, 1H), 7.08-7.06 (m, 1H), 7.00 (s, 1H), 6.48-6.43 (m, 2H), 6.11 (s, 2H), 4.98 (s, 2H), 4.79-4.75 (m, 1H), 4.32-4.28 (m, 1H), 3.83-3.80 (m, 1H), 2.83-2.78 (m, 1H), 2.65-2.59 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of I Ib-15 calculated: 352.1,founded: 352.0.

Example 16

Synthesis of Compound **I Ib-16**

The synthesis method for the preparation of compound **I Ib-16** was the same as that of Example 1, and the product **I Ib-16** was obtained by etherification and hydrolysis reaction, wherein compound **RM-Ib-15** (0.39 mmol) was used in the reaction instead of compound **RM-Ib-1** to obtain intermediate RM-I Ib-16, and the white solid product **I Ib-16** (0.032 g) was obtained from the intermediate by hydrolysis in two steps. Yield: 19.7%.

¹H-NMR for the**I Ib-16** hydrochloride (400 MHz, CDCl₃)δ: 7.51-7.48 (m, 2H), 7.29-7.27 (m, 2H), 7.06-7.04 (m, 1H), 6.47-6.43 (m, 2H), 4.97 (s, 2H), 4.78-4.74 (m, 1H), 4.30-4.27 (m, 1H), 3.83-3.78 (m, 1H), 2.83-2.77 (m, 1H), 2.65-2.58 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of I Ib-16 calculated: 405.0,founded: 404.8.

Example 17

Synthesis of Compound **I Ib-17**

The synthesis of compound **I Ib-17** was carried out in the same way as in Example 1,

and the product **Ib-17** was obtained by etherification and hydrolysis reaction, in which compound **RM-Ib-16** (0.28 mmol) was used in the reaction instead of compound **RM-Ib-1** to obtain intermediate **RM-Ib-17**, and the white solid product **Ib-17** (0.020 g) was obtained from the intermediate by hydrolysis in two steps. Yield: 15.8%.

¹H-NMR for the **Ib-17**hydrochloride (400 MHz, CDCl₃) δ: 7.30-7.18 (m, 1H), 7.10-7.06 (m, 1H), 7.03-7.01 (m, 1H), 6.50-6.46 (m, 2H), 5.07 (s, 2H), 4.79-4.75 (m, 1H), 4.31-4.27 (m, 1H), 3.84-3.79 (m, 1H), 2.83-2.78 (m, 1H), 2.65-2.58 (m, 1H). ¹⁹F NMR (376 MHz, CDCl₃): δ -49.74.

ESI-MS [(M-H)⁺]: m/z of **Ib-17** calculated: 363.1, founded: 362.9.

Example 18

Synthesis of Compound **Ib-18**

The synthesis of compound **Ib-18** was carried out in the same way as in Example 1, and the product **Ib-18** was obtained by etherification and hydrolysis, wherein compound **RM-Ib-16** (0.22 mmol) was used instead of compound **RM-Ib-1** and compound **SM1-2** (0.22 mmol) was used instead of **SM1-1** to obtain intermediate **RM-Ib-18**. The intermediate was hydrolyzed to give the white solid product **Ib-18** (0.009 g) in two-step yield: 11.4%.

¹H-NMR for the **Ib-18** hydrochloride (400 MHz, CDCl₃) δ: 7.20-7.18 (m, 1H), 7.10-7.03 (m, 2H), 7.01 (m, 1H), 6.51-6.46 (m, 2H), 5.07 (s, 2H), 4.79-4.75 (m, 1H), 4.32-4.28 (m, 1H), 3.84-3.79 (m, 1H), 2.85-2.79 (m, 1H), 2.66-2.60 (m, 1H); ¹⁹F NMR (376 MHz, CDCl₃): δ -49.63.

ESI-MS [(M-H)⁺]: m/z of **Ib-18** calculated: 363.1, founded: 363.1.

Example 19

Synthesis of Compound **Ib-19**

The synthesis of compound **Ib-19** was carried out in the same way as in Example 1, and the product **Ib-19** was obtained by etherification and hydrolysis, in which compound **RM-Ib-17** (0.91 mmol) was used in the reaction instead of compound **RM-Ib-1** to obtain intermediate **RM-Ib-19**, and the light yellow solid product **Ib-19** (0.231 g) was obtained from the intermediate by hydrolysis. Two-step yield: 72.8%.

¹H-NMR for the **Ib-19**hydrochloride (400 MHz, CDCl₃) δ: 7.06-7.04 (m, 1H), 7.00-6.98 (m, 1H), 6.84-6.83 (m, 2H), 6.53-6.48 (m, 2H), 5.02 (s, 2H), 4.78-4.74 (m, 1H),

4.30-4.23 (m, 5H), 3.83-3.77 (m, 1H), 2.82-2.77 (m, 1H), 2.63-2.57 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of Iib-19 calculated: 341.1, founded: 341.0.

Example 20

5 Synthesis of compound **Iib-20**

The synthesis of compound **Iib-20** was carried out in the same way as in Example 1, and the product **Iib-20** was obtained by etherification and hydrolysis reaction, in which compound **RM-Ib-18** (0.64 mmol) was used in the reaction instead of compound **RM-Ib-1** to obtain intermediate **RM-Iib-20**, and the light yellow solid product **Iib-20** (0.166 g) was
10 obtained from the intermediate by hydrolysis. Two-step yield: 71.2%.

¹H-NMR for the **Iib-20**hydrochloride (400 MHz, CDCl₃)δ: 7.05-7.03 (m, 1H), 6.91-6.89 (m, 1H), 6.80-6.76 (m, 1H), 6.70-6.68 (m, 1H), 6.52-6.49 (m, 2H), 4.98 (s, 2H), 4.77-4.73 (m, 1H), 4.30-4.26 (m, 1H), 3.82-3.77 (m, 1H), 2.81-2.76 (m, 1H), 2.62-2.55 (m, 1H), 1.69 (s, 6H).

15 ESI-MS [(M-H)⁺]: m/z of Iib-20 calculated: 355.1, founded: 355.0.

Example 21

Synthesis of Compound **Iib-21**

The synthesis of compound **Iib-21** was carried out in the same way as in Example 1,
20 and the product **Iib-21** was obtained by etherification and hydrolysis, in which compound **RM-Ib-19** (0.57 mmol) was used in the reaction instead of RM-Ib-1 to obtain intermediate **RM-Iib-21**, and the light yellow solid product Iib-21 (0.072 g) was obtained from the intermediate by hydrolysis. Two-step yield: 28.4%.

ESI-MS [(M-H)⁺]: m/z of Iib-21calculated: 441.0, founded: 440.8.

25

Example 22

Synthesis of Compound **Iib-22**

The synthesis method for the preparation of compound Iib-22 was the same as that of Example 1, and the product **Iib-22** was obtained by etherification and hydrolysis reaction,
30 in which compound **RM-Ib-20** (0.43 mmol) was used in the reaction instead of compound **RM-Ib-1** to obtain intermediate **RM-Iib-22**, and light yellow oil **Iib-22** (0.150 g) was obtained from the intermediate by hydrolysis in two steps. Yield: 80%.

ESI-MS [(M-H)⁺]: m/z ofIib-22calculated: 435.0, founded: 434.9.

Example 23

Synthesis of Compound **Ib-23**

The synthesis of compound **Ib-23** was carried out in the same way as in Example 1, and the product **Ib-23** was obtained by etherification and hydrolysis, wherein compound **RM-Ib-21** (0.13 mmol) was used in the reaction instead of compound **RM-Ib-1** to obtain intermediate **RM-Ib-23**, and the light yellow solid product **Ib-23** (0.0033 g) was obtained from the intermediate by hydrolysis. Two-step yield: 6.2%.

ESI-MS [(M-H)⁺]: m/z of **Ib-23** calculated: 397.0, founded: 397.3.

Example 24

Synthesis of compound **Ib-24**

The synthesis of compound **Ib-24** was carried out in the same way as in Example 1, and the product **Ib-24** was obtained by etherification and hydrolysis, in which compound **RM-Ib-22** (0.38 mmol) was used in the reaction instead of **RM-Ib-1** to obtain intermediate **RM-Ib-24**, and the light yellow solid product **Ib-24** (0.028 g) was obtained from the intermediate by hydrolysis. Two-step yield: 18.8%.

ESI-MS [(M-H)⁺]: m/z of **Ib-24** calculated: 397.0, founded: 397.4.

Example 25

Synthesis of compound **Ib-25**

The synthesis of compound **Ib-25** was prepared in the same way as in Example 1, and the product **Ib-25** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-5** (0.54 mmol) was used in the reaction instead of compound **RM-Ib-1**, and compound **SM1-2** (0.54 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-Ib-25**, which was obtained from intermediate **Ib-25** (0.085 g) was obtained as a white solid product by hydrolysis in two-step yield: 44%.

¹H-NMR for the **Ib-25**hydrochloride (400 MHz, CDCl₃)δ: 7.07-7.05 (m, 1H), 6.94 (m, 1H), 6.78-6.77 (m, 1H), 6.50-6.45 (m, 2H), 6.02 (s, 2H), 4.95 (s, 2H), 4.79-4.74 (m, 1H), 4.31-4.27 (m, 1H), 3.84-3.79 (m, 1H), 2.84-2.79 (m, 1H), 2.66-2.59 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of **Ib-25** calculated: 361.0, founded: 361.0.

Example 26

Synthesis of Compound **Ib-26**

The synthesis of compound **Ib-26** was prepared as in Example 1, and the product **Ib-26** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-16** (0.22 mmol) was used in the reaction instead of compound **RM-Ib-1**, and compound **SM1-3** (0.22 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-Ib-26**, which was obtained from The intermediate was hydrolyzed to give the white solid product **Ib-26** (0.045 g) in two-step yield: 21%.

¹H-NMR for the **Ib-26** hydrochloride (400 MHz, CDCl₃) δ: 7.33 (s, 1H), 7.32-7.30 (m, 1H), 7.13-7.09 (m, 1H), 7.06-7.02 (m, 1H), 6.50 (s, 1H), 5.13 (s, 2H), 4.80-4.76 (m, 1H), 4.32-4.29 (m, 1H), 3.845-3.81 (m, 1H), 2.83-2.78 (m, 1H), 2.67-2.60 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of **Ib-26** calculated: 441.0, founded: 441.0.

Sample 27

Synthesis of Compound **Ib-27**

The synthesis of compound **Ib-27** was prepared as in Example 1, and the product **Ib-27** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-16** (0.22 mmol) was used in the reaction instead of compound **RM-Ib-1**, and compound **SM1-4** (0.22 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-Ib-27**, which was obtained from The intermediate was hydrolyzed to give the white solid product **Ib-27** (0.070 g) in two-step yield: 28%.

¹H-NMR for the **Ib-27** hydrochloride (400 MHz, CDCl₃) δ: 7.43-7.41 (m, 1H), 7.32 (s, 1H), 7.14-7.10 (m, 1H), 7.06-7.04 (m, 1H), 5.08 (s, 2H), 4.93-4.88 (m, 1H), 4.45-4.41 (m, 1H), 4.00-3.96 (m, 1H), 2.88-2.82 (m, 2H), 2.74-2.67 (m, 1H); ¹⁹F NMR (376 MHz, CDCl₃): δ -49.68.

ESI-MS [(M-H)⁺]: m/z of **Ib-27** calculated: 520.9, founded 520.9.

Example 28

Synthesis of Compound **Ib-28**

The synthesis of compound **Ib-28** was prepared in the same way as in Example 1, and the product **Ib-28** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-16** (0.48 mmol) was used in the reaction instead of compound **RM-Ib-1**, and compound **SM1-5** (0.48 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-Ib-28**, which was obtained from The intermediate was hydrolyzed to

give the white solid product **Iib-28** (0.070 g) in two-step yield: 36%.

¹H-NMR for the **Iib-28**hydrochloride (400 MHz, CDCl₃) δ: 7.29-7.26 (m, 1H), 7.18 (s, 1H), 7.12-7.08 (m, 1H), 7.04-7.02 (m, 1H), 6.50 (s, 2H), 5.14 (s, 2H), 4.80-4.75 (m, 1H), 4.32-4.28 (m, 1H), 3.84-3.80 (m, 1H), 2.83-2.77 (m, 1H), 2.66-2.60 (m, 1H).

5 ESI-MS [(M-H)⁺]:m/z of Iib-28 calculated:397.0, founded397.0.

Example 29

Synthesis of Compound **Iib-29**

The synthesis method for the preparation of compound **Iib-29** was the same as that of Example 1, and the product **Iib-29** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-16** (1.0 mmol) was used in the reaction instead of compound **RM-Ib-1** and compound **SM1-6** (1.0 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-Iib-29**, which was obtained from the intermediate by The white solid product **Iib-29** (0.070 g) was obtained by hydrolysis, two-step yield: 18%.

15 ESI-MS [(M-H)⁺]:m/z of Iib-29 calculated: was 381.1, founded 381.0.

Example 30

Synthesis of compound **Iib-30**

The reaction was completed by adding LiHMDS (1M, 1.4mL, 1.4mmol) dropwise and stirring for 0.5h. TMSCl (0.14g, 1.3mmol) was added dropwise and stirred for 0.5h, Add NBS (0.11g, 0.62mmol), slowly raise to room temperature, and react for 5h, the reaction is completed.HPLC showed that after the reaction, the reaction solution was added with water, neutralized with hydrochloric acid (1N) to a pH of about 2, extracted with ethyl acetate, and then the organic phases were combined, washed with saline and dried, and finally purified by column chromatography to give the light yellow solid product Iib-30 (0.055 g) in one-step yield: 24%.

25 ESI-MS [(M-H)⁺]: m/z of Iib-30calculated: 441.0, founded: 440.9.

Example 31

30 Synthesis of compound **Iib-31**

The reaction was completed by adding the raw material **Iib-30** (0.035 g, 0.08 mmol) to concentrated ammonia (3 mL) and heating in an oil bath at 100 degrees C for 2.5h.HPLC showed that the reaction solution was concentrated and dried under reduced

pressure to obtain the light yellow solid product **I Ib-31** (0.025g) with one-step yield: 83%.

ESI-MS [(M+H)⁺]: m/z of **I Ib-31** calculated: 380.1, founded: 380.0.

Example 32

5 Synthesis of Compound **I Ib-32**

Step 1:

RM-I Ib-17 (0.29 g, 0.78 mmol) was dissolved in 2 mL of THF, protected by nitrogen and cooled to -70°C in an acetone bath on dry ice, LiHMDS (1 M, 0.8 mL, 0.8 mmol) was added dropwise and stirred for 0.5 h. (PhSO₂)NF/THF (0.29 g, 0.93 mmol) solution was added dropwise, and the reaction was slowly increased to room temperature. The reaction was completed at room temperature for 2h. HPLC showed that after the reaction was finished, the reaction solution was added with water, neutralized with hydrochloric acid (1N) to a pH of about 2, extracted with ethyl acetate, then the organic phases were combined, washed with saline and dried, and finally purified by column chromatography to obtain intermediate RM-I Ib-30 (0.043g).

Step 2:

MeOH (2 mL), THF (1 mL) and aqueous LiOH (1N, 1 mL) were added sequentially to intermediate **RM-I Ib-30** and stirred at room temperature for 1 h. After the reaction was completed, the reaction solution was neutralized with hydrochloric acid (1N) to pH about 4, water was added, ethyl acetate was added for extraction, and then the organic phases were combined, washed with saline and dried, and finally purified by column chromatography. A white solid product **I Ib-32** (0.018 g) was obtained in two-step yield: 6%.

ESI-MS [(M-H)⁺]: m/z of **I Ib-32** calculated: 379.1, founded: 379.0.

25 Example 33

Synthesis of Compound **I Ib-33**

Step 1:

RM-I Ib-17 (0.24 g, 0.65 mmol) was dissolved in 2 mL of THF, protected by nitrogen and cooled to -70°C in an acetone bath on dry ice, LiHMDS (1 M, 0.8 mL, 0.8 mmol) was added dropwise, stirred for 0.5 h, and (PhSO₂)NF/THF (0.24 g, 0.78 mmol) solution was added dropwise, and the reaction was slowly increased to room temperature. The reaction was completed at room temperature for 2h. HPLC showed that after the reaction was finished, the reaction solution was added with water, neutralized with

hydrochloric acid (1N) to pH about 2, extracted by adding ethyl acetate, and then the organic phases were combined, washed with saline and dried, and finally purified by column chromatography to obtain intermediates **RM-IIb-31** and **RM-IIb-32** (0.045g).

Step 2:

5 MeOH (2 mL), THF (1 mL) and aqueous LiOH (1N, 1 mL) were added to intermediates **RM-IIb-31** and **RM-IIb-32** in turn and stirred at room temperature for 1 h. After the reaction was completed, the reaction solution was neutralized with hydrochloric acid (1N) to a pH of about 4, water was added, ethyl acetate was added for extraction, and then the organic phases were combined, washed with saline and dried. Finally, the reaction
10 was purified by column chromatography, and the light yellow solid products **I Ib-33** and **I Ib-34** (0.030 g) were obtained in two-step yield:12%.

ESI-MS [(M-H)⁺]: m/z of **I Ib-33** calculated 381.1, found: 381.0.

Example 34

15 Synthesis of compound **I Ib-34**

The synthesis of compound **I Ib-34** was prepared as in Example 1, and the product **I Ib-34** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-16** (0.36 mmol) was used in the reaction instead of compound **RM-Ib-1**, and compound **SM1-7** (0.36 mmol) was used instead of compound **SM1-1** to obtain intermediate
20 **RM-IIb-32**, which was obtained from intermediate was hydrolyzed to give the white solid product **I Ib-34** (0.021 g) in two-step yield: 14%.

ESI-MS [(M-H)⁺]: m/z of **I Ib-34** calculated: 399.0, founded: 399.0.

Example 35

25 Synthesis of compound **I Ib-35**

The synthesis method for the preparation of compound **I Ib-35** was the same as that of Example 1, and the product **I Ib-35** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-6** (0.27 mmol) was used in the reaction instead of compound **RM-Ib-1** and compound **SM1-8** (0.27 mmol) was used instead of compound
30 **SM1-1** to obtain intermediate **RM-IIb-33**, and from intermediate **I Ib-35** (0.056 g) was obtained as a light yellow solid product by hydrolysis in two-step yield: 60%.

¹H-NMR for the **I Ib-35** hydrochloride (400 MHz, CDCl₃) δ: 7.12-7.10 (m, 1H), 6.85 (m, 1H), 6.80-6.78 (m, 1H), 6.69-6.66 (m, 1H), 6.57-6.54 (m, 1H), 6.01 (s, 2H), 4.99 (s,

2H) 3.56-3.52 (m, 1H), 2.92-2.77 (m, 3H), 2.51-2.41 (m, 2H), 1.81-1.74 (m, 1H).

ESI-MS [(M-H)⁺]: m/z of **IIb-35** calculated 343.1 founded 343.0.

Example 36

5 Synthesis of compound **IIb-36**

The synthesis method for the preparation of compound **IIb-36** was the same as that of Example 1, and the product **IIb-36** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-6** (0.27 mmol) was used in the reaction instead of compound **RM-Ib-1**, and compound **SM1-9** (0.27 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-IIb-34**, and from intermediate **IIb-36** (0.072 g) was obtained as a light yellow solid product by hydrolysis in two-step yield: 75%.

¹H-NMR for the **IIb-36**hydrochloride (400 MHz, CDCl₃)δ: 7.31-7.26 (m, 2H), 6.93-6.91 (m, 2H), 6.69-6.68 (m, 1H), 6.66 (m, 1H), 6.01 (s, 2H), 4.99 (s, 2H), 2.64 (s, 2H), 1.45 (s, 6H).

15 ESI-MS [(M-H)⁺]: m/z of **IIb-36**calculated345.1, founded 345.0.

Example 37

Synthesis of compound **IIb-37**

The synthesis of compound **IIb-37** was prepared in the same way as in Example 1, and the product **IIb-37** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-6** (0.27 mmol) was used in the reaction instead of compound **RM-Ib-1**, and compound **SM1-10** (0.27 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-IIb-35**, which was obtained from The intermediate was hydrolyzed to give the light yellow solid product **IIb-37** (0.055 g) in two-step yield: 53%.

25 ¹H-NMR for the **IIb-37**hydrochloride (400 MHz, CDCl₃) δ: 7.41-7.39 (m, 2H), 6.95-6.92 (m, 2H), 6.67-6.63 (m, 1H), 6.57-6.55 (m, 1H), 6.03-6.01 (m, 2H), 4.99 (s, 2H), 4.13-4.10 (m, 2H), 3.83-3.82 (m, 2H), 2.98 (s, 2H).

ESI-MS [(M-H)⁺]: m/z of **IIb-37**calculated:375.1,founded:374.9.

30 Example 38

Synthesis of compound **IIb-38**

The synthesis of compound **IIb-38** was prepared as in Example 2, and the product **IIb-38** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-23**

(0.16 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-2** (0.16 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-IIb-36**, which was obtained from intermediate was hydrolyzed to give the white solid product **Ib-38** (0.040 g) in two-step yield:44%.

5 ESI-MS [(M-H)⁺]: m/z of **Ib-38** calculated:567.2,founded 567.0.

Example 39

Synthesis of Compound **Ib-39**

The synthesis of compound **Ib-39** was prepared as in Example 1, and the product
10 **Ib-39** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-5** (0.35 mmol) was used in the reaction instead of compound **RM-Ib-1**, and compound **SM1-7** (0.35 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-IIb-37**, which was obtained from intermediate **Ib-39** (0.021 g) was obtained as a white solid product by hydrolysis, two-step yield:15%.

15 ESI-MS [(M-H)⁺]: m/z of **Ib-39** calculated:397.0, founded:397.0.

Example 40

Synthesis of compound **Ib-40**

The synthesis of compound **Ib-40** was prepared as in Example 2, and the product
20 **Ib-40** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-24** (0.31 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-2** (0.31 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-IIb-38**, which was obtained from intermediate was hydrolyzed to give the white solid product **Ib-40** (0.067 g) in two-step yield:47%.

25 ESI-MS [(M-H)⁺]: m/z of **Ib-40** calculated:453.0, founded: 453.0.

Example 41

Synthesis of compound **Ib-41**

The synthesis of compound **Ib-41** was prepared as in Example 2, and the product
30 **Ib-41** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-25** (0.10 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-2** (0.10 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-IIb-39**, which was obtained from intermediate was hydrolyzed to give the white solid

product **Ib-41** (0.014 g) in two-step yield:36%.

ESI-MS [(M+H)⁺]: m/z of **Ib-41**calculated:372.1, founded 372.1.

Example 42

5 Synthesis of compound **Ib-42**

The synthesis of compound **Ib-42** was prepared as in Example 2, and the product **Ib-42** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-26** (0.13 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-2** (0.13 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-Ib-40**, which was obtained from intermediate was hydrolyzed to give the white solid product **Ib-42** (0.029 g) in two-step yield:58%.

ESI-MS [(M+H)⁺]: m/z of **Ib-42** calculated:372.1, founded:372.1

Example 43

15 Synthesis of compound **Ib-43**

The synthesis of compound **Ib-43** was prepared in the same way as in Example 2, and the product **Ib-43** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-27** (0.09 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-2** (0.09 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-Ib-41**, which was obtained from intermediate was hydrolyzed to give the white solid product **Ib-43** (0.012 g) in two-step yield: 31%.

¹H-NMR for the **Ib-43** hydrochloride (400 MHz, CDCl₃) δ: 7.06-7.03 (m, 2H), 6.73 (s, 1H), 6.49-6.46 (m, 2H), 5.96 (s, 2H), 4.95 (s, 2H), 4.78-4.73 (m, 1H), 4.30-4.26 (m, 1H), 3.82-3.78 (m, 1H), 2.83-2.77 (m, 1H), 2.65-2.58 (m, 1H) , 1.49 (s, 9H).

25 ESI-MS [(M-H)⁺]: m/z of **Ib-43**calculated: 442.1,founded: 442.2.

Example 44

Synthesis of Compound **Ib-44**

Compound **Ib-43** (0.017g,0.04mmol) was added to HCl/MeOH (6N,2mL) and stirred at room temperature for 1h, and the reaction was detected to be complete, and the white solid product **Ib-44** (0.015g) was obtained by concentration in one-step yield:91%.

ESI-MS [(M+H)⁺]: m/z of **Ib-44**calculated: 344.1, founded: 344.1.

Example 45

Synthesis of compound **I Ib-45**

The synthesis of compound **I Ib-45** was prepared as in Example 2, and the product **I Ib-45** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-28** (0.14 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-2** (0.14 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-I Ib-42**, which was obtained from intermediate was hydrolyzed to give the white solid product **I Ib-45** (0.035 g) in two-step yield:66%.

ESI-MS [(M-H)⁺]: m/z of **I Ib-45** calculated: 357.3,founded: 357.2.

Example 46

Synthesis of compound **I Ib-46**

The synthesis of compound **I Ib-46** was prepared as in Example 2, and the product **I Ib-46** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-29** (0.12 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-2** (0.12 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-I Ib-43**, which was obtained from intermediate was hydrolyzed to give the white solid product **I Ib-46** (0.030 g) in two-step yield:52%.

ESI-MS [(M-H)⁺]: m/z of **I Ib-46**calculated: 478.4,founded: 478.3.

Example 47

Synthesis of Compound **I Ib-47**

Step 1:

The compound **RM-I Ib-41** (0.6 g,1.31 mmol) was added to HCl/MeOH (6N,12 mL) and stirred overnight at room temperature, and the reaction was assayed to completion and concentrated to give the white solid intermediate **RM-I Ib-44** (0.5 g).

Step 2:

Dissolve the raw material **RM-I Ib-44** (0.050g, 0.13mmol) in 2mL of DCM, add DMAP (0.039g, 0.32mmol), and add methyl chloroformate (0.015g, 0.15mmol) under the cooling of ice water bath, stir the reaction overnight at room temperature, and the reaction was completed. the reaction solution was post-treated after the HPLC showed the end of the reaction The intermediate **RM-I Ib-45** (0.049g) was purified by column chromatography.

Step 3:

To intermediate **RM-IIb-45**, MeOH (2mL), THF (1mL) and aqueous LiOH (1N, 1mL) were added sequentially and stirred at room temperature for 1h, HPLC showed that the reaction solution was neutralized with hydrochloric acid (1N) to pH about 4, water was added, ethyl acetate was added for extraction, then the organic phases were combined, washed with saline and dried, and finally purified by column chromatography. The white solid product **Ib-47** (0.047 g) was obtained in two-step yield:92%.

ESI-MS [(M-H)⁺]: m/z of **Ib-47**calculated: 400.4,founded: 400.3.

Example 48

Synthesis of Compound **Ib-48**

Step 1:

RM-IIb-44 (0.050 g, 0.13 mmol) was dissolved in 2 mL of DCM, DMAP (0.039 g, 0.32 mmol) was added and cyclopentyl chloroformate (0.023 g, 0.15 mmol) was added under cooling in an ice water bath and the reaction was completed by stirring at room temperature overnight.The HPLC showed that the reaction solution was purified by post-treatment column chromatography after the reaction was finished, and intermediate **RM-IIb-46** (0.050g) was obtained.

Step 2:

To intermediate **RM-IIb-46**, MeOH (2mL), THF (1mL) and aqueous LiOH (1N, 1mL) were added sequentially and stirred at room temperature for 1h, HPLC showed that the reaction solution was neutralized with hydrochloric acid (1N) to pH about 4, water was added, ethyl acetate was added for extraction, then the organic phases were combined, washed with saline and dried, and finally purified by column chromatography. A white solid product **Ib-48** (0.040 g) was obtained in two-step yield:70%.

ESI-MS [(M-H)⁺]: m/z of **Ib-48** calculated:454.5, founded:454.5.

Example 49

Synthesis of Compound **Ib-49**

Step 1:

The reaction was completed by dissolving **RM-IIb-44** (0.050g, 0.13mmol) in 2mL of DCM, adding DMAP (0.039g, 0.32mmol), cooling in an ice water bath, adding phenyl chloroformate (0.024g, 0.15mmol), and stirring the reaction overnight at room temperature.

The intermediate (0.050g) was obtained by column chromatography separation and purification.

Step 2:

5 The intermediate was redissolved in 2 mL of DMF, DMAP (0.025 g, 0.2 mmol) and tert-butylamine (0.015 g, 0.2 mmol) were added, and the reaction was stirred at room temperature for 3 h. The reaction was purified by post-treatment column chromatography after the reaction was shown by HPLC, and intermediate **RM-IIb-47** (0.040 g) was obtained.

Step 3:

10 To intermediate **RM-IIb-47**, MeOH (2mL), THF (1mL) and aqueous LiOH (1N, 1mL) were added sequentially and stirred at room temperature for 1h, HPLC showed that the reaction solution was neutralized with hydrochloric acid (1N) to pH about 4, water was added, ethyl acetate was added for extraction, and then the organic phases were combined, washed with saline and dried, and finally purified by column chromatography. A white
15 solid product **Ib-49** (0.020 g) was obtained in three-step yield:35%.

ESI-MS [(M-H)⁺]: m/z of **Ib-49**calculated: 441.5, founded:441.4.

Example 50

Synthesis of Compound **Ib-50**

20 Step 1:

RM-IIb-44 (0.050g, 0.13mmol) was dissolved in 2mL of DCM, DMAP (0.039g, 0.32mmol) was added, and cyclopentyl chloroformate (0.023g, 0.15mmol) was added under cooling in an ice water bath, and the reaction was stirred at room temperature overnight, and the reaction was completed.After the reaction was completed as indicated
25 by HPLC, the reaction solution was post-treated The intermediate **RM-IIb-48** (0.045g) was purified by column chromatography.

Step 2:

MeOH (2mL), THF (1mL) and aqueous LiOH (1N, 1mL) were added sequentially to intermediate **RM-IIb-48** and stirred at room temperature for 1h, HPLC showed that the
30 reaction solution was neutralized with hydrochloric acid (1N) to pH about 4, water was added, ethyl acetate was added for extraction, then the organic phases were combined, washed with saline and dried, and finally purified by column chromatography. A white solid product **Ib-50** (0.020 g) was obtained in two-step yield:46%.

ESI-MS [(M-H)⁺]: m/z of I Ib-50 calculated:426.7, founded: 426.4.

Example 51

Synthesis of Compound **I Ib-51**

Step 1:

RM-I Ib-44 (0.050 g, 0.13 mmol) was dissolved in 2 mL of DCM, DMAP (0.039 g, 0.32 mmol) was added, and the reaction was completed by stirring at room temperature overnight under cooling in an ice water bath. The intermediate **RM-I Ib-49** (0.015g) was purified by column chromatography.

Step 2:

MeOH (2mL), THF (1mL) and aqueous LiOH (1N, 1mL) were added sequentially to intermediate **RM-I Ib-49** and stirred at room temperature for 1h, HPLC showed that the reaction solution was neutralized with hydrochloric acid (1N) to pH about 4, water was added, ethyl acetate was added for extraction, then the organic phases were combined, washed with saline and dried, and finally purified by column chromatography. The white solid product **I Ib-51** (0.009 g) was obtained in two-step yield:16%.

ESI-MS [(M-H)⁺]: m/z of I Ib-51 calculated: 448.5, founded:448.3.

Example 52

Synthesis of Compound **I Ib-52**

The synthesis of compound **I Ib-52** was prepared as in Example 2, and the product **I Ib-52** was obtained by etherification and hydrolysis reaction, where compound **RM-I Ib-30** (0.06 mmol) was used in the reaction instead of compound **RM-I Ib-2**, and compound **SM1-2** (0.06 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-I Ib-50**, which was obtained from intermediate was hydrolyzed to give the white solid product **I Ib-52** (0.004g) in two-step yield:15%.

ESI-MS [(M-H)⁺]: m/z of I Ib-52 calculated: 411.1, founded: 411.2.

Example 53

Synthesis of compound **I Ib-53**

The synthesis of compound **I Ib-53** was prepared as in Example 2, and the product **I Ib-53** was obtained by etherification and hydrolysis reaction, where compound **RM-I Ib-31** (0.24 mmol) was used in the reaction instead of compound **RM-I Ib-2**, and compound

SM1-2 (0.24 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-IIb-51**, which was obtained from intermediate was hydrolyzed to give the white solid product **Ib-53** (0.018 g) in two-step yield:19%.

ESI-MS [(M-H)⁺]: m/z of Ib-53 calculated: 395.1, founded: 395.2.

5

Example 54

Synthesis of Compound **Ib-54**

The synthesis of compound **Ib-54** was prepared as in Example 2, and the product **Ib-54** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-32** (0.24 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-2** (0.24 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-IIb-52**, which was obtained from intermediate was hydrolyzed to give the white solid product **Ib-54** (0.028 g) in two-step yield:27%.

ESI-MS [(M-H)⁺]: m/z of Ib-54 calculated: 431.0, founded:431.1.

15

Example 55

Synthesis of compound **Ib-55**

The synthesis of compound **Ib-55** was prepared as in Example 2, and the product **Ib-55** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-33** (0.24 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-2** (0.24 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-IIb-53**, which was obtained from intermediate was hydrolyzed to give the white solid product **Ib-55** (0.020 g) in two-step yield:22%.

ESI-MS [(M-H)⁺]: m/z of Ib-55 calculated: 381.1, founded:381.2.

25

Example 56

Synthesis of Compound **Ib-56**

The synthesis of compound **Ib-56** was prepared in the same way as in Example 2, and the product **Ib-56** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-34** (0.24 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-2** (0.24 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-IIb-54**, which was obtained from intermediate was hydrolyzed to give the white solid product **Ib-56** (0.028 g) in two-step yield:30%.

30

ESI-MS [(M-H)⁺]: m/z of Iib-56 calculated: 397.0, founded:397.1.

Example 57

Synthesis of Compound **Iib-57**

- 5 The synthesis of compound **Iib-57** was prepared in the same way as in Example 2, and the product **Iib-57** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-35** (0.23 mmol) was used in the reaction instead of compound **RM-Ib-2** and compound **SM1-11** (0.23 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-Iib-55** from intermediate was hydrolyzed to give the white solid product
- 10 **Iib-57** (0.018 g) in two-step yield:21%.

ESI-MS [(M-H)⁺]: m/z of Iib-57 calculated363.1, founded:363.2.

Example 58

Synthesis of compound **Iib-58**

- 15 The synthesis of compound Iib-58 was prepared as in Example 2, and the product Iib-58 was obtained by etherification and hydrolysis reaction, where compound RM-Ib-36 (0.23 mmol) was used in the reaction instead of compound RM-Ib-2, and compound SM1-11 (0.23 mmol) was used instead of compound SM1-1 to obtain intermediate RM-Iib-56, which was obtained from intermediate was hydrolyzed to give the white solid
- 20 product Iib-58 (0.026 g) in two-step yield:30%.

ESI-MS [(M-H)⁺]: m/z of Iib-58 calculated:379.0, founded: 379.1.

Example 59

Synthesis of Compound **Iib-59**

- 25 The synthesis of compound **Iib-59** was prepared as in Example 2, and the product **Iib-59** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-27** (0.23 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-11** (0.23 mmol) was used instead of compound SM1-1 to obtain intermediate **RM-Iib-57**, which was obtained from intermediate was hydrolyzed to give the white solid
- 30 product **Iib-59** (0.016 g) in two-step yield:30%.

ESI-MS [(M-H)⁺]: m/z of Iib-59 calculated:460.1, founded: 460.2.

Example 60

Synthesis of compound **I Ib-60**

The synthesis of compound **I Ib-60** was prepared in the same way as in Example 2, and the product **I Ib-60** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-27** (0.24 mmol) was used in the reaction instead of compound **RM-Ib-2** and compound **SM1-8** (0.24 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-I Ib-58**, which was obtained from intermediate was hydrolyzed to give the white solid product **I Ib-60** (0.021 g) in two-step yield:20%.

ESI-MS [(M-H)⁺]: m/z of I Ib-60 calculated: 442.2, founded: 442.1.

Example 61

Synthesis of compound **I Ib-61**

The synthesis of compound **I Ib-61** was prepared in the same way as in Example 1, and the product **I Ib-61** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-16** (0.24 mmol) was used in the reaction instead of compound **RM-Ib-1**, and compound **SM1-8** (0.24 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-I Ib-59**, which was obtained from The intermediate was hydrolyzed to give the white solid product **I Ib-61** (0.026 g) in two-step yield: 30%.

ESI-MS [(M-H)⁺]: m/z of I Ib-61 calculated: 361.1, founded: 361.2.

Example 62

Synthesis of compound **I Ib-62**

The synthesis of compound **I Ib-62** was prepared in the same way as in Example 1, and the product **I Ib-62** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-16** (0.24 mmol) was used in the reaction instead of compound **RM-Ib-1**, and compound **SM1-12** (0.24 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-I Ib-60**, which was obtained from The intermediate was hydrolyzed to give the white solid product **I Ib-62** (0.015 g) in two-step yield: 16%.

ESI-MS [(M-H)⁺]: m/z of I Ib-62 calculated:373.1, founded: 373.1.

Example 63

Synthesis of compound **I Ib-63**

The synthesis of compound **I Ib-63** was prepared in the same way as in Example 2, and the product **I Ib-63** was obtained by etherification and hydrolysis reaction, where

compound **RM-Ib-29** (0.24 mmol) was used in the reaction instead of compound **RM-Ib-2**, and compound **SM1-8** (0.24 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-Ib-61**, which was obtained from The intermediate was hydrolyzed to give the white solid product **Ib-63** (0.020 g) in two-step yield: 17%.

5 ESI-MS [(M-H)⁺]: m/z of Ib-63 calculated: 476.2, founded 476.2.

Sample 64

Synthesis of compound **Ib-64**

The synthesis method for the preparation of compound Ib-64 was the same as that of
10 Example 2, and the product **Ib-64** was obtained by etherification and hydrolysis reaction, where compound **RM-Ib-29** (0.24 mmol) was used in the reaction instead of compound **RM-Ib-2** and compound **SM1-12** (0.24 mmol) was used instead of compound **SM1-1** to obtain intermediate **RM-Ib-62**, which was obtained from The intermediate was hydrolyzed to give the white solid product Ib-64 (0.015 g) in two-step yield: 13%.

15 ESI-MS [(M-H)⁺]: m/z of Ib-64 calculated: 488.2, founded:488.3.

Example 65

Assay for GPR40 activity of compounds:

The compounds prepared herein can be initially screened for their effect on GPR40
20 by the following preclinical in vitro inhibition assays, and then further confirmed in clinical trials for efficacy and safety. Other methods will be apparent to those of ordinary skill in the art. The preclinical and clinical trial results study of TAK-875, a GPR40 agonist previously developed by Takeda, Japan, showed that the results of the ex vivo test were consistent with the results of the relevant in vivo activity test.

25 The compounds of the present invention, or stereoisomers, tautomer, esterified or amidated precursor drugs, or pharmaceutically acceptable salts thereof, and mixtures thereof, were tested by evaluation experiments of the activity of compounds Ib-01 to Ib-64 and a reference compound Ref-1 (TAK-875) against GPR40, and comparison of the test results revealed that a number of compounds showed better inhibitory activity (EC50)
30 than the reference compound.

Inoculate CHO-K1/GPR40 cells in 384-well plates at a density of 10,000 cells/well. Cells were incubated at 37°C for 24 hours in a 5% CO₂ incubator. After the experiment, discard the cell culture medium, quickly add 30 µl of 1X FLIPR® Calcium 6 dye

containing 2.5 mM Probenecid to each well and incubate at 37°C for 2 hours protected from light. For the determination, different concentrations of drug (15 µl/well) were added to the wells and the fluorescence values were read. The fluorescence excitation wavelength was 494 nm and emission wavelength was 516 nm. FLIPR instrument data, EC₅₀ was derived from the formula $Y = \text{Bottom} + (\text{Top} - \text{Bottom}) / (1 + (\text{EC}_{50}/X)^{\text{HillSlope}})$.

The results of the GPR40 agonist activity tests for each compound of novel structure of Formula IIb are listed in Table 3 and Table 2 below; where the GPR40 agonist activity effect range (EC₅₀) of the compounds of the present invention is labeled "A" for <10 nM; "B" for the activity range of 10-100 nM; and "C" for the activity range >100 nM.

Table 3: Results of GPR40 agonist activity assay for compounds of formula IIb

No.	Compound	GPR40 activity	No.	Compound	GPR40 activity
1	IIb-01	C	2	IIb-02	C
3	IIb-03	C	4	IIb-04	B
5	IIb-05	B	6	IIb-06	B
7	IIb-07	B	8	IIb-08	C
9	IIb-09	C	10	IIb-10	C
11	IIb-11	C	12	IIb-12	C
13	IIb-13	C	14	IIb-14	B
15	IIb-15	B	16	IIb-16	C
17	IIb-17	B	18	IIb-18	A
19	IIb-19	C	20	IIb-20	B
21	IIb-21	B	22	IIb-22	C
23	IIb-23	B	24	IIb-24	C
25	IIb-25	A	26	IIb-26	C
27	IIb-27	C	28	IIb-28	C
29	IIb-29	C	30	IIb-30	C
31	IIb-31	C	32	IIb-32	C
33	IIb-33	B	34	IIb-34	C

35	I Ib-35	B	36	I Ib-36	C
37	I Ib-37	C	38	I Ib-38	B
39	I Ib-39	C	40	I Ib-40	B
41	I Ib-41	C	42	I Ib-42	C
43	I Ib-43	A	44	I Ib-44	C
45	I Ib-45	C	46	I Ib-46	A
47	I Ib-47	C	48	I Ib-48	B
49	I Ib-49	C	50	I Ib-50	C
51	I Ib-51	C	52	I Ib-52	C
53	I Ib-53	B	54	I Ib-54	B
55	I Ib-55	B	56	I Ib-56	B
57	I Ib-57	B	58	I Ib-58	B
59	I Ib-59	B	60	Ref-1	B
61	I Ib-61	B	62	I Ib-62	B
63	I Ib-63	B	64	I Ib-64	B
65	Ref-1	B			

Example 66

Compound toxicity screening test

In order to test the toxicity of some of the more active (EC₅₀ <10nM) compounds (I Ib-18, I Ib-25 and I Ib-43) of the new compounds (I Ib-01 to I Ib-64), the Acute Toxicity Study (MTD) test was conducted using a single dose of 600mg/kg administered by gavage to healthy mice at 18-22g for 5 consecutive days to assess the toxicity of the subject to the organism, the results showed that the overall toxicity of the compounds was low (LD₅₀: >600), the survival rate of the mice after administration was 100%, no abnormalities in body weight and external and internal observations occurred during the dosing and 5-day recovery period, and the autopsy results did not reveal any abnormal changes in various organs such as heart, liver, lung, kidney and intestine. To this end, several highly active compounds of the present invention have been determined by in vitro experiments to be not only more effective in type 2 diabetes, but also safe and reliable.

Example 67

BSEP (Bile Salt Export Pump) assay for bile salt transport effect of compounds:

In this experiment, we performed a preliminary study on the inhibition of the uptake
5 of the substrate taurocholic acid (TCA) by the BSEP bile salt transporter by LC/MS/MS. The BSEP protein microvesicles with reverse uptake ability were obtained by transfecting Hi5 cells with the BSEP gene, which is capable of TCA uptake and uptake of taurocholic acid. The principle of this experiment is to perform substrate uptake by BSEP protein microcapsules in the presence of energy ATP, the substrate TCA and the candidate
10 compound to be tested, after a certain transit time, the energy, substrate and candidate compound were completely washed out of the microvesicle, and the absorbed substrate TCA was released by lysis of the microvesicle.

Sample processing:

1. Dissolve compounds individually to 100 mM using 100% DMSO and store in a
15 -20 degree refrigerator with all compounds completely dissolved.

2. The starting concentration of the sample was 100 mM, diluted in 2-fold gradient with Bravo apparatus, with 11 concentration gradients; the lowest point concentration was 97.65 μ M.

3. Control compound (Glyburide) starting at 20 mM, diluted in 2-fold gradient with
20 Bravo instrument, with 11 concentration gradients; the lowest point concentration was 19.53 μ M.

4. 300nl of compound was transferred to the middle plate using the ECHO autosampler.

5. Positive control (HPE) wells and negative control (ZPE) wells were also
25 transferred 300nl DMSO each.

1. Add 14.7 μ l of ATP buffer to the corresponding wells of the compound and ZPE, respectively.

2. Add 14.7 μ l of AMP buffer to the corresponding wells of the HPE.

3. Shake the plate at 25 degrees C for 10 minutes.

4. Add 15ul of BSEP-Hi5-VT Vesicle solution to all wells separately and shake the
30 plate for 40 minutes at 25 degrees Celsius.

5. Immediately add 5ul of 0.5M EDTA solution to all wells, and then add 65 μ l of Buffer B. All reactions are completed.

6. Transfer 95 µl of liquid from the end of reaction compound plate to the filter plate using the spiking apparatus.

7. After placing the liquid receiving plate under the filter plate, use a centrifuge to remove the liquid and discard the liquid received in the receiving plate.

5 8. Add 90 µl of buffer B to the filter plate and repeat step 7. Repeat steps 7 and 8 to wash the filter plate a total of three times.

9. Let the filter plate dry overnight.

10. Add 81 µl of 80% methanol/water solution to the filter plate the next day.

11. Apply membrane to the filter plate and vibrate the plate for 15 minutes.

10 12. Place a new liquid receiver plate under the filter plate and centrifuge for 5 min to filter all the liquid from the filter plate into the receiver plate.

13. Add 13.5 µl of internal standard solution to each well of the liquid receiver plate and seal the plate with a sealing film.

14. Using LC/MS/MS, detect the content of taurocholic acid in the receiver plate.

15 Table 4: Comparison of optimized GPR40 agonist activity and its potential side effects

No.	Compound	GPR40 Activity (hEC ₅₀ , nM)	Potential inhibitory side effects BSEP (IC ₅₀ , µM)
1	I Ib-18	7.9	161.7
2	I Ib-25	9.2	57.7
3	I Ib-43	6.8	66.8
4	Ref-1 (TAK-875)	13.7	7.1

Note: The BSEP inhibition results (IC₅₀) for the compounds shown in Table 4 are below 25 µM [e.g., Ref-1 (TAK-875) has a value of 7.1, which is significantly below 25],
 20 then the BSEP inhibition side effects of the compound will likely lead to a potential risk of liver damage in vivo.

The results of the activity assays in Tables 3 and 4 above show that: (1) the activity

(EC50) of the five-membered heterocyclic compounds containing 1-2 oxygen atoms among the various novel benzo-heterocyclic compounds of the present invention is significantly better than that of the six-membered heterocyclic compounds of similar structures containing 1-2 oxygen atoms;; (2) The activity of the benzoxygen-containing five-membered heterocyclic compounds of the present invention when Z1 is "Oxygen(O)" is better than that of the corresponding compound when Z1 is "CH2" in the same structure;(3) The benzoxygen-containing five-membered heterocyclic compounds I Ib-18, I Ib-25, I Ib-43 and I Ib-46 of the present invention show good activity (EC50: <10nM) against the GPR40 target, which is better than that of the similar control compound TAK-875, and are novel GPR40 agonists with good activity and safety advantages in this research field.

The three benzo oxygen-containing five-membered heterocyclic compounds listed in Table 4 above not only showed better activity but also had BSEP side effect results greater than 25 that could lead to liver damage, which was significantly better than the new drug TAK-875, which was terminated from the Phase III clinical trial by Takeda, Japan, because of liver toxicity problems, and the results of the completed Acute Toxicity Study (MTD) test showed that the three highly active compounds I Ib-18, I Ib-25 and I Ib-43 listed in Table 4 had a better safety profile, the survival rate of the mice was 100% after 5 days of instillation of the new compound at a dose of 600 mg/kg, and no abnormalities occurred during the dosing and 5-day recovery period, and the autopsy results did not reveal any abnormalities. Therefore, several highly active compounds in the novel benzo oxygen-containing five-membered heterocyclic compound I Ib designed and synthesized by the present invention are valuable for further animal and clinical testing and promotion of application.

As a result, several compounds with better activity (e.g., **I Ib-18**, **I Ib-25** and **I Ib-43**) were identified in the innovative study of GPR40 target agonists, and the acute toxicity of the TMD high dose (600 mg/kg/day) test was very low, and the results of BSEP (>25uM), hERG (>30uM) and Ames assays showed no relevant side effects, and the relevant results were all superior to the currently known reference compound TAK-875, the results were better than those of the currently known reference compound TAK-875 and provided a basis for the invention of a new GPR40 target agonists with better GPR40 target selectivity and safe and efficient treatment of type II diabetic patients.


These and other changes to the embodiments may be made in accordance with the

detailed description above. In general, the terms used in the claims should not be considered as limiting the claims to the specific embodiments disclosed in this specification and claims, but rather as including all possible embodiments that follow the full scope of the equivalent of the listed claims. Accordingly, the claims are not limited by
5 this disclosure.

CLAIMS:

What is claimed is:

1. A compound represented by formula **Ib**, or its cis-trans isomer, enantiomer, diastereoisomer, racemate, tautomer, solvate, hydrate, or pharmaceutically acceptable salt, or mixtures thereof;



wherein, $n = 0, 1$ or 2 ;

when $n = 0$, Y does not exist, and Y^1 is directly single bonded to Z^1 adjacent (ortho-) to Y to form a five-membered heterocyclic group;

When E is not directly connected to G^1 to form a cyclic compound, E is selected from: hydrogen, deuterium(D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, carboxyl, amino(NH_2), aminocarbonyl(H_2NCO), alkyl, alkoxy, alkoxycarbonyl, alkylcarbonylamino, alkylaminocarbonyl, aryl, aryloxy, or heterocyclic aryl group; G^1 is CH, or C(Rb), wherein Rb is hydrogen, deuterium(D), alkyl, cycloalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, alkoxycarbonyl, alkylaminyl, cycloalkylaminyl, alkylaminylcarbonyl, cycloalkylaminylcarbonyl, aryl, aryloxy, heterocyclic, heterocyclic aryl, or heterocyclic aryloxy, or Rb and R^4 may be interconnected to form a cycloalkyl, or heterocyclic group;

When E is linked to G^1 to form a cyclic compound, G^1 is -C-; E is -O-, -C(RcRd)-, -OC(RcRd)-, -C(RcRd)O-, or -NRe-; wherein Rc and Rd are hydrogen, deuterium(D), alkyl, cycloalkyl, alkenyl, hydrocarbon, alkoxy, cycloalkoxy, alkoxycarbonyl, alkylaminyl, cycloalkylaminyl, alkylaminylcarbonyl, cycloalkylaminylcarbonyl, aryl, aryloxy, heterocyclic, heterocyclic aryl, or heterocyclic aryloxy, respectively, and Rc and Rd may be interconnected to form cycloalkyl, or heterocyclic groups; Re is hydrogen, deuterium(D), alkyl, cycloalkyl, alkylcarbonyl, alkoxycarbonyl, cycloalkoxycarbonyl, alkylaminylcarbonyl, cycloalkylaminylcarbonyl, alkylsulfonyl, or aryl sulfonyl group; L^1 and L^2 are each independently -O-, -S-, -C(O)-, -S(O) $_2$ -, -CH $_2$ -, -C(RfRg)-, -OC(RfRg)-, -C(RfRg)O-, -N(Re)-, -N(Rc)C(RfRg)-, or -C(RfRg)N(Re)-; wherein Rf and Rg are hydrogen, deuterium(D), alkyl, cycloalkyl, alkenyl, alkynyl, alkoxy, cycloalkoxy, alkoxycarbonyl, alkylaminyl, cycloalkylaminyl, alkylaminylcarbonyl, cycloalkylaminylcarbonyl, aryl,

aryloxy, heterocyclic, heterocyclic aryl, or heterocyclic aryloxy, respectively, and Re is defined as described above for Re in E;

R¹, R² and R³ are each independently hydrogen, deuterium(D), halogen, trifluoromethyl, trifluoromethoxy, nitrile, hydroxyl, aminocarbonyl (H₂NCO), alkyl, alkoxy, alkoxy carbonyl, alkylaminocarbonyl, alkylcarbonylamino, aryl, aryloxy, or heterocyclic aryl group; wherein R² and L¹ in formula **1b** can be interconnected to form a 4-8-membered heterocyclic compound, and L¹ is -CH-;

R⁴, R⁵ and R^{5b} are each independently hydrogen, deuterium(D), halogen, hydroxyl, amino, nitrile, alkyl, alkoxy, cycloalkyl, heterocycloalkyl, cycloalkoxy, optionally substituted alkenyl, optionally substituted alkynyl, alkoxy carbonyl, alkylaminocarbonyl, alkylcarbonylamino, alkoxy carbonylamino, aryl, aryloxy, or heterocyclic aryl; wherein R⁵ and R^{5b} may be interconnected to become cycloalkyl, heterocyclic, or heterocyclic aryl group;

R⁶ is a carboxyl, alkoxy carbonyl, aryloxy carbonyl, alkylaminyl carbonyl, cycloalkylaminyl carbonyl, alkylsulfonamidocarbonyl, cycloalkylsulfonamido carbonyl, heterocyclic, or heterocyclic aryl; or R⁶ and the adjacent (ortho-) substituent R⁵ may be interconnected to become heterocyclic, or heterocyclic aryl;


X⁵, X⁶ and X⁷ are each independently hydrogen, deuterium(D), halogen, nitrile, amino, trifluoromethyl, trifluoromethoxy, aminocarbonyl (H₂NCO), alkyl, heterocycloalkyl, alkoxy, heteroatomically substituted alkoxy, alkylamino (NR_iR_j), heteroatomically substituted alkylamino, alkoxy carbonyl, alkylaminocarbonyl, alkylcarbonylamino, alkoxy carbonylamino, cyclo alkoxy carbonylamino group, alkyl sulfonamido group, cycloalkyl sulfonamido group, aryl, aryloxy, aryl amido carbonyl, aryl carbonyl amido, aryloxy carbonyl amido, heterocyclic aryl, heterocyclic aryloxy, or heterocyclic aryl amido group; wherein R_i and R_j are each independently hydrogen, deuterium(D), alkyl, heterocycloalkyl, alkylcarbonyl, alkoxy carbonyl, cycloalkoxy carbonyl, alkylaminocarbonyl, alkylsulfonyl, cycloalkylsulfonyl, aryl, aryloxy carbonyl, arylaminocarbonyl, heterocycloaryl group, or R_i and R_j are interconnected to form a 3-8-membered heterocyclic group containing 1-3 heteroatoms;

Y and Y¹ are each independently -O-, -S-, -CH₂-, -CHF-, -CF₂-, -CCl₂-, -C(R_fR_g-, or -N(Re)-; wherein R_f and R_g are defined as described above for R_f and R_g in L¹, respectively, and Re is defined as described above for Re in E;

Z and Z¹ are each independently selected from: -O-, -S-, -CH₂-, -CHF-, -CF₂-, -C

(R_fR_g)-, -N(Re)-, or -C(O)-; wherein R_f and R_g are defined as described above for R_f and R_g in L¹, respectively, and Re is defined as described above for Re in E.

2. The compound according to claim 1, or its cis-trans isomer, enantiomer,
5 diastereoisomer, racemate, tautomer, solvate, hydrate, or a pharmaceutically acceptable salt, or a mixture thereof, wherein the compound has the structure of **IIb**;



wherein,

- The definitions of n, E, G¹, L¹, R¹, R², R³, R⁴, R⁵, R^{5b}, X⁵, X⁶, X⁷, Y, Y¹, Z are the
10 same as the definitions of n, E, G¹, L¹, R¹, R², R³, R⁴, R⁵, R^{5b}, X⁵, X⁶, X⁷, Y, Y¹, Z in claim 1;

- R⁷ is a hydroxyl, an alkoxy, an alkyl amino, a cycloalkyl amino, a heterocyclic amino,
an alkyl sulfonamido, a cycloalkyl sulfonamido, an aryloxy, a heterocyclic aryloxy, an aryl
amino, or a heterocyclic aryl amino; or R⁷ and the adjacent (ortho-) substituent R⁵ can be
15 interconnected to become a heterocyclic group.

3. The compound according to claim 2, wherein,

n = 0 or 1;

- When n = 0, Y does not exist and Y¹ is directly single bonded to the oxygen adjacent
20 (ortho-) to Y to form a five-membered heterocyclic group;

when n = 1, Y is -CH₂-;

- When E is not directly connected to G¹ to form a cyclic compound, E is hydrogen,
halogen, trifluoromethyl, trifluoromethoxy, or alkoxy; and G¹ is -CH-, or -C(Rb)-, wherein
Rb is hydrogen, alkyl, optionally substituted alkenyl, optionally substituted alkynyl, alkoxy,
25 or Rb and R⁴ are interconnected to form an oxygen-containing heterocyclic group; R⁴ is
hydrogen, alkyl, alkoxy, optionally substituted alkynyl, or R⁴ and Rb are interconnected
into an oxygen-containing heterocyclic group;

When E is directly connected to G¹ to form a cyclic compound, E is -OC(RcRd)-, G¹
is -C-, and R⁴ is hydrogen, where Rc and Rd are each independently hydrogen;

L¹ is -CH₂-, or L¹ is -OCH- when R² and L¹ in Formula **IIb** are interconnected to form a 5~6-membered heterocyclic compound;

R¹, R² and R³ are each independently hydrogen, halogen, alkyl or alkoxy group;

R⁵ is hydrogen, halogen, hydroxyl, amino, alkylamino, alkyl or alkoxy group;

5 R^{5b} is hydrogen, halogen, alkyl or alkoxy group;

R⁷ is an alkoxy, hydroxy, alkyl sulfonamido, or cycloalkyl sulfonamido group;

X⁵ is hydrogen, deuterium(D), halogen, nitrile, amino, trifluoromethyl, trifluoromethoxy, alkyl, alkoxy, alkylamino (NR_iR_j), alkylcarbonylamino, alkylaminocarbonylamino, alkoxycarbonylamino, cycloalkoxycarbonylamino, alkylsulfonamido, cycloalkylsulfonamido, aryl, or aryloxycarbonylamino group; wherein R_i and R_j are each independently hydrogen, alkyl, alkylcarbonyl, alkoxycarbonyl, or cycloalkoxycarbonyl group;

X⁶ and X⁷ are each independently hydrogen, deuterium(D), halogen, C₁-C₈ alkylamine group or C₁-C₈ alkoxycarbonylamino group;

15 Y is -CH₂-, or Y does not exist when n = 0;

Y¹ is -CH₂-, -CHF-, -CF₂-, or -C(CH₃)₂-;

Z is -O-, or -CH₂-.

4. The compound according to claim 3, wherein,

20 When n = 0, Y does not exist;

When E is not directly connected to G¹ to form a cyclic compound, E is hydrogen, or halogen; G¹ is -CH-, or -C(Rb)-, wherein Rb is alkyl, alkenyl, alkynyl, or alkoxy; and R⁴ is hydrogen, alkyl, or alkoxy group;

When E is directly connected to G¹ to form a cyclic compound, E is -OC(RcRd)-, G¹ is -C-, and R⁴ is hydrogen, wherein Rc and Rd are each independently hydrogen;

L¹ is -CH₂-, or L¹ is -OCH- when R² and L¹ in Formula **IIb** are interconnected to form a 5-6-membered heterocyclic compound;

R¹, R² and R³ are each independently hydrogen, halogen, or alkoxy group;

R⁵ is hydrogen, halogen, hydroxyl, amino, alkyl or alkoxy group;

30 R^{5b} is hydrogen, halogen, alkyl or alkoxy;

R⁷ is selected from: alkoxy, hydroxy, alkyl sulfonamido, or cycloalkyl sulfonamido group;

X⁵ is hydrogen, deuterium(D), halogen, nitrile, amino, trifluoromethyl,

trifluoromethoxy, alkyl, alkoxy, alkylamino (NR_iR_j), alkylcarbonylamino, alkylaminocarbonylamino, alkoxycarbonylamino, cycloalkoxycarbonylamino, alkylsulfonamido, cycloalkylsulfonamido, aryl, or aryloxy carbonylamino group; wherein R_i and R_j are each independently hydrogen, alkyl, alkylcarbonyl, alkoxycarbonyl, or cycloalkoxycarbonyl group;


X^6 is hydrogen, deuterium(D), halogen, $\text{C}_1\text{-C}_8$ alkylamino or $\text{C}_1\text{-C}_8$ alkoxycarbonylamino group;

X^7 is hydrogen;


Y^1 is $-\text{CH}_2-$, $-\text{CHF}-$, $-\text{CF}_2-$, or $-\text{C}(\text{CH}_3)_2-$;

Z is $-\text{O}-$.


5. The compound according to claim 2, which is represented by one of the following structures:




IIb-18



IIb-25



IIb-43



IIb-46.

6. A composition comprising a compound according to one of claims 1-5 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable diluent and/or excipient.

5 7. The use of the compound according to one of claims 1-5 or the composition according to claim 6, which is used as a GPR40 agonist.

8. A method of treating or preventing diabetes or a related metabolic syndrome, characterized by administering to a patient an effective dose of the compound according to
10 one of claims 1-5 or a composition according to claim 6.

9. The use of a compound according to one of claims 1-5 or the composition according to claim 6 in the preparation of a medicament for the treatment or prevention of diabetes or a related metabolic syndrome.

15

10. The use according to claim 9, the diabetes are type II diabetes.