

(II),

where

R^1, R^2, R^3, R^4 are each independently selected from: -H, -(C_1-C_{12})-alkyl;

V and W are each independently selected from:

-H, -(C_1-C_{12})-alkyl, -O-(C_1-C_{12})-alkyl, -(C_1-C_{12})-heteroalkyl, -(C_4-C_{20})-aryl, -(C_4-C_{20})-aryl-(C_1-C_{12})-alkyl, -(C_4-C_{20})-aryl-O-(C_1-C_{12})-alkyl, -(C_1-C_{12})-alkyl-(C_4-C_{20})-aryl, -(C_4-C_{20})-aryl-COO-(C_1-C_{12})-alkyl, -(C_4-C_{20})-aryl-CONH-(C_1-C_{12})-alkyl, , -(C_4-C_{20})-aryl-CON[(C_1-C_{12})-alkyl]$_2$, -(C_4-C_{20})-heteroaryl, -(C_4-C_{20})-heteroaryl-(C_1-C_{12})-alkyl, -(C_3-C_{12})-cycloalkyl, -(C_3-C_{12})-cycloalkyl-(C_1-C_{12})-alkyl, -(C_3-C_{12})-heterocycloalkyl, -(C_3-C_{12})-heterocycloalkyl-(C_1-C_{12})-alkyl, -CO-(C_1-C_{12})-alkyl, -CO-(C_4-C_{20})-aryl, -CO-(C_4-C_{20})-heteroalkyl, -(C_4-C_{20})-aryl-CO-(C_1-C_{12})-alkyl, -(C_4-C_{20})-aryl-CO-(C_4-C_{20})-aryl;

X and Y are each independently selected from O and N;

where, in the case that X is O, n = 1,

and, in the case that Y is O, m = 1,

and, in the case that X is N, n = 2,

and, in the case that Y is N, m = 2;