


PROCESS FOR THE MANUFACTURE OF
HYDROXYALKOXYBENZOPHENONES

The invention relates to a process for the
5 manufacture of hydroxyalkoxybenzophenones of general
formula:



in which

- R represents a hydrogen atom, a hydroxyl radical or an optionally substituted C₁ to C₂₀ alkyl or aralkyl group,
- 15 - R' represents a hydrogen atom or an optionally substituted C₁ to C₂₀, preferably C₁, C₈ or C₁₂, alkyl or aralkyl group.

It is known to prepare the hydroxyalkoxybenzophenones of formula (I) by reaction of an α,α,α -trihalo-
20 toluene with a polyhydroxyphenol or a hydroxyalkoxyphenol in the presence of a specific catalyst.


In this regard,

- patent US-A-3,769,349 proposes to perform the process in an aqueous-acetic medium,
- 25 - patent US-A-3,639,483 proposes to perform the process in an aqueous-alcoholic medium in the presence of alkyl halide,
- patent BE 795,782 proposes to work in two successive steps in different solvents, namely, an aqueous-alcoholic solvent first, followed by a ketonic solvent,
- 30 - patent FR-A-2,473,506 involves a more or less heavy alcohol in a stoichiometric amount,
- patent ZA 7,602,667 priority US 589,967 proposes to work in the presence of dimethylsulphoxide,
- 35 - patents EP-A-154,092 and US-A-4,568,429 propose to use N-methylpyrrolidone as catalyst,

- patent DE 39 04 371 of 16/08/90 proposes to use alcoholic or polyhydroxylated emulsifying agents or surfactants,
- patent HU 37,588 proposes to involve Lewis acids of the type ZnCl_2 , MgCl_2 , AlCl_3 .

The abovementioned known processes have the drawback of using considerable amounts of catalysts and/or of co-solvent; moreover, they are complicated and generate very considerable amounts of non-recyclable effluents which it is necessary to destroy at great expense.

Finally, in all these processes, the benzophenones obtained starting with trichlorotoluene, and more particularly 2,4-dihydroxybenzophenone, are contaminated with considerable amounts of xanthenones which constitute coloured impurities, and especially up to approximately 90% by weight with 6-hydroxy-9-phenylfluorone, also referred to as 6-hydroxy-9-phenyl-3H-xanthen-3-one, of formula:



as well as with other similar xanthenic ring systems.

The undesired xanthenones are formed by the action of a second polyhydroxyphenol molecule on the trihalogenated carbon; one of the aims of the invention is to prevent their formation.

The abovementioned coloured impurities constituted by the xanthenones represent a major drawback, since it is extremely difficult to get rid of them subsequently. Moreover, the main product released by the reaction of the trichloromethylbenzene is hydrochloric acid, and it is not

possible, with any of the catalysts previously used, to work in a concentrated hydrochloric medium without generating considerable amounts of these highly coloured xanthenones.

5 Industrially, this therefore leads to the products formed being distilled under very weak vacuums and at very high temperatures, thereby resulting in expensive industrial constraints.

10 However, the benzophenones of formula (I), and in particular 2,4-dihydroxybenzophenone, have the property of absorbing ultraviolet solar radiation and, in this respect, they are used as anti-UV additives for polymers or in the cosmetics industry.

15 In view of these industrial uses, the user consequently wishes to have available benzophenones of the greatest possible purity and of the least possible coloration.

20 The objective of the invention is thus, above all, to provide a process which not only no longer has the drawbacks of the prior processes but which, in addition, leads to hydroxyalkoxybenzophenones which are virtually free of coloured impurities.

25 As a result of extensive research, the Applicant Company has had the merit of finding that this objective was achieved when, in the processes of the type in question, use is made of a sulphur-containing or selenium-containing catalyst consisting of a mercaptan or of a seleno-mercaptan acting as a nucleophile in a very acidic medium, this being all the more unexpected since
30 sulphur-containing products other than mercaptans do not give satisfactory results.


35 It follows that the process for the preparation of the hydroxyalkoxybenzophenones of formula (I) in accordance with the invention, in which an α,α,α -trihalo-toluene is reacted with a polyhydroxyphenol or a hydroxy-alkoxyphenol, is characterized in that the reaction is

carried out in the presence of a sulphur-containing or selenium-containing catalyst consisting of a compound of the mercaptan or seleno-mercaptan type and preferably containing a short C₁ to C₂₀, preferably C₂ to C₁₂, carbon chain which is largely sterically unhindered.


It is possible, although the Applicant Company does not wish to be bound by this theory, that the catalyst used in accordance with the invention adds itself temporarily to the product formed as an intermediate during the reaction of the α,α,α -trihalotoluene (preferably trichlorotoluene) with the polyhydroxyphenol (for example resorcinol).

By virtue of this process, the action of a second molecule of polyhydroxyphenol would then be slowed down, or even completely prevented, thereby avoiding the final formation of the xanthenones.

The simplified reaction diagram representing this process might be as follows:




The abovementioned non-isolable intermediate would then rearrange, again losing HCl acid, leading to the following probable product (III):



The intermediate (III) would in turn form, with the catalyst used in accordance with the invention, a very

temporary addition compound of the type:



in which S represents the nucleophilic atom of the catalyst group.

5 The sulphur atom is replaced by a selenium atom when the catalyst used is a seleno-mercaptan.

 The temporary addition of the catalyst limits the attack of a second molecule of resorcinol on the compound (III), by virtue of the formation of a compound of
10 type (IV).

 The latter compound leads, on acidic hydrolysis, to the corresponding benzophenone (I), the hydrolysis releasing the catalyst.

 One of the advantages of the process according to
15 the present invention is to reduce the amounts of xanthenones formed secondarily during the synthesis to a very low level, the hydroxyalkoxybenzophenones obtained having a xanthene content which is especially less than 0.5% by weight.

20 These yields of pure isolated product, in this instance 2,4-dihydroxybenzophenone, are excellent, of the order of 95 to 99% relative to the two main constituents, and are thus considerably superior to those described in the prior art.

25 According to an advantageous embodiment of the process in accordance with the invention, the compound of mercaptan type is of the formula



in which R represents a hydrogen atom or an optionally substituted C_1 to C_{30} , preferably C_2 or C_3 , alkyl, cycloalkyl or aralkyl group, n being an integer equal to 1 or 2.

5 According to an advantageous embodiment of the process in accordance with the invention, the catalyst used is represented by the abovementioned formula (V) in which n is greater than 2 and may take a value tending towards infinity, the catalysts in question then being in
10 the form of a resin or a polymer.

According to another advantageous embodiment of the process in accordance with the invention, the catalyst used and consisting of a compound of mercaptan type is an acid-mercaptan of formula



in which

- R represents an optionally substituted C_1 to C_{30} , preferably C_2 or C_3 , alkyl, cycloalkyl or aralkyl group, and
- 20 - R' represents a hydrogen atom, an optionally substituted C_1 to C_{20} alkyl, cycloalkyl or aralkyl group or a metal atom, preferably an alkali metal or alkaline-earth metal atom,
- m and p, which are identical to or different from each
25 other, are integers from 1 to 5.

According to another embodiment of the process in accordance with the invention, the catalyst of formula (V) or (VI) is chosen from the group comprising:

- lauryl mercaptan,
- 30 - n-hexyl mercaptan,
- cyclohexyl mercaptan,
- 1,2-ethanedithiol,
- 1,6-hexanedithiol,
- mercaptoacetic (thioglycolic) acid,
- 35 - 3-mercaptopropionic acid,
- thiolactic (2-mercaptopropionic) acid,

- mercaptosuccinic acid (thiomalic acid),
- thiosalicylic acid (mercaptobenzoic acid),
- mercaptonicotinic acid and
- mercaptopyruvic acid

5 as well as the corresponding esters and salts.

According to another advantageous embodiment of the process in accordance with the invention, the catalyst used and consisting of a compound of mercaptan type is a mercaptosulphonic acid of formula



in which R and R' have the meanings given in relation to the formula (VI).

Among the preferred compounds of formula (VII) are

- mercaptoethanesulphonic acid and
- 15 - 1-mercaptopropanesulphonic acid

as well as the alkali metal salts, especially the sodium salt, thereof.

According to another advantageous embodiment of the process in accordance with the invention, the reaction is carried out in the presence of a sulphur-containing compound, which is introduced before the trihalo derivative is run in, and which is capable of giving a compound of the mercaptan type by prior reaction with the acidic reaction medium, the said reaction possibly being:

- 25 - of the "hydrolysis" type, the sulphur-containing compound then being chosen from the group comprising thioacetals and hemithioacetals, thiolactones such as γ -thiobutyrolactone, dimers, ortho esters and products of self-esterification of the acid-thiols corresponding to
- 30 the formula (VI) and in particular those of thioglycolic acid, trithiocarbonates such as vinylene trithiocarbonate and ethylene trithiocarbonate, dithiocarboxylic acids, dithiophosphoric acids and dithiophosphinic acids,

- 35 - or of the "decomposition" type, the sulphur-containing compound then being chosen from the group comprising thioacids, thioamides such as thiourea,

thioacetamide, thioacetanilide, thiobenzamide, thiouracil, thiosemicarbazide and thiophosgene,

- or of the "reduction" type with the secondary involvement of a reducing agent, the sulphur-containing compound then being chosen from the group comprising organic (cyclo)-di- or polysulphides and metal di- or polysulphides.

According to another embodiment of the process in accordance with the invention, the reaction is carried out in the presence of an amount of catalyst from 0.1 mol% to 100 mol%, preferably from 5 to 10 mol%, relative to the α,α,α -trihalotoluene, the slight gain obtained beyond this limit not justifying the use of greater amounts.

According to another embodiment of the process in accordance with the invention, the reaction is carried out in a solvent comprising an acidic aqueous phase which is concentrated and, preferably, saturated with hydrochloric acid; this is a considerable simplification.

By virtue of the purity of the hydroxyalkoxybenzophenones prepared using the process according to the invention, the distillation treatments necessary in the case of the products obtained by the processes of the prior art are eliminated.

In other terms, the hydroxyalkoxybenzophenones obtained in accordance with the invention are pure enough to be employed as they are in the synthesis of the alkylated derivatives used as UV stabilizers in the polymer and cosmetics industries.

As a result of the presence of the mercaptan-type catalyst, it is possible to perform the process in a concentrated, or even saturated (at the saturation corresponding to the temperature at which the reaction is carried out) hydrochloric phase without generating the xanthenones in question.

The aqueous hydrochloric medium may consequently be recycled, which constitutes a considerable advantage since

the amount of effluents needing to be destroyed is decreased, or even eliminated altogether; it follows therefrom that the process according to the invention is particularly economical.

5 Furthermore, by recycling the hydrochloric aqueous phase and by readjusting the amount of catalyst, hydroxyalkoxybenzophenones of increasing purity, that is to say of increasingly lower optical density, are obtained.

10 Finally, on account of the presence of the catalyst, the amount of aqueous phase used during the process in accordance with the invention may be greatly reduced in comparison with the amounts necessary in the processes of the prior art, which required large volumes
15 of water; the production efficiency of the process according to the present invention is thus much greater than that of the known processes.

 For economic reasons, it is preferred to work as closely as possible to stoichiometric conditions, a very
20 slight excess of trihalide nevertheless being desirable; the presence of secondary products, which would risk accumulating in the course of the recyclings if the differences were large, is thus limited.

 The temperature at which the process is carried out
25 is controlled to be between 0 and 100°C, preferably between 20 and 40°C, which makes it possible to limit the premature hydrolysis of the trihalo derivative and the lowering of yield which would result.

 A subsequent gradual increase of the temperature up
30 to a value of 95 to 100°C makes it possible to complete the reaction without rushing the gradual precipitation of the benzophenone.

 A recrystallization may then be performed if a
35 purer product than that produced by simple draining and washing of the reaction mass is desired. To do this, the crystalline mass is taken up in known solvents for benzo-

phenone, under hot conditions, and washes are performed with water while hot, followed by a crystallization.

Since the main drawback of the mercaptans is their very strong, repulsive odour which is detrimental as far as the process is concerned, it is desirable to increase their molar mass in order to limit these odour problems. In this regard, the mercaptans constituting the preferred catalysts in the sense of the invention are those which contain a hydroxylated or carboxylated polar radical, which limits the vapour pressure of the product and, in the case of the carboxyl and sulfonic group, the odour.

The presence of this polar radical furthermore gives the mercaptan better solubility in the aqueous phase, which is very advantageous.

Conversely, the presence of a radical which is too hydrophilic, such as the NH_3Cl and NR_2HCl group, makes the mercaptan which bears it totally ineffective from the point of view of its catalysing power.

According to an advantageous embodiment of the process in accordance with the invention, 2,4-dihydroxybenzophenone is prepared in a first step and is subjected without isolation, in a second step, to methylation using methyl sulphate, which leads to 2-hydroxy-4-methoxybenzophenone.

According to an advantageous embodiment of the process in accordance with the invention, 2,4-dihydroxybenzophenone is prepared in a first step and is subjected without isolation, in a second step, to octylation using octyl chloride, which leads to 2-hydroxy-4-octyloxybenzophenone.

In order to assess the effectiveness of a given mercaptan as a catalyst, the optical density of a solution of the benzophenone obtained may be measured.

Indeed, the optical density is closely linked to the degree of contamination of the benzophenone by the xanthenones.

This measurement of the optical density is made by application of the Beer-Lambert law, at the benzophenone maximum absorption, with the dilutions and the methods usually used by those skilled in the art. In the case of 2,4-dihydroxybenzophenone, the measurement is made at 450 nm, the dry benzophenone being dissolved to an amount of 10% in acetone, the cell having a thickness of 4 cm.

The invention may be better understood with the aid of the non-limiting examples which follow and which relate

- to advantageous embodiments of the process in accordance with the invention,

- to comparisons between the catalysts of the prior art and those used in accordance with the invention,

- to comparisons between various sulphur-containing compounds showing the superiority of those used in accordance with the invention,

- to the determination of the influence of the amount of catalyst used, and

- to the influence on the optical density of the benzophenone obtained from the recycling of the hydrochloric aqueous phase.

EXAMPLE 1

Preparation of 2,4-dihydroxybenzophenone

110 g of resorcinol, 750 g of water, 7.5 g of 32% HCl and 2.5 g of lauryl mercaptan are mixed together.

200 g (1.02 mol) of α,α,α -trichlorotoluene are run into this mixture over 5 hours, at a temperature of 35 to 40°C.

The temperature is then maintained at 60°C for 2 hours.

The orange-yellow crystalline mass obtained is drained on a sinter funnel at 50°C.

This product is washed three times with 100 ml of water.

It is dried in a ventilated oven at 80°C and 205 g of crude 2,4-dihydroxybenzophenone, equivalent to a 95.8%

yield, are obtained as a product having a melting point of 139-143°C and an optical density of 490 (10% solution in acetone at 450 nm), which corresponds to a total xanthenone content of 0.5%.

5 EXAMPLE 2

Preparation of 2,4-dihydroxybenzophenone

110 g of resorcinol, 600 ml of 32% HCl and 10 g of mercaptoethanol are mixed together.

200 g (1.02 mol) of α,α,α -trichlorotoluene are run
10 in over 3 hours, at a temperature of 35 to 40°C.

The temperature is then maintained at 70-80°C for 1 hour.

The orange-yellow crystalline mass obtained is drained on a sinter funnel and is washed three times with
15 100 ml of water.

The product is dried in a ventilated oven at 80°C and 210 g of crude 2,4-dihydroxybenzophenone, equivalent to a 98% yield, are obtained as a product having a melting point of 139-143°C and an optical density of 220 (10%
20 solution in acetone at 450 nm), which corresponds to a total xanthenone content of 0.35%.

EXAMPLE 3

Preparation of 2,4-dihydroxybenzophenone

110 g of resorcinol, 700 ml of 37% of HCl and 15 g
25 of thioglycolic acid (2-mercaptoacetic acid) are mixed together.

200 g (1.02 mol) of α,α,α -trichlorotoluene are run in over 3 hours, at a temperature of 50°C.

The temperature is then gradually raised to 95°C.

30 The crystalline mass may then be drained or taken up in a good water-immiscible solvent, followed by washes of the organic phase according to the usual practices of those skilled in the art.

After recrystallization, followed by draining and
35 drying in a ventilated oven, 207 g of 2,4-dihydroxybenzophenone, equivalent to a 96.5% yield, are obtained as a

product having a melting point of 139-143°C and an optical density of 500 (10% solution in acetone at 450 nm, 4 cm cell), which corresponds to a total xanthenone content of approximately 0.5%.

5 **EXAMPLE 4**

 The procedure is performed in the same way as for example 3, but the catalyst is made of 10 g of methylthioglycolic ester (formula VI). After work up, 169 g of 2,4-dihydroxy-benzophenone are obtained, equivalent to a 79% yield, having an optical density of 420.

10 **EXAMPLE 5**

 The procedure is performed in the same way as for example 3, but the catalyst is made of 15 g of buthylmercaptopropionic ester (formula VI). After work up, 194 g of 2,4-dihydroxybenzophenone are obtained, equivalent to a 90.7% yield, having an optical density of 120.

15 **EXAMPLE 6**

 This example shows that in order to obtain low coloured benzophenones, it is not absolutely essential to perform the synthesis at high temperature, provided a catalyst is used in accordance with the invention.

 110 g of resorcinol, 600 g of 32% HCl and 12 g of 3-mercepto-1-propanesulfonic acid sodium salt (formula VII) are mixed together. 200 g (1.02 mol) of α,α,α -trichlorotoluene are run into the mixture over 3 hours, at a temperature of 30°C. This same temperature is then maintained for 3 hours more, while stirring. The product is then filtered off, washed with water, dried in a ventilated oven. 205 g of 2,4-dihydroxybenzophenone, equivalent to a 96% yield, are obtained as a product having a melting point of 139-143°C and an optical density of 100 to 140, which correspond to a total xanthenon content of 0.4 to 0.6%.

EXAMPLE 7**Preparation of 2,4-dihydroxybenzophenone and of 2-hydroxy-4-methoxybenzophenone**

110 g of resorcinol, 560 ml of 32% HCl and 10 g of
5 3-mercaptopropionic acid are mixed together.

200 g (1.02 mol) of α,α,α -trichlorotoluene are run
in over 3 hours, at a temperature of 35 to 40°C.

The temperature is then maintained at 70-80°C for
1 hour.

600 g of solvent are run in to dissolve the crys-
talline mass and the temperature in the mass is brought to
95°C.

The lower acidic aqueous phase is separated out
after settling of the phases has taken place, and this
15 acidic phase may be recycled.

The organic phase is washed with 150 ml of water
containing 30 g of sodium hydrogen carbonate, and then
with 100 ml of water.

The organic phase containing dihydroxybenzophenone
20 is dried azeotropically.

If it is desired to obtain 2-hydroxy-4-methoxy-
benzophenone, the methylation may be performed directly by
salification using K_2CO_3 and methyl sulphate without
isolating the intermediate benzophenone.

190 g of 2-hydroxy-4-methoxybenzophenone are
25 obtained, having a melting point of 62.5-63.8°C and an
optical density of 0.8 (20% solution of dry benzophenone;
measurement at 450 nm in a 1 cm cell).

If a higher purity is sought, the dihydroxybenzo-
phenone may be recrystallized beforehand by cooling.

The dark yellow crystalline mass is then drained on
a sinter funnel at 20°C and is washed with 100 g of the
same solvent.

The product is dried in a ventilated oven at 80°C
35 and 205 g of purified 2,4-dihydroxybenzophenone, corres-
ponding to a yield of 95.8%, are obtained as a product

having a melting point of 139-144°C and an optical density of 80-120 (10% solution in acetone at 450 nm), the total xanthenone content consequently being from 0.20 to 0.25%.

5 EXAMPLE 8

This example shows that the way of getting the chemicals reacted is not very important, provided a catalyst is used in accordance with the invention. The procedure is performed with the same materials and same general process as for example 7, but the run of reactants is reversed: a mixture of 110 g of resorcine and 560 g of 32% HCl are fed in two hours on a mixture of 200 g α,α,α -trichlorotoluene and 10 g of mercaptopropionic acid, at a temperature of 50°C. The following procedure is then exactly performed as in example 7. It is also possible to combine this with the synthesis of 2-hydroxy-4-methoxybenzophenone as already mentioned in example 7, or to perform the alkylation with octyl chloride after the salification is being obtained with K_2CO_3 in a similar manner, without the intermediate benzophenone to be isolated. If not, 192 g of 2,4-dihydroxybenzophenone are obtained after work up, having an optical density of 50, which corresponds to a total xanthenon content of less than 0.2%.

EXAMPLE 9

25 Preparation of 2,4-dihydroxybenzophenone and of 2-hydroxy-4-methoxybenzophenone

The lower aqueous phase isolated in Example 7 (or acidic mother liquors) is taken up and 110 g of resorcinol are added, along with 5 g of 3-mercaptopropionic acid, then the α,α,α -trichlorotoluene is run in, the procedure being performed exactly as in Example 7.

208 g of 2,4-dihydroxybenzophenone, corresponding to a yield of 97%, are obtained as a product having an optical density of 50 to 60 (10% solution in acetone at 450 nm), the total xanthenone content consequently being from 0.15 to 0.20%.

It is also possible, in the same manner as in

Example 7, to combine this with the synthesis of 2-hydroxy-4-methoxybenzophenone by taking the crystalline mass up beforehand in a solvent, followed by performing washings, salification and then the methylation with methyl sulphate. 188 g of 2-hydroxy-4-methoxybenzophenone are then obtained, having a melting point of 62.5 to 63.8°C and an optical density of 0.5 to 1.

EXAMPLE 10

Preparation of 2-hydroxy-4-methoxybenzophenone

10 124 g of 3-methoxyphenol, 600 ml of 32% HCl and 10 g of 3-mercaptopropionic acid are mixed together.

200 g (1.02 mol) of α,α,α -trichlorotoluene are run in over 3 hours, at a temperature between 35 and 40°C.

15 The temperature is then maintained at 70-80°C for 1 hour.

The lower acidic aqueous phase is separated out after settling of the phases has taken place.

20 The organic phase is washed with 150 ml of water containing 30 g of sodium hydrogen carbonate, and then with 100 ml of water.

200 ml of a low-molecular-weight alcohol are run in and the mixture is cooled in order to crystallize the 2-hydroxy-4-methoxybenzophenone.

25 The product is filtered off and is then washed with a minimum of this same alcohol.

30 It is dried in a ventilated oven at 50°C and 114 g of purified 2-hydroxy-4-methoxybenzophenone, corresponding to a yield of 50%, are obtained as a product having an optical density of 2 (20% solution in acetone, measurement at 450 nm) and a melting point of 62.5-63.8°C.

EXAMPLE 11

Preparation of 2-hydroxy-4-octyloxybenzophenone

222 g of 3-octoxyphenol, 600 ml of 32% HCl and 10 g of 3-mercaptopropionic acid are mixed together.

35 200 g (1.02 mol) of α,α,α -trichlorotoluene are run in over 3 hours, at a temperature of 35 to 40°C.

The temperature is then maintained at 70-80°C for 1 hour.

The lower acidic aqueous phase is separated out after settling of the phases has taken place.

5 The organic phase is washed with 150 ml of water containing 30 g of sodium hydrogen carbonate, and then with 100 ml of water.

10 200 ml of lower alcohol are run in and the mixture is cooled to crystallize the 2-hydroxy-4-methoxybenzophenone.

The product is filtered off and washed with a minimum of the same lower alcohol.

15 It is dried in a ventilated oven at 40°C and 146 g of 2-hydroxy-4-octyloxybenzophenone, corresponding to a yield of 45%, are obtained as a product having a melting point of 47-59°C and an optical density of 0.25.

EXAMPLE 12

20 This example shows the superiority of the catalysts used in accordance with the invention relative to nucleophilic compounds which may theoretically serve as catalysts and relative to catalysts used according to the prior art.

25 To do this, 20 samples of 2,4-dihydroxybenzophenone were prepared using a proportion of 2 to 10 mol% of the following catalysts:

TABLE I

Sample No.	Nature of the catalyst used	Optical density
1	Methanesulphonic acid	- 10 800
2	1,4-thioxane	- 10 000
3	NONE	- 7 000
4	carbon disulfide	- 7 200
5	Phenol	- 6 500
6	dimethylamino pyridine hydrochloride	- 6 500
7	Triflic acid	- 6 000
8	2-Mercaptoethylamine hydrochloride	- 5 600
9	Methyldialkyl(C ₁₈)amine	- 5 000
10	Sodium dioctyl- sulphosuccinate	- 4 800
11	dimethylsulfoxide	- 4 000
12	Polyvinyl alcohol	- 4 000
13	γ -Mercaptotrimethoxysilane	- 3 000
14	n-Hexanol	1 700
15	Lauryl mercaptan	1 500
16	Thioglycolic acid	500
17	2-Mercaptoethanol	220
18	3-Mercaptopropionic acid	150
19	3-mercapto-1-propanesulfonic acid, sodium salt	100-140
20	butyl-3-mercaptopropionate	100-120

The results collected in Table I show that the samples prepared in accordance with the invention exhibit a very significant enhancement in the optical density and thus in the quality of the product.

EXAMPLE 13

In this example, the influence of the amount of catalyst used is studied.

Six samples of 2,4-dihydroxybenzophenone were prepared using increasing amounts of catalyst consisting of 3-mercaptopropionic acid, namely:

<u>Sample No.</u>	<u>Amount of 3-mercaptopropionic acid per mole of trichlorotoluene</u>
1	2.5 g
2	5 g
3	10 g
4	20 g
5	30 g
6	200 g

Measurement of the optical density for the six samples under the conditions indicated above gives the following results:

<u>Sample No.</u>	<u>Optical density</u>
1	2 500
2	300
3	100
4	80
5	80
6	55

It follows therefrom that, from an economic point of view, there is no point in exceeding 10 g of catalyst per mole of α,α,α -trihalotoluene.

EXAMPLE 14

The influence of recycling the hydrochloric aqueous phase was studied.

To do this, five samples of 2,4-dihydroxybenzophenone were prepared successively by working as indicated in Example 4 for the first sample and by

- recycling the hydrochloric aqueous phase for the preparation of each of the samples 2 to 5, while at the same time

- readjusting the amount of catalyst.


Measurement of the optical density on each of the five samples gives the following results:

	<u>Sample No.</u>	<u>Optical density</u>
5	1	120
	2	70
	3	60
	4	55
	5	40

10 The advantage derived by recycling the hydrochloric aqueous phase is clearly apparent.

CLAIMS

1. Process for the manufacture of hydroxyalkoxybenzophenones of general formula:



in which

- 10
- R represents a hydrogen atom, a hydroxyl radical or an optionally substituted C_1 to C_{20} alkyl or aralkyl group,
 - R' represents a hydrogen atom or an optionally substituted C_1 to C_{20} , preferably C_1 , C_8 or C_{12} , alkyl or aralkyl group,


15 by reaction of an α,α,α -trihalotoluene with a polyhydroxyphenol or a hydroxyalkoxyphenol, characterized in that the reaction is carried out in the presence of a sulphur-containing or selenium-containing catalyst consisting of a compound of the mercaptan or seleno- mercaptan type and preferably containing a short C_1 to C_{20} , preferably C_2 to C_{12} , carbon chain which is largely sterically unhindered.

2. Process according to Claim 1, characterized in that the reaction is carried out in the presence of a compound of mercaptan type of formula



in which R represents a hydrogen atom or an optionally substituted C_1 to C_{30} , preferably C_2 or C_3 , alkyl, cycloalkyl or aralkyl group, n being an integer equal to 1 or 2.

30 3. Process according to Claim 1, characterized in that the reaction is carried out in the presence of a compound of acid-mercaptan type of formula



in which

- 35
- R represents an optionally substituted C_1 to C_{30} , preferably C_2 or C_3 , alkyl, cycloalkyl or aralkyl group, and



- R' represents a hydrogen atom, an optionally substituted C₁ to C₂₀ alkyl, cycloalkyl or aralkyl group or a metal atom, preferably an alkali metal or alkaline-earth metal atom,

5 - m and p, which are identical to or different from each other, are integers from 1 to 5.

4. Process according to one of Claims 1 to 3, characterized in that the reaction is carried out in the presence of a catalyst chosen from the group comprising:

- 10 - lauryl mercaptan,
- n-hexyl mercaptan,
- cyclohexyl mercaptan,
- 1,2-ethanedithiol,
- 1,6-hexanedithiol,
15 - mercaptoacetic (thioglycolic) acid,
- 3-mercaptopropionic acid,
- thiolactic (2-mercaptopropionic) acid,
- mercaptosuccinic acid (thiomalic acid),
- thiosalicylic acid (mercaptobenzoic acid),
20 - mercaptonicotinic acid and
- mercaptopyruvic acid

as well as their esters and salts.

5. Process according to Claim 1, characterized in that the reaction is carried out in the presence of a
25 mercaptosulphonic acid of formula



in which R and R' represent a hydrogen atom, an optionally substituted C₁ to C₂₀ alkyl, cycloalkyl or aralkyl group or a metal atom, preferably an alkali metal or alkaline-
30 earth metal atom.

6. Process according to Claim 1, characterized in that the reaction is carried out in the presence of a sulphur-containing compound, which is introduced before the trihalo derivative is run in, and which is capable of
35 giving a compound of the mercaptan type by prior reaction with the acidic reaction medium, the said reaction



possibly being:

5 - of the "hydrolysis" type, the sulphur-containing compound then being chosen from the group comprising thioacetals and hemithioacetals, thiolactones such as γ -thiobutyrolactone, dimers, ortho esters and products of self-esterification of the acid-thiols corresponding to the formula (VI) and in particular those of thioglycolic acid, trithiocarbonates such as vinylene trithiocarbonate and ethylene trithiocarbonate, dithiocarboxylic acids, 10 dithiophosphoric acids and dithiophosphinic acids,

15 - or of the "decomposition" type, the sulphur-containing compound then being chosen from the group comprising thioacids, thioamides such as thiourea, thioacetamide, thioacetanilide, thiobenzamide, thiouracil, thiosemicarbazide and thiophosgene,

20 - or of the "reduction" type with the secondary involvement of a reducing agent, the sulphur-containing compound then being chosen from the group comprising organic (cyclo)-di- or polysulphides and metal di- or polysulphides.

25 7. Process according to one of Claims 1 to 6, characterized in that the reaction is carried out in the presence of an amount of catalyst from 0.1 mol% to 100 mol%, preferably from 5 to 10 mol%, relative to the α,α,α -trihalotoluene.

30 8. Process according to one of Claims 1 to 7, characterized in that the reaction is carried out in a solvent consisting of a concentrated acidic aqueous phase, which is preferably saturated with hydrochloric acid.

 9. Process according to Claim 8, characterized in that the acidic aqueous solvent is recycled after recovery of the hydroxyalkoxybenzophenone formed.