Modified Release 1-[(3-Hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile Formulation

Background of the Invention

The present invention relates to a sustained release preparation containing a dipeptidyl peptidase IV inhibitor (DPP-IV) useful for preventing or treating diabetes mellitus, non-insulin-dependent diabetes mellitus, obesity, arthritis, osteoporosis, and other diseases.

DPP-IV is a useful drug for treating diabetes mellitus, non-insulin-dependent diabetes mellitus, obesity, arthritis, osteoporosis, and other diseases because it inhibits inactivation of glucagons-like peptide-I (GLP-I) in blood plasma and increases its incretion action (WO 02/062,764, WO 01/55105, WO 02/02560).

However, as described in detail below, there were some cases in which strong inhibition of DPP-IV activity in a living body was not always preferable.

For example, it has been reported that the DPP-IV inhibitor enhances vasodialation action by substance O and that the DPP-IV activity was elevated when treating chronic rhinosinusitis because there was an inverse relationship between DPP-IV activity of nasal mucosa of a patient with chronic rhinosinusitis and density of inflammatory cells on the nasal mucosa (for example, *FASEB* J., 2002, 16:1132-1134). Accordingly, it is considered undesirable to strongly inhibit DPP-IV activity in a patient with diabetes complicating chronic inflammation because it results in aggravation of inflammation.

It has been reported that selective DPP-IV activity is important for glucose homeostasis since GLP-I (9-36) amide, a metabolite from GLP-I with DPP-IV, reduces blood glucose level (*Am. J. Physiol. Endocrinol. Metab.*, 2002, 282:E873-E879).

Further it has been reported that in treatment of a patient with type C Hepatitis using interferon-α, severity of depression and anxiety correlated with decrease of serum DPP-IV activity (for example, *Mol. Psychiatry*, 2001, 6:475-480).

In view of the current circumstances that there are some cases in which strong inhibition of *in vivo* DPP-IV activity is not always preferable for the living body, the present inventors considered it necessary to develop a preparation that could appropriately inhibit DPP-IV activity and also afford excellent convenience or compliance.

The inventors herein disclose a modified/sustained release 1-[(3-Hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile (also known as LAF237 or vildagliptin (INN)) formulation for these purposes. The development of a modified/sustained release (MR) formulation comprising vildagliptin is necessary to improve the treatment quality of the treated patients. vildagliptin is an orally active, dipeptidylpepidase IV inhibitor indicated for improvement of glycemic control in type II diabetes.

LAF237 is specifically disclosed in Example 1 of WO 00/34241. LAF237 can be formulated as described in WO 00/34241, or in the International Patent Application No. EP2005/000400 (application number).

Since vildagliptin is sensitive to moisture, one of the main emphases during the Market Formulation (MF) development was product stability. Hydrolysis is the main degradation pathway for vildagliptin. The stability of vildagliptin is known to be affected by initial moisture content of the tablet, excipients with high intrinsic moisture and the amount of excipients present in the tablet (% drug load). The current validated vildagliptin formulation is a DC (direct compression) tablet with a 25% drug load.

Considering the water-sensitivity of vildagliptin, Clinical service form (CSF) and Market Formulation (MF) development was

planned to focus on techniques that would not require the use of water, such as direct compression.

vildagliptin is classified as Category II, therefore, no special handling practices are required.

The vildagliptin drug substance (DS) is sensitive to moisture. Very tight packaging is necessary. Current retest date is 30 months with storage at 25°C.

It is well known that patient adherence to a drug regimen indirectly correlates with frequency of dosing i.e. greater adherence is seen with a once daily (OD) dosing regimen compared to a twice daily (BID) dosing regimen. The majority of data currently available for vildagliptin is derived from studies based on the immediate release (IR) formulation of vildagliptin given BID.


Replacing the 50mg IR BID dose with a OD modified release dose will provide the dual benefit of dosing convenience to the patient and a inovative OD product for treating patients.

The present invention provides a pharmaceutical tablet formulation, capable of being compressed, preferably directly compressed, into a tablet having adequate hardness/friability, an acceptable disintegration time, low sensitivity to moisture, improved stability, an acceptable dissolution pattern and improved pharmacokinetic profile in treated patients. The pharmaceutical tablet formulation, can also be used in e.g. a capsule, tablet, compressed table, direct compressed tablets, granule.

Summary of the Invention

The subject invention provides a pharmaceutical tablet formulation comprising per unit dosage form e.g. per tablet the following ingredients:

(a) a compound as an active ingredient, wherein the compound has a formula:



wherein R is substituted adamantyl and n is an integer from 0 to 3; or a pharmaceutically acceptable salt thereof;

(b) a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) a microcrystalline cellulose; and

(d) a magnesium stearate

The subject invention also provides a pharmaceutical tablet formulation, comprising per 400 mg tablet the following ingredients:

(a) vildagliptin or a pharmaceutically acceptable salt thereof in an amount of about 100 mg;

(b) a hydroxypropyl methylcellulose in an amount of about 160 mg, wherein the hydroxypropyl methylcellulose has an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;


(c) a microcrystalline cellulose in an amount of 120 mg;

(d) a lactose in an amount of about 16 mg; and

(e) a magnesium stearate in an amount of 4 mg.

The subject invention also provides a pharmaceutical tablet formulation comprising per unit dosage form e.g. per tablet the following ingredients:

(a) a compound as an active ingredient, wherein the compound has a formula:



wherein R is substituted adamantyl and n is an integer from 0 to 3; or a pharmaceutically acceptable salt thereof; and

(b) a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution.

Brief Description of the Figures

Figure 1. Process Flow. This figure shows the selected manufacturing process.

Figure 2. Dissolution Profile for Fast and Slow Release.

Figure 3. vildagliptin MR 100mg tablet compression profile for CSF formulation containing 40% HPMC.

Figure 4. vildagliptin MR 100 mg tablet friability for CSF Formulation containing 40% HPMC.

Figure 5. Effect of roller compaction and dwell time, 30% HPMC, 100mg tablet. 30% HPMC was evaluated to improve the compression profile. The effect of the tablet dwell time was simultaneously evaluated at speeds from 40 to 80rpm.

Figure 6. Effect of roller compaction on 100mg tablet hardness. A Fitzpatrick (Chilsonator, model IR220) was evaluated with the interim powder blend without the addition of magnesium stearate, at increasing compaction forces.

Figure 7. Effect of roller compaction on 150mg tablet hardness. A Fitzpatrick (Chilsonator, model IR220) was evaluated with the interim powder blend without the addition of magnesium stearate, at increasing compaction forces.

Figure 8. Effect of roller compaction on the 100mg MR dissolution profile. The effect of roller compaction on the dissolution profile was studied with 40% HPMC K100M and compared to the dissolution profile of the clinical batch.

Figure 9. Effect of roller compaction on the 150mg MR dissolution profile. The effect of roller compaction on the dissolution profile was studied with 40% HPMC K100M and compared to the dissolution profile of the clinical batch.

Figure 10. Effect of roller compaction on vildagliptin 100mg hardness using a Bepex roller compactor. The effect of increasing compaction forces was studied.

Figure 11. Effect of roller compaction on vildagliptin 150 mg tablets using Bepex roller compactor. The effect of increasing compaction forces was studied.

Figure 12. Effect of roller compaction on 100mg tablet. The effect of Fitzpatrick (Chilsonator, model IR220) roller compactor at compaction forces from 500 to 10,000 lb/in on a powder blend containing 40% HPMC.

Figure 13. Effect of roller compaction on 150mg tablets. The effect of Fitzpatrick (Chilsonator, model IR220) roller compactor at compaction forces from 500 to 10,000 lb/in on a powder blend containing 40% HPMC.

Figure 14. Effect of roller compaction on 100mg tablet friability. Roller compaction forces greater than 5,000 lb/in (or 43.75 KN) produced tablets with a compression profile worse than the CSF without roller compaction or pre-compression.

Figure 15. Effect of roller compaction on HPMC concentration and dwell time with 100mg tablet.

Figure 16. Effect of roller compaction on HPMC concentration and dwell time with 150mg tablet.

Figure 17. Dissolution profile for 100mg tablet.

Figure 18. Dissolution profile for 150mg tablet.

Figure 19. Effect of roller compaction on sieve analysis.

Figure 20. Effect of roller compaction on sieve analysis.


Detailed Description of the Invention

Terms

"DPP-IV inhibitor" means a compound that inhibits enzymatic activity of DPP-IV [Classification by IUBMB; EC3.4.14.5]. The compound may be either peptidyl or non-peptidyl. As long as the DPP-IV inhibitor retains its inhibitory activity, the form may be different before and after administration into the living body. Namely, the DPP-IV inhibitor may be "active metabolites" which have DPP-IV inhibitory activity after undergoing structural changes through metabolism in the living body. Further, the DPP-IV inhibitor may be a "prodrug" that is changed into an activated form by reactions with enzymes, stomach acids, etc. under *in vivo* physiological conditions. In addition, the DPP-IV inhibitory activity can be confirmed by utilizing, for example, the method by Raymond et al. (*Diabetics*, 1998, 47:1253-1258).

Specific examples of the DPP-IV inhibitor include

(1) A compound of the formula:



(I)

(wherein ring A is a 5- to 10-membered aromatic ring which may be substituted; R^1 and R^2 are each a hydrocarbon group which may be the same or different and may be substituted, or a heterocyclic group which may be substituted; X is a bond, -O-, -S-, -SO-, $-SO_2-$, or $-NR^3-$ (R^3 is a hydrogen atom or a hydrocarbon

group which may be substituted); and L is a divalent hydrocarbon group) described in WO 02/062764.

The compound represented by formula (I) may be an anhydrous or hydrous substance, as well as a prodrug.

Suitable examples of the compound represented by formula (I) include the following compounds.

Compound I-a

A compound wherein the ring A is a benzene ring which may have 1 or 2 substitutions selected from:

1) a cyano group;

2) a C_{1-10} alkyl group (preferably, ethyl) or a C_{2-10} alkenyl group (preferably, ethenyl) which may be substituted with a carbamoyl group or a carboxyl group, respectively;

3) a hydroxyl group whch may be substituted (preferably, a 1-10 C alkoxy group (preferably, methoxy, isopropoxy) which may have 1 to 3 substituents selected from a carbomoyl group, a carboxyl group and a 2-5 C alkoxycarbonyl group (preferably, methoxycarbonyl); a hydroxyl group; a 7-13 C aralkyloxy group (preferably, benzyloxy)) (more preferably carbamoylmethoxy);

4) an acyl group (preferably, a C_{1-6} alkyl-carbonyl (preferably acetyl), carbamoyl, mono- or di-(C_{1-6} alkyl which may have 1 to 3 subsitutents selected from a halogen atom and a C_{1-6} alkoxy-carbonyl)-carbamoyl (preferably, methycarbamoyl, ethycarbamoyl, propylcarbamoyl, dimethylcarbamoyl, trifluoroethylcarbamoyl, ethoxycarbonylmethylcarbamoyl, etc.) C_{3-10} cycloalkyl-carbamoyl (preferably, cyclopropylcarbamoyl), C_{7-13} aralkyl-carbamoyl (preferably, benzylcarbamoyl), a nitrogen-containing heterocycle-carbonyl which may be substituted with hydroxyl (preferably, pyrrolidinylcarbonyl, piperidinocarbonyl), C_{1-6} alkylsulfonyl (preferably, methylsulfonyl), a C_{1-6} alkylsulfinyl (preferably, methylsulfinyl), carboxyl, C_{1-6} alkoxy-carbonyl (preferably, methoxycarbonyl), and thiocarbamoyl);

5) an amino group which may be substituted (preferably,

carbamoylamnino);

6) a thiol group (preferably, a 1-10 C alkylthio group which may be substituted with a carbamoyl group (preferably methylthio);

7) a heterocyclic group which may be substituted (an aromatic theterocyclic group (preferably, furyl, thienyl, oxazolyl, oxadiazolyl, thiazolyl, tetrasolyl, pyridyl, pyrrolyl, triazolyl) or a non-aromatic heterocyclic group (preferably, dioxoisoindole, 5-oxooxadiazole-3-yl, 5-oxothiaxialole 3-yl), respectively, which may have 1 or 2 substituents preferably selected from a C_{1-6} alkyl group which may be substituted with 1 to 3 halogen atoms (preferably, methyl, trifluoromethyl), a carboxyl group, a 2-8 calkoxy carbonyl (preferably, ethoxycarbonyl), a cyano group, a carboamoyl group, anamino group, a mono- or di-C_{2-10} alkanoylamnio group (e.g. acetylamnio isopentanoylamnio), C_{1-10} alkoxy-carbonylamnio group (e.g. methoxycarbonylamnio), a carbamoylamino group, a mono- or di-C_{1-10} alkyl-carbamoylamnio group (e.g., methylcarbmoylamino, dimethylcaramoylamino), a C_{6-14} aryl-carbonylaminogroup (e.g., benzoylamnio), a C_{3-10} cycloalkyl-carbonylamnio group, a C_{7-13} aralkyloxy-carbonylamnio, dimethylsulfonylamnio), a mono- or di-C_{1-10} alkyl sulfonyl amino group (e.g., methyl sulfonyl amino, diemthyl sulfonylamino), a C_{6-14} arylsulfonylamnio group and a C_{1-6} alkoxy-carbanoylamnino group (e.g., methoxycarbamoylamnio);

8) a benzene ring, which may have 1 or 2 substituents slected from amidino groups; R^1 is a 4-10 C alkyl group (preferably, isobutyl, neopentyl) or a 4-10 cycloalkylalkyl group (preferably, cyclopropylmethyl); R^2 is a 6-14 C aryl group (preferably phenyl) which may have 1 or 2 substituents selected from a halogen atom (preferably, fluorine, chlorine) and C_{1-6} alkyl (preferably methyl); X is a bond; and L is C_{1-10}alkylene (preferably, $-CH_2-$).

Compound I-b

A compound wherein the ring A is a benzene ring which may have 1 or 2 substituents selected from:

1) a C_{1-10} alkyl group (preferably, ethyl) or a C_{2-10} alkenyl group (preferably, ehthenyl) which may be substituted with a 2-8 C alkoxycarbonyl group (preferably, ethoxycarbonyl) or a carbamoyl group, respectively;

2) a hydroxyl group which may be substituted (preferably, a 1-10 C alkoxy group (preferably, methoxy) which may be substituted with a carbamoyl group; more preferably, carbamoylmethoxy);


3) an acyl group (preferably, carbamoyl, thiocarbamoyl, carboxyl);

4) a heterocyclic group which may be substituted (an aromatic heterocyclic group (preferably, furyl, rhienyl, oxazoyl, oxadiazolyl, thiazolyl, tetrasolyl, pyridyl, pyrrolyl, triazoyl) or a non-aromatic heterocyclic group (preferably, 5-oxooxadiazole-3-yl), which may have 1 or 2 substituents preferably selected from a C_{1-6} alkyl group (preferably, methyl), a carboxyl group, a 2-8 C alkoxycarbonyl group (preferably, ethoxycarbonyl), a cyano group, a carbamoyl group, an amnio group, a mono- or di-C_{2-10} alkanoylamnino group (e.g., acetylamnio, isopentanoylamino), C_{1-10} alkoxy-carbonylamnio group (e.g. methoxycarbonylamnio), a carbamoylamnio group, a mono- or di-C_{1-10} alkyl-carbamoylamnio group (e.g., methylcarbamoylamnio, dimethylcarbamoylamino group), a C_{6-14} aryl-carbonylamnio group (e.g., benzoylamino), a C_{3-10} cycloalkyl-carbonylamino group (e.g., methylsulfonylamino, dimethylsulfonylamino), C_{6-14} arylsulfonylamnio group and a C_{1-6} alkoxy-carbamoylamino group (e.g., mehtoxycarbamoylamino)); R^1 is a 4-10 C alkyl group (preferably, isobutyl, neopentyl) or a 4-10 C cycloalkylalkyl group (preferably, cyclopropylmethyl); R^2 is a 1-10 C alkyl group (preferably, butyl) which amy be substituted with 1 to 3 halogen atoms; X is -O-; and L is C_{1-10} alkylene (preferably, -CH_3-).

Among the compounds represented by formula (I), preferably are, especially, 1-[2-[(5-cyanopyridin-2-yl)amino]ethylamino]acetyl-

2-cyano(S)-pyrrolidone; 1-[(3-hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile; 2-[3-(aminomethyl)-4-butoxy-2-isobuthyl-1-oxo-1,2-dihydro-6-isoquinolyl[-1,3-thiazole-4-carbonitrile; 2-[3-(aminomethyl)-4-butoxy-2-isobuthyl-1-oxo-1,2-dihydro-6-isoquinolyl]-1,3-thiazole-4-carboxylic acid; 2-[3-(aminomethyl)-4-butoxy-2-isobuthyl-1-oxo-1,2-dihydro-6-isoquinolyl]-1,3-thiazole-4-carboxamide; ethyl 2-[3-(aminomethyl)-4-butoxy-2-isobuthyl-1-oxo-1,2-dihydro-6-isoquinolyl]-1,3-thiazole-4-carboxylate; (E) 3-[3-(aminomethyl)-4-butoxy-2-isobuthyl-1-oxo-1,2-dihydro-6-isoquinolyl]-2-propenamide; (E)-3-[3-(aminomethyl)-2-isobuthyl-4-phenyl-1-oxo-1,2-dihydro-6-isoquinolyl]-2-propenamide; 3-(aminomethyl)-2-isobuthyl-1-oxo-4-phenyl-1,2-dihydro-6-isoquinolinecarboxamide; 2-{[3-(aminomethyl)-2-isobuthyl-4-phenyl-1-oxo-1,2-dihydro-6-isoquinolyl]oxy}acctamide.

(2) A compound represented by a formula



(II)

(wherein f is 1 or 2; g is 0, 1 or 2; X is $-CH_2-$, $-O-$, $-S-$, $-SO-$, $-SO_2-$ or $-NR_3-$(R^3 is a hydrogen atom or a C_{1-6} alkyl group); R is a hydrogen atom, a cyano group, $-CHO$, $-B(OH)_2$, $P(O)(O-R^3)$, $CC-R^4$, or $CH=N-R^5$(R^4 is a hydrogen atom, a fluorine atom, C_{1-6} alkyl group, cyano group, nitro group, $-OR^3$, $-CO_2R^3$, or $-COR^3$ (R^3 indicates the same meaning as aforementioned); R^5 is a phenyl group, hydroxyl group, $-OR^3$, $-OCOR^3$, or benzyloxy group (R^3 indicates the same meaning as aforementioned); and A indicates an amino acid residue which may be substituted) described in WO 95/15309 etc.

In the formula, the C_{1-6} alkyl group indicated by the R^3 includes, for example, methyl, ethyl, propyl, isopropyl, butyl,

isobutyl, sec-butyl, t-butyl, pentyl, isopentyl, neopentyl, 1-ethylpropyl, hexyl, isohexyl, 1,1-dimethyl, 2,2-dimentylbutyl, 3,3-dimethylbutyl, 2-ethylbutyl and the like.

The amino acid residue in "amino acid residue which may be substituted" indicated by A includes a group in which OH of a carboxyl group constituting α- or β-amino acids has been removed from these amino acids.

Herein, α-amino acids include, for example, alanine, arginine, asparagine, aspartic acid, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lycine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, citrulline, ornithine, homocystein and the like.

β-Amino acids include, for example, β-alanine, β-aminocyclopropanoic acid, β-aminocyclobutanoic acid, β-aminocyclopentanoic acid, β-aminocyclohexanoic acid, β-aminocycloheptanoic acid, β-aminocyclooctanoic acid. The β-amino acids may have unsaturated bonds in a carbon chain constituting the amino acids.

The α- and β-amino acids as described above may be in any of D-, L-, and DL-forms, preferably in the natural type of L form.

The said amino acid residue may have 1 or 2 substituents on an amino group constituting the amino acid or on the side chain of the amino acid.

The said "substituted on the amino group" includes preferably a hydrocarbon group which may be substituted and a piperidinyl group which may be substituted.

A hydrocarbon group in the "hydrocarbon group which may be substituted" includes, for example, a C_{1-6} alkyl group, a C_{3-12} cycloalkyl group, a C_{2-6} alkenyl group, a C_{3-12} cycloalkenyl group, a C_{2-6} alkenyl group, a C_{4-12} cycloalkadienyl group, a C_{6-14} aryl group (preferably, phenyl group), a C_{7-15} aralkyl group (preferably, benzyl group, phenethyl group), an adamantly group,

a bicycle [2,2,1]heptyl group, a bicyclo[3,1,1]heptyl group and the like.

The hydrocarbon group may have 1 to 3 substituents at substitutable positions, and such substitutions include, for example, a halogen atom (preferably, fluorine, chlorine); a cyano group; a hydroxyl group which may be substituted with an acyl group; a hydroxymethyl group; a C_{1-6} alkoxy group which may be substituted with 1 to 3 halogen atoms (preferably fluorine); and an amino group which may be mono- or di-substituted with a C_{6-14} aryl group which may be substituted or a heterocyclic group which may be substituted.

Herein, an acyl group on the "hydroxyl group which may be substituted with an acyl group" includes, for example, the one exemplified as a substitutent on ring A in the said compound I-a.

A C_{6-14} aryl group in the "C_{6-14} aryl group which may be substituted" includes, for example, a phenyl group or a naphthyl group.

A heterocyclic group in the "heterocyclic group which may be substituted" includes, for example, a pyridiyl group, a pyrimidyl group, a pyrazyl group, a quinolyl group, an isoquinolyl group, a quinoxalyl group and the like.

The C_{6-14} aryl group and heterocyclic group may have 1 to 3 substituents at substitutable positions and such substituents include, for example, a halogen atom (preferably, fluorine, chlorine, bromine); a cyano group; a nitro group; a C_{1-6} alkyl group; a C_{1-6} alkoxy group which may be substituted with 1 to 3 halogen atoms (preferably fluorine); a carboxyl group; a carbamoyl group; a C_{1-6} alkylsulfonyl group (preferably, methanesulfonyl group); an aminosulfonyl group which may be mono- or di-substituted with C_{1-6} alkyl group (preferably diemethylaminosulfonyl group).

Especially preferable for a substiutent in the said "hydrocarbon group which may be substituted" is a 5-nitro-2-pyridylamino

group, 5-cyano-2-pyridylamino group, 2-pyrimidylamino group, 2-pyrazylamino group and the like.

A substutent in the said "piperidinyl group which may be substituted" includes, for example, a C_{1-6} alkyl group; a hydroxymethyl group, "a C_{6-14} aryl group which may be substituted" and "a heterocyclic group which may be substituted" exemplified in the said "amino group which may be mono- or di-substituted with C_{6-14} aryl group which may be substituted or the heterocyclic group which may be substituted." The number of substituents is, for example, 1 to 3.

The said "substituent on the side chain of an amino acid" includes, for example, a hydrocarbon group which may be substituted, a hydroxyl group, a C_{1-10} alkoxy group which may be substituted with 1 to 3 halogen atoms (preferably fluorine), an acyl group, and amino group which may be substituted and the like.

Herein, a hydrocarbon in the said "hydrocarbon group which may be substituted" includes, for example, a C_{1-10} alkyl group, a C_{3-12} cycloalkyl group, a C_{2-10} alkenyl group, a C_{3-12} cycloalkenyl group and the like.


The hydrocarbon group may have 1 to 3 substituents at substitutable positions, and such substituents include, for example, an amino group, a C_{1-6} alkyl-carbonylamino group (preferably, acetylamino group), a hydroxyl group, a C_{1-6} alkoxy group, a heterocyclic group (preferably, pyridyl group) and the like.

The said "acyl group" is preferably a nitrogen-containing heterocyclic-carbonyl group which may be substituted. The "nitrogen-containing heterocyclic which may be substituted" includes nitrogen-containing heterocycles ((preferably, pyridine, pyridazine, pyrimidine, pyrazine, imidazole, pyrazole, thiazole, isothiazole, oxazole, isooxazole, etc.) which may have 1 to 3 substituents selected from, for example, a halogen atom (preferably, fluorine, chlorine, bromine), a cyano group, a


nitro goup, a C_{1-6} alkyl group (e.g., trifluoromethyl group) which may be substituted with 1 to 3 halogen atoms (preferably fluorine), a C_{1-6} alkoxy group, an amino group which may be mono- or di-substituted with C_{1-6} alkyl group, a hydroxyl group, a carboxy group and a C_{1-6} alkyl-oxycarbonyl group.

A substituent in the said "amino group which may be substituted" includes a C_{1-6} alkyl group, etc. which may have 1 tom 3 substituents selected from, for example, a carboxyl group, a carbamoyl group, a C_{1-6} alkyl-oxycarbonyl group and a nitrogen-containing heterocyclic group (preferably, pyridyl). These substituents may be coupled with a hydroxyl group, a carboxyl group, an amino group and the like which is on the side chain of an amino acid.

A suitable example of the compound represented by formula (II) includes N-(N'-substituted glycyl)-2-cyano-pyrrolidine derivatives such as (2S)-1-{{{2-[(5-cyanopyridine-2-yl)amino]ethyl}amino}acetyl}-2-cyano-pyrrolidine (DPP-728) (described in WO 98/19998) represented by the formula:




(2S)-1-{[(3-hydroxy-1-adamantyl)amino]acetyl}-2-cyano-pyrrolidine (1-[(3-Hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile) (also known as LAF237 or vildagliptin)(described in WO 00/34241 represented by the formula:



(2S)-1-{{{2-[(pyrimidine-2-ylpiperidine-4-yl}amino}-acetyl-2-cyano-pyrrolidine (described in WO 02/30890).

(2S)-1-{{{2-[(pyridine-2-yl)amino]ethyl}amino}acetyl}-2-cyano-pyrrolidine and (S)-1-{1-[5-(N,N'-dimethylaminosulfonyl)-2-pyridylamino]-2-methyl-propylamino}acetyl-2-pyrrolidinecarbonitrile (K-361) (described in WO 02/51836); thiazoline or pyrrolidine derivatives such as L-threo-isoleucylthiazoline (P32/98), L-allo-isolencylthiazoline, L-threo-isoeuculpryrrolidine, L-allo-isoleucylpyrrolidine and L-valylpyrrolidine (described in WO 01/72290) represented by the formula



(3) N-substituted 2-cyanopyrrole and 2-cyanopyrroline derivatives described in WO 01/55105. Preferably, (S,S)-1-(2-amino-3,3-dimethylbutylyl)-2,5-dihydro-1H-pyrrole-2-carbonitrile.

(4) Heterocyclic compounds described in WO 02/02560. Preferably, 7-benzyl-8-[6-hydroxymethyl)-1,4-diazepam-1-yl]-1,3-dimethyl-3,7-dihydropurine-2,6-dione.

(5) Pyrrolidine derivatives of a condensed ring with cyclopropane described in WO 01/68603. Preferably, (1S,3S,5S)-2-[(2S)-2-amino-3,3-dimethylbutyryl]-3-cyano-2-azabicyclo[3,1,0]hexane.

(6) Proline derivatives described in WO 02/14271. Preferably, (2S)-1-[(2S,4S)-4-(3-chloro-4-cyanophenyl)amino-2-pyrrolidinylcarbonyl]-2-cyanopyrrolidine.

(7) Cyanopyrrolidine derivatives described in WO 02/38541. Preferably, (2S,4S)-1-[(2S,3S)-2-amino-3-methyl-pentanoyl]-2-cyano-4-fluoropyrrolidine, (2S,4S)-2-cyano-4-fluoro-1-[(1-hydroxymethyl)cyclopentylamino]acetypyrrolidine, and (2S,4S)-2-


cyano-4-fluoro-1-[(1-hydroxy-3-adamanylamino]acetylpyrrolidine.

(8) Compounds such as P93/01 described in WO 02/02560, WO 03/055881, WO 03/040174, WO 03/037327, WO 03/035057, WO 03/035067, WO 03/024942, WO 03/024965, WO 03/004498, WO 03/004496, WO 03/000250, WO 03/002530, WO 03/002531, WO 03/002553, WO 03/000180, WO 03/000181, EP 1258476, WO 02/51836, WO 02/68420, US 6,432,969, etc.

(9) Another preferred DPP-4 inhibitor is the compound BMS-477118 disclosed in WO 2001068603 or U.S. Patent No. 6,395,767 (compound of example 60) also known as is (1S,3S,5S)-2-[(2S)-2-amino-2-(3-hydroxytricyclo[3.3.1.$1^{3,7}$]dec-1-yl)-1-oxoethyl]-2-azabicyclo[3.1.0]hexane-3-carbonitrile, benzoate (1:1) as depicted in Formula M of the patent application WO 2004/052850 on page 2, and the corresponding free base, (1S,3S,5S)-2-[(2S)-2-amino-2- (3-hydroxy-tricyclo[3.3.1.$1^{3,7}$]dec-1-yl)-1-oxoethyl]-2-azabicyclo-[3.1.0]hexane-3-carbonitrile (M') and its monohydrate (M") as depicted in Formula M of the patent application WO 2004/052850 on page 3. The compound BMS-477118 is also known as saxagliptin.

(10) Another preferred inhibitor is the compound GSK23A disclosed in WO 03/002531 (example 9) also known as (2S,4S)- 1-((2R)-2-Amino-3-[(4-methoxybenzyl)sulfonyl]-3-methylbutanoyl)-4-fluoropyrrolidine-2-carbonitrile hydrochloride.

(11) Other preferred DPP-IV inhibitors are described in the patent application WO 03/004498 especially examples 1 to 33 and most preferably the compound of the formula



MK-0431

described by the example 7 and also known as MK-0431 or Sitagliptin (INN).

(12) Other preferred DPP-IV inhibitors are described in the patent applications WO 2004/037169 especially those described in the examples 1 to 48 and WO 02/062764 especially the described examples 1 to 293, even preferred are the compounds 3-(aminomethyl)-2-isobuthyl-1-oxo-4-phenyl-1,2-dihydro-6-isoquinolinecarboxamide and 2-{[3-(aminomethyl)-2-isobuthyl-4-phenyl-1-oxo-1,2-dihydro-6-isoquinolyl]oxy}acetamide described on page 7 and also in the patent application WO2004/024184 especially in the reference examples 1 to 4.

Specific examples of hydroxypropyl methylcellulose include: Metholose SB-4 (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solution at 20°C: about 4 mPa·s), TC-5 RW (commercial name, Sinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solution at 20°C: about 6 mPa·s), TC-5 S (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solutionat 20°C: about 15 mPa·s), Metholose 60SH-50 (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity at 20°C: about 50 mPa·s), Metholose 65SH-50 (commercial name, Shinetsu Chemical Industry Co. Ltd. -made) (viscosity of 2 wt.% aqueous solution at 20°C: about 50 mPa·s), Metholose 90SH-100 (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solution at 20°C: about 100 mPa·s), Metholose 90SH-100SR (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) viscosity of 2 wt.% aqueous solution at 20°C: about 100 mPa·s), Metholose 65SH-400 (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solution at 20°C: 400 mPa·s), Metholose 65SH-1500 (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solution at 20°C: about 1500 mPa·s), Metholose 65SH-4000 (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous at 20°C: about 1000 mPa·s), Metholose 65SH-1000 (commercial name,

Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solution at 20 °C: about 1000 mPa·s), Metholose 90SH-1000 (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solution at 20°C: 4000 mPa·s), Metholose 90SH 1000SR (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solution at 20 °C: about 4000 mPa·s), Metholose 90SH-30000 (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solution at 20°C: about 30000 mPa·s), Metholose 90SH-100000 (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solution at 20°C: about 100000 mPa·s), and Metholose 90SH-100000SR (commercial name, Shinetsu Chemical Industry Co. Ltd. - made) (viscosity of 2 wt.% aqueous solution at 20 °C: about 100000 mPa·s).

"Sustained release preparation" or "modified release preparation" herein means a preparation wherein "the elution rate of the drug at 30 min after starting the test" is less than 85% when conducting, for example, the Second Method (paddle method) of the Elution Test Method in Japan Pharmacopocia using 900 mL of an appropriate test solution when the rotating counts of the drug in the preparation eluted by 100% into the test solution is below 1/3 the saturation solubility of the drug. Also, the conventional test solutions in the pharmaceutical technical field, e.g., water, a butter solution and the like are used. "preparation" means a pharmaceutical formulation or unit dosage form e.g. tablet, granule.

In addition, a preparation wherein the elution rate of a drug at 30 min after starting the test is not less than 85% when conducing the Second Method (paddle method) of the Elution Test Method in Japan Pharmacopoeia under conditions similar to the above is referred to as an immediate release preparation in this specification.

Dosage forms or unit dosage forms of the present sustained release preparation include, for example, an oral formulation such as a tablet, capsule (including a microcapsule), granule, or powder; and a parenteral formulation such as a suppository (e.g., rectal suppository, vaginal suppository, etc.), and these can be safely administered orally or parenterally, respectively. Of these forms, an oral formulation such as a tablet, capsule, or granule is preferred.

The sustained release preparation of the invention can be manufactured by mixing the DPP-IV inhibitor and hydrophilic polymer and molding them. Herein, the mixing and molding are performed according to conventional pharmaceutical methods. Also, when performing the said mixing and molding, a pharmacologically acceptable carrier may be used.

Herein, the pharmacologically acceptable carriers include different types of conventional organic or inorganic carrier substances, for example, excipients, lusters, binders, disintegrators, and the like as base materials for the preparation. Further, additives for a preparation such as preservatives, antioxidants, coloring agents, and sweeteners may also be used as needed.

When the DPP-IV inhibitor used in the sustained release preparation of the invention is basic, an organic acid may be added to regulate elution behavior of the sustained release preparation. Generally, since solubility of a basic drug is greater in the acidic condition than in the neutral condition, the drug elution from the sustained release preparation may vary depending on the surrounding pH. In such a case, change of the drug elution property based on the surrounding pH can be reduced by use of an organic acid. Since the pH in the body may vary in individual patients, reducing changes of the drug elution based on the surrounding pH is extremely significant for obtaining a uniform drug effect for various patients.

The organic acids include, for example, citric acid, tartaric acid, ascorbic acid, malic acid, fumaric acid, malonic acid, succinic acid, maleic acid, aspartic acid, glutamic acid, etc. Of these, citric acid, tartaric acid, ascorbic acid, etc. are preferred.

The sustained release preparation of the invention has less toxicity and fewer side effects, accordingly it can be administered to mammals (e.g. humans, bovines, horses, dogs, cats, monkeys, mice, and rats) as a preventative and therapeutic agent for various diseases.

The sustained release preparation of the invention may be used as a preventative and therapeutic agent, for example, for diseases such as diabetes (e.g. type 1 diabetes, type 2 diabetes, pregnancy diabetes), hyperlipidemia (e.g. hypertriglyceridemia, hypercholesterolemia, hypoHDLemia, postprandial hyperlipidemia), arterial sclerosis, Impaired Glucose Tolerance (IGT), Impaired Fasting Glucose (IFG), Impaired Fasting Glycemia (IFG), and diabetic complications (e.g., neuropathy, nephropathy, retinopathy, cataract, angiopathy of large vessel, osteopenia, diabetic hyperosmolar coma, infectious diseases (e.g., respiratory infection, urinary tract infection, digestive system infection, dermal soft tissue infection, lower limb infection), diabetic gangrene, dry oral cavity, decrease of audition, cerebrovascular disease, peripheral circulation disorder) and as an agent for reducing blood sugar and the like.

The sustained release preparation of the invention may prevent Abnormal Tolerance, Impaired Fasting Glucose (IFG) or Impaired Fasting Glycemeia (IFG) from developing into diabetes.

Further, the sustained release preparation of the invention can be used to improve pancreatic (β-cell) function, pancreatic (β-cell) regeneration, acceleration of pancreatic (β-cell) regeneration, etc.

Since the DPP-IV inhibitor is an agent accelerating glucose-depending insulin secretion which exhibits selectively accelerating action on insulin secretion in a patient with high blood sugar (e.g., patients who have not less than 126 mg/dL of blood sugar value at fasting or not less than 140 mg/dL of 2-hour value after 75 g Oral Glucose Tolerance Test (75 g OGTT)). The sustained release preparation of the invention is useful as a safe preventative and therapeutic agent for diabetes with less risk for blood vessel complication, induction of hypoglycemia, etc., which are harmful effects of insulin.

While the dosing amount of the sustained release preparation of the present invention varies depending on the subject, administration route, target disease, etc., for oral administration to an adult patient with diabetes, for example, a single dose of the DPP-IV inhibitor, which is the active ingredient, should usually be about 0.01-100 mg/kg weight, preferably 0.05-30 mg/kg weight, more preferably 0.1-10 mg/kg weight, and ideally be administered once or twice a day. Preferably, the sustained release preparation should be administered to continuously obtain DPP-IV inhibitory action in the living body during from before a meal to at least 2 hours after a meal (preferably 4 hours after a meal).

The sustained release preparation's release duration of the DPP-IV inhibitor in the living body should preferably be 1 to 24 hours; more preferably, 2 to 14 hours.

Usually, when using a DPP-IV inhibitor to prevent or treat diabetes, it is necessary to take the DPP-IV inhibitor before every meal since GLP-1, which is a substrate of DPP-IV, is secreted upon intake of food. However, since the sustained release preparation of the invention can release the DPP-IV inhibitor over a long time, it exhibits sufficient inhibitory effect of DPP-IV, even when taken once a day.

The sustained release preparation of the invention may be used in combination with drugs (hereinafter, referred to as

concomitant drugs) such as a therapeutic agent for diabetes, a therapeutic agent for diabetic complications, an antihyperlipidemia agent, a depressor, and anti-obesity agent, a diuretic, and an antithromotic agent. In this case, the medication time of the sustained release preparation of the invention and the comcomitant drug is not limited. Thus, these may be simultaneously administered to a subject, or administered at a different time, respectively. Further, the sustained release preparation of the invention and the concomitant drug may be administered as two preparations containing one active ingredient each, or administered as a single preparation containing both active ingredients.

The dosing amount of the concomitant drug can be appropriately selected based on the dose clinically used. The ratio of the sustained release preparation of the invention to the concomitant drug can be appropriately selected based on the subject, administration route, target disease, symptom, combination, etc. For example, when the subject is human, 0.01-100 parts by weight to 1 part by weight of the concomitant drug based on the DPP-IV inhibitor, which is an active ingredient of the sustained release preparation, may be used.

The said therapeutic agents for diabetes (antidiabetic) include, for example, insulin preparations (e.g., animal insulin preparations extracted from the pancreas of a bovine or swine; human insulin preparations synthesized in genetic engineering sing E. coli or yeast; zinc insulin; zinc protamine insulin; fragments or derivatives of insulin (e.g., INS-1, etc.)), insulin-resistance-improving agents (e.g. Pioglitazone hydrochloride, Lociglytazone (maleate), GI-262570, Reglixane (JTT-501), Netoglitazone (MCC-555), YM-440, KRP-297, CS-011, FK-614, Ragaglitazar (NN-622), Tesaglitazar (AZ-242), DMS-298585, EML-16336, compounds described in WO 99/58510 (e.g., (E)-4-[4-(5-methyl-2-phenyl-4-oxazolylmehtoxy)benzyloxyimino]-4-phenylbutyric acid)), PPARγ agonist, PPARγ antagonist, PPARγ/αdual agonist, α-glucosidase inhibitors (e.g. Voglibose,

Acarbose, Migritol, Emiglytate, biguanide agents (e.g., Phenformin, Metformin, Buformine or their salts) (e.g. hydrochloride, furmarate, succinate)), accelerators of insulin secretion (sulfonylurea agents (e.g., Tolbutamide, Glibenclamide, Gliclazide, Chlorpropamide, Tolazamide, Acetohexamide, Glyclopyramide, Glymepyride, Glypizide, Glybuzole, etc.), Repaglinide, Senaglinide, Nateglinide, Mitiglinide or their calcium salt hydrates), GLP-1 receptor agonists (e.g., GLP-1, NN-2211, AC-2993 (exedin-4), BIM-51077, Alb (8,35) h GLP-1 (7,37) NH_2), amyline agonists (e.g., Plamlintide, phosphotyrosine phosphatase inhibitors (e.g. vanadic acid), β3 agonists (e.g., CL-316243, SR-58611-A, UL-TG-307, SB-226552, AJ-9677, BMS-196085, AZ40140), inhibitors of neosugar (e.g., glycogen phosphorylase ihnhibitor, glucose-6-phosphatase inhibitor, glucagon antagonist, somatostatin receptor agonist), sodium-glucose cotransporter (SGLT) inhibitors (e.g., T-1095), etc.

The therapeutic agents for diabetic complications include inhibitors of aldose reductase (e.g., Tolrestat, Epalrestat, Zenarestat, Zopolrestat, Minalrestat, Phydarestat, SNK-860, CT-112), neurotrophy factor and its increasing agents (e.g., NGF, NT-3, BDNF, accelerators of neurotrophin production/secretion (e.g., 4-(4-chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-[3-(2-methylphenoxy)propyl]oxazole etc.)) described in WO 01/14372), ACE inhibitors (e.g., ALT946, Pimagedine, N-phenacylthiazolium bromide (ALT766), (EXO-226), active oxygen-crasing agents (e.g., thiocitic acid), cerebrovascular dilators (e.g., Tiapride, Mexiletine).

The anti-hyperdilidemia agents include statin compounds which are inhibitors of cholesterol synthesis (e.g., Cerivastatin, Pravastatin, Simvastatin, Rovastatin, Atorvastatin, Fluevastatin, Itavastatin, or their salts (e.g., sodium salts)), inhibitors or squalene synthase (e.g., compounds described in WO 97/10224, for example, N-[[(3R,5S)-1-(3-acetoxy-2,2-dimethylpropyl)-7-chloro-5-(2,3-dimethoxyphenyl-2-oxo-1,2,3,5-

terahydro-4,1-benzooxazepine-3-yl]acetyl]piperidine-4-azetic acid etc.), fibrate compounds (e.g., Benzafibrate, Clofibrate, Simfibrate, Clinofibrate), ACAT inhibitors (e.g., Avasimibe, Eflucimibe), anion exchange resins (e.g., Cholestyramine), Probucol, nicotinic acid drugs (e.g., nicomol, niceritrol), ethyl icosapentoate, plant sterols (soysterol), γ-oryzanol, etc.

The depressors include angiotensin converting enzyme inhibitors (e.g., Captropril, Enalapril, Delapril), angiotensin II antagonists (e.g., Candesartan, Cilexetil, Losartan, Eprosartan, Valsartan, Telmisartan, Irbesartan, Tasosartin), calcium antagonists (e.g., Manidipine, Nifedipine, Amlodipine, Efonidipine, Nicardipine), potassium channel opening drugs(e.g., Levcromakalim, L-27152, AL671, NIP-121), Clonidine, etc.

The anti-obesity agents include, for example, central anti-obesity agents (e.g., Dexphenfluramine, Phenfluramine, Phentelmine, Sibutramine, Anfepramone, Dexanfetamine, Mazindol, Phenylpropanolamine, Kobenzorex, pancreatic lipase inhibitors (e.g., Orlistat), β3 agonists (e.g., CL 316243, SR-58611-A, UL-TG-307, SB-226552, AJ-9677, BMS-196085, AZ40140), peptidyl anorectics (e.g., Leptin, Ciliary Neurotrophic Factor (CNTF)), cholecystokinin agonists (e.g., FPL-15849)), etc.

The diruetics include, for example, xanthine derivative (e.g., sodium salicylate theobromine, calcium salicylate theobromine), thiazide preparations (e.g. ethiazide, cyclopentiazide, trichlormethiazide, hydrochlorothiazide, hydroflumethiazide, bentylhydrochlorothiazide, penfluthizide, polythiaside, methichlothiazide), anti-aldosterone preparations (e.g., spylonoacetone, trimetellen), carbonic anhydrase inhibitors (e.g., acetazolamide), chlorbenzenesufonamide preparations (e.g., chlortalidone, meflucide, idapamide), azocernide, isosorbide, ethacrynic acid, pyrcladnide, bromethanide, flocemide, etc.

The antithrombotic agents include, for example, heparin (e.g., heparin sodium, heparin calcium, dalteparin sodium), warfarin

(e.g., warfarin potasium), antithrombin agents (e.g., aragatroban), thrombolytics (e.g. urokinase, tisokinase, alteplase, nateplase, monteplase, pamiteplase), platelet aggregation inhibitors (e.g., flolopidine hydrochloride, cllostazol, ethyl icosapentoate, beraprost sodium, sarpogrelate hudrochloride), etc.

The concomitant drugs are preferably insulin preparations, insulin-resistance improving agents, α-glucosidase inhibitors, biguanide agents, insulin secretion accelerators (preferably sulonyl urea), etc.

Further, the present invention relates to "a drug comprising a combination of two or more preparations with DPP-IV inhibitor and different releasing rates fo the DPP-IV inhibitor."

Herein, "the DPP-IV inbibitor-containing preparation" may be any preparation containing the DPP-IV inhibitor, either a sustained release or immediate release preparation. In addition, the release control mechanism of the DPP-IV inhibitor in the "DPP-IV inhibitor-containing preparation" has no particular limitations, and the preparation may be any that releases the DPP-IV inhibitor; by passive diffusion accompanying degradation of the prepartion; by response to change of the surrounding pH; by the inner pressure of the swollen inside of the preparation caused by intake of the surrounding water; immediate release by the preparation's disintegration or dissolution, and the like.

Herein, "preparations that release the DPP-IV inhibitor by passive diffusion" include, for example, the sustained release preparation of the invention, (preferably, a matrix tablet using hydorphylic polymers (e.g., hyroxypropylmethylcellulose, hydroxypropylcellulose, and polyethyleneoxixde), a matrix tablet using lipophilic base materials (e.g., carnauba wax, hydrogenated castor oil, hydrogenated rape oil, polyglycerin fatty acid esters), a tablet or a granule coated with sustained release base materials (e.g., cellulose polymers such as ethylcellulose, acrylic acid polymers such as aminoalkyl

methacrylate copolymer RS (Eudragit RS (commercial name, Rohm Pharma Co. Ltd. - made), ethyl acrylate-methyl methacrylate copolymer suspension (Eudragit NE (commercial name, Rohm Pharma Co., Ltd. - made), etc.

The "preparations that release DPP-IV inhibitor from the preparation by degradation of the preparation" include, for example, a capsule containing polyglycolated glycerides (e.g. Gelucire10/13 (commercial name, GATTEFOSSE Co. Ltd. - made)), etc.

The "preparations that release DPP-IV inhibitor in response to change of the surrounding pH" include, for example, a tablet or granule coated with enteric base materials (acrylic acid polymers such as methacrylate copolymer (Eudragit L (commercial name, Rohm Pharma Co. Ltd. - made)), methacrylate copolymer LD (Eudragit L-30D55 (commercial name, Rohm Pharma Co. Ltd. - made)), methyacrylate copolymer S (Eudragit S (commercial name, Rohm Pharma Co. Ltd. - made)), etc.

The "preparations that release DPP-IV inhibitor by the inner pressure of the swollen inside of the preparation caused by intake of the surrounding water" include, for example an osmotic system, i.e., Concerta™ (Alza Co. Ltd., Fort Washington, PA), etc.

The "preparations that release DPP-IV inhibitor immediately by the preparation's disintegration or dissolution" include, for example, those that are obtained by mixing the DPP-IV inhibitor with a pharmacologically acceptable carrier and then carrying out molding. Herein, the pharmacologically acceptable carrier includes those similar to the sustained release preparation of the invention. Also, mixing and molding can be performed according to conventional pharmaceutical techniques.

The release control mechanism of the "two or more preparations containing DPP-IV inhibitors" constituting the present drug of the invention may be either same or different from each other. The "two ore more preparations containing DPP-IV inhibitors" may

be a single preparation or plural preparations independent to each other. The single preparation includes a single capsule containing two ore more preparations containing the DPP-IV inhibitor; a multi-layer tablet (preferably a 2-layer tablet) or nucleated tablet having multiple release controlling parts, and the like.

The present drug of the invention should preferably be composed of a combination of a sustained release preparation containing a DPP-IV inhibitor and an immediate release preparation containing a DPP-IV inhibitor, and by adopting such a combination, the excellent DPP-IV inhibitory action can be obtained immediately after administration and be maintained over a long term.

While the content of the DPP-IV inhibitor in the sustained release preparation containing a DPP-IV inhibitor varies depending on the type of DPP-IV inhibitor, amount of preparation, etc., the content is, for example 20-30% by weight, preferably 25-35% by weight, and ideally 25% by weight.

Dosage forms of the DPP-IV inhibitor-containing preparation are similar to those of the sustained release preparation of the present invention.

The present drug of the invention has lower toxicity and fewer side effects, accordingly it can be administered to mammals (e.g., humans, bovines, horses, dogs, cats, monkeys, mice, and rats) as preventative and therapeutic agents of various diseases similarly to the said sustained release preparation of the invention.

The dosage of the present drug of the invention may be combined with the two or more preparations containing DPP-IV inhibitors at administration. Such a dosage form includes, for example, 1) administration as a single preparation of two ore more preparations containing DPP-IV inhibitors, 2) simultaneous administration as multiple preparations of two or more preparations containing DPP-IV inhibitors, 3) administration at

different times of multiple preparations of two or more preparations contining DPP-IV inhibitors, etc.

While the dosing amount of the drug of the invention varies depending on the subject, administration route, target disease, etc., when orally administering it to an adult patient with diabetes, for example, the usual does of DPP-IV inhibitor, which is the active ingredient should be about 0.01-100 mg/kg weight, preferably 0.5 mg/kg weight, more preferably .1-10 mg/kg weight, and these amounts should ideally be administered once or twice a day. Preferably, the drug of the invention should be administered to continuously obtain action of the DPP-IV inhibitor in the living body from before a meal to at least 2 hours after a meal preferably to 4 hours after a meal.

The present drug of the invention may be used in combination with the concomitant drug similar to the case of the sustained release preparation of the present invention.

The present invention further relates to "the release-controlled preparation containing the DPP-IV inhibitor, which may decrease DPP-IV activity in blood plasma by 10 to 90% (preferably, 10 to 85%) 1 hr after administration", "the release-controlled preparation containing the DPP-IV inhibitor which may decrease DPP-IV activity in blood plasma by 10 to 90% (preferably, 10 to 85%) 8 hrs after administration", "the release-controlled preparation contining the DPP-IV inhibitor which may decrease DPP-IV activity in blood plasma by 10 5o 90% (preferably, 10 to 85%) 12 hrs after administration", "the release-controlled preparation containing the DPP-IV inhibitor, which may decrease DPP-IV activity in the blood plasma by 10 to 90% (preferably, 10 to 85% over 1 to 8 hrs after administration", "the release-controlled preparation containing the DPP-IV inhibitor, which may decrease DPP-IV activity in blood plasma by 10 to 90% (preferably, 10 to 85%) over 1 to 12 hrs after administration", and the like.

Herein, the blood plasma in "DPP-IV activity in blood plasma" means peripheral venous blood plasma. Although the DPP-IV activity and its decrease rate may vary depending on the type of blood plasma (e.g., venous, arterial or portal blood plasma), any release-controlled preparations belong to the present invention of "release-controlled preparation containing the DPP-IV inhibitor" as long as they can decrease DPP-IV activity in peripheral venous blood plasma by 10 to 90% (preferably 10 to 85%).

The present invention of the release-controlled preparation containing the DPP-IV inhibitor can decrease DPP-IV activity in blood plasma by 10 to 90%, preferably 10 to 85%, more preferably 10 to 80%, most preferably 15 to 75%.

The DPP-IV activity in blood plasma can be measured according to, for example, "the method of Raymond et al., Diabetes, Vol. 47, p. 1253- 1258, 1998." As long as the said decrease rate of the DPP-IV activity in blood plasma is within normal errors, it may be different from the said values (10, 15, 75, 80, 85, 90%). Further, the decrease rate of the DPP-IV activity in blood plasma may be different from the said values depending on the method of measuring DPP-IV activity in blood plasma. For example, if measuring conditions such as substrate type, substrate concentration, reaction time, diluted multiples of blood plasma, and the like of DPP-IV activity in blood plasma are different from those in the method described in the above literature, the decrease rate of the DPP-IV activity in blood plasma may be greater than those described above, for example, the value of 90% may be over 95%.

Among preparations containing the DPP-IV inhibitor of the present invention, the "release-controlled preparation containing the DPP-IV inhibitor" of the present invention is a preparation wherein release of the DPP-IV inhibitor is controlled.

Such a preparation is preferably the inventive sustained release preparation. Also, among the drugs of the invention, "a drug comprising a combination of sustained release preparation containing a DPP-IV inhibitor and an immediate release preparation containing a DPP-IV inhibitor" is preferable.

The release-controlled preparation containing the DPP-IV inhibitor of the present invention has lower toxicity and fewer side effects, accordingly it can be administered to mammals (e.g., humans, bovines, horses, dogs, cats, monkeys, mice, and rats) as preventative and therapeutic agents of various diseases similarly to the sustained release preparation of the invention.

While the dosing amount of the release-controlled preparation containing the DPP-IV inhibitor of the present invention varies depending on he subject, administration route, target disease, etc., when orally administering it to an adult patient with diabetes, for example, the usual does of DPP-IV inhibitor, which is the active ingredient, should be about 0.01-100 mg/kg weight, preferably 0.05-30 mg/kg weight, more preferably 0.1-10 mg/kg weight, and ideally administered once or twice a day. Preferably, the release-controlled preparation containing the DPP-IV inhibitor of the present invention should be administered so that the action of the DPP-IV inhibitor in the living body is continuously obtained from before a meal to at least about 2 hours after a meal (preferably 4 hours after a meal).

The release-controlled preparation containing the DPP-IV inhibitor of the present invention may be used in combination with a concomitant drug as in the case of the sustained release preparation of the invention.

Specific examples of a microcrystalline cellulose include, for example Avicel® PH102 and 101 (FMC Biopolymer, Philadelphia, PA).

Specific examples of a magnesium stearate include, for example Synpro® Magnesium Stearate NF-Vegetable Grade (Ferro Corp., Walton Hills, OH).

Specific examples of a lactose include, for example, Lactose DT and Lactose SD.

In addition, tablets often contain a diluent or filler which are added to increase the bulk weight of the blend resulting in a practical size for compression. One, two, three or more diluent and/or filler can be selected. Examples of pharmaceutically acceptable fillers and pharmaceutically acceptable diluents include, but are not limited to, confectioner's sugar, compressible sugar, dextrates, dextrin, dextrose, lactose, mannitol, microcrystalline cellulose, powdered cellulose, sorbitol, sucrose and talc. The filler and/or diluent, e.g., may be present in an amount from about 15% to about 55% or from about 15% to about 45% by weight of the composition. Most preferably the above described compositions comprise one or two fillers selected from microcrystalline cellulose such as Avicel PH 102 and lactose.

Lubricants are typically added to prevent the tableting materials from sticking to punches, minimize friction during tablet compression and allow for removal of the compressed tablet from the die. Such lubricants are commonly included in the final tablet mix in amounts usually about or less than 1% by weight. The lubricant component may be hydrophobic or hydrophilic. Examples of such lubricants include stearic acid, talc and magnesium stearate. Magnesium stearate reduces the friction between the die wall and tablet mix during the compression and ejection of the tablets. It helps prevent adhesion of tablets to the punches and dies. Magnesium stearate also aids in the flow of the powder in the hopper and into the die. It has a particle size range of 450-550 microns and a density range of 1.00-1.80 g/mL. It is stable and does not polymerize within the tableting mix. The preferred lubricant, magnesium stearate is also employed in the formulation. Preferably, the lubricant is present in the tablet formulation in an amount of from about 0.25% to about 6%; also preferred is a level of about 0.5% to about 4% by weight; and most preferably

from about 0.1% to about 1% by weight. Other possible lubricants include talc, polyethylene glycol, silica and hardened vegetable oils. In an optional embodiment of the invention, the lubricant is not present in the formulation, but is sprayed onto the dies or the punches rather than being added directly to the formulation.

While the present invention is described in more detail in the following Experimental Details, the present invention is not limited to these and the uses of the invention may be varied as long as the scope is not exceeded.

Embodiments of the Invention

The subject invention provides a pharmaceutical tablet formulation (Formulation C) comprising per unit dosage form e.g. per tablet the following ingredients:

(a) a DPP-IV inhibitor preferably vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(b) a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution,

(c) and optionally a filler and/or a lubricant.

The subject invention provides a pharmaceutical tablet formulation (Formulation D) comprising per unit dosage form e.g. per tablet the following ingredients:

(a) 10-50% preferably 15-35% by weight on a dry weight basis of a DPP-IV inhibitor preferably vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(b) 20-60% preferably 30-50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution,

(c) and optionally a filler and/or a lubricant.

The subject invention provides a pharmaceutical tablet formulation (Formulation E) comprising per unit dosage form e.g. per tablet the following ingredients:

(a) 10-50% preferably 15-35% by weight on a dry weight basis of a DPP-IV inhibitor preferably vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(b) 20-60% preferably 30-50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) a filler; and

(d) a lubricant.

The subject invention provides a pharmaceutical tablet formulation (Formulation F) comprising per unit dosage form e.g. per tablet the following ingredients:

(a) 10-50% preferably 15-35% by weight on a dry weight basis of a DPP-IV inhibitor preferably vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(b) 20-60% preferably 30-50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) 15-55% preferably 25-45% by weight on a dry weight basis of a pharmaceutically acceptable filler; and optionally

(d) 0.1-10% preferably 0.1-3% by weight on a dry weight basis of a pharmaceutically acceptable lubricant.

The subject invention provides a pharmaceutical tablet formulation (Formulation G) comprising per unit dosage form e.g. per tablet the following ingredients:

(a) 10-50% preferably 15-35% by weight on a dry weight basis of a DPP-IV inhibitor preferably vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(b) 20-60% preferably 30-50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) 15-55% preferably 25-45% by weight on a dry weight basis of one or two pharmaceutically acceptable fillers selected from lactose and microcrystalline cellulose ; and optionally

(d) 0.1-10% preferably 0.1-3% by weight on a dry weight basis of a pharmaceutically acceptable lubricant such as magnesium stearate.

The subject invention provides a pharmaceutical tablet formulation (Formulation H) comprising per unit dosage form e.g. per tablet the following ingredients:

(a) 10-50% preferably 15-35% by weight on a dry weight basis of a DPP-IV inhibitor preferably vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(b) 20-60% preferably 30-50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000

cP (nominal value 100,000 cP) when present in a 1% solution;

(c) 25-40% by weight on a dry weight basis of one or two pharmaceutically acceptable fillers selected from lactose and microcrystalline cellulose ; and optionally

(d) 0.1-10% preferably 0.1-3% by weight on a dry weight basis of a pharmaceutically acceptable lubricant.

A pharmaceutical tablet formulation as described herein wherein the filler is selected from lactose and microcrystalline cellulose.

A pharmaceutical tablet formulation (Formulation I) as described herein comprising at least two fillers preferably lactose and microcrystalline cellulose. In an embodiment, the lactose is present in an amount from 1 to 8% preferably 1 to 5% by weight and microcrystalline cellulose is present in an amount from 25 to 35% by weight

A pharmaceutical tablet formulation (Formulation J) as described herein wherein 20-30% by weight on a dry weight basis of a DPP-IV inhibitor especially LAF237 is contained in the formulation.

A pharmaceutical tablet formulation (Formulation K) as described herein wherein the hydroxypropyl methyl cellulose is present in an amount from 34% to 46% preferably from 38% to 42% by weight.

A formulation as described herein wherein the compound (DPP-IV inhibitor) is 1-[(3-hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile (LAF237 or vildagliptin).


A formulation (Formulation L), capsule, granule or tablet as described herein wherein vildagliptin (LAF237) is a crystalline form of vildagliptin (crystal "Form A"), characterized by an X-ray diffraction pattern with peaks at about 16.6°, 17.1°, 17.2°

+/- 0.3 degrees 2-theta or characterized by an X-ray diffraction pattern with peaks at about 12.0°, 13.5°, 16.6°, 17.1°, 17.2°, 20.1°, 22.5°, 27.4°, 28.1° , +/- 0.3 degrees 2-theta. Such a crystal form is described in the International patent application No. PCT/US2006/001473

The term "vildagliptin" covers any crystalline form, preferably the crystal form "A" of vildagliptin.

The subject invention provides a pharmaceutical tablet formulation comprising per unit dosage form e.g. per tablet the following ingredients:

(a) a compound as an active ingredient, wherein the compound has a formula:



wherein R is substituted adamantyl and n is an integer from 0 to 3, preferably vildagliptin; or a pharmaceutically acceptable salt thereof;

(b) a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) a microcrystalline cellulose; and

(d) a magnesium stearate.

In an embodiment, relative to the weight of the formulation:

(a) the compound e.g. vildagliptin is present in an amount from 20% to 30% by weight;

(b) the hydroxypropyl methylcellulose is present in an amount from 30% to 50% by weight;

(c) the microcrystalline cellulose is present in an amount from 25% to 35% by weight; and

(d) the magnesium stearate is present in an amount from 0.1% to 3% by weight.

In another embodiment, relative to the weight of the formulation:

(a) the compound e.g. vildagliptin is present in an amount of about 25% by weight;

(b) the hydroxypropyl methylcellulose is present in an amount of about 40% by weight;

(c) the microcrystalline cellulose is present in an amount of about 30% by weight; and

(d) the magnesium stearate is present in an amount of about 1% by weight.

In a further embodiment, the tablet formulation further comprising a lactose. In an embodiment, the lactose is present in an amount from 1% to 8% by weight. In an additional embodiment the lactose is in an amount of about 4% by weight.

In an embodiment, the DPP-IV inhibitor is selected from [*S*]-1-[2-(5-cyano-2-pyridinylamino)ethylamino]acetyl-2-pyrolidine carbonitrile monohydrochloride, vildagliptin, L-threo-isoleucyl thiazolidine, Sitagliptin, Saxagliptin, 3-(aminomethyl)-2-isobuthyl-1-oxo-4-phenyl-1,2-dihydro-6-isoquinolinecarboxamide and 2-{[3-(aminomethyl)-2-isobuthyl-4-phenyl-1-oxo-1,2-dihydro-6-isoquinolyl]oxy}acetamide and optionally pharmaceutical salts thereof.

In an embodiment, the compound (DPP-IV inhibitor) is 1-[2-[(5-cyanopyridin-2-yl)amino]ethylamino]acetyl-2-cyano(S)-pyrrolidone or a pharmaceutically acceptable salt thereof.

In a preferred embodiment, the invention concerns a composition or unit dosage form as described herein wherein the compound (DPP-IV inhibitor) is vildagliptin or a salt thereof.

In yet another embodiment, the compound vildagliptin is present in the formulation or a pharmaceutically acceptable salt thereof.

The subject invention also provides a pharmaceutical tablet formulation (Formulation A), comprising per 400 mg tablet the following ingredients:

(a) vildagliptin or a pharmaceutically acceptable salt thereof in an amount of about 100 mg;

(b) a hydroxypropyl methylcellulose in an amount of about 160 mg, wherein the hydroxypropyl methylcellulose has an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) a microcrystalline cellulose in an amount of 120 mg;

(d) a lactose in an amount of about 16 mg; and

(e) a magnesium stearate in an amount of 4 mg.

The subject invention also provides a pharmaceutical tablet formulation (Formulation B), comprising per 600 mg tablet the following ingredients:

(a) vildagliptin or a pharmaceutically acceptable salt thereof in an amount of about 150 mg;

(b) a hydroxypropyl methylcellulose in an amount of about 240 mg, wherein the hydroxypropyl methylcellulose has an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) a microcrystalline cellulose in an amount of 180 mg;

(d) a lactose in an amount of about 24 mg; and

(e) a magnesium stearate in an amount of 6 mg.

In an embodiment, the pharmaceutical tablet formulation comprises combining the ingredients in the amounts recited above.

The subject invention also provides a method of inhibiting dipeptidyl peptidase IV activity in a subject comprising administering to the subject an amount of the above-mentioned pharmaceutical tablet formulations effective to inhibit the activity of dipeptidyl peptidase IV in the subject.

In an embodiment, the subject is a human being.

The subject invention also provides a method of treating a condition alleviated by dipeptidyl peptidase IV inhibition in a subject suffering from the condition comprising administering to the subject a therapeutically effective dose of any of the herein-mentioned pharmaceutical tablet formulations, unit dosage forms e.g. capsule, tablet, compressed table, direct compressed tablets, granule.

The subject invention also covers the use of any of the herein-mentioned pharmaceutical tablet formulations, unit dosage forms e.g. capsule, tablet, compressed table, direct compressed tablets, granule, for the manufacture of a medicament for the treatment of a condition alleviated by dipeptidyl peptidase IV inhibition in a subject suffering from the condition comprising administering to the subject a therapeutically effective dose of any of the above-mentioned pharmaceutical tablet formulations.

The subject invention also covers the use of any of the herein-mentioned pharmaceutical tablet formulations, unit dosage forms e.g. capsule, tablet, compressed table, direct compressed tablets, granule for the treatment of a condition alleviated by dipeptidyl peptidase IV inhibition in a subject suffering from the condition comprising administering to the subject a therapeutically effective dose of any of the above-mentioned pharmaceutical tablet formulations.

Preferably the condition alleviated by dipeptidyl peptidase IV inhibition is selected from non-insulin-dependent diabetes

mellitus, arthritis, obesity, allograft transplantation, calcitonin-osteoporosis, Heart Failure, Impaired Glucose Metabolism), IGT (Impaired Glucose Tolerance), neurodegenerative diseases such as Alzheimer's and Parkinson disease, modulating hyperlipidemia, modulating conditions associated with hyperlipidemia or for lowering VLDL, LDL and Lp(a) levels, cardiovascular or renal diseases e.g. diabetic cardiomyopathy, left or right ventricular hypertrophy, hypertrophic medial thickening in arteries and/or in large vessels, mesenteric vasculature hypertrophy, mesanglial hypertrophy, neurodegenerative disorders and cognitive disorders, to produce a sedative or anxiolytic effect, to attenuate post-surgical catabolic changes and hormonal responses to stress, to reduce mortality and morbidity after myocardial infarction, the treatment of conditions related to the above effects which may be mediated by GLP-1 and/or GLP-2 levels.

In an embodiment, the condition is non-insulin-dependent diabetes mellitus.


In another embodiment, the condition is obesity, arthritis, or osteoporosis. In a further embodiment, the subject is a human being.

The subject invention further provides a method of treating a condition alleviated by dipeptidyl peptidase IV inhibition in a subject suffering from the condition comprising administering to subject a therapeutically effective amount of any of the above-mentioned pharmaceutical tablet formulations in combination with a therapeutically effective dose of an anti-diabetic or arthritis drug.

In an embodiment, the subject is a human being.

The subject invention also provides a pharmaceutical tablet formulation comprising per unit dosage form e.g. per tablet the following ingredients:

(a) a compound as an active ingredient, wherein the compound has a formula:



wherein R is substituted adamantyl and n is an integer from 0 to 3; or a pharmaceutically acceptable salt thereof; and

(b) a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution.

In one embodiment, the herein described pharmaceutical tablet formulations also contains a filler. In one embodiment, the filler is lactose. In another embodiment, the filler is microcrystalline cellulose. In a further embodiment the filler is microcrystalline cellulose and lactose.

In another embodiment, the herein described pharmaceutical tablet formulations also contains a lubricant. In one embodiment, the lubricant is preferably magnesium stearate.

In yet another embodiment, relative to the weight of the formulation, the hydroxypropyl methyl cellulose is present in the herein described formulations in an amount from 30% to 50% by weight.

In an embodiment, the hydroxypropyl methyl cellulose is present in the herein described formulations in an amount from 34% to 46% preferably from 38% to 42% by weight.

In a further embodiment, the present invention also concerns the herein described pharmaceutical tablet formulation, wherein in the unit dosage form, the ratio of the weight of vildagliptin to the weight of hydroxypropyl methylcellulose is of 0.16 to 2.5, preferably 0.3 to 1.16 or 0.4 to 1.

Preferably the herein expressed ratios or % are expressed on a dry weight basis.

In a further embodiment, the present invention also concerns the herein described pharmaceutical tablet formulation, wherein in the unit dosage form, vildagliptin is present in an amount of 100 mg to 200 mg, preferably 100 mg, 150 mg, or 200 mg or the corresponding amount of any of its salt.

The herein described pharmaceutical tablet formulation can also be used in any unit dosage form, for example, an oral formulation such as a tablet (including multilayer tablets), capsule (including a microcapsule, capsules containing several e.g. two compartments), granule, or powder; or a parenteral formulation such as a suppository or any other pharmaceutical delivery system. Preferably the herein described pharmaceutical tablet formulation are used in the form of capsule, granule or tablet.

In a further embodiment, the present invention also concerns a pharmaceutical multilayer tablet wherein the herein described pharmaceutical tablet formulation, represents one of the tablet layers. This can be done as a multilayer tablet (one over the other) or as a coating one layer surrounding the other one.

Preferably the present invention also concerns a pharmaceutical multilayer tablet wherein the herein described pharmaceutical tablet formulation, represents one layer and a formulation comprising a further active ingredient represent the second layer and/or third layer. Preferably the further active ingredient is a glitazone (e.g. pioglitazone or rosiglitazone) or metformin. Preferably the further or second layer is an immediate release formulation. The herein described pharmaceutical tablet formulation can represent the middle layer and the immediate release formulation comprising a further active ingredient can represent the two external layers. The herein described pharmaceutical tablet formulation can represent the central layer and the immediate release formulation

comprising a further active ingredient can represent the surrounding coating layer.

In a further embodiment, the present invention also concerns a pharmaceutical tablet obtained by compression of the herein described pharmaceutical tablet formulation.

In a further embodiment, the present invention also concerns the above pharmaceutical tablet obtained by compression of the herein described pharmaceutical tablet formulation, wherein the herein described pharmaceutical tablet formulation is subject to roller compaction before compression into tablet.

In a further embodiment, the present invention also concerns the above pharmaceutical tablet wherein the hardness range is of between 10 to 13Kp for the 100 mg vildagliptin tablet, and between 11 to 25Kp for the 150 mg vildagliptin tablet.

In the development of the herein described pharmaceutical compositions, the applicant has discovered that the compressed tablets especially direct compressed tablet is particularly advantageous if the particles comprising the DPP-IV inhibitor especially LAF237, have a particle size distribution of less than 250 μm preferably between 10 to 250 μm.

Thus in a further embodiment the present invention concerns the herein described pharmaceutical tablet formulations, wherein the dispersion contains particles comprising DPP-IV inhibitor preferably LAF237, in free form or in acid addition salt form, and wherein at least 60%, preferably 80% and most preferably 90% of the particle size distribution in the formulation is less than 250 μm or preferably between 10 to 250 μm.

The present invention concerns the herein described pharmaceutical tablet formulations, wherein the dispersion contains particles comprising DPP-IV inhibitor preferably LAF237, in free form or in acid addition salt form, and wherein at least 60%, preferably 80% and most preferably 90% of the

particle size distribution in the formulation is greater than 10 μm.

The term "wherein at least 60%, preferably 80% and most preferably 90%" means at least 60%, preferably at least 80% and most preferably at least 90%.

The term "wherein at least at least 25%, preferably 35% and most preferably 45%" means at least 25%, preferably at least 35% and most preferably at least 45%.

In particular the present invention concerns the herein described pharmaceutical tablet formulations, wherein the dispersion contains particles comprising DPP-IV inhibitor preferably LAF237, in free form or in acid addition salt form, and wherein at least 25%, preferably 35% and most preferably 45% of the particle size distribution in the formulation is between 50 to 150 μm.

The selected particle size distribution provides an acceptable dissolution profile, as well as acceptable degrees of hardness, friability, Tablet breaking strength, Dispersion Quality and resistance to chipping, as well as a proper disintegration time, required flow and cohesive properties necessary to obtain an acceptable solid dosage form, Improved manufacturing robustness, required compactibility.

Particle size of drug, e.g. LAF237 particles size, is controlled by crystallisazion, drying and/or milling/sieving (non limiting examples are described below). Particle size can also be comminuted using roller compaction and milling/sieving. Producing the right particle size is well known and described in the art such as in "Pharmaceutical dosage forms: volume 2, 2nd edition, Ed.: H.A.Lieberman, L.Lachman, J.B.Schwartz (Chapter 3: SIZE REDUCTION)".

Multiple particle sizes have been studied and it has been discovered that the herein described specific size range provides unexpected good results for direct compaction.

PARTICLE SIZE DISTRIBUTION ESTIMATION BY ANALYTICAL SIEVING: Particle size distribution is measured using Sieve analysis, Photon Correlation Spectroscopy or laser diffraction (international standart ISO 13320-1), or electronic sensing zone, light obstruction, sedimentation or microscopy which are procedures well known by the person skilled in the art. Sieving is one of the oldest methods of classifying powders by particle size distribution. Such methods are well known and described in the art such as in any analytical chemistry text book or by the United State Pharmacopeia's (USP) publication USP-NF (2004 - Chapter 786 - (The United States Pharmacopeial Convention, Inc., Rockville, MD)) which describes the US Food and Drug Administration (FDA) enforceable standards. The used techniques are e.g. described in Pharmaceutical dosage forms: volume 2, 2nd edition, Ed.: H.A.Lieberman, L.Lachman, J.B.Schwartz is a good example. It also mentions (page 187) additional methods: Electronic sensing zone, light obstruction, air permeation, sedimentation in gas or liquid.

In an air jet sieve measurement of particle size, air is drawn upwards, through a sieve, from a rotating slit so that material on the sieve is fluidised. At the same time a negative pressure is applied to the bottom of the sieve which removes fine particles to a collecting device. Size analyses and determination of average particle size are performed by removal of particles from the fine end of the size distribution by using single sieves consecutively. See also "Particle Size Measurement", 5th Ed. , p 178, vol. 1; T. Allen, Chapman & Hall, London, UK, 1997, for more details on this. For a person skilled in the art, the size measurement as such is thus of conventional character.

The herein described formulations can also be in the form of a layer in a multi or 2-layer tablet.

The herein described formulations can also be in the form of capsules, tablets, compressed tables, direct compressed tablets, granules.

The DPP-IV inhibitor compounds especially vildagliptin, useful in this invention are hygroscopic, presents stability problems, and are not inherently compressible into tablets. Consequently, there is a need to provide a free-flowing and cohesive composition capable of being compressed, preferably directly, into strong tablets with an acceptable *in vitro* dissolution, pharmacokinetic, and pharmacodynamic profile.

The Paddle method to measure the drug dissolution rate (% of release) is used with 1000ml of 0.01N HCl. Such methods are well known and described in the art such as in any analytical chemistry text book or by the United State Pharmacopeia's (USP) publication USP-NF (2004 - Chapter 711) which describes the US Food and Drug Administration (FDA) enforceable standards.

In a further embodiment, the present invention concerns two pharmaceutical tablet formulations comprising respectively 100 mg or 150 mg of vildagliptin and providing a vildagliptin release profile particularly adapted for providing an improved pharmacokinetic profile. The resulting pharmacokinetic profile result in an improved treatment quality of the patient.

Thus, the present invention concerns;

1) a pharmaceutical tablet formulation comprising 100 mg of vildagliptin or a salt thereof, preferably compressed in the form of a sustained release tablet, wherein;

- between 10% and 16% preferably between 11% and 15% of vildagliptin is released after 0.5 hour,

- between 18% and 24% preferably between 19% and 23% of vildagliptin is released after 1 hour,

- between 30% and 36% preferably between 31% and 35% of vildagliptin is released after 2 hours,

- between 46% and 52% preferably between 47% and 51% of vildagliptin is released after 4 hour,

- between 58% and 64% preferably between 59% and 63% of vildagliptin is released after 6 hours,

- between 67% and 73% preferably between 68% and 72% of vildagliptin is released after 8 hours,

- between 74% and 80% preferably between 75% and 79% of vildagliptin is released after 10 hours,

- between 80% and 86% preferably between 81% and 85% of vildagliptin is released after 12 hours,

- between 91% and 97% preferably between 92% and 96% of vildagliptin is released after 18 hours,

- between 95% and 100% preferably between 96% and 100% of vildagliptin is released after 24 hours.

2) a pharmaceutical tablet formulation comprising 150 mg of vildagliptin or a salt thereof, preferably compressed in the form of a sustained release tablet, wherein;

- between 3.8% and 9.8% preferably between 4.8% and 8.8% of vildagliptin is released after 0.25 hour,

- between 8.1% and 14.1% preferably between 9.1% and 13.1% of vildagliptin is released after 0.5 hour,

- between 14.7% and 20.7% preferably between 15.7% and 19.7% of vildagliptin is released after 1 hours,

- between 25.3% and 31.3% preferably between 26.3% and 30.3% of vildagliptin is released after 2 hour,

- between 40.9% and 46.9% preferably between 41.9% and 45.9% of vildagliptin is released after 6 hours,

- between 62.1% and 68.1% preferably between 63.1% and 67.1% of vildagliptin is released after 8 hours,

- between 76.5% and 82.5% preferably between 77.5% and 81.5% of vildagliptin is released after 10 hours,

- between 83.5% and 89.5% preferably between 84.5% and 88.5% of vildagliptin is released after 12 hours,

- between 88.5% and 94.5% preferably between 89.5% and 93.5% of vildagliptin is released after 18 hours.

For the above described two pharmaceutical tablet formulations the Paddle method to measure the drug dissolution rate (% of release)has been used.

Preferably the above described two pharmaceutical tablet formulations comprise a hydroxypropyl methylcellulose, preferably between 20% and 60%, between 30% and 50% by weight on a dry weight basis of a hydroxypropyl methylcellulose.

Preferably the above described two pharmaceutical tablet formulations comprise a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution, preferably between 20% and 60%, or between 30% and 50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution.

Preferably the above described two pharmaceutical tablet formulations comprising 150 mg or 100 mg of vildagliptin are in the form of a tablet or multilayer tablet.

Preferably the above described two pharmaceutical tablet formulations comprising 150 mg or 100 mg of vildagliptin comprise any of the herein described pharmaceutical tablet formulations.

Preferably the above described two pharmaceutical tablet formulations are in the form of a tablet obtained by compression of a pharmaceutical tablet formulation as herein described,

wherein the pharmaceutical tablet formulation is subject to roller compaction before compression into tablet.

Preferably the above described pharmaceutical tablets comprise 100 mg of vildagliptin or a salt thereof wherein the tablet hardness range is of between 10 to 13Kp.

Preferably the above described pharmaceutical tablets comprise 150 mg of vildagliptin or a salt thereof wherein the tablet hardness range is of between 11 to 25Kp.

In a further aspect, the present invention concerns the use of the herein described pharmaceutical formulations, capsules, tablets, compressed tables, direct compressed tablets, granules for the treatment of conditions, such as non-insulin-dependent diabetes mellitus, arthritis, obesity, allograft transplantation, calcitonin-osteoporosis, Heart Failure, Impaired Glucose Metabolism), IGT (Impaired Glucose Tolerance), neurodegenerative diseases such as Alzheimer's and Parkinson disease, modulating hyperlipidemia, modulating conditions associated with hyperlipidemia or for lowering VLDL, LDL and Lp(a) levels, cardiovascular or renal diseases e.g. diabetic cardiomyopathy, left or right ventricular hypertrophy, hypertrophic medial thickening in arteries and/or in large vessels, mesenteric vasculature hypertrophy, mesanglial hypertrophy, neurodegenerative disorders and cognitive disorders, to produce a sedative or anxiolytic effect, to attenuate post-surgical catabolic changes and hormonal responses to stress, to reduce mortality and morbidity after myocardial infarction, the treatment of conditions related to the above effects which may be mediated by GLP-1 and/or GLP-2 levels.

In a further aspect, the present invention concerns a method of treating conditions, such as non-insulin-dependent diabetes mellitus, arthritis, obesity, allograft transplantation, calcitonin-osteoporosis, Heart Failure, Impaired Glucose Metabolism), IGT (Impaired Glucose Tolerance), neurodegenerative diseases such as Alzheimer's and Parkinson disease, modulating

hyperlipidemia, modulating conditions associated with hyperlipidemia or for lowering VLDL, LDL and Lp(a) levels, cardiovascular or renal diseases e.g. diabetic cardiomyopathy, left or right ventricular hypertrophy, hypertrophic medial thickening in arteries and/or in large vessels, mesenteric vasculature hypertrophy, mesanglial hypertrophy, neurodegenerative disorders and cognitive disorders, to produce a sedative or anxiolytic effect, to attenuate post-surgical catabolic changes and hormonal responses to stress, to reduce mortality and morbidity after myocardial infarction, the treatment of conditions related to the above effects which may be mediated by GLP-1 and/or GLP-2 levels, comprising administering to a warm-blooded animal in need thereof a therapeutically effective amounts of a pharmaceutical tablet formulation, capsules, tablets, compressed tables, direct compressed tablets, granules as herein described.

Preferably, the condition is selected from such as non-insulin-dependent diabetes mellitus, obesity, calcitonin-osteoporosis, Heart Failure, Impaired Glucose Metabolism, IGT (Impaired Glucose Tolerance), neurodegenerative diseases such as Alzheimer's and Parkinson disease, modulating hyperlipidemia, modulating conditions associated with hyperlipidemia or for lowering VLDL, LDL and Lp(a) levels.

In a further aspect, the present invention concerns a process for preparing a tablet, in unit dosage form, which comprises:

(a) blending a pharmaceutical tablet formulation as described herein above,

(b) compressing the formulation prepared during step (a) to form the compressed tablet in unit dosage form.

In a further aspect, the present invention concerns a process for preparing a tablet, in unit dosage form, which comprises:

(a) blending a pharmaceutical tablet formulation as described herein above,

(b) roller compacting the formulation prepared during step (a)

(c) compressing the formulation prepared during step (b) to form the compressed tablet in unit dosage form.

In a further aspect, the present invention concerns a process for preparing a tablet, in unit dosage form, which comprises:

(d) blending a pharmaceutical tablet formulation as described herein above,

(e) roller compacting the formulation prepared during step (a) with a compaction force comprised between 10 and 16 KN or any of the herein described preferred compaction forces,

(f) compressing the formulation prepared during step (b) to form the compressed tablet in unit dosage form.

In another embodiment, the present invention provides a 100 mg vildagliptin sustained release solid oral pharmaceutical dosage form which is;

i-1) a solid oral pharmaceutical dosage form comprising about 100 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma concentration of vildagliptin ranging from about 15.8 ng/mL +/- 6.85 ng/mL to about 173 ng/mL +/- 52 ng/mL between about 0.5 and about 16 hours following oral administration of said dosage form in a patient not treated with vildagliptin before said administration and wherein patient is under fasted conditions (day 1 of the above described study) , and/or

i-2) a solid oral pharmaceutical dosage form comprising about 100 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma

concentration of vildagliptin ranging from about 26.3 ng/mL +/- 13.1 ng/mL to about 175 ng/mL +/- 62.5 ng/mL between about 0.5 and about 16 hours following administration of said dosage form on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

i-3) a solid oral pharmaceutical dosage form comprising about 100 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma concentration of vildagliptin ranging from about 26.9 ng/mL +/- 14.1 ng/mL to about 186 ng/mL +/- 80.6 ng/mL between about 0.5 and about 16 hours following administration of said dosage form on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions (day 10 of the above described study) , and/or

ii-1) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-inf)}$ of vildagliptin ranging from about 1073 to about 1825 ng•h/mL i.e. 1449 ng•h/mL +/- 376 ng•h/mL following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions (day 1 of the above described study) , and/or

ii-2) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-24)}$ of vildagliptin ranging from about 1001 to about 1977 ng•h/mL i.e. 1489 ng•h/mL +/- 488 ng•h/mL following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has

been served an ADA breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

ii-3) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-24)}$ of vildagliptin ranging from about 1103 to about 2173 ng•h/mL i.e. 1638 ng•h/mL +/- 535 ng•h/mL following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions (day 10 of the above described study) , and/or

iii-1) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 3.61 hr +/- 1.44 hr following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions (day 1 of the above described study) , and/or

iii-2) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 2.59 hr +/- 1.4 hr following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

iii-3) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a

pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 3.74 hr +/- 1.44 hr following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions (day 10 of the above described study) , and/or

iv-1) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean C_{max} of vildagliptin of 205 ng/ml +/- 47 ng/ml following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions (day 1 of the above described study) , and/or

iv-2) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean C_{max} of vildagliptin of 200 ng/ml +/- 64 ng/ml following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

iv-3) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean C_{max} of vildagliptin of 245 ng/ml +/- 68 ng/ml following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein

said patient is under fasted conditions (day 10 of the above described study) , and/or

v-1) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean % inhibition of DPP-IV activity over 24 hours of 85.64% +/- 12.76% following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions (day 1 of the above described study) , and/or

v-2) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean % inhibition of DPP-IV activity over 24 hours of 87.78% +/- 16.37% following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

v-3) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean % inhibition of DPP-IV activity over 24 hours of 90.20% +/- 7.35% following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions (day 10 of the above described study) , and/or

vi-1) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as

substantially depicted in figure 24, following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions (day 1 of the above described study) , and/or

vi-2) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as substantially depicted in figure 25, following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

vi-3) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as substantially depicted in figure 26, following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions (day 10 of the above described study).

Preferably the 100 mg vildagliptin sustained release solid oral pharmaceutical dosage form comprises a hydroxypropyl methylcellulose, preferably between 20% and 60%, between 30% and 50% or between 34% and 46% by weight on a dry weight basis of a hydroxypropyl methylcellulose.

Preferably the 100 mg vildagliptin sustained release solid oral pharmaceutical dosage form comprises a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution, preferably between 20% and 60%, or between 30% and 50%

or between 34% and 46% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution.

Preferably the 100 mg vildagliptin sustained release solid oral pharmaceutical dosage form comprises any of the hereinabove described pharmaceutical tablet formulations.

Preferably the 100 mg vildagliptin sustained release solid oral pharmaceutical dosage form comprises a pharmaceutical tablet formulations selected from the herein described formulations A, B, C, D, E, F, G, H, I, J, K, or L.

In another embodiment, the present invention provides a 150 mg vildagliptin sustained release solid oral pharmaceutical dosage form which is;

i-a) a solid oral pharmaceutical dosage form comprising about 150 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma concentration of vildagliptin ranging from about 30.7 ng/mL +/- 21.9 ng/mL to about 223 ng/mL +/- 77.3 ng/mL between about 0.5 and about 16 hours following oral administration of said dosage form in a patient not treated with vildagliptin before said administration and wherein patient is under fasted conditions (day 1 of the above described study), and/or

i-b) a solid oral pharmaceutical dosage form comprising about 150 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma concentration of vildagliptin ranging from about 48.7 ng/mL +/- 25.8 ng/mL to about 223 ng/mL +/- 99.7 ng/mL between about 0.5 and about 16 hours following administration of said dosage form on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA

breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

i-c) a solid oral pharmaceutical dosage form comprising about 150 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma concentration of vildagliptin ranging from about 44.6 ng/mL +/- 28.5 ng/mL to about 263 ng/mL +/- 84.4 ng/mL between about 0.5 and about 16 hours following administration of said dosage form on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions (day 10 of the above described study) , and/or

ii-a) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-inf)}$ of vildagliptin ranging from about 1346 to about 3196 ng•h/mL i.e. 2271 ng•h/mL +/- 925 ng•h/mL following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions (day 1 of the above described study) , and/or

ii-b) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-24)}$ of vildagliptin ranging from about 1277 to about 3207 ng•h/mL i.e. 2242 ng•h/mL +/- 965 ng•h/mL following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

ii-c) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a

pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-24)}$ of vildagliptin ranging from about 1643 to about 3273 ng•h/mL i.e. 2458 ng•h/mL +/- 815 ng•h/mL following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions (day 10 of the above described study) , and/or

iii-a) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 3.57 hr +/- 1.17 hr following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions (day 1 of the above described study) , and/or

iii-b) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 2.87 hr +/- 1.59 hr following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

iii-c) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 4.13 hr +/- 1.24 hr following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein

said patient is under fasted conditions (day 10 of the above described study) , and/or

iv-a) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean C_{max} of vildagliptin of 257 ng/ml +/- 59 ng/ml following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions (day 1 of the above described study) , and/or

iv-b) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean C_{max} of vildagliptin of 272 ng/ml +/- 111 ng/ml following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

iv-c) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean C_{max} of vildagliptin of 308 ng/ml +/- 91 ng/ml following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions (day 10 of the above described study) , and/or

v-a) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean % inhibition of

DPP-IV activity over 24 hours of 90.04% +/- 11.91% following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions (day 1 of the above described study) , and/or

v-b) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean % inhibition of DPP-IV activity over 24 hours of 90.4% +/- 17.50% following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

v-c) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean % inhibition of DPP-IV activity over 24 hours of 91.64% +/- 8.47% following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions (day 10 of the above described study) , and/or

vi-a) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as substantially depicted in figure 24, following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions (day 1 of the above described study) , and/or

vi-b) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as substantially depicted in figure 25, following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form (day 9 of the above described study) , and/or

vi-c) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as substantially depicted in figure 26, following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions (day 10 of the above described study).

Preferably the 150 mg vildagliptin sustained release solid oral pharmaceutical dosage form comprises a hydroxypropyl methylcellulose, preferably between 20% and 60%, between 30% and 50% or between 34% and 46% by weight on a dry weight basis of a hydroxypropyl methylcellulose.

Preferably the 150 mg vildagliptin sustained release solid oral pharmaceutical dosage form comprises a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution, preferably between 20% and 60%, or between 30% and 50% or between 34% and 46% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution.

Preferably the 150 mg vildagliptin sustained release solid oral pharmaceutical dosage form comprises any of the hereinabove described pharmaceutical tablet formulations.

Preferably the 150 mg vildagliptin sustained release solid oral pharmaceutical dosage form comprises a pharmaceutical tablet formulations selected from the herein described formulations A, B, C, D, E, F, G, H, I, J, K or L.

Any of the above described 100 mg or 150 mg vildagliptin sustained release solid oral pharmaceutical dosage form wherein;

i) the formulation is a matrix formulation containing a pharmaceutically acceptable hydrophilic polymer which can retard diffusion of vildagliptin,

ii) the solid oral pharmaceutical dosage form is a compressed tablet, and optionally

iii) the elution rate of vildagliptin at 30 minutes after starting the test is less than 30% when conducting the Paddle method.

"pharmaceutically acceptable hydrophilic polymer which can retard diffusion of an active ingredient" are known in the art e.g. Hydroxypropyl-methylcelluose.

Experimental Details

I. Pharmaceutical Development and Manufacture

A. Selected Formulation and Process Composition

A dry blend process with roller compaction was developed for direct compression, as shown in the process flow diagram (Figure 1). Two dosage strengths, 100 mg and 150 mg of vildagliptin, were selected for further development and are shown in Tables 2-1 and 2-2. This process utilizes a common dry blend for the two strengths.

Table 2-1 Composition per 100mg dosage form unit for Market Formulation containing 40% HPMC K100M

Product name: vildagliptin Tab 100mg				
Basis number/ Variant: 3768652.003				
Shape /size /color /imprint: 11mm FFBE (round)/white to off-white				
Global material no.	**Component**	**Composition per unit (mg)**	**Testing monograph**	**Function**
146082	vildagliptin	100	Novartis	Drug Substance
103266	Microcrystalline cellulose, PH102	120	Ph. Eur., NF	Tablet filler
103282	Lactose, anhydrous DT	16	Ph. Eur., NF	Tablet filler
151580	Hydroxypropyl-methylcelluose K100M	160	Ph. Eur., NF	Controlled release polymer
100217	Magnesium stearate	4	Ph. Eur., NF	Tablet lubricant
Total weight		**400**		

Table 2-2 Composition per 150mg dosage form unit for Market Formulation containing 40% HPMC K 100M

Product name: vildagliptin Tab 150mg				
Basis number/ Variant: 6001732.001				
Shape /size /color /imprint: 17 x 6.7mm ovaloid/white to off-white				
Global material no.	**Component**	**Composition per unit (mg)**	**Testing monograph**	**Function**
146082	vildagliptin	150	Novartis	Drug Substance
103266	Microcrystalline cellulose, PH102	180	Ph. Eur., NF	Tablet filler
103282	Lactose, anhydrous DT	24	Ph. Eur., NF	Tablet filler
151580	Hydroxypropyl-methylcellulose K100M	240	Ph. Eur., NF	Controlled release polymer
100217	Magnesium stearate	6	Ph. Eur., NF	Tablet lubricant
Total weight		**600**		

Process

The selected manufacturing process is shown in figure 2-1.

B. Formulation and Alternative Variants

Several formulation variants were evaluated as potential clinical service forms (CSF) in order to provide the requested release rate profile. The term clinical service forms (CSF) means a formulation adapted for administration to a patient. Several types of release rate profiles were studied; in particular a slow and fast. The profile rate was determined by the amount and type of polymer selected and/or tablet technology. The Market Formulation (MF) selected is a specific slow release rate profile (see Section I.A.)particularly adapted for a once a day dosage regimen and production of compressed tablets. Examples of the fast release rate formulations are shown in Tables 2-3 to 2-7 and alternate formulations with slow release profiles are shown in Tables 2-8 to 2-10.

Table 2-3 Composition per 100mg dosage form unit for fast release profile containing 20% HPMC K100LV

Product name: vildagliptin Tab 100mg				
Basis number/ Variant: 3768652.001				
Shape /size /color /imprint: /white to off-white				
Global material no.	**Component**	**Composition per unit (mg)**	**Testing monograph**	**Function**
146082	vildagliptin	100	Novartis	Drug Substance
103266	Microcrystalline cellulose, PH102	170	Ph. Eur., NF	Tablet filler
103282	Lactose, anhydrous DT	46	Ph. Eur., NF	Tablet filler
151580	Hydroxypropyl-methylcellulose K100LVP	80	Ph. Eur., NF	Controlled release polymer
100217	Magnesium Stearate	4	Ph. Eur., NF	Tablet lubricant
Total weight		**400**		

Table 2-4 Composition per 200mg dosage form unit for fast release profile containing 20% HPMC K100LV

Product name: vildagliptin Tab 200mg				
Basis number/ Variant: 3768660.002				
Shape /size /color /imprint: /white to off-white				
Global material no.	**Component**	**Composition per unit (mg)**	**Testing monograph**	**Function**
146082	vildagliptin	200	Novartis	Drug Substance
103266	Microcrystalline cellulose, PH102	340	Ph. Eur., NF	Tablet filler
103282	Lactose, anhydrous DT	92	Ph. Eur., NF	Tablet filler
151580	Hydroxypropyl-methylcellulose K100LVP	160	Ph. Eur., NF	Controlled release polymer
100217	Magnesium Stearate	8	Ph. Eur., NF	Tablet lubricant
Total weight		**800**		

Table 2-5 Composition per 200mg dosage form unit for fast release profile containing 20% HPMC E10M

Product name: vildagliptin Tab 200mg				
Basis number/ Variant:				
Shape /size /color /imprint: /white to off-white				
Global material no.	**Component**	**Composition per unit (mg)**	**Testing monograph**	**Function**
146082	vildagliptin	200	Novartis	Drug Substance
103266	Microcrystalline cellulose, PH102	242	Ph. Eur., NF	Tablet filler
103282	Lactose, anhydrous DT	32	Ph. Eur., NF	Tablet filler
151580	Hydroxypropyl-methylcelluose, E10M	120	Ph. Eur., NF	Controlled release polymer
100217	Magnesium stearate	6	Ph. Eur., NF	Tablet lubricant
Total weight		**600**		

The concentration of HPMC E10M was varied from 20% to 35% to match the target release rate profile. A combination of HPMC E10 (15%) and HPMC K100M (10%) was also evaluated for release rate.

Table 2-6 Composition per 100mg dosage form unit for fast release profile containing X% polyox

Product name: vildagliptin Tab 100mg				
Basis number/ Variant:				
Shape /size /color /imprint: /white to off-white				
Global material no.	**Component**	**Composition per unit (mg)**	**Testing monograph**	**Function**
146082	vildagliptin	100	Novartis	Drug Substance
103266	Microcrystalline cellulose, PH102	170	Ph. Eur., NF	Tablet filler
103282	Lactose, anhydrous DT	46	Ph. Eur., NF	Tablet filler
	Polyox ??			Controlled release polymer
100217	Magnesium Stearate	4	Ph. Eur., NF	Tablet lubricant
Total weight		**400**		

Bi-layer tablet technology was evaluated using a 50mg immediate release (IR) tablet in one layer, and an 150mg MR as the second layer as shown in Table 2-7.

Table 2-7 Composition per 200mg dosage form unit for bi-layer tablet containing HPMC K100M

Product name: vildagliptin Tab 200mg				
Basis number/ Variant: NA				
Shape /size /color /imprint: /white to off-white				
Global material no.	**Component**	**Composition per unit (mg)**	**Testing monograph**	**Function**
First Layer (IR)				
146082	vildagliptin	50	Novartis	Drug Substance
103266	Microcrystalline cellulose, PH102	96	Ph. Eur., NF	Tablet filler
103282	Lactose, anhydrous DT	48	Ph. Eur., NF	Tablet filler
	Explotab	4	Ph. Eur., NF	Disintegrant
	Magnesium Stearate	2	Ph. Eur., NF	Lubricant
Total		200		
Second Layer (MR)				
146082	1-[(3-Hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile	150	Novartis	Drug Substance
151580	Hydroxypropyl-methylcellulose K100M	130	Ph. Eur., NF	Controlled release polymer
103266	Microcrystalline cellulose, PH102	120	Ph. Eur., NF	Tablet filler
103282	Lactose, anhydrous DT	46	Ph. Eur., NF	Tablet filler
100217	Magnesium stearate	4	Ph. Eur., NF	Tablet lubricant
Total weight		**450**		

The amount of vildagliptin DS and HPMC K100M were varied in each layer to provide a suitable release rate profile. For example, the DS was varied from 25mg to 50mg (immediate release layer) and 150mg to 175mg (modified release layer). The amount of HPMC K100M was varied from 29% to 40% (modified release layer).

Another exploratory slow release variant was the HPMC concentration was reduced to 30%, with and without roller compaction, as shown in Tables 2-8 and 2-9 . This change did not have an effect on the dissolution profile. However, 30% HPMC K100M exhibited a lower flow quality, 0.21 and 0.37, for both non-roller compacted and roller compacted material, respectively, when compared to 40% HPMC K100M. Filming was also observed during compression with 30% HPMC K100M.

Table 2-8 Composition per 100mg dosage form unit with 30% HPMC

Product name: vildagliptin Tab 100mg				
Basis number/ Variant: NA				
Shape /size /color /imprint: 11mm FFBE (round)/white to off-white				
Global material no.	**Component**	**Composition per unit (mg)**	**Testing monograph**	**Function**
146082	vildagliptin	100	Novartis	Drug Substance
103266	Microcrystalline cellulose, PH102	120	Ph. Eur., NF	Tablet filler
103282	Lactose, anhydrous DT	58	Ph. Eur., NF	Tablet filler
151580	Hydroxypropyl-methylcelluose K100M	120	Ph. Eur., NF	Controlled release polymer
100217	Magnesium stearate	4	Ph. Eur., NF	Tablet lubricant
Total weight		**400**		

Table 2-9 Composition per 150mg dosage form unit with 30% HPMC

Product name: vildagliptin Tab 150mg				
Basis number/ Variant: NA				
Shape /size /color /imprint: 17 x 6.7mm ovaloid/white to off-white				
Global material no.	**Component**	**Composition per unit (mg)**	**Testing monograph**	**Function**
146082	vildagliptin	150	Novartis	Drug Substance
103266	Microcrystalline cellulose, PH102	180	Ph. Eur., NF	Tablet filler
103282	Lactose, anhydrous DT	24	Ph. Eur., NF	Tablet filler
151580	Hydroxypropyl-methylcelluose K100M	180	Ph. Eur., NF	Controlled release polymer
100217	Magnesium stearate	6	Ph. Eur., NF	Tablet lubricant

Table 2-10 Composition per 200mg dosage form unit for slow release profile containing 40% HPMC K100M

Product name: vildagliptin Tab 200mg				
Basis number/ Variant: 3768660.003				
Shape /size /color /imprint: /white to off-white				
Global material no.	**Component**	**Composition per unit (mg)**	**Testing monograph**	**Function**
146082	vildagliptin	200	Novartis	Drug Substance
103266	Microcrystalline cellulose, PH102	240	Ph. Eur., NF	Tablet filler
103282	Lactose, anhydrous DT	32	Ph. Eur., NF	Tablet filler
151580	Hydroxypropyl-methylcelluose K100M	320	Ph. Eur., NF	Controlled release polymer
100217	Magnesium Stearate	8	Ph. Eur., NF	Tablet lubricant
Total weight		**800**		

Selection of Formulations for the Manufacture of Clinical Variants

The four formulations selected for further clinical evaluation comprised of both types of release rate profiles (slow and fast)

at two dosage strengths (100mg and 200mg). The batch size was 4,000 tablets (or 3.1kg). Pre-compression was required to meet the hardness and/or friability requirements. The slow formulation contained 40% HPMC K100M as shown in Tables 2-1 and 2-10. The fast formulation contained 20% HPMC K100 LVP as shown in Tables 2-3 and 2-4.

Table 2-11 Clinical batch summary

Rate Profile	Strength	Polymer	Composition	KN Number/ Variant	Batch Number
Slow	100mg	40% HPMC K100M	Table 2-1	3768652.002	AEUS20020081
Slow	200mg	40% HPMC K100M	Table 2-10	3768660.003	AEUS20020083
Fast	100mg	20% HPMC K100LVP	Table 2-3	3768652.001	AEUS20020080
Fast	200mg	20% HPMC K100LVP	Table 2-4	3768660.002	AEUS20020082

Formulation Rationale

1-[(3-Hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile i.e. vildagliptin DS (Drug Substance) is highly water soluble (up to 125mg/mL) and so requires polymers that can act as substantial diffusion barriers for a dramatic reduction in drug dissolution and diffusion. A once a day MR dosage form is likely to require a significant level of polymer load for retarding drug diffusion. During the development, the applicant has selected monolithic formulation with low risk for dose dumping, eliminating coating technologies in favor of a matrix formulation. After intensive studies on compatibility, ability to retard diffusion pharmacokinetic and pharmacodynamic studies, and regulatory acceptance requirements the applicant narrowed the polymer choice to hydrophilic polymers, especially hydroxpropyl methyl cellulose (HPMC).

Major and Critical Findings During Formulation Development

A critical excipient is the control release polymer, HPMC, as the K100M grade. During the CSF development, various control

release polymers were evaluated alone, or in combination, in order to achieve the desired release profile. HPMC, K100M grade, provided the best match to the desired release profile. The "slow" formulation was selected after evaluation in man.

As previously mentioned, pre-compression was utilized during the clinical batch manufacture. Experiments were conducted to further evaluate the CSF formulation and the need for pre-compression.

The compression profile for the CSF formulation containing 40% HPMC is shown in Figures 3 and 4.

The data indicated that the compression profile and friability could be improved with pre-compression. Additional studies were conducted to improve the compression profile and to avoid the use of pre-compression.

As shown in Figure 5, 30% HPMC K100M was evaluated to improve the compression profile. The effect of tablet dwell time was simultaneously evaluated at speeds from 40 to 80rpm.

It was determined that an improvement in the compression profile was achieved using roller compaction. As shown in Figures 6 and 7, a Fitzpatrick (Chilsonator®, model IR220) was evaluated with the interim powder blend without the addition of magnesium stearate, at increasing compaction forces.

As shown in Figures 8 and 9, the effect of roller compaction on the dissolution profile was studied with 40% HPMC K100M and compared to the dissolution profile of the clinical batch. It was concluded that roller compaction did not impact the dissolution profile.

The roller compaction process was scaled to a 50mm Bepex roller compactor using a linear relationship developed for a 50mm Fitzpatrick roller compactor (in lb/in units, model IR520) and a 50mm Bepex (in KN units) roller compactor. As shown in Figures 10 and 11, the effect of increasing compaction force was studied. It was concluded that the process was scaleable to a

Bepex roller compactor, and the tablet hardness was not significantly affected with increasing roller compaction force (13-31KN), and roller compactor roll speed (4-8rpm).

Overview of Compositions Including CSF in Clinical Studies

See section "*B. Formulation and Alternative Variants.*"

C. Manufacturing Process

All variants were manufactured as detailed in the process flow chart shown in Figure 1, except variants without roller compaction. A batch size of up to 12 Kg was used. Drug substance from lots; 0223007, 0223008, 0223009 were used for all batches manufactured

Description and Assessment of the Processes Tested

The effect of roller compaction was evaluated with 30% and 40% HPMC. Roller compaction was utilized to densify the powder blend prior to compression. The effect of Fitzpatrick (Chilsonator®, model IR220) roller compactor at compaction forces from 500 to 10,000 lb/in on a powder blend containing 40% HPMC, are shown in Figures 12 and 13.

The data indicated that with increasing roller compaction force, led to a decrease in tablet hardness. As shown in Figure 14, roller compaction forces greater than 5,000 lb/in (or 43.75 KN) produced tablets with a compression profile worst than the CSF without roller compaction or pre-compression.

The effect of roller compaction on tablet friability is shown in Figure 14.

Key Section Criteria and Rationale for Final Process

The key criteria for the final process was based on the dissolution profile. Since a PK/safety study was conducted *in vivo* using the 100mg, 150mg and 200mg tablets, the formulation containing 40% HPMC K100M was desired. Roller compaction was added to the process to improve the compression profile. As shown in Figures 17 and 18, the dissolution profile for roller

compacted material did not differ from the CSF dissolution profile. The Japanese Ministry of Health has imposed 40mg/kg as a daily intake limit for HPMC K100M. It is not anticipated that the daily intake limits will become a regulatory hurdle for the 1-[(3-Hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile MF.

The physical properties of the various powder blends are shown in Tables 2-12 and 2-13.

Table 2-12 Physical properties of roller compacted powder blend (Fitzpatrick)

Sample	Roller Compaction Force (lb/in) [calculated KN]	Powder Flow Quality (Sotax)	Bulk Density/Tap Density (g/mL)
TRD-1926-78 40% HPMC	NA	0.38	0.38/0.576
TRD-1971-53 40% HPMC	NA	0.33	0.38/0.66
TRD-1971-55 40% HPMC	NA	0.36	0.36/ 0.59
TRD-1926-79 30% HPMC	NA	0.21	0.385/ 0.60
TRD-1926-79 30% HPMC	1,000 [8.75]	0.37	0.432/ 0.645
TRD-1926-01 40% HPMC	500 [4.37]	0.25	0.398/ 0.588
TRD-1926-78 40% HPMC	1,000 [8.75]	0.49	0.448/ 0.652
TRD-1926-01 40% HPMC	1,500 [13.1]	0.53	0.448/ 0.666
TRD-1771-123B 40% HPMC	5,000 [43.75]	0.76	0.526/ 0.714
TRD-1771-123C 40% HPMC	10,000 [87.5]	ND (not tabletable)	0.566/ 0.732
Avicel PH102	NA	0.64	0.29/ 0.35

Table 2-13 Physical properties of roller compacted powder blend (Bepex)

Sample	Roller Compaction Force (Kn) [calculated lb/in]	Roll Speed (rpm)	Powder Flow Quality (Sotax)	Bulk Density/Tap Density (g/mL)
TRD-1926-78 40% HPMC	NA	NA	0.38	0.38/0.576
TRD-1971-53 40% HPMC	NA	NA	0.33	0.38/0.66
TRD-1971-55 40% HPMC	NA	NA	0.36	0.36/ 0.59
TRD-1971-53C 40% HPMC	13 [1500 lb/in]	4	0.25	0.398/ 0.588
TRD-1971-53A 40% HPMC	22 [2500 lb/in]	4	0.49	0.448/ 0.652
TRD-1971-53B2 40% HPMC	31 [3500 lb/in]	4	0.53	0.448/ 0.666
TRD-1971-55A 40% HPMC	13	4	0.76	0.526/ 0.714
TRD-1971-55B 40% HPMC	13	6	ND (not tabletable)	0.566/ 0.732
TRD-1971-55C 40% HPMC	13	8		
Avicel PH102	NA	NA	0.64	0.29/ 0.35

The use of roller compaction indicated an increase in the powder flow quality and an increase in bulk density. This was limited by the roller compaction force, where tablets from 5,000 lb/in and 10,000 lb/in force exhibited a poor friability of 0.6% and > 10%, respectively (100mg tablets).

The effect of roller compaction on the sieve analysis is shown in Figures 19 and 20. All roller compacted material was screened using an 18 mesh screen. Selection of screen size as a means to

particle size control, and will be further evaluated during the Lab Phase FMI.

Equipment Used

Table 2-14 Process equipment

Description	Model
Bin Blender	LB Bohle, 10L, 25L, 40L
Oscillating mill	Ferwitt
Roller compactor	Fitzpatrick, Chilsonator, IR220
Roller Compactor	Bepex, 50mm
Tablet Press	Manesty Beta Press, 16 sta.

Statement on the Up-Scaling Potential and Robustness of the final process

Table 2-15 IPC summary for vildagliptin 100mg MR containing 40% HPMC with roller compaction (400mg tablet weight, 11mm FFBE)

Sample	Thickness (mm) IPC: 2.8-4.2mm	Hardness (Kp)	%Friability (100 drops)
TRD-1926-78 1,000 lb/in RC (FP)	3.95 - 3.72	40 rpm: 11.5 - 13.2 60 rpm: 9.76 - 12.69 80 rpm: 9.76 - 12.84	40 rpm: < 0.5 60 rpm: < 0.5 80 rpm: < 0.5
TRD-1771-123A 1,000 lb/in RC (FP)	4.07 – 3.68	40 rpm: 8 – 18.3	40 rpm: < 0.5
TRD-1926-01A 500 lb/in RC (FP)	4.0 – 3.81	40 rpm: 9.3 -14.8	40 rpm: 0.1
TRD-1926-01B 1,500 lb/in RC (FP)	4.02 – 3.66	40 rpm: 7.6 – 19.9	40 rpm: 0.1
TRD-1771-31 Not RC	4.13 – 3.94	40 rpm: 5.1 – 7.75	40 rpm: 6.3, capping

Table 2-16 IPC summary for vildagliptin 150mg MR containing 40% HPMC with roller compaction (600mg tablet weight, 17 x 6.7mm ovaloid)

Sample	Thickness (mm) IPC: 2.8-4.2mm	Hardness (Kp)	%Friability (100 drops)
TRD-1926-78 1,000 lb/in RC (FP)	40 rpm: 6.5 – 6.08 60 rpm: 6.4 – 6.03 80 rpm: 6.6 – 6.07	40 rpm: 17.5 – 25.9 60 rpm: 15.5 – 24.8 80 rpm: 11.2 – 24.4	40 rpm: < 0.2 60 rpm: < 0.2 80 rpm: < 0.2
TRD-1771-123A 1,000 lb/in RC (FP)	6.4 – 5.84	40 rpm: 12.3 – 30.4	40 rpm: < 0.25
TRD-1926-01A 500 lb/in RC (FP)	6.56 – 5.99	40 rpm: 14.6 – 30.3	40 rpm: 0.5, 1 tablet capped at 20Kn
TRD-1926-01B 1,500 lb/in RC (FP)	6.5 – 5.84	40 rpm: 13.9 – 32	40 rpm: 0.25
TRD-1771-33 Not RC	6.59 – 5.97	40 rpm: 18.3 – 27.2	40 rpm: 0.6, lamination at 20Kn

Roller compaction improved bulk powder flow and compression profile without the use of pre-compression. The acceptable hardness range was 10 to 13Kp and 11 to 25Kp for the 100mg and 150mg tablets, respectively. Lamination was observed at compression forces greater than 18 KN for the 150mg tablet. The formulation does not appear to be dwell time sensitive. The lab phase FMI activities will start pending the confirmation of the FMI tablet shape and size for the 100mg strength.

II. Analytical Investigations for Stress Tests of Variants

A. *Elaboration of Analytical Methods*
Assay and Related Substances

The assay of vildagliptin and the quantitation of degradation products of vildagliptin were performed by a gradient HPLC method [AM54001B(AS6105)]. The sample was extracted with methanol and diluted to target concentration using 10:90 methanol/acetonitrile and then chromatographed using a reverse phase HPLC method with UV detection at 210 nm.

Drug Release

Dissolution of vildagliptin from six 100mg vildagliptin tablets was determined under the test conditions elaborated in Table 3-1

Samples obtained were analyzed by using an isocratic reverse phase HPLC method [AM50161A(AS6105)] with UV detection at 2010 nm. Same experimental have been done for the 150mg and 200mg tablets.

B. Analytical Results

Stability Protocol

The technical and clinical batches slow and fast formulations of CSF were put on stability and the stability was monitored up to 12 months. The stability of technical batches of MF variants, 100mg and 150mg is currently ongoing as shown in the protocol, Table 3-2.

Table 3-2 Drug product stability – storage conditions and testing intervals batch TRD-1926-078A(100mg), and TRD-1926-078B (150mg); [packaging: HDPE, CR cap, 1g desiccant].

Storage condition	Time (months)
5°C	[24]
25°C/60%RH	3, 6, 9, 12, 24
30°C/65%RH	[3], [6], [9], [12]
40°C/75%RH	1.5, 3, 6

[ ] Optional testing

Stability Results

Table 3-3 Summary of stability for vildagliptin 100 mg tablets, TRD-0739-0113

					Degradation products						
Conditions	**Time**	**Appearance**	**%Assay**	**% Amide**	**%Cyclic**	**%Diket.**	**%Unk. (RRT = 0.26)**	**%Unk. (RRT = 0.77)**	**%Unk. (RRT = 1.10)**	**%Total deg.**	**% Moisture**
	Initial	Complies	100.8	<LOQ	<LOD	ND	ND	<LOQ	<LOQ	-	3.52
25°C/60%RH	3M	Complies	102.6	<LOQ	<LOQ	<LOQ	ND	<LOQ	<LOQ	-	2.40
	6M	Complies	101.0	0.06	0.05	<LOQ	ND	<LOQ	<LOQ	0.11	2.66
	9M	Complies	100.1	0.08	0.05	<LOQ	ND	<LOQ	<LOQ	0.13	2.74
	12M	Complies	102.5	0.11	0.05	<LOQ	ND	<LOQ	<LOQ	0.16	3.16
30°C/60%RH	3M	Complies	102.3	0.06	<LOQ	<LOQ	ND	<LOQ	<LOQ	0.06	2.51
	6M	Complies	104.3	0.12	0.05	<LOQ	ND	<LOQ	<LOQ	0.17	2.79
	9M	Complies	100.3	0.17	0.05	<LOQ	ND	<LOQ	<LOQ	0.22	3.07
	12M	Complies	100.7	0.21	0.05	0.07	ND	<LOQ	<LOQ	0.33	3.59
40°C/75%RH	6W	Complies	101.4	0.10	0.05	<LOQ	ND	<LOQ	<LOQ	0.15	2.73
	3M	Complies	103.3	0.20	0.05	0.07	ND	<LOQ	<LOQ	0.32	2.90
	6M	Complies	100.7	0.39	0.05	0.12	<LOQ	<LOQ	<LOQ	0.56	3.35

NT = not tested, Cyclic = cyclic amidine, Diketo = diketopiperazine, LOQ = 0.05% for cyclic, diketo and amide, 0.06% for vildagliptin; LOD = 0.02%; ND = Not Detected

Table 3-4 Summary of dissolution data for vildagliptin 100mg tablets, TRD-0739-0113

		Percent released									
Conditions	**Time**	**0.5 hr**	**1 hr**	**2 hr**	**4 hr**	**6 hr**	**8 hr**	**10 hr**	**12 hr**	**18 hr**	**24 hr**
	Initial	23	36	56	82	95	102	104	104	104	102
25°C/60%RH	3M	NT									
	6M	NT									
	9M	NT									
	12M	NT									
30°C/60%RH	3M	NT									
	6M	NT									
	9M	NT									
	12M	24		57		94				100	
40°C/75%RH	6W	NT									
	3M	23	38	57	84	98	103	103	104	102	101
	6M	24	-	57	-	95	-	-	-	100	-

NT = Not tested

Note: Dissolution was tested only at accelerated conditions as these represent worst case situation.

Table 3-5 Summary of stability for vildagliptin 100mg tablets, TRD-0739-0121

Conditions	Time	Appearance	%Assay	Degradation products							
				% Amide	%Cyclic	%Diket.	%Unk. (RRT= 0.26)	%Unk. (RRT= 0.77)	%Unk. (RRT= 1.10)	%Total	% Moisture
	Initial	complies	103.1	<LOQ	<LOD	ND	ND	<LOQ	0.07	0.07	3.19
25°C/60%RH	3M	Complies	103.3	<LOQ	<LOQ	<LOQ	ND	<LOQ	<LOQ	0.00	2.53
	6M	Complies	104.3	0.05	<LOQ	<LOD	ND	<LOQ	<LOQ	0.05	2.70
	9M	Complies	102.2	0.07	<LOQ	<LOQ	ND	<LOQ	<LOQ	0.07	3.03
	12M	Complies	101.4	0.08	<LOQ	<LOQ	ND	<LOQ	<LOQ	0.08	3.31
30°C/60%RH	3M	Complies	104.7	0.05	<LOQ	<LOQ	ND	<LOQ	<LOQ	0.05	2.61
	6M	Complies	103.9	0.09	<LOQ	<LOQ	ND	<LOQ	<LOQ	0.09	2.88
	9M	Complies	100.8	0.11	<LOQ	<LOQ	ND	<LOQ	<LOQ	0.11	3.33
	12M	Complies	102.4	0.14	<LOQ	0.05	ND	<LOQ	<LOQ	0.19	3.65
40°C/75%RH	6W	Complies	102.1	0.08	<LOQ	<LOQ	ND	<LOQ	<LOQ	0.08	2.81
	3M	Complies	104.8	0.13	<LOQ	0.05	ND	<LOQ	<LOQ	0.18	2.97
	6M	Complies	102.9	0.25	<LOQ	0.08	<LOQ	<LOQ	<LOQ	0.33	3.52

NT = not tested, Cyclic = cyclic amidine, Diketo = diketopiperazine, LOQ = 0.05% for cyclic, diketo and amide, 0.06% for vildagliptin; LOD = 0.02%; ND = Not Detected

Table 3-6 Summary of dissolution data for vildagliptin 100mg tablets, TRD-0739-0121

Conditions	Time	Percent released									
		0.5 hr	1 hr	2 hr	4 hr	6 hr	8 hr	10 hr	12 hr	18 hr	24 hr
	Initial	12	20	30	45	56	65	72	78	89	95
25°C/60%RH	3M	NT									
	6M	NT									
	9M	NT									
	12M	NT									
30°C/60%RH	3M	NT									
	6M	NT									
	9M	NT									
	12M	13	-	32	-	60	-	-	-	94	
40°C/75%RH	6W	NT									
	3M	13	21	32	49	59	70	77	83	93	98
	6M	13	-	33	-	61	-	-	-	95	-

NT = Not tested

Note: Dissolution was tested only at accelerated conditions as these represent worst case situation.

Table 3-7 References for the stability data in Table 3-3 through Table 3-6

	Assay & Purity	% Moisture Content	Dissolution
Initial	TRD0958/057 (XG)	TRD0958/109 (XG)	WS18431
6W	TRD0958/057 (XG)	TRD0958/109 (XG)	NA
3M	TRD0983/082 (XG)	TRD0978/110 (MZ)	WS20247
6M	TRD1514/110 (XG)	TRD1514/110 (XG)	WS21349
9M	TRD1645/129 (MZ)	TRD1645/101 (MZ)	NA
12M	TRD1642/069 (JT)	TRD1642/145 (JT)	WS22591

Table 3-8 Summary of stability for vildagliptin 100mg tablets, AEUS/2002-0080

							Degradation products				
Conditions	**Time**	**Appearance**	**%Assay**	**% Amide**	**%Cyclic**	**%Diket.**	**%Unk. (RRT = 0.26)**	**%Unk. (RRT = 0.77)**	**%Unk. (RRT = 1.10)**	**% Total Deg.**	**% Moisture**
Initial	0	complies	102.0	ND	ND	ND	-			-	2.62
5°C	3M	complies	104.3	0.02	0.02	ND	ND	0.04	0.03	0.00	2.17
	6M	complies	101.8	0.02	0.02	ND	ND	0.05	0.03	0.00	2.32
	9M	complies	102.3	0.02	0.03	ND	ND	0.04	ND	0.00	2.39
	12M	NT									
25°C/60%RH	3M	complies	101.1	0.04	0.05	ND	ND	0.04	0.03	0.05	2.06
	6M	complies	103.0	0.07	0.06	0.02	ND	0.04	0.04	0.13	2.34
	9M	complies	100.6	0.08	0.06	0.02	ND	0.04	ND	0.14	2.43
	12M	complies	101.9	0.09	0.05	0.02	ND	0.03	0.02	0.14	2.72
25°C/60%RH	3M	complies	101.4	0.04	0.05	ND	ND	0.04	0.03	0.05	2.22
(With Cotton)	6M	complies	102.0	0.06	0.06	ND	ND	0.04	0.05	0.12	2.67
	9M	complies	102.2	0.09	0.07	ND	ND	0.04	ND	0.16	2.69
	12M	complies	101.7	0.11	0.07	0.03	0.02	0.04	0.02	0.18	2.85

						Degradation products					
°C/60%RH	3M	complies	102.6	0.06	0.06	ND	ND	0.04	0.04	0.12	2.12
	6M	complies	102.7	0.11	0.06	0.03	ND	0.04	0.05	0.17	2.60
	9M	complies	101.7	0.15	0.07	0.03	ND	0.03	ND	0.22	2.78
	12M	complies	101.3	0.19	0.06	0.05	ND	0.03	0.02	0.30	2.80
40°C/75%RH	6W	complies	101.7	0.10	0.06	0.03	ND	0.04	0.06	0.22	2.28
	3M	complies	101.9	0.20	0.07	0.05	ND	0.03	0.05	0.32	2.59
	6M	complies	102.5	0.40	0.07	0.11	0.03	0.04	0.06	0.64	3.41

NT = not tested, Cyclic = cyclic amidine, Diketo = diketopiperazine, LOQ = 0.05% for cyclic, diketo and amide, 0.06% for vildagliptin; LOD = 0.02%; ND = Not detected

Table 3-9 Summary of dissolution data for vildagliptin 100mg tablets, AEUS/2002-0080

		Percent released									
Conditions	**Time**	**0.5 hr**	**1 hr**	**2 hr**	**4 hr**	**6 hr**	**8 hr**	**10 hr**	**12 hr**	**18 hr**	**24 hr**
	Initial	24	37	55	79	92	99	101	101	100	99
25°C/60%RH	3M	NT									
	6M	NT									
	9M	NT									
	12M	NT									
25°C/60%RH	3M	NT									
(With cotton)	6M	NT									
	9M	NT									
30°C/60%RH	3M	NT									
	6M	NT									
	9M	22		53		90				102	
	12M	NT									
40°C/75%RH	6W	NT									
	3M	23		54		92				101	
	6M	24		56		93				104	

NT = Not tested

Note: Dissolution was tested only at accelerated conditions as these represent worst case situation.

Table 3-10 Summary of stability data for vildagliptin 100mg tablets, AEUS/2002-0081

						Degradation products				
Conditions	**Time**	**Appearance**	**%Assay**	**% Amide**	**%Cyclic**	**%Diket**	**%Unk. (RRT = 0.26)**	**%Unk. (RRT = 0.77)**	**%Unk. (RRT = 1.10)**	**% Total deg.**	**% Moisture**
Initial	0	complies	102.5	ND	ND	ND	-	-		-	2.45
5°C	3M	complies	103.4	0.02	0.01	ND	ND	0.04	0.04	0.00	2.35
	6M	complies	102.7	0.02	0.02	ND	ND	0.04	0.05	0.00	2.19
	9M	complies	101.4	0.02	ND	ND	ND	0.04	0.02	0.00	2.24
	12M	NT									
25°C/60%RH	3M	complies	103.1	0.03	0.03	ND	ND	0.04	0.05	0.00	2.01
	6M	complies	102.9	0.05	0.03	ND	ND	0.05	0.07	0.12	2.33
	9M	complies	100.8	0.06	0.03	ND	ND	0.04	ND	0.06	2.44
	12M	complies	101.7	0.07	0.03	0.02	0.02	0.04	0.04	0.07	2.57
25°C/60%RH	3M	complies	101.8	0.03	0.03	ND	ND	0.04	0.04	0.00	2.14
(With Cotton)	6M	complies	104.1	0.05	0.04	ND	ND	0.04	0.06	0.11	2.61
	9M	complies	102.9	0.06	0.04	ND	ND	0.04	ND	0.06	2.58
	12M	complies	104.9	0.08	0.03	0.02	ND	0.03	0.03	0.08	2.57
30°C/60%RH	3M	complies	101.0	0.05	0.03	0.01	ND	0.04	0.05	0.05	2.06
	6M	complies	103.4	0.08	0.04	0.03	ND	0.04	0.07	0.15	2.64
	9M	complies	102.3	0.10	0.04	0.03	ND	0.03	ND	0.10	2.75
	12M	complies	101.0	0.13	0.04	0.04	0.02	0.04	0.03	0.13	2.62
40°C/75%RH	6W	complies	102.8	0.07	0.03	0.03	ND	0.04	0.07	0.14	2.24
	3M	complies	102.6	0.13	0.04	0.03	ND	0.04	0.05	0.13	2.63
	6M	complies	103.1	0.25	0.05	0.08	0.02	0.04	0.07	0.45	3.58

NT = not tested, Cyclic = cyclic amidine, Diketo = diketopiperazine, LOQ = 0.05% for cyclic, diketo and amide, 0.06% for vildagliptin; LOD = 0.02%; ND = Not detected

Table 3-11 Summary of dissolution data for vildagliptin 100mg tablets, AEUS/2002-0081, 30HDPE90

		Percent released									
Conditions	**Time**	**0.5 hr**	**1 hr**	**2 hr**	**4 hr**	**6 hr**	**8 hr**	**10 hr**	**12 hr**	**18 hr**	**24 hr**
	Initial	13	21	33	49	61	70	77	83	94	98
25°C/60%RH	3M	NT									
	6M	NT									
	9M	NT									
	12M	NT									
25°C/60%RH	3M	NT									
(With cotton)	6M	NT									
	9M	NT									
30°C/60%RH	3M	NT									
	6M	NT									
	9M	13		33		62				95	
	12M	NT									
40°C/75%RH	6W	NT									
	3M	12		31		57				89	
	6M	12		30		56				89	

NT = Not tested Note: Dissolution was tested only at accelerated conditions as these represent worst case situation.

Table 3-12 Summary of stability data for vildagliptin 200mg tablets, AEUS/2002-0082

							Degradation products				
Conditions	**Time**	**Appearance**	**%Assay**	**% Amide**	**%Cyclic**	**%Diket.**	**%Unkn (RRT = 0.26)**	**%Unk (RRT = 0.77)**	**%Unk (RRT = 1.10)**	**%Total**	**% Moisture**
Initial	0	complies	100.6	ND	ND	ND	-			-	2.61
5°C	3M	complies	100.7	0.02	0.02	ND	ND	0.04	0.02	0.00	2.23
	6M	complies	100.3	0.02	0.02	ND	ND	0.04	0.03	0.00	2.54
	9M	complies	99.9	0.02	0.02	ND	ND	0.04	ND	0.00	2.46
	12M	NT									
25°C/60%RH	3M	complies	100.6	0.04	0.04	ND	ND	0.04	0.02	0.00	2.16
	6M	complies	100.1	0.06	0.05	ND	ND	0.04	0.04	0.11	2.50
	9M	complies	99.9	0.08	0.06	0.02	ND	0.03	ND	0.14	2.56
	12M	complies	99.9	0.09	0.05	0.03	0.02	0.03	0.02	0.14	2.53
25°C/60%RH	3M	complies	100.8	0.04	0.05	ND	ND	0.04	0.02	0.05	2.22
(With Cotton)	6M	complies	99.3	0.06	0.05	ND	ND	0.04	0.04	0.11	2.59
	9M	complies	99.1	0.08	0.05	0.02	ND	0.03	ND	0.14	2.64
	12M	complies	100.3	0.10	0.06	0.03	0.02	0.03	0.02	0.16	2.63
30°C/60%RH	3M	complies	99.5	0.06	0.06	ND	ND	0.04	0.03	0.12	2.18
	6M	complies	100.8	0.11	0.05	0.03	ND	0.04	0.04	0.16	2.59
	9M	complies	100.4	0.16	0.06	0.03	ND	0.03	ND	0.22	2.75
	12M	complies	99.2	0.19	0.05	0.05	0.02	0.03	0.02	0.29	2.77
40°C/75%RH	6W	complies	101.2	0.11	0.05	0.03	ND	0.03	0.06	0.22	2.31
	3M	complies	99.3	0.20	0.05	0.05	ND	0.03	0.03	0.30	2.51
	6M	complies	99.5	0.36	0.05	0.10	0.02	0.04	0.05	0.51	3.16

NT = not tested, Cyclic = cyclic amidine, Diketo = diketopiperazine, LOQ = 0.05% for cyclic, diketo and amide, 0.06% for vildagliptin; LOD = 0.02%; ND = Not detected

Table 3-13 Summary of dissolution data for 200mg tablets, AEUS/2002-0082

Conditions	Time	Percent released									
		0.5 hr	1 hr	2 hr	4 hr	6 hr	8 hr	10 hr	12 hr	18 hr	24 hr
	Initial	24	36	52	73	86	93	98	99	99	97
5°C	3M	NT									
	6M	NT									
	9M	NT									
	12M	NT									
25°C/60%RH	3M	NT									
	6M	NT									
	9M	NT									
	12M	NT									
25°C/60%RH (With cotton)	3M	NT									
	6M	NT									
	9M	NT									
30°C/60%RH	3M	NT									
	6M	NT									
	9M	23		51		84				99	
	12M	NT									
40°C/75%RH	6W	NT									
	3M	24		52		87				99	
	6M	25		54		87				99	

NT = Not tested
Note: Dissolution was tested only at accelerated conditions as these represent worst case situation.

Table 3-14 Summary of dissolution data for 200mg tablets, AEUS/2002-0083

							Degradation products				
Conditions	**Time**	**Appearance**	**%Assay**	**% Amide**	**%Cyclic**	**%Diket.**	**%Unkn (RRT = 0.26)**	**%Unkn (RRT = 0.77)**	**%Unkn (RRT = 1.10)**	**%Total**	**% Moisture**
Initial	0	complies	100.8	ND	ND	ND	-			-	2.36
5°C	3M	complies	100.8	0.02	ND	ND	ND	0.04	0.03	0.00	2.06
	6M	complies	100.5	ND	ND	ND	ND	0.04	0.04	0.00	2.41
	9M	complies	101.1	ND	ND	ND	ND	0.04	0.03	0.00	2.36
	12M	NT									
25°C/60%RH	3M	complies	99.9	0.03	0.02	ND	ND	0.04	0.03	0.00	2.04
	6M	complies	100.6	0.05	0.03	ND	ND	0.04	0.05	0.05	2.32
	9M	complies	99.6	0.06	0.03	ND	ND	0.03	0.02	0.06	2.49
	12M	complies	100.2	0.08	0.02	0.02	0.02	0.04	0.04	0.08	2.50
25°C/60%RH	3M	complies	100.5	0.03	0.03	ND	ND	0.04	0.04	0.05	2.09
(With Cotton)	6M	complies	100.2	0.05	0.03	ND	ND	0.04	0.05	0.05	2.50
	9M	complies	100.1	0.07	0.03	ND	ND	0.03	ND	0.07	2.66
	12M	complies	100.0	0.08	0.03	0.02	ND	0.04	0.03	0.08	2.54
30°C/60%RH	3M	complies	100.4	0.05	0.03	ND	ND	0.04	0.04	0.05	2.07
	6M	complies	100.3	0.08	0.03	0.02	ND	0.04	0.08	0.16	2.52
	9M	complies	100.2	0.11	0.03	0.03	ND	0.04	ND	0.11	2.67
	12M	complies	100.3	0.13	0.02	0.05	0.02	0.04	0.03	0.18	2.71
40°C/75%RH	6W	complies	101.2	0.08	0.03	0.03	ND	0.03	0.08	0.16	2.24
	3M	complies	100.7	0.14	0.03	0.03	ND	0.03	0.04	0.14	2.47
	6M	complies	99.8	0.23	0.03	0.07	0.02	0.04	0.06	0.36	3.14

NT = not tested, Cyclic = cyclic amidine, Diketo = diketopiperazine, LOQ = 0.05% for cyclic, diketo and amide, 0.06% for vildagliptin; LOD = 0.02%; ND = Not detected

Table 3-15 **Summary of dissolution data for 200mg tablets, AEUS/2002-0083**

		Percent released									
Conditions	**Time**	**0.5 hr**	**1 hr**	**2 hr**	**4 hr**	**6 hr**	**8 hr**	**10 hr**	**12 hr**	**18 hr**	**24 hr**
	Initial	11	18	27	40	50	59	66	71	84	91
5°C	3M	NT									
	6M	NT									
	9M	NT									
	12M	NT									
25°C/60%RH	3M	NT									
	6M	NT									
	9M	NT									
	12M	NT									
25°C/60%RH (With cotton)	3M	NT									
	6M	NT									
	9M	NT									
30°C/60%RH	3M	NT									
	6M	NT									
	9M	11		27		51				84	
	12M	10		25		48				81	
40°C/75%RH	6W	NT									
	3M	10		25		49				81	
	6M	10		25		48				81	

NT = Not tested

Note: Dissolution was tested only at accelerated conditions as these represent worst case situation.

Table 3-16 References for the stability data in Table 3-8 through Table 3-15

Reference	Assay & Purity	Moisture Content	Dissolution
Initial	WS17548	TRD0958/109XG	WS19437,19427,19449,19455
6W	WS20721	TRD0983/99XG	NA
3M	WS20768	WS20768	WS21307, 20772, 21321
6M	WS21855	WS21855	WS22761, 22771, 22775
9M	WS23831	WS23831	WS22591, 22599
12M	WS25593	WS25593	WS25471, 25507

Table 3-17 Summary of stability data for 100mg tablets, TRD-1926-078A (packaging: HDPE, 1g desiccant and CR cap)

					Degradation products						
Conditions	**Time**	**Appearance**	**%Assay**	**% Amide**	**%Cyclic**	**%Diket.**	**%Unk. (RRT = 0.26)**	**%Unk. (RRT = 0.77)**	**%Unk. (RRT = 1.10)**	**%Total deg.**	**% Moisture**
	Initial	Complies	102.0	<LOQ	<LOQ	ND	ND	<LOQ	<LOQ	<LOQ	2.39
40°C/75%RH	6W	Complies	98.3	0.10	0.06	<LOQ	<LOQ	<LOQ	0.06	0.22	

NT = not tested, Cyclic = cyclic amidine, Diketo = diketopiperazine, LOQ = 0.05% for cyclic, diketo and amide, 0.06% for vildagliptin; LOD = 0.02%; ND = Not Detected.

Ref: Initial – TRD 1928/127 JT: 6W – TRD 1980/

Table 3-18 Summary of dissolution data for 100mg tablets, TRD-1926-078A (packaging: HDPE, 1g desiccant and CR cap)

		Percent released									
Conditions	**Time**	**0.25 hr**	**0.5hr**	**1 hr**	**2 hr**	**4 hr**	**6 hr**	**8 hr**	**10 hr**	**12 hr**	**18 hr**
	Initial	7.6	12.3	19.5	30.5	46.5	-	68.4	82.3	89.1	93.8
40°C/75%RH	6W Not tested										

Ref: WS35331 KE

5

Table 3-19 Summary of stability data for 150mg tablets, TRD-1926-078B (packaging: HDPE, 1g desiccant and CR cap)

				Degradation products							
Conditions	**Time**	**Appearance**	**%Assay**	**% Amide**	**%Cyclic**	**%Diket.**	**%Unk. (RRT = 0.26)**	**%Unk. (RRT = 0.77)**	**%Unk. (RRT = 1.10)**	**%Total deg.**	**% Moisture**
	Initial	Complies	101.7	<LOQ	<LOQ	ND	ND	<LOQ	<LOQ	<LOQ	2.31
40°C/75%RH	6W	Complies	99.1	0.10	0.07	<LOQ	ND	<LOQ	0.06	0.23	

NT = not tested, Cyclic = cyclic amidine, Diketo = diketopiperazine, LOQ = 0.05% for cyclic, diketo and amide, 0.06% for vildagliptin; LOD = 0.02%; ND = Not Detected

Ref: Initial – TRD1928/127 JT: 6W – TRD

Table 3-20 Summary of dissolution data for 150mg tablets, TRD-1926-078B(packaging: HDPE, 1g desiccant and CR cap)

		Percent released									
Conditions	**Time**	**0.25hr**	**0.5 hr**	**1 hr**	**2 hr**	**4 hr**	**6 hr**	**8 hr**	**10 hr**	**12 hr**	**18 hr**
	Initial	6.8	11.1	17.7	28.3	-	43.9	65.1	79.5	86.5	91.5
40°C/75%RH	6W Not tested										

Ref: WS35331 KE

Table 3-21 Summary of assay and related substances data for the initial analysis of 1-[(3-Hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile MR tablets (technical batches)

				Degradation products						
Batch no.	**Appearance**	**%Assay**	**% Amide**	**%Cyclic**	**%Diket.**	**%Unk. (RRT = 0.26)**	**%Unk. (RRT = 0.77)**	**%Unk. (RRT = 1.10)**	**%Total deg.**	**% Moisture**
TRD-1971-053C 100mg	Complies	100.9	ND	ND	ND	ND	<LOQ	ND	<LOQ	2.44
TRD-1971-055A 150mg	Complies	120.4	ND	ND	ND	ND	<LOQ	ND	<LOQ	2.31

NT = not tested, Cyclic = cyclic amidine, Diketo = diketopiperazine, LOQ = 0.05% for cyclic, diketo and amide, 0.06% for vildagliptin; LOD = 0.02%; ND = Not Detected
Ref: TRD 1980/096 JT

Table 3-22 Summary of dissolution data for the initial analysis of vildagliptin MR tablets (technical batches)

	Percent released									
Batch no.	**0.25hr**	**0.5 hr**	**1 hr**	**2 hr**	**4 hr**	**6 hr**	**8 hr**	**10 hr**	**12 hr**	**18 hr**
TRD-1971-053C 100mg										
TRD-1971-055A 150mg										
Ref:										

Specifications

Tentative analytical specifications were laid down in specifications during early development of vildagliptin 25mg, 50mg and 100mg tablets [AS6105B].

III. Conclusions

A Clinical Service Form (CSF) was developed to provide a "slow" release profile containing 40% HPMC K100M as the controlled release polymer. A direct compression powder blend was developed for the 100mg and 150mg strengths. It was determined that the use of roller compaction improved the compression profile and bulk powder flow of the CSF. Roller compaction exhibits a dissolution profile similar to the CSF (which was tested in vivo in humans). The formulation does not appear to be effected by roller compaction roll speed, or dwell time on compression.

For the CSF 100mg, 150mg and 200mg strengths, the stability data are available up to 12 months at 25°C/60%RH and 30°60%RH. The appearance, assay, purity and dissolution data were found to within specification limits after 12 months at 30°C/60%RH. These data support a 24 month re-test period with specification "do not store above 25°C" for 100mg, 150mg and 200mg modified release tablets. The composition of vildagliptin 150mg modified release tablets is bracketed by 100mg and 200mg strengths. Therefore, the stability of 150mg tablets is expected to be same as 100mg and 200mg strengths. Thus a re-test period similar to that of 100 and 200mg strengths is applied for 150mg tablets. The stability profile of 100mg and 150mg tablets manufactured by roller compaction after 6W, 40°C/75%RH was found to be comparable to that of CSF formulations. Thus, the stability does not appear to be affected by the inclusion of roller compaction step in the manufacturing process.

The roller compacted formulation containing 40% or 30% HPMC K100M has been selected.

Preferred formulations are described in the below table:

	100		150	
Component	mg/ unit	%	mg/ unit	%
Vildagliptin	100	25.0	150	25.0
Microcrystalline cellulose, PH102	120	30.0	180	30.0
Lactose, anhydrous DT	16	4.0	24	4.0
Hydroxypropyl-methylcelluose K100M	160	40.0	240	40.0
Magnesium stearate	4	1.0	6	1.0
Total tablet weight (mg)	400		600	
KN number	3768652		6001732	
Tablet diameter, shape	11 mm, FFBE		17x6.7mm, ovaloid	

The invention has been described in an illustrative manner, and it is to be understood that the terminology which has been used is intended to be in the nature of words of description rather than of limitation.

Obviously, many modifications and variations of the present invention are possible in light of the above teachings. It is, therefore, to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described.

Example A : Mechanical stress (particle size distribution)

The material in the desired particle size range can be produced from any form of LAF237 e.g. amorphous form or crystal form, by mechanical stress. This stress can be mediated by impact, shear or compression. In most commercially available grinding equipment a combination of these principles occurs. For LAF237 preferably a mechanical impact or jet mill is used. The most preferable mechanical impact mill can be equipped with different kind of beaters, screens, liners or with pin plates. For our process preferably an impact mill with plate beater and a slit screen 5 * 2.5 cm is used. The impact speed should be variable between 20 and 100 m/s (as peripheral speed) to adapt to any

batch to batch variation. In our case a peripheral speed of the beater of about 40 - 50 m/s is used.

Pharmacokinetic and Pharmacodynamic study:

AUC	area under the concentration time curve
AUC_{0-t}	The area under the plasma concentration-time curve from time zero to the last quantifiable data point t [ng*hr/mL]
AUC_{0-inf} or $AUC_{(0-\infty)}$	The area under the plasma concentration-time curve from time zero to infinity [ng*hr/mL]
BAPK	Bioanalytics and Pharmacokinetics section
C_{max}	maximum plasma concentration
CRF	case report / record form
CRO	Clinical Research Organization
CV	Coefficient of variation
ECG	Electrocardiogram
DPP-4	dipeptidyl peptidase 4; dipeptidyl peptidase IV
FMI	Final Market Image
GLP-1	glucagon-like peptide 1
ICH	International Council on Harmonization
IRB	Institutional Review Board
LAF237	Vildagliptin
LC-MS/MS	Liquid chromatography-mass spectrometry/mass spectrometry
LOQ	limit of quantitation
o.d.	once a day
PD	Pharmacodynamics
PK	Pharmacokinetics
p.o	per os / by mouth / orally
QC	Quality Control
SOP	Standard Operating Procedures
SD	standard deviation
t_{max}	time to reach C_{max}
$t_{½}$	elimination half-life
Vd/f	volume of distribution, corrected for the absolute bioavailability

Study synopsis and assessment schedule

Title of study: A randomized, open-label, three period, cross-over study to evaluate the pharmacokinetics, pharmacodynamics, safety, and tolerability, of two new LAF237 100mg and 150mg modified release formulations in patients with type 2 diabetes.

Objectives:

Primary objective

- To evaluate the pharmacokinetic profiles (single and multiple dose) after oral administration of two dosage strengths (100mg and 150mg) of a modified release formulation of LAF237 in patients with type 2 diabetes.
- To evaluate the effect of a modified release formulation of LAF237 at two dosage strengths (100mg and 150mg) (single and multiple dose) on DPP-IV activity.

- To evaluate the safety and tolerability of the modified release formulations after oral administration for 7 days to patients with type 2 diabetes.

Secondary objective(s)

- To evaluate the effect of a modified release formulation of LAF237 at two dosage strengths (100mg and 150mg) on the glucose profile after 7 days of treatment.
- To compare the pharmacokinetic profiles of the modified release formulations of LAF237 at two dosage strengths when given with and without an ADA breakfast.

Design: This study is a single center, randomized, open-label, three-period, crossover design study. A total of 27 patients with type 2 diabetes (both male and female) are enrolled. Each subject participate in a 28 day screening period, three base-line periods, three 11-day treatment periods separated by two 7-day washout periods and a study completion evaluation.

Subjects are randomized to the following 3 treatment sequence groups under William's design.

Sequence	Period 1	Period 2	Period 3
I	A	B	C
II	B	C	A
III	C	A	B

- Treatment A:
  - Day 1: 100mg LAF237 MR OD, single dose
  - Days 3-10: 100mg LAF237 MR OD, for 8 days
  - Day 11-17: Wash out
- Treatment B:
  - Day 1: 150mg LAF237 MR OD, single dose
  - Days 3-10: 150mg LAF237 MR OD, for 8 days
  - Day 11-17: Wash out
- Treatment C:
  - Day 1: 50mg LAF237 BID, two doses
  - Days 3-10: 50mg LAF237 BID, for 8 days
  - Day 11-17: Wash out

Patients receiving metformin therapy are required to stop taking metformin for at least 21 days prior to dosing on Day 1. On the evening of Day -1, patients who satisfy all inclusion/exclusion criteria at screening (Days -28 to -2) enter the study center for verification of inclusion/exclusion criteria, baseline evaluations and subsequent

randomization to one of three sequences. Patients remain domiciled until the morning of Day 3 (48 h after the first dose of LAF237 is given).

On Day 1 of each treatment period, patients receive the first dose of LAF237 following an over night fast of 10-12 hours. They continue to fast for 5 hours post-dose. Patients are discharged from the site after the Day 3 morning dose of LAF237 with study drug medication, instructions on how to administer drug at home during the out-patient treatment days (Days 3 to 7) and a dosing diary.

The modified release formulation of LAF237 (100mg and 150mg) are administered once daily before breakfast and the 50mg LAF237 is administered twice daily, once before breakfast and once before dinner. A dosing diary is kept by the patients for all non-domiciled doses.

Patients return to the site on the morning of Day 5 for pre-dose pharmacokinetic sample collection. They are admitted to the site at least 10 h prior to the Day 8 morning dose of LAF237 and remain domiciled until the morning of Day 12 (48 h after the Day 10 dose is given). On Day 9, patients are served an ADA breakfast within 30 minutes of the morning dose of LAF237.

Meal times on all domiciled days are strictly enforced. Breakfast are served within 30 minutes of the morning dose, except on days 1 and 10 when patients fast for 5 hours after dosing. Lunch, dinner and a snack are served 5, 10 and 12 hours post-dose, respectively.

<table>
<tr><td>Screening</td><td colspan="6">Periods One, Two, and Three</td></tr>
<tr><td>Days -28 to -2</td><td>Day -1</td><td>Days 1-2</td><td>Days 3-8</td><td>Day 9</td><td>Day 10-11</td><td>Days 12-17</td></tr>
<tr><td rowspan="4">Patients receiving metformin required to stop therapy on Day -21</td><td rowspan="4">B/L</td><td>48 hr Sampling under fasted conditions</td><td>Day 3: Start of multiple dosing</td><td>24 hr Sampling, ADA Breakfast served</td><td>48 hr Sampling under fasted conditions</td><td rowspan="4">w/o
(Not required after Period 3)</td></tr>
<tr><td>PK profile</td><td>Days 5 and 8: Pre-dose PK sampling</td><td>PK profiling</td><td>PK profiling</td></tr>
<tr><td rowspan="2">DPP-IV profile</td><td rowspan="2">Days 3-7: Out patient Day 8: Domiciled</td><td>DPP-IV profiling</td><td rowspan="2">DPP-IV profiling</td></tr>
<tr><td>Glucose profiling</td></tr>
</table>

Number of subjects: 27 patients with type 2 diabetes (male and female)

Criteria for inclusion:

- Male and female (postmenopausal, surgically sterile or using double-barrier method of contraception) type two diabetic patients between 20 and 65 years of age.
- Hb1Ac range between 6.5 and 9%.
- Body mass index between 22 – 40 kg/m2.
- Patients must be drug-naive to diabetic medications or on metformin treatment.
- Patients must be able to complete a 3 week wash-out of current metformin therapy. Patients will monitor FPG at home and call PI if values exceed 250mg/dL.
- Patients must not be taking prescription or non-prescription medications.
- Patients must have a fasting blood glucose of 7.0 – 12.2 mmol/dL (100-240 mg/dL) at screening and at predose on Day 1.
- Patients on diuretics or cyclic hormone therapy must be on stable dose (at least 3 months prior to screening) and the maintenance dose may not be adjusted during the study.
- All subjects must be able to provide written informed consent prior to participation the study.

Criteria for exclusion:

- History of type 1 diabetes mellitus, diabetes that is a result of pancreatic injury, or secondary forms of diabetes e.g. Cushing's syndrome and acromegaly.
- Use of thiazolindiones in the previous 3 months.
- History of ketoacidosis.
- Need for insulin within the previous 3 months.
- Significant concomitant diseases or complications of diabetes.
- Fasting triglycerides >5.1 mmol/L (>450 mg/dL) within the past 4 weeks.
- Treatment with systemic steroids.
- Patients with any history of gastrointestinal surgery e.g. partial bowel resections, partial gastric resections, etc.

Investigational drug:

LAF237 100mg MR (herein described preferred modified release MR formulation)

LAF237 150mg MR (herein described preferred modified release MR formulation)

LAF237 50mg (as described in the US provisional patent application No. 60/604274)

Duration of treatment: 46 days total study length, including washout periods.

Treatment A (LAF237 100mg MR OD): Single dose treatment on Day 1and 8 days of multiple dose treatment starting on Day 3.

Treatment B (LAF237 150mg MR OD): Single dose treatment on Day 1 and 8 days of multiple dose treatment starting on Day 3.

Treatment C (50mg BID): BID dose treatment on Day 1 and 8 days of multiple dose treatment starting on Day 3.

Two 7 day wash out periods between above treatments.

Assessments and evaluations:

Background, demographic and administrative assessments

- Inclusion/exclusion criteria; Relevant medical history/Current medical conditions: screening, review at baselines
- Demography: screening
- Physical examination: screening, baselines, end of study
- Hepatitis screen, HIV screen: screening
- Alcohol test, Drug screen, Urine cotinine: screening, baselines
- Pregnancy test: screening, baselines, end of study
- Drug administration record: each time study drug is administered
- Meal record: days 1, 2, 8, 9, 10 and 11.
- Study completion information: end of study
- Comments: as required

Safety and tolerability assessments

- Vital signs and body measurements
  - Body height: screening
  - Body weight: screening, baselines, end of study
  - Body temperature: screening, baselines, end of study
  - Blood pressure, pulse rate: screening*, baseline*, predose and 1, 6, 12, and 24 h postdose on Days 1, 2, 8, 9, and 10; end of study (Note: measurements are supine, or supine and after 3 min standing if indicated by an *asterisk)
- ECG evaluation: screening, baselines, end of study
- Hematology; Blood chemistry; Urinalysis: screening, baselines, end of study
- Adverse events; Concomitant medications/Significant non-drug therapies: from time of first administration of study drug until end of study

Pharmacokinetic assessments

- Blood collection (2 mL blood per sample, heparin tubes (plasma)):

On Days 1, 9*,10: pre-dose, 0.5, 1, 1.5, 2, 2.5, 3, 4, 6, 8,10,12,14,16,24,36, and 48 hr post-dose

On Days 5 and 8: predose.

*Samples collected to 24 hr postdose on Day 9. The Day 10 predose sample will also serve as the Day 9 24 hr sample

- Analytes, media and methods: LAF237 in plasma will be determined by LC-MS-MS with LOQ at 2ng/mL.
- PK parameters: AUC_{0-t}, $AUC_{(0-\infty)}$, C_{max}, t_{max}, $t_{½}$, and $C_{max}/AUC_{(0-t)}$ will be evaluated for each treatment period.

Pharmacodynamic assessments

DPP-IV activity will be measured on Days 1,, 9 and 10.

- Blood collection (1 mL blood per sample, EDTA tubes (plasma)):

On Days 1, 9* and 10: pre-dose, 0.25, 0.5, 0.75, 1, 2, 4, 6, 8, 10, 12, 14, 16, 24, 36 and 48 hr post-dose

*Samples collected to 24 hr post-dose on Day 9. The Day 10 predose sample will also serve as the Day 9 24 hr sample.

Glucose samples on Day 9.

- Blood Collection (2 mL blood per sample, sodium fluoride tubes (plasma)) Following morning dose- 0.75, 1, 1.25, 1.5, 2, 2.5, 3, 4, 5 (pre-lunch meal), 5.5, 5.75, 6, 6.5, 7, 8, 9.75 (prior to evening dose and pre-dinner meal), 10.25, 10.75, 11, 11.25, 11.5, 12, 12.5, 13, 14, 16 and 24 hr post-dose.

Statistical methods

Sample Size: The sample size is determined based on a two one-sided t-test at 5% significance level (Chen et. al. 1997) for PK parameters. When test-reference ratio is equal to 0.95, a sample size of 24 subjects allow at least 88% power to meet the bioequivalence criterion, i.e., to have the 90% CI for test-reference ratio of a bioavailability measure contained in the bioequivalence range of (0.8, 1.25) if intrasubject CV is not greater than 0.20, or 72% power if the intra-subject CV is equal to 0.25. The intrasubject CV values used in the sample size determination are derived from a previous study of similar formulation (LAF237A2214) in healthy subjects in which the largest observed intrasubject CV is around 0.20 for Cmax.

Statistical analysis: An analysis of variance (ANOVA) are performed on log-transformed AUC and Cmax data using the PROC MIXED SAS procedure. The

sources of variation included in the ANOVA model are sequence, subject (sequence), period, and treatment, with subject (sequence) as random effect. Using the ESTIMATE statement of the PROC MIXED SAS procedure, the contrast are constructed between the test and the reference treatments to obtain the p-value, the estimated mean difference, an the 90% confidence interval (CI) for the log-scale test-reference difference. The anti-logs of the estimated mean difference and the 90% CI constitute the ratio of geometric means and the 90% CI for the true test-reference ratio. The outputs from the comparison are tabulated.

For comparison of bioavailability between MR OD and IR BID, the test will be the MR OD and the reference are IR BID. The LAF237 AUC and Cmax are compared between the two treatments within the ANOVA.

The PD endpoints are analyzed using the same ANOVA model as described above.

Drug administration

On Days 1, 3, 5, 8, 9 and 10 study medication arre administered by the study center personnel with 240 mL of water between 0700 and 0800, after at least a 10-hour fast. All subjects are dosed within a maximum of a 1-hour interval. Subjects must be instructed not to chew the medication, but to swallow it whole. The investigator has to check each subject's mouth to ensure that the medication was swallowed. Unless performing a study assessment, subjects have to rest quietly in the upright position for the next 4 hours. With the exception of Days 1 and 10 when the first meal is served 5 hours post-dose, an ADA breakfast is served within 30 minutes of dosing.

During Treatment C, patients receive the evening dose of 50mg LAF237 30 minutes prior to dinner.

General study conditions and restrictions

Definition of study phases

Screening

"Screening" is defined as assessments made for 2 to 28 days prior to initiation of the study (i.e. prior to the day of first dosing, Day 1) to determine eligibility for entry into this study. Potentially eligible subjects whose screening examinations show any disqualifying abnormality may have the examination repeated ONCE if the investigator wishes to enroll the subject.

Baseline

If not yet domiciled, subjects should be admitted to the study site at least 10-hours prior to dosing in each treatment period. "Baseline" is defined as the period of a subject's continuous presence at the study facility to 30 minutes prior to the administration of the study drug (i.e.

Day -1). Baseline activities include safety assessments, as outlined in the Study synopsis and on the Assessment schedule.

Treatment period(s)

Each treatment period includes the predose activities (within 30 minutes prior to dosing), drug administration, the postdose evaluations until 24-hours after dosing, and - if applicable - the washout period until predose of the following treatment period.

End of study

End of study evaluations will be performed on the last day of the last treatment period, prior to discharge from the study site.

Domiciling and physician coverage

Patients are confined to the study center for at least 10 hours before administration of study drug on Day 1 until one hour after study drug is administered on Day 3. Patients return to the study center on the morning of day 5 for pre-dose blood collection. Patients are domiciled at least 10 hours before study drug is administered on Day 8 until 48 hours after the Day 10 morning dose of LAF237 is received.

Presence of a physician on site is expected at a minimum for the first 2 hours after dosing during each period. A physician shall be available by pager at all other times throughout the study.

Dietary, fluid and other restrictions

During recruitment, informed consent review, and baseline .period, the subjects is informed and reminded of the following restrictions:

- No strenuous physical exercise (e.g., weight training, aerobics, football) for 7 days before dosing until after the end of study evaluation.
- Diet should be free of read meat for at least 72 hours prior to the study entry until the end of study.
- No alcohol for 72 hours before dosing until after the end of study evaluation.
- Intake of xanthine (e.g. caffeine) containing food or beverage must be discontinued 49 hours prior to dosing. Consumption of such food and beverage (i.e. coffee, tea, soda, chocolate) is not permitted at any time while the subjects are domiciled. If a deviation occurs during the domiciled period, it must be noted on the Comments CRF page.

On all domiciled days, except Days 1 and 10, patients receive the study drug 30 minutes prior to an ADA breakfast and an overnight fast of 10 hours. On Day 1 and 10, no breakfast is served.

During Treatment C, patients also receive the evening dose of study drug 30 minutes prior to dinner. The tested drug should be administered with 240 mL (8 fl oz) of water. No fluid intake

apart from the fluid given at the time of drug intake is allowed from 2 h before until 2 h after dosing. Otherwise, subjects should have a fluid intake of at least 200 mL every 4 hours during waking hours in addition to fluid taken with meals and medication.

During domiciled days, lunch and dinner are served at 5 hours and 10 hours post dose, respectively, and a large snack is served at 12 hours post dose.

Subjects follow a standard weight maintaining diet while domiciled on non-dosing days. No other food is consumed at any time during confinement. Subjects should consume the entire contents of the meal. Meals should be similar in caloric content and distribution for all subjects on the day of dosing.

When meal and blood draw times coincide, blood is drawn BEFORE the meal is provided.

Intake of xanthine (e.g., caffeine) containing food or beverages must be discontinued 48 hours before dosing. Consumption of such foods and beverages (i.e., coffee, tea, and soda, chocolate) is not permitted at any time while the subjects are domiciled. If a deviation occurs during the domicile period, it must be noted on the Comments CRF page.

Background, demographic and administrative assessments

Inclusion/exclusion criteria

Subject selection is to be established by checking through all standard inclusion/exclusion criteria. A relevant record (e.g. checklist) must be stored with the source documentation at the study site. Violation of any entry criterion excludes a subject from enrollment into the study unless granted specific clearance by the sponsor.

Drug administration record

Date and time of dose administration is recorded in the Dosage administration record section of the CRF and - if applicable - on the Blood collection CRF page for pharmacokinetics and pharmacodynamic evaluations.

Meal record

The date and start time of meal consumption is recorded in the appropriate section of the CRF for all domiciled days.

Study completion information

Information on the date the subject last took drug, the subject's completion or discontinuation of the study and the reason for discontinuation of the study is recorded on the Study completion CRF page.

Pharmacokinetic assessments

Blood collection : All blood samples is taken by either direct venipuncture or an indwelling cannula inserted in a forearm vein.

LAF237: For each scheduled LAF237 sample, collect a 2 mL blood sample into a sodium or lithium heparin tube at the times specified in the assessment schedule.

Handling of blood samples

Immediately after each tube of blood is drawn, it should be inverted gently several times to insure the mixing of tube contents (e.g., anticoagulant). Avoid prolonged sample contact with the rubber stopper. Place the tube upright in a test tube rack (e.g. surrounded by ice or at room temperature) until centrifugation. Within 15 minutes, centrifuge the sample between 3 and 5°C for 15 minutes at approximately 2500 rpm. Transfer all available plasma to a polypropylene screw-cap tube and freeze at -70°C or below within 60 minutes of venipuncture.

Analytical method(s)

Analytes, media and methods: LAF237 in plasma by LC-MS-MS; LLOQ at 2 ng/mL.

Pharmacodynamic assessments: All blood samples are taken either by direct venipuncture or an indwelling cannula inserted in a forearm vein from each patient.

Glucose: For each scheduled glucose sample, collect a 2 mL blood sample into a sodium flouride (grey top) tube at the times specified in the assessment schedule. The central laboratory to be used for analysis of plasma glucose samples has not been determined at this point. Once determined, a protocol supplement do outline the sample handling and data transfer procedures.

DPP-IV: DPP-IV measurements is conducted by Novartis Laboratory. For each scheduled sample collection for DPP-4 enzymatic analysis, collect a 1 ml blood sample into a tube containing potassium EDTA. Invert gently several times to mix the contents of the tube. Avoid prolonged sample contact with the rubber stopper. Place the tube upright in rack surrounded by ice until centrifugation. Within 15 minutes after collection, centrifuge the sample between 3 and 5°C for 15 mins at approximately 2500 rpm. Transfer all available plasma to a polypropylene screw-cap micro tube and freeze at -70°C or below within 60 minutes of venipuncture

For labeling and shipment instructions, see Part B, Section 9.

Data analysis

Analysis of the data will be under the direction of Novartis personnel.

General statistical considerations

Sample size determination

The sample size is determined based on a two one-sided t-test at 5% significance level (Chen et. al. 1997 - Chen KW, Chow SC, Li G (1997). A note on sample size determination for bioequivalence studies with higher-order crossover designs. J Parmarcokin, Biopharm. 25: 753-765.) for PK parameters. When test-reference ratio is equal to 0.95, a sample size of 24 subjects allows at least 88% power to meet the bioequivalence criterion, i.e., to have the 90% CI for test-reference ratio of a bioavailability measure contained in the bioequivalence range of (0.8, 1.25) if intrasubject CV is not greater than 0.20, or 72% power if the intra-subject CV is equal to 0.25. The intrasubject CV values used in the sample size determination are derived from a previous study of similar formulation [LAF237A2214] in healthy subjects in which the largest observed intrasubject CV is around 0.17 for C_{max}.

Background, demographic and administrative data analysis

Descriptive statistics are provided for background and demographic variables such as age, weight, height, gender and race.

Relevant medical history, current medical conditions, results of laboratory screens, drug tests and any other relevant information are listed.

Safety and tolerability data analysis

All subjects who received at least one treatment are included in the safety and tolerability evaluation.

Pharmacokinetic data analysis

All completed subjects are included in the pharmacokinetic data analysis.

Pharmacokinetic variables

The following pharmacokinetic parameters are evaluated for each treatment period: AUC_{0-t}, $AUC_{(0-\infty)}$, C_{max}, t_{max}, $t_{½}$, and $C_{max}/AUC_{(0-t)}$.

Biofluid concentrations are expressed in mass per volume units. All concentrations below the limit of quantitation or missing data are labeled as such in the concentration data listings. Concentrations below the limit of quantitation are treated as zero in summary statistics and for the calculation of pharmacokinetic parameters.

Descriptive statistics of pharmacokinetic parameters include mean, SD, and CV, min and max. When a geometric mean is presented it is stated as such. A range of values is presented for selected variables. Since t_{max} is generally evaluated by a nonparametric method, median values and ranges are given for this parameter.

Pharmacokinetic parameters are determined based on non-compartmental method(s) using WinNonlin Pro.

Statistical methods for pharmacokinetic analyses

An analysis of variance (ANOVA) are performed on log-transformed AUC and Cmax data using the PROC MIXED SAS procedure. The sources of variation included in the ANOVA model are sequence, subject (sequence), period, and treatment, with subject (sequence) as random effect. Using the ESTIMATE statement of the PROC MIXED SAS procedure, the contrast are constructed between the test and the reference treatments to obtain the p-value, the estimated mean difference, an the 90% confidence interval (CI) for the log-scale test-reference difference. The anti-logs of the estimated mean difference and the 90% CI constitute the ratio of geometric means and the 90% CI for the true test-reference ratio. The outputs from the comparison are tabulated.

For comparison of bioavailability between MR OD and IR BID, the test treatment is the MR OD and the reference is IR BID. The LAF237 AUC and Cmax are compared between the two treatments within the ANOVA.

Pharmacodynamic data analysis

All subjects who complete the trial with evaluable pharmacodynamic (PD) measurements are included in the data analysis.

Pharmacodynamic variables

Following glucose parameters may be derived for treatment comparison:

AUE area under the effect-time curve from time 0 to last available observation by the linear-trapezoidal rule

E_{max} maximal effect (minimum value)

t_{max} time to the first occurrence of the maximal effect.

In order not to lose data from subjects due to missing values, the algorithm below are employed to replace missing values:

- If the first or the last observation of a subject's record is missing then they are replaced by the second or the previous observation.
- If an observation is missing within a series of non-missing observations, then the missing values are substituted by the result of a linear regression based on its neighbors.
- If two consecutive observations are missing, then the record is completely ignored.

Statistical methods for pharmacodynamic analyses

The glucose parameters are analyzed using the same ANOVA model as described above for statistical analysis of PK parameters.

In addition, timepoint-wise comparison of glucose concentration are performed using the same ANOVA model as described above.

Results:

Pharmacokinetic results (described on figures 24, 25, 26)

Treatment	Tmax (h) median (min, max)	Cmax (ng/mL) mean ± SD (CV%)	AUC a (h•ng/mL) mean ± SD (CV%)	CL/F (L/h) mean ± SD (CV%)	t½ (h) mean ± SD (CV%)
Day 1 (Single Dose Fasted) (N=27)					
LAF237 100 mg MR OD	4.00 (1.50, 6.00)	205 ± 47 (23)	1449 ± 376 (26)	73.60 ± 19.60 (27)	11.69 ± 5.68 (49)
LAF237 150 mg MR OD	4.00 (1.50, 6.00)	257 ± 59 (23)	2271 ± 925 (41)	75.63 ± 28.54 (38)	10.75 ± 5.01 (47)
LAF237 50 mg MR BID	1.00 (0.50, 4.00)	278 ± 75 (27)	2159 ± 425 (20)	45.14 ± 7.91 (18)	2.43 ± 1.12 (46)
Day 9 (Multiple Dose Fed) (N=27)					
LAF237 100 mg MR OD	2.00 (1.00, 6.00)	200 ± 64 (32)	1489 ± 488 (33)	75.45 ± 29.07 (39)	8.70 ± 3.75 (43)
LAF237 150 mg MR OD	2.50 (1.00, 6.00)	272 ± 111 (41)	2242 ± 965 (43)	78.84 ± 31.30 (40)	10.34 ± 5.06 (49)
LAF237 50 mg BID	1.00 (1.00, 4.00)	285 ± 91 (32)	1992 ± 461 (23)	50.92 ± 16.51 (32)	2.83 ± 0.62 (22)
Day 10 (Multiple Dose Fasted) (N=27)					
LAF237 100 mg MR OD	3.00 (1.00, 6.00)	215 ± 68 (31)	1638 ± 535 (33)	66.77 ± 20.87 (31)	11.80 ± 5.69 (48)
LAF237 150 mg MR OD	4.00 (2.50, 6.00)	309 ± 91 (29)	2458 ± 815 (33)	67.19 ± 21.27 (32)	10.54 ± 5.39 (51)
LAF237 50 mg MR BID	2.00 (0.50, 9.97)	257 ± 60 (23)	2198 ± 492 (22)	44.92 ± 10.97 (24)	3.38 ± 2.62 (77)

a: AUC0-inf on Day 1, AUC0-24 on Days 9 and 10, AUC0-24 on Day 1 for LAF237 50 mg BID

Results:

Pharmacodynamic results (described on figures 21, 22, 23)


Treatment	DPP-4-AUC24 (%.h) mean ± SD (CV%	DPP-4-MRT (h) mean ± SD (CV%)	DPP-4-Avg (%) mean ± SD (CV%)	DPP-4-24h (%) mean ± SD (CV%)
Day 1 (Single Dose Fasted) (N=27)				
LAF237 100 mg MR	2208 ± 73 (3)	11.8 ± 0.3 (3)	92.01 ± 3.03	85.64 ± 12.76

OD			(3)	(15)
LAF237 150 mg MR OD	2242 ± 70 (3)	11.9 ± 0.3 (3)	93.43 ± 2.93 (3)	90.04 ± 11.91 (13)
LAF237 50 mg MR BID	2225 ± 48 (2)	11.7 ± 0.2 (2)	92.69 ± 2.01 (2)	80.92 ± 9.03 (11)
Day 9 (Multiple Dose Fed) (N=27)				
LAF237 100 mg MR OD	2240 ± 98 (4)	11.8 ± 0.5 (4)	93.35 ± 4.09 (4)	87.78 ± 16.37 (19)
LAF237 150 mg MR OD	2270 ± 86 (4)	11.8 ± 0.5 (4)	94.59 ± 3.57 (4)	90.40 ± 17.50 (19)
LAF237 50 mg BID	2252 ± 33 (1)	11.8 ± 0.1 (1)	93.85 ± 1.36 (1)	86.15 ± 4.78 (6)
Day 10 (Multiple Dose Fasted) (N=27)				
LAF237 100 mg MR OD	2261 ± 40 (2)	11.9 ± 0.2 (1)	94.21 ± 1.66 (2)	90.20 ± 7.35 (8)
LAF237 150 mg MR OD	2281 ± 49 (2)	11.9 ± 0.2 (2)	95.06 ± 2.03 (2)	91.64 ± 8.47 (9)
LAF237 50 mg MR BID	2243 ± 46 (2)	11.7 ± 0.2 (1)	93.44 ± 1.94 (2)	82.80 ± 7.39 (9)

What is claimed is:

1. A pharmaceutical tablet formulation comprising per unit dosage form e.g. per tablet the following ingredients:

(a) a compound as an active ingredient, wherein the compound has a formula:



wherein R is substituted adamantyl and n is an integer from 0 to 3; or a pharmaceutically acceptable salt thereof;

(b) a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) a microcrystalline cellulose; and

(d) a magnesium stearate.

2. The pharmaceutical tablet formulation of claim 1, wherein relative to the weight of the formulation:

(a) the compound is present in an amount from 20% to 30% by weight;

(b) the hydroxypropyl methylcellulose is present in an amount from 30% to 50% by weight;

(c) the microcrystalline cellulose is present in an amount from 25% to 35% by weight; and

(d) the magnesium stearate is present in an amount from 0.1% to 3% by weight.

3. The pharmaceutical tablet formulation of claim 1 or 2, wherein relative to the weight of the formulation:

(a) the compound is present in an amount of about 25% by weight;

(b) the hydroxypropyl methylcellulose is present in an amount of about 40% by weight;

(c) the microcrystalline cellulose is present in an amount of about 30% by weight; and

(d) the magnesium stearate is present in an amount of about 1% by weight.

4. The pharmaceutical tablet formulation of any of claims 1-3, further comprising a lactose.

5. The pharmaceutical tablet formulation of claim 4, wherein the lactose is present in an amount from 1% to 8% by weight.

6. The pharmaceutical tablet formulation of claim 3, further comprising a lactose in an amount of about 4% by weight.

7. The pharmaceutical tablet formulation of any of claims 1-6, wherein the compound is 1-[2-[(5-cyanopyridin-2-yl)amino]ethylamino]acetyl-2-cyano(S)-pyrrolidone or a pharmaceutically acceptable salt thereof.

8. The pharmaceutical tablet formulation of any of claims 1-6, wherein the compound is vildagliptin or a pharmaceutically acceptable salt thereof.

9. The pharmaceutical tablet formulation of claim 8, wherein the compound is a crystal form of vildagliptin preferably the crystal form "A" or a pharmaceutically acceptable salt thereof.

10. A pharmaceutical tablet formulation, comprising per 400 mg tablet the following ingredients:

(a) vildagliptin or a pharmaceutically acceptable salt thereof.

(b) a hydroxypropyl methylcellulose in an amount of about 160 mg, wherein the hydroxypropyl methylcellulose has an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) a microcrystalline cellulose in an amount of 120 mg;

(d) a lactose in an amount of about 16 mg; and

(e) a magnesium stearate in an amount of 4 mg.

11. A method of producing the pharmaceutical tablet formulation of any of claims 1-10 which comprises combining the ingredients in the amounts recited in such claim.

12. A method of inhibiting dipeptidyl peptidase IV activity in a subject comprising administering to the subject an amount of the pharmaceutical tablet formulation of any of claims 1-10 effective to inhibit the activity of dipeptidyl peptidase IV in the subject.

13. The method of claim 12, wherein the subject is a human being.

14. A method of treating a condition alleviated by dipeptidyl peptidase IV inhibition in a subject suffering from the condition comprising administering to the subject a therapeutically effective dose of the pharmaceutical tablet formulation of any of claims 1-10.

15. The method of claim 14, wherein the condition is non-insulin-dependent diabetes mellitus.

16. The method of claim 14, wherein the condition is obesity, arthritis, or osteoporosis.

17. The method of any of claims 14-16, wherein the subject is a human being.


18. A method of treating a condition alleviated by dipeptidyl peptidase IV inhibition in a subject suffering from the

condition comprising administering to subject a therapeutically effective amount of the pharmaceutical tablet formulation of any of claims 1-10 in combination with a therapeutically effective dose of an anti-diabetic or arthritis drug.

19. The method of claim 18, wherein the subject is a human being.

20. A pharmaceutical tablet formulation comprising per unit dosage form e.g. per tablet the following ingredients:

(a) a compound as an active ingredient, wherein the compound has a formula:



wherein R is substituted adamantyl and n is an integer from 0 to 3; or a pharmaceutically acceptable salt thereof; and

(b) a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution.

21. The pharmaceutical tablet formulation of claim 20, wherein it also contains a filler.

22. The pharmaceutical tablet formulation of claim 21, wherein the filler is lactose.

23. The pharmaceutical tablet formulation of claim 21, wherein the filler is microcrystalline cellulose.

24. The pharmaceutical tablet formulation of claim 20, wherein it also contains a lubricant.

25. The pharmaceutical tablet formulation of claim 24, wherein the lubricant is magnesium stearate.

26. The pharmaceutical tablet formulation of claim 20, wherein relative to the weight of the formulation, the hydroxypropyl methyl cellulose is present in an amount from 30% to 50% by weight.

27. The pharmaceutical tablet formulation of claim 20, wherein the hydroxypropyl methyl cellulose is present in an amount from 34% to 46% preferably from 38% to 42% by weight.

28. A pharmaceutical tablet formulation comprising per unit dosage form e.g. per tablet the following ingredients:

(d) vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(e) a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution,

(f) and optionally a filler and/or a lubricant.

29. A pharmaceutical tablet formulation according to claim 28, wherein in the unit dosage form, the ratio of the weight of vildagliptin to the weight of hydroxypropyl methylcellulose is of 0.16 to 2.5, preferably 0.3 to 1.16 or 0.4 to 1.

30. A pharmaceutical tablet formulation according to claim 28 or claim 29 comprising;

(a) 15-55% preferably 25-45% by weight on a dry weight basis of a pharmaceutically acceptable filler; and optionally

(b) 0.1-10% preferably 0.1-3% by weight on a dry weight basis of a pharmaceutically acceptable lubricant.

31. A pharmaceutical tablet formulation according to claim 28 or claim 29 comprising;

(a) 15-55% preferably 25-45% by weight on a dry weight basis of one or two pharmaceutically acceptable fillers selected from lactose and microcrystalline cellulose ; and optionally

(b) 0.1-10% preferably 0.1-3% by weight on a dry weight basis of a pharmaceutically acceptable lubricant.

32. A pharmaceutical tablet formulation according to claim 28 comprising per unit dosage form:

(a) 10-50% preferably 15-35% by weight on a dry weight basis of vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(b) 20-60% preferably 30-50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution,

and optionally a filler and/or a lubricant.

33. A pharmaceutical tablet formulation according to claim 28 comprising per unit dosage form:

(a) 10-50% preferably 15-35% by weight on a dry weight basis of vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(b) 20-60% preferably 30-50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) a filler; and

(d) a lubricant.

34. A pharmaceutical tablet formulation according to claim 28 comprising per unit dosage form:

(c) 10-50% preferably 15-35% by weight on a dry weight basis of vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(d) 20-60% preferably 30-50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(e) 15-55% preferably 25-45% by weight on a dry weight basis of a pharmaceutically acceptable filler; and optionally

(f) 0.1-10% preferably 0.1-3% by weight on a dry weight basis of a pharmaceutically acceptable lubricant.

35. A pharmaceutical tablet formulation according to claim 28 comprising per unit dosage form:

(c) 10-50% preferably 15-35% by weight on a dry weight basis of vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(d) 20-60% preferably 30-50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(e) 15-55% preferably 25-45% by weight on a dry weight basis of one or two pharmaceutically acceptable fillers selected from lactose and microcrystalline cellulose ; and optionally

(f) 0.1-10% preferably 0.1-3% by weight on a dry weight basis of a pharmaceutically acceptable lubricant.

36. A pharmaceutical tablet formulation according to claim 28 comprising per unit dosage form:

(a) 10-50% preferably 15-35% by weight on a dry weight basis of vildagliptin, or a pharmaceutically acceptable salt thereof as an active ingredient,

(b) 20-60% preferably 30-50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) 25-40% by weight on a dry weight basis of one or two pharmaceutically acceptable fillers selected from lactose and microcrystalline cellulose ; and optionally

(d) 0.1-10% preferably 0.1-3% by weight on a dry weight basis of a pharmaceutically acceptable lubricant.

37) A pharmaceutical tablet formulation according to any of claims 28-36 wherein the filler is selected from lactose and microcrystalline cellulose.

38) A pharmaceutical tablet formulation according to any of claims 28-37 comprising at least two fillers.

39) A pharmaceutical tablet formulation according to claim 38 wherein the filers are lactose and microcrystalline cellulose.

40) A pharmaceutical tablet formulation according to claim 39 wherein the lactose is present in an amount from 1 to 8% preferably 1 to 5% by weight and microcrystalline cellulose is present in an amount from 25 to 35% by weight

41) A pharmaceutical tablet formulation according to any of claims 28-40 wherein 20-30% by weight on a dry weight basis of vildagliptin is contained in the formulation.

42) A pharmaceutical tablet formulation according to any of claims 28-41 wherein the hydroxypropyl methyl cellulose is present in an amount from 34% to 46% preferably from 38% to 42% by weight.

43) A pharmaceutical tablet formulation according to any of claims 28-42 wherein the lubricant is magnesium stearate.

44) A pharmaceutical tablet formulation according to any of the previous claims wherein in the unit dosage form, vildagliptin is present in an amount of 100 mg to 200 mg, or the corresponding amount of any of its salt.

45) A pharmaceutical tablet formulation according to any of the previous claims wherein in the unit dosage form, vildagliptin is present in an amount of 100 mg, 150 mg or 200 mg or the corresponding amount of any of its salt.

46) A pharmaceutical tablet formulation, comprising per 600 mg tablet the following ingredients:

(a) 1-[(3-hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile or a pharmaceutically acceptable salt thereof in an amount of about 150 mg;

(b) a hydroxypropyl methylcellulose in an amount of about 240 mg, wherein the hydroxypropyl methylcellulose has an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) a microcrystalline cellulose in an amount of 180 mg;

(d) a lactose in an amount of about 24 mg; and

(e) a magnesium stearate in an amount of 6 mg.

47) A pharmaceutical tablet formulation, comprising per 400 mg tablet the following ingredients:

(a) 1-[(3-hydroxy-adamant-1-ylamino)-acetyl]-pyrrolidine-2(S)-carbonitrile or a pharmaceutically acceptable salt thereof in an amount of about 100 mg;

(b) a hydroxypropyl methylcellulose in an amount of about 160 mg, wherein the hydroxypropyl methylcellulose has an apparent viscosity of

80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution;

(c) a microcrystalline cellulose in an amount of 120 mg;

(d) a lactose in an amount of about 16 mg; and

(e) a magnesium stearate in an amount of 4 mg.

48) A pharmaceutical multilayer tablet wherein the pharmaceutical tablet formulation of any of claims 28 to 47, represents one of the tablet layers.

49) A pharmaceutical multilayer tablet according to claim 48 wherein the further layer contains a glitazone (e.g. pioglitazone or rosiglitazon) or metformin.

50) A pharmaceutical multilayer tablet according to claim 48 wherein the further layer is an immediate release formulation which contains a glitazone (e.g. pioglitazone or rosiglitazon) or metformin.

51) A pharmaceutical tablet obtained by compression of a pharmaceutical tablet formulation of any of the claims 1 to 47.

52) A pharmaceutical tablet obtained by compression of a pharmaceutical tablet formulation of any of the claims 1 to 47, wherein the pharmaceutical tablet formulation is subject to roller compaction before compression into tablet.

53) A pharmaceutical tablet according to claim 52 comprising 100 mg of vildagliptin or a salt thereof wherein the tablet hardness range is of between 10 to 13Kp.

54) A pharmaceutical tablet according to claim 52 comprising 150 mg of vildagliptin or a salt thereof wherein the tablet hardness range is of between 11 to 25Kp.

55) A pharmaceutical tablet formulation comprising 100 mg of vildagliptin or a salt thereof, preferably compressed in the form of a sustained release tablet, wherein;

- between 10% and 16% preferably between 11% and 15% of vildagliptin is released after 0.5 hour,

- between 18% and 24% preferably between 19% and 23% of vildagliptin is released after 1 hour,

- between 30% and 36% preferably between 31% and 35% of vildagliptin is released after 2 hours,

- between 46% and 52% preferably between 47% and 51% of vildagliptin is released after 4 hour,

- between 58% and 64% preferably between 59% and 63% of vildagliptin is released after 6 hours,

- between 67% and 73% preferably between 68% and 72% of vildagliptin is released after 8 hours,

- between 74% and 80% preferably between 75% and 79% of vildagliptin is released after 10 hours,

- between 80% and 86% preferably between 81% and 85% of vildagliptin is released after 12 hours,

- between 91% and 97% preferably between 92% and 96% of vildagliptin is released after 18 hours,

- between 95% and 100% preferably between 96% and 100% of vildagliptin is released after 24 hours.

56) A pharmaceutical tablet formulation comprising 150 mg of vildagliptin or a salt thereof, preferably compressed in the form of a sustained release tablet, wherein;

- between 3.8% and 9.8% preferably between 4.8% and 8.8% of vildagliptin is released after 0.25 hour,

- between 8.1% and 14.1% preferably between 9.1% and 13.1% of vildagliptin is released after 0.5 hour,

- between 14.7% and 20.7% preferably between 15.7% and 19.7% of vildagliptin is released after 1 hours,

- between 25.3% and 31.3% preferably between 26.3% and 30.3% of vildagliptin is released after 2 hour,

- between 40.9% and 46.9% preferably between 41.9% and 45.9% of vildagliptin is released after 6 hours,

- between 62.1% and 68.1% preferably between 63.1% and 67.1% of vildagliptin is released after 8 hours,

- between 76.5% and 82.5% preferably between 77.5% and 81.5% of vildagliptin is released after 10 hours,

- between 83.5% and 89.5% preferably between 84.5% and 88.5% of vildagliptin is released after 12 hours,

- between 88.5% and 94.5% preferably between 89.5% and 93.5% of vildagliptin is released after 18 hours.

57) A pharmaceutical tablet formulation according to claims 55 or 56, which comprises a hydroxypropyl methylcellulose, preferably between 20% and 60%, between 30% and 50% by weight on a dry weight basis of a hydroxypropyl methylcellulose.

58) A pharmaceutical tablet formulation according to claims 55 or 56, which comprises a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution, preferably between 20% and 60%, or between 30% and 50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution.

59) A pharmaceutical tablet formulation according to any of claims 55 to 56 which is a pharmaceutical tablet formulation according to any of claims 1 to 47.

60) A pharmaceutical tablet obtained by compression of a pharmaceutical tablet formulation of any of claims 55 to 59.

61) A pharmaceutical tablet according to claim 60 obtained by compression of a pharmaceutical tablet formulation of any of the claims 1 to 47 or 55 to 59, wherein the pharmaceutical tablet

formulation is subject to roller compaction before compression into tablet.

62) A pharmaceutical tablet according to claim 60 or claim 61 comprising 100 mg of vildagliptin or a salt thereof wherein the tablet hardness range is of between 10 to 13Kp.

63) A pharmaceutical tablet according to claim 60 or claim 61 comprising 150 mg of vildagliptin or a salt thereof wherein the tablet hardness range is of between 11 to 25Kp.

64) A pharmaceutical capsule, tablet, compressed table, direct compressed tablets, granule comprising a pharmaceutical tablet formulation of any of the claims 1 to 47 or 55 to 59.

65) A pharmaceutical tablet formulation or a pharmaceutical tablet according to any of claims 1 to 64, wherein the dispersion contains particles comprising DPP-IV inhibitor preferably vildagliptin, in free form or in acid addition salt form, and wherein at least 60%, preferably 80% and most preferably 90% of the particle size distribution in the formulation is less than 250 μm or preferably between 10 to 250 μm.

66) A pharmaceutical tablet formulation or a pharmaceutical tablet according to any of claims 1 to 64, wherein the dispersion contains particles comprising DPP-IV inhibitor preferably vildagliptin, in free form or in acid addition salt form, and wherein at least 60%, preferably 80% and most preferably 90% of the particle size distribution in the tablet is greater than 10 μm.

67) A pharmaceutical tablet formulation or a pharmaceutical tablet according to any of claims 1 to 66, wherein the dispersion contains particles comprising DPP-IV inhibitor preferably LAF237, in free form or in acid addition salt form, and wherein at least 25%, preferably 35% and most preferably 45% of the particle size distribution in the formulation is between 50 to 150 μm.

68) A pharmaceutical tablet formulation or a pharmaceutical tablet according to any of claims 1 to 67, comprising at least one further therapeutic agent.

69) A pharmaceutical tablet formulation or a pharmaceutical tablet according to claim 68, comprising at least one further therapeutic agent selected from an antidiabetic, an angiotensin II antagonist or a statin.

70) A pharmaceutical tablet formulation or a pharmaceutical tablet according to any of claims 68 to 69, wherein the antidiabetic agent is selected from pioglitazone, rosiglitazone or metformin.

71) A pharmaceutical tablet formulation or a pharmaceutical tablet according to any of claims 1 to 70, wherein the pharmaceutical tablet formulation is in the form of a layer in a multi or 2-layer tablet.

72) Use of a pharmaceutical formulation, capsule, tablet, compressed table, direct compressed tablet, granule according to any of the previous claims for the manufacture of a medicament for the treatment of conditions, such as non-insulin-dependent diabetes mellitus, arthritis, obesity, allograft transplantation, calcitonin-osteoporosis, Heart Failure, Impaired Glucose Metabolism), IGT (Impaired Glucose Tolerance), neurodegenerative diseases such as Alzheimer's and Parkinson disease, modulating hyperlipidemia, modulating conditions associated with hyperlipidemia or for lowering VLDL, LDL and Lp(a) levels, cardiovascular or renal diseases e.g. diabetic cardiomyopathy, left or right ventricular hypertrophy, hypertrophic medial thickening in arteries and/or in large vessels, mesenteric vasculature hypertrophy, mesanglial hypertrophy, neurodegenerative disorders and cognitive disorders, to produce a sedative or anxiolytic effect, to attenuate post-surgical catabolic changes and hormonal responses to stress, to reduce mortality and morbidity after myocardial

infarction, the treatment of conditions related to the above effects which may be mediated by GLP-1 and/or GLP-2 levels.

73) A method of treating conditions, such as non-insulin-dependent diabetes mellitus, arthritis, obesity, allograft transplantation, calcitonin-osteoporosis, Heart Failure, Impaired Glucose Metabolism), IGT (Impaired Glucose Tolerance), neurodegenerative diseases such as Alzheimer's and Parkinson disease, modulating hyperlipidemia, modulating conditions associated with hyperlipidemia or for lowering VLDL, LDL and Lp(a) levels, cardiovascular or renal diseases e.g. diabetic cardiomyopathy, left or right ventricular hypertrophy, hypertrophic medial thickening in arteries and/or in large vessels, mesenteric vasculature hypertrophy, mesanglial hypertrophy, neurodegenerative disorders and cognitive disorders, to produce a sedative or anxiolytic effect, to attenuate post-surgical catabolic changes and hormonal responses to stress, to reduce mortality and morbidity after myocardial infarction, the treatment of conditions related to the above effects which may be mediated by GLP-1 and/or GLP-2 levels, comprising administering to a warm-blooded animal in need thereof a therapeutically effective amounts of a pharmaceutical formulation, capsule, tablet, compressed table, direct compressed tablet, granule according to any of the previous claims.

74) A process for preparing a tablet, in unit dosage form, which comprises:

(a) blending a pharmaceutical tablet formulation according to any of the previous claims,

(b) compressing the formulation prepared during step (a) to form the compressed tablet in unit dosage form.

75) A process for preparing a tablet, in unit dosage form, which comprises:

(a) blending a pharmaceutical tablet formulation according to any of the previous claims,

(b) roller compacting the formulation prepared during step (a)

(c) compressing the formulation prepared during step (b) to form the compressed tablet in unit dosage form.

76) A process for preparing a tablet, in unit dosage form, which comprises:

(a) blending a pharmaceutical tablet formulation according to any of the previous claims,

(b) roller compacting the formulation prepared during step (a) with a compaction force comprised between 10 and 16 KN,

(c) compressing the formulation prepared during step (b) to form the compressed tablet in unit dosage form.

77) A sustained release solid oral pharmaceutical dosage form which is;

i-1) a solid oral pharmaceutical dosage form comprising about 100 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma concentration of vildagliptin ranging from about 15.8 ng/mL +/- 6.85 ng/mL to about 173 ng/mL +/- 52 ng/mL between about 0.5 and about 16 hours following oral administration of said dosage form in a patient not treated with vildagliptin before said administration and wherein patient is under fasted conditions, and/or

i-2) a solid oral pharmaceutical dosage form comprising about 100 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma concentration of vildagliptin ranging from about 26.3 ng/mL +/- 13.1 ng/mL to about 175 ng/mL +/- 62.5 ng/mL between about 0.5 and about 16 hours following administration of said dosage form on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

i-3) a solid oral pharmaceutical dosage form comprising about 100 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma concentration of vildagliptin ranging from about 26.9 ng/mL +/- 14.1 ng/mL to about 186 ng/mL +/- 80.6 ng/mL between about 0.5 and about 16 hours following administration of said dosage form on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions, and/or

ii-1) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-inf)}$ of vildagliptin ranging from about 1073 to about 1825 ng•h/mL i.e. 1449 ng•h/mL +/- 376 ng•h/mL following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions, and/or

ii-2) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-24)}$ of vildagliptin ranging from about 1001 to about 1977 ng•h/mL i.e. 1489 ng•h/mL +/- 488 ng•h/mL following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

ii-3) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-24)}$ of vildagliptin ranging from about 1103 to about 2173 ng•h/mL i.e.

1638 ng•h/mL +/- 535 ng•h/mL following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions, and/or

iii-1) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 3.61 hr +/- 1.44 hr following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions, and/or

iii-2) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 2.59 hr +/- 1.4 hr following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

iii-3) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 3.74 hr +/- 1.44 hr following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions , and/or

iv-1) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean C_{max} of vildagliptin of 205 ng/ml +/- 47 ng/ml following oral

administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions, and/or

iv-2) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean C_{max} of vildagliptin of 200 ng/ml +/- 64 ng/ml following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

iv-3) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean C_{max} of vildagliptin of 245 ng/ml +/- 68 ng/ml following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions, and/or

v-1) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean % inhibition of DPP-IV activity over 24 hours of 85.64% +/- 12.76% following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions, and/or

v-2) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean % inhibition of DPP-IV activity over 24 hours of 87.78% +/- 16.37% following oral administration of said dosage form , on day 9 in a patient

treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

v-3) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean % inhibition of DPP-IV activity over 24 hours of 90.20% +/- 7.35% following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions, and/or

vi-1) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as substantially depicted in figure 24, following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions, and/or

vi-2) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as substantially depicted in figure 25, following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

vi-3) a solid oral dosage form comprising about 100 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as substantially depicted in figure 26, following oral administration of said dosage form , on day 10 in a patient

treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions.

78) A sustained release solid oral pharmaceutical dosage form according to claim 77, which comprises a pharmaceutical tablet formulations according to any of the previous claims.

79) A sustained release solid oral pharmaceutical dosage form which is;

i-a) a solid oral pharmaceutical dosage form comprising about 150 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma concentration of vildagliptin ranging from about 30.7 ng/mL +/- 21.9 ng/mL to about 223 ng/mL +/- 77.3 ng/mL between about 0.5 and about 16 hours following oral administration of said dosage form in a patient not treated with vildagliptin before said administration and wherein patient is under fasted conditions, and/or

i-b) a solid oral pharmaceutical dosage form comprising about 150 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma concentration of vildagliptin ranging from about 48.7 ng/mL +/- 25.8 ng/mL to about 223 ng/mL +/- 99.7 ng/mL between about 0.5 and about 16 hours following administration of said dosage form on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

i-c) a solid oral pharmaceutical dosage form comprising about 150 mg of vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean maximum plasma concentration of vildagliptin ranging from about 44.6 ng/mL +/- 28.5 ng/mL to about 263 ng/mL +/- 84.4 ng/mL between about 0.5

and about 16 hours following administration of said dosage form on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions, and/or

ii-a) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-inf)}$ of vildagliptin ranging from about 1346 to about 3196 ng•h/mL i.e. 2271 ng•h/mL +/- 925 ng•h/mL following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions, and/or

ii-b) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-24)}$ of vildagliptin ranging from about 1277 to about 3207 ng•h/mL i.e. 2242 ng•h/mL +/- 965 ng•h/mL following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

ii-c) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean $AUC_{(0-24)}$ of vildagliptin ranging from about 1643 to about 3273 ng•h/mL i.e. 2458 ng•h/mL +/- 815 ng•h/mL following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions, and/or

iii-a) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a

pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 3.57 hr +/- 1.17 hr following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions, and/or

iii-b) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 2.87 hr +/- 1.59 hr following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

iii-c) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean t_{max} of vildagliptin of 4.13 hr +/- 1.24 hr following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions, and/or

iv-a) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean C_{max} of vildagliptin of 257 ng/ml +/- 59 ng/ml following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions, and/or

iv-b) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium,

said dosage form providing an arithmetic mean C_{max} of vildagliptin of 272 ng/ml +/- 111 ng/ml following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

iv-c) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean C_{max} of vildagliptin of 308 ng/ml +/- 91 ng/ml following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions, and/or

v-a) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean % inhibition of DPP-IV activity over 24 hours of 90.04% +/- 11.91% following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions, and/or

v-b) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing an arithmetic mean % inhibition of DPP-IV activity over 24 hours of 90.4% +/- 17.50% following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

v-c) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium,

said dosage form providing an arithmetic mean % inhibition of DPP-IV activity over 24 hours of 91.64% +/- 8.47% following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions, and/or

vi-a) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as substantially depicted in figure 24, following oral administration of said dosage form , in a patient not treated with vildagliptin before said administration and wherein said patient is under fasted conditions, and/or

vi-b) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as substantially depicted in figure 25, following oral administration of said dosage form , on day 9 in a patient treated with said dosage form once a day since day 1 and wherein said patient has been served an ADA breakfast within 30 minutes of the morning administration of said dosage form, and/or

vi-c) a solid oral dosage form comprising about 150 mg vildagliptin free base, or a respective amount of a pharmaceutically acceptable salt thereof, and a carrier medium, said dosage form providing a pharmacokinetic profile as substantially depicted in figure 26, following oral administration of said dosage form , on day 10 in a patient treated with said dosage form once a day since day 1 and wherein said patient is under fasted conditions.

80) A sustained release solid oral pharmaceutical dosage form according to claim 79, which comprises a pharmaceutical tablet formulations according to any of the previous claims.

81) A sustained release solid oral pharmaceutical dosage form according to claim 79 or claim 77, which comprises a hydroxypropyl methylcellulose, preferably between 20% and 60%, between 30% and 50% by weight on a dry weight basis of a hydroxypropyl methylcellulose.

82) A sustained release solid oral pharmaceutical dosage form according to claim 79 or claim 77, which comprises a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution, preferably between 20% and 60%, or between 30% and 50% by weight on a dry weight basis of a hydroxypropyl methylcellulose with an apparent viscosity of 80,000 cP to 120,000 cP (nominal value 100,000 cP) when present in a 1% solution.

83) A sustained release solid oral pharmaceutical dosage form according to claim 79 or claim 77, wherein;

i) the formulation is a matrix formulation containing a pharmaceutically acceptable hydrophilic polymer which can retard diffusion of vildagliptin,

ii) the solid oral pharmaceutical dosage form is a compressed tablet, and optionally

iii) the elution rate of vildagliptin at 30 minutes after starting the test is less than 30% when conducting the Paddle method.