(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

G00002

(43) International Publication Date
25 July 2002 (25.07.2002)

PCT

(10) International Publication Number
WO 02/057468 A3

(51) International Patent Classification[7]: C12N 15/82, 5/10, C07K 14/00, A01H 5/00

(21) International Application Number: PCT/JP02/00361

(22) International Filing Date: 18 January 2002 (18.01.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2001-12519 19 January 2001 (19.01.2001) JP

(71) Applicant *(for all designated States except US)*: **THE DOW CHEMICAL COMPANY** [US/US]; 2030 Dow Center, Midland, MI 48674 (US).

(72) Inventors; and

(75) Inventors/Applicants *(for US only)*: **FUJIYAMA, Kazuhito** [JP/JP]; A18-308, 1-28, Yamadanishi, Suita-shi, Osaka 565-0824 (JP). **SEKI, Tatsuji** [JP/JP]; 2-1-1-1015, Shinsenrinishimachi, Toyonaka-shi, Osaka 565-0083 (JP). **YOSHIDA, Toshiomi** [JP/JP]; A1-505, 2-4, Yamadanishi, Suita-shi, Osaka 565-0824 (JP).


(74) Agents: **ISHIDA, Takashi** et al.; A. Aoki, Ishida & Associates, Toranomon 37 Mori Bldg., 5-1, Toranomon 3-chome, Minato-ku, Tokyo 105-8423 (JP).

(81) Designated States *(national)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States *(regional)*: ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: METHOD FOR SECRETORY PRODUCTION OF GLYCOPROTEIN HAVING HUMAN-TYPE SUGAR CHAIN USING PLANT CELL



162162

(57) Abstract: A method for the secretory production of a glycoprotein having a human-type sugar chain, comprising a step of introducing a gene of an enzyme capable of performing a transfer reaction of a galactose residue to a non-reducing terminal acetylglucosamine residue, and a gene of heterologous glycoprotein, to obtain a transformed plant cell, a step of culturing the plant cell, and a step of recovering the culture medium of the plant cell.

WO 02/057468 A3

Published:
— *with international search report*

(88) Date of publication of the international search report:
3 July 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.