



(51) International Patent Classification:  
*A61B 1/06* (2006.01) *A61B 18/20* (2006.01)

(21) International Application Number:  
PCT/US2019/027007

(22) International Filing Date:  
11 April 2019 (11.04.2019)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
62/656,639 12 April 2018 (12.04.2018) US  
62/688,913 22 June 2018 (22.06.2018) US

(71) Applicant: AVAVA, INC. [US/US]; 185 Cambridge Street, Second Floor, Boston, MA 02114 (US).

(72) Inventors: DRESSER, Charles, Holland; 185 Cambridge Street, Second Floor, Boston, MA 02114 (US).  
BHAWALKAR, Jayant; 185 Cambridge Street, Second Floor, Boston, MA 02114 (US).  
KATKAM, Rajender; 185 Cambridge Street, Second Floor, Boston, MA 02114 (US).

(74) Agent: MATTONI, Michael, A. et al.; Mintz Levin Cohn Ferris Glovsky and Popeo, P.C., One Financial Center, Boston, MA 02111 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: DIFFRACTIVE OPTICS FOR EMR-BASED TISSUE TREATMENT




FIG. 11

(57) Abstract: A method includes generating a plurality of primary beams from a laser beam, and generating, from a primary beam one or more secondary beams. The method also includes focusing the first secondary beam to a first focal region in the target tissue and the second secondary beam to a second focal region in the target tissue. The first focal region and the second focal region can be located at different depths in the target tissue.

[Continued on next page]

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
KM, ML, MR, NE, SN, TD, TG).

**Published:**

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments (Rule 48.2(h))*