

(Piperidinylalkyl)quinazoline derivatives, process for their preparation and pharmaceutical compositions containing them

Background of the invention

There are known in the art a number of pharmacologically active quinazolinones which are substituted on their heterocyclic ring with a piperidinylalkyl side chain. Such compounds may be found in the following references:

US. Pat. Nos. 3,322,766; 3,528,982; 3,635,976; 3,812,257; 3,865,827; 4,096,144; and 4,099,002;

Fr. Pat. No. 1,431,815; and


J. Med. Chem., 8, 807 (1965).

The compounds of the present invention differ from the foregoing essentially by the presence of particular substituent groups on the piperidine ring of the piperidinyl-alkyl side chain.

In U.S. Pat. No. 4,035,369 there are described a series of 1-(benzazolylalkyl)-4-substituted piperidines, from which the subject compound of this invention differ essentially by the replacement of the benzazole group by a quinazoline group.

Description of the preferred embodiments:


This invention is concerned with a novel series of quinazoline derivatives which are structurally represented by the formula



and the pharmaceutically acceptable acid addition salts thereof, wherein:

Ar is an aryl radical;

X is a member selected from the group consisting of



wherein said R_q is hydrogen or lower alkyl and said q is the integer 2 or 3;

R is a member selected from the group consisting of hydrogen, hydroxy and lower alkyl;

Alk is an alkylene chain having from 1 to 4 carbon atoms; and

Q is a quinazolinyl radical, the 1-, 2-, 3- or 4-position of which is connected with the alkylene side chain, said quinazolinyl radical bearing in one or both of its 2- and 4-positions a carbonyl or thiocarbonyl group, wherein the benzene ring of said quinazolinyl radical is optionally substituted with 1 to 3 substituents each independently selected from the group consisting of halo, lower alkyl, lower alkyloxy, trifluoromethyl, nitro and cyano, and wherein the pyrimidino ring of said quinazolinyl radical may be partly or fully saturated, said pyrimidino ring being optionally substituted with 1 to 3 substituents independently selected from the group consisting of lower alkyl, aryl and aryl(lower alkyl);

wherein said aryl as used in the definition of said Ar and of said Q is a member selected from the group consisting of phenyl, substituted phenyl, thienyl and pyridinyl, wherein said substituted phenyl has from 1 to 3 substituents each independently selected from the group consisting of halo, lower alkyl, lower alkyloxy, trifluoromethyl and amino.

As used in the foregoing definitions the term "lower alkyl" is meant to include straight and branched hydrocarbon radicals having from 1 to 6 carbon atoms, such as, for example, methyl, ethyl, 1-methylethyl, 1,1-dimethylethyl, propyl, butyl, pentyl, hexyl and the like; "alkylene" as used in the definition of Alk comprises straight and branched alkylene chains having from 1 to 4 carbon atoms; and the term "halo" is generic to fluoro, chloro, bromo and iodo.

Examples of quinazolinyl radicals within the scope of Q are optionally substituted 1,4 - dihydro - 2,4 - dioxo - 3(2H) - quinazolinyl, 3,4 - dihydro - 2,4 - dioxo - 1(2H) - quinazolinyl, 3,4 -


dihydro - 4 - oxo - 1(2H) - quinazolinyl, 1,2,3,4 - tetrahydro - 4 - oxo - 2 - quinazolinyl, 1,4 - dihydro - 4 - oxo - 3(2H) - quinazolinyl and the like radicals.

Preferred compounds within the scope of formula (I) are those wherein X represents CO. Particularly preferred compounds are those wherein X is CO and Alk represents an 1,2-ethanediyl radical. The compound 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,4 - (1H,3H) - quinazolinedione is especially preferred.

The compounds of formula (I) have basic properties and, consequently, they may be converted to their therapeutically active non-toxic acid addition salt forms by treatment with appropriate acids, such as, for example, inorganic acids, such as hydrohalic acid, e.g., hydrochloric, hydrobromic and the like, and sulfuric acid, nitric acid, phosphoric acid and the like; or organic acids, such as, for example, acetic, propanoic, hydroxyacetic, 2-hydroxypropanoic, 2-oxopropanoic, propanedioic, butanedioic, (Z)-2-butenedioic, (E)-2-butenedioic, 2-hydroxybutanedioic, 2,3-dihydroxybutanedioic, 2-hydroxy-1,2,3-propanetricarboxylic, benzoic, 3-phenyl-2-propenoic, α -hydroxybenzeneacetic, methanesulfonic, ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxybenzoic and the like acids.


Conversely the salt form can be converted by treatment with alkali into the free base form.

The compounds of formula (I) can generally be prepared by the reaction of an appropriate reactive ester of the formula (II), wherein Q and Alk are as previously defined and W is a reactive ester residue such as, for example, halo, particularly chloro, bromo or iodo, or a sulfonyloxy radical such as methylsulfonyloxy, 4-methylphenylsulfonyloxy and the like, with an appropriate piperidine derivative of the formula (III) wherein R, X and Ar have the previously defined meanings.



The foregoing reaction may be carried out following standard N-alkylating procedures. Said reaction is preferably carried out in an appropriate reaction-inert organic solvent such as, for example, a lower alkanol, e.g., methanol, ethanol, propanol, butanol and the like alkanols; an aromatic hydrocarbon, e.g., benzene, methylbenzene, dimethylbenzene, and the like; an ether, e.g., 1,4-dioxane, 1,1'-oxybispropane and the like; a ketone, e.g., 4-methyl-2-pentanone; N,N-dimethylformamide; nitrobenzene; and the like. The addition of an appropriate base such as, for example, an alkali or earth alkaline metal carbonate or hydrogen carbonate, may be utilized to pick up the acid which is liberated during the course of the reaction. A small amount of an appropriate metal iodide, e.g., sodium or potassium iodide may be added as a reaction promoter. Somewhat elevated temperatures are appropriate to enhance the rate of the reaction and preferably the reaction is carried out at the reflux temperature of the reaction mixture.


The compounds of formula (I) may also be prepared starting from a corresponding intermediate of the formula (IV)



wherein Alk, R, X and Ar are as previously defined and Q' is an appropriate precursor of Q, by converting said Q' into said Q following art-known procedures.

Such procedures are described, for example, in "The Chemistry of Heterocyclic Compounds — Fused Pyrimidines; Part I: Quinazolines", Ed. D. J. Brown, Interscience Publishers; New York, London, Sidney (1967), p. 74—102 and p. 116—128. Examples of such conversions of Q' into Q are given hereafter.


In order to simplify the structural representation of the compounds of formula (I) and of certain precursors and intermediates thereof the appropriately substituted piperidinylalkyl radical of the formula



will hereinafter be represented by the symbol D.

The compounds of formula (I) wherein Q represents a 1,2,3,4-tetrahydro 4-oxo- or 4-thioxo-quinazolinyl radical, said compounds being represented by the formula (I-a), can be prepared by cyclizing an appropriately substituted 2-aminobenzamide or -benzenethioamide of formula (IV-a) with an appropriately substituted carbonyl or thiocarbonyl compound of formula (V).

In the following reaction-equation one of R^2 , R^3 , R^4 and R^5 is D, each of the remaining being independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryllower alkyl; R^1 is selected from the group consisting of halo, lower alkyl, lower alkyloxy, trifluoromethyl, nitro and cyano; each Y is O or S; and n is an integer of from 0 to 3 inclusive.




Said cyclization-reaction may conveniently be carried out by stirring the reactants together in the presence of an appropriate solvent such as, for example, a lower alkanol, e.g., methanol, ethanol, propanol and the like. Somewhat elevated temperature and the addition of a catalytic amount of an appropriate strong acid, e.g., 4-methylbenzenesulfonic acid, hydrochloric acid and the like, may be used to enhance the rate of the reaction. Most preferably the reaction is carried out at the reflux temperature of the reaction mixture.

In the foregoing reaction the reagent of formula (V) may be replaced by an appropriate functionally equivalent derivative thereof such as, for example, a di(lower alkyl)acetal; a cyclic acetal and the like.

The compounds of formula (I) wherein Q represents a 3,4-dihydro-4-oxo-2-quinazolinyl radical, a 4-oxo-3(4H)-quinazolinyl radical or a corresponding thioxo analog thereof, said compounds being represented by the formula (I-b), can be derived from an appropriately substituted 2-aminobenzamide or -benzenethioamide of formula (IV-a) wherein R^3 is hydrogen, (IV-a-1), by converting (IV-a-1) into the corresponding 2-(acylamino)benzamide or -benzenethioamide of formula (VI) and cyclizing the latter following art-known cyclizing procedures.

In the following reaction-equations either of R^2 and R^4 is D, the other being a member selected from the group consisting of hydrogen, lower alkyl, aryl and aryllower alkyl.



The foregoing cyclization reaction may be carried out by stirring and heating, preferably refluxing, the 2-(acylamino)benzamide or -benzenethioamide (VI) in a suitable reaction-inert solvent such as, for example, a lower alkanol, e.g., methanol, ethanol, 2-propanol and the like; an aliphatic or aromatic hydrocarbon, e.g., hexane, benzene, methylbenzene and the like; a halogenated hydrocarbon, e.g., trichloromethane and the like. In order to enhance the rate of the reaction there may be added an appropriate base such as, for example, an alkali or earth alkaline metal hydroxide, carbonate or hydrogencarbonate, e.g., sodium hydroxide, potassium carbonate, sodium hydrogen carbonate and the like.


The conversion of the 2-aminobenzamide or -benzenethioamide (IV-a-1) into the 2-(acylamino)-benzamide or -benzenethioamide (VI) may be carried out by reacting (IV-a-1) with an appropriate acylating agent derived from the acid



wherein R^4 has the previously defined meaning. Appropriate acylating agents include anhydride and acyl halides derived from (VII). The acid (VII) itself may optionally be used as an acylating agent. In the latter case it is appropriate to remove the water formed during the course of the reaction by azeotropic distillation.

The compounds of formula (I) wherein Q is a 1,4-dihydro-2,4-dioxo-3(2H)-quinazolinyl radical, a 3,4-dihydro-2,4-dioxo-1(2H)-quinazolinyl radical or a mono- or dithio derivative thereof, said compounds being represented by the formula (I-c), may be derived from an appropriate 2-amino-benzamide or -benzenethioamide of formula (IV-a), by cyclizing the latter with urea, thiourea or a functionally equivalent derivative thereof.


In the following reaction scheme one of R^2 and R^3 is D, the remaining being selected from the group consisting of hydrogen, lower alkyl, aryl and aryl lower alkyl, R^1 and n are as previously described and each Y is independently selected from the group consisting of O and S.



Said cyclization may be carried out by stirring the reactants together, preferably at an elevated temperature and, if desired, in the presence of an appropriate solvent such as, for example, an aromatic hydrocarbon, e.g., benzene and the like, a cyclic ether, e.g., tetrahydrofuran and the like. A suitable derivative of urea and thiourea are, for example, 1,1'-bis[1H-imidazol-1-yl]methanone and its corresponding thione.


The compounds of formula (I-c) may also be prepared by cyclizing an appropriately substituted compound of formula (IV-b) with an appropriate primary amine of the formula (VIII).

In the following reaction-scheme, Y, R^1 and n have the previously defined meanings, one of R^2 and R^3 is D, the other hydrogen, lower alkyl or aryl lower alkyl and R^6 and R^6_a each represent an appropriate leaving group such as, for example, lower alkyloxy, amino and mono- and di(lower alkyl)amino.




The compounds of formula (I-c) wherein R^3 is hydrogen and R^2 is D, said compounds being represented by the formula (I-c-1), can also be prepared by cyclizing an isocyanate or an isothiocyanate of formula (IV-c) with a primary amine of formula (VIII) wherein R^2 is D, (VIII-a).

In the following reaction-equation R^6 represents an appropriate leaving group and R^1 , n and Y are as previously described.



15 The compounds of formula (I-c) wherein R^3 is D and R^2 is hydrogen, said compounds being represented by the formula (I-c-2), can also be prepared by cyclizing an appropriately substituted benzenamine or benzenethioamine of formula (IV-d), wherein D , Y and R^6 are as previously described, with an alkali metal cyanate or thiocyanate, e.g., potassium cyanate, sodium thiocyanate and the like.



30 Said cyclization-reactions are conveniently carried out by stirring and heating the reactants together, optionally in a suitable reaction-inert solvent having a relatively high boiling point such as aliphatic and aromatic hydrocarbons, e.g., petroleum ether, dimethylbenzene and the like.

The compounds of formula (I) wherein Q is a 1,4-dihydro-2-oxo-3(2H)-quinazolinyl radical and wherein X is as previously described but other than

35

CO_2

40 said X being represented by X' and said compounds by the formula (I-d), may also be prepared by acylating an intermediate of formula (IV-e) wherein R^7 is hydrogen, lower alkyl, aryl or aryl lower alkyl and R^1 , n and R^3 are as previously described, with a trihalo-acetic or -ethanethioic acid, or an acid halide thereof, subsequently reacting the thus obtained acylate (IX) with an amine of formula (VIII-b) and cyclizing the thus obtained Schiff's base (X) in the presence of an appropriate reducing agent, e.g., sodium borohydride and the like.


45

50

55


60

65




The acylation is carried out following art-known acylating procedures. The Schiff's base is prepared by stirring and heating the intermediates (IX) and (VIII-b) together in the presence of a suitable solvent, such as, for example, an alcohol, e.g., methanol and the like. The reductive cyclization is carried out by stirring the Schiff's base (X) and the reducing agent, e.g., sodium borohydride and the like, together in the presence of a suitable solvent, e.g., N,N-dimethylformamide and the like. Sometimes external cooling may be appropriate.

The compounds of formula (I) may also be derived from a compound of formula (XI) wherein P represents a precursor of the corresponding Ar-X-radical by converting said P into the desired Ar-X following methods known in the art.




For example, the compounds of formula (I) wherein Ar-X- is an optionally substituted 2-amino-benzoyl radical, said compounds being represented by the formula (I-e), may be prepared by the oxidative cleavage of the double bond in the corresponding indole-derivative (XII) and subsequent

hydrolysis of the thus formed formamide (XIII). Said oxidative cleavage may be carried out by the reaction of (XII) with an appropriate oxidizing agent, such as, for example, sodium periodate in the presence of a catalytic amount of osmiumtetroxide in a suitable solvent, e.g. 1,4-dioxane and the like. The oxidation may equally well be carried out by bubbling ozonized oxygen through a solution of (XII) in acetic acid and subsequently decomposing the intermediately formed ozonide with water. The thus obtained (formylamino)phenylcarbonyl intermediate (XIII) is then converted into (I-e) by hydrolysis in acidic medium. In the following reaction-equations R^8 and R^9 are each independently selected from the group consisting of hydrogen, halo, lower alkyl, lower alkyloxy, trifluoromethyl and amino.



The compounds of formula (I) wherein X is a CHOH radical, (I-f), may generally be prepared starting from the corresponding carbonyl-derivatives, (I-g), by reducing the carbonyl group of the latter with an appropriate reducing agent, e.g. sodium borohydride, sodium cyanoborohydride and the like following art-known methodologies.




When, for example, sodium borohydride is used as a reducing agent the reaction may conveniently be carried out in alkaline aqueous medium, if desired, in admixture with a water-miscible organic solvent such as, for example, an alicyclic ether, e.g. tetrahydrofuran, 1,4-dioxane and the like; or a lower alkanol, e.g. methanol, propanol and the like.

The compounds of formula (I) wherein X represent a radical




wherein R_a has the previously defined meaning, said compounds being represented by the formula (I-h), may be derived from the corresponding alcohols (I-f) by acylating the latter with an appropriate acylating agent according to art-known procedures. Appropriate acylating agents which may be used for this purpose include lower alkanolic acids and acyl halides and anhydrides derived therefrom.



The compounds of formula (I) wherein X is a methylene radical, (I-j), may be derived from the

corresponding carbonyl derivatives (I-g) by the reduction of the carbonyl group of (I-g) to a methylene group, e.g. by the Clemmensen reduction, using amalgated zinc and hydrochloric acid, or by the Wolff-Kishner reduction, using hydrazine and alkali in a high-boiling polar solvent, such as, 1,2-ethanediol and the like.



The compounds of formula (I) wherein X represents a di(lower alkyl)ketal or a cyclic lower alkylene ketal, wherein the lower alkylene chain has from 2 to 4 carbon atoms, may be derived from the corresponding carbonyl compounds by subjecting the latter to a ketalization-reaction following methodologies generally known in the art. Cyclic lower alkylene ketals, for example, may be prepared following methodologies analogous to those described in *Synthesis*, 1974, (I) 23-26.

The compounds of formula (I) wherein X represents a radical of the formula




can easily be derived from the corresponding carbonyl compounds by reacting the latter with respectively hydroxylamine hydrochloride or hydrazine hydrochloride according art-known procedures of preparing oximes and hydrazones.

Certain of the intermediates and starting materials used in the foregoing preparations are known compounds, others may be prepared according to art-known methodologies of preparing similar compounds and some of them are novel and consequently their preparation will be described hereafter.


The intermediates of formula (II) can be prepared by converting the hydroxyl function of the corresponding alcohol (XIV) into a reactive leaving group, e.g., by reacting the alcohol (XIV) with thionyl chloride, sulfonyl chloride, phosphor pentabromide, phosphoryl chloride, methanesulfonyl chloride, 4-methylbenzenesulfonyl chloride and the like. The alcohols (XIV) used as starting materials herein, can be derived from an alcohol (XV) wherein Q' is a precursor of Q following similar procedures as previously described herein for the preparation of (I-a), (I-b), (I-c) and (I-d) starting from appropriate intermediates of formula (IV).

Alternatively, the intermediates of formula (II) can be derived from an alcohol (XV) by converting the hydroxyl function into a reactive leaving group and subsequently cyclizing the precursor Q' in the thus obtained intermediate (XVI) following the previously described procedures.

In the following reaction-scheme Q, Q', Alk and W have the previously defined meanings.



The intermediates of the formula (III) may be derived from an intermediate of formula (XVII) by eliminating the protective group Z following art-known procedures.




Said elimination of Z can be carried out following art-known procedures, depending upon the

nature of Z, e.g., by catalytically hydrogenating a benzyl analog of (XVII), in case Z is benzyl, or by hydrolyzing a lower alkyloxycarbonyl analog of formula (XVII) in acidic medium, in case Z is lower alkyloxycarbonyl.


The intermediates of formula (XVII) may be prepared following art-known procedures.

For example, the intermediates of formula (XVII) wherein X is CO, (XVII-a), may be prepared starting from an appropriately substituted 4-piperidinone (XVIII) and an appropriate arylmethylcyanide (XIX) as shown in the following reaction-scheme.



The reaction of (XVIII) and (XIX) can be carried out by stirring and, if desired, heating the reactants together in a suitable reaction-inert solvent and in the presence of an appropriate base such as, for example, an alkali metal alkanolate, e.g., sodium methanolate and the like. The catalytic hydrogenation of (XX) is desirably conducted in a suitable reaction-inert solvent such as, for example, a lower alkanol, e.g., methanol and the like, in the presence of an appropriate catalyst, e.g., palladium-on-charcoal and the like. In order to avoid hydrogenation of the nitrile group it may be appropriate to add an appropriate catalyst-poison, e.g., thiophene and the like. The oxidation of the thus obtained (XXI) can be carried out following art-known oxidative procedures, e.g., as described in *Journal of Organic Chemistry* 40, 267 (1975).

The intermediates of formula (XVII-a), wherein R is other than hydroxy, said R being represented by R' and said intermediates by (XVII-a-1), can also be prepared by the Grignard reaction of an appropriately substituted 4-piperidynyl magnesium halide (XXII) with an appropriate aryl cyanide (XXIII).




Said Grignard-reaction is carried out by stirring and, if desired heating the reactants together in the presence of a suitable reaction-inert solvent, e.g., 1,1'-oxybisethane, tetrahydrofuran and the like.

The intermediates of formula (III) wherein X is other than CO, (III-b), can be derived from the corresponding arylcarbonylpiperidines, (III-a), following the same procedures as described hereinbefore for the preparation of compounds (I) wherein X is other than


C=O, starting from (I-g).

The intermediates of formula (IV) can generally be prepared by N-alkylating a piperidine (III) with an appropriately substituted reactive ester (XVI) following art-known N-alkylating procedures, as


previously described herein for the preparation of compounds (I) starting from (II) and (III).




The intermediates of formula (IV-a-1) can also be prepared by reducing the corresponding nitro-derivatives (XXIV) following art-known nitro-to-amine reduction reactions as known in the art, e.g., by catalytically hydrogenating the nitro-derivatives (XXIV) in the presence of an appropriate catalyst, e.g., platinum-on-charcoal and the like, in the presence of a suitable reaction-inert solvent, e.g., a lower alkanol such as methanol and the like. A catalyst poison, e.g., thiophene and the like may be appropriate to avoid undesired reductions of other functional groups.



The intermediates of formula (IV-a) can generally be derived from the intermediates (IV-a-1) by N-alkylating the latter with an appropriate reactive ester $\text{R}^3\text{-W}$, (XXV), following art-known N-alkylating procedures.




The nitro-derivatives (XXIV) may themselves be prepared by N-alkylating an appropriately substituted 2-nitrobenzamide (XXVI) with an appropriate compound of the formula (XXVII) wherein W and W' are each a reactive leaving group, provided that W' has better leaving capacity than W. The thus obtained (XXVIII) is subsequently reacted with an appropriately substituted piperidine derivative (III).




Both N-alkylations are carried out following art-known procedures as previously described herein for the preparation of compounds (I) starting from (II) and (III).

The intermediates of formula (XII) may be prepared by N-alkylating a piperidine (XXIX), wherein R,

R^8 and R^9 are as previously defined, with an appropriate reactive ester of the formula (II) following standard N-alkylating procedures.



15 The starting compounds (XXIX) herein are described in Belg. Pat. No. 858,101 and can be prepared by condensating benzoyl chloride with an appropriately substituted pyridine (XXX) and an appropriately substituted 1H-indole (XXXI), subsequently reducing the thus obtained dihydro-pyridine (XXXII), e.g., by catalytically hydrogenating the latter in the presence of an appropriate catalyst, such as, for example, palladium-on-charcoal and the like, and hydrolyzing the benzoyl derivative (XXXIII) in alkaline medium.



The compounds of formula (I); the intermediates of formula (XII) and their pharmaceutically active acid addition salts have useful pharmacological properties. They are very potent serotonin-antagonists and as such they can be used in the treatment of a variety of diseases in which serotonin release is of predominant importance.

The potency of the subject compounds as serotonin-antagonists is clearly evidenced by the results obtained in the following tests wherein the antagonistic activity of the compounds (I) on the effect of serotonin is examined.

Test 1

Antagonistic activity on the effect of serotonin on the caudal artery of the rat.

Caudal arteries from fasted male rats (210—235 g) are used in the test. Two helical strips having a length of 5—6 cm and a width of 2 mm are obtained from each artery and mounted vertically in a 100 ml organ bath containing an oxygenated Krebs-Henseleit solution. Submaximal contractions of the arterial strips are produced by adding single doses of serotonin (40 ng/ml) to the organ bath for 2 minutes with each time an interval of 10 minutes. The amplitude of the contraction is measured before and 5 minutes after adding the drug. After washing out, the agonist is added again three times in order to see whether the contraction is restored and normalized.

The first column of table 1 shows the ED_{50} -values in ng/ml for a number of compounds of formula (I) in the above test. In this connection the ED_{50} -values are the minimal concentrations of the concerned drugs which reduce the amplitude of the contraction to at least 50% of its normal value.

Test 2

Effects in gastric lesion tests.

a. Lesions induced by compound 48/80:

Compound 48/80 (a mixture of oligomers obtained by condensation of 4-methoxy-N-methyl-benzenethanamine and formaldehyde) is a potent releaser of vasoactive amines from endogenous stores such as, for example, histamine and serotonin. Rats injected with compound 48/80 exhibit consistent changes of blood flow in different vascular beds: cyanosis of the ears and the extremities are prominent within five minutes after injection of the compound; the rats die from shock within 30 minutes. The shock, followed by dead, can be avoided if the rats are pretreated with a classical H₁ antagonist. However the stimulatory effects on gastric secretion are not suppressed so that rats treated with compound 48/80 and protected from shock by an H₁ antagonist may exhibit all signs of intensive gastric gland activity: gross autopsy shows distended stomachs with abnormal contents and rough bright red patches all over the mucosa, corresponding to areas of disintegrated glands. A number of known serotonin antagonists such as, for example, methylsergide, cyproheptadine, cinanserin, mianserin, pipamperone, spiperone, pizotifen and metergoline, prevent completely the cyanosis of ears and extremities as well as the lesions in the glandular area of the stomach and the abnormal gastric distension.

b. Method:


Male rats of a Wistar inbred strain, weighing 220—250 g, are starved overnight, water being available ad libitum. The test compounds are administered orally as a solution or as a suspension in aqueous medium. A control rat and a "blank" rat receive the test compound. One hour later 5 - [4 - (diphenylmethyl) - 1 - piperazinylmethyl] - 1 - methyl - 1H - benzimidazole - 2 - methanol is administered subcutaneously to all rats at the dose of 2.5 mg/kg. Two hours after the oral administration of the test compound the compound 48/80 (freshly solved in water at a concentration of 0.25 mg/ml) is injected intravenously into all rats (dose: 1 mg/kg) except the "blank" rats. Five minutes after the injection the intensity of purple-blue coloration (cyanosis) of the extremities is scored as 0 (absent), + (moderate) or ++ (intense). Four hours after the intravenous injection of compound 48/80 the rats are decapitated and the stomachs are removed. Subsequently the stomachs are inspected for distension and contents (blood, fluid, food) and thoroughly rinsed. The macroscopic lesions are scored from 0 to +++, 0 corresponding to complete absence of visible lesions and the highest score corresponding to reddish rough patches covering more than half the glandular area.

The second column of table 1 shows for a number of compounds of formula (I) the doses (in mg/kg body weight) at which the distension of the stomach as well as the lesions in the glandular area of the stomach are completely absent in 50% of the test rats (ED_{50} -values).


The third column of table 1 shows for a number of compounds of formula (I) the doses (in mg/kg body weight) at which the cyanosis of ears and extremities is completely absent in 50% of the test rats (ED_{50} -values).

The compounds listed in table 1 are not given for the purpose of limiting the invention thereto but only to exemplify the useful pharmacological activities of all the compounds within the scope of formula (I).


TABLE 1




(R') _n	R ₃	Y	Alk	R	X	Ar	Caudal artery ng/ml	G. lesion ED ₅₀ in mg/kg	Cyanosis ED ₅₀ in mg/kg
—	H	O	—(CH ₂) ₃ —	—	—CO—	4-F—C ₆ H ₄	1.25	2.5	—
—	H	O	—(CH ₂) ₂ —	—	—CO—	4-F—C ₆ H ₄	0.7	0.16	0.08
—	H	O	—(CH ₂) ₂ —	—	—CO—	4-Cl—C ₆ H ₄	2.5	—	—
—	H	O	—(CH ₂) ₂ —	—	—CO—	4-OCH ₃ —C ₆ H ₄	2.5	—	—
—	H	S	—(CH ₂) ₃ —	—	—CO—	4-F—C ₆ H ₄	2	2.5	0.63
—	H	S	—(CH ₂) ₂ —	—	—CO—	4-F—C ₆ H ₄	0.6	0.08	0.04
7-Cl	H	O	—(CH ₂) ₂ —	—	—CO—	4-F—C ₆ H ₄	2	1.25	1.25
—	CH ₃	O	—(CH ₂) ₂ —	—	—CO—	4-F—C ₆ H ₄	0.7	1.25	0.16
—	H	O	—(CH ₂) ₂ —	—	—CO—	2-NH ₂ , 4-F—C ₆ H ₃	0.6	1.25	0.16
7-F	H	O	—(CH ₂) ₂ —	—	—CO—	4-F—C ₆ H ₄	0.7	1.25	0.16
7-OCH ₃	H	O	—(CH ₂) ₂ —	—	—CO—	4-F—C ₆ H ₄	0.7	0.31	0.16
6-CH ₃	H	O	—(CH ₂) ₂ —	—	—CO—	4-F—C ₆ H ₄	1.3	2.5	0.63
—	H	O	—(CH ₂) ₂ —	—	—CHOH—	4-F—C ₆ H ₄	20	1.25	1.25
—	H	O	—(CH ₂) ₂ —	—	—C—	4-F—C ₆ H ₄	11	0.31	0.16




(R ¹) _n	R ₄	Y	Alk	R	X	Ar	Caudal artery ng/ml	G. lesion ED ₅₀ in mg/kg	Cyanosis ED ₅₀ in mg/kg
—	H	O	—(CH ₂) ₂ —	—	—CO—	4-F-C ₆ H ₄	0.6	0.31	0.31
—	H	O	—(CH ₂) ₃ —	—	—CO—	4-F-C ₆ H ₄	5	2.5	—
—	CH ₃	O	—(CH ₂) ₂ —	—	—CO—	4-F-C ₆ H ₄	0.6	0.16	0.16




(R ¹) _n	R ₂	R ₃	R ₄	Alk	X	Ar	Caudal artery ng/ml	G. lesion ED ₅₀ in mg/kg	Cyanosis ED ₅₀ in mg/kg
—	H	H	H	—(CH ₂) ₂ —	CO	4-F-C ₆ H ₄	0.7	2.5	1.25



(R ¹) _n	R ₃	R ₄	R ₅	Alk	X	Ar	Caudal artery ng/ml	G. lesion ED ₅₀ in mg/kg	Cyanosis ED ₅₀ in mg/kg
—	H	H	H	—(CH ₂) ₂ —	—	4-F-C ₆ H ₄	5	≥2.5	—
—	H	H	H	—(CH ₂) ₃ —	—	4-F-C ₆ H ₄	10	≥2.5	—



(R ¹) _n	R ⁶	Alk	Ar	Caudal artery ng/ml	G. lesion ED ₅₀ in mg/kg	Cyanosis ED ₅₀ in mg/kg
—	C ₆ H ₅	(CH ₂) ₂	4-F-C ₆ H ₄	0.6	0.08	0.08
6-Cl	C ₆ H ₅	(CH ₂) ₂	4-F-C ₆ H ₄	—	0.63	0.63



(R ¹) _n	R ³	R ⁴	R ⁵	Alk	Caudal artery ng/ml	G. lesion ED ₅₀ in mg/kg	Cyanosis ED ₅₀ in mg/kg
—	H	H	H	(CH ₂) ₂	0.7	0.31	0.16

The compounds of formula (I) and the intermediates of formula (XII) prevent completely the lesions which are caused by excessive serotonin release and they also block the serotonin-induced contractions of bronchial tissues and of blood vessels, arteries as well as veins, and, consequently, the compounds of the present invention can be used in the treatment of gastrointestinal ulcer, bronchial spasm, hemorrhoids, varises and the like diseases, all of which are caused by congestion.

In view of their useful anti-congestive properties, the subject compounds may be formulated into various pharmaceutical forms for administration purposes. To prepare the pharmaceutical compositions of this invention, an effective anti-congestive amount of the particular compound, in base or acid-addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, preferably, for administration orally, rectally or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example to aid solubility, may be included. Injectable solutions, for example may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Acid addition salts of (I), due to their increased water solubility over the corresponding base form, are obviously more suitable in the preparation of aqueous compositions.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages; each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

Although the amount of the active ingredient to be administered may vary within rather wide limits depending on the particular circumstances, such as the nature and the severity of the disease, doses of from about 0.04 to about 4 mg of active ingredient per kg of body weight, and particularly from about 0.1 to about 2 mg per kg of body weight, administered once or repeatedly, are in general satisfactory.

The following formulations exemplify typical anti-convulsant pharmaceutical compositions in dosage unit form suitable for systemic administration to animal and human subjects in accordance with the present invention. These examples are given to illustrate and not to limit the scope of the present invention.

Oral drops.

The following formulation provides 50 liters of an oral-drop solution comprising 10 mg of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidiny] - ethyl] - 2,4(1H,3H) - quinazolinedione as the active ingredients (A.I.) per milliliter.

	A.I.	500 grams
50	2-hydroxypropanoic acid	0.5 liters
	sodium saccharin	1750 grams
55	cocoa flavor	2.5 liters
	purified water	2.5 liters
60	polyethylene glycol q.s. ad	50 liters

The A.I. is dissolved in the 2-hydroxypropanoic acid and 1.5 liters of the polyethylene glycol at 60—80°C. After cooling to 30—40°C there are added 35 liters of polyethylene glycol and the mixture is stirred well. Then there is added a solution of the sodium saccharin in 2.5 liters of purified water and while stirring there are added the cocoa flavor and polyethylene glycol q.s. ad volume. The resulting solution is filled into suitable containers.

0013612

Oral solution.

The following formulation provides 20 liters of an oral solution comprising 20 mg of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinedione as the active ingredient (A.I.) per teaspoonful (5 milliliters).

5	A.I.	20	grams
	2,3-dihydroxybutanedioic acid	10	grams
10	sodium saccharin	40	grams
	1,2,3-propanetriol	12	liters
	Sorbitol 70% solution	3	liters
15	Methyl 4-hydroxybenzoate	9	grams
	Propyl 4-hydroxybenzoate	1	gram
20	Raspberry essence	2	milliliters
	Gooseberry essence	2	milliliters
	Purified water q.s. ad	20	liters.

25 The methyl and propyl 4-hydroxybenzoates are dissolved in 4 liters of boiling purified water. In 3 liters of this solution are dissolved first the 2,3-dihydroxybutanedioic acid and thereafter the A.I.. The latter solution is combined with the remaining part of the former solution and the 1,2,3-propanetriol and the sorbitol solution are added thereto. The sodium saccharine is dissolved in 0.5 liters of water and the raspberry and gooseberry essences are added. The latter solution is combined with the former, 30 water is added q.s. ad volume and the resulting solution is filled in suitable containers.

Capsules:

The following formulation provides 1000 capsules comprising each 20 mg of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinedione as the active ingredient (A.I.). 35

	A.I.	20	grams
40	Sodium lauryl sulfate	6	grams
	Starch	56	grams
	Lactose	56	grams
45	Colloidal silicone dioxide	0.8	grams
	Magnesium stearate	1.2	grams

50 The composition is prepared by stirring the ingredients vigorously together. The resulting mixture is subsequently filled into suitable hardened gelatine capsules.

Film-coated tablets:

10,000 compressed tablets, each containing as the active ingredient 10 mg of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinedione, are prepared from the 55 following formulation:

Tablet core:

	A.I.	100	grams
	Lactose	570	grams
5	Starch	200	grams
	Polyvinylpyrrolidone (Kollidon K 90)	10	grams
10	Microcrystalline cellulose (Avicel)		
	Sodium dodecyl sulfate	5	grams
	Hydrogenated vegetable oil (Sterotex)	15	grams
15	Coating:		
	Methyl cellulose (Methocel 60 HG)	10	grams
20	Ethyl cellulose (Ethocel 22 cps)	5	grams
	1,2,3-propanetriol	2.5	milliliters
	Polyethylene glycol 6000	10	grams
25	Concentrated colour suspension (Opaspray K-1-2109)	30	milliliters
	Polyvinylpyrrolidone (Povidone)	5	grams
30	Magnesium octadecanoate	2.5	grams

Preparation of tablet core:

A mixture of the A.I., the lactose and the starch is mixed well and thereafter humidified with a solution of the sodium dodecyl sulfate and the polyvinylpyrrolidone in about 200 milliliters of water. The wet powder mixture is sieved, dried and sieved again. Then there is added the microcrystalline cellulose and the hydrogenated vegetable oil. The whole is mixed well and compressed into tablets.

Coating:

To a solution of the methyl cellulose in 75 milliliters of denatured ethanol there is added a solution of the ethyl cellulose in 150 milliliters of dichloromethane. Then there are added 75 milliliters of dichloromethane and the 1,2,3-propanetriol. The polyethylene glycol is molten and dissolved in 75 milliliters of dichloromethane. The latter solution is added to the former and then there are added the magnesium octadecanoate, the polyvinylpyrrolidone and the concentrated colour suspension and the whole is homogenated.

The tablet cores are coated with the thus obtained mixture in a coating apparatus.

Injectable solution:

The following formulation provides 1 liter of a parenteral solution comprising 4 mg of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinedione as the active ingredient per milliliter.

	A.I.	4	grams
55	Lactic acid	4	grams
	Propylene glycol	0.05	grams
	Methyl 4-hydroxybenzoate	1.8	grams
60	Propyl 4-hydroxybenzoate	0.2	grams
	Purified water q.s. ad	1	liter.

The methyl and propyl 4-hydroxybenzoates are dissolved in about 0.5 liters of boiling water for injection. After cooling to about 50°C there are added while stirring the lactic acid, the propylene glycol and the A.I.. The solution is cooled to room temperature and supplemented with water for injection q.s. ad volume. The solution is sterilized by filtration (U.S.P. XVII p. 811) and filled in sterile containers.

Suppositories:

100 Suppositories each containing 20 mg of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidiny]ethyl] - 2,4(1H,3H) - quinazolinedione as the active ingredient are prepared from the following formulations:

A.I.	3 grams
2,3-Dihydroxybutanedioic acid	3 grams
Polyethylene glycol 400	25 milliliters
Surfactant (Span)	12 grams
Triglycerides (Witepsol 555) q.s. ad	300 grams.

The A.I. is dissolved in a solution of the 2,3-dihydroxybutanedioic acid in the polyethylene glycol 400. The surfactant and the triglycerides are molten together. The latter mixture is mixed well with the former solution. The thus obtained mixture is poured into moulds at a temperature of 37—38°C to form the suppositories.

In view of the anti-congestive activity of the subject compounds, it is evident that the present invention provides a method of treating congestive diseases of warm-blooded animals by the systemic administration of an effective anti-congestive amount of a compound of formula (I) or of an intermediate of formula (XII) or a pharmaceutically acceptable acid addition salt thereof in admixture with a pharmaceutical carrier.

The following examples are intended to illustrate but not to limit the scope of the present invention. Unless otherwise stated all parts herein are by weight and all temperatures are in the centigrade scale.

A) Preparation of Intermediates.

Example I

To 5 parts of magnesium are added 2.18 parts of 1,2-dibromoethane and a small amount of iodine to initiate the reaction. Then there is added dropwise a solution of 28 parts of 4-chloro-1-methylpiperidine in 180 parts of tetrahydrofuran while the mixture is heated to 70°C. After cooling, there is added dropwise a solution of 14 parts of 3-methylbenzonitrile in 90 parts of tetrahydrofuran. Upon completion, stirring is continued for 1 hour at reflux temperature. The reaction mixture is cooled and poured onto a solution of 75 parts of ammonium chloride in water. The product is extracted with 2,2'-oxybispropane. The extract is washed with water, dried, filtered and evaporated, yielding 35 parts of (3-methylphenyl) (1-methyl-4-piperidinyl)methanone as an oily residue.

Following the same procedure and using equivalent amounts of the appropriately substituted benzonitriles and 4-chloro-1-methylpiperidine there are also prepared: (4-bromophenyl) (1-methyl-4-piperidinyl)methanone as a residue; and (2-chlorophenyl) (1-methyl-4-piperidinyl)methanone as an oily residue.

Example II

To 7 parts of magnesium is added dropwise a solution of 50 parts of 1-bromo-2-methylbenzene in 140 parts of 1,1'-oxybisethane so that the mixture is refluxing. The whole is stirred for 15 minutes at reflux. The Grignard-complex is cooled to 10°C and there is added dropwise a solution of 30 parts of 1-(phenylmethyl)-4-piperidinecarbonitrile in 70 parts of 1,1'-oxybisethane. Upon completion, stirring is continued for 4 hours at room temperature. The reaction mixture is decomposed with a solution of 40 parts of ammonium chloride in 400 parts of water. The organic phase is separated, dried, filtered and evaporated, yielding 31 parts of (2-methylphenyl) [1-(phenylmethyl)-4-piperidinyl]methanone as an oily residue.

Example III

To a stirred mixture of 35 parts of (3-methylphenyl) (1-methyl-4-piperidinyl)methanone, 1 part of sodium carbonate and 225 parts of dimethylbenzene are added dropwise 22 parts of ethyl carbonylchloride at 20°C. Upon completion, stirring is continued for 6 hours at reflux temperature. The reaction mixture is evaporated, yielding 12 parts of ethyl 4-(3-methylbenzoyl)-1-piperidinecarboxylate as an oily residue.

In a similar manner there are also prepared starting from the corresponding phenyl(1-methyl or 1-phenylmethyl-4-piperidiny)methanone:

ethyl 4-(4-bromobenzoyl)-1-piperidinecarboxylate as a residue;

ethyl 4-(2-chlorobenzoyl)-1-piperidinecarboxylate as an oily residue; and

ethyl 4-(2-methylbenzoyl)-1-piperidinecarboxylate as an oily residue.

Example IV

A mixture of 12 parts of ethyl 4-(3-methylbenzoyl)-1-piperidinecarboxylate and 225 parts of a hydrobromic acid solution 48% in water is stirred and refluxed for 3 hours. The reaction mixture is evaporated and the residue is crystallized from 2-propanol, yielding 7.5 parts of (3-methylphenyl) (4-piperidiny)methanone hydro-bromide.

Following the same hydrolysis-procedure there are also prepared:

(4-bromophenyl) (4-piperidiny)methanone hydrobromide;

(2-chlorophenyl) (4-piperidiny)methanone hydrobromide; mp. 200°C; and

(2-methylphenyl) (4-piperidiny)methanone hydrobromide.

Example V

A mixture of 25 parts of 2-chloropropanenitrile, 61 parts of (4-fluorophenyl) (4-piperidiny)methanone hydrochloride, 63 parts of sodium carbonate and 160 parts of acetonitrile is stirred and refluxed (100°C) overnight. The reaction mixture is cooled, water is added and the layers are separated. The aqueous phase is extracted with 4-methyl-2-pentanone. The combined organic phases are dried, filtered and evaporated. The residue is purified by column-chromatography over silica gel using a mixture of trichloromethane, hexane and methanol (50:49:1 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The solid residue is crystallized from 2,2'-oxybispropane. The product is filtered off, and dried yielding 17 parts of 4-(4-fluorobenzoyl)- α -methyl-1-piperidine-acetonitrile; mp. 126.7°C.

A mixture of 32 parts of 4-(4-fluorobenzoyl)- α -methyl-1-piperidineacetonitrile and 400 parts of methanol, saturated with ammonia is hydrogenated in the Parr-apparatus at 25°C with 5 parts of Raney-nickel catalyst. After the calculated amount of hydrogen is taken up, the catalyst is filtered off and the filtrate is evaporated. The residue is taken up in methylbenzene and the latter is evaporated again. The residue is dissolved in 2,2'-oxybispropane and the solution is filtered till clear. The solvent is evaporated, yielding 32 parts (100%) of [1-(2-amino-1-methylethyl)-4-piperidiny] (4-fluorophenyl)methanone as an oily residue.

Example VI

A mixture of 4.5 parts of 1-chloro-3,3-diethoxypropane, 12.15 parts of (4-fluorophenyl) (4-piperidiny)methanone hydrochloride, 10.6 parts of sodium carbonate and 120 parts of 4-methyl-2-pentanone is stirred and refluxed overnight. The reaction mixture is cooled, water is added and the layers are separated. The organic phase is dried, filtered and evaporated. The residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The residue is crystallized from a mixture of methylbenzene and 2,2'-oxybispropane, yielding 9 parts (53%) of [1-(3,3-diethoxypropyl)-4-piperidiny] (4-fluorophenyl)methanone.

Example VII

A mixture of 50 parts of 2-amino-3,5-dichlorobenzoic acid and 240 parts of ethanol is saturated with gaseous hydrogen chloride. The whole is stirred and refluxed for 10 hours. The reaction mixture is allowed to cool and the solvent is evaporated. To the solid residue are added water and sodium hydroxide. The precipitated product is filtered off and dried, yielding 25 parts of ethyl 2-amino-3,5-dichlorobenzoate.

To a stirred mixture of 25 parts of ethyl 2-amino-3,5-dichlorobenzoate and 180 parts of dimethylbenzene are added dropwise 30 parts of ethyl carbonochloridate. Upon completion, stirring is continued for 8 hours at reflux temperature. The reaction mixture is evaporated and the residue is crystallized from petroleumether. The product is filtered off and dried, yielding 30 parts of ethyl 3,5-dichloro-2-(ethoxycarbonylamino)benzoate.

Example VIII

A mixture of 40 parts of ethyl 4-chloro-2-[(2-ethoxycarbonyl)amino]benzoate and 10 parts of 2-aminoethanol is stirred and heated to 160—170°C while the formed ethanol is distilled off. After stirring for about 30 minutes, the mixture is cooled and 2-propanol is added. The solid product is filtered off and dried, yielding 23 parts (64%) of 7-chloro-3-(2-hydroxyethyl)-2,4-(1H,3H)-quinazoline-dione.

Following the same cyclization procedure and using equivalent amounts of the appropriately substituted 2-[(ethoxycarbonyl)amino]benzoates and 2-aminoethanol there are also prepared:

3-(2-hydroxyethyl)-1-methyl-2,4-(1H,3H)-quinazolinone;

6-chloro-3-(2-hydroxyethyl)-2,4(1H,3H)-quinazolinedione; and
 6,8-dichloro-3-(2-hydroxyethyl)-2,4(1H,3H)-quinazolinedione; and
 3-(2-hydroxyethyl)-6-methyl-2,4(1H,3H)-quinazolinedione.

Example IX

A mixture of 14 parts of 6-chloro-3,4-dihydro-3-(2-hydroxyethyl)-4-phenyl-2(1H)-quinazolinone, 5 parts of sodium acetate and 200 parts of methanol is hydrogenated at normal pressure and at room temperature with 2 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen is taken up, the catalyst is filtered off and the filtrate is evaporated. The residue is stirred in 2,2'-oxybispropane. The solid product is filtered off and stirred in 200 parts of water. It is filtered off again and dried, yielding 11.2 parts (91%) of 3,4-dihydro-3-(2-hydroxyethyl)-4-phenyl-2(1H)-quinazolinone; mp. 141°C.

Example X

A mixture of 23 parts of 7-chloro-3-(2-hydroxyethyl)-2,4(1H,3H)-quinazolinedione, 32 parts of thionyl chloride and 150 parts of trichloromethane is stirred and refluxed for 4 hours. The reaction mixture is cooled. The precipitated product is filtered off, washed with trichloromethane and with petroleum ether, and dried, yielding 22 parts (88%) of 7-chloro-3-(2-chloroethyl)-2,4(1H,3H)-quinazolinedione.

In a similar manner there are also prepared:
 3-(2-chloroethyl)-1-methyl-2,4(1H,3H)-quinazolinedione;
 3-(2-chloroethyl)-3,4-dihydro-4-phenyl-2(1H)-quinazolinone; mp. 179.5°C;
 6-chloro-3-(2-chloroethyl)-3,4-dihydro-4-phenyl-2(1H)-quinazolinone;
 6-chloro-3-(2-chloroethyl)-2,4(1H,3H)-quinazolinedione;
 6,8-dichloro-3-(2-chloroethyl)-2,4(1H,3H)-quinazolinedione; mp. 207°C; and
 3-(2-chloroethyl)-6-methyl-2,4(1H,3H)-quinazolinedione.

Example XI

A mixture of 13.6 parts of 2-aminobenzamide, 31.5 parts of 1-bromo-3-chloropropane, 21 parts of sodium carbonate and 200 parts of ethanol is stirred and refluxed over week-end. The reaction mixture is filtered and the filtrate is evaporated. The oily residue is stirred in methylbenzene. The mixture is filtered till clear and the solvent is evaporated. The oily residue is purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The solid residue is stirred in 2,2'-oxybispropane. The product is filtered off and dried, yielding 7 parts (33%) of 2-[(3-chloropropyl)amino]benzamide; mp. $\pm 100^\circ\text{C}$.

Example XII

A mixture of 10 parts of 2-[(3-chloropropyl)amino]benzamide, 16 parts of 2-propanone, 1 part of 4-methylbenzenesulfonic acid and 40 parts of ethanol is stirred and refluxed overnight. The reaction mixture is evaporated and the oily residue is stirred in 2,2'-oxybispropane. The product is filtered off and dried, yielding 10 parts of 1-(3-chloropropyl)-2,3-dihydro-2,2-dimethyl-4(1H)-quinazolinone.

Example XIII

To a stirred and cooled (ice-bath) solution of 27.5 parts of 2-bromoethanamine hydrobromide in 200 parts of water is added a solution of 28.9 parts of 4-methoxy-2-nitrobenzoyl chloride in 63 parts of benzene at 5—10°C. While stirring vigorously, there is added dropwise a solution of 10.8 parts of sodium hydroxide in 250 parts of water at 5—10°C. Upon completion, stirring is continued for 2 hours at this temperature. The supernatant phase is decanted and the oily residue is stirred in 2-propanol. The product is filtered off and dried, yielding 27.5 parts (68%) of N-(2-bromoethyl)-4-methoxy-2-nitrobenzamide; mp. 133.8°C.

Example XIV

To a stirred and cooled mixture of 25 parts of 2-aminoethanol and 135 parts of methylbenzene is added dropwise a solution of 29 parts of 4-fluoro-2-nitrobenzoyl chloride in methylbenzene. Upon completion, stirring is continued for 30 minutes at room temperature. The precipitated product is filtered off, taken up in water and extracted with 4-methyl-2-pentanone. The extract is dried, filtered and evaporated, yielding 28 parts (64%) of 4-fluoro-N-(2-hydroxyethyl)-2-nitrobenzamide as a residue.

A mixture of 28 parts of 4-fluoro-N-(2-hydroxyethyl)-2-nitrobenzamide, 40 parts of thionyl chloride and 150 parts of trichloromethane is stirred and refluxed for 2 hours. After cooling, the precipitated product is filtered off and dried, yielding 24.5 parts (83%) of N-(2-chloroethyl)-4-fluoro-2-nitrobenzamide.

Example XV

A mixture of 14.4 parts of N-(3-bromopropyl)-2-nitrobenzamide, 12.2 parts of (4-fluorophenyl)

(4-piperidinyl)methanone hydrochloride, 16 parts of sodium carbonate and 200 parts of 4-methyl-2-pentanone is stirred and refluxed for 3 hours using a water-separator. The reaction mixture is cooled, water is added and the layers are separated. The organic phase is dried, filtered and evaporated. The oily residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions are collected and the eluent is evaporated yielding 18 parts (87%) of N-[3-[4-(4-fluorobenzoyl)-1-piperidinyl]propyl]-2-nitrobenzamide as an oily residue.

Following the same procedure and using equivalent amounts of (4-fluorophenyl)(4-piperidinyl)-methanone hydrochloride and an appropriately substituted 2-nitrobenzamide there are also prepared: N-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]-2-nitrobenzamide mp. 150°C; 4-fluoro-N-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]-2-nitrobenzamide; mp. 163.5°C; and N-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]-4-methoxy-2-nitrobenzamide; mp. $\pm$ 134°C.

Example XVI

To 1 part of a solution of 2 parts of thiophene in 40 parts of ethanol are added 18 parts of N-[3-[4-(4-fluorobenzoyl)-1-piperidinyl]propyl]-2-nitrobenzamide and 200 parts of methanol. The whole is hydrogenated at normal pressure and at room temperature with 2 parts of platinum-on-charcoal catalyst 5%. After the calculated amount of hydrogen is taken up, the catalyst is filtered off and the filtrate is evaporated. The oily residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The oily residue is crystallized from 2-propanol. The product is filtered off and dried, yielding 7.5 parts (45%) of 2-amino-N-[3-[4-(4-fluorobenzoyl)-1-piperidinyl]propyl]benzamide; mp. 113.2°C.

The mother liquor is evaporated, yielding a second fraction of 6 parts (36%) of 2-amino-N-[3-[4-(4-fluorobenzoyl)-1-piperidinyl]propyl]benzamide as an oily residue.

Following the same hydrogenating procedure there are also prepared:

2-amino-N-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]benzamide; mp. 142.5°C

2-amino-4-fluoro-N-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]benzamide; mp. 124.1°C; and

2-amino-N-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]-4-methoxybenzamide; mp. 168.1°C.

Example XVII

A mixture of 7.2 parts of 2-[(4-chloro-1-oxobutyl)amino]benzamide, 7.3 parts of (4-fluorophenyl)(4-piperidinyl)methanone hydrochloride, 10.1 parts of N,N-diethylethanamine and 120 parts of acetonitrile is stirred and refluxed for 24 hours. The reaction mixture is evaporated and water is added to the residue. The oily product is extracted with trichloromethane. The extract is dried, filtered and evaporated, yielding 10 parts (81%) of N-[2-(aminocarbonyl)phenyl]-4-(4-fluorobenzoyl)-1-piperidinebutanamide as a residue.

Example XVIII

A. To a stirred mixture of 250 parts of pyridine and 39 parts of 6-fluoro-1H-indole are added dropwise 42 parts of benzoyl chloride at 22°C. Upon completion, stirring is continued for 2 hours at room temperature. The reaction mixture is diluted with water and a hydrochloric acid solution 2N is added. The product is extracted twice with 350 parts of 1,1'-oxybisethane. The combined extracts are dried, filtered and evaporated. The residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions are collected and the eluent is evaporated, yielding 95 parts of 1-benzoyl-4-(6-fluoro-1H-indol-3-yl)-1,4-dihydropyridine as an oily residue.

B. A mixture of 95 parts of 1-benzoyl-4-(6-fluoro-1H-indol-3-yl)-1,4-dihydropyridine and 540 parts of N,N-dimethylacetamide is hydrogenated in the Parr-apparatus with 5 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen is taken up, the catalyst is filtered off over Hyflo and the filtrate is poured onto water while stirring; an oil is precipitated. The supernatant aqueous phase is decanted and the residual oil is washed with water. The oily product is extracted with trichloromethane. The extract is dried, filtered and evaporated. The residue is washed successively with a mixture of petroleum ether and 2,2'-oxybispropane, with a hydrochloric acid solution 2N and with water, and dissolved in trichloromethane. The latter is washed with a sodium hydroxide solution 1N and with water, dried, filtered and evaporated. The residue is heated in a small amount of N,N-dimethylacetamide at 80°C and the mixture is allowed to stand over week-end. The solid product is filtered off, washed with a mixture of water and ethanol, and dried in vacuo at 80°C, yielding 25 parts of 1-benzoyl-4-(6-fluoro-1H-indol-3-yl)-piperidine; mp. 220°C.

C. A mixture of 24 parts of 1-benzoyl-4-(6-fluoro-1H-indol-3-yl)-piperidine, 70 parts of potassium hydroxide, 495 parts of 1,2-ethanediol and 80 parts of water is stirred and refluxed for 6 hours. The reaction mixture is cooled and 500 parts of water are added while stirring. The precipitated product is filtered off, washed with water and petroleum ether and dried in vacuo at 80°C, yielding 16 parts of 6-fluoro-3-(4-piperidinyl)-1H-indole; mp. 224°C.

D. A mixture of 7.5 parts of 3-(2-chloroethyl)-2,4(1H,3H)-quinazolinedione, 6.8 parts of 6-fluoro-

3-(4-piperidiny)-1H-indole, 10 parts of sodium carbonate, 0.1 parts of potassium iodide and 240 parts of 4-methyl-2-pentanone is stirred and refluxed overnight. Water is added and the layers are separated. The aqueous phase is set aside. The organic phase is dried, filtered and evaporated. The solid residue is boiled in 2-propanol. The product is filtered off and dried, yielding a first fraction of 4 parts of 3 - [2 - [4 - (6 - fluoro - 1H - indol - 3 - yl) - 1 - piperidiny]ethyl] - 2,4(1H,3H) - quina-

zolinone. From the aqueous phase (see above), a solid product is filtered off and boiled in 2-propanol. The product is filtered off and dried, yielding a second fraction of 3 parts of 3 - [2 - [4 - (6 - fluoro - 1H - indol - 3 - yl) - 1 - piperidiny]ethyl] - 2,4(1H,3H) - quiazolinone; mp. 300.6°C.

10 E. Ozonised oxygen is bubbled through a stirred mixture of 6 parts of 3 - [2 - [4 - (6 - fluoro - 1H - indol - 3 - yl) - 1 - piperidiny]ethyl] - 2,4 - (1H,3H) - quiazolinone and 100 parts of acetic acid till a clear solution is obtained. Then there are added 200 parts of crushed ice and 100 parts of water and the whole is alkalinized with a sodium hydroxide solution. The product is extracted with tri-
15 parts of N - [2 - [1 - [2 - (1,4 - dihydro - 2,4 - dioxo - 3(2H) - quiazolinyl)ethyl] - 4 - piperidiny]carbonyl] - 5 - fluorophenyl]formamide as a residue.

Example XIX

20 A mixture of 13.5 parts of 1-(2-aminophenyl)ethanone, 12.1 parts of N,N-diethylethanamine and 72 parts of benzene are stirred and cooled at 5—10°C. Then there is added dropwise a solution of 18.2 parts of trichloroacetyl chloride in 36 parts of benzene while still cooling at 1—10°C. Upon completion, stirring at that temperature is continued for 30 minutes. The reaction mixture is filtered, the filter-cake is washed with benzene and the filtrate is evaporated. The solid residue is boiled in methanol. After cooling, the product is filtered off and dried, yielding 19 parts (68%) of N-(2-acetylphenyl)-2,2,2-tri-
25 chloroacetamide; mp. 112.7°C.

A mixture of 60 parts of N-(2-acetylphenyl)-2,2,2-trichloroacetamide, 14 parts of 2-amino-ethanol and 200 parts of ethanol is stirred first for 5.50 hours at reflux and further overnight while the temperature is allowed to cool to room temperature. The formed precipitate is filtered off. The product is washed with 2,2'-oxybispropane and dried in vacuo, yielding 40 parts (59%) of 2,2,2 - trichloro -
30 N - [2 - [1 - [(2 - hydroxyethyl)imino]ethyl]phenyl]acetamide; mp. 182°C.

To 135 parts of N,N-dimethylformamide are added first 16.2 parts of 2,2,2 - trichloro - N - [2 - [1 - [(2 - hydroxyethyl)imino]ethyl]phenyl]acetamide and then portionwise 5.7 parts of sodium borohydride while stirring: exothermic reaction. Upon completion, stirring is continued overnight at room temperature. The reaction mixture is poured onto water and the product is extracted with tri-
35 chloromethane. The extract is dried, filtered and evaporated. The residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions are collected and the eluent is evaporated, yielding 8.5 parts of 3,4-dihydro-3-(2-hydroxyethyl)-4-methyl-2(1H)-quinazolinone as a solid residue.

A mixture of 8 parts of 3,4-dihydro-3-(2-hydroxyethyl)-4-methyl-2(1H)-quinazolinone, 8 parts of thionyl chloride and 90 parts of trichloromethane is stirred and refluxed for 1.50 hours. The reaction mixture is evaporated. The residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The residue is stirred in 2,2'-oxybispropane. The product is filtered off and dried, yielding 6.6 parts of 3-(2-chloroethyl)-3,4-dihydro-4-methyl-2(1H)-quinazolinone; mp. 126.4°C.
45

Example XX

To a stirred mixture of 10 parts of methyl 2-[(2-hydroxyethyl)amino]benzoate and 100 parts of acetic acid is added dropwise a solution of 4.5 parts of potassium cyanate in 25 parts of water. Upon completion, stirring at room temperature is continued overnight. The reaction mixture is evaporated. The solid residue is stirred with water. The product is filtered off, washed with water and ethanol, and dried, yielding 5 parts (48%) of 1-(2-hydroxyethyl)-2,4(1H,3H)-quinazolinone; mp. 273.6°C.
50

A mixture of 3.6 parts of 1-(2-hydroxyethyl)-2,4(1H,3H)-quinazolinone and 40 parts of thionyl chloride is stirred and refluxed overnight. The reaction mixture is allowed to cool to room temperature. The precipitated product is filtered off, washed with 2,2'-oxybispropane and dried, yielding 3.4 parts (86%) of 1-(2-chloroethyl)-2,4(1H,3H)-quinazolinone; mp. 215.3°C.
55

Following the same procedure and using equivalent amounts of the appropriate starting materials there are also prepared:

1-(3-hydroxypropyl)-2,4(1H,3H)-quinazolinone; mp. 240°C; and
1-(3-chloropropyl)-2,4(1H,3H)-quinazolinone; mp. 187.1°C.
60

Example XXI

To a stirred mixture of 18 parts of 2-phenyl-4(3H)-quinazolinone and 225 parts of N,N-dimethylformamide are added portionwise at room temperature 3.2 parts of sodium hydride dispersion 60% (exothermic reaction: the temperature rises to 34°C). After the addition is complete, the whole is stirred for 10 minutes whereafter 12.4 parts of 2-bromoethanol are added dropwise (slightly exothermic).
65

Upon completion, stirring is continued first for one hour at room temperature, then for 2 hours at 80°C and further overnight at room temperature. The reaction mixture is cooled, poured onto water and the solid product is sucked off. It is washed with water and 2,2'-oxybispropane and dried in vacuo at 60°C, yielding 15 parts of 3-(2-hydroxyethyl)-2-phenyl-4(3H)-quinazolinone.

To a stirred mixture of 15 parts of 3-(2-hydroxyethyl)-2-phenyl-4(3H)-quinazolinone and 375 parts of trichloromethane are added dropwise 24 parts of thionyl chloride at room temperature. Upon completion, stirring is continued for 2 hours at reflux. The reaction mixture is evaporated. Water is added to the residue and the whole is neutralized with a sodium hydrogen carbonate solution. The product is extracted with trichloromethane. The extract is washed with water, dried, filtered and evaporated. The residue is crystallized from a mixture of 2,2'-oxybispropane and petroleum ether. The product is filtered off and dried, yielding 11.6 parts of 3-(2-chloroethyl)-2-phenyl-4(3H)-quinazolinone.

A mixture of 5.5 parts of 3-(2-chloroethyl)-2-phenyl-4(3H)-quinazolinone, 3.2 parts of 3-(4-piperidinyl)-1H-indole, 7 parts of sodium carbonate, 0.1 parts of potassium iodide and 200 parts of 4-methyl-2-pentanone is stirred and refluxed for 22 hours. The reaction mixture is filtered over Hyflo and the filtrate is evaporated. The residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (92:8 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The residue is crystallized from a mixture of ethanol and 1,1'-oxybisethane, yielding 5 parts of 3-[2-[4-(1H-indol-3-yl)-1-piperidinylethyl]-2-phenyl-4(3H)-quinazolinone; mp. 191.3°C.

20

Example XXII

To a stirred mixture of 80 parts of sodium methoxide and 160 parts of methanol are added successively 50 parts of 2-thiopheneacetonitrile and then dropwise 66 parts of 1-(phenylmethyl)-4-piperidinone. Upon completion, the whole is heated to reflux and stirring at reflux temperature is continued for one hour. The reaction mixture is cooled and evaporated. The residue is distilled in a molecular distillation-apparatus, yielding about 70 parts of α -[1-(phenylmethyl)-4-piperidinylidene]-2-thiopheneacetonitrile as a residue.

A mixture of 70 parts of α -[1-(phenylmethyl)-4-piperidinylidene]-2-thiopheneacetonitrile in 800 parts of methanol is hydrogenated at normal pressure and at room temperature with 10 parts of palladium-on-charcoal catalyst 10%. Upon the calculated amount of hydrogen is taken up, the catalyst is filtered off and the filtrate is evaporated, yielding 70 parts of 1-(phenylmethyl)- α -(2-thienyl)-4-piperidineacetonitrile as a residue.

To a stirred mixture of 29.6 parts of 1-(phenylmethyl)- α -(2-thienyl)-4-piperidineacetonitrile in 100 parts of dimethyl sulfoxide are added portionwise 4 parts of sodium hydride dispersion 60%. Upon completion, stirring is continued overnight. The reaction mixture is poured onto water. The precipitated product is sucked off and extracted with trichloromethane. The extract is dried, filtered and evaporated. The residue is crystallized from 2,2'-oxybispropane, yielding 10 parts (35%) of [1-(phenylmethyl)-4-piperidinyl](2-thienyl)methanone; mp. 100.5°C.

To a stirred mixture of 15 parts of [1-(phenylmethyl)-4-piperidinyl](2-thienyl)methanone in 120 parts of benzene are added dropwise 8 parts of ethyl carbonochloridate. Upon completion, the whole is heated to reflux and stirred for 6 hours at reflux temperature. The reaction mixture is cooled, filtered and evaporated, yielding 13 parts of ethyl 4-(2-thienylcarbonyl)-1-piperidinecarboxylate as a residue.

A mixture of 20 parts of ethyl 4-(2-thienylcarbonyl)-1-piperidinecarboxylate and 120 parts of hydrobromic acid solution 48% in water is stirred and refluxed for 2 hours. The reaction mixture is cooled and the precipitated product is filtered off. It is washed with 2-propanol and dried, yielding 17 parts (85%) of (4-piperidinyl)(2-thienyl)methanone hydrobromide.

B) Preparation of Final Compounds.


Example XXIII


A mixture of 4.5 parts of 3-(2-chloroethyl)-2,4(1H,3H)-quinazolinedione, 4.9 parts of (4-fluorophenyl)(4-piperidinyl)methanone hydrochloride, 8 parts of sodium carbonate and 80 parts of 4-methyl-2-pentanone is stirred and refluxed overnight. The reaction mixture is cooled and water is added. The precipitated product is filtered off and crystallized from 4-methyl-2-pentanone, yielding, after drying, 2.2 parts (27%) of 3-[2-[4-(4-fluorobenzoyl)-1-piperidinylethyl]-2,4(1H,3H)-quinazolinedione; mp. 227—235°C.

Following the same procedure and using equivalent amounts of the appropriate starting materials there are also prepared:

60

65



	(R ¹) _n	R ³	Alk	R	X	Ar	Base or Salt	Melting Point in °C
10	—	H	(CH ₂) ₃	H	CO	4-F—C ₆ H ₄	—	187.2
15	—	H	(CH ₂) ₂	H	CO	4-Cl—C ₆ H ₄	—	259
	—	H	(CH ₂) ₂	H	CO	4-OCH ₃ —C ₆ H ₄	—	236.5
	—	H	(CH ₂) ₂	H	CO	4-CH ₃ —C ₆ H ₄	—	211.7
20	7-Cl	H	(CH ₂) ₂	H	CO	4-F—C ₆ H ₄	—	235.2
	—	CH ₃	(CH ₂) ₂	H	CO	4-F—C ₆ H ₄	HCl	248.1
25	—	H	(CH ₂) ₂	H	CO	3-CF ₃ —C ₆ H ₄	—	182.6—184.6
	—	H	(CH ₂) ₂	H	CO	3-CH ₃ —C ₆ H ₄	—	171
	—	H	(CH ₂) ₂	H	CO	4-Br—C ₆ H ₄	—	251.5
30	6,8-Cl ₂	H	(CH ₂) ₂	H	CO	4-F—C ₆ H ₄	—	211
	—	H	(CH ₂) ₂	H	CO	2-Cl—C ₆ H ₄	—	162.5
	—	H	(CH ₂) ₂	H		4-F—C ₆ H ₄	—	204.5
40	—	H	(CH ₂) ₂	H	CH ₂	4-F—C ₆ H ₄	—	193.4
	—	H	(CH ₂) ₂	4-OH	CO	4-F—C ₆ H ₄	—	255.2
45	6-Cl	H	(CH ₂) ₂	H	CO	4-F—C ₆ H ₄	—	222.2—223
	6-CH ₃	H	(CH ₂) ₂	H	CO	4-F—C ₆ H ₄	—	205.5
50	—	H	(CH ₂) ₂	H	CO	C ₆ H ₅	—	214.5
	—	H	(CH ₂) ₂	H	CO	2-CH ₃ —C ₆ H ₄	—	193.8
	—	H	(CH ₂) ₂	H	CO	2-thienyl	—	266.9

Example XXIV

55 A mixture of 6 parts of 3-(2-chloroethyl)-1-phenyl-2,4(1H,3H)-quinazolinedione, 8.3 parts of (4-fluorophenyl)(4-piperidyl)methanone and 48 parts of ethanol is stirred and refluxed. N,N-dimethylformamide is added at reflux temperature till all solid enters solution. Stirring is continued overnight at
 60 reflux. The reaction mixture is evaporated. The residue is purified twice by column-chromatography over silica gel using first a mixture of trichloromethane and methanol (95:5 by volume) and then a mixture of trichloromethane and methanol 98:2 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The residue is crystallized from 2-propanol, yielding 2.2 parts (23%) of
 65 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidyl] - ethyl] - 1 - phenyl - 2,4(1H,3H) - quinazolinedione; mp. 166.1°C.

Example XXV

A mixture of 5.8 parts of 6 - chloro - 3 - (2 - chloroethyl) - 3,4 - dihydro - 4 - phenyl - 2 - (1H) - quinazolinone, 7.4 parts of (4 - fluorophenyl)(4 - piperidinyl)methanone and 56 parts of ethanol is stirred and refluxed overnight. The reaction mixture is evaporated. The residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The oily residue solidifies on triturating in 2,2'-oxybispropane. The product is filtered off and dried, yielding 5 parts (57%) of 6 - chloro - 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 3,4 - dihydro - 4 - phenyl - 2(1H) - quinazolinone; mp. 196°C.

In a similar manner there are also prepared:

3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 3,4 - dihydro - 4 - phenyl - 2(1H) - quinazolinone; mp. 173.3°C; and

3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 3,4 - dihydro - 4 - methyl - 2(1H) - quinazolinone; mp. 172°C.

Example XXVI

A mixture of 3.6 parts of 1 - (3 - chloropropyl) - 2,4(1H,3H) - quinazolinone, 3.02 parts of (4 - fluorophenyl)(4 - piperidinyl)methanone hydrochloride, 3.7 parts of sodium carbonate, 0.1 parts of potassium iodide and 120 parts of 4 - methyl - 2 - pentanone is stirred and refluxed overnight using a water-separator. The reaction mixture is cooled, water is added and the layers are separated. The organic phase is dried, filtered and concentrated to a small volume. After cooling the concentrate, the precipitated product is filtered off and crystallized from 2-propanol, yielding 1.3 parts (18.5%) of 1 - [3 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]propyl] - 2,4(1H,3H) - quinazolinone; mp. 192.9°C.

Following the same procedure and starting from 1 - (2 - chloroethyl) - 2,4(1H,3H) - quinazolinone there is also prepared:

1 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,4 - (1H, 3H) - quinazolinone; mp. 219.7°C.

Example XXVII

A mixture of 5.6 parts of 1 - (3 - chloropropyl) - 2,3 - dihydro - 2,2 - dimethyl - 4(1H) - quinazolinone, 4.9 parts of (4 - fluorophenyl)(4 - piperidinyl)methanone hydrochloride, 10 parts of sodium carbonate, 0.1 parts of potassium iodide and 200 parts of 4 - methyl - 2 - pentanone is stirred and refluxed overnight with water-separator. The reaction mixture is cooled, water is added and the layers are separated. The 4 - methyl - 2 - pentanone phase is dried, filtered and evaporated. The solid residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The solid residue is stirred in 2,2'-oxybispropane. The product is filtered off and dried, yielding 6.6 parts (78%) of 1 - [3 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]propyl] - 2,3 - dihydro - 2,2 - dimethyl - 4(1H) - quinazolinone; mp. 185.7°C.

Example XXVIII

A mixture of 5.5 parts of 3 - (2 - chloroethyl) - 2 - phenyl - 4(3H) - quinazolinone, 4 parts of (4 - fluorophenyl)(4 - piperidinyl)methanone hydrochloride, 7 parts of sodium carbonate, 0.1 parts of potassium iodide and 200 parts of 4 - methyl - 2 - pentanone is stirred and refluxed for 24 hours using a water-separator. The reaction mixture is filtered while hot. The filtrate is evaporated. The residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The residue is crystallized from a mixture of ethanol and 1,1'-oxybisethane. The product is filtered off and dried, yielding 5 parts (69%) of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2 - phenyl - 4(3H) - quinazolinone; mp. 143.8°C.

Example XXIX

A mixture of 10 parts of N - [2 - (aminocarbonyl)phenyl] - 4 - (4 - fluorobenzoyl) - 1 - piperidinebutanamide, 60 parts of sodium hydroxide solution 2N and 80 parts of ethanol is stirred and refluxed for 1.50 hours. The reaction mixture is evaporated and water is added to the residue. The oily product is dissolved in trichloromethane. The solution is dried, filtered and evaporated. The residue is crystallized twice from 4 - methyl - 2 - pentanone. The product is filtered off and dried, yielding 1 part (10.5%) of 2 - [3 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]propyl] - 4(3H) - quinazolinone; mp. 184°C.

Example XXX

A mixture of 7 parts of 2 - amino - N - [3 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]propyl]benzamide, 12 parts of formic acid and 135 parts of methylbenzene is stirred and refluxed for 1 hour. Stirring at reflux is continued for 1 hour using a water-separator. The solvent is evaporated. The oily residue is purified by column-chromatography over silica gel using a mixture of

trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The residue is converted into the (E) - 2 - butene - dioate salt in 2 - propanol. The salt is filtered off and dried, yielding 5 parts (54%) of 3 - [3 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]propyl] - 4(3H)-quinazolinone (E)-2-butenedioate (1:1); mp. 201.1°C.

- 5 Following the same procedure and starting from 2 - amino - N - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl]benzamide there is also prepared.
3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 4(3H) - quinazolinone; mp. 139.2°C.

Example XXXI

- 10 A mixture of 6.5 parts of 2 - amino - N - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl]benzamide, 0.5 parts of paraformaldehyde, 3 drops of sodium hydroxide solution 50%, 80 parts of ethanol and 10 parts of water is stirred and refluxed overnight. The reaction mixture is allowed to cool to room temperature while stirring. The precipitated product is filtered off, washed with ethanol and crystallized from 200 parts of ethanol, yielding 3.3 parts (50%) of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,3 - dihydro - 4(1H) - quinazolinone; mp. 203.4°C.

- 15 Following the same procedure and starting from 2 - amino - N - [3 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]propyl]benzamide there is also prepared:

3 - [3 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]propyl] - 2,3 - dihydro - 4 - (1H) - quinazolinone (E) - 2 - butenedioate (1:1); imp. 188.2°C.

20

Example XXXII

- A mixture of 6 parts of 2 - amino - N - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl]benzamide, 10 parts of acetic acid anhydride and 108 parts of methylbenzene is stirred and refluxed overnight. The reaction mixture is evaporated and water is added to the oily residue.
25 The whole is alkalinized with a dilute sodium hydroxide solution and extracted with trichloromethane. The extract is dried, filtered and evaporated. The oily residue is crystallized from a mixture of 4 - methyl - 2 - pentanone and 2,2' - oxybispropane. The product is filtered off and recrystallized from 2-propanol, yielding 4.4 parts (69%) of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2 - methyl - 4(3H) - quinazolinone; mp. 164.9°C.

30

Example XXXIII

- A mixture of 9 parts of 2-aminobenzenecarboxamide, 12 parts of concentrated hydrochloric acid and 120 parts of absolute ethanol is stirred and refluxed for 2 hours. Then there are added 3.6 parts of [1 - (3,3 - diethoxypropyl) - 4 - piperidinyl] (4-fluorophenyl) methanone and stirring at reflux is continued overnight. The reaction mixture is evaporated and the residue is stirred with water. The whole is alkalinized with ammonia and the product is extracted with trichloromethane. The extract is dried, filtered and evaporated. The residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The residue is crystallized twice from 4 - methyl - 2 - pentanone,
40 yielding 1.9 parts (19%) of 2 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,3 - dihydro - 4(1H) - quinazolinone; mp. 153.4°C.

Example XXXIV

- A mixture of 6.7 parts of methyl 2-isothiocyanatobenzoate, 8 parts of [1 - (2 - amino - 1 - methylethyl) - 4 - piperidinyl] (4 - fluorophenyl)methanone and 108 parts of tetrahydrofuran is stirred overnight at room temperature. The reaction mixture is evaporated. The residue is purified twice by column-chromatography over silica gel using first a mixture of trichloromethane and methanol (90:10 by volume) and then a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The residue is crystallized twice from 2-propanol,
50 yielding 1.6 parts (13%) of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]propyl] - 2,3 - dihydro - 2 - thioxo - 4(1H) - quinazolinone; mp. 160.6°C.

Example XXXV

- A mixture of 2.2 parts of 2 - amino - 4 - fluoro - N - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl]benzamide, 1.2 parts of 1,1'-bis[1H-imidazol-1-yl]methanone and 45 parts of tetrahydrofuran is stirred and refluxed for 3 hours. The reaction mixture is cooled and the solvent is evaporated. The residue is boiled in 4 - methyl - 2 - pentanone. The solid product is filtered off and crystallized from N,N-dimethylformamide, yielding 0.8 parts (30%) of 7 - fluoro - 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinedione; mp. 248°C.

- 60 In a similar manner there is also prepared:

3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 7 - methoxy - 2,4(1H,3H) - quinazolinedione; mp. 245.4°C.

Example XXXVI

- 65 To a stirred solution of 6.5 parts of 2 - amino - N - [3 - [4 - (4 - fluorobenzoyl) - 1 -

piperidiny]propyl]benzamide in 63 parts of tetrahydrofuran are added 3 parts of 1,1' - bis[1H - imidazol - 1 - yl]methanethione at room temperature. Stirring is continued for 1 hour at room temperature. The reaction mixture is filtered till clear and the solvent is evaporated. The oily residue is stirred in a mixture of water and 2,2'-oxybispropane. The precipitated product is filtered off and crystallized from a mixture of N,N-dimethylformamide, water and a small amount of ethanol, yielding 5.6 parts (78%) of 3 - [3 - [4 - (4 - fluorobenzoyl) - 1 - piperidiny]propyl] - 2,3 - dihydro - 2 - thioxo - 4(1H) - quinazolinone, mp. 200.2°C.

In a similar manner there is also prepared:

3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidiny]ethyl] - 2,3 - dihydro - 2 - thioxo - 4(1H) - quinazolinone; mp. 225.5°C.

Example XXXVII

A mixture of 2.3 parts of N - [2 - [1 - [2 - (1,4 - dihydro - 2,4 - dioxo - 3(2H) - quinazolinyl) - ethyl] - 4 - piperidiny]carbonyl] - 5 - fluorophenyl] - formamide, 33 parts of concentrated sulfuric acid and 144 parts of ethanol is stirred and refluxed for 1 hour. The reaction mixture is poured onto crushed ice and alkalized with a dilute sodium hydroxide solution. The precipitated product is filtered off and crystallized from 4 - methyl - 2 - pentanone, yielding 0.4 parts of 3 - [2 - [4 - (2 - amino - 4 - fluorobenzoyl) - 1 - piperidiny]ethyl] - 2,4(1H,3H) - quinazolinone; mp. 269.5°C.

Example XXXVIII

To a stirred mixture of 2.6 parts of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidiny]ethyl] - 2,4(1H,3H) - quinazolinone and 120 parts of methanol are added portionwise 2 parts of sodium borohydride. Upon completion, stirring is continued overnight at room temperature. Water is added and the hole is evaporated. The residue is stirred in water. The solid product is filtered off and crystallized from 2-propanone, yielding 1.2 parts of 3 - [2 - [4 - [(4 - fluorophenyl)hydroxymethyl] - 1 - piperidiny]ethyl] - 2,4(1H,3H) - quinazolinone; mp. 212.8°C.

Example XXXIX

A mixture of 5 parts of 3 - (2 - chloroethyl) - 2,4(1H,3H) - quinazolinone, 6 parts of α - (4 - fluorophenyl) - 4 - piperidinemethanol acetate (ester) hydrochloride, 8 parts of sodium carbonate and 280 parts of 4 - methyl - 2 - pentanone is stirred and refluxed overnight. The reaction mixture is filtered and the filtrate is evaporated. The oily residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions are collected and the eluent is evaporated. The residue is crystallized from 4 - methyl - 2 - pentanone, yielding 2.3 parts of 1 - [2 - (1,4 - dihydro - 2,4 - dioxo - 3(2H) - quinazolinyl)ethyl] - α - (4 - fluorophenyl) - 4 - piperidinemethanol acetate (ester); mp. 185.2°C.

Example XL

Following the procedure described in Example XXV and using equivalent amounts of the appropriate starting materials there is also prepared:

3,4-dihydro - 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidiny]ethyl] - 2(1H) - quinazolinone; mp. 182. 182.1°C.

Example XLI

Following the procedure described in Example XXIII and using equivalent amounts of the appropriate starting materials there is also prepared:

3 - [2 - [4 - (3 - pyridinylcarbonyl) - 1 - piperidiny]ethyl] - 2,4(1H,3H) - quinazolinone.


Example XLII

Following the procedure described in Example XVIII—D and using equivalent amounts of the appropriate starting materials there is also prepared:

3 - [2 - [4 - (1H-indol - 3 - yl) - 1 - piperidiny]ethyl] - 2,4(1H,3H) - quinazolinone; mp. 291.2°C.

Claims

1. A chemical compound selected from the group consisting of a quinazoline derivative having the formula



and the pharmaceutically acceptable acid addition salts thereof, wherein:



0013612

piperidinyl]propyl]benzamide in 63 parts of tetrahydrofuran are added 3 parts of 1,1' - bis[1H - imidazol - 1 - yl]methanethione at room temperature. Stirring is continued for 1 hour at room temperature. The reaction mixture is filtered till clear and the solvent is evaporated. The oily residue is stirred in a mixture of water and 2,2'-oxybispropane. The precipitated product is filtered off and
5 crystallized from a mixture of N,N-dimethylformamide, water and a small amount of ethanol, yielding 5.6 parts (78%) of 3 - [3 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]propyl] - 2,3 - dihydro - 2 - thioxo - 4(1H) - quinazolinone, mp. 200.2°C.

In a similar manner there is also prepared:

3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,3 - dihydro - 2 - thioxo - 4(1H) -
10 quinazolinone; mp. 225.5°C.

Example XXXVII

A mixture of 2.3 parts of N - [2 - [1 - [2 - (1,4 - dihydro - 2,4 - dioxo - 3(2H) - quinazolinyl)- ethyl] - 4 - piperidinylcarbonyl] - 5 - fluorophenyl] - formamide, 33 parts of concentrated sulfuric
15 acid and 144 parts of ethanol is stirred and refluxed for 1 hour. The reaction mixture is poured onto crushed ice and alkalized with a dilute sodium hydroxide solution. The precipitated product is filtered off and crystallized from 4 - methyl - 2 - pentanone, yielding 0.4 parts of 3 - [2 - [4 - (2 - amino - 4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinone; mp. 269.5°C.

Example XXXVIII

To a stirred mixture of 2.6 parts of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinone and 120 parts of methanol are added portionwise 2 parts of sodium
20 borohydride. Upon completion, stirring is continued overnight at room temperature. Water is added and the hole is evaporated. The residue is stirred in water. The solid product is filtered off and crystallized from 2-propanone, yielding 1.2 parts of 3 - [2 - [4 - [(4 - fluorophenyl)hydroxymethyl] - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinone; mp. 212.8°C.

Example XXXIX

A mixture of 5 parts of 3 - (2 - chloroethyl) - 2,4(1H,3H) - quinazolinone, 6 parts of α - (4 -
30 fluorophenyl) - 4 - piperidinemethanol acetate (ester) hydrochloride, 8 parts of sodium carbonate and 280 parts of 4 - methyl - 2 - pentanone is stirred and refluxed overnight. The reaction mixture is filtered and the filtrate is evaporated. The oily residue is purified by column-chromatography over silica gel using a mixture of trichloromethane and methanol (95:5 by volume) as eluent. The pure fractions
35 are collected and the eluent is evaporated. The residue is crystallized from 4 - methyl - 2 - pentanone, yielding 2.3 parts of 1 - [2 - (1,4 - dihydro - 2,4 - dioxo - 3(2H) - quinazolinyl)ethyl] - α - (4 - fluorophenyl) - 4 - piperidinemethanol acetate (ester); mp. 185.2°C.

Example XL

Following the procedure described in Example XXV and using equivalent amounts of the
40 appropriate starting materials there is also prepared:

3,4-dihydro - 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2(1H) - - quinazolinone; mp. 182. 182.1°C.

Example XLI

Following the procedure described in Example XXIII and using equivalent amounts of the
45 appropriate starting materials there is also prepared:

3 - [2 - [4 - (3 - pyridinylcarbonyl) - - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinone.


Example XLII

Following the procedure described in Example XVIII—D and using equivalent amounts of the
50 appropriate starting materials there is also prepared:

3 - [2 - [4 - (1H-indol - 3 - yl) - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinone; mp. 291.2°C.

Claims

1. A chemical compound selected from the group consisting of a quinazoline derivative having the
60 formula




65 and the pharmaceutically acceptable acid addition salts thereof, wherein:



Ar is an aryl radical;

X is a member selected from the group consisting of



15 wherein said R_a is hydrogen or lower alkyl and q is the integer 2 or 3;


R is a member selected from the group consisting of hydrogen, hydroxy and lower alkyl;

Alk is an alkylene chain having from 1 to 4 carbon atoms; and

Q is a quinazoliny radical, the 1-, 2-, 3- or 4-position of which is connected with the alkylene side chain, said quinazoliny radical bearing in one or both of its 2- and 4-positions a carbonyl or thiocarbonyl group, wherein the benzene ring of said quinazoliny radical is optionally substituted with 1 to 3 substituents each independently selected from the group consisting of halo, lower alkyl, lower alkyloxy, trifluoromethyl, nitro and cyano, and wherein the pyrimidino ring of said quinazoliny radical may be partly or fully saturated, said pyrimidino ring being optionally substituted with 1 to 3 substituents independently selected from the group consisting of lower alkyl, aryl and aryl(lower alkyl);
 20 wherein said aryl as used in the definition of said Ar and of said Q is a member selected from the group consisting of phenyl, substituted phenyl, thienyl and pyridinyl, wherein said substituted phenyl has from 1 to 3 substituents each independently selected from the group consisting of halo, lower alkyl, lower alkyloxy, trifluoromethyl and amino, and wherein said lower alkyl contains from 1 to 6 carbon atoms.

2. A chemical compound selected from the group consisting of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinone and the pharmaceutically acceptable acid addition salts thereof.


3. A chemical compound selected from the group consisting of a quinazoline derivative having the formula



45 and the pharmaceutically acceptable acid addition salts thereof, wherein R , Q and Alk are as defined in claim 1 and

R^8 and R^9 are each independently selected from the group consisting of hydrogen, halo, C_1-C_6 alkyl, C_1-C_6 alkyloxy, trifluoromethyl and amino.

4. An anti-congestive composition comprising an inert carrier and as an active ingredient an effective anti-congestive amount of a chemical compound selected from the group consisting of a quinazoline derivative having the formula




60 and the pharmaceutically acceptable acid addition salts thereof, wherein Q , Alk , R , X and Ar are as defined in claim 1.

5. An anti-congestive composition comprising an inert carrier and as an active ingredient an effective anti-congestive amount of a chemical compound selected from the group consisting of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl]ethyl] - 2,4(1H,3H) - quinazolinone and the pharmaceutically acceptable acid-addition salts thereof.




6. An anti-congestive composition comprising an inert carrier and as an active ingredient an effective anti-congestive amount of a chemical compound selected from the group consisting of a quinazoline derivative having the formula



and the pharmaceutically acceptable acid addition salts thereof, wherein

Q, Alk, R, R⁸ and R⁹ are as defined in claim 3.


7. A chemical compound selected from the group consisting of a quinazoline derivative having the formula



and the pharmaceutically acceptable acid addition salts thereof, wherein Q, Alk, R, X and Ar are as defined in claim 1 for use in treating congestion in warm-blooded animals.


8. A chemical compound selected from the group consisting of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidiny]ethyl] - 2,4(1H,3H) - quinazolinedione and the pharmaceutically acceptable acid addition salts thereof, for use in treating congestion in warm blooded animals.

9. A chemical compound selected from the group consisting of a quinazoline derivative having the formula



and the pharmaceutically acceptable acid addition salts thereof, wherein Q, Alk, R, R⁸ and R⁹ are as defined in claim 3 for use in treating congestion in warm-blooded animals.

10. A process for preparing a chemical compound selected from the group consisting of a quinazoline derivative having the formula



(I)


and the pharmaceutically-acceptable acid addition salts thereof, wherein Q, Alk, R, X and Ar are as defined in claim 1, characterized by

a) reacting a compound of the formula



(II)

wherein W is a reactive ester residue, such as, halo, or a sulfonyloxy radical such as methylsulfonyloxy, or 4-methylphenylsulfonyloxy with a compound of the formula




(III)




at elevated temperatures in an appropriate reaction-inert organic solvent and in the presence of an appropriate base;

b) preparing a compound of the formula




wherein one of R^2 , R^3 , R^4 , and R^5 is D, D being a radical of the formula



and each of the remaining radicals of R^2 , R^3 , R^4 , and R^5 being independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-lower alkyl; R^1 being selected from the group consisting of halo, lower alkyl, lower alkyloxy, trifluoromethyl, nitro and cyano; each Y being O or S; and

25 n being an integer of from 0 to 3 inclusive, by cyclizing a compound of the formula




with a compound of the formula




said cyclization-reaction being carried out by stirring the reactants together in the presence of an appropriate solvent at elevated temperatures; and if desired, in the foregoing reaction, the reagent of formula (V) may be replaced by an appropriate functionally equivalent derivative thereof such as, a

45 di(lower alkyl)acetal or a cyclic acetal; or

c) preparing a compound of the formula



55 by cyclizing a compound of the formula






0013612


wherein either of R^2 and R^4 is D, the other being a member selected from the group consisting of hydrogen, lower alkyl, aryl and aryl lower alkyl, said cyclization reaction being carried out by stirring and heating the 2-(acylamino)benzamide or -benzene-thioamide (VI) in a suitable reaction-inert solvent and in the presence of an appropriate base; or

d) preparing a compound of the formula



15 wherein one of R^2 and R^3 is D, as defined above, the remaining being selected from the group consisting of hydrogen, lower alkyl, aryl and aryl lower alkyl, R^1 and n being as previously described and each Y being independently selected from the group consisting of O and S, by cyclizing a compound of the formula (IV-a) as defined above, with urea or thiourea, said cyclization being carried out by stirring the reactants together at an elevated temperature and, in the presence of an appropriate solvent; or

20 e) preparing a compound of the formula (I-c) as defined above by cyclizing a compound of the formula




with a compound of the formula




wherein Y, R^1 and n have the previously-defined meanings, one of R^2 and R^3 is D, as defined above, the other being hydrogen, lower alkyl or aryl lower alkyl and R^6 and R^6a each representing an appropriate leaving group such as, lower alkyloxy, amino or mono- and di(lower alkyl)amino, said cyclization being carried out by stirring the reactants together at an elevated temperature and, in the presence of an appropriate solvent; or

40 f) preparing a compound of the formula



by cyclizing a compound of the formula



with a compound of the formula




wherein R^6 represents an appropriate leaving group and R^1 , n and Y are as previously described, said cyclizing being carried out by stirring the reactants together at an elevated temperature and, in the presence of an appropriate solvent; or




42

0013612

g) preparing a compound of the formula (1-c-2)




10 by cyclizing a compound of the formula




20 wherein D, Y and R⁶ are as previously described, with an alkali metal cyanate or thiocyanate, said cyclization being carried out by stirring the reactants together at an elevated temperature and, in the presence of an appropriate solvent; or


h) preparing a compound of the formula




35 wherein X' is the same as X as previously described by other than



40 by cyclizing a compound of the formula




55 in the presence of an appropriate reducing agent, and a suitable solvent at room temperature; or
k) preparing a compound of the formula




65




wherein R^8 and R^9 are each independently selected from the group consisting of hydrogen, halo, lower alkyl, lower alkyloxy, trifluoromethyl or amino, by hydrolyzing a compound of the formula



10 in acidic medium; or
j) preparing a compound of the formula




20 by reducing the carbonyl group of a compound of the formula




with an appropriate reducing agent in alkaline aqueous medium with a water-miscible organic solvent;
30 or

k) preparing a compound of the formula




40 by acylating the corresponding alcohol (I-f) as defined above, with an appropriate acylating agent; or
l) preparing a compound of the formula



50 from the corresponding carbonyl derivatives (I-g) by the reduction of the carbonyl group of (I-g) to a methylene group, such as, by the Clemmensen reduction, using amalgated zinc and hydrochloric acid, or by the Wolff-Kishner reduction, using hydrazine and alkali in a high-boiling polar solvent; or

m) preparing compounds of formula (I) wherein X represents a di(loweralkyl)ketal or a cyclic lower alkylene ketal, wherein the lower alkylene chain has from 2 to 4 carbon atoms, from the corresponding carbonyl compounds by subjecting the latter to a ketalization-reaction; or

55 n) preparing compounds of formula (I) wherein X represents a radical of the formula




60 from the corresponding carbonyl compounds by reacting the latter with respectively hydroxylamine hydrochloride or hydrazine hydrochloride; and, if desired, preparing pharmaceutically acceptable acid addition salts of the products of steps a) through n).

65 11. A process for preparing a chemical compound selected from the group consisting of 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - piperidinyl] - ethyl] - 2,4(1H,3H) - quinazolin-2-one and the



pharmaceutically-acceptable acid addition salts thereof, characterized by reacting 3-(2-(chloroethyl)-2,4(1H,3H)-quinazolidinedione with (4-fluorophenyl) (4-piperidiny) methanone hydrochloride, and, if desired, preparing a pharmaceutically-acceptable acid addition salt thereof.

12. A process for preparing a chemical compound selected from the group consisting of a quinazoline derivative having the formula




and the pharmaceutically-acceptable acid addition salts thereof, wherein R, Q, Alk, R⁸ and R⁹ are as defined in claim 3 characterized by

a) reacting a compound of the formula



wherein W is a reactive ester residue such as halo, or a sulfonyloxy radical such as methylsulfonyloxy, or 4-methylphenylsulfonyloxy with a compound of the formula



(XXIX)


at elevated temperatures, in an appropriate reaction inert organic solvent and in the presence of an appropriate base, and if desired, preparing a pharmaceutically-acceptable acid addition salts thereof.

13. A compound according to any one of claims 1 to 3, or a composition according to any one of claims 4 to 6, for use as a serotonin antagonist.

14. A process for preparing an anti-congestive composition according to any one of claims 4 to 6 comprising combining said active ingredient in intimate admixture with said inert carrier.

Patentansprüche


1. Eine chemische Verbindung, ausgewählt aus der ein Chinazolinderivat der Formel



und die pharmazeutisch annehmbaren Säureadditionssalze hiervon umfassenden Gruppe, worin:

Ar ein Arylradikal darstellt;

X ein Vertreter aus der





umfassenden Gruppe ist, worin das erwähnte R_q Wasserstoff oder Niederalkyl darstellt und das erwähnte q die ganze Zahl 2 oder 3 bedeutet;


R ein Vertreter aus der aus Wasserstoff, Hydroxy und Niederalkyl bestehenden Gruppe ist; Alk eine Alkylkette mit von 1 bis 4 Kohlenstoffatomen bedeutet; und

Q ein Chinazolinrest ist, dessen 1-, 2-, 3- oder 4-Stellung mit der Alkylseitenkette verknüpft ist, welcher Chinazolinrest in einer oder in beiden seiner 2- und 4-Stellungen eine Carbonyl- oder Thio-carbonylgruppe trägt, wobei der Benzolring des genannten Chinazolinrestes gewünschtenfalls mit 1 bis 3 Substituenten substituiert ist, die jeweils unabhängig voneinander ausgewählt sind aus einer Gruppe umfassend Halogen, Niederalkyl, Niederalkyl-oxo, Trifluormethyl, Nitro und Cyano, und worin der Pyrimidinring des genannten Chinazolinrestes teilweise oder vollständig gesättigt sein kann, welcher Pyrimidinring gewünschtenfalls mit 1 bis 3 Substituenten substituiert ist, die unabhängig voneinander ausgewählt sind aus der Gruppe bestehend aus Niederalkyl, Aryl und Aryl(niederalkyl);

worin das genannte Aryl, wie es in der Definition von Ar und Q verwendet wird, ein Vertreter aus der Gruppe, bestehend aus Phenyl, substituiertes Phenyl, Thienyl und Pyridinyl ist, wobei genanntes substituiertes Phenyl 1 bis 3 Substituenten aufweist, die jeweils unabhängig voneinander aus der Gruppe, bestehend aus Halogen, Niederalkyl, Niederalkyl-oxo, Trifluormethyl und Amino, ausgewählt sind und worin genanntes Niederalkyl 1 bis 6 Kohlenstoffatome aufweist.


2. Eine chemische Verbindung, ausgewählt aus der 3-[2-[4-(4-Fluorbenzoyl)-1-piperidinyl]-ethyl]-2,4-(1H,3H)-chinazolin-1-one und die pharmazeutisch annehmbaren Säureadditionssalze hiervon umfassenden Gruppe.

3. Eine chemische Verbindung, ausgewählt aus der ein Chinazolin-derivat der Formel



un die pharmazeutisch annehmbaren Säureadditionssalze hiervon umfassenden Gruppe, wobei R , Q und Alk wie in Anspruch 1 definiert sind und R^8 und R^9 jeweils unabhängig voneinander ausgewählt sind aus der Gruppe bestehend aus Wasserstoff, Halogen, C_1-C_6 -Alkyl, C_1-C_6 -Alkyl-oxo, Trifluormethyl und Amino.


4. Eine antikongestive Zusammensetzung, umfassend einen inerten Träger und als einen wirksamen Bestandteil eine antikongestiv wirksame Menge einer chemischen Verbindung, ausgewählt aus der ein Chinazolin-derivat der Formel



und deren pharmazeutisch annehmbaren Säureadditionssalze umfassenden Gruppe, worin Q , Alk, R , X und Ar wie in Anspruch 1 definiert sind.


5. Eine antikongestive Zusammensetzung, umfassend einen inerten Träger und als einen wirksamen Bestandteil eine antikongestiv wirksame Menge einer chemischen Verbindung, ausgewählt aus der 3-[2-[4-(4-Fluorbenzoyl)-1-piperidinyl]-ethyl]-2,4-(1H,3H)-chinazolin-1-one und die pharmazeutisch annehmbaren Säureadditionssalze hiervon umfassenden Gruppe.

6. Eine antikongestive Zusammensetzung, umfassend einen inerten Träger und als einen wirksamen Bestandteil eine antikongestiv wirksame Menge einer chemischen Verbindung, ausgewählt aus der ein Chinazolin-derivat der Formel



und die pharmazeutisch annehmbaren Säureadditionssalze hievon umfassenden Gruppe, worin Q, Alk, R, R⁸ und R⁹ wie in Anspruch 3 definiert sind.


7. Eine chemische Verbindung, ausgewählt aus der aus einem Chinazolinderivat der Formel



und den pharmazeutisch annehmbaren Säureadditionssalzen hievon bestehenden Gruppe, worin Q, Alk, R, X und Ar wie in Anspruch 1 definiert sind, zur Anwendung in der Behandlung von Kongestionen bei warmblütigen Tieren.


8. Eine chemische Verbindung, ausgewählt aus der aus 3-[2-[4-(4-Fluorbenzoyl)-1-piperidiny]-ethyl]-2,4(1H,3H)-chinazolindion und den pharmazeutisch annehmbaren Säureadditionssalzen hievon bestehenden Gruppe, zur Verwendung in der Behandlung von Kongestionen in warmblütigen Tieren.

9. Eine chemische Verbindung, ausgewählt aus der aus einem Chinazolinderivat der Formel



und den pharmazeutisch annehmbaren Säureadditionssalzen hievon bestehenden Gruppe, worin Q, Alk, R, R⁸ und R⁹ wie in Anspruch 3 definiert sind, zur Anwendung in der Behandlung von Kongestionen in warmblütigen Tieren.

10. Ein Verfahren zur Herstellung einer chemischen Verbindung, ausgewählt aus der aus einem Chinazolinderivat der Formel




und den pharmazeutisch annehmbaren Säureadditionssalzen hievon bestehenden Gruppe, worin Q, Alk, R, X und Ar wie in Anspruch 1 definiert sind, gekennzeichnet durch

a) Umsetzen einer Verbindung der Formel



worin W einen reaktionsfähigen Esterrest, wie Halogen, oder einen Sulfonyloxyrest, wie Methylsulfonyloxy, oder 4-Methylsulfonyloxy bedeutet, mit einer Verbindung der Formel




bei erhöhten Temperaturen in einem geeigneten reaktionsinerten organischen Lösungsmittel und in Gegenwart einer geeigneten Base;

60


65




b) Bereiten einer Verbindung der Formel



worin einer der Reste R^2 , R^3 , R^4 und R^5 für D steht, wobei D ein Rest der Formel



ist und jeder der verbleibenden Reste R^2 , R^3 , R^4 und R^5 unabhängig voneinander ausgewählt ist aus der Gruppe, bestehend aus Wasserstoff, Niederalkyl, Aryl und Arylniederalkyl; R^1 aus der aus Halogen, Niederalkyl, Niederalkyloxy, Trifluormethyl, Nitro und Cyano bestehenden Gruppe ausgewählt ist; jedes Y für Sauerstoff oder Schwefel steht; und n eine ganze Zahl von Null bis einschließlich 3 bedeutet, durch Cyclisieren einer Verbindung der Formel




mit einer Verbindung der Formel




welche Cyclisierungsreaktion durch gemeinsames Rühren der Reaktanten in Gegenwart eines geeigneten Lösungsmittels bei erhöhten Temperaturen bewirkt wird; und wobei gewünschtenfalls in der vorstehenden Umsetzung das Reaktionsmittel der Formel (V) durch ein entsprechendes funktionell äquivalentes Derivat hiervon, wie ein Di-(niederalkyl)-acetal oder ein cyclisches Acetal, ersetzt werden kann; oder

c) Bereiten einer Verbindung der Formel




durch Cyclisieren einer Verbindung der Formel






- worin entweder R^2 oder R^4 für D steht, während der andere Rest einen Vertreter, ausgewählt aus der aus Wasserstoff, Niederalkyl, Aryl und Arylniederalkyl bestehenden Gruppe, bedeutet, welche Cyclisierungsreaktion durch Rühren und Erhitzen des 2-(Acylamino)-benzamids oder -benzthioamids (VI) in einem geeigneten reaktionsinerten Lösungsmittel und in Gegenwart einer entsprechenden Base bewirkt wird; oder
- d) Bereiten einer Verbindung der Formel




- 15 worin einer der Reste R^2 und R^3 für D steht, wie vorstehend definiert, während der andere Rest aus der aus Wasserstoff, Niederalkyl, Aryl und Arylniederalkyl bestehenden Gruppe ausgewählt ist, R^1 und n wie zuvor beschrieben definiert sind und jedes Y unabhängig voneinander ausgewählt ist aus der aus Sauerstoff und Schwefel bestehenden Gruppe, durch Cyclisieren einer Verbindung der Formel (IV-a) gemäß vorstehender Definition mit Harnstoff oder Thioharnstoff, welche Cyclisierung ausgeführt wird
- 20 durch gemeinsames Rühren der Reaktanten bei einer erhöhten Temperatur und in Gegenwart eines geeigneten Lösungsmittels; oder
- e) Bereiten einer Verbindung der Formel (I-c) gemäß obiger Definition durch Cyclisieren einer Verbindung der Formel




mit einer Verbindung der Formel



- worin Y, R^1 und n die zuvor angegebenen Bedeutungen aufweisen, einer der Reste R^2 und R^3 für D, wie zuvor definiert, steht, während der andere Rest Wasserstoff, Niederalkyl oder Arylniederalkyl ist, und R^6 und R^6_a jeweils eine geeignete Leaving-Gruppe wie Niederalkyloxy, Amino oder Mono- und Di-(niederalkyl)-amino bedeuten, welche Cyclisierung ausgeführt wird durch gemeinsames Rühren der Reaktanten bei einer erhöhten Temperatur und in Gegenwart eines geeigneten Lösungsmittels; oder
- f) Bereiten einer Verbindung der Formel



durch Cyclisieren einer Verbindung der Formel




mit einer Verbindung der Formel



- worin R^6 eine geeignete Leaving-Gruppe bedeutet und R^1 , n und Y wie zuvor beschrieben definiert sind,
- 65 welche Cyclisierung ausgeführt wird durch gemeinsames Rühren der Reaktanten bei einer erhöhten




Temperatur und in Gegenwart eines geeigneten Lösungsmittels; oder
g) Bereiten einer Verbindung der Formel



(I-c-2)


durch Cyclisieren einer Verbindung der Formel



(IV-d)


worin D, Y und R^6 wie zuvor beschrieben definiert sind, mit einem Alkalimetallcyanat oder -thiocyanat, welche Cyclisierung ausgeführt wird durch gemeinsames Rühren der Reaktanten bei einer erhöhten Temperatur und in Gegenwart eines geeigneten Lösungsmittels; oder

h) Bereiten einer Verbindung der Formel




(I-d)

worin X' die gleiche Bedeutung wie X gemäß vorstehender Definition, ausgenommen die Gruppe




aufweist, durch Cyclisieren einer Verbindung der Formel



(X)

in Gegenwart eines entsprechenden Reduktionsmittels und eines geeigneten Lösungsmittels bei Raumtemperatur; oder

i) Bereiten einer Verbindung der Formel




(I-e)




57


worin R^8 und R^9 jeweils unabhängig voneinander ausgewählt sind aus der aus Wasserstoff, Halogen, Niederalkyl, Niederalkyloxy, Trifluormethyl und Amino bestehenden Gruppe, durch Hydrolyse einer Verbindung der Formel



10 in saurem Medium; oder
j) Bereiten einer Verbindung der Formel




20 durch Reduzieren der Carbonylgruppe einer Verbindung der Formel




mit einem entsprechenden Reduktionsmittel in alkalischem wässrigem Medium mit einem wasser-

30 k) Bereiten einer Verbindung der Formel



40 durch Acylieren des entsprechenden Alkohols (I-f) gemäß vorstehender Definition mit einem entsprechenden Acylierungsmittel; oder


l) Bereiten einer Verbindung der Formel



50 aus den entsprechenden Carbonylderivaten (I-g) durch die Reduktion der Carbonylgruppe von (I-g) zu einer Methylengruppe, beispielsweise nach der Clemmensen-Reduktion unter Verwendung von Zinkamalga-
m und Chlorwasserstoffsäure, oder nach der Wolff-Kishner-Reduktion unter Verwendung von Hydrazin und Alkali in einem hochsiedenden polaren Lösungsmittel; oder

m) Bereiten von Verbindungen der Formel (I), worin X ein Di-(niederalkyl)-ketal oder ein cyclisches Niederalkylenketal bedeutet, worin die Niederalkylenkette 2 bis 4 Kohlenstoffatome aufweist, aus den
55 entsprechenden Carbonylverbindungen durch Ausführung einer Ketalisierungsreaktion an den letztgenannten Verbindungen; oder

n) Bereiten von Verbindungen der Formel (I), worin X einen Rest der Formel




darstellt, aus den entsprechenden Carbonylverbindungen durch Umsetzen der letztgenannten mit Hydroxylaminhydrochlorid bzw. Hydrazinhydrochlorid; und, gewünschtenfalls Bereiten von pharmazeutisch annehmbaren Säureadditionssalzen der Produkte der Stufen a) bis n).



0013612

11. Ein Verfahren zur Herstellung einer chemischen Verbindung, ausgewählt aus der 3-[2-[4-(4-Fluorbenzoyl)-1-piperidiny]-ethyl]-2,4-(1H,3H)-chinazolidindion und die pharmazeutisch annehmbaren Säureadditionssalze hievon umfassenden Gruppe, gekennzeichnet durch Umsetzen von 3-(2-Chlor-ethyl)-2,4-(1H,3H)-chinazolidindion mit (4-Fluorphenyl)-(4-piperidiny)-methanon-hydrochlorid und

12. Ein Verfahren zur Herstellung einer chemischen Verbindung, ausgewählt aus der ein Chinazolinderivat der Formel




(XII)

und die pharmazeutisch annehmbaren Säureadditionssalze hievon umfassenden Gruppe, worin R, Q, Alk, R⁸ und R⁹ wie in Anspruch 3 definiert sind, gekennzeichnet durch Umsetzen einer Verbindung der Formel



(II)

worin W einen reaktionsfähigen Esterrest wie Halogen oder einen Sulfonyloxyrest wie Methylsulfonyloxy oder 4-Methylphenylsulfonyloxy bedeutet, mit einer Verbindung der Formel



(XXIX)


bei erhöhten Temperaturen in einem entsprechenden reaktionsinerten organischen Lösungsmittel und in Gegenwart einer geeigneten Base und gewünschtenfalls Bereiten von pharmazeutisch annehmbaren Säureadditionssalzen hievon.

13. Eine Verbindung gemäß einem der Ansprüche 1 bis 3, oder eine Zusammensetzung gemäß einem der Ansprüche 4 bis 6, zur Anwendung als ein Serotonin-Antagonist.

14. Ein Verfahren zur Herstellung einer antikongestiven Zusammensetzung nach einem der Ansprüche 4 bis 6, umfassend das Überführen des genannten wirksamen Bestandteiles in ein inniges Gemisch mit dem genannten inerten Träger.

Revendications

1. Un composé chimique choisi dans le groupe consistant en un dérivé de quinazoline de formule




(I)

et ses sels formés par addition avec des acides acceptables pour l'usage pharmaceutique, dans lesquels:

Ar représente un radical aryle;

X est choisi dans le groupe consistant en



R_a représentant l'hydrogène ou un groupe alkyle inférieur et q étant égal à 2 ou 3;

15 R est choisi dans le groupe formé par l'hydrogène, les groupes hydroxy et alkyle inférieurs;


Alk représente une chaîne alkylène contenant de 1 à 4 atomes de carbone; et

Q représente un radical quinazolinyle dont la position 1, 2, 3 ou 4 est reliée avec la chaîne latérale alkylène, ce radical quinazolinyle portant sur l'une de ses positions 2 et 4 ou sur les deux un groupe carbonyle ou thiocarbonyle, le noyau benzénique de ce radical quinazolinyle portant facultativement 1 à 3 substituants choisis chacun, indépendamment les uns des autres, parmi les atomes d'halogènes, les groupes alkyle inférieurs, alkyloxy inférieurs, trifluorométhyle, nitro et cyano, le cycle pyrimidino de ce radical quinazolinyle pouvant être saturé en totalité ou en partie, ce cycle pyrimidino portant facultativement 1 à 3 substituants choisis indépendamment parmi les groupes alkyle inférieurs, aryle et aryl-

25 les groupes aryle dont il est question dans la définition de Ar et de Q étant choisis parmi les groupes phényle, phényle substitué, thiényl et pyridinyle, le groupe phényle substitué portant 1 à 3 substituants choisis chacun, indépendamment, parmi les atomes d'halogènes, les groupes alkyle inférieurs, alkyloxy inférieurs, trifluorométhyle et amino, les groupes alkyle inférieurs contenant de 1 à 6 atomes de carbone.

30 2. Un composé chimique choisi dans le groupe formé par la 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - pipéridinyl]éthyl] - 2,4(1H,3H) - quinazoline - dione et ses sels formés par addition avec des acides acceptables pour l'usage pharmaceutique.

3. Un composé chimique choisi dans le groupe consistant en un dérivé de quinazoline répondant à la formule



45 et ses sels formés par addition avec des acides acceptables pour l'usage pharmaceutique, dans lesquels R, Q et Alk ont les significations indiquées dans la revendication 1 et R^8 et R^9 sont choisis chacun, indépendamment, parmi les atomes d'hydrogène, les atomes d'halogènes, les groupes alkyle en C1-C6, alkyloxy en C1-C6, trifluorométhyle et amino.

50 4. Une composition anti-congestive comprenant un véhicule inerte et, en tant que composant actif, en quantité anti-congestive efficace, un composé chimique choisi dans le groupe consistant en un dérivé de quinazoline répondant à la formule



60 et ses sels formés par addition avec des acides acceptables pour l'usage pharmaceutique, dans lesquels Q, Alk, R, X et Ar ont les significations indiquées dans la revendication 1.


5. Une composition anti-congestive comprenant un véhicule inerte et, en tant que composant actif, en proportion anti-congestive efficace, un composé chimique choisi dans le groupe consistant en la 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - pipéridinyl] - éthyl] - 2,4(1H,3H) - quinazoline - dione et ses sels formés par addition avec des acides acceptables pour l'usage pharmaceutique.

6. Une composition anti-congestive comprenant un véhicule inerte et, en tant que composant




0013 612

actif, en proportion anti-congestive efficace, un composé chimique choisi dans le groupe consistant en un dérivé de quinazoline de formule



et ses sels formés par/addition avec des acides acceptables pour l'usage pharmaceutique, dans lesquels Q, Alk, R, R⁸ et R⁹ ont les significations indiquées dans la revendication 3.


15 7. Un composé chimique choisi dans le groupe consistant en un dérivé de quinazoline répondant à la formule



et ses sels formés par addition avec des acides acceptables pour l'usage pharmaceutique, dans lesquels Q, Alk, R, X et Ar ont les significations indiquées dans la revendication 1, pour l'utilisation dans le traitement de la congestion chez les animaux à sang chaud.


25 8. Un composé chimique choisi dans le groupe consistant en la 3 - [2 - [4 - (4 - fluoro-benzoyl) - 1 - pipéridinyl] - éthyl] - 2,4(1H,3H) - quinazoline - dione et ses sels formés par addition avec des acides acceptables pour l'usage pharmaceutique, pour l'utilisation dans le traitement de la congestion chez les animaux à sang chaud.

30 9. Un composé chimique choisi dans le groupe consistant en un dérivé de quinazoline répondant à la formule



et ses sels formés par addition avec des acides acceptables pour l'usage pharmaceutique, dans lesquels Q, Alk, R, R⁸ et R⁹ ont les significations indiquées dans la revendication 3, pour l'utilisation dans le traitement de la congestion chez les animaux à sang chaud.

45 10. Un procédé de préparation d'un composé chimique choisi dans le groupe consistant en un dérivé de quinazoline répondant à la formule



(I)


et ses sels formés par addition avec des acides acceptables pour l'usage pharmaceutique, dans lesquels Q, Alk, R, X et Ar ont les significations indiquées dans la revendication 1, caractérisé en ce que

55 a) on fait réagir un composé de formule


Q-Alk-W

(II)

60 dans laquelle W est un reste d'ester réactif tel qu'un reste halogéno, ou un radical sulfonyloxy tel que méthylsulfonyloxy ou 4-méthylphénylsulfonyloxy, avec un composé de formule




(III)




0013612

à température élevée, dans un solvant organique approprié, inerte dans la réaction, et en présence d'une base appropriée;

b) on prépare un composé de formule




dans laquelle l'un des symboles R^2 , R^3 , R^4 et R^5 représente D, D étant un radical de formule



chacun des symboles R^2 , R^3 , R^4 et R^5 restants étant choisis indépendamment parmi les atomes d'hydrogène, les groupes alkyle inférieurs, aryle et aryl-(alkyle inférieur); R^1 étant choisi parmi les atomes d'halogènes, les groupes alkyle inférieurs, alkyloxy inférieurs, trifluorométhyle, nitro et cyano, chacun des symboles Y représentant O ou S; et n représentant un nombre de 0 à 3 inclus, en cyclisant

25 un composé de formule




avec un composé de formule




cette réaction de cyclisation étant effectuée par agitation des réactifs ensemble en présence d'un solvant approprié à température élevée; et si on le désire, dans la réaction ci-dessus, on peut remplacer le réactif de formule V par un dérivé fonctionnel équivalent approprié tel qu'un di-(alkyle inférieur)-acétal ou un acétal cyclique; ou bien

c) on prépare un composé de formule



en cyclisant un composé de formule






0013612

86

dans laquelle l'un ou l'autre des symboles R^2 et R^4 représente D, l'autre étant choisi parmi les atomes d'hydrogène, les groupes alkyle inférieurs, aryle et aryl-(alkyle inférieur), cette réaction de cyclisation étant effectuée par agitation et chauffage du 2-(acylamino)-benzamide ou -benzène-thioamide (VI) dans un solvant approprié inerte dans la réaction et en présence d'une base appropriée; ou bien


5 d) on prépare un composé de formule



15 dans laquelle l'un des symboles R^2 et R^3 représente D tel que défini ci-dessus, l'autre étant choisi parmi les atomes d'hydrogène, les groupes alkyle inférieurs, aryle et aryl-(alkyle inférieur), R^1 et n ayant les significations indiquées précédemment et chacun des symboles Y étant choisi indépendamment dans le groupe consistant en O et S, en cyclisant un composé de formule IV-a tel que défini ci-dessus, à l'aide de l'urée ou de la thiourée, cette cyclisation étant effectuée par agitation des réactifs ensemble à


20 température élevée et en présence d'un solvant approprié; ou bien

e) on prépare un composé de formule I-c tel que défini ci-dessus en cyclisant un composé de formule



30

avec un composé de formule




(VIII),

35

Y, R^1 et n ayant les significations indiquées précédemment, l'un des symboles R^2 et R^3 représentant D tel que défini ci-dessus, l'autre consistant en un atome d'hydrogène, un groupe alkyle inférieur ou aryl-(alkyle inférieur) et R^6 et R^6_a représentant chacun un groupe éliminable approprié tel qu'un groupe alkyloxy inférieur, amino ou mono- et di-(alkyle inférieur)-amino, cette cyclisation étant effectuée par


40 agitation des réactifs ensemble à température élevée et en présence d'un solvant approprié; ou bien

f) on prépare un composé de formule



50

en cyclisant un composé de formule



60

avec un composé de formule



(VIII-a),


65

R^6 représentant un groupe éliminable approprié et R^1 , n et Y ayant les significations indiquées précédemment, cette cyclisation étant effectuée par agitation des réactifs ensemble et à température élevée et en présence d'un solvant approprié; ou bien

65




g) on prépare un composé de formule



(I-c-2)


10 en cyclisant un composé de formule IV-d



(IV-d)


20 D, Y et R⁶ ayant les significations indiquées précédemment, avec un cyanate ou thiocyanate de métal alcalin, cette cyclisation étant effectuée par agitation des réactifs ensemble à température élevée et en présence d'un solvant approprié; ou bien

h) on prépare un composé de formule




(I-d)

35 dans laquelle X' a la même signification que X ci-dessus à l'exception de




40 en cyclisant un composé de formule



(X)

55 en présence d'un agent réducteur approprié et d'un solvant approprié à température ambiante; ou bien
k) on prépare un composé de formule




(I-e)

65 dans laquelle R⁸ et R⁹ sont choisis chacun, indépendamment, parmi les atomes d'hydrogène, les




0013611


atomes d'halogènes; les groupes alkyle inférieurs, alkyloxy inférieurs, trifluorométhyle ou amino, en hydrolysant un composé de formule




10 en milieu acide; ou bien
j) on prépare un composé de formule




20 en réduisant le groupe carbonyle d'un composé de formule



à l'aide d'un agent réducteur approprié en milieu aqueux alcalin avec un solvant organique miscible à l'eau; ou bien
30 k) on prépare un composé de formule




en acylant l'alcool correspondant (I-f) tel que défini ci-dessus, à l'aide d'un agent acylant approprié; ou bien
40 l) On prépare un composé de formule



50 à partir des dérivés carbonylés correspondants I-g en réduisant le groupe carbonyle des composés I-g en un groupe méthylène, par exemple par réduction de Clemmensen, à l'aide de zinc amalgamé et d'acide chlorhydrique ou par la réduction de Wolff-Kishner, à l'aide d'hydrazine et d'un alcali dans un solvant polaire à haut point d'ébullition; ou bien

55 m) on prépare les composés de formule I dans laquelle X représente un di-(alkyle inférieur)-acétal ou un (alkylène inférieur)-acétal cyclique dans lequel la chaîne alkylène inférieure contient de 2 à 4 atomes de carbone, à partir des composés carbonylés correspondants en soumettant ces derniers à une réaction d'acétalisation; ou bien

n) on prépare les composés de formule I dans laquelle X représente un radical de formule



à partir des composés carbonylés correspondants en faisant réagir ces derniers respectivement avec le chlorhydrate d'hydroxylamine ou le chlorhydrate d'hydrazine; et si on le désire, on prépare les sels des
65




0013612

produits obtenus dans les opérations a) à n) par addition avec des acides acceptables pour l'usage pharmaceutique.

11. Un procédé de préparation d'un composé chimique choisi dans le groupe consistant en la 3 - [2 - [4 - (4 - fluorobenzoyl) - 1 - pipéridinyl] - éthyl - 2,4(1H,3H) - quinazoline - dione et ses sels formés par addition avec des acides acceptables pour l'usage pharmaceutique, caractérisé en ce que l'on fait réagir la 3 - (2 - chloréthyl) - 2,4(1H,3H) - quinazolidine - dione avec le chlorhydrate de la (4 - fluorophényl) (4 - pipéridinyl) - méthanone, et si on le désire, on prépare un sel du composé obtenu par addition avec un acide acceptable pour l'usage pharmaceutique.

12. Un procédé de préparation d'un composé chimique choisi dans le groupe consistant en un dérivé de quinazoline répondant à la formule




(XII)

et ses sels formés par addition avec des acides acceptables pour l'usage pharmaceutique, dans lesquels R, Q, Alk, R⁸ et R⁹ ont les significations indiquées dans la revendication 3, caractérisé en ce que a) on fait réagir un composé de formule

Q-Alk-W

(II)

dans laquelle W représente le reste d'un ester réactif tel qu'un reste halogéno, ou un radical sulfonyloxy tel que méthylsulfonyloxy ou 4-méthyl-phénylsulfonyloxy, avec un composé de formule



(XXIX)

à température élevée, dans un solvant organique approprié, inerte dans la réaction et en présence d'une base appropriée, et si on le désire, on prépare des sels des composés obtenus par addition avec des acides acceptables pour l'usage pharmaceutique.

13. Un composé selon l'une quelconque des revendications 1 à 3, ou une composition selon l'une quelconque des revendications 4 à 6, pour l'utilisation en tant qu'antagoniste de la sérotonine.

14. Un procédé de préparation d'une composition anti-congestive selon l'une quelconque des revendications 4 à 6, qui consiste à combiner le composant actif en mélange intime avec le véhicule inerte.