

CLAIMS

1. A composition for vagina, wherein the composition comprises an antimicrobial agent, a total content of which, calculated as 2-phenylethanol, is 0.05 to 0.60 percent (weight/volume), and an antimicrobial agent, a total content of which, calculated as sodium propionate, is 0.25 to 0.70 percent (weight/volume);

the antimicrobial agent comprises 2-phenylethanol; the antimicrobial agent comprises at least one of propionic acid, sodium propionate and calcium propionate; and

wherein a dosage form of the composition is a solution, a water-soluble gel, or an emulsive ointment, and wherein a pH value of the composition ranges from 3.0 to 4.3.

2. The composition according to claim 1, wherein, of a total amount of the composition, the total content of the antimicrobial agent, calculated as 2-phenylethanol, is 0.10 to 0.40 percent (weight/volume), and the total content of the antimicrobial agent, calculated as sodium propionate, is 0.525 to 0.575 percent (weight/volume).

3. The composition according to claim 1 or 2, wherein the composition further comprises an antimicrobial agent, a total content of which, calculated as sodium benzoate, is 0.05 to 0.15 percent (weight/volume); wherein the antimicrobial agent comprises one or both of benzoic acid and sodium benzoate.

4. The composition according to claim 3, wherein, of a total amount of the composition, the total content of the antimicrobial agent, calculated as sodium benzoate, is 0.065 to 0.090 percent (weight/volume).

5. The composition according to claim 4, wherein the antimicrobial agent is 2-phenylethanol, the antimicrobial agent is one or both of propionic acid, and sodium propionate, and the antimicrobial agent is one or both of benzoic acid and sodium benzoate.

6. The composition according to any one of claims 1 to 5, wherein the composition

further comprises one or more of substances selected from a group consisting of diethylstilbestrol, estradiol, estriol, daidzein, aglycone of daidzein, genistin, genistein, glycitein, aglycone of glycitein, biochanin, coumestrol, and formononetin, with a total content of the one or more of substances ranging from 0.001 percent to 1.0 percent (weight/volume).

7. The composition according to any one of claims 1 to 5, wherein the composition further comprises one or more of saccharides selected from a group consisting of glucose, fructose, mannose, sucrose, isomaltulose, 1-kestose, nistose trihydrate, 1F-fructofuranosylmaltose, maltose, isomaltose, isomaltotriose, isomaltotetraose, isomaltopentaose, trehalose, cellobiose, melibiose, gentiobiose, gentiooligosaccharide, raffinose, panose, maltooligosaccharide, isomaltulooligosaccharide, fructooligosaccharide, glucomannan, dextrin, starch, and glycogen, with a total content of the one or more of saccharides ranging from 1.0 percent to 9.0 percent (weight/volume).

8. The composition according to claim 7, wherein the saccharides are one or more of saccharides selected from a group consisting of glucose, fructose, mannose, sucrose, maltose, isomaltulose, isomaltose, trehalose, maltooligosaccharide, with a total content of the one or more of saccharides ranging from 2.0 percent to 6.5 percent (weight/volume).

9. The composition according to any one of claims 1 to 5, wherein the composition further comprises one or more of substances selected from a group consisting of 0.01 to 5.0 percent (weight/volume) of aloe extract, 0.01 to 5.0 percent (weight/volume) of lavender extract, 0.001 to 1.0 percent (weight/volume) of vitamin E, 0.001 to 1.0 percent (weight/volume) of vitamin A, 0.001 to 1.0 percent (weight/volume) of vitamin D, and 0.001 to 1.0 percent (weight/volume) of vitamin C.

10. The composition according to any one of claims 1 to 5, wherein the composition further comprises one or more of amino acids and/or salts thereof, selected from a group consisting of glutamic acid, glutamine, aspartic acid, asparagine, isoleucine, phenylalanine, valine, leucine, proline, and threonine, with a total content of the one or more of amino acids ranging from 0.1 percent to 10.0 percent (weight/volume).

11. The composition according to claim 10, wherein the total content of the amino acids ranges from 1.0 percent to 5.0 percent (weight/volume), and wherein the amino acid is glutamic acid and/or aspartic acid.

12. A method for preparing a composition for vagina, wherein the method comprises adding

an antimicrobial agent, a total content of which, calculated as 2-phenylethanol, is 0.05 to 0.60 percent (weight/volume), and an antimicrobial agent, a total content of which, calculated as sodium propionate, is 0.25 to 0.70 percent (weight/volume); or

an antimicrobial agent, a total content of which, calculated as 2-phenylethanol, is 0.05 to 0.60 percent (weight/volume), an antimicrobial agent, a total content of which, calculated as sodium propionate, is 0.25 to 0.70 percent (weight/volume), and an antimicrobial agent, a total content of which, calculated as sodium benzoate, is 0.05 to 0.15 percent (weight/volume);

into at least one of water, water-soluble gel matrix and emulsive ointment matrix; wherein the antimicrobial agent comprises 2-phenylethanol, the antimicrobial agent comprises at least one of propionic acid, sodium propionate and calcium propionate, and the antimicrobial agent comprises one or both of benzoic acid and sodium benzoate.