

A Pharmaceutical Composition Regulating the Function
of a Living Body ✓

The present invention relates to silicate polymers which regulate the physiological function of a living body, a method for manufacturing thereof, and a pharmaceutical composition containing said polymer as an active ingredient.

A living body maintains its life as an independent organism by controlling physical and chemical conditions of the body within a certain physiological state, and responding to alterations of the internal and external environment.

Such a mechanism maintaining the homeostasis of a living body is effected by the living cells of the body, particularly through their cell membranes. The cell membrane, as is well known, is made up of a phospholipid bilayer, and recognizes various molecules with its surface protein, which acts as a receptor, and selectively transmits chemicals and ions to achieve intra- and extra-cellular equilibria of substances and chemical conditions. These mechanisms maintain the physiological balance of the cell and cause it to act normally.

If said balance is disturbed for any reason, however, various troubles may arise; for instance, a change in the amount of fatty acid components of phospholipids, i.e. a decrease of unsaturated fatty acids or an increase in the amount of unsaturated and saturated fatty acids, may make the cell membrane solid and lower the fluidity of the membrane.

The alteration of the membrane fluidity may result in the hypofunction of the membrane receptors as well as of channels for sodium, potassium, calcium and other ions, which eventually brings about functional disturbance of the cell.

The present inventors have conducted continued studies on the substances which enhance the naturally acquired healing



THIS SHEET HAS BEEN
REPLACED

- 2 -

power of a living body and contribute to normalization of functions of the body, focusing on the homeostatic mechanisms regulating and restoring neurological, endocrinological and immunological disturbances due to functional abnormality of cells in a morbid state, and, as a result thereof, have accomplished the present invention.

The present invention relates to water-soluble silicates which are activated by polymerization. It also relates to biologically active substances comprising said silicate polymers which regulate and maintain the function of a living body by restoring and normalizing the lowered cell functions due to diseases. It further relates to a pharmaceutical composition containing said silicate polymer as an active ingredient, and to a method for manufacturing the same.

Silicon is a natural element widely spread in organisms of the animal and plant kingdoms. In particular, it exists as silicate in animal tissues like hair, feather, bone and skin and is known as an essential element in osteogenesis. In animal tissues, it is involved in cross linkage of collagen tissues and is present in acidic mucopolysaccharide as one of the components. Silicon is thus an essential element for the living body. It is known, additionally, that silicon has pharmacological activities such as immuno-suppression through its anti-macrophage activity and anti-diabetic activity, when administered to animals.

Some silicates like magnesium silicate and aluminum silicate are widely used as anti-acid pharmaceutical products. However, it has not been reported that a silicate polymer has any specific pharmacological activities.

The objects of the present invention are to provide novel silicate polymers which regulate the function of a living body, a method for manufacturing the same and a pharmaceutical composition containing said polymer as an active ingredient.



THIS SHEET HAS BEEN REPLACED

- 3 -

The substances of the invention, which regulate the function of a living body by restoring and normalizing the lowered cell functions due to diseases, are silicate polymers. The silicate polymers of the invention are polymerized substances of silicic acids or silicates and, preferably, are water-soluble silicate polymers.

The silicic acids include orthosilicic acid, metasilicic acid, mesodisilicic acid, mesotrisilicic acid, etc. The silicates are the salts of said silicic acids, for example, salts with alkali metals such as sodium and potassium. A material containing silicic acid, for instance, water glass which is a concentrated aqueous solution of alkali salts of silicic acid, is also used. Further, a silicate solution prepared by heating and dissolving silicon oxide in alkaline aqueous solution may be utilized.

The silicate polymers of the invention have a molecular weight in the range of 4,800 to 2,000,000, preferably, 20,000 to 1,000,000, which are determined by gel-filtration, ultrafiltration, electrophoresis and the like.

The said silicate polymers have a degree of polymerization in the range of 75 to 33,000, preferably, 490 to 16,500.

An example of the manufacturing methods for silicate polymers of the present invention will be given below.

Water-soluble silicate such as sodium orthosilicate, sodium metasilicate, potassium orthosilicate or potassium metasilicate, or silicic acid containing material such as water glass is dissolved in an aqueous solution. Since said aqueous solution of silicate displays a high pH value, it is preferable to adjust the solution to pH 2-10, more preferably, pH 4-9.5, by use of a conventional acid such as hydrochloric acid, sulfuric acid or acetic acid. It is desirable to add a saccharide such as lactose, mannitol, sorbitol, sucrose, glucose, fructose or galactose to the aqueous solution. A salt

such as sodium chloride, potassium chloride or sodium sulfate may also be added.

For the purpose of manufacturing a pharmaceutical composition of the invention, said aqueous solution of silicate is preferably dried to a powder. The powderization may be carried out according to a conventional method such as heating or lyophilization. To give a preferred powder, for example, the solution is dried by heating at from 150 to 250°C. Also a conventional lyophilization under reduced pressure may be used to powderize the solution.

To detect the production of silicate polymers, for example, a molybdenum blue coloration is measured. Namely, a solution of ammonium molybdate is added to the aqueous solution of silicate and then sulfite solution is added to produce a blue coloration. The blue coloration reduces as the polymerization of silicate proceeds. Also, the silicate polymers so produced can be separated by gel-filtration under acidic conditions at high molecular bands. An aliquot of this separated fraction is decomposed under alkaline conditions to detect silicate by the molybdenum blue reaction.

Examples

Example 1

7.6 g of water glass (silicon; 1.2 g) was dissolved in 100 ml of water. 97.5 g of lactose was dissolved in 300 ml of water with heating. The water glass solution was mixed with the lactose solution, and then the pH was adjusted to 8.0 with diluted hydrochloric acid. The reaction mixture was dried at 200°C to give 90 g of a powder. The resulting silicate polymers of the present invention have the following physical properties and contain 12 mg of silicon per 1 g.

Molecular weight:	13,000 to 1,000,000
Degree of Polymerization:	210 to 16,500

Example 2

12.9 g of sodium metasilicate (silicon; 1.2 g) was dissolved in 100 ml of water. 95.8 g of lactose was dissolved in 300 ml of water with heating. The aqueous solution of sodium metasilicate was mixed with the lactose solution, and then the pH was adjusted to 8.0 with diluted hydrochloric acid. The reaction mixture was dried at 200°C to give 90 g of a powder. The resulting silicate polymers of the present invention have the following physical properties and contain 12 mg of silicon per 1 g.

Molecular weight:	15,000 to 900,000
Degree of Polymerization:	250 to 15,000

Example 3

5.9 g of sodium orthosilicate (silicon; 0.6 g) was dissolved in 100 ml of water. 99 g of mannitol was dissolved in 300 ml of water with heating. The aqueous solution of sodium orthosilicate was mixed with the mannitol solution, and then the pH was adjusted to 8.0 with diluted hydrochloric acid. The reaction mixture was dried at 200°C to give 88 g of a powder. The resulting silicate polymers of the present invention have the following physical properties and contain 6 mg of silicon per 1 g.

Molecular weight:	20,000 to 1,000,000
Degree of Polymerization:	330 to 16,500

Example 4

1.89 g of water glass (silicon; 0.3 g) was dissolved in 100 ml of water. 500 ml of 20 % aqueous solution of lactose was added thereto, and then the pH was adjusted to 8.0 with diluted hydrochloric acid. Each 1.2 ml of the solution was pipetted into a vial and lyophilized. The resulting dry-powdered

silicate polymers of this invention have the following physical properties and contain 0.6 mg of silicon per vial.

Molecular weight:	30,000 to 1,500,000
Degree of Polymerization:	490 to 25,000

In the following experiments, the amount of substance of the present invention was indicated as the amount of silicon (Si) contained therein, such as Si g/ml or Si mg/kg.

1. Inhibitory effect on mast cell degranulation:

An inhibitory effect of a substance of the invention on mast cell degranulation was determined according to a usual method. A solution of the substances was added to a cell suspension of rat peritoneal mast cells and incubated at 37°C for 10 minutes. To this cell suspension, a solution of Compound 48/80, a histamine releaser, was added. The reaction mixture was incubated for 10 minutes, and then the reaction was stopped by adding a cold buffer solution. The supernatant was separated by centrifuging, and the amount of histamine in it was measured by fluorometric methods. The inhibition rate of the substance on histamine release was determined according to a conventional method. The substance significantly inhibited the histamine release from mast cells.

An example of the results is shown in Table 1.

Table 1

Concentration of the substance of Example 1 (Si g/ml)	Inhibition rate on histamine release (%)
1×10^{-8}	6.2 ± 2.0
1×10^{-7}	7.6 ± 8.5
1×10^{-6}	37.6 ± 0.7
1×10^{-5}	83.0 ± 11.9
1×10^{-4}	96.2 ± 0.9

2. Inhibitory effect on histamine release by antigen-antibody reaction:

Rats were passively sensitized by intraperitoneal injection of anti-ovalbumin serum (IgE-like antibody). Twenty-four hours thereafter, a suspension of peritoneal mast cells of the rats was prepared. To this cell suspension, a solution of the substance of the invention was added, and incubated for 10 minutes at 37°C. A solution of ovalbumin was then added to the mixture to induce antigen-antibody reaction. The reaction was stopped after 10 minutes and the inhibition rate of histamine release by the substance was measured, according to the methods described above, in 1. The substance significantly inhibited the histamine release from mast cells caused by antigen-antibody reaction.

An example of the results is shown in Table 2---

Table 2

Concentration of the substance of Example 2 (Si g/ml)	Inhibition rate on histamine release (%)
1×10^{-8}	3.1 ± 1.1
1×10^{-7}	16.6 ± 12.2
1×10^{-6}	85.1 ± 5.9
1×10^{-5}	96.5 ± 4.5
1×10^{-4}	98.4 ± 2.8

3. Inhibitory effect on hyaluronidase activity:

A substance of the present invention was added to a solution containing hyaluronidase, and incubated at 37°C for 10 minutes. A solution of hyaluronic acid as a substrate was added to the solution to react with the enzyme for 15 minutes, and then the reaction was terminated by addition of a succinic acid-buffer solution. The absorbance of the reaction solution at 540 nm was measured. The inhibition rate of the substance to hyaluronidase was calculated in comparison with the absorbance of a control solution wherein a physiological saline solution was used instead of the tested substance. The substance showed inhibition of hyaluronidase activity dose-dependently.

An example of the results is shown in Table 3.

Table 3

Concentration of the substance of Example 2 (Si μ g/ml)	Inhibition rate to hyaluronidase (%)
50	10.9
100	24.5
250	51.7
500	89.1

4. Inhibitory effect on vascular permeability:

Guinea pig nasal cavity was perfused with histamine solution, and the increase of nasal membrane permeability and of nasal secretion was investigated by measuring the amount of dye extravasated at the site according to the method of Kojima and Tsutsumi [Allergy, vol. 35, 180-187 (1986)]. A polyethylene tube was inserted into the nasal cavity from an incision in the trachea for perfusion, and a solution of dye was intravenously injected. A physiological saline solution was perfused at 1 ml per minute. The initial perfusate for 10 minutes was drained off from the nostrils for washing, and the perfusate of the following 20 minutes was collected (Perfusate A). A physiological saline solution containing histamine was then perfused for 10 minutes to collect the perfusate (Perfusate B). A normal saline solution was again perfused for 10 minutes to collect the perfusate (Perfusate C). Each perfusate was centrifuged and the absorbance of the supernatant was measured at 620 nm.

A substance of the invention, which was intraperitoneally administered 30 minutes before histamine perfusion, significantly suppressed the permeability of the nasal membrane to the dye.

An example of the results is shown in Table 4. The amount of dye in the perfusate was expressed as the absorbance.

Table 4

Amount of dye in the perfusate			
		Dose of the substance of Example 2 (Si mg/kg)	
Perfusate	Saline	12.5	25.0
A	0.88 ± 0.44	0.45 ± 0.13	0.33 ± 0.09
B	4.07 ± 2.40	1.34 ± 1.12	0.75 ± 0.46
C	3.82 ± 2.05	1.72 ± 1.75	0.88 ± 0.56

5. Effect on disturbance of peripheral blood circulation:

The effect of a substance of the invention on the disturbance of peripheral blood circulation was determined, as an index of improving paresthesia. Quinoform was administered to rats intraperitoneally for 27 days in progressively increased amounts to cause disturbance of peripheral circulation. The hind limbs of the quinoform-treated rats were then exposed to cold water at 5°C for 2 minutes. The effect of the substance in improving peripheral blood circulation was evaluated by monitoring the recovering process of the limb temperature with thermographical image analysis. The substance was intravenously administered to rats consecutively for 7 days from the 21st day of quinoform treatment.

An example of the results is shown in Table 5. The dose of the substance is expressed as Si mg/kg and is shown in parenthesis.

Table 5

	Average skin temperature (°C) of the hind limb at 15 minutes after exposure to cold water
Untreated group	27.7 ± 0.7
Control: quinoform-treated	23.9 ± 0.5
The substance of Example 1	
(5)	24.2 ± 0.2
(10)	26.3 ± 0.2
(20)	27.0 ± 0.3

6. Analgesic effect:

Analgesic effects of a substance of the present invention was studied by Randall-Selitto method using SART-Stressed mice prepared according to Kita, et al. [Folia Pharmacol. Jap., vol. 71, 195-210 (1975)]. The SART-stressed animal is an animal model of hyperalgesia, with lowered threshold to pain sensation, and highly responsive when applying pressure stimulus to the tail root of the animal by a Randall-Selitto device.

The effect of the substance was evaluated by an analgesic index: the quotient of the pain threshold after treatment divided by that before treatment. The lowered pain threshold of SART-stressed mice was elevated dose-dependently up to normal level, when the substance of the present invention was intraperitoneally administered.

An example of the results is shown in Table 6.

Table 6

Dose of the substance of Example 3 (Si mg/kg)	Average pain threshold
2.5	1.35
5.0	1.51
10.0	1.70

7. Acute toxicity:

Acute toxicity of a substance of the present invention was measured by administering the substance intravenously or intraperitoneally to a group of 10 ddY male mice. The LD₅₀ values of the substance were 60.4 Si mg/kg (i.v.) and 71.8 Si mg/kg (i.p.), respectively.

As apparently shown by the above-mentioned results of pharmacological studies, the silicate polymers of the present invention have biological activities which restore and normalize the abnormal functions of a living body due to diseases. For example, the substances of this invention show a therapeutic effect on diseased conditions such as allergy or inflammation, and elevate the lowered pain threshold in animals under stress.

The substances of the present invention have, as demonstrated, an effect on neurological, endocrinological and immunological systems and show an excellent activity in restoring and normalizing the impaired functions of living body due to diseases. They are very useful as drugs such as anti-allergic agents, anti-inflammatory agents, improving agents for peripheral blood circulation and paresthesia, and analgesic agents. Consequently, a pharmaceutical composition containing a substance of the invention as an active ingredient is useful for treating or preventing various diseases accompanying allergy, inflammation or pain, including bronchial asthma,

pollinosis, allergic rhinitis, urticaria, eczema, contact dermatitis, allergic conjunctivitis, gastritis, peptic ulcer, enterocolitis, idiopathic ulcerative colitis, rheumatoid arthritis, herpes, systemic lupus erythematosus, gastrointestinal neurosis, autonomic dystonia, vertigo, neuralgia, frozen shoulder, low back pain, headache, etc.

The substances of the present invention can be made into pharmaceuticals by combining with suitable carriers or diluents for medical use and can be made into pharmaceutical preparations by any conventional methods giving solid, semisolid, liquid or gaseous forms for oral or parenteral administration.

In manufacturing such preparations, the substances of the present invention may be used either solely or jointly in the form of a suitable combination. Alternatively, the substances may be combined with other pharmaceutically active components.

In the case of preparations for oral use, the substances of the present invention may be made into tablets, diluted powders, granules or capsules with or without suitable additives such as conventional fillers (e.g. lactose, mannitol, corn starch and potato starch) as well as binders (e.g. crystalline cellulose, cellulose derivatives, gum arabic, corn starch and gelatin), lubricants (e.g. talc and magnesium stearate), extenders, moisturizers, preservatives and perfumes.

Alternatively, the substances of the present invention may be mixed with a base material of fat/oil type (e.g. cacao butter), emulsifying base material, water-soluble base material (e.g. Macrogol), hydrophilic base material, etc. to give a suppository.

In the case of injections, the substances may be made into a solution or a suspension in aqueous solvents or nonaqueous ones such as distilled water for injection, physiological

saline liquid, Ringer solution, plant oil, synthetic fatty acid glycerides, higher fatty acid esters and propylene glycol.

Further, depending upon the state of the patient, or the type of the disease, the substances may be made into other preparation forms such as ointments and eye drops which are most suitable for the therapy.

A desired dose of the substances of the present invention may vary depending upon the patient to be treated, preparation form, method of administration, period for administration, etc. In general, 1 μ g/kg to 10 mg/kg per day may be given to an adult for achieving the desired effect.

In the case of parenteral administration such as injections, the desired dose may be 1/3 to 1/10 as large as an oral dose in general because of the difference of absorption.

Some prescriptions of the pharmaceutical compositions which contain the substances of the present invention as active ingredients are shown below as examples. These examples, however, do not limit the present invention.

Prescription example 1 (tablet)	
Component	Content in a tablet (mg)
substance of this invention	100
lactose	58
Crystalline cellulose	40
magnesium stearate	2
Total	200

Prescription example 2 (capsule)	
Component	Content in a capsule (mg)
substance of this invention	50
lactose	78
potato starch	15
magnesium stearate	2
talc	5
Total	150

Prescription example 3 (injection)	
Component	Content in an ampule (mg)
substance of this invention	10
saline	proper amount
Total	1 ml

Prescription example 4 (ointment)	
Component	(g)
substance of this invention	1
glycerol monostearate	5
white vaseline	94
Total	100

Prescription example 5 (suppository)	
Component	(g)
substance of this invention	0.1
witepsol H-15	1.9
Total	2.0

This Sheet Has Been Replaced



- 17 -


Claims

THIS SHEET HAS BEEN
REPLACED

1. A pharmaceutical composition containing a silicate polymer as an active ingredient, which regulates the functions of a living body.
2. The pharmaceutical composition is an anti-allergic agent according to claim 1.
3. The pharmaceutical composition is an anti-inflammatory agent according to claim 1.
4. The pharmaceutical composition is an improving agent for peripheral blood circulation according to claim 1.
5. The pharmaceutical composition is an improving agent for paresthesia according to claim 1.
6. The pharmaceutical composition is an analgesic agent according to claim 1.
7. The pharmaceutical composition according to any of the preceding claims wherein the silicate polymer consists of water-soluble silicates.
8. The pharmaceutical composition according to claim 7 wherein the water-soluble silicates are salts with alkali metals.
9. The pharmaceutical composition according to claims 7 or 8 wherein the silicates are sodium silicates.
10. The pharmaceutical composition according to any of claims 1 to 7 wherein the silicate polymer consists of water glass.



This Sheet Has Been Replaced



- 18 -

11. The pharmaceutical composition according to any of the preceding claims wherein the silicate polymer has a molecular weight in the range of 4,800 to 2,000,000.
12. The pharmaceutical composition according to any of the preceding claims wherein the silicate polymer has a degree of polymerization in the range of 75 to 16,500.
13. A process for manufacturing a silicate polymer wherein silicic acid or silicate is dissolved in aqueous solvent, adjusted to pH 4-10, and dried.
14. The process according to claim 13 wherein a saccharide is added to the silicate solution.
15. The process according to claims 13 or 14 wherein the silicate aqueous solution is dried to a powder.
16. The process according to claims 13, 14 or 15 wherein the drying method is lyophilization.
17. The use of a silicate polymer in the manufacture of a medicament effective in regulating the function of a living body.
18. The use according to claim 17 wherein the medicament is an effective anti-allergy agent, anti-inflammatory agent, agent for improving peripheral blood circulation, agent for improving paresthesia or an analgesic agent.

THIS SHEET HAS BEEN
REPLACED



* C L R S *

Abstract

In the present invention, a pharmaceutical composition containing a silicate polymer as an active ingredient which regulates the function of a living body is provided. The silicate polymer of the composition may be manufactured in a process wherein silicic acid or silicate is dissolved in an aqueous solvent, adjusted to pH 4 to 10, and dried. The silicate polymer thus produced is used in the manufacture of medicines effective in regulating the function of living bodies.

THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



A PHARMACEUTICAL COMPOSITION REGULATING THE FUNCTION
OF A LIVING BODY

ABSTRACT

In the present invention, a pharmaceutical composition containing a silicate polymer as an active ingredient which regulates the function of a living body is provided. The silicate polymer of the composition may be manufactured in a process wherein silicic acid or silicate is dissolved in an aqueous solvent, adjusted to pH 4 to 10, and dried. The silicate polymer thus produced is used in the manufacture of medicines effective in regulating the function of living bodies.

THIS SHEET HAS BEEN
REPLACED

REPLACEMENT SHEET

07 OCT 1996

This Sheet Has Been Replaced



power of a living body and contribute to normalization of functions of the body, focusing on the homeostatic mechanisms regulating and restoring neurological, endocrinological and immunological disturbances due to functional abnormality of cells in a morbid state, and, as a result thereof, have accomplished the present invention.

The present invention relates to water-soluble silicates which are activated by polymerization. It also relates to biologically active substances comprising said silicate polymers which regulate and maintain the function of a living body by restoring and normalizing the lowered cell functions due to diseases. It further relates to a pharmaceutical composition containing said silicate polymer as an active ingredient, and to a method for manufacturing the same.

Silicon is a natural element widely spread in organisms of the animal and plant kingdoms. In particular, it exists as silicate in animal tissues like hair, feather, bone and skin and is known as an essential element in osteogenesis. In animal tissues, it is involved in cross linkage of collagen tissues and is present in acidic mucopolysaccharide as one of the components. Silicon is thus an essential element for the living body. It is known, additionally, that silicon has pharmacological activities such as immuno-suppression through its anti-macrophage activity and anti-diabetic activity, when administered to animals.

Some silicates like magnesium silicate and aluminum silicate are widely used as anti-acid pharmaceutical products. However, it has not been reported that a silicate polymer has any specific pharmacological activities.

The objects of the present invention are to provide novel silicate polymers which regulate the function of a living body, a method for manufacturing the same and a pharmaceutical composition containing said polymer as an active ingredient.

According to one aspect of the present invention there is provided a pharmaceutical composition for use in treating diseases accompanied by or associated with allergy, inflammation or pain or for use in improving peripheral blood circulation or paresthesia comprising a pharmaceutically effective amount of a water-soluble silicate polymer as an active ingredient, and an inert carrier or diluent comprising at least one member selected from the group consisting of saccharides and sugar alcohols, said silicate polymer having a molecular weight in the range of 4,800 to 2,000,000 as determined by gel-filtration and a degree of polymerization in the range of 75 to 33,000 wherein the monomer unit is —(SiO₂)— .

According to another aspect of the present invention there is provided a pharmaceutical composition comprising a pharmaceutically effective amount of a water-soluble silicate polymer having a molecular weight in the range of 4,800 to 2,000,000 as determined by gel-filtration and a degree of polymerization in the range of 75 to 33,000,

REPLACEMENT SHEET

11 DEC 1997

wherein the monomer unit is $\text{---}(\text{SiO}_2)\text{---}$, as an active ingredient, and an inert carrier or diluent comprising a saccharide.

An example of the manufacturing method for silicate polymers of the present invention will be given below.

Water-soluble silicate such as sodium orthosilicate, sodium metasilicate, potassium orthosilicate or potassium metasilicate, or silicic acid containing material such as water glass is dissolved in an aqueous solution. Since said aqueous solution of silicate displays a high pH value, it is preferable to adjust the solution to pH 2-10, more preferably, pH 4-9.5, by use of a conventional acid such as hydrochloric acid, sulfuric acid or acetic acid. It is desirable to add a saccharide such as lactose, mannitol, sorbitol, sucrose, glucose, fructose or galactose to the aqueous solution. A salt

REPLACEMENT SHEET


11 DEC 1997

CLAIMS

1. A pharmaceutical composition for use in treating diseases accompanied by or associated with allergy, inflammation or pain or for use in improving peripheral blood circulation or paresthesia comprising a pharmaceutically effective amount of a water-soluble silicate polymer as an active ingredient, and an inert carrier or diluent comprising at least one member selected from the group consisting of saccharides and sugar alcohols, said silicate polymer having a molecular weight in the range of 4,800 to 2,000,000 as determined by gel-filtration and a degree of polymerization in the range of 75 to 33,000 wherein the monomer unit is $\text{---}(\text{SiO}_2)\text{---}$.
2. A pharmaceutical composition as claimed in claim 1 which is an anti-allergic agent.
3. A pharmaceutical composition as claimed in claim 1 which is an anti-inflammatory agent.
4. A pharmaceutical composition as claimed in claim 1 which is an improving agent for peripheral blood circulation.
5. A pharmaceutical composition as claimed in claim 1 which is an improving agent for paresthesia.
6. A pharmaceutical composition as claimed in claim 1 which is an analgesic agent.
7. A pharmaceutical composition comprising a pharmaceutically effective amount of a water-soluble silicate polymer having a molecular weight in the range of 4,800 to 2,000,000 as determined by gel-filtration and a degree of polymerization in the range of 75 to 33,000, wherein the monomer unit is $\text{---}(\text{SiO}_2)\text{---}$, as an active ingredient, and an inert carrier or diluent comprising a saccharide.
8. A pharmaceutical composition as claimed in claim 7 wherein the silicate polymer is a polymerization product of water-soluble silicates.
9. A pharmaceutical composition as claimed in claim 8 wherein the water-soluble silicates are salts of silicic acid with alkali metals.
10. A pharmaceutical composition as claimed in claim 9 wherein the silicates are sodium silicates.
11. A pharmaceutical composition as claimed in claim 9 wherein the silicate polymer is a polymerization product of water glass.
12. A pharmaceutical composition as claimed in claim 7 wherein the saccharide is lactose.



* C L H S *



A PHARMACEUTICAL COMPOSITION REGULATING THE FUNCTION
OF A LIVING BODY

ABSTRACT

In the present invention, a pharmaceutical composition containing a silicate polymer as an active ingredient which regulates the function of a living body is provided. The silicate polymer of the composition may be manufactured in a process wherein silicic acid or silicate is dissolved in an aqueous solvent, adjusted to pH 4 to 10, and dried. The silicate polymer thus produced is used in the manufacture of medicines effective in regulating the function of living bodies.

