

FN3 Domain-siRNA Conjugates and Uses Thereof

RELATED APPLICATIONS

[001] The present application claims priority to U.S. Provisional Patent Application No. 63/174,776, filed April 14, 2021, U.S. Provisional Patent Application No. 63/203,776, filed July 30, 2021, and U.S. Provisional Application No. 63/324,437, filed March 28, 2022, which are hereby incorporated by reference in their entirety.

FIELD

[002] The present embodiments relate to siRNA molecules that can be conjugated fibronectin type III domains (FN3) and methods of making and using the molecules.

BACKGROUND

[003] Therapeutic nucleic acids include, e.g., small interfering RNA (siRNA), micro RNA (miRNA), antisense oligonucleotides, ribozymes, plasmids, immune stimulating nucleic acids, antisense, antagomir, antimir, microRNA mimic, supermir, U1 adaptor, and aptamer. In the case of siRNA or miRNA, these nucleic acids can down-regulate intracellular levels of specific proteins through a process termed RNA interference (RNAi). The therapeutic applications of RNAi are extremely broad, since siRNA and miRNA constructs can be synthesized with any nucleotide sequence directed against a target protein. To date, siRNA constructs have shown the ability to specifically down-regulate target proteins in both in vitro and in vivo models. In addition, siRNA constructs are currently being evaluated in clinical studies and have been approved for a variety of diseases..

[004] However, two problems currently faced by siRNA constructs are, first, their susceptibility to nuclease digestion in plasma and, second, their limited ability to gain access to the intracellular compartment where they can bind the RISC (RNA-induced Silencing Complex) when administered systemically as the free siRNA or miRNA. Certain delivery systems, such as lipid nanoparticles formed from cationic lipids with other lipid components, such as cholesterol and PEG lipids, carbohydrates (such as GalNac trimers) have been used to facilitate the cellular uptake of the oligonucleotides. However, these have not been shown to be successful in efficiently and effectively delivering siRNA to its intended target in tissues other than the liver.

[005] There remains a need for compositions and methods for delivering siRNA to its intended cellular target. Further, what is needed is a FN3 domain with optimized properties for clinical use that can specifically bind to CD71, and methods of using such molecules for novel

therapeutics that enable intracellular access via receptor mediated internalization of CD71. The present embodiments fulfill these needs as well as others.

SUMMARY

[006] In some embodiments, siRNA conjugated to FN3 domains that bind CD71 protein are provided.

[007] In some embodiments, FN3 domains are provided that comprise the amino acid sequence of any FN3 domain provided herein. In some embodiments, the FN3 domains bind to CD71. In some embodiments, the FN3 domains specifically bind to CD71.

[008] In some embodiments, the composition comprises two FN3 domains connected by a linker, such as a flexible linker. In some embodiments, the two FN3 domains bind to different targets. In some embodiments, a first FN3 domain binds to CD71. In some embodiments, a second FN3 domain binds to a different target that is not CD71.

[009] In some embodiments, oligonucleotides, such as dsRNA or siRNA molecules are provided herein. In some embodiments, the oligonucleotides have the sequences as provided herein, with or without the modifications provided herein. In some embodiments, the oligonucleotides are provided in a composition, such as a pharmaceutical composition. In some embodiments, the oligonucleotides are conjugated to a polypeptide.

[0010] In some embodiments, composition comprising one or more FN3 domains conjugated to a siRNA molecule are provided.

[0011] In some embodiments, a composition having a formula of $(X1)_n-(X2)_q-(X3)_y-L-X4$, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; and X4 is an oligonucleotide molecule, wherein n, q, and y are each independently 0 or 1, are provided.

[0012] In some embodiments, a composition having a formula of $C-(X1)_n-(X2)_q-(X3)_y-L-X4$, wherein C is a polymer or albumin binding domain (ABD); X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; and X4 is an oligonucleotide molecule, wherein n, q, and y are each independently 0 or 1, are provided.

[0013] In some embodiments, a composition having a formula of $(X1)_n-(X2)_q-(X3)_y-L-X4-C$, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; X4 is an oligonucleotide molecule; and C is a polymer or albumin binding domain (ABD), wherein n, q, and y are each independently 0 or 1, are provided.

[0014] In some embodiments, a composition having a formula of $X4-L-(X1)_n-(X2)_q-(X3)_y$, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; and X4 is an oligonucleotide molecule, wherein n, q, and y are each independently 0 or 1, are provided.

[0015] In some embodiments, a composition having a formula of $C-X4-L-(X1)_n-(X2)_q-(X3)_y$, wherein C is a polymer or albumin binding domain (ABD); X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; and X4 is an oligonucleotide molecule, wherein n, q, and y are each independently 0 or 1, are provided.

[0016] In some embodiments, a composition having a formula of $X4-L-(X1)_n-(X2)_q-(X3)_y-C$, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; X4 is an oligonucleotide molecule; and C is a polymer or albumin binding domain (ABD), wherein n, q, and y are each independently 0 or 1, are provided.

[0017] In some embodiments, a composition having a formula of $C-(X1)_n-(X2)_q[L-X4]-(X3)_y$, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; X4 is an oligonucleotide molecule; and C is a polymer or albumin binding domain (ABD), wherein n, q, and y are each independently 0 or 1, are provided.

[0018] In some embodiments, a composition having a formula of $(X1)_n-(X2)_q[L-X4]-(X3)_y-C$, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; X4 is an oligonucleotide molecule; and C is a polymer or albumin binding domain (ABD), wherein n, q, and y are each independently 0 or 1, are provided.

[0019] In some embodiments, pharmaceutical compositions comprising one or more of the compositions provided herein are provided.

[0020] In some embodiments, methods of treating Pompe Disease (GSD2, acid alpha-glucosidase (GAA) deficiency) in a subject in need thereof, the method comprising administering a composition provided herein are provided.

[0021] In some embodiments, methods of treating glycogen storage disease in a subject in need thereof, the method comprising administering a composition provided herein are provided. In some embodiments, the glycogen storage disease is selected from the group consisting of Cori's disease or Forbes' disease (GSD3, Glycogen debranching enzyme (AGL) deficiency), McArdle disease (GSD5, Muscle glycogen phosphorylase (PYGM) deficiency), type II Diabetes/diabetic

nephropathy, Aldolase A Deficiency GSD12, Lafora Disease, hypoxia, Andersen disease (GSD4, Glycogen debranching enzyme (GBE1) deficiency), Tarui's Disease (GSD7, Muscle phosphofructokinase (PFKM) deficiency), and adult polyglucosan body disease. In some embodiments, the glycogen storage disease is selected from the group consisting of Glycogen synthase (GYS2) deficiency (GSD0), Glucose-6-phosphatase (G6PC / SLC37A4) deficiency (GSD1, von Gierke's disease), Hers' disease (GSD6, Liver glycogen phosphorylase (PYGL) or Muscle phosphoglycerate mutase (PGAM2) deficiency), Phosphorylase kinase (PHKA2 / PHKB / PHKG2 / PHKA1) deficiency (GSD9), Phosphoglycerate mutase (PGAM2) deficiency (GSD10), Muscle lactate dehydrogenase (LDHA) deficiency (GSD11), Fanconi-Bickel syndrome (GSD 11, Glucose transporter (GLUT2) deficiency, Aldolase A deficiency (GSD 12), β -enolase (ENO3) deficiency (GSD13), and Glycogenin-1 (GYG1) deficiency (GSD15).

[0022] In some embodiments, methods of treating cancer in a subject in need thereof, the method comprising administering to the subject a composition provided herein are provided.

[0023] In some embodiments, methods of treating a neurological condition and/or a brain tumor in a subject in need thereof, the method comprising administering to the subject a composition provided herein are provided. In some embodiments, the neurological condition is selected from the group consisting of Alzheimer's Disease, Amyotrophic Lateral Sclerosis, Parkinson's Disease, Lafora Disease, Pompe Disease, adult polyglucosan body disease, stroke, spinal cord injury, ataxia, Bell's Palsy, cerebral aneurysm, epilepsy, seizures, Guillain-Barre Syndrome, multiple sclerosis, muscular dystrophy, neurocutaneous syndromes, migraine, encephalitis, septicemia, and myasthenia gravis.

[0024] In some embodiments, methods of treating an autoimmune disease in a subject in need thereof, the method comprising administering to the subject a composition provided herein are provided. In some embodiments, the autoimmune disease is selected from the group consisting of rheumatoid arthritis, Hashimoto's autoimmune thyroiditis, celiac disease, diabetes mellitus type 1, vitiligo, rheumatic fever, pernicious anemia/atrophic gastritis, alopecia areata, and immune thrombocytopenic purpura.

[0025] In some embodiments, a use of a composition as provided herein or of any of in the preparation of a pharmaceutical composition or medicament for treating cancer are provided. In some embodiments, the cancer is selected from the group consisting of acute myeloid leukemia, myelodysplastic syndromes, gastric cancer, clear cell renal cell carcinoma, clear cell carcinomas of the breast, clear cell carcinomas of the endometrium, clear cell carcinomas of the ovary, clear cell carcinomas of the uterus, hepatocellular carcinoma, pancreatic cancer, prostate cancer, soft tissue cancer, Ewings sarcoma, and non-small cell lung cancer

[0026] In some embodiments, methods of reducing the expression of a target gene in a cell, the method comprising contacting the cell with a composition as provided herein are provided. In some embodiments, the cell is a tumor cell, a liver cell, a muscle cell, an immune cell, a dendritic cell, a heart cell, or a cell of the CNS.

[0027] In some embodiments, methods of selectively reducing GYS1 mRNA and protein in skeletal muscle. In certain embodiments, GYS1 mRNA and protein is not reduced in the liver and/or the kidney.

[0028] In some embodiments, isolated polynucleotides encoding the FN3 domains described herein are provided.

[0029] In some embodiments, a vector comprising the polynucleotides described herein are provided.

[0030] In some embodiments, a host cell comprising the vectors described herein are provided.

[0031] In some embodiments, methods of producing the FN3 domains are provided. In some embodiments, the method comprises culturing a host cell comprising a vector encoding or expressing the FN3 domain. In some embodiments, the method further comprises purifying the FN3 domain. In some embodiments, the FN3 domain binds CD71.

[0032] In some embodiments, pharmaceutical compositions comprising a FN3 domain that binds to CD71 linked to an oligonucleotide molecule and a pharmaceutically acceptable carrier are provided. In some embodiments, kits comprising one or more of the FN3 domains with or without the oligonucleotide molecules are provided.

DESCRIPTION OF THE DRAWINGS

[0033] FIG. 1 is a flow chart representing the properties assessed and considered for siRNA screening.

[0034] FIG. 2 is a graph of an RNA sequencing experiment which identifies the changes in the transcriptome after cells are transfected with siRNA pair HHH, wherein the arrow identifies the significant decrease in GYS1 transcript.

[0035] FIG. 3 provides the results of a target binding assay using over 6,000 receptors in the proteome array, wherein the data demonstrates that CD71 is the exclusive binding target of the FN3 domain.

[0036] FIG. 4A demonstrates the knockdown of GYS1 mRNA in mouse gastrocnemius muscle using 3 difference FN3 domain-siRNA conjugates compared with vehicle alone. FIG. 4B demonstrates the knockdown of GYS1 protein in mouse gastrocnemius muscle using 3 difference FN3 domain-siRNA conjugates compared with vehicle alone.

[0037] FIG. 5 demonstrates the GYS1 knockdown is highly specific for skeletal muscle using 3 different FN3 domain-siRNA conjugates compared with a siRNA to a different target (AHA-1).

[0038] FIG. 6 is an example of Histrap chromatogram for purification of conjugates (Tagged proteins).

[0039] FIG. 7 is an example of HIC chromatogram for purification of conjugate (tagless proteins).

[0040] FIG. 8 is an example of Ion-exchange chromatogram for purification of conjugate (Tagged/tagless protein).

[0041] FIG. 9 is an example of an analytical SEC of the centyrin-oligonucleotide conjugate.

[0042] FIG. 10 is an example of SDS PAGE Gel of the conjugate.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0043] As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to “a cell” includes a combination of two or more cells, and the like.

[0044] “Fibronectin type III (FN3) domain” (FN3 domain) refers to a domain occurring frequently in proteins including fibronectins, tenascin, intracellular cytoskeletal proteins, cytokine receptors and prokaryotic enzymes (Bork and Doolittle, *Proc Nat Acad Sci USA* 89:8990-8994, 1992; Meinke *et al.*, *J Bacteriol* 175:1910-1918, 1993; Watanabe *et al.*, *J Biol Chem* 265:15659-15665, 1990). Exemplary FN3 domains are the 15 different FN3 domains present in human tenascin C, the 15 different FN3 domains present in human fibronectin (FN), and non-natural synthetic FN3 domains as described for example in U.S. Pat. No. 8,278,419. Individual FN3 domains are referred to by domain number and protein name, e.g., the 3rd FN3 domain of tenascin (TN3), or the 10th FN3 domain of fibronectin (FN10). As used throughout, “centyrin” also refers to a FN3 domain. Further, FN3 domains as described herein are not antibodies as they do not have the structure of a variable heavy (V_H) and/or light (V_L) chain.

[0045] As used herein, “autoimmune disease” refers to disease conditions and states wherein the immune response of an individual is directed against the individual's own constituents, resulting in an undesirable and often debilitating condition. As used herein, “autoimmune disease” is intended to further include autoimmune conditions, syndromes, and the like. Autoimmune diseases include, but are not limited to, Addison's disease, allergy, allergic rhinitis, ankylosing spondylitis, asthma, atherosclerosis, autoimmune diseases of the ear, autoimmune diseases of the eye, autoimmune atrophic gastritis, autoimmune hepatitis, autoimmune hemolytic anemia, autoimmune parotitis, autoimmune uveitis, celiac disease, primary biliary cirrhosis, benign lymphocytic anitis, COPD, colitis, coronary heart disease, Crohn's disease, diabetes (Type I),

depression, diabetes, including Type 1 and/or Type 2 diabetes, epididymitis, glomerulonephritis, Goodpasture's syndrome, Graves' disease, Guillain-Barre syndrome, Hashimoto's disease, hemolytic anemia, idiopathic thrombocytopenic purpura, inflammatory bowel disease (IBD), immune response to recombinant drug products, e.g., factor VII in hemophilia, juvenile idiopathic arthritis, systemic lupus erythematosus, lupus nephritis, male infertility, mixed connective tissue disease, multiple sclerosis, myasthenia gravis, oncology, osteoarthritis, pain, primary myxedema, pemphigus, pernicious anemia, polymyositis, psoriasis, psoriatic arthritis, reactive arthritis, rheumatic fever, rheumatoid arthritis, sarcoidosis, scleroderma, Sjogren's syndrome, spondyloarthropathies, sympathetic ophthalmia, T-cell lymphoma, T-cell acute lymphoblastic leukemia, testicular antio-centric T-cell lymphoma, thyroiditis, transplant rejection, ulcerative colitis, autoimmune uveitis, and vasculitis. Autoimmune diseases include, but are not limited to, conditions in which the tissue affected is the primary target, and in some cases, the secondary target. Such conditions include, but are not limited to, AIDS, atopic allergy, bronchial asthma, eczema, leprosy, schizophrenia, inherited depression, transplantation of tissues and organs, chronic fatigue syndrome, Alzheimer's disease, Parkinson's disease, myocardial infarction, stroke, autism, epilepsy, Arthus's phenomenon, anaphylaxis, and alcohol and drug addiction.

[0046] The term “capture agent” refers to substances that bind to a particular type of cells and enable the isolation of that cell from other cells. Exemplary capture agents are magnetic beads, ferrofluids, encapsulating reagents, molecules that bind the particular cell type and the like.

[0047] “Sample” refers to a collection of similar fluids, cells, or tissues isolated from a subject, as well as fluids, cells, or tissues present within a subject. Exemplary samples are tissue biopsies, fine needle aspirations, surgically resected tissue, organ cultures, cell cultures and biological fluids such as blood, serum and serosal fluids, plasma, lymph, urine, saliva, cystic fluid, tear drops, feces, sputum, mucosal secretions of the secretory tissues and organs, vaginal secretions, ascites fluids, fluids of the pleural, pericardial, peritoneal, abdominal and other body cavities, fluids collected by bronchial lavage, synovial fluid, liquid solutions contacted with a subject or biological source, for example, cell and organ culture medium including cell or organ conditioned medium and lavage fluids and the like.

[0048] “Substituting” or “substituted” or “mutating” or “mutated” refers to altering, deleting or inserting one or more amino acids or nucleotides in a polypeptide or polynucleotide sequence to generate a variant of that sequence.

[0049] “Variant” refers to a polypeptide or a polynucleotide that differs from a reference polypeptide or a reference polynucleotide by one or more modifications for example, substitutions, insertions or deletions.

[0050] “Specifically binds” or “specific binding” refers to the ability of a FN3 domain to bind to its target, such as CD71, with a dissociation constant (K_D) of about 1×10^{-6} M or less, for example about 1×10^{-7} M or less, about 1×10^{-8} M or less, about 1×10^{-9} M or less, about 1×10^{-10} M or less, about 1×10^{-11} M or less, about 1×10^{-12} M or less, or about 1×10^{-13} M or less. Alternatively, “specific binding” refers to the ability of a FN3 domain to bind to its target (*e.g.* CD71) at least 5-fold above a negative control in standard solution ELISA assay. Specific binding can also be demonstrated using the proteome array as described herein and shown in FIG. 3. In some embodiments, a negative control is an FN3 domain that does not bind CD71. In some embodiment, an FN3 domain that specifically binds CD71 may have cross-reactivity to other related antigens, for example to the same predetermined antigen from other species (homologs), such as *Macaca Fascicularis* (cynomolgous monkey, cyno) or *Pan troglodytes* (chimpanzee).

[0051] “Library” refers to a collection of variants. The library may be composed of polypeptide or polynucleotide variants.

[0052] “Stability” refers to the ability of a molecule to maintain a folded state under physiological conditions such that it retains at least one of its normal functional activities, for example, binding to a predetermined antigen such as CD71.

[0053] “CD71” refers to human CD71 protein having the amino acid sequence of SEQ ID NOs: 2 or 5. In some embodiments, SEQ ID NO: 2 is full length human CD71 protein. In some embodiments, SEQ ID NO: 5 is the extracellular domain of human CD71.

[0054] “Tencon” refers to the synthetic fibronectin type III (FN3) domain having the consensus sequence shown in SEQ ID NO:1

LPAPKNLVVSEVTEDSLRLSWTAPDAAFDSEFLIQYQSEKVGGEAINLTPGSEERSYD
LTGLKPGTEYTVSIYGVKGGHRSNPLSAEFTT

and described in U.S. Pat. Publ. No. 2010/0216708.

[0055] A “cancer cell” or a “tumor cell” refers to a cancerous, pre-cancerous or transformed cell, either *in vivo*, *ex vivo*, and in tissue culture, that has spontaneous or induced phenotypic changes that do not necessarily involve the uptake of new genetic material. Although transformation can arise from infection with a transforming virus and incorporation of new genomic nucleic acid, or uptake of exogenous nucleic acid, it can also arise spontaneously or following exposure to a carcinogen, thereby mutating an endogenous gene. Transformation/cancer is exemplified by, *e.g.*, morphological changes, immortalization of cells, aberrant growth control, foci formation,

proliferation, malignancy, tumor specific markers levels, invasiveness, tumor growth or suppression in suitable animal hosts such as nude mice, and the like, *in vitro*, *in vivo*, and *ex vivo* (Freshney, Culture of Animal Cells: A Manual of Basic Technique (3rd ed. 1994)).

[0056] A “dendritic cell” refers to a type of antigen-presenting cell (APC) that form an important role in the adaptive immune system. The main function of dendritic cells is to present antigens to T lymphocytes, and to secrete cytokines that may further modulate the immune response directly or indirectly. Dendritic cells have the capacity to induce a primary immune response in the inactive or resting naïve T lymphocytes.

[0057] An “immune cell” refers to the cells of the immune system categorized as lymphocytes (T-cells, B-cells and NK cells), neutrophils, or monocytes/macrophages. These are all types of white blood cells.

[0058] “Vector” refers to a polynucleotide capable of being duplicated within a biological system or that can be moved between such systems. Vector polynucleotides typically contain elements, such as origins of replication, polyadenylation signal or selection markers that function to facilitate the duplication or maintenance of these polynucleotides in a biological system. Examples of such biological systems may include a cell, virus, animal, plant, and reconstituted biological systems utilizing biological components capable of duplicating a vector. The polynucleotide comprising a vector may be DNA or RNA molecules or a hybrid of these.

[0059] “Expression vector” refers to a vector that can be utilized in a biological system or in a reconstituted biological system to direct the translation of a polypeptide encoded by a polynucleotide sequence present in the expression vector.

[0060] “Polynucleotide” refers to a synthetic molecule comprising a chain of nucleotides covalently linked by a sugar-phosphate backbone or other equivalent covalent chemistry. cDNA is a typical example of a polynucleotide.

[0061] “Polypeptide” or “protein” refers to a molecule that comprises at least two amino acid residues linked by a peptide bond to form a polypeptide. Small polypeptides of less than about 50 amino acids may be referred to as “peptides”.

[0062] “Valent” refers to the presence of a specified number of binding sites specific for an antigen in a molecule. As such, the terms “monovalent”, “bivalent”, “tetravalent”, and “hexavalent” refer to the presence of one, two, four and six binding sites, respectively, specific for an antigen in a molecule.

[0063] “Subject” includes any human or nonhuman animal. “Nonhuman animal” includes all vertebrates, *e.g.*, mammals and non-mammals, such as nonhuman primates, sheep, dogs, cats,

horses, cows chickens, amphibians, reptiles, etc. Except when noted, the terms “patient” or “subject” are used interchangeably.

[0064] “Isolated” refers to a homogenous population of molecules (such as synthetic polynucleotides or a polypeptide such as FN3 domains) which have been substantially separated and/or purified away from other components of the system the molecules are produced in, such as a recombinant cell, as well as a protein that has been subjected to at least one purification or isolation step. “Isolated FN3 domain” refers to an FN3 domain that is substantially free of other cellular material and/or chemicals and encompasses FN3 domains that are isolated to a higher purity, such as to 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% purity.

[0065] In some embodiments, a composition comprising a polypeptide, such as a polypeptide comprising a FN3 domain, linked to an oligonucleotide molecule are provided. The oligonucleotide molecule can be, for example, a siRNA molecule.

[0066] Accordingly, in some embodiments, the siRNA is a double-stranded RNAi (dsRNA) agent capable of inhibiting the expression of a target gene. The dsRNA agent comprises a sense strand (passenger strand) and an antisense strand (guide strand). In some embodiments, each strand of the dsRNA agent can range from 12-40 nucleotides in length. For example, each strand can be from 14-40 nucleotides in length, 17-37 nucleotides in length, 25-37 nucleotides in length, 27-30 nucleotides in length, 17-23 nucleotides in length, 17-21 nucleotides in length, 17-19 nucleotides in length, 19-25 nucleotides in length, 19-23 nucleotides in length, 19-21 nucleotides in length, 21-25 nucleotides in length, or 21-23 nucleotides in length.

[0067] In some embodiments, the sense strand and antisense strand typically form a duplex dsRNA. The duplex region of a dsRNA agent may be from 12-40 nucleotide pairs in length. For example, the duplex region can be from 14-40 nucleotide pairs in length, 17-30 nucleotide pairs in length, 25-35 nucleotides in length, 27-35 nucleotide pairs in length, 17-23 nucleotide pairs in length, 17-21 nucleotide pairs in length, 17-19 nucleotide pairs in length, 19-25 nucleotide pairs in length, 19-23 nucleotide pairs in length, 19-21 nucleotide pairs in length, 21-25 nucleotide pairs in length, or 21-23 nucleotide pairs in length. In another example, the duplex region is selected from 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, and 27 nucleotide pairs in length.

[0068] In some embodiments, the dsRNA comprises one or more overhang regions and/or capping groups of dsRNA agent at the 3'-end, or 5'-end or both ends of a strand. The overhang can be 1-10 nucleotides in length, 1-6 nucleotides in length, for instance 2-6 nucleotides in length, 1-5 nucleotides in length, 2-5 nucleotides in length, 1-4 nucleotides in length, 2-4 nucleotides in length, 1-3 nucleotides in length, 2-3 nucleotides in length, or 1-2 nucleotides in

length. The overhangs can be the result of one strand being longer than the other, or the result of two strands of the same length being staggered. The overhang can form a mismatch with the target mRNA or it can be complementary to the gene sequences being targeted or can be other sequence. The first and second strands can also be joined, e.g., by additional bases to form a hairpin, or by other non-base linkers.

[0069] In some embodiments, the nucleotides in the overhang region of the dsRNA agent can each independently be a modified or unmodified nucleotide including, but not limited to 2'-sugar modified, such as, 2-F, 2'-O-methyl, 2'-O-(2-methoxyethyl), 2'-O-(2-methoxyethyl), 2'-O-(2-methoxyethyl), and any combinations thereof. For example, TT (UU) can be an overhang sequence for either end on either strand. The overhang can form a mismatch with the target mRNA or it can be complementary to the gene sequences being targeted or can be other sequence.

[0070] The 5'- or 3'-overhangs at the sense strand, antisense strand or both strands of the dsRNA agent may be phosphorylated. In some embodiments, the overhang region contains two nucleotides having a phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate between the two nucleotides, where the two nucleotides can be the same or different. In one embodiment, the overhang is present at the 3'-end of the sense strand, antisense strand or both strands. In one embodiment, this 3'-overhang is present in the antisense strand. In one embodiment, this 3'-overhang is present in the sense strand.

[0071] The dsRNA agent may comprise only a single overhang, which can strengthen the interference activity of the dsRNA, without affecting its overall stability. For example, the single-stranded overhang is located at the 3'-terminal end of the sense strand or, alternatively, at the 3'-terminal end of the antisense strand. The dsRNA may also have a blunt end, located at the 5'-end of the antisense strand (or the 3'-end of the sense strand) or vice versa. Generally, the antisense strand of the dsRNA has a nucleotide overhang at the 3'-end, and the 5'-end is blunt. While not bound by theory, the asymmetric blunt end at the 5'-end of the antisense strand and 3'-end overhang of the antisense strand favor the guide strand loading into RISC. For example the single overhang comprises at least two, three, four, five, six, seven, eight, nine, or ten nucleotides in length.

[0072] In some embodiments, the dsRNA agent may also have two blunt ends, at both ends of the dsRNA duplex.

[0073] In some embodiments, every nucleotide in the sense strand and antisense strand of the dsRNA agent may be modified. Each nucleotide may be modified with the same or different modification which can include one or more alteration of one or both of the non-linking

phosphate oxygens and/or of one or more of the linking phosphate oxygens; alteration of a constituent of the ribose sugar, e.g., of the 2 hydroxyl on the ribose sugar; wholesale replacement of the phosphate moiety with “dephospho” linkers; modification or replacement of a naturally occurring base; and replacement or modification of the ribose-phosphate backbone.

[0074] In some embodiments all or some of the bases in a 3' or 5' overhang may be modified, e.g., with a modification described herein. Modifications can include, e.g., the use of modifications at the 2' position of the ribose sugar with modifications that are known in the art, e.g., the use of deoxyribonucleotides, 2'-deoxy-2'-fluoro (2'-F) or 2'-O-methyl modified instead of the ribosugar of the nucleobase, and modifications in the phosphate group, e.g., phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate modifications. Overhangs need not be homologous with the target sequence.

[0075] In some embodiments, each residue of the sense strand and antisense strand is independently modified with LNA, HNA, CeNA, 2'-methoxyethyl, 2'-O-methyl, 2'-O-allyl, 2'-C-allyl, 2'-deoxy, or 2'-fluoro. The strands can contain more than one modification. In one embodiment, each residue of the sense strand and antisense strand is independently modified with 2'-O-methyl or 2'-fluoro.

[0076] In some embodiments, at least two different modifications are typically present on the sense strand and antisense strand. Those two modifications may be the 2'-deoxy, 2'-O-methyl or 2'-fluoro modifications, acyclic nucleotides or others.


[0077] In one embodiment, the sense strand and antisense strand each comprises two differently modified nucleotides selected from 2'-fluoro, 2'-O-methyl or 2'-deoxy.

[0078] The dsRNA agent may further comprise at least one phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, mesyl phosphoramidate, or methylphosphonate internucleotide linkage. The phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, mesyl phosphoramidate, or methylphosphonate internucleotide linkage modification may occur on any nucleotide of the sense strand or antisense strand or both in any position of the strand. For instance, the internucleotide linkage modification may occur on every nucleotide on the sense strand and/or antisense strand; each internucleotide linkage modification may occur in an alternating pattern on the sense strand or antisense strand; or the sense strand or antisense strand comprises both internucleotide linkage modifications in an alternating pattern. The alternating pattern of the internucleotide linkage modification on the sense strand may be the same or different from the antisense strand, and the alternating pattern of the internucleotide linkage modification on the sense strand may have a shift relative to the alternating pattern of the internucleotide linkage modification on the antisense strand.

[0079] In some embodiments, the dsRNA agent comprises the phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, mesyl phosphoramidate, or methylphosphonate internucleotide linkage modification in the overhang region. For example, the overhang region comprises two nucleotides having a phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, mesyl phosphoramidate, or methylphosphonate internucleotide linkage between the two nucleotides. Internucleotide linkage modifications also may be made to link the overhang nucleotides with the terminal paired nucleotides within duplex region. For example, at least 2, 3, 4, or all the overhang nucleotides may be linked through phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, mesyl phosphoramidate, or methylphosphonate internucleotide linkage, and optionally, there may be additional phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, mesyl phosphoramidate, or methylphosphonate internucleotide linkages linking the overhang nucleotide with a paired nucleotide that is next to the overhang nucleotide. For instance, there may be at least two phosphorothioate internucleotide linkages between the terminal three nucleotides, in which two of the three nucleotides are overhang nucleotides, and the third is a paired nucleotide next to the overhang nucleotide. In some embodiments, these terminal three nucleotides may be at the 3'-end of the antisense strand.

[0080] In some embodiments, the dsRNA composition is linked by a modified base or nucleoside analogue as described in U.S. Patent No. 7,427,672, which is incorporated herein by reference. In some embodiments, the modified base or nucleoside analogue is referred to as the linker or L in formulas described herein.

[0081] In some embodiments, the modified base or nucleoside analogue has the structure as shown in Chemical Formula I and a salt thereof:



where Base represents an aromatic heterocyclic group or aromatic hydrocarbon ring group optionally having a substituent, R₁ and R₂ are identical or different, and each represent a

hydrogen atom, a protective group for a hydroxyl group for nucleic acid synthesis, an alkyl group, an alkenyl group, a cycloalkyl group, an aryl group, an aralkyl group, an acyl group, a sulfonyl group, a silyl group, a phosphate group, a phosphate group protected with a protective group for nucleic acid synthesis, or $--P(R_4)R_5$ where R_4 and R_5 are identical or different, and each represent a hydroxyl group, a hydroxyl group protected with a protective group for nucleic acid synthesis, a mercapto group, a mercapto group protected with a protective group for nucleic acid synthesis, an amino group, an alkoxy group having 1 to 5 carbon atoms, an alkylthio group having 1 to 5 carbon atoms, a cyanoalkoxy group having 1 to 6 carbon atoms, or an amino group substituted by an alkyl group having 1 to 5 carbon atoms, R_3 represents a hydrogen atom, an alkyl group, an alkenyl group, a cycloalkyl group, an aryl group, an aralkyl group, an acyl group, a sulfonyl group, or a functional molecule unit substituent, and m denotes an integer of 0 to 2, and n denotes an integer of 0 to 3. In some embodiments, m and n are 0.

[0082] In some embodiments, the modified base or nucleoside analogue has the structure as shown in Chemical Formula I and salts thereof, wherein R_1 is a hydrogen atom, an aliphatic acyl group, an aromatic acyl group, an aliphatic or aromatic sulfonyl group, a methyl group substituted by one to three aryl groups, a methyl group substituted by one to three aryl groups having an aryl ring substituted by a lower alkyl, lower alkoxy, halogen, or cyano group, or a silyl group.

[0083] In some embodiments, the modified base or nucleoside analogue has the structure as shown in Chemical Formula I and salts thereof, wherein R_1 is a hydrogen atom, an acetyl group, a benzoyl group, a methanesulfonyl group, a p-toluenesulfonyl group, a benzyl group, a p-methoxybenzyl group, a trityl group, a dimethoxytrityl group, a monomethoxytrityl group, or a tert-butyldiphenylsilyl group.

[0084] In some embodiments, the modified base or nucleoside analogue has the structure as shown in Chemical Formula I and salts thereof, wherein R_2 is a hydrogen atom, an aliphatic acyl group, an aromatic acyl group, an aliphatic or aromatic sulfonyl group, a methyl group substituted by one to three aryl groups, a methyl group substituted by one to three aryl groups having an aryl ring substituted by a lower alkyl, lower alkoxy, halogen, or cyano group, a silyl group, a phosphoroamidite group, a phosphonyl group, a phosphate group, or a phosphate group protected with a protective group for nucleic acid synthesis.

[0085] In some embodiments, the modified base or nucleoside analogue has the structure as shown in Chemical Formula I and salts thereof, wherein R_2 is a hydrogen atom, an acetyl group, a benzoyl group, a methanesulfonyl group, a p-toluenesulfonyl group, a benzyl group, a p-

methoxybenzyl group, a tert-butyldiphenylsilyl group, $--P(OC_2H_4CN)(N(i-Pr)_2)$, $--P(OCH_3)(N(i-Pr)_2)$, a phosphonyl group, or a 2-chlorophenyl- or 4-chlorophenylphosphate group.

[0086] In some embodiments, the modified base or nucleoside analogue has the structure as shown in Chemical Formula I and salts thereof, wherein R_3 is a hydrogen atom, a phenoxyacetyl group, an alkyl group having 1 to 5 carbon atoms, an alkenyl group having 1 to 5 carbon atoms, an aryl group having 6 to 14 carbon atoms, a methyl group substituted by one to three aryl groups, a lower aliphatic or aromatic sulfonyl group such as a methanesulfonyl group or a p-toluenesulfonyl group, an aliphatic acyl group having 1 to 5 carbon atoms such as an acetyl group, or an aromatic acyl group such as a benzoyl group.

[0087] In some embodiments, the modified base or nucleoside analogue has the structure as shown in Chemical Formula I and salts thereof, wherein the functional molecule unit substituent as R_3 is a fluorescent or chemiluminescent labeling molecule, a nucleic acid incision activity functional group, or an intracellular or nuclear transfer signal peptide.


[0088] In some embodiments, the modified base or nucleoside analogue has the structure as shown in Chemical Formula I and salts thereof, wherein Base is a purin-9-yl group, a 2-oxypyrimidin-1-yl group, or a purin-9-yl group or a 2-oxypyrimidin-1-yl group having a substituent selected from the following α group: α group: A hydroxyl group, a hydroxyl group protected with a protective group for nucleic acid synthesis, an alkoxy group having 1 to 5 carbon atoms, a mercapto group, a mercapto group protected with a protective group for nucleic acid synthesis, an alkylthio group having 1 to 5 carbon atoms, an amino group, an amino group protected with a protective group for nucleic acid synthesis, an amino group substituted by an alkyl group having 1 to 5 carbon atoms, an alkyl group having 1 to 5 carbon atoms, and a halogen atom.

[0089] In some embodiments, the modified base or nucleoside analogue has the structure as shown in Chemical Formula I and salts thereof, wherein Base is 6-aminopurin-9-yl (i.e., adeninyl), 6-aminopurin-9-yl having the amino group protected with a protective group for nucleic acid synthesis, 2,6-diaminopurin-9-yl, 2-amino-6-chloropurin-9-yl, 2-amino-6-chloropurin-9-yl having the amino group protected with a protective group for nucleic acid synthesis, 2-amino-6-fluoropurin-9-yl, 2-amino-6-fluoropurin-9-yl having the amino group protected with a protective group for nucleic acid synthesis, 2-amino-6-bromopurin-9-yl, 2-amino-6-bromopurin-9-yl having the amino group protected with a protective group for nucleic acid synthesis, 2-amino-6-hydroxypurin-9-yl (i.e., guaninyl), 2-amino-6-hydroxypurin-9-yl having the amino group protected with a protective group for nucleic acid synthesis, 6-amino-2-methoxypurin-9-yl, 6-amino-2-chloropurin-9-yl, 6-amino-2-fluoropurin-9-yl, 2,6-

dimethoxypurin-9-yl, 2,6-dichloropurin-9-yl, 6-mercaptopurin-9-yl, 2-oxo-4-amino-1,2-dihydropyrimidin-1-yl (i.e., cytosinyl), 2-oxo-4-amino-1,2-dihydropyrimidin-1-yl having the amino group protected with a protective group for nucleic acid synthesis, 2-oxo-4-amino-5-fluoro-1,2-dihydropyrimidin-1-yl, 2-oxo-4-amino-5-fluoro-1,2-dihydropyrimidin-1-yl having the amino group protected with a protective group for nucleic acid synthesis, 4-amino-2-oxo-5-chloro-1,2-dihydropyrimidin-1-yl, 2-oxo-4-methoxy-1,2-dihydropyrimidin-1-yl, 2-oxo-4-mercapto-1,2-dihydropyrimidin-1-yl, 2-oxo-4-hydroxy-1,2-dihydropyrimidin-1-yl (i.e., uraciny), 2-oxo-4-hydroxy-5-methyl-1,2-dihydropyrimidin-1-yl (i.e., thyminy), 4-amino-5-methyl-2-oxo-1,2-dihydropyrimidin-1-yl (i.e., 5-methylcytosinyl), or 4-amino-5-methyl-2-oxo-1,2-dihydropyrimidin-1-yl having the amino group protected with a protective group for nucleic acid synthesis.

[0090] In some embodiments, the modified base or nucleoside analogue has the structure as shown in Chemical Formula I and salts thereof, wherein m is 0, and n is 1.

[0091] In some embodiments, the modified base or nucleoside analogue is a DNA oligonucleotide or RNA oligonucleotide analogue, containing one or two or more of one or more types of unit structures of nucleoside analogues having the structure as shown in Chemical Formula II, or a pharmacologically acceptable salt thereof, provided that a form of linking between respective nucleosides in the oligonucleotide analogue may contain one or two or more phosphorothioate bonds [--OP(O)(S⁻)O--], phosphorodithioate bonds [--O₂PS₂--], phosphonate bonds [--PO(OH)₂--], phosphoramidate bonds [--O=P(OH)₂--], or mesyl phosphoramidate bonds [--OP(O)(N)(SO₂)(CH₃)O--] aside from a phosphodiester bond [--OP(O₂⁻)O--] identical with that in a natural nucleic acid, and if two or more of one or more types of these structures are contained, Base may be identical or different between these structures:



(Chemical Formula II)

where Base represents an aromatic heterocyclic group or aromatic hydrocarbon ring group optionally having a substituent, R_3 represents a hydrogen atom, an alkyl group, an alkenyl group, a cycloalkyl group, an aryl group, an aralkyl group, an acyl group, a sulfonyl group, a silyl group, or a functional molecule unit substituent, and m denotes an integer of 0 to 2, and n denotes an integer of 0 to 3. In some embodiments, m and n are 0.

[0092] In some embodiments, the oligonucleotide analogue or the pharmacologically acceptable salt thereof has the structure as shown in Chemical Formula II, wherein R_1 is a hydrogen atom, an aliphatic acyl group, an aromatic acyl group, an aliphatic or aromatic sulfonyl group, a methyl group substituted by one to three aryl groups, a methyl group substituted by one to three aryl groups having an aryl ring substituted by a lower alkyl, lower alkoxy, halogen, or cyano group, or a silyl group.

[0093] In some embodiments, the oligonucleotide analogue or the pharmacologically acceptable salt thereof has the structure as shown in Chemical Formula II, wherein R_1 is a hydrogen atom, an acetyl group, a benzoyl group, a methanesulfonyl group, a p-toluenesulfonyl group, a benzyl group, a p-methoxybenzyl group, a trityl group, a dimethoxytrityl group, a monomethoxytrityl group, or a tert-butyldiphenylsilyl group.

[0094] In some embodiments, the oligonucleotide analogue or the pharmacologically acceptable salt thereof has the structure as shown in Chemical Formula II, wherein R_2 is a hydrogen atom, an aliphatic acyl group, an aromatic acyl group, an aliphatic or aromatic sulfonyl group, a methyl group substituted by one to three aryl groups, a methyl group substituted by one to three aryl groups having an aryl ring substituted by a lower alkyl, lower alkoxy, halogen, or cyano group, a silyl group, a phosphoroamidite group, a phosphonyl group, a phosphate group, or a phosphate group protected with a protective group for nucleic acid synthesis.

[0095] In some embodiments, the oligonucleotide analogue or the pharmacologically acceptable salt thereof has the structure as shown in Chemical Formula II, wherein R_2 is a hydrogen atom, an acetyl group, a benzoyl group, a benzyl group, a p-methoxybenzyl group, a methanesulfonyl group, a p-toluenesulfonyl group, a tert-butyldiphenylsilyl group, $--P(OC_2H_4CN)(N(i-Pr)_2)$, $--P(OCH_3)(N(i-Pr)_2)$, a phosphonyl group, or a 2-chlorophenyl- or 4-chlorophenylphosphate group.

[0096] In some embodiments, the oligonucleotide analogue or the pharmacologically acceptable salt thereof has the structure as shown in Chemical Formula II, wherein R_3 is a hydrogen atom, a phenoxyacetyl group, an alkyl group having 1 to 5 carbon atoms, an alkenyl group having 1 to 5 carbon atoms, an aryl group having 6 to 14 carbon atoms, a methyl group substituted by one to three aryl groups, a lower aliphatic or aromatic sulfonyl group such as a methanesulfonyl group

or a p-toluenesulfonyl group, an aliphatic acyl group having 1 to 5 carbon atoms such as an acetyl group, or an aromatic acyl group such as a benzoyl group.

[0097] In some embodiments, the oligonucleotide analogue or the pharmacologically acceptable salt thereof has the structure as shown in Chemical Formula II, wherein the functional molecule unit substituent as R₃ is a fluorescent or chemiluminescent labeling molecule, a nucleic acid incision activity functional group, or an intracellular or nuclear transfer signal peptide.

[0098] In some embodiments, the oligonucleotide analogue or the pharmacologically acceptable salt thereof has the structure as shown in Chemical Formula II, wherein Base is a purin-9-yl group, a 2-oxypyrimidin-1-yl group, or a purin-9-yl group or a 2-oxypyrimidin-1-yl group having a substituent selected from the following α group: α group: A hydroxyl group, a hydroxyl group protected with a protective group for nucleic acid synthesis, an alkoxy group having 1 to 5 carbon atoms, a mercapto group, a mercapto group protected with a protective group for nucleic acid synthesis, an alkylthio group having 1 to 5 carbon atoms, an amino group, an amino group protected with a protective group for nucleic acid synthesis, an amino group substituted by an alkyl group having 1 to 5 carbon atoms, an alkyl group having 1 to 5 carbon atoms, and a halogen atom.

[0099] In some embodiments, the oligonucleotide analogue or the pharmacologically acceptable salt thereof has the structure as shown in Chemical Formula II, wherein Base is 6-aminopurin-9-yl (i.e. adeninyl), 6-aminopurin-9-yl having the amino group protected with a protective group for nucleic acid synthesis, 2,6-diaminopurin-9-yl, 2-amino-6-chloropurin-9-yl, 2-amino-6-chloropurin-9-yl having the amino group protected with a protective group for nucleic acid synthesis, 2-amino-6-fluoropurin-9-yl, 2-amino-6-fluoropurin-9-yl having the amino group protected with a protective group for nucleic acid synthesis, 2-amino-6-bromopurin-9-yl, 2-amino-6-bromopurin-9-yl having the amino group protected with a protective group for nucleic acid synthesis, 2-amino-6-hydroxypurin-9-yl (i.e., guaninyl), 2-amino-6-hydroxypurin-9-yl having the amino group protected with a protective group for nucleic acid synthesis, 6-amino-2-methoxypurin-9-yl, 6-amino-2-chloropurin-9-yl, 6-amino-2-fluoropurin-9-yl, 2,6-dimethoxypurin-9-yl, 2,6-dichloropurin-9-yl, 6-mercaptopurin-9-yl, 2-oxo-4-amino-1,2-dihydropyrimidin-1-yl (i.e., cytosinyl), 2-oxo-4-amino-1,2-dihydropyrimidin-1-yl having the amino group protected with a protective group for nucleic acid synthesis, 2-oxo-4-amino-5-fluoro-1,2-dihydropyrimidin-1-yl, 2-oxo-4-amino-5-fluoro-1,2-dihydropyrimidin-1-yl group having the amino group protected with a protective group for nucleic acid synthesis, 4-amino-2-oxo-5-chloro-1,2-dihydropyrimidin-1-yl, 2-oxo-4-methoxy-1,2-dihydropyrimidin-1-yl, 2-oxo-4-mercapto-1,2-dihydropyrimidin-1-yl, 2-oxo-4-hydroxy-1,2-dihydropyrimidin-1-yl (i.e.,

uraciny], 2-oxo-4-hydroxy-5-methyl-1,2-dihydropyrimidin-1-yl (i.e., thyminy], 4-amino-5-methyl-2-oxo-1,2-dihydropyrimidin-1-yl (i.e., 5-methylcytosiny], or 4-amino-5-methyl-2-oxo-1,2-dihydropyrimidin-1-yl having the amino group protected with a protective group for nucleic acid synthesis.

[00100] In some embodiments, the oligonucleotide analogue or the pharmacologically acceptable salt thereof has the structure as shown in Chemical Formula II, wherein m is 0, and n is 1.

[00101] In some embodiments, compositions described herein further comprises a polymer (polymer moiety C). In some instances, the polymer is a natural or synthetic polymer, consisting of long chains of branched or unbranched monomers, and/or cross-linked network of monomers in two or three dimensions. In some instances, the polymer includes a polysaccharide, lignin, rubber, or polyalkylene oxide (e.g., polyethylene glycol). In some instances, the at least one polymer includes, but is not limited to, alpha-, omega-dihydroxypolyethyleneglycol, biodegradable lactone-based polymer, e.g. polyacrylic acid, polylactide acid (PLA), poly(glycolic acid) (PGA), polypropylene, polystyrene, polyolefin, polyamide, polycyanoacrylate, polyimide, polyethyleneterephthalat (PET, PETG), polyethylene terephthalate (PETE), polytetramethylene glycol (PTG), or polyurethane as well as mixtures thereof. As used herein, a mixture refers to the use of different polymers within the same compound as well as in reference to block copolymers. In some cases, block copolymers are polymers wherein at least one section of a polymer is build up from monomers of another polymer. In some instances, the polymer comprises polyalkylene oxide. In some instances, the polymer comprises PEG. In some instances, the polymer comprises polyethylene imide (PEI) or hydroxy ethyl starch (HES).

[00102] In some instances, C is a PEG moiety. In some instances, the PEG moiety is conjugated at the 5' terminus of the oligonucleotide molecule while the binding moiety is conjugated at the 3' terminus of the oligonucleotide molecule. In some instances, the PEG moiety is conjugated at the 3' terminus of the oligonucleotide molecule while the binding moiety is conjugated at the 5' terminus of the oligonucleotide molecule. In some instances, the PEG moiety is conjugated to an internal site of the oligonucleotide molecule. In some instances, the PEG moiety, the binding moiety, or a combination thereof, are conjugated to an internal site of the oligonucleotide molecule. In some instances, the conjugation is a direct conjugation. In some instances, the conjugation is via native ligation.

[00103] In some embodiments, the polyalkylene oxide (e.g., PEG) is a polydispers or monodispers compound. In some instances, polydispers material comprises disperse distribution of different molecular weight of the material, characterized by mean weight (weight average)

size and dispersity. In some instances, the monodisperse PEG comprises one size of molecules. In some embodiments, C is poly- or monodispersed polyalkylene oxide (e.g., PEG) and the indicated molecular weight represents an average of the molecular weight of the polyalkylene oxide, e.g., PEG, molecules.

[00104] In some embodiments, the molecular weight of the polyalkylene oxide (e.g., PEG) is about 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1450, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, 2500, 2600, 2700, 2800, 2900, 3000, 3250, 3350, 3500, 3750, 4000, 4250, 4500, 4600, 4750, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 10,000, 12,000, 20,000, 35,000, 40,000, 50,000, 60,000, or 100,000 Da.

[00105] In some embodiments, C is polyalkylene oxide (e.g., PEG) and has a molecular weight of about 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1450, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, 2500, 2600, 2700, 2800, 2900, 3000, 3250, 3350, 3500, 3750, 4000, 4250, 4500, 4600, 4750, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 10,000, 12,000, 20,000, 35,000, 40,000, 50,000, 60,000, or 100,000 Da. In some embodiments, C is PEG and has a molecular weight of about 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1450, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, 2500, 2600, 2700, 2800, 2900, 3000, 3250, 3350, 3500, 3750, 4000, 4250, 4500, 4600, 4750, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 10,000, 12,000, 20,000, 35,000, 40,000, 50,000, 60,000, or 100,000 Da. In some instances, the molecular weight of C is about 200 Da. In some instances, the molecular weight of C is about 300 Da. In some instances, the molecular weight of C is about 400 Da. In some instances, the molecular weight of C is about 500 Da. In some instances, the molecular weight of C is about 600 Da. In some instances, the molecular weight of C is about 700 Da. In some instances, the molecular weight of C is about 800 Da. In some instances, the molecular weight of C is about 900 Da. In some instances, the molecular weight of C is about 1000 Da. In some instances, the molecular weight of C is about 1100 Da. In some instances, the molecular weight of C is about 1200 Da. In some instances, the molecular weight of C is about 1300 Da. In some instances, the molecular weight of C is about 1400 Da. In some instances, the molecular weight of C is about 1450 Da. In some instances, the molecular weight of C is about 1500 Da. In some instances, the molecular weight of C is about 1600 Da. In some instances, the molecular weight of C is about 1700 Da. In some instances, the molecular weight of C is about 1800 Da. In some instances, the molecular weight of C is about 1900 Da. In some instances, the molecular weight of C is about 2000 Da. In some instances, the molecular weight of C is about 2100 Da. In some instances, the molecular weight of C is about 2200 Da. In some instances, the molecular weight of C is about 2300 Da. In some instances, the molecular

weight of C is about 2400 Da. In some instances, the molecular weight of C is about 2500 Da. In some instances, the molecular weight of C is about 2600 Da. In some instances, the molecular weight of C is about 2700 Da. In some instances, the molecular weight of C is about 2800 Da. In some instances, the molecular weight of C is about 2900 Da. In some instances, the molecular weight of C is about 3000 Da. In some instances, the molecular weight of C is about 3250 Da. In some instances, the molecular weight of C is about 3350 Da. In some instances, the molecular weight of C is about 3500 Da. In some instances, the molecular weight of C is about 3750 Da. In some instances, the molecular weight of C is about 4000 Da. In some instances, the molecular weight of C is about 4250 Da. In some instances, the molecular weight of C is about 4500 Da. In some instances, the molecular weight of C is about 4600 Da. In some instances, the molecular weight of C is about 4750 Da. In some instances, the molecular weight of C is about 5000 Da. In some instances, the molecular weight of C is about 5500 Da. In some instances, the molecular weight of C is about 6000 Da. In some instances, the molecular weight of C is about 6500 Da. In some instances, the molecular weight of C is about 7000 Da. In some instances, the molecular weight of C is about 7500 Da. In some instances, the molecular weight of C is about 8000 Da. In some instances, the molecular weight of C is about 10,000 Da. In some instances, the molecular weight of C is about 12,000 Da. In some instances, the molecular weight of C is about 20,000 Da. In some instances, the molecular weight of C is about 35,000 Da. In some instances, the molecular weight of C is about 40,000 Da. In some instances, the molecular weight of C is about 50,000 Da. In some instances, the molecular weight of C is about 60,000 Da. In some instances, the molecular weight of C is about 100,000 Da.

[00106] In some embodiments, the polyalkylene oxide (e.g., PEG) is a discrete PEG, in which the discrete PEG is a polymeric PEG comprising more than one repeating ethylene oxide units. In some instances, a discrete PEG (dPEG) comprises from 2 to 60, from 2 to 50, or from 2 to 48 repeating ethylene oxide units. In some instances, a dPEG comprises about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 24, 26, 28, 30, 35, 40, 42, 48, 50 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 2 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 3 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 4 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 5 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 6 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 7 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 8 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 9 or more repeating ethylene oxide units. In some instances, a dPEG comprises

about 10 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 11 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 12 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 13 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 14 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 15 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 16 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 17 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 18 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 19 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 20 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 22 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 24 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 26 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 28 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 30 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 35 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 40 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 42 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 48 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 50 or more repeating ethylene oxide units. In some cases, a dPEG is synthesized as a single molecular weight compound from pure (e.g., about 95%, 98%, 99%, or 99.5%) starting material in a step-wise fashion. In some cases, a dPEG has a specific molecular weight, rather than an average molecular weight. In some cases, a dPEG described herein is a dPEG from Quanta Biodesign, LMD.

[00107] In some embodiments, C is an albumin binding domain. In certain aspects, the albumin binding domain specifically binds to serum albumin, e.g., human serum albumin (HSA) to prolong the half-life of the domain or of another therapeutic to which the albumin-binding domain is associated or linked with. In some embodiments, the human serum albumin-binding domain comprises an initiator methionine (Met) linked to the N-terminus of the molecule. In some embodiments, the human serum albumin-binding domain comprise a cysteine (Cys) linked to a C-terminus or the N-terminus of the domain. The addition of the N-terminal Met and/or the C-terminal Cys may facilitate expression and/or conjugation to another molecule, which can be another half-life extending molecules, such as PEG, a Fc region, and the like.

[00108] In some embodiments, the albumin binding domain comprises the amino acid sequence of SEQ ID NOs: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114,

115, 116, 117, 118, or 119, provided in Table 8. In some embodiments, the albumin binding domain (protein) is isolated. In some embodiments, the albumin binding domain comprises an amino acid sequence that is at least, or is, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence of SEQ ID NO: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119. In some embodiments, the albumin binding domain comprises an amino acid sequence that is at least, or is, 85%, 86%, 87%, 88%, 89%, 90%, 901%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence of SEQ ID NO: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119 provided that the protein has a substitution that corresponds to position 10 of SEQ ID NO: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119. In some embodiments, the substitution is A10V. In some embodiments, the substitution is A10G, A10L, A10I, A10T, or A10S. In some embodiments, the substitution at position 10 is any naturally occurring amino acid. In some embodiments, the isolated albumin binding domain comprises an amino acid sequence that has 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14 substitutions when compared to the amino acid sequence of SEQ ID NOs: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119. In some embodiments, the substitution is at a position that corresponds to position 10 of SEQ ID NOs: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119. In some embodiments, FN3 domains provided comprises a cysteine residue in at least one residue position corresponding to residue positions 6, 11, 22, 25, 26, 52, 53, 61, 88 or positions 6, 8, 10, 11, 14, 15, 16, 20, 30, 34, 38, 40, 41, 45, 47, 48, 53, 54, 59, 60, 62, 64, 70, 88, 89, 90, 91, or 93 of SEQ ID NO: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119, or at a C-terminus. Although the positions are listed in a series, each position can also be chosen individually. In some embodiments, the cysteine is at a position that corresponds to position 6, 53, or 88. In some embodiments, additional examples of albumin binding domains can be found in U.S. Patent No. 10,925,932, which hereby incorporated by reference.

Table 8

SEQ ID NO:	SEQUENCE
101	MLPAPKLNLVASRVTEDSARLSWTAPDAAFDSFNIAIWEPGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSPPLIARFTT
102	MLPAPKLNLVVSRVTEDESARLSWTAPDAAFDSFNIAIWEPGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSPPLIARFTT
103	MLPAPKLNLVVSRVTEDESARLSWTAPDAAFDSFAIAIWEPGIGGEAIWLRVPGSERSYD


	LTGLKPGTEYKVWIHGVKGGASSPPLIARFTT
104	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNISYWEPGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSPPLIARFTT
105	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YAEPEGIGGEAIWLRVPGSRSYDL TGLKPGTEYKVWIHGVKGGASSPPLIARFTT
106	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEAGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSPPLIARFTT
107	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPAGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSPPLIARFTT
108	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGAGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSPPLIARFTT
109	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGIAGEAIWLRVPGSERSYDL TGLKPGTEYKVWIHGVKGGASSPPLIARFTT
110	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGIGGEAIALRVPGSERSYDL TGLKPGTEYKVWIHGVKGGASSPPLIARFTT
111	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGIGGEAIWLAVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSPPLIARFTT
112	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSPPLIARFTT
113	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSPPLIARFTT
114	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIAGVKGGASSPPLIARFTT
115	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGSSPPLIARFTT
116	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASAPPLIARFTT
117	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSAPPLIARFTT
118	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSPLAARFTT
119	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFNIA YWEPGIGGEAIWLRVPGSERSYD LTGLKPGTEYKVWIHGVKGGASSPLIAAFTT

[00109] In some embodiments, the dsRNA agent comprises mismatch(es) with the target, within the duplex, or combinations thereof. The mismatch can occur in the overhang region or the duplex region. The base pair can be ranked on the basis of their propensity to promote dissociation or melting (e.g., on the free energy of association or dissociation of a particular pairing, the simplest approach is to examine the pairs on an individual pair basis, though next neighbor or similar analysis can also be used). In terms of promoting dissociation: A:U is preferred over G:C; G:U is preferred over G:C; and I:C is preferred over G:C (I=inosine). Mismatches, e.g., non-canonical or other than canonical pairings (as described elsewhere herein) are preferred over canonical (A:T, A:U, G:C) pairings; and pairings which include a universal base are preferred over canonical pairings.


[00110] In some embodiments, the dsRNA agent can comprise a phosphorus-containing group at the 5'-end of the sense strand or antisense strand. The 5'-end phosphorus-containing group can be 5'-end phosphate (5'-P), 5'-end phosphorothioate (5'-PS), 5'-end phosphorodithioate (5'-PS₂), 5'-end vinylphosphonate (5'-VP), 5'-end methylphosphonate (MePhos), 5'-end mesyl phosphoramidate (5'-MsPA), or 5'-deoxy-5'-C-malonyl. When the 5'-end phosphorus-containing group is 5'-end vinylphosphonate (5'-VP), the 5'-VP can be either 5'-E-VP isomer, such as trans-vinylphosphate or cis-vinylphosphate, or mixtures thereof. Representative structures of these modifications can be found in, for example, U.S. Patent No. 10,233,448, which is hereby incorporated by reference in its entirety.

[00111] In some embodiments, nucleotide analogues or synthetic nucleotide base comprise a nucleic acid with a modification at a 2' hydroxyl group of the ribose moiety. In some instances, the modification includes an H, OR, R, halo, SH, SR, NH₂, NHR, NR₂, or CN, wherein R is an alkyl moiety. Exemplary alkyl moiety includes, but is not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, C₁-C₁₀ chain lengths both linear and branched. In some instances, the alkyl moiety further comprises a modification. In some instances, the modification comprises an azo group, a keto group, an aldehyde group, a carboxyl group, a nitro group, a nitroso group, a nitrile group, a heterocycle (e.g., imidazole, hydrazine or hydroxylamino) group, an isocyanate or cyanate group, or a sulfur containing group (e.g., sulfoxide, sulfone, sulfide, and disulfide). In some instances, the alkyl moiety further comprises additional hetero atom such as O, S, N, Se and each of these hetero atoms can be further substituted with alkyl groups as described above. In some instances, the carbon of the heterocyclic group is substituted by a nitrogen, oxygen or sulfur. In some instances, the heterocyclic substitution includes but is not limited to, morpholino, imidazole, and pyrrolidino.

[00112] In some instances, the modification at the 2' hydroxyl group is a 2'-O-methyl modification or a 2'-O-methoxyethyl (2'-O-MOE) modification. Exemplary chemical structures of a 2'-O-methyl modification of an adenosine molecule and 2'-O-methoxyethyl modification of an uridine are illustrated below.




[00113] In some instances, the modification at the 2' hydroxyl group is a 2'-O-aminopropyl modification in which an extended amine group comprising a propyl linker binds the amine group to the 2' oxygen. In some instances, this modification neutralizes the phosphate derived overall negative charge of the oligonucleotide molecule by introducing one positive charge from the amine group per sugar and thereby improves cellular uptake properties due to its zwitterionic properties. An exemplary chemical structure of a 2'-O-aminopropyl nucleoside phosphoramidite is illustrated below.



2'-O-aminopropyl nucleoside phosphoramidite


[00114] In some instances, the modification at the 2' hydroxyl group is a locked or bridged ribose modification (e.g., locked nucleic acid or LNA) in which the oxygen molecule bound at the 2' carbon is linked to the 4' carbon by a methylene group, thus forming a 2'-C,4'-C-oxy- methylene-linked bicyclic ribonucleotide monomer. Exemplary representations of the chemical structure of LNA are illustrated below. The representation shown to the left highlights the chemical connectivities of an LNA monomer. The representation shown to the right highlights the locked 3'-endo (3E) conformation of the furanose ring of an LNA monomer.



LNA (Locked Nucleic Acids)

[00115] In some instances, the modification at the 2' hydroxyl group comprises ethylene nucleic acids (ENA) such as for example 2'-4'-ethylene-bridged nucleic acid, which locks the sugar conformation into a C3'-endo sugar puckering conformation. ENA are part of the bridged

nucleic acids class of modified nucleic acids that also comprises LNA. Exemplary chemical structures of the ENA and bridged nucleic acids are illustrated below.




[00116] In some embodiments, additional modifications at the 2' hydroxyl group include 2'- deoxy, 2'-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O- dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA).


[00117] In some embodiments, nucleotide analogues comprise modified bases such as, but not limited to, 5-propynyluridine, 5-propynylcytidine, 6- methyladenine, 6-methylguanine, N, N, - dimethyladenine, 2-propyladenine, 2-propylguanine, 2-aminoadenine, 1-methylinosine, 3-methyluridine, 5-methylcytidine, 5-methyluridine and other nucleotides having a modification at the 5 position, 5- (2- amino) propyl uridine, 5-halocytidine, 5-halouridine, 4-acetylcytidine, 1-methyladenosine, 2-methyladenosine, 3-methylcytidine, 6-methyluridine, 2- methylguanosine, 7-methylguanosine, 2, 2-dimethylguanosine, 5- methylaminoethyluridine, 5-methoxyuridine, deazanucleotides such as 7-deaza- adenosine, 6-azouridine, 6-azocytidine, 6-azothymidine, 5-methyl-2-thiouridine, other thio bases such as 2-thiouridine and 4-thiouridine and 2-thiocytidine, dihydrouridine, pseudouridine, queuosine, archaeosine, naphthyl and substituted naphthyl groups, any O-and N-alkylated purines and pyrimidines such as N6-methyladenosine, 5-methylcarbonylmethyluridine, uridine 5-oxyacetic acid, pyridine-4-one, pyridine-2-one, phenyl and modified phenyl groups such as aminophenol or 2,4, 6-trimethoxy benzene, modified cytosines that act as G-clamp nucleotides, 8-substituted adenines and guanines, 5-substituted uracils and thymines, azapyrimidines, carboxyhydroxyalkyl nucleotides, carboxyalkylaminoalkyl nucleotides, and alkylcarbonylalkylated nucleotides. Modified nucleotides also include those nucleotides that are modified with respect to the sugar moiety, as well as nucleotides having sugars or analogs thereof that are not ribosyl. For example, the sugar moieties, in some cases are or be based on, mannoses, arabinoses, glucopyranoses, galactopyranoses, 4'-thioribose, and other sugars, heterocycles, or carbocycles. The term nucleotide also includes what are known in the art

as universal bases. By way of example, universal bases include but are not limited to 3-nitropyrrole, 5-nitroindole, or nebularine.

[00118] In some embodiments, nucleotide analogues further comprise morpholinos, peptide nucleic acids (PNAs), methylphosphonate nucleotides, thiolphosphonate nucleotides, 2'-fluoro N3-P5'-phosphoramidites, 1', 5'- anhydrohexitol nucleic acids (HNAs), or a combination thereof. Morpholino or phosphorodiamidate morpholino oligo (PMO) comprises synthetic molecules whose structure mimics natural nucleic acid structure by deviates from the normal sugar and phosphate structures. In some instances, the five-member ribose ring is substituted with a six member morpholino ring containing four carbons, one nitrogen and one oxygen. In some cases, the ribose monomers are linked by a phosphordiamidate group instead of a phosphate group. In such cases, the backbone alterations remove all positive and negative charges making morpholinos neutral molecules capable of crossing cellular membranes without the aid of cellular delivery agents such as those used by charged oligonucleotides.



[00119] In some embodiments, peptide nucleic acid (PNA) does not contain sugar ring or phosphate linkage and the bases are attached and appropriately spaced by oligoglycine-like molecules, therefore, eliminating a backbone charge.



[00120] In some embodiments, one or more modifications optionally occur at the internucleotide linkage. In some instances, modified internucleotide linkage include, but is not limited to, phosphorothioates, , mesyl phosphoramidate, phosphorodithioates, methylphosphonates, 5'- alkylenephosphonates, 5'-methylphosphonate, 3'-alkylene phosphonates, borontrifluoridates, borano phosphate esters and selenophosphates of 3'-5' linkage

or 2'-5' linkage, phosphotriesters, thionoalkylphosphotriesters, hydrogen phosphonate linkages, alkyl phosphonates, alkylphosphonothioates, arylphosphonothioates, phosphoroselenoates, phosphorodiselenoates, phosphinates, phosphoramidates, 3'-alkylphosphoramidates, aminoalkylphosphoramidates, thionophosphoramidates, phosphoropiperazidates, phosphoroanilothioates, phosphoroanilidates, ketones, sulfones, sulfonamides, carbonates, carbamates, methylenehydrazos, methylenedimethylhydrazos, formacetals, thioformacetals, oximes, methyleneiminos, methylenemethyliminos, thioamides, linkages with riboacetyl groups, aminoethyl glycine, silyl or siloxane linkages, alkyl or cycloalkyl linkages with or without heteroatoms of, for example, 1 to 10 carbons that are saturated or unsaturated and/or substituted and/or contain heteroatoms, linkages with morpholino structures, amides, polyamides wherein the bases are attached to the aza nitrogens of the backbone directly or indirectly, and combinations thereof. Phosphorothioate antisense oligonucleotides (PS ASO) are antisense oligonucleotides comprising a phosphorothioate linkage. Mesyl phosphoramidate antisense oligonucleotides (MsPA ASO) are antisense oligonucleotides comprising a mesyl phosphoramidate linkage.

[00121] In some instances, the modification is a methyl or thiol modification such as methylphosphonate, mesyl phosphoramidate, or thiolphosphonate modification. In some instances, a modified nucleotide includes, but is not limited to, 2'-fluoro N3- P5'-phosphoramidites.

[00122] In some instances, a modified nucleotide includes, but is not limited to, hexitol nucleic acid (or 1', 5'- anhydrohexitol nucleic acids (HNA)).

[00123] In some embodiments, one or more modifications further optionally include modifications of the ribose moiety, phosphate backbone and the nucleoside, or modifications of the nucleotide analogues at the 3' or the 5' terminus. For example, the 3' terminus optionally include a 3' cationic group, or by inverting the nucleoside at the 3'-terminus with a 3'-3' linkage. In another alternative, the 3'-terminus is optionally conjugated with an aminoalkyl group, e.g., a 3' C5-aminoalkyl dT. In an additional alternative, the 3'-terminus is optionally conjugated with an abasic site, e.g., with an apurinic or apyrimidinic site. In some instances, the 5'-terminus is conjugated with an aminoalkyl group, e.g., a 5'-O-alkylamino substituent. In some cases, the 5'-terminus is conjugated with an abasic site, e.g., with an apurinic or apyrimidinic site.

[00124] In some embodiments, the oligonucleotide molecule comprises one or more of the synthetic nucleotide analogues described herein. In some instances, the oligonucleotide molecule comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 25, or more of the synthetic nucleotide analogues described herein. In some embodiments, the synthetic nucleotide

analogues include 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, 2'-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified, LNA, ENA, PNA, HNA, morpholino, methylphosphonate nucleotides, thiolphosphonate nucleotides, 2'-fluoro N3-P5'-phosphoramidites, or a combination thereof. In some instances, the oligonucleotide molecule comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 25, or more of the synthetic nucleotide analogues selected from 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, 2'-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified, LNA, ENA, PNA, HNA, morpholino, methylphosphonate nucleotides, thiolphosphonate nucleotides, 2'-fluoro N3-P5'-phosphoramidites, or a combination thereof. In some instances, the oligonucleotide molecule comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 25, or more of 2'-O-methyl modified nucleotides. In some instances, the oligonucleotide molecule comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 25, or more of 2'-O-methoxyethyl (2'-O-MOE) modified nucleotides. In some instances, the oligonucleotide molecule comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 25, or more of thiolphosphonate nucleotides.

[00125] In some instances, the oligonucleotide molecule comprises at least one of: from about 5% to about 100% modification, from about 10% to about 100% modification, from about 20% to about 100% modification, from about 30% to about 100% modification, from about 40% to about 100% modification, from about 50% to about 100% modification, from about 60% to about 100% modification, from about 70% to about 100% modification, from about 80% to about 100% modification, and from about 90% to about 100% modification. In some instances, the oligonucleotide molecule comprises 100% modification

[00126] In some cases, the oligonucleotide molecule comprises at least one of: from about 10% to about 90% modification, from about 20% to about 90% modification, from about 30% to about 90% modification, from about 40% to about 90% modification, from about 50% to about 90% modification, from about 60% to about 90% modification, from about 70% to about 90% modification, and from about 80% to about 100% modification.

[00127] In some cases, the oligonucleotide molecule comprises at least one of: from about 10% to about 80% modification, from about 20% to about 80% modification, from about 30% to about 80% modification, from about 40% to about 80% modification, from about 50% to about

80% modification, from about 60% to about 80% modification, and from about 70% to about 80% modification.

[00128] In some instances, the oligonucleotide molecule comprises at least one of: from about 10% to about 70% modification, from about 20% to about 70% modification, from about 30% to about 70% modification, from about 40% to about 70% modification, from about 50% to about 70% modification, and from about 60% to about 70% modification.

[00129] In some instances, the oligonucleotide molecule comprises at least one of: from about 10% to about 60% modification, from about 20% to about 60% modification, from about 30% to about 60% modification, from about 40% to about 60% modification, and from about 50% to about 60% modification.

[00130] In some cases, the oligonucleotide molecule comprises at least one of: from about 10% to about 50% modification, from about 20% to about 50% modification, from about 30% to about 50% modification, and from about 40% to about 50% modification.

[00131] In some cases, the oligonucleotide molecule comprises at least one of: from about 10% to about 40% modification, from about 20% to about 40% modification, and from about 30% to about 40% modification.

[00132] In some cases, the oligonucleotide molecule comprises at least one of: from about 10% to about 30% modification, and from about 20% to about 30% modification.

[00133] In some cases, the oligonucleotide molecule comprises from about 10% to about 20% modification.

[00134] In some cases, the oligonucleotide molecule comprises from about 15% to about 90%, from about 20% to about 80%, from about 30% to about 70%, or from about 40% to about 60% modifications.

[00135] In additional cases, the oligonucleotide molecule comprises at least about 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 99% modification.

[00136] In some embodiments, the oligonucleotide molecule comprises at least about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, or about 40 modifications.

[00137] In some instances, the oligonucleotide molecule comprises at least about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about

22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, or about 40 modified nucleotides.

[00138] In some instances, from about 5 to about 100% of the oligonucleotide molecule comprise the synthetic nucleotide analogues described herein. In some instances, about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% of the oligonucleotide molecule comprise the synthetic nucleotide analogues described herein. In some instances, about 5% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 10% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 15% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 20% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 25% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 30% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 35% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 40% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 45% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 50% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 55% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 60% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 65% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 70% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 75% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 80% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 85% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 90% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 95% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 96% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In

some instances, about 97% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 98% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 99% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some instances, about 100% of the oligonucleotide molecule comprises the synthetic nucleotide analogues described herein. In some embodiments, the synthetic nucleotide analogues include 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, 2'-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified, LNA, ENA, PNA, HNA, morpholino, methylphosphonate nucleotides, thiolphosphonate nucleotides, 2'-fluoro N3-P5'-phosphoramidites, or a combination thereof.

[00139] In some embodiments, the oligonucleotide molecule comprises from about 1 to about 25 modifications in which the modification comprises a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 1 modification in which the modification comprises a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 2 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 3 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 4 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 5 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 6 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 7 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 8 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 9 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 10 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 11 modifications in which the modifications

[illegible]

herein. In some embodiments, the oligonucleotide molecule comprises about 30 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 31 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 32 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 33 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 34 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 35 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 36 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 37 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 38 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 39 modifications in which the modifications comprise a synthetic nucleotide analogue described herein. In some embodiments, the oligonucleotide molecule comprises about 40 modifications in which the modifications comprise a synthetic nucleotide analogue described herein.

[00140] In some embodiments, an oligonucleotide molecule is assembled from two separate polynucleotides wherein one polynucleotide comprises the sense strand and the second polynucleotide comprises the antisense strand of the oligonucleotide molecule. In other embodiments, the sense strand is connected to the antisense strand via a linker molecule, which in some instances is a polynucleotide linker or a non-nucleotide linker.

[00141] In some embodiments, an oligonucleotide molecule comprises a sense strand and antisense strand, wherein pyrimidine nucleotides in the sense strand comprises 2'-O-methylpyrimidine nucleotides and purine nucleotides in the sense strand comprise 2'-deoxy purine nucleotides. In some embodiments, an oligonucleotide molecule comprises a sense strand and antisense strand, wherein pyrimidine nucleotides present in the sense strand comprise 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein purine nucleotides present in the sense strand comprise 2'-deoxy purine nucleotides.

[00142] In some embodiments, an oligonucleotide molecule comprises a sense strand and antisense strand, wherein the pyrimidine nucleotides when present in said antisense strand are 2'-deoxy-2'-fluoro pyrimidine nucleotides and the purine nucleotides when present in said antisense strand are 2'-O-methyl purine nucleotides.

[00143] In some embodiments, an oligonucleotide molecule comprises a sense strand and antisense strand, wherein the pyrimidine nucleotides when present in said antisense strand are 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein the purine nucleotides when present in said antisense strand comprise 2'-deoxy-purine nucleotides.


[00144] In some embodiments, an oligonucleotide molecule comprises a sense strand and antisense strand, and at least one of sense strand and antisense strands has a plurality of (e.g., two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, etc) 2'-O-methyl or 2'-deoxy-2'-fluoro modified nucleotides. In some embodiments, at least two, three, four, five, six, or seven out of the a plurality of 2'-O-methyl or 2'-deoxy-2'-fluoro modified nucleotides are consecutive nucleotides. In some embodiments, consecutive 2'-O-methyl or 2'-deoxy-2'-fluoro modified nucleotides are located at the 5'-end of the sense strand and/or the antisense strand. In some embodiments, consecutive 2'-O-methyl or 2'-deoxy-2'-fluoro modified nucleotides are located at the 3'-end of the sense strand and/or the antisense strand. In some embodiments, the sense strand of oligonucleotide molecule includes at least four, at least five, at least six consecutive 2'-O-methyl modified nucleotides at its 5' end and/or 3' end, or both. Optionally, in such embodiments, the sense strand of oligonucleotide molecule includes at least one, at least two, at least three, at least four 2'-deoxy-2'-fluoro modified nucleotides at the 3' end of the at least four, at least five, at least six consecutive 2'-O-methyl modified nucleotides at the polynucleotides' 5' end, or at the 5' end of the at least four, at least five, at least six consecutive 2'-O-methyl modified nucleotides at polynucleotides' 3' end. Also optionally, such at least two, at least three, at least four 2'-deoxy-2'-fluoro modified nucleotides are consecutive nucleotides.

[00145] In some embodiments, an oligonucleotide molecule comprises a sense strand and antisense strand, and at least one of sense strand and antisense strand has 2'-O-methyl modified nucleotide located at the 5'-end of the sense strand and/or the antisense strand. In some embodiments, at least one of sense strand and antisense strands has 2'-O-methyl modified nucleotide located at the 3'-end of the sense strand and/or the antisense strand. In some embodiments, the 2'-O-methyl modified nucleotide located at the 5'-end of the sense strand and/or the antisense strand is a purine nucleotide. In some embodiments, the 2'-O-methyl

modified nucleotide located at the 5'-end of the sense strand and/or the antisense strand is a pyrimidine nucleotide.

[00146] In some embodiments, an oligonucleotide molecule comprises a sense strand and antisense strand, and one of sense strand and antisense strand has at least two consecutive 2'-deoxy-2'-fluoro modified nucleotides located at the 5'-end, while another strand has at least two consecutive 2'-O-methyl modified nucleotides located at the 5'-end. In some embodiments, where the strand has at least two consecutive 2'-deoxy-2'-fluoro modified nucleotides located at the 5'-end, the strand also includes at least two, at least three consecutive 2'-O-methyl modified nucleotides at the 3'-end of the at least two consecutive 2'-deoxy-2'-fluoro modified nucleotides. In some embodiments, one of sense strand and antisense strand has at least two, at least three, at least four, at least five, at least six, or at least seven consecutive 2'-O-methyl modified nucleotides that are linked to a 2'-deoxy-2'-fluoro modified nucleotide on its 5'-end and/or 3'-end. In some embodiments, one of sense strand and antisense strand has at least four, at least five nucleotides that have alternating 2'-O-methyl modified nucleotide and 2'-deoxy-2'-fluoro modified nucleotide.

[00147] In some embodiments, the oligonucleotide molecule, such as a siRNA, has the formula as illustrated in Formula I:



wherein each nucleotide represented by N, is independently, A, U, C, or G or a modified nucleotide base, such as those provided for herein. The N₁ nucleotides of the sense strand and the antisense strand represent the 5' end of the respective strands. For clarity, although Formula I utilizes N₁, N₂, N₃, etc. in both the sense and the antisense strand, the nucleotide bases do not need to be the same and are not intended to be the same. The siRNA that is illustrated in Formula I would be complementary to a target sequence.

[00148] For example, in some embodiments, the sense strand comprises a 2'-O-methyl modified nucleotide with a phosphorothioate (PS) modified backbone at N₁ and N₂, a 2'-fluoro modified nucleotide at N₃, N₇, N₈, N₉, N₁₂, and N₁₇, and a 2'-O-methyl modified nucleotide at N₄, N₅, N₆, N₁₀, N₁₁, N₁₃, N₁₄, N₁₅, N₁₆, N₁₈, and N₁₉.

[00149] In some embodiments, the antisense strand comprises a vinylphosphonate moiety attached to N₁, a 2'-fluoro- modified nucleotide with a phosphorothioate (PS) modified backbone at N₂, a 2'-O-methyl modified nucleotide at N₃, N₄, N₅, N₆, N₇, N₈, N₉, N₁₀, N₁₁, N₁₂, N₁₃, N₁₅,

N₁₆, N₁₇, N₁₈, and N₁₉, a 2'-fluoro- modified nucleotide at N₁₄, and a 2'-O-methyl modified nucleotide with a phosphorothioate (PS) modified backbone at N₂₀ and N₂₁.

[00150] In some embodiments, an oligonucleotide molecule comprises a sense strand and an antisense strand, wherein the sense strand includes a terminal cap moiety at the 5'-end, the 3'-end, or both of the 5' and 3' ends of the sense strand. In other embodiments, the terminal cap moiety is an inverted deoxy abasic moiety.

[00151] In some embodiments, an oligonucleotide molecule comprises a sense strand and an antisense strand, wherein the antisense strand comprises a glyceryl modification at the 3' end of the antisense strand.

[00152] In some embodiments, an oligonucleotide molecule comprises a sense strand and an antisense strand, in which the sense strand comprises one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or about one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and in which the antisense strand comprises about 1 to about 10 or more, specifically about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In other embodiments, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, pyrimidine nucleotides of the sense and/or antisense strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

[00153] In some embodiments, an oligonucleotide molecule comprises a sense strand and an antisense strand, in which the sense strand comprises about 1 to about 25, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide

linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and in which the antisense strand comprises about 1 to about 25 or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In other embodiments, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, pyrimidine nucleotides of the sense and/or antisense strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without about 1 to about 25 or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

[00154] In some embodiments, an oligonucleotide molecule comprises a sense strand and an antisense strand, in which the antisense strand comprises one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages, and/or about one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand and/or antisense strand, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand. In some embodiments, the antisense strand comprises about 1 to about 10 or more, specifically about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In other embodiments, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more pyrimidine nucleotides of the sense and/or antisense strand are

chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'- end, or both of the 3' and 5'-ends, being present in the same or different strand.

[00155] In some embodiments, an oligonucleotide molecule comprises a sense strand and an antisense strand, in which the antisense strand comprises about 1 to about 25 or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and the antisense strand comprises about 1 to about 25 or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In other embodiments, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more pyrimidine nucleotides of the sense and/or antisense strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without about 1 to about 5, for example about 1, 2, 3, 4, 5 or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'- end, or both of the 3'- and 5'-ends, being present in the same or different strand.

[00156] In some embodiments, an oligonucleotide molecule described herein is a chemically- modified short interfering nucleic acid molecule having about 1 to about 25, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate internucleotide linkages in each strand of the oligonucleotide molecule. In some embodiments, an oligonucleotide molecule comprises a sense strand and an antisense strand, and the antisense strand comprises a phosphate backbone modification at the 3' end of the antisense strand. Alternatively and/or additionally, an oligonucleotide molecule comprises a

sense strand and an antisense strand, and the sense strand comprises a phosphate backbone modification at the 5' end of the antisense strand. In some instances, the phosphate backbone modification is a phosphorothioate. In some instances, the phosphate backbone modification is a phosphorodithioate. In some instances, the phosphate backbone modification is a phosphonate. In some instances, the phosphate backbone modification is a phosphoramidate. In some instances, the phosphate backbone modification is a mesyl phosphoramidate. In some embodiments, the sense or antisense strand has three consecutive nucleosides that are coupled via two phosphorothioate backbone. In some embodiments, the sense or antisense strand has three consecutive nucleosides that are coupled via two phosphorodithioate backbone. In some embodiments, the sense or antisense strand has three consecutive nucleosides that are coupled via two phosphonate backbone. In some embodiments, the sense or antisense strand has three consecutive nucleosides that are coupled via two phosphoramidate backbone. In some embodiments, the sense or antisense strand has three consecutive nucleosides that are coupled via two mesyl phosphoramidate backbone.

[00157] In another embodiment, an oligonucleotide molecule described herein comprises 2'-5' internucleotide linkages. In some instances, the 2'-5' internucleotide linkage(s) is at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of one or both sequence strands. In addition instances, the 2'-5' internucleotide linkage(s) is present at various other positions within one or both sequence strands, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more including every internucleotide linkage of a pyrimidine nucleotide in one or both strands of the oligonucleotide molecule comprise a 2'-5' internucleotide linkage, or about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more including every internucleotide linkage of a purine nucleotide in one or both strands of the oligonucleotide molecule comprise a 2'-5' internucleotide linkage.

[00158] In some embodiments, an oligonucleotide molecule is a single stranded molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the oligonucleotide molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the oligonucleotide molecule are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the oligonucleotide molecule are 2'-deoxy purine nucleotides (e.g., wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), and a terminal cap modification, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the oligonucleotide molecule optionally

further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the oligonucleotide molecule, wherein the terminal nucleotides further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate or mesyl phosphoramidate internucleotide linkages, and wherein the oligonucleotide molecule optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

[00159] In some cases, one or more of the synthetic nucleotide analogues described herein are resistant toward nucleases such as for example ribonuclease such as RNase H, deoxyribonuclease such as DNase, or exonuclease such as 5'-3' exonuclease and 3'-5' exonuclease when compared to natural polynucleic acid molecules and endonucleases. In some instances, synthetic nucleotide analogues comprising 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, 2'-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified, LNA, ENA, PNA, HNA, morpholino, methylphosphonate nucleotides, thiophosphonate nucleotides, 2'-fluoro N3-P5'-phosphoramidites, or combinations thereof are resistant toward nucleases such as for example ribonuclease such as RNase H, deoxyribonuclease such as DNase, or exonuclease such as 5'-3' exonuclease and 3'-5' exonuclease. In some instances, 2'-O-methyl modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, 2'-O-methoxyethyl (2'-O-MOE) modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, 2'-O-aminopropyl modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, 2'-deoxy modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, 2'-deoxy-2'-fluoro modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, 2'-O-aminopropyl (2'-O-AP) modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, 2'-O-dimethylaminoethyl (2'-O-DMAOE) modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, 2'-O-dimethylaminopropyl (2'-O-DMAP) modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE) modified oligonucleotide molecule is nuclease

resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, 2'-O-N-methylacetamido (2'-O-NMA) modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, LNA modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, ENA modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, HNA modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, morpholinos is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, PNA modified oligonucleotide molecule is resistant to nucleases (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, methylphosphonate nucleotides modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, thiophosphonate nucleotides modified oligonucleotide molecule is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, oligonucleotide molecule comprising 2'-fluoro N3-P5'-phosphoramidites is nuclease resistance (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistance). In some instances, the 5' conjugates described herein inhibit 5'-3' exonucleolytic cleavage. In some instances, the 3' conjugates described herein inhibit 3'-5' exonucleolytic cleavage.

[00160] In some embodiments, one or more of the synthetic nucleotide analogues described herein have increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. The one or more of the synthetic nucleotide analogues comprising 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, 2'-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified, LNA, ENA, PNA, HNA, morpholino, methylphosphonate nucleotides, thiophosphonate nucleotides, or 2'-fluoro N3-P5'-phosphoramidites have increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-methyl modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-methoxyethyl (2'-O-MOE) modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-aminopropyl modified oligonucleotide molecule has increased binding affinity toward their

mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-deoxy modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-deoxy-2'-fluoro modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-aminopropyl (2'-O-AP) modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-dimethylaminoethyl (2'-O-DMAOE) modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-dimethylaminopropyl (2'-O-DMAP) modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE) modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-N-methylacetamido (2'-O-NMA) modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, LNA modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, ENA modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, PNA modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, HNA modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, morpholino modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, methylphosphonate nucleotides modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, thiolphosphonate nucleotides modified oligonucleotide molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, oligonucleotide molecule comprising 2'-fluoro N3-P5'-phosphoramidites has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some cases, the increased affinity is illustrated with a lower K_d, a higher melt temperature (T_m), or a combination thereof.

[00161] In some embodiments, an oligonucleotide molecule described herein is a chirally pure (or stereo pure) polynucleic acid molecule, or a polynucleic acid molecule comprising a single enantiomer. In some instances, the oligonucleotide molecule comprises L-nucleotide. In some instances, the oligonucleotide molecule comprises D-nucleotides. In some instance, an oligonucleotide molecule composition comprises less than 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1%, or less of its mirror enantiomer. In some cases, an oligonucleotide molecule composition comprises less than 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1%, or less of a racemic mixture.

[00162] In some embodiments, an oligonucleotide molecule described herein is further modified to include an aptamer conjugating moiety. In some instances, the aptamer conjugating moiety is a DNA aptamer conjugating moiety. In some instances, the aptamer conjugating moiety is Alphamer, which comprises an aptamer portion that recognizes a specific cell-surface target and a portion that presents a specific epitopes for attaching to circulating antibodies.

[00163] In additional embodiments, an oligonucleotide molecule described herein is modified to increase its stability. In some embodiment, the oligonucleotide molecule is RNA (e.g., siRNA). In some instances, the oligonucleotide molecule is modified by one or more of the modifications described above to increase its stability. In some cases, the oligonucleotide molecule is modified at the 2' hydroxyl position, such as by 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, 2'-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modification or by a locked or bridged ribose conformation (e.g., LNA or ENA). In some cases, the oligonucleotide molecule is modified by 2'-O-methyl and/or 2'-O-methoxyethyl ribose. In some cases, the oligonucleotide molecule also includes morpholinos, PNAs, HNA, methylphosphonate nucleotides, thiolphosphonate nucleotides, and/or 2'-fluoro N3-P5'-phosphoramidites to increase its stability. In some instances, the oligonucleotide molecule is a chirally pure (or stereo pure) oligonucleotide molecule. In some instances, the chirally pure (or stereo pure) oligonucleotide molecule is modified to increase its stability. Suitable modifications to the RNA to increase stability for delivery will be apparent to the skilled person.

[00164] In some embodiments, the oligonucleotide molecule comprises 2' modifications. In some embodiments, the nucleotides of the oligonucleotide molecule at positions 3, 7, 8, 9, 12, and 17 from the 5' end of the sense strand are not modified with a 2'-O-methyl modification. In some embodiments, the nucleotides of the oligonucleotide molecule at positions 3, 7, 8, 9, 12, and 17 from the 5' end of the sense strand are modified with a 2'-fluoro modification. In some

embodiments, the nucleotides of the oligonucleotide molecule at positions 2 and 14 from the 5' end of the anti-sense strand are not modified with a 2'O-methyl modification. In some embodiments, the nucleotides of the oligonucleotide molecule at positions 2 and 14 from the 5' end of the anti-sense strand are modified with a 2'fluoro modification. In some embodiments, any of the nucleotides may further comprise a 5'-phosphorothioate group modification. In some embodiments, the nucleotides of the oligonucleotide molecule at positions 1 and 2 from the 5' end of the sense strand are modified with a 5'-phosphorothioate group modification. In some embodiments, the nucleotides of the oligonucleotide molecule at positions 1, 2, 20, and 21 from the 5' end of the antisense strand are modified with a 5'-phosphorothioate group modification. In some embodiments, the 5' end of the sense or antisense strand of the oligonucleotide molecule may further comprise a vinylphosphonate modification. In some embodiments, the nucleotide of the oligonucleotide molecule at position 1 from the 5' end of the antisense strand is modified with a vinylphosphonate modification.

[00165] In some instances, the oligonucleotide molecule is a double-stranded polynucleotide molecule comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. In some instances, the oligonucleotide molecule is assembled from two separate polynucleotides, where one strand is the sense strand and the other is the antisense strand, wherein the antisense and sense strands are self-complementary (e.g., each strand comprises nucleotide sequence that is complementary to nucleotide sequence in the other strand; such as where the antisense strand and sense strand form a duplex or double stranded structure, for example wherein the double stranded region is about 19, 20, 21, 22, 23, or more base pairs); the antisense strand comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense strand comprises nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. Alternatively, the oligonucleotide molecule is assembled from a single oligonucleotide, where the self-complementary sense and antisense regions of the oligonucleotide molecule are linked by means of a nucleic acid based or non-nucleic acid-based linker(s).

[00166] In some cases, the oligonucleotide molecule is a polynucleotide with a duplex, asymmetric duplex, hairpin or asymmetric hairpin secondary structure, having self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a separate target nucleic acid molecule

or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. In other cases, the oligonucleotide molecule is a circular single-stranded polynucleotide having two or more loop structures and a stem comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof, and wherein the circular polynucleotide is processed either in vivo or in vitro to generate an active oligonucleotide molecule capable of mediating RNAi. In additional cases, the oligonucleotide molecule also comprises a single-stranded polynucleotide having nucleotide sequence complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof (for example, where such oligonucleotide molecule does not require the presence within the oligonucleotide molecule of nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof), wherein the single stranded polynucleotide further comprises a terminal phosphate group, such as a 5'-phosphate, or 5', 3'-diphosphate.

[00167] In some instances, an asymmetric hairpin is a linear oligonucleotide molecule comprising an antisense region, a loop portion that comprises nucleotides or non-nucleotides, and a sense region that comprises fewer nucleotides than the antisense region to the extent that the sense region has enough complimentary nucleotides to base pair with the antisense region and form a duplex with loop. For example, an asymmetric hairpin oligonucleotide molecule comprises an antisense region having length sufficient to mediate RNAi in a cell or in vitro system (e.g., about 19 to about 22 nucleotides) and a loop region comprising about 4 to about 8 nucleotides, and a sense region having about 3 to about 18 nucleotides that are complementary to the antisense region. In some cases, the asymmetric hairpin oligonucleotide molecule also comprises a 5'-terminal phosphate group that is chemically modified. In additional cases, the loop portion of the asymmetric hairpin oligonucleotide molecule comprises nucleotides, non-nucleotides, linker molecules, or conjugate molecules.

[00168] In some embodiments, an asymmetric duplex is an oligonucleotide molecule having two separate strands comprising a sense region and an antisense region, wherein the sense region comprises fewer nucleotides than the antisense region to the extent that the sense region has enough complimentary nucleotides to base pair with the antisense region and form a duplex. For example, an asymmetric duplex oligonucleotide molecule comprises an antisense region having length sufficient to mediate RNAi in a cell or in vitro system (e.g., about 19 to about 22

nucleotides) and a sense region having about 3 to about 19 nucleotides that are complementary to the antisense region.

[00169] In some cases, a universal base refers to nucleotide base analogs that form base pairs with each of the natural DNA/RNA bases with little discrimination between them. Non-limiting examples of universal bases include C-phenyl, C-naphthyl and other aromatic derivatives, inosine, azole carboxamides, and nitroazole derivatives such as 3-nitropyrrole, 4-nitroindole, 5-nitroindole, and 6-nitroindole as known in the art.

[00170] In some embodiments, the dsRNA agents are 5' phosphorylated or include a phosphoryl analog at the 5' prime terminus. 5'-phosphate modifications include those which are compatible with RISC mediated gene silencing. Suitable modifications include: 5'-monophosphate ((HO)₂(O)P--O-5'); 5'-diphosphate ((HO)₂(O)P--O--P(HO)(O)--O-5'); 5'-triphosphate ((HO)₂(O)P--O--(HO)(O)P--O--P(HO)(O)--O-5'); 5'-guanosine cap (7-methylated or non-methylated) (7m-G-O-5'-(HO)(O)P--O--(HO)(O)P--O--P(HO)(O)--O-5'); 5'-adenosine cap (Appp), and any modified or unmodified nucleotide cap structure (N--O-5'-(HO)(O)P--O--(HO)(O)P--O--P(HO)(O)--O-5'); 5'-monothiophosphate (phosphorothioate; (HO)₂(S)P--O-5'); 5'-monodithiophosphate (phosphorodithioate; (HO)(HS)(S)P--O-5'), 5'-phosphorothiolate ((HO)₂(O)P--S-5'); phosphorodithioate [--O₂PS₂--]; phosphonate [--PO(OH)₂--]; phosphoramidate [--O=P(OH)₂--]; mesyl phosphoramidate (CH₃)(SO₂)(N)P(O)₂--O-5'); any additional combination of oxygen/sulfur replaced monophosphate, diphosphate and triphosphates (e.g. 5'-alpha-thiotriphosphate, 5'-gamma-thiotriphosphate, etc.), 5'-phosphoramidates ((HO)₂(O)P--NH-5', (HO)(NH₂)(O)P--O-5'), 5'-alkylphosphonates (R=alkyl=methyl, ethyl, isopropyl, propyl, etc., e.g. RP(OH)(O)--O-5'-, 5'-alkenylphosphonates (i.e. vinyl, substituted vinyl), (OH)₂(O)P-5'-CH₂-), 5'-alkyletherphosphonates (R=alkylether=methoxymethyl (MeOCH₂-), ethoxymethyl, etc., e.g. RP(OH)(O)--O-5'-). In some embodiments, the modification can be placed in the antisense strand of a dsRNA agent.

[00171] In some embodiments, the sequence of the oligonucleotide molecule is at least 40%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, or 99.5% complementary to a target sequence of GYS1. In some embodiments, the target sequence of GYS1 is a nucleic acid sequence of about 10-50 base pair length, about 15-50 base pair length, 15-40 base pair length, 15-30 base pair length, or 15-25 base pair length sequences in GYS1, in which the first nucleotide of the target sequence starts at any nucleotide in GYS1 mRNA transcript in the coding region, or in the 5' or 3'-untranslated region (UTR). For example, the first nucleotide of the target sequence can be selected so that it starts at the nucleic acid location (nal, number starting from the 5'-end of the full length of GYS1 mRNA, e.g., the 5'-end first

nucleotide is nal.1) 1, nal 2, nal 3, nal 4, nal 5, nal 6, nal 7, nal 8, nal 9, nal 10, nal 11, nal 12, nal 13, nal 14, nal 15, nal 15, nal 16, nal 17, or any other nucleic acid location in the coding or noncoding regions (5' or 3'-untranslated region) of GYS1 mRNA. In some embodiments, the first nucleotide of the target sequence can be selected so that it starts at a location within, or between, nal 10- nal 15, nal 10- nal 20, nal 50- nal 60, nal 55- nal 65, nal 75- nal 85, nal 95- nal 105, nal 135- nal 145, nal 155- nal 165, nal 225- nal 235, nal 265- nal 275, nal 275- nal 245, nal 245- nal 255, nal 285- nal 335, nal 335- nal 345, nal 385- nal 395, nal 515- nal 525, nal 665- nal 675, nal 675- nal 685, nal 695- nal 705, nal 705- nal 715, nal 875- nal 885, nal 885- nal 895, nal 895- nal 905, nal 1035- nal 1045, nal 1045- nal 1055, nal 1125- nal 1135, nal 1135- nal 1145, nal 1145- nal 1155, nal 1155- nal 1165, nal 1125- nal 1135, nal 1155- nal 1165, nal 1225- nal 1235, nal 1235- nal 1245, nal 1275- nal 1245, nal 1245- nal 1255, nal 1265- nal 1275, nal 1125- nal 1135, nal 1155- nal 1165, nal 1225- nal 1235, nal 1235- nal 1245, nal 1275- nal 1245, nal 1245- nal 1255, nal 1265- nal 1275, nal 1275- nal 1285, nal 1335- nal 1345, nal 1345- nal 1355, nal 1525- nal 1535, nal 1535- nal 1545, nal 1605- nal 1615, nal 1615- c.1625, nal 1625- nal 1635, nal 1635- 1735, nal 1735-1835, nal 1835-1935, nal. 1836-1856, nal 1935-2000, nal 2000 -2100, nal 2100 -2200, nal 2200 -2260, nal 2260 -2400, nal 2400 -2500, nal 2500 -2600, nal 2600 -2700, nal 2700 -2800, nal 2800 -2500, nal 2500 -2600, nal 2600 -2700, nal 2700 -2800, nal 2800 -2860, etc. In some embodiments, the sequence of GYS1 mRNA is provided as NCBI Reference Sequence: NM_002103.

[00172] In some embodiments, the antisense strand of the dsRNA agent is 100% complementary to a target RNA to hybridize thereto and inhibits its expression through RNA interference. The target RNA can be any RNA expressed in a cell. In another embodiment, the cell is a tumor cell, a liver cell, a muscle cell, an immune cell, a dendritic cell, a heart cell, or a cell of the central nervous system. In another embodiment, the antisense strand of the dsRNA agent is at least 99%, at least 98%, at least 97%, at least 96%, 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, or at least 50% complementary to a target RNA. In some embodiments, the target RNA is GYS1 RNA. In some embodiments, the siRNA molecule is a siRNA that reduces the expression of GYS1. In some embodiments, the siRNA molecule is a siRNA that reduces the expression of GYS and does not reduce the expression of other RNAs by more than 50% in an assay described herein at a concentration of no more than 200 nM as described herein.

[00173] The siRNA can be targeted against any gene or RNA (*e.g.* mRNA) transcript of interest.

[00174] Other modifications and patterns of modifications can be found in, for example, U.S. Patent No. 10,233,448, which is hereby incorporated by reference.

[00175] Other modifications and patterns of modifications can be found in, for example, Anderson et al. **Nucleic Acids Research** 2021, 49 (16), 9026-9041, which is hereby incorporated by reference.

[00176] Other modifications and patterns of modifications can be found in, for example, PCT Publication No. WO2021/030778, which is hereby incorporated by reference.

[00177] Other modifications and patterns of modifications can be found in, for example, PCT Publication No. WO2021/030763, which is hereby incorporated by reference.

[00178] In some embodiments, the siRNA is linked to a protein, such as a FN3 domain. The siRNA can be linked to multiple FN3 domains that bind to the same target protein or different target proteins. In some embodiments, the linker is attached to the sense strand, which is used to facilitate the linkage of the sense strand to the FN3 domain.

[00179] In some embodiments, compositions are provided herein having a formula of $(X1)_n-(X2)_q-(X3)_y-L-X4$, wherein X1 is a first FN3 domain, X2 is second FN3 domain, X3 is a third FN3 domain or half-life extender molecule, L is a linker, and X4 is a nucleic acid molecule, such as, but not limited to a siRNA molecule, wherein n, q, and y are each independently 0 or 1. In some embodiments, X1, X2, and X3 bind to different target proteins. In some embodiments, y is 0. In some embodiments, n is 1, q is 0, and y is 0. In some embodiments, n is 1, q is 1, and y is 0. In some embodiments, n is 1, q is 1, and y is 1. In some embodiments, the third FN3 domain increases the half-life of the molecule as a whole as compared to a molecule without X3. In some embodiments, the half-life extending moiety is a FN3 domain that binds to albumin. Examples of such FN3 domains include, but are not limited to, those described in U.S. Patent Application Publication No. 20170348397 and U.S. Patent No. 9,156,887, which is hereby incorporated by reference in its entirety. The FN3 domains may incorporate other subunits for example via covalent interaction. In some embodiments, the FN3 domains further comprise a half-life extending moiety. Exemplary half-life extending moieties are albumin, albumin variants, albumin-binding proteins and/or domains, transferrin and fragments and analogues thereof, and Fc regions. Amino acid sequences of the human Fc regions are well known, and include IgG1, IgG2, IgG3, IgG4, IgM, IgA and IgE Fc regions. In some embodiments, the FN3 domains may incorporate a second FN3 domain that binds to a molecule that extends the half-life of the entire molecule, such as, but not limited to, any of the half-life extending moieties described herein. In some embodiments, the second FN3 domain binds to albumin, albumin variants, albumin-binding proteins and/or domains, and fragments and analogues thereof.

[00180] In some embodiments, compositions are provided herein having a formula of (X1)-(X2)-L-(X4), wherein X1 is a first FN3 domain, X2 is second FN3 domain, L is a linker, and X4 is a nucleic acid molecule. In some embodiments, X4 is a siRNA molecule. In some embodiments, X1 is a FN3 domain that binds to one of CD71. In some embodiments, X2 is a FN3 domain that binds to one of CD71. In some embodiments X1 and X2 do not bind to the same target protein. In some embodiments, X1 and X2 bind to the same target protein, but at different binding sites on the protein. In some embodiments, X1 and X2 bind to the same target protein. In some embodiments, X1 and X2 are FN3 domains that bind to CD71. In some embodiments, the composition does not comprise (e.g. is free of) a compound or protein that binds to ASGPR.

[00181] In some embodiments, compositions are provided herein having a formula of C-(X1)_n-(X2)_q[L-X4]-(X3)_y, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; X4 is an oligonucleotide molecule; and C is a polymer, wherein n, q, and y are each independently 0 or 1, are provided.

[00182] In some embodiments, compositions are provided herein having a formula of (X1)_n-(X2)_q[L-X4]-(X3)_y-C, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; X4 is an oligonucleotide molecule; and C is a polymer, wherein n, q, and y are each independently 0 or 1, are provided.

[00183] In some embodiments, compositions are provided herein having a formula of C-(X1)_n-(X2)_q[L-X4]L-(X3)_y, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; X4 is an oligonucleotide molecule; and C is a polymer, wherein n, q, and y are each independently 0 or 1, are provided.

[00184] In some embodiments, compositions are provided herein having a formula of (X1)_n-(X2)_q[L-X4]L-(X3)_y-C, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; X4 is an oligonucleotide molecule; and C is a polymer, wherein n, q, and y are each independently 0 or 1, are provided.

[00185] In some embodiments, compositions or complexes are provided having a formula of A₁-B₁, wherein A₁ has a formula of C-L₁-X_S and B₁ has a formula of X_{AS}-L₂-F₁, wherein:

C is a polymer, such as PEG;

L₁ and L₂ are each, independently, a linker;

X_S is a 5' to 3' oligonucleotide sense strand of a double stranded siRNA molecule;

X_{AS} is a 3' to 5' oligonucleotide antisense strand of a double stranded siRNA molecule;

F₁ is a polypeptide comprising at least one FN3 domain;

wherein X_S and X_{AS} form a double stranded oligonucleotide molecule to form the composition/complex.

[00186] In some embodiments, C can be a molecule that extends the half-life of the molecule. Examples of such moieties are described herein. In some embodiments, C can also be Endopporter, INF-7, TAT, polyarginine, polylysine, or an amphipathic peptide. These moieties can be used in place of or in addition to other half-life extending moieties provided for herein. In some embodiments, C can be a molecule that delivers the complex into the cell, the endosome, or the ER; said molecules are selected from those peptides listed in Table 9:

SEQ ID NO:	Name	Sequence
1043	TAT	RKKRRQRRR
1044	Penetratin	RQIKIWFQNRRMKWKK
1045	Transportan	GWTLSAGYLLGKINKALAALAKKIL
1046	MAP	KLALKLALKALKAALKLA
1047	Pep-1	KETWWETWWTEWSQPKKKRKV
1048	KDEL	KDEL
1049	GALA	WEAALAEALAEHLAEALAEALAEALAA
1050	HA2	GDIMGEWGNEIFGAIAGFLGC
1051	Aurine 1.2	GLFDIHKIAESF
1052	MPG	GALFLGWLGAAGSTMGAPKSKRKV
1053	TP-10	AGYLLGKINKALAALAKKIL
1054	EB-1	LIRLWSHLIHIWFQNRRLLKWKKK
1055	HA2-Penetratin	GLFGAIAGFIENGWEGMIDGRQIKIWFQNRRMKWKK
1056	Endosomolytic	FFKKLAHALHLLALLALHLAHLKKA
1057	Endosomolytic	LFEAIEGFIENGWGMIDGWYG
1058	Endosomolytic	LFEAIEGFIENGWEGMIDGWYGRKKRRQRRR
1059	Endosomolytic	IGAVLKVLTTGLPALISWIKRKRQQ
1060	ER Targeting	MKLAVTLTLVTLALSSSSASA
1061	ER Targeting	RLIEDICLPRWGCLWEDDKDEL
1062	ER Targeting	MIRTLTLLSTLVAGALSK
1063	ER Targeting	ILSSLTVTQLLRRLHQWIK
1064	ER Targeting	MIRTLTLLSTLVAGALSKDEL

[00187] In some embodiments, compositions or complexes are provided having a formula of A_1-B_1 , wherein A_1 has a formula of X_S and B_1 has a formula of $X_{AS}-L_2-F_1$.

[00188] In some embodiments, compositions or complexes are provided having a formula of A_1-B_1 , wherein A_1 has a formula of $C-L_1-X_S$ and B_1 has a formula of X_{AS} .

[00189] In some embodiments, the sense strand is a sense strand as provided for herein.

[00190] In some embodiments, the antisense strand is an antisense strand as provided for herein.

[00191] In some embodiments, the sense and antisense strand form a double stranded siRNA molecule that targets GYS1. In some embodiments, the double stranded oligonucleotide is about 21-23 nucleotides base pairs in length. In certain embodiments, C is optional.

[00192] In some embodiments, compositions or complexes are provided having a formula of A_1-B_1 , wherein A_1 has a formula of $F_1-L_1-X_S$ and B_1 has a formula of $X_{AS}-L_2-C$, wherein:

F_1 is a polypeptide comprising at least one FN3 domain;

L_1 and L_2 are each, independently, a linker;

C is a polymer, such as PEG;

X_S is a 5' to 3' oligonucleotide sense strand of a double stranded siRNA molecule;

X_{AS} is a 3' to 5' oligonucleotide antisense strand of a double stranded siRNA molecule;

wherein X_S and X_{AS} form a double stranded oligonucleotide molecule to form the composition/complex. In certain embodiments, C is optional.

In some embodiments, compositions or complexes are provided having a formula of A_1-B_1 , wherein A_1 has a formula of X_S and B_1 has a formula of $X_{AS}-L_2-C$.

In some embodiments, compositions or complexes are provided having a formula of A_1-B_1 , wherein A_1 has a formula of $F_1-L_1-X_S$ and B_1 has a formula of X_{AS} .

[00193] In some embodiments, C is a natural or synthetic polymer, consisting of long chains of branched or unbranched monomers, and/or cross-linked network of monomers in two or three dimensions. In some instances, the polymer includes a polysaccharide, lignin, rubber, or polyalkylene oxide, which can be for example, polyethylene glycol. In some instances, the at least one polymer includes, but is not limited to, alpha-, omega-dihydroxypolyethyleneglycol, biodegradable lactone -based polymer, e.g. polyacrylic acid, polylactide acid (PLA), poly(glycolic acid) (PGA), polypropylene, polystyrene, polyolefin, polyamide, polycyanoacrylate, polyimide, polyethyleneterephthalat (PET, PETG), polyethylene -B-terephthalate (PETE), polytetramethylene glycol (PTG), or polyurethane as well as mixtures thereof. As used herein, a mixture refers to the use of different polymers within the same compound as well as in reference to block copolymers. In some cases, block copolymers are

polymers wherein at least one section of a polymer is build up from monomers of another polymer. In some instances, the polymer comprises polyalkylene oxide. In some instances, the polymer comprises PEG. In some instances, the polymer comprises polyethylene imide (PEI) or hydroxy ethyl starch (HES).

[00194] In some embodiments, the polyalkylene oxide (e.g., PEG) is a polydispers or monodispers compound. In some instances, polydispers material comprises disperse distribution of different molecular weight of the material, characterized by mean weight (weight average) size and dispersity. In some instances, the monodisperse PEG comprises one size of molecules. In some embodiments, C is poly- or monodispersed polyalkylene oxide (e.g., PEG) and the indicated molecular weight represents an average of the molecular weight of the polyalkylene oxide, e.g., PEG, molecules.

[00195] In some embodiments, the molecular weight of the polyalkylene oxide (e.g., PEG) is about 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1450, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, 2500, 2600, 2700, 2800, 2900, 3000, 3250, 3350, 3500, 3750, 4000, 4250, 4500, 4600, 4750, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 10,000, 12,000, 20,000, 35,000, 40,000, 50,000, 60,000, or 100,000 Da.













[00196] In some embodiments, C is polyalkylene oxide (e.g., PEG) and has a molecular weight of about 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1450, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, 2500, 2600, 2700, 2800, 2900, 3000, 3250, 3350, 3500, 3750, 4000, 4250, 4500, 4600, 4750, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 10,000, 12,000, 20,000, 35,000, 40,000, 50,000, 60,000, or 100,000 Da. In some embodiments, C is PEG and has a molecular weight of about 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1450, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, 2500, 2600, 2700, 2800, 2900, 3000, 3250, 3350, 3500, 3750, 4000, 4250, 4500, 4600, 4750, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 10,000, 12,000, 20,000, 35,000, 40,000, 50,000, 60,000, or 100,000 Da. In some instances, the molecular weight of C is about 200 Da. In some instances, the molecular weight of C is about 300 Da. In some instances, the molecular weight of C is about 400 Da. In some instances, the molecular weight of C is about 500 Da. In some instances, the molecular weight of C is about 600 Da. In some instances, the molecular weight of C is about 700 Da. In some instances, the molecular weight of C is about 800 Da. In some instances, the molecular weight of C is about 900 Da. In some instances, the molecular weight of C is about 1000 Da. In some instances, the molecular weight of C is about 1100 Da. In some instances, the molecular weight of C is about 1200 Da. In some instances, the molecular weight of C is about 1300 Da. In some instances, the molecular weight of C is about 1400 Da. In

some instances, the molecular weight of C is about 1450 Da. In some instances, the molecular weight of C is about 1500 Da. In some instances, the molecular weight of C is about 1600 Da. In some instances, the molecular weight of C is about 1700 Da. In some instances, the molecular weight of C is about 1800 Da. In some instances, the molecular weight of C is about 1900 Da. In some instances, the molecular weight of C is about 2000 Da. In some instances, the molecular weight of C is about 2100 Da. In some instances, the molecular weight of C is about 2200 Da. In some instances, the molecular weight of C is about 2300 Da. In some instances, the molecular weight of C is about 2400 Da. In some instances, the molecular weight of C is about 2500 Da. In some instances, the molecular weight of C is about 2600 Da. In some instances, the molecular weight of C is about 2700 Da. In some instances, the molecular weight of C is about 2800 Da. In some instances, the molecular weight of C is about 2900 Da. In some instances, the molecular weight of C is about 3000 Da. In some instances, the molecular weight of C is about 3250 Da. In some instances, the molecular weight of C is about 3350 Da. In some instances, the molecular weight of C is about 3500 Da. In some instances, the molecular weight of C is about 3750 Da. In some instances, the molecular weight of C is about 4000 Da. In some instances, the molecular weight of C is about 4250 Da. In some instances, the molecular weight of C is about 4500 Da. In some instances, the molecular weight of C is about 4600 Da. In some instances, the molecular weight of C is about 4750 Da. In some instances, the molecular weight of C is about 5000 Da. In some instances, the molecular weight of C is about 5500 Da. In some instances, the molecular weight of C is about 6000 Da. In some instances, the molecular weight of C is about 6500 Da. In some instances, the molecular weight of C is about 7000 Da. In some instances, the molecular weight of C is about 7500 Da. In some instances, the molecular weight of C is about 8000 Da. In some instances, the molecular weight of C is about 10,000 Da. In some instances, the molecular weight of C is about 12,000 Da. In some instances, the molecular weight of C is about 20,000 Da. In some instances, the molecular weight of C is about 35,000 Da. In some instances, the molecular weight of C is about 40,000 Da. In some instances, the molecular weight of C is about 50,000 Da. In some instances, the molecular weight of C is about 60,000 Da. In some instances, the molecular weight of C is about 100,000 Da.

[00197] In some embodiments, the polyalkylene oxide (e.g., PEG) is a discrete PEG, in which the discrete PEG is a polymeric PEG comprising more than one repeating ethylene oxide units. In some instances, a discrete PEG (dPEG) comprises from 2 to 60, from 2 to 50, or from 2 to 48 repeating ethylene oxide units. In some instances, a dPEG comprises about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 24, 26, 28, 30, 35, 40, 42, 48, 50 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 2 or more repeating

ethylene oxide units. In some instances, a dPEG comprises about 3 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 4 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 5 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 6 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 7 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 8 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 9 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 10 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 11 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 12 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 13 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 14 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 15 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 16 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 17 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 18 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 19 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 20 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 22 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 24 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 26 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 28 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 30 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 35 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 40 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 42 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 48 or more repeating ethylene oxide units. In some instances, a dPEG comprises about 50 or more repeating ethylene oxide units. In some cases, a dPEG is synthesized as a single molecular weight compound from pure (e.g., about 95%, 98%, 99%, or 99.5%) starting material in a step-wise fashion. In some cases, a dPEG has a specific molecular weight, rather than an average molecular weight. In some cases, a dPEG described herein is a dPEG from Quanta Biodesign, LMD.

[00198] In some embodiments, L_1 is any linker that can be used to link the polymer C to the sense strand X_S or to link the polypeptide of F_1 to the sense strand X_S . In some embodiments, L_1 has a formula of:









wherein X_S, X_{AS}, and F₁ are as defined above.

[00199] In some embodiments, n = 0-20. In some embodiments, R and R₁ are independently methyl. In some embodiments, R and R₁ are independently present or both are absent. In some embodiments, X and Y are independently S. In some embodiments, X and Y are independently present or absent. In some embodiments, Peptide is an enzymatically cleavable peptide, such as, but not limited to, Val-Cit, Val-Ala etc.

[00200] In some embodiments, L₂ is any linker that can be used to link the polypeptide of F₁ to the antisense strand X_{AS} or to link the polymer C to the antisense strand X_{AS}.







[00201] In some embodiments, L₂ has a formula of in the complex of:




wherein X_{AS} and F_1 are as defined above.

[00202] In some embodiments, $n = 0-20$. In some embodiments, R and R_1 are independently methyl. In some embodiments, R and R_1 are independently present or both are absent. In some embodiments, X and Y are independently S . In some embodiments, X and Y are independently present or absent. In some embodiments, Peptide is an enzymatically cleavable peptide, such as, but not limited to, Val-Cit, Val-Ala etc.

[00203] In some embodiments, the linker is covalently attached to F_1 through a cysteine residue present on F_1 , which can be illustrated as follows:



In some embodiments, A1-B1 has a formula of:



wherein C is the polymer, such as PEG, Endoport, INF-7, TAT, polyarginine, polylysine, an amphipathic peptide, or peptides listed in Table 9 as provided for herein, X_S is a 5' to 3' oligonucleotide sense strand of a double stranded siRNA molecule; X_{AS} is a 3' to 5' oligonucleotide antisense strand of a double stranded siRNA molecule; and F_1 is a polypeptide comprising at least one FN3 domain, wherein X_S and X_{AS} form a double stranded siRNA molecule. The sense and antisense strands are represented by the "N" notations, wherein each nucleotide represented by N, is independently, A, U, C, or G or a modified nucleobase, such as those provided for herein. The N_1 nucleotides of the sense strand and the antisense strand represent the 5' end of the respective strands. For clarity, although Formula I utilizes N_1 , N_2 , N_3 , etc. in both the sense and the antisense strand, the nucleotide bases do not need to be the same and are not intended to be the same. The siRNA that is illustrated in Formula I would be complementary to a target sequence.

[00204] For example, in some embodiments, the sense strand comprises a 2'-O-methyl modified nucleotide with a phosphorothioate (PS) modified backbone at N_1 and N_2 , a 2'-fluoro modified nucleotide at N_3 , N_7 , N_8 , N_9 , N_{12} , and N_{17} , and a 2'-O-methyl modified nucleotide at N_4 , N_5 , N_6 , N_{10} , N_{11} , N_{13} , N_{14} , N_{15} , N_{16} , N_{18} , and N_{19} .

[00205] In some embodiments, the antisense strand comprises a vinylphosphonate moiety attached to N_1 , a 2'-fluoro- modified nucleotide with a phosphorothioate (PS) modified backbone at N_2 , a 2'-O-methyl modified nucleotide at N_3 , N_4 , N_5 , N_6 , N_7 , N_8 , N_9 , N_{10} , N_{11} , N_{12} , N_{13} , N_{15} , N_{16} , N_{17} , N_{18} , and N_{19} , a 2'-fluoro- modified nucleotide at N_{14} , and a 2'-O-methyl modified nucleotide with a phosphorothioate (PS) modified backbone at N_{20} and N_{21} .

[00206] In some embodiments, a compound having a formula of:

[00207] In some embodiments, F₁ comprises polypeptide having a formula of (X₁)_n-(X₂)_q-(X₃)_y, wherein X₁ is a first FN3 domain; X₂ is second FN3 domain; X₃ is a third FN3 domain or half-life extender molecule; wherein n, q, and y are each independently 0 or 1, provided that at least one of n, q, and y is 1. In some embodiments, n, q, and y are each 1. In some embodiments, n and q are 1 and y is 0. In some embodiments n and y are 1 and q is 0.

[00209] In some embodiments, compositions are provided herein having a formula of (X1)_n-(X2)_q-(X3)_y-L-X4-C, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; X4 is a nucleic acid molecule; and C is a polymer, wherein n, q, and y are each independently 0 or 1.

[00210] In some embodiments, compositions are provided herein having a formula of X4-L-(X1)_n-(X2)_q-(X3)_y, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; and X4 is a nucleic acid molecule, wherein n, q, and y are each independently 0 or 1.

[00211] In some embodiments, compositions are provided herein having a formula of C-X4-L-(X1)_n-(X2)_q-(X3)_y, wherein C is a polymer; X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; and X4 is a nucleic acid molecule, wherein n, q, and y are each independently 0 or 1.

[00212] In some embodiments, compositions are provided herein having a formula of X4-L-(X1)_n-(X2)_q-(X3)_y-C, wherein X1 is a first FN3 domain; X2 is second FN3 domain; X3 is a third FN3 domain or half-life extender molecule; L is a linker; X4 is a nucleic acid molecule; and C is a polymer, wherein n, q, and y are each independently 0 or 1.

[00213] In some embodiments, the GYS1 siRNA pair may follow the sequence: sense strand (5'-3') nsnsnnnnNfNfNfnnnnnnnsnsa and antisense strand (5'-3') UfsNfsnnnNfnnnnnnNfNfNfnnnsusu, wherein (n) is 2'-O-Me (methyl), (Nf) is 2'-F (fluoro), (s) is phosphorothioate backbone modification. Each nucleotide in both sense and antisense strands are modified independently or in combination at ribosugar and nucleobase positions.

[00214] In some embodiments, the siRNA molecule comprises a sequence pair from Tables 1A or 1B.

Table 1A: siRNA Sense and Anti-sense sequences

siRNA Pair	SEQ ID NO:	Sense Strand 5'-3'	SEQ ID NO:	Antisense Strand 5'-3'
A	10	[fC][*mU][*fG][mG][fG][mA][fG][fG][fA][mU][fG][mA][mA][mU][fU][mC][fG][mA][mA][T]	11	[mU][*fU][*mC][fG][mA][fA][fU][fU][mC][fA][mU][mC][mC][fU][mC][fC][mC][fA][mG][*fU][*mU]
B	12	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mG][mC][mA][fU][mC][fU][mA][mA][T]	13	[mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]
C	14	[fU][*mC][*fC][mG][fC][mA][fG][fC][fC][mU][fG][mG][mA][mU][fG][mA][fU][mU][mA][T]	15	[mU][*fA][*mA][fU][mC][fA][fU][fC][mC][fA][mG][mG][mC][fU][mG][fC][mG][fG][mA][*fU][*mU]
D	16	[mG][*mC][*fG][mA][mC][mU][fG][fC][fC][mU][mG][fU][mA][mG][mC][mA][fA][mC][mA][idT]	17	[fU][*fG][*mU][mU][mG][mC][mU][mA][mC][mA][mG][mG][mC][fA][mG][mU][mC][mG][mC][*mU][*mU]
E	18	[mU][*mU][*fU][mA][mU][mG][fG][fG][fC][mA][mU][fC][mU][m	19	[fU][*fA][*mG][mU][mC][mC][mA][mG][mA][mU][mG][mC][mC][fC][mA][mU][mA][mA][mA][*mU][*m

		G][mG][mA][fC][mU][mA][idT]		U]
F	20	[mG][*mC][*fG][mC][mG][mG][fA][fC][fC][mA][mA][fC][mA][mA][mU][mU][fU][mC][mA][idT]	21	[fU][*fG][*mA][mA][mA][mU][mU][mG][mU][mG][mG][mU][fC][mC][mG][mC][mG][mC][*mU][*mU]
G	22	[mG][*mG][*fA][mC][mC][mA][fA][fC][fA][mA][mU][fU][mU][mC][mA][mA][fC][mG][mA][idT]	23	[fU][*fC][*mG][mU][mU][mG][mA][mA][mA][mU][mU][mG][mU][fU][mG][mG][mU][mC][mC][*mU][*mU]
H	24	[mA][*mC][*fC][mA][mA][mC][fA][fA][fU][mU][mU][fC][mA][mA][mC][mG][fU][mG][mA][idT]	25	[fU][*fC][*mA][mC][mG][mU][mU][mG][mA][mA][mA][mU][mU][fG][mU][mU][mG][mG][mU][*mU][*mU]
I	26	[mG][*mC][*fG][mC][mA][mA][fA][fC][fA][mG][mC][fU][mU][mU][mG][mG][fG][mA][mA][idT]	27	[fU][*fU][*mC][mC][mC][mA][mA][mA][mG][mC][mU][mG][mU][fU][mU][mG][mC][mG][mC][*mU][*mU]
J	28	[mA][*mG][*fA][mG][mC][mC][fA][fU][fC][mU][mU][fU][mG][mC][mA][mA][fC][mG][mA][idT]	29	[fU][*fC][*mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][mU][fG][mG][mC][mU][mC][mU][*mU][*mU]
K	30	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mC][fG][mC][mA][idT]	31	[fU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]
L	32	[mA][*mG][*fC][mC][mA][mU][fC][fU][fU][mU][mG][fC][mA][mA][mC][mG][fC][mA][mA][idT]	33	[fU][*fU][*mG][mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][fA][mU][mG][mG][mC][mU][*mU][*mU]
M	34	[mC][*mC][*fA][mU][mC][mU][fU][fU][fG][mC][mA][fA][mC][mG][mC][mA][fG][mC][mA][idT]	35	[fU][*fG][*mC][mU][mG][mC][mG][mU][mU][mG][mC][mA][mA][fA][mG][mA][mU][mG][mG][*mU][*mU]
N	36	[mA][*mC][*fG][mC][mA][mG][fC][fG][fG][mC][mA][fG][mU][mC][mU][mU][fU][mC][mA][idT]	37	[fU][*fG][*mA][mA][mA][mG][mA][mC][mU][mG][mC][mC][mG][fC][mU][mG][mC][mG][mU][*mU][*mU]
O	38	[mU][*mG][*fA][mC][mC][mA][fC][fC][fA][mU][mC][fC][mG][mC][mC][mG][fA][mA][mA][idT]	39	[fU][*fU][*mU][mC][mG][mG][mC][mG][mG][mA][mU][mG][mG][fU][mG][mG][mU][mC][mA][*mU][*mU]
P	40	[mC][*mG][*fA][mA][mU][mC][fG][fG][fC][mC][mU][fC][mU][mU][mC][mA][fA][mU][mA][idT]	41	[fU][*fA][*mU][mU][mG][mA][mA][mG][mA][mG][mC][mC][fG][mA][mU][mU][mC][mG][*mU][*mU]
Q	42	[mC][*mU][*fU][mC][mA][mA][fU][fA][fG][mC][mA][fG][mU][mG][mC][mC][fG][mA][mA][idT]	43	[fU][*fU][*mC][mG][mG][mC][mA][mC][mU][mG][mC][mU][mA][fU][mU][mG][mA][mA][mG][*mU][*mU]

				U]
R	44	[mC][*mA][*fG][mU][mA][mU][fC][fU][fC][mC][mA][fC][mC][mA][mA][mU][fC][mU][mA][idT]	45	[fU][*fA][*mG][mA][mU][mU][mG][mG][mU][mG][mG][mA][mG][fA][mU][mA][mC][mU][mG][*mU][*mU]
S	46	[mC][*mC][*fA][mC][mC][mA][fA][fU][fC][mU][mC][fU][mC][mC][mG][mG][fC][mU][mA][idT]	47	[fU][*fA][*mG][mC][mC][mG][mG][mA][mG][mA][mG][mA][fU][mG][mG][mU][mG][mG][*mU][*mU]
T	48	[mC][*mA][*fC][mC][mA][mA][fU][fC][fU][mC][mU][fC][mC][mG][mG][mC][fU][mU][mA][idT]	49	[fU][*fA][*mA][mG][mC][mC][mG][mG][mA][mG][mA][mG][mA][fU][mU][mG][mG][mU][mG][*mU][*mU]
U	50	[mC][*mC][*fC][mC][mU][mC][fA][fG][fC][mU][mU][fA][mC][mG][mG][mU][fA][mU][mA][idT]	51	[fU][*fA][*mU][mA][mC][mC][mG][mU][mA][mA][mG][mC][mU][fG][mA][mG][mG][mG][mG][*mU][*mU]
V	52	[mC][*mC][*fC][mU][mC][mA][fG][fC][fU][mU][mA][fC][mG][mG][mU][mA][fU][mC][mA][idT]	53	[fU][*fG][*mA][mU][mA][mC][mC][mG][mU][mA][mA][mG][mC][fU][mG][mA][mG][mG][mG][*mU][*mU]
W	54	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fC][mU][mA][idT]	55	[fU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]
X	56	[mG][*mG][*fA][mA][mC][mC][fG][fC][fA][mC][mG][fG][mA][mG][mC][mG][fC][mC][mA][idT]	57	[fU][*fG][*mG][mC][mG][mC][mU][mC][mC][mG][mU][mG][mC][fG][mG][mU][mU][mC][mC][*mU][*mU]
Y	58	[mA][*mC][*fG][mG][mA][mG][fC][fG][fC][mC][mU][fC][mU][mC][mC][mG][fA][mC][mA][idT]	59	[fU][*fG][*mU][mC][mG][mG][mA][mG][mA][mG][mC][mC][mC][fC][mU][mC][mC][mG][mU][*mU][*mU]
Z	60	[mA][*mG][*fC][mA][mA][mG][fC][fG][fC][mA][mA][fC][mU][mC][mU][mG][fU][mG][mA][idT]	61	[fU][*fC][*mA][mC][mA][mG][mA][mG][mU][mU][mG][mC][mG][fC][mU][mU][mG][mC][mU][*mU][*mU]
AA	62	[mC][*mA][*fG][mA][mU][mG][fG][fU][fC][mC][mU][fC][mC][mA][mU][mU][fU][mC][mA][idT]	63	[fU][*fG][*mA][mA][mA][mU][mG][mG][mA][mG][mG][mA][mC][fC][mA][mU][mC][mU][mG][*mU][*mU]
BB	64	[mU][*mC][*fU][mA][mG][mA][fU][fC][fU][mG][mG][fA][mA][mC][mC][mU][fU][mA][mA][idT]	65	[fU][*fU][*mA][mA][mG][mG][mU][mU][mC][mC][mA][mG][mA][fU][mC][mU][mA][mG][mA][*mU][*mU]
CC	66	[mC][*mC][*fC][mA][mC][mA][fU][fU][fU][mA][mC][fC][mU][mC][mG][mA][fG][mG][mA][idT]	67	[fU][*fC][*mC][mU][mC][mG][mA][mG][mG][mU][mA][mA][mA][fU][mG][mU][mG][mG][mG][*mU][*mU]

				U]
DD	68	[mA][*mC][*fC][mG][mC][mU][fU][fG][fG][mU][mU][fU][mG][mC][mA][mG][fU][mU][mA][idT]	69	[fU][*fA][*mA][mC][mU][mG][mC][mA][mA][mA][mC][mC][mA][fA][mG][mC][mG][mG][mU][*mU][*mU]
EE	600	[fC][*mU][*fG][mG][fG][mA][fG][fG][fA][mU][fG][mA][mA][mU][fU][mC][fG][mG][mA][T]	601	[mU][*fU][*mC][fG][mA][fA][fU][fU][mC][fA][mU][mC][mC][fU][mC][fC][mC][fA][mG][*fU][*mU]
FF	602	[fC][*mU][*fG][mG][fG][mA][fG][fG][fA][mU][fG][mA][mA][mU][fU][mA][fG][mG][mA][T]	603	[mU][*fU][*mC][fG][mA][fA][fU][fU][mC][fA][mU][mC][mC][fU][mC][fC][mC][fA][mG][*fU][*mU]
GG	604	[fC][*mU][*fG][mG][fG][mA][fG][fG][fA][mU][fG][mA][mA][mU][fG][mA][fG][mG][mA][T]	605	[mU][*fU][*mC][fG][mA][fA][fU][fU][mC][fA][mU][mC][mC][fU][mC][fC][mC][fA][mG][*fU][*mU]
HH	606	[fC][*mU][*fG][mG][fG][mA][fG][fG][fA][mU][fG][mA][mA][mU][fU][mC][fG][mA][mA][T]	607	[fU][*fU][*fC][fG][fA][fA][fU][fU][fC][fA][fU][fC][fC][fU][fC][fC][fC][fA][fG][*fU][*fU]
II	608	[fC][*mU][*fG][mG][fG][mA][fG][fG][fA][mU][fG][mA][mA][mU][fU][mC][fG][mA][mA][T]	609	[mU][*fU][*fC][fG][fA][fA][fU][fU][mC][fA][mU][fC][fC][fU][fC][fC][fC][fA][fG][*fU][*mU]
JJ	610	[fC][*mU][*fG][mG][fG][mA][fG][fG][fA][mU][fG][mA][mA][*fU][idT]	611	[mU][*fU][*mC][fG][mA][fA][fU][fU][mC][fA][mU][mC][mC][fU][mC][fC][mC][fA][*mG][*fU][*mU]
KK	612	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mC][mA][fU][mC][fU][mA][mA][idT]	613	[mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]
LL	614	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mC][mA][fU][mC][fU][mA][mG][idT]	615	[mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]
MM	616	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mC][mA][fU][mC][fU][mG][mA][idT]	617	[mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]
NN	618	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mC][mA][fU][mC][fG][mA][mA][idT]	619	[mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]
OO	620	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mC][mA][fU][mA][fU][mA][mA][idT]	621	[mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]
PP	622	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mC][mA][fG][mC][fU][mA][mA][idT]	623	[mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]
QQ	624	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mC][mA][fG][mC][fU][mA][mG][idT]	625	[mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]
RR	626	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][m	627	[mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*m

		A][mA][mC][fG][mC][mA][idT]		U]
SS	628	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mC][fG][mC][mG][idT]	629	[mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]
TT	630	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mC][fG][mA][mA][idT]	631	[mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]
UU	632	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mC][fA][mC][mA][idT]	633	[mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]
VV	634	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mA][fG][mC][mA][idT]	635	[mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]
WW	636	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mG][mC][fG][mC][mA][idT]	637	[mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]
XX	638	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mG][mC][fG][mC][mG][idT]	639	[mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]
YY	640	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fC][mU][mA][idT]	641	[mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]
ZZ	642	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fC][mU][mG][idT]	643	[mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]
AAA	644	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fC][mG][mA][idT]	645	[mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]
BBB	646	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fG][mU][mA][idT]	647	[mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]
CCC	648	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mG][fC][mU][mA][idT]	649	[mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]
DDD	650	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mG][fC][mU][mA][idT]	651	[mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]

				mU]
EEE	652	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mG][mU][fC][mU][mA][idT]	653	[mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]
FFF	654	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mG][mU][fC][mU][mG][idT]	655	[mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]
GGG	656	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mG][mC][mA][fU][mC][fU][mA][mG][idT]	657	[mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]
HHH	704	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fC][mU][mA]	705	[mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]

[00215] Abbreviations Key: (n/N= any nucleotide) mN= 2'-O-methyl residues, fN= 2'-F residues, * = phosphorothioate and (idT)= inverted Dt, (VP) 2'-O methyl vinyl phosphonate uridine. The brackets indicate the individual bases.

Table 1B

SEQ ID NO:	Sense strand (5'-3')	SEQ ID NO:	Antisense strand (5'-3')
801	caggggugcg gucuugcaa	861	uugcaagacc gcacccugu u
802	ggugcgguu ugc aaauagg	862	ccuauugcaa gaccgcaccu u
803	gugcggguu gca auagga	863	uccuauugca agaccgcacu u
804	agccaugccu uuaaacgc	864	gcgguuuaa ggcauggcuu u
805	augccuuua accgcacuu	865	aagugcgguu uaaaggcau u
806	ugggaggaug aaau cgacc	866	ggugcaauu auccuccau u
807	aacgcagugc ucuucgaag	867	cuucgaagag cacugcguu u
808	ggacgaugg ggcgacaac	868	guugcgccc cauucguccu u
809	ggcgacugcc uguagcaac	869	guugcuacag gcagugccu u
810	cacgcugcug gggcgcuac	870	guagcgccc agcagcgugu u
811	ugcgucacg ucuucacua	871	uagugaagac gugagcgcau u
812	ccuauacuu cuuuaucgc	872	gcgaauaaga aguauaaggu u
813	cuuuaucgcc ggccgcua u	873	auagcgccg gcgaauaagu u
814	ucgccggccg cuagaguu	874	aacucauagc ggccggcgau u
815	gccgcuauga guucuccaa	875	uuggagaacu cauagcggu u

816	ccgcuaugag uucuccaac	876	guuggagaac ucauagcgg u
817	ggcucggcuc aacuaucug	877	cagauaguug agccgagccu u
818	gcucggcuca acuaucugc	878	gcagauagu u gagccgagcu u
819	ucggcucaac uaucugcuc	879	gagcagauag uugagccgau u
820	ccagcgcgga ccaacaauu	880	aauguuuggu ccgcgcuggu u
821	agcgcggacc aacaauuuc	881	gaaauguug guccgcgcuu u
822	ucgggaggaa gcuuuuga	882	ucauaagcu uccucccgau u
823	acgcagcggc agucuuucc	883	ggaaagacug ccgcugcguu u
824	gaccaccauc cgccgauc	884	gauucgcggc augguggucu u
825	ccgaucggc cucuucuu	885	auugaaggc ccgaucgg u
826	aaucggccuc uucauagc	886	gcuauugaag aggccgauu u
827	caauagcagu gccgacagg	887	ccugucggc cugcuauugu u
828	aggaguuugu ccguggcug	888	cagccacgga caaacuccu u
829	cggcugagug cacgguuau	889	auaacgugc acucagccgu u
830	ugggaauccc caguauuc	890	gagauacugg ggauuccau u
831	cagacccuc agcuuacgg	891	ccguaagcug aggggucugu u
832	ccccucagu uacgguauc	892	gauaccguu gcuagggg u
833	cucagcuuac gguauuac	893	guagauaccg uaagcugagu u
834	gcuuacggu ucuacauuc	894	gaauguagau accguuagcu u
835	ccggcggcag cguauauc	895	gaugauacgc ugccgccggu u
836	agcguaucau ccagcggaa	896	uuccgcugga ugauacgcu u
837	gcguauauc cagcggaa	897	guuccgcug augauacgu u
838	uaccuaggcc gguacuaua	898	uauaguaccg gccuagguu u
839	cuaggccggu acuaauugu	899	acauauagua ccggccuagu u
840	aggccgguac uauaugucu	900	agacauauag uaccggccu u
841	ggccgguacu auaugucug	901	cagacauaua guaccggccu u
842	gccgguacua uaugucugc	902	gcagacauau aguaccggcu u
843	cgguaucua uugucgcgc	903	gcgcagacau auaguaccgu u
844	gguacuauau gucugcgcg	904	cgcgagaca uauaguaccu u
845	cuauaugucu gcgcgccac	905	guggcgcgca gacauauagu u
846	uauugucgcg gccacaug	906	caugugcgc gcagacauu u
847	augucgcgc gccacaugg	907	ccaugugcgc gcagacauu u

848	ucgcccucgc ugucacgac	908	gucgugacag cgaggcgcau u
849	cggcgagcgc uacgaugag	909	cucaucguag cgcucgccgu u
850	caaggaccgg cgcaacauc	910	gauguugcgc cgguccuugu u
851	acauccgugc accagagug	911	cacucuggug cacggauguu u
852	cuggcgagg agcguaacu	912	aguuacgcuc cugccccagu u
853	gggcgaggag cguaacuaa	913	uuaguuaacgc uccucgccc u
854	gcgaggagcg uaacuaagu	914	acuuaguuaac gcuccucgcu u
855	gaggagcgua acuaagucc	915	ggacuuaugu acgcuccucu u
856	aguccgcaa acacuccac	916	guggaguguu uggcggaucu u
857	ggcgaucaag uccagagcc	917	ggcucuggac uugaucgccu u
858	cccuaacug gcuaauucc	918	ggaauaagcc agguuaggg u
859	ugugaaacca cuagguucu	919	agaaccuagu gguuucacau u
860	accacuaggu ucuaggucc	920	ggaccuagaa ccuagugguu u
921	caggggugcg gucuugcaa	981	uugcaagacc gcacccugu u
922	ggugcgguu ugcauaga	982	ucuauugcaa gaccgcaccu u
923	gugcgguuu gcaauagga	983	uccuauugca agaccgcacu u
924	agccaugccu uuaaacga	984	ucgguuuaa ggcaugccu u
925	augccuuuaa accgcacua	985	uagugcgguu uaaaggcau u
926	ugggaggaug aaucgaca	986	ugucgaauuc auccuccau u
927	aacgcagugc ucuucgaaa	987	uuucgaagag cacugcguuu u
928	ggacgaau ggcgacaaa	988	uuugcgccc cauucgucu u
929	ggcgacugcc uguagcaaa	989	uuugcuacag gcagucgccu u
930	cacgcugcug gggcgcuua	990	uuagcgccc agcagcgugu u
931	ugcgucacg ucuucacua	991	uagugaagac gugagcgau u
932	ccuuauacuu cuuuauca	992	ucgauaaaga aguauaaggu u
933	cuuuauccg gccgcuaa	993	uuagcgccg gcgaauaagu u
934	ucgccggccg cuaugagua	994	uacucauagc ggccggcgau u
935	gccgcuauga guucuccaa	995	uuggagaacu cauagcggu u
936	ccgcuaugag uucuccaaa	996	uuuggagaac ucauagcggu u
937	ggcucggcuc aacuaucua	997	uagauaguug agccgagccu u
938	gcucggcuca acuaucuga	998	ucagauagu gagccgagcu u
939	ucggcucaac uaucugcua	999	uagcagauag uugagccgau u

940	ccagcgcgga ccaacaaua	1000	uauuguuggu ccgcgcuggu u
941	agcgcggacc aacaauua	1001	uaauuguug guccgcgcuu u
942	ucgggaggaa gcuuuaua	1002	ucauaaagcu uccucccgau u
943	acgcagcggc agucuuuca	1003	ugaaagacug ccgcgcgcuu u
944	gaccaccauc cgccgaaua	1004	uauucggcgg augguggucu u
945	ccgaaucggc cucuucaaa	1005	uuugaagagg ccgauucggu u
946	aaucggccuc uucaauaga	1006	ucuauugaag aggccgauuu u
947	caauagcagu gccgacaga	1007	ucugucggca cugcuauugu u
948	aggaguugu ccguggcua	1008	uagccacgga caaacuccu u
949	cggcugagug cacgguaa	1009	uuaaccgugc acucagccgu u
950	ugggaauc cagauucua	1010	uagauacugg ggauuccau u
951	cagaccucc agcuuacga	1011	ucguaagcug aggggucugu u
952	ccccucagcu uacgguaa	1012	uauaccguaa gcugaggggu u
953	cucagcuuac gguaucuaa	1013	uuagauaccg uaagcugagu u
954	gcuuacggua ucuacaua	1014	uaauguagau accguaagcu u
955	ccggcggcag cguaucua	1015	uauauacgc ugccgccggu u
956	agcguauc cagcggaa	1016	uuccgcugga ugauacgcu u
957	gcguauauc cagcggaaa	1017	uuuccgcugg augauacgu u
958	uaccuaggcc gguacuua	1018	uauaguaccg gccuagguau u
959	cuaggccggu acuaauaga	1019	ucauauagua ccggccuagu u
960	aggccgguac uauauguca	1020	ugacauauag uaccggccu u
961	ggccgguacu auaugucua	1021	uagacauua guaccggccu u
962	gccgguacua uaugucuga	1022	ucagacauau aguaccggcu u
963	cgguaucua ugucugcga	1023	ucgcagacau auaguaccgu u
964	gguacuauau gucugcgca	1024	ugcgagaca uauaguaccu u
965	cuauaugucu gcgcgccaa	1025	uuggcgcgca gacauauagu u
966	uauugucgc gccacaua	1026	uauuggcgc gcagacauu u
967	augucugcg gccacaua	1027	ucaugugcg cgagacauu u
968	ucgcccucgc ugucacgaa	1028	uucgugacag cgaggcgau u
969	cggcgagcg uacgaugaa	1029	uucaucguag cgcucgccgu u
970	caaggaccgg cgcaacua	1030	uauugcgc cgguccuugu u
971	acauccgugc accagagua	1031	uacucuggug cacggauguu u

972	cugggcgagg agcguaaca	1032	uguuacgcuc cucgccagu u
973	gggcgaggag cguaacuaa	1033	uuaguuacgc uccucgccu u
974	gcgaggagcg uaacuaaga	1034	ucuuaguac gcuccucgu u
975	gaggagcgua acuaaguca	1035	ugacuuaguu acgcuccu u
976	aguccgccaa acacuccaa	1036	uuggaguguu uggcggacu u
977	ggcgaucaag uccagagca	1037	ugcucuggac uugaucgccu u
978	cccuaccug gcuuuuca	1038	ugaauagcc agguuaggu u
979	ugugaaacca cuagguuca	1039	ugaaccuagu gguuacau u
980	accacuaggu ucuagguca	1040	ugaccuagaa ccuaguggu u
1041	uggacuuaaccuagacaa	1042	uugucuagguugaaguccau
1043	cugggaggaugaauucgaa	1044	uucgaaaucauccuccagu
1045	gggugacaacuacuaucua	1046	uagauaguaguugucacccu
1047	cugggaggaugaauucgaa	1048	uucgaaaucauccuccagu
1049	caagguggguggcaucuaa	1050	uuagaugccaccaccuugu
1051	uccgcagccuggaugauua	1052	uaaucauccaggcugcggau

[00216] In some embodiments, the polynucleotides illustrated above include those that do not include a 2'-O methyl vinyl phosphonate uridine as the 5' nucleotide on the antisense strand of the siRNA.

[00217] In some embodiments, a polynucleotide is as provided for herein. In some embodiments, the polynucleotide comprises a first strand and a second strand to form a portion that comprises a duplex. In some embodiments, the polynucleotide comprises a sense strand and an antisense strand. In some embodiments, comprises the sequences as illustrated in Tables 1A or 1B. In some embodiments, comprises the sequences as illustrated in Tables 1A or 1B but without the base modifications. In some embodiments, a pharmaceutical composition comprises a siRNA pair as provided herein. In some embodiments, the siRNA pair is not conjugated to a FN3 domain.

[00218] In some embodiments, an oligonucleotide molecule described herein is constructed using chemical synthesis and/or enzymatic ligation reactions using procedures known in the art. For example, an oligonucleotide molecule is chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed

between the oligonucleotide molecule and target nucleic acids. Alternatively, the oligonucleotide molecule is produced biologically using an expression vector into which a oligonucleotide molecule has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted oligonucleotide molecule will be of an antisense orientation to a target polynucleic acid molecule of interest).

[00219] In some embodiments, an oligonucleotide molecule is synthesized via a tandem synthesis methodology, wherein both strands are synthesized as a single contiguous oligonucleotide fragment or strand separated by a cleavable linker which is subsequently cleaved to provide separate fragments or strands that hybridize and permit purification of the duplex.

[00220] In some instances, an oligonucleotide molecule is also assembled from two distinct nucleic acid strands or fragments wherein one fragment includes the sense region and the second fragment includes the antisense region of the molecule.

[00221] In some instances, while chemical modification of the oligonucleotide molecule internucleotide linkages with phosphorothioate, phosphorodithioate, phosphonate, phosphoramidate, or mesyl phosphoramidate, linkages improves stability. Excessive modifications sometimes cause toxicity or decreased activity. Therefore, when designing nucleic acid molecules, the amount of these internucleotide linkages in some cases is minimized. In such cases, the reduction in the concentration of these linkages lowers toxicity, increases efficacy and higher specificity of these molecules.

[00222] As described herein, in some embodiments, the nucleic acid molecules can be modified to include a linker at the 5' end of the of the sense strand of the dsRNA. In some embodiments, the nucleic acid molecules can be modified to include a vinyl phosphonate or modified vinyl phosphonate at the 5' end of the of the anti-sense strand of the dsRNA. In some embodiments, the nucleic acid molecules can be modified to include a linker at the 3' end of the of the sense strand of the dsRNA. In some embodiments, the nucleic acid molecules can be modified to include a vinyl phosphonate at the 3' end of the of the anti-sense strand of the dsRNA. The linker can be used to link the dsRNA to the FN3 domain. The linker can covalently attach, for example, to a cysteine residue on the FN3 domain that is there naturally or that has been substituted as described herein, and for example, in U.S. Patent No. 10,196,446, which is hereby incorporated by reference in its entirety. Non-limiting examples of such modified strands of the dsRNA are illustrated in Table 2.

Table 2: Pairs with Linker and/or vinyl phosphonate

siRNA Pair	SEQ ID NO:	Sense Strand 5'-3'	SEQ ID NO:	Antisense Strand 5'-3'	L=linker
III	70	[mU][*mC][*fU][mC][fG][mU][fG][mG][fC][mC][fU][mU][fA][mA][fU][mG][fA][mA]-L	71	[VP][*fU][*mU][fC][mA][fU][mU][fA][mA][fG][mG][fC][mC][fA][mC][fG][mA][fG][mA][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
JJJ	72	[fC][*mU][*fG][mG][fG][mA][fG][fG][fA][mU][fG][mA][mA][mU][fU][mC][fG][mA][mA]-L	73	[VP][*fU][*mC][mG][mA][mA][mU][mU][mC][mA][mU][mC][mC][fU][mC][mC][mC][mA][mG][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
KKK	74	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mG][mC][mA][fU][mC][fU][mA][mA]-L	75	[VP][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
LLL	76	[mG][*mC][*fG][mC][mG][mG][fA][fC][fC][mA][mA][fC][mA][mA][mU][mU][fU][mC][mA]-L	77	[VP][*fG][*mA][mA][mU][mU][mG][mU][mU][mG][mG][mU][fC][mC][mG][mC][mG][mC][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
MMM	78	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mC][fG][mC][mA]-L	79	[VP][*fG][*mC][mG][mU][mU][mG][mC][mC][mA][mA][mG][mG][mU][fU][mG][mG][mC][mU][mC][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
NNN	80	[mA][*mG][*fC][mC][mA][mU][fC][fU][fU][mU][mG][fC][mA][mA][mC][mG][fC][mA][mA]-L	81	[VP][*fU][*mG][mC][mG][mU][mU][mG][mC][mC][mA][mA][mG][mU][fA][mU][mG][mG][mC][mU][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
OOO	82	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fC][mU][mA]-L	83	[VP][*fA][*mG][mA][mU][mU][mG][mU][mU][mG][mA][mU][mU][mA][fC][mC][mG][mU][mU][mU][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
OPPP	84	[mC][*mC][*fC][mA][mC][mA][fU][fU][fU][mU][mC][fC][mU][mC][mG][mU][fG][mG][mU]-L	85	[VP][*fC][*mC][mU][mC][mG][mU][mG][mG][mU][mU][mU][mU][mU][mU][fU][mG][mU][mG][mG][mG][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
QQQ	86	[mU][*mC][*fU][mC][fG][mU][fG][mG][fC][mC][fU][mU][fA][mU][fU][mG][mG][mU]-L	87	[VP][*fU][*mU][fC][mU][fU][mU][fA][mU][fG][mG][fC][mC][fA][mC][fG][mU][fG][mU][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
RRR	658	[fC][*mA][*fA][mG][fG][mU][f	659	[VP][mU][*fU][*mU][fG][mU][f	Mal-

		G][fG][fG][mU][fG][mG][mC][mA][fU][mC][fU][mA][mA][idT]-L		U][fG][fC][mC][fA][mC][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]	C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
SSS	660	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mG][mC][mA][fU][mC][fU][mA][mG][idT]-L	661	[VP][mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]	Mal-C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
TTT	662	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mG][mC][mA][fU][mC][fU][mA][mA][idT]-L	663	[VP][mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]	Mal-C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
UUU	664	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mG][mC][mA][fU][mC][fU][mA][mA][idT]-L	665	[VP][mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]	Mal-C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
VVV	666	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mG][mC][mA][fU][mA][fU][mA][mA][idT]-L	667	[VP][mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]	Mal-C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
WWW	668	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mG][mC][mA][fG][mC][fU][mA][mA][idT]-L	669	[VP][mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]	Mal-C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
XXX	670	[fC][*mA][*fA][mG][fG][mU][fG][fG][fG][mU][fG][mG][mC][mA][fG][mC][fU][mA][mG][idT]-L	671	[VP][mU][*fU][*mA][fG][mA][fU][fG][fC][mC][fA][mC][mC][mC][fA][mC][fC][mU][fU][mG][*fU][*mU]	Mal-C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
YYY	672	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mC][fG][mC][mA][idT]-L	673	[VP][mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
ZZZ	674	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mC][fG][mC][mG][idT]-L	675	[VP][mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
AAAA	676	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mC][fG][mA][mA][idT]-L	677	[VP][mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -

BBBB	678	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mC][fA][mC][mA][id T]-L	679	[VP][mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
CCCC	680	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mC][fG][mC][mA][id T]-L	681	[VP][mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
DDDD	682	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mG][mC][fG][mC][mA][id T]-L	683	[VP][mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
EEEE	684	[mG][*mA][*fG][mC][mC][mA][fU][fC][fU][mU][mU][fG][mC][mA][mA][mG][mC][fG][mC][mG][id T]-L	685	[VP][mU][*fG][*mC][mG][mU][mU][mG][mC][mA][mA][mA][mG][mA][fU][mG][mG][mC][mU][mC][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
FFFF	686	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fC][mU][mA][id T]-L	687	[VP][mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
GGGG	688	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fC][mU][mG][id T]-L	689	[VP][mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
HHHH	690	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fC][mG][mA][id T]-L	691	[VP][mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
IIII	692	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fG][mU][mA][id T]-L	693	[VP][mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
JJJJ	694	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mG][fC][mU][mA][id T]-L	695	[VP][mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆ -
KKKK	696	[mU][*mU][*fA][mC][mG][mG][fU][fA][fU][mC][mU][fA][mC][mA][mU][mG][fC][mU][mA][id T]-L	697	[VP][mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]	Mal-C ₂ H ₄ C(O)(NH)-





					(CH ₂) ₆ -
LLLL	698	[mU][*mU][*fA][mC][mG][mG] [fU][fA][fU][mC][mU][fA][mC][mA][mG][mU][fC][mU][mA][id T]-L	699	[VP][mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][m A][mA][*mU][*mU]	Mal- C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
MMM M	700	[mU][*mU][*fA][mC][mG][mG] [fU][fA][fU][mC][mU][fA][mC][mA][mG][mU][fC][mU][mG][id T]-L	701	[VP][mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][mU][mA][fC][mC][mG][mU][m A][mA][*mU][*mU]	Mal- C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
NNNN	702	[fC][*mA][*fA][mG][fG][mU][f G][fG][fG][mU][fG][mG][mC][mA][fU][mC][fU][mA][mG][idT]-L	703	[VP][mU][*fU][*mA][fG][mA][f U][fG][fC][mC][fA][mC][mC][m C][fA][mC][fC][mU][fU][mG][*f U][*mU]	Mal- C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
OOOO	706	[mU][*mU][*fA][mC][mG][mG] [fU][fA][fU][mC][mU][fA][mC][mA][mU][mU][fC][mU][mA]-L	707	[VP][mU][*fA][*mG][mA][mA][mU][mG][mU][mA][mG][mA][m U][mA][fC][mC][mG][mU][mA][mA][*mU][*mU]	Mal- C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
PPPP	711	[mU][*mG][*fG][mA][mC][mU] [fU][fC][fA][mA][mC][fC][mU][mA][mG][mA][fC][mA][mA]-L	712	[VP][mU][*fU][*mG][mU][mC][mU][mA][mG][mG][mU][mU][m G][mA][fA][mG][mU][mC][mC][mA][*mU][*mU]	Mal- C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
QQQQ	1053	[mG][*mG][*fG][mU][mG][mA] [fC][fA][fA][mC][mU][fA][mC][mU][mA][mU][fC][mU][mA]	1054	[VP][mU][fA][*mG][*mA][mU][mA][mG][mU][mA][mG][mU][mU][mG][fU][mC][mA][mC][m C][mC][*mU][*mU]	Mal- C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
RRRR	1055	[fC][*mU][*fG][mG][fG][mA][f G] [mG][fA][mU][fG][mA][fA][mU] [fU][mC][fG][mA][fA]	1056	[VP][mU][*fU][*mC][fG][mA][f A][mU][fU][mC][fA][mU][fC][m C][fU][mC][fC][mC][fA][mG][*f U][*mU]	Mal- C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
SSSS	1057	[fC][*mA][*fA][mG][fG][mU][f G] [mG][fG][mU][fG][mG][fC][mA] [fU][mC][fU][mA][fA]	1058	[VP][mU][*fU][*mA][fG][mA][f U][mG][fC][mC][fA][mC][fC][m C][fA][mC][fC][mU][fU][mG][*f U][*mU]	Mal- C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -
TTTT	1059	[fU][*mC][*fC][mG][fC][mA][f G] [mC][fC][mU][fG][mG][fA][mU] [fG][mA][fU][mU][fA]	1060	[VP][mU][*fA][*mA][fU][mC][f A][mU][fC][mC][fA][mG][fG][m C][fU][mG][fC][mG][fG][mA][* fU][*mU]	Mal- C ₂ H ₄ C(O)(NH) - (CH ₂) ₆ -

[00223] In some embodiments, the siRNA pairs of A to PPPP provided for above comprise a linker at the 3' end of the sense strand. In some embodiments, the siRNA pairs of A to PPPP provided for above comprise a vinyl phosphonate at the 5' end of the sense strand.

[00224] **Abbreviations Key:** (n/N= any nucleotide) mN= 2'-O-methyl residues, fN= 2'-F residues, * = phosphorothioate, (idT)= inverted Dt, (VP) 2'-O methyl vinyl phosphonate uridine, BMPS = propyl maleimide,


[00225] Structure of the linkers (L) are as follows in Table 3.

Table 3: Representative examples of Linkers (L)

Linker Structure	Linker Name
	Mal-C ₂ H ₄ C(O)(NH)-(CH ₂) ₆
	Mal-(PEG) ₁₂ (NH)(CH ₂) ₆
	Mal-NH-(CH ₂) ₆ or Aminoethyl linker-(CH ₂) ₆ -
	Val-Cit-PABA

[00226] Other linkers can also be used, such as, linkers formed with click chemistry, amide coupling, reductive amination, oxime, enzymatic couplings such as transglutaminase and sortase conjugations. The linkers provided here are exemplary in nature and other linkers made with other such methods can also be used.

[00227] When connected to the siRNA, the structures, L-(X4) can be represented by the following formulas:



[00228] Although certain siRNA sequences are illustrated herein with certain modified nucleobases, the sequences without such modifications are also provided herein. That is, the sequence can comprise the sequences illustrated in the tables provided herein without any modifications. The unmodified siRNA sequences can still comprise, in some embodiments, a linker at the 5' end of the of the sense strand of the dsRNA. In some embodiments, the nucleic acid molecules can be modified to include a vinyl phosphonate at the 5' end of the of the anti-sense strand of the dsRNA. In some embodiments, the nucleic acid molecules can be modified to include a linker at the 3' end of the of the sense strand of the dsRNA. In some embodiments, the nucleic acid molecules can be modified to include a vinyl phosphonate at the 3' end of the of the anti-sense strand of the dsRNA. The linker can be as provided herein.

[00229] In some embodiments, the FN3 proteins comprise a polypeptide comprising a polypeptide that binds CD71 are provided. In some embodiments, the polypeptide comprises a FN3 domain that binds to CD71. In some embodiments, the polypeptide comprises a sequence of SEQ ID NOs: 273, 288-291, 301-310, 312-572, 592-599, or 708-710 are provided. In some embodiments, the polypeptide that binds CD71 comprises a sequence of SEQ ID NOs: 301-301, 310, 312-572, 592-599, or 708-710. The sequence of CD71 protein that the polypeptides can bind to can be, for example, SEQ ID Nos: 2 or 3. In some embodiments, the FN3 domain that binds to CD71 specifically binds to CD71.

[00230] In some embodiments, the FN3 domain that binds CD71 is based on Tencon sequence of SEQ ID NO:1 or Tencon 27 sequence of SEQ ID NO:4
(LPAPKNLVVSRVTEDSARLSWTAPDAAFDSFLIQYQSEKVGGEAIVLTVPGSERSYDLT

GLKPGTEYTVSIYGVKGGHRSNPLSAIFTT), optionally having substitutions at residues positions 11, 14, 17, 37, 46, 73, or 86 (residue numbering corresponding to SEQ ID NO:4).

[00231] In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NOs: 273, 288-291, 301-310, 312-572, 592-599, or 708-710.

[00232] In some embodiments, proteins comprising a polypeptide comprising an amino acid sequence of SEQ ID NO: 273. SEQ ID NO: 273 is a consensus sequence based on the sequences of SEQ ID NO: 288, SEQ ID NO: 289, SEQ ID NO: 290, and SEQ ID NO: 291. The sequence of SEQ ID NO: 273 is

MLPAPKNLVVSRTEDSARLSWTAPDAAFDSEFX₁IX₂YX₃EX₄X₅X₆X₇GEAIX₈LX₉VPGSE
RSYDLTGLKPGTEYX₁₀VX₁₁IX₁₂X₁₃VKGGX₁₄X₁₅SX₁₆PLX₁₇AX₁₈FTT

wherein X₈, X₉, X₁₇, and X₁₈ are each, independently, any amino acid other than methionine or proline, and

X₁ is selected from D, F, Y, or H,

X₂ is selected from Y, G, A, or V,

X₃ is selected from I, T, L, A, or H,

X₄ is selected from S, Y or P,

X₅ is selected from Y, G, Q, or R,

X₆ is selected from G or P,

X₇ is selected from A, Y, P, D, or S,

X₁₀ is selected from W, N, S, or E,

X₁₁ is selected from L, Y, or G,

X₁₂ is selected from D, Q, H, or V,

X₁₃ is selected from G or S,

X₁₄ is selected from R, G, F, L, or D,

X₁₅ is selected from W, S, P, or L, and

X₁₆ is selected from T, V, M, or S.

[00233] In some embodiments:

X₁ is selected from D, F, Y, or H,

X₂ is selected from G, A, or V,

X₃ is selected from T, L, A, or H,

X₄ is selected from Y or P,

X₅ is selected from G, Q, or R,

X₆ is selected from G or P,

X₇ is selected from Y, P, D, or S,

X₁₀ is selected from W, N, S, or E,

X₁₁ is selected from L, Y, or G,

X₁₂ is selected from Q, H, or V,

X₁₃ is selected from G or S,

X₁₄ is selected from G, F, L, or D,

X₁₅ is selected from S, P, or L, and

X₁₆ is selected from V, M, or S.

[00234] In some embodiments, X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, and X₁₆ are as shown in the sequence of SEQ ID NO: 288. In some embodiments, X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, and X₁₆ are as shown in the sequence of SEQ ID NO: 289. In some embodiments, X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, and X₁₆ are as shown in the sequence of SEQ ID NO: 290. In some embodiments, X₁, X₂, X₃, X₄, X₅, X₆, X₇, X₁₀, X₁₁, X₁₂, X₁₃, X₁₄, X₁₅, and X₁₆ are as shown in the sequence of SEQ ID NO: 291.

[00235] In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, phenylalanine, serine, threonine, tryptophan, tyrosine, or valine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, not alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, phenylalanine, serine, threonine, tryptophan, tyrosine, or valine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, alanine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, arginine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, asparagine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, aspartic acid. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, cysteine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, glutamine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, glutamic acid. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, glycine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, histidine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, isoleucine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, leucine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, lysine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, phenylalanine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, serine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, threonine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, tryptophan. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, tyrosine. In some embodiments, X₈, X₉, X₁₇, and X₁₈ is, independently, valine.

[00236] In some embodiments, the sequence is set forth as shown in in the sequence of SEQ ID NO: 288, except that the positions that correspond to the positions of X₈, X₉, X₁₇, and X₁₈ can be any other amino acid residue as set forth above, except that in some embodiments, X₈ is not V, X₉ is not T, X₁₇ is not S, and X₁₈ is not I.

[00237] In some embodiments, the sequence is set forth as shown in in the sequence of SEQ ID NO: 289, except that the positions that correspond to the positions of X₈, X₉, X₁₇, and X₁₈ can be any other amino acid residue as set forth above, except that in some embodiments, X₈ is not V, X₉ is not T, X₁₇ is not S, and X₁₈ is not I.

[00238] In some embodiments, the sequence is set forth as shown in in the sequence of SEQ ID NO: 290, except that the positions that correspond to the positions of X₈, X₉, X₁₇, and X₁₈ can be any other amino acid residue as set forth above, except that in some embodiments, X₈ is not V, X₉ is not T, X₁₇ is not S, and X₁₈ is not I.

[00239] In some embodiments, the sequence is set forth as shown in in the sequence of SEQ ID NO: 291, except that the positions that correspond to the positions of X₈, X₉, X₁₇, and X₁₈ can be any other amino acid residue as set forth above, except that in some embodiments, X₈ is not V, X₉ is not T, X₁₇ is not S, and X₁₈ is not I.

In some embodiments, proteins comprising a polypeptide comprising an amino acid sequence that is at least 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a sequence of SEQ ID NO: 273. In some embodiments, the protein is at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a sequence of SEQ ID NO: 273. In some embodiments, the protein is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to a sequence of SEQ ID NO: 273. In some embodiments, the protein is at least 95%, 96%, 97%, 98% or 99% identical to a sequence of SEQ ID NO: 273.

[00240] Percent identity can be determined using the default parameters to align two sequences using BlastP available through the NCBI website.

[00241] In some embodiments, fibronectin type III (FN3) domains that bind or specifically bind human CD71 protein (SEQ ID Nos: 2 or 5) are provided. As provided herein, the FN3 domains can bind to the CD71 protein. Also provided, even if not explicitly stated is that the domains can also specifically bind to the CD71 protein. Thus, for example, a FN3 domain that binds to CD71 would also encompass a FN3 domain protein that specifically binds to CD71. These molecules can be used, for example, in therapeutic and diagnostic applications and in imaging. In some embodiments, polynucleotides encoding the FN3 domains disclosed herein or

complementary nucleic acids thereof, vectors, host cells, and methods of making and using them are provided.

[00242] In some embodiments, an isolated FN3 domain that binds or specifically binds CD71 is provided.

[00243] In some embodiments, the FN3 domain comprises two FN3 domains connected by a linker. The linker can be a flexible linker. The linker can be a short peptide sequence, such as those described herein. For example, the linker can be a G/S linker and the like.

[00244] In some embodiments, the FN3 domain comprising two FN3 domains connected by a linker, such as those provided for herein. Exemplary linker include, but are not limited to, (GS)₂, (SEQ ID NO: 720), (GGGS)₂ (SEQ ID NO: 721), (GGGS)₁₋₅ (SEQ ID NO: 722), (AP)₁₋₂₀; (AP)₂ (SEQ ID NO: 723), (AP)₅ (SEQ ID NO: 724), (AP)₁₀ (SEQ ID NO: 725), (AP)₂₀ (SEQ ID NO: 726), A(EAAAK)₅AAA (SEQ ID NO: 727), or (EAAAK)₁₋₅ (SEQ ID NO: 728). In some embodiments, the linker comprises or is an amino acid sequence of: EAAAKEAAAKEAAAKEAAAK (SEQ ID NO: 729); GGGSGGGSGGGSGGGGS (SEQ ID NO: 730); APAPAPAPAP (SEQ ID NO: 731); or EAAAK (SEQ ID NO: 732).

[00245] In some embodiments, the FN3 domain may bind CD71 with a dissociation constant (K_D) of less than about 1×10^{-7} M, for example less than about 1×10^{-8} M, less than about 1×10^{-9} M, less than about 1×10^{-10} M, less than about 1×10^{-11} M, less than about 1×10^{-12} M, or less than about 1×10^{-13} M as determined by surface plasmon resonance or the Kinexa method, as practiced by those of skill in the art. The measured affinity of a particular FN3 domain-antigen interaction can vary if measured under different conditions (e.g., osmolarity, pH). Thus, measurements of affinity and other antigen-binding parameters (e.g., K_D , K_{on} , K_{off}) are made with standardized solutions of protein scaffold and antigen, and a standardized buffer, such as the buffers described herein.

[00246] In some embodiments, the FN3 domain may bind CD71 at least 5-fold above the signal obtained for a negative control in a standard solution ELISA assay.

[00247] In some embodiments, the FN3 domain that binds or specifically binds CD71 comprises an initiator methionine (Met) linked to the N-terminus of the molecule. In some embodiments, the FN3 domain that binds or specifically binds CD71 comprises a cysteine (Cys) linked to a C-terminus of the FN3 domain. The addition of the N-terminal Met and/or the C-terminal Cys may facilitate expression and/or conjugation to extend half-life and to provide other functions of molecules.

[00248] The FN3 domain can also contain cysteine substitutions, such as those that are described in U.S. Patent No. 10,196,446, which is hereby incorporated by reference in its

entirety. Briefly, in some embodiments, the polypeptides provided herein can comprise at least one cysteine substitution at a position selected from the group consisting of residues 6, 8, 10, 11, 14, 15, 16, 20, 30, 34, 38, 40, 41, 45, 47, 48, 53, 54, 59, 60, 62, 64, 70, 88, 89, 90, 91, and 93 of the FN3 domain based on SEQ ID NO: 6 or SEQ ID NO: 1 of U.S. Patent No. 10,196,446, and the equivalent positions in related FN3 domains. In some embodiments, the substitution is at residue 6. In some embodiments, the substitution is at residue 8. In some embodiments, the substitution is at residue 10. In some embodiments, the substitution is at residue 11. In some embodiments, the substitution is at residue 14. In some embodiments, the substitution is at residue 15. In some embodiments, the substitution is at residue 16. In some embodiments, the substitution is at residue 20. In some embodiments, the substitution is at residue 30. In some embodiments, the substitution is at residue 34. In some embodiments, the substitution is at residue 38. In some embodiments, the substitution is at residue 40. In some embodiments, the substitution is at residue 41. In some embodiments, the substitution is at residue 45. In some embodiments, the substitution is at residue 47. In some embodiments, the substitution is at residue 48. In some embodiments, the substitution is at residue 53. In some embodiments, the substitution is at residue 54. In some embodiments, the substitution is at residue 59. In some embodiments, the substitution is at residue 60. In some embodiments, the substitution is at residue 62. In some embodiments, the substitution is at residue 64. In some embodiments, the substitution is at residue 70. In some embodiments, the substitution is at residue 88. In some embodiments, the substitution is at residue 89. In some embodiments, the substitution is at residue 90. In some embodiments, the substitution is at residue 91. In some embodiments, the substitution is at residue 93.

[00249] A cysteine substitution at a position in the domain or protein comprises a replacement of the existing amino acid residue with a cysteine residue. In some embodiments, instead of a substitution a cysteine is inserted into the sequence adjacent to the positions listed above. Other examples of cysteine modifications can be found in, for example, U.S. Patent Application Publication No. 20170362301, which is hereby incorporated by reference in its entirety. The alignment of the sequences can be performed using BlastP using the default parameters at, for example, the NCBI website.

[00250] In some embodiments, a cysteine residue is inserted at any position in the domain or protein.

[00251] In some embodiments, the FN3 domain that binds CD71 is internalized into a cell. In some embodiments, internalization of the FN3 domain may facilitate delivery of a detectable label or therapeutic into a cell. In some embodiments, internalization of the FN3 domain may

facilitate delivery of a cytotoxic agent into a cell. The cytotoxic agent can act as a therapeutic agent. In some embodiments, internalization of the FN3 domain may facilitate the delivery of any detectable label, therapeutic, and/or cytotoxic agent disclosed herein into a cell. In some embodiments, internalization of the FN3 domain may facilitate delivery of an oligonucleotide into a cell. In some embodiments, the cell is a tumor cell. In some embodiments, the cell is a liver cell. In some embodiments, the cell is a muscle cell. In some embodiments, the cell is an immune cell. In some embodiments, the cell is a dendritic cell. In some embodiments, the cell is a cell of the central nervous system. In some embodiments, the cell is a heart cell.

[00252] In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NOs: 273, 288-291, 301-310, 312-572, 592-599, or 708-710.

[00253] In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:301. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:302. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:303. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:304. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:305. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:306. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:307. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:310. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:312. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:313. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:314. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:315. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:316. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:317. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:318. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:319. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:320. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:321. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO:322. In

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 564. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 565. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 566. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 567. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 568. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 569. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 570. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 571. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 572. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 708. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 709. In some embodiments, an isolated FN3 domain that binds CD71 comprises the amino acid sequence of SEQ ID NO: 710.

[00254] In some embodiments, the isolated FN3 domain that binds CD71 comprises an initiator methionine (Met) linked to the N-terminus of the molecule.

In some embodiments, the isolated FN3 domain that binds CD71 comprises an amino acid sequence that is 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to one of the amino acid sequences of SEQ ID NOs: 273, 288-291, 301-310, 312-572, 592-599, or 708-710. Percent identity can be determined using the default parameters to align two sequences using BlastP available through the NCBI website. The sequences of the FN3 domains that bind to CD71 can be found, for example, in Table 4.

Table 4: CD71-binding FN3 domain sequences

SEQ ID	Amino Acid sequence of FN3 domains that bind to CD71
301	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFPITYIEAVVLGEAIVLTVPGSE RSYDLTGLKPGTEYPVGISGVKGGHNSMPLSAIFTT
302	MLPAPKNLVVSEVTEDSARLSWQGVARAFDSFMINYSELFWMGEAIVLTVPG SERSYDLTGLKPGTEYVVRIGVKGGKGSWPLHAHFTT
303	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFNIEYAETRWYGEAIVLTVPGS ERSYDLTGLKPGTEYVVPIDGVKGGIASKPLSAIFTT

304	MLPAPKPNLVVSEVTEDSARLSWQGVARAFDSFLITYRDQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAESTT
305	MLPAPKPNLVVSEVTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYWVYIWGVKGGKPSFPLRAGFTT
306	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFPIIYMETFSRGEAIVLTVPGSE RSYDLTGLKPGTEYRVPPIGGVKGGSSCPLSAIFTT
307	MLPAPKPNLVVSDVTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAEFTT
310	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFKIAYIETATRGEAIVLTVPGSE RSYDLTGLKPGTEYVVPPIGVKGGNTSSPLSAIFTT
312	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFFIPYAEPSPTGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGHLSPLSAISTT
313	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFLIAYPEDGFRGEAIVLTVPGSE RSYDLTGLKPGTEYPVPILGVKGGGSGPLSAIFTT
314	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFPIIYYVENVWGEAIVLTVPGS ERSYDLTGLKPGTEYWEVIIGVKGGQCSRPLSAIFTT
315	MLPAPKPNLVVSRVTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTECPVWIQGVKGGSPSAPLSAEFTT
316	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFGIAYREFRPSGEAIVLTVPGSE RSYDLTVETGYRNEVVICGVKGGPWSGPLSAIFTT
317	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFPILYTECVYRGEAIVLTVPGSE RSYDLTGLKPGTEYHVPITGVKGGGGSWPLSAIFTT
318	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFNIMYHEIIYVGEAIVLTVPGSE RSYDLTGLKPGTEYPVPIEGVKGGGTSGPLSAIFTT
319	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFAITYTEAALCGEAIVLTVPGSE RSYDLTGLKPGTEYPVPINGVKGGGTSGPLSAIFTT
320	MLPAPKPNLVVARVTEDSARLSWTAPDAAIDSFPIDYSEYWWGGEAIVLTVPGS ERSYDLTGLKPGTEYPVLITGVKGGYRSGPLSAIFTT
321	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFSIRYNEFIVAGEAIVLTVPGSE RSYDLTGLKPGTEYDVPIAGVKGGGASWPLSAIVTT
322	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFWYLELQFAGEAIVLTVPGSER SYDLTGLKPGTEYNVPITGVKGGHISFPLSAIFTT
323	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFAIWYHEWYGDGEAIVLTVPG SERSYDLTGPKPGTEYRVRISGVKGGFESGPLSAIFTT
324	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFMIRYQEGTRWGEAIVLTVPGS ERSYDLTGLKPGTEYIVMIAGVKGGQISLPLSAIFTT
325	MLPAPKPNLVVSRVTEDSARLSWTAPDAAFDSFPIWYLEKSYQGEAIVLTVPGS ERSYDLTGLKPGTEYVPIIGVKGGRDSCPLSAIFTT
326	MLPAPKPNLVVSEVTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAEFTT

327	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFRISYAETVRQGEAIVLTVPGSE RSYDLTVETGYRNWVMILGVKGGPGSLPLSAIFTT
328	MLPAPKNLVVSEVTEDSARLSWQGVVRAFDNFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWVQGVKGGSPSAPLSAEFTT
329	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFWIEYWEAVGFGEAIVLTVPGS ERSYDLTGLKPGTEYFVGIYGVKGGYLSAPLSAIFTT
330	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFIIHYVEQQLIGEIVLTVPGSE RSYDLTGLKPGTEYPVPITGVKGGACSWPLSAIFTT
331	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFTIEYSEHPIDGEAIVLTVPGSER SYDLTGLKPGTEYVRIHGVKGGWFSHPLWAFFTT
332	MLPAPKNLVVSRVTEDSARLSWQGVVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYGVTIAGVKGGWRSKPLNAESTT
333	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFGIAYVESYWYGEAIVLTVPGS ERSYDLTGLKPGTEYNVPIYGVKGGDGSGLPLSAIFTT
334	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYITYVELNLAGEAIVLTVPGS ERSYDLTGLKPGTEYPVPILGVKGGSLSQPLSAIFTT
335	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFPIYSIESIADGEAIVLTVPGSER SYDLTGLKPGTEYWVAIVGVKGGPFSWSLSAIVTT
336	MLPAPKNLVVSEVTEDSARLSWQGVVARAFDSFLITYREQIFAGEVIVPTVPGSE RSYDLTGLKPGTEYPVPIAGVKGGGPSAPLSAIFTT
337	MLPAPKNLVVSRVTEDSARLSWTTPDAAFDSFPIYYWEVTITGEAIYLSVPGSE RSYDLTGLKPGTEYPVDIPGVKGGAAASPPLSAIFTT
338	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFPILYLEHTVRSEAIVLTVPGSE RSYDLTDLKPGTEYCVPIDGVKGGRLSRPLSAIFTT
339	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFPIPYTEPPDPGEAIVLTVPGSE RSYDLTGLKPGTEYLVITLGVKGGSMVPLSAIFTT
340	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFTIDYWENRCPGEAIVLTVPGS ERSYDLTGLKPGTEYCVWISGVKGGYSSWPLSAIFTT
341	MLPAPKNLVVSRVTEDSARLSWQGVVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWVQGVKGGHLSLPLSAIVTT
342	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFLIPYAETSPSGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGDYSEPLSAIFTT
343	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFMIVYYEYTRFGEAIVLTVPGS ERSYDLTGLKPGTEYTPIDGVKGGGRSSPLSAIFTT
344	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFFIPYAEPSTGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGHLSLPLSAIVTT
345	MLPAPKNLVVSEVTEDSARLSWQGVVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWVQGVKGGSPSAPLSAEFTT
346	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFIIPYAEVRPDGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGKLSLPLSAIFTT

347	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFGIVYLEMMVTGEAIVLTVPGS ERSYDLTGLKPGTEYDVPILGVKGGTRSVPLSAIFTT
348	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFLIYYEEGYLEYEYYSGEAIVLT VPGSERSYDLTGLKPGTEYYVGIVGVKGGGLSGPLSAISTT
349	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFPIAYAEPRPDGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGDWLPLSAIFTT
350	MSPAPKNLVVSRTEDSARLSWTAPDAAAFDSFTIHYREFQLSGEAIVLTVPGS ERSYDLTGLKPGTEYDVPIEGVKGGPGSRPLSAIFTT
351	MLPAPKNLVVSRTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE CSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAEFTT
352	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFEIDYDELAIFYGEAIVLTVPGSE RSYDLTGLKPGTEYGVRIPIGVKGGMPSLPLSAIVTT
353	MLPAPENLVVSRTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAESTT
354	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFAIAYGEHIVIGEIVLTVPGSE RSYDLTGLKPGTEYMPVPIAGVKGGPISLPLSAIFTT
355	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFFIPYAEPSPTGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGHLSLPLSAIFTT
356	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFSIGYVELVLLGEAIVLTVPGSE RSYDLTGLKPGTEYDVLIPGVKGGSLSRPLSAIFTT
357	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFAIPYAELSRNGEAIVLTVPGSE RSYDLTGLKPGTEYTVLIHGVKGGCLSDPLSAIFTT
358	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFHIEYLELSRHGEAIVLTVPGSE RSYDLTGLKPGTEYWVMIFGVKGGGPSKPLSAIFTT
359	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFVYNEVHWIGEIVLTVPGSER SYDLTGLKPGTEYFVGIVGVKGGHWSKPLSAIFTT
360	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFEIDYDELAIFYGEAIVLTVPGSE RSYDLTGLKPGTEYGVRIPIGVKGGMPSLPLSAIVTT
361	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFQIVYSELWIKGEAIVLTVPGSE RSYDLTGLKPGTEYQVPIPIGVKGGGRNSFPLSAIFTT
362	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFKIRYTETRSIGEIVLTVPGSE RSYDLTGLKPGTEYCVPIGGVKGGDSSWPLSAISTT
363	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFCISYYERMGRGEAIVLTVPGS ERSYDLTGLKPGTEYMVYIFGVKGGGLNSLPLSAIFTT
364	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFYIVYAEPIPNGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGGRNSDPLSAIFTT
365	MLPAPKNLVVSRTKDSARLSWTAPDAAAFDSFPIAYAEPRPDGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAIFTT
366	MLPAPKNLVVSRTEDSARLSWTAPDAAAFDSFTIDYDEPRSPGEAIVLTVPGSE RSYDLTGLKPGTEYRVFIWGIKGGDTSFPLSAIFTT

367	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFTILYAEQAQFGEAIVLTVPGSE RSYDLTGLKPGTEYPITGVKGGTRSGPLSAISTT
368	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFIIPYAEVRPDGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGHLSPLSAISTT
369	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFHIA YEETATSGEAIYLRVPGSE RSYDLTGLKPGTEYGVEIEGVKGGARSRLYADFTT
370	MLPAPKNLVVSRTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWIQGVKGGDLNPLSAIFTT
371	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFPISYLELSLYGEAIVLTVPGSE RSYDLTGLKPGTEYPVGIAGVKGGVSRPLSAIFTT
372	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFTIGYREWYWGGEAIVLTVPG SERSYDLTGLKPGTEYNVPISGVKGGDLDFPLSAIFTT
373	MLPAPKNLVVSRTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAESTT
374	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFSITYLEWWNLGEAIVLTVPGS ERSYDLTGLKPGTEYMTIPGVKGGMSSYPLSAIFTT
375	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFTISYGEEALIGEAIYLRVPGSE RSYDLTGLKPGTEYYVHIEGVKGGWSQPLAAFTT
376	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFTIEYYENIGIGEIVLTVPGSE RSYDLTGLKPGTEYSVPIGVKGGPYSHPLSAIFTT
377	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFPIAYAEPRPDGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAIFTT
378	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFGIGYYEHKRFGEAIQLSVPGS ERSYDLTGLKPGTEYEV DIEGVKGGVLSWPLFAEFTT
379	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFVIEYTERFWSGEAIVLTVPGSE RSYDLTGLKPGTEYSVPIDGVKGGQCSTPLSAIFTT
380	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFWIDYEEEGVIGEAIYLVHPGS ERSYDLTGLKPGTEYVVKIHGVKGGHPSHPLVAVFTT
381	MLPAPKNLVVSRTEDSARLSWQGVARAFDSFLITYVELRHLGEAIVLTVPGS ERSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAEFTT
382	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFLIPYAETSPSGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGDYSSPLSAIFTT
383	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFPIAYAEPRPDGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAISTT
384	MLPAPKNLVVSRTEDSARLSWQGVARAFDSFSILYLELTPKGEAIVLTVPGSE RSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAEFTT
385	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFIIIEYFEPIPIGEAIVLTVPGSERS YDLTGLKPGTEYAVNIYGVKGGYLSHPLSAIFTT
386	MLPAPKNLVVSRTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE CSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAEFTT

387	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFAIEYTEFLYSGEAIVLTVPGSE RSYDLTGLKPGTEYGVPIINGVKGGFVSPPLSAIFTT
388	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFPIKYREVLRCGEAIVLTVPGSE RSYDLTGLKPGTEYTPITGVKGGFGSSPLSAIFTT
389	MLPAPENLVVSRTEDSARLSWTAPDAAFDSFWIEYYEGVIQGEAIVLTVPGSE RSYDLTGLKPGTEYFVAIWGVKGGKWSVPLSAIFTT
390	MLPAPKNLVVSRTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWIQGVKGGSPSAEFTT
391	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFQIHYWETQGFGEAIVLTVPGS ERSYDLTGLKPGTEYPVLIPGVKGGPSSLPLSAIFTT
392	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIPYAEPSTGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGHLSPLSAIFTT
393	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFAIEYYEPVPAGEAIYLDVPGS ERSYDLTGLKPGTEYDVTIYGVKGGYSHPLFASFTT
394	MLPAPKNLVVSRTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAIFTT
395	MLPAPKNLVVSRTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAEFTT
396	MSPAPKNLVVSRTEDSARLSWTAPDAAFDSFPIAYLEV FYEGEAIVLTVPGS ERSYDLTGLKPGTEYQVPIEGVKGGAMSLPLSAIFTT
397	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFHIWYEEETTIGEAIYLHVPGE RSYDLTGLKPGTEYEVHITGVKGGPYSRPLFANFTT
398	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFGIAYDEWPEFGEAIVLTVPGS ERSYDLTGLKPDTEYIVEIYGVKGGWFSWPLSAIFTT
399	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIPYAEPSTGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGHLSPLSVIFTT
400	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIWEYEEVMYLGEAIVLTVPGS ERSYDLTGLKPGTEYNVPIPGVKGGHSSPPLSAIFTT
401	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFHILYEELFLVGEAIVLTVPGSE RSYDLTGLKPGTEYKVPISGVKGGPVSRLPLSAIFTT
402	MLPAPKNLVVSRTEDSARLSWQGVARAFDSFLITYREQIFAGEVIVLTVPGSE RSYDLTGLKPGTEYPVWIQGVKGGSPSAPLSAEFTT
403	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFHIVYHEPRPSGEAIWLHVPGE ERSYDLTGLKPGTEYEVGIVSVKGGDLVPLVAFFTT
404	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAISLLVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLYAVFTT
405	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIFLVVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLHANFTT
406	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLDVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLYASFTT

407	MSLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAISLYVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAIFTT
408	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIQLRVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFSPMPLSAIFTT
409	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFHISYEEDYTFGEAIYLRVPGSE RSYDLTGLKPGTEYRVVIGGVKGGWFSEPLLAFTT
410	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIYLTVPGESE RSYDLTGLKPGTEYNVTIQGVKGGFSPSYPLDASFTT
411	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIDLGVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFSPMPLDPLEAYFTT
412	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLLVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAIFTT
413	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAINLQVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLVAFFTT
414	MLPAPKNLVVSRTEDSARLSWTTPDAAFDSFFIGYLEPQPPGEAISLQVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSSPLFAVFTT
415	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIELHVPGESE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLFTT
416	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLVVPGESE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAIFTT
417	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAITLDVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAIFTT
418	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIWLVPGESE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLVASFTT
419	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAINLDVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLVAEFTT
420	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIHLSVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLVAIFTT
421	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIALWVPGESE RSYDLTGLKPGTEYNVTIQGVKGGFSPMPLSAIFTT
422	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAAILVVPGESE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAHFTT
423	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIQLWVPGESE RSYDLTGLKPGTEYNVTIQGVKGGFSPHPLGAVFTT
424	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLHVPGESE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLLASFTT
425	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIALHVPGESE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLVAFFTT
426	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIHLHVPGESE RSYDLTGLKPGTEYNVTIQGVKGGFSPILHANFTT

427	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIFLGVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAIFTT
428	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIVLRVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLIASFTT
429	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAINLWVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLDASFTT
430	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFVIEYFEWTLNGEAIVLTVPGS ERSYDLTGLKPGTEYSVQIYGVKGGLSRPLSAIFTT
431	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIHLWVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLIAHFTT
432	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIPYAEPSPTGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAHFTT
433	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIYLYVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLDAFTT
434	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIGLQVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAIFTT
435	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLAVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLHAFTT
436	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIWLHVPGE RSYDLTGLKPGTEYNVTIQGVKGGFPSIPLIAFTT
437	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIVLDVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLVAEFTT
438	MLPTPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLRVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLHASFTT
439	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIQLGVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSHPLNANFTT
440	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIQLEVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSMPLSAIFTT
441	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIFLGVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLIAFTT
442	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIGLQVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSHPLKAQFTT
443	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLFVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLVAHFTT
444	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIGLYVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLGAFTT
445	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLQVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLTAFTT
446	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAITLHVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAIFTT

447	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIVLEVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLDAHFTT
448	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIALHVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLRAVFTT
449	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLWVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAIFTT
450	MLPAPKNLVVSRTEDSARLSRTAPDAAFDSFYIAYAEPRPDGEAIVLIVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAIFTT
451	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIQLWVPGE RSYDLTGLKPGTEYNVTIQGVKGGFPSRPLQAHFTT
452	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAITLDVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLYAFHTT
453	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIALHVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAIFTT
454	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIGLWVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLIAHFTT
455	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIWLVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLHARFTT
456	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIFLQVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLAAVFTT
457	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLHVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAISTT
458	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIIHQVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAVFTT
459	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIYLKVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLVAHFTT
460	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIVLTVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLLAYFTT
461	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIIHLVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLEAKFTT
462	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIKLEVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLAIFTT
463	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIYLEVPGE RSYDLTGLKPGTEYNVTIQGVKGGFPSFPLKAAFTT
464	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIIHRVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLVAIFTT
465	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIVLQVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLAAWFTT
466	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIFLQVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLNAFTT

467	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAILGVPGSER RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLHAYSTT
468	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLDVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAIFTT
469	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLLVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAVFTT
470	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIHLLVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLLAHFTT
471	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLWVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLDAYFTT
472	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIVLTVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFSPMPLASFTT
473	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFHIVYHEPRPSGEAIHLQVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFSPSYPLSAFTT
474	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIQLWVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFSPMPLSAIFTT
475	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFRISYCETFYHGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLIAKFTT
476	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIWLKVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLQANFTT
477	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIWLKVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLQANFTT
478	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLQVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAIVTT
479	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFPIAYAEPRPDGEAIVLTVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAFTT
480	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIALLVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLVAQFTT
481	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILHVPGE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLEAKFTT
482	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIDLHVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLHALFTT
483	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLDVPGSE RSYDLTGLKPGTEYSVLIHGVKGGFSPMPLSAIFTT
484	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIDLAVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSFTT
485	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIYLGVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLRAKFTT
486	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLGVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAIFTT

487	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAISLLVPDSE RSYDLTGLKPGTEYNVTIQGVKGGFPSMPLKFTT
488	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIGLVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLDASFTT
489	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLTVPGSE RSYDLTGPKPGTEYWVLIQGVKGGSSVPLVAYFTT
490	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAISLDVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLEASFTT
491	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAII LAVPGSER SYDLTGLKPGTEYNVTIQGVKGGFPSPLVASFTT
492	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFDIGYTEYGGYGEAIYLSVPGS ERSYDLTGLKPGTEYWVLIQGVKGGSSVPLSAIFTT
493	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAISLSVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLIANFTT
494	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIALLVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAIVTT
495	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAII LDVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLSIFTT
496	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIVLVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLRASFTT
497	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIKLDVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLVAFFTT
498	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAII LEVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLVAYFTT
499	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIIHLVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLHADFTT
500	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIWLEVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLVADFTT
501	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAISLVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLLAHFTT
502	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFDIGYTEYGGYGEAILHVPGE RSYDLTGLKPGTEYWVLIQGVKGGSSVPLSAIFTT
503	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIVLLVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSVPLAAFTT
504	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAII LLVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSQFTT
505	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAII LGVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSMPLHPLVALFTT
506	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIGLDVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAIFTT

507	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIHLSVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLAAYFTT
508	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIVLAVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSYPLVAAFTT
509	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLQVPGSCRSYDLTGLKP GTEYSVLIHGVKGGLSSPLTAIFTT
510	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAINLQVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSFPLSAVFTT
511	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLHVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLVAIFTT
512	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIWLAVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLHAQFTT
513	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLGVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLFTT
514	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIGLQVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLCAEFTT
515	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIVLWVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLIAEFTT
516	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAISLSVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPPKFTT
517	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIILEVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLRAVFTT
518	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIHLVVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLSAIFTT
519	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAISLKVPGE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLEAIFTT
520	MLPAPKNPVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIHLLVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLKRLSPPVTTITITMAVCRKPVAEN LSQTLS
521	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIFLDVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSPLTAFTT
522	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIVLDVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSHPLAAFTT
523	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIGLAVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSVPLQANFTT
524	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLRVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLVAEFTT
525	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLQVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLSASFTT
526	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIGLHVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLTASFTT

527	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAIGLRVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAIFTT
528	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAIVLRVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLAASFTT
529	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAIQLLVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLVAHFTT
530	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAIWLLVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLVAFHTT
531	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIHLYVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFSPDPLDAVFTT
532	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIYLDVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFPSTFTT
533	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAIQLFVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLKAYFTT
534	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAIVLVVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAIFTT
535	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIQLTVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFSPSLPLSADFTT
536	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAILLQVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLDAEFTT
537	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAIVLAVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLYASFTT
538	MLPAPKNLVVSRTEDSARLSWTTPDAAFDSFYIAYAEPDPGEAIRLQVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLGFTT
539	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAIQLVVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLYAIFTT
540	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAISLSVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLHAKFTT
541	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIHLGVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFSPISPLFASFTT
542	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAIQLLVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLYAAFTT
543	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIHLAVPGSE RSYDLTGLKPGTEYNVTIQGVKGGFSPVPLAAVFTT
544	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAISLQVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLGAHFTT
545	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAIALWVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLVASFTT
546	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPDPGEAIQLHVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLYAFHTT

547	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLHVPGE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLRASFTT
548	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIWLGVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLHATFTT
549	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIVLEVPGE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLHANFTT
550	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLRVPGE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLYAKFTT
551	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIGLWVPGE RSYDLTGLKPGTEYNVTIQGVKGGFSPDPLQAVFTT
552	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIVLHVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLDAFTT
553	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILHVPGE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLDAYFTT
554	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLAVPGE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLSAKFTT
555	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAILLFVPGE RSYDLTGLKPGTEYNVTIQGVKGGFPSTPLSASFTT
556	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLTVPGE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLHAYFTT
557	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLGVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLRAYFTT
558	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLEVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLVAFFTT
559	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIQLGVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLLAFTT
560	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIHLRVPGE ERSYDLTGLKPGTEYNVTIQGVKGGFPMPLSAFTT
561	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAILLQVPGE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLIAKFTT
562	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIVLHVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLQAIFTT
563	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIALVVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLAANFTT
564	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAINLSVPGE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLDAYFTT
565	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIVLEVPGE RSYDLTGLKPGTEYSVLIHGVKGGLLSSPLTASFTT
566	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIRLQVPGE ERSYDLTGLKPGTEYSVLIHGVKGGLLSSPLGASFTT

567	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIGLWVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLVAYFTT
568	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIYLEVPGSE RSYDLTGLKPGTEYSVLIHGVKGGLSSPLFTT
569	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFYIAYAEPRPDGEAIWLDVPGS ERSYDLTGLKPGTEYSVLIHGVKGGLSSPLDAYFTT
570	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFAIEYCETKMCGEAIVLTVPGS ERSYDLTGLKPGTEYRVPPIGVKGGTASLPLSAIFTT
571	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIYYIESYPAGEAIVLTVPGSE RSYDLTGLKPGTEYWVGIDGVKGGRWSTPLSAIFTT
572	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFFIYYIESYPAGEAIVLTVPGSC RSYDLTGLKPGTEYWVGIDGVKGGRWSTPLSAIFTT
592	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFAIVYHEPRPDGEAIVLTVPGS CRSYDLTGLKPGTEYEVVILGVKGGVHSYPLSAIFTTAPAPAPAPAPLPAPKNL VVSRTEDSARLSWTAPDAAFDSFAIVYHEPRPDGEAIVLTVPGSERSYDLTGL KPGTEYEVVILGVKGGVHSYPLSAIFTT
593	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFAIVYHEPRPDGEAIVLTVPGS CRSYDLTGLKPGTEYEVVILGVKGGVHSYPLSAIFTTGGGGSGGGGSGGGGSG GGGSLPAPKNLVVSRTEDSARLSWTAPDAAFDSFAIVYHEPRPDGEAIVLTV PGSERSYDLTGLKPGTEYEVVILGVKGGVHSYPLSAIFTT
594	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFAIVYHEPRPDGEAIVLTVPGS CRSYDLTGLKPGTEYEVVILGVKGGVHSYPLSAIFTTEAAAKEAAAKEAAAKE AAAKLPAPKNLVVSRTEDSARLSWTAPDAAFDSFAIVYHEPRPDGEAIVLTV PGSERSYDLTGLKPGTEYEVVILGVKGGVHSYPLSAIFTT
595	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFAIVYHEPRPDGEAIVLTVPGS CRSYDLTGLKPGTEYEVVILGVKGGVHSYPLSAIFTTEAAAKLPAPKNLVVS RTEDSARLSWTAPDAAFDSFAIVYHEPRPDGEAIVLTVPGSERSYDLTGLKPGT EYEVVILGVKGGVHSYPLSAIFTT
596	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFKIEYFEYVGYGEAIVLTVPGS CRSYDLTGLKPGTEYYVAIYGVKGGWYSRPLSAIFTTAPAPAPAPAPLPAPKNL VVSRTEDSARLSWTAPDAAFDSFKIEYFEYVGYGEAIVLTVPGSERSYDLTGL KPGTEYYVAIYGVKGGWYSRPLSAIFTT
597	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFKIEYFEYVGYGEAIVLTVPGS CRSYDLTGLKPGTEYYVAIYGVKGGWYSRPLSAIFTTGGGGSGGGGSGGGGS GGGSLPAPKNLVVSRTEDSARLSWTAPDAAFDSFKIEYFEYVGYGEAIVLT VPGSERSYDLTGLKPGTEYYVAIYGVKGGWYSRPLSAIFTT
598	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFKIEYFEYVGYGEAIVLTVPGS CRSYDLTGLKPGTEYYVAIYGVKGGWYSRPLSAIFTTEAAAKEAAAKEAAAK EAAAKLPAPKNLVVSRTEDSARLSWTAPDAAFDSFKIEYFEYVGYGEAIVLT VPGSERSYDLTGLKPGTEYYVAIYGVKGGWYSRPLSAIFTT
599	MLPAPKNLVVSRTEDSARLSWTAPDAAFDSFKIEYFEYVGYGEAIVLTVPGS CRSYDLTGLKPGTEYYVAIYGVKGGWYSRPLSAIFTTEAAAKLPAPKNLVVS RTEDSARLSWTAPDAAFDSFKIEYFEYVGYGEAIVLTVPGSERSYDLTGLKPGT

	EYYVAIYGVKGGWYSRPLSAIFTT
708	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFAIVYHEPRPDGEAIVLTVPGSCRSYDLTGLKP GTEYEVVILGVKGGVHSYPLSAIFTT
709	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFAIVYHEPRPDGEAIVLTVPGSERSYDLTGLKP GTEYEVVILGVKGGVHSYPLSAIFTT
710	MLPAPKNLVVSRVTEDSARLSWTAPDAAFDSFFIGYLEPQPPGEAIHLGVPGSCRSYDLTGLKPG TEYNVTIQGVKGGFPSIPLFASFTT

[00255] As provided herein, in some embodiments, the FN3 domain that binds to CD71 binds to SEQ ID NO: 2 (human mature CD71) or SEQ ID NO: 5 (human mature CD71 extracellular domain), sequence of each provided below:

2	MTKEYQDLQHLDNAEEDHHQLRKGPPLQRLCSGPRLLLLSLGLSLLLLV VCVIGSQNSQLQEELRGLRETFSNFTASTEAVKGLSTQGGNVGRKMKSLESQLE KQKDLSEDHSSLLLHVVKQFVSDLRSLSCQMAALQGNGSERTCCPVNWVEHERS CYWFSRSGKAWADADNYCRLEDAHLVVVTSWEEQKFVQHHIGPVNTWMGLHD QNGPWKWVDGTDYETGFKNWRPEQPDDWYGHGLGGGEDCAHFTDDGRWNDD VCQRPYRWVCETELDKASQEPPLL
5	QNSQLQEELRGLRETFSNFTASTEAVKGLSTQGGNVGRKMKSLESQLEKQKQD LSEDHSSLLLHVVKQFVSDLRSLSCQMAALQGNGSERTCCPVNWVEHERSCYWFS RSGKAWADADNYCRLEDAHLVVVTSWEEQKFVQHHIGPVNTWMGLHDQNGPW KWVDGTDYETGFKNWRPEQPDDWYGHGLGGGEDCAHFTDDGRWNDDVCQRP YRWVCETELDKASQEPPLL

[00256] In some embodiments, the FN3 domain comprises two FN3 domains connected by a linker. The linker can be a flexible linker. The linker can be a short peptide sequence, such as those described herein. For example, the linker can be a G/S or G/A linker and the like. As provided herein, the linker can be, for example, (GS)₂, (SEQ ID NO:720), (GGGS)₂ (SEQ ID NO:721), (GGGS)₅ (SEQ ID NO:722), (AP)₂₋₂₀, (AP)₂ (SEQ ID NO:723), (AP)₅ (SEQ ID NO:724), (AP)₁₀ (SEQ ID NO:725), (AP)₂₀ (SEQ ID NO:726) and A(EAAAK)₅AAA (SEQ ID NO:727) or (EAAAK)₁₋₅ (SEQ ID NO: 728). These are non-limiting examples and other linkers can also be used. The number of GGGGS or GGGGA repeats can also be 1, 2, 3, 4, or 5. In some embodiments, the linker comprises one or more GGGGS repeats and one or more GGGGA repeats. In some embodiments, the linker comprises EAAAKEAAKEAAKEAAK (SEQ ID NO: 729); GGGSGGGSGGGSGGGGS (SEQ ID NO: 730); APAPAPAPAP (SEQ ID NO: 731); or EAAAK (SEQ ID NO: 732).

[00257] In some embodiments, the FN3 domain comprising two FN3 domains connected by a linker have the amino acid sequence of SEQ ID NO: 592. In some embodiments, the FN3 domain comprising two FN3 domains connected by a linker have the amino acid sequence of SEQ ID NO: 593. In some embodiments, the FN3 domain comprising two FN3 domains

connected by a linker have the amino acid sequence of SEQ ID NO: 594. In some embodiments, the FN3 domain comprising two FN3 domains connected by a linker have the amino acid sequence of SEQ ID NO: 595. In some embodiments, the FN3 domain comprising two FN3 domains connected by a linker have the amino acid sequence of SEQ ID NO: 596. In some embodiments, the FN3 domain comprising two FN3 domains connected by a linker have the amino acid sequence of SEQ ID NO: 597. In some embodiments, the FN3 domain comprising two FN3 domains connected by a linker have the amino acid sequence of SEQ ID NO: 598. In some embodiments, the FN3 domain comprising two FN3 domains connected by a linker have the amino acid sequence of SEQ ID NO: 599. In some embodiments, the FN3 domain comprising two FN3 domains connected by a linker have the amino acid sequence of one of SEQ ID NOs: 592-599.

[00258] In some embodiments, the FN3 domains may bind CD71, as applicable, with a dissociation constant (K_D) of less than about 1×10^{-7} M, for example less than about 1×10^{-8} M, less than about 1×10^{-9} M, less than about 1×10^{-10} M, less than about 1×10^{-11} M, less than about 1×10^{-12} M, or less than about 1×10^{-13} M as determined by surface plasmon resonance or the Kinexa method, as practiced by those of skill in the art. The measured affinity of a particular FN3 domain-antigen interaction can vary if measured under different conditions (e.g., osmolarity, pH). Thus, measurements of affinity and other antigen-binding parameters (e.g., K_D , K_{on} , K_{off}) are made with standardized solutions of protein scaffold and antigen, and a standardized buffer, such as the buffers described herein.

[00259] In some embodiments, the FN3 domain may bind to its target protein at least 5-fold above the signal obtained for a negative control in a standard solution ELISA assay.

[00260] In some embodiments, the FN3 domain that binds or specifically binds its target protein comprises an initiator methionine (Met) linked to the N-terminus of the molecule. In some embodiments, the FN3 domain that binds or specifically binds to its target protein comprises a cysteine (Cys) linked to a C-terminus of the FN3 domain. The addition of the N-terminal Met and/or the C-terminal Cys may facilitate expression and/or conjugation of half-life extending molecules.

[00261] The FN3 domain can also contain cysteine substitutions, such as those that are described in U.S. Patent No. 10,196,446, which is hereby incorporated by reference in its entirety. Briefly, in some embodiments, the polypeptide comprising an FN3 domain can have an FN3 domain that has a residue substituted with a cysteine, which can be referred to as a cysteine engineered fibronectin type III (FN3) domain. In some embodiments, the FN3 domain comprises at least one cysteine substitution at a position selected from the group consisting of

residues 6, 8, 10, 11, 14, 15, 16, 20, 30, 34, 38, 40, 41, 45, 47, 48, 53, 54, 59, 60, 62, 64, 70, 88, 89, 90, 91, and 93 of the FN3 domain based on SEQ ID NO: 1

(LPAPKNLVVSEVTEDSLRLSWTAPDAAFDSEKVGAINLTVPGSERSYDLTG LKPGTEYTVSIYGVKGGHRSNPLSAEFTT) of U.S. Patent No. 10,196,446, which is hereby incorporated by reference in its entirety, and the equivalent positions in related FN3 domains. A cysteine substitution at a position in the domain or protein comprises a replacement of the existing amino acid residue with a cysteine residue. Other examples of cysteine modifications can be found in, for example, U.S. Patent Application Publication No. 20170362301, which is hereby incorporated by reference in its entirety. The alignment of the sequences can be performed using BlastP using the default parameters at, for example, the NCBI website.

In some embodiments, the FN3 domain that binds to the target protein is internalized into a cell. In some embodiments, internalization of the FN3 domain may facilitate delivery of a detectable label or therapeutic into a cell. In some embodiments, internalization of the FN3 domain may facilitate delivery of a cytotoxic agent into a cell. The cytotoxic agent can act as a therapeutic agent. In some embodiments, internalization of the FN3 domain may facilitate the delivery of any detectable label, therapeutic, and/or cytotoxic agent disclosed herein into a cell. In some embodiments, the cell is a tumor cell. In some embodiments, the cell is a liver cell, a lung cell, muscle cell, an immune cell, a dendritic cell, a cell of the CNS, or a heart cell. In some embodiments, the therapeutic is a siRNA molecule as provided for herein. The FN3 domains that bind CD71 conjugated to a detectable label can be used to evaluate expression of CD71 on samples such as tumor tissue *in vivo* or *in vitro*. The FN3 domains that bind CD71 conjugated to a detectable label can be used to evaluate expression of CD71 on samples blood, immune cells, muscle cells, or dendritic cells *in vivo* or *in vitro*.

[00262] As provided herein, the different FN3 domains that are linked to the siRNA molecule can also be conjugated or linked to another FN3 domain that binds to a different target. This would enable the molecule to be multi-specific (e.g. bi-specific, tri-specific, etc.), such that it binds to a first target and another, for example, target. In some embodiments, the first FN3 binding domain is linked to another FN3 domain that binds to an antigen expressed by a tumor cell (tumor antigen).

[00263] In some embodiments, FN3 domains can be linked together by a linker to form a bivalent FN3 domain. The linker can be a flexible linker. In some embodiments, the linker is a G/S linker. In some embodiments the linker has 1, 2, 3, or 4 G/S repeats. A G/S repeat unit is four glycines followed by a serine, e.g. GGGGS. Other examples of linkers are provided herein and can also be used.

[00264] In some embodiments, the linker is a polypeptide of (GS)₂ (SEQ ID NO:720), (GGGS)₂ (SEQ ID NO:721), (GGGS)₅ (SEQ ID NO:722), (AP)₂₋₂₀, (AP)₂ (SEQ ID NO:723), (AP)₅ (SEQ ID NO:724), (AP)₁₀ (SEQ ID NO:725), (AP)₂₀ (SEQ ID NO:726) and A(EAAAK)₅AAA (SEQ ID NO:727) or (EAAAK)₁₋₅ (SEQ ID NO: 728). These are non-limiting examples and other linkers can also be used. The number of GGGGS or GGGGA repeats can also be 1, 2, 3, 4, or 5. In some embodiments, the linker comprises one or more GGGGS repeats and one or more GGGGA repeats. In some embodiments, the linker comprises one or more GGGGS repeats and one or more EAAAK repeats. In some embodiments, the linker comprises one or more GGGGS repeats and one or more "AP" repeats. In some embodiments, the linker comprises EAAAKEAAAKEAAAKEAAK (SEQ ID NO: 729); GGGSGGGSGGGSGGGGS (SEQ ID NO: 730); APAPAPAPAP (SEQ ID NO: 731); or EAAAK (SEQ ID NO: 732).

[00265] Without being bound to any particular theory, in some embodiments, the FN3 domains that are linked to the nucleic acid molecule may be used in the targeted delivery of the therapeutic agent to cells that express the binding partner of the one or more FN3 domains (*e.g.* tumor cells), and lead intracellular accumulation of the nucleic acid molecule therein. This can allow the siRNA molecule to properly interact with the cell machinery to inhibit the expression of the target gene, improve efficacy, and also avoid, in some embodiments, toxicity that may arise with untargeted administration of the same siRNA molecule.

[00266] The FN3 domain described herein that bind to their specific target protein may be generated as monomers, dimers, or multimers, for example, as a means to increase the valency and thus the avidity of target molecule binding, or to generate bi- or multispecific scaffolds simultaneously binding two or more different target molecules. The dimers and multimers may be generated by linking monospecific, bi- or multispecific protein scaffolds, for example, by the inclusion of an amino acid linker, for example a linker containing poly-glycine, glycine and serine, or alanine and proline. Exemplary linker include (GS)₂ (SEQ ID NO:720), (GGGS)₂ (SEQ ID NO:721), (GGGS)₅ (SEQ ID NO:722), (AP)₂₋₂₀, (AP)₂ (SEQ ID NO:723), (AP)₅ (SEQ ID NO:724), (AP)₁₀ (SEQ ID NO:725), (AP)₂₀ (SEQ ID NO:726) and A(EAAAK)₅AAA (SEQ ID NO:727) or (EAAAK)₁₋₅ (SEQ ID NO: 728). In some embodiments, the linker comprises or is an amino acid sequence of: EAAAKEAAAKEAAAKEAAK (SEQ ID NO: 729); GGGSGGGSGGGSGGGGS (SEQ ID NO: 730); APAPAPAPAP (SEQ ID NO: 731); or EAAAK (SEQ ID NO: 732).

[00267] The dimers and multimers may be linked to each other in a N-to C-direction. The use of naturally occurring as well as synthetic peptide linkers to connect polypeptides into

novel linked fusion polypeptides is well known in the literature (Hallewell *et al.*, *J Biol Chem* 264, 5260-5268, 1989; Alfthan *et al.*, *Protein Eng.* 8, 725-731, 1995; Robinson & Sauer, *Biochemistry* 35, 109-116, 1996; U.S. Pat. No. 5,856,456). The linkers described in this paragraph may be also be used to link the domains provided in the formula provided herein and above.

[00268] Half-life extending moieties

[00269] The FN3 domains may also, in some embodiments, incorporate other subunits for example via covalent interaction. In some embodiments, the FN3 domains that further comprise a half-life extending moiety. Exemplary half-life extending moieties are albumin, albumin variants, albumin-binding proteins and/or domains, transferrin and fragments and analogues thereof, and Fc regions. Amino acid sequences of the human Fc regions are well known, and include IgG1, IgG2, IgG3, IgG4, IgM, IgA and IgE Fc regions. In some embodiments, the FN3 domain binds to albumin, albumin variants, albumin-binding proteins and/or domains, and fragments and analogues thereof. extending the half-life of the entire molecule.

[00270] In some embodiments, the albumin binding domain comprises the amino acid sequence of SEQ ID NOs: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119. In some embodiments, the albumin binding domain (protein) is isolated. In some embodiments, the albumin binding domain comprises an amino acid sequence that is at least, or is, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence of SEQ ID NO: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119. In some embodiments, the albumin binding domain comprises an amino acid sequence that is at least, or is, 85%, 86%, 87%, 88%, 89%, 90%, 901%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence of SEQ ID NO: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119 provided that the protein has a substitution that corresponds to position 10 of SEQ ID NO: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119. In some embodiments, the substitution is A10V. In some embodiments, the substitution is A10G, A10L, A10I, A10T, or A10S. In some embodiments, the substitution at position 10 is any naturally occurring amino acid. In some embodiments, the isolated albumin binding domain comprises an amino acid sequence that has 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14 substitutions when compared to the amino acid sequence of SEQ ID NOs: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119. In some embodiments, the substitution is at a position that corresponds to position 10 of SEQ ID NOs: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118,

or 119. In some embodiments, FN3 domains provided comprises a cysteine residue in at least one residue position corresponding to residue positions 6, 11, 22, 25, 26, 52, 53, 61, 88 or positions 6, 8, 10, 11, 14, 15, 16, 20, 30, 34, 38, 40, 41, 45, 47, 48, 53, 54, 59, 60, 62, 64, 70, 88, 89, 90, 91, or 93 of SEQ ID NO: 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, or 119, or at a C-terminus. Although the positions are listed in a series, each position can also be chosen individually. In some embodiments, the cysteine is at a position that corresponds to position 6, 53, or 88. In some embodiments, additional examples of albumin binding domains can be found in U.S. Patent No. 10,925,932, which hereby incorporated by reference.

[00271] All or a portion of an antibody constant region may be attached to the FN3 domain to impart antibody-like properties, especially those properties associated with the Fc region, such as Fc effector functions such as C1q binding, complement dependent cytotoxicity (CDC), Fc receptor binding, antibody-dependent cell-mediated cytotoxicity (ADCC), phagocytosis, down regulation of cell surface receptors (e.g., B cell receptor; BCR), and may be further modified by modifying residues in the Fc responsible for these activities (for review; see Strohl, *Curr Opin Biotechnol.* 20, 685-691, 2009).

[00272] Additional moieties may be incorporated into the FN3 domains such as polyethylene glycol (PEG) molecules, such as PEG5000 or PEG20,000, fatty acids and fatty acid esters of different chain lengths, for example laurate, myristate, stearate, arachidate, behenate, oleate, arachidonate, octanedioic acid, tetradecanedioic acid, octadecanedioic acid, docosanedioic acid, and the like, polylysine, octane, carbohydrates (dextran, cellulose, oligo- or polysaccharides) for desired properties. These moieties may be direct fusions with the protein scaffold coding sequences and may be generated by standard cloning and expression techniques. Alternatively, well known chemical coupling methods may be used to attach the moieties to recombinantly produced molecules disclosed herein.

[00273] A PEG moiety may for example be added to the FN3 domain t by incorporating a cysteine residue to the C-terminus of the molecule, or engineering cysteines into residue positions that face away from the binding face of the molecule, and attaching a PEG group to the cysteine using well known methods.

[00274] FN3 domains incorporating additional moieties may be compared for functionality by several well-known assays. For example, altered properties due to incorporation of Fc domains and/or Fc domain variants may be assayed in Fc receptor binding assays using soluble forms of the receptors, such as the Fc γ RI, Fc γ RII, Fc γ RIII or FcRn receptors, or using

well known cell-based assays measuring for example ADCC or CDC, or evaluating pharmacokinetic properties of the molecules disclosed herein in *in vivo* models.

[00275] The compositions provided herein can be prepared by preparing the FN3 proteins and the nucleic acid molecules and linking them together. The techniques for linking the proteins to a nucleic acid molecule are known and any method can be used. For example, in some embodiments, the nucleic acid molecule is modified with a linker, such as the linker provided herein, and then the protein is mixed with the nucleic acid molecule comprising the linker to form the composition. For example, in some embodiments, a FN3 domains is conjugated to a siRNA a cysteine using thiol-maleimide chemistry. In some embodiments, a cysteine-containing FN3 domain can be reduced in, for example, phosphate buffered saline (or any other appropriate buffer) with a reducing agent (*e.g.* tris(2-carboxyethyl) phosphine (TCEP)) to yield a free thiol. Then, in some embodiments, the free thiol containing FN3 domain was mixed with a maleimide linked-modified siRNA duplex and incubated under conditions to form the linked complex. In some embodiments, the mixture is incubated for 0-5 hr, or about 1, 2, 3, 4 or 5 hr at RT. The reaction can be, for example, quenched with N-ethyl maleimide. In some embodiments, the conjugates can be purified using affinity chromatography and ion exchange. Other methods can also be used and this is simply one non-limiting embodiment.

[00276] Methods of making FN3 proteins are known and any method can be used to produce the protein. Examples are provided in the references incorporated by reference herein.

[00277] In some embodiments, the FN3 domain specifically binding CD71 comprises the amino acid sequence of SEQ ID NOs: 301-301, 310, 312-519, 521-572, 592-599, or 708-710, wherein a histidine tag has been appended to the N-terminal or C-terminal end of the polypeptide for ease of purification. In some embodiments, the histidine tag (His-tag) comprises six histidine residues. In further embodiments, the His-tag to connected to the FN3 domain by at least one glycine residue or about 2 to about 4 glycine residues. Accordingly, after purification of the FN3 domain and cleavage of the His-tag from the polypeptide one or more glycine may be left on the N-terminus or C-terminus. In some embodiments, if the His-tag is removed from the N-terminus all of the glycines are removed. In some embodiments, if the His-tag is removed from the C-terminus one or more of the glycines are retained.

[00278] In some embodiments, the FN3 domain specifically binding CD71 comprises the amino acid sequence of SEQ ID NOs: 301-301, 310, 312-519, 521-572, 592-599, or 708-710, wherein the N-terminal methionine is retained after purification of the FN3 domain.

[00279] Kits

[00280] In some embodiments, a kit comprising the compositions described herein are provided.

[00281] The kit may be used for therapeutic uses and as a diagnostic kit.

[00282] In some embodiments, the kit comprises the FN3 domain conjugated to the nucleic acid molecule.

[00283] **Uses of the Conjugates FN3 domains**

[00284] The compositions provided for herein may be used to diagnose, monitor, modulate, treat, alleviate, help prevent the incidence of, or reduce the symptoms of human disease or specific pathologies in cells, tissues, organs, fluid, or, generally, a host.

[00285] In some embodiments, methods of selectively reducing GYS1 mRNA and protein in skeletal muscle. In certain embodiments, GYS1 mRNA and protein is not reduced in the liver and/or the kidney.

[00286] In some embodiments, the reduction in the GYS1 mRNA and protein is sustained for about 1 week, about 2 weeks, about 3 weeks, about 4 weeks, about 5 weeks, or greater than 5 weeks after administration of the conjugate described herein.

[00287] In some embodiments, the FN3 domain can facilitate delivery into CD71 positive tissues (eg. skeletal muscle, smooth muscle) for treatment of muscle diseases.

[00288] In some embodiments, the FN3 domain can facilitate delivery to activated lymphocytes, dendritic cells, or other immune cells for treatment of immunological diseases.

[00289] In some embodiments, the polypeptide that binds to CD71 is directed to immune cells. In some embodiments, the polypeptide that binds to CD71 is directed to dendritic cells. In some embodiments, methods of treating an autoimmune disease in a subject in need thereof are provided. In some embodiments, the methods comprise administering to the subject a polypeptide or the pharmaceutical composition that binds to CD71. In some embodiments, that the polypeptide is a FN3 domain that binds to CD71. In some embodiments, the polypeptide comprises a sequence such as SEQ ID Nos: 301-301, 310, 312-519, 521-572, 592-599, or 708-710, or a polypeptide as provided herein that is linked to or conjugated to a therapeutic agent. In some embodiments, the autoimmune disease is selected from the group consisting of rheumatoid arthritis, Hashimoto's autoimmune thyroiditis, celiac disease, diabetes mellitus type 1, vitiligo, rheumatic fever, pernicious anemia/atrophic gastritis, alopecia areata, and immune thrombocytopenic purpura.

[00290] In some embodiments, a method of treating a subject having Pompe Disease (GSD2, acid alpha-glucosidase (GAA) deficiency) is provided, the method comprising administering to the subject a composition provided for herein. In some embodiments, the

methods comprise administering to the subject a polypeptide or the pharmaceutical composition that binds to CD71. In some embodiments, that the polypeptide is a FN3 domain that binds to CD71. In some embodiments, the polypeptide comprises a sequence such as SEQ ID Nos: 301-301, 310, 312-519, 521-572, 592-599, or 708-710, or a polypeptide as provided herein that is linked to or conjugated to a therapeutic agent.

[00291] In some embodiments, methods of treating glycogen storage disease in a subject in need thereof, the method comprising administering a composition provided herein are provided. In some embodiments, the glycogen storage disease is selected from the group consisting of Cori's disease or Forbes' disease (GSD3, Glycogen debranching enzyme (AGL) deficiency), McArdle disease (GSD5, Muscle glycogen phosphorylase (PYGM) deficiency), type II Diabetes/diabetic nephropathy, Aldolase A Deficiency GSD12, Lafora Disease, hypoxia, Andersen disease (GSD4, Glycogen debranching enzyme (GBE1) deficiency), Tarui's Disease (GSD7, Muscle phosphofructokinase (PFKM) deficiency), and adult polyglucosan body disease. In some embodiments, the glycogen storage disease is selected from the group consisting of Glycogen synthase (GYS2) deficiency (GSD0), Glucose-6-phosphatase (G6PC / SLC37A4) deficiency (GSD1, von Gierke's disease), Hers' disease (GSD6, Liver glycogen phosphorylase (PYGL) or Muscle phosphoglycerate mutase (PGAM2) deficiency), Phosphorylase kinase (PHKA2 / PHKB / PHKG2 / PHKA1) deficiency (GSD9), Phosphoglycerate mutase (PGAM2) deficiency (GSD10), Muscle lactate dehydrogenase (LDHA) deficiency (GSD11), Fanconi-Bickel syndrome (GSD 11, Glucose transporter (GLUT2) deficiency, Aldolase A deficiency (GSD 12), β -enolase (ENO3) deficiency (GSD13), and Glycogenin-1 (GYG1) deficiency (GSD15).

[00292] In some embodiments, a use of a composition as provided herein or of any of in the preparation of a pharmaceutical composition or medicament for treating cancer are provided. In some embodiments, the cancer is selected from the group consisting of acute myeloid leukemia, myelodysplastic syndromes, gastric cancer, clear cell renal cell carcinoma, clear cell carcinomas of the breast, clear cell carcinomas of the endometrium, clear cell carcinomas of the ovary, clear cell carcinomas of the uterus, hepatocellular carcinoma, pancreatic cancer, prostate cancer, soft tissue cancer, Ewings sarcoma, and non-small cell lung cancer

[00293] In some embodiments, the CD71 cell is a cell involved in a CNS diseases, inflammatory/immune diseases, such as MS & infectious diseases of the brain. In some embodiments, the polypeptide that binds to CD71 is directed to the central nervous system. In some embodiments, methods of treating a neurological condition and/or a brain tumor in a subject in need thereof are provided. In some embodiments, the methods comprise administering

to the subject a polypeptide or the pharmaceutical composition that binds to CD71. In some embodiments, that the polypeptide is a FN3 domain that binds to CD71. In some embodiments, the polypeptide comprises a sequence such as SEQ ID Nos: 301-301, 310, 312-519, 521-572, 592-599, or 708-710, or a polypeptide as provided herein that is linked to or conjugated to a therapeutic agent. In some embodiments, the brain tumor is selected from the group consisting of nonmalignant, benign, and malignant brain tumors. In some embodiments, the neurological condition is selected from the group consisting of Alzheimer's Disease, Amyotrophic Lateral Sclerosis, Parkinson's Disease, Lafora Disease, Pompe Disease, adult polyglucosan body disease, stroke, spinal cord injury, ataxia, Bell's Palsy, cerebral aneurysm, epilepsy, seizures, Guillain-Barre Syndrome, multiple sclerosis, muscular dystrophy, neurocutaneous syndromes, migraine, encephalitis, septicemia, and myasthenia gravis.

[00294] In some embodiments, a method of treating a subject having cancer is provided, the method comprising administering to the subject a composition provided for herein. In some embodiments, the method comprises administering to the subject a polypeptide or the pharmaceutical composition that binds to CD71. In some embodiments, that the polypeptide is a FN3 domain that binds to CD71. In some embodiments, the polypeptide comprises a sequence such as SEQ ID Nos: 301-301, 310, 312-519, 521-572, 592-599, or 708-710, or a polypeptide as provided herein that is linked to or conjugated to a therapeutic agent.

[00295] In some embodiments, the subject has a solid tumor.

[00296] In some embodiments, the solid tumor is a melanoma.

[00297] In some embodiments, the solid tumor is a lung cancer. In some embodiments, the solid tumor is a non-small cell lung cancer (NSCLC). In some embodiments, the solid tumor is a squamous non-small cell lung cancer (NSCLC). In some embodiments, the solid tumor is a non-squamous NSCLC. In some embodiments, the solid tumor is a lung adenocarcinoma.

[00298] In some embodiments, the solid tumor is a renal cell carcinoma (RCC).

[00299] In some embodiments, the solid tumor is a mesothelioma.

[00300] In some embodiments, the solid tumor is a nasopharyngeal carcinoma (NPC).

[00301] In some embodiments, the solid tumor is a colorectal cancer.

[00302] In some embodiments, the solid tumor is a prostate cancer. In some embodiments, the solid tumor is castration-resistant prostate cancer.

[00303] In some embodiments, the solid tumor is a stomach cancer.

[00304] In some embodiments, the solid tumor is an ovarian cancer.

[00305] In some embodiments, the solid tumor is a gastric cancer.

[00306] In some embodiments, the solid tumor is a liver cancer.

- [00307] In some embodiments, the solid tumor is pancreatic cancer.
- [00308] In some embodiments, the solid tumor is a thyroid cancer.
- [00309] In some embodiments, the solid tumor is a squamous cell carcinoma of the head and neck.
- [00310] In some embodiments, the solid tumor is a carcinomas of the esophagus or gastrointestinal tract.
- [00311] In some embodiments, the solid tumor is a breast cancer.
- [00312] In some embodiments, the solid tumor is a fallopian tube cancer.
- [00313] In some embodiments, the solid tumor is a brain cancer.
- [00314] In some embodiments, the solid tumor is an urethral cancer.
- [00315] In some embodiments, the solid tumor is a genitourinary cancer.
- [00316] In some embodiments, the solid tumor is an endometriosis.
- [00317] In some embodiments, the solid tumor is a cervical cancer.
- [00318] In some embodiments, the solid tumor is a metastatic lesion of the cancer.
- [00319] In some embodiments, the subject has a hematological malignancy.
- [00320] In some embodiments, the hematological malignancy is a lymphoma, a myeloma or a leukemia. In some embodiments, the hematological malignancy is a B cell lymphoma. In some embodiments, the hematological malignancy is Burkitt's lymphoma. In some embodiments, the hematological malignancy is Hodgkin's lymphoma. In some embodiments, the hematological malignancy is a non-Hodgkin's lymphoma.
- [00321] In some embodiments, the hematological malignancy is a myelodysplastic syndrome.
- [00322] In some embodiments, the hematological malignancy is an acute myeloid leukemia (AML). In some embodiments, the hematological malignancy is a chronic myeloid leukemia (CML). In some embodiments, the hematological malignancy is a chronic myelomonocytic leukemia (CMML).
- [00323] In some embodiments, the hematological malignancy is a multiple myeloma (MM).
- [00324] In some embodiments, the hematological malignancy is a plasmacytoma.
- [00325] In some embodiments, the cancer is a soft tissue cancer. In some embodiments, the soft tissue cancer is Ewings sarcoma.
- [00326] In some embodiments, methods of treating cancer in a subject in need thereof are provided. In some embodiments, the methods comprise administering to the subject any composition provided herein. In some embodiments, a use of a composition as provided herein

are provided in the preparation of a pharmaceutical composition or medicament for treating cancer. In some embodiments, the composition can be used for treating cancer.

[00327] In some embodiments, methods of treating Pompe Disease (GSD2, acid alpha-glucosidase (GAA) deficiency) in a subject in need thereof are provided. In some embodiments, the methods comprise administering to the subject any composition provided herein. In some embodiments, a use of a composition as provided herein are provided in the preparation of a pharmaceutical composition or medicament for treating Pompe Disease (GSD2, acid alpha-glucosidase (GAA) deficiency). In some embodiments, the composition can be used for treating Pompe Disease (GSD2, acid alpha-glucosidase (GAA) deficiency).

[00328] In some embodiments, methods of treating glycogen storage disease in a subject in need thereof are provided. In some embodiments, the methods comprise administering to the subject any composition provided herein. In some embodiments, a use of a composition as provided herein are provided in the preparation of a pharmaceutical composition or medicament for treating glycogen storage disease. In some embodiments, the composition can be used for treating glycogen storage disease.

[00329] In some embodiments, methods of treating glycogen storage disease in a subject in need thereof, the method comprising administering a composition provided herein are provided. In some embodiments, the glycogen storage disease is selected from the group consisting of Cori's disease or Forbes' disease (GSD3, Glycogen debranching enzyme (AGL) deficiency), McArdle disease (GSD5, Muscle glycogen phosphorylase (PYGM) deficiency), type II Diabetes/diabetic nephropathy, Aldolase A Deficiency GSD12, Lafora Disease, hypoxia, Andersen disease (GSD4, Glycogen debranching enzyme (GBE1) deficiency), Tarui's Disease (GSD7, Muscle phosphofructokinase (PFKM) deficiency), and adult polyglucosan body disease. In some embodiments, the glycogen storage disease is selected from the group consisting of Glycogen synthase (GYS2) deficiency (GSD0), Glucose-6-phosphatase (G6PC / SLC37A4) deficiency (GSD1, von Gierke's disease), Hers' disease (GSD6, Liver glycogen phosphorylase (PYGL) or Muscle phosphoglycerate mutase (PGAM2) deficiency), Phosphorylase kinase (PHKA2 / PHKB / PHKG2 / PHKA1) deficiency (GSD9), Phosphoglycerate mutase (PGAM2) deficiency (GSD10), Muscle lactate dehydrogenase (LDHA) deficiency (GSD11), Fanconi-Bickel syndrome (GSD 11, Glucose transporter (GLUT2) deficiency, Aldolase A deficiency (GSD 12), β -enolase (ENO3) deficiency (GSD13), and Glycogenin-1 (GYG1) deficiency (GSD15).

[00330] In some embodiments, the polypeptide that binds to CD71 is directed to the central nervous system. In some embodiments, methods of treating a neurological condition

and/or a brain tumor in a subject in need thereof are provided. In some embodiments, the methods comprise administering to the subject a polypeptide or the pharmaceutical composition that binds to CD71. In some embodiments, that the polypeptide is a FN3 domain that binds to CD71. In some embodiments, the polypeptide comprises a sequence such as SEQ ID Nos: 301-301, 310, 312-519, 521-572, 592-599, or 708-710, or a polypeptide as provided herein that is linked to or conjugated to a therapeutic agent. In some embodiments, the brain tumor is selected from the group consisting of nonmalignant, benign, and malignant brain tumors. In some embodiments, the neurological condition is selected from the group consisting of Alzheimer's Disease, Amyotrophic Lateral Sclerosis, Parkinson's Disease, Lafora Disease, Pompe Disease, adult polyglucosan body disease, stroke, spinal cord injury, ataxia, Bell's Palsy, cerebral aneurysm, epilepsy, seizures, Guillain-Barre Syndrome, multiple sclerosis, muscular dystrophy, neurocutaneous syndromes, migraine, encephalitis, septicemia, and myasthenia gravis. In some embodiments, a method of treating a neurological condition and/or a brain tumor in a subject, the method comprising administering to the subject a FN3 domain that binds CD71 and the FN3 domain is conjugated to a therapeutic agent (*e.g.* cytotoxic agent, an oligonucleotide, such as a siRNA, ASO, and the like, a FN3 domain that binds to another target, and the like).

[00331] In some embodiments, methods of treating Pompe disease in a subject in need thereof are provided. In some embodiments, the methods comprise administering to the subject a polypeptide or the pharmaceutical composition that binds to CD71. In some embodiments, that the polypeptide is a FN3 domain that binds to CD71. In some embodiments, the polypeptide comprises a sequence such as SEQ ID Nos: 301-301, 310, 312-519, 521-572, 592-599, or 708-710, or a polypeptide as provided herein that is linked to or conjugated to a therapeutic agent. In some embodiments, a method of treating a Pompe disease in a subject, the method comprising administering to the subject a FN3 domain that binds CD71 and the FN3 domain is conjugated to a therapeutic agent (*e.g.* cytotoxic agent, an oligonucleotide, such as a siRNA, ASO, and the like, a FN3 domain that binds to another target, and the like).

[00332] In some embodiments, the polypeptide that binds to CD71 is directed to immune cells. In some embodiments, the polypeptide that binds to CD71 is directed to dendritic cells. In some embodiments, methods of treating an autoimmune disease in a subject in need thereof are provided. In some embodiments, the methods comprise administering to the subject a polypeptide or the pharmaceutical composition that binds to CD71. In some embodiments, that the polypeptide is a FN3 domain that binds to CD71. In some embodiments, the polypeptide comprises a sequence such as SEQ ID Nos: 301-301, 310, 312-519, 521-572, 592-599, or 708-710, or a polypeptide as provided herein that is linked to or conjugated to a therapeutic agent. In

some embodiments, the autoimmune disease is selected from the group consisting of rheumatoid arthritis, Hashimoto's autoimmune thyroiditis, celiac disease, diabetes mellitus type 1, vitiligo, rheumatic fever, pernicious anemia/atrophic gastritis, alopecia areata, and immune thrombocytopenic purpura. In some embodiments, a method of treating an autoimmune disease in a subject, the method comprising administering to the subject a FN3 domain that binds CD71 and the FN3 domain is conjugated to a therapeutic agent (*e.g.* cytotoxic agent, an oligonucleotide, such as a siRNA, ASO, and the like, a FN3 domain that binds to another target, and the like).

[00333] In some embodiments, methods of reducing the expression of a target gene in a cell are provided. In some embodiments, the methods comprise delivering to the cell with a composition or a pharmaceutical composition as provided herein. In some embodiments, the cell is *ex-vivo*. In some embodiments, the cell is *in-vivo*. In some embodiments, the target gene is GYS1. The target gene, however, can be any target gene as the evidence provided herein demonstrates that siRNA molecules can be delivered efficiently when conjugated to a FN3 domain. In some embodiments, the siRNA targeting GYS1 is linked to a FN3 domain. In some embodiments, the FN3 polypeptide (domain) is one that binds to CD71. In some embodiments, the FN3 polypeptide is as provided for herein or as provided for in PCT Application No. PCT/US20/55509, U.S. Application No. 17/070,337, PCT Application No. PCT/US20/55470, or U.S. Application No. 17/070,020, each of which is hereby incorporated by reference in its entirety. In some embodiments, the siRNA is not conjugated to a FN3 domain.

[00334] In some embodiments, methods of reducing the expression of a target gene in a cell are provided. In some embodiments, the methods comprise delivering to the cell with a composition or a pharmaceutical composition as provided herein. In some embodiments, the cell is *ex-vivo*. In some embodiments, the cell is *in-vivo*. In some embodiments, a method of reducing the expression of a target gene results in a reduction of about 99%, 90-99%, 50-90%, or 10-50% in the expression of the target gene.

[00335] In some embodiments, a method of reducing the expression of GYS1 is provided. In some embodiments, the reduced expression is the expression (amount) of GYS1 mRNA. In some embodiments, a method of reducing the expression of GYS1 results in a reduction of about 99%, 90-99%, 50-90%, or 10-50% in the expression of GYS1. In some embodiments, the reduced expression is the expression (amount) of GYS1 protein. In some embodiments, the reduced protein is glycogen. In some embodiments, reduction of glycogen occurs in muscle cells. In some embodiments, reduction of glycogen occurs in heart cells. In some embodiments, the method comprises delivering to a cell with a siRNA molecule as provided herein that targets

GYS1. In some embodiments, the siRNA is conjugated to a FN3 domain. In some embodiments, the FN3 domain is a FN3 domain that binds to CD71. In some embodiments, the FN3 domain is as provided for herein. In some embodiments, the FN3 domain is a dimer of two FN3 domains that bind to CD71. In some embodiments, the FN3 domains are the same. In some embodiments, the two FN3 domains are different, i.e., bind to different regions or amino acid residues of CD71, i.e. a different epitope. In some embodiments, the method comprises administering to a subject (patient) a GYS1 siRNA molecule, such as those provided herein. In some embodiments, the GYS1 siRNA administered to the subject is conjugated or linked to a FN3 domain. In some embodiments, the FN3 domain is a FN3 domain that binds to CD71. In some embodiments, the FN3 domain is as provided for herein. In some embodiments, the FN3 domain is a dimer of two FN3 domains that bind to CD71. In some embodiments, the FN3 domains are the same. In some embodiments, the two FN3 domains are different, i.e., bind to different regions or amino acid residues of CD71, i.e. a different epitope. In some embodiments, the CD71 binding domain is a polypeptide as provided for herein.

[00336] In some embodiments, methods of delivering a siRNA molecule to a cell in a subject are provided. In some embodiments, the methods comprise administering to the subject a pharmaceutical composition comprising a composition as provided for herein. In some embodiments, the cell is a CD71 positive cell. The term “positive cell” in reference to a protein refers to a cell that expresses the protein. In some embodiments, the protein is expressed on the cell surface. In some embodiments, the cell is a tumor cell, a liver cell, an immune cell, a dendritic cell, a heart cell, a muscle cell, a cell of the CNS, or a cell inside the blood brain barrier. In some embodiments, the siRNA downregulates the expression of a target gene in the cell. In some embodiments, the target gene is GYS1.

[00337] In some embodiments, the compositions provided herein may be used to diagnose, monitor, modulate, treat, alleviate, help prevent the incidence of, or reduce the symptoms of human disease or specific pathologies in cells, tissues, organs, fluid, or, generally, a host, also exhibit the property of being able to cross the blood brain barrier. The blood-brain barrier (BBB) prevents most macromolecules (e.g., DNA, RNA, and polypeptides) and many small molecules from entering the brain. The BBB is principally composed of specialized endothelial cells with highly restrictive tight junctions, consequently, passage of substances, small and large, from the blood into the central nervous system is controlled by the BBB. This structure makes treatment and management of patients with neurological diseases and disorders (e.g., brain cancer) difficult as many therapeutic agents cannot be delivered across the BBB with desirable efficiency. Additional conditions that involve disruptions of the BBB include: stroke,

diabetes, seizures, hypertensive encephalopathy, acquired immunodeficiency syndrome, traumatic brain injuries, multiple sclerosis, Lafora Disease, Pompe Disease, adult polyglucosan body disease, Parkinson's disease (PD) and Alzheimer disease. This ability is especially useful for treating brain cancers including for example: astrocytoma, medulloblastoma, glioma, ependymoma, germinoma (pinealoma), glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, and congenital tumors; or a cancer of the spinal cord, e.g., neurofibroma, meningioma, glioma, and sarcoma. This is further useful in treating Pompe Disease and/or glycogen storage disease. In certain embodiments, the compositions provided for herein can be used to deliver a therapeutic or cytotoxic agent, for example, across the blood brain barrier. In certain embodiments, the compositions provided for herein can be used to deliver a therapeutic or cytotoxic agent, for example, into the muscle.

[00338] In some embodiments, methods of treating glycogen storage disease in a subject in need thereof, the method comprising administering a composition provided herein are provided. In some embodiments, the glycogen storage disease is selected from the group consisting of Cori's disease or Forbes' disease (GSD3, Glycogen debranching enzyme (AGL) deficiency), McArdle disease (GSD5, Muscle glycogen phosphorylase (PYGM) deficiency), type II Diabetes/diabetic nephropathy, Aldolase A Deficiency GSD12, Lafora Disease, hypoxia, Andersen disease (GSD4, Glycogen debranching enzyme (GBE1) deficiency), Tarui's Disease (GSD7, Muscle phosphofructokinase (PFKM) deficiency), and adult polyglucosan body disease. In some embodiments, the glycogen storage disease is selected from the group consisting of Glycogen synthase (GYS2) deficiency (GSD0), Glucose-6-phosphatase (G6PC / SLC37A4) deficiency (GSD1, von Gierke's disease), Hers' disease (GSD6, Liver glycogen phosphorylase (PYGL) or Muscle phosphoglycerate mutase (PGAM2) deficiency), Phosphorylase kinase (PHKA2 / PHKB / PHKG2 / PHKA1) deficiency (GSD9), Phosphoglycerate mutase (PGAM2) deficiency (GSD10), Muscle lactate dehydrogenase (LDHA) deficiency (GSD11), Fanconi-Bickel syndrome (GSD 11, Glucose transporter (GLUT2) deficiency, Aldolase A deficiency (GSD 12), β -enolase (ENO3) deficiency (GSD13), and Glycogenin-1 (GYG1) deficiency (GSD15).

[00339] In some embodiments, the compositions or pharmaceutical compositions provided herein can be used to treat muscle diseases, such as muscular dystrophy, DMD, and the like.

[00340] In some embodiments, the compositions or pharmaceutical compositions provided herein may be administered alone or in combination with other therapeutics, that is, simultaneously or sequentially. In some embodiments, the other or additional therapeutics are other anti-tumor agent or therapeutics. Different tumor types and stages of tumors can require the

use of various auxiliary compounds useful for treatment of cancer. For example, the compositions provided herein can be used in combination with various chemotherapeutics such as taxol, tyrosine kinase inhibitors, leucovorin, fluorouracil, irinotecan, phosphatase inhibitors, MEK inhibitors, among others. The composition may also be used in combination with drugs which modulate the immune response to the tumor such as anti-PD-1 or anti-CTLA-4, among others. Additional treatments can be agents that modulate the immune system, such antibodies that target PD-1 or PD-L1.

[00341] In some embodiments, the compositions or pharmaceutical compositions provided herein may be administered in combination with GAA enzyme replacement therapy (ERT).

[00342] “Treat” or “treatment” refers to the therapeutic treatment and prophylactic measures, wherein the object is to prevent or slow down (lessen) an undesired physiological change or disorder, such as the development or spread of cancer. In some embodiments, beneficial or desired clinical results include, but are not limited to, alleviation of symptoms, diminishment of extent of disease, stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable. “Treatment” can also mean prolonging survival as compared to expected survival if not receiving treatment. Those in need of treatment include those already with the condition or disorder as well as those prone to have the condition or disorder or those in which the condition or disorder is to be prevented.

[00343] A “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired therapeutic result. A therapeutically effective amount of the compositions provided herein may vary according to factors such as the disease state, age, sex, and weight of the individual. Exemplary indicators of an effective amount is improved well-being of the patient, decrease or shrinkage of the size of a tumor, arrested or slowed growth of a tumor, and/or absence of metastasis of cancer cells to other locations in the body.

[00344] Administration/ Pharmaceutical Compositions

[00345] In some embodiments, pharmaceutical compositions of the compositions provided herein and a pharmaceutically acceptable carrier, are provided. For therapeutic use, the compositions may be prepared as pharmaceutical compositions containing an effective amount of the domain or molecule as an active ingredient in a pharmaceutically acceptable carrier. “Carrier” refers to a diluent, adjuvant, excipient, or vehicle with which the active compound is administered. Such vehicles can be liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and

the like. For example, 0.4% saline and 0.3% glycine can be used. These solutions are sterile and generally free of particulate matter. They may be sterilized by conventional, well-known sterilization techniques (*e.g.*, filtration). The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, stabilizing, thickening, lubricating and coloring agents, etc. The concentration of the molecules disclosed herein in such pharmaceutical formulation can vary widely, *i.e.*, from less than about 0.5%, usually at least about 1% to as much as 15 or 20% by weight and will be selected primarily based on required dose, fluid volumes, viscosities, etc., according to the particular mode of administration selected. Suitable vehicles and formulations, inclusive of other human proteins, *e.g.*, human serum albumin, are described, for example, in *e.g.* Remington: The Science and Practice of Pharmacy, 21st Edition, Troy, D.B. ed., Lipincott Williams and Wilkins, Philadelphia, PA 2006, Part 5, Pharmaceutical Manufacturing pp 691-1092, See especially pp. 958-989.

[00346] The mode of administration for therapeutic use of the compositions disclosed herein may be any suitable route that delivers the agent to the host, such as parenteral administration, *e.g.*, intradermal, intramuscular, intraperitoneal, intravenous or subcutaneous, pulmonary; transmucosal (oral, intranasal, intravaginal, rectal), using a formulation in a tablet, capsule, solution, powder, gel, particle; and contained in a syringe, an implanted device, osmotic pump, cartridge, micropump; or other means appreciated by the skilled artisan, as well known in the art. Site specific administration may be achieved by for example intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelial, intracerebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intracardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravascular, intravesical, intralesional, vaginal, rectal, buccal, sublingual, intranasal, or transdermal delivery.

[00347] Pharmaceutical compositions can be supplied as a kit comprising a container that comprises the pharmaceutical composition as described herein. A pharmaceutical composition can be provided, for example, in the form of an injectable solution for single or multiple doses, or as a sterile powder that will be reconstituted before injection. Alternatively, such a kit can include a dry-powder disperser, liquid aerosol generator, or nebulizer for administration of a pharmaceutical composition. Such a kit can further comprise written information on indications and usage of the pharmaceutical composition.

EXAMPLES

[00348] The following examples are illustrative of the embodiments disclosed herein. These examples are provided for the purpose of illustration only and the embodiments should in no way be construed as being limited to these examples, but rather should be construed to encompass any and all variations which become evidence as a result of the teaching provided herein. Those of skill in the art will readily recognize a variety of non-critical parameters that could be changed or modified to yield essentially similar results.

[00349] EXAMPLE 1: GYS1 siRNA sequence ID and characterization

[00350] siRNA In Silico Screening: In silico siRNA screening was performed to identify a human siRNA complementary to the human *GYS1* mRNA, see FIG. 1. All possible 19-mer antisense sequences were generated from the human *GYS1* siRNA sequence (NM_001161587) and each 19-mer was assessed for activity against other human *GYS1* isoforms as well as potential cross-reactivity mouse, rat, and *Cynomolgus macaque*. Human siRNA target sites were assessed for common human SNPs (MAF > 1%) using dbSNP (b155 v2). Sequences that targeted a common allele were discarded. Next, siRNA off-target genes were assessed for sense and antisense strands in all relevant model organisms. This selection yielded 200 potential candidates, that were further defined by and *in vitro* knockdown screen. Lead siRNA candidates are in Table 1A and 1B. GYS1 siRNA linkers for conjugation to cysteine engineered Centyrins were prepared as described in Table 3.

[00351] HEK293T cells were lipofected with 10nM ABXO-HHH (siRNA Pair HHH) for 24 hours alongside untreated control cells (6 replicates per treatment group). mRNA was polyA+ selected from cells and subjected to unstranded 2x150bp paired end sequencing on the Illumina HiSeq platform to an average depth of > 30 million reads.

[00352] FIG. 2 Volcano plot showing the DESeq2 results of ABXO-HHH versus untreated cells. The X axis represents the log₂ fold change in gene expression (ABXO-HHH divided by untreated) for all genes. The Y axis shows the negative log₁₀ transformed adjusted P value. Colored dots represent genes with significant complementarity to ABXO-HHH detected via BLAST.

[00353] RNA-seq library quality was inspected using FastQC. Libraries were pseudo aligned to the GrCH38 human transcriptome using Kallisto. We achieved exceptional read alignment with an average of > 90% of reads aligning to the transcriptome. Differential expression was then assessed using DESeq2 comparing ABXO-HHH treated cells to lipofectamine only treated cells.

[00354] DESeq results showed substantial and significant knockdown of the *GYS1* target. *GYS1* expression in ABXO-HHH treated samples was downregulated to 31% of untreated samples (FIG. 2). *GYS1* was the most downregulated protein coding gene for all genes that were differentially expressed (DE) under ABXO-HHH treatment. *GYS1* was also by far the DE gene with the lowest p-value, with an adjusted p-value of $1e-87.7$, over 30 orders of magnitude smaller than the next lowest adjusted p-value.

[00355] In silico prediction of potential off-target effects was performed using BLAST. The ABXO-HHH sense and antisense sequences were BLAST aligned against an internal BLAST database built from the GRCh38 transcriptome. Of all the potential off-target effects identified by BLAST, only RAP2C was significantly downregulated in the presence of ABXO-HHH (FIG. 2) and failed to surpass the DE threshold of $\log_2$ fold change < -1 .

[00356] Together these data demonstrate that ABXO-HHH is a highly specific siRNA even at relatively high concentrations in the human transcriptome.

[00357] Oligonucleotide synthesis, deprotection and annealing protocol:

[00358] Synthesis of oligonucleotides was performed on Mermade 12 synthesizer using standard phosphoramidite chemistry on 500Å controlled pore glass (CPG) with phosphoramidites at a 0.1 M concentration in acetonitrile. I2 in THF/pyridine/water (0.02M) was the oxidizing agent with 0.6 M ETT (5-ethylthiotetrazole) as the activation agent. N,N-dimethyl-N'-(3-thioxo-3H-1,2,4-dithiazol-5-yl)methanimidamide (DDTT), 0.09 M in pyridine, was used as the sulfurizing reagent for the introduction of phosphorothioate (PS) bonds. 3% (v/v) dichloroacetic acid in dichloromethane was used the deblocking solution. All single strands without maleimide were purified by ion exchange chromatography (IEX) with 20 mM phosphate at pH 8.5 as Buffer A and 20 mM phosphate pH 8.5 and 1 M sodium bromide as Buffer B. After purification, the oligonucleotide fractions were pooled, concentrated, and desalted. Desalted samples were then lyophilized to dryness and stored at -20°C .

[00359] Deprotection of antisense strands

[00360] After synthesis, the support was washed with acetonitrile (ACN) and dried in the column under vacuum and transferred into a 1 mL screw cap that could be tightly sealed and shaken with a solution of 5% diethylamine in aqueous ammonia at 65°C for 5 h. Cleavage and deprotection of crude oligo was checked by liquid chromatography-mass spectrometry (LC-MS) and was subsequently purified by IEX-HPLC.

[00361] Synthesis and deprotection of maleimide-containing oligonucleotides

[00362] Maleimide containing oligos were made using either a 3' amino-modified CPG solid support or a 5' amino modifier phosphoramidite. The support was transferred into a 1 mL

vial that could be tightly sealed and incubated with 50/50 v/v 40% aqueous methyl amine and aqueous ammonia (AMA) at room temperature for 2 h or 65°C for 10 minutes to cleave and deprotect. The single strand was purified by IEX chromatography and desalted the same conditions as the antisense before maleimide addition.

[00363] Approximately 20 mg/mL of the amine-modified sense strand in 0.05 M phosphate buffer at pH 7.1 was made to which 10 equivalents of the maleimide N-hydroxysuccinimide (NHS) ester, dissolved in ACN was added. The NHS ester solution was added to the aqueous oligonucleotide solution and shaken for 3 h at room temperature. The now maleimide conjugated oligonucleotides were purified by reverse-phase chromatography (20 mM triethyl ammonium acetate with 80% acetonitrile in Buffer B) to prevent the maleimide hydrolysis under ion exchange buffer conditions.

[00364] After purification, the oligonucleotide fractions were pooled, concentrated, and desalted.

[00365] To avoid hydrolysis of maleimide, duplexing of the sense and antisense strands was performed via freeze-drying using equimolar amounts of each desalted single strand.

[00366] Centyrin conjugation to siRNA, conjugate purification and analysis: Centyrins were conjugated to siRNA through cysteine-specific chemistry via maleimide. Cysteine-containing Centyrin in PBS at 50-200 μ M were reduced with 10 mM tris(2-carboxyethyl) phosphine (TCEP) at room temperature (30 mins) to yield a free thiol. The free thiol containing Centyrin is then mixed with maleimide containing siRNA duplex in water immediately prior at a molar ratio of ~1.5:1 Centyrin:siRNA. After 2 hr incubation at RT or 37°C, reaction was quenched with N-ethyl maleimide (1mM final NEM concentration in the reaction mixture) The conjugate was purified in two steps. Step I either immobilized metal affinity chromatography (for tagged proteins) or Hydrophobic interaction chromatography (for tagless proteins); to remove un-reacted SiRNA linker. Step II-Capto-DEAE; to remove un-reacted centyrin. Fractions containing conjugate were pooled, exchanged into HBS by desalting using dialysis, and concentrated if necessary.

[00367] Analytical Characterization of Centyrin-siRNA conjugates: Centyrin -siRNA conjugates were characterized by a combination of analytical techniques. SDS-PAGE was used to compare amounts of conjugate to free protein. For SDS-PAGE, 4-20% Mini-PROTEAN® TGX Stain-Free™ Protein Gels (BioRad) were run in SDS buffer for one hour at 100 V. Gels were visualized under UV light. Analytical SEC (Superdex-75 5/150 GL column-GE) was used to analyze purity and aggregation state of Centyrin-siRNA conjugates. Liquid chromatography/mass spectrometry (LCMS) was used to confirm identity and purity of the

conjugates. Samples were analyzed using a Waters Acuity UPLC/Xevo G2-XS TOF mass spectrometer system. The instrument was operated in negative electro-spray ionization mode and scanned from m/z 200 to 3000. Conjugate was seen as two fragments; Antisense and Sense-Centyrin.

[00368] Centyryn conjugation to siRNA, conjugate purification and analysis: Centyrins were conjugated to siRNA through cysteine-specific chemistry via maleimide. Cysteine-containing Centyryn in PBS at 50-200 μ M were reduced with 10 mM tris(2-carboxyethyl) phosphine (TCEP) at room temperature (30 mins) to yield a free thiol. The free thiol containing Centyryn is then mixed with maleimide containing siRNA duplex in water immediately prior at a molar ratio of ~1.5:1 Centyryn:siRNA. After 2 hr incubation at RT or 37°C, reaction was quenched with N-ethyl maleimide (1mM final NEM concentration in the reaction mixture) The conjugate was purified in two steps. Step I either immobilized metal affinity chromatography (for tagged proteins) or Hydrophobic interaction chromatography (for tagless proteins); to remove un-reacted SiRNA linker. Step II-Capto-DEAE; to remove un-reacted centyryn. Fractions containing conjugate were pooled, exchanged into HBS by desalting using dialysis, and concentrated if necessary.

[00369] FIG. 6: IMAC chromatography for Tagged protein. For IMAC, Histrap HP Columns (1ml, 5ml) from Cytiva were used. Histrap Buffer A (Binding Buffer) was 50mM Tris pH7.4, 500mM NaCl and 10mM imidazole in Type 1 H₂O, Histrap Buffer B (Elution Buffer): 50mM Tris pH7.4, 500mM NaCl and 250mM imidazole in Type 1 H₂O. Reaction sample was directly injected onto the column through sample loop or sample pump. Followed by sample application, column is washed with 5-10 CV of Buffer A. Elution is typically started with step gradient (0-100%). Fractions were collected when UV reading is 50 mAU and above using fraction collector and peak fragmentation.

[00370] FIG. 7: HIC-for tagless proteins (removal of excess of siRNA). For HIC, HiTrap Butyl HP Columns (1ml, 5ml) from Cytiva were used. HIC Buffer A (Binding Buffer) was 2M Ammonium Sulfate, 25mM Sodium Phosphate pH 7.0 in Type 1 H₂O, while, HIC Buffer B (Elution Buffer): 25mM Sodium Phosphate pH 7.0 in Type 1 H₂O. Conjugation reaction sample is diluted with Buffer A in ratio of 1:1. Sample prepared above was injected onto the column through sample loop or sample pump. Followed by sample application, column is washed with 5 CV of Buffer A. Elution is typically started with 0.0 %B and then a gradient as Table 10 below. Fractions collected when UV reading is 50 mAU and above using fraction collector and peak fragmentation.

Table 10

	Type	%B	Length (CV)
1	Linear	30 to 90	3.00
2	Linear	35.0	10.00
3	Linear	100.0	10.00
4	Linear	100.0	5.00

[00371] Ring-opening-To avoid loss of cargo via retro-Michael rection, maleimide ring hydrolysis is performed. Pooled fractions from either histrap (tagged protein) or from HIC (tagless protein) is dialyzed into 25 mM TRIS pH 8.9 buffer. In this buffer the reaction is placed in an incubator shaker at 37 degrees C for 72 hrs. Reaction monitored for completion by LC-MS.

[00372] FIG. 8: Ion-exchange chromatography (IEX)-for tagged and tagless proteins (removal of excess unreacted centyrin). For IEX chromatography, HiTrap Capto DEAE Columns (1ml, 5ml) from Cytiva were used. Capto DEAE Buffer A (Binding Buffer) was 25mM Tris pH 8.8 in Type 1 H₂O, while, Capto DEAE Buffer B (Elution Buffer) used was 25mM Tris pH 8.8, 1M NaCl in Type 1 H₂O.

[00373] Sample after ring opening was directly injected onto the column through sample loop or sample pump. Followed by sample application, column is washed with 5-10 CV of Buffer A. Elution is typically started with 0.0 %B and then a gradient as Table 11 below.

[00374] Unreacted protein is eluted in the flowthrough which is typically the first peak and second peak is typically pure conjugate. All the fractions are collected and pooled. Concentration of the pool is determined by measuring A260 using Nanodrop for yield calculation.

Table 11

1	Linear	20	5
2	Linear	100	15
3	Linear	100	5

[00375] FIG. 9: Example of an analytical SEC of the centyrin-oligonucleotide conjugate. Centyrin-siRNA conjugates were characterized by a combination of analytical techniques. Analytical SEC was used to analyze purity and aggregation state of Centyrin-siRNA conjugates. Waters H-class UPLC along with Waters ACQUITY UPLC Protein BEH SEC Column, 125Å, 1.7 µm, 4.6 X 150mm was used for routine SEC analysis. Typically, 2-5ul sample was injected

using the autosampler, flow rate was 0.25 ml/min and mobile phase used was either 1X PBS or 100mM, pH 7.2 phosphate buffer.

[00376] FIG. 10: Example of SDS PAGE Gel of the conjugate. SDS-PAGE was used to compare amounts of conjugate to free protein. For performing the SDS-PAGE Gel; Invitrogen Mini Gel tank. PowerEase™ Touch 600W Power Supply and 115 VAC was used. Invitrogen - NuPAGE™ 4-12% Bis-Tris Protein Gels, 1.0 mm were used along with 20 × NuPAGE® MES SDS Running Buffer and SeeBlue™ Pre-stained Protein Standard. Samples normalized to around 0.5 mg/mL by ultrapure water. For non-reducing gel, 10 µL of normalized samples are mixed with 10 µL of 2x Laemmli Sample Buffer in 1 to 1 ratio. For reducing gel, 2x Laemmli Sample Buffer is mixed with β-Mercaptoethanol in 95 to 1 ratio and then mixed with 10 µL of samples in 1 to 1 ratio. The resulting sample mixtures were boiled at 95 °C for 5 minutes and then cool down in the Thermocycler. 15 µL of samples and Protein Ladder into appropriate wells were loaded, the voltage was set at 200V and run time was set for around 30 minutes. Once the gel run was finished, it was visualized by Coomassie blue or SYBR Green or methylene blue. Left most lane is Protein Ladder, next is Centyrin by itself and last is the centyrin-siRNA conjugate.

[00377] Candidate siRNA sequences were transfected into human cells (H358) at a range of concentrations. RNA was harvested 24 hours post-transfection and analyzed for GYS1 levels via quantitative reverse transcription polymerase chain reaction (RT-PCR). 18S ribosomal RNA was used as the RT-PCR endogenous control gene. Levels of knockdown were compared to untreated cells. EC50 values were calculated using Graphpad Prism software and Emax values represent the maximum percent knockdown of GYS1 mRNA observed (Table 5).

Table 5

siRNA	Ref.	EC50 (pM)	Emax (%)
A	ABXO8	3.841	70
B	ABXO9	10.02	70.2
C	ABXO12	12.62	71.5
D	ABXO110	21.73	87.9
E	ABXO113	23.74	82.1
F	ABXO114	6.716	75.5
G	ABXO115	4.5	75.2
H	ABXO116	0.7542	69.4
I	ABXO117	20.87	93.8
J	ABXO122	8.93	65.6
K	ABXO123	4.049	75.5

L	ABXO124	2.642	70.5
M	ABXO125	0.8761	53.3
N	ABXO128	2.549	76.3
O	ABXO129	0.4961	64.3
P	ABXO139	0.6143	70.6
Q	ABXO142	23.19	80.3
R	ABXO150	58.88	96.1
S	ABXO151	1.502	75.6
T	ABXO152	24.76	91.1
U	ABXO155	5.893	60.2
V	ABXO156	0.6893	57.6
W	ABXO160	0.787	80.1
X	ABXO168	8.662	75.7
Y	ABXO171	38.82	69.2
Z	ABXO191	54.2	75.4
AA	ABXO197	19.8	77.1
BB	ABXO201	7.1	75.6
CC	ABXO203	5.7	74.9
DD	ABXO205	9.8	80.4

Other siRNAs have also been tested as described above and their EC50 are provided in Table 6.

Table 6

siRNA Pair	EC50 (pM)
EE	1.8
FF	6
GG	5.9
HH	15.1
II	8.2
JJ	25.2
KK	
LL	1.35
MM	3.74
NN	5.66
OO	5.56
PP	6.34
QQ	11.76
RR	2.85
SS	3.35
TT	5.67
UU	0.4
VV	3.72
WW	3.15
XX	2.57
YY	1.64
ZZ	1.56
AAA	ND

BBB	4.87
CCC	2.65
DDD	2.66
EEE	1.21
FFF	1.31
GGG	NA

[00378] Selectivity of candidate siRNA sequences was evaluated by transfection of siRNAs into human cells (HEK-293) at a range of concentrations. Cell viability was assessed 72 hours post-transfection using CellTiterGlo. Emax values are reported as the maximum percent reduction in cell viability at the highest siRNA concentration tested (10 nM) (Table 7).

Table 7

siRNA Pair	Ref.	Emax, % (at 10nM)
A	ABXO8	5.52
B	ABXO9	-0.3
C	ABXO12	31.67
D	ABXO110	25.68
E	ABXO113	22.83
F	ABXO114	-1.1
G	ABXO115	30.85
H	ABXO116	29.27
I	ABXO117	35.19
J	ABXO122	28.25
K	ABXO123	17.21
L	ABXO124	24.09
M	ABXO125	71.45
N	ABXO128	42.41
O	ABXO129	58.02
P	ABXO139	34.44
Q	ABXO142	8.94
R	ABXO150	16.06
S	ABXO151	60.95
T	ABXO152	42.84
U	ABXO155	54.4
V	ABXO156	34.1
W	ABXO160	-4.59
X	ABXO168	51.66
Y	ABXO171	4.79
Z	ABXO191	46.41
AA	ABXO197	25.01
BB	ABXO201	46.92

CC	ABXO203	15.18
DD	ABXO205	60.13

[00379] EXAMPLE 2: Selection of fibronectin type III (FN3) domains that bind CD71

[00380] Panning and Biochemical Screening Methods for Identifying FN3 domains that bind to CD71 that do not inhibit transferrin binding to CD71. To screen for FN3 domains that specifically bind CD71 and do not inhibit transferrin binding to CD71, streptavidin-coated Maxisorp plates (Nunc catalog 436110) are blocked for 1 hour in Starting Block T20 (Pierce) and then are coated with biotinylated CD71 (using same antigen as in panning) or negative controls (an unrelated Fc-fused recombinant protein and human serum albumin) for 1 hour in the presence of transferrin or with FN3 protein that binds to the CD71 transferrin binding site. The concentration of transferrin is up to 35 μ M. Without being bound to any particular theory, the inclusion of the transferrin or the FN3 protein that binds to the CD71 transferrin binding site pushes the selection of the FN3 domains to those that do not compete or inhibit with transferrin binding to CD71. Plates are rinsed with TBST and diluted lysate is applied to plates for 1 hour. Following additional rinses, wells are treated with HRP-conjugated anti-V5 tag antibody (Abcam, ab1325), for 1 hour and then are assayed with POD (Roche, 11582950001). The DNA from FN3 domain lysates with signals at least 10-fold ELISA signal above that of streptavidin controls are sequenced resulting in FN3 domain sequences isolated from the screening.

[00381] EXAMPLE 3: Selection of fibronectin type III (FN3) domains that bind CD71 and are not competitive with transferrin

[00382] To identify CD71 binding FN3 domains that were either not competitive or minimally competitive with transferrin a biased CIS-display strategy was designed. In short, using the output recovered after 5 rounds of panning on the ECD of human CD71 (Example 3). Additional rounds of off-rate selection were performed as described in Example 3 with the addition of either 1) a wash step with human holo transferrin to elute Centyrins that bound at the same site as transferrin before the final elution step or 2) elution of FN3 domain binders with monoclonal antibody OKT9. FN3 domains recovered from the transferrin wash strategy and the OKT9 elution strategy were PCR amplified and cloned into pET vector as previously described. 228 FN3 domains that specifically bound huCD71 were confirmed by ELISA for binding to huCD71 ECD. A subset of the unique binders was analyzed by SEC, conjugated to MMAF and assessed for internalization via cell viability assay in SKBR-3 cells +/- holo human transferrin. The polypeptides were found to be internalized by the receptor.

[00383] Integral Molecular performed Membrane Proteome Array (MPA) assay to profile the specificity of ABX1198 (SEQ ID NO: 509) and ABX1100 (SEQ ID NO: 509 plus siRNA pair with linker number OOOO) against the library of human membrane proteins. The MPA library contains over 6000 human membrane proteins, including 94% of all single-pass, multi-pass and GPI anchored proteins including GPCRs, ion channels and transporters with each membrane protein uniquely expressed in an avian QT6 cell background. Flow cytometry is used to directly detect ligand (FN3 domain) binding to membrane proteins individually expressed in unfixed cells.

[00384] ABX1198 (SEQ ID NO: 509) and ABX1100 (SEQ ID NO: 509 plus siRNA pair with linker number OOOO) were screened at the concentration with optimal signal/background noise ratio, 1.25 ug/ml, 1.25 ug/ml and 0.31 ug/ml respectively, against the MPA. Membrane protein targets identified in screening were followed up in validation procedure using ligand serial dilution and cells individually transfected with identified targets.

[00385] EXAMPLE 4. Knockdown of mRNA in muscle cells using CD71 FN3 domain-oligonucleotide conjugates.

[00386] muCD71 binding FN3 domains are conjugated to siRNA oligonucleotides or antisense oligonucleotides (ASOs) using maleimide chemistry via a cysteine that is uniquely engineered into the FN3 domain. The cysteine substitutions can be one such as those provided for herein and also as provided for in U.S. Patent Application Publication No. 20150104808, which is hereby incorporated by reference in its entirety. siRNAs or ASOs are modified with standard chemical modifications and confirmed to enable knockdown of the targeted mRNA in vitro. FN3 domain-oligonucleotide conjugates are dosed intravenously in mice at doses up to 10 mg/kg oligonucleotide payload. At various time points following dosing, mice are sacrificed; skeletal muscle, heart muscle and various other tissues will be recovered and stored in RNeasyTM (Sigma Aldrich) until needed. Target gene knockdown is assessed using standard qPCR $\Delta\Delta C_T$ methods and primers specific for the target gene and a control gene. The target gene is found to be knock downed in the muscles and such knockdown is enhanced by conjugating the siRNA or ASO to the CD71 FN3 binding domain.

[00387] FN3-siRNA conjugates tested are as described in table 8

Table 8

Conjugate	CD71 FN3 domain (Centyrin)	GYS1 siRNA Pair
ABXC-1	SEQ ID NO: 572	RRRR
ABXC-2	SEQ ID NO: 572	SSSS
ABXC-3	SEQ ID NO: 572	TTTT

ABX1005	SEQ ID NO: 572	Non-GYS1 target/control
ABXC-27	SEQ ID NO: 572	A
ABXC-28	SEQ ID NO: 572	QQQQ
ABXC-29	SEQ ID NO: 572	PPPP

[00388] FIG. 4A demonstrates the knockdown of GYS1 mRNA in mouse gastrocnemius muscle using 3 different FN3 domain-siRNA conjugates compared with vehicle alone. For efficacy studies, male GAA^{-/-} mice (at the ages of 4-5 weeks) were obtained from Jackson Laboratories. All animals were treated in accordance with IACUC protocols. Five animals received a single tail vein intravenous bolus injection of either 5.4 mg/kg of three different FN3 domain-siRNA conjugates (3 mpk Gys1siRNA) or vehicle. Four weeks after the single dose, the mice were euthanized, gastrocnemius muscle were collected in RNAlater, stored at 4C overnight and were frozen at -80C. Total RNA was isolated from the gastrocnemius using Qiagen's RNeasy Fibrous Tissue kit. Expression levels of the target Gys1 and the endogenous controls (Pfkfb3, Ubc, Hprt1 and Aha1) were analyzed using real-time, quantitative PCR. Data were analyzed using the $\Delta\Delta C_t$ method normalizing to control animals dosed with vehicle alone. The percentage knockdown of Gys1 mRNA in the FN3 domain-siRNA conjugate treatment groups were measured by subtracting the percentage remaining Gys1 mRNA levels by 100. Statistical significance was calculated using One-way ANOVA with Dunnett's multiple comparison tests in the GraphPad Prism software. Statistical significance is displayed on the figure with asterisk ***p<0.001.

[00389] FIG. 4B demonstrates the knockdown of GYS1 protein in mouse gastrocnemius muscle using 3 different FN3 domain-siRNA conjugates compared with vehicle alone. Gys1 protein quantification in gastrocnemius was performed by homogenizing gastrocnemius in RIPA buffer. Protein concentrations in the gastrocnemius were measured using the Bradford assay. Gys1 levels were quantified using the manufacturer's standard method for 12-230 kDa Jess separation modules (SM-W004). The proteins were separated by immobilizing on capillaries using protein Simple's proprietary photoactivated capture chemistry. Anti-Gys1 primary antibody were used at 1:100 dilution. The chemiluminescent revelations were established using peroxide/luminol-S. A digital image of the capillaries' chemiluminescence was captured using Compass' Simple Western software, which automatically measures height (chemiluminescence intensity), area, and signal/noise ratio. An internal system was included in each run. The peak area values of FN3 domain-siRNA conjugate treatment groups were normalized to the vehicle treated tissues and the percentage knockdown of Gys1 protein in the treatment groups were

measured by subtracting the percentage remaining Gys1 protein levels by 100. Statistical significance was calculated using One-way ANOVA with Dunnett's multiple comparison tests in the GraphPad Prism software. Statistical significance is displayed on the figure with asterisk *** $p < 0.001$.

[00390] FIG. 5 demonstrates the GYS1 knockdown is highly specific for skeletal muscle using 3 different FN3 domain-siRNA conjugates compared with a siRNA to a different target (AHA-1). Male GAA^{-/-} mice (at the ages of 8-9 weeks) were obtained from Jackson Laboratories. All animals were treated in accordance with IACUC protocols. Three animals received a single tail vein intravenous bolus injection of either 17.9 mg/kg of three different FN3 domain-siRNA conjugates (10 mpk Gys1siRNA), 17.9 mg/kg of one FN3 domain-siRNA conjugate (10 mpk Aha1 siRNA) or vehicle. Two weeks after the single dose, the mice were euthanized. Gastrocnemius, quadriceps, diaphragm, heart, liver, and kidney tissues were collected in RNAlater, stored at 4°C overnight and were frozen at -80°C. Total RNA was isolated from the tissues using Qiagen's RNeasy Fibrous Tissue kit. Expression levels of the target Gys1/Aha1 and the endogenous control (Pfkfb3) were analyzed using real-time, quantitative PCR. Data were analyzed using the $\Delta\Delta C_t$ method normalizing to control animals dosed with vehicle alone.

[00391] EXAMPLE 5: RNA-seq experiments comparing ABX-HHH treated and untreated cells.

[00392] mRNA sequencing was performed on polyA⁺ selected mRNA after HEK293T cells were lipofected with 10 nM ABXO-HHH for 24h. 6 replicates per treatment were sequenced to an average depth > 30 million reads. Libraries were pseudoaligned to the GRCh38 transcriptome using Kallisto with a mapping rate > 90%. The values in this volcano plot were generated using DESeq2 to compare ABXO-HHH treated RNA-seq libraries versus untreated cells. Each dot in the plot in FIG. 2 represents the measured expression change in a gene. The X axis represents the $\log_2$ fold change in gene expression (ABXO-HHH divided by untreated) for all genes. The Y axis shows the negative $\log_{10}$ transformed adjusted P value. The black arrow points to GYS1 on the plot.

[00393] EXAMPLE 6: CD71 FN3 domain siRNA conjugate binding specificity.

[00394] Integral Molecular (www.integralmolecular.com) performed their proprietary Membrane Proteome Array (MPA) assay to profile the specificity of CD71 FN3 domain and CD71 FN3 domain siRNA conjugate against the library of human membrane proteins (FIG. 3). The MPA contains over 6000 human membrane proteins, covering 94% of all single-pass, multi-pass and GPI anchored proteins including GPCRs, ion channels and transporters with each

membrane protein uniquely expressed in an avian QT6 cell background. Flow cytometry was used to directly detect FN3 domain binding to membrane proteins individually expressed in unfixed cells.

[00395] FN3 domain and FN3 domain-siRNA conjugate were screened at the concentration with optimal signal/background noise ratio, 1.25 ug/ml or 0.31 ug/ml respectively, against the MPA. Membrane protein targets identified in screening were validated using ligand serial dilution on cells uniquely expressing the identified targets.

[00396] GENERAL METHODS

[00397] Standard methods in molecular biology are described Sambrook, Fritsch and Maniatis (1982 & 1989 2nd Edition, 2001 3rd Edition) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Sambrook and Russell (2001) *Molecular Cloning, 3rd ed.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Wu (1993) *Recombinant DNA*, Vol. 217, Academic Press, San Diego, CA). Standard methods also appear in Ausbel, *et al.* (2001) *Current Protocols in Molecular Biology, Vols.1-4*, John Wiley and Sons, Inc. New York, NY, which describes cloning in bacterial cells and DNA mutagenesis (Vol. 1), cloning in mammalian cells and yeast (Vol. 2), glycoconjugates and protein expression (Vol. 3), and bioinformatics (Vol. 4).

[00398] Methods for protein purification including immunoprecipitation, chromatography, electrophoresis, centrifugation, and crystallization are described (Coligan, *et al.* (2000) *Current Protocols in Protein Science, Vol. 1*, John Wiley and Sons, Inc., New York). Chemical analysis, chemical modification, post-translational modification, production of fusion proteins, glycosylation of proteins are described (see, *e.g.*, Coligan, *et al.* (2000) *Current Protocols in Protein Science, Vol. 2*, John Wiley and Sons, Inc., New York; Ausubel, *et al.* (2001) *Current Protocols in Molecular Biology, Vol. 3*, John Wiley and Sons, Inc., NY, NY, pp. 16.0.5-16.22.17; Sigma-Aldrich, Co. (2001) *Products for Life Science Research*, St. Louis, MO; pp. 45-89; Amersham Pharmacia Biotech (2001) *BioDirectory*, Piscataway, N.J., pp. 384-391). Production, purification, and fragmentation of polyclonal and monoclonal antibodies are described (Coligan, *et al.* (2001) *Current Protocols in Immunology, Vol. 1*, John Wiley and Sons, Inc., New York; Harlow and Lane (1999) *Using Antibodies*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Harlow and Lane, *supra*). Standard techniques for characterizing ligand/receptor interactions are available (see, *e.g.*, Coligan, *et al.* (2001) *Current Protocols in Immunology, Vol. 4*, John Wiley, Inc., New York).

[00399] All references cited herein are incorporated by reference to the same extent as if each individual publication, database entry (e.g. Genbank sequences or GeneID entries), patent

application, or patent, was specifically and individually indicated to be incorporated by reference. This statement of incorporation by reference is intended by Applicants, pursuant to 37 C.F.R. §1.57(b)(1), to relate to each and every individual publication, database entry (e.g. Genbank sequences or GeneID entries), patent application, or patent, each of which is clearly identified in compliance with 37 C.F.R. §1.57(b)(2), even if such citation is not immediately adjacent to a dedicated statement of incorporation by reference. The inclusion of dedicated statements of incorporation by reference, if any, within the specification does not in any way weaken this general statement of incorporation by reference. Citation of the references herein is not intended as an admission that the reference is pertinent prior art, nor does it constitute any admission as to the contents or date of these publications or documents.

[00400] The present embodiments are not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the embodiments in addition to those described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims.

[00401] The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the embodiments. Various modifications of the embodiments in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.


WHAT IS CLAIMED IS:

1. A composition comprising a siRNA comprising a sense strand and antisense strand as provided herein.
2. The composition of claim 1, wherein the siRNA does not contain any modified nucleobases.
3. The composition of claims 1 or 2, wherein the siRNA further comprises a linker covalently attached to the sense strand or the anti-sense strand of the siRNA.
4. The composition of claim 3, wherein the linker is attached to the 5' end or the 3' end of the sense strand or the anti-sense strand.
5. The composition of any one of claims 1-4, wherein the siRNA further comprises a vinyl phosphonate modification on the sense strand or the anti-sense strand.
6. The composition of 5, wherein the vinyl phosphonate modification is attached to the 5' end or the 3' end of the sense strand or the anti-sense strand.
7. The composition of any one of claims 1-6, wherein the sense strand comprises a sequence of SEQ ID NO: 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 600, 602, 604, 606, 608, 610, 612, 614, 616, 618, 620, 622, 624, 626, 628, 630, 632, 634, 636, 638, 640, 642, 644, 646, 648, 650, 652, 654, 656, 658, 660, 662, 664, 666, 668, 670, 672, 674, 676, 678, 680, 682, 684, 686, 688, 690, 692, 694, 696, 698, 700, 702, 704, 706, 801-860, 921-980, or as set forth in Table 1A, Table 1B, or Table 2.
8. The composition of any one of claims 1-7, wherein the anti-sense strand comprises a sequence of SEQ ID NO: 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 601, 603, 605, 607, 609, 611, 613, 615, 617, 619, 621, 623, 625, 627, 629, 631, 633, 635, 637, 639, 641, 643, 645, 647, 649, 651, 653, 655, 657, 659, 661, 663, 665, 667, 669, 671, 673, 675, 677, 679, 681, 683, 685, 687, 689, 690, 691, 693, 695, 697, 699, 701, 703, 705, 707, 861-920, 981-1042, or as set forth in Table 1A, Table 1B, or Table 2.
9. The composition of any one of the preceding claims, wherein the siRNA molecule comprises the siRNA pair of A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, S, T, U, V, W, X, Y, Z, AA, BB, CC, DD, EE, FF, GG, HH, II, JJ, KK, LL, MM, NN, OO, PP, QQ, RR, SS, TT, UU, VV, WW, XX, YY, ZZ, AAA, BBB, CCC, DDD, EEE, FFF, GGG, HHH, III, JJJ, KKK, LLL, MMM, NNN, OOO, PPP, QQQ, RRR, SSS, TTT, UUU, VVV, WWW, XXX,

YYY, ZZZ, AAAA, BBBB, CCCC, DDDD, EEEE, FFFF, GGGG, HHHH, IIII, JJJJ, KKKK, LLLL, MMMM, NNNN, OOOO, PPPP, or as set forth in Table 1A, Table 1B, or Table 2.

10. The composition of any one of the preceding claims, wherein the sense strand comprises 19 nucleotides.
11. The composition of any one of the preceding claims, wherein the antisense strand comprises 21 nucleotides.
12. The composition of any one of claims 1-11, wherein the composition comprises the siRNA pair as provided for in Table 2, with a linker and/or vinyl phosphonate modification as set forth in Table 3.
13. The composition of any one of the preceding claims, wherein the siRNA molecule has the formula as illustrated in Formula I:

N ₁ N ₂ N ₃ N ₄ N ₅ N ₆ N ₇ N ₈ N ₉ N ₁₀ N ₁₁ N ₁₂ N ₁₃ N ₁₄ N ₁₅ N ₁₆ N ₁₇ N ₁₈ N ₁₉	Sense-strand (SS)
N ₂₁ N ₂₀ N ₁₉ N ₁₈ N ₁₇ N ₁₆ N ₁₅ N ₁₄ N ₁₃ N ₁₂ N ₁₁ N ₁₀ N ₉ N ₈ N ₇ N ₆ N ₅ N ₄ N ₃ N ₂ N ₁	Antisense-strand (AS)
- wherein each nucleotide represented by N, is independently, A, U, C, or G or a modified nucleotide base, such as those provided for herein.
14. The composition of claim 13, wherein the sense strand comprises a 2'-O-methyl modified nucleotide with a phosphorothioate (PS) modified backbone at N₁ and N₂, a 2'-fluoro modified nucleotide at N₃, N₇, N₈, N₉, N₁₂, and N₁₇, and a 2'-O-methyl modified nucleotide at N₄, N₅, N₆, N₁₀, N₁₁, N₁₃, N₁₄, N₁₅, N₁₆, N₁₈, and N₁₉.
15. The composition of claim 13, wherein the antisense strand comprises a vinylphosphonate moiety attached to N₁, a 2'-fluoro- modified nucleotide with a phosphorothioate (PS) modified backbone at N₂, a 2'-O-methyl modified nucleotide at N₃, N₄, N₅, N₆, N₇, N₈, N₉, N₁₀, N₁₁, N₁₂, N₁₃, N₁₅, N₁₆, N₁₇, N₁₈, and N₁₉, a 2'-fluoro- modified nucleotide at N₁₄, and a 2'-O-methyl modified nucleotide with a phosphorothioate (PS) modified backbone at N₂₀ and N₂₁.
16. The composition of claim 13, wherein a vinylphosphonate moiety attached to N₁ of the antisense strand.
17. The composition of any one of the preceding claims, wherein the siRNA molecule has the formula as illustrated in Formula I:



wherein F_1 is a polypeptide comprising at least one FN3 domain and is conjugated to a linker, L_1 , L_1 is linked to X_S , wherein X_S is a 5' to 3' oligonucleotide sense strand of a double stranded siRNA molecule and X_{AS} is a 3' to 5' oligonucleotide antisense strand of a double stranded siRNA molecule; and wherein X_S and X_{AS} form a double stranded siRNA molecule.

18. The composition of claim 17, wherein F_1 comprises polypeptide having a formula of $(X_1)_n-(X_2)_q-(X_3)_y$, wherein X_1 is a first FN3 domain; X_2 is second FN3 domain; X_3 is a third FN3 domain or half-life extender molecule; wherein n , q , and y are each independently 0 or 1, provided that at least one of n , q , and y is 1.

19. A composition comprising one or more FN3 domains conjugated to a siRNA comprising a sense strand and an anti-sense strand.

20. The composition of claim 19, wherein the siRNA does not contain any modified nucleobases.

21. The composition of claims 19 or 20, wherein the siRNA further comprises a linker.

22. The composition of the claim 21, wherein the linker is covalently attached to the sense strand or the anti-sense strand.

23. The composition of claims 21 or 22, wherein the linker is attached to the 5' end or the 3' end of the sense strand or the anti-sense strand.

24. The composition of any one of claims 19-23, wherein the siRNA further comprises a vinyl phosphonate modification on the sense strand or the anti-sense strand.

25. The composition of 24, wherein the vinyl phosphonate modification is attached to the 5' end or the 3' end of the sense strand or the anti-sense strand.

26. The composition of any one of claims 19-25, wherein the sense strand comprises a sequence of SEQ ID NO: 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 600, 602, 604, 606, 608, 610, 612, 614, 616, 618, 620, 622, 624, 626, 628, 630, 632, 634, 636, 638, 640, 642, 644, 646, 648, 650, 652, 654, 656, 658, 660, 662, 664, 666, 668, 670, 672, 674, 676, 678, 680, 682, 684, 686, 688, 690, 692,

694, 696, 698, 700, 702, 704, 706, 801-860, 921-980, or as set forth in Table 1A, Table 1B, or Table 2.

27. The composition of any one of claims 19-25, wherein the anti-sense strand comprises a sequence of SEQ ID NO: 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 601, 603, 605, 607, 609, 611, 613, 615, 617, 619, 621, 623, 625, 627, 629, 631, 633, 635, 637, 639, 641, 643, 645, 647, 649, 651, 653, 655, 657, 659, 661, 663, 665, 667, 669, 671, 673, 675, 677, 679, 681, 683, 685, 687, 689, 690, 691, 693, 695, 697, 699, 701, 703, 705, 707, 861-920, 981-1042, or as set forth in Table 1A, Table 1B, or Table 2.

28. The composition of any one of claims 19-25, wherein the siRNA molecule comprises the siRNA pair of A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, S, T, U, V, W, X, Y, Z, AA, BB, CC, DD, EE, FF, GG, HH, II, JJ, KK, LL, MM, NN, OO, PP, QQ, RR, SS, TT, UU, VV, WW, XX, YY, ZZ, AAA, BBB, CCC, DDD, EEE, FFF, GGG, HHH, III, JJJ, KKK, LLL, MMM, NNN, OOO, PPP, QQQ, RRR, SSS, TTT, UUU, VVV, WWW, XXX, YYY, ZZZ, AAAA, BBBB, CCCC, DDDD, EEEE, FFFF, GGGG, HHHH, IIII, JJJJ, KKKK, LLLL, MMMM, NNNN, OOOO, PPPP, or as set forth in Table 1A, Table 1B, or Table 2.

29. The composition of any one of claims 19-25, wherein the composition comprises the siRNA pair of as provided for in Table 2, with a linker and/or vinyl phosphonate modification as set forth in Table 3.

30. The composition of any one of claims 19-29, wherein the FN3 domain is conjugated to the siRNA molecule through a cysteine on the FN3 domain.

31. The composition of claim 30, wherein the cysteine is at a position as described herein.

32. The composition of claims 30 or 31, wherein the cysteine in the FN3 domain is at a position that corresponds to residue 6, 8, 10, 11, 14, 15, 16, 20, 30, 34, 38, 40, 41, 45, 47, 48, 53, 54, 59, 60, 62, 64, 70, 88, 89, 90, 91, or 93 of the FN3 domain based on SEQ ID NO: 1 of U.S. Patent No. 10,196,446.

33. The composition of claim 32, wherein the cysteine is located at a position that corresponds to residue 6, 53, or 88.

34. The composition of any one of claims 19-33, wherein the one or more FN3 domains comprises a FN3 domain that binds to CD71.

35. The composition of any one of claims 19-34, wherein the FN3 domain has a sequence selected from the group consisting of SEQ ID NOs: 509, 708, and 710.

36. The composition of any one of claims 19-35, wherein the one or more FN3 domains comprises a FN3 domain that binds to CD71 comprises at least two FN3 domains linked by a peptide linker.
37. The composition of claim 36, wherein the composition comprises a first FN3 domain and a second FN3 domain.
38. The composition of claim 37, wherein the first FN3 domain and the second FN3 domain bind to different proteins.
39. The composition of claim 37, wherein the first FN3 domain and the second FN3 domain bind to the same protein.
40. The composition of any one of claims 36-39, wherein the first FN3 domain binds to CD71.
41. The composition of any one of claims 36-39, wherein the second FN3 domain binds to a different target that does not bind CD71.
42. The composition of any one of claims 19-41, wherein the FN3 domain comprises a sequence that is at least 87% identical to or is identical to a sequence of SEQ ID NO: 273, 288-291, 301-310, 312-572, 592-599, or 708-710.
43. The composition of any one of claim 19-42, further comprising a third FN3 domain.
44. The composition of claim 43, wherein the third FN3 domain is a FN3 domain that binds to CD71 or albumin.
45. The composition of claim 44, wherein the FN3 domain that binds CD71 has an amino acid sequence as provided herein, including but not limited to SEQ ID NO: 273, 288-291, 301-310, 312-572, 592-599, or 708-710, or a binding fragment thereof.
46. The composition of claim 44, wherein the FN3 that binds albumin has an amino acid sequence as provided herein, including but not limited to SEQ ID NO: 101-119, or a binding fragment thereof.
47. The composition of claims 45-46, wherein the FN3 domain has a cysteine substitution as provided herein.
48. A composition having a formula selected from $(X_1)_n-(X_2)_q-(X_3)_y-L-X_4$, $C-(X_1)_n-(X_2)_q-L-X_4-(X_3)_y$, $(X_1)_n-(X_2)_q-L-X_4-(X_3)_y-C$, $C-(X_1)_n-(X_2)_q-L-X_4-L-(X_3)_y$, or $(X_1)_n-(X_2)_q-L-X_4-L-(X_3)_y-C$, wherein:
- X₁ is a first FN3 domain;
 - X₂ is second FN3 domain;
 - X₃ is a third FN3 domain or half-life extender molecule;
 - L is a linker; and


X₄ is a nucleic acid molecule,

C is a polymer, such as PEG, albumin binding protein,



wherein n, q, and y are each independently 0 or 1.

49. The composition of claim 48, wherein X₁, X₂, and X₃ bind to the same or different target proteins.
50. The composition of claims 48 or 49, wherein y is 0.
51. The composition of claims 48 or 49, wherein n is 1, q is 0, and y is 0.
52. The composition of claims 48 or 49, wherein n is 1, q is 1, and y is 0.
53. The composition of claims 48 or 49, wherein n is 1, q is 1, and y is 1.
54. The composition of any one of claims 48-53, wherein the third FN3 domain increases the half-life of the molecule as a whole as compared to a molecule without X₃.
55. The composition of claim 48, wherein the third FN3 domain is a FN3 domain that binds to albumin.
56. The composition of any one of claims 48-55, wherein the linker is a linker as provided herein.
57. The composition of any one of claims 48-56, wherein the FN3 domains are connected by a peptide linker.
58. The composition of claim 57, wherein the peptide linker is (GS)₂, (SEQ ID NO:720), (GGGS)₂ (SEQ ID NO:721), (GGGS)₅ (SEQ ID NO:722), (AP)₂₋₂₀, (AP)₂ (SEQ ID NO:723), (AP)₅ (SEQ ID NO:724), (AP)₁₀ (SEQ ID NO:725), (AP)₂₀ (SEQ ID NO:726) and A(EAAAK)₅AAA (SEQ ID NO:727) or (EAAAK)₁₋₅ (SEQ ID NO: 728), or any combination thereof.
59. The composition of any one of claims 48-58, wherein the first, second, or third FN3 domain has an amino acid sequence as provided herein.
60. The composition of any one of claims 48-59, wherein X₄ is a siRNA molecule.
61. The composition of claim 60, wherein the siRNA molecule is a siRNA molecule provided herein.
62. The composition of claim 61, wherein the siRNA molecule is a siRNA that reduces the expression of GYS1.
63. The composition of claim 61, wherein the siRNA molecule is a siRNA that specifically reduces the expression of GYS1.
64. The composition of claim 61, wherein the siRNA molecule is a siRNA that reduces the expression of GYS1 and does not significantly reduce the expression of other RNAs.

65. The composition of claim 61, wherein the siRNA molecule is a siRNA that reduces the expression of GYS1 and does not reduce the expression of other RNAs by more than 50% in an assay described herein at a concentration of no more than 200 nM as described herein.
66. The composition of claim 61, wherein the siRNA molecule is a siRNA that reduces the expression of GYS1 and reduces the concentration of GYS1 protein.
67. The composition of claim 61, wherein the siRNA molecule is a siRNA that reduces the expression of GYS1 and reduces the concentration of glycogen in a cell.
68. The composition of claim 61, wherein the cell is a muscle cell or a heart cell.
69. The method of any one of claims 54-74, wherein the siRNA is a siRNA pair as provided in the following formula:



70. The composition of claim 69, wherein N₁ of the anti-sense strand comprises a vinyl phosphonate modification.
71. The composition of claim 69, wherein maleimide is hydrolyzed to form the following mixture of compounds, or one or both of each compound, or exclusively one of the compounds



72. The composition of any one of claims 48-71, wherein the siRNA is a siRNA Pair as provided herein or a siRNA pair selected from the group consisting as provided for in Table 1A, Table 1B, or Table 2.
73. A composition having a formula A₁-B₁, wherein A₁ has a formula of (C)_n-(L₁)_t-X_s and B₁ has a formula of X_{AS}-(L₂)_q-(F₁)_y, wherein:

C is a polymer, such as PEG, albumin binding protein;

L₁ and L₂ are each, independently, a linker;

X_S is a 5' to 3' oligonucleotide sense strand of a double stranded siRNA molecule;

X_{AS} is a 3' to 5' oligonucleotide antisense strand of a double stranded siRNA molecule;

F_1 is a polypeptide comprising at least one FN3 domain;

wherein n , t , q , and y are each independently 0 or 1;

wherein X_S and X_{AS} form a double stranded oligonucleotide molecule to form the composition/complex.

74. A composition having a formula A_1-B_1 , wherein A_1 has a formula of $(F_1)_n-(L_1)_t-X_S$ and B_1 has a formula of $X_{AS}-(L_2)_q-(C)_y$, wherein:

C is a polymer, such as PEG, albumin binding protein;

L_1 and L_2 are each, independently, a linker;

X_S is a 5' to 3' oligonucleotide sense strand of a double stranded siRNA molecule;


X_{AS} is a 3' to 5' oligonucleotide antisense strand of a double stranded siRNA molecule;

F_1 is a polypeptide comprising at least one FN3 domain;


wherein n , t , q , and y are each independently 0 or 1;

wherein X_S and X_{AS} form a double stranded oligonucleotide molecule to form the composition/complex.


75. The composition of claim 73 or 74, wherein L_1 has the formula:




76. The composition of claim 73 or 74, wherein L_2 has the formula:



77. The composition of claim 73 or 74, wherein A_1-B_1 has a formula of:



78. The composition of claim 73 or 74, where in A_1-B_1 has a formula of:



79. The composition of claim 73 or 74, wherein F₁ comprises polypeptide having a formula of (X₁)_n-(X₂)_q-(X₃)_y, wherein X₁ is a first FN3 domain; X₂ is second FN3 domain; X₃ is a third FN3 domain or half-life extender molecule; wherein n, q, and y are each independently 0 or 1, provided that at least one of n, q, and y is 1.
80. The composition of claim 73 or 74, wherein X₁ is a CD71 FN3 binding domain.
81. The composition of claim 73 or 74, wherein X₂ is a CD71 FN3 binding domain.
82. The composition of claim 73 or 74, wherein X₃ is a FN3 domain that binds to human serum albumin.
83. The composition of claim 73 or 74, wherein X₃ is a Fc domain without effector function that extends the half-life of a protein.
84. The composition of any one of claims 73-83, wherein X_S comprises a sequence of SEQ ID NO: 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 600, 602, 604, 606, 608, 610, 612, 614, 616, 618, 620, 622, 624, 626, 628, 630, 632, 634, 636, 638, 640, 642, 644, 646, 648, 650, 652, 654, 656, 658, 660, 662, 664, 666, 668, 670, 672, 674, 676, 678, 680, 682, 684, 686, 688, 690, 692, 694, 696, 698, 700, 702, 704, 706, 801-860, 921-980, or as set forth in Table 1A, Table 1B, or Table 2.
85. The composition of any one of claims 73-83, wherein X_{AS} comprises a sequence of SEQ ID NO: 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 601, 603, 605, 607, 609, 611, 613, 615, 617, 619, 621, 623, 625, 627, 629, 631, 633, 635, 637, 639, 641, 643, 645, 647, 649, 651, 653, 655, 657, 659, 661, 663, 665, 667, 669, 671, 673, 675, 677, 679, 681, 683, 685, 687, 689, 690, 691, 693, 695, 697, 699, 701, 703, 705, 707, 861-920, 981-1042, or as set forth in Table 1A, Table 1B, or Table 2.
86. The composition of any one of claims 73-83, wherein X_S and X_{AS} form a siRNA pair selected from the group consisting of A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, S, T, U, V, W, X, Y, Z, AA, BB, CC, DD, EE, FF, GG, HH, II, JJ, KK, LL, MM, NN, OO, PP, QQ, RR,

SS, TT, UU, VV, WW, XX, YY, ZZ, AAA, BBB, CCC, DDD, EEE, FFF, GGG, HHH, III, JJJ, KKK, LLL, MMM, NNN, OOO, PPP, QQQ, RRR, SSS, TTT, UUU, VVV, WWW, XXX, YYY, ZZZ, AAAA, BBBB, CCCC, DDDD, EEEE, FFFF, GGGG, HHHH, IIII, JJJJ, KKKK, LLLL, MMMM, NNNN, OOOO, PPPP, or as set forth in Table 1A, Table 1B, or Table 2.

87. The composition of any one of claims 73-83, wherein F₁ comprises a sequence that is at least 87% identical to or is identical to a sequence of SEQ ID NO: 273, 288-291, 301-310, 312-572, 592-599, or 708-710.

88. The composition of any one of claims 73-83, wherein F₁ comprises a polypeptide that binds to albumin.

89. A pharmaceutical composition comprising a composition of any one of claims 1-88.

90. A kit comprising a composition of any of claims 1-88.

91. A method of treating muscle diseases, immunological diseases, cancer, Pompe Disease, or glycogen storage disease in a subject in need thereof, the method comprising administering to the subject composition of any of claims 1-88 or any composition provided herein.

92. A use of a composition as provided herein or of any of claims 1-88 in the preparation of a pharmaceutical composition or medicament for treating muscle diseases, immunological diseases, cancer, Pompe Disease, or glycogen storage disease.

93. Use of a composition as provided herein or any of claims 1-88 for treating muscle diseases, immunological diseases, cancer, Pompe Disease, or glycogen storage disease.

94. A method of reducing the expression of a target gene in a cell, the method comprising contacting the cell with a composition of any of claims 1-88 or a composition as provided herein.

95. The method of claim 94, wherein the target gene is GYS1.

96. The method of claim 95, wherein the GYS1 has a mutation.

97. A method of delivering a siRNA molecule to a cell in a subject, the method comprising administering to the subject a pharmaceutical composition comprising a composition of any of claims 1-88 to the subject.

98. The method of claim 97, wherein the cell is a CD71 positive cell.

99. The method of any one of claims 97-98, wherein the cell is a tumor cell, a liver cell, a muscle cell, an immune cell, a dendritic cell, a heart cell, a cell of the CNS, or a cell inside the blood brain barrier.

100. The method of any one of claims 97-99, wherein the siRNA downregulates the expression of a target gene in the cell.

101. The method of claim 100, wherein the downregulation of the expression of the target gene results in a reduction of about 99%, 90-99%, 50-90%, or 10-50%.

102. The method of claim 100, wherein the target gene is GYS1.
103. The method of claim 102, wherein the GYS1 has a mutation.
104. The method of claim 100-103, wherein the reduced target gene and protein results in reduction in glycogen.
105. The method of claim 104, wherein the glycogen reduction leads to improvement in disease status such as Pompe disease and any glycogen storage diseases.
106. The methods of claims 91-93, wherein the glycogen storage disease is selected from the group consisting of Cori's disease or Forbes' disease (GSD3, Glycogen debranching enzyme (AGL) deficiency), McArdle disease (GSD5, Muscle glycogen phosphorylase (PYGM) deficiency), type II Diabetes/diabetic nephropathy, Aldolase A Deficiency GSD12, Lafora Disease, hypoxia, Andersen disease (GSD4, Glycogen debranching enzyme (GBE1) deficiency), Tarui's Disease (GSD7, Muscle phosphofructokinase (PFKM) deficiency), adult polyglucosan body disease, Glycogen synthase (GYS2) deficiency (GSD0), Glucose-6-phosphatase (G6PC / SLC37A4) deficiency (GSD1, von Gierke's disease), Hers' disease (GSD6, Liver glycogen phosphorylase (PYGL) or Muscle phosphoglycerate mutase (PGAM2) deficiency), Phosphorylase kinase (PHKA2 / PHKB / PHKG2 / PHKA1) deficiency (GSD9), Phosphoglycerate mutase (PGAM2) deficiency (GSD10), Muscle lactate dehydrogenase (LDHA) deficiency (GSD11), Fanconi-Bickel syndrome (GSD 11, Glucose transporter (GLUT2) deficiency, Aldolase A deficiency (GSD 12), β -enolase (ENO3) deficiency (GSD13), and Glycogenin-1 (GYG1) deficiency (GSD15).