




FIG. 1A

2/28




FIG. 1B




FIG. 2

4/28




FIG. 3A




FIG. 3B

5/28




FIG. 4A




FIG. 4B




FIG. 5A




FIG. 5B

7/28




FIG. 6A




FIG. 6B

8/28




FIG. 7A




FIG. 7B

9/28




FIG. 8

10/28




FIG. 9

U87 MG Tumor Volume




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 10D




FIG. 10E

12/28




FIG. 11

13/28




FIG. 12

14/28





FIG. 13

15/28



Group	Median Survival	P Value
Vehicle	52 days	
Cmpd 1 (20 mg/kg)	59 days	<0.05
Cmpd 1 (10 mg/kg)	63 days	<0.05
Cmpd 1 (5 mg/kg)	50 days	NS
Cmpd 1 (2.5 mg/kg)	55 days	NS
Temozolomide	>75 days	TBD

FIG. 14

16/28





FIG. 15

17/28



Group	Median Survival	P Value (Log-rank test)
Vehicle	45 days	
Compound 2 (5 mg/kg)	58 days	0.0018
Compound 2 (2 mg/kg)	52 days	0.018
Compound 2 (1 mg/kg)	52 days	0.0088
Compound 2 (0.5 mg/kg)	50 days	NS
Temozolomide	TBD	TBD

FIG. 16

18/28




FIG. 17


Compound 1 Dose (mg/kg)

FIG. 18

19/28

Tumor Growth Delay (Median time to endpoint versus control group)

Compound1 Dose (mg/kg)	% Tumor Growth Inhibition (Day 12)	% Tumor Growth Delay
0	-	-
0.3	-8	4
3	44	25
10	73	76

* statistically significant

FIG. 19




FIG. 20


20/28

Age Mean (Range)	50	25–80
Gender (n,%)		
M	9	32
F	19	68
ECOG (n, %)		
0	16	57
1	11	39
2	1	4
Tumor Type (n, %)		
CRC	6	21
Breast	3	11
Pancreas	3	11
NSCLC	2	7
GBM	2	7
HCC	2	7
Salivary	2	7
Other* (1 each)	8	29
No. Prior Therapis (n, %)		
1–3	14	50
>3	14	50

FIG. 21

21/28

Part A: Accelerated (1+5) Dose Escalation



DLT Definition:

grade ≥ 3 Comp. 1 related AE, except rash ≤ 4 days, diarrhea/vomiting < 3 days
 fasting hyperglycemia: grade 2 > 14 days, grade 3 > 4 days
 heme: F&N, grade 4 ANC or platelets > 7 days, grade 3/4 platelets with bleeding
 liver: grade 3 not associated with PD in liver, and grade 4

FIG. 22

22/28




FIG. 23




FIG. 24

24/28




FIG. 25

25/28




FIG. 26

26/28




FIG. 27B




FIG. 27A




FIG. 28

28/28

Treatment Cycles		C1	C2	C3	C4	C5	C6	C7	CB-C12
7.5 mg	1-001	Escopagus		PD					
15 mg	1-002	Breast						SD	
15 mg	1-007	GBM		PD					
30 mg	1-003	NSCLC		SD					
30 mg	2-002	Pancreas	PD (Hyperglycemia DLT)						
30 mg	1-004	Renal	PD						
30-45 mg	1-005	Breast							PR
30 mg	2-004	Lelomyosarcoma		PD					
30 mg	1-006	HCC		NE					
30 mg	1-010	HCC		PD					
30/45 mg	1-008	Salivary						SD	
45 mg	2-005	CRC		PD					
45 mg	2-006	Paraganglioma		PD					
45 mg	1-009	Neuroendocrine		PD					
45 mg	2-007	Pancreas						SD (Rash DLT)	
45 mg	2-008	MM	NE						
45 mg	1-012	NSCLC						SD	
45 mg	2-009	CRC	NE						
45 mg	2-015	Adenocystic		SD					
45 mg	2-016	GBM	NE						
60/45 mg	2-010	CRC		PD (Fatigue DLT)					
60 mg	2-011	CRC	NE						
60/45 mg	1-013	CRC		NE					
60 mg	1-014	Pancreas		PD					
60/30 mg	1-017	Breast		PD (Mucositis DLT)					
60/45 mg	2-013	Adrenal						SD	
60 mg	2-014	CRC		SD					

*Patient 2-01 excluded
(protocol violation after single dose)

FIG. 29