

SULFONYL-QUINOLINE DERIVATIVES

FIELD OF THE INVENTION

The present invention relates to new mGluR1 and mGluR5 receptor subtype preferring ligands of formula (I) and/or salts and/or hydrates and/or solvates thereof, to the processes and intermediates for their preparation, to pharmaceutical compositions containing these compounds and to their use in therapy and/or prevention of a condition which requires modulation of mGluR1 and mGluR5 receptors.

BACKGROUND OF THE INVENTION

A major excitatory neurotransmitter in the mammalian central nervous system (CNS) is the glutamate molecule, which binds to neurons, thereby activating cell surface receptors. These receptors can be divided into two major classes, ionotropic and metabotropic glutamate receptors, based on the structural features of the receptor proteins, the means by which the receptors transduce signals into the cell, and pharmacological profiles.

The metabotropic glutamate receptors (mGluRs) are G protein-coupled receptors that activate a variety of intracellular second messenger systems following the binding of glutamate. Activation of mGluRs in intact mammalian neurons elicits one or more of the following responses: activation of phospholipase C; increases in phosphoinositide (PI) hydrolysis; intracellular calcium release; activation of phospholipase D; activation or inhibition of adenylyl cyclase; increases or decreases in the formation of cyclic adenosine monophosphate (cAMP); activation of guanylyl cyclase; increases in the formation of cyclic guanosine monophosphate (cGMP); activation of phospholipase A₂; increases in arachidonic acid release; and increases or decreases in the activity of voltage- and ligand-gated ion channels. (*Trends Pharmacol. Sci.*, **1993**, 14, 13; *Neurochem. Int.*, **1994**, 24, 439; *Neuropharmacology*, **1995**, 34, 1; *Prog. Neurobiol.*, **1999**, 59, 55).

Eight distinct mGluR subtypes, termed mGluR1 through mGluR8, have been identified by molecular cloning (*Neuron*, **1994**, 13, 1031; *Neuropharmacology*, **1995**, 34, 1; *J. Med. Chem.*, **1995**, 38, 1417). Further receptor diversity occurs via expression of alternatively spliced forms of certain mGluR subtypes (*PNAS*, **1992**, 89, 10331; *BBRC*, **1994**, 199, 1136; *J. Neurosci.*, **1995**, 15, 3970).

Metabotropic glutamate receptor subtypes may be subdivided into three groups, Group I, Group II, and Group III mGluRs, based on amino acid sequence homology, the second

messenger systems utilized by the receptors, and by their pharmacological characteristics. Group I mGluR comprises mGluR1, mGluR5 and their alternatively spliced variants.

Attempts at elucidating the physiological roles of Group I mGluRs suggest that activation of these receptors elicits neuronal excitation. Evidence indicates that this excitation is due to direct activation of postsynaptic mGluRs, but it also has been suggested that activation of presynaptic mGluRs occurs, resulting in increased neurotransmitter release (*Trends Pharmacol. Sci.*, **1992**, 15, 92; *Neurochem. Int.*, **1994**, 24, 439; *Neuropharmacology*, **1995**, 34, 1; *Trends Pharmacol. Sci.*, **1994**, 15, 33).

Metabotropic glutamate receptors have been implicated in a number of normal processes in the mammalian CNS. Activation of mGluRs has been shown to be required for induction of hippocampal long-term potentiation and cerebellar long-term depression (*Nature*, **1993**, 363, 347; *Nature*, **1994**, 368, 740; *Cell*, **1994**, 79, 365; *Cell*, **1994**, 79, 377). A role for mGluR activation in nociception and analgesia also has been demonstrated (*Neuroreport*, **1993**, 4, 879; *Brain Res.*, **1999**, 871, 223).

Group I metabotropic glutamate receptors and mGluR5 in particular, have been suggested to play roles in a variety of pathophysiological processes and disorders affecting the CNS. These include stroke, head trauma, anoxic and ischemic injuries, hypoglycemia, epilepsy, neurodegenerative disorders such as Alzheimer's disease, acute and chronic pain, substance abuse and withdrawal, obesity and gastroesophageal reflux disease (GERD) and irritable bowel syndrome (*Trends Pharmacol. Sci.*, **1993**, 14, 13; *Life Sci.*, **1994**, 54, 135; *Ann. Rev. Neurosci.*, **1994**, 17, 31; *Neuropharmacology*, **1995**, 34, 1; *J. Med. Chem.*, **1995**, 38, 1417; *Trends Pharmacol. Sci.*, **2001**, 22, 331; *Curr. Opin. Pharmacol.*, **2002**, 2, 43; *Pain*, **2002**, 98, 1, *Curr Top Med Chem.*, **2005**, 5(9):897-911). Much of the pathology in these conditions is thought to be due to excessive glutamate-induced excitation of CNS neurons. As Group I mGluRs appear to increase glutamate-mediated neuronal excitation via postsynaptic mechanisms and enhanced presynaptic glutamate release, their activation probably contributes to the pathology. Accordingly, selective antagonists of Group I mGluR receptors could be therapeutically beneficial, specifically as neuroprotective agents, analgesics or anticonvulsants.

WO 2006 120573 relates to a method for inhibiting the proliferation of cancer cells in mammal, comprising administering to said mammal a therapeutically effective amount of heterocyclic mono-N-oxides, among them N-oxides of compounds of formula (I) of the present invention.

The preparation of quinolines was stated by the application of general method that can be found in WO 2005 070890, *J. Med. Chem.*, **2003**, 46, 49 and *J. Med. Chem.*, **2005**, 48, 1107. In the cited publications 4-amino-3-cyano-quinoline derivatives are prepared by condensation of anilines with 2-cyano-3-ethoxy-acrylic acid ethyl ester followed by the ring closure of the obtained intermediates. Thermal ring closure afforded 3-cyano-4-hydroxy-quinoline derivatives that was converted into 3-cyano-4-chloro-quinoline derivatives by phosphorous(V) oxychloride. Ring closure with phosphorous(V) oxyhalogenides directly resulted in 3-cyano-4-halogen-quinoline derivatives. From these intermediates no 3-arylsulfonyl-4-aryl-quinoline derivatives of formula (I) of the present invention were prepared or characterized by physical properties either in WO 2006 120573 or in the cited publications.


WO 2005 58834 provides novel quinoline derivatives for use in treating liver X receptor (LXR) mediated diseases particularly multiple sclerosis, rheumatoid arthritis, inflammatory bowel disease and atherosclerosis, which compounds suppress Th-1 type lymphokine production, resulting in increased HDL levels, and cholesterol metabolism. The general formula of WO 2005058834 covers some of the compounds of formula (I) of the present invention, but only 3-benzenesulfonyl-4-phenyl-8-trifluoromethyl-quinoline was prepared by the reaction of the appropriate aniline derivative (described in Scheme 9 of WO 2005 58834) with 1,2-bis(benzenesulfonyl)-ethylen. This compound is not covered by the general formula of the present invention and proved to be inactive on mGluR1 and mGluR5 receptors.

WO 2005 30129 relates to compounds useful as potassium channel inhibitors. Compounds in this class may be useful as Kv1.5 antagonists for treating and preventing cardiac arrhythmias, and the like, and as Kv1.3 inhibitors for treatment of immunosuppression, autoimmune diseases, and the like. The general formula of WO 2005 30129 covers some of the compounds of formula (I) of the present invention, but no 3-arylsulfonyl-4-aryl-quinoline derivatives of formula (I) of the present invention only 2-substituted-6-methoxy-quinolin-3-carbonitrile derivatives were prepared or characterized by physical properties.

The compounds mentioned in the above publications are not declared or even not suggested having activity on the mGluR1 and mGluR5 receptors.

SUMMARY OF THE INVENTION

The present invention relates to new mGluR1 and mGluR5 receptor subtype preferring ligands of formula (I):



(I)

wherein


Ar₁ represents an optionally substituted phenyl or heteroaryl group;

Ar₂ represents a substituted phenyl or an optionally substituted heteroaryl group;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, halogen, cyano, alkyl, alkoxy, hydroxy, trifluoromethyl, amino, alkylamino, dialkylamino, aminomethyl, alkylaminomethyl, dialkylaminomethyl, and/or salts and/or hydrates and/or solvates thereof, to the processes and intermediates for producing the same, to pharmaceutical compositions containing the same and to their use in therapy and/or prevention of pathological conditions which require the modulation of mGluR1 and mGluR5 receptors such as neurological disorders, psychiatric disorders, acute and chronic pain and neuromuscular dysfunctions of the lower urinary tract.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to new mGluR1 and mGluR5 receptor subtype preferring ligands of formula (I):



(I)

wherein

Ar₁ represents an optionally substituted phenyl or heteroaryl group;

Ar₂ represents a substituted phenyl or an optionally substituted heteroaryl group;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, halogen, cyano, alkyl, alkoxy, hydroxy, trifluoromethyl, amino, alkylamino, dialkylamino, aminomethyl, alkylaminomethyl, dialkylaminomethyl, and/or salts and/or hydrates and/or solvates thereof.

When Ar₁ represents phenyl, the phenyl group may be optionally substituted with one or more substituent(s) selected from hydrogen, halogen, cyano, alkyl, alkoxy, trifluoromethyl, dialkylamino. When Ar₂ represents phenyl, the phenyl group is substituted with one or more substituent(s) selected from halogen, cyano, alkyl, alkoxy, trifluoromethyl, dialkylamino.

When Ar₁ and/or Ar₂ represent heteroaryl, the heteroaryl group may be an aromatic 5 to 6 membered heterocyclic ring containing 1 to 2 heteroatom(s) selected from O, N or S, such as pyridyl, thiazolyl, oxazolyl. The heteroaryl group may be optionally substituted with one or more substituent(s) selected from hydrogen, alkyl, alkoxy, halogen.

When R₁ and/or R₂ and/or R₃ and/or R₄ represent represents alkyl, the alkyl group contains 1 to 4 carbon atom(s) with straight or branched chain.

When R₁ and/or R₂ and/or R₃ and/or R₄ represent alkoxy, alkylamino, dialkylamino, alkylaminomethyl, dialkylaminomethyl the alkyl moiety inside the group contains 1 to 4 carbon atom(s) with straight or branched chain.

The term "halogen" includes fluorine, chlorine, bromine and iodine atoms.

In this specification the term "halo" may be fluoro, chloro, bromo or iodo.

Compounds of formula (I) contain basic function(s) so may form salts with acids. The invention relates also to the salts of compounds of formula (I) formed with acids, especially the salts formed with pharmaceutically acceptable acids. The meaning of compound of formula (I) is either the free base or the salt even if it is not referred separately.

Both organic and inorganic acids can be used for the formation of acid addition salts. Suitable inorganic acids can be for example hydrochloric acid, sulfuric acid, nitric acid and phosphoric acid. Representatives of monovalent organic acids can be for example formic acid, acetic acid, propionic acid, and different butyric acids, valeric acids and capric acids. Representatives of bivalent organic acids can be for example oxalic acid, malonic acid, maleic acid, fumaric acid and succinic acid. Other organic acids can also be used, such as hydroxy acids for example citric acid, tartaric acid, or aromatic carboxylic acids for example benzoic acid or salicylic acid, as well as aliphatic and aromatic sulfonic acids for example methanesulfonic acid, naphthalenesulfonic acid and p-toluenesulfonic acid. Especially valuable group of the acid addition salts is in which the acid component itself is

physiologically acceptable and does not have therapeutical effect in the applied dose or it does not have unfavourable influence on the effect of the active ingredient. These acid addition salts are pharmaceutically acceptable acid addition salts. The reason why acid addition salts, which do not belong to the pharmaceutically acceptable acid addition salts belong to the present invention is, that in given case they can be advantageous in the purification and isolation of the desired compounds.

Solvates and/or hydrates of compounds of formula (I) are also included within the scope of the invention.

Preferred compounds of the invention are those compounds of formula (I), wherein

Ar₁ represents phenyl or heteroaryl group, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

Ar₂ represents phenyl, substituted with one or more substituent(s) selected from fluoro, chloro, cyano, methyl, methoxy; or

heteroaryl, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy, hydroxy, trifluoromethyl, amino, methylamino, dimethylamino, aminomethyl, methylaminomethyl, dimethylaminomethyl, and/or salts and/or hydrates and/or solvates thereof.

Particular preferred compounds of the invention are those compounds of formula (I), wherein

Ar₁ represents phenyl, pyridyl, thienyl or oxazolyl group, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

Ar₂ represents phenyl, substituted with one or more substituent(s) selected from fluoro, chloro, cyano, methyl, methoxy; or

pyridyl, thienyl or oxazolyl, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy, hydroxy, trifluoromethyl, amino, methylamino, dimethylamino, aminomethyl, methylaminomethyl, dimethylaminomethyl, and/or salts and/or hydrates and/or solvates thereof.

Especially important compounds of formula (I) of the present invention are the following:

4-(4-chloro-phenyl)-3-(4-methyl-benzenesulfonyl)-quinoline,
7-chloro-3-(4-chloro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
8-chloro-4-(3-chloro-phenyl)-3-(3,4-dichloro-benzenesulfonyl)-quinoline,
7-fluoro-3-(4-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
4-(4-chloro-phenyl)-7-fluoro-3-(4-methoxy-benzenesulfonyl)-quinoline,
7-fluoro-3-(4-methoxy-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
7-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
7-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-4-(3-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-8-fluoro-3-(4-fluoro-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-8-fluoro-3-(4-fluoro-benzenesulfonyl)-quinoline,
8-fluoro-3-(4-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
8-fluoro-3-(4-fluoro-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
4-(4-chloro-phenyl)-6-fluoro-3-(4-methoxy-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-3-(3,4-dimethyl-benzenesulfonyl)-6-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(4-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-7-fluoro-quinoline,
3-(3-cyano-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
7-fluoro-3-(4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
4-(4-chloro-phenyl)-7-fluoro-3-(3-fluoro-benzenesulfonyl)-quinoline,
4-(3-chloro-phenyl)-7-fluoro-3-(3-methoxy-benzenesulfonyl)-quinoline,
7-fluoro-4-(4-fluoro-phenyl)-3-(3-methoxy-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-chloro-4-methoxy-benzenesulfonyl)-7-fluoro-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,

8-fluoro-3-(4-fluoro-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-8-fluoro-3-(4-methoxy-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-8-fluoro-3-(4-methoxy-benzenesulfonyl)-quinoline,
8-fluoro-4-(4-fluoro-phenyl)-3-(4-methoxy-benzenesulfonyl)-quinoline,
8-fluoro-3-(4-methoxy-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
8-fluoro-3-(4-methoxy-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-8-fluoro-3-(3-fluoro-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-8-fluoro-3-(3-fluoro-benzenesulfonyl)-quinoline,
8-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-6-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-6-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-6-fluoro-3-(3-methoxy-benzenesulfonyl)-quinoline,
6-fluoro-4-(4-fluoro-phenyl)-3-(3-methoxy-benzenesulfonyl)-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-6-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-6-fluoro-4-(3-methoxy-phenyl)-quinoline,
4-(4-chloro-phenyl)-7-fluoro-3-(4-fluoro-benzenesulfonyl)-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-4-(4-chloro-phenyl)-6-fluoro-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-7-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-7-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-quinoline,
7-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
7-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-4-(3-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-7-fluoro-quinoline,

3-(3-cyano-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-7-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3-chloro-4-methyl-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-7-fluoro-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-4-(3-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-quinoline,
8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(4-chloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-6-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-6-fluoro-4-(3-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-6-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-quinoline,

6-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
6-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
6-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
6-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-6-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-6-fluoro-quinoline,
3-(3-cyano-benzenesulfonyl)-6-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3-chloro-4-methyl-benzenesulfonyl)-4-(3-chloro-phenyl)-6-fluoro-quinoline,
3-(3-chloro-4-methyl-benzenesulfonyl)-4-(4-chloro-phenyl)-6-fluoro-quinoline,
3-(3-chloro-4-methyl-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-chloro-4-methyl-benzenesulfonyl)-6-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3-chloro-4-methyl-benzenesulfonyl)-6-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(4-chloro-phenyl)-6-fluoro-3-(3-methoxy-benzenesulfonyl)-quinoline,
6-fluoro-3-(3-methoxy-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-6-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-6-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-6-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-6-fluoro-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-6-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-6-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-7-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-7-fluoro-quinoline,
4-(3-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-quinoline,
4-(3-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-6-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-6-fluoro-quinoline,
6-fluoro-3-(4-methoxy-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
7-chloro-4-(4-chloro-phenyl)-3-(3,5-difluoro-benzenesulfonyl)-quinoline,
7-chloro-4-(4-chloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-quinoline,
7-chloro-4-(4-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-quinoline,
7-chloro-3-(3,5-dichloro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,

7-chloro-3-(4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3,4-difluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
6-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-7-fluoro-quinoline,
6-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-7-cyano-4-(2-fluoro-phenyl)-quinoline,
7-chloro-3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-8-fluoro-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
7-chloro-4-(4-chloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline.

3-(3-cyano-4-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-8-fluoro-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-8-fluoro-4-(2-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-8-fluoro-4-(2-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-4-fluoro-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-4-fluoro-benzenesulfonyl)-quinoline,
3-(3-cyano-5-fluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3-cyano-4-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-8-fluoro-quinoline,
7-chloro-3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-4-(2-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(2-fluoro-phenyl)-quinoline,

4-(4-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-8-fluoro-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-5-fluoro-benzenesulfonyl)-quinoline,
7-chloro-3-(3-cyano-benzenesulfonyl)-4-(2-fluoro-phenyl)-quinoline,
3-(3-cyano-5-fluoro-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(3-chloro-phenyl)-8-fluoro-quinoline,
3-(3-cyano-benzenesulfonyl)-7-fluoro-4-(2-fluoro-phenyl)-quinoline,
3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-7-fluoro-quinoline,
3-(3-cyano-4-fluoro-benzenesulfonyl)-4-(3,4-dichloro-phenyl)-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(2-fluoro-phenyl)-quinoline,
7-chloro-3-(3-cyano-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-4-fluoro-benzenesulfonyl)-7-fluoro-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-4-(3,5-difluoro-phenyl)-8-fluoro-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-8-fluoro-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-7-fluoro-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
7-chloro-3-(3-cyano-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-chloro-4-(3-chloro-phenyl)-3-(3-cyano-4-fluoro-benzenesulfonyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-4-(2-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-5-fluoro-benzenesulfonyl)-7-fluoro-quinoline,
3-(3-cyano-benzenesulfonyl)-4-(3,5-difluoro-phenyl)-7-fluoro-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-8-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-quinoline,
7-chloro-3-(3,4-dichloro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-quinoline,
3-(3,5-dicyano-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(3-chloro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-7-fluoro-quinoline,
7-chloro-4-(3-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-quinoline,
7-chloro-4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-8-fluoro-quinoline,
7-chloro-4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-quinoline,
7-chloro-3-(3,5-dichloro-benzenesulfonyl)-4-(2-fluoro-phenyl)-quinoline,
3-(3,5-dicyano-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-7-fluoro-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-7-fluoro-4-(2-fluoro-phenyl)-quinoline,

3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(3-chloro-phenyl)-7-fluoro-quinoline,
4-(3-chloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-quinoline,
4-(3,4-dichloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-4-(2-fluoro-phenyl)-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(3,5-difluoro-phenyl)-quinoline,
7-chloro-4-(3-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-quinoline,
7-amino-3-(3,4-difluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-quinoline,
3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-quinoline,
3-(3-cyano-5-fluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(4-chloro-phenyl)-3-(3-cyano-5-fluoro-benzenesulfonyl)-7-fluoro-quinoline,
7-chloro-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-quinoline,
7-chloro-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-7-fluoro-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-quinoline,
7-amino-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-amino-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline.

Pharmaceutical formulations

The invention also relates to the pharmaceutical compositions containing the compounds of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof as active ingredient and one or more physiologically acceptable carriers.

The compounds of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof may be administered by any convenient method, for example by oral, parenteral (including subcutaneous, intramuscular, and intravenous), buccal, sublingual, nasal, rectal or transdermal administration and the pharmaceutical compositions adapted accordingly.

The compounds of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof which are active when given orally can be formulated as liquids or solids, for example syrups, suspensions or emulsions, tablets, capsules and lozenges.

A liquid formulation of the compounds of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof generally consist of a suspension or solution of the compound of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof in a suitable liquid carrier(s) for example an aqueous solvent, such as water and ethanol or glycerine, or a non-aqueous solvent, such as polyethylene glycol or an oil. The formulation may also contain a suspending agent, preservative, flavouring or colouring agent.

A composition in the solid form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of solid carriers include lactose, terra alba, sucrose, talc, gelatine, agar, pectin, acacia, magnesium stearate, stearic acid etc. Optionally, tablets may be coated by standard aqueous or nonaqueous techniques.

A composition in the solid form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then these are filled into a hard gelatine capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then is filled into a soft gelatine capsule.

Typical parenteral compositions consist of a solution or suspension of the compound of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

Compositions of the present invention for nasal administration containing a compound of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof may conveniently be formulated as aerosols, drops, gels and powders. Aerosol formulations of the present invention typically comprise a solution or fine suspension of the compound of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates in a physiologically acceptable aqueous or non-aqueous solvent and are usually presented in a single or multidose quantities in sterile form in a sealed container, which can take the form of

a cartridge or refill for use with an atomizing device. Alternatively, the sealed container may be a unitary dispensing device, such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve which is intended for disposal once the contents of the container have been exhausted. If the dosage form comprises an aerosol dispenser, it will contain a propellant which can be a compressed gas, such as compressed air or an organic propellant, such as a fluorochlorohydrocarbon. The aerosol dosages form can also take the form of a pump-atomiser.

Compositions of the present invention containing a compound of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates are suitable for buccal or sublingual administration including tablets, lozenges and pastilles, wherein the active ingredient is formulated with a carrier, such as sugar and acacia, tragacanth, or gelatine, glycerin etc.

Compositions of the present invention containing a compound of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof for rectal administration are conveniently in the form of suppositories containing a conventional suppository base, such as cocoa butter and other materials commonly used in the art. The suppositories may be conveniently formed by first admixing the composition with the softened or melted carrier(s) followed by chilling and shaping in moulds.

Compositions of the present invention containing a compound of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof for transdermal administration include ointments, gels and patches.

The compositions of the present invention containing a compound of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof is preferably in the unit dose form, such as tablet, capsule or ampoule.

Each dosage unit of the present invention for oral administration contains preferably from 0.1 to 500 mg of a compound of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof calculated as a free base.

Each dosage unit of the present invention for parenteral administration contains preferably from 0.1 to 500 mg of a compound of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof calculated as a free base.

The compounds of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates thereof can normally be administered in a daily dosage regimen. In the treatment of mGluR1 and mGluR5 mediated disorders, such as schizophrenia, anxiety,

depression, panic, bipolar disorders, and circadian disorders or chronic and acute pain disorders the dosage levels from about 0,01 mg/kg to about 140 mg/kg of body weight per day are useful or alternatively about 0.5 mg to about 7 g per patient per day.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for the oral administration to humans may conveniently contain from about 0.5 mg to about 5 g of active agent, compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95 percent of the total composition. Unit dosage forms will generally contain between from about 1 mg to about 1000 mg of the active ingredient, typically 25 mg, 50 mg, 100 mg, 200 mg, 250-300 mg, 400 mg, 500 mg, 600 mg, 800 mg or 1000 mg.

It is understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

Medical use

The compounds of formula (I) and/or physiologically acceptable salts and/or hydrates and/or solvates of the present invention have been found to exhibit biological activity at mGluR1 and mGluR5 receptors and are expected to be useful in the treatment of mGluR1 and mGluR5 mediated disorders.

It has been found that the compounds according to the present invention or salts thereof, exhibit a high degree of potency and selectivity for individual metabotropic glutamate receptor (mGluR) subtypes. In particular there are compounds according to the present invention that are potent and selective for mGluR1 and mGluR5 receptors. Accordingly, the compounds of the present invention are expected to be useful in the prevention and/or treatment of conditions associated with excitatory activation of mGluR1 and mGluR5 receptor and for inhibiting neuronal damage caused by excitatory activation of mGluR1 and mGluR5 receptor. The compounds may be used to produce an inhibitory effect of mGluR1 and mGluR5, in mammals, including human.

Thus, it is expected that the compounds of the invention are well suited for the prevention and/or treatment of mGluR1 and mGluR5 receptor-mediated disorders such as

acute and chronic neurological and psychiatric disorders, chronic and acute pain disorders and neuromuscular dysfunctions of the lower urinary tract.

The dose required for the therapeutic or preventive treatment of a particular disorder will necessarily be varied depending on the host treated and the route of administration.

The invention relates to compounds of formula (I) as defined hereinbefore, for use in therapy.

The invention relates to compounds of formula (I) as defined hereinbefore, for use in prevention and/or treatment of mGluR1 and mGluR5 receptor-mediated disorders.

The invention relates to compounds of formula (I) as defined hereinbefore, for use in prevention and/or treatment of neurological disorders.

The invention relates to compounds of formula (I) as defined hereinbefore, for use in prevention and/or treatment of psychiatric disorders.

The invention relates to compounds of formula (I) as defined hereinbefore, for use in prevention and/or treatment of chronic and acute pain disorders.

The invention relates to compounds of formula (I) as defined hereinbefore, for use in prevention and/or treatment of neuromuscular dysfunctions of the lower urinary tract.

The invention relates to compounds of formula (I) as defined hereinbefore, for use in prevention and/or treatment of pain related to migraine, inflammatory pain, neuropathic pain disorders such as diabetic neuropathies, arthritis and rheumatoid diseases, low back pain, post-operative pain and pain associated with various conditions including angina, in renal or biliary colic, menstruation, migraine and gout.

The invention relates to compounds of formula (I) as defined hereinbefore, for use in prevention and/or treatment of Alzheimer's disease senile dementia, AIDS-induced dementia Parkinson's disease, amyotrophic lateral sclerosis, Huntington's Chorea, migraine, epilepsy, schizophrenia, depression, anxiety, acute anxiety, obesity, obsessive compulsive disorder, ophthalmological disorders such as retinopathies, diabetic retinopathies, glaucoma, auditory neuropathic disorders such as tinnitus, chemotherapy induced neuropathies, post-herpetic neuralgia and trigeminal neuralgia, tolerance, substance abuse and withdrawal, Fragile X, autism, mental retardation, schizophrenia and Down's Syndrome.

The invention relates to compounds of formula (I) as defined hereinbefore, for use in prevention and/or treatment of stroke, head trauma, anoxic and ischemic injuries, hypoglycemia, cardiovascular diseases and epilepsy.

The compounds are also well suited for the treatment of neuromuscular dysfunction of the lower urinary tract, such as urinary urgency, overactive bladder, greater urinary frequency, reduced urinary compliance, cystitis, incontinence, enuresis and dysuria.

The compounds are also well suited for the treatment of gastrointestinal disorders, such as transient lower esophageal sphincter relaxation (TLESR), gastrointestinal reflux disease and irritable bowel syndrome.

The present invention relates also to the use of a compound of formula (I) as defined hereinbefore, in the manufacture of a medicament for the prevention and/or treatment of mGluR1 and mGluR5 receptor-mediated disorders and any disorder listed above.

The invention also provides a method of treatment and/or prevention of mGluR1 and mGluR5 receptor mediated disorders and any disorder listed above, in a patient suffering from, or at risk of, said condition, which comprises administering to the patient an effective amount of a compound of formula (I), as hereinbefore defined.

In the context of the present specification, the term "therapy" includes treatment as well as prevention, unless there are specific indications to the contrary. The terms "therapeutic" and "therapeutically" should be construed accordingly.

In this specification, unless stated otherwise, the term "antagonist" means a compound that by any means, partly or completely blocks the transduction pathway leading to the production of a response by the ligand.


The term "disorder", unless stated otherwise, means any condition and disease associated with metabotropic glutamate receptor activity.

Methods of preparation

The reactions are carried out by conventional methods. If necessary, protecting groups can be used for protect functional groups as it apparent to one skilled in the art.

According to the present invention a process for the preparation of compounds of formula (I):

19



(I)


wherein

Ar₁ represents an optionally substituted phenyl or heteroaryl group;

Ar₂ represents a substituted phenyl or an optionally substituted heteroaryl group;


R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, halogen, cyano, alkyl, alkoxy, hydroxy, trifluoromethyl, amino, alkylamino, dialkylamino, aminomethyl, alkylaminomethyl, dialkylaminomethyl,

reacting a compound of formula (II):




(II)

wherein R₁, R₂, R₃ and R₄ are as defined above for a compound of formula (I), with a compound of formula (III):



(III)


wherein M is selected from alkali metals or alkaline-earth metals, Ar₁ is as defined above for compounds of formula (I), to give a compound of formula (IV):



(IV)


wherein R₁, R₂, R₃, R₄ and Ar₁ are as defined above for compounds of formula (I), thereafter oxidizing a compound of formula (IV) to obtain a compound of formula (V):

20




(V)

wherein R_1 , R_2 , R_3 , R_4 and Ar_1 are as defined above for compounds of formula (I), thereafter oxidizing a compound of formula (V) to obtain a compound of formula (VI):




(VI)

wherein R_1 , R_2 , R_3 , R_4 and Ar_1 are as defined above for compounds of formula (I), thereafter converting the obtained compound of formula (VI) to a compound of formula (VII):



(VII)

wherein X is selected from chloro, bromo, benzenesulfonyloxy, 4-fluoro-benzenesulfonyloxy, 4-methyl-benzenesulfonyloxy, methanesulfonyloxy or trifluoromethanesulfonyloxy groups and R_1 , R_2 , R_3 , R_4 and Ar_1 are as defined above for compounds of formula (I), thereafter reacting the obtained compound of formula (VII) with a boronic acid derivative of formula (VIII):



(VIII)


in the presence of base (e.g. sodium carbonate) and catalyst (e.g. tetrakis(triphenylphosphine)-palladium(0)) in a solvent, and optionally thereafter forming salts and/or hydrates and/or solvates of compounds of formula (I).

According to Scheme 1, 3-Bromo-quinolin-4-ol derivatives of formula (II) can be reacted by alkali or alkaline-earth metal salts (e.g. sodium salt) of thiophenols of formula (III) to provide a compound of formula (IV) (e.g. *Bioorg. Med. Chem. Lett.*, **2001**, 9, 1141). Advantageously the reaction can be carried out in the presence of palladium catalyst and under microwave conditions. 3-Bromo-quinolin-4-ol derivatives of formula (II) are known (e.g. 3-bromo-6-chloro-quinolin-4-ol: *J. Chem. Soc.*, **1950**, 384; 3-bromo-7-trifluoromethyl-quinolin-4-ol: *Synthesis*, **1977**, 865) or can be synthesised by conventional methods. Alkali or alkaline-earth metal salts of thiophenols of formula (III) can be purchased or can be prepared by conventional methods.


Oxidation of 3-arenesulfanyl-quinolin-4-ol derivatives of formula (IV) can be accomplished in a suitable acid (e.g. trifluoroacetic acid) with hydrogen peroxid to give sulfoxides of formula (V) and sulfones of formula (VI), respectively.

Conversion of 3-arenesulfonyl-quinolin-4-ol derivatives of formula (VI) to compounds of formula (VII) can be carried out by known halogenation methods with suitable reagents (e.g. POCl_3 , SOCl_2 , PCl_5 , POBr_3 , PBr_3), or by acylation with the appropriate sulfonic acid halogenid or sulfonic acid anhydride derivatives.


Scheme 1




Another method to prepare intermediates of formula (VI) comprises reacting a compound of formula (IX):




wherein Ar_1 is as defined above for a compound of formula (I), with α -halogen-acetic acid esters of formula (X):




wherein Hlg is halogen and R_5 is an ethyl or methyl group, to obtain a compound of formula (XI):




wherein Ar_1 is as defined above for a compound of formula (I) and R_5 is as defined above for compounds of formula (X); reacting a compound of formula (XI) with a trialkyl orthoformate of formula (XII):



wherein R_6 is an ethyl or methyl group, to obtain a compound of formula (XIII):




wherein Ar_1 is as defined above for a compound of formula (I), R_5 is as defined above for compounds of formula (X) and R_6 is as defined above for compounds of formula (XII); reacting a compound of formula (XIII) with an aniline derivative of formula (XIV):



wherein R_1 , R_2 , R_3 and R_4 are as defined above for a compound of formula (I), to obtain a compound of formula (VI).

23




(VI)

wherein R_1 , R_2 , R_3 and R_4 are as defined above for a compound of formula (I), thereafter converting the obtained compound of formula (VI) to a compound of formula (I) as described above.

According to Scheme 2 compounds of formula (IX) are reacted with α -halogen-acetic acid esters of formula (X) in a suitable solvent (e.g. DMF, water). Compounds of formula (IX) can be purchased or can be prepared from the appropriate benzenesulfonyl chloride derivatives by known methods (e.g. *Org. Lett.*, **2003**, 5(21), 3895). Compounds of formula (XIII) can be prepared by the reaction of compounds of formula (XI) and compounds of formula (XII) in the presence of acetic anhydride [*J. Org. Chem. USSR (Engl. Transl.*, **1980**, 16(7), 1275; *Zh. Org. Khim.*, **1980**, 16(7), 1483]. Reaction of compounds of formula (XIII) with aniline derivatives of compounds of formula (XIV) gives benzenesulfonyl-phenylamino-acrylic acid esters [e.g. *J. Org. Chem. USSR (Engl. Transl.*, **1980**, 16(7), 1275; *Zh. Org. Khim.*, **1980**, 16(7), 1483-1487] that can be *in situ* convert into quinolin-4-ol derivatives of formula (VI) (see analogous reaction: *J. Chem. Soc. Perkin Trans.*, 1, **1994**, 4, 387-392).

Scheme 2



Compounds of formula (IV), compounds of formula (V), compounds of formula (VI) and compounds of formula (VII) and/or enantiomers and/or racemates and/or diastereomers and/or pharmaceutically acceptable salts thereof formed with acids or bases are new.

Compounds of formula (I) contain basic function(s) so can be transformed into the salts thereof with acids and/or can be liberated from the obtained acid addition salts by treatment with a base.

Compounds of formula (I) can be transformed into hydrates and/or solvates.

The compounds of formula (I) can optionally be interconverted to a different compound of formula (I) by conventional synthetic methods.

Biological test methods

MGluR1 receptor binding test

MGluR1 receptor binding testes were performed according to modified method of Lavreysen et al. (Mol.Pharm., 2003, 63, 1082). Based on the high homology between the human and rat mGluR1 receptors, rat cerebellar membrane preparation was used to determine the binding characteristics of reference compounds and novel compounds to the rat mGluR1. As radioligand [³H]R214127 (3 nM) was used and the nonspecific binding was determined in the presence of 1 μ M of R214127.

IC-50 values were determined from displacement curves by nonlinear regression analysis and were converted by equation method of Cheng and Prusoff (Biochem. Pharmacol., 1973, 22, 3099) to K_i values.

MGluR5 receptor binding tests

MGluR5 receptor binding was determined according to Gasparini et.al. (Bioorg. Med. Chem. Lett. 2000, 12:407-409) with modifications. Rat cerebro-cortical membrane preparation was used to determine the binding characteristics of reference compounds and novel compounds to the rat mGluR5. The A18 cell line expressing hmGluR5a (purchased from Euroscreen) was used to determine binding characteristics of the chemical compounds to the human mGluR5a receptor. As radioligand [³H]-M-MPEP (2 nM) was used. The nonspecific binding was determined in the presence of 10 μ M M-MPEP.

Assessment of functional activity

Cell cultures for native rat mGluR5 and mGluR1 receptors

Functional potency at native rat mGluR5 and mGluR1 receptors was estimated using primary neocortical cell cultures derived from 17 day old Charles River rat embryos and primary cerebellar cell cultures derived from 4-day old Wistar rats, respectively (for the details on the preparation of neural cell cultures see Johnson, M.I.; Bunge, R.P. (1992): Primary cell cultures of peripheral and central neurons and glia. In: Protocols for Neural Cell Culture, eds: Fedoroff, S.; Richardson A., The Humana Press Inc., 51-77). After isolation the cells were plated onto standard 96-well microplates and the cultures were maintained in an atmosphere of 95% air-5% CO₂ at 37 °C. The neocortical and cerebellar cultures were used for the calcium measurements after 5-7 and 3-4 days in vitro, respectively.

Cell cultures for recombinant human mGluR5a receptors

Chinese hamster ovary (CHO) cells stably expressing recombinant human mGluR5a (CHO-mGluR5a, purchased from Euroscreen) receptors were cultured in F12 medium containing 10% FCS, 1% antibiotic antimycotic solution, 400 µg/ml G418, 250 µg/ml zeocin, 5 µg/ml puromycin. Cells were kept at 37 °C in a humidified incubator in an atmosphere of 5% CO₂/95% air and were passaged three times a week. Cells were plated at 2.5-3.5×10⁴ cell/well on standard 96-well microplates, receptor expression was induced by adding 600 ng/ml doxycycline on the next day. The calcium measurements were carried out 16-24 hours after the addition of the inducing agent.

Fluorimetric measurement of cytosolic calcium concentration

Measurements of cytosolic calcium concentration ($[Ca^{2+}]_i$) were carried out on primary neocortical and cerebellar cultures, and on CHO-mGluR5a cells stably expressing human mGluR5a receptors. Cells were grown in standard 96-well microplates and before the measurement were loaded with a fluorescent Ca²⁺-sensitive dye, fluo-4/AM (2 µM): the neural cultures were loaded in their growth medium, CHO-mGluR5a cells were loaded in assay buffer (145 mM NaCl, 5 mM KCl, 2 mM MgCl₂, 2 mM CaCl₂, 10 mM HEPES, 20 mM D-glucose, 2 mM probenecid, pH=7.4) supplemented with 2 mM Na-pyruvate and 30 µg/ml glutamate-pyruvate transaminase (in case of CHO-mGluR5a cells these supplements were also present during the course of the $[Ca^{2+}]_i$ measurements). Loading was done by incubating the cells with 100 µl/well dye solution at 37 °C in a humidified incubator in an atmosphere of

5% CO₂/95% air for 40-120 min. To stop dye loading cells were washed twice with assay buffer. After washing, various concentrations of the test compounds (diluted in assay buffer from a DMSO or a dimethylformamide (DMF) stock solution, final DMSO/DMF concentration was <0.1%) or buffer were added to each well depending on the experimental setup. In the case of neocortical cultures the assay buffer also contained TTX (0.5 μ M, to suppress spontaneous oscillations of [Ca²⁺]_i, in the case of cerebellar cultures probenecid was substituted with sulfinpyrazone (0.25 mM).

After incubation at 37 °C for 10-20 min. baseline and agonist-evoked changes of [Ca²⁺]_i were measured column by column with a plate reader fluorimeter (FlexStation II, Molecular Devices). Excitation and detection of emission was carried out from the bottom of the plate. The whole measurement process was performed at 37 °C and was controlled by custom software. Inhibitory potency of the test compounds was assessed by measuring the reduction in the agonist-evoked [Ca²⁺]_i -elevation in the presence of different concentrations of the compounds. DHPG was used as agonist for all three cultures, the concentration was 20 and 100 μ M for neocortical and cerebellar cultures, respectively. In the case of CHO-mGluR5a cells DHPG was applied at an EC₈₀ concentration, the EC₈₀-values were derived from daily determined dose-response curves. Fluorescence data were expressed as $\Delta F/F$ (fluorescence change normalized to baseline).

All treatments on a single plate were measured in multiple wells. Data from all wells with the same treatment were averaged and the average values were used for analysis. Inhibitory potency of a compound at a single concentration point was expressed as percent inhibition of the control agonist response. Sigmoidal concentration-inhibition curves were fitted to the data (derived from at least three independent experiments) and IC₅₀-values were determined as the concentration that produces half of the maximal inhibition caused by the compound. Raw fluorescence data were analyzed using Soft Max Pro (Molecular Devices), curve fitting was done with GraphPad Prism.

Results

Compounds of formula (I) of the present invention showed affinity for both rat and human mGluR1 and mGluR5 receptors and proved to be functional antagonists that are they inhibited functional responses elicited by stimulation of mGluR5 receptors.

The invention is further illustrated by the following non-limiting examples.

Unless specifically stated otherwise, all operation were carried out at room temperature, that is at a temperature range of 18-25 °C. The course of reactions was followed by thin layer chromatography (TLC) and reaction times are given for illustration only. The structure of all intermediates and end products were elucidated by IR, NMR and MS spectroscopy. When given yields are for illustration only. When given, NMR data are in the form of delta (δ) values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as internal standard, using the indicated solvent. Conventional abbreviations are used for signal shape.

Examples

All starting materials are either commercially available or can be synthesized by different known methods described in the literature.

Example 1

4-(4-chloro-phenyl)-3-(4-methyl-benzenesulfonyl)-quinoline; Table I, compound 1

3-(4-Methyl benzenesulfanyl)-quinolin-4-ol

A mixture of 3-bromo-quinolin-4-ol (0.448 g, 2 mmol; *J. Am. Chem. Soc.* 1946, 68, 1229-1231), 4-methylbenzenethiol (0.30 g, 2.4 mmol), tetrakis-(triphenylphosphin)palladium(0) (0.115 g, 0.1 mmol), sodium-*tert*-butylate (0.23 g, 2.4 mmol) and DMF (2.0 ml) was stirred and irradiated at 142 °C for 3 hours in a 8-ml microwave vial. The solvent was evaporated *in vacuo* and the residue was purified by gradient silica gel flashchromatography (80 g silica gel, eluent A: chloroform, eluent B: chloroform : methanol = 95:5) to give 0.33 g of the title compound in 62% yield.

MS (EI) M^+ = 268.2

3-(4-Methyl-benzenesulfonyl)-quinolin-4-ol

To a mixture of 3-(4-Methyl-benzenesulfanyl)-quinolin-4-ol (0.25 g, 0.936 mmol) and trifluoroacetic acid (5.0 ml) a solution of hydrogen peroxide in trifluoroacetic acid (c = 3.0M, 3.7 ml) was added dropwise. The solution was stirred for 8 hours at room temperature. To the reaction mixture 6ml of water was added dropwise. The precipitate was filtered off, washed with water and dried *in vacuo* to give 0.24 g of the title compound in 86%.

MS (EI) M^+ = 300.1

4-Chloro-3-(4-methyl-benzenesulfonyl)-quinoline

A mixture of 3-(4-methyl-benzenesulfonyl)-quinolin-4-ol (0.24 g, 0.8 mmol) and phosphorus(V) oxychloride (15 ml) was refluxed for 5 hours. Phosphorus(V) oxychloride was distilled off, and the residue was poured onto ice. The slurry was stirred for 2 hours at 0-5 °C, neutralized with sodium carbonate and extracted with chloroform (50 ml). The organic layer was dried over anhydrous sodium sulfate, filtered and the solvent was removed *in vacuo* to give 0.23 g of the title compound in 90% yield..

MS (EI) $M^+ = 318.2$

4-(4-chloro-phenyl)-3-(4-methyl-benzenesulfonyl)-quinoline

A mixture of 4-chloro-3-(4-methyl-benzenesulfonyl)-quinolin-4-ol (0.2 g, 0.67 mmol) and 4-chlorophenylboronic acid (0.16 g, 1.0 mmol) in 8 ml dioxane was stirred for 20 hours at 90 °C with potassium carbonate (0.5 g, 3.6 mmol) and tetrakis-(triphenylphosphin)palladium(0) (0.04 g, 0.035 mmol). After cooling the mixture was concentrated *in vacuo*. The crude residue was purified by column chromatography on silica gel (Kieselgel 60, eluent: chloroform) to obtain 0.21 g of the title compound in 80% yield.

^1H NMR (500 MHz, DMSO- d_6): 9.61 (s, 1H); 8.21 (dm, $J = 8.6$ Hz, 1H); 7.98 (ddd, $J = 8.5, 6.8, 1.4$ Hz, 1H); 7.64 (ddd, $J = 8.5, 6.8, 1.2$ Hz, 1H); 7.49-7.44 (m, 2H); 7.37-7.32 (m, 2H); 7.30-7.24 (m, 3H); 7.04-6.98 (m, 2H), 2.36 (s, 3H).

MS (EI) $M^+ = 393.8$

4-(4-chloro-phenyl)-3-(4-methyl-benzenesulfonyl)-quinoline hydrochloride

4-(4-chloro-phenyl)-3-(4-methyl-benzenesulfonyl)-quinoline (40 mg, 0.102 mmol) was dissolved in ethyl acetate (1.5 ml) and a solution of hydrogen chloride in ethyl acetate ($c = 1.6\text{M}$, 0.14 ml, 0.224 mmol) was added dropwise to the solution. The solid was filtered off, washed with ethyl acetate and dried *in vacuo* to give 35 mg of the title compound in 80% yield.

^1H NMR (500 MHz, DMSO- d_6): 9.61 (s, 1H); 8.22 (dm, $J = 8.6$ Hz, 1H); 7.98 (ddd, $J = 8.6, 6.8, 1.4$ Hz, 1H); 7.64 (ddd, $J = 8.6, 6.8, 1.2$ Hz, 1H); 7.50-7.43 (m, 2H); 7.37-7.31 (m, 2H); 7.31-7.23 (m, 3H); 7.05-6.98 (m, 2H), 2.36 (s, 3H).

MS (EI) $M^+ = 393.8$

Example 2**(4-Chloro-benzenesulfonyl)-acetic acid methyl ester (intermediate)**

A mixture of methyl bromoacetate (11.25 ml, 116 mmol) and sodium 4-chloro-benzenesulfinate (25.2g, 116 mmol) in DMF (120 ml) was stirred and heated at 80°C for 2 h. The solution was diluted with water (360 ml). The separated oil was extracted with chloroform (200 ml) and washed with water (3 X 80 ml). The organic phase was evaporated *in vacuo* to obtain 22.4 g of the title compound in 77.7% yield. MS (EI) $M^+ = 249.1$.

Applying the above procedure the following compounds were prepared: e.g.

(3-chloro-benzenesulfonyl)-acetic acid methyl ester (MS (EI) $M^+ = 249.1$);
(3,5-dichloro-benzenesulfonyl)-acetic acid methyl ester (MS (EI) $M^+ = 283.1$);
(4-methoxy-benzenesulfonyl)-acetic acid methyl ester (MS (EI) $M^+ = 245.1$); (3-chloro-4-fluoro-benzenesulfonyl)-acetic acid methyl ester (MS (EI) $M^+ = 267.1$);
(3,5-difluoro-benzenesulfonyl)-acetic acid methyl ester (MS (EI) $M^+ = 251.1$);
(3-fluoro-benzenesulfonyl)-acetic acid methyl ester (MS (EI) $M^+ = 233.2$).

Example 3**2-(4-Chloro-benzenesulfonyl)-3-ethoxy-acrylic acid methyl ester (intermediate)**

The mixture of (4-chloro-benzenesulfonyl)-acetic acid methyl ester (22.4 g, 90 mmol), triethyl orthoformate (33.2 ml, 216 mmol) and acetic anhydride (19.1 ml, 203 mmol) was refluxed for 3 h with simultaneous distillation of ethanol, triethyl orthoformate and acetic anhydride, and then evaporated to dryness. The crude material (22.7 g, 82.8%) was used in the next step without purification. MS (EI) $M^+ = 305.1$.

Applying the above procedure the following compounds were prepared: e.g. 2-(3-chloro-benzenesulfonyl)-3-ethoxy-acrylic acid methyl ester (MS (EI) $M^+ = 305.1$); 2-(3-chloro-4-fluoro-benzenesulfonyl)-3-ethoxy-acrylic acid methyl ester (MS (EI) $M^+ = 323.1$); 2-(3,5-dichloro-benzenesulfonyl)-3-ethoxy-acrylic acid methyl ester (MS (EI) $M^+ = 340.1$); 2-(4-fluoro-benzenesulfonyl)-3-ethoxy-acrylic acid methyl ester (MS (EI) $M^+ = 289.1$); 2-(3-cyano-5 fluoro-benzenesulfonyl)-3-ethoxy-acrylic acid methyl ester (MS (EI) $M^+ = 314.2$).

Example 4**7-Chloro-3-(4-chloro-benzenesulfonyl)-quinolin-4-ol (intermediate)**

The mixture of 2-(4-chloro-benzenesulfonyl)-3-ethoxy-acrylic acid methyl ester (7.62 g, 25 mmol) and 3-chloroaniline (3.19 g, 25 mmol) in diphenyl ether (20 ml) was heated at near reflux for 1 h. After cooling the precipitate was filtered and washed with ether to obtain 4.25g of the title compound in 48.0% yield. MS (EI) $M^+ = 355.1$.

Applying the above procedure the following compounds were prepared: e.g. 8-chloro-3-(4-chloro-benzenesulfonyl)-quinolin-4-ol (MS (EI) $M^+ = 355.1$); 6-chloro-3-(4-fluoro-benzenesulfonyl)-quinolin-4-ol (MS (EI) $M^+ = 338.1$); 6-cyano-3-(4-fluoro-benzenesulfonyl)-quinolin-4-ol (MS (EI) $M^+ = 329.2$); 8-chloro-3-(3,4-dichloro-benzenesulfonyl)-quinolin-4-ol (MS (EI) $M^+ = 390.1$); 7-chloro-3-(3,5-difluoro-benzenesulfonyl)-quinolin-4-ol (MS (EI) $M^+ = 356.1$).

Example 5

3-(4-chloro-benzenesulfonyl)-4,7-dichloroquinoline (intermediate)

7-Chloro-3-(4-chloro-benzenesulfonyl)-4-hydroxyquinoline (4.25 g, 12 mmol) in phosphorus(V) oxychloride (5.6 ml, 60 mmol) was refluxed for 3 h. The reaction mixture was poured into water (50 ml) and was alkalized with 5M sodium hydroxide solution. After cooling the precipitate was filtered and washed with water to obtain 3.8 g of the title compound in 85.0% yield. MS (EI) $M^+ = 373.2$.

Applying the above procedure the following compounds were prepared: e.g. 3-(3-chloro-benzenesulfonyl)-4,6-dichloroquinoline (MS (EI) $M^+ = 373.2$); 3-(4-chloro-benzenesulfonyl)-4,8-dichloroquinoline (MS (EI) $M^+ = 373.2$); 4-chloro-6-cyano-3-(4-fluoro-benzenesulfonyl)-quinoline (MS (EI) $M^+ = 347.1$); 4-chloro-3-(4-chloro-benzenesulfonyl)-6-fluoro-quinoline (MS (EI) $M^+ = 357.1$).

Example 6

4-bromo-7-chloro-3-(4-chloro-benzenesulfonyl)-quinoline (intermediate)

A mixture of 7-chloro-3-(4-chloro-benzenesulfonyl)-quinolin-4-ol (0.5 g, 1.41 mmol) and phosphorus(V) oxybromide (1.2 g, 4.2 mmol) in chloroform (30 ml) and triethylamine (1 ml) was refluxed for 6 hours. The reaction mixture was diluted with water (30 ml) and the pH was adjusted to 10 with aqueous sodium hydroxide solution. The organic layer was separated, dried over sodium sulfate, filtered, and concentrated *in vacuo*. The obtained crude product

was purified by crystallization from diethyl ether to obtain 0.45 g of the title compound in 77% yield. MS (EI) M^+ = 418.1.

Applying the above procedure the following compounds were prepared: e.g. 4-bromo-3-(4-chloro-benzenesulfonyl)-6-fluoro-quinoline (MS (EI) M^+ = 347.1); 4-bromo-7-chloro-3-(3,5-difluoro-benzenesulfonyl)-quinoline (MS (EI) M^+ = 420.1); 4-bromo-7-chloro-3-(4-chloro-benzenesulfonyl)-6-fluoro-quinoline (MS (EI) M^+ = 436.1); 4-bromo-7-chloro-3-(4-fluoro-benzenesulfonyl)-quinoline (MS (EI) M^+ = 402.1); 4-bromo-7-chloro-3-(3-fluoro-benzenesulfonyl)-quinoline (MS (EI) M^+ = 402.1); 4-bromo-7-chloro-3-(3-cyano-benzenesulfonyl)-quinoline (MS (EI) M^+ = 409.2); 4-bromo-7-chloro-3-(3-cyano-5-fluoro-benzenesulfonyl)-quinoline (MS (EI) M^+ = 427.1).

Example 7

7-chloro-3-(4-chloro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline

Table I compound 4

A mixture of 4-bromo-7-chloro-3-(4-chloro-benzenesulfonyl)-quinoline (0.42 g, 1 mmol), 4-fluorophenylboronic acid (0.21 g, 1.5 mmol) and tetrakis-(triphenylphosphin)palladium(0) (0.08 g, 0.07 mmol) in 30 ml 1,2-dimethoxyethane and 8 ml of 2M aqueous sodium carbonate solution was stirred for 2 hours at 70 °C. After cooling the mixture was concentrated *in vacuo*. The crude residue was purified by column chromatography on silica gel (Kieselgel 60, eluent: chloroform) and crystallized from methanol to obtain 0.29 g of the title compound in 67% yield..

^1H NMR (400 MHz, DMSO- d_6): 6.92-6.98 m (2H) [H-14,18]; 7.06-7.13 m (2H) [H-15,17]; 7.25-7.32 m (5H) [H-6, 22,23,25, 26]; 7.44 dd (1H) [H-7]; 9.23 d (1H) [H-9]; 9.77 s (1H) [H-2]

MS (EI) M^+ = 433.2

Applying the above procedure the following compounds were prepared: e.g. 7-fluoro-3-(4-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline (Table I compound 9, MS (EI) M^+ = 400.2), 4-(3-chloro-phenyl)-3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-quinoline (Table I compound 17, MS (EI) M^+ = 426.2),

4-(4-chloro-phenyl)-3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-quinoline (Table I compound 35, MS (EI) $M^+ = 434.2$),

4-(3-chloro-phenyl)-8-fluoro-3-(4-methoxy-benzenesulfonyl)-quinoline (Table I compound 53, MS (EI) $M^+ = 428.2$),

3-(3-cyano-benzenesulfonyl)-6-fluoro-4-(3-methoxy-phenyl)-quinoline (Table I compound 68, MS (EI) $M^+ = 419.3$),




3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline (Table I compound 101, MS (EI) $M^+ = 418.2$),








7-chloro-3-(3,4-difluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline (Table I compound 170, MS (EI) $M^+ = 434.2$),










7-chloro-3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline (Table I compound 186, MS (EI) $M^+ = 441.2$).








Examples of compounds of formula (I) and their affinity for rat mGluR5 and mGluR5 receptors are given in the table below.









Table








Comp. No.	Structure	(M+H) ⁺ or M ⁺	mGlu5 K _i (nM)	mGlu1 K _i (nM)
1.		393.8	***	**
2.		433.2	*	*
3.		433.2	*	*









4.		433.2	***	**
5.		450.1	**	*
6.		450.1	**	*
7.		484.1	***	***
8.		433.2	*	*
9.		400.2	***	**
10.		412.2	**	*








11.		428.2	***	*
12.		412.2	**	*
13.		424.2	***	*
14.		412.2	**	*
15.		400.2	***	**
16.		400.2	***	*
17.		426.2	***	*
18.		410.2	***	*
19.		422.3	***	*







20.	 <chem>Fc1ccc2nc(ccc2c1)S(=O)(=O)c3ccc(Cl)cc3</chem>	416.2	***	***
21.	 <chem>Fc1ccc2nc(ccc2c1)S(=O)(=O)c3ccc(Cl)cc3</chem>	416.2	***	**
22.	 <chem>Fc1ccc2nc(ccc2c1)S(=O)(=O)c3ccc(F)cc3</chem>	400.2	***	**
23.	 <chem>Fc1ccc2nc(ccc2c1)S(=O)(=O)c3ccc(OC)cc3</chem>	412.2	***	*
24.	 <chem>Fc1ccc2nc(ccc2c1)S(=O)(=O)c3ccc(Cl)cc3</chem>	416.2	*	*
25.	 <chem>Fc1ccc2nc(ccc2c1)S(=O)(=O)c3ccc(Cl)cc3</chem>	416.2	**	*
26.	 <chem>Fc1ccc2nc(ccc2c1)S(=O)(=O)c3ccc(F)cc3</chem>	400.2	**	*









27.		412.2	*	*
28.		400.2	*	*
29.		428.2	***	*
30.		412.2	**	*
31.		416.2	*	*
32.		416.2	*	*
33.		426.2	***	*
34.		428.2	*	*










35.		434.2	***	*
36.		418.2	***	*
37.		423.2	***	**
38.		407.2	***	**
39.		407.2	***	**
40.		400.2	***	*
41.		416.2	***	*









42.		412.2	**	*
43.		428.2	***	*
44.		412.2	***	*
45.		426.2	***	*
46.		410.2	***	*
47.		463.2	***	**
48.		446.2	***	*
49.		446.2	***	*









50.		451.2	***	***
51.		418.2	***	*
52.		412.2	***	*
53.		428.2	***	*
54.		428.2	***	***
55.		412.2	***	*
56.		424.2	***	*









57.		424.2	***	*
58.		416.2	***	**
59.		416.2	***	**
60.		400.2	***	**
61.		412.2	***	*
62.		412.2	***	*








63.		400.2	***	**
64.		434.2	*	*
65.		418.2	**	**
66.		418.2	**	*
67.		407.2	***	**
68.		419.3	***	*
69.		407.2	***	**
70.		458.3	*	*







71.		454.3	*	*
72.		428.2	***	*
73.		412.2	***	*
74.		458.3	***	*
75.		458.3	***	*
76.		416.2	***	*
77.		422.3	***	*
78.		463.2	***	*
79.		446.2	***	*







80.		451.2	***	**
81.		463.2	***	*
82.		463.2	***	*
83.		451.2	***	**
84.		430.2	***	*
85.		414.2	***	*
86.		414.2	***	*
87.		434.2	**	*









88.		418.2	***	**
89.		430.3	***	*
90.		430.3	***	*
91.		423.2	***	***
92.		419.3	***	**
93.		419.3	***	*
94.		430.2	***	*
95.		412.2	**	*









96.		458.3	**	*
97.		434.2	**	**
98.		418.2	***	**
99.		430.3	***	*
100.		430.3	***	*
101.		418.2	***	**
102.		451.2	***	**
103.		434.2	***	***









104.		446.2	***	**
105.		434.2	***	***
106.		451.2	***	*
107.		463.2	***	*
108.		463.2	***	*
109.		430.2	***	*
110.		430.2	***	*









111.		414.2	***	*
112.		426.3	***	*
113.		426.3	***	*
114.		414.2	***	*
115.		434.2	***	**
116.		434.2	***	**








117.		430.3	***	*
118.		430.3	***	*
119.		418.2	***	***
120.		434.2	***	**
121.		418.2	***	***
122.		430.3	***	*










123.		418.2	***	***
124.		451.2	***	**
125.		463.2	***	*
126.		463.2	***	*
127.		430.2	***	*
128.		414.2	***	*
129.		426.3	***	*
130.		426.3	***	*









131.		414.2	***	*
132.		434.2	**	*
133.		430.3	**	*
134.		430.3	**	*
135.		423.2	***	**
136.		423.2	***	**
137.		419.3	***	*
138.		447.2	***	*








139.		447.2	***	*
140.		430.2	***	*
141.		442.2	***	*
142.		430.2	***	*
143.		412.2	**	*
144.		428.2	***	*
145.		424.2	***	*
146.		410.2	***	*









147.		422.3	***	*
148.		410.2	***	*
149.		434.2	**	**
150.		418.2	***	**
151.		430.3	**	*
152.		418.2	***	**
153.		451.2	***	**
154.		446.2	***	*








155.		434.2	***	**
156.		468.1	***	**
157.		468.1	***	**
158.		468.1	***	**
159.		468.1	***	**
160.		468.1	***	*
161.		468.1	***	*







162.		458.3	*	*
163.		424.2	***	*
164.		458.3	*	*
165.		446.2	*	*
166.		468.1	***	*
167.		416.2	***	*
168.		416.2	***	*
169.		434.2	***	*
170.		434.2	***	*








171.		451.2	***	*
172.		433.2	***	*
173.		433.2	***	*
174.		451.2	***	*
175.		451.2	***	*
176.		468.1	***	*
177.		440.2	***	*
178.		468.1	***	*








179.		416.2	***	*
180.		416.2	***	*
181.		434.2	***	*
182.		434.2	***	*
183.		451.2	***	*
184.		484.1	*	*
185.		484.1	*	*








186.		441.2	***	*
187.		439.2	**	*
188.		469.2	**	*
189.		485.2	***	*
190.		469.2	***	*
191.		430.2	**	*
192.		425.2	*	*
193.		414.3	**	*








194.		469.2	*	*
195.		441.2	***	*
196.		452.2	***	*
197.		485.2	***	*
198.		469.2	*	*
199.		425.3	**	*
200.		458.2	*	*







201.		469.2	*	*
202.		452.2	**	*
203.		469.2	***	*
204.		469.2	***	*
205.		469.2	***	*
206.		407.3	***	*








207.		407.3	***	*
208.		451.1	***	**
209.		434.2	***	**
210.		434.2	***	**
211.		407.3	***	*
212.		434.2	***	**
213.		407.2	***	*








214.		389.1	***	*
215.		407.2	***	*
216.		425.1	***	*
217.		423.2	***	*
218.		425.1	***	*
219.		441.0	***	***
220.		469.4	***	*








221.		441.0	***	**
222.		407.2	***	*
223.		451.2	***	*
224.		434.2	***	**
225.		423.2	***	*
226.		423.2	***	*
227.		423.2	***	***







228.		425.1	***	*
229.		451.2	***	**
230.		407.2	***	*
231.		443.4	***	*
232.		457.3	***	*
233.		451.2	***	***
234.		423.3	***	***







235.		441.2	***	*
236.		436.2	***	**
237.		400.3	***	*
238.		436.2	***	*
239.		468.8	***	*
240.		400.1	***	*







241.		423.3	***	**
242.		458.0	***	***
243.		418.2	***	*
244.		441.2	***	*
245.		425.2	***	**
246.		423.2	***	*
247.		434.2	***	*

248.		486.2	***	**
249.		432.3	***	*
250.		458.0	***	**
251.		436.2	***	*
252.		503.8	***	**
253.		458.2	***	**
254.		440.1	***	**

255.		468.1	***	**
256.		450.3	***	*
257.		450.8	***	*
258.		451.1	***	**
259.		434.2	***	*
260.		451.3	***	*
261.		417.9	***	*

262.		469.4	***	***
263.		484.2	***	**
264.		415.1	***	**
265.		405.0	***	*
266.		425.1	***	*
267.		425.2	***	*

268.		441.2	***	*
269.		458.2	***	*
270.		441.2	***	*
271.		407.3	***	*
272.		423.2	***	*
273.		423.2	***	*

274.		441.2	***	*
275.		425.2	***	*
276.		441.2	***	**
277.		458.2	***	**
278.		422.1	***	*
279.		422.1	***	*

*** $K_i < 200 \text{ nM}$

** $200\text{nM} < K_i < 500 \text{ nM}$

* $500 \text{ nM} < K_i$

Example 8**Preparation of pharmaceutical compositions:****a) Tablets:**

0.01-50 % of active ingredient of formula (I), 15-50 % of lactose, 15-50 % of potato starch, 5-15 % of polyvinyl pyrrolidone, 1-5 % of talc, 0.01-3 % of magnesium stearate, 1-3 % of colloid silicon dioxide and 2-7 % of ultraamylopectin were mixed, then granulated by wet granulation and pressed to tablets.

b) Dragées, film coated tablets:

The tablets made according to the method described above were coated by a layer consisting of entero- or gastrosolvent film, or of sugar and talc. The dragées were polished by a mixture of beeswax and carnuba wax.

c) Capsules:

0.01-50 % of active ingredient of formula (I), 1-5 % of sodium lauryl sulfate, 15-50 % of starch, 15-50 % of lactose, 1-3 % of colloid silicon dioxide and 0.01-3 % of magnesium stearate were thoroughly mixed, the mixture was passed through a sieve and filled in hard gelatin capsules.

d) Suspensions:

Ingredients: 0.01-15 % of active ingredient of formula (I), 0.1-2 % of sodium hydroxide, 0.1-3 % of citric acid, 0.05-0.2 % of nipagin (sodium methyl 4-hydroxybenzoate), 0.005-0.02 % of nipasol, 0.01-0.5 % of carbopol (polyacrylic acid), 0.1-5 % of 96 % ethanol, 0.1-1 % of flavoring agent, 20-70 % of sorbitol (70 % aqueous solution) and 30-50 % of distilled water.

To solution of nipagin and citric acid in 20 ml of distilled water, carbopol was added in small portions under vigorous stirring, and the solution was left to stand for 10-12 h. Then the sodium hydroxide in 1 ml of distilled water, the aqueous solution of sorbitol and finally the ethanolic raspberry flavor were added with stirring. To this carrier the active ingredient was added in small portions and suspended with an immersing homogenizator. Finally the suspension was filled up to the desired final volume with distilled water and the suspension syrup was passed through a colloid milling equipment.

e) Suppositories:

For each suppository 0.01-15% of active ingredient of formula (I) and 1-20% of lactose were thoroughly mixed, then 50-95% of adeps pro suppository (for example Witepsol

4) was melted, cooled to 35 °C and the mixture of active ingredient and lactose was mixed in it with homogenizator. The obtained mixture was mould in cooled forms.

f) Lyophilized powder ampoule compositions:

A 5 % solution of mannitol or lactose was made with bidistilled water for injection use, and the solution was filtered so as to have sterile solution. A 0.01-5 % solution of the active ingredient of formula (I) was also made with bidistilled water for injection use, and this solution was filtered so as to have sterile solution. These two solutions were mixed under aseptic conditions, filled in 1 ml portions into ampoules, the content of the ampoules was lyophilized, and the ampoules were sealed under nitrogen. The contents of the ampoules were dissolved in sterile water or 0.9 % (physiological) sterile aqueous sodium chloride solution before administration.

Example 9

3-(3,4-Difluoro-benzenesulfonyl)-7-nitro-quinolin-4-ol (intermediate)

The title compound was prepared applying the procedure described in Example 4 from 2-(3,4-difluoro-benzenesulfonyl)-3-ethoxy-acrylic acid methyl ester (3.49 g, 11.4 mmol) and 3-nitroaniline (1.57 g, 11.4 mmol). The yield was 1.6 g (38.3%).

MS (EI) $M^+ = 367.2$.

In the same way was prepared: e.g. 3-(3-cyano-5-fluoro-benzenesulfonyl)-8-nitro-quinolin-4-ol (MS (EI) $M^+ = 374.3$).

Example 10

4-Bromo-3-(3,4-difluoro-benzenesulfonyl)-7-nitro-quinoline (intermediate)

A mixture of 3-(3,4-difluoro-benzenesulfonyl)-7-nitro-quinolin-4-ol (1.6 g, 4.37 mmol) and phosphorus(V) oxybromide (2.5 g, 8.72 mmol) in DMF (16 ml) was stirred at 65°C for 1 hour. The reaction mixture was diluted with water (100 ml) and the pH was adjusted to 10 with aqueous sodium hydroxide solution. After cooling the precipitate was filtered and washed with water. The obtained crude product was purified by crystallization from ethanol to obtain 1.25 g of the title compound in 67% yield.

MS (EI) $M^+ = 430.9$.

Applying the above procedure the following compound was prepared: e.g. 4-bromo-3-(3-cyano-5-fluoro-benzenesulfonyl)-8-nitro-quinoline (MS (EI) $M^+ = 437.2$).

Example 11

3-(3,4-Difluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-7-nitro-quinoline (intermediate)

The title compound was prepared applying the procedure described in Example 7 from 4-Bromo-3-(3,4-difluoro-benzenesulfonyl)-7-nitro-quinoline (0.53 g, 1.23 mmol) and 3-fluorophenylboronic acid (0.21 g, 1.5 mmol). The crude product was purified by column chromatography on silica gel (Kieselgel 60, eluent: chloroform) and crystallized from methanol to obtain 0.31 g of the title compound in 57% yield.

MS (EI) $M^+ = 445.3$.

In the same way were prepared: e.g. 3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-8-nitro-quinoline (MS (EI) $M^+ = 452.3$); 3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-8-nitro-quinoline (MS (EI) $M^+ = 452.3$).

Example 12

7-Amino-3-(3,4-difluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline

Table I compound 264


A mixture of 3-(3,4-difluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-7-nitro-quinoline (0.31 g, 0.69 mmol) and iron powder (0.15 g, 2.6 mmol) in acetic acid was stirred for 30 minutes at 60°C. The reaction mixture was diluted with water (5 ml). The precipitate was filtered and washed with water (2x5 ml). The obtained crude product was purified by crystallization from methanol to obtain 0.12 g of the title compound in 42% yield.

MS (EI) $M^+ = 415.1$.

Applying the above procedure the following compounds were prepared: e.g. 7-amino-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline (MS (EI) $M^+ = 422.1$); 7-amino-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline (MS (EI) $M^+ = 422.1$).

CLAIMS

1. A compound having the formula (I):



(I)


wherein

Ar₁ represents an optionally substituted phenyl or heteroaryl group;

Ar₂ represents a substituted phenyl or an optionally substituted heteroaryl group;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, halogen, cyano, alkyl, alkoxy, hydroxy, trifluoromethyl, amino, alkylamino, dialkylamino, aminomethyl, alkylaminomethyl, dialkylaminomethyl, and/or salts and/or hydrates and/or solvates thereof.

2. A compound having the formula (I):



(I)

wherein


Ar₁ represents phenyl or heteroaryl group, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

Ar₂ represents phenyl, substituted with one or more substituent(s) selected from fluoro, chloro, cyano, methyl, methoxy; or

heteroaryl, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy, hydroxy, trifluoromethyl, amino, methylamino, dimethylamino, aminomethyl, methylaminomethyl, dimethylaminomethyl, and/or salts and/or hydrates and/or solvates thereof.

3. A compound having the formula (I):



(I)

wherein

Ar₁ represents phenyl, pyridyl, thienyl or oxazolyl group, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

Ar₂ represents phenyl, substituted with one or more substituent(s) selected from fluoro, chloro, cyano, methyl, methoxy; or

pyridyl, thienyl or oxazolyl, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy, hydroxy, trifluoromethyl, amino, methylamino, dimethylamino, aminomethyl, methylaminomethyl, dimethylaminomethyl, and/or salts and/or hydrates and/or solvates thereof.

4. A compound according to claim 1 selected from the group consisting of
 4-(4-chloro-phenyl)-3-(4-methyl-benzenesulfonyl)-quinoline,
 7-chloro-3-(4-chloro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
 8-chloro-4-(3-chloro-phenyl)-3-(3,4-dichloro-benzenesulfonyl)-quinoline,
 7-fluoro-3-(4-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
 4-(4-chloro-phenyl)-7-fluoro-3-(4-methoxy-benzenesulfonyl)-quinoline,
 7-fluoro-3-(4-methoxy-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
 7-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
 7-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,

4-(3-chloro-phenyl)-3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-4-(3-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-8-fluoro-3-(4-fluoro-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-8-fluoro-3-(4-fluoro-benzenesulfonyl)-quinoline,
8-fluoro-3-(4-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
8-fluoro-3-(4-fluoro-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
4-(4-chloro-phenyl)-6-fluoro-3-(4-methoxy-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-3-(3,4-dimethyl-benzenesulfonyl)-6-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(4-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-7-fluoro-quinoline,
3-(3-cyano-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
7-fluoro-3-(4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
4-(4-chloro-phenyl)-7-fluoro-3-(3-fluoro-benzenesulfonyl)-quinoline,
4-(3-chloro-phenyl)-7-fluoro-3-(3-methoxy-benzenesulfonyl)-quinoline,
7-fluoro-4-(4-fluoro-phenyl)-3-(3-methoxy-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-chloro-4-methoxy-benzenesulfonyl)-7-fluoro-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
8-fluoro-3-(4-fluoro-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-8-fluoro-3-(4-methoxy-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-8-fluoro-3-(4-methoxy-benzenesulfonyl)-quinoline,
8-fluoro-4-(4-fluoro-phenyl)-3-(4-methoxy-benzenesulfonyl)-quinoline,
8-fluoro-3-(4-methoxy-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
8-fluoro-3-(4-methoxy-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-8-fluoro-3-(3-fluoro-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-8-fluoro-3-(3-fluoro-benzenesulfonyl)-quinoline,

8-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-6-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-6-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-6-fluoro-3-(3-methoxy-benzenesulfonyl)-quinoline,
6-fluoro-4-(4-fluoro-phenyl)-3-(3-methoxy-benzenesulfonyl)-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-6-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-6-fluoro-4-(3-methoxy-phenyl)-quinoline,
4-(4-chloro-phenyl)-7-fluoro-3-(4-fluoro-benzenesulfonyl)-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-4-(4-chloro-phenyl)-6-fluoro-quinoline,
3-(3-chloro-4-methoxy-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-7-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-7-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-quinoline,
7-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
7-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-7-fluoro-4-(3-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-7-fluoro-quinoline,
3-(3-cyano-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-7-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3-chloro-4-methyl-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-7-fluoro-quinoline,

3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-4-(3-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-quinoline,
4-(4-chloro-phenyl)-8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-quinoline,
8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
8-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3,5-difluoro-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(4-chloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-6-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-6-fluoro-4-(3-methoxy-phenyl)-quinoline,
4-(3-chloro-phenyl)-6-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-quinoline,
6-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
6-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
6-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
6-fluoro-3-(3-fluoro-4-methyl-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-6-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-6-fluoro-quinoline,
3-(3-cyano-benzenesulfonyl)-6-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3-chloro-4-methyl-benzenesulfonyl)-4-(3-chloro-phenyl)-6-fluoro-quinoline,

3-(3-chloro-4-methyl-benzenesulfonyl)-4-(4-chloro-phenyl)-6-fluoro-quinoline,
3-(3-chloro-4-methyl-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-chloro-4-methyl-benzenesulfonyl)-6-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3-chloro-4-methyl-benzenesulfonyl)-6-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(4-chloro-phenyl)-6-fluoro-3-(3-methoxy-benzenesulfonyl)-quinoline,
6-fluoro-3-(3-methoxy-benzenesulfonyl)-4-(3-methoxy-phenyl)-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-6-fluoro-4-(4-methoxy-phenyl)-quinoline,
3-(3,4-dimethyl-benzenesulfonyl)-6-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-6-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-6-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-6-fluoro-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-6-fluoro-4-(3-methoxy-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-6-fluoro-4-(4-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-7-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-7-fluoro-quinoline,
4-(3-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-quinoline,
4-(3-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-6-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-6-fluoro-quinoline,
6-fluoro-3-(4-methoxy-benzenesulfonyl)-4-(4-methoxy-phenyl)-quinoline,
7-chloro-4-(4-chloro-phenyl)-3-(3,5-difluoro-benzenesulfonyl)-quinoline,
7-chloro-4-(4-chloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-quinoline,
7-chloro-4-(4-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-quinoline,
7-chloro-3-(3,5-dichloro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3,4-difluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline and
7-chloro-3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline
6-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-7-fluoro-quinoline,
6-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-7-cyano-4-(2-fluoro-phenyl)-quinoline,

7-chloro-3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-8-fluoro-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
7-chloro-4-(4-chloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline.


3-(3-cyano-4-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-8-fluoro-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-8-fluoro-4-(2-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-8-fluoro-4-(2-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-8-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-4-fluoro-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-4-fluoro-benzenesulfonyl)-quinoline,
3-(3-cyano-5-fluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3-cyano-4-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-8-fluoro-quinoline,
7-chloro-3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
3-(3-cyano-benzenesulfonyl)-8-fluoro-4-(4-fluoro-phenyl)-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-4-(2-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(2-fluoro-phenyl)-quinoline,
4-(4-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-8-fluoro-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-5-fluoro-benzenesulfonyl)-quinoline,
7-chloro-3-(3-cyano-benzenesulfonyl)-4-(2-fluoro-phenyl)-quinoline,
3-(3-cyano-5-fluoro-benzenesulfonyl)-7-fluoro-4-(3-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(3-chloro-phenyl)-8-fluoro-quinoline,
3-(3-cyano-benzenesulfonyl)-7-fluoro-4-(2-fluoro-phenyl)-quinoline,
3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-7-fluoro-quinoline,
3-(3-cyano-4-fluoro-benzenesulfonyl)-4-(3,4-dichloro-phenyl)-quinoline,

7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(2-fluoro-phenyl)-quinoline,
7-chloro-3-(3-cyano-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-4-fluoro-benzenesulfonyl)-7-fluoro-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-4-(3,5-difluoro-phenyl)-8-fluoro-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-8-fluoro-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-7-fluoro-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
7-chloro-3-(3-cyano-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
7-chloro-4-(3-chloro-phenyl)-3-(3-cyano-4-fluoro-benzenesulfonyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-4-(2-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-5-fluoro-benzenesulfonyl)-7-fluoro-quinoline,
3-(3-cyano-benzenesulfonyl)-4-(3,5-difluoro-phenyl)-7-fluoro-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-8-fluoro-quinoline,
4-(4-chloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-quinoline,
7-chloro-3-(3,4-dichloro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-quinoline,
3-(3,5-dicyano-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
7-chloro-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(3-chloro-phenyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-7-fluoro-quinoline,
7-chloro-4-(3-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-8-fluoro-quinoline,
7-chloro-4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-8-fluoro-quinoline,
7-chloro-4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-quinoline,
7-chloro-3-(3,5-dichloro-benzenesulfonyl)-4-(2-fluoro-phenyl)-quinoline,
3-(3,5-dicyano-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-7-fluoro-quinoline,
3-(3,5-dichloro-benzenesulfonyl)-7-fluoro-4-(2-fluoro-phenyl)-quinoline,
3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(3-chloro-phenyl)-7-fluoro-quinoline,
4-(3-chloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-8-fluoro-quinoline,
4-(3,4-dichloro-phenyl)-3-(3,4-difluoro-benzenesulfonyl)-quinoline,
3-(3,4-difluoro-benzenesulfonyl)-7-fluoro-4-(2-fluoro-phenyl)-quinoline,
7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(3,5-difluoro-phenyl)-quinoline,
7-chloro-4-(3-chloro-phenyl)-3-(3,5-dichloro-benzenesulfonyl)-quinoline,
7-amino-3-(3,4-difluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline,
4-(3-chloro-phenyl)-3-(3-cyano-benzenesulfonyl)-quinoline,

3-(3-cyano-5-fluoro-benzenesulfonyl)-4-(3,4-difluoro-phenyl)-quinoline,
 3-(3-cyano-5-fluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
 4-(4-chloro-phenyl)-3-(3-cyano-5-fluoro-benzenesulfonyl)-7-fluoro-quinoline,
 7-chloro-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-quinoline,
 7-chloro-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
 3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
 3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-quinoline,
 3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-quinoline,
 3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-7-fluoro-quinoline,
 3-(3-chloro-4-fluoro-benzenesulfonyl)-7-fluoro-4-(4-fluoro-phenyl)-quinoline,
 7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
 7-chloro-3-(3-chloro-4-fluoro-benzenesulfonyl)-4-(4-chloro-phenyl)-quinoline,
 7-amino-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(4-fluoro-phenyl)-quinoline,
 7-amino-3-(3-chloro-5-fluoro-benzenesulfonyl)-4-(3-fluoro-phenyl)-quinoline.

and/or salts and/or hydrates and/or solvates thereof.


5. mGluR1 and mGluR5 receptor subtype preferring ligands of formula (I):



(I)

wherein Ar₁, Ar₂, R₁, R₂, R₃ and R₄ are as defined in any one of claims 1 to 3,
 and/or salts and/or hydrates and/or solvates thereof.

6. A process for the preparation of a compound of formula (I):




(I)

wherein Ar₁, Ar₂, R₁, R₂, R₃ and R₄ are as defined in any one of claims 1 to 3, and/or salts and/or hydrates and/or solvates thereof,


a,

a1. by reacting a compound of formula (II):




(II)

wherein R₁, R₂, R₃ and R₄ are as defined above for a compound of formula (I), with a compound of formula (III):




(III)

wherein Ar₁ is as defined above for compounds of formula (I), M is selected from alkali metals or alkaline-earth metals, to give a compound of formula (IV):



(IV)


wherein Ar₁, R₁, R₂, R₃ and R₄ and are as defined above for compounds of formula (I), thereafter oxidizing a compound of formula (IV) to obtain a compound of formula (V):



(V)

wherein Ar₁, R₁, R₂, R₃ and R₄ are as defined above for compounds of formula (I), thereafter oxidizing a compound of formula (V) to obtain a compound of formula (VI):

85



(VI)


wherein Ar₁, R₁, R₂, R₃ and R₄ are as defined above for compounds of formula (I), or a2.

by reacting a compound of formula (IX):



(IX)

wherein Ar₁ is as defined above for a compound of formula (I), with α -halogen-acetic acid esters of formula (X):




(X)

wherein Hlg is halogen and R₅ is an ethyl or methyl group, to obtain a compound of formula (XI):




(XI)

wherein Ar₁ is as defined above for a compound of formula (I) and R₅ is as defined above for compounds of formula (X), thereafter reacting a compound of formula (XI) with a trialkyl orthoformate of formula (XII):



(XII)


wherein R₆ is an ethyl or methyl group, to obtain a compound of formula (XIII):



(XIII)


wherein Ar₁ is as defined above for a compound of formula (I), R₅ is as defined above for compounds of formula (X) and R₆ is as defined above for compounds of formula (XII), thereafter reacting a compound of formula (XIII) with an aniline derivative of formula (XIV):

86




(XIV)

wherein R_1 , R_2 , R_3 and R_4 are as defined above for a compound of formula (I), to obtain a compound of formula (VI):



(VI)


wherein Ar_1 , R_1 , R_2 , R_3 and R_4 are as defined above for a compound of formula (I),
thereafter converting a compound of formula (VI) to a compound of formula (VII):



(VII)

wherein Ar_1 , R_1 , R_2 , R_3 and R_4 are as defined above for compounds of formula (I), X is selected from chloro, bromo, benzenesulfonyloxy, 4-fluoro-benzenesulfonyloxy, 4-methyl-benzenesulfonyloxy, methanesulfonyloxy or trifluoromethanesulfonyloxy groups,

thereafter reacting the obtained compound of formula (VII) with a boronic acid derivative of formula (VIII):




(VIII)

wherein Ar_2 is as defined above for a compound of formula (I), in the presence of base and catalyst in a solvent,

and optionally thereafter forming salts and/or hydrates and/or solvates of compounds of formula (I), or

b, interconverting one compound of formula (I), wherein Ar₁, Ar₂, R₁, R₂, R₃ and R₄ are as defined above for the formula (I), to a different compound of formula (I), wherein Ar₁, Ar₂, R₁, R₂, R₃ and R₄ are as defined as described above for the formula (I), and optionally thereafter forming salts and/or hydrates and/or solvates of compounds of formula (I).


7. An intermediate of the formula (IV):



wherein

Ar₁, R₁, R₂ and R₃ are as defined above for compounds of formula (I), R₄ is selected from hydrogen, halogen, cyano, alkyl, alkoxy, hydroxy, amino, alkylamino, dialkylamino, aminomethyl, alkylaminomethyl, dialkylaminomethyl, and/or salts and/or hydrates and/or solvates thereof.


8. An intermediate of the formula (V):



wherein

Ar₁, R₁, R₂, R₃ and R₄ are as defined above for compounds of formula (I), and/or salts and/or hydrates and/or solvates thereof.

9. An intermediate of the formula (VI):




(VI)

wherein

Ar₁, R₁, R₂, R₃ and R₄ are as defined above for a compound of formula (I), and/or salts and/or hydrates and/or solvates thereof.

10. An intermediate of the formula (VII):




(VII)

wherein

Ar₁, R₁, R₂, R₃ and R₄ are as defined above for compounds of formula (I), X is selected from chloro, bromo, benzenesulfonyloxy, 4-fluoro-benzenesulfonyloxy, 4-methyl-benzenesulfonyloxy, methanesulfonyloxy or trifluoromethanesulfonyloxy groups, and/or salts and/or hydrates and/or solvates thereof.

11. A pharmaceutical formulation comprising a therapeutically effective amount of a compound of formula (I):



(I)


wherein

Ar₁ represents an optionally substituted phenyl or heteroaryl group;

Ar₂ represents a substituted phenyl or an optionally substituted heteroaryl group;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, halogen, cyano, alkyl, alkoxy, hydroxy, trifluoromethyl, amino, alkylamino, dialkylamino, aminomethyl, alkylaminomethyl, dialkylaminomethyl, and/or physiologically acceptable salts and/or hydrates and/or solvates thereof in association with one or more physiologically acceptable diluents, excipients and/or inert carriers.

12. A pharmaceutical formulation comprising a therapeutically effective amount of a compound of formula (I):



(I)

wherein


Ar₁ represents phenyl or heteroaryl group, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

Ar₂ represents phenyl, substituted with one or more substituent(s) selected from fluoro, chloro, cyano, methyl, methoxy; or

heteroaryl, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy, hydroxy, trifluoromethyl, amino, methylamino, dimethylamino, aminomethyl, methylaminomethyl, dimethylaminomethyl, and/or physiologically acceptable salts and/or hydrates and/or solvates thereof in association with one or more physiologically acceptable diluents, excipients and/or inert carriers.

13. A pharmaceutical formulation comprising a therapeutically effective amount of a compound of formula (I):



(I)

wherein

Ar₁ represents phenyl, pyridyl, thienyl or oxazolyl group, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;


Ar₂ represents phenyl, substituted with one or more substituent(s) selected from fluoro, chloro, cyano, methyl, methoxy; or

pyridyl, thienyl or oxazolyl, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy, hydroxy, trifluoromethyl, amino, methylamino, dimethylamino, aminomethyl, methylaminomethyl, dimethylaminomethyl, and/or physiologically acceptable salts and/or hydrates and/or solvates thereof in association with one or more physiologically acceptable diluents, excipients and/or inert carriers.

14. A pharmaceutical composition according to any claims of 10 to 12 for use in the prevention and/or treatment of mGluR1 and mGluR5 receptor mediated disorders.

15. The use of a compound of formula (I):



(I)


wherein

Ar₁ represents an optionally substituted phenyl or heteroaryl group;

Ar₂ represents a substituted phenyl or an optionally substituted heteroaryl group;

R_1 , R_2 , R_3 and R_4 represent independently a substituent selected from hydrogen, halogen, cyano, alkyl, alkoxy, hydroxy, trifluoromethyl, amino, alkylamino, dialkylamino, aminomethyl, alkylaminomethyl, dialkylaminomethyl, and/or physiologically acceptable salts and/or hydrates and/or solvates thereof in the manufacture of a medicament for the treatment and/or prevention of mGluR1 and mGluR5 receptor mediated disorders.

16. The use of a compound of formula (I):



(I)

wherein


Ar_1 represents phenyl or heteroaryl group, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

Ar_2 represents phenyl, substituted with one or more substituent(s) selected from fluoro, chloro, cyano, methyl, methoxy; or

heteroaryl, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

R_1 , R_2 , R_3 and R_4 represent independently a substituent selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy, hydroxy, trifluoromethyl, amino, methylamino, dimethylamino, aminomethyl, methylaminomethyl, dimethylaminomethyl, and/or physiologically acceptable salts and/or hydrates and/or solvates thereof in the manufacture of a medicament for the treatment and/or prevention of mGluR1 and mGluR5 receptor mediated disorders.

17. The use of a compound of formula (I):



(I)

wherein

Ar₁ represents phenyl, pyridyl, thienyl or oxazolyl group, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

Ar₂ represents phenyl, substituted with one or more substituent(s) selected from fluoro, chloro, cyano, methyl, methoxy; or

pyridyl, thienyl or oxazolyl, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy, hydroxy, trifluoromethyl, amino, methylamino, dimethylamino, aminomethyl, methylaminomethyl, dimethylaminomethyl, and/or physiologically acceptable salts and/or hydrates and/or solvates thereof in the manufacture of a medicament for the treatment and/or prevention of mGluR1 and mGluR5 receptor mediated disorders.

18. The use of a compound according to any claims of 14 to 16 wherein said mGluR1 and mGluR5 receptor mediated disorders are psychiatric disorders.

19. The use of a compound according to any claims of 14 to 16 wherein said mGluR1 and mGluR5 receptor mediated disorders are neurological disorders.


20. The use of a compound according to any claims of 14 to 16 wherein said mGluR1 and mGluR5 receptor mediated disorders are chronic and acute pain.

21. The use of a compound according to any claims of 14 to 16 wherein said mGluR1 and mGluR5 receptor mediated disorders are neuromuscular dysfunctions of the lower urinary tract.

22. The use of a compound according to any claims of 14 to 16 wherein said mGluR1 and mGluR5 receptor mediated disorders are gastrointestinal reflux disease and irritable bowel syndrome.

23. The use of a compound according to any claims of 14 to 16 wherein said mGluR1 and mGluR5 receptor mediated disorders are substance abuse and withdrawal.

24. A method of prevention and/or treatment of mGluR1 and mGluR5 receptor mediated disorders, comprising administering to a mammal in need of such prevention and/or treatment, a therapeutically effective amount of a compound of formula (I):



(I)


wherein

Ar₁ represents an optionally substituted phenyl or heteroaryl group;

Ar₂ represents a substituted phenyl or an optionally substituted heteroaryl group;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, halogen, cyano, alkyl, alkoxy, hydroxy, trifluoromethyl, amino, alkylamino, dialkylamino, aminomethyl, alkylaminomethyl, dialkylaminomethyl, and/or physiologically acceptable salts and/or hydrates and/or solvates thereof.

25. A method of prevention and/or treatment of mGluR1 and mGluR5 receptor mediated disorders, comprising administering to a mammal in need of such prevention and/or treatment, a therapeutically effective amount of a compound of formula (I):



(I)

wherein


Ar₁ represents phenyl or heteroaryl group, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

Ar₂ represents phenyl, substituted with one or more substituent(s) selected from fluoro, chloro, cyano, methyl, methoxy; or

heteroaryl, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy, hydroxy, trifluoromethyl, amino, methylamino, dimethylamino, aminomethyl, methylaminomethyl, dimethylaminomethyl, and/or physiologically acceptable salts and/or hydrates and/or solvates thereof.

26. A method of prevention and/or treatment of mGluR1 and mGluR5 receptor mediated disorders, comprising administering to a mammal in need of such prevention and/or treatment, a therapeutically effective amount of a compound of formula (I):



(I)

wherein

Ar₁ represents phenyl, pyridyl, thienyl or oxazolyl group, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

Ar₂ represents phenyl, substituted with one or more substituent(s) selected from fluoro, chloro, cyano, methyl, methoxy; or

pyridyl, thienyl or oxazolyl, optionally substituted with one or more substituent(s) selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy;

R₁, R₂, R₃ and R₄ represent independently a substituent selected from hydrogen, fluoro, chloro, cyano, methyl, methoxy, hydroxy, trifluoromethyl, amino, methylamino, dimethylamino, aminomethyl, methylaminomethyl, dimethylaminomethyl, and/or physiologically acceptable salts and/or hydrates and/or solvates thereof.

27. A method according to any claims of 23 to 25, wherein said mammal is a human.
28. A method according to any claims of 23 to 25, wherein said mGluR1 and mGluR5 receptor mediated disorders are psychiatric disorders.
29. A method according to any claims of 23 to 25, wherein said mGluR1 and mGluR5 receptor mediated disorders are neurological disorders.
30. A method according to any claims of 23 to 25, wherein said mGluR1 and mGluR5 receptor mediated disorders are chronic and acute pain disorders.
31. A method according to any claims of 23 to 25, wherein said mGluR1 and mGluR5 receptor mediated disorders are neuromuscular dysfunctions of the lower urinary tract.
32. A method according to any claims of 23 to 25, wherein said mGluR1 and mGluR5 receptor mediated disorders are gastrointestinal reflux disease and irritable bowel syndrome.
33. A method according to any claims of 23 to 25, wherein said mGluR1 and mGluR5 receptor mediated disorders are substance abuse and withdrawal.