


TITLE OF INVENTION
RETINOID-LIKE COMPOUNDS



This is a continuation-in-part application of U.S.S.N. 08/306,092 filed September 19, 1994, which is a
5 continuation-in-part application of U.S.S. 08/216,740
filed March 23, 1994, which in turn is a continuation-
in-part of U.S.S.N. 08/176,746 filed January 3, 1994;
all previous applications are herein incorporated by
reference in their entirety.


10

The present invention provides compounds having
retinoid-like activity. More specifically, the
15 compounds of the present invention are useful for
preventing and/or treating various skin disorders,
such as, but not limited to, acne, psoriasis and
damage from irradiation. Further, they have antitumor
and antiarthritic activities.

20

Retinoic acid and its natural and synthetic analogues
(retinoids) exert a wide array of biological effects.

25




They have been shown to affect cellular growth and
differentiation and are promising drugs for the
30 treatment of several cancers. Roberts, A.B. and
Sporn, M.B. in "The Retinoids," Sporn, M.B., Roberts,


- A.B., and Goodman, D.S., eds, 1984, 2, pp. 209-286, Academic Press, New York; Lippman, S.M., Kessler, J.F., and Meyskens, F.L., *Cancer Treat. Rep.*, 1987, 71, p. 391; *ibid.*, p. 493; Hong, W.K. et al., *N. Engl. J. Med.*, 1990, 323, p. 795; Huang, M. et al., *Blood*, 1988, 72, p. 567. Retinods have also been shown to be useful in treating rheumatic diseases, see for example: J.W. Coffey et al., *Retinoids as Potential Antirheumatic Agents*, Chemistry and Biology of Synthetic Retinoids, pp 520-537, CRC Press Inc., M. I. Dawson and W.H. Okamura Ed.(1990)

A few retinoids are already in clininal use in the treatment of dermatological diseases such as acne and psoriasis. For example, isotretinoin is used clinically for oral therapy of severe acne, and etretinate is particularly useful in the treatment of psoriasis. Orfanos, C.E., Ehlert, R., and Gollnick, H., *Drugs*, 1987, 34, pp. 459-503.


20




ISOTRETINOIN



ETRETINATE



II



III


Other examples of retinoid compounds include aroatinoid of formula II and retinobenzoic acid of formula III, in which Q equals -NHCO-, -CONH-, -COCH=CH-,

-CH=CHCO-, -COCH₂-, etc.

See for example: Loeliger, P., Bollag, W., and Mayer,
 5 H., *Eur. J. Med. Chem.* 1980, 15, pp. 9-15; Kagechika,
 H. et al., *J. Med. Chem.*, 1988, 31, No. 11, pp. 2182-
 2192.

10

The present invention relates to a compound of formula
 I



15

or a nontoxic pharmaceutically acceptable salt,
 physiologically hydrolyzable ester or solvate thereof,
 in which

20 X is -O-CO-, -NH-CO-, -CS-NH-, -CO-O-, -CO-NH-,
 -COS-, -SCO-, -SCH₂-, -CH₂-CH₂-, -C≡C-,
 -CH₂-NH-, -COCH₂-, -NHCS-, -CH₂S-, -CH₂O-,
 -OCH₂-, -NHCH₂- or -CR⁵=CR⁶-;

R^m and R^k are independently hydrogen, halogen,
 25 C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy or nitro;
 n is zero or one;


R⁴ is -(CH₂)_t-Y, C₁₋₆alkyl, or C₃₋₆cycloalkyl;

R¹ is -CO₂Z, C₁₋₆alkyl, CH₂OH, -CONHR^y, or CHO;

R² and R³ are independently hydrogen or C₁₋₆alkyl;

30 R^a and R^b are independently hydrogen or C₁₋₆alkyl;

but when n is one, R^a and R^b together can form a radical of the formula



5

Y is naphthyl or phenyl, both radicals can be optionally substituted with one to three same or different C₁₋₆alkyl or halogen;

Z is hydrogen or C₁₋₆alkyl;

10 R⁵, R⁶ and R⁷ are independently hydrogen or C₁₋₆alkyl; and

t is zero to six.


Also provided by this invention are methods for
 15 preventing and/or treating tumors, arthritis, and non-malignant skin disorders comprising administering a compound of formula I to a mammal. Further provided is a pharmaceutical formulation (composition) comprising a compound of formula I in admixture with
 20 (a) pharmaceutically acceptable excipient(s).

Figure 1, 2 and 3 are the cytotoxicity dose response
 25 curves for lung line L2987.

Figure 4 is a dose response curve for collagen-induced arthritis.

The present invention relates to a compound of formula
I

5




or a nontoxic pharmaceutically acceptable salt,
10 physiologically hydrolyzable ester or solvate thereof,
in which

X is -O-CO-, -NH-CO-, -CS-NH-, -CO-O-, -CO-NH-,
-COS-, -SCO-, -SCH₂-, -CH₂-CH₂-, -C≡C-,
15 -CH₂-NH-, -COCH₂-, -NHCS-, -CH₂S-, -CH₂O-,
-OCH₂-, -NHCH₂- or -CR⁵=CR⁶-;

R^m and R^k are independently hydrogen, halogen,
C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy or nitro;
n is zero or one;

20 R⁴ is -(CH₂)_t-Y, C₁₋₆alkyl, or C₃₋₆cycloalkyl;
R¹ is -CO₂Z, C₁₋₆alkyl, CH₂OH, -CONHR^y, or CHO;
R² and R³ are independently hydrogen or C₁₋₆alkyl;
R^a and R^b are independently hydrogen or C₁₋₆alkyl;
but when n is one, R^a and R^b together can
25 form a radical of the formula



Y is naphthyl or phenyl, both radicals can be

optionally substituted with one to three
same or different C₁₋₆alkyl or halogen;

Z is hydrogen or C₁₋₆alkyl;

R⁵, R⁶ and R⁷ are independently hydrogen or

5 C₁₋₆alkyl; and

t is zero to six.

In the instant application, the numbers in subscript
after the symbol "C" define the number of carbon atoms
10 a particular group can contain. For example, C₁₋₆alkyl
refers to straight and branched chain alkyl groups
with one to six carbon atoms and such groups include
methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl,
n-pentyl, n-hexyl, 3-methylpentyl, or the like alkyl
15 groups; C₃₋₆cycloalkyl refers to cyclopropyl,
cyclobutyl, cyclopentyl, or cyclohexyl; and halogen
refers to fluorine, chlorine, bromine, or iodine. In
the instant application all symbols once defined
retain the same meaning until they are redefined.

20

Some compounds of formula I may also form
pharmaceutically acceptable metal and amine salts in
which the cation does not contribute significantly to
the toxicity or biological activity of the salt.

25 These salts are also part of the present invention.

Suitable metal salts include the sodium, potassium,
calcium, barium, zinc, and aluminum salts. The sodium
or potassium salts are preferred. Amines which are
capable of forming stable salts group include

30 trialkylamines such as triethylamine, procaine,
dibenzylamine, N-benzyl-β-phenethylamine, 1-
ephenamine, N,N'-dibenzylethylenediamine,
dehydroabietylamine, N-ethylpiperidine, benzylamine,
dicyclohexylamine, or the like pharmaceutically

acceptable amines.

When compounds of formula I contains carboxy groups, it can form physiologically hydrolyzable esters which
 5 serve as prodrugs by being hydrolyzed in the body to yield formula I compounds per se. They are preferably administered orally since hydrolysis in many instances occurs principally under the influence of the digestive enzymes. Parenteral administration may be
 10 used where the ester per se is active, or in those instances where hydrolysis occurs in the blood. Examples of physiologically hydrolyzable esters of compounds of formula I include C₁₋₆alkyl, benzyl, 4-methoxybenzyl, indanyl, phthalidyl, methoxymethyl,
 15 C₁₋₆alkanoyloxyC₁₋₆alkyl, e.g. acetoxymethyl, pivaloyloxymethyl or propionyloxymethyl, C₁₋₆alkoxycarbonyloxyC₁₋₆alkyl, e.g. methoxycarbonyloxymethyl or ethoxycarbonyloxymethyl, glycyloxymethyl, phenylglycyloxymethyl, (5-methyl-2-
 20 oxo-1-3-dioxolen-4-yl)-methyl and other well-known physiologically hydrolyzable esters used, for example, in the penicillin and cephalosporin arts. Such esters are prepared by conventional techniques known in the art.

25 The structural formulae as drawn in the instant application are believed to best represent the structures of compounds of the present invention. However, some compounds within the scope of the
 30 invention may exist as other tautomeric forms, in which hydrogen atoms are transposed to other parts of the molecules and the chemical bonds between the atoms of the molecules are consequently rearranged. It should be understood that the structural formulae

represent all tautomeric forms, insofar as they may exist.

The synthesis of a compound of formula I can be
5 accomplished by a wide variety of methods using
conventional starting materials and processes. The
synthetic descriptions and specific examples that
follow are only intended for the purpose of
illustration, and are not to be construed as limiting
10 in any manner ways to make compounds of the present
invention by other methods.

Typically a compound of formula I can be made by
employing one of the processes or obvious variations
15 thereof as depicted in Schemes I to XXII. All the
steps in Schemes I to XXII are standard processes
which can be easily practiced by anyone skilled in the
art. The specific examples that are provided after
the Schemes are intended to illustrate specific
20 conditions which may be employed to carry out certain
steps in the Schemes and are not to be construed as
limiting the conditions in any way.

In the Schemes, R⁷ is a conventional carboxy protecting
25 group; it is preferably C₁₋₆alkyl or phenyl; even more
preferably R⁷ is phenyl, methyl, ethyl or t-butyl. When
R⁷ radical is t-butyl, it can be removed by
trifluoroacetic acid.


30 In Step (a) of Scheme IV, a compound of formula XVIII
is reacted with at least two equivalents of R⁵Li in
which R⁵ is as previously defined, but preferably
primary C₁₋₆alkyl. (When a compound of formula XIX in
which R⁵ is hydrogen is desired, it is preferable to

employ a reducing agent which converts a carboxylic acid residue to an aldehyde on a compound of formula XVIII. Many such reducing agents are well known in the art.) Subsequently, an anion of p-


- 5 [(diethoxyphosphoryl)methyl]benzene derivative of formula XX can be reacted with a compound XIX in a routine Horner-Wadsworth-Emmons reaction (see: *Org. React.*, 25, 73-253 (1977); *Stec, Acc. Chem. Res.*, 411-417 (1983)) to afford additional compounds of formula
- 10 I⁵. Subsequent hydrolysis yields a compound of formula I⁶. Alternatively, a compound within the scope of formula I⁶ can be made by a process of Scheme IVa. In Scheme V, R⁸ is a phenolic hydroxy protecting group, such as t-butyldimethylsilyl which can be removed by
- 15 tetrabutylammonium fluoride (TBAF).

- Starting compounds of general formula XVIII in Schemes III, IV, VI, VII, XI, XII and XIII can be prepared by a wide variety of methods using conventional starting
- 20 materials and processes. The syntheses of certain compounds within the scope of formula XVIII are illustrated in Schemes XIX to XXII.


SCHEME I




SCHEME II




SCHEME III




SCHEME IV




SCHEME IVa




SCHEME V




SCHEME VI




Scheme VII




Scheme VIII




Scheme IX




Scheme X




Scheme XI




Scheme XII




Scheme XIII




Scheme XIV




Scheme XV




SCHEME XVI




SCHEME XVII




SCHEME XVIII




SCHEME XIX




SCHEME XX



SCHEME XXI



SCHEME XXII



DESCRIPTION OF SPECIFIC EMBODIMENTS

The specific examples which follow illustrate the synthesis of representative compounds of the instant invention and are not to be construed as limiting the invention in sphere or scope. The methods may be adapted to variations in order to produce compounds embraced by this invention but not specifically disclosed. Further, variations of the methods to produce the same compounds in somewhat different fashion will also be evident to one skilled in the art.

All temperatures are understood to be in Centigrade (C) when not specified. The nuclear magnetic resonance (NMR) spectral characteristics refer to chemical shifts (δ) expressed in parts per million (ppm) versus tetramethylsilane (TMS) as reference standard. The relative area reported for the various shifts in the proton NMR spectral data corresponds to the number of hydrogen atoms of a particular functional type in the molecule. The nature of the shifts as to multiplicity is reported as broad singlet (bs), broad doublet (bd), broad triplet (bt), broad quartet (bq), singlet (s), multiplet (m), doublet (d), quartet (q), triplet (t), doublet of doublet (dd), doublet of triplet (dt), and doublet of quartet (dq). The solvents employed for taking NMR spectra are DMSO- d_6 (perdeuterodimethylsulfoxide), D_2O (deuterated water), $CDCl_3$ (deuteriochloroform) and other conventional deuterated solvents. The infrared (IR) spectral description include only absorption wave numbers (cm^{-1}) having functional group identification value.

Celite is a registered trademark of the Johns-Manville Products Corporation for diatomaceous earth.

- 5 The abbreviations used herein are conventional abbreviations widely employed in the art. Some of which are:

	MS	:	mass spectrometry
10	HRMS	:	high resolution mass spectrometry
	Ar	:	aryl
	DCI	:	desorption (or direct) chemical ionization
	Hex	:	hexane(s)
15	tBu	:	tertiarybutyl
	h	:	hour(s)
	min	:	minute(s)
	Ph	:	phenyl
	Y	:	yield
20	THF	:	tetrahydrofuran
	Tf ₂ O	:	triflic anhydride (trifluoromethanesulfonic anhydride)
	SEMCl	:	2-(trimethylsilyl)ethoxymethyl
25	chloride		

EXAMPLE 1

5,5-Dimethyl-dihydrofuran-2-one (IV)

30

A solution of ethyl levulinate (50.0 g, 0.345 mole) in anhydrous ethyl ether (200 mL) and anhydrous benzene (200 mL) was treated with methylmagnesium bromide (3.0 M solution in diethyl ether, 121.0 mL, 0.365 mole)

dropwise at 0°C over 30 minutes. At this time the ether was removed by slow distillation and the resulting benzene solution was heated at reflux for 2 hours. An ice cold solution of 20% phosphoric acid (500 mL) and ethyl acetate (500 mL) were then added at 0°C. The organic phase was then separated, washed with brine (1 x 300 mL), dried over anhydrous magnesium sulfate, filtered and concentrated in vacuo. The resulting crude oil was purified by distillation (b.p., 43°C, 0.35 mm Hg) yielding 26.0 g of the title furan-2-one (Y: 66%); ¹H-NMR (CDCl₃): δ 2.62 (t, J=8.5 Hz, 2H), 2.05 (t, J=8.5 Hz, 2H), 1.42 (s, 6H).

15

EXAMPLE 2

4,4-Dimethyl-1-tetralone (V)

To a solution of anhydrous aluminum chloride (0.446 mmol, 59.36g) in anhydrous benzene (94.0 mL) at 5°C was added over a period of 45 minutes 5,5-dimethyl-dihydrofuran-2-one (0.149 mole, 17.0 g). The reaction mixture was then slowly warmed to 90-100°C. After 3 hours, the mixture was quenched with ice water, 1N HCl and ethyl acetate at 0°C. The organic phase was then separated and concentrated in vacuo. The residue was chromatographed (eluted with 3% ethyl acetate in hexane) on silica gel to give 18.70g (Y: 72%) of 4,4-dimethyl-1-tetralone; ¹H-NMR (CDCl₃): δ 8.01 (m, 1H), 7.50 (m, 1H), 7.41 (m, 1H), 7.28 (m, 1H), 2.72 (t, J=7.0 Hz, 2H), 2.02 (t, J=7.0 Hz, 2H), 1.39 (s, 6H); MS (DCI) m/e: 174 (MH⁺)

Anal. calcd. for C₁₂H₁₄O₁: C, 81.77; H, 9.14. Found: C,

81.70; H, 9.12.

EXAMPLE 3

5

4,4-Dimethyl-7-nitro-1-tetralone (VI)

The title compound was prepared by Procedure of Heck & Winstein, *J. Org. Chem.*, Vol. 37, No. 6, 1972, p. 825;

10 ¹H-NMR (CDCl₃): δ 8.84 (d, J=2.5 Hz, 1H); 8.36 (dd, J=7.0 Hz, 2.5 Hz, 1H), 7.62 (d, J=7.0 Hz, 2H), 2.80 (t, J=7.0 Hz, 3H), 2.08 (t, J=7.0 Hz, 3H), 1.45 (s, 6H); MS (DCI) m/e: 220 (MH⁺).

15

EXAMPLE 4


4,4-Dimethyl-7-amino-1-tetralone (VII)

20 The title compound was prepared by procedure of Heck & Winstein, *J. Org. Chem.*, Vol. 37, No. 6, 1972, p. 825;

¹H-NMR (CDCl₃): δ 7.28 (d, J=8.5 Hz, 1H), 7.21 (d, J=2.5 Hz, 1H), 6.86 (dd, J=8.5, 2.5 Hz, 1H), 3.73 (bs, 2H), 2.70 (t, J=7.0 Hz, 2H), 1.97 (t, J=7.0 Hz, 2H),
25 1.37 (s, 6H); MS (DCI) m/e: 190 (MH⁺).

EXAMPLE 5

30 4-[[[5,6,7,8-Tetrahydro-5,5-dimethyl-8-oxo-2-naphthalenyl]amino]carbonyl]benzoic acid, methyl ester (IXa)




A solution of monomethyl terephthalate (VIIIa) (2.86g, 15.89 mmol) in thionylchloride (50 mL) with 2 drops of N,N-dimethylformamide was allowed to stir at room temperature. The mixture became homogeneous within 30 minutes and was then concentrated in vacuo. The residue was then taken up in 30 mL of anhydrous pyridine and treated with 4,4-dimethyl-7-amino-1-tetralone (3.00 g, 15.9 mmol). After 16 hours at room temperature 1N HCl was added to the mixture. It was extracted with ethyl acetate, washed with 1N HCl (4 x 200 mL) and washed with saturated sodium bicarbonate (2 x 200 mL). The organic phase was then separated, dried over magnesium sulfate, and concentrated in vacuo to give 5.07g (Y: 91%) of the title compound; ¹H-NMR (CDCl₃): δ 8.27 (dd, J=8.7, 2.5 Hz, 1H), 8.19 (bs, 1H), 8.14 (d, J=8.4 Hz, 2H), 7.95 (d, J=8.4 Hz, 2H), 7.91 (d, J=2.5 Hz, 1H), 7.46 (d, J=8.7 Hz, 1H), 3.94 (s, 3H), 2.70 (t, J=7.0 Hz, 3H), 2.00 (t, J= 7.0 Hz, 3H), 1.38 (s, 6H); MS (DCI) m/e: 352 (MH⁺).

25

EXAMPLE 6


N-[2-(Trimethylsilyl)ethoxymethyl]-4-[[5,6,7,8-tetrahydro-5,5-dimethyl-8-oxo-2-naphthalenyl]aminocarbonyl]benzoic acid, methyl ester
 30 (Xa)



A solution of 4-[[[5,6,7,8-tetrahydro-5,5-dimethyl-8-oxo-2-naphthalenyl)amino]carbonyl]benzoic acid, methyl ester (5.07 g, 14.4 mmol) in anhydrous N,N-dimethylformamide (75 mL) at 0°C was treated with 80% sodium hydride (477 mg, 15.9 mmoles). When hydrogen evolution ceased 2-(trimethylsilyl)ethoxymethyl chloride (3.61 g, 21.7 mmol) was slowly added. After 16 h at room temperature, the mixture was diluted with a 10% sodium bicarbonate solution (100 mL) and extracted with diethyl ether. The organic phase was concentrated in vacuo and the residue chromatographed (eluted with 20% ethyl acetate in hexane) over silica to give 4.24 g (Y: 61%) of the title product; MS (DCI) m/e: 482 (MH⁺); ¹H-NMR (CDCl₃): δ 7.90 (d, J=7.5 Hz, 2H), 7.82 (s, 1H), 7.45 (d, J=7.5 Hz, 2H), 7.25 (m, 2H), 5.22 (bs, 2H), 3.93 (s, 3H), 3.67 (t, J=8.0 Hz, 2H), 2.69 (t, J=7.0 Hz, 2H), 1.98 (t, J=7.0 Hz, 2H), 1.33 (s, 6H), 0.96 (t, J=8.0 Hz, 2H), 0.00 (s, 9H).

EXAMPLE 7

N-[2-(Trimethylsilyl)ethoxymethyl]-4-[[[5,6,7,8-tetrahydro-5,5-dimethyl-8-ethyl-8-hydroxy-2-naphthalenyl)amino]carbonyl]benzoic acid, methyl ester (XIa)




To a solution of compound Xa (1.07g, 2.22 mmol) in anhydrous tetrahydrofuran (15 mL) at -78°C was added 5 ethyl magnesium bromide (3.0 M solution in diethyl ether, 1.05 mL, 3.15 mmol). After 10 minutes, the reaction mixture was allowed to warm to room temperature. After 3 hours at room temperature, the reaction mixture was diluted with ethyl acetate (100 10 mL) and washed with water (100 mL). The organic phase was evaporated and the residue chromatographed on silica gel (eluted with 20% ethyl acetate in hexane) to give 103 mg (Y: 9%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.87 (m, 2H), 7.40 (m, 2H), 7.20 (m, 1H), 15 7.10 (m, 2H), 5.40 (m, 1H), 5.20 (m, 1H), 3.90 (s, 3H), 3.75 (m, 2H), 2.00 (m, 1H), 1.78-1.42 (m, 5H), 1.27 (s, 3H), 1.25 (s, 3H), 1.05 (t, $J=8.0$ Hz, 3H), 0.95 (m, 2H), 0.00 (s, 9H); MS (DCI) m/e : 494 ($\text{MH}^+ - \text{H}_2\text{O}$).

20

EXAMPLE 8

4-[[[5,6-Dihydro-5,5-dimethyl-8-ethyl-2-
25 naphthalenyl)amino]carbonyl]benzoic acid, methyl ester
(I^{1a})



To a solution of compound XIa (169 mg, 0.33 mmol) in toluene (10 mL) was added p-toluenesulfonic acid (pTsOH) monohydrate (a few crystals). After heating at 75°C for 5 minutes, the reaction mixture was concentrated in vacuo and the residue chromatographed on silica gel (eluted with 15% ethyl acetate in hexane) to give 34 mg (Y: 28%) of the title compound;


10 ¹H-NMR (CDCl₃): δ 8.20 (d, J=8.5 Hz, 2H), 7.95 (d, J=8.5 Hz, 2H), 7.50 (m, 2H), 7.35 (d, J=7.0 Hz, 1H), 5.80 (t, J=4.4 Hz, 1H), 3.95 (s, 3H), 2.50 (q, J=7.0 Hz, 2H), 2.20 (d, J=4.4 Hz, 2H), 1.25 (s, 6H), 1.10 (t, J=7.0 Hz, 3H).

15

EXAMPLE 9

4-[[[5,6-Dihydro-5,5-dimethyl-8-ethyl-2-naphthalenyl]aminocarbonyl]benzoic acid (I²a)

20




To a stirred solution of compound I¹a (34 mg, 0.094 mmol) in an ethanol and tetrahydrofuran solution (5 mL, 1:1) was added 10 N NaOH (1.0 mmol, 0.1 mL) at

- room temperature. After 72 hours an excess of 1N HCl (20 mL) was added. The precipitate was collected by vacuum filtration, washed with 1N HCl, water, and air dried to give 21 mg (Y: 64%) of the title compound; ¹H-NMR (DMSO-d₆): δ 10.32 (s, 1H), 8.05 (s, 4H), 7.67 (m, 2H), 7.27 (d, J=9.0 Hz, 1H), 5.78 (t, J=4.4 Hz, 1H), 2.41 (q, J=7.0 Hz, 2H), 2.13 (d, J=4.4 Hz, 2H), 1.18 (s, 6H), 1.12 (t, J=7.0 Hz, 3H); ¹³C-NMR (DMSO-d₆): 166.79, 164.63, 164.55, 140.30, 138.64, 136.99, 136.75, 133.50, 133.26, 129.26, 127.84, 123.94, 122.48, 119.13, 115.43, 38.06, 32.86, 28.23, 25.13, 13.22; MS (DCI) m/e: 350 (MH⁺); IR (KBr): 2962, 1700, 1652, 1532.
- Anal. calcd. for C₂₂H₂₃O₃N₁·0.74 H₂O: C, 72.80; H, 6.80; N, 3.86. Found: C, 72.49; H, 6.56; N, 3.74.

EXAMPLE 10

20

N-[2-(Trimethylsilyl)ethoxymethyl]-4-[[[5,6,7,8-tetrahydro-5,5,8-trimethyl-8-hydroxy-2-naphthalenyl]-aminolcarbonyl]benzoic acid, methyl ester (XIb)



25


- To a solution of compound Xa (438 mg, 0.91 mmol) in anhydrous tetrahydrofuran (10 mL) at -78°C was added methylmagnesium bromide (3.0 M solution in diethyl ether, 0.43 mL, 1.30 mmol). After 10 minutes at -78°C the reaction mixture was allowed to warm to room

temperature. After 3 hours at room temperature, the reaction mixture was diluted with ethyl acetate (100 mL) and washed with water (100 mL). The organic phase was evaporated and the residue chromatographed on silica gel (eluted with 20% ethyl acetate in hexane) to give 90 mg (Y: 20%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 7.90 (m, 2H), 7.40 (m, 2H), 7.25 (m, 2H), 7.10 (m, 1H), 5.25 (m, 2H), 3.90 (s, 3H), 3.70 (m, 2H), 1.90 (t, $J=7.0$ Hz, 2H), 1.75 (t, $J=7.0$ Hz, 2H), 1.30 (s, 3H), 1.29 (s, 3H), 1.27 (s, 3H), 1.00 (t, $J=8.0$ Hz, 3H), 0.00 (s, 9H); MS (DCI) m/e : 480 ($\text{MH}^+ - \text{H}_2\text{O}$).

15

EXAMPLE 11

4-[[[5,6-Dihydro-5,5,8-trimethyl-2-naphthalenyl]-aminocarbonyl]benzoic acid, methyl ester (I^b)



20


To a solution of compound XIb (218 mg, 0.44 mmol) in toluene (10 mL) was added p-toluene sulfonic acid monohydrate (70 mg, 0.37 mmol). After heating at 75°C for 15 minutes, the reaction mixture was concentrated in vacuo and the residue chromatographed on silica gel (eluted with 20% ethyl acetate in hexane) to give 28 mg (Y: 18%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 8.20 (d, $J=9.0$ Hz, 2H), 7.95 (d, $J=9.0$ Hz, 2H), 7.90 (bs, 1H), 7.50 (m, 2H), 7.35 (d, $J=8.5$ Hz, 1H), 5.80 (m, 1H), 3.95 (s, 3H), 2.25 (m, 2H), 2.10 (d, $J=1.4$

Hz, 3H), 1.30 (s, 6H); MS (DCI) m/e: 350 (MH⁺).

EXAMPLE 12

5

4-[[[5,6-Dihydro-5,5,8-trimethyl-2-naphthalenyl]-aminolcarbonyl]benzoic acid (I^{2b})



10


To a stirred solution of compound I^{1b} (0.097 mmol, 34 mg) in a 1:1 ethanol and tetrahydrofuran solution (5 mL) was added 10N NaOH (1.0 mmol, 0.1 mL) at room temperature. After 72 hours, an excess of 1N HCl (20 mL) was added. The precipitate was collected by vacuum filtration, washed with 1N HCl, water, and dried to give 14 mg (Y: 43%) of the title compound; ¹H-NMR (DMSO-d₆): δ 10.32 (s, 1H), 8.04 (s, 4H), 7.66 (dd, J=8.3, 2.1 Hz, 1H), 7.62 (d, J=2.1 Hz, 1H), 7.27 (d, J=8.3 Hz, 1H), 5.80 (m, 1H), 2.14 (m, 2H), 2.01 (d, J=1.4 Hz, 3H), 1.18 (s, 6H); ¹³C-NMR (DMSO-d₆): δ 166.79, 164.64, 139.95, 136.96, 134.15, 130.84, 129.27, 127.85, 124.41, 123.84, 119.30, 115.72, 38.25, 32.96, 28.42, 19.18; MS (DCI) m/e: 336(MH⁺); IR (KBr): 3422, 2962, 1700, 1652, 1532.

25

Anal. calcd for C₂₁H₂₁N₁O₃·1.5 H₂O: C, 69.60; H, 6.68; N, 3.86. Found: C, 69.53; H, 6.97; N, 3.76.

EXAMPLE 13

N-[2-(Trimethylsilyl)ethoxymethyl]-4-[[[5,6,7,8-tetrahydro-5,5-dimethyl-8-phenyl-8-hydroxy-2-naphthalenyl]aminolcarbonyl]benzoic acid, methyl ester
(XIc)




10 To a solution of compound Xa (595 mg, 1.24 mmol) in anhydrous tetrahydrofuran at -78°C was added phenylmagnesium bromide (3.0 M solution in diethyl ether, 0.59 mL, 1.76 mmol). After 10 minutes at -78°C , the reaction mixture was diluted with ethyl
15 acetate (100 mL) and washed with water (100 mL). The organic phase was evaporated and the residue chromatographed on silica gel (eluted with 20% ethyl acetate in hexane) to give 449 mg (Y: 67%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 7.85 (d, 2H), 7.20-
20 6.80 (m, 10H), 5.20 (m, 2H), 3.95 (s, 3H), 3.65 (m, 2H), 2.20-2.00 (m, 2H), 1.85 (m, 1H), 1.50 (m, 1H), 1.40 (s, 3H), 1.30 (s, 3H), 0.95 (m, 2H), 0.00 (s, 9H); MS (DCI) m/e: 542 ($\text{MH}^+ - \text{H}_2\text{O}$).

25

EXAMPLE 14

4-[[[5,6,8-Dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl]aminolcarbonyl]benzoic acid, methyl ester
30 (I'c)




To a solution of compound XIc (as monohydrate, 449 mg, 0.83 mol) in toluene (10 mL) was added p-toluene
 5 sulfonic acid (190 mg, 1.0 mmol). After heating at 75°C for 0.5 hour, the reaction mixture was diluted with ethyl acetate (100 mL) and washed with saturated sodium bicarbonate solution (100 mL). The organic phase was concentrated in vacuo and the residue
 10 chromatographed on silica gel (eluted with 15% ethyl acetate in hexane) to give 162 mg (Y: 48%) of the title compound; ¹H-NMR (CDCl₃): δ 8.15 (d, J=8.5 Hz, 2H), 7.90 (d, J=8.5 Hz, 2H), 7.90 (m, 1H), 7.65 (m, 1H), 7.40 (m, 5H), 6.90 (m, 1H), 6.05 (t, J=4.6 Hz,
 15 1H), 3.95 (s, 3H), 2.40 (d, J=4.6 Hz, 2H), 1.40 (s, 6H); MS (DCI) m/e: 412 (MH⁺).

EXAMPLE 15

20

4-[[5,6-Dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl]aminocarbonyl]benzoic acid (I²c)



25

To a stirred solution of compound I¹c (0.21 mmol, 86

mg) in a 1:1 ethanol and tetrahydrofuran solution (5 mL) was added 10 N NaOH (2.1 mmol, 0.21 mL) at room temperature. After 72 hours, an excess of 1N HCl (20 mL) was added, the precipitate collected by vacuum
 5 filtration, washed with 1N HCl, water, and dried to give 74 mg (Y: 89%) of the title compound; $^1\text{H-NMR}$ (DMSO- d_6): δ 10.29 (s, 1H), 7.98 (m, 4H), 7.76 (dd, $J=8.4$, 2.1 Hz, 1H), 7.35 (m, 7H), 5.97 (t, $J=4.6$ Hz, 1H), 2.89 (d, $J=4.6$ Hz, 2H), 1.27 (s, 6H); $^{13}\text{C-NMR}$
 10 (DMSO- d_6): 166.70, 164.58, 140.46, 140.33, 138.81, 138.62, 136.78, 133.39, 133.10, 129.16, 128.42, 128.35, 127.89, 127.18, 126.83, 124.04, 119.84, 118.03, 38.41, 32.99, 28.04; MS (DCI) m/e : 398 (MH^+); IR (KBr): 3056, 2958, 1700, 1652, 1532.
 15
 Anal. calcd for $\text{C}_{26}\text{H}_{23}\text{O}_3\text{N}_1 \cdot 1.54 \text{ H}_2\text{O}$: C, 73.45; H, 5.81; N, 3.29. Found: C, 73.05; H, 5.53; N, 3.22.

20

EXAMPLE 16

4,4-Dimethyl-7-diazotetrafluorborate-1-tetralone
(XIII)

25 To 4,4-dimethyl-7-amino-1-tetralone (15.10g, 79.89 mmol) was added fluoboric acid (27.86 mL) diluted with water (27.86 mL) at 0°C. A cold solution of sodium nitrate (13.75 g, 199 mmol) in water (27.86 mL) was added slowly while keeping the temperature at about
 30 10°C. The mixture was then cooled to 0°C, filtered and washed with 5% fluoboric acid (200 mL) and dried in vacuo to give 20.5 g (Y: 89%) of the title compound; $^1\text{H-NMR}$ (DMSO- d_6): δ 9.15 (d, $J=2.5$ Hz, 1H), 8.75 (dd, $J=8.5$, 2.5 Hz, 1H), 8.20 (d, $J=8.5$ Hz, 1H), 2.87 (t,

J=7.0 Hz, 2H), 2.07 (t, J=7.0 Hz, 2H), 1.43 (s, 6H);
MS (DCI) m/e: 193 (MH⁺ - N₂BF₄).

5

EXAMPLE 17

4,4-Dimethyl-7-hydroxy-1-tetralone (XIV)

Compound XIII (1.19g, 4.13 mmol) was added to an
10 already boiling solution of sulfuric acid (3.0 mL) and
water (30 mL). After 1 hour at reflux the reaction
mixture was cooled and extracted with ethyl acetate (2
x 50 mL). The combined organic phases were
concentrated in vacuo and the residue chromatographed
15 on silica gel (eluted with 20% ethyl acetate in
hexane) to give 690mg (Y: 88%) of the title compound;
¹H-NMR (CDCl₃): δ 7.48 (d, J=2.5 Hz, 1H), 7.33 (d,
J=8.5 Hz, 1H), 7.05 (dd, J=8.5, 2.5 Hz, 1H), 2.75
(t, J= 7.0 Hz, 2H), 2.00 (t, J=7.0 Hz, 2H), 1.38 (s,
20 6H).

EXAMPLE 18

25 4,4-Dimethyl-7-trifluoromethanesulfonate-1-tetralone
(XV)


To a solution of compound XIV (690 mg, 3.63 mmol) in
anhydrous pyridine (10 mL) was added
30 trifluoromethanesulfonic anhydride (4.42 mmol, 0.74
mL) at 0°C. The reaction mixture was then allowed to
warm to room temperature. After 16 hours, 1N HCl (25
mL) was added and the mixture was extracted with ethyl
acetate (2 x 50 mL). The combined organic phases were

dried over anhydrous magnesium sulfate and concentrated in vacuo to give 1.17g (Y: 100%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 7.88 (d, $J=2.8$ Hz, 1H), 7.52 (d, $J=8.7$ Hz, 1H), 7.40 (dd, $J=8.7$, 2.8 Hz, 1H), 2.76 (t, $J=7.0$ Hz, 2H), 2.04 (t, $J=7.0$ Hz, 2H), 1.40 (s, 6H); MS (DCI) m/e : 323 (MH^+).

EXAMPLE 19

10

5,5-Dimethyl-8-oxo-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid, methyl ester (XVIa)



15

To a solution of compound XV (1.15g, 3.57 mmol) in methanol (10.8 mL) and dimethyl sulfoxide (10.8 mL) was added triethylamine (1.09 mL, 7.82 mmol) palladium (II) acetate (24 mg, 0.11 mmol) and 1,3-bis(diphenylphosphino)propane (44 mg, 0.11 mmol). The reaction mixture was then saturated with carbon monoxide at room temperature and heated to 70°C under a balloon of carbon monoxide for 3 hours. After cooling to room temperature, the reaction mixture was poured into water and extracted with ethyl acetate. The organic phase was then concentrated in vacuo and the residue chromatographed on silica gel (eluted with 15% ethyl acetate in hexane) to give 692 mg (Y: 93%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 8.66 (d, $J=2.0$ Hz, 1H), 8.18 (dd, $J=8.3$, 2.0 Hz, 1H), 7.52 (d, $J=8.3$ Hz, 1H), 3.92 (s, 3H), 2.76 (t, $J=7.0$ Hz, 2H), 2.04


30

(t, J=7.0 Hz, 2H), 1.41 (s, 6H); MS (DCI) m/e: 233 (MH⁺).

5

EXAMPLE 20

5,5-Dimethyl-8-hydroxy-8-phenyl-5,6,7,8-tetrahydro-naphthalene-2-carboxylic acid, methyl ester (XVIIa)



10


To a solution of compound XVIa (167 mg 0.72 mmol,) in tetrahydrofuran (5 mL) at -78°C was added phenylmagnesium bromide (3.0 M solution in diethyl
15 ether, 1.08 mmol, 0.35 mL). After warming to room temperature (2 hours), the reaction mixture was concentrated and the residue chromatographed on silica gel (eluted with 10% ethyl acetate in hexane) to give 152 mg (Y: 68%) of the title product; ¹H-NMR (CDCl₃):
20 δ 7.95 (dd, J=8.3, 2.0 Hz, 1H), 7.85 (d, J=2.0 Hz, 1H), 7.50 (d, J=8.3 Hz, 1H), 7.25 (m, 5H), 3.82 (s, 3H), 2.20 (m, 2H), 1.85 (m, 1H), 1.60 (m, 1H), 1.43 (s, 3H), 1.38 (s, 3H); MS (DCI) m/e: 311 (MH⁺).

25

EXAMPLE 21

5,5-Dimethyl-5,6-dihydro-8-phenyl-naphthalene-2-carboxylic acid (XVIIIa)


30



To a solution of compound XVIIa (150 mg, 0.484 mmol) in toluene (7 mL) was added a few milligrams (2-4 mg) of p-toluenesulfonic acid. After heating at 70°C for 5 minutes, the reaction mixture was cooled and concentrated in vacuo. The residue was then dissolved in ethyl alcohol (7 mL) and treated with 10 N NaOH (7.5 mmol, 0.74 mL) at room temperature. After 16 hours, an excess of 1N HCl (30 mL) was added and the precipitate collected by vacuum filtration to give 135 mg (Y: 99%) of the title compound; ¹H-NMR (DMSO-d₆): δ 7.95 (dd, J=2.0, 8.5 Hz, 1H), 7.75 (d, J=2.0 Hz, 1H), 7.47 (d, J=8.5 Hz, 1H), 7.38 (m, 5H), 6.05 (t, J=4.6 Hz, 1H), 2.40 (d, J=4.6 Hz, 2H), 1.40 (s, 6H); MS (DCI) m/e: 279 (MH⁺).

EXAMPLE 22

4-[[[5,6-Dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl]carbonyl]aminobenzic acid, methyl ester (I^{3a})




A solution of compound XVIIIa (135 mg, 0.485 mmol) in

thionylchloride (5 mL) with 2 drops of N,N-dimethylformamide was allowed to stir at room temperature. The mixture became homogeneous within 1 hour and was then concentrated in vacuo. The residue
 5 was dissolved in anhydrous pyridine (5 mL) to which was added methyl 4-aminobenzoate (Aldrich, 0.534 mmol, 81 mg). After 16 hours at room temperature, the mixture was diluted with 1N HCl, extracted with ethyl acetate (100 mL), washed with 1N HCl (3 x 100 mL) and
 10 washed with saturated sodium bicarbonate (2 x 100 mL). The organic phase was then separated, dried over magnesium sulfate and concentrated in vacuo to give 68 mg (Y: 34%) of the title compound; ¹H-NMR (CDCl₃): δ 8.05 (d, J=7.0 Hz, 2H), 7.75 (m, 2H), 7.65 (d, J=7.0
 15 Hz, 2H), 7.50 (m, 1H), 7.40 (m, 5H), 6.08 (t, J=4.6 Hz, 1H), 3.90 (s, 3H), 2.45 (d, J=4.6 Hz, 2H), 1.40 (s, 6H); MS (DCI) m/e: 412 (MH⁺).

20

EXAMPLE 23

4-[[5,6-Dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl]carbonyl]aminobenzoic acid (I⁴a)



25

To a stirred solution of compound I³a (0.165 mmol, 68 mg) in ethanol (5 mL) was added 10N NaOH (.165 mL, 1.65 mmol) at room temperature. After 72 hours, an
 30 excess of 1N HCl (30 mL) was added. The precipitate was collected by vacuum filtration, washed with 1N HCl


and water, and air dried to give 45 mg (Y: 69%) of the title compound; $^1\text{H-NMR}$ (DMSO-d_6): δ 10.42 (s, 1H), 7.83 (m, 5H), 7.52 (d, $J=8.0$ Hz, 1H), 7.38 (m, 6H), 6.05 (t, $J=4.6$ Hz, 1H), 2.33 (d, $J=4.6$ Hz, 2H), 1.30 (s, 6H); $^{13}\text{C-NMR}$ (DMSO-d_6): 166.91, 166.00, 148.26, 143.25, 143.14, 139.86, 138.29, 133.36, 132.48, 130.17, 128.55, 128.31, 127.41, 127.31, 126.84, 125.40, 125.00, 123.95, 119.38, 119.30, 37.99, 33.52, 27.75; MS (DCI) m/e : 398 (MH^+); IR (KBr): 2958, 1688, 1596, 1522.

Anal. calcd for $\text{C}_{26}\text{H}_{23}\text{N}_1\text{O}_3 \cdot 0.5 \text{ H}_2\text{O}$: C, 76.83; H, 5.95; N, 3.45. Found: C, 76.47; H, 6.00; N, 3.22.

15

EXAMPLE 24

5,5-Dimethyl-8-ethyl-8-hydroxy-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid, methyl ester
20 (XVIIb)




Using a method analogous to the preparation of the 8-phenyl derivative XVIIa, 490 mg (2.11 mmol) of compound XVIa gave 130 mg (Y: 24%) of the title compound.

EXAMPLE 25

5,6-Dihydro-8-ethyl-5,5-dimethylnaphthalene-2-
carboxylic acid (XVIIIb)

5




Using a method analogous to the the preparation of 8-
 10 phenyl derivative XVIIIa, 130 mg (0.49 mmol) of
 compound XVIIb gave 113 mg (Y: 100%) of the title
 product; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$): δ 7.98 (d, $J=2.0$ Hz, 1H),
 7.95 (dd, $J=8.5$, 2.0 Hz, 1H), 7.40 (d, $J=8.5$ Hz, 1H),
 5.83 (t, $J=4.6$ Hz, 1H), 2.55 (q, $J=7.5$ Hz, 2H), 2.15
 15 (d, $J=4.6$ Hz, 2H), 1.27 (s, 6H), 1.18 (t, $J=7.5$ Hz,
 3H).

EXAMPLE 26

20

4-[[[5,6-Dihydro-5,5-dimethyl-8-ethyl-2-
naphthalenyl)carbonyl]aminobenzoyl]methyl ester
(I³b)



25

Using a method analogous to the preparation of the 8-
 phenyl derivative I³a, 113 mg (0.491 mmol) of compound


XVIIIb gave 163 mg (Y: 92%) of the title compound;
¹H-NMR (CDCl₃): δ 8.08 (d, J=8.5 Hz, 2H), 7.95 (bs, 1H), 7.79 (d, J=2.0 Hz, 1H), 7.75 (d, J=8.5 Hz, 2H), 7.65 (dd, J=8.0, 2.0 Hz, 1H), 7.40 (d, J=8.0 Hz, 1H),
 5 5.88 (t, J=4.6 Hz, 1H), 3.90 (s, 3H), 2.55 (q, J=7.5 Hz, 2H), 2.25 (d, J=4.6 Hz, 2H), 1.30 (s, 6H), 1.18 (t, J=7.5 Hz, 3H); MS (DCI) m/e: 364 (MH⁺).

10

EXAMPLE 27

4-[[[(5,6-Dihydro-5,5-dimethyl-8-ethyl-2-naphthalenyl)carbonyl]aminol]benzoic acid
 (I⁴b)

15



Using a method analogous to the preparation of the 8-phenyl derivative I⁴a, 163 mg (0.45 mmol) of compound
 20 I³b gave 128 mg (Y: 92%) of the title compound; ¹H-NMR (DMSO-d₆): δ 10.46 (s, 1H), 7.91 (m, 4H), 7.77 (m, 2H), 7.45 (d, J=7.9 Hz, 1H), 5.84 (t, J=4.6 Hz, 1H), 2.52 (q, J=7.4 Hz, 2H), 2.17 (d, J=4.6 Hz, 2H), 1.21 (s, 6H), 1.11 (t, J=7.4 Hz, 3H); ¹³C-NMR (DMSO-d₆): 166.97
 25 166.133, 148.61, 143.33, 136.36, 133.27, 132.54, 130.19, 126.72, 125.52, 123.95, 122.80, 122.11, 119.50, 119.41, 37.62, 33.42, 27.94, 24.65, 12.86; MS (DCI) m/e: 350 (MH⁺); IR (KBr): 2964, 1690, 1596, 1524.

30

Anal. calcd. for C₂₂H₂₃O₃N₁·0.75 H₂O: C, 72.81; H, 6.80;


N, 3.86. Found: C, 72.84; H, 6.61; N, 3.86.

5

EXAMPLE 28

8-Hydroxy-5,5,8-trimethyl-5,6,7,8-
tetrahydronaphthalene-2-carboxylic acid, methyl ester
(XVIIc)

10




Using a method analogous to the preparation of the 8-phenyl derivative XVIIa, 292 mg (1.26 mmol) of
 15 compound XVIa gave 188 mg (Y: 60%) of the title
 product; $^1\text{H-NMR}$ (CDCl_3): δ 8.29 (d, $J=2.0$ Hz, 1H), 7.85
 (dd, $J=8.0$ Hz, 2.0 Hz, 1H), 7.38 (d, $J=8.0$ Hz, 1H),
 3.90 (s, 3H), 2.0 (t, $J=7.0$ Hz, 2H), 1.87 (t, $J=7.0$
 Hz, 2H), 1.60 (d, $J=1.3$ Hz, 3H), 1.38 (s, 6H).

20

EXAMPLE 29

5,6-Dihydro-5,5,8-trimethylnaphthalene-2-carboxylic
 25 acid (XVIIIc)




Using a method analogous to the preparation of the 8-phenyl derivative XVIIIa, 188 mg (0.76 mmol) of compound XVIIc gave 154 mg (Y: 94%) of the title compound; $^1\text{H-NMR}$ (DMSO-d_6): δ 7.96 (m, 2H), 7.40 (d, J=8.0 Hz, 1H), 5.87 (m, 1H), 2.27 (m, 2H), 2.17 (d, J=1.3 Hz, 3H), 1.27 (s, 6H).

EXAMPLE 30

10

4-[[[5,6-Dihydro-5,5,8-trimethyl-2-naphthalenyl)carbonyl]aminolbenzoic acid, methyl ester (I³c)



15


Using a method analogous to the preparation of the 8-phenyl derivative I³a, 154 mg (0.713 mmol) of compound XVIIc gave 90 mg (Y: 36%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 8.05 (d, J=8.5 Hz, 2H), 7.90 (bs, 1H), 7.75 (d, J=8.5 Hz, 2H), 7.75 (d, J=2.0 Hz, 1H), 7.65 (dd, J=8.0, 2.0 Hz, 1H), 7.43 (d, J=8.0 Hz, 1H), 5.87 (m, 1H), 3.92 (s, 3H), 2.27 (m, 2H), 2.18 (d, J=1.3 Hz, 3H), 1.30 (s, 6H); MS (DCI) m/e: 350 (MH^+).

25

EXAMPLE 31

4-[[[5,6-Dihydro-5,5,8-trimethyl-2-naphthalenyl)carbonyl]aminolbenzoic acid (I⁴c)

30




- Using a method analogous to the preparation of the 8-phenyl derivative I⁴a, 90 mg (0.26 mmol) of compound
- 5 I³c gave 70 mg (Y: 81%) of the title compound; ¹H-NMR (DMSO-d₆): δ 12.72 (bs, 1H), 10.46 (s, 1H), 7.89 (m, 5H), 7.79 (dd, J=8.0, 2.0 Hz, 1H), 7.72 (d, J=2.0 Hz, 1H), 7.45 (d, J=8.0 Hz, 1H), 5.85 (m, 1H), 2.17 (m, 1H), 2.09 (d, J=1.3 Hz, 3H), 1.22 (s, 6H); MS (DCI)
- 10 m/e: 336 (MH⁺); IR (KBr): 2958, 1674, 1656, 1416; ¹³C-NMR: 166.93, 166.13, 148.21, 143.38, 133.98, 132.58, 130.63, 130.22, 126.89, 125.36, 124.90, 123.83, 122.48, 119.48, 119.39, 37.84, 33.54, 28.14, 19.12.
- 15 Anal. calcd for C₂₁H₂₁N₁O₃: C, 75.20; H, 6.31; N, 4.1. Found C, 74.90; H, 6.36; N, 3.99.

20

EXAMPLE 32

1-(5,6-Dihydro-5,5-dimethyl-8-phenyl-naphthalen-2-yl)-ethanone (XIXa)



25


To a stirred solution of compound XVIIIa (816 mg, 2.94 mmol) in diethyl ether (15.0 mL) at -78°C was added methyllithium (1.4 M solution in diethyl ether, 4.19 mL, 5.88 mmol). After 1 hour at room temperature, 1N HCl (50 mL) was added. The organic phase was separated, washed with brine (50 mL), 1N NaOH (50 mL), dried over anhydrous magnesium sulfate and concentrated to give 600 mg (Y: 74%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 7.83 (dd, $J=8.0, 1.7$ Hz, 1H), 7.63 (d, $J=1.7$ Hz, 1H), 7.46 (d, $J=8.0$ Hz, 1H), 7.37 (m, 5H), 6.05 (t, $J=4.7$ Hz, 1H), 2.44 (s, 3H), 2.37 (d, $J=4.7$ Hz, 2H), 1.35 (s, 6H); MS (DCI) m/e : 277 (MH^+).

15

EXAMPLE 33

4-(E,Z)-[2-(5,6-dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)-1-propenyl]benzoic acid, ethyl ester (I^5a)

20




Methyl *p*-[(diethoxyphosphoryl)methyl]benzoate (1.02 g, 3.40 mmol, prepared as in Liebigs Ann. Chem. 1985, 929) was added to a 1M dimsyl anion solution in dimethylsulfoxide (3.09 mL, 3.09 mmol, prepared by warming sodium hydride in dimethylsulfoxide for 1 hour at 65°C). After 30 minutes, the mixture was added to a solution of compound XIXa (595 mg, 2.16 mmol) in dimethylsulfoxide (6.50 mL) at room temperature. After

30

3 hours at room temperature, a 2 M solution of sodium ethoxide in ethanol (1.74 mL, 3.48 mmol) was added. After 16 hours at room temperature, the mixture was diluted with 5% sodium bicarbonate (50 mL) and
 5 extracted with diethyl ether (50 mL x 3). The combined organic phases were concentrated in vacuo and the residue chromatographed on silica gel (5% ethyl acetate in hexane) to give 159 mg (Y: 18%) of the title compound. NMR indicated a mixture of isomers
 10 (E:Z) (4:1).

EXAMPLE 34

15 4-(E)-[2-(5,6-dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)-1-propenyl]benzoic acid (I⁵a)



20 To a stirred solution of compound I⁵a (159 mg, 0.388 mmol) in ethanol (15.0 mL) at room temperature was added a 10 N NaOH solution (0.39 mL, 3.90 mmol). After 48 hours, the mixture was diluted with an excess of 1N HCl (40.0 mL) and then filtered. After washing with
 25 water and drying, the solid was recrystallized from ethanol to provide 60 mg (Y: 40%) of the title compound; ¹H-NMR (DMSO-d₆): δ 7.91 (d, J=8.3 Hz, 2H), 7.42 (m, 9H), 7.06 (d, J=1.7 Hz, 1H), 6.70 (s, 1H), 6.03 (t, J=4.6 Hz, 1H), 2.34 (d, J=4.6 Hz, 2H), 2.12
 30 (s, 3H), 1.31 (s, 6H); ¹³C-NMR (DMSO-d₆): 144.42,

140.58, 140.14, 138.94, 138.72, 133.12, 129.26,
129.07, 128.47, 128.35, 127.32, 126.71, 125.89,
125.32, 124.07, 123.06, 38.25, 33.16, 27.94, 17.38;
MS (DCI) m/e: 395 (MH⁺); IR (KBr): 2958, 1680, 1602,
5 1292.


Anal. calcd. for C₂₇H₂₆O₂: C, 84.78; H, 6.85. Found: C,
84.98; H, 6.66.

10

EXAMPLE 35

4,4-Dimethyl-7-[(t-butyltrimethylsilyl)oxy]-1-tetralone
(XXIa)

15




To a solution of compound XIV (2.00 g, 10.5 mmol) in
dimethylformamide (16 mL) was added tert-
20 butyldimethylsilyl chloride (1.90 g, 12.6 mmol) and
imidazole (1.79 g, 26.3 mmol) at room temperature.
After 5 hours, 5% NaHCO₃ (50 mL) was added and the
mixture was extracted with hexane (2 x 75 mL). The
combined organic phases were dried over anhydrous
25 magnesium sulfate and concentrated in vacuo to give
3.22 g (Y: 99%) of the title compound; ¹H-NMR (CDCl₃):
δ 7.44 (d, J=2.8 Hz, 1H), 7.29 (d, J=8.7 Hz, 1H), 7.01
(dd, J=8.7, 2.8 Hz, 1H), 2.71 (t, J=7.0 Hz, 2H), 1.99
(t, J=7.0 Hz, 2H), 1.36 (s, 6H), 0.98 (s, 9H), 0.20
30 (s, 6H); MS (DCI) m/e: 305 (MH⁺).

EXAMPLE 36

2-[(t-Butyldimethylsilyl)oxyl]-5,6-dihydro-5,5-
dimethyl-8-trifluoromethanesulfonylnaphthalene (XXIIa)


5



To a solution of compound XXIa (2.73 g, 8.98 mmol) in tetrahydrofuran (80.0 mL) at -78°C was added sodium
 10 bis(trimethylsilyl)amide (1.0 M solution in tetrahydrofuran, 9.88 mmol, 9.88 mL) and N-(2-pyridyl)triflimide (9.88 mmol, 3.54 g). After stirring for 1 hour at -78°C and 1.5 hours at room temperature, the reaction mixture was concentrated in
 15 vacuo and the residue chromatographed on silica gel (eluted with 10% ethyl acetate in hexane) to give 2.26 g (Y: 58%) of the title product; ¹H-NMR (CDCl₃): δ 7.15 (d, J=8.4 Hz, 1H), 6.85 (d, J=2.5 Hz, 1H), 6.78 (dd, J=8.4 Hz, 2.5 Hz, 1H), 5.94 (t, J=4.8 Hz, 1H), 2.38
 20 (d, J=4.8 Hz, 2H), 1.27 (s, 6H), 0.98 (s, 9H), 0.20 (s, 6H); MS (DCI) m/e: 437 (MH⁺).

EXAMPLE 37

25 2-(t-Butyldimethylsilyloxy)-5,6-dihydro-5,5-dimethyl-
8-phenyl-naphthalene (XXIIIa)




To a solution of compound XXIIa (2.26 g, 5.18 mmol) in 1-methyl-2-pyrrolidinone (25.0 mL) at room temperature
 5 was added triphenylarsine (300 mg, 0.980 mmol), tris(dibenzylideneacetone) dipalladium (0) (147 mg, 0.16 mmol), and tributylphenyltin (4.18 g, 11.4 mmol). After stirring at 85°C for 16 hours, water (50 mL) and ethyl acetate (50 mL) were added. The organic phase
 10 was separated and stirred over an aqueous saturated potassium fluoride solution for 30 minutes. The organic phase was again separated, concentrated in vacuo and the residue chromatographed on silica gel (eluted with 100% hexane) to give 1.10 g (Y: 58%) of
 15 the title product; ¹H-NMR (CDCl₃): δ 7.10 (m, 5H), 6.97 (d, J=8.4 Hz, 1H), 6.41 (dd, J=8.4 Hz, 2.5 Hz, 1H), 6.20 (d, J=2.5 Hz, 1H), 5.68 (t, J=4.8 Hz, 1H), 2.05 (d, J=4.8 Hz, 2H), 1.02 (s, 6H), 0.60 (s, 9H), 0.00 (s, 6H); MS (DCI) m/e: 365 (MH⁺).

20

EXAMPLE 38

5,6-Dihydro-5,5-dimethyl-8-phenyl-naphthalen-2-ol
 (XXVa)


25



To a solution of compound XXIIIa 1.10 g, 3.02 mmol) in tetrahydrofuran (10.0 mL) at room temperature was
 5 added tetrabutylammonium fluoride (1.0 M solution in tetrahydrofuran, 3.32 mL, 3.32 mmol). After 5 minutes the reaction mixture was concentrated in vacuo and the residue chromatographed on silica gel (eluted with 10% ethyl acetate in hexane) to give 575 mg (Y: 76%) of
 10 the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.36 (m, 5H), 7.23 (d, $J=8.3$ Hz, 1H), 6.71 (dd, $J=8.3, 2.7$ Hz, 1H), 6.49 (d, $J=2.7$ Hz, 1H), 5.99 (t, $J=4.6$ Hz, 1H), 2.34 (d, $J=4.6$ Hz, 2H), 1.32 (s, 6H); MS (DCI) m/e : 251 (MH^+).

15

EXAMPLE 39


Di-tert-butyl terephthalate (XXVI)

20

To a solution of terephthaloylchloride (2.00 g, 9.85 mmol) in dry pyridine (25.0 mL) was added tert-butyl alcohol (803 mg, 10.8 mmol). After 16 hours at 85°C water (75.0 mL) was added and the solid filtered. The

solid was dissolved in diethyl ether (40 mL) and washed with saturated sodium bicarbonate (2 x 75 mL). The organic phase was then dried over anhydrous magnesium sulfate and concentrated in vacuo to give 1.15 g (Y: 42%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 8.01 (s, 4H), 1.60 (s, 18H).

EXAMPLE 40

10 Mono-tert-butyl terephthalate (XXVII)


To a solution of potassium hydroxide (255 mg, 4.55 mmol) in 4.0 mL tert-butyl alcohol with 0.5 mL of water added for solubility was added compound XXVI (1.15 g, 4.14 mmol) in tert-butyl alcohol (5.5 mL). After 3 hours at 50°C, diethyl ether (35 mL) was added and the reaction mixture filtered. The solid was dissolved in water (35.0 mL), extracted with methylene chloride (2 x 50 mL), and then acidified with 1N HCl. The precipitate was then collected to give 270 mg (Y: 30%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 8.02 (m, 4H), 1.56 (s, 9H); MS (DCI) m/e: 223 (MH^+).

25

EXAMPLE 41

4-[[[5,6-Dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl]oxy]carbonyl]benzoic acid, tert-butyl ester (I'a)

30




To a solution of compound XXVII (107 mg, 0.480 mmol) in methylene chloride (5 mL) at 0°C was added oxalyl chloride (0.15 mL) and dimethylformamide (2 drops). After 2 hours at room temperature, the reaction mixture was concentrated in vacuo to give the corresponding acid chloride (compound XXIVa). To a solution of compound XXIVa (0.480 mmol) in dry pyridine (5 mL) was then added compound XXVa (120 mg, 0.48 mmol). After 16 hours at room temperature, the mixture was diluted with 1N HCl, extracted with ethyl acetate (100 mL), washed with 1N HCl (4 x 100 mL), and washed with saturated sodium bicarbonate (2 x 100 mL). The organic phase was then separated, dried over magnesium sulfate and concentrated in vacuo to give 137 mg (Y: 63%) of the title compound; ¹H-NMR (CDCl₃) δ 8.16 (d, J=8.4 Hz, 2H), 8.06 (d, J=8.4 Hz, 2H), 7.35 (m, 6H), 7.07 (dd, J=8.4 Hz, 2.4 Hz, 1H), 6.84 (d, J=2.4 Hz, 1H), 6.04 (t, J=4.6 Hz, 1H), 2.38 (d, J=4.6 Hz, 2H), 1.61 (s, 9H), 1.36 (s, 6H); MS (DCI) m/e: 455 (MH⁺).

EXAMPLE 42

25

4-(((5,6-Dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)oxycarbonyl)benzoic acid (I^a)




To a stirred solution of compound I'a (137 mg, 0.30 mmol) in methylene chloride was added trifluoroacetic acid (0.270 mL) at room temperature. After 72 hours, 1N HCl (30 mL) was added. The precipitate was collected by vacuum filtration, washed with 1N HCl, and air dried to give 85 mg (Y: 71%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 13.39 (bs, 1H), 8.15 (d, J=7.6 Hz, 2H), 8.05 (d, J=7.6 Hz, 2H), 7.38 (m, 6H), 7.18 (d, J=8.4 Hz, 1H), 6.73 (s, 1H), 6.06 (t, J=4.2 Hz, 1H), 2.35 (d, J=4.2 Hz, 2H), 1.32 (s, 6H); $^{13}\text{C-NMR}$ (CDCl_3): 166.48, 163.99, 148.51, 142.55, 139.77, 138.17, 35.24, 134.59, 132.45, 131.61, 130.01, 129.66, 129.48, 128.57, 128.28, 127.63, 127.41, 125.24, 120.74, 118.36, 67.39, 38.24, 38.07, 33.12, 28.37; MS (DCI) m/e: 399 (MH^+); IR (KBr): 2964, 1738, 1698, 1246.

Anal calcd. for $\text{C}_{24}\text{H}_{22}\text{O}_4 \cdot 0.5 \text{ H}_2\text{O}$: C, 76.64; H, 5.69.
 Found: C, 76.66; H, 5.89.

EXAMPLE 43


4-Hydroxybenzoic acid tert-butyl ester (XXVIIIa)



To a solution of 4-hydroxybenzoic acid (2.00 g, 14.5 mmol) in 1,4-dioxane (10.0 mL) saturated with isobutylene at -78°C in a metal bomb was added concentrated sulfuric acid (0.150 mL). The bomb was sealed and warmed to room temperature. After 72 hours at room temperature, the reaction mixture was cooled to -78°C , poured into saturated sodium bicarbonate (30.0 mL) and extracted with diethyl ether (2 x 50.0 mL). The combined organic phases were concentrated and the residue chromatographed on silica gel (eluted with 5% methanol in methylene chloride to give 290 mg (Y: 10%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.89 (d, $J=8.8$ Hz, 2H), 6.83 (d, $J=8.8$ Hz, 2H), 1.57 (s, 9H).

EXAMPLE 44


4-[[[5,6-Dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl]carbonyloxy]benzoic acid, tert-butyl ester (I^{9a})



To a stirred solution of compound XVIIIa (200 mg, 0.720 mmol) in dry methylene chloride (7.00 mL) at 0°C was added oxalyl chloride (0.075 mL) and dimethylformamide (2 drops). After 2 hours at room temperature, the mixture was concentrated in vacuo. To the residue in anhydrous pyridine (5.00 mL) was added compound XXVIIIa (154 mg, 0.792 mmol). After 2 hours at room temperature, the mixture was diluted with 1N HCl, extracted with ethyl acetate (50 mL), washed with 1N HCl (4 x 100 mL) and washed with saturated sodium bicarbonate (2 x 100 mL). The organic phase was then separated, concentrated in vacuo and the residue chromatographed on silica gel (eluted with 3% ethyl acetate in hexane) to give 203 mg (Y: 61%) of the title product; ¹H-NMR (CDCl₃): δ 8.05 (m, 3H), 7.85, (d, J=1.7 Hz, 1H), 7.51 (d, J=8.1 Hz), 7.38 (m, 6H), 7.18 (d, J=8.6 Hz, 1H), 6.07 (t, J=4.6 Hz, 1H), 2.40 (d, J=4.6 Hz, 2H), 1.59 (s, 9H), 1.38 (s, 6H); MS (DCI) m/e: 455 (MH⁺).

EXAMPLE 45

4-[[[5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyloxy]benzoic acid (I^{10a})




To a solution of compound I^{9a} (203 mg, 0.480 mmol) in methylene chloride (5.00 mL) was added trifluoroacetic acid (0.400 mL) at room temperature. After 16 hours,

1N HCl (30 mL) was added. The precipitate was collected by vacuum filtration, washed with 1N HCl and air dried to give 133 mg (Y: 75%) of the title compound; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$): δ 8.05-7.31 (m, 12H), 6.10 (t, $J=4.6$ Hz, 1H), 2.39 (d, $J=4.6$ Hz, 2H), 1.34 (s, 6H); IR (KBr): 2962, 1742, 1684, 1602; MS (DCI) m/e : 399 (MH^+).

Anal. calcd. for $\text{C}_{24}\text{H}_{22}\text{O}_4 \cdot 0.850 \text{ H}_2\text{O}$: C, 75.47; H, 5.77.
10 Found: C, 75.52; H, 5.43.

EXAMPLE 46

15 5,5-Dimethyl-8-hydroxy-8-(2-fluorophenyl)-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid, methyl ester (XVIIId)




20 To a solution of compound XVIa (297 mg, 1.28 mmol) in tetrahydrofuran (7.0 mL) at -78°C was added 2-fluorophenyllithium [0.35 M solution in THF, 1.93 mmol, 5.50 mL; prepared by treating 2-fluoro-bromobenzene (552 mg, 3.15 mmol) in tetrahydrofuran (5.0 mL) with
25 t-butyllithium (1.7 M solution in pentane, 2.41 mmol, 4.09 mL) at -78°C . After 5 minutes at -78°C , 5.50 mL was transferred via syringe to the solution of compound XVIa]. After warming to room temperature (1 hour), the reaction mixture was concentrated and the
30 residue chromatographed on silica gel (eluted with 10%

ethyl acetate in hexane) to give 280 mg (Y: 67%) of the title product; $^1\text{H-NMR}$ (CDCl_3): (dd, $J=8.4$, 1.7 Hz, 1H), 7.72 (d, $J=1.7$ Hz, 1H), 7.50 (m, 2H), 7.18 (m, 1H), 6.96 (m, 2H), 3.80 (s, 3H), 2.50 (m, 1H), 2.40 (d, $J=2.8$ Hz, 1H), 2.05 (m, 2H), 1.60 (m, 1H), 1.43 (s, 3H), 1.35 (s, 3H). MS (DCI) m/e : 329 (MH^+).

EXAMPLE 47

10

5,5-Dimethyl-5,6-dihydro-8-(2-fluorophenyl)-naphthalene-2-carboxylic acid (XVIIId)




15

Using the method given for the preparation of 8-phenyl derivative XVIIIa, 280 mg (0.85 mmol) of compound XVIIId gave 209 mg (Y: 83%) of the title product; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$): δ 7.81 (dd, $J=8.0$, 1.8 Hz, 1H), 7.55-7.25 (m, 6H), 6.09 (t, $J=4.5$ Hz, 1H), 2.38 (d, $J=4.5$ Hz, 2H), 1.32 (s, 6H); MS (DCI) m/e 297 (MH^+).

20

EXAMPLE 48

4-[[[5,6-Dihydro-5,5-dimethyl-8-(2-fluorophenyl)-2-naphthalenyl]carbonyl]amino]benzoic acid, methyl ester
 5 (I³g)




A solution of compound XVIIIId (200 mg, .676 mmol) in
 10 anhydrous methylene chloride (5.0 mL) was treated with
 oxalyl chloride (0.20 mL, 2.29 mmol) and 2 drops of
 N,N-dimethylformamide at 0°C. The reaction mixture was
 then allowed to stir at room temperature. After 2
 hours, the mixture was concentrated in vacuo. The
 15 residue was dissolved in anhydrous pyridine (5.0 mL)
 to which was added methyl 4-aminobenzoate (Aldrich,
 102 mg, 0.68 mmol). After 2 hours at room temperature,
 the mixture was diluted with 1N HCl, extracted with
 ethyl acetate (100 mL), washed with 1N HCl (3 x 100
 20 mL) and washed with saturated sodium bicarbonate (2 x
 100 mL). The organic phase was then separated and
 concentrated in vacuo. The residue chromatographed on
 silica gel (eluted with 20% ethyl acetate in hexane)
 to give 226 mg (Y: 78%) of the title compound; ¹H-NMR
 25 (CDCl₃): δ 8.02 (d, J=8.7 Hz, 1H), 7.71 (m, 1H), 7.64
 (d, J=8.7 Hz, 1H), 7.48 (d, J=8.0 Hz, 1H), 7.40-7.10
 (m, 7H), 6.11 (t, J=4.5 Hz, 1H), 3.90 (s, 3H), 2.42
 (d, J=4.5 Hz, 2H), 1.39 (s, 6H). MS (DCI) m/e: 430
 (MH⁺).

EXAMPLE 49

4-[[[5,6-Dihydro-5,5-dimethyl-8-(2-fluorophenyl)-2-naphthalenyl]carbonyl]amino]benzoic acid (I⁴g)

5



Using the method given for the preparation of the 8-phenyl derivative I⁴a, 226 mg (0.527 mmol) of compound
 10 I³g gave 180 mg (Y: 82%) of the title compound; ¹H-NMR (DMSO-d₆) δ 12.73 (s, 1H), 10.43 (s, 1H), 7.89-7.78 (m, 5H), 7.53 (d, J=8.0 Hz, 1H), 7.48-7.19 (m, 5H), 6.10 (t, J=4.5 Hz, 1H), 2.38 (d, J=4.5 Hz, 2H), 1.33 (s, 6H); ¹³C-NMR (DMSO-d₆): δ 166.89, 165.80, 161.12,
 15 147.85, 143.24, 133.08, 132.95, 132.57, 131.54, 130.18, 129.93, 129.48, 126.91, 125.37, 124.88, 124.29, 123.86, 119.42, 119.33, 115.75, 115.46, 37.89, 33.48, 27.75.

20 MS (DCI) m/e: 416 (MH⁺).
 IR (KBr): 2962, 1688, 1594, 1520.


Anal. calcd. for C₂₆H₂₂N₁O₃F₁ · 0.25 H₂O: C, 74.36; H, 5.40; N, 3.34. Found: C, 74.50; H, 5.40; N, 3.17

25

EXAMPLE 50

5,5-Dimethyl-8-oxo-5,8-dihydronaphthalene-2-carboxylic acid, methyl ester (XXXa)

30




A mixture of a compound of formula XVI (Scheme XXI) in
 5 which R⁷ is methyl (2.00 g, 8.60 mmol),
 chlorotrimethylsilane (TMS-Cl, 1.12 g, 10.3 mmol) and
 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU, 1.70 g, 11.2
 mmol) in 10 mL of anhydrous methylene chloride was
 stirred at reflux for 1 hour. The mixture was diluted
 10 with 75 mL of diethyl ether, washed with 20 mL of 0.1
 N of HCl, 20 mL of water, and dried over magnesium
 sulfate. The solvent was evaporated and the residue
 was dried in vacuum. To the residue was added 30 mL
 of acetonitrile and palladium acetate (2.12 g, 9.50
 15 mmol). After stirring for 16 hours, additional
 palladium acetate (1.06 g, 4.75 mmol) was added, and
 the resulting mixture was stirred for 16 hours. The
 mixture was filtered through a pad of celite, and
 washed with acetonitrile. The filtrate collected was
 20 evaporated and the residue was diluted with water (30
 mL) and extracted with ether (30 mL x 3). The
 combined extracts were dried over magnesium sulfate,
 filtered, and evaporated. The residue was purified by
 flash chromatography on silica gel (eluted with EtOAc
 25 : hexane, 1:20 to 1: 5) to give 1.42 g (64% yield) of
 the title compound as a solid; ¹H-NMR (CDCl₃) δ 1.50
 (s, 6 H), 3.94 (s, 3 H), 6.41 (d, J = 10.2 Hz, 1 H),
 6.94 (d, J = 10.2 Hz, 1 H), 7.62 (d, J = 8.2 Hz, 1 H),
 8.22 (dd, J = 2.0, 8.2 Hz, 1 H), 8.82 (d, J = 2.0 Hz,
 30 1 H); MS (DCI) m/e: 321 (MH⁺).

EXAMPLE 51

5,5,6-Trimethyl-8-oxo-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid, methyl ester (XXXIa)

5




To a suspension of cuprous cyanide (724 mg, 8.08 mmol) in 40 mL of anhydrous ether was slowly added methyl
10 lithium (1.4 M in ether, 11.5 mL, 16.2 mmol) at -50 °C. After stirring for 1 h, the solution was cooled to -78 °C and boron trifluoride etherate (573 mg, 4.04 mmol) was added followed by slow addition of enone ester
15 XXXa (930 mg, 4.04 mmol) in 10 mL of ether. After stirring for 45 min, the reaction was quenched with saturated ammonium chloride solution (30 mL). The mixture was extracted with ether (30 mL x 3). The combined ether extracts were dried over magnesium sulfate, filtered, and evaporated. The residue was
20 purified by flash chromatography on silica gel (eluted with hexane : EtOAc, 20:1 to 5:1) to give 251 mg (25% yield) of the title compound as a solid; ¹H-NMR (CDCl₃) δ 1.05 (d, J = 6.9 Hz, 3 H), 1.30, 1.44 (s, 3 H each), 2.15-2.30 (m, 1H), 2.55 (dd, J = 9.3, 17.5 Hz, 1 H),
25 2.81 (dd, J = 4.5, 17.5 Hz, 1 H), 3.93 (s, 3 H), 7.55 (d, J = 8.3 Hz, 1 H), 7.69 (dd, J = 1.9, 8.3 Hz, 1 H), 8.66 (d, J = 1.9 Hz, 1 H); MS (DCI) m/e: 247 (MH⁺).

Anal. Calcd. for C₁₅H₁₈O₃ : C, 73.15; H, 7.37. Found: C
30 72.78; H, 7.31.

EXAMPLE 52

5,5,6-Trimethyl-8-phenyl-5,6-dihydronaphthalene-2-
carboxylic acid, methyl ester (XXXIIa)

5




- To a solution of compound XXXIa (390 mg, 1.58 mmol) in 10 mL of anhydrous tetrahydrofuran was added
- 10 phenylmagnesium bromide (3.0 M in ether, 1.05 mL, 3.15 mmol) at -30 °C. After stirring for 10 min, the solution was raised to 0 °C and stirred for 20 min. The reaction was quenched with 20 mL of saturated ammonium chloride and extracted with ethyl acetate (25
- 15 mL x 3). The combined extracts were washed with water (20 mL), dried over magnesium sulfate, filtered and evaporated. To the residue was added 20 mL of anhydrous benzene and p-toluensulfonic acid (50 mg). The solution was stirred at reflux for 30 min and then
- 20 evaporated. The residue was purified by flash chromatography on silica gel (eluted with EtOAc : hexane, 1:20 to 1: 10) to give 380 mg (78% yield) of the title product as a light yellow oil; ¹H-NMR (CDCl₃)
- 25 δ 1.01 (d, J = 7.1 Hz, 3 H), 1.29, 1.36 (s, 3 H each), 2.35-2.43 (m, 1 H), 3.82 (s, 3 H), 5.94 (d, J = 5.0 Hz, 1 H), 7.33-7.42 (m, 5 H), 7.43 (d, J = 8.1 Hz, 1 H), 7.69 (d, J = 1.8 Hz, 1 H), 7.90 (dd, J = 1.8, 8.1 Hz, 1 H); MS (DCI) m/e: 307 (MH⁺).
- 30 Anal. Calcd. for C₂₁H₂₂O₂ : C, 82.32; H, 7.24. Found: C 82.10; H, 7.32.

EXAMPLE 53

5,5,6-Trimethyl-8-phenyl-5,6-dihydronaphthalene-2-carboxylic acid (XVIIIe)

5




A solution of compound XXXIIa (224 mg, 0.73 mmol) and 2N NaOH (3.7 ml, 7.4 mmol) in 7 mL of tetrahydrofuran and 7 mL of methanol was stirred at 60 °C for 1.5 hours. The mixture was concentrated under reduced pressure and acidified with 10 mL of 1 N HCl, extracted with EtOAc (20 mL x 3). The combined extracts were washed with water (10 mL), dried over magnesium sulfate, filtered, and evaporated. The residue was crystallized from ether-hexane to give 220 mg of the product (quantative yield) as a solid; ¹H-NMR (CDCl₃) δ 1.02 (d, J = 7.1 Hz, 3 H), 1.31, 1.37 (s, 3 H each), 2.35-2.45 (m, 1 H), 5.95 (d, J = 5.1 Hz, 1 H), 7.30-7.40 (m, 5 H), 7.46 (d, J = 8.1 Hz, 1 H), 7.73 (d, J = 1.9 Hz, 1 H), 7.94 (dd, J = 1.9, 8.1 Hz, 1 H); MS (DCI) m/e: 293 (MH⁺).

Anal. Calcd. for C₂₀H₂₀O₂·0.125 H₂O: C, 81.53; H, 6.93.
Found: C 81.68; H, 6.86.

EXAMPLE 54

4-[[[(5,6-Dihydro-5,5,6-trimethyl-8-phenyl-2-naphthalenyl)carbonyl]amino]benzoic acid, methyl ester (I³d)




To a solution of compound XVIIIe (206 mg, 0.71 mmol) in 4 mL of methylene chloride was added oxalyl chloride (179 mg, 1.41 mmol) and 1 drop of dimethylformamide at 0 °C. The solution was stirred at 0 °C for 2 hours and at room temperature for 1 hour. The solvent was evaporated and the residue was dried in vacuum. To the residue was added methyl p-aminobenzoate (128 mg, 0.85 mmol) and 4 mL of anhydrous pyridine. The mixture was stirred at room temperature for 18 hours. The solvent was evaporated, and the residue was diluted with 1 N HCl (20 mL) and extracted with EtOAc (20 mL X 3). The combined extracts were dried over magnesium sulfate, filtered, and evaporated. The residue was purified by flash chromatography on silica gel (eluted with hexane : EtOAc, 10:1 to 5:1) to give 234 mg (78% yield) of the title compound as a foam; ¹H-NMR (CDCl₃) δ 1.03 (d, J = 7.1 Hz, 3 H), 1.30, 1.37 (s, 3 H each), 2.35-2.45 (m, 1 H), 3.90 (s, 3 H), 5.98 (d, J = 5.0 Hz, 1 H), 7.30-7.41 (m, 5 H), 7.46 (d, J = 8.1 Hz, 1 H), 7.51 (d, J = 1.8 Hz, 1 H), 7.65 (d, J = 8.7 Hz, 1 H), 7.72 (dd, J = 1.8, 8.1 Hz, 1 H), 7.97 and 8.00 (s over d, J = 8.7 Hz, 3 H); MS (DCI) m/e: 426 (MH⁺)

Anal. Calcd. for C₂₈H₂₇NO₃·0.5 H₂O : C, 77.39, H, 6.50; N, 3.22. Found: C, 77.34; H, 6.19; N, 3.33.

EXAMPLE 55

4-[[[(5,6-Dihydro-5,5,6-trimethyl-8-phenyl-2-naphthalenyl)carbonyl]amino]benzoic acid (I⁴d)

5



A mixture of compound I³d (224 mg, 0.53 ml) and 2 N NaOH (2.65 mL, 5.30 mmol) in 5 mL of tetrahydrofuran and 5 mL of methanol was stirred at room temperature for 16 hours. The resulting solution was acidified with 1 N HCl and concentrated under reduced pressure. The residue was diluted with 20 mL of water and filtered. The collected solids were triturated in ether to give 127 mg (59% yield) of the title product as light yellow powder; ¹H-NMR (DMSO-d₆) δ 0.97 (d, J = 7.1 Hz, 3 H), 1.26, 1.31 (s, 3 H each), 2.37-2.41 (m, 1 H), 5.98 (d, J = 5.0 Hz, 1 H), 7.31-7.45 (m, 6 H), 7.54 (d, J = 8.1 Hz, 1 H), 7.79-7.89 (m, 6 H), 10.43 (s, 1 H), 12.72 (s, 1 H); MS (DCI) m/e: 411 (MH⁺).


Anal. Calcd. for C₂₇H₂₅NO₃·0.5 H₂O: C, 77.12, H, 6.23; N, 3.33. Found: C, 77.41; H, 6.05; N, 3.26.

25

EXAMPLE 56

5,5,7-Trimethyl-8-oxo-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid, methyl ester (XXXVa)

30




- To a solution of a compound of formula XVI (Scheme XXII) in which R⁷ is methyl (0.60 g, 2.58 mmol) in 10 mL of anhydrous tetrahydrofuran and 1 mL of hexamethylphosphoramide (HMPA) was added lithium bis(trimethylsilyl)amide (LHMDS, 1 M in hexane, 2.71 mL, 2.71 mmol) at -78 °C under a nitrogen atmosphere. After stirring for 2 hours, the solution was warmed to
- 10 -40 °C and methyl iodide (0.73 g, 5.16 mmol) was added. The reaction was stirred for 2.5 hours, and quenched with saturated ammonium chloride solution (20 mL). The mixture was extracted with EtOAc (20 mL x 3) and the combined extracts were dried over magnesium
- 15 sulfate, and evaporated. The residue was purified by flash chromatography on silica gel (eluted with EtOAc : hexane, 1:30 to 1:10) to give 516 mg (81% yield) of the title product as a white solid; ¹H-NMR (CDCl₃) δ
- 20 1.27 (d, J = 6.6 Hz, 3 H), 1.40, 1.45 (s, 3 H each), 1.91 (d, J = 2.0 Hz, 1 H), 1.94 (s, 1H), 2.78-2.91 (m, 1 H); 3.93 (s, 3 H), 7.50 (d, J = 8.2 Hz, 1 H), 8.16 (dd, J = 1.9; 8.2 Hz, 1 H), 8.64 (d, J = 1.9 Hz, 1 H); MS (DCI) m/e: 247.
- 25 Anal. Calcd. for C₁₅H₁₈O₃: C, 73.15, H, 7.37. Found: C, 73.15; H, 7.51.

EXAMPLE 57

5,5,7-Trimethyl-8-phenyl-5,6-dihydronaphthalene-2-
carboxylic acid, methyl ester (XXXVIa)

5




To a solution of ketone XXXVa (500 mg, 2.03 mmol) in 20 mL of tetrahydrofuran was slowly added phenyl
 10 magnesium bromide (3.0 M in ether, 1.35 mL, 4.06 mmol) at 0 °C. The red mixture was stirred at 0 °C for 1 hour, and quenched with saturated ammonium chloride solution (20 mL). The mixture was extracted with EtOAc (25 mL x 3). The combined extracts were dried over
 15 magnesium sulfate, and evaporated. To the residue was added benzene (10 mL) and p-toluenesulfonic acid (0.1 g), and stirred at reflux for 30 min. The solvent was evaporated and the residue was purified by flash chromatography on silica gel (eluted with EtOAc :
 20 hexane, 1:30 to 1:10) to give 380 mg (61% yield) of the title product; ¹H-NMR (CDCl₃) δ 1.35 (s, 6 H), 1.73 (s, 3 H), 2.31 (s, 2H), 3.77 (s, 3H), 7.17 (d, J = 6.8 Hz, 2 H), 7.29-7.45 (m, 5 H), 7.80 (dd, J = 1.8, 8.0 Hz, 1 H); MS (DCI) m/e: 307.

25

EXAMPLE 58

5,5,7-Trimethyl-8-phenyl-5,6-dihydronaphthalene-2-
carboxylic acid (XVIIIIf)

30



A solution of ester XXXVla (180 mg, 0.62 mmol) and 1 N NaOH (1.22 mL, 2 mmol) in 5 mL of tetrahydrofuran and 5 2 mL of MeOH was stirred at 60 °C for 3 hours. The solution was concentrated to approximate 3 mL under reduced pressure and diluted with 1 N HCl (10 mL), extracted with EtOAc (30 mL x 2). The combined 10 extracts were washed with water (10 mL), dried over magnesium sulfate, and evaporated to give 150 mg of the title acid (83% yield) as a solid;


¹H-NMR (CDCl₃) δ 1.35 (s, 6 H), 1.73 (s, 3 H), 2.32 (s 2H), 7.15 (d, J = 6.7 Hz, 2 H), 7.32 (d, J = 1.6 Hz, 1 H); 7.35-7.45 (m, 4 H), 7.86 (dd, J = 1.6, 8.0 Hz, 15 1 H); MS (DCI) m/e: 293 (MH⁺).

Anal. Calcd. for C₂₀H₂₀O₂·0.25 H₂O C, 80.91, H, 6.96.
Found: C, 81.01; H, 6.82.

20

EXAMPLE 59

4-(((5,6-Dihydro-5,5,7-trimethyl-8-phenyl-2-naphthyl)carboxyl)amino)benzoic acid, methyl ester
25 (I^{3e})




To a solution of acid XVIIIf (171 mg, 0.59 mmol) in 4 mL of methylene chloride was added oxalyl chloride (150 mg, 1.18 mmol) and 1 drop of dimethylformamide at 0 °C. The solution was stirred at 0 °C for 1 hour and at room temperature for 16 hours. The solvent was evaporated and the residue was dried in vacuum. To the residue was added methyl p-aminobenzoate (98 mg, 0.65 mmol) and 3 mL of anhydrous pyridine. The mixture was stirred at room temperature for 18 hours. The solvent was evaporated, and the residue was diluted with 1 N HCl (20 mL) and extracted with EtOAc (20 mL X 3). The combined extracts were dried over magnesium sulfate, filtered, and evaporated. The residue was purified by flash chromatography on silica gel (eluted with hexane : EtOAc, 20:1 to 5:1) to give 149 mg (78% yield) of the title compound as an oil; ¹H-NMR (CDCl₃) δ 1.37 (s, 6 H), 1.75 (s, 3 H), 2.34 (s, 2H), 3.90 (s, 3 H), 7.09 (d, J = 1.8 Hz, 1 H), 7.18 (bd, J = 6.8 Hz, 2 H); 7.33-7.46 (m, 4 H), 7.60 (d, J = 8.8 Hz, 2 H), 7.65 and 7.67 (d over bs, J = 1.9 Hz, 1 H), 8.00 (d, J = 8.7 Hz, 2 H); MS (DCI) m/e: 426 (MH⁺).

25

EXAMPLE 60

4-[[[5,6-Dihydro-5,5,7-trimethyl-8-phenyl-2-naphthanenyl]carbonyl]aminolbenzoic acid (I'e)




30

A mixture of compound I³e (97 mg, 0.23 ml) and 2 N NaOH (1.15 mL, 2.30 mmol) in 5 mL of tetrahydrofuran and 5 mL of MeOH was stirred at room temperature for 12 hours. The resulting solution was acidified with 1 N HCl and concentrated under reduced pressure. The residue was diluted with 10 mL of water and extracted with EtOAc (15 mL X2). The combined extracts were dried over magnesium sulfate, filtered, and evaporated. The residue was triturated with Et₂O-hexane to give 77 mg (82% yield) of the product as white solids; ¹H-NMR (DMSO-d₆) δ 1.33 (s, 6 H), 1.69 (s, 3 H), 2.31 (s, 2H), 7.02 (s, 1 H), 7.15 (d, J = 7.9 Hz, 2 H), 7.33-7.49 (m, 4 H), 7.75 (d, J = 8.4 Hz, 1 H), 7.79 (d, J = 8.6 Hz, 2 H), 7.87 (d, J = 8.6 Hz, 2 H); MS (DCI) m/e: 412 (MH⁺).

Anal. Calcd. for C₂₇H₂₅NO₃·0.25 H₂O: C, 77.95, H, 6.18; N, 3.37. Found: C, 77.93; H, 6.01; N, 3.27.

EXAMPLE 61

10,10-Dimethyl-9-oxo-5,8,8a,9,10,10a-hexahydro-anthracene-2-carboxylic acid, methyl ester (XXXVIIa)




A solution of enone XXXa (500 mg, 2.17 mmol) and aluminum chloride (260 mg, 1.95 mmol) in 10 mL of anhydrous toluene was saturated with 1,3-butadiene at -78 °C. After stirring at room temperature for 18 hours, the mixture was diluted with 20 mL of 1 N HCl

- and extracted with EtOAc (30 mL X3). The combined extracts were washed with water (20 mL), saturated sodium chloride and dried over magnesium sulfate. The solvent was evaporated and the residue was purified by
- 5 flash chromatography on silica gel (eluted with EtOAc:hexane, 1:20 to 1:5) to give 638 mg (quantative yield) of the title product as an oil; $^1\text{H-NMR}$ (CDCl_3) δ 1.45, 1.49 (s, 3 H each), 1.60-1.75 (m, 1 H), 2.10-2.30 (m, 3 H), 2.99-3.06 (bd, $J = 18$ Hz, 1 H), 3.29
- 10 (bs, 1 H), 3.91 (s, 3 H), 5.53-5.58 (m, 1 H), 5.70-5.76 (m, 1 H), 7.46 (d, $J = 8.2$ Hz, 1 H), 8.16 (dd, $J = 1.8, 8.2$ Hz, 1 H), 8.64 (d, $J = 1.8$ Hz, 1 H); MS (DCI) m/e : 285 (MH^+).
- 15 Anal. Calcd. for $\text{C}_{18}\text{H}_{20}\text{O}_3$: C, 76.03, H, 7.09. Found: C, 75.73; H, 7.25.

EXAMPLE 62

20

10,10-Dimethyl-9-phenyl-5,8,10,10a-tetrahydro-anthracene-2-carboxylic acid, methyl ester (XXXVIIIa)




25

- To a solution of compound XXXVIIa (600 mg, 2.11 mmol) in 5 mL of anhydrous tetrahydrofuran was slowly added phenylmagnesium bromide (3.0 M in ether, 1.76 mL, 5.28 mmol) at -30 $^{\circ}\text{C}$. After stirring for 15 min, the
- 30 mixture was slowly warmed to 0 $^{\circ}\text{C}$ in 20 min. The reaction was quenched with saturated ammonium chloride solution (30 mL) and extracted with EtOAc (30 mL X 3).

The combined extracts were dried over magnesium sulfate, filtered, and evaporated. The residue was partially purified by flash chromatography on silica gel (eluted with EtOAc:hexane, 1:20 to 1:5) to give 520 mg of a crude alcohol. To the alcohol was added anhydrous benzene (10 mL) and p-toluenesulfonic acid (50 mg). The resulting solution was stirred at reflux for 3 hours. The solvent was evaporated and the residue was purified by flash chromatography on silica gel (eluted with EtOAc:hexane, 1:20 to 1:5) to give 265 mg (36% yield) of the title compound as an oil; $^1\text{H-NMR}$ (CDCl_3) δ 1.34, 1.44 (s, 3 H each), 1.92-2.05 (m, 1 H), 2.25-2.45 (m, 2 H), 2.65-2.80 (m, 2 H), 3.78 (s, 3 H), 5.55-5.78 (m, 2 H), 7.16 (d, $J = 6.8$ Hz, 2 H), 7.31 (d, $J = 1.7$ Hz, 1 H), 7.30-7.50 (m, 6 H), 7.82 (dd, $J = 1.7, 8.0$ Hz, 1 H).

EXAMPLE 63

10,10-Dimethyl-9-phenyl-5,8,10,10a-tetrahydroanthracene-2-carboxylic acid (XVIIIg)




A solution of compound XXXVIIIa (260 mg, 0.75 mmol) and 3 N NaOH (2.5 mL, 7.5 mmol) in 10 mL of methanol was stirred at 80 °C for 3 hours. The solvent was evaporated, and the residue was acidified with 1 N HCl, extracted with EtOAc (30 mL X3). The combined extracts were washed with water (20 mL), dried over magnesium sulfate, filtered and evaporated to give 235

mg (95% yield) of the title compound; $^1\text{H-NMR}$ (CDCl_3) δ 1.34, 1.44 (s, 3 H each), 1.93-2.05 (m, 1 H), 2.25-2.37 (m, 1 H), 2.41 (dd, $J = 5.0, 11.9$ Hz, 1 H), 2.65-2.80 (m, 2 H), 5.60-5.70 (m, 2 H), 7.14 (d, $J = 6.7$ 5 Hz, 2 H), 7.32 (d, $J = 1.8$ Hz, 1 H), 7.36-7.46 (m, 4 H), 7.85 (dd, $J = 1.8, 8.0$ Hz, 1 H); MS (DCI) m/e : 331 (MH^+).

Anal. Calcd. for $\text{C}_{23}\text{H}_{22}\text{O}_2$: C, 83.61, H, 6.71. Found: C, 10 83.25; H, 6.86.

EXAMPLE 64

15 4-[[[(5,8,10,10a-Tetrahydro-10,10-dimethyl-9-phenyl-2-anthracenyl)carbonyl]aminol]benzoic acid, methyl ester (I^3f)




20 To the solution of acid XVIIIg (225 mg, 0.68 mmol) in 4 mL of methylene chloride was added oxalyl chloride (172 mg, 1.36 mmol) and 1 drop of dimethylformamide at 0 °C. The solution was stirred at 0 °C for 2 hour and 25 at room temperature for 1 hour. The solvent was evaporated and the residue was dried in vaccum. To the residue was added methyl p-aminobenzoate (122 mg, 0.81 mmol) and 2 mL of anhydrous pyridine. The mixture was stirred at room temperature for 16 hours. The 30 solvent was evaporated, and the residue was diluted with 1 N HCl (20 mL) and extracted with EtOAc (30 mL X

3). The combined extracts were dried over maganesium sulfate, filtered, and evaporated. The residue was purified by flash chromatography on silica gel (eluted with hexane : EtOAc, 20:1 to 5:1) to give 210 mg (67%
 5 yield) of the title compound; $^1\text{H-NMR}$ (CDCl_3) δ 1.36, 1.45 (s, 3 H each), 1.95-2.05 (m, 1 H), 2.33 (dt, J = 4.6, 16.1 Hz, 1 H), 2.44 (dd, J = 5.1, 12.0 Hz, 1 H), 2.65-2.82 (m, 2 H), 3.89(s, 3 H), 5.60-5.73 (m, 2 H), 7.09 (d, J = 1.9 Hz, 1 H), 7.18 (bd, J = 6.7 Hz, 2 H),
 10 7.33-7.47 (m, 4 H), 7.59-7.65 (m, 3H), 7.70 (bs, 1 H), 8.00 (d, J = 8.8 Hz, 2 H); MS (DCI) m/e : 464 (MH^+).

EXAMPLE 65

15

4-[[[5,8,10,10a-Tetrahydro-10,10-dimethyl-9-phenyl-2-anthracenyl]carbonyl]aminolbenzoic acid (I^4f)



20

A solution of ester I^3f (210 mg, 0.45 mmol) and dibutyltin oxide(228 mg, 0.92 mmol) in 5 mL of anhydrous toluene was stirred at reflux for 2 days. The mixture was added 1 N HCl (20 mL) and extracted
 25 with EtOAc (20 mL X3). The combined extracts were washed with water (10 mL), and concentrated to approximate 20 mL. The solution was then stirred with 10 mL of 20% potassium fluoride solution for 30 min. The mixture was extracted with EtOAc (20 mL X2). The
 30 combined extracts were dried over magnesium sulfate, filtered, and evaporated. The residue was purified by

flash chromatography on silica gel (eluted with MeOH: CH₂Cl₂, 1:100 to 1:5) to give 72 mg (35% yield) of the title product as a yellow solid; ¹H-NMR (DMSO-d₆) δ 1.30, 1.41 (s, 3 H each), 1.85 (bt, J = 12.1 Hz, 1 H),
 5 2.28-2.35 (m, 1H), 2.41 (dd, J = 5.0, 11.5 Hz, 1 H), 2.55 (bd, J = 19.1 Hz, 1 H), 2.75 (bd, J = 19.1 Hz, 1H), 5.60-5.80 (m, 2 H), 7.02 (d, J = 1.9 Hz, 1 H), 7.12 (d, J = 6.9 Hz, 2 H), 7.35-7.48 (m, 4 H), 7.73 (dd, J = 1.9, 8.2 Hz, 1 H), 7.77 (d, J = 8.8 Hz, 2 H),
 10 7.86 (d, J = 8.8 Hz, 2 H), 10.35 (s, 1 H); MS (DCI) m/e: 450 (MH⁺).

Anal. Calcd. for C₃₀H₂₇NO₃·0.75 H₂O: C, 77.81, H, 6.20; N, 3.02. Found: C, 78.09; H, 5.99; N, 3.14.

15


EXAMPLE 66

20 4,4-Dimethyl-7-iodo-1-tetralone (XXXIX)

To a solution of 4,4-dimethyl-7-amino-1-tetralone (XII) (1.82 g, 10.0 mmol) in concentrated hydrochloric acid (4.69 mL) was added ice cold water (3.13 mL). The
 25 reaction mixture was then cooled to 0°C by use of an ice-salt bath. The reaction mixture was then diazotized by the dropwise addition with stirring of a solution of sodium nitrite (0.76 g, 11.0 mmol) in water (3.13 mL) keeping temperature between 0-5°C.
 30 After stirring for 15 minutes, the reaction mixture was added to a solution of potassium iodide (3.63 g, 21.9 mmol) in water (18.8 mL). After standing for 30 minutes, the dark gum was extracted with ethyl acetate (1 x 100 mL). The organic phase was then concentrated
 35 in vacuo and the residue chromatographed on silica gel

(eluted with 5% ethyl acetate in hexane) to give 1.56 g (Y: 54%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 8.33 (d, $J=2.0$ Hz, 1H), 7.82 (dd, $J=8.3$, 2.0 Hz, 1H), 7.17 (d, $J=8.4$ Hz, 1H), 2.72 (t, $J=6.8$ Hz, 2H), 2.01 (t, $J=6.8$ Hz, 2H), 1.37 (s, 6H); MS (DCI) m/e : 301 (MH^+).

EXAMPLE 67


10 Methyl 4-vinylbenzoate

To a solution of 4-vinylbenzoic acid (Aldrich, 2.18 g, 14.7 mmol) in anhydrous acetonitrile (14.0 mL) was added 1.8-diazabicyclo [5.4.0]undec-7-ene (Aldrich, 2.46 g, 16.2 mmol) and iodomethane (Aldrich, 3.13g, 22.1 mmol) at 0°C. The reaction mixture was then warmed to room temperature and allowed to stir for 3 h. Ethyl acetate (100 mL) was added to the mixture and the solution was washed with brine (50 mL). The organic phase was then separated and concentrated in vacuo. The residue was chromatographed on silica gel (eluted with 5% ethyl acetate in hexane) to give 1.05g (Y: 44%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 8.00 (d, $J=8.4$ Hz, 2H), 7.47 (d, $J=8.4$ Hz, 2H), 6.76 (m, 1H), 5.87 (d, $J=17.6$ Hz, 1H), 5.38 (d, $J=11.0$ Hz, 1H), 3.91 (s, 3H); MS (DCI) m/e : 163 (MH^+).

EXAMPLE 68

4-[[[(E)-(5,6,7,8-Tetrahydro-5,5-dimethyl-8-oxo)-2-naphthalenyl]vinyl]benzoic acid, methyl ester (XLa)


5



To a solution of 4,4-dimethyl-7-iodo-1-tetralone
 (~~XXXIX~~) (1.55 g, 5.17 mmol) and methyl 4-vinylbenzoate
 10 (1.67 g, 10.34 mmol) in dimethylformamide (16.0 mL)
 was added palladium (II) acetate (Aldrich, 58 mg,
 0.259 mmol), tetrabutylammonium chloride hydrate
 (Aldrich, 1.49 g, 5.17 mmol) and sodium bicarbonate
 (Mallinckrodt, 1.09 g, 12.9 mmol). The reaction
 15 mixture was heated to 70°C for 4 h and then allowed to
 stir at room temperature for 16 h. Ethyl acetate (50
 mL) was added to the mixture and the solution was
 washed with brine (50 mL). The organic phase was
 concentrated in vacuo and the residue chromatographed
 20 on silica gel (eluted with 10% ethyl acetate in
 hexane) to give 1.29 g (Y: 75%) of the title compound;
¹H-NMR (CDCl₃): δ 8.19 (d, J=1.7 Hz, 1H), 8.04 (d,
 J=8.2 Hz, 2H), 6.69 (dd, J=8.1, 1.7 Hz, 1H), 7.57 (d,
 J=8.2 Hz, 2H), 7.45 (d, J=8.1 Hz, 1H), 7.21 (s, 2H),
 25 3.93 (s, 3H), 2.76 (t, J=6.8 Hz, 2H), 2.05 (t, J=6.8
 Hz, 2H), 1.41 (s, 6H); MS (DCI) m/e: 335 (MH⁺).

EXAMPLE 69


4-[[(E)-(5,6,7,8-Tetrahydro-5,5-dimethyl-8-phenyl-8-hydroxy)-2-naphthalenyl]]vinyl]benzoic acid, methyl ester (XLIa)



Using the method for the preparation of the 8-phenyl derivative XVIIa, 1.29 g, (3.86 mmol) of compound XLa gave 1.39 g (Y: 87%) of the title compound; ¹H-NMR (CDCl₃): δ 7.98 (d, J=8.1 Hz, 2H), 7.48 (m, 5H), 7.31 (s, 5H), 7.03 (m, 2H), 3.91 (s, 3H), 2.22 (m, 2H), 1.85 (m, 1H), 1.60 (m, 1H), 1.40 (s, 3H), 1.35 (s, 3H).

EXAMPLE 70

4-[[(E)-(5,6,-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]]vinyl]benzoic acid (I¹¹a)




To a solution of compound XLIa (1.38 g, 3.35 mmol) in toluene (20 mL) was added p-toluenesulfonic acid (p-

TsOH, 20 mg). After heating at 70°C for 0.5 h, the reaction mixture was cooled and concentrated in vacuo. The residue was then dissolved in a 1:1 ethyl alcohol tetrahydrofuran solution (20.0 mL) and treated with 10
 5 N NaOH (33.7 mmol, 3.37 mL) at room temperature. After 16 hours, an excess of 1N HCl (75 mL) was added and the precipitate collected by vacuum filtration to give 1.23 g (Y: 97%) of the title compound; ¹H-NMR (DMSO-d₆): δ 12.87 (s, 1H), 7.85 (d, J=8.4 Hz, 2H), 7.63 (d, J=8.4 Hz, 2H), 7.58 (dd, J=8.1, 1.7 Hz, 1H), 7.47-7.24
 10 (m, 7H), 7.07 (m, 2H), 6.00 (t, J=4.5 Hz, 1H), 2.30 (d, J=4.5 Hz, 2H), 1.28 (s, 6H); ¹³C-NMR (DMSO-d₆) 167.02, 145.09, 141.35, 140.16, 138.59, 134.22, 133.52, 130.96, 129.66, 129.23, 128.52, 128.33,
 15 127.31, 126.87, 126.65, 126.45, 125.30, 124.69, 124.46, 38.23, 33.29, 27.89; MS (DCI) m/e: 381 (MH⁺); IR (KBr): 2922, 2818, 1684, 1604 cm⁻¹.

Anal. calcd for C₂₇H₂₄O₂·0.65 H₂O: C, 82.69; H, 6.50.
 20 Found: C, 83.00; H, 6.41.

EXAMPLE 71

25 4-[[[(5,6,-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]sulfamyl]benzoic acid (I^{12a})



30 To a solution of compound XVIIIa (119 mg, 0.430 mmol)


in anhydrous methylene chloride (5.0 mL) was added oxalyl chloride (0.13 mL, 1.5 mmol) and 2 drops of N,N-dimethylformamide at 0°C. The reaction mixture was allowed to stir at room temperature for 2 hours. The
 5 reaction mixture was then concentrated in vacuo. The resultant residue was dissolved in anhydrous pyridine (5.0 mL) to which was added 4-mercaptobenzoic acid (Apin, 66 mg, 0.43 mmol). After 3 hours at room temperature, the mixture was diluted with 1 N HCl,
 10 extracted with ethyl acetate (50 mL), washed with 1N HCl (3 x 50 mL) and washed with saturated sodium bicarbonate (2 x 100 mL). The organic phase was then separated, concentrated in vacuo and the residue chromatographed on silica gel (eluted with 5% methanol
 15 in methylene chloride) to give 40 mg (Y: 23%) of the title product; ¹H-NMR (DMSO): δ 7.98 (d, J=8.3 Hz, 2H), 7.91 (dd, J=8.1, 2.0 Hz, 1H), 7.62 (d, J=8.2 Hz, 1H), 7.56 (d, J=8.2 Hz, 2H), 7.48-7.27 (m, 6H), 6.11 (t, J=4.5 Hz, 1H), 2.38 (d, J=4.5 Hz, 2H), 1.33 (s, 6H);
 20 MS (DCI) m/e: 415 (MH⁺); IR (KBr): 3440, 2962, 1684, 1594 cm⁻¹.

Anal. calcd for C₂₆H₂₂O₃S₁·1.2 H₂O: C, 71.60; H, 5.64.
 Found: C, 71.65, H, 5.44.

25

EXAMPLE 72

4-[[[(5,6,7,8-Tetrahydro-5,5-dimethyl-8-oxo)-2-
 30 naphthalenyl]sulfamyl]carbonyl]benzoic acid, t-butyl
ester (XLIVa)




To a solution of mono-tert-butylterephthalate (122 mg, 0.550 mmol) in anhydrous methylene chloride (5.0 mL) 5 was added oxalyl chloride (0.17 mL, 1.95 mmol) and 2 drops of N,N-dimethylformamide at 0°C. After being stirred at room temperature for 2 hours, the mixture was concentrated in vacuo. The residue was dissolved in anhydrous pyridine (5.0 mL) to which was added 4,4- 10 dimethyl-7-mercapto-1-tetralone (XLIII) (113 mg, 0.550 mmol). After 2 hours at room temperature, the mixture was diluted with 1N HCl, extracted with ethyl acetate (50 mL), washed with 1N HCl (3 x 50 mL) and washed with saturated sodium bicarbonate (2 x 50 mL). The 15 organic phase was then separated and concentrated in vacuo. The residue was chromatographed on silica gel (eluted with 10% ethyl acetate in hexane) to give 80 mg (Y: 35%) of the title product; ¹H-NMR (CDCl₃): δ 8.16 (d, J=1.8 Hz, 1H), 8.11-8.03 (m, 4H), 7.67 (dd, J=8.3, 2.1 Hz, 1H), 7.54 (d, J=8.5 Hz, 1H), 2.77 (t, J=6.8 Hz, 2H), 2.06 (t, J=6.8 Hz, 2H), 1.63 (s, 9H), 1.44 (s, 6H); MS (DCI) m/e: 355 (M-C₄H₉ + H)⁺.

EXAMPLE 73

25

4[[[(5,6,7,8-Tetrahydro-5,5-dimethyl-8-phenyl-8-hydroxy)-2-naphthalenyl]sulfamoyl]carbonyl]benzoic acid, t-butyl ester (XLVa)




Using the method for the preparation of the 8-phenyl derivative XVIIa, 80 mg (0.19 mmol) of compound XLIVa
 5 gave 46 mg (Y: 50%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 8.03 (m, 4H), 7.52 (d, $J=8.2$ Hz, 1H), 7.43 (dd, $J=8.2$, 2.1 Hz, 1H), 7.20 (m, 6H), 2.30-2.10 (m, 3H), 1.90 (m, 1H), 1.63 (s, 9H), 1.45 (s, 3H), 1.37 (s, 3H); MS (DCI) m/e : 471 (MH^+).

10

EXAMPLE 74

4-[[[5,6-Dihydro-5,5-dimethyl-8-phenyl]-2-
 15 naphthalenyl]sulfamoyl]benzoic acid (I^{13}a)



20 To a solution of compound XLVa (46 mg, 0.094 mmol) in toluene (5.0 mL) was added a few milligrams (~ 3 mg) of p-toluenesulfonic acid. After being heated at 70°C for 0.5 h, the reaction mixture was cooled and concentrated in vacuo. The residue was then dissolved
 25 in methylene chloride (2.0 mL) to which was added

trifluoroacetic acid (0.08 mL) at room temperature. After 16 hours, the reaction mixture was diluted with ethyl acetate (20 mL) and washed with 1N HCl (20 mL). The organic phase was separated, dried over anhydrous magnesium sulfate and concentrated in vacuo to give 34 mg (Y: 87%) of the title product; $^1\text{H-NMR}$ (DMSO-d_6): δ 8.03 (m, 4H), 7.53 (d, $J=8.1$ Hz, 1H), 7.43-7.30 (m, 6H), 6.96 (d, $J=1.9$ Hz, 1H), 6.07 (t, $J=4.5$ Hz, 1H), 2.36 (d, $J=4.5$ Hz, 2H), 1.18 (s, 6H); MS (DCI) m/e : 415 (MH^+). IR (KBr): 3432, 2962, 1680, 1202.


Anal. calcd. for $\text{C}_{26}\text{H}_{22}\text{O}_3\text{S}_1 \cdot 0.35 \text{ H}_2\text{O}$: C, 74.21; H, 5.44. Found: C, 74.18, H, 5.22.

15

EXAMPLE 75

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]ethyl]benzoic acid, ethyl ester

20



To a stirred solution of compound I¹⁵a (118 mg, 0.290 mmol) in toluene (7.0 mL) was added 5% palladium on calcium carbonate, poisoned with lead (Aldrich, Lindlar catalyst, 15 mg) at 45 psi of H_2 . After 16 h, the reaction mixture was filtered through a pad of celite and the filtrate concentrated in vacuo to give 60 mg (Y: 50%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 7.93 (d, $J=8.1$ Hz, 2H), 7.30 (m, 6H), 7.18 (d, $J=8.1$


30

Hz, 2H), 7.05 (dd, $J=7.9, 1.8$ Hz, 1H), 6.81 (d, $J=1.8$ Hz, 1H), 5.97 (t, $J=4.5$ Hz, 1H), 4.38 (q, $J=7.1$ Hz, 2H), 2.84 (m, 4H), 2.34 (d, $J=4.5$ Hz, 2H), 1.42 (t, $J=7.1$ Hz, 3H), 1.33 (s, 6H); MS (DCI) m/e : 411 (MH^+).

5

EXAMPLE 76

4-[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]]ethyl]benzoic acid ($I^{17}a$)




To a solution of compound 4-[[(5,6-dihydro-5,5-
 15 dimethyl-8-phenyl)-2-naphthalenyl]]ethyl]benzoic acid,
 ethyl ester (60mg, 0.146 mmol) in ethyl alcohol (5.0
 mL) was added 10 N NaOH (2.0 mmol, 0.20 mL). After 1h
 at 70°C an excess of 1N HCl (20 mL) was added and the
 precipitate collected by vacuum filtration to give 59
 20 mg (Y: 99%) of the title product; 1H -NMR ($DMSO-d_6$); δ
 7.79 (d, $J=8.2$ Hz, 2H), 7.38-7.29 (m, 3H), 7.25 (d,
 $J=8.2$ Hz, 2H), 7.21-7.10 (m, 4H), 6.62 (d, $J=1.7$ Hz,
 1H), 5.93 (t, $J=4.5$ Hz, 1H), 2.78 (m, 4H), 2.25 (d,
 $J=4.5$ Hz, 2H), 1.24 (s, 6H). MS (DCI) m/e : 383 (MH^+);
 25 IR (KBr): 2956, 1688, 1610, 1422 cm^{-1} .

Anal. calcd. $C_{27}H_{26}O_2$: C, 84.78; H, 6.85. Found: C,
 84.53; H, 6.81.

30

EXAMPLE 77

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]thiocarbonyl]aminolbenzoic acid, methyl ester




To a solution of compound I³a (285 mg, 0.693 mmol) in tetrahydrofuran (6.0 mL) was added phosphorus pentasulfide (Aldrich, 205 mg, 0.461 mmol). After 0.75 h at reflux, the mixture was concentrated down, diluted with methylene chloride (50 mL) and washed with 5% sodium carbonate (2 x 50 mL). The organic phase was then concentrated in vacuo and the residue chromatographed on silica gel (eluted with 10% ethyl acetate in hexane) to give 88 mg (Y: 30%) of the title product; ¹H-NMR (CDCl₃): δ 8.92 (s, 1H), 8.05 (d, J=8.6 Hz, 2H), 7.74 (dd, J=8.2, 1.8 Hz, 1H), 7.43 (d, J=8.1 Hz, 1H), 7.33 (m, 8H), 6.07 (t, J=4.5 Hz, 1H), 3.92 (s, 3H), 2.38 (d, J=4.5 Hz, 2H), 1.36 (s, 6H); MS (DCI) m/e: 428 (MH⁺).

25

EXAMPLE 78

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]thiocarbonyl]aminolbenzoic acid (I¹⁸a)




Using the method given for the preparation of the 8-phenyl derivative I⁴a, 80 mg (0.186 mmol) of 4-[[[(5,6-
 5 dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]thiocarbonyl]amino]benzoic acid, methyl
 ester gave 69 mg (Y: 90%) of the title product; ¹H-NMR
 (DMSO-d₆): δ 12.88 (s, 1H), 11.89 (s, 1H), 7.87 (m,
 4H), 7.60 (dd, J=8.0, 1.7 Hz, 1H), 7.47 (d, J=8.1 Hz,
 10 1H), 7.35 (m, 6H), 6.07 (t, J=4.5 Hz, 1H), 2.35 (d,
 J=4.5 Hz, 2H), 1.32 (s, 6H); MS (DCI) m/e: 414 (MH⁺);
 IR (KBr): 2922, 1688, 1606, 1514 cm⁻¹.

Anal. calcd. for C₂₆H₂₃O₂N₁S₁·0.5 H₂O: C, 73.91; H, 5.73;
 15 N, 3.31. Found: C, 73.60; H, 5.76; N, 3.21.

EXAMPLE 79

5,6-Dihydro-5,5-dimethyl-2-[1-(trimethylsilyl)oxy]ethenyl-8-phenylnaphthalene (LVia)

5




To a solution of compound XIXa of EXAMPLE 32 (2.73 g, 9.89 mmol) in anhydrous methylene chloride (10.0 mL) was added 1,8-diazabicyclo [5.4.0]undec-7-ene (Aldrich, 1.80 g, 11.87 mmol) and chlorotrimethylsilane (1.18 g, 10.9 mmol). The reaction mixture is allowed to reflux gently for 2h and stirred at room temperature for 16 h. Pentane (100 mL) is added to the mixture and the solution is washed with 0.1 N HCl (50 mL) and dilute sodium bicarbonate (50 mL). The organic phase is then dried over anhydrous magnesium sulfate and concentrated in vacuo to give 3.17 g (Y: 92%) of the title product; ¹H-NMR (CDCl₃): δ 7.45 (dd, J=8.0, 2.0 Hz, 1H), 7.39-7.24 (m, 7H), 5.99 (t, J=4.5 Hz, 1H), 4.74 (d, J=1.6 Hz, 1H), 4.30 (d, J=1.6 Hz, 1H), 2.35 (d, J=4.5 Hz, 2H), 1.34 (s, 6H), 0.14 (s, 9H).

25

EXAMPLE 80

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)carbonyl]methyl]benzoic acid, methyl ester (I^{22a})


30



To a solution of compound LVia (864 mg, 2.48 mmol) and methyl 4-bromobenzoate (Aldrich, 355 mg, 1.65 mmol) in 5 anhydrous benzene (10.0 mL) was added tributyltin fluoride (Aldrich, 806 mg, 2.61 mmol) and $\text{PdCl}_2(\text{P}(o\text{-CH}_3\text{C}_6\text{H}_4)_3)_2$ [(59 mg, 0.074 mmol) made from bis(acetonitrile)palladium (II) chloride (Aldrich, 0.074 mmol, 19.3 mg) and tri-*o*-tolylphosphine 10 (Aldrich, 0.149 mmol, 45.3 mg)]. After 4 h at reflux, the reaction mixture was diluted with ethyl acetate (50 mL) and washed with 1N NaOH (50 mL). The organic phase is then chromatographed on silica gel (eluted with 5% ethyl acetate in hexane) to give 265 mg (Y: 15 39%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.92 (d, $J=8.5$ Hz, 2H), 7.87 (dd, $J=8.2$, 2.0 Hz, 1H), 7.63 (d, $J=2.0$ Hz, 1H), 7.44 (d, $J=8.2$ Hz, 1H), 7.41 (m, 3H), 7.29 (m, 2H), 7.17 (d, $J=8.0$ Hz, 2H), 6.03 (t, $J=4.5$ Hz, 1H), 4.12 (s, 2H), 3.90 (s, 3H), 2.37 (d, $J=4.5$ 20 Hz, 2H), 1.35 (s, 6H); MS (DCI) m/e : 411 (MH^+).

EXAMPLE 81

25 4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]methyl]benzoic acid (I^{23}a)




To a solution of compound I²²a (200 mg, 0.488 mmol) in anhydrous toluene was added dibutyltin oxide (Aldrich, 364 mg, 1.46 mmol). After 16 h at reflux, the reaction mixture was concentrated in vacuo and the residue chromatographed on silica gel (eluted with 5% methanol in methylene chloride) to give 70 mg of impure product (tin by-products present). A second purification by preparative thin layer chromatography afforded 10 mg (Y: 5%) of the title product; ¹H-NMR (DMSO-d₆): δ 7.99 (dd, J=7.9, 1.9 Hz, 1H), 7.82 (d, J=8.0 Hz, 2H), 7.54 (d, J=8.0 Hz, 1H), 7.49 (s, 1H), 7.42 (m, 3H), 6.05 (t, J=4.5 Hz, 1H), 4.30 (s, 1H), 4.28 (s, 1H), 2.34 (d, J=4.5 Hz, 2H), 1.31 (s, 6H); MS (DCI) m/e: 397 (MH⁺); High res. M.S. Calcd.: 397.1804. Found: 397.1816, 3.0 ppm deviation.

20

EXAMPLE 82

5,6-Dihydro-5,5-dimethyl-2-hydroxymethyl-8-phenylnaphthalene (LVIIa)



25


A solution of 95% lithium aluminum hydride powder

(LAH, 40 mg, 1.0 mmol) in anhydrous diethyl ether (5.0 mL) was allowed to reflux until most of the hydride dissolved. A solution of compound XVIIIa (485 mg, 1.66 mmol) in anhydrous diethyl ether (5.0 mL) was then
 5 slowly added. After 0.5 h, ethyl acetate (20 mL) and 1 N hydrochloric acid (20 mL) were added. The organic phase was separated, dried over anhydrous magnesium sulfate and concentrated in vacuo to give 368 mg (Y: 84%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 7.36 (m, 6H), 7.26 (m, 1H), 7.02 (d, $J=1.8$ Hz, 1H), 5.99 (t, $J=4.5$ Hz, 1H), 4.55 (s, 2H), 2.05 (d, $J=4.5$ Hz, 2H), 1.34 (s, 6H); MS (DCI) m/e : 247 (M^+-OH).

15

EXAMPLE 83

5,6-Dihydro-5,5-dimethyl-2-bromomethyl-8-phenylnaphthalene (LVIIIa)



20

To a solution of carbon tetrabromide (Aldrich, 1.36g, 4.09 mmol) and triphenylphosphine (Aldrich, 1.07g, 4.09 mmol) in anhydrous diethyl ether (20 mL) was
 25 added compound LVIIa (900 mg, 3.41 mmol). After 16 h at room temperature, the reaction mixture was filtered and the filtrate was concentrated in vacuo. The residue was chromatographed on silica gel (eluted with 3% ethyl acetate in hexane) to give 790 mg (Y: 71%) of
 30 the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.39 (m, 6H), 7.27 (dd, $J=8.1, 1.8$ Hz, 1H), 7.04 (d, $J=1.8$ Hz, 1H), 6.01


(t, J=4.5 Hz, 1H), 4.39 (s, 2H), 2.35 (d, J=4.5 Hz, 2H), 1.34 (s, 6H); MS (DCI) m/e: 327 (MH⁺).

5

EXAMPLE 84

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]methyloxy]benzoic acid, ethyl ester
(I^{25a})

10




To a solution of ethyl 4-hydroxybenzoate (Aldrich, 152 mg, 0.917 mmol) in ethylene glycol dimethyl ether (6.0 mL) was added 60% sodium hydride (1.05 mmol, 43 mg) at room temperature. After 0.15 h, compound LVIIIa (330 mg, 1.01 mmol) was added. After 16 h the reaction mixture was diluted with ethyl acetate (50 mL) and washed with 1N hydrochloric acid (2 x 50 mL). The organic phase was concentrated in vacuo and the residue chromatographed on silica gel (eluted with 10% ethyl acetate in hexane) to give 64 mg (Y: 17%) of the title product; ¹H-NMR (CDCl₃): δ 7.97 (d, J=8.8 Hz, 2H), 7.36 (m, 7H), 7.06 (s, 1H), 6.92 (d, J=8.8 Hz, 2H), 6.01 (t, J=4.5 Hz, 1H), 4.96 (s, 2H), 4.34 (q, J=7.2 Hz, 2H), 2.36 (d, J=4.5 Hz, 2H), 1.40 (t, J=7.2 Hz, 3H), 1.35 (s, 6H); MS (DCI) m/e 413 (MH⁺).

EXAMPLE 85

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]methyloxy]benzoic acid (I²⁶a)

5



Using the method given for the preparation of the 8-phenyl derivative I⁴a, 64 mg (0.155 mmol) of compound
 10 I²⁵a gave 50 mg (Y: 84%) of the title product; ¹H-NMR
 (DMSO-d₆): δ 12.59 (s, 1H), 7.83 (d, J=8.8 Hz, 2H),
 7.33 (m, 7H), 6.97 (d, J=8.9 Hz, 2H), 6.96 (s, 1H),
 5.99 (t, J=4.5 Hz, 1H), 5.04 (s, 2H), 2.30 (d, J=4.5
 Hz, 2H), 1.27 (s, 6H); MS (DCI) m/e: 385 (MH⁺); IR
 15 (KBr): 2958, 1682, 1604, 1252 cm⁻¹.


Anal. calcd. for C₂₆H₂₄O₃·0.25 H₂O: C, 80.28; H, 6.35.
 Found: C, 80.41; H, 6.23.

20

EXAMPLE 86

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]oxy]methyl]benzoic acid (I²⁷a)

25




To a solution of compound XXVa OF EXAMPLE 38 (193 mg,

0.772 mmol) in ethylene glycol dimethyl ether (5.0 mL) was added 60% sodium hydride (36 mg, 0.888 mmol) at room temperature. Methyl 4-(bromomethyl)benzoate (Aldrich, 195 mg, 0.848 mmol) was added after 15 minutes. After 16 h at room temperature, 60% sodium hydride (36 mg, 0.888 mmol) was added again. After 24 h. at reflux, the reaction mixture was diluted with ethyl acetate (50 mL) and washed with 1 N hydrochloric acid (2 x 50 mL). The organic phase was concentrated in vacuo and the residue chromatographed on silica gel (eluted with 5% methanol in methylene chloride) to give 170 mg of impure material. The crude material was dissolved in ethyl alcohol (3 mL) and 1N hydrochloric acid (20 mL) was added. The precipitate was collected by vacuum filtration. 50 mg (Y: 16%) of the title product was isolated; $^1\text{H-NMR}$ (DMSO-d_6): δ 12.95 (s, 1H), 7.89 (d, $J=8.3$ Hz, 2H), 7.42 (d, $J=8.3$ Hz, 2H), 7.39-7.21 (m, 6H), 6.88 (dd, $J=8.5, 2.7$ Hz, 1H), 6.44 (d, $J=2.7$ Hz, 1H), 5.97 (t, $J=4.5$ Hz, 1H), 5.04 (s, 2H), 2.27 (d, $J=4.5$ Hz, 2H), 1.24 (s, 6H); MS (DCI) m/e : 385 (MH^+); IR (KBr): 3442, 1682, 1282 cm^{-1} .

Anal. calcd. for $\text{C}_{26}\text{H}_{24}\text{O}_3 \cdot 0.5 \text{ H}_2\text{O}$: C, 80.59; H, 6.50.
Found: C, 80.31; H, 6.26.

EXAMPLE 87


5,5-Dimethyl-8-hydroxy-8-(2,4-dimethylphenyl)-5,6,7,8-tetrahydro-naphthalene-2-carboxylic acid, methyl ester (XVIIe)



Using the method given for the preparation of the 8-(2-fluorophenyl) derivative XVIId, 311 mg (1.34 mmol) of compound XVIa gave 284 mg (Y: 63%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 7.91 (dd, $J=8.3$, 1.8 Hz, 1H), 7.66 (s, 1H), 7.49 (d, $J=8.3$ Hz, 2H), 7.02 (d, $J=8.0$ Hz, 1H), 6.93 (s, 1H), 3.80 (s, 3H), 2.76 (m, 1H), 2.34 (m, 1H), 2.32 (s, 3H), 2.08 (m, 1H), 1.63 (m, 1H), 1.55 (s, 3H), 1.42 (s, 3H), 1.35 (s, 3H); MS (DCI) m/e : 339 (MH^+).

EXAMPLE 88

5,5-Dimethyl-5,6-dihydro-8-(2,4-dimethylphenyl)-naphthalene-2-carboxylic acid (XVIIIh)




Using the method given for the preparation of 8-phenyl derivative XVIIIe, 311 mg (1.34 mmol) of compound XVIIa gave 285 mg (Y: 63%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.78 (dd, $J=8.0$, 1.7 Hz, 1H), 7.48 (d, $J=8.0$ Hz, 1H), 7.18 (d, $J=1.6$ Hz, 1H), 7.06 (m, 3H), 5.86 (t, $J=4.5$ Hz, 1H), 2.35 (m, 2H), 2.33 (s, 3H), 1.99 (s, 3H), 1.39 (s, 3H), 1.28 (s, 3H); MS (DCI) m/e : 307

(MH⁺).

EXAMPLE 89

5

4-[[[5,6-Dihydro-5,5-dimethyl-8-(2,4-
dimethylphenyl)]-2-naphthalenyl]carbonyl]aminolbenzoic
acid, methyl ester (I^{3h})




10

Using the method given for the preparation of the 8-
 (2-fluorophenyl) derivative I^{3g}, 221 mg (0.722 mmol) of
 15 compound XVIIIh gave 67 mg (Y: 21%) of the title
 product; ¹H-NMR (CDCl₃): δ 8.00 (d, J=8.7 Hz, 2H), 7.77
 (s, 1H), 7.68 (dd, J=8.2, 1.7 Hz, 1H), 7.63 (d, J=8.7
 Hz, 2H), 7.47 (d, J=8.0 Hz, 1H), 7.15 (d, J=1.7 Hz,
 1H), 7.06 (m, 3H), 5.93 (t, J=4.5 Hz, 1H), 3.90 (s,
 20 3H), 2.43 (m, 2H), 2.36 (s, 3H), 2.08 (s, 3H), 1.45
 (s, 3H); MS (DCI) m/e: 440 (MH⁺).

EXAMPLE 90

25

4-[[[5,6-Dihydro-5,5-dimethyl-8-(2,4-
dimethylphenyl)]-2-naphthalenyl]carbonyl]aminolbenzoic
acid (I^{4h})




Using the method given for the preparation of the 8-phenyl derivative I⁴a, 67 mg (0.153 mmol) of compound
 5 I³h gave 61 mg (Y: 94%) of the title product; ¹H-NMR (DMSO-d₆): δ 12.72 (s, 1H), 10.40 (s, 1H), 7.87 (d, J=8.8 Hz, 2H), 7.80 (m, 3H), 7.52 (d, J=8.0 Hz, 1H), 7.04 (m, 4H), 5.88 (t, J=4.5 Hz, 1H), 2.38 (m, 2H), 2.31 (s, 3H), 1.99 (s, 3H), 1.41 (s, 3H), 1.29
 10 (s, 3H); MS (DCI) m/e: 426 (MH⁺); IR (KBr): 2960, 1688, 1594, 1518 cm⁻¹.

Anal. calcd. for C₂₈H₂₇N₁O₃·0.5 H₂O: C, 77.39; H, 6.50; N, 3.22. Found: C, 77.57; H, 6.50; N, 3.22.

15

EXAMPLE 91

5,5-Dimethyl-8-hydroxy-(4-methylphenyl)-5,6,7,8-
 20 tetrahydro-naphthalene-2-carboxylic acid, methyl ester
(XVIIIf)




25 Using the method given for the preparation of the 8-(2-fluorophenyl) derivative XVIIId, 260 mg (1.12 mmol) of compound XVIa gave 189 mg (Y: 52%) of the title

product; $^1\text{H-NMR}$ (CDCl_3): δ 7.92 (dd, $J=8.4$, 1.7 Hz, 1H), 7.84 (d, $J=1.7$ Hz, 1H), 7.47 (d, $J=8.4$ Hz, 1H), 7.10 (s, 4H), 3.82 (s, 3H), 2.34 (s, 3H), 2.18 (m, 2H), 1.80 (m, 1H), 1.60 (m, 1H), 1.39 (s, 3H), 1.34 (s, 3H); MS (DCI) m/e : 325 (MH^+).

EXAMPLE 92

10 5,5-Dimethyl-5,6-dihydro-8-(4-methylphenyl)-naphthalene-2-carboxylic acid (XVIIIii)




15 Using the method for the preparation of 8-phenyl derivative XVIIIa, 189 mg (0.583 mmol) of compound XVIIIf gave 155 mg (Y: 91%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.81 (dd, $J=8.0$, 1.7 Hz, 1H), 7.51 (d, $J=1.7$ Hz, 1H), 7.49 (d, $J=8.0$ Hz, 1H), 7.23 (m, 4H), 6.00 (t, $J=4.5$ Hz, 1H), 2.35 (s, 3H), 2.33 (d, $J=4.5$ Hz, 1H), 1.30 (s, 6H); MS (DCI) m/e : 293 (MH^+).

EXAMPLE 93

25

4-[[[5,6-Dihydro-5,5-dimethyl-8-(4-methylphenyl)]-2-naphthalenyl]carbonyl]aminobenzoic acid, methyl ester (I³ⁱ)




Using the method given for the preparation of the 8-
 (2-fluorophenyl) derivative I³g, 155 mg (0.531 mmol) of
 5 compound XVIIIi gave 143 mg (Y: 63%) of the title
 product; ¹H-NMR (CDCl₃): δ 8.02 (d, J=8.8 Hz, 2H), 7.74
 (s, 1H), 7.72 (dd, J=8.0, 1.8 Hz, 1H), 7.66 (d, J=8.8
 Hz, 2H), 7.53 (d, J=1.8 Hz, 1H), 7.49 (d, J=8.0 Hz,
 1H), 7.23 (m, 4H), 6.06 (t, J=4.5 Hz, 1H), 3.90 (s,
 10 3H), 2.40 (s, 3H), 2.39 (d, J=4.5 Hz, 2H), 1.54 (s,
 3H), 1.37 (s, 3H); MS (DCI) m/e: 426 (MH⁺).

EXAMPLE 94

15

4-[[[5,6-Dihydro-5,5-dimethyl-8-(4-methylphenyl)]-2-
naphthalenyl]carbonyl]aminobenzoic acid (I⁴i)



20

Using the method given for the preparation of the 8-
 phenyl derivative I⁴a, 143 mg (0.336mmol) of compound
 I³i gave 123 mg (Y: 89%) of the title product; ¹H-NMR
 25 (DMSO-d₆): δ 12.72 (s, 1H), 10.42 (s, 1H), 7.85 (m,
 5H), 7.52 (d, J=8.0 Hz, 1H), 7.44 (d, J=1.8 Hz, 1H),
 7.21 (s, 4H), 6.02 (t, J=4.5 Hz, 1H), 2.33 (s, 3H),

2.33 (d, $J=4.5$ Hz, 2H), 1.31 (s, 6H); ^{13}C -NMR (DMSO- d_6): 166.90, 166.06, 148.77, 143.28, 138.16, 136.93, 136.60, 133.49, 132.48, 130.19, 129.12, 128.21, 126.75, 125.35, 125.03, 123.92, 119.38, 119.29, 37.98, 33.52, 27.75, 20.79; MS (DCI) m/e : 412 (MH^+); IR (KBr): 2960, 1688, 1594, 1518 cm^{-1} .

Anal. calcd for $\text{C}_{27}\text{H}_{25}\text{N}_3\text{O}_3 \cdot 0.25 \text{H}_2\text{O}$: C, 77.96; H, 6.18, N, 3.37. Found: C, 77.73; H, 6.09; N, 3.29.

EXAMPLE 95

4,4-Dimethyl-7-bromo-1-tetralone (XLVIII)

To a cooled (10°C) stirred solution of sodium nitrate (2.18 g, 13.92 mmol) in concentrated sulfuric acid (28.4 mL) and glacial acetic acid (26.27 mL) [prepared by adding sodium nitrate to cooled (10°C) concentrated sulfuric acid, heating to dissolve, recooling and then adding glacial acetic acid] was added a solution of 4,4-dimethyl-7-amino-1-tetralone (XII) (5.0 g, 26.46 mmol) in glacial acetic acid (89 mL) over 10 minutes. The resulting solution was added slowly (over 10 minutes) to a heated (60°C) solution of copper (I) bromide (16.66 g) in concentrated hydrobromic acid (159 mL). The mixture was warmed to 90°C for 10 minutes, cooled, diluted with water (250 mL) and extracted with ethyl acetate (2 x 200 mL). The combined organic phases were concentrated in vacuo and the residue chromatographed on silica gel (eluted with 3% ethyl acetate in hexane) to give 1.95 g (Y: 29%) of the title compound; ^1H -NMR (CDCl_3): δ 8.13 (d, $J=2.3$ Hz, 1H), 7.62 (dd, $J=8.5$, 2.3 Hz, 1H), 7.31 (d, $J=8.5$ Hz,

1H), 2.73 (t, J=7.0 Hz, 2H), 2.01 (t, J=7.0 Hz, 2H), 1.38 (s, 6H); MS (DCI) m/e: 253 (MH⁺).

5

EXAMPLE 96

4,4-Dimethyl-7-trimethylsilyl-ethynyl-1-tetralone
(IL)

10 To a deaerated solution of compound XLVIII (1.53 g, 6.04 mmol), palladium (II) acetate (Aldrich, 15.4 mg, 0.975 mmol) and triphenylphosphine (31 mg, 0.118 mmol) in anhydrous triethylamine (30 mL) was added (trimethylsilyl)acetylene (Aldrich, 3.0 mL, 21.23
15 mmol) and heated to 100°C over 0.5 h. After 4 h at 100°C the reaction mixture was cooled to room temperature (16 h). The reaction mixture was then filtered and the filtrate concentrated in vacuo. The residue was chromatographed on silica gel (eluted with
20 3% ethyl acetate in hexane) to give 1.24 g (Y: 76%) of the title product; ¹H-NMR (CDCl₃): δ 8.11 (d, J=1.8 Hz, 1H), 7.58 (dd, J=8.2, 1.8 Hz, 1H), 7.36 (d, J=8.2 Hz, 1H), 2.72 (t, J=7.0 Hz, 2H), 2.01 (t, J=7.0 Hz, 2H), 1.37 (s, 6H), 0.24 (s, 9H); MS (DCI) m/e 271 (MH⁺).

25

EXAMPLE 97

4,4-Dimethyl-7-ethynyl-1-tetralone (L)


30

To a solution of compound IL (1.24 g, 4.59 mmol), in absolute ethyl alcohol (8.0 mL) was added anhydrous potassium carbonate (170 mg, 1.21 mmol) at room temperature. After 2 h the reaction mixture was

concentrated in vacuo, diluted with saturated sodium bicarbonate (25 mL) and extracted with methylene chloride (2 x 25 mL). The organic phases were combined, concentrated in vacuo and the residue chromatographed on silica gel (eluted with 5% ethyl acetate in hexane) to give 694 mg (Y: 76%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 8.15 (d, $J=1.9$ Hz, 1H), 7.62 (dd, $J=8.2$, 1.9 Hz, 1H), 7.39 (d, $J=8.2$ Hz, 1H), 3.07 (s, 1H), 2.73 (t, $J=7.0$ Hz, 2H), 2.02 (t, $J=7.0$ Hz, 2H), 1.39 (s, 6H); MS (DCI) m/e : 199 (MH^+).

EXAMPLE 98

15 4-[[[(5,6,7,8-Tetrahydro-5,5-dimethyl-8-oxo)-2-naphthalenyl]ethynyl]benzoic acid, ethyl ester (LIa)




20 A solution of compound L (153 mg, 0.773 mmol), ethyl 4-iodobenzoate (Lancaster, 160 mg, 0.58 mmol), bis(triphenylphosphine) palladium (II) chloride (Aldrich, 923 mg, 0.013 mmol) and copper (I) iodide (Aldrich, 4.5 mg, 0.02 mmol) in anhydrous triethylamine (3.0 mL) was allowed to stir at room temperature for 2 h. The reaction mixture was diluted with ethyl acetate (25 mL) and washed with water (1 x 25 mL). The organic phase was concentrated in vacuo and the residue chromatographed on silica gel (eluted with 10% ethyl acetate in hexane) to give 183 mg (Y: 91%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 8.20 (d, $J=1.8$ Hz, 1H), 8.04 (d, $J=8.3$ Hz, 2H), 7.67 (dd,

J=8.2, 1.8 Hz, 1H), 7.58 (d, J=8.3 Hz, 2H), 7.44 (d, J=8.2 Hz, 1H), 4.39 (q, J=7.1 Hz, 2H), 2.76 (t, J=6.8 Hz, 2H), 2.04 (t, J=6.8 Hz, 2H), 1.41 (t, J=7.1 Hz, 3H), 1.41 (s, 6H); MS (DCI) m/e: 347 (MH⁺).

5

EXAMPLE 99

4-[[[(5,6,7,8-Tetrahydro-5,5-dimethyl-8-phenyl-8-hydroxy)-2-naphthalenyl]ethynyl]benzoic acid, ethyl ester (LIIa)




Using the method for the preparation of the 8-phenyl derivative XVIIa, 183 mg (0.529 mmol) of compound LIa gave 177 mg (Y: 79%) of the title compound; ¹H-NMR (CDCl₃): δ 7.98 (d, J=8.3 Hz, 2H), 7.50 (d, J=8.3 Hz, 2H), 7.47-7.26 (m, 8H), 4.37 (q, J=7.0 Hz, 2H), 2.20 (m, 2H), 1.89 (m, 1H), 1.60 (m, 1H), 1.41 (s, 6H), 1.39 (t, J=7.0 Hz, 3H); MS (DCI) m/e: 425 (MH⁺).

25

EXAMPLE 100

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]ethynyl]benzoic acid (I¹⁶a)



To a solution of compound LIIa (177 mg, 0.42 mmol) was added a few milligrams (~ 10-20 mg) of p-toluenesulfonic acid. After heating at 70°C for 0.5 h, the reaction mixture was cooled and concentrated in vacuo to afford 4-[[5,6-dihydro-5,5-dimethyl-8-phenyl]-2-naphthalenyl]ethynyl]benzoic acid, ethyl ester (I^{15a}). The residue was then dissolved in ethyl alcohol (5.0 mL) and treated with 10 N NaOH (6.5 mmol, 0.65 mL) at 70°C. After 0.5 h, an excess of 1N HCl (20 mL) was added and the precipitate collected by vacuum filtration to give 142 mg (Y: 90%) of the title product; ¹H-NMR (DMSO-d₆): δ 13.13 (s, 1H), 7.90 (d, J=8.4 Hz, 2H), 7.58 (d, J=8.4 Hz, 2H), 7.39 (m, 6H), 7.30 (d, J=6.6 Hz, 1H), 6.99 (s, 1H), 6.03 (t, J=4.5 Hz, 1H), 2.33 (d, J=4.5 Hz, 2H), 1.29 (s, 6H); ¹³C-NMR (DMSO-d₆): 166.64, 146.14, 139.79, 137.96, 133.75, 131.50, 130.98, 130.37, 129.48, 128.63, 128.38, 127.88, 127.60, 127.49, 126.54, 124.69, 119.23, 91.96, 88.11, 37.98, 33.40, 27.75; MS (DCI) m/e: 379 (MH⁺); IR (KBr): 2958, 1684, 1604, 1420 cm⁻¹.

Anal. calcd. for C₂₇H₂₂O₂ · 0.15 H₂O: C, 85.08; H, 5.90.
 Found: C, 85.06; H, 5.97.

EXAMPLE 101

4,4-Dimethyl-7-[(dimethylamino)thiocarbonyloxy]-1-tetralone (XLII)

5 To a solution of 4,4-dimethyl-7-hydroxy-1-tetralone (XIV) (3.07 g, 16.16 mmol) in water (10.77 mL) containing potassium hydroxide (904 mg) cooled below 10°C was added N,N-dimethylthiocarbamoyl chloride
10 (2.67 g, 21.6 mmol) in tetrahydrofuran (4.30 mL); the reaction temperature kept below 12°C. After 10 minutes, reaction mixture is made basic by the addition of a 10% aqueous KOH solution (5.39 mL). Product is extracted with ethyl acetate (2 x 50 mL),
15 organic phases concentrated in vacuo and the residue chromatographed on silica gel (eluted with 10% ethyl acetate in hexane) to give 3.74 g (Y: 84%) of title product; ¹H-NMR (CDCl₃): δ 7.68 (d, J=2.6 Hz, 1H), 7.45 (d, J=8.5 Hz, 1H), 7.27 (dd, J=8.5, 2.6 Hz, 1H), 3.46
20 (s, 3H), 3.34 (s, 3H), 2.73 (t, J=6.8 Hz, 2H), 2.04 (t, J=6.8 Hz, 2H), 1.41 (s, 6H); MS (DCI) m/e: 278 (MH⁺).

EXAMPLE 102

25


4,4-Dimethyl-7-mercapto-1-tetralone (XLIII)

Compound XLII (3.74 g 13.50 mmol) under nitrogen with
30 a gas outlet was heated at 270-275°C for 0.75 h in a salt bath (1:1 mole mixture of potassium nitrate and sodium nitrite). After cooling, potassium hydroxide (1.13 g) in water (1.35 mL) and ethylene glycol (10.0 mL) was added to the reaction mixture and it was
35 refluxed for 1 h. The cooled reaction mixture was then

poured onto ice (30 g). The mixture was washed with methylene chloride (2 x 50 mL), made acidic with 1 N HCl and extracted with methylene chloride (2 x 50 mL). The organic phase is concentrated in vacuo and the residue chromatographed on silica gel (eluted with 5% ethyl acetate in hexane) to give 1.22 g (Y: 44%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.91 (d, $J=2.2$ Hz, 1H), 7.41 (dd, $J=8.2, 2.2$ Hz, 1H), 7.30 (d, $J=8.2$ Hz, 1H), 3.48 (s, 1H), 2.71 (t, $J=6.8$ Hz, 2H), 2.00 (t, $J=6.8$ Hz, 2H), 1.37 (s, 6H); MS (DCI) m/e : 207 (MH^+).

EXAMPLE 103

15 4-[[[(5,6,7,8-Tetrahydro-5,5-dimethyl-8-oxo)-2-naphthalenyl]sulfamyl]methyl]benzoic acid, methyl ester (XLVIA)




20 A solution of 4,4-dimethyl-7-mercapto-1-tetralone (XLIII) (262 mg, 1.27 mmol), methyl 4-(bromomethyl)benzoate (Aldrich, 291 mg, 1.27 mmol) and diisopropylethylamine (Aldrich, 179 mg, 1.38 mmol) in anhydrous methylene chloride was allowed to stir at room temperature. After 1.5 h the reaction mixture was concentrated in vacuo and the residue chromatographed on silica gel (eluted with 5% ethyl acetate in hexane) to give 402 mg (Y: 89%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.98 (d, $J=2.1$ Hz, 1H), 7.95 (d, $J=8.5$ Hz, 2H), 7.38 (dd, $J=8.3, 2.1$ Hz, 1H), 7.38 (d, $J=8.5$ Hz, 2H), 7.29 (d, $J=8.3$ Hz, 1H), 4.17 (s, 2H), 3.90 (s,

3H), 2.71 (t, J=6.8 Hz, 2H), 1.99 (t, J=6.8 Hz, 2H), 1.36 (s, 6H); MS (DCI) m/e: 355 (MH⁺).

EXAMPLE 104

5

4-[[[(5,6,7,8-Tetrahydro-5,5-dimethyl-8-phenyl-8-hydroxy)-2-naphthalenyl]sulfamyl]methyl]benzoic acid, methyl ester (XLVIIa)



10

Using the method for the preparation of the 8-phenyl derivative XVIIa, 400 mg (1.13 mmol) of compound XLVIa gave 415 mg (Y: 85%) of the title compound; ¹H-NMR


15 (CDCl₃): δ 7.87 (d, J=8.2 Hz, 2H), 7.30-7.18 (m, 9H), 7.02 (d, J=2.0 Hz, 1H), 3.97 (s, 2H), 3.91 (s, 3H), 2.22-2.08 (m, 3H), 1.85 (m, 1H), 1.36 (s, 3H), 1.28 (s, 3H); MS (DCI) m/e: 433 (MH⁺).

20

EXAMPLE 105

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]sulfamyl]methyl]benzoic acid (I¹⁴a)

25




Using the method for the preparation of compound 4-

[(E)-(5,6,-dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)vinyl]benzoic acid (I^{11a}), 415 mg (0.96 mmol) of compound XLVIIa gave 365 mg (Y: 95%) of the title compound; ¹H-NMR (DMSO-d₆): δ 12.87 (s, 1H), 7.78 (d, J=8.2 Hz, 2H), 7.37 (m, 3H), 7.29 (s, 1H), 7.25 (d, J=8.2 Hz, 2H), 7.25 (d, J=8.1 Hz, 1H), 7.13 (m, 2H), 6.67 (d, J=1.9 Hz, 1H), 5.95 (t, J=4.5 Hz, 1H), 4.09 (s, 2H), 2.27 (d, J=4.5 Hz, 2H), 1.24 (s, 6H); MS (DCI) m/e: 401 (MH⁺); IR (KBr): 2956, 1684, 1610, 1422, 1286 cm⁻¹.

Anal. calcd. for C₂₆H₂₄O₂S₁ · 0.5 H₂O: C, 76.25; H, 6.15. Found: C, 76.20; H, 6.15.

EXAMPLE 106

5,5-Dimethyl-5,6-dihydro-8-phenyl-naphthalene-2-carboxylic acid, N,O-dimethyl hydroxy amide (LIIIIa)



To a solution of compound XVIIIa (1.04 g, 3.74 mmol) in anhydrous methylene chloride (15.0 mL) was added oxalyl chloride (0.39 mL, 4.49 mmol) and 2 drops of N,N-dimethylformamide at 0°C. The reaction mixture was then allowed to stir at room temperature. After 2 hours, N,O-dimethylhydroxylamine hydroxychloride (Aldrich, 401 mg, 4.11 mmol) and anhydrous pyridine (650 mg, 0.665 mL) were added at 0°C. After 2 hours at room temperature, the reaction mixture was


concentrated in vacuo, diluted with ethyl acetate (50 mL) washed with brine (2 x 50 mL), dried over anhydrous magnesium sulfate and concentrated in vacuo to give 1.13 (Y: 94%) of the title product; $^1\text{H-NMR}$ (CDCl₃): δ 7.55 (dd, J=8.0, 1.9 Hz, 1H), 7.39 (d, J=8.0 Hz, 1H), 7.37 (m, 5H), 7.32 (d, J=1.8 Hz, 1H), 6.01 (t, J=4.5 Hz, 1H), 3.49 (s, 3H), 3.27 (s, 3H), 2.37 (d, J=4.5 Hz, 2H), 1.35 (s, 6H); MS (DCI) m/e: 322 (MH⁺).

10

EXAMPLE 107

5,5-Dimethyl-5,6-dihydro-2-formyl-8-phenylnaphthalene (LIIVa)

15




To a solution of compound LIIIIa (1.13 g, 3.52 mmol) in tetrahydrofuran (32 mL) at -78°C was added diisobutylaluminum hydride (DIBAL, 1.0 M solution in hexanes, 5.28 mmol, 5.28 mL). After 1 h, the reaction mixture was diluted with 5% HCl in ethyl alcohol (10 mL) and brine (10 mL). Mixture was then extracted with ethyl acetate (2 x 40 mL). The organic phases were combined, dried over anhydrous magnesium sulfate and concentrated in vacuo to give 720 mg (Y: 78%) of the title product; $^1\text{H NMR}$ (CDCl₃): δ 9.86 (s, 1H), 7.76 (dd, J=8.0, 1.7 Hz, 1H), 7.54-7.52 (m, 2H), 7.52-7.26 (m, 5H), 6.07 (t, J=4.5 Hz, 1H), 2.40 (d, J=4.5 Hz, 2H), 1.38 (s, 6H); MS (DCI) m/e: 263 (MH⁺).

30

EXAMPLE 108

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]methyl]amino]benzoic acid, methyl ester

5 (I²⁰a)




To a solution of methyl 4-aminobenzoate (505 mg, 3.35
10 mmol) in a 1% glacial acetic acid methyl alcohol
solution (6.0 mL) was added LIVa (720 mg, 2.75 mmol)
in a 1% glacial acetic acid methyl alcohol solution
(6.0 mL). Sodium cyanoborohydride (174 mg, 2.75 mmol)
was then added over an one hour period (6 x 29 mg
15 portions every 10 minutes). After 16 h at room
temperature, the reaction mixture was concentrated in
vacuo and the residue chromatographed on silica gel
(eluted with 10% ethyl acetate in hexane) to give 856
mg (Y: 78%) of the title product; ¹H-NMR (CDCl₃): δ
20 7.83 (d, J=8.9 Hz, 2H), 7.34 (m, 6H), 7.21 (dd, J=7.8,
2.0 Hz, 1H), 6.99 (d, J=2.0 Hz, 1H), 6.52 (d, J=8.9
Hz, 2H), 5.99 (t, J=4.5 Hz, 1H), 4.37 (m, 1H), 4.23
(d, J=5.5 Hz, 2H), 3.84 (s, 3H), 2.35 (t, J=4.5 Hz,
2H), 1.34 (s, 6H); MS (DCI) m/e: 398 (MH⁺).

25

EXAMPLE 109

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-

30 naphthalenyl]methyl]amino]benzoic acid, hydrochloride
(I²¹a)




To a solution of compound I²⁰a (197 mg, 0.496 mmol) in ethyl alcohol (5.0 mL) was added 10 N NaOH (22.0 mmol, 2.2 mL). After 4 h at reflux, an excess of 1N HCl (40 mL) was added and the precipitate collected by vacuum filtration to give 175 mg (Y: 92%) of the title product; ¹H-NMR (DMSO-d₆): δ 7.60 (d, J=8.8 Hz, 2H), 7.32 (m, 4H), 7.20 (m, 3H), 6.90 (d, J=1.6 Hz, 1H), 6.49 (d, J=8.8 Hz, 2H), 5.96 (t, J=4.5 Hz, 1H), 4.16 (s, 2H), 2.27 (d, J=4.5 Hz, 2H), 1.24 (s, 6H); MS (DCI) m/e: 384 (MH⁺); IR (KBr): 3418, 2960, 1672, 1602, 1176 cm⁻¹.

Anal. calcd. for C₂₆H₂₅O₂N₁·1.0 HCl: C, 74.36; H, 6.24; N, 3.34. Found: C, 74.39; H, 6.19; N, 3.29.

EXAMPLE 110

20

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]aminolthiocarbonyl]benzoic acid, methyl ester




25

Using the method given for the preparation of 4-
 [[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-
 naphthalenyl]thiocarbonyl]amino]benzoic acid, methyl
 ester, 190 mg (0.462 mmol) of compound I¹c gave 104 mg
 5 (Y: 52%) of the title product; ¹H-NMR (CDCl₃): δ 8.84
 (s, 1H), 8.05 (d, J=8.2 Hz, 2H), 7.91 (dd, J=8.8, 2.4
 Hz, 1H), 7.81 (d, J=8.2 Hz, 2H), 7.44 (d, J=8.8 Hz,
 1H), 7.35 (m, 5H), 7.13 (d, J=2.4 Hz, 1H), 6.04 (t,
 J=4.5 Hz, 1H), 3.94 (s, 3H), 2.38 (d, J=4.5 Hz, 2H),
 10 1.35 (s, 6H); MS (DCI) m/e: 428 (MH⁺).

EXAMPLE 111

15 4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-
 naphthalenyl]amino]thiocarbonyl]benzoic acid (I¹⁹a)




20 Using the method given for the preparation of the 8-
 phenyl derivative I⁴a, 104 mg (0.240 mmol) of compound
 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-
 naphthalenyl]amino]thiocarbonyl]benzoic acid, methyl
 ester gave 80 mg (Y: 80%) of the title product; ¹H-NMR
 25 (DMSO-d₆): δ 11.80 (s, 1H), 7.93 (d, J=8.4 Hz, 2H),
 7.77 (d, J=8.4 Hz, 2H), 7.75 (s, 1H), 7.43 (d, J=8.4
 Hz, 1H), 7.34 (m, 6H), 6.02 (t, J=4.5 Hz, 1H), 2.33

(d, $J=4.5$ Hz, 2H), 1.30 (s, 6H); MS (DCI) m/e : 414 (MH^+); IR (KBr): 2958, 1694, 1490, 1410 cm^{-1} .

Anal. calcd for $C_{26}H_{23}O_2N_1S_1 \cdot 0.5 H_2O$: C, 73.91; H, 5.73, 5 N, 3.31. Found: C, 73.78,; H, 5.55; N, 3.22.

EXAMPLE 112

10 Methyl 4-mercaptobenzoate




To a solution of 4-mercaptobenzoic acid (Apin, 1.72 g, 15 11.17 mmol) in anhydrous methyl alcohol (20.0 mL) was added concentrated sulfuric acid (0.43 mL). The reaction mixture is then warmed to reflux for 16 h, concentrated in vacuo, diluted with ethyl acetate (100 mL) and washed with saturated sodium bicarbonate (2 x 20 100 mL). The organic phase is then separated, dried over anhydrous magnesium sulfate and concentrated in vacuo to give 712 mg (Y: 38%) of the title product; 1H -NMR ($CDCl_3$): δ 7.89 (d, $J=8.7$ Hz, 2H), 7.29 (d, $J=8.7$ Hz, 2H), 3.90 (s, 3H), 3.60 (s, 1H).

25

EXAMPLE 113


4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]methyl]sulfamyl]benzoic acid, methyl ester
30 (I^{24a})



To a solution of methyl 4-mercaptobenzoate (89 mg, 0.528 mmol) in ethylene glycol dimethyl ether (3.0 mL) was added 60% sodium hydride (0.607 mmol, 24 mg) at room temperature. After 0.15 h compound LVIIIa (190 mg, 0.581 mmol) was added. After 2 h the reaction mixture was diluted with ethyl acetate (50 mL) and washed with 1N hydrochloric acid (2 x 50 mL). The organic phase was concentrated in vacuo and the residue chromatographed on silica gel (eluted with 10% ethyl acetate in hexane) to give 30 mg (Y: 13%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.88 (d, $J=8.5$ Hz, 2H), 7.28 (m, 9H), 6.98 (d, $J=1.4$ Hz, 1H), 5.98 (t, $J=4.5$ Hz, 1H), 4.07 (s, 2H), 3.90 (s, 3H), 2.33 (d, $J=4.5$ Hz, 2H), 1.32 (s, 6H); MS (DCI) m/e : 415 (MH^+).

EXAMPLE 114

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]methyl]sulfamyl]benzoic acid (I^{34}a)



Using the method given for the preparation of the 8-phenyl derivative I^4a , 30 mg (0.07 mmol) of compound


I²⁴a gave 21 mg (Y: 72%) of the title product; ¹H-NMR (DMSO-d₆): δ 12.87 (s, 1H), 7.77 (d, J=8.5 Hz, 2H), 7.35 (m, 7H), 7.28 (d, J=1.8 Hz, 1H), 7.19 (dd, J=7.9, 1.8 Hz, 1H), 6.90 (d, J=1.5 Hz, 1H), 5.95 (t, J=4.5 Hz, 1H), 4.19 (s, 2H), 2.27 (d, J=4.5 Hz, 2H), 1.24 (s, 6H); MS (DCI) m/e: 401 (MH⁺); IR (KBr): 3436, 2960, 1688, 1592 cm⁻¹.

Anal. calcd. for C₂₆H₂₄O₂S₁·H₂O: C, 76.93; H, 6.11.

10 Found: C, 76.91; H, 6.03.

EXAMPLE 115

15 1,2-Dihydro-1,1-dimethyl-6-nitro-8-(trifluoromethanesulfonyloxy)naphthalene (LXI)




- 20 To a solution of 4,4-dimethyl-7-nitro-1-tetralone (VI) (2.4 g, 10.96 mmol) in tetrahydrofuran (10 mL) at -78°C was added lithium bis(trimethylsilyl)amide (1.0 M solution in hexane, 12.05 mmol, 12.05 mL) and then N-(2-pyridyl)triflimide (12.05 mmol, 4.30 g) at -78°C.
- 25 After 0.5 h at -78°C the reaction mixture was diluted with water and extracted with ethyl acetate (2 x 50 mL). The organic phases were combined and concentrated in vacuo. The residue was chromatographed on silica gel (eluted with 5% ethyl acetate in hexane) to give
- 30 2.68 g (Y: 70%) of the title product; ¹H-NMR (CDCl₃): δ 8.20 (m, 2H), 7.49 (d, J=8.2 Hz, 1H), 6.15 (t, J=4.5

Hz, 1H), 2.50 (d, J=4.5 Hz, 2H), 1.37 (s, 6H); MS (DCI) m/e: 352 (MH⁺).

5

EXAMPLE 116

1,2-Dihydro-1,1-dimethyl-6-nitro-4-phenylnaphthalene (LXIa)




10

To a solution of compound LXI (2.68 g, 7.64 mmol) in anhydrous 1-methyl-2-pyrrolidinone (30.0 ml) was added triphenylarsine (442 mg, 1.44 mmol), tris (dibenzylideneacetone) dipalladium (0) (217 mg, 0.237 mmol) and tributylphenyltin (6.17 g, 16.78 mmol) in anhydrous 1-methyl-2-pyrrolidinone (5.0 ml). After 16 h. at 60° C the mixture was diluted with water and extracted with ethyl acetate (2 x 100 ml). The organic phases were combined, stirred over a saturated aqueous solution of potassium fluoride for 0.5 h., separated and concentrated in vacuo. The residue was chromatographed on silica gel (eluted with 20% ethyl acetate in hexane) to give 1.93 g (Y: 90%) of the title product; ¹H-NMR (CDCl₃): δ 8.07 (dd, J=8.5, 2.5 Hz, 1H), 7.87 (d, J=2.5 Hz, 1H), 7.50 (d, J=8.5 Hz, 1H), 7.36 (m, 5H), 6.12 (t, J=4.5 Hz, 1H), 2.41 (d, J=4.5 Hz, 2H), 1.39 (s, 6H); MS (DCI) m/e: 280 (MH⁺).

30

EXAMPLE 117


5,5-Dimethyl-8-phenyl-5,6-dihydro-naphthalene-2-amine (LXIIa)



- 5 To a solution of compound LXIIa (1.75 g, 6.27 mmol) in benzene (15 mL) was added ferric chloride (33%, 2.94 mL) and water (11.9 mL). The reaction mixture was heated to reflux and iron (3.50 g) was added over a
- 10 period of 1.5 h in four equal aliquots. The resulting mixture was then refluxed for 16 h. After cooling to room temperature, the reaction mixture was filtered through a pad of celite. The organic phase was separated, dried over anhydrous magnesium sulfate and
- 15 concentrated in vacuo to give 1.56 g (Y: 99%) of the title product; ¹H-NMR (CDCl₃): δ 7.35 (m, 5H), 7.15 (d, J=8.1 Hz, 1H), 6.57 (dd, J=8.1, 2.5 Hz, 1H), 6.37 (d, J=2.5 Hz, 1H), 5.95 (t, J=4.5 Hz, 1H), 3.45 (bs, 2H), 2.31 (d, J=4.5 Hz, 2H), 1.30 (s, 6H); MS (DCI) m/z:
- 20 250 (MH⁺).

EXAMPLE 118

- 25 4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]amino]methyl]benzoic acid, methyl ester (I^{28a})



Using the method for the preparation of 4-[[[(5,6-


5 dihydro-5,5-dimethyl-8-phenyl)-2-
 5 naphthalenyl]methyl]amino]benzoic acid, methyl ester
 (I^{20a}), 1.35 g (5.38 mmol) of compound LXIIIa and
 methyl 4-formylbenzoate (Aldrich, 723 mg, 4.41 mmol)
 gave 1.38 g (Y: 79%) of the title product; $^1\text{H-NMR}$
 (CDCl_3): δ 7.95 (d, $J=8.5$ Hz, 2H), 7.29 (m, 7H), 7.16
 10 (d, $J=8.3$ Hz, 1H), 6.47 (dd, $J=8.3$, 2.6 Hz, 1H), 6.29
 (d, $J=2.6$ Hz, 1H), 5.94 (t, $J=4.5$, 1H), 4.53 (s, 2H),
 3.91 (s, 3H), 2.30 (d, $J=4.5$ Hz, 2H), 1.29 (s, 6H); MS
 (DCI) m/e : 398 (MH^+).

15

EXAMPLE 119

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-
naphthalenyl]amino]methyl]benzoic acid (I^{29a})

20




To a solution of compound I^{28a} (1.38 g, 3.48 mmol) in a
 1:1 mixture of ethyl alcohol/tetrahydrofuran (15.0 mL)
 25 was added 10 N NaOH (35.0 mmol, 3.5 mL) at room
 temperature. After 16 h an excess of 1N HCl (100 mL)
 was added and the precipitate collected by vacuum

filtration to give 1.28 g (Y: 96%) of the title product as a hydrochloride salt; $^1\text{H-NMR}$ (DMSO-d_6): δ 7.84 (d, $J=8.2$ Hz, 2H), 7.36 (d, $J=8.2$ Hz, 1H), 7.31 (m, 5H), 7.15 (m, 4H), 6.99 (bs, 1H), 6.36 (bs, 1H), 5.90 (t, $J=4.5$ Hz, 1H), 4.32 (s, 2H), 2.21 (d, $J=5.5$ Hz, 2H), 1.19 (s, 6H); MS (DCI) m/e : 384 (MH^+); IR (KBr): 3420, 2958, 1694, 1606 cm^{-1} .

Anal. calcd. for $\text{C}_{26}\text{H}_{25}\text{N}_1\text{O}_2 \cdot 1.0 \text{ HCl}$: C, 72.80; H, 6.34; N, 3.27. Found: C, 72.98; H, 6.09; N, 3.14.

EXAMPLE 120

15 3,3-Dimethyl-6-nitro-indan-1-one




To a solution of 3,3-dimethylindan-1-one (LXIV) (16.3 g, 50.9 mmol) (Harms, W. M. and Eisenbraun, E. J. *Org. Prep. Proc. Int.* 1972, 4, 67-72) in 56 mL of sulfuric acid was slowly added 4.18 mL of 70% nitric acid in 20 mL of sulfuric acid at 0-5 °C. After being stirred for 1 hour, the reaction mixture was poured into 600 g of ice. The precipitates was filtered and washed with water. The solids collected were dissolved in 600 mL of ethyl acetate and washed with NaHCO_3 (80 mL X 2), dried over magnesium sulfate, and evaporated. The residue was triturated with methanol to give 22.2 g of the title compound as yellow solids; $^1\text{H-NMR}$ (CDCl_3) δ 1.49 (s, 6 H), 2.71 (s, 2 H), 7.68 (d,

$J = 8.4$ Hz, 1 H), 8.48 (dd, $J = 2.0, 8.4$ Hz, 1 H), 8.53 (d, $J = 2.0$ Hz, 1 H); MS m/e 206 (MH^+).

Anal. calcd. for $C_{11}H_{11}NO_3$: C, 64.38; H, 5.40; N, 6.83.
5 Found: C, 64.33; H, 5.33; N, 6.86.

EXAMPLE 121

10 6-Amino-3,3-dimethyl-indan-1-one (LXV)

3,3-Dimethyl-6-nitro-indan-1-one (22.2 g, 54.1 mmol)
15 and platinum oxide (0.44 g, 1.94 mmol) in 30 mL of ethyl acetate and 60 mL of methanol was hydrogenated on a Parr Shaker for 40 min. The mixture was filtered through a pad of celite. The filtrate was evaporated and the residue was triturated with hexane to give
20 18.4 g (97% yield) of the title product as yellow solids; 1H -NMR ($CDCl_3$) δ 1.38 (s, 6 H), 2.56 (s, 2 H), 6.98 (d, $J = 2.3$ Hz, 1 H), 7.02 (dd, $J = 2.3, 8.2$ Hz, 1 H), 7.30 (d, $J = 8.2$ Hz, 1 H); MS m/e 176 (MH^+).

25 Anal. calcd. for $C_{11}H_{11}NO$: C, 75.40; H, 7.48; N, 7.99.
Found: C, 75.07; H, 7.45; N, 7.94.

EXAMPLE 122

30


6-Hydroxy-3,3-dimethyl-indan-1-one (LXVI)

A solution of sodium nitrite (14.8 g, 214 mmol) in 30 mL of water was slowly added to the solution of 6-amino-3,3-dimethyl-indan-1-one (14.0 g, 85.8 mmol) in 30 mL of 48% tetrafluoroboric acid and 30 mL of water at 0-5 °C. After being stirred for 30 min, the mixture was filtered, and the solids were washed with cold 5% tetrafluoroboric acid and dried in vacuum. The solids were then added in several portions into a boiling solution of 50 mL of sulfuric acid in 0.5 liter of water. After refluxing for 1 hour, the solution was cooled to room temperature and extracted with ethyl acetate (100 mL X 3). The combined extracts were dried over magnesium sulfate and evaporated. The residue was purified by flash chromatography (EtOAc:hexane = 1:10 to 1:2) to give 7.60 g (50% yield) of the title compound as a solid; ¹H-NMR (CDCl₃) δ 1.40 (s, 6 H), 2.64 (s, 2 H), 7.13 (s, 1 H), 7.18 (d, J = 8.2 Hz, 1 H), 7.39 (d, J = 8.2 Hz, 1 H); MS m/e 177 (MH⁺).

Anal. calcd. for C₁₁H₁₂O₂: C, 74.98; H, 6.86.
Found: C, 74.81; H, 6.76.

EXAMPLE 123

Trifluoromethanesulfonic acid 1,1-dimethyl-3-oxo-indan-5-yl ester



To a solution of 6-hydroxy-3,3-dimethyl-indan-1-one (4.01 g, 24.4 mmol) and 4-dimethylaminopyridine (DMAP, 5.97 g, 48.9 mmol) in 30 mL of methylene chloride was slowly added triflic acid anhydride (9.65 g, 34.2 mmol) at -78 °C. After stirring at room temperature for 1 hour, the mixture was diluted with methylene chloride (60 mL), washed with 1 N HCl (10 mL X 2) and water (10 mL), dried over magnesium sulfate, and evaporated. The residue was purified by flash chromatography (EtOAc:hexane = 1:10 to 1:5) to give 5.58 g (74% yield) of the title compound; ¹H-NMR (CDCl₃) δ 1.46 (s, 6 H), 2.67 (s, 2 H), 7.51 (dd, J = 2.3, 8.4 Hz, 1 H), 7.57 (d, J = 2.3 Hz, 1 H), 7.60 (d, J = 8.4 Hz, 1 H); MS m/e 309 (MH⁺).

Anal. calcd. for C₁₂H₁₁F₃O₄S: C, 46.75; H, 3.60. Found: C, 46.83; H, 3.58.

EXAMPLE 124

Methyl 1,1-dimethyl-3-oxo-indan-5-carboxylate (LXVII)

A solution of trifluoromethanesulfonic acid 1,1-dimethyl-3-oxo-indan-5-yl ester (3.08 g, 10.0 mmol), triethylamine (2.02 g, 20.0 mmol), palladium acetate (0.11 g, 0.50 mmol) and 1,3-bis(diphenylphosphino)propane (dppp, 0.21 g, 0.50 mmol) in 30 mL of anhydrous dimethylsulfoxide and 20 mL of methanol was saturated with carbon monoxide for 10 min, and then stirred under a balloon filled with carbon monoxide at 65-70 °C for 2 hours. Methanol was evaporated, and the resulting solution was diluted with 50 mL of water, extracted with ethyl ether (40 mL


X 4). The combined extracts were dried over magnesium sulfate and evaporated. The residue was purified by flash chromatography (EtOAc:hexane = 1:20 to 1:5) to give 1.98 g (91 % yield) of the title compound as
 5 light yellow solids; $^1\text{H-NMR}$ (CDCl_3) δ 1.45 (s, 6 H), 2.65 (s, 2 H), 3.94 (s, 3 H), 7.58 (dd, $J = 8.1$ Hz, 1 H), 8.30 (dd, $J = 1.6, 8.1$ Hz, 1 H), 8.36 (d, $J = 1.6$ Hz, 1 H); MS m/e 219 (MH^+).

10 Anal. calcd. for $\text{C}_{13}\text{H}_{14}\text{O}_3$: C, 71.54; H, 6.47.
 Found: C, 71.71; H, 6.46.

EXAMPLE 125

15

Methyl 1,1-dimethyl-3-(trifluoromethanesulfonyloxy)-
1H-indene-5-carboxylate



20


To a solution of methyl 1,1-dimethyl-3-oxo-indan-5-carboxylate (0.90 g, 4.12 mmol) and 2,6-di-tert-butyl-4-methylpyridine (1.10 mg, 5.36 mmol) in 10 mL of methylene chloride was added triflic acid anhydride
 25 (1.39 g, 4.94 mmol) at -78°C . After being stirred at room temperature for 16 hours, the mixture was diluted with 70 mL of ethyl ether, washed with 1 N HCl (20 mL), dried over magnesium sulfate, and evaporated. The residue was purified by flash chromatography
 30 (EtOAc:hexane = 1: 20 to 1:5) to give 1.23 g (85%

yield) of the title compound; ^1H NMR (CDCl_3) δ 1.41 (s, 6 H), 3.95 (s, 3 H), 6.30 (s, 1 H), 7.44 (d, $J = 7.8$ Hz, 1 H), 7.99 (s, 1 H), 8.06 (d, $J = 7.8$ Hz, 1 H); MS m/e 351 (MH^+).

5

EXAMPLE 126

10 Methyl 1,1-dimethyl-3-phenyl-1H-indene-5-carboxylate
(LXVIIIa)




Methyl 1,1-dimethyl-3-(trifluoromethanesulfonyloxy)-
15 1H-indene-5-carboxylate (580 mg, 1.86 mmol),
tris(dibenzylideneacetone)dipalladium (0) (Pd_2dba_3 , 17
mg, 0.02 mmol), triphenylarsine (46 mg, 0.15 mmol),
lithium chloride (240 mg, 5.59 mmol) and
phenyltributyltin (680 mg, 1.86 mmol) in 5 mL of 2-
20 methyl pyrrolidinone were stirred at 55 °C for 1.5
days. The mixture was diluted with water (30 mL), and
extracted with ethyl ether (30 mL X 3). The combined
extracts were dried over magnesium sulfate and
evaporated. The residue was purified by flash
25 chromatography (EtOAc:hexane = 1: 20 to 1: 5) to give
0.43 g (83 % yield) of the title compound as a white
solid; ^1H -NMR (CDCl_3) δ 1.41 (s, 6 H), 3.91 (s, 3 H),
6.48 (s, 1 H), 7.39-7.62 (m, 6 H), 7.99 (d, $J = 6.8$
Hz, 1 H), 8.14 (s, 1 H); MS m/e 279 (MH^+).

30

Anal. calcd. for $\text{C}_{19}\text{H}_{18}\text{O}_2$. 0.25 H_2O : C, 80.68; H, 6.49.
Found: C, 80.76; H, 6.41.

EXAMPLE 127

Methyl 4-[(1,1-Dimethyl-3-phenyl-1H-indene-5-
 5 carbonyl)aminobenzoate



- 10 Methyl 1,1-dimethyl-3-phenyl-1H-indene-5-carboxylate
 (315 mg, 1.13 mmol) was stirred with 10 N NaOH (1.1
 mL, 11.0 mmol) in 5 mL of methanol and 10 mL of
 tetrahydrofuran at 60 °C for 1.5 hours. The solution
 was concentrated under reduced pressure and acidified
 15 with 1 N HCl (15 mL), extracted with ethyl acetate (15
 mL X 3). The combined extracts were washed with water
 (10 mL), dried over magnesium sulfate, and evaporated.
 The residue was dried in vacuum and dissolved in 2.5
 mL of methylene chloride. To the solution was added
 20 oxalyl chloride (348 mg, 2.74 mmol) and 2 drops of
 dimethylformamide at 0 °C. The solution was stirred at
 0 °C for 30 min and at room temperature for 30 min.
 The mixture was evaporated and dried in vacuum. The
 residue was stirred with methyl 4-aminobenzoate (181
 25 mg, 1.32 mmol) in 2 mL of anhydrous pyridine for 18
 hours. Excess pyridine was evaporated and the residue
 was diluted with 20 mL of 2 N HCl, extracted with
 ethyl ether (30 mL X 4). The combined extracts were
 washed with water (10 mL), dried over magnesium
 30 sulfate and evaporated. The residue was purified by
 flash chromatography (EtOAc: hexane = 1: 20 to 1: 5)


to give 372 mg (85% yield) of the title compound as a glassy mass; $^1\text{H-NMR}$ (CDCl_3) δ 1.43 (s, 6 H), 3.91 (s, 3 H), 6.52 (s, 1 H), 7.40-7.50 (m, 4 H), 7.59-7.61 (m, 2 H), 7.72-7.75 (m, 3 H), 7.96 (s, 1 H), 8.00 (bs, 1 H), 8.05 (d, $J = 8.7$ Hz, 2 H); MS m/e 398 (MH^+).

Anal. calcd. for $\text{C}_{26}\text{H}_{23}\text{NO}_3 \cdot 0.125 \text{ H}_2\text{O}$: C, 78.12; H, 5.93; N, 3.50.

Found: C, 77.97; H, 5.50; N, 3.24

EXAMPLE 128

4-[(1,1-Dimethyl-3-phenyl-1H-indene-5-
carbonyl)amino]benzoic acid (I^{30}a)



Methyl 4-[(1,1-dimethyl-3-phenyl-1H-indene-5-carbonyl)amino]benzoate (359 mg, 0.90 mmol) was stirred with 4.5 mL of 2 N NaOH in 5 mL of methanol and 5 mL of tetrahydrofuran for 10 hours. The solution was acidified with 1N HCl and concentrated under reduced pressure. After addition of 15 mL of water, the mixture was extracted with ethyl acetate (15 mL X 3). The combined extracts were washed with water (10 mL), dried over magnesium sulfate, and evaporated. The residue was triturated in ether-hexane to give 326 mg (94 % yield) of the title compound as solids; $^1\text{H-NMR}$ (DMSO-d_6) δ 1.39 (s, 6 H),

6.68 (s, 1 H), 7.39-7.53 (m, 3 H), 7.64-7.68 (m, 3 H),
7.85-7.97 (m, 6 H), 10.54 (s, 1 H), 12.75 (bs, 1 H);
MS m/e 384 (MH⁺).


5 Anal. calcd. for C₂₅H₂₁NO₃ · 0.25 H₂O: C 77.39; H, 5.46;
N, 3.61.

Found: C, 77.26; H, 5.51; N, 3.52

10

EXAMPLE 129

1,1-Dimethyl-5-nitro-3-(trifluoromethanesulfonyloxy)-
1H-indene



15

To a solution of 3,3-dimethyl-6-nitro-indan-1-one
(LXIX) (1.00 g, 4.65 mmol) and 2,6-di-tert-butyl-4-
methylpyridine (1.19 g, 5.80 mmol) in 10 mL of
20 methylene chloride was added triflic acid anhydride
(1.57 g, 5.58 mmol) at -78 °C. After being stirred at
room temperature for 16 hours, the mixture was diluted
with 1 N HCl (20 mL) and extracted with ethyl ether
(30 mL X 3). The combined extracts were washed with
25 water (20 mL), dried over magnesium sulfate, and
evaporated. The residue was purified by flash
chromatography (EtOAc: hexane = 1: 20 to 1:10) to give
1.37 g (87% yield) of the title compound as an oil; ¹H-
NMR (CDCl₃) δ 1.45 (s, 6 H), 6.43 (s, 1 H), 7.53 (d, J
30 = 8.3 Hz, 1 H), 8.15 (d, J = 2.0 Hz, 1 H), 8.24 (dd, J


= 2.0, 8.3 Hz, 1 H); MS m/e 338 (MH⁺).

Anal. calcd. for C₁₂H₁₀F₃NO₅S: C, 42.73; H, 2.99; N, 4.15.

5 Found: C, 42.70; H, 2.80; 4.10.

EXAMPLE 130

10 1,1-Dimethyl-5-nitro-3-phenyl-1H-indene (LXXa)



1,1-Dimethyl-5-nitro-3-(trifluoromethanesulfonyloxy)-
 15 1H-indene (1.32 g, 2.89 mmol),
 tris(dibenzylideneacetone)dipalladium (0) (21 mg, 0.02
 mmol), triphenylarsine (70 mg, 0.23 mmol), lithium
 chloride (0.37 g, 8.67 mmol) and phenyltributyltin
 (1.06 g, 2.89 mmol) were stirred at 60 °C for 48 hours.
 20 The mixture was diluted with water (30 mL), extracted
 with ethyl ether (30 mL X 3). The combined extracts
 were dried over magnesium sulfate and evaporated. The
 residue was purified by flash chromatography (EtOAc:
 hexane = 1: 25 to 1: 5) to give a solid which
 25 recrystallized from hexane to give 575 mg (75 % yield)
 of the title compound as light yellow crystals;
¹H-NMR (CDCl₃) δ 1.44 (s, 6 H), 6.58 (s, 1 H), 7.40-
 7.60 (m, 6 H), 8.18 (d, J = 2.0, 8.2 Hz, 1 H), 8.32
 (d, J = 2.0 Hz, 1 H); MS m/e 266 (MH⁺).


30

Anal. calcd. for C₁₇H₁₅NO₂: C, 76.96; H, 5.70; N, 5.28.
 Found: C, 76.71; H, 5.69; N, 5.22.

EXAMPLE 131

5-Amino-1,1-dimethyl-3-phenyl-1H-indene (LXXIa)

5




To a mixture of 1,1-dimethyl-5-nitro-3-phenyl-1H-indene (0.56 g, 2.11 mmol) in 5 mL of benzene was
 10 added 33% ferric chloride (1 mL) and 4 mL of water. The mixture was stirred at reflux and iron powder (1.00 g, 5.28 mmol) was added in four portions during 2 hours. The resulting mixture was stirred at reflux for additional 18 hours. The mixture was diluted with
 15 20 mL of water and 20 mL of ethyl acetate, and filtered through a pad of celite. The filtrate was extracted with ethyl acetate (20 mL X 3). The combined extracts were dried over magnesium sulfate to give 0.49 g (98% yield) of the crude title compound
 20 which was not further purified and used in the next reaction; ¹H-NMR (CDCl₃) δ 1.37 (s, 6 H), 6.40 (s, 1 H), 6.71 (dd, J = 2.0, 7.9 Hz, 1 H), 6.93 (d, J = 2.0 Hz, 1 H), 7.19 (d, J = 7.9 Hz, 1 H), 7.36-7.58 (m, 5 H); MS m/e 236 (MH⁺).

25

EXAMPLE 132

Methyl 4-(1,1-dimethyl-3-phenyl-1H-inden-5-
 30 yl)aminolcarbonylbenzoate




5-Amino-1,1-dimethyl-3-phenyl-1H-indene (0.49 g, 2.06 mmol) and terephthalic monomethyl ester chloride (0.49 g, 2.47 mmol) were stirred in 5 mL of pyridine for 18 hours. The solvent was evaporated and the residue was diluted with 1 N HCl (20 mL), extracted with ethyl acetate (20 mL X 3). The combined extracts were washed with saturated NaCl solution, dried over magnesium sulfate, and evaporated. The crude product was purified by flash chromatography (EtOAc:hexane =1:10 to 1:5) to give 772 mg (94% yield) of the title compound which crystallized from EtOAc-hexane to give 720 mg of crystals; ¹H-NMR (CDCl₃) δ 1.41 (s, 6 H), 3.96 (s, 3 H), 6.47 (s, 1 H), 7.37-7.60 (m, 7 H), 7.68 (s, 1 H), 7.84 (s, 1 H), 7.94 (d, J = 8.3 Hz, 2 H), 8.16 (d, J = 8.3 Hz, 2 H); MS m/e 398 (MH⁺).

Anal. calcd. for C₂₆H₂₃NO₃: C, 78.57; H, 5.83; N, 3.52.
Found: C, 78.45; H, 5.76; N, 3.41.

EXAMPLE 133

4-[[[1,1-Dimethyl-3-phenyl-1H-inden-5-yl]amino]carbonyl]benzoic acid (I^{31a})




Methyl 4-[[[(1,1-dimethyl-3-phenyl-1H-inden-5-yl)amino]carbonyl]benzoate (325 mg, 0.82 mmol), 4.1 mL of 2 N NaOH in 10 mL of tetrahydrofuran and 10 mL of methanol were stirred for 2 hours. The mixture was concentrated and acidified with 1 N HCl (20 mL), extracted with ethyl acetate (35 mL X 2). The combined extracts were washed with water (10 mL), dried over magnesium sulfate, and evaporated. The residue was triturated in ether-hexane to give 246 mg (78 % yield) of the title compound as white solids; ¹H-NMR (DMSO-d₆) δ 1.35 (s, 6 H), 6.58 (s, 1 H), 7.38-7.70 (m, 7 H), 7.73 (d, J = 7.9 Hz, 1 H), 7.90 (s, 1H), 8.04 (s, 4 H), 10.38 (bs, 1 H), 13.25 (bs, 1 H); MS m/e 384 (MH⁺).

Anal. calcd. for C₂₅H₂₁NO₃ · 0.25 H₂O: C, 77.40; H, 5.59; N, 3.61.
Found: C, 77.24; H, 5.33; N, 3.41.

EXAMPLE 134

25

Methyl 4-(1,1-dimethyl-3-oxo-indan-5-ylloxymethyl)-benzoate (LXXIIa)



6-Hydroxy-3,3-dimethyl-indan-1-one (0.72 g, 4.37 mmol) in 5 mL of dimethylformamide was added sodium hydride (0.16 g, 6.55 mmol) at 0 °C. After stirring for 30 min, methyl 4-bromomethylbenzoate (1.00 g, 4.37 mmol) was added. The resulting mixture was stirred at 0° C for 1 hour and at room temperature for 2 hours. The mixture was cooled to 0 °C and acidified with 1 N HCl.

10 The solvent was evaporated and the residue was acidified with 20 mL of 1 N HCl and extracted with ethyl ether (30 mL X 3). The combined extracts were dried over magnesium sulfate and evaporated. The residue was purified by flash chromatography (EtOAc: hexane = 1: 20 to 1: 4) to give 0.68 g (48% yield) of the title compounds as a solid; ¹H-NMR (CDCl₃) δ 1.41 (s, 6 H), 2.61 (s, 2 H), 3.93 (s, 1 H), 5.15 (s, 2 H), 7.18 (d, J = 2.5 Hz, 1 H), 7.30 (d, J = 2.5, 8.4 Hz, 1 H), 7.42 (d, J = 8.4 Hz, 1 H), 7.51 (d, J = 8.3 Hz, 2 H), 8.07 (d, J = 8.3 Hz, 2 H); MS m/e 325 (MH⁺).

20


Anal. calcd. for C₂₀H₂₀O₄: C, 74.06; H, 6.21.
Found: C, 73.99; H, 6.24.

25

EXAMPLE 135

Methyl 4-[1,1-dimethyl-3-(trifluoromethanesulfonyloxy)-1H-indene-5-yloxy]methylbenzoate

30



To a solution of methyl 4-(1,1-dimethyl-3-oxo-indan-5-
 5 yloxymethyl)benzoate (675 mg, 2.08 mmol) and 2,6-di-
 tert-butyl-4-methylpyridine (534 mg, 2.60 mmol) in 5
 mL of methylene chloride was added triflic acid
 anhydride (705 mg, 2.50 mmol) at -78 °C. After being
 stirred at room temperature for 16 hours, the mixture
 10 was diluted with 1 N HCl (20 mL), extracted with ethyl
 ether (20 mL X 3). The combined extracts were dried
 over magnesium sulfate and evaporated. The residue
 was purified by flash chromatography (EtOAc: hexane =
 1: 20 to 1:10) to give 849 mg (95% yield) of the title
 15 compound as a solid; ¹H-NMR (CDCl₃) δ 1.37 (s, 6 H),
 3.93 (s, 1 H), 5.15 (s, 2 H), 6.24 (s, 1 H), 6.90-
 6.94 (m, 2 H), 7.25 (d over CHCl₃, J = 8.8 Hz, 1 H),
 7.53 (d, J = 8.7 Hz, 2 H), 8.08 (d, J = 8.7 Hz, 2 H);
 MS m/e 457 (MH⁺).


20

Anal. calcd. for C₂₁H₁₉F₃O₆S: C, 55.26; H, 4.20.
 Found: C, 55.39; H, 4.08.

25

EXAMPLE 136

Methyl 4-(1,1-dimethyl-3-phenyl-1H-inden-5-
yloxymethyl)benzoate




Methyl 4-[(1,1-dimethyl-3-
 (trifluoromethanesulfonyloxy)-1H-indene-5-
 5 yloxymethyl]benzoate (838 mg, 1.84 mmol),
 tris(dibenzylideneacetone) dipalladium (0) (17 mg,
 0.02 mmol), triphenylarsine (64 mg, 0.20 mmol),
 lithium chloride (234 mg, 5.51 mmol) and
 phenyltributyltin (676 mg, 1.84 mmol) were stirred at
 10 60 °C for 18 hours. The mixture was diluted with water
 (30 mL), extracted with ethyl ether (30 mL X 3). The
 combined extracts were dried over magnesium sulfate
 and evaporated. The residue was purified by flash
 chromatography (EtOAc: hexane = 1: 25 to 1: 10) to
 15 give a product which recrystallized from EtOAc-hexane
 (1:20 to 1:10) to give 454 mg (64 % yield) of the
 title compound as crystals; ¹H-NMR (CDCl₃) δ 1.38 (s, 6
 H), 3.93 (s, 1 H), 5.13 (s, 2 H), 6.44 (s, 1 H), 6.86
 (dd, J = 2.2, 8.1 Hz, 1 H), 7.11 (d, J = 2.2 Hz, 1 H),
 20 7.27-7.55 (m, 8 H), 8.06 (d, J = 8.1 Hz, 2 H); MS m/e
 385 (MH⁺).

Anal. calcd. for C₂₆H₂₄O₃ · 0.125 H₂O: C, 80.75; H, 6.32.
 Found: C, 80.76; H, 6.19.

25

EXAMPLE 137

4-[(1,1-Dimethyl-3-phenyl-1H-inden-5-yloxymethyl)-
 30 benzoic acid (I^{32a})




Methyl 4-(1,1-dimethyl-3-phenyl-1H-inden-5-yloxymethyl)benzoate (300 mg, 0.78 mmol) and 3.9 mL of 2 N NaOH in 5 mL of tetrahydrofuran and 5 mL of methanol were stirred at room temperature for 16 hours. The mixture was concentrated and acidified with 1 N HCl (10 mL), extracted with ethyl acetate (20 mL X 3). The combined extracts were washed with water (10 mL), dried over magnesium sulfate, and evaporated. The residue was crystallized from ether-hexane to give 247 mg (85% yield) of the title compound as white crystals; $^1\text{H-NMR}$ (DMSO- d_6) δ 1.31 (s, 6 H), 5.20 (s, 2 H), 6.57 (s, 1 H), 6.89 (dd, $J = 1.1, 8.2$ Hz, 1 H), 7.02 (d, $J = 1.1$ Hz, 1 H), 7.36-7.57 (m, 8 H), 7.95 (d, $J = 8.3$ Hz, 2 H), 12.94 (bs, 1 H); MS m/e 371 (MH^+).

Anal. calcd. for $\text{C}_{25}\text{H}_{22}\text{O}_3 \cdot 0.25 \text{ H}_2\text{O}$: C, 80.08; H, 6.05.
Found: C, 80.17; H, 5.86.

EXAMPLE 138

6-Bromo-3,3-dimethyl-indan-1-one (LXXIV)




To aluminum chloride (4.16 g, 31.2 mmol) in a flask was added 3,3-dimethyl-1-indan-1-one (2.00 g, 12.5 mmol) at 90-100 °C. After stirring for 15 min, bromine (2.39 g, 15.0 mmol) was slowly added. The mixture was stirred at 100 °C for 1 hour and then quenched with ice-water (200 g), extracted with ethyl acetate (40 mL X 3). The combined extracts were dried over magnesium sulfate and evaporated. The residue was purified by flash chromatography (EtOAc: hexane = 1: 20 to 1:15) to give a crude product which recrystallized from MeOH-EtOAc to give 1.23 g (41 % yield) of the title compound as colorless crystals; ¹H-NMR (CDCl₃) δ 1.42 (s, 6 H), 2.61 (s, 2 H), 7.39 (d, J = 8.2 Hz, 1 H), 7.72 (dd, J = 2.0, 8.2 Hz, 1H), 7.82 (d, J = 2.0 Hz, 1 H); MS m/e 239 (MH⁺).

Anal. calcd. for C₁₁H₁₁BrO₃: C, 55.26; H, 4.64.
Found: C, 55.19; H, 4.61.

EXAMPLE 139

Methyl 4-[2-(1,1-dimethyl-3-oxo-indan-5-yl)vinyl]-benzoate (LXXVa)




6-Bromo-3,3-dimethyl-indan-1-one (1.20 g, 5.02 mmol), methyl 4-vinylbenzoate (1.63 g, 10.0 mmol), palladium acetate (56 mg, 0.25 mmol), tetrabutylammonium chloride hydrate (1.53 g, 5.52 mmol), and sodium bicarbonate (1.05 g, 12.6 mmol) were stirred in 10 mL of anhydrous N,N-dimethylformamide at 80-100 °C for 8 hours. The mixture was diluted with water (100 mL),
 10 extracted with methylene chloride (50 mL X 3). The combined extracts were dried over magnesium sulfate and evaporated. The residue was purified by flash chromatography (CH₂Cl₂: hexane = 1: 1 to 1:0, then CH₂Cl₂:EtOAc = 10:1) to give 1.12 g (70 % yield) of the
 15 title compound as a yellow solid; ¹H-NMR (CDCl₃) δ 1.45 (s, 6 H), 2.64 (s, 2 H), 5.30 (s, 3 H), 7.18 (d, J = 16.5 Hz, 1 H), 7.26 (d, J = 16.5 Hz, 1 H), 7.52 (d, J = 8.1 Hz, 1 H), 7.58 (d, J = 8.5 Hz, 2 H), 7.78 (d, J = 1.7, 8.1 Hz, 1 H), 7.86 (d, J = 1.7 Hz, 1 H), 8.05
 20 (d, J = 8.5 Hz, 2 H); MS m/e 321 (MH⁺).

Anal. calcd. for C₂₁H₂₀O₃: C, 78.18; H, 6.32.
 Found: C, 78.22; H, 6.26.

25

EXAMPLE 140

Methyl [2-[1,1-dimethyl-3-
(trifluoromethanesulfonyloxy)-1H-indene-5-
 30 yl]vinyl]benzoate



To a solution of methyl 4-[2-(1,1-dimethyl-3-oxo-indan-5-yl)vinyl]benzoate (450 mg, 1.40 mmol) and 2,6-
 5 di-tert-butyl-4-methylpyridine (345 mg, 1.68 mmol) in
 5 mL of methylene chloride was added triflic acid
 anhydride (434 mg, 1.54 mmol) at -78 °C. After
 stirring at room temperature for 16 hours, the mixture
 was diluted with 1 N HCl (20 mL), extracted with ethyl
 10 ether (20 mL X 3). The combined extracts were dried
 over magnesium sulfate and evaporated. The residue
 was purified by flash chromatography (CH₂Cl₂; hexane =
 1: 2 to 1:1) to give 606 mg (95% yield) of the title
 compound as a solid; ¹H-NMR (CDCl₃) δ 1.41 (s, 6 H),
 15 3.94 (s, 3 H), 6.27 (s, 1 H), 7.15 (d, J = 16.3 Hz, 1
 H), 7.27 (d, J = 16.3 Hz, 1 H), 7.37 (d, J = 8.2 Hz, 1
 H), 7.49 and 7.50 (d over s, J = 8.2 Hz, 2 H), 7.59
 (d, J = 8.4 Hz, 2 H), 8.05 (d, J = 8.4 Hz, 2 H); (MS
 m/e 453 (MH⁺)).


20

Anal. calcd. for C₂₂H₁₉F₃O₅S: C, 58.40; H, 4.23.
 Found: C, 58.38; H, 4.00.

25

EXAMPLE 141


Methyl 4-[2-(1,1-dimethyl-3-phenyl-1H-inden-5-yl)-
vinyl]benzoate



- Methyl [2-[1,1-dimethyl-3-(trifluoromethanesulfonyloxy)-1H-indene-5-yl]vinyl]benzoate (590 mg, 1.30 mmol), tris(dibenzylideneacetone)dipalladium (0) (48 mg, 0.05 mmol), triphenylarsine (64 mg, 0.20 mmol), lithium chloride (166 mg, 3.91 mmol) and phenyltributyltin (525 mg, 1.43 mmol) were stirred at 95 °C for 16 hours.
- The mixture was diluted with water (75 mL), extracted with ethyl acetate (20 mL X 2) and ethyl ether (20 mL X 2). The combined extracts were dried over magnesium sulfate and evaporated. The residue was purified by flash chromatography (EtOAc: hexane = 1: 25 to 1: 10) to give a product which recrystallized from EtOAc-hexane to give 311 mg (63 % yield) of the title compound as yellow solids; ¹H NMR (CDCl₃) δ 1.42 (s, 6 H), 3.93 (s, 3 H), 6.45 (s, 1 H), 7.11 (d, J = 16.3 Hz, 1 H), 7.29 (d, J = 16.3 Hz, 1 H), 7.41-7.65 (m, 10 H), 8.02 (d, J = 8.4 Hz, 2 H).

EXAMPLE 142

- 4-[2-(1,1-dimethyl-3-phenyl-1H-inden-5-yl)vinyl]benzoic acid (I^{33a})




Methyl 4-[2-(1,1-dimethyl-3-phenyl-1H-inden-5-yl)-
vinyl]benzoate (294 mg, 0.77 mmol) and 0.77 mL of 10 N
5 NaOH in 7 mL of tetrahydrofuran and 5 mL of methanol
were stirred at reflux for 1 hour. The mixture was
concentrated and acidified with 1 N HCl (10 mL),
extracted with ethyl acetate (30 mL X 2). The
combined extracts were washed with water (10 mL),
10 dried over magnesium sulfate, and evaporated. The
residue was triturated in hot methanol to give 276 mg
(90 % yield) of the title compound which contained a
molecule of methanol; ¹H-NMR (DMSO-d₆) δ 1.34 (s, 6 H),
3.14 (s, 3 H), 6.58 (s, 1 H), 7.25 (d, J = 16.4 Hz, 1
15 H), 7.37-7.71 (m, 11H), 7.90 (d, J = 8.3 Hz, 2 H); MS
m/e 367 (MH⁺).

Anal. calcd. for C₂₆H₂₆O₃ · CH₃OH: C, 81.38; H, 6.58.
Found: C, 81.18; H, 6.78.

20

EXAMPLE 143

N-(2-pyridyl)triflimide




The title compound was prepared by procedure of Comins & Dehghani, *Tetrahedron Lett.*, Vol. 33, No. 42, 1992, p. 6299. ^1H NMR (CDCl_3): δ 8.66 (m, 1H), 7.95 (m, 1H), 7.56 (m, 1H), 7.48 (d, 1H). MS (DCI) m/e : 359 (MH^+).

Example 144

10

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]amino]-2-hydroxybenzoic acid, phenyl ester (I^3j)



15

Using the method given for the preparation of the 8-(2-fluorophenyl) derivative I^3g , 230 mg (0.83 mmol) of compound XVIIIa and phenyl 4-aminosalicylate (Aldrich; 190 mg, 0.83 mmol) gave 193 mg (Y: 48%) of the title product; ^1H -NMR (CDCl_3): δ 8.03 (d, $J=9.3$ Hz, 1H),


20

7.75-7.71 (m, 2H), 7.51-7.19 (m, 13H), 6.09 (t, J=4.5 Hz, 1H), 2.40 (d, J=4.5 Hz, 2H), 1.38 (s, 6H); MS (DCI) m/e: 490 (MH⁺).

5

Example 145

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]aminol-2-hydroxybenzoic acid


10 (I^{4j})

Using the method for the preparation of the 8-phenyl derivative I^{4a}, 193 mg (0.39 mmol) of compound I^{3j} gave 135 mg (Y: 83%) of the title compound; ¹H-NMR (DMSO-d₆): δ 11.33 (bs, 1H), 10.38 (s, 1H), 7.83 (dd, J=8.0, 1.7 Hz, 1H), 7.71 (d, J=8.7 Hz, 1H), 7.54 (d, J=8.0 Hz, 1H), 7.45-7.29 (m, 7H), 7.24 (dd, J=8.7, 1.8 Hz, 1H), 6.06 (t, J=4.5 Hz, 1H), 2.35 (d, J=4.5 Hz, 2H), 1.32 (s, 6H); MS (DCI) m/e: 400 (MH⁺); IR (KBr): 2960, 1668, 1600, 1508.

Anal. calcd. for C₂₆H₂₃N₁O₄ · 0.5 H₂O: C, 73.92, H, 5.73; N, 3.32. Found: C, 73.91; H, 5.73; N, 2.99.

25


Example 146

2-Fluoro-4-nitrobenzoic acid

5

A solution of 2-fluoro-4-nitrotoluene (3.95 g, 25.5 mmol, Janssen) in water (200 mL) was treated portionwise with potassium permanganate (24.6 g, 0.15 mmol) at 80°C. After 2h the reaction mixture was
10 filtered hot, acidified with 5N hydrochloric acid and extracted with ethyl acetate (2 x 100 mL). The extract was dried over anhydrous sodium sulfate and concentrated in vacuo to give 1.15 g (Y: 24%) of the title compound; ¹H-NMR (DMSO-d₆): δ 8.21 (d, J=10.5 Hz,
15 1H), 8.16-8.08 (m, 2H); MS (DCI) m/e: 186 (MH⁺).

Example 147


20 2-Fluoro-4-nitrobenzoic acid ethyl ester

To a solution of 2-fluoro-4-nitrobenzoic acid (1.15 g, 6.22 mmol) in absolute ethyl alcohol (20 mL) was added
25 p-toluenesulfonic acid monohydrate (100 mg). After 16

h at reflux, the reaction mixture was concentrated in vacuo and the residue chromatographed on silica gel (eluted with 5% ethyl acetate in hexane) to give 770 mg (Y: 58%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 8.16-8.00 (m, 3H), 4.45 (q, $J=7.1$ Hz, 2H), 1.43 (t, $J=7.1$ Hz, 3H). MS (DCI) m/e : 214 (MH^+).

Example 148

10


2-Fluoro-4-aminobenzoic acid ethyl ester

A solution of 2-fluoro-4-nitrobenzoic acid ethyl ester (770 mg, 3.62 mmol) in methyl alcohol (15 mL) was treated with platinum (IV) oxide (75 mg) at 40 psi of H_2 . After 0.5 h the reaction mixture was filtered through celite and concentrated in vacuo to give 700 mg (Y: 99%) of the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.76 (t, $J=8.3$ Hz, 1H), 6.42 (d, $J=8.5$ Hz, 1H), 6.35 (d, $J=12.9$ Hz, 1H), 4.33 (q, $J=7.1$ Hz, 2H), 4.20 (bs, 2H), 1.36 (t, $J=7.1$ Hz, 3H); MS (DCI) m/e : 184 (MH^+).

25

Example 149

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]amino]-2-fluorobenzoic acid ethyl ester (I^3k)




Using the method given for the preparation of the 8-(2-fluorophenyl) derivative I³g, 250 mg (0.90 mmol) of compound XVIIIa and 2-fluoro-4-aminobenzoic acid ethyl ester (181 mg, 0.99 mmol) gave 240 mg (Y: 60%) of the title product; ¹H-NMR (CDCl₃): δ 7.91 (t, J=8.5 Hz, 1H), 7.72 (d, J=10.1 Hz, 1H), 7.66 (d, J=1.8 Hz, 1H), 7.51-7.48(m, 1H), 7.44-7.35 (m, 6H), 7.20 (d, J=8.6 Hz, 1H), 6.08 (t, J=4.5 Hz, 1H), 4.37 (q, J=7.1 Hz, 2H), 2.40 (d, J=4.5 Hz, 2H), 1.38 (t, J=7.1 Hz, 3H), 1.37 (s, 6H); MS (DCI) m/e: 444 (MH⁺).

15

Example 150

4-[[[5,6-Dihydro-5,5-dimethyl-8-phenyl]-2-naphthalenyl]carbonylamino]-2-fluorobenzoic acid (I⁴k)



Using the method given for the preparation of the 8-phenyl derivative I⁴a, 240 mg (0.54 mmol) of compound I³k gave 175 mg (Y: 78%) of the title compound; ¹H-NMR

5 (DMSO-d₆): δ 12.98 (bs, 1H), 10.58 (s, 1H), 7.89-7.77 (m, 2H), 7.73 (d, J=1.5 Hz, 1H), 7.54 (d, J=8.1 Hz, 1H), 7.44-7.30 (m, 7H), 6.06 (t, J=4.5 Hz, 1H), 2.34 (d, J=4.5 Hz, 2H), 1.31 (s, 6H). MS (DCI) m/e: 416 (MH⁺); IR (KBr): 2960, 1692, 1596, 1526.


10

Anal. calcd for C₂₆H₂₂N₁O₃F₁ · 0.25 H₂O: C, 74.36; H, 5.40; N, 3.34. Found: C, 74.16; H, 5.74; N, 3.13.

15

Example 151

N-(4-Methyl-3-nitro-phenyl)acetamide




20 A solution of 4-methyl-3-nitroaniline (3.60 g, 23.7

mmol) in acetic anhydride (28 mL) was allowed to stir at room temperature. After 16 h, the reaction mixture was concentrated in vacuo, diluted with ethyl acetate (25 mL), washed with saturated sodium bicarbonate (2 x 5 50 mL), dried over anhydrous sodium sulfate and concentrated in vacuo to give 4.20 g (Y: 99%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 8.10 (d, $J=1.7$ Hz, 1H), 7.75 (dd, $J=8.5$, 1.7 Hz, 1H), 7.38 (bs, 1H), 7.29 (d, $J=8.5$ Hz, 1H), 2.56 (s, 3H), 2.21 (s, 3H); MS 10 (DCI) m/e : 195 (MH^+).

Example 152

15 4-Acetylamino-2-nitrobenzoic acid




A solution of N-(4-methyl-3-nitro-phenyl)acetamide (4.20 g, 23.6 mmol) in water (200 mL) was treated 20 portionwise with potassium permanganate (22.77 g, 0.144 mmol) at 80°C. After 2 h the reaction mixture was filtered hot, acidified with 5N hydrochloric acid and extracted with ethyl acetate (2 x 100 mL). The extract was dried over anhydrous sodium sulfate and 25 concentrated in vacuo to give 1.80 g (Y: 34%) of the title compound; $^1\text{H-NMR}$ (CDCl_3): δ 10.63 (s, 1H), 8.19 (d, $J=1.7$ Hz, 1H), 7.85 (d, $J=8.5$ Hz, 1H), 7.78 (dd, $J=8.5$, 1.7 Hz, 1H), 2.10 (s, 3H); MS (DCI) m/e : 225 (MH^+).

Example 153

4-Amino-2-nitrobenzoic acid ethyl ester

5




A solution of 4-acetylamino-2-nitrobenzoic acid (1.80 g, 8.0 mmol) in 12 N hydrochloric acid (14 mL) and absolute ethyl alcohol (10 mL) was heated to 90-100°C for 5 h. Concentrated down in vacuo to remove ethyl alcohol only, adjusted pH to 4 with 1N sodium hydroxide and filtered off precipitate to give 140 mg (Y: 8%) of the title compound; ¹H-NMR (CDCl₃): δ 7.61 (d, J=8.5 Hz, 1H), 6.82 (d, J=2.1 Hz, 1H), 6.74 (dd, J=8.5, 2.1 Hz, 1H), 6.52 (bs, 2H), 4.17 (q, J=7.1 Hz, 2H), 1.21 (t, J=7.1 Hz, 3H); MS (DCI) m/e: 211 (MH⁺).

Example 154

20

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]amino]-2-nitrobenzoic acid ethyl ester (I³¹)




Using the method given for the preparation of the 8-(2-fluorophenyl) derivative I³g, 199 mg (0.71 mmol) of compound XVIIIa and 140 mg (0.78 mmol) of 4-amino-2-nitrobenzoic acid ethyl ester gave 155 mg (Y: 46%) of the title product; ¹H-NMR (CDCl₃): δ 8.15 (s, 1H), 7.84 (s, 1H), 7.79 (s, 1H), 7.73 (dd, J=8.1, 1.9 Hz, 1H), 7.52-7.49 (m, 2H), 7.41-7.26 (m, 5H), 6.09 (t, J=4.5 Hz, 1H), 4.36 (q, J=7.2 Hz, 2H), 2.40 (d, J=4.5 Hz, 2H), 1.37 (s, 6H), 1.34 (t, J=7.2 Hz, 3H); MS (DCI) m/e: 471 (MH⁺).

Example 155

15

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]amino]-2-nitrobenzoic acid (I⁴l)



Using the method given for the preparation of the 8-phenyl derivative I⁴a, 155 mg (0.33 mmol) of compound I³l gave 105 mg (Y: 72%) of the title compound; ¹H-NMR (DMSO-d₆): δ 13.65 (bs, 1H), 10.76 (s, 1H), 8.31 (d, J=1.7 Hz, 1H), 7.99 (dd, J=8.3, 1.7 Hz, 1H), 7.90-7.84 (m, 2H), 7.55 (d, J=8.3 Hz, 1H), 7.47 (d, J=1.7 Hz, 1H), 7.44-7.30 (m, 5H), 6.07 (t, J=4.5 Hz, 1H), 2.33 (d, J=4.5 Hz, 2H), 1.31 (s, 6H); MS (DCI) m/e: 443 (MH⁺); IR (KBr): 2960, 1702, 1542, 1518.

10


Anal. calcd. for C₂₆H₂₂N₂O₅ · 0.5 H₂O: C, 69.17, H, 5.14, N, 6.20. Found: C, 69.55, H, 4.93, N, 5.94.

15

Example 156

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]amino-2-methoxybenzoic acid, methyl ester (I³m)

20




Using the method for the preparation of the 8-(2-fluorophenyl) derivative I³g, 415 mg (1.49 mmol) of compound XVIIIa and 297 mg (1.64 mmol) of methyl 4-amino-2-methoxybenzoate (Apin) gave 570 mg (Y: 90%) of

the title product; $^1\text{H-NMR}$ (CDCl_3): δ 7.82 (d, $J=8.5$ Hz, 1H), 7.77 (d, $J=1.7$ Hz, 1H), 7.73 (s, 1H), 7.70 (d, $J=1.9$ Hz, 1H), 7.51 (m, 1H), 7.48 (s, 1H), 7.41-7.33 (m, 4H), 6.80 (dd, $J=8.5, 1.9$ Hz, 1H), 6.08 (t, $J=4.5$ Hz, 1H), 3.92 (s, 3H), 3.87 (s, 3H), 2.39 (d, $J=4.5$ Hz, 2H), 1.38 (s, 6H); MS (DCI) m/e : 442 (MH^+).

Example 157

10

4-[[[(5,6-Dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]amino]-2-methoxybenzoic acid
(I^4m)



15

Using the method given for the preparation of the 8-phenyl derivative I^4a , 135mg (0.31 mmol) of compound I^3m gave 100 mg (Y: 77%) of the title compound; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$): δ 12.32 (bs, 1H), 10.36 (s, 1H), 7.85 (dd, $J=8.1, 1.7$ Hz, 1H), 7.79 (s, 1H), 7.64 (d, $J=1.7$ Hz, 1H), 7.53 (d, $J=8.1$ Hz, 1H), 7.44-7.30 (m, 7H), 6.06 (t, $J=4.5$ Hz, 1H), 3.76 (s, 3H), 2.34 (d, $J=4.5$ Hz, 2H), 1.31 (s, 6H); MS (DCI) m/e : 428 (MH^+); IR (KBr): 2960, 1718, 1592, 1524.


25

Anal. calcd. for $C_{27}H_{25}N_1O_4 \cdot 0.25 H_2O$: C, 75.07; H, 5.95; N, 3.24. Found: C, 75.04; H, 5.86; N, 3.04.

5

Example 158

4-[[[(5,6-Dihydro-5,5-dimethyl-8-(2-naphthalene)]-2-naphthalenyl]carbonyl]amino]benzoic acid (I⁴ⁿ)



10


Starting from compound XVIa the title compound was made from a method analogous to the preparation of the 8-(2-fluorophenyl) derivative I^{4g}; ¹H-NMR (DMSO-d₆): δ 12.69 (s, 1H), 10.41 (s, 1H), 7.95-7.88 (m, 5H), 7.85 (d, J=8.7 Hz, 2H), 7.77 (d, J=8.7 Hz, 2H), 7.57-7.40 (m, 5H), 6.20 (t, J=4.5 Hz, 1H), 2.40 (d, J=4.5 Hz, 2H), 1.35 (s, 6H); MS (DCI) m/e: 448 (MH⁺); IR (KBr): 2960, 1686, 1596, 1518.

20 Anal. calcd. for $C_{30}H_{25}O_3N_1 \cdot 0.90 H_2O$: C, 77.70; H, 5.83; N, 3.02. Found: C, 78.10; H, 5.55; N, 3.13.

Example 159

4-[[[(5,6-Dihydro-8-phenyl-2-naphthalenyl)carbonyl]aminolbenzoic acid (I^{4o})

5



Starting from 6-methoxy tetralone the title compound was made from a method analogous to the preparation of the 8-(2-fluorophenyl) derivative I^{4g}; ¹H-NMR (DMSO-d₆): δ 12.73 (s, 1H), 10.42 (s, 1H), 7.91-7.81 (m, 5H), 7.55-7.25 (m, 7 H), 6.18 (t, J=4.5 Hz, 1H), 2.86 (t, J=7.6 Hz, 2H), 2.41-2.34 (m, 2H); MS (DCI) m/e: 370 (MH⁺); IR (KBr): 2950, 1680, 1648, 1518.


15

Anal. calcd. for C₂₄H₁₉N₁O₃ : C, 78.03; H, 5.18; N, 3.79. Found: C, 77.61; H, 5.14; N, 3.81.

Example 160

20

Methyl 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)carbonyl]amino]-3-fluorobenzoate (I^{3p})




Using the method given for the preparation of the 8-(2-fluorophenyl) derivative I³g, compound XVIIIa (300
 5 mg) and methyl 3-fluoro-4-aminobenate (245 mg) gave 320 mg (69% yield) of the title compound as a glassy mass; ¹H-NMR (CDCl₃) δ 1.38 (s, 6 H), 2.40 (d, J = 4.7 Hz, 2 H), 3.90 (s, 3 H), 6.08 (t, J = 4.7 Hz, 1 H), 7.32-7.42 (m, 5 H), 7.49 (d, J = 8.0 Hz, 1 H), 7.56
 10 (d, J = 1.9 Hz, 1 H), 7.73 (d, J = 1.9, 8.0 Hz, 1 H), 7.74 (d, J = 1.8, 11.5 Hz, 1 H), 7.84 (dd, J = 1.8, 8.5 Hz, 1 H), 8.11 (bd, J = 3.8 Hz, 1 H), 8.55 (t, J = 8.5 Hz, 1 H).

15

Example 161

4-[[5,6-Dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl]carbonylamino]-3-fluorobenzoic acid (I⁴p)



Using the same method given for the preparation of the 8-phenyl derivative I⁴a, 310 mg of compound I³p gave 170 mg (54% yield) of the title compound; ¹H-NMR(CDCl₃)

5 δ 1.23 (s, 6 H), 2.39 (d, J = 4.7 Hz, 2 H), 6.07 (t, J = 4.7 Hz, 1 H), 7.30-7.42 (m, 5 H), 7.49 (d, J = 8.1 Hz, 1 H), 7.52 (d, J = 2.0 Hz, 1 H), 7.72 (d, J = 2.0, 8.1 Hz, 1 H), 7.79 (d, J = 1.8, 11.4 Hz, 1 H), 7.90 (d, J = 8.5 Hz, 1 H), 8.06 (bd, J = 4.2 Hz, 1 H), 8.59 (t,


10 J = 8.6 Hz, 1 H); MS m/e 416 (MH⁺).

Anal. Calcd. for C₂₆H₂₂FNO₃: C, 75.17; H, 5.34; N, 3.37. Found: C, 74.96; H, 5.53; N, 3.33.

15

Example 162

Methyl 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)carbonyl]amino]-3-methylbenzoate (I³g)




Using the method given for the preparation of the 8-(2-fluorophenyl) derivative I³g, compound XVIIIa (300 mg) and methyl 3-methyl-4-aminobenzoate (319 mg) gave 305 mg (66 % yield) of the title compound as a glassy mass; ¹H-NMR (CDCl₃) δ 1.39 (s, 6 H), 2.20 (s, 3 H), 2.41 (d, J = 4.7 Hz, 2 H), 3.89 (s, 3 H), 6.09 (t, J = 4.7 Hz, 1 H), 7.30-7.45 (m, 5 H), 7.48 (d, J = 2.0 Hz, 1 H), 7.51 (d, J = 8.0 Hz, 1 H), 7.68 (s, 1 H), 7.79 (d, J = 2.0, 8.0 Hz, 1 H), 7.85 (s, 1 H), 7.91 (dd, J = 2.0, 8.5 Hz, 1 H), 8.31 (d, J = 8.5 Hz, 1 H); MS m/e 426 (MH⁺).

15 Anal. Calcd. for C₂₈H₂₇NO₃·0.125 H₂O: C, 78.61; H, 6.42; N, 3.27. Found: C, 78.51; H, 6.46; N, 3.26.

Example 163

20 4-[[[5,6-Dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl]carbonyl]amino]-3-methylbenzoic acid (I⁴g)



Using the same method given for the preparation of the 8-phenyl derivative I4a, 280 mg of compound I³q gave
 5 223 mg (82 % yield) of the title compound; ¹H-NMR (CDCl₃) δ 1.37 (s, 6 H), 2.20 (s, 3 H), 2.39 (d, J = 4.7 Hz, 2 H), 6.07 (t, J = 4.7 Hz, 1 H), 7.30-7.42 (m, 5 H), 7.45 (d, J = 2.0 Hz, 1 H), 7.51 (d, J = 8.1 Hz, 1 H), 7.68 (s, 1 H), 7.78 (d, J = 2.0, 8.1 Hz, 1 H),
 10 7.80 (s, 1 H), 7.96 (dd, J = 1.9, 8.6 Hz, 1 H), 8.36 (d, J = 8.6 Hz, 1 H); MS m/e 412 (MH⁺).

Anal. Calcd. for C₂₇H₂₅NO₃: C, 78.81; H, 6.12; N, 3.40.
 Found: C, 78.68; H, 6.12; N, 3.40.

15

This invention is further illustrated by the following biological tests, which are illustrative only.

20 Rhino Mouse Study

Representatives from compounds of formula I were tested for their effect on utriculi reduction on rhino mouse and directly compared to all-trans retinoic
 25 acid.

Rhino mouse utriculi reduction assay

Six to nine week old female hairless rhino mice (hr^{rh}/hr^{rh}) were produced in the Bristol-Myers Squibb colony. Test retinoids in ethanol vehicle (50ul) were applied to the dorsal area (approximately 1.5 x 3 cm²) of rhino mice once daily for 5 days (Monday to Friday). For various retinoids, a dose response was obtained with concentrations ranging from 0.00033 mM to 16.5 mM. The animals were sacrificed on the following Monday by CO₂ inhalation. A 7/8" full thickness punch was taken from the central dorsal area of each animal. The epidermis of the biopsy was removed from the dermis after incubation in 0.5% acetic acid overnight at 4°C. The separated epidermis was then fixed in formalin, dehydrated with ethanol, and cleared in xylene. To determine the utriculi diameter, each epidermis sheet was placed on a glass slide in xylene. For each specimen, the diameter of 40 utricules was measured with an image analysis system (IBM PC, Image Measure program and Olympus microscope with video camera). % Utriculi reduction was calculated as

$$\left(1 - \frac{\text{utriculi diameter in the test group}}{\text{utriculi diameter in the ethanol control group}} \right) \times 100\%$$

Since the maximum effect in this assay is approximately 60% utriculi reduction, the activity of various test compounds is reported as ED₃₀ in Table 1, the concentration to reach 30% (half-maximum) utriculi

reduction.

Table 1

Compound*	ED ₃₀ (mM)
I ⁴ a	1.25
I ⁴ d	0.931
I ⁴ e	0.123
I ⁴ f	0.038
I ³⁰ a	2.35
I ³³ a	0.86
I ⁴ k	3.12

* The following compounds were not active in this Rhino mouse model: I²c, I¹⁰a, I¹¹a, I¹⁴a, I¹⁵a, I¹²a, I¹p and I¹q

The following biological test indicates that the compounds of the instant invention possess cytotoxicity activity normally associated with retinoids. Thus in one aspect, the invention provides a method of treating various tumors.

Cytotoxicity Result

The cytotoxicity assay was set up similar to those run by the National Cancer Institute (D.A. Scudiero, et al, "Evaluation of a Soluble Tetrazolium/Formazan Assay for Cell Growth and Drug Sensitivity in Culture Using Human and Other Tumor Cell Lines", Cancer

Research, 48, 4827-4833, September 1, 1988; M. C. Alley, et al, "Feasibility of Drug Screening with Panels of Human Tumor Cell Lines Using a Microculture Tetrazolium Assay", *Cancer Research*, 48, 589-601, February 1, 1988) with the exception that the new vital stain alamarBlue™ was used to determine cellular viability. Briefly, the assay involved plating 1000 cells per well in a volume of 120 μ L in a 96 well flat-bottom dish (Corning) on day -1. Twenty-four hours later the appropriate dilution of a compound of formula I was added in a volume of 30 μ L complete medium (final volume 150 μ L). The plates were sealed with a plate sealer (Dynatech Labs) to prevent evaporation. On day 5 the mylar film was removed and 15 μ L of sterile alamarBlue was added to each well and the cells were incubated 37°C 5% CO₂ for two hours. Using a Vmax plate reader the optical density for each well was determined having the OD₅₇₀ subtracted from the OD₆₀₀. The 100% signal was determined for cells grown in complete medium containing only 0.5% DMSO. All wells were set-up in triplicate and the mean values were plotted in Figures 1, 2 and 3. The IC₅₀ values were determined for the second experiment and are listed in table 2.

TABLE 2

IC₅₀ values for L2987 experiment 2

5	Compound	IC ₅₀ (μM)
	All trans retinoic acid	68
	I ⁴ a	51
	I ¹⁰ a	47
	I ⁴ g	38
10	I ⁴ d	28
	I ³⁰ a	60
	I ²⁶ a	48
	I ¹² a	>100
15	I ⁶ a	28

Antiarthritic Test

Compound I¹¹a was also evaluated in a classical animal
 20 model for rheumatoid arthritis, i.e. in the collagen
 induced arthritis (CIA) model as described by Trentham
 et al in Collagen arthritis as a relevant model for
 rheumatoid arthritis. Evidence pro and con, *Arthritis*
Rheum 25: 911-916. (1982); and Autoimmunity to type II
 25 collagen: an experimental model of arthritis. *J. Exp.*
Med 146: 857-868 (1977).

In our protocol, drug treatment began three days prior
 to the first intradermal injection of 100 μg of type

II collagen in Freund's complete adjuvant. At day 7, a second booster injection of collagen was given. Typically, animals begin to manifest symptoms of arthritis, such as joint swelling, around day 30.

- 5 Daily i.p. administration of compound I¹¹a at a concentration of 10 mg/kg and 30 mg/kg in a volume of 10 ml/kg were given. Clinical score by visual assessment of three parameters (swelling, joint deformation, and ankylosis) was performed daily to
10 determine the onset of symptoms (see Figure 4). At the end of the experiment period (day 70), animals were sacrificed and the joints were evaluated for histological changes. These changes were graded as 1 for synovial hypertrophy, 2 for pannus erosion into
15 cartilage, 3 for pannus erosion into bone, and 4 for loss of joint integrity.

As shown in Figure 4, the mean clinical score per animal remains normal until about day 30. In vehicle
20 treated animals, the clinical score increases linearly to about 2.5 (maximum of 4.0) at day 65 indicating well established arthritis in these animals. By contrast, animals treated with compound I¹¹a showed a marked reduction in the mean clinical score over the
25 same time period. At day 65, the mean score was 0.6 with a small number of animals exhibiting joint swelling (grade 1) as the only obvious clinical symptom. However, most of the limbs evaluated were normal in appearance.

30

Preliminary data in this model with 30 mg/kg showed complete absence of disease at the visual and histological levels.

The compounds of formula I may be used topically or systemically, as anticancer agents and in the treatment, amelioration, or prevention of the skin disorders for which retinoic acid and other retinoids are useful. In this regard, they may be used for therapy in animals, including humans, for premalignant epithelial cell lesions, as a prophylaxis against tumor promotion in epithelial cells and treatment for dermatoses such as ichthyoses, follicular disorders, benign epithelial disorders, and other proliferative skin diseases (nonmalignant conditions of the skin that are characterized by epidermal cell proliferation or incomplete cell differentiation) such as acne, psoriasis, eczema, atopic dermatitis, nonspecific dermatitis and the like. The compounds of formula I may also be used in reversing and preventing the effects of irradiation damage to skin. The compounds of formula I may also be used to treat rheumatic illnesses, including but not limited to, rheumatoid arthritis, psoriatic arthropathy, Reiter's disease, and cutaneous lupus erythematosus. The compounds of formula I have an additional use in the treatment of osteoarthritis.

When used for the above treatments they will usually be formulated with a pharmaceutically acceptable liquid, semi-solid, or solid carrier. A pharmaceutically acceptable carrier is a material that is nontoxic and generally inert and does not affect the functionality of the active ingredients adversely. Such materials are well known and include those materials sometimes referred to as diluents or vehicles (excipients) in the pharmaceutical formulation art. The carrier may be organic or

- inorganic in nature. Examples of pharmaceutically acceptable carriers that may be used to formulate a compound of formula I are water, gelatin, lactose, starch, mineral oil, cocoa butter, dextrose, sucrose, sorbitol, mannitol, gum acacia, alginates, cellulose, talc, magnesium stearate, polyoxyethylene sorbitan monolaurate, and other commonly used pharmaceutical carriers. In addition to a compound of formula I and carrier, the formulation may contain minor amounts of additives such as flavoring agents, coloring agents, thickening or gelling agents, emulsifiers, wetting agents, buffers, stabilizers, and preservatives such as antioxidants.
- The dosages and dosage regimen in which the compounds of formula I are administered will vary according to the dosage form, mode of administration, the condition being treated and particulars of the patient being treated. Accordingly, optimal therapeutic concentrations will be best determined at the time and place through routine experimentation.

In the treatment of dermatoses, it will generally be preferred to administer the drug topically, though in certain cases such as treatment of severe cystic acne, oral administration may also be used. If the compounds according to the invention are used topically, it will be found that they exhibit a good activity over a very broad range of dilution; in particular, concentrations of the active compound or compounds ranging from 0.0005% to 2% by weight can generally be used. It is of course possible to use higher concentrations if this should become necessary for a particular application; however, the preferred concentration of

active principle are from 0.002% to 1% by weight.

For topical administration the compounds of formula I are conveniently provided in the form of unguents, gels, creams, ointments, powders, dyeing compositions, solutions, suspensions, emulsions, lotions, sprays, adhesive plasters and impregnated pads. The compounds according to the invention can be mixed with inert nontoxic, generally liquid or pasty, bases suitable for topical treatment. Preparation of such topical formulations are well described in the art of pharmaceutical formulations as exemplified, for example, Remington's Pharmaceutical Science, Edition 17, Mack Publishing Company, Easton, Pa. Other medicaments can be added to such topical formulation for such secondary purposes as treating skin dryness, providing protection against light; other medications for treating dermatoses, preventing infection, reducing irritation, inflammation and the like.

The compounds according to the invention can also be used enterally. Orally, the compounds according to the invention are suitably administered at the rate of 2 μ g to 100 mg per day per kg of body weight. The required dose can be administered in one or more portions. For oral administration, suitable forms are, for example, tablets, pills, dragees, syrups, suspensions, emulsions, solutions, powders and granules; a preferred method of administration consists in using pills containing from 0.1 mg to about 1 mg of active substance.

U.S. Patent No.4,876,381 issued on October, 24, 1989 to Lang et al. provides examples of formulations

constituting gel, unguent, powder, cream, etc. for a retinoid compound. The aforesaid U.S. Patent can be used as a guide to formulate a compound of formula I and is herein incorporated by reference in its
5 entirety.

Isotretinoin (Accutane®) and etretinate (Tegison®) are used clinically to treat severe recalcitrant cystic acne and severe recalcitrant psoriasis, including the
10 erythrodermica and generalized pustular types, respectively. Their mode of use is amply illustrated in the Physician's Desk Reference, 47th Edition, 1993, published by Medical Economics Data. The compounds of formula I may also be used to treat severe
15 recalcitrant cystic acne or severe recalcitrant psoriasis. In so doing, the compounds of the present invention may be used in a similar fashion to isotretinoin and etretinate; thus, the relevant sections on isotretinoin and etretinate in the
20 Physician's Desk Reference will serve as a convenient guide which will obviate the need for any undue experimentation.

The compounds according to the invention can also be
25 administered parenterally in the form of solutions or suspensions for intravenous, intraperitoneal or intramuscular perfusions or injections. In that case, the compounds according to the invention are generally administered at the rate of about 2 µg to 100 mg per
30 day per kg of body weight; a preferred method of administration consists of using solutions or suspensions containing approximately from 0.01 mg to 1 mg of active substance per ml.


- Several retinoids have been found to possess anti-tumor properties. Roberts, A.B. and Sporn, M.B. in "The Retinoids," Sporn, M.B., Roberts, A.B., and Goodman, D.S., eds, 1984, 2 pp. 209-286, Academic Press, New York; Lippman, S.M., Kessler, J.F., and Meyskens, F.L., *Cancer Treat. Rep.*, 1987, 71, p. 391; *ibid.*, p. 493. As used herein, the term "anti-tumor" includes both chemopreventative (prophylactic or tumor promotion inhibiting) and therapeutic (curative) use.
- 10 For example, all-trans retinoic acid can be used to treat acute promyelocytic leukemia. Huang, M. et al., *Blood*, 1988, 72, p. 567. Isotretinoin has been shown to be useful in prevention of second primary tumors in squamous-cell carcinoma of the head and neck. Hong,
- 15 W.K. et al., *N. Engl. J. Med.*, 1990, 323, p. 795.

- The compounds of formula I can also be used in substantially the similar manner to retinoids for treating (both chemopreventively and therapeutically)
- 20 various tumors. For the compounds of this invention, the anti-tumor dose to be administered, whether a single dose, multiple dose, or a daily dose, will of course vary with the particular compound employed because of the varying potency of the compound, the
- 25 chosen route of administration, the size of the recipient, the type of tumor, and the nature of the patient's condition. The dosage to be administered is not subject to definite bounds, but it will usually be an effective amount, or the equivalent on a molar
- 30 basis of the pharmacologically active free form produced from a dosage formulation upon the metabolic release of the active drug to achieve its desired pharmacological and physiological effects. An oncologist skilled in the art of cancer treatment will

- be able to ascertain, without undue experimentation, appropriate protocols for the effective administration of the compounds of this present invention, such as by referring to the earlier published studies on
- 5 retinoids found to have anti-tumor properties. For example, for the prevention of second primary tumors with a compound of formula I in squamous-cell carcinoma of the head and neck, an oncologist may refer to the study by Hong, W.K. et al. in *N. Engl.*
- 10 *J. Med.*, 1990, 323, p. 795. For treating acute promyelocytic leukemia, s/he may refer to the study by Huang, M. et al. in *Blood*, 1988, 72, p. 567.

What we claim is:

1. A compound of formula I



or a nontoxic pharmaceutically acceptable salt,
physiologically hydrolyzable ester or solvate thereof,
in which

X is -O-CO-, -NH-CO-, -CS-NH-, -CO-O-, -CO-NH-,
-COS-, -SCO-, -SCH₂-, -CH₂-CH₂-, -C≡C-,
-CH₂-NH-, -COCH₂-, -NHCS-, -CH₂S-, -CH₂O-,
-OCH₂-, -NHCH₂- or -CR⁵=CR⁶-;

R^m and R^k are independently hydrogen, halogen,
C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy or nitro;
n is zero or one;


R⁴ is -(CH₂)_t-Y, C₁₋₆alkyl, or C₃₋₆cycloalkyl;

R¹ is -CO₂Z, C₁₋₆alkyl, CH₂OH, -CONHR^y, or CHO;

R² and R³ are independently hydrogen or C₁₋₆alkyl;

R^a and R^b are independently hydrogen or C₁₋₆alkyl;

but when n is one, R^a and R^b together can
form a radical of the formula



Y is naphthyl or phenyl, both radicals can be
optionally substituted with one to three same



or different C₁₋₆alkyl or halogen;
 Z is hydrogen or C₁₋₆alkyl;
 R⁵, R⁶ and R⁷ are independently hydrogen or
 C₁₋₆alkyl; and
 5 t is zero to six.

2. A compound of claim 1 in which R¹ is -CO₂H; n is
 one; R² and R³ are independently methyl or hydrogen;
 and R^a and R^b are independently hydrogen or C₁₋₆alkyl.
 10
3. The compound of claim 2 that is 4-[[[(5,6-dihydro-
 5,5-dimethyl-8-ethyl-2-
 naphthalenyl)amino]carbonyl]benzoic acid.
- 15 4. The compound of claim 2 that is 4-[[[(5,6-dihydro-
 5,5,8-trimethyl-2-naphthalenyl)amino]carbonyl]benzoic
 acid.
5. The compound of claim 2 that is 4-[[[(5,6-dihydro-
 20 5,5-dimethyl-8-phenyl-2-
 naphthalenyl)amino]carbonyl]benzoic acid.
6. The compound of claim 2 that is 4-[[[(5,6-dihydro-
 5,5-dimethyl-8-ethyl-2-
 25 naphthalenyl)carbonyl]amino]benzoic acid.
7. The compound of claim 2 that is 4-[[[(5,6-dihydro-
 5,5,8-trimethyl-2-naphthalenyl)carbonyl]amino]benzoic
 acid.
 30
8. The compound of claim 2 that is 4-(E)-[2-(5,6-
 dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)-1-
 propenyl]benzoic acid.



9. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)oxy]carbonyl]benzoic acid.
- 5 10. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)carbonyl]amino]benzoic acid.
11. The compound of claim 2 that is 4-[[[(5,6-
10 dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)carbonyl]oxy]benzoic acid.
12. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-(2-fluorophenyl)-2-
15 naphthalenyl)carbonyl]amino]benzoic acid.
13. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5,6-trimethyl-8-phenyl-2-naphthalenyl)carbonyl]amino]benzoic acid.
20
14. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5,7-trimethyl-8-phenyl-2-naphthanenyl)carbonyl]amino]benzoic acid.
- 25 15. The compound of claim 2 that is 4-[[[(E)-(5,6,-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]vinyl]benzoic acid.
16. The compound of claim 2 that is 4-[[[(5,6,-
30 dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]sulfamyl]benzoic acid.
17. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-



naphthalenyl]sulfamyl]carbonyl]benzoic acid.

18. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]ethyl]benzoic
5 acid.

19. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]thiocarbonyl]amino]benzoic acid.
10

20. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]methyl]benzoic acid.

15 21. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]methyl]oxy]benzoic acid.

22. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]oxy]methyl]benzoic acid.
20

23. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-(2,4-dimethylphenyl))-2-naphthalenyl]carbonyl]amino]benzoic acid.
25

24. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-(4-methylphenyl))-2-naphthalenyl]carbonyl]amino]benzoic acid.
30

25. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]ethynyl]benzoic acid.



26. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]sulfamyl]methyl]benzoic acid.
- 5 27. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]methyl]amino]benzoic acid.
28. The compound of claim 2 that is 4-[[[(5,6-
10 dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]amino]thiocarbonyl]benzoic acid.
29. The compound of claim 2 that is 4-[[[(5,6-
15 naphthalenyl]methyl)sulfamyl]benzoic acid.
30. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]amino]methyl]benzoic acid.
20
31. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]amino]-2-hydroxybenzoic acid.
- 25 32. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]amino]-2-fluorobenzoic acid.
33. The compound of claim 2 that is 4-[[[(5,6-dihydro-
30 5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]amino]-2-nitrobenzoic acid.
34. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]carbonyl]amino]-



2-methoxybenzoic acid.


35. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-(2-naphthalene)]-2-naphthalenyl]carbonyl]amino]benzoic acid.

36. The compound of claim 2 that is 4-[[[(5,6-dihydro-8-phenyl-2-naphthalenyl)carbonyl]amino]benzoic acid.

37. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)carbonyl]amino]-3-fluorobenzoic acid.

38. The compound of claim 2 that is 4-[[[(5,6-dihydro-5,5-dimethyl-8-phenyl-2-naphthalenyl)carbonyl]amino]-3-methylbenzoic acid.

39. A compound of claim 1 in which R^1 is $-CO_2H$; X is $-CONH-$; R^2 and R^3 are methyl; and R^a and R^b together form a radical of the formula



40. The compound of claim 39 that is 4-[[[(5,8,10,10a-tetrahydro-10,10-dimethyl-9-phenyl-2-anthracenyl)carbonyl]amino]benzoic acid.

41. A compound of claim 1 in which R^1 is $-CO_2H$; n is zero; and R^2 and R^3 are methyl.

42. The compound of claim 41 that is 4-[[[(1,1-dimethyl-3-phenyl-1H-inden-5-yl)amino]carbonyl]benzoic



acid.

43. The compound of claim 41 that is 4-[(1,1-dimethyl-3-phenyl-1H-inden-5-yloxymethyl)benzoic acid.

5

44. The compound of claim 41 that is 4-[2-(1,1-dimethyl-3-phenyl-1H-inden-5-yl)vinyl]benzoic acid.

45. The compound of claim 41 that is 4-[(1,1-dimethyl-3-phenyl-1H-indene-5-carbonyl)amino]benzoic acid.

10

46. A pharmaceutical composition comprising a compound as claimed in claim 1.

15

47. A method of treating tumor in a mammal comprising administering a therapeutically effective amount of a compound as claimed in claim 1.

20 48. A method of treating rheumatoid arthritis in a mammal comprising administering a therapeutically effective amount of a compound as claimed in claim 1.

49. A method of treating rheumatoid arthritis as
25 claimed in claim 48 for 4-[(E)-(5,6,-dihydro-5,5-dimethyl-8-phenyl)-2-naphthalenyl]vinyl]benzoic acid.

This Sheet Has Been Replaced



* C L M S *