Claims

1. Use of a tumor-effective oligonucleotide for the manufacture of a medicament for the treatment ~~in~~ of a tumor in a patient after complete or partial resection or ablation of a solid tumor of this patient, characterized in that the oligonucleotide medicament ~~is applied~~when used in a body cavity created during the resection or ablation of the solid tumor of the patient ~~in order to~~ fights tumor cells remaining in the tumor bed and within its vicinity after surgery or ablation or ~~to~~ fights metastases, and ~~in order to~~ counteracts recurrences or the formation of new metastases in this area.

2. Use of an oligonucleotide according to claim 1, characterized in that the oligonucleotide used has a pleiotropic effect in at least one type of tumor cell.

3. Use of an oligonucleotide according to ~~anyone~~any one of claims 1 or 2, characterized in that th~~it is applied~~e oligonucleotide medicament is in the form of a gel, ~~in~~ a liquid, or ~~an~~ elastic material.

4. Use of an oligonucleotide according to any one of claims 1, 2, or 3 ~~as a load of a carrier in the form of gel, liquid or an elastic material~~, characterized in that ~~the~~ a carrier in the form of a gel, liquid, or elastic material comprises the oligonucleotides preferably at a concentration of 15 or 70 micrograms per milliliter, and ~~particularly preferred~~ more preferably at a concentration of more than 250 or 1000 micrograms, or 5, 25 or 100 milligrams per milliliter.

5. Use of an oligonucleotide according to claim 4, characterized in that the carrier consists of resorbable elastic or gel-like material and at least partially reproduces the shape of the resected organ.

6. Use of an oligonucleotide according to any one of ~~the~~ claims 1, 2, or 3, ~~as a load of a flat carrier such as membranes, foils or gauze,~~ wherein the oligonucleotides are carried directly, packaged in particles such as liposomes, conjugated or formulated as a gel on a flat carrier such as membranes, foils, or gauze, characterized in that ~~that~~ the loading densities are at least 10

micrograms per square centimeter, particularly preferred 50 or 200 micrograms~~,~~ or 1, 5 or 20 milligrams of oligonucleotide per square centimeter.

7. Use of an oligonucleotide according to any one of claims 4 to 6, characterized in that the carrier contains materials with low immunogenicity, in particular collagen, atelocollagen, gelatin, chitosan, or hyaluronic acid.

8. Use of an oligonucleotide according to ~~claim~~claims 6 or 7, characterized in that the carrier additionally has markers for specific detection, in particular absorptive or fluorescent dyes.

9. Use of an oligonucleotide according to any one of claims 4 to 8, characterized in that the crosslinking ~~or other stabilization~~ of the carrier is stopped such that the carrier material is degraded in the patient's body ~~at a preferred rate, in particular~~ with an *in vivo* half-life of more than a week, and ~~particularly preferred~~ preferably with a half-life of more than 2 weeks, 4 weeks and 3 months.

10. Use of an oligonucleotide according to any one of claims 4 to 9, characterized in that in addition to the oligonucleotides, other tumor-effective molecules, in particular cytostatics, are to be applied to the carrier in therapeutically relevant dosages.

11. Use of an oligonucleotide according to any one of claims 4 to 10, characterized in that carriers are combined with each another, in particular gel or gauze in combination with a liquid formulation, such as to reach optimally tumor cells which are located at a distance from ~~the~~ a wound edge.

12. Use of an oligonucleotide according to any one of claims 4 to 11, characterized in that ~~the~~ diagnostic elements are integrated ~~in~~into the carrier, in particular sensors for heart rate, LDL, or HDL concentrations.

~~13. Use of the oligonucleotides according to any one of claims 1 to 12, characterized in that standard treatments, such as operations, aftercare and adjuvant chemotherapy, are carried out in parallel with or almost simultaneously with the method according to the invention.~~

134. Use of oligonucleotides according to any one of claims 1 to 13 ~~for the adjuvant therapy of tumors, in particular of~~characterized in that the tumor is selected from the group consisting of adrenocortical carcinoma, ovarian cancer, mesothelioma, and glioblastoma.