

BICYCLO [2.2.1] ACID GPR120 MODULATORS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application Serial No.
5 61/782,469, filed March 14, 2013; the entire content of which is incorporated herein by reference.

FIELD OF THE INVENTION

[0002] The present invention provides novel bicyclo [2.2.1] compounds, and their
10 analogues thereof, which are GPR120 G protein-coupled receptor modulators, compositions containing them, and methods of using them, for example, for the treatment of diabetes and related conditions.

BACKGROUND OF THE INVENTION

15 [0003] Diabetes mellitus is a progressively debilitating disorder of epidemic proportions leading to various micro- and macrovascular complications and morbidity. The most common type of diabetes, type 2 diabetes, is characterized by increasing insulin resistance associated with inadequate insulin secretion after a period of compensatory hyperinsulinemia. Polyunsaturated fatty acids (PUFAs) such as omega-3 fatty acids are
20 known to improve sensitivity to insulin. Insulin sensitivity can be improved by exerting anti-inflammatory effects in monocytes and/or macrophages and/or by enhancing glucose uptake in adipose and muscle. GPR120 is a membrane-bound receptor responsive to PUFAs which is preferentially expressed in adipose tissue and monocytes/macrophages. To decrease the medical burden of type 2 diabetes through enhanced glycemic control,
25 GPR120 modulator compounds hold the promise of exerting a sensitizing effect to insulin as well as potential combination with a broad range of antidiabetic drugs.

[0004] The present invention relates to novel substituted bicyclic acid compounds which have the ability to modulate GPR120. Such compounds are therefore potentially useful for the treatment or prophylaxis of diabetes and related conditions.

30

SUMMARY OF THE INVENTION

[0005] The present invention provides bicyclo [2.2.1] compounds, and their analogues thereof, which are useful as GPR120 modulators, including stereoisomers, tautomers, pharmaceutically acceptable salts, or solvates thereof.

5 [0006] The present invention also provides processes and intermediates for making the compounds of the present invention or stereoisomers, tautomers, pharmaceutically acceptable salts, or solvates thereof.

[0007] The present invention also provides pharmaceutical compositions comprising a pharmaceutically acceptable carrier and at least one of the compounds of the present
10 invention or stereoisomers, tautomers, pharmaceutically acceptable salts, or solvates thereof.

[0008] The compounds of the invention may be used in the treatment of multiple diseases or disorders associated with GPR120, such as diabetes and related conditions, microvascular complications associated with diabetes, the macrovascular complications
15 associated with diabetes, cardiovascular diseases, Metabolic Syndrome and its component conditions, disorders of glucose metabolism, obesity and other maladies.

[0009] The compounds of the invention may be used in therapy.

[0010] The compounds of the invention may be used for the manufacture of a medicament for the treatment of multiple diseases or disorders associated with GPR120.


20 [0011] The compounds of the invention can be used alone, in combination with other compounds of the present invention, or in combination with one or more other agent(s).

[0012] Other features and advantages of the invention will be apparent from the following detailed description and claims.

25 DETAILED DESCRIPTION OF THE INVENTION

I. COMPOUNDS OF THE INVENTION

[0013] In a first aspect, the present disclosure provides, *inter alia*, a compound of Formula (I):



30

or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof, wherein:

L_1 is independently L_4 -O or O- L_4 ;

L_2 is independently a hydrocarbon linker substituted with 0-2 R^c ,

- 5 a hydrocarbon-heteroatom linker substituted with 0-2 R^c , or $-(CH_2)_{1-2}-(C_{3-4}$ cycloalkyl substituted with 0-2 R^c)-(CH₂)₀₋₁-; wherein said hydrocarbon linker has one to six carbon atoms and may be straight or branched, saturated or unsaturated; and said hydrocarbon-heteroatom linker has one to four carbon atoms and one group selected from O, -CO-, S, -SO-, -SO₂-, NH, N(C₁₋₄ alkyl), -CONH-, and -NHCO-;

- 10 L_4 is independently a bond or a hydrocarbon linker; wherein said hydrocarbon linker has one to four carbon atoms and may be straight or branched;

R^1 is independently selected from: C₆₋₁₀ carbocycle and a 5- to 10-membered heterocycle comprising carbon atoms and 1-4 heteroatoms selected from N, NR^b, O, and S(O)_p; wherein said carbocycle and heterocycle are substituted with 0-4 R^3 and 0-1 R^4 ;

- 15 R^2 independently selected from: OH, CO₂H, CO₂(C₁₋₄ alkyl), CONR^eR^f, and -CONHSO₂R^f;

R^3 , at each occurrence, is independently selected from: halogen, C₁₋₆ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, C₁₋₄ haloalkylthio, and NO₂;

- 20 R^4 is independently - L_3 - R^5 ;

L_3 is independently selected from: a bond, O, and C(=O);

R^5 is independently selected from: phenyl and a 5- to 6-membered heteroaryl comprising carbon atoms and 1-4 heteroatoms selected from N, NR^b, O, and S(O)_p; wherein each ring moiety is substituted with 0-2 R^a ;

- 25 R^a , at each occurrence, is independently selected from: halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkyl;

R^b , at each occurrence, is independently selected from: H, C₁₋₄ alkyl, and $-(CH_2)_{0-2}-(phenyl \text{ substituted with } 0-3 R^d)$;

R^c , at each occurrence, is independently selected from: =O, halogen, OH, C_{1-4} alkyl, C_{1-4} haloalkyl, and C_{1-4} alkoxy;

R^d , at each occurrence, is independently selected from: halogen, C_{1-4} alkyl, C_{1-4} alkoxy, and C_{1-4} haloalkyl;

5 R^e , at each occurrence, is independently selected from: H and C_{1-4} alkyl;

R^f , at each occurrence, is independently selected from: H, C_{1-4} alkyl, $-(CH_2)_{0-2}$ -phenyl, and C_{3-6} cycloalkyl substituted with 1-2 OH;

R^e and R^f may be combined with the nitrogen atom to which they are attached to form a 4- to 6-membered heterocyclic ring comprising carbon atoms and 1 additional
10 heteroatom selected from N, NR^b , O, and S; wherein said heterocycle is substituted with and 0-1 OH; and

p is, independently at each occurrence, selected from 0, 1, and 2.

[0014] In a second aspect, the present invention includes a compound of Formula (I)
15 or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof, within the scope of the first aspect, wherein:


L_1 is independently L_4 -O;

L_2 is independently a hydrocarbon linker substituted with 0-1 R^c ,
a hydrocarbon-heteroatom linker substituted with 0-1 R^c , or $-(CH_2)_{1-2}$ -(C_{3-4} cycloalkyl
20 substituted with 0-1 R^c)- $(CH_2)_{0-1}$ -; wherein said hydrocarbon linker has one to six carbon atoms and may be straight or branched, saturated or unsaturated; and said hydrocarbon-heteroatom linker has one to four carbon atoms and one group selected from O and S;

R^1 is independently selected from: phenyl, indanyl, naphthyl, and a 5- to
25 10-membered heterocycle comprising carbon atoms and 1-4 heteroatoms selected from N, NR^b , O, and $S(O)_p$; wherein each ring moiety substituted with 0-4 R^3 and 0-1 R^4 ; and

R^4 is independently selected from: thienyl, oxadiazolyl, and $-L_3$ -phenyl; wherein each ring moiety is substituted with 0-2 R^a .


30 **[0015]** In a third aspect, the present invention includes a compound of Formula (II):




or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof, within the scope of the first or second aspect, wherein:

- L_2 is independently a hydrocarbon linker a hydrocarbon-heteroatom linker, or
 5 $-(CH_2)_{1-2}$ -(cyclopropyl substituted with 0-1 R^c)- $(CH_2)_{0-1}$ -; wherein said hydrocarbon linker has one to five carbon atoms and may be straight or branched, saturated or unsaturated; and said hydrocarbon-heteroatom linker has one to three carbon atoms and one O;

L_4 is independently selected from: a bond, CH_2 and $CH(C_{1-4} \text{ alkyl})$;

- 10 R^1 is independently selected from: , and a ring moiety substituted with 0-2 R^3 and selected from the group consisting of thienyl,

isoxazolyl, pyrimidinyl, indanyl, naphthyl, benzothiophenyl, and ;

R^3 , at each occurrence, is independently selected from: NO_2 , halogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkylthio, C_{1-4} haloalkyl, C_{1-4} haloalkoxy, and C_{1-4} haloalkylthio;

- 15 R^4 is independently selected from: thienyl, oxadiazolyl, and $-L_3$ -phenyl; wherein each ring moiety is substituted with 0-2 R^a ;

L_3 is independently selected from: a bond, O, and $C(=O)$;

R^a , at each occurrence, is independently selected from: halogen and C_{1-4} alkyl;

and


- 20 R^b is independently phenyl substituted with 0-2 halo.

[0016] In a fourth aspect, the present invention includes a compound of Formula (I) or (II), or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof, within the scope of any of the above aspects, wherein:

L_2 is independently selected from: CH_2OCH_2 , OCH_2CH_2 , $CH_2CH_2CH_2$,

5 $CH_2CH(C_{1-2} \text{ alkyl})CH_2$, $CH_2CH_2CH(C_{1-2} \text{ alkyl})$, $CH_2CH=CH$, and ;

R^{1-L_4-} is independently selected from:



R^3 , at each occurrence, is independently selected from: halogen, C_{1-4} alkyl,

10 C_{1-4} alkoxy, C_{1-4} alkylthio, C_{1-4} haloalkyl, C_{1-4} haloalkoxy, and C_{1-4} haloalkylthio;

R^4 is independently selected from: thienyl and $-L_3$ -(phenyl substituted with 0-2 halo); and


L_3 is independently selected from: a bond, O, and $C(=O)$.

15 [0017] In a fifth aspect, the present invention includes a compound of Formula (I) or (II), or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof, within the scope of any of the above aspects, wherein:

L_2 is independently selected from the group consisting of CH_2OCH_2 , OCH_2CH_2 ,

$CH_2CH_2CH_2$, $CH_2CH(Me)CH_2$, $CH_2CH_2CH(Me)$, $CH_2CH=CH$, and ; and

20 R^{1-L_4-} is independently selected from:



[0018] In a sixth aspect, the present invention includes a compound of Formula (I) or (II), or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate


10 thereof, within the scope of any of the above aspects, wherein:

L_2 is independently selected from: OCH_2CH_2 , $CH_2CH_2CH_2$, and




and


R^1-L_4- is independently selected from:



[0019] In a seventh aspect, the present invention includes a compound of Formula (I) or (II), or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof, within the scope of any of the above aspects, wherein:

R^1-L_4- is independently selected from:





[0020] In an eighth aspect, the present invention provides a compound selected from the exemplified examples within the scope of the first aspect, or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

[0021] In another aspect, the present invention provides a compound selected from any subset list of compounds within the scope of any of the above aspects.

[0022] In another embodiment, R^1 is independently phenyl substituted with 0-4 R^3 and 0-1 R^4 .

[0023] In another embodiment, R^e , at each occurrence, is independently selected from: H and C_{1-4} alkyl; and R^f , at each occurrence, is independently selected from: H, C_{1-4} alkyl, $-(CH_2)_{0-2}$ -phenyl, and C_{3-6} cycloalkyl substituted with 1-2 OH.

15

[0024] In another embodiment, the compounds of the present invention have hGPR120 EC_{50} values $\leq 10 \mu M$.

[0025] In another embodiment, the compounds of the present invention have hGPR120 EC_{50} values $\leq 5 \mu M$.

[0026] In another embodiment, the compounds of the present invention have hGPR120 EC_{50} values $\leq 1 \mu M$.

[0027] In another embodiment, the compounds of the present invention have hGPR120 EC_{50} values $\leq 0.5 \mu M$.

II. OTHER EMBODIMENTS OF THE INVENTION

[0028] In another embodiment, the present invention provides a composition comprising at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

5 [0029] In another embodiment, the present invention provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

[0030] In another embodiment, the present invention provides a pharmaceutical
10 composition, comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

[0031] In another embodiment, the present invention provides a process for making a
15 compound of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

[0032] In another embodiment, the present invention provides an intermediate for making a compound of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

[0033] In another embodiment, the present invention provides a pharmaceutical
20 composition further comprising additional therapeutic agent(s). Examples of additional therapeutic agent(s), according to the present invention include, but are not limited to, anti-diabetic agents, anti-hyperglycemic agents, anti-hyperinsulinemic agents, anti-retinopathic agents, anti-neuropathic agents, anti-nephropathic agents, anti-atherosclerotic agents, anti-ischemic agents, anti-hypertensive agents, anti-obesity
25 agents, anti-dyslipidemic agents, anti-hyperlipidemic agents, anti-hypertriglyceridemic agents, anti-hypercholesterolemic agents, anti-pancreatitis agents, lipid lowering agents, anorectic agents and appetite suppressants.

[0034] In a preferred embodiment, the present invention provides pharmaceutical
30 composition, wherein the additional therapeutic agents are, for example, a dipeptidyl peptidase-IV (DPP4) inhibitor (for example a member selected from saxagliptin, sitagliptin, vildagliptin, linagliptin and alogliptin), a sodium-glucose transporter-2 (SGLT2) inhibitor (for example a member selected from dapagliflozin, canagliflozin,

empagliflozin and remagliflozin), a GPR40/FFAR1 (Free fatty acid receptor 1) agonist (for example, TAK-875), and/or an MGAT2 (monoacylglycerol transferase 2) inhibitor (for example, compounds from WO 2012/124744, or compound (S)-10 from *Bioorg. Med. Chem. Lett.* (2013), doi: <http://dx.doi.org/10.1016/j.bmcl.2013.02.084>).

5 **[0035]** In a preferred embodiment, the present invention provides pharmaceutical composition, wherein the additional therapeutic agents are, for example, a dipeptidyl peptidase-IV inhibitor, a sodium-glucose transporter-2 inhibitor and an 11b-HSD-1 inhibitor.

10 **[0036]** In another embodiment, the present invention provides a method for the treatment of multiple diseases or disorders associated with GPR120, comprising administering to a patient in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention, alone, or, optionally, in combination with another compound of the present invention and/or at least one other type of therapeutic agent.

15 **[0037]** Examples of diseases or disorders associated with the activity of the GPR120 that can be prevented, modulated, or treated according to the present invention include, but are not limited to, diabetes, hyperglycemia, impaired glucose tolerance, gestational diabetes, insulin resistance, hyperinsulinemia, retinopathy, neuropathy, nephropathy, diabetic kidney disease, acute kidney injury, cardiorenal syndrome, delayed wound
20 healing, atherosclerosis and its sequelae, abnormal heart function, congestive heart failure, myocardial ischemia, stroke, Metabolic Syndrome, hypertension, obesity, fatty liver disease, dyslipidemia, dyslipidemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), lipid disorders and liver diseases such as NASH (Non-Alcoholic SteatoHepatitis),
25 NAFLD (Non-Alcoholic Fatty Liver Disease) and liver cirrhosis.

30 **[0038]** In another embodiment, the present invention provides a method for the treatment of diabetes, hyperglycemia, gestational diabetes, obesity, dyslipidemia and hypertension, comprising administering to a patient in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention, alone, or, optionally, in combination with another compound of the present invention and/or at least one other type of therapeutic agent.

[0039] In another embodiment, the present invention provides a method for the treatment of diabetes, comprising administering to a patient in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention, alone, or, optionally, in combination with another compound of the present invention and/or at least one other type of therapeutic agent.

[0040] In another embodiment, the present invention provides a method for the treatment of hyperglycemia, comprising administering to a patient in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention, alone, or, optionally, in combination with another compound of the present invention and/or at least one other type of therapeutic agent.

[0041] In another embodiment, the present invention provides a method for the treatment of obesity, comprising administering to a patient in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention, alone, or, optionally, in combination with another compound of the present invention and/or at least one other type of therapeutic agent.

[0042] In another embodiment, the present invention provides a method for the treatment of dyslipidemia, comprising administering to a patient in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention, alone, or, optionally, in combination with another compound of the present invention and/or at least one other type of therapeutic agent.

[0043] In another embodiment, the present invention provides a compound of the present invention for use in therapy.

[0044] In another embodiment, the present invention provides a compound of the present invention for use in therapy for the treatment of multiple diseases or disorders associated with GPR120.

[0045] In another embodiment, the present invention also provides the use of a compound of the present invention for the manufacture of a medicament for the treatment of multiple diseases or disorders associated with GPR120.

[0046] In another embodiment, the present invention provides a method for the treatment of multiple diseases or disorders associated with GPR120, comprising administering to a patient in need thereof a therapeutically effective amount of a first and second therapeutic agent, wherein the first therapeutic agent is a compound of the present

invention. Preferably, the second therapeutic agent, for example, dipeptidyl peptidase-IV (DPP4) inhibitor (for example a member selected from saxagliptin, sitagliptin, linagliptin, vildagliptin and alogliptin).

[0047] In another embodiment, the present invention provides a combined
5 preparation of a compound of the present invention and additional therapeutic agent(s) for simultaneous, separate or sequential use in therapy.

[0048] In another embodiment, the present invention provides a combined
preparation of a compound of the present invention and additional therapeutic agent(s) for
simultaneous, separate or sequential use in the treatment of multiple diseases or disorders
10 associated with GPR120.

[0049] Where desired, the compound of the present invention may be used in
combination with one or more other types of antidiabetic agents and/or one or more other
types of therapeutic agents which may be administered orally in the same dosage form, in
a separate oral dosage form or by injection. The other type of antidiabetic agent that may
15 be optionally employed in combination with the GPR120 receptor modulator of the
present invention may be one, two, three or more antidiabetic agents or
antihyperglycemic agents which may be administered orally in the same dosage form, in
a separate oral dosage form, or by injection to produce an additional pharmacological
benefit.

[0050] The antidiabetic agents used in the combination with the GPR120 receptor
modulator of the present invention include, but are not limited to, insulin secretagogues
or insulin sensitizers, other GPR120 receptor modulators, or other antidiabetic agents.
These agents include, but are not limited to, dipeptidyl peptidase IV (DP4) inhibitors (for
example, sitagliptin, saxagliptin, linagliptin, alogliptin and vildagliptin), biguanides (for
25 example, metformin and phenformin), sulfonyl ureas (for example, glyburide, glimepiride
and glipizide), glucosidase inhibitors (for example, acarbose, miglitol), PPAR γ agonists
such as thiazolidinediones (for example, rosiglitazone and pioglitazone), PPAR α/γ dual
agonists (for example, muraglitazar, peliglitazar, tesaglitazar and aleglitazar), glucokinase
activators (for example, PF-04937319 and AMG-151), GPR119 receptor modulators (for
30 example, MBX-2952, PSN821, and APD597), sodium-glucose transporter-2 (SGLT2)
inhibitors (for example, dapagliflozin, canagliflozin, empagliflozin and remagliflozin),

GPR40 receptor agonists (*e.g.*, TAK-875), amylin analogs such as pramlintide, and/or insulin.

[0051] The GPR120 receptor modulator of the present invention may also be optionally employed in combination with agents for treating complication of diabetes.

5 These agents include PKC inhibitors and/or AGE inhibitors.

[0052] The GPR120 receptor modulator of the present invention may also be optionally employed in combination with one or more hypophagic and/or weight-loss agents such as diethylpropion, phendimetrazine, phentermine, orlistat, sibutramine, lorcaserin, pramlintide, topiramate, MCHR1 receptor antagonists, oxyntomodulin, naltrexone, Amylin peptide, NPY Y5 receptor modulators, NPY Y2 receptor modulators, NPY Y4 receptor modulators, cetilistat, 5HT2c receptor modulators, MGAT2 inhibitors and the like. The GPR120 receptor modulator of the present invention may also be employed in combination with an agonist of the glucagon-like peptide-1 receptor (GLP-1 R), such as exenatide, liraglutide, GLP-1(1-36) amide, GLP-1(7-36) amide, GLP-1(7-37), which may be administered via injection, intranasal, or by transdermal or buccal devices.

[0053] The present invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof. This invention encompasses all combinations of preferred aspects of the invention noted herein. It is understood that any and all embodiments of the present invention may be taken in conjunction with any other embodiment or embodiments to describe additional embodiments. It is also understood that each individual element of the embodiments is its own independent embodiment. Furthermore, any element of an embodiment is meant to be combined with any and all other elements from any embodiment to describe an additional embodiment.

25 III. CHEMISTRY

[0054] Throughout the specification and the appended claims, a given chemical formula or name shall encompass all stereo and optical isomers and racemates thereof where such isomers exist. The term "stereoisomer(s)" refers to compound(s) which have identical chemical constitution, but differ with regard to the arrangement of the atoms or groups in space. Unless otherwise indicated, all chiral (enantiomeric and diastereomeric) and racemic forms are within the scope of the invention. The term "chiral" refers to molecules which have the property of non-superimposability of the mirror image partner,

while the term "achiral" refers to molecules which are superimposable on their mirror image partner. The terms "racemic mixture" and "racemate" refer to an equimolar mixture of two enantiomeric species, devoid of optical activity.

[0055] Many geometric isomers of C=C double bonds, C=N double bonds, ring systems, and the like can also be present in the compounds, and all such stable isomers are contemplated in the present invention. *Cis*- and *trans*- (or *E*- and *Z*-) geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. The present compounds can be isolated in optically active or racemic forms. Optically active forms may be prepared by resolution of stereoisomeric forms or by synthesis from optically active starting materials. All processes used to prepare compounds of the present invention and intermediates made therein are considered to be part of the present invention. When enantiomeric or diastereomeric products are prepared, they may be separated by conventional methods, for example, by chromatography or fractional crystallization. Depending on the process conditions the end products of the present invention are obtained either in free (neutral) or salt form. Both the free form and the salts of these end products are within the scope of the invention. If so desired, one form of a compound may be converted into another form. A free base or acid may be converted into a salt; a salt may be converted into the free compound or another salt; a mixture of isomeric compounds of the present invention may be separated into the individual isomers. Compounds of the present invention, free form and salts thereof, may exist in multiple tautomeric forms, in which hydrogen atoms are transposed to other parts of the molecules and the chemical bonds between the atoms of the molecules are consequently rearranged. It should be understood that all tautomeric forms, insofar as they may exist, are included within the invention.

[0056] Unless otherwise indicated, any heteroatom with unsatisfied valences is assumed to have hydrogen atoms sufficient to satisfy the valences.

[0057] As used herein, the term "alkyl" or "alkylene" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, "C₁ to C₆ alkyl" or "C₁₋₆ alkyl" denotes alkyl having 1 to 6 carbon atoms. Alkyl group can be unsubstituted or substituted with at least one hydrogen being replaced by another chemical group. Example alkyl groups include, but are not limited to, methyl (Me), ethyl (Et), propyl (*e.g.*, *n*-propyl and isopropyl), butyl

(*e.g.*, *n*-butyl, isobutyl, *t*-butyl), and pentyl (*e.g.*, *n*-pentyl, isopentyl, neopentyl). When "C₀ alkyl" or "C₀ alkylene" is used, it is intended to denote a direct bond.

[0058] "Alkenyl" or "alkenylene" is intended to include hydrocarbon chains of either straight or branched configuration having the specified number of carbon atoms and one or more, preferably one to two, carbon-carbon double bonds that may occur in any stable point along the chain. For example, "C₂ to C₆ alkenyl" or "C₂₋₆ alkenyl" (or alkenylene), is intended to include C₂, C₃, C₄, C₅, and C₆ alkenyl groups. Examples of alkenyl include, but are not limited to, ethenyl, 1-propenyl, 2-propenyl, 2-butenyl, 3-butenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 2-methyl-2-propenyl, and 4-methyl-3-pentenyl.

[0059] The term "alkoxy" or "alkyloxy" refers to an -O-alkyl group. For example, "C₁ to C₆ alkoxy" or "C₁₋₆ alkoxy" (or alkyloxy), is intended to include C₁, C₂, C₃, C₄, C₅, and C₆ alkoxy groups. Example alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy (*e.g.*, *n*-propoxy and isopropoxy), and *t*-butoxy. Similarly, "alkylthio" or "thioalkoxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through a sulphur bridge; for example methyl-S- and ethyl-S-.

[0060] "Halo" or "halogen" includes fluoro, chloro, bromo, and iodo. "Haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogens. Examples of haloalkyl include, but are not limited to, fluoromethyl, difluoromethyl, trifluoromethyl, trichloromethyl, pentafluoroethyl, pentachloroethyl, 2,2,2-trifluoroethyl, heptafluoropropyl, and heptachloropropyl. Examples of haloalkyl also include "fluoroalkyl" that is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more fluorine atoms.

[0061] "Haloalkoxy" or "haloalkyloxy" represents a haloalkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. For example, "C₁₋₆ haloalkoxy", is intended to include C₁, C₂, C₃, C₄, C₅, and C₆ haloalkoxy groups. Examples of haloalkoxy include, but are not limited to, trifluoromethoxy, 2,2,2-trifluoroethoxy, and pentafluoroethoxy. Similarly, "haloalkylthio" or "thiohaloalkoxy" represents a haloalkyl group as defined above with the indicated number

of carbon atoms attached through a sulphur bridge; for example trifluoromethyl-S-, and pentafluoroethyl-S-.

[0062] The term "cycloalkyl" refers to cyclized alkyl groups, including mono-, bi- or poly-cyclic ring systems. For example, "C₃ to C₆ cycloalkyl" or "C₃₋₆ cycloalkyl" is

5 intended to include C₃, C₄, C₅, and C₆ cycloalkyl groups. Example cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and norbornyl. Branched cycloalkyl groups such as 1-methylcyclopropyl and 2-methylcyclopropyl are included in the definition of "cycloalkyl". The term "cycloalkenyl" refers to cyclized alkenyl groups. C₄₋₆ cycloalkenyl is intended to include
10 C₄, C₅, and C₆ cycloalkenyl groups. Example cycloalkenyl groups include, but are not limited to, cyclobutenyl, cyclopentenyl, and cyclohexenyl.

[0063] As used herein, "carbocycle", "carbocyclyl", or "carbocyclic residue" is intended to mean any stable 3-, 4-, 5-, 6-, 7-, or 8-membered monocyclic or bicyclic or 7-, 8-, 9-, 10-, 11-, 12-, or 13-membered bicyclic or tricyclic ring, any of which may be

15 saturated, partially unsaturated, unsaturated or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclobutenyl, cyclopentyl, cyclopentenyl, cyclohexyl, cycloheptenyl, cycloheptyl, cycloheptenyl, adamantyl, cyclooctyl, cyclooctenyl, cyclooctadienyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl,
20 adamantyl, anthracenyl, and tetrahydronaphthyl (tetralin). As shown above, bridged rings are also included in the definition of carbocycle (*e.g.*, [2.2.2]bicyclooctane). Preferred carbocycles, unless otherwise specified, are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, indanyl, and tetrahydronaphthyl. When the term "carbocycle" is used, it is intended to include "aryl". A bridged ring occurs when one or more, preferably
25 one to three, carbon atoms link two non-adjacent carbon atoms. Preferred bridges are one or two carbon atoms. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge.

[0064] As used herein, the term "bicyclic carbocycle" or "bicyclic carbocyclic group" is intended to mean a stable 9- or 10-membered carbocyclic ring system that contains two
30 fused rings and consists of carbon atoms. Of the two fused rings, one ring is a benzo ring fused to a second ring; and the second ring is a 5- or 6-membered carbon ring which is

saturated, partially unsaturated, or unsaturated. The bicyclic carbocyclic group may be attached to its pendant group at any carbon atom which results in a stable structure. The bicyclic carbocyclic group described herein may be substituted on any carbon if the resulting compound is stable. Examples of a bicyclic carbocyclic group are, but not limited to, naphthyl, 1,2-dihydronaphthyl, 1,2,3,4-tetrahydronaphthyl, and indanyl.

[0065] "Aryl" groups refer to monocyclic or bicyclic aromatic hydrocarbons, including, for example, phenyl, and naphthyl. Aryl moieties are well known and described, for example, in Lewis, R.J., ed., *Hawley's Condensed Chemical Dictionary*, 13th Edition, John Wiley & Sons, Inc., New York (1997). "C₆₋₁₀ aryl" refers to phenyl and naphthyl.

[0066] The term "benzyl", as used herein, refers to a methyl group on which one of the hydrogen atoms is replaced by a phenyl group.

[0067] As used herein, the term "heterocycle", "heterocyclyl", or "heterocyclic group" is intended to mean a stable 3-, 4-, 5-, 6-, or 7-membered monocyclic or bicyclic or 7-, 8-, 9-, 10-, 11-, 12-, 13-, or 14-membered polycyclic heterocyclic ring that is saturated, partially unsaturated, or fully unsaturated, and that contains carbon atoms and 1, 2, 3 or 4 heteroatoms independently selected from the group consisting of N, O and S; and including any polycyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized (*i.e.*, N→O and S(O)_p, wherein p is 0, 1 or 2). The nitrogen atom may be substituted or unsubstituted (*i.e.*, N or NR wherein R is H or another substituent, if defined). The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. A nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. When the term "heterocycle" is used, it is intended to include heteroaryl.

[0068] Examples of heterocycles include, but are not limited to, acridinyl, azetidiny, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzoxazoliny, benzthiazolyl, benztriazolyl, benztetrazolyl,

benzisoxazolyl, benzisothiazolyl, benzimidazolyl, carbazolyl, 4a*H*-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolyl, decahydroquinolyl, 2*H*,6*H*-1,5,2-dithiazinyl, dihydrofuro[2,3-*b*]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolyl, imidazolyl, 1*H*-indazolyl, imidazolopyridinyl, indolenyl, 5 indolyl, indolizyl, indolyl, 3*H*-indolyl, isatinoyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolyl, isoindolyl, isoquinolyl, isothiazolyl, isothiazolopyridinyl, isoxazolyl, isoxazolopyridinyl, methylenedioxyphenyl, morpholyl, naphthyridinyl, octahydroisoquinolyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxazolopyridinyl, 10 oxazolidinylperimidinyl, oxindolyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, piperidonyl, 4-piperidonyl, piperonyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolidinyl, pyrazolyl, pyrazolopyridinyl, pyrazolyl, pyridazinyl, pyridooxazolyl, pyridoimidazolyl, pyridothiazolyl, pyridinyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 15 2-pyrrolidinonyl, 2*H*-pyrrolyl, pyrrolyl, quinazolyl, quinolyl, 4*H*-quinolizyl, quinoxalyl, quinuclidinyl, tetrazolyl, tetrahydrofuranyl, tetrahydroisoquinolyl, tetrahydroquinolyl, 6*H*-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thiazolopyridinyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 20 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

[0069] Examples of 5- to 10-membered heterocycles include, but are not limited to, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrazinyl, piperazinyl, piperidinyl, imidazolyl, imidazolidinyl, indolyl, tetrazolyl, isoxazolyl, morpholyl, oxazolyl, 25 oxadiazolyl, oxazolidinyl, tetrahydrofuranyl, thiadiazinyl, thiadiazolyl, thiazolyl, triazinyl, triazolyl, benzimidazolyl, 1*H*-indazolyl, benzofuranyl, benzothiofuranyl, benzotetrazolyl, benzotriazolyl, benzisoxazolyl, benzoxazolyl, oxindolyl, benzoxazolyl, benzthiazolyl, benzisothiazolyl, isatinoyl, isoquinolyl, octahydroisoquinolyl, tetrahydroisoquinolyl, tetrahydroquinolyl, isoxazolopyridinyl, quinazolyl, 30 quinolyl, isothiazolopyridinyl, thiazolopyridinyl, oxazolopyridinyl, imidazolopyridinyl, and pyrazolopyridinyl.

[0070] Examples of 5- to 6-membered heterocycles include, but are not limited to, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrazinyl, piperazinyl, piperidinyl, imidazolyl, imidazolidinyl, indolyl, tetrazolyl, isoxazolyl, morpholinyl, oxazolyl, oxadiazolyl, oxazolidinyl, tetrahydrofuranyl, thiadiazinyl, thiadiazolyl, thiazolyl, triazinyl, and triazolyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

[0071] As used herein, the term "bicyclic heterocycle" or "bicyclic heterocyclic group" is intended to mean a stable 9- or 10-membered heterocyclic ring system which contains two fused rings and consists of carbon atoms and 1, 2, 3, or 4 heteroatoms independently selected from the group consisting of N, O and S. Of the two fused rings, one ring is a 5- or 6-membered monocyclic aromatic ring comprising a 5-membered heteroaryl ring, a 6-membered heteroaryl ring or a benzo ring, each fused to a second ring. The second ring is a 5- or 6-membered monocyclic ring which is saturated, partially unsaturated, or unsaturated, and comprises a 5-membered heterocycle, a 6-membered heterocycle or a carbocycle (provided the first ring is not benzo when the second ring is a carbocycle).

[0072] The bicyclic heterocyclic group may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The bicyclic heterocyclic group described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1.

[0073] Examples of a bicyclic heterocyclic group are, but not limited to, quinolinyl, isoquinolinyl, phthalazinyl, quinazolinyl, indolyl, isoindolyl, indolinyl, 1H-indazolyl, benzimidazolyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, 5,6,7,8-tetrahydro-quinolinyl, 2,3-dihydro-benzofuranyl, chromanyl, 1,2,3,4-tetrahydro-quinoxalyl, and 1,2,3,4-tetrahydro-quinazolinyl.

[0074] As used herein, the term "aromatic heterocyclic group" or "heteroaryl" is intended to mean stable monocyclic and polycyclic aromatic hydrocarbons that include at least one heteroatom ring member such as sulfur, oxygen, or nitrogen. Heteroaryl groups include, without limitation, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, furyl, quinolyl, isoquinolyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl,

benzofuryl, benzothienyl, benzthiazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, indazolyl, 1,2,4-thiadiazolyl, isothiazolyl, purinyl, carbazolyl, benzimidazolyl, indolinyl, benzodioxolanyl, and benzodioxane. Heteroaryl groups are substituted or unsubstituted. The nitrogen atom is substituted or unsubstituted (*i.e.*, N or NR wherein R is H or another
5 substituent, if defined). The nitrogen and sulfur heteroatoms may optionally be oxidized (*i.e.*, N→O and S(O)_p, wherein p is 0, 1 or 2).

[0075] Examples of 5- to 6-membered heteroaryls include, but are not limited to, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrazinyl, imidazolyl, imidazolidinyl, tetrazolyl, isoxazolyl, oxazolyl, oxadiazolyl, oxazolidinyl, thiadiazinyl, thiadiazolyl,
10 thiazolyl, triazinyl, and triazolyl.

[0076] As used herein, the term "aromatic heterocyclic group" or "heteroaryl" is intended to mean stable monocyclic and polycyclic aromatic hydrocarbons that include at least one heteroatom ring member such as sulfur, oxygen, or nitrogen. Heteroaryl groups include, without limitation, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, furyl,
15 quinolyl, isoquinolyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuryl, benzothienyl, benzthiazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, indazolyl, 1,2,4-thiadiazolyl, isothiazolyl, purinyl, carbazolyl, benzimidazolyl, indolinyl, benzodioxolanyl, and benzodioxane. Heteroaryl groups are substituted or unsubstituted. The nitrogen atom is substituted or unsubstituted (*i.e.*, N or NR wherein R is H or another
20 substituent, if defined). The nitrogen and sulfur heteroatoms may optionally be oxidized (*i.e.*, N→O and S(O)_p, wherein p is 0, 1 or 2).

[0077] Examples of 5- to 6-membered heteroaryls include, but are not limited to, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrazinyl, imidazolyl, imidazolidinyl, tetrazolyl, isoxazolyl, oxazolyl, oxadiazolyl, oxazolidinyl, thiadiazinyl, thiadiazolyl,
25 thiazolyl, triazinyl, and triazolyl.

[0078] Bridged rings are also included in the definition of heterocycle. A bridged ring occurs when one or more, preferably one to three, atoms (*i.e.*, C, O, N, or S) link two non-adjacent carbon or nitrogen atoms. Examples of bridged rings include, but are not limited to, one carbon atom, two carbon atoms, one nitrogen atom, two nitrogen atoms,
30 and a carbon-nitrogen group. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge.

[0079] The term "counter ion" is used to represent a negatively charged species such as chloride, bromide, hydroxide, acetate, and sulfate or a positively charged species such as sodium (Na⁺), potassium (K⁺), calcium (Ca²⁺) ammonium (R_nNH_m⁺ where n=0-4 and m=0-4) and the like.

5 [0080] When a dotted ring is used within a ring structure, this indicates that the ring structure may be saturated, partially saturated or unsaturated.

[0081] As used herein, the term "amine protecting group" means any group known in the art of organic synthesis for the protection of amine groups which is stable to an ester reducing agent, a disubstituted hydrazine, R4-M and R7-M, a nucleophile, a hydrazine
10 reducing agent, an activator, a strong base, a hindered amine base and a cyclizing agent. Such amine protecting groups fitting these criteria include those listed in Wuts, P.G.M. et al., *Protecting Groups in Organic Synthesis*, 4th Edition, Wiley (2007) and *The Peptides: Analysis, Synthesis, Biology*, Vol. 3, Academic Press, New York (1981), the disclosure of which is hereby incorporated by reference. Examples of amine protecting groups include,
15 but are not limited to, the following: (1) acyl types such as formyl, trifluoroacetyl, phthalyl, and p-toluenesulfonyl; (2) aromatic carbamate types such as benzyloxycarbonyl (Cbz) and substituted benzyloxycarbonyls, 1-(p-biphenyl)-1-methylethoxycarbonyl, and 9-fluorenylmethyloxycarbonyl (Fmoc); (3) aliphatic carbamate types such as *tert*-butyloxycarbonyl (Boc), ethoxycarbonyl, diisopropylmethoxycarbonyl, and
20 allyloxycarbonyl; (4) cyclic alkyl carbamate types such as cyclopentyloxycarbonyl and adamantyloxycarbonyl; (5) alkyl types such as triphenylmethyl and benzyl; (6) trialkylsilane such as trimethylsilane; (7) thiol containing types such as phenylthiocarbonyl and dithiasuccinoyl; and (8) alkyl types such as triphenylmethyl, methyl, and benzyl; and substituted alkyl types such as 2,2,2-trichloroethyl,
25 2-phenylethyl, and *t*-butyl; and trialkylsilane types such as trimethylsilane.

[0082] As referred to herein, the term "substituted" means that at least one hydrogen atom is replaced with a non-hydrogen group, provided that normal valencies are maintained and that the substitution results in a stable compound. Ring double bonds, as used herein, are double bonds that are formed between two adjacent ring atoms (*e.g.*,
30 C=C, C=N, or N=N).

[0083] In cases wherein there are nitrogen atoms (*e.g.*, amines) on compounds of the present invention, these may be converted to N-oxides by treatment with an oxidizing

agent (*e.g.*, mCPBA and/or hydrogen peroxides) to afford other compounds of this invention. Thus, shown and claimed nitrogen atoms are considered to cover both the shown nitrogen and its N-oxide (N→O) derivative.

[0084] When any variable occurs more than one time in any constituent or formula
5 for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-3 R, then said group may optionally be substituted with up to three R groups, and at each occurrence R is selected independently from the definition of R.

[0085] When a bond to a substituent is shown to cross a bond connecting two atoms
10 in a ring, then such substituent may be bonded to any atom on the ring. When a substituent is listed without indicating the atom in which such substituent is bonded to the rest of the compound of a given formula, then such substituent may be bonded via any atom in such substituent.

[0086] Combinations of substituents and/or variables are permissible only if such
15 combinations result in stable compounds.

[0087] The phrase "pharmaceutically acceptable" is employed herein to refer to those
compounds, materials, compositions, and/or dosage forms that are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, and/or other problem or
20 complication, commensurate with a reasonable benefit/risk ratio.

[0088] Compounds of the present invention can form salts which are also within the
scope of this invention. Unless otherwise indicated, reference to an inventive compound is understood to include reference to one or more salts thereof. Pharmaceutically acceptable salts are preferred. However, other salts may be useful, *e.g.*, in isolation or
25 purification steps which may be employed during preparation, and thus, are contemplated within the scope of the invention.

[0089] As used herein, "pharmaceutically acceptable salts" refer to derivatives of the
disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited
30 to, mineral or organic acid salts of basic groups such as amines; and alkali or organic salts of acidic groups such as carboxylic acids. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent

compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, and nitric; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, and isethionic, and the like.

[0090] The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound that contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Allen, L.V., Jr., ed., *Remington: The Science and Practice of Pharmacy*, 22nd Edition, Pharmaceutical Press, London, UK (2012), the disclosure of which is hereby incorporated by reference.

[0091] In addition, compounds of formula I may have prodrug forms. Any compound that will be converted *in vivo* to provide the bioactive agent (*i.e.*, a compound of formula I) is a prodrug within the scope and spirit of the invention. Various forms of prodrugs are well known in the art. For examples of such prodrug derivatives, see:

a) Bundgaard, H., ed., *Design of Prodrugs*, Elsevier (1985), and Widder, K. et al., eds., *Methods in Enzymology*, 112:309-396, Academic Press (1985);

b) Bundgaard, H., Chapter 5, "Design and Application of Prodrugs", Krosgaard-Larsen, P. et al., eds., *A Textbook of Drug Design and Development*, pp.

113-191, Harwood Academic Publishers (1991);

c) Bundgaard, H., *Adv. Drug Deliv. Rev.*, 8:1-38 (1992);

d) Bundgaard, H. et al., *J. Pharm. Sci.*, 77:285 (1988);

e) Kakeya, N. et al., *Chem. Pharm. Bull.*, 32:692 (1984);

f) Rautio, J. et al., *Nature Rev. Drug Discovery*, 7: 255-270 (2008), and

g) Rautio, J., ed., *Prodrugs and Targeted Delivery (Methods and Principles in Medicinal Chemistry)*, Vol. 47, Wiley-VCH (2011).

- [0092] Compounds containing a carboxy group can form physiologically hydrolyzable esters that serve as prodrugs by being hydrolyzed in the body to yield formula I compounds *per se*. Such prodrugs are preferably administered orally since hydrolysis in many instances occurs principally under the influence of the digestive enzymes. Parenteral administration may be used where the ester *per se* is active, or in those instances where hydrolysis occurs in the blood. Examples of physiologically hydrolyzable esters of compounds of formula I include C₁₋₆alkyl, C₁₋₆alkylbenzyl, 4-methoxybenzyl, indanyl, phthalyl, methoxymethyl, C₁₋₆alkanoyloxy-C₁₋₆alkyl (*e.g.*, acetoxymethyl, pivaloyloxymethyl or propionyloxymethyl), C₁₋₆alkoxycarbonyloxy-C₁₋₆alkyl (*e.g.*, methoxycarbonyl-oxymethyl or ethoxycarbonyloxymethyl, glycyloxymethyl, phenylglycyloxymethyl, (5-methyl-2-oxo-1,3-dioxolen-4-yl)-methyl), and other well known physiologically hydrolyzable esters used, for example, in the penicillin and cephalosporin arts. Such esters may be prepared by conventional techniques known in the art.
- [0093] Preparation of prodrugs is well known in the art and described in, for example, King, F.D., ed., *Medicinal Chemistry: Principles and Practice*, The Royal Society of Chemistry, Cambridge, UK (Second Edition, reproduced, 2006); Testa, B. et al., *Hydrolysis in Drug and Prodrug Metabolism. Chemistry, Biochemistry and Enzymology*, VCHA and Wiley-VCH, Zurich, Switzerland (2003); Wermuth, C.G., ed., *The Practice of Medicinal Chemistry*, Third Edition, Academic Press, San Diego, CA (2008).
- [0094] The present invention is intended to include all isotopes of atoms occurring in the present compounds. isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include deuterium and tritium. isotopes of carbon include ¹³C and ¹⁴C. isotopically-labeled compounds of the invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described herein, using an appropriate isotopically-labeled reagent in place of the non-labeled reagent otherwise employed.
- [0095] The term "solvate" means a physical association of a compound of this invention with one or more solvent molecules, whether organic or inorganic. This physical association includes hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in

the crystal lattice of the crystalline solid. The solvent molecules in the solvate may be present in a regular arrangement and/or a non-ordered arrangement. The solvate may comprise either a stoichiometric or nonstoichiometric amount of the solvent molecules.

"Solvate" encompasses both solution-phase and isolable solvates. Exemplary solvates

5 include, but are not limited to, hydrates, ethanolates, methanolates, and isopropanolates.

Methods of solvation are generally known in the art.

[0096] Abbreviations as used herein, are defined as follows: "1 x" for once, "2 x" for twice, "3 x" for thrice, "°C" for degrees Celsius, "eq" for equivalent or equivalents, "g" for gram or grams, "mg" for milligram or milligrams, "L" for liter or liters, "mL" for milliliter or milliliters, "μL" for microliter or microliters, "N" for normal, "M" for molar, "mmol" for millimole or millimoles, "min" for minute or min, "h" for hour or h, "rt" for room temperature, "RT" for retention time, "atm" for atmosphere, "psi" for pounds per square inch, "conc." for concentrate, "aq" for "aqueous", "sat" or "sat'd" for saturated, "MW" for molecular weight, "mp" for melting point, "MS" or "Mass Spec" for mass spectrometry, "ESI" for electrospray ionization mass spectroscopy, "HR" for high resolution, "HRMS" for high resolution mass spectrometry, "LCMS" for liquid chromatography and mass spectrometry, "HPLC" for high pressure liquid chromatography, "[M-H]" for parent mass minus a proton, "RP HPLC" for reverse phase HPLC, "TLC" or "tlc" for thin layer chromatography, "NMR" for nuclear magnetic resonance spectroscopy, "nOe" for nuclear Overhauser effect spectroscopy, "¹H" for proton, "δ" for delta, "s" for singlet, "d" for doublet, "t" for triplet, "q" for quartet, "m" for multiplet, "br" for broad, "Hz" for hertz, and "α", "β", "R", "S", "E", and "Z" are stereochemical designations familiar to one skilled in the art.

Me	methyl
Et	ethyl
Pr	propyl
<i>i</i> -Pr	isopropyl
Bu	butyl
<i>i</i> -Bu	isobutyl
<i>t</i> -Bu	<i>tert</i> -butyl
Ph	phenyl

Bn	benzyl
Hex	hexanes
MeOH	methanol
EtOH	ethanol
i-PrOH or IPA	isopropanol
AcOH or HOAc	acetic acid
Ag ₂ CO ₃	silver carbonate
AgOAc	silver acetate
AgOTf	silver triflate
CDCl ₃	deutero-chloroform
CHCl ₃	chloroform
cDNA	complimentary DNA
Cu(OTf)	copper triflate
DMF	dimethyl formamide
DMSO	dimethyl sulfoxide
EDC or EDCI	1-ethyl-3-(3-dimethylaminopropyl) carbodiimide
EDTA	ethylenediaminetetraacetic acid
EtOAc	ethyl acetate
Et ₂ O	diethyl ether
AlCl ₃	aluminum chloride
Boc	<i>tert</i> -butoxycarbonyl
CH ₂ Cl ₂	dichloromethane
CH ₃ CN or ACN	acetonitrile
Cs ₂ CO ₃	cesium carbonate
HCl	hydrochloric acid
H ₂ SO ₄	sulfuric acid
K ₂ CO ₃	potassium carbonate
KCN	potassium cyanide
mCPBA or m-CPBA	<i>meta</i> -chloroperbenzoic acid
Pd/C	palladium on carbon
PhSO ₂ Cl	benzenesulfonyl chloride

<i>i</i> -Pr ₂ NEt	diisopropylethylamine
PS	polystyrene
SiO ₂	silica oxide/silica gel
SnCl ₂	tin(II) chloride
TEA	triethylamine
TFA	trifluoroacetic acid
TFAA	trifluoroacetic anhydride
THF	tetrahydrofuran
TMSCHN ₂	trimethylsilyldiazomethane
TMSI	trimethylsilyl iodide
KOAc	potassium acetate
MgSO ₄	magnesium sulfate
MsCl	methanesulfonyl chloride
MsOH or MSA	methylsulfonic acid
NaCl	sodium chloride
NaH	sodium hydride
NaHCO ₃	sodium bicarbonate
NaOH	sodium hydroxide
Na ₂ SO ₃	sodium sulfite
Na ₂ SO ₄	sodium sulfate
NH ₃	ammonia
NH ₄ Cl	ammonium chloride
NH ₄ OH	ammonium hydroxide

[0097] The compounds of the present invention can be prepared in a number of ways known to one skilled in the art of organic synthesis. The compounds of the present invention can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or by variations thereon as appreciated by those skilled in the art. Preferred methods include, but are not limited to, those described below. The reactions are performed in a solvent or solvent mixture appropriate to the reagents and materials employed and suitable for the transformations

being effected. It will be understood by those skilled in the art of organic synthesis that the functionality present on the molecule should be consistent with the transformations proposed. This will sometimes require a judgment to modify the order of the synthetic steps or to select one particular process scheme over another in order to obtain a desired compound of the invention.

[0098] The novel compounds of this invention may be prepared using the reactions and techniques described in this section. Also, in the description of the synthetic methods described below, it is to be understood that all proposed reaction conditions, including choice of solvent, reaction atmosphere, reaction temperature, duration of the experiment and workup procedures, are chosen to be the conditions standard for that reaction, which should be readily recognized by one skilled in the art. Restrictions to the substituents that are compatible with the reaction conditions will be readily apparent to one skilled in the art and alternate methods must then be used.


SYNTHESIS

[0099] The compounds of Formula (I) may be prepared by the exemplary processes described in the following schemes and working examples, as well as relevant published literature procedures that are used by one skilled in the art. Exemplary reagents and procedures for these reactions appear hereinafter and in the working examples.

Protection and de-protection in the processes below may be carried out by procedures generally known in the art (see, for example, Wuts, P.G.M. et al., *Protecting Groups in Organic Synthesis*, 4th Edition, Wiley (2007)). General methods of organic synthesis and functional group transformations are found in: Trost, B.M. et al., eds., *Comprehensive Organic Synthesis: Selectivity, Strategy & Efficiency in Modern Organic Chemistry*, Pergamon Press, New York, NY (1991); Smith, M.B. et al., *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 6th Edition, Wiley & Sons, New York, NY (2007); Katritzky, A.R. et al., eds., *Comprehensive Organic Functional Groups Transformations II*, 2nd Edition, Elsevier Science Inc., Tarrytown, NY (2004); Larock, R.C., *Comprehensive Organic Transformations*, VCH Publishers, Inc., New York, NY (1999), and references therein.


- [00100] A general synthesis of compounds of Formula 1 is described in Scheme 1. The alcohol 3 is alkylated with alkyl halide 2 under silver triflate-mediated conditions (Burk et al., *Tetrahedron Lett.*, 35:8111 (1994)), followed by hydrolysis with a base (*i.e.*, 1 N NaOH or 1 N LiOH) to afford the acid 1. Further derivatization of the acid 1 to amides 4 and acylsulfonamides 5 can be carried out with standard literature conditions, such as: (1) the use of oxalyl chloride with catalytic DMF to form the acid chloride intermediate followed by subsequent reaction with amines in the presence of pyridine or triethylamine; or (2) the treatment of a mixture of the acid 1 and an amine or alkylsulfonamide (or arylsulfonamide or sulfamide) with a coupling reagent such as BOP/Et₃N, EDCI/HOAt/Et₃N, or DEPBT (Li et al., *Org. Lett.*, 1:91 (1999)).

Scheme 1



- [00101] The alkyl halides 2 can be prepared from their corresponding alcohols according to literature procedures. Scheme 2 depicts methods to synthesize the alkyl halides 2a and the corresponding homologated halide 2b.


Scheme 2



[00102] Schemes 3 to 8 illustrate the syntheses of the intermediated alcohols **3a-e**.

- 5 Reduction of acid **8** (Adcock et al., *J. Org. Chem.*, 49:1387 (1984)) with BH_3 -THF provides the corresponding alcohol, which is treated with PPh_3 and I_2 to afford the iodide **9**. Nickel[0]-mediated conjugate addition of alkyl iodide **9** to methyl acrylate affords the ester **10** (Sustmann et al., *Tetrahedron Lett.*, 30:689 (1989)). Reaction of **10** with trimethylsilyl iodide (TMSI) provides a mixture of alcohol **3a** and acid **11**, which is
- 10 converted to the **3a** upon treatment with trimethylsilyldiazomethane.

Scheme 3




- 15 [00103] Alternatively, **3a** and **3b** can be synthesized from the acid **12** (synthesized from 4-hydroxybicyclo[2.2.1]heptane-1-carboxylic acid (prepared according to the procedure of Adcock et al., *J. Org. Chem.*, 49:1387 (1984)) as shown in Scheme 4. Reduction of acid **12** with BH_3 -THF provides the corresponding alcohol, which is oxidized with Dess-Martin periodinane to afford aldehyde **13**. Wittig reaction with methyl 2-(triphenylphosphoranylidene)acetate and then hydrogenation affords the ester
- 20 **14**. Methyl ester cleavage with LiI /pyridine provides the acid **15**, which is converted to

its one carbon homolog **16** by an Arndt-Eistert reaction (Bachmann, W.E. et al., "The Arndt-Eistert Reaction", *Org. React.*, 1:38 (1942)). Hydrolysis of the benzoyl ester followed by esterification of the acid (with TMSCHN₂) furnishes the methyl ester **3a**.

Similarly, the acid **15** can also be converted to intermediate **3b** using an identical


5 sequence.

Scheme 4




- 10 [00104] Alternatively, compounds **20** and **21** can also be synthesized by the sequence described in Scheme 5. Borane reduction of the acid **15** provides an alcohol, which is protected with a THP group. Further hydrolysis of the 3,5-difluorobenzoyl ester with $\text{LiOH-H}_2\text{O}$ affords alcohol **17**. The alcohol **17** can be alkylated with the conditions described in Scheme 1 ($\text{ArCH}_2\text{X}/\text{AgOTf}/2,6\text{-di-}t\text{-butylpyridine}$) or ($\text{NaH}/\text{ArCH}_2\text{X}$) to
- 15 provide ether **18**. The THP group is then removed to give the primary alcohol **19**, which is oxidized to the acid **20** in a two-step sequence (Dess-Martin periodinane oxidation followed by sodium chlorite oxidation). One-carbon homologation of acid **20** using the Arndt-Eistert reaction sequence as described previously provides acid **21**.

Scheme 5



- 5 [00105] Scheme 6 illustrates the synthesis of the hydroxy cyclopropane-ester **3c**. Alcohol **22** is alkylated with an alkyl halide such as 1-(chloromethyl)-3-phenoxybenzene and the resulting ester is reduced to the corresponding alcohol **23**. The alcohol is then converted to the nitrile **24** by a two-step sequence (conversion to mesylate, followed by NaCN displacement of the mesylate). Nitrile **24** is reduced (DIBAL-H) to the
- 10 corresponding aldehyde **25**, which then undergoes a Horner-Emmons reaction with an appropriate phosphonate-ester to give the α,β -unsaturated ester **26**. The α,β -unsaturated ester **26** is cyclopropanated with diazomethane to give the corresponding α,β -cyclopropyl ester, which is then deprotected (hydrogenolysis conditions) to give the alcohol **3c**.


Scheme 6



[00106] Scheme 7 illustrates the synthesis of the alcohol intermediate **3d**. Alcohol **22**


- 5 is alkylated with 1-(chloromethyl)-2-phenoxybenzene and the resulting ether-ester is reduced to the corresponding alcohol **27**. The alcohol is then alkylated with *tert*-butyl 2-bromoacetate under basic conditions to give the corresponding ester which is hydrolyzed to the acid **28**. Esterification of the acid **28** with trimethylsilyldiazomethane and subsequent hydrogenolysis of the benzyl ether affords the alcohol intermediate **3d**.

Scheme 7



- [00107] Scheme 8 illustrates the syntheses of the alcohol intermediate **3e**. The acid **8**
- 5 is converted to the ketone **29** through the reaction of the corresponding acid chloride with (3,5-difluorophenyl)magnesium bromide in the presence of acetylacetonate iron (III) salt. Baeyer-Villiger oxidation of the ketone **29** with hydrogen peroxide provides the methyl ester **30**. Treatment of **30** with iodotrimethylsilane furnishes the alcohol **31**. This alcohol is alkylated with methyl 3-bromopropionate to give the corresponding diester, which is
- 10 hydrolyzed under basic conditions to give the acid-alcohol **32**. Further treatment with trimethylsilyldiazomethane affords the alcohol intermediate **3e**.

Scheme 8



IV. BIOLOGY

- 5 **[00108]** Diabetes mellitus is a serious disease afflicting over 100 million people worldwide. It is diagnosed as a group of disorders characterized by abnormal glucose homeostasis resulting in elevated blood glucose. Diabetes is a syndrome with interrelated metabolic, vascular, and neuropathic components. The metabolic abnormality is generally characterized by hyperglycemia and alterations in carbohydrate, fat and protein
- 10 metabolism caused by absent or reduced insulin secretion and/or ineffective insulin secretion. The vascular syndrome consists of abnormalities in the blood vessels leading to cardiovascular, retinal and renal complications. Abnormalities in the peripheral and autonomic nervous systems are also part of diabetic syndrome. Strikingly, diabetes is the fourth leading cause of global death by disease, the largest cause of kidney failure in
- 15 developed countries, the leading cause of vision loss in industrialized countries and has the greatest prevalence increase in developing countries.

- [00109] Type 2 diabetes, which accounts for 90% of diabetes cases, is characterized by increasing insulin resistance associated with inadequate insulin secretion after a period of compensatory hyperinsulinemia. The reasons for β cell secondary failure are not completely understood. Acquired pancreatic islet damage or exhaustion and/or genetic factors causing susceptibility to islet secretory insufficiency have been hypothesized.
- [00110] Recently, five GPCRs (FFAR1 (GPR40), FFAR2 (GPR43), FFAR3 (GPR41), GPR84, and GPR120) were reported to recognize free fatty acids FFAR1, recognizes medium-long chainfatty acids like palmitic acid and linoleic acid FFAR2 and FFAR3 recognize short-chain fatty acids like acetate and butyrate whereas GPR84 recognizes medium-chain fatty acid like lauric acid. GPR120 recognizes long-chain fatty acids, especially EPA and DHA (Im, *Progress in Lipid Research*, 51 232-237(2012)). GPR120 has been detected in macrophages, dendritic cells, adipocytes, clara cells in bronchiole epithelium, and enteroendocrine L cells in colon (Miyauchi et al., *Naunyn-Schmiedebergs Arch Pharmacol.*, 379:427-434 (2009)). The anti-inflammatory mechanism of omega-3 fatty acids using GPR120 knock-out mice was investigated (Oh et al., *Cell*, 142:687-698 (2010)). They suggested GPR120 activation by DHA interacts with TAB1 via b-arrestin-2, and that this interaction interrupts TAK1 activation by LPS or TNF-alpha, suppressing inflammatory responses via NF- κ B and JNK in macrophages and dendritic cells (Oh et al., *Cell*, 142:687-698 (2010)). Furthermore, GPR120 activation was shown to enhance insulin-induced glucose uptake in adipose tissues through Gq/11 proteins and PI 3-kinase.
- [00111] Similarly, GPR120-deficient mice fed a high-fat diet develop obesity, glucose intolerance and fatty liver with decreased adipocyte differentiation and lipogenesis and enhanced hepatic lipogenesis (Ichimura et al., *Nature*, 483(7389):350-354 (2012). Insulin resistance in such mice was shown to be associated with reduced insulin signalling and enhanced inflammation in adipose tissue. In humans, GPR120 expression in adipose tissue was shown to be significantly higher in obese individuals than in lean controls. GPR120 gene sequencing in obese subjects revealed a deleterious non-synonymous mutation (p.R270H) that inhibits GPR120 signalling activity. Furthermore, the p.R270H variant was associated with increased risk of obesity in European populations.
- [00112] Given the increase in the worldwide patient population afflicted by type 2 diabetes, there is a need for novel therapies which are effective with minimal adverse events. To decrease medical burden of type 2 diabetes through enhanced glycemic

control, GPR120 modulator compounds of the present invention are being investigated here for their ability to increase glucose tolerance as well as the potential combination with a broad range of anti-diabetic drugs.

[00113] The term "modulator" refers to a chemical compound with capacity to either
5 enhance (*e.g.*, "agonist" activity) or partially enhance (*e.g.*, "partial agonist" activity) or inhibit (*e.g.*, "antagonist" activity or "inverse agonist" activity) a functional property of biological activity or process (*e.g.*, enzyme activity or receptor binding); such enhancement or inhibition may be contingent on the occurrence of a specific event, such as activation of a signal transduction pathway, receptor internalization, and/or may be
10 manifest only in particular cell types.

[00114] It is also desirable and preferable to find compounds with advantageous and improved characteristics compared with known anti-diabetic agents, in one or more of the following categories that are given as examples, and are not intended to be limiting:
(a) pharmacokinetic properties, including oral bioavailability, half life, and clearance;
15 (b) pharmaceutical properties; (c) dosage requirements; (d) factors that decrease blood drug concentration peak-to-trough characteristics; (e) factors that increase the concentration of active drug at the receptor; (f) factors that decrease the liability for clinical drug-drug interactions; (g) factors that decrease the potential for adverse side-effects, including selectivity versus other biological targets; and (h) improved therapeutic
20 index with less propensity for hypoglycemia.

[00115] As used herein, the term "patient" encompasses all mammalian species.

[00116] As used herein, the term "subject" refers to any human or non-human organism that could potentially benefit from treatment with a GPR120 modulator. Exemplary subjects include human beings of any age with risk factors for metabolic
25 disease. Common risk factors include, but are not limited to, age, sex, weight, family history, or signs of insulin resistance such as acanthosis nigricans, hypertension, dislipidemia, or polycystic ovary syndrome (PCOS).

[00117] As used herein, "treating" or "treatment" cover the treatment of a disease-state in a mammal, particularly in a human, and include: (a) inhibiting the disease-state, *i.e.*,
30 arresting its development; (b) relieving the disease-state, *i.e.*, causing regression of the disease state; and/or (c) preventing the disease-state from occurring in a mammal, in

particular, when such mammal is predisposed to the disease-state but has not yet been diagnosed as having it.

[00118] As used herein, "preventing" or "prevention" cover the preventive treatment (*i.e.*, prophylaxis and/or risk reduction) of a subclinical disease-state in a mammal, particularly in a human, aimed at reducing the probability of the occurrence of a clinical disease-state. Patients are selected for preventative therapy based on factors that are known to increase risk of suffering a clinical disease state compared to the general population. "Prophylaxis" therapies can be divided into (a) primary prevention and (b) secondary prevention. Primary prevention is defined as treatment in a subject that has not yet presented with a clinical disease state, whereas secondary prevention is defined as preventing a second occurrence of the same or similar clinical disease state. As used herein, "risk reduction" covers therapies that lower the incidence of development of a clinical disease state. As such, primary and secondary prevention therapies are examples of risk reduction.

[00119] "Therapeutically effective amount" is intended to include an amount of a compound of the present invention that is effective when administered alone or in combination to modulate GPR120 and/or to prevent or treat the disorders listed herein. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the preventive or therapeutic effect, whether administered in combination, serially, or simultaneously.

[00120] GPR120 activity was monitored by measuring phosphorylation of ERK (pERK), since G protein receptors are known to activate the ERK signaling cascade either directly and/or through recruitment of arrestin that serves as a scaffold for downstream signaling events. Molecules that activated GPR120 with sufficient potency and efficacy in the pERK assay that also possessed desirable pharmacokinetic properties were evaluated in mice for glucose lowering by monitoring disposition of an oral glucose load by an oral glucose tolerance test (oGTT).

GPR120 pERK AlphaScreen SureFire Assay

[00121] The human and mouse GPR120-mediated intracellular phosphorylated ERK assays were established using CHO A12 cells stably transfected with the short form of human or mouse GPR120 receptor. Cells were cultured in growth medium consisting of

F-12 media (Invitrogen Cat. #11765) with 5% Charcoal/Dextran FBS (Invitrogen Cat. #12676-029), 500µg/mL GENETICIN® (Life Technologies Cat. #10131-027) and 250µg/mL Zeocin (Invitrogen Cat. #R250-01). Cells were cryo preserved at a concentration of 2×10^7 cells/mL, in 90% Charcoal/Dextran FBS and 10% DMSO, and frozen in liquid nitrogen at a low passage number.

[00122] For the pERK assay, 2×10^7 cells/mL cryopreserved human and mouse cells were thawed rapidly in a 37 °C water bath and added to a T-225 flask containing 50 mL growth medium. The flasks were placed in a tissue culture incubator overnight (37 °C, 5%CO₂). The next day, cells were harvested with trypsin (Gibco Cat. #25300-054), resuspended in serum-containing growth medium and counted using a Cellometer and volume adjusted to a concentration of 0.6×10^6 cells/mL. Cells were plated into 384-well clear bottom tissue culture plates (BD Cat. #353962) at 50 µL/well, for a density of 30,000 cells/well using a MULTIDROP® and incubated for 16-18 hours (overnight) at 37 °C with 5% CO₂. The next day, cells were serum starved in 30 µL of F-12 media without any serum or antibiotics for 2 hours at 37 °C.

[00123] Test compounds were 3-fold, 11-point serially diluted in DMSO in a REMP assay plate (Matrix Cat. #4307) by Tecan and 5 µL was transferred into an ECHO source plate (LabCyte Cat. #LC-0200). Cells were then stimulated with 50 nL of compound dilutions using ECHO liquid handler for 7 minutes at 37 °C. Compounds ranged from final assay concentrations of 33.33 µM to 0.56 nM.

[00124] The media was then dumped and cells lysed with 20 µL of 1x Lysis buffer from the AlphaScreen SureFire Phospho-ERK 1/2 Kit (Perkin Elmer Cat. #6760617M). The lysis buffer was diluted 5-fold with water before use. The plate was agitated on a shaker for 10 minutes after which 2 µL was transferred into a 384-well white proxiplate (Perkin Elmer Cat. #6008289). The SureFire assay reagent mix was prepared by mixing 60 parts Reaction Buffer, 10 parts Activation Buffer, 1 part Donor Beads, 1 part Acceptor Beads (Perkin Elmer Cat. #TGRES10K). 3.5 µL/well of this reagent mix was manually added to the proxiplate with a multichannel pipettor. Plates were spun down at 1000 rpm for 2 minutes, followed by light-protected incubation at room temperature for 2 hours. The plates were read on the Alpha-technology compatible Envision multilabel plate reader using AlphaScreen protocol according to manufacturer's specifications. The agonist effect of compounds was expressed as 100 x (average sample-average

blank)/(average total- average blank) where sample is the luminescence activity in the presence of test compound, blank is equal to the luminescence activity in the presence of DMSO control and the total is signal induced by 50 μ M linolenic acid as reference compound. Activation data for the test compound over a range of concentrations was plotted as percentage activation of the test compound (100% = maximum response). After correcting for background, EC₅₀ values were determined. The EC₅₀ is defined as the concentration of test compound which produces 50% of the maximal response and was quantified using the 4 parameter logistic equation to fit the data.

10 [00125] The human and mouse GPR120-mediated intracellular phosphorylated ERK assays were also established using CHO-K1 cells stably transfected with the short form of human or mouse GPR120 receptor. Cells were cultured in growth medium consisting of F-12 media (Invitrogen Cat. #11765) with 5% Charcoal/Dextran FBS (Invitrogen Cat. #12676-029) and 500 μ g/mL GENETICIN® (Life Technologies Cat. #10131-027). Cells were cryo preserved at a concentration of 3x10⁶ cells/mL, in 70% F-12, 20% Charcoal/Dextran FBS and 10% DMSO, and frozen in liquid nitrogen at a low passage number.

[00126] For the pERK assay, 3x10⁶ cells/mL cryopreserved human and mouse cells were thawed rapidly in a 37 °C water bath and added to a T-225 flask containing 50 mL growth medium. The flasks were placed in a tissue culture incubator overnight (37 °C, 5%CO₂). The next day, cells were harvested with trypsin (Gibco Cat. #25300-054), resuspended in serum-containing growth medium and counted using a Cellometer and volume adjusted to a concentration of 0.5x10⁶cells/mL. Cells were plated into 384-well clear bottom tissue culture plates (BD Cat. #353962) at 50 μ L/well, for a density of 25,000 cells/well using a MULTIDROP® and incubated for 16-18 hours (overnight) at 37 °C with 5% CO₂. The next day, cells were washed once with 50 μ L of PBS without Ca⁺⁺/Mg⁺⁺ (Gibco Cat. #14190-036) and serum starved in 25 μ L of F-12 media without any serum or antibiotics for 2 hours at 37 °C.

[00127] Test compounds were 3-fold, 11-point serially diluted in DMSO in a REMP assay plate (Matrix Cat. #4307) by Tecan and 5 μ L was transferred into an ECHO source plate (LabCyte Cat. #LC-0200). Cells were then stimulated with 40 nL of compound

dilutions using ECHO liquid handler for 7 minutes at 37 °C. Compounds ranged from final assay concentrations of 32 μ M to 0.54 nM.

[00128] The media was then dumped and cells lysed with 20 μ L of 1x Lysis buffer from the AlphaScreen SureFire Phospho-ERK 1/2 Kit (Perkin Elmer Cat. #6760617M).

5 The lysis buffer was diluted 5-fold with water before use. The plate was agitated on a shaker for 10 minutes after which 2 μ L was transferred into a 384-well white proxiplate (Perkin Elmer Cat. #6008289). The SureFire assay reagent mix was prepared by mixing 60 parts Reaction Buffer, 10 parts Activation Buffer, 1 part Donor Beads, 1 part Acceptor Beads (Perkin Elmer Cat. #TGRES10K). 3.5 μ L/well of this reagent mix was manually
10 added to the proxiplate with a multichannel pipettor. Plates were spun down at 1000 rpm for 2 minutes, followed by light-protected incubation at room temperature for 2 hours. The plates were read on the Alpha-technology compatible Envision multilabel plate reader using AlphaScreen protocol according to manufacturer's specifications. The agonist effect of compounds was expressed as 100 x (average sample-average
15 blank)/(average total- average blank) where sample is the luminescence activity in the presence of test compound, blank is equal to the luminescence activity in the presence of DMSO control and the total is signal induced by 50 μ M linolenic acid as reference compound.

20 [00129] Activation data for the test compound over a range of concentrations was plotted as percentage activation of the test compound (100% = maximum response). After correcting for background, EC₅₀ values were determined. The EC₅₀ is defined as the concentration of test compound which produces 50% of the maximal response and was quantified using the 4 parameter logistic equation to fit the data.

25

[00130] The exemplified Examples disclosed below were tested in the GPR120 *in vitro* assays described above and were found having GPR120 agonist activity. Table 1 below lists the EC₅₀ values measured in the human GPR120 pERK assay for the following examples.

30

Example Number	(EC ₅₀ μ M)
1	1.56
2	1.82
3	0.53
4	0.65
5	1.02
6	0.45
7	0.23
8	2.34
9	0.60
10	2.64
11	1.09
12	6.72
13	7.67
14	0.27
15	0.36
16	0.64
17	0.25
18	0.25
19	0.52
20	0.71
21	1.92
22	0.73
23	1.15
24	2.80
25	0.64
26	0.62
27	1.56
28	0.39
29	5.07
30	0.85

Example Number	(EC ₅₀ μM)
31	0.90
32	0.59
33	0.26
34	0.40
35	0.44
36	1.55
37	3.12
38	1.68
39	0.82
40	2.30
41	1.10
42	9.10
43	0.32
44	0.93
45	0.71
46	0.29
47	7.81
48	0.32
49	0.66
50	0.54
51	2.39
52	0.69
53	6.12
54	1.69
55	0.17
56	1.88
57	0.77
58	1.72
59	0.61
60	4.15

Example Number	(EC ₅₀ μ M)
61	0.75
62	0.81
63	2.47
64	1.51
65	1.27
66	0.37
67	4.96
68	1.00
69	0.87
70	7.25
71	6.64
72	1.12
73	1.62
74	1.43
75	1.85
76	3.04
77	4.53
78	0.94
79	0.27
80	2.75
81	0.35
82	0.98
83	0.40
84	3.03
85	2.40
86	6.99
87	1.31
88	2.64
89	2.64
90	1.22

Example Number	(EC ₅₀ μ M)
91	0.28
92	0.60
93	0.88
94	0.63
95	0.51
96	0.38
97	2.48
98	2.68
99	6.36
100	0.57

In vivo GPR120 Assays

1) Acute Oral Glucose Tolerance Test

[00131] C57BL/6 mice were housed individually and fed a standard rodent chow diet.

- 5 At approximately 11 weeks age, after a 5 h fast, these mice were orally treated with vehicle or test compounds 60 min before a glucose challenge (2 g/kg). Blood glucose levels were determined from tail bleeds taken at -60, 0, 15, 30, 60 and 120 min after the glucose challenge. The blood glucose excursion profile from t = 0-120 min was used to calculate an area under the curve (AUC) for compound treatment. This AUC for
- 10 compound treatment is compared to vehicle treatment.

[00132] In an oral glucose tolerance test in mice at dose of 30 mg/kg, Examples 73 and 92 reduced glucose AUC levels by 21% and 25% respectively.

2) Acute Intraperitoneal Insulin Tolerance Test

- 15 **[00133]** C57BL/6 mice were housed individually and fed a standard rodent chow diet. At approximately 11 weeks age, after 5 h fast, these mice were orally treated with vehicle or test compounds 30 min before an insulin challenge (0.1 U/kg). Blood glucose levels were determined from tail bleeds taken at -30, 0, 15, 30, 60, 90 and 120 min after insulin injection. The blood glucose excursion profile from t = 0-120 min was used to calculate a
- 20 negative area under the curve (AUC) for compound treatment. This AUC for compound treatment is compared to vehicle treatment.

[00134] The compounds of the present invention possess activity as modulators of GPR120, and, therefore, may be used in the treatment of diseases associated with GPR120 activity. Via modulation of GPR120, the compounds of the present invention
5 may preferably be employed to modulate the production/secretion of insulin and/or gut hormones, such as GLP-1, GIP, PYY, CCK and amylin.

[00135] Accordingly, the compounds of the present invention can be administered to mammals, preferably humans, for the treatment of a variety of conditions and disorders, including, but not limited to, treating, preventing, or slowing the progression of diabetes
10 and related conditions, microvascular complications associated with diabetes, macrovascular complications associated with diabetes, cardiovascular diseases, Metabolic Syndrome and its component conditions, inflammatory diseases and other maladies. Consequently, it is believed that the compounds of the present invention may be used in preventing, inhibiting, or treating diabetes, hyperglycemia, impaired glucose tolerance,
15 gestational diabetes, insulin resistance, hyperinsulinemia, retinopathy, neuropathy, nephropathy, diabetic kidney disease, acute kidney injury, cardiorenal syndrome, delayed wound healing, atherosclerosis and its sequelae (acute coronary syndrome, myocardial infarction, angina pectoris, peripheral vascular disease, intermittent claudication, myocardial ischemia, stroke, heart failure), Metabolic Syndrome, hypertension, obesity,
20 fatty liver disease, dyslipidemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, low HDL, high LDL, lipid disorders, lipodystrophy, liver diseases such as NASH (Non-Alcoholic SteatoHepatitis), NAFLD (Non-Alcoholic Fatty Liver Disease) and liver cirrhosis, and treatment of side-effects related to diabetes.

[00136] Metabolic Syndrome or "Syndrome X" is described in Ford et al., *J. Am. Med. Assoc.*, 287:356-359 (2002) and Arbeeny et al., *Curr. Med. Chem. - Imm., Endoc. & Metab. Agents*, 1:1-24 (2001).

V. PHARMACEUTICAL COMPOSITIONS, FORMULATIONS AND COMBINATIONS

[00137] The compounds of this invention can be administered for any of the uses
30 described herein by any suitable means, for example, orally, such as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders,

granules, elixirs, tinctures, suspensions (including nanosuspensions, microsuspensions, spray-dried dispersions), syrups, and emulsions; sublingually; buccally; parenterally, such as by subcutaneous, intravenous, intramuscular, or intrasternal injection, or infusion techniques (*e.g.*, as sterile injectable aqueous or non-aqueous solutions or suspensions);
5 nasally, including administration to the nasal membranes, such as by inhalation spray; topically, such as in the form of a cream or ointment; or rectally such as in the form of suppositories. They can be administered alone, but generally will be administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

- 10 **[00138]** The term "pharmaceutical composition" means a composition comprising a compound of the invention in combination with at least one additional pharmaceutically acceptable carrier. A "pharmaceutically acceptable carrier" refers to media generally accepted in the art for the delivery of biologically active agents to animals, in particular, mammals, including, *i.e.*, adjuvant, excipient or vehicle, such as diluents, preserving
15 agents, fillers, flow regulating agents, disintegrating agents, wetting agents, emulsifying agents, suspending agents, sweetening agents, flavoring agents, perfuming agents, antibacterial agents, antifungal agents, lubricating agents and dispensing agents, depending on the nature of the mode of administration and dosage forms.

- [00139]** Pharmaceutically acceptable carriers are formulated according to a number of
20 factors well within the purview of those of ordinary skill in the art. These include, without limitation: the type and nature of the active agent being formulated; the subject to which the agent-containing composition is to be administered; the intended route of administration of the composition; and the therapeutic indication being targeted. Pharmaceutically acceptable carriers include both aqueous and non-aqueous liquid media,
25 as well as a variety of solid and semi-solid dosage forms. Such carriers can include a number of different ingredients and additives in addition to the active agent, such additional ingredients being included in the formulation for a variety of reasons, *e.g.*, stabilization of the active agent, binders, etc., well known to those of ordinary skill in the art. Descriptions of suitable pharmaceutically acceptable carriers, and factors involved in
30 their selection, are found in a variety of readily available sources such as, for example, Allen, L.V., Jr. et al., *Remington: The Science and Practice of Pharmacy* (2 Volumes), 22nd Edition, Pharmaceutical Press (2012).

[00140] The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and hepatic function of the patient, and the effect desired.

[00141] By way of general guidance, the daily oral dosage of each active ingredient, when used for the indicated effects, will range between about 0.001 to about 5000 mg per day, preferably between about 0.01 to about 1000 mg per day, and most preferably between about 0.1 to about 250 mg per day. Intravenously, the most preferred doses will range from about 0.01 to about 10 mg/kg/minute during a constant rate infusion.

Compounds of this invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily.

[00142] The compounds are typically administered in admixture with suitable pharmaceutical diluents, excipients, or carriers (collectively referred to herein as pharmaceutical carriers) suitably selected with respect to the intended form of administration, *e.g.*, oral tablets, capsules, elixirs, and syrups, and consistent with conventional pharmaceutical practices.

[00143] Dosage forms (pharmaceutical compositions) suitable for administration may contain from about 1 milligram to about 2000 milligrams of active ingredient per dosage unit. In these pharmaceutical compositions the active ingredient will ordinarily be present in an amount of about 0.1-95% by weight based on the total weight of the composition.

[00144] A typical capsule for oral administration contains at least one of the compounds of the present invention (250 mg), lactose (75 mg), and magnesium stearate (15 mg). The mixture is passed through a 60 mesh sieve and packed into a No. 1 gelatin capsule.

[00145] A typical injectable preparation is produced by aseptically placing at least one of the compounds of the present invention (250 mg) into a vial, aseptically freeze-drying and sealing. For use, the contents of the vial are mixed with 2 mL of physiological saline, to produce an injectable preparation.

[00146] The present invention includes within its scope pharmaceutical compositions comprising, as an active ingredient, a therapeutically effective amount of at least one of the compounds of the present invention, alone or in combination with a pharmaceutical carrier. Optionally, compounds of the present invention can be used alone, in
5 combination with other compounds of the invention, or in combination with one or more other therapeutic agent(s), *e.g.*, an antidiabetic agent or other pharmaceutically active material.

[00147] The compounds of the present invention may be employed in combination with other GPR120 modulators or one or more other suitable therapeutic agents useful in
10 the treatment of the aforementioned disorders including: anti-diabetic agents, anti-hyperglycemic agents, anti-hyperinsulinemic agents, anti-retinopathic agents, anti-neuropathic agents, anti-nephropathic agents, anti-atherosclerotic agents, anti-ischemic agents, anti-hypertensive agents, anti-obesity agents, anti-dyslipidemic agents, anti-hyperlipidemic agents, anti-hypertriglyceridemic agents,
15 anti-hypercholesterolemic agents, anti-restenotic agents, anti-pancreatitis agents, lipid lowering agents, anorectic agents, appetite suppressants, treatments for heart failure, treatments for peripheral arterial disease and anti-inflammatory agents.

[00148] Where desired, the compound of the present invention may be used in combination with one or more other types of antidiabetic agents and/or one or more other
20 types of therapeutic agents which may be administered orally in the same dosage form, in a separate oral dosage form or by injection. The other type of antidiabetic agent that may be optionally employed in combination with the GPR120 receptor modulator of the present invention may be one, two, three or more antidiabetic agents or antihyperglycemic agents which may be administered orally in the same dosage form, in
25 a separate oral dosage form, or by injection to produce an additional pharmacological benefit.

[00149] The antidiabetic agents used in the combination with the compound of the present invention include, but are not limited to, insulin secretagogues or insulin sensitizers, other GPR120 receptor modulators, or other antidiabetic agents. These agents
30 include, but are not limited to, dipeptidyl peptidase IV (DP4) inhibitors (for example, sitagliptin, saxagliptin, linagliptin, alogliptin, vildagliptin), biguanides (for example, metformin, phenformin), sulfonyl ureas (for example, gliburide, glimepiride, glipizide),

glucosidase inhibitors (for example, acarbose, miglitol), PPAR γ agonists such as thiazolidinediones (for example, rosiglitazone, pioglitazone), PPAR α/γ dual agonists (for example, muraglitazar, peliglitazar, tesaglitazar, aleglitazar), glucokinase activators (e.g., PF-04937319 and AMG-151, as well as other compounds described in Fyfe, M.C.T. et al., *Drugs of the Future*, 34(8):641-653 (2009) and incorporated herein by reference), GPR40 receptor modulators (e.g., TAK-875), GPR119 receptor modulators (e.g., MBX-2952, PSN821, APD597), other GPR120 receptor modulators (e.g., compound 43 from *J. Med. Chem.*, 55:4511-4515 (2012)), sodium-glucose transporter-2 (SGLT2) inhibitors (for example dapagliflozin, canagliflozin, remagliflozin), 11 β -HSD-1 inhibitors (for example MK-0736, BI35585, BMS-823778, and LY2523199), amylin analogs such as pramlintide, and/or insulin. Reviews of current and emerging therapies for the treatment of diabetes can be found in: Mohler, M.L. et al., *Medicinal Research Reviews*, 29(1):125-195 (2009), and Mizuno, C.S. et al., *Current Medicinal Chemistry*, 15:61-74 (2008).

[00150] The GPR120 receptor modulator of the present invention may also be optionally employed in combination with agents for treating complication of diabetes. These agents include PKC inhibitors and/or AGE inhibitors.

[00151] The GPR120 receptor modulator of the present invention may also be optionally employed in combination with one or more hypophagic agents such as diethylpropion, phendimetrazine, phentermine, orlistat, sibutramine, lorcaserin, pramlintide, topiramate, MCHR1 receptor antagonists, oxyntomodulin, naltrexone, Amylin peptide, NPY Y5 receptor modulators, NPY Y2 receptor modulators, NPY Y4 receptor modulators, cetilistat, 5HT2c receptor modulators, MGAT2 (monoacylglycerol transferase 2) inhibitors (for example, compounds from WO 2012/124744, or compound (S)-10 from *Bioorg. Med. Chem. Lett.* (2013),

doi: <http://dx.doi.org/10.1016/j.bmcl.2013.02.084>) and the like. The compound of structure I may also be employed in combination with an agonist of the glucagon-like peptide-1 receptor (GLP-1 R), such as exenatide, liraglutide, GLP-1(1-36) amide, GLP-1(7-36) amide, GLP-1(7-37) (as disclosed in U.S. Patent No. 5,614,492 to Habener, the disclosure of which is incorporated herein by reference), which may be administered via injection, intranasal, or by transdermal or buccal devices. Reviews of current and emerging therapies for the treatment of obesity can be found in: Melnikova, I. et al., *Nature Reviews Drug Discovery*, 5:369-370 (2006); Jones, D., *Nature Reviews: Drug*

Discovery, 8:833-834 (2009); Obici, S., *Endocrinology*, 150(6):2512-2517 (2009); and Elangbam, C.S., *Vet. Pathol.*, 46(1):10-24 (2009).

[00152] The above other therapeutic agents, when employed in combination with the compounds of the present invention may be used, for example, in those amounts
5 indicated in the *Physicians' Desk Reference*, as in the patents set out above, or as otherwise determined by one of ordinary skill in the art.

[00153] Particularly when provided as a single dosage unit, the potential exists for a chemical interaction between the combined active ingredients. For this reason, when the compound of the present invention and a second therapeutic agent are combined in a
10 single dosage unit they are formulated such that although the active ingredients are combined in a single dosage unit, the physical contact between the active ingredients is minimized (that is, reduced). For example, one active ingredient may be enteric coated. By enteric coating one of the active ingredients, it is possible not only to minimize the contact between the combined active ingredients, but also, it is possible to control the
15 release of one of these components in the gastrointestinal tract such that one of these components is not released in the stomach but rather is released in the intestines. One of the active ingredients may also be coated with a material that affects a sustained-release throughout the gastrointestinal tract and also serves to minimize physical contact between the combined active ingredients. Furthermore, the sustained-released component can be
20 additionally enteric coated such that the release of this component occurs only in the intestine. Still another approach would involve the formulation of a combination product in which the one component is coated with a sustained and/or enteric release polymer, and the other component is also coated with a polymer such as a low viscosity grade of hydroxypropyl methylcellulose (HPMC) or other appropriate materials as known in the
25 art, in order to further separate the active components. The polymer coating serves to form an additional barrier to interaction with the other component.

[00154] These as well as other ways of minimizing contact between the components of combination products of the present invention, whether administered in a single dosage form or administered in separate forms but at the same time by the same manner, will be
30 readily apparent to those skilled in the art, once armed with the present disclosure.

[00155] The compounds of the present invention can be administered alone or in combination with one or more additional therapeutic agents. By "administered in

combination" or "combination therapy" it is meant that the compound of the present invention and one or more additional therapeutic agents are administered concurrently to the mammal being treated. When administered in combination, each component may be administered at the same time or sequentially in any order at different points in time.

- 5 Thus, each component may be administered separately but sufficiently closely in time so as to provide the desired therapeutic effect.

[00156] The compounds of the present invention are also useful as standard or reference compounds, for example as a quality standard or control, in tests or assays involving the GPR120 receptor. Such compounds may be provided in a commercial kit,
10 for example, for use in pharmaceutical research involving GPR120 or anti-diabetic activity. For example, a compound of the present invention could be used as a reference in an assay to compare its known activity to a compound with an unknown activity. This would ensure the experimenter that the assay was being performed properly and provide a basis for comparison, especially if the test compound was a derivative of the reference
15 compound. When developing new assays or protocols, compounds according to the present invention could be used to test their effectiveness.

[00157] The compounds of the present invention may also be used in diagnostic assays involving GPR120.

[00158] The present invention also encompasses an article of manufacture. As used
20 herein, article of manufacture is intended to include, but not be limited to, kits and packages. The article of manufacture of the present invention, comprises: (a) a first container; (b) a pharmaceutical composition located within the first container, wherein the composition, comprises: a first therapeutic agent, comprising a compound of the present invention or a pharmaceutically acceptable salt form thereof; and, (c) a package
25 insert stating that the pharmaceutical composition can be used for the treatment of multiple diseases or disorders associated with GPR120 (as defined previously). In another embodiment, the package insert states that the pharmaceutical composition can be used in combination (as defined previously) with a second therapeutic agent for the treatment of multiple diseases or disorders associated with GPR120. The article of
30 manufacture can further comprise: (d) a second container, wherein components (a) and (b) are located within the second container and component (c) is located within or outside

of the second container. Located within the first and second containers means that the respective container holds the item within its boundaries.

[00159] The first container is a receptacle used to hold a pharmaceutical composition. This container can be for manufacturing, storing, shipping, and/or individual/bulk selling.

5 First container is intended to cover a bottle, jar, vial, flask, syringe, tube (*e.g.*, for a cream preparation), or any other container used to manufacture, hold, store, or distribute a pharmaceutical product.

[00160] The second container is one used to hold the first container and, optionally, the package insert. Examples of the second container include, but are not limited to,

10 boxes (*e.g.*, cardboard or plastic), crates, cartons, bags (*e.g.*, paper or plastic bags), pouches, and sacks. The package insert can be physically attached to the outside of the first container via tape, glue, staple, or another method of attachment, or it can rest inside the second container without any physical means of attachment to the first container. Alternatively, the package insert is located on the outside of the second container. When
15 located on the outside of the second container, it is preferable that the package insert is physically attached via tape, glue, staple, or another method of attachment. Alternatively, it can be adjacent to or touching the outside of the second container without being physically attached.

[00161] The package insert is a label, tag, marker, etc. that recites information relating

20 to the pharmaceutical composition located within the first container. The information recited will usually be determined by the regulatory agency governing the area in which the article of manufacture is to be sold (*e.g.*, the United States Food and Drug Administration). Preferably, the package insert specifically recites the indications for which the pharmaceutical composition has been approved. The package insert may be
25 made of any material on which a person can read information contained therein or thereon. Preferably, the package insert is a printable material (*e.g.*, paper, plastic, cardboard, foil, adhesive-backed paper or plastic, etc.) on which the desired information has been formed (*e.g.*, printed or applied).

[00162] Other features of the invention will become apparent in the course of the

30 following descriptions of exemplary embodiments that are given for illustration of the invention and are not intended to be limiting thereof.

VI. EXAMPLES

[00163] The following Examples are offered as illustrative, as a partial scope and particular embodiments of the invention and are not meant to be limiting of the scope of the invention. Abbreviations and chemical symbols have their usual and customary meanings unless otherwise indicated. Unless otherwise indicated, the compounds described herein have been prepared, isolated and characterized using the schemes and other methods disclosed herein or may be prepared using the same.

[00164] As appropriate, reactions were conducted under an atmosphere of dry nitrogen (or argon). For anhydrous reactions, DRISOLV[®] solvents from EM were employed. For other reactions, reagent grade or HPLC grade solvents were utilized. Unless otherwise stated, all commercially obtained reagents were used as received.

HPLC/MS and Preparatory/Analytical HPLC Methods Employed in Characterization or Purification of Examples

[00165] Analytical HPLC (unless otherwise noted) was performed to determine compound purity on a Shimadzu SIL-10A using the following method:

HPLC-1: SunFire C18 (4.6 x 150 mm) 3.5 μ , gradient 10 to 100% B:A for 12 min, then 3 min hold at 100% B

Mobile Phase A: 0.05% TFA in water : CH₃CN (95:5)

Mobile Phase B: 0.05% TFA in CH₃CN : water (95:5)

TFA Buffer pH = 2.5

Flow rate: 1 mL/min

Wavelength: 254 nm, 220 nm

HPLC-2: XBridge Phenyl (4.6 x 150 mm) 3.5 μ , gradient 10 to 100% B:A for 12 min, then 3 min hold at 100% B

Mobile Phase A: 0.05% TFA in water : CH₃CN (95:5)

Mobile Phase B: 0.05% TFA in CH₃CN : water (95:5)

TFA Buffer pH = 2.5

Flow rate: 1 mL/min

Wavelength: 254 nm, 220 nm

HPLC-3: Waters BEH, 2.0 x 50 mm, 1.7 μ m C18, gradient 0 to 100% B:A for 4 min, then 0.5 min hold at 100% B

Mobile Phase A: water : MeOH (95:5) + 10 μ M NH₄OAc

5 Mobile Phase B: MeOH : water (95:5) + 10 μ M NH₄OAc

Temperature = 40 °C

Flow rate: 1 mL/min.

10 HPLC-4: Waters BEH, 2.0 x 50 mm, 1.7 μ m C18, gradient 0 to 100% B:A for 4 min, then 0.5 min hold at 100% B

Mobile Phase A: water : CH₃CN (95:5) + 10 μ M NH₄OAc

Mobile Phase B: CH₃CN : water (95:5) + 10 μ M NH₄OAc

Temperature = 40 °C

Flow rate: 1 mL/min

15

HPLC- 5: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles

Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate

Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate

20 Temperature: 50 °C

Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B

Flow: 1.11 mL/min

25 HPLC-6: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles

Mobile Phase A: 5:95 acetonitrile:water with 0.1% TFA

Mobile Phase B: 95:5 acetonitrile:water with 0.1% TFA

Temperature: 50 °C

Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B

30 Flow: 1.11 mL/min.


NMR Employed in Characterization of Examples

[00166] NMR (nuclear magnetic resonance) spectra were typically obtained on Bruker or JEOL[®] 400 MHz and 500 MHz instruments in the indicated solvents. All chemical shifts are reported in ppm from tetramethylsilane with the solvent resonance as the internal standard. ¹H NMR spectral data are typically reported as follows: chemical shift, multiplicity (s = singlet, br s = broad singlet, d = doublet, dd = doublet of doublets, t = triplet, q = quartet, sep = septet, m = multiplet, app = apparent), coupling constants (Hz), and integration.


[00167] Spectral data are reported as chemical shift (multiplicity, number of hydrogens, coupling constants in Hz) and are reported in ppm (δ units) relative to either an internal standard (tetramethyl silane = 0 ppm) for ¹H NMR spectra, or are referenced to the residual solvent peak (2.49 ppm for CD₃SOCD₂H, 3.30 ppm for CD₂HOD, 1.94 for CHD₂CN, 7.26 ppm for CHCl₃, 5.32 ppm for CDHCl₂).

15 Example 1

4-(4-((1-(4-Chlorophenyl)-3-methyl-1*H*-pyrazol-5-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid




20 1A. 4-(Hydroxymethyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate



[00168] Borane .THF complex (3.38 mL, 3.38 mmol) was added dropwise to a -15 °C solution of 4-((3,5-difluorobenzoyl)oxy)bicyclo[2.2.1]heptane-1-carboxylic acid (1 g, 3.38 mmol) in THF (5 mL) and the resulting solution was stirred at -15 to 0 °C for 1.5 h.


The reaction mixture was then neutralized with acetic acid (1.5 mL) and the organic solvent was partially removed *in vacuo*. The residue was diluted with 5% aq. NaHCO₃ and extracted with EtOAc (3 x 5 mL). The combined organic phases were washed with brine, dried over MgSO₄, filtered, and concentrated *in vacuo* to afford the title compound (954 mg, 100% yield). LCMS, [M+H]⁺ = 283.0. ¹H NMR (400 MHz, CDCl₃) δ 7.56 - 7.48 (m, 2H), 7.00 (tt, *J* = 8.6, 2.4 Hz, 1H), 3.69 (s, 2H), 2.27 - 2.19 (m, 2H), 2.05 - 1.95 (m, 2H), 1.91 - 1.77 (m, 4H), 1.54 - 1.48 (m, 2H).

1B. 4-Formylbicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate



[00169] To a solution of 4-(hydroxymethyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate (954 mg, 3.38 mmol) in CH₂Cl₂ (30 mL) was added Dess-Martin periodinane (1.72 g, 4.06 mmol). The reaction mixture was stirred at rt until the oxidation was complete (about 1 h). The reaction mixture was filtered through a plug of CELITE®, washed with saturated aq. NaHCO₃, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The crude was purified by flash chromatography (SiO₂; gradient from 0 to 100% EtOAc/hexanes) to afford the title compound (887 mg, 94% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.79 (s, 1H), 7.56 - 7.46 (m, 2H), 7.00 (tt, *J* = 8.5, 2.2 Hz, 1H), 2.30 - 2.16 (m, 4H), 2.14 - 2.02 (m, 4H), 1.71 - 1.60 (m, 2H).


1C. (*E*)-4-(3-Methoxy-3-oxoprop-1-enyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate



[00170] A solution of 4-formylbicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate (590 mg, 2.11 mmol) and methyl(triphenylphosphoranylidene)acetate (1.41 g, 4.21 mmol) in THF (10 mL) was heated to 100 °C for 60 min in a microwave vial. The reaction was


cooled to RT and concentrated *in vacuo*. The residue was purified by flash chromatography (SiO₂; gradient from 0 to 100% EtOAc/hexanes) to afford the title compound (673 mg, 95% yield) as a white solid. LCMS, [M+H]⁺ = 337.1. ¹H NMR (400 MHz, CDCl₃) δ 7.56 - 7.49 (m, 2H), 7.15 (d, *J* = 15.9 Hz, 1H), 7.01 (tt, *J* = 8.5, 2.5 Hz, 1H), 5.84 (d, *J* = 15.4 Hz, 1H), 3.76 (s, 3H), 2.30 - 2.20 (m, 2H), 2.10 - 1.99 (m, 4H), 1.97 - 1.87 (m, 2H), 1.73 - 1.63 (m, 2H).

1D. 4-(3-Methoxy-3-oxopropyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate



- 10 **[00171]** A solution of (*E*)-4-(3-methoxy-3-oxoprop-1-enyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate (0.336 g, 1.0 mmol) and Pd/C (0.106 g, 0.100 mmol) in MeOH (8 mL) and THF (1 mL) was stirred under 1 atm of H₂ for 1 h. The reaction was then diluted with CH₂Cl₂ (10 mL) and filtered through a plug of CELITE®. The filtrate was concentrated *in vacuo* to afford the title compound (648 mg, 91% yield) as a white solid.
- 15 LCMS, [M+H]⁺ = 338.9. ¹H NMR (400 MHz, CDCl₃) δ 7.55 - 7.47 (m, 2H), 6.99 (tt, *J* = 8.5, 2.2 Hz, 1H), 3.69 (s, 3H), 2.38 - 2.30 (m, 2H), 2.23 - 2.13 (m, 2H), 2.03 - 1.93 (m, 2H), 1.90 - 1.83 (m, 2H), 1.80 (s, 2H), 1.73 - 1.62 (m, 2H), 1.57 - 1.48 (m, 2H).


1E. 3-(4-(3,5-Difluorobenzoyloxy)bicyclo[2.2.1]heptan-1-yl)propanoic acid



- 20 **[00172]** A mixture of 4-(3-methoxy-3-oxopropyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate (648 mg, 1.92 mmol) and LiI (1.28 g, 9.58 mmol) in pyridine (10 mL) was heated at 120 °C under Ar for 96 h, then was cooled to RT and concentrated *in vacuo*. The residue was taken up in EtOAc (20 mL) and washed with 1 N aq. HCl (10 mL) and water (10 mL). The organic layer was separated. The aqueous layer was
- 25


extracted with EtOAc (3 x 20 mL). The combined organic extracts were dried over MgSO₄, filtered and concentrated *in vacuo*. The crude material was purified by flash chromatography (SiO₂; gradient from 0 to 100% EtOAc/hexanes) to afford the title compound (500 mg, 80% yield) as a white solid. LCMS, [M+H]⁺ = 325.1. ¹H NMR (500 MHz, CDCl₃) δ 7.55 - 7.48 (m, 2H), 7.00 (tt, *J* = 8.6, 2.3 Hz, 1H), 2.44 - 2.35 (m, 2H), 2.23 - 2.14 (m, 2H), 2.05 - 1.95 (m, 2H), 1.92 - 1.86 (m, 2H), 1.83 (s, 2H), 1.74 - 1.64 (m, 2H), 1.60 - 1.50 (m, 2H).

1F. 4-(4-Diazo-3-oxobutyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate




[00173] To a solution of 3-(4-(3,5-difluorobenzoyloxy)bicyclo[2.2.1]heptan-1-yl) propanoic acid (490 mg, 1.511 mmol) in CH₂Cl₂ (15 mL) at 0 °C was added oxalyl chloride (2.27 mL, 4.53 mmol), followed by 1 drop of DMF. After 10 min, the mixture was warmed up to rt and stirred at rt for 1 h. The reaction was concentrated *in vacuo* and further azeotroped with toluene. The acid chloride intermediate was taken up in THF (5 mL)/MeCN (5 mL) under N₂, cooled to 0 °C, and added trimethylsilyldiazomethane (3.78 mL, 7.56 mmol). The resulting mixture was allowed to warm up to rt slowly and stirred at rt overnight. The reaction mixture was concentrated *in vacuo* and purified by flash chromatography (SiO₂; gradient from 0 to 30% EtOAc/hexanes) to afford the title compound (346 mg, 66% yield) as a light brown oil. LCMS, [M+H]⁺ = 349.3. ¹H NMR (500 MHz, CDCl₃) δ 7.54 - 7.49 (m, 2H), 7.00 (tt, *J* = 8.5, 2.5 Hz, 1H), 2.34 (br. s., 2H), 2.18 (td, *J* = 10.2, 4.5 Hz, 2H), 2.05 (s, 1H), 2.03 - 1.96 (m, 2H), 1.88 - 1.83 (m, 2H), 1.81 (s, 2H), 1.71 - 1.64 (m, 2H), 1.54 - 1.49 (m, 2H).

1G. 4-(4-(3,5-Difluorobenzoyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid



[00174] To a solution of 4-(4-diazo-3-oxobutyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate (346 mg, 0.99 mmol) in THF (20 mL) and water (10 mL) was added
 5 silver nitrate (177 mg, 1.04 mmol). The yellow/green mixture was stirred at rt overnight, then was concentrated *in vacuo* to remove THF, and the resulting slurry was partitioned between H₂O and CH₂Cl₂. The aqueous layer was extracted with CH₂Cl₂ (5 x 20 mL). The combined organic extracts were washed with brine (10 mL), dried over MgSO₄, filtered and concentrated *in vacuo* to afford the title compound (336 mg, 100% yield) as a
 10 white solid. LCMS, [M+Na]⁺ = 361.3.

1H. Methyl 4-(4-hydroxybicyclo[2.2.1]heptan-1-yl)butanoate



[00175] LiOH·H₂O (0.167 g, 3.97 mmol) was added to a mixture of 4-(4-(3,5-difluorobenzoyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid (0.336 g, 0.993 mmol) in
 15 THF (6 mL) and water (3 mL) at rt. The reaction was stirred at rt overnight and then diluted with EtOAc (30 mL) and H₂O (20 mL). The aqueous layer was washed with EtOAc (2 x 10 mL). The organic layer was extracted with H₂O (3 x 20 mL). The combined aqueous extracts were adjusted with 1 N aq. HCl to pH ~3 and extracted with
 20 EtOAc (3 x 50 mL). The combined organic extracts were washed with brine (10 mL), dried over MgSO₄, filtered and concentrated *in vacuo*. The crude product was dissolved in a mixture of CH₂Cl₂ (2 mL) and MeOH (2 mL). Trimethylsilyl(diazomethane) (1.99 mL of a 2 M solution in hexanes, 3.97 mmol) was added dropwise at 0 °C under Ar and the reaction was allowed to warm to RT and stirred at RT for 5 h. The reaction mixture
 25 was concentrated *in vacuo* and the residue was purified by flash chromatography (SiO₂;

gradient from 0 to 50% EtOAc/hexanes) to afford the title compound (178 mg, 85% yield) as colorless oil. LCMS, $[M+H]^+ = 213.3$. 1H NMR (500 MHz, $CDCl_3$) δ 3.67 (s, 3H), 2.30 (t, $J = 7.5$ Hz, 2H), 1.75 - 1.53 (m, 10H), 1.50 - 1.40 (m, 5H).


5 Example 1

[00176] To a 0 °C suspension of methyl 4-(4-hydroxybicyclo[2.2.1]heptan-1-yl) butanoate (10 mg, 0.047 mmol), 2,6-di-*tert*-butylpyridine (0.016 mL, 0.071 mmol), and AgOTf (13 mg, 0.052 mmol) in CH_2Cl_2 (0.5 mL) was added 5-(bromomethyl)-1-(4-chlorophenyl)-3-methyl-1*H*-pyrazole (16.14 mg, 0.057 mmol); a yellow precipitate formed within a few minutes. The reaction was slowly warmed up to rt and stirred at RT overnight. The reaction was diluted with CH_2Cl_2 and filtered through a plug of CELITE®. The filtrate was concentrated *in vacuo*. The residue was taken up in THF (1 mL) and water (0.5 mL), and LiOH·H₂O (10 mg, 0.24 mmol) was added. The reaction was stirred at rt overnight and diluted with EtOAc (30 mL) and H₂O (20 mL). The aqueous layer was washed with EtOAc (2 x 10 mL). The organic layer was extracted with H₂O (3 x 20 mL). The combined aqueous extracts were adjusted with 1 N aq. HCl to pH ~3 and extracted with EtOAc (3 x 50 mL). The combined organic extracts were washed with brine (10 mL), dried over MgSO₄, filtered and concentrated *in vacuo*. The residue was purified by preparative HPLC (PHENOMENEX® Synergi ODS-A-5 μ 21.2 x 250 mm column; flow rate = 25 mL/min, 15 to 100% solvent B over 20 min, hold to 21 min, where solvent A = 90:10:0.1 H₂O:MeCN:TFA and solvent B = 90:10:0.1 MeCN:H₂O:TFA) to give the title compound (10 mg, 50% yield) as a white solid. LCMS, $[M+H]^+ = 403.3$. 1H NMR (500 MHz, $CDCl_3$) δ 7.55 (d, $J = 8.8$ Hz, 2H), 7.43 (d, $J = 8.8$ Hz, 2H), 6.27 (s, 1H), 4.37 (s, 2H), 2.36 (t, $J = 7.4$ Hz, 2H), 2.33 (s, 3H), 1.84 - 1.75 (m, 2H), 1.67 - 1.55 (m, 6H), 1.52 - 1.45 (m, 4H), 1.42 (s, 2H). HPLC-1: RT = 11.3 min, purity = 100%.




[00177] The following Examples (Table 2) were prepared in a manner analogous to Example 1.




30




Table 2










Example No.	Name	R ¹ -L ₄ -	LCMS, [M-H] ⁺	¹ H NMR (500 MHz, CDCl ₃) δ	HPLC-1: RT, purity; HPLC-2: RT, purity
2	4-(4-(2-fluoro-5-methoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid			6.99 (dd, <i>J</i> = 5.8, 3.3 Hz, 1H), 6.93 (t, <i>J</i> = 9.2 Hz, 1H), 6.74 (dt, <i>J</i> = 8.9, 3.6 Hz, 1H), 4.55 (s, 2H), 3.79 (s, 3H), 2.36 (t, <i>J</i> = 7.5 Hz, 2H), 1.94 - 1.82 (m, 2H), 1.75 - 1.56 (m, 6H), 1.54 - 1.44 (m, 6H)	NA
3	4-(4-((5-chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		391.2	7.22 - 7.19 (m, 1H), 6.99 (d, <i>J</i> = 1.1 Hz, 1H), 4.45 (s, 2H), 2.97 (s, 2H), 2.36 (t, <i>J</i> = 7.4 Hz, 2H), 1.91 - 1.83 (m, 2H), 1.71 - 1.57 (m, 6H), 1.52 - 1.44 (m, 12H)	12.5 min, 100% 10.7 min, 100%





4	4-(4-(3,5-dichlorobenzyloxy) bicyclo[2.2.1]heptan-1-yl) butanoic acid		not shown	7.24 (s, 1H), 7.23 - 7.22 (m, 2H), 4.45 (s, 2H), 2.35 (t, $J = 7.3$ Hz, 2H), 1.88 - 1.80 (m, 2H), 1.70 - 1.57 (m, 6H), 1.53 - 1.44 (m, 6H)	13.1 min, 100% 10.7 min, 95.0%
5	4-(4-(5-chloro-2-isopropoxybenzyloxy) bicyclo[2.2.1]heptan-1-yl) butanoic acid		379.1	7.41 (d, $J = 2.8$ Hz, 1H), 7.14 (dd, $J = 8.8$, 2.8 Hz, 1H), 6.76 (d, $J = 8.8$ Hz, 1H), 4.53 - 4.49 (m, 1H), 4.48 (s, 2H), 2.39 - 2.31 (m, 2H), 1.92 - 1.83 (m, 2H), 1.73 - 1.57 (m, 6H), 1.53 - 1.44 (m, 6H), 1.34 (d, $J = 6.1$ Hz, 6H)	12.8 min, 100% 10.7 min, 96.5%
6	4-(4-(3-chloro-5-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		$[M+Na]^+$ 437.2	7.38 - 7.33 (m, 2H), 7.15 (t, $J = 7.4$ Hz, 1H), 7.08 (s, 1H), 7.02 (dd, $J = 8.7$, 1.0 Hz, 2H), 6.88 (s, 1H), 6.85 (t, $J = 1.9$ Hz, 1H), 4.44 (s, 2H), 2.38 (br. s., 2H), 1.87 - 1.78 (m, 2H), 1.70 - 1.54 (m, 6H), 1.52 - 1.40 (m, 6H)	13.5 min, 100% 11.5 min, 92.4%





7	4-(4-(3-methyl-5-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		[M+Na] ⁺ 409.2	7.09 (s, 1H), 7.01 (s, 1H), 6.92 (s, 1H), 4.48 (s, 2H), 2.40 - 2.32 (m, 5H), 1.86 (dd, <i>J</i> = 3.3, 2.5 Hz, 2H), 1.62 (br. s., 6H), 1.54 - 1.44 (m, 6H)	12.6 min, 100% 10.4 min, 100%
8	4-(4-(1-(3-(trifluoromethyl)phenyl)ethoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		[M+Na] ⁺ 393.1	7.61 (s, 1H), 7.53 (d, <i>J</i> = 7.4 Hz, 1H), 7.51 - 7.48 (m, 1H), 7.46 - 7.41 (m, 1H), 4.70 (q, <i>J</i> = 6.4 Hz, 1H), 2.31 (t, <i>J</i> = 7.4 Hz, 2H), 1.77 - 1.67 (m, 2H), 1.67 - 1.45 (m, 6H), 1.44 - 1.36 (m, 7H), 1.27 (s, 2H)	12.3 min, 100% 10.3 min, 100%
9	4-(4-(4,6-dichloro-2,3-dihydro-1 <i>H</i> -inden-1-yloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		381.0	7.22 (d, <i>J</i> = 1.7 Hz, 1H), 7.18 (s, 1H), 5.08 (t, <i>J</i> = 6.9 Hz, 1H), 3.03 (ddd, <i>J</i> = 16.6, 9.2, 3.3 Hz, 1H), 2.78 - 2.68 (m, 1H), 2.74 (dt, <i>J</i> = 16.6, 8.1 Hz, 1H), 2.50 - 2.41 (m, 1H), 2.37 (t, <i>J</i> = 7.3 Hz, 2H), 2.05 - 1.95 (m, 1H), 1.95 - 1.73 (m, 3H), 1.72 - 1.59 (m, 5H), 1.57 - 1.43 (m, 6H)	14.3 min, 98.5% 11.6 min, 94.4%





10	4-(4-(2-chlorobenzyloxy) bicyclo[2.2.1]heptan-1-yl) butanoic acid		[M+H] ⁺ 323.3	7.51 (dd, <i>J</i> = 7.4, 1.4 Hz, 1H), 7.31 (dd, <i>J</i> = 8.0, 1.4 Hz, 1H), 7.26 (br. s., 1H), 7.23 - 7.15 (m, 1H), 4.59 (s, 2H), 2.34 (t, <i>J</i> = 7.4 Hz, 2H), 1.95 - 1.82 (m, 2H), 1.74 - 1.58 (m, 6H), 1.52 - 1.43 (m, 6H)	11.7 min, 100% 9.9 min, 93.5%
11	4-(4-(3,4-dichlorobenzyloxy) bicyclo[2.2.1]heptan-1-yl) butanoic acid		[M+Na] ⁺ 379.3	7.42 (d, <i>J</i> = 1.6 Hz, 1H), 7.36 (d, <i>J</i> = 8.2 Hz, 1H), 7.14 (dd, <i>J</i> = 8.2, 2.2 Hz, 1H), 4.43 (s, 2H), 2.33 (t, <i>J</i> = 7.4 Hz, 2H), 1.88 - 1.76 (m, 2H), 1.70 - 1.55 (m, 6H), 1.51 - 1.41 (m, 6H)	12.7 min, 100% 10.5 min, 92.0%
12	4-(4-(3-fluoro-4-methylbenzyloxy)bicyclo [2.2.1]heptan-1-yl)butanoic acid		[M+Na] ⁺ 343.4	7.09 (t, <i>J</i> = 7.7 Hz, 1H), 7.01 - 6.94 (m, 2H), 4.43 (s, 2H), 2.33 (t, <i>J</i> = 7.1 Hz, 2H), 2.22 (s, 3H), 1.89 - 1.76 (m, 2H), 1.70 - 1.52 (m, 6H), 1.50 - 1.40 (m, 6H)	11.3 min, 100% 9.5 min, 92.1%




13	4-(4-(4-fluoro-3-methylbenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		[M+Na] ⁺ 343.4	7.18 (dd, <i>J</i> = 7.4, 1.7 Hz, 1H), 7.13 (ddd, <i>J</i> = 7.8, 5.2, 2.1 Hz, 1H), 6.96 (dd, <i>J</i> = 9.4, 8.5 Hz, 1H), 4.45 (s, 2H), 2.40 - 2.36 (m, 2H), 2.28 (d, <i>J</i> = 1.7 Hz, 3H), 1.92 - 1.84 (m, 2H), 1.71 - 1.60 (m, 6H), 1.54 - 1.47 (m, 6H)	11.5 min, 100% 9.6 min, 96.8%
14	4-(4-(2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		[M+Na] ⁺ 403.3	7.55 (d, <i>J</i> = 7.4 Hz, 1H), 7.31 (t, <i>J</i> = 7.8 Hz, 2H), 7.23 (t, <i>J</i> = 7.7 Hz, 1H), 7.17 - 7.12 (m, 1H), 7.07 (t, <i>J</i> = 7.3 Hz, 1H), 6.95 (d, <i>J</i> = 8.0 Hz, 2H), 6.88 (d, <i>J</i> = 8.0 Hz, 1H), 4.58 (s, 2H), 2.34 (t, <i>J</i> = 7.3 Hz, 2H), 1.85 - 1.77 (m, 2H), 1.68 - 1.52 (m, 6H), 1.49 - 1.38 (m, 6H)	12.1 min, 100% 10.5 min, 100%
15	4-(4-(3-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		379.2	7.36 - 7.28 (m, 3H), 7.13 - 7.07 (m, 2H), 7.05 - 6.98 (m, 3H), 6.89 (dd, <i>J</i> = 8.0, 1.9 Hz, 1H), 4.49 (s, 2H), 2.36 (t, <i>J</i> = 7.4 Hz, 2H), 1.89 - 1.80 (m, 2H), 1.72 - 1.54 (m, 6H), 1.52 - 1.41 (m, 6H)	12.6 min, 100% 10.9 min, 100%




16	4-(4-(3-(2-fluorophenoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		[M+Na] ⁺ 421.3	7.31 - 7.23 (m, 1H), 7.21 - 7.15 (m, 1H), 7.14 - 7.03 (m, 4H), 7.01 (br. s., 1H), 6.85 (d, <i>J</i> = 6.6 Hz, 1H), 4.49 (s, 2H), 2.36 (t, <i>J</i> = 7.3 Hz, 2H), 1.89 - 1.81 (m, 2H), 1.70 - 1.56 (m, 6H), 1.51 - 1.43 (m, 6H)	11.9 min, 100% 10.5 min, 100%
17	4-(4-(3-(4-fluorophenoxy)benzyloxy)bicyclo[2.2.1]-1-yl)butanoic acid		not shown	7.30 - 7.25 (m, 1H), 7.08 (d, <i>J</i> = 7.4 Hz, 1H), 7.05 - 6.96 (m, 5H), 6.85 (dd, <i>J</i> = 8.1, 2.1 Hz, 1H), 4.49 (s, 2H), 2.36 (t, <i>J</i> = 7.4 Hz, 2H), 1.84 (d, <i>J</i> = 2.2 Hz, 2H), 1.70 - 1.57 (m, 6H), 1.52 - 1.44 (m, 6H)	12.1 min, 100% 10.6 min, 91.7%
18	4-(4-(3-chloro-5-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		[M+H] ⁺ 407.3	7.29 (s, 1H), 7.13 (s, 1H), 7.11 (s, 1H), 4.50 (s, 2H), 2.36 (t, <i>J</i> = 7.4 Hz, 2H), 1.89 - 1.80 (m, 2H), 1.72 - 1.58 (m, 6H), 1.54 - 1.44 (m, 6H)	12.9 min, 100% 10.8 min, 100%
19	4-(4-(5-chloro-2-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		[M+Na] ⁺ 429.2	7.57 (d, <i>J</i> = 2.8 Hz, 1H), 7.28 - 7.24 (m, 1H), 7.15 (dq, <i>J</i> = 8.8, 1.5 Hz, 1H), 4.54 (s, 2H), 2.36 (s, 2H), 1.90 - 1.83 (m, 2H), 1.63 (d, <i>J</i> = 3.9 Hz, 6H), 1.54 - 1.45 (m, 6H)	12.8 min, 100% 10.7 min, 100%




20	4-(4-(2-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		[M+Na] ⁺ 395.3	7.59 - 7.55 (m, 1H), 7.31 - 7.28 (m, 2H), 7.22 (ddt, <i>J</i> = 5.4, 3.6, 1.8 Hz, 1H), 4.58 (s, 2H), 2.36 (t, <i>J</i> = 7.4 Hz, 2H), 1.92 - 1.84 (m, 2H), 1.74 - 1.58 (m, 6H), 1.53 - 1.45 (m, 6H)	11.7 min, 100% 10.0 min, 93.9%
21	4-(4-(3,5-dimethoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		347.1	6.52 (d, <i>J</i> = 2.5 Hz, 2H), 6.37 (t, <i>J</i> = 2.2 Hz, 1H), 4.46 (s, 2H), 3.79 (s, 6H), 2.36 (t, <i>J</i> = 7.4 Hz, 2H), 1.91 - 1.82 (m, 2H), 1.72 - 1.56 (m, 6H), 1.52 - 1.44 (m, 6H)	10.3 min, 100% 8.9 min, 100%
22	4-(4-(3-chloro-5-fluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		339.1	7.13 (s, 1H), 6.98 (s, 1H), 6.97 (s, 1H), 4.48 (s, 2H), 2.36 (t, <i>J</i> = 7.4 Hz, 2H), 1.84 (d, <i>J</i> = 1.7 Hz, 2H), 1.71 - 1.58 (m, 6H), 1.53 - 1.44 (m, 6H)	12.2 min, 100% 10.0 min, 95.0%
23	4-(4-(3-chlorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		321.1	7.36 (s, 1H), 7.28 - 7.25 (m, 1H), 7.25 - 7.23 (m, 1H), 7.23 - 7.19 (m, 1H), 4.49 (s, 2H), 2.36 (t, <i>J</i> = 7.4 Hz, 2H), 1.90 - 1.82 (m, 2H), 1.73 - 1.57 (m, 6H), 1.53 - 1.45 (m, 6H)	11.7 min, 100% 9.7 min, 100%




24	4-(4-(4-chlorobenzyloxy) bicyclo[2.2.1]heptan-1-yl) butanoic acid		321.1	7.32 - 7.28 (m, 4H), 4.48 (s, 2H), 2.36 (t, J = 7.4 Hz, 2H), 1.89 - 1.82 (m, 2H), 1.72 - 1.56 (m, 6H), 1.52 - 1.44 (m, 6H)	11.7 min, 97.3% 9.0 min, 95.0%
25	4-(4-(3-fluoro-5-(trifluoromethyl)benzyloxy) bicyclo[2.2.1]heptan-1-yl) butanoic acid		373.1	7.39 (s, 1H), 7.30 - 7.25 (m, 1H), 7.23 - 7.19 (m, 1H), 4.56 (s, 2H), 2.36 (t, J = 7.4 Hz, 2H), 1.90 - 1.82 (m, 2H), 1.73 - 1.59 (m, 6H), 1.55 - 1.45 (m, 6H)	12.4 min, 100% 10.5 min, 100%
26	4-(4-(3-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1] heptan-1-yl)butanoic acid		371.1	7.37 - 7.32 (m, 1H), 7.28 - 7.25 (m, 1H), 7.22 (s, 1H), 7.11 (d, J = 8.0 Hz, 1H), 4.53 (s, 2H), 2.36 (t, J = 7.4 Hz, 2H), 1.90 - 1.82 (m, 2H), 1.73 - 1.55 (m, 6H), 1.53 - 1.45 (m, 6H)	12.2 min, 100% 10.3 min, 100%
27	4-(4-(3-fluoro-5-methoxybenzyloxy)bicyclo [2.2.1]heptan-1-yl)butanoic acid		335.2	6.70 - 6.64 (m, 2H), 6.51 (dt, J = 10.7, 2.3 Hz, 1H), 4.47 (s, 2H), 3.80 (s, 3H), 2.36 (s, 2H), 1.90 - 1.82 (m, 2H), 1.72 - 1.57 (m, 6H), 1.53 - 1.43 (m, 6H)	11.0 min, 100% 9.5 min, 100%





28	4-(4-(3-fluoro-5-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		389.3	7.05 - 7.02 (m, 1H), 7.01 (s, 1H), 6.85 (d, $J = 8.8$ Hz, 1H), 4.52 (s, 2H), 2.36 (t, $J = 7.4$ Hz, 2H), 1.90 - 1.81 (m, 2H), 1.71 - 1.57 (m, 6H), 1.54 - 1.44 (m, 6H)	12.1 min, 100% 10.0 min, 95.5%
29	4-(4-(2-methoxy-5-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		401.4	7.36 - 7.32 (m, 1H), 7.10 - 7.05 (m, 1H), 6.80 (d, $J = 8.8$ Hz, 1H), 4.53 (s, 2H), 3.83 (s, 3H), 2.36 (t, $J = 7.4$ Hz, 2H), 1.92 - 1.83 (m, 2H), 1.73 - 1.57 (m, 6H), 1.53 - 1.44 (m, 6H)	12.2 min, 100% 10.3 min, 100%
30	4-(4-((5-chlorobenzo[<i>b</i>]thiophen-3-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		377.3	7.80 (d, $J = 1.9$ Hz, 1H), 7.75 (d, $J = 8.5$ Hz, 1H), 7.44 (s, 1H), 7.32 (dd, $J = 8.5$, 1.9 Hz, 1H), 4.72 (s, 2H), 2.39 (t, $J = 7.4$ Hz, 2H), 1.98 - 1.90 (m, 2H), 1.76 (td, $J = 10.0$, 3.7 Hz, 2H), 1.70 - 1.61 (m, 4H), 1.57 - 1.48 (m, 6H)	12.5 min, 100% 10.7 min, 84.9%
31	4-(4-(3-chloro-5-methoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		351.3	6.94 (s, 1H), 6.81 - 6.76 (m, 2H), 4.45 (s, 2H), 3.80 (s, 3H), 2.36 (t, $J = 7.4$ Hz, 2H), 1.89 - 1.81 (m, 2H), 1.71 - 1.56 (m, 6H), 1.52 - 1.44 (m, 6H)	11.7 min, 100% 9.9 min, 100%





32	4-(4-(3-(4-chlorophenoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		413.0	7.33 - 7.27 (m, 3H), 7.11 (d, $J = 7.7$ Hz, 1H), 7.01 (d, $J = 1.7$ Hz, 1H), 6.96 - 6.91 (m, 2H), 6.88 (dd, $J = 8.1, 1.8$ Hz, 1H), 4.49 (s, 2H), 2.36 (t, $J = 7.4$ Hz, 2H), 1.89 - 1.80 (m, 2H), 1.71 - 1.56 (m, 6H), 1.52 - 1.44 (m, 6H)	13.2 min, 100% 11.1 min, 100%
33	4-(4-(2-fluoro-5-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		389.0	7.36 (dd, $J = 5.2, 3.0$ Hz, 1H), 7.12 - 7.08 (m, 1H), 7.06 - 7.00 (m, 1H), 4.57 (s, 2H), 2.37 (t, $J = 7.4$ Hz, 2H), 1.91 - 1.83 (m, 2H), 1.74 - 1.59 (m, 6H), 1.55 - 1.46 (m, 6H)	12.1 min, 100% 10.1 min, 100%
34	4-(4-(3-fluoro-5-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		397.0	7.40 - 7.34 (m, 2H), 7.18 - 7.12 (m, 1H), 7.05 - 7.01 (m, 2H), 6.82 (d, $J = 9.1$ Hz, 1H), 6.78 (s, 1H), 6.60 - 6.54 (m, 1H), 4.46 (s, 2H), 2.36 (t, $J = 7.4$ Hz, 2H), 1.88 - 1.79 (m, 2H), 1.71 - 1.56 (m, 6H), 1.53 - 1.42 (m, 6H)	12.7 min, 100% 10.8 in, 95.0%





35	4-(4-(3-fluoro-5-isopropoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		363.4	6.66 (s, 1H), 6.64 (d, $J = 9.1$ Hz, 1H), 6.48 (d, $J = 10.7$ Hz, 1H), 4.54 - 4.48 (m, 1H), 4.45 (s, 2H), 2.36 (t, $J = 7.3$ Hz, 2H), 1.90 - 1.82 (m, 2H), 1.72 - 1.56 (m, 6H), 1.54 - 1.44 (m, 6H), 1.33 (d, $J = 6.1$ Hz, 6H)	12.1 min, 100% 10.1 min, 95.0%
36	4-(4-(2-isopropoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		345.5	7.43 (d, $J = 6.9$ Hz, 1H), 7.20 (t, $J = 7.6$ Hz, 1H), 6.92 (t, $J = 7.4$ Hz, 1H), 6.85 (d, $J = 8.3$ Hz, 1H), 4.60 - 4.54 (m, 1H), 4.53 (s, 2H), 2.36 (t, $J = 7.4$ Hz, 2H), 1.90 (br. s., 2H), 1.75 - 1.56 (m, 6H), 1.54 - 1.44 (m, 6H), 1.34 (d, $J = 6.1$ Hz, 6H)	11.8 min, 100% 9.8 min, 95.0%
37	4-(4-(3-(5-methyl-1,2,4-oxadiazol-3-yl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		369.0	8.04 (s, 1H), 7.97 (d, $J = 7.7$ Hz, 1H), 7.51 (d, $J = 7.7$ Hz, 1H), 7.48 - 7.42 (m, 1H), 4.58 (s, 2H), 2.67 (s, 3H), 2.36 (t, $J = 7.4$ Hz, 2H), 1.94 - 1.83 (m, 2H), 1.76 - 1.57 (m, 6H), 1.55 - 1.42 (m, 6H)	10.5 min, 100% 8.8 min, 96.2%




38	4-(4-(3-isopropoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		345.1	7.22 (t, $J = 8.1$ Hz, 1H), 6.92 - 6.85 (m, 2H), 6.78 (d, $J = 6.9$ Hz, 1H), 4.59 - 4.52 (m, 1H), 4.48 (s, 2H), 2.35 (br. s., 2H), 1.91 - 1.82 (m, 2H), 1.73 - 1.54 (m, 6H), 1.46 (br. s., 6H), 1.33 (d, $J = 6.1$ Hz, 6H)	11.6 min, 100% 9.9 min, 100%
39	4-(4-(3-benzoylbenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		391.1	7.85 - 7.79 (m, 3H), 7.69 (d, $J = 7.7$ Hz, 1H), 7.64 - 7.58 (m, 2H), 7.49 (ddd, $J = 18.6, 11.6, 4.5$ Hz, 3H), 4.60 (s, 2H), 2.37 (t, $J = 7.4$ Hz, 2H), 1.97 - 1.82 (m, 2H), 1.78 - 1.57 (m, 6H), 1.57 - 1.43 (m, 6H)	11.1 min, 92.8% 9.5 min, 91.8%
40	4-(4-(4-chloro-2-methoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		[M+Na] ⁺ 375.2	7.37 (d, $J = 8.1$ Hz, 1H), 6.95 (dd, $J = 8.1, 1.9$ Hz, 1H), 6.84 (d, $J = 1.9$ Hz, 1H), 4.51 (s, 2H), 3.83 (s, 3H), 2.37 (t, $J = 7.4$ Hz, 2H), 1.93 - 1.84 (m, 2H), 1.75 - 1.57 (m, 6H), 1.55 - 1.46 (m, 6H)	NA





41	4-(4-(2,4-dichlorobenzyloxy) bicyclo[2.2.1]heptan-1-yl) butanoic acid		355.1	7.49 (d, $J = 8.3$ Hz, 1H), 7.36 (d, $J = 2.1$ Hz, 1H), 7.26 (dd, $J = 8.3, 2.1$ Hz, 1H), 4.57 (s, 2H), 2.37 (t, $J = 7.4$ Hz, 2H), 1.97 - 1.84 (m, 2H), 1.77 - 1.57 (m, 6H), 1.56 - 1.44 (m, 6H)	13.4 min, 96.2% 10.8 min, 94.0%
42	4-(4-(2-fluoro-4-methoxybenzyloxy)bicyclo [2.2.1]heptan-1-yl)butanoic acid		335.4	7.33 (t, $J = 8.5$ Hz, 1H), 6.69 (dd, $J = 8.5, 2.3$ Hz, 1H), 6.61 (dd, $J = 11.8, 2.5$ Hz, 1H), 4.51 (s, 2H), 3.80 (s, 3H), 2.37 (t, $J = 7.4$ Hz, 2H), 1.96 - 1.83 (m, 2H), 1.78 - 1.56 (m, 6H), 1.57 - 1.44 (m, 6H)	10.6 min, 93.6% 9.2 min, 94.8%
43	4-(4-(4-methoxy-2-(trifluoromethoxy)benzyloxy) bicyclo[2.2.1]heptan-1-yl) butanoic acid		401.4	7.44 (d, $J = 8.6$ Hz, 1H), 6.84 (dd, $J = 8.6, 2.5$ Hz, 1H), 6.82 - 6.77 (m, 1H), 4.51 (s, 2H), 3.82 (s, 3H), 2.37 (t, $J = 7.4$ Hz, 2H), 1.94 - 1.82 (m, 2H), 1.75 - 1.57 (m, 6H), 1.56 - 1.44 (m, 6H)	11.9 min, 93.6% 9.8 min, 94.6%





44	4-(4-(5-fluoro-2-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		389.3	7.44 - 7.39 (m, 1H), 7.35 (dd, $J = 9.2, 3.4$ Hz, 1H), 7.27 (td, $J = 8.5, 3.2$ Hz, 1H), 4.50 (s, 2H), 2.18 - 2.08 (m, 2H), 1.82 - 1.73 (m, 2H), 1.62 - 1.50 (m, 4H), 1.38 (s, 8H)*	4.1 min, 100% 3.1 min, 100%**
45	4-(4-(3-chloro-2-fluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		339.2	7.54 - 7.48 (m, 1H), 7.41 (t, $J = 6.4$ Hz, 1H), 7.21 (t, $J = 7.9$ Hz, 1H), 4.54 (s, 2H), 2.17 (t, $J = 7.0$ Hz, 2H), 1.83 - 1.73 (m, 2H), 1.63 - 1.50 (m, 4H), 1.39 (s, 8H)*	3.94 min, 100% 2.7 min, 98.8%**
46	4-(4-(2-fluoro-3-(trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		373.2	7.78 (t, $J = 7.2$ Hz, 1H), 7.71 (t, $J = 7.2$ Hz, 1H), 7.40 (t, $J = 7.6$ Hz, 1H), 4.58 (s, 2H), 2.17 (t, $J = 7.2$ Hz, 2H), 1.84 - 1.74 (m, 2H), 1.64 - 1.50 (m, 4H), 1.40 (s, 8H)*	4.0 min, 99.2% 2.8 min, 98.3%**
47	4-(4-(2,3,6-trifluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		341.2	7.54 - 7.46 (m, 1H), 7.17 - 7.11 (m, 1H), 4.51 (s, 2H), 2.17 (t, $J = 7.2$ Hz, 2H), 1.80 - 1.71 (m, 2H), 1.63 - 1.50 (m, 4H), 1.39 (s, 8H)*	3.7 min, 94.0% 2.5 min, 90.1%**


48	4-(4-(biphenyl-2-ylmethoxy) bicyclo[2.2.1]heptan-1-yl) butanoic acid		363.3	7.50 - 7.41 (m, 3H), 7.41 - 7.33 (m, 5H), 7.24 (dd, $J = 5.3, 3.5$ Hz, 1H), 4.31 (s, 2H), 2.15 (t, $J = 7.2$ Hz, 2H), 1.71 - 1.61 (m, 2H), 1.51 - 1.39 (m, 6H), 1.39 - 1.30 (m, 4H), 1.25 (s, 2H)*	4.1 min, 100% 3.0 min, 100%**
49	4-(4-(2-(trifluoromethylthio) benzyloxy)bicyclo[2.2.1] heptan-1-yl)butanoic acid		387.2	7.70 (d, $J = 7.6$ Hz, 1H), 7.64 - 7.55 (m, 2H), 7.45 (td, $J = 7.5, 1.8$ Hz, 1H), 4.67 (s, 2H), 2.16 (t, $J = 7.2$ Hz, 2H), 1.85 - 1.76 (m, 2H), 1.63 - 1.51 (m, 4H), 1.40 (s, 8H)*	4.2 min, 100% 2.9 min, 100%**
50	4-(4-(3-(trifluoromethylthio) benzyloxy)bicyclo[2.2.1] heptan-1-yl)butanoic acid		387.2	7.65 (s, 1H), 7.61 (d, $J = 7.6$ Hz, 1H), 7.52 (dt, $J = 16.0, 7.9$ Hz, 2H), 4.54 (s, 2H), 2.18 (t, $J = 7.2$ Hz, 2H), 1.83 - 1.74 (m, 2H), 1.62 - 1.50 (m, 4H), 1.50 - 1.33 (m, 8H)*	4.2 min, 97.9% 3.0 min, 100%**
51	4-(4-(2-chloro-5- nitrobenzyloxy)bicyclo[2.2.1] heptan-1-yl)butanoic acid		366.1	8.29 (d, $J = 2.7$ Hz, 1H), 8.16 (dd, $J = 8.7,$ 2.9 Hz, 1H), 7.75 (d, $J = 8.5$ Hz, 1H), 4.63 (s, 2H), 2.18 (t, $J = 7.0$ Hz, 2H), 1.88 - 1.80 (m, 2H), 1.68 - 1.52 (m, 4H), 1.51 - 1.36 (m, 8H)*	3.9 min, 98.5% 2.7 min, 98.7%**

52	4-(4-(5-chloro-2-fluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		339.2	¹ H NMR (400 MHz, CDCl ₃): 7.46 (dd, <i>J</i> = 6.3, 2.6 Hz, 1H), 7.42 - 7.37 (m, 1H), 7.23 (t, <i>J</i> = 9.2 Hz, 1H), 4.50 (s, 2H), 2.17 (t, <i>J</i> = 7.2 Hz, 2H), 1.83 - 1.73 (m, 2H), 1.57 (s, 4H), 1.39 (s, 8H)*	4.0 min, 95.8% 2.7 min, 91.9%**
53	4-(4-(2,3-difluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		323.2	¹ H NMR (400 MHz, CDCl ₃): 7.25 - 7.20 (m, 1H), 7.19 - 7.10 (m, 2H), 4.60 (s, 2H), 2.22 (t, <i>J</i> = 7.5 Hz, 2H), 1.91 - 1.81 (m, 2H), 1.72 - 1.54 (m, 6H), 1.47 (s, 6H)	4.8 min, 98.4% 2.4 min, 90.2%**
54	4-(4-((2-bromothiophen-3-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		373.1	¹ H NMR (400 MHz, CDCl ₃): 7.55 (d, <i>J</i> = 5.5 Hz, 1H), 7.01 (d, <i>J</i> = 5.8 Hz, 1H), 4.36 (s, 2H), 2.16 (t, <i>J</i> = 7.0 Hz, 2H), 1.81 - 1.72 (m, 2H), 1.62 - 1.49 (m, 4H), 1.37 (s, 8H)*	3.8 min, 96.7% 2.7 min, 95.4%**
55	4-(4-(2-chloro-3-(trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		389.2	¹ H NMR (400 MHz, CDCl ₃): 7.84 (d, <i>J</i> = 7.6 Hz, 1H), 7.81 (d, <i>J</i> = 7.6 Hz, 1H), 7.58 (t, <i>J</i> = 7.8 Hz, 1H), 4.64 (s, 2H), 2.19 (t, <i>J</i> = 7.2 Hz, 2H), 1.89 - 1.80 (m, 2H), 1.68 - 1.53 (m, 4H), 1.53 - 1.37 (m, 8H)*	3.8 min, 93.3% 3.1 min, 100%**

56	4-(4-(2-chloro-3,6-difluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		357.2	7.48 (dt, $J = 8.9, 4.5$ Hz, 1H), 7.31 (td, $J = 9.0, 4.3$ Hz, 1H), 4.55 (s, 2H), 2.20 - 2.14 (m, 2H), 1.79 (br. s., 2H), 1.64 - 1.51 (m, 4H), 1.50 - 1.35 (m, 8H)*	3.9 min, 100% 2.6 min, 97.9%**
57	4-(4-(biphenyl-3-ylmethoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		363.3	7.64 (d, $J = 7.6$ Hz, 2H), 7.57 (s, 1H), 7.54 (d, $J = 7.6$ Hz, 1H), 7.47 (t, $J = 7.8$ Hz, 2H), 7.42 (t, $J = 7.5$ Hz, 1H), 7.39 - 7.34 (m, 1H), 7.31 (d, $J = 7.6$ Hz, 1H), 4.54 (s, 2H), 2.17 (t, $J = 7.2$ Hz, 2H), 1.86 - 1.76 (m, 2H), 1.65 - 1.51 (m, 4H), 1.50 - 1.35 (m, 8H)*	4.2 min, 100% 3.2 min, 100%**
58	4-(4-(4-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		379.3	7.38 (t, $J = 7.9$ Hz, 2H), 7.32 (d, $J = 8.5$ Hz, 2H), 7.15 - 7.09 (m, 1H), 6.97 (t, $J = 8.7$ Hz, 4H), 4.44 (s, 2H), 2.17 (t, $J = 7.2$ Hz, 2H), 1.83 - 1.74 (m, 2H), 1.63 - 1.51 (m, 4H), 1.49 - 1.35 (m, 8H)*	4.2 min, 98.6% 3.2 min, 100%**

59	4-(4-(2,3,6-trichlorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		389.1	7.67 (d, $J = 8.5$ Hz, 1H), 7.52 (d, $J = 8.9$ Hz, 1H), 4.68 (s, 2H), 2.17 (t, $J = 7.0$ Hz, 2H), 1.86 - 1.77 (m, 2H), 1.62 (br. s., 4H), 1.51 - 1.35 (m, 8H)*	4.2 min, 100% 3.2 min, 100%**
60	4-(4-(5-bromo-2-methoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		397.2	7.43 - 7.37 (m, 2H), 6.94 (d, $J = 8.2$ Hz, 1H), 4.43 (s, 2H), 3.77 (s, 3H), 2.16 (t, $J = 7.2$ Hz, 2H), 1.81 - 1.72 (m, 2H), 1.63 - 1.50 (m, 4H), 1.50 - 1.38 (m, 8H)*	4.0 min, 100% 3.0 min, 96.9%**
61	4-(4-(3-(thiophen-2-yl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		369.2	7.58 - 7.53 (m, 3H), 7.50 (d, $J = 3.4$ Hz, 1H), 7.37 (t, $J = 7.6$ Hz, 1H), 7.25 (d, $J = 7.6$ Hz, 1H), 7.14 (dd, $J = 5.2, 3.7$ Hz, 1H), 4.51 (s, 2H), 2.15 (t, $J = 7.0$ Hz, 2H), 1.85 - 1.76 (m, 2H), 1.64 - 1.51 (m, 4H), 1.50 - 1.35 (m, 8H)*	4.2 min, 98.5% 3.1 min, 100%**
62	4-(4-(5-bromo-2-methylbenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		381.2	7.47 (d, $J = 1.8$ Hz, 1H), 7.35 (dd, $J = 7.9, 2.1$ Hz, 1H), 7.11 (d, $J = 8.2$ Hz, 1H), 4.43 (s, 2H), 2.20 (s, 3H), 2.18 (t, $J = 7.0$ Hz, 2H), 1.86 - 1.77 (m, 2H), 1.63 - 1.51 (m, 4H), 1.50 - 1.35 (m, 8H)*	4.2 min, 100% 3.1 min, 100%**

63	4-(4-(2-chloro-5-(trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		389.2	7.80 (s, 1H), 7.71 - 7.66 (m, 2H), 4.60 (s, 2H), 2.15 (t, $J = 7.2$ Hz, 2H), 1.86 - 1.77 (m, 2H), 1.65 - 1.51 (m, 4H), 1.50 - 1.34 (m, 8H)*	4.2 min, 95.0% 3.2 min, 95.4%**
64	4-(4-(2-(trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		355.2	7.72 - 7.64 (m, 3H), 7.52 - 7.47 (m, 1H), 4.61 (s, 2H), 2.17 (t, $J = 7.2$ Hz, 2H), 1.84 - 1.75 (m, 2H), 1.64 - 1.50 (m, 4H), 1.50 - 1.35 (m, 8H)*	4.0 min, 100% 2.7 min, 100%**
65	4-(4-(3-(trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid		355.2	7.65 - 7.60 (m, 3H), 7.59 - 7.54 (m, 1H), 4.57 (s, 2H), 2.16 (t, $J = 7.0$ Hz, 2H), 1.84 - 1.75 (m, 2H), 1.64 - 1.50 (m, 4H), 1.50 - 1.33 (m, 8H)*	4.0 min, 94.2% 2.7 min, 100%**
66	methyl 4-(4-((4-fluoro-4'-methyl-[1,1'-biphenyl]-2-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)butanoate		395.1	7.29 - 7.22 (m, 6H), 7.20 - 7.14 (m, 1H), 4.32 (s, 2H), 2.35 (s, 3H), 2.14 (t, $J=7.2$ Hz, 2H), 1.72 - 1.63 (m, 2H), 1.52 - 1.30 (m, 10H), 1.26 (s, 2H)*	2.20 min, 99% 2.52 min, 99%***

67	4-(4-((2-(methylthio) pyrimidin-4-yl)methoxy) bicyclo[2.2.1]heptan-1-yl) butanoic acid		335.2	8.61 (d, $J = 5.1$ Hz, 1H), 7.22 (d, $J = 5.1$ Hz, 1H), 4.52 (s, 2H), 2.50 (s, 3H), 2.15 (t, $J = 7.1$ Hz, 2H), 1.84 - 1.76 (m, 2H), 1.64 - 1.51 (m, 4H), 1.50-1.35 (m, 8H)*	3.51 min, 95% 1.95 min, 90%**
----	---	--	-------	--	----------------------------------


* ^1H NMR (500 MHz, DMSO- d_6) δ

** HPLC-3: RT, purity; HPLC-4: RT, purity

*** HPLC-5: purity; HPLC-4: RT, purity

Example 68

2-Methyl-4-(4-(2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid




5 68A. 4-Methoxybicyclo[2.2.1]heptane-1-carbaldehyde



[00178] 4-Methoxybicyclo[2.2.1]heptane-1-carbaldehyde was prepared using a procedure analogous to 4-formylbicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate except that 4-((3,5-difluorobenzoyl)oxy)bicyclo[2.2.1]heptane-1-carboxylic acid was replaced
 10 with 4-methoxybicyclo[2.2.1]heptane-1-carboxylic acid. ¹H NMR (500 MHz, CDCl₃) δ 9.75 (s, 1H), 3.34 (s, 3H), 2.15 - 2.07 (m, 2H), 1.93 - 1.85 (m, 2H), 1.73 - 1.66 (m, 4H), 1.63 - 1.56 (m, 2H).


68B. 2-(4-Methoxybicyclo[2.2.1]heptan-1-yl)acetaldehyde



15 [00179] To a -78 °C stirred suspension of (methoxymethyl)triphenylphosphonium chloride (2.74 g, 8.01 mmol) in THF (10 mL) was added a solution of potassium bis(trimethylsilyl)amide (21.4 mL of a 0.5 M solution in toluene, 10.7 mmol). The resulting yellow mixture was stirred at -78 °C for 1.5 h, and a solution of 4-methoxy-
 20 bicyclo[2.2.1]heptane-1-carbaldehyde (0.823 g, 5.34 mmol) in THF (2 mL) was added over a period of 2 min. The mixture was held at -78 °C for 2 h and allowed to warm to RT and stirred at RT for 20 min. 1 N aq. HCl (5 mL) was added and the reaction was stirred at rt for 48 h. The reaction mixture was extracted with EtOAc (3 x 10 mL). The combined organic extracts were washed with brine (10 mL) and concentrated *in vacuo*.
 25 The resulting orange oil was purified by flash chromatography (SiO₂; gradient from 0 to


50% EtOAc/hexanes) to afford the title compound (400 mg, 45% yield) as a colorless oil. LCMS, $[M+H]^+ = 169.1$.

68C. (*E*)-Ethyl 4-(4-methoxybicyclo[2.2.1]heptan-1-yl)-2-methylbut-2-enoate



[00180] A solution of 2-(4-methoxybicyclo[2.2.1]heptan-1-yl)acetaldehyde (400 mg, 2.38 mmol) and ethyl 2-(triphenylphosphoranylidene)propanoate (862 mg, 2.39 mmol) in THF (8 mL) was heated to 100 °C for 1 h, and then 130 °C for 1 h in a microwave reactor. The reaction was cooled to rt and concentrated *in vacuo*. The residue was purified by
10 flash chromatography (SiO₂; gradient from 0 to 50% EtOAc/hexanes) to afford the title compound (242 mg, 40% yield). LCMS, $[M+H]^+ = 253.2$.


68D. Ethyl 4-(4-methoxybicyclo[2.2.1]heptan-1-yl)-2-methylbutanoate



15 [00181] Ethyl 4-(4-methoxybicyclo[2.2.1]heptan-1-yl)-2-methylbutanoate was prepared using a procedure analogous to the synthesis of 4-(3-methoxy-3-oxopropyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate except that (*E*)-4-(3-methoxy-3-oxoprop-1-enyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate was replaced with (*E*)-ethyl 4-(4-methoxy bicyclo[2.2.1]heptan-1-yl)-2-methylbut-2-enoate. LCMS, $[M+H]^+ = 255.2$.

20

68E. Ethyl 4-(4-hydroxybicyclo[2.2.1]heptan-1-yl)-2-methylbutanoate



[00182] To a solution of impure ethyl 4-(4-methoxybicyclo[2.2.1]heptan-1-yl)-2-methylbutanoate (200 mg, 0.250 mmol) in MeCN (1 mL) at 0 °C was added TMSI (0.194
25 mL, 1.43 mmol). The mixture was warmed to rt and stirred overnight. The reaction was


cooled -40 °C, quenched with sat'd aq. NaHCO₃ (5 mL) and then concentrated *in vacuo*. The residue was diluted with EtOAc (5 mL) and 10% aq. Na₂S₂O₃ (3 mL), and then acidified with 1 N aq. HCl to pH 2-3. The organic layer was washed with brine (2 mL), dried over MgSO₄, filtered and concentrated *in vacuo*. The residue was purified by flash chromatography (SiO₂; gradient from 0 to 50% EtOAc/hexanes) to afford the title compound (200 mg, 26% yield) as a colorless oil. LCMS, [M+H]⁺ = 241.2.

Example 68

[00183] The title compound was prepared using a procedure analogous to Example 1 except that methyl 4-(4-hydroxybicyclo[2.2.1]heptan-1-yl)butanoate was replaced with ethyl 4-(4-hydroxybicyclo[2.2.1]heptan-1-yl)-2-methylbutanoate. LCMS, [M-H]⁺ = 393.1. ¹H NMR (500 MHz, CDCl₃) δ 7.55 (dd, *J* = 7.4, 1.4 Hz, 1H), 7.34 - 7.29 (m, 2H), 7.25 - 7.20 (m, 1H), 7.17 - 7.12 (m, 1H), 7.10 - 7.05 (m, 1H), 6.97 - 6.93 (m, 2H), 6.88 (dd, *J* = 8.1, 1.0 Hz, 1H), 4.58 (s, 2H), 1.86 - 1.77 (m, 2H), 1.69 - 1.49 (m, 7H), 1.47 - 1.39 (m, 6H), 1.19 (d, *J* = 6.9 Hz, 3H). HPLC-1: RT = 13.2 min, purity = 92.8%; HPLC-2: RT = 11.1 min, purity = 100%.

Example 69

3-Methyl-4-(4-(3-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid




[00184] The title compound was synthesized similarly as for Example 68 through the following sequence: (1) reaction of 4-methoxybicyclo[2.2.1]heptane-1-carbaldehyde with the anion of (methoxymethyl)triphenylphosphonium chloride to afford the corresponding ester; (2) TMSI treatment to give the alcohol as in 68E; (3) AgOTf-mediated alkylation of the alcohol as in Example 1H; (4) LiOH hydrolysis of the ethyl ester as in Example 1H; (5) one-carbon elongation of the resulting acid using the Arndt-Eistert reaction sequence provided the title compound. LCMS, [M-H]⁺ = 393.1. ¹H NMR (500 MHz, CDCl₃) δ 7.57 (dd, *J* = 7.55, 1.51 Hz, 1H), 7.36 - 7.31 (m, 2H), 7.24 (td, *J* = 7.84, 1.72 Hz, 1H),


7.16 (td, $J = 7.48, 1.09$ Hz, 1H), 7.10 - 7.06 (m, 1H), 6.99 - 6.95 (m, 2H), 6.90 (dd, $J = 8.1, 1.0$ Hz, 1H), 4.59 (s, 2H), 2.36 (dd, $J = 15.2, 5.97$ Hz, 1H), 2.18 (dd, $J = 15.2, 8.0$ Hz, 1H), 2.1 - 2.0 (m, 1H), 1.87 - 1.78 (m, 2H), 1.69 - 1.56 (m, 4H), 1.54 - 1.43 (m, 5H), 1.36 (dd, $J = 14.2, 7.4$ Hz, 1H), 1.01 (d, $J = 6.6$ Hz, 3H). HPLC-1: RT = 13.2 min, purity = 95%; HPLC-2: RT = 11.1 min, purity = 95%.

Example 70

3-(4-((5-Chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)propan-1-ol




70A. 4-(3-Hydroxypropyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate



[00185] 4-(3-Hydroxypropyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate was prepared using a procedure analogous to 4-(hydroxymethyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate except that 4-((3,5-difluorobenzoyl)oxy)bicyclo[2.2.1]heptane-1-carboxylic acid was replaced with 3-(4-(3,5-difluorobenzoyloxy)bicyclo[2.2.1]heptan-1-yl)propanoic acid. LCMS, $[M+H]^+ = 310.9$. 1H NMR (500 MHz, $CDCl_3$) δ 7.55 - 7.49 (m, 2H), 6.99 (tt, $J = 8.5, 2.3$ Hz, 1H), 3.67 (t, $J = 6.2$ Hz, 2H), 2.22 - 2.14 (m, 2H), 2.02 - 1.94 (m, 2H), 1.81 (s, 2H), 1.72 - 1.66 (m, 2H), 1.62 - 1.51 (m, 6H).


70B. 4-(3-(Tetrahydro-2H-pyran-2-yloxy)propyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate



[00186] To a solution of 4-(3-hydroxypropyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate (145 mg, 0.467 mmol) and 3,4-dihydro-2H-pyran (0.085 mL, 0.93 mmol) in CH₂Cl₂ (2 mL) at 0 °C was added *p*-toluenesulfonic acid monohydrate (1 mg, 4.8 μmol) and the solution stirred at rt under N₂ for 16 h. The reaction was washed with water (1 mL), and sat. aq. NaHCO₃ (1 mL), dried over anhydrous MgSO₄, filtered, and concentrated *in vacuo*. The crude product was purified by flash chromatography (SiO₂; gradient from 0 to 30% EtOAc/hexanes) to afford the title compound (184 mg, 100% yield) as a clear oil. LCMS, [M+Na]⁺ = 416.9.

10


70C. 4-(3-(Tetrahydro-2H-pyran-2-yloxy)propyl)bicyclo[2.2.1]heptan-1-ol



[00187] To a solution of 4-(3-(tetrahydro-2H-pyran-2-yloxy)propyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate in THF (3 mL) was added LiOH·H₂O (66 mg, 1.61 mmol) in water (3 mL). The mixture was stirred at rt for 2 h. MeOH (3 mL) was added and the reaction was stirred for additional 1 h, then was concentrated *in vacuo*. The residue was extracted with CH₂Cl₂ (3 x 10 mL). The combined organic extracts were dried over MgSO₄, filtered and concentrated *in vacuo*. The residue was purified by flash chromatography (SiO₂; gradient from 0 to 100% EtOAc/hexanes) to afford the title compound (107 mg, 78% yield) as a clear oil. LCMS, [M+Na]⁺ = 277.0.

20

70D. 5-Chloro-2,2-dimethyl-7-((4-(3-(tetrahydro-2H-pyran-2-yloxy)propyl)bicyclo[2.2.1]heptan-1-yloxy)methyl)-2,3-dihydrobenzofuran




[00188] To a solution of 4-(3-(tetrahydro-2*H*-pyran-2-yloxy)propyl)bicyclo[2.2.1]heptan-1-ol (107 mg, 0.421 mmol) in DMF (1 mL) at 0 °C was added NaH (19 mg, 0.46 mmol). The mixture was stirred at 0 °C for 45 min. 5-Chloro-7-(chloromethyl)-2,2-dimethyl-2,3-dihydrobenzofuran (97 mg, 0.42 mmol) was added and the reaction was stirred at rt for 3 days. The reaction was diluted with water (2 mL) and CH₂Cl₂ (5 mL). The organic layer was separated, washed with 10% aq. LiCl (3 mL), dried over MgSO₄, filtered, and concentrated *in vacuo*. The residue was purified by flash chromatography (SiO₂; gradient from 0 to 45% EtOAc/hexanes) to afford the title compound (150 mg, 72% yield) as a white solid. LCMS, [M+H₂O]⁺ = 466.3. ¹H NMR (500 MHz, CDCl₃) δ 7.21 (s, 1H), 6.99 (s, 1H), 4.58 (dd, *J* = 4.3, 2.9 Hz, 1H), 4.45 (s, 2H), 3.88 (ddd, *J* = 11.1, 7.5, 3.2 Hz, 1H), 3.72 (dt, *J* = 9.5, 6.8 Hz, 1H), 3.54 - 3.48 (m, 1H), 3.39 (dt, *J* = 9.5, 6.7 Hz, 1H), 2.97 (s, 2H), 1.90 - 1.82 (m, 3H), 1.77 - 1.64 (m, 3H), 1.64 - 1.52 (m, 8H), 1.50 - 1.43 (m, 12H).

Example 70


[00189] To a solution of 5-chloro-2,2-dimethyl-7-((4-(3-(tetrahydro-2H-pyran-2-yloxy)propyl)bicyclo[2.2.1]heptan-1-yloxy)methyl)-2,3-dihydrobenzofuran (150 mg, 0.334 mmol) in MeOH (2 mL) was added *p*-toluenesulfonic acid monohydrate (6 mg, 0.033 mmol) and the reaction stirred at rt under N₂ for 1 h. The reaction was diluted with CH₂Cl₂ (10 mL) and washed with water (5 mL). The organic layer was dried (Na₂SO₄) and concentrated *in vacuo*. The crude product was purified by flash chromatography (SiO₂; gradient from 0 to 100% EtOAc/hexanes) to afford the title compound (125 mg, 97% yield) as a colorless oil. LCMS, [M+Na]⁺ = 387.2. ¹H NMR (500 MHz, CDCl₃) δ 7.21 (s, 1H), 6.98 (s, 1H), 4.45 (s, 2H), 3.63 (t, *J* = 6.5 Hz, 2H), 2.96 (s, 2H), 1.91 - 1.83 (m, 2H), 1.72 - 1.65 (m, 2H), 1.64 - 1.52 (m, 4H), 1.51 - 1.43 (m, 12H). HPLC-1: RT = 13.2 min, purity = 95.4%; HPLC-2: RT = 11.0 min, purity = 95.6%.

Example 71

3-(4-((5-Chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)propanoic acid



71A. 3-(4-((5-Chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)propanal



- 10 **[00190]** 3-(4-((5-Chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)propanal was prepared using a procedure analogous to the synthesis of 4-formyl-bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate except that 4-(hydroxymethyl)bicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate was replaced with 3-(4-((5-chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)propan-1-ol.
- 15 LCMS, $[M-H]^+ = 361.2$. 1H NMR (500 MHz, $CDCl_3$) δ 9.79 (s, 1H), 7.20 (s, 1H), 6.99 (s, 1H), 4.44 (s, 2H), 2.96 (s, 2H), 2.47 - 2.38 (m, 2H), 1.93 - 1.84 (m, 2H), 1.82 - 1.75 (m, 2H), 1.73 - 1.64 (m, 2H), 1.62 - 1.53 (m, 2H), 1.53 - 1.38 (m, 10H).


Example 71

- 20 **[00191]** To a 0 °C suspension of 3-(4-((5-chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)propanal (40 mg, 0.11 mmol) in 2-methylpropan-2-ol (1 mL, 0.11 mmol) and 2-methylbut-2-ene (1 mL, 0.11 mmol) at was added sodium chlorite (199 mg, 2.205 mmol) and $NaH_2PO_4 \cdot H_2O$ (258 mg, 1.65 mmol) in water (1 mL). The reaction was stirred at rt for 2 h, then was extracted with
- 25 EtOAc (3 x 3 mL), dried over $MgSO_4$, filtered and concentrated *in vacuo*. The residue


was purified by preparative HPLC (YMC ODS-A-5 μ 30 x 100 mm column; flow rate = 40 mL/min, 0 to 100% solvent B over 30 min, hold to 40 min, where solvent A = 90:10:0.1 H₂O:CH₃CN:TFA and solvent B = 90:10:0.1 CH₃CN:H₂O:TFA) to afford the title compound (8 mg, 19% yield) as a colorless oil. LCMS, [M-H]⁺ = 377.2. ¹H NMR (500 MHz, CDCl₃) δ 7.20 (s, 1H), 6.99 (s, 1H), 4.44 (s, 2H), 2.97 (s, 2H), 2.40 - 2.32 (m, 2H), 1.92 - 1.85 (m, 2H), 1.84 - 1.79 (m, 2H), 1.69 (td, *J* = 9.8, 2.9 Hz, 2H), 1.64 - 1.56 (m, 2H), 1.51 - 1.41 (m, 10H). HPLC-1: RT = 12.5 min, purity = 100%; HPLC-2: RT = 10.7 min, purity = 100%.

Example 72

2-((4-(2-Phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid




72A. Methyl 4-hydroxybicyclo[2.2.1]heptane-1-carboxylate




[00192] Methyl 4-hydroxybicyclo[2.2.1]heptane-1-carboxylate was prepared using a procedure analogous to the synthesis of methyl 4-(4-hydroxybicyclo[2.2.1]heptan-1-yl)butanoate except that 4-(4-(3,5-difluorobenzoyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid was replaced with 4-((3,5-difluorobenzoyl)oxy)bicyclo[2.2.1]heptane-1-carboxylic acid. LCMS, [M+H]⁺ = 171.1. ¹H NMR (500 MHz, CDCl₃) δ 3.69 (s, 3H), 2.16 - 2.08 (m, 2H), 1.82 - 1.68 (m, 9H).

72B. Methyl 4-(2-phenoxybenzyloxy)bicyclo[2.2.1]heptane-1-carboxylate



[00193] To a 0 °C suspension of methyl 4-hydroxybicyclo[2.2.1]heptane-1-carboxylate (100 mg, 0.588 mmol), 2,6-di-*tert*-butylpyridine (0.198 mL, 0.88 mmol), and AgOTf (166 mg, 0.65 mmol) in CH₂Cl₂ (5 mL) was added 1-(chloromethyl)-2-phenoxybenzene (128 mg, 0.59 mmol); a yellow precipitate formed within a few minutes. The reaction
 5 was slowly warmed up to RT and stirred overnight at RT, then was diluted with CH₂Cl₂ and filtered through a plug of CELITE®. The filtrate was concentrated *in vacuo* and purified by flash chromatography (SiO₂; gradient from 0 to 20% EtOAc/hexanes) to afford the title compound (126 mg, 61% yield) as a colorless oil. LCMS, [M+Na]⁺ = 375.2. ¹H NMR (500 MHz, CDCl₃) δ 7.54 (d, *J* = 7.2 Hz, 1H), 7.32 (t, *J* = 7.8 Hz, 2H),
 10 7.24 (d, *J* = 7.7 Hz, 1H), 7.15 (t, *J* = 7.4 Hz, 1H), 7.08 (t, *J* = 6.9 Hz, 1H), 6.95 (d, *J* = 8.5 Hz, 2H), 6.89 (d, *J* = 8.0 Hz, 1H), 4.60 (s, 2H), 3.67 (s, 3H), 2.12 - 2.02 (m, 2H), 1.93 - 1.85 (m, 2H), 1.82 (br. s., 2H), 1.77 - 1.70 (m, 2H), 1.67 (d, *J* = 9.6 Hz, 2H).


72C. (4-(2-Phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)methanol



15

[00194] To a solution of methyl 4-((2-phenoxybenzyl)oxy)bicyclo[2.2.1]heptane-1-carboxylate (125 mg, 0.36 mmol) in THF (2 mL) at -78 °C was added dropwise LiAlH₄ (1 M in Et₂O) (0.36 mL, 0.36 mmol) over a period of 3-4 min. The solution was allowed to warm to rt and kept at rt for 30 min. The reaction was cooled to 0 °C and quenched
 20 with EtOAc (20 mL) followed by 1 N aq. HCl (10 mL). The aqueous layer was extracted with EtOAc (3 x 20 mL). The combined organic extracts were concentrated *in vacuo* and the residue was purified by flash chromatography (SiO₂; gradient from 0 to 50% EtOAc/hexanes) to afford the title compound (101 mg, 88% yield) as a clear oil. LCMS, [M+Na]⁺ = 347.2. ¹H NMR (500 MHz, CDCl₃) δ 7.56 (d, *J* = 7.7 Hz, 1H), 7.35 - 7.29
 25 (m, 2H), 7.23 (t, *J* = 7.7 Hz, 1H), 7.18 - 7.12 (m, 1H), 7.10 - 7.04 (m, 1H), 6.98 - 6.92 (m, 2H), 6.88 (d, *J* = 8.3 Hz, 1H), 4.60 (s, 2H), 3.60 (d, *J* = 5.5 Hz, 2H), 1.90 - 1.82 (m, 2H), 1.73 - 1.64 (m, 4H), 1.47 (s, 2H), 1.46 - 1.39 (m, 2H), 1.24 (t, *J* = 5.9 Hz, 1H).

72D. *tert*-Butyl 2-((4-(2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetate



- [00195] To a 0 °C solution of (4-(2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl) methanol (100 mg, 0.31 mmol) in toluene (6 mL) was added 35% aq. NaOH (2 g in 5 mL of H₂O) was added, followed by tetrabutylammonium hydrogen sulfate (44 mg, 0.13 mmol). The mixture was stirred at 0 °C for 30 min. *tert*-Butyl 2-bromoacetate (0.091 mL, 0.62 mmol) was added and the mixture was stirred for 14 h at rt. The reaction was neutralized with conc. aq. HCl and extracted with EtOAc. The combined organic extracts were washed with brine (5 mL), dried over MgSO₄, filtered and concentrated *in vacuo*.
- 10 The residue was purified by flash chromatography (SiO₂; gradient from 0 to 100% EtOAc/hexanes) to afford the title compound (100 mg, 74% yield) as colorless oil. LCMS, [M+Na]⁺ = 641.3.

Example 72


- 15 [00196] LiOH·H₂O (48 mg, 1.14 mmol) was added to a solution of *tert*-butyl 2-((4-(2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetate (100 mg, 0.23 mmol) in THF (2 mL), water (1 mL), and MeOH (2 mL) at rt. The reaction was stirred at rt overnight and diluted with EtOAc (30 mL) and H₂O (20 mL). The aqueous layer was washed with EtOAc (2 x 10 mL). The organic layer was extracted with H₂O (3 x 20 mL). The combined aqueous extracts were adjusted with 1 N aq. HCl to pH ~3 and extracted with EtOAc (3 x 50 mL). The combined organic extracts were washed with brine (10 mL), dried over MgSO₄, filtered and concentrated *in vacuo*. The crude product was purified by preparative HPLC (PHENOMENEX® Synergi ODS-A-5μ 21.2 x 250 mm column; flow rate = 25 mL/min, 30 to 100% solvent B over 20 min, hold to 22 min, where solvent A = 90:10:0.1 H₂O:ACN:TFA and solvent B = 90:10:0.1 ACN:H₂O:TFA) to give the title compound (51 mg, 59% yield) as a white solid. LCMS, [M-H]⁺ = 381.0.
- 25 ¹H NMR (500 MHz, CDCl₃) δ 7.55 (d, *J* = 7.4 Hz, 1H), 7.35 - 7.29 (m, 2H), 7.26 - 7.21

(m, 1H), 7.17 - 7.12 (m, 1H), 7.10 - 7.05 (m, 1H), 6.98 - 6.92 (m, 2H), 6.88 (d, $J = 8.3$ Hz, 1H), 4.60 (s, 2H), 4.10 (s, 2H), 3.55 (s, 2H), 1.91 - 1.81 (m, 2H), 1.75 - 1.63 (m, 4H), 1.53 - 1.43 (m, 4H). HPLC-1: RT = 11.3 min, purity = 100%; HPLC-2: RT = 9.9 min, purity = 100%.




5


[00197] The following Examples (Table 3) were prepared in a manner analogous to Example 72.

Table 3




Example	Name	R ¹ -L ₄ -	LCMS, [M-H] ⁺	¹ H NMR (500 MHz, CDCl ₃) δ	HPLC-1: RT min, purity; HPLC-2: RT min, purity
73	2-((4-(3-fluoro-5-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid		399.0	7.37 (dd, <i>J</i> = 8.5, 7.4 Hz, 2H), 7.19 - 7.12 (m, 1H), 7.04 (dd, <i>J</i> = 8.5, 1.1 Hz, 2H), 6.82 (d, <i>J</i> = 9.1 Hz, 1H), 6.79 - 6.75 (m, 1H), 6.60 - 6.55 (m, 1H), 4.48 (s, 2H), 4.12 (s, 2H), 3.57 (s, 2H), 1.93 - 1.82 (m, 2H), 1.80 - 1.66 (m, 4H), 1.58 - 1.46 (m, 4H).	11.9 min, 100% 9.9 min, 93.2%
74	2-((4-(3-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid		381.0	7.34 (s, 3H), 7.13 - 7.07 (m, 2H), 7.05 - 6.98 (m, 3H), 6.92 - 6.88 (m, 1H), 4.51 (s, 2H), 4.12 (s, 2H), 3.57 (s, 2H), 1.93 - 1.85 (m, 2H), 1.79 - 1.68 (m, 4H), 1.58 - 1.45 (m, 4H).	11.4 min, 96.8% 9.6 min, 100%

75	2-((4-(3-fluoro-5-(4-fluorophenoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid		417.1	7.09 - 6.97 (m, 4H), 6.80 (dd, $J = 9.1, 0.6$ Hz, 1H), 6.73 (s, 1H), 6.56 - 6.50 (m, 1H), 4.47 (s, 2H), 4.13 (s, 2H), 3.56 (s, 2H), 1.92 - 1.82 (m, 2H), 1.80 - 1.66 (m, 4H), 1.57 - 1.47 (m, 4H).	11.8 min, 100% 9.9 min, 92.2%
76	2-((4-(1-(3-fluoro-5-(4-fluorophenoxy)phenyl)ethoxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid		431.1	7.06 (d, $J = 8.0$ Hz, 2H), 7.04 - 6.98 (m, 2H), 6.79 (dd, $J = 9.1, 1.9$ Hz, 1H), 6.74 (s, 1H), 6.53 - 6.47 (m, 1H), 4.59 (q, $J = 6.6$ Hz, 1H), 4.09 (s, 2H), 3.51 (s, 2H), 1.80 - 1.60 (m, 5H), 1.54 - 1.40 (m, 3H), 1.40 - 1.32 (m, 5H)	12.2 min, 100% 10.1 min, 95.0%
77	2-((4-((5-chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid		393.1	^1H NMR (500MHz, CDCl_3) δ 7.20 (s, 1H), 7.00 (s, 1H), 4.46 (s, 2H), 4.13 (s, 2H), 3.57 (s, 2H), 2.97 (s, 2H), 1.96 - 1.86 (m, 2H), 1.79 - 1.67 (m, 4H), 1.58 - 1.48 (m, 4H), 1.46 (s, 6H).	11.8 min, 100% 9.7 min, 100%


78	2-((4-((3-chloro-5-(trifluoromethoxy)benzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid		407.0	¹ H NMR (500MHz, CDCl ₃) 7.29 (s, 1H), 7.12 (d, <i>J</i> =13.5 Hz, 2H), 4.52 (s, 2H), 4.13 (s, 2H), 3.58 (s, 2H), 1.93 - 1.84 (m, H), 1.82 - 1.68 (m, 4H), 1.59 - 1.49 (m, 4H).	11.2 min, 100% 9.7 min, 100%
----	---	--	-------	--	---------------------------------

Example 79

3-(4-(3-Fluoro-5-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yloxy)propanoic acid




5 79A. (3,5-Difluorophenyl)(4-methoxybicyclo[2.2.1]heptan-1-yl)methanone



- [00198] To a solution of 4-methoxybicyclo[2.2.1]heptane-1-carboxylic acid (2 g, 11.75 mmol) in CH₂Cl₂ (30 mL) was added oxalyl chloride (8.81 mL of a 2.0 M solution in CH₂Cl₂; 17.6 mmol) followed by 1 drop of DMF. The reaction mixture was stirred at rt
- 10 for 1 h, then was concentrated *in vacuo* and dried *in vacuo* for 1 h to give the crude acid chloride product. To a solution of this crude acid chloride (2.22 g, 11.8 mmol) in THF (25 mL) was added acetylacetone iron (III) salt (0.124 g, 0.35 mmol). The orange mixture was cooled to 0 °C and (3,5-difluorophenyl) magnesium bromide (30.6 mL of a 0.5 M solution in THF, 15.3 mmol) was added dropwise. The mixture was stirred for at 0
- 15 °C for 2 h. The mixture was added 1 N aq. HCl and extracted with EtOAc. The combined organic layers were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by flash chromatography (SiO₂; gradient from 0 to 20% EtOAc/hexanes) to afford the title compound (2.30 g, 74% yield) as a light yellow oil. LCMS, [M+Na]⁺ = 289.1. ¹H NMR (500 MHz, CDCl₃) δ 7.35 - 7.29 (m, 2H), 6.97 (tt, *J* = 8.5, 2.4 Hz, 1H),
- 20 3.36 (s, 3H), 2.17 - 2.07 (m, 2H), 2.05 - 1.91 (m, 6H), 1.80 - 1.72 (m, 2H).


79B. 4-Methoxybicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate



- [00199] To a solution of (3,5-difluorophenyl)(4-methoxybicyclo[2.2.1]heptan-1-yl) methanone (2.3 g, 8.64 mmol) in TFA (25 mL) was added 30% aq. H₂O₂ (3.53 mL, 34.5 mmol) and the solution was stirred for 4 h at 50 °C. The mixture was cooled to 0 °C and sodium metabisulfite (1.64 g, 8.64 mmol) in water (5 mL) was added. The solvent was evaporated *in vacuo* and the residue was partitioned between EtOAc (100 mL) and H₂O (10 mL). The aqueous layer was extracted with EtOAc (2 x 20 mL). The combined organic extracts were washed with H₂O (2 x 10 mL) and brine (10 mL), dried over MgSO₄, filtered and concentrated *in vacuo*. The residue was purified by flash chromatography (SiO₂; gradient from 0 to 20% EtOAc/hexanes) to afford the title compound (2.21 g, 91% yield) as a colorless oil. LCMS, [M+Na]⁺ = 304.4. ¹H NMR (500 MHz, CDCl₃) δ 7.54 - 7.48 (m, 2H), 7.00 (tt, *J* = 8.5, 2.5 Hz, 1H), 3.34 (s, 3H), 2.32 - 2.23 (m, 2H), 2.17 - 2.08 (m, 4H), 2.06 - 1.96 (m, 2H), 1.81 - 1.72 (m, 2H).

15

79C. 4-Hydroxybicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate




- [00200] To a 0 °C solution of 4-methoxybicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate (2.2 g, 7.79 mmol) in MeCN (20 mL) at was added TMSI (1.59 mL, 11.7 mmol). The mixture was warmed to rt and was stirred overnight. The mixture was cooled to -78 °C, and sat'd aq. NaHCO₃ (10 mL) was added. The mixture was allowed to warm to rt, and 10% aqueous Na₂S₂O₃ was added. The mixture was extracted with EtOAc (3 x 100 mL). The combined organic extracts were washed with brine (20 mL), dried over MgSO₄, filtered and concentrated *in vacuo*. The residue was purified by flash chromatography (SiO₂; gradient from 0 to 30% EtOAc/hexanes) to afford the title compound (1.59 g, 76% yield) as a white waxy solid. LCMS, [M+Na]⁺ = 291.1. ¹H

20

25

NMR (500 MHz, CDCl_3) δ 7.54 - 7.48 (m, 2H), 7.00 (tt, J = 8.5, 2.5 Hz, 1H), 2.30 - 2.22 (m, 2H), 2.20 - 2.10 (m, 4H), 1.98 - 1.90 (m, 2H), 1.89 - 1.81 (m, 2H), 1.68 (s, 1H).


79D. 3-(4-Hydroxybicyclo[2.2.1]heptan-1-yloxy)propanoic acid



[00201] 3-(4-Hydroxybicyclo[2.2.1]heptan-1-yloxy)propanoic acid was prepared using a procedure analogous to the synthesis of Example 1 except that methyl 4-(4-hydroxybicyclo[2.2.1]heptan-1-yl)butanoate was replaced with 4-hydroxybicyclo[2.2.1]heptan-1-yl 3,5-difluorobenzoate and 5-(bromomethyl)-1-(4-chlorophenyl)-3-methyl-1*H*-pyrazole was replaced with methyl 3-bromopropanoate. LCMS, $[M+H]^+ = 201.1$.

10

79F. Methyl 3-(4-hydroxybicyclo[2.2.1]heptan-1-yloxy)propanoate




[00202] To a solution of 3-(4-hydroxybicyclo[2.2.1]heptan-1-yloxy)propanoic acid (480 mg, 2.40 mmol) in CH_2Cl_2 (6 mL) and MeOH (6 mL) at 0 °C under Ar was added (diazomethyl)trimethylsilane (3.60 mL of a 2 M solution in hexanes, 7.20 mmol) dropwise. The reaction was allowed to warm to rt and stirred for 5 h. The reaction mixture was concentrated *in vacuo* and purified by flash chromatography (SiO_2 ; gradient from 0 to 50% EtOAc/hexanes) to afford the title compound (370 mg, 72% yield) as a colorless oil. LCMS, $[M+H]^+ = 215.1$. ^1H NMR (500 MHz, CDCl_3) δ 3.73 - 3.66 (m, 5H), 2.57 (t, J = 6.5 Hz, 2H), 1.97 - 1.88 (m, 2H), 1.85 - 1.66 (m, 8H).

15


20

79G. 3-Fluoro-5-phenoxybenzoic acid



- [00203] A flask containing a mixture of 3-fluoro-5-hydroxybenzoic acid (2.0 g, 12.8 mmol), Cs₂CO₃ (10.4 g, 32.0 mmol), Cu(I)Br (0.184 g, 1.28 mmol), and ethyl
- 5 2-oxocyclohexanecarboxylate (0.436 g, 2.56 mmol) was evacuated and backfilled with N₂. DMSO (30 mL) was added by syringe and pre-stirred for 10 min. at RT. Then a solution of iodobenzene in DMSO (0.5 mL) was added via syringe under a counter flow of N₂. The tube was sealed, and the mixture was allowed to stir at 65 °C for 24 h. The reaction was cooled to RT, and passed through CELITE®. After being rinsed with
- 10 another 20 mL of EtOAc, the combined filtrates were washed with brine and dried by Na₂SO₄, then was concentrated *in vacuo*. The residue was purified by column chromatography (SiO₂; 10% MeOH/CH₂Cl₂) to give the title compound (2.15 g, 9.26 mmol, 72% yield) as a beige solid. LCMS, [M-H]⁺ = 231.1. ¹H NMR (500 MHz, CDCl₃) δ 7.50 - 7.44 (m, 2H), 7.38 (t, *J*=8.0 Hz, 2H), 7.21 - 7.15 (m, 1H), 7.04 (d, *J*=7.7
- 15 Hz, 2H), 6.89 (d, *J*=9.6 Hz, 1H).


79H. (3-Fluoro-5-phenoxyphenyl)methanol



- [00204] 3-Fluoro-5-phenoxybenzoic acid (2.15 g, 9.26 mmol) was dissolved in THF
- 20 (30 mL) and then borane-THF complex (18.5 mL, 18.5 mmol) was added. After stirring overnight, the reaction was quenched with 1N aq. HCl and then stirred for 30 min. The reaction was extracted with EtOAc. The organic extract was washed with brine, dried over MgSO₄ and concentrated *in vacuo*. The crude product was chromatographed (silica gel column (120 g); continuous gradient from 0 to 30% solvent B over 25 min, hold at

30% solvent B for 10 min, where solvent A = hexanes and solvent B = EtOAc) to give the title compound (1.88 g, 8.62 mmol, 93% yield) as a colorless oil. LCMS, $[M-H]^+ = 217.1$.

5 79I. 1-(Bromomethyl)-3-fluoro-5-phenoxybenzene




[00205] To a 0 °C solution of (3-fluoro-5-phenoxyphenyl)methanol (1.8 g, 8.25 mmol) and CBr₄ (3.01 g, 9.07 mmol) in CH₂Cl₂ (30 mL) was added Ph₃P (2.38 g, 9.07 mmol) was added portionwise. The reaction was allowed to slowly warm to RT over 3 h. TLC
 10 showed complete conversion of the starting material. The reaction was concentrated *in vacuo* and the residue was purified by flash chromatography (SiO₂; 80 g cartridge; A = Hex, B = EtOAc; 30 min. grad.; 0% B to 15% B; flow rate = 30 mL/min). The pure fractions were concentrated *in vacuo* to give the title compound (2.26 g, 8.04 mmol, 97% yield) as a colorless oil. LCMS, $[M+H]^+ = 279.1$. ¹H NMR (500 MHz, CDCl₃) δ 7.42 -
 15 7.35 (m, 2H), 7.22 - 7.17 (m, 1H), 7.08 - 7.01 (m, 2H), 6.86 - 6.79 (m, 2H), 6.62 (dt, *J*=9.9, 2.2 Hz, 1H), 4.39 (s, 2H).

Example 79



[00206] The title compound was prepared using a procedure analogous to the synthesis
 20 of Example 1 except that methyl 4-(4-hydroxybicyclo[2.2.1]heptan-1-yl)butanoate was replaced with methyl 3-(4-hydroxybicyclo[2.2.1]heptan-1-yloxy)propanoate and 5-(bromomethyl)-1-(4-chlorophenyl)-3-methyl-1*H*-pyrazole was replaced with 1-(bromomethyl)-3-fluoro-5-phenoxybenzene. LCMS, $[M+H]^+ = 401.1$. ¹H NMR (500 MHz, CDCl₃) δ 7.40 - 7.34 (m, 2H), 7.19 - 7.13 (m, 1H), 7.07 - 6.99 (m, 2H), 6.83 - 6.78 (m,
 25 1H), 6.77 (s, 1H), 6.58 (dt, *J* = 9.9, 2.3 Hz, 1H), 4.45 (s, 2H), 3.72 (t, *J* = 6.3 Hz, 2H), 2.63 (t, *J* = 6.2 Hz, 2H), 2.01 - 1.88 (m, 4H), 1.81 - 1.70 (m, 6H). HPLC-1: RT = 11.4 min, purity = 97.1%; HPLC-2: RT = 9.6 min, purity = 97.6%.

[00207] The following Examples (Table 4) were prepared in a manner analogous to Example 79.

Table 4




Example	Name	R ¹ -L ₄	LCMS, [M+H] ⁺	¹ H NMR (500 MHz, CDCl ₃) δ	HPLC-1: RT min, purity; HPLC-2: RT min, purity
80	3-(4-(2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yloxy)propanoic acid		381.1	7.53 (dd, <i>J</i> = 7.6, 1.5 Hz, 1H), 7.35 - 7.29 (m, 2H), 7.26 - 7.22 (m, 1H), 7.15 (td, <i>J</i> = 7.4, 1.1 Hz, 1H), 7.10 - 7.05 (m, 1H), 6.98 - 6.92 (m, 2H), 6.89 (dd, <i>J</i> = 8.1, 1.0 Hz, 1H), 4.56 (s, 2H), 3.70 (t, <i>J</i> = 6.2 Hz, 2H), 2.62 (t, <i>J</i> = 6.2 Hz, 2H), 1.99 - 1.84 (m, 4H), 1.77 - 1.64 (m, 6H).	10.9 min, 100% 9.2 min, 97.9%

81	3-(4-(3-fluoro-5-(4-fluorophenoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yloxy)propanoic acid		417.1	7.10 - 7.04 (m, 2H), 7.03 - 6.97 (m, 2H), 6.79 (d, $J = 9.1$ Hz, 1H), 6.72 (s, 1H), 6.53 (dt, $J = 9.9, 2.3$ Hz, 1H), 4.45 (s, 2H), 3.72 (t, $J = 6.2$ Hz, 2H), 2.63 (t, $J = 6.2$ Hz, 2H), 2.00 - 1.88 (m, 4H), 1.82 - 1.70 (m, 6H).	11.4 min, 100% 9.6 min, 95.0%
82	3-(4-(1-(3-fluoro-5-(4-fluorophenoxy)phenyl)ethoxy)bicyclo[2.2.1]heptan-1-yloxy)propanoic acid		431.1	7.10 - 7.04 (m, 2H), 7.04 - 6.98 (m, 2H), 6.77 (dt, $J = 9.0, 1.8$ Hz, 1H), 6.73 (s, 1H), 6.50 (dt, $J = 9.9, 2.3$ Hz, 1H), 4.57 (q, $J = 6.6$ Hz, 1H), 3.66 (t, $J = 6.3$ Hz, 2H), 2.59 (t, $J = 6.2$ Hz, 2H), 1.93 - 1.76 (m, 4H), 1.73 - 1.53 (m, 6H), 1.36 (d, $J = 6.6$ Hz, 3H).	11.7 min, 100% 9.8 min, 95.0%

Example 83


4-(4-(5-Fluoro-2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)-N-(methanesulfonyl)butanamide



- 5 **[00208]** A mixture of 4-(4-(5-fluoro-2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid (20 mg, 0.050 mmol), methanesulfonamide (14 mg, 0.151 mmol), DMAP (19 mg, 0.156 mmol), and EDC (19 mg, 0.100 mmol) in CH_2Cl_2 (0.5 mL) was stirred at rt for 2 h. The reaction was concentrated *in vacuo*. The residue was purified by preparative HPLC (YMC ODS-A-5 μ 30 x 100 mm column; flow rate = 40 mL/min, 20 to 100% solvent B over 12 min, hold to 14 min, where solvent A = 90:10:0.1 H_2O :MeOH:TFA and solvent B = 90:10:0.1 MeOH: H_2O :TFA) to give the title compound (14 mg, 60% yield) as a clear oil. LCMS, $[\text{M}-\text{H}]^+ = 474.0$. ^1H NMR (500 MHz, CDCl_3) δ 7.93 (br. s., 1H), 7.40 - 7.34 (m, 2H), 7.19 - 7.12 (m, 1H), 7.07 - 7.01 (m, 2H), 6.81 (d, $J = 9.1$ Hz, 1H), 6.78 (s, 1H), 6.57 (dd, $J = 9.9, 1.9$ Hz, 1H), 4.47 (s, 2H), 3.32 (s, 3H), 2.33 (t, $J = 7.3$ Hz, 2H), 1.89 - 1.79 (m, 2H), 1.73 - 1.55 (m, 6H), 1.54 - 1.40 (m, 6H). HPLC-1: RT = 12.2 min, purity = 100%; HPLC-2: RT = 10.8 min, purity = 100%.

Example 84

- 20 N-Methyl-4-(4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)butanamide



- [00209]** To a solution of 4-(4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid (10 mg, 0.026 mmol) and methylamine (10 mg of a 33% solution in EtOH, 0.105 mmol) in CH_2Cl_2 (0.8 mL) was added EDCI (10 mg, 0.053 mmol) and DMAP (10 mg, 0.081 mmol). The reaction mixture was stirred overnight at RT, then was partitioned


between CH₂Cl₂ (20 mL) and H₂O (5 mL). The aqueous layer was extracted with CH₂Cl₂ (3 x 5 mL). The combined organic extracts were washed with brine (5 mL), dried over MgSO₄, filtered and evaporated *in vacuo* to afford the crude product. The crude material was purified via preparative LC/MS with the following conditions:

- 5 Column: Waters XBridge C18, 19 x 100 mm, 5-μm particles; Mobile Phase A: 5:95 MeCN:water with 0.1% formic acid; Mobile Phase B: 95:5 MeCN:water with 0.1% formic acid; Gradient: 45-85% B over 10 minutes, then a 5-minute hold at 100% B; Flow: 20 mL/min. Fractions containing the desired product were combined and dried via centrifugal evaporation to give the title compound (9 mg, 0.023 mmol, 88% yield).
- 10 LCMS, [M+H]⁺ = 394.3. ¹H NMR (500M Hz, DMSO-d₆) δ 7.65 (br. s., 1H), 7.39 (t, *J*=7.6 Hz, 2H), 7.33 (t, *J*=7.8 Hz, 1H), 7.14 (t, *J*=7.4 Hz, 1H), 7.08 (d, *J*=7.7 Hz, 1H), 6.99 (d, *J*=8.0 Hz, 2H), 6.94 (s, 1H), 6.88 (d, *J*=8.0 Hz, 1H), 4.46 (s, 2H), 2.54 (d, *J*=4.7 Hz, 3H), 2.01 (t, *J*=7.3 Hz, 2H), 1.79 - 1.70 (m, 2H), 1.59 - 1.29 (m, 12H). HPLC-5: RT = 2.19 min, purity = 100%; HPLC-6: RT = 2.22 min, purity = 100%.

15

Example 85


1-(3-Hydroxyazetidin-1-yl)-4-(4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)butan-1-one




- 20 [00210] Example 85 was prepared in a manner analogous to Example 84 except that 3-hydroxyazetidine was used instead of methylamine. LCMS, [M+H]⁺ = 436.3. ¹H NMR (500M Hz, DMSO-d₆) δ 7.39 (t, *J*=7.4 Hz, 2H), 7.33 (t, *J*=7.8 Hz, 1H), 7.16 - 7.12 (m, 1H), 7.08 (d, *J*=7.7 Hz, 1H), 7.00 (d, *J*=8.3 Hz, 2H), 6.94 (s, 1H), 6.88 (d, *J*=8.0 Hz, 1H), 5.66 (d, *J*=6.1 Hz, 1H), 4.46 (s, 2H), 4.44 - 4.38 (m, 1H), 4.23 (t, *J*=7.7 Hz, 1H), 4.01 -
- 25 3.95 (m, 1H), 3.79 (dd, *J*=8.8, 4.4 Hz, 1H), 3.53 (dd, *J*=10.0, 4.3 Hz, 1H), 1.99 (t, *J*=7.2 Hz, 2H), 1.80 - 1.70 (m, 2H), 1.59 - 1.47 (m, 4H), 1.46 - 1.32 (m, 8H). HPLC-5: RT = 2.09 min, purity = 95.9%; HPLC-6: RT = 2.12 min, purity = 95.5%.

Example 86

3-(4-(2-Phenoxyphenethoxy)bicyclo[2.2.1]heptan-1-yl)propanoic acid




5 86A. 4-((3-Phenoxybenzyl)oxy)bicyclo[2.2.1]heptane-1-carbaldehyde



[00211] To a solution of (4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl) methanol (600 mg, 1.85 mmol) (prepared as described for the synthesis of Example 72) in CH_2Cl_2 (10 mL) was added Dess-Martin periodinane (941 mg, 2.22 mmol). The reaction mixture was stirred at RT until the oxidation was complete (~1h). The reaction solution was filtered through a plug of CELITE[®], washed with sat'd aq. NaHCO_3 and brine and dried over MgSO_4 . Volatiles were removed *in vacuo* to give the crude product, which was chromatographed (silica gel column (40 g); continuous gradient from 0 to 35% solvent B over 30 min, hold at 35% solvent B for 10 min, where solvent A = hexanes and solvent B = EtOAc) to give the title compound (290 mg, 0.90 mmol, 49% yield) as a colorless oil. LCMS, $[\text{M}+\text{CH}_3\text{OH}]^+ = 354$. ^1H NMR (400 MHz, CDCl_3) δ 9.76 (s, 1H), 7.37 - 7.30 (m, 3H), 7.14 - 7.07 (m, 2H), 7.04 - 6.99 (m, 3H), 6.91 (dd, $J=7.9$, 1.8 Hz, 1H), 4.52 (s, 2H), 2.17 - 2.08 (m, 2H), 2.00 - 1.91 (m, 2H), 1.81 - 1.73 (m, 4H), 1.67 - 1.57 (m, 2H).

20

86B. (E)-Methyl 3-(4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)acrylate




[00212] Trimethyl phosphonoacetate (0.195 mL, 1.35 mmol) and DBU (0.203 mL, 1.35 mmol) was added to a suspension of LiCl (57 mg, 1.35 mmol) in MeCN (4 mL) at 0 °C under N_2 . The reaction was stirred for 30 min, and then 4-((3-phenoxybenzyl) oxy)

25

bicyclo[2.2.1]heptane-1-carbaldehyde (290 mg, 0.90 mmol) was added. The reaction was stirred for 2 h at RT, then was concentrated *in vacuo*. The residue was diluted with EtOAc, and then washed in succession with 1N aq. HCl, sat'd aq. NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered, and the filtrate was concentrated *in vacuo*. The crude product was purified by flash chromatography (SiO₂; 40 g column gradient of EtOAc/hexanes (0% to 20% over 20 min.) to give the title compound (322 mg, 0.85 mmol, 95% yield) as a colorless oil. LCMS, [M+H]⁺ = 379.2. ¹H NMR (500 MHz, CDCl₃) δ 7.37 - 7.28 (m, 3H), 7.15 - 7.07 (m, 3H), 7.04 - 6.99 (m, 3H), 6.91 (d, *J*=9.4 Hz, 1H), 5.81 - 5.76 (m, 1H), 4.51 (s, 2H), 3.75 (s, 3H), 1.97 - 1.89 (m, 2H), 1.88 - 1.79 (m, 2H), 1.78 - 1.71 (m, 2H), 1.68 - 1.58 (m, 4H).


86C. Methyl 3-(4-hydroxybicyclo[2.2.1]heptan-1-yl)propanoate



[00213] A solution of (E)-Methyl 3-(4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)acrylate (250 mg, 0.661 mmol) in MeOH (10 mL) was evacuated and flushed with Ar. 10% Pd/C (40 mg, 0.066 mmol) was added and the reaction was evacuated and flushed with an atmosphere of hydrogen. After stirred for overnight, the reaction was filtered and the catalyst was rinsed with EtOAc. The filtrate was concentrated *in vacuo*. The crude product was chromatographed (silica gel column (12 g); continuous gradient from 0 to 40% solvent B over 15 min, hold at 40% solvent B for 10 min, where solvent A = hexanes and solvent B = EtOAc) to give the desired compound (123 mg, 0.62 mmol, 94% yield) as a colorless oil. LCMS, [M+H]⁺ = 199.2. ¹H NMR (500 MHz, CDCl₃): δ 3.67 (s, 3H), 2.34 - 2.26 (m, 2H), 1.83 - 1.77 (m, 2H), 1.76 - 1.54 (m, 8H), 1.48 - 1.41 (m, 2H).

25

86D. 1-(2-Bromoethyl)-2-phenoxybenzene



[00214] To a 0 °C solution of 2-(2-phenoxyphenyl)ethanol (990 mg, 4.62 mmol) and CBr₄ (1.69 g, 5.08 mmol) in CH₂Cl₂ (15 mL) was added Ph₃P (1.33 g, 5.08 mmol) portionwise. The reaction was warmed to RT over 3 h, then was concentrated *in vacuo* and the residue was purified by flash chromatography (SiO₂; A = Hex, B = EtOAc; 15 min. gradient from 0% B to 15% B; flow rate = 30 mL/min) to give the title compound (1.19 g, 4.29 mmol, 93% yield) as a colorless oil. ¹H NMR (500 MHz, CDCl₃) δ 7.37 - 7.32 (m, 2H), 7.30 (dd, *J*=7.6, 1.5 Hz, 1H), 7.23 (td, *J*=7.7, 1.7 Hz, 1H), 7.13 - 7.07 (m, 2H), 6.99 - 6.94 (m, 2H), 6.88 (dd, *J*=8.0, 0.8 Hz, 1H), 3.62 (t, *J*=7.4 Hz, 2H), 3.23 (t, *J*=7.6 Hz, 2H).

Example 86


[00215] To a 0 °C suspension of methyl 3-(4-hydroxybicyclo[2.2.1]heptan-1-yl)propanoate (15 mg, 0.076 mmol), 2,6-di-*tert*-butylpyridine (0.059 mL, 0.265 mmol), and AgOTf (58 mg, 0.23 mmol) in CH₂Cl₂ (1.0 mL) was added 1-(2-bromoethyl)-2-phenoxybenzene (67 mg, 0.24 mmol); a yellow precipitate formed within a few minutes. The reaction was stirred overnight, then was diluted with CH₂Cl₂ and filtered through a plug of CELITE®. The filtrate was concentrated *in vacuo*. The residue was taken up in THF (1 mL) and water (0.5 mL) and MeOH (1 mL) and LiOH.H₂O (13 mg, 0.30 mmol) was added. The reaction was stirred at RT overnight, then was diluted with EtOAc (20 mL) and H₂O (5 mL). The aqueous layer was adjusted with 1N aq. HCl to pH ~3 and extracted with EtOAc (3 x 5 mL). The combined organic extracts were washed with brine (5 mL), dried over MgSO₄, filtered and evaporated *in vacuo* to afford the crude product which was purified via preparative LC/MS (Column: Waters XBridge C18, 19 x 100 mm, 5-μm particles; Guard Column: Waters XBridge C18, 19 x 10 mm, 5-μm particles; Mobile Phase A: 5:95 MeCN:water with 0.1% NH₄OH; Mobile Phase B: 95:5 MeCN:water with 0.1% NH₄OH; Gradient: 10-50% B over 10 min, then a 5-min hold at

100% B; Flow: 20 mL/min) to give the title compound (23 mg, 0.060 mmol, 79% yield).
LCMS, $[M+H]^+ = 381.3$. ^1H NMR (500 MHz, DMSO- d_6) δ 7.39 - 7.33 (m, 3H), 7.27 -
7.21 (m, 1H), 7.15 - 7.05 (m, 2H), 6.89 (d, $J=8.3$ Hz, 3H), 3.52 (t, $J=7.3$ Hz, 2H), 2.74 (t,
 $J=7.2$ Hz, 2H), 2.13 - 2.06 (m, 2H), 1.59 (t, $J=8.1$ Hz, 4H), 1.47 - 1.38 (m, 4H), 1.35 -
5 1.19 (m, 4H). HPLC-5: RT = 1.80 min, purity = 100%; HPLC-6: RT = 1.97 min, purity
= 100%.



[00216] The following Examples (Table 5) were prepared in a manner analogous to
Example 86.


10

Table 5




Example	Name	R ¹ -L ₄ -	LCMS, [M-H] ⁺	¹ H NMR (500 MHz, CDCl ₃) δ	HPLC-5: RT, purity; HPLC-6: RT, purity
87	3-(4-(2-(naphthalen-1-yl)ethoxy) bicyclo[2.2.1]heptan-1-yl) propanoic acid		337.1	¹ H NMR (500MHz, DMSO-d ₆) δ 8.06 (d, <i>J</i> =8.3 Hz, 1H), 7.91 (d, <i>J</i> =8.0 Hz, 1H), 7.78 (d, <i>J</i> =7.7 Hz, 1H), 7.58 - 7.48 (m, 2H), 7.45 - 7.38 (m, 2H), 3.69 (t, <i>J</i> =7.3 Hz, 2H), 3.26 - 3.22 (t, <i>J</i> =7.3 Hz, 2H), 2.15 - 2.09 (m, 2H), 1.71 - 1.58 (m, 4H), 1.53 - 1.43 (m, 4H), 1.38 - 1.27 (m, 4H).	1.72 min, 98.9% 1.93 min, 99.5%

88	3-(4-(3-phenoxyphenethoxy) bicyclo[2.2.1]heptan-1-yl) propanoic acid		379.1	¹ H NMR (500MHz, DMSO-d ₆) δ 7.38 (t, <i>J</i> =8.0 Hz, 2H), 7.28 (t, <i>J</i> =7.8 Hz, 1H), 7.16 - 7.10 (m, 1H), 7.03 - 6.96 (m, 3H), 6.90 (s, 1H), 6.83 (dd, <i>J</i> =8.1, 2.1 Hz, 1H), 3.56 (t, <i>J</i> =6.6 Hz, 2H), 2.74 (t, <i>J</i> =6.7 Hz, 2H), 2.17 - 2.10 (m, 2H), 1.66 - 1.57 (m, 4H), 1.51 - 1.41 (m, 4H), 1.38 - 1.21 (m, 4H).	1.94 min, 99.2% 2.37 min, 100%
89	3-((1S,4R)-4-(2-fluoro-5-(trifluoromethoxy) phenethoxy)bicyclo[2.2.1] heptan-1-yl)propanoic acid		389.2	¹ H NMR (500MHz, CDCl ₃) δ 7.19 - 7.15 (m, 1H), 7.07 - 6.99 (m, 2H), 3.64 (t, <i>J</i> =6.7 Hz, 2H), 2.89 (t, <i>J</i> =6.7 Hz, 2H), 2.35 - 2.30 (m, 2H), 1.82 - 1.69 (m, 4H), 1.62 - 1.51 (m, 4H), 1.48 - 1.40 (m, 2H), 1.35 (s, 2H).	HPLC-1: 11.7 min, 100% HPLC-2: 9.5 min, 100%

90	3-(4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)propanoic acid		383.2	¹ H NMR (500MHz, DMSO-d ₆) δ 7.43 (t, <i>J</i> =8.0 Hz, 2H), 7.23 - 7.17 (m, 1H), 7.06 (d, <i>J</i> =8.0 Hz, 2H), 6.88 (d, <i>J</i> =9.4 Hz, 1H), 6.77 - 6.70 (m, 2H), 4.46 (s, 2H), 2.18 - 2.12 (m, 2H), 1.78 - 1.69 (m, 2H), 1.67 - 1.61 (m, 2H), 1.58 - 1.46 (m, 4H), 1.41 - 1.31 (m, 4H).	1.96 min, 99.4% 2.26 min, 100%
----	--	--	-------	--	-----------------------------------


Example 91

2-((4-((3-Fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methyl)
cyclopropanecarboxylic acid




5

91A. (4-((3-Phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methyl methanesulfonate



[00217] MsCl (0.203 mL, 2.60 mmol) was added dropwise to a 0 °C solution of (4-((3-
phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methanol (650 mg, 2.0 mmol) and Et₃N
10 (0.84 mL, 6.0 mmol) in CH₂Cl₂ (6 mL) under N₂. The reaction mixture was then stirred
at 0 °C for 2 h and then concentrated *in vacuo*. The residue was dissolved in EtOAc (50
mL), washed with 1N aq. HCl, water and brine. The organic layer was dried over
MgSO₄ and filtered. The filtrate was concentrated *in vacuo* to provide the title compound
(801 mg, 1.99 mmol, 99% yield) as a light yellowish oil. It was used directly in the next
15 reaction without further purification. LCMS, [M+NH₄]⁺ = 420.2.


91B. 2-(4-((3-Phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)acetonitrile



[00218] A mixture of (4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methyl
20 methanesulfonate (807 mg, 2.0 mmol), tetrabutylammonium iodide (7 mg, 0.020 mmol)
and NaCN (0.98 g, 20 mmol) in DMSO (10 mL) was stirred at 80 °C under N₂ for 18 h.
The reaction was cooled to RT and diluted with EtOAc (50 mL). The organic phase was
washed with water (3 x 10 mL), dried over MgSO₄, filtered, and concentrated *in vacuo*.
The crude oil was purified by flash chromatography (SiO₂; EtOAc/hexane (gradient from
25 0% to 30% over 30 min.) to give the title compound (625 mg, 1.87 mmol, 94% yield) as a

colorless oil. LCMS, $[M+NH_4]^+ = 351.2$. 1H NMR (500 MHz, $CDCl_3$) δ 7.37 - 7.28 (m, 3H), 7.13 - 7.06 (m, 2H), 7.04 - 6.99 (m, 3H), 6.91 (dd, $J=8.1, 2.1$ Hz, 1H), 4.51 (s, 2H), 2.48 (s, 2H), 2.00 - 1.91 (m, 2H), 1.81 - 1.71 (m, 4H), 1.67 - 1.60 (m, 4H).

5 91C. 2-(4-((3-Phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)acetaldehyde




[00219] DIBAL-H (2.42 mL of a 1.0 M solution in CH_2Cl_2 , 2.42 mmol) was added dropwise to a $-78^\circ C$ solution of 2-(4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-

10 yl)acetonitrile (620 mg, 1.86 mmol) in CH_2Cl_2 (6 mL) under N_2 . The reaction mixture was stirred at $-78^\circ C$ for 2 h and then quenched with 1N aq. HCl at $0^\circ C$. The mixture was extracted with EtOAc. The organic layer was washed with brine, dried over $MgSO_4$, and concentrated *in vacuo*. The crude oil was purified by flash chromatography (SiO_2 ; EtOAc/hexane (gradient from 0% to 30% over 20 min.) to give the title compound (470 mg, 1.40 mmol, 75% yield) as a colorless oil. LCMS, $[M-H]^+ = 335.2$. 1H NMR (500

15 MHz, $CDCl_3$) δ 9.80 (t, $J=2.6$ Hz, 1H), 7.37 - 7.28 (m, 3H), 7.13 - 7.07 (m, 2H), 7.04 - 6.97 (m, 3H), 6.91 (d, $J=1.7$ Hz, 1H), 4.51 (s, 2H), 2.53 (s, 2H), 1.94 - 1.85 (m, 2H), 1.77 - 1.60 (m, 8H).

91D. (E)-Methyl 4-(4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)but-2-enoate




20

[00220] Trimethyl phosphonoacetate (0.302 mL, 2.10 mmol) and DBU (0.32 mL, 2.1 mmol) were added to a $0^\circ C$ suspension of LiCl (89 mg, 2.1 mmol) in MeCN (4 mL) under N_2 . The reaction was stirred for 30 min, then 2-(4-((3-phenoxybenzyl)oxy)bicyclo

25 [2.2.1]heptan-1-yl)acetaldehyde (470 mg, 1.40 mmol) was added. The reaction was stirred for 2 h at room RT, then was concentrated *in vacuo*. The residue was diluted with EtOAc, and then washed in succession with 1N aq. HCl, sat'd aq. $NaHCO_3$ and brine,


dried (MgSO_4) and concentrated *in vacuo*. The residual crude oil was purified by flash chromatography (SiO_2 ; EtOAc/hexanes (gradient from 0% to 20% over 20 min.) to give the title compound (490 mg, 1.25 mmol, 89% yield) as a colorless oil. LCMS, $[\text{M}+\text{H}]^+ = 393.2$. ^1H NMR (500 MHz, CDCl_3) δ 7.36 - 7.28 (m, 3H), 7.13 - 7.06 (m, 2H), 7.03 - 6.99 (m, 3H), 6.99 - 6.87 (m, 2H), 5.86 - 5.80 (m, 1H), 4.49 (s, 2H), 3.74 (d, $J=1.1$ Hz, 3H), 2.33 (d, $J=8.0$ Hz, 2H), 1.91 - 1.82 (m, 2H), 1.72 - 1.53 (m, 8H).

91E. Methyl 2-((4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methyl)cyclopropanecarboxylate




[00221] To a vigorously stirred 0 °C mixture of Et_2O (5 mL) and aq. 40% KOH (2 mL) was added N-methyl-N'-nitro-N-nitrosoguanidine (1.0 g, 6.8 mmol) portionwise over 15 min. Upon completed of addition, the aqueous layer was separated. The ether solution was dried with KOH at 0 °C for 5 min, decanted into KOH pellets again and then poured onto a 0 °C solution of (E)-methyl 4-(4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)but-2-enoate (460 mg, 1.17 mmol) in THF (5 mL). $\text{Pd}(\text{OAc})_2$ (26 mg, 0.12 mmol) was added and the reaction was allowed to warm to RT and stirred for 1 h. The reaction was concentrated *in vacuo* and the residual crude oil was purified by flash chromatography (SiO_2 ; EtOAc/ hexane; gradient from 0% to 20% over 20 min) to give the title compound (451 mg, 1.11 mmol, 95% yield) as a colorless oil. LCMS, $[\text{M}+\text{H}]^+ = 407.2$. ^1H NMR (500 MHz, CDCl_3) δ 7.36 - 7.28 (m, 3H), 7.13 - 7.07 (m, 2H), 7.04 - 6.99 (m, 3H), 6.89 (d, $J=8.3$ Hz, 1H), 4.50 (s, 2H), 3.68 (s, 3H), 1.90 - 1.81 (m, 2H), 1.72 - 1.15 (m, 14H).

91F. Methyl 2-((4-hydroxybicyclo[2.2.1]heptan-1-yl)methyl)cyclopropanecarboxylate



- [00222] A solution of methyl 2-((4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methyl)cyclopropanecarboxylate (425 mg, 1.045 mmol) in MeOH (10 mL) was
 5 evacuated and flushed with Ar, after which 10% Pd/C (20 mg, 0.11 mmol) was added. The reaction was evacuated and flushed with an atmosphere of hydrogen and stirred at RT overnight. The mixture was filtered and the catalyst was washed with EtOAc. The combined filtrates were concentrated *in vacuo* to give the crude product which was
 10 chromatographed (SiO₂; continuous gradient from 0 to 40% solvent B over 15 min, hold at 40% solvent B for 10 min, where solvent A = hexanes and solvent B = EtOAc) to give the title compound (226 mg, 1.01 mmol, 96% yield) as a colorless oil. LCMS, [M+H]⁺ = 225.2. ¹H NMR (500 MHz, CDCl₃) δ 3.68 (s, 3H), 1.77 - 1.30 (m, 14H), 1.21 - 1.15 (m, 1H), 0.68 (td, *J*=6.9, 4.1 Hz, 1H).

15 91G. Methyl 2-((4-hydroxybicyclo[2.2.1]heptan-1-yl)methyl)cyclopropanecarboxylate




- [00223] To a 0 °C suspension of methyl 2-((4-hydroxybicyclo[2.2.1]heptan-1-yl)methyl)cyclopropanecarboxylate (25 mg, 0.111 mmol), 2,6-di-*tert*-butylpyridine (0.045 mL, 0.201 mmol), and AgOTf (43 mg, 0.17 mmol) in CH₂Cl₂ (1.2 mL) was added 1-
 20 (bromomethyl)-3-fluoro-5-phenoxybenzene (50 mg, 0.18 mmol); a yellow precipitate formed within a few minutes. The reaction was stirred overnight at RT, then diluted with CH₂Cl₂ and filtered through a plug of CELITE[®]. The filtrate was concentrated *in vacuo* and used directly without further purification for the next step.


Example 91

[00224] LiOH.H₂O (30 mg, 0.71 mmol) was added to methyl 2-((4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methyl)cyclopropanecarboxylate (76 mg, 0.18 mmol) in THF (1 mL) and water (0.5 mL) at RT. The reaction was stirred overnight, then was diluted with EtOAc (30 mL) and H₂O (5 mL). The aqueous layer was adjusted with 1N aq. HCl to pH ~3 and extracted with EtOAc (3 x 10 mL). The combined organic extracts were washed with brine (5 mL), dried (MgSO₄), filtered and concentrated *in vacuo* to afford the crude product which was purified via preparative LC/MS [Column: Waters XBridge C18, 19 x 200 mm, 5-μm particles; Guard Column: Waters XBridge C18, 19 x 10 mm, 5-μm particles; Mobile Phase A: 5:95 MeCN:water with 0.1% TFA; Mobile Phase B: 95:5 MeCN:water with 0.1% TFA; Gradient: 50-90% B over 20 minutes, then a 5-minute hold at 100% B; Flow: 20 mL/min] to give the title compound (32 mg, 0.077 mmol, 43% yield). LCMS, [M+Na]⁺ = 433.2. ¹H NMR (500 MHz, DMSO-d₆) δ 7.45 - 7.40 (m, 2H), 7.22 - 7.18 (m, 1H), 7.06 (d, *J*=7.7 Hz, 2H), 6.89 (d, *J*=9.1 Hz, 1H), 6.76 (s, 1H), 6.72 (dd, *J*=10.0, 2.1 Hz, 1H), 4.46 (s, 2H), 1.79 - 1.69 (m, 2H), 1.63 - 1.52 (m, 4H), 1.47 - 1.35 (m, 6H), 1.23 (dt, *J*=8.1, 4.2 Hz, 1H), 1.17 - 1.08 (m, 1H), 0.95 (dt, *J*=8.6, 4.1 Hz, 1H), 0.65 - 0.59 (m, 1H). HPLC-5: RT = 2.17 min, purity = 100%; HPLC-6: RT = 2.37 min, purity = 100%.

20 Example 92



and Example 93



[00225] The two enantiomers of racemic Example 91 were separated by preparatory chiral SFC chromatography with a Berger Multigram II SFC chromatograph using the following method: UV visualization at 220 nm; Column: CHIRALPAK[®] AD-H SFC,

250 x 30 mm ID, 5 μ m; Flow rate: 55.0 mL/min, 100 bar backpressure, 40 °C and Mobile Phase: 17% methanol / 83% CO₂. Injection Details: 0.5 mL of ~12 mg/mL in CAN.


[00226] Analytical chiral SFC chromatography was performed on a Thar Analytical SFC chromatography instrument using the following method: UV visualization at 220 nm; Column: CHIRALPAK[®] AD-H, 250 x 4.6 mm ID, 5 μ m; Flow rate: 2.0 mL/min, 100 bar backpressure, 35 °C; and Mobile Phase: 20% MeOH/ 80% CO₂. Injection Details: 10 μ L of 1 mg/mL in MeCN.

[00227] Example 92 (Enantiomer 1): LCMS, [M-H]⁺ = 409.3. ¹H NMR (500 MHz, CDCl₃) δ 7.36 (t, *J*=7.6 Hz, 2H), 7.18 - 7.12 (m, 1H), 7.07 - 7.00 (m, 2H), 6.82 (d, *J*=9.1 Hz, 1H), 6.78 (s, 1H), 6.57 (d, *J*=9.9 Hz, 1H), 4.47 (s, 2H), 1.90 - 1.80 (m, 2H), 1.67 (d, *J*=6.3 Hz, 4H), 1.59 - 1.32 (m, 8H), 1.28 - 1.22 (m, 1H), 0.79 - 0.72 (m, 1H). HPLC-1: RT = 12.8 min, purity = 100%; HPLC-2: RT = 10.5 min, purity = 100%.

[00228] Example 93 (Enantiomer 2): LCMS, [M-H]⁺ = 409.3. ¹H NMR (500 MHz, CDCl₃) δ 7.40 - 7.34 (m, 2H), 7.18 - 7.13 (m, 1H), 7.06 - 7.01 (m, 2H), 6.83 (s, 1H), 6.78 (s, 1H), 6.57 (dt, *J*=9.9, 2.2 Hz, 1H), 4.47 (s, 2H), 1.90 - 1.80 (m, 2H), 1.73 - 1.62 (m, 4H), 1.59 - 1.32 (m, 8H), 1.28 - 1.22 (m, 1H), 0.79 - 0.72 (m, 1H). HPLC-1: RT = 12.8 min, purity = 100%; HPLC-2: RT = 10.5 min, purity = 100%.

20 Example 94

(E)-4-(4-((3-Phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)but-2-enoic acid




[00229] LiOH.H₂O (11 mg, 0.26 mmol) was added to (E)-methyl 4-(4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)but-2-enoate (25 mg, 0.064 mmol) in THF (1 mL) and water (0.5 mL) and the reaction was stirred at RT overnight. The reaction was diluted with EtOAc (20 mL) and H₂O (5 mL). The aqueous layer was adjusted with 1N aq. HCl to pH ~3 and extracted with EtOAc (3 x 5 mL). The combined organic layers were washed with brine (5 mL), dried over MgSO₄, filtered and evaporated *in vacuo*. The residual crude product was purified by preparative LC/MS [Column: Waters XBridge C18, 19 x 100 mm, 5- μ m particles; Guard Column: Waters XBridge C18, 19 x 10 mm, 5-

μm particles; Mobile Phase A: 5:95 MeCN:water with 0.1% TFA; Mobile Phase B: 95:5 MeCN:water with 0.1% TFA; Gradient: 45-85% B over 10 min, then a 7-min hold at 100% B; Flow: 20 mL/min] to give the title compound (21 mg, 0.053 mmol, 84% yield). LCMS, $[M-H]^+ = 377.2$. 1H NMR (500 MHz, DMSO- d_6) δ 7.41 - 7.36 (m, 2H), 7.33 (t, $J=7.8$ Hz, 1H), 7.16 - 7.11 (m, 1H), 7.07 (d, $J=8.0$ Hz, 1H), 7.01 - 6.97 (m, 2H), 6.93 (s, 1H), 6.90 - 6.85 (m, 1H), 6.80 - 6.72 (m, 1H), 5.77 (d, $J=15.4$ Hz, 1H), 4.46 (s, 2H), 2.29 (d, $J=7.7$ Hz, 2H), 1.80 - 1.71 (m, 2H), 1.60 - 1.50 (m, 4H), 1.42 - 1.35 (m, 4H). HPLC-5: RT = 1.88 min, purity = 96.4%; HPLC-6: RT = 2.18 min, purity = 99.2%.

10 Example 95

2-((4-((3-Phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methyl)cyclopropanecarboxylic acid




[00230] LiOH.H₂O (10 mg, 0.246 mmol) was added to methyl 2-((4-((3-


- 15 phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methyl)cyclopropanecarboxylate (25 mg, 0.061 mmol) in THF (1 mL) and water (0.5 mL) and the reaction was stirred at RT overnight. The reaction was diluted with EtOAc (20 mL) and H₂O (5 mL). The aqueous layer was adjusted with 1N aq. HCl to pH ~3 and extracted with EtOAc (3 x 5 mL). The combined organic extracts were washed with brine (10 mL), dried over MgSO₄, filtered
 20 and concentrated *in vacuo* to afford the crude product which was purified by preparative LC/MS [Column: Waters XBridge C18, 19 x 100 mm, 5-μm particles; Guard Column: Waters XBridge C18, 19 x 10 mm, 5-μm particles; Mobile Phase A: 5:95 MeCN:water with 0.1% TFA; Mobile Phase B: 95:5 MeCN:water with 0.1% TFA; Gradient: 45-85% B over 10 min, then a 5-min hold at 100% B; Flow: 20 mL/min] to give the title
 25 compound (18 mg, 0.045 mmol, 73% yield). LCMS, $[M-H]^+ = 391.2$. 1H NMR (500 MHz, CD₃OD) δ 7.33 - 7.24 (m, 3H), 7.10 - 7.03 (m, 2H), 6.98 - 6.94 (m, 3H), 6.86 (dd, $J=8.1, 2.1$ Hz, 1H), 4.48 (s, 2H), 1.88 - 1.78 (m, 2H), 1.71 - 1.62 (m, 4H), 1.56 - 1.38 (m, 6H), 1.33 - 1.23 (m, 2H), 1.12 (dt, $J=8.6, 4.4$ Hz, 1H), 0.66 (ddd, $J=8.0, 6.4, 4.0$ Hz, 1H). HPLC-5: RT = 2.07 min, purity = 100%; HPLC-6: RT = 2.29 min, purity = 95.0%.

30

Example 96 (S-Enantiomer) and Example 97 (R-Enantiomer). 4-(4-((3-Fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)butane-1,2-diol




5 96A. 1-(Bromomethyl)-4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptane



[00231] To a 0 °C solution of 4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methanol (150 mg, 0.44 mmol; prepared as described for the synthesis of Example 72) and CBr₄ (160 mg, 0.48 mmol) in CH₂Cl₂ (3 mL) was added Ph₃P (126 mg,

10 0.48 mmol) portionwise. The reaction was allowed to warm to RT over 3 h, then was concentrated *in vacuo*. The residue was purified by flash chromatography [SiO₂; A = hexanes, B = EtOAc; 25 min gradient; 0% B to 20% B; flow rate = 30 mL/min] to give the title compound (160 mg, 0.40 mmol, 90% yield) as a colorless oil. LCMS, [M+Na]⁺ = 428.9. ¹H NMR (500 MHz, CDCl₃) δ 7.40 - 7.34 (m, 2H), 7.18 - 7.13 (m, 1H), 7.04 (d, *J*=7.7 Hz, 2H), 6.84 - 6.75 (m, 2H), 6.58 (dt, *J*=10.0, 2.3 Hz, 1H), 4.47 (s, 2H), 3.51 (s, 15 *J*=7.7 Hz, 2H), 1.96 - 1.86 (m, 2H), 1.81 - 1.71 (m, 4H), 1.64 - 1.52 (m, 4H).

96B. 1-(But-3-en-1-yl)-4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptane



20 [00232] A solution of 1-(bromomethyl)-4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptane (80 mg, 0.20 mmol) in diisopropyl ether (0.5 mL) was added to Cu(OTf)₂ (4 mg, 10 μmol) under Ar followed by allylmagnesium bromide (0.59 mL, 0.59 mmol) dropwise. The reaction mixture was stirred at RT for 2 h. The reaction was quenched

with saturated aq. NH_4Cl and extracted with EtOAc. The organic were combined, washed with brine, dried over MgSO_4 , filtered and concentrated *in vacuo*. The crude product was chromatographed (SiO_2 ; continuous gradient from 0 to 10% solvent B over 15 min, held at 10% solvent B for 10 min, where solvent A = hexanes and solvent B = EtOAc) to give the title compound (52 mg, 0.14 mmol, 71% yield) as a colorless oil. LCMS, $[\text{M}+\text{Na}]^+ = 389.2$.

96C. Example 96 (S-Enantiomer) and Example 97 (R-Enantiomer)

- [00233] To a 5 °C mixture of $(\text{DHQ})_2\text{PHAL}$ (AD-mix- α) (191 mg, 0.25 mmol), *t*-BuOH (1 mL) and water (1 mL) was added 1-(but-3-en-1-yl)-4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptane (50 mg, 0.14 mmol) and the reaction was stirred at 5 °C for 2 days. To this mixture was added sodium sulfite (206 mg, 1.64 mmol) and the reaction was warmed to RT and stirred at RT, upon which the yellow color disappeared. The reaction was diluted with water (10 mL) and extracted with EtOAc (2 x 30 mL). The combined organic extracts were washed with brine, dried over Na_2SO_4 , filtered and concentrated *in vacuo*. The residue was purified via preparative LC/MS [Column: Waters XBridge C18, 19 x 200 mm, 5- μm particles; Guard Column: Waters XBridge C18, 19 x 10 mm, 5- μm particles; Mobile Phase A: 5:95 MeCN:water with 0.1% TFA; Mobile Phase B: 95:5 MeCN:water with 0.1% TFA; Gradient: 50-90% B over 20 min, then a 5-min hold at 100% B; Flow: 20 mL/min]. The racemate was separated into the 2 individual enantiomers (Example 96 and 97) by preparatory chiral SFC chromatography [Berger Multigram II SFC chromatograph using the following method: UV visualization at 220 nm; Column: CHIRALPAK[®] AD-H SFC, 250 x 30 mm ID, 5 μm ; Flow rate: 85.0 mL/min, 150 bar backpressure, 40 °C; Mobile Phase: 20% IPA / 80% CO_2 ; Injection Details: 0.5 mL of ~14 mg/mL (4:1) ACN:IPA; Analytical chiral SFC chromatography was performed on a Thar Analytical SFC chromatography instrument using the following method: UV visualization at 220 nm; Column: CHIRALPAK[®] AD-H, 250 x 4.6 mm ID, 5 μm ; Flow rate: 2.0 mL/min, 100 bar backpressure, 35 °C; and Mobile Phase: 25% IPA / 75% CO_2].
- [00234] Example 96: (S)-Enantiomer (14 mg, 0.033 mmol, 25% yield): LCMS, $[\text{M}+\text{Na}]^+ = 423.3$. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ : 7.43 (t, $J=7.7$ Hz, 2H), 7.23 - 7.17


(m, 1H), 7.23 - 7.17 (m, 1H), 7.06 (d, $J=7.7$ Hz, 2H), 6.89 (d, $J=9.4$ Hz, 1H), 6.77 - 6.69 (m, 2H), 4.46 (s, 2H), 4.44 - 4.34 (m, 2H), 3.28 - 3.17 (m, 1H), 1.78 - 1.68 (m, 2H), 1.61 - 1.10 (m, 12H). HPLC-5: RT = 2.10 min, purity = 97%; HPLC-6: RT = 2.13 min, purity = 100%.

- 5 **[00235]** Example 97: (R)-Enantiomer (3 mg, 7.9 μ mol, 6% yield): LCMS, $[M+Na]^+ = 423.3$. 1H NMR (500 MHz, DMSO- d_6) δ 7.43 (t, $J=7.4$ Hz, 2H), 7.23 - 7.17 (m, 1H), 7.06 (d, $J=8.3$ Hz, 2H), 6.89 (d, $J=9.1$ Hz, 1H), 6.77 - 6.70 (m, 2H), 4.46 (s, 2H), 4.44 - 4.34 (m, 2H), 3.23 (d, $J=14.9$ Hz, 1H), 1.73 (br. s., 2H), 1.61 - 1.10 (m, 12H). HPLC-5: RT = 2.08 min, purity = 99%; HPLC-6: RT = 2.11 min, purity = 100%.

10

Example 98

N-(2-Amino-2-oxoethyl)-4-(4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)butanamide




15

- [00236]** Example 98 (8 mg; 70% yield; colorless oil) was prepared in a manner analogous to Example 84 except that 4-(4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid was used instead of 4-(4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid and 2-aminoacetamide hydrochloride was used instead of methylamine. LCMS, $[M+Na]^+ = 477.3$. 1H NMR (500 MHz, DMSO- d_6) δ 7.90 (t, $J=5.8$ Hz, 1H), 7.45 - 7.39 (m, 2H), 7.25 - 7.17 (m, 2H), 7.08 - 7.04 (m, 2H), 6.96 (br. s., 1H), 6.89 (d, $J=8.5$ Hz, 1H), 6.77 - 6.70 (m, 2H), 4.46 (s, 2H), 3.60 (d, $J=5.8$ Hz, 2H), 2.10 (t, $J=7.3$ Hz, 2H), 1.77 - 1.69 (m, 2H), 1.57 - 1.30 (m, 12H). HPLC-5: RT = 2.03 min, purity = 98%; HPLC-6: RT = 2.03 min, purity = 100%.

25

Example 99


2-(4-((3-Fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)ethanol



- [00237] $\text{BH}_3 \cdot \text{THF}$ (0.227 mL, 0.227 mmol) was added dropwise (gas evolution observed) to a -15°C solution of 2-(4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)acetic acid (70 mg, 0.189 mmol) in THF (1 mL) and the resulting solution was allowed to warm to 0°C over 2 h. The reaction was carefully quenched (gas evolution observed) with MeOH (~ 5 mL) and volatiles were removed *in vacuo*. The reaction mixture was then neutralized with 1N aq. HCl. The mixture was diluted with 5% aq. NaHCO_3 and extracted with EtOAc (3 x 10 mL). The combined organic extracts were washed with brine, dried (MgSO_4) and filtered. The filtrate was concentrated *in vacuo* to afford the crude product, which was chromatographed (SiO_2 ; 8 g; continuous gradient from 0 to 60% EtOAc/hexanes over 15 min, then held at 60% EtOAc/40% hexane for 10 min) to give the title compound (66 mg, 0.18 mmol, 96% yield) as a colorless oil. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 7.46 - 7.38 (m, 2H), 7.19 (t, $J=7.3$ Hz, 1H), 7.05 (d, $J=7.6$ Hz, 2H), 6.87 (d, $J=9.2$ Hz, 1H), 6.76 - 6.66 (m, 2H), 4.44 (s, 2H), 3.49 - 3.40 (m, 2H), 2.05 (s, 1H), 1.75 - 1.66 (m, 2H), 1.59 - 1.46 (m, 6H), 1.43 - 1.32 (m, 4H). HPLC-5: RT = 2.25 min, purity = 98%; HPLC-6: RT = 2.19 min, purity = 100%.

20 Example 100

2-(2-(4-((3-Fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)ethoxy)acetic acid




- [00238] To a 0°C solution of 2-(4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)ethanol (60 mg, 0.168 mmol) in toluene (1.2 mL) was added a solution of NaOH (0.7 g in 1.2 mL of H_2O), followed by Bu_4NHSO_4 (44 mg, 0.13 mmol). The mixture was stirred at 0°C for 30 min, after which *tert*-butyl 2-bromoacetate (0.05 mL,

0.337 mmol) was added and the mixture was stirred for 14 h at rt. The reaction was neutralized using concentrated HCl and then extracted with EtOAc (2 x 5 mL). The combined organic layers were washed with brine (5 mL), dried (MgSO₄), filtered and concentrated *in vacuo* to afford the crude *t*-butyl ester product, which was dissolved in THF (1 mL). To this solution were added LiOH·H₂O (35 mg, 0.84 mmol), water (0.5 mL) and MeOH (1 mL), after which the reaction was stirred at RT overnight. The reaction was diluted with EtOAc (5 mL) and H₂O (5 mL). The aqueous layer was extracted with EtOAc (2 x 5 mL). The combined organic layers were washed with H₂O (3 x 5 mL). The combined aqueous layers were adjusted to pH ~3 with 1N aq. HCl and extracted with EtOAc (3 x 20 mL). The combined organic layers were washed with brine (10 mL), dried (MgSO₄), filtered and evaporated *in vacuo*. The residual crude product was purified by preparative LC/MS [Column: Waters XBridge C18, 19 x 200 mm, 5-μm particles; Guard Column: Waters XBridge C18, 19 x 10 mm, 5-μm particles; Mobile Phase A: 5:95 MeCN:water with 0.1% TFA; Mobile Phase B: 95:5 MeCN:water with 0.1% TFA; Gradient: 45-90% B over 20 min, then a 5-min hold at 100% B; Flow: 20 mL/min] to give the title compound (44 mg, 0.104 mmol, 62% yield). LCMS, [M-H]⁺ = 413.0. ¹H NMR (500 MHz, DMSO-d₆) δ 7.45 - 7.38 (m, 2H), 7.19 (t, *J*=7.3 Hz, 1H), 7.05 (d, *J*=7.9 Hz, 2H), 6.87 (d, *J*=8.8 Hz, 1H), 6.77 - 6.67 (m, 2H), 4.44 (s, 2H), 3.90 (s, 2H), 3.44 (m, 2H), 1.76 - 1.60 (m, 4H), 1.59 - 1.47 (m, 4H), 1.45 - 1.33 (m, 4H). HPLC-5: RT = 1.69 min, purity = 98%; HPLC-6: RT = 2.21 min, purity = 98%.

Claims

1. A compound of Formula (I):



5 or a stereoisomer, a tautomer, a solvate or a pharmaceutically acceptable salt thereof, wherein:

L_1 is independently L_4 -O or O- L_4 ;

L_2 is independently a hydrocarbon linker substituted with 0-2 R^c , a hydrocarbon-heteroatom linker substituted with 0-2 R^c , or $-(CH_2)_{1-2}$ -(C_{3-4} cycloalkyl substituted with 0-2 R^c)- $(CH_2)_{0-1}$ -; wherein said hydrocarbon linker has one to six carbon atoms and may be straight or branched, saturated or unsaturated; and said hydrocarbon-heteroatom linker has one to four carbon atoms and one group selected from O, -CO-, S, -SO-, -SO₂-, NH, N(C_{1-4} alkyl), -CONH-, and -NHCO-;

15 L_4 is independently a bond or a hydrocarbon linker; wherein said hydrocarbon linker has one to four carbon atoms and may be straight or branched;

R^1 is independently selected from: C_{6-10} carbocycle and a 5- to 10-membered heterocycle comprising carbon atoms and 1-4 heteroatoms selected from N, NR^b , O, and $S(O)_p$; wherein said carbocycle and heterocycle are substituted with 0-4 R^3 and 0-1 R^4 ;

20 R^2 independently selected from: OH, CO₂H, CO₂(C_{1-4} alkyl), $CONR^eNR^f$, and -CONHSO₂ R^f ;

R^3 , at each occurrence, is independently selected from: halogen, C_{1-6} alkyl, C_{1-4} alkoxy, C_{1-4} alkylthio, C_{1-4} haloalkyl, C_{1-4} haloalkoxy, C_{1-4} haloalkylthio, and NO₂;

R^4 is independently - L_3 - R^5 ;

25 L_3 is independently selected from: a bond, O, and C(=O);

R^5 is independently selected from: phenyl and a 5- to 6-membered heteroaryl comprising carbon atoms and 1-4 heteroatoms selected from N, NR^b , O, and $S(O)_p$;

wherein each ring moiety is substituted with 0-2 R^a;

R^a, at each occurrence, is independently selected from: halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkyl;

R^b, at each occurrence, is independently selected from: H, C₁₋₄ alkyl, and
5 -(CH₂)₀₋₂-(phenyl substituted with 0-3 R^d);

R^c, at each occurrence, is independently selected from: =O, halogen, OH, C₁₋₄ alkyl, C₁₋₄ haloalkyl, and C₁₋₄ alkoxy;

R^d, at each occurrence, is independently selected from: halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, and C₁₋₄ haloalkyl;

10 R^e, at each occurrence, is independently selected from: H and C₁₋₄ alkyl;

R^f, at each occurrence, is independently selected from: H, C₁₋₄ alkyl, -(CH₂)₀₋₂-phenyl, and C₃₋₆ cycloalkyl substituted with 1-2 OH;

R^e and R^f may be combined with the nitrogen atom to which they are attached to form a 4- to 6-membered heterocyclic ring comprising carbon atoms and 1 additional
15 heteroatom selected from N, NR^b, O, and S; wherein said heterocycle is substituted with and 0-1 OH; and

p is, independently at each occurrence, selected from 0, 1, and 2.

2. A compound according to claim 1, wherein:

20 L₁ is independently L₄-O;


L₂ is independently a hydrocarbon linker substituted with 0-1 R^c,
a hydrocarbon-heteroatom linker substituted with 0-1 R^c, or -(CH₂)₁₋₂-(C₃₋₄ cycloalkyl substituted with 0-1 R^c)-(CH₂)₀₋₁-; wherein said hydrocarbon linker has one to six carbon atoms and may be straight or branched, saturated or unsaturated; and said
25 hydrocarbon-heteroatom linker has one to four carbon atoms and one group selected from O and S;

R¹ is independently selected from: phenyl, indanyl, naphthyl, and a 5- to 10-membered heterocycle comprising carbon atoms and 1-4 heteroatoms selected from N, NR^b, O, and S(O)_p; wherein each ring moiety substituted with 0-4 R³ and 0-1 R⁴; and

R⁴ is independently selected from: thienyl, oxadiazolyl, and -L₃-phenyl; wherein each ring moiety is substituted with 0-2 R^a.

3. A compound according to claim 1 or claim 2, wherein the compound is of

5 Formula (II):




or a stereoisomer, a tautomer, or a pharmaceutically acceptable salt thereof, wherein:

L₂ is independently a hydrocarbon linker a hydrocarbon-heteroatom linker, or -(CH₂)₁₋₂-(cyclopropyl substituted with 0-1 R^c)-(CH₂)₀₋₁-; wherein said hydrocarbon


10 linker has one to five carbon atoms and may be straight or branched, saturated or unsaturated; and said hydrocarbon-heteroatom linker has one to three carbon atoms and one O;

L₄ is independently selected from: a bond, CH₂ and CH(C₁₋₄ alkyl);



R¹ is independently selected from:

15 moiety substituted with 0-2 R³ and selected from the group consisting of thienyl,



isoxazolyl, pyrimidinyl, indanyl, naphthyl, benzothiophenyl, and

R³, at each occurrence, is independently selected from: NO₂, halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, and C₁₋₄ haloalkylthio;

R⁴ is independently selected from: thienyl, oxadiazolyl, and -L₃-phenyl; wherein
20 each ring moiety is substituted with 0-2 R^a;


L₃ is independently selected from: a bond, O, and C(=O);

R^a, at each occurrence, is independently selected from: halogen and C₁₋₄ alkyl;
and


R^b is independently phenyl substituted with 0-2 halo.

5 4. A compound according to claims 1 to 3, wherein:

L₂ is independently selected from: CH₂OCH₂, OCH₂CH₂, CH₂CH₂CH₂,

CH₂CH(C₁₋₂ alkyl)CH₂, CH₂CH₂CH(C₁₋₂ alkyl), CH₂CH=CH, and ;

R¹-L₄- is independently selected from:



10


R³, at each occurrence, is independently selected from: halogen, C₁₋₄ alkyl,
C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, and C₁₋₄ haloalkylthio;

R⁴ is independently selected from: thienyl and -L₃-(phenyl substituted with
0-2 halo); and


15 L₃ is independently selected from: a bond, O, and C(=O).

5. A compound according to claims 1 to 4, wherein:

L₂ is independently selected from the group consisting of CH₂OCH₂, OCH₂CH₂,

CH₂CH₂CH₂, CH₂CH(Me)CH₂, CH₂CH₂CH(Me), CH₂CH=CH, and ; and

20 R¹-L₄- is independently selected from:



6. A compound according to claims 1 to 5, wherein:

L₂ is independently selected from: OCH₂CH₂, CH₂CH₂CH₂, and ;

10 and

R^1-L_4 - is independently selected from:

8. A compound according to claim 1, wherein the compound is selected from 4-(4-((1-(4-Chlorophenyl)-3-methyl-1H-pyrazol-5-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-fluoro-5-methoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-((5-chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3,5-dichlorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(5-chloro-2-isopropoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-chloro-5-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-methyl-5-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(1-(3-(trifluoromethyl)phenyl)ethoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(4,6-dichloro-2,3-dihydro-1H-inden-1-yloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-chlorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3,4-dichlorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-fluoro-4-methylbenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(4-fluoro-3-methylbenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-(2-fluorophenoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-(4-fluorophenoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-chloro-5-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(5-chloro-2-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3,5-dimethoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-chloro-5-fluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-chlorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(4-chlorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-fluoro-5-(trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-fluoro-5-methoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-fluoro-5-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-methoxy-5-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-((5-

- chlorobenzo[b]thiophen-3-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-chloro-5-methoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-(4-chlorophenoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-fluoro-5-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-fluoro-5-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-fluoro-5-isopropoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-isopropoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-(5-methyl-1,2,4-oxadiazol-3-yl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-isopropoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-benzoyloxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(4-chloro-2-methoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2,4-dichlorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-fluoro-4-methoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(4-methoxy-2-(trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-5-fluoro-2-(trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-chloro-2-fluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-fluoro-3-(trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2,3,6-trifluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(biphenyl-2-ylmethoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-(trifluoromethylthio)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-(trifluoromethylthio)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-chloro-5-nitrobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(5-chloro-2-fluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2,3-difluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-((2-bromothiophen-3-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-chloro-3-(trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-chloro-3,6-difluorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(biphenyl-3-ylmethoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(4-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2,3,6-trichlorobenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(5-bromo-2-methoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(3-(thiophen-2-yl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(5-bromo-2-

methylbenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-chloro-5-
 (trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(2-
 (trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 4-(4-(
 (trifluoromethyl)benzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, methyl 4-(4-((4-
 5 fluoro-4'-methyl-1-[1,1'-biphenyl]-2-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid,
 4-(4-((2-(methylthio)pyrimidin-4-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 2-
 Methyl-4-(4-(2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 3-Methyl-4-
 (4-(3-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)butanoic acid, 3-(4-((5-Chloro-2,2-
 dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.1]heptan-1-yl)propan-1-ol, 3-
 10 (4-((5-Chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-yl)methoxy)bicyclo[2.2.1]heptan-1-
 yl)propanoic acid, 2-((4-(2-Phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic
 acid, 2-((4-(3-fluoro-5-Phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic
 acid, 2-((4-(3-Phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid, 2-((4-
 (3-fluoro-5-(4-fluorophenoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid,
 15 2-((4-(1-(3-fluoro-5-(4-fluorophenoxy)phenyl)ethoxy)bicyclo[2.2.1]heptan-1-
 yl)methoxy)acetic acid, 2-((4-((5-chloro-2,2-dimethyl-2,3-dihydrobenzofuran-7-
 yl)methoxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid, 2-((4-((3-chloro-5-
 (trifluoromethoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yl)methoxy)acetic acid, 3-(4-(3-
 Fluoro-5-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yloxy)methoxy)propanoic acid, 3-
 20 (4-(2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yloxy)methoxy)propanoic acid, 3-(4-(3-
 fluoro-5-(4-fluorophenoxy)benzyloxy)bicyclo[2.2.1]heptan-1-yloxy))propanoic acid, 3-
 (4-(1-(3-fluoro-5-(4-fluorophenoxy)phenyl)ethoxy)bicyclo[2.2.1]heptan-heptan-1-
 yloxy)propanoic acid, 4-(4-(5-Fluoro-2-phenoxybenzyloxy)bicyclo[2.2.1]heptan-1-yl)N-
 (methylsulfonyl) butanamide, N-methyl-4-(4-((3-
 25 phenoybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)butanamide, 1-(3-Hydroxyazetidin-1-yl)-4-
 (4-((3-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)butan-1-one, 3-(4-(2-
 Phenoxyphenethoxy)bicyclo[2.2.1]heptan-1-yl)propanoic acid, 3-(4-(2-(naphthalen-1-
 yl)ethoxy)bicyclo[2.2.1]heptan-1-yl)propanoic acid, 3-(4-(3-
 phenoxyphenethoxy)bicyclo[2.2.1]heptan-1-yl)propanoic acid, 3-((1S,4R)-4-(2-fluoro-5-
 30 (trifluoromethoxy)phenethoxy)bicyclo[2.2.1]heptan-1-yl)propanoic acid, 3-(4-((3-fluoro-
 5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)propanoic acid, 2-((4-((3-Fluoro-5-
 phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methyl)cyclopropanecarboxylic acid, (E)-

4-(4-((3-Phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)but-2-enoic acid, 2-((4-((3-Phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)methyl)cyclopropanecarboxylic acid, N-(2-Amino-2-oxoethyl)-4-(4-((3-fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)butanamide, 2-(4-((3-Fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)ethanol,
5 2-(2-(4-((3-Fluoro-5-phenoxybenzyl)oxy)bicyclo[2.2.1]heptan-1-yl)ethoxy)acetic acid or a stereoisomer, a tautomer, a solvate or a pharmaceutically acceptable salt thereof.

9. A pharmaceutical composition, comprising a pharmaceutically acceptable carrier and a compound of any one of claims 1 to 8, or a stereoisomer, a tautomer, a solvate or a
10 pharmaceutically acceptable salt thereof.

10. The pharmaceutical composition according to claim 9, further comprising one or more other suitable therapeutic agents useful in the treatment of the aforementioned disorders including: anti-diabetic agents, anti-hyperglycemic agents,
15 anti-hyperinsulinemic agents, anti-retinopathic agents, anti-neuropathic agents, anti-nephropathic agents, anti-atherosclerotic agents, anti-ischemic agents, anti-hypertensive agents, anti-obesity agents, anti-dyslipidemic agents, anti-hyperlipidemic agents, anti-hypertriglyceridemic agents, anti-hypercholesterolemic agents, anti-pancreatitis agents, lipid lowering agents, anorectic agents and appetite
20 suppressants.

11. The pharmaceutical composition according to claim 10, further comprising one or more other suitable therapeutic agents selected from: a dipeptidyl peptidase-IV inhibitor, a sodium-glucose transporter-2 inhibitor and a 11b-HSD-1 inhibitor.
25

12. The compound of any one of claims 1 to 8, or a stereoisomer, a tautomer, a solvate or a pharmaceutically acceptable salt thereof for use in therapy.

13. The compound of any one of claims 1 to 8, or a stereoisomer, a tautomer, a
30 solvate or a pharmaceutically acceptable salt thereof for use in treating diabetes, hyperglycemia, impaired glucose tolerance, gestational diabetes, insulin resistance, hyperinsulinemia, retinopathy, neuropathy, nephropathy, diabetic kidney disease, acute

kidney injury, cardiorenal syndrome, delayed wound healing, atherosclerosis and its sequelae, abnormal heart function, congestive heart failure, myocardial ischemia, stroke, Metabolic Syndrome, hypertension, obesity, fatty liver disease, dislipidemia, dyslipidemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, low

5 high-density lipoprotein (HDL), high low-density lipoprotein (LDL), lipid disorders and liver diseases such as NASH (Non-Alcoholic SteatoHepatitis), NAFLD (Non-Alcoholic Fatty Liver Disease) and liver cirrhosis.

14. A compound for use according to claim 13, wherein the compound of any one of

10 claims 1 to 8 is used simultaneously, separately or sequentially with one or more additional therapeutic agents.

15