

HUMANIZED AND DE-IMMUNIZED ANTIBODIES**Field of the Invention**

The invention relates to humanized and de-immunized antibodies
5 that bind to an epitope at the N-terminus of pyroglutamated amyloid
beta (A β N3pE) peptide and to preventive and therapeutic treatment
of diseases and conditions that are related to accumulation and
deposition of amyloid peptides, such as amyloidosis, a group of
10 disorders and abnormalities associated with pyroglutamated amyloid
peptide, like Alzheimer's disease, Down's syndrome, cerebral
amyloid angiopathy and other related aspects. More specifically,
it pertains to the use of monoclonal antibodies of the invention
to bind pyroglutamated amyloid beta peptide in plasma, brain, and
15 cerebrospinal fluid to prevent accumulation or to reverse
deposition of A β N3pE within the brain and in various tissues in
the periphery, and to alleviate amyloidosis. The present invention
further pertains to diagnostic assays for the diagnosis of
amyloidosis using the antibodies of the invention.

Background Art

Amyloidosis is not a single disease entity but rather a diverse
group of progressive disease processes characterized by
extracellular tissue deposits of a waxy, starch-like protein
called amyloid, which accumulates in one or more organs or body
25 systems. As the amyloid deposits accumulate, they begin to
interfere with the normal function of the organ or body system.
There are at least 15 different types of amyloidosis. The major
forms are primary amyloidosis without known antecedent, secondary
amyloidosis following some other condition, and hereditary
30 amyloidosis.

Secondary amyloidosis occurs during chronic infection or
inflammatory disease, such as tuberculosis, a bacterial infection
called familial Mediterranean fever, bone infections
35 (osteomyelitis), rheumatoid arthritis, inflammation of the small
intestine (granulomatous ileitis), Hodgkin's disease and leprosy.

Amyloid deposits include amyloid P (pentagonal) component (AP), a glycoprotein related to normal serum amyloid P (SAP), and sulphated glycosaminoglycans (GAG), complex carbohydrates of connective tissue. Amyloid protein fibrils, which account for about 90% of the amyloid material, comprise one of several different types of proteins. These proteins are capable of folding into so-called "beta-pleated" sheet fibrils, a unique protein configuration which exhibits binding sites for Congo red resulting in the unique staining properties of the amyloid protein.

Many diseases of aging are based on or associated with amyloid-like proteins and are characterized, in part, by the buildup of extracellular deposits of amyloid or amyloid-like material that contribute to the pathogenesis, as well as the progression of the disease. These diseases include, but are not limited to, neurological disorders such as mild cognitive impairment (MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome, , Lewy body dementia, hereditary cerebral hemorrhage with amyloidosis (Dutch type); the Guam Parkinson-Dementia complex. Other diseases which are based on or associated with amyloid-like proteins are progressive supranuclear palsy, multiple sclerosis; Creutzfeldt Jacob disease, Parkinson's disease, HIV-related dementia, ALS (amyotrophic lateral sclerosis), Adult Onset Diabetes; senile cardiac amyloidosis; endocrine tumors, and others, including macular degeneration.

Although pathogenesis of these diseases may be diverse, their characteristic deposits often contain many shared molecular constituents. To a significant degree, this may be attributable to the local activation of pro-inflammatory pathways thereby leading to the concurrent deposition of activated complement components, acute phase reactants, immune modulators, and other inflammatory mediators (McGeer et al., Tohoku J Exp Med. 174(3): 269-277 (1994)).

Recently, accumulating evidence demonstrates involvement of N-terminal modified A β peptide variants in Alzheimer's disease. Aiming biopsies display a presence of A β 1-40 and A β 1-42 not only in the brain of Alzheimer's patients but also in senile plaques of unaffected individuals. However, N-terminal truncated and pyroGlu modified A β N3pE-40/A β N3pE-42 is almost exclusively engrained within plaques of Alzheimer's disease patients, making this A β variant an eligible diagnostic marker and a potential target for drug development.

At present, several commercial manufacturers offer ELISA kits which allow a detection of A β 1-40 / 1-42 and A β N3pE-40/A β N3pE-42 in the low picogram (pg) range.

The brains of Alzheimer's disease (AD) patients are morphologically characterized by the presence of neurofibrillary tangles and by deposits of A β peptides in neocortical brain structures (Selkoe, D.J. & Schenk, D. Alzheimer's disease: molecular understanding predicts amyloid-based therapeutics. Annu. Rev. Pharmacol. Toxicol. 43, 545-584 (2003)). A β peptides are liberated from the amyloid precursor protein (APP) after sequential cleavage by β - and γ -secretase. The γ -secretase cleavage results in the generation of A β 1-40 and A β 1-42 peptides, which differ in their C-termini and exhibit different potencies of aggregation, fibril formation and neurotoxicity (Shin, R.W. et al. Amyloid beta-protein (Abeta) 1-40 but not Abeta 1-42 contributes to the experimental formation of Alzheimer disease amyloid fibrils in rat brain. J. Neurosci. 17, 8187-8193 (1997); Iwatsubo, T. et al. Visualization of Abeta 42(43) and Abeta 40 in senile plaques with end-specific Abeta monoclonals: evidence that an initially deposited species is Abeta 42(43). Neuron 13, 45-53 (1994); Iwatsubo, T., Mann, D.M., Odaka, A., Suzuki, N. & Ihara, Y. Amyloid beta protein (Abeta) deposition: Abeta 42(43) precedes Abeta 40 in Down syndrome. Ann. Neurol. 37, 294-299 (1995); Hardy, J.A. & Higgins, G.A. Alzheimer's disease: the amyloid cascade hypothesis. Science 256, 184-185 (1992); Roßner, S., Ueberham, U., Schliebs, R., Perez-Polo, J.R. & Bigl, V. The regulation of amyloid precursor

protein metabolism by cholinergic mechanisms and neurotrophin receptor signaling. *Prog. Neurobiol.* 56, 541-569 (1998)).

The majority of A β peptides deposited in diffuse plaques are N-terminal truncated or modified. Studies of Piccini and Saido have shown that the core structure of senile plaques and vascular deposits consist of 50 % pyroglutamate (pyroGlu) modified peptides (Piccini *et al.*, *J Biol Chem.* 2005 Oct 7;280(40):34186-92; Saido *et al.*, *Neuron.* 1995 Feb; 14(2): 457-66). PyroGlu modified peptides are more strongly cytotoxic than other A β species and stable against aminopeptidases (Russo *et al.*, *J Neurochem.* 2002 Sep;82(6):1480-9). Thus, pyroGlu A β species have a longer half-life whereby the accumulation of these species and the formation of neurotoxic oligomers as well as aggregates are beneficial (Saido, *Neurobiol Aging.* 1998 Jan-Feb;19(1 Suppl):S69-75). Due to the cyclization of glutamate to pyroGlu, charged amino acids will be lost which strongly reduces the solubility of the peptide and causes an increased aggregation tendency. *In vitro* studies have shown that the initial oligomerisation of e.g. A β 3(pE) is much faster compared to non-modified peptides (Schilling *et al.*, *Biochemistry.* 2006 Oct 17;45(41):12393-9). The A β N3pE-42 peptides coexist with A β 1-40/1-42 peptides (Saido, T.C. *et al.* Dominant and differential deposition of distinct beta-amyloid peptide species, Abeta N3pE, in senile plaques. *Neuron* 14, 457-466 (1995) ; Saido, T.C., Yamao, H., Iwatsubo, T. & Kawashima, S. Amino- and carboxyl-terminal heterogeneity of beta-amyloid peptides deposited in human brain. *Neurosci. Lett.* 215, 173-176 (1996)), and, based on a number of observations, could play a prominent role in the pathogenesis of AD. For example, a particular neurotoxicity of A β N3pE-42 peptides has been outlined (Russo, C. *et al.* Pyroglutamate-modified amyloid beta-peptides--AbetaN3(pE)--strongly affect cultured neuron and astrocyte survival. *J. Neurochem.* 82, 1480-1489 (2002) and the pE-modification of N-truncated A β peptides confers resistance to degradation by most aminopeptidases as well as A β -degrading endopeptidases (Russo, C. *et al.* Pyroglutamate-modified amyloid beta-peptides--AbetaN3(pE)--strongly affect cultured neuron and astrocyte survival. *J. Neurochem.* 82, 1480-

1489 (2002); Saido, T.C. Alzheimer's disease as proteolytic disorders: anabolism and catabolism of beta-amyloid. *Neurobiol. Aging* 19, S69-S75 (1998)). The cyclization of glutamic acid into pE leads to a loss of N-terminal charge resulting in accelerated aggregation of A β N3pE compared to the unmodified A β peptides (He, W. & Barrow, C.J. The Abeta 3-pyroglutamyl and 11-pyroglutamyl peptides found in senile plaque have greater beta-sheet forming and aggregation propensities in vitro than full-length A beta. *Biochemistry* 38, 10871-10877 (1999); Schilling, S. et al. On the seeding and oligomerization of pGlu-amyloid peptides (in vitro). *Biochemistry* 45, 12393-12399 (2006)). Thus, reduction of A β N3pE-42 formation should destabilize the peptides by making them more accessible to degradation and would, in turn, prevent the formation of higher molecular weight A β aggregates and enhance neuronal survival.

However, for a long time it was not known how the pE-modification of A β peptides occurs. Recently, it was discovered that glutaminyl cyclase (QC) is capable to catalyze A β N3pE-42 formation under mildly acidic conditions and that specific QC inhibitors prevent A β N3pE-42 generation *in vitro* (Schilling, S., Hoffmann, T., Manhart, S., Hoffmann, M. & Demuth, H.-U. Glutaminyl cyclases unfold glutamyl cyclase activity under mild acid conditions. *FEBS Lett.* 563, 191-196 (2004) ; Cynis, H. et al. Inhibition of glutaminyl cyclase alters pyroglutamate formation in mammalian cells. *Biochim. Biophys. Acta* 1764, 1618-1625 (2006)).

All facts suggest that pyroGlu A β is a kind of germ for the initialization of fibril formation. In a further study (Piccini et al., 2005, *supra*) volunteers with plaque depositions but without AD specific pathology could be distinguished from AD patients due to the characteristic amount of A β -species. Thereby the amount of N-terminal truncated, pyroGlu modified peptides was significant higher in the brain of AD patients.

The posttranslational formation of pyroGlu at position 3 or 11 of A β -peptide implies cyclization of an N-terminal glutamate residue.

Glutaminyl cyclase (QC) plays an important role in the generation of pyroGlu peptides. QC is wide-spread in the plant- and animal kingdom and *inter alia*, is involved in the maturation of peptide hormones. Both the cyclisation of glutamine by release of ammonia and of glutamate by release of water to pyroGlu is performed by QC. In contrast to the glutamine cyclization the glutamate cyclisation occurs not spontaneously. QC catalyses the efficient (unwanted) side reaction from glutamate to pyroGlu. The generated pyroGlu residue protects the protein against proteolytic degradation. There are several references which shows that QC plays an important role in the generation of pyroGlu A β :

1. In several studies it was shown that QC catalyses the formation of pyroGlu residues from glutamate at N-terminus of A β (Cynis *et al.*, Biochim Biophys Acta. 2006 Oct;1764(10):1618-25, Schilling *et al.*, FEBS Lett. 2004 Apr 9;563(1-3):191-6);

2. Both A β peptides and QC are expressed in large quantities in hippocampus and cortex. These brain areas are at particular risk in AD (Pohl *et al.*, Proc Natl Acad Sci U S A. 1991 Nov 15;88(22):10059-63, Selkoe, Physiol Rev. 2001 Apr;81(2):741-66);

3. The APP is cleaved by β -secretase during the transport to the plasma membrane whereby the N-terminus of A β with the free glutamate residue can be produced (Greenfield *et al.*, Proc Natl Acad Sci U S A. 1999 Jan 19;96(2):742-7). In the secretory vesicles a co-localisation of processed APP and the QC was determined. So in the mild acid milieu of the vesicles an accelerated modification of glutamate residue to pyroglutamate can occur.

4. Also other neurodegenerative diseases (familial Danish (FDD) or British dementia (FBD)) are related with N-terminal pyroGlu modified peptides e.g. Bri2, but in contrast they are not related to A β in terms of their primary structure (Vidal R. *et al.*, 1999 Proc. Natl. Acad. Sci. U.S.A. 97, 4920-4925).

Possibly the QC-catalysed formation of pyroGlu A β is involved in the development and progression of neurodegenerative diseases. The formation of N-terminal modified amyloid peptides certainly represents a fundamental factor in the process of A β aggregation and could be the onset of disease. The suppression of the pyroGlu

A β formation by inhibition of QC, might represent a therapeutic approach. QC inhibitors would be able to prevent the formation of pyroGlu A β , reduce the concentration of pyroglutamate A β in the brain and so delay the oligomerisation of A β -peptides. Schilling
5 *et al.* show, that QC expression was up regulated in the cortex of AD patients and correlated with the appearance of pyroGlu-modified A β -peptide. Oral application of a QC inhibitor resulted in reduced pyroglutamate modified A β pE(3-42) level in two different transgenic mouse models of AD and in a new *Drosophila* model
10 (Schilling *et al.*, 2008 *Biol. Chem.* (389), 983-991).

Lewy body dementia (LBD) is a neurodegenerative disorder that can occur in persons older than 65 years of age, and typically causes symptoms of cognitive (thinking) impairment and abnormal
15 behavioral changes. Symptoms can include cognitive impairment, neurological signs, sleep disorder, and autonomic failure. Cognitive impairment is the presenting feature of LBD in most cases. Patients have recurrent episodes of confusion that progressively worsen. The fluctuation in cognitive ability is
20 often associated with shifting degrees of attention and alertness. Cognitive impairment and fluctuations of thinking may vary over minutes, hours, or days. Lewy bodies are formed from phosphorylated and nonphosphorylated neurofilament proteins; they contain the synaptic protein alpha-synuclein as well as ubiquitin, which is
25 involved in the elimination of damaged or abnormal proteins. In addition to Lewy Bodies, Lewy neurites, which are inclusion bodies in the cell processes of the nerve cells, may also be present. Amyloid plaques may form in the brains of patients afflicted with DLB, however they tend to be fewer in number than seen in patients
30 with Alzheimer's disease. Neurofibrillary tangles, the other micropathological hallmark of AD, are not a main characteristic of LBD but are frequently present in addition to amyloid plaques.

Amyotrophic lateral sclerosis (ALS) is characterized by
35 degeneration of upper and lower motor neurons. In some ALS patients, dementia or aphasia may be present (ALS-D). The dementia is most commonly a frontotemporal dementia (FTD), and many of these

cases have ubiquitin- positive, tau-negative inclusions in neurons of the dentate gyrus and superficial layers of the frontal and temporal lobes.

Inclusion-body myositis (IBM) is a crippling disease usually found
5 in people over age 50, in which muscle fibers develop inflammation and begin to atrophy – but in which the brain is spared and patients retain their full intellect. Two enzymes involved in the production of amyloid- β protein were found to be increased inside the muscle cells of patients with this most common, progressive muscle disease
10 of older people, in which amyloid- β is also increased.

Another disease that is based on or associated with the accumulation and deposit of amyloid-like protein is macular degeneration. Macular degeneration is a common eye disease that
15 causes deterioration of the macula, which is the central area of the retina (the paper-thin tissue at the back of the eye where light-sensitive cells send visual signals to the brain). Sharp, clear, "straight ahead" vision is processed by the macula. Damage to the macula results in the development of blind spots and blurred
20 or distorted vision. Age-related macular degeneration (AMD) is a major cause of visual impairment in the United States and for people over age 65 it is the leading cause of legal blindness among Caucasians. Approximately 1.8 million Americans of age 40 and older have advanced AMD, and another 7.3 million people with intermediate
25 AMD are at substantial risk for vision loss. The government estimates that by 2020 there will be 2.9 million people with advanced AMD. Victims of AMD are often surprised and frustrated to find out how little is known about the causes and treatment of this blinding condition.

30

There are two forms of macular degeneration: dry macular degeneration and wet macular degeneration. The dry form, in which the cells of the macula slowly begin to break down, is diagnosed in 85 percent of macular degeneration cases. Both eyes are usually
35 affected by dry AMD, although one eye can lose vision while the other eye remains unaffected. Drusen, which are yellow deposits under the retina, are common early signs of dry AMD. The risk of

developing advanced dry AMD or wet AMD increases as the number or size of the drusen increases. It is possible for dry AMD to advance and cause loss of vision without turning into the wet form of the disease; however, it is also possible for early-stage dry AMD to suddenly change into the wet form.

The wet form, although it only accounts for 15 percent of the cases, results in 90 percent of the blindness, and is considered advanced AMD (there is no early or intermediate stage of wet AMD). Wet AMD is always preceded by the dry form of the disease. As the dry form worsens, some people begin to have abnormal blood vessels growing behind the macula. These vessels are very fragile and will leak fluid and blood (hence 'wet' macular degeneration), causing rapid damage to the macula.

The dry form of AMD will initially often cause slightly blurred vision. The center of vision in particular may then become blurred and this region grows larger as the disease progresses. No symptoms may be noticed if only one eye is affected. In wet AMD, straight lines may appear wavy and central vision loss can occur rapidly.

Diagnosis of macular degeneration typically involves a dilated eye exam, visual acuity test, and a viewing of the back of the eye using a procedure called fundoscopy to help diagnose AMD, and – if wet AMD is suspected – fluorescein angiography may also be performed. If dry AMD reaches the advanced stages, there is no current treatment to prevent vision loss. However, a specific high dose formula of antioxidants and zinc may delay or prevent intermediate AMD from progressing to the advanced stage. Macugen® (pegaptanib sodium injection), laser photocoagulation and photodynamic therapy can control the abnormal blood vessel growth and bleeding in the macula, which is helpful for some people who have wet AMD; however, vision that is already lost will not be restored by these techniques. If vision is already lost, low vision aids exist that can help improve the quality of life.

One of the earliest signs of age-related macular degeneration (AMD) is the accumulation of extracellular deposits known as drusen between the basal lamina of the retinal pigmented epithelium (RPE) and Bruch's membrane (BM). Recent studies conducted by Anderson et al. have confirmed that drusen contain amyloid beta. (Experimental Eye Research 78 (2004) 243 - 256).

Pyroglutamated A β peptides have been shown to play a key role in accumulation of A β peptides and in plaque formation in Alzheimer's diseases. Due to their hydrophobic potential it has been shown that these peptides promote aggregation and plaque formation. It has further been shown in a transgenic mouse model expressing A β N3pE-42 in neurons that this peptide is neurotoxic *in vivo* and leads to loss of neurons (Wirhth et al. (2009) Acta Neuropatho/118, 487-496).

Antibodies with specificities against the N-terminal pyroglutamate of A β peptides are believed to be advantageous because of their specificity towards only the pathogenic species of A β , which carry a pyroglutamate at the N-terminus, but not detecting APP or other A β species w/o the N-terminal pyroglutamate. It is thus believed that the risk of potential side effects, such as uncontrollable cerebral inflammation, will be reduced by use of the antibodies of the invention compared to antibodies directed to other A β species that the pyroglutamated variants.

Antibodies targeting A β N3pE peptides are known (Acero et al (2009) J Neuroimmunol 213, 39-46; Saido et al. (1996) Neuron 14, 457-466; US 7,122,374 and WO 2012/136552).

However, there is a need for humanized and de-immunized antibodies with specificity for A β N3pE peptides that can be used in human treatment and that positively affect amyloidosis, in particular cognition in diseases and conditions where A β N3pE may be involved, such as clinical or pre-clinical Alzheimer's disease, Down's syndrome, and clinical or pre-clinical cerebral amyloid angiopathy.

The parental antibody of the antibody of the present invention is the clone#6 variant disclosed in WO 2017/009459, which has the light chain variable region with the amino acid sequence:

5

DVVMTQSPLSLPVTLGQPASISCKSSQSLLYSDGKTYLNWFQQRPGQSPRRLTYLVSKLDSGVP
DRFSGSGSGTDFTLKISRVEAEDVGVYYCVQGTHFPFTFGGGTKVEIK (SEQ ID NO: 1),
which is disclosed as SEQ ID NO: 14 in WO 2017/009459;

10 and

which has the heavy chain variable region with the amino acid sequence:

15 QVQLVQSGAEVKKSGASVKVSCASGYSFTGHTMNWVRQAPGQGLEWMGLINPSNGVTRYNQKF
QGRVTITRDITSTTTVHMEITSLTSEDATYYCTREAKREWDETYWGQGTILVTVSS (SEQ ID
NO: 49); which is disclosed as SEQ ID NO: 27 in WO 2017/009459.

However, with this clone#6 variant as disclosed in WO 2017/009459,
20 it was not possible to establish a CMC manufacturing. In particular, and despite several attempts, no stable cell clones could be established in CHO-DG44 cells, only a weak transient expression (yielding insufficient antibody amounts) of this clone#6 variant was observed, and no stable expression, as a
25 prerequisite for CMC manufacturing, could be established.

Summary of the invention

It was therefore the purpose of the invention to provide humanized and de-immunized antibodies with improved properties to overcome
30 the disadvantages of the prior art antibodies.

In general, the invention provides novel methods and compositions comprising highly specific and highly effective antibodies, including chimeric antibodies and fragments thereof, including
35 partially or fully humanized antibodies and fragments thereof, having the ability to specifically recognize and bind to specific epitopes from a range of β -amyloid antigens, in particular A β N3pE

peptides, which may be presented to the antibody in a monomeric, dimeric, trimeric, etc, or a polymeric form, in form of an aggregate, fibers, filaments or in the condensed form of a plaque.

5 The purpose of the invention is in particular solved by an antibody or a functional variant thereof, wherein the variable part of the light chain of said antibody comprises, consists essentially of or consists of an amino acid sequence of:

10 DVVMTQSPLSLPVTLGQPASISCKSSQSLLYSDGKTYLNWFQQRPQGSPRRLTYLVSKLDSGVP
DRFSGSGSGTDFTLKISRVEAEDVGVYYCVQGTHFPFTFGGGTKVEIK (SEQ ID NO: 1),

and

15 wherein the variable part of heavy chain of said antibody comprises, consists essentially of or consists of an amino acid sequence of:

QVQLVQSGAEVVKPGASVKVSCKASGYSFTGHTMNWVRQAPGQGLEWMGLINPSDGVTRYNQKF
20 QGRVTITRDTSTTTVHMEITSLTSEDATYYCTREAKREWDETYWGQGTILVTVSS (SEQ ID
NO: 2).

The variable part of the heavy chain (SEQ ID NO: 2) contains three mutations at positions K12V, S14P and N55D compared to the parental
25 antibody disclosed in WO 2017/009459.

Surprisingly, it was found that the introduction of these three point mutations led to a humanized and de-immunized antibody, which is capable for CMC manufacturing with high yields. It was possible
30 to establish antibody producing cell lines in CHO-DG44 cells with the antibody of the invention comprising these three point mutations. It was further possible to establish an improved transient expression in CHO-DG44, resulting in higher yields of the antibody of the invention comprising K12V, S14P and N55D
35 mutations compared to the parental antibody, while maintaining the favorable binding characteristics of the antibody. Finally, a stable expression could be established with the antibody of the

invention comprising K12V, S14P and N55D mutations with high expression levels, enabling CMC production.

5 The invention provides humanized and de-immunized antibodies, or fragments thereof, that positively affect diseases and conditions of amyloidosis, where A β N3pE may be involved.

10 In another embodiment, the invention provides antibodies and fragments thereof that bind to A β N3pE peptides in the circulation and tissue, in particular in the brain. The antibodies of the invention are capable of binding free A β N3pE peptide molecules or even bound forms of A β N3pE peptides.

15 Thus, the present invention further provides antibodies that alter clearance of soluble and bound forms of A β N3pE peptides in the central nervous system, such as the brain, and the circulation, such as plasma.

20 In a further embodiment, the invention provides antibodies and fragments thereof, wherein the antibodies specifically bind to the pyroglutamate carrying N-terminus of A β N3pE.

25 In a further embodiment, the invention provides antibodies and fragments thereof, wherein the antibodies show an increased selectivity towards oligomers and/or fibrils of A β peptides. The antibodies of the present invention show a manifold, such as 10 times, 25 times, 50 times, 100 times, 150 times, 200 times, 250 times or more than 250 times lower binding constant (K_D value) for binding to oligomers and/or fibrils of A β (1-42) than comparable
30 monoclonal antibodies known in the prior art, in particular compared to the parental antibody disclosed in WO 2017/009459. Accordingly, the antibodies of the present invention, which were established to selectively bind to A β N3pE peptides, are more specific for A β N3pE peptides and show a decreased cross-reactivity
35 against A β peptides other than A β N3pE.

In yet a further embodiment, the present invention also relates to the host cells transformed with the vectors or incorporating the nucleic acid molecules that express the antibodies of the invention or fragments thereof.

5

Moreover, the present invention provides pharmaceutical compositions comprising the antibodies of the invention and fragments thereof.

10 The invention further relates to the use of the antibodies of the invention and fragments thereof are useful for binding to and clearing or removing of A β N3pE in humans and thereby for diagnosing, preventing and treating diseases and conditions characterized by amyloidosis or A β N3pE toxicity.

15

In a particular embodiment, the antibodies of the invention, which are capable of binding to and clearing or removing of A β N3pE peptides in biological fluids and tissues, are useful for the prevention and/or treatment of conditions associated with the
20 formation of A β N3pE-containing plaques, such as diffuse, neuritic, and cerebrovascular plaques in the brain.

The administration of the antibodies of the invention, including immunologically reactive fragments thereof, may lead to the
25 clearance or removal of A β N3pE from the aforementioned plaques or other biological complexes. Thus, the antibody of the invention will readily be transport in the circulation, other body fluids and to sites where the aforementioned plaques and/or other biological complexes are formed or elsewhere where A β N3pE exhibits
30 damaging effects.

In addition, removal of A β N3pE from plaques or other biological complexes by the antibodies of the invention may lead to the solubilization of insoluble forms of plaques and thus lead to the
35 removal of complete plaques from the affected tissue, such as brain tissue. This, in turn, may lead to improvement of cognition in patients diagnosed with a neurodegenerative disease, such as mild

cognitive impairment (MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD) or others, neurodegeneration in
5 Down Syndrome, Lewy body dementia, hereditary cerebral hemorrhage with amyloidosis (Dutch type); the Guam Parkinson-Dementia complex. In particular, the present invention provides an antibody of the present invention for use in treatment of a condition selected from prodromal AD, mild AD, moderate AD and severe AD. In
10 another embodiment, the present invention provides an antibody of the present invention for use in slowing cognitive decline in a patient diagnosed with a condition selected from clinical or pre-clinical Alzheimer's disease, Down's syndrome, and clinical or pre-clinical cerebral amyloid angiopathy.

15 The binding of the antibodies of the invention to A β N3pE in the circulation or other body fluids may further result to the removal of the circulating or soluble forms of A β N3pE. As discussed above, A β N3pE exhibits a high hydrophobicity and has a high affinity to
20 other, e.g. nonpyroglutamated A β peptides, which results in the formation of oligomeric and supermolecular structures, such as amyloid plaques. It has been shown that in particular these oligomeric structures are highly neurotoxic. The formation of oligomeric structures leads to cell damage and death of neuronal
25 cells. Thus, the removal of circulating or soluble forms of A β N3pE or even of oligomers comprising A β N3pE leads to the prevention of cell damage and/or neurotoxicity. Thus, the invention also provides methods of preventing of neurodegenerative disease, such as mild cognitive impairment (MCI), Alzheimer's
30 disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD) or others, neurodegeneration in Down Syndrome, Lewy body dementia, hereditary cerebral hemorrhage with amyloidosis (Dutch type), the Guam
35 Parkinson-Dementia complex. In particular, the present invention provides methods of treatment of a condition selected from prodromal AD, mild AD, moderate AD and severe AD. In another

embodiment, the present invention provides a method of slowing cognitive decline in a patient diagnosed with a condition selected from clinical or pre-clinical Alzheimer's disease, Down's syndrome, and clinical or pre-clinical cerebral amyloid angiopathy.

The invention further provides methods of preventing and /or treating of other diseases which are based on or associated with amyloid-like proteins, in particular A β N3pE, such as progressive supranuclear palsy, multiple sclerosis, Creutzfeldt Jacob disease, Parkinson's disease, HIV-related dementia, ALS (amyotrophic lateral sclerosis), dementia related to Adult Onset Diabetes; senile cardiac amyloidosis, and others, including macular degeneration.

The invention further provides a highly sensitive and concomitantly robust detection technique that allows quantitative determination of A β variants, in particular A β N3pE, in biological samples, e.g. liquor or serum samples, preferably serum samples, or tissue samples. This is a tremendous challenge, taking the low abundance of these A β N3pE peptides in blood into account. Having such a detection technique available is, however, a prerequisite for studying efficacy of small molecule inhibitors in drug screening and drug development programs.

The antibodies enabled by the teaching of the present invention are particularly useful for diagnosis of amyloidosis, a group of diseases and disorders associated with amyloid plaque formation including secondary amyloidosis and age-related amyloidosis including, but not limited to, neurological disorders such as Alzheimer's Disease (AD), Lewy body dementia, Down's syndrome, hereditary cerebral hemorrhage with amyloidosis (Dutch type), the Guam Parkinson- Dementia complex, as well as other diseases which are based on or associated with amyloid-like proteins such as progressive supranuclear palsy, multiple sclerosis; Creutzfeldt Jacob disease, hereditary cerebral hemorrhage with amyloidosis Dutch type, Parkinson's disease, HIV-related dementia, ALS (amyotrophic lateral sclerosis), dementia related to Adult Onset

Diabetes, senile cardiac amyloidosis, and others, including macular degeneration, to name just a few.

5 Description of the Figures

Figure 1 shows the influence of the point mutations in the heavy chain on transient expression (A) and target binding (B) of individual variants of the antibody of the invention. All antibodies tested contain the variable part of the light chain of SEQ ID NO. 1 and the K324A mutation in the heavy chain.

Ab 1: contains the two mutations K12V and S14P in the variable part of the heavy chain compared to the parental sequence SEQ ID NO: 49;

Ab 2: contains the mutation N55D in the variable part of the heavy chain compared to the parental sequence SEQ ID NO: 49;

Ab 3: represents the parental sequence SEQ ID NO: 49 of the variable part of the heavy chain w/o any mutations.

Figure 2 shows the influence of the point mutations in the heavy chain on transient expression (A) and target binding (B) of individual variants of the antibody of the invention. All antibodies tested contain the variable part of the light chain of SEQ ID NO. 1 and the K324A mutation in the heavy chain.

Ab 1: contains the two mutations K12V and S14P in the variable part of the heavy chain compared to the parental sequence SEQ ID NO: 49;

Ab 2: contains the mutation N55D in the variable part of the heavy chain compared to the parental sequence SEQ ID NO: 49;

Ab 3: contains the three mutations K12V, S14P and N55D in the variable part of the heavy chain compared to the parental sequence SEQ ID NO: 49

5

Figure 3 shows CHO-DG44 clones expressing the antibody with the variable part of the light chain of SEQ ID NO: 1 and the variable part of the heavy chain of SEQ ID NO: 2, seven, ten and 13 days after seeding wells of a 24-well plate from colonies picked from a 96-well plate.

10

Figure 4 shows CHO-DG44 expression titers after gene amplification through MTX exposure (Day 7).

15 Detailed description of the Invention

Definitions

The term "antibody" is used in the broadest sense and specifically covers intact monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g. bispecific antibodies) formed from at least two intact antibodies, and antibody fragments so long as they exhibit the desired biological activity. The antibody may be an IgM, IgG (e.g. IgG1, IgG2, IgG3 or IgG4), IgD, IgA or IgE, for example. Preferably however, the antibody is not an IgM antibody.

25

"Antibody fragments" comprise a portion of an intact antibody, generally the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

30

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e. the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal

35

antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to "polyclonal antibody" preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies can frequently be advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Köhler *et al.*, Nature, 256:495 (1975), or may be made by generally well known recombinant DNA methods. The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, Nature, 352:624-628 (1991) and Marks *et al.*, J. Mol. Biol., 222:581-597 (1991), for example.

The monoclonal antibodies herein specifically include chimeric antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity.

"Humanized" forms of non-human (e.g., murine) antibodies are immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain a minimal sequence derived from a non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from

a complementarity-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv
5 framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences.

10

These modifications are made to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR
15 regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further
20 details, see Jones *et al.*, Nature, 321:522-525 (1986), Reichmann *et al.*, Nature. 332:323-329 (1988): and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992).

In addition to "humanization", "de-immunization" includes other
25 changes such as removing T cell epitopes.

The term "therapeutically effective amount" as used herein and in the appended claims means that the amount of antibody administered is of sufficient quantity to achieve the intended purpose such as,
30 in this case, at least the removal of circulating or soluble forms of pyroglutamated amyloid beta (A β N3pE) peptide and variants thereof, but preferably the clearance or removal of A β N3pE peptide from plaques or other biological complexes. Or more preferably the reduction of the plaque load and/or the removal of complete plaques
35 from the affected tissue, such as brain tissue.

- "Single-chain Fv" or "sFv" antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Generally, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv see Plückthun in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).
- The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_D) in the same polypeptide chain ($V_H - V_D$). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in Hollinger et al., Proc. Natl. Acad. Sol. USA, 90:6444-6448 (1993).
- An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

As used herein, the expressions "cell", "cell line," and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and culture derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Mutant progeny that have the same function or biological activity as screened for in the originally transformed cell are included. Where distinct designations are intended, this will be clear from the context.

The terms "polypeptide", "peptide", and "protein", as used herein, are interchangeable and are defined to mean a biomolecule composed of amino acids linked by a peptide bond.

If peptide or amino acid sequences are mentioned herein, each amino acid residue is represented by a one-letter or a three-letter designation, corresponding to the trivial name of the amino acid, in accordance with the following conventional list:

<u>Amino Acid</u>	<u>One-Letter Symbol</u>	<u>Three-Letter Symbol</u>
Alanine	A	Ala
Arginine	R	Arg
25 Asparagine	N	Asn
Aspartic acid	D	Asp
Cysteine	C	Cys
Glutamine	Q	Gln
Glutamic acid	E	Glu
30 Glycine	G	Gly
Histidine	H	His
Isoleucine	I	Ile
Leucine	L	Leu
Lysine	K	Lys
35 Methionine	M	Met
Phenylalanine	F	Phe
Proline	P	Pro

	Serine	S	Ser
	Threonine	T	Thr
	Tryptophan	W	Trp
	Tyrosine	Y	Tyr
5	Valine	V	Val

The terms "a", "an" and "the" as used herein are defined to mean "one or more" and include the plural unless the context is inappropriate.

10

The language "diseases and disorders which are caused by or associated with amyloid or amyloid-like proteins" includes, but is not limited to, diseases and disorders caused by the presence or activity of amyloid-like proteins in monomeric, fibril, or polymeric state, or any combination of the three. Such diseases and disorders include, but are not limited to, amyloidosis, endocrine tumors, and macular degeneration.

The term "amyloidosis" refers to a group of diseases and disorders associated with amyloid plaque formation including, but not limited to, secondary amyloidosis and age-related amyloidosis such as diseases including, but not limited to, neurological disorders such as Alzheimer's Disease (AD), including diseases or conditions characterized by a loss of cognitive memory capacity such as, for example, mild cognitive impairment (MCI), sporadic Alzheimer's disease, Lewy body dementia, Down's syndrome, hereditary cerebral hemorrhage with amyloidosis (Dutch type); the Guam Parkinson-Dementia complex, familial forms of Alzheimer's disease like Familial British Dementia (FBD) and Familial Danish Dementia (FDD); as well as other diseases which are based on or associated with amyloid-like proteins such as progressive supranuclear palsy, multiple sclerosis; Creutzfeldt Jacob disease, Parkinson's disease, HIV-related dementia, ALS (amyotrophic lateral sclerosis), inclusion-body myositis (IBM), Adult Onset Diabetes, and senile cardiac amyloidosis; and various eye diseases including macular degeneration, drusen-related optic neuropathy, and cataract due to beta-amyloid deposition.

"Amyloid β , A β or β -amyloid" is an art recognized term and refers to amyloid β proteins and peptides, amyloid β precursor protein (APP), as well as modifications, fragments and any functional equivalents thereof. In particular, by amyloid β as used herein is meant any fragment produced by proteolytic cleavage of APP but especially those fragments which are involved in or associated with the amyloid pathologies including, but not limited to, A β_{1-38} , A β_{1-40} , A β_{1-42} . The amino acid sequences of these A β peptides are as follows:

A β 1-42 (SEQ ID NO. 37):

Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val-Ile-Ala

A β 1-40 (SEQ ID NO. 38):

Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val

A β 1-38 (SEQ ID NO. 39):

Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly

"pGlu-A β " or "A β N3pE" refers to N-terminally truncated forms of A β , that start at the glutamic acid residue at position 3 in the amino acid sequence of A β , and wherein said glutamic acid residue is cyclized to form a pyroglutamic acid residue. In particular, by pGlu-A β or A β N3pE as used herein are meant those fragments which are involved in or associated with the amyloid pathologies including, but not limited to, pGlu-A β_{3-38} , pGlu-A β_{3-40} , p-Glu-A β_{3-42} .

The sequences of the N-terminally truncated forms of A β , A β_{3-38} , A β_{3-40} , A β_{3-42} are as follows:

Aβ 3-42 (SEQ ID NO. 40) :

Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-
 Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-
 5 Met-Val-Gly-Gly-Val-Val-Ile-Ala

Aβ 3-40 (SEQ ID NO. 41) :

Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-
 Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-
 10 Met-Val-Gly-Gly-Val-Val

Aβ 3-38 (SEQ ID NO. 42) :

Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-
 Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-
 15 Met-Val-Gly-Gly

The present invention pertains to antibodies specific for human Aβ peptides that are N-terminally truncated by cleaving off or losing amino acids no. 1 and 2 of the N-terminus and in which the so
 20 uncovered N-terminal amino acid no. 3 is modified by pyroglutamate formation and which thus bear a pyroglutamate residue at position 3 of the N-terminus (further referred to as Aβ N3pE peptides or N3pE-Aβ peptides or pyroglutamated Aβ peptides).

25 In a first aspect, the present invention pertains to an antibody, wherein the variable part of the light chain of said antibody comprises, consists essentially of or consists of an amino acid sequence of:

30 DVVMTQSPLSLPVTLGQPASISCKSSQSLLYSDGKTYLNWFQQRPQGQSPRRLTYLVSKLDSGVP
 DRFSGSGSGTDFTLKISRVEAEDVGVYYCVQGTHTFPFTFGGGTKVEIK (SEQ ID NO: 1),

and

35 wherein the variable part of heavy chain of said antibody comprises, consists essentially of or consists of an amino acid sequence of:

QVQLVQSGAEVVKPGASVKVSCASGYSFTGHTMNWVRQAPGGLEWMGLINPSDGVTRYNQKF
QGRVTITRDTSTTTVHMELTSLTSEDATYYCTREAKREWDETYWGQGTLLTVSS (SEQ ID
NO: 2)

5

In a preferred embodiment of the present invention, the antibody
having the variable part of the light chain of said antibody, which
comprises, consists essentially of or consists of the amino acid
sequence of SEQ ID NO: 1, comprises the following CDR regions in
10 the light chain:

V_L CDR1: SSQSLLYSDGKTYLN (SEQ ID NO: 3);

V_L CDR2: LVSKLDS (SEQ ID NO: 4); and

V_L CDR3: VQGTHFP (SEQ ID NO: 5).

15

In a further preferred embodiment of the present invention, the
antibody having the variable part of the heavy chain of said
antibody, which comprises, consists essentially of or consists of
20 the amino acid sequence of SEQ ID NO: 2, comprises the following
CDR regions in the heavy chain:

V_H CDR1: GYSFTGHTMN (SEQ ID NO: 6);

V_H CDR2: LINPSDGVTRYNQKFQG (SEQ ID NO: 7); and

25 V_H CDR3: EAKREWDETY (SEQ ID NO: 8).

The invention further provides the light chains and the heavy
chains of the antibody of the invention.

30 In a preferred embodiment, the antibody of the invention has a
light chain, wherein the light chain comprises, consists
essentially of or consists of the amino acid sequence of:

DVVMTQSPLSLPVTLGQPASISCKSSQSLLYSDGKTYLNWFQQRPGQSPRRLTYLVSKLDSGVP
35 DRFSGSGSGTDFTLKISRVEAEDVGVYYCVQGTHFPFTFGGGTKVEIKRTVAAPSVFIFPPSDE
QLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYLSSTLTLSKADYE
KHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 17).

In a further preferred embodiment, the antibody of the invention has a heavy chain, wherein the heavy chain comprises, consists essentially of or consists of the amino acid sequence of:

5

QVQLVQSGAEVVKPGASVKVSCASGYSFTGHTMNWVRQAPGQGLEWMGLINPSDGVTRYNQKF
QGRVTITRDTSTTTVHMELTSLTSEDATATYYCTREAKREWDETYWGQGTLLTVSSASTKGPSVF
PLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS
SSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTLMIS
10 RTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE
YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE
SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQKSLSLSPG
K (SEQ ID NO: 19).

15 Clq and two serine proteases, C1r and C1s, form the complex C1, the first component of the complement dependent cytotoxicity (CDC) pathway. Clq is a hexavalent molecule with a molecular weight of approximately 460,000 and a structure likened to a bouquet of tulips in which six collagenous "stalks" are connected to six
20 globular head regions (Burton and Woof, Advances in Immunol 51:1-84; 1992). Binding of IgG1 molecules to Clq initiates complement activation and subsequently leads to complement-mediated cell lysis. The antibodies of the present invention shall be used in treatment of inflammatory diseases and conditions, i.e. the
25 antibodies of the present invention shall have anti-inflammatory properties.

Effector functions of the antibodies of the invention can also be mediated by the interaction of the Fc region of an antibody with
30 Fc receptors (FcRs), which are specialized cell surface receptors on hematopoietic cells. Fc receptors belong to the immunoglobulin superfamily, and have been shown to mediate both the removal of antibody-coated pathogens by phagocytosis of immune complexes, and the lysing of erythrocytes and various other cellular targets (e.g.
35 tumor cells) coated with the corresponding antibody, via antibody dependent cell mediated cytotoxicity (ADCC) (Van de Winkel and Anderson, J. Leuk. Biol. 49:511-24; 1991).

Therefore, the present invention further provides antibodies that still bind to the Fc receptors to fulfill their effector functions. But, preferably the antibodies of the invention do not show a complement dependent cytotoxicity. More preferably, the antibodies of the invention do not activate the complement system, but rather inhibit the complement-mediated cell lysis.

Thus, in a preferred embodiment, the antibodies of the present invention have a human IgG Fc region, which comprises one or more amino acid substitutions, preferably the substitution of 3 or 2 amino acids, most preferably the substitution of one amino acid. The amino acid substitutions can be achieved by conventional methods, such as site-directed mutagenesis of the human IgG1 Fc region of the antibodies of the present invention.

In a more preferred embodiment, the antibodies of the present invention have a human IgG Fc region which comprises an amino acid substitution at position 324 as shown in SEQ ID NO: 18 [position 324 corresponds to position 322 according to EU numbering scheme, Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed., US Department of Health and Human Services, NIH Publication No. 91-3242, National Institutes of Health, Bethesda, MD (1991); Edelman et al., PNAS USA 63:78-85 (1969)]. The amino acid substitution is preferably K324A.

In a most preferred embodiment, the antibody of the present invention has a heavy chain, which comprises, consists essentially of or consists of an amino acid sequence:

QVQLVQSGAEVVKPGASVKVSCASGYSFTGHTMNWVRQAPQGQLEWMGLINPSDGVTRYNQKF
QGRVTITRDTSTTTVHMEITSLTSEDATYYCTREAKREWDETYWGQGLTVTVSSASTKGPSVF
PLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS
SSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKHTCTPPCPAPELLGGPSVFLFPPKPKDTLMIS
RTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE
YKCAVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE

SNGQPENNYKTTTPVLDSGSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQKSLSLSPG
K (SEQ ID NO: 18).

Further according to the invention, the antibodies comprising,
5 essentially consisting of or consisting of the following
combinations of the variable parts of the light chain and heavy
chain and the light chain and the heavy chain are preferred:

- | | | |
|----|-------------------------------|---------------|
| | A) Light chain variable part: | SEQ ID NO: 1 |
| 10 | Heavy chain variable part: | SEQ ID NO: 2 |
| | Light chain: | SEQ ID NO: 17 |
| | Heavy chain: | SEQ ID NO: 19 |
| | B) Light chain variable part: | SEQ ID NO: 1 |
| 15 | Heavy chain variable part: | SEQ ID NO: 2 |
| | Light chain: | SEQ ID NO: 17 |
| | Heavy chain: | SEQ ID NO: 18 |

Antibody according to B) is most preferred. The heavy chain
20 contains the K324A amino acid exchange.

Preferred antibodies according to the invention are humanized
forms of monoclonal mouse antibodies that are produced by hybridoma
cell line A β 6-1-6 (Deposit No. DSM ACC 2924), which is described
25 in WO 2010/009987.

The sequences of the light and heavy chains for the antibodies of
the present invention can vary. The immunoglobulins can have two
pairs of light chain/heavy chain complexes, at least one chain
30 comprising one or more mouse complementarity determining regions
(CDRs) functionally joined to human framework region segments.

In a preferred embodiment, the antibody of the invention comprises
two light chains and two heavy chains, wherein the amino acid
35 sequence of each light chain is SEQ ID NO:17, and the amino acid
sequence of each heavy chain is SEQ ID NO: 18 or SEQ ID NO: 19.

More preferably, the antibody of the invention comprises two light chains and two heavy chains, wherein the amino acid sequence of each light chain is SEQ ID NO:17, and the amino acid sequence of each heavy chain is SEQ ID NO: 19.

Most preferably, the antibody of the invention comprises two light chains and two heavy chains, wherein the amino acid sequence of each light chain is SEQ ID NO:17, and the amino acid sequence of each heavy chain is SEQ ID NO: 18.

In another embodiment, the present invention is directed to recombinant nucleic acid molecules encoding the antibodies of the invention comprising the heavy and light chain CDRs as set forth herein.

The human framework region of the antibodies of the invention is determined by comparison of a framework or variable region amino acid sequence of a CDR-providing non-human immunoglobulin with corresponding sequences in a sequence collection comprising human immunoglobulin variable regions. A sequence having a high percentage of identical amino acids is selected.

In a preferred embodiment of the invention, the variable part of the light chain having the amino acid sequence according to SEQ ID NO: 1 is encoded by a nucleic acid molecule which comprises, consists essentially of or consists of the nucleic acid sequence of:

Gacgtggtgatgactcagtctccactctccctgcccgtcacccttggacagccggcctccatct
cctgcaagtcaagtcagagcctcctgcactccgacggcaagacctacttgaactggttccagca
gaggccaggccagtctccaaggcgctgacctatctggtgtctaagctggactctgggggtccca
gacagattcagcggcagtggttcaggcactgacttcacactgaagatcagcagggtggaggctg
aggatgtcggagtgctactactgcgtgcaaggtacacacttcccattcacgttcggcgaggaggac
caaggtggaaatcaaa (SEQ ID NO: 9).

In a further preferred embodiment of the invention, the variable part of the heavy chain having the amino acid sequence according to SEQ ID NO: 2 is encoded by a nucleic acid molecule which comprises, consists essentially of or consists of the nucleic acid sequence of:

cagggtgcagctcgtgcagtcctggggctgaggtggtgaagccagggtgcctcagtgaaggtctcct
gcaaggcatctggttactcattcactggtcacaccatgaactgggtgacagggccctggaca
agggttgagtggatgggactcatcaatccttccgatggtggttactaggtacaaccagaagttc
10 cagggcagagtcaccatcaccagggacacgtccacgaccaccggttcacatggagctgaccagcc
tgacatctgaggacacggccacctaactgtacgagagaggcgaaacgggagtgaggacgagac
ttactggggccagggaaccctgggtcaccgtctcctca (SEQ ID NO: 10).

In a further preferred embodiment of the invention, the CDR regions of the light chain of the antibody of the invention are encoded by a nucleic acid molecule having the nucleic acid sequence:

V_L CDR1: tcaagtcagagcctcctgcactccgacggcaagacctaacttgaac (SEQ ID NO: 11);
20 V_L CDR2: ctggtgtctaagctggactct (SEQ ID NO: 12); and
V_L CDR3: gtgcaaggtacacacttccca (SEQ ID NO: 13).

In a further preferred embodiment of the invention, the CDR regions of the heavy chain of the antibody of the invention are encoded by a nucleic acid molecule having the nucleic acid sequence:

V_H CDR1: gggttactcattcactggtcacaccatgaac (SEQ ID NO: 14);
V_H CDR2: ctcatcaatccttccgatggtggttactaggtacaaccagaagttcc
Agggc (SEQ ID NO: 15); and
30 V_H CDR3: gaggcgaaacgggagtgaggacgagacttac (SEQ ID NO: 16).

In a further preferred embodiment of the invention, the light chain is encoded by a nucleic acid molecule which comprises, consists essentially of or consists of the amino acid sequence of:

35 Gacgtggtgatgactcagtcctccactctccctgcccgtcacccttggacagccggcctccatct
cctgcaagtcagtcagagcctcctgcactccgacggcaagacctaacttgaactggttccagca

gaggccaggccagtcctccaaggcgccctgacctatctgggtgtctaagctggactctgggggtccca
gacagattcagcggcagtggggtcaggcactgacttcacactgaagatcagcaggggtggaggctg
aggatgtcggagtctactactgcgtgcaaggtacacacttcccattcacgttcggcggaggggac
caaggtggaaatcaaaaggaccgtggccgcaccctctgtgttcattcttccccccagcgacgag
5 cagctgaagagcggcactgcatctgtcgtgtgtctgctgaacaacttctaccaaggaggcga
aagtgcagtggaaggtagacaacgccttgcaatccggcaactcccaggagagcgtgaccgagca
ggacagcaaagactcaacctacagcctgagcagctactttgaccctgtctaaggccgattacgag
aagcacaaggtgtacgcctgcgaggttaaccaccagggaactgagctctcccgtagaccaagagct
tcaacaggggagcagtg (SEQ ID NO: 20).

10

In a further preferred embodiment of the invention, the heavy
chain is encoded by a nucleic acid molecule which comprises,
consists essentially of or consists of the amino acid sequence
of:

15

caggtgcagctcgtgcagtcctgggggtgaggtggtgaagccaggtgcctcagtgaaaggtctcct
gcaaggcatctgggtactcattcactgggtcacaccatgaactgggtgagcagggccctggaca
agggtcttgagtggtgggactcatcaatccttccgatgggtgttactaggtacaaccagaagttc
cagggcagagtcaccatcaccagggaacagtcacgaccaccgttcacatggagctgaccagcc
20 tgacatctgaggacacggccacctactactgtacgagagaggcgaaacgggagtgaggacgagac
ttactggggccagggaaccctgggtcaccgtctcctcagccagcactaagggcccagcgtgttc
ccctcgccctagcagtaagagcaccagcgggtggcacggcggaacttggctgcttgggttaagg
actacttcccagagcccgtgaccgtgtcctggaactctggggcacttaccagtggcgtgcacac
cttccccgctgtactgcagagcagcggcttgtacagcttgtcttccgtcgtaacggtgccacg
25 agcagcttgggaaccagacctacatctgcaacgtaaacccacaagccatccaacaccaaggtag
acaaaaaggtcgaacccaagtcctgcgacaagacccacacctgtccacctgtcctgcacccga
gctcctgggaggtcccagcgttttctgttccctccaaagccaaaggataccctgatgatcagc
aggacccccgaggtgacctgcgtgggtgggtgagctgagccacgaagaccctgaggtcaagttca
actggtacgttgatggggtggaggtacacaatgccaaagaccaaactcgagaggagcaatacaa
30 cagcacctaccgagttgtgagcgtgcttaccgtgctgcaccaggactgggtgaacggcaaggag
tacaagtgcgctgtgagcaacaaggctctgccggctcccatcgagaagaccatcagcaaggcca
aggggccagcccaggaggccacaggtttacacgttgccccctcaaggagcaggttgaccaagaa
ccaggtttccctcacgtgccttgtgaagggttctaccccagcgacatcgccgtggaatgggag
agcaacgggagcccagagaacaactacaagacgacccccctgttctggacagcgacggctctt

tcttcctgtattcaaagctcacctggacaaaagcaggtggcagcagggtaatgtgttctcctg
cagcgtgatgcacgagggccctgcataaaccactacacccaaaagagcttgagcctctcccccggt
aag (SEQ ID NO: 21).

5 In a further preferred embodiment of the invention, the heavy
chain is encoded by a nucleic acid molecule which comprises,
consists essentially of or consists of the amino acid sequence
of:

10 caggtgcagctcgtgcagtctggggctgaggtggtgaagccaggtgcctcagtgaaggtctcct
gcaaggcatctggttactcattcactggtcacaccatgaactgggtgcgacaggccctggaca
agggttgagtggatgggactcatcaatccttccgatggtgttactaggtacaaccagaagttc
cagggcagagtcaccatcaccagggacacgtccacgaccaccgttcacatggagctgaccagcc
tgacatctgaggacacggccacctaactactgtacgagagaggcgaaacgggagtgaggacgagac
15 ttactggggccagggaaccctggtcacctgtcctcagccagcactaaggggcccagcgtgttc
cccctcgcccctagcagtaagagcaccagcgggtggcacggcggcacttggctgcttggttaagg
actacttcccagagcccgtgaccgtgtcctggaactctggggcacttaccagtggcgtgcacac
cttccccgctgtactgcagagcagcggcttgtagcgttgtcttccgtcgtaacggtgcccagc
agcagcttggaacccagacctaactctgcaacgtaaacacaaagccatccaacaccaaggtag
20 acaaaaaggctgaacccaagtctctgcgacaagaccacacctgtccacctgtcctgcacccga
gctcctgggaggtcccagcgttttctgttccctccaaagccaaaggataccctgatgatcagc
aggacccccgaggtgacctgcgtgggtggagctgagccacgaagaccctgaggtcaagttca
actggtacgttgatggggtggaggtacacaatgccaaagaccaaactcgagaggagcaatacaa
cagcacctaaccaggttgtagcgtgcttaccgtgctgcaccaggactggctgaacggcaaggag
25 tacaagtgaaggtgagcaacaaggtctgccggtcccacgagaagaccatcagcaaggcca
agggccagcccagggagccacaggtttacacgttgccccctcaaggagcaggttgaccaagaa
ccaggtttccctcacgtgccttgtagaagggttctaccccagcgacatcgccgtggaatgggag
agcaacgggcagcccagagaacaactacaagacgacccccctgttctggacagcgagcgtctt
tcttcctgtattcaaagctcacctggacaaaagcaggtggcagcagggtaatgtgttctcctg
30 cagcgtgatgcacgagggccctgcataaaccactacacccaaaagagcttgagcctctcccccggt
aag (SEQ ID NO: 22)

Further according the invention, antibodies are preferred, which
are encoded by combination of nucleic acid molecules comprising,
35 essentially consisting of or consisting of:

C) Light chain variable part: SEQ ID NO: 9
Heavy chain variable part: SEQ ID NO: 10
Light chain: SEQ ID NO: 20
Heavy chain: SEQ ID NO: 22

5

D) Light chain variable part: SEQ ID NO: 9
Heavy chain variable part: SEQ ID NO: 10
Light chain: SEQ ID NO: 20
Heavy chain: SEQ ID NO: 21

10

Combination D) is most preferred since it encodes for a humanized and de-immunized antibody which contains the K324A amino acid exchange in the heavy chain.

15 The aforementioned nucleic acid molecules can be integrated into expression vectors well known in the art. Transfection of these expression vectors in an appropriate host, the selection of the host as well as the expression collection and purification of the light chains, heavy chains, light/heavy chain dimers or intact
20 antibodies, binding fragments or other immunoglobulin forms are well-known procedures in the art.

One skilled in the art can select a vector based on desired properties, for example, for production of a vector in a particular
25 cell such as a mammalian cell or a bacterial cell.

Any of a variety of inducible promoters or enhancers can be included in the vector for expression of an antibody of the invention or nucleic acid that can be regulated. Such inducible
30 systems, include, for example, tetracycline inducible System (Gossen & Bizard, Proc. Natl. Acad. Sci. USA, 89:5547-5551 (1992); Gossen et al., Science, 268:17664769 (1995); Clontech, Palo Alto, Calif.); metallothionein promoter induced by heavy metals; insect steroid hormone responsive to ecdysone or related steroids such as
35 muristerone (No et al., Proc. Natl. Acad. Sci. USA, 93:3346-3351 (1996); Yao et al., Nature, 366:476-479 (1993); Invitrogen,

Carlsbad, Calif.); mouse mammary tumor virus (MMTV) induced by steroids such as glucocorticoid and estrogen (Lee et al., *Nature*, 294:228-232 (1981); and heat shock promoters inducible by temperature changes; the rat neuron specific enolase gene promoter (Forss-Petter, et al., *Neuron* 5; 197-197 (1990)); the human β -actin gene promoter (Ray, et al., *Genes and Development* (1991) 5:2265-2273); the human platelet derived growth factor B (PDGF-B) chain gene promoter (Sasahara, et al., *Cell* (1991) 64:217-227); the rat sodium channel gene promoter (Maue, et al., *Neuron* (1990) 4:223-231); the human copper-zinc superoxide dismutase gene promoter (Ceballos-Picot, et al., *Brain Res.* (1991) 552:198-214); and promoters for members of the mammalian POU-domain regulatory gene family (Xi et al., (1989) *Nature* 340:35-42).

Regulatory elements, including promoters or enhancers, can be constitutive or regulated, depending upon the nature of the regulation. The regulatory sequences or regulatory elements are operatively linked to one of the nucleic acid molecule sequences of the invention such that the physical and functional relationship between the nucleic acid molecule sequence and the regulatory sequence allows transcription of the nucleic acid molecule sequence. Vectors useful for expression in eukaryotic cells can include, for example, regulatory elements including the CAG promoter, the SV40 early promoter, the cytomegalovirus (CMV) promoter, the mouse mammary tumor virus (MMTV) steroid-inducible promoter, Pgtf, Moloney murine leukemia virus (MMLV) promoter, thy-1 promoter and the like.

If desired, the vector can contain a selectable marker. As used herein, a "selectable marker" refers to a genetic element that provides a selectable phenotype to a cell in which the selectable marker has been introduced. A selectable marker is generally a gene whose gene product provides resistance to an agent that inhibits cell growth or kills a cell. A variety of selectable markers can be used in the DNA constructs of the invention, including, for example, Neo, Hyg, hisD, Gpt and Ble genes, as described, for example in Ausubel et al. (*Current Protocols in*

Molecular Biology (Supplement 47), John Wiley & Sons, New York (1999)) and U.S. Patent No. 5,981,830. Drugs useful for selecting for the presence of a selectable marker include, for example, G418 for Neo, hygromycin for Hyg, histidinol for hisD, xanthine for Gpt, and bleomycin for Ble (see Ausubel et al, supra, (1999); U.S. Patent No. 5,981,830). DNA constructs of the invention can incorporate a positive selectable marker, a negative selectable marker, or both (see, for example, U.S. Patent No. 5,981,830).

Various mammalian cell culture systems can also be employed to express a recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, Cell, 23: 175 (1981). Other cell lines capable of expressing a compatible vector include, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will generally comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide required nontranscribed genetic elements.

The polypeptides can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Recovery can be facilitated if the polypeptide is expressed at the surface of the cells, but such is not a prerequisite. Recovery may also be desirable of cleavage products that are cleaved following expression of a longer form of the polypeptide. Protein refolding steps as known in this art can be used, as necessary, to complete

configuration of the mature protein. High performance liquid chromatography (HPLC) can be employed for final purification steps.

- 5 Human constant region DNA sequences can be isolated in accordance with well-known procedures from a variety of human cells.

The present invention pertains in particular to antibodies which are characterized in that they bind to A β N3pE peptides with a high affinity. The present invention also pertains to antibodies
10 which are characterized in that they bind to A β N3pE peptides or immunologically active fragments thereof with a high affinity. Said high affinity means in the context of the present invention an affinity of a K_D value of 10^{-5} M, 10^{-6} M or 10^{-7} M or better,
15 preferably a K_D value of 10^{-8} M or better, and even more preferably a K_D value of 10^{-9} M - 10^{-12} M. Thereby, the inventive antibodies bind to monomeric A β N3pE with a higher affinity than previously known antibodies.

- 20 Preferably, the binding epitope of the antibodies of the present invention in A β N3pE binds is an epitope, which carries a pyroglutamate at the N-terminus. More preferably, the binding epitope of the antibody of the invention is selected from the group consisting of

25 pEFRHDSGYEVHHQKLV (SEQ ID NO: 23),
pEFRHDSGYEVHHQKL (SEQ ID NO: 24),
pEFRHDSGYEVHHQK (SEQ ID NO: 25),
pEFRHDSGYEVHHQ (SEQ ID NO: 26),
30 pEFRHDSGYEVHH (SEQ ID NO: 27),
pEFRHDSGYEVH (SEQ ID NO: 28),
pEFRHDSGYEV (SEQ ID NO: 29),
pEFRHDSGYE (SEQ ID NO: 30),
pEFRHDSGY (SEQ ID NO: 31),
35 pEFRHDSG (SEQ ID NO: 32),

pEFRHDS (SEQ ID NO: 33),
pEFRHD (SEQ ID NO: 34)
pEFRH (SEQ ID NO: 35), and
pEFR (SEQ ID NO: 36).

5

Most preferably, the antibodies of the invention do not bind to binding epitopes that do not carry a pyroglutamate at the N-terminus.

10 Even most preferably, when binding to the aforementioned and subsequently mentioned binding epitopes, the antibodies of the invention always bind to sequences or parts of sequences, which contain the pyroglutamate at the N-terminus. The antibodies of the invention do not bind to sequences or parts of sequences, which do
15 not contain the pyroglutamate at the N-terminus.

Further, the antibody of the invention can also bind to an A β N3pE variant.

20 In the context of the present invention, an A β N3pE variant is in particular

pE-A β_{3-38} ,
pE-A β_{3-40} ,
25 pE-A β_{3-42}

Further variants of A β N3pE peptides are all A β N3pE variants, which have been shown to accumulate in the brain as a consequence of Alzheimer's disease or preceding Alzheimer's disease. These are
30 the pE-A β_{3-x} peptides, wherein x is defined as an integer between 19 and 42, e.g. in the above pE-A β_{3-42} , "42" would be the integer for "x".

In the context of the present invention a "functional variant" of
35 the inventive antibody is an antibody which retains the binding capacities, in particular binding capacities with high affinity to

a pE-A β_{3-x} peptide. The provision of such functional variants is known in the art and encompasses the above-mentioned possibilities, which were indicated under the definition of antibodies and fragments thereof.

5

In a further embodiment, the antibody is an antibody fragment, as defined above.

10 In a further preferred embodiment, the antibody of the invention is a humanized and de-immunized antibody which has the complementarity-determining regions (CDRs) of the above-defined antibodies. Preferably, the antibody can be labeled; possible labels are those as mentioned above and all those known to a person skilled in the art of diagnostic uses of antibodies in particular.

15

In another embodiment, the antibodies may be immobilized on a solid phase.

20 In another embodiment, the antibodies according to the invention and as described herein before or a fragments thereof, exhibit a binding affinity to an A β N3pE oligomer, fiber, fibril or filament which is at least 2 times, particularly at least 4 times, particularly at least 10 times, particularly at least 15 times, more particularly at least 20 times, but especially at least 25
25 times higher than the binding affinity to an A β N3pE monomer.

In still another embodiment, antibodies or fragments thereof are provided as described herein before, which substantially bind to aggregated A β , including A β plaques, which contain A β N3pE, in the
30 mammalian, particularly the human brain but, preferably, do not show any significant cross-reactivity with amyloid precursor protein (APP).

35 In another aspect of the invention, antibodies or fragments thereof are provided as described herein before, which antibodies substantially bind to oligomeric or polymeric amyloid, which contains A β N3pE, particularly amyloid β (A β) in the mammalian,

particularly the human brain but, preferably, do not show any significant cross-reactivity with amyloid precursor protein (APP).

The present invention relates also to compositions comprising said
5 antibodies and the use of said compositions for the treatment of amyloidosis, especially for the treatment of neurodegenerative disease in a mammal, in particular in a human. Said neurodegenerative disease is in particular selected from the group consisting of mild cognitive impairment (MCI), Alzheimer's disease
10 (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome. Preferably, said neurodegenerative disease is Alzheimer's disease.

15 Thus, in a preferred embodiment, the present invention is directed to a method of treating and/or preventing conditions characterized by the formation of plaques comprising A β N3pE in mammals, preferably in humans, which method comprises administering,
20 preferably peripherally, to a human in need of such treatment a therapeutically or prophylactically effective amount of an monoclonal antibody of the invention or a immunologically reactive fragment thereof, which antibody specifically binds to an epitope of the A β N3pE peptide that carries pyroglutamate at the N-
25 terminus.

In another embodiment, the invention is directed to a method to inhibit the formation of amyloid plaques and to clear or remove amyloid plaques in mammals, preferably in humans, which method
30 comprises administering to a human subject in need of such inhibition an effective amount of an antibody that binds to A β N3pE in the circulation, body fluids or tissues, especially in the brain and further preferably, leads to the clearance of A β N3pE in plasma and the brain.

35 Accordingly, the invention also provides methods of reversing cognitive decline, improving cognition, treating cognitive

decline, and preventing cognitive decline in a subject diagnosed with mild cognitive impairment (MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome and clinical or pre-clinical cerebral amyloid angiopathy, preferably Alzheimer's disease comprising administering to the subject an effective amount of an antibody of the invention.

10 The invention also provides the use of an antibody of the invention for the manufacture of a medicament, for treating, preventing, or reversing mild cognitive impairment (MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome and clinical or pre-clinical cerebral amyloid angiopathy, preferably Alzheimer's disease; or to reverse cognitive decline, improve cognition, treat cognitive decline, and prevent cognitive decline in a subject diagnosed with mild cognitive impairment (MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome and clinical or pre-clinical cerebral amyloid angiopathy, preferably Alzheimer's disease.

The invention further provides the antibodies disclosed herein for use in the prevention, treatment, or the reversion of mild cognitive impairment (MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome and clinical or pre-clinical cerebral amyloid angiopathy, preferably Alzheimer's disease; for treating, preventing, or the reversion of cognitive decline, improvement of cognition, treatment of cognitive decline, and prevention of cognitive decline in a subject diagnosed with mild cognitive impairment

(MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome and clinical or pre-clinical cerebral amyloid angiopathy, preferably Alzheimer's disease; or the inhibition of the formation of amyloid plaques or the effects of A β N3pE in mammals, preferably in humans.

In a specific embodiment the invention provides a method for retaining or increasing cognitive memory capacity but, particularly, for restoring the cognitive memory capacity of a mammal, particularly a human, suffering from memory impairment by administering to an animal, particularly a mammal or a human, an antibody of the invention, or a pharmaceutical composition comprising an antibody according to the invention and as described herein before.

The invention further provides methods to assess the response of a human subject to treatment with an antibody that binds A β N3pE or a variant thereof, comprising:

- a) administering an antibody of the invention or a fragment thereof to the subject; and
- b) measuring the concentration of A β N3pE in a biological sample taken from the subject.

The invention also provides a method of treating a human subject with an antibody that binds A β N3pE or a variant thereof, comprising:

- a) administering a first amount of the antibody or fragment thereof to the subject;
- b) within 3 hours to two weeks after administering the first dose, measuring the concentration of A β N3pE in a biological sample taken from the subject;
- c) if necessary, calculating a second amount of antibody or fragment thereof based on the result of step b), which second amount is the same as or different than the first amount; and
- d) administering the second amount of the antibody or fragment.

The invention also includes a method of assessing in a mammalian, preferably a human subject the efficacy of an antibody that binds to A β N3pE, or a fragment thereof, for inhibiting or preventing A β N3pE related amyloid plaque formation, for reducing the load of A β N3pE containing plaques, for reducing the effects of toxic A β N3pE and variants thereof, or for treating a condition or a disease associated with plaques containing A β N3pE, comprising:

- a) obtaining a first biological sample from the subject;
- 10 b) measuring a baseline concentration of A β N3pE in the first sample;
- c) administering an antibody of the invention or fragment thereof to the subject;
- d) within 3 hours to two weeks after administering the antibody or
- 15 fragment thereof, obtaining a second biological sample from the subject; and
- e) measuring the concentration of A β N3pE in the second biological sample; wherein, efficacy is related to the quantity of A β N3pE bound to the antibody in the blood and the concentration of A β N3pE, in particular the reduction of the concentration thereof, in
- 20 the second biological sample compared to the first biological sample.

The biological sample may be any sample, for example from a human.

25 In one specific example, the sample is a tissue sample, a body fluid sample or a cell sample. In one embodiment, the biological sample is selected from the group consisting of blood, serum, urine, cerebrospinal fluid (CSF), plasma, lymph, saliva, sweat, pleural fluid, synovial fluid, tear fluid, bile and pancreas

30 secretion. In a further embodiment, the biological sample is plasma. In a preferred embodiment, the biological sample is CSF.

The biological sample can be obtained from a subject in a manner well-known to a person skilled in the art. In particular, a blood

35 sample can be obtained from a subject and the blood sample can be separated into serum and plasma by conventional methods. The subject, from which the biological sample is obtained is preferably

a subject suspected of being afflicted with a disease or condition of amyloidosis, preferably Alzheimer's disease, at risk of developing Alzheimer's disease and/or being at risk of or having any other kind of dementia. In particular, the sample is obtained
5 from a subject suspected of having Mild Cognitive Impairment (MCI) and/or being in the early stages of Alzheimer's disease.

The efficacy of the antibodies of the invention in the diagnosis, prevention and/or treatment of amyloidosis, such as mild cognitive
10 impairment, Alzheimer's Disease, Familial British Dementia or Familial Danish Dementia and, e.g. neurodegeneration in Down Syndrome can be tested in existing animal models of Alzheimer's disease.

15 Suitable animal models of Alzheimer's Disease are reviewed in McGowan et al. TRENDS in Genetics, Vol. 22, No. May 2006, pp 281-289, and are selected from PDAPP, Tg2576, APP23, TgCRND8, PSEN₁M146V or PSEN₁M146L, PSAPP, APP_{Dutch}, BRI-Aβ40 and BRI-Aβ42, JNPL3, TauP301S, TauV337M, TauR406W, rTg4510, H_{tau}, TAPP, 3 x TgAD, as described below.

20

PDAPP: First mutant APP transgenic model with robust plaque pathology. Mice express a human APP cDNA with the Indiana mutation (APP_{V717F}). Plaque pathology begins between 6-9 months in hemizygous PDAPP mice. There is synapse loss but no overt cell loss and not
25 NFT pathology is observed. This model has been used widely in vaccination therapy strategies.

Tg2576: Mice express mutant APP_{SWE} under control of the hamster prion promoter. Plaque pathology is observed from 9 months of age.
30 These mice have cognitive deficits but no cell loss or NFT pathology. This model is one of the most widely used transgenic models in the field of Alzheimer's disease.

APP23: Mice express mutant APP_{SWE} under control of the Thyl promoter. Prominent cerebrovascular amyloid, amyloid deposits are observed from 6 months of age and some hippocampal neuronal loss is associated with amyloid plaque formation.
35

TgCRND8: Mice express multiple APP mutations (Swedish plus Indiana). Cognitive deficits coincide with rapid extracellular plaque development at ~ 3 months of age. The cognitive deficits
5 can be reversed by A β vaccination therapy.

PSEN_{1M146V} or PSEN_{1M146L} (lines 6.2 and 8.9, respectively): These models where the first demonstration *in vivo* that mutant *PSEN1* selectively elevates A β 42. No overt plaque pathology is observed.
10

PSAPP (Tg2576 x PSEN_{1M146L}, PSEN1-A246E + APP_{SWE}): Bigenic transgenic mice, with the addition of the mutant *PSEN1* transgene which markedly accelerated amyloid pathology compared with singly transgenic mutant APP mice, demonstrating that the PSEN1-driven
15 elevation of A β 42 enhances plaque pathology.

APP_{Dutch}: Mice express APP with the Dutch mutation that causes hereditary cerebral hemorrhage with amyloidosis-Dutch type in humans. APP_{Dutch} mice develop severe congophilic amyloid angiopathy.
20 The addition of a mutant *PSEN1* transgene redistributes the amyloid pathology to the parenchyma indicating differing roles for A β 40 and A β 42 in vascular and parenchymal amyloid pathology.

BRI-A β 40 and BRI-A β 42: Mice express individual A β isoforms without
25 APP over-expression. Only mice expressing A β 42 develop senile plaques and CAA, whereas BRI-A β 40 mice do not develop plaques, suggesting that A β 42 is essential for plaque formation.

JNPL3: Mice express 4R0N MAPT with the P301L mutation. This is the
30 first transgenic model, with marked tangle pathology and cell loss, demonstrating that MAPT alone can cause cellular damage and loss. JNPL3 mice develop motor impairments with age owing to severe pathology and motor neuron loss in the spinal cord.

Tau_{P301S}: Transgenic mice expressing the shortest isoform of 4R MAPT with the P301S mutation. Homozygous mice develop severe paraparesis at 5-6 months of age with widespread neurofibrillary
35

pathology in the brain and spinal cord and neuronal loss in the spinal cord.

Tau_{V337M}: Low level synthesis of 4R MAPT with the V337M mutation (1/10 endogenous MAPT) driven by the promoter of platelet-derived growth factor (PDGF). The development of neurofibrillary pathology in these mice suggests the nature of the MAPT rather than absolute MAPT intracellular concentration drives pathology.

Tau_{R406W}: Mice expressing 4R human MAPT with the R406W mutation under control of the CAMKII promoter. Mice develop MAPT inclusions in the forebrain from 18 months of age and have impaired associative memory.

rTg4510: Inducible *MAPT* transgenic mice using the TET-off system. Abnormal MAPT pathology occurs from one month of age. Mice have progressive NFT pathology and severe cell loss. Cognitive deficits are evident from 2.5 months of age. Turning off the transgene improves cognitive performance but NT pathology worsens.

H_{tau}: Transgenic mice expressing human genomic MAPT only (mouse MAPT knocked-out). Htau mice accumulate hyperphosphorylated MAPT from 6 months and develop Thio-S-positive NFT by the time they are 15 months old.

TAPP (Tg2576 x JNPL3): Increased MAPT forebrain pathology in TAPP mice compared with JNPL3 suggesting mutant APP and/or A β can affect downstream MAPT pathology.

3xTgAD: Triple transgenic model expressing mutant APP_{SWE}, MAPT_{P301L} on a PSEN1_{M146V} 'knock-in' background (PSEN1-KI). Mice develop plaques from 6 months and MAPT pathology from the time they are 12 months old, strengthening the hypothesis that APP or A β can directly influence neurofibrillary pathology.

Moreover, WO 2009/034158 discloses non-human transgenic animal models, wherein the transgene encodes at least one amyloid beta

(A β) peptide selected from the group consisting of A β N3E-42, A β N3Q-42, A β N3E-40 and A β N3Q-40. These A β peptides are substrates of QC and QPCTL, resulting in the cyclization of the N-terminal glutamine (Q) or glutamate (N) to pyroglutamate (pGlu). Thus, these
5 transgenic animal models provide a model system for the investigation of the effect of pGlu- A β peptides on the course of the development of neurodegeneration.

Anti-A β pN3pE antibodies may also be useful in diagnostic assays
10 for A β pN3pE, e.g. detecting its occurrence in specific cells, tissues, or serum. Thus, the antibodies according to the present invention are especially useful in a diagnostic method to detect amyloidosis, in particular a neurodegenerative disease selected from the group consisting of mild cognitive impairment (MCI),
15 Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome; preferably Alzheimer's disease.

20

For diagnostic applications, the antibody typically will be labelled with a detectable moiety. Numerous labels are available which can be generally grouped into the following categories:

25 (a) Radioisotopes, such as ^{35}S , ^{14}C , ^{125}I , ^3H , and ^{131}I . The antibody can be labeled with the radioisotope using the techniques described in *Current Protocols in Immunology*, Volumes 1 and 2, Gütigen et al., Ed., Wiley-Interscience. New York, New York. Pubs., (1991) for example and radioactivity can be measured using
30 scintillation counting.

(b) Fluorescent labels such as rare earth chelates (europium chelates) or fluorescein and its derivatives, rhodamine and its derivatives, dansyl, Lissamine, phycoerythrin and Texas Red are
35 available. The fluorescent labels can be conjugated to the antibody using the techniques disclosed in *Current Protocols in Immunology*,

supra for example. Fluorescence can be quantified using a fluorimeter.

(c) Various enzyme-substrate labels are available. The enzyme
5 generally catalyses a chemical alteration of the chromogenic
substrate which can be measured using various techniques. For
example, the enzyme may catalyze a color change in a substrate,
which can be measured spectrophotometrically. Alternatively, the
enzyme may alter the fluorescence or chemiluminescence of the
10 substrate. Techniques for quantifying a change in fluorescence are
described above. The chemiluminescent substrate becomes
electronically excited by a chemical reaction and may then emit
light which can be measured (using a chemiluminometer, for example)
or donates energy to a fluorescent acceptor. Examples of enzymatic
15 labels include luciferases (e.g, firefly luciferase and bacterial
luciferase; U.S. Patent No, 4,737,456), luciferin, 2,3-
dihydrophthalazinediones, malate dehydrogenase, urease,
peroxidase such as horseradish peroxidase (HRPO), alkaline
phosphatase. 0-galactosidase, glucoamylase, lysozyme, saccharide
20 oxidases (e.g., glucose oxidase, galactose oxidase, and glucose-
6-phosphate dehydrogenase), heterocyclic oxidases (such as uricase
and xanthine oxidase), lactoperoxidase, microperoxidase, and the
like. Techniques for conjugating enzymes to antibodies are
described in O'Sullivan et al., Methods for the Preparation of
25 Enzyme-Antibody Conjugates for use in Enzyme Immunoassay, in
Methods in Enzym (ed Langone & H. Van Vunakis), Academic Press,
New York, 73: 147-166 (1981).

Examples of enzyme-substrate combinations include, for example:
30

- (i) Horseradish peroxidase (HRPO) with hydrogen peroxidase as a
substrate, wherein the hydrogen peroxidase oxidizes a dye
precursor (e.g. orthophenylene diamine (OPD) or 3,3',5,5'-
tetramethyl benzidine hydrochloride (TMB));
- 35 (ii) alkaline phosphatase (AP) with para-Nitrophenyl phosphate as
chromogenic substrate; and

(iii) β -D-galactosidase (β -D-Gal) with a chromogenic substrate (e.g. p-nitrophenyl- β -D-galactosidase) or the fluorogenic substrate 4-methylumbelliferyl- β -D-galactosidase.

5 Numerous other enzyme-substrate combinations are available to those skilled in the art.

Sometimes, the label is indirectly conjugated with the antibody. The skilled artisan will be aware of various techniques for
10 achieving this. For example, the antibody can be conjugated with biotin and any of the three broad categories of labels mentioned above can be conjugated with avidin, or vice versa. Biotin binds selectively to avidin and thus, the label can be conjugated with the antibody in this indirect manner. Alternatively, to achieve
15 indirect conjugation of the label with the antibody, the antibody is conjugated with a small hapten (e.g. digoxin) and one of the different types of labels mentioned above is conjugated with an anti-hapten antibody (e.g. anti-digoxin antibody). Thus, indirect conjugation of the label with the antibody can be achieved.

20

The antibodies of the present invention may be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, *Monoclonal Antibodies A Manual of Techniques*, pp.147-158 (CRC
25 Press. Inc., 1987)

Competitive binding assays rely on the ability of a labeled standard to compete with the test sample analyte for binding with a limited amount of antibody. The amount of A β N3pE in the test
30 sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies generally are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may
35 conveniently be separated from the standard and analyte which remain unbound.

Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected. In a sandwich assay, the test sample analyte is bound by a first antibody which is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three-part complex. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assay). For example, one preferable type of sandwich assay is an ELISA assay, in which case the detectable moiety is an enzyme.

For immunohistochemistry, the tissue sample may be fresh or frozen or may be embedded in paraffin and fixed with a preservative such as formalin, for example.

The present invention also relates to a composition which comprises the antibodies as defined above, wherein said composition is a composition for a diagnostic use, especially for the diagnosis of a neurodegenerative disease selected from the group consisting of mild cognitive impairment (MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome; preferably Alzheimer's disease; in particular by detection of A β N3pE or variants thereof in a biological sample.

Diagnostic Kits

As a matter of convenience, the antibody of the present invention can be provided in a kit, i.e., a packaged combination of reagents in predetermined amounts with instructions for performing the diagnostic assay. Where the antibody is labelled with an enzyme, the kit will include substrates and cofactors required by the enzyme (e.g. a substrate precursor which provides the detectable chromophore or fluorophore). In addition, other additives may be included such as stabilizers, buffers (e.g. a block buffer or lysis

buffer) and the like. The relative amounts of the various reagents may be varied widely to provide for concentrations in solution of the reagents which substantially optimize the sensitivity of the assay. Particularly, the reagents may be provided as dry powders, usually lyophilized, including excipients which on dissolution will provide a reagent solution having the appropriate concentration.

The diagnostic kit according to the invention may contain a further biologically active substance as described below. Especially preferred for the use in the diagnostic kit as said further further biologically active substance is an inhibitor of glutaminyl cyclase.

The diagnostic kit of the invention is especially useful for the detection and diagnosis of amyloid- associated diseases and conditions, in particular neurodegenerative diseases selected from the group consisting of mild cognitive impairment (MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome; preferably Alzheimer's disease.

The present invention also pertains to the antibody of the invention or the composition comprising the antibody, both as defined above, for use in an *in vitro* diagnostic method. In particular, this diagnostic method is directed to diagnosis of a neurodegenerative disease selected from the group consisting of mild cognitive impairment (MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome; preferably Alzheimer's disease; especially by detecting an A β N3pE or variants thereof in a biological sample.

In a particularly preferred embodiment, the present invention pertains to the following method:

In vitro or *in situ* diagnostic method for the diagnosis of an amyloid-associated disease or condition, preferably Alzheimer's disease, comprising the following steps:
5 contacting an antibody according to the invention with a sample, preferably selected from a serum, liquor or CSF sample, most preferably a serum sample; or a specific body part or body area of
10 a subject suspected to be afflicted with said condition or disease, and
detecting binding of the antibody to A β N3pE, from the sample.

More particularly, the invention relates to a method of diagnosis
15 of an amyloid-associated disease or condition, preferably Alzheimer's disease, comprising detecting the immunospecific binding of an antibody of the invention or an immunologically active fragment thereof to A β N3pE, in a sample or *in situ* which includes the steps of
20 (a) bringing the sample or a specific body part or body area suspected to contain the amyloid protein into contact with an antibody of the invention, or a fragment thereof;
(b) allowing the antibody and/or a functional part thereof, to bind to A β N3pE to form an immunological complex;
25 (c) detecting the formation of the immunological complex; and
(d) correlating the presence or absence of the immunological complex with the presence or absence of A β N3pE in the sample or specific body part or area.

30 Also comprised is a method of determining the extent of amyloidogenic plaque burden in a tissue and/or body fluids comprising
(a) obtaining a sample representative of the tissue and/or body fluids under investigation;
35 (b) testing said sample for the presence of amyloid protein with an antibody according to the invention, or a chimeric antibody or a fragment thereof;

(c) determining the amount of antibody bound to the protein; and
(d) calculating the plaque burden in the tissue and/or body fluids.

5 In particular, the invention relates to a method of determining the extent of amyloidogenic plaque burden in a tissue and/or body fluids, wherein the formation of the immunological complex in step c) is determined such that presence or absence of the immunological complex correlates with presence or absence of amyloid protein, in
10 particular A β N3pE.

In still another embodiment, the invention relates to a composition comprising the antibody according to the invention, or a chimeric antibody or a fragment thereof, and as described herein before
15 including any functionally equivalent antibody or any derivative or functional parts thereof, in particular a composition which is a pharmaceutical composition optionally further comprising a pharmaceutically acceptable carrier.

20 In another embodiment of the invention, said composition comprises the antibody of the invention in a therapeutically effective amount.

Further comprised by the invention is a mixture comprising an
25 antibody of the invention, or a chimeric antibody or a fragment thereof, and as described herein before including any functionally equivalent antibody or any derivative or functional parts thereof, in a therapeutically effective amount and, optionally, a further biologically active substance and/or a pharmaceutically acceptable
30 carrier and/or a diluent and/or an excipient.

In particular, the invention relates to a mixture, wherein the further biologically active substance is a compound used in the medication of amyloidosis, a group of diseases and disorders
35 associated with amyloid or amyloid-like protein such as A β N3pE involved in neurodegenerative diseases selected from the group consisting of mild cognitive impairment (MCI), Alzheimer's disease

(AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome; preferably Alzheimer's disease.

In another embodiment of the invention, the other biologically active substance or compound may also be a therapeutic agent that may be used in the treatment of amyloidosis caused by A β N3pE or may be used in the medication of other neurological disorders.

The other biologically active substance or compound may exert its biological effect by the same or a similar mechanism as the antibody according to the invention or by an unrelated mechanism of action or by a multiplicity of related and/or unrelated mechanisms of action.

Generally, the other biologically active compound may include neutron-transmission enhancers, psychotherapeutic drugs, acetylcholine esterase inhibitors, calcium-channel blockers, biogenic amines, benzodiazepine tranquillizers, acetylcholine synthesis, storage or release enhancers, acetylcholine postsynaptic receptor agonists, monoamine oxidase-A or -B inhibitors, N-methyl- D-aspartate glutamate receptor antagonists, non-steroidal anti-inflammatory drugs, antioxidants, and serotonergic receptor antagonists.

More particularly, the invention relates to a mixture comprising at least one compound selected from the group consisting of compounds effective against oxidative stress, anti-apoptotic compounds, metal chelators, inhibitors of DNA repair such as pirenzepin and metabolites, 3- amino-1-propanesulfonic acid (3 APS), 1,3-propanedisulfonate (1,3PDS), α -secretase activators, β - and γ -secretase inhibitors, tau proteins, neurotransmitter, β -sheet breakers, attractants for amyloid beta clearing / depleting cellular components, inhibitors of N-terminal truncated amyloid beta including pyroglutamated amyloid beta 3-42, such as

inhibitors of glutaminy cyclase, anti-inflammatory molecules, or cholinesterase inhibitors (ChEIs) such as tacrine, rivastigmine, donepezil, and/or galantamine, M1 agonists and other drugs including any amyloid or tau modifying drug and nutritive
5 supplements, and nutritive supplements, together with an antibody according to the present invention and, optionally, a pharmaceutically acceptable carrier and/or a diluent and/or an excipient.

10 The invention further relates to a mixture, wherein the compound is a cholinesterase inhibitor (ChEIs), particularly a mixture, wherein the compound is one selected from the group consisting of tacrine, rivastigmine, donepezil, galantamine, niacin and memantine.

15 In a further embodiment, the mixtures according to the invention may comprise niacin or memantine together with an antibody according to the present invention and, optionally, a pharmaceutically acceptable carrier and/or a diluent and/or an
20 excipient.

In a further embodiment, the mixtures according to the invention may comprise a glutaminy cyclase inhibitor together with an antibody according to the present invention and, optionally, a
25 pharmaceutically acceptable carrier and/or a diluent and/or an excipient.

Preferred inhibitors of glutaminy cyclase are described in WO 2005/075436, WO 2008/055945, WO 2008/055947, WO 2008/055950,
30 WO 2008/065141, WO 2008/110523, WO 2008/128981, WO 2008/128982, WO 2008/128983, WO 2008/128984, WO 2008/128985, WO 2008/128986, WO 2008/128987, WO 2010/026212, WO 2011/131748, WO 2011/029920, WO 2011/107530, WO 2011/110613, WO 2012/123563 and WO 2014/140279, the disclosure of which is incorporated herein by reference.

35 In still another embodiment of the invention mixtures are provided that comprise "atypical antipsychotics" such as, for example

clozapine, ziprasidone, risperidone, aripiprazole or olanzapine for the treatment of positive and negative psychotic symptoms including hallucinations, delusions, thought disorders (manifested by marked incoherence, derailment, tangentiality), and bizarre or disorganized behavior, as well as anhedonia, flattened affect, apathy, and social withdrawal, together with an antibody, particularly a monoclonal antibody according to the invention, but particularly a chimeric antibody or a fragment thereof, or an antibody or a fragment thereof according to the invention and as described herein and, optionally, a pharmaceutically acceptable carrier and/or a diluent and/or an excipient.

In a specific embodiment of the invention, the compositions and mixtures according to the invention and as described herein before comprise the antibody of the invention and the biologically active substance, respectively, in a therapeutically effective amount.

Other compounds that can be suitably used in mixtures in combination with the antibody according to the present invention are described in WO2008/065141 (see especially pages 37/38), including PEP-inhibitors (pp. 43/44), LiCl, inhibitors of dipeptidyl aminopeptidases, preferably inhibitors of DP IV or DP IV-like enzymes (see pp. 48/49); acetylcholinesterase (ACE) inhibitors (see p. 47), PIMT enhancers, inhibitors of beta secretases (see p. 41), inhibitors of gamma secretases (see pp. 41/42), inhibitors of neutral endopeptidase, inhibitors of phosphodiesterase-4 (PDE-4) (see pp. 42/43), TNFalpha inhibitors, muscarinic M1 receptor antagonists (see p. 46), NMDA receptor antagonists (see pp. 47/48), sigma-1 receptor inhibitors, histamine H3 antagonists (see p. 43), immunomodulatory agents, immunosuppressive agents or an agent selected from the group consisting of antegren (natalizumab), Neurelan (fampridine-SR), campath (alemtuzumab), IR 208, NBI 5788/MSP 771 (tiplimotide), paclitaxel, Anergix.MS (AG 284), SH636, Differin (CD 271, adapalene), BAY 361677 (interleukin-4), matrix-metalloproteinase-inhibitors (e.g. BB 76163), interferon-tau (trophoblastin) and SAIK-MS; beta-amyloid antibodies (see p.44), cysteine protease

inhibitors (see p. 44); MCP-1 antagonists (see pp. 44/45), amyloid protein deposition inhibitors (see 42) and beta amyloid synthesis inhibitors (see p. 42), which document is incorporated herein by reference.

5

In another embodiment, the invention relates to a mixture comprising the antibody according to the invention, or a chimeric antibody or a fragment thereof and as described herein before and/or the biologically active substance in a therapeutically effective amount.

10

The pharmaceutical compositions may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington : The Science and Practice of Pharmacy, 21th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 2005.

15

The pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients should be suitable for the chosen antibody of the present invention and the chosen mode of administration. Suitability for carriers and other components of pharmaceutical compositions is determined based on the lack of significant negative impact on the desired biological properties of the chosen antibody or pharmaceutical composition of the present invention (e.g., less than a substantial impact (10% or less relative inhibition, 5% or less relative inhibition, etc.)) on antigen binding.

20

25

A pharmaceutical composition of the present invention may also include diluents, fillers, salts, buffers, detergents (e.g., a nonionic detergent, such as Tween-20 or Tween- 80), stabilizers (e.g., sugars or protein-free amino acids), preservatives, tissue fixatives, solubilizers, and/or other materials suitable for inclusion in a pharmaceutical composition.

30

35

The actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, or the amide thereof, the route of administration, the time of administration, the rate of excretion of the particular antibody being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

The pharmaceutical composition may be administered by any suitable route and mode. Suitable routes of administering an antibody of the present invention in vivo and in vitro are well known in the art and may be selected by those of ordinary skill in the art.

In one embodiment, a pharmaceutical composition of the present invention is administered parenterally.

The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and include epidermal, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, intratendinous, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, intracranial, intrathoracic, epidural and intrasternal injection and infusion.

In one embodiment that pharmaceutical composition is administered by intravenous or subcutaneous injection or infusion.

Pharmaceutically acceptable carriers include any and all suitable solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonicity agents, antioxidants and absorption delaying agents, and the like that are physiologically compatible with an
5 antibody of the present invention.

Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the present
10 invention include water, saline, phosphate buffered saline, ethanol, dextrose, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, corn oil, peanut oil, cottonseed oil, and sesame oil, carboxymethyl cellulose colloidal solutions,
15 tragacanth gum and injectable organic esters, such as ethyl oleate, and/or various buffers. Other carriers are well known in the pharmaceutical arts.

Pharmaceutically acceptable carriers include sterile aqueous
20 solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the active
25 antibody, use thereof in the pharmaceutical compositions of the present invention is contemplated.

Proper fluidity may be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the
30 required particle size in the case of dispersions, and by the use of surfactants.

Pharmaceutical compositions of the present invention may also comprise pharmaceutically acceptable antioxidants for instance (1)
35 water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as

ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid,
5 phosphoric acid, and the like.

Pharmaceutical compositions of the present invention may also comprise isotonicity agents, such as sugars, polyalcohols, such as mannitol, sorbitol, glycerol or sodium chloride in the
10 compositions.

The pharmaceutical compositions of the present invention may also contain one or more adjuvants appropriate for the chosen route of administration such as preservatives, wetting agents, emulsifying
15 agents, dispersing agents, preservatives or buffers, which may enhance the shelf life or effectiveness of the pharmaceutical composition. The antibodies of the present invention may be prepared with carriers that will protect the antibody against rapid release, such as a controlled release formulation, including
20 implants, transdermal patches, and microencapsulated delivery systems. Such carriers may include gelatin, glyceryl monostearate, glyceryl distearate, biodegradable, biocompatible polymers such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid alone or with a
25 wax, or other materials well known in the art. Methods for the preparation of such formulations are generally known to those skilled in the art. See, e.g., Sustained and Controlled Release Drug Delivery Systems, J. R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

30

In one embodiment, the antibodies of the present invention may be formulated to ensure proper distribution in vivo. Pharmaceutically acceptable carriers for parenteral administration include sterile aqueous solutions or dispersions and sterile powders for the
35 extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. Except insofar as any

conventional media or agent is incompatible with the antibody, use thereof in the pharmaceutical compositions of the present invention is contemplated.

5 Pharmaceutical compositions for injection must typically be sterile and stable under the conditions of manufacture and storage. The composition may be formulated as a solution, microemulsion, liposome, or other ordered structure suitable to high drug concentration. The carrier may be a aqueous or non-aqueous solvent
10 or dispersion medium containing for instance water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. The proper fluidity may be maintained, for example, by the use of
15 a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as glycerol, mannitol, sorbitol, or sodium chloride in the
20 composition. Prolonged absorption of the injectable compositions may be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin. Sterile injectable solutions may be prepared by incorporating the antibody in the required amount in an appropriate solvent with one
25 or a combination of ingredients e.g. as enumerated above, as required, followed by sterilization microfiltration. Generally, dispersions are prepared by incorporating the antibody into a sterile vehicle that contains a basic dispersion medium and the required other ingredients e.g. from those enumerated above. In
30 the case of sterile powders for the preparation of sterile injectable solutions, examples of methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

35

Sterile injectable solutions may be prepared by incorporating the antibody in the required amount in an appropriate solvent with one

or a combination of ingredients enumerated above, as required, followed by sterilization microfiltration. Generally, dispersions are prepared by incorporating the antibody into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, examples of methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Dosage regimens in the above methods of treatment and uses are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. Parenteral compositions may be formulated in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of antibody calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the present invention are dictated by and directly dependent on (a) the unique characteristics of the antibody and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an antibody for the treatment of sensitivity in individuals.

The effective dosages and the dosage regimens for the antibodies of the invention depend on the disease or condition to be treated and may be determined by the persons skilled in the art. An exemplary, non-limiting range for a therapeutically effective amount of an antibody of the present invention is about 0.1 -10 mg/kg/body weight, such as about 0.1-5 mg/kg/body weight, for example about 0.1-2 mg/kg/body weight, such as about 0.1-1

mg/kg/body weight, for instance about 0.15, about 0.2, about 0.5, about 1, about 1.5 or about 2 mg/kg/body weight.

A physician or veterinarian having ordinary skill in the art may readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the 8ηίί-AβpE3 antibody employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. In general, a suitable daily dose of a composition of the present invention will be that amount of the antibody which is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. Administration may e.g. be intravenous, intramuscular, intraperitoneal, or subcutaneous, and for instance administered proximal to the site of the target. If desired, the effective daily dose of a pharmaceutical composition may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms. While it is possible for an antibody of the present invention to be administered alone, it is preferable to administer the antibody as a pharmaceutical composition as described above.

25 **EXAMPLES**

1. Humanization approach to generate N3pE-Aβ specific antibodies

The humanized and de-immunized antibodies according to the invention are humanized and de-immunized forms of monoclonal mouse antibodies that are produced by hybridoma cell line Aβ 6-1-6 (Deposit No. DSM ACC 2924), which is described in WO 2010/009987. Humanization was performed as described in WO 2017/009459. Working Example 1 disclosed on pp. 58-60 in WO 2017/009459 is hereby incorporated by reference.

2. RNA isolation and cDNA synthesis

As source for constant sequences, RNA of human B cells was isolated by lysis of 500 μ l whole blood with 5ml 1x FACS Lysis Solution (Becton Dickinson) for 10 minutes at room temperature. The lysate was centrifuged at 300g for 5 min; the pellet was washed two times
 5 with PBS and was then resolved in 350 μ l RA1 Buffer of Nucleo Spin® RNA II (Macherey-Nagel) and added with 3.5 μ l 0,5M TCEP (SIGMA). The RNA was isolated by manufacturers' instructions. 10 μ l of RNA was first incubated with 1 μ l 0.5 μ g/ μ l OligodT Primer (Invitrogen) and 1 μ l 10mM dNTPs for 5 min at 65°C. Then 4 μ l of 5x First Strand
 10 Buffer (Invitrogen), 2 μ l of 100mM DTT and 0.5 μ l SuperScript III Reverse Transcriptase (Invitrogen) was added to 20 μ l and mix was incubated for 5 minutes at 25 °C, 50min at 50°C and 15 min at 70°C. By PCR with primer pairs shown in Table 2, synthesized cDNA of constant region of light and heavy chain could be amplified.

15

Table 2: Primer for cloning of constant region

SEQ ID NO	Name	Sequence
43	hkappa5'	ACTGTGGCTGCACCATCTGTCTTC
44	hkappa3'	CTAACTCTCCCTGTTGAAGCTC
45	hlgG1Hc5'1	AGGGAACCTGGTCACCGTCTCC
46	hlgG1Hc3'	TCATTACCCGGAGACAGGGAGAGG

For amplification of the PCR product of the light chain of clone#6
 20 following forward and reverse primers were used:

RT_chim_humKl6f: CAAGTCAGAGCCTCTTATATAGTG (SEQ ID NO: 47);

RT_chim_humKl6r: GTACCTTGACGCAGTAATAAAC (SEQ ID NO: 48).

For amplification of the reference gene mouse HPRT Primer were used.

25 To perform the amplification 7.5 μ l Sybergreen (Firma), 1 μ l Primer forward (25 pmol/ μ l), 1 μ l Primer reverse (25 pmol/ μ l), 5.5 μ l ddH₂O and 1 μ l cDNA were used in cycler.

3. Expression of recombinant antibody in CHO cells by separately cloning LC and HC into two different expression plasmids

The sequences of the light and heavy chain of the antibodies were
5 separately cloned into two different mammalian expression vectors,
pCDNA3.1 and HC-pOptiVEC respectively. To identify the optimal
combination of vectors to express the recombinant antibody in CHO
cell culture, different plasmid combinations were used to perform
transient expressions in adherent CHO cells. In a second step, it
10 was investigated whether different DNA ratios between LC and HC
plasmid influences the expression level. With transfection of 3 µg
LC-pCDNA3.1 and 1 µg HC-pOptiVEC, an increased expression level
was found.

15 For further adherent CHO cells expression of antibody, plasmid
combination of LC-pCDNA3.1 and 1 µg HC-pOptiVEC and a plasmid DNA
ratio of LC 3:1 HC was used.

Freestyle™ CHO suspension cells were used in the following
20 transfections to cultivate a higher amount of transient expressing
cells which to generate recombinant antibodies. First was tested
whether an excess of LC plasmid could improve the expression of
antibody like in case of the adherent cells. Like in adherent CHO
cells a LC to HC plasmid DNA ratio of 1:1 and 3.1 was used. Western
25 blot analysis revealed that an excess of LC plasmid increases the
expression of antibody as in the case of adherent CHO cells. By
measurement of cell viability it become obvious, that the cell
viability decreases to about 50% of transfected cells after 6 days.
After day six, no further increase of antibody level in supernatant
30 was detectable. Consequently, culture supernatants were harvest at
day six in the case of following transfections.

To investigate if the produced antibodies are efficient
transported into the cell supernatant, a cell lysate sample was
35 applied to SDS PAGE, analyzed by Western blot. GAPDH, a
housekeeping cytoplasmic protein was used for reference loading
comparable amounts of cell lysate protein to the SDS gel. In cell

lysate of antibody expressing CHO cells a strong band of 120 kDa occurs migrating at the same size as detected in the cell supernatant.

5 **4. Purification of recombinant antibody by Protein G chromatography**

The antibody clone#6 was purified to investigate the antigen binding property of the protein in comparison with the original murine antibody. Therefore 300 ml supernatant with expressed
10 chimeric and humanized antibody was produced and purified by Protein G chromatography. Because the amount of expressed antibody was very low, the yield was less than 0.1 µg/ml, in total 25 µg purified protein. The eluted antibody was concentrated to about 200 µg/ml and 2 µg protein was applied to SDS-PAGE following
15 Coomassie Blue staining.

5. Stable cell line generation of humanized antibody

The parental antibody of the humanized antibody of the present invention is the clone#6 variant disclosed in WO 2017/009459, which
20 has the light chain variable region with the amino acid sequence:

DVVMTQSPLSLPVTLGQPASISCKSSQSLLYSDGKTYLNWFQQRPGQSPRRLTYLVSKLDSGVP
DRFSGSGSGTDFTLKISRVEAEDVGVYYCVQGTHFPFTFGGGTKVEIK (SEQ ID NO: 1),
which is disclosed as SEQ ID NO: 14 in WO 2017/009459;

25

and

which has the heavy chain variable region with the amino acid sequence:

30

QVQLVQSGAEVKKSGASVKVSCASGYSFTGHTMNWVRQAPGQGLEWMGLINPSNGVTRYNQKF
QGRVTITRDSTTTVHMEITSLTSEDATYYCTREAKREWDETYWGQGLTVTVSS (SEQ ID
NO: 49); which is disclosed as SEQ ID NO: 27 in WO 2017/009459.

35

Two clones, C06.08 and C06.09, of the parental clone#6 variant were chosen for stable cell line generation. For the generation

of C06.08- and C06.09-producing Chinese hamster ovary (CHO) DG44 cells expression plasmids containing the sequences of the heavy and light chain genes for both antibodies were first generated. The plasmids were linearized, purified by isopropanol precipitation, reconstituted in 10 mM Tris pH 8.0 and used in transfections as follows: 1 x 10⁶ CHO-DG44 cells were suspended in 100 µL Nucleofector solution V (Lonza) and mixed with 10 µg linearized vector DNA. The suspension was pulsed with an electroporator Amaxa Nucleofector II (Lonza). Subsequently, 500 µL CD CHO medium (Invitrogen) was added and the transfected cell pools were cultivated for 24 hours.

For the generation of stable mini pools (MPs) transfected cell pools were combined one day after transfection, transferred either into CD CHO medium and seeded at 2000 cells/well and 4000 cells/well in 96-well plates, respectively, or transferred into CD CHO medium with 2.5 nM Methotrexate (MTX) and seeded at 4000 cells/well in 96-well plates. After a cultivation period of 21 (C06.08) or 27 (C06.09) days, 20 MPs were transferred to 24-well plates. On day 22 or day 28 post transfection, the MPs were transferred to 12-well plates and the MPs were expanded to 6-well plates on day 34 or day 48. On day 34 (C06.08) or 48 (C06.09), pools were transferred into selective medium with 30 nM MTX to induce the amplification process and antibody expression titers were determined by Octet measurements.

25

Results

Table 2. Analysis of C06.08 stable Mini Pools		
Day 1	Day 21	Day 34 to 72
	Selected MPs	MTX amplification

2000 cells/well	#1-10	No cell growth; very low or no antibody production
4000 cells/well	#11-15	
4000 cells/well & 2.5 nM MTX	#15-20	

Table 3. Analysis of C06.09 stable Mini Pools		
Day 1	Day 27	Day 48 to 52
	Selected MPs	MTX amplification
2000 cells/well	#1-10	No cell growth; very low or no antibody production
4000 cells/well	#11-15	
4000 cells/well & 2.5 nM MTX	#15-20	

As summarized in Tables 2 and 3, all MPs displayed poor cells growth, very low to no detectable antibody production (<10
5 pg/cell/day) and many MPs did not survive the MTX amplification. Repeat transfection experiments with a new set of reagents lead to similar results, while transfection with control vectors resulted in normal protein expression levels. These results suggested that genome-integrated C06.08 and C06.09 proteins may
10 have inherent toxic effects on CHO-DG44 cells, e.g. these proteins may trigger cell death due to the overexpression of misfolded C06.08 and C06.09 proteins in the nucleus. Further

modifications to the primary sequence of C06.08 and C06.09 are warranted, which correct protein misfolding and eliminate the induction of CHO-DG44 cell death.

5 PBD-C06 modifications to improve protein folding and expression

In silico analysis of the C06 amino acid sequences revealed several stretches of amino acids (spots) in the framework regions of the heavy chain that were proposed to negatively influence protein
10 folding. To examine whether these spots could be exchanged with alternative amino acids that improve protein expression, four individual heavy chain mutations (K12V, S14P, N55D and F64V) were first synthesized and tested individually in transient transfections for protein expression together with the light chain
15 of SEQ ID NO: 17. Target binding studies revealed that the F64V modification prevented antibody binding. In a second set of experiments, combinations of each mutation were integrated in pairs and triplets into individual heavy chain genes and tested in combination with the light chain of SEQ ID NO: 17 for expression
20 and target binding studies (Figs. 1 and 2).

Results

The combination of the K12V, S14P and N55D mutations resulted in the best transient C06 expression titers and optimal target binding
25 properties (Figs 1 and 2) and a respective clone comprising all these three mutations was chosen for CHO-DG44 stable transfection studies.

Stable expression of the humanized antibody in CHO-DG44 cells

30 The parental CHO-DG44 cells were originally obtained from Gibco, Life Technologies (Freedom™ DG44 Kit) now managed by Thermo Fisher

Scientific. The CHO cells line is DHFR-deficient and cGMP banked with documentation.

Transfection of CHO-DG44 cells with vectors expressing humanized antibody comprising the light chain of SEQ ID NO: 1 and heavy chain of SEQ ID NO: 2 (which comprises the K12V, S14P and N55D mutations) resulted in several-fold improved number of CHO-DG44 clones compared to prior transfection attempts (see above). Furthermore, after MTX exposure multiple clones with high target antibody expression titers were identified, revealing that the K12V, S14P and N55D mutations improved antibody folding and expression after stable integration into the CHO-DG44 genome. Prolonged stable expression was confirmed under fed-batch conditions over extended periods of cultivation confirming that the chosen alterations (K12V, S14P and N55D) in the heavy chain of parental clone#6 facilitated proper folding and expression in CHO-DG44 cells.

Additional stable cell lines were obtained after further vector and electroporation optimization. Initial transfectants, prior to gene amplification, expressed high levels of antibody early after isolation and expansion to 0.5 ml cultures in a 24-well plate (Fig. 3). Production titers at day 7 further increased after the first MTX exposure (Fig. 4), suggesting that these clones, in particular c17, will be a much more efficient producer.

25 Summary

Four heavy chain mutations were tested individually to see if they would increase antibody expression in transiently transfected HEK293 cells. Three of these mutations significantly increased antibody expression while the fourth decreased expression. Combinations of the individual mutations were also tried in pairs and the triple mutation, all of which were expressed well and purified by protein A chromatography. Upon Biacore analysis it was found that the combination of four mutations decreased binding while the combination of specific mutations K12V, S14P and N55D

were expressed well and had optimal binding properties. No changes in the light chain sequences were necessary for high expression.

Studies were performed in stably transfected CHO DG44 cells and higher levels of antibody expression were obtained after initial selection and following gene amplification induced by step-wise increases of methotrexate (MTX). It was further observed that higher levels of antibody expression did not prevent good cell growth. In summary, the transfection campaign lead to the identification of several new candidates for CMC manufacturing.

5. Surface Plasmon Resonance measurement to analyze the binding of the humanized and de-immunized antibody to monomeric A β peptides

Surface Plasmon Resonance measurement was used to investigate the binding efficacy of the antibody, which comprises the variable part of the light chain with SEQ ID NO. 1 and the the variable part of the heavy chain with SEQ ID NO. 2. To prevent mass transfer and avidity effects during measurement, the following procedure was used.

First a polyclonal α -human antibody was coupled to an SPR-Chip subsequent loaded with the antibody until the Response Unit was more than 1000.

Kinetic measurements were performed at different concentrations (of 5 to 1000 nM) of A β -peptide. The graphs of the measured series are shown as an overlay plot with the sensorgrams, corrected by the sensogram measuring the running buffer, aligned at the time of injection and the baseline adjusted to zero before injection. The results are evaluated according to a simple 1:1 interaction model (Langmuir fit), which promote the k_{off} and k_{on} rate constants.

Apparent kinetic constants according to the 1:1 Langmuir fitting are listed in Table 4. Due to the fact that the antibody is non-covalently bound to the chip surface, small amounts of antibody molecules were washed out during the measurement. Therefore the R_{max} values were fitted locally for every single sensogram.

Table 4: Statistics of Langmuir fit in kinetics of humanized and de-immunized antibody clone#6

Aβ peptide	K_D	k_a, (s⁻¹M⁻¹)	k_d, (s⁻¹)	R_{max}	RI	Chi²
A β (pE3-18)	8.22 nM	4.79 x 10 ⁵	3.94 x 10 ⁻³	Global (43.9 RU)	local	0.462
A β (3-18)	1.21 μ M	3.09 x 10 ³	3.74 x 10 ⁻³	Global (40.8 RU)	local	0.243
A β (1-18)	23.1 μ M	173	3.99 x 10 ⁻³	Global (57.4 RU)	local	0.400
A β (4-18)	28.3 μ M	–	–	48.9 RU	–	0.0815

5

Binding to A β (pE3-18)

The qualitative and visual inspection of the recorded sensorgrams shows a good shape for the association as well as the dissociation phase. Furthermore, a concentration dependent elevation of the response signal and an increase of the initial slope in the association phase were observed. The maximal SPR signals were ~44 RU, which was within the optimal range measuring binding kinetics without or negligible mass transport effects. No baseline drift was observed, therefore the data were evaluated using the Langmuir 1:1 binding model obtaining the binding properties of this interaction and quality of the fit (Tab. 4).

10

15

Binding to A β (3-18)

The qualitative and visual inspection of the recorded sensorgrams shows a good shape for the association as well as the dissociation phase. Furthermore, a concentration dependent elevation of the response signal and an increase of the initial slope in the association phase were observed. The maximal SPR signals were ~44 RU, which was within the optimal range measuring binding kinetics

20

without or negligible mass transport effects. No baseline drift was observed, therefore the data were evaluated using the Langmuir 1:1 binding model obtaining the binding properties of this interaction and quality of the fit (Tab. 4).

5

Binding to A β (1-18)

The qualitative and visual inspection of the recorded sensorgrams shows a typical shape for a weak binding appropriate to the used peptide concentrations. A slight baseline drift was observed; therefore, the data were fitted using the Langmuir 1:1 binding with drifting baseline model. Evaluation of the kinetic data (Tab. 4) yielded a dissociation constant of 23.1 μ M, which was above the highest measured A β (1-18) concentration. Furthermore, the calculated Rmax value was significantly higher than the response values of the highest concentration. Both facts indicate that the obtained binding properties be definitely in the range $K_D > 10 \mu$ M.

Binding to A β (4-18)

The qualitative and visual inspection of the recorded sensorgrams shows a typical shape for a very weak binding appropriate to the used peptide concentrations. Because of a very fast association and dissociation the data cannot be evaluated kinetically, therefore a steady-state model was used. The evaluation of the response signals in steady-state (Tab. 4) yielded a dissociation constant of 28.3 μ M, which was above the highest measured A β (4-18) concentration. Furthermore, the calculated Rmax value was significantly higher than the response values of the highest concentration. Both facts indicate that the obtained binding properties must be definitely in the range $K_D > 10 \mu$ M.

30

6. Binding to A β fibrils and A β oligomers

Fibrils of A β (1-42) peptides were generated according standard protocols at pH 8.7 and 37°C. After complete fibrillation the structures were aspirated in 60 μ l, diluted in 140 μ l running buffer and loaded onto a sensor chip by use of capture antibodies with a flow rate of 1 μ l/min. Then, the system was washed until

35

getting a stable baseline. About 100 RU of the A β (1-42)_fibrils were captured onto the sensor chip.

Oligomers of A β (1-42) peptides_were generated according to the
5 ACUMEN protocol in Ham's F12 medium overnight at 4°C and separated from aggregated A β by centrifugation. Afterwards the oligomers were aspirated in 20 μ l, diluted in 60 μ l running buffer and loaded onto a sensor chip by use of capture antibodies with a flow rate of 1 μ l/min. Then, the system was washed overnight with 100 μ l/min
10 getting a stable baseline. About 300 RU of the of A β (1-42) oligomers were captured onto the sensor chip.

For fibrils of A β (1-42) peptides: After getting a stable baseline the interaction analysis was performed by 11 consecutive
15 injections (10 pM, 30 pM, 90 pM, 270 pM, 810 pM, 2.43 nM, 7.29 nM, 21.87 nM, 65.61 nM, 196.83 nM and 590.49 nM) of the parental antibody clone #6 as disclosed in WO 2017/009459 and the humanized and de-immunized antibody clone#6 of the present invention with 30 μ l/min, 480 sec contact time and 1200 sec dissociation time. The
20 obtained sensorgrams were evaluated using five consecutive concentrations (starting with the first injection showing binding) and the Single Cycle Kinetics model with a global fit of Rmax and baseline drift and local fits of the bulk effect of every injection.

25
For oligomers of A β (1-42) peptides: After getting a stable baseline the interaction analysis was performed by 11 consecutive injections (10 pM, 30 pM, 90 pM, 270 pM, 810 pM, 2.43 nM, 7.29 nM, 21.87 nM, 65.61 nM, 196.83 nM and 590.49 nM) the parental antibody
30 clone #6 as disclosed in WO 2017/009459 and the humanized and de-immunized antibody clone#6 of the present invention with 30 μ l/min, 480 sec contact time and 3600 sec dissociation time. The obtained sensorgrams were evaluated using five consecutive concentrations (starting with the first injection showing binding) and the Single
35 Cycle Kinetics model with a global fit of Rmax and baseline drift and local fits of the bulk effect of every injection.

Results**Table 5: Binding to fibrils A β (1-42) peptides**

Antibody	K _D	k _a M ⁻¹ s ⁻¹	k _d 10 ⁻⁵ s ⁻¹	R _{max}	Chi ²
A	65.9 pM	9.9·10 ⁵	6.53·	26 RU	0.336
B	1.67 nM	5.97·10 ⁴	9.99·	19.2 RU	0.602

5

Table 6: Binding to oligomers of A β (1-42) peptides

Antibody	K _D	k _a M ⁻¹ s ⁻¹	k _d 10 ⁻⁴ s ⁻¹	R _{max}	Chi ²
A	7.61 nM	1.86·10 ⁴	1.42·	442 RU	6.31
B	269 nM	868	2.33·	216 RU	0.284

In tables 5 and 6,

10

A is the parental antibody clone #6 disclosed in WO 2017/009459 with a variable region of the light chain of SEQ ID NO: 1 and a variable region of the heavy chain of SEQ ID NO: 49; and

15

B is the humanized and de-immunized antibody clone #6 of the present invention with a variable region of the light chain of SEQ ID NO: 1 and a variable region of the heavy chain of SEQ ID NO: 2.

20

As can be seen from these results, the invention provides antibodies and fragments thereof, wherein the antibodies show an increased selectivity towards oligomers and/or fibrils of A β peptides. The antibodies of the present invention show a manifold, such as 10 times, 25 times, 50 times, 100 times, 150 times, 200 times, 250 times or more than 250 times lower binding constant (K_D value) for binding to oligomers and/or fibrils of A β (1-42) than comparable monoclonal antibodies known in the prior art, in particular compared to the parental antibody disclosed in WO 2017/009459. Accordingly, the antibodies of the present

30

invention, which were established to selectively bind to A β N3pE peptides, are more specific for A β N3pE peptides and show a decreased cross-reactivity against A β peptides other than A β N3pE.

5 **7. Binding to Fc gamma receptors**

The binding of two antibodies, which either comprised the heavy chain of SEQ ID NO: 19 or the K324A mutant variant thereof (SEQ ID NO: 18), to different Fc gamma receptors (CD16A, CD32A, CD32B, and CD64) was compared.

10

The K324A mutant was produced by site-directed mutagenesis. The binding was measured in a FACS based bioassay to Chinese Hamster Ovary (CHO) cells stably expressing full length human CD16A, CD32A, CD32B, or CD64. Both antibodies were incubated with each cell line at 7 different concentrations for one hour followed by washing. Receptor-bound H6 or H67 was detected with fluorochrome conjugated-anti-Fab'. Binding capacity was measured by FACS and the Kd and Bmax were calculated by non-linear regression.

15

20 Results: Both antibodies showed comparable binding to all receptors.

8. Binding to Clq

The binding of two antibodies, which either comprised the heavy chain of SEQ ID NO: 19 or the K324A mutant variant thereof (SEQ ID NO: 18), to Clq was compared in order to better characterize the effector functions of the antibodies.

25

A number of assay formats of binding of the antibodies to Clq was tested, including

30

- a) direct binding of the two antibodies to the plate and then binding to Clq in solution; and
- b) streptavidin coated plates first incubated with biotinylated pE-A β peptide, binding to antibodies and then Clq.

35

In summary, format a) produced best results. The procedure is summarized below:

The ELISA plate was coated with the antibody, comprising the heavy chain of SEQ ID NO: 19, the K324A mutant variant thereof (SEQ ID NO: 18), and a K324A control that does not bind Clq at 10, 8, 6, 5 4, 3, 2, 1 and 0 µg/ml in triplicate and incubated at 4°C overnight. Next day, the plate was washed three times with 1x PBS and then blocked with 1% BSA in 1x PBS at 50µl/well. Clq (Sigma, Cat.# C1740) was added to each well at 2µg/ml in blocking buffer and incubated for 1hour at room temperature. The plate was then washed 10 three times with 200µl of 1x PBS. Anti-Clq-HRP (Thermo, Cat. # PA1-84324) was added to the plate to detect the binding at a 1:250 dilution in blocking buffer (50µl/well) for 1hour. The plate was washed again three times with 200µl of 1x PBS. 50 µl of TMB (Invitrogen, Cat.# 002023) was added to each well to visualize the 15 interaction (Invitrogen, Cat.# 002023) for 2 min. 50µl of stop solution ((1M Sulfuric Acid) was added to each well before reading the absorbance at 450nm.

Results: The antibody, which comprised the wild-type heavy chain 20 of SEQ ID NO: 19 did bind to Clq. The K324A mutant variant thereof (comprising the heavy chain of SEQ ID NO: 18), did not bind to Clq.

9. Immunohistochemistry

25 With IHC the antigen Aβ N3pE can be localized in cerebral tissue sections. Therefore the antibodies of the invention were used for detection of Aβ N3pE.

For the IHC human cerebral tissue sections of the hippocampus and the frontal cortex from AD patients and furthermore cerebral tissue 30 sections of hippocampus from existing animal models for Alzheimer's disease as described herein can be used. These mouse models show increased brain Aβ levels followed by development of neuritic plaques. The tissue sections were paraffin-embedded and serial cut. The sections were stained with hematoxylin to colored 35 nuclei of cells and then immunostained with the anti Aβ N3pE antibodies of the invention. The tissue section preparation and staining were performed in accordance with standard methodology.

10. Treatment of Alzheimer mice in vivo

A total of 62 male mice were utilized in this study. Prior to the start of immunization, four mice of an existing mouse model for Alzheimer's disease (avg. 5.6 mo $\pm$ 0.45) mice were sacrificed as baseline controls to assess cerebral A β plaque burden at the commencement of treatment. The remaining mice were divided into four groups and received the following treatment: 250 μ l sterile PBS (n= 12; avg. 5.89 mo $\pm$ 0.13), 200 μ g of an antibody of the invention. A group of age- and gender-matched Wt littermates were injected with 250 μ l PBS (n=12; avg. 5.80 mo $\pm$ 0.12) and served as behavioral controls. Mice were treated with a total volume of 250 μ l (antibody or PBS) via intraperitoneal injection for 28 weeks.

15 Euthanasia and tissue preparation

Mice were euthanized, perfused and plasma harvested at 6 months (baseline) or 13 months of age. The brain was extracted and divided sagittally. The hippocampus, cortex and cerebellum were dissected from one hemisphere and snap frozen for biochemical analyses. The other hemisphere was drop-fixed in 4% paraformaldehyde (Electron Microscopy Sciences) for 24 h at 4°C, cryoprotected in graded sucrose solutions at 4°C and embedded in OCT compound (Tissue Tek).

Claims

1. An antibody or a functional variant thereof, wherein the variable part of the light chain of said antibody comprises,
5 consists essentially of or consists of an amino acid sequence of:

DVVMTQSPLSLPVTLGQPASISCKSSQSLLYSDGKTYLNWFQQRPGQSPRRLTYLVSKLDSGVP
DRFSGSGSGTDFTLKISRVEAEDVGVYYCVQGTHFPFTFGGGTKVEIK (SEQ ID NO: 1),

10 and

wherein the variable part of heavy chain of said antibody comprises, consists essentially of or consists of an amino acid sequence of:

15

QQQLVQSGAEVVKPGASVKVSKASGYSFTGHTMNWVRQAPGQGLEWMGLINPSDGVTRYNQKF
QGRVTITRDTSTTTVHMEITSLTSEDATATYYCTREAKREWDETYWGQGTILVTVSS (SEQ ID
NO: 2).

20 2. The antibody of claim 1, comprising the CDR regions:

V_L CDR1: SSQSLLYSDGKTYLN (SEQ ID NO: 3);

V_L CDR2: LVSKLDS (SEQ ID NO: 4); and

V_L CDR3: VQGTHFP (SEQ ID NO: 5)

25 in the light chain.

3. The antibody of claim 1, comprising the CDR regions:

V_H CDR1: GYSFTGHTMN (SEQ ID NO: 6);

30 V_H CDR2: LINPSDGVTRYNQKFQG (SEQ ID NO: 7); and

V_H CDR3: EAKREWDETY (SEQ ID NO: 8)

in the heavy chain.

35 4. The antibody according to any one of the preceding claims, wherein the light chain comprises, consists essentially of or consists of the amino acid sequence of:

DVVMTQSPLSLPVTLGQPASISCKSSQSLLYSDGKTYLNWFQQRPGQSPRRLTYLVSKLDSGVP
DRFSGSGSGTDFTLKISRVEAEDVGVYYCVQGTHFPFTFGGGTKVEIKRTVAAPSVFIFPPSDE
QLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYLSSTLTLSKADYE
5 KHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 17).

5. The antibody according to any one of the preceding claims,
wherein the heavy chain comprises, consists essentially of or
consists of the amino acid sequence of:

10 QVQLVQSGAEVVKPGASVKVSCKASGYSFTGHTMNWVRQAPGQGLEWMGLINPSDGVTRYNQKF
QGRVTITRDTSTTTVHMELTSLTSEDATYYCTREAKREWDETYWGQGTLTVTVSSASTKGPSVF
PLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS
SSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMIS
15 RTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE
YKCAVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE
SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVSMHEALHNHYTQKSLSLSPG
K (SEQ ID NO: 18).

20 6. The antibody according to any one of the preceding claims,
wherein the heavy chain comprises, consists essentially of or
consists of the amino acid sequence of:

25 QVQLVQSGAEVVKPGASVKVSCKASGYSFTGHTMNWVRQAPGQGLEWMGLINPSDGVTRYNQKF
QGRVTITRDTSTTTVHMELTSLTSEDATYYCTREAKREWDETYWGQGTLTVTVSSASTKGPSVF
PLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS
SSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMIS
RTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE
YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE
30 SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVSMHEALHNHYTQKSLSLSPG
K (SEQ ID NO: 19).

7. The antibody according to any one of the preceding claims,
wherein said humanized antibody specifically binds to the
35 pyroglutamate carrying N-terminus of the A β N3pE epitope.

8. The antibody according to any one of the preceding claims, wherein said humanized antibody binds specifically to an epitope selected from the group consisting of

5 pEFRHDSGYEVHHQKLV (SEQ ID NO: 23),
 pEFRHDSGYEVHHQKL (SEQ ID NO: 24),
 pEFRHDSGYEVHHQK (SEQ ID NO: 25),
 pEFRHDSGYEVHHQ (SEQ ID NO: 26),
 pEFRHDSGYEVHH (SEQ ID NO: 27),
10 pEFRHDSGYEVH (SEQ ID NO: 28),
 pEFRHDSGYEV (SEQ ID NO: 29),
 pEFRHDSGYE (SEQ ID NO: 30),
 pEFRHDSGY (SEQ ID NO: 31),
 pEFRHDSG (SEQ ID NO: 32),
15 pEFRHDS (SEQ ID NO: 33),
 pEFRHD (SEQ ID NO: 34)
 pEFRH (SEQ ID NO: 35), and
 pEFR (SEQ ID NO: 36).

20

9. The antibody according to any one of claims 1 to 6, wherein said humanized antibody of the invention binds to an A β N3pE variant, wherein the A β N3pE variant is defined as pE-A β _{3-x}, wherein x is defined as an integer between 19 and 42.

25

10. The antibody of claim 9, wherein the A β N3pE variant is selected from:

 pE-A β ₃₋₃₈,
30 pE-A β ₃₋₄₀,
 pE-A β ₃₋₄₂.

11. The antibody according to any one of the preceding claims, wherein said antibody does not bind to epitopes that do not carry
35 a pyroglutamate at the N-terminus.

12. A pharmaceutical composition comprising the antibody according to any one of the preceding claims.

13. The pharmaceutical composition of claim 12, further comprising
5 a further biologically active substance and/or a pharmaceutically acceptable carrier and/or a diluent and/or an excipient.

14. The pharmaceutical composition of claim 13, wherein said
10 further biologically active substance is selected from neutron-transmission enhancers, psychotherapeutic drugs, acetylcholine esterase inhibitors, calcium-channel blockers, biogenic amines, benzodiazepine tranquillizers, acetylcholine synthesis, storage or release enhancers, acetylcholine postsynaptic receptor agonists, monoamine oxidase-A or -B inhibitors, N-methyl- D-aspartate
15 glutamate receptor antagonists, non-steroidal anti-inflammatory drugs, antioxidants, and serotonergic receptor antagonists.

15. The pharmaceutical composition of claim 14, wherein said
20 further biologically active substance is selected from the group consisting of compounds effective against oxidative stress, anti-apoptotic compounds, metal chelators, inhibitors of DNA repair such as pirenzepin and metabolites, 3- amino-1-propanesulfonic acid (3 APS), 1,3-propanedisulfonate (1,3PDS), α -secretase activators, β - and γ -secretase inhibitors, tau proteins,
25 neurotransmitter, β -sheet breakers, attractants for amyloid beta clearing / depleting cellular components, inhibitors of N-terminal truncated amyloid beta including pyroglutamated amyloid beta 3-42, such as inhibitors of glutaminy cyclase, anti-inflammatory molecules, or cholinesterase inhibitors (ChEIs) such as tacrine,
30 rivastigmine, donepezil, and/or galantamine, M1 agonists and other drugs including any amyloid or tau modifying drugs and nutritive supplements, cholinesterase inhibitors (ChEIs), memantine or glutaminy cyclase inhibitors.

35 16. The antibody or the pharmaceutical composition according to any one of the preceding claims for use in the diagnosis,

treatment of amyloidosis, especially for use in the treatment of a neurodegenerative disease selected from the group consisting of mild cognitive impairment (MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome and clinical or pre-clinical cerebral amyloid angiopathy; and/or the treatment of a condition selected from prodromal AD, mild AD, moderate AD and severe AD and/or the slowing of cognitive decline in a patient diagnosed with a condition selected from clinical or pre-clinical Alzheimer's disease, Down's syndrome, and clinical or pre-clinical cerebral amyloid angiopathy.

17. The antibody or the pharmaceutical composition for use according to claim 24, wherein the administration of said antibody or said pharmaceutical composition leads to the reversion of cognitive decline, improvement of cognition or the prevention of cognitive decline in a subject diagnosed with mild cognitive impairment (MCI), Alzheimer's disease (AD), like for instance sporadic Alzheimer's disease (SAD) or Familial Alzheimer's dementias (FAD) like Familial British Dementia (FBD) and Familial Danish Dementia (FDD), neurodegeneration in Down Syndrome and clinical or pre-clinical cerebral amyloid angiopathy and/or of a condition selected from prodromal AD, mild AD, moderate AD and severe AD; or

wherein the administration of said antibody or said pharmaceutical composition leads to the clearance or removal of A β N3pE from plaques or other biological complexes; or

wherein the administration of said antibody or said pharmaceutical composition leads to the reduction of the plaque load and/or the removal of complete plaques from the affected tissue, such as brain tissue.