

PHARMACEUTICAL FORMULATION OF N-[5-[2-(3,5-DIMETHOXYPHENYL)ETHYL]-2H-PYRAZOL-3-YL]-4-[(3R,5S)-3,5-DIMETHYLPIPERAZIN-1-YL]BENZAMIDE


FIELD OF THE INVENTION

The present invention relates to pharmaceutical/formulation chemistry. The invention is understood to apply generally to formulations of compounds which contain an increased percent loading of the active ingredient. As a preferred aspect, provided herein are formulations of N-[5-[2-(3,5-dimethoxyphenyl)ethyl]-2H-pyrazol-3-yl]-4-[(3R,5S)-3,5-dimethylpiperazin-1-yl]benzamide (Compound I) which exhibit satisfactory manufacturability, stability and *in vitro* dissolution. The formulations are useful for treating of cancer.

BACKGROUND OF THE INVENTION

In the manufacture of pharmaceutical formulations for oral administration, it may be desirable for the drug to dissolve rapidly soon after administration. However, it is known that certain physic-chemical properties of the drug, such as particle size, wettability, or solubility, may lead to a pharmaceutical formulation which exhibits unsatisfactory and/or variable dissolution or to a formulation which exhibits unsatisfactory and/or variable bioavailability. Such formulations may be unsuitable for use by patients.

Compound I (below) is disclosed in international patent application WO2008/075068.



Compound I

for use in the treatment of cancer.


Javaid et al (J. Pharm. Sci. 61 (9) 1972 pp 1370-1373) studied the effect of various
5 classes of buffering agents on the dissolution of aspirin from tablet formulations.

Compound I is currently in clinical studies for the treatment of cancer, in particular cancers
of the lung, breast, gastric and bladder. Dosing is currently done with orally delivered
tablets with tablet strengths of 20 and 100mg. These tablets exhibit satisfactory dissolution
across the physiological pH range. However, the manufacturing process used for clinical
10 batches cannot be operated at commercial scale, due to a high incidence of filming which
cannot be corrected using conventional means. It is desirable, therefore, to produce
new pharmaceutical formulations of Compound I which overcome at least in part the
above problems.

15 DESCRIPTION OF THE INVENTION

This invention is generally directed to formulations of compounds with improved
manufacturability, in particular to formulations which contain an alkaline effervescent
excipient and which exhibit satisfactory dissolution across the physiological pH range.

The compound of formula (I) (known hereafter as "Formula (I)") is shown below:



20 **Formula (I)**

The compound of Formula (I) is a base and exhibits pH dependent solubility,
having a solubility in simulated gastric fluid (pH1.2) of approximately 5mg/mL ('slightly
25 soluble', using the definition given in the United States Pharmacopeia / National
Formulary, USP35-NF30), which reduces in fasted simulated intestinal fluid (pH 6.5) to
approximately 0.25mg/mL ('very slightly soluble' by the USP definition). Furthermore, we

have observed that the compound of Formula (I) may form a viscous material at low pH, which has the effect of reducing the rate at which the drug dissolves. In order to achieve an acceptable rate and extent of dissolution across the physiological pH range, the earlier clinical formulation was manufactured using conditions designed to give a fine granule which, when compressed into tablet form would disperse rapidly upon administration. While this approach did result in an improvement in dissolution performance, 'filming' problems were experienced during manufacture. In addition, it was observed that the use of conventional lubricants such as magnesium stearate and sodium stearyl fumarate led to chemical degradation including impurity formation and / or complexation.

It was unexpectedly found that alkaline effervescent excipients were effective in both improving the rate and extent of dissolution at low pH, despite the reduction in solubility under alkaline conditions, and in ameliorating the "filming" issues. A further unexpected finding was that the use of an alternative lubricant, glyceryl dibehenate, was effective in ameliorating chemical degradation.

In particular, this invention is directed at least in part to the unexpected result that the use of an alkaline effervescent excipient with Formula (I) in the formulation allows the manufacture of tablets with improved manufacturability and / or a satisfactory dissolution across the physiological pH range; and, at least in part, to the unexpected result that an alternative lubricant allows the manufacture of tablets with improved stability.

In a further aspect, this invention provides the use of an alkaline effervescent excipient with Formula (I) in the formulation allowing the manufacture of tablets with improved manufacturability and/or a satisfactory dissolution across the physiological pH range.

In a still further aspect, this invention provides the use of magnesium carbonate with Formula (I) in the formulation allowing the manufacture of tablets with improved manufacturability and/or a satisfactory dissolution across the physiological pH range.

In a still further aspect, this invention provides the use of calcium carbonate with Formula (I) in the formulation allowing the manufacture of tablets with improved manufacturability and/or a satisfactory dissolution across the physiological pH range.

In a still further aspect, this invention provides the use of sodium bicarbonate with Formula (I) in the formulation allowing the manufacture of tablets with improved manufacturability and/or a satisfactory dissolution across the physiological pH range.

In a further aspect, this invention provides the use of an alternative lubricant with Formula (I) in the formulation allowing the manufacture of tablets with improved stability.

In a still further aspect, this invention provides the use of glyceryl dibehenate with Formula (I) in the formulation allowing the manufacture of tablets with improved stability.

5 In a further aspect of the invention, there is provided a pharmaceutical composition comprising greater than 10% w/w of Formula (I) and an amount of an alkaline effervescent excipient that is sufficient to provide satisfactory *in vitro* dissolution; and further comprising one or more pharmaceutically acceptable ingredients.

In another aspect of the invention, there is provided a pharmaceutical composition
10 in unit dosage form comprising from 10 mg to 200 mg of Formula (I) (for example 20 mg, 30 mg, 40 mg, 50 mg, 60 mg, 70mg, 80 mg, 90 mg, 100 mg, 110 mg, 120 mg, 130 mg, 140 mg, 150 mg, 160 mg, 170mg, 180mg, 190mg or 200mg) and an amount of an alkaline effervescent excipient that is sufficient to provide satisfactory *in vitro* dissolution; and further comprising one or more pharmaceutically acceptable ingredients. For the avoidance
15 of doubt, each of the previous integers represents a separate and independent aspect of the invention.

In another aspect of the invention a unit dosage form of the pharmaceutical composition comprises between about 10mg to about 160 mg of Formula (I).

In another aspect of the invention a unit dosage form of the pharmaceutical
20 composition comprises between about 10 mg to about 140 mg of Formula (I).

In a still further aspect, a unit dosage form of the pharmaceutical composition comprises between about 10mg to about 130 mg of Formula (I).

In a yet further aspect, a unit dosage form of the pharmaceutical composition comprises between about 15 mg to about 110 mg of Formula (I).

25 In a specific aspect of the invention, a unit dosage form of the pharmaceutical composition comprises 20 mg $\pm$ 1 mg of Formula (I).

In a further specific aspect of the invention, a unit dosage form of the pharmaceutical composition comprises 80 mg $\pm$ 4 mg of Formula (I).

In a further specific aspect of the invention, a unit dosage form of the
30 pharmaceutical composition comprises 100 mg $\pm$ 5 mg of Formula (I).

In a further specific aspect of the invention, a unit dosage form of the pharmaceutical composition comprises 160 mg $\pm$ 8 mg of Formula (I).

In a further specific aspect of the invention, a unit dosage form of the pharmaceutical composition comprises 200 mg $\pm$ 10 mg of Formula (I).

In another aspect of the invention the pharmaceutical composition comprises between 10% w/w to 60% w/w of Formula (I).

5 In a further aspect, the pharmaceutical composition comprises between 15% w/w to 50% w/w of Formula (I).

In a still further aspect, the pharmaceutical composition comprises between 15% w/w to 45% w/w of Formula (I).

10 In a still further aspect, the pharmaceutical composition comprises between 15% w/w to 40% w/w of Formula (I).

In a still further aspect, the pharmaceutical composition comprises between 15% w/w to 25% w/w of Formula (I).

In another aspect of the invention the pharmaceutical composition comprises about 20% w/w of Formula (I).

15 In a specific aspect of the invention, the pharmaceutical composition comprises 21.33% $\pm$ 5% w/w of Formula (I).

In a further aspect, the pharmaceutical composition comprises between 20.26% w/w to 22.40% w/w of Formula (I).

20 In a still further aspect of the invention, the pharmaceutical composition comprises from 1% w/w to 50% w/w of an alkaline effervescent excipient

In a still further aspect of the invention, the pharmaceutical composition comprises from 1% w/w to 40% w/w of an alkaline effervescent excipient

In a further aspect, the pharmaceutical composition comprises from 10% w/w to 30% w/w of an alkaline effervescent excipients.

25 In a still further aspect, the pharmaceutical composition comprises about 20% w/w of an alkaline effervescent excipients.

In a further aspect, the pharmaceutical composition comprises from 15% w/w to 20% w/w of an alkaline effervescent excipients.

30 In a still further aspect of the invention, the pharmaceutical composition comprises less than or equal to 6% w/w of a conventional lubricant

Alternatively the use of an alternative lubricant may improve stability.

In a still further aspect of the invention, the pharmaceutical composition comprises less than or equal to 5% w/w of an alternative lubricant

In a still further aspect of the invention, the pharmaceutical composition comprises less than or equal to 4% w/w of an alternative lubricant

5 In a further aspect, the pharmaceutical composition comprises less than or equal to 3% w/w of an alternative lubricant.

In a still further aspect of the invention, the pharmaceutical composition comprises between 0.25% w/w and 8% w/w of an alternative lubricant.

10 In a still further aspect of the invention, the pharmaceutical composition comprises between 0.5% w/w and 5% w/w of an alternative lubricant.

In a still further aspect of the invention, the pharmaceutical composition comprises between 1% w/w and 4% w/w of an alternative lubricant.

In a still further aspect of the invention, the pharmaceutical composition comprises between 2.5% w/w and 3.5% w/w of an alternative lubricant.

15 In a further aspect, the pharmaceutical composition comprises about 3% w/w of an alternative lubricant.

In a further aspect of the invention, there is provided a pharmaceutical composition comprising greater than 10% w/w of Formula (I) and less than or equal to 50% w/w of an alkaline effervescent excipient ; and further comprising one or more pharmaceutically acceptable ingredients.

20 In a further aspect of the invention, there is provided a pharmaceutical composition comprising greater than 15% w/w of Formula (I) and less than or equal to 40% w/w of an alkaline effervescent excipient ; and further comprising one or more pharmaceutically acceptable ingredients.

25 In a further aspect of the invention, there is provided a pharmaceutical composition comprising greater than 15% w/w of Formula (I) and less than or equal to 30% w/w of an alkaline effervescent excipient; and further comprising one or more pharmaceutically acceptable ingredients.

30 In a further aspect of the invention, there is provided a pharmaceutical composition comprising greater than 15% w/w of Formula (I) and less than or equal to 20% w/w of an alkaline effervescent excipient; and further comprising one or more pharmaceutically acceptable ingredients.

In a further aspect of the invention, there is provided a pharmaceutical composition comprising from 10% w/w to 50%w/w of Formula (I) and from 1% w/w to 50% w/w of an alkaline effervescent excipient ; and optionally further comprising one or more pharmaceutically acceptable ingredients.

5 In a further aspect of the invention, there is provided a pharmaceutical composition comprising from 15% w/w to 35% w/w of Formula (I) and from 10% to 40% w/w of an alkaline effervescent excipient ; and further comprising one or more pharmaceutically acceptable ingredients.

10 In a further aspect of the invention, there is provided a pharmaceutical composition comprising from 15% w/w to 25% w/w of Formula (I) and from 15% w/w to 25% w/w of an alkaline effervescent excipient; and further comprising one or more pharmaceutically acceptable ingredients.

15 In a further aspect of the invention, there is provided a pharmaceutical composition comprising about 20% w/w of Formula (I) and about 20% w/w of an alkaline effervescent excipient; and further comprising one or more pharmaceutically acceptable ingredients.

In a further aspect of the invention, there is provided a unit dosage form comprising from 15% w/w to 45% w/w of Formula (I) and from 10% to 40% w/w of an alkaline effervescent excipients and further comprising one or more pharmaceutically acceptable ingredients, wherein the unit comprises from 10 to 200 mg of Formula (I)

20 In a further aspect of the invention, there is provided a unit dosage form comprising from 15% w/w to 40% w/w of Formula (I) and from 10% to 40% w/w of an alkaline effervescent excipients and further comprising one or more pharmaceutically acceptable ingredients, wherein the unit comprises from 10 to 200 mg of Formula (I).

25 In a further aspect of the invention, there is provided a unit dosage form comprising from 15% w/w to 25% w/w of Formula (I) and from 15% to 25% w/w of an alkaline effervescent excipients and further comprising one or more pharmaceutically acceptable ingredients, wherein the unit comprises 20 mg of Formula (I).

30 In a further aspect of the invention, there is provided a unit dosage form comprising from 15% w/w to 25% w/w of Formula (I) and from 15% to 25% w/w of an alkaline effervescent excipients and further comprising one or more pharmaceutically acceptable ingredients, wherein the unit comprises 80 mg of Formula (I).

In a further aspect of the invention, there is provided a unit dosage form comprising from 15% w/w to 25% w/w of Formula (I) and from 15% to 25% w/w of an alkaline effervescent excipients and further comprising one or more pharmaceutically acceptable ingredients, wherein the unit comprises 160 mg of Formula (I).

5

In a further aspect of the invention, optional ingredients which can be added to the pharmaceutical composition include one or more of the following:

- a) fillers;
- b) binding agents;
- 10 c) lubricants; and
- d) disintegrants.

Where optional ingredients are added to make up the remainder of the pharmaceutical composition, the remainder may optionally include one or more of the following:

- 15 a) fillers which, when employed, range between for example about 10 to about 75 weight percent (e.g. about 15 to about 70 weight percent) of the remainder of the dry formulation;
- b) binding agents which, when employed range between for example about 2 to about 8 weight percent of the remainder of the dry formulation;
- 20 c) lubricants which, when employed, range from between about 0.25 and 5 weight percent of the remainder of the dry formulation; and
- d) disintegrants which, when employed, range from between about 0.5 and 10.0 weight percent (e.g. about 5 weight percent) of the remainder of the dry formulation.

25 In a further aspect of the invention, the pharmaceutical composition further comprises one or more additional ingredients independently selected from, for example

- a) fillers such as mannitol (e.g. Pearlitol 50c, Peralitol 120c or Pearlitol 160c) or microcrystalline celluloses (e.g. MCC Avicel PH 101, Emcocel 90M, etc.);
- b) binding agents such as Plasdane K29/32, Povidone, microcrystalline celluloses
30 or Kollidon K30;
- c) lubricants such as glyceryl dibehenate; and

d) disintegrants such as sodium starch glycolate, for example ExploTab or Glycolys LV.

In a further aspect of the invention, there is provided a pharmaceutical composition comprising from 15% w/w to 25% w/w of Formula (I), from 15% w/w to 25% w/w of an
5 alkaline effervescent excipients, from 2.5% w/w to 3.5% w/w of an alternative lubricant; and further comprising from 40% w/w to 60% filler w/w, from 1% w/w to 3% w/w binder and 5% w/w to 9% w/w disintegrant.

In another aspect of the invention, there is provided a tablet comprising greater than 10% w/w of Formula (I) and an amount of an alkaline effervescent excipient that is
10 sufficient to provide satisfactory *in vitro* dissolution; and further comprising one or more pharmaceutically acceptable ingredients.

In another aspect of the invention, there is provided a tablet comprising from 10mg to 200 mg of Formula (I) (for example 20 mg, 30 mg, 40 mg, 50 mg, 60 mg, 70 mg, 80 mg, 90 mg, 100 mg, 110 mg, 120 mg, 130 mg, 140 mg, 150 mg, 160 mg, 170mg, 180mg
15 190mg or 200mg) and an amount of an alkaline effervescent excipient that is sufficient to provide satisfactory *in vitro* dissolution; and further comprising one or more pharmaceutically acceptable ingredients. For the avoidance of doubt, each of the previous integers represents a separate and independent aspect of the invention.

In another aspect of the invention, the tablet comprises between about 10 mg to
20 about 160 mg of Formula (I).

In another aspect of the invention the tablet comprises between about 10 mg to about 140 mg of Formula (I).

In a still further aspect, the tablet comprises between about 10mg to about 130 mg of Formula (I).

25 In a still further aspect, the tablet comprises between about 15 mg to about 110 mg of Formula (I).

In a specific aspect of the invention, the tablet comprises 20 mg $\pm$ 1 mg of Formula (I).

In a further specific aspect of the invention, the tablet comprises 80 mg $\pm$ 4 mg of
30 Formula (I).

In a further specific aspect of the invention, the tablet comprises 100 mg $\pm$ 5 mg of Formula (I).

In a further specific aspect of the invention, the tablet comprises 160 mg $\pm$ 8 mg of Formula (I).

In a further specific aspect of the invention, the tablet comprises 200 mg $\pm$ 10 mg of Formula (I).

5 In a still further aspect of the invention, the tablet comprises from 1% w/w to 50% w/w of an alkaline effervescent excipient

In a still further aspect of the invention, the tablet comprises from 1% w/w to 40% w/w of an alkaline effervescent excipient

10 In a further aspect, the tablet comprises from 10% w/w to 30% w/w of an alkaline effervescent excipients.

In a still further aspect, the tablet comprises about 20% w/w of an alkaline effervescent excipients.

In a specific aspect of the invention, the tablet comprises 21.33% $\pm$ 5% w/w of Formula (I).

15 In a further aspect, the tablet comprises between 20.26% w/w to 22.40% w/w of Formula (I).

In a still further aspect, the tablet comprises between about 15% w/w to about 25% w/w of Formula (I).

20 In a still further aspect of the invention, the tablet comprises less than or equal to 50% w/w of an alkaline effervescent excipient.

In a further aspect, the tablet comprises less than or equal to 40% w/w of an alkaline effervescent excipient.

In a further aspect, the tablet comprises less than or equal to 30% w/w of an alkaline effervescent excipient.

25 In a still further aspect, the tablet comprises less than or equal to 20% w/w of an alkaline effervescent excipient.

In a further aspect of the invention, the tablet comprises less than or equal to 6% w/w of a conventional lubricant

Alternatively the use of an alternative lubricant may improve stability.

30 In a still further aspect of the invention, the tablet comprises less than or equal to 5% w/w of an alternative lubricant

In a still further aspect of the invention, the tablet comprises less than or equal to 4% w/w of an alternative lubricant

In a further aspect, the tablet comprises less than or equal to 3% w/w of an alternative lubricant.

5 In a still further aspect of the invention, the tablet comprises between 0.25% w/w and 8% w/w of an alternative lubricant.

In a still further aspect of the invention, the tablet comprises between 0.5% w/w and 5% w/w of an alternative lubricant.

10 In a still further aspect of the invention, the tablet comprises between 1% w/w and 4% w/w of an alternative lubricant.

In a still further aspect of the invention, the tablet comprises between 2.5% w/w and 3.5% w/w of an alternative lubricant.

In a further aspect, the tablet comprises about 3% w/w of an alternative lubricant.

15 In a further aspect of the invention, there is provided a tablet comprising from 10% w/w to 50% w/w of Formula (I) and from 1% w/w to 50% w/w of an alkaline effervescent excipient ; and optionally further comprising one or more pharmaceutically acceptable ingredients.

In a further aspect of the invention, there is provided a tablet comprising from 15% w/w to 35% w/w of Formula (I) and from 10% to 40% w/w of an alkaline effervescent excipient ; and further comprising one or more pharmaceutically acceptable ingredients.

In a further aspect of the invention, there is provided a tablet comprising from 15% w/w to 25% w/w of Formula (I) and from 15% w/w to 25% w/w of an alkaline effervescent excipient; and further comprising one or more pharmaceutically acceptable ingredients.

25 In a further aspect of the invention, there is provided a tablet comprising about 20% w/w of Formula (I) and about 20% w/w of an alkaline effervescent excipient; and further comprising one or more pharmaceutically acceptable ingredients.

In a further aspect of the invention, there is provided a tablet comprising from 15% w/w to 45% w/w of Formula (I) and from 10% to 40% w/w of an alkaline effervescent excipients and further comprising one or more pharmaceutically acceptable ingredients, wherein the tablet comprises from 10 to 200 mg of Formula (I)

In a further aspect of the invention, there is provided a tablet comprising from 15% w/w to 40% w/w of Formula (I) and from 10% to 40% w/w of an alkaline effervescent excipients and further comprising one or more pharmaceutically acceptable ingredients, wherein the tablet comprises from 10 to 200 mg of Formula (I).

5 In a further aspect of the invention, there is provided a tablet comprising from 15% w/w to 25% w/w of Formula (I) and from 15% to 25% w/w of an alkaline effervescent excipients and further comprising one or more pharmaceutically acceptable ingredients, wherein the tablet comprises 20 mg of Formula (I).

10 In a further aspect of the invention, there is provided a tablet comprising from 15% w/w to 25% w/w of Formula (I) and from 15% to 25% w/w of an alkaline effervescent excipients and further comprising one or more pharmaceutically acceptable ingredients, wherein the tablet comprises 80 mg of Formula (I).

15 In a further aspect of the invention, there is provided a tablet comprising from 15% w/w to 25% w/w of Formula (I) and from 15% to 25% w/w of an alkaline effervescent excipients and further comprising one or more pharmaceutically acceptable ingredients, wherein the tablet comprises 160 mg of Formula (I).

20 The dosage forms of this invention may include one or more pharmaceutically acceptable excipients which may be selected, for example, from adjuvants, carriers, binders, lubricants, diluents, stabilising agents, buffering agents, emulsifying agents, viscosity-regulating agents, surfactants, preservatives, flavourings or colorants. It will be understood that an individual excipient may be multifunctional. Examples of pharmaceutically acceptable excipients are described in the Handbook of Pharmaceutical Excipients (Fifth Edition, 2005, edited by Ray C. Rowe, Paul J. Sheskey and Sian C. Owen, published by the American Pharmaceutical Association and the Pharmaceutical Press). As will be understood by those skilled in the art, the most appropriate method of administering the active ingredients is dependent on a number of factors.

25 It will be understood that the therapeutic dose of each active ingredient administered in accordance with the present invention will vary depending upon the particular active ingredient employed, the mode by which the active ingredient is to be administered, and the condition or disorder to be treated.

30 In a further aspect of the invention, optional ingredients which can be added to make up the remainder of the tablet include one or more of the following:

- a) fillers which, when employed, range between for example about 10 to about 75 weight percent (e.g. about 15 to about 70 weight percent) of the remainder of the tablet formulation;
- b) binding agents which, when employed range between for example about 2 to about 8 weight percent of the remainder of the tablet formulation;
- c) lubricants which, when employed, range from between about 0.25 and 3.5 weight percent of the remainder of the tablet formulation; and
- d) disintegrants which, when employed, range from between about 0.5 and 10.0 weight percent (e.g. about 5 weight percent) of the remainder of the tablet formulation.

In a further aspect of the invention, the tablet further comprises one or more additional ingredients independently selected from, for example:

- a) fillers such as mannitol (e.g. Pearlitol 50c, Peralitol 120c or Pearlitol 160c) or microcrystalline celluloses (e.g. MCC Avicel PH 101, Emcocel 90M, etc.);
- b) binding agents such as Plasdone K29/32, Povidone, microcrystalline celluloses or Kollidon K30;
- c) lubricants such as glyceryl dibehenate;
- d) disintegrants such as sodium starch glycolate, for example ExploTab or Glycolys LV;

In a further aspect of the invention, the tablet optionally further comprises a suitable coating, for example a film coating. A coating can be used to provide protection against, for example, moisture ingress or degradation by light, to colour the formulation, or to modify or control the release of Formula (I) from the formulation.

In a yet further aspect of the invention, the pharmaceutical composition comprises the following components by weight:

Composition A (mg)		Composition B (mg)	
Formula (I)	20.00	Formula (I)	80.00
Microcrystalline cellulose	14.06	Microcrystalline cellulose	56.25
Mannitol	29.22	Mannitol	116.87

Magnesium carbonate	18.75	Magnesium carbonate	75.00
Hydroxypropyl cellulose	1.88	Hydroxypropyl cellulose	7.50
Sodium starch glycollate	7.03	Sodium starch glycollate	28.13
Glyceryl dibehenate	2.81	Glyceryl dibehenate	11.25

5 In a yet further aspect of the invention, the pharmaceutical composition comprises the following components (% w/w):

Compositions A and B (% w/w)

Formula (I)	21.33
10 Microcrystalline cellulose	15.00
Mannitol	31.17
Magnesium carbonate	20.00
Hydroxypropyl cellulose	2.00
Sodium starch glycollate	7.50
15 Glyceryl dibehenate	3.00

In a still further aspect, the invention comprises a tablet formed from the pressing of composition A and/or composition B into tablet form.

In a further aspect of the invention, there is provided a process for the preparation of a pharmaceutical composition which process comprises the following steps:

20 Step A – comprises mixing Formula (I) with an alkaline effervescent excipient optionally in the presence of one or more pharmaceutically acceptable ingredients . In a further aspect, Step A is carried out in the presence of one or more additional fillers (such as mannitol) and optionally in the presence of one or more pharmaceutically acceptable ingredients. In a still further aspect, Step A is carried out in the presence of one or more additional fillers (such as mannitol) and optionally in the presence of one or more binding agents and/or one or more disintegrants.

25 Step B - comprises adding purified water and/or binder solution into the powder mixture from Step A above and mixing to form granules and optionally passing through a filter screen to break-up agglomerates. In a further aspect between about 10% and 45% by weight of purified water is added into the powder mixture.

Step C - comprises drying the granules produced in Step B above until an LOD of less than 10% (e.g. less than 5%) is achieved, to provide dried granules.

Step D – comprises optionally milling the dried granules from Step C.

Step E – optionally, comprises mixing the milled granules from Step D with an alkaline effervescent excipient.

In a further aspect of the invention there is provided a process for the preparation of a pharmaceutical composition which process (wet granulation process) comprises:

- a) blending Formula (I) with an effervescent agent, one or more additional fillers (such as mannitol) and optionally in the presence of one or more binding agents and/or one or more disintegrants and/or one or more other excipients;
- b) adding between about 10% and 45% by weight of purified water and/or binder solution into the powder mixture of a) above and mixing to form enlarged granules and optionally passing through a filter screen to break-up large agglomerates; and
- c) drying the enlarged granules produced in b) above until an LOD of less than 10% (e.g. less than 5%) is achieved, to provide dried granules.

Alternatively, in another aspect of the invention, there is provided a process for the preparation of a pharmaceutical composition which process comprises the following steps

Step A - mixing Formula (I) optionally with an alkaline effervescent excipient optionally in the presence of one or more pharmaceutically acceptable ingredients.

In a further aspect, Step A is carried out in the presence of one or more additional fillers (such as mannitol) and optionally in the presence of one or more pharmaceutically acceptable ingredients. In a still further aspect, Step A is carried out in the presence of one or more additional fillers (such as mannitol) and optionally in the presence of one or more binding agents and/or one or more disintegrants.

Step B - comprises adding purified water and/or binder solution into the powder mixture from Step A above and mixing to form granules and optionally passing through a filter screen to break-up agglomerates. Typically, between about 10% and 45% by weight of purified water and/or binder solution is added into the powder mixture.

Step C - comprises drying the granules produced in Step B above until an LOD of less than 10% (e.g. less than 5%) is achieved, to provide dried granules.

Step D – comprises milling the dried granules from Step C to give milled granules

Step E – comprises mixing the milled granules from Step D with an effective amount of an alkaline effervescent excipient.

In a further aspect of the invention there is provided a process for the preparation of a pharmaceutical composition which process (wet granulation process) comprises:

- a) blending Formula (I) with one or more additional fillers (such as mannitol) and optionally in the presence of one or more binding agents and/or one or more disintegrants and/or one or more other excipients;
- b) adding between about 10% and 45% by weight of purified water and/or binder solution into the powder mixture of a) above and mixing to form granules and optionally passing through a filter screen to break-up agglomerates;
- c) drying the granules produced in b) above until an LOD of less than 10% (e.g. less than 5%) is achieved, to provide dried granules;
- d) milling the dried granules produce in c) to give milled granule; and
- e) mixing the milled granules from d) with an alkaline effervescent excipient.

In another of its method aspects, this invention further comprises milling the dried granules. In one aspect, the dried granules are milled so that about 90 weight percent have a particle size between about 25 μm to about 3500 μm in diameter.

In yet another aspect, the dried, milled, granules are mixed with a conventional and/or alternative lubricant, and then the resulting pharmaceutical composition is tableted. Conventional and alternative lubricants include glyceryl dibehenate, sodium stearyl fumarate, magnesium stearate, colloidal silica and talc.

In a further aspect of the invention, the alternative lubricant (such as glyceryl dibehenate) can be added to the dry granules prior to milling, and then the resulting pharmaceutical composition is milled and then tableted.

In another aspect, this invention provides a wet granulated formulation comprising between 10% w/w to 60% w/w of Formula (I) and an amount of an alkaline effervescent

excipient that is sufficient to provide satisfactory *in vitro* dissolution; and further comprising one or more pharmaceutically acceptable ingredients.

In another aspect of the invention the wet granulated formulation comprises between 15% w/w to 50% w/w of Formula (I).

5 In a further aspect, the wet granulated formulation comprises between 15% w/w to 40% w/w of Formula (I).

In a further aspect, the wet granulated formulation comprises between 15% w/w to 25% w/w of Formula (I).

10 In another aspect of the invention the wet granulated formulation comprises about 20% w/w of Formula (I).

In a specific aspect of the invention, the wet granulated formulation contains 21.33% $\pm$ 5% w/w of Formula (I).

In a further aspect, the wet granulated formulation comprises between 20.26% w/w to 22.40% w/w of Formula (I).

15 In a still further aspect of the invention, the wet granulated formulation comprises from 1% w/w to 50% w/w of an alkaline effervescent excipient.

In a further aspect, the wet granulated formulation comprises from 1% w/w to 40% w/w of an alkaline effervescent excipient.

20 In a further aspect, the wet granulated formulation comprises from 10% w/w to 30% w/w of an alkaline effervescent excipient

In a still further aspect, the wet granulated formulation comprises from 15% w/w to 25% w/w of an alkaline effervescent excipient

In a still further aspect, the wet granulated formulation comprises about 20% w/w of an alkaline effervescent excipient

25 In a further aspect of the invention, there is provided a wet granulation formulation comprising from 10% w/w to 50% w/w of Formula (I) and from 1% w/w to 50% w/w of an effervescent agent; and further comprising one or more pharmaceutically acceptable ingredients.

30 In a further aspect of the invention, there is provided a wet granulation formulation comprising greater from 10% w/w to 45% w/w of Formula (I) and from 10% w/w to 45% w/w of an alkaline effervescent excipient; and further comprising one or more pharmaceutically acceptable ingredients.

In a further aspect of the invention, there is provided a wet granulation formulation comprising from 15% w/w to 25% w/w of Formula (I) and from 15% to 25% w/w of an alkaline effervescent excipient; and further comprising one or more pharmaceutically acceptable ingredients.

5 In a further aspect of the invention, there is provided a wet granulation formulation comprising about 20% w/w of Formula (I) and about 20% w/w of an alkaline effervescent excipient; and further comprising one or more pharmaceutically acceptable ingredients.

In another aspect of the invention the wet granulation formulation comprises Formula (I), water, an alkaline effervescent excipient, additional filler(s), binding agent(s)
10 and disintegrant(s).

In another aspect, this invention provides a tablet formed by compressing the wet granulated formulation.

In a further aspect of the invention, there is provided a further process for the preparation of a pharmaceutical composition as defined above which process comprises
15 passing the mixture of Step A above through a compactor to produce dry granules (Step D).

In a further aspect of the present invention there is provided a process for the manufacture of a pharmaceutical composition which process (roller compaction process) comprises:

20 (a) blending Formula (I) with an alkaline effervescent excipient, one or more additional fillers (such as mannitol) and optionally in the presence of one or more binding agents and/or one or more disintegrants and/or one or more other excipients;

(b) passing the mixture of (a) above through a compactor to produce dry granules.

25 In another of its method aspects, this invention further comprises milling the dried granules. In one aspect, the dried granules are milled so that about 90 weight percent have a particle size between about 25 μm to about 3500 μm in diameter.

In yet another aspect, the dried, milled, granules are mixed with a lubricant, and then the resulting pharmaceutical composition is tableted. Suitable lubricants include
30 glyceryl dibehenate, sodium stearyl fumarate, magnesium stearate, colloidal silica and talc.

In yet another aspect of the present invention there is provided a process for the manufacture of a pharmaceutical composition which process (roller compaction process) comprises:

- 5 (a) blending Formula (I) with an alkaline effervescent excipient, one or more additional fillers (such as mannitol) and optionally in the presence of one or more binding agents and/or one or more disintegrants and/or one or more other excipients;
- (b) passing the mixture of (a) above through a compactor to produce dry granules.

10 In another of its method aspects, this invention further comprises milling the dried granules. In one aspect, the dried granules are milled so that about 90 weight percent have a particle size between about 25 μm to about 3500 μm in diameter.

In yet another aspect of the present invention there is provided a process for the manufacture of a pharmaceutical composition which process (roller compaction process) comprises:

- 15 (a) blending Formula (I) optionally with an alkaline effervescent excipient, one or more additional fillers (such as mannitol) and optionally in the presence of one or more binding agents and/or one or more disintegrants and/or one or more other excipients;
- (b) passing the mixture of (a) above through a compactor to produce dry granules.

20 The dried, milled, granules are then mixed with an alkaline effervescent excipient.

In yet another aspect, the dried, milled, granules are mixed with a lubricant, and then the resulting pharmaceutical composition is tableted. Suitable lubricants include glyceryl dibehenate, sodium stearyl fumarate, magnesium stearate, colloidal silica and talc.

25 In an alternative aspect of the invention, the lubricant (such as glyceryl dibehenate) can be added to the dry granules prior to milling, and then the resulting pharmaceutical composition is milled and then tableted.

In another aspect, this invention provides a roller compaction formulation comprising greater than 10% w/w of Formula (I) and an amount of an alkaline effervescent excipient that is sufficient to provide satisfactory *in vitro* dissolution; and further
30 comprising one or more pharmaceutically acceptable ingredients.

In another aspect of the invention the roller compaction formulation comprises between 10% w/w to 60% w/w of Formula (I).

In a further aspect, the roller compaction formulation comprises between 15% w/w to 50% w/w of Formula (I).

In a still further aspect, the roller compaction formulation comprises between 15% w/w to 45% w/w of Formula (I).

5 In a still further aspect, the roller compaction formulation comprises between 15% w/w to 40% w/w of Formula (I).

In a still further aspect, the roller compaction formulation comprises between 15% w/w to 25% w/w of Formula (I).

10 In another aspect of the invention the roller compaction formulation comprises about 20% w/w of Formula (I).

In a specific aspect of the invention, the roller compaction formulation contains 21.33% $\pm$ 5% w/w of Formula (I).

In a further aspect, the roller compaction formulation contains 20.26% w/w to 22.40% w/w of Formula (I).

15 In a still further aspect of the invention, the roller compaction formulation comprises from 1% w/w to 50% w/w of an alkaline effervescent excipient.

In a further aspect, the roller compaction formulation comprises from 1% w/w to 40% w/w of an alkaline effervescent excipient.

20 In a still further aspect, the roller compaction formulation comprises from 10% w/w to 30% w/w of an alkaline effervescent excipient.

In a still further aspect, the roller compaction formulation comprises from 15% w/w to 25% w/w of an alkaline effervescent excipient.

In a still further aspect, the roller compaction formulation comprises about 20% w/w of an alkaline effervescent excipient.

25 In a further aspect of the invention, there is provided a roller compaction formulation comprising from 15% w/w to 45% of Formula (I) and from 10% w/w to 40% w/w of an alkaline effervescent excipient; and further comprising one or more pharmaceutically acceptable ingredients.

30 In a further aspect of the invention, there is provided a roller compaction formulation comprising from 15% w/w to 25% w/w of Formula (I) and from 15% w/w to 25% w/w of an alkaline effervescent excipient; and further comprising one or more pharmaceutically acceptable ingredients.

In another aspect of the invention the roller compaction formulation comprises Formula (I), an alkaline effervescent excipient, additional filler(s), binding agent(s) and disintegrant(s).

5 In another aspect, this invention provides a tablet formed by compressing the roller compaction formulation.

In a further aspect of the invention there is provided a process for the manufacture of a pharmaceutical composition which process (direct compression process) comprises:

- (a) blending Formula (I) with an alkaline effervescent excipient, one or more additional fillers (such as mannitol) and optionally in the presence of one or more binding agents and/or one or more disintegrants and/or one or more lubricants and/or one or more other excipients;
- 10 (b) compressing the mixture of (a) above.

In another aspect of the invention the direct compression formulation comprises Formula (I), an alkaline effervescent excipient, additional filler(s), binding agent(s), lubricant(s) and disintegrant(s).

15 In another aspect, this invention provides a tablet formed directly by compressing the mixture of (a) above.

In another aspect, this invention provides a direct compression formulation comprising greater than 10% w/w of Formula (I) and an amount of an alkaline effervescent excipient that is sufficient to provide satisfactory *in vitro* dissolution; and further comprising one or more pharmaceutically acceptable ingredients.

In another aspect of the invention the direct compression formulation comprises between 10% w/w to 60% w/w of Formula (I).

25 In a further aspect, the direct compression formulation comprises between 10% w/w to 50% w/w of Formula (I).

In a still further aspect, the direct compression formulation comprises between 15% w/w to 40% w/w of Formula (I).

In a still further aspect, the direct compression formulation comprises between 15% w/w to 25% w/w of Formula (I).

30 In another aspect of the invention the direct compression formulation comprises about 20% w/w of Formula (I).

In a specific aspect of the invention, the direct compression formulation contains 21.33% $\pm$ 5% w/w of Formula (I).

In a further aspect, the direct compression formulation contains 20.26% w/w to 22.40% w/w of Formula (I).

5 In a still further aspect of the invention, the direct compression formulation comprises from 1% w/w to 50% w/w of an alkaline effervescent excipient.

In a further aspect, the direct compression formulation comprises from 1% w/w to 40% w/w of an alkaline effervescent excipient.

10 In a still further aspect, the direct compression formulation comprises from 10% to 30% w/w of an alkaline effervescent excipient.

In a still further aspect, the direct compression formulation comprises from 15% to 25% w/w of an alkaline effervescent excipient.

In a still further aspect, the direct compression formulation comprises about 20% w/w of an alkaline effervescent excipient.

15 In a further aspect of the invention, there is provided a direct compression formulation comprising from 10% w/w to 50% w/w of Formula (I) and from 1% w/w to 50% w/w of an effervescent agent; and further comprising one or more pharmaceutically acceptable ingredients.

20 In a further aspect of the invention, there is provided a direct compression formulation comprising from 15% w/w to 45% w/w of Formula (I) and from 10% w/w to 40% w/w of an effervescent agent; and further comprising one or more pharmaceutically acceptable ingredients.

25 In a further aspect of the invention, there is provided a direct compression formulation comprising from 15% w/w to 25% w/w of Formula (I) and from 15% w/w to 25% w/w of an effervescent agent; and further comprising one or more pharmaceutically acceptable ingredients.

30 The pharmaceutical composition and/or tablet and/or wet granulation formulation and/or roller compaction formulation and/or direct compression formulation can additionally and optionally include a colourant, as long as it is approved and certified by the FDA. For example, exemplary colours include allura red, acid fuschin D, naphthalone red B, food orange 8, eosin Y, phyloxine B, erythrosine, natural red 4, carmine, red iron oxide, yellow iron oxide, black iron oxide, titanium dioxide and the like.

Sweetening agents can also be added to the pharmaceutical composition and/or tablet and/or wet granulation formulation and/or roller compaction formulation and/or direct compression formulation or to the the outer core of the tablet to create or add to the sweetness. Saccharide fillers and binders, e.g. mannitol, lactose, and the like, can add to this effect. For example, cyclamates, saccharin, aspartame, acesulfame K (Mukherjee (1997) Food Chem. Toxicol. 35:1177-1179), or the like (Rolls (1991) Am. J. Clin. Nutr. 53:872-878), can be used. Sweeteners other than sugars have the advantage of reducing the bulk volume of the pharmaceutical composition and/or tablet (core tablet and/or coat) and/or wet granulation formulation and/or roller compaction formulation and/or direct compression formulation and not effecting the physical properties of the tablet.

The pharmaceutical composition and/or tablet and/or wet granulation formulation and/or roller compaction formulation and/or direct compression formulation can additionally and optionally be coated using a conventional pan coater. The film coat may be applied by spraying an aqueous suspension of the coating ingredients onto the tablet cores.

Definitions

As used herein, the term “effervescent excipient” refers to any pharmaceutically acceptable material which evolves a gas in response to a stimulus, for example the evolution of carbon dioxide on acidification. An example of an effervescent excipient is a carbonate, for example a metal carbonate (such as sodium carbonate, potassium carbonate, magnesium carbonate, calcium carbonate or aluminium carbonate) or an organic carbonate (such as disodium glycine carbonate, dimethyl carbonate or ethylene carbonate). A further example of an effervescent excipient is a bicarbonate, for example a metal bicarbonate (such as sodium hydrogen carbonate or potassium hydrogen carbonate).

As used herein, the term “alkaline” refers to a material which induces an increase in pH when added to an aqueous system. The term “alkaline excipient” refers to any pharmaceutically acceptable material which is alkaline, for example an inorganic base such as disodium hydrogen phosphate or sodium hydroxide.

An alkaline effervescent excipient is a pharmaceutically acceptable material having both effervescent activity and alkaline properties, for example sodium hydrogen carbonate, potassium hydrogen carbonate, magnesium carbonate and sodium carbonate. For the

avoidance of doubt, each of the alkaline effervescent excipients referred to above represents a separate and independent aspect of the invention. In one particular aspect of the invention, the alkaline effervescent excipient is selected from a metal carbonate or a metal bicarbonate. In another particular aspect of the invention, the alkaline effervescent excipient is selected from magnesium carbonate, sodium hydrogen carbonate, potassium hydrogen carbonate, or sodium carbonate. In a further particular aspect of the invention, the alkaline effervescent excipient is magnesium carbonate.

As used herein, the term “binding agent” refers to a pharmaceutically acceptable compound or composition added to a formulation to hold the active pharmaceutical ingredient and inactive ingredients together in a cohesive mix. Dry binders used for direct compaction must exhibit cohesive and adhesive forces so that when compacted the particles agglomerate. Binders used for wet granulation are hydrophilic and soluble in water and are usually dissolved in water to form a wet mass that is then granulated. Examples of suitable binding agents includes, but are not limited to, Povidone, Plasdone K29/32, Plasdone S-630, hydropropyl cellulose, methylcellulose, polyvinylpyrrolidone, aluminium stearate, hydroxypropylmethylcellulose and the like. It is possible for such binding agents to additionally act as water sequestering agents (e.g. Povidone).

As used herein, the term “filler” refers to any pharmaceutically acceptable material or composition added to a formulation to add bulk. Suitable fillers include, but are not limited to, mannitol, lactose, microcrystalline cellulose, silified microcrystalline cellulose and dicalcium phosphate.

As used herein, the term “lubricant” refers to any pharmaceutically acceptable agent which reduces surface friction, lubricates the surface of the granule, decreases tendency to build-up of static electricity, and/or reduces friability of the granules. Thus, lubricants can serve as anti-agglomeration agents. Conventional lubricants include stearic acid and related compounds such as magnesium stearate and sodium stearyl fumarate. Alternative lubricants include glyceryl dibehenate, colloidal silica, talc, other hydrogenated vegetable oil or triglycerides. Examples of suitable alternative lubricants include, but are not limited to, glyceryl dibehenate.

As used herein, the term “disintegrant” refers to materials added to the composition to help it break apart (disintegrate) and release the medicaments. Examples of disintegrants include, but are not limited to, non-saccharide water soluble polymers, such as cross-linked

povidine. Other disintegrants that can also be used include, e.g. croscarmellose sodium, sodium starch glycolate, and the like, e.g. see Khattab (1992) J. Pharm. Pharmacol. 45:687-691.

5 The term “drying” and “dried” refer to a process which decreases the water content of a composition to a desired level.

The terms “compressing”, “molding” and “pressing” refer to the process of applying compressive force to a formulation (powder or granules), as within a die, to form a tablet. The terms “compressed tablet” and “pressed tablet” mean any tablet formed by such a process.

10 The term “filming” refers to the adhesion of material to tablet punch surfaces. If sufficient material is allowed to build on punch surfaces then, among other defects, tablet weights may reduce below acceptable limits. (Journal of Pharmaceutical Sciences, Vol. 93(2), 2004).

15 The term “tablet” is used in its common context, and refers to a solid composition made by compressing and/or molding a mixture of compositions in a form convenient for swallowing or application to any body cavity.

As used herein, “tablet strength” is calculated based upon the amount of Compound I.

20 As used herein, “percent loading” is calculated by reference to the percentage by weight of Compound I

The term “low pH” refers to a measured pH of less than 5, such as less than 3, for example between 0 and 3.

25 The term “satisfactory *in vitro* dissolution” refers to a percent dissolution of greater than or equal to 70% within 30 minutes in a suitable dissolution medium at 37°C ±0.5°C as measured using the general procedure of the United States Pharmacopeia (Apparatus 2).

30 The term “stable formulation” refers to a formulation which, following storage for 4 weeks at elevated temperature and humidity, such as 40°C and 75% relative humidity, exhibits water absorption of less than 10%, such as less than 5%, for example between 0 and 5%; and / or chemical degradation of less than 3%, such as less than 2.5 %, for example between 0 and 2.5%; and / or which exhibits satisfactory *in vitro* dissolution.

The term “manufacturability” means the extent to which a product can be manufactured with relative ease at minimum cost and maximum reliability.

DESCRIPTION OF FIGURES

Figure 1 shows a plot of the percentage dissolution using pH1.3 hydrochloric acid /
5 sodium chloride buffer of ten alternative tablet formulations.

Figure 2 shows a plot of the percentage dissolution using pH1.3 hydrochloric acid /
sodium chloride buffer of a further nineteen alternative tablet formulations.

Figure 3 shows a plot of the percentage dissolution using pH6.8 phosphate buffer of five
alternative tablet formulations in which the lubricant content was varied.

10 **Figure 4** shows a plot of the percentage dissolution using pH1.3 hydrochloric acid /
sodium chloride buffer for ten alternative tablet formulations in which Formula (I),
magnesium carbonate and lubricant was varied.

Figure 5 shows a plot of the percentage dissolution using pH6.8 phosphate buffer for ten
alternative tablet formulations in which Formula (I), magnesium carbonate and lubricant
15 was varied.

EXAMPLES

The invention is further understood by reference to the following examples, which are intended to be purely exemplary of the invention. The present invention is not limited in scope by the exemplified aspects, which are intended as illustrations of single aspects of the invention only. Various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications fall within the scope of the appended claims.

In the examples below as well as throughout the application, the following abbreviations have the following meanings. If not defined, the terms have their generally accepted meanings.

	API	=	Active Pharmaceutical Ingredient
	CCS	=	croscarmellose sodium
15	CroSPov	=	crospovidone
	BP	=	British Pharmacopoeia 2012
	DCPA	=	dicalcium Phosphate (anhydrous)
	DCPD	=	dicalcium Phosphate (dihydrate)
	Glydb	=	glyceryl dibehenate
20	HPC	=	hydroxypropylcellulose
	L-HPC	=	hydroxypropylcellulose, low-substituted
	LOD	=	loss on drying
	Mag carb	=	magnesium carbonate
	MCC	=	cellulose, microcrystalline
25	MgSt	=	magnesium stearate
	min	=	minute
	ml	=	milliliter
	nm	=	nanometer
	JP	=	Japanese Pharmacopoeia 15 th Edition, English Version (Society of Japanese Pharmacopoeia) 2006
30	PhEur	=	European Pharmacopoeia 6 th Edition (Directorate for the Quality of Medicines of the Council of Europe) 2009

	rpm	=	revolutions per minute
	SLS	=	sodium lauryl sulphate
	SSF	=	sodium stearyl fumarate
	SSG	=	sodium starch glycolate
5	USP/USP-NF	=	United States Pharmacopeia 31 / National Formulary 26 (The United States Pharmacopeia Convention) 2008
	UV	=	ultraviolet
	w/w	=	weight for weight

10 Table 1 below shows materials used, pharmacopeial status, grade and supplier.

Table 1

Material	Pharmacopeia	Grade	Supplier
Mannitol	PhEur USP-NF JP	Pearlitol 160c	Roquette Freres S.A. (France)
Cellulose, microcrystalline	PhEur USP-NF JP	Avicel® PH-101	FMC Biopolymer (Ireland)
Salicified cellulose, microcrystalline	USP-NF	Prosolv® 90	Rettenmaier UK Ltd (UK)
Dicalcium phosphate (anhydrous)	PhEur BP JP USP	Calipharm A	Innophos (USA)
Dicalcium phosphate (dihydrate)	PhEur BP JP USP	Calipharm D	Innophos (USA)
Sodium bicarbonate	PhEur BP	N/A	Dr Paul Lohmann (Germany)

Material	Pharmacopeia	Grade	Supplier
	JP USP		
Calcium carbonate (heavy)	PhEur BP JP USP	N/A	Dr Paul Lohmann (Germany)
Magnesium carbonate (heavy)	PhEur BP JP USP	N/A	Dr Paul Lohmann (Germany)
Disodium phosphate (dibasic)	PhEur BP JP USP-NF	N/A	Budenheim (USA)
Sodium starch glycolate	Ph Eur USP-NF	Glycolys LV	Roquette Freres S.A. (France)
Hydroxypropylcellulose, low-substituted	JP USP-NF	L-HPC	Shin Etsu, (Japan)
Croscarmellose sodium	Ph Eur USP JP	Ac-di-Sol	FMC Biopolymer (Ireland)
Crospovidone	PhEur BP USP-NF	Polyplasdone XL	Ashland Speciality Ingredients, (UK)
Hydroxypropylcellulose	PhEur BP USP-NF JP	Klucel EXF	Ashland Speciality Ingredients (UK)
Hydroxypropylmethylcellulose (hypromellose)	PhEur BP	Pharmacoat 603	Shin Etsu, (Japan)

Material	Pharmacopeia	Grade	Supplier
	USP-NF JP		
Sodium lauryl sulphate (Sodium dodecyl sulfate)	USP NF	N/A	Sigma Aldrich (UK)
Magnesium stearate	PhEur USP-NF JP	NF Non Bovine	Mallinckrodt (USA)
Sodium stearyl fumurate	PhEur BP USP-NF	Pruv	JRS Pharma, (Germany)
Glyceryl dibehenate	PhEur USP	Compritol 888 ATO	Gattefosse (France)
Opadry II Biege	N/A	N/A	Colorcon (USA)

Table 2 below shows equipment used, model and supplier.

Table 2

Make	Model	Supplier
Pro-C-ept	Mi-pro	Pro-C-ept, Belgium
Diosna	P1/6	Dierks & Söhne Gmbh, Osnabrück, Germany
Collette	Gral 10 & Gral 25	Collette Machines, Belgium
Quadro	Comil U3 & Comil 194	Quadro Engineering, Waterloo, Canada
WAB	Turbula T2F	Willy A. Bachofen AG, Muttens, Switzerland
Copley Mobile Blender	Mobile Blender	Copley Scientific, Nottingham, UK
Aeromatic	Strea 1	Casburt Pharmaceutical Equipment, Stoke-on-Trent, UK
Aeromatic- Fielder	MP1	Aeromatic Fielder, Eastleigh, UK

Make	Model	Supplier
Vector	MFL.01	Vector Corporation, Marion, IA, U.S.A
Glatt	59P	Glatt GmbH, Binzen, Germany
Riva	Piccola-Nova,	RivaSA, Buenos Aires, Argentina
Manesty	F3	Manesty, Knowsley, UK
Korsch	Korsch XL100	Korsch AG, Berlin, Germany
Riva	Riva mini-press	RivaSA, Buenos Aires, Argentina
Riva	Piccola W.I.P	RivaSA, Buenos Aires, Argentina
O'Hara	Labcoat II-X	O'hara technologies inc, Ontario, Canada

Example 1: Assessment of dissolution performance of ten alternative tablet forms

It has been found that the rheology of Formula (I) can change under certain conditions. In particular, Formula (I) can convert from a crystalline powder to a highly viscous material under low pH and at high concentration (both conditions need to be met simultaneously). Theoretically, these conditions will be met in the microenvironment of the tablet matrix either using a low pH dissolution method (e.g. pH1.3) or in the stomach. The relative surface area of Formula (I) reduces when the viscous material forms and this is associated with a reduced rate of solubilisation of Formula (I). This can be observed as a reduced rate of dissolution using a low pH method.

Based on this fundamental understanding, there are two hypothetical mechanisms to avoid the rheological transformation; first, to not allow Formula (I) to solubilise in a low pH environment (Hypothesis I); second, if dispersion in a low pH environment cannot be avoided, to disperse Formula (I) rapidly before the transformation can occur (Hypothesis II). Hypothesis II is dependent on the concentration of Formula (I) in the tablet matrix as higher concentrations of Formula (I) reduce the likelihood of rapid dispersion.

Ten different prototype tablets were prepared from a wet granulation formulation using methods well known to those skilled in the art. The composition of each of these tablets is set out in Table 3.

Table 3

Run	Formula (I), % w/w	MCC type (level, % w/w)	Mannitol level (% w/w)	pH Buffer type (% w/w)	Disintegrant Type (% w/w)	HPC (% w/w)
1	20	Avicel PH101 (48.5)	20	N/A	SSG (7.5)	3
2	40	Avicel PH101 (34.3)	14.2	N/A	SSG (7.5)	3
3	40	Avicel PH101 (16.7)	11.8	NaHCO ₃ (20.0)	SSG (7.5)	3
4	40	Avicel PH101 (16.7)	11.8	CaCO ₃ (20.0)	SSG (7.5)	3
5	40	Avicel PH101 (16.7)	11.8	Na ₂ HPO ₄ (20.0)	SSG (7.5)	3
6	40	Avicel PH101 (16.7)	11.8	MgCO ₃ (20.0)	SSG (7.5)	3
7	40	Prosolv® SMCC (34.3)	14.2	N/A	SSG (7.5)	3
8	40	N/A	31.0	NaHCO ₃ (20.0)	SSG (5)	3
9	20	Avicel PH101 (48.5)	20	N/A	CCS (7.5)	3
10	20	Avicel PH101 (48.5)	20	N/A	CrosPov (7.5)	3

Run 1 is comparable to the Phase 1 clinical formulation and is the positive control. Run 2 is a negative control as it contains a high concentration of Formula (I) and has no alkalisng agent. Run 5 tests Hypothesis I (Na₂HPO₄ is an alkalisng agent). Runs 7, 9 and 10 test Hypothesis II (no alkalisng agent, but varying disintegrants). Runs 3, 4, 6 and 8 test both Hypotheses I and II (they contain carbonate/bicarbonate alkalisng agents which both increase the pH microenvironment and liberate carbon dioxide in acidic conditions; carbon dioxide liberation can help to disperse Formula (I)).

Formula (I) and the excipients (except lubricant) described in Table 1 (total batch size approximately 250g) were charged to a mixer-granulator (Diosna, 1 litre bowl, P1/6) and mixed. Purified water was added to the powders with further mixing until a suitable wet mass was formed. The resultant granules were dried to appropriate moisture content (≤2% w/w LOD) using a fluid bed dryer (Vector, MFL.01) with an inlet air temperature of

65°C. The dried granules were milled using an appropriately sized screen (1 mm, Quadro Comil U3).

SSF was then added to the granules (Table 4), which were then blended (WAB turbula) for 10 mins at 55 rpm before compressing into tablet cores using conventional tableting

5 equipment (Manesty F3 tablet press)Table 4

Milled granules from composition variant (Table 1)	SSF Addition (% w/w)
1	1
2	1
3	2
4	2
5	2
6	2
7	2
8	2
9	2
10	2

Concentration of SSF in the compositions was increased after Run 2 to allow viable processability during compression. Theoretically, this would reduce adhesion of material to tablet punches and dies (Pharmaceutical Powder Compaction Technology, edited by Goran Alderborn and Christer Nyström, Informa Healthcare, New York, 2008). However,

10 increasing level of lubricant also typically reduces rate of dissolution due to the hydrophobic nature of the lubricant.

Further process conditions are given in Table 5.

Table 5

Run	Chopper (rpm)	Impellor (rpm)	Total Water Added (ml)	Total granulation time (min)
1	1000	300	81	4.1
2	1000	300	40	2.0
3	1500	750	65	6.5
4	1500	750	80	8.0
5	1500	750	80	8.0

Run	Chopper (rpm)	Impellor (rpm)	Total Water Added (ml)	Total granulation time (min)
6	1500	750	65	6.5
7	1500	750	70	7.0
8	1500	750	25	2.5
9	1500	750	80	8.0
10	1500	750	70	7.0

Impellor and chopper speeds were increased after Run 2 to allow viable processability during compression. Theoretically, increasing these process conditions increases granule density (Powder Technology, 117, pp 3-39, 2001) which aids granule flow and reduces punch filming. However, increasing granule density also typically reduces rate of dissolution.

Similarly, increasing total water added (Powder Technology, 88, pp 15-20, 1996) and granulation time (granule densification is a rate process) are also likely to increase granule density and thus reduce dissolution rate.

Dissolution was determined according to the general procedure of the United States Pharmacopeia using Apparatus 2 with pH1.3 hydrochloric acid and sodium chloride buffered solution at 37°C ±0.5°C and stirrer speed of 50 rpm. At 15, 30 and 60 minutes dissolution media was withdrawn and the concentration of Formula (I) in solution was determined by UV spectroscopy at a wavelength of 311 nm against an external standard solution. Dissolution profiles are shown in Figure 1, the dissolution data is presented in Table 6.

Table 6

Run	15 minutes	30 minutes	60 minutes
1	67.4	76.1	85.8
2	42.9	50.7	62.6
3	100.6	99.9	100.3
4	65.3	77.0	86.1
5	29.4	43.8	53.1
6	92.4	95.1	96.7
7	48.1	55.5	65.7
8	100.1	99.3	99.7

Run	15 minutes	30 minutes	60 minutes
9	53.2	63.4	78.0
10	54.9	63.4	74.4

Example 2: Assessment of tablet punch filming of ten alternative tablet forms

Ten different prototype tablets were prepared from a wet granulation formulation using methods well known to those skilled in the art. The composition and manufacturing process of each of these tablets is described in Example 1.

- 5 Material adhesion to tablet punch surfaces (described below as ‘filming’) is a well known tableting process defect (Journal of Pharmaceutical Sciences, Vol. 93(2), 2004). Extent of filming was visually assessed for each formulation and reported in Table 7.

Table 7

Run	Filming
1	★ ★
2	★ ★ ★
3	★
4	No filming
5	No filming
6	★
7	★ ★
8	★ ★ ★
9	No filming
10	No filming

★ = minor ★ ★ = moderate ★ ★ ★ = severe

10

Example 3: Assessment of tablet water absorption of ten alternative tablet forms

Ten different prototype tablets were prepared from a wet granulation formulation using methods well known to those skilled in the art. The composition and manufacturing process of each of these tablet compositions is described in Example 1.

- 15 Extent of water absorption was measured for each formulation (see Table 8). The tablets were exposed to a controlled environment (40°C and 75% relative humidity) for one month.

Table 8

Run	Water Absorption (% w/w)
1	8.5
2	7.9
3	32.2
4	13.5
5	27.9
6	4.4
7	7.6
8	32.2
9	9.1
10	8.8

Example 4: Assessment of dissolution performance of a further nineteen alternative tablet forms

Run 6 (Examples 1, 2 and 3) was selected for further development because, unlike other compositions, it showed a marked improvement in pH1.3 dissolution (Figure 1), an improvement in punch filming (Table 7) and low water absorption (Table 8).

A further nineteen different prototype tablets were prepared from a wet granulation formulation using methods well known to those skilled in the art. The composition of each of these tablets is qualitatively similar to Run 6. Quantitative compositions are set out in Table 9.

Table 9

Run	Formula (I) (% w/w)	MCC (% w/w)	Mannitol (% w/w)	Mag Carb (% w/w)	HPC (% w/w)	SSG (% w/w)	SSF (% w/w)	Water Addition (% w/w)
1	21.3	0.0	59.7	10.0	1.0	5.0	3.0	20.10
2	21.3	30.0	29.7	10.0	1.0	5.0	3.0	26.20
3	21.3	0.0	39.7	30.0	1.0	5.0	3.0	15.00
4	21.3	34.8	0.0	34.8	1.0	5.0	3.0	45.00
5	21.3	0.0	57.7	10.0	3.0	5.0	3.0	20.00
6	21.3	30.0	27.7	10.0	3.0	5.0	3.0	35.00

Run	Formula (I) (% w/w)	MCC (% w/w)	Mannitol (% w/w)	Mag Carb (% w/w)	HPC (% w/w)	SSG (% w/w)	SSF (% w/w)	Water Addition (% w/w)
7	21.3	0.0	37.7	30.0	3.0	5.0	3.0	15.00
8	21.3	33.8	0.0	33.8	3.0	5.0	3.0	40.20
9	21.3	0.0	54.7	10.0	1.0	10.0	3.0	20.00
10	21.3	30.0	24.7	10.0	1.0	10.0	3.0	30.00
11	21.3	0.0	34.7	30.0	1.0	10.0	3.0	30.40
12	21.3	32.3	0.0	32.3	1.0	10.0	3.0	40.00
13	21.3	0.0	52.7	10.0	3.0	10.0	3.0	20.30
14	21.3	30.0	22.7	10.0	3.0	10.0	3.0	32.50
15	21.3	0.0	32.7	30.0	3.0	10.0	3.0	32.40
16	21.3	31.3	0.0	31.3	3.0	10.0	3.0	35.30
17	21.3	15.0	31.2	20.0	2.0	7.5	3.0	40.00
18	21.3	15.0	31.2	20.0	2.0	7.5	3.0	40.00
19	21.3	15.0	31.2	20.0	2.0	7.5	3.0	40.30

Formula (I) and the excipients described in Table 9 (total batch size approximately 1.5 kg) were charged to a mixer-granulator (Colette Gral 10) and mixed. Purified water (Ranging from 15% w/w to 45% w/w as set out in Table 9) was added to the powders with further mixing until a suitable wet mass was formed (ranging from approximately 3 to 14 mins) at 420 rpm. The resultant granules were dried to appropriate moisture content ($\leq 2\%$ LOD) using a fluid bed dryer (Aeromatic Strea 1) with an inlet air temperature of 80°C. The dried granules were milled using an appropriately sized screen (1.4 mm, Quadro Comil U3). SSF was then added to the granules, which were then blended (Copley, Mobile Blender 7.5 litre drum) for 5 mins at 25 rpm before compressing into tablet cores using conventional tableting equipment (Korch XL100).

Dissolution was determined according to the general procedure of the United States Pharmacopeia using Apparatus 2 with pH1.3 hydrochloric acid and sodium chloride buffered solution at 37°C $\pm 0.5^\circ\text{C}$ and stirrer speed of 50 rpm. At 15, 30 and 60 minutes dissolution media was withdrawn and the concentration of Formula (I) in solution was determined by UV spectroscopy at a wavelength of 311nm against an external standard solution. Dissolution profiles are shown in Figure 2, the dissolution data is presented in Table 10.

Table 10

Run	15 minutes	30 minutes	60 minutes
1	82.6	91.3	91.8
2	68.9	78.6	89.2
3	67.5	77	86.3
4	101.1	100.7	100.2
5	91.7	97.8	99.8
6	96.9	97	97.5
7	61.8	73.9	89.9
8	107.2	106.9	107.9
9	54.5	69.4	84.9
10	98.6	93.6	98
11	92.9	97.9	101.3
12	104.3	108.7	112.8
13	106.3	108.1	108.5
14	101.1	100.8	101.1
15	100	99.5	100.4
16	97.5	99.9	101.3
17	97.9	88.7	98.3
18	86	94.5	100.4
19	82.6	91.3	91.8

Example 5: Assessment of chemical stability of a sixteen alternative tablet forms

- 5 Sixteen different prototype tablets were prepared from a wet granulation formulation using methods well known to those skilled in the art. The composition and manufacturing process of each of these tablets is described in Table 11.

Table 11

Run Order	Main Filler	Secondary Filler	Disintegrant	Binder	Surfactant	Lubricant
1	Mannitol	DCPA	L-HPC	HPC	None	SSF
2	Mannitol	DCPD	SSG	HPC	None	SSF
3	Mannitol	DCPD	L-HPC	HPC	SLS	MgST

Run Order	Main Filler	Secondary Filler	Disintegrant	Binder	Surfactant	Lubricant
4	Mannitol	DCPA	SSG	HPMC	SLS	SSF
5	Mannitol	DCPA	L-HPC	HPMC	None	MgST
6	MCC	DCPA	SSG	HPMC	None	SSF
7	MCC	DCPA	L-HPC	HPC	SLS	SSF
8	Mannitol	DCPD	L-HPC	HPMC	SLS	SSF
9	MCC	DCPD	L-HPC	HPMC	None	SSF
10	MCC	DCPA	L-HPC	HPMC	SLS	MgST
11	MCC	DCPA	SSG	HPC	None	MgST
12	MCC	DCPD	L-HPC	HPC	None	MgST
13	Mannitol	DCPA	SSG	HPC	SLS	MgST
14	Mannitol	DCPD	SSG	HPMC	None	MgST
15	MCC	DCPD	SSG	HPMC	SLS	MgST
16	MCC	DCPD	SSG	HPC	SLS	SSF

Formula (I) and the excipients (except lubricant) described in Table 11 (total batch size approximately 50g) were charged to a mixer-granulator (Mi-Pro, 500 ml bowl) and mixed. Purified water was added (approximately 10 ml/min) to the powders with further mixing until a suitable wet mass was formed. The resultant granules were dried to appropriate moisture content ($\leq 2\%$ w/w LOD) using a fluid bed dryer (Vector, MF1.01). The dried granules were milled using an appropriately sized screen (1 mm, Quadro Comil U3). Lubricant was then added to the granules, which were then blended (WAB turbula) for 4 mins at 24 rpm before compressing into tablet cores using conventional tableting equipment (Manesty F3 tablet press).

Total impurities were measured by injection of the prepared sample and standard solutions onto an LC system selected to ensure the separation of Formula (I) from organic impurities and excipients. The chromatographic responses due to Formula (I) and organic

impurities are measured on a UV detector at wavelength 245nm. The response due to Formula (I) present in the sample was compared to that of a standard and its assay was calculated. The level of organic impurities was calculated as %w/w. Equivalent response was assumed between Formula (I) and organic impurities.

Samples were stored in a controlled environment for four weeks at 60°C and 80% relative humidity. After analysis, samples with SSF in their composition contained 0.99±0.36% (mean±standard deviation%) and samples with MgSt in their composition contained 1.93±1.34% (mean±standard deviation%).

Example 6: Assessment of dissolution performance when varying SSF concentration and inclusion of an alternative lubricant

Full extent of release was not achieved for each of the prototype formulations presented in Example 4 in pH 6.8 phosphate buffered solution. The affect of lubricant was investigated and five different prototype tablets were prepared by a wet granulation process using methods well known to those skilled in the art. The composition of each of these tablets is described in Table 12.

Table 12

Run	Milled granules from composition variant (Table 5)	SSF (% w/w)	Glydb (% w/w)
0% SSF	17 and 18	0.0	N/A
1% SSF	17 and 18	1.0	N/A
2% SSF	17 and 18	2.0	N/A
5% SSF	17 and 18	5.0	N/A
3% Glybd	17 and 18	N/A	3.0

Milled unlubricated granules from the prototype variants 17 and 18 (equivalent compositions), presented in example 4 (Table 9), were combined with the relevant level of lubricant, presented in Table 12, and blended (WAB turbula) for 5 mins at 25 rpm before compressing into tablet cores using conventional tableting equipment (RIVA mini-press). Dissolution was determined according to the general procedure of the United States Pharmacopeia using Apparatus 2 with pH6.8 phosphate buffered solution at 37°C ±0.5°C and stirrer speed of 75 rpm. At 15, 30, 45 and 60 minutes dissolution media was withdrawn

and the concentration of Formula (I) in solution was determined by UV spectroscopy at a wavelength of 298 nm against an external standard solution. Dissolution results are shown in Figure 3 and the dissolution data is presented in Table 13.

5 Table 13

Run	15 minutes	30 minutes	45 minutes	60 minutes
0% SSF	85	94	97	99
1% SSF	83	91	94	96
2% SSF	74	81	86	89
5% SSF	65	71	74	76
3% Glybd	83	93	97	99

Example 7: Assessment of tablets of Formula (I) prepared with different lubricants

Three different prototype tablets were prepared from a wet granulation formulation using methods well known to those skilled in the art. The composition of each of these tablets (including a range of lubricants i.e. MgSt, SSF and Glydb) is described below.

The pharmaceutical composition comprises the following components (% w/w):

	<u>Composition A (% w/w)</u>		<u>Composition B (% w/w)</u>		<u>Composition C (% w/w)</u>	
<u>API</u>	Formula (I)	10.0	Formula (I)	21.3	Formula (I)	21.3
<u>Primary Filler</u>	Mannitol	60.0	Mannitol	31.2	Mannitol	31.2
<u>Secondary Filler</u>	DCPD	20.0	MCC	15.0	MCC	15.0
<u>Tertiary Filler</u>			MgCO ₃	20.0	MgCO ₃	20.0
<u>Disintegrant</u>	SSG	5.0	SSG	7.5	SSG	7.5
<u>Binder</u>	HPMC	4.0	HPC	2.0	HPC	2.0
<u>Lubricant</u>	MgSt	1.0	SSF	3.0	Glydb	3.0

Manufacturing processes for Compositions A, B and C are given in Example 5, Example 4 and Example 6, respectively.

An evaluation of tablet dissolution performance was made (Table 14). Dissolution was determined according to the general procedure of the United States Pharmacopeia using Apparatus 2 with pH6.8 phosphate buffered solution at 37°C ±0.5°C and stirrer speed of 50 rpm. At 15, 30, and 60 minutes dissolution media was withdrawn and the concentration

of Formula (I) in solution was determined by UV spectroscopy at a wavelength of 298nm against an external standard solution.

Table 14

Composition	15 minutes	30 minutes	60 minutes
A	55.8	82.5	88.3
B	69.3	76.3	78.1
C	79.0	88.2	93.7

- 5 Composition C was selected for further study because, unlike other compositions, it (i) did not contain MgSt, which on average gave more impurities than SSF (Example 5); (ii) did not contain SSF, inclusion of which can affect extent of dissolution release (Figure 3); and (iii), demonstrated acceptable dissolution performance throughout the physiologically relevant pH range (Table 15).

10

Table 15

Composition C	15 minutes	30 minutes	60 minutes
pH1.3	98.2	101.5	102.0
pH6.8	79.0	88.2	93.7

Example 8: Assessment of dissolution performance of ten alternative tablet formulations

- 15 Ten alternative prototype tablets were prepared from a wet granulation using methods well known to those skilled in the art. The composition of each of these tablets is qualitatively similar to Composition C (Example 7). Quantitative compositions are set out in Table 16.

Table 16

Composition	API (% w/w)	MgCO ₃ (% w/w)	Glydb (% w/w)	Mannitol (% w/w)	MCC (% w/w)	SSG (% w/w)	HPC (% w/w)
1	10	15	8	38.72	18.78	7.5	2
2	40	25	8	11.67	5.83	7.5	2
3	10	25	1	36.79	17.71	7.5	2
4	40	15	1	23.29	11.21	7.5	2

Composition	API (% w/w)	MgCO ₃ (% w/w)	Glydb (% w/w)	Mannitol (% w/w)	MCC (% w/w)	SSG (% w/w)	HPC (% w/w)
5	40	25	8	11.67	5.83	7.5	2
6	10	15	1	43.55	20.95	7.5	2
7	40	25	1	16.54	7.96	7.5	2
8	10	25	8	31.67	15.83	7.5	2
9	21.33	20	3	31.17	15	7.5	2
10	21.33	20	3	31.17	15	7.5	2

Formula (I) and the excipients (except lubricant) described in Table 16 (total batch size approximately 250g) were charged to a mixer-granulator (Diosna, 1 litre bowl, P1/6) and mixed. Purified water was added (approximately 10 ml/min) to the powders with further mixing until a suitable wet mass was formed. The resultant granules were dried to appropriate moisture content ($\leq 2\%$ w/w LOD) using a fluid bed dryer (Aeromatic Strea 1). The dried granules were milled using an appropriately sized screen (1.4 mm, Quadro Comil U3). Lubricant was then added to the granules, which were then blended (WAB turbula) for 10 mins at 55 rpm. The granules were then compressed into tablet cores (each core normalised to 80mg of formula 1) using conventional tableting equipment (Riva Piccola (W.I.P) tablet press) at a normalised pressure of 100MPa.

An evaluation of tablet dissolution performance was made (Table 17 and 18). Dissolution was determined according to the general procedure of the United States Pharmacopeia using Apparatus 2 with both pH6.8 phosphate solution and pH1.3 hydrochloric acid and sodium chloride solution at $37^{\circ}\text{C} \pm 0.5^{\circ}\text{C}$ and stirrer speed of 50 rpm. At 15, 30, and 60 minutes dissolution media was withdrawn and the concentration of Formula (I) in solution was determined by UV spectroscopy at a wavelength of 311 nm (for pH 1.3 solution) or 298 nm (for pH 6.8 solution) against an external standard solution. Dissolution results are shown in Table 17/Figure 4 (pH 1.3) and Table 18/Figure 5 (pH 6.8).

Table 17

Composition	15 minutes	30 minutes	60 minutes
1	104	102	102
2	84	90	93
3	95	94	94
4	72	79	87
5	78	90	95
6	111	110	110
7	91	93	95
8	100	98	98
9	98	97	98
10	96	95	96

Table 18

Composition	15 minutes	30 minutes	60 minutes
1	67	86	97
2	38	56	73
3	55	72	84
4	38	58	77
5	36	54	72
6	83	98	105
7	56	75	87
8	71	86	93
9	56	77	91
10	47	68	84

5

Example 9: Assessment of tablet punch filming of ten alternative tablet formulations

Ten alternative prototype tablets were prepared from a wet granulation formulation using methods well known to those skilled in the art. The composition and manufacturing process of each of these tablets is described in Example 8.

10

Material adhesion to tablet punch surfaces (described below as ‘filming’) is a well known tableting process defect (Journal of Pharmaceutical Sciences, Vol. 93(2), 2004). Extent of filming was visually assessed for each formulation and reported in Table 19.

Table 19

Composition	Filming
1	No filming
2	No filming
3	★★
4	★★
5	No filming
6	★★
7	★★★
8	No filming
9	No filming
10	No filming

★ = minor ★★ = moderate ★★★ = severe

No punch filming was observed for the compositions with 3-8% w/w glydb. Moderate to severe levels of punch filming was observed for the compositions containing 1% w/w glydb under the process conditions applied.

Example 11: Assessment of dissolution of an alternative tablet formulation

One alternative tablet formulation was prepared from a wet granulation using methods well known to those skilled in the art. The composition of the tablet core is quantitatively similar to Composition C (Example 7) and a film coat was applied using a conventional film coating method to enhance the tablet appearance. The quantitative composition of the tablet core formulation is presented in Table 20.

Table 20

Component	Composition (% w/w)
AZD4547	21.33
Mannitol	31.17
Micro crystalline cellulose (Avicel PH101)	15
Magnesium Carbonate (Heavy)	20
Sodium Starch Glycolate	7.5
Hydroxy Propyl Cellulose	2
Glyceryl dibehenate	3

A film coat was applied using a propriety mixture of coating excipients, Opadry II Biege, supplied by Colorcon.

5

Formula (I) and the excipients (except lubricant) described in Table 20 (total batch size approximately 5kg) were charged to a mixer-granulator (Gral 25) and mixed. Purified water was added (approximately 166 ml/min) to the powders with further mixing until a suitable wet mass was formed. The resultant granules were dried to appropriate moisture content ($\leq 2\%$ w/w LOD) using a fluid bed dryer (Glatt 59P). The dried granules were milled using an appropriately sized screen (1.4 mm, Quadro Comil U3). Lubricant was then added to the granules, which were then blended (Copley Mobile Blender, 7.5L container) for 5 mins at 25 rpm. The granules were then compressed into tablet cores using conventional tableting equipment (Riva Piccola-Nova, tablet press) to achieve a target compression weight of 375mg. The tablet cores were over coated with a film coat using conventional pan coating equipment (O'Hara Labcoat II-X) to achieve a tablet weight gain of 3% w/w.

10

15

20

An evaluation of tablet dissolution performance was made (Table 22). Dissolution was determined according to the general procedure of the United States Pharmacopeia using


Apparatus 2 with both pH6.8 phosphate solution and pH1.3 hydrochloric acid and sodium chloride solution at $37^{\circ}\text{C} \pm 0.5^{\circ}\text{C}$ and stirrer speed of 50 rpm. At 15, 30 and 60 minutes dissolution media was withdrawn and the concentration of Formula (I) in solution was determined by UV spectroscopy at a wavelength of 311 nm (for pH 1.3 solution) or 298 nm (for pH 6.8 solution) against an external standard solution.

Table 22

	15 minutes	30 minutes	60 minutes
pH1.3	95	101	103
pH6.8	75	87	93

CLAIMS

1. A pharmaceutical composition comprising greater than 10% w/w of the compound
of Formula (I):



and an amount of an alkaline effervescent agent that is sufficient to provide satisfactory *in vitro* dissolution; and further comprising one or more pharmaceutically acceptable ingredients.

2. A pharmaceutical composition according to claim 1 comprising 15% w/w to 40% w/w of the compound of Formula (I).

3. A pharmaceutical composition according to any of the preceding claims comprising 15% to 25% w/w of the alkaline effervescent agent.

4. A pharmaceutical composition according to any of the preceding claims comprising 15% w/w to 25 % w/w of the compound of Formula (I) and 15% w/w to 25% w/w of the effervescent agent; and further comprising one or more pharmaceutically acceptable ingredients.

5. A pharmaceutical composition according to any of the preceding claims wherein the effervescent agent is magnesium carbonate.

6. A pharmaceutical composition according to any of the preceding claims wherein the composition comprises between 0.25% w/w and 8% w/w of an alternative lubricant.

7. A pharmaceutical composition according to any of the preceding claims wherein the composition comprises between 2.5% w/w and 3.5% w/w of an alternative lubricant

8. A pharmaceutical composition according to claims 7 and 8 wherein the alternative
5 lubricant is glyceryl dibehenate.

9. A wet granulation formulation comprising the compound of Formula (I), water, an alkaline effervescent agent, additional filler(s), binding agent(s) and disintegrant(s).

10 10. A wet granulation formulation comprising 15% w/w to 25% w/w of the compound of Formula (I) and 15% w/w to 25% w/w of an alkaline effervescent agent; and further comprising one or more pharmaceutically acceptable ingredients.

11. A roller compaction formulation comprising 15% w/w to 25%w/w of the
15 compound of Formula (I) and 15% w/w to 25% w/w of an alkaline effervescent agent; and further comprising one or more pharmaceutically acceptable ingredients.

12. A direct compression formulation comprising 15% w/w to 25% w/w of the compound of Formula (I) and/or hydrate thereof and 15% w/w to 25% w/w of an alkaline
20 effervescent agent; and further comprising one or more pharmaceutically acceptable ingredients.