GENE-BASED ALGORITHMIC CANCER PROGNOSIS

Field of the Invention

[0001] The present invention is related to new method and tools for improving cancer prognosis.

Background of the Invention

[0002] Microarray profiling, or the assessment of the mRNA expression levels of hundreds and thousands of genes, has shown that cancer can be divided into distinct molecular subgroups by the expression levels of certain genes. These subgroups seem to have distinct clinical outcomes and also may respond differently to different therapeutic agents used in cancer treatment. But the current understanding of the underlying biology does not permit "individualization" of a particular cancer patients' care. As a result for breast cancer, for example, many women today are given systemic treatments such as chemotherapy or endocrine therapy in an attempt to reduce her risk of the breast cancer recurring after initial diagnosis. Unfortunately, this systemic treatment only benefits a minority of women who will relapse, hence exposing many women to unnecessary and potentially toxic treatment. New prognostic tools developed using microarray technology show potential in allowing us to facilitate tailored treatment of breast cancer patients (Paik et al, New England Journal of Medicine 351:27(2004); Van de Vijver et al, New England Journal of Medicine 347:199(2002); Wang et al, Lancet 365: 671 (2005)). These genomic tools may be

a much needed improvement over currently used clinical methods.

[0003] Histological grading of breast carcinomas has long been recognised to provide significant clinical prognostic information (1). However, despite recommendations by the College of American Pathologists (2) for use of tumor grade as a prognostic factor in breast cancer, the latest Breast Task Force serving the American Joint Committee on Cancer (AJCC) did not include it in its staging criteria, citing insurmountable inconsistencies between institutions and lack of data (3). This may be in part related to inter-observer variability and the various grading approaches used, resulting in poor reproducibility across institutions. With the advent of standardized methods such as those developed by Elston and Ellis (1), concordance between institutions has been improved. Nevertheless, whilst grade 1 (low risk) and 3 (high risk) are clearly associated with different prognoses, tumors classified as intermediate grade present a difficulty in clinical decision making for treatment because their survival profile is not different from that of the total (non-graded) population and their proportion is large (40%-50%). A more accurate grading system would allow for better prognostication and improved selection of women for further breast cancer treatment.

[0004] The majority of breast cancers diagnosed today are hormone responsive. Tamoxifen is the most common anti-estrogen agent prescribed today in the adjuvant treatment of these patients. Yet up to 40% of these patients will relapse when given tamoxifen in this setting. At present, due to the positive results of several large trials evaluating the use of aromatase inhibitors instead of, or in combination or sequence with tamoxifen in the adjuvant setting, there are many options available for post

menopausal women with hormone responsive breast cancer. Furthermore, it is unclear which treatment option is the best especially given that the long term health costs of aromatase inhibitor use are unknown. The ability to identify a group at high risk of relapse when given tamoxifen could aid in identifying patients for whom tamoxifen is probably not the best option. These patients could then be specifically targeted for alternative treatment strategies.

[0005] Particularly relating to the issue of predicting relapse for women treated with adjuvant tamoxifen, two publications have been reported claiming gene sets that can predict clinical outcome (Ma et al, Cancer Cell 5:607(2004), Jansen et al. Journal of Clinical Oncology 23:732(2005). These studies involved small numbers of patients and hence are not thoroughly validated to be widely used clinically.

[0006] Accordingly need exists for methods and systems that can accurately assess prognosis and hence help oncologists tailor their treatment decisions for the individual cancer patient. In particular, a need exists for methods and systems directed to breast cancer patients.

Aims of the Invention

[0007] The present invention aims to provide new methods and tools for improving cancer prognosis that do not present the drawbacks of the methods of the state of the art.

Summary of the Invention

[0008] In one aspect, there is provided a diagnostic kit or device which is a computerized system comprising :

a) a bio assay module configured for detecting gene expression for a tumor sample based on a gene set comprising at least 4 genes selected from the genes of table 3 designated as "up regulated genes in grade 3 tumors" and,

b) a processor module configured to calculate gene-expression grade index (GGI) or relapse score(RS) based on the gene expression and to generate a risk assessment for the tumor sample.

[0008a] In another aspect, there is provided a method comprising the step of :

a) measuring gene expression in a tumor sample,

b) calculating gene-expression grade index (GGI) of the tumor sample by using the formula :

$$\sum_{j \in G_3} x_j - \sum_{j \in G_1} x_j$$

wherein:

x is the gene expression level of mRNA,

G1 and G_3 are sets of genes up-regulated in HG1 and HG3, respectively, and

j refers to a probe or probe set, wherein the gene set comprises or corresponds the gene set comprising at least 4 genes selected from the genes of table 3 designated as "up regulated genes in grade 3 tumors".

[0008b] Disclosed is a method, comprising the steps of

(a) measuring gene expression in a tumor sample submitted to an analysis and obtained from

a mammal subject,

preferably a human patient;

(b) calculating the gene-expression grade index (or genomic grade)(GGI) of the tumor sample using the formula:

$$\sum_{j \in G_3} x_j - \sum_{j \in G_1} x_j$$

wherein: x is the gene expression level of mRNA, G_1 and G_3 are sets of genes up-regulated in histological grade 1 (HG1) and histological grade 3 (HG3), respectively, and j refers to a probe or probe set.

[0009] The tumor sample may be from tissue afflicted by a cancer selected from the group consisting of breast cancer, colon cancer, lung cancer, prostate cancer, hepatocellular cancer, gastric cancer, pancreatic cancer, cervical cancer, ovarian cancer, liver cancer, bladder cancer, cancer of the urinary tract, thyroid cancer, renal cancer, carcinoma, melanoma, or brain cancer. Preferably, the tumor sample is a histological grade HG2 breast tumor sample.

[0010] This embodiment may further comprise designating the tumor sample as low risk (GG1) or high risk (GG3) based on the gene expression grade index (GGI). This embodiment may further comprise providing a breast cancer treatment regimen for a patient consistent with the low risk or high risk designation of the breast tumor sample submitted to the analysis.

[0011] The gene expression grade index GGI may include cutoff and scale values chosen so that the mean GGI of the HG1 cases is about -1 and the mean GGI of the HG3 cases is about +1. The cutoff value is required for calibration of the data obtained from different platforms applying different scales:

$$GGI = scale\left[\sum_{j\in G_3} x_j - \sum_{j\in G_1} x_j - cutoff\right]$$

[0012] The G_1 gene set may comprise at least one gene selected from the genes in Table 3 designated as "Up-regulated in grade 1 tumors". Preferably, the G_1 gene set comprises at least 4 of those genes, and may include the entire set. The G_3 gene set may comprise at least one gene selected from the genes in Table 3 designated as "Up-regulated in grade 3 tumors." Preferably, the G_3 gene set comprises at least 4 those genes, and may include the entire set.

[0013] Also disclosed is the method comprising the steps of

(a) measuring gene expression in a tumor sample;

(b) calculating a relapse score (RS) for the tumor sample using the formula:

$$\sum_{i\in G} w_i \sum_{j\in P_i} \frac{x_{ij}}{n_i}$$

wherein: G is a gene set that is associated with distant recurrence of cancer, P_i is the probe or probe set, i identifies the specific cluster or group of genes, w_i is the weight of the cluster i, j is the specific probe set value, x_{ij} is the intensity of the probe set j in cluster i, and n_i is the number of probe sets in cluster i.

[0014] This embodiment may further comprises the step of classifying the said tumor sample based on the relapse score as low risk or high risk for cancer relapse. The cutoff for distinguishing low risk from high risk may be a relapse score (RS) of from -100 to +100 or a relapse score (RS) of from -10 to +10. The relapse may be relapse after treatment with tamoxifen or other chemotherapy, endocrine therapy, antibody therapy or any other treatment method

used by the person skilled in the art. Preferably, the relapse is after treatment with tamoxifen.

[0015] The tumor sample may be from tissue afflicted by a cancer selected from the group consisting of breast cancer, colon cancer, lung cancer, prostate cancer, hepatocellular cancer, gastric cancer, pancreatic cancer, cervical cancer, ovarian cancer, liver cancer, bladder cancer, cancer of the urinary tract, thyroid cancer, renal cancer, carcinoma, melanoma, or brain cancer. Preferably, the tumor sample is a breast tumor sample.

[0016] The patient's treatment regimen may be adjusted based on the tumor sample's cancer relapse risk status. For example (a) if the patient is classified as low risk, treating the low risk patient sequentially with tamoxifen and sequential aromatase inhibitors (AIs), or (b) if the patient is classified as high risk, treating the high risk patient with an alternative endocrine treatment other than tamoxifen. For a patient classified as high risk, the patient's treatment regimen may be adjusted to chemotherapy treatment or specific molecularly targeted anti-cancer therapies.

[0017] The gene set may be generated from an estrogen receptor (or another marker specific of the cancer tissue sample) positive population. The gene set may be generated by a variety of methods and the component genes may vary depending on the patient population and the specific disorder.

[0018] Another embodiment of the present disclosure provides a computerized system or diagnostic device (or kit), comprising: (a) a bioassay module, preferably a bioarray, configured for detecting gene expression for a tumor sample based on a gene set; and (b) a processor module configured to calculate GGI or RS of the tumor sample based on the gene expression and to generate a risk assessment for the

breast tumor sample. The bioassay module may include at least one gene chip (microarray) comprising the gene set. The gene set may include at least one gene, preferably at least 4 genes, selected from the genes in Table 3 designated as "Up-regulated in grade 1 tumors" or may include the entire set. The gene may include at least 4 genes selected from the genes in Table 3 designated as "Up-regulated in grade 3 tumors" or may include the entire set.

Brief Description of the Drawings

[0019] Figure 1 is representing heatmaps showing the pattern of gene expression in the training (panel a) and the validation sets (panel b). The horizontal axis corresponds to the tumors sorted first by HG and then by GGI as the secondary criterion. The vertical axis corresponds to the genes. The GGI values of each tumor and the relapse free survival are indicated underneath. Two groups of genes are found: those that are highly expressed in grade 1 (16 probe sets; highlighted in red) and, reciprocally, those highly expressed in grade 3 (112 probe sets). The GGI values for HG2 tumors cover the range of values for HG1 and HG3, and those with high GGI tend to relapse earlier (red dots).

[0020] Figure 2 shows Kaplan-Meier RFS analysis based on the HG (panel a) and the GG (panel b) for data pooled from the validation datasets 2-5 (table 11). HG1, HG2 and HG3 can be split further into low and high risk subsets by GG, indicating that GG is an improvement over HG (panel c, d and e respectively). ER status identifies some, but not all, of the patients with poor prognosis (panel f).

[0021] Figure 3 shows Kaplan-Meier RFS analysis based on the NPI (a) and the NPI-GG (b) classification. NPI-GG improves the prognostic discrimination in both low (panel c) and high (panel d) risk NPI subsets, but not vice

versa (panels e and f). The Sorlie et al. dataset was excluded from this analysis because of incomplete tumor size information.

[0022] Figure 4 shows a Forest plot for hazard ratios for HG2 patients split into GG1 and GG3, showing consistent results in different datasets Hazard ratios were estimated with Cox proportional hazard regressions, horizontal lines are 95% confidence intervals for the hazard ratio. P values were determined by the log rank test.

[0023] Figure 5 shows distant metastasis free survival (DMFS) analysis based on the 70-gene expression signature (left row, panels a, c and e) and on GGI (right row, panels b, d and f) for data from the Van de Vijver et al. validation study. a) and b) are all patients, c) and d) are node-negative, and e) and f) are node-positive patients. Note that the node-negative subset includes patients used to derive the 70-gene signature.

[0024] Figure 6 represents a genomic grade applied to previously reported molecular subtypes.

[0025] Figure 7 represents Kaplan Meyer survival curves for distant metastasis free survival for GGI (high vs. low).

Detailed Description of the Invention

Definitions

[0026] Most terms scientific, medical and technical terms are commonly understood to one skilled in the art.

[0027] The term "microarray" refers to an ordered arrangement of hybridizable array elements, preferably polynucleotide probes, on a substrate (an insoluble solid support).

[0028] The terms "differentially expressed gene", "differential gene expression" and their synonyms, which

are used interchangeably, refer to a gene whose expression is activated to a higher or lower level in a subject suffering from a disease, specifically cancer, such as breast cancer, relative to its expression in a normal or control subject. The terms also include genes whose expression is activated to a higher or lower level at different stages of the same disease. It is also understood that a differentially expressed gene may be either activated or inhibited at the nucleic acid level or protein level, or may be subject to alternative splicing to result in a different polypeptide product. Such differences may be evidenced by a change in mRNA levels, surface expression, secretion or other partitioning of a polypeptide, for example. Differential gene expression may include a comparison of expression between two or more genes or their gene products, or a comparison of the ratios of the expression between two or more genes or their gene products, or even a comparison of two differently processed products of the same gene, which differ between normal subjects and subjects suffering from a disease, specifically cancer, or between various stages of the same disease. Differential expression includes both quantitative, as well as qualitative, differences in the temporal or cellular expression pattern in a gene or its expression products among, for example, normal and diseased cells, or among cells which have undergone different disease events or disease stages. For the purpose of this invention, "differential gene expression" is considered to be present when there is at least an about two-fold, preferably at least about four-fold, more preferably at least about six-fold, most preferably at least about ten-fold difference between the expression of a given gene in normal and diseased subjects, or in various stages of disease development in a diseased subject.

[0029] Gene expression profiling: includes all methods of quantification of mRNA and/or protein levels in a biological sample.

The term "prognosis" is used herein to refer to the prediction of the likelihood of cancer-attributable death or progression, including recurrence, metastatic spread, and drug resistance, of a neoplastic disease, such as breast cancer.

[0030] The term "prediction" is used herein to refer to the likelihood that a patient will respond either favorably or unfavorably to a drug or set of drugs, and also the extent of those responses, or that a patient will survive, following surgical removal or the primary tumor and/or chemotherapy for a certain period of time without cancer recurrence. The predictive methods of the present invention are valuable tools in predicting if a patient is likely to respond favorably to a treatment regimen, such as, chemotherapy with a given drug or drug combination, and/or radiation therapy, or whether long-term survival of the patient, following surgery and/or termination of chemotherapy or other treatment modalities is likely.

[0031] The term "high risk" means the patient is expected to have a distant relapse in less than 5 years, preferably in less than 3 years.

[0032] The term "low risk" means the patient is expected to have a distant relapse after 5 years, preferably in less than 3 years.

[0033] The term "tumor," as used herein, refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues.

[0034] The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth.

Examples of cancer include but are not limited to, breast cancer, colon cancer, lung cancer, prostate cancer, hepatocellular cancer, gastric cancer, pancreatic cancer, cervical cancer, ovarian cancer, liver cancer, bladder cancer, cancer of the urinary tract, thyroid cancer, renal cancer, carcinoma, melanoma, and brain cancer.

[0035] Raw "GGI" (Gene expression grade index) is the sum of the log expression (or log ratio) of all genes high-in-HG3 - sum of the log expression (or log ratio) of all genes high-in-HG1 and can be written as:

$$\sum_{j \in G_3} x_j - \sum_{j \in G_1} x_j$$

wherein:

x is the gene expression level of mRNA,

[0036] G_1 and G_3 are sets of genes up-regulated in HG1 and HG3, respectively, and j refers to a probe or probe set.

[0037] GGI may include cutoff and scale values chosen so that the mean GGI of the HG1 cases is about -1 and the mean GGI of the HG3 cases is about +1:

$$GGI = scale\left[\sum_{j \in G_3} x_j - \sum_{j \in G_1} x_j - cutoff \right]$$

The cutoff in GGI is 0 and corresponds to the mean of means.

GGI ranges in value from -4 to +4.

Example 1

Material and Methods for development of grade index (GGI)

Patient demographics

[0038] Six datasets of primary breast cancer were used, four of which were publicly available (Table 11) (4, 5, 10, 11). No patient received adjuvant chemotherapy and

some had received adjuvant tamoxifen treatment. Histological grade (HG) was based on the Elston-Ellis grading system. Each institutional ethics board approved the use of the tissue material.

Table 1: Microarray datasets used in this study

Identifier	Institution	N	Microarray Platform	Systemic Treatment	Reference
1. Training set (KJX64)	Karolinska John Radcliffe	24 40	Affymetrix U133A	yes (tamoxifen only)	this paper
2. Validation set (KJ129)	Karolinska John Radcliffe	68 61	Affymetrix U133A	No	this paper
3. Sotiriou *et al.* (NCI)	John Radcliffe	99	cDNA (NCI)	Yes	10
4. Sorlie *et al.* (STNO)	Stanford	80	cDNA (Stanford)	Yes	11
5. van't Veer *et al.* (NKI)	Netherlands Cancer Institute	97	Agilent	No	4
6. Van de Vijver *et al.* (NKI2)	Netherlands Cancer Institute	295 [61 also in 5)]	Agilent	No	5
Total		703			

[0039] The samples from Oxford were processed at the Jules Bordet Institute in Brussels, Belgium, and those from Sweden at the Genome Institute of Singapore in Singapore. RNA extraction, amplification, hybridization and scanning were done according to standard Affymetrix protocols. Affymetrix U133A Genechips (Affymetrix, Santa Clara, CA). Gene expression values from the CEL files were normalized using RMA (12).

[0040] The default options (with background correction and quantile normalization) were used. The output were in logarithmic scale.

[0041] The normalizations were done separately for .CEL files from different institutions and batch of measurements. In subsequent analysis, the expression data matrices were treated as if they were "blocks" of separate studies. The training set KJX64 consisted of two blocks (corresponding to two different institutions), and so did the validation set KJ129.

[0042] **STNO** The Stanford/Norway dataset (Sorlie et al., 2001) was downloaded from http://genome-www.stanford.edu/breast.cancer/mopo.clinical/data.shtml

[0043] It consists of 85 arrays, with several different chip designs. Only the probes that are common to all were used. The gene expression values used are from the column LOG RAT2N MEAN in the array data files. No further transformation is applied prior to computing the GGI. When more than one spot corresponds to a probe, their average was used.

[0044] All 85 patients were used in the heatmap, but only those with non missing and non zero follow up time were used in survival analysis. This dataset was excluded from analysis involving tumor size, since this information

was not available (Only TNM category was given, but the conversion to tumor size is not straightforward, particularly when one is concerned with what is appropriate for the NPI formula).

[0045] **NKI/NKI2** The data set NKI (van't Veer et al., 2002) and NKI2 (van de Vijver et al., 2002) were downloaded from Rosetta website www.rii.com. The log ratio was used without further transformation. For NKI2, flagged expression values were considered missing. Age, tumor size, and histological grade were not available for NKI2.

[0046] The field 'conservFlag' in the clinical data table were used to stratify the dataset into two groups. Each group had its own threshold for deciding 'good' vs 'poor' prognosis, as was done for in the original results in van de Vijver et al. (2002).

[0047] **NCI** This dataset from Sotiriou et al. (2003) was downloaded from the PNAS web site http://www.pnas.org/cgi/content/full/1732912100/DC1. The expression values were not modified.

Statistical analysis

[0048] Gene selection was done only on the KJX64 dataset, which are all estrogen receptor (ER)-positive and either HG1 or HG3. Dataset KJ129 (43 ER-negative, all node-negative, no systemic treatment) was used as the validation set, along with other previously published data (see table 11). ER-positive tumor s were used for the training set, because ER-status and grade were not independent, with very few ER-negative, HG1 tumor s. Using all HG1 and HG3 tumor s regardless of the ER status would have resulted in spurious associations.

[0049] The standardized mean difference of Hedges and Olkin (13), was used to rank genes based on their differential expression with respect to HG1 or HG3. This

meta-analytical score is similar to the t-statistic, but better suited for our training set which consisted of array data originating from two different centres.

[0050] To control for multiple testing, the *maxT* algorithm of Westfall and Young (14), with an extension proposed by Korn *et al.* (15), was applied to compute false discovery counts (FDC). All 22,283 probe sets were considered. Probe sets having a family-wise error rate *p*-value lower than 0.05 with FDC>2 were identified. Mapping of probes between platforms was done through Unigene (build #180), according to the method in Praz *et al.* (16).

[0051] The gene-expression grade index (GGI) is defined as:

$$GGI = scale\left[\sum_{j \in G_3} x_j - \sum_{j \in G_1} x_j - cutoff\right]$$

where x is the logarithmic gene expression measure, and G_1 and G_3 are the sets of genes up-regulated in HG3 and HG1, respectively. These sets differed across platforms. For convenience, the cutoff and the scale were chosen so that the mean GGI of the HG1 cases was -1 and that of the HG3 cases was +1. This rescaling was done separately for each data source.

[0052] The Nottingham Prognostic Index (NPI) was calculated according to Todd *et al.* (17):

NPI= 0.2 x size [cm] + lymph node status + histological grade.

[0053] An index called NPI/GG was defined, where HG was replaced by GG. Cases with NPI ≥ 3.4 to be high risk in both NPI and NPI/GG were considered. Survival data were visualized using Kaplan-Meier plot. The hazard ratios (HR) were estimated using Cox regression, stratified by the data source. Assumption-free comparisons were done using the stratified log rank test.

Heat maps

[0054] For visualization, the values used in the heatmaps for each probe were meancentered across patients. No genespecific scaling (standardization) was done, in order to keep the information about the relative signal strength of all probes. The color tone were calibrated such that saturated red and green were reached at the values three times the standard deviation of the expression values of the entire matrix. Note that the scaled GGI values were not affected by genespecific centering.

Survival Analysis

[0055] The survival package for R was used by Terry Therneau and a custom program for the KaplanMeier plots, which was checked against the output of the survival package for correctness.

Mapping across microarray platforms

[0056] The approach of CleanEx database (http://www.cleanex.isbsib.ch), described in Praz et al. (2004) was used. Probe identifies were first mapped into sequence accession number. Unigene (build 180) were then used to map the correspondence between platforms. For Affymetrix chips, probesets which contain oligos that were ambiguously mapped to more than one Unigene id were excluded.

Results

Differentially expressed genes between high and low grade subsets

[0057] 242 Probe sets corresponding to 183 unique genes with FDC>2 at family-wise error rate *p*-value of 0.05, corresponding to a low false discovery proportion of 0.008 were identified (Table 3). Of these, a list of 128 probe

sets (97 genes) based on a more conservative criterion (FDC>0 at *p*-value of 0.05) was used in all subsequent analyses, except for checking common genes with signatures published by others, where we used the 183-gene list.

[0058] Figure 1a shows two strong and reciprocal patterns of expression clearly associated with HG1 and HG3. Many genes up-regulated in HG3 were mostly associated with cell cycle progression and proliferation (Table 3). The same gene selection algorithm to contrast HG2 tumor s with a pool combining HG1 and HG3 tumor s were applied. This yielded no differentially expressed genes. Thus, the HG2 population as a whole has no peculiar characteristics of its own that are independent from the HG1 and HG3 distinction.

[0059] The list of 128 probe sets was then applied to untreated breast cancer patients (dataset KJ129). As shown in Fig. 1b, visual inspection revealed an expression pattern for HG1 and HG3 similar to that which was observed on the training set (Fig.1a). The GEP of the grade 2 population looked like a mixture of grade 1 and grade 3 cases, rather than intermediate between the two. To make this observation more objective, the GGI (which essentially summarizes the differences in the GEP of the reporting genes by averaging their expression levels) was defined. As shown under the heat maps in Fig. 1, the GGI distribution of HG2 covered the range of the GGI values of HG1 and HG3, confirming the visual impression. A similar observation was made on the three previously published datasets, despite differences in the clinical populations and microarray platforms (see figures 6a, b, and c).

Histological grade, gene-expression grade (GG) and prognosis

[0060] These findings lead to showing that intermediate histological grade can be replaced by low and high grade based on gene expression. Gene-expression grade (GG) based on the GGI score was defined. Patients were classified as GG1 (low grade) if their GGI value was negative or as GG3 (high grade) otherwise. Note that the GGI score of zero corresponds to the midpoint between the average GGI values of HG1 and HG3 (see methods). This choice might not be clinically optimal and could be improved based on the trade-off between the cost of treatment and risk, but it would be sufficient for evaluating the prognostic value of GGI.

[0061] For this purpose, breast cancer samples derived from a pool of our own validation population (KJ129) and additional datasets STNO, NCI and NKI (table 11) were used. In figure 2a, the association between histological grade and relapse-free survival (RFS) was examined. As expected, HG3 tumor s had significantly worse RFS than HG1, while HG2 tumor s had an intermediate risk and constituted 38% of the population. In figure 2b, GG1 and GG3 subgroups showed distinct RFS, similar to the RFS of HG1 and HG3 tumor s, respectively. To examine how the discordance between GG and HG are related to prognosis, GG was split for each of the histological categories (figure 2c, 2d and 2e). The most striking result was that GG split HG2 into two groups, namely HG2/GG1 and HG2/GG3, whose RFS were also respectively similar to those of HG1 and HG3 (Fig. 2d). The log rank test failed to reveal any significant difference in survival between HG1 and HG2/GG1, as well as between HG3 and HG2/GG3 (see figure 7). For comparison, ER status also had prognostic power in HG2 tumor s (Fig, 2f), although the hazard ratio was less than

that of GG (Fig. 2d). Notably, the ER-positive group showed similar RFS as the total population.

[0062] While GG was better than HG by classifying some patients with poor prognosis in the HG1 population (fig. 2c), the reverse seems to be the case in HG3 population: it classified some patients as low-risk despite their poor prognosis (fig. 2d). Thus, in the case of discordance involving low and high grade categories, neither GG nor HG were consistently outperform the other. It seemed that whichever decided to classify as high grade tended to be more accurate prognostically. This suggests that for both HG and GG, correctly detecting any indication of high grade was easier than accurately declaring it absent. If this observation is confirmed by future studies, corrections should be done in clinical practice, for example by using a rule which substitutes HG1 and HG2, but not HG3, by GG. However, the frequency of this type of discordance in the data used here was relatively small and such modifications were not used in this study, which aims to characterize GG purely on its own.

Table 12: Multivariate analysis of breast cancer prognostic factors (*N* = 302)

	Univariate analysis		Multivariate analysis	
	Hazard ratio (95%CI)	***p***	**Hazard ratio (95%CI)**	***p***
Gene-Expression Grade				
GG3 vs GG1	2.97 (2.03 – 4.37)	0.0001	2.29 (1.44 – 3.63)	0.0004
Histological Grade				
2 + 3 vs 1	1.93 (1.15 – 3.28)	0.0150	0.85 (0.46 – 1.57)	0.61
3 vs 1 + 2	2.03 (1.41 – 2.92)	0.0001	1.25 (0.80 – 1.94)	0.33
Estrogen Receptor				
Negative vs Positive	1.76 (1.24 – 2.49)	0.0016	1.19 (0.81 – 1.76)	0.38
Nodal Status				
Positive vs Negative	2.53 (1.34 – 4.78)	0.0040	1.95 (1.01 – 3.73)	0.045
Tumor Size				
> 2cm vs ≤2cm	2.06 (1.41 – 3.03)	0.0002	1.63 (1.10 – 2.43)	0.015
Age (years)				
≤50 vs >50	0.99 (0.69 – 1.42)	0.97	1.13 (0.78 – 1.63)	0.53

Prognostic value of GG in multivariate model

[0063] Almost all clinicopathological variables were significantly associated with clinical outcome in univariate analysis (Table 12). GG and HG status had the strongest effect. However, in multivariate analysis, only GG, nodal status and tumor size kept their significance, with GG having the largest hazard ratio. In accordance with figure 2, GG replaced HG when both were considered, and GG considerably reduced the prognostic impact of ER.

GG and the Nottingham Prognostic Index

[0064] The independence of GG, nodal status and tumor size in explaining the disease outcome mirrored the Nottingham Prognostic Index (NPI), which combines HG, nodal status and size. To test whether GG can be used to improve this well-characterized risk score, we propose a score called NPI/GG, which is analogous to NPI except that HG is replaced by GG, with only two possible values (either 1 or 3). As shown in Fig. 3a and 3b, NPI/GG was significantly more discriminative than classical NPI. Moreover, NPI/GG was able to split both the NPI low and high risk groups into subgroups with significantly different clinical outcome (Fig. 3c, 3d), while the reverse was not true (Fig. 3e, 3f).

Example 2

Consistent prognostic value of GG in different populations and microarray platforms

[0065] The results of the pooled analysis above were consistently present in the individual datasets, as shown by the forest plot of hazard ratios in Figure 4. More complete results are shown in figure 8. Figure 4 shows that in each independent validation dataset, GG divided the grade 2 populations into two distinct groups with

statistically different clinical outcomes. There was no significant heterogeneity between the hazard ratios, even though the different datasets included heterogeneous patient populations, were graded by various pathologists and used different microarray platforms.

Relationship with the 70-gene signature

[0066] In their pioneering work, van't Veer *et al.* identified a 70-gene expression signature significantly correlated with distant metastasis in node negative breast cancer patients (5). The present list of 97 genes (128 probe sets) could be mapped to 93 genes (113 probes) in their Agilent arrays. To allow comparison under the same trade-off between risk and the cost of treatment as the Netherlands Cancer Institute (NKI) classification, cutoffs for GGI that gave the same numbers of patients in high- and low-risk groups were selected (see methods). Figure 5 shows the comparisons between the NKI prognostic signature and the GGI on distant-metastases-free survival for the overall population (fig.5a, b), as well as for the node negative (fig.5c, d) and positive subgroups (fig.5e, f). Despite the fact that our probes were selected without using clinical outcome and had to be mapped across platforms, the results were strikingly close. Similar results were found when considering overall survival (see figures 9). Data were unavailable to compare relapse-free survival.

[0067] Low and high grade breast cancers were unexpectedly associated with many differentially expressed genes, the majority being involved in cell cycle and proliferation. For these genes, HG2 tumor s had heterogeneous transcriptional profiles that covered the range of variation of HG1 and HG3 tumor s. A similar observation was made in at least one previous report (18). Here, the clinical implications of this finding and

discovered that the grade-related GEPs were also correlated with disease outcome are investigated.

[0068] As demonstrated by Figure 4 improvements by GG were consistent across the different datasets which would have not been the case if the grading quality differed significantly between these studies. Similarly, figure 2a shows good prognostic separation between HG1 and HG3, indicating that the histological grading was of high quality. Furthermore, central pathologist review would still result in a significant portion of tumor s being classified as HG2. Finally, these results were more reflective of clinical reality, since grading by a central pathologist is rarely done in practice.

[0069] The approach in identifying GEP associated with prognosis is quite different from that used by other investigators. Instead of selecting the prognostic genes directly through their correlation with survival, one may identify them indirectly through histological grade, a well-established prognostic factor rooted in cell biology. This may explain the robustness and reproducibility of GGI across independent and heterogeneous validation sets and different microarray platforms. Furthermore, since the GGI can be interpreted as "molecular grade", it can be integrated easily into existing prognostic systems which uses histological grade, such as the NPI.

[0070] This gene selection process was not meant to define a specific set of genes to be used as a prognostic "signature". The present invention aims to build a comprehensive "catalogue" where different sets of signatures could be chosen from. This was illustrated by the cross-platform applicability of the catalogue. Although the actual sets of probes used in various platforms differed in numbers and gene compositions, the results were still reproducible. It is remarkable to obtain good

prognostic discrimination in very different datasets with a linear classifier where the weights of the genes were simply +1 or -1, based on their association with grade on a training set of 64 patients. Thus, the "grade signal" identified was not bound to a particular set of genes nor to any special combination of their expression levels, since the genes were highly correlated and the GGI effectively behaves as a single prognostic factor. It is still beneficial to use many genes, if only to provide redundancy against noise. The consequence for the development of practical diagnostic systems is that arbitrary subsets of the "grade gene catalogue" of the invention might be used, constrained only by technical considerations.

[0071] Jenssen and Hovig (19) recently discussed two issues regarding the use of gene-expression signatures for prognosis. These were 1) the lack of agreement between genes included in different signatures and 2) the difficulty in understanding the biological basis of the correlation between the signatures and survival. The present gene catalogue is rich in genes with likely roles in cell cycle progression and proliferation. This class of genes is one important –if not the most important– component of any existing profile-based risk prediction method for breast cancer. In Paik *et al.*(7), the "proliferation set", whose five genes are all in our 183-gene catalogue (Table 3), was the one that had the largest hazard ratios in their extensive training and validation sets and has the highest weight in the "recurrence score" formula. The application to the NKI data in figure 5 also lends support to the idea that grade-related genes may constitute a significant portion of the prognostic power of the NKI 70-gene signature. When compared against our 183-gene catalogue, the following numbers of genes in common

with other prognostic signatures: 11/70 and 30/231 genes (van't Veer et al.), 5/15 (Paik et al) and 7/76 (Wang et al.)(4, 7, 8) were found.

[0072] In summary, gene-expression based grading could significantly improve current grading systems for the prognostic assessment of cancer, in particular breast cancer.

[0073] Reproduction of these findings across multiple independent datasets and across different platforms suggests our conclusions are robust. The GGI score does not require a specific set of genes nor is it bound to a particular detection platform. Grading based on the GGI can be incorporated into existing prognostic systems, by substituting HG with GG. Refined grading based on gene expression measurements could have important clinical application for breast cancer management in the future.

Example 3

Definition of clinically distinct subtypes within Estrogen receptor positive breast carcinoma

Materials and methods

Tumor samples

[0074] Three hundred and thirty five early-stage breast carcinoma samples comprised our own dataset. Eighty-six of these samples have been previously used in another study and the raw data are available at the Gee Expression Omnibus repository database (http://www.ncbi.nlm.nih.gov/geo), with accession code GSE2990. These samples had received no adjuvant systemic therapy. Two hundred and forty-nine samples, previously unpublished, had received adjuvant tamoxifen only (tam-treated dataset). All samples were required to be ER-positive by protein ligand binding assay.

[0075] Microarray analysis was performed with Affymetrix™ U113A Genechips® (Affymetrix, Santa Clara, CA). This dataset contained samples from the John Radcliffe Hospital, Oxford, U.K., Guys Hospital, London, U.K. and Uppsala University Hospital, Uppsala, Sweden. Samples from Oxford and London were processed at the Jules Bordet Institute in Brussels, Belgium. For the samples from Uppsala, RNA was extracted at the Karolinska Institute and hybridized at the Genome Institute of Singapore in Singapore. The quality of the RNA obtained from each tumour sample was assessed via the RNA profile generated by the Agilent bioanalyzer. RNA extraction, amplification, hybridization, and scanning were done according to standard Affymetrix protocols. Gene expression values from the CEL were normalized by use of RMA [12]. Each population was normalised separately. Each hospital's institutional ethics board approved the use of the tissue material and written informed consent was obtained. The raw data for the tam-treated dataset are available at the Gene Expression Omnibus repository database (http://www.ncbi.nlm.nih.gov/geo/), with accession code GSE XXX.

[0076] The inventors also used four other publically available datasets, described in recent publications: van de Vijver[5] (n=295), Wang[8] (n=286), Sotiriou[10] (n=99), Sorlie[11] (n=78), in the analysis. For the survival analysis, we used tumors classified as ER-positive only (van de Vijver[5] (n=122), Wang[8] (n=209)). For the survival analysis involving patients who had received no systemic adjuvant treatment, patients from the van de Vijver *et al.*[5], Wang *et al.*[8] and previously published dataset were combined (n=417 ER-positive patients, hereby referred to as the "untreated" dataset)

All clinical data are shown in Table S1 of the Supplementary Information.

Data Analysis

Estrogen (ER) and Progesterone receptor (PgR) level

[0077] Patients were initially selected at their institutions according to a positive ER status which was determined by protein ligand-binding assay. The inventors subsequently confirmed a positive ER level by using the microarray data. The ER level was measured by probe set (a 30-mer oligonucleotide) on our human Affymetrix™ GeneChip® U133 A&B microarray. The inventors have used the probe set "205225_at" for ER. PgR was represented by the probe set "208305_at". The immunohistochemical measurement of ER is known to correlate with mRNA levels of ER [4]. Tumours with any positive expression level of ER and PgR were considered.

Histological grade

[0078] Histological grade was based on the Elston-Ellis grading system. A central pathologist reviewed the histological grade and ER status for all samples from Uppsala, Sweden, Guys Hospital, London, UK and the Van de Vijver *et al.* dataset [5].

An index based on the expression of proliferation-related genes to quantify genomic grade: gene expression grade index (GGI)

[0079] "Gene expression grade index" (GGI) is a linear combination of the expression of 128 probe sets (97 genes) that were found to be differentially expressed between histological grade 1 and 3 (see definitions). The index is effectively, a quantification of the degree of similarity between the tumour expression profile and tumour

grade. A high gene-expression grade index corresponds to a high grade and vice versa. This index was used to divide each data set into high and low grade sub-groups.

[0080] Mapping of probes between microarray platforms was done through Unigene (build #180), according to the method in Praz *et al.*[16].

Hierarchical clustering

The "Cluster" program was used to perform average linkage hierarchical cluster analysis [28] after median centering of each gene using an uncentered Pearson correlation as similarity measurement. The cluster results were viewed using "TreeView". Expression data was downloaded and extracted from datasets Sorlie *et al.*[11] and Sotiriou *et al.* [10]. The samples were ordered according to subtype as in the original publications [10, 11] to investigate the relation between the expression of the genes in the GGI and the subtypes.

Statistical Analysis

[0081] In order to assess the relation between survival and some continuous variable, a variant of a method introduced to compute the expected survival for individual was used: "Rate of distant recurrence" plots [29] (ref: Terry M. Therneau and Patricia M. grambsch, 2000, "Modeling Survival Data: Extending the Cox Model", chapter 10). The expected proportion of distant metastasis with respect to the GGI, ER and PgR was plotted using a Cox model fitted with only the variable under study.

[0082] Survival curves were visualized using Kaplan-Meier plots and compared using log-rank tests. The univariate and multivariate hazard ratios (HR) were estimated using Cox regression analysis. All statistical

tests were two-sided. Statistical analysis was performed using SPSS statistical software package, version 11.5.

RESULTS

Applying genomic grade to the previously reported molecular subtypes

[0083] To investigate the expression of the gene expression grade index (GGI) in relation to the subtypes, expression data were extracted from data sets Sorlie and Sotiriou *et al.*, the original and confirmatory publications respectively [11, 13]. The genes were clustered using average-linkage clustering and the samples were ordered according to the subtypes as presented in the published manuscripts [11, 13]. Applying genomic grade to the previously reported molecular subtypes (6a: Sorlie et *al.*; 6b: Sotiriou et *al.*) Subtypes are ordered the same as in the original publications. The heatmap of GGI genes is placed below the dendrogram. Boxplots of the GGI score (median and range) are placed below each subtype. High grade is indicated by a GGI score >1 and vice versa.

[0084] Figure 6 shows the results of this analysis. In general, the ER-negative subtypes, the basal and the erbB2 subtypes, had high expression of GGI, or were of high grade. However, the ER-positive subtypes showed a diverse range of GGI levels, particularly the luminal C or 3 subtype both highly expressing these proliferation-associated genes, whereas luminal A or 1, and the normal-like were mostly negative for the expression of the GGI, or low grade. This confirmed the hypothesis that there are varying degrees of contribution of cell cycle genes to the biological makeup of ER-positive tumours, whereas ER-negative tumours seem to consistently have over-expression of these genes. It is interesting to note the similarity in

expression profiles of the GGI genes between the high grade ER-positive subtype and the ER-negative subtypes.

Clinical relevance of ER-positive luminal subtypes as defined by genomic grade

[0085] Genomic grade could distinguish clinically subtypes within the ER-positive tumours and the prognostic value of these genomic grade defined subtypes were an improvement over current traditional methods, such as that based on quantitative levels of estrogen and progesterone receptor levels. A Kaplan-Meier survival analysis was performed comparing classes of ER-positive tumours according to GGI score (high vs. low grade) and expression levels of estrogen and progesterone receptor (rich vs. poor expression) with respect to time to distant metastasis (TDM), which is often used as a surrogate for breast cancer specific survival (Figure 7- KM and Cox). Kaplan Meier survival curves for distant metastasis free survival for GGI (high vs. low), ER expression levels and PgR expression levels (rich vs. poor). Figure 7a displays the results for the untreated dataset (n=417). Figure 7b for the tamoxifen-treated dataset (n=249). For the untreated dataset, results shown were combined from multiple datasets involving 417 ER-positive samples hybridized using two popular commercially available oligonucleotide microarray platforms- Affymetrix™ and Agilent™ (see methods). As shown, for both untreated and tamoxifen-treated populations, the expression levels of the ER did not have any prognostic value ($p=0.74$ and 0.51 respectively). In contrast, both the GGI and expression levels of the PgR had prognostic value (untreated: $p<0.0001$ for both GGI and PgR; tam-treated: GGI $p<0.0001$, PgR $p=0.0058$). The luminal low grade subtype had a much better 10-year estimate of TDM compared with the luminal high grade subtype.

[0086] Table 13 shows the univariate and multivariate analysis with other standard prognostic covariates of age, grade, tumour size as well as genomic grade. In the multivariate Cox regression analysis, only the GGI retained significant prognostic value (untreated: HR 2.3 (95%CI: 1.2-4.3; p=0.008; tam-treated: HR 2.14 (95%CI: 1.04-4.02; p=0.0038), subsuming those factors that were significant at the univariate level, including the progesterone receptor expression levels (p=0.3). For the untreated population, tumour size also retained significance in the multivariate model (HR 2.2 (95% CI:1.2-3.8, p=0.0068). This suggests that genomic grade, as measured by the GGI, can distinguish clinically distinct groups of patients within those that express positive levels of estrogen receptor. Furthermore, the GGI had highly significant prognostic value, suggesting a better ability to discriminate clinical outcome over these traditional factors. The ER-positive high grade subgroup's worse disease outcome in the tamoxifen-treated dataset seems to suggest that adjuvant tamoxifen does not alter this subtype's natural disease history despite having a positive ER status. This could potentially flag a group of tumours worthy of further investigation from both a biological and therapeutic standpoint.

[0087] As further demonstration of the GGI's prognostic value in ER-positive tumours, the inventors generated figures displaying the rate of distant recurrence as continuous function of the GGI and compared this to continuous levels of ER and PgR for both untreated and tam-treated populations.

[0088] Two subtypes of tumours can be distinguished within patients whose breast cancers express at least some level of estrogen receptor. In patients whose tumours express a high level of the genes that comprise the GGI,

i.e. corresponding to high genomic grade, their disease outcome was clearly different, with a higher incidence of relapses compared with tumours of low genomic grade. Furthermore, their worse disease outcome seemed unchanged even when given adjuvant tamoxifen, suggesting that this group of women do not seem to benefit from adjuvant tamoxifen despite their positive estrogen receptor values. Note that none of the patients in this study had received adjuvant chemotherapy, so it is unclear if chemotherapy can alter this group's natural disease history. The potential clinical significance of this finding is also underscored by the similarities between the high grade ER-positive group and the high grade ER-negative tumours (basal and erbB2), further suggesting that high levels of expression of the genes associated with high genomic grade is associated with a poor prognosis. The GGI can consistently identify these two groups across multiple datasets which were hybridized using several microarray platforms, involving 666 ER-positive samples, suggesting our conclusions are robust and highly reproducible than that produced previously by hierarchical cluster analysis [1,3].

[0089] The genes present in the GGI are associated with cell cycle progression and proliferation: among the top 20 overexpressed genes were UBE2C, KPNA2, TPX2, FOXM1, STK6, CCNA2, BIRC5, and MYBL2; see Supplemental Table 14). For ER-positive tumours, genomic grade was associated with differing relapse-free survival, but for ER-negative tumours, as almost all are associated with high genomic grade, the GGI had no prognostic value. Therefore, cell-cycle related genes seem to have prognostic value only in breast cancer patients with positive expression of ER. Within this group, the incidence of distant metastases seems to be predominantly driven by this set of proliferation and grade-derived genes. However, in ER-

negative tumours, there may be further factors driving the underlying biology of metastasis besides cell-cycle associated genes. The prognostic ability of a "cell proliferation signature" in a subset of patients has been reported previously in women who express relatively high estrogen receptor expression for their age [5]. The analysis of the ER-positive subgroups was divided by genomic grade to the previously described luminal subgroups and this concept was validated in over 650 patients. Furthermore, genomic grade remains the strongest variable in univariate and multivariate analysis (Table 4) that takes clinical prognostic factors into consideration.

[0090] Currently there are several molecular signatures derived from microarray technology that claim to be able to predict prognosis in breast cancer patients [8, 4, 7, 24] Some of these gene signatures reported can predict clinical outcome in ER-positive tumours treated with adjuvant tamoxifen [7, 24, 30]. In the recurrence score developed by Paik *et al.* [7] the proliferation set of five genes had the largest hazard ratios in their large training and validation sets and the highest "weight" or coefficient in their recurrence score formula indicating their high importance in deriving a prognosis classification for women with early stage breast cancer treated with adjuvant tamoxifen. Proliferation-related genes appear to be an important—if not the most important—component of many existing prognostic gene signatures for breast cancer that are based on gene-expression profiles. By using the 11 genes in common between the GGI and a 70-gene prognostic gene classifier for women with early stage breast cancer under the age of 55 [4], similar survival curves to the validation publication [5] were obtained, suggesting that grade-related genes constitute a significant amount of the prognostic power of this signature. The subgroups achieved

by these prognostic signatures and that obtained by the classification of ER-positive tumours by genomic grade overlap significantly because of a strong dependence on cell-cycle genes to drive metastasis and relapse. The advantage of this approach is that the biological mechanism that is responsible for the poor outcome is obvious, rather than a gene set that likely represents a variety of molecular functions and biological processes [8, 4]. Because antiestrogens such as tamoxifen have a cell cycle-specific action on breast cancer cells and influence the expression and activity of several cell cycle-regulatory molecules, the development of aberrant cell cycle control mechanisms is an obvious mechanism by which cells might develop resistance to antiestrogens. It is currently incompletely understood why up to 30-40% of ER-positive breast cancers develop resistance to tamoxifen when positive expression of the ER is the best predictor predictors of tamoxifen response in the clinical setting [31]. Over-expression of cyclin D1, a critical controller of the cell cycle, has been associated with tamoxifen resistance and can reverse the growth-inhibitory effect of antiestrogens in estrogen receptor-positive breast cancer cells [32]. Further investigation into the oncogenic pathways that drive the cell cycle machinery will be beneficial in developing new agents to treat the high grade subgroup.

[0091] Definition of clinically relevant tumour subclasses within ER-positive breast cancers is of great importance to the treating oncologist today. The emergence of new strategies of adjuvant anti-estrogen therapy [33-37] as well as new chemotherapeutic and biological agents has made treatment decision making for women with early stage breast cancer sometimes a difficult task. Previously, tamoxifen was the mainstay of anti-estrogen therapy, with significant reductions in the risk or relapse, death and contralateral

breast cancer for women with early stage, ER-positive breast cancer [38]. However, since the advent of aromatase inhibitors and the reporting of several trials finding them to be more effective than tamoxifen in postmenopausal women, the American Society of Clinical Oncology has recommended that an aromatase inhibitor be included in the therapy of postmenopausal women with early stage hormone responsive breast cancers [39]. However, it is still unclear the best combination and sequencing of aromatase inhibitors and tamoxifen, and whether all women with ER-positive tumours derive the same or differing benefit from these agents. The elucidation of clinically relevant and biological distinct hormone responsive breast tumour phenotypes can help facilitate the optimization of such therapy as they may require different therapeutic strategies.

[0092] In conclusion, the use of genomic grade can distinguish two subtypes with ER-positive breast cancers in a reproducible manner across multiple datasets and microarray platforms. This is validated ept in over 650 ER-positive breast cancer samples. These subgroups have statistically distinct clinical outcome in both systemically untreated and tamoxifen-only treated populations. Stratification by subtype in clinical trials may provide important information on the potentially diverse effect of endocrine therapies, chemotherapies and biological agents on these subgroups. A focussed biological investigation into these distinct phenotypes may result in identification of separate and different therapeutic targets.

[0093] The genes identified herein may be used to generate a model capable of predicting the breast cancer grade of an unknown breast cell sample based on the expression of the identified genes in the sample. Such a

model may be generated by any of the algorithms described herein or otherwise known in the art as well as those recognized as equivalent in the art using gene(s) (and subsets thereof) disclosed herein for the identification of whether an unknown or suspicious breast cancer sample is normal or is in one or more stages and/or grades of breast cancer. The model provides a means for comparing expression profiles of gene(s) of the subset from the sample against the profiles of reference data used to build the model. The model can compare the sample profile against each of the reference profiles or against model defining delineations made based upon the reference profiles. Additionally, relative values from the sample profile may be used in comparison with the model or reference profiles.

[0094] In a preferred embodiment of the invention, breast cell samples identified as normal and non-normal and/or atypical from the same subject may be analyzed for their expression profiles of the genes used to generate the model. This provides an advantageous means of identifying the stage of the abnormal sample based on relative differences from the expression profile of the normal sample. These differences can then be used in comparison to differences between normal and individual abnormal reference data which was also used to generate the model. The detection of gene expression from the samples may be by use of a single microarray able to assay gene expression. One method of analyzing such data would be from all pairwise comparisons disclosed herein for convenience and accuracy.

[0095] Other uses of the present invention include providing the ability to identify breast cancer cell samples as being those of a particular stage and/or grade of cancer for further research or study. This provides a particular advantage in many contexts requiring the

identification of breast cancer stage and/or grade based on objective genetic or molecular criteria rather than cytological observation. It is of particular utility to distinguish different grades of a particular breast cancer stage for further study, research or characterization.

[0096] The materials for use in the methods of the present invention are ideally suited for preparation of kits produced in accordance with well known procedures. The invention thus provides kits comprising agents for the detection of expression of the disclosed genes for identifying breast cancer stage. Such kits optionally comprising the agent with an identifying description or label or instructions relating to their use in the methods of the present invention, is provided. Such a kit may comprise containers, each with one or more of the various reagents (typically in concentrated form) utilized in the methods, including, for example, pre-fabricated microarrays, buffers, the appropriate nucleotide triphosphates (e.g., dATP, dCTP, dGTP and dTTP; or rATP, rCTP, rGTP and UTP), reverse transcriptase, DNA polymerase, RNA polymerase, and one or more primer complexes of the present invention (e.g., appropriate length poly(T) or random primers linked to a promoter reactive with the RNA polymerase). A set of instructions will also typically be included.

[0097] The methods provided by the present invention may also be automated in whole or in part. All aspects of the present invention may also be practiced such that they consist essentially of a subset of the disclosed genes to the exclusion of material irrelevant to the identification of breast cancer stages in a cell containing sample.

[0098] An exemplary system for implementing the overall system or portions of the invention might include a general purpose computing device in the form of a computer,

including a processing unit, a system memory, and a system bus that couples various system components including the system memory to the processing unit. The system memory may include read only memory (ROM) and random access memory (RAM). The computer may also include a magnetic hard disk drive for reading from and writing to a magnetic hard disk, a magnetic disk drive for reading from or writing to a removable magnetic disk, and an optical disk drive for reading from or writing to a removable optical disk such as a CD ROM or other optical media. The drives and their associated machine-readable media provide nonvolatile storage of machine-executable instructions, data structures, program modules and other data for the computer.

[0099] Embodiments of the present invention may be practiced in a networked environment using logical connections to one or more remote computers having processors. Logical connections may include a local area network (LAN) and a wide area network (WAN) that are presented here by way of example and not limitation. Such networking environments are commonplace in office-wide or enterprise-wide computer networks, intranets and the Internet and may use a wide variety of different communication protocols. Those skilled in the art will appreciate that such network computing environments will typically encompass many types of computer system configurations, including personal computers, hand-held devices, multi-processor systems, microprocessor-based or programmable consumer electronics, network PCs, minicomputers, mainframe computers, and the like. Embodiments of the invention may also be practiced in distributed computing environments where tasks are performed by local and remote processing devices that are linked (either by hardwired links, wireless links, or by a

combination of hardwired or wireless links) through a communications network. In a distributed computing environment, program modules may be located in both local and remote memory storage devices.

[0100] Different embodiments of the present invention have been described according to the present invention. Many modifications and variations may be made to the techniques and structures described and illustrated herein without departing from the spirit and scope of the invention. Accordingly, it should be understood that the apparatuses described herein are illustrative only and are not limiting upon the scope of the invention.

Table 3: Up-regulated in grade 3 tumors

No.	D	p-values FD>0	FD>1	FD>2	probeset	gene symbol	description
1	2.1162	0.0001	0.0001	0.0001	202954_at	UBE2C	ubiquitin-conjugating enzyme E2C
2	1.9037	0.0001	0.0001	0.0001	222077_s_at	RACGAP1	Rac GTPase activating protein 1
3	1.7292	0.0001	0.0001	0.0001	201088_at	KPNA2	karyopherin alpha 2 (RAG cohort 1, importin alpha 1)
4	1.7264	0.0001	0.0001	0.0001	218542_at	C10orf3	chromosome 10 open reading frame 3
5	1.7259	0.0001	0.0001	0.0001	203554_x_at	PTTG1	pituitary tumor-transforming 1
6	1.7053	0.0001	0.0001	0.0001	218355_at	KIF4A	kinesin family member 4A
7	1.6600	0.0001	0.0001	0.0001	210052_s_at	TPX2	TPX2, microtubule-associated protein homolog (Xenopus laevis)
8	1.6598	0.0001	0.0001	0.0001	202580_x_at	FOXM1	forkhead box M1
9	1.6548	0.0001	0.0001	0.0001	208079_s_at	STK6	serine/threonine kinase 6
10	1.6513	0.0001	0.0001	0.0001	204092_s_at	STK6	serine/threonine kinase 6
11	1.6495	0.0001	0.0001	0.0001	218755_at	KIF20A	kinesin family member 20A
12	1.6387	0.0001	0.0001	0.0001	201584_s_at	DDX39	DEAD (Asp-Glu-Ala-Asp) box polypeptide 39
13	1.6347	0.0001	0.0001	0.0001	203764_at	DLG7	discs, large homolog 7 (Drosophila)
14	1.6223	0.0001	0.0001	0.0001	204825_at	MELK	maternal embryonic leucine zipper kinase
15	1.6213	0.0001	0.0001	0.0001	203418_at	CCNA2	cyclin A2
16	1.6095	0.0001	0.0001	0.0001	204766_s_at	NUDT1	nudix (nucleoside diphosphate linked moiety X)-type motif 1
17	1.6057	0.0001	0.0001	0.0001	206102_at	KIAA0186	KIAA0186 gene product
18	1.5986	0.0001	0.0001	0.0001	202095_s_at	BIRC5	baculoviral IAP repeat-containing 5 (survivin)
19	1.5957	0.0001	0.0001	0.0001	201710_at	MYBL2	v-myb myeloblastosis viral oncogene homolog (avian)-like 2
20	1.5879	0.0002	0.0001	0.0001	211762_s_at	KPNA2	karyopherin alpha 2 (RAG cohort 1, importin alpha 1)
21	1.5816	0.0002	0.0001	0.0001	209680_s_at	KIFC1	Kinesin family member C1
22	1.5785	0.0002	0.0001	0.0001	209408_at	KIF2C	kinesin family member 2C
23	1.5671	0.0002	0.0001	0.0001	219918_s_at	ASPM	asp (abnormal spindle)-like, microcephaly associated (Drosophila)
24	1.5650	0.0003	0.0001	0.0001	203145_at	SPAG5	sperm associated antigen 5
25	1.5595	0.0003	0.0001	0.0001	204962_s_at	CENPA	centromere protein A, 17kDa
26	1.5551	0.0003	0.0001	0.0001	202870_s_at	CDC20	CDC20 cell division cycle 20 homolog (S. cerevisiae)
27	1.5446	0.0003	0.0001	0.0001	38158_at	ESPL1	extra spindle poles like 1 (S. cerevisiae)
28	1.5376	0.0003	0.0001	0.0001	202107_s_at	MCM2	MCM2 minichromosome maintenance deficient 2, mitotin (S. cerevisiae)
29	1.5236	0.0004	0.0001	0.0001	204767_s_at	FEN1	flap structure-specific endonuclease 1
30	1.5226	0.0004	0.0001	0.0001	203046_s_at	TIMELESS	timeless homolog (Drosophila)
31	1.5221	0.0004	0.0001	0.0001	221677_s_at	DONSON	downstream neighbor of SON
32	1.5134	0.0005	0.0001	0.0001	210559_s_at	CDC2	cell division cycle 2, G1 to S and G2 to M
33	1.5047	0.0006	0.0001	0.0001	221520_s_at	CDCA8	cell division cycle associated 8
34	1.5017	0.0007	0.0001	0.0001	214710_s_at	CCNB1	cyclin B1
35	1.4945	0.0007	0.0001	0.0001	209714_s_at	CDKN3	cyclin-dependent kinase inhibitor 3 (CDK2-associated dual specificity phosphatase)
36	1.4933	0.0008	0.0001	0.0001	204444_at	KIF11	kinesin family member 11
37	1.4927	0.0008	0.0001	0.0001	210821_x_at	CENPA	centromere protein A, 17kDa
38	1.4915	0.0008	0.0001	0.0001	218726_at	DKFZp762E1312	hypothetical protein DKFZp762E1312
39	1.4895	0.0009	0.0001	0.0001	220651_s_at	MCM10	MCM10 minichromosome maintenance deficient 10 (S. cerevisiae)
40	1.4865	0.0010	0.0001	0.0001	201475_x_at	MARS	methionine-tRNA synthetase
41	1.4715	0.0014	0.0001	0.0001	204033_at	TRIP13	thyroid hormone receptor interactor 13
42	1.4672	0.0014	0.0001	0.0001	202705_at	CCNB2	cyclin B2
43	1.4624	0.0014	0.0001	0.0001	204649_at	TROAP	trophinin associated protein (tastin)
44	1.4603	0.0014	0.0001	0.0001	220060_s_at	FLJ20641	hypothetical protein FLJ20641
45	1.4534	0.0016	0.0001	0.0001	209836_x_at	LAT1-3TM	LAT1-3TM protein
46	1.4533	0.0016	0.0001	0.0001	203276_at	LMNB1	lamin B1
47	1.4471	0.0018	0.0001	0.0001	205034_at	CCNE2	cyclin E2
48	1.4455	0.0018	0.0001	0.0001	203213_at	CDC2	cell division cycle 2, G1 to S and G2 to M
49	1.4384	0.0019	0.0001	0.0001	209464_at	AURKB	aurora kinase B
50	1.4381	0.0019	0.0001	0.0001	205046_at	CENPE	centromere protein E, 312kDa
51	1.4373	0.0019	0.0001	0.0001	203755_at	BUB1B	BUB1 budding uninhibited by benzimidazoles 1 homolog beta (yeast)
52	1.4336	0.0019	0.0001	0.0001	203214_x_at	CDC2	cell division cycle 2, G1 to S and G2 to M
53	1.4236	0.0022	0.0002	0.0001	214804_at	FSHPRH1	FSH primary response (LRPR1 homolog, rat) 1
54	1.4167	0.0026	0.0003	0.0001	212949_at	BRRN1	barren homolog (Drosophila)
55	1.4134	0.0027	0.0003	0.0001	204318_s_at	GTSE1	G-2 and S-phase expressed 1

Table 3: ... continued

No.	D	p-values FDR>0	FDR>1	FDR>2	probeset	gene symbol	description
56	1.4105	0.0030	0.0003	0.0001	207165_at	RHMMR	hyaluronan-mediated motility receptor (RHAMM)
57	1.4079	0.0031	0.0003	0.0001	212022_s_at	MKI67	antigen identified by monoclonal antibody Ki-67
58	1.4051	0.0031	0.0003	0.0001	213226_at	CCNA2	cyclin A2
59	1.3931	0.0041	0.0004	0.0001	219510_at	POLQ	polymerase (DNA directed), theta
60	1.3890	0.0044	0.0004	0.0001	204026_s_at	ZWINT	ZW10 interactor
61	1.3890	0.0044	0.0004	0.0001	203432_at	TMPO	thymopoietin
62	1.3872	0.0046	0.0004	0.0001	204768_s_at	FEN1	flap structure-specific endonuclease 1
63	1.3855	0.0047	0.0004	0.0001	209773_s_at	RRM2	ribonucleotide reductase M2 polypeptide
64	1.3847	0.0047	0.0004	0.0001	214431_at	GMPS	guanine monphosphate synthetase
65	1.3842	0.0048	0.0004	0.0001	212023_s_at	MKI67	antigen identified by monoclonal antibody Ki-67
66	1.3752	0.0052	0.0004	0.0002	218883_s_at	MLP1IP	MLP1 interacting protein
67	1.3541	0.0077	0.0006	0.0003	211519_s_at	KIF2C	kinesin family member 2C
68	1.3503	0.0083	0.0006	0.0003	202240_at	PLK1	polo-like kinase 1 (Drosophila)
69	1.3460	0.0089	0.0007	0.0003	205733_at	BLM	Bloom syndrome
70	1.3457	0.0092	0.0008	0.0003	222039_at	LOC146909	hypothetical protein LOC146909
71	1.3443	0.0096	0.0008	0.0003	209642_at	BUB1	BUB1 budding uninhibited by benzimidazoles 1 homolog (yeast)
72	1.3376	0.0102	0.0010	0.0003	213599_at	OIP5	Opa-interacting protein 5
73	1.3372	0.0102	0.0010	0.0003	214096_s_at	SHMT2	serine hydroxymethyltransferase 2 (mitochondrial)
74	1.3348	0.0105	0.0012	0.0003	211072_x_at	K-ALPHA-1	tubulin, alpha, ubiquitous
75	1.3237	0.0130	0.0017	0.0004	202779_s_at	UBE2S	ubiquitin-conjugating enzyme E2S
76	1.3226	0.0133	0.0017	0.0004	218447_at	DC13	DC13 protein
77	1.3215	0.0138	0.0017	0.0004	213911_s_at	H2AFZ	H2A histone family, member Z
78	1.3211	0.0138	0.0017	0.0004	212141_at	MCM4	MCM4 minichromosome maintenance deficient 4 (S. cerevisiae)
79	1.3156	0.0153	0.0019	0.0005	221591_s_at	FLJ10156	hypothetical protein FLJ10156
80	1.3139	0.0162	0.0019	0.0005	204822_at	TTK	TTK protein kinase
81	1.3121	0.0165	0.0020	0.0005	209251_x_at	TUBA6	tubulin alpha 6
82	1.3086	0.0173	0.0023	0.0006	217835_x_at	C20orf24	chromosome 20 open reading frame 24
83	1.3081	0.0176	0.0023	0.0006	201890_at	RRM2	ribonucleotide reductase M2 polypeptide
84	1.3059	0.0184	0.0024	0.0006	213671_s_at	MARS	methionine-tRNA synthetase
85	1.3053	0.0185	0.0024	0.0006	218009_s_at	PRC1	protein regulator of cytokinesis 1
86	1.3010	0.0197	0.0026	0.0007	207828_s_at	CENPF	centromere protein F, 350/400ka (mitosin)
87	1.3002	0.0198	0.0026	0.0007	219555_s_at	BM039	uncharacterized bone marrow protein BM039
88	1.2969	0.0206	0.0026	0.0007	204695_at	CDC25A	cell division cycle 25A
89	1.2953	0.0214	0.0026	0.0009	212021_s_at	MKI67	antigen identified by monoclonal antibody Ki-67
90	1.2898	0.0229	0.0028	0.0009	201090_x_at	K-ALPHA-1	tubulin, alpha, ubiquitous
91	1.2885	0.0233	0.0033	0.0010	218039_at	NUSAP1	nucleolar and spindle associated protein 1
92	1.2851	0.0246	0.0034	0.0012	204603_at	EXO1	exonuclease 1
93	1.2846	0.0248	0.0034	0.0012	203362_s_at	MAD2L1	MAD2 mitotic arrest deficient-like 1 (yeast)
94	1.2845	0.0248	0.0034	0.0012	202094_at	BIRC5	baculoviral IAP repeat-containing 5 (survivin)
95	1.2840	0.0249	0.0034	0.0012	204162_at	KNTC2	kinetochore associated 2
96	1.2825	0.0254	0.0037	0.0012	222036_s_at	MCM4	MCM4 minichromosome maintenance deficient 4 (S. cerevisiae)
97	1.2780	0.0272	0.0039	0.0014	204252_at	CDK2	cyclin-dependent kinase 2
98	1.2775	0.0274	0.0039	0.0014	219000_s_at	DCC1	defective in sister chromatid cohesion homolog 1 (S. cerevisiae)
99	1.2772	0.0277	0.0041	0.0014	201524_x_at	UBE2N	ubiquitin-conjugating enzyme E2N (UBC13 homolog, yeast)
100	1.2694	0.0294	0.0044	0.0018	204817_at	ESPL1	extra spindle poles like 1 (S. cerevisiae)
101	1.2657	0.0313	0.0046	0.0019	218662_s_at	HCAP-G	chromosome condensation protein G
102	1.2620	0.0324	0.0052	0.0022	206364_at	KIF14	Kinesin family member 14
103	1.2612	0.0329	0.0054	0.0022	221436_s_at	CDCA3	cell division cycle associated 3
104	1.2609	0.0331	0.0054	0.0022	201195_s_at	SLC7A5	solute carrier family 7 (cationic amino acid transporter, y+ system), member 5
105	1.2602	0.0332	0.0055	0.0022	208696_at	CCT5	chaperonin containing TCP1, subunit 5 (epsilon)
106	1.2553	0.0358	0.0059	0.0023	218556_at	ORMDL2	ORM1-like 2 (S. cerevisiae)
107	1.2476	0.0400	0.0069	0.0024	211058_x_at	K-ALPHA-1	tubulin, alpha, ubiquitous
108	1.2468	0.0407	0.0072	0.0024	212723_at	PTDSR	phosphatidylserine receptor
109	1.2447	0.0414	0.0075	0.0024	203022_at	RNASEH2A	ribonuclease H2, large subunit
110	1.2402	0.0450	0.0084	0.0029	210334_x_at	BIRC5	baculoviral IAP repeat-containing 5 (survivin)
111	1.2383	0.0461	0.0090	0.0031	203744_at	HMGB3	high-mobility group box 3
112	1.2360	0.0478	0.0095	0.0034	219306_at	KNSL7	kinesin-like 7
113	1.2332	0.0509	0.0102	0.0038	220865_s_at	TPRT	trans-prenyltransferase
114	1.2331	0.0510	0.0103	0.0038	204641_at	NEK2	NIMA (never in mitosis gene a)-related kinase 2

Table 3: ... continued

No.	D	p-values FD>0	FD>1	FD>2	probeset	gene symbol	description
115	1.2319	0.0518	0.0104	0.0038	203358_s_at	EZH2	enhancer of zeste homolog 2 (Drosophila)
116	1.2249	0.0581	0.0119	0.0044	213088_s_at	DNAJC9	DnaJ (Hsp40) homolog, subfamily C, member 9
117	1.2233	0.0593	0.0123	0.0047	214516_at	HIST1H4B	histone 1, H4b
118	1.2226	0.0602	0.0124	0.0049	202110_at	COX7B	cytochrome c oxidase subunit VIIb
119	1.2214	0.0610	0.0128	0.0050	218982_s_at	MRPS17	mitochondrial ribosomal protein S17
120	1.2207	0.0615	0.0129	0.0051	205339_at	SIL	TAL1 (SCL) interrupting locus
121	1.2206	0.0615	0.0129	0.0051	201342_at	SNRPC	small nuclear ribonucleoprotein polypeptide C
122	1.2199	0.0618	0.0131	0.0051	201678_s_at	DC12	DC12 protein
123	1.2192	0.0622	0.0132	0.0052	218875_s_at	FBXO5	F-box protein 5
124	1.2160	0.0650	0.0139	0.0055	218663_at	HCAP-G	chromosome condensation protein G
125	1.2155	0.0652	0.0141	0.0055	212020_s_at	MKI67	antigen identified by monoclonal antibody Ki-67
126	1.2065	0.0738	0.0163	0.0065	217755_at	HN1	hematological and neurological expressed 1
127	1.2028	0.0777	0.0176	0.0067	202635_s_at	POLR2K	polymerase (RNA) II (DNA directed) polypeptide K, 7.0kDa
128	1.2003	0.0802	0.0183	0.0069	202397_at	NUTF2	nuclear transport factor 2
129	1.1993	0.0811	0.0183	0.0071	201930_at	MCM6	MCM6 minichromosome maintenance deficient 6 (MIS5 homolog, S. pombe) (S. cerevisiae)
130	1.1960	0.0849	0.0194	0.0078	222037_at	MCM4	MCM4 minichromosome maintenance deficient 4 (S. cerevisiae)
131	1.1914	0.0900	0.0210	0.0083	205024_s_at	RAD51	RAD51 homolog (RecA homolog, E. coli) (S. cerevisiae)
132	1.1884	0.0932	0.0217	0.0090	211750_x_at	TUBA6	tubulin alpha 6
133	1.1880	0.0938	0.0221	0.0092	203856_at	VRK1	vaccinia related kinase 1
134	1.1861	0.0961	0.0229	0.0093	204267_x_at	PKMYT1	membrane-associated tyrosine- and threonine-specific cdc2-inhibitory kinase
135	1.1807	0.1036	0.0256	0.0107	219787_s_at	ECT2	epithelial cell transforming sequence 2 oncogene
136	1.1800	0.1045	0.0258	0.0109	219494_at	RAD54B	RAD54 homolog B (S. cerevisiae)
137	1.1790	0.1050	0.0262	0.0110	219990_at	FLJ23311	FLJ23311 protein
138	1.1770	0.1078	0.0268	0.0114	219061_s_at	DXS9879E	DNA segment on chromosome X (unique) 9879 expressed sequence
139	1.1767	0.1083	0.0269	0.0114	203832_at	SNRPF	small nuclear ribonucleoprotein polypeptide F
140	1.1757	0.1094	0.0274	0.0115	213646_x_at	K-ALPHA-1	tubulin, alpha, ubiquitous
141	1.1749	0.1107	0.0277	0.0117	201518_at	TOMM70A	translocase of outer mitochondrial membrane 70 homolog A (yeast)
142	1.1728	0.1131	0.0287	0.0121	202824_s_at	TCEB1	transcription elongation factor B (SIII), polypeptide 1 (15kDa, elongin C)
143	1.1725	0.1134	0.0290	0.0122	222029_x_at	HKE2	HLA class II region expressed gene KE2
144	1.1714	0.1142	0.0296	0.0127	205644_s_at	SNRPG	small nuclear ribonucleoprotein polypeptide G
145	1.1664	0.1222	0.0317	0.0146	204170_s_at	CKS2	CDC28 protein kinase regulatory subunit 2
146	1.1658	0.1228	0.0321	0.0147	205394_at	CHEK1	CHK1 checkpoint homolog (S. pombe)
147	1.1630	0.1270	0.0340	0.0155	204023_at	RFC4	replication factor C (activator 1) 4, 37kDa
148	1.1619	0.1289	0.0345	0.0155	218151_x_at	GPR172A	G protein-coupled receptor 172A
149	1.1616	0.1290	0.0345	0.0155	202352_s_at	PSMD12	proteasome (prosome, macropain) 26S subunit, non-ATPase, 12
150	1.1597	0.1322	0.0362	0.0158	202188_at	NUP93	nucleoporin 93kDa
151	1.1548	0.1420	0.0390	0.0175	201291_s_at	TOP2A	topoisomerase (DNA) II alpha 170kDa
152	1.1528	0.1459	0.0404	0.0179	219978_s_at	NUSAP1	nucleolar and spindle associated protein 1
153	1.1525	0.1462	0.0405	0.0182	201266_at	TXNRD1	thioredoxin reductase 1
154	1.1514	0.1487	0.0415	0.0186	204126_s_at	CDC45L	CDC45 cell division cycle 45-like (S. cerevisiae)
155	1.1508	0.1497	0.0418	0.0189	209709_s_at	HMMR	hyaluronan-mediated motility receptor (RHAMM)
156	1.1501	0.1513	0.0421	0.0189	219512_at	C20orf172	chromosome 20 open reading frame 172
157	1.1466	0.1583	0.0446	0.0204	218408_at	TIMM10	translocase of inner mitochondrial membrane 10 homolog (yeast)
158	1.1444	0.1616	0.0457	0.0215	201555_at	MCM3	MCM3 minichromosome maintenance deficient 3 (S. cerevisiae)
159	1.1413	0.1670	0.0479	0.0223	218238_s_at	GTPBP4	GTP binding protein 4
160	1.1412	0.1674	0.0479	0.0223	200783_s_at	STMN1	stathmin 1/oncoprotein 18
161	1.1389	0.1729	0.0498	0.0228	214095_at	SHMT2	serine hydroxymethyltransferase 2 (mitochondrial)
162	1.1385	0.1735	0.0503	0.0231	200853_at	H2AFZ	H2A histone family, member Z
163	1.1346	0.1818	0.0545	0.0248	203931_s_at	MRPL12	mitochondrial ribosomal protein L12
164	1.1332	0.1840	0.0554	0.0254	209744_x_at	ITCH	itchy homolog E3 ubiquitin protein ligase (mouse)
165	1.1320	0.1846	0.0560	0.0256	212639_x_at	TUBA3	tubulin, alpha 3
166	1.1316	0.1873	0.0576	0.0259	204044_at	QPRT	quinolinate phosphoribosyltransferase (nicotinate-nucleotide pyrophosphorylase (carboxylating))

Table 3: ... *continued*

No.	D	p-values FD>0	FD>1	FD>2	probeset	gene symbol	description
167	1.1254	0.2017	0.0638	0.0296	208864_s_at	TXN	thioredoxin
168	1.1233	0.2063	0.0661	0.0304	201114_x_at	PSMA7	proteasome (prosome, macropain) subunit, alpha type, 7
169	1.1228	0.2073	0.0666	0.0311	209172_s_at	CENPF	centromere protein F, 350/400ka (mitosin)
170	1.1224	0.2080	0.0672	0.0314	201577_at	NME1	non-metastatic cells 1, protein (NM23A) expressed in
171	1.1204	0.2120	0.0699	0.0324	213330_s_at	STIP1	stress-induced-phosphoprotein 1 (Hsp70/Hsp90-organizing protein)
172	1.1197	0.2142	0.0701	0.0331	218238_at	GTPBP4	GTP binding protein 4
173	1.1192	0.2155	0.0708	0.0335	214437_s_at	SHMT2	serine hydroxymethyltransferase 2 (mitochondrial)
174	1.1181	0.2180	0.0726	0.0343	218027_at	MRPL15	mitochondrial ribosomal protein L15
175	1.1178	0.2193	0.0728	0.0346	203612_at	BYSL	bystin-like
176	1.1173	0.2209	0.0733	0.0347	202487_s_at	H2AFV	H2A histone family, member V
177	1.1099	0.2410	0.0815	0.0399	218308_at	TACC3	transforming, acidic coiled-coil containing protein 3
178	1.1089	0.2449	0.0823	0.0408	208511_at	PTTG3	pituitary tumor-transforming 3
179	1.1070	0.2509	0.0849	0.0421	212160_at	XPOT	exportin, tRNA (nuclear export receptor for tRNAs)
180	1.1061	0.2541	0.0863	0.0429	2028_s_at	E2F1	E2F transcription factor 1
181	1.1037	0.2608	0.0900	0.0449	203746_s_at	HCCS	holocytochrome c synthase (cytochrome c heme-lyase)
182	1.1018	0.2655	0.0934	0.0464	219004_s_at	C21orf45	chromosome 21 open reading frame 45
183	1.1010	0.2681	0.0940	0.0473	206632_s_at	APOBEC3B	apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3B
184	1.1006	0.2691	0.0946	0.0478	219588_s_at	MTB	more than blood homolog
185	1.1000	0.2706	0.0953	0.0483	205393_s_at	CHEK1	CHK1 checkpoint homolog (S. pombe)

Table 3: Up-regulated in grade 1 tumors

No.	D	p-values FD>0	FD>1	FD>2	probeset	gene symbol	description
1	-1.4739	0.0014	0.0001	0.0001	213103_at	STARD13	START domain containing 13
2	-1.4647	0.0014	0.0001	0.0001	204703_at	TTC10	tetratricopeptide repeat domain 10
3	-1.4196	0.0024	0.0002	0.0001	218346_s_at	SESN1	sestrin 1
4	-1.4084	0.0031	0.0003	0.0001	218471_s_at	BBS1	Bardet-Biedl syndrome 1
5	-1.3840	0.0048	0.0004	0.0001	205898_at	CX3CR1	chemokine (C-X3-C motif) receptor 1
6	-1.3482	0.0084	0.0007	0.0003	204072_s_at	13CDNA73	hypothetical protein CG003
7	-1.3235	0.0131	0.0017	0.0004	219455_at	FLJ21062	hypothetical protein FLJ21062
8	-1.2840	0.0249	0.0034	0.0012	217889_s_at	CYBRD1	cytochrome b reductase 1
9	-1.2835	0.0252	0.0034	0.0012	219238_at	FLJ20477	hypothetical protein FLJ20477
10	-1.2663	0.0312	0.0045	0.0018	216264_s_at	LAMB2	laminin, beta 2 (laminin S)
11	-1.2656	0.0314	0.0046	0.0019	221562_s_at	SIRT3	sirtuin (silent mating type information regulation 2 homolog) 3 (S. cerevisiae)
12	-1.2628	0.0322	0.0049	0.0022	216520_s_at	TPT1	tumor protein, translationally-controlled 1
13	-1.2568	0.0351	0.0056	0.0023	220141_at	FLJ23554	hypothetical protein FLJ23554
14	-1.2557	0.0356	0.0059	0.0023	218483_s_at	FLJ21827	hypothetical protein FLJ21827
15	-1.2548	0.0364	0.0060	0.0023	221771_s_at	HSMPP8	M-phase phosphoprotein, mpp8
16	-1.2399	0.0450	0.0084	0.0029	220917_s_at	WDR19	WD repeat domain 19
17	-1.2259	0.0574	0.0117	0.0044	212695_at	CRY2	cryptochrome 2 (photolyase-like)
18	-1.2233	0.0593	0.0124	0.0047	213340_s_at	KIAA0495	KIAA0495
19	-1.2156	0.0652	0.0140	0.0055	213444_at	KIAA0543	KIAA0543 protein
20	-1.2144	0.0662	0.0141	0.0057	220173_at	C14orf45	chromosome 14 open reading frame 45
21	-1.2139	0.0666	0.0144	0.0059	201384_s_at	M17S2	membrane component, chromosome 17, surface marker 2 (ovarian carcinoma antigen CA125)
22	-1.2098	0.0711	0.0152	0.0062	203156_at	AKAP11	A kinase (PRKA) anchor protein 11
23	-1.2064	0.0740	0.0163	0.0065	209407_s_at	DEAF1	deformed epidermal autoregulatory factor 1 (Drosophila)
24	-1.2017	0.0789	0.0179	0.0067	219469_at	DNCH2	dynein, cytoplasmic, heavy polypeptide 2
25	-1.2003	0.0802	0.0183	0.0069	203984_s_at	CASP9	caspase 9, apoptosis-related cysteine protease
26	-1.1973	0.0837	0.0190	0.0071	217844_at	CTDSP1	CTD (carboxy-terminal domain, RNA polymerase II, polypeptide A) small phosphatase 1
27	-1.1906	0.0914	0.0212	0.0084	213397_x_at	RNASE4	ribonuclease, RNase A family, 4
28	-1.1896	0.0919	0.0214	0.0088	206197_at	NME5	non-metastatic cells 5, protein expressed in (nucleoside-diphosphate kinase)
29	-1.1878	0.0941	0.0221	0.0093	219922_s_at	LTBP3	latent transforming growth factor beta binding protein 3
30	-1.1829	0.1003	0.0247	0.0102	201383_s_at	M17S2	membrane component, chromosome 17, surface marker 2 (ovarian carcinoma antigen CA125)
31	-1.1827	0.1007	0.0249	0.0102	206081_at	SLC24A1	solute carrier family 24 (sodium/potassium/calcium exchanger), member 1
32	-1.1709	0.1153	0.0296	0.0129	213266_at	76P	Gamma tubulin ring complex protein (76p gene)
33	-1.1707	0.1156	0.0297	0.0129	209189_at	FOS	v-fos FBJ murine osteosarcoma viral oncogene homolog
34	-1.1679	0.1201	0.0307	0.0142	214829_at	AASS	aminoadipate-semialdehyde synthase
35	-1.1633	0.1268	0.0337	0.0153	221123_x_at	ZNF395	zinc finger protein 395
36	-1.1625	0.1279	0.0344	0.0155	200810_s_at	CIRBP	cold inducible RNA binding protein
37	-1.1612	0.1298	0.0354	0.0157	210365_at	RUNX1	runt-related transcription factor 1 (acute myeloid leukemia 1; aml1 oncogene)
38	-1.1602	0.1314	0.0358	0.0158	212842_x_at	RANBP2L1	RAN binding protein 2-like 1
39	-1.1595	0.1324	0.0362	0.0158	213364_s_at	SNX1	Sorting nexin 1
40	-1.1586	0.1339	0.0369	0.0161	220911_s_at	KIAA1305	KIAA1305
41	-1.1539	0.1444	0.0398	0.0175	201335_s_at	ARHGEF12	Rho guanine nucleotide exchange factor (GEF) 12
42	-1.1499	0.1519	0.0421	0.0190	221276_s_at	SYNCOILIN	intermediate filament protein syncoilin
43	-1.1453	0.1600	0.0451	0.0209	221824_s_at	MIR	c-mir, cellular modulator of immune recognition
44	-1.1437	0.1625	0.0461	0.0217	211943_x_at	TPT1	tumor protein, translationally-controlled 1
45	-1.1408	0.1681	0.0481	0.0223	218552_at	FLJ10946	hypothetical protein FLJ10946
46	-1.1382	0.1743	0.0510	0.0232	220326_s_at	FLJ10357	hypothetical protein FLJ10357
47	-1.1254	0.2014	0.0638	0.0296	212869_x_at	TPT1	tumor protein, translationally-controlled 1
48	-1.1184	0.2172	0.0722	0.0342	218648_at	TORC3	transducer of regulated cAMP response element-binding protein (CREB) 3
49	-1.1171	0.2215	0.0736	0.0348	212549_at	STAT5B	signal transducer and activator of transcription 5B
50	-1.1165	0.2233	0.0743	0.0356	219951_s_at	C20orf12	chromosome 20 open reading frame 12
51	-1.1147	0.2277	0.0763	0.0364	212678_at	NF1	Neurofibromin 1 (neurofibromatosis, von Recklinghausen disease, Watson disease)
52	-1.1113	0.2369	0.0807	0.0386	210852_s_at	AASS	aminoadipate-semialdehyde synthase

Table 3: ... *continued*

		p-values					
No.	*D*	FD>0	FD>1	FD>2	probeset	gene symbol	description
53	-1.1112	0.2370	0.0807	0.0387	202962_at	KIF13B	kinesin family member 13B
54	-1.1102	0.2404	0.0813	0.0394	214724_at	DIXDC1	DIX domain containing 1
55	-1.1089	0.2450	0.0823	0.0408	206542_s_at	SMARCA2	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 2
56	-1.1085	0.2464	0.0827	0.0411	207757_at	FLJ21628	hypothetical protein FLJ21628
57	-1.1085	0.2464	0.0827	0.0411	218466_at	TBC1D17	TBC1 domain family, member 17

Table 3: Up-regulated in grade 3 tumors

No.	D	p-values FD>0	FD>1	FD>2	probeset	gene symbol	description
1	2.1162	0.0001	0.0001	0.0001	202954_at	UBE2C	ubiquitin-conjugating enzyme E2C
2	1.9037	0.0001	0.0001	0.0001	222077_s_at	RACGAP1	Rac GTPase activating protein 1
3	1.7292	0.0001	0.0001	0.0001	201088_at	KPNA2	karyopherin alpha 2 (RAG cohort 1, importin alpha 1)
4	1.7264	0.0001	0.0001	0.0001	218542_at	C10orf3	chromosome 10 open reading frame 3
5	1.7259	0.0001	0.0001	0.0001	203554_x_at	PTTG1	pituitary tumor-transforming 1
6	1.7053	0.0001	0.0001	0.0001	218355_at	KIF4A	kinesin family member 4A
7	1.6600	0.0001	0.0001	0.0001	210052_s_at	TPX2	TPX2, microtubule-associated protein homolog (Xenopus laevis)
8	1.6598	0.0001	0.0001	0.0001	202580_x_at	FOXM1	forkhead box M1
9	1.6548	0.0001	0.0001	0.0001	208079_s_at	STK6	serine/threonine kinase 6
10	1.6513	0.0001	0.0001	0.0001	204092_s_at	STK6	serine/threonine kinase 6
11	1.6495	0.0001	0.0001	0.0001	218755_at	KIF20A	kinesin family member 20A
12	1.6387	0.0001	0.0001	0.0001	201584_s_at	DDX39	DEAD (Asp-Glu-Ala-Asp) box polypeptide 39
13	1.6347	0.0001	0.0001	0.0001	203764_at	DLG7	discs, large homolog 7 (Drosophila)
14	1.6223	0.0001	0.0001	0.0001	204825_at	MELK	maternal embryonic leucine zipper kinase
15	1.6213	0.0001	0.0001	0.0001	203418_at	CCNA2	cyclin A2
16	1.6095	0.0001	0.0001	0.0001	204766_s_at	NUDT1	nudix (nucleoside diphosphate linked moiety X)-type motif 1
17	1.6057	0.0001	0.0001	0.0001	206102_at	KIAA0186	KIAA0186 gene product
18	1.5986	0.0001	0.0001	0.0001	202095_s_at	BIRC5	baculoviral IAP repeat-containing 5 (survivin)
19	1.5957	0.0001	0.0001	0.0001	201710_at	MYBL2	v-myb myeloblastosis viral oncogene homolog (avian)-like 2
20	1.5879	0.0002	0.0001	0.0001	211762_s_at	KPNA2	karyopherin alpha 2 (RAG cohort 1, importin alpha 1)
21	1.5816	0.0002	0.0001	0.0001	209680_s_at	KIFC1	Kinesin family member C1
22	1.5785	0.0002	0.0001	0.0001	209408_at	KIF2C	kinesin family member 2C
23	1.5671	0.0002	0.0001	0.0001	219918_s_at	ASPM	asp (abnormal spindle)-like, microcephaly associated (Drosophila)
24	1.5650	0.0003	0.0001	0.0001	203145_at	SPAG5	sperm associated antigen 5
25	1.5595	0.0003	0.0001	0.0001	204962_s_at	CENPA	centromere protein A, 17kDa
26	1.5551	0.0003	0.0001	0.0001	202870_s_at	CDC20	CDC20 cell division cycle 20 homolog (S. cerevisiae)
27	1.5446	0.0003	0.0001	0.0001	38158_at	ESPL1	extra spindle poles like 1 (S. cerevisiae)
28	1.5376	0.0003	0.0001	0.0001	202107_s_at	MCM2	MCM2 minichromosome maintenance deficient 2, mitotin (S. cerevisiae)
29	1.5236	0.0004	0.0001	0.0001	204767_s_at	FEN1	flap structure-specific endonuclease 1
30	1.5226	0.0004	0.0001	0.0001	203046_s_at	TIMELESS	timeless homolog (Drosophila)
31	1.5221	0.0004	0.0001	0.0001	221677_s_at	DONSON	downstream neighbor of SON
32	1.5134	0.0005	0.0001	0.0001	210559_s_at	CDC2	cell division cycle 2, G1 to S and G2 to M
33	1.5047	0.0006	0.0001	0.0001	221520_s_at	CDCA8	cell division cycle associated 8
34	1.5017	0.0007	0.0001	0.0001	214710_s_at	CCNB1	cyclin B1
35	1.4945	0.0007	0.0001	0.0001	209714_s_at	CDKN3	cyclin-dependent kinase inhibitor 3 (CDK2-associated dual specificity phosphatase)
36	1.4933	0.0008	0.0001	0.0001	204444_at	KIF11	kinesin family member 11
37	1.4927	0.0008	0.0001	0.0001	210821_x_at	CENPA	centromere protein A, 17kDa
38	1.4915	0.0008	0.0001	0.0001	218726_at	DKFZp762E1312	hypothetical protein DKFZp762E1312
39	1.4895	0.0009	0.0001	0.0001	220651_s_at	MCM10	MCM10 minichromosome maintenance deficient 10 (S. cerevisiae)
40	1.4865	0.0010	0.0001	0.0001	201475_x_at	MARS	methionine-tRNA synthetase
41	1.4715	0.0014	0.0001	0.0001	204033_at	TRIP13	thyroid hormone receptor interactor 13
42	1.4672	0.0014	0.0001	0.0001	202705_at	CCNB2	cyclin B2
43	1.4624	0.0014	0.0001	0.0001	204649_at	TROAP	trophinin associated protein (tastin)
44	1.4603	0.0014	0.0001	0.0001	220060_s_at	FLJ20641	hypothetical protein FLJ20641
45	1.4534	0.0016	0.0001	0.0001	209836_x_at	LAT1-3TM	LAT1-3TM protein
46	1.4533	0.0016	0.0001	0.0001	203276_at	LMNB1	lamin B1
47	1.4471	0.0018	0.0001	0.0001	205034_at	CCNE2	cyclin E2
48	1.4455	0.0018	0.0001	0.0001	203213_at	CDC2	cell division cycle 2, G1 to S and G2 to M
49	1.4384	0.0019	0.0001	0.0001	209464_at	AURKB	aurora kinase B
50	1.4381	0.0019	0.0001	0.0001	205046_at	CENPE	centromere protein E, 312kDa
51	1.4373	0.0019	0.0001	0.0001	203755_at	BUB1B	BUB1 budding uninhibited by benzimidazoles 1 homolog beta (yeast)
52	1.4336	0.0019	0.0001	0.0001	203214_x_at	CDC2	cell division cycle 2, G1 to S and G2 to M
53	1.4236	0.0022	0.0002	0.0001	214804_at	FSHPRH1	FSH primary response (LRPR1 homolog, rat) 1
54	1.4167	0.0026	0.0003	0.0001	212949_at	BRIN1	barren homolog (Drosophila)
55	1.4134	0.0027	0.0003	0.0001	204318_s_at	GTSE1	G-2 and S-phase expressed 1

Table 3: ... continued

No.	D	p-values FDR>0	FDR>1	FDR>2	probeset	gene symbol	description
56	1.4105	0.0030	0.0003	0.0001	207165_at	HMMR	hyaluronan-mediated motility receptor (RHAMM)
57	1.4079	0.0031	0.0003	0.0001	212022_s_at	MKI67	antigen identified by monoclonal antibody Ki-67
58	1.4061	0.0031	0.0003	0.0001	213226_at	CCNA2	cyclin A2
59	1.3931	0.0041	0.0004	0.0001	219510_at	POLQ	polymerase (DNA directed), theta
60	1.3890	0.0044	0.0004	0.0001	204026_s_at	ZWINT	ZW10 interactor
61	1.3890	0.0044	0.0004	0.0001	203432_at	TMPO	thymopoietin
62	1.3872	0.0046	0.0004	0.0001	204768_s_at	FEN1	flap structure-specific endonuclease 1
63	1.3855	0.0047	0.0004	0.0001	209773_s_at	RRM2	ribonucleotide reductase M2 polypeptide
64	1.3847	0.0047	0.0004	0.0001	214431_at	GMPS	guanine monphosphate synthetase
65	1.3842	0.0048	0.0004	0.0001	212023_s_at	MKI67	antigen identified by monoclonal antibody Ki-67
66	1.3752	0.0052	0.0004	0.0002	218883_s_at	MLF1IP	MLF1 interacting protein
67	1.3541	0.0077	0.0006	0.0003	211519_s_at	KIF2C	kinesin family member 2C
68	1.3503	0.0083	0.0006	0.0003	202240_at	PLK1	polo-like kinase 1 (Drosophila)
69	1.3460	0.0089	0.0007	0.0003	205733_at	BLM	Bloom syndrome
70	1.3457	0.0092	0.0008	0.0003	222039_at	LOC146909	hypothetical protein LOC146909
71	1.3443	0.0096	0.0008	0.0003	209642_at	BUB1	BUB1 budding uninhibited by benzimidazoles 1 homolog (yeast)
72	1.3376	0.0102	0.0010	0.0003	213599_at	OIP5	Opa-interacting protein 5
73	1.3372	0.0102	0.0010	0.0003	214096_s_at	SHMT2	serine hydroxymethyltransferase 2 (mitochondrial)
74	1.3348	0.0105	0.0012	0.0003	211072_x_at	K-ALPHA-1	tubulin, alpha, ubiquitous
75	1.3237	0.0130	0.0017	0.0004	202779_s_at	UBE2S	ubiquitin-conjugating enzyme E2S
76	1.3226	0.0133	0.0017	0.0004	218447_at	DC13	DC13 protein
77	1.3215	0.0138	0.0017	0.0004	213911_s_at	H2AFZ	H2A histone family, member Z
78	1.3211	0.0138	0.0017	0.0004	212141_at	MCM4	MCM4 minichromosome maintenance deficient 4 (S. cerevisiae)
79	1.3156	0.0153	0.0019	0.0005	221591_s_at	FLJ10156	hypothetical protein FLJ10156
80	1.3139	0.0162	0.0019	0.0005	204822_at	TTK	TTK protein kinase
81	1.3121	0.0165	0.0020	0.0005	209251_x_at	TUBA6	tubulin alpha 6
82	1.3086	0.0173	0.0023	0.0006	217835_x_at	C20orf24	chromosome 20 open reading frame 24
83	1.3081	0.0176	0.0023	0.0006	201890_at	RRM2	ribonucleotide reductase M2 polypeptide
84	1.3059	0.0184	0.0024	0.0006	213671_s_at	MARS	methionine-tRNA synthetase
85	1.3053	0.0185	0.0024	0.0006	218009_s_at	PRC1	protein regulator of cytokinesis 1
86	1.3010	0.0197	0.0026	0.0007	207828_s_at	CENPF	centromere protein F, 350/400ka (mitosin)
87	1.3002	0.0198	0.0026	0.0007	219555_s_at	BM039	uncharacterized bone marrow protein BM039
88	1.2969	0.0206	0.0026	0.0007	204695_at	CDC25A	cell division cycle 25A
89	1.2953	0.0214	0.0026	0.0009	212021_s_at	MKI67	antigen identified by monoclonal antibody Ki-67
90	1.2898	0.0229	0.0028	0.0009	201090_x_at	K-ALPHA-1	tubulin, alpha, ubiquitous
91	1.2885	0.0233	0.0033	0.0010	218039_at	NUSAP1	nucleolar and spindle associated protein 1
92	1.2851	0.0246	0.0034	0.0012	204603_at	EXO1	exonuclease 1
93	1.2846	0.0248	0.0034	0.0012	203362_s_at	MAD2L1	MAD2 mitotic arrest deficient-like 1 (yeast)
94	1.2845	0.0248	0.0034	0.0012	202094_at	BIRC5	baculoviral IAP repeat-containing 5 (survivin)
95	1.2840	0.0249	0.0034	0.0012	204162_at	KNTC2	kinetochore associated 2
96	1.2825	0.0254	0.0037	0.0012	222036_s_at	MCM4	MCM4 minichromosome maintenance deficient 4 (S. cerevisiae)
97	1.2780	0.0272	0.0039	0.0014	204252_at	CDK2	cyclin-dependent kinase 2
98	1.2775	0.0274	0.0039	0.0014	219000_s_at	DCC1	defective in sister chromatid cohesion homolog 1 (S. cerevisiae)
99	1.2772	0.0277	0.0041	0.0014	201524_x_at	UBE2N	ubiquitin-conjugating enzyme E2N (UBC13 homolog, yeast)
100	1.2694	0.0294	0.0044	0.0018	204817_at	ESPL1	extra spindle poles like 1 (S. cerevisiae)
101	1.2657	0.0313	0.0046	0.0019	218662_s_at	HCAP-G	chromosome condensation protein G
102	1.2620	0.0324	0.0052	0.0022	206364_at	KIF14	Kinesin family member 14
103	1.2612	0.0329	0.0054	0.0022	221436_s_at	CDCA3	cell division cycle associated 3
104	1.2609	0.0331	0.0054	0.0022	201195_s_at	SLC7A5	solute carrier family 7 (cationic amino acid transporter, y+ system), member 5
105	1.2602	0.0332	0.0055	0.0022	208696_at	CCT5	chaperonin containing TCP1, subunit 5 (epsilon)
106	1.2553	0.0358	0.0059	0.0023	218556_at	ORMDL2	ORM1-like 2 (S. cerevisiae)
107	1.2476	0.0400	0.0069	0.0024	211058_x_at	K-ALPHA-1	tubulin, alpha, ubiquitous
108	1.2468	0.0407	0.0072	0.0024	212723_at	PTDSR	phosphatidylserine receptor
109	1.2447	0.0414	0.0076	0.0024	203022_at	RNASEH2A	ribonuclease H2, large subunit
110	1.2402	0.0450	0.0084	0.0029	210334_x_at	BIRC5	baculoviral IAP repeat-containing 5 (survivin)
111	1.2383	0.0461	0.0090	0.0031	203744_at	HMGB3	high-mobility group box 3
112	1.2360	0.0478	0.0096	0.0034	219306_at	KNSL7	kinesin-like 7
113	1.2332	0.0509	0.0102	0.0038	220865_s_at	TPRT	trans-prenyltransferase
114	1.2331	0.0510	0.0103	0.0038	204641_at	NEK2	NIMA (never in mitosis gene a)-related kinase 2

Table 3: ... continued

No.	D	p-values FDR>0	FDR>1	FDR>2	probeset	gene symbol	description
115	1.2319	0.0518	0.0104	0.0038	203358_s_at	EZH2	enhancer of zeste homolog 2 (Drosophila)
116	1.2249	0.0581	0.0119	0.0044	213088_s_at	DNAJC9	DnaJ (Hsp40) homolog, subfamily C, member 9
117	1.2233	0.0593	0.0123	0.0047	214516_at	HIST1H4B	histone 1, H4b
118	1.2225	0.0602	0.0124	0.0049	202110_at	COX7B	cytochrome c oxidase subunit VIIb
119	1.2214	0.0610	0.0128	0.0050	218982_s_at	MRPS17	mitochondrial ribosomal protein S17
120	1.2207	0.0615	0.0129	0.0051	205339_at	SIL	TAL1 (SCL) interrupting locus
121	1.2206	0.0615	0.0129	0.0051	201342_at	SNRPC	small nuclear ribonucleoprotein polypeptide C
122	1.2199	0.0619	0.0131	0.0051	201678_s_at	DC12	DC12 protein
123	1.2192	0.0622	0.0132	0.0052	218875_s_at	FBXO5	F-box protein 5
124	1.2160	0.0650	0.0139	0.0055	218663_at	HCAP-G	chromosome condensation protein G
125	1.2155	0.0652	0.0141	0.0055	212020_s_at	MKI67	antigen identified by monoclonal antibody Ki-67
126	1.2085	0.0738	0.0163	0.0065	217755_at	HN1	hematological and neurological expressed 1
127	1.2028	0.0777	0.0176	0.0067	202635_s_at	POLR2K	polymerase (RNA) II (DNA directed) polypeptide K, 7.0kDa
128	1.2003	0.0802	0.0183	0.0069	202397_at	NUTF2	nuclear transport factor 2
129	1.1993	0.0811	0.0183	0.0071	201930_at	MCM6	MCM6 minichromosome maintenance deficient 6 (MIS5 homolog, S. pombe) (S. cerevisiae)
130	1.1960	0.0848	0.0194	0.0073	222037_at	MCM4	MCM4 minichromosome maintenance deficient 4 (S. cerevisiae)
131	1.1914	0.0900	0.0210	0.0083	205024_s_at	RAD51	RAD51 homolog (RecA homolog, E. coli) (S. cerevisiae)
132	1.1884	0.0932	0.0217	0.0090	211750_x_at	TUBA6	tubulin alpha 6
133	1.1880	0.0938	0.0221	0.0092	203856_at	VRK1	vaccinia related kinase 1
134	1.1861	0.0961	0.0229	0.0093	204267_x_at	PKMYT1	membrane-associated tyrosine- and threonine-specific cdc2-inhibitory kinase
135	1.1807	0.1036	0.0256	0.0107	219787_s_at	ECT2	epithelial cell transforming sequence 2 oncogene
136	1.1800	0.1045	0.0258	0.0109	219494_at	RAD54B	RAD54 homolog B (S. cerevisiae)
137	1.1790	0.1050	0.0262	0.0110	219990_at	FLJ23311	FLJ23311 protein
138	1.1770	0.1078	0.0268	0.0114	219061_s_at	DXS9879E	DNA segment on chromosome X (unique) 9879 expressed sequence
139	1.1767	0.1083	0.0269	0.0114	203832_at	SNRPF	small nuclear ribonucleoprotein polypeptide F
140	1.1757	0.1094	0.0274	0.0115	213646_x_at	K-ALPHA-1	tubulin, alpha, ubiquitous
141	1.1749	0.1107	0.0277	0.0117	201519_at	TOMM70A	translocase of outer mitochondrial membrane 70 homolog A (yeast)
142	1.1728	0.1131	0.0287	0.0121	202824_s_at	TCEB1	transcription elongation factor B (SIII), polypeptide 1 (15kDa, elongin C)
143	1.1725	0.1134	0.0290	0.0122	222029_x_at	HKE2	HLA class II region expressed gene KE2
144	1.1714	0.1142	0.0296	0.0127	205644_s_at	SNRPG	small nuclear ribonucleoprotein polypeptide G
145	1.1664	0.1222	0.0317	0.0146	204170_s_at	CKS2	CDC28 protein kinase regulatory subunit 2
146	1.1658	0.1228	0.0321	0.0147	205394_at	CHEK1	CHK1 checkpoint homolog (S. pombe)
147	1.1630	0.1270	0.0340	0.0155	204023_at	RFC4	replication factor C (activator 1) 4, 37kDa
148	1.1619	0.1289	0.0345	0.0155	218151_x_at	GPR172A	G protein-coupled receptor 172A
149	1.1616	0.1290	0.0345	0.0155	202352_s_at	PSMD12	proteasome (prosome, macropain) 26S subunit, non-ATPase, 12
150	1.1597	0.1322	0.0362	0.0158	202188_at	NUP93	nucleoporin 93kDa
151	1.1548	0.1420	0.0390	0.0175	201291_s_at	TOP2A	topoisomerase (DNA) II alpha 170kDa
152	1.1528	0.1459	0.0404	0.0179	219978_s_at	NUSAP1	nucleolar and spindle associated protein 1
153	1.1525	0.1462	0.0405	0.0182	201266_at	TXNRD1	thioredoxin reductase 1
154	1.1514	0.1487	0.0415	0.0186	204126_s_at	CDC45L	CDC45 cell division cycle 45-like (S. cerevisiae)
155	1.1508	0.1497	0.0418	0.0189	209709_s_at	HMMR	hyaluronan-mediated motility receptor (RHAMM)
156	1.1501	0.1513	0.0421	0.0189	219512_at	C20orf172	chromosome 20 open reading frame 172
157	1.1466	0.1583	0.0446	0.0204	218408_at	TIMM10	translocase of inner mitochondrial membrane 10 homolog (yeast)
158	1.1444	0.1616	0.0457	0.0215	201555_at	MCM3	MCM3 minichromosome maintenance deficient 3 (S. cerevisiae)
159	1.1413	0.1670	0.0479	0.0223	218239_s_at	GTPBP4	GTP binding protein 4
160	1.1412	0.1674	0.0479	0.0223	200783_s_at	STMN1	stathmin 1/oncoprotein 18
161	1.1389	0.1720	0.0498	0.0228	214095_at	SHMT2	serine hydroxymethyltransferase 2 (mitochondrial)
162	1.1365	0.1735	0.0503	0.0231	200853_at	H2AFZ	H2A histone family, member Z
163	1.1346	0.1818	0.0545	0.0248	203931_s_at	MRPL12	mitochondrial ribosomal protein L12
164	1.1332	0.1840	0.0554	0.0254	209744_x_at	ITCH	itchy homolog E3 ubiquitin protein ligase (mouse)
165	1.1329	0.1846	0.0560	0.0256	212639_x_at	TUBA3	tubulin, alpha 3
166	1.1316	0.1873	0.0576	0.0259	204044_at	QPRT	quinolinate phosphoribosyltransferase (nicotinate-nucleotide pyrophosphorylase (carboxylating))

Table 3: ... continued

No.	D	p-values FD>0	FD>1	FD>2	probeset	gene symbol	description
167	1.1254	0.2017	0.0638	0.0296	208864_s_at	TXN	thioredoxin
168	1.1233	0.2063	0.0661	0.0304	201114_x_at	PSMA7	proteasome (prosome, macropain) subunit, alpha type, 7
169	1.1228	0.2073	0.0666	0.0311	209172_s_at	CENPF	centromere protein F, 350/400ka (mitosin)
170	1.1224	0.2080	0.0672	0.0314	201577_at	NME1	non-metastatic cells 1, protein (NM23A) expressed in
171	1.1204	0.2129	0.0699	0.0324	213330_s_at	STIP1	stress-induced-phosphoprotein 1 (Hsp70/Hsp90-organizing protein)
172	1.1197	0.2142	0.0701	0.0331	218238_at	GTPBP4	GTP binding protein 4
173	1.1192	0.2156	0.0708	0.0335	214437_s_at	SHMT2	serine hydroxymethyltransferase 2 (mitochondrial)
174	1.1181	0.2180	0.0726	0.0343	218027_at	MRPL15	mitochondrial ribosomal protein L15
175	1.1178	0.2193	0.0728	0.0346	203612_at	BYSL	bystin-like
176	1.1173	0.2209	0.0733	0.0347	202487_s_at	H2AFV	H2A histone family, member V
177	1.1099	0.2418	0.0815	0.0399	218308_at	TACC3	transforming, acidic coiled-coil containing protein 3
178	1.1089	0.2449	0.0823	0.0408	208511_at	PTTG3	pituitary tumor-transforming 3
179	1.1070	0.2509	0.0849	0.0421	212160_at	XPOT	exportin, tRNA (nuclear export receptor for tRNAs)
180	1.1061	0.2541	0.0863	0.0429	2028_s_at	E2F1	E2F transcription factor 1
181	1.1037	0.2608	0.0900	0.0449	203746_s_at	HCCS	holocytochrome c synthase (cytochrome c heme-lyase)
182	1.1018	0.2655	0.0934	0.0464	219004_s_at	C21orf45	chromosome 21 open reading frame 45
183	1.1010	0.2681	0.0940	0.0473	206632_s_at	APOBEC3B	apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3B
184	1.1006	0.2691	0.0946	0.0478	219588_s_at	MTB	more than blood homolog
185	1.1000	0.2706	0.0953	0.0483	205393_s_at	CHEK1	CHK1 checkpoint homolog (S. pombe)

Table 3: Up-regulated in grade 1 tumors

No.	D	p-value FD>0	p-value FD>1	p-value FD>2	probeset	gene symbol	description
1	-1.4739	0.0014	0.0001	0.0001	213103_at	STARD13	START domain containing 13
2	-1.4647	0.0014	0.0001	0.0001	204703_at	TTC10	tetratricopeptide repeat domain 10
3	-1.4196	0.0024	0.0002	0.0001	218346_s_at	SESN1	sestrin 1
4	-1.4084	0.0031	0.0003	0.0001	218471_s_at	BBS1	Bardet-Biedl syndrome 1
5	-1.3840	0.0048	0.0004	0.0001	205898_at	CX3CR1	chemokine (C-X3-C motif) receptor 1
6	-1.3482	0.0084	0.0007	0.0003	204072_s_at	13CDNA73	hypothetical protein CG003
7	-1.3235	0.0131	0.0017	0.0004	219455_at	FLJ21062	hypothetical protein FLJ21062
8	-1.2840	0.0249	0.0034	0.0012	217889_s_at	CYBRD1	cytochrome b reductase 1
9	-1.2835	0.0252	0.0034	0.0012	219238_at	FLJ20477	hypothetical protein FLJ20477
10	-1.2663	0.0312	0.0045	0.0018	216264_s_at	LAMB2	laminin, beta 2 (laminin S)
11	-1.2656	0.0314	0.0046	0.0019	221562_s_at	SIRT3	sirtuin (silent mating type information regulation 2 homolog) 3 (S. cerevisiae)
12	-1.2628	0.0322	0.0049	0.0022	216520_s_at	TPT1	tumor protein, translationally-controlled 1
13	-1.2568	0.0351	0.0055	0.0023	220141_at	FLJ23554	hypothetical protein FLJ23554
14	-1.2557	0.0356	0.0059	0.0023	218483_s_at	FLJ21827	hypothetical protein FLJ21827
15	-1.2548	0.0364	0.0060	0.0023	221771_s_at	HSMPP8	M-phase phosphoprotein, mpp8
16	-1.2399	0.0450	0.0084	0.0029	220917_s_at	WDR19	WD repeat domain 19
17	-1.2259	0.0574	0.0117	0.0044	212695_at	CRY2	cryptochrome 2 (photolyase-like)
18	-1.2233	0.0593	0.0124	0.0047	213340_s_at	KIAA0495	KIAA0495
19	-1.2156	0.0652	0.0149	0.0055	213444_at	KIAA0543	KIAA0543 protein
20	-1.2144	0.0662	0.0141	0.0057	220173_at	C14orf45	chromosome 14 open reading frame 45
21	-1.2139	0.0666	0.0144	0.0059	201384_s_at	M17S2	membrane component, chromosome 17, surface marker 2 (ovarian carcinoma antigen CA125)
22	-1.2098	0.0711	0.0152	0.0062	203166_at	AKAP11	A kinase (PRKA) anchor protein 11
23	-1.2064	0.0740	0.0163	0.0065	209407_s_at	DEAF1	deformed epidermal autoregulatory factor 1 (Drosophila)
24	-1.2017	0.0789	0.0179	0.0067	219469_at	DNCH2	dynein, cytoplasmic, heavy polypeptide 2
25	-1.2003	0.0802	0.0183	0.0069	203984_s_at	CASP8	caspase 8, apoptosis-related cysteine protease
26	-1.1973	0.0837	0.0190	0.0071	217844_at	CTDSP1	CTD (carboxy-terminal domain, RNA polymerase II, polypeptide A) small phosphatase 1
27	-1.1906	0.0914	0.0212	0.0084	213397_x_at	RNASE4	ribonuclease, RNase A family, 4
28	-1.1896	0.0919	0.0214	0.0088	206197_at	NME5	non-metastatic cells 5, protein expressed in (nucleoside-diphosphate kinase)
29	-1.1878	0.0941	0.0221	0.0093	219922_s_at	LTBP3	latent transforming growth factor beta binding protein 3
30	-1.1829	0.1003	0.0247	0.0102	201383_s_at	M17S2	membrane component, chromosome 17, surface marker 2 (ovarian carcinoma antigen CA125)
31	-1.1827	0.1007	0.0249	0.0102	206081_at	SLC24A1	solute carrier family 24 (sodium/potassium/calcium exchanger), member 1
32	-1.1709	0.1153	0.0296	0.0129	213266_at	76P	Gamma tubulin ring complex protein (76p gene)
33	-1.1707	0.1156	0.0297	0.0129	209189_at	FOS	v-fos FBJ murine osteosarcoma viral oncogene homolog
34	-1.1679	0.1201	0.0307	0.0142	214829_at	AASS	aminoadipate-semialdehyde synthase
35	-1.1633	0.1268	0.0337	0.0153	221123_x_at	ZNF395	zinc finger protein 395
36	-1.1625	0.1279	0.0344	0.0155	200810_s_at	CIRBP	cold inducible RNA binding protein
37	-1.1612	0.1298	0.0354	0.0157	210365_at	RUNX1	runt-related transcription factor 1 (acute myeloid leukemia 1; aml1 oncogene)
38	-1.1602	0.1314	0.0358	0.0158	212842_x_at	RANBP2L1	RAN binding protein 2-like 1
39	-1.1596	0.1324	0.0362	0.0158	213364_s_at	SNX1	Sorting nexin 1
40	-1.1586	0.1339	0.0369	0.0161	220911_s_at	KIAA1305	KIAA1305
41	-1.1539	0.1444	0.0398	0.0175	201335_s_at	ARHGEF12	Rho guanine nucleotide exchange factor (GEF) 12
42	-1.1499	0.1519	0.0421	0.0190	221276_s_at	SYNCOILIN	intermediate filament protein syncollin
43	-1.1453	0.1609	0.0451	0.0209	221824_s_at	MIR	c-mir, cellular modulator of immune recognition
44	-1.1437	0.1626	0.0461	0.0217	211943_x_at	TPT1	tumor protein, translationally-controlled 1
45	-1.1408	0.1661	0.0481	0.0223	218552_at	FLJ10948	hypothetical protein FLJ10948
46	-1.1382	0.1743	0.0510	0.0232	220326_s_at	FLJ10357	hypothetical protein FLJ10357
47	-1.1254	0.2014	0.0638	0.0296	212869_x_at	TPT1	tumor protein, translationally-controlled 1
48	-1.1184	0.2172	0.0722	0.0342	218648_at	TORC3	transducer of regulated cAMP response element-binding protein (CREB) 3
49	-1.1171	0.2215	0.0735	0.0348	212549_at	STAT5B	signal transducer and activator of transcription 5B
50	-1.1165	0.2233	0.0743	0.0366	219951_s_at	C20orf12	chromosome 20 open reading frame 12
51	-1.1147	0.2277	0.0763	0.0364	212678_at	NF1	Neurofibromin 1 (neurofibromatosis, von Recklinghausen disease, Watson disease)
52	-1.1113	0.2360	0.0807	0.0386	210852_s_at	AASS	aminoadipate-semialdehyde synthase

Table 3: ... continued

No.	D	p-values FD>0	FD>1	FD>2	probeset	gene symbol	description
53	-1.1112	0.2370	0.0807	0.0387	202962_at	KIF13B	kinesin family member 13B
54	-1.1102	0.2404	0.0813	0.0394	214724_at	DIXDC1	DIX domain containing 1
55	-1.1089	0.2450	0.0823	0.0408	206542_s_at	SMARCA2	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 2
56	-1.1085	0.2464	0.0827	0.0411	207757_at	FLJ21628	hypothetical protein FLJ21628
57	-1.1085	0.2464	0.0827	0.0411	218466_at	TBC1D17	TBC1 domain family, member 17

Table 4: Univariate and Multivariate analysis of breast cancer prognostic markers (N=417*)

	Univariate Analysis		Multivariate Analysis	
	Hazard ratio (95%CI)	p¶	Hazard ratio (95%CI)	p¶
Age (years)				
≤50 vs >50	1.055 (0.556-2.004)	0.869	0.906 (0.416-1.975)	0.8040
Size				
>2cm vs ≤2cm	2.694 (1.618-4.485)	**0.0001**	2.153 (1.235-3.755)	**0.0068**
Histological grade				
1 vs 2 vs 3	2.102 (1.461-3.024)	**0.00006**	1.446 (0.963-2.171)	0.0754
Estrogen Receptor				
Rich vs Poor	0.937 (0.671-1.307)	0.937	1.212 (0.667-2.202)	0.5275
Progesterone Receptor				
Rich vs Poor	0.536 (0.381-0.754)	**0.00034**	0.755 (0.430-1.328)	0.3300
Genomic Grade				
High vs Low	2.610 (1.833-3.717)	**0.0000001**	2.302 (1.241-4.271)	**0.0081**

* Only patients with complete information in all variables were included in the multivariate analysis (N=208)

¶ Based on Cox regression, stratified according to the datasets

Table 5: Univariate and Multivariate analysis of breast cancer prognostic markers (N=249*)

	Univariate Analysis		Multivariate Analysis	
	Hazard ratio (95%CI)	p¶	Hazard ratio (95%CI)	p¶
Age (years)				
≤50 vs >50	0.926 (0.328-2.612)	0.8840	0.807 (0.223-2.916)	0.7440
Size				
>2cm vs ≤2cm	2.002 (1.157-3.463)	**0.0130**	1.712 (0.897-3.268)	0.1030
Histological grade				
1 vs 2 vs 3	1.728 (1.128-2.647)	**0.0120**	1.071 (0.624-1.839)	0.8040
Nodal status				
Positive vs Negative	1.444 (0.836-2.493)	0.1870	1.053 (0.554-2.001)	0.8760
Estrogen Receptor				
Rich vs Poor	0.839 (0.512-1.376)	0.4860	0.982 (0.547-1.764)	0.9530
Progesterone Receptor				
Rich vs Poor	0.485 (0.291-0.806)	**0.0050**	0.751 (0.409-1.381)	0.3570
Genomic Grade				
High vs Low	3.119 (1.861-5.228)	**<0.000001**	2.147 (1.042-4.422)	**0.0380**

* Only patients with complete information in all variables were included in the multivariate analysis

¶ Based on Cox regression, stratified according to the datasets

References

1. Elston CW, Ellis IO. Pathological prognostic factors in breast cancer. I. The value of histological grade in breast cancer: experience from a large study with long-term follow-up. Histopathology 1991;19(5):403-10.

2. Elston CW, Ellis IO. Pathological prognostic factors in breast cancer. I. The value of histological grade in breast cancer: experience from a large study with long-term follow-up. C. W. Elston & I. O. Ellis. Histopathology 1991; 19; 403-410. Histopathology 2002;41(3A):151.

3. Galea MH, Blamey RW, Elston CE, Ellis IO. The Nottingham Prognostic Index in primary breast cancer. Breast Cancer Res Treat 1992;22(3):207-19.

4. Paik S, Shak S, Tang G, Kim C, Baker J, Cronin M, et al. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. N Engl J Med 2004;351(27):2817-26.

5. Robbins P, Pinder S, de Klerk N, Dawkins H, Harvey J, Sterrett G, et al. Histological grading of breast carcinomas: a study of interobserver agreement. Hum Pathol 1995;26(8):873-9.

6. Hopton DS, Thorogood J, Clayden AD, MacKinnon D. Observer variation in histological grading of breast cancer. Eur J Surg Oncol 1989;15(1):21-3.

7. Theissig F, Kunze KD, Haroske G, Meyer W. Histological grading of breast cancer. Interobserver, reproducibility and prognostic significance. Pathol Res Pract 1990;186(6):732-6.

8. Fitzgibbons PL, Page DL, Weaver D, Thor AD, Allred DC, Clark GM, et al. Prognostic factors in breast cancer. College of American Pathologists Consensus Statement 1999. Arch Pathol Lab Med 2000;124(7):966-78.

9. Singletary SE, Allred C, Ashley P, Bassett LW, Berry D, Bland KI, et al. Revision of the American Joint Committee on Cancer staging system for breast cancer. J Clin Oncol 2002;20(17):3628-36.

10. Perou CM, Sorlie T, Elsen MB, van de Rijn M, Jeffrey SS, Ross CA, et al. Molecular portraits of human breast tumours. Nature 2000;406:747-52.

11. Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, et al. Gene expression patterns of breast carcinomas distinguish tumour subclasses with clinical implications. Proc Natl Acad Science 2001;98(19):10869-74.

12. Sorlie T, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, et al. Repeated observation of breast tumour subtypes in independent gene expression data sets. Proc Natl Acad Science 2003;100(14):8418-23.

13. Sotiriou C, Neo SY, McShane LM, Korn EL, Long PM, Jazaeri A, et al. Breast cancer classification and prognosis based on gene expression profiles from a population-based study. Proc Natl Acad Sci U S A 2003;100(18):10393-8.

14. van de Vijver MJ, He YD, van't Veer LJ, Dai H, Hart AA, Voskuil DW, et al. A gene-expression signature as a predictor of survival in breast cancer. N Engl J Med 2002;347(25):1999-2009.

15. Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U, et al. Exploration, normalization, and summaries of high density oligonucleotide array probe level data. Biostatistics 2003;4(2):249-64.

16. Hedges L, Olsen I. Statistical methods for meta-analysis: Academic Press, London; 1985.

17. Korn EL, Troendle J, McShane LM, Simon R. Controlling the number of false discoveries: application to high dimensional genomic data. J Statist Plann Inference 2004;124:379-398.

18. Praz V, Jagannathan V, Bucher P. CleanEx: a database of heterogeneous gene expression data based on a consistent gene nomenclature. Nucleic Acids Res 2004;32(Database issue):D542-7.

19. Ma XJ, Salunga R, Tuggle JT, Gaudet J, Enright E, McQuary P, et al. Gene expression profiles of human breast cancer progression. Proc Natl Acad Sci U S A 2003;100(10):5974-9.

20. van 't Veer LJ, Dai H, van de Vijver MJ, He YD, Hart AA, Mao M, et al. Gene expression profiling predicts clinical outcome of breast cancer. Nature 2002;415(6871):530-6.

21. Wang Y, Klijn J, Zhang Y, Sieuwerts A, Look MP, Atkins D, et al. Gene-expression profiles to predict distant metastasis of lymph-node-negative primary breast cancer. Lancet 2005;365:671-79.

22. Ein-Dor L, Kela I, Getz G, Givol D, Domany E. Outcome signature genes in breast cancer: is there a unique set? Bioinformatics 2004;21(2):171-8.

23. Michiels S, Koscielny S, Hill C. Prediction of cancer outcome with microarrays: a multiple random validation strategy. Lancet 2005;365(9458):488-92.

24. Jenssen TK, Hovig E. Gene-expression profiling in breast cancer. Lancet 2005;365(9460):634-5.

25. Sorlie T, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, Deng S, O'Brown P, Borresen-Dale AL, Botstein D. *Proc Natl Acad Science.* 2003;100:8418-23.

26. Dai H, van't Veer L, Lamb J, He YD, Mao M, Fine BM, Bernards R, van de Vijver M, Deutsch P, Sachs A, Stoughton R, Friend S. *Cancer Res.* 2005;65:4059-66.

27. Sorlie T. *Eur J Cancer.* 2004;40:2667-75.

28. Eisen MB, Spellman PT, Brown PO, Botstein D. *Proc Natl Acad Sci U S A.* 1998;95:14863-8.

29.Therneau TM, Grambasch PM. Modeling Survival Data: Extending the Cox Model". In; 2000.

30.Loi S, Piccart MJ, Desmedt C, Haibe-Kains B, Harris A, Tutt A, Sotiriou C. Prediction of early distant relapses on tamoxifen in early-stage breast cancer (BC). *Proc Am Soc Clin Oncol*. 2005;23:6s.

31.Clarke R, Liu MC, Bouker KB, Gu Z, Lee RY, Zhu Y, Skaar TC, Gomez B, O'Brien K, Wang Y, Hilakivi-Clarke LA. *Oncogene*. 2003;22:7316-39.

32.Wilcken NR, Prall OW, Musgrove EA, Sutherland RL. *Clin Cancer Res*. 1997;3:849-54.

33.Baum M, Buzdar A, Cuzick J, Forbes J, Houghton J, Howell A, Sahmoud T. *Cancer*. 2003;98:1802-10.

34.Boccardo F, Rubagotti A, Puntoni M, Guglielmini P, Porpiglia M, Mesiti M, Rinaldini M, Paladini G, Distante V, Franchi R. *American Society of Clinical Oncology*. Orlando, Florida abstract (526); 2005.

35.Goss PE, Ingle JN, Martino S. *Proc Am Soc Clin Oncol*. 2004;22:88s.

36.Coombes RC, Hall E, Gibson LJ, Paridaens R, Jassem J, Delozier T, Jones SE, Alvarez I, Bertelli G, Ortmann O, Coates AS, Bajetta E, Dodwell D, Coleman RE, Fallowfield LJ, Mickiewicz E, Andersen J, Lonning PE, Cocconi G, Stewart A, Stuart N, Snowdon CF, Carpentieri M, Massimini G, Bliss JM. *N Engl J Med*. 2004;350:1081-92.

37.Jakesz R, Jonat W, Gnant M, Mittlboeck M, Greil R, Tausch C, Hilfrich J, Kwasny W, Menzel C, Samonigg H, Seifert M, Gademann G, Kaufmann M, Wolfgang J. *Lancet*. 2005;366:455-62.

38.Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. *Lancet*. 2005;365:1687-717.

39.Winer EP, Hudis C, Burstein HJ, Wolff AC, Pritchard KI, Ingle JN, Chlebowski RT, Gelber R, Edge SB, Gralow J, Cobleigh MA, Mamounas EP, Goldstein LJ, Whelan TJ, Powles TJ, Bryant J, Perkins C, Perotti J, Braun S, Langer AS, Browman GP, Somerfield MR. *J Clin Oncol.* 2005;23:619-29.

CLAIMS

1. A diagnostic kit or device which is a computerized system comprising :

a) a bio assay module configured for detecting gene expression for a tumor sample based on a gene set comprising at least 4 genes selected from the genes of table 3 designated as "up regulated genes in grade 3 tumors" and,

b) a processor module configured to calculate gene-expression grade index (GGI) or relapse score(RS) based on the gene expression and to generate a risk assessment for the tumor sample.

2. The kit or device according to the claim 1, wherein the genes are proliferation-related genes.

3. The kit or device according to the claim 2, wherein the gene set comprises 5 proliferation-related genes.

4. The kit or device according to any of the preceding claims 1 to 3 wherein the genes are selected from group consisting of UBE2C, KPNA2, TPX2, FOXM1, STK6, CCNA2, BIRC5, MYBL2.

5. The kit or device according to any of the preceding claims 1 to 4 which further comprises at least 4 genes selected from the genes in table 3 designated as "up regulated genes in grade 1 tumors".

6. The kit or device according to any of the preceding claims wherein the gene sequences are bounded to a solid support surface as an array.

7. The kit or device according to any of the claims 1 to 6, wherein the tumor sample is from tissue affected by a cancer selected from the group consisting of breast cancer, colon cancer, lung cancer, prostate cancer, hepatocellular cancer, gastric cancer, pancreatic cancer, cervical cancer, ovarian cancer, liver cancer, bladder cancer, cancer of the urinary tract, thyroid cancer, renal cancer, carcinoma, melanoma, or brain cancer.

8. The kit or device according to any of the claims 1 to 7, wherein the tumor sample is a breast tumor sample.

9. A method comprising the step of :

a) measuring gene expression in a tumor sample,

b) calculating gene-expression grade index (GGI) of the tumor sample by using the formula :

$$\sum_{j \in G_3} x_j - \sum_{j \in G_1} x_j$$

wherein:

x is the gene expression level of mRNA,

G1 and G_3 are sets of genes up-regulated in HG1 and HG3, respectively, and

j refers to a probe or probe set, wherein the gene set comprises or corresponds the gene set comprising at least 4 genes selected from the genes of table 3 designated as "up regulated genes in grade 3 tumors"..

10. The method according to the claim 9, wherein the tumor sample is from tissue afflicted by a cancer selected from a group consisting of breast cancer, colon cancer, lung cancer, prostate cancer, hepatocellular cancer, gastric

cancer, pancreatic cancer, cervical cancer, ovarian cancer, liver cancer, bladder cancer, cancer of the urinary tract, thyroid cancer, renal cancer, carcinoma, melanoma and brain cancer.

11. The method according to claim 10, wherein the tumor sample is a breast tumor sample.

12. The method according to the claim 11, wherein the breast tumor sample is histological grade HG2.

13. The method according to claim 11, further comprising the step of designating the breast tumor sample as low risk (GG1) or high risk (GG3) based on the GGI index obtained.

14. The method according to any of the preceding claims 9 to 13, further comprising the step of providing a breast cancer treatment regimen for a patient consistent with the low risk or high risk designation of the breast tumor sample.

15. The method according to any of the preceding claims 9 to 14, wherein the GGI includes cutoff and scale values chosen so that the mean GGI of the HG1 cases is about −1 and the mean GGI of the HG3 cases is about +1:

$$GGI = scale\left[\sum_{j \in G_3} x_j - \sum_{j \in G_1} x_j - cutoff \right]$$

16. The method according to any of the preceding claims 9 to 15, wherein the G_1 and G_3 gene sets are generated from an estrogen receptor positive population.

17. The method according to any of the preceding claims 9 to 16, which further comprises a step of designating a breast tumor sample as different subtypes within ER-positive tumors.

18. The method according to any of the preceding claims 9 to 17, which further comprises a step of designating

a tumor sample as a subtype to be submitted to a different treatment than the other subtype.

19. The method according to any of the preceding claims 9 to 18, which is combined to oestrogen receptor and/or progesterone receptor gene expression detection.